

## Index 1: Subject

abundance, of Sciomyzidae, 174  
 adaptations, to habitat, 171  
 adult behavior  
     copulatory posture, 128–129  
     diapausing, 107, 122  
     dispersal, experimental studies, 374  
     emergence, 120, 202–203  
     flight period, 109, 110, 119, 175  
     mating  
         assault type, 126  
         cues, 126, 128  
         genetic compatibility, 269  
         preference, 126  
         pre-mating, 128  
         trophallaxis, 126–128  
     migration, 120  
     mimicry, 204  
     pharate, 107  
     seasonal dynamics, 110  
     species richness and diversity, 110  
 adult physiology  
     hypertrophy, of fat bodies, 143  
     pre-oviposition period, 148  
     protein, in diet, 151  
     reproductive status, females, 123  
     thermoregulation, 120  
 anatomy, of gastropods, 13  
 ancestral forms, of behavior, 33  
 assemblages, of Sciomyzidae, 173, 175  
 Baltic amber, 338  
 behavioral attributes  
     adults, 34  
     egg, 34  
     larvae, 34–35  
     puparia and pupae, 35  
 behavioral groups, 26, 29, 30, 31, 40, 72, 77  
 biological control (= biocontrol), 21  
     acclimatization, 405  
     *Achatina fulica*, 412–413  
     approaches  
         augmentation, 397, 403  
         ecosystem service, 397  
         inoculative release, 397, 400, 405–406  
         integrated control, 399

biological control and ecological theory, 26, 398  
 biological control, of millipedes, 45  
 cryopreservation, 407  
 dispersal along rivers, 191  
 experimental studies, 400  
 greenhouse situations, 415  
 habitat exploitation, 405  
 molluscicides, impact on  
     Sciomyzidae, 410  
 natural enemies, reviews, 23, 397–398  
 natural prey, 399  
 need for biocontrol  
     increase in schistosomiasis, 397  
     resistance to antihelminthics, 397  
 optimal rearing conditions, 409–410  
 population dynamics, 189  
 potential impact, of Sciomyzidae, 416  
 rates of increase, of Sciomyzidae, 408–409  
 requirements for success, 399–400  
 research needs, 398, 399, 401  
*Salticella fasciata*, as biocontrol agent, 411–413  
*Schistosoma*, life cycle, 398  
 sex, prediction of, 407–408  
 sex ratio, 407  
 snail egg masses, 409  
 snail infection, by trematodes, 408  
 South Africa, trials, 403–404  
*Subulina octona*, 414  
 terrestrial snails, review, 410  
 bivalve molluscs, 30  
 brain, of larvae, function, 106, 124, 125, 159  
 categories of behavior, 26  
 co-adaptation, 126  
 competition  
     interspecific, 94  
     intraspecific, 36, 94, 190  
     niche exploitation, 172–173  
 conservation value, 177  
 cross-mating studies, 269  
 dead molluscs  
     clams, 29  
 snails 27, 68  
 development studies  
     affect of temperature on size and weight, 148, 155  
     duration of egg, larval, pupal stages, 142  
     phenological groups, 140  
     rates of development, 142  
     rearing conditions, 136–137  
     temperature thresholds, 143  
     thermal constants, 143  
     under controlled conditions, 140  
 diapause  
     estival, 109, 120  
      facultative, 119  
     hibernal, 120  
     obligatory, 119  
     photoperiodic response curve, 120  
     reproductive, 109  
 dispersal, of flies, 28, 103, 168, 190, 191  
 DNA, 270  
 dry-season survival, 115  
 Eco-Groups, 26  
 ecological assemblage, of snails, 97  
 ectoparasite, 10, 29  
 ectoparasitoid, 39  
 eggs  
     as alternate hosts of pests, 180  
     hatching stimuli, 148  
     in temporary aquatic habitats, 148–150  
     mortality, 192  
     number produced, 134, 143  
     on snails, natural populations, 197  
     submergence, 135, 150  
     thermal reaction norm, 150  
     water loss, 213  
 egg masses, of snails, 39, 63, 75  
 encapsulation, of larvae, 78, 100, 132, 185–186  
 endoparasite, 10, 29  
 endoparasitoid, 12, 45  
 evolution  
     cladistics  
         cladogram/behavioral groups, 387  
         cladogram/immature morphology, 385

- cladogram/sciomyzids-molluscs, 26, 50, 391–392  
 co-evolution, 392  
 concordance, phylogeny/behavior, 394–395  
 explosive speciation, 115, 374–377  
 extinction, 374  
 marine origin, 379–380  
 niche widths, 381  
 non-adaptive characters, 394, 395  
 parallel evolution, 374, 382, 383–384  
 residual adaptations, 392  
 scenarios, parasitoids  
     Diptera, 373  
     Hymenoptera, 373  
     Sciomyzidae, 36, 374, 386  
 selective pressure, 379  
 sister-group comparisons, 376–377
- fingernail clams, 40, 58, 75  
 feeding by larvae  
     anatomical part, 72  
     biomass, of snails killed, 54, 58, 59  
     coefficient, of food for growth, 73, 88  
     commensal feeding, 29, 30, 76, 187  
     competitive exclusion, 100, 102  
     consumption capacities, 85  
     curves  
         consumption, of snails, 59  
         nutrition, 62  
         predation, 59  
     energy returned, from feeding, 61, 87–88  
     factorial laboratory trials, 95  
     feeding, during winter, 106  
     field enclosure studies, 96  
     food consumption curves, 154  
     food consumption rate and growth rate, 77, 91, 94  
     foraging strategies, 92, 94  
     gregarious or multiple infestations, 76, 99–100  
     in gastropod mouth, 187  
     physiological state of host/prey, 72  
     prey  
         choice and consumption, 141  
         choice tests, 56  
         density, 153–155  
         specificity, experiments, 85  
     relative sizes, prey/larva, 76, 85  
     resource partitioning, by larvae, 98, 125, 173  
     salivary gland toxin, 75, 184  
     sites, categorized, 73–75  
     solitary feeding, 76, 99  
     subsurface foraging, 75, 79–81  
     time spent, 72–73  
     unfed larvae, 95
- functional response, 58, 93, 94, 133, 154, 155  
 ghost of competition past, 98, 392  
 Glöger's Rule, exception to, 200  
 guilds, 22, 28, 30, 36, 97, 98  
 gut contents, analyses of, 23, 25, 44, 75–76
- habitats  
     disturbed, 171–172  
     ephemeral wetland, 39, 147, 162, 176  
     farm, case study, 175–177  
     horizontal distribution, 169  
     hydrological regimes, 177  
     ice cover, 106  
     ideal, 171  
     insolation of, 120, 172  
     intertidal zone, 169  
     moisture, influence of, 115  
     quantitative studies, of snails, 163  
     seasonal studies, 162–163  
     shoreline, 38, 40, 56, 172  
     spatial heterogeneity, 177  
     thermal waters, 170  
     water quality, 169–170
- host/prey finding  
     by microhabitat location, 67  
     by search for host/prey, 67, 69
- ice-nucleation, in larvae, 106  
 immigrant snails, 63
- karyotypes, 267–268
- labile behavior, 33–34, 40, 75, 76  
 larval behavior  
     air in gut, 187  
     ambush, 69  
     anterior spiracles, function, 264  
     dispersal, 101, 187  
     dispersal, experimental studies, 374  
     freezing tolerance, 105  
     swimming behavior, 187  
     wandering phase, 103, 106, 125
- life cycles, types of, 103
- malacophagy, overview, 381, 382  
 malacophagy, defined, 26  
 marine molluscs, 29  
 mesoparasite, 10, 12  
 mesoparasitoid, 39, 75  
 microhabitats  
     preferences, 126  
     surface tension, of water, 170  
     watery and earthy, 161
- molecular systematics  
     *Sepedon fuscipennis* group, 194, 268–269  
     *Tetanocera*, 374, 382, 383
- morphology, adults  
     aedeagal filaments, 200  
     cochleate vesicle, 199, 200, 278  
     color forms, 120, 200  
     corpora allata, 199  
     female abdomen, 279  
     labellar hooks, 204, 259  
     palpi, loss of, 200  
     ptilinum, 160, 201–204, 273  
     tibial comb, 211  
     wings, absence, 275
- morphology, eggs  
     aeropyles, 213  
     chorion structure, 214  
     classification, by structure, 213–214  
     plastron, 212
- morphology, larvae  
     anterior spiracles, number papillae, 291–292  
     cibarial ridges, 2, 29, 74, 99, 247  
     comparative tables of terminology, 281  
     ecomorphological classification, 223–224  
     fat bodies, 158  
     habitat-adaptive classification, 228  
     indentation index, of pharyngeal sclerite, 282  
     internal anatomy, 217  
     Keilin organs, 222, 282  
     Malpighian tubules, 258, 283, 287  
     oral grooves, 2, 29, 99  
     postanal portion, segment XII, 228
- morphology, puparia  
     calcareous septum, 106, 187, 251, 267  
     false chamber, 251  
     posterior spiracles, elevation, 252–255  
     tracheal trunks, 247
- multivoltine species, 105
- natural enemies, as limiting factors, 180  
 numerical response, 25, 88
- overwintered adults  
     experiments, in screen cages, 121, 122, 192  
     reproductive capacity, 143
- overwintered larvae  
     as free larvae, 157  
     as third-instar larvae, 158  
     collections of, 147, 158  
     parasitoid Hymenoptera, 181
- overwintered pupae, 147  
 overwintering, 105, 106, 120  
 primitive pattern, 125

## 500 INDEX 1: SUBJECT

- oviposition
  - curves, 134
  - on egg masses, 130
  - on snail flesh, 68–69, 130
  - on snail shells, 68, 130, 132
  - on snails, experiments, 132
  - on substrate, 130, 132
  - sites, 65, 132
- parasitoid Hymenoptera
  - families of, 180
  - field studies, 180–181
  - host recognition, 267
  - laboratory experiments, 180, 181
  - patchy food resources, 186 *see* snails, dispersion
  - phenological diversity, evolution of, 125, 126
  - population
    - emergence abundance, productivity, 174
    - estimation, 176, 190
    - experimental design, of sampling, 197
    - functional response, 93, 94, 155
    - genetics, 194
    - model, 192
    - most dense population, 195
    - natural populations, 192
    - net production, 189
    - rate of increase, 192
    - tracer snails, 195
  - predator–prey dynamics, 152
  - pre-pupal stage, 103
  - pre-pupariation pause, 158
  - pupae
    - diapause, 109
    - duration, 121
    - respiration, 248, 266
  - pupariation site, 157, 159, 162, 257
  - puparium, phases within, 159–160
  - puparium, in shells, 187–188, 249–252
  - quiescence, 105
  - r* and *K* strategies, 67
  - recapture of, adults, 105, 120, 122, 189, 191, 192
  - searching, by larvae, 69, 70
  - seasonality, in tropics, 116–117
  - slugs
    - host preference, 414–415
    - mucus, 185
    - radula, 184
  - snails
    - abundance, 20
    - alarm pheromones, 67, 183
    - brain, 72
    - database of gastropods, 18, 162
    - dispersion, 20, 62, 78, 169
    - ecological assemblage, 380
    - estivating, 19, 68, 100
    - extrapallial fluid, 74, 265
    - hemocoel, 73, 74
    - hemolymph, 73
    - migration, 18, 20, 115
    - mortality rates, 93
    - movement, 18
    - mucus, 57, 71, 95, 184–185
    - mucus trail, 24, 67, 70–71
    - operculate, 29, 39, 41, 51, 56, 151, 170, 183–184
    - pallial cavity, 101
    - populations, 98, 113, 117
    - prey avoidance behavior, 95, 184
    - radula, 184
    - stranded, 18, 40, 51
    - survival, of larvae, 54, 79, 189, 190
    - switching phenomenon, 62
  - synoecological indexes and statistical tests, 175
  - taxonomy and nomenclature
    - cladistics, 324–327, 350
    - comparison of suprageneric classifications, 341
    - discussion of Sciomyzoidea, 339–341
    - earliest available family group name, 338
    - historical, suprageneric classifications, 341–342
    - Phaemyiidae/Sciomyzidae, 348
    - segregation, of *Sepedon*, 331
    - Sepedon*, suprageneric position, 345–348
    - Tetanura*, suprageneric position, 345
  - trophic guilds, 41
  - tropical insects, dormancy, 119
  - univoltine species, 105, 125
  - vegetation, influences in habitats, 170–171, 174, 177
  - voltinism, 105
  - wandering-overwintering, distinction, 159
  - wasteful feeding, by larvae, 73, 74, 84, 89, 93, 96
  - water
    - depth, 54, 82, 94, 96
    - film, 62
  - wet/dry seasonal phenomena, 114
  - zoogeography
    - distributional patterns, Palearctic, 356
    - maritime distributions, 358
    - Nearctic/Palearctic dispersal, 363
    - Neotropical, 359
    - Palearctic/Afrotropical dispersal, 364
    - vicariance, 361

## Index 2: Genera and species cited other than Sciomyzidae, Phaeomyiidae, and Mollusca

- Ablabesmyia* Johannsen, 30
- Ablattaria* Reitter, 24
  - laevigata* (Fabricius), 24
- Aclerixenous* Frauenfeld, 251
- Aeschna* Fabricius, 25
- Alisma* L., 130
- Anax* Rambur, 25
- Angioneura cyrtoneurina* (Zetterstedt), 29, 241
- Anopheles* Meigen, 170
- Antilocnus* Stål, 22
- Aphaereta* Förster, 180
- Arundo donax* L., 177
- Aspilota* Förster, 180
- Atractodes* Gravenhorst, 180
- Aulophorus furcatus* (Müller), 82
- Baeoconus bicolor* Saether, 30
- Bellardia* Robineau-Desvoidy, 44
- Belostoma flumineum* Say, 22
- Bidens* L., 130
- Blaesoxiphia* Loew, 376, 377
  - beamери* (Hall), 46
- Calliphora* Robineau-Desvoidy, 44, 259
  - vicina* Robineau-Desvoidy, 159
- Callitricha* L., 1
- Caltha* L., 258
- Carabus* (*Procerus*) Dejean, 23
- Cardamine* L., 258
- Carex* L., 255
  - lacustris* Willdenow, 174
  - stricta* Lambert, 174
  - vesicaria* L., see Fig. 10.2
- Chaetogaster limnaei* Baer, 417
- Chironomus acutus* Goetghebuer, 30
  - niveipennis* Fabricius, 29
- Chortoicetes terminifera* Walker, 103
- Chrysops* Meigen, 258
- Cobergius* Barnes, see Table 15.14
- Coelopa frigida* (Fabricius), 341
- Coenosia agromyzina* (Fallén), 44
- Colopha praecusta* Eschscholtz, 24
- Cordilura* Fallén, 241
- Corixa* Geoffroy, 22
- Cremifania* Czerny, 341
- Cricotopus sylvestris* (Fabricius), 44
- Cryptochironomus* Kieffer, 29, 44
- Cyprhus* Fabricius, 23
- Diaphanes* Motschulsky, 24
- Dindymus* Stål, 22
  - pulcher* Stål, 22
  - rubiginosus* var. *sanguineus* (Fabricius), 22
  - versicolor* (Herrick-Schaffer), 22
- Diplonychus annulatus* Fabricius, 22
  - rusticum* Fabricius, 22
- Drilus* Olivier, 24
  - mauritanicus* Lucas, 24
- Drosophila melanogaster* Meigen, 209, 267
- Dryomyza* Fallén, 282, 375, 377, 418
- Dytiscus marginalis* L., 25
- Eginia* Robineau-Desvoidy, 44
  - ocypterata* (Meigen), 46
- Entomophthora* Fresnius, 178
- Equisetum* Prôle, 161
  - arvense* L., 167
- Eriplanus* Förster, 180
- Eukiefferiella aencyla* Svensson, 30
- Euleia heraclei* (L.), 251
- Eupteronotus* Kurdjumov, 180
- Eurychaeta muscaria* (Meigen), 410
- Eurytocus* Steyskal, 375
- Fasciola* L., 263, 397, 404, 408, 409, 417
  - gigantica* Cobbold, 400, 405
  - hepatica* L., 400, 407, 408
- Fontinalis antipyretica* L. ex Hedw., 263
- Gambusia* Poey, 182
- Glossina brevipalpus* (Newstead), see Fig. 9.45
- Glyptotendipes paripes* (Edwards), 30
- Helosciomyza* Hendel, 349, 375
- Hemiteles* (Meyrick), 180
- Heterocheila* Rondani, 341
- Hiltonius* Chamberlain, 46
- Hirsutella citriformis* Speare, 178
- Hybomitra* Enderlein, 26
  - schineri* Lyneborg, 182
- Hydrophilus* Müller, 25
- Iulus sabulosus* L., 46
- Lamprophorus tenebrosus* (Walker), 24
- Lampyris noctiluca* (L.), 24
- Lemna* L., 1, 424
  - trisulca* L., 130
- Limnogeton sieberi* Mayr, 22
- Limnophora* Robineau-Desvoidy, 44
- Limosina* Macquart, 46
- Lucilia sericata* (Meigen), 159
- Luciola cruciata* Motschulsky, 24
  - lateralis* Motschulsky, 24
- Macquartia chalconota* (Meigen), 187
- Mastrus* Förster, 180,
- Megaselia* Rondani, 46, 375
  - (Aphiochaeta) equitans* Schmitz, 46
  - aqualis* (Wood), 30, 63
  - ciliata* (Zetterstedt), 30, 63
  - elongata* (Wood), 46
  - fuscinervis* (Wood), 29, 375
  - tertia* Brues, 30
- Melinda viridicyanea* (Robineau-Desvoidy), 29
- Meromyza* Meigen, 241
- Mesoleptus* Gravenhorst, 180, 181
  - ripicola* Thomson, 180
- Mesovelia mulsanti* White, 181
- Metroclemens* Wulp, 30
  - knabi* Coquillett, 124
- Molinia* Schrank, 176
- Musca* L., 251
  - domestica* L., 100, 178, 271
  - fergusoni* Johnson et Bancroft, 251
  - umbrarum* L., 418
- Natalimyza* Barraclough et McAlpine, 341
- Naucoris* Geoffroy, 22
- Neohelina* Malloch, 46
- Notochaeta aurata* Walker, 44
  - (=*Lepidodexia* (*N.*) *cognata* Walker)
- Notonecta* L., 182

## 502 INDEX 2: GENERA AND SPECIES

- Ocypus olens* Müller, 25  
*Oedoparena* Curran, 375, 377  
*Ommatoiulus* Latzel, 45  
 moreleti (Lucas), 44, 45, 46  
 oliveirae (Verhoeff), 45  
*Oncocladosoma* Jeekel, 46  
*Onesia* Robineau-Desvoidy, 44  
*Oniscus asellus* L., 411  
*Orconectes obscurus* (Hagen), 185  
*Orthizema* Förster, 180  
*Orygma* Meigen, 200, 251  
 luctuosum Meigen, 380
- Parachironomus limnaei* Guibé, 375  
 varus (Goetghebuer), 30  
 varus *limnaei* Guibé, 29  
 varus varus (Goetghebuer), 29, 375
- Paratrichina* Motschulsky, 25  
*Parnassia palustris* L., 259  
*Pericallimyia greataedi* Zumpt, 26  
*Phaenocarpa* Förster, 180  
 antichaetae Fischer, 180  
*Phaenopria popei* Muesebeck, Plate 4  
*Phasmarhabditis hermaphrodita* (Schneider), 416  
*Phausis splendidula* (LeConte), 24  
*Pheidole megacephala* (Fabricius), 181  
*Pheidologeton* Mayr, 25  
*Phormia regina* (Meigen), 248, 264  
*Phosphuga atrata* (L.), 24  
*Photuris* LeConte, 24  
*Phragmites* Trinius, 170, 171  
*Phygadeuon* Gravenhorst, 180, 181  
*Pinus pinaster* Soland, 45  
*Plastophora* Brues, 46  
 juli (Brues), 46  
*Pollenia* Robineau-Desvoidy, 44
- Polypedilum anale* Freeman, 30  
 kibatiense Goetghebuer, 30  
*Polytocus* Lamb, 349, 375  
*Potamogeton* L., 427  
*Pseudocella* Filipjev, 178  
*Pterostichus cupreus* L., 416  
*Pyraustomera lucifera* (Melsheimer), 24
- Rheotanytarsus* Bause, 30
- Salicornia* L., 168  
*Saltella* Robineau-Desvoidy, 251  
*Salvinia* Adanson, 24  
*Sarcophaga* Meigen, 44, 101, 187, 396  
 (*Heteronychia*) *penicillata* Villeneuve, 29  
 (*Pierretia*) *iulicida* Pape, 46  
 (*S.*) *meiosiflosia* Pape et al., 29  
 (*Sarcorohdendorfia*) *megafloria* Pape et al., 29  
*bullata* Parker, 78  
*muscaria* Meigen, 410  
*Scaphinotus* Dejean, 23  
*Schistosoma* Weinland, 22, 56, 58, 93, 260, 397, 398, 403, 404, 405, 406  
*intercalatum* Fischer, 408  
*japonicum* Katsurada, 397  
*Sciogriphoneura* Malloch, 349  
*Selasia unicolor* (Guérin), 24  
*Solenopsis* Westwood, 25  
*Spalangia* Latreille, 180  
*Sphagnum* L., 255  
*Spilomicrus* Westwood, 181  
*Spirobolomyia* Townsend, 46  
*flavipalpis* (Aldrich), 46  
*singularis* (Aldrich), 46
- Spirobolus* Brant, 46  
*Spirogyra* Link, 258, 259
- Tabanus* L., 26  
*Theroscopus* Förster, 180  
*Thienemannimyia festiva* (Meigen), 44  
*Toxorhynchites rutilus* (Coquillett), 44  
*Trichogramma* Westwood, 179, 180, 192  
*japonicum* Ashmead, 180  
*julianoi* Platner et Oatman, 179  
*semblidis* (Aurivillius), 179  
 sp. near *californicum* Nagaraja et Nagarkatti, 179
- Trichogrammatoides* Giraud, 179  
*Trichomalopsis* Crawford, 180  
*Trichopria* Ashmead, 181, 427  
*atrichomelinae* Muesebeck, 181  
*drosophilae* Ashmead, 267
- Trixoscelis puncticornis* (Becker), 204  
*Tropisternus* Solier, 182  
*Tylobolus* Cook, 46  
*Typha* L., 130, 170
- Ula macroptera* (Macquart), 44  
*Urolepis* Walker, 180
- Vespa* L., 28
- Wandolleckia* Cook, 29, 375  
*Wyeomyia smithii* (Coquillett), 124
- Xenochironomus canterburyensis* (Freeman), 30  
*Xenosciomyza* Tonnoir et Malloch, 349

## Index 3: Group names cited other than Sciomyzidae, Phaeomyiidae, and Mollusca

- Acalyptatae, 126, 205, 271, 338–342, 375, 376, 428
- Agromyzidae, 173, 271, 428
- amphibians, 21
- Anthomyiidae, 26, 271
- Anthomyzoidea, 338
- ants, 25, 181
- Aphrophoridae, 259
- Aschiza, 266, 271, 375
- Asilidae, 182
- bacteria, 21, 22, 106, 178, 241, 247, 379, 416
- barnacles, 375, 377
- beetles, 21, 23, 24
- Belostomatidae, 22, 172
- birds, 21, 182, 183
- Bombyliidae, 259, 373, 376
- Brachycera, 175, 199, 271
- brachylaemid flatworms, 21
- Braconidae, 180, 181, 187
- Calliphoridae, 26, 28, 29, 44, 156, 157, 159, 160, 172, 375, 376
- Calyptatae, 211, 271, 375
- Canaceidae, 169
- Cantharidae, 25
- Carabidae, 22, 23, 173, 376, 415
- Celyphidae, 375
- Cercopidae, 376
- Chamaemyiidae, 204, 341, 375
- Chironomidae, 26, 29, 30, 375, 376, 427, 496
- Chloropidae, 174, 241, 376
- Chrysomelidae, 179, 212
- Clusiidae, 338
- Coccinellidae, 179
- Coelopidae, 126, 273, 338, 339, 341, 375, 380
- Coleoptera, 21, 22, 172, 376
- Conopidae, 202, 271, 376
- copepods, 44
- Corixidae, 22
- Corydalidae, 179
- Cremifaniidae, 339, 340, 342
- Culicidae, 44, 150, 421
- Culicomorpha, 375
- Curculionidae, 358
- Cyclorrhapha, 199, 206, 211, 230, 240, 241, 259, 264, 266, 267, 271, 279, 395
- decapods, 183
- delphacid, 178
- Dermaptera, 376
- Diapriidae, 181, 267, 427
- Diopsidae, 259, 338
- diplopods, 31, 34, 381
- Diplopoda, 44, 46, 348, 375, 376, 382
- Dolichoceridae, 341
- Dolichopodidae, 44, 173, 199, 205
- Drilidae, 22, 23, 24, 75, 173, 376
- Drosophilidae, 156, 159, 160, 206, 209, 240, 251, 376, 394
- Dryomyzidae, 26, 126, 243, 245, 271, 273, 282, 338, 340, 341, 349, 375, 377, 380, 383
- Dryomyzinae, 342, 344
- Dytiscidae, 25, 179, 427
- earthworms, 44, 258, 259, 375, 411
- Echinostomatidae, 22
- Eginiidae, 44, 46, 375
- Empididae, 28, 182
- Enchytraeidae, 44
- Ephydriidae, 26, 97, 174, 179, 202, 240, 271, 273, 376, 427, 430
- Eurychoromyiidae, 341, 375
- Fanniidae, 26, 375, 376
- fish, 21, 182, 397
- fungi, 21, 28, 106, 183, 379
- Glossinidae, 156, 159, 160
- Gordidae, 178
- Haplostomata, 339
- Helcomyzidae, 273, 339–341, 349, 350, 375, 376, 380
- Helomyzinae, 338, 341
- Heleomyzidae, 28, 273
- helminths, 21, 22
- Helomyzoidea, 338
- Helosciomyzidae, 273, 340, 341, 349, 375
- Helosciomyzinae, 340, 341, 349,
- Hemiptera, 22, 376
- Heterocheilidae, 341, 375, 376, 380
- Hippoboscidae, 202
- Hippoboscoidea, 376
- Holometopa, 342
- Hydrophilidae, 25, 172, 182
- Hymenoptera, 11, 125, 173, 187, 376, 381, 395
- Hypocreales, 178
- Ichneumonidae, 180, 181, 187
- Isopoda, 411
- Iulidae, 44
- Keroplatidae, 373
- Lampyridae, 22, 23, 24, 97, 173, 185, 376
- Lauxaniidae, 251, 273, 341, 375
- Lauxanioidea, 204, 212, 339, 341, 343, 375, 376, 380
- leeches, 397
- Lepidoptera, 373, 395
- Leptosomatidae, 178
- Lonchaeidae, 260
- Lonchopteridae, 271
- lung worms, 21
- mammals, 21
- Marginidae, 376, 382
- Megamerinidae, 338–341, 348
- micro-organisms, 2, 21, 29, 74, 99
- Micropezidae, 273
- microsporidia, 179
- millipedes, 44, 45, 46, 220, 249, 320, 348, 374, 375, 386, 391
- mites, 327
- mosquitoes, 170, 395, 397
- Muscidae, 156, 157, 160, 174, 241, 375, 376, 413, 425, 431, 435, 447
- Muscidi, 342
- Muscii, 342
- Muscini, 342
- Muscoidea, 343, 346
- Muscomorpha, 125, 209, 260, 271, 278, 338, 340, 341, 385

## 504 INDEX 3: GROUP NAMES

- Naididae, 44, 82, 382
- Natalimyzidae, 341, 376
- Nematocera, 199, 230
- Nematoda, 178
- nematode, 416
- Nemestrinoidea, 376
- Neottiophilidae, 338
- Neuroptera, 373, 376
- Notonectidae, 172
- Nycteribiidae, 160, 199, 201
- Odonata, 25, 182
- Oligochaeta, 82, 376
- oligochaete, 29, 31, 40, 41, 44, 47, 75, 117, 139, 165, 235, 238, 348, 374, 375, 377, 381, 382, 383, 386, 387, 411, 413, 416, 417, 434, 435
- Opomyzidae, 241, 425
- Opomyzidae, 341, 345
- Opomyzoidea, 376
- Orthoptera, 182
- Orthorrhapha, 125, 230, 251
- Pallopteridae, 260
- pathogens, 137, 178
- Phoridae, 22, 26, 28, 29, 30, 46, 64, 97, 172, 183, 271, 373, 376, 379, 381, 413, 416
- Phthiraptera, 376
- Piophilidae, 28, 97, 338, 381
- Piophilini, 342
- Pipunculidae, 183, 376
- Platypezidae, 271
- protozoans, 21
- Pseudococcidae, 376
- Psilidae, 241, 259, 338, 425
- Psychodidae, 26, 28
- psyllid, 178
- Pteromalidae, 180, 187
- Pyralidae, 179
- Pyrgotidae, 183
- Pyrrhocoridae, 22
- reptiles, 21
- Rhinophoridae, 26, 373, 375, 376
- Rhinophorinae, 241
- Rhinotoridae, 376
- Rhopalocera, 421
- rice-stem borer, 197
- Risidae, 376
- Ropalomeridae, 211, 338, 339, 375
- Sarcophagidae, 26, 28, 29, 39, 44, 46, 78, 97, 156, 159, 160, 172, 373, 375, 376, 381, 413, 441
- Scathophagidae, 175, 241, 271, 425
- Scatomyzini, 342
- Schizophora, 160, 199, 202, 251, 266, 278, 339, 348, 379
- Sciomyzoidea, 126, 200, 204, 208, 209, 211, 212, 270, 282, 338–341, 343, 346, 348, 350, 375, 376, 380, 385, 425
- Sciomyzoinea, 339, 340, 348
- Sciomyzomorphae, 338
- Sepsidae, 26, 200, 204, 211, 230, 251, 338, 339, 341, 342, 347, 376, 380, 395
- Sepsoidea, 338
- Sialidae, 179
- Silphidae, 22, 24, 173, 376
- Siphonaptera, 376
- Sphaeroceridae, 26, 28, 46
- Sphaeroceroidea, 376
- spiders, 182, 205, 376
- Spirobolidae, 46
- Staphylinidae, 22, 173, 376
- Stratiomyidae, 179, 251, 429
- Stripsiptera, 376
- Stylommatophora, 51
- stylommatophorans, 379
- Syrphidae, 16, 162, 175, 176, 177, 179, 279, 373, 432
- Tabanidae, 26, 179, 199, 258
- Tachinidae, 183, 187, 373, 374, 376, 429
- Tanypezidae, 240, 241
- Tanypezinae, 342, 345
- Tenthredoidea, 173
- Tephritidae, 156, 199, 205, 259, 273, 374, 408, 428, 430
- Thyreophoridae, 338
- Tipulidae, 44, 150, 174, 175
- trematodes, 22
- Trichogrammatidae, 179, 427
- Trichoptera, 165, 259, 427
- Trixoscelididae, 204, 376
- vertebrates, 21, 22, 375
- viruses, 21
- yeasts, 241, 247, 379

## Index 4: Suprageneric names in Sciomyzidae and Phaeomyiidae (see especially Table 15.14)

- Chaetomacerini, 342  
Ctenulina, see Table 15.14  
Ditaeniinae, 342  
Euthycerina, 342  
Euthycerinae, 342  
Euthycerini, 342  
Huttonininae, 314, 331, 341, 348, 349, 350, 361, 362, 379, 382, 418, 422  
Huttoninidae, 341, 379  
Huttoninini, 291, 382  
Neolimniinae, see Table 15.14  
Oidematopsini, 342
- Pelidnopterinae, 341  
Phaeomyiina, see Table 15.14  
Phaeomyiinae, 338, 341, 348, 422  
Prosochaetini, 331, 349  
Renocerinae, 329, 341  
Salticellina, 342  
Salticellinae, 2, 12, 35, 41, 67, 100, 206, 211, 273, 338, 358, 362, 383, 385, 418, 423  
Sciomyziden, 342  
Sciomyzides, 338, 341, 345, 418  
Sciomyzina, see Table 15.14  
Sciomyzinae, 2, 287, 361, 362, 376, 383, 385, 418  
Sciomyzinæ (“subsectio”), 291  
Sepedonidae, 339, 342
- Sepedonina, 342  
Sepedoninae, 118, 338, 341, 425  
Sepedonini, 342  
Sepedontini, 342  
Tetanoceratidae, 342  
Tetanoceratoidea, 339, 342  
Tetanoceriden, 342  
Tetanocerina, 342  
Tetanocerinae, 291, 329, 342, 418  
Tetanocerites, 338, 342  
Tétanocéroides, 339  
Tetanurinae, 342, 345