

---

# THAILAND RED DATA : MAMMALS, REPTILES AND AMPHIBIANS

---

compiled by

Jarujin Nabhitabhata  
Tanya Chan-ard



Office of Natural Resources and  
Environmental Policy and Planning



United Nations Development  
Programme

2005

**First published :** November 2005  
by Office of Natural Resources and  
Environmental Policy and Planning (ONEP),  
Thailand.

**ISBN :** 974-9929-85-3

This publication is financially supported by ONEP and may be reproduced in whole or in part and in any form for educational or non-profit purposes without special permission from ONEP, providing that acknowledgment of the source is made.

No use of this publication may be made for resale or for any other commercial purposes.

**Citation :** Nabhitabhata J. and T. Chan-ard. 2005.  
*Thailand Red Data : Mammals, Reptiles  
and Amphibians.*  
Office of Natural Resources  
and Environmental Policy and Planning,  
Bangkok, Thailand. 234 p.

**Authors :** **Jarujin Nabhitabhata**  
*Director, Natural History Museum,  
National Science Museum*

With 35 years of working experience in the study of vertebrate and invertebrate taxonomy in Thailand, Dr. Jarujin has also participated in the development of master plans for several national parks in Thailand. In addition to experience on terrestrial ecosystems and entomology, Dr. Jarujin also carry out study on the effects of various projects.

**Tanya Chan-ard**  
*Technical Official 6, Natural History Museum,  
National Science Museum*

Mr. Tanya is an expert in reptile and amphibian study and research. With over 20 years of experience, Mr. Tanya is very knowledgeable and has come across a great deal of reptiles and amphibians species samples from all across Thailand. He is presently conducting surveys and collecting reptile and amphibian samples continuously.

**Available from :** Biological Diversity Division  
Office of Natural Resources and Environmental  
Policy and Planning  
Ministry of Natural Resources and Environment  
60/1 Rama VI Rd. Bangkok 10400  
THAILAND  
Telephone (66) 2265 6638-39  
Facsimile (66) 2265 6638  
Website: <http://chm-thai.onep.go.th>  
E-mail: [chm\\_thai@onep.go.th](mailto:chm_thai@onep.go.th)

**Designed & Printed :** Integrated Promotion Technology Co., Ltd.  
Telephone (66) 2158 1312-6  
Facsimile (66) 2158 1319





- |                        |                              |                       |                         |
|------------------------|------------------------------|-----------------------|-------------------------|
| 1. Mae Hong Son        | 20. Nakhon Sawan             | 39. Udon Thani        | 58. Chachoengsao        |
| 2. Chiang Mai          | 21. Uthai Thani              | 40. Sakon Nakhon      | 59. Chon Buri           |
| 3. Chiang Rai          | 22. Chai Nat                 | 41. Nong Khai         | 60. Rayong              |
| 4. Lamphun             | 23. Suphan Buri              | 42. Nakhon Phanom     | 61. Chanthaburi         |
| 5. Lampang             | 24. Ang Thong                | 43. Mukda Han         | 62. Trat                |
| 6. Phayao              | 25. Sing Buri                | 44. Kalasin           | 63. Chumphon            |
| 7. Nan                 | 26. Lop Buri                 | 45. Khon Kaen         | 64. Ranong              |
| 8. Phrae               | 27. Saraburi                 | 46. Chaiyaphum        | 65. Surat Thani         |
| 9. Uttaradit           | 28. Phra Nakhon Si Ayutthaya | 47. Maha Sarakham     | 66. Phangnga            |
| 10. Tak                | 29. Nakhon Pathom            | 48. Roi Et            | 67. Phuket              |
| 11. Kanchanaburi       | 30. Nonthaburi               | 49. Yaso Thon         | 68. Krabi               |
| 12. Ratchaburi         | 31. Pathum Thani             | 50. Umnajareon        | 69. Nakhon Si Thammarat |
| 13. Phetchaburi        | 32. Nakhon Nayok             | 51. Nakhon Ratchasima | 70. Trang               |
| 14. Prachuap Khirikhan | 33. Samut Songkhram          | 52. Buri Ram          | 71. Phatthalung         |
| 15. Sukhothai          | 34. Samut Sakhon             | 53. Surin             | 72. Satun               |
| 16. Phitsanulok        | 35. Bangkok                  | 54. Si Sa Ket         | 73. Songkhla            |
| 17. Kamphaeng Phet     | 36. Samut Prakan             | 55. Ubon Ratchathani  | 74. Pattani             |
| 18. Phichit            | 37. Loei                     | 56. Prachin Buri      | 75. Yala                |
| 19. Phetchabun         | 38. Nong Bua Lumpoo          | 57. Srakaew           | 76. Narathiwat          |

# FOREWORD

In 1996, the Office of Natural Resources and Environmental Policy and Planning (ONEP) organized a meeting with a major purpose to assess the status of the biological resources in Thailand, including mammals, birds, reptiles, amphibians, and fish. The IUCN Red List Categories was used as a guiding document. The initial IUCN List, which provided a set of criteria to evaluate the extinction risk of thousands of species and subspecies, was created under the 2.3 : IUCN (1994) version numbering system. Its identification criteria was later modified and upgraded to the 3.1 : IUCN (2001) version. In 2004, the IUCN released a Red List of Threatened Species. Which is considered as the world's most comprehensive inventory of the global conservation status of plant and animal species.

ONEP as the National Focal Point to the CBD, found it necessary to update the inventory and status of threatened species in Thailand. Thus in October 2004, the Thailand Red Data: Vertebrates List and Project was initiated and underwent a series of data collection, analyses, and meetings amongst involved experts.

ONEP sincerely hopes that this manual will be helpful in the identification of mammals, reptiles and amphibians in Thailand as well as provide basic information for research use in development projects, and related policy development that would all lead to sustainable biological management. Furthermore, ONEP had also published the books titled "Thailand Red Data: Birds" and "Thailand Red Data: Fishes", which will be helpful in the same way.

ONEP would also like to thanks all experts and academics involved who have helped in the creation of this Red List. Lastly, ONEP would like to pay gratitude to the United Nations Development Programme (UNDP) for their financial support in publishing this manual.

*N. Kosiratna*

Mrs. Nisakorn Kosiratna  
Secretary General  
ONEP  
2005



# ABSTRACT



Thailand is a country with a wide range of biological diversity, and high in forest resources and wildlife. Concluding from the status study of mammals, reptiles and amphibians, there are 13 orders, 42 families, 147 genera, and 302 species of mammals; 3 orders, 23 families, 139 genera, and at least 350 species (366 forms) of reptiles; 3 orders, 8 families, 40 genera, and 137 species (138 forms) of amphibians. However new records of these three groups of animals are continuously being reported.

The above three groups of animals have been categorized and listed as either extinct, extinct in the wild, critically endangered, endangered, vulnerable, near threatened, least concern, data deficient or endemic. Out of 159 species of mammals classified, one is an extinct species, the Schomburgk's Deer (*Cervus schomburgki*). This deer has been extinct from Thailand and the world for over 70 years. In the past, the Schomburgk's Deer was prevalent in the central plains of Thailand, but presently only antler remains are found. There are also 4 species that are extinct in the wild, 12 critically endangered, 35 endangered, 69 vulnerable, 15 near threatened, 10 species of least concern, 13 with deficient data, and 5 endemic.

A total of 350 species (366 forms) of reptiles have been classified, all of which have been found and reported in Thailand. The false gavia (*Tomistoma schlegelii*) is listed as extinct in the wild due to intense hunting, and since its eggs require a long incubation period, they are at greater risk to harm from various predators. Presently there is no existing record of false gavials found in the wild. Within the remaining reptile species, eleven are classified as critically endangered, 5 (6 forms) as endangered, 16 as vulnerable, 48 (50 forms) as near threatened, 183 (190 forms) as least concern, 89 (92 forms) as data deficient, and 47 species (49 forms) as endemic. It is note that certain subspecies of the same species are categorized separately.

There are 137 species (138 forms) of amphibians that have been categorized, all of which have been found and reported in Thailand, including 5 species that are vulnerable, 33 species that are near threatened, 64 species (65 forms) that are of least concern, 35 species that are deficient in data and finally, 7 species are endemic.

# ACKNOWLEDGEMENT

We would like to thank all those who provided the valuable information in creating this book; from the brainstorming session with participants from the meeting on the status of Thailand's biological resources, to the information received through personal communication on various occasions. Additionally, we would like to give thanks to Thailand Institute of Scientific and Technological Research, Wildlife Fund Thailand, Wildlife Conservation Society (WCS Thailand), Phuket Marine Biological Center, Kanchanaburi Inland Fisheries Research and Development Center, Thung Yai Naresuan Wildlife Sanctuary, Prateep Duengkae, Nanthachai Pongpattananurak, Pongsakorn Pattapong, Kriangsak Sribuarod, Obhas Khobkhet, Naris Bhumpakphan, Sawai Wanghongsa, Kaset Sutecha, Budsabong Kanchanasaka, Dome Pratumtong, Surachit Waengsothorn (Environment, Ecology and Energy Department, Thailand Institute of Scientific and Technological Research), Somying Thunhikorn, Jonathan Murray, Dusit Arthitayawan, Ronayuth Sribanyaranond, Advanced Thailand Geographic Magazine and Narumon Kritsanachandee. This study and publication is sponsored by UNDP and well organized by Biological Diversity Division of ONEP.

Lastly, we sincerely hope that this book is educational and useful in the identification and status assessment of mammals, reptiles and amphibians, as well as provide basic information for research use in development projects, and related policy creation that would all lead to sustainable biological management.



# C ONTENTS



## Mammals

<i>Cervus schomburgki</i> .....	10
<i>Cervus eldii</i> .....	11
<i>Bos sauveli</i> .....	12
<i>Rhinoceros sondaicus</i> .....	13
<i>Dicerorhinus sumatrensis</i> .....	14
<i>Callosciurus prevostii</i> .....	15
<i>Felis chaus</i> .....	16
<i>Ictailurus planiceps</i> .....	17
<i>Cynogale bennettii</i> .....	18
<i>Lutra sumatrana</i> .....	19
<i>Craseonycteris thonglongyai</i> .....	20
<i>Hylobates agilis</i> .....	21
<i>Mesoplodon ginkgodens</i> .....	22
<i>Orcaella brevirostris</i> .....	23
<i>Naemorhedus caudatus</i> .....	24
<i>Bos javanicus</i> .....	25
<i>Dugong dugon</i> .....	26
<i>Sundasciurus hippurus</i> .....	27
<i>Niviventer hinpoon</i> .....	28
<i>Manis pentadactyla</i> .....	29
<i>Pardofelis marmorata</i> .....	30
<i>Panthera tigris</i> .....	31
<i>Viverra megaspila</i> .....	32
<i>Prionodon pardicolor</i> .....	33
<i>Hemigalus derbyanus</i> .....	34
<i>Lutra lutra</i> .....	35
<i>Mustela strigidorsa</i> .....	36
<i>Rhinolophus marshalli</i> .....	37
<i>Hipposideros halophyllus</i> .....	38
<i>Eptesicus demissus</i> .....	39
<i>Macaca assamensis</i> .....	40
<i>Hylobates pileatus</i> .....	41
<i>Balaenoptera physalus</i> .....	42
<i>Balaenoptera edeni</i> .....	43
<i>Physeter macrocephalus</i> .....	44
<i>Kogia simus</i> .....	45
<i>Kogia breviceps</i> .....	46
<i>Orcinus orca</i> .....	47
<i>Delphinus capensis</i> .....	48
<i>Steno bredanensis</i> .....	49
<i>Stenella attenuata</i> .....	50
<i>Stenella coeruleoalba</i> .....	51
<i>Globicephala macrorhynchus</i> .....	52
<i>Peponocephala electra</i> .....	53

<i>Feresa attenuata</i> .....	54
<i>Neophocaena phocaenoides</i> .....	55
<i>Tragulus napu</i> .....	56
<i>Muntiacus feae</i> .....	57
<i>Axis porcinus</i> .....	58
<i>Bubalus bubalis</i> .....	59
<i>Tapirus indicus</i> .....	60
<i>Elephas maximus</i> .....	61
<b>Status: Vulnerable</b> .....	62
<b>Status: Near Threatened</b> .....	75
<b>Status: Least concern</b> .....	79
<b>Status: Data Deficient</b> .....	82
<b>Status: Endemic</b> .....	84

### Reptiles

<i>Tomistoma schlegelii</i> .....	86
<i>Crocodylus porosus</i> .....	87
<i>Crocodylus siamensis</i> .....	88
<i>Chelonia mydas</i> .....	89
<i>Eretmochelys imbricata</i> .....	90
<i>Lepidochelys olivacea</i> .....	91
<i>Dermochelys coriacea</i> .....	92
<i>Batagur baska</i> .....	93
<i>Callagur borneensis</i> .....	94
<i>Chitra chitra</i> .....	95
<i>Chitra vandijki</i> .....	96
<i>Pelochelys cantorii</i> .....	97
<i>Platysternon megacephalum</i> .....	98
<i>Indotestudo elongata</i> .....	99
<i>Manouria emys emys</i> .....	100
<i>Manouria emys phayrei</i> .....	101
<i>Manouria impressa</i> .....	102
<i>Physignathus cocincinus</i> .....	103
<b>Status: Vulnerable</b> .....	104
<b>Status: Near Threatened</b> .....	107
<b>Status: Least concern</b> .....	115
<b>Status: Data Deficient</b> .....	157
<b>Status: Endemic</b> .....	168

### Amphibians

<i>Paa bourreti</i> .....	176
<i>Paa fasciculispina</i> .....	177
<i>Theloderma gordonii</i> .....	178
<i>Theloderma horridum</i> .....	179
<i>Theloderma stellatum</i> .....	180
<b>Status: Near Threatened</b> .....	181
<b>Status: Least concern</b> .....	187
<b>Status: Data Deficient</b> .....	201
<b>Status: Endemic</b> .....	206
Reference .....	208
Index .....	215
Sources .....	233







# MAMMALS

EX = Extinct  
EW = Extinct in the Wild  
CR = Critically Endangered  
EN = Endangered  
VU = Vulnerable  
NT = Near Threatened  
LC = Least Concern  
DD = Data Deficient  
Endemic

WS = Wildlife Sanctuary  
NH = Non-hunting area  
NP = National Park



<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Cervidae
<b>Scientific Name:</b>	<i>Cervus schomburgki</i>
<b>Common Name:</b>	เนื้อสมัน Schomburgk's deer

**Habitat:**

Lowland floodplain of rivers.

**Locality:**

Formerly throughout the Central Chao Phraya River Floodplain, west to Uthai Thani, east to Nakhon Nayok, Prachin Buri, Chachoengsao, Bangkok and Samut Prakarn. Now entirely extirpated from Thailand.

**Description:**

This spectacular deer is endemic to low-lying plains of the central part, but at present already extinct. Back uniformly dark brown with lighter underpart. Forehead and legs of reddish shades of brown. Short tail has silvery white under surface. Distinctive antlers of the bucks often described as basket-like, formed by the forking at every point of branching; beam relatively short; and brow tine always forked.

*Cervus eldii*

<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Cervidae
<b>Scientific Name:</b>	<i>Cervus eldii</i>
<b>Common Name:</b>	ละอง, ละมั่ง Eld's deer

**Habitat:**

Clearings in mixed deciduous forest.

**Locality:**

Formerly seen in small herds in Lop Buri, Chaiyaphum, Nakhon Ratchasima and adjoining provinces.

**Description:**

Shoulder height 1.2 m.

This medium-sized deer has variable coat colours according to the season, light brown with pale underpart in dry season and dark brown with whitish underpart in wet season. White colour appears on chin, around eyes and along ear margins. Only bucks have antlers, which are bow-shaped with long brow tine forming a continuous curve with the beam, with many small tines clustered at the upper end of the beam. Does lighter in colour than bucks; fawn distinctly spotted.



**Class:** Mammalia  
**Order:** Artiodactyla  
**Family:** Bovidae  
**Scientific Name:** *Bos sauveli*  
**Common Name:** กูปรี่  
 Kouprey

**Habitat:**

Clearings in mixed deciduous forest.


**Locality:**

Extinct from Thailand. Formerly from Ubon Ratchathani (Yod Dom) and Buri Ram (Dong Yai).

**Description:**

Head-body 2.1-2.2 m; tail 100-110 cm; height 1.7-1.9 m, weight 700-900 kg.

A large wild cattle with peculiar horn shape and in adult bulls long hanging dewlaps. Old bulls almost black or very dark brown; cows and young bulls grey with lighter underpart. Both sexes have white stockings. Horns of bulls curving forwards and round, then up with frayed tips, while those of cows are lyre-shaped. Dorsal ridge in bulls weakly developed.

*Rhinoceros sondaicus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Perissodactyla
<b>Family:</b>	Rhinocerotidae
<b>Scientific Name:</b>	<i>Rhinoceros sondaicus</i>
<b>Common Name:</b>	แรด, ระมาด Javan rhinoceros

**Habitat:**

Evergreen forest with hills and streams.

**Locality:**

Now already extinct from Thailand. Formerly from Kanchanaburi (Thung Yai).

**Description:**

Height up to 1.7 m; weight up to 1.4 t.

This massive herbivore, of armour-plated appearance, has only one horn on the end of the nose, grey hairless skin and short stout legs. Unlike the cattle horns, rhino horns have no bony core like those of cattle and sheep, and consisting entirely of keratin fibres fusing compacted together on a roughened area on the skull. Each foot has only three toes. Three skin folds crossing body on neck, behind front legs and before hind legs.



<b>Class:</b>	Mammalia
<b>Order:</b>	Perissodactyla
<b>Family:</b>	Rhinocerotidae
<b>Scientific Name:</b>	<i>Dicerorhinus sumatrensis</i>
<b>Common Name:</b>	ពិន្ទេង Sumatran rhinoceros

**Habitat:**

Evergreen forest with hills and streams.

**Locality:**

Now already extinct from Thailand. Formerly from Chaiyaphum (Phu Khieo-1983); Narathiwat (Hala-Bala-1985-1986); Phetchaburi (Kaeng Krachan-1987); Surat Thani (Khao Sok-1987, Khlong Saeng-1987); Kanchanaburi (Thung Yai Naresuan-1988); Chaiyaphum (Phu Khieo-1996); Narathiwat/Yala (Hala-Bala-1997).

**Description:**

Head-body 2.5-3.2 m; tail 40-50 cm; height 1.2-1.4 m; anterior horn up to 38 cm; weight 800-1,000 kg.

A two-horned hairy herbivore with large stocky body, short stout legs and three toes on each foot. Skin grey or reddish grey, sparsely covered with long hair. Young somewhat shaggy. Anterior horn much longer than posterior one; both on top of its nose. Only one skin fold seen across the body behind the front legs.



**Class:** Mammalia  
**Order:** Rodentia  
**Family:** Sciuridae  
**Scientific Name:** *Callosciurus prevostii*  
**Common Name:** กระรอกสามสี  
Prevost's squirrel



**Habitat:**

Lowland swamp forest, rubber plantations.

**Locality:**

Narathiwat (Pa Phru Toh Dang).

**Description:**

Head-body 20-27 cm; tail 20-27 cm; weight 250-500 g.

A strikingly-coloured arboreal squirrel of the extreme south rainforest. Upperpart black, underpart and lower part of both legs chestnut red, between which there is a broad white side stripe running from the snout to hind quarter, with a thin black line below it. Tail bushy black, as long as the head-body length.



**Class:** Mammalia  
**Order:** Carnivora  
**Family:** Felidae  
**Scientific Name:** *Felis chaus*  
**Common Name:** เสือกระทาย  
 Jungle cat

**Habitat:**

Dry savana and scrub forest.

**Locality:**

Loei (Phu Kradung); Chaiyaphum (Phu Khieo); Maha Salakham (Na Chuak); Ratchaburi.

**Description:**

This scrub-dwelling cat is larger than a domestic cat with long legs and a short ringed tail. Body sandy brown to yellow grey, paler on the belly; sometimes with dark stripes on face and legs. Ears tapered and tufted with a light spot at the base, having longer ears than other wild cats. Young have close-set striped pattern which disappears in the adult.



*Ictailurus planiceps*

<b>Class:</b>	Mammalia
<b>Order:</b>	Carnivora
<b>Family:</b>	Felidae
<b>Scientific Name:</b>	<i>Ictailurus planiceps</i>
<b>Common Name:</b>	แมวป่าหัวแบน Flat-headed cat

**Habitat:**

Evergreen forest near streams.

**Locality:**

Narathiwat (Pa Phru Toh Dang).

**Description:**

Head-body 41-50 cm; tail 13-17 cm; weight 5.5-8.0 kg.

A small dark greyish to reddish brown cat, of the size of a domestic cat, with slightly flattened head, small rounded ears and very short tail. Hair of upperpart with grey and pale buff speckling; those of underpart usually white spotted sparsely with dark marking on throat, belly and inner sides of legs. Face with distinct white tear streaks. Legs short with small paws and semi-retractile claws.



<b>Class:</b>	Mammalia
<b>Order:</b>	Carnivora
<b>Family:</b>	Viverridae
<b>Scientific Name:</b>	<i>Cynogale bennettii</i>
<b>Common Name:</b>	อีเห็นน้ำ Otter civet

**Habitat:**

Lowland swamp forest.

**Locality:**

Narathiwat (Pa Phru Toh Dang).

**Description:**

Head-body 50-65; tail 30-36 cm; weight 2-5 kg.

This semi-aquatic dark carnivore is otter-like in general appearance, though it has broader snout and smaller shorter tail. Body dark brown, finely grizzled by lighter spots; belly paler brown. Head broad and flat; thick lips adorned by very long whiskers. Its first three upper premolars unusually large with high, compressed and pointed crown, remaining cheek teeth broad. Ears small, rounded and well-separated. Feet partially webbed.



**Class:** Mammalia

**Order:** Carnivora

**Family:** Mustelidae

**Scientific Name:** *Lutra sumatrana*

**Common Name:** นากใหญ่หัวปลาคูก, นากใหญ่จมูกขน  
Hairy-nosed otter



**Habitat:**

Lowland riverine forest to mangrove forest.

**Locality:**

Trang (Kao Bantad); Narathiwat (Pa Phru Toh Dang).

**Description:**

Head-body 50-82 cm; tail 35-38 cm; weight 5-6 kg.

The presence of entirely hairy nose and rhinarium (moist part of nose), which being naked in other coexisting otter species makes this otter easy to be recognized at close range, and its dark chocolate pelage with slightly paler underpart in the field. Upper lip, chin and throat sharply-demarcated white. Head flattened considerably than those of other otters. Tail flattened and oval in cross section. Feet well-webbed and armed with strong claws.



**Class:** Mammalia  
**Order:** Chiroptera  
**Family:** Craseonycteridae  
**Scientific Name:** *Craseonycteris thonglongyai*  
**Common Name:** คางคาวคุณกิตติ  
 Kitti's hog-nosed bat

**Habitat:**

Limestone caves in mixed deciduous forest.

**Locality:**

Kanchanaburi (Sai Yok).

**Description:**

Head-body 2.9-3.3 cm; tail none; weight 1.5-3.0 g.

The world smallest flying mammal, sometimes called “Bumblebee bat” has a pig-like snout. Upperpart brownish red or grey; underpart paler and wings darker. Ears relatively large with a prominence (tragus) at the base of outer ear. Tailless, but interfemoral membrane stretched between its thin hind legs.

*Hylobates agilis*

<b>Class:</b>	Mammalia
<b>Order:</b>	Primates
<b>Family:</b>	Hominidae or Hylobatidae
<b>Scientific Name:</b>	<i>Hylobates agilis</i>
<b>Common Name:</b>	ชะนีมือดำ Agile gibbon

**Habitat:**

Lowland evergreen forest.

**Locality:**

Narathiwat (Hala-Bala).

**Description:**

Head-body 42-47 cm; tail none; weight 5.0-6.4 kg.

Can be told apart from the common Lar gibbons by having black hands and feet, not white. It occurs in two colour phases, one black and the other grey to buff, which are not related either to age or sex. Males have white brows and cheeks, while females only white brows.



<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Hyperoodontidae or Ziphiidae
<b>Scientific Name:</b>	<i>Mesoplodon ginkgodens</i>
<b>Common Name:</b>	วาฬฟันซี่ยาว Ginkgo-toothed beaked whale

**Habitat:**

Offshore; both in the Gulf of Thailand and the Andaman Sea.

**Locality:**

Phuket (Patong Beach - 1988).

**Description:**

Maximum length 5.2 m; weight 1.5-3.6 t.

A beaked whale having slender head, slightly bulbous forehead and an upturned beak. Body black, paler on ventral surface; often adorned with oval white scars on belly and sides. Dorsal fin small and curved. Flippers lobate and pointed. Tail flukes without any notch. Mandibular teeth distinctively shaped like ginkgo-tree leaves and situated upon raised areas in the rear half of each side of the lower jaw; teeth widest antero-posteriorly of any *Mesoplodon*, i.e. 100 mm. Juveniles 2-2.5 m long at birth.

*Orcaella brevirostris*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Monodontidae or Delphinidae
<b>Scientific Name:</b>	<i>Orcaella brevirostris</i>
<b>Common Name:</b>	โลมาอิรวดี, โลมาหัวบาตรคريبหลัง Irrawaddy dolphin

**Habitat:**

Along both shores; also in rivers and lagoon.

**Locality:**

Chachoengsao (Bangprakong); Songkhla (Songkhla Lake); Surat Thani (Kanchanadit); Phatthalung (Lumpam, Khlong Luang, Pak Prayun); Phangnga (Phangnga Bay); Krabi (Krabi Bay); Bangkok (Chao Praya River); Chon Buri (Bang Saen, Naklua); Rayong (Koh Samet, Ban Phe and Rayong Districts); Pattani (Tanjong Mas); Trat (Klong Yai, Trat and Laem Ngop Districts); Chantaburi (Laem Sing).

**Description:**

Adult length : 1.8-2.75 m; body weight about 100 kg.

A blunt-headed dolphin with a small dorsal fin. Body uniformly slaty grey or slaty blue, slightly paler on the underside. Upper jaws bearing 34-40 teeth, while the lower ones 30-36 teeth. Dorsal fin small, sickle-shaped with a rounded tip, located more on the rear half of the body. Flippers broadly triangular. Juveniles weigh 3.8 to 6 kg at birth.



**Class:** Mammalia

**Order:** Artiodactyla

**Family:** Bovidae

**Scientific Name:** *Naemorhedus caudatus*

**Common Name:** กวางผา  
Chinese goral



### Habitat:

Hill evergreen forest, especially isolated precipitous peaks.

### Locality:

Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Phu Moen, Mae Lao-Mae Sah); Tak (Doi Mon Chong, Doi Dukado, Mae Ping).

### Description:

Head-body 1.0-1.2 m; shoulder height 50-75 cm; weight 22-35 kg.

This inhabitant of steep cliffs is looked like a small goat. Having rough, shaggy coat of greyish to grey brown in colour; underpart lighter, it always has a distinct spinal line, most pronounced at the rump. Pale throat patch usually quite faint or even indistinct in some individuals. Tail with a terminal tuft. Preorbital gland rather rudimentary.



*Bos javanicus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Bovidae
<b>Scientific Name:</b>	<i>Bos javanicus</i>
<b>Common Name:</b>	วัวแดง Banteng

**Habitat:**

Mixed deciduous forest, along streams.

**Locality:**

Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra, Thung Yai); Ratchaburi (Maenam Pachi); Phetchaburi (Kaeng Krachan); Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jonk-Na Yoi, Yot Dom); Chachoengsao (Khao Ang Rue Nai).

**Description:**

Head-body 1.9-2.5 m; tail 65-70 cm; overall height 1.7 m; weight 600-800 kg.

In all proportions, it is similar to domestic cattle, but larger. Adult bulls dark chestnut, some southern individuals almost black; young bulls and cows reddish brown. All have white band around muzzle, white patch over eyelids, white stockings and white rump patch or buttock. Bulls have dorsal ridge and dewlap, and also horny bald skin patch between horn bases, Both sexes have horns, though much larger in bulls.



**Class:** Mammalia  
**Order:** Uranotheria or Sirenia  
**Family:** Dugongidae  
**Scientific Name:** *Dugong dugon*  
**Common Name:** พะยูง  
 Dugong

**Habitat:**

Along shores of both the Gulf and Andaman Sea.

**Locality:**

Trang (Hat Chao Mai, Koh Libong-1991); Rayong (Khlung-2003); Trat (Ban Mai Rut-Ban Saphan Hin-2003); Phuket (2004); Surat Thani (2004); Chon Buri (Sattahip-2005); Krabi (Koh Sriboya, Koh Pu, Koh Chom-2005); Phangnga (2005); Seagrass beds of Chanthaburi-Trat (2005).

**Description:**

Head-tail 1-4 m; weight 230-900 kg.

An aquatic non-ruminant herbivore of the coastal shallows. Body dull brownish grey with short sensory bristles at the intervals of 2-3 cm. Front limbs modified into paddle-like flippers. Head large with a pair of valued nostrils on the top of the head; below equipped with numerous long whiskers. Tail flattened, fluke-like with a concave trailing edge. This animal is evolutionarily related to elephants and ungulates.

*Sundasciurus hippurus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Rodentia
<b>Family:</b>	Sciuridae
<b>Scientific Name:</b>	<i>Sundasciurus hippurus</i>
<b>Common Name:</b>	กระรอกหางมาใหญ่ Horse-tailed squirrel

**Habitat:**

Evergreen forest, near streams.

**Locality:**

Nakhon Si Thammarat (Khao Luang); Narathiwat (Pa Phru Toh Dang); Yala (Hala-Bala).

**Description:**

Head-body 21-25 cm; tail 17-25 cm; weight 260-365 g.

Easily recognized by its bushy, entirely dark grey to black tail, this diurnal medium-sized squirrel is reddish brown on back with grey head and shoulders. Underpart dark reddish.



**Class:** Mammalia

**Order:** Rodentia

**Family:** Muridae

**Scientific Name:** *Niviventer hinpoon*

**Common Name:** หนูชนเสี้ยนเขาหินปูน  
Limestone rat



#### Habitat:

Limestone caves.

#### Locality:

Saraburi (Kaeng Khoi); Lop Buri; Uthai Thani (Huai Kha Kaeng).

#### Description:

This endemic mammal is small, about the size of a house mouse, with long uniformly dark tail at least 125 % of the head-body length, and thinly tufted with hairs at the tip. Coat dark reddish brown with numerous stiff spines, especially at the hind part; belly much paler than back or sometimes whitish.

*Manis pentadactyla*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cimolesta or Pholidota
<b>Family:</b>	Manidae
<b>Scientific Name:</b>	<i>Manis pentadactyla</i>
<b>Common Name:</b>	ลิ่นจีน Chinese pangolin

**Habitat:**

Hill evergreen forest.

**Locality:**

Chiang Mai (Doi Inthanon); Mae Hong Son (Lumnam Pai); Loei (Chiang Khan).

**Description:**

It can be recognized instantly by having body covered almost wholly by horny scales. Body elongate, tapering to a stout tail. Head small and conical with a reduced outer ears. Limbs short, powerful, terminate in 5 clawed digits, of which three middle ones on the front foot long and curved. This one has 14-17 scales along the edge of the tail. Few hairs present between the scales, lacking in African species.



<b>Class:</b>	Mammalia
<b>Order:</b>	Carnivora
<b>Family:</b>	Felidae
<b>Scientific Name:</b>	<i>Pardofelis marmorata</i>
<b>Common Name:</b>	แมวลายหินอ่อน Marbled cat

**Habitat:**

Evergreen forest along waterways.

**Locality:**

Tak (Thung Yai); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Kaeng Krachan); Loei (Chiang Khan); Chaiyaphum (Phu Khieo); Ubon Ratchathani (Yot Dom); Sakaew (Ta Phraya); Surat Thani (Khlong Saeng); Yala (Hala-Bala).

**Description:**

Head-body 40-60 cm; tail 45-54 cm; weight 2.5-5.5 kg.

An arboreal long-furred wild cat with cloud-like body patterns, similar to that of the Clouded leopard, but smaller in size, and possessing a much bushy tail. Black-edged blotches on body less distinctive, rather blurred in overall appearance. Spots on legs much smaller and more elongate in shape. Behind each ear with a white spot.

*Panthera tigris*

<b>Class:</b>	Mammalia
<b>Order:</b>	Carnivora
<b>Family:</b>	Felidae
<b>Scientific Name:</b>	<i>Panthera tigris</i>
<b>Common Name:</b>	เสือโคร่ง Tiger

**Habitat:**

Evergreen forest, to deciduous forest.

**Locality:**

Tak (Thung Yai Naresuan); Chiang Mai (Om Koi); Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Khao Yai); Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jong-Na Yoi, Yot Dom); Chachoengsao (Khao Ang Rue Nai); Prachin Buri (Pang Sida); Sakaew (Ta Phraya); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra, Thung Yai, Sai Yok, Thong Pha Phum); Phetchaburi (Kaeng Krachan); Ratchaburi (Menam Pachi); Prachuap Khirikhan (Kui Buri); Ranong (Khlong Naka); Surat Thani (Khlong Saeng).

**Description:**

An unmistakable large, massively built carnivore with brownish orange coat patterned all over from head to tail by dark vertical stripes. Underparts paler. Its front limbs and shoulders heavily muscled, and paws equipped with long, sharp retractile claws; white the hind limbs, though longer, less muscled.



**Class:** Mammalia

**Order:** Carnivora

**Family:** Viverridae

**Scientific Name:** *Viverra megaspila*

**Common Name:** ชะมดแผงสันหางดำ  
Large-spotted civet



### Habitat:

Scrub forest around swamps, and along waterways.

### Locality:

Ranong (Ban Bang Non); Uthai Thani (Huai Kha Khaeng); Srakaew (Ta Phraya); Chachoengsao (Khao Ang Rue Nai).

### Description:

Head-body 76 cm; tail 37 cm; weight 6.6 kg.

A large terrestrial civet with spinal crest of erectile black hairs from shoulder to rump, flanked by a longitudinal row of black spots. Coat greyish brown with small indistinct dark spots on the foreparts down to legs. Flanks adorned with large dark spots often fusing into stripes and bars. Neck with pronounced black and white stripes. Tail ringed with 5-7 white bands, but most of them do not circle the tail completely and black-tipped. Claws semiretractile; sole naked, scantily haired between toes and foot pads. A perineal scent gland occurs between the anus and genitalia.



*Prionodon pardicolor*

**Class:** Mammalia

**Order:** Carnivora

**Family:** Viverridae

**Scientific Name:** *Prionodon pardicolor*

**Common Name:** ชะมดแปดงลายจุด  
Spotted linsang

**Habitat:**

Hill evergreen forest to lowland evergreen forest.

**Locality:**

Mae Hong Son (Lum Nam Pai); Nan (Phu Ka).

**Description:**

Head-body 39 cm; tail 34 cm; weight 600 g.

This small, trimly built secretive forest carnivore can be readily recognized by its light yellow coat with dark spots on back, which decreasing in sizes down the legs; and neck stripes. Tail with 8-9 dark bands alternating with thin light bands. It has no second upper molar and civet gland; but with fully retractile claws.



<b>Class:</b>	Mammalia
<b>Order:</b>	Carnivora
<b>Family:</b>	Viverridae
<b>Scientific Name:</b>	<i>Hemigalus derbyanus</i>
<b>Common Name:</b>	อีเห็นลายพาด Banded palm civet

**Habitat:**

Lowland evergreen forest.

**Locality:**

Nakhon Si Thammarat (Thung Song, Thung Yai); Surat Thani (Khlung Saeng, Khao Sok); Yala (Hala-Bala).

**Description:**

Head-body 50-65 cm; tail 30-36 cm; weight 2-3 kg.

A nocturnal terrestrial civet with varying shades of coat colour, from pale yellow, greyish buff, to reddish brown. Body transversely crossed by 5 dark brown bands, extending halfway down flank. Longitudinal stripes present on face and shoulders. Tail mostly dark brown, only with distinct 2 dark rings at the base.



**Class:** Mammalia

**Order:** Carnivora

**Family:** Mustelidae

**Scientific Name:** *Lutra lutra*

**Common Name:** นากใหญ่ภูเขา

Common otter



**Habitat:**

Mountain streams in evergreen forest.

**Locality:**

Chiang Rai (Nong Bong Khai); Mae Hong Son (Lum Nam Pai, Huai Nam Dung); Tak (Thung Yai, Umphang); Kamphaeng Phet (Khlong Lan, Khlong Wang Chao); Nakhon Sawan (Mae Wong); Kanchanaburi (Thung Yai, Sri Nakarin); Uthai Thani (Huai Kha Kaeng).

**Description:**

Head-body 57-70 cm; tail 35-40 cm.

An amphibious carnivore with elongated, sinuous and lithe body, short limbs, and a long tail. Its coat light brownish-grey to dusky brown; lighter on the underside, except the buff to cream chest. Tail thick at the base, tapering to a point; with flattened underside. Feet well-webbed, equipped with strong claws. There are numerous stiff whiskers around the snout and in tufts on the elbows.



**Class:** Mammalia

**Order:** Carnivora

**Family:** Mustelidae

**Scientific Name:** *Mustela strigidorsa*

**Common Name:** เพียงพอนเสนหลังขาว

Back-striped weasel



### Habitat:

Hill evergreen forest.

### Locality:

Nan (Ban Na Kha); Chiang Mai (Doi Pha Hom Pok, Doi Ang Khang); Loei (Phu Luang); Khamphaeng Phet (Doi Mokoju); Chaiyaphum (Phu Khieo).

### Description:

Head-body 29 cm; tail 15 cm.

Medium-sized weasel with deep chocolate brown body, tail and limbs. Head colouration may be very slightly paler. It has a diagnostic silvery white stripe running down the centre of the back. Also has a yellowish white stripe on underside. Upper lips, cheeks, chin and throat are pale yellow. Tail bushy. Feet naked at all seasons.

*Rhinolophus marshalli*

**Class:** Mammalia

**Order:** Chiroptera

**Family:** Rhinolophidae

**Scientific Name:** *Rhinolophus marshalli*

**Common Name:** คางคาวมงกุฎหูโตมาร์แชล  
Marshall's horseshoe bat

**Habitat:**

Dry evergreen forest to moist evergreen forest.

**Locality:**

Chiang Mai (Doi Chang Khian); Chanthaburi (Khao Soi Dao); Phetchubun (Nam Nao); Phitsanulok (Phu Hin Rong Klar).

**Description:**

A large-eared insectivorous bat with strange complex nose-leaf; the front part horseshoe-shaped with a sella projects forward and a generally pointed lancet running lengthwise. Recognized by having a cup-shaped structure between the nostrils, and its sides expanding into trapezoid flaps.



<b>Class:</b>	Mammalia
<b>Order:</b>	Chiroptera
<b>Family:</b>	Rhinolophidae
<b>Scientific Name:</b>	<i>Hipposideros halophyllus</i>
<b>Common Name:</b>	ค้างคาวหน้ายักษ์จมูกปุ่ม Thai roundleaf bat

**Habitat:**

Limestone caves in agricultural areas.

**Locality:**

Sakaew (Tham Khao Sing-to); Lop Buri (Khao Samokhon); Uthai Thani (Huai Kha Kaeng); Khon Kaen (Phu Khao-Phu Pan Kum).

**Description:**

A small grey to pale grey insectivorous bat, endemic to Thailand, has an elaborate nose-leaf. The anterior part rounded and horseshoe-shaped; the middle part a low cushion-like structure expanding laterally; the posterior part low and rounded. The diagnostic character is a pair of small kidney-like nodules between nostrils. Ears very large and broad.

*Eptesicus demissus*

**Class:** Mammalia

**Order:** Chiroptera

**Family:** Vespertilionidae

**Scientific Name:** *Eptesicus demissus*

**Common Name:** คางคาวทองสีน้ำตาลสุราษฎร์  
Surat brown bat

**Habitat:**


Evergreen forest.

**Locality:**

Surat Thani (Khao Nong); Phangnga (Khao Sok).

**Description:**

One of Thai endemic mammals from the southern forests. It has short chestnut brown fur dorsally and light brown ventrally; some hairs have paler tips distributing sparsely all over the back. Fur extends onto the interfemoral membrane at the base of the tail in a triangular pattern. Ears short, narrow and rounded.



**Class:** Mammalia  
**Order:** Primates  
**Family:** Cercopithecidae  
**Scientific Name:** *Macaca assamensis*  
**Common Name:** ลิงอายเงี้ยว  
 Assamese macaque

**Habitat:**

Hill evergreen forest to dry evergreen forest.

**Locality:**

Chiang Rai; Tak (Huai Nua Pla, Ban Mae Lamao, Umphang); Kanchanaburi (Si Sawat, Thong Pha Phum); Uthai Thani (Huai Kha Kaeng); Chiang Mai (Weng Hang).

**Description:**

Head-body 53-68 cm; tail 19-38 cm; weight 7.8 kg (male) 6.1 kg (female). A large macaque of varying shades of coat colour: yellowish brown, grey brown to dark brown; always running lighter from front part to rear part. Face and skin around anus naked and red in adult. Tail about half the length of head-body. Some individuals have rather long beard.



*Hylobates pileatus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Primates
<b>Family:</b>	Hominidae or Hylobatidae
<b>Scientific Name:</b>	<i>Hylobates pileatus</i>
<b>Common Name:</b>	ชะนีมงกุฏ Pileated gibbon

**Habitat:**

Lowland evergreen forest.

**Locality:**

Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jong-Na Yoi, Yot Dom); Chachoengsao (Khao Ang Rue Nai); Prachin Buri (Khao Yai, Tub Lan); Sakaew (Ta Phraya); Chon Buri (Ban Nong Khor); Chanthaburi (Khao Soi Dao); Trat (Khlung Yai, Ban Huang Som).

**Description:**

Head-body 45-65 cm; weight 5.5-6.7 kg.

This lesser ape, though both sexes similar in size, shows sexual dimorphism in coat colour. Males black with white hands and feet, also possessing a white circles around the face and the crown. Females silvery-grey with black crown, cheeks and breast. Young grey at birth.



<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Balaenopteridae
<b>Scientific Name:</b>	<i>Balaenoptera physalus</i>
<b>Common Name:</b>	วาฬแกลบครีบน้ำดำ, วาฬฟิน Fin whale

**Habitat:**

Gulf of Thailand.

**Locality:**

Chachoengsao (Muang).

**Description:**

Maximum length 17-27 m, weight 75-80 t.

A huge whale with flat-topped, v-shaped head. Body dark grey to brownish black on back and sides; underside white, including underside of flippers. Series of 56-100 ventral grooves extending from head to at least the naval. Left front baleen white or yellow white, the rest striped with alternate bands of white and bluish grey. Dorsal fin more than 60 cm tall, located closed to the tail flukes. Flippers relatively short and narrow. Flukes having straight trailing edge with median notch. Females larger than males.

*Balaenoptera edeni*

**Class:** Mammalia

**Order:** Cetacea

**Family:** Balaenopteridae

**Scientific Name:** *Balaenoptera edeni*

**Common Name:** วาฬแถบ, วาฬบรูดา, วาฬซิทตั้ง  
Bryde's whale

**Habitat:**

Gulf of Thailand and Andaman Sea, along shores and around islands.

**Locality:**

Prachuap Khirikhan (Bang Saphan, Khlong Wan, Sam Roi Yod, Hua Hin); Chumphon (Koh Tao, Khanthuli, Bo Mao Bay, Pathio); Trang (Koh Sukorn); Phuket (Sapum Bay, Para Bay, Koh Racha, Phuket Harbour); Phangnga (Natai Beach); Rayong (Ban Pae); Nakhon Si Thammarat (Tha Sala, Si Chon); Bangkok (Bang Na); Chon Buri (Bang Saen, Ban Bang Lamung); Surat Thani (Ao Ban Don); Phetchaburi (Had Chao Samran); Satun (Koh Tarutao); Krabi (Koh Lanta Yai, Koh Lanta Boh); Chon Buri (Koh Si Chang); Samut Prakarn (Samut Prakarn Harbour).

**Description:**

Maximum length 13-15.5 m; weight 26-30 t.

A large-sized whale with 3 prominent ridges on the head from blowholes to snout, one medial and one on each side of it. Body overall dark grey. Baleen slaty grey with coarse dark bristles. Only 40-50 ventral grooves extending beyond or at least to navel. Dorsal fin extremely falcate and often pointed apically, located at a point one-thirds of the body length. Flippers long, rectangular and pointed at tips. Tail flukes broad, with straight trailing edge and slight median notch. Females larger than males.



**Class:** Mammalia

**Order:** Cetacea

**Family:** Physeteridae

**Scientific Name:** *Physeter macrocephalus*

**Common Name:** วาฬหัวทุย  
Sperm whale



### Habitat:

Off-shores in Andaman Sea.

### Locality:

Phuket (Cherng Tale Beach, Surin Beach, Phuket Harbour); Phangnga (Kha Pilai Beach, Koh Similan); Satun (Koh Bu Boi).

### Description:

Length maximum males 18 m, females 12 m.

A massive, boxlike-shaped whale with head comprising up to 40% of the body length. Snout bluntly squarish extends beyond tip of lower jaw. Body tapering rapidly after the dorsal hump to robust caudal peduncle. Skin corrugated and wrinkled in appearance. Body usually dark brownish grey, more paler on belly. Flippers broad, lobate. Flukes broad, triangular, with straight trailing edge and deeply notched. Teeth thick, conical, 18-25, fitting into sockets in a toothless upper jaw. Males considerably larger than females.

*Kogia simus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Physeteridae
<b>Scientific Name:</b>	<i>Kogia simus</i>
<b>Common Name:</b>	วาฬหัวทุยแคระ Dwarf sperm whale

**Habitat:**

Off-shores in Andaman Sea.

**Locality:**

Phuket (Patong Beach); Satun; Phangnga.

**Description:**

Maximum length 2.7 m, weight 280 kg.

In general appearance, similar to *Kogia breviceps* but smaller in size, with more erect dorsal fin showing as a much longer base. Body dark bluish grey dorsally, lighter grey on sides and dull white or pink on ventral surface. Skin wrinkled. False gill present. Dorsal fin tall and falcate at mid-body, height more than 5% of body length. Flippers originating below and behind the false gill. Tail flukes with concave trailing edge and distinct median notch. Mandibular teeth 7-12; all teeth shorter than 30 mm and less than 4.5 mm in diameter.



<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Physeteridae
<b>Scientific Name:</b>	<i>Kogia breviceps</i>
<b>Common Name:</b>	วาฬหัวทุยเล็ก Pygmy sperm whale

**Habitat:**

Off-shores in Andaman sea.

**Locality:**

Rayong; Chumphon; Songkhla; Phangnga; Phuket.

**Description:**

Maximum length 3.4 m, body weight 400 kg.

A small robust shark-like whale having squarish or conical head with underslung lower jaw. False gill mark on side of head set off by dark line posteriorly. Body colour dark bluish-grey dorsally, lighter grey laterally and white ventrally. Flippers long, up to 14% of body length, located just behind the false gill. Dorsal fin low, strongly falcate, generally less than 5% of body length. Tail flukes with concave trailing edge and distinct median notch. Teeth thin and sharply pointed, 12-16 pairs in lower jaw.

*Orcinus orca*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Delphinidae
<b>Scientific Name:</b>	<i>Orcinus orca</i>
<b>Common Name:</b>	วาฬเพชฌฆาต Killer whale

**Habitat:**

Gulf of Thailand and Andaman Sea.

**Locality:**

Ranong (Similan); Surat Thani; Phangnga; Phuket; Satun.

**Description:**

Maximum length males 9.5 m, females 7 m; body weight males 8 t., females 4 t.

A striking black and white dolphin with extremely tall and wide dorsal fin, triangular in male and falcate in female. Head with indistinct beak and conical snout. Body color black dorsally except for oval white patch above and behind each eye and grey patch behind dorsal fin. Chin and throat areas white continuing to white flank patch to white anal patch. Flippers broad and paddle-shaped with blunt tips. Flukes having concave trailing edge and median notch. Juveniles at birth 2.1-2.4 m long and 180 kg in weight.



<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Delphinidae
<b>Scientific Name:</b>	<i>Delphinus capensis</i>
<b>Common Name:</b>	โลมาปากยาว Long-beaked dolphin

**Habitat:**

Along shores; both waters.

**Locality:**

Prachuap Khirikhan; Surat Thani (Koh Samui).

**Description:**

Maximum length males 2.6 m, females 2.5 m; weight about 235 kg.

A slender and streamlined dolphin has a long beak sharply demarcated from the gently sloping forehead. Colour very complex and distinctive. Generally a crisscross colour pattern on sides, being tawny-brown to yellow anteriorly and greyish posteriorly. Dorsal fin large and located at mid-body, slightly falcate. Flippers tapering to a point. Flukes with concave trailing edge and well-defined median notch. Jaw with 40-58 small sharply pointed teeth.



*Steno bredanensis*

**Class:** Mammalia

**Order:** Cetacea

**Family:** Delphinidae

**Scientific Name:** *Steno bredanensis*

**Common Name:** โลมาฟันห่าง  
Rough-toothed dolphin

**Habitat:**

Off-shores in Andaman Sea and in the Gulf.

**Locality:**

Phuket (Chalong Bay); Satun; Nakhon Si Thammarat.

**Description:**

Maximum length 2.8 m.

Generally slim; body cone-shaped ahead of flippers; long slender beak with no demarcation from head. Forehead slopes smoothly to long beak. Gape straight. Eyes large and black. Body usually grey to purplish black dorsally, white or pinkish-washed ventrally; yellowish white blotches often present on body. Lips and lower jaws often white, with 20-27 teeth whose crowns marked by series of vertical wrinkles.



<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Delphinidae
<b>Scientific Name:</b>	<i>Stenella attenuata</i>
<b>Common Name:</b>	โลมาลายจุด Bridled dolphin

**Habitat:**

Off-shores in the Gulf of Thailand and Andaman Sea.

**Locality:**

Songkhla; Phuket; Samut Sakhon; Samut Prakan; Chon Buri; Krabi (Ban Lanpong).

**Description:**

Medium-sized, slender animal with varied spotting depending on age and geographical area. Upper back (cape) and dorsal fin dark grey, lateral band of mid-grey and underside light grey/white. The cape dips down below the dorsal fin. A very falcate, narrow-based, dorsal fin, normally pointed to the tip. A long rostrum with a white-tipped beak. Tooth counts range from 34 to 48 pairs of sharply pointed teeth in each jaw.

*Stenella coeruleoalba*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Delphinidae
<b>Scientific Name:</b>	<i>Stenella coeruleoalba</i>
<b>Common Name:</b>	โลมาแถบ Striped dolphin

**Habitat:**


Gulf of Thailand and Andaman Sea.

**Locality:**

Trat (Khlong Yai); Chantaburi (Laem Sing); Rayong (Ban Phe); Chon Buri (Sattahip); Phuket (Patong Beach); Phangnga; Krabi; Satun.

**Description:**

Medium-sized dolphin. Distinctive dark stripe from eye to anus and dark flipper stripe. Occasionally a small auxiliary stripe can be seen behind the eye. Characteristic dark cape with shoulder blaze, pale grey sides and tailstock, and white (bright pink in some individuals) undersides. Beak is dark and prominent. Dorsal fin is dark and strongly falcate. Tooth counts range from 40 to 55 pairs of small teeth in each jaw.



**Class:** Mammalia

**Order:** Cetacea

**Family:** Delphinidae

**Scientific Name:** *Globicephala macrorhynchus*

**Common Name:** โลมาหัวกลม, วาฬน้ำร่องกริบสั้น  
Short-finned pilot whale



#### Habitat:

Off-shores.

#### Locality:

Narathiwat (Naratat Beach); Nakhon Si Thammarat; Phuket (5°55' N 97°5' E).

#### Description:

Maximum length males 5.4 m, females 4.0 m.

A robust black dolphin with thick and bulbous head. Dorsal fin located in anterior half of body. Flippers long and sickle-shaped. Tail flukes broad with deep median notch. Body black with pronounced white 'saddle' on dorsal surface behind dorsal fin; ventral with short white area from chin to upper breast, usually not beyond the level of dorsal fin. Only 7-9 peg-like teeth present in each side of both jaws.

*Peponocephala electra*

<b>Class:</b>	Mammalia
<b>Order:</b>	Cetacea
<b>Family:</b>	Delphinidae
<b>Scientific Name:</b>	<i>Peponocephala electra</i>
<b>Common Name:</b>	โลมาหัวแตงโม Melon-headed whale

**Habitat:**

Gulf of Thailand.

**Locality:**

Songkhla waters; Chon Buri.

**Description:**

Maximum length 2.7 m.

Large black dolphin having bluntly pointed beak and pointed-tipped flippers. Head triangular when viewed from above. Body overall black, except whitish on lips, around genitals and anus; ventrally a light grey patch present on chest between flippers. Dorsal fin tall, up to 30 cm, and back-swept with rounded tip. Flippers very long and pointed distally. Flukes with concave trailing edge, median notch and pointed at tips; Series of 21-24 small teeth in each side of lower jaw and 22-26 of upper jaw.



**Class:** Mammalia

**Order:** Cetacea

**Family:** Delphinidae

**Scientific Name:** *Feresa attenuata*

**Common Name:** วาฬเพชรฆาตเล็ก  
Pygmy killer whale



#### Habitat:

Andaman Sea.

#### Locality:

Trang; Phetchaburi

#### Description:

Long, slender, all black, body with a small, tapered head. Lips and snout sometimes white. Cape dark and straight along the body. Dorsal fin situated just behind the midpoint of the back and can range from clearly rounded to sharply pointed at the tip. Pectoral fins long and rounded. Tooth counts range from 8 to 13 pairs of teeth in each jaw.

*Neophocaena phocaenoides*

**Class:** Mammalia

**Order:** Cetacea

**Family:** Phocoenidae

**Scientific Name:** *Neophocaena phocaenoides*

**Common Name:** โลมาไร้ครีบทิ้ง, โลมาหัวบาตรหลังเรียบ  
Finless porpoise

**Habitat:**

Along shores, mainly Gulf of Thailand.

**Locality:**

Trang (Chaomai Beach); Ranong; Bangkok; Songkhla; Phangnga; Chon Buri (Bang Saen, Bang Sarae, Koh Si Chang, Naklua); Prachuap Khirikhan; Pattani (Thepa); Trat (Trat and Laem Ngop Districts); Chantaburi (Laem Sing District); Chumphon (Khanduli, Tha Chana Bay).

**Description:**

Total length 1.4-1.7 m.

One of the smaller cetaceans. The only true porpoise with no dorsal fin. In its place a series of tubercles runs down the back. Body colour is pale grey with a blue tinge on the back and sides, with off-white ventral regions. (The body colour only becomes black after death.) Tail flukes have slightly rounded tips and a well-defined notch. Flippers are long, broad and pointed at the tips. Tooth counts range from 13 to 22 pairs in each jaw.



<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Tragulidae
<b>Scientific Name:</b>	<i>Tragulus napu</i>
<b>Common Name:</b>	กระเจิงควาย Larger mouse deer

**Habitat:**

Lowland evergreen forest.

**Locality:**

Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Kui Buri); Trang (Khao Chong); Sakaew (Ta Phraya); Satun (Koh Tarutao); Yala (Hala-Bala); Surat Thani (Khleng Saeng).

**Description:**

Head-body 50-60 cm; tail 6-10 cm; weight 4-6 kg.

This forest-dwelling ruminant, about the size of a small dog, has slender legs and hooves, walking in a peculiar gait with high hindquarter. Coat a shade of grey buff to orange buff, giving the coarsely flecked overall appearance; underparts paler with a pale median brownish line and dark markings on the chest. White neck pattern usually appears as two separate bars. Males have enlarged canines, but always smaller in the females.



*Muntiacus feae*

<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Cervidae
<b>Scientific Name:</b>	<i>Muntiacus feae</i>
<b>Common Name:</b>	เก้งหม้อ, เก้งดำ Fea's muntjac

**Habitat:**

Evergreen forest along streams.

**Locality:**

Kanchanaburi (Thung Yai, Sai Yok, Thong Pha Phum); Phetchaburi (Kaeng Krachan); Surat Thani (Khlung Saeng); Krabi; Narathiwat (Hala-Bala).

**Description:**

A seldomly-seen barking deer has dark yellowish brown to dark brown coat; usually darker on face and legs. Blackish legs each with a whitish line down to hoof. Face has a distinct tuft of black hairs, more pronounced around the antler bases. Tail has a striking black upper part and white under part.



<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Cervidae
<b>Scientific Name:</b>	<i>Axis porcinus</i>
<b>Common Name:</b>	เนื้อทราย Hog deer

**Habitat:**

Swamp forest.

**Locality:**

Chaiyaphum (Phu Khieo); Chon Buri (Khao Chi-on); Rayong (Koh Kradat); all sites are kept in semi-captive condition.

**Description:**

Shoulder height 66-74 cm; weight 36-45 kg.

A medium-sized, low and heavily built deer with short face and legs. Coat yellow brown to dark brown according to the season, but with darker underpart. Only males have antlers, which can be recognized by the inner tine of upper fork curve more or less downwards. Fawns spotted.

*Bubalus bubalis*

<b>Class:</b>	Mammalia
<b>Order:</b>	Artiodactyla
<b>Family:</b>	Bovidae
<b>Scientific Name:</b>	<i>Bubalus bubalis</i>
<b>Common Name:</b>	ควายป่า, มหิงษา Wild water buffalo

**Habitat:**

Clearings and swamps in evergreen forest.

**Locality:**

Uthai Thani (Huai Kha Kaeng).

**Description:**

Head-body 2.4-2.8 m; tail 60-85 cm; shoulder height 1.6-1.9 m; weight 1,200 kg. (male), 800 kg (female).

Similar to its domestic counterpart, but more stout and with longer and slender horns, this dark herbivore has no dewlap or hump. A white chevron present on chest and dirty white stockings down from knees and hocks. Horns, triangular in cross section, heavy and backswept. This animal has flexible fetlock joints make it nimble in the mud.



<b>Class:</b>	Mammalia
<b>Order:</b>	Perissodactyla
<b>Family:</b>	Tapiridae
<b>Scientific Name:</b>	<i>Tapirus indicus</i>
<b>Common Name:</b>	สมเสร็จ Malayan Tapir

**Habitat:**

Evergreen forest along streams.

**Locality:**

Tak (Thung Yai, Umphang); Uthai thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra); Phetchaburi (Kaeng Krachan); Ranong (Khlong Naka); Surat Thani (Khlong Saeng, Khao Nun); Trang (Khao Bantad); Yala (Hala-Bala).

**Description:**

Head-body 2.2-2.5 m; tail 5-10 cm; weight 250-300 kg.

A stout-bodied forest herbivore with peculiar colouration of being white in middle part and black in the front and hind parts; also when walking its rump slightly higher than the shoulder. Limbs short and sturdy; front feet having four toes and the hind feet three. Neck short; snout extends into a short fleshy trunk derived from the nose and the upper lip, with nostrils at the tip. Ears often tipped with white.

*Elephas maximus*

<b>Class:</b>	Mammalia
<b>Order:</b>	Uranotheria or Proboscidea
<b>Family:</b>	Elephantidae
<b>Scientific Name:</b>	<i>Elephas maximus</i>
<b>Common Name:</b>	ช้างป่า Asian elephant

**Habitat:**

Evergreen forest to mixed deciduous forest.

**Locality:**

Mae Hong Son (San Pandan, Huai Nam Dung); Chiang Mai (Om Koi, MaeTuen); Nan (Doi Phu Kha); Tak (Doi Mon Chong, Thun Yai Naresuan); Kamphaeng Phet (Khlung Lan); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra, Sai Yok, Erawan, Thong Pha Phum, Khao Laem); Phetchaburi (Kaeng Krachan); Uttaradit (Phu Miang-Phu Thong); Phitsanulok (Thung Salaeng Luang); Phetchabun (Nam Nao); Chaiyaphum (Phu Khieo, Ta Bak-Huai Yai); Nakhon Ratchasima (Khao Yai); Prachin Buri (Pang Sida); Sakaew (Ta Praya); Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jong-Na Yoi); Prachuap Khirikhan (Kui Buri); Surat Thani (Khlung Saeng, Tai Rom Yen); Phangnga (Khao Sok); Songkhla (Ton Nga Chang, Saba Yoi); Narathiwat/Yala (Hala-Bala, Bang Lang, Huai Sai Khao, Khao Chi Po); Loei (Phu Luang, Phu Kradung); Nong Khai (Phu Wua); Sakon Nakhon (Phu Phan); Buriram (Dong Yai); Chachoengsao (Khao Ang Rue Nai); Rayong (Khlung Krue Whai, Khao Chamao-Khao Wong); Chanthaburi (Khao Soi Dao).

**Description:**

The largest terrestrial mammal in Thailand with a very large, round body and long trunk. The skin dark and thick with short hairs which cover the body and large ears. Both sexes can have tusks, but those of males always larger than those of females.

# Status VU

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ratufa affinis</i>	พญากระรอกเหลือง Cream-coloured Giant Squirrel	Lowland swamp forest to evergreen forest.	Chumphon (Phru Kantuli); Trang; Narathiwat (Pa Phru Toh Dang); Phattalung (Khao Phap Pha); Nakhon Si Thammarat (Lan Saka); Satun (Thale Bun).
<i>Rhinosciurus laticaudatus</i>	กระรอกหน้ากระแต Shrew-faced Ground Squirrel	Evergreen forest, mainly on forest floors.	Nakhon Si Thammarat (Khao Luang); Surat Thani (Khlung Saeng); Narathiwat (Pa Phru Toh Dang).
<i>Lariscus insignis</i>	กระรอกดินหลังลาย Three-striped Ground Squirrel	Evergreen forest.	Trang (Khao Chong); Yala.
<i>Hylopetes spadiceus</i>	กระรอกบินแกมสีแดง Red-cheeked Flying Squirrel	Dry evergreen forest to mixed deciduous forest.	Sakon Nakhon (Phu Phan); Satun (Koh Terutau).
<i>Hylopetes alboniger</i>	กระรอกบินเล็กขาวสูง Particoloured Flying Squirrel	Hill evergreen forest.	Chiang Mai (Doi Inthanon); Loei (Dan Sai); Chaiyaphum (Phu Khieo).
<i>Petinomys vordermanni</i>	กระรอกบินจิวมลายู Vordermann's Flying Squirrel	Lowland swamp forest.	Narathiwat (Pa Phru Toh Dang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Petaurista elegans</i>	พญากระรอกบินหูดำ Spotted Giant Flying Squirrel	Evergreen forest.	Mae Hong Son (Salween); Chiang Mai (Doi Pui, Doi Inthanon); Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Chawang); Yala (Hala-Bala).
<i>Belomys pearsonii</i>	กระรอกบินเท้าขน Hairy-footed Flying Squirrel	Dry evergreen forest to human habitations.	Nakhon Ratchasima (Sakaerat).
<i>Eothenomys miletus</i>	หนูน้ำคอกอ่างกา South Chinese Vole	Hill evergreen forest along mountain streams.	Chiang Mai (Doi Inthanon, Doi Pha Hom Pok).
<i>Hapalomys longicaudatus</i>	หนูไฟเล็บแม่มือแบน Marmoset Rat	Evergreen forest.	Nakhon Sawan; Kanchanaburi; Ratchaburi (Lumnam Pachi).
<i>Rattus remotus</i>	หนูเกาะ Island Rat	Near waterfalls in dry evergreen forest.	Chumphon (Koh Tao); Surat Thani (Koh Pangan, Koh Samui); Nakhon Si Thammarat (Koh Kra); Chon Buri (Koh Kram, Koh Kra); Trat (Koh Kradat).
<i>Maxomys whiteheadi</i>	หนูฟานเล็ก Whitehead's Rat	Evergreen forest.	Trang.
<i>Leopoldamys edwardsi</i>	หนูเขาสูง Edwards' Rat	Hill evergreen forest.	Chiang Mai, Tak.
<i>Leopoldamys neilli</i>	หนูถ้ำ Neill's Rat	Limestone caves.	Saraburi (Kaeng Khoi); Kanchanaburi (Sai Yok); Surat Thani (Khlung Saeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Neofelis nebulosa</i>	เสือลายเมฆ Clouded Leopard	Evergreen forest.	Loei (Phu Luang); Nakhon Ratchasima (Khao Yai); Chaiyaphum (Phu Khieo); Kanchanaburi (Thung Yai); Surat Thani (Klong Saeng); Srakaew (Ta Phraya); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Kaeng Krachan).
<i>Prionailurus viverrinus</i>	เสือปลา Fishing Cat	Riverine forest.	Nakhon Ratchasima (Khao Yai); Kanchanaburi (Salak Phra); Samut Sakhon (Mahachai); Phatthalung (Thale Noi); Satun (Kuan Kalong).
<i>Profelis temminckii</i>	เสือไฟ Asiatic Golden Cat	Evergreen forest.	Mae Hong son (Lumnam Pai, Namtok Mae Surin); Chiang Mai (Chiang Dao); Chaiyaphum (Phu Khieo); Kanchanaburi (Salak Phra, Sri Sawat, Thung Yai); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Kaeng Krachan); Surat Thani (Khlung Saeng).
<i>Panthera pardus</i>	เสือดาว, เสือดำ Leopard, Panther	Evergreen forest.	Tak (Thung Yai); Chaiyaphum (Phu Khieo), Ubon Ratchathani (Phu Jong-Na Yoi, Yot Dom); Kanchanaburi (Salak Phra); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Kaeng Krachan); Prachuab Khirikhan (Kui Buri); Surat thani (Khlung Saeng); Yala (Hala-Bala).
<i>Prionodon linsang</i>	ชะมดแปดลายแถบ Banded Linsang	Evergreen forest.	Uthai Thani (Huai Kha Kaeng); Yala (Than To, Hala-Bala); Phetchaburi (Kaeng Krachan); Surat Thani (Khlung Saeng).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Canis aureus</i>	หมาจิ้งจอก Golden Jackal	Mixed deciduous forest to dry evergreen forest, also around human dwellings.	Tak (Thung Yai, Muang, Om Koi); Kanchanaburi (Salak Phra, Thung Yai); Phetchaburi (Kaeng Krachan); Chaiyaphum (Phu Khieo); Phetchabun (Nam Nao); Nakhon Ratchasima (Khao Yai, Sakaerat); Nakhon Phanom; Chachoengsao (Khao Ang Rue Nai); Srakaew (Ta Phraya); Phitsanulok (Wang Thong).
<i>Cuon alpinus</i>	หมาใน Dhole	Evergreen forest.	Tak (Thung Yai Naresuan); Kamphaeng Phet (Khlung Lan); Loei (Phu Luang); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra); Phetchaburi (Kaeng Krachan); Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Khao Yai); Chachoengsao (Khao Ang Rue Nai); Prachin Buri (Tub Lan) Srakaew (Ta Phraya); Prachuap Khirikhan (Kui Buri); Surat Thani (Khlung Saeng).
<i>Ursus thibetanus</i>	หมีควาย Asian Black Bear	Evergreen forest.	Tak (Thung Yai Naresuan); Nakhon Sawan (Mae Wong); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra, Thung Yai); Phetchaburi (Kaeng Krachan); Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Khao Yai); Srakaew (Ta Phraya); Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jong-Na Yoi, Yot Dom); Prachuap Khirikhan (Kui Buri); Surat Thani (Khlung Saeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Helarctos malayanus</i>	หมีหมา Sun Bear	Evergreen forest.	Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Khao Yai); Ubon Ratchathani (Bun Thrik-Yot Mon, Phu Jong-Na Yoi, Yot Dom); Chanthaburi (Khao Soi Dao); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Thung Yai, Salak Phra); Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Kui Buri); Surat Thani (Khlung Saeng); Narathiwat (Pa Phru Toh Dang).
<i>Lutrogale perspicillata</i>	นากใหญ่ขนเรียบ Indian Smooth-coated Otter	Evergreen forest along streams.	Uthai Thani (Huai Kha Kaeng); Kampaeng Phet (Khlung Lan); Nakhon Sawan (Mae Wong); Ubon Ratchathani (Mekong River); Chachoengsao (Khao Ang Rue Nai); Kanchanaburi (Salak Phra, Thung Yai, Sri Nakarin); Nakhon Ratchasima (Khao Yai); Phetchaburi (Kaeng Krachan); Surat Thani (Nong Thung Tong, Mookoh Ang Thong); Krabi; Phangnga (Phangnga Bay); Trang (Hat Chao Mai); Yala (Bang Lang, Hala-Bala).
<i>Mustela sibirica</i>	เพียงพอนเหลื่อง Kolinsky	Dry evergreen forest to savana.	Chaiyaphum (Phu Khieo); Loei (Phu Luang); Nakhon Ratchasima (Sakaerat).
<i>Mustela nudipes</i>	เพียงพอนเล็กตีน้าตาล Barefoot Weasel	Lowland evergreen forest.	Ranong (Ban Bang Non); Surat Thani (Khlung Saeng); Krabi (Khao Nor Chuchi); Trang (Khao Chong); Narathiwat (Pa Phru Toh Dang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Echinosorex gymmurus</i>	หนูเหม็น Greater Moonrat	Lowland evergreen forest.	Trang (Khao Chong); Surat Thani (Khlong Saeng); Phatthalung (Ban Tamod); Krabi.
<i>Anourosorex squamipes</i>	หนูผีป่าหางจู้ Mole-shrew	Hill evergreen forest on forest floors.	Chiang Mai (Doi Inthanon, Doi Pha Hom Pok).
<i>Pteropus vampyrus</i>	ค้างคาวแม่ไก่ป่าฝน Large Flying Fox	Evergreen forest on islands.	Chon Buri (Koh Kram Yai); Trat (Koh Kut, Koh Mesan, Koh Chang); Nakhon Ratchasima; Nakhon Si Thammarat (Chawang); Chumphon (Lang Suan); Krabi (Muang).
<i>Dyacopterus spadiceus</i>	ค้างคาวด้ายค Dayak Fruit Bat	Lowland swamp forest.	Narathiwat (Pa Phru Toh Dang).
<i>Sphaerias blanfordi</i>	ค้างคาวดอย Blanford's Fruit Bat	Hill evergreen forest in caves.	Chiang Mai (Doi Inthanon, Doi Pha Hom Pok, Doi Ang Khang).
<i>Taphozous saccolaimus</i>	ค้างคาวปีกถุงปลอม Blyth's Tomb Bat		Surat Thani (Khlong Saeng).
<i>Rhinopoma microphyllum</i>	ค้างคาวหางหนู Greater Mouse-tailed Bat	Seaside caves.	Krabi.
<i>Nycteris tragata</i>	ค้างคาวหน้าร่อง Slit-faced Bat	Lowland evergreen forest.	Yala (Bang Lang); Narathiwat (Hala-Bala).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rhinolophus paradoxolophus</i>	ค้างคาวมงกุฎหูโตใหญ่ Bouret's Horseshoe Bat	Dry evergreen forest.	Chiang Mai (Doi Ang Khang); Chaiyaphum (Phu Khieo).
<i>Rhinolophus robinsoni</i>	ค้างคาวมงกุฎเดือนมลายูเล็ก Robinson's Horseshoe Bat	Evregreen forest in limestone caves.	Surat Thani (Khao Nawng); Chiang Mai (Doi Ang Khang).
<i>Hipposideros turpis</i>	ค้างคาวหน้ายักษ์กุ่มกกรวม Lesser Leaf-nosed Bat	Lowland evergreen forest in caves with streams.	Nakhon Si Thammarat (Khao Luang); Krabi (Ban Tap Plick).
<i>Hipposideros lekaguli</i>	ค้างคาวหน้ายักษ์หมอบูญสูง Lekagul's Leaf-nosed Bat	Evergreen forest in limestone caves.	Saraburi (Kaeng Khoi); Nan (Tham Song Satang); Phrae (Rong Kwang); Nakhon Sawan (Nong Ben); Chaiyaphum (Tung Luei Lai); Srakaew (Tham Khao Sing-to); Chanthaburi (Khao Wong Kot); Pattani (Sai Khao).
<i>Coelops frithi</i>	ค้างคาวอัยแหวง Tailless Leaf-nosed Bat	Evergreen forest.	Phetchabun (Nam Nao); Chaiyaphum (Phu Khieo).
<i>Cheiromeles torquatus</i>	ค้างคาวขนช้าง Hairless Bat	Lowland evergreen forest.	Satun (Koh Tarutao); Narathiwat (Hala-Bala).
<i>Tadarida teniotis</i>	ค้างคาวหูหนา European Free-tailed Bat	Hill evergreen forest to valleys	Loei (Phu Luang).
<i>Myotis chinensis</i>	ค้างคาวหูหนูยักษ์ Large Myotis	Hill evergreen forest with extensive savana.	Nan (Muang); Chiang Mai (Doi Ang Khang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Myotis altarium</i>	ค้างคาวหูหนูคอดอขาง South Chinese Myotis	Hill evergreen forest in limestone caves.	Chiang Mai (Doi Ang Khang).
<i>Nyctalus noctula</i>	ค้างคาวกินแมลงนิ้วสั้น Noctule	Hill evergreen forest.	Chiang Mai (Doi Pha Hom Pok).
<i>Ia io</i>	ค้างคาวอี้ออี้อ Great Evening Bat	Lowland evergreen forest.	
<i>Eptesicus serotinus</i>	ค้างคาวทองสีน้ำตาลใหญ่ Serotine	Hill evergreen forest to valleys.	Chiang Mai (Hot, Doi Chiang Dao); Tak (Thung Yai Naresuan); Uthai Thani (Huai Kha Kaeng).
<i>Eptesicus pachyotis</i>	ค้างคาวทองสีน้ำตาลหูหนา Thick-eared Bat	Evergreen forest, mainly hill evergreen forest.	Chiang Mai (Doi Par Sakeng).
<i>Hesperoptenus tickelli</i>	ค้างคาวพันทนาซอนใหญ่ Tickell's Bat	Evergreen forest.	Phetchaburi (Kaeng Krachan).
<i>Hesperoptenus blanfordi</i>	ค้างคาวพันทนาซอนเล็ก Blanford's Bat	Evergreen forest.	Phetchaburi (Kaeng Krachan).
<i>Scotomanes ornatus</i>	ค้างคาวคอดหลังลายขาว Harlequin Bat	Hill evergreen forest.	Chiang Mai (Doi Inthanon).
<i>Murina leucogaster</i>	ค้างคาวจมูกหลอดทองขาว Greater Tube-nosed Bat	Evergreen forest, mainly dry evergreen forest.	Uthai Thani (Huai Kha Kaeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Harpiocephalus harpia</i>	ค้างคาวปีกขนไต Hairy-winged Bat	Dry evergreen forest.	Prachin Buri (Thap Lan); Chanthaburi (Khao Soi Dao).
<i>Kerivoula whiteheadi</i>	ค้างคาวยอดคดกล้วยป่า Whitehead's Forest Bat	Lowland evergreen forest.	Yala (Jalor).
<i>Kerivoula minuta</i>	ค้างคาวยอดคดกล้วยเล็ก Least Forest Bat	Evergreen forest.	Trang.
<i>Kerivoula papillosa</i>	ค้างคาวยอดคดกล้วยปีกปุ่ม Papillose Bat	Lowland evergreen forest.	Surat Thani (Tha Chang).
<i>Kerivoula hardwickei</i>	ค้างคาวยอดคดกล้วยปีกใส Hardwicke's Forest Bat	Lowland evergreen forest.	Surat Thani (Khlung Saeng).
<i>Phoniscus atrox</i>	ค้างคาวฟันร่อง Groove-toothed Bat	Evergreen forest.	Unknow.
<i>Macaca arctoides</i>	ลิงเสน Stump-tailed Macaque	Dry evergreen forest.	Tak (Thung Yai Naresuan); Kanchanaburi (Salak Phra); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Khao Tao Mor); Chanthaburi; Ubon Ratchathani (Phu Jong-Na Yoi); Surat Thani (Khlung Saeng, Khao Sok, Khao Nun); Trang (Khao Chong, Angut, Rum Phura, Ban Palian); Krabi (Ban Nong Kok).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Presbytis femoralis</i>	ค่างดำ Banded Sureli	Evergreen forest.	Trang (Ko Khau, Khao Chong, Ban Palian); Chumphon (Ban Tha San); Nakhon Si Thammarat (Ron Phibun, Khao Luang, Khao Wang Hip); Phatthalung (Khao Phap Pha); Phuket (Khlong Tung Sai); Phetchaburi (Kaeng Krachan); Surat Thani (Khao Nawng); Narathiwat (Pa Phru Toh Dang).
<i>Semnopithecus phayrei</i>	ค่างแว่นถิ่นเหนือ Phayre's Leaf Monkey	Evergreen forest with limestone hills	Chiang Mai (Maenam Ping rapids, Chiang Dao, Doi Inthanon); Tak (Mae Sod, Thung Yai Naresuan, Tha Chang Tai, Ban Mae Lamao, Umphang); Kamphaeng Phet (Ban Pong Nam Ron, Ko Keow, Khlong Khlung); Uthai Thani (Huai Kha Kaeng, Kata Taek); Nakhon Sawan (Pak Nam Pho, Me Wong); Kanchanaburi (Ban Muang Bow Ngam, Sisawat, Salak Phra); Phetchabun (Nam Nao); Phisanulok (Thung Salaeng Luang); Chiyaphum (Phu Khieo); Nakhon Ratchasima (Lat Bua Kao); Loei (Muang, Tham Pha Pu, Phu Luang).
<i>Semnopithecus obscurus</i>	ค่างแว่นถิ่นใต้ Dusky Leaf Monkey	Evergreen forest with limestone hills.	Chumphon (Ban Pak Nam, Ban Tha San, Khlong Bang Lai, Moo Koh Chumphon); Prachuap Khiri Khan (Bang Saphan, Sam Roi Yot); Phetchaburi (Kaeng Krachan); Ratchaburi (Hat Sanuk); Ranong (Koh Kam Yai); Surat Thani (Khlong Saeng, Ban Khok Klap, Ban Makhm Tia, Koh Phangan, Moo Koh

Scientific Name	Thai name/Common name	Habitat	Site
<i>Hylobates lar</i>	ชะนีมือขาว Lar Gibbon	Evergreen forest.	<p>Ang Thong, Tai Rom Yen); Nakhon Si Thammarat (Khao Luang, Ban Krung Khayan, Khao Wang Hip, Ban Bha Ngan); Songkhla (Ton Nga Chang, Khao Plak Phaya); Trang (Khao Chong, Ban Palian); Phangnga (Koh Yao Yai, Takau Thung, Telok Poh); Krabi (Ban Nong Kok, Koh Lanta, Ban Thap Plik, Kuan Ying Mee, Khao Na Nong, Ban Kuan Daeng); Satun (Koh Tarutao); Narathiwat.</p> <p>Mae Hong Son (Nam Pai, San Pan Dan, Namtok Mae Surin); Chiang Mai (Doi Inthanon, Doi Chiang Dao); Tak (Thung Yai Naresuan, Huai Nua Pla, Sam Ngao, Umphang); Kamphaeng Phet (Me Wong); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra, Si Sawat); Ratchaburi (Maenam Phachi); Phetchaburi (Kaeng Krachan); Phitsanulok (Thung Salaeng Luang); Phetchabun (Nam Nao); Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Khao Yai); Ranong (Tabli, Ban Klong Wan); Chumphon (Ban Tha San); Nakhon Si Thammarat (Khao Luang); Surat Thani (Khlung Saeng, Tha Chana); Trang (Khao Chong); Krabi (Ban Nong Kok, Ban Thap Plik); Phangnga (Takua Thung); Phuket (Khlung Tung Sai).</p>



Scientific Name	Thai name/Common name	Habitat	Site
<i>Dendrogale murina</i>	กระแตหางหนู Northern Smooth-tailed Treeshrew	Lowland evergreen forest.	Trat (Khao Saming, Khlong Yai); Chanthaburi.
<i>Ptilocercus lowii</i>	กระแตหางขนนก Pen-tailed Treeshrew	Lowland evergreen forest.	Satun; Yala; Narathiwat.
<i>Pseudorca crassidens</i>	โลมาเพชรฆาตแปดง, วาฬเพชรฆาตดำ False Killer Whale	Off-shores.	Prachuap Khirikhan (Bang Saphan Noi); Trat (Laem Ngop); Surat Thani (Kanchanadit); Phangnga (Similan Islands); Krabi (Ban Pak Khlong); Phangnga (Similan); Phuket.
<i>Tursiops aduncus</i>	โลมาปากขวด Indo-Pacific Bottle-nosed Dolphin	Along shores; both waters.	Chachoengsao (Bangpakong River); Rayong (Ban Pae Beach, Laem Mae Phim); Nakhon Si Thammarat (Tha Sala, Si Chon); Ranong; Trang (Chao Mai Beach); Phangnga (Khai Nok Island); Phuket (Maiton Islands); Chonburi (Bang Saen, Sattahip); Songkhla; Prachuab Khirikhan (Hua Hin); Satun (Tarutao); Trat; Chantaburi (Laem Sing, Paknam Prasae).
<i>Stenella longirostris</i>	โลมากระโดด Spinner Dolphin	Gulf of Thailand.	Bangkok Harbour; Chonburi (Bang Saen, Sattahip); Samut Songkram; Samut Sakhon; Phuket (7°52'N 9°27'E); Trat (Laem Ngop); Chantaburi (Laem Sing); Rayong (Ban Phe and Rayong Districts, Koh Samet); Songkhla (Songkhla water).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Sousa chinensis</i>	โลมาขาวเทา, โลมาเผือก, โลมาหลังโหนก Indo-Pacific Humpbacked Dolphin	Along shores, mainly Gulf of Thailand.	Chachoengsao (Bangprakong); Rayong (Ban Phe); Chonburi (Laem Chabang, Sattahip, Naklua, Bang Lamung); Samut Sakon; Phetchaburi; Chumphon; Krabi (Ao Luk); Trang (Trang river, Kantang); Prachuap Khirikhan; Rayong (Koh Samet); Trat (near Khlong Yai, Trat and Laem Ngop Districts); Bangkok (Bangkok Harbour); Chantaburi (Laem Sing, Pak Nam Prasae).
<i>Bos gaurus</i>	กระทิง Gaur	Evergreen forest.	Tak (Thung Yai Naresuan); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Salak Phra); Phetchaburi (Kaeng Krachan); Ratchaburi (Maenam Pachi); Phisanulok (Thung Salaeng Luang); Chaiyaphum (Phu Khieo); Phetchabun (Nam Nao); Nakhon Ratchasima (Khao Yai, Tab Lan, Khao Phang Ma); Prachin Buri (Pang Sida); Chachoengsao (Khao Aug Ru Nai); Chanthaburi (Khao Soi Dao); Srakaew (Ta Phraya); Surat Thani (Khlong Saeng); Ubon Ratchathani (Yot Dom, Phu Jong-Na Yoi).

# Status NT

Scientific Name	Thai name/Common name	Habitat	Site
<i>Manis javanica</i>	คินชาวา Malayan Pangolin	All forest types to human habitations and agricultural areas.	Chiang Mai (Doi Suthep), Tak (Thung Yai), Kanchanaburi (Salak Phra, Erawan, Sai Yok); Uthai Thani (Huai Kha Kaeng); Phetchaburi (Kaeng Krachan); Phitsanulok (Thung Salaeng Luang); Prachub Khiri Khan (Kui Buri); Ranong (Khlong Naka); Surat Thani (Khlong Saeng); Pangnga (Khao Sok, Khao Lak-Lum Ru); Songkhla (Ton Nga Chang); Yala (Bang Lang, Hala-Bala); Sakon Nakhon (Phu Phan); Nakhon Ratchasima (Khao Yai, Sakaerat); Chachoengsao (Khao Ang Ru Nai); Chanthaburi (Khao Soi Dao).
<i>Crocidura pullata</i>	หนูสีอ่างกา White-toothed Shrew	Evergreen forest along streams.	Chiang Mai (Doi Inthanon).
<i>Pteropus hypomelanus</i>	ค้างคาวแม่ไก่เกาะ Small Flying Fox	Evergreen forest on mainland and islands.	Surat Thani (Koh Samui, Koh Pangan, Koh Tan); Chumphon (Koh Tao); Krabi (Plai Phraya); Chon Buri (Koh Kram); Chanthaburi (Tha Mai); Trat (Koh Chang); Rayong (Ban Phe, Koh Samet, Koh Nom Sao).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Pteropus lylei</i>	ค้างคาวแม่ไก่ภาคกลาง Lyle's Flying Fox	Around human dwelling and temples.	Bangkok (Yanawa); Chachoengsao (Bang Khla); Ang Thong (Chai Yo); Nakhon Nayok (Ongkharak); Chon Buri (Ang Sila, Si Racha, Phanat Nikhom); Suphan Buri (Sam Chuk); Phra Nakhon Si Ayutthaya (Wat Tan En).
<i>Cherephon johorensis</i>	ค้างคาวปากขนมลายู Dato Meldrum's Bat	Lowland evergreen forest.	Narathiwat (Hala-Bala).
<i>Tadarida mops</i>	ค้างคาวหูหนามมลายู Malayan Free-tailed Bat	Lowland evergreen forest.	Narathiwat (Hala-Bala).
<i>Myotis montivagus</i>	ค้างคาวหูหนูปม่า Burmese Whiskered Bat	Hill evergreen forest to dry evergreen forest.	Chiang Mai (Doi Pha Hom Pok); Loei (Phu Luang).
<i>Myotis annectans</i>	ค้างคาวหูหนูหนาน Hairy-faced Bat	Hill evergreen forest in limestone caves.	Chiang Mai (Doi Pui).
<i>Pipistrellus pulveratus</i>	ค้างคาวลูกหนูถ้ำ Chinese Pipistrelle	Hill evergreen forest in limestone caves.	Chiang Mai (Doi Ang Khang, Mae Taeng); Khon Kaen (Chum Phae); Lop Buri (Tham Wat Phra That).
<i>Pipistrellus cadornae</i>	ค้างคาวกรมหนานบนถ้ำ Thomas' Pipistrelle	Hill evergreen forest in limestone caves.	Phetchabun (Thung Salaeng Luang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Murina huttonii</i>	ค้างคาวจมูกหลอดหูขาว Hutton's Tube-nosed Bat	Hill evergreen forest.	Chiang Mai (Doi Pui, Doi Inthanon).
<i>Macaca nemestrina</i>	ลิงกังใต้ Pig-tailed Macaque	Lowland evergreen forest.	Chumphon (Ban Tha San, Phru Kanthuri); Surat Thani (Khlung Saeng); Trang (Lam Phura, Khao Chong); Phatthalung (Huai Lian); Phangnga (Koh Yao Yai); Krabi (Ban Thap Plik, Ban Nong Kok); Phuket (Klong Tung Sai); Narathiwat (Pa Phru Toh Dang, Hala-Bala).
<i>Semnopithecus germaini</i>	ค้างหงอก Silvered Leaf Monkey	Evergreen forest.	Chaiyaphum (Phu Khieo); Nakhon Ratchasima (Pak Chong, Lat Bua Khao); Ubon Ratchathani (Yot Dom, Phu Jong-Na Yoi, Bun Thrik-Yot Mon); Trat (Koh Chang, Khlung Yai); Chachoengsao (Ban Pakong); Kanchanaburi (Huai Maenam Noi, Sangkla Buri); Ratchaburi (Mae Nam Phachi).
<i>Hylobates syndactylus</i>	ชะนีใหญ่ Siamang	Moist evergreen forest.	Narathiwat (Hala-Bala).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Capricornis sumatraensis</i>	เลียงผา Mainland Serow	Evergreen forest with hills.	Tak (Thung Yai Naresuan); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Sai Yok, Salak Phra); Phetchaburi (Kaeng Krachan); Chaiyaphum (Phu Khieo); Loei (Phu Kradung, Phu Luang); Ubon Ratchathani (Yot Dom, Bun Thrik-Yot Mon, Phu Jong-Na Yoi); Surat Thani (Khlung Saeng); Nakhon Ratchasima (Khao Yai, Sakaerat); Chachoengsao (Khao Ang Rue Nai); Chanthaburi (Khao Soi Dao, Khao Sabab); Rayong (Khao Chamao); Prachuap Khirikhan (Sam Roi Yot); Nakhon Si Thammarat (Khao Luang); Surat Thani (Khao Nun); Trang (Khao Chong); Songkhla (Ton Nga Chang).

# Status LC

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ratufa bicolor</i>	พญากระรอกดำ Black Giant Squirrel	Evergreen forest.	Chiang Mai (Doi Suthep-Doi Pui, Doi Hua Mot, Doi Nangka, Doi Nangkeo, Doi Inthanon, Doi Chiang Dao, Mae Taeng); Lampang (Koon Tarn); Nan (Doi Phu Kha); Phrae (Mae Yom Valley); Tak (Mae Taw, Umphang, Tee Lor Su); Kamphangphet (Khlong Klung, Me Ping river, Wang Pratart); Nakhon Sawan (Mae Wong); Uthai Thani (Huai Kha Kaeng); Kanchanaburi (Sai Yok, Baw Ploy, Salak Phra); Ratchaburi (Suan Phung); Chaiyaphum (Phu Khieo); Phisanulok (Thung Salaeng Luang); Phetchabun (Nam Nao); Nakhon Ratchasima (Khao Yai, Sakaerat); Ubon Ratchathani (Phibun Mangsahan, Chong Mek); Saraburi (Hip Lap); Prachin Buri (Krabin Buri); Chon Buri (Siracha, Nong Khor, Hup Bon); Chanthaburi (Khao Sabab); Trat (Laem Ngop, Khlong Yai, Koh Kut); Surat Thani (Koh Samut, Koh Pengan, Khlong Saeng); Nakhon Si Thammarat (Chawang, Ban Ta-Phae); Trang (Telibon, Lay Song Hong); Narathiwat (Pa Phru Toh Dang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Petaurista petaurista</i>	พญากระรอกบินหูแดง Red Giant Flying Squirrel	Evergreen forest.	Chiang Mai (Fang, Mae Taeng); Mae Hong Son (Pai, Mae Sariang); Loei (Dan Sai, Phu Luang); Chaiyaphum (Phu Khieo); Saraburi (Tub Kwang); Nakhon Ratchasima (Khao Yai); Chon Buri (Siracha); Surat Thani (Ban Don, Khlong Saeng, Khao Ta Phet); Nakhon Si Thammarat (Thung Song); Satun (Koh Teratau); Pattani (Sai Khao).
<i>Petaurista philippensis</i>	พญากระรอกบินหูดำหางสีเข้ม Black-tailed Giant Flying Squirrel	Evergreen forest to lowland dry evergreen forest.	Mae Hong Son (Mae Sariang); Tak ( Melamung); Chiang Mai (Doi Suthep).
<i>Petaurista alborufus</i>	พญากระรอกบินหูขาว Red and White Giant Flying Squirrel	Hill evergreen forest.	Pitsanulok (Thung Salaeng Luang); Loei (Dan Sai).
<i>Rattus annandalei</i>	หนูป่าพรุ Annandale's Rat	Lowland swamp forest.	Narathiwat (Pa Phru Toh Dang); Yala (Hala-Bala).
<i>Chiromyschiropus</i>	หนูไม้เล็บแม่มือแบน Fea's Tree Rat	Clearing in hill evergreen forest.	Chiang Mai (Doi Inthanon, Doi Nangkeo); Loei (Dan Sai, Phu Luang).
<i>Rhizomys pruinosus</i>	อนกลาง Hoary Bamboo Rat	Evergreen forest.	Chanthaburi.



Scientific Name	Thai name/Common name	Habitat	Site
<i>Hystrix brachyura</i>	เม่นใหญ่ Malayan Porcupine	Evergreen forest to deciduous forest.	Chiang Mai (Doi Inthanon, Doi Suthep); Tak (Thung Yai); Nakhon Ratchasima (Khao Yai); Chaiyaphum (Phu Khieo); Kanchanaburi (Salak Phra); Phetchaburi (Kaeng Krachan); Ranong (Khlung Naka); Phangnga (Khao Sok); Trang (Khao Chong); Phattalung (Khao Pu-Khao Ya).
<i>Atherurus macrourus</i>	เม่นหางพวง Asiatic Brush-tailed Porcupine	Evergreen forest in limestone caves or holes.	Chiang Mai (Doi Suthep); Nong Khai (Phu Wua); Chaiyaphum (Phu Khieo); Kanchanaburi (Thung Yai); Sutun (Koh Terutao); Tak (Thung Yai); Surat Thani (Khlung Saeng, Khao Nun); Narathiwat (Pa Phru Toh dang); Phetchaburi (Kaeng Krachan); Loei (Phu Luang).
<i>Cynopterus horsfieldii</i>	ค้างคาวขอบหูขาวใหญ่ Horsfield's Fruit Bat	Lowland evergreen forest.	Surat Thani (Khao Tha-pet); Nakhon Si Thammarat (Thung Yai, Khao Luang); Trang (Khao Chong); Satun (Muang); Songkhla (Ton Nga Chang); Yala (Bang Lang).

**Status DD**

Scientific Name	Thai name/Common name	Habitat	Site
<i>Callosciurus nigrovittatus</i>	กระรอกขางลายทองเทา Black-banded Squirrel	Lowland evergreen forest around swamps.	Narathiwat (Pa Phru Toh Dang); Yala (Hala-Bala).
<i>Pteromyscus pulverulentus</i>	กระรอกบินสีเขม่า Smoky Flying Squirrel	Lowland swamp forest	Narathiwat (Pa Phru Toh Dang).
<i>Rattus osgoodi</i>	หนูภูหลวง Osgood's Rat	Hill evergreen forest in clearings.	Loei (Phu Luang).
<i>Mustela nivalis</i>	เพียงพอนดอยผาหม่นปก European Common Weasel	Hill evergreen forest.	Chiang Mai (Doi Pha Hom Pok).
<i>Soriculus parca</i>	หนูตีหางขาวฟันแดง Brown-toothed Shrew	Evergreen forest along streams.	Chiang Mai (Doi Inthanon).
<i>Crocidura monticola</i>	หนูผีภูเขา Sunda Shrew	Hill evergreen forest.	Nakhon Si Thammarat (Khao Luang).
<i>Suncus malayanus</i>	หนูตีป่า Malay Shrew	Lowland evergreen forest.	Pattani (Bang Nara).
<i>Pteropus intermedius</i>	ค้างคาวแม่ไก่นครสวรรค์ Intermediate Flying Fox	Around human dwelling near evergreen forest.	Nakhon Sawan (Ban Yang En).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cynopterus</i> sp.	ค้างคาวขอบหูขาวเล็กป่า Dog-faced Fruit Bat	Evergreen forest.	Yala (Hala-Bala).
<i>Rhinolophus rouxii</i>	ค้างคาวมงกุฎอินเดีย Indian Horseshoe Bat	Evergreen forest clearings and edges.	Chon Buri (Siracha).
<i>Rhinolophus siamensis</i>	ค้างคาวมงกุฎหูโตเล็ก Siamese Horseshoe Bat	Evergreen forest in limestone caves.	Chiang Mai (Tham Tap Tao); Chanthaburi (Khao Soi Dao).
<i>Coelops robinsoni</i>	ค้างคาวอายุหางเล็ก Malayan Tailless Leaf-nosed Bat	Evergreen forest on islands.	Satun (Koh Tarutao).
<i>Miniopterus medius</i>	ค้างคาวปีกพับกลาง Southeast Asian Bent-winged Bat	Evergreen forest.	Nakhon Si Thammarat (Thung Yai); Satun (Koh Tarutao).

# Status Endemic

Scientific Name	Thai name/Common name	Habitat	Site
<i>Leopoldamys neilli</i>	หนูถ้ำ Niell's Rat	Limestone caves.	Saraburi (kaeng Khoi); Kanchanaburi (Sai Yok); Surat Thani (Khleng Saeng).
<i>Niviventer hinpoon</i>	หนูเขินเขินเขาหินปูน Limestone Rat	Limestone caves.	Saraburi (Kaeng Khoi).
<i>Hipposideros halophyllus</i>	ค้างคาวหน้ายักษ์งูมกุ่ม Thai Horseshoe Bat	Limestone caves in agricultural areas.	Srakaew (Tham Khao Sing- to); Lop Buri (Khao Samokhon); Uthai Thani (Huai Kha Kaeng).
<i>Eptesicus demissus</i>	ค้างคาวท้องสีน้ำตาลสุราษฎร์ Surat Serotine	Evergreen forest.	Surat Thani (Ban Don); Phangnga (Khao Sok).
<i>Cervus schomburgki</i>	เนื้อสมัน Schomburgk's Deer	Lowland floodplain of rivers.	Formerly throughout the Central Chao Phraya River Floodplain, west to Uthai Thani, east to Nakhon Nayok, Prachin Buri, Chachoengsao, Bangkok and Samut Prakarn. Now entirely extirpated from Thailand.



# REPTILES

EX = Extinct  
EW = Extinct in the Wild  
CR = Critically Endangered  
EN = Endangered  
VU = Vulnerable  
NT = Near Threatened  
LC = Least Concern  
DD = Data Deficient  
Endemic

WS = Wildlife Sanctuary  
NH = Non-hunting Area  
NP = National Park



**Class:** Reptilia  
**Order:** Crocodylia  
**Family:** Crocodylidae  
**Scientific Name:** *Tomistoma schlegelii*  
**Common Name:** ตะโขง  
 False Gavia

**Habitat:**

Inhabits short and large rivers.

**Locality:**

Prachuap Khirikhan (formerly in Pranburi); Phatthalung (formerly); Narathiwat (formerly in Maenam Bang Nara).

**Description:**

Size (total length) 4.75 m.

Snout long, narrow; 4 longitudinal series of juxtaposed keeled body scute on back. Generally dull olive or greenish, with dark brown spots; posterior dorsum and tail sometimes with transverse bands and spots; whitish beneath; iris yellow brown.



**Class:** Reptilia  
**Order:** Crocodylia  
**Family:** Crocodylidae  
**Scientific Name:** *Crocodylus porosus*  
**Common Name:** จระเข้น้ำเค็ม  
Estuarine Crocodile



**Habitat:**

Inhabits mouth of muddy rivers.

**Locality:**

Chumphon (formerly in Pa To); Surat Thani (formerly in Tapi River); Nakhon Si Thammarat (Kanom); Songkhla (formerly in Lake Songkhla); Narathiwat (Maenam Bang Nara, Klong Sungi Padi); Satun (Koh Tarutao).

**Description:**

Size (total length) 9.7 m.

Head somewhat triangular, relatively short; a pair of ridges extending forward on the skull between eyes, then curving slightly outward and then inward, the discontinuous ridges converging slightly; 3 large nuchals, often not contiguous, on each side; usually no enlarged occipital scutes proceeding nuchal scutes. Generally dark olive dorsally; small dots on head, venter yellowish.



<b>Class:</b>	Reptilia
<b>Order:</b>	Crocodylia
<b>Family:</b>	Crocodylidae
<b>Scientific Name:</b>	<i>Crocodylus siamensis</i>
<b>Common Name:</b>	จระเข้น้ำจืด Siamese Crocodile

**Habitat:**

Inhabits swamps, forest streams and rivers.

**Locality:**

Uttaradit (formerly); Phitsanulok (Maenam Kwa Noi); Pichit (formerly in Bung Sifi); Nakhon Sawan (Paknam Po, Bung Borapet); Nong Khai (formerly in Bung Khong Long); Ubon Ratchathani (Lam Dome Noi, Lam Dome Yai); Nakhon Ratchasima (Lam Chae, Lum Mun Bon); Srakaew (Pang Sida); Chachoengsao (Kwa Noi); Kanchanaburi (formerly in Kwa Noi); Phetchaburi (Kaeng Krachan, Maenam Phetchaburi, Huai Mae Prachan).

**Description:**

Size (total length) 3.5-4 m.

Snout less than twice as long as wide at base; two large pairs of scute each bordered laterally by a single enlarged seute on nuchal region; 4 occipital scutes proceeding nuchal scutes. Olive to brownish above; whitish or yellowish below.





**Class:** Reptilia  
**Order:** Testudines  
**Family:** Cheloniidae  
**Scientific Name:** *Chelonia mydas*  
**Common Name:** เต่าตนุ  
Green Sea Turtle



**Habitat:**

Inhabits seas and oceans, eggs laying in hole-nests on sand beaches.

**Locality:**

Nesting sites of this species are known from Chon Buri (Koh Kra and Koh Kram); Rayong (Koh Man Ni); Chumphon (Chumphon Is).

**Description:**

Size (carapace length) 73.5 cm.

Limbs paddle-shaped, with 1 or 2 claws; shell covered with horny shields; costal plates fused to ribs and carapace, not extending to edge of carapace, Greenish to greyish brown above, yellow below.



<b>Class:</b>	Reptilia
<b>Order:</b>	Testudines
<b>Family:</b>	Cheloniidae
<b>Scientific Name:</b>	<i>Eretmochelys imbricata</i>
<b>Common Name:</b>	เต่ากระ Hawksbill Sea Turtle

**Habitat:**

Inhabits seas and oceans, eggs laying in hole-nests on sand beaches.

**Locality:**

Nesting sites of this species are known from Chon Buri (Koh Kra and Koh Kram).

**Description:**

Size (carapace length) 1 m.

Four pairs of costal shields; head covered with regular shields consisting of a frontal preceded by 2 pairs of scales, a pair of supraorbitals, parietals, occipitals, 2 large supratemporals and 8 or 9 temporals; arms flipperlike; leg similar, shorter, widen, carapace covered with strongly imbricating shields, all keeled with accessory none parallel ridges; jaw hooked; usually 2 claws on hand and foot, Above brown, streaked with amber; plastron light yellowish.

*Lepidochelys olivacea*

**Class:** Reptilia

**Order:** Testudines

**Family:** Cheloniidae

**Scientific Name:** *Lepidochelys olivacea*

**Common Name:** เต่าหญ้า

Olive Ridley Sea Turtle

**Habitat:**

Inhabits seas and oceans, eggs laying in hole-nests on sand beaches.

**Locality:**

Nesting sites of this species are known from Phuket (Hat Kra Ron, Hat Suan Maprao); Phangnga (Hat Ta Nun, Hat Tai Muang, Koh Pra Tong).

**Description:**

Size (carapace length) 1 m. or more.

Normally 6 pairs of costals; intergular shield usually present, sometimes divided; usually 27 marginals. Drab olive or gray-olive, somewhat lighter at scutes; side and underside of neck whitish; plastron whitish with somewhat darker areas on plastral scutes.



<b>Class:</b>	Reptilia
<b>Order:</b>	Testudines
<b>Family:</b>	Dermochelyidae
<b>Scientific Name:</b>	<i>Dermochelys coriacea</i>
<b>Common Name:</b>	เต่ามะเฟือง Leatherback Sea Turtle

**Habitat:**

Inhabits seas and oceans, eggs laying in hole-nests on sand beaches.

**Locality:**

Nesting sites of this species are known from Phuket (Hat Pa Tong).

**Description:**

Size (carapace length) 2 m.

Large sea turtles, the young covered with small polygonal shields, largest ones on the soft leathery carapace and plastron. Adult with smooth skin. Young with 7 dorsal keels on carapace and 5 on plastron, the keels covered by raised quadrangular scutes. More or less symmetrical plates or scutes on head, the occipital usually the largest. Tail very short. Arms large flattened paddles, legs shorter flattened paddles. Dorsally a slaty black with 3 or 4 longitudinal rows of small white spots extending, between each pair of carapace ridges. Ventrally pinkish white or white, usually with dark reticulation representing scale marks.

*Batagur baska*

**Class:** Reptilia  
**Order:** Testudines  
**Family:** Geoemydidae  
**Scientific Name:** *Batagur baska*  
**Common Name:** เต่ากระอาน  
 River Terrapin

**Habitat:**

Inhabits large rivers, eggs laying on sand beach near mouth of the river.

**Locality:**

Nesting sites of this species is known from Prachuap Khirikhan (Khao Sam Roi Yot); Songkhla (formerly in Lake Songkhla); Ranong (Laem Son); Satun (Khlong La Ngu).

**Description:**

Size (carapace length) 60 cm.

Digits fully webbed with 4 claws only; tail very short; snout projecting, and some what turned up at end. Shell brownish olive to olive-green, whitish or yellowish below.



<b>Class:</b>	Reptilia
<b>Order:</b>	Testudines
<b>Family:</b>	Geoemydidae
<b>Scientific Name:</b>	<i>Callagur borneensis</i>
<b>Common Name:</b>	เต่าลายตีนเป็ด Painted Terrapin

**Habitat:**

Inhabits large rivers in coastal area, eggs laying on sand beach near mouth of the river.

**Locality:**

Nesting site of this species is known only from Satun (Khlung La Ngu).

**Description:**

Size (carapace length) 41 cm.

Oval carapace, light brown to olive with three broad, black longitudinal stripes. Both plastron and bridge uniformly yellow or cream colored. During the mating season the male's head become white and a red stripe develops between the eyes.



**Class:** Reptilia  
**Order:** Testudines  
**Family:** Trionichidae  
**Scientific Name:** *Chitra chitra*  
**Common Name:** ตะพาบมานลาย  
Thai Giant softshell



**Habitat:**

Inhabits Mae Klong River system and Mae Ping River.

**Locality:**

Tak (Bhumipol Dam); Kanchanaburi (Mae Khlong River and streamlets); Ratchaburi (Maenam Pachi).

**Description:**

Size (carapace length) 75 cm.

Carapace varied in shade of greenish yellow with a median vertebral stripe originating over the anterior bony carapace, no indistinct “shadow” markings, and the adornments of the vertebral area are not set off or distinct in any way from those on the costals, sometime two distinct discontinuous paramediam vertebral stripes also present. Plastron without pigments.



**Class:** Reptilia

**Order:** Testudines

**Family:** Trionychidae

**Scientific Name:** *Chitra vandijki*

**Common Name:** ตะพาบมันลายพม่า  
Burmese Giant Softshell



#### Habitat:

Inhabits Salween River system.

#### Locality:

Tak (Mae Sod).

#### Description:

Size (carapace length) 41.2 cm.

Carapace chocolate-brown and yellow-ochre colors predominating with a relatively complex vertebral and costal pattern, a neck “v” with divergence point located on the anterior half of the extended neck; paramedian neck stripes blending into the carapacial pattern and not forming a bell-like figure near the front of the carapace; 3-4 forelimb lamellae; no continuous light rim around the carapace; and no distinct naso-orbital triangular “figure” plastron whitish or pinkish.



*Pelochelys cantorii*

**Class:** Reptilia

**Order:** Testudines

**Family:** Trionychidae

**Scientific Name:** *Pelochelys cantorii*

**Common Name:** ตะพาบหัวกบ  
Asian Giant Softshell

**Habitat:**

Inhabits the large rivers, estuaries and mudflat areas.

**Locality:**

Bangkok (Maenam Chao Phraya); Phetchaburi (Maenam Phetchaburi); Yala (Banglang Dam).

**Description:**

Size (carapace length) 65 cm.

Head large, much wider in temporal region; proboscis short, tips thick; eyes small with a dark line in front and behind pupil; tail short behind vent. Head above olive with minute black dots; carapace olive, with a few darker and lighter striations along median dorsal part; outer edge olive with small spots of darker and lighter color; plastron flesh-colored with a few white dots on anterior parts; chin and throat with minute dots of black and white.



**Class:** Reptilia  
**Order:** Testudines  
**Family:** Platysternidae  
**Scientific Name:** *Platysternon megacephalum*  
**Common Name:** เต่าปูลู  
 Big-headed Turtle

**Habitat:**

Inhabits hill streams in forest areas.

**Locality:**

Mae Hong Son (Khun Yuam); Chiang Mai (Doi Ang Kang, Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Loei (Dan Sai, Phu Kradung, Phu Luang); Chaiyaphum (Phu Kieo); Phetchabun (Nam Nao); Phrae; Tak; Kanchanaburi (Sai Yok, Thong Pa Phum).

**Description:**

Size (carapace length) 18 cm.

Plastron much smaller than the opening of carapace; head large, with powerful jaws covered with a horny shield; 5 fingers with claws; 4 toes with claws, the webbing not quite reaching base of claw; tail greatly elongated, covered above and below with heavy scutes. Nearly uniform light brown; head in same color; underside of marginals and plastron uniform yellowish or yellow brown.

*Indotestudo elongata*

**Class:** Reptilia

**Order:** Testudines

**Family:** Testudinidae

**Scientific Name:** *Indotestudo elongata*

**Common Name:** เต่าเหลื่อม

Elongated Tortoise

**Habitat:**

Inhabits hilly areas in deciduous and evergreen forests.

**Locality:**

Chiang Mai (Doi Chiang Dao; Doi Inthanon); Phayao (Doi Pachang); Lampang ; Phrae; Nan; Tak (Thung Yai); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok, Thong Pa Phum); Surat Thani (Klong Saeng); Nakhon Si Thammarat (Khao Luang); Krabi (Klong Praya, Ban Kha Nom); Narathiwat (Waeng); Loei (Phu Luang); Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida, Ta Phraya); Chanthaburi (Khao Sebab, Khao Soi Dao).

**Description:**

Size of carapace 30 cm.

Carapace elongate, strongly arched. Generally yellow above and below, with large dark spots or flecks on dorsum, larger dark spots on marginal scales. Top of head with symmetrical scales.



**Class:** Reptilia  
**Order:** Testudines  
**Family:** Testudinidae  
**Scientific Name:** *Manouria emys emys*  
**Common Name:** เต่าหกเหลือง  
 Asian Brown Tortoise

**Habitat:**

Inhabits hilly areas in evergreen forests.

**Locality:**

Songkhla (Ton Ngachang); Trang (Khao Bantat); Narathiwat (Waeng).

**Description:**

Carapace size 47 cm.

Carapace broadly arching, dorsal surface somewhat flattened; part of shield concave, especially the posterior costals and some marginals. Above, carapace a shade of olive-black; plastron largely yellowish with blackish areas, flecks or clouding.

*Manouria emys phayrei*

**Class:** Reptilia

**Order:** Testudines

**Family:** Testudinidae

**Scientific Name:** *Manouria emys phayrei*

**Common Name:** เต่าหกค้ำ

Burmese Giant Tortoise

**Habitat:**

Inhabits hilly areas in evergreen forests.

**Locality:**

Tak (Um Phang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip).

**Description:**

Similar to *M.e. emys* but differs by pectoral shields meet together at the median of plastron (in *M.e. emys*, the pectoral shields widely separated from each other). Blackish above and below.



<b>Class:</b>	Reptilia
<b>Order:</b>	Testudines
<b>Family:</b>	Testudinidae
<b>Scientific Name:</b>	<i>Manouria impressa</i>
<b>Common Name:</b>	เต่าเตี๋ย Impressed Tortoise

**Habitat:**

Inhabits hilly areas in high altitude forests.

**Locality:**

Chiang Mai (Doi Inthanon); Tak (Mae Sot, Umphang); Loei (Phu Luang); Narathiwat (Waeng).

**Description:**

Size of carapace 30 cm.

Carapace much flattened on dorsal surface, vertebral and costal scutes somewhat concave; Top of head with symmetrical shields. Light brown of varying shades, lighter near sutures; about periphery of each scale, short dark brown radiating lines; laterally on marginals some indefinite larger dark spots; plastron yellow-brown becoming pale yellowish near sutures, distal remnants of radiating lines.

*Physignathus cocincinus*

**Class:** Reptilia

**Order:** Squamata

**Family:** Agamidae

**Scientific Name:** *Physignathus cocincinus*

**Common Name:** ตะกอลง, ลีง

Indochinese Water Dragon

**Habitat:**

Inhabits along hill streams in evergreen forest.

**Locality:**

Sakon Nakhon (Phu Phan); Ubon Ratchathani (Sanoi River); Nakhon Ratchasima (Khao Yai, Sakaerat); Saraburi (Muak Lek); Nakhon Nayok (Sarika Falls); Srakaew (Pang Sida) Prachin Buri (Tap Lan); Chachoengsao (Khao Ang Rue Ni); Rayong (Khao Chamao); Chanthaburi (Khao Sebab, Khao Soidao); Trat (Koh Chang); Chon Buri (Khao Kieo).

**Description:**

Size (snout-vent length) 20 cm.

Body compressed; nuchal and dorsal crest continuous, caudal crest separated by a space; a nuchal fold; femoral pores in short series; tail strongly compressed. Above, head and body light greenish olive, the enlarged scales on chin bluish bordered with rose-lavender; venter surface green.

**Status VU**

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cuora amboinensis</i>	เต่าห้า Malayan Box Turtle	Inhabits lowland wetlands such as swamp, wet paddy fields, canals, rivers including hill streams.	Chiang Mai; Nakhon Sawan; Kanchanaburi; Bangkok; Pathum Thani; Narathiwat.
<i>Cyclemys atripons</i>	เต่าห้วยทองเหลือง Southeastern Leaf Turtle	Inhabits hill streams in evergreen forest.	Nakhon Ratchasima (Kon Buri, Wang Namkieo, Khaoyai); Chachoengsao (Khao Ang Rue Ni); Chon Buri (Khao Khieo); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat.
<i>Cyclemys dentata</i>	เต่าใบไม้ Asian Leaf Turtle	Inhabits hill streams in evergreen forest.	Surat Thani (Samui); Yala (Bhetong); Narathiwat (Phru To Daeng).
<i>Cyclemys oldhami</i>	เต่าใบไม้พม่า Burmese Leaf Turtle	Inhabits hill streams in evergreen forest.	Uthai Thani (Hui Khakhaeng); Kanchanaburi (Thung Yai); Nakhon Si Thammarat (Khao Luang); Yala (Banangstars).
<i>Cyclemys shanensis</i>	เต่าห้วยคดลย Indochinese Leaf Turtle	Inhabits hill streams in evergreen forest.	Chiang Mai (Doi Chiang Dao), Tak (Umpang), Phayao (Doi Pa Chang), Loei (Phu Luang); Ubon Ratchathani (Sanoi River).
<i>Heosemys grandis</i>	เต่าห้วย Giant Asian Pond Turtle	Inhabits hill streams in evergreen forest.	Uthai Thani (Hui Khakhaeng); Kanchanaburi (Thung Yai); Narathiwat (Waeng).
<i>Heosemys spinosa</i>	เต่าจ๊กกร Indochinese Leaf Turtle	Inhabits hill streams and forest floor in	Surat Thani (Khlung Saeng); Krabi (Khao Pra Bang Kram);



Scientific Name	Thai name/Common name	Habitat	Site
<i>Hieremys annandalei</i>	เต่าบัว Yellow-headed Temple Turtle	evergreen forest. Inhabits lowland wetlands such as swamp, wet rice field, canals, rivers.	Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Yala (Banangstars, Bhetong); Narathiwat (Waeng). Bangkok (Bangkhen, Bung Kum); Pathum Thani (Rangsit, Khlung Luang); Samut Prakan (Maung); Ratchaburi; Pattani (Yee Ngo).
<i>Malayemys subtrijuga</i>	เต่านา Malaysian Snail-eating Turtle	Inhabits lowland wetlands such as swamp, wet rice field, canals, rivers.	Ching Mai (Maung); Phitsanulok (Wat Bos); Nhong Bua Lumpoo (Maung); Nakhon Sawan (Bung Borapet); Bangkok (Bangkhen, Bung Kum); Pathum Thani (Rangsit, Khlung Luang); Samut Prakan (Maung); Ratchaburi; Kanchanaburi; Ubon Ratchathani; Nakhon Ratchasima; Chachoengsao; Phangnga; Narathiwat.
<i>Notochelys platynota</i>	เต่าทับทิม Malaysian Flat-shelled Turtle	Inhabits hill streams in evergreen forest.	Songkhla (Ton Ngachang); Narathiwat (Waeng).
<i>Siebenrockiella crassicollis</i>	เต่าดำ Black Pond Turtle	Inhabits lowland wetlands such as swamp, ditches, canals, rivers.	Bangkok; Pathum Thani; Surat Thani; Narathiwat (Sungi Kolok).
<i>Amyda cartilaginea</i>	ตะพาบธรรมดา Common Asiatic Softshell	Inhabits from forest streams to canals and rivers in lowland wetlands.	Mae Hong Son (Mae Sariang); Kamphaeng Phet (Khlung Lan); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Mae

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gonocephalus abbotti</i>	กิ้งก่าดงคิ้วสั้น Abbolt's Angle-headed Lizard	Inhabits along hill streams in evergreen forest.	Khlong River); Loei (Phu Luang); Srakaew (Pang Sida); Chanthaburi (Khao Sebab); Bangkok; Pathum Thani (Rangsit); Surat Thani (Khlong Saeng); Nakhon Si Thammarat (Thung Song, Thung Taa); Satun (Taleban); Yala (Bannangstars).
<i>Gonocephalus bellii</i>	กิ้งก่าดงคอฟ้า Blue-necked Angle-headed Lizard	Inhabits along hill streams in evergreen forest.	Trang (Khao Chong); Yala (Bannangstars); Narathiwat (Waeng).
<i>Gonocephalus grandis</i>	กิ้งก่าดงใหญ่ Great Angle-headed Lizard	Inhabits along hill streams in evergreen forest.	Surat Thani (Khao Nang, Ban Kok Khlab); Yala (Bannangstars); Narathiwat (Waeng).
<i>Mantheyus phuwuaensis</i>	กิ้งก่าภูว Phu Wua Lizard	Inhabits along stream bank in evergreen forest.	Songkhla (Na Twi); Satun (Tale Ban); Yala (Bannangstars, Bhetong); Narathiwat (Waeng). Nong Khai (Phu Wua).

# Status NT

Scientific Name	Thai name/Common name	Habitat	Site
<i>Dogania subplana</i>	ตะพาบดำ Malayan Soffshell	Inhabits forest streams in evergreen forest.	Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok, Thong Phaphum); Ratchaburi (Maenam Pachi); Phetchaburi (Kaeng Krachaan); Surat Thani (khlong Saeng); Phangnga, Krabi (Khao Panombencha); Songkhla (Ton Ngachang); Satun (Taleban, Tarutao); Yala (Bannangstars).
<i>Pseudocalotes floweri</i>	กิ้งก่าหัวยาว Flower's Long-headed Lizard	Inhabits evergreen forest.	Chanthaburi (Khao Soi Dao).
<i>Pseudocalotes microlepis</i>	กิ้งก่าเขาสอง Boulenger's Long-headed Lizard	Inhabits sub-montane forest.	Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Loei (Phu Luang); Chaiyaphum (Phu Kieo).
<i>Leiolepis bohmei</i>	แยงซงขลา Bohme's Butterfly Lizard	Inhabits sand beach area.	Songkhla.
<i>Leiolepis belliana belliana</i>	แยงใต้ Southern Butterfly Lizard	Inhabits coastal areas, Melaleuca forest, deciduous forest and dry evergreen forest.	Chon Buri (Bang Pra); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Ratchaburi (Maenam Pachi); Phetchaburi (Cha-Um); Trang (Kantang, Koh Libong); Pattani (Napradoo); Narathiwat (Takbai).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Leiolepis belliana ocellata</i>	แย้เหนือ Northern Butterfly Lizard	Inhabits deciduous forest.	Chiang Mai; Phayao.
<i>Leiolepis reevesi rubritaeniata</i>	แย้อีสาน Indochinese Butterfly lizard	Inhabits deciduous forest and dry evergreen forest.	Chaiyaphum (Ban Mak Yang); Khon Kaen (Phu Wiang); Nakhon Ratchasima (Pak Chong, Sakaerat); Si Sa Ket; Ubon Ratchathani (Nam Yun).
<i>Ophisaurus gracilis</i>	กิ้งก่า Serpent Lizard	Inhabits sub-montane forest.	Chiang Mai (Doi Pahompok); Loei (Phu Luang).
<i>Cyrtodactylus angularis</i>	ตุ๊กแกป่าดงพญาเย็น Angular spotted Gecko	Inhabits deciduous and dry evergreen forest, usually found at night on the rock of forest floor.	Nakhon Ratchasima (Lad Bua Khao, Pak Chong Sakaerat); Saraburi (Hin Lab, Muak Lek, Pha Sadet).
<i>Cyrtodactylus brevipalmatus</i>	ตุ๊กแกป่าโคนไม้ Short-webber Gecko	Inhabits evergreen forest, usually found at night on tree trunks.	Nakhon Si Thammarat (Khao Luang).
<i>Cyrtodactylus chanhomeae</i>	ตุ๊กแกป่าสระบุรี Chanhome's Gecko	Inhabits limestone caves.	Saraburi.
<i>Cyrtodactylus jarujini</i>	ตุ๊กแกป่าจระจก Jarujin's Gecko	Inhabits evergreen forest and caves nearby.	Nong Khai (Phu Wua).
<i>Cyrtodactylus sumonthai</i>	ตุ๊กแกป่าเขาชะเมา Sumontha's Gecko	Inhabits caves.	Rayong (Khao Cha Mao).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cyrtodactylus tigroides</i>	ตุ๊กแก่ป่าไทรโยค Sai Yok Gecko	Inhabits evergreen forest and nearby caves.	Tak (Umphang); Kanchanaburi (Sai Yok).
<i>Gekko siamensis</i>	ตุ๊กแกสยาม Siamese Gecko	Found in rock crevices and cave in degraded forest.	Nakhon Ratchasima (Pak Chong); Phetchabun (Nam Nao).
<i>Davewakeum miriamae</i>	จิ้งเหลนคางบักธงชัย Miriam's Legless Skink	Inhabits evergreen forest, usually found under ground of tree buttress.	Nakhon Ratchasima (Sakaerat).
<i>Eumeces quadrilineatus</i>	จิ้งเหลนหางสีฟ้า Blue-tailed Skink	Inhabits evergreen and submontane forest.	Chiang Mai (Doi Angka, Doi Chiang Dao); Loei (Dan Sai, Phu Kradung); Saraburi (Muak Lek).
<i>Isopachys borealis</i>	จิ้งเหลนคางป่าตะวันตก Western Legless Skink	Inhabits evergreen forest, usually found under humus soil.	Khon Kaen; Phetchabun (Nam Len); Tak (Mae Sot); Nakhon Sawan (Taling Sung); Uthai Thani (Hau Khakhaeng, Lan Sak).
<i>Isopachys gyldenstolpei</i>	จิ้งเหลนคางกลาย Gyldenstolpe's Legless Skink	Inhabits beach and evergreen forest, usually found under humus soil.	Uthai Thani (Ban Rai), Kanchanaburi, Phetchaburi (Cha-Um); Prachuap Khirikhan (Hua Hin, Koh Lak, Muang).
<i>Isopachys roulei</i>	จิ้งเหลนคางชลบุรี Roule's Legless Skink	Inhabits beach forest, usually found under humus soil.	Chon Buri (Ang Sila).
<i>Leptoceps osellai</i>	จิ้งเหลนภูเขาสามนิ้ว Osella's Skink	Inhabits evergreen forest, usually found under humus soil and rotten logs.	Chiang Mai (Mae Kung); Phetchabun (Nam Nao).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Lygosoma corpulentum</i>	จิ้งเหลนเรียวจันทบุรี Annamese Slender Skink	Inhabits lowland evergreen forest, usually found under rotten logs on forest ground.	Chachoengsao (Khao Takrup); Chanthaburi (Khao Sebab, Khao Soi Dao).
<i>Lygosoma haroldyoungi</i>	จิ้งเหลนเรียวลาย Banded Slender Skink	Inhabits evergreen forest, usually found in termite mound or under humus soil.	Chiang Mai (Doi Suthep-Pui, Fang); Chiang Rai; Phitsanulok (Thung Salaeng Luang); Loei (Phu Luang); Nong Khai (Bunh Kan, Rattana Wapi); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo).
<i>Lygosoma isodactylum</i>	จิ้งเหลนเรียวภาคกลาง Central Slender Skink	Inhabits shrub and evergreen forest, usually found under rotten log.	Nakhon Sawan; Phra Nakhon Si Ayutthaya; Saraburi (Chong Kae, Muak Lek).
<i>Lygosoma koratense</i>	จิ้งเหลนเรียวโคราช Khorat Slender Skink	Inhabits evergreen forest.	Saraburi (Muak Lek); Nakhon Ratchasima (Dong Lek, Lad Bua Khao, Non Thai).
<i>Scincella melanosticta kohtaoensis</i>	จิ้งเหลนดินเกาะเต่า Koh Tao Ground Skink	Inhabits evergreen forest, found under rotten logs and humus soil.	Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Loei (Phu Luang).
<i>Varanus dumerilii</i>	ตุ๊กตุ๊ก Dumeril's Moutior	Inhabits evergreen forest.	Kanchanaburi (Thong Paphum); Ratchaburi (Mae Pradon, Maenam Pachi); Surat Thani (Khlung Saeng, Ta Chana); Nakhon Si Thammarat (Tai Rom Yen, Thung Song); Trang (Khao Chong); Phatthalung (Tamod, Khao Pu- Khao Ya); Songkhla (Ton Ngachang); Yala (Bannangstars); Narathiwat (Sungai Kolok, Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Varanus nebulosus</i>	ตะกวด Indomalayan Monitor	Inhabits deciduous and evergreen forest.	Kanchanaburi (Thong Paphum); Ratchaburi (Mae Pradon, Maenam Pachi); Surat Thani (Khlung Saeng, Ta Chana); Nakhon Si Thammarat (Tai Rom Yen, Thung Song); Trang (Khao Chong); Phatthalung (Tamod, Khao Pu- Khao Ya); Songkhla (Ton Ngachang); Yala (Bannangstars); Narathiwat (Sungai Kolok, Waeng).
<i>Varanus rudicollis</i>	เหาซาง Rough-necked Monitor	Inhabits evergreen forest.	Tak (Um Phang); Uthai Thani (Hau Khakhaeng); Ranong (Khlung Naka); Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Tai Rom Yen, Thung Song); Krabi (Khao Pra Bang Kram); Trang (Khao Chong); Phatthalung (Khao Pu- Khao Ya); Songkhla (Ton Ngachang); Satun (Tale Ban); Yala (Bannangstars); Narathiwat (Waeng).
<i>Acrochordus javanicus</i>	งูวงช้าง Javan File Snake	Inhabits freshwater wetlands.	Bangkok (Bangkhen, Pathumwan); Prachin Buri (Prachantakham); Narathiwat (Sungai Kolok).
<i>Boiga sangsomi</i>	งูเขียวดงลาย Banded Cat Snake	Inhabits evergreen forest, found on palm tree.	Krabi (Ban Khanom); Nakhon Si Thammarat (Ron Pibun).
<i>Chrysopelea paradisi paradisi</i>	งูเขียวรอน Paradise Tree Snake	Inhabits evergreen forest.	Phuket; Phangnga (Khao Lak); Narathiwat (Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Chrysopelea pelias</i>	งูดอกหมากแดง Red Tree Snake	Inhabits evergreen forest.	Songkhla (Ton Ngachang); Narathiwat (Waeng) (Chan-ard, per.observ.).
<i>Dryocalamus thungsongensis</i>	งูปล่องฉนวนทุ่งสง Thungsong Wolf Snake	Inhabits lowland evergreen forest.	Nakhon Si Thammarat (Khao Luang, Thung Song).
<i>Lycodon cadamomensis</i>	งูปล่องฉนวนจันทบุรี Cadamon Mountain Wolf Snake	Inhabits lowland evergreen forest.	Chanthaburi (Khao Sebab, Khao Soi Dao) (Pauwels, per.observ.; Chan-ard, per. observ.).
<i>Oreophis porphyraceus coxi</i>	งูทางมะพร้าวแดงภูหลวง Cox's Racer	Inhabits evergreen forest. In Loei, it was observed while consuming a mole.	Loei (Phu Luang); Chaiyaphum (Phu Kieo).
<i>Oreophis porphyraceus porphyraceus</i>	งูทางมะพร้าวแดงลาขวัน Red Mountain Racer	Inhabits hill forest and caves nearby.	Loei (Phu Luang); Kanchanaburi (Sai Yok, Salak Pra); Phetchaburi (Khao Yoi).
<i>Pseudorhabdion longiceps</i>	งูพงอห้วยยาว Long-headed Reed Snake	Inhabits lowland evergreen forest.	Yala (Betong); Narathiwat (Sukirin, Waeng).
<i>Ptyas fuscus</i>	งูสิงทอง Gunther's Rat Snake	Inhabits evergreen forest.	Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang).
<i>Opisthotropis praemaxillaris</i>	งูลายสอลาเหนือ Angel's Mountain Keelback	Inhabits hill streams in evergreen forest.	Chiang Mai (Doi Saket).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Opisthotropis spenceri</i>	งูลายสอสองสี Spencer's Mountain Keelback	Inhabits hill streams in evergreen forest.	Lampang (Ngao).
<i>Parahelicops boonsongi</i>	งูลายสอหมอบนูนสูง Boonsong's keelback	Inhabits hill streams in evergreen forest.	Loei (Dansai, Phu Rua).
<i>Bungarus flaviceps flaviceps</i>	งูสามเหลี่ยมหัวแดง Red-headed Krait	Inhabits lowland evergreen forest, usually found near forest stream.	Ratchaburi (Chom Bung); Ranong (Khlong Naka); Phangnga (Khao Lak); Surat Thani (Khao Sok, Khlong Saeng); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Yala (Betong); Narathiwat (Waeng).
<i>Calliophis bivirgatus flaviceps</i>	งูพริกทองแดง Blue Long-glanded Coral Snake	Inhabits evergreen forest.	Trang (Khao Bantad, Khao Chong); Songkhla (Ton Ngachang); Narathiwat (Waeng).
<i>Calliophis intestinalis lineata</i>	งูพริกสีน้ำตาล Brown Long-glanded Coral Snake	Inhabits evergreen forest.	Surat Thani; Nakhon Si Thammarat (Khao Luang); Pattani (Napradoo); Narathiwat (Waeng).
<i>Calliophis maculiceps</i>	งูปล่องหาวหัวดำ Small spotted Coral Snake	Inhabits evergreen forest.	Chiang Mai (Doi Chiang Dao); Phetchabun (Nam Nao); Nong Khai (Phu Wua); Nakhon Ratchasima (Pak Chong); Saraburi (Muak Lek); Chon Buri (Siracha); Kanchanaburi; Phetchaburi

Scientific Name	Thai name/Common name	Habitat	Site
<i>Sinomicrurus macclelandi macclelandi</i>	งูปล้องหวายลายขั้วนาค้า Maclelland's Coral Snake	Inhabits evergreen forest up to 1,200 m.	(Kaengkrachan); Surat Thani (Khao Tapetch, Koh Tao); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Phuket; Pattani (Napradoo). Chiang Mai (Doi Suthep-Pui); Loei (Phu Luang); Nakhon Ratchasima (Sakaerat); Chachoengsao (Khao Ang Rue Ni); Phetchaburi (Kaengkrachan).
<i>Trimeresurus (Trimeresurus) borneensis</i>	งูปล่าม Borneaw Pit Viper	Inhabits lowland evergreen forest, including peat swamp forest. It is usually found on forest ground or the base of tree buttress.	Nakhon Si Thammarat (Khao Luang, Ron Pibun, Thung Song); Narathiwat (Sungai Kolok).
<i>Trimeresurus (Parias) sumatranus</i>	งูเขียวหางไหม้สุมาตรา Sumatran Pit Viper	Inhabits lowland evergreen forest.	Pattani; Narathiwat (Waeng).
<i>Tropidolaemus wagleri</i>	งูเขียวตุ๊กแก Wagler's Pit Viper	Inhabits lowland evergreen forest, including swampy forest.	Surat Thani (Khlung Saeng); Phangnga (Khao Lak); Pattani (Napradoo); Yala; Narathiwat (Sungai Kolok, Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Acanthosaura armata</i>	กิ้งก่าเขาหนามขาว Greater Spiny Lizard	Inhabits evergreen forest.	Pattani (Napradoo); Yala (Bannangstars); Narathiwat (Waeng).
<i>Acanthosaura crucigera</i>	กิ้งก่าเขาหนามดำ Masked Spiny Lizard	Inhabits evergreen forest.	Phrae; Tak (Thung Yai); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Ratchaburi (Maenam Pachi); Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Pa-La-U); Ranong (Khlung Nakaa); Surat Thani (Khao Sok, Khlung Saeng); Phangnga (Khao Lak Lam Ru); Krabi (Khao Panom Bencia, Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Bantat); Songkhla (Ton Ngachang); Nakhon Ratchasima (Khao Yai, Sakaerat); Roi Et (Phu Pa Namtip); Srakaew (Pang Sida); Chachoengsao (Khao Ang Rue Ni); Chon Buri (Khao Kieo); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (Koh Chang).
<i>Acanthosaura lepidogaster</i>	กิ้งก่าเขาลูก Cuvier's Spiny Lizard	Inhabits evergreen forest.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Lampang (Doi Khun Tan); Tak (Doi Mon Chong); Loei (Phu Luang); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Aphaniotis fusca</i>	กิ้งก่าพระอินทร์ Malayan Blue-Mouthed Lizard	Inhabits evergreen forest.	Yala (Banangstara); Narathiwat (Waeng).
<i>Bronchocele cristatella</i>	กิ้งก่าเขียวโต Green Crested Lizard	Inhabits evergreen forest.	Kanchanaburi (Sai Yok); Chumphon (Pathiu); Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Khao Luang); Trang (Koh Libong); Pattani (Naapradoo); Yala (Bannangstars, Tharn To); Narathiwat (Tak Bi, Sungai Kolok, Waeng).
<i>Bronchocele smaragdina</i>	กิ้งก่าเขียวจันทบูรณ Chantabun Green Tree Lizard	Inhabits evergreen forest.	Nakhon Nayok (Khao Yai); Chanthaburi (Khao Sebab, Khao Soi Dao).
<i>Calotes emma alticristata</i>	กิ้งก่าแกวเหนือ Northern Forest Crested Lizard	Inhabits evergreen forest.	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Phrae (Mae Yom); Tak (Doi Mon Chong, Thung Yai, Um Phang); Kamphaeng Phet (Khlung Lan); Uthai Thani (Hau Kha Khaeng); Kanchanaburi (Erawan, Sai Yok, Thong Pa Phum); Ratchaburi (Maenam Pachi); Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Pa-La-U); Loei (Phu Luang); Nong Khai (Phu Wua); Sakon Nakhon (Phu Phan); Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida,

Scientific Name	Thai name/Common name	Habitat	Site
<i>Calotes emma emma</i>	กิ้งก่าแก้วใต้ Southern Forest Crested Lizard	Inhabits evergreen forest.	Taphraya); Prachinburi (Tap Lan); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (koh Chang); Chon Buri (Khao Kieo). Surat Thani (Khao Sok, Khlong Saeng); Phangnga (Khao Lak Lam Ru); Ranong (Khlong Naka); Nakhon Si Thammarat (Khao Luang); Krabi (Khao Panom Bencha, Khao Pra Bang Kram); Trang (Khao Bantat, Koh Libong); Songkhla (Ton Ngachang); Yala (Bhetong, Bannangstars, Than To); Narathiwat (Sungai Kolok, Waeng).
<i>Calotes mystaceus</i>	กิ้งก่าสวน, กิ้งก่าหัวฟ้า Blue Crested Lizard	Inhabits evergreen forest.	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon); Phrae (Mae Yom); Nakhon Sawan; Uthai Thani (Hau Kha Khaeng); Kanchanaburi (Sai Yok); Ratchaburi (Maenam Pachi); Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Bang Saphan, Hua Hin); Loei (Wang Saphung); Nong Khai (Phu Wua); Sakon Nakhon (Phu Phan); Chaiyaphum (Phu Kieo); Udon Thani; Khon Kaen; Ubon Ratchathani (Nam Yun); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida, Taphraya).
<i>Calotes versicolor</i>	กิ้งก่าริ้ว, กิ้งก่าหัวแดง Changeable Lizard	Inhabits from beach forest to deciduous forest, including urban and agricultural areas.	All provinces.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Draco blanfordi</i>	กิ้งก่าบินคอแดง Blanford's Flying Lizard	Inhabits evergreen forest.	Chiang Rai; Chiang Mai (Doi Suthep-Pui); Phrae; Loei (Phu Luang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Sai Yok); Prachuap Khirikhan (Pa-La-U); Chunphon (Tapli, Tasan); Ranong (Khlung Naka); Surat Thani (Khao Sok, Khlung Saeng); Phuket; Krabi (Khao Pra Bangkram; Nong Kok); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Songkhla (Ton Ngachang); Satun (Tale Ban); Yala (Banangstars); Narathiwat (Waeng).
<i>Draco fimbriatus</i>	กิ้งก่าบินคอส้ม Orange-bearded Flying Lizard	Inhabits evergreen forest.	Nakhon Si Thammarat (Khao Luang, Nabon); Trang (Khao Chong); Yala (Banangstars); Narathiwat (Waeng).
<i>Draco haematopogon</i>	กิ้งก่าบินปีกแดง Red-bearded Flying Lizard	Inhabits evergreen forest.	Surat Thani (Koh Pa Ngan).
<i>Draco maculatus divergens</i>	กิ้งก่าบินปีกสมคอดอยสุเทพ Doi Suthep Flying Lizard	Inhabits evergreen forest.	Chiang Mai (Doi Suthep-Pui).
<i>Draco maculatus haasei</i>	กิ้งก่าบินปีกสมจันทบูรณ Chantabun Flying Lizard	Inhabits evergreen forest.	Ubon Ratchathani (Sanoi River); Nakhon Ratchasima (Khao Yai); Srakaew (Pang Sida); Chanthaburi (Khao Kitchakud, Khao Soi Dao); Trat (Koh Chang); Chon Buri (Khao Kieo).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Draco maculatus maculatus</i>	กิ้งก่าบินปีกสมมลาชู Southern Flying Lizard	Inhabits evergreen forest.	Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Sai Yok); Prachuap Khirikhan (Pa-La-U); Ranong (Maung); Surat Thani (Khao Sok, Khlong Saeng, Koh Samui); Nakhon Si Thammarat (Had Nai Ploy, Khao Luang, Namtok Yong); Trang (Khao Chong); Songkhla (Ton Ngachang); Phatthalung (Tale Noi); Krabi (Khao Pra Bang Kram).
<i>Draco maculatus whiteheadi</i>	กิ้งก่าบินปีกสมไหหลำ Whitehead's Flying Lizard	Inhabits evergreen forest.	Chiang Mai; Phrae; Loei (Phu Luang); Nong Khai (Phu Wua); Chaiyaphum (Phu Kieo).
<i>Draco melanopogon</i>	กิ้งก่าบินคอดำ Black-bearded Flying Lizard	Inhabits evergreen forest.	Surat Thani, Nakhon Si Thammarat (Khao Luang, Nabon, Ronpibun); Krabi; Trang (Khao Chong); Phatthalung; Yala (Banangstars, Bhetong); Narathiwat (Bangnara, Waeng).
<i>Draco obscurus formosus</i>	กิ้งก่าบินมลาชู Malayan Flying Lizard	Inhabits evergreen forest.	Chumphon (Pathieu, Mabrit); Nakhon Si Thammarat (Khao Wang Hip); Trang (Khao Chong); Pattani (Napradoo); Yala (Banangstars, Bhetong); Narathiwat (Waeng).
<i>Draco quinquefasciatus</i>	กิ้งก่าบินปีกห้าแถบ Five-banded Flying Lizard	Inhabits evergreen forest.	Surat Thani (Khlong Saeng); Trang (Khao Chong); Pattani; Yala (Bhetong); Narathiwat (Sungai Kolok, Waeng).
<i>Draco taeniopterus</i>	กิ้งก่าบินปีกกลาย	Inhabits evergreen forest.	Phrae (Song); Chiang Mai; Nakhon Ratchasima; Chanthaburi

Scientific Name	Thai name/Common name	Habitat	Site
	Banded Winged Flying Lizard		(Khao Sebab, Khao Soi Dao); Trat (Koh Chang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Thong Phaphum); Chumphon (Tapi); Ranong (Mamoh, Maung, Pak Chan); Phuket (Thung Phran); Krabi (Khao Pra Bang Kram, Koh Ra, Nong Kok, Prai Phraya); Surat Thani (Khao Sok); Nakhon Si Thammarat (Khao Luang); Trang (Kantang, Khao Chong, Koh Libong).
<i>Draco volans sumatranus</i>	กิ้งก่าบินคอฟ้า	Inhabits evergreen forest.	Kanchanaburi (Sai Yok); Nakhon Si Thammarat; Yala (Bhetong, Bannangstars); Narathiwat (Waeng).
	Blue-necked Flying Lizard		
<i>Cnemaspis affinis</i>	จิ้งจกนิ้วขาวมลายู	Found around tree buttress and under rocks closed to forest streams.	Pattani (Khok Po).
	Malayan Rock Gecko		
<i>Cnemaspis chanthaburiensis</i>	จิ้งจกนิ้วขาวจันทบุรี	Found in tree buttress and under log on forest floor.	Chanthaburi (Khao Sebab, Khao Soi Dao).
	Dring's Rock Gecko		
<i>Cnemaspis flavolineata</i>	จิ้งจกนิ้วขาวหลังขีด	Found in tree buttress along stream bank in evergreen forest.	Narathiwat (Waeng).
	Striped Rock Gecko		
<i>Cnemaspis kandianus</i>	จิ้งจกนิ้วขาวอินเดีย	Found in tree buttress on forest floor.	Phetchaburi (Kaengkrachan); Chumphon (Namtok Krapo, Tasan); Ranong (Koh Lan, Koh Yan Yan).
	Indian Rock Gecko		
<i>Cnemaspis kumpoli</i>	จิ้งจกนิ้วขาวกัมพล	Inhabits evergreen forest, usually found in tree buttress and under rock along	Prachuap Khirikhan (Hau Yang); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Trang (Khao
	Kumpol's Gecko		



Scientific Name	Thai name/Common name	Habitat	Site
<i>Cnemaspis siamensis</i>	จิ้งจกนิ้วขาวไทย Siamese Rock Gecko	stream bank. Inhabits evergreen forest, usually found in tree buttress and under rock along stream bank.	Chong); Satun (Tale Ban); Narathiwat (Waeng). Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Chumphon (Mab Rit, Pa Tieu, Tasan); Phangnga (Khao Lak Lam Ru); Phuket; Krabi (Khao Pra Bang Kram); Surat Thani (Koh Samui); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip); Trang (Khao Chong); Satun (Tale Ban); Yala (Than To).
<i>Cosymbotus craspedotus</i>	จิ้งจกหางแบนมลายู Malayan Frilly Gecko	Inhabits evergreen forest, usually found under barks of dead tree trunk.	Nakhon Si Thammarat (Khao Luang, Khao Ron Pibun); Trang (Khao Chong); Songkhla (Ton Ngachang); Narathiwat (Waeng).
<i>Cosymbotus platyurus</i>	จิ้งจกบ้าน Common Frilly Gecko	Inhabits all types of forest, crop plantation, also houses and gardens.	All provinces.
<i>Cyrtodactylus consobrinus</i>	ตุ๊กแกป่ามลายู Peter's Gecko	Inhabits evergreen forest, usually found at night on tree trunks.	Yala (Bannangstars); Narathiwat (Waeng).
<i>Cyrtodactylus interdigitalis</i>	ตุ๊กแกป่าห้าหนวด Ulber's Gecko	Inhabits evergreen forest, usually found at night on tree trunks.	Loei (Phu Luang); Phetchabun (Nam Nao); Tak (Um Phang); Uthai Thani (Hau Khakhaeng, Khao Pla Ra); Phetchaburi (Kaengkrachan).
<i>Cyrtodactylus intermedius</i>	ตุ๊กแกป่าลายพาดตะวันออก	Inhabits evergreen forest, usually found	Nakhon Ratchasima (Khao Yai, Sakaerat, Tap Lan); Srakaew

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cyrtodactylus oldhami</i>	Eastern Banded Gecko ตุ๊กแกป่าคอขวั้น Oldham's Gecko	at night on tree trunks. Inhabits evergreen forest, usually found at night on tree trunks.	(Pang Sida, Taphraya); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (Koh Chang); Chon Buri (khao Kieo). Tak (Umphang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Lin Tin, Pilok, Sai Yok, Thong Phaphum); Ratchaburi (Maenam Pachi); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Klong Bang Lai, Pa-La-U); Chumphon (Mab Ammarit, Pa Tieu, Tasan, Tung Wua-laen); Ranong (Klong Naka); Surat Thani (Khao Sok, Klong Saeng); Krabi (Khao Panom Bencha); Phangnga (Khao Lak Lam Ru).
<i>Cyrtodactylus papilionoides</i>	ตุ๊กแกป่าลายจุดผีเสื้อ Butterfly Gecko	Inhabits evergreen forest.	Nakhon Ratchasima (Pak Chong); Chaiyaphum (Phu Kieo, Thep Satit); Roi Et (Phu Pa nam-tip).
<i>Cyrtodactylus peguensis peguensis</i>	ตุ๊กแกป่าลายจุดเหนือ Common spotted Gecko	Inhabits evergreen forest, usually found at night on tree buttress or forest floor.	Tak (Umphang); Uthai Thani (Ban Rai, Hau Khakhaeng); Kanchanaburi (Sai Yok).
<i>Cyrtodactylus peguensis zebraicus</i>	ตุ๊กแกป่าลายจุดใต้ Zebra Spotted Gecko	Inhabits evergreen forest, usually found at night on tree buttress or forest floor.	Krabi (Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong).
<i>Cyrtodactylus pulchellus</i>	ตุ๊กแกป่าลายพาดป่าใต้ Southern Banded Gecko	Inhabits evergreen forest, usually found at night on rocks along hill streams.	Surat Thani (Klong Saeng); Krabi (Khao Panom Bencha, Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip); Trang (Khao Chong); Phatthalung (Khao Pu

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cyrtodactylus quadrivirgatus</i> (Sungi)	ตุ๊กแกป่าลายสี่ขีด Four-lined Gecko	Inhabits evergreen forest, usually found at night on tree buttress and tree trunk.	Khao Ya); Songkhla (Ton Ngachang); Pattani (Khok Po); Yala (Bannangstaars, Be Tong); Narathiwat (Waeng). Trang (Khao Chong); Yala (Banangstars); Narathiwat Kolok, Waeng).
<i>Cyrtodactylus variegatus</i> Dao,	ตุ๊กแกป่าพม่า Varigated Gecko	Inhabits evergreen forest, usually found at night on rocks along hill streams and in caves.	Mae Hong Son (Pang Ma Pa); Chiang Mai (Doi Chiang Doi Suthep-Pui); Tak (Doi Mon Jong, Um Phang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Thong Paphum).
<i>Dixoneus siamensis</i>	จิ้งจกดินลายจุด Spotted Ground Gecko	Inhabits deciduous and evergreen forest, including beach forest, usually found at night on forest ground.	Chiang Mai (Doi Inthanon); Phrae (Mae Yom); Chaiyaphum (Muang); Loei (Phu Luang); Udon Thani (Phu Phan Kam); Nong Khai (Phu Wua); Kalasin (Khao Wong, Phu Si Than); Nakhon Ratchasima (Sakaerat); Si Sa Ket (Ban Mak Yang); Ubon Ratchathani (Yod Dome); Buriram; Srakaew (Ko Klan, Pang Sida, Taphraya, Wattana Nakhon); Prachin Buri (Prachantakham); Chon Buri (Bangsaen); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Pilok, Salakpra); Phetchaburi (Cha-Um, Kaengkrachan); Prachuap Khirikhan (Bang Saphan, Had Ao Noi, Muang); Chumphon (Lang Suan).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gehyra angusticaudata</i>	จิ้งจกหินหางเรียวยาว Slender-tailed Four-clawed Gecko	This species was reported from old nipa-hut that was being turned down.	Chon Buri (Sriracha).
<i>Gehyra fehlmanni</i>	จิ้งจกหินลายกระ Fehlman's Four-clawed Gecko	This species inhabits disturbed and primary forest, usually found on forest ground and tree buttress.	Nakhon Ratchasima (Khao Yai, Pak Chong); Kanchanaburi; Nakhon Si Thammarat (Khao Luang, Khao Ron Pibun).
<i>Gehyra lacerata</i>	จิ้งจกหินเมืองกาญจน์ Western Four-clawed Gecko	Known from beach forest and deciduous forest, usually found under rock on forest floor.	Nakhon Ratchasima (Sakaerat); Srakaew (Ko Klan, Pang Sida); Chon Buri (Angsila); Kanchaburi; Phetchaburi (Cha-Um).
<i>Gehyra mutilata</i>	จิ้งจกหินสี่จาง Common Four-clawed Gecko	Inhabits all types of forests and other plantations including houses and gardens.	All provinces.
<i>Gekko gekko</i>	ตุ๊กแกบ้าน Tokay Gecko	This species dwells in all habitats of forest and other plantations, including houses and gardens in towns.	All Provinces.
<i>Gekko monarchus</i>	ตุ๊กแกหลังจุดคู่ Malaysian House Gecko	Found in houses.	Phangnga (Khao Lak Lam Ru); Phuket; Pattani (Muang); Narathiwat (Muang, Sungai Kolok, Waeng).
<i>Gekko petricolus</i>	ตุ๊กแกเขาหินทราย	Inhabits deciduous forest, found in	Mukda Han; Kalasin (Khao Wong); Umnajjareon (Phu Singh

Scientific Name	Thai name/Common name	Habitat	Site
	Sandstone Gecko	sandstone crevices and also in houses.	Phu Pa Pung); Ubon (Kaeng Ta Na, Sanoi River; Yod Dome); Roi Et (Nong Pok, Phu Pa Nam Tip); Srakaew (Ko Klan, Tapraya).
<i>Gekko smithi</i>	<sup>๓</sup> ตุ๊กแกสี่เท้า Smith's Gecko	Found in tree hollow in evergreen forest and peat swamp forest, including houses.	Phangnga (Khao Lak Lam Ru); Krabi (Khao Pra Bang Kram); Satun (Tale Ban); Yala (Bannangstars, Be Tong); Narathiwat (Sungai Kolok, Waeng).
<i>Hemidactylus frenatus</i>	<sup>๓</sup> จิ้งจกหางหนาม Spiny-tailed House Gecko	Found in all types of forests and other plantations, also in human houses.	All Provinces.
<i>Hemidactylus garnotii</i>	<sup>๓</sup> จิ้งจกหางเรียบ Garnot's House Gecko	Found in all types of forests and other plantations, also in human houses.	All Provinces.
<i>Hemiphyllodactylus typus</i>	<sup>๓</sup> จิ้งจกเขาสองมลายู Malayan Dwarf Gecko	Found on tree trunk of evergreen forest.	Trat (Klong Mano); Ranong (Muang); Trang (Khao Chong); Yala (Bannangstars).
<i>Hemiphyllodactylus yunnanensis</i>	<sup>๓</sup> จิ้งจกเขาสองยูนนาน Yunnan Dwarf Gecko	Found on tree trunk of evergreen forest and submontane forest.	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Tak (Umphang); Loei (Phu Kradung, Phu Luang); Nakhon Ratchasima (Khao Yai, Sakaerat); Surat Thani (Koh Samui).
<i>Ptychozoon kuhli</i>	<sup>๓</sup> ตุ๊กแกบินหางแบน Kuhl's Parachute Gecko	Found on tree trunk of evergreen forest.	Nakhon Si Thammarat (Pak Panang); Pattani (Napradoo); Satun (Tale Ban); Yala (Betong, Bannangstars); Narathiwat

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ptychozoon lionatum</i>	ตุ๊กแกบินทางหยัก Common Parachute Gecko	Found on tree trunk in evergreen forest.	(Waeng). Mae Hong Son (Pang Tong); Chiang Mai (Doi Suthep-Pui); Tak (Umphang); Uthai Thani (Hau Khakhaeng); Phetchaburi (Kaengkrachan); Ranong Muang); Phuket; Phangnga (Khao Lak Lam Ru); Surat Thani (Khao Sok, Khlong Saeng); Nakhon Si Thammarat (Khao Luang); Krabi (Khao Panom Bencha); Trang (Khao Chong, Koh Libong); Songkhla (Ton Ngachang); Phatthalung (Ta Moat); Satun (Tale Ban); Yala (Bannangstars); Nakhon Ratchasima (Sakaerat, Khao Yai); Chanthaburi (Khao Kitchakud, Khao Sebab, Khao Soi Dao); Trat (Koh Chang); Chon Buri (Hup Bon).
<i>Takydromus sexlineatus ocellatus</i>	กิ้งก่านอขทางยาว Long-tailed Lizard	Inhabits grassland, open woodland and crop plantation.	Mae Hong Son; Chiang Mai; Loei; Roi Et (Phu Pa Nam Tip); Ubon Ratchathani; Nakhon Ratchasima (Sakaerat); Chachoengsao (Khao Ang Rue Ni); Srakaew (Pang Sida); Chanthaburi (Khao Sebab); Pathum Thani (Rang Sit); Bangkok; Chumphon (Thung Wua Lan); Nakhon Si Thammarat; Pattani; Narathiwat (Takbai).
<i>Dasia olivacea</i>	จิ้งเหลนต้นไม้ Olivaceous Tree Skink	Inhabits beach and evergreen forest.	Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida); Chanthaburi (Khao Sebab); Kanchanaburi (Sai Yok); Surat

Scientific Name	Thai name/Common name	Habitat	Site
<i>Isopachys anguinoides</i>	จิ้งเหลนคางประจวบ Striped Legless Skink	Beach and evergreen forest, usually found under humus soil.	Thani (Koh Samui, Kho Tao, Muang); Phangnga (Phangnga Bay); Nakhon Si Thammarat (Khao Luang); Phuket; Trang (Koh Libong); Satun (Tale Ban); Narathiwat (Sukirin). Phetchaburi (Cha-Um, Kaengkrachan); Prachuap Khirikhan (Bang Saphan, Hua Hin, Pa-La-U, Pa Tiu, Tap Sakae, Wanakorn Beach).
<i>Lipinia vittigera</i>	จิ้งเหลนลายอินโดจีน Common Striped Skink	Inhabits deciduous and evergreen forest, usually found on tree trunk.	Chiang Mai (Doi Inthanon); Phrae (Mae Yom); Tak; Lampang; Loei (Tham Pa Pu); Ubon Ratchathani (Lam Dome Yai, Sa Noi River, Yod Dome); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida); Chanthaburi (Khao Sebab, Khao Soi Dao); Nakhon Nayok (Muang); Uthai Thani (Hauai Khakhaeng); Kanchanaburi (Sai Yok); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Klong Banglai, Pa-La-U, Pran Buri); Chumphon (Tasan); Narathiwat (Waeng).
<i>Lygosoma quadrupes</i>	จิ้งเหลนเรียวขาเล็ก Common Slender Skink	Inhabits beach, shrub, deciduous and evergreen forest, including crop plantation and gardens, usually found under rotten logs or humus soil.	Loei (Phu Luang); Nakhon Ratchasima (Sakaerat, Tap Lan); Srakaew (Ta Phraya); Chachoengsao (Khao Ang Rue Ni); Chon Buri (Ang Sila, Sriracha); Chanthaburi (Khao Sebab); Pathum Thani (Rang Sit); Bangkok (Bang Khen); Kanchanaburi; Phetchaburi (Cha-Um, Kaeng Krachan);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Mabuya longicaudata</i>	จิ้งเหลนหางยาว Long-tailed Skink	Inhabits shrub, deciduous and evergreen forest, including coconut plantation.	Prachuap Khirikhan (Pa-La-U); Nakhon Si Thammarat; Trang; Songkhla; Pattani; Yala (Bannangstars); Narathiwat (Waeng). Chiang Mai (Doi Chiang Dao); Loei (Phu Luang); Nong Khai (Phu Wua); Udon Thani; Nakhon Phanom; Chaiyaphum (Phu Kieo); Ubon Rachathani (Yod Dome); Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida); Chon Buri (Bang Saen); Bangkok (Thonburi); Saraburi; Kanchanaburi (Sai Yok); Ratchaburi; Phatthalung (Tale Noi); Songkhla (Tale Sap); Narathiwat (Takbai).
<i>Mabuya macularia malcolmi</i>	จิ้งเหลนหลากลายมัลคอล์ม Malcolm's Skink	Inhabits deciduous and evergreen forest.	Chiang Mai (Doi Suthep-Pui); Loei (Phu Pak Kinak); Uthai Thani (Hau Khakhaeng); Ratchaburi (Maenam Pachi); Phetchaburi (Kaengkrachan); Phatthalung.
<i>Mabuya macularia quadrifasciata</i>	จิ้งเหลนหลากลายสี่ขีด Four-lined Skink	Inhabits deciduous and evergreen forest.	Pattani (Bukit Besar).
<i>Mabuya multifasciata</i>	จิ้งเหลนบาน Common Sun Skink	Inhabits all types of forests, plantations and gardens, including urban areas.	Loei (Phu Kradung); Nong Khai (Phu Wua); Chaiyaphum (Phu Kieo); Roi Et (Phu Pa Nam Tip); Ubon Ratchathani (Yod Dome); Nakhon Ratchasima (Sakaerat); Srakaew (Koklan, Pang Sida); Chanthaburi (Khao Soi Dao).
<i>Mabuya novemcarinata</i>	จิ้งเหลนเกล็ดสั้น Inhabits evergreen forest.	Inhabits evergreen forest.	All provinces.



Scientific Name	Thai name/Common name	Habitat	Site
<i>Mabuya rugifera</i>	Keeled Skink จิ้งเหลนลายขีด	Inhabits lowland evergreen forest, including peat swamp forest.	Nakhon Si Thammarat (Khaoluang, Khao Wang Hip); Trang (Khao Chong); Yala (Bukit Bersa); Narathiwat (Waeng).
<i>Riopa bowringi</i>	Nicobar Skink จิ้งเหลนเรียวทองเหลือง Bowring's Slender Skink	Inhabits beach, deciduous and evergreen forest, including urban areas as parks and home gardens.	Yala (Bannangstars); Narathiwat (Sungai Kolok, Waeng).
<i>Riopa herberti</i>	Herbert's Slender Skink จิ้งเหลนเรียวใต้	Inhabits evergreen forest, usually found under rotten logs on forest floor.	Chiang Mai (Doi Inthanon); Phrae (Mae Yom); Nong Khai (Phu Wua); Roi Et (Phu Pa Namtip); Ubon Ratchathani (Yod Dome); Nakhon Ratchasima (Sakaerat); Srakaew (Koklan, Pang Sida); Chanthaburi (Khao Kittchakud, Khao Sebab, Khao Soi Dao); Pathum Thani (Rang Siit), Bangkok (Pathumwan); Kanchanaburi (Sai Yok); Phetchaburi (Cha-Um, Kaengkrachan); Prachuap Khirikhan (Pa-La-U, Tap Saka); Surat Thani (Koh Samui); Phuket; Trang (Had Chao Mi, Khao Chong, Koh Libong).
<i>Scincella doriae</i>	Doria's Ground Skink จิ้งเหลนดินดอร์เรีย	Inhabits evergreen forest, usually found under rotten logs.	Ranong (Pak Chan River); Phangnga (Khao Lak); Phuket; Surat Thani (Khlung Saeng, Koh Tao, Ta Chana); Nakhon Si Thammarat (Khao Luang); Krabi (Khao Panom Bencha).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Scincella melanosticta melanosticta</i>	จิ้งเหลนดินจุดดำ Black-spotted Ground Skink	Inhabits evergreen forest, found under rotten logs and in humus soil of tree buttress.	Chiang Mai (Doi Inthanon); Nakhon Sawan (Chong Kae); Loei (Phu Luang); Udon Thani (Phuphankam); Nong Khai (Phu Wua); Nakhon Panom (Nakae); Kalasin (Phu Sithan); Sakon Nakhon (Phu Phan); Nakhon Ratchasima (Sakaerat); Ubon Ratchathani (Sa Noi River, Si Muang Mai, Yod Dome); Roi Et (Phu Pa Namtip, Nong Pok); Si Sa Ket (Khao Pra Wihan); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (Koh Chang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Saiyok); Nakhon Si Thammarat (Khao Wang Hip).
<i>Scincella tavesae</i>	จิ้งเหลนดินเมืองกาญจน์ Taves's Ground Skink	Inhabits deciduous and evergreen forest, found under rotten logs and in humus soil of tree buttress.	Surat Thani (Koh Tao).
<i>Scincella reevesi</i>	จิ้งเหลนดินดงพญาเย็น Reeves's Ground Skink	Inhabits evergreen forest, found under rotten logs and in humus soil.	Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Ratchaburi (Chaum Bung).
<i>Sphenomorphus indicus indicus</i>	จิ้งเหลนภูเขาอินเดีย Indian Forest Skink	Inhabits hill forest.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Lampang (Doi Khun Tan, Tak (Thung Yai)); Loei (Phu Kradung, Phu Luang); Nakhon Ratchasima (Sakaerat); Ubon Ratchathani (Sa Noi River); Chanthaburi (Khao Sebab); Uthai Thani (Hau Khakhaeng); Phetchaburi (Kaengkrachan).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Sphenomorphus maculatus</i>	จิ้งเหลนภูเขาเกดสีดเรียบ Common Forest Skink	Inhabits hill and lowland forest.	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Phrae (Mae Yom); Nong Khai (Phu Wua); Nakhon Phanom; Sakon Nakhon (Phu Phan); Kalasin (Phu Si Than); Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida); Chanthaburi (Khao Kitchakud, Khao Sebab, Khao Soi Dao); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Pilok, Saiyok, Salak Pra, Thong Paphum); Phetchaburi (Kaengkrachan); Chumphon; Ranong (Muang); Surat Thani (Khlung Saeng, Khao Sok); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip); Trang (Khao Chong); Narathiwat (Takbai).
<i>Sphenomorphus tersus</i>	จิ้งเหลนภูเขาสีจาง Pale Forest Skink	Inhabits hill forest.	Phetchaburi (Kaengkrachan); Chumphon (Tasan); Ranong (Pak Chan Estuary); Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip).
<i>Tropidophorus bermorei</i>	จิ้งเหลนห้วยทองแดง Bermore's Stream Skink	Inhabits hill streams in evergreen forest.	Mae Hong Son (Lamnam Pai, Namtok Mae Surin); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Phrae (Khao Phlueng); Tak (Umpang, Thung Yai); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Lin Thin, Tan Lod Cave).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Tropidophorus laotus</i>	จิ้งเหลนห้วยลาว Laotian Stream Skink	Inhabits hill streams in evergreen forest.	Loei (Phu Kradung, Phu Luang).
<i>Tropidophorus microlepis</i>	จิ้งเหลนห้วยเล็ก Cambodian Stream Skink	Inhabits hill streams in evergreen forest.	Chanthaburi (Khao Sebab, Khao Soi Dao).
<i>Tropidophorus robinsoni</i>	จิ้งเหลนห้วยชุมพร Robinson's Stream Skink	Inhabits hill streams in evergreen forest.	Chumphon (Tasan); Phangnga (Khao Lak).
<i>Tropidophorus thai</i>	จิ้งเหลนห้วยไทย Thai Stream Skink	Inhabits hill streams in evergreen forest.	Mae Hong Son; Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui, Mae Wang, Pa Miang).
<i>Varanus salvator</i>	เหี้ย Common Water Monitor	Inhabits lowland wetlands and lowland forests, including urban areas. Usually found in canals, swamps, streams and rivers.	All provinces.
<i>Ramphotyphlops albiceps</i>	งูดินหัวขาว White-headed Blind Snake	Inhabits deciduous and evergreen forest, including parks and gardens. This species can be introduced by pot soil.	Chiang Mai (Doi Pui); Nakhon Ratchasima (San Kamphaeng Mts.); Chanthaburi (Khao Sebab, Khao Soi Dao); Bangkok; Ratchaburi (Chaum Bung); Phangnga (Koh Similan); Nakhon Si Thammarat; Pattani; Yala (Betong).
<i>Ramphotyphlops braminus</i>	งูดินบาน Common Blind Snake	Inhabits deciduous and evergreen forest, including parks and gardens. This species can be introduced by pot soil.	Chiang Mai (Doi Suthep-Pui); Phrae (Mae Yom); Udon Thani; Nakhon Ratchasima (Sakaerat); Prachin Buri (Tap Lan); Chanthaburi (Khao Sebab, Khao Soi Dao); Chon Buri

Scientific Name	Thai name/Common name	Habitat	Site
<i>Typhlops diardi</i>	งูดินใหญ่อินโดจีน Diard's Blind Snake	Inhabits evergreen forest, including parks and gardens.	(Siracha); Saraburi; Pathum Thani (Rangsit); Bangkok (Lad Krabang); Kanchanaburi (Salak Pra); Phetchaburi (Cha-Um); Prachuap Khirikhan (Muang); Phangnga (Khao Lak); Phuket; Phatthalung; Nakhon Si Thammarat (Had Nai Phloi); Trang (Koh Libong). Chiang Mai (Doi Chiang Dao); Tak (Umpang); Uthai Thani (Hau Khakhaeng); Loei (Phu Luang); Chaiyaphum (Phu Kieo); Nakhon Panom; Nakhon Ratchasima (Sakaerat); Prachin Buri (Tap Lan); Chon Buri (Koh Lan); Bangkok; Trang.
<i>Typhlops floweri</i>	งูดินหัวเหลือง Flower's Blind Snake	Inhabits evergreen forest, including parks and gardens.	Bangkok; Pathum Thani (Rangsit).
<i>Typhlops khoratensis</i>	งูดินโคราช Khorat Blind Snake	Inhabits evergreen forest, usually found under rotten logs.	Chiang Mai; Nakhon Ratchasima (Sakaerat); Saraburi (Muak Lek); Prachuap Khirikhan (Hua Hin).
<i>Typhlops muelleri</i>	งูดินใหญ่มลายู Mueller's Blind Snake	Inhabits evergreen forest, usually found under rotten logs.	Nakhon Ratchasima (Sakaerat, Tap Lan); Surat Thani (Khao Sok, Koh Samui); Nakhon Si Thammarat (Khao Luang); Phatthalung (Tale Noi); Pattani, Yala (Betong).
<i>Typhlops ozakiae</i>	งูดินปักธงชัย Ozakia's Blind Snake	Inhabits evergreen forest, usually found under rotten logs.	Nakhon Ratchasima (Sakaerat).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Typhlops porrectus</i>	งูดินดอขยุ่ย Doi Pui Blind Snake	Inhabits evergreen forest and parks, usually found under rotten logs.	Chiang Mai (Doi Suthep-Pui); Pathum Thani (Rangsit); Bangkok (Lad Plakao, Pathumwan, Sukumvit).
<i>Cylindrophis ruffus</i>	งูกนขบ Red-tailed Pipe Snake	Inhabits from lowland wetlands to evergreen forest, usually found near water bodies. Consumes on water snakes and eels.	Chiang Mai (Muang); Nong Khai (Phu Wua); Nong Bua Lampu (Muang); Nakhon Ratchasima (Sakaerat); Pathum Thani (Rangsit); Prachin Buri (Prachantakham); Chon Buri (Bang Saen); Bangkok (Bangkhen, Lad Krabang); Phetchaburi (Kaengkrachan); Ranong (Khlong Naka); Surat Thani (Koh Samui); Phangnga (Khao Lak): Phuket; Nakhon Si Thammarat (Khao Luang); Phatthalung (Tale Noi); Songkhla (Ranod); Narathiwat (Sungai Kolok, Waeng).
<i>Xenopeltis unicolor</i>	งูแสงอาทิตย์ Sunbeam Snake	Inhabits from lowland wetlands to evergreen forest.	Chiang Mai; Phrae (Mae Yom); Nakhon Ratchasima (Sakaerat); Saraburi; Pathum Thani (Rangsit); Prachin Buri (Prachantakham); Chon Buri (Bang Saen); Phetchaburi (Cha-Um); Prachuap Khirikhan (Hua Hin); Surat Thani (Koh Samui, Khao Tapetch); Phangnga (Khao Sok): Phuket; Nakhon Si Thammarat (Khao Luang); Yala (Bannangstar); Narathiwat (Sungai Kolok).
<i>Python brongersmai</i>	งูหลามปากเป็ด Brongersma's Python	Inhabits lowland evergreen forest.	Kanchanaburi (Sai Yok); Phetchaburi (Kaengkrachan); Ranong (Khlong Naka); Surat Thani (Khlong Saeng);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Python molurus bivittatus</i>	งูหลาม Burmese Python	Inhabits from lowland wetlands to evergreen forest.	Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Narathiwat (Waeng). Tak; Phetchabun (Nam Nao); Loei (Phu Luang); Nakhon Sawan (Ta Kli); Lop Buri; Pathum Thani (Rang Sit); Prachin Buri (Prachantakham); Nakhon Nayok (Ongkarak); Chachoengsao (Khao Ang Rue Ni); Chon Buri (Siracha); Uthai Thani (Hau Khakhaeng); Surat Thani.
<i>Python reticulatus</i>	งูเห่ล้อม Reticulated Python	Inhabits from lowland wetlands to evergreen forest.	All provinces.
<i>Acrochordus granulatus</i>	งูฝาขี้วัว Granular File Snake	Inhabits estuary areas.	Trat; Samut Prakan; Bangkok; Prachuap Khirikhan (Muang); Songkhla (Ranod); Satun (Tammalang); Phangnga (Phang Nga Bay); Phuket.
<i>Ahaetulla nasuta</i>	งูเขี้ยวปากเหน็บ Long-nosed Whip Snake	Inhabits from parks and gardens to evergreen forest.	Chiang Mai (Doi Suthep); Phrae (Mae Yom); Sing Buri (Prom Buri); Bangkok; Srakaew (Pang Sida); Prachuap Khirikhan (Hua Hin).
<i>Ahaetulla prasina</i>	งูเขี้ยวหัวจิ้งจกป่า Oriental Whip Snake	Inhabits deciduous and evergreen forest, found on bushes and trees. Consumed agama lizards and birds.	Mae Hong Son, Chiang Mai (Doi Inthanon); Phrae (Mae Yom); Chaiyaphum (Phu Kieo); Loei (Phu Luang); Nong Khai (Phu Wua); Nakhon Ratchasima (Khao Yai, Pak Chong,

Scientific Name	Thai name/Common name	Habitat	Site
<i>Boiga cyanea</i>	งูเขียวบอน Green Cat Snake	Inhabits evergreen forest, usually found on bushes and trees. Consumed birds.	Sakaerat); Srakaew (Pang Sida); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab, Khao Soi Dao); Uthai Thani (Hau Khakhaeng); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Surat Thani (Khao Ta Petch, Koh Samui, Koh Tao, Khlong Saeng); Phangnga (Koh Lak); Phuket; Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Songkhla (Ton Ngachang), Yala (Bannangstar Narathiwat (Waeng). Lampang; Loei (Dan Sai, Na Haeo); Nong Khai (Phu Wua); Khon Kaen; Phu Wiang); Sakon Nakhon (Phu Phan); Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Sakaerat); Ubon Ratchathani (Kaeng Tana); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (Klong Manao); Uthai Thani (Hau Khakhaeng); Phetchaburi (Kaengkrachan); Surat Thani (Ban Na San, Ban Takun, Khao Tapetch, Koh Pa Ngan, Tung Tao); Trang (Kan Tang); Phuket; Nakhon Si Thammarat (Khao Luang).
<i>Boiga cynodon</i>	งูแส้หางมา Dog-toothed Cat Snake	Inhabits evergreen forest, usually found on bushes and trees. Consumed birds.	Phangnga (Khao Lak); Surat Thani (Khao Tapetch, Khlong Saeng, Koh Samui), Trang (Khao Chong); Krabi (Khao Pra Bang Kram); Pattani (Napradoo); Narathiwat (Sungi Golok,



Scientific Name	Thai name/Common name	Habitat	Site
<i>Boiga dendrophila melanota</i>	งูปล้องทอง Mangrove Snake	Inhabits evergreen forest, usually found on bushes and trees. Consumed birds and small mammals.	Waeng). Chumphon, Phangnga (Khao Lak); Surat Thani (Khlung Saeng, Koh Samui); Nakhon Si Thammarat; Krabi (Khao Panombencha); Trang (Khao Chong); Phatthalung (Muang); Songkhla (Ton Ngachang); Pattani (Napradoo); Narathiwat (Sungai Kolok, Waeng).
<i>Boiga drapiezii</i>	งูดงคา White-spotted Cat Snake	Inhabits evergreen forest, usually found on bushes and trees. Consumed small geckos.	Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Pattani (Napradoo); Narathiwat (Sungai Kolok, Waeng).
<i>Boiga jaspidea</i>	งูกระ Jasper Cat Snake	Inhabits evergreen forest, usually found on bushes and trees. Consumed small geckos.	Phangnga (Khao Lak); Phuket; Pattani (Napradoo); Narathiwat (Waeng).
<i>Boiga multomaculata</i>	งูแมวตะจาวรังนก Many-spotted Cat Snake	Inhabits deciduous and evergreen forest, usually found on grasses on over ground vegetation.	Chiang Mai (Doi Suthep-Pui); Loei (Phu Kradung, Phu Luang); Kalasin (Phu Sithan); Nakhon Ratchasima (Tap Lan, Wang Namkiao); Srakaew (Pang Sida); Chanthaburi (Khao Soi Dao); Chon Buri (Muang, Ang Sila); Uthai Thani (Hau Khakhaeng).
<i>Boiga nigriceps</i>	งูตองไฟ	Inhabits evergreen forest, has been found	Ratchaburi (Mae Pradone); Trang (Khao Chong); Songkhla

Scientific Name	Thai name/Common name	Habitat	Site
<i>Boiga ocellata</i>	Red Cat Snake งูแสมทางเทา Grey Cat Snake	once in bamboo trunk. Inhabits deciduous and evergreen forest.	(Khao Namkang); Pattani (Bukit Besar); Narathiwat (Waeng). Kamphaeng Phet (Khlung Lan); Uthai Thani (Hau Khakhaeng); Nakhon Ratchasima (Wang Namkio); Ubon Ratchathani (Kaeng Tana); Srakaew (Pang Sida); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab).
<i>Calamaria pavementata</i>	งูพงอท้องเหลือง Collared Reed Snake	Inhabits lowland evergreen forest.	Chiang Mai (Doi Suthep-Pui); Nakhon Ratchasima (Pak Chong, Wang Namkio); Roi Et (Nong Pok); Chanthaburi (Khao Soi Dao); Prachin Buri (Tap Lan); Pattani (Napradoo).
<i>Chrysopelea ornata ornatissima</i>	งูเขียวดอกหมาก Golden Tree Snake	Inhabits from parks and gardens to evergreen forest, including houses.	All provinces.
<i>Coelognathus flavolineatus</i>	งูทางมะพร้าวดำ Common Malayan Racer	Inhabits evergreen forest.	Nakhon Ratchasima (Pak Chong, Wang Namkio); Surat Thani (Khlung Saeng); Phangnga; Nakhon Si Thammarat (Khao Luang); Phuket; Satun; Yala (Bannangstar); Narathiwat (Bang Nara).
<i>Coelognathus radiatus</i>	งูทางมะพร้าวลายขีด Copperheaded Racer	Inhabits from beach forest to hill forest, including urban areas such as parks and gardens, paddy fields and other crop plantations.	All provinces.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Dendrelaphis caudolineatus</i>	งูสายमानหลังลาย Striped Bronzeback	Inhabits evergreen forest.	Chumphon (Khlung Aon, Tapli); Phangnga (Khao Lak); Phuket; Pattani (Napradoo); Yala (Bannangstar); Narathiwat (Waeng).
<i>Dendrelaphis cyanochloris</i>	งูสายमानฟ้าเขียว Blue Bronzeback	Inhabits evergreen forest (including mangrove).	Chiang Mai (Doi Inthanon); Phetchaburi (Kaengkrachan); Trang (Kantang); Phuket; Yala (Bannangstar); Narathiwat (Waeng).
<i>Dendrelaphis formosus</i>	งูสายमानหลังทอง Elegant Bronzeback	Inhabits evergreen forest.	Phuket; Pattani; Yala (Bannangstar); Narathiwat (Waeng).
<i>Dendrelaphis pictus</i>	งูสายमानพระอินทร์ Common Bronzeback	Inhabits from home gardens to hill forests.	All provinces.
<i>Dendrelaphis subocularis</i>	งูสายमानเกล็ดไตดำใหญ่ Mountain Bronzeback	Inhabits deciduous and evergreen forests.	Chiang Mai (Chiang Dao); Nakhon Ratchasima (Sakaerat); Uthai Thani (Hau Khakhaeng); Chumphon (Tasan).
<i>Dryocalamus davisonii</i>	งูปล้องฉนวนอินเดีย Common Bridle Snake	Inhabits from beach forest to hill forest.	Nakhon Ratchasima (Sakaerat); Chanthaburi (Khao Seab, Khao Soi Dao); Chon Buri (Siracha); Bangkok; Uthai Thani (Hau Khakhaeng); Phetchaburi (Cha-Um); Surat Thani (Khlung Saeng).
<i>Dryocalamus subannulatus</i>	งูปล้องฉนวนมลายู Malayan Bridle Snake	Inhabits lowland evergreen forest.	Narathiwat (Waeng).
<i>Dryophiops rubescens</i>	งูเถา	Inhabits evergreen forest.	Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Khao

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gongylosoma scriptus</i>	Brown Whip Snake งูสายทองคอแหวน Common Ringneck	Inhabits lowland evergreen forest.	Luang); Phuket; Trang (Kan Tang, Muang); Pattani (Napradoo, Sai Khao); Narathiwat (Waeng). Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Sakaerat); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Phetchaburi (Phetchaburi basin); Nakhon Si Thammarat; Phuket (Ta Lo Po).
<i>Gonyosoma oxycephalum</i>	งูเขียวกาบหมวก Red-tailed Racer	Inhabits lowland evergreen forest including swamp forest.	Chiang Mai (Doi Suthep-Pui, Doi Inthanon); Phetchabun (Nam Nao); Loei (Phu Luang); Kalasin (Phu Si Than); Roi Et (Nong Pok); Ubon Ratchathani (Yod Dome); Nakhon Ratchasima (Khao Yai, Sakaerat); Chanthaburi (Khao Kitchakud, Khao Soi Dao); Kanchanaburi (Sai Yok); Phetchaburi (Kaengkrachan); Surat Thani (Khlong Saeng, Koh Tan); Krabi (Khao Panombencha); Phangnga; Trang (Khao Chong); Satun (Tale Ban); Pattani; Narathiwat (Sungai Kolok, Waeng).
<i>Gonyosoma prasina</i>	งูทางมะพร้าวเขียว Green Mountain Racer	Inhabits hill and submontane forest.	Chiang Mai (Doi Chiang Dao); Loei (Phu Luang).
<i>Lycodon capucinus</i>	งูสรอยเหลือง Common Wolf Snake	Inhabits lowland evergreen forest including parks and gardens.	Chiang Mai (Doi Suthep-Pui, Doi Inthanon); Phrae (Mae Yom); Phetchabun (Nam Nao); Loei (Phu Luang); Roi Et

Scientific Name	Thai name/Common name	Habitat	Site
<i>Lycodon effraensis</i>	งูปล้องจนวนแดง Brown Wolf Snake	Inhabits lowland evergreen forest.	(Phu Pa Namtip); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida); Chon Buri; Lop Buri; Bangkok; Uthai Thani (Hauai Khakhaeng); Kanchanaburi (Salak Pra); Phetchaburi (Kaengkrachan); Ranong; Phangnga; Phuket. Nakhon Si Thammarat (Thung Song); Yala (Bannangstar); Narathiwat (Waeng).
<i>Lycodon fasciatus</i>	งูปล้องจนวนเมืองเหนือ Indian Banded Wolf Snake	Inhabits lowland evergreen forest.	Mae Hon Son (Pang Tong); Chiang Mai (Doi Suthep-Pui, Doi Inthanon); Tak (Umpang); Phrae (Mae Yom); Phetchabun (Nam Nao); Loei (Phu Luang); Nakhon Ratchasima (Khao Yai, Sakaerat); Chachoengsao (Khao Ang Rue Ni); Uthai Thani (Hauai Khakhaeng); Phetchaburi (Kaengkrachan).
<i>Lycodon laotus</i>	งูปล้องจนวนลาว Laotian Wolf Snake	Inhabits lowland evergreen forest.	Chiang Mai (Doi Suthep-Pui, Doi Inthanon); Nong Khai (Phu Wua); Nakhon Ratchasima (Khao Yai, Sakaerat); Si Sa Ket (Ban Mak Yang); Srakaew (Pang Sida); Trat (Koh Chang); Rayong; Saraburi (Muak Lek); Prachuap Khirikhan (Hauai Yang, Thap Sakae); Surat Thani (Khao Tapetch); Krabi (Khao Panombencha); Nakhon Si Thammarat (Ron Pibun, Thung Song); Trang (Had Chao Mai); Pattani (Napradoo).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Lycodon subcinctus</i>	งูปลอดจนวนนพาน Malaysian Banded Wolf Snake	Inhabits lowland evergreen forest.	Phrae (Mae Yom); Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida); Surat Thani (Khlung Saeng); Nakhon Si Thammarat (Ron Pibun, Thung Song); Pattani (Napradoo); Narathiwat (Tak Bai, Waeng).
<i>Oligodon barroni</i>	งูบีแก้วลายหัวใจ Barron's Kukri Snake	Inhabits lowland evergreen forest, parks and gardens.	Saraburi (Muak Lek); Chon Buri (Siracha); Rayong (Muang).
<i>Oligodon catenatus</i>	งูบีแก้วภูหลวง Phuluang Kukri snake	Inhabits submontane forest.	Loei (Phu Luang).
<i>Oligodon cinereus</i>	งูบีแก้วลายกระ Ashy Kukri Snake	Inhabits lowland and submontane forests.	Chiang Mai (Chiang Dao, Muang); Nakhon Ratchasima (Sakaerat).
<i>Oligodon fasciolatus</i>	งูบีแก้วลายแถม Banded Kukri Snake	Inhabits lowland forest, including parks and gardens.	Chiang Mai (Doi Suthep); Nakhon Ratchasima (Sakaerat); Ubon Ratchathani (Yod Dome); Srakaew (Tapraya); Chanthaburi (Khao Sebab, Khao Soi Dao); Chon Buri (Bang La Mung); Bangkok; Chumphon; Nakhon Si Thammarat (Lan Saka).
<i>Oligodon inornatus</i>	งูบีแก้วสีจาง Inornate Kukri Snake	Inhabits evergreen and deciduous forest.	Loei; Trat; Chon Buri; Tak; Uthai Thani (Hau Khakhaeng).
<i>Oligodon joynsoni</i>	งูบีแก้วใหญ่	Inhabits deciduous and evergreen forests.	Lampang (Mae Wang, Ngao); Kamphaeng Phet (Khlung Lan).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Oligodon purpurascens</i>	Joynson's Kukri Snake งูกุด Brown Kukri Snake	Inhabits lowland and swampy forests.	Surat Thani (Khao Tapetch); Pattani (Napradoo); Narathiwat (Tak Bai, Waeng).
<i>Oligodon taeniatus</i>	งูจอกไทย Striped Kukri Snake	Inhabits lowland forest including parks and gardens.	Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Sakaerat); Si Sa Ket; Bangkok; Pathum Thani (Rangsit); Saraburi; Nonthaburi; Prachuap Khirikhan (Hua Hin); Pattani (Napradoo).
<i>Orthriophis taeniura yunnanensis</i>	งูกาบหมากยูนนาน Yunnan Racer	Inhabits hill forest and caves nearby.	Loei (Phu Luang); Kanchanaburi (Sai Yok, Salak Pra); Phetchaburi (Khao Yoi).
<i>Orthriophis taeniura ridleyi</i>	งูกาบหมากดำ Ridley's Racer	Inhabits hill forest and limestone caves.	Surat Thani (Khlong Saeng); Nakhon Si Thammarat (Kaeo Surakan Cave); Songkhla (Ton Ngachang); Satun; Yala (Bannangstar, Thanto); Narathiwat (Waeng).
<i>Psammodynastes pulverulentus</i>	งูหมอก Mock Viper	Inhabits evergreen forest.	Chiang Mai (Doi Suthep-Pui, Kaeng Pan Tao); Lampang (Doi Khun Tan); Loei (Phu Luang); Nong Khai (Phu Wua); Nakhon Ratchasima (Khao Yai, Pak Chong; Wang Namkieo); Srakaew (Pang Sida); Chanthaburi (Khao Sebab); Trat (Khao Saming); Chon Buri (Siracha); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Phetchaburi (Kaengkrachan);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ptyas carinatus</i>	งูสิงหางดำ Keeled Rat Snake	Inhabits evergreen forest.	Prachuap Khirikhan (Pran Buri); Surat Thani (Khao Sok, Khlong Saeng, Thung Tao); Nakhon Si Thammarat (Khao Luang, Thung Song); Phuket; Pattani; Narathiwat. Chiang Mai; Lampang (Ngao); Tak (Umpang); Kanchanaburi (Salak Pra); Nakhon Si Thammarat (Khao Luang); Krabi (Khao Panombencha); Trang (Khao Chong); Pattani; Yala (Bannangstar).
<i>Ptyas korros</i>	งูสิงบ้าน Indochinese Rat Snake	Inhabits from agricultural areas to hill forest.	All provinces.
<i>Ptyas mucosus</i>	งูสิงหางลาย Oriental Rat Snake	Inhabits from agricultural areas to hill forest.	All provinces.
<i>Sibynophis collaris</i>	งูคอกวันดำ Common Blackhead	Inhabits evergreen forest.	Chiang Mai (Doi Chiang Dao, Doi Suthep-Pui, Khaeng Pan Tao); Loei (Phu Luang); Chaiyaphum (Phu Kieo); Kanchanaburi (Sai Yok).
<i>Sibynophis melanocephalus</i>	งูคอกวันหัวดำ Malayan Blackhead	Inhabits from beach forest to hill forest.	Nakhon Si Thammarat (Khao Luang Mts.); Pattani (Napradoo); Narathiwat (Tak Bai).
<i>Sibynophis triangularis</i>	งูคอกวันหัวลายสามเหลี่ยม Triangle Blackhead	Inhabits evergreen forest.	Nakhon Ratchasima (Pak Chong, Wang Namkiao); Chachoengsao (Khao Ang Rue Ni); Trat (Koh Chang);



Scientific Name	Thai name/Common name	Habitat	Site
<i>Xenelaphis hexagonotus</i>	งูควนขนุน Malayan Brown Snake	In Thailand this species has been found inhabiting peat swamp forest.	Chon Buri (Siracha). Phatthalung (Tale Noi); Narathiwat (Sungai Kolok).
<i>Cerberus rynchops</i>	งูปากกว้างน้ำเค็ม Dog-faced Water Snake	Inhabits mud flat and mangrove area.	Chon Buri (Ang Sila); Bangkok; Samut Prakan (Bang Pu); Phetchaburi (Ban Laem); Chumphon (estuarine area); Nakhon Si Thammarat (Pak Panang); Pattani (Yaring); Narathiwat (Tak Bai, estuarine area); Satun (Tammalang); Trang (Hat Chao Mi, Kan Tang, Koh Libong); Phangnga (Phangnga Bay); Phuket.
<i>Enhydryis bocourti</i>	งูไซ Bocourt's Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Nakhon Sawan (Bung Borapet); Prachin Buri (Prachantakham); Bangkok; Surat Thani (Nong Tung Tong); Nakhon Si Thammarat (Prom Kiri); Phatthalung (Tale Noi); Pattani.
<i>Enhydryis enhydryis</i>	งูสายรุ้ง Rainbow Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Chiang Mai (Hauai Kaeo); Chaiyaphum (Phu Kieo); Khon Kaen (Ubonratana Reservoir); Kalasin (Phu Sitan); Nakhon Ratchasima (Pak Thonchai, Lam Kakong Reservoir); Srakaew (Pang Sida); Prachin Buri (Prachantakham); Pathum Thani (Rangsit); Nakhon Sawan (Bung Borapet); Bangkok; Nakhon Si Thammarat (Prom Kiri); Phatthalung (Tale Noi); Songkhla (Tale Sap); Pattani; Narathiwat (Tak Bai; Sungai Kolok).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Enhydryis plumbea</i>	งูปลิง Plumbeous Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Mae Hong Son (Pang Tong); Lampang (Doi Pa Muang); Phrae (Mae Yom); Nakhon Sawan (Bung Borapet); Loei (Wang Saphung); Nong Khai (Phu Wua); Khon Kaen (Sum Sung); Ubon Ratchathani (Kaeng Tana); Nakhon Ratchasima (Pak Tongchai, Wang Namkiao); Srakaew (Tapraya); Prachin Buri (Prachantakham); Chachoengsao (Khao Ang Rue Ni); Chon Buri (Bang Pra); Bangkok; Phetchaburi (Kaengkrachan); Nakhon Si Thammarat (Thung Yai); Phatthalung (Tale Noi); Songkhla (Tale Sap); Narathiwat (Sungai Kolok).
<i>Enhydryis subtaeniata</i>	งูสายรุ้งลาย Striped Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Nakhon Sawan (Bung Borapet); Nakhon Ratchasima (Pak Tongchai); Khon Kaen; Udon Thani; Prachin Buri (Prachantakham).
<i>Erpeton tentaculatum</i>	งูกระดาง Tentacled Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Bangkok; Pathum Thani (Rangsit); Nakhon Pathom (Bang Laen); Prachin Buri (Prachantakham); Nakhon Si Thammarat (Prom Kiri); Phatthalung (Tale Noi).
<i>Homalopsis buccata</i>	งูหัวกะโหลก Puff-faced Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Chaiyaphum (Phu Kiao); Ubon Ratchathani (Kaeng Tana); Prachin Buri (Prachantakham); Chachoengsao (Khao Ang Rue Ni); Pathum Thani (Rangsit); Bangkok; Samut Prakan (Bang Pli); Ratchaburi; Ranong; Phangnga (Khao Lak); Surat Thani

Scientific Name	Thai name/Common name	Habitat	Site
<i>Amphiesma stolatum</i>	งูลายสาบดอกหญ้า White-Striped Keelback	Inhabits from scrub to hill forests, including agricultural areas.	(Khlung Saeng, Koh Samui); Nakhon Si Thammarat (Lan Saka, Prom Kiri); Trang (Khao Chong); Phatthalung (Tale Noi); Songkhla (Tale Sap); Pattani (Napradoo); Narathiwat (Tak Bai, Sungai Kolok). Chiang Mai (Doi Suthep, Haui Kaeo); Lamphun (Muang); Nakhon Sawan; Phetchabun (Nam Nao); Udon Thani; Nakhon Phanom (Na Kae); Khon Kaen; Nakhon Ratchasima (Sakaerat); Si Sa Ket (Ban Mak Taeng); Lop Buri; Chai Nat; Saraburi; Bangkok; Ratchaburi (Khao Pratapchang).
<i>Rhabdophis chrysargos</i>	งูลายสาบจุดดำขาว Speckled-bellied Keelback	Inhabits hill deciduous and evergreen forests.	Chiang Mai; Tak (Thung Yai); Nakhon Ratchasima (Khao Yai, Tap Lan); Roi Et (Nak Prok); Chanthaburi (Khao Sebab, Khao Soi Dao); Trat (Klong Manao, Koh Chang); Rayong (Khao Chamao); Kachanaburi (Sai Yok); Prachuap Khirikhan (Klong Bang Lai); Chumphon (Mab Ammarit); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip); Phangnga (Khao Lak); Trang (Khao Chong); Krabi (Sa Bokkorani); Narathiwat (Waeng).
<i>Rhabdophis nigrocinctus</i>	งูลายสาบเขียววันดำ Green Keelback	Inhabits deciduous and evergreen forests.	Chiang Mai (Doi Inthanon, Doi Suthep, Doi Pahompok); Loei (Phu Luang); Nakhon Ratchasima (Khao Yai);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rhabdophis subminiatus subminiatus</i>	งูลายสาบคอแดง Red-necked Keelback	Inhabits deciduous and evergreen forests, including agricultural areas.	Srakaew (Pang Sida); Chanthaburi (Khao Sebab); Kanchanaburi; Phetchaburi (Kaengkrachan); Chumphon (Tapli, Tasan); Phangnga (Khao Lak); Surat Thani (Khlong Saeng); Trang (Khao Chong); Phuket. Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Loei (Phu Luang); Chaiphaphum (Phu Kieo); Phetchabun (Nam Nao); Nong Khai (Phu Wua); Kalasin (Phu Si Tan); Ubon Ratchathani (Kaeng Tana, Yod Dome); Buri Ram (Hau Sala); Nakhon Ratchasima (Khao Yai); Chanthaburi (Khao Sebab, Khao Soi Dao); Bangkok; Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok, Thong Pa Phum); Phetchaburi (Kaengkrachan); Ranong (Muang); Phangnga (Hat Taimuang; Khao Lak); Surat Thani (Khlong Saeng); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Phuket; Narathiwat.
<i>Sinonatrix percarinata</i>	งูลายสอจีน Chinese Keelback	Inhabits hill stream in evergreen forests.	Chiang Mai (Doi Saket; Doi Sutep-Pui); Loei (Phu Luang); Nakhon Ratchasima (Pak Tongchai).
<i>Xenochrophis flavipunctatus</i>	งูลายสอสวน Common Keelback	Inhabits from lowland wetlands to lowland evergreen forest.	Thai localities.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Xenochrophis piscator</i>	งูลายสอใหญ่ Checkedred Keelback	Inhabits from lowland wetlands to lowland evergreen forest.	Mae Hong Son (Salawin); Chiang Mai (Muang); Tak (Thung Yai); Nakhon Ratchasima (Pak Tongchai); Srakaew (Pang Sida); Trat (Koh Chang); Bangkok (Bang Khen); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok); Prachuap Khirikhan (Pa-La-U).
<i>Xenochrophis punctulatus</i>	งูลายสอพม่า Burmese Keelback	Unknown.	Mae Hong Son (Muang).
<i>Xenochrophis trianguligerus</i>	งูลายสอลายสามเหลี่ยม Triangle Keelback	Inhabits lowland evergreen forest.	Kanchanaburi (Sangkhlaburi, Tong Pa Phum); Prachuap Khirikhan (Pa-La-U); Chumphon (Patyu); Ranong (Muang); Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip); Pattani (Napradoo); Narathiwat (Waeng).
<i>Aplopeltura boa</i>	งูกินทากหัวโทนก Blunt-headed Tree Snake	Inhabits lowland evergreen forest.	Chumphon (Phuket Mts.); Nakhon Si Thammarat (Khao Luang); Pattani (Napradoo); Narathiwat (Waeng).
<i>Asthenodipsas laevis</i>	งูกินทากเกล็ดเรียบ Smooth Slug Snake	Inhabits lowland evergreen forest.	Nakhon Si Thammarat (Khao Luang, Khao Ram); Narathiwat (Waeng).
<i>Asthenodipsas malaccanus</i>	งูกินทากมลายู Malaccan Slug Snake	Inhabits lowland evergreen forest, including swampy forest.	Yala (Betong, Than To); Narathiwat (Waeng).
<i>Pareas carinatus</i>	งูกินทากเกล็ดสัน	Inhabits lowland evergreen and	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao,

Scientific Name	Thai name/Common name	Habitat	Site
	Keeled Slug Snake	submontane forest.	Doi Inthanon, Doi Suthep-Pui); Loei (Phu Luang); Nakhon Ratchasima (Pak Chong, Pak Tongchai); Srakaew (Pang Sida); Chanthaburi (Khao Sebab, Khao Soi Dao); Chachoengsao (Khao Ang Rue Ni); Phetchaburi (Kaengkrachan); Chumphon (Khlong Ton Si); Nakhon Si Thammarat (Khao Luang); Narathiwat (Waeng).
<i>Pareas macularius</i>	งูกินทากจุดดำ Black-spotted Slug Snake	Inhabits lowland evergreen and submontane forest.	Chiang Mai (Doi Inthanon); Loei (Phu Luang).
<i>Pareas margaritophorus</i>	งูกินทากจุดขาว White-spotted Slug Snake	Inhabits lowland evergreen forest.	Chiang Mai (Doi Ang Kang, Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Phrae (Mae Yom); Phetchabun (Nam Nao); Loei (Phu Kradung, Phu Luang); Nakhon Ratchasima (Pak Chong, Pak Tongchai); Chanthaburi (Khao Sebab, Khao Soi Dao); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Tong Pa Phum); Ratchaburi (Suan Pung); Phetchaburi (Kaengkrachan); Ranong (Khlong Naka); Nakhon Si Thammarat (Khao Luang); Pattani (Napradoo); Narathiwat (Waeng).
<i>Psammophis indochinensis</i>	งูมานทอง Indochinese Sand Snake	Inhabits lowland areas from grassland to evergreen forest.	Chiang Mai (Doi Suthep-Pui); Chaiyaphum (Phu Kieo); Saraburi; Lop Buri; Pathum Thani (Rangsit); Bangkok; Ratchaburi (Potharam).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Plagiopholis nuchalis</i>	งูหัวลายสร Assamese Mountain Snake	Inhabits submontane forest, usually found on forest ground.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Pahompok, Doi Suthep-Pui); Lampang (Doi Khun Tan); Loei (Phu Luang).
<i>Pseudoxenodon macrops</i>	งูลายสาบตาโต Big-eyed Mountain Keelback	Inhabits submontane forest, usually found on forest ground or temporary ponds.	Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Lampang (Doi Pa Muang); Phitsanulok (Phu Hin Rong Kla); Loei (Dan Sai, Phu Luang); Phetchabun (Nam Nao); Nakhon Ratchasima (Khao Yai); Chanthaburi (Khao Soi Dao).
<i>Bungarus candidus</i>	งูทับสมิงคลา Blue Krait	Inhabits from lowland wetland to lowland evergreen forest, usually found near water bodies.	Chiang Mai (Doi Chiang Dao, Haui Kaeo); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Loei (Phu Luang); Nong Khai (Phu Wua); Mukda Han; Kalasin (Phu Si Than); Ubon Ratchathani (Kaeng Tana, Yod Dome); Buri Ram (Nang Rong); Nakhon Ratchasima (Khao Yai); Chanthaburi (Khao Sebab); Trat (Koh Kut); Chon Buri (Siracha); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Pattani (Napradoo); Narathiwat (Waeng).
<i>Bungarus fasciatus</i>	งูสามเหลี่ยม Banded Krait	Inhabits from lowland wetland to lowland evergreen forest, usually found near water bodies. It consumes water snakes.	Chiang Mai (Muang); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Loei (Phu Luang); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Ubon Ratchathani (Kaeng Tana, Yod Dome); Roi Et (Nong Pok); Nakhon Ratchasima (Khao Yai); Chanthaburi (Khao Kitchakud, Khao Sebab);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Naja kaouthia</i>	งูเห่าหม้อ Monocled Cobra	Inhabits from lowland wetland to lowland evergreen forest. usually found in paddy field.	Bangkok (Bangkok Noi, Bangkok Yai, Nong Jok); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Khao Laem); Ranong (Khlong Naka); Surat Thani (Khao Tapetch, Nong Tung Tong, Thung Tao); Nakhon Si Thammarat (Prom Kiri); Narathiwat (Tak Bai). All provinces.
<i>Naja siamensis</i>	งูเห่าสยาม Siamese Cobra	Inhabits from agricultural areas to deciduous and evergreen forest.	Phrae (Mae Yom); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Ubon Ratchathani (Yod Dome); Si Sa Ket (Ban Mak Yang); Nakhon Sawan (Ta Kli); Uthai Thani (Lan Sak); Chai Nat (Manorom); Lop Buri (Chaibadan, Kok Samrong); Saraburi (Muak Lek); Chanthaburi (Khao Kitchakud); Chon Buri (Bang Pra).
<i>Naja sumatrana</i>	งูเห่าสุมาตรา Sumatran Cobra	Inhabits hill forest.	Phangnga (Khao Lak); Krabi (Khao Nor Chuchi); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Yala (Bannangstar); Narathiwat (Waeng).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Ophiophagus hannah</i>	งูจงอาง King Cobra	Inhabits evergreen forest.	Mae Hong Son; Ching Mai (Doi Inthanon); Tak (Thung Yai); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Loei (Phu Luang); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Ubon Ratchathani (Yod Dome); Buri Ram (Nang Rong); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida); Prachin Buri (Tap Lan); Chanthaburi (Khao Kitchakud, Khao Soi Dao); Uthai Thani (Hauai Khakhaeng); Kanchanaburi (Sai Yok, Tong Pa Phum); Surat Thani (Khleng Saeng); Phetchaburi (Kaengkrachan); Krabi (Khao Panombencha); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Yala (Khao Kampan); Narathiwat (Waeng).
<i>Hydrophis brookii</i>	งูแสมรังทองเหลือง Brook's Sea Snake	Inhabits Lagoon, sea and ocean. It is also found in coral reefs.	Gulf of Thailand Waters and Songkhla Lake.
<i>Pelamis platura</i>	งูขาซงหลังดำ Yellow-bellied Sea Snake	Gulf of Thailand and Andaman Waters.	Gulf of Thailand and Andaman Sea.
<i>Caloselasma rhodostoma</i>	งูกะปะ Malayan Pit Viper	Inhabits from beach forest to lowland evergreen forest, including rubber plantations closed to the forests.	Chiang Rai; Chiang Mai; Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Nong Khai (Phu Wua); Sakon Nakhon; Udon Thani (Kumpawapi); Kalasin (Phu Si Than); Ubon Ratchathani (Kaeng Tana, Yod Dome); Roi Et (Nong Pok); Nakhon

Scientific Name	Thai name/Common name	Habitat	Site
<i>Trimeresurus (Cryptelytrops) albolabris albolabris</i>	งูเขียวหางไหม้ทองเหลือง Yellow-lipped Green Pit Viper	Inhabits from lowland evergreen to submontane forest, including rubber plantations closed to the forests.	Ratchasima (Sakaerat); Srakaew (Pang Sida); Chanthaburi (Khao Soi Dao); Rayong (Khao Chamao); Chon Buri (Siracha); Lop Buri; Uthai Thani (Hau Khakhaeng); Kanchanaburi; Phetchaburi (Cha-Um); Prachuap Khirikhan (Bang Saphan); Chumphon (Thung Wualan); Nakhon Si Thammarat Ron Phibun); Trang; Phuket; Pattani; Narathiwat (Takbai). Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon; Doi Suthep); Phetchabun (Nam Nao); Loei (Phu Luang); Nong Khai (Phu Wua); Ubon Ratchathani (Kaeng Tana); Roi Et (Phu Pa Namtip); Nakhon Ratchasima (Sakaerat); Chanthaburi (Khao Seab); Phra Nakhon Si Ayutthaya; Bangkok; Phetchaburi (Kaeng Krachan); Prachuap Khirikhan (Hau Yang, Pa-La-U); Surat Thani (Khlung Saeng, Thung Tao); Nakhon Si Thammarat (Thung Yai).
<i>Trimeresurus (Cryptelytrops) kanburiensis</i>	งูหางเหี้ยมกาญจนบุรี Kanburi Green Pit Viper	Inhabits deciduous and evergreen forest. This species was rediscovered by Warrell <i>et al.</i> (1992) from the bamboo forest at	Kanchanaburi (Sai Yok).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Trimeresurus (Cryptelytrops) macrops</i>	งูเขียวหางไหม้ตาโต Large-eyed Green Pit Viper	Nong Bu Wa after the first discovery 50 years. Inhabits from parks and gardens to evergreen forest, including flower pots in backyard of town houses.	Nakhon Ratchasima (Pak Chong, Sakaerat); Chanthaburi (Khao Sebab, Khao Soi Dao); Sing Buri (Prom Buri); Bangkok (Bangkhen); Pathum Thani (Rangsit); Samut Prakan (Muang).
<i>Trimeresurus (Cryptelytrops) purpureomaculatus</i>	งูพังกา Mangrove Pit Viper	Inhabits mangrove forest.	Bangkok (Bang Kuntien); Surat Thani (Bandon); Nakhon Si Thammarat (Pak Panang); Satun (Tammalang); Trang (Kantang, Koh Libong); Krabi; Ranong (Maenam La Un; Koh Surin).
<i>Trimeresurus (Cryptelytrops) venustus</i>	งูเขียวหางไหม้ทู่สง Brown-spotted Green Pit Viper	Inhabits evergreen forests.	Surat Thani (Khao Tapetch, Tai Rom Yen N.P.); Nakhon Si Thammarat (Khao Luang, Thungsong); Krabi (Khao Pra Bang Kram).
<i>Ovophis monticola convictus</i>	งูหางหมึกภูเขา Mountain Pit Viper	Inhabits submontane forest.	Loei (Phu Luang); Chiang Mai (Doi Angkang); Tak (Umpang); Kanchanaburi (Thong Pa Phum); Ranong (Khlung Naka).
<i>Trimeresurus (Parias) hageni</i>	งูเขียวหางไหม้ฮาเจน Hagen's Green Pit Viper	Inhabits lowland evergreen forest.	Surat Thani (Ron Pibun); Songkhla (Ton Ngachang); Trang (Khao Chong); Narathiwat (Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Trimeresurus (Popeia) fugatus</i>	งูเขียวหางไหม้ทองเขียวโต Banded Green Pit Viper	Inhabits lowland evergreen forest.	Chumphon (Pa To); Phangnga (Khao Lak); Surat Thani (Khlung Saeng; Krabi (Khao Nor Chuchi); Nakhon Si Thammarat (Lan Saka, Thungsong).
<i>Trimeresurus (Popeia) nebularis</i>	งูเขียวหางไหม้ทองเขียวมลายู Green-eyed Pit Viper	Inhabits submountain forest.	Narathiwat (Waeng).
<i>Trimeresurus (Popeia) popeiorum</i>	งูเขียวหางไหม้ทองเขียวเหนือ Red-eyed Pit Viper	Inhabits lowland evergreen forest.	Chiang Rai; Chiang Mai (Doi Inthanon); Uthai Thani (Hau Khakhaeng); Phetchaburi (Kaengkrachan).
<i>Trimeresurus (Viridovipera) gumprechtii</i>	งูเขียวไผ่ Bamboo Pit Viper	Inhabits lowland evergreen and submontane forest. It is usually found on bamboo branches along forest streams.	Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo); Loei (Phu Luang).
<i>Trimeresurus (Viridovipera) vogeli</i>	งูเขียวหางเทา Grey-tailed Pit Viper	Inhabits lowland evergreen forest.	Nakhon Ratchasima (Khao Yai); Chanthaburi (Khao Soi Dao).
<i>Daboia russelii siamensis</i>	งูแมวเซา Siamese Russell's Viper	Inhabits lowland paddy fields.	Nakhon Sawan (Muang, Takli); Chai Nat (Manorom); Lop Buri; Saraburi; Ang Thong; Phra Nakhon Si Ayutthaya (Bang Pa-in); Pathum Thani (Ladlumkaeo, Rangsit), Nonthaburi (Sai Noi); Nakhon Pathom (Bang Lane); Samut Prakan (Bang Pli, Samrong).

# Status DD

Scientific Name	Thai name/Common name	Habitat	Site
<i>Caretta caretta</i>	เตาหัวโต Logger-headed Sea Turtle	Inhabits seas and oceans, nested on sand beaches.	No record of nesting sites in Thailand sea, but one individual was formerly caught in the Andaman Sea near Phuket Waters and another one was caught in Gulf of Thailand near Rayong Waters.
<i>Pyxidea mouhoti</i>	เตาจัน Mouhot's Turtle	Mainly reported from forested hill areas and lowland swamps.	Uncertainly along Thai-Laos Border.
<i>Dibamus alfredi</i>	จิ้งเหลนดวงสีม่วง Alfred's Snake Skink	Found under rotten log and debris on forest floor.	Pattani (Napradoo); Yala (Thanto).
<i>Dibamus somsaki</i>	จิ้งเหลนดวงเขาสอยดาว Somsak's Snake Skink	Found under rotten log and debris on forest floor.	Chanthaburi (Khao Soi Dao).
<i>Aeluroscalabotes felinus</i>	ตุ๊กแกหัวโต Fox Gecko	Unknown.	Yala (Bannangstar); Narathiwat (Waeng).
<i>Cnemaspis phuketensis</i>	จิ้งจกนิ้วขาวภูเก็ต Phuket Gecko	Unknown.	Phuket.
<i>Cyrtodactylus thirakhupti</i>	ตุ๊กแกปากำธร, ตุ๊กแกป่าสุราษฎร์ Thirakeept's Gecko	Inhabits caves.	Surat Thani.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Dixoneus hangseesom</i>	จิ้งจกตีนหางสีส้ม Orange-tailed Ground Gecko	Inhabits bamboo forest, usually found at night on forest floor.	Kanchanaburi (Sai Yok).
<i>Dixoneus melanostictus</i>	จิ้งจกตีนข้างดำ Black-sided Ground Gecko	Inhabits evergreen forest, usually found at night on forest ground.	Nakhon Ratchasima (Pak Chong); Saraburi (Muak Lek).
<i>Ptychozoon horsfieldi</i>	ตุ๊กแกบินหางเฟิน Horsfield's Parachute Gecko	Unknown.	Unknown.
<i>Ptychozoon trinotatera</i>	ตุ๊กแกบินลายสามแถบ Three-banded Parachute Gecko	Found on tree trunk in evergreen forest.	Nakhon Ratchasima (Sakaerat).
<i>Lipinia quadrivittata</i>	จิ้งเหลนลายสี่ขีด Four-lined Skink	Inhabits evergreen forest, may be found under rotten log.	Pattani (Bukit Besar).
<i>Lipinia surda</i>	จิ้งเหลนลายมลายู Malayan Strip Skink	Inhabits evergreen forest.	Nakhon Si Thammarat (Khao Luang); Yala (Betong).
<i>Lygosoma anguinum</i>	จิ้งเหลนเรียวพม่า Burmese Sleuder Skink	Inhabits deciduous and evergreen forest.	Mae Hong Son (Lamnam Pai); Prachuapkhirikhan (Bang Saphan); Chumphon (Mab Ammarit).
<i>Mabuya macularia postnasalis</i>	จิ้งเหลนหลากลายดานซาย Dansai Skink	Inhabits evergreen forest.	Loei (Phu Lom Lo, Dan Sai).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Sphenomorphus butleri</i>	จิ้งเหลนภูเขามลายู Butler's Forest Skink	Unknown.	Mae Hong Son; Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep); Lampang (Doi Khun Tan); Phrae (Mae Yom); Loei (Phu Luang, Ta Li); Ubon Ratchathani (Sa Noi River); Nakhon Ratchasima (Sakaerat); Prachin Buri (Kabin Buri).
<i>Sphenomorphus grandisonae</i>	จิ้งเหลนภูเขาแกรนิตินัน Grandison's Forest Skink	Unknown.	Northern (Ban Tong Pheung).
<i>Sphenomorphus helenae</i>	จิ้งเหลนภูเขาเฮเลน Helen's Forest Skink	Unknown.	Nonthaburi.
<i>Sphenomorphus lineopunctulatus</i>	จิ้งเหลนภูเขาอุบล Ubon Forest Skink	Unknown.	Ubon Ratchathani (Sa Noi River).
<i>Sphenomorphus mimicus</i>	จิ้งเหลนภูเขาเล็ก Minute Forest Skink	Inhabits hill forest.	Loei (Phu Luang); Nakhon Ratchasima (Pak Chong).
<i>Sphenomorphus presignis</i>	จิ้งเหลนภูเขาตาขกระ Larut Forest Skink	Inhabits hill forest.	Nakhon Si Thammarat (Khao Wang Hip).
<i>Sphenomorphus scotophilus</i>	จิ้งเหลนภูเขาตาขแฉก Selangnor Forest Skink	Inhabits hill forest.	Trang (Khao Chong).
<i>Sphenomorphus stellatus</i>	จิ้งเหลนภูเขาจูดาว Cambodian Forest Skink	Inhabits hill forest.	Chanthaburi (Khao Sebab).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Tropidophorus laticutatus</i>	จิ้งเหลนหัวกุ้ว Phu Wua Depressed - bodied Skink	Inhabits hill streams, found in crevices along stream bank.	Nong Khai (Phu Wua).
<i>Tropidophorus matsui</i>	จิ้งเหลนหัวกุ่มาน้ำทิพย์ Matsui's Depressed- bodied Skink	Inhabits hill streams, found in crevices along stream bank.	Roi Et (Phu Pa Namtip).
<i>Ramphotyphlops lineatus</i>	งูดินลายขีด Striped Blind Snake	Inhabits evergreen forest.	Pattani (Napradoo); Narathiwat (Muang).
<i>Typhlops roxanae</i>	งูดินกรุงเทพฯ Roxan's Blind Snake	Unknown.	Bangkok.
<i>Typhlops siamensis</i>	งูดินสยาม Siamese Blind Snake	Unknown.	The type specimens was collected by Mouhot in Siam, the exact locality is unknown.
<i>Typhlops trangensis</i>	งูดินตรัง Trang Blind Snake	Inhabits evergreen forest, found under rotten logs.	Trang (Khao Chong).
<i>Ahaetulla fasciolata</i>	งูเขี้ยวหัวลายกระ Speckle-headed Whip Snake	Inhabits evergreen forest.	Nakhon Si Thammarat (Thung Song).
<i>Ahaetulla mycterizans</i>	งูเขี้ยวจิ้งจกมลายู	Inhabits evergreen forest.	Chumphon (Muang, Tasan); Krabi (Khao Pra Bang Kram);



Scientific Name	Thai name/Common name	Habitat	Site
	Malayan Green Whip Snake		Trang.
<i>Calamaria lumbricoidea</i>	งูพองออหลากลาย Variable Reed Snake	Inhabits lowland evergreen forest.	Pattani (Napradoo); Narathiwat (Bang Nara, Waeng).
<i>Calamaria schlegelii</i>	งูพองออหัวแดง Red-headed Reed Snake	Inhabits lowland evergreen forest.	Pattani (Napradoo).
<i>Dendrelaphis ngansaoensis</i>	งูสายमानอันนัม Annamese Bronzeback	Inhabits evergreen forest.	Chanthaburi (Khao Sebab) (Chan-ard, Chauynkern and Sukprakarn, per. observ.)
<i>Dendrelaphis striatus</i>	งูสายमानลายเฉียง Oblique-striped Bronzeback	Inhabits evergreen forest.	Phangnga; Nakhon Si Thammarat (Thung Song).
<i>Dinodon septentrionale</i>	งูปลองฉนวนภูเขา Hill Wolf Snake	Inhabits hill forest.	Chiang Mai (Doi Inthanon, Doi Suthep-Pui).
<i>Gongylosoma baliodeirus baliodeirus</i>	งูสายทองลายแถบ Striped Ringneck	Inhabits lowland evergreen forest.	Narathiwat (Sungi Waeng).
<i>Gongylosoma baliodeirus cochrae</i>	งูสายทองจันทบุรี Cochran's Ringneck	Inhabits lowland evergreen forest.	Chanthaburi (Khao Soi Dao).
<i>Gongylosoma longicauda</i>	งูสายทองหางยาว Long-tailed Ringneck	Inhabits lowland evergreen forest.	Narathiwat (Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gonyosoma floweri</i>	งูกาบหมากมลายู Flower's Racer	Unknown .	Trang.
<i>Liopeltis tricolor</i>	งูสายทองมลายู Malayan Ringneck	Inhabits lowland evergreen forest.	South (unknown).
<i>Lepturophis albofuscus</i>	งูปล่องฉนวนมลายู Slender Wolf Snake	Inhabits lowland evergreen forest.	Nakhon Si Thammarat (Thung Song).
<i>Lycodon butleri</i>	งูปล่องฉนวนทุ่งสง Butler's Wolf Snake	Inhabits lowland evergreen forest.	Nakhon Si Thammarat (Thung Song).
<i>Macrocalamus lateralis</i>	งูพงออกภูเขา Mountain Reed Snake	Inhabits lowland evergreen forest.	Narathiwat (Waeng).
<i>Oligodon dorsolateralis</i>	งูบี๊แกวลายซิด Dorsolateral-lined Kukri Snake	This species has been found once previously in the vicinity of a hospital in Chiang Mai (Taylor, 1965).	Chiang Mai.
<i>Oligodon mouhoti</i>	งูอดเขมร Mouhot's Kukri Snake	Inhabit evergreen forest, including beach forest.	Phetchaburi (Cha-Um); Songkhla (Hat Yai).
<i>Stegonotus borneensis</i>	งูแดงบอร์เนียว Malayan Black Snake	Unknown.	South (Unknown).
<i>Bitia hydroides</i>	งูปากกว้างทองสัน Keel-bellied Water Snake	Inhabits mud flat and mangrove area.	Bangkok.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cantoria violacea</i>	งูปากกว้างลาย Cantor's Water Snake	Inhabits mud flat and mangrove area.	Phuket (Saphan Hin).
<i>Enhydria jagorii</i>	งูสายรุ้งดำ Black Water Snake	Inhabits canals, rivers, swamps, ponds and lakes including reservoirs.	Nakhon Sawan (Bung Borapet); Bangkok.
<i>Fordonia leucobalia</i>	งูปลาหัวเทา Crab-eating Water Snake	Inhabits mud flat and mangrove area.	Phuket (Saphan Hin); Satun (Tammalang).
<i>Gerarda prevostiana</i>	งูปลาตาแมว Cat-eyed Water Snake	Inhabits coastal area.	Chon Buri (Ang Sila).
<i>Homalopsis nigroventralis</i>	งูหัวกะโหลกท้องดำ Black-bellied Water Snake	Inhabits forest streams.	Sakon Nakhon (Phu Phan).
<i>Amphiesma bitaeniatum</i>	งูลายสาบยูนนาน Yunnan Keelback	Inhabits submontane forest, found near hill stream.	Chiang Mai (Doi Inthanon).
<i>Amphiesma deschaunseei</i>	งูลายสาบทองสามขีด Northern Keelback	Inhabits hill streams in evergreen forest.	Chiang Mai (Doi Chiang Dao); Loei (Phu Luang); Uthai Thani (Hau Khakhaeng).
<i>Amphiesma groundwateri</i>	งูลายสาบท่าสาร Groundwater's Keelback	Inhabits hill streams in evergreen forest.	Chumphon (Tasan).
<i>Amphiesma inas</i>	งูลายสาบมลายู Malayan Mountain Keelback	Inhabits hill streams in evergreen forest.	Prachuap Khirikhan (Hau Tam Pra); Nakhon Si Thammarat (Khao Luang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Amphiesma khasiense</i>	งูลายสาบเขาสูง Khasi Mountain Keelback	Inhabits hill streams in evergreen forest.	Chumphon (Tasan).
<i>Macropisthodon flaviceps</i>	งูริ้งแห้วแดง Red-headed Keelback	Inhabits hill streams in evergreen forest.	Surat Thani (Ban Don).
<i>Macropisthodon rhodomelas</i>	งูริ้งแห้วดำ Blue-necked Keelback	Inhabits hill streams in evergreen forest.	Nakhon Si Thammarat; Trang (Khao Chong); Satun (Thale Ban); Narathiwat (Waeng).
<i>Rhabdophis subminiatus helleri</i>	งูลายสาบชุนนาน Heller's Keelback	Unknown.	Chiang Mai.
<i>Pareas hamptoni</i>	งูกินทากลายขี้ Hampton's Slug Snake	Inhabits lowland evergreen forest.	Narathiwat (Bangnara).
<i>Xenodermus javanicus</i>	งูทองขาว Javan White-bellied Snake	Inhabits forest stream, found under log in stream.	Pattani (Sai Khao Falls).
<i>Calliophis gracilis</i>	งูปล้องหวายมลายู Grey Coral snake	Unknown.	Pattani (Napradoo).
<i>Laticauda colubrina</i>	งูส้มิงทะเลเหลือง Yellow-lipped Sea Krait	Inhabits coastal areas.	Gulf of Thailand and Andaman Sea.
<i>Laticauda laticaudata</i>	งูส้มิงทะเลปากดำ Black Banded Sea Krait	Inhabits coastal areas.	Gulf of Thailand and Andaman Sea.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Aipysurus eydouxii</i>	งูชายฆนวนล White-spotted Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand Waters.
<i>Acalyptophis peronii</i>	งูทากลายทองขาว Peron's Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand Waters.
<i>Astrotia stokesii</i>	งูทากลาย Stoke's Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand Waters; Chon Buri (Ang Sila); Pattani.
<i>Enhydrina schistosa</i>	งูคออ่อนปากจะงอย Beaked Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand and Andaman Sea.
<i>Hydrophis atriceps</i>	งูแสมรังหัวดำ Black-headed Sea Snake	Inhabits sea and ocean.	Gulf of Thailand Sea.
<i>Hydrophis belcheri</i>	งูแสมรังเกี๋ยดเบลเชอร์ Belcher's Sea Snake	Inhabits sea and ocean.	Gulf of Thailand Sea.
<i>Hydrophis bituberculatus</i>	งูแสมรังเกี๋ยดสองตุ่ม Bitubercled Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Andaman Sea.
<i>Hydrophis caeruleascens</i>	งูแสมรังลายเขือง Dark Blue-banded Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand and Andaman Sea.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Hydrophis cantoris</i>	งูแสมรังแคนเทอร์ Gunther's Sea Snake	Inhabits mouth of river.	Gulf of Thailand Waters: Chanthaburi Estaurine and Koh Chang.
<i>Hydrophis cyanocinctus</i>	งูแสมรังเหลืองลายคราม Blue-banded Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand and Andaman Sea: Songkhla, Pattani and Phuket.
<i>Hydrophis fasciatus</i>	งูแสมรังหัวดำ Striped Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand and Andaman Sea.
<i>Hydrophis gracilis</i>	งูคออ่อนหัวเข็ม Graceful Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand and Andaman Sea: Phuket.
<i>Hydrophis klossi</i>	งูฝักมะรุม Kloss's Sea Snake	Inhabits sea. It is also found in coral reefs.	Gulf of Thailand Waters.
<i>Hydrophis lamberti</i>	งูแสมรังแลมเบิร์ต Lambert's Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Gulf of Thailand Waters.
<i>Hydrophis lapemoides</i>	งูแสมรังอ่าวเปอร์เซีย Persian Gulf Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Andaman Sea: Phuket.
<i>Hydrophis melanosoma</i>	งูแสมรังดำ Black Sea Snake	Inhabits sea and ocean.	Gulf of Thailand Waters.
<i>Hydrophis ornatus ornatus</i>	งูแสมรังหางขาว Ornate Sea Snake	Inhabits sea and ocean. It is also found in coral reefs.	Andaman Sea: Phuket.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Hydrophis spiralis</i>	งูแสมรังลายเหลือง Yellow Sea Snake	Inhabits sea and ocean.	Andaman Sea: Phuket.
<i>Hydrophis torquatus aagaardi</i>	งูแสมรังมลายู Black-headed Sea Snake	Inhabits mouth of rivers.	Gulf of Thailand Waters: Songkhla; Narathiwat (Bangnara estuarine).
<i>Hydrophis torquatus diadema</i>	งูแสมรังเทา Grey Sea Snake	Inhabits mouth of rivers.	Gulf of Thailand Waters: Chanthaburi (estuarine); Trat (Koh Chang).
<i>Kerilia jerdonii jerdonii</i>	งูชายธงขาวทลายมตัด Jerdon's Sea Snake	Inhabits sea and ocean.	Andaman Waters: Phuket. Gulf of Thailand Waters: Songkhla; Pattani.
<i>Kerilia jerdonii siamensis</i>	งูชายธงสยาม Siamese Sea Snake	Inhabits sea and ocean.	Gulf of Thailand Waters: Songkhla; Pattani.
<i>Kolpophis annandalei</i>	งูชายธงหัวโต Annandale's Sea Snake	Inhabits lagoon and mouth of rivers.	Gulf of Thailand Waters: Songkhla; Pattani.
<i>Lapemis curtus</i>	งูอายจิ้ง Hardwicke's Sea Snake	Inhabits sea and ocean.	Gulf of Thailand and Andaman Sea.
<i>Thalassophina viperina</i>	งูชายธงทองขาว Viperine Sea Snake	Inhabits sea and ocean.	Gulf of Thailand and Andaman Sea: Phuket.
<i>Thalassophis anomalus</i>	งูเสมีขนรังหัวสั้น Anomalous Sea Snake	Inhabits coastal areas.	Gulf of Thailand Waters.

## Status Endemic

Scientific Name	Thai name/Common name	Habitat	Site
<i>Cnemaspis chanthaburiensis</i>	จิ้งจกนิ้วยาวจันทบุรี Dring's Rock Gecko	Found in tree buttress and under log on forest floor.	Chanthaburi (Khao Sebab, Khao Soi Dao).
<i>Cnemaspis phuketensis</i>	จิ้งจกนิ้วยาวภูเก็ต Phuket Rock Gecko	Unknown.	Phuket.
<i>Cyrtodactylus angularis</i>	ตุ๊กแกป่าดงพญาเย็น Angular-spotted Gecko	Inhabits deciduous and dry evergreen forest, usually found at night on the rock of forest floor.	Nakhon Ratchasima (Lad Bua Khao, Pak Chong Sakaerat); Saraburi (Hin Lab, Muak Lek, Pha Sadet).
<i>Cyrtodactylus brevipalmatus</i>	ตุ๊กแกป่าโคนนิ้วติด Short-webbed Gecko	Inhabits evergreen forest, usually found at night on tree trunks.	Nakhon Si Thammarat (Khao Luang).
<i>Cyrtodactylus chanhomeae</i>	ตุ๊กแกป่าสระบุรี Chanhome's Gecko	Inhabits limestone caves.	Saraburi.
<i>Cyrtodactylus peguensis zebraicus</i>	ตุ๊กแกป่าลายจุดไต้ Zebra-spotted Gecko	Inhabits evergreen forest, usually found at night on tree buttress or forest floor.	Krabi (Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong).
<i>Cyrtodactylus sumonthai</i>	ตุ๊กแกป่าเขาชะเมา Sumontha's Gecko	Inhabits caves.	Rayong (Khao Cha Mao).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Cyrtodactylus thirakhupti</i>	ตุ๊กแกป่าท่าซธร, ตุ๊กแกป่าสุราษฎร์ Thirakupt's Gecko	Inhabits caves.	Surat Thani.
<i>Cyrtodactylus tigroides</i>	ตุ๊กแกป่าไทรโยค Sai Yok Gecko	Inhabits evergreen forest and nearby caves.	Tak (Umphang); Kanchanaburi (Sai Yok).
<i>Dixoneus hangseesom</i>	จิ้งจกดินหางสีส้ม Orange-tailed Ground Gecko	Inhabits bamboo forest, usually found at night on forest floor.	Kanchanaburi (Sai Yok).
<i>Dixoneus melanostictus</i>	จิ้งจกดินข้างดำ Dark-sided Ground Gecko	Inhabits evergreen forest, usually found at night on forest ground.	Nakhon Ratchasima (Pak Chong); Saraburi (Muak Lek).
<i>Gehyra angusticaudata</i>	จิ้งจกหินหางเรียว Slender-tailed Four-clawed Gecko	This species was reported from old nipa-hut that was being turned down.	Chon Buri (Sriracha).
<i>Gehyra fehlmanni</i>	จิ้งจกหินลาขกระ Fehlman's Four-clawed Gecko	This species inhabits disturbed and primary forest, usually found on forest ground and tree buttress.	Nakhon Ratchasima (Khao Yai, Pak Chong); Kanchanaburi; Nakhon Si Thammarat (Khao Luang, Khao Ron Pibun).
<i>Gehyra lacerata</i>	จิ้งจกหินเมืองกาญจน์ Western Four-clawed Gecko	Known from beach forest and deciduous forest, usually found under rock on forest floor.	Nakhon Ratchasima (Sakaerat); Srakaew (Ko Klan, Pang Sida); Chon Buri (Angsila); Kanchaburi; Phetchaburi (Cha-Um).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gekko siamensis</i>	ตุ๊กแกสยาม Siamese Gecko	Found in rock crevices and cave in degraded forest.	Nakhon Ratchasima (Pak Chong); Phetchabun (Nam Nao).
<i>Isopachys anguinoides</i>	จิ้งเหลนคางประจวบ Striped Legless Skink	Beach and evergreen forest, usually found under humus soil.	Phetchaburi (Cha-Um, Kaengkrachan); Prachuap Khirikhan (Bang Saphan, Hua Hin, Pa-La-U, Pa Tiu, Tap Sakae, Wanakorn Beach).
<i>Isopachys gyldenstolpei</i>	จิ้งเหลนคางลาย Gyldenstolpe's Legless Skink	Inhabits beach and evergreen forest, usually found under humus soil.	Uthai Thani (Ban Rai), Kanchanaburi, Phetchaburi (Cha-Um); Prachuap Khirikhan (Hua Hin, Koh Lak, Muang).
<i>Isopachys roulei</i>	จิ้งเหลนคางชลบุรี Roule's Legless Skink	Inhabits beach forest, usually found under humus soil.	Chon Buri (Ang Sila).
<i>Leptoceps osellai</i>	จิ้งเหลนภูเขาสามันัว Osella's Larut Skink	Inhabits evergreen forest, usually found under humus soil and rotten logs.	Chiang Mai (Mae Kung); Phetchabun (Nam Nao).
<i>Lygosoma haroldyoungi</i>	จิ้งเหลนเรียวลาย Banded Slender Skink	Inhabits evergreen forest, usually found in termite mound or under humus soil.	Chiang Mai (Doi Suthep-Pui, Fang); Chiang Rai; Phitsanulok (Thung Salaeng Luang); Loei (Phu Luang); Nong Khai (Bunh Kan, Rattana Wapi); Phetchabun (Nam Nao); Chaiyaphum (Phu Kieo).
<i>Lygosoma koratense</i>	จิ้งเหลนเรียวโคราช Khorat Slender Skink	Inhabits evergreen forest.	Saraburi (Muak Lek); Nakhon Ratchasima (Dong Lek, Lad Bua Khao, Non Thai).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Mabuya macularia malcolmi</i>	จิ้งเหลนหลากลายมัลคอล์ม Malcolm's Skink	Inhabits deciduous and evergreen forest.	Chiang Mai (Doi Suthep-Pui); Loei (Phu Pak Kinak); Uthai Thani (Hau Khakhaeng); Ratchaburi (Maenam Pachi); Phetchaburi (Kaengkrachan); Phatthalung.
<i>Mabuya macularia postnasalis</i>	จิ้งเหลนหลากลายดานซาย Dansai Skink	Inhabits evergreen forest.	Loei (Phu Lom Lo, Dan Sai).
<i>Mabuya macularia quadrifasciata</i>	จิ้งเหลนหลากลายสี่ขีด Four-lined Skink	Inhabits deciduous and evergreen forest.	Pattani (Bukit Besar).
<i>Riopa herberti</i>	จิ้งเหลนเรียวไต Herbert's Slender Skink	Inhabits evergreen forest, usually found under rotten logs on forest floor.	Chiang Mai (Doi Inthanon); Phrae (Mae Yom); Nong Khai (Phu Wua); Roi Et (Phu Pa Namtip); Ubon Ratchathani (Yod Dome); Nakhon Ratchasima (Sakaerat); Srakaew (Koklan, Pang Sida); Chanthaburi (Khao Kittchakud, Khao Sebab, Khao Soi Dao); Pathum Thani (Rang Sit), Bangkok (Pathumwan); Kanchanaburi (Sai Yok); Phetchaburi (Cha-Um, Kaengkrachan); Prachuap Khirkhan (Pa-La-U, Tap Sakae); Surat Thani (Koh Samui); Phuket; Trang (Had Chao Mi, Khao Chong, Koh Libong).
<i>Scincella melanosticta kohtaoensis</i>	จิ้งเหลนดินเกาะเต่า Koh Tao Slender Skink	Inhabits evergreen forest, found under rotten logs and humus soil.	Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Loei (Phu Luang).
<i>Scincella tavesae</i>	จิ้งเหลนดินเมืองกาญจน์	Inhabits deciduous and evergreen forest,	Surat Thani (Koh Tao).

Scientific Name	Thai name/Common name	Habitat	Site
	Taves's Slender Skink	found under rotten logs and in humus soil of tree buttress.	
<i>Sphenomorphus grandisonae</i>	จิ้งเหลนภูเขาแกรนิตสีน้ำตาล	Unknown.	Northern (Ban Tong Pheung).
	Grandison's Forest Skink		
<i>Sphenomorphus helenae</i>	จิ้งเหลนภูเขาเฮเลน	Unknown.	Nonthaburi.
	Helen's Forest Skink		
<i>Sphenomorphus lineopunctulatus</i>	จิ้งเหลนภูเขาอุบล	Unknown.	Ubon Ratchathani (Sanoi River).
	Ubon Forest Skink		
<i>Sphenomorphus mimicus</i>	จิ้งเหลนภูเขาเล็ก	Inhabits hill forest.	Loei (Phu Luang); Nakhon Ratchasima (Pak Chong).
	Minute Forest Skink		
<i>Sphenomorphus tersus</i>	จิ้งเหลนภูเขาสีจาง	Inhabits hill forest.	Phetchaburi (Kaengkrachan); Chumphon (Tasan); Ranong (Pak Chan Estuarine); Phangnga (Khao Lak); Nakhon Si Thammarat (Khao Luang, Khao Wang Hip).
	Pale Forest Skink		
<i>Tropidophorus latiscutatus</i>	จิ้งเหลนห้วยภูว้า	Inhabits hill streams, found in crevices along stream bank.	Nong Khai (Phu Wua).
	Phu Wua Depressed-bodied Skink		
<i>Tropidophorus matsuii</i>	จิ้งเหลนห้วยภูพาน้ำทิพย์	Inhabits hill streams, found in crevices along stream bank.	Roi Et (Phu Pa Namtip).
	Matsui's Depressed-bodied Skink		

Scientific Name	Thai name/Common name	Habitat	Site
<i>Tropidophorus thai</i>	จิ้งเหลนห้วยไทย Thai Stream Skink	Inhabits hill streams in evergreen forest.	Mae Hong Son; Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui, Mae Wang, Pa Miang).
<i>Typhlops floweri</i>	งูดินหัวเหลือง Flower's Blind Snake	Inhabits evergreen forest, including parks and gardens.	Bangkok, Pathum Thani (Rangsit).
<i>Typhlops khoratensis</i>	งูดินโคราช Khorat Blind Snake	Inhabits evergreen forest, usually found under rotten logs.	Chiang Mai; Nakhon Ratchasima (Sakaerat); Saraburi (Muak Lek); Prachuap Khirikhan (Hua Hin).
<i>Typhlops ozakiae</i>	งูดินปักธงชัย Ozakia's Blind Snake	Inhabits evergreen forest, usually found under rotten logs.	Nakhon Ratchasima (Sakaerat).
<i>Typhlops roxanae</i>	งูดินกรุงเทพฯ Roxan's Blind Snake	Unknown.	Bangkok.
<i>Typhlops siamensis</i>	งูดินสยาม Siamese Blind Snake	Unknown.	The type specimens was collected by Mouhot in Siam, the exact locality is unknown.
<i>Typhlops trangensis</i>	งูดินตรัง Trang Blind Snake	Inhabits evergreen forest, found under rotten logs.	Trang (Khao Chong).
<i>Boiga sangsomi</i>	งูเขียวดงลาย Banded Cat Snake	Inhabits evergreen forest, found on palm tree.	Krabi (Ban Khanom); Nakhon Si Thammarat (Ron Pibun).
<i>Dryocalamus thungsongensis</i>	งูปล้องฉนวนทุ่งสง Thungsong Bridle Snake	Inhabits lowland evergreen forest.	Nakhon Si Thammarat (Khao Luang, Thung Song).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Gongylosoma baliodeirus cochranae</i>	งูสายทองจันทบุรี Cochran's Ringneck	Inhabits lowland evergreen forest.	Chanthaburi (Khao Soi Dao).
<i>Oreophis porphyraceus coxi</i>	งูทางมะพร้าวแดงภูหลวง Lox's Mountain Racer	Inhabits evergreen forest. In Loei, it was observed while consuming a mole.	Loei (Phu Luang); Chaiyaphum (Phu Kieo).
<i>Sibynophis triangularis</i>	งูคอขี้วันหัวลายสามเหลี่ยม Triangle Blackhead	Inhabits evergreen forest.	Nakhon Ratchasima (Pak Chong, Wang Namkiao); Chachoengsao (Khao Ang Rue Ni); Trat (Koh Chang); Chon Buri (Siracha).
<i>Amphiesma deschaunseei</i>	งูลายสาบทองสามขีด Deschaunsee's Keelback	Inhabits hill streams in evergreen forest.	Chiang Mai (Doi Chiang Dao); Loei (Phu Luang); Uthai Thani (Hau Khakhaeng).
<i>Amphiesma groundwateri</i>	งูลายสาบทาสาร Groundwater's Keelback	Inhabits hill streams in evergreen forest.	Chumphon (Tasan).
<i>Trimeresurus (Cryptelytrops) kanburiensis</i>	งูหางหมัดกาญจนบุรี Kanburi Pit Viper	Inhabits deciduous and evergreen forest. This species was rediscovered by Warrell <i>et al.</i> (1992) from the bamboo forest at Nong Bu Wa after the first discovery 50 years.	Kanchanaburi (Sai Yok).



# A MPHIBIANS

EX = Extinct  
EW = Extinct in the Wild  
CR = Critically Endangered  
EN = Endangered  
VU = Vulnerable  
NT = Near Threatened  
LC = Least Concern  
DD = Data Deficient  
Endemic

WS = Wildlife Sanctuary  
NH = Non-hunting Area  
NP = National Park



**Class:** Amphibia

**Order:** Anura

**Family:** Ranidae

**Scientific Name:** *Paa bourreti*

**Common Name:** กบอกหนามเหนือ

Bourret's Spiny-breasted Frog



### Habitat:

Inhabits mountain streams in high altitude forest.

### Locality:

Mae Hong Son (Pa Sue Falls); Chiang Mai (Doi Pahom Pok).

### Description:

Size (snout to vent) 91 mm.

Back with short, thick ridges interspersed with circular warts. Yellowish brown above with dark flecks.



*Paa fasciculispina*

**Class:** Amphibia

**Order:** Anura

**Family:** Ranidae

**Scientific Name:** *Paa fasciculispina*

**Common Name:** กบอกหนามจันทบุรี

Chanthaburi Spiny-breasted Frog

**Habitat:**

Inhabits mountain streams in high altitude forest.

**Locality:**

Chanthaburi (Khao Kitchakud, Khao Sebab, Khao Soi Dao).

**Description:**

Size (snout to vent) 106 mm.

Back with numerous short, thick ridges, none as long as diameter of eye, interspersed with circular warts; webbing complete, not emarginate; tips of toes swollen to twice width of phalanges; proximal subarticular tubercles of fingers twice as long as distal ones; male with ventral spines in groups of 5-10 clusters distributed across chest and throat. Brown with obscure darker spots on back.



**Class:** Amphibia  
**Order:** Anura  
**Family:** Rhacophoridae  
**Scientific Name:** *Theلودerma gordoni*  
**Common Name:** ปาดตะปุ่มใหญ่  
 Large Warted Tree Frog

**Habitat:**

Inhabits submontane forest.

**Locality:**

Chiang Mai (Doi Suthep-Pui); Loei (Phu Luang).

**Description:**

Size (snout to vent) 48 mm.

Dorsal and lateral parts of body with very numerous large warts covered with small pearly granular asperities; fingers entirely free, toes little less than two-thirds webbed; tips of all digits dilated but all considerably smaller than tympanum. Dark wood brown above with darker spots on head; granules on tubercles light grey or cream; an irregular darker area in groin; venter lavender with dark lavender flecks and reticulations.



**Class:** Amphibia  
**Order:** Anura  
**Family:** Rhacophoridae  
**Scientific Name:** *Theلودerma horridum*  
**Common Name:** ปาดตะปุ่มมลายู  
Thorny Warted Tree Frog



**Habitat:**

Inhabits lowland evergreen forest.

**Locality:**

Nakhon Si Thammarat (Khao Luang); Pattani (Kok Po); Narathiwat (Waeng).

**Description:**

Size (snout to vent) 40 mm.

Prominent irregular warts studded with granular asperities; fingers half webbed; toes fully webbed. Dark brown above with rather indistinct blackish spots on the body and regular cross-bars on limbs; granular asperities greyish; a large black lumbar spot; Lower surface blue-grey, largely spotted and marbled with black.



**Class:** Amphibia

**Order:** Anura

**Family:** Rhacophoridae

**Scientific Name:** *Theلودerma stellatum*

**Common Name:** ปาดตะปุ่มจันทบุรี

Taylor's Warted Tree Frog



### Habitat:

Inhabits lowland forest.

### Locality:

Nakhon Ratchasima (Sakaerat); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Kitchakud, Khao Sebab).

### Description:

Size (snout to vent) 34 mm.

Body with very numerous tubercles covered with granular asperities; fingers about one-third webbed, toes about four-fifths webbed; digits with large terminal discs, those on fingers as large as tympanum. Brown or lavender gray above with cream speckles; black spots on snout and one or two between eyes; atrifoliate spot across shoulders, followed by 4-5 smaller dark spots; regular cross bars on limbs; lower surface cream with purplish-brown flecks or spots, partially reticulated.

# Status NT

Scientific Name	Thai name/Common name	Habitat	Site
<i>Tylototriton verrucosus</i>	กะท่าง Boulenger's Tree Frog	Breeds in pools in mountain forest . streams; terrestrial phase in montane forest. and possibly scrub. Occurred above 1,000 m.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui, Doi Ang Khang); Mae Hong Son (Nam Tok Mae Surin NP); Nan (Doi Phu Ka); Loei (Phu Luang WS).
<i>Brachytarsophrys carinense</i>	อึ่งกรายขางแถบ Burmese Horned Frog	Inhabits evergreen forests. Male was found guarding the fertilized eggs which attached the crevice roof under stream current (Stuart, per.com.).	Chiang Mai (Doi Chiang Dao, Doi Suthep-Pui); Lampang (Mae Toh); Kamphaeng Phet (Khlong Lan NP); Uthai Thani (Hau Khakhaeng WS); Phetchaburi (Kaengkrachan NP); Surat Thani (Khlong Saeng WS) and Phangnga (Khao Lak-Lam Ru NP).
<i>Megophrys nasuta</i>	อึ่งกรายหัวแหลม Long-nosed Horned Frog	Inhabits lowland moist evergreen forest.	Songkhla (Ton Ngachang); Yala (Betong, Bannangstar); Narathiwat (Tanjong Mas, Waeng).
<i>Ophryophryne microstoma</i>	อึ่งกรายปากแคบ Narrow-mouthed Horned Frog	The specimens from Thailand have been reported from the high elevation of mountain forest.	Nan (Doi Phu Kha).
<i>Xenophrys aceras</i>	อึ่งกรายหัวมน Malayan Horned Frog	Inhabits lowland moist evergreen forest.	Surat Thani (Khlong Saeng, Khao Sok); Phangnga; Nakhon Si Thammarat (Khao Luang, Thung Song); Trang (Khao Bantat, Khao Chong); Pattani (Nam Tok Sai Khao); Yala (Betong, Bannangstar); Narathiwat (Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Xenophrys longipes</i>	อึ่งกรายขายาว Long-legged Horned Frog	Inhabits evergreen forest in high altitude.	Ranong (Khlung Naka); Phangnga; Nakhon Si Thammarat (Khao Luang); Yala (Betong, Bannangstar); Narathiwat (Waeng).
<i>Xenophrys major</i>	อึ่งกรายห้วยใหญ่ Greater Stream Horned Frog	Inhabits evergreen forest.	Mae Hong Son (Muang); Chiang Mai (Doi Inthanon, Doi Suthep-Pui, Tha Ton, Mae Ai); Loei (Phu Luang); Tak (Umpang); Kanchanaburi (Tan Lot Cave, Thong Pa Phum); Phetchaburi (Kaengkrachan).
<i>Xenophrys minor</i>	อึ่งกรายห้วยแคระ Dwarf Horned Frog	Inhabits high altitude evergreen forest.	Chiang Mai (Doi Inthanon).
<i>Xenophrys parva</i>	อึ่งกรายเล็ก Lesser Stream Horned Frog	Inhabits evergreen forest.	Mae Hong Son (Pha Sue Falls); Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Uthai Thani (Hau Khakhaeng); Phuket.
<i>Ansonia inthanon</i>	คางคกห้วยอินทนนท์ Inthanon Stream Toad	The endemic species inhabits in high altitude of mountain forest, but the population in Kanchanaburi is probably swampy area in lowland evergreen forest.	Chiang Mai (Doi Inthanon); Kanchanaburi (Thong Pa Phum).
<i>Ansonia siamensis</i>	คางคกห้วยไทย Siamese Stream Toad	Inhabits near streams of evergreen forest.	Trang (Khao Chong).
<i>Glyphoglossus molossus</i>	อึ่งเพา, อึ่งปากขวด	Inhabits deciduous and secondary forests.	Sakon Nakhon (Phu Phan); Nong Bua Lumpoo (Muang, Si

Scientific Name	Thai name/Common name	Habitat	Site
	Truncat-snouted Spadefoot Frog		Boonruang, Na Klang, Noan Sang); Nakhon Ratchasima (Wang Nam Kieo); Srakaew (Ko Klan, Tapraya); Kamphaeng Phet; Uthai Thani (Hau Khakhaeng); Phra Nakhon Si Ayutthaya; Chon Buri, Chanthaburi; Ratchaburi.
<i>Kaloula mediolineata</i>	อึ่งอ่างก้นขีด Mediam-striped Bullfrog	Inhabits deciduous and secondary forests. Usually found in mass breeding in the same areas with other edible species.	Known from Thailand and Laos.
<i>Bourretia kohchangae</i>	กบเกาะช้าง Koh Chang Frog	Inhabits evergreen forest along mountain streams.	Trat (Koh Chang).
<i>Ingerana tasanae</i>	กบท่าสาร Tasan Frog	Inhabits mountain streams in evergreen forest.	Chumphon (Tasan); Ranong (Khlong Naka); Surat Thani (Khao Sok, Khlong Saeng); Phangnga (Khao Lak Lam Ru).
<i>Ingerana tenasserimensis</i>	กบตะนาวศรี Tenasserim Frog	Inhabits the forest floor in vicinity of mountain streams.	Kanchanaburi (Pilok, Sai Yok, Thung Yai); Ratchaburi (Mae Nam Pachi); Nakhon Sithammarat (Khao Ram, Khao Ron Pibun); Trang (Khao Chong).
<i>Limnnectes blythii</i>	กบพุด, เขียดแฉว Blyth's Frog	Inhabits mountain streams in evergreen forest.	Mae Hong Son (Salawin); Tak (Thung Yai); Kamphaeng Phet (Khlong Lan); Uthai Thani (Hau Khakhaeng); Kachanaburi (Erawan, Lin Tin, Sangkhlaburi, Sai Yok, Thong Pa Phum); Ratchaburi (Mae Nam Pa Chi); Phetchaburi (Kaengkrachan);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Limnectes kuhlii</i>	กบหัวขานุ่ม, กบมีน Kuhl's Stream Frog	Inhabits mountain streams in evergreen forest.	Prachuap Khirikhan (Pa-La-U); Ranong (Khlung Naka, Muang); Surat Thani (Koh Samui, Khao Sok, Khlung Saeng); Phangnga (Khao Lak Lum Ru); Krabi (Khao Panom Bencha, Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong, Koh Libong); Songkhla (Ton Nga Chang); Yala (Banangstar, Betong); Narathiwat (Waeng). Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Pa Hom Pok, Doi Suthep-Pui, Mae Ai, Tha Ton); Pa Yao (Doi Pa Chang); Phrae; Tak (Thung Yai); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Pi Lok, Sai Yok, Sangkhlaburi, Tham Tan Lot); Phetchaburi (Kaengkrachan); Surat Thani (Khao Sok, Khlung Saeng); Nakhon Si Thammarat (Khao Luang); Yala (Bang Lang, Betong); Narathiwat (Waeng); Loei (Phu Luang, Phu Kradung); Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Kao Yai, Sakaerat).
<i>Limnectes malesianus</i>	กบตุ้มตาชู Peat Swamp Frog	Inhabits small swampy patches in lowland evergreen forest.	Narathiwat (Waeng).
<i>Limnectes paramacrodon</i>	กบป่าพรุ Masked Swamp Frog	Inhabits scattered areas of peat swamp forest.	Narathiwat (Tak Bai and Sungai Kolok).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Rana chapaensis</i>	กบภูเขาหลวง Chapa Frog	Inhabits mountain streams and shallow wetlands in hill evergreen forest.	Loei (Phu Luang).
<i>Rana lateralis</i>	กบหลังไพล, เขียดเหลือง Yellow Frog	Inhabits dry deciduous forest. Breeding in deep rainpools in rainy season.	Chiang Mai (Doi Suthep-Pui, Kaeng Pan Tao); Kamphaeng Phet (Khlung Lan); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Muang); Phitsanulok (Wat Bos); Loei (Dan Sai, Phu Rue); Sakon Nakhon (Phu Phan); Nakhon Ratchasima (Sakaerat); Ubon Ratchathani (Yod Dome); Umnajjareon (Phu Singh); Srakaew (Wattana Nakhon).
<i>Rana leptoglossa</i>	กบสังขละ Cope's Frog	Inhabits evergreen forest along mountain streams.	Kanchanaburi (Sangkhlaburi, Thong Pa Phum); Tak (Thung Yai).
<i>Rana miopus</i>	กบหลังลายเขียว Three Striped Frog	Inhabits lowland evergreen forest.	Phangnga (Khao Lak Lam Ru); Krabi (Ao Luk, Khao Panombencha, Khao Pra Bang Kram); Nakhon Si Thammarat; Songkhla (Ton Ngachang); Yala (Banangstar).
<i>Rana faber</i>	กบเขาสอยดาว Smith's Frog	Inhabits mountain streams in high altitude forest.	Chanthaburi (Khao Soi Dao).
<i>Rana signata</i>	กบหลังจุด Spotted Stream Frog	Mainly found among tangled vegetation and root mass along muddy streams in lowland evergreen forest.	Chumphon (Tasan); Surat Thani (Khlung Saeng); Trang (Khao Bantat); Songkhla (Khao Nam Kang, Ton Ngachang); Yala (Banangstar); Narathiwat (Sukirin, Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rhacophorus feae</i>	ปาดคอดอยอินทนนท์ Fea's Tree Frog	Inhabits submontane forest.	Mae Hong Son (Pa Sue Falls); Chiang Mai (Doi Inthanon); Loei (Phu Luang).
<i>Rhacophorus maximus</i>	ปาดยักษ์ Giant Tree Frog	Inhabits lowland and submontane forest.	Kanchanaburi (Thong Pha Phum); Ratchaburi (Tako Pidtong); Phetchaburi (Kaengkrachan).
<i>Rhacophorus nigropalmatus</i>	ปาดเขียวตีนดำ Wallace's Tree Frog	Inhabits primary evergreen forest.	Ranong (Ka Pur), Pattani (Kok Po); Yala (Banangstar); Narathiwat (Waeng).
<i>Rhacophorus pardalis</i>	ปาดตีนแดง Harlequin Tree Frog	Inhabits lowland evergreen forest.	Narathiwat (Waeng).
<i>Rhacophorus prominanus</i>	ปาดเขียวตีนแดง Malayan Tree Frog	Inhabits lowland evergreen forest.	Narathiwat (Waeng); Yala (Bannangstar).
<i>Rhacophorus reinwardtii</i>	ปาดเขียวตีนลาย Reinwardt's Tree Frog	Inhabits lowland and submontane forests.	Chiang Mai (Doi Chiang Dao); Kanchanaburi (Thung Yai); Phetchaburi (Kaengkrachan); Narathiwat (Waeng).
<i>Rhacophorus robinsoni</i>	ปาดขาลาย Robinson's Tree Frog	Inhabits closed canopy forest.	Pattani (Bakit Besar), reported since 1903.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Leptobrachium hendricksoni</i>	อึ่งกรายลายจุด Spotted Litter Frog	Inhabits lowland evergreen forests, degraded forest and plantations adjoining forest.	Trang (Khao Bantat WS); Songkhla (Ton Ngachang WS); Yala and Narathiwat (Hala-Bala WS).
<i>Leptobrachium smithi</i>	อึ่งกรายลายเลอะ Dark Litter Frog	Evergreen forest as well as moist deciduous forest. Tadpoles inhabit in forest streams with little current.	Mae Hong Son (Lam Nam Pai, Pa Sue Falls); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Saket, Doi Suthep-Pui); Tak (Thung Yai WS); Uthai Thani (Hau Khakhaeng WS); Kanchanaburi (Erawan, Piloc, Sangkhlaburi, Thong Pa Phum); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Hau Yang Falls, Pa-La-U Falls); Loei (Na Haeo, Dan Sai, Phu Luang, Phu Rue, Phu Kradung); Phangnga; Nakhon Si Thammarat (Khao Luang); Trang (Khao Bantat, Khao Chong); Krabi (Khao Panombencha, Khao Pra Bang Kram); Songkhla (Ton Ngachang WS); Yala and Narathiwat (Hala-Bala WS).
<i>Leptolalax heteropus</i>	อึ่งกรายหลากลาย Variable Litter Frog	Inhabits mid-altitude rainforest between 400-1,000 m. elevation.	Narathiwat (Waeng District).
<i>Leptolalax pelodytoides</i>	อึ่งกรายหนังปุ่ม Mud Slender Frog	Streamside evergreen forest in mid altitude.	Mae Hong Son (Muang); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Loei (Wang Sapung, Phu Luang); Chaiyaphum (Phu Kieo); Uthai Thani (Hau Khakhaeng);

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ansonia malayana</i>	คางคกหัวขมลาชู Malayan Stream Toad	Inhabits near streams of evergreen forest.	Kanchanaburi (Erawan); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U Falls); Surat Thani (Khlung Saeng). Chumphon (Tasan); Ranong (La-Un); Phangnga (Khao Lak Lam Ru); Phuket; Nakhon Si Thammarat (Thung Song); Pattani; Satun (Koh Tarutao); Yala (Betong); Narathiwat (Waeng).
<i>Bufo asper</i>	จิ้งโคร่ง Giant Asiatic Toad	Inhabits rain forest streams as well as the caves in vicinity areas.	Mae Hong Son (Pha Sue Falls); Tak (Umpang); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Sai Yok, Salak Pra, Sangkhlaburi, Thong Pha Phum); Ratchaburi (Mae Nam Pa Chi); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Chumphon (Ta San); Ranong (Muang); Surat Thani (Khao Sok, Khlung Saeng); Phangnga (Khao Lak Lam Ru); Nakhon Si Thammarat (Na Bon, Khao Luang); Trang (Khao Bantat, Khao Chong); Krabi (Hat Thai Muang, Khoa Panombencha, Khao Pra Bang Kram); Songkhla (Ton Ngachang); Yala (Banangstar, Betong, Than To); Narathiwat (Waeng). Yala (Banangstar, Betong, Than To); Narathiwat (Waeng).
<i>Bufo divergens</i>	คางคกแคระขมลาชู Malayan Dwarf Toad	Inhabits moist evergreen forest.	Yala (Banangstar, Betong, Than To); Narathiwat (Waeng).
<i>Bufo macrotis</i>	คางคกหัวراب Large-eared Toad	Almost found in deciduous forest.	Chiang Mai (Doi Chiang Dao, Doi Kram, Mong Huad); Phrae (Mae Yom); Prachin Buri (Kabin Buri); Nakhon Nayok (Nong

Scientific Name	Thai name/Common name	Habitat	Site
<i>Bufo melanostictus</i>	คางคกบ้าน Black-spined Toad	Mainly of disturbed areas, uncommon in closed forests.	Ree); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Salak Pra, Sai Yok); Phangnga (Hat Tai Muang); Pattani (Sai Buri). All provinces throughout the country.
<i>Bufo parvus</i>	คางคกแคระอินโดจีน Dwarf Toad	Mainly inhabits in evergreen forests.	Chanthaburi (Khao Kitchakud, Khao Sebab, Khao Soi Dao ); Uthai Thani (Hau Khakakaeng); Kanchanaburi (Huai Mae La Mun, Pilok, Sangkhlaburi); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Hau Yang Falls, Pa-La-U); Chumphon (Pa Tiu); Ranong (Muang, Khlong Naka); Surat Thani (Khao Sok, Khlong Saeng); Nakhon Si Thammarat (Khao Luang); Trang (Khao Bantat, Khao Chong); Songkhla (Ton Ngachang).
<i>Leptophryne borbonica</i>	คางคกขาขาว Long-legged Slender Toad	Inhabits streams in primary evergreen forests.	Trang (Khao Bantat, Khao Chong); Phatthalung (Kuan Nieng); Songkhla (Ton Ngachang); Narathiwat (Waeng).
<i>Pedostibes hosii</i>	คางคกต้นไม้ Brown Tree Toad	Found in moist evergreen forest. Calling males in breeding season are easily seen on the trees along forest streams.	Trang (Khao Chong); Satun (Tha Le Ban); Yala (Bannangstar); Narathiwat (Waeng).
<i>Calluella guttulata</i>	อึ่งลาย Striped Spadefoot Frog	Most populations were observed from secondary and mixed deciduous forests.	Phrae (Mae Yom); all provinces in western, northeast and southeast regions; Surat Thani (Muang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Kalophrynus interlineatus</i>	อึ่งปทุมหลังลาย Striped Sticky Frog	Inhabits montane forest.	Loei (Phu Luang); Chaiyaphum (Phu Kieo); Phitsanulok (Thung Salaeng Luang); Chanthaburi (Khao Kitchakud, Khao Sebab).
<i>Kalophrynus pleurostigma</i>	อึ่งปทุมมลายู Red-sided Sticky Frog	Inhabits lowland forest.	Chumphon (Pa Tiu); Phangnga (Khao Lak Lam Ru).
<i>Kaloula baleata</i>	อึ่งอ่างมลายู Brown Bullfrog	Inhabits evergreen forest.	Ratchaburi (Tako Pid Tong); Surat Thani (Khao Sok, Khlong Saeng); Krabi (Khao Pra Bang Kram, Koh Lan Ta); Trang (Khao Chong); Nakhon Si Thammarat (Khao Luang); Phuket.
<i>Kaloula pulchra</i>	อึ่งอ่างบ้าน Banded Bullfrog	Inhabits all habitats including beach forest and urban areas.	All provinces of the country.
<i>Microhyla berdmorei</i>	อึ่งแม่หนาว Berdmore's Chorus Frog	Inhabits the forest floor closed to the streams. Breeds in side pools of streams.	Mae Hong Son (Pang Tong); Chiang Mai (Mae Rim); Loei (Phu Luang); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Chaiyaphum (Phu Kieo); Nakhon Ratchasima (Khao Yai, Sakaerat); Srakaew (Pang Sida); Chachengsao (Khao Ang Rue Ni); Tak (Thung Yai, Um Pang); Phrae (Mae Yom); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Surat Thani (Khao Sok); Krabi (Khao Panom Bencha); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Songkhla (Ton Ngachang); Yala (Banangstar); Narathiwat (Waeng).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Microhyla butleri</i>	อึ่งลายแตรม Noisy Chorus Frog	Inhabits forest edge to the primary forest.	Chiang Mai (Doi Chiang Dao, Doi Suthep-Pui); Phrae (Mae Yom); Nakhon Ratchasima (Skaerat); Prachin Buri (Pra Chantakhom); Kanchanaburi (Pilok, Sankhlaburi); Surat Thani (Khlong Saeng); Trang (Kan Tang); Satun; Narathiwat (Tak Bai, Waeng).
<i>Microhyla heymonsi</i>	อึ่งข้างดำ Dark-sided Chorus Frog	Inhabits disturbed areas such as gardens, crop plantations, secondary forests and primary forests.	All provinces.
<i>Microhyla ornata</i>	อึ่งน้ำเต้า Ornate Chorus Frog	Inhabits most of disturbed areas, gardens, crop plantations, secondary forests and primary forests.	Mae Hog Son (Muang); Chiang Mai (Doi Chiang Dao); Phrae (Mae Yom); Tak (Um Pang, Thung Yai); Phitsanulok (Wat Bos); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Roi Et (Phu Pha Namthip); Ubon Ratchathani (Yod Dome); Nakhon Ratchasima (Sakaerat); Chachoengsao (Khao Ang Rue Ni); Srakaew (Ko Klan, Pang Si Da, Ta Praya); Chanthaburi (Khao Sebab); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Erawan, Pilok, Thong Pha Phum); Phetchaburi (Kaengkrachan); Surat Thani (Khlong Saeng, Koh Samui); Ranong (Muang); Krabi (Khao Pra Bang Kram); Nakhon Si Thammarat (Muang, Khao Luang); Songkhla (Ton Ngachang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Microhyla pulchra</i>	อึ่งขาคำ Painted Chorus Frog	Inhabits deciduous forest.	Chiang Mai (Doi Chiang Dao); Phrae (Mae Yom); Phitsanulok (Wat Bos); Loei (Phu Rue); Nong Khai (Phu Wua); Kalasin (Phu Si Than); Nakhon Ratchasima (Sakaerat); Chachoengsao (Khao Ang Rue Ni); Srakaew (Ko Klan, Pang Sida, Ta Praya); Chanthaburi (Khao Sebab); Uthai Thani (Hau Khakhaeng); Nakhon Si Thammarat.
<i>Micryletta inornata inornata</i>	อึ่งหลังจุด Inornate Chorus Frog	Inhabits evergreen forest and disturbed Areas.	Chiang Mai (Doi Chiang Dao, Doi Suthep-Pui); Phrae (Mae Yom); Tak (Thung Yai); Loei (Phu Luang); Nong Khai (Phu Wua); Nakhon Ratchasima (Sakaerat); Prachin Buri (Prachantakham); Chachoengsao (Khao Ang Rue Ni); Srakaew (Pang Sida, Ta Praya); Chanthaburi (Khao Sebab, Khao Soi Dao); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Thong Pha Phum).
<i>Micryletta inornata lineata</i>	อึ่งหลังขีด Striped Chorus Frog	Inhabits evergreen forest and disturbed Areas.	Ratchaburi (Chaum Bung); Phetchaburi (Kaengkrachan); Ranong, Surat Thani; Phangnga, Nakhon Si Thammarat; Trang; Pattani; Narathiwat.
<i>Amolops larutensis</i>	กบลายหินเมืองใต้ Larut Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing water forest streams.	Trang (Khao Bantat); Pattani, Yala (Bang Lang, Betong); Narathiwat (Waeng).



Scientific Name	Thai name/Common name	Habitat	Site
<i>Amolops marmoratus</i>	กบลายหินเมืองเหนือ Marbled Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing water forest streams.	Mae Hong Son (Muang, Pa Sue Falls, Namtok Mae Surin); Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Kamphang Phet (Khlung Lan).
<i>Limnnectes gyldenstolpei</i>	กบทอง Gyldeustolpe's Frog	Inhabits evergreen forest along mountain streams.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Lam Pang (Doi Pa Muang, Doi Khun Tan); Phrae (Khao Pluang, Mae Song, Mae Yom); Kamphaeng Phet (Khlung Lan); Uttaradit; Loei (Phu Luang); Chaiyaphum (Phu Kieo); Udon Thani; Nong Khai (Phu Wua); Sakon Nakhon (Phu Phan); Kalasin (Phu Si Than); Nakhon Ratchasima (Khao Yai, Sakaerat); Roi Et (Phu Pa Nam Tip); Srakaew (Pang Sida, Tapraya); Prachin Buri (Tap Lan); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab, Khao Soidao); Rayong (Khao Cha Mao); Chon Buri (Khao Kieo); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Thung Yai).
<i>Limnnectes macrognathus</i>	กบหัวโต Big-headed Frog	Inhabits evergreen forest along mountain streams.	Mae Hong Son (Salawin); Chiang Mai (Doi Inthanon, Doi Suthep-Pui); Tak (Um Pang); Ranong (Muang); Surat Thani (Khao Sok; Khlung Saeng); Krabi (Khao Panombencha, Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Songkhla (Ton Ngachang).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Limnonectes plicatellus</i>	กบแรด Rhinoceros Frog	Inhabits evergreen forest along mountain streams.	Yala (Betong, Banangstar); Narathiwat Waeng).
<i>Limnonectes laticeps</i>	กบหลังตาพับ Rivulet Frog	Inhabits mountain streams in evergreen forest.	Chumphon (Mamoh); Yala (Bang Lang, Betong); Narathiwat (Waeng).
<i>Fejervarya cancrivora</i>	กบน้ำเค็ม Mangrove Frog	Inhabits mangrove forest, estuarine habitats, swamp and open wet areas.	Chon Buri (Ang Sila); Chachoengsao (Bang Pakong); Samut Prakan (Bang Pu, Khlong Dan); Bangkok (Bang Khun Tien); Prachuap Khirikhan (Sam Roi Yod); Chumphon (Swi); Nakhon Si Thammarat (Pak Panang); Sonkhla (Sating Pra); Narathiwat (Tak Bai); Trang (Kan Tang, Koh Libong, Hat Chao Mai); Phangnga (Ao Phangnga).
<i>Hoplobatrachus chinensis</i>	กบนา Chinese Bullfrog	Inhabits plain wetlands.	All provinces.
<i>Occidozyga laevis</i>	เขียดคลื่น Spotted Puddle Frog	Inhabits shallow muddy puddles and pools near small forest streams and forest edge.	Songkhla (Kor Hong); Pattani; Narathiwat (Sungai Kolok, Waeng)
<i>Occidozyga lima</i>	เขียดจระนา Common Puddle Frog	Inhabits lowland wetlands including wet paddy areas.	All provinces.
<i>Occidozyga magnapustulosa</i>	เขียดหลังปุ่ม Tubercled Frog	Occurs in seepages, puddles and shallow waters in high altitude forest.	Chiang Mai; Loei; Nakhon Phanom; Ubon Ratchathani.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Occidozyga martensii</i>	เขียดทราย Marten's Puddle Frog	Primarily occurs in streamside puddles, seepages and other damp patches along streams and rivers, as well as rain puddles and pools.	All provinces.
<i>Rana cf. alticola</i>	กบเขาสูง Hillstream Frog	Inhabits mountain streams in evergreen forest.	Uthai Thani (Hauai Khakhaeng); Kanchanaburi (Sai Yok, Sangkhlaburi, Thong Pa Phum); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Chumphon (Tasan); Ranong (Khlung Naka); Surat Thani (Khao Sok, Khlung Saeng); Phangnga (Khao Lak Lum Ru).
<i>Rana archotaphus</i>	กบชะง่อนผาอินทนนท์ Doi Inthanon Cascade Frog	Inhabits mountain streams fast water flows in high altitude forest.	Chiang Mai (Doi Inthanon); Chiang Rai (Mae Tum).
<i>Rana baramica</i>	กบวากเล็ก Masked Frog	Inhabits swampy areas lowland evergreen forest.	Narathiwat (Waeng).
<i>Rana cubitalis</i>	กบหูดำ Dark-eared Frog	Inhabits mountain streams in evergreen forest.	Chiang Mai (Doi Chiang Dao, Doi Nga Chang); Uthai Thani (Hauai Khakhaeng); Ratchaburi (Suan Pung); Phetchaburi (Kaengkrachan); Ranong (Khlung Naka); Nakhon Si Thammarat; Loei (Dan Sai, Na Haeo, Phu Luang); Phetchabun (Nam Nao).
<i>Rana erythraea</i>	เขียดจิก Green Paddy Frog	Inhabits lowlands wetlands including garden ponds.	All provinces.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rana glandulosa</i>	กบวากใหญ่ Rough-sided Frog	Inhabits swampy areas in forests and crops plantations.	Chumphon (Lang Suan); Phangnga (Khao Lak Lam Ru); Surat Thani (Pru Kantuli); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Trang; Satun (Tale Ban); Yala (Bannangstar, Betong); Narathiwat (Sungai Kolok, Waeng).
<i>Rana hosii</i>	กบชะงอนผาใต้ Hose's Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing water forest streams.	Chumphon (Ta San); Surat Thani (Khao Sok, Khlong Saeng); Phangnga (Khao Lak Lam Ru); Krabi (Khao Panombencha); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Yala (Banangstar); Narathiwat (Waeng).
<i>Rana johnsi</i>	กบสี่ชมพู Johns's Frog	Inhabits the high altitude forest.	Loei (Phu Luang).
<i>Rana livida</i>	กบชะงอนผาเหนือ Burmese Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing water forest streams.	Mae Hong Son (Pha Sue Falls); Chiang Mai (Doi Suthep-Pui); Kamphaeng Phet (Khlong Lan); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Hau Mae Kamin; Pilok, Salak Pra, Thong Pha Phum); Prachuap Khirikhan (Kui Buri, Pa-La-U).
<i>Rana macrodactyla</i>	กบหลังขี้ด Long-toed Frog	Known from lowland wetlands.	All provinces.
<i>Rana nicobariensis</i>	กบนิโคบาร์ Nicobar Cricket Frog	Occurs in disturbed areas and crop plantations.	Nakhon Si Thammarat, Yala.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rana nigrovittata</i>	กบอองเล็ก Dark-sided Frog	Inhabits gentle streams in evergreen forest.	All provinces except the Central Plain.
<i>Rana raniceps</i>	กบเขาหลังตอง White-lipped Frog	Individuals are seen on rocks or vegetation along small lowland forest streams.	Prachuap Khirikhan (Bang Sapan Noi); Chumphon (Mamoh, Tasae, Tasan); Ranong (Muang, Khlong Naka); Surat Thani (Khao Sok, Khlong Saeng, Na San, Ta Chana); Krabi (Khao Panom Bencha, Khao Pra Bang Kram); Nakhon Si Thammarat (Khao Luang); Trang (Khao Bantat, Khao Chong); Songkhla (Ton Ngachang); Yala (Bannangstar, Betong); Narathiwat (Sukirin, Sungai Kolok, Tak Bai, Waeng).
<i>Rana taipehensis</i>	กบไต้หวัน Taiwanese Frog	Inhabits grassy wetland areas, forest ponds and streams of deciduous and evergreen forests.	Phitsanulok (Thung Salaeng Luang); Loei (Phu Luang); Nong Khai (Phu Wua); Nakhon Ratchasima (Khao Yai).
<i>Chirixalus doriae</i>	ปาดจิวหลังซิด Doria's Bush Frog	Inhabits open grassy areas. Breeds in swampy pools.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Loei (Phu Kradung, Phu Luang); Chaiyaphum (Phu Kieo); Uthai Thani (Ban Rai); Kanchanaburi (Sangkhlaburi, Thung Yai).
<i>Chirixalus nongkhorensis</i>	ปาดจิวลายเต็ม Nongkhor Bush Frog	Inhabits open grassy areas surrounding with bamboo forest. Breeds in swampy pools.	Nakhon Ratchasima (Sakaerat); Srakaew (Pang Sida); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab); Chon Buri (Siracha).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Chirixalus vittatus</i>	ปาดจิวพม่า Burmese Bush Frog	Inhabits open grassy areas. Breeds in swampy pools.	Uthai Thani (Ban Rai); Kanchanaburi (Sangkhlaburi); Phra Nakhon Si Ayutthaya.
<i>Nyctixalus pictus</i>	ปาดป่าจูดขาว Cinnamon Tree Frog	Inhabits evergreen forest as well as bamboo forest.	Surat Thani (Khao Sok, Khlong Saeng); Phangnga (Khao Lak Lam Ru); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Yala (Bang Lang, Betong).
<i>Philautus parvulus</i>	ปาดแคระป่า Dwarf Bush Frog	Inhabits evergreen forest.	Mae Hong Son (Pang Tong); Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui); Pa Yao (Doi Pa Chang); Loei (Phu Kradung, Phu Luang); Chanthaburi (Khao Soi Dao, Khao Sebab); Tak (Thung Yai); Uthai Thani (Ban Rai); Phetchaburi (Kaengkrachan); Nakhon Si Thammarat (Khao Luang).
<i>Philautus petersi</i>	ปาดแคระทองเหลือง Peters's Bush Frog	Inhabits evergreen forest.	Narathiwat (Waeng).
<i>Polypedates colleti</i>	ปาดนิ้วแถมกลาย Collett's Tree Frog	Inhabits evergreen forest as well as Peat Swamp Forest.	Yala (Banangstar); Narathiwat (Sungai Kolok).
<i>Polypedates leucomystax</i>	ปาดบาน Four-lined Tree Frog	Found widespread in all habitats, from beach vegetation to primary evergreen forest as well as urban areas.	All Provinces.

Scientific Name	Thai name/Common name	Habitat	Site
<i>Polypedates macrotis</i>	ปาดหูดำ Dark-eared Tree Frog	Inhabits evergreen forest.	Nakhon Si Thammarat (Khao Luang); Yala (Bannagstar); Narathiwat (Waeng).
<i>Polypedates mutus</i>	เขียดตะปาด Northern Tree Frog	Inhabits primary and secondary forests.	Main provinces in the North and Northeast of the country.
<i>Rhacophorus appendiculatus</i>	ปาดลายเลอะไต Fringed Tree Frog	Inhabits evergreen forest.	Nakhon Si Thammarat (Khao Luang); Narathiwat (Waeng).
<i>Rhacophorus bipunctatus</i>	ปาดตีนเหลือง Twin-spotted Tree Frog	Inhabits lowland evergreen forest up to submontane forest.	Chiang Mai (Doi Chiang Dao, Doi Inthanon, Doi Suthep-Pui, Doi Pahom Pok, Kaeng Pan Tao); Loei (Phu Luang); Sakon Nakhon (Phu Phan); Kalasin (Phu Si Than); Nakhon Ratchasima (Sakaerat); Kamphaeng Phet (Khlung Lan); Kanchanaburi (Thung Yai); Phetchaburi (Kaengkrachan); Nakhon Si Thammarat (Khao Luang).
<i>Rhacophorus bisacculus</i>	ปาดลายเลอะอีสาน Taylor's Tree Frog	Inhabits lowland forest as well as rubber plantation surrounding the forest. Breeding in large temporary pool.	Phetchabun (Nam Nao); Loei ((Phu Kradung, Phu Luang); Nong Khai (Phu Wua); Nakhon Ratchasima (Sakaerat, Khao Yai); Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rhacophorus cyanopunctatus</i>	ปาดจุดฟ้า Blue-spotted Tree Frog	Inhabits lowland and submontane forests along mountain streams.	Surat Thani (Khao Sok); Nakhon Si Thammarat (Khao Luang); Songkhla (Ton Ngachang); Satun; Narathiwat (Waeng).
<i>Theلودerma asperum</i>	ปาดตะปุ่มตะป่ำ Pied Warted Tree Frog	Inhabits lowland and submontane forest.	Mae Hong Son (Pang Ung); Chiang Mai (Doi Chiang Dao, Doi Inthanon); Loei (Phu Luang); Tak (Thung Yai); Uthai Thani (Hau Khakhaeng); Ratchaburi (Maenam Pachi); Kanchanaburi (Si Nakharin); Srakaew (Pang Sida); Chanthaburi (Khao Kitchakud, Khao Sebab).



## Status **DD**

Scientific Name	Thai name/Common name	Habitat	Site
<i>Caudacaecilia asplenica</i>	เขียดจุมถลาย Malayan Caecilian	Fossorial dwells underground of moist evergreen forest.	Yala (La Doo Tin Mine).
<i>Caudacaecilia larutensis</i>	เขียดจุมดำ, เขียดจุมบาลา Larut Hill Caecilian	Fossorial dwells underground of moist evergreen forest.	Narathiwat (Khlong Tan Tong, Su Kirin District; Sirindhorn Waterfall, Waeng District).
<i>Ichthyophis acuminatus</i>	เขียดจุมหัวแหลม Pointed-snouted Caecilian	Fossorial dwells underground of moist evergreen forest.	Chiang Mai (Mae Wang Valley).
<i>Ichthyophis kohtaoensis</i>	เขียดจุมเกาะเต่า Koh Tao Caecilian	Fossorial dwells underground of evergreen forest and agricultural areas.	Chiang Mai (Doi Suthep); Uthai Thani (Hau Kha Khaeng WS); Sing Buri (Prom Buri District); Loei (Phu Luang WS); Chaiyaphum (Phu Kieo WS); Nakhon Ratchasima (Wang Nam Kieo District); Srakaew (Pang Sida NP); Surat Thani (Koh Tao, Koh Samui).
<i>Ichthyophis supachaii</i>	เขียดจุมศุภชัย Supachai's Caecilian	Fossorial dwells underground of evergreen forest and agricultural areas.	Nakhon Si Thammarat (Khao Luang NP); Trang (Khao Bantat WS); Phthalung (Tale Noi WS); Yala (Than To Falls); Narathiwat (Waeng District).
<i>Ichthyophis youngorum</i>	เขียดจุมดอยสุเทพ Doi Suthep Caecilian	Fossorial dwells underground of montane forest about 1,000 m elevation.	Chiang Mai (Doi Suthep).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Brachytarsophrys feae</i>	อึ่งกรายพม่า Fea's Horned Frog	Inhabits evergreen forests in high elevation. Only one specimens collected from Angka Luang 2,800 m elevation in Doi Inthanon now is depicted in FMNH Collection.	Chiang Mai (Doi Inthanon).
<i>Leptobrachium chapaense</i>	อึ่งกรายตาขาว White-eyed Litter Frog	Inhabits evergreen forests in high elevation.	Chiang Mai (Doi Inthanon).
<i>Leptolalax bourreti</i>	อึ่งกรายคอดยอินทนนท์ Bourret's Litter Frog	Hill evergreen forest in high elevation closed to 2,800 m.	Chiang Mai (Doi Inthanon).
<i>Leptolalax cf. gracilis</i>	อึ่งกรายทองผาภูมิ Thong Pa Pum Litter Frog	Ecological information of this species is confined to primary and old secondary rain forests. Adults and juveniles live in the floor litter and underground covered vegetation.	Doubtful record of tadpole from Thong Pa Phum in Kanchanaburi.
<i>Hyla annectans</i>	ป่าดเมืองจีน Chinese Tree Frog	Probably found in moist rain forest.	Mae Hong Son.
<i>Chaperina cf. fusca</i>	อึ่งทองเหล็อง Spiny-heeled Froglet	Inhabits primary evergreen forest.	Surat Thani (Khlung Saeng).
<i>Microhyla annamensis</i>	อึ่งอันนัม Annam Chorus Frog	Inhabits swampy areas in evergreen forest.	Chanthaburi (Khao Sebab).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Microhyla annectens</i>	อึ่งน้ำตามลาชู Boulenger's Chorus Frog	Inhabits the forest floor.	Chumphon (Pa Tiu).
<i>Microhyla borneensis</i>	อึ่งบอร์เนียว Bornean Chorus Frog	Inhabits the forest floor closed to the streams.	Narathiwat (Waeng).
<i>Phrynella pulchra</i>	อึ่งทองแดง Red-bellied Frog	Inhabits evergreen forest.	Narathiwat (Waeng).
<i>Limnonectes doriae</i>	กบดอร์เรีย Doria's Frog	Inhabits evergreen forest.	Tak; Uthai Thani (Hau Khakhaeng); Kanchanaburi (Pi Lok, Sai Yok, Sangkhlaburi, Thong Pha Phum); Prachuap Khirikhan (Huai Yang Falls); Chumphon (Ta Sae).
<i>Chaparana aenea</i>	กบดอยช้าง Doi Chang Frog	Inhabits evergreen forest.	Chiang Mai (Doi Chang, Doi Inthanon).
<i>Fejervarya limnocharis</i>	กบหนอง Paddy Frog	Inhabits most open wetlands, rarely found in closed canopy forest.	All provinces.
<i>Fejervarya raja</i>	กบราชา Raja Frog	Inhabits marsh areas.	Phatthalung (Thale Noi); Songkhla (Thale Sap); Pattani (Ru Samilae); Narathiwat (Tak Bai).
<i>Huia melasma</i>	กบชะง่อนผาไทย Siamese Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing.	Chiang Mai (Doi Chiang Dao); Kanchanaburi (Tan Lot Cave); Prachuap Khirikhan (Pa-La-U).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Rana banjarana</i>	กบหลังจุดมลายู Banja Frog	Inhabits mountain streams in high altitude forest.	Yala (Banangstar).
<i>Rana chloronota</i>	กบชะง่อนผาอีสัม Assamese Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing water forest streams.	Prachuap Khirikhan (Pa-La-U); Surat Thani (Khao Sok); Krabi (Khao Panombencha).
<i>Rana luctuosa</i>	กบเขาทองลาย Mahogany Frog	Inhabits forest floor of lowland evergreen forest. Breeding in wet wild boar hollows.	Nakhon Si Thammarat (Khao Ram).
<i>Rana milleti</i>	กบเขาใหญ่ Millet's Frog	Known from mountain streams in degraded and primary forests.	Umnajjareon (Phu Sing); Nakhon Ratchasima (Khao Yai), Chachoengsao (Khao Ang Rue Ni); Chanthaburi (Khao Sebab).
<i>Rana mortenseni</i>	กบอ่องใหญ่ Mortensen's Frog	Inhabits forest streams in evergreen forest.	Trat (Koh Chang); Chanthaburi (Khao Sebab).
<i>Rana schmackeri</i>	กบคอยอินทนนท์ Schmacker's Cascade Frog	Known from mountain stream in hill evergreen forest.	Chiang Mai (Doi Inthanon).
<i>Taylorana hascheana</i>	กบป่าไผ่ Hasche's Frog	Inhabits bamboo forest, usually found in dense leaf litter layer on soil.	Mae Hong Son (Salawin); Chiang Mai (Doi Suthep-Pui); Kamphaeng Phet (Khlung Lan, Mae Wong); Uthai Thani (Hau Khakhaeng); Kanchanaburi (Thong Pa Phum); Phetchaburi (Kaengkrachan); Prachuap Khirikhan (Pa-La-U); Ranong (Maung); Surat Thani (Khao Sok); Krabi (Khao Panombencha).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Taylorana limborgii</i>	กบกา Stoliczlea's Frog	Inhabits bamboo forest, usually found in dense leaf litter layer on soil.	Nakhon Si Thammarat (Khao Luang); Trang (Khao Chong); Songkhla (Ton Ngachang); Narathiwat (Waeng).
<i>Chirixalus hansenae</i>	ปาดจ้าวตะวันออก Hansen Bush Frog	Inhabits open grassy areas. Breeds in swampy pools.	Chiang Mai (Kaeng Paeng Tao, Nong Pung); Loei (Phu Kradung, Phu Luang); Nakhon Ratchasima (Sakaerat); Chon Buri (Siracha).
<i>Philautus carinensis</i>	ปาดกระพม่า Burmese Bush Frog	Inhabits tropical submontane forest.	Chiang Mai (Doi Chiang Dao, Doi Inthanon).
<i>Philautus gracilipes</i>	ปาดกระขาเรียว Slender-legged Bush Frog	Inhabits high altitude hill evergreen forest.	Chiang Mai (Doi Inthanon).
<i>Philautus vermiculatus</i>	ปาดกระทองลาย Vermiculated Bush Frog	Inhabits evergreen forest.	Narathiwat (Waeng).
<i>Rhacophorus orlovi</i>	ปาดตีนเหลืองอีสาน Orlov's Tree Frog	Inhabits along sandstone mountain streams in deciduous and evergreen forests.	Umnajjareon (Muang); Ubon Ratchathani (Nachaluai).
<i>Rhacophorus verrucosus</i>	ปาดลายเลอะเหนื่อ Boulenger's Tree Frog	Inhabits lowland evergreen forest.	Tak (Um Pang); Prachuap Khirikhan (Pa-La-U).

## Status **Endemic**

Scientific Name	Thai name/Common name	Habitat	Site
<i>Ichthyophis acuminatus</i>	เขียดคูดหัวแหลม Pointed-snouted Caecilian	Fossorial dwells underground of moist evergreen forest.	Chiang Mai (Mae Wang Valley).
<i>Ichthyophis kohtaoensis</i>	เขียดคูดเกาะเต่า Koh Tao Caecilian	Fossorial dwells underground of evergreen forest and agricultural areas.	Chiang Mai (Doi Suthep); Uthai Thani (Hau Kha Khaeng WS); Sing Buri (Prom Buri Distric); Loei (Phu Luang WS); Chaiyaphum (Phu Kieo WS); Nakhon Ratchasima (Wang Nam Kieo District); Srakaew (Pang Sida NP); Surat Thani (Koh Tao, Koh Samui).
<i>Ichthyophis youngorum</i>	เขียดคูดอยสุเทพ Doi Suthep Caecilian	Fossorial dwells underground of montane forest about 1,000 m elevation.	Chiang Mai (Doi Suthep).
<i>Ansonia inthanon</i>	คางคกห้วยอินทนนท์ Inthanon Stream Toad	The endemic species inhabits in high altitude of mountain forest, but the population in Kanchanaburi is probably swampy area in lowland evergreen forest.	Chiang Mai (Doi Inthanon); Kanchanaburi (Thong Pa Phum).
<i>Ansonia siamensis</i>	คางคกห้วยไทย Siamses Stream Toad	Inhabits near streams of evergreen forest.	Trang (Khao Chong).

Scientific Name	Thai name/Common name	Habitat	Site
<i>Huia melasma</i>	กบชะง่อนผาไทย Siamses Cascade Frog	Exclusively occurs on boulders and bedrock in fast flowing.	Chiang Mai (Doi Chiang Dao); Kanchanaburi (Tan Lot Cave); Prachuap Khirikhan (Pa-La-U).
<i>Ingerana tasanae</i>	กบทาสาร Tasan Frog	Inhabits mountain streams in evergreen forest.	Chumphon (Tasan); Ranong (Khlong Naka); Surat Thani (Khao Sok, Khlong Saeng); Phangnga (Khao Lak Lam Ru).

## References

- Adulyanukosol, K. and K. Kittiwattanawong. 2004. Field Guide to Cetaceans of Thailand. Phuket Marine Biological Center, Department of Marine and Coastal Resources, Ministry of Natural Resources and Environment. (in Thai)
- Ananjeva, N.B. and B.L. Stuart. 2001. The Agamid Lizard *Ptyctolaemus phuwuaensis* Manthey and Nabhitabhata, 1991 from Thailand and Laos Represent a New Genus. *Russian J. Herpetol.* 8(3): 165-170.
- Andersen, M. and C.C. Kinze. 2000. Annotated checklist and identification key to the whales, dolphins, and porpoises (Order Cetacea) of Thailand and adjacent water. *Nat. Hist. Bull.* 47: 27-62.
- Andersen, M. and C.C. Kinze. 2000. Review and new records of the marine mammals and sea turtles of indochinese waters. *Nat. Hist. Siam Soc.* 48: 177-184.
- Bauer, A.M. and I. Das. 1998. New Species of *Cnemaspis* (Reptilia: Gekkonidae) from Southeastern Thailand. *Copeia* 1998(2): 439-444.
- Bauer, A.M., M. Sumontha and O.S.G. Pauwels. 2003. Two new species of *Cyrtodactylus* (Reptilia: Squamata: Gekkonidae) from Thailand. *Zootaxa* 376: 1-18.
- Bauer, A.M., M. Sumnontha, W. Grossmann, O.S.G. Pauwels and G. Vogels. 2004. A New Species of *Dixoneus* (Squamata: Gekkonidae) from Kanchanaburi Province, Western Thailand. *Current Herpetol.* 23(1): 17-26.
- Bauer, A.M., O.S.G. Pauwels and L. Chanhomes. 2002. A New Species of Cave-dwelling *Cyrtodactylus* (Squamata: Gekkonidae) from Thailand. *Nat. Hist. J. Chula. Univ.* 2(2): 19-29.
- Bhompakphan, N., P. Thammaroekrid, P. Thanchai, P. Manee-ngam and T. Puangsri. 2003. Siamese Crocodile known from the Lum Dom Yai River, Ubon Rachatani. *Wild. J.* 11(1): 148-155.
- Bohme, W. 1981. A new lygosomine skink from Thailand (Reptilia: Scincidae). *Bull. Mus. Civ. St. Nat. Verona* 8(1981): 375-382.
- Bohme, W. 1982. Concerning Butterfly Agamas, *Leiolepis b. belliana* (Gray, 1827) from the Malay Peninsula and Their Parthenogenetic Lines (Sauria: Uromasticidae). *Zool. Jb. Syst.* 109 (1982): 157-169.
- Boulenger, G.A. 1912. A Vertebrate Fauna of the Malay Peninsula from Isthmus of Kra to Singapore including the adjacent islands. p. 64.
- Brown, R.M. 1999. New Species of Parachute Gecko (Squamata: Gekkonidae: Genus *Ptychozoon*) from Northeastern Thailand and Central Vietnam. *Copeia* 1999(4): 990-1001.



- Campden-Main, S.M. 1969. The status of *Oligodon taeniatus* (Gunther), 1861, and *Oligodon mouhoti* (Boulenger), 1914 (serpents, Colubridae). *Herpetologica* 25: 295-299.
- Chan-ard, T. 2003. A photographic Guide of Amphibians in Thailand. Darnsutha Press Co., Ltd., Bangkok.
- Chan-ard, T., W. Grossmann, A. Gumprecht and K.D. Schulz. 1999. Amphibians and reptiles of Peninsular Malaysia and Thailand- an illustrated checklist. Bush Master, Wurselen, Germany.
- Chan-ard, T., Y. Chauynkern and S. Thong-aree. 2003. The Diversity of Herpetofauna in Hala-Bala Wildlife Sanctuary, Yala and Narathiwat Provinces. *BRT 2003 Research Report*, 245-258.
- Cochran, D.M. 1930. The Herpetological Collections made by Dr. Hugh M. Smith in Siam from 1923 to 1929. *Proc. U.S. Nat. Mus.* 77: 1-39.
- Corbet, G.B. and J.E. Hill. 1991. A World List of Mammalian Species. 3<sup>rd</sup>. ed. Oxford University Press, Oxford.
- Cox, M.J. 1991. The Snakes of Thailand and Their Husbandry. Krieger Publishing Comp., Malabar, Florida.
- Daltry, J.C. and W. Wuster. 2002. A New Species of Wolf Snake (Serpentes: Colubridae: *Lycodon*) from the Cardamon Mountains, Southwestern Cambodia. *Herpetologica* 58(4): 498-504.
- Darevsky, I.S. and L.A. Kupriyanova. 1993. Two new all-female lizard species of the genus *Leiolepis* Cuvier, 1829 from Thailand and Vietnam (Squamata: Sauria: Uromastycinae). *Herpetozoa* 6(1/2): 3-20.
- David, P. and I. Ineich. 1999. Les serpents venimeux du monde: systematique et repartition. *Dumerilia* 3: 1-300.
- David, P., G. Vogel, M. Sumontha, O.S.G. Pauwels and L. Chanhome. 2004. Expanded description of the poorly known pitviper *Trimeresurus kanburiensis* Smith, 1943, with confirmation of the validity of *Trimeresurus venustus* Vogel, 1991 (Reptilia: Serpentes: Crotalidae). *Russian J. Herpetol.* 11(2): 81-90.
- David, P., M.J. Cox, O.S.G. Pauwels, L. Chanhome and K. Thirakupt. 2004. When a bookreview is not sufficient to say all: an in-depth analysis of a recent book on the snakes of Thailand, with an updated checklist of the snakes of the Kingdom. *Nat. Hist. J. Chula. Univ.* 4(1): 47-80.
- David, P., N. Vidal and O.S.G. Pauwels. 2001. A morphological study of Stejneger's pit viper *Trimeresurus stejnegeri* Serpentes, Viperidae, Crotalinae, with a description of a new species from Thailand. *Russian J. Herpetol.* 8: 205-222.
- Deuve, J. 1970. Serpent du Laos. O.R.S.T.O.M., Paris.

- Dowling, H.G. and J.V. Jenner. 1988. Rediscovery of the type specimen of *Oligodon mcdougalli* with a description of its relationships. In The Snakes of Burma. Smithsonian Herpetological Information Service, no.76, 1988. Dept. Biol., New York Univ.
- Dring, J.C.M. 1979. Amphibians and reptiles from northern Tregganu, Malaysia, with the descriptions of two new geckos: *Cnemaspis* and *Cyrtodactylus*. *Bull. Br. Mus. nat. Hist. (Zool.)* 34(5): 181-241.
- Dubois, A. and A. Ohler. 2000. Systematics of *Fejervarya limnocharis* (Gravenhorst, 1829)(Amphibia, Anura, Ranidae) and related species. 1. Nomenclatural status and type- specimens of the nominal species *Rana limnocharis* Gravenhorst, 1829. *Alytes* 18(1,2): 15-50.
- Duengkae, P. 1998. Wild Mammals in Thailand. Office of Environment Policy and Planning, Bangkok, Thailand. (in Thai)
- Frith, C.B. and D. MacIver. 1977. The Crab-eating Water Snake, *Fordonia leucobalia* (Schleg.); Another Snake New to Thailand. *Nat. Hist. Bull. Siam Soc.* 27:189-191.
- Frith, C.B. and J. Boswall. 1978. Cantor's Water Snake, *Cantoria violacea* Girard; a Vertebrate New to the Fauna of Thailand. *Nat. Hist. Bull. Siam Soc.* 27: 187-188.
- Fritz, U. and T. Ziegler. 1999. Contribution to the knowledge of *Cyclemys tcheponensis* (Bourret, 1939) and the distribution of *Cyclemys* in the Indochinese region (Reptilia: Testudines: Bataguridae). *Revue fr. Aquariol.* 26(1999): 71-78.
- Grandison, A.C.G. 1972. The Gunong Benom Expedition 1967. 5. Reptiles and Amphibians of Gunong Benom with a Description of a New Species of *Macrocalamus*. *Bull. Br. Mus. Nat. Hist. Zool.* 23(4): 45-101.
- Greer, A.E. 1997. *Leptoseps*: A New Genus of Scincid Lizards from Southeast Asia. *J. Herpetol.* 31(3): 393-398.
- Grossmann, W. and F. Tillack. 1998. Bemerkungen zur Verbreitung von *Elaphe flavolineata* (Schlegel, 1837). *Sauria, Berl.* 20(3): 43-45.
- Grossmann, W. and T. Ulber. 1990. Ein neuer Gecko aus Zentral-Thailand: *Gekko siamensis* sp.nov. (Reptilia: Sauria: Gekkonidae). *Sauria, Berl.* 12(3): 9-18.
- Guicking, D., U. Fritz, M. Wink, and E. Lehr. 2002. New data on the diversity of the Southeast Asian leaf turtle genus *Cyclemys* Bell, 1834. Molecular results (Reptilia: Testudines: Geoemydidae). *Faun. Abh. Mus. Tierkd. Dresden* 23(4): 75-86
- Heyer, W.R. 1972. A New Limbless Skink (Reptilia: Scincidae) From Thailand With Comments On the Generic Status of the Limbless Skinks. *Fieldiana Zoology* 58(10): 109-129.

- Hikida, T. 1990. Bornean Gekkonid Lizards of the Genus *Cyrtodactylus* (Lacertilia: Gekkonidae) with Descriptions of Three New Species. *Jpn. J. Herpetol.* 13(3): 91-107.
- Hikida, T., N.L. Orlov, J. Nabhitabhata and H. Ota. 2002. Three New Depressed-bodied Water Skinks of the Genus *Tropidophorus* (Lacertilia: Scincidae) from Thailand and Vietnam. *Current Herpetol.* 21(1): 9-23.
- Honda, M., J. Nabhitabhata, H. Ota and T. Hikida. 1997. A new species of *Dibamus* (Squamata: Dibamidae) from Thailand. *Raffles Bull. Zool.* 45(2): 275-279.
- Horton, D.R. 1973. A New Scincid Genus from Southeast Asia. *J. Herpetol.* 7(3): 283-287.
- Iskandar, D.T. and E. Colijn. 2001. Checklist of the Southeast Asian and New Guinian Reptiles. Part I: Serpents. Kodokijo- Binamitra, Jakarta.
- IUCN. 2001. IUCN Red List Categories and Criteria: Version 3.1. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, U.K.
- Iverson, J.B. and W.P. McCord. 1997. A new species of *Cyclemys* (Testudines: Bataaaguridae) from Southeast Asia. *Proc. Biol. Soc. Washington* 110(4): 629-639.
- Janthrapornsil, S. 1985. Experimental cultivation of sea turtles in styrofoam boxes. Phuket Marine Biological Center. Department of Fisheries. Academic Document. (in Thai).
- Kiew, B.H. 1984. A New Species of Toad, *Ansonia siamensis* (Bufonidae), from the Isthmus of Kra, Thailand. *Nat. Hist. Bull. Siam Soc.* 32(2): 111-115.
- Kitimasak, W. and K. Therakhupt. 2002. New Record of *Chitra chitra* Nutphand, 1986 in Mae Ping River, Thailand. *Nat. Hist. Chula. Univ.* 2(1): 62-63.
- Kosuch, J., M. Vences, A. Dubois, A. Ohler and W. Bohme. 2001. Out of Asia: Mitochondrial DNA Evidence for an Oriental Origin of Tiger Frogs, Genus *Hoplobatrachus*. *Molecular Phylogenetics and Evolution* 21(3): 398-407.
- Lang, M. and W. Bohme. 1990. Description and phylogenic position of a new species of *Isopachys* from central Thailand and southern Burma (Squamata: Scincidae). *Bull. Ins. Roy. Sci Nat Bel. Biologie* 60: 231-240.
- Laohajinda, V., S. Wongtirawat and P. Vewattana. 1999. Amphibians in Forest of Thon Pa Phum District, Kanchanaburi Province. *Thailand's Wild Animal Journal* 7(1): 24-29. (in Thai)
- Lekagul, B. and J.A. McNeely. 1977. Mammals of Thailand. Kurusapha Ladprao Press, Bangkok.
- Leong, T.M. and B.L. Lim. 2003. A New Species of *Rana* (Amphibia: Anura: Ranidae) from the Highlands of the Malay Peninsula, With Diagnostic Larval Descriptions. *Raffles Bull. Zool.* 51(1): 115-122.

- Leong, T.M., T. Chan-ard and Y. Chaynkern. 2003. Additional Anuran and Saurian Records for Phuket, South Thailand. *Nat. Hist. J. Chula. Univ.* 3(1): 17-21.
- Malholtra, A. and R.S. Thorpe. 2004. A phylogeny of four mitochondrial gene regions suggests a revised taxonomy for Asian pitvipers (*Trimeresurus* and *Ovophis*). *Molecular Phylogenetics and Evolution* 32 (2004): 83-100.
- Manthey, U. and J. Nabhitabhata. 1991. An Agamid lizard, *Ptyctolaemus phuwuaensis* Manthey and Nabhitabhata 1991 (Sauria: Agamidae), from Northeast Thailand. *Sauria (E)* 1(2): 3-6.
- Matsui, M., J. Nabhitabhata and S. Panha. 1998. A New *Ansonia* from Northern Thailand (Anura: Bufonidae). *Herpetologica* 54(4): 448-454.
- Matsui, M., J. Nabhitabhata, T. Chan-ard and K. Thirakupt. 1996. Amphibian fauna of Thailand observed in 1994 and 1995, pp. 28-63. In M.Matsui (Ed). *Evolutionary Studies of Small Animals Living in Asian Tropics 1994-1995*. Graduate School of Human and Environmental Studies, Kyoto University.
- McCord, W.P. and P.C.H. Pritchard. 2002. A Review of the Softshell Turtles of the Genus *Chitra*, With the Description of New Taxa from Myanmar and Indonesia (Java). *Hamadryad* 27(1): 11-56.
- McKenna, M.C. and S.K. Bell. 1997. *Classification of Mammals above the Species Level*. Columbia University Press, New York.
- Musters, C.J.M. 1983. Taxonomy of the genus *Draco* L. (Agamidae, Lacertilia, Reptilia). *Zoologische Verhandelingen* 199: 1-120.
- Nabhitabhata, J., T. Chan-ard and Y. Chauyngern. 2000. Checklist of Amphibians and Reptiles in Thailand. Office of Environmental Policy and Planning, Bangkok.Thailand.
- Nicholls, L. 1949. A New Gekkonid from the Malay Peninsula. *Bull. Raffles Mus.* 1949, 47-49.
- Niyomwan, P. 1999. Species diversity, morphology and habitat types of blind snakes (Family typhlopidae) in Thailand. M.Sc. Thesis, Chulalongkorn University.
- Nowak, R.M. 1991. *Mammals of the Worlds. Volume 1*. The Johns Hopkins University Press, Baltimore and London.
- Nussbaum, R.A., E.D. Broadie and D. Yang. 1995. A Taxonomic Review of *Tylostotriton verrucosus* Anderson (Amphibia: Caudata: Salamandridae). *Herpetologica* 51(3): 257-268.
- Nutphand, W. 1976. Wolf Snakes. *Bull. Thai Zool.* 1(9): 1-7.
- Office of Environmental Policy and Planning. 1997. *Proceeding on Status of Biological Resources in Thailand*. Ministry of Science, Technology and Environment. Bangkok. (in Thai).

- Ohler, A. and A. Dubois. 1999. The identity of *Elachy glossa gyldenstolpei* Andersson, 1916 (Amphibia, Ranidae), with comments on some aspects of statistical support to taxonomy. *Zoologica Scripta* 1999: 269-279.
- Ota, H. and J. Nabhitabhata. 1991. A New Species of Gekko (Gekkonidae: Squamata) from Thailand. *Copeia* 1991(2): 503-509.
- Ouboter, P.E. 1986. A revision of the genus *Scincella* (Reptilia: Sauria: Scincidae) of Asia, with some notes on its evolution. *Zoologische Verhandelingen* 129: 1-66.
- Parr, J.W.K. 2003. Large mammals of Thailand. Sarakadee Press, Bangkok.
- Pauwels, O.S.G., A.M. Bauer, M. Sumontha and L. Chanhom. 2004. *Cyrtodactylus thirakhupti* (Squamata: Gekkonidae), a new cave-dwelling gecko from southern Thailand. *Zootaxa* 772: 1-11.
- Pauwels, O.S.G., O. Laohawat, P. David, R. Bour, P. Dangsee, C. Puanggit and C. Chimsunchart. 2000. Herpetological investigations in Phang-Nga Province, southern Peninsular Thailand, with a list of reptile species and notes on their biology. *Dumerilia* 4(2): 123-154.
- Pauwels, O.S.G., O. Laohawat, W. Naaktae, C. Puangjit, T. Wisatharom, C. Chimsunchart and P. David. 2002. Reptile and Amphibian Diversity in Phang-nga Province, Southern Thailand. *Nat. Hist. J. Cula. Univ.* 2(1): 25-30.
- Pauwels, O.S.G., P. David, W. Nutphand and C. Chimsunchart. 2001. First Record of *Xenochrophis punctulatus* (Gunther, 1858) (Serpentes: Colubridae: Natricinae) from Thailand. *Hamadryad* 26(2): 247-252.
- Rasmussen, J.B. 1982. A new record of the rare *Opisthotropis praemaxillaris* (Serpentes: Colubridae). *Amphibia-Reptilia* 3: 279-280.
- Schultz, K.D. and N. Helfenberger. 1998. Ein Revision des Unterarten-Komplexes der Roten Bambusnatter *Elaphe porphyraea* (Cantor, 1839). *Sauria, Berl.* 20(1): 25-45.
- Smith, M.A. 1919. *Crocodylus siamensis*. *J. Nat. Hist. Soc. Siam.* 3(3): 217-221.
- Smith, M.A. 1935. The Fauna of British India including Ceylon and Burma. Reptilia and Amphibia. Vol.II- Sauria. pp.158-160.
- Taksintum, V. Lauhachinda and B. Kanchanasaka. 2003. First Record of Froglet, Genus *Chaperina*, in Thailand. *Wild. J.* 11(1): 156-163.
- Taylor, E.H. 1934. Zoological of Results of the Third De Schauensee Siamese Expedition, Part III.- Amphibians and Reptiles. *Proc. Akad. Nat. Sci. Phila.* 1934: 281-310.
- Taylor, E.H. 1962. The amphibian fauna of Thailand. *Univ. Kansas Sci. Bull.* 43: 583-586.
- Taylor, E.H. 1963. The lizards of Thailand. *Univ. Kansas Sci. Bull.* 44: 866-870.

- Taylor, E.H. 1965. New Asiatic and African Caecilians with Redescriptions of Related Species. *Univ. Kansas Sci. Bull.* 46: 253-302.
- Taylor, E.H. 1965. The surpents of Thailand and adjacent waters. *Univ. Kansas Sci. Bull.* 45: 776-779.
- Taylor, E.H. 1968. The caecilians of the world: a taxonomic review. Univ. Kansas Press, Lawrence. pp. 167-171.
- Taylor, E.H. 1970. The turtles and crocodiles of Thailand and adjacent waters, with a synoptic herpetological bibliography. *Univ. Kansas Sci. Bull.* 49: 87-179.
- Ulber, T. 1993. Notes on Cyrtodactyline Geckos from Thailand with Descriptions of Two New Species (Reptilia: Gekkonidae). *Mitt. Zool. Mus. Berl.* (1993)2, 187-200.
- Utiger, U., N. Helfenberger, B. Schatti, C. Schmidt, M. Ruf and Z. Ziswiler. 2002. Molecular systematics and phylogeny of Old and New World Ratsnakes, *Elaphe* Auct., and related genera (Reptilia, Squamata, Colubridae). *Russ. J. Herp.* 9(2): 105-124.
- Van Dijk, P.P. 1998. A Review of the Conservation Status of Tortoises and Freshwater Turtles in Thailand. Final Report submitted to The Nature Conservation Bureau, Environmental Agency of Japan. 54 p.
- Vogel, G. 1990. Die Bronzenattern Thailands. *Herpetofauna* 12(65): 10-16.
- Vogel, G., P. David and O.S.G. Pauwels. 2004. A review of morphological variation in *Trimeresurus popeiorum* (Serpentes: Viperidae: Crotalinae), with the description of two new species. *Zootaxa* 727: 1-63.
- Wagner, F.W. 1973. A revision of the Asian colubrid snakes *Oligodon cinereus* (Gunther), *Oligodon joynsoni* (Smith), and *Oligodon cyclurus* (Cantor). M.Sc. Thesis, Arizona State Univ.
- Wallach, V. 2001. *Typhlops roxanae*, a new species of Thai blindsnake from the *T. diardii* species group, with a synopsis of the Typhlopidae of Thailand (Serpentes: Scolecophidia). *Raffles Bull. Zool.* 49(1): 39-49.
- Warrell, D.A., S. Looreesuwan, A.F. Stimson and R.A. Hutton. 1992. Rediscovery and redefinition of Malcolm Smith's *Trimeresurus kanburiensis* in Thailand, with a report of envenoming. *Trans. Roral Soc. Trop. Med. Hygiene* 86: 95-99.
- Welch, K.R.G., P.S. Cooke and A.S. Wright. 1990. Lizards of the Orient: a checklist. R.E.Krieger Pub. Comp., Malabar. 162 p.
- Wongratana, T. 1984. Range extention of the crocodile salamander, *Tylototriton verrucosus*, to Phu Luang, Thailand. *Nat. Hist. Bull. Siam Soc.* 32: 107-110.

# Index

## Mammal

### Sort by Scientific Name

<i>Anourosorex squamipes</i>	67	<i>Felis chaus</i>	16
<i>Atherurus macrourus</i>	81	<i>Feresa attenuata</i>	54
<i>Axis porcinus</i>	58	<i>Globicephala macrorhynchus</i>	52
<i>Balaenoptera edeni</i>	43	<i>Hapalomys longicaudatus</i>	63
<i>Balaenoptera physalus</i>	42	<i>Harpiocephalus harpia</i>	70
<i>Belomys pearsonii</i>	63	<i>Helarctos malayanus</i>	66
<i>Bos gaurus</i>	74	<i>Hemigalus derbyanus</i>	34
<i>Bos javanicus</i>	25	<i>Hesperoptenus blanfordi</i>	69
<i>Bos sauveli</i>	12	<i>Hesperoptenus tickelli</i>	69
<i>Bubalus bubalis</i>	59	<i>Hipposideros halophyllus</i>	38, 84
<i>Callosciurus nigrovittatus</i>	82	<i>Hipposideros lekaguli</i>	68
<i>Callosciurus prevostii</i>	15	<i>Hipposideros turpis</i>	68
<i>Canis aureus</i>	65	<i>Hylobates agilis</i>	21
<i>Capricornis sumatraensis</i>	78	<i>Hylobates lar</i>	72
<i>Cervus eldii</i>	11	<i>Hylobates pileatus</i>	41
<i>Cervus schomburgki</i>	10, 84	<i>Hylobates syndactylus</i>	77
<i>Cheiromeles torquatus</i>	68	<i>Hylopetes alboniger</i>	62
<i>Cherephon johorensis</i>	76	<i>Hylopetes spadiceus</i>	62
<i>Chironomyscus chiropus</i>	80	<i>Hystrix brachyura</i>	81
<i>Coelops frithi</i>	68	<i>Ia io</i>	69
<i>Coelops robinsoni</i>	83	<i>Ictailurus planiceps</i>	17
<i>Craseonycteris thonglongyai</i>	20	<i>Kerivoula hardwickei</i>	70
<i>Crocidura monticola</i>	82	<i>Kerivoula minuta</i>	70
<i>Crocidura pullata</i>	75	<i>Kerivoula papillosa</i>	70
<i>Cuon alpinus</i>	65	<i>Kerivoula whiteheadi</i>	70
<i>Cynogale bennettii</i>	18	<i>Kogia breviceps</i>	46
<i>Cynoptyerus horsfieldii</i>	81	<i>Kogia simus</i>	45
<i>Cynoptyerus sp.</i>	83	<i>Lariscus insignis</i>	62
<i>Delphinus capensis</i>	48	<i>Leopoldamys edwardsi</i>	63
<i>Dendrogale murina</i>	73	<i>Leopoldamys neilli</i>	63, 84
<i>Dicerorhinus sumatrensis</i>	14	<i>Lutra lutra</i>	35
<i>Dugong dugon</i>	26	<i>Lutra sumatrana</i>	19
<i>Dyacopterus spadiceus</i>	67	<i>Lutrogale perspicillata</i>	66
<i>Echinosorex gymnurus</i>	67	<i>Macaca arctoides</i>	70
<i>Elephas maximus</i>	61	<i>Macaca assamensis</i>	40
<i>Eothenomys miletus</i>	63	<i>Macaca nemestrina</i>	77
<i>Eptesicus demissus</i>	39, 84	<i>Manis javanica</i>	75
<i>Eptesicus pachyotis</i>	69	<i>Manis pentadactyla</i>	29
<i>Eptesicus serotinus</i>	69	<i>Maxomys whiteheadi</i>	63

<i>Mesoplodon ginkgodens</i>	22	<i>Pteropus hypomelanus</i>	75
<i>Miniopterus medius</i>	83	<i>Pteropus intermedius</i>	82
<i>Muntiacus feae</i>	57	<i>Pteropus lylei</i>	76
<i>Murina huttonii</i>	77	<i>Pteropus vampyrus</i>	67
<i>Murina leucogaster</i>	69	<i>Ptilocercus lowii</i>	73
<i>Mustela nivalis</i>	82	<i>Rattus annandalei</i>	80
<i>Mustela nudipes</i>	66	<i>Rattus osgoodi</i>	82
<i>Mustela sibirica</i>	66	<i>Rattus remotus</i>	63
<i>Mustela strigidorsa</i>	36	<i>Ratufa affinis</i>	62
<i>Myotis altarium</i>	69	<i>Ratufa bicolor</i>	79
<i>Myotis annectans</i>	76	<i>Rhinoceros sondaicus</i>	13
<i>Myotis chinensis</i>	68	<i>Rhinolophus marshalli</i>	37
<i>Myotis montivagus</i>	76	<i>Rhinolophus paradoxolophus</i>	68
<i>Naemorhedus caudatus</i>	24	<i>Rhinolophus robinsoni</i>	68
<i>Neofelis nebulosa</i>	64	<i>Rhinolophus rouxii</i>	83
<i>Neophocaena phocaenoides</i>	55	<i>Rhinolophus siamensis</i>	83
<i>Niviventer hinpoon</i>	28, 84	<i>Rhinopoma microphyllum</i>	67
<i>Nyctalus noctula</i>	69	<i>Rhinosciurus laticaudatus</i>	62
<i>Nycteris tragata</i>	67	<i>Rhizomys pruinosus</i>	80
<i>Orcaella brevirostris</i>	23	<i>Scotomanes ornatus</i>	69
<i>Orcinus orca</i>	47	<i>Semnopithecus germaini</i>	77
<i>Panthera pardus</i>	64	<i>Semnopithecus obscurus</i>	71
<i>Panthera tigris</i>	31	<i>Semnopithecus phayrei</i>	71
<i>Pardofelis marmorata</i>	30	<i>Soriculus parca</i>	82
<i>Peponocephala electra</i>	53	<i>Sousa chinensis</i>	74
<i>Petaurista alborufus</i>	80	<i>Sphaerias blanfordi</i>	67
<i>Petaurista elegans</i>	63	<i>Stenella attenuata</i>	50
<i>Petaurista petaurista</i>	80	<i>Stenella coeruleoalba</i>	51
<i>Petaurista philippensis</i>	80	<i>Stenella longirostris</i>	73
<i>Petinomys vordermanni</i>	62	<i>Steno bredanensis</i>	49
<i>Phoniscus atrox</i>	70	<i>Suncus malayanus</i>	82
<i>Physeter macrocephalus</i>	44	<i>Sundasciurus hippurus</i>	27
<i>Pipistrellus cadornae</i>	76	<i>Tadarida mops</i>	76
<i>Pipistrellus pulveratus</i>	76	<i>Tadarida teniotis</i>	68
<i>Presbytis femoralis</i>	71	<i>Taphozous saccolaimus</i>	67
<i>Prionailurus viverrinus</i>	64	<i>Tapirus indicus</i>	60
<i>Prionodon linsang</i>	64	<i>Tragulus napu</i>	56
<i>Prionodon pardicolor</i>	33	<i>Tursiops aduncus</i>	73
<i>Profelis temminckii</i>	64	<i>Ursus thibetanus</i>	65
<i>Pseudorca crassidens</i>	73	<i>Viverra megaspila</i>	32
<i>Pteromyscus pulverulentus</i>	82		



Sort by Common Name

กระจุยควาย	56	คางคางมงกุฎหูโตใหญ่	68
กระซู่	14	คางคางมงกุฎหูโตเล็ก	83
กระแตหางขนนก	73	คางคางมงกุฎอินเดีย	83
กระแตหางหนู	73	คางคางแม่ไก่เกาะ	75
กระทิง	74	คางคางแม่ไก่นครสวรรค์	82
กระรอกขางลายทองเทา	82	คางคางแม่ไก่ป่าฝน	67
กระรอกดินหลังลาย	62	คางคางแม่ไก่ภาคกลาง	76
กระรอกบินแกมสีแดง	62	คางคางยอคอกกล้วยป่า	70
กระรอกบินจิ้งจก	62	คางคางยอคอกกล้วยปึกป้อม	70
กระรอกบินเทาขน	63	คางคางยอคอกกล้วยปึกใส	70
กระรอกบินเล็กเขาสูง	62	คางคางยอคอกกล้วยเล็ก	70
กระรอกบินสีเขม่า	82	คางคางลูกหนูดำ	76
กระรอกสามสี	15	คางคางหนูยักษ์กุ่มภักกรรม	68
กระรอกหนากุระแต่	62	คางคางหนูยักษ์จุกป้อม	38, 84
กระรอกหางมาใหญ่	27	คางคางหนูยักษ์หมอนบุญสง	68
กวางผา	24	คางคางหนารอง	67
กูปรี	12	คางคางหางหนู	67
เก้งดำ	57	คางคางหูหนา	68
เก้งหม้อ	57	คางคางหูหนามลาย	76
ควายป่า	59	คางคางหูหนูดอยอย่างขาง	69
คางคางกรมหนามูนเล็ก	76	คางคางหูหนูพม่า	76
คางคากินแมลงนิวตัน	69	คางคางหูหนูยักษ์	68
คางคางขอบหูขาวเล็กป่า	83	คางคางหูหนูหนาน	76
คางคางขอบหูขาวใหญ่	81	คางคางอายุแหวง	68
คางคางขุนขาง	68	คางคางอายุแหวงเล็ก	83
คางคากุญทิด	20	คางคางอ้ออีไอ	69
คางคางจุกหลอดทองขาว	69	คางดำ	71
คางคางจุกหลอดหูขาว	77	คางแวนถิ่นใต้	71
คางคางคอย	67	คางแวนถิ่นเหนือ	71
คางคางคอยหลังลายขาว	69	คางหงอก	77
คางคางค้ายัค	67	ชะนีมงกุฎ	41
คางคางทองสีน้ำตาลสุราษฎร์	39, 84	ชะนีมือขาว	72
คางคางทองสีน้ำตาลหูหนา	69	ชะนีมือดำ	21
คางคางทองสีน้ำตาลใหญ่	69	ชะนีใหญ่	77
คางคางปากขนมลาย	76	ชะมดแปลงลายจุด	33
คางคาวปีกขนไต่	70	ชะมดแปลงลายแถบ	64
คางคาวปีกกล้วยป้อม	67	ชะมดแผงสันหางดำ	32
คางคาวปีกพับกลาง	83	ช่างป่า	61
คางคาวพื้นรอง	70	นากใหญ่ขนเรียบ	66
คางคาวพื้นหนาขนเล็ก	69	นากใหญ่จุกขน	19
คางคาวพื้นหนาขนใหญ่	69	นากใหญ่ภูเขา	35
คางคางมงกุฎเขินมลายูเล็ก	68	นากใหญ่หัวปลาคูก	19
คางคางมงกุฎหูโตมารแชด	37	เนื้อทราย	58

เนื้อสมัน	10, 84	วาพแกลบ	43
พญากระรอกดำ	79	วาพแกลบครีบบัวดำ	42
พญากระรอกบินหูขาว	80	วาพพิทตั้ง	43
พญากระรอกบินหูดำ	63	วาพนาร่องครีบลิ้น	52
พญากระรอกบินหูแดง	80	วาพบรูดา	43
พญากระรอกบินหูดำหางสีเข้ม	80	วาพเพชรฆาต	47
พญากระรอกเหลือง	62	วาพเพชรฆาตดำ	73
พะยูน	26	วาพเพชรฆาตเล็ก	54
เพียงพอนคอยผ้าหมูปก	82	วาพพื้นเขียว	22
เพียงพอนเล็กสีน้ำตาล	66	วาพพื้น	42
เพียงพอนเสนหลังขาว	36	วาพหัวทุย	44
เพียงพอนเหลือง	66	วาพหัวทุยแคระ	45
มัญญา	59	วาพหัวทุยเล็ก	46
เม่นหางพวง	81	สมเสร็จ	60
เม่นใหญ่	81	เสือกระต่าย	16
แมวป่าหัวแบน	17	เสือโคร่ง	31
แมวลายหินอ่อน	30	เสือดาว	64
ระมัด	13	เสือด้า	64
แรด	13	เสือปลา	64
ละมั่ง	11	เสือไฟ	64
ละอง	11	เสือดายเมฆ	64
ลิงกังใต้	77	หนูเกาะ	63
ลิงเสน	70	หนูขนสั้นเขานินปุ่น	28, 84
ลิงอายุเงาะ	40	หนูเขาสูง	63
ลิงจีน	29	หนูถ้ำ	63
ลิงชวา	75	หนูน้ำดอยอ่างกา	63
เลียงผา	78	หนูป่าพรุ	80
โลมากระโดด	73	หนูผีป่า	82
โลมาขาวเทา	74	หนูผีป่าหางจู้	67
โลมาแถบ	51	หนูผีภูเขา	82
โลมาปากขวด	73	หนูผีหางยาวพื้นแดง	82
โลมาปากขาว	48	หนูผีอ่างกา	75
โลมาเผือก	74	หนูไฟเล็บแม่มือแบน	63
โลมาเพชรฆาตแปลง	73	หนูฟันเล็ก	63
โลมาพื้นหาง	49	หนูภูหลวง	82
โลมาไรครีบล้าง	55	หนูไฟเล็บแม่มือแบน	80
โลมาลายจุด	50	หนูเหม็น	67
โลมาหลังโหนก	74	หมาจิ้งจอก	65
โลมาหัวกลม	52	หมาใน	65
โลมาหัวแดงโม	53	หมึกควาย	65
โลมาหัวบาตรครีบล้าง	23	หมึกหมา	66
โลมาหัวบาตรหลังเรียบ	55	อนกลาง	80
โลมาอริวดี	23	อิเห็นน้ำ	18
วัวแดง	25	อิเห็นลายพาด	34

## Reptile

### Sort by Scientific Name

<i>Acalyptophis peronii</i>	165	<i>Calliophis maculiceps</i>	113
<i>Acanthosaura armata</i>	115	<i>Calloselasma rhodostoma</i>	153
<i>Acanthosaura crucigera</i>	115	<i>Calotes emma alticristata</i>	116
<i>Acanthosaura lepidogaster</i>	115	<i>Calotes emma emma</i>	117
<i>Acrochordus granulatus</i>	135	<i>Calotes mystaceus</i>	117
<i>Acrochordus javanicus</i>	111	<i>Calotes versicolor</i>	117
<i>Aeluroscalabotes felinus</i>	157	<i>Cantoria violacea</i>	163
<i>Ahaetulla fasciolata</i>	160	<i>Caretta caretta</i>	157
<i>Ahaetulla mycterizans</i>	160	<i>Cerberus rynchops</i>	145
<i>Ahaetulla nasuta</i>	135	<i>Chelonia mydas</i>	89
<i>Ahaetulla prasina</i>	135	<i>Chitra chitra</i>	95
<i>Aipysurus eydouxii</i>	165	<i>Chitra vandijki</i>	96
<i>Amphiesma bitaeniatum</i>	163	<i>Chrysopelea ornata ornatissima</i>	138
<i>Amphiesma deschaunseei</i>	163, 174	<i>Chrysopelea paradisi paradisi</i>	111
<i>Amphiesma groundwateri</i>	163, 174	<i>Chrysopelea pelias</i>	112
<i>Amphiesma inas</i>	163	<i>Cnemaspis affinis</i>	120
<i>Amphiesma khasiense</i>	164	<i>Cnemaspis chanthaburiensis</i>	120, 168
<i>Amphiesma stolatum</i>	147	<i>Cnemaspis flavolineata</i>	120
<i>Amyda cartilaginea</i>	105	<i>Cnemaspis kandianus</i>	120
<i>Aphaniotis fusca</i>	116	<i>Cnemaspis kumpoli</i>	120
<i>Aplopeltura boa</i>	149	<i>Cnemaspis phuketensis</i>	157, 168
<i>Asthenodipsas laevis</i>	149	<i>Cnemaspis siamensis</i>	121
<i>Asthenodipsas malaccanus</i>	149	<i>Coelognathus flavolineatus</i>	138
<i>Astrotia stokesii</i>	165	<i>Coelognathus radiatus</i>	138
<i>Batagur baska</i>	93	<i>Cosymbotus craspedotus</i>	121
<i>Bitia hydroides</i>	162	<i>Cosymbotus platyurus</i>	121
<i>Boiga cyanea</i>	136	<i>Crocodylus porosus</i>	87
<i>Boiga cynodon</i>	136	<i>Crocodylus siamensis</i>	88
<i>Boiga dendrophila melanota</i>	136	<i>Cuora amboinensis</i>	104
<i>Boiga drapiezii</i>	137	<i>Cyclemys atripons</i>	104
<i>Boiga jaspidea</i>	137	<i>Cyclemys dentata</i>	104
<i>Boiga multomaculata</i>	137	<i>Cyclemys oldhami</i>	104
<i>Boiga nigriceps</i>	137	<i>Cyclemys shanensis</i>	104
<i>Boiga ocellata</i>	137	<i>Cylindrophis ruffus</i>	134
<i>Boiga sangsomi</i>	111, 173	<i>Cyrtodactylus angularis</i>	108, 168
<i>Bronchocela cristatella</i>	116	<i>Cyrtodactylus brevipalmatus</i>	108, 168
<i>Bronchocela smaragdina</i>	116	<i>Cyrtodactylus chanhomeae</i>	108, 168
<i>Bungarus candidus</i>	151	<i>Cyrtodactylus consobrinus</i>	121
<i>Bungarus fasciatus</i>	151	<i>Cyrtodactylus interdigitalis</i>	121
<i>Bungarus flaviceps flaviceps</i>	113	<i>Cyrtodactylus intermedius</i>	121
<i>Calamaria lumbricoidea</i>	161	<i>Cyrtodactylus jarujini</i>	108
<i>Calamaria pavimentata</i>	138	<i>Cyrtodactylus oldhami</i>	122
<i>Calamaria schlegelii</i>	161	<i>Cyrtodactylus papilionoides</i>	122
<i>Callagur borneensis</i>	94	<i>Cyrtodactylus peguensis peguensis</i>	122
<i>Calliophis bivirgatus flaviceps</i>	113	<i>Cyrtodactylus peguensis</i>	
<i>Calliophis gracilis</i>	164	<i>zebraicus</i>	122, 168
<i>Calliophis intestinalis lineata</i>	113	<i>Cyrtodactylus pulchellus</i>	122

<i>Cyrtodactylus quadrivirgatus</i>	123	<i>Gehyra fehlmanni</i>	124, 169
<i>Cyrtodactylus sumonthai</i>	108, 168	<i>Gehyra lacerata</i>	124, 169
<i>Cyrtodactylus thirakhupti</i>	157, 169	<i>Gehyra mutilata</i>	124
<i>Cyrtodactylus tigroides</i>	109, 169	<i>Gekko gecko</i>	124
<i>Cyrtodactylus variegatus</i>	123	<i>Gekko monarchus</i>	124
<i>Daboia russelii siamensis</i>	156	<i>Gekko petricolus</i>	124
<i>Dasia olivacea</i>	126	<i>Gekko siamensis</i>	109, 170
<i>Davewakeum miriamae</i>	109	<i>Gekko smithi</i>	125
<i>Dendrelaphis caudolineatus</i>	138	<i>Gerarda prevostiana</i>	163
<i>Dendrelaphis cyanochloris</i>	139	<i>Gongylosoma baliodeirus</i>	
<i>Dendrelaphis formosus</i>	139	<i>baliodeirus</i>	161
<i>Dendrelaphis ngansaoensis</i>	161	<i>Gongylosoma baliodeirus</i>	
<i>Dendrelaphis pictus</i>	139	<i>cochranae</i>	161, 174
<i>Dendrelaphis striatus</i>	161	<i>Gongylosoma longicauda</i>	161
<i>Dendrelaphis subocularis</i>	139	<i>Gongylosoma scriptus</i>	140
<i>Dermochelys coriacea</i>	92	<i>Gonocephalus abbotti</i>	106
<i>Dibamus alfredi</i>	157	<i>Gonocephalus bellii</i>	106
<i>Dibamus somsaki</i>	157	<i>Gonocephalus grandis</i>	106
<i>Dinodon septentrionale</i>	161	<i>Gonyosoma floweri</i>	162
<i>Dixoneus hangseesom</i>	158, 169	<i>Gonyosoma oxycephalum</i>	140
<i>Dixoneus melanostictus</i>	158, 169	<i>Gonyosoma prasina</i>	140
<i>Dixoneus siamensis</i>	123	<i>Hemidactylus frenatus</i>	125
<i>Dogania subplana</i>	107	<i>Hemidactylus garnotii</i>	125
<i>Draco blanfordi</i>	118	<i>Hemiphyllodactylus typus</i>	125
<i>Draco fimbriatus</i>	118	<i>Hemiphyllodactylus yunnanensis</i>	125
<i>Draco haematopogon</i>	118	<i>Heosemys grandis</i>	104
<i>Draco maculatus divergens</i>	118	<i>Heosemys spinosa</i>	104
<i>Draco maculatus haasei</i>	118	<i>Hieremys annandalei</i>	105
<i>Draco maculatus maculatus</i>	119	<i>Homalopsis buccata</i>	146
<i>Draco maculatus whiteheadi</i>	119	<i>Homalopsis nigroventralis</i>	163
<i>Draco melanopogon</i>	119	<i>Hydrophis atriceps</i>	165
<i>Draco obscurus formosus</i>	119	<i>Hydrophis belcheri</i>	165
<i>Draco quinquefasciatus</i>	119	<i>Hydrophis bituberculatus</i>	165
<i>Draco taeniopterus</i>	119	<i>Hydrophis brookii</i>	153
<i>Draco volans sumatranus</i>	120	<i>Hydrophis caeruleus</i>	165
<i>Dryocalamus davisonii</i>	139	<i>Hydrophis cantoris</i>	166
<i>Dryocalamus subannulatus</i>	139	<i>Hydrophis cyanocinctus</i>	166
<i>Dryocalamus thungsongensis</i>	112, 174	<i>Hydrophis fasciatus</i>	166
<i>Dryophiops rubescens</i>	139	<i>Hydrophis gracilis</i>	166
<i>Enhydrina schistosa</i>	165	<i>Hydrophis klossi</i>	166
<i>Enhydria bocourti</i>	145	<i>Hydrophis lamberti</i>	166
<i>Enhydria enhydria</i>	145	<i>Hydrophis lapemoides</i>	166
<i>Enhydria jagorii</i>	163	<i>Hydrophis melanosoma</i>	166
<i>Enhydria plumbea</i>	145	<i>Hydrophis ornatus ornatus</i>	166
<i>Enhydria subtaeniata</i>	146	<i>Hydrophis spiralis</i>	167
<i>Eretmochelys imbricata</i>	90	<i>Hydrophis torquatus aagaardi</i>	167
<i>Erpeton tentaculatum</i>	146	<i>Hydrophis torquatus diadema</i>	167
<i>Eumeces quadrilineatus</i>	109	<i>Indotestudo elongata</i>	99
<i>Fordonia leucobalia</i>	163	<i>Isopachys anguinoides</i>	127, 170
<i>Gehyra angusticaudata</i>	123, 169	<i>Isopachys borealis</i>	109

<i>Isopachys gyldenstolpei</i>	109, 170	<i>Naja siamensis</i>	152
<i>Isopachys roulei</i>	109, 170	<i>Naja sumatrana</i>	152
<i>Kerilia jerdonii jerdonii</i>	167	<i>Notochelys platynota</i>	105
<i>Kerilia jerdonii siamensis</i>	167	<i>Oligodon barroni</i>	142
<i>Kolpophis annandalei</i>	167	<i>Oligodon catenatus</i>	142
<i>Lapemis curtus</i>	167	<i>Oligodon cinereus</i>	142
<i>Laticauda colubrina</i>	164	<i>Oligodon dorsolateralis</i>	162
<i>Laticauda laticaudata</i>	164	<i>Oligodon fasciolatus</i>	142
<i>Leiolepis belliana belliana</i>	107	<i>Oligodon inornatus</i>	142
<i>Leiolepis belliana ocellata</i>	108	<i>Oligodon joynsoni</i>	142
<i>Leiolepis bohmei</i>	107	<i>Oligodon mouhoti</i>	162
<i>Leiolepis reevesi rubritaeniata</i>	108	<i>Oligodon purpurascens</i>	142
<i>Lepidochelys olivacea</i>	91	<i>Oligodon taeniatus</i>	143
<i>Leptoceps osellai</i>	109, 170	<i>Ophiophagus hannah</i>	152
<i>Lepturophis albofuscus</i>	162	<i>Ophisaurus gracilis</i>	108
<i>Liopeltis tricolor</i>	162	<i>Opisthotropis praemaxillaris</i>	112
<i>Lipinia quadrivittata</i>	158	<i>Opisthotropis spenceri</i>	113
<i>Lipinia surda</i>	158	<i>Oreophis porphyraceus coxi</i>	113, 174
<i>Lipinia vittigera</i>	127	<i>Oreophis porphyraceus porphyraceus</i>	112
<i>Lycodon butleri</i>	162	<i>Orthriophis taeniura ridleyi</i>	143
<i>Lycodon cadamomensis</i>	112	<i>Orthriophis taeniura yunnanensis</i>	143
<i>Lycodon capucinus</i>	140	<i>Ovophis monticola convictus</i>	155
<i>Lycodon effraensis</i>	141	<i>Parahelicops boonsongi</i>	113
<i>Lycodon fasciatus</i>	141	<i>Pareas carinatus</i>	149
<i>Lycodon laotus</i>	141	<i>Pareas hamptoni</i>	164
<i>Lycodon subcinctus</i>	141	<i>Pareas macularius</i>	150
<i>Lygosoma anguinum</i>	158	<i>Pareas margaritophorus</i>	150
<i>Lygosoma corpulentum</i>	110	<i>Pelamis platura</i>	153
<i>Lygosoma haroldyoungi</i>	110, 170	<i>Pelochelys cantorii</i>	97
<i>Lygosoma isodactylum</i>	110	<i>Physignathus cocincinus</i>	103
<i>Lygosoma koratense</i>	110, 170	<i>Plagiopholis nuchalis</i>	150
<i>Lygosoma quadrupes</i>	127	<i>Platysternon megacephalum</i>	98
<i>Mabuya longicaudata</i>	128	<i>Psammodynastes pulverulentus</i>	143
<i>Mabuya macularia malcolmi</i>	128, 171	<i>Psammophis indochinensis</i>	150
<i>Mabuya macularia postnasalis</i>	158, 171	<i>Pseudocalotes floweri</i>	107
<i>Mabuya macularia quadrifasciata</i>	128, 171	<i>Pseudocalotes microlepis</i>	107
<i>Mabuya multifasciata</i>	128	<i>Pseudorhabdion longiceps</i>	112
<i>Mabuya novemcarinata</i>	128	<i>Pseudoxenodon macrops</i>	150
<i>Mabuya rugifera</i>	129	<i>Ptyas carinatus</i>	144
<i>Macrocalamus lateralis</i>	162	<i>Ptyas fuscus</i>	112
<i>Macropisthodon rhodomelas</i>	164	<i>Ptyas korros</i>	144
<i>Macropisthodon flaviceps</i>	164	<i>Ptyas mucosus</i>	144
<i>Malayemys subtrijuga</i>	105	<i>Ptychozoon horsfieldi</i>	158
<i>Manouria emys emys</i>	100	<i>Ptychozoon kuhli</i>	125
<i>Manouria emys phayrei</i>	101	<i>Ptychozoon lionatum</i>	126
<i>Manouria impressa</i>	102	<i>Ptychozoon trinotatera</i>	158
<i>Mantheyus phuwuaensis</i>	106	<i>Python brongersmai</i>	134
<i>Naja kaouthia</i>	151	<i>Python molurus bivittatus</i>	135
		<i>Python reticulatus</i>	135

<i>Pyxidea mouhoti</i>	157	<i>Trimeresurus (Cryptelytrops)</i>	
<i>Ramphotyphlops albiceps</i>	132	<i>macrops</i>	154
<i>Ramphotyphlops braminus</i>	132	<i>Trimeresurus (Cryptelytrops)</i>	
<i>Ramphotyphlops lineatus</i>	160	<i>purpureomaculatus</i>	154
<i>Rhabdophis chrysargos</i>	147	<i>Trimeresurus (Cryptelytrops)</i>	
<i>Rhabdophis nigrocinctus</i>	147	<i>venustus</i>	154
<i>Rhabdophis subminiatus helleri</i>	164	<i>Trimeresurus (Parias) hageni</i>	155
<i>Rhabdophis subminiatus</i>		<i>Trimeresurus (Parias) sumatranus</i>	114
<i>subminiatus</i>	148	<i>Trimeresurus (Popeia) fugatus</i>	155
<i>Riopa bowringi</i>	129	<i>Trimeresurus (Popeia) nebularis</i>	155
<i>Riopa herberti</i>	129, 171	<i>Trimeresurus (Popeia) popeiorum</i>	155
<i>Scincella doriae</i>	129	<i>Trimeresurus (Trimeresurus)</i>	
<i>Scincella melanosticta</i>		<i>borneensis</i>	114
<i>kohtaoensis</i>	110, 171	<i>Trimeresurus (Viridovipera)</i>	
<i>Scincella melanosticta</i>		<i>gumprechtii</i>	155
<i>melanosticta</i>	129	<i>Trimeresurus (Viridovipera) vogeli</i>	155
<i>Scincella reevesi</i>	130	<i>Tropidolaemus wagleri</i>	114
<i>Scincella tavesae</i>	130, 172	<i>Tropidophorus berdmorei</i>	131
<i>Sibynophis collaris</i>	144	<i>Tropidophorus laotus</i>	131
<i>Sibynophis melanocephalus</i>	144	<i>Tropidophorus latiscutatus</i>	160, 172
<i>Sibynophis triangularis</i>	144, 174	<i>Tropidophorus matsuii</i>	160, 172
<i>Siebenrockiella crassicollis</i>	105	<i>Tropidophorus microlepis</i>	132
<i>Sinomicrurus macclelandi</i>		<i>Tropidophorus robinsoni</i>	132
<i>maclelandi</i>	114	<i>Tropidophorus thai</i>	132, 173
<i>Sinonatrix percarinata</i>	148	<i>Typhlops diardi</i>	133
<i>Sphenomorphus butleri</i>	159	<i>Typhlops floweri</i>	133, 173
<i>Sphenomorphus grandisonae</i>	159, 172	<i>Typhlops khoratensis</i>	133, 173
<i>Sphenomorphus helenae</i>	159, 172	<i>Typhlops muelleri</i>	133
<i>Sphenomorphus indicus indicus</i>	130	<i>Typhlops ozakiae</i>	133, 173
<i>Sphenomorphus</i>		<i>Typhlops porrectus</i>	133
<i>lineopunctulatus</i>	159, 172	<i>Typhlops roxanae</i>	160, 173
<i>Sphenomorphus maculatus</i>	130	<i>Typhlops siamensis</i>	160, 173
<i>Sphenomorphus mimicus</i>	159, 172	<i>Typhlops trangensis</i>	160, 173
<i>Sphenomorphus presignis</i>	159	<i>Typhlops ozakiae</i>	133
<i>Sphenomorphus scotophilus</i>	159	<i>Varanus dumerilii</i>	110
<i>Sphenomorphus stellatus</i>	159	<i>Varanus nebulosus</i>	111
<i>Sphenomorphus tersus</i>	131, 172	<i>Varanus rudicollis</i>	111
<i>Stegonotus borneensis</i>	162	<i>Varanus salvator</i>	132
<i>Takydromus sexlineatus ocellatus</i>	126	<i>Xenelaphis hexagonotus</i>	144
<i>Thalassophina viperina</i>	167	<i>Xenochrophis flavipunctatus</i>	148
<i>Thalassophis anomalus</i>	167	<i>Xenochrophis piscator</i>	148
<i>Tomistoma schlegelii</i>	86	<i>Xenochrophis punctulatus</i>	149
<i>Trimeresurus (Cryptelytrops)</i>		<i>Xenochrophis trianguligerus</i>	149
<i>albolabris albolabris</i>	153	<i>Xenodermus javanicus</i>	164
<i>Trimeresurus (Cryptelytrops)</i>		<i>Xenopeltis unicolor</i>	134
<i>kanburiensis</i>	154, 174		

Sort by Common Name

กิ่งก้านแก้วไต้	117	งูคาบหมากมลายู	162
กิ่งก้านแก้วเหนือ	116	งูคาบหมากยูนนาน	143
กิ่งก้านเขาลี้ก	115	งูกินทากเกล็ดเรียบ	149
กิ่งก้านเขาสูง	107	งูกินทากเกล็ดสัน	149
กิ่งก้านเขาหนามยาว	115	งูกินทากจุดขาว	150
กิ่งก้านเขาหนามสั้น	115	งูกินทากจุดดำ	150
กิ่งก้านเขียวจันทบูรณ	116	งูกินทากมลายู	149
กิ่งก้านเขียวไต้	116	งูกินทากลายขี้วัน	164
กิ่งก้านงู	108	งูกินทากหัวโหนด	149
กิ่งก้านดงคอฟ้า	106	งูเขียวคาบหมาก	140
กิ่งก้านดงคว้าน	106	งูเขียวจิ้งจกมลายู	160
กิ่งก้านดงใหญ่	106	งูเขียวดงลาย	111, 173
กิ่งก้านนอยหางยาว	126	งูเขียวดอกหมาก	138
กิ่งก้านมินคอดำ	119	งูเขียวตุ๊กแก	114
กิ่งก้านมินคอดแดง	118	งูเขียวบอน	136
กิ่งก้านมินคอฟ้า	120	งูเขียวปากแหนบ	135
กิ่งก้านมินคอส้ม	118	งูเขียวไผ่	155
กิ่งก้านมินปิกแดง	118	งูเขียวร้อน	111
กิ่งก้านมินปิกลาย	119	งูเขียวหัวจิ้งจกป่า	135
กิ่งก้านมินปิกส้มจันทบูรณ	118	งูเขียวหัวลายกระ	160
กิ่งก้านมินปิกส้มดอยสุเทพ	118	งูเขียวหางเทา	155
กิ่งก้านมินปิกส้มมลายู	119	งูเขียวหางไหม้ตาโต	154
กิ่งก้านมินปิกส้มไหหลำ	119	งูเขียวหางไหม้ทองเขียว	155
กิ่งก้านมินปิกหาแถบ	119	งูเขียวหางไหม้ทองเขียวโต	155
กิ่งก้านมินมลายู	119	งูเขียวหางไหม้ทองเหลือง	153
กิ่งก้านพระอินทร	116	งูเขียวหางไหม้ทุ่งสง	154
กิ่งก้านภูว	106	งูเขียวหางไหม้สุมาตรา	114
กิ่งก้านรว	117	งูเขียวหางไหม้ฮาเจน	155
กิ่งก้านสวน	117	งูควนขนุน	144
กิ่งก้านหัวแดง	117	งูคอขี้วันดำ	144
กิ่งก้านหัวฟ้า	117	งูคอขี้วันหัวดำ	144
กิ่งก้านหัวยาว	107	งูคอขี้วันหัวลายสามเหลี่ยม	144, 174
งูกนขบ	134	งูคออ่อนปากจะงอย	165
งูกระ	137	งูคออ่อนหัวเขี้ยว	166
งูกระดาง	146	งูกุด	142
งูกะปะ	153	งูวงช้าง	111
งูคาบหมากดำ	143	งูอดเขมร	162

งูจอกไทย	143	งูปล้องฉนวนบ้าน	141
งูจงอาง	152	งูปล้องฉนวนภูเขา	161
งูชายธงขาวหลามดัด	167	งูปล้องฉนวนมลายู	139, 162
งูชายธงทองขาว	167	งูปล้องฉนวนเมืองเหนือ	141
งูชายธงขาว	165	งูปล้องฉนวนลาว	141
งูชายธงสยาม	167	งูปล้องฉนวนอินเดีย	139
งูชายธงหลังดำ	153	งูปล้องทอง	136
งูชายธงหัวโต	167	งูปล้องหวายมลายู	164
งูไซ	145	งูปล้องหวายลายขวันดำ	114
งูดงคา	137	งูปล้องหวายหัวดำ	113
งูดอกหมากแดง	112	งูปลาตาแมว	163
งูดินกรุงเทพฯ	160, 173	งูปลาหัวเทา	163
งูดินโคราช	133, 173	งูปลิง	145
งูดินคอยปุย	133	งูปากกว้างทองสัน	162
งูดินตรัง	160, 173	งูปากกว้างน้ำเค็ม	145
งูดินบ้าน	132	งูปากกว้างลาย	163
งูดินบึงกระชาย	133, 173	งูปาลม	114
งูดินลายซิด	160	งูปีแก้วกุหลวง	142
งูดินสยาม	160, 173	งูปีแก้วลายกระ	142
งูดินหัวขาว	132	งูปีแก้วลายซิด	162
งูดินหัวเหลือง	133, 173	งูปีแก้วลายเต็ม	142
งูดินใหญ่มลายู	133	งูปีแก้วลายหัวใจ	142
งูดินใหญ่อินโดจีน	133	งูปีแก้วสีจาง	142
งูแดงบอร์เนียว	162	งูปีแก้วใหญ่	142
งูคองไฟ	135	งูผาขี้รี	135
งูเถา	139	งูฝักมะรุม	166
งูทองขาว	164	งูพงอทองเหลือง	138
งูทับสมิงคลา	151	งูพงอภูเขา	162
งูทากลาย	165	งูพงอหลากลาย	161
งูทากลายทองขาว	165	งูพงอหัวแดง	161
งูทางมะพร้าวเขียว	140	งูพงอหัวขาว	112
งูทางมะพร้าวดำ	138	งูพริกทองแดง	113
งูทางมะพร้าวแดงกุหลวง	112, 174	งูพริกสีน้ำตาล	113
งูทางมะพร้าวแดงลายขวัน	112	งูพังกา	154
งูทางมะพร้าวลายซิด	138	งูมานทอง	150
งูปล้องฉนวนจันทบุรี	112	งูแมะตะจาวรังนก	137
งูปล้องฉนวนแดง	141	งูแมวเซา	156
งูปล้องฉนวนทุ่งสง	112, 162, 174	งูริ้งเหหลังศร	164



รู้งิ้วแห้วแดง	164	รูสายรุ้งลาย	146
รูลายสอจีน	148	รูสิงทอง	112
รูลายสอพมา	149	รูสิงบาน	144
รูลายสอลายสามเหลี่ยม	149	รูสิงหางดำ	144
รูลายสอลาวเหนือ	112	รูสิงหางลาย	144
รูลายสอสวน	148	รูเสมียนรังหัวสั้น	167
รูลายสอสองสี	113	รูแสงอาทิตย์	134
รูลายสอหมอบุญสง	113	รูเสมวังเกิ้ลด์เบลเซอร์	165
รูลายสอใหญ่	148	รูเสมวังเกิ้ลด์สองตุ้ม	165
รูลายสาบเขาสูง	164	รูเสมวังแคนเทอร์	166
รูลายสาบเขี้ยววันดำ	147	รูเสมวังดำ	166
รูลายสาบคอแดง	148	รูเสมวังทองเหลือง	153
รูลายสาบจุดดำขาว	147	รูเสมวังเทา	167
รูลายสาบคอกหนู	147	รูเสมวังมลายู	167
รูลายสาบคาโต	150	รูเสมวังลายเยื้อง	165
รูลายสาบทองสามขีด	163, 174	รูเสมวังลายเหลือง	167
รูลายสาบท่าสาร	163, 174	รูเสมวังแลมเบิร์ต	166
รูลายสาบมลายู	163	รูเสมวังหัวดำ	165, 166
รูลายสาบขุนนาน	163, 164	รูเสมวังหางขาว	166
รูสมิงทะเลปากดำ	164	รูเสมวังเหลืองลายคราม	166
รูสมิงทะเลเหลือง	164	รูเสมวังอ่าวเปอร์เซีย	166
รูสรอยเหลือง	140	รูเสมหางเทา	137
รูสามเหลี่ยม	151	รูเสมหางมา	136
รูสามเหลี่ยมหัวแดง	113	รูหมอก	143
รูสายทองคอแหวน	140	รูหลาม	135
รูสายทองจันทบุรี	161, 174	รูหลามปากเป็ด	134
รูสายทองมลายู	162	รูหวักะโหลก	146
รูสายทองลายแถบ	161	รูหวักะโหลกทองคำ	163
รูสายทองหางยาว	161	รูหัวลายศร	150
รูสายม่านเกิ้ลด์ไตดาใหญ่	139	รูหางแห้มกาญจน์	154, 174
รูสายม่านพระอินทร์	139	รูหางแห้มภูเขา	155
รูสายม่านฟ้าเขียว	139	รูเหลือง	135
รูสายม่านลายเฉียง	161	รูเห่าสยาม	152
รูสายม่านหลังทอง	139	รูเห่าสุมาตรา	152
รูสายม่านหลังลาย	138	รูเห่าหมอ	151
รูสายม่านอันนัม	161	รูอัยยั้ว	167
รูสายรุ้ง	145	จรเข้หน้าเค็ม	87
รูสายรุ้งดำ	163	จรเข้หน้าจืด	88

จิงจกเขาสูงมลายู	125	จิงเหลนภูเขาจุดดาว	159
จิงจกเขาสูงขุนนาน	125	จิงเหลนภูเขาหลาย	159
จิงจกดินขวางคำ	158, 169	จิงเหลนภูเขาหลายกระ	159
จิงจกดินลายจุด	123	จิงเหลนภูเขาหลายแฉก	159
จิงจกดินหางสี่สม	158, 169	จิงเหลนภูเขาเล็ก	159, 172
จิงจกนิ้วยาวกัมพล	120	จิงเหลนภูเขาสามนิ้ว	109, 170
จิงจกนิ้วยาวจันทบุรี	120, 168	จิงเหลนภูเขาสี่ข้าง	131, 172
จิงจกนิ้วยาวไทย	121	จิงเหลนภูเขาอินเดีย	130
จิงจกนิ้วยาวภูเก็ต	157, 168	จิงเหลนภูเขาอุบล	159, 172
จิงจกนิ้วยาวมลายู	120	จิงเหลนภูเขาเฮเลน	159, 172
จิงจกนิ้วยาวหลังซิด	120	จิงเหลนเรียวกาเล็ก	127
จิงจกนิ้วยาวอินเดีย	120	จิงเหลนเรียวกโรราช	110, 170
จิงจกบาน	121	จิงเหลนเรียวกจันทบุรี	110
จิงจกหางแบนมลายู	121	จิงเหลนเรียวกไต	129, 171
จิงจกหางเรียบ	125	จิงเหลนเรียวกทองเหลือง	129
จิงจกหางหนาม	125	จิงเหลนเรียวกพมา	158
จิงจกหินเมืองกาญจน	124, 169	จิงเหลนเรียวกภาคกลาง	110
จิงจกหินลายกระ	124, 169	จิงเหลนเรียวกลา	110, 170
จิงจกหินสี่ข้าง	124	จิงเหลนลายซิด	129
จิงจกหินหางเรียวก	123, 169	จิงเหลนลายมลายู	158
จิงเหลนเกล็ดสัน	128	จิงเหลนลายสี่ซิด	158
จิงเหลนควงเขาซอชดาว	157	จิงเหลนลายอิน โดจีน	127
จิงเหลนควงชลบุรี	109, 170	จิงเหลนหลากลายดานชาย	158, 171
จิงเหลนควงประจวบ	127, 170	จิงเหลนหลากลายมัลคอล์ม	128, 171
จิงเหลนควงปักธงชัย	109	จิงเหลนหลากลายสี่ซิด	128, 171
จิงเหลนควงป่าตะวันตก	109	จิงเหลนห้วยชุมพร	132
จิงเหลนควงลา	109, 170	จิงเหลนห้วยทองแดง	131
จิงเหลนควงสี่มวง	157	จิงเหลนห้วยไทย	132, 173
จิงเหลนดินเกาะเต่า	110, 171	จิงเหลนห้วยภูผาน้ำทิพย์	160, 172
จิงเหลนดินจุดคำ	129	จิงเหลนห้วยภูว	160, 172
จิงเหลนดินคองพญาเย็น	130	จิงเหลนห้วยลาว	131
จิงเหลนดินคอร์เรียวก	129	จิงเหลนห้วยเล็ก	132
จิงเหลนดินเมืองกาญจน	130, 172	จิงเหลนหางยาว	128
จิงเหลนต้นไม้	126	จิงเหลนหางสี่ฟ้า	109
จิงเหลนบาน	128	ตะกวด	111
จิงเหลนภูเขาเกล็ดเรียบ	130	ตะกอง	103
จิงเหลนภูเขาแกรนดิสัน	159, 172	ตะโขง	86

ตะพานคำ	107	ตุ๊กแกหลังจุดคู่	124
ตะพานธรรมดา	105	ตุ๊กแกหัวโต	157
ตะพานมานลาย	95	ตุ๊กตุ๊ก	110
ตะพานมานลายพม่า	96	เตากระ	90
ตะพานหัวกบ	97	เตากระอัน	93
ตุ๊กแกเขาคินทราย	124	เตาจักร	104
ตุ๊กแกบาน	124	เตาจีน	157
ตุ๊กแกบินลายสามแถบ	158	เตาคำ	105
ตุ๊กแกบินหางแบน	125	เตาเค็ย	102
ตุ๊กแกบินหางเพิน	158	เตาคนุ	89
ตุ๊กแกบินหางหยัก	126	เตาทับทิม	105
ตุ๊กแกปากำธร	157, 169	เตานา	105
ตุ๊กแกปากเขาชะเมา	108, 168	เตาบัว	105
ตุ๊กแกปากอชัน	122	เตาใบไม้	104
ตุ๊กแกปากโคนนิ้วคิด	108, 168	เตาใบไม้พม่า	104
ตุ๊กแกปากจารุจินต์	108	เตาปลู	98
ตุ๊กแกปากงพญาเย็น	108, 168	เตามะเฟือง	92
ตุ๊กแกปากไทรโยค	109, 169	เตาลายดินเปิด	94
ตุ๊กแกปากน้ำหนาว	121	เตาหกคำ	101
ตุ๊กแกปากพม่า	123	เตาหกเหลี่ยม	100
ตุ๊กแกปากมลายู	121	เตาหญา	91
ตุ๊กแกปากลายจุดโต	122, 168	เตาห้วยคูลอย	104
ตุ๊กแกปากลายจุดสี่เสื่อ	122	เตาห้วยทองเหลี่ยม	104
ตุ๊กแกปากลายจุดเหนือ	122	เตาห้วย	104
ตุ๊กแกปากลายพาดตะวันออก	121	เตาหีบ	104
ตุ๊กแกปากลายพาดป่าไค้	122	เตาหัวโต	157
ตุ๊กแกปากลายสี่ซึก	123	เตาเหลี่ยม	99
ตุ๊กแกปากสุราษฎร์	157, 169	แย้โต	107
ตุ๊กแกปากสระบุรี	108, 168	แย้สงขลา	107
ตุ๊กแกสยาม	109, 170	แย้เหนือ	108
ตุ๊กแกสีเทา	125	แย้ฮิสาน	108
		ลั้ง	103
		เห่าซาง	111
		เหี้ย	132

## Amphibian

### Sort by Scientific Name

<i>Amolops larutensis</i>	192	<i>Leptobrachium hendricksoni</i>	187
<i>Amolops marmoratus</i>	193	<i>Leptobrachium smithi</i>	187
<i>Ansonia inthanon</i>	182, 206	<i>Leptolalax bourreti</i>	202
<i>Ansonia malayana</i>	188	<i>Leptolalax cf. gracilis</i>	202
<i>Ansonia siamensis</i>	182, 206	<i>Leptolalax heteropus</i>	187
<i>Bourretia kohchangae</i>	183	<i>Leptolalax pelodytoides</i>	187
<i>Brachytarsophrys carinense</i>	181	<i>Leptophryne borbonica</i>	189
<i>Brachytarsophrys feae</i>	202	<i>Limnonectes blythii</i>	183
<i>Bufo asper</i>	188	<i>Limnonectes doriae</i>	203
<i>Bufo divergens</i>	188	<i>Limnonectes gyldenstolpei</i>	193
<i>Bufo macrotis</i>	188	<i>Limnonectes kuhlii</i>	184
<i>Bufo melanostictus</i>	189	<i>Limnonectes laticeps</i>	194
<i>Bufo parvus</i>	189	<i>Limnonectes macrognathus</i>	193
<i>Calluella guttulata</i>	189	<i>Limnonectes malesianus</i>	184
<i>Caudacaecilia asplenia</i>	201	<i>Limnonectes paramacrodon</i>	184
<i>Caudacaecilia larutensis</i>	201	<i>Limnonectes plicatellus</i>	194
<i>Chaparana aenea</i>	203	<i>Megophrys nasuta</i>	181
<i>Chaperina cf. fusca</i>	202	<i>Microhyla annamensis</i>	202
<i>Chirixalus doriae</i>	197	<i>Microhyla annectens</i>	203
<i>Chirixalus hansenae</i>	205	<i>Microhyla berdmorei</i>	190
<i>Chirixalus nongkhorensis</i>	197	<i>Microhyla borneensis</i>	203
<i>Chirixalus vittatus</i>	198	<i>Microhyla butleri</i>	191
<i>Fejervarya cancrivora</i>	194	<i>Microhyla heymonsii</i>	191
<i>Fejervarya limnocharis</i>	203	<i>Microhyla ornata</i>	191
<i>Fejervarya raja</i>	203	<i>Microhyla pulchra</i>	192
<i>Glyphoglossus molossus</i>	182	<i>Micryletta inornata inornata</i>	192
<i>Hoplobatrachus chinensis</i>	194	<i>Micryletta inornata lineata</i>	192
<i>Huia melasma</i>	203, 207	<i>Nyctixalus pictus</i>	198
<i>Hyla annectans</i>	202	<i>Occidozyga laevis</i>	194
<i>Ichthyophis acuminatus</i>	201, 206	<i>Occidozyga lima</i>	194
<i>Ichthyophis kohtaoensis</i>	201, 206	<i>Occidozyga magnapustulosa</i>	194
<i>Ichthyophis supachaii</i>	201	<i>Occidozyga martensii</i>	195
<i>Ichthyophis youngorum</i>	201, 206	<i>Ophryophryne microstoma</i>	181
<i>Ingerana tasanae</i>	183, 207	<i>Paa bourreti</i>	176
<i>Ingerana tenasserimensis</i>	183	<i>Paa fasciculispina</i>	177
<i>Kalophrynus interlineatus</i>	190	<i>Pedostibes hosii</i>	189
<i>Kalophrynus pleurostigma</i>	190	<i>Philautus carinensis</i>	205
<i>Kaloula baleata</i>	190	<i>Philautus gracilipes</i>	205
<i>Kaloula mediolineata</i>	183	<i>Philautus parvulus</i>	198
<i>Kaloula pulchra</i>	190	<i>Philautus petersi</i>	198
<i>Leptobrachium chapaense</i>	202	<i>Philautus vermiculatus</i>	205

<i>Phrynella pulchra</i>	203	<i>Rana schmackeri</i>	204
<i>Polypedates colleti</i>	198	<i>Rana signata</i>	185
<i>Polypedates leucomystax</i>	198	<i>Rana taipehensis</i>	197
<i>Polypedates macrotis</i>	199	<i>Rhacophorus appendiculatus</i>	199
<i>Polypedates mutus</i>	199	<i>Rhacophorus bipunctatus</i>	199
<i>Rana archotaphus</i>	195	<i>Rhacophorus bisacculus</i>	199
<i>Rana banjarana</i>	204	<i>Rhacophorus cyanopunctatus</i>	200
<i>Rana baramica</i>	195	<i>Rhacophorus feae</i>	186
<i>Rana cf. alticola</i>	195	<i>Rhacophorus maximus</i>	186
<i>Rana chapaensis</i>	185	<i>Rhacophorus nigropalmatus</i>	186
<i>Rana chloronota</i>	204	<i>Rhacophorus orlovi</i>	205
<i>Rana cubitalis</i>	195	<i>Rhacophorus pardalis</i>	186
<i>Rana erythraea</i>	195	<i>Rhacophorus prominans</i>	186
<i>Rana faber</i>	185	<i>Rhacophorus reinwardtii</i>	186
<i>Rana glandulosa</i>	196	<i>Rhacophorus robinsoni</i>	186
<i>Rana hosii</i>	196	<i>Rhacophorus verrucosus</i>	205
<i>Rana johnsi</i>	196	<i>Taylorana hascheana</i>	204
<i>Rana lateralis</i>	185	<i>Taylorana limborgii</i>	205
<i>Rana leptoglossa</i>	185	<i>Theloderma asperum</i>	200
<i>Rana livida</i>	196	<i>Theloderma gordonii</i>	178
<i>Rana luctuosa</i>	204	<i>Theloderma horridum</i>	179
<i>Rana macrodactyla</i>	196	<i>Theloderma stellatum</i>	180
<i>Rana milleti</i>	204	<i>Tylototriton verrucosus</i>	181
<i>Rana miopus</i>	185	<i>Xenophrys parva</i>	182
<i>Rana mortenseni</i>	204	<i>Xenophrys aceras</i>	181
<i>Rana nicobariensis</i>	196	<i>Xenophrys longipes</i>	182
<i>Rana nigrovittata</i>	197	<i>Xenophrys major</i>	182
<i>Rana raniceps</i>	197	<i>Xenophrys minor</i>	182

Sort by Common Name

กบกา	205	กบหลังจุดมลายู	204
กบเกาะช้าง	183	กบหลังคาพับ	194
กบเขาทองลาย	204	กบหลังไพล	185
กบเขาสอยดาว	185	กบหลังลายเขียว	185
กบเขาสูง	195	กบห้วยข่าม	184
กบเขาหลังคอง	197	กบหัวโต	193
กบเขาใหญ่	204	กบหูดำ	195
กบชะง่อนผาใต้	196	กบอกหนามจันทบุรี	177
กบชะง่อนผาไทย	203, 207	กบอกหนามเหนือ	176
กบชะง่อนผาเหนือ	196	กบอ่องเล็ก	197
กบชะง่อนผาสัตม	204	กบอ่องใหญ่	204
กบชะง่อนผาอินทนนท์	195	กะทาง	181
กบคอยช้าง	203	เขียดงูเกาะเต่า	201, 206
กบคอยอินทนนท์	204	เขียดงูคอยสุเทพ	201, 206
กบคอร์เรีย	203	เขียดงูดำ	201
กบตะนาวศรี	183	เขียดงูบลา	201
กบใต้ห้วย	197	เขียดงูมลายู	201
กบท่าสาร	183, 207	เขียดงูสุกซัย	201
กบทุด	183	เขียดงูหัวแหลม	201, 206
กบทุดมลายู	184	เขียดจะนา	194
กบนา	194	เขียดจิก	195
กบน้ำเค็ม	194	เขียดตะปาด	199
กบนิโคบาร์	196	เขียดทราย	195
กบป่าไผ่	204	เขียดลิ้น	194
กบป่าพรุ	184	เขียดแลว	183
กบภูหลวง	185	เขียดหลังป้อม	194
กบมัน	184	เขียดเหลืออง	185
กบทราย	203	คางคกขาขาว	189
กบแรด	194	คางคกแคะมะลายู	188
กบลายหินเมืองใต้	192	คางคกแคะอินโดจีน	189
กบลายหินเมืองเหนือ	193	คางคกตุนไม้	189
กบวากเล็ก	195	คางคกบาน	189
กบวากใหญ่	196	คางคกห้วยไทย	182, 206
กบสังขละ	185	คางคกห้วยมลายู	188
กบสีชมพู	196	คางคกห้วยอินทนนท์	182, 206
กบหนอง	193	คางคกหัวราบ	188
กบหนอง	203	จิ้งโคร่ง	188
กบหลังขี้ด	196	ปาดขาลาย	186
กบหลังจุด	185	ปาดเขี้ยวดินดำ	186

ปาดเขียวดินแดง	186	อิงกรายทองผาภูมิ	202
ปาดเขียวดินลาย	186	อิงกรายปากแคบ	181
ปาดแคระขาเขียว	205	อิงกรายพม่า	202
ปาดแคระทองลาย	205	อิงกรายลายจุด	187
ปาดแคระทองเหลือง	198	อิงกรายลายละเอียด	187
ปาดแคระป่า	198	อิงกรายเล็ก	182
ปาดแคระพม่า	205	อิงกรายหนังปุ่ม	187
ปาดจิวตะวันออก	205	อิงกรายหลากลาย	187
ปาดจิวพม่า	198	อิงกรายห้วยแคระ	182
ปาดจิวลายเต็ม	197	อิงกรายห้วยใหญ่	182
ปาดจิวหลังซัด	197	อิงกรายห้วยมน	181
ปาดจุดฟ้า	200	อิงกรายหัวแหลม	181
ปาดคอยอินทนนท์	186	อิงขาคำ	192
ปาดตะปุมจันทบุรี	180	อิงขาคำ	191
ปาดตะปุมมลายู	179	อิงทองแดง	203
ปาดตะปุมเล็ก	200	อิงทองเหลือง	202
ปาดตะปุมใหญ่	178	อิงน้ำเตา	191
ปาดดินแดง	186	อิงน้ำเต้ามลายู	203
ปาดดินเหลือง	199	อิงบอร์เนียว	203
ปาดดินเหลืองอีสาน	205	อิงปากขวด	182
ปาดนิ้วแยกมลายู	198	อิงปุมมลายู	190
ปาดบาน	198	อิงปุมหลังลาย	190
ปาดป่าจูดขาว	198	อิงเพา	182
ปาดเมืองจีน	202	อิงแม่หนาว	190
ปาดยักษ์	186	อิงลาย	189
ปาดลายละเอียด	199	อิงลายเต็ม	191
ปาดลายละเอียดเหนือ	205	อิงหลังซัด	192
ปาดลายละเอียดอีสาน	199	อิงหลังจุด	192
ปาดหูคำ	199	อิงอันนัม	202
อิงกรายขวางแถบ	181	อิงอังกนซัด	183
อิงกรายขาขาว	182	อิงอังกน	190
อิงกรายคอยอินทนนท์	202	อิงอังกนลาย	190
อิงกรายตาขาว	202		

## Sources

Photo Credit	Page
สถาบันวิจัยวิทยาศาสตร์และเทคโนโลยีแห่งประเทศไทย Thailand Institute of Scientific and Technological Research	10, 12, 13, 18, 27, 33, 39
มูลนิธิคุ้มครองสัตว์ป่าและพรรณพืชแห่งประเทศไทย Wildlife Fund Thailand	
นันทชัย พงศ์พัฒนานุรักษ์ Nanthachai Pongpattananurak	11, 16, 25
นริศ ภูมิภาคพันธ์ Naris Bhumpakphan	14
ไสว วังหงษา Sawai Wanghongsa	15
นายสัตวแพทย์เกษตร สุเตชะ Kaset Sutecha	17
บุษบง กาญจนสาขา Budsabong Kanchanasaka	19
พงศกร บัณฑิตพงศ์ Pongsakorn Pattapong	20, 58, 59
โดม ประทุมทอง Dome Pratumtong	21
สถาบันวิจัยและพัฒนาทรัพยากรทางทะเล ชายฝั่งทะเล และป่าชายเลน Phuket Marine Biological Center	22, 23 26 42-55
โอภาส ขอบเขตต์ Obhas Khobkhet	24
สุรชิต แวงโสธรณ์ (ฝ่ายวิจัยสิ่งแวดล้อม นิเวศวิทยา และพลังงาน) สถาบันวิจัยวิทยาศาสตร์และเทคโนโลยีแห่งประเทศไทย Surachit Waengsothorn (Environment, Ecology and Energy Department), Thailand Institute of Scientific and Technological Research	28
สมหญิง ทัพหิกรณ Somying Thunhikorn	29



Photo Credit	Page
โจนาธาน เมอเรย์ Jonathan Murray	30
เขตรักษาพันธุ์สัตว์ป่าทุ่งใหญ่นเรศวร Thung Yai Naresuan Wildlife Sanctuary	31
เกรียงศักดิ์ ศรีบัวรอด Kriangsak Sribuarod	32, 34
ดุสิต อาทิตยवार Dusit Arthitayawan	35
รณยุทธ ศรีบุญยรานนท์ Ronayuth Sribanyaranond	36
ประทีป ดวงแคะ Prateep Duengkae	37, 38, 40, 41, 57, 61
นิตยสาร Advanced Thailand Geographic Advanced Thailand Geographic Magazine	56
สมาคมอนุรักษ์สัตว์ป่า (WCS) ประเทศไทย Wildlife Conservation Society (WCS Thailand)	60
ธัญญา จันอาจ Tanya Chan-ard	86-95, 97, 99-103, 176-180
ศูนย์วิจัยและพัฒนาประมงน้ำจืดกาญจนบุรี Kanchanaburi Inland Fisheries Research and Development Center	96
นฤมล กฤษณชาญดี Narumon Kritsanachandee	98

**Not for sale :** If you have purchased or witnessed purchasing of this document, kindly inform such activity to  
Natural Resources and Environmental Management  
and Coordination Division  
Office of Natural Resources and Environmental Policy  
and Planning  
60/1 Rama VI Road Bangkok 10400 THAILAND  
Telephone (66 2) 265 6638-39  
Facsimile (66 2) 265 6638  
Website: <http://chm-thai.onep.go.th>  
E-mail: [chm\\_thai@onep.go.th](mailto:chm_thai@onep.go.th)

**ห้ามซื้อขาย :** หากท่านได้ซื้อ หรือพบเห็นการซื้อขายเอกสารเล่มนี้ โปรดแจ้ง  
ฝ่ายความหลากหลายทางชีวภาพ  
สำนักงานนโยบายและแผนทรัพยากรธรรมชาติและสิ่งแวดล้อม  
60/1 ซอยพินุลวิวัฒนา 7 ถนนพระราม 6 กรุงเทพฯ 10400  
โทรศัพท์ : (66 2) 265 6638-39  
โทรสาร : (66 2) 265 6638