## **5. SOULIEA** Franchet, J. Bot. (Morot) 12: 69. 1898.

黄三七属 huang san qi shu

Li Liangqian (李良千); Michio Tamura

Herbs perennial. Rhizome robust. Stems with membranous scales at base. Leaves petiolate, 2 or 3 × ternately sect. Inflorescence racemose. Flowers actinomorphic. Sepals 5, petaloid, white, obovate or obovate-elliptic. Petals 5, flabellate-obovate, denticulate. Stamens numerous; filaments narrowly linear; anthers broadly ellipsoid. Follicles 1–3, broadly linear, shortly stalked, reticulate; persistent style short. Seeds many, black, narrowly ovoid-globose. One species: Bhutan, China, Myanmar, Sikkim.

Elsewhere (Compton et al., Taxon 47: 593–634. 1998), *Souliea* has been transferred to *Actaea*. However, for the present account, as in FRPS, the authors have maintained *Souliea* as distinct, and regional advisor Rudolf V. Kamelin (pers. comm.) agrees.

**1. Souliea vaginata** (Maximowicz) Franchet, J. Bot. (Morot) 12: 70. 1898.

黄三七 huang san qi

Isopyrum vaginatum Maximowicz, Fl. Tangut. 18. 1889; Actaea vaginata (Maximowicz) J. Compton; Coptis ospriocarpa Brühl.

Rhizome robust, creeping, 4–9 cm in diam., branched, with fibrous roots. Stems more than 15 cm tall, to 75 cm at fruiting, glabrous or subglabrous, with 2-4 membranous broad scales at base. Cauline leaves ca. 2, glabrous, long petiolate; petiole 5–34 cm; leaf blade triangular, to 24 cm; primary seg-ments ovate to broadly so, central secondary segments ovate-triangular,  $4-7.5 \times 3.5-6.5$  cm, central tertiary segments rhomboid, 1- or 2-pinnatifid, unequally dentate at margin, lateral tertiary segments similar to central tertiary ones, oblique. Raceme 4-6-flowered; bracts ovate, membranous. Pedicel 1.2-1.4 cm. Flowers 1.2-1.4 cm in diam. Sepals  $8-11 \times 4-7$  mm, 3-veined, margin irregularly shallowly wavy at apex, apex rounded. Petals 2.5-5.5 mm, multiveined. Stamens 4-7 mm. Follicles 1 or 2(or 3),  $3.5-7 \times 0.3-0.7$  cm. Seeds 12–16, black when ripe, 3-4 mm. Fl. May-Jun, fr. Jul-Sep. 2n = 16\*.

Forests, forest margins, grassy slopes; 2800–4000 m. S Gansu, E Qinghai, S Shaanxi, W Sichuan, SE Xizang, NW Yunnan [Bhutan, India, N Myanmar, Sikkim].

Flora of China 6: 143. 2001.