

Trematoda

Rediae, Cercariae encyst, Metacercariae, mature adult

Developmental stages of *Ribeiroia ondatrae*.

Commonly referred to as "Flukes"
 2 Dominant species Responsible for Amphibian declines:
Ribeiroia ondatrae,
Echinostomatidae trivolvis

Both of these trematodes have complex life cycles requiring them to infect three different host species, namely, planorbid snails, larval amphibians or fish, and birds or mammals

Kingdom: Animalia
 Phylum: Platyhelminthes
 Class: Trematoda
 Order: Echinostomida

<http://www.science-art.com/2008/07/10/3553262.html>

<http://youtu.be/AcQxChHTGg>

Ribeiroia ondatrae


Conservationwildlife.org


amphibiaweb.org


Pieter T.J. Johnson

Echinostomatidae trivolvis









References

Kiesecker, J. M. (2011). Global stressors and the global decline of amphibians: tipping the stress immunocompetency axis. *Ecological research*, 26(5), 897-908.

Goodman, B. A., & Johnson, P. T. (2011). Disease and the extended phenotype: parasites control host performance and survival through induced changes in body plan. *PLoS One*, 6(5), e20193.

Johnson, P. T., Chase, J. M., Dosch, K. L., Hartson, R. B., Gross, J. A., Larson, D. J., ... & Carpenter, S. R. (2007). Aquatic eutrophication promotes pathogenic infection in amphibians. *Proceedings of the National Academy of Sciences*, 104(40), 15781-15786.

Johnson, P. T., & Sutherland, D. R. (2003). Amphibian deformities and *Ribeiroia* infection: an emerging helminthiasis. *Trends in Parasitology*, 19(8), 332-335.
