

## Minoru Tsukagoshi<sup>1,\*</sup>, Arata Momohara<sup>2</sup> and Mutsuhiko Minaki<sup>3</sup>: *Metasequoia* and the life and work of Dr. Shigeru Miki

**Abstract** Dr. Shigeru Miki (1901–1974) made great contributions to the fields of paleobotany, paleoecology, and plant taxonomy, including the discovery of *Metasequoia* Miki and the study of extant water plants. Details of Dr. Miki's biography, the original description of *Metasequoia*, his collection of fossil and extant plants that are stored in the Osaka Museum of Natural History, the introduction of living *Metasequoia* to Japan, episodes of Dr. Miki's life, and a list of Dr. Miki's publications are presented in this paper. Dr. Miki was a scientist and colleague who spent his career being enthused with research of the natural world.

**Key words:** bibliography, Dr. Shigeru Miki, *Metasequoia*

### Introduction

Dr. Shigeru Miki (1901–1974; Fig. 1) established the genus *Metasequoia* Miki in his paper entitled “On the change of flora in eastern Asia since Tertiary period (I). The clay or lignite beds flora in Japan with special reference to the *Pinus trifolia* beds in central Hondo.” This paper was published on September 18, 1941 in the *Japanese Journal of Botany*. Accordingly 2010 is the 70th anniversary of the establishment of the genus *Metasequoia*, and as a tribute to Dr. Miki the Third International *Metasequoia* Symposium was held at the Osaka Museum of Natural History.

In this paper we summarize the life and work of Dr. Miki, describe the history of living *Metasequoia* following its introduction to Japan in 1950, and provide a list of Dr. Miki's publications.

### Biography of Dr. Shigeru Miki

Biographies of Dr. Miki have been published by Kōkawa (1990), Saito (1995), and Tsukagoshi (2010). We summarize them in this paper and introduce aspects of his life with a photographic collage.

Dr. Miki was born in the town of Miki, Kida Province, Kagawa Prefecture in 1901. Following graduation at the Morioka Higher Professional School of Agriculture and Forestry (later Faculty of Agriculture, Iwate University), he became a teacher at the Ishikawa Prefectural School of Agriculture and Forestry. A short time later he left Ishikawa Prefectural School and en-




Fig. 1 Dr. Shigeru Miki (1901–1974). Photographed at Osaka City University in 1963.

<sup>1</sup> Osaka Museum of Natural History, 1-23 Nagai Park Higashi-sumiyoshi Osaka 546-0034, Japan

<sup>2</sup> Graduate School of Horticulture Chiba University, Matsudo 648 Chiba 271-8510, Japan

<sup>3</sup> University of Marketing and Distribution Sciences, 3-1 Gakuen-Nishimachi Nishi-ku Kobe Hyogo 651-2188, Japan

\* Corresponding author (e-mail: mtsuka@mus-nh.city.osaka.jp)

tered the Department of Botany, Faculty of Science, Kyoto University to study plant taxonomy. He began the study of the taxonomy and ecology of water plants under Professor Kwan Koriba (Figs. 2, 3). In 1925, he graduated from Kyoto University and became an assistant at Kyoto University. In 1931, he began working on fossil plants. His thesis was founded on better understanding changes in aquatic floras in time and in space. In 1949, he was promoted to Professor of the Faculty of Science and Technology, Osaka City University (Figs. 4, 5). After retiring from the Osaka City University, he was appointed Professor of the Faculty of Pharmacy Mukogawa Women's University.

From 1971 to 1974 he was the president of a support group for Osaka Museum of Natural History and here he focused on disseminating information about natural history to children and the community (Fig. 6).

#### Highlights of Dr. Miki's life

- 1901. Born in the town of Miki, Kida Province Kagawa Prefecture.
- 1918. Graduated from the Kida School of Agriculture and Forestry (later the Faculty of Agriculture Kagawa University).
- 1921. Graduated from the Morioka Higher Profes-




Fig. 3 A photograph showing Dr. Miki collecting specimens of *Nuphar* Sm.



Fig. 2 Dr. Miki as a student in the Department of Botany, Faculty of Science, Kyoto University.



Fig. 4 Dr. Miki in the laboratory of Osaka City University in 1957.

sional School of Agriculture and Forestry (later the Faculty of Agriculture, Iwate University). He became a teacher at the Ishikawa Prefectural School of Agriculture and Forestry.

1922. Entrance into the Department of Botany Faculty of Science, Kyoto University.
1925. Graduated from Kyoto University and hired as an Assistant Professor at Kyoto University.
1937. Published “The water phanerogams in Japan, with special reference to those of Prov. Yamashiro”.
1938. Completed his doctoral dissertation at Kyoto University “On the Pleistocene Flora in Prov. Yamashiro with the descriptions of 3 new species and 1 new variety”.
1939. Promoted to Lecturer at Kyoto University.
1941. Established the new fossil genus *Metasequoia* in the paper “On the change of flora in eastern Asia since Tertiary Period (I). The clay or lignite beds flora in Japan with special reference to the *Pinus trifolia* beds in Central Hondo”.
1944. Appointed as a researcher, Makassar Institute at Borneo.
1947. Promoted to Professor, Osaka University of Liberal Art and Education (later the Osaka University of Education).
1949. Promoted to Professor of the Faculty of Science and Technology, Osaka City University.
1951. Won the Asahi Shimbun Company prize for culture.
1964. Retired from the Osaka City University and appointed Professor of the Faculty of Pharmacy Mukogawa Women’s University.
1974. Deceased (73 years old).



Fig. 5 Dr. Miki (left) at the laboratory, Osaka City University laboratory in 1964 with Dr. Shohei Kikawa (center) and Dr. Baki Kasapliligil (right).



Fig. 6 Dr. Miki discussing *Convallaria* L. (lily of the valley) to elementary and junior high school students at an educational event organized by the Osaka Museum of Natural History.

### Bibliography of Dr. Shigeru Miki

Dr. Miki published 120 papers and wrote and edited two books (Appendix 1). Dr. Miki had great achievements in late Cenozoic paleobotany and paleoecology and the taxonomy of aquatic plants. Dr. Miki described the floral change of the late Pliocene to middle Pleistocene in Japan (Miki, 1948), collected many plant fossils throughout Japan, studied the taxonomy and phylogeny of fossil plants, and published a number of monographs on fossil plants. Important monographs include the Taxodiaceae (Miki, 1949c, 1950c), Pinaceae (Miki, 1957a), *Trapa* L. (Miki, 1952b), Nymphaeaceae (Miki, 1960a), Juglandaceae (Miki, 1955d), Vitaceae (Miki,

1956b), Alangiaceae-Cornaceae-Nyssaceae (Miki, 1956c), aquatic plants (Miki, 1961), and Lauraceae (Miki, 1970a).

Dr. Miki described many new species of fossil plants. However, in some cases he did not specify their type specimens. Some new species were described without any figures and photographs, and some were invalidly published names without description (e.g., some fossil species named by Miki, 1963a). As such, future work on Dr. Miki's collection will require lectotypification of parts of his collection and publication of valid descriptions for some of the fossil species that he named.




Fig. 7 Prepared specimens of shoot, cone (longitudinal section, axis), and seed remains of *Metasequoia disticha* (Heer) Miki. In the top row, the first and third shoots from the left (top left) are the specimens figured in Plate 5B (left) and Figure 8Da and in Plate 5B (centre) and Figure 8Db, respectively, of Miki (1941) (see Fig. 10). In the top row, the left shoot in the sixth slide from the left is the specimen illustrated in Figure 8Gc in Miki (1941). Graduations on the scale bar are 1 cm.

**Dr. Miki's collections  
of Osaka Museum of Natural History**

After his death in 1974, his collection was donated to the Osaka Museum of Natural History and the Mukogawa Women's University. Plant fossils and aquatic plant specimens reside in the Osaka Museum of Natural History, while extant plant specimens (excluding aquatic plants) are curated at the Mukogawa Women's University. The Osaka Museum of Natural History has published lists of these specimens (Nasu & Taruno, 1999; Kokawa et al., 2006; Shiga et al., 2009).

The specimens donated to the Osaka Museum of Natural History consist of plant fossil specimens that are stored in alcohol, prepared slide specimens, and extant plant materials that include herbarium sheets composed mainly of aquatic plants, prepared slides, and specimens preserved in alcohol. Catalogs of water plants (Shiga et al., 2009) and fossils of *Archaeozostera* Koriba et Miki (Nasu & Taruno, 1999) have been published.

The fossil and extant collection, excluding aquatic



**Fig. 8** Prepared specimens of fruits with cupule (longitudinal section), fruits (adhering part), cupules (juvenile), pericarps, male inflorescences, and leaves of extant plants (upper three rows and left half of the bottom row) and plant remains (right half of the bottom row) of *Quercus variabilis* Blume var. *brevipetiolata* Nakai. Graduations on the scale bar are 1 cm.

plants and *Archaeozostera*, was re-examined and arranged by Dr. Miki's favorite pupil, Dr. Shohei Kokawa, together with Minoru Tsukagoshi, Mutsuhiko Minaki, and Arata Momohara. The work included taxonomic verification of the specimens, label transcription and documentation, and replacement of the sample vials and preservative. Following Dr. Kokawa's death in 2001 Tsukagoshi, Minaki, and Momohara published a catalog listing the specimens held at the Osaka Museum of Natural History (Kokawa et al., 2006).

This catalog lists 24,779 plant fossil specimens and 7250 extant plants (Figs. 7–9). The fossil specimens include 27 fungi, 5 algae, 164 bryophytes, 68 pteridophytes, 7214 gymnosperms, and 15,975 angiosperms with 25 galls, and 1326 unidentified plants. The extant specimens include 42 bryophytes, 118 pteridophytes, 662 gymnosperms, and 6289 angiosperms with 139 unidentified plants. The collection is composed mainly of various sizes of glass slides mounted with Canada balsam and vials of materials stored in 70% ethanol. The collection includes many of the type specimens and slides used by Dr. Miki to describe new species and materials of the extant plant specimens that illustrate aspects of plant morphology and anatomy that were used to facilitate the taxonomic identity of the fossil materials.



**Fig. 9** Plant remains of *Pinus trifolia* Miki preserved in alcohol used in Plate 4E and Figure 3C of Miki (1939a). Graduations on the scale bar are 1 cm.

### Original description of *Metasequoia*

Although Dr. Miki established *Metasequoia* in 1941 (Miki, 1941; Fig. 10), he had already described the

character of shoots bearing distichous leaves and cones bearing decussate scales for a fossil taxon assigned to *Sequoia disticha* Heer (Miki, 1937; Figs. 1E-G, Pl. N-

#### *Metasequoia* n. g. (Pl. V A-D, Fig. 8 A-H)

The remains have usually been referred to *Sequoia* or *Taxodium*, indeed the cone is like that of *Sequoia* and the foliated shoot is somewhat like those of *Taxodium*.

The cones were never found connected to branches, but as the leaf-scars on the peduncle are also distichous, it is conceivable that the cones and the shoots belonged to the same plant. The foliated shoots seem to be lateral branches shedding in autumn, because their length is usually constant and the proximal end is covered by scaly leaves, although they have no scaly bud on the top and the branches two or more years old have two or more bud scars on the nearly same point as in Fig. 8 Ge.

The cone is distinguished from that of *Sequoia* by the decussate arrangement of scales and by the delicate peduncle having scale leaves at the base. The foliated shoot differs from *Sequoia* by distichous arrangement of leaves and by the brittle petiole. At a glance the shape of the shoot of fragmental remain seems to be *Taxodium* or *Cephalotaxus* but it differs from *Taxodium*<sup>(1)</sup> by distichous leaf and parallel arrangement of stomata on it and from *Cephalotaxus* by short delicate shoot without scaly bud at the terminal and by the obtuse top of leaf.

The decussate arrangement of cone-scales is not found in living Taxodiaceae, but a common character in Cupressaceae. The shedding of lateral foliated shoot with linear leaves is common in *Glyptostrobus* and *Taxodium*. So it is sure that the remains belong to Taxodiaceae but as the characters do not harmonize with those of the living forms, a new genus *Metasequoia* is established.

Character: Cone pedunculate, scale decussate, shield-form; peduncle with distichous scars of leaves and scaly leaves at the base. Shoot deciduous; leaf distichous, linear, obtuse, petiolate; stomata parallel to the midrib.

Two species were distinguished by the number of cone scales.

(1) The arrangement of stomata in *Taxodium* is transversal to the midrib (Fig. 8 I) being exceptional in Coniferae.

#### 10. *Metasequoia disticha* n. g. n. com. (Pl. V A-Ca, Fig. 8 A-G)

*Sequoia disticha* HEER (1876) 63, pl. 12, fig. 2 a, pl. 13, fig. 9-11; NATHORST (1888) 5, pl. 1, fig. 1; MIKI (1937) 306, pl. 8 N-O, fig. 1 E-G.

*Sequoia Onuki* ENDO (1936) 173 fig. 6.

*Taxodium distichum* RICH. in FLORIN (1920) 16, 30, pl. 1, fig. 1-2; KONNO (1931) pl. 8, fig. 1-2.

Remains of cone and foliated shoot were found from Osusawa. Cone elliptic, 15-20 mm long, 1.5-2 cm broad, pedunculate about 1-3 cm long. Scale 3-5 mm high, 11-14 mm broad, number of scale on the cone about 16-20; seed 4-5 mm long, 3-3.5 mm broad including the wings; twig with distichous pectinate leaves. Leaf linear, obtuse, 12 mm long, 1.5 mm broad, stomata parallel to the midrib on the ventral side.

The remains identified to the genus by decussate arrangement of cone scales and distichous arrangement of leaves and to the species by the size and shape of the leaf. This species was described by HEER as *Sequoia disticha*, but after his illustrations he did not note the scale leaves on the base of lateral branches. The difference is, however, not essential, because the destitution of scale leaves on the lateral branch of terminal shoot in *Taxodium* and *Sequoia* is quite usual.

Other localities in Japan: Cliff of Asiyagawa, Pref. Hyogo; Clay Beds of Huke and Tutimaru in Sennangun, Pref. Osaka; Clay Beds of Hasimoto, Pref. Wakayama; Clay Beds of Sidatani in Simagahara, Pref. Mie; Cliff of Syurakuen in Titagun, Pref. Aiti; Lignite Beds of Hanataka near Takasaki, Pref. Gunma.

#### 11. *Metasequoia japonica* n. g. n. com. (Pl. V D, Fig. 8 Ab, H)

*Sequoia japonica* ENDO (1936) 172, fig. 5, 7-13; (1939) 337, pl. 23, fig. 15.

A few remains of cone occurred at Osusawa. The cone compressed, 1.2 cm long, 1.2 cm wide. Number of scale about 12-16, decussate; size of scale 10-11 mm broad, 4-5 mm high.

The remain identified to the genus by decussate arrangement of cone-scales and to the species by narrow and fewer number of cone-scales than the former.





Fig. 8. *Metasequoia* excl. I.

A-G (excl. Ab) *Metasequoia disticha* n. g. n. com.  
A Cones: a from Osusawa, c from Hanataka, Pref. Gunma, d-f from Hasimoto, Pref. Wakayama  $\times 1$ .  
B Axial parts of cone  $\times 1$  from Sidatani, Pref. Mie.  
C Seeds from Sidatani, Pref. Mie  $\times 2$ .  
D Foliated shoots from Osusawa  $\times 1$ .  
E Leaf from Osusawa: a top  $\times 10$ ; b arrangement of stomata  $\times 43$ .  
F Epidermis of the leaf: a from upper side  $\times 263$ ; b stomata from underside  $\times 335$ .  
G Twigs: a young terminal shoot from Osusawa  $\times 2$ ; b-c two years old shoot from Hanataka, Pref. Gunma  $\times 1$ ; d-e shoot from Hasimoto, Pref. Wakayama  $\times 1$ .  
Ab, H *Metasequoia japonica* n. g. n. com. from Osusawa  $\times 1$ .  
I Arrangement of stomata on the living leaf of *Taxodium distichum* RICH.  $\times 43$ .



Pl. V. Taxodiaceae.

A-Ca *Metasequoia disticha* n. g. n. com.  
A Cones from Hasimoto, Pref. Wakayama. B Foliated shoots from Osusawa.  
Ca Seeds from Hanataka, Pref. Gunma.  
D *Metasequoia japonica* n. g. n. com. from Osusawa.  
Eb, E-F *Sequoia sempervirens* ENDL. from Osusawa excl. Eb from Sidatani, Pref. Mie. E Cones. F Foliated shoots.

Fig. 10 The original description of the fossil genus *Metasequoia* (Miki, 1941).

O). It is assumed that Miki had realized the oppositely-arranged leaves and the opposite and decussate seed cones were distinct and probably belonged to a new genus at the time of the 1937 publication, but held off describing a new genus until he had obtained more well preserved specimens. Miki (1937) also discussed the synonymy of the taxon with *Sequoia japonica* Endo and *Taxodium distichum* Richard and pointed out their wide distribution in the polar regions. *Sequoia disticha* was assigned to *Metasequoia disticha* (Heer) Miki and *Sequoia japonica* to *Metasequoia japonica* (Endo) Miki, respectively in Miki (1941).

Miki (1941)'s description of *Metasequoia* was based on various isolated parts of the plants including shoots, leaves, cuticles, seeds, and cones and was aimed to reconstruct the whole plant and its paleoecology. Among the four fossil localities that provided the fossil materials used for the description, the *Pinus trifolia* bed assigned to the early Pliocene by Miki (1941) is now correlated with the early late Miocene based on the fission track dating of 12–10 Ma from tephra in the Tokiguchi and Seto Porcelain Clay Formations (Todo Collaborative Research Group, 1999).

The establishment of *Metasequoia* and the discovery of living *Metasequoia* caused paleobotanists to re-examine fossil *Sequoia* and *Taxodium* that had been previously described (Chaney, 1950). Following re-examination of the fossil record, temporal and spatial distribution patterns of *Metasequoia* and their biogeography throughout the temperate regions of the Northern Hemisphere were clarified (Florin, 1963; LePage et al., 2005).

### Introduction of living trees of *Metasequoia* into Japan

The *Metasequoia* Preservation Society was established on December 19, 1949. Initially, the Secretariat was set up by the Osaka University of the Liberal Art and Education, but was removed to the Osaka City University when Miki transferred institutions. Dr. Hitoshi Kihara, who was Dr. Koriba's senior pupil, became chairman of the *Metasequoia* Preservation Society (Fig. 11).

On February 28, 1950, one hundred seedlings of *Metasequoia glyptostroboides* Hu et Cheng were presented to the *Metasequoia* Preservation Society by Dr. Ralph W. Chaney, Berkeley University, USA. These seedlings were grown by Chaney from seeds planted in February 1949.

Many of these seedlings were planted throughout Japan to study their growth in various climate and soil types. The locations and number of seedlings planted throughout Japan are listed in Table 1 (*Metasequoia* Preservation Society, 1951a). In 1951, the tallest *Metasequoia* seedling was 125 cm in height and 2.3 cm in diameter and was growing at the Osaka City University (*Metasequoia* Preservation Society, 1951b). Records indicate that growth of the tree occurred very early in the growing season.

The first seed cones produced were from trees growing at the Yashiroda Botanical Garden, Shodo-shima Island, Kagawa Prefecture in 1952 (Yoshikawa, 1957). Although the tree growing at the Yashiroda Botanical Garden is not included in Table 1, it is assumed that this tree was initially sent to Osaka City University



Fig. 11 Dr. Ralph W. Chaney (left), Dr. Shigeru Miki (center), and Dr. Hitoshi Kihara (right). Dr. Kihara was a senior pupil in Dr. Koriba's laboratory and president of the *Metasequoia* Preservation Society. They took part in the planting of a small *Metasequoia* tree. The location and planting date are unknown.

**Table 1** Distribution list of seedlings of *Metasequoia* sent by Dr. Ralph W. Chaney to the *Metasequoia* Preservation Society of Japan in 1950 (*Metasequoia* Preservation Society, 1951a).

Locations of <i>Metasequoia</i> seedlings planted in Japan in 1950	No.
Hokkaido (Sapporo City, Hokkaido Pref.)	3
Tohoku University (Sendai City, Miyagi Pref.)	3
The University of Tokyo	
University Forests of Amagi (Minami- Izu Town, Shizuoka Pref.)	4
University Forests of Chichibu (Chichibu City, Saitama Pref.)	4
University Forests of Kiyosumi (Kamogawa City, Chiba Pref.)	4
Faculty of Agriculture (Tokyo, Tokyo Pref.)	3
Koishikawa Botanical Gardens (Tokyo, Tokyo Pref.)	3
Nikko Botanical Garden (Nikko City, Tochigi Pref.)	3
Faculty of Science (Tokyo, Tokyo Pref.)	8
Tokyo Pref.	3
Shinjuku Imperial Garden (Tokyo, Tokyo Pref.)	2
The Government Forestry Experiment Station (Meguro, Tokyo, Tokyo Pref.)	4
Asakawa Forest, the Government Forestry Experiment Station (Hachioji City, Tokyo Pref.)	4
National Diet Building (Tokyo, Tokyo Pref.)	2
Kyoto University (Kyoto City, Kyoto Pref.)	12
Kyoto City (Kyoto Pref.)	3
Nara City (Nara Pref.)	2
Kobe City (Hyogo Pref.)	3
Nishinomiya City (Hyogo Pref.)	2
Osaka City (Osaka Pref.)	2
Osaka City University (Osaka City, Osaka Pref.)	3
Takarazuka Botanical Garden (Takarazuka City, Hyogo Pref.)	2
Nagoya City (Aichi Pref.)	3
Gifu City (Gifu Pref.)	3
Kochi University (Kochi City, Kochi Pref.)	3
Kyusyu University (Fukuoka City, Fukuoka Pref.)	3
Miyazaki niversity (Miyazaki City, Miyazaki Pref.)	2
Doshisya University (Kyoto City, Kyoto Pref.)	1
Tottori University (Tottori City, Tottori Pref.)	3
The Asahi Shinbun (Newspaper Company)	3
	100

or Kyoto University and was later transferred to the Yashiroda Botanical Garden. In 1953 seed cones were also confirmed at the Botanical Garden at the Faculty of Science, Kyoto University, and Kamigamo Experimental Forest Station, Faculty of Agriculture, Kyoto University. In 1972 a tree grown from cuttings of original trees at the Kamigamo Experimental Forest Station bore the first pollen cones. However, it is believed that the cuttings were not cultivated from any of the original trees sent to the Kamigamo Experimental forest Station. In 1973 *Metasequoia* seeds collected from the trees growing at the Kamigamo Experimental Forest Station were germinated (Nakai & Furuno, 1974).

The purpose of the *Metasequoia* Preservation Society was to grow and increase the number of *Metasequoia* trees in Japan and to study the ancient environments under which *Metasequoia* grew. Besides these activities the *Metasequoia* Preservation Society provided reprints of Dr. Miki's (1950b) and distributed this paper to the community as an educational activity.

The *Metasequoia* Preservation Society propagated the species through cuttings and then sold them to schools, companies, and the public. These activities facilitated the distribution of the tree throughout Japan. Currently, large *Metasequoia* trees can be observed and enjoyed throughout the country because the trees grow very well in Japan. Thus trees introduced in 1950 cannot be distinguished from those propagated from cuttings of original trees. Unfortunately some trees introduced in 1950 were cut down unintentionally. We could confirm the *Metasequoia* trees that were introduced in 1950 at only six institutes as follows: Botanical Garden Faculty of Science Osaka City University;



**Fig. 12** Dr. Miki's students and staff during a laboratory outing. Front row, left, Dr. Hideo Takada, center Dr. Jitsuro Ueno, right, Dr. Shigeru Miki. Middle row, Dr. Matsuo Tsukada.

Botanical Garden Faculty of Science Kyoto University; Kamigamo Experimental Forest Station, Faculty of Agriculture, Kyoto University; Kobe Municipal Arboretum; The Government Forestry Experiment Station (Meguro); Tohoku University. These trees should be preserved as a monument to the international collaboration of scientists and exchange of ideas since the introduction of *Metasequoia* to Japan.

The discovery of living *Metasequoia* that Dr. Miki had initially established on the basis fossil materials brought very good news for Japan following World War II. His contribution was honored with the 1950 Asahi Prize for distinguished researchers. *Metasequoia* and the word “living fossil” became popular throughout the world because of the discovery of the living plants and Dr. Miki’s description of a living fossil. In Japan nursery trees were supplied to each school as an example of a “living fossil”. This is one reason that *Metasequoia* is perhaps one of the best known trees in Japan.

#### Anecdotes of Dr. Shigeru Miki

Dr. Miki immersed himself in study and is remembered by a number of great achievements, including the discovery and description of fossil *Metasequoia*. The following anecdotes of Dr. Miki were provided by Mrs.

Tamiko Miki, who was Dr. Miki’s wife, and Dr. Shohei Kokawa, who was Dr. Miki’s pupil (Figs. 12–15):

- There was a person in a pond who wore a suit. It was Dr. Miki who was collecting aquatic plants.
- When it rained heavily, Dr. Miki went to the fossil localities by train, only leaving a message to his wife, just saying going to the west or to the east. When he found a cliff from the train, he got off at the next station and collected plant fossils.
- After collecting plant fossils by Dr. Miki, there were no plant fossils left at the outcrop. Another paleobotanist had to wait for the next heavy rain to destruct



Fig. 13 Dr. Miki crouching on a boardwalk in the pond where *Trapa* and *Nuphar* had been growing.



Fig. 14 Shigeru Miki with his wife Tamiko Miki. She had visited the first discovered *Metasequoia* tree in China in 1988.



Fig. 15 Two trees of *Metasequoia* and one tree of *Sequoia* at the home of Dr. Miki in Takarazuka City, Hyogo Prefecture. The word Miki means three trees in Japanese.

the cliff.

- There was no fingerprints on Dr. Miki's fingers because he ground plant fossils on whetstones without gloves to make prepared slides.
- Dr. Miki instructed students to observe plants by using five senses, "Sight, sense of touch, olfaction, taste, and hearing."

Dr. Miki also served as a member of the editorial board for science textbooks published by Osaka Shyoseki Co. Ltd. Miki's work and discovery of *Metasequoia* were published in a school textbook of morality by Osaka Shyoseki Co. Ltd. Each school that adopted the textbook was provided with *Metasequoia* seedlings by Osaka Shyoseki Co. Ltd. (Saito, 1995; Sakazaki & Ando, 1998).

In looking back on his research (Miki, 1964c, d), Dr. Miki wrote, "My life was like that of a mole, because I studied aquatic plants and plant fossils, which were obtained from under the horizon or from underground," and "I could keep researching by the help of everybody. I make an effort to complete my incomplete research." Dr. Miki was a researcher who was enthusiastic with the study of nature.

#### Acknowledgements

We thank Mr. Hiroshi Miki for his offer of the photographs of Dr. Shigeru Miki, Mrs. Sachiyo Kokawa for her offer of photographs of Dr. Shohei Kokawa, Dr. Naoko Sasaki, Research Institute for Humanity and Nature and Dr. Ben A. LePage, PECO Energy Company for their valuable comments that improved the manuscript, and Dr. Ashley B. Lamb, University of Tennessee for giving us useful comments to the manuscript of the early stage.

#### References (Dr. Miki's papers are cited in Appendix 1)

- Chaney, R. 1950. A revision of fossil *Sequoia* and *Taxodium* in western North America based on the recent discovery of *Metasequoia*. *Transactions of American Philosophical Society, N. S.* 40: 171–239.
- Florin, R. 1963. The distribution of conifer and taxad genera in time and space. *Acta Horti Bergiani* 20: 121–312.
- Kokawa S. 1990. New genus *Metasequoia* and Prof. Shigeru Miki. *Japanese Journal of Historical Botany* 5: 49–55.
- Kokawa S., Tsukagoshi, M., Minaki, M. & Momohara, A. 2006. A list of fossil and extant plant specimens collected by Dr. Shigeru Miki. *Special Publications from the Osaka Museum of Natural History* 38: 1–254 (in Japanese with English list).
- LePage, B.A., Yang, H. & Matsumoto, M. 2005. The evolution and biogeographic history of *Metasequoia*. In: LePage, B. A., Williams, C. J. & Yang, H., eds., *The Geobiology and Ecology of Metasequoia*, 3–114. Springer, Dordrecht.
- Metasequoia* Preservation Society. 1951a. Report of outline of *Metasequoia* Preservation Society. 1 p.
- Metasequoia* Preservation Society. 1951b. Report of passage of *Metasequoia* Preservation Society 5: 1.
- Nakai, I. & Furuno, T. 1974. On the florescence and the fruitification of *Metasequoia glyptostroboides* Hu et Cheng. *Journal of Japanese Forest Research* 56: 294–296.
- Nasu, T. & Taruno, H. 1999. Fossils of *Archaeozostera* collected by Professor Shigeru Miki. *Special Publications from the Osaka Museum of Natural History* 31: 1–11. (In Japanese).
- Saito, S. 1995. *Metasequoia; Tree that emperor Hirohito loved*. Chuokoron-sha, Tokyo, 238 pp.
- Sakazaki, N. & Ando, M. 1998. *Metasequoia*, introduced to Japan after World War II. *Bulletin of Japan Association of Botanical Gardens* 32: 99–104.
- Shiga, T., Fujii, S. & Seto, K. 2009. A list of water plants collected by Dr. Shigeru Miki. *Special Publications from the Osaka Museum of Natural History* 41: 1–42 (in Japanese).
- Todo Collaborative Research Group. 1999. Fault bounded inland basin of multiple blocks: An example from the sedimentary basin of the Tokai Group around Tajimi City in Gifu Prefecture, central Japan. *Earth Science (Chikyu Kagaku)* 53: 291–306 (in Japanese with English abstract).
- Tsukagoshi, M. 2010. Great paleobotanist Dr. Shigeru Miki. In: Tsukagoshi, M., supervised, *Fossil Plants; Memories of 500 million years ago*, 65–72. INAX Publisher, Tokyo.
- Yoshikawa, K. 1957. Flowering of *Metasequoia glyptostroboides* Hu et Cheng. *Journal of Japanese Forest Research* 39: 357–360.

(Accepted: 9 Mar. 2011)

### Appendix 1. Bibliography of Dr. Shigeru Miki

- Miki, S. 1927a. Über das Verzweigungsmodus und die Blattanordnung des Rhizoms von *Nelumbo nucifera* Gaertn. *Botanical Magazine, Tokyo* 41: 522–526, 18 plates (in Japanese with German abstract).
- Miki, S. 1927b. Oekologische Studien über die Sumpf- und Wassergewächse sowie ihre Formationen im Ogra-Teiche. *Kyoto-fu Shiseki-meisyo Tennen-kinenbutu Chyosahoukoku [Report of the Natural Monument in Kyoto Prefecture]* 8: 81–145, plates 26–50 (in Japanese).
- Miki, S. 1928. Über das Regenerationsvermögen der Blätter von *Pinellia ternata*, Breitenb. *Morioka Koutou Nourin Gakkou Souritsu 25-shu-nen Kinen Ronsou [Report of 25th Anniversary of Morioka Higher Professional School of Agriculture and Forestry]*, 1–8, 1 plate (in Japanese).
- Miki, S. 1929. Oekologische Studien von Mizoro-Teiche. Die Entstehung der Schwimmseln und die Schwankung des Sauerstoffes der Kohlensäure und der Wasserstoffionenkonzentration im Laife des Tages und der Jahreszeiten (Vorlaufiege Mitteilung). *Kyoto-fu Shiseki-meisyo Tennen-kinenbutu Chosa-houkoku [Report of the Natural Monument in Kyoto Prefecture]* 8: 81–145, plates 24–33 (in Japanese).
- Miki, S. & Koriba K. 1931. On *Archeozostera* from Izumi Sandstone. *Chikyū [The Globe]* 15: 165–204, plates 24–33 (in Japanese).
- Miki, S. 1932a. On sea-grasses new to Japan. *Botanical Magazine, Tokyo* 46: 774–788, 13 plates.
- Miki, S. 1932b. Ecological study of plant in Shinto shrine. *Jingū-shityō Rimen-ka Houkoku-shō [Report of Forestry Section of Shinto Shrine Branch]*, 1–75, plates 1–6 (in Japanese).
- Miki, S. 1932c. Morphology of lotus flower with special reference to the double headed flower. *Morioka Koutou Norin Gakkou Dousou-kai Gakujutu Shu-u-hō [Science Report of the Alumni Association of the Morioka Higher Professional School of Agriculture and Forestry]* 7: 1–10, plates 1–3 (in Japanese).
- Miki, S. & Ohwi, J. 1932. On the occurrence of *Loropetalum chinense* Oliver in Prov. Ise, Japan Proper. *Botanical Magazine, Tokyo* 46: 435–436 (in Japanese).
- Miki, S. 1933a. Water and marsh plants from Iturup Island. *Japanese Journal of Limnology* 3: 10–15 (in Japanese).
- Miki, S. 1933b. On the regeneration of leaves and roots of some water and marsh plants in Japan. *Transactions of the Tottori Society of Agricultural Science* 4: 183–194 (in Japanese with English abstract).
- Miki, S. 1933c. On the Pleistocene flora in Prov. Yamashiro. *Kyoto-fu Shiseki-meisyo Tennen-kinenbutu Chosahoukoku [Report of the Natural Monument in Kyoto Prefecture]* 14: 1–27, plates 1–5 (in Japanese).
- Miki, S. 1933d. On the Pleistocene flora in Prov. Yamashiro with the descriptions of 3 new species and 1 new variety. *Botanical Magazine, Tokyo* 47: 619–631, 1 plate.
- Miki, S. 1933e. On the sea-grasses in Japan (I). *Zostera* and *Phyllospadix*, with special reference to morphological and ecological characters. *Botanical Magazine, Tokyo* 47: 842–862, 3 plates.
- Miki, S. 1934a. On the sea-grasses in Japan (II). Cymodoceaceae and marine Hydrocharitaceae. *Botanical Magazine, Tokyo* 48: 131–142.
- Miki, S. 1934b. On the sea-grasses in Japan (III). General consideration on the Japanese sea-grasses. *Botanical Magazine, Tokyo* 48: 171–178. (Japanese abstract of “On the sea-grasses in Japan I – III”: pp. 215–219 in volume 48.).
- Miki, S. 1934c. On fresh water plants new to Japan. *Botanical Magazine, Tokyo* 48: 326–337, 1 plate. (Japanese abstract: pp. 361–363 in volume 48).
- Miki, S. 1934d. *Potamogeton* from Kurile Islands. *Japanese Journal of Limnology* 3: 122–128 (in Japanese).
- Miki, S. 1935a. New water plants in Asia Orientalis I. *Botanical Magazine, Tokyo* 49: 687–693 (Japanese abstract: pp. 739–740 in volume 49).
- Miki, S. 1935b. New water plants in Asia Orientalis II. *Botanical Magazine, Tokyo* 49: 773–780.
- Miki, S. 1935c. New water plants in Asia Orientalis III. *Botanical Magazine, Tokyo* 49: 847–852 (Japanese abstract of “New water plants in Asia Orientalis II and III”: pp. 827–829 in volume 49).
- Miki, S. 1936. Plant fossils from *Stegodon* and *Elephas* beds near Akashi. *Chikyū [The Globe]* 26: 155–190, plates 3 and 4 (in Japanese).
- Miki, S. 1937a. The origin of *Najas* and *Potamogeton*. *Botanical Magazine, Tokyo* 51: 472–480, 10 plates. (Japanese abstract: pp. 619 in volume 51).
- Miki, S. 1937b. Plant Fossil from the *Stegodon* Beds and the *Elephas* Beds near Akashi. *Japanese Journal of Botany* 8: 303–341, plates 8 and 9.
- Miki, S. 1937c. The water phanerogams in Japan with special reference to those of Prov. Yamashiro. *Kyoto-fu Shiseki-meisyo Tennen-kinenbutu Chosa-houkoku [Report of the Natural Monument in Kyoto Prefecture]* 17: 1–127, 4 plates (in Japanese).
- Miki, S. 1938a. On the change of flora of Japan since the upper Pliocene and the floral composition at the present. *Japanese Journal of Botany* 9: 213–251, plates 3 and 4.
- Miki, S. 1938b. On the remains of the water plants in Asia. *Japanese Journal of Limnology* 8: 410–416 (in Japanese).
- Miki, S. 1939. On the remains of *Pinus trifolia* n. sp. in the Upper Tertiary from central Honshū in Japan. *Botanical Magazine, Tokyo* 53: 239–246, 4 plates. (Japanese abstract: pp. 279–280 in volume 53).
- Koriba, K. & Miki, S. 1939. On *Paleodictyon* and fossil *Hydrodictyon*. *Jubilee Publication in the Commemoration of Professor Yabe's 60th Birthday* 1: 55–68, plates 4 and 5.
- Miki, S. 1940. Water plants of Manchuria. *Kanto-shū oyobi Manshū-koku Rikusui-seibutsu Chosa-shō [Report of the Fresh Water Flora and Fauna of Kwantung Leased Territory and Manchukuo]*, 470–476.
- Miki, S. 1941a. On the change of flora in Eastern Asia since Tertiary Period (I). The clay or lignite beds flora in Japan with special reference to the *Pinus trifolia* beds in Central Hondo. *Japanese Journal of Botany* 11: 237–303, plates 4–7.
- Miki, S. 1941b. Floral remains of conifer age at Manzidani near Nishinomiya, Japan (Preliminary note). *Japanese Journal of Botany* 11: 377–383.

- Miki, S. & Yano. 1941. Medicinal plant of Shangdong Province. *Kousei Chosa Shiryo [Report of the East Asia Development Board]* 46: 1–52 (in Japanese).
- Miki, S. & Kawase, I. 1942. Report of pasture resources in North China. *Kouain Chyosa Shiryo [Report of the East Asia Development Board]*, 214, 1–72, (includes floral list: 1–31)(in Japanese).
- Miki, S. 1945. Factor of healthy place and medical use of sediments in Banjarmasin. 3 pp. Banjarmasin Branch of Makassar Institute (in Japanese).
- Iseki, T. & Miki, S. 1945. Miki liquid as ulcer-healing agent. 6 pp. Banjarmasin Branch of Makassar Institute (in Japanese).
- Miki, S. 1948a. For the systematic position of *Hemitrapa* and some fossil *Trapa*. *Botanical Magazine, Tokyo* 61: 74–77 (in Japanese).
- Miki, S. 1948b. *Metasequoia*, “a living fossil.” *Botanical Magazine, Tokyo* 61: 108 (in Japanese).
- Miki, S. 1948c. Origin of Kibushi clay, Shibutsuchi-clay and Tennouji-clay and sedimentary environment of Kinki district. *Kobutsu to Chishitsu [Mineralogy & Geology]* 2: 16–18 (in Japanese).
- Miki, S. 1948d. Floral Remains in Kinki and adjacent district since the Pliocene with description 8 new species. *Kobutsu to Chishitsu [Mineralogy & Geology]* 9: 105–144 (in Japanese with English abstract and description of new species).
- Miki, S. 1948e. On *Metasequoia*, with special reference to the discovery of living species. *Seibutsu [Biology]* 4: 146–149 (in Japanese).
- Miki, S. 1949a. Consideration about origin of laterite. *Kobutsu to Chishitsu [Mineralogy & Geology]* 10: 152–153.
- Miki, S. 1949b. Question and consideration about origin of laterite. *Osaka Hakubutsu Gakkai-shi [Journal of Osaka Natural Science Association]* 9: 9–10.
- Miki, S. 1949c. Taxodiaceae in Japan, with special reference to the plant remains in lignite or clay beds. *Japanese Journal of Botany* 24: 34–40 (in Japanese with English abstract).
- Miki, S. 1949d. Tertiary environmental change based on differentiation of plant species. *Osaka Hakubutsu Gakkai-shi [Journal of Osaka Natural Science Association]* 9: 11–14.
- Miki, S. 1950a. A study on the floral remains in Japan since the Pliocene. “Nature and Culture”, *Science Report of Osaka Liberal Arts University* 1: 69–116 (in Japanese with English abstract).
- Miki, S. 1950b. *Metasequoia*. In: Komai, T. & Kihara, H., eds., *Saikin no Seibutsu-gaku [Recent Biology]* 1: 300–312. Baifukan Co., Ltd., Tokyo (in Japanese).
- Miki, S. 1950c. Taxodiaceae of Japan, with special reference to its remains. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 1: 63–77.
- Miki, S. & Hikita, S. 1950. The probable chromosome number on the remains of *Metasequoia* and *Sequoia* in Japan. *Botanical Magazine, Tokyo* 63: 119–123, 5 plates (in Japanese with English abstract).
- Miki, S. 1951a. Discovery of *Metasequoia*. *Kagaku Asahi* 11: 31–34 (in Japanese).
- Miki, S. 1951b. On *Metasequoia*. *Hana to Shokubutsu [Flower and Plant]* 1: 2–3 (in Japanese).
- Miki, S. & Hikita, S. 1951. Probable chromosome number of fossil *Sequoia* and *Metasequoia* found in Japan. *Science* 113: 3–4.
- Miki, S. 1952a. On the systematic position of *Hemitrapa* and some other fossil *Trapa*. *Palaeobotanist* 1: 346–350.
- Miki, S. 1952b. *Trapa* of Japan with special reference to its remains. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 3: 1–30.
- Miki, S. 1952c. *Metasequoia*. *Nihon Seibutsu Kyoiku Taikai Dai-6-Kai Osaka Taikai-shi [Report of Sixth Meeting of Japanese Society of Biological Sciences Education]*, 9–12, 1 plate (in Japanese).
- Miki, S. 1953a. Biological character and adaptation to the water life of *Trapa*, with special reference to its remains. *Bulletin of the Society of Plant Ecology* 2: 111–116 (in Japanese).
- Miki, S. 1953b. *Metasequoia*, *Fossil and Living*. 141 pp. Nippon Kobutsu Shumi-no-Kai, Kyoto (in Japanese).
- Miki, S. 1954a. The occurrence of the remain of *Taiwania* and *Palaeotsuga* (n. subg.) from Pliocene beds in Japan. *Proceedings of the Japanese Academy* 30: 976–981.
- Miki, S. 1954b. A few consideration for palaeontological problem in younger deposits of Kinki-district. *Chigaku Kenkyu [Geoscience Magazine]* 7: 5–8 (in Japanese).
- Miki, S. 1954c. Plant remains from construction ground of Osaka Common Building for Government Offices. In: The Society for the Study of the Site of the Castle in Osaka, ed., *Memoir upon the site of the Castle in Osaka II, Studies on the Site of the Old Forbidden City in Naniwa, the Capital of Japan in the Fifth and Eighth centuries*, 1–7, 1 plate. Osaka City University, Osaka (in Japanese).
- Miki, S. 1955a. *Five important timber trees and their historical change from aspect of plant fossils*. 17 pp, 2 plates. Nagano Regional Forestry Office, Nagano (in Japanese).
- Miki, S. 1955b. Water plants of “Sokonashi-ike” of the Tokara Islands. *Japanese Journal of Limnology* 17: 39–43, 1 plate (in Japanese).
- Miki, S. 1955c. Plants of Shodo-shima Island. *Nature Study (Special Issue, Nature of Shodo-shima Island)*, 6–8 (in Japanese).
- Miki, S. 1955d. Nut remains of Juglandaceae in Japan. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 6:131–144, plates 1–3.
- Miki, S. 1955e. Climate change since the Pliocene based on composition of plant remains. *News Bulletin, Japanese Branch of INQUA* 11: 34–39 (in Japanese).
- Miki, S. 1956a. Remains of *Pinus koraiensis* S. et Z. and associated remains in Japan. *Botanical Magazine, Tokyo* 69: 447–454, 13 plates.
- Miki, S. 1956b. Seed remains of Vitaceae in Japan. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 7: 247–271, 1 plate.
- Miki, S. 1956c. Endocarp remains of Alangiaceae, Cornaceae and Nyssaceae. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 7: 275–295, 1 plate.
- Miki, S. 1956d. Water plants and Environment. *Iden [Heredity]* 10: 32–35 (in Japanese).

- Miki, S. 1956e. *Greening of Rokko mountains and demand*. 2 pp, 1 plate (in Japanese).
- Miki, S. 1957a. Pinaceae of Japan with special reference to its remains. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 8: 221–272, plates 1–10.
- Miki, S. 1957b. *Metasequoia*. *Lecture Record of Japanese Association of Bamboo and Bamboo Grass* 1: 3–19 (in Japanese).
- Miki, S. 1957c. Remains of bamboo and bamboo grass. *Reports of Fuji Bamboo Garden* 2: 12–14 (in Japanese).
- Miki, S., Huzita, K. & Kokawa, S. 1957. On the occurrence of many broad-leaved evergreen tree remains in the Pleistocene bed of Uegahara, Nishinomiya City, Japan. *Proceedings of the Japanese Academy* 33: 41–46.
- Miki, S. 1958a. Gymnosperms in Japan, with special reference to the remains. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 9: 125–152, plates 1–3.
- Miki, S. 1958b. Estimation of topography, climate and age by paleontological method. *Issuikai Kaihou [News Bulletin of Issuikai]* 1: 67–72 (in Japanese).
- Miki, S. & Koriba, K. 1958. *Archeozostera*, a new genus from Upper Cretaceous in Japan. *Palaeobotanist* 7: 107–110, plates 1–2.
- Miki, S. 1959a. Evolution of *Trapa* from ancestral *Lythrum* through *Hemitrapa*. *Proceedings of the Japanese Academy* 35: 289–294.
- Miki, S. 1959b. Floral composition in the remains of Gymnosperms and Gamopetalae from Japan. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 10: 89–93.
- Miki, S. 1959c. Report of the Ninth International Botanical Congress. *Nature Study* 5: 5–7 (in Japanese).
- Miki, S. 1960a. Nymphaeaceae remains in Japan, with new fossil genus *Eoeryale*. *Journal of the Institute of Polytechnics, Osaka City University, Series D* 11: 63–78.
- Miki, S. 1960b. Seed-remains of *Euryale* in Japan. *Hokuriku no Shokubutsu [Journal of Geobotany]* 8: 53–57 (in Japanese with English abstract).
- Miki, S. 1961. Aquatic floral remains in Japan. *Journal of Biology, Osaka City University* 12: 91–121, plates 1–3.
- Miki, S. & Sakamoto, T. 1961. Neogene flora from Sasazu in Toyama Pref., Japan. **Professor J. Makiyama Memorial Volume**, 256–264, plates 1–2 (in Japanese with English abstract).
- Miki, S. 1962a. The phylogenetical considerations on the aquatic fossil genera, *Eoeryale*, *Eotrapa* and *Hemitrapa* from Akazu, Seto City, Japan. *Acta Phytotaxonomica et Geobotanica* 20: 139–144 (in Japanese with English abstract).
- Miki, S. 1962b. Fossil exhibition and *Pinus protodiphylla* of national treasure class. *Chigaku Kenkyu [Geoscience Magazine]* 12: 382–383 (in Japanese).
- Miki, S. 1962c. Plants and animals in Ogura-ike pond. *Ogura-ike Kantaku-shi [Report of Land Reclamation of Ogura-ike Pond]*, 254–265 (in Japanese).
- Miki, S. 1962d. Memories when *Metasequoia* was established. *Nature Study* 8: 26–29 (in Japanese).
- Miki, S. 1962e. *Archeozostera*. *Nature Study* 8: 15–16 (in Japanese).
- Miki, S. 1962f. *Archeozostera*. In: Hori, M., ed., *Flora of Osaka*, 172–173. Rokugatsu-shya, Osaka (in Japanese).
- Miki, S. & Kokawa, S. 1962. Late Cenozoic floras of Kyushu, Japan. *Journal of Biology, Osaka City University* 13: 65–85, plates 1–12.
- Miki, S., Kokawa, S. & Hikita, S. 1962. Floral change in Osaka. *Nature Study* 8: 16–18 (in Japanese).
- Miki, S. 1963a. Further study of plant remains in *Pinus trifolia* beds, central Hondo, Japan. *Chigaku Kenkyu [Geoscience Magazine]*, 31st Anniversary Volume, 80–93 (in Japanese).
- Miki, S. 1963b. Fossil plants. *Osaka and Science Education* 1: 41–43 (in Japanese).
- Miki, S. 1964a. Living fossil *Metasequoia*. *Iden [Heredity]* 18: 8–11.
- Miki, S. 1964b. Thirty years of Paleontology. *Kaseki [Fossil]* 8: 118–133 (in Japanese).
- Miki, S. 1964c. My life was like a mole, looking back on my research. *Nature Study* 10: 22–24 (in Japanese).
- Miki, S. 1964d. Pursuing the dream of natural research. *Nature Study* 10: 34–36 (in Japanese).
- Miki, S. 1964e. Mesozoic flora of *Lycoptera* beds in South Manchuria. *Bulletin of the Mukogawa Women's University (Natural Science)* 12: 13–22 (in Japanese with English abstract).
- Miki, S. ed. 1964f. *Biology*. 290 pp. Sogensha, Osaka (in Japanese).
- Miki, S. 1965. *Sequoiadendron primarium* Miki n. sp. and *Sequoia couttisii* Heer from Tertiary beds in Japan. *Bulletin of the Mukogawa Women's University (Natural Science)* 13: 1–7 (in Japanese with English abstract).
- Miki, S. 1966a. Difference of the floral composition and their origin between both side of Pacific Basin since Upper Tertiary beds in Japan. *Bulletin of the Mukogawa Women's University (Natural Science)* 14: 7–16 (in Japanese with English abstract).
- Miki, S. 1966b. Plant remains of *Camellia*. *Kyoto Engei [Gardening in Kyoto]* 53: 6–9 (in Japanese).
- Miki, S. & Maeda, Y. 1966. Fossil cone of *Keteleeria cretacea* n. sp. from Upper Cretaceous beds in Awaji, Japan. *Chigaku Kenkyu [Geoscience Magazine]* 17: 248–251 (in Japanese).
- Miki, S. 1967. Morphology and genus relation of fossil *Eotrapa*. *Bulletin of the Mukogawa Women's University (Natural Science)* 15: 267–272 (in Japanese with English abstract).
- Miki, S., Tsuchida, A. & Kishida, K. 1967. Floral list of medical herb garden and campus of Mukogawa Women's University. *Bulletin of the Mukogawa Women's University (Natural Science)* 15: 301–309 (in Japanese).
- Miki, S. 1968a. Morphology and evolutionary relationship between *Hemitrapa* and *Trapa*. *Bulletin of the Mukogawa Women's University (Natural Science)* 16: 281–286 (in Japanese with English abstract).
- Miki, S. 1968b. *Paleodavidia*, synonym of *Melliodendron* and fossil remains in Japan. *Bulletin of the Mukogawa Women's University (Natural Science)* 16: 287–291 (in Japanese with English abstract).

- Miki, S. 1969a. Evolution of water plants. *Shin-Kaki [New Ornamental Plants]* 62: 48–51.
- Miki, S. 1969b. *Protosequoia* (n.g.) in Taxodiaceae from *Pinus trifolia* beds in central Honshu, Japan. *Proceedings of the Japanese Academy* 45: 727–732.
- Miki, S. 1969c. Leaf adhered to the Sakuragaoka No.7 Bronze Bell. *Hyogo-ken Bunkazai Chyosa Houkoku-syo [Report of Cultural Asset Investigation of Hyogo Prefecture]* 1: 224–226 (in Japanese).
- Miki, S. 1970a. Lauraceae remains in Japan since the Pliocene, with description of 3 new species. *Bulletin of the Mukogawa Women's University (Natural Science)* 18: 231–248 (in Japanese with English abstract).
- Miki, S. 1970b. Trees in fossil age. *Ringyo Gijyutsu [Forestry Technology]* 334: 2–5 (in Japanese).
- Miki, S., Kishida, K. & Otsuka, K. 1970. Herbarial list of seed plants in Faculty of Pharmaceutical Sciences, Mukogawa Women's University. *Bulletin of the Mukogawa Women's University (Natural Science)* 18: 249–324 (in Japanese with English abstract).
- Miki, S. & Mizuno, M. 1970. *Osaka-fu Bunkazai Zuzetsu Tennen Kinenbutsu Syokubutsu-hen [Cultural Asset of Osaka Natural Monument, Plants]*. 31 pp., plates 1–48. The Board of Education of Osaka Prefecture (in Japanese).
- Miki, S. 1971. Plant remains from Kawashima and Tateoka sites. *Kawashima-Tateoka Iseki [Report of Kawasima and Tateoka Sites]*, 208–210. The Board of Education of Taishi Town (in Japanese).
- Miki, S. 1972a. On the fossil of *Pinus trifolia* Miki from Paleogene Formation in Kyushu. *Geological Report Faculty of Science Kumamoto University* 2: 72–73 (in Japanese)
- Miki, S. 1972b. Plant remains from moat in Shironoyama-Ikeda site. *Shironoyama-Ikeda kofun [Report of Shironoyama-Ikeda Site]*, 98–100. The Board of Education of Wadayama Town (in Japanese).
- Miki, S. 1972c. Plant remains from Gongen Site. *Harima-Gongen Iseki [Report of Harima-Gongen Iseki]* 6: 60–63. Association for Society and Culture of Hyogo Prefecture (in Japanese).
- Miki, S. 1973. Relationship between some semi-fossil occurring from Japanese Pliocene beds and the living in Asia. *Bulletin of the Mukogawa Women's University (Pharmacy)* 20–21: 1–9 (in Japanese with English abstract).
- Miki, S. 1995. Plant remains in and around of storage pit. *Minamikata-mae Iseki [Report of Minamikata-mae Site]*, 31–40. The Board of Education of Sanyo Town (in Japanese).