

The echinoderm (Deuterostomia) fauna of the Aegean Sea, and comparison with those of the neighbouring seas

ATHANASIOS KOUKOURAS*, APOSTOLOS I. SINIS, DIMITRA BOBORI,
SAVAS KAZANTZIDIS and MILTIADIS-SPYRIDON KITSOS

Department of Zoology, School of Biology, Aristotle University, Thessaloniki, 541 24, Greece

Received: 28 July 2006

Accepted after revision: 18 October 2006

Earlier and recent sampling carried out in the Aegean Sea revealed the presence of 20 asteroid, 14 ophiuroid, 21 echinoid, and 1 holothuroid species. The ophiuroid *Monaphiura apicula*, and the echinoids *Arbaciella elegans*, *Echinus melo* and *Hemaster exasperatus* are recorded for the first time in the Eastern Mediterranean; the latter has previously been known only from the Western Mediterranean. The ophiuroid *Amphiura (Acrocnida) brachiata* is recorded for the first time in the Aegean Sea. For all species, information on their distribution and habitat is given. A checklist of the Mediterranean and Black Sea echinoderms, as well as their distribution in the Mediterranean territorial areas and the Black Sea are also presented. Furthermore, the faunas of the Mediterranean territorial areas are compared. According to the considered data, the number of species decreases as follows: Western Mediterranean, Aegean Sea, Adriatic Sea, Central Mediterranean, Levantine Sea and Black Sea. Species with an Atlanto-Mediterranean distribution dominate in all areas and are followed in numbers by the Mediterranean endemics and cosmopolitans. Four species of Indo-Pacific origin are located in the Levantine and SE Aegean seas.

Key words: Echinodermata, Mediterranean Sea, Aegean Sea.

INTRODUCTION

Specific or scattered information on the echinoderm fauna of the Aegean Sea is mainly included in: Spratt & Forbes (1842), Forbes (1844, 1845), Raulin (1870), Carus (1885), Steindachner (1891), Marenzeller (1893, 1895), Ostroumoff (1896), Marion (1898), Panagiotopoulos (1916), Athanassopoulos (1917, 1921), Issel (1928), Tortonese (1946, 1947, 1965), Belloc (1948), Pérès & Picard (1958), Laborel (1960), Tortonese & Demir (1960), Jacquotte (1962), Kisseleva (1963, 1983), Makkavieva (1963), Caspers (1968), Vamvakas (1971), Geldiay & Koçatas (1972), Chardy *et al.* (1973), Koukouras & Sinis (1981), Bianchi & Morri (1983), Ünsal (1985), Pancucci & Zenetos (1990), Fiege & Yulin (1994) and Pancucci (1994).

Pancucci-Papadopoulou (1996), mainly based on a survey of the literature, recorded the presence of 103 echinoderm species (2 crinoids, 33 holothuroids, 25 asteroids, 23 ophiuroids and 20 echinoids) in the

Aegean Sea. More recent information on the Aegean echinoderm fauna has been provided only by Özaydin *et al.* (1995).

According to the information included in the above papers, the echinoderm fauna of the Aegean Sea (including the Sea of Marmara) consists of 103 species (2 crinoids, 33 holothuroids, 25 asteroids, 22 ophiuroids and 21 echinoids).

The present paper focuses on: i) the new information on the echinoderm fauna of the Aegean Sea and, ii) the comparison of the Aegean fauna with the faunas of the neighbouring seas.

MATERIALS AND METHODS

A total of 6725 echinoderm specimens, collected from 181 stations located in the Aegean Sea (Fig. 1) were examined. Information on the characteristics of the sampling stations is given in the presentation of the species found. The samples were obtained using fishing nets, various types of dredges and grabs and by free or SCUBA diving at depths up to 1250 m. The specimens have been deposited at the Museum of the

* Corresponding author: tel.: +30 2310 998363, fax: +30 2310 998269, e-mail: akoukour@bio.auth.gr




FIG. 1. Map of the Aegean Sea, indicating the sampling stations.

Department of Zoology, Aristotle University of Thessaloniki (MDZAUT).

RESULTS AND DISCUSSION

Taxonomic list

1. Eleutherozoa – Asteroidea

The following 20 species were found in the Aegean Sea during the present study.

Anseropoda placenta (Pennant, 1777)

Material: 31 specimens; stas 2, 16, 85, 104, C, E, I, J, M, V and W; depth 5-190 m; on rocks, biogenic detritus, and sandy, sandy silt and silty bottoms; Dmax. (maximum diameter) = 145.6 mm.

Distribution: A species known from various localities of the Aegean Sea (Forbes, 1844 as *Palmipes membranaceus*; Tortonese, 1947 as *Palmipes placenta*; Pérès & Picard, 1958 as *Anseropoda membrana-*

cea; Vamvakas, 1971; Geldiay & Koçatas, 1972).

An Atlanto-Mediterranean species (Table 1), known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1960; Tortonese, 1965; Kaspiris & Tortonese, 1982; Özaydin *et al.*, 1995).

Asterina gibbosa (Pennant, 1777)

Material: 152 specimens; stas 20, 25, 29, 42, 102, 105, 123, 127, 129, 130, 135, 140, 142, 146, 147, 151, 159, 162, 163, 168, 172, 173, 174, 186, 192, 194, 199, 200, 203, 205, 208 and 209; depth 0-27 m; on rocks, boulders, cobbles, and meadows of *Zostera* and *Posidonia*; Dmax. = 43.3 mm.

Distribution: Known from many localities of the Aegean Sea (Athanassopoulos, 1917; Tortonese, 1947; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1), known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1961, 1965; Zavodnik, 1976; Pancucci & Zenetos, 1989).

Astropecten aranciacus (Linnaeus, 1758)

Material: 54 specimens; stas 20, 50, 78, 80, 85, 87, 89, 99, 104, 105, 108, 109, 110, 127, 130, 134, 147, 153, 156, 162, 176, 189, 191, 193, 196, 201, A, C, D and E; depth 1-110 m; on biogenic detritus, sandy and sandy silt bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 289.2 mm.

Distribution: Known from various localities of the Aegean Sea (Tortonese, 1947; Tortonese & Demir, 1960; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1), known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Tortonese, 1965; Kaspiris & Tortonese, 1982; Özaydin *et al.*, 1995).

Astropecten bispinosus (Otto, 1823)

Material: 63 specimens; stas 97, 104, 105, 109, 115, 152, 159, 161, 162, 168, 192, A, C, D, E, V and W; depth 1-105 m; on sandy, sandy silt and silty bottoms, and meadows of *Posidonia*; Dmax. = 212.0 mm.

Distribution: This species has been known in the Aegean Sea, from the Sea of Marmara (Ostroumoff, 1896; Tortonese & Demir, 1960), Rhodos island (Tortonese, 1946) and the Izmir gulf (Geldiay & Koçatas, 1972).

An Atlanto-Mediterranean species (Table 1), known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1965; Zavodnik, 1981; Pancucci & Zenetos, 1989; Özaydin *et al.*, 1995).

Astropecten irregularis pentacanthus (Delle Chiaje, 1825)

Material: 632 specimens; stas 33, 40, 41, 43, 49, 79, 85, 89, 95, 103, 105, 109, 110, 112, 129, 133a, 155, 162, A, C, D, E, F, G, J, L, M, N, Q, R, S and W; depth 1-460 m; on biogenic detritus, sandy, sandy silt and silty bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 197.6 mm.

Distribution: Known in the Aegean Sea, from Antimilos islet (Marenzeller, 1893 as *A. pentacanthus*), the Sea of Marmara (Ostroumoff, 1896; Tortonese & Demir, 1960) and the Izmir gulf (Geldiay & Koçatas, 1972; Ünsal, 1985).

An Atlanto-Mediterranean species (Table 1), known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Zavodnik, 1977; Pancucci & Zenetos, 1989).

Astropecten jonstoni (Delle Chiaje, 1825)

Material: 24 specimens; stas 20, 54, 95, 104, 125, 132, 146, 160, 163, 168, 196 and F; depth 1-90 m; on biogenic detritus, sandy and sandy silt bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 75.2 mm.

Distribution: This species has been known from the Sea of Marmara (Tortonese & Demir, 1960), the Izmir gulf (Geldiay & Koçatas, 1972) and Kos island (Bianchi & Morri, 1983).

A Mediterranean endemic species (Table 1), known from all over the Mediterranean region (Mortensen & Steuer, 1937; Tortonese, 1965; Zavodnik, 1972; Pancucci & Zenetos, 1989).

Astropecten platyacanthus (Philippi, 1837)

Material: 7 specimens; stas 76, 104, 130, 157, 188 and 199; depth 1-16 m; on sandy and sandy silt bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 160.0 mm.

Distribution: This species has been known in the Aegean Sea only from the Sea of Marmara (Ostroumoff, 1896; Tortonese & Demir, 1960), Antiparos islet (Tortonese, 1965), and the Izmir gulf (Ünsal, 1985).

A Mediterranean endemic species (Table 1) known from all over the Mediterranean region (Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Rodríguez, 1979; Pancucci & Zenetos, 1989).

Astropecten spinulosus (Philippi, 1837)

Material: 7 specimens; stas 20, 127, 156 and 157; depth 1-28 m; on sandy and sandy silt bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 98.5 mm.

Distribution: This species has been known in the area only from the Sea of Marmara (Demir, 1952/1954) and the Izmir gulf (Geldiay & Koçatas, 1972; Ünsal, 1985).

A Mediterranean endemic species (Table 1) known from all over the Mediterranean region (Mortensen & Steuer, 1937; Cherbonnier, 1958; Tortonese, 1961; Zavodnik, 1967; Kaspiris & Tortonese, 1982).

Brisingella coronata (O. Sars, 1871)

Material: 2 specimens; stas Q and R; depth 120-320 m; on sandy silt and silty bottoms; LMax. (maximum arm length) = 310.0 mm.

Distribution: Known from various localities of the Aegean Sea (Steindachner, 1891; Marenzeller, 1893, 1895; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1893, 1895; Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976).

Chaetaster longipes (Retzius, 1805)

Material: 1 specimen; sta 79; depth 40-50 m; on sandy silt bottoms; D = 240.0 mm.

Distribution: This species has been known in the Aegean Sea only from the SW coast of Turkey (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Ludwig, 1897; Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Gjiknuri, 1981; Kaspiris & Tortonese, 1982).

Coscinasterias tenuispina (Lamarck, 1816)

Material: 49 specimens; stas 5, 15, 19, 20, 30, 68, 71, 89, 123, 138, 162, 175, 179, 181, 202, 205, 207, D and E; depth 0-50 m; on rocks, biogenic detritus, sandy

and silty sand bottoms, and meadows of *Zostera* and *Posidonia*; Dmax. = 165.0 mm.

Distribution: Known in the Aegean from Rhodos island (Tortonese, 1946), the Tainaron cape (Pérès & Picard, 1958), the Izmir gulf (Geldiay & Koçatas, 1972; Ünsal, 1985) and the Güllük gulf (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Ludwig, 1897; Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1958; Tortonese, 1961; Zavodnik, 1972; Kaspiris & Tortonese, 1982).

Echinaster (Echinaster) sepositus sepositus (Retzius, 1783)

Material: 100 specimens; stas 20, 30, 40, 79, 89, 110, 112, 129, 132, 134, 156, 168, 174, 183, 191, 194, 196, A, D, E, I, K, M, P, T, V and W; depth 0-210 m; on rocks, biogenic detritus and sandy, silty sand and sandy silt substrata and meadows of *Zostera* and *Posidonia*; Dmax. = 260.5 mm.

Distribution: Known from various localities of the Aegean Sea and the Sea of Marmara (Forbes, 1844; Ostroumoff, 1896; Tortonese, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Vamvakas, 1971; Geldiay & Koçatas, 1972; Ünsal, 1985; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Koehler, 1924; Cherbonnier, 1958; Tortonese, 1961; Zavodnik, 1972; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989; Özaydin et al., 1995).

Haelia attenuata (Gray, 1840)

Material: 6 specimens; stas 175, 176, 177, 186 and M; depth 1-190 m; on rocks, biogenic detritus, and sandy, silty sand and sandy silt substrata; Dmax. = 120.5 mm.

Distribution: Known from various localities of the Aegean Sea (Marenzeller, 1895; Pérès & Picard, 1958; Tortonese & Demir, 1960; Vamvakas, 1971; Geldiay & Koçatas, 1972; Kisseleva, 1983; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Ludwig, 1897; Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1972, 1981; Kaspiris & Tortonese, 1982).

Luidia ciliaris (Philippi, 1837)

Material: 6 specimens; stas 185, 189, 191, 192, A and T; depth 1-90 m; on rocks, biogenic detritus and sandy, silty sand and sandy silt substrata; Dmax. = 545.5 mm.

Distribution: Known in the Aegean Sea from the Sea of Marmara (Demir, 1952/1954; Tortonese & Demir, 1960), from Rhodos island (Tortonese, 1946, 1947) from the south of Santorini (Jacquotte, 1962), and from the SE of Limnos island and the Edremit gulf (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Ludwig, 1897; Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Gjiknuri, 1981; Kaspiris & Tortonese, 1982).

Luidia sarsi sarsi Düben & Koren, in Düben, 1845

Material: 6 specimens; stas 79, Q, S and V; depth 20-190 m; on sandy, silty sand and sandy silt substrata; Dmax. = 290.5 mm.

Distribution: This species has been known in the Aegean Sea only from the SE of the cape Maleas and between the cape Maleas and Milos (Marenzeller, 1893, 1895; as *L. paucispina*), from between Paros and Naxos islands (Pérès & Picard, 1958) and from the NE of Samothraki island (Jacquotte, 1962).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1893; Mortensen, 1927; Tortonese, 1957; Cherbonnier, 1958; Zavodnik, 1972; Gjiknuri, 1981).

Marthasterias glacialis (Linnaeus, 1758)

Material: 96 specimens; stas 18, 62, 78, 79, 85, 87, 109, 116, 125, 129, 133, 137, 146, 152, 153, 156, 162, 167, 174, 177, 168b, 179, 180, 183, 192, 193, 203, 205, A, C, L, Q, V and W; depth 1-105 m; on rocks, biogenic detritus and sandy, silty sand and sandy silt substrata, and meadows of *Zostera* and *Posidonia*; Dmax. = 360.6 mm.

Distribution: Known from various localities of the Aegean Sea (Forbes, 1844, as *Uraster glacialis*; Ostromoff, 1896, as *Asterias glacialis*; Athanassopoulos, 1917; Pérès & Picard, 1958; Tortonese & Demir, 1960; Geldiay & Koçatas, 1972; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic

regions (Mortensen, 1927; Cherbonnier, 1958; Tortonese, 1965; Zavodnik, 1972, 1981; Özaydin et al., 1995) as well as from the Black Sea – the Bosphorus and off the Bosphorus (Marion, 1898, as *Asterias glacialis*; Tortonese & Demir, 1960).

Odontaster mediterraneus (Marenzeller, 1893)

Material: 2 specimens; stas S and T; depth 80-250 m; on sandy silt and silty substrata; Dmax. = 80.5 mm.

Distribution: This species has been known in the Aegean Sea only from the west of Milos island (Marenzeller, 1893, as *Gnathaster mediterraneus*), between Serifos and Milos islands (Marenzeller, 1895) and from the west of Lesvos island, the SW of Skyros island, the north of Samos island and the north of Rhodos island (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from the European Atlantic coast (Mortensen, 1927), the western Mediterranean (Ludwig, 1897; Piras, 1972; Sibuet, 1974), the Adriatic Sea (Zavodnik, 1972) and the Levantine Basin; south coast of Turkey (Özaydin et al., 1995). It is not known from the Central Mediterranean.

Ophidiaster ophidianus (Lamarck, 1816)

Material: 1 specimen; sta R; depth 40 m; on silty sand substratum; D = 205.0 mm.

Distribution: This species has been known in the Aegean (Forbes, 1844, as *O. laevigata*) only from Rhodos island (Tortonese, 1946, 1947), Kos island (Bianchi & Morri, 1983), the south of Limnos island and the SE of Chios island (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Cherbonnier, 1958; Tortonese, 1965; Zavodnik, 1972, 1981; Özaydin et al., 1995; Pancucci-Papadopoulou, 1996).

Peltaster placenta (J. Müller & Troschel, 1842)

Material: 5 specimens; stas A, C, G and S; depth 40-180 m; on sandy and silty sand substrata; Dmax. = 90.7 mm.

Distribution: Known from various localities of the Aegean Sea (Marenzeller, 1895, as *Pentagonaster placenta*; Tortonese, 1946, 1947, as *Ceramaster placenta*; Pérès & Picard, 1958, as *Ceramaster placenta*; Tortonese & Demir, 1960, as *Sphaeriodiscus placenta*; Özaydin et al., 1995, as *Sphaeriodiscus placenta*).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Ludwig, 1897; Mortensen, 1927; Cherbonnier, 1958; Tortonese, 1961; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Kaspiris & Tortonese, 1982; Özaydin *et al.*, 1995).

Tethyaster subinermis (Philippi, 1837)

Material: 3 specimens; stas 79, R and S; depth 40-320 m; on silty sand and sandy silt substrata; Dmax. = 360.2 mm.

Distribution: This species has been known in the Aegean Sea only from Rhodos island (Tortonese, 1946, 1947), the S and SE of Limnos island, the SW and S of Lesvos island and the Gökova gulf (Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known in the Atlantic Ocean from the Bay of Biscay to Morocco and Liberia (Mortensen, 1927). Also known from the western Mediterranean (Cherbonnier, 1958), the Adriatic (Zavodnik, 1972), the central Mediterranean, the Taranto gulf (Tortonese, 1961), Peloponnisos (Kaspiris & Tortonese, 1982), the Levantine basin, Cyprus (Demetropoulos & Hadjichristophorou, 1976), and the south coast of Turkey (Özaydin *et al.*, 1995).

2. Eleutherozoa – Cryptosyringida – Ophiuroidea

The following 14 species were found in the Aegean Sea during the present study.

Amphipholis squamata (Delle Chiaje, 1828)

Material: 31 specimens; stas 16, 115, 120, 123, 141, 175, 188, 203, C, Q and V1; depth 0-210 m; on rocks, organic detritus, sandy and sandy silt substrata, as well as associated with sponges. Ddmax. (Maximum disc diameter) = 3.4 mm.

Distribution: Known from many localities of the Aegean Sea (Forbes, 1844, as *Amphiura neglecta*; Ostromouff, 1896, as *Amphiura squamata*; Tortonese, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Caspers, 1968; Vamvakas, 1971; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

A cosmopolitan species (Table 1) known from all the Mediterranean areas (Mortensen & Steuer, 1937; Tortonese, 1956, 1961, 1966; Cherbonnier, 1958; Zavodnik, 1972; Pancucci & Zenetos, 1989) and the Black Sea – the Bosphorus (Tortonese & Demir, 1960), the Atlantic Ocean (Mortensen, 1927) and the Pacific Ocean (Clark & Rowe, 1971).

Amphiura (Acrocnida) brachiata (Montagu, 1804)

Material: 1 specimen; sta 96; depth 5.3 m; on sandy substratum; Ddmax. = 8.5 mm.

Distribution: This species is reported for the first time from the Aegean Sea.

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Tortonese, 1965; Cherbonnier & Guille, 1967; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Pancucci & Zenetos, 1989; Özaydin *et al.*, 1995).

Amphiura chiajei Forbes, 1843

Material: 2815 specimens; stas 77, 78, 84, 96, 97, 101, 106, 107, 112, 113, 152, 156, 198, 209, C, D, I and V1; depth 2-250 m; on sandy, silty sand and sandy silt substrata; Ddmax. = 9.7 mm.

Distribution: Known from many localities of the Aegean Sea (Forbes, 1844, 1845, as *A. florifera*; Marenzeller, 1895; Ostromouff, 1896; Pérès & Picard, 1958; Tortonese & Demir, 1960; Caspers, 1968; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Spratt & Forbes, 1842; Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1952, 1956, 1961; Cherbonnier, 1958; Zavodnik, 1972, 1981; Demetropoulos & Hadjichristophorou, 1976; Pancucci & Zenetos, 1990) as well as from the Black Sea – off the Bosphorus (Tortonese & Demir, 1960).

Amphiura filiformis (O. F. Müller, 1776)

Material: 132 specimens; stas 77, 78, 153, A, C, E, H, I, J, R and W; depth 22-42 m; on sandy and silty sand substrata; Ddmax. = 8.4 mm.

Distribution: Also known from many localities of the Aegean Sea (Forbes, 1844; Raulin, 1870; Marenzeller, 1893, 1895; Ostromouff, 1896; Tortonese & Demir, 1960; Caspers, 1968; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1956, 1965; Cherbonnier, 1958, 1969; Zavodnik, 1972, 1981; Pancucci & Zenetos, 1989).

Amphiura mediterranea Lyman, 1882

Material: 4 specimens; stas 96 and C; depth 0-25 m; on sandy and silty sand substrata; Ddmax. = 6.2 mm.

Distribution: This species has been known in the Aegean Sea only from the SW of Lesvos island, the SW of Limnos island and off the SE coast of Evvoia island (Kisseleva, 1963) and from Lesvos island – Geras gulf (Pancucci & Zenetos, 1990).

A Mediterranean endemic species (Table 1) known from all over the Mediterranean regions (Mortensen & Steuer, 1937; Tortonese, 1956, 1961, 1965; Cherbonnier, 1958; Zavodnik, 1972; Pancucci & Zenetos, 1989).

Monamphiura apicula (Cherbonnier, 1957)

Material: 6 specimens; stas 33 and 207; depth 2-8 m; on sandy substrata with cobbles and meadows of *Zostera*; Ddmax. = 2.4 mm.

Distribution: This species is reported for the first time from the Aegean Sea.

A Mediterranean endemic species (Table 1) known until now only from the western Mediterranean (Cherbonnier, 1957, 1958, as *Amphiura apicula*; Tortonese, 1959, 1965, as *A. apicula*), and the Adriatic Sea (Zavodnik, 1995).

Ophiacantha setosa (Retzius, 1805)

Material: 114 specimens; stas 1, 45, 87, L and S; depth 5-255 m; on sandy, silty sand and sandy silt substrata; Ddmax. = 8.5 mm.

Distribution: This species has been known in the Aegean Sea only from the Sea of Marmara (Ostroumoff, 1896) and Syros island (Pérès & Picard, 1958).

An Atlanto-Mediterranean species (Table 1) known from the Atlantic – Bay of Biscay to the south of the Canaries (Mortensen, 1927), the western Mediterranean (Cherbonnier, 1956, 1958; Tortonese, 1956, 1965; Cherbonnier & Guille, 1967), the central Mediterranean (Marenzeller, 1893; Kaspiris & Tortonese, 1982), and the Adriatic (Zavodnik, 1972). It is not known in the Levantine basin.

Ophioderma longicaudum (Retzius, 1805)

Material: 147 specimens; stas 2, 17, 20, 22, 29, 68, 83, 107, 110, 116, 125, 126, 131, 137, 142, 146, 154, 167, 169, 174, 188a, 192, 205, 209, 219b, C and I; depth 0-30 m; on rocks, biogenic detritus, and sandy and silty sand substrata, and meadows of *Zostera* and *Posidonia*; Ddmax. = 28.8 mm.

Distribution: Known from various localities of the Aegean Sea (Forbes, 1844, 1845, as *O. lacertosa*; Tortonese, 1946, 1947; Tortonese & Demir, 1960).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1956, 1961, 1966; Cherbonnier, 1958; Zavodnik, 1972, 1981; Demetropoulos & Hadjichristophorou, 1976; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989).

Ophiomyxa pentagona (Lamarck, 1816)

Material: 50 specimens; stas 92, 101, 102, 128, 131, 135, 155, 196, 203, 209, A, C, E, I, R and V; depth 0-260 m; on rocks, biogenic detritus, and sandy, silty sand and sandy silt substrata; Ddmax. = 24.2 mm.

Distribution: Known from various localities of the Aegean Sea (Forbes, 1844, 1845, as *O. lubrica*; Tortonese, 1946, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquette, 1962).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1958; Tortonese, 1961, 1966; Zavodnik, 1972; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989).

Ophiopsila aranea Forbes, 1843

Material: 2 specimens; stas C, and I; depth 16-25 m; on silty sand substratum; Ddmax. = 7.4 mm.

Distribution: Known from certain localities of the Aegean Sea (Forbes, 1844, 1845; Ostroumoff, 1896; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquette, 1962; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1895; Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1956, 1961, 1965; Cherbonnier, 1958; Zavodnik, 1967).

Ophiothrix fragilis (Abildgaard, in O.F. Müller, 1789)

Material: 1102 specimens; stas 1, 2, 15, 18, 20, 25, 26, 42, 45, 52, 54, 66, 68, 85, 87, 89, 108, 112, 117, 120, 123, 126, 127, 128, 129, 130, 131, 132, 134, 135, 137, 138, 139, 140, 141, 142, 143, 146, 147, 151, 153, 155, 157, 165, 168a, 168b, 171, 175, 178, 179, 180, 181, 182, 192, 194, 200, 203, 204, 205, 205a, 207, 208, 209, 219b, A, C, D, E, H, I, R, S, V, V1 and W; depth 0-380 m; on sandy, silty sand and sandy silt substrata, and associated with algae and sponges; Ddmax. = 17.2 mm.

Distribution: Known from certain localities of the Aegean Sea, under the names *O. rosula*, *O. fragilis*, *O.*

TABLE 1. Checklist of the Mediterranean Echinodermata and their distribution in certain geographical areas of the Mediterranean and the Black Seas, with reference to their presence in the Atlantic and the Indo-Pacific Oceans. WM = Western Mediterranean, CM = Central Mediterranean, AD = Adriatic Sea, AS = Aegean Sea (including the Sea of Marmara), LB = Levantine Basin, BS = Black Sea, AO = Atlantic Ocean, IP = Indo-Pacific Ocean (the Red Sea). Zoogeographical characterization (ZC): AM, Atlantic-Mediterranean; C, Cosmopolitan; E, Possibly endemic; LM, Lessepsian migrants. VD = Vertical distribution. Species marked with * were found in the present study. Species marked with + are reported for the first time from the Aegean Sea. Species marked with ++ are reported for the first time from the Eastern Mediterranean (Aegean Sea and Levantine Basin)

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
Crinoidea											
<i>Antedon bifida</i> (Pennant, 1777)	+	+	+	+	+	+	+	+	AM	5-200	
<i>Antedon mediterranea</i> (Lamarck, 1816)	+	+	+	+	+	+	+	E	1-420		
<i>Leptometra celtica</i> (Barrett & McAndrew, 1858)	+	+	+	+	+	+	+	AM	46-1279		
<i>Leptometra phalangium</i> (J. Müller, 1841)	+	+	+	+	+	+	+	E	40-1300		
<i>Neocomatella europea</i> (A.H. Clark, 1931)	+					+	+	AM	400-1710		
Eleutherozoa											
Asteroidea											
<i>Anseropoda lobiancoi</i> (Ludwig, 1897)	+	+	+	+	+	+	+	E	40-100		
* <i>Anseropoda placentia</i> (Pennant, 1777)								AM	10-600	5-190	
<i>Aquilonaster burtoni</i> (Gray, 1840)								LM	8-10		
* <i>Asterina gibbosa</i> (Pennant, 1777)	+	+	+	+	+	+	+	AM	0-126	0-27	
<i>Asterina ocellifera</i> (Gray, 1847)								AM	6-200		
<i>Asterina panceri</i> (Gasco, 1870)								E	0-40		
<i>Asterina phylactica</i> Emson & Crump, 1979								AM	0-2		
* <i>Astropecten aranciacus</i> (Linnaeus, 1758)								AM	1-80	1-110	
* <i>Astropecten bipinnosus</i> (Otto, 1823)								AM	1-245	1-105	
* <i>Astropecten irregularis pentacanthus</i> (Delle Chiaje, 1825)	+	+	+	+	+	+	+	AM	3-1829	1-460	
* <i>Astropecten irregularis irregularis</i> (Pennant, 1777)	+	+	+	+	+	+	+	AM	3-900		
* <i>Astropecten jonstoni</i> (Delle Chiaje, 1825)	+	+	+	+	+	+	+	E	2-12	1-90	
* <i>Astropecten platyacanthus</i> (Philippi, 1837)	+	+	+	+	+	+	+	E	2-64	1-16	
* <i>Astropecten spinulosus</i> (Philippi, 1837)	+	+	+	+	+	+	+	E	3-55	1-28	
* <i>Brisingella coronata</i> (O. Sars, 1871)	+	+	+	+	+	+	+	AM	100-2904	120-320	
<i>Ceramaster grenadensis</i> (Perrier, 1881)	+	+	+	+	+	+	+	AM	200-2210		
* <i>Chonetes longipes</i> (Retzius, 1805)	+	+	+	+	+	+	+	AM	30-1139	40	
* <i>Coscinasterias tenuispina</i> (Lamarcq, 1816)	+	+	+	+	+	+	+	AM	0-79	0-50	
* <i>Echinaster (Echinaster) sepositus sepositus</i> (Retzius, 1783)	+	+	+	+	+	+	+	AM	0-250	0-210	
* <i>Hæcilia attenuata</i> (Gray, 1840)	+	+	+	+	+	+	+	AM	1-150	1-190	
<i>Henricia cylindrella</i> (Sladen, 1883)								AM	530-1400		

TABLE 1. continued

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
* <i>Luidia ciliaris</i> (Philippi, 1837)	+	+	+	+	+	+	AM	4-400	1-90		
* <i>Luidia sarsi sarsi</i> Düben & Koren, in Düben, 1845	+	+	+	+	+	+	AM	10-1292	20-190		
<i>Marginaster capreensis</i> (Gasco, 1876)	+	+	+	+	+	+	AM	49-2487			
* <i>Marthasterias glacialis</i> (Linnaeus, 1758)	+	+	+	+	+	+	AM	0-180	1-105		
* <i>Nymphaster arenatus</i> (Perrier, 1881)	+	+	+	+	+	+	AM	225-2790			
* <i>Odontaster mediterraneus</i> (Marenzeller, 1893)	+	+	+	+	+	+	AM	24-1804	80-250		
* <i>Ophidiaster ophidianus</i> (Lamarck, 1816)	+	+	+	+	+	+	AM	0-100	40		
* <i>Peltaster placenta</i> (J. Müller & Troschel, 1842)	+	+	+	+	+	+	AM	10-500	40-180		
<i>Plutonaster bifrons</i> (W. Thompson, 1873)	+	+	+	+	+	+	AM	106-2525			
<i>Sclerasterias neglecta</i> (Perrier, 1891)	+	+	+	+	+	+	AM	160-485			
<i>Sclerasterias richardi</i> (Perrier, 1882)	+	+	+	+	+	+	AM	100-710			
* <i>Tethyaster subinermis</i> (Philippi, 1837)	+	+	+	+	+	+	AM	50-1400	40-320		
Cryptosyringida											
Ophiuroidea											
<i>Amphilepis norvegica</i> (Ljungman, 1865)							+		AM	70-2940	
<i>Amphioplus (Lymanella) laevis</i> (Lyman, 1874)							+	+	LM	0-40	
* <i>Amphipholis squamata</i> (Delle Chiaje, 1828)							+	+	C	0-740	0-210
*+ <i>Amphiura (Acrocnida) brachiatia</i> (Montagu, 1804)							+	+	AM	5-40	5.3
<i>Amphiura cherbonnierii</i> Guille, 1972							+	+	E	12-130	2-250
* <i>Amphiura chiaiei</i> Forbes, 1843							+	+	AM	9-1200	
<i>Amphiura delamarei</i> Cherbonnier, 1958							+	+	E	43-200	
* <i>Amphiura filiformis</i> (O. F. Müller, 1776)							+	+	AM	5-1200	22-42
<i>Amphiura incana</i> Lyman, 1879							+	+	AM	18-58	
<i>Amphiura lacazei</i> Guille, 1976							+	+	E	20-175	
* <i>Amphiura mediterranea</i> Lyman, 1882							+	+	AM	9-90	0-25
<i>Amphiura (Ophiopeltis) securigena</i> (Düben & Koren, 1846)	+	+	+	+	+	+	+	+	E	40-600	
* <i>Amphiura stepanovi</i> Dyakonov, 1954							+	+	AM	70-205	
<i>Astrospartus mediterraneus</i> (Risso, 1826)							+	+	E	50-188	
<i>Cryptopelta brevispina</i> (Ludwig, 1879)							+	+	E	20-145	
*+ <i>Monamphiura apicula</i> (Cherbonnier, 1957)							+	+	E	15-170	2-8
* <i>Ophiacantha setosa</i> (Retzius, 1805)							+	+	AM	50-1480	5-255

TABLE 1. continued

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
<i>Ophiacts balli</i> (W. Thompson, 1840)	+							+	AM	50-1765	
<i>Ophiacts macrolepidota</i> Marktanner-Turneretscher, 1887					+	+	+	+	LM	0-200	
<i>Ophiacts savignyi</i> (J. Müller & Troschel, 1842)	+				+	+	+	+	C	2-518	
<i>Ophiacts virens</i> (M. Sars, 1857)	+	+						AM	0-90		
<i>Ophioconina nigra</i> (Abildgaard, in O. F. Müller, 1789)	+							AM	0-400		
<i>Ophioconis forbesi</i> (Heller, 1863)	+							AM	20-200		
<i>Ophiocten abyssicolum</i> Marenzeller, 1893	+							AM	40-620		
* <i>Ophioderma longicaudum</i> (Retzius, 1805)								AM	0-70	0-30	
* <i>Ophiomyxa pentagona</i> (Lamarck, 1816)								AM	0-250	0-260	
<i>Ophiopsis annulosa</i> (M. Sars, 1857)								AM	10-100		
* <i>Ophiopsis aranea</i> Forbes, 1843								AM	9-185	16-25	
* <i>Ophiopsis aff. guineensis</i> Kochler, 1914								AM	27-175		
* <i>Ophiothrix fragilis</i> (Abildgaard, in O. F. Müller, 1789)								AM	0-1244	0-380	
<i>Ophiura aifica</i> (Koehler, 1923)								AM	20-60		
* <i>Ophiura albida</i> Forbes, 1839								AM	4-850	2-26	
<i>Ophiura (Dichtenophiura) carnea</i> Lütken, 1858 ex M. Sars MS	+							AM	30-1260		
* <i>Ophiura grisea</i> Heller, 1863								AM	10-187	1-30	
* <i>Ophiura ophiura</i> (Linnaeus, 1758)								AM	0-685	0-350	
<i>Pectinura vestita</i> Forbes, 1843								E	160		
Echinozoa											
Echinoidea											
* <i>Arbacia lixula</i> (Linnaeus, 1758)								AM	0-40	0-50	
*+ <i>Arbaciella elegans</i> Mortensen, 1910								AM	3-40	1-15	
* <i>Briassopsis lyra</i> (Forbes, 1841)								AM	5-1500	4-20	
* <i>Briassopsis atlantica mediterranea</i> Mortensen, 1913								AM	100-3200	2-105	
* <i>Briissus unicolor</i> (Leske, 1778)								AM	0-250	1-12	
* <i>Centrostephanus longispinus</i> (Philippi, 1845)								AM	40-208	40-60	
* <i>Cidaris cidaris</i> (Linnaeus, 1758)								AM	50-2000	20-250	
* <i>Echinocardium cordatum</i> (Pennant, 1777)								C	0-230	1-70	
<i>Echinocardium fennaxi</i> Péquignat, 1963								E	20-50		
<i>Echinocardium flavescens</i> (O. F. Müller, 1776)								AM	5-360		
* <i>Echinocardium mediterraneum</i> (Forbes, 1843)								AM	3-40	2-35	

TABLE 1, continued

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
<i>Echinocardium mortenseni</i> Thiéry, 1909	+	+	+	+	+	+	+	+	E	10-70	1-70
* <i>Echinocytamus pusillus</i> (O. F. Müller, 1776)								AM	0-1250		
* <i>Echinus acutus</i> Lamarck, 1816								AM	20-1280	3-350	
* + <i>Echinus melo</i> Olivi, 1792								AM	25-1100	20-350	
* <i>Genocidaris maculata</i> A. Agassiz, 1869								AM	12-500	20-22	
* + <i>Hemimaster expurgatus</i> Lovén, 1874								AM	400-3120	1230-1249	
<i>Neolampris rostellata</i> A. Agassiz, 1869								AM	95-1260		
* <i>Paracentrotus lividus</i> (Lamarck, 1816)								AM	0-80	0-90	
* <i>Plagiotrius costai</i> (Gasco, 1876)								AM	25-200	20	
* <i>Psammechinus microtuberculatus</i> (Blainville, 1825) Heller, 1868								E	4-685	1-110	
* <i>Schizaster canaliferus</i> (Lamarck, 1816)								E	9-100	10-105	
<i>Spatangus inermis</i> Mortensen, 1913								E	20-350		
* <i>Spatangus purpureus</i> (O. F. Müller, 1776)								AM	15-969	18-50	
* <i>Sphaerechinus granularis</i> (Lamarck, 1816)								AM	3-100	2-120	
* <i>Stylocidaris affinis</i> (Philippi, 1845)								AM	30-1000	5-180	
Holothuroidea											
<i>Aslia lefevrei</i> (Barrois, 1882)								AM	6-20		
<i>Havelockia inermis</i> (Heller, 1868)								AM	6-180		
<i>Hedingia mediterranea</i> (Bartolini-Baldelli, 1914) Tortonese, 1965	+	+	+	+	+	+	+	E	800-1005		
<i>Holothuria (Panningothuria) forstkali</i> Delle Chiaje, 1823	+	+	+	+	+	+	+	AM	1-100		
<i>Holothuria (Holothuria) helleri</i> Marenzeller, 1878	+	+	+	+	+	+	+	E	0-80		
<i>Holothuria (Thymioscyta) impatiens</i> (Forskål, 1775)	+	+	+	+	+	+	+	+ C	0-30		
<i>Holothuria (Roweothuria) lenitignosa</i> Marenzeller, 1893	+	+	+	+	+	+	+	AM	100-250		
<i>Holothuria (Holothuria) mammata</i> Grube, 1840	+	+	+	+	+	+	+	AM	1-77		
<i>Holothuria (Lessonothuria) politi</i> Delle Chiaje, 1823	+	+	+	+	+	+	+	AM	0-250		
<i>Holothuria (Platyperona) sanctori</i> Delle Chiaje, 1823	+	+	+	+	+	+	+	AM	2-30		
<i>Holothuria (Holothuria) stellata</i> Delle Chiaje, 1823	+	+	+	+	+	+	+	E	3-65		
<i>Holothuria (Holothuria) tubulosa</i> Gmelin, 1788	+	+	+	+	+	+	+	AM	0-100		
<i>Labidoplax digitata</i> (Montagu, 1815)	+	+	+	+	+	+	+	AM	10-540		
<i>Labidoplax media</i> Ostergren, 1905	+	+	+	+	+	+	+	AM	0-268		
<i>Labidoplax thomsoni</i> (Herapath, 1865)	+	+	+	+	+	+	+	AM	5-95		
								E	7-70		

TABLE 1. continued

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
<i>Leptopentacta elongata</i> (Düben & Koren, 1844)	+	+	+	+	+			+		AM 0-150	
<i>Leptopentacta tergestina</i> (M. Sars, 1857)	+	+	+	+				+	AM 8-170		
<i>Leptosynapta decaria</i> (Östergren, 1905)			+	+				+	AM 2-70		
<i>Leptosynapta gallienii</i> (Herapath, 1865)		+	+	+				+	AM 5-30		
<i>Leptosynapta inhaerens</i> (O. F. Müller, 1776)	+	+	+	+	+	+		+	AM 2-173		
<i>Leptosynapta makranryra</i> (Ludwig, 1898)		+	+	+	+				E 1-36		
<i>Leptosynapta minuta</i> (Becher, 1906)		+	+	+	+			+	AM 3-50		
<i>Mesothuria intestinalis</i> (Ascanius, 1805) Östergren, 1896	+	+	+	+	+	+	+	AM 18-4255			
<i>Mesothuria verrilli</i> (Théel, 1886)	+	+	+	+	+	+	+	AM 280-2520			
<i>Molpadiia musculus</i> Risso, 1826								AM 25-2098			
<i>Myioirochus geminiradiatus</i> Salvini-Plawen, 1972								E 70-225			
<i>Neocnus incubans</i> Cherbonnier, 1972	+	+	+	+				E 1-?			
<i>Neocucumis atlantica</i> (Ludwig & Heding, 1935)		+	+	+				AM 50-300			
<i>Neocucumis marioni</i> (Marenzeller, 1878)		+	+	+	+	+	+	AM 25-560			
<i>Ocnus grubei</i> (Marenzeller, 1874)					+	+	+	AM 3-40			
<i>Ocnus koellikeri</i> (Semper, 1867)					+	+	+	AM 50-685			
<i>Ocnus lacteus</i> (Forbes & Goodsir, 1839)							+	AM 0-100			
<i>Ocnus petitii</i> (Cherbonnier, 1958)							+	AM 30-35			
<i>Ocnus planci</i> (Brandt, 1835)		+	+	+	+	+	+	AM 5-250			
<i>Ocnus syracusanus</i> (Grube, 1840) Panning, 1949		+	+	+	+	+	+	AM 7-100			
<i>Panningia hyndmani</i> (W. Thompson, 1840)		+	+	+	+	+	+	AM 7-1152			
<i>Parastichopus regalis</i> (Cuvier, 1817)		+	+	+	+	+	+	AM 5-1200			
<i>Pawsonia saxicola</i> (Brady & Robertson, 1871)		+	+	+	+	+	+	AM 0-130			
<i>Peniliptida ludwigi</i> (Marenzeller, 1893)					+	+	+	AM 755-4766			
<i>Phyllophorus drachi</i> Cherbonnier & Guille, 1968							+	E ?90			
<i>Phyllophorus granulatus</i> (Grube, 1840)							+	E 3-15			
<i>Phyllophorus (Phyllophorus) urma</i> Grube, 1840							+	E 2-150			
<i>Prototrochus meridionalis</i> (Salvini-Plawen, 1977)							+	E ?540			
<i>Pseudostichopus occultatus</i> Marenzeller, 1893							+	C 232-5300			
<i>Pseudohyone raphanus</i> (Düben & Koren, 1846)							+	AM 7-1050			
<i>Pseudohyone sculptorea</i> Cherbonnier, 1928							+	E 25-120			
<i>Stereoderma kirschbergi</i> (Heller, 1868)							+	AM 30-80			

TABLE 1. continued

Mediterranean species	WM	CM	AD	AS	LB	BS	AO	IP	ZC	VD (m) (literature)	VD (m) (present data)
* <i>Synaptula reciprocans</i> (Forskål, 1775)					+	+		+	LM	1-20	0-4
<i>Thyone cherbonnierii</i> Reys, 1960	+		+	+					E	3-63	
<i>Thyone fusus mediterranea</i> Madsen, 1941	+	+	+	+					E	20-150	
<i>Thyone gadeana</i> R. Perrier, 1902	+							+	AM	562-1045	
<i>Trochodota fuscipraedita</i> Salvini-Plawen, 1972	+								E	4-5	
<i>Trochodota venusta</i> (Semon, 1887)	+								E	1-13	
Total number of species: 154	144	91	100	108	72	10	110	9			

fragilis echinata and *O. quinquemaculata* (Forbes, 1844; Ostroumoff, 1896; Panagiotopoulos, 1916; Athanasioupolous, 1917; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquette, 1962; Caspers, 1968; Vamvakas, 1971; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1956, 1961, 1966; Cherbonnier, 1958; Zavodnik, 1972; Pancucci & Zenetos, 1989), as well as from the Black Sea (D'yakonov, 1954; Tortonese & Demir, 1960).

Ophiura albida Forbes, 1839

Material: 16 specimens; stas 97, 109, 113, 154, 156, 157, and I; depth 2-26 m; on sandy and silty sand substrata; Ddmax. = 8.3 mm.

Distribution: Known from many localities of the Aegean Sea, (Forbes, 1844, 1845; Ostroumoff, 1896; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquette, 1962; Kisseeleva, 1963; Makkavieva, 1963; Vamvakas, 1971; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Tortonese, 1956, 1961; Cherbonnier, 1958; Zavodnik, 1972; Kaspiris & Tortonese, 1982; Özaydin *et al.*, 1995).

Ophiura grubei Heller, 1863

Material: 15 specimens; stas 154, C, I and J; depth 1-30 m; on sandy and silty sand substrata; Ddmax. = 5.2 mm.

Distribution: This species has been known in the Aegean Sea only from the Sea of Marmara (Tortonese & Demir, 1960), the Saronikos gulf (Vamvakas, 1971) and Lesvos island (Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean region (Tortonese, 1956, 1965; Zavodnik, 1972; Pancucci-Papadopoulou, 1996) as well as from the Atlantic coasts of Morocco (Mortensen, 1927).

Ophiura ophiura (Linnaeus, 1758)

Material: 46 specimens; stas 78, 87, 107, 201, 222, A, E, F and W; depth 0-350 m; on rocks, organic detritus, sandy and sandy silt substrata, and meadows of *Zostera* and *Posidonia*; Ddmax. = 27.6 mm.

Distribution: Known from various localities of the Aegean Sea under the name *O. lacertosa*, *O. texturata* and *O. ophiura* (Forbes, 1844, 1845; Ostroumoff, 1896; Tortonese, 1946; Pérès & Picard, 1958; Tortonese & Demir, 1960, Kisseeleva, 1961; Jacquette, 1962; Vamvakas, 1971; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1956, 1958; Tortonese, 1956, 1961, 1965; Zavodnik, 1972; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989; Özaydin *et al.*, 1995).

3. Eleutherozoa – Cryptosyringida – Echinozoa – Echinoidea

The following 21 species were found in the Aegean Sea during the present study.

Arbacia lixula (Linnaeus, 1758)

Material: 115 specimens; stas 15, 25, 29, 32, 42, 45, 76, 83, 85, 86, 88, 89, 109, 110, 114, 115, 117, 120, 122, 128, 133, 135, 137, 138, 139, 140, 141, 146, 159, 165, 169, 172, 173, 174, 175, 177, 178, 179, 188, 189, 196, 203, 208, 209 and C; depth 0-50 m; on rocks, organic detritus, as well as on live leaves of *Posidonia*; Dmax. = 56.5 mm.

Distribution: Known from various localities of the Aegean Sea under the name *A. aequituberculata*, *A. pustulosa* and *A. lixula* (Ostroumoff, 1896; Issel, 1928; Tortonese, 1946; Pérès & Picard, 1958; Geldiay & Koçatas, 1972; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Cherbonnier, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Kaspiris & Tortonese, 1982).

Arbaciella elegans Mortensen, 1910

Material: 20 specimens; stas 15, 20, 41, 130, 142, 147, 172, 173, 188a, 203, 205, and 208; depth 1-15 m; under rocks and on organic detritus; Dmax. = 7.6 mm.

Distribution: This species is reported for the first time from the Eastern Mediterranean (Aegean Sea and Levantine Basin). Not known from the Adriatic Sea.

An Atlanto-Mediterranean species (Table 1). In the Atlantic it occurs on the African west coast, but it is not known to the north of Cape Blanc (Mortensen, 1927). It is known from a few localities of the

Western Mediterranean (Gautier-Michaz, 1958; Tortonese, 1965; Galán-Novella & López-Ibor Alíño, 1981) and from Marsaxlokk Bay, Malta, in the Central Mediterranean (Schembri, 1978).

Brissopsis lyrifera (Forbes, 1841)

Material: 4 specimens; stas 45, 109 and 210; depth 4-20 m; in sandy substrata; Dmax. = 76.2 mm.

Distribution: Known from various localities of the Aegean Sea (Marenzeller, 1895; Ostroumoff, 1896; Pérès & Picard, 1958; Tortonese & Demir, 1960; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1958; Pérès & Picard, 1958; Tortonese, 1958, 1965; Zavodnik, 1972).

Brissopsis atlantica mediterranea Mortensen, 1913

Material: 17 specimens; stas 23, 109, C, D, E, J, and M; depth 2-105 m; in sandy, silty sand and sandy silt substrata; Dmax. = 72.0 mm.

Distribution: The presence of this species in the Aegean Sea has been documented only from the north of Lesvos island (Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from the Atlantic (Mortensen, 1927; Tortonese, 1965), the western Mediterranean (Tortonese, 1965; Rodríguez, 1979) and the Adriatic (Zavodnik, 1980). It is not known from the central Mediterranean and the Levantine basin.

Brissus unicolor (Leske, 1778)

Material: 20 specimens; stas 15, 34, 68a, 110 and 171; depth 1-12 m; in sandy substrata; Dmax. = 102.0 mm.

Distribution: This species has been known in the Aegean Sea only from the Messiniakos gulf-Kalamata and the Kyklades islands – Nata (Pérès & Picard, 1958).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1956, 1958; Tortonese, 1965; Zavodnik, 1972; Schembri, 1978; Rodríguez, 1979; Kaspiris & Tortonese, 1982).

Centrostephanus longispinus (Philippi, 1845)

Material: 3 specimens; sta 77; depth 40-60 m; on biogenic detritus on silty sand substratum; Dmax. = 35.4 mm.

Distribution: This species has been known in the Aegean Sea from various localities (Tortonese, 1946, 1947; Pérès & Picard, 1958; Laborel, 1960; Tortonese & Demir, 1960; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Cherbonnier, 1956; Tortonese, 1965; Zavodnik, 1972; Schembri, 1978; Rodríguez, 1979; Kaspiris & Tortonese, 1982; Özaydin et al., 1995).

Cidaris cidaris (Linnaeus, 1758)

Material: 30 specimens; stas 77, 175, M, Q and S; depth 20-250 m; on rocks, biogenic detritus, and silty sand and sandy silt substrata; Dmax. = 38.8 mm.

Distribution: Known from various localities of the Aegean Sea, under the names *C. hystrix*, *Dorocidaris papillata* and *C. cidaris* (Spratt & Forbes, 1842; Forbes, 1844; Steindachner, 1891; Marenzeller, 1893, 1895; Ostroumoff, 1896; Pérès & Picard, 1958; Kisileva, 1983; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1893; Mortensen, 1927; Cherbonnier, 1956, 1958; Tortonese, 1965; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Schembri, 1978; Özaydin et al., 1995).

Echinocardium cordatum (Pennant, 1777)

Material: 59 specimens; stas 23, 68a, 92, 95, 147, 159, 163, 171, J and K; depth 1-70 m; in sandy and silty sand substrata; Dmax. = 48.5 mm.

Distribution: This species has been known in the Aegean Sea from the Sea of Marmara (Ostroumoff, 1896), Syros island (Marion, 1898), the Izmir gulf (Geldiay & Koçatas, 1972), and Lesvos island (Pancucci & Zenetos, 1990).

A cosmopolitan species (Table 1) known from all the Mediterranean areas (Mortensen & Steuer, 1937; Tortonese, 1956, 1965; Cherbonnier, 1958; Pérès & Picard, 1958; Zavodnik, 1972; Rodríguez, 1979; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989), the Atlantic (Mortensen, 1927; Tortonese, 1965) and the Pacific Oceans (Mortensen, 1927; Tortonese, 1965).

Echinocardium mediterraneum (Forbes, 1843)

Material: 24 specimens; stas 125, 147, 207 and K; depth 2-35 m; in sandy and silty sand substrata; Dmax. = 42.5 mm.

Distribution: This species has been known in the Aegean Sea only from Poros island (Forbes, 1844) and the Sea of Marmara (Tortonese & Demir, 1960).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Tortonese, 1956, 1965; Cherbonnier, 1958; Zavodnik, 1972; Rodríguez, 1979; Pancucci & Zenetos, 1989).

Echinocyamus pusillus (O.F. Müller, 1776)

Material: 21 specimens; stas 107, 108, J and V; depth 1-70 m; in sandy and silty sand substrata; Dmax. = 8.6 mm.

Distribution: Known from many localities of the Aegean Sea (Forbes, 1844; Ostroumoff, 1896; Marion, 1898; Tortonese, 1946; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Vamvakas, 1971; Kisseleva, 1983; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Raulin, 1870, as *E. tarentinus*; Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1956, 1958, 1969; Tortonese, 1956, 1965; Zavodnik, 1972, 1981; Pancucci & Zenetos, 1989).

Echinus acutus Lamarck, 1816

Material: 53 specimens; stas 79, 130, B, E, G, H, K, Q, R, S and V; depth 3-350 m; on organic detritus, sandy and sandy silt substrata; Dmax. = 152.0 mm.

Distribution: Known from certain localities of the Aegean Sea under the names *E. norvegicus*, *E. melo* and *E. acutus* (Steindachner, 1891; Marenzeller, 1893, 1895; Tortonese & Demir, 1960; Makkavieva, 1963).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1893; Mortensen, 1927; Cherbonnier, 1956, 1958; Tortonese, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Schembri, 1978; Rodríguez, 1979; Kaspiris & Tortonese, 1982).

Echinus melo Olivi, 1792

Material: 52 specimens; stas 71, 79, 109, B, E, G, H, K, M, Q, S and V; depth 20-350 m; on rocks, organic detritus and silty sand substrata; Dmax. = 130.4 mm.

Distribution: This species is reported for the first time from the Eastern Mediterranean (Aegean Sea and Levantine basin).

An Atlanto-Mediterranean species (Table 1) also known from the European Atlantic coasts (Mortensen, 1927), the western Mediterranean (Cherbonnier, 1958; Tortonese, 1965; Rodríguez, 1979), the central Mediterranean (Marenzeller, 1893; Schembri, 1978) and the Adriatic (Zavodnik, 1972).

Genocidaris maculata A. Agassiz, 1869

Material: 1 specimen; sta 197; depth 20-22 m; on calcareous algae; D = 8.0 mm.

Distribution: Known from various localities of the Aegean Sea (Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Vamvakas, 1971).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Tortonese, 1956; Cherbonnier, 1958; Zavodnik, 1972; Pancucci & Zenetos, 1989).

Hemaster expurgatus Lovén, 1874

Material: 3 specimens; sta Z1, 39° 16.58' N, 23° 42.49' E & 39° 15.51' N, 23° 42.51' E; depths 1249 m & 1230 m; in silty substratum; Dmax. = 32.6 mm.

Distribution: This species is reported for the first time from the Eastern Mediterranean (Aegean Sea and Levantine basin).

An Atlanto-Mediterranean species (Table 1) known in the Atlantic from the Norwegian coast and the south of Iceland to the Azores and the Cape Verde island and from the Davies Strait to the West Indies (Mortensen, 1927). It has been known in the Mediterranean only from Mallorca – Baleares (Cherbonnier, 1958) and the gulf of Genova (Tortonese, 1972).

Paracentrotus lividus (Lamarck, 1816)

Material: 331 specimens; stas 4, 15, 16, 20, 23, 25, 28, 29, 31, 34, 35, 42, 45, 47, 48, 49, 53, 59, 60, 68, 76, 85, 88, 89, 91, 93, 95, 102, 104, 108, 109, 110, 112, 115, 118, 122, 123, 127, 128, 130, 131, 135, 138, 140, 141, 142, 146, 147, 151, 156, 157, 159, 161, 162, 163, 168a, 169, 171, 172, 173, 174, 175, 177, 178, 179, 188, 189, 192, 193, 196, 199, 200, 203, 205, 207, 208, 209, C, I and K; depth 0-90 m; on rocks, organic detritus and silty sand substrata, as well as on live leaves of *Posidonia* (personal observation); Dmax. = 62.4 mm.

Distribution: Known from many localities of the Aegean Sea, under the names *Echinus (Toxopneustes) lividus*, *Strongylocentrotus lividus* and *P. lividus* (Spratt & Forbes, 1842; Forbes, 1844; Raulin, 1870; Ostrou-

moff, 1896; Panagiotopoulos, 1916; Issel, 1928; Tortonese, 1946, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Vamvakas, 1971; Pancucci & Zenetos, 1990).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1958; Tortonese, 1965; Zavodnik, 1972; Schembri, 1978; Pastore, 1981; Pancucci & Zenetos, 1989).

Plagiobrissus costai (Gasco, 1876)

Material: 1 specimen; sta 109; depth 20 m; in silty sand substratum; D = 80.2 mm.

Distribution: This species has been known in the Aegean Sea only from the Messiniakos gulf-Kalamata (Jacquotte, 1962; as *Metalia costae*) and the west of Karavi islet, close to Kythira island (Pérès & Picard, 1958; as *Melita costae*).

An Atlanto-Mediterranean species (Table 1) known from the eastern Atlantic (Tortonese, 1965), the western Mediterranean (Tortonese, 1965), the central Mediterranean (Cherbonnier, 1956; Tortonese, 1961), and the Levantine basin (Mortensen & Steuer, 1937; Tortonese, 1956).

Psammechinus microtuberculatus (Blainville, 1825) Heller, 1868

Material: 131 specimens; stas 20, 22, 77, 79, 93, 95, 96, 105, 106, 109, 110, 128, 156, 194, 196, 199, 200, 201, 207, 208, C, E, V and V1; depth 1-110 m; on organic detritus, silty sand and sandy silt substrata, as well as on leaves of *Posidonia* and algae; Dmax. = 29.8 mm.

Distribution: Known from certain localities of the Aegean Sea, under the names *Echinus microtuberculatus* and *Psammechinus microtuberculatus* (Ostroumoff, 1896; Marion, 1898; Tortonese & Demir, 1960; Kisileva, 1983; Pancucci & Zenetos, 1990; Özaydin et al., 1995).

A Mediterranean endemic species (Table 1) known from all over the Mediterranean region (Cherbonnier, 1956, 1958; Tortonese, 1961; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Schembri, 1978; Kaspiris & Tortonese, 1982; El-Beshbeeshy, 1995).

Schizaster canaliferus (Lamarck, 1816)

Material: 10 specimens; stas 15, 108, C, E and I; depth 10-105 m; in sandy and silty sand substrata; Dmax. = 63.2 mm.

Distribution: Known from certain localities of the Aegean Sea (Forbes, 1844, as *Brissus atropos*; Ostroumoff, 1896; Marion, 1898; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Pancucci & Zenetos, 1990; Özaydin et al., 1995).

A Mediterranean endemic species (Table 1) known from all the Mediterranean regions (Mortensen & Steuer, 1937; Cherbonnier, 1956, 1958; Tortonese, 1956, 1965; Zavodnik, 1972; Schembri, 1978; Kaspiris & Tortonese, 1982).

Spatangus purpureus (O.F. Müller, 1776)

Material: 2 specimens; stas 80 and 175; depth 18-50 m; in silty sand and sandy silt substrata; Dmax. = 108.5 mm.

Distribution: Known from certain localities of the Aegean Sea (Forbes, 1844; Marenzeller, 1893, 1895; Ostroumoff, 1896; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Vamvakas, 1971; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Marenzeller, 1893; Mortensen, 1927; Cherbonnier, 1956, 1958; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Schembri, 1978; Kaspiris & Tortonese, 1982).

Sphaerechinus granularis (Lamarck, 1816)

Material: 47 specimens; stas 40, 60, 82, 83, 85, 92, 105, 109, 110, 180, 191, 196, 200, 201, 203, C, M and T; depth 2-120 m; on rocks and organic detritus, as well as on sandy (meadows of *Posidonia*) and silty sand substrata; Dmax. = 79.6 mm.

Distribution: Known from various localities of the Aegean Sea (Ostroumoff, 1896; Athanassopoulos, 1921; Tortonese, 1946, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquotte, 1962; Vamvakas, 1971; Pancucci & Zenetos, 1990; Özaydin et al., 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Cherbonnier, 1956, 1958; Tortonese, 1965; Zavodnik, 1972; Demetropoulos & Hadjichristophorou, 1976; Schembri, 1978; Kaspiris & Tortonese, 1982; Özaydin et al., 1995).

Stylocidaris affinis (Philippi, 1845)

Material: 43 specimens; stas 45, 77, 78, 130, 175, H, S and T; depth 5-180 m; on organic detritus, calcareous

algae, and silty sand and sandy silt substrata; Dmax. = 46.8 mm.

Distribution: Known from various localities of the Aegean Sea (Tortonese, 1946, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Jacquette, 1962; Makkavieva, 1963; Vamvakas, 1971; Özaydin *et al.*, 1995).

An Atlanto-Mediterranean species (Table 1) known from all over the Mediterranean and the Atlantic regions (Mortensen, 1927; Mortensen & Steuer, 1937; Cherbonnier, 1956; Tortonese, 1956, 1961, 1965; Zavadnik, 1972; Schembri, 1978; Kaspiris & Tortonese, 1982).

4. Eleutherozoa – Cryptosyringida – Echinozoa – Holothuroidea

The results of intensive samplings in the Aegean concerning crinoids and holothuroids have been presented by Koukouras & Sinis (1981) and are included in Table 1.

Pancucci-Papadopoulou (1996) when listing the Echinodermata of Greece, included among the holothuroids the species *Echinocucumis hispida* (Barrett, 1857) (as *E. typical*, Sars, 1861), based on a record from the gulf of Kavala, at a depth of 40 m (Papazacharias, 1991). However, the specimen by Papazacharias (1991) which is mentioned as deposited at the Zoological Museum of the Department of Zoology, Aristotle University of Thessaloniki has been lost. Furthermore, this species is known only from the Central Skagerrak; Norway, between Arendal and N Finnmark; off SW Faeroes; off S and W Iceland; off W Ireland and the Biscay gulf; at depths 50-1430 m (Mortensen, 1927; Hansson, 1999). Fredj (1974) included *E. hispida* (as *E. typica*) in his list of the western Mediterranean benthic invertebrates, but without any documentation. For the above reasons, the record of *E. hispida* from the gulf of Kavala has to be considered as erroneous.

During recent sampling, the following holothroid species was found in the Aegean Sea, at the coast of Megisti island (Kastellorizo) and the coasts of Cyprus.

Synaptula reciprocans (Forskål, 1775)

Material: 10 specimens; stas 218b, 219b, 219c, 239 (Megisti Island) and Cyprus – Lemesos – Zigi; depth 0-4 m; on sandy substrata; Dmax. = 450.0 mm.

Distribution: This species was known from the Red Sea and the SE Arabia (Clark & Rowe, 1971). Tortonese (1947) reported *Synaptula reciprocans*

from the Suez Canal (Lake Timsah). Later, Tortonese (1979) noted “*In Antalya (South Anatolia) I was recently told that dark and rather large synaptids were seen near the shore but unfortunately were not collected: they might be Synaptula reciprocans, a Red Sea species quite recently observed along the Israeli shores*”. However, Cherbonnier (1986) reported for the first time its presence in the Mediterranean (two specimens from the Akrotiri gulf, Cyprus, and two more from Nachsholin and Dor, Israel), although he did not attribute it to Lessepsian migration, as Zibrowius (1991) noted. Recently, this species was reported from the coast of Lebanon (Zibrowius & Bitar, 2003), the SW coast of Turkey (Çinar *et al.*, 2006) and the SE Aegean Sea (Zaitsev & Öztürk, 2001).

In any case, the specimens collected from the area Zigi – Lemesos confirm the presence of *S. reciprocans* in the south coasts of Cyprus.

According to Koukouras & Sinis (1981) two species of Crinoidea are known from the Aegean Sea (Table 1).

Concerning Asteroidea, five more species are known from the Aegean Sea (Table 1): *Asterina panceri* (Tortonese & Demir, 1960), *Ceramaster grenadensis grenadensis* (Marenzeller, 1893, 1895; as *Pentagonaster hystricis*), *Marginaster capreensis* (Marenzeller, 1893, 1895), *Sclerasterias neglecta* (Marenzeller, 1895; as *Stolasterias neglecta*) and *Sclerasterias richardi* (Marenzeller, 1893, 1895; as *Asterias richardi*). Including the above five species, the number of the known Asteroidea from the Aegean Sea is 25.

In respect of Ophiuroidea, ten more species are known from the area (Table 1): *Amphilepis norvegica* (Forbes, 1845; as *Amphiura florifera*), *Amphiura cherbonnieri* (Pancucci & Zenetos, 1990), *Amphiura lacazei* (Pancucci-Papadopoulou, 1996), *Astrospartus mediterraneus* (Belloc, 1948), *Ophiactis savignyi* (Pancucci-Papadopoulou, 1996), *Ophioconis forbesi* (Pérès & Picard, 1958), *Ophiocten abyssicolum* (Forbes, 1845, as *Ophiura abyssicola*; Marenzeller, 1893), *Ophiopsila annulosa* (Pérès & Picard, 1958), *Ophiura (Dictenophiura) carnea* (Marenzeller, 1893; as *Ophioglypha carnea*), and *Pectinura vestita* (Forbes, 1844). Including the above ten species, the number of the known Ophiuroidea from the Aegean Sea is 24.

Concerning Echinoidea, three more species are known from the Aegean Sea (Table 1): *Echinocardium fenauxi* (Borghi, 1994), *Echinocardium flavescent* (Pérès & Picard, 1958) and *Spatangus inermis* (Özaydin *et al.*, 1995). Thus, with these three species, the number of the known Echinoidea from the Aegean

Sea is 24.

Koukouras & Sinis (1981) showed the presence of 30 Holothuroidea species in the Aegean Sea. Pancucci & Zenetos (1990) reported *Neocucumis marioni* and Pancucci (1994), *Leptosynapta minuta* from this area. In this study, *Synaptula reciprocans* is also reported (Table 1). Thus, the total number of the known Holothuroidea species from the Aegean Sea rises to 33.

Comparison of the Aegean fauna with the faunas of other neighbouring seas

Tortonese (1979) has numbered 143 Mediterranean echinoderm species (5 Crinoidea, 30 Asteroidea, 34 Ophiuroidea, 26 Echinoidea, 48 Holothuroidea). The review of the relevant literature showed that, up to day, 154 valid species (5 Crinoidea, 33 Asteroidea, 36 Ophiuroidea, 26 Echinoidea, 54 Holothuroidea) have been known from the Mediterranean and the Black Seas. Their distribution over the geographical areas of the Mediterranean and their presence in the Black and Red Seas, and the Atlantic Ocean, as well as their depth range according to the literature and present data, are given in Table 1. The distribution of the known echinoderm species in the main geographical

areas of the Mediterranean Sea and the Black Sea (as real numbers and percentages of the total Mediterranean species) as it results from the present study, is given in Table 2 and Fig. 2. Taking into account Tables 1 and 2, Fig. 2, data on the Mediterranean water masses and circulation (Ovchinnikov, 1966; The POEM Group, 1992; Perivoliotis *et al.*, 1997) along with data on temperature and salinity variations (Lacombe *et al.*, 1958; Özsoy *et al.*, 1993) and geographical aspects (Bianchi & Morri, 2000; Pinaridi & Masetti, 2000), the following considerations can be made:

Western Mediterranean (WM): 144 species, 93.5% of the known Mediterranean fauna (Ludwig, 1897; Koehler, 1927; Bartolini-Baldelli, 1914; Clark, 1931; Panning, 1949; Tortonese, 1952, 1965; Cherbonnier, 1956, 1958, 1968, 1972; Cherbonnier & Guille, 1967, 1968; Salvini-Plawen, 1972, 1977; Lopez-Ibor Aliño & Galan-Novella, 1982; Guille *et al.*, 1983; Pérez-Ruzafa & López-Ibor, 1986; Burke *et al.*, 1988). In the western Mediterranean, only ten Mediterranean species have not been found: *Aquilonaster burtoni* (asteroid, Lessepsian migrant), *Amphioplus (Lymanella) laevis* (ophiuroid, Lessepsian migrant), *Amphiura stepanovi* (ophiuroid, Black Sea endemic), *Ophiactis macrolepidota* (ophiuroid, Lessepsian migrant), *Pect-*




FIG. 2. Distribution of the known species of echinoderms in the main geographical areas of the Mediterranean and the Black Seas, as real numbers and percentages of the total Mediterranean species. Concerning the Aegean Sea, the blank part of the column shows the number of species added by the present study (abbreviations as in Table 1).

TABLE 2. Species number for each one of the five echinoderm taxa separately, as well as total echinoderm species number and Mediterranean species percentage for each one of the geographical areas (abbreviations as in Table 1)

	Geographical areas and species numbers					
	WM	CM	AD	AS	LB	BS
Crinoidea	5	2	2	2	1	0
Asteroidea	32	24	23	25	22	1
Ophiouroidea	32	19	23	24	16	4
Echinoidea	26	21	19	24	17	0
Holothuroidea	49	25	33	33	16	5
Total number	144	91	100	108	72	10
(%)	(93.5)	(59.1)	(64.9)	(70.1)	(46.8)	(6.5)

inura vestita (ophiuroid, Aegean Sea endemic), *Synaptula reciprocans* (holothuroid, Lessepsian migrant), and *Leptosynapta decaria*, *L. gallienii*, *Myriotrochus geminiradiatus*, *Penilpida ludwigi* (holothuroid, known from other Mediterranean areas). The highest species number of the echinoderm fauna in the western basin of the Mediterranean (Fig. 2, Table 2) could be attributed to the fact that the influx of Atlantic species is initially limited in this large basin, which, having a wide range of physicochemical parameters, permits the settlement of both cold and warm water species in its northern and southern parts, respectively. This general trend does not seem to be strongly based on the more intensive sampling carried out in this area.

Central Mediterranean (CM): 91 species, 59.1% of the known Mediterranean fauna (Marenzeller, 1893; Cherbonnier, 1956; Pérès & Picard, 1956; Gauquier-Michaz, 1958; Tortonese, 1965; Schembri, 1978; Kaspiris & Tortonese, 1982; Pancucci & Zenetos, 1989; Fiege & Liao, 1996). Central Mediterranean comes fourth in echinoderm species number (Fig. 2, Table 2), although it should have a higher species number compared with the Aegean and Adriatic Seas, due to its direct neighbouring with the western Mediterranean. This lower number of species should be attributed to the limited sampling effort carried out in this area.

Adriatic Sea (AD): 100 species, 64.9% of the known Mediterranean fauna (Mayer, 1937; Tortonese, 1956, 1965; Zavodnik, 1960, 1972, 1979, 1980, 1995; Salvini-Plawen, 1972, 1977). The Adriatic Sea (Fig. 2, Table 2), although intensively sampled, displays a relatively low species number. This should mainly be attributed to a) its considerably restricted communica-

cation with the western basin (Ovchinnikov, 1966; Theocharis *et al.*, 1993), b) the larger amplitude of temperature variations (Delépine *et al.*, 1987), and c) the shallow waters of its northern part with relatively low winter temperatures and low salinity (Lacombe & Tchernia, 1960).

Aegean Sea (AS): 108 species, 70.1% of the known Mediterranean fauna (Forbes, 1844, 1845; Marenzeller, 1893, 1895; Ostroumoff, 1896; Tortonese, 1946, 1947; Pérès & Picard, 1958; Tortonese & Demir, 1960; Koukouras & Simis, 1981; Pancucci & Zenetos, 1990; Özaydin *et al.*, 1995; present study). Although the Aegean Sea is more distant from Gibraltar (the main pathway of enrichment for the Mediterranean fauna) (Ekman, 1967) than the Adriatic, the Aegean is inhabited by a larger number of species (Fig. 2, Table 2). The main reasons for the presence of a higher species number in the Aegean could be a) its more direct communication with the western basin (Ovchinnikov, 1996) and b) its higher habitat variability (Pérès, 1967; Koukouras *et al.*, 2001).

Levant Basin (LB): 72 species, 46.8% of the known Mediterranean fauna (Forbes, 1845; Mortensen & Steuer, 1937; Tortonese, 1956, 1965, 1966, 1979; Achituv, 1969, 1973; Demetropoulos & Hadjichristophorou, 1976; Por, 1978; Cherbonnier, 1986; Achituv & Sheer, 1991; Zibrowius, 1991; Özaydin *et al.*, 1995; Zaitsev & Öztürk, 2001; Zibrowius & Bitar, 2003). The lowest species numbers of the Levantine basin in comparison with those of other Mediterranean areas (Fig. 2, Table 2) should mainly be attributed to its impoverished fauna, a fact that is without doubt a result of the fluctuations resulting from the last glacial cycle, and much less of the present restricting circumstances, as for example the oligotro-

phic conditions (Por & Dimentman, 1989), as well as to the less intensive sampling effort carried out in the area. Moreover, from the Atlantic species inflowing in the Mediterranean through the Gibraltar Straits, few species are able to reach and settle in the Levantine Sea, where particularly unfavourable conditions prevail (Por & Dimentman, 1989; Koukouras & Russo, 1991; Koukouras *et al.*, 2001; Arvanitidis *et al.*, 2002). Conversely, the fauna of the eastern part of the Levantine Basin became enriched with four Lessepsian migrants, one of which, *Synaptula reciprocans*, has extended its distribution up to SE Aegean Sea (present study).

Black Sea (BS): 10 species, 6.5% of the known Mediterranean fauna (Caspers, 1968; Tortonese & Demir, 1960; Tortonese, 1979). The extremely low

species number of the Black Sea fauna is a result of the very peculiar oceanographic conditions prevailing in the area, especially the low salinities and temperatures (Caspers, 1957; Tortonese & Demir, 1960; Longhurst, 1998).

The total of the 154 Mediterranean echinoderm species is distributed within the five taxa as it is shown in Fig. 3 and Table 3. In Fig. 4 (based on Tables 1 and 3), the participation of the four zoogeographical categories as percentages of the total Mediterranean species is given. As it is demonstrated, most species (68.2%) have an Atlanto-Mediterranean distribution, while 26% are possibly Mediterranean endemics and only five species (3.2%) are cosmopolitan. Four species (2.6%) are Lessepsian immigrants in the Mediterranean Sea. Only one of them, *Synaptula recipro-*

TABLE 3. Species number for each one of the five echinoderm taxa separately, as well as total echinoderm species number and percentage for each zoogeographical category. The number and percentage of Mediterranean species per taxon is also given (abbreviations as in Table 1)

	Zoogeographical categories				All Mediterranean species and (%)
	AM	E	C	LM (IP)	
Crinoidea	3	2	0	0	5 (3.2)
Asteroidea	27	5	0	1	33 (21.4)
Ophiuroidea	24	8	2	2	36 (23.4)
Echinoidea	20	5	1	0	26 (16.9)
Holothuroidea	31	20	2	1	54 (35.1)
Total number (%)	105 (68.2)	40 (26.0)	5 (3.2)	4 (2.6)	154




FIG. 3. Echinoderm fauna composition in the Mediterranean and the Black Seas (percentages and real numbers) regarding the contribution of the five taxa.




FIG. 4. Echinoderm fauna composition in the Mediterranean and the Black Seas (percentages and real numbers), regarding the zoogeographical characterization of the species (abbreviations as in Table 1).




FIG. 5. Percentages of the four zoogeographical categories in the Mediterranean territorial areas and the Black Sea; calculations were made for the total of species found from each area (abbreviations as in Table 1).

cans has extended its distribution beyond the Levantine Basin to the SE Aegean Sea. The percentage of endemic species (26%) is relatively very high probably due to the sluggish locomotion of echinoderms and their relatively short pelagic stage of development. Tortonese (1985) has listed 144 Mediterranean species, 35 of which (24.3%) were characterized as endemic.

In Fig. 5 (based also on Tables 1 and 3), the percentages of the four zoogeographical categories, for the total of species known from each Mediterranean area and the Black Sea, are shown. From this Figure it is obvious that in each Mediterranean area, Atlanto-Mediterranean species dominate followed by endemic and cosmopolitan species. Furthermore, the species number (and percentage) of each zoogeographical category seems to decline from west to east, as it also applies to the total known species numbers from each Mediterranean area. The above considerations are supported by those of Koukouras *et al.* (2001) and Koukouras & Karachle (2005).

REFERENCES

- Achituv Y, 1969. Studies on the reproduction and distribution of *Asterina burtoni* Gray and *A. wega* Perrier (Asteroidea) in the Red Sea and the Eastern Mediterranean. *Israel journal of zoology*, 18: 329-342.
- Achituv Y, 1973. On the distribution and variability of the Indo-Pacific sea star *Asterina wega* (Echinodermata: Asteroidea) in the Mediterranean Sea. *Marine biology*, 18: 333-336.
- Achituv Y, Sheer E, 1991. Sexual reproduction and fission in the sea star *Asterina burtoni* from the Mediterranean coast of Israel. *Bulletin of marine science*, 48: 670-678.
- Arvanitidis Ch, Bellan G, Dracopoulos P, Valavanis V, Dounas C, Koukouras A, Eleftheriou A, 2002. Seascape biodiversity patterns along the Mediterranean and the Black Sea: lessons from the biogeography of benthic polychaetes. *Marine ecology progress series*, 244: 139-152.
- Athanassopoulos GD, 1917. Quelques éléments de recherches hydrobiologiques en Grèce: quelques crustacés. *Bulletin station hydrobiologique marine Grèce*, 2: 31-34.

- Athanassopoulos GD, 1921. Ajouts aux tableaux de faune. *Bulletin station hydrobiologique marine Grèce*, 3: 12-13.
- Bartolini-Baldelli C, 1914. Asteroidi, Ophiuroidi, Crinoidi, Olothuroidi raccolti nel Mediterraneo dalla R. N. "Washington" (1981-1982). *Archivio zoologico italiano*, 7: 81-113.
- Belloc G, 1948. *Inventory of the fishery resources of Greek waters. Appendix B*. UNRRA-FAO: 65-76.
- Bianchi CN, Morri C, 1983. Note sul benthos marino costiero dell' isola di Kos (Egeo Sud-Orientale). *Natura, Milano*, 74: 96-114.
- Bianchi CN, Morri C, 2000. Marine biodiversity of the Mediterranean Sea: Situation, Problems and Prospects for future research. *Marine pollution bulletin*, 40: 367-376.
- Borghi E, 1994. Osservazioni su un ritrovamento di *Echinocardium fenauxi* Péquignat, 1963 nel Mediterraneo orientale. *Museo regionale di scienze naturali bollettino (Torino)*, 14: 1-6.
- Burke RS, Mladenov PV, Lambert P, Parsley RL, 1988. Echinoderm Biology. In: *Proceedings of the sixth international Echinoderm conference, Victoria, 23-28 August 1987*. Balkema, Rotterdam: 337.
- Carus JV, 1885. *Prodromus faunae Mediterraneae. Echinodermata. I*. Stuttgart: 84-111.
- Caspers H, 1957. Black Sea and Sea of Azov. *The geological society of America memoir*, 67: 801-890.
- Caspers H, 1968. La macrofaune benthique du Bosphore et les problèmes de l'infiltration des éléments méditerranéens dans la mer Noire. *Rapport de la commission internationale pour la mer Méditerranée*, 19: 107-115.
- Chardy P, Laubier L, Reyss D, Sibuet M, 1973. Dragages profonds en mer Egée. *Rapport de la commission internationale pour la mer Méditerranée*, 22: 107-108.
- Cherbonnier G, 1956. Les échinodermes de Tunisie. *Bulletin de la station océanographique de Salammbô*, 53: 1-23.
- Cherbonnier G, 1957. Une Ophiure nouvelle de Méditerranée: *Amphiura apicula* nov. sp. *Vie et milieu*, 8: 200-204.
- Cherbonnier G, 1958. *Faune marine des Pyrénées-Orientales. 2. Echinodermes*. Université de Paris: 1-67.
- Cherbonnier G, 1968. Considérations sur l'Holothurie apo de *Leptosynapta makrankyra* (Ludwing). *Bulletin du muséum national d'histoire naturelle, Paris*, 39: 1214-1218.
- Cherbonnier G, 1969. Echinodermes récoltés par la "Thalassa" au large des côtes ouest de Bretagne et du golfe de Gascogne (3-12 Aout 1967). *Bulletin du muséum national d'histoire naturelle, Paris*, 41: 343-361.
- Cherbonnier G, 1972. *Neocnus incubans* nouveau genre et nouvelle espèce d'Holothurie dendrochirote incubatrice de Méditerranée. *Comptes rendus des séances de l'académie de science de Paris*, 275: 225-227.
- Cherbonnier G, 1986. Holothuries de Méditerranée et du nord de la mer Rouge. *Bulletin du muséum national d'histoire naturelle, Paris*, 8: 43-46.
- Cherbonnier G, Guille A, 1967. Complément à la faune des échinodermes de la mer de Banyuls. *Vie et milieu*, XVIII(2B): 317-330.
- Cherbonnier G, Guille A, 1968. Sur une nouvelle holothurie dendrochirote de Méditerranée: *Phyllophorus drachi* nov. sp. *Bulletin du muséum national d'histoire naturelle, Paris*, 40: 630-633.
- Çinar ME, Bilecenoglu M, Öztürk B, Can A, 2006. New records of alien species on the Levantine coast of Turkey. *Aquatic invasions*, 1: 84-90.
- Clark AH, 1931. A monograph of the existing crinoids. Vol. 1 (part 3). Superfamily Comasterida. *Bulletin of the United States national museum*, 82: 145-155.
- Clark AM, Rowe FWE, 1971. *Monograph of shallow-water Indo-West Pacific echinoderms*. British Museum (Natural History) London.
- Delépine R, Boudouresque CF, Frada-Orestano C, Noailles MC, Asensi A, 1987. Algues et autres végétaux marins. In: Fischer W, Bauchot ML, Schneider M, eds. *Fishes FAO d'identification des espèces pour les besoins de la pêche (Revision, 1), Méditerranée et mer noire, zone de pêche 37*. FAO, Rome: 1-136.
- Demetropoulos A, Hadjichristophorou M, 1976. Echinodermata of Cyprus. *Fisheries bulletin of the fisheries department, Cyprus*, 4: 1-83.
- Demir M, 1952/54. *The invertebrate benthos of the Bosphorus and of the littoral of the sea of Marmara, closer to the Bosphorus* (in Turkish). İstanbul üniversitesi fen fakültesi hidrobioloji araştırma enstitüsü. Yayınlarından, İstanbul.
- D'yakonov AM, 1954. *Ophiuroids of the USSR seas*. Israel program for scientific translations, Jerusalem.
- Ekman S, 1967. *Zoogeography of the sea*. Sidgwick and Jackson, London.
- El-Beshbeeshy M, 1995. Some echinoderms from the Egyptian Mediterranean waters. *Bulletin of the national institute of oceanography and fisheries (Egypt)*, 21: 367-396.
- Fiege D, Yulin Y, 1994. *Penilpidia ludwigi* (Marenzeller, 1893) (Holothurioidea: Elpidiidae) rediscovered from the Eastern Mediterranean. *Seventh deep sea biology symposium (poster session), Crete, September 1994*.
- Fiege D, Liao Y, 1996. *Penilpidia ludwigi* (Marenzeller, 1893) (Holothurioidea: Elpidiidae) rediscovered from the Eastern Mediterranean Sea. In: Uiblein F, Ott J, Stachowitsch M, eds. *Deep-sea and extreme shallow-water habitats: affinities and adaptations. Biosystematics and ecology series*, 11: 61-66.
- Forbes E, 1844. Report on the Mollusca and Radiata of the Aegean Sea, and on their distribution, considered as bearing on Geology. *British association on advance science*, 13: 130-193.

- Forbes E, 1845. On the Radiata of the Eastern Mediterranean, I. Ophiuridae. *Transactions of the Linnean society of London*, 19: 143-153.
- Fredj G, 1974. Stockage et exploitation des données en écologie marine. Considérations biogéographiques sur le peuplement benthique de la Méditerranée. *Mémoires de l'institut océanographique, Monaco*, 7: 1-88.
- Galán-Novella C, López-Ibor Alíño A, 1981. Nota preliminar sobre faunística y biogeografía de los equinoideos y ofiuroideos de la Península Ibérica y Baleares. *Boletín de la real sociedad española de historia natural. Sección biológica*, 79: 293-309.
- Gautier-Michaz M, 1958. Résultats scientifiques des campagnes de la "Calypso". III. 5. Echinodermes. *Annales de l'institut océanographique*, 34: 145-155.
- Geldiay R, Koçatas A, 1972. Note préliminaire sur les peuplements benthiques du Golfe d'Izmir. *Scientific monographs of the faculty of science, Ege university*, 12: 4-34.
- Gjiknuri L, 1981. Contribution à l'étude des échinodermes (Crinoidea et Asteroidea) du littoral Albanais. 2^e Congrès international sur la zoogéographie et l'écologie de la Grèce et des régions avoisinantes, Athens: 61-70.
- Guille A, Monteiro-Marques V, O'Connor B, 1983. Présence d'*Amphiura incana* (Ophiuroidea, Echinodermata) le long des côtes nord-est Atlantiques. *Cahiers de Biologie Marine*, XXIV: 383-390.
- Hansson HG, 1999. European Echinodermata check-list. A draft for the European register of marine species. Part of "Species 2000" compiled at TMBL (Tjärnö Marine Biological Laboratory). Web site: <http://erms.biol.soton.ac.uk/>.
- Issel R, 1928. Cenni sui risultati ottenuti dalla Missione Zoologica nel Dodecaneso (1926) per quanto concerne la fauna e la flora marine, con alcune osservazioni generali. In: *Ricerche Faunistiche nelle isole italiane dell'Egeo. Archivio zoologico italiano*, 12: 259-271.
- Jacquotte R, 1962. Etude des fonds de la mer Méditerranée. *Recueil des travaux de la station marine d'Endoume*, 26: 141-235.
- Kaspiris P, Tortonese E, 1982. Echinoderms from the western seas of Greece. *Thalassographica*, 2: 27-32.
- Kisseleva MI, 1961. Répartition qualitative et quantitative du benthos dans la région des Dardanelles en Mer Égée (in Russian). *Travaux de la station biologique de Sébastopol*, 14: 135-146.
- Kisseleva MI, 1963. La distribution qualitative et quantitative du benthos dans la mer Egée (in Russian). *Travaux de la station biologique de Sébastopol*, 16: 192-200.
- Kisseleva MI, 1983. Comparative characteristics of the benthos at some banks in the Aegean Sea. *Thalassographica*, 6: 107-118.
- Koehler R, 1924. *Les échinodermes des mers d'Europe I*. Encyclopédie scientifique, Doin, Paris.
- Koehler R, 1927. *Les échinodermes des mers d'Europe II*. Encyclopédie scientifique, Doin, Paris.
- Koukouras AS, Sinis AI, 1981. Benthic fauna of the North Aegean Sea II. Crinoidea and Holothuroidea (Echinodermata). *Vie milieu*, 31: 271-281.
- Koukouras A, Russo A, 1991. Midlittoral soft substratum macrofaunal assemblages in the North Aegean Sea. *Pubblicazioni della stazione zoologica di Napoli I: Marine ecology*, 12: 293-316.
- Koukouras A, Karachle P, 2005. The polyplacophoran (Eumollusca, Mollusca) fauna of the Aegean Sea with the description of a new species, and comparison with those of the neighbouring seas. *Journal of biological research*, 3: 23-28.
- Koukouras A, Voultsiadou E, Kitsos MS, Doulgeraki S, 2001. Macrofaunal diversity in the Aegean Sea, affinities with other Mediterranean regions and the Black Sea. *Bios (Macedonia, Greece)*, 6: 61-76.
- Laborel J, 1960. Contribution à l'étude directe des peuplements benthiques sciaphiles sur substrat rocheux en Méditerranée. *Recueil des travaux de la station marine d'Endoume*, 33: 117-173.
- Lacombe H, Tchernia P, 1960. Quelques traits généraux de l'hydrologie Méditerranéenne. *Cahiers océanographiques*, 12: 527-547.
- Lacombe H, Tchernia P, Benoist G, 1958. Contribution à l'étude hydrologique de la mer Egée en période d'été. *Bulletin d'information du COEC*, 8: 454-468.
- Longhurst A, 1998. *Ecological geography of the sea*. Academic Press, London.
- Lopez-Ibor Alíño A, Galan-Novella C, 1982. Primera cita de *Aslia lefevrei* (Barrois, 1882) (Echinodermata, Holothuroidea) en el Mediterraneo. *Cahiers de biologie marine*, 23: 269-273.
- Ludwig H, 1897. *Die Seesterne des Mittelmeeres. Fauna und Flora des Golfes von Neapel*. Monogr. XXIV, Berlin.
- Makkavieva E, 1963. Quelques peuplements des zones sableuses de la mer Égée (in Russian). *Travaux de la station biologique de Sébastopol*, 14: 211-214.
- Marenzeller E, 1893. Berichte der Commission für Erforschung des östlichen Mittelmeeres. Zoologische Ergebnisse. 1. Echinodermen gesammelt 1890, 1891 und 1892. *Denkschriften der kaiserlichen akademie der wissenschaften*, 60: 1-24.
- Marenzeller E, 1895. Berichte der Commission für Erforschung des östlichen Mittelmeeres. Zoologische Ergebnisse. 5. Echinodermen gesammelt 1893 und 1894. *Denkschriften der kaiserlichen akademie der wissenschaften*, 16: 1-21.
- Marion AF, 1898. Notes sur la faune des Dardanelles et du Bosphore. *Annales du musée d'histoire naturelle de Marseille*, 2: 163-182.
- Mayer B, 1937. Die Holothurien der Adria. *Thalassia*, 2: 1-58.

- Mortensen Th, 1927. *Handbook of the echinoderms of the British Isles*. Oxford University Press, Oxford.
- Mortensen Th, Steuer A, 1937. The fishery grounds near Alexandria. XIII-Echinodermata. *Hydrobiology and fisheries directorate, notes and memoirs*, 21: 1-37.
- Ostroumoff AD, 1896. Comptes rendus des dragages et du plancton de l'expédition de "Salianik" (in Russian). *Bulletin de l'académie impériale des sciences de St. Pétersbourg*, 5: 33-92.
- Ovchinnikov IM, 1966. Circulation in the surface and intermediate layer of the Mediterranean. *Oceanology*, 6: 48-59.
- Özaydin O, Katagan T, Ünsal S, 1995. The Echinoderms of the Turkish seas. *Israel journal of zoology*, 41: 57-68.
- Özsoy E, Hecht A, Ünlüta Ü, Brenner S, Sur HI, Bishop J, Latif MA, Rozentraub Z, Ogur T, 1993. A synthesis of the Levantine Basin circulation and hydrography, 1895-1990. *Deep sea research part II: topical studies in oceanography*, 40: 1075-1119.
- Panagiotopoulos P, 1916. Report on the hydrobiological station, Athens (in Greek). Ministry of national economy: 559-602.
- Pancucci MA, 1994. New records of Echinoderm species in Greek waters. *Bios (Macedonia, Greece)*, 2: 31-33.
- Pancucci-Papadopoulou MA, 1996. *Fauna Graeciae. VI. The Echinodermata of Greece*. Hellenic zoological society, Athens.
- Pancucci MA, Zenetos A, 1989. Infralittoral macrobenthos of the Patras gulf and Ionian Sea. II-Echinodermata. *Cahiers de biologie marine*, 30: 217-226.
- Pancucci MA, Zenetos A, 1990. On the echinoderm fauna of the Geras Gulf (Lesvos Island, Greece). *Acta adriatica*, 31: 293-300.
- Panning A, 1949. Versuch einer Neuordnung des familie Cucumariidae (Holothuroidea, Dendrochiota). *Zoologische Jahrbücher*, 78: 404-470.
- Papazacharias AS, 1991. Ecological study of the soft substratum macrofauna in the infralittoral zone of the gulf of Kavala (in Greek). Ph. D. Thesis, University of Thessaloniki.
- Pastore M, 1981. Osservazioni preliminari sull' infralitorale di substrato roccioso lungo la costa Salentina (Golfo di Taranto). *Thalassia salentina*, 11: 81-104.
- Pérès JM, 1967. The Mediterranean Benthos. *Oceanography and marine biology. An annual review*, 5: 449-533.
- Pérès JM, Picard J, 1956. Notes préliminaires sur les résultats de la campagne de recherches benthiques de la "Calypso" dans la Méditerranée nord-orientale. *Recueil des travaux de la station marine d'Endoume*, 18: 5-13.
- Pérès JM, Picard J, 1958. Recherches sur les peuplements benthiques de la Méditerranée nord-orientale. *Annales de l'institut océanographique*, 34: 213-281.
- Pérez-Ruzafa A, López-Ibor A, 1986. Presencia de *Holothuria (Vaneyothuria) lentiginosa* *lentiginosa* Marenzeller, 1983 (Echinodermata: Holothuroidea) en el mar de Alborán (Mediterráneo Occidental). *Boletín del instituto español de oceanografía*, 3: 105-109.
- Perivoliotis L, Lascaratos A, Nittis K, 1997. Seasonal variability of the general circulation and water mass formation in the Aegean Sea: A numerical study. *Proceedings of the 5th Hellenic symposium on oceanography and fisheries*, 1: 339-342.
- Pinardi N, Masetti E, 2000. Variability of the large scale general circulation of the Mediterranean Sea from observations and modelling: a review. *Paleogeography, paleoclimatology, paleoecology*, 158: 153-173.
- Piras A, 1972. *Odontaster mediterraneus* Mar. nelle acque della Sardegna (Echinodermata Asteroidea). *Doria*, 5: 1-3.
- Por FD, 1978. *Lessepsian Migration. The influx of Red Sea biota into the Mediterranean by way of the Suez Canal. Ecological studies* 23. Springer-Verlag, Berlin.
- Por FD, Dimentman Ch, 1989. The legacy of Tethys: an aquatic biogeography of the Levant. In: Dumont J, Wergel A, eds. *Monographiae biologicae*. Kluwer Academic Publishers, Dordrecht: 1-214+xi.
- Raulin V, 1870. Description physique de l'île de Crète. *Actes de la société Linnéenne de Bordeaux*, 22: 109-425, 23: 1-444, 23: 491-584, 24: 353-770.
- Rodríguez J, 1979. Echinoderms (except Holothuroidea) of the Southern Mediterranean coast of Spain. *Proceedings of the european colloquium on echinoderms, Brussels*: 127-131.
- Salvini-Plawen L, 1972. Die nordatlantische *Labidoplax buski* (Holothuroidea-Synaptidae) in der Adria. *Zoologischer anzeiger, zugleich organ der deutschen zoologischen gesellschaft*. Leipzig, 188S: 301-304.
- Salvini-Plawen L, 1977. Caudofoveata (Mollusca), Priapulida und apode Holothurien (*Labidoplax, Myriotrochus*) bei Banyuls und im Mittelmeer allgemein. *Vie milieu*, 27: 55-81.
- Schembri PJ, 1978. Recent echinoids (Echinodermata: Echinoidea) from the Maltese Islands and surrounding waters. *Animalia*, 5: 123-132.
- Sibuet M, 1974. Echinodermes de la mer d'Alboran. *Bulletin du muséum national d'histoire naturelle. 3e série, zoologie*, 231: 789-798.
- Spratt LTAB, Forbes E, 1842. *Travels in Lycia, Milyas, and the Cibyritis, in company with the late Rev. E. T. Daniell*. Bangor House, Shoe Lane, London.
- Steindachner F, 1891. Veröffentlichungen der Commission für Erforschung des östlichen Mittelmeeres. Vorläufiger Bericht über die Zoologischen Arbeiten in Sommer 1891. *Sitzungsberichte der mathematisch-naturwissenschaftlichen classe der kaiserlichen akademie der wissenschaften*, 100: 435-443.
- Theocharis A, Georgopoulos D, Lascaratos A, Nittis K, 1993. Water masses and circulation in the central region of the Eastern Mediterranean: Eastern Ionian,

- South Aegean and Northwest Levantine, 1986-1987. *Deep sea research part II: topical studies in oceanography*, 40: 1121-1142.
- Tortonese E, 1946. Echinoderms from the Eastern Mediterranean (Island of Rhodes). *The annali and magazine of natural history*, 13: 716-719.
- Tortonese E, 1947. Note inforno alla fauna e flora marine dell'isola di Rodi (Mara Egeo). *Bulletino di pesca, piscicolturae idrobiologia (note e memorie scientifiche)*, 2: 13-20.
- Tortonese E, 1952. Gli Echinodermi del Mar Ligure e delle zone vicine. *Atti dell'accademia ligure di scienze e lettere*, 8: 163-242.
- Tortonese E, 1956. Catalogo degli Echinodermi della collezione E. Tortonese. *Annali di museo civico di storia naturale di Genova*, LXVIII: 177-233.
- Tortonese E, 1957. On the Echinoderm fauna of Haifa bay. *The bulletin of the research council of Israel*, 6B: 189-192.
- Tortonese E, 1958. Il popolamento di Echinodermi nelle zone profonde del Mediterraneo. *Rapport commission international pour la mer méditerranée*, XIV: 485-491.
- Tortonese E, 1959. Campagnes de la "Calypso": Golfe de Gênes. 2. Echinodermes. *Annales de l'institut océanographique, Paris*, 37: 289-294.
- Tortonese E, 1961. Echinodermi di Taranto (Mar Jonio). *Thalassia jonica*, IV: 191-194.
- Tortonese E, 1965. *Fauna d'Italia. Echinodermata*. Edizioni Calderini, Bologna.
- Tortonese E, 1966. Echinoderms from the coast of Lebanon. *Miscellaneous papers on the natural science of the american university of Beirut*, 5: 1-5.
- Tortonese E, 1972. L' Echinofauna del piano batiale nel golfo di Genova. *Doriana*, 5: 1-7.
- Tortonese E, 1979. Review of present status of knowledge of the Mediterranean Echinoderms. *Proceedings of the european colloquium on echinoderms, Brussels*: 141-149.
- Tortonese E, 1985. Echinodermi di Torraldaliga (Civitareccchia-Roma). *Oebalia*, XI: 799-802.
- Tortonese E, Demir M, 1960. The Echinoderm fauna of the Sea of Marmara and the Bosphorus. *Hydrobiologia Istanbul*, 5: 1-16.
- The POEM Group (Robinson RA, Malanote-Rizzoli P, Hecht A, Michelato A, Roether W, Theocharis A, Ünlüta Ü, Pinardi N, Artegiani A, Bishop J, Brenner S, Christianidis S, Garic M, Georgopoulos D, Golnaraghi M, Hausmann M, Junghaus H-G, Laskaratos A, Latif A, Leslie G, Oguz T, Özsoy E, Papageorgiou E, Pachini E, Rosentroub Z, Sansone E, Scarazzato P, Schlitzer R, Spezie G-C, Zodiatis G, Athanassiadou L, Gerges M, Osman M), 1992. General circulation of the Eastern Mediterranean. *Earth-science reviews*, 32: 285-309.
- Ünsal S, 1985. Observations on the effects of pollutions of Izmir Bay on the Sea Stars (Echinodermata Asteroidae). *Rapport de la commission internationale pour la mer Méditerranée*, 29: 293-294.
- Vamvakas CN, 1971. Contribution to the study of soft bottom benthic biocoenoses in soft substrata around the Greek waters (W. Saronikos Gulf) (in Greek). Ph. D. thesis. University of Athens.
- Zaitsev Y, Öztürk B, 2001. Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas. *Turkish marine research foundation, Istanbul, publication No. 8*.
- Zavodnik D, 1960. Echinodermata der Insel Krk. *Acta adriatica*, 9: 1-20.
- Zavodnik D, 1967. Adriatic Echinoderms inhabiting the phytal. *Thalassia jugoslavica*, 3: 11-22.
- Zavodnik D, 1972. Peculiarities of geographical distribution of Adriatic echinoderms. *Thalassia jugoslavica*, 8: 321-330.
- Zavodnik D, 1976. Adriatic echinoderms inhabiting benthic organisms. *Thalassia jugoslavica*, 12: 375-380.
- Zavodnik D, 1977. Echinodermata of the Island Vir (Adriatic Sea). *Biosistemata*, 3: 69-78.
- Zavodnik D, 1979. Ergänzungen zur Echinodermenfauna des Adriatischen Meeres. *Zoologischer anzeiger, Jena*, 203: 122-128.
- Zavodnik D, 1980. Distribution of Echinodermata in the north Adriatic insular region. *Acta adriatica*, 21: 437-468.
- Zavodnik D, 1981. Report on Echinoderms from Malta. *Rapport commission international pour la mer méditerranée*, 27: 225-226.
- Zavodnik D, 1995. Additions to Adriatic Sea ophiuroid fauna and its diversity. *Natura croatica*, 4: 107-111.
- Zibrowius H, 1991. Ongoing modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Mésogée*, 51: 83-107.
- Zibrowius H, Bitar G, 2003. Invertébrés marins exotiques sur la côte du Liban. *Lebanese science journal*, 4: 67-74.