

- Acids, 219–220
- Acoustic cavitation, 20
- Aesthetic aromatherapy, 298
- African citrus essential oils, volatile components of, 50–57
- African citrus fruits
- Burundian, 67–68
 - compositional analysis of, 47–69
 - Ethiopian, 68–69
 - Kenyan, 48–64
 - Rwandan, 66–67
 - Ugandan, 64–66
- Albedo, 29, 30
- Albedo extracts, inhibitory activities of, 243
- Alchemy, 9
- Alcohols, 57–58, 218–219
- in American lemon essential oils, 71–72
 - in American lime essential oils, 84
 - in American mandarin essential oils, 79–80
 - in American orange essential oils, 75
 - in Anliu oranges, 92
 - in Hongjian oranges, 92
 - in Japanese citrus essential oils, 109, 113, 118–119, 122–123
 - in Kenyan mandarins, 60–61
 - in mandarin oil, 334
 - in Venezuelan orange oil, 74
 - in Vietnamese citrus fruits, 133, 140, 142
 - in Washington navel oranges, 93
- Aldehyde C-6, 350
- Aldehyde C-10, 350
- Aldehydes, 218. *See also* Aliphatic aldehydes
- in American lemon essential oils, 71
 - in American lime essential oils, 83–84
 - in American mandarin essential oils, 78–79
 - in American orange essential oils, 75
 - antimicrobial activities of, 287–288
 - in European citrus fruits, 146
 - in Japanese citrus essential oils, 109, 113, 118, 120–122
 - mandarin flavor and, 77
 - in Venezuelan orange oil, 74
 - in Vietnamese citrus fruits, 133, 140, 142
- Aliphatic aldehydes, 49–56, 59, 63–64, 68, 218
- in Chinese sweet oranges, 94
 - in mandarin oil, 331
 - in Uruguayan mandarins, 77
 - in Vietnamese citrus fruits, 142
 - in Washington navel oranges, 93
- Aliphatic hydrocarbons, 217. *See also* Aliphatics
- Aliphatics, in Vietnamese citrus essential oils, 134
- American citrus fruits
- compositional analysis of, 70–85
 - lemons, 70–74

- American citrus fruits (*cont'd*)
 limes, 83–85
 sweet oranges, 74–76
- American lime essential oils, volatile components of, 83–85
- American orange essential oils, volatile components in, 75–76, 78–82
- Analysis of variance (ANOVA), 307
- Analytical methods, 38–46
- Animal-derived aroma substances, 346
- Anliu oranges, 86, 92
- Antibacterial activity, of volatile terpenes, 286
- Antifungal activities
 of lemon oil, 282–283
 of mandarin oil, 283–284
- Anti-inflammatory activity, of essential oils, 300
- Antimicrobial activities
 of citrus oils, 264–277, 278–279
 of essential oils, 301
 of volatile compounds, 271–277
- Antimicrobial effects, assessment methods for, 277–278
- Antimicrobials, 230
- Antimicrobiology, 264–277
- Antioxidant activity, 231–244
- Antioxidant compounds, 231. *See also* Antioxidants
- Antioxidants, 4–5, 103, 230
 effectiveness of, 231–232
- Antioxidative compounds, natural, 5
- Antioxidative studies, alternative, 243–244
- AQUAWAX column, 42
- Aroma-active compounds
 identification of, 210
 in Pontianak citrus peel oil, 99, 100
- Aroma attributes, 354
- Aroma component analysis, 357
- Aroma components, in Korean Shiranui, 131
- Aroma compounds, 3–4, 37
 antimicrobial activities of, 279
- Aroma Extract Dilution Analysis (AEDA) techniques, 99, 203, 204–206, 209, 212, 213, 214
- Aromagrams, 205, 206
- Aroma materials
 for flavor/fragrance formulation, 345–351
 natural, 349
- Aroma models, of citrus essential oils, 216–217
- Aroma research, 3
- Aroma-resource plants, 38
- Aroma substances, synthetic, 349–350
- Aroma systems, analyzing the constituents of, 38–39
- Aromatherapeutic effect, 4
- Aromatherapy, 5, 297–337
 basic fields of, 298
 history of, 298–300
 importance of, 299–300
 symptoms and disorders affected by, 298
 therapeutic uses of essential oils in, 304–306
- Aromatic baths, therapeutic use of, 305
- Aromatic plants, functional properties of, 229–231
- Aromatic substances, industrial view of, 343, 344
- Ascorbic acid, 246
 effect on NDMA formation, 259
- Asian citrus fruits
 compositional analysis of, 86–145
 Indonesian, 96–105
 Japanese, 105–124
 Korean, 124–132
 Vietnamese, 132–145
- Atomic emission detector (AED), 43
- Atomic mass unit measurement, by GC-TOFMS, 177
- Aucuba japonica*, 252
- Authentic compounds, effect on NDMA formation, 249–251
- Authentic flavor compounds, antioxidative activities of, 239–240
- Authenticity control, 192
- Autonomic nerve activity, citrus essential oils and, 306–311
- Bacteria, foodborne, 277. *See also* Antibacterial activity; Antimicrobial entries
- Baked confectionery flavors, 355

- Banhakuyu, 115
Base note of flavors, 356
Base peaks, isotope ratio values for, 194
Beauty (aesthetic) aromatherapy, 298
Bergamot, 32
 European, 148–149
Bergamot aroma model, 216–217
Bergamot essential oil, ageing of, 323–330.
 See also Bergamot oils
Bergamot oil extractor, 32–33
Bergamot oils. *See also* Bergamot essential oil
 antimicrobial activities of, 266, 280–282
 compositional changes under differing storage conditions, 324–325
 comprehensive changes under storage conditions, 329–330
Bergapten, 311–314
Beverage flavors, 355
Beverages, yuzu flavor formulations for, 363
Biologically active compounds, 230
Bitter orange oils, antimicrobial activities of, 270
Bitter oranges, European, 150
Bitter/sour oranges, Japanese, 111
Body temperature, effect of olfactory stimulation on, 307–308
Brazilian lime essential oils, 83–85
Brazilian mandarins, 76–77
 volatile components in, 78–82
Brown adipose tissue (BAT), 306–311
Brown adipose tissue sympathetic nerve activity (BAT-SNA), 306–311
Brown juice extractor, 31
Buntan, antimicrobial activities of, 270
Burundian citrus essential oils, volatile components of, 51–57
Burundian mandarin peel oil, 68
Burundian mandarins, 67–68

Cai mandarin oils, 77
Calabrian Machine, The, 33
Californian lemon oils, 70, 71–73
Capillary column, 39
Carbon, photosynthetic fixation of, 168
Carbonyl compounds, 69
Carbonyl volatile compounds, 49
Carcinogenic factors, types of, 244–245
Carcinogens
 foodstuffs that inhibit, 246
 inhibitory formation of, 244–263
 strong, 252
Carvacrol, 285
Carvone, 141, 219, 327
(*E*)-Carvone, 59–60
(*Z*)-Carvone, 56–57, 59–60
Character-impact compounds, 201
Charm value, 206. *See also* Combined Hedonic Aroma Response Measurements analysis (CharmAnalysis)
Chemat, Farid, xi, 9, 11, 14, 20
Chemical microcapsulization methods, 365
Chemical structure, aroma characteristics and, 343
Chemistry, development of, 299
Chewing gum flavors, 355
Chinese citrus fruits
 compositional analysis of, 86–96
 cultivation area and production of, 87
 mandarins, 89–91, 94–96
 sweet oranges, 89–91
Chiral compounds, 167
Chiral flavor/fragrance components, 166
Chiral fruit flavor compounds, 166
Choi, Hyang-Sook, xi, 229, 231
Citral, 285, 350
Citrange oil, antimicrobial activities of, 270
Citron, European, 152
Citronellal, 102
 separating chiral isomers of, 46
Citronellol, antimicrobial activities of, 273
Citrus ampullacea, 111
Citrus aroma substances, synthetic, 350–351
Citrus aurantifolia, 63, 83–85, 107, 143–145, 180, 187, 213–214, 219, 249, 280–282
Citrus aurantium, 88, 111, 211, 216, 243, 358
Citrus bergamia, 32, 216–217
Citrus clementine, 76–83
Citrus components, pharmaceutical activities of, 5

- Citrus deliciosa*, 65, 76–83
Citrus depressa, 124
Citrus essential oil(s), ix, 3. *See also* Citrus oil(s); Essential oil(s)
antioxidative activities of, 240–241
aroma models of, 216–217
autonomic nerve activity and, 306–311
bergapten in, 312–313
chemical composition of, 19
Chinese, 86
chiral compounds of, 167
discrimination among, 192–196
effect on NDMA formation, 248–251
European, 146
flavor and odor/aroma impact compounds in, 208–217
information about, 6
inhibition of NDMA, 246
isotope fingerprint of, 186–190
isotope ratio of monoterpene hydrocarbons in, 171–174
Japanese, 314
oxidation of, 102–103
properties of, ix
relative antioxidative activities of, 238, 239
reports on, 37–38
scavenging effects of, 235, 236
storage conditions and analysis of, 315–316
tocopherol contents in, 242
types of, 5
volatile chemical compounds of, 48
volatile components of, 249–251
Citrus essential oil analysis, types of columns used in, 39–42
Citrus essential oil based on modified isotope ratio, isotope fingerprint of, 188–190
Citrus essential oil profiles, 304–305
Citrus extracts, inhibitory effects of, 238–239
Citrus flaviculpus, 217
Citrus flavor activity, functional groups related to, 217–220
Citrus flavor quality, evaluation of, 197
Citrus flavors. *See also* Flavors in food products, 97
manufacturing and formulation of, 358–361
Citrus fruit(s), ix
African, 47–69
alternative antioxidative studies of, 243–244
American, 70–85
Asian, 86–145
cold-pressed oil from, 30–33
European, 145–152
Indonesian, 96–105
industrial view of, 344–345
major, 37
popularity of, ix, 3
production of, 2–3
spread of, 37
structure of, 29
Citrus funadoko, 111
Citrus genus, 3
samples of, 247
Citrus glaberrima, 116
Citrus grandis, 64, 115–116, 141–143, 180, 187, 218
Citrus hanaju, 107, 186
Citrus hybrid oils, flavor and odor/aroma impact compounds in, 215–216
Citrus hystrix, 101–105
Citrus hystrix oil, odor profiles of, 105
Citrus ichangensis, 106–107, 214–215, 249, 367
Citrus indica, 94, 124
Citrus inflata, 107, 217
Citrus iyo, 111
Citrus junos, 31, 252, 306, 360, 367
Citrus kawachinesis, 116
Citrus kinokuni, 111
Citrus latifolia, 107
Citrus limon, 70–74, 107, 186, 214, 249, 282–283
Citrus limonia, 143–145
Citrus lumia, 243
Citrus materials, industrial view of, 343–378
Citrus maxima, 284
Citrus medioglobosa, 116
Citrus natsudaikai, 116, 186
Citrus nobilis, 65, 98–101, 213
Citrus note, 353
Citrus obovoidea, 116

- Citrus oil(s). *See also* Citrus essential oil(s)
antifungal activity of, 264, 278–279
antimicrobial activities of, 264–277, 278–279, 288–289
cold-pressed, 33
effect on fungal growth, 283
as flavor/fragrance aroma materials, 346–347
industrial view of, 345
processing method of, 349
weight percent of, 34
- Citrus oil monoterpene hydrocarbons, MS fragment isotope ratio of, 184
- Citrus otachibana*, 116
- Citrus ozu*, 116–124
- Citrus paradisi*, 58, 62, 63, 116, 211–212, 284, 348
- Citrus peel essential oils, 368
antioxidative activities of, 238–241
radical-scavenging activities of, 232–238
tocopherol analysis of, 241–243
- Citrus peel tissues, 20
- Citrus peel wastes, eco-conscious oil extraction from, 367–378
- Citrus recovery essence, 348–349
- Citrus reticulata*, 58, 76–83, 94–95, 105, 111–115, 130, 139–141, 180, 187, 212–213, 236, 243, 358, 367
- Citrus sinensis*, 74–76, 86, 93–94, 111, 116–124, 130, 133–139, 180, 187, 236, 243, 279–280, 281, 358
- Citrus sphaerocarpa*, 107, 216
- Citrus sudachi*, 107, 215
- Citrus sulcata*, 116, 216
- Citrus tachibana*, 59, 106
- Citrus taguma-sudachi*, 107
- Citrus tamurana*, 111, 216
- Citrus tangerina*, 58
- Citrus taxonomies, 1
- Citrus tengu*, 116
- Citrus tree valorization, 10
- Citrus ujukitsu*, 111
- Citrus unshiu*, 76–83, 95, 105, 111, 116–124, 130, 186, 212, 283–284
- Citrus varieties
propagation of, 1
spread of, 1
- Citrus yamabuki*, 116
- Citrus yuko*, 107, 124
- Clementine mandarin oils, 77–83
flavor and odor/aroma impact compounds in, 212–213
- Clementine mandarins, 76, 94
European, 151
- Cluster analysis, 192, 193, 195, 196
- Coefficient of variation (CV), 173–174
- Cold-pressed (CP) citrus essential oils, adulteration of, 165
- Cold-pressed essential oils, 148
- Cold-pressed lime peel oil (CPO), 176
- Cold-pressed oil(s) (CPO), 346, 359, 369, 374
from Chinese mandarins, 95
from Chinese oranges, 88–92
from citrus fruits, 30–33
extraction of, 28–34
quantitative determination of, 33–34
- Cold-pressed peel oil (CPO), 171–172
- Cold-pressing (CP), 11, 19
of Chinese sweet oranges, 34
technique for, 9
- Colombian lemon oils, 71–73, 74
- Colombian mandarin oils, 77
- Colombian mandarins, 76–77
volatile components in, 78–82
- Colombian orange essential oils, volatile composition of, 74–76
- Columns
in citrus essential oil analysis, 39–42
principal groups of, 40–41
- Combined Hedonic Aroma Response Measurements analysis (CharmAnalysis), 203, 204, 206, 210
- Component identification, 44
- Components, isotope content of, 170
- Compositional analysis, 37–152
of African citrus fruits, 47–69
of American citrus fruits, 70–85
analytical methods in, 38–46
of Asian citrus fruits, 86–145
of European citrus fruits, 145–152
- Compound stereochemistry, antimicrobial activities and, 288
- Compresses, therapeutic use of, 306
- Comune mandarins, volatile components in, 78–82
- Constituents, isotope values of, 168

- Conventional solvent extraction, 21
 Conventional steam-distilled oil (CSDO), 370–374
 Correspondence analysis, 210
 Cost of extraction, for MHG method, 20
 Coumarins, 149, 150
 Cross-matching test, 210, 215
 Cyclodextrins (CDs), 166–167
p-Cymene, 327
 interconversion of, 329–330
- Daidai, 211. *See also* Kiyookadaidai
 Daidai aroma model, 216
 Dairy product flavors, 355
 DB-WAX, 41, 42
 Deans switch, 45
trans-4-Decenal, 350
 Dekopon, 215. *See also* Hallabong
 Dessert flavors, 355
 Detection frequency technique, 203, 207
 Detector response, 43
 Detector types, 42–43
 8,9-Didehydronootkatone, 350
 Diffuser, therapeutic use of, 306
 Dihydromyrcenol, 351
 Dihydromyrcenyl acetate, 351
 Dilution analysis, 203, 204
 Dilution techniques, 204
 Dilution-to-threshold method, 203
Diospyros kaki, 252
 Diphenyl groups, 42
 Direct steam distillation, 12
 Distillation column, 38
 Distillation methods, 11–13
 Distillation residue, treatment of, 374–377
 Distillation technique, 9
 Distillation technologies, 13
 Distilled oil, 359
trans-2-Dodecenal, 350
 DPPH method, 232–236
 DPPH radical, 235, 237
 Dressing flavors, 355
 Dry distillation, 11, 12
- Eco-conscious oil extraction, from citrus peel wastes, 367–378
 Egg-box model, 376
 Egyptian medicine, aromatic plants used in, 298–299
- Electrical theory, 20
 Electrolytic conductivity detector (ELCD), 43
 Electron capture detector (ECD), 42
 Emulsified flavors, 354, 364
 Enantiomeric analysis, 165, 166–168
 Enantiomeric component distribution, GC analysis of, 45
 Enantiomeric distribution, 167–168
 Enantiomeric excess, 167
 Enantiomeric purity, 167
 Enantioselective capillary gas chromatography (enantio-cGC), 166
 Environmental burden, 15
 Environmental impact of MHG method, 20
 of soxhlet versus ultrasound-accelerated extraction, 23–24
 Epoxides, 58, 220
 “The era of fragrance,” 3
 Essential oil(s), 5, 9, 106. *See also* Citrus essential oil(s)
 ageing of, 315–323
 antimicrobial activities of, 279, 288
 in aromatherapy, 297
 authenticity control of, 165–166
 bergamot, 32, 148–149
 biologically active compounds in, 230–231
 bitter orange, 150
 “character-impact odorants” in, 201
 citron, 152
 composition of, 37
 compositional changes in, 337
 defined, 11
 as functional ingredients, 38
 mandarin, 150–151
 mechanism of action of, 279
 occurrence of, 29–30
 organoleptic properties of, 19–20
 pharmacological effects of, 300–304
 processing technologies of, 349
 separations of, 38–39
 sweet orange, 149–150
 therapeutic uses of, 304–306
 of Vietnamese citrus fruits, 133
 Essential oil compounds, chemotaxonomic value of, 62

- Essential oil constituents, pharmacological properties of, 301–303
- Essential oil industry, gas chromatograph impact on, 39
- Essential oil origins, authenticity control of, 179–196
- Essential oil safety, 311–314
- Essential oil spray, therapeutic use of, 306
- Esters, 58, 219
- in American lemon essential oils, 72–73
 - in American lime essential oils, 84
 - in American mandarin essential oils, 81
 - in American orange essential oils, 76
 - in Japanese citrus essential oils, 110, 114, 119, 122–123
 - in Kenyan mandarins, 61
 - odor-active, 209
 - in Vietnamese citrus fruits, 140–141, 142–143
- Ethiopian citrus essential oils, volatile components of, 51–57
- Ethiopian citrus fruits, 58–69
- Ethyl anthranilate, 351
- Eureka lemon oil,
- European citrus fruits, 145–152
- bergamot, 148–149
 - bitter orange, 150
 - citron, 152
 - grapefruit, 151–152
 - lemon, 146–148
 - sweet orange, 149–150
- European Union regulations, 6
- Extract, 346
- Extraction techniques, drawbacks of, 12
- Extraction technologies, 13
- in European countries, 148
- “Fantastico” bergamot variety, 33
- FD factors, 99–101
- Fiber, selectivity of, 26
- Fiber assembly, 25
- Flame ionization detector (FID), 42
- Flame ionization detector scheme, 43, 44
- Flame photometric detector (FPD), 43
- Flavado, 29, 30
- bergamot, 149
 - lemon, 148
 - mechanical pressing of, 369
- Flavado extract, inhibitory activities of, 243
- Flavor, analyzing volatile compounds in, 171. *See also* Flavors
- Flavor creation, 355–356
- Flavor design, 354
- Flavor dilution factors (FD-factors), 206, 213, 216, 217
- Flavor evaluation, 356–357
- Flavor formulation, 353–367
- aroma component analysis for, 357
 - aroma materials for, 345–351
- Flavor materials, 356
- Flavor recipes, 353
- Flavor research, 3–4, 166
- flowchart of, 4
- Flavors. *See also* Citrus flavors
- application of, 354–355
 - authenticity control of, 165–166
 - classifying by form, 361–365
 - industrial view of, 344
 - verifying the origin of, 169
- Florentine flask, 12, 18
- Floridian lime essential oils, 83–85
- FMC in-line juice extractor, 30–31
- Food
- analyzing volatile compounds in, 171
 - antioxidative activity of, 231–244
- Food antioxidants, 230
- Foodborne bacteria, 277
- Food constituents, isotope analysis of, 6
- Food flavors, microcapsulization of, 365
- Food safety/reliability concerns, 6
- Formulation. *See also* Fragrance
- formulation
 - of essential oils for processed foods, 345–367
 - process of, 344
- Fortunella*, 243
- Fortunella japonica*, 124, 216
- Fragrance formulation, 352
- aroma materials for, 345–351
- Fragrance research, 166
- Fragrances
- industrial view of, 344
 - verifying the origin of, 169
- Frankincense, 299
- Free-fatty acid (FFA) levels, 306, 307, 309, 311
- Free radicals, 231, 238
- Frozen food flavors, 355

- Fruiting position, effects on isotope ratio values, 175
- Fruit maturity, effects on isotope ratio, 175
- Fruit types, major, 2–3
- Functional properties, 229–289
- Fungi, citrus oil effect on, 283. *See also* Antifungal activities
- Fungitoxicity, of orange oil, 279–280
- Furanocoumarin, 314
- Gas chromatograms, 27, 205
- Gas chromatographic separation, 211
- Gas chromatography (GC), 24, 33, 34, 38–39. *See also* Enantioselective capillary gas chromatography (enantio-cGC); GC detector types; High-resolution gas chromatography (HRGC); Multidimensional GC analysis; Two-dimensional GC (GC H GC)
- Gas chromatography and flame ionization detection (GC-FID), 42–44
- Gas chromatography–isotope ratio mass spectrometry (GC-IRMS), 169–170. *See also* Isotope ratio mass spectrometry (IRMS)
- Gas chromatography–mass spectrometry (GC-MS), 44–45, 166, 172. *See also* Isotope ratio analysis by GC-MS
- isotope ratio accuracy by, 178–179
- isotope ratio of monoterpene hydrocarbons by, 171–174, 180–182
- isotope ratio of oxygenated compounds by, 178
- Gas chromatography–olfactometry (GC-O), 3–4, 99, 201–203, 209. *See also* GC-O entries
- performance of, 202
- prospects for, 220–221
- Gas chromatography–time-of-flight mass spectrometry (GC-TOFMS), isotope analysis by, 176–177
- Gas–liquid chromatograph, 38
- Gattefossé, René Maurice, 297, 299
- GC detector types, 42–43. *See also* Gas chromatography entries
- GC-O analysis, factors affecting, 202–203. *See also* Gas chromatography–olfactometry (GC-O)
- GC-O techniques, 203–204
- Germacrene D, 49
- Grapefruit(s). *See also* Redblush grapefruit
- effect on NDMA formation, 250
- European, 151–152
- Japanese, 116
- Kenyan, 61–64
- Grapefruit flavors, manufacturing and formulation of, 359–360
- Grapefruit juice, flavor and odor/aroma impact compounds in, 211–212
- Grapefruit oil(s)
- antimicrobial activities of, 269–270, 284
- therapeutic use of, 311
- Grapefruit peel oil, flavor and odor/aroma impact compounds in, 212
- Grapefruit production, 348
- Green extraction techniques, 15
- Green techniques, 10, 13
- Hallabong, 130–131
- Hallabong oil, flavor and odor/aroma impact compounds in, 215
- Hanayu, 187
- Headspace gas extraction, 25
- Headspace-SPME analysis method, 131. *See also* Solid-phase microextraction (SPME)
- Herbal medicine, 299
- High-performance liquid chromatography (HPLC), 26, 46, 246
- High-resolution gas chromatography (HRGC), 165
- Holder assembly, 25
- Holistic aromatherapy, 298
- Home-care aromatherapy, 298
- Hongjian oranges, 86, 88–92
- Hotspot theory, 20
- HPLC-DPPH screening method, 232
- Hybrid citrus fruits, Japanese, 116–124
- Hydrocarbons
- in Burundian mandarins, 67–68
- in Chinese mandarins, 96
- in Ethiopian citrus fruits, 69
- in Kenyan mandarins, 59
- in Sihui oranges, 93
- in Vietnamese citrus fruits, 133, 139
- in Washington navel oranges, 93

- Hydrodiffusion, 16–18
- Hydrodistillation, 12
- Hydrodistilled (HD) essential oil, 20
- Hyuganatsu, 111
- Hyuganatsu aroma model, 216
- Hyuganatsu oil, 218, 219
flavor and odor/aroma impact
compounds in, 216
- Ichang-papeda, 106–107
- Ichang-papeda oil, flavor and odor/aroma
impact compounds in, 214–215
- Incremental dilution techniques, 204
- Indirect steam distillation, 12
- Indonesian citrus families, odor-active
compounds of, 105
- Indonesian citrus fruits, 96–105
- Indonesian lime, 101–105
- Infrared detector (IRD), 43
- Inhalation, therapeutic use of, 304
- In-line juice extractor, 30–31
- Inside-needle technique, 28
- International Fragrance Association
(IFRA), 314
- Ion-trap mass spectrometer chemical
sensor, 211
- Ir discrimination, 192. *See also* Isotope
ratio entries
- Ir values, 173–174, 179, 190–191. *See also*
Isotope ratio values
- Isotope analysis, by gas chromatography–
time-of-flight mass spectrometry,
176–177
- Isotope dilution assays, 209
- “Isotope effect,” 6, 168
- Isotope fingerprint, of citrus essential oil,
186–190
- Isotope ratio(s). *See also* Ir entries
accuracy of, 178–179
determination of, 172
effects of fruit maturity on, 175
effects of variable factors on, 175–176
influence of storage on, 175–176
of linalool, 178, 190–191
of monoterpene hydrocarbons,
180–182
multivariate analysis of, 192–195
of oxygenated compounds, 178
- Isotope ratio analysis, 165, 168–179
- Isotope ratio analysis by GC-MS. *See also*
Gas chromatography–mass
spectrometry (GC-MS)
accuracy of, 173–174
repeatability and precision of, 172–173
- Isotope ratio mass spectrometry (IRMS),
166, 169–170
- Isotope ratio values, 173–174. *See also*
Ir values
effects on, 175–176, 191
- Isotopic Afingerprint, @ 169
- Italian citrus essential oils, volatile
components of, 147
- Italy, citrus production in, 145
- Iwabuchi, Hisakatsu, xi, 343
- Japanese Agricultural Standard (JAS), 6
- Japanese citrus essential oils, 314. *See also*
Japanese yuzu essential oils
MS fragment isotope ratio analysis
of monoterpene hydrocarbons in,
183
volatile components of, 108–110,
112–114, 117–123
- Japanese citrus fruits, 105–124
bitter/sour orange, 111
hybrids and unidentified species,
116–124
ichang-papeda, 106–107
kumquat, 124
lemon, 107
lime, 107
mandarins, 111–115
pummelo, 115–116
sweet orange, 111
tachibana, 106
- Japanese Washington navel oil, 93
- Japanese yuzu essential oils, isotope ratio
analysis of monoterpene
hydrocarbons in, 180–182
- Jeruk keprok, 96–97, 98
- Jeruk pontianak, 98–101
- Jeruk purut, 97, 98, 101–105
oxidized, 103
- Jeruk purut flavor powder, 103–105
- Jeruk purut leaves, 101–102
- Jeruk purut oil, 102–103
- Jeruk siam, 98
- Juice-extraction methods, yuzu, 125

- Kabosu, 107
Kabosu aroma model, 216
Kaffir lime, 101–105
Kaffir lime oil, oxidized, 102, 104
Kara mandarins, 58–61
 volatile components of, 56
Kashiwagi, Takehiro, xi, 367
Kenyan citrus essential oils, volatile components of, 50–57
Kenyan citrus fruits
 compositional analysis of, 48–64
 grapefruit, 61–64
 mandarins, 58–61
 pummelo, 63–64
Kenyan sweet orange oils
 monoterpene hydrocarbons in, 48–49
 oxygenated compounds in, 49–58
 sesquiterpene hydrocarbons in, 49
 volatile components of, 58
Ketones, 219–220
 in American lemon essential oils, 72
 in American lime essential oils, 84
 in American mandarin essential oils, 80–81
 in American orange essential oils, 75
 in Japanese citrus essential oils, 110, 114, 119
Key lime oil, volatile components of, 83–85
Kiyomi, Korean, 132
Kiyookadaidai, 116–124. *See also* Daidai
Korean citrus essential oils, 125
 volatile flavor components of, 126–127
 volatile profile characteristics of, 128–129
Korean citrus fruits, 124–132
 Kiyomi, 132
 kumquat, 131
 satsuma mandarin, 130
 Setoka, 132
 Shiranui, 130–131
 yuzu, 125–130
Kumagai, Chizu, xi, 297, 298
Kumquat
 Japanese, 124
 Korean, 131
Kumquat oil, flavor and odor/aroma impact compounds in, 216
Laboratory cold-pressed oil (CPO)
 extraction, 30
Lan-Phi, Nguyen Thi, xi, 105, 132, 201
Latin American citrus fruits, 70
Lavandula angustifolia, 307–308
Lemon essential oil(s)
 ageing of, 316–323
 compositional changes under differing storage conditions, 316–322
 volatile components in, 71–73
Lemon flavors, manufacturing and formulation of, 358–359
Lemon oil(s). *See also* Californian lemon oils; Colombian lemon oils; Eureka lemon oil; Uruguayan lemon oils; Venezuelan lemon oils
 antimicrobial activities of, 267–268, 282–283
 antioxidative activities of, 240–241
 flavor and odor/aroma impact compounds in, 214
 SPME assay for, 26–28
Lemon production, 347–348
Lemons. *See also* Lisbon lemon
 American, 70–74
 European, 146–148
 Japanese, 107
Lime essential oils, isotope fingerprint of, 188
Lime extract, antimicrobial activities of, 282
Lime flavors, manufacturing and formulation of, 359
Lime oil(s). *See also* American lime essential oils; Brazilian lime essential oils; Cold-pressed lime peel oil (CPO); Floridian lime essential oils; Key lime oil; Mexican lime essential oils; Persian lime oil
 antimicrobial activities of, 266–267
 flavor and odor/aroma impact compounds in, 213–214
Limes. *See also* Indonesian lime; Kaffir lime; Persa lime; Tahiti lime
 American, 83–85
 Japanese, 107
 Vietnamese, 143–145

- Limonene, 10, 48, 59, 62, 64–65, 74, 106, 107, 144, 146, 217
antimicrobial activities of, 271
as a dilution solvent, 208
in Japanese citrus essential oils, 124
in mandarin oil, 331
- Linalool, 95, 149, 350
antimicrobial activities of, 272
in bergamot oil, 326, 329–330
as a dilution solvent, 208
effects of, 311
isotope ratios of, 178, 179, 190–191
- Linalyl acetate, 149, 326, 328, 350
in bergamot oil, 329–330
- Linoleic acid peroxidation, antioxidative activities against, 238–241
- Liquid chromatography (LC), 28. *See also* High-performance liquid chromatography (HPLC)
- Liquid chromatography–mass spectrometry (LC-MS), 253
- Liquid flavor preparation, extraction methods in, 103
- Liquid fragrance formulations, 353
- Lisbon cold-pressed oil, 319
- Lisbon lemon, 107, 241
- Lisbon lemon oil, 241
- Mandarin essential oil(s)
ageing of, 330–337
compositional changes under differing storage conditions, 331–336
composition of, 330–331
comprehensive changes under storage conditions, 336–337
isotope fingerprint of, 189
- Mandarin hybrids, volatile components of, 51–57
- Mandarin oils(s). *See also* Cai mandarin oils
antimicrobial activities of, 268–269, 283–284
flavor and odor/aroma impact compounds in, 212–213
- Mandarins. *See also* Brazilian mandarins; Clementine mandarins; Colombian mandarins; Comune mandarins; Kara mandarins; Minneola mandarins; Miyagawa-wase mandarins; Miyamoto mandarins; Nova mandarins; Nules mandarins; Okitsu-wase mandarins; Oranges; Ponkan mandarins; Sabine mandarins; Satsuma mandarin(s); Shatang mandarins; Unshiu mandarins; Uruguayan mandarins; Venezuelan mandarins
American, 76–83
Burundian, 67–68
Chinese, 86, 94–96
European, 150–151
Japanese, 111–115
Kenyan, 58–61
Korean, 130
Mediterranean, 94
Ugandan, 64–66
Vietnamese, 139–141
volatile components of, 51–57, 89–91
- Massage oils, therapeutic use of, 305
- Mass selective detector (MSD), 43
- Mass spectrometry (MS). *See* MS
fragment entries
- Mediterranean mandarins, 94
- 1-*p*-Menthen-8-thiol, 351
- Methyl dihydrojasmonate, 351
- Methyl *N*-methlanthranilate, 77
- Methyltrisulfide, 220
- Mexican lime essential oils, 83–85
- Microcapsulization, of food flavors, 365–366
- Microextraction, solid-phase, 24–28
- Microextraction in a packed syringe (MEPS) technique, 28
- Microorganisms, antimicrobial activities of citrus oils on, 264, 277
- Microwave-assisted extraction (MAE), 18.
See also Microwave extraction
- Microwave Assisted Process (MAP), 14
- Microwave-assisted solvent extraction (MASE), 14
- Microwave dielectric heating, 14
- Microwave extraction, 14–20
- Microwave hydrodiffusion, 18
- Microwave hydrodiffusion and gravity (MHG), 15–20
- Middle note of flavors, 356

- Minh-Tu, Nguyen Thi, xi, 38
- Minimum detectable level (MDL), 43
- Minneola mandarins, volatile components of, 50–56
- Minneola tangelo, 58–61
- Miyagawa-wase mandarins, 212
- Miyamoto mandarins, 212
- Monoterpene alcohols, 285–287
- Monoterpene aldehydes, 56
- Monoterpene hydrocarbons, 28, 67–68, 69, 70. *See also* Monoterpenes
 in American lemon essential oils, 71
 in American lime essential oils, 83
 in American mandarin essential oils, 78
 in Anliu oranges, 92
 in Chinese sweet oranges, 96
 in citrus essential oils, 179
 in clementine oils, 77
 determination of, 46
 in European citrus fruits, 151
 in Hongjian oranges, 88
 isotope ratio by GC-MS, 171–174
 isotope ratio in citrus essential oils, 171–174
 isotope ratios of, 175–176, 177, 180–182
 in Japanese citrus essential oils, 124
 in Kenyan sweet orange oils, 48–49
 in mandarin oils, 59, 336
- MS fragment isotope ratio analysis of, 182–186, 194–195
- multivariate analysis of, 192–195
- in pummelo, 63
- in Ugandan mandarins, 64–65
- in Ugandan sweet oranges, 66
- in Venezuelan orange oil, 74
- in Vietnamese citrus fruits, 142, 144
- in Washington navel oranges, 93
- in yuzu oil, 371–374
- Monoterpene hydrocarbon standards, isotope ratio values of, 173
- Monoterpenes, 19, 217
 in American orange essential oils, 75
 composition analysis of, 41–42
 difference in the isotope ratio values among, 183–186
 in Japanese citrus essential oils, 108, 112, 117, 120–121
 in Vietnamese citrus essential oils, 134
 in Vietnamese citrus fruits, 139
- MS fragment isotope ratio analysis, of
 monoterpene hydrocarbons, 182–186
- MS fragment ratio, 45
- Multidimensional GC analysis, 203. *See also* Gas chromatography (GC)
- Multidimensional HPLC/GC, 46. *See also* High-performance liquid chromatography (HPLC)
- Multivariate analysis, 45
 of isotope ratio, 192–195
- Myrcene, 48–49, 63, 177, 217, 331
 effect on NDMA formation, 249
 inhibition of NDMA formation, 249
- Myrtenol, 327
- Naoshichi, 107
- Nasal impact frequency (NIF), 207
- Natural antioxidants, 5
- Natural aroma materials, processing technologies of, 349
- NDMA assay, 246–248, 253. *See also* N-Nitrosodimethylamine (NDMA)
- NDMA formation
 effect of ascorbic acid, nitrite, nitrate, and saliva on, 259–260
 effect of terpene hydrocarbons on, 260–263
 effects of vegetable extracts on, 253–257
 effects of vegetable extracts with yuzu oil and saliva on, 255–256, 257–258
- NDMA formation chromatograms, 262
- Neroli essential oil, 47
- Nitrate(s), 252
 effect on NDMA formation, 260
- Nitrate content, determination of, 253
- Nitrite(s), 252
 effect on NDMA formation, 259
- Nitrite content, determination of, 253
- Nitrogen isotope fractionation, 169
- Nitrogen phosphorus detector (NPD), 42
- Nitrosoamines, 246, 252
- Nitroso compounds, inhibition of, 252
- N-Nitroso compounds, 252
 carcinogenesis and, 244
- N-Nitrosodialkylamine, metabolic activation of, 245
- N-Nitrosodimethylamine (NDMA), 246.
See also NDMA entries
 inhibitory effects on, 248–253

- Njoroge, Simon Muhoho, xi, 47
- Nonvolatile compounds, under storage conditions, 322–323
- Nootkatone, 60, 62, 64, 69, 151, 152, 211, 350
detection of, 94
in Vietnamese citrus fruits, 143
- Nova mandarin oils, 77
- Nova mandarins, 76
volatile components in, 78–82
- Nuclear magnetic resonance (NMR), 170–171
- Nules mandarins, volatile components in, 78–82
- Nutraceuticals, 229
- Octyl acetate, 219
- Odor-active compounds, 221
- Odor-active volatiles, 221
- Odor-activity value (OAV), 101
- Odor/aroma key compounds, 201
- Odor cluster, 220
- Odor detection threshold, 204
- Odor peaks, 206–207
- Odor spectrum values (OSV), 212
- Oil extraction. *See also* Citrus essential oil(s); Essential oil(s)
eco-conscious, 367–378
techniques for, 9–36
- Oil-soluble liquid flavors, 364
- Oily flavor preparations, 355
- Okitsu-wase mandarins, 95
- Oleoresin, 346
- Olfactory stimulation
with grapefruit scent, 311
with yuzu scent, 307–311
- Omission test, 207–208
- On-column injection, 39
- Open tubular capillary columns, 39–40
- Oral-care product flavors, 355
- Orange essential oils, isotope fingerprint of, 188. *See also* American orange essential oils; Colombian orange essential oils
- Orange flavors, manufacturing and formulation of, 358
- Orange juices, antioxidant efficiency of, 243
- Orange oil(s). *See also* Kenyan sweet orange oils; Pera orange oil; Pontianak orange oil; Sour orange oil; Venezuelan orange oil
antimicrobial activities of, 265–266, 278, 279–280
inhibition effect on spore germination, 280
- Oranges. *See also* Anliu oranges; Bitter oranges; Hongjian oranges; Mandarins; Rwandan sweet oranges; Salustiana oranges; Sihui oranges; Sweet oranges; Ugandan sweet oranges; Valencia oranges; Washington navel oranges
production of, 347
volatile components of, 51–57
- Organoleptic effects, 4
- Orlando tangelo, 68–69
- OSME time-intensity technique, 203, 206–207
- Oxidation reactions, 230
- Oxides, 58, 220
in American lemon essential oils, 72
in American lime essential oils, 84
in American mandarin essential oils, 80–81
in Japanese citrus essential oils, 110, 114, 119, 122–123
- Oxygenated compounds, 15, 19
in American mandarin essential oils, 77
in Ethiopian citrus fruits, 69
isotope ratio of, 178
in Kenyan mandarins, 59–61
in Kenyan sweet orange oils, 49–58
in Ugandan mandarins, 65–66
- Oxygenated monoterpenes, 285
- Oyu essential oil, 249
- Packed columns, 39–40
- Peel oil(s)
lime, 144
pummelo, 143
- Pelatrice system, 33
- Pera orange oil, Charm values of, 210
- Perfumery, 9
- Perillene, 63, 64
- Persa lime, 143

- Persian lime oil, volatile components of, 83–85
- Pharmaceutical activities, of citrus components, 5
- Pharmacological effects, of essential oils, 300–304
- Pharmacologically active compounds, 229
- Pharmacological properties, of essential oil constituents, 301–303
- Phase separation method, 366
- Phenolic compounds, 232
- Phenolic content, influence on antioxidant effectiveness, 243
- Photoionization detector (PID), 43
- Photosynthesis, 168–169
- Physical microcapsulization methods, 366
- Physicochemical microcapsulization methods, 365–366
- Phytochemicals, 229
 - antioxidant activity of, 231
- Piper hispidinervum*, 66
- Plant-derived aroma substances, 345–346
- Plant essential oils, antimicrobial activity of, 278
- Plants, functional properties of, 229
- Plasma FFA levels, effect of olfactory stimulation of, 308–309
- Poiana, Marco, xi, 145
- Polyethylene glycol (PEG), 41
- Polygonum longisetum*, 252
- Poncirus*, 243
- Ponkan essential oil, 96
- Ponkan mandarins, 94, 95, 96
- Pontianak orange oil, flavor and odor/aroma impact compounds in, 213
- Posterior-intensity technique, 203, 207
- Postharvest pathogens, 264
- Potent odorants, 201
- Powdered flavors, 354, 364–365
- Power ultrasound, 20
- Pummelo, 61
 - Japanese, 115–116
 - Kenyan, 63–64
 - Vietnamese, 141–143
 - volatile components of, 51–57
- Pummelo essential oils, isotope fingerprint of, 189
- Pummelo oil(s), antimicrobial activities of, 270, 284
- Quinta essentia, 9
- Radical-scavenging activities/effects, 233
 - Citrus* genus samples for, 234
 - of citrus peel essential oils, 232–238
- Recovery essence, 346, 359
- Redblush grapefruit
 - Kenyan, 61, 62
 - volatile components of, 51–57
- Regulations, food safety, 6
- Relative antioxidative activities, 238, 239
- Relative flavor activity (RFA), 101, 213, 216
- Relative lipid peroxidation rate (RLPR), 238, 239–241, 242
- Rwandan citrus essential oils, 51–57
- Rwandan sweet oranges, 66–67
- S-1 storage condition
 - for bergamot oil, 326–327
 - for lemon oil, 320–321
 - for mandarin essential oil, 331–334
- S-2 storage condition
 - for bergamot oil, 328
 - for lemon oil, 321–322
 - for mandarin essential oil, 334–335
- S-3 storage condition
 - for bergamot oil, 328
 - for lemon oil, 322
 - for mandarin essential oil, 336
- S-4 storage condition
 - for bergamot oil, 328–329
 - for lemon oil, 322
 - for mandarin essential oil, 336
- Sabine mandarins, 58–61
 - volatile components of, 50–56
- Saliva, NDMA formation in the presence of, 258, 260
- Salustiana oranges, volatile components of, 50–56
- Salustian navel fruits, 48–58
- Sanbokan, 116
- Sanbokan oil, flavor and odor/aroma impact compounds in, 216
- Satake, Atsushi, xi, 168
- Satsuma mandarin(s), 76, 77, 94, 95, 105, 111, 130
 - effect on NDMA formation, 250
 - volatile components in, 78–82

- Satsuma mandarin oil(s)
 flavor and odor/aroma impact
 compounds in, 212
 Korean, 140
- Sawamura, Masayoshi, ix, xii, 1, 24, 28, 37,
 47, 86, 105, 315, 367
- Scanning electron microscopy, 21
- Seafood product flavors, 355
- Selected ion monitoring (SIM) acquisition
 mode, 171, 172, 175, 180
- Selective detection, limited, 44–45
- Sensory analysis, 24, 202, 203
- Separation techniques, 13
- Sesquiterpene alcohols, 58
- Sesquiterpene hydrocarbons. *See also*
 Sesquiterpenes
 in American lemon essential oils, 73
 in American lime essential oils, 85
 in American mandarin essential oils,
 81–82
 in Kenyan mandarins, 59
 in Kenyan sweet orange oils, 49
 in Ugandan mandarins, 65
 in Ugandan sweet oranges, 66
 in Vietnamese citrus fruits, 142
- Sesquiterpenes, 19, 217–218
 in American orange essential oils, 75
 in Japanese citrus essential oils, 108–109,
 112–113, 117–118, 120–121
 in Vietnamese citrus essential oils,
 134–138
 in Vietnamese citrus fruits, 139
 in yuzu oil, 374
- Setoka, Korean, 132
- Shaddock, Japanese, 115–116
- Shatang mandarins, 96
- Shimada, Masaki, xii, 345
- Shiranui, 116
 Korean, 130–131
- Signal-to-noise ratio, 45
- Sihui oranges, 86, 93
- α -Sinensal, 56
- β -Sinensal, 56
- Sinensals, 93
- Site-specific natural isotope fractionation
 (SNIF-NMR), 170–171
- Skin disease microorganisms, growth
 inhibition of, 284
- Slanted-soil-chamber system, 376–377
- Smell, sense of, 300. *See also* Olfactory
 stimulation
- Softer ionization energy, 179
- Solid-phase dynamic extraction (SPDE)
 technique, 28
- Solid-phase microextraction (SPME),
 24–28, 99. *See also* Headspace-SPME
 analysis method; SPME entries
 features of, 24
 principle of, 25–26
- Solvent, selection of, 21–22
- Solvent-extracted oil (SEO), 131
- Solvent extraction, 368–369
 ultrasound-accelerated, 20–24
 method of, 125
- Solvent-free microwave extraction
 (SFME), 14, 15, 17, 18
- Son, U-Sun, xii, 315
- Song, Hee-Sun, xii, 124, 264
- Sono-extraction, 22
- Sour citrus fruit, 3, 31, 111
- Sour orange oil, flavor and odor/aroma
 impact compounds in, 211–212
- Soxhlet, 22, 23
- Spain, citrus production in, 145
- Spanish clementine oil, 212
- Spikenard, 299
- Spirit, 9
- Split injection, 39
- SPME assay procedure, 26
- SPME extraction, 24. *See also* Solid-phase
 microextraction (SPME)
- SPME sampling equipment, 25
- Spray-drying method, 366
- Stable isotope ratio analysis, 165, 168–179
- Static headspace aroma (SHA) dilution
 analysis, 213
- Stationary phases, 40
- Steam distillation, 9–10, 12
 of yuzu, 369–370
- Stored citrus oils, antioxidative activities
 of, 241
- Sudachi, 107
- Sudachi oil, flavor and odor/aroma impact
 compounds in, 215
- Sulfur-containing compounds, 220
- Surface of nasal impact frequency (SNIF),
 207
- Sustainable growth, 13

- Sweet citrus fruit, 3
- Sweet orange oil, flavor and odor/aroma impact compounds in, 208–211
- Sweet oranges
- African, 47
 - American, 74–76
 - Chinese, 86–94
 - European, 149–150
 - Japanese, 111
 - Kenyan, 48–58
 - Rwandan, 66–67
 - Ugandan, 66
 - Vietnamese, 133–139
- Synthetic antioxidants, 5, 232
- Synthetic aroma substances, 349–350
- Synthetic citrus aroma substances, 350–351
- Tachibana, 106
- Tahiti lime, 107
- Tangerines. *See* Mandarins
- Temple tangor, 68–69
- Terpene aldehydes, 59, 62, 65, 66
- Terpene component, antimicrobial activities of, 285–289
- Terpene hydrocarbons
- in chemoprevention of cancers, 250, 252
 - effect on NDMA formation, 260–263
 - in mandarin oil, 334
- Terpenes, 5
- radical-scavenging activity and, 236–237
- Terpene thiols, 220
- Terpenoids, antibioactivity of, 288
- α -Terpinene, 261–263
- antimicrobial activities of, 271
- α -Terpinene derivative, 261, 263
- γ -Terpinene, 77, 96, 115–116, 124, 236, 239, 241, 320
- interconversion of, 329–330
- Tetradecane, 93
- Thao, Nguyen Thi, xii, 70, 165, 166, 168, 179
- Thermal conductivity detector (TCD), 42
- Thin-layer chromatography (TLC), 165
- Thymol, 285
- antimicrobial activities of, 274
- Time-intensity techniques, 203, 206–207
- Tisserand, Robert, 299–300
- Tocopherol analysis, of citrus peel essential oils, 241–243
- Tocopherols, 246
- Top note of flavors, 356
- Tosa-buntan, 115
- Total ion chromatogram (TIC), 360–361
- Treated wastewater, qualities of, 378
- Triacetin, 326
- Trolox, antioxidant activity of, 233
- Turbodistillation, 12, 13
- Two-dimensional GC (GC H GC), 45–46
- Ugandan citrus essential oils, volatile components of, 51–57
- Ugandan citrus fruits, compositional analysis of, 64–66
- Ugandan sweet oranges, 66
- Ujukitsu oil, 111
- Ultrasonic cleaning bath, 22
- Ultrasonic distillation, 370
- Ultrasonic irradiation, 370
- Ultrasonic reactors, 22, 23
- Ultrasonic steam-distilled oil (USDO), 370–374
- Ultrasound-accelerated extraction (UAE), 21, 22–24
- Ultrasound-accelerated solvent extraction, 20–24
- Ultrasound cavitation, 21
- Unidentified citrus species, Japanese, 116–124
- Unshiu mandarins, 95
- “Upside-down” microwave alembic, 15, 18
- Uruguayan clementine mandarins, volatile components in, 78–82
- Uruguayan lemon oils, 70, 71–73, 77–83
- Uruguayan mandarins, 76, 77
- volatile components in, 78–82
- Vacuum microwave hydro distillation (VMHD), 15
- Valencene, 88–92, 133, 150, 211
- Valencia navel fruits, 48–58
- Valencia orange oil, Charm values of, 210
- Valencia oranges
- sabinene content of, 88
 - volatile components of, 50–56
- Vegetable essential oils, 5
- Vegetable extracts, effects on NDMA formation, 253–257

- Vegetable extracts with yuzu oil and saliva, effects on NDMA formation, 255–256, 257–258
- Vegetables, inhibition of NDMA formation in, 252–253
- Vegetable samples
for NDMA assay, 253
nitrate, nitrite, and ascorbic acid contents in, 255–256
- Venezuelan lemon oils, 70, 71–73
- Venezuelan mandarins, 76
- Venezuelan orange oil, volatile composition of, 74
- Vietnamese citrus essential oil(s)
analyzing, 44
composition analysis of, 43–44
isotope fingerprint of, 187–188
isotope ratio of monoterpene hydrocarbons in, 180, 181
isotope ratio of the base peak for monoterpene hydrocarbons in, 183–186
volatile components of, 134–138
- Vietnamese citrus fruits, 132–145
lime, 143–145
mandarin, 139–141
orange, 133–139
pummelo, 141–143
- Volatile aroma component, antimicrobial activities of, 285–289
- Volatile chemical compounds, 48
- Volatile components. *See also* Volatile aroma component; Volatile compounds; Volatile flavor components
in African citrus essential oils, 50–57
in American lemon essential oils, 71–73
in American lime essential oils, 83–85
in American mandarin essential oils, 78–82
in American orange essential oils, 75–76
in Chinese citrus fruits, 89–91, 94–96
of citrus essential oils, 249–251
extraction problems related to, 29
in Indonesian citrus, 97–98
in Italian citrus essential oils, 147
in Japanese citrus essential oils, 108–110, 112–114, 117–123
in Vietnamese citrus essential oils, 134–138
in Vietnamese citrus fruits, 143–144
- Volatile compounds, 10
analytical studies on, 38, 171
antimicrobial activities of, 271–277
in Japanese citrus essential oil, 106
in Pontianak citrus juice, 99
under storage conditions, 322–323
in Vietnamese citrus fruits, 133
yuzu peel, 125–130
- Volatile flavor components
of Korean citrus essential oils, 126–127
of USDO, CSDO, and CPO, 372–373
- Volatile oils, 9
- Volatile profile characteristics, of Korean citrus essential oils, 128–129
- Volatile terpenes, antibacterial activity of, 286
- Washington navel fruits, 48–58
- Washington navel oranges
Chinese, 86, 93–94
volatile components of, 50–56
- Water-cleaning system, 376–377
- Water distillation, 12
- Water-free microwave extraction, 14
- Water-soluble liquid flavors, 354, 364
- White adipose tissue (WAT), 306–311
- White adipose tissue sympathetic nerve activity (WAT-SNA), 307–311
- Wijayi, C. Hanny, xii, 96
- Yeast capsules, 366–367
- Yeasts, use in microbial assay, 264–277
- Yu, Xiaolin, xii, 86
- Yukawa, Chiyoki, xii, 345
- Yuko, 107
- Yuko oil, 240
- Yuzu, 31
Chinese, 86
commercial value of, 367–368
inhibitory effect of, 263
Japanese, 106
Korean, 125–130
post-squeezing waste from, 370

- Yuzu buri, 378
- Yuzu essential oil(s), 5, 125, 368
 - effect on NDMA Formation in Vegetables, 252–253
 - effects of isotope ratio values in, 175
 - isotope fingerprint of, 186–187
 - isotope ratio of, 195–196
 - isotope ratios of linalool in, 190–191
 - products of, 378
- Yuzu flavor formulations, for beverages, 363
- Yuzu flavors, manufacturing and formulation of, 360–361
- Yuzu fruit juice extractor, 31–32
- Yuzu juice, 368
- Yuzunone, 215
- Yuzu oil
 - autonomic nerve activity and, 306–311
 - extraction of, 368–374
 - flavor and odor/aroma impact
 - compounds in, 214–215
 - NDMA formation and, 253–257, 258
- Yuzu peel, 125–130, 368
- Yuzu peel cold-pressed oil, components identified in, 362
- Yuzu peel oil, 106
- Yuzu residue, treatment of, 374–377
- Yuzu samples, linalool content of, 179
- Yuzu scent, olfactory stimulation with, 307–311
- Yuzu waste residue, practical uses of, 377–378
- Yuzu wastewater treatment, post-distilled, 375–377