



August 26, 2015
Revised: October 12, 2016

Ana Straabe
Baldwin Hills Regional Conservation Authority
Los Angeles River Center and Gardens
570 West Avenue 26, Suite 100
Los Angeles, California 90065

VIA EMAIL
Ana.Straabe@mrca.ca.gov

Subject: Results of Special Status Plant Surveys for the Blair Hills Corridor (Segment C) Portion of the Proposed Park to Playa Trail Project, Los Angeles County, California

Dear Ms. Straabe:

This Letter Report presents the findings of special status plant surveys conducted for the Blair Hills Corridor (Segment C) portion of the proposed Park to Playa Trail project (hereinafter referred to as the “proposed Segment C project”) located in Los Angeles County, California (Exhibit 1).

PROJECT DESCRIPTION AND LOCATION

The objective of the proposed Segment C project is to link existing trails in public parks and open spaces in the Baldwin Hills area. The proposed Segment C project includes a new trail leading down from the parking lot of the Baldwin Hills Scenic Overlook; along the northern end of a retention basin on the parcel owned by the Baldwin Hills Regional Conservation Authority (BHRCA) and on the slopes around the parcel that were previously developed with a school and where the Stoneview Nature Center is under construction; down to the northeastern section of BHRCA-owned surface property that is part of the active Inglewood Oilfield, and the proposed pedestrian bridge over La Cienega Boulevard to connect to an existing trail at the Kenneth Hahn State Recreation Area (KHSRA).

The trail improvements on the project site would include an at-grade compacted earth trail; identification, wayfinding and regulation signs; an interpretive node (e.g., information kiosk, shade structure, benches, and trash can); landscaping with native plants and restoring habitat in disturbed areas and adjacent to the trail; relocation of the access road, installation of new fencing and relocation of the fence, relocation of the water line, and installation of a drip irrigation system; undergrounding of existing utility lines along La Cienega Boulevard; and construction of a pedestrian bridge over La Cienega Boulevard.

The approximate 18-acre project site is located in the Baldwin Hills area of the western section of Los Angeles County in the City of Culver City. It is bound by La Cienega Boulevard on the east; private properties under active oil and gas exploration, production, processing, and associated activities on the south and southwest; the Baldwin Hills Scenic Overlook to the northwest; and single-family homes, Blair Hills Park, and the Stoneview Nature Center site (under construction) on the north. The site was formerly used for oil operations. Vegetation in this area is primarily a mix of sage scrub and grassland, with a small amount of chaparral and mulefat thicket (Exhibit 2).

METHODS

Botanical surveys were floristic in nature and consistent with the protocols created by the California Department of Fish and Wildlife (CDFW) (CDFG 2009). Prior to the field surveys, a literature search was conducted to identify special status plant species reported from the vicinity of the survey area. Sources reviewed include the U.S. Geological Survey's (USGS') Beverly Hills, Hollywood, Los Angeles, South Gate, Inglewood, and Venice 7.5-minute quadrangles in the California Native Plant Society's (CNPS') Locational Inventory of Rare and Endangered Vascular Plants of California (CNPS 2015) and the CDFW's California Natural Diversity Database (CNDDDB) (CDFW 2015).

Rainfall received in the winter and spring determines the germination of many annual and perennial herb species. According to the California Irrigation Management Information System (CIMIS), the region (data taken from Santa Monica) received 8.5 inches of precipitation between October 1, 2014 and May 31, 2015; the average for that time period (data taken from October 1993 to May 2015) was approximately 13.3 inches (CIMIS 2015).

Reference populations were monitored for annual and difficult-to-detect target species to ensure that the surveys were comprehensive. This is especially relevant during periods of unusual rainfall patterns or below average rainfall. If conditions at a nearby reference population are suitable for germination and growth, then it can be inferred that conditions would also be suitable in the survey area. Table 1 summarizes the flowering status of known reference populations monitored during the 2015 special status plant survey period. Reference populations were not monitored for species with a California Rare Plant Rank (CRPR) of 3 or 4; large perennials (e.g., Nuttall's scrub oak [*Quercus dumosa*]), which would be identifiable throughout the year; or for species with no extant, publically accessible reference population in the project region.

TABLE 1
REFERENCE POPULATION BLOOMING DATES

Species	Area Monitored for Blooming	Date Observed Blooming
<i>California macrophylla</i> round-leaved filaree	Tehachapi Mountains	March 23, 2015
<i>Chorizanthe parryi</i> var. <i>fernandina</i> San Fernando Valley spineflower	West of the San Fernando Valley	April 24, 2015
<i>Dudleya multicaulis</i> many-stemmed dudleya	University of California, Irvine	March 24, 2015

The survey area was surveyed by BonTerra Psomas Senior Biologist Allison Rudalevige on April 16 and June 25, 2015. The total number of person-hours spent was approximately eight hours. Potentially suitable habitat for special status plant species in the survey area was systematically surveyed during the site visits. All plant species observed were recorded in field notes. Plant species were identified in the field or collected for later identification. Plants were identified to the taxonomic level necessary to determine whether or not they are a special status species. Plants were identified using taxonomic keys, descriptions, and illustrations in Baldwin et al. (2012), Hickman (1993), and Munz (1974). Taxonomy and nomenclature follows the Jepson Herbarium (2014), Hickman (1993), and current scientific journals for scientific and common names.

Special status plant species were mapped either with a Global Positioning System (GPS) unit or onto a one inch equals 150 feet (1" = 150') scale aerial when the population was not directly accessible. Data was collected on the number and phenology of individuals; microsite characteristics such as slope, aspect, soil texture, and surrounding habitat; and associated species.

SURVEY RESULTS

Table 2 identifies the special status plant species with potential to occur in the survey area and the survey results. One special status species, Southern California black walnut (*Juglans californica*), was observed during the survey effort and is discussed below. A list of all plant species observed in the survey area during the surveys can be found in Attachment A. Representative photographs are provided in Attachment B.

TABLE 2
SPECIAL STATUS PLANT SPECIES REPORTED
FROM THE SURVEY AREA VICINITY

Species	Status			Habitat Suitability and Survey Results
	USFWS	CDFW	CRPR	
<i>Abronia maritima</i> red sand-verbena	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Arenaria paludicola</i> marsh sandwort	FE	SE	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Astragalus brauntonii</i> Braunton's milkvetch	FE	–	1B.1	Marginally suitable habitat, but not observed during focused surveys.
<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i> Ventura marsh milkvetch	FE	SE	1B.1	No suitable habitat; only known from one occurrence and not observed during focused surveys; not expected to occur.
<i>Astragalus tener</i> var. <i>titi</i> coastal dunes milkvetch	FE	SE	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Atriplex parishii</i> Parish's brittle-scale	–	–	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Atriplex serenana</i> var. <i> davidsonii</i> Davidson's salt-scale	–	–	1B.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>California macrophylla</i> round-leaved filaree	–	–	1B.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Calochortus catalinae</i> Catalina mariposa lily	–	–	4.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Calochortus plummerae</i> Plummer's mariposa lily	–	–	4.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Calystegia felix</i> lucky morning-glory*	–	–	3.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Calystegia sepium</i> ssp. <i>binghamiae</i> Santa Barbara morning-glory	–	–	1A	No suitable habitat; considered extinct and not observed during focused surveys; not expected to occur.

TABLE 2
SPECIAL STATUS PLANT SPECIES REPORTED
FROM THE SURVEY AREA VICINITY

Species	Status			Habitat Suitability and Survey Results
	USFWS	CDFW	CRPR	
<i>Camissoniopsis lewisii</i> [<i>Camissonia l.</i>] Lewis' evening-primrose	–	–	3	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Centromadia parryi</i> ssp. <i>australis</i> southern tarplant	–	–	1B.1	Potentially suitable habitat, but not observed during focused surveys.
<i>Chaenactis glabriuscula</i> var. <i>orcuttiana</i> Orcutt's pincushion	–	–	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Chenopodium littoreum</i> coastal goosefoot	–	–	1B.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Chloropyron maritimum</i> ssp. <i>maritimum</i> salt marsh bird's-beak	FE	SE	1B.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Chorizanthe parryi</i> var. <i>fernandina</i> San Fernando Valley spineflower	FC	SE	1B.1	Marginal potentially suitable habitat, but not observed during focused surveys.
<i>Convolvulus simulans</i> small-flowered morning-glory	–	–	4.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Deinandra paniculata</i> paniculate tarplant	–	–	4.2	Suitable habitat, but not observed during focused surveys.
<i>Dichondra occidentalis</i> western dichondra	–	–	4.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Dithyrea maritima</i> beach spectaclepod	–	ST	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Dudleya multicaulis</i> many-stemmed dudleya	–	–	1B.2	Potentially suitable habitat, but not observed during focused surveys.
<i>Eryngium aristulatum</i> var. <i>parishii</i> San Diego button-celery	FE	SE	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Erysimum insulare</i> island wallflower	–	–	1B.3	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Erysimum suffrutescens</i> suffrutescent wallflower	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Frankenia palmeri</i> Palmer's frankenia	–	–	2B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Helianthus nuttallii</i> ssp. <i>parishii</i> Los Angeles sunflower	–	–	1A	No suitable habitat; considered extinct and not observed during focused surveys; not expected to occur.
<i>Hesperocyparis macrocarpa</i> Monterey cypress	–	–	1B.2	No suitable habitat and not observed during focused surveys; not expected to occur.

TABLE 2
SPECIAL STATUS PLANT SPECIES REPORTED
FROM THE SURVEY AREA VICINITY

Species	Status			Habitat Suitability and Survey Results
	USFWS	CDFW	CRPR	
<i>Hordeum intercedens</i> bobtail barley	–	–	3.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Horkelia cuneata</i> var. <i>puberula</i> mesa horkelia	–	–	1B.1	Marginal potentially suitable habitat, but not observed during focused surveys.
<i>Juglans californica</i> Southern California black walnut	–	–	4.2	Suitable habitat present. Observed during focused surveys.
<i>Juncus acutus</i> ssp. <i>leopoldii</i> southwestern spiny rush	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Lasthenia glabrata</i> ssp. <i>coulteri</i> Coulter's goldfields	–	–	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Lepidium virginicum</i> var. <i>robinsonii</i> Robinson's pepper-grass	–	–	4.3	Potentially suitable habitat, but not observed during focused surveys.
<i>Mucronea californica</i> California spineflower	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Nama stenocarpum</i> mud nama	–	–	2B.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Nasturtium gambelii</i> Gambel's water cress	FE	ST	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Navarretia fossalis</i> spreading navarretia	FT	–	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Navarretia prostrata</i> prostrate vernal pool navarretia	–	–	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Orcuttia californica</i> California Orcutt grass	FE	SE	1B.1	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Phacelia hubbyi</i> Hubby's phacelia	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Phacelia ramosissima</i> var. <i>austrolitoralis</i> * south coast branching phacelia	–	–	3.2	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Phacelia stellaris</i> Brand's star phacelia	–	–	1B.1	Potentially suitable habitat, but not observed during focused surveys.
<i>Potentilla multijuga</i> Ballona cinquefoil	–	–	1A	No suitable habitat and not observed during focused surveys; not expected to occur.
<i>Pseudognaphalium leucocephalum</i> white rabbit-tobacco	–	–	2B.2	No suitable habitat and not observed during focused surveys; not expected to occur.

TABLE 2
SPECIAL STATUS PLANT SPECIES REPORTED
FROM THE SURVEY AREA VICINITY

Species	Status			Habitat Suitability and Survey Results																
	USFWS	CDFW	CRPR																	
<i>Quercus dumosa</i> Nuttall's scrub oak	–	–	1B.1	Potentially suitable habitat, but not observed during focused surveys.																
<i>Ribes divaricatum</i> var. <i>parishii</i> Parish's gooseberry	–	–	1A	No suitable habitat; considered extinct and not observed during focused surveys; not expected to occur.																
<i>Sidalcea neomexicana</i> salt spring checkerbloom	–	–	2B.2	No suitable habitat and not observed during focused surveys; not expected to occur.																
<i>Suaeda esteroa</i> estuary seablite	–	–	1B.2	No suitable habitat and not observed during focused surveys; not expected to occur.																
<i>Suaeda taxifolia</i> woolly seablite	–	–	4.2	No suitable habitat and not observed during focused surveys; not expected to occur.																
<i>Symphyotrichum defoliatum</i> San Bernardino aster	–	–	1B.2	Potentially suitable habitat, but not observed during focused surveys.																
<i>Symphyotrichum greatae</i> Greata's aster	–	–	1B.3	No suitable habitat and not observed during focused surveys; not expected to occur.																
USFWS: U.S. Fish and Wildlife Service; CDFW: California Department of Fish and Wildlife; CRPR: California Rare Plant Rank LEGEND: <table style="width: 100%; border: none;"> <tr> <td colspan="2">Federal (USFWS)</td> <td colspan="2">State (CDFW)</td> </tr> <tr> <td>FE</td> <td>Endangered</td> <td>SE</td> <td>Endangered</td> </tr> <tr> <td>FT</td> <td>Threatened</td> <td>ST</td> <td>Threatened</td> </tr> <tr> <td>FC</td> <td>Candidate</td> <td></td> <td></td> </tr> </table> CRPR 1A Plants presumed extirpated in California and either rare or extinct elsewhere 1B Plants Rare, Threatened, or Endangered in California and elsewhere 2B Plants Rare, Threatened, or Endangered in California but more common elsewhere 3 Plants about which we need more information – A Review List 4 Plants of limited distribution – A Watch List CRPR Threat Code Extensions None Plants lacking any threat information .1 Seriously threatened in California (over 80% of occurrences threatened; high degree and immediacy of threat) .2 Fairly threatened in California (20–80% of occurrences threatened; moderate degree and immediacy of threat) .3 Not very threatened in California (<20% of occurrences threatened; low degree and immediacy of threat or no current threats known) Note: Taxa observed in the survey area are shown in boldface type. * Taxon not currently recognized by the Jepson Herbarium (2014), but it is still tracked by the CDFW and/or CNPS.					Federal (USFWS)		State (CDFW)		FE	Endangered	SE	Endangered	FT	Threatened	ST	Threatened	FC	Candidate		
Federal (USFWS)		State (CDFW)																		
FE	Endangered	SE	Endangered																	
FT	Threatened	ST	Threatened																	
FC	Candidate																			

Southern California Black Walnut

Southern California black walnut has a CRPR of 4.2. This deciduous tree occurs on hillsides and canyons at elevations between approximately 100 and 2,950 feet above msl (Baldwin et al. 2012). It is known from the Outer South Coast Ranges and is cultivated in the Santa Lucia Range and southwestern California, excluding the Channel Islands and the San Bernardino Mountains (Baldwin et al. 2012).


Four Southern California black walnut trees were observed adjacent to the parking lot at the west end of the survey area (Exhibit 3). The trees were all young and primarily in fruit. Associated species include coyote brush (*Baccharis pilularis* ssp. *consanguinea*), mule fat (*Baccharis salicifolia* ssp. *salicifolia*), and California brittlebush (*Encelia californica*).

CONCLUSIONS/RECOMMENDATIONS

One special status plant species was observed in the survey area during the 2015 survey. It is not anticipated that the Southern California black walnuts in the survey area would be impacted by proposed project activities. Although reference populations and regional rainfall amounts were monitored to ensure the scientific adequacy of these focused surveys, there is always a minimal potential for false negative survey results as species could possibly be present on a site but may not be detectable at the time of the surveys.

Sincerely,
BonTerra Psomas


Marc T. Blain
Senior Project Manager


Allison D. Rudalevige
Senior Biologist

Attachments: Exhibits 1–3
A – Plant Compendium
B – Site Photographs

R:\Projects\ALT\3ALT000302\Revised Plant Report_SegC-101216.docx

REFERENCES


- Baldwin, B.G., D.J. Keil, R. Patterson, T.J. Rosatti, and D.H. Wilken (Eds.). 2012. *The Jepson Manual: Vascular Plants of California* (Second ed.). Berkeley, CA: University of California Press.
- California Department of Fish and Game (CDFG). 2009 (November 24). *Protocols for Surveying and Evaluating Impacts to Special Status Native Plant Populations and Natural Communities*. Sacramento, CA: CDFG.
- California Department of Fish and Wildlife (CDFW). 2015 (April). California Natural Diversity Database. Records of Occurrence for Beverly Hills, Hollywood, Los Angeles, South Gate, Inglewood, and Venice U.S. Geological Survey 7.5-minute quadrangles. Sacramento, CA: CDFW, Natural Heritage Division.
- California Native Plant Society (CNPS). 2015. Locational Inventory of Rare and Endangered Vascular Plants of California (online edition, v8-02). Records of Occurrence for Beverly Hills, Hollywood, Los Angeles, South Gate, Inglewood, and Venice U.S. Geological Survey 7.5-minute quadrangles. Sacramento, CA: CNPS. <http://www.rareplants.cnps.org/advanced.html>.
- California Irrigation Management Information System (CIMIS). 2015. CIMIS Monthly Report for Santa Monica Station 99. Sacramento, CA: California Department of Water Resources, CIMIS. <http://www.cimis.water.ca.gov>.

Ana Straabe
August 26, 2015 (revised October 12, 2016)
Page 8

Hickman, J.C., Ed. 1993. *The Jepson Manual of Higher Plants of California*. Berkeley, CA: University of California Press.

Jepson Herbarium. 2014 (December 19, Revision 2). Jepson eFlora. Berkeley, CA: Jepson Herbarium.
<http://ucjeps.berkeley.edu/IJM.html>.

Munz, P.A. 1974. *A Flora of Southern California*. Berkeley, CA: University of California Press




 Survey Area

Aerial Source: LAR-IAC 2011


Project Location

Exhibit 1

Park to Playa Trail Project – Segment C



D:\Projects\Alta\003\MXD\PlantReport\SegmentC\Ex_Project_Location_20150821.mxd



Aerial Source: LAR-IAC 2011

Vegetation Types and Other Areas

Park to Playa Trail Project – Segment C




Exhibit 2





Survey Area
● Southern California Black Walnut

Aerial Source: LAR-IAC 2011

Special Status Species
 Park to Playa Trail Project – Segment C

Exhibit 3



D:\Projects\Alta\003\Map\PlantReport\SegmentC\Ex3_SSPlants_20150821.mxd

ATTACHMENT A
PLANT COMPENDIUM

PLANT SPECIES OBSERVED IN THE SURVEY AREA DURING SURVEYS

Species	
Scientific Name	Common Name
GYMNOSPERMS	
PINACEAE - PINE FAMILY	
<i>Pinus</i> sp.	pine
EUDICOTS	
ADOXACEAE - MUSKROOT FAMILY	
<i>Sambucus nigra</i> ssp. <i>caerulea</i>	blue elderberry
AIZOACEAE - FIG-MARIGOLD FAMILY	
<i>Carpobrotus edulis</i> *	freeway iceplant
ANACARDIACEAE - SUMAC FAMILY	
<i>Schinus molle</i> *	Peruvian pepper tree
<i>Schinus terebinthifolius</i> *	Brazilian pepper tree
<i>Toxicodendron diversilobum</i>	western poison oak
APIACEAE - CARROT FAMILY	
<i>Foeniculum vulgare</i> *	fennel
ARALIACEAE - GINSENG FAMILY	
<i>Hedera helix</i> *	English ivy
ASTERACEAE - SUNFLOWER FAMILY	
<i>Ambrosia acanthicarpa</i>	annual bur-sage
<i>Ambrosia psilostachya</i>	western ragweed
<i>Artemisia californica</i>	California sagebrush
<i>Baccharis pilularis</i> ssp. <i>consanguinea</i>	coyote brush
<i>Baccharis salicifolia</i> ssp. <i>salicifolia</i>	mule fat
<i>Carduus pycnocephalus</i> ssp. <i>pycnocephalus</i> *	Italian thistle
<i>Centaurea melitensis</i> *	toocalote
<i>Deinandra fasciculata</i>	fascicled tarplant
<i>Encelia californica</i>	California brittlebush
<i>Ericameria</i> sp.	goldenbush
<i>Erigeron canadensis</i>	horseweed
<i>Glebionis coronaria</i> *	crown daisy
<i>Grindelia camporum</i>	field gumplant
<i>Helminthotheca echioides</i> *	bristly ox-tongue
<i>Heterotheca grandiflora</i>	telegraph weed
<i>Hypochaeris glabra</i> *	smooth cat's-ear
<i>Lactuca serriola</i> *	prickly lettuce
<i>Malacothrix saxatilis</i>	cliff desert dandelion
<i>Pseudognaphalium californicum</i>	California everlasting
<i>Pseudognaphalium canescens</i>	hairy everlasting
<i>Pseudognaphalium luteoalbum</i> *	white lamb everlasting
<i>Pseudognaphalium microcephalum</i>	white head everlasting
<i>Silybum marianum</i> *	blessed milk thistle
<i>Sonchus oleraceus</i> *	common sow thistle
<i>Stephanomeria exigua</i>	small wire-lettuce
<i>Uropappus lindleyi</i>	silver puffs

PLANT SPECIES OBSERVED IN THE SURVEY AREA DURING SURVEYS

Species	
Scientific Name	Common Name
BORAGINACEAE - BORAGE FAMILY	
<i>Phacelia tanacetifolia</i>	lacy phacelia
BRASSICACEAE - MUSTARD FAMILY	
<i>Hirschfeldia incana</i> *	shortpod mustard
<i>Raphanus sativus</i> *	radish
CACTACEAE - CACTUS FAMILY	
<i>Opuntia ficus-indica</i> *	mission prickly pear
CARYOPHYLLACEAE - PINK FAMILY	
<i>Silene</i> sp.	catchfly
CHENOPODIACEAE - GOOSEFOOT FAMILY	
<i>Atriplex semibaccata</i> *	Australian saltbush
<i>Salsola tragus</i> *	Russian thistle
CONVOLVULACEAE - MORNING-GLORY FAMILY	
<i>Calystegia macrostegia</i>	coast morning-glory
CRASSULACEAE - STONECROP FAMILY	
<i>Crassula connata</i>	pygmyweed
CUCURBITACEAE - GOURD FAMILY	
<i>Cucurbita foetidissima</i>	calabazilla
<i>Marah macrocarpa</i>	chilicothe
EUPHORBIACEAE - SPURGE FAMILY	
<i>Euphorbia peplus</i> *	petty spurge
<i>Ricinus communis</i> *	castor bean
FABACEAE - LEGUME FAMILY	
<i>Acmispon glaber</i>	deerweed
<i>Cytisus multiflorus</i> *	Spanish broom
<i>Medicago polymorpha</i> *	California burclover
<i>Melilotus indicus</i> *	sourclover
FAGACEAE - OAK FAMILY	
<i>Quercus agrifolia</i>	coast live oak
GERANIACEAE - GERANIUM FAMILY	
<i>Erodium botrys</i> *	longbeak filaree
<i>Geranium rotundifolium</i> *	roundleaf geranium
JUGLANDACEAE - WALNUT FAMILY	
<i>Juglans californica</i>	southern California black walnut
MALVACEAE - MALLOW FAMILY	
<i>Malva parviflora</i> *	cheeseweed
MYRSINACEAE - MYRSINE FAMILY	
<i>Lysimachia arvensis</i> *	scarlet pimpernel
OLEACEAE - OLIVE FAMILY	
<i>Olea europaea</i> *	olive
ONAGRACEAE - EVENING PRIMROSE FAMILY	
<i>Oenothera elata</i> ssp. <i>hirsutissima</i>	great marsh evening primrose
PAPAVERACEAE - POPPY FAMILY	
<i>Eschscholzia californica</i>	California poppy

PLANT SPECIES OBSERVED IN THE SURVEY AREA DURING SURVEYS

Species	
Scientific Name	Common Name
POLYGONACEAE - BUCKWHEAT FAMILY	
<i>Eriogonum fasciculatum</i>	California buckwheat
ROSACEAE - ROSE FAMILY	
<i>Heteromeles arbutifolia</i>	toyon
RUBIACEAE - COFFEE FAMILY	
<i>Galium angustifolium</i>	narrow leaved bedstraw
<i>Galium aparine</i>	goose grass
SCROPHULARIACEAE - FIGWORT FAMILY	
<i>Verbascum virgatum*</i>	wand mullein
SOLANACEAE - NIGHTSHADE FAMILY	
<i>Datura wrightii</i>	Wright's jimson weed
<i>Nicotiana glauca*</i>	tree tobacco
<i>Solanum xanti</i>	chaparral nightshade
ZYGOPHYLLACEAE - CALTROP FAMILY	
<i>Tribulus terrestris*</i>	puncturevine
MONOCOTS	
ARECACEAE - PALM FAMILY	
<i>Phoenix canariensis*</i>	Canary Island palm
<i>Washingtonia robusta*</i>	Mexican fan palm
POACEAE - GRASS FAMILY	
<i>Arundo donax*</i>	giant reed
<i>Avena barbata*</i>	slender wild oat
<i>Avena fatua*</i>	wild oat
<i>Bromus diandrus*</i>	ripgut grass
<i>Bromus hordeaceus*</i>	soft chess
<i>Cortaderia selloana*</i>	pampas grass
<i>Cynodon dactylon*</i>	bermuda grass
<i>Distichlis spicata</i>	salt grass
<i>Elymus condensatus</i>	giant wild-rye
<i>Festuca perennis*</i>	rye grass
<i>Hordeum murinum*</i>	wall barley
<i>Pennisetum setaceum*</i>	crimson fountain grass
<i>Stipa miliacea</i> var. <i>miliacea*</i>	smilo grass
THEMIDACEAE - BRODIAEA FAMILY	
<i>Dichelostemma capitatum</i>	blue dicks
* non-native	

ATTACHMENT B
SITE PHOTOGRAPHS



Southern California black walnut (*Juglans californica*) observed in the survey area.



Southern California black walnut (*Juglans californica*) observed in the survey area.

D:\Projects\Alta\003\Graphics\Plant_Report\SegC\Ex_SP1_20150821.ai

Site Photographs

Park to Playa Trail Project – Segment C

Attachment B

