

## **Appendix A.8.19**

### **Habitat Survey Results - Species Lists**

## A.8.19

---

This appendix presents the species lists for habitats recorded along, and adjacent to, the proposed road development.

**Section A.8.19.1** presents the species lists for those habitat areas within the ecological sites. **Section A.8.19.2** presents the species lists for habitat areas between the ecological sites but excluding the area between the River Corrib and Lackagh Quarry (which is described separately in **Section A.8.19.3**). These areas are shown on **Figures 1-4 (Section A.8.19.5)**.

**Section A.8.19.4** presents the relevé data collected along and adjacent to the proposed road development, as referred to in **Sections A.8.19.1 to A.8.19.3**. The locations of the relevés, and the reference codes, are also shown on **Figures 1 to 4**.

Where referred to, the habitat codes for non-Annex I habitat types are after *A Guide to Habitats in Ireland* (Fossitt, 2000). Annex I habitat codes are as per the *Interpretation Manual of European Union Habitats* (CEC, 2013). Where vegetation community codes are referred to in the sections below, they are as per the following reports (as relevant):

- *Turlough Hydrology, Ecology and Conservation* (Waldren, S. 2015, Ed.)
- *The Irish semi-natural grasslands survey 2007-2012. Irish Wildlife Manuals, No. 78* (O'Neill et al., 2013)
- *Results of monitoring survey of old sessile oak woods and alluvial forests. Irish Wildlife Manuals, No. 71* (O'Neill, F.H. & Barron, S.J., 2013)
- *National survey of limestone pavement and associated habitats in Ireland. Irish Wildlife Manuals, No. 73* (Wilson, S. and Fernández, F., 2013)
- *Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland. Version 2.0. Irish Wildlife Manuals, No. 79* (Perrin et al., 2014)
- *National Survey of Native Woodlands 2003-2008. Volume II: Woodland Classification* (Perrin et al., 2008)

Nomenclature for vascular plant species follows *New Flora of the British Isles*, 3rd Edition (Stace, 2010) and for bryophyte species follows the *Checklist of British and Irish bryophytes* (British Bryological Society, 2009<sup>1</sup>).

---

<sup>1</sup> <http://rbg-web2.rbge.org.uk/bbs/Resources/Downloads.htm>

## A.8.19.1

---

## Habitat Species Lists – Ecological Sites

**Table A.8.19.1** presents the species lists for those habitat areas within the ecological sites. It does not include species lists for the residential habitat category (see **Section 8.3.4.34** of **Chapter 8, Biodiversity** of the EIA Report) and it also does not include species lists for the aquatic habitats, which are described in full in **Section 8.3.4** of **Chapter 8, Biodiversity** and in **Appendix A.8.20 (Results of the 2014 N6 Galway City Transport Project Aquatic Habitat Surveys)**.

Where relevés were taken in the field, the reference codes are provided in the Key Species column (e.g. EC03 R1). These relevé codes refer to the full species lists which are provided in **Appendix A.8.19.4**.

**Table A.8.19.1: Habitat Species Lists Relating to Ecological Sites within, or adjacent to, the Proposed Road Development**

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC03</b>		
CB1	1220	<i>Elytrigia repens</i> , <i>Senecio jacobaea</i> , <i>Leontodon autumnalis</i> , <i>Lotus corniculatus</i> , <i>Rumex crispus</i> and <i>Plantago lanceolata</i> .
CM2	1330	<i>Agrostis stolonifera</i> , <i>Festuca rubra</i> , <i>Eleocharis uniglumis</i> , <i>Triglochin maritima</i> , <i>Leontodon autumnalis</i> , <i>Potentilla anserina</i> , <i>Lotus corniculatus</i> and <i>Cochlearia officinalis</i> .
FS1	n/a	<i>Phragmites australis</i> and <i>Filipendula ulmaria</i> .
GA1	n/a	<i>Lolium perenne</i> , <i>Holcus lanatus</i> , <i>Trifolium repens</i> , <i>Ranunculus repens</i> and <i>Rumex</i> sp.
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Trifolium pratense</i> , <i>Holcus lanatus</i> , <i>Centaurea nigra</i> , <i>Trifolium repens</i> , <i>Agrostis capillaris</i> , <i>Poa pratensis</i> , <i>Trifolium repens</i> , <i>Senecio jacobaea</i> and <i>Cynosurus cristatus</i> .
GS3	n/a	<i>Anthoxanthum odoratum</i> , <i>Agrostis capillaris</i> , <i>Potentilla erecta</i> , <i>Trifolium repens</i> and <i>Poa pratensis</i> .
GS4	n/a	<i>Juncus effusus</i> and <i>Juncus acutiflorus</i> .
GM1	n/a	<i>Iris pseudacorus</i> , <i>Filipendula ulmaria</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Anagallis tenella</i> , <i>Carex nigra</i> , <i>Hydrocotyle vulgaris</i> and <i>Senecio aquaticus</i> .
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Molinia caerulea</i> , <i>Anthoxanthum odoratum</i> , <i>Potentilla erecta</i> , <i>Succisa pratensis</i> and <i>Daboecia cantabrica</i> <b>EC03 R1</b>
WS1	n/a	<i>Ulex europaeus</i> , <i>Prunus spinosa</i> and <i>Rubus fruticosus</i> .

<sup>2</sup> Fossitt (2000)

<sup>3</sup> Where applicable, vegetation community codes are given in parenthesis.

\*8240 (LPE) = exposed limestone pavement; \*8240 (LPW) = wooded limestone pavement

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC05</b>		
FW1/GM1	n/a	<i>Iris pseudacorus</i> , <i>Apium nodiflorum</i> , <i>Potentilla palustris</i> , <i>Filipendula ulmaria</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Salix cinerea</i> ,
GS3	n/a	<i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Cynosurus cristatus</i> , <i>Lolium perenne</i> , <i>Hypochaeris radicata</i> , <i>Achillea millefolium</i> , <i>Trifolium pratense</i> , <i>Veronica officinalis</i> , <i>Prunella vulgaris</i> , <i>Trifolium repens</i> , <i>Rumex obtusifolius</i> , <i>R acetosella</i> , <i>Rhytidiadelphus squarrosus</i> . <i>Sedum anglicum</i> , <i>Dicranum</i> sp., <i>Succisa pratensis</i> , <i>Euphrasia</i> sp., <i>Hypnum cupressiforme</i>
GS4	n/a	<i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Holcus lanatus</i> , <i>Agrostis stolonifera</i> , <i>Ranunculus repens</i> , <i>Hydrocotyle vulgaris</i> , <i>Lotus pedunculatus</i> , <i>Cirsium palustre</i> , <i>Lythrum salicaria</i> , <i>Rumex acetosa</i> , <i>Galium palustre</i> , <i>Potentilla erecta</i> , <i>Calliergonella cuspidata</i> , <i>Rhytidiadelphus squarrosus</i>
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Salix cinerea</i> , <i>Rubus fruticosus</i> agg. <i>Calystegia sepium</i> .
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Erica cinerea</i> , <i>Daboecia cantabrica</i> , <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Agrostis capillaris</i> . <i>Succisa pratensis</i> , <i>Teucrium scorodonia</i> , <i>Hylocomium splendens</i> , <i>Solidago virgaurea</i> and <i>Viola</i> sp. <b>EC05 R1</b>
PF2	n/a	<i>Juncus articulatus</i> , <i>Juncus effusus</i> , <i>Hypericum elodes</i> , <i>Hydrocotyle vulgaris</i> , <i>Anagallis tenella</i> , <i>Carex echinata</i> , <i>Carex nigra</i> , <i>Ranunculus flammula</i> and <i>Sphagnum</i> sp.
WS1	n/a	<i>Prunus spinosa</i> , <i>Salix cinerea</i> , <i>Crataegus monogyna</i> , <i>Ulex europaeus</i> , <i>Hedera helix</i> , <i>Pteridium aquilinum</i> .
WS1/HD1/ GS3	n/a	<i>Prunus spinosa</i> , <i>Salix cinerea</i> , <i>Crataegus monogyna</i> , <i>Ulex europaeus</i> , <i>Hedera helix</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Calystegia sepium</i> , <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Cynosurus cristatus</i> , <i>Lolium perenne</i> , <i>Hypochaeris radicata</i> , <i>Achillea millefolium</i> , <i>Trifolium pratense</i> , <i>Veronica officinalis</i> , <i>Prunella vulgaris</i> , <i>Trifolium repens</i> , <i>Rumex obtusifolius</i> , <i>R acetosella</i> , <i>Rhytidiadelphus squarrosus</i> . <i>Sedum anglicum</i> , <i>Dicranum</i> sp., <i>Succisa pratensis</i> , <i>Euphrasia</i> sp., <i>Hypnum cupressiforme</i> . Dense bracken tends to shade out much of the grassland species beneath
<b>EC09</b>		
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i> .
ED3	n/a	<i>Tussilago farfara</i> , <i>Ulex europaeus</i> , <i>Ulex gallii</i> , <i>Centaurea nigra</i> , <i>Hypochaeris radiata</i> , <i>Daucus carota</i> , <i>Lotus corniculatus</i> , <i>Carex disticha</i> , <i>Hypericum pulchellum</i> , <i>Achillea millefolium</i> , <i>Carex panacea</i> , <i>Potentilla anserina</i> , <i>Juncus effusus</i> , <i>Salix cinerea</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
FS1	n/a	<i>Phragmites australis</i> , <i>Agrostis stolonifera</i> , <i>Filipendula ulmaria</i> , <i>Juncus articulatus</i> and <i>Potentilla palustris</i> .
GA1	n/a	<i>Lolium perenne</i> , <i>Urtica dioica</i> , <i>Dactylis glomerata</i> , <i>Rumex obtusifolius</i> , <i>Ranunculus repens</i> , <i>Agrostis stolonifera</i> , <i>Cirsium arvense</i> , <i>Polygonum aviculare</i> , <i>Epilobium montanum</i> , <i>Plantago lanceolata</i> , <i>Holcus lanatus</i> , <i>Calystegia sepium</i> , <i>Rubus fruticosus</i> agg., <i>Persicaria maculosa</i> , <i>Sonchus asper</i> , <i>Brassica napus</i> , <i>Trifolium repens</i> and <i>Poa annua</i> .
GS2	n/a	<i>Dactylis glomerata</i> , <i>Arrhenatherum elatius</i> , <i>Rumex acetosa</i> , <i>Festuca rubra</i> , <i>Lotus corniculatus</i> , <i>Anthoxanthum odoratum</i> , <i>Agrostis stolonifera</i> , <i>Succisa pratensis</i> , <i>Centaurea nigra</i> and <i>Veronica chamaedrys</i> . Some <i>Thymus polytrichus</i> on ant hills only. Recolonized ground has some nutrient enrichment indicators such as <i>Cirsium arvense</i> and <i>Rumex obtusifolius</i> .
GS3	n/a	Typical species are <i>Agrostis capillaris</i> , <i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Plantago lanceolata</i> , <i>Anthoxanthum odoratum</i> , <i>Nardus stricta</i> , <i>Rumex acetosa</i> , <i>Succisa pratensis</i> , <i>Lotus corniculatus</i> , <i>Ranunculus repens</i> , <i>Leontodon autumnalis</i> , <i>Centaurea nigra</i> , <i>Potentilla erecta</i> and <i>Centaureum erythraea</i> .
	n/a	Highly species-rich variant recorded with abundant <i>Succisa pratensis</i> , on infertile ground.
	n/a	Slightly calcareous variant also recorded with <i>Lotus corniculatus</i> , <i>Prunella vulgaris</i> , <i>Carex disticha</i> and <i>Linum catharticum</i> .
GS4	n/a	More species poor variant containing species such as <i>Juncus effusus</i> , <i>Holcus lanatus</i> , <i>Lythrum salicaria</i> , <i>Ranunculus repens</i> , <i>Ranunculus acris</i> , <i>Filipendula ulmaria</i> , <i>Potentilla anserina</i> , <i>Agrostis stolonifera</i> , <i>Juncus articulatus</i> , <i>Galium palustre</i> , <i>Lolium perenne</i> , <i>Dactylis glomerata</i> , <i>Lotus pedunculatus</i> and <i>Cirsium palustre</i> .
	6410 (GL1c)	<i>Molinia</i> meadow in mosaic with acid grassland. A high-quality habitat with high species-richness (38 spp per 2m <sup>2</sup> ). Typical species include <i>Cirsium dissectum</i> , <i>Rhinanthus minor</i> , <i>Ranunculus flammula</i> , <i>Prunella vulgaris</i> , <i>Carex nigra</i> , <i>Carex demissa</i> , <i>Pedicularis sylvatica</i> , <i>Lophocolea bidentata</i> , <i>Riccardia</i> cf. <i>multifida</i> and <i>Calliergonella cuspidata</i> . <b>EC09 R1</b>
	6410 (GL1d)	<i>Molinia</i> meadow in mosaic with wet heath and has a low species-richness and forb cover. Typical species include <i>Molinia caerulea</i> , <i>Juncus conglomeratus</i> , <i>Juncus articulatus</i> , <i>Anthoxanthum odoratum</i> , <i>Succisa pratensis</i> , <i>Carex echinata</i> , <i>Carex panacea</i> , <i>Potentilla erecta</i> and <i>Pleurozium schreberi</i> . <b>EC09 R4</b>
GM1	n/a	<i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>angelica sylvestris</i> , <i>Filipendula ulmaria</i> , <i>Arrhenatherum elatius</i> , <i>Agrostis stolonifera</i> , <i>Lathyrus pratensis</i> , <i>Stellaria graminea</i> , <i>Equisetum fluviatile</i> , <i>Hydrocotyle vulgaris</i> , <i>Juncus effusus</i> and <i>Juncus articulatus</i> .
HD1	n/a	<i>Pteridium aquilinum</i> with patches of <i>Rubus fruticosus</i> agg. <i>Epilobium hirsutum</i> . Often grading into <i>Ulex europaeus</i> scrub.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
HH1	4030 (DH3, DH1)	<i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Cladonia portentosa</i> , <i>Carex binervis</i> , <i>Potentilla erecta</i> , <i>Cladonia uncialis</i> , <i>Carex panicea</i> , <i>Danthonia decumbens</i> , <i>Anthoxanthum odoratum</i> , <i>Succisa pratensis</i> , <i>Polygala serpyllifolia</i> and <i>Molinia caerulea</i> . Small patches of poor-quality <i>Ulex gallii</i> HH1 with <i>Erica cinerea</i> and <i>Rubus fruticosus</i> agg. in parts. <b>EC09 R2</b>
HH3	4010 (WH3)	<i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Molinia caerulea</i> , <i>Succisa pratensis</i> , <i>Carex echinata</i> , <i>Potentilla erecta</i> , <i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Sphagnum capillifolium</i> ssp. <i>rubellum</i> , <i>Pleurozium schreberi</i> , <i>Eriophorum angustifolium</i> , <i>Hylocomium splendens</i> and <i>Narthecium ossifragum</i> . <b>EC09 R3</b>
WL2	n/a	<i>Fraxinus excelsior</i> , <i>Crataegus monogyna</i> , <i>Pteridium aquilinum</i> and <i>Rubus fruticosus</i> agg.
WS1	n/a	<i>Ulex europaeus</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Salix cinerea</i> , <i>Rubus fruticosus</i> , <i>Pteridium aquilinum</i> , <i>Ulex gallii</i> and <i>Erica cinerea</i> .
<b>EC11</b>		
ED3	n/a	<i>Poa annua</i> , <i>Tussilago farfara</i> , <i>Ranunculus repens</i> , <i>Papaver rhoeas</i> , <i>Persicaria maculosa</i> , <i>Atriplex patula</i> , <i>Veronica chamaedrys</i> , <i>Agrostis stolonifera</i> , <i>Sonchus asper</i> , <i>Cerastium fontanum</i> , <i>Fallopia japonica</i> and <i>Fumaria muralis</i> .
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i> .
FS1	n/a	<i>Phragmites australis</i> and <i>Typha latifolia</i> .
FS2	n/a	<i>Apium nodiflorum</i> , <i>Sparganium erectum</i> , <i>Epilobium hirsutum</i> , <i>Juncus articulatus</i> , <i>Hydrocotyle vulgaris</i> and <i>Glyceria fluitans</i> .
GA1	n/a	<i>Rumex obtusifolius</i> , <i>Ranunculus repens</i> , <i>Senecio jacobaea</i> , <i>Persicaria maculosa</i> , <i>Urtica dioica</i> and <i>Plantago major</i> .
GS1	n/a	<i>Centaurea nigra</i> , <i>Festuca rubra</i> , <i>Linum catharticum</i> , <i>Festuca rubra</i> , <i>Dactylis glomerata</i> , <i>Arrhenatherum elatius</i> , <i>Lotus corniculatus</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Leucanthemum vulgare</i> , <i>Euphrasia officinalis</i> , <i>Holcus lanatus</i> , <i>Achillea millefolium</i> and <i>Trifolium pratensis</i> .
GS2	n/a	<i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Arrhenatherum elatius</i> , <i>Anthoxanthum odoratum</i> and <i>Agrostis stolonifera</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS3	n/a	<i>Anthoxanthum odoratum</i> , <i>Agrostis capillaris</i> , <i>Nardus stricta</i> , <i>Carex disticha</i> , <i>Succisa pratensis</i> , <i>Ranunculus repens</i> , <i>Prunella vulgaris</i> , <i>Hypochaeris radicata</i> , <i>Potentilla erecta</i> , <i>Hylocomium splendens</i> , <i>Juncus bulbosus</i> , <i>Galium saxatile</i> , <i>Danthonia decumbens</i> , <i>Plantago lanceolata</i> and <i>Rhytidiadelphus squarrosus</i> .  Well-drained areas around rock outcrops also support <i>Lotus corniculatus</i> , <i>Euphrasia officinalis</i> , <i>Festuca rubra</i> , <i>Succisa pratensis</i> , <i>Carex panicea</i> and <i>Achillea millefolium</i> .
GS4	n/a	Species-poor <i>Juncus effusus</i> dominated vegetation. Typical species are <i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Juncus conglomeratus</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Lythrum salicaria</i> , <i>Lotus pedunculatus</i> , <i>Rubus fruticosus</i> , <i>Arrhenatherum elatius</i> , <i>Dryopteris dilatata</i> , <i>Dactylis glomerata</i> , <i>Anthoxanthum odoratum</i> , <i>Agrostis stolonifera</i> and <i>Carex echinata</i> .
GS4	n/a (GL1diii)  6410 (GL1c)	Tussocky <i>Molinia</i> species-poor type occurring between and grading to heath habitats. Not considered annex habitat but of higher value than species-poor <i>Juncus effusus</i> dominated vegetation. Typical species are <i>Molinia caerulea</i> , <i>Ranunculus repens</i> , <i>Ranunculus acris</i> , <i>Succisa pratensis</i> , <i>Potentilla erecta</i> , <i>Cirsium vulgare</i> , <i>Vicia sepium</i> and <i>Anthoxanthum odoratum</i> . <b>EC11 R12</b>  Species include <i>Molinia caerulea</i> , <i>Angelica sylvestris</i> , <i>Juncus articulatus</i> , <i>Juncus conglomeratus</i> , <i>Lythrum salicaria</i> , <i>Filipendula ulmaria</i> , <i>Carex echinata</i> , <i>Carex flacca</i> , <i>Potentilla erecta</i> , <i>Lotus pedunculatus</i> , <i>Centaurea nigra</i> , <i>Galium palustre</i> and <i>Mentha aquatica</i> . <b>EC11 R1</b>
GM1	n/a	<i>Typha latifolia</i> , <i>Osmunda regalis</i> , <i>Angelica sylvestris</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Hydrocotyle vulgare</i> , <i>Potentilla palustris</i> , <i>Galium palustre</i> , <i>Juncus effusus</i> , <i>Agrostis stolonifera</i> and <i>Equisetum fluviatile</i> .
HD1	n/a	<i>Pteridium aquilinum</i> , with patches of <i>Rubus fruticosus</i> agg. <i>Calystegia sepium</i> , <i>Osmunda regalis</i> , <i>Epilobium hirsutum</i> and <i>Ulex europaeus</i> . Often grading into scrub.
HH1	4030 (DH1/DH3)	Generally occurs in small patches round rock outcrops within a peatland or grassland mosaic. Typical species are <i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Potentilla erecta</i> , <i>Carex binervis</i> , <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Carex demissa</i> , <i>Molinia caerulea</i> , <i>Agrostis viminalis</i> , <i>Nardus stricta</i> , <i>Succisa pratensis</i> , <i>Campylopus flexuosus</i> , <i>Pleurozium schreberi</i> , <i>Scleropodium purum</i> and <i>Hypnum jutlandicum</i> . <b>EC11 R4, EC11 R7 and EC11 R10</b>



Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
HH3	4010 (WH4a, Wh4b, WH7)	The habitat is generally found in mosaic with blanket bog and dry heath. Typical species include <i>Trichophorum germanicum</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Molinia caerulea</i> , <i>Carex binervis</i> , <i>Potentilla erecta</i> , <i>Eriophorum angustifolium</i> , <i>Sphagnum capillifolium</i> ssp. <i>rubellum</i> , <i>Sphagnum fallax</i> , <i>Hypnum jutlandicum</i> , <i>Campylopus atrovirens</i> and <i>Cladonia portentosa</i> . In flushed areas <i>Myrica gale</i> , <i>Molinia caerulea</i> , <i>Schoenus nigricans</i> and <i>Sphagnum tenellum</i> . Grasses and sedges are frequent in heavily poached areas. Some encroachment from <i>Ulex</i> spp. and <i>Salix</i> spp. <b>EC11 R6, EC11 R8, EC11 R9 and EC11 R11</b>
PB3	*7130 (BB2, BB3)	Typical species include <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Schoenus nigricans</i> , <i>Rhynchospora alba</i> , <i>Erica tetralix</i> , <i>Carex panicea</i> , <i>Eriophorum angustifolium</i> , <i>Eriophorum vaginatum</i> , <i>Narthecium ossifragum</i> , <i>Trichophorum germanicum</i> , <i>Sphagnum capillifolium</i> ssp. <i>rubellum</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum tenellum</i> , <i>Sphagnum cuspidatum</i> , <i>Aulacomnium palustre</i> , <i>Drosera rotundifolia</i> , <i>Odontoschisma sphagni</i> and <i>Potentilla erecta</i> . Recovering hollows mostly have 100% <i>Sphagnum</i> cover with <i>Eriophorum angustifolium</i> , some <i>Eriophorum vaginatum</i> , <i>Erica tetralix</i> and <i>Calluna vulgaris</i> . <i>Rhynchospora alba</i> is locally abundant. <b>EC11 R3 and EC11 R5</b>
PB3	7150 (HW3)	<i>Rhynchospora</i> lawns occur in natural hollows and some of the recovering cutover hollows. The habitat is mostly found in the northern parts of the site but small patches present throughout. <i>Rhynchospora alba</i> , <i>Menyanthes trifoliata</i> , <i>Narthecium ossifragum</i> , <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum magellanicum</i> , <i>Sphagnum cuspidatum</i> , <i>Sphagnum denticulatum</i> and <i>Eriophorum angustifolium</i> . <b>EC11 R2</b>
PB4	n/a	A high cover of bare peat with occasional <i>Molinia caerulea</i> , <i>Juncus effusus</i> , <i>Juncus bulbosus</i> and <i>Carex panicea</i> .
PF2	n/a	Small linear drainage feature within acid grassland. <i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Sphagnum denticulatum</i> , <i>Ranunculus flammula</i> , <i>Potamogeton polygonifolius</i> , <i>Juncus bulbosus</i> , <i>Philonotis fontana</i> , <i>Agrostis stolonifera</i> , <i>Lythrum salicaria</i> and <i>Calliergonella cuspidata</i> .
WL1	n/a	<i>Crataegus monogyna</i> , <i>Salix cinerea</i> , <i>Prunus spinosa</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Lonicera periclymenum</i> and <i>Rubus fruticosus</i> agg.
WL2	n/a	Variable habitat type. Species typically found include <i>Pinus contorta</i> , <i>Cupressus</i> sp., <i>Picea sitchensis</i> , <i>Salix cinerea</i> , <i>Fraxinus excelsior</i> , <i>Acer pseudoplatanus</i> , <i>Crataegus monogyna</i> , <i>Rubus fruticosus</i> agg. and <i>Hedera helix</i>
WS1	n/a	Mostly taller scrub. Typical species <i>Ulex europaeus</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> and <i>Salix cinerea</i> . Low scrub is dominated by <i>Rubus fruticosus</i> agg. Usually where it has encroached upon grassland.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC12</b>		
GA1	n/a	<i>Lolium perenne</i> , <i>Senecio jacobaea</i> , <i>Odontites verna</i> , <i>Rumex obtusifolius</i> , <i>Cerastium fontanum</i> , <i>Agrostis stolonifera</i> , <i>Ranunculus repens</i> , <i>Prunella vulgaris</i> and <i>Plantago lanceolata</i>
GS3	n/a	<i>Agrostis capillaris</i> , <i>Cynosurus cristatus</i> , <i>Achillea millefolium</i> , <i>Plantago lanceolata</i> , <i>Anthoxanthum odoratum</i> , <i>Potentilla erecta</i> , <i>Ranunculus repens</i> and <i>Leontodon autumnalis</i>
GS4	n/a	<i>Juncus effusus</i> , <i>Lythrum salicaria</i> , <i>Agrostis stolonifera</i> , <i>Juncus articulatus</i> , <i>Holcus lanatus</i> , <i>Arrhenatherum elatius</i> , <i>Ranunculus repens</i> , <i>Stachys palustris</i> , <i>Angelica sylvestris</i> , <i>Succisa pratensis</i> and <i>Filipendula ulmaria</i> .
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i>
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Daboecia cantabrica</i> , <i>Erica cinerea</i> , <i>Agrostis capillaris</i> , <i>Molinia caerulea</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Epilobium hirsutum</i> and <i>Ulex europaeus</i> . <b>EC12 R1 and EC12 R3</b>
HH3	4010 (WH3)	<i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Narthecium ossifragum</i> , <i>Succisa pratensis</i> , <i>Potentilla erecta</i> , <i>Juncus articulatus</i> , <i>Drosera rotundifolia</i> , <i>Sphagnum capillifolium</i> , <i>S. denticulatum</i> and <i>Polygala serpyllifolia</i> . <b>EC12 R2</b>
WS1	n/a	<i>Rubus fruticosus</i> agg., <i>Ulex europaeus</i> and <i>Salix cinerea</i> .
<b>EC13</b>		
ED3	n/a	<i>Ulex europaeus</i> , <i>Salix cinerea</i> , <i>Rubus fruticosus</i> agg. <i>Molinia caerulea</i> , <i>Rumex acetosa</i> , <i>Succisa pratensis</i> , <i>Erica cinerea</i> , <i>Centaurea nigra</i> and <i>Anthoxanthum odoratum</i> .
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i> .
FS1	n/a	<i>Phragmites australis</i> .
GA1	n/a	<i>Rumex obtusifolius</i> , <i>Ranunculus repens</i> , <i>Cirsium arvense</i> , <i>Senecio jacobaea</i> , <i>Urtica dioica</i> , <i>Plantago major</i> , <i>Holcus lanatus</i> , <i>Dactylis glomerata</i> and <i>Poa annua</i> .
GM1	n/a	<i>Angelica sylvestris</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Hydrocotyle vulgare</i> , <i>Galium palustre</i> , <i>Juncus effusus</i> , <i>Agrostis stolonifera</i> , <i>Potentilla palustris</i> <i>Equisetum fluviatile</i> , <i>Plantago lanceolata</i> , <i>Succisa pratensis</i> and <i>Juncus articulatus</i> .
GS2	n/a	<i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Holcus lanata</i> , <i>Arrhenatherum elatius</i> , <i>Anthoxanthum odoratum</i> , <i>Agrostis stolonifera</i> , <i>Rumex crispus</i> and <i>Rubus fruticosus</i> agg.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS3	n/a	Relatively species-rich dry acid grasslands in mosaic with dry heath. <i>Anthoxanthum odoratum</i> , <i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Carex panicea</i> , <i>Succisa pratensis</i> , <i>Euphrasia officinalis</i> , <i>Pedicularis sylvatica</i> , <i>Hypochaeris radicata</i> , <i>Potentilla erecta</i> , <i>Hylocomium splendens</i> , <i>Centaurea nigra</i> , <i>Galium saxatile</i> , <i>Festuca rubra</i> , <i>Cynosurus cristatus</i> , <i>Prunella vulgaris</i> , <i>Trifolium pratense</i> , <i>Leontodon autumnalis</i> , <i>Achillea millefolium</i> , <i>Nardus stricta</i> and <i>Lotus corniculatus</i> .
GS4	n/a  6410 (GL1c)	Typical soft rush wet agricultural grasslands and patches of higher value (but non-Annex I) <i>Molinia</i> grassland. Merges to mosaic with <i>Molinia</i> meadow, marsh and poor flush. Species include <i>Juncus effusus</i> , <i>Molinia caerulea</i> , <i>Angelica sylvestris</i> , <i>Juncus articulatus</i> , <i>Holcus lanatus</i> , <i>Juncus conglomeratus</i> , <i>Lythrum salicaria</i> , <i>Filipendula ulmaria</i> , <i>Carex echinata</i> , <i>Carex flacca</i> , <i>Potentilla erecta</i> , <i>Galium palustre</i> , <i>Mentha aquatica</i> and <i>Arrhenatherum elatius</i> .  <i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Juncus conglomeratus</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Lythrum salicaria</i> , <i>Anthoxanthum odoratum</i> , <i>Agrostis stolonifera</i> , <i>Carex echinata</i> , <i>Potentilla erecta</i> , <i>Carex flacca</i> , <i>Eriophorum angustifolium</i> , <i>Galium palustre</i> , <i>Angelica sylvestris</i> , <i>Carex panicea</i> and <i>Dactylorhiza maculata</i> .  <b>EC13 R1</b>
HD1	n/a	<i>Pteridium aquilinum</i> , with patches of <i>Rubus fruticosus</i> agg. and <i>Ulex europaeus</i> .
HH1	4030 (DH1)  4030 (DH3)	Two variants occur at the site. DH1 variant is found around rock outcrops in the peatland areas. It also generally occurs in isolated patches. Dominated by a combination of <i>Ulex gallii</i> and <i>Calluna vulgaris</i> . <i>Campylopus introflexus</i> is common on peat that has been burned in the past.  <b>EC13 R5</b>  The DH3 variant occurs in mosaic with relatively species-rich GS3 grassland. Typical species are <i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Potentilla erecta</i> , <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Carex demissa</i> , <i>Carex panicea</i> , <i>Succisa pratensis</i> , <i>Lotus corniculatus</i> , <i>Hylocomium splendens</i> , <i>Scleropodium purum</i> , <i>Hypnum jutlandicum</i> , <i>Festuca rubra</i> and <i>Trifolium pratense</i> .  <b>EC13 R2</b>

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
HH3	4010 (WH4a, WH3)	Vegetation variable across the site but the shallow peat of the rocky areas mainly supports <i>Trichophorum</i> WH4 wet heath while the deeper peats in sections are more typical of WH3. Typical species are <i>Trichophorum germanicum</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Molinia caerulea</i> , <i>Potentilla erecta</i> , <i>Eriophorum angustifolium</i> , <i>Sphagnum capillifolium</i> ssp. <i>rubellum</i> , <i>Hypnum jutlandicum</i> , <i>Campylopus atrovirens</i> , <i>Schoenus nigricans</i> , <i>Aulacomnium palustre</i> , <i>Narthecium ossifragum</i> , <i>Sphagnum compactum</i> , <i>Sphagnum subnitens</i> , <i>Cladonia portentosa</i> and <i>Sphagnum tenellum</i> . <i>Juncus effusus</i> , <i>Juncus articulatus</i> and grasses frequent in degraded areas. <i>Salix repens</i> . <b>EC13 R3 and EC13 R8</b>
PB3	*7130 (BB2, BB3)	Restricted to small patches in deeper hollows between rock outcrops in an area mostly covered by wet heath. <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Schoenus nigricans</i> , <i>Rhynchospora alba</i> , <i>Erica tetralix</i> , <i>Eriophorum angustifolium</i> , <i>Eriophorum vaginatum</i> , <i>Narthecium ossifragum</i> , <i>Sphagnum capillifolium</i> ssp. <i>rubellum</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum magellanicum</i> , <i>Sphagnum cuspidatum</i> , <i>Drosera rotundifolia</i> , <i>Odontoschisma spheni</i> , <i>Potentilla erecta</i> and <i>Eleocharis multicaulis</i> . <b>EC13 R4 and EC13 R7</b>
PB3	7150 (HW3)	<i>Rhynchospora</i> lawns are scattered throughout the wetter parts of the bog, in natural hollow. <i>Rhynchospora alba</i> , <i>Narthecium ossifragum</i> , <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum cuspidatum</i> and <i>Eriophorum angustifolium</i> . <b>EC13 R6</b>
PF2	n/a	<i>Juncus articulatus</i> , <i>Sphagnum denticulatum</i> , <i>Sphagnum palustre</i> , <i>Ranunculus flammula</i> , <i>Agrostis stolonifera</i> , <i>Lythrum salicaria</i> .
WS1	n/a	Taller scrub dominated by <i>Ulex europaeus</i> or <i>Prunus spinosa</i> . The low scrub, dominated by <i>Rubus fruticosus</i> agg., has colonized areas of neglected/abandoned land.
<b>EC14</b>		
ED3	n/a	<i>Rubus fruticosus</i> agg., <i>Ulex europaeus</i> , <i>Chamerion angustifolium</i> , <i>Persicaria maculosa</i> , <i>Lythrum salicaria</i> , <i>Pteridium aquilinum</i> and <i>Calystegia sepium</i> .
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> and <i>Dactylis glomerata</i> .
GS2	n/a	<i>Dactylis glomerata</i> , <i>Centaurea nigra</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Lathyrus pratensis</i> , <i>Odontites verna</i> , <i>Anthoxanthum odoratum</i> , <i>Senecio jacobaea</i> and <i>Rumex crispus</i> .
GS3	n/a	<i>Agrostis capillaris</i> , <i>Nardus stricta</i> , <i>Potentilla erecta</i> , <i>Succisa pratensis</i> , <i>Hylocomium splendens</i> and <i>Juncus articulatus</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS4	n/a	<i>Lythrum salicaria</i> , <i>Juncus articulatus</i> , <i>Filipendula ulmaria</i> , <i>Agrostis stolonifera</i> , <i>Potentilla anserine</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> and <i>Cardamine pratensis</i> . <b>EC14 R5</b>
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Daboecia cantabrica</i> , <i>Molinia caerulea</i> , <i>Potentilla erecta</i> and <i>Hypnum jutlandicum</i> . <b>EC14 R1</b>
HH3	4010 (WH3/WH7)	<i>Trichophorum germanicum</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Ulex gallii</i> , <i>Eriophorum angustifolium</i> , <i>Molinia caerulea</i> , <i>Sphagnum capillifolium</i> , <i>Schoenus nigricans</i> and <i>Carex binervis</i> . <b>EC14 R2 and EC14 R4</b>
PB3	*7130 (BB1b/BB1a)	<i>Schoenus nigricans</i> , <i>Rhynchospora alba</i> , <i>R. fusca</i> , <i>Trichophorum germanicum</i> , <i>Molinia caerulea</i> , <i>Calluna vulgaris</i> , <i>Drosera rotundifolia</i> , <i>Erica tetralix</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum denticulatum</i> and <i>Cladonia portentosa</i> . <b>EC14 R3</b>
PF2	n/a	There are a couple of small flushes within an area of wet grassland. <i>Menyanthes trifoliata</i> , <i>Juncus articulatus</i> , <i>Potentilla anserine</i> , <i>Sphagnum palustre</i> , <i>Ranunculus flammula</i> , <i>Juncus effusus</i> , <i>Succisa pratensis</i> , <i>Polytrichum formosum</i> , <i>Agrostis stolonifera</i> and <i>Persicaria maculosa</i> .
WS1	n/a	<i>Ulex europaeus</i> , <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> , <i>Betula pubescens</i> , <i>Sorbus aucuparia</i> and <i>Ilex aquifolium</i> .
<b>EC18</b>		
ED2	n/a	Cleared vegetation with pockets of wet grassland (GS4) and scrub (WS1) remaining. <i>Arrhenatherum elatius</i> , <i>Juncus articulatus</i> , <i>Iris pseudacorus</i> , <i>Epilobium hirsutum</i> , <i>Filipendula ulmaria</i> , <i>Rubus fruticosus</i> agg. and <i>Ulex europaeus</i> .
GS3	n/a	<i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Prunella vulgaris</i> , <i>Plantago lanceolata</i> , <i>Rumex acetosa</i> , <i>Trifolium pratense</i> , <i>T. Repens</i> , <i>Hypochaeris radicata</i> , <i>Centaurea nigra</i> , <i>Festuca rubra/ovina</i> and <i>Rhytidiadelphus squarrosus</i> . Other species if near dry heath include <i>Succisa pratensis</i> , <i>Galium saxatile</i> , <i>Potentilla erecta</i> and <i>Pleurozium schreberi</i> .
GS3a	n/a (UG2a)	<i>Nardus stricta</i> , <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Cynosurus cristatus</i> , <i>Juncus effusus</i> , <i>J. articulatus</i> , <i>Festuca rubra/ovina</i> , <i>Galium saxatile</i> , <i>Trifolium repens</i> , <i>Rumex acetosa</i> , <i>Pedicularis sylvatica</i> , <i>Plantago lanceolata</i> , <i>Potentilla erecta</i> , <i>Prunella vulgaris</i> , <i>Rhytidiadelphus squarrosus</i> , <i>Hylocomium splendens</i> and <i>Pleurozium schreberi</i> . <b>EC18 R6</b>

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS4	n/a	<i>Arrhenatherum elatius</i> , <i>Molinia caerulea</i> , <i>Juncus effusus</i> , <i>Filipendula ulmaria</i> , <i>Lythrum salicaria</i> , <i>Juncus articulatus</i> , <i>Mentha aquatica</i> , <i>Potentilla erecta</i> , <i>Galium palustre</i> , <i>Rubus fruticosus</i> agg., <i>Cirsium palustre</i> , <i>Salix cinerea</i> and <i>Betula pubescens</i> .
GS4	6410 (GL1c)	Localised areas of <i>Molinia</i> dominated grasslands on peaty ground and often associated with wet grassland or near heath. <i>Molinia caerulea</i> , <i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Anthoxanthum odoratum</i> , <i>Agrostis capillaris</i> , <i>Danthonia decumbens</i> , <i>Nardus stricta</i> , <i>Juncus articulatus</i> , <i>Succisa pratensis</i> , <i>Luzula multiflora</i> , <i>Potentilla erecta</i> , <i>Carex panacea</i> , <i>Carex echinata</i> , <i>Dactylorhiza</i> sp., <i>Filipendula ulmaria</i> , <i>Mentha aquatic</i> , <i>Pleurozium schreberi</i> , <i>Rhytidiadelphus squarrosus</i> , <i>Sphagnum</i> sp., <i>Hylocomium splendens</i> and <i>Calliergonella cuspidata</i> . <b>EC18 R2 and EC18 R5</b>
GM1	n/a	<i>Glyceria fluitans</i> , <i>Agrostis stolonifera</i> , <i>Equisetum fluviatile</i> , <i>Holcus lanatus</i> , <i>Filipendula ulmaria</i> , <i>Mentha aquatic</i> , <i>Apium nodiflorum</i> , <i>Lythrum salicaria</i> and <i>Iris pseudacorus</i> . No bryophytes.
HD1	n/a	<i>Pteridium aquilinum</i>
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Daboecia cantabrica</i> , <i>Succisa pratensis</i> , <i>Arctostaphylos uva-ursi</i> , <i>Agrostis capillaris</i> , <i>Hylocomium splendens</i> , <i>Solidago virgaurea</i> , <i>Potentilla erecta</i> , <i>Molinia caerulea</i> and <i>Succisa pratensis</i> . <b>EC18 R3 and EC18 R7</b>
HH3	4010 (WH3)	<i>Calluna vulgaris</i> , <i>Molinia caerulea</i> , <i>Erica tetralix</i> , <i>Narthecium ossifragum</i> , <i>Juncus articulatus</i> , <i>Potentilla erecta</i> , <i>Succisa pratensis</i> , <i>Dactylorhiza</i> sp., <i>Carex echinata</i> , <i>C. panacea</i> , <i>Rhytidiadelphus squarrosus</i> , <i>Aulacomnium palustre</i> , <i>Hylocomium splendens</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum capillifolium</i> , <i>Thuidium tamariscinum</i> and <i>Cladonia portentosa</i> . Localised areas at south with <i>Schoenus nigricans</i> , <i>Trichophorum germanicum</i> , <i>Eleocharis multicaulis</i> and <i>Myrica gale</i> <b>EC18 R1 and EC18 R4</b>
WD1/WN5	n/a	<i>Alnus glutinosa</i> , <i>Fraxinus excelsior</i> , <i>Betula pubescens</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Hedera helix</i> , <i>Ranunculus repens</i> , <i>Veronica chamaedrys</i> , <i>Phyllitis scolopendrium</i> , <i>Solidago virgaurea</i> , <i>Plagiomnium undulatum</i> , <i>Lophocolea bidentata</i> , <i>Fagus sylvatica</i> , <i>Quercus robur</i> , <i>Sanicula</i> sp., <i>Brachypodium sylvaticum</i> , <i>Geum urbanum</i> , <i>Circaea lutetiana</i> and <i>Rhododendron ponticum</i> .
WS1	n/a	<i>Prunus spinosa</i> , <i>Ulex europaeus</i> , <i>Crataegus monogyna</i> , <i>Acer pseudoplatanus</i> , <i>Sorbus aucuparia</i> , <i>Rubus fruticosus</i> agg. and <i>Pteridium aquilinum</i> . <i>Salix cinerea</i> in hollows.
WS1/GS3	n/a	<i>Prunus spinosa</i> (F-A), <i>Crataegus monogyna</i> , <i>Rubus fruticosus</i> agg. and <i>Pteridium aquilinum</i> . Grassland species as described above under GS3.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC20</b>		
ED3	n/a	<i>Senecio jacobaea</i> , <i>Tussilago farfara</i> , <i>Bellis perennis</i> , <i>Ulex europaeus</i> , <i>Holcus lanatus</i> , <i>Taraxacum officinale</i> , <i>Centaurea nigra</i> and <i>Rumex obtusifolius</i> .
ER1	n/a	Although the rock outcrops are generally devoid of vegetation some occasional clumps of <i>Calluna vulgaris</i> and <i>Blechnum spicant</i> were recorded growing in narrow fissures.
HD1	n/a	<i>Pteridium aquilinum</i> overwhelmingly dominant. Few other species apart from <i>Rubus fruticosus</i> agg.
HH1	4030 (DH1 & DH3)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> , <i>Erica cinerea</i> , <i>Molinia caerulea</i> , <i>Potentilla erecta</i> , <i>Carex binervis</i> , <i>Carex panicea</i> , <i>Hypnum jutlandicum</i> and <i>Daboecia cantabrica</i> .
HH3	4010 (WH4a)	Confined to shallow peat. Typical/dominant species are <i>Molinia caerulea</i> , <i>Trichophorum germanicum</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Sphagnum capillifolium</i> , <i>Narthecium ossifragum</i> and <i>Racomitrium lanuginosum</i> . <b>EC20 R1, EC20 R4 and EC20 R5</b>
FS1	n/a	A small lake completely infilled with dense <i>Phragmites australis</i> reed swamp. <i>Phragmites</i> cover is generally >75%. Other flora includes <i>Equisetum fluviatile</i> , <i>Cammarum palustre</i> , <i>Galium palustre</i> and <i>Calystegia sepium</i> .
FW1	n/a	Very little associated submerged vegetation occurs apart from some sparse <i>Juncus bulbosus</i> , <i>Callitriche stagnalis</i> and <i>Glyceria fluitans</i> .
GS2	n/a	<i>Holcus lanatus</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Trifolium repens</i> , <i>Cynosurus cristatus</i> and <i>Dactylis glomerata</i> .
GS3	n/a	In place vegetation grades in GS2 dry meadow. Typical species are <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Potentilla erecta</i> , <i>Succisa pratensis</i> , <i>Galium saxatile</i> , <i>Lotus corniculatus</i> , <i>Plantago lanceolata</i> and <i>Rhytidadelphus squarrosus</i> .
GS4	n/a  6410	Species-poor <i>Juncus effusus</i> dominated grasslands. Typical species are <i>Juncus effusus</i> , <i>Juncus acutiflorus</i> , <i>Holcus lanatus</i> , <i>Anthoxanthum odoratum</i> , <i>Cirsium palustre</i> , <i>Agrostis stolonifera</i> , <i>Ranunculus repens</i> , <i>Ranunculus flammula</i> , <i>Trifolium repens</i> , <i>Galium palustre</i> and the wetland moss <i>Calliergonella cuspidata</i> .  <i>Molinia caerulea</i> dominated wet grassland type the dominant species generally however in some areas a more open, species-rich wetland vegetation occurs. Associated species include <i>Juncus acutiflorus</i> , <i>Agrostis stolonifera</i> , <i>Lythrum salicaria</i> , <i>Potentilla erecta</i> and <i>Succisa pratensis</i> .  <b>EC20 R3</b>

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
PB3	7130* (Active blanket bog) (BB2 and BB3)	<i>Schoenus nigricans</i> , <i>Molinia caerulea</i> , <i>Erica tetralix</i> , <i>Calluna vulgaris</i> , <i>Narthecium ossifragum</i> , <i>Rhynchospora alba</i> , <i>Eriophorum vaginatum</i> , <i>Eriophorum angustifolium</i> and <i>Trichophorum germanicum</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum capillifolium</i> , <i>Sphagnum cuspidatum</i> and <i>Racomitrium lanuginosum</i> . <b>EC20 R2</b>
PF3	7140 (PFLU5)	<i>Schoenus nigricans</i> , <i>Sphagnum papillosum</i> , <i>Molinia caerulea</i> , <i>Rhynchospora alba</i> , <i>Menyanthes trifoliata</i> , <i>Carex limosa</i> , <i>Carex rostrata</i> , <i>Eleocharis multicaulis</i> , <i>Potamogeton polygonifolius</i> and <i>Aulacomnium palustris</i> .
WD1	n/a	<i>Fraxinus excelsior</i> , <i>Fagus sylvatica</i> , <i>Acer pseudoplatanus</i> and <i>Ilex aquifolium</i> are the main tree species.
WS1	n/a	<i>Ulex europaeus</i> is the dominant species with <i>Rubus fruticosus</i> , <i>Pteridium aquilinum</i> and <i>Molinia caerulea</i> along the margins of scrub. There is also an area of scrub vegetation dominated by <i>Salix cinerea</i> .
<b>EC21</b>		
ER1	n/a	Although the rock outcrops are generally devoid of some occasional plants of <i>Sedum anglicum</i> , <i>Calluna vulgaris</i> and <i>Aira praecox</i> were recorded growing in small rock cracks.
ED3	n/a	Mineral soil dominates these areas with a sparse, weedy, recolonizing flora which includes <i>Senecio jacobaea</i> , <i>Holcus lanatus</i> , <i>Ulex europaeus</i> , <i>Cirsium arvense</i> , <i>Plantago major</i> , <i>Centaurea nigra</i> , <i>Rumex crispus</i> and <i>Leucanthemum vulgare</i> .
HD1	n/a	<i>Pteridium aquilinum</i> is overwhelmingly dominant and there are few associated species apart from <i>Rubus fruticosus</i> and <i>Arrhenatherum elatius</i> .
HH1	4030 (DH1 & DH3)	<i>Ulex gallii</i> , <i>Calluna vulgaris</i> and <i>Erica cinerea</i> , <i>Potentilla erecta</i> , <i>Carex binervis</i> , <i>Agrostis canina</i> , <i>Hypnum jutlandicum</i> , <i>Scleropodium purum</i> , <i>Nardus stricta</i> and <i>Daboecia cantabrica</i> . <b>EC21 R1</b>
HH3	4010 (WH4a)	Areas of wet heath occur on shallow peat. Typical species are <i>Trichophorum germanicum</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Molinia caerulea</i> , <i>Sphagnum capillifolium</i> , <i>Narthecium ossifragum</i> , <i>Eriophorum angustifolium</i> , <i>Potentilla erecta</i> , <i>Succisa pratensis</i> and <i>Racomitrium lanuginosum</i> . <b>EC21 R2 and EC21 R3</b>
GA1	n/a	The dominant plant species tend to be the agricultural grasses <i>Lolium perenne</i> and <i>Holcus lanatus</i> , <i>Senecio jacobaea</i> , <i>Cirsium arvense</i> , <i>Cynosurus cristatus</i> , <i>Trifolium repens</i> and <i>Taraxacum officinale</i> .
GS2	n/a	<i>Arrhenatherum elatius</i> tends to dominate the grassy vegetation with <i>Centaurea nigra</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Trifolium repens</i> and <i>Anthoxanthum odoratum</i> .



Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS3	n/a	Generally species-poor. <i>Agrostis capillaris</i> , <i>Potentilla erecta</i> , <i>Anthoxanthum odoratum</i> , <i>Holcus lanatus</i> , <i>Cynosurus cristatus</i> and <i>Rhytidiadelphus squarrosus</i> .
GS4	n/a	<i>Juncus effusus</i> species-poor grassland. Typical species are <i>Juncus effusus</i> , <i>Juncus acutiflorus</i> , <i>Holcus lanatus</i> , <i>Cirsium palustre</i> , <i>Ranunculus repens</i> , <i>Ranunculus flammula</i> , <i>Galium palustre</i> , <i>Lythrum salicaria</i> , <i>Potentilla erecta</i> and the wetland moss <i>Calliergonella cuspidata</i> .
WS1	n/a	<i>Ulex europaeus</i> is the dominant species with <i>Rubus fruticosus</i> , <i>Pteridium aquilinum</i> and <i>Molinia caerulea</i> along the margins.
<b>EC23</b>		
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg.
GM1	n/a	<i>Juncus effusus</i> , <i>Juncus acutiflorus</i> , <i>Agrostis stolonifera</i> , <i>Epilobium hirsutum</i> , <i>Typha latifolia</i> , <i>Epilobium palustre</i> , <i>Hydrocotyle vulgaris</i> , <i>Lotus pedunculatus</i> , <i>Filipendula ulmaria</i> and <i>Calliergonella</i> sp.
GS3	n/a	<i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Cynosurus cristatus</i> , <i>Festuca rubra/ovina</i> , <i>Hypochaeris radicata</i> , <i>Trifolium repens</i> , <i>Lotus corniculatus</i> , <i>Plantago lanceolata</i> , <i>Trifolium repens</i> , <i>Rumex acetosa</i> and <i>Rhytidiadelphus squarrosus</i> . <b>EC23 R1</b>
GS4	n/a	<i>Juncus effusus</i> , <i>Ranunculus repens</i> , <i>Rumex acetosa</i> , <i>Calystegia soldanella</i> , <i>Cirsium palustre</i> , <i>Filipendula ulmaria</i> , <i>Stachys palustris</i> , <i>Holcus lanatus</i> , <i>Agrostis stolonifera</i> , <i>Lotus pedunculatus</i> , <i>Lythrum salicaria</i> , <i>Calliergonella cuspidata</i> and <i>Rhytidiadelphus squarrosus</i> .
<b>EC25</b>		
ED3	n/a	<i>Hypochaeris radicata</i> , <i>Senecio jacobaea</i> , <i>Agrostis capillaris</i> , <i>Achillea millefolium</i> , <i>Trifolium pratensis</i> , <i>Centaureum erythraea</i> , <i>Anthoxanthum odoratum</i> , <i>Trifolium dubium</i> , <i>Anagallis arvensis</i> , <i>Plantago lanceolata</i> , <i>Cirsium arvensis</i> , <i>Ulex europaeus</i> , <i>Juncus effusus</i> , <i>Potentilla anserine</i> , <i>Rubus fruticosus</i> , <i>Leontodon autumnalis</i> , <i>Linum catharticum</i> and <i>Hypericum tetrapterum</i> .
FL1	3160	Two small ponds, both surrounded by reed swamp with adjacent blanket bog and scrub. Typical species are <i>Phragmites australis</i> , <i>Potamogeton polygonifolius</i> , <i>Hypericum elodes</i> and <i>Menyanthes trifoliata</i> .
FL4	n/a	<i>Carex lasiocarpa</i> , <i>Hypericum elodes</i> , <i>Menyanthes trifoliata</i> , <i>Schoenoplectus lacustris</i> and <i>Potamogeton polygonifolius</i>
FS1	n/a	<i>Phragmites australis</i> , <i>Mentha aquatica</i> and <i>Myrica gale</i> .
GA1	n/a	<i>Lolium perenne</i> , <i>Rumex crispus</i> , <i>Plantago major</i> , <i>Senecio jacobaea</i> , <i>Trifolium repens</i> , <i>Bellis perennis</i> , <i>Cerastium fontanum</i> , <i>Cirsium arvensis</i> and <i>Achillea millefolium</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS2	6150 (GL3e)	<i>Arrhenatherum elatius</i> , <i>Dactylis glomerata</i> , <i>Epilobium hirsutum</i> , <i>Galium saxatile</i> , <i>Cirsium arvensis</i> , <i>Urtica dioica</i> , <i>Rumex obtusifolius</i> , <i>Euphrasia</i> spp., <i>Plantago lanceolata</i> , <i>Cynosurus cristatus</i> , <i>Prunus vulgaris</i> , <i>Lathyrus pratensis</i> , <i>Anthoxanthum odoratum</i> , <i>Helictotrichon pubescens</i> , <i>Ranunculus acris</i> , <i>Heracleum sphondylium</i> and <i>Pimpinella major</i> . <b>EC25 R6</b>
GS3	n/a	<i>Leontodon autumnalis</i> , <i>Succisa pratensis</i> , <i>Achillea millefolium</i> , <i>Festuca rubra</i> , <i>Prunus vulgaris</i> , <i>Plantago lanceolata</i> , <i>Senecio jacobaea</i> , <i>Rumex acetosella</i> , <i>Potentilla erecta</i> , <i>Euphrasia</i> spp., <i>Trifolium pratensis</i> , <i>Agrostis capillaris</i> and <i>Anthoxanthum odoratum</i> , <i>Holcus lanatus</i> , <i>Agrostis canina</i> , <i>Pteridium aquilinum</i> , <i>Centaurea nigra</i> , <i>Lolium perenne</i> , <i>Leontodon hispidus</i> , <i>Plantago major</i> , <i>Hylocomium splendens</i> , <i>Dactylis glomerata</i> and <i>Solidago virgaurea</i> . <b>EC25 R7</b>
GS3/ED2/ED3	n/a	Disturbed area with mosaic of GS3/ED2 (i.e. bare soil) and ED3., <i>Agrostis canina</i> , <i>Arrhenatherum elatius</i> , <i>Dactylis glomerata</i> , <i>Festuca</i> spp., <i>Leontodon autumnalis</i> , <i>Odontites vernus</i> , <i>Potentilla reptans</i> , <i>Prunella vulgaris</i> , <i>Pteridium aquilinum</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Succisa pratensis</i> , <i>Tussilago farfara</i> and <i>Ulex europaeus</i> .
GS4	n/a	<i>Juncus effusus</i> , <i>Rumex acetosa</i> , <i>Stachys palustris</i> , <i>Rumex acetosella</i> , <i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Rumex crispus</i> , <i>Juncus acutiflorus</i> , <i>Lythrum salicaria</i> , <i>Rubus fruticosus</i> agg., <i>Arrhenatherum elatius</i> , <i>Rumex obtusifolius</i> and <i>Galium palustre</i> . <b>EC25 R8</b>
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Ulex europaeus</i> , <i>Ulex gallii</i> and <i>Erica cinerea</i> .
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Daboecia cantabrica</i> , <i>Calluna vulgaris</i> , <i>Molinia caerulea</i> and <i>Cladonia portentosa</i> .
HH3	4010 (WH1b)	<i>Molinia caerulea</i> , <i>Narthecium ossifragum</i> , <i>Succisa pratensis</i> , <i>Calluna vulgaris</i> , <i>Erica tetralix</i> , <i>Drosera rotundifolia</i> , <i>Pedicularis sylvatica</i> , <i>Potentilla erecta</i> , <i>Dactylis maculate</i> , <i>Sphagnum capillifolium</i> , <i>Carex panicea</i> , <i>Agrostis stolonifera</i> and <i>Schoenus nigricans</i> . <b>EC25 R1 and EC25 R3</b>
PB3	*7130 (BB1b)	<i>Molinia caerulea</i> , <i>Erica tetralix</i> , <i>Narthecium ossifragum</i> , <i>Rhynchospora alba</i> , <i>Sphagnum papillosum</i> , <i>Sphagnum capillifolium</i> , <i>Eriophorum angustifolium</i> , <i>Potentilla erecta</i> , <i>Drosera rotundifolia</i> and <i>Sphagnum cuspidatum</i> . <b>EC25 R2</b>
PB3	7150 (HW3)	<i>Rhynchospora</i> hollows are abundant within the extensive area of blanket bog. <i>Rhynchospora alba</i> , <i>Sphagnum papillosum</i> , <i>Narthecium ossifragum</i> , <i>Drosera rotundifolia</i> , <i>Eriophorum angustifolium</i> , <i>Sphagnum cuspidatum</i> and <i>Erica tetralix</i> . <b>EC25 R4</b>

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
PF2	n/a	There are several small flushes throughout the site in association with blanket bog, wet heath, transition mire and wet grassland. <i>Hypericum elodes</i> , <i>Menyanthes trifoliata</i> , <i>Ranunculus flammula</i> , <i>Potamogeton polygonifolius</i> , <i>Sphagnum denticulatum</i> , <i>Molinia caerulea</i> , <i>Carex echinata</i> , <i>Schoenus nigricans</i> and <i>Anagallis arvensis</i> .
PF3	7140 (PFLU5)	<i>Phragmites australis</i> , <i>Schoenus nigricans</i> , <i>Menyanthes trifoliata</i> , <i>Narthecium ossifragum</i> , <i>Equisetum hyemale</i> , <i>Myrica gale</i> , <i>Hypericum elodes</i> , <i>Potamogeton polygonifolius</i> , <i>Eriophorum angustifolium</i> , <i>Potentilla palustris</i> , <i>Erica tetralix</i> , <i>Sphagnum palustre</i> , <i>Carex rostrata</i> , <i>Campylium stellatum</i> and <i>Vaccinium oxycoccus</i> . <b>EC25 R5</b>
WS1	n/a	<i>Ulex europaeus</i> , <i>Crataegus monogyna</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Pteridium aquilinum</i> , <i>Urtica dioica</i> and <i>Salix cinerea</i> .
<b>EC29</b>		
FS1	n/a	Species-poor, tall reed swamp vegetation dominates wide, wet drains which occur intermittently on both sides of the railway embankment. <i>Phragmites australis</i> , <i>Epilobium hirsutum</i> , <i>Urtica dioica</i> , <i>Menyanthes trifoliata</i> and <i>Equisetum fluviatile</i> .
GA1	n/a	<i>Holcus lanatus</i> , <i>Ranunculus repens</i> , <i>Lolium perenne</i> , <i>Cynosurus cristatus</i> , <i>Bellis perennis</i> and <i>Plantago lanceolata</i> .
GS1	6210 (GL1b)	Relatively species-rich calcareous grassland is frequent on shallow soil which overlies the limestone of the old railway embankment. Parts correspond to annex habitat, while other areas do not meet criteria. Typical species are <i>Centaurea nigra</i> , <i>Trifolium pratense</i> , <i>Festuca rubra</i> , <i>Plantago lanceolata</i> , <i>Lotus corniculatus</i> , <i>Briza media</i> , <i>Anthoxanthum odoratum</i> , <i>Leontodon hispidus</i> , <i>Rhinanthus minor</i> , <i>Linum catharticum</i> , <i>Vicia cracca</i> and <i>Prunella vulgaris</i> . <b>EC29 R1</b>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Agrostis capillaris</i> , <i>Trifolium repens</i> , <i>Ranunculus repens</i> , <i>Trifolium pratense</i> and <i>Anthoxanthum odoratum</i> .
GS4/GM1	n/a	<i>Juncus effusus</i> , <i>Iris pseudacorus</i> , <i>Holcus lanatus</i> , <i>Lythrum salicaria</i> , <i>Cirsium palustre</i> , <i>Ranunculus acris</i> , <i>Juncus articulatus</i> , <i>Agrostis stolonifera</i> and <i>Filipendula ulmaria</i> .
PF1	7230	<i>Succisa pratensis</i> , <i>Juncus acutiflorus</i> , <i>Potentilla erecta</i> , <i>Prunus vulgaris</i> , <i>Anagallis tenella</i> , <i>Equisetum</i> sp., <i>Anthoxanthum odoratum</i> , <i>Cirsium palustre</i> , <i>Carex panicea</i> , <i>Call cusp</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Schoenus nigricans</i> and <i>Hydrocotyle vulgaris</i> .
WS1	n/a	<i>Prunus spinosa</i> , <i>Rubus fruticosus</i> , <i>Salix cinerea</i> , <i>Urtica dioica</i> and <i>Arrhenatherum elatius</i> . The scrub is generally impenetrable and dense.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC30</b>		
WD1	n/a	<i>Fraxinus excelsior</i> , <i>Acer pseudoplatanus</i> , <i>Alnus cordata</i> and <i>Populus</i> sp, <i>Hedera helix</i> , <i>Urtica dioica</i> , <i>Rubus fruticosus</i> , <i>Galium aparine</i> , <i>Dactylis glomerata</i> , <i>Ranunculus repens</i> , <i>Cirsium vulgare</i> and <i>Geranium robertianum</i> .
WN2	n/a	<i>Fraxinus excelsior</i> , <i>Salix cinerea</i> , <i>Salix caprea</i> , <i>Rubus fruticosus</i> , <i>Fagus sylvatica</i> , <i>Hedera helix</i> , <i>Urtica dioica</i> , <i>Acer pseudoplatanus</i> , <i>Calystegia sepium</i> , <i>Fallopia japonica</i> , <i>Crataegus monogyna</i> , <i>Betula pubescens</i> , <i>Geranium robertianum</i> , <i>Arum maculatum</i> , <i>Polystichum setiferum</i> , <i>Thuidium tamariscinum</i> and <i>Eurhynchium</i> spp.
<b>EC31</b>		
GS1	6210* (GL3a)	Species-rich calcareous grassland occurs in mosaic with lowland hay meadow. Orchids are abundant in places. <i>Briza media</i> , <i>Dactylorhiza fuchsii</i> , <i>Lotus corniculatus</i> , <i>Anthyllis vulneraria</i> , <i>Blackstonia perfoliata</i> , <i>Carex flacca</i> , <i>Linum catharticum</i> , <i>Listera ovata</i> and <i>Pilosella officinarum</i> . <b>EC31 R2</b>
GS2	n/a  6510 (GL3e)	Species-poor meadow and verge. Typical species include <i>Dactylis glomerata</i> , <i>Helictotrichon pubescens</i> , <i>Centaurea nigra</i> , <i>Trifolium repens</i> , <i>Rumex crispus</i> , <i>Cirsium arvensis</i> and <i>Plantago lanceolata</i> .  Occurs on the site in mosaic with calcareous grassland (6210). <i>Plantago lanceolata</i> , <i>Trifolium pratense</i> , <i>Heracleum sphondylium</i> , <i>Helictotrichon pubescens</i> , <i>Lathyrus pratensis</i> , <i>Holcus lanatus</i> , <i>Centaurea nigra</i> , <i>Anthoxanthum odoratum</i> , <i>Dactylis glomerata</i> and <i>Agrostis capillaris</i> , <i>Dactylorhiza fuchsia</i> and <i>Dactylorhiza incarnata</i> . <b>EC31 R1</b>
GS4	n/a	<i>Filipendula ulmaria</i> , <i>Urtica dioica</i> , <i>Juncus effusus</i> , <i>Iris pseudacorus</i> , <i>Convolvulus arvensis</i> , <i>Galium aparine</i> , <i>Festuca arundinacea</i> , <i>Juncus conglomeratus</i> , <i>Epilobium hirsutum</i> and <i>Valeriana officinalis</i> .
WD1	n/a	<i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Rubus fruticosus</i> , <i>Ilex aquifolium</i> , <i>Primula vulgaris</i> , <i>Acer pseudoplatanus</i> and <i>Conopodium majus</i> .
WN2	n/a	<i>Fraxinus excelsior</i> , <i>Acer pseudoplatanus</i> , <i>Hedera helix</i> , <i>Crataegus monogyna</i> , <i>Circaea lutetiana</i> , <i>Geranium robertianum</i> , <i>Rubus fruticosus</i> agg. <i>Urtica dioica</i> , <i>Geum urbanum</i> , <i>Rumex sanguineus</i> and <i>Ranunculus ficaria</i> .
WN6	*91EO	Typical species include <i>Alnus glutinosa</i> , <i>Salix cinerea</i> , <i>Crataegus monogyna</i> , <i>Listera ovata</i> , <i>Holcus mollis</i> , <i>Carex remota</i> and <i>Agrostis stolonifera</i> . <b>EC31 R3</b>
WS1	n/a	Scrub composition varies but dominant species include <i>Crataegus monogyna</i> , <i>Prunus spinosa</i> and <i>Rubus fruticosus</i> agg.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC33</b>		
ED3	n/a	<i>Sonchus oleraceus</i> , <i>Arrhenatherum elatius</i> , <i>Rubus fruticosus</i> , <i>Senecio jacobaea</i> , <i>Potentilla sterilis</i> , <i>Epilobium montana</i> , <i>Chamerion angustifolium</i> , <i>Hedera helix</i> , <i>Lonicera periclymenum</i> and <i>Cirsium arvensis</i> .
ER2	*8240 (LPE_3a)	<i>Hedera helix</i> , <i>Rosa spinosissima</i> , <i>Asplenium ruta-muraria</i> , <i>Ctenidium molluscum</i> , <i>Geranium robertianum</i> , <i>Sesleria caerulea</i> , <i>Ilex aquilinum</i> , <i>Prunus spinosa</i> , <i>Sorbus hibernica</i> , <i>Corylus avellana</i> , <i>Teucrium scorodonia</i> , <i>A. Trichomanes</i> , <i>Thymus polytrichus</i> , <i>Fragaria vesca</i> and <i>Viburnum opulus</i> . <b>EC33 R3</b>
FS1	n/a	<i>Phragmites australis</i> , <i>Lythrum salicaria</i> and <i>Filipendula ulmaria</i> .
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Arrhenatherum elatius</i> , <i>Sonchus oleraceus</i> , <i>Holcus lanatus</i> and <i>Galium aparine</i> .
GS1	6210 (GL3a)	<i>Briza media</i> , <i>Galium verum</i> , <i>Prunella vulgaris</i> , <i>Succisa pratensis</i> , <i>Dactylis fuchsia</i> , <i>Thymus polytrichus</i> , <i>Trifolium pratensis</i> , <i>Sesleria albicans</i> , <i>Anthoxanthum odoratum</i> and <i>Lotus corniculatus</i> . Also some wet elements in grassland adjacent to wetland e.g. <i>Schoenus nigricans</i> , <i>Molinia caerulea</i> and <i>Galium boreale</i> . <b>EC33 R2 and EC33 R4</b>
GS2	n/a	Species-poor variant. Typical species include <i>Lolium perenne</i> , <i>Odontites verna</i> , <i>Ranunculus repens</i> , <i>Cirsium arvensis</i> , <i>Centaurea nigra</i> , <i>Holcus lanatus</i> and <i>Agrostis capillaris</i> .
	6510 (GL3e)	Typical species include <i>Plantago lanceolata</i> , <i>Odontites verna</i> , <i>Ranunculus acris</i> , <i>Prunus vulgaris</i> , <i>Senecio jacobaea</i> , <i>Lolium perenne</i> , <i>Cerastium fontanum</i> , <i>R. repens</i> , <i>Centaurea nigra</i> , <i>Leucanthemum vulgare</i> and <i>Trifolium repens</i> . <b>EC33 R1</b>
WN2	n/a	<i>Corylus avellana</i> , <i>Hedera helix</i> , <i>Fraxinus excelsior</i> , <i>Veronica chamaedrys</i> , <i>Rubus fruticosus</i> , <i>Primula vulgaris</i> , <i>Rhytidadelphus triquetrus</i> , <i>Geranium robertianum</i> and <i>Brachythecium rutabulum</i> .
WN6	*91E0	Occurs in mosaic with scrub and reed swamp (FS1). <i>Salix cinerea</i> , <i>Fraxinus excelsior</i> , <i>Iris pseudacorus</i> , <i>Phragmites australis</i> and <i>Filipendula ulmaria</i> .
WS1	n/a	<i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Rubus fruticosus</i> , <i>Arrhenatherum elatius</i> , <i>Euonymus europaeus</i> , <i>Lonicera periclymenum</i> and <i>Calystegia sepium</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC36</b>		
ER2	*8240 (LPE_1b/1c)	<i>Acer pseudoplatanus</i> , <i>Asplenium ruta-muraria</i> , <i>Buddleja davidii</i> , <i>Calluna vulgaris</i> , <i>Carex flacca</i> , <i>Carlina vulgaris</i> , <i>Centranthus ruber</i> , <i>Ceterach officinalis</i> , <i>Corylus avellana</i> , <i>Cotoneaster</i> sp., <i>Festuca ovina</i> , <i>Fissidens dubius</i> , <i>Fraxinus excelsior</i> , <i>Geranium robertianum</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Linum catharticum</i> , <i>Lotus corniculatus</i> , <i>Mycelis muralis</i> , <i>Potentilla erecta</i> , <i>Prunus spinosa</i> , <i>Rosa spinosissima</i> , <i>Rubus fruticosus</i> agg., <i>Sesleria albicans</i> , <i>Sorbus Hibernica</i> , <i>Succisa pratensis</i> , <i>Teucrium scorodonia</i> , <i>Thalictrum minus</i> , <i>Thymus praecox</i> , <i>Tortella tortuosa</i> and <i>Trichostomum crispulum</i> . <b>EC36 R6</b>
GA1	n/a	<i>Holcus lanatus</i> , <i>Trifolium repens</i> , <i>Ranunculus repens</i> , <i>Cirsium arvense</i> , <i>Ranunculus acris</i> , <i>Rumex crispus</i> , <i>Agrostis stolonifera</i> , <i>Heracleum sphondylium</i> with <i>Plantago major</i> , <i>Ononis repens</i> , <i>Rumex obtusifolius</i> . Occasional GS4 species e.g. <i>Juncus effusus</i> , <i>Potentilla anserina</i> and <i>Carex hirta</i> .
GS1	n/a  6210/6210* (GL3a)	Non-annex variant. Typical species include <i>Ranunculus acris</i> , <i>Holcus lanatus</i> , <i>Dactylis glomerata</i> , <i>Centaurea nigra</i> , <i>Plantago lanceolata</i> , <i>Cirsium arvense</i> , <i>Senecio jacobaea</i> , <i>Cynosurus cristatus</i> , <i>Prunella vulgaris</i> , <i>Festuca rubra</i> , <i>Trifolium repens</i> , <i>Lolium perenne</i> , <i>Agrostis stolonifera</i> , <i>Trifolium pratense</i> , <i>Taraxacum officinale</i> agg. Some <i>Daucus carota</i> , <i>Lotus corniculatus</i> , <i>Galium verum</i> , <i>Linum catharticum</i> . Some GS4 species locally e.g. <i>Filipendula ulmaria</i> and <i>Carex hirta</i> .  Annex Habitat sometimes with orchids. Typical species include <i>Briza media</i> , <i>Sesleria caerulea</i> , <i>Anthyllis vulneraria</i> , <i>Succisa pratensis</i> , <i>Lotus corniculatus</i> , <i>Leontodon hispidus</i> , <i>Carex flacca</i> , <i>Centaurea nigra</i> , <i>Linum catharticum</i> , <i>Ctenidium molluscum</i> , <i>Pilosella officinarum</i> , <i>Primula veris</i> , <i>Galium verum</i> , <i>Carlina vulgaris</i> , <i>Rhinanthus minor</i> , <i>Solidago virgaurea</i> , with orchids <i>Gymnadenia conopsea</i> and <i>Dactylorhiza fuchsii</i> and <i>Neottia ovata</i> , <i>Primula veris</i> , <i>Leucanthemum vulgare</i> , <i>Daucus carota</i> , <i>Crepis capillaris</i> , <i>Avenula pubescens</i> and <i>Homalothecium lutescens</i> . Some encroaching scrub of <i>Pteridium aquilinum</i> , <i>Prunus spinosa</i> and <i>Crataegus monogyna</i> . <b>EC36 R1 and EC36 R3</b>
GS1/ GS2b	n/a	Species-poor GS2b <i>Arrhenatherum elatius</i> , <i>Ranunculus repens</i> , <i>Cirsium arvense</i> , <i>Rumex obtusifolius</i> , <i>Rumex crispus</i> , <i>Potentilla anserina</i> , <i>Heracleum sphondylium</i> , <i>Cerastium fontanum</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Alopecurus pratensis</i> , <i>Urtica dioica</i> , <i>Potentilla anserina</i> , with occasional GS1 species e.g. <i>Prunella vulgaris</i> , <i>Centaurea nigra</i> , <i>Trifolium pratensis</i> , <i>Achillea millefolium</i> , <i>Leontodon hispidus</i> , <i>Rosa spinosissima</i> , <i>Centaurea nigra</i> , <i>Anthyllis vulneraria</i> , <i>Linum catharticum</i> , <i>Prunella vulgaris</i> , <i>Lotus corniculatus</i> and <i>Euphrasia officinalis</i> agg.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS1/ HD1	n/a	Typical GS1 species with <i>Pteridium aquilinum</i> . Some areas highly grazed, others with low grazing.
GS2	n/a	Low species-diversity variant. Typical species: <i>Arrhenatherum elatius</i> , <i>Ranunculus repens</i> , <i>Cirsium arvense</i> , <i>Rumex obtusifolius</i> , <i>Rumex crispus</i> , <i>Potentilla anserina</i> , <i>Heracleum sphondylium</i> , <i>Cerastium fontanum</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Alopecurus pratensis</i> and <i>Urtica dioica</i> . Occasional GS1 species e.g. <i>Prunella vulgaris</i> , <i>Centaurea nigra</i> , <i>Trifolium pratensis</i> , <i>Achillea millefolium</i> , in shorter vegetation.
GS2	n/a	More species-rich variant. Typical species are <i>Heracleum sphondylium</i> (LD), <i>Stachys sylvatica</i> , <i>Phleum pratense</i> , <i>Calystegia sepium</i> , <i>Lolium perenne</i> , <i>Rumex obtusifolius</i> , <i>Lathyrus pratensis</i> , <i>Tussilago farfara</i> , <i>Arrhenatherum elatius</i> (LA), <i>Cirsium arvense</i> (LA), <i>Dactylis glomerata</i> and <i>Lapsana communis</i> . Occasional GS1/ 6210 indicators e.g. <i>Knautia arvensis</i> , <i>Galium verum</i> , <i>Centaurea nigra</i> (LA), <i>Rhinanthus minor</i> , <i>Rosa spinosissima</i> , <i>Leucanthemum vulgare</i> , <i>Daucus carota</i> , <i>Pimpinella saxifraga</i> , <i>Dactylorhiza fuchsii</i> , <i>Succisa pratense</i> (LA), <i>Carex flacca</i> , <i>Thymus polytrichus</i> , <i>Lotus corniculatus</i> . Scattered scrub frequent e.g. <i>Rubus fruticosus</i> agg. <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Corylus avellana</i> , <i>Sorbus</i> sp.
GS2	6510	Typical species are (but not limited to) <i>Cynosurus cristatus</i> , <i>Festuca rubra/ovina</i> , <i>Dactylis glomerata</i> , <i>Phleum pratense</i> , <i>Agrostis capillaris</i> , <i>Briza media</i> , <i>Holcus lanatus</i> , <i>Anthoxanthum odoratum</i> , <i>Rhinanthus minor</i> , <i>Centaurea nigra</i> , <i>Plantago lanceolata</i> , <i>Lotus corniculatus</i> , <i>Achillea millefolium</i> , <i>Daucus carota</i> , <i>Agrimonia eupatoria</i> , <i>Thymus praecox</i> , <i>Hypochaeris radicata</i> , <i>Prunella vulgaris</i> , <i>Ranunculus acris</i> , <i>Cerastium fontanum</i> , <i>Prunus spinosus</i> and <i>Filipendula ulmaria</i> . <b>EC36 R7</b>
GS2/ WS1	n/a	GS2 species: <i>Centaurea nigra</i> , <i>Leucanthemum vulgare</i> , <i>Arrhenatherum elatius</i> , <i>Centranthus ruber</i> , <i>Plantago lanceolata</i> , <i>Polygala vulgaris</i> , <i>Calliargonella cuspidata</i> , <i>Agrostis stolonifera</i> , <i>Potentilla erecta</i> , <i>Hypochaeris radicata</i> , <i>Brachypodium sylvaticum</i> , with frequent WS1 species: <i>Rubus fruticosus</i> agg., <i>Fraxinus excelsior</i> , <i>Acer pseudoplatanus</i> , <i>Chamerion angustifolium</i> , <i>Salix caprea</i> and <i>Corylus avellana</i> . 6210 species in shorter vegetation e.g. <i>Briza media</i> , <i>Lotus corniculatus</i> , <i>Carex flacca</i> , <i>Galium verum</i> , <i>Carlina vulgaris</i> and <i>Tortella tortuosa</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS4	n/a	High species-diversity wet grassland. Typical species include <i>Juncus conglomeratus</i> , <i>Agrostis stolonifera</i> , <i>Potentilla erecta</i> , <i>Dactylorhiza fuchsii</i> , <i>Prunella vulgaris</i> , <i>Juncus bufonius</i> , <i>Filipendula ulmaria</i> (LD), <i>Lotus corniculatus</i> , <i>Lathyrus linifolius</i> , <i>Juncus effusus</i> , <i>Holcus lanatus</i> , <i>Carex flacca</i> , <i>Succisa pratensis</i> (LA), <i>Anthoxanthum odoratum</i> , <i>Molinia caerulea</i> , <i>Hypochaeris radicata</i> , <i>Vicia cracca</i> , <i>Centaurea nigra</i> , <i>Carex panicea</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Vicia sepium</i> , <i>Cirsium arvense</i> (LD), <i>Lathyrus pratensis</i> , <i>Persicaria amphibia</i> and <i>Alopecurus pratensis</i> . Occasional 6210 species e.g. <i>Lotus corniculatus</i> , <i>Rhinanthus minor</i> , <i>Euphrasia officinalis</i> agg. and <i>Gymnadenia conopsea</i> .
GS4/ FS2	n/a	Central wet channel supports tall-herb swamp FS2 (with local areas of reed and large sedge swamp FS1): <i>Juncus effusus</i> , <i>Nasturtium aquaticum</i> (LD), <i>Glyceria fluitans</i> , <i>Juncus articulatus</i> , <i>Typha latifolia</i> , <i>Eleocharis palustris</i> , <i>Potamogeton natans</i> , <i>Iris pseudacorus</i> , <i>Lemna minor</i> and <i>Lemna minuta</i> . Vegetation cover decreases in channel to southeast. Wet grassland with typical GS4 species (above) with additions such as <i>Trifolium pratense</i> , <i>Ranunculus acris</i> , <i>Hypericum pulchrum</i> , <i>Equisetum palustre</i> , <i>Epilobium hirsutum</i> , <i>Cirsium palustre</i> , <i>Mentha aquatica</i> , <i>Carex nigra</i> , <i>Carex disticha</i> , <i>Galium palustre</i> , <i>Epilobium palustre</i> , <i>Dactylorhiza fuchsii</i> , <i>Lythrum salicaria</i> , <i>Daucus carota</i> and <i>Senecio aquatica</i> .
HD1	n/a	<i>Pteridium aquilinum</i> (D), <i>Cirsium arvense</i> and <i>Rumex crispus</i> .
WL1	n/a	<i>Crataegus monogyna</i> , <i>Prunus spinosa</i> , <i>Fraxinus excelsior</i> , <i>Rubus fruticosus</i> agg. <i>Pteridium aquilinum</i> (at edges), <i>Corylus avellana</i> , <i>Calystegia sepium</i> , <i>Urtica dioica</i> , <i>Hedera helix</i> and <i>Ilex aquifolium</i> .
WL2	n/a	<i>Fraxinus excelsior</i> dominant (mature and young trees) with <i>Prunus spinosa</i> , <i>Hedera helix</i> , <i>Asplenium scolopendrium</i> , <i>Rubus fruticosus</i> agg. <i>Thamnobryum alopecurum</i> , <i>Crataegus monogyna</i> , <i>Calystegia sepium</i> , <i>Urtica dioica</i> , <i>Corylus avellana</i> and <i>Populus tremula</i> .
WN2	n/a	<i>Fraxinus excelsior</i> with <i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Acer pseudoplatanus</i> , <i>Populus tremula</i> , <i>Ilex aquifolium</i> , <i>Lonicera periclymenum</i> , <i>Dryopteris dilatata</i> , <i>Carex remota</i> , <i>Eurhynchium striatum</i> and <i>Neckera complanata</i> , <i>Prunus spinosa</i> , <i>Ilex aquilinum</i> , <i>Corylus avellana</i> , <i>Fragaria vesca</i> , <i>Hedera helix</i> , <i>Rubus fruticosus</i> , <i>Viola</i> spp., <i>Geranium robertianum</i> , <i>Rosa canina</i> , <i>Circaea lutetiana</i> and <i>Brachypodium sylvaticum</i> .
WN2	*8240 (LPW_2a)	<i>Corylus avellana</i> , <i>Ilex aquifolium</i> , <i>Euonymus europaeus</i> , <i>Fraxinus excelsior</i> , <i>Crataegus monogyna</i> and <i>Rubus fruticosus</i> agg. Ground flora includes <i>Thamnobryum alopecurum</i> , <i>Eurhynchium striatum</i> , <i>Neckera crispa</i> <i>Dryopteris filix-mas</i> , <i>Asplenium scolopendrium</i> , <i>Melica uniflora</i> , <i>Circaea lutetiana</i> , <i>Potentilla sterilis</i> , <i>Primula vulgaris</i> and <i>Geranium robertianum</i> .  <b>EC36 R2</b>



Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WS1	n/a	<i>Corylus avellana</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Rubus fruticosus</i> agg. <i>Pteridium aquilinum</i> and <i>Hedera helix</i> . Occasional mature <i>Fraxinus excelsior</i> trees.
<b>EC37</b>		
ED3	n/a	<i>Cirsium arvense</i> , <i>Tripleurospermum inodorum</i> , <i>Persicaria maculata</i> , <i>Rumex crispus</i> , <i>Stellaria media</i> , <i>Euphorbia helioscopia</i> , <i>Plantago major</i> , <i>Rumex obtusifolius</i> , <i>Sonchus oleraceus</i> , <i>Sisymbrium officinale</i> , <i>Cirsium vulgare</i> , <i>Centaurea nigra</i> , <i>Potentilla anserina</i> with scattered <i>Rubus fruticosus</i> agg. scrub.
ER2	*8240 (LPE_1b)	Typical species include <i>Teucrium scorodonia</i> , <i>Geranium robertianum</i> , <i>Asplenium ruta-muraria</i> , <i>Asplenium scolopendrium</i> , <i>Tortella tortuosa</i> , <i>Fissidens dubius</i> , <i>Fragaria vesca</i> , <i>Sesleria caerulea</i> , <i>Mycelia muralis</i> , <i>Hedera helix</i> , <i>Ctenidium molluscum</i> , <i>Erica cinerea</i> , <i>Asperula cynanchica</i> , <i>Plantago maritima</i> , <i>Thymus polytrichus</i> , <i>Briza media</i> , <i>Calluna vulgaris</i> and <i>Hedera helix</i> . Invading scrub comprises <i>Corylus avellana</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Euonymus europaeus</i> and <i>Rubus fruticosus</i> agg. <b>EC37 R3</b>
GS1	n/a	Low species-diversity variant. Typical species include <i>Trifolium repens</i> , <i>Cirsium arvense</i> , <i>Calliargonella cuspidata</i> , <i>Ranunculus repens</i> , <i>Dactylis glomerata</i> , <i>Lolium perenne</i> , <i>Cynosurus cristatus</i> , <i>Odontites verna</i> , <i>Cerastium fontanum</i> , <i>Trifolium pratense</i> , <i>Prunella vulgaris</i> , <i>Pteridium aquilinum</i> , <i>Plantago lanceolata</i> , <i>Brachythecium rutabulum</i> , <i>Holcus lanatus</i> , <i>Festuca rubra</i> , <i>Filipendula ulmaria</i> and <i>Bellis perennis</i> .
	n/a	Higher species-diversity variant. Typical species include <i>Cynosurus cristatus</i> , <i>Alopecurus pratensis</i> , <i>Trifolium repens</i> , <i>Cirsium palustre</i> , <i>Ranunculus repens</i> , <i>Ranunculus acris</i> , <i>Festuca rubra</i> , <i>Cerastium fontanum</i> , <i>Lolium perenne</i> , <i>Taraxacum officinale</i> , <i>Rumex acetosa</i> , <i>Trifolium pratense</i> , <i>Holcus lanatus</i> and <i>Plantago lanceolata</i> . <i>Leucanthemum vulgare</i> , <i>Trisetum flavescens</i> , <i>Thymus polytrichus</i> , <i>Lotus corniculatus</i> <i>Briza media</i> and <i>Cirsium arvense</i> . <b>EC37 R5 and EC37 R7</b>
GS1	6210 and *6210 (GL3a)	Typical GS1 species (mentioned above) with calcareous indicator species frequent e.g. <i>Lotus corniculatus</i> , <i>Centaurea nigra</i> , <i>Briza media</i> , <i>Carex flacca</i> , <i>Daucus carota</i> , <i>Sesleria caerulea</i> , <i>Anthyllis vulneraria</i> and <i>Thymus polytrichus</i> ; and frequent orchids <i>Anacamptis pyramidalis</i> and <i>Dactylorhiza fuchsia</i> . <b>EC37 R1, EC37 R4, EC37 R6 and EC37 R8</b>
GS1/ ER2	6210 (GL3A LPE_4B)	Species include <i>Sesleria albicans</i> , <i>Anacamptis pyramidalis</i> , <i>Briza media</i> , <i>Carex flacca</i> , <i>Thymus polytrichus</i> , <i>Teucrium scorodonia</i> , <i>Linum catharticum</i> , <i>Pilosella officinarum</i> , <i>Carlina vulgaris</i> , <i>Lotus corniculatus</i> , <i>Blackstonia perfoliata</i> , <i>Plantago maritima</i> , <i>Tortella tortuosa</i> , <i>Fissidens dubius</i> and <i>Ctenidium molluscum</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
GS2	n/a	<i>Arrhenatherum elatius</i> generally dominant with <i>Vicia cracca</i> , <i>Plantago lanceolata</i> , <i>Taraxacum officinale</i> , <i>Potentilla anserina</i> , <i>Potentilla reptans</i> , <i>Dactylis glomerata</i> , <i>Trifolium repens</i> , <i>Rhynchospora squarrosus</i> , <i>Rubus fruticosus</i> agg., <i>Ranunculus acris</i> , <i>Festuca rubra</i> , <i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Calystegia sepium</i> , <i>Cirsium arvense</i> , <i>Lathyrus pratensis</i> , <i>Heracleum sphondylium</i> . Where it occurs in a mosaic with *8240, there are some calcareous indicators (e.g. <i>Carex flacca</i> , <i>Briza media</i> , <i>Galium verum</i> , <i>Sesleria albicans</i> , <i>Leucanthemum vulgare</i> , <i>Dactylorhiza fuchsii</i> ) but not enough for 6210 and <i>A. elatius</i> still abundant in tall vegetation.
WN2	n/a	<i>Corylus avellana</i> (D), <i>Crataegus monogyna</i> , <i>Hedera helix</i> (LD), <i>Primula veris</i> , <i>Ilex aquifolium</i> , <i>Rubus fruticosus</i> agg., <i>Arum maculatum</i> , <i>Primula vulgaris</i> , <i>Prunus spinosa</i> , <i>Kindbergia praelonga</i> , <i>Thamnobryum alopecurum</i> and <i>Eurhynchium striatum</i> . Grades to WS1 at edges.
WN2/ WS1	*8240 (LPW_2a in mosaic with WS1)	<i>Corylus avellana</i> dominant (mostly <3m in height) with typical *8240 (LPW) species in understorey e.g. <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Sorbus aria</i> , <i>Sorbus hibernica</i> , <i>Fragaria vesca</i> , <i>Neckera crispa</i> , <i>Rosa spinosissima</i> , <i>Thuidium tamariscinum</i> , <i>Ilex aquifolium</i> , <i>Brachypodium sylvaticum</i> and <i>Hedera helix</i> . <b>EC37 R2</b>
WS1	n/a	Scrub with occasional mature <i>Fraxinus excelsior</i> . <i>Pteridium aquilinum</i> often abundant at edges. Additional species: <i>Rubus fruticosus</i> , <i>Prunus spinosa</i> , <i>Corylus avellana</i> (particularly on limestone pavement), <i>Epilobium hirsutum</i> , <i>Chamerion angustifolium</i> , <i>Ulex europaeus</i> and <i>Hedera helix</i> .
<b>EC39</b>		
ED3	n/a	<i>Cirsium arvense</i> , <i>Tripleurospermum inodorum</i> , <i>Polygonum aviculare</i> , <i>Plantago major</i> , <i>officinale</i> , <i>Cirsium arvensis</i> , <i>Plantago major</i> , <i>Potentilla anserina</i> with scattered <i>Rubus fruticosus</i> agg. scrub.
ER2	*8240 (LPE_1b)	Exposed limestone pavement: <i>Hedera helix</i> , <i>Corylus avellana</i> , <i>Prunus spinosa</i> , <i>Geranium robertianum</i> , <i>Sesleria albicans</i> , <i>Asplenium ruta-muraria</i> , <i>Teucrium scorodonia</i> , <i>Mycelis muralis</i> , <i>Epipactis helleborine</i> , <i>Fissidens dubius</i> , <i>Ctenidium molluscum</i> , <i>Tortella tortuosa</i> and <i>Scapania asper</i> in grykes. <b>EC39 R1 and EC39 R2</b>
FL5	n/a	<i>Nymphaea alba</i> , <i>Elodea Canadensis</i> , <i>Lemna</i> sp. The smallest water body near the eastern edge had <i>Potamogeton natans</i> , <i>Hippuris vulgaris</i> , <i>Elodea canadensis</i> , <i>Alisma plantago-aquatica</i> and <i>Ranunculus trichophyllus</i> on exposed mud at edge.
FS1	n/a	<i>Phragmites australis</i> , <i>Schoenoplectus lacustris</i> and <i>Cladium mariscus</i> are the main species. <b>EC39 R12</b>
	*7210	<i>Cladium mariscus</i> and <i>Phragmites australis</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
FW4	n/a	Mostly no emergent vegetation. Some drains with <i>Sparganium natans</i> , <i>Utricularia vulgaris</i> and <i>Myriophyllum</i> sp.
GS4	6410 (GL1d and GL1c)	<i>Molinia caerulea</i> , <i>Carex panacea</i> , <i>Carex nigra</i> , <i>Carex elata</i> , <i>Carex echinata</i> , <i>Carex pulicaris</i> , <i>Carex disticha</i> , <i>Carex ovalis</i> , <i>Juncus articulatus</i> , <i>Filipendula ulmaria</i> , <i>Potentilla erecta</i> , <i>Cirsium dissectum</i> and <i>Achillea ptarmica</i> . <i>Calliergonella cuspidata</i> is the main bryophyte, but <i>Climacium dendroides</i> and <i>Rhytidiadelphus squarrosus</i> occur towards grazed edges. At the edges of the wet meadows, where <i>Molinia caerulea</i> is very sparse or absent, other species occurring more frequently include <i>Potentilla anserina</i> , <i>Carex nigra</i> , <i>Holcus lanatus</i> , <i>Ranunculus repens</i> , <i>Galium palustre</i> , <i>Festuca arundinacea</i> and <i>Eleocharis palustris</i> . <b>EC39 R6, EC39 R7, EC39 R8, EC39 R10, EC39 R14 and EC39 R15</b>
GA1	n/a	<i>Lolium perenne</i> and <i>Trifolium repens</i> sward. <i>Cirsium arvense</i> and <i>Rumex obtusifolius</i> . Also <i>Poa Pratensis</i> , <i>Odontites verna</i> and <i>Polygonum persicaria</i> .
GS1	n/a  n/a	<i>Trifolium repens</i> , <i>Cirsium arvense</i> , <i>Calliergonella cuspidata</i> , <i>Ranunculus repens</i> , <i>Dactylis glomerata</i> , <i>Lolium perenne</i> , <i>Cynosurus cristatus</i> , <i>Anthoxanthum odoratum</i> , <i>Cerastium fontanum</i> , <i>Trifolium repens</i> , <i>Prunella vulgaris</i> , <i>Pteridium aquilinum</i> , <i>Plantago lanceolata</i> , <i>Brachythecium rutabulum</i> , <i>Holcus lanatus</i> , <i>Festuca rubra</i> , <i>Filipendula ulmaria</i> and <i>Bellis perennis</i> . High-diversity calcareous grassland. Aforementioned species with calcareous indicator species frequent e.g. <i>Lotus corniculatus</i> , <i>Centaurea nigra</i> , <i>Briza media</i> , <i>Carex flacca</i> , <i>Daucus carota</i> , <i>Sesleria caerulea</i> , <i>Anthyllis vulneraria</i> and <i>Thymus polytrichus</i> ; and frequent orchids <i>Anacamptis pyramidalis</i> , <i>Dactylorhiza fuchsii</i> . <b>EC39 R5</b>
GS1	6210 (GL3a)	Aforementioned species and <i>Agrostis capillaris</i> , <i>Trifolium pratense</i> , <i>Odontites verna</i> , <i>Ranunculus acris</i> , <i>Festuca rubra</i> , <i>Leucanthemum vulgare</i> , <i>Centaurea nigra</i> , <i>Hypochaeris radicata</i> , <i>Prunella vulgaris</i> , <i>Plantago lanceolata</i> , <i>Blackstonia perfoliata</i> and <i>Galium verum</i> . <b>EC39 R11</b>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Vicia cracca</i> , <i>Plantago lanceolata</i> , <i>Taraxacum officinale</i> , <i>Potentilla anserina</i> , <i>Dactylis glomerata</i> , <i>Trifolium repens</i> , <i>Rubus fruticosus</i> agg., <i>Ranunculus acris</i> , <i>Festuca rubra</i> , <i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Calystegia sepium</i> , <i>Cirsium arvense</i> , <i>Lathyrus pratensis</i> , <i>Heracleum sphondylium</i> . and <i>Rhytidiadelphus squarrosus</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
	6510	Typical species: <i>Leucanthemum vulgare</i> , <i>Trifolium pratense</i> , <i>Potentilla anglica</i> , <i>Plantago lanceolata</i> , <i>Helictotrichon pubescens</i> , <i>Odontites verna</i> , <i>Calliergonella cuspidata</i> , <i>Trifolium dubium</i> , <i>Trisetum flavescens</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Leontodon hispidus</i> , <i>Cerastium fontanum</i> , <i>Pimpinella major</i> , <i>Carex flacca</i> and <i>Daucus carota</i> . <b>EC39 R13</b>
PF1	n/a  7230 (RFLU4)	Non-annex habitat as the vegetation does not contain brown mosses. Typical species are <i>Phragmites australis</i> , <i>Schoenus nigricans</i> , <i>Myrica gale</i> , <i>Carex panicea</i> , <i>Carex nigra</i> , <i>Carex elata</i> , <i>Molinia caerulea</i> , <i>Hydrocotyle vulgaris</i> , <i>Menyanthes trifoliata</i> , <i>Filipendula ulmaria</i> , <i>Juncus acutiflorus</i> , <i>Agrostis stolonifera</i> , <i>Epilobium palustre</i> and <i>Ranunculus flammula</i> . <b>EC39 R3</b>  <i>Schoenus nigricans</i> , <i>Juncus subnodulosus</i> , <i>Molinia caerulea</i> , <i>Hydrocotyle vulgaris</i> , <i>Carex panicea</i> , <i>C. peliocarpa</i> , <i>Juncus articulatus</i> , <i>Ranunculus flammula</i> , <i>Cirsium dissectum</i> , <i>Parnassia palustris</i> , <i>Drosera anglica</i> , <i>Calliergonella cuspidata</i> , <i>Scorpidium scorpioides</i> , <i>S. cossonii</i> , <i>Campylium stellatum</i> , <i>Myrica gale</i> <b>EC39 R9</b>
WN2/ER2	*8240 (LPE_2a)	Wooded limestone pavement: dominated by <i>Corylus avellana</i> with <i>Crataegus monogyna</i> and <i>Prunus spinosa</i> scrub. Bryophytes include <i>Neckera complanata</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Eurhynchium striatum</i> , <i>Neckera complanata</i> and <i>Thuidium tamariscinum</i> as well as <i>Ctenidium molluscum</i> and <i>Thamnobryum alopecurum</i> .
WN2	n/a	<i>Corylus avellana</i> dominated canopy with scrub ( <i>Prunus spinosa</i> & <i>Crataegus monogyna</i> ). Understorey with typical WN2 woodland flora such as <i>Circaea lutetiana</i> , <i>Phyllitis scolopendrium</i> , <i>Viola</i> sp., <i>Primula vulgaris</i> , <i>Hedera helix</i> , <i>Geum urbanum</i> , <i>Rumex sanguineus</i> , <i>Arum maculatum</i> , <i>Fragaria vesca</i> , <i>Brachypodium sylvaticum</i> and <i>Thamnobryum alopecurum</i> .
WS1	n/a	<i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Salix cinerea</i> and <i>Salix caprea</i> trees. <i>Prunus spinosa</i> is frequent in the more open areas with <i>Rubus fruticosus</i> agg. and <i>Pteridium aquilinum</i> locally abundant. <i>Fraxinus excelsior</i> . <i>Pteridium aquilinum</i> often abundant at edges. Additional species: <i>Rubus fruticosus</i> , <i>Prunus spinosa</i> , (particularly on limestone pavement), <i>Epilobium hirsutum</i> , <i>Chamerion angustifolium</i> , <i>Ulex europaeus</i> and <i>Hedera helix</i> .
WS1/WN6	n/a	Scrub grading to wet woodland at north-western edge of Ballindooly Lough. <i>Salix cinerea</i> , <i>Phragmites australis</i> , <i>Mentha aquatica</i> , <i>Filipendula ulmaria</i> , <i>Lysimachia</i> sp. and <i>Carex</i> sp.

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
<b>EC42</b>		
ER2	*8240 (LPE_1B)	Open pavement areas have a characteristic sparse flora. The most conspicuous species include <i>Rosa spinosissima</i> , <i>Corylus avellana</i> , <i>Prunus spinosa</i> , <i>Mycelis muralis</i> , <i>Asplenium ruta-muraria</i> , <i>Senecio jacobaea</i> , <i>Teucrium scorodonia</i> , <i>Thymus polytrichus</i> , <i>Lotus corniculatus</i> , <i>Plantago maritima</i> , <i>Geranium robertianum</i> , <i>Phyllitis scolopendrium</i> , <i>Sesleria albicans</i> and <i>Carex flacca</i> .  EC42 R1, EC42 R2 and EC42 R5
ER2	*8240 (Wooded) (LPE_2B)	The dominant shrub/tree species is <i>Corylus avellana</i> with <i>Fraxinus excelsior</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Ilex aquifolium</i> and <i>Sorbus aria</i> also frequent in the low canopy. Common ground layer species include <i>Brachypodium sylvaticum</i> , <i>Rubus fruticosus</i> , <i>Hedera helix</i> , <i>Potentilla sterilis</i> , <i>Neckera crispa</i> , <i>Fragaria vesca</i> , <i>Thuidium tamariscinum</i> , <i>Primula vulgaris</i> and <i>Thamnobryum alopecurum</i> .
GA1	n/a	The characteristic species noted included <i>Lolium perenne</i> , <i>Holcus lanatus</i> , <i>Cirsium arvense</i> , <i>Cirsium vulgare</i> , <i>Trifolium repens</i> , <i>Taraxacum officinale</i> and <i>Rumex obtusifolius</i> .
GM1	n/a	The main species in the vegetation include <i>Juncus articulatus</i> , <i>Juncus effusus</i> , <i>Mentha aquatica</i> , <i>Galium palustre</i> , <i>Ranunculus repens</i> , <i>Comarum palustre</i> , <i>Carex nigra</i> , <i>Carex panacea</i> , <i>Carex disticha</i> , <i>Carex echinata</i> , <i>Carex rostrata</i> , <i>Cirsium palustre</i> , <i>Epilobium</i> sp., <i>Potentilla anserina</i> , <i>Luzula campestris</i> , <i>Menyanthes trifoliata</i> , <i>Hypericum tetrapterum</i> , <i>Filipendula ulmaria</i> , <i>Molinia caerulea</i> , <i>Salix aurita</i> , <i>Anthoxanthum odoratum</i> , <i>Hydrocotyle vulgaris</i> , <i>Persicaria amphibia</i> , <i>Eleocharis palustris</i> , <i>Glyceria fluitans</i> , <i>Apium nodiflorum</i> and the wetland moss <i>Calliergonella cuspidata</i> .
GS2	n/a	<i>Arrhenatherum elatius</i> tends to dominate the tall grassy vegetation with <i>Centaurea nigra</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Trifolium repens</i> and <i>Anthoxanthum odoratum</i> .
GS1	n/a	Common calcareous grassland species noted include <i>Lotus corniculatus</i> , <i>Briza media</i> , <i>Carex flacca</i> , <i>Galium verum</i> , <i>Linum catharticum</i> , <i>Thymus polytrichus</i> and <i>Leontodon</i> sp.
GS1	6210	Typical species include <i>Lotus corniculatus</i> , <i>Briza media</i> , <i>Molinia caerulea</i> , <i>Carex flacca</i> , <i>Lathyrus pratensis</i> , <i>Scleropodium purum</i> , <i>Calliergonella cuspidata</i> , <i>Euphrasia</i> sp., <i>Centaurea nigra</i> , <i>Holcus lanatus</i> , <i>Hypericum maculatum</i> , <i>Potentilla erecta</i> , <i>Thuidium tamariscinum</i> , <i>Hypochaeris radicata</i> , <i>Trifolium pratense</i> , <i>Ctenidium molluscum</i> , <i>Leucanthemum vulgare</i> , <i>Festuca rubra</i> , <i>Hypericum pulchrum</i> , <i>Potentilla sterilis</i> , <i>Danthonia decumbens</i> , <i>Linum catharticum</i> , <i>Thymus polytrichus</i> , <i>Prunella vulgaris</i> , <i>Veronica chamaedrys</i> , <i>Daucus carota</i> and <i>Galium verum</i> .  EC42 R4

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WN2	*8240	<i>Corylus avellana</i> , <i>Ilex aquifolium</i> , <i>Hedera helix</i> , <i>Brachypodium sylvaticum</i> , <i>Fraxinus excelsior</i> , <i>Potentilla sterilis</i> , <i>Rubus fruticosus</i> , <i>Crataegus monogyna</i> , <i>Fragaria vesca</i> , <i>Primula vulgaris</i> , <i>Taraxacum</i> agg., <i>Brachythecium rutabulum</i> , <i>Geum urbanum</i> , <i>Lonicera periclymenum</i> , <i>Prunus spinosa</i> , <i>Thamnobryum alopecurum</i> , <i>Viola</i> sp., <i>Thuidium tamariscinum</i> and <i>Phyllitis scolopendrium</i> . <b>EC42 R3</b>
WS1	n/a	<i>Corylus avellana</i> , <i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Ilex aquifolium</i> , <i>Sorbus aria</i> , <i>Rubus fruticosus</i> , <i>Hedera helix</i> , <i>Fragaria vesca</i> , <i>Thuidium tamariscinum</i> , <i>Geranium robertianum</i> , <i>Primula vulgaris</i> and <i>Kindbergia praelonga</i> .
<b>EC44</b>		
ER2	*8240 (LPE_1e)	<i>Hedera helix</i> , <i>Corylus avellana</i> , <i>Sesleria albicans</i> , <i>Teucrium scorodonia</i> , <i>Pteridium aquilinum</i> , <i>Neckera crispa</i> , <i>Ctenidium molluscum</i> and <i>Rubia peregrina</i>
GS1	n/a	<i>Briza media</i> , <i>Dactylis glomerata</i> , <i>Anthoxanthum odoratum</i> , <i>Holcus lanatus</i> , <i>Festuca rubra</i> , <i>Centaurea nigra</i> , <i>Linum catharticum</i> , <i>Lotus corniculatus</i> , <i>Euphrasia</i> sp., <i>Carex flacca</i> , <i>Rubus fruticosus</i> , <i>Blackstonia perfoliata</i> , <i>Daucus carota</i> , <i>Leucanthemum vulgare</i> , <i>Hypericum androsaemum</i> and <i>Potentilla reptans</i> . <b>EC44 R4</b>
GS2/GA1	n/a	<i>Rumex sanguineus</i> , <i>Cirsium arvense</i> , <i>Holcus lanatus</i> , <i>Plantago lanceolata</i> , <i>Senecio jacobaea</i> , <i>Cerastium fontanum</i> , <i>Urtica dioica</i> , <i>Heracleum sphondylium</i> , <i>Ranunculus repens</i> and <i>Potentilla anserina</i>
WN2/ER2	*8240 (LPW_2a)	<i>Corylus avellana</i> dominated canopy (4-5m high) with occasional <i>Crataegus monogyna</i> , <i>Ilex aquifolium</i> and <i>Sorbus aucuparia</i> . <i>Prunus spinosa</i> localised <2m high. Ground flora dominated by bryophytes notably <i>Eurhynchium striatum</i> , but also <i>Rhytidiadelphus triquetrus</i> and <i>Thuidium tamariscinum</i> and <i>Neckera</i> spp. common on base of trees. <i>Hedera helix</i> locally frequent. <b>EC44 R1, EC44 R2</b>
WN2	n/a	<i>Corylus avellana</i> dominated canopy (6-9m high) with occasional <i>Prunus spinosa</i> & <i>Crataegus monogyna</i> . Understorey with typical WN2 woodland flora such as <i>Veronica montana</i> , <i>Circaea lutetiana</i> , <i>Sanicula europaea</i> , <i>Viola</i> sp., <i>Primula vulgaris</i> , <i>Hedera helix</i> , <i>Carex sylvatica</i> , <i>Brachypodium sylvaticum</i> , <i>Fragaria vesca</i> , <i>Epipactis helleborine</i> , <i>Thamnobryum alopecurum</i> and <i>Plagiomnium undulatum</i> .
WS1/WN2	n/a	<i>Prunus spinosa</i> dominant scrub species with occasional <i>Crataegus monogyna</i> and <i>Corylus avellana</i> . Also <i>Rubus fruticosus</i> agg. and occasionally <i>Pteridium aquilinum</i> and <i>Urtica dioica</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WS1	*8240	<i>Corylus avellana</i> , <i>Hedera helix</i> , <i>Eurhynchium striatum</i> , <i>Brachypodium sylvaticum</i> , <i>Thamnobryum alopecurum</i> , <i>Crataegus monogyna</i> , <i>Kindbergia praelonga</i> , <i>Potentilla sterilis</i> , <i>Viola riviniana</i> , <i>Lonicera periclymenum</i> , <i>Ctenidium molluscum</i> , <i>Plagiochila asplenioides</i> , <i>Rubus fruticosus</i> , <i>Fragaria vesca</i> , <i>Solidago virgaurea</i> , <i>Sorbus aucuparia</i> , <i>Prunus spinosa</i> , <i>Primula vulgaris</i> , <i>Fissidens</i> sp., <i>Carex</i> sp., <i>Phyllitis scolopendrium</i> , <i>Epipactis helleborine</i> and <i>Sanicula europaea</i> . <b>EC44 R3</b>
WS1	n/a	<i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Urtica dioica</i> , <i>Rubus fruticosus</i> and <i>Pteridium aquilinum</i> .
<b>EC47</b>		
GS1	n/a	<i>Pseudoscleropodium purum</i> , <i>Rhytidiadelphus squarrosus</i> , <i>Anthoxanthum odoratum</i> , <i>Holcus lanatus</i> , <i>Cynosurus cristatus</i> , <i>Trifolium repens</i> (O-LF), <i>Geranium robertianum</i> , <i>Rubus fruticosus</i> agg., <i>Fragaria vesca</i> , <i>Lotus corniculatus</i> , <i>Galium verum</i> and <i>Cirsium arvense</i> .
WN2/ER2	*8240 (LPW_2b)	<i>Fraxinus excelsior</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Rubus fruticosus</i> agg., <i>Asplenium scolopendrium</i> , <i>Viola</i> sp., <i>Dryopteris filix mas</i> , <i>Fragaria vesca</i> , <i>Primula veris</i> , <i>Veronica chamaedrys</i> , <i>Eurhynchium striatum</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Thamnobryum alopecurum</i> and <i>Scapania aspera</i> . <b>EC47 R1</b>
WS1/WN2/ER2	n/a and *8240	<i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Fraxinus excelsior</i> , <i>Corylus avellana</i> , <i>Ilex aquifolium</i> , <i>Urtica dioica</i> , <i>Geranium robertianum</i> , <i>Hedera helix</i> , <i>Veronica chamaedrys</i> , <i>Urtica dioica</i> . Bryophytes 70% cover: <i>Thamnobryum alopecurum</i> and <i>Eurhynchium striatum</i> . Some small areas of *8240 (LPW), within the woodland, but too small to map.
<b>EC48</b>		
WS1/WN2/ER2	n/a	<i>Prunus spinosa</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Urtica dioica</i> , <i>Geranium robertianum</i> , <i>Primula veris</i> , <i>Thamnobryum alopecurum</i> , <i>Eurhynchium striatum</i> and <i>Thuidium tamariscinum</i> .
<b>EC49</b>		
ED3/GS1/W S1	n/a	Bare flat disturbed ground with Pockets of grassland: <i>Centaurea nigra</i> , <i>Achillea millefolium</i> , <i>Linum catharticum</i> , <i>Blackstonia perfoliata</i> and <i>Centaureum erythraea</i> . Piles of spoil with <i>Buddleja davidii</i> , <i>Rubus fruticosus</i> agg, <i>Prunus spinosa</i> , <i>Centranthus ruber</i> , <i>Petasites</i> sp. and <i>Cotoneaster</i> sp.
GS1	n/a	<i>Anthoxanthum odoratum</i> , <i>Cynosurus cristatus</i> , <i>Trifolium repens</i> , <i>Ranunculus acris</i> , <i>Carex flacca</i> , <i>Linum catharticum</i> , <i>Rhinanthus minor</i> , <i>Euphrasia</i> sp. and <i>Odontites verna</i> .

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WN2	n/a	<i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Rubus fruticosus</i> agg. <i>Phyllitis scolopendrium</i> , <i>Arum maculatum</i> , <i>Lonicera periclymenum</i> , <i>Primula veris</i> , <i>Dryopteris filix mas</i> , <i>Thamnobryum alopecurum</i> and <i>Thuidium tamariscinum</i> .
<b>EC56</b>		
ER2	*8240 (LPE_1b)	<i>Corylus avellana</i> , <i>Ulex europaeus</i> , <i>Juniperus communis</i> , <i>Hedera helix</i> , <i>Rosa spinosissima</i> , <i>Teucrium scorodonia</i> , <i>Sesleria albicans</i> , <i>Mycelis muralis</i> , <i>Asplenium ruta-muraria</i> , <i>Geranium robertianum</i> , <i>Carlina vulgaris</i> , <i>Neckera crispa</i> , <i>Fissidens dubius</i> , <i>Tortella tortuosa</i> , <i>Ctenidium molluscum</i> , <i>Festuca rubra</i> , <i>Koeleria macrantha</i> , <i>Thymus praecox</i> , <i>Antennaria dioica</i> , <i>Lotus corniculatus</i> , <i>Hypericum pulchrum</i> , <i>Gentiana verna</i> , <i>Carex caryophyllea</i> , <i>Dactylorhiza fuchsia</i> , <i>Succisa pratensis</i> , <i>Rubus fruticosus</i> , <i>Phyllitis scolopendrium</i> , <i>Fragaria vesca</i> , <i>Dactylis glomerata</i> and <i>Solanum dulcamara</i> . <b>EC56 R2 and EC56 R5</b>
FL8/FS1	n/a	Wetland forming in excavated attenuation pond area adjacent to west side of M6/N6 junction. <i>Phragmites australis</i> , <i>Typha latifolia</i> , <i>Sparganium erectum</i> , <i>Schoenoplectus lacustris</i> , <i>Apium nodiflorum</i> , <i>Nasturtium officinale</i> , <i>Lemna minor</i> and <i>Epilobium hirsutum</i>
GA1	n/a	<i>Lolium perenne</i> , <i>Trifolium repens</i> , <i>Juncus bufonius</i> , <i>Rumex obtusifolius</i> and <i>Poa pratensis</i> .
GS1	n/a	<i>Anthoxanthum odoratum</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Cynosurus cristatus</i> , <i>Plantago lanceolata</i> , <i>Centaurea nigra</i> , <i>Trifolium repens</i> , <i>Trifolium pratense</i> , <i>Lotus corniculatus</i> , <i>Ranunculus bulbosus</i> , <i>Achillea millefolium</i> , <i>Luzula campestris</i> , <i>Pteridium aquilinum</i> , <i>Hypochaeris radicata</i> , <i>Dactylorhiza fuchsii</i> , <i>Danthonia decumbens</i> , <i>Potentilla anserina</i> , <i>Senecio jacobaea</i> , <i>Bellis perennis</i> , <i>Agrostis canina</i> , <i>Prunella vulgaris</i> , <i>Taraxacum</i> agg., <i>Leucanthemum vulgare</i> , <i>Euphrasia</i> sp., <i>Briza media</i> , <i>Sonchus</i> sp., <i>Urtica dioica</i> , <i>Succisa pratensis</i> , <i>Rumex obtusifolius</i> , <i>Odontites verna</i> , <i>Pimpinella saxifraga</i> and <i>Primula vulgaris</i> . <b>EC56 R8 and EC56 R9</b>
GS1	6210 (GS1_3a)	<i>Succisa pratensis</i> , <i>Briza media</i> , <i>Danthonia decumbens</i> , <i>Festuca ovina</i> , <i>Schoenus nigricans</i> , <i>Cynosurus cristatus</i> , <i>Sesleria albicans</i> , <i>Carex pulicaris</i> , <i>Carex panacea</i> , <i>Carex demissa</i> , <i>Carex flacca</i> , <i>Pinguicula vulgaris</i> , <i>Trifolium repens</i> , <i>Antennaria dioica</i> , <i>Blackstonia perforata</i> , <i>Plantago maritima</i> , <i>Anagallis tenella</i> , <i>Selaginella</i> sp., <i>Linum catharticum</i> , <i>Cirsium dissectum</i> , <i>Molinia caerulea</i> , <i>Gentiana verna</i> , <i>Euphrasia</i> sp., <i>Polygala vulgaris</i> , <i>Prunella vulgaris</i> , <i>Lathyrus linifolius</i> , <i>Dactylorhiza fuchsii</i> , <i>Platanthera bifolia</i> , <i>Calliargonella cuspidata</i> , <i>Ctenidium molluscum</i> , <i>Fissidens adianthoides</i> and <i>Didymodon ferrugineus</i> . <b>EC56 R1, EC56 R3, EC56 R4 and EC56 R7</b>
GS4/GS1	n/a	<i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Juncus articulatus</i> , <i>Carex flacca</i> , <i>Carex panacea</i> , <i>Potentilla anserina</i> , <i>Epilobium hirsutum</i> and <i>Ranunculus repens</i> . On drier patches <i>Centaurea nigra</i> , <i>Leucanthemum vulgare</i> and <i>Daucus carota</i> .



Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WN2/ER2	*8240 (LPW_2a)	<i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Rosa spinosissima</i> , <i>Fragaria vesca</i> , <i>Viola</i> sp. <i>Rubus fruticosus</i> agg., <i>Neckera crispa</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Ctenidium molluscum</i> , <i>Eurhynchium striatum</i> , <i>Thamnobryum alopecurum</i> and <i>Thuidium tamariscinum</i> . <b>EC56 R6</b>
WN2	n/a	Similar to WN2/ER2, and with similar species, as well as the following: <i>Acer pseudoplatanus</i> , <i>Prunus spinosa</i> . Other ground flora species include <i>Arum maculatum</i> , <i>Circaea lutetiana</i> and <i>Hedera helix</i> . Bryophyte cover <50% <i>Thamnobryum alopecurum</i> and <i>Thuidium tamariscinum</i> .
WS1	n/a	<i>Crataegus monogyna</i> , <i>Prunus spinosa</i> , <i>Ilex aquifolium</i> , <i>Sorbus hibernica</i> , <i>S. aucuparia</i> , <i>Rubus fruticosus</i> agg., <i>Corylus avellana</i> , <i>Acer pseudoplatanus</i> , <i>Ilex aquifolium</i> , <i>Salix caprea</i> , <i>Hedera helix</i> , <i>Lonicera periclymenum</i> , <i>Fragaria vesca</i> , <i>Pteridium aquilinum</i> , <i>Brachypodium sylvaticum</i> , <i>Fraxinus excelsior</i> and <i>Ulex europaeus</i> .
<b>EC57</b>		
ED3	n/a	<i>Brassica rapa</i> , <i>Urtica dioica</i> , <i>Polygonum persicaria</i> , <i>Cirsium arvense</i> , <i>Potentilla anserina</i> and <i>Matricaria discoidea</i> .
GS1	n/a  6210 (GS1_3A)	<i>Cynosurus cristatus</i> , <i>Holcus lanatus</i> , <i>Poa pratensis</i> , <i>Cirsium arvense</i> , <i>Trifolium repens</i> , <i>T. Pratense</i> , <i>Rumex acetosa</i> , <i>Pteridium aquilinum</i> and <i>Prunus spinosa</i> .  <i>Anthyllis vulneraria</i> , <i>Briza media</i> , <i>Danthonia decumbens</i> , <i>Anthoxanthum odoratum</i> , <i>Festuca rubra</i> , <i>Koeleria macrantha</i> , <i>Leontodon hispidus</i> , <i>Potentilla erecta</i> , <i>Sesleria albicans</i> , <i>Carex caryophylla</i> , <i>C. panacea</i> , <i>C. flacca</i> , <i>Antennaria dioica</i> , <i>Linum catharticum</i> , <i>Galium verum</i> , <i>Plantago maritima</i> , <i>Succisa pratensis</i> , <i>Centaurea nigra</i> , <i>Euphrasia</i> sp. <i>Polygala vulgaris</i> , <i>Prunella vulgaris</i> and <i>Lotus corniculatus</i> . <b>EC57 R1 and EC57 R2</b>
GS1	*6210 (GS1_3A)	Includes a small area of orchid rich grassland (Priority Annex I *6210). Includes the orchids <i>Dactylorhiza fuchsii</i> , <i>Platanthera bifolia</i> and <i>Listera ovata</i> and without the <i>Anthyllis vulneraria</i> listed above, plus <i>Molinia caerulea</i> and <i>Lathyrus linifolius</i> . <b>EC57 R4</b>
GS2	n/a	<i>Holcus lanatus</i> , <i>Poa pratensis</i> , <i>Lolium perenne</i> , <i>Dactylis glomerata</i> , <i>Rumex acetosa</i> , <i>Ranunculus repens</i> and <i>Cynosurus cristatus</i> .
WN2/ER2	*8240 (LPW_2a)	<i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Prunus spinosa</i> , <i>Fragaria vesca</i> , <i>Viola</i> sp., <i>Rubus fruticosus</i> agg., <i>Hedera helix</i> , <i>Lonicera periclymenum</i> , <i>Brachypodium sylvaticum</i> , <i>Arum maculatum</i> , <i>Circaea lutetiana</i> , <i>Geum urbanum</i> , <i>Phyllitis scolopendrium</i> , <i>Dryopteris filix-mas</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Eurhynchium striatum</i> , <i>Thamnobryum alopecurum</i> and <i>Thuidium tamariscinum</i> . <b>EC57 R3</b>

Habitat code <sup>2</sup>	Annex I habitat <sup>3</sup>	Key species
WS1/ER2	n/a	<i>Prunus spinosa</i> , <i>Crataegus monogyna</i> , <i>Corylus avellana</i> , <i>Rubus fruticosus</i> agg., <i>Pteridium aquilinum</i> , <i>Hedera helix</i> , <i>Rosa spinosissima</i> , <i>Teucrium scorodonia</i> , <i>Solidago virgaurea</i> , <i>Brachypodium sylvaticum</i> , <i>Fragaria vesca</i> , <i>Lathyrus linifolius</i> and <i>Molinia caerulea</i> .
WS1/ER2	*8240	<i>Prunus spinosa</i> , <i>Corylus avellana</i> , <i>Fraxinus excelsior</i> , <i>Crataegus monogyna</i> , <i>Ulex europaeus</i> , <i>Rosa spinosissima</i> , <i>Rubus fruticosus</i> agg., <i>Pteridium aquilinum</i> , <i>Hedera helix</i> , <i>Teucrium scorodonia</i> , <i>Lonicera periclymenum</i> , <i>Solidago virgaurea</i> , <i>Sesleria albicans</i> , <i>Brachypodium sylvaticum</i> , <i>Fragaria vesca</i> , <i>Thymus praecox</i> , <i>Molinia caerulea</i> , <i>Ctenidium molluscum</i> and <i>Scleropodium purum</i> . <b>EC57 R5</b>
<b>EC63</b>		
ED3	n/a	Variable but includes species such as <i>Holcus lanatus</i> , <i>Fumaria officinalis</i> , <i>Trifolium pratensis</i> , <i>T. repens</i> , <i>Plantago lanceolata</i> and <i>Sisymbrium officinale</i> .
HD1	n/a	<i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg. and <i>Chamerion angustifolium</i> .
GA2	n/a	<i>Lolium perenne</i> , <i>Bellis perennis</i> and <i>Trifolium repens</i> .
GS2	n/a	<i>Holcus lanatus</i> , <i>Plantago lanceolata</i> , <i>Trifolium repens</i> , <i>Senecio jacobaea</i> , <i>Dactylis glomerata</i> , <i>Arrhenatherum elatius</i> , <i>Cirsium arvensis</i> and occasional <i>Leucanthemum vulgare</i> .
GS4	n/a	<i>Juncus effusus</i> , <i>Filipendula ulmaria</i> , <i>Rumex obtusifolius</i> , <i>Chamerion angustifolium</i> , <i>Arrhenatherum elatius</i> , <i>Holcus lanatus</i> and <i>Ranunculus repens</i> .
WL1	n/a	<i>Cupressus x leylandii</i> , <i>Griselinia</i> sp., <i>Fuchsia magellanica</i> , <i>Fagus sylvatica</i> F. <i>purpurea</i> ., <i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Fraxinus excelsior</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Viburnum opulus</i> , <i>Buddleja davidii</i> , <i>Alnus glutinosa</i> , <i>Ulex europaeus</i> , <i>Betula pubescens</i> , <i>Cornus</i> sp. <i>Ulex europaeus</i> , <i>Alnus incana</i> , <i>Salix caprea</i> and <i>Calystegia sepium</i> .
WS1	n/a	<i>Ulex europaeus</i> , <i>Rubus fruticosus</i> agg., <i>Urtica dioica</i> and some <i>Salix cinerea</i> on damp ground.

## A.8.19.2

---

## Habitat Species Lists – Areas between Ecological Sites and Excluding the Lough Corrib cSAC Habitat Survey Area

**Table A.8.19.2** presents the species lists for those habitat areas between the ecological sites; excluding the area between the River Corrib and Lackagh Quarry (which is described separately in **Appendix A.8.19.3**). It does not include species lists for the residential habitat category (see **Section 8.3.4.34** of **Chapter 8, Biodiversity**) and it also does not include species lists for the aquatic habitats, which are described in full in **Section 8.3.4** of **Chapter 8, Biodiversity** and in **Appendix A.8.20 (Results of the 2014 N6 Galway City Transport Project Aquatic Habitat Surveys)**.

Where relevés were taken in the field, the reference codes are provided in the Key Species column (e.g. EC03 R1). These relevé codes refer to the full species lists which are provided in **Appendix A.8.19.4**.

**Table A.8.19.2: Habitat Species Lists Relating to habitat areas within, or adjacent to, the Proposed Road Development that lie between the Ecological Sites**

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
R336 to the Foráí Maola Road Ch. 0+000 – Ch. 1+150 Habitats between EC03, EC05 and EC11 as far as the Foráí Maola Road		
ED3	n/a	<i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Anagallis tenella</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Bellis perennis</i> , <i>Campylopus introflexus</i> (A/F), <i>Centaurea erythraea</i> , <i>Erica cinerea</i> , <i>Hieracium</i> agg., <i>Holcus lanatus</i> , <i>Leucanthemum vulgare</i> , <i>Lotus corniculatus</i> , <i>Nardus stricta</i> , <i>Potentilla erecta</i> , <i>Potentilla reptans</i> , <i>Senecio jacobaea</i> , <i>Succisa pratensis</i> and <i>Ulex gallii</i>
ED3a	n/a	<i>Agrostis stolonifera</i> , <i>Calluna vulgaris</i> , <i>Carex panicea</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> and <i>Molinia caerulea</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Cirsium vulgare</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago major</i> , <i>Poa annua</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Sonchus asper</i> , <i>Taraxacum officinale</i> agg. and <i>Trifolium repens</i>
GM1	n/a	<i>Angelica sylvestris</i> , <i>Cirsium palustre</i> , <i>Filipendula ulmaria</i> (A), <i>Galium palustre</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> (F), <i>Lotus pedunculatus</i> , <i>Lythrum salicaria</i> (F), <i>Mentha aquatica</i> , <i>Potentilla anserina</i> , <i>Ranunculus acris</i> , <i>Scrophularia auriculata</i> and <i>Typha latifolia</i>

<sup>4</sup> Fossitt (2000)

<sup>5</sup> Where applicable, vegetation community codes are given in parenthesis.

<sup>6</sup> Species abundance described (where relevant) using DAFOR scale: D = dominant; A = Abundant; F = Frequent; O = Occasional; R = Rare. L = Locally (e.g. LF = Locally Frequent)

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
GS2	n/a	<i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Cerastium fontanum</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Leontodon autumnalis</i> , <i>Lotus corniculatus</i> , <i>Plantago lanceolata</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rumex acetosa</i> and <i>Trifolium pratensis</i>
GS3	n/a	<i>Achillea millefolium</i> , <i>Agrostis</i> sp., <i>Lolium</i> sp., <i>Ranunculus repens</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> and <i>Urtica dioica</i>
GS4	n/a	<i>Angelica sylvestris</i> , <i>Calystegia sepium</i> , <i>Cirsium palustre</i> , <i>Festuca rubra/ovina</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Iris pseudacorus</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Lythrum salicaria</i> , <i>Molinia caerulea</i> , <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Pteridium aquilinum</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex obtusifolius</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> and <i>Trifolium pratense</i>
HD1	n/a	<i>Pteridium aquilinum</i> and <i>Rubus fruticosus</i> agg.
HH1	4030 (DH1)	<i>Ulex gallii</i> , <i>Erica cinerea</i> , <i>Hypnum jutlandicum</i> , <i>Agrostis stolonifera</i> , <i>Calluna vulgaris</i> , <i>Pteridium aquilinum</i> , <i>Daboecia cantabrica</i> , <i>Hylocomium splendens</i> , <i>Holcus lanatus</i> , <i>Campylopus introflexus</i> , <i>Deschampsia flexuosa</i> , <i>Molinia caerulea</i> , <i>Rubus fruticosus</i> agg., <i>Cladonia</i> sp., <i>Danthonia decumbens</i> , <i>Hypnum jutlandicum</i> , <i>Scleropodium purum</i> , <i>Campylopus flexuosus</i> , <i>Carex binervis</i> , <i>Hylocomium splendens</i> , <i>Juncus squarrosus</i> , <i>Nardus stricta</i> , <i>Potentilla erecta</i> , <i>Racomitrium lanuginosum</i> , <i>Sedum anglicum</i> , <i>Succisa pratensis</i> and <i>Teucrium scorodonia</i> <b>R336-NFMR R1 and R336-NFMR R2</b>
WD1	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus cordata</i> , <i>Alnus glutinosa</i> , <i>Blechnum spicant</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Dryopteris felix-mas</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Phyllitis scolopendrium</i> , <i>Prunus</i> sp., <i>Quercus robur</i> , <i>Rubus fruticosus</i> agg., <i>Salix</i> sp., <i>Sorbus aria</i> and <i>Urtica dioica</i>
WD2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus cordata</i> , <i>Alnus glutinosa</i> , <i>Betula pubescens</i> , <i>Fagus sylvatic</i> , <i>Fraxinus excelsior</i> , <i>Picea sitchensis</i> , <i>Pinus</i> spp. and <i>Quercus robur</i>
WL1	n/a	<i>Cotoneaster</i> sp., <i>Crococsmia x crocosmiiflora</i> , <i>Griselinia</i> sp. and <i>Lonicera nitida</i>
WL1a	n/a	<i>Calystegia sepium</i> , <i>Epilobium hirsutum</i> , <i>Filipendula ulmaria</i> , <i>Geranium robertianum</i> , <i>Hedera helix</i> , <i>Lonicera periclymenum</i> , <i>Phyllitis scolopendrium</i> , <i>Prunus spinosa</i> and <i>Pteridium aquilinum</i>
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus glutinosa</i> , <i>Cupressus</i> sp., <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Pinus contorta</i> , <i>Pinus radiata</i> , <i>Prunus</i> sp. and <i>Sorbus aucuparia</i>
WS1	n/a	<i>Acer pseudoplatanus</i> , <i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Dactylis glomerata</i> , <i>Epilobium</i> sp., <i>Lythrum salicaria</i> , <i>Rubus fruticosus</i> agg. and <i>Vicia cracca</i>
WS1/WN6	n/a	<i>Salix cinerea</i> , <i>Rubus fruticosus</i> agg., <i>Angelica sylvestris</i> and <i>Arrhenatherum elatius</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
The Foraf Maola Road to Troscaigh Thiar Ch. 1+150 – Ch. 1+650 Habitats between EC12, EC13, EC14 and the Foraf Maola Road		
BL1	n/a	<i>Cladonia</i> spp., <i>Dicranum scoparium</i> , <i>Hedera helix</i> , <i>Homalothecium sericeum</i> , <i>Hypnum cupressiforme</i> , <i>Isothecium myosuroides</i> , <i>Mnium hornum</i> , <i>Polypodium vulgare</i> and <i>Racomitrium aciculare</i>
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i>
FS2	n/a	<i>Apium nodiflorum</i> , <i>Equisetum fluviatile</i> , <i>Glyceria fluitans</i> , <i>Iris pseudacorus</i> , <i>Mentha aquatica</i> and <i>Veronica beccabunga</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Lolium perenne</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> and <i>Urtica dioica</i>
GS2	n/a	<i>Agrostis stolonifera</i> , <i>Angelica sylvestris</i> , <i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Cirsium palustre</i> , <i>Hieracium</i> agg., <i>Holcus lanatus</i> , <i>Leucanthemum vulgare</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Ranunculus repens</i> , <i>Rumex crispus</i> , <i>Scrophularia auriculata</i> , <i>Sonchus asper</i> and <i>Trifolium dubium</i>
GS3	n/a	<i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Calliargonella cuspidatum</i> , <i>Cerastium fontanum</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Lotus corniculatus</i> , <i>Pilosella officinarum</i> , <i>Plantago lanceolata</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rhytidadelphus squarrosus</i> , <i>Rumex acetosa</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratensis</i> , <i>Trifolium repens</i> , <i>Veronica chamaedrys</i> , <i>Veronica serpyllifolia</i> and <i>Viola riviniana</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Centaurea nigra</i> , <i>Cirsium palustre</i> , <i>Dactylis glomerata</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Iris pseudacorus</i> , <i>Juncus effusus</i> , <i>Lythrum salicaria</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rumex</i> spp. and <i>Succisa pratensis</i>
HD1	n/a	<i>Pteridium aquilinum</i>
HH1	4030 (DH1)	<i>Calluna vulgaris</i> , <i>Cladonia</i> sp., <i>Daboecia cantabrica</i> , <i>Danthonia decumbens</i> , <i>Deschampsia flexuosa</i> , <i>Erica cinerea</i> , <i>Hypnum jutlandicum</i> , <i>Molinia caerulea</i> , <i>Scleropodium purum</i> and <i>Ulex gallii</i>
WD2	n/a	<i>Acer pseudoplatanus</i> , <i>Betula pubescens</i> , <i>Cupressus</i> sp., <i>Fraxinus excelsior</i> , <i>Pinus</i> spp., <i>Prunus</i> sp. and <i>Rhododendron ponticum</i>
WL1	n/a	<i>Crataegus monogyna</i> , <i>Cupressus x leylandii</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rosa rugosa</i> and <i>Rubus fruticosus</i> agg.
WL2	n/a	<i>Cupressus</i> sp.
WS1	n/a	<i>Calystegia sepium</i> , <i>Crataegus monogyna</i> , <i>Dryopteris felix-mas</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Lonicera periclymenum</i> , <i>Phyllitis scolopendrium</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Ulex europaeus</i> , <i>Umbilicus rupestris</i> , <i>Veronica chamaedrys</i> and <i>Vicia sepium</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
Ann Gibbons Road (L13215) Ch. 2+400 – Ch. 2+500 Habitats between EC14 and EC09		
ED3	n/a	<i>Plantago major</i> , <i>Poa annua</i> , <i>Potentilla anserina</i>
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Lolium perenne</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> and <i>Urtica dioica</i>
GS3	n/a	<i>Agrostis capillaris</i> , <i>Cynosurus cristatus</i> , <i>Holcus lanatus</i> , <i>Leontodon autumnalis</i> , <i>Plantago lanceolata</i> , <i>Potentilla erecta</i> and <i>Ranunculus repens</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Lythrum salicaria</i> and <i>Potentilla anserina</i>
WS1	n/a	<i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg. and <i>Ulex europaeus</i>
Ballard West to Ballard East An Chloch Scoilte Junction, Ch. 3+050 – Ch. 3+450 Habitats between EC09 and EC18		
ER1	n/a	The exposed rock is mostly bare. It has few crevices or clefts to harbour significant bryophyte or pteridophyte communities. Where rock outcrops occur in grassland habitats, they are often surrounded by species that prefer good drainage such as <i>Sedum anglica</i> , <i>Pilosella officinarum</i> and <i>Veronica serpyllifolia</i>
FS2	n/a	<i>Apium nodiflorum</i> , <i>Epilobium hirsutum</i> and <i>Iris pseudacorus</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Calystegia sepium</i> , <i>Cirsium arvense</i> , <i>Galium aparine</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex obtusifolius</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Urtica dioica</i>
GM1	n/a	<i>Carex panicea</i> , <i>Hydrocotyle vulgare</i> , <i>Hypericum perforatum</i> , <i>Juncus articulatus</i> , <i>Leontodon autumnalis</i> , <i>Lythrum salicaria</i> , <i>Mentha aquatica</i> , <i>Myosotis scorpioides</i> , <i>Potentilla palustris</i> , <i>Ranunculus acris</i> and <i>Solidago virgaurea</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Arrhenatherum odoratum</i> , <i>Cirsium arvensis</i> , <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Filipendula ulmaria</i> , <i>Geranium robertianum</i> , <i>Holcus lanatus</i> , <i>Iris pseudacorus</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Lythrum salicaria</i> , <i>Plantago lanceolata</i> , <i>Poa annua</i> , <i>Potentilla anserina</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex acetosa</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Urtica dioica</i> and <i>Veronica persica</i>
WD1	n/a	<i>Acer pseudoplatanus</i> , <i>Cotoneaster</i> sp., <i>Crataegus monogyna</i> , <i>Dryopteris felix-mas</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Heracleum sphondylium</i> , <i>Phyllitis scolopendrium</i> , <i>Rubus fruticosus</i> agg., <i>Symphoricarpos alba</i> and <i>Urtica dioica</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
WS1	n/a	<i>Prunus spinosa</i> , <i>Ulex europaeus</i> , <i>Crataegus monogyna</i> , <i>Rubus fruticosus</i> agg. A small area scrub associated with an abandoned residential property has several non-native garden shrubs including <i>Cotoneaster</i> sp. and <i>Symphoricarpos alba</i> .
Cappagh Ch. 3+900 – Ch. 4+650 Habitats between EC18 and EC20		
BL1	n/a	Few epiphytes for the most part. Significant bryophyte communities in some places but few higher plant species. <i>Brachythecium rutabulum</i> , <i>Cladonia</i> spp., <i>Dicranum scoparium</i> , <i>Frullania dilatata</i> , <i>Hedera helix</i> , <i>Homalothecium</i> , <i>Hypnum cupressiforme</i> , <i>Isothecium myosuroides</i> , <i>Mnium hornum</i> and <i>Racomitrium aciculare</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Cirsium arvense</i> , <i>Cynosurus cristatus</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago major</i> , <i>Ranunculus repens</i> , <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Urtica dioica</i>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Ranunculus repens</i> and <i>Rumex crispus</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Lythrum salicaria</i> , <i>Potentilla anserina</i> and <i>Ranunculus acris</i>
PF2	n/a	<i>Calliergonella cuspidata</i> , <i>Carex echinata</i> , <i>Carex panicea</i> , <i>Hypericum elodes</i> , <i>Juncus articulatus</i> , <i>Juncus bulbosus</i> , <i>Ranunculus flammula</i> and <i>Sphagnum palustre</i>
WL1	n/a	<i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg. and <i>Urtica dioica</i>
WL2	n/a	Varying species compositions across the site. <i>Pinus</i> sp. and <i>Alnus cordata</i> are dominant in some treelines but the majority are composed of <i>Fraxinus excelsior</i> , <i>Alnus glutinosa</i> and <i>Acer pseudoplatanus</i>
WS1	n/a	<i>Crataegus monogyna</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg. and <i>Ulex europaeus</i> (D)
Ballymoneen Road Ballymoneen Road Junction, Ch. 5+550 – Ch. 5+700 Habitats between EC20 and EC21		
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Cirsium arvense</i> , <i>Lolium perenne</i> , <i>Plantago major</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Trifolium repens</i> and <i>Urtica dioica</i>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Prunella vulgaris</i> and <i>Succisa pratensis</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Lythrum salicaria</i> and <i>Ranunculus acris</i>
WL1	n/a	<i>Crataegus monogyna</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg. and <i>Urtica dioica</i>
WL2	n/a	<i>Cupressus</i> sp., <i>Fraxinus excelsior</i> , <i>Picea sitchensis</i> and <i>Pinus</i> sp.



Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
WS1	n/a	<i>Crataegus monogyna</i> , <i>Prunus spinosa</i> and <i>Ulex europaeus</i>
Keeraun Bóithrín to Knocknafroska and the N59 Link Road Ch. 5+850 – Ch. 7+550 and the N59 Link Road Site Habitats between EC21, EC23, EC25 and EC63, and between EC25 and EC29 along the N59 Link Road		
BL1	n/a	<i>Asplenium trichomanes</i> , <i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Hedera helix</i> and <i>Rubus fruticosus</i> agg.
ED2	n/a	Beginning to recolonise but vegetation cover is considered less than 50%. Species present include early colonisers – <i>Epilobium</i> spp., <i>Fallopia japonica</i> , <i>Matricaria discoidea</i> , <i>Plantago major</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg. and <i>Tussilago farfara</i>
ED3	n/a	<i>Arrhenatherum elatius</i> , <i>Bellis perennis</i> , <i>Centaurea nigra</i> , <i>Dactylis glomerata</i> , <i>Leucanthemum vulgare</i> , <i>Plantago lanceolata</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex</i> spp., <i>Senecio jacobaea</i> , <i>Tussilago farfara</i> and <i>Ulex europaeus</i>
FW4	n/a	<i>Angelica sylvestris</i> , <i>Potamogeton polygonifolius</i> , <i>Ranunculus flammula</i> and <i>Ranunculus repens</i>
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> ., <i>Lolium perenne</i> , <i>Persicaria maculosa</i> , <i>Plantago lanceolata</i> , <i>Poa annua</i> , <i>Polygonum aviculare</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Trifolium repens</i> and <i>Urtica dioica</i>
GA1/GS4	n/a	<i>Agrostis stolonifera</i> , <i>Cirsium</i> sp., <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Trifolium pratense</i> and <i>Urtica dioica</i>
GA2	n/a	<i>Arrhenatherum elatius</i> , <i>Bellis perennis</i> , <i>Dactylis glomerata</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Ranunculus repens</i> and <i>Trifolium repens</i>
GS1	n/a	<i>Achillea millefolium</i> , <i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Centaurea nigra</i> , <i>Cynosurus cristatus</i> , <i>Holcus lanatus</i> , <i>Lotus corniculatus</i> , <i>Prunella vulgaris</i> , <i>Trifolium pratense</i> and <i>Trifolium repens</i>
GS2	n/a	<i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Festuca rubra</i> , <i>Lolium perenne</i> , <i>Lotus corniculatus</i> , <i>Rumex acetosa</i> , <i>Rumex obtusifolius</i> , <i>Trifolium pratense</i> , <i>Trifolium repens</i> and <i>Veronica chamaedrys</i>
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Centaurea nigra</i> , <i>Cirsium palustre</i> , <i>Filipendula ulmaria</i> , <i>Galium palustre</i> , <i>Holcus lanatus</i> , <i>Juncus articulatus</i> , <i>Juncus effusus</i> , <i>Lotus pedunculatus</i> , <i>Lythrum salicaria</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> and <i>Senecio jacobaea</i>
WD1	n/a	<i>Acer pseudoplatanus</i> , <i>Aesculus pseudoplatanus</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> and <i>Hedera helix</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
WL1	n/a	Dominated by <i>Rubus fruticosus</i> agg. <i>Hedera helix</i> also common. <i>Crataegus monogyna</i> is present and has matured to form small shrubs/trees along the hedgerow. Occasional <i>Fraxinus excelsior</i> .
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Fraxinus excelsior</i> and <i>Populus</i> spp.
WS1	n/a	<i>Acer pseudoplatanus</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Fraxinus excelsior</i> , <i>Fuchsia magellanica</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rhododendron ponticum</i> , <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> , <i>Sorbus</i> sp., <i>Ulex europaeus</i> (D and <i>Urtica dioica</i>
Upper Dangan/NUIG Ch. 8+150 – Ch. 9+300 Habitats between EC25 and the west bank of River Corrib (excluding EC30)		
BL1	n/a	<i>Crataegus monogyna</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> and <i>Rubus fruticosus</i> agg.
ED2	n/a	<i>Bellis perennis</i> , <i>Lolium perenne</i> and <i>Taraxacum officinale</i> agg.
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Bellis perennis</i> , <i>Cirsium</i> sp., <i>Holcus lanatus</i> , <i>Lathyrus pratensis</i> , <i>Lolium perenne</i> , <i>Plantago major</i> , <i>Ranunculus repens</i> , <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg. and <i>Urtica dioica</i>
GA2	n/a	<i>Bellis perennis</i> , <i>Cirsium arvense</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Lolium</i> spp., <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Senecio vulgaris</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Vicia cracca</i>
GS1	n/a	A more species poor variant of calcareous/neutral grassland supported the following species: <i>Centaurea nigra</i> , <i>Dactylis glomerata</i> , <i>Leontodon hispidus</i> , <i>Lolium perenne</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> and <i>Urtica dioica</i>
	n/a	Species-rich variant of calcareous grassland which did not meet criteria for EU Annex I habitat. Species recorded include <i>Anthoxanthum odoratum</i> , <i>Calliargonella cuspidata</i> , <i>Holcus lanatus</i> , <i>Briza media</i> , <i>Festuca ovina</i> , <i>Linum catharticum</i> , <i>Trifolium pratense</i> , <i>Homalothecium lutescens</i> , <i>Lathyrus pratensis</i> , <i>Lolium perenne</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Carex flacca</i> , <i>Cerastium fontanum</i> , <i>Euphrasia</i> spp., <i>Plantago lanceolata</i> , <i>Ranunculus acris</i> , <i>Agrostis stolonifera</i> , <i>Alchemilla mollis</i> , <i>Carex panacea</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Festuca</i> spp., <i>Leontodon autumnalis</i> , <i>Leontodon hispidus</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Rumex crispus</i> , <i>Taraxacum officinale</i> agg. and <i>Trifolium repens</i> <b>UD-NUIG R1</b>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Crataegus monogyna</i> , <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Heracleum sphondylium</i> , <i>Holcus lanatus</i> , <i>Lathyrus pratensis</i> , <i>Plantago lanceolata</i> , <i>Prunella vulgaris</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Rumex</i> spp., <i>Salix cinerea</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> and <i>Urtica dioica</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
GS4	n/a	Species-poor <i>Juncus effusus</i> wet grassland with <i>Agrostis stolonifera</i> , <i>Cirsium palustre</i> , <i>Galium palustre</i> , <i>Holcus lanatus</i> , <i>Potentilla anserina</i> , <i>Ranunculus acris</i> and <i>Ranunculus repens</i>
GS4	n/a	A more species rich wet grassland variant included <i>Agrostis stolonifera</i> , <i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Epilobium montanum</i> , <i>Festuca rubra</i> , <i>Filipendula ulmaria</i> , <i>Holcus lanatus</i> , <i>Iris pseudacorus</i> , <i>Juncus articulatus</i> , <i>Juncus effusus</i> , <i>Lolium perenne</i> , <i>Lythrum salicaria</i> , <i>Phragmites australis</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rhinanthus minor</i> , <i>Rubus fruticosus</i> agg., <i>Rumex acetosa</i> , <i>Rumex</i> spp., <i>Succisa pratensis</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> , <i>Urtica dioica</i> and <i>Vicia cracca</i> .
WD1	n/a	<i>Acer pseudoplatanus</i> , <i>Aesculus hippocastanum</i> , <i>Anthriscus sylvestris</i> , <i>Arrhenatherum elatius</i> , <i>Brachypodium sylvaticum</i> , <i>Crataegus monogyna</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Geranium robertianum</i> , <i>Geum urbanum</i> , <i>Hedera helix</i> , <i>Heracleum sphondylium</i> , <i>Pinus</i> spp., <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> , <i>Tilia cordata</i> , <i>Torilis japonica</i> , <i>Ulmus glabra</i> and <i>Vicia sepium</i>
WL1	n/a	<i>Acer pseudoplatanus</i> , <i>Crataegus monogyna</i> , <i>Cupressus x leylandii</i> , <i>Epilobium ciliatum</i> , <i>Escallonia</i> spp., <i>Fagus purpurea</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Fuchsia magellanica</i> , <i>Geranium robertianum</i> , <i>Griselinia</i> sp., <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Plantago lanceolata</i> , <i>Prunus laurocerasus</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Rumex obtusifolius</i> , <i>Ulmus glabra</i> , <i>Urtica dioica</i> and <i>Vicia</i> sp.
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Aesculus hippocastanum</i> , <i>Alnus glutinosa</i> , <i>Betula</i> sp., <i>Cornus</i> sp., <i>Crataegus monogyna</i> , <i>Cupressus x Leylandii</i> , <i>Fagus sylvatica</i> , <i>Fallopia japonica</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Prunus lusitanica</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Sorbus</i> sp., <i>Symphoricarpos albus</i> , <i>Tilia cordata</i> and <i>Urtica dioica</i>
WN2	n/a	<i>Hedera helix</i> , <i>Fraxinus excelsior</i> , <i>Rubus fruticosus</i> agg., <i>Acer pseudoplatanus</i> , <i>Crataegus monogyna</i> , <i>Ilex aquifolium</i> , <i>Phyllitis scolopendrium</i> and <i>Thuidium</i> sp. <b>UD-NUIG R3</b>
WS1	n/a	<i>Betula pubescens</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Epilobium hirsutum</i> , <i>Fraxinus excelsior</i> , <i>Galium aparine</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> , <i>Salix cinerea</i> subsp. <i>oleifolia</i> , <i>Ulex europaeus</i> and <i>Urtica dioica</i>
WN6	*91E0	<i>Ranunculus repens</i> , <i>Hedera helix</i> , <i>Salix cinerea</i> , <i>Scrophularia auriculata</i> , <i>Alnus glutinosa</i> , <i>Urtica dioica</i> , <i>Crataegus monogyna</i> , <i>Filipendula ulmaria</i> , <i>Circaea lutetiana</i> , <i>Geranium robertianum</i> , <i>Heracleum sphondylium</i> , <i>Phyllitis scolopendrium</i> , <i>Taraxacum officinale</i> agg. and <i>Holcus lanatus</i> <b>UD-NUIG R5</b>



Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
GS1/GS2	n/a	Small area of grassland on rocky mound, waste land that is becoming encroached by bramble and scrub; Small patch of <i>Fallopia japonica</i> . <i>Angelica sylvestris</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Epipactis helleborine</i> , <i>Festuca rubra</i> , <i>Festuca vivipara</i> , <i>Homalothecium sericeum</i> , <i>Hylocomium splendens</i> , <i>Lathyrus pratensis</i> , <i>Leontodon hispidus</i> , <i>Leucanthemum vulgare</i> , <i>Linum catharticum</i> , <i>Lotus corniculatus</i> , <i>Lythrum salicaria</i> , <i>Petasites fragrans</i> , <i>Pimpinella major</i> , <i>Pimpinella saxifraga</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Potentilla reptans</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Senecio jacobaea</i> , <i>Trifolium dubium</i> , <i>Trifolium pratense</i> and <i>Vicia cracca</i>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Centaurea nigra</i> , <i>Dactylis glomerata</i> , <i>Epilobium montanum</i> , <i>Filipendula ulmaria</i> , <i>Fraxinus excelsior</i> , <i>Lathyrus pratensis</i> , <i>Leucanthemum vulgare</i> , <i>Plantago lanceolata</i> , <i>Prunus spinosa</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex crispus</i> , <i>Salix cinerea</i> , <i>Senecio jacobaea</i> , <i>Tussilago farfara</i> and <i>Ulex europaeus</i>
HD1	n/a	<i>Dactylis glomerata</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> and <i>Rubus fruticosus</i> agg.
WL1	n/a	<i>Calystegia sepium</i> , <i>Cirsium arvense</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Euonymus europaeus</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Lathyrus pratensis</i> , <i>Lonicera periclymenum</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Quercus robur</i> , <i>Rosa canina</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Sorbus aria</i> , <i>Urtica dioica</i> and <i>Viburnum opulus</i>
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus glutinosa</i> , <i>Alnus incana</i> , <i>Cupressus x leylandii</i> , <i>Fraxinus excelsior</i> , <i>Populus</i> sp., <i>Prunus lusitanica</i> , <i>Salix cinerea</i> subsp. <i>oleifolia</i> and <i>Tilia cordata</i>
WS1	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus glutinosa</i> , <i>Alnus incana</i> , <i>Betula pubescens</i> , <i>Buddleja davidii</i> , <i>Buxus sempervirens</i> , <i>Calystegia sepium</i> , <i>Centaurea nigra</i> , <i>Corylus avellana</i> , <i>Cotoneaster</i> spp., <i>Crataegus monogyna</i> , <i>Euonymus europaeus</i> , <i>Fallopia japonica</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Leucanthemum vulgare</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Salix viminalis</i> , <i>Sorbus aria</i> and <i>Ulex europaeus</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
Ballinooly to Parkmore Industrial Estate/Galway Racecourse Ch. 12+550 – Ch. 14+400 Habitats between EC39, EC42, EC44, EC47, EC49 and the Parkmore Industrial Estate/Galway Racecourse		
ED3	n/a	<i>Achillea millefolium</i> , <i>Anagallis arvensis</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Bellis perennis</i> , <i>Blackstonia perfoliata</i> , <i>Buddleja davidii</i> , <i>Carex flacca</i> , <i>Centaurea nigra</i> , <i>Centaureum erythraea</i> , <i>Crepis</i> sp., <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Dactylorhiza fuchsia</i> , <i>Epilobium ciliatum</i> , <i>Epilobium montanum</i> , <i>Epilobium parviflorum</i> , <i>Euphrasia officinalis</i> agg., <i>Festuca rubra</i> , <i>Hedera helix</i> , <i>Holcus lanatus</i> , <i>Hypericum perforatum</i> , <i>Leucanthemum vulgare</i> , <i>Linum catharticum</i> , <i>Lotus corniculatus</i> , <i>Medicago lupulina</i> , <i>Odontites vernus</i> , <i>Plantago lanceolata</i> , <i>Poa pratensis</i> , <i>Potentilla anserina</i> , <i>Potentilla reptans</i> , <i>Primula veris</i> , <i>Prunella vulgaris</i> , <i>Rubus fruticosus</i> agg., <i>Rumex</i> spp., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Scrophularia auriculata</i> , <i>Senecio jacobaea</i> , <i>Sonchus asper</i> , <i>Stachys palustris</i> , <i>Tussilago farfara</i> and <i>Veronica chamaedrys</i>
GA1	n/a	<i>Achillea millefolium</i> , <i>Bellis perennis</i> , <i>Cerastium fontanum</i> , <i>Cerastium glomeratum</i> , <i>Cirsium arvense</i> , <i>Cirsium</i> spp., <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Elytrigia repens</i> , <i>Heracleum sphondylium</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Poa trivialis</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Urtica dioica</i>
GA2	n/a	<i>Achillea millefolium</i> , <i>Bellis perennis</i> , <i>Cerastium fontanum</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Ranunculus repens</i> , <i>Taraxacum officinale</i> agg. and <i>Trifolium pratense</i>
GS1	n/a	<i>Bellis perennis</i> , <i>Cerastium fontanum</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Heracleum sphondylium</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Lolium</i> spp., <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Ranunculus repens</i> , <i>Rumex</i> spp., <i>Taraxacum officinale</i> agg. and <i>Urtica dioica</i>
GS2	n/a	Species-poor variant. <i>Arrhenatherum elatius</i> , <i>Agrostis capillaris</i> , <i>Anthoxanthum odoratum</i> , <i>Holcus lanatus</i> , <i>Phleum pratense</i> , <i>Dactylis glomerata</i> , <i>Plantago lanceolata</i> , <i>Cirsium arvense</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Cerastium fontanum</i> , <i>Leucanthemum vulgare</i> , <i>Epilobium ciliatum</i> , <i>Potentilla anserina</i> , <i>Scorzoneroides autumnalis</i> and <i>Taraxacum officinale</i> agg.
	n/a	More species-rich variant. Species include: <i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Alopecurus pratensis</i> , <i>Anthoxanthum odoratum</i> , <i>Arrhenatherum elatius</i> , <i>Calystegia sepium</i> , <i>Centaurea nigra</i> , <i>Cerastium fontanum</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Heracleum sphondylium</i> , <i>Holcus lanatus</i> , <i>Phleum pratense</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Ranunculus acris</i> , <i>Rumex</i> spp., <i>Scorzoneroides autumnalis</i> , <i>Trifolium repens</i> and <i>Urtica dioica</i>

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
GS4	n/a	<i>Calliergonella cuspidate</i> , <i>Carex disticha</i> , <i>Carex flacca</i> , <i>Carex hirta</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Equisetum arvense</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Juncus inflexus</i> , <i>Lathyrus pratensis</i> , <i>Persicaria amphibia</i> , <i>Phleum pratense</i> , <i>Plantago lanceolata</i> , <i>Poa trivialis</i> , <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Scorzoneroides autumnalis</i> , <i>Taraxacum officinale</i> agg. and <i>Trifolium dubium</i>
WD1	n/a	<i>Acer pseudoplatanus</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Rubus fruticosus</i> agg. and <i>Urtica dioica</i>
WL1	n/a	<i>Alnus incana</i> , <i>Anthriscus sylvestris</i> , <i>Arrhenatherum elatius</i> , <i>Asplenium scolopendrium</i> , <i>Brachypodium sylvaticum</i> , <i>Calystegia sepium</i> , <i>Circaea lutetiana</i> , <i>Convolvulus arvensis</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Dryopteris filix-mas</i> , <i>Euonymus europaeus</i> , <i>Fraxinus excelsior</i> , <i>Fuchsia magellanica</i> , <i>Galium aparine</i> , <i>Geranium robertianum</i> , <i>Geum urbanum</i> , <i>Hedera helix</i> , <i>Heracleum sphondylium</i> , <i>Lathyrus pratensis</i> , <i>Lonicera periclymenum</i> , <i>Prunus domestica</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Ribes</i> spp., <i>Rosa canina</i> , <i>Rosa</i> spp., <i>Rubus fruticosus</i> agg., <i>Rumex sanguineus</i> , <i>Salix cinerea</i> , <i>Salix</i> spp., <i>Sambucus nigra</i> , <i>Urtica dioica</i> , <i>Vicia sepium</i> and <i>X Cuprocypris leylandii</i>
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus incana</i> , <i>Betula</i> sp., <i>Cornus</i> sp., <i>Cupressus x leylandii</i> , <i>Lonicera nitida</i> , <i>Picea glauca</i> , <i>Pinus radiata</i> and <i>Sorbus aucuparia</i>
WN2	n/a	<i>Acer pseudoplatanus</i> , <i>Corylus avellana</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> and <i>Salix cinerea</i>
WS1	n/a	<i>Alnus glutinosa</i> , <i>Alnus incana</i> , <i>Arrhenatherum elatius</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Fraxinus excelsior</i> , <i>Hebe</i> sp., <i>Hedera helix</i> , <i>Plantago lanceolata</i> , <i>Prunus laurocerasus</i> , <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> and <i>Sorbaria</i> sp.
Galway Racecourse to Ardaun Ch. 14+400 – Ch. 18+350 and along the R446 tie-in Habitats between EC56, EC57 and EC59		
ED3	n/a	<i>Buddleja davidii</i> , <i>Hypericum</i> spp., <i>Leucanthemum vulgare</i> , <i>Reseda luteola</i> and <i>Senecio jacobaea</i>
GA1	n/a	<i>Bellis perennis</i> , <i>Cirsium arvense</i> , <i>Cirsium vulgare</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Poa annua</i> , <i>Potentilla anserina</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Urtica dioica</i>
GA2	n/a	<i>Bellis perennis</i> , <i>Cirsium arvense</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Lathyrus pratensis</i> , <i>Lolium</i> spp., <i>Plantago lanceolata</i> , <i>Poa annua</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Senecio jacobaea</i> , <i>Senecio vulgaris</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium repens</i> and <i>Trifolium</i> spp.

Habitat code <sup>4</sup>	Annex I habitat <sup>5</sup>	Key species <sup>6</sup>
GS1	n/a	<i>Achillea millefolium</i> , <i>Arctium lappa</i> , <i>Bellis perennis</i> , <i>Centaurea nigra</i> , <i>Cerastium fontanum</i> , <i>Cirsium arvense</i> , <i>Cirsium vulgare</i> , <i>Epilobium hirsutum</i> , <i>Holcus lanatus</i> , <i>Leucanthemum vulgare</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> , <i>Trifolium repens</i> , <i>Urtica dioica</i> and <i>Veronica montana</i>
GS2	n/a	<i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Dactylis glomerata</i> , <i>Epilobium parviflorum</i> , <i>Epilobium</i> sp., <i>Festuca rubra</i> , <i>Heracleum sphondylium</i> , <i>Plantago lanceolata</i> , <i>Pteridium aquilinum</i> , <i>Ranunculus repens</i> , <i>Rubus fruticosus</i> agg., <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> , <i>Ulex europaeus</i> , <i>Urtica dioica</i> , <i>Vicia cracca</i> and <i>Vicia sepium</i>
WD1/ WS1	n/a	<i>Acer pseudoplatanus</i> , <i>Arrhenatherum elatius</i> , <i>Arum maculatum</i> , <i>Asplenium scolopendrium</i> , <i>Athyrium filix-femina</i> , <i>Brachypodium sylvaticum</i> , <i>Chamaecyparis lawsoniana</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Dactylis glomerata</i> , <i>Dryopteris filix-mas</i> , <i>Kindbergia praelonga</i> , <i>Eurhynchium striatum</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Hedera hibernica</i> , <i>Picea</i> sp., <i>Pinus sylvestris</i> , <i>Polystichum setiferum</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Senecio jacobaea</i> , <i>Veronica hederacea</i> , <i>Veronica montana</i> and <i>Viola canina</i>
WL1	n/a	<i>Cornus</i> sp., <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Escallonia</i> sp., <i>Fagus sylvatica</i> , <i>Fuchsii</i> sp., <i>Hedera helix</i> , <i>Prunus laurocerasus</i> , <i>Prunus spinosa</i> , <i>Rubus fruticosus</i> agg., <i>Salix caprea</i> , <i>Salix cinerea</i> , <i>Sorbaria</i> , <i>Sorbus aucuparia</i> and <i>Ulex europaeus</i>
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus glutinosa</i> , <i>Alnus incana</i> , <i>Betula pubescens</i> , <i>Cornus</i> spp., <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Prunus spinosa</i> , <i>Rosa rugosa</i> , <i>Rubus fruticosus</i> agg., <i>Sorbus aucuparia</i> , <i>Symphoricarpos albus</i> and <i>Viburnum opulus</i>
WN2	n/a	<i>Corylus avellana</i> , <i>Cupressus x Leylandii</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Ilex aquifolium</i> and <i>Sorb aucuparia</i>
WS1	n/a	<i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Trifolium pratense</i> and <i>Trifolium repens</i>



### **A.8.19.3**

---

## Habitat Species Lists – Areas between the River Corrib and Lackagh Quarry

**Table A.8.19.3** presents the species lists for those habitat areas along the proposed road development that lie between the River Corrib and Lackagh Quarry. This excludes habitat areas within the ecological sites EC33 and EC36 (which are described separately in **Appendix A.8.19.1**). It does not include species lists for the residential habitat category (see **Section 8.3.4.34** of **Chapter 8, Biodiversity**) and it also does not include species lists for the aquatic habitats, which are described in full in **Section 8.3.4** of **Chapter 8, Biodiversity** and in **Appendix A.8.20 (Results of the 2014 N6 Galway City Transport Project Aquatic Habitat Surveys)**.

Where relevés were taken in the field, the reference codes are provided in the Key Species column (e.g. EC03 R1). These relevé codes refer to the full species lists which are provided in **Appendix A.8.19.4**.

**Table A.8.19.3: Habitat Species Lists Relating to areas Affected by the Proposed Road Development between the Ecological Sites (excluding habitats within the Lough Corrib cSAC Survey Area**

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
River Corrib to Lackagh Quarry and Surrounding areas (which includes part of Lough Corrib cSAC) Ch. 9+400 – Ch. 11+400 Habitats between the River Corrib and Lackagh Quarry		
BL1	n/a	<i>Crataegus monogyna</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Prunus spinosa</i> and <i>Rubus fruticosus</i> agg.
ED2	n/a	<i>Lolium perenne</i> and <i>Potentilla anserina</i>
ED3	n/a	<i>Agrostis stolonifera</i> , <i>Lolium perenne</i> , <i>Plantago major</i> , <i>Potentilla anserina</i> , <i>Ranunculus repens</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> , <i>Stellaria media</i> , <i>Trifolium pratense</i> and <i>Veronica persica</i>
ER2	*8240 LPE	The most frequently recorded species across these 39 relevés were: <i>Anthoxanthum odoratum</i> , <i>Asperula cynanchica</i> , <i>Asplenium ruta-muraria</i> , <i>Brachythecium rutabulum</i> , <i>Briza media</i> , <i>Bryum capillare</i> , <i>Calliargonella cuspidata</i> , <i>Carex flacca</i> , <i>Carlina vulgaris</i> , <i>Centaurea nigra</i> , <i>Ceterach officinarum</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Ctenidium molluscum</i> , <i>Dicranum scoparium</i> , <i>Encalypta streptocarpa</i> , <i>Epilobium montanum</i> , <i>Epilobium parviflorum</i> , <i>Euphrasia salisburgensis</i> , <i>Festuca ovina</i> , <i>Festuca rubra</i> , <i>Fissidens dubius</i> , <i>Fragaria vesca</i> , <i>Fraxinus excelsior</i> , <i>Galium verum</i> , <i>Geranium robertianum</i> , <i>Grimmia pulvinata</i> , <i>Hedera helix</i> , <i>Homalothecium lutescens</i> , <i>Homalothecium sericeum</i> , <i>Hypericum pulchrum</i> , <i>Hypochaeris radicata</i> , <i>Ilex aquifolium</i> , <i>Koeleria macrantha</i> , <i>Lapsana communis</i> , <i>Leontodon hispidus</i> , <i>Leucanthemum vulgare</i> , <i>Linum catharticum</i> , <i>Lotus corniculatus</i> , <i>Mycelis muralis</i> , <i>Neckera complanata</i> , <i>Neckera crispa</i> , <i>Phyllitis scolopendrium</i> , <i>Pilosella</i>

<sup>7</sup> \*8240 (E) = exposed limestone pavement; \*8240 (W) = wooded limestone pavement

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
		<i>officinatum</i> , <i>Plantago lanceolata</i> , <i>Plantago maritima</i> , <i>Polygala vulgaris</i> , <i>Potentilla erecta</i> , <i>Prunus spinosa</i> , <i>Ranunculus acris</i> , <i>Rosa spinosissima</i> , <i>Rubus fruticosus</i> agg., <i>Rubus idaeus</i> , <i>Scleropodium purum</i> , <i>Senecio jacobaea</i> , <i>Sesleria caerulea</i> , <i>Solidago virgaurea</i> , <i>Sonchus asper</i> , <i>Sorbus aria</i> , <i>Succisa pratensis</i> , <i>Taraxacum officinale</i> agg., <i>Teucrium scorodonia</i> , <i>Thymus polytrichus</i> , <i>Tortella nitida</i> , <i>Tortella tortuosa</i> , <i>Trifolium pratense</i> and <i>Veronica chamaedrys</i> .  Relevés BEC 115, BEC 118, BEC 343 and BEC 345
FL6	*3180 <i>Potentilla anserina</i> - <i>Carex nigra</i> vegetation community (Waldren, 2015, Ed.).	Typical FL6/ GM1 vegetation with <i>Agrostis stolonifera</i> , <i>Apium inundatum</i> , <i>Caltha palustris</i> , <i>Cinclidotus fontinaloides</i> , <i>Eleocharis palustris</i> , <i>Equisetum palustre</i> , <i>Filipendula ulmaria</i> , <i>Glyceria fluitans</i> , <i>Mentha aquatica</i> , <i>Myosotis aquatica</i> , <i>Persicaria amphibia</i> , <i>Persicaria maculata</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus repens</i> , <i>Rumex crispus</i> , <i>Rumex obtusifolius</i> , <i>Senecio aquaticus</i> , <i>Urtica dioica</i> and <i>Veronica catenata</i>
FS2	6430	<i>Epilobium palustre</i> , <i>Epilobium hirsutum</i> , <i>Galium palustre</i> , <i>Valeriana dioica</i> , <i>Menyanthes trifoliata</i> , <i>Phragmites australis</i> , <i>Calliergonella cuspidata</i> , <i>Equisetum fluviatile</i> , <i>Carex rostrata</i> , <i>Ranunculus acris</i> , <i>Prunus spinosa</i> , <i>Iris pseudacorus</i> , <i>Holcus lanatus</i> and <i>Agrostis stolonifera</i> .  EC36 R4
GA1	n/a	<i>Agrostis stolonifera</i> , <i>Arrhenatherum elatius</i> , <i>Centaurea nigra</i> , <i>Cirsium arvense</i> , <i>Crepis</i> sp., <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Odontites vernus</i> , <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rumex obtusifolius</i> , <i>Rumex</i> spp., <i>Senecio jacobaea</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> , <i>Trifolium repens</i> and <i>Veronica</i> sp.
GA2	n/a	<i>Bellis perennis</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Lolium perenne</i> , <i>Plantago lanceolata</i> , <i>Rumex acetosa</i> , <i>Rumex obtusifolius</i> , <i>Taraxacum officinale</i> agg. and <i>Trifolium repens</i>
GS1	n/a (GL2C, GL3B, GL3C, and GL3E)	The following species were commonly encountered in non-Annex I calcareous grassland: <i>Agrostis capillaris</i> , <i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Brachythecium rutabulum</i> , <i>Calliergonella cuspidata</i> , <i>Carex flacca</i> , <i>Centaurea nigra</i> , <i>Cerastium fontanum</i> , <i>Cirsium arvense</i> , <i>Cynosurus cristatus</i> , <i>Dactylis glomerata</i> , <i>Daucus carota</i> , <i>Festuca rubra</i> , <i>Holcus lanatus</i> , <i>Hypochaeris radicata</i> , <i>Leontodon autumnalis</i> , <i>Lolium perenne</i> , <i>Lotus corniculatus</i> , <i>Odontites vernus</i> , <i>Plagiomnium undulatum</i> , <i>Plantago lanceolata</i> , <i>Plantago major</i> , <i>Potentilla anserina</i> , <i>Prunella vulgaris</i> , <i>Ranunculus acris</i> , <i>Ranunculus repens</i> , <i>Rhinanthus minor</i> , <i>Rubus fruticosus</i> agg., <i>Rumex acetosa</i> , <i>Taraxacum officinale</i> agg., <i>Trifolium pratense</i> , <i>Trifolium repens</i> and <i>Veronica chamaedrys</i>  Relevés BEC 175, BEC 258, BEC 363, RC-LQ R1, RC-LQ R6, RC-LQ R7 and RC-LQ R8.

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
GS1	6210	<p>The following species were commonly encountered in calcareous grassland which corresponded with the Annex I Calcareous grassland [6210] habitat:</p> <p><i>Achillea millefolium, Agrostis capillaris, Anthoxanthum odoratum, Brachythecium rutabulum, Briza media, Calliergonella cuspidata, Campanula rotundifolia, Carex caryophylla, Carex flacca, Centaurea nigra, Ctenidium molluscum, Cynosurus cristatus, Dactylis glomerata, Daucus carota, Festuca ovina, Festuca rubra, Galium verum, Helictotrichon pubescens, Hypericum pulchrum, Hypochaeris radicata, Leontodon autumnalis, Leontodon hispidus, Leucanthemum vulgare, Linum catharticum, Lotus corniculatus, Pilosella officinarum, Plantago lanceolata, Plantago maritima, Polygala vulgaris, Potentilla erecta, Prunella vulgaris, Ranunculus acris, Rosa spinosissima, Rubus fruticosus agg., Scleropodium purum, Sesleria caerulea, Solidago virgaurea, Succisa pratensis, Taraxacum officinale agg., Thymus polytrichus, Trifolium pratense and Trifolium repens</i></p> <p><b>Relevé BEC 7</b></p>
GS1	*6210	<p>The following species were commonly encountered in calcareous grassland which corresponded with the priority Annex I calcareous grassland [*6210] habitat:</p> <p><i>Agrostis capillaris, Anthoxanthum odoratum, Anthyllis vulneraria, Asperula cynanchica, Briza media, Calliergonella cuspidata, Campanula rotundifolia, Carex caryophylla, Carex flacca, Carex panicea, Centaurea nigra, Ctenidium molluscum, Cynosurus cristatus, Dactylis glomerata, Dactylorhiza fuchsii, Danthonia decumbens, Daucus carota, Euphrasia officinalis ag., Festuca rubra, Galium verum, Gymnadenia conopsea, Helictotrichon pubescens, Holcus lanatus, Hypericum pulchrum, Koeleria macrantha, Leontodon hispidus, Leucanthemum vulgare, Linum catharticum, Listera ovata, Lotus corniculatus, Pilosella officinarum, Plantago lanceolata, Plantago maritima, Polygala vulgaris, Potentilla erecta, Prunella vulgaris, Rhinanthus minor, Rhytidadelphus squarrosus, Rosa spinosissima, Scleropodium purum, Sesleria caerulea, Solidago virgaurea, Succisa pratensis, Thymus polytrichus, Trifolium pratense and Trifolium repens</i></p> <p><b>Relevés BEC 15, BEC 17 and BEC 102.</b></p>
GS2	n/a	<p><i>Agrostis stolonifera, Arrhenatherum elatius, Briza media, Calystegia sepium, Carex flacca, Carex hirta, Centaurea nigra, Chamerion angustifolium, Cirsium arvense, Cirsium vulgare, Crepis sp., Cynosurus cristatus, Dactylis glomerata, Epilobium ciliatum, Epilobium hirsutum, Equisetum fluviatile, Festuca rubra, Filipendula ulmaria, Galium aparine, Heracleum sphondylium, Holcus lanatus, Juncus inflexus, Lathyrus pratensis, Leontodon hispidus, Lolium perenne, Plantago lanceolata, Poa annua, Prunella vulgaris, Pteridium aquilinum, Ranunculus acris, Ranunculus repens, Rubus fruticosus agg., Rumex acetosa, Rumex obtusifolius, Sonchus asper, Stellaria media, Succisa pratensis, Taraxacum officinale agg., Trifolium pratense, Trifolium repens, Urtica dioica and Vicia sepium</i></p>

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
GS4	n/a	<i>Agrostis stolonifera</i> , <i>Anthoxanthum odoratum</i> , <i>Calliergonella cuspidata</i> , <i>Cirsium arvense</i> , <i>Cirsium palustre</i> , <i>Cynosurus cristatus</i> , <i>Epilobium montanum</i> , <i>Filipendula ulmaria</i> , <i>Galium palustre</i> , <i>Juncus articulatus</i> , <i>Mentha aquatica</i> , <i>Molinia caerulea</i> , <i>Juncus effusus</i> , <i>Arrhenatherum elatius</i> , <i>Holcus lanatus</i> , <i>Lychnis flos-cuculi</i> , <i>Lythrum salicaria</i> , <i>Plantago lanceolata</i> , <i>Potentilla anserina</i> , <i>Potentilla erecta</i> , <i>Trifolium repens</i> , <i>Ranunculus flammula</i> , <i>Ranunculus repens</i> , <i>Ranunculus acris</i> , <i>Rumex obtusifolius</i> , <i>Senecio jacobaea</i> , <i>Trifolium pratense</i> and <i>Valeriana officinalis</i> <b>Relevé: RC-LQ R3</b>
WD1	n/a	The most frequently recorded species within this habitat were: <i>Acer pseudoplatanus</i> , <i>Arum maculatum</i> , <i>Circaea lutetiana</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Eurhynchium striatum</i> , <i>Fagus sylvatica</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Isoethecium alopecuroides</i> , <i>Isoethecium myosuroides</i> , <i>Kindbergia praelonga</i> , <i>Lejeunea cavifolia</i> , <i>Metzgeria furcata</i> , <i>Neckera complanata</i> , <i>Phyllitis scolopendrium</i> , <i>Radula complanata</i> , <i>Rhynchostegiella tenella</i> , <i>Rubus fruticosus</i> agg., <i>Thamnobryum alopecurum</i> , <i>Thuidium tamariscinum</i> , <i>Tortella tortuosa</i> and <i>Urtica dioica</i> , <b>Relevés: BEC 174, and RC-LQ R2</b>
WL1	n/a	<i>Calystegia sepium</i> , <i>Circaea lutetiana</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Euonymus europaeus</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Heracleum sphondylium</i> , <i>Ilex aquifolium</i> , <i>Lathyrus pratensis</i> , <i>Lonicera periclymenum</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Quercus robur</i> , <i>Rubus fruticosus</i> agg., <i>Sambucus nigra</i> , <i>Sorbus aria</i> , <i>Sorbus aucuparia</i> , <i>Urtica dioica</i> and <i>Viburnum opulus</i>
WL2	n/a	<i>Acer pseudoplatanus</i> , <i>Alnus glutinosa</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Fraxinus excelsior</i> , <i>Hedera helix</i> , <i>Ilex aquifolium</i> , <i>Prunus spinosa</i> , <i>Pteridium aquilinum</i> , <i>Rubus fruticosus</i> agg., <i>Salix cinerea</i> and <i>Sambucus nigra</i>
WN2	n/a (WL2A WL2E)	Oak-ash-hazel woodland in this area was typically either <i>Corylus avellana</i> or <i>Fraxinus excelsior</i> woodlands. The following species were recorded in areas that did not conform to any Annex I habitat type: <i>Acer pseudoplatanus</i> , <i>Arum maculatum</i> , <i>Brachypodium sylvaticum</i> , <i>Circaea lutetiana</i> , <i>Corylus avellana</i> , <i>Crataegus monogyna</i> , <i>Eurhynchium hians</i> , <i>Eurhynchium striatum</i> , <i>Fissidens taxifolius</i> , <i>Fraxinus excelsior</i> , <i>Frullania dilatata</i> , <i>Geranium robertianum</i> , <i>Geum urbanum</i> , <i>Hedera helix</i> , <i>Homalothecium sericeum</i> , <i>Hypnum resupinatum</i> , <i>Ilex aquifolium</i> , <i>Isoethecium alopecuroides</i> , <i>Isoethecium myosuroides</i> , <i>Kindbergia praelonga</i> , <i>Lonicera periclymenum</i> , <i>Metzgeria furcata</i> , <i>Neckera complanata</i> , <i>Phyllitis scolopendrium</i> , <i>Plagiomnium undulatum</i> , <i>Potentilla sterilis</i> , <i>Primula vulgaris</i> , <i>Prunus spinosa</i> , <i>Quercus robur</i> , <i>Radula complanata</i> , <i>Rhytidiadelphus triquetrus</i> , <i>Rubus fruticosus</i> agg., <i>Rumex sanguineus</i> , <i>Thamnobryum alopecurum</i> , <i>Thuidium tamariscinum</i> , <i>Ulota bruchii</i> , <i>Veronica chamaedrys</i> and <i>Viola</i> spp. <b>Relevés: BEC 139, BEC 336, BEC 340, BEC 366, BEC 404, RC-LQ R9, and RC-LQ R10.</b>

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
WN2	*8240 (LPW)	<p>Woodland which corresponded with the Annex I Limestone pavement [*8240] habitat included the following species:</p> <p><i>Asplenium trichomanes</i>, <i>Brachypodium sylvaticum</i>, <i>Carex flacca</i>, <i>Corylus avellana</i>, <i>Crataegus monogyna</i>, <i>Ctenidium molluscum</i>, <i>Euonymus europaeus</i>, <i>Eurhynchium striatum</i>, <i>Fissidens dubius</i>, <i>Fragaria vesca</i>, <i>Fraxinus excelsior</i>, <i>Frullania dilatata</i>, <i>Geranium robertianum</i>, <i>Hedera helix</i>, <i>Homalothecium sericeum</i>, <i>Ilex aquifolium</i>, <i>Isothecium alopecuroides</i>, <i>Kindbergia praelonga</i>, <i>Lonicera periclymenum</i>, <i>Metzgeria furcata</i>, <i>Neckera complanata</i>, <i>Neckera crispa</i>, <i>Phyllitis scolopendrium</i>, <i>Plagiomnium undulatum</i>, <i>Potentilla sterilis</i>, <i>Prunus spinosa</i>, <i>Pteridium aquilinum</i>, <i>Radula complanata</i>, <i>Rhytidiadelphus triquetrus</i>, <i>Rosa spinosissima</i>, <i>Rubus fruticosus</i> agg., <i>Scapania aspera</i>, <i>Scleropodium purum</i>, <i>Sesleria caerulea</i>, <i>Solidago virgaurea</i>, <i>Thamnobryum alopecurum</i>, <i>Thuidium tamariscinum</i>, <i>Tortella tortuosa</i>, <i>Ulotia bruchii</i>, <i>Ulotia crispa</i>, <i>Veronica chamaedrys</i> and <i>Viola</i> spp.</p> <p>Relevés BEC 9, BEC 42, BEC 120, BEC 121, BEC 144, BEC 145, BEC 240, BEC 246, BEC 333, BEC 335, and RC-LQ R4.</p>
WN6	*91E0 (WN6_3c)	<p><i>Agrostis stolonifera</i>, <i>Calliargonella cuspidatum</i>, <i>Corylus avellana</i>, <i>Crataegus monogyna</i>, <i>Epilobium hirsutum</i>, <i>Equisetum fluviatile</i>, <i>Eurhynchium striatum</i>, <i>Filipendula ulmaria</i>, <i>Fraxinus excelsior</i>, <i>Galium aparine</i>, <i>Galium palustre</i>, <i>Geranium robertianum</i>, <i>Hedera helix</i>, <i>Juncus effusus</i>, <i>Kindbergia praelonga</i>, <i>Lythrum salicaria</i>, <i>Prunus spinosa</i>, <i>Ranunculus repens</i>, <i>Rubus fruticosus</i> agg., <i>Rumex sanguineus</i>, <i>Salix cinerea</i> subsp. <i>oleifolia</i> and <i>Thamnobryum alopecurum</i></p>
WS1	n/a	<p>The following species were recorded within scrub habitat that does not meet the criteria for the Annex I Limestone pavement [*8240] habitat:</p> <p><i>Agrostis capillaris</i>, <i>Agrostis stolonifera</i>, <i>Anthoxanthum odoratum</i>, <i>Arrhenatherum elatius</i>, <i>Brachypodium sylvaticum</i>, <i>Calliargonella cuspidata</i>, <i>Centaurea nigra</i>, <i>Circaea lutetiana</i>, <i>Corylus avellana</i>, <i>Crataegus monogyna</i>, <i>Dactylis glomerata</i>, <i>Festuca rubra</i>, <i>Filipendula ulmaria</i>, <i>Fragaria vesca</i>, <i>Fraxinus excelsior</i>, <i>Geum urbanum</i>, <i>Hedera helix</i>, <i>Holcus lanatus</i>, <i>Ilex aquifolium</i>, <i>Juncus conglomeratus</i>, <i>Kindbergia praelonga</i>, <i>Molinia caerulea</i>, <i>Myrica gale</i>, <i>Odontites vernus</i>, <i>Oxyrrhynchium hians</i>, <i>Plagiomnium undulatum</i>, <i>Plantago lanceolata</i>, <i>Potentilla erecta</i>, <i>Potentilla palustris</i>, <i>Primula vulgaris</i>, <i>Prunus spinosa</i>, <i>Pteridium aquilinum</i>, <i>Radula complanata</i>, <i>Ranunculus repens</i>, <i>Rhytidiadelphus squarrosus</i>, <i>Rosa canina</i>, <i>Rubus fruticosus</i> agg., <i>Rumex sanguineus</i>, <i>Salix cinerea</i>, <i>Symphoricarpos albus</i>, <i>Thamnobryum alopecurum</i>, <i>Thuidium tamariscinum</i>, <i>Trifolium pratense</i>, <i>Trifolium repens</i>, <i>Ulex europaeus</i>, <i>Urtica dioica</i>, <i>Veronica montana</i> and <i>Vicia sepium</i></p>

Habitat code (Fossitt)	Annex I habitat code <sup>7</sup>	Key species
WS1	*8240 (LPE)	<p>The following species were typical of areas of scrub that corresponded with the Annex I Limestone pavement [*8240] habitat:</p> <p><i>Agrimonia eupatoria, Asplenium adiantum-nigrum, Brachythecium rutabulum, Calliergonella cuspidata, Campylium stellatum v. protensum, Cladonia pocillum, Didymodon insulanus, Dryopteris filix-mas, Epipactis atrorubens, Euphrasia salisburgensis, Euphrasia species, Fissidens taxifolius, Galium sternerii, Galium verum, Holcus lanatus, Homalothecium sericeum, Hylocomium brevirostre, Hypnum lacunosum, Hypochaeris radicata, Isothecium myosuroides, Lathyrus pratensis, Leontodon hispidus, Leontodon saxatilis, Plagiomnium species, Ranunculus acris, Schistidium apocarpum, Schistidium crassipilum, Trichostomum brachydontium, Trifolium pratense, Ulota bruchii, Veronica chamaedrys and Veronica montana</i></p> <p>Relevés BEC 8, BEC 18, BEC 35, BEC 148 and BEC 242</p>
WS2	n/a	<p><i>Anthriscus sylvestris, Corylus avellana, Crataegus monogyna, Fraxinus excelsior, Geranium robertianum, Hedera helix, Rubus fruticosus agg. and Salix spp.</i></p>

## **A.8.19.4**

---



## Habitat Relevé Data

This section presents the relevé data collected along and adjacent to the proposed route, as referenced in **Sections A.8.19.1 to A.8.19.3**.

The recorders referred to in the tables below are as follows:

JC	John Conaghan	JM	Jim Martin
KD	Kathy Duff	FON	Fionnuala O'Neill
EOS	Eamonn O'Sullivan	SB	Simon Barron
JF	Janice Fuller	FD	Fiona Devaney
JD	Joanne Denyer	ML	Maria Long
CC	Colm Clarke	KMN	Kate McNutt
KMOC	Kate-Marie O'Connor		

In some instances, the plant abundance scores are given as percentage cover, for others the plant abundance scores are as per the Domin Scale:

Cover	Domin Value
91–100%	10
76–90%	9
51–75%	8
34–50%	7
26–33%	6
11–25%	5
4–10%	4
<4% (many individuals)	3
<4% (several individuals)	2
<4% (few individuals)	1

Relevé code	EC03 R1
Date	28/07/2014
Recorder	JC
Species recording scale	Percentage
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis canina</i>	8
<i>Anthoxanthum odoratum</i>	3
<i>Calluna vulgaris</i>	30
<i>Carex binervis</i>	0.5
<i>Carex panicea</i>	8
<i>Danthonia decumbens</i>	3
<i>Erica cinerea</i>	18
<i>Molinia caerulea</i>	3
<i>Potentilla erecta</i>	18
<i>Succisa pratensis</i>	18
<i>Ulex gallii</i>	30

Relevé code	EC05 R1
Date	10/09/2014
Recorder	KD
Species recording scale	Percentage
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Percentage
<i>Arctostaphylos uva-ursi</i>	3
<i>Calluna vulgaris</i>	42
<i>Daboecia cantabrica</i>	3
<i>Erica cinerea</i>	3
<i>Erica tetralix</i>	3
<i>Hypnum cupressiforme.</i>	3
<i>Molinia caerulea</i>	18
<i>Potentilla erecta</i>	0.5
<i>Pteridium aquilinum</i>	0.3
<i>Rhytidiadelphus squarrosus</i>	3
<i>Ulex gallii</i>	42

Relevé code	EC09 R1
Date	16/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/GS4/HH1
Habitat - Annex I	6410
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.5
<i>Carex nigra</i>	3
<i>Carex panicea</i>	0.5
<i>Hylocomium splendens</i>	42
<i>Juncus articulatus</i>	30
<i>Juncus conglomeratus</i>	0.5

Relevé code	EC09 R1
Species	Percentage
<i>Molinia caerulea</i>	42
<i>Pleurozium schreberi</i>	42
<i>Potentilla erecta</i>	30
<i>Rhytidadelphus squarrosus</i>	18
<i>Sphagnum papillosum</i>	0.5
<i>Succisa pratensis</i>	8

Relevé code	EC09 R2
Date	16/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/GS4/HH1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis vinealis</i>	0.5
<i>Anthoxanthum odoratum</i>	.3
<i>Calluna vulgaris</i>	63
<i>Carex nigra</i>	0.5
<i>Carex panicea</i>	0.5
<i>Cladonia portentosa</i>	8
<i>Danthonia decumbens</i>	0.5
<i>Erica cinerea</i>	18
<i>Galium saxatile</i>	0.3
<i>Hylocomium splendens</i>	42
<i>Hypnum jutlandicum</i>	3
<i>Hypochaeris radicata</i>	0.3
<i>Lotus corniculatus</i>	0.3
<i>Molinia caerulea</i>	0.5
<i>Pleurozium schreberi</i>	42
<i>Racomitrium heterostichum</i>	0.5
<i>Succisa pratensis</i>	3

Relevé code	EC09 R3
Date	16/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3 /GS4 /HH1
Habitat - Annex I	4010
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.3
<i>Calluna vulgaris</i>	42
<i>Carex echinata</i>	18
<i>Erica tetralix</i>	8
<i>Eriophorum angustifolium</i>	3
<i>Hylocomium splendens</i>	30
<i>Juncus articulatus</i>	0.5
<i>Juncus effusus</i>	0.5
<i>Molinia caerulea</i>	18
<i>Narthecium ossifragum</i>	0.3

Relevé code	EC09 R3
Species	Percentage
<i>Pettigera</i> sp.	0.2
<i>Pleurozium schreberi</i>	63
<i>Potentilla erecta</i>	0.5
<i>Rhytidiadelphus squarrosus</i>	3
<i>Sphagnum capillifolium</i> s. <i>rubellum</i>	18
<i>Succisa pratensis</i>	8
<i>Thuidium tamariscinum</i>	0.3

Relevé code	EC09 R4
Date	17/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/GS4/HH1
Habitat - Annex I	6410
Species	Percentage
<i>Agrostis vinealis</i>	0.5
<i>Anagallis tenella</i>	8
<i>Aneura pinguis</i>	0.3
<i>Anthoxanthum odoratum</i>	18
<i>Bryum</i> sp.	0.3
<i>Calliergonella cuspidata</i>	0.3
<i>Carex echinata</i>	3
<i>Carex nigra</i>	3
<i>Carex panicea</i>	18
<i>Carex viridula</i>	0.5
<i>Cirsium dissectum</i>	18
<i>Danthonia decumbens</i>	0.5
<i>Euphrasia officinalis</i> ag.	0.3
<i>Filipendula ulmaria</i>	0.5
<i>Hypnum jutlandicum</i>	0.5
<i>Juncus articulatus</i>	0.5
<i>Leontodon autumnalis</i>	8
<i>Lophocolea bidentata</i>	0.3
<i>Luzula multiflora</i>	1
<i>Lythrum salicaria</i>	0.5
<i>Molinia caerulea</i>	8
<i>Mnium hornum</i>	0.3
<i>Nardus stricta</i>	18
<i>Pettigera</i> sp.	0.3
<i>Plantago lanceolata</i>	0.5
<i>Potentilla erecta</i>	3
<i>Prunella vulgaris</i>	3
<i>Ranunculus flammula</i>	8
<i>Ranunculus repens</i>	2
<i>Rhinanthus minor</i>	0.5
<i>Rhytidiadelphus squarrosus</i>	0.3
<i>Riccardia multifida</i>	0.3
<i>Scapania</i> sp.	0.3
<i>Scleropodium purum</i>	0.5
<i>Sphagnum denticulatum</i>	42
<i>Succisa pratensis</i>	18

Relevé code	EC09 R4
Species	Percentage
<i>Thuidium tamariscinum</i>	18

Relevé code	EC11 R1
Date	09/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Percentage
<i>Angelica sylvestris</i>	8
<i>Anthoxanthum odoratum</i>	3
<i>Calliergonella cuspidata</i>	8
<i>Carex echinata</i>	3
<i>Carex flacca</i>	8
<i>Centaurea nigra</i>	0.3
<i>Filipendula ulmaria</i>	18
<i>Galium palustre</i>	0.3
<i>Holcus lanatus</i>	0.3
<i>Hypericum</i> sp.	8
<i>Juncus articulatus</i>	30
<i>Juncus conglomeratus</i>	3
<i>Juncus effusus</i>	0.5
<i>Luzula multiflora</i>	0.3
<i>Lythrum salicaria</i>	18
<i>Mentha aquatica</i>	8
<i>Molinia caerulea</i>	8
<i>Plantago lanceolata</i>	0.5
<i>Potentilla erecta</i>	0.5
<i>Ranunculus acris</i>	0.5
<i>Diplophyllum albicans</i>	0.5

Relevé code	EC11 R2
Date	09/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	PB3/HH3
Habitat - Annex I	7150
Species	Percentage
<i>Calluna vulgaris</i>	3
<i>Cephalozia connivens</i>	0.3
<i>Drosera rotundifolia</i>	0.3
<i>Erica tetralix</i>	8
<i>Eriophorum angustifolium</i>	18
<i>Menyanthes trifoliata</i>	3
<i>Molinia caerulea</i>	8
<i>Myliia taylorii</i>	0.3
<i>Narthecium ossifragum</i>	3
<i>Odontoschisma sphagni</i>	0.3
<i>Rhynchospora alba</i>	30

Relevé code	EC11 R2
Species	Percentage
<i>Sphagnum cuspidatum</i>	42
<i>Sphagnum denticulatum</i>	18
<i>Sphagnum magellanicum</i>	30
<i>Sphagnum papillosum</i>	30
<i>Sphagnum tenellum</i>	3

Relevé code	EC11 R3
Date	09/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	PB3/HH3
Habitat - Annex I	*7130
Species	Percentage
<i>Aulacomnium palustre</i>	0.5
<i>Calluna vulgaris</i>	30
<i>Calypogeia muelleriana</i>	0.3
<i>Cephalozia connivens</i>	0.3
<i>Drosera rotundifolia</i>	0.3
<i>Erica tetralix</i>	8
<i>Eriophorum angustifolium</i>	30
<i>Eriophorum vaginatum</i>	18
<i>Hypnum jutlandicum</i>	18
<i>Kurzia</i> sp.	0.5
<i>Molinia caerulea</i>	30
<i>Narthecium ossifragum</i>	3
<i>Odontoschisma sphagni</i>	3
<i>Potentilla erecta</i>	0.5
<i>Sphagnum capillifolium</i> s. <i>rubellum</i>	42
<i>Sphagnum fallax</i>	0.3
<i>Sphagnum magellanicum</i>	8
<i>Sphagnum papillosum</i>	42
<i>Sphagnum subnitens</i>	0.3

Relevé code	EC11 R4
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/HH1/PB3/GS3 /ER1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis capillaris</i>	3
<i>Anthoxanthum odoratum</i>	0.3
<i>Calluna vulgaris</i>	42
<i>Campylopus flexuosus</i>	8
<i>Carex binervis</i>	18
<i>Carex panicea</i>	8
<i>Carex viridula</i>	0.3
<i>Cephalozia</i> sp.	0.3
<i>Cladonia portentosa</i>	30

Relevé code	EC11 R4
Species	Percentage
<i>Erica cinerea</i>	18
<i>Hylocomium splendens</i>	8
<i>Hypnum jutlandicum</i>	0.3
<i>Hypochaeris radicata</i>	0.5
<i>Molinia caerulea</i>	3
<i>Pedicularis sylvatica</i>	0.3
<i>Pleurozium schreberi</i>	63
<i>Potentilla erecta</i>	3

Relevé code	EC11 R5
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	PB3
Habitat - Annex I	*7130
Species	Percentage
<i>Calluna vulgaris</i>	8
<i>Calypogeia muelleriana</i>	0.3
<i>Drosera rotundifolia</i>	0.3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	8
<i>Molinia caerulea</i>	18
<i>Narthecium ossifragum</i>	0.5
<i>Odontoschisma sphagni</i>	0.3
<i>Rhynchospora alba</i>	3
<i>Schoenus nigricans</i>	63
<i>Sphagnum capillifolium s. rubellum</i>	8
<i>Sphagnum denticulatum</i>	42
<i>Sphagnum papillosum</i>	18

Relevé code	EC11 R6
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3 /HH1 /PB3 /WS1 /ER1
Habitat - Annex I	4010
Species	Percentage
<i>Aulacomnium palustre</i>	18
<i>Calluna vulgaris</i>	30
<i>Campylopus flexuosus</i>	0.3
<i>Carex panicea</i>	3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	18
<i>Hypnum jutlandicum</i>	18
<i>Molinia caerulea</i>	8
<i>Odontoschisma sphagni</i>	3
<i>Polygala serpyllifolia</i>	0.3
<i>Sphagnum capillifolium s. rubellum</i>	42
<i>Trichophorum germanicum</i>	42

Relevé code	EC11 R6
Species	Percentage
<i>Ulex gallii</i>	8

Relevé code	EC11 R7
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/ HH1/ PB3/ WS1/ ER1
Habitat - Annex I	4030
Species	Percentage
<i>Calluna vulgaris</i>	63
<i>Campylopus flexuosus</i>	0.3
<i>Cladonia portentosa</i>	0.3
<i>Cladonia sp.</i>	0.3
<i>Daboecia cantabrica</i>	0.3
<i>Erica cinerea</i>	0.3
<i>Hypnum jutlandicum</i>	42
<i>Molinia caerulea</i>	3
<i>Pteridium aquilinum</i>	0.3
<i>Ulex gallii</i>	42

Relevé code	EC11 R8
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	PF2
Habitat - Annex I	4010
Species	Percentage
<i>Agrostis vinealis</i>	0.5
<i>Anthoxanthum odoratum</i>	0.3
<i>Aulacomnium palustre</i>	0.3
<i>Calluna vulgaris</i>	8
<i>Erica tetralix</i>	3
<i>Hypnum jutlandicum</i>	8
<i>Juncus articulatus</i>	8
<i>Molinia caerulea</i>	42
<i>Potentilla erecta</i>	0.5
<i>Rhynchospora squarrosus</i>	3
<i>Succisa pratensis</i>	0.5
<i>Thuidium tamariscinum</i>	18
<i>Ulex gallii</i>	42

Relevé code	EC11 R9
Date	13/10/2017
Recorder	CC
Species recording scale	Percentage
Habitat - Fossitt	HD1 /WS1 /GS3 /GS4
Habitat - Annex I	4010
Species	Percentage
<i>Calluna vulgaris</i>	30
<i>Carex panicea</i>	3
<i>Erica tetralix</i>	8



Relevé code	EC11 R9
Species	Percentage
<i>Eriophorum angustifolium</i>	18
<i>Molinia caerulea</i>	18
<i>Narthecium ossifragum</i>	0.5
<i>Potentilla erecta</i>	0.5
<i>Rhytidiadelphus loreus</i>	8
<i>Schoenus nigricans</i>	8
<i>Sphagnum fallax</i>	42
<i>Trichophorum germanicum</i>	8
<i>Ulex gallii</i>	8

Relevé code	EC11 R10
Date	13/10/2017
Recorder	CC
Species recording scale	Percentage
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Percentage
<i>Calluna vulgaris</i>	83
<i>Molinia caerulea</i>	0.5
<i>Rhytidiadelphus loreus</i>	3
<i>Ulex gallii</i>	18

Relevé code	EC11 R11
Date	13/10/2017
Recorder	CC
Species recording scale	Percentage
Habitat - Fossitt	HH3/ER1
Habitat - Annex I	4010
Species	Percentage
<i>Calluna vulgaris</i>	30
<i>Carex panicea</i>	42
<i>Erica tetralix</i>	8
<i>Hylocomium splendens</i>	3
<i>Molinia caerulea</i>	42
<i>Potentilla erecta</i>	0.5
<i>Ulex europaeus</i>	8
<i>Ulex gallii</i>	18

Relevé code	EC11 R12
Date	13/10/2017
Recorder	CC
Species recording scale	Percentage
Habitat - Fossitt	GS4
Habitat - Annex I	n/a
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.5
<i>Cirsium vulgare</i>	0.3
<i>Molinia caerulea</i>	83

Relevé code	EC11 R12
Species	Percentage
<i>Potentilla erecta</i>	3
<i>Ranunculus acris</i>	0.5
<i>Ranunculus repens</i>	0.5
<i>Succisa pratensis</i>	3
<i>Vicia sepium</i>	0.3

Relevé code	EC12 R1
Date	08/09/2014
Recorder	JF
Species recording scale	Percentage
Habitat - Fossitt	HH1/HD1
Habitat - Annex I	4030
Species	Percentage
<i>Calluna vulgaris</i>	42
<i>Carex binervis</i>	3
<i>Daboecia cantabrica</i>	0.5
<i>Erica cinerea</i>	3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	3
<i>Hypnum jutlandicum</i>	8
<i>Molinia caerulea</i>	3
<i>Potentilla erecta</i>	0.3
<i>Pseudoscleropodium purum</i>	3
<i>Trichophorum germanicum</i>	3
<i>Ulex gallii</i>	42

Relevé code	EC12 R2
Date	08/09/2014
Recorder	JF
Species recording scale	Percentage
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Percentage
<i>Calluna vulgaris</i>	3
<i>Carex echinata</i>	0.3
<i>Carex panicea</i>	3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	3
<i>Festuca ovina</i>	3
<i>Hypnum jutlandicum</i>	0.3
<i>Juncus bufonius</i>	3
<i>Molinia caerulea</i>	42
<i>Narthecium ossifragum</i>	3
<i>Pedicularis sylvatica</i>	0.5
<i>Polygala serpyllifolia</i>	0.5
<i>Potentilla erecta</i>	0.5
<i>Sphagnum affine</i>	18
<i>Sphagnum capillifolium s. rubellum</i>	8
<i>Sphagnum denticulatum</i>	18

Relevé code	EC12 R2
Species	Percentage
<i>Sphagnum papillosum</i>	8
<i>Succisa pratensis</i>	3
<i>Ulex gallii</i>	0.5

Relevé code	EC12 R3
Date	07/09/2015
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis stolonifera</i>	18
<i>Anthoxanthum odoratum</i>	3
<i>Blechnum spicant</i>	0.1
<i>Calluna vulgaris</i>	3
<i>Carex flacca</i>	0.5
<i>Danthonia decumbens</i>	0.5
<i>Erica cinerea</i>	8
<i>Festuca rubra</i>	8
<i>Holcus lanatus</i>	3
<i>Hypnum jutlandicum</i>	42
<i>Potentilla erecta</i>	3
<i>Rhytidiadelphus squarrosus</i>	18
<i>Rubus fruticosus</i> agg.	8
<i>Solidago virgaurea</i>	3
<i>Taraxacum officinale</i> agg.	8
<i>Ulex gallii</i>	63
<i>Viola riviniana</i>	0.3

Relevé code	EC13 R1
Date	04/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Percentage
<i>Agrostis stolonifera</i>	18
<i>Angelica sylvestris</i>	0.5
<i>Anthoxanthum odoratum</i>	3
<i>Carex echinata</i>	8
<i>Carex panicea</i>	3
<i>Carex viridula s. brachyrrhyncha</i>	0.5
<i>Festuca rubra</i>	3
<i>Filipendula ulmaria</i>	18
<i>Holcus lanatus</i>	0.3
<i>Juncus articulatus</i>	18
<i>Juncus conglomeratus</i>	18
<i>Luzula multiflora</i>	0.3
<i>Lythrum salicaria</i>	8

Relevé code	EC13 R1
Species	Percentage
<i>Molinia caerulea</i>	42
<i>Plantago lanceolata</i>	8
<i>Potentilla erecta</i>	8
<i>Potentilla palustris</i>	18
<i>Succisa pratensis</i>	18

Relevé code	EC13 R2
Date	10/09/2016
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3 /HH1 /GS3 /WS1 /ER1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis vinealis</i>	8
<i>Calluna vulgaris</i>	42
<i>Campylopus introflexus</i>	18
<i>Carex binervis</i>	3
<i>Carex panicea</i>	3
<i>Erica cinerea</i>	8
<i>Hylocomium splendens</i>	0.3
<i>Hypnum jutlandicum</i>	18
<i>Hypochaeris radicata</i>	0.3
<i>Molinia caerulea</i>	0.3
<i>Pleurozium schreberi</i>	18
<i>Potentilla erecta</i>	0.3
<i>Succisa pratensis</i>	0.5
<i>Ulex gallii</i>	42

Relevé code	EC13 R3
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3 /PB3 /ER1
Habitat - Annex I	4010
Species	Percentage
<i>Aulacomnium palustre</i>	18
<i>Calluna vulgaris</i>	6
<i>Campylopus flexuosus</i>	3
<i>Cladonia portentosa</i>	0.3
<i>Drosera rotundifolia</i>	0.5
<i>Erica tetralix</i>	8
<i>Hypnum jutlandicum</i>	8
<i>Molinia caerulea</i>	8
<i>Narthecium ossifragum</i>	8
<i>Potentilla erecta</i>	0.3
<i>Sphagnum capillifolium s. rubellum</i>	42
<i>Sphagnum papillosum</i>	0.3
<i>Trichophorum germanicum</i>	30

Relevé code	EC13 R4
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3 /PB3 /ER1
Habitat - Annex I	*7130
Species	Percentage
<i>Calluna vulgaris</i>	8
<i>Drosera rotundifolia</i>	0.3
<i>Erica tetralix</i>	8
<i>Eriophorum angustifolium</i>	8
<i>Hypnum jutlandicum</i>	0.3
<i>Molinia caerulea</i>	30
<i>Odontoschisma sphagni</i>	3
<i>Potentilla erecta</i>	0.3
<i>Schoenus nigricans</i>	42
<i>Sphagnum capillifolium s. rubellum</i>	0.5
<i>Sphagnum denticulatum</i>	18
<i>Sphagnum magellanicum</i>	0.3
<i>Sphagnum papillosum</i>	83
<i>Trichophorum germanicum</i>	3

Relevé code	EC13 R5
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	GS3 /GS4 /HH1 /GM1
Habitat - Annex I	4030
Species	Percentage
<i>Agrostis vinealis</i>	8
<i>Calluna vulgaris</i>	42
<i>Campylopus flexuosus</i>	0.5
<i>Carex panicea</i>	18
<i>Carex viridula</i>	0.3
<i>Cladonia portentosa</i>	0.5
<i>Dactylorhiza maculata</i>	0.3
<i>Danthonia decumbens</i>	3
<i>Erica cinerea</i>	8
<i>Festuca rubra</i>	0.5
<i>Hylocomium splendens</i>	63
<i>Lotus corniculatus</i>	8
<i>Pleurozium schreberi</i>	8
<i>Potentilla erecta</i>	3
<i>Rhytidiadelphus squarrosus</i>	18
<i>Succisa pratensis</i>	8
<i>Trifolium pratense</i>	0.3

Relevé code	EC13 R6
Date	10/09/2014
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3/PB3/ER1

Relevé code	EC13 R6
Habitat - Annex I	7150
Species	Percentage
<i>Calluna vulgaris</i>	8
<i>Cladonia sp.</i>	0.3
<i>Drosera rotundifolia</i>	0.5
<i>Erica tetralix</i>	8
<i>Eriophorum vaginatum</i>	0.5
<i>Molinia caerulea</i>	8
<i>Narthecium ossifragum</i>	3
<i>Odontoschisma sphagni</i>	0.3
<i>Potentilla erecta</i>	0.3
<i>Racomitrium lanuginosum</i>	0.3
<i>Rhynchospora alba</i>	63
<i>Sphagnum cuspidatum</i>	18
<i>Sphagnum papillosum</i>	63
<i>Succisa pratensis</i>	0.3

Relevé code	EC13 R7
Date	07/09/2015
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	PB3
Habitat - Annex I	*7130
Species	Percentage
<i>Aulacomnium palustre</i>	3
<i>Calluna vulgaris</i>	18
<i>Calypogeia muelleriana</i>	0.3
<i>Carex panicea</i>	3
<i>Erica tetralix</i>	8
<i>Eriophorum angustifolium</i>	3
<i>Eriophorum vaginatum</i>	0.5
<i>Hylocomium splendens</i>	8
<i>Hypnum jutlandicum</i>	18
<i>Molinia caerulea</i>	63
<i>Narthecium ossifragum</i>	0.5
<i>Pedicularis sylvatica</i>	0.3
<i>Potentilla erecta</i>	0.3
<i>Sphagnum capillifolium s. rubellum</i>	8
<i>Sphagnum papillosum</i>	63
<i>Succisa pratensis</i>	3
<i>Thuidium tamariscinum</i>	0.5

Relevé code	EC13 R8
Date	07/09/2015
Recorder	EOS
Species recording scale	Percentage
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.3
<i>Calluna vulgaris</i>	8

Relevé code	EC13 R8
Species	Percentage
<i>Calypogeia muelleriana</i>	3
<i>Dactylorhiza maculata</i>	0.3
<i>Erica tetralix</i>	18
<i>Eriophorum angustifolium</i>	8
<i>Eriophorum vaginatum</i>	3
<i>Hypnum jutlandicum</i>	8
<i>Leucobryum glaucum</i>	3
<i>Molinia caerulea</i>	63
<i>Pleurozium schreberi</i>	0.5
<i>Potentilla erecta</i>	3
<i>Riccardia palmata</i>	0.3
<i>Schoenus nigricans</i>	42
<i>Sphagnum subnitens</i>	63
<i>Succisa pratensis</i>	0.3

Relevé code	EC14 R1
Date	04/09/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Domin
<i>Calluna vulgaris</i>	4
<i>Erica cinerea</i>	4
<i>Erica tetralix</i>	1
<i>Hypnum jutlandicum</i>	4
<i>Molinia caerulea</i>	4
<i>Potentilla erecta</i>	3
<i>Pseudoscleropodium purum</i>	1
<i>Succisa pratensis</i>	1
<i>Ulex gallii</i>	8

Relevé code	EC14 R2
Date	04/09/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	5
<i>Campylopus introflexus</i>	2
<i>Carex panicea</i>	3
<i>Cladonia portentosa</i>	3
<i>Dicranum scoparium</i>	4
<i>Erica tetralix</i>	3
<i>Hypnum jutlandicum</i>	4
<i>Molinia caerulea</i>	5
<i>Potentilla erecta</i>	2
<i>Schoenus nigricans</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	4

Relevé code	EC14 R2
Species	Domin
<i>Trichophorum germanicum</i>	6
<i>Ulex gallii</i>	3
<i>Diplophyllum albicans</i>	1

Relevé code	EC14 R3
Date	04/09/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	PB3
Habitat - Annex I	*7130
Species	Domin
<i>Aulacomnium palustre</i>	2
<i>Calluna vulgaris</i>	3
<i>Carex panicea</i>	1
<i>Carex pulicaris</i>	1
<i>Drosera rotundifolia</i>	2
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	3
<i>Juncus articulatus</i>	3
<i>Molinia caerulea</i>	3
<i>Narthecium ossifragum</i>	2
<i>Odontoschisma sphagni</i>	1
<i>Rhynchospora alba</i>	3
<i>Schoenus nigricans</i>	7
<i>Sphagnum capillifolium s. rubellum</i>	3
<i>Sphagnum papillosum</i>	7
<i>Sphagnum tenellum</i>	1
<i>Trichophorum germanicum</i>	3

Relevé code	EC14 R4
Date	08/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Anthoxanthum odoratum</i>	1
<i>Calluna vulgaris</i>	2
<i>Cirsium dissectum</i>	4
<i>Erica tetralix</i>	3
<i>Holcus lanatus</i>	1
<i>Juncus articulatus</i>	3
<i>Molinia caerulea</i>	7
<i>Myrica gale</i>	6
<i>Narthecium ossifragum</i>	2
<i>Potentilla erecta</i>	2
<i>Scleropodium purum</i>	1
<i>Thuidium tamariscinum</i>	1
<i>Ulex gallii</i>	1
<i>Warnstorfia fluitans</i>	1



Relevé code	EC14 R4
Species	Domin
<i>Sphagnum</i> sp.	1

Relevé code	EC14 R5
Date	13/10/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	-
Species	Domin
<i>Anthoxanthum odoratum</i>	3
<i>Calluna vulgaris</i>	1
<i>Hylocomium splendens</i>	4
<i>Juncus acutiflorus</i>	7
<i>Molinia caerulea</i>	8
<i>Potentilla erecta</i>	4
<i>Rhynchospora alba</i>	3
<i>Rumex acetosa</i>	2
<i>Sphagnum fallax</i>	5
<i>Succisa pratensis</i>	5
<i>Thuidium</i> sp.	3

Relevé code	EC18 R1
Date	04/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Aulacomnium palustre</i>	4
<i>Calluna vulgaris</i>	7
<i>Hylocomium splendens</i>	7
<i>Juncus articulatus</i>	3
<i>Molinia caerulea</i>	6
<i>Potentilla erecta</i>	2
<i>Rhynchospora alba</i>	3
<i>Sphagnum papillosum</i>	5
<i>Succisa pratensis</i>	2
<i>Thuidium tamariscinum</i>	3

Relevé code	EC18 R2
Date	04/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Agrostis capillaris</i>	2
<i>Agrostis stolonifera</i>	4

Relevé code	EC18 R2
<i>Anthoxanthum odoratum</i>	4
<i>Calliergonella cuspidata</i>	2
<i>Carex panicea</i>	3
<i>Dactylorhiza</i> sp.	2
<i>Filipendula ulmaria</i>	3
<i>Holcus lanatus</i>	7
<i>Hylocomium splendens</i>	2
<i>Kindbergia praelonga</i>	2
<i>Luzula multiflora</i>	2
<i>Lythrum salicaria</i>	2
<i>Mentha aquatica</i>	3
<i>Molinia caerulea</i>	7
<i>Plantago lanceolata</i>	3
<i>Potentilla erecta</i>	3
<i>Succisa pratensis</i>	5

Relevé code	EC18 R3
Date	04/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Domin
<i>Arctostaphylos uva-ursi</i>	3
<i>Calluna vulgaris</i>	6
<i>Carex binervis</i>	2
<i>Carex echinata</i>	2
<i>Carex panicea</i>	4
<i>Daboecia cantabrica</i>	3
<i>Hylocomium splendens</i>	3
<i>Juncus effusus</i>	3
<i>Molinia caerulea</i>	5
<i>Potentilla erecta</i>	2
<i>Sphagnum papillosum</i>	4
<i>Succisa pratensis</i>	2

Relevé code	EC18 R4
Date	04/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	6
<i>Carex echinata</i>	2
<i>Carex panicea</i>	3
<i>Cladonia portentosa</i>	3
<i>Dactylorhiza</i> sp.	1
<i>Erica tetralix</i>	4
<i>Hylocomium splendens</i>	4
<i>Juncus articulatus</i>	3

Relevé code	EC18 R4
Species	Domin
<i>Molinia caerulea</i>	5
<i>Narthecium ossifragum</i>	4
<i>Sphagnum capillifolium s. rubellum</i>	8
<i>Succisa pratensis</i>	3
<i>Trichophorum germanicum</i>	5

Relevé code	EC18 R5
Date	08/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Agrostis capillaris</i>	5
<i>Agrostis stolonifera</i>	5
<i>Anagallis tenella</i>	2
<i>Anthoxanthum odoratum</i>	4
<i>Carex echinata</i>	3
<i>Carex panicea</i>	3
<i>Danthonia decumbens</i>	2
<i>Hydrocotyle vulgaris</i>	2
<i>Juncus articulatus</i>	5
<i>Luzula multiflora</i>	3
<i>Molinia caerulea</i>	5
<i>Nardus stricta</i>	3
<i>Pedicularis sylvatica</i>	2
<i>Pleurozium schreberi</i>	5
<i>Polygala serpyllifolia</i>	2
<i>Potentilla erecta</i>	3
<i>Rhynchospora squarrosus</i>	5
<i>Succisa pratensis</i>	5
<i>Sphagnum sp.</i>	4

Relevé code	EC18 R6
Date	08/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS3
Habitat - Annex I	-
Species	Domin
<i>Agrostis capillaris</i>	4
<i>Anthoxanthum odoratum</i>	4
<i>Carex sp.</i>	2
<i>Cynosurus cristatus</i>	2
<i>Festuca ovina</i>	4
<i>Galium saxatile</i>	4
<i>Hylocomium splendens</i>	5
<i>Hypochaeris radicata</i>	3
<i>Juncus articulatus</i>	3
<i>Juncus effusus</i>	3

Relevé code	EC18 R6
Species	Domin
<i>Nardus stricta</i>	6
<i>Pedicularis sylvatica</i>	2
<i>Plantago lanceolata</i>	3
<i>Pleurozium schreberi</i>	5
<i>Potentilla erecta</i>	5
<i>Prunella vulgaris</i>	3
<i>Rhytidadelphus squarrosus</i>	6
<i>Rumex acetosa</i>	3
<i>Trifolium repens</i>	4

Relevé code	EC18 R7
Date	09/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Domin
<i>Agrostis capillaris</i>	3
<i>Anthoxanthum odoratum</i>	2
<i>Arrhenatherum elatius</i>	1
<i>Calluna vulgaris</i>	7
<i>Cladonia portentosa</i>	2
<i>Deschampsia cespitosa</i>	3
<i>Dicranum scoparium</i>	1
<i>Erica cinerea</i>	7
<i>Festuca rubra</i>	2
<i>Hylocomium splendens</i>	8
<i>Hypnum jutlandicum</i>	2
<i>Lathyrus pratensis</i>	1
<i>Pleurozium schreberi</i>	4
<i>Polypodium vulgare</i>	1
<i>Potentilla erecta</i>	3
<i>Rumex acetosa</i>	1
<i>Sedum anglicum</i>	1
<i>Succisa pratensis</i>	1

Relevé code	EC20 R1
Date	29/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	4
<i>Carex panicea</i>	3
<i>Cladonia portentosa</i>	3
<i>Cladonia uncialis</i>	3
<i>Dicranum scoparium</i>	1
<i>Erica cinerea</i>	3
<i>Erica tetralix</i>	5

Relevé code	EC20 R1
Species	Domin
<i>Hypnum jutlandicum</i>	5
<i>Molinia caerulea</i>	5
<i>Narthecium ossifragum</i>	3
<i>Pedicularis sylvatica</i>	3
<i>Potentilla erecta</i>	4
<i>Racomitrium lanuginosum</i>	4
<i>Sphagnum capillifolium s. rubellum</i>	3
<i>Sphagnum compactum</i>	2
<i>Trichophorum germanicum</i>	6

Relevé code	EC20 R2
Date	29/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	PB3
Habitat - Annex I	*7130
Species	Domin
<i>Calluna vulgaris</i>	4
<i>Carex panicea</i>	3
<i>Cladonia portentosa</i>	3
<i>Cladonia uncialis</i>	1
<i>Drosera rotundifolia</i>	1
<i>Eleocharis multicaulis</i>	2
<i>Erica tetralix</i>	5
<i>Eriophorum angustifolium</i>	4
<i>Eriophorum vaginatum</i>	3
<i>Molinia caerulea</i>	5
<i>Narthecium ossifragum</i>	5
<i>Odontoschisma sphagni</i>	4
<i>Potentilla erecta</i>	2
<i>Rhynchospora alba</i>	6
<i>Sphagnum capillifolium s. rubellum</i>	5
<i>Sphagnum cuspidatum</i>	5
<i>Sphagnum magellanicum</i>	2
<i>Sphagnum papillosum</i>	4
<i>Sphagnum tenellum</i>	3

Relevé code	EC20 R3
Date	29/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Agrostis stolonifera</i>	5
<i>Anthoxanthum odoratum</i>	4
<i>Calluna vulgaris</i>	3
<i>Carex echinata</i>	3
<i>Carex panicea</i>	3
<i>Dactylorhiza sp.</i>	1

Relevé code	EC20 R3
Species	Domin
<i>Erica tetralix</i>	1
<i>Holcus lanatus</i>	3
<i>Hylocomium splendens</i>	5
<i>Juncus acutiflorus</i>	5
<i>Lythrum salicaria</i>	4
<i>Molinia caerulea</i>	6
<i>Potentilla erecta</i>	5
<i>Scleropodium purum</i>	2
<i>Sphagnum palustre</i>	4
<i>Succisa pratensis</i>	5
<i>Thuidium tamariscinum</i>	3

Relevé code	EC20 R4
Date	29/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	6
<i>Campylopus introflexus</i>	4
<i>Carex panicea</i>	4
<i>Cladonia portentosa</i>	3
<i>Cladonia</i> sp.	3
<i>Erica cinerea</i>	4
<i>Molinia caerulea</i>	5
<i>Pedicularis sylvatica</i>	2
<i>Polygala serpyllifolia</i>	1
<i>Potentilla erecta</i>	2
<i>Ulex gallii</i>	6

Relevé code	EC20 R5
Date	09/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	5
<i>Calypogeia muelleriana</i>	1
<i>Campylopus flexuosus</i>	3
<i>Campylopus introflexus</i>	1
<i>Carex panicea</i>	5
<i>Cladonia portentosa</i>	2
<i>Cladonia uncialis</i>	1
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	2
<i>Hypnum jutlandicum</i>	3
<i>Kurzia</i> sp.	1
<i>Leucobryum glaucum</i>	3

Relevé code	EC20 R5
Species	Domin
<i>Narthecium ossifragum</i>	5
<i>Odontoschisma sphagni</i>	1
<i>Pleurozia purpurea</i>	3
<i>Racomitrium lanuginosum</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	2
<i>Sphagnum compactum</i>	4
<i>Sphagnum cuspidatum</i>	2
<i>Sphagnum denticulatum</i>	6
<i>Sphagnum tenellum</i>	2
<i>Trichophorum germanicum</i>	5
<i>Ulex gallii</i>	2
<i>Diplophyllum albicans</i>	1

Relevé code	EC21 R1
Date	07/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
Species	Domin
<i>Agrostis canina</i>	3
<i>Calluna vulgaris</i>	7
<i>Carex binervis</i>	1
<i>Cladonia portentosa</i>	1
<i>Daboecia cantabrica</i>	4
<i>Erica cinerea</i>	4
<i>Hypnum jutlandicum</i>	3
<i>Molinia caerulea</i>	5
<i>Nardus stricta</i>	2
<i>Scleropodium purum</i>	3
<i>Ulex gallii</i>	5
<i>Diplophyllum albicans</i>	2

Relevé code	EC21 R2
Date	07/08/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	5
<i>Carex panicea</i>	1
<i>Cladonia portentosa</i>	3
<i>Dicranum scoparium</i>	1
<i>Erica tetralix</i>	5
<i>Eriophorum angustifolium</i>	4
<i>Molinia caerulea</i>	5
<i>Narthecium ossifragum</i>	4
<i>Odontoschisma sphagni</i>	1
<i>Polygala serpyllifolia</i>	1

Relevé code	EC21 R2
Species	Domin
<i>Potentilla erecta</i>	3
<i>Racomitrium lanuginosum</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	5
<i>Sphagnum tenellum</i>	2
<i>Succisa pratensis</i>	3
<i>Trichophorum germanicum</i>	5
<i>Ulex gallii</i>	3

Relevé code	EC21 R3
Date	23/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Agrostis stolonifera</i>	1
<i>Aulacomnium palustre</i>	4
<i>Calluna vulgaris</i>	4
<i>Carex panicea</i>	4
<i>Cladonia portentosa</i>	1
<i>Erica tetralix</i>	1
<i>Eriophorum angustifolium</i>	1
<i>Hylocomium splendens</i>	1
<i>Hypnum jutlandicum</i>	4
<i>Juncus bulbosus</i>	3
<i>Molinia caerulea</i>	4
<i>Mylia taylorii</i>	1
<i>Narthecium ossifragum</i>	3
<i>Odontoschisma sphagni</i>	1
<i>Poa trivialis</i>	1
<i>Potentilla erecta</i>	1
<i>Racomitrium lanuginosum</i>	4
<i>Sphagnum capillifolium s. rubellum</i>	2
<i>Sphagnum cuspidatum</i>	1
<i>Sphagnum denticulatum</i>	2
<i>Succisa pratensis</i>	2
<i>Trichophorum germanicum</i>	2
<i>Trifolium repens</i>	1

Relevé code	EC23 R1
Date	09/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS3
Habitat - Annex I	-
Species	Domin
<i>Achillea millefolium</i>	2
<i>Agrostis capillaris</i>	8 combined
<i>Anthoxanthum odoratum</i>	8 combined
<i>Arrhenatherum elatius</i>	8 combined



Relevé code	EC23 R1
Species	Domin
<i>Centaurea nigra</i>	5
<i>Cerastium fontanum</i>	3
<i>Cynosurus cristatus</i>	8 combined
<i>Dactylis glomerata</i>	8 combined
<i>Festuca rubra</i>	8 combined
<i>Holcus lanatus</i>	8 combined
<i>Hypericum pulchrum</i>	1
<i>Leucanthemum vulgare</i>	4
<i>Linum catharticum</i>	4
<i>Lotus corniculatus</i>	5
<i>Pilosella officinarum</i>	2
<i>Plantago lanceolata</i>	4
<i>Prunella vulgaris</i>	4
<i>Pseudoscleropodium purum</i>	4
<i>Rhynchospora squarrosus</i>	5
<i>Senecio jacobaea</i>	2
<i>Trifolium pratense</i>	4
<i>Veronica officinalis</i>	2
<i>Viola</i> sp.	2

Relevé code	EC25 R1
Date	05/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Calluna vulgaris</i>	5
<i>Campylopus introflexus</i>	2
<i>Carex binervis</i>	4
<i>Carex panicea</i>	3
<i>Cladonia portentosa</i>	2
<i>Daboecia cantabrica</i>	4
<i>Dicranum scoparium</i>	3
<i>Erica cinerea</i>	4
<i>Hypnum jutlandicum</i>	4
<i>Molinia caerulea</i>	6
<i>Pedicularis sylvatica</i>	2
<i>Ulex gallii</i>	5

Relevé code	EC25 R2
Date	05/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	PB3
Habitat - Annex I	*7130
Species	Domin
<i>Calluna vulgaris</i>	2
<i>Campylopus atrovirens</i>	3
<i>Carex panicea</i>	1

<b>Relevé code</b>	<b>EC25 R2</b>
Species	Domin
<i>Drosera rotundifolia</i>	2
<i>Erica tetralix</i>	4
<i>Eriophorum angustifolium</i>	4
<i>Hypnum jutlandicum</i>	5
<i>Menyanthes trifoliata</i>	1
<i>Molinia caerulea</i>	8
<i>Narthecium ossifragum</i>	3
<i>Polygala serpyllifolia</i>	1
<i>Potentilla erecta</i>	3
<i>Rhynchospora alba</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	5
<i>Sphagnum denticulatum</i>	3
<i>Sphagnum papillosum</i>	3
<i>Trichophorum germanicum</i>	3

<b>Relevé code</b>	<b>EC25 R3</b>
Date	06/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	HH3
Habitat - Annex I	4010
Species	Domin
<i>Agrostis stolonifera</i>	3
<i>Aulacomnium palustre</i>	2
<i>Calluna vulgaris</i>	4
<i>Campylopus atrovirens</i>	1
<i>Carex panicea</i>	2
<i>Drosera rotundifolia</i>	3
<i>Erica tetralix</i>	4
<i>Eriophorum angustifolium</i>	3
<i>Hypnum jutlandicum</i>	3
<i>Juncus acutiflorus</i>	3
<i>Molinia caerulea</i>	7
<i>Narthecium ossifragum</i>	3
<i>Pedicularis sylvatica</i>	2
<i>Polygala serpyllifolia</i>	3
<i>Potentilla erecta</i>	3
<i>Pseudoscleropodium purum</i>	3
<i>Rhynchospora fusca</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	7
<i>Sphagnum denticulatum</i>	3
<i>Sphagnum papillosum</i>	4
<i>Succisa pratensis</i>	4
<i>Trichophorum germanicum</i>	2

<b>Relevé code</b>	<b>EC25 R4</b>
Date	06/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	PB3

<b>Relevé code</b>	<b>EC25 R4</b>
Habitat - Annex I	7150
Species	Domin
<i>Calluna vulgaris</i>	4
<i>Cladonia portentosa</i>	2
<i>Drosera rotundifolia</i>	3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	3
<i>Hypnum jutlandicum</i>	2
<i>Molinia caerulea</i>	4
<i>Narthecium ossifragum</i>	3
<i>Potentilla erecta</i>	2
<i>Rhynchospora alba</i>	8
<i>Sphagnum capillifolium s. rubellum</i>	5
<i>Sphagnum cuspidatum</i>	4
<i>Sphagnum papillosum</i>	5

<b>Relevé code</b>	<b>EC25 R5</b>
Date	13/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	PF3
Habitat - Annex I	7140
Species	Domin
<i>Aulacomnium palustre</i>	3
<i>Calluna vulgaris</i>	2
<i>Carex lasiocarpa</i>	2
<i>Carex rostrata</i>	3
<i>Drosera rotundifolia</i>	2
<i>Equisetum fluviatile</i>	3
<i>Erica tetralix</i>	3
<i>Eriophorum angustifolium</i>	4
<i>Hylocomium splendens</i>	3
<i>Juncus bulbosus</i>	2
<i>Menyanthes trifoliata</i>	3
<i>Molinia caerulea</i>	3
<i>Myrica gale</i>	4
<i>Phragmites australis</i>	4
<i>Potentilla palustris</i>	4
<i>Pseudoscleropodium purum</i>	3
<i>Schoenus nigricans</i>	3
<i>Sphagnum capillifolium s. rubellum</i>	6
<i>Sphagnum palustre</i>	3
<i>Succisa pratensis</i>	3
<i>Vaccinium oxycoccos</i>	2

<b>Relevé code</b>	<b>EC25 R6</b>
Date	06/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS2
Habitat - Annex I	6510

<b>Relevé code</b>	<b>EC25 R6</b>
Species	Domin
<i>Achillea millefolium</i>	2
<i>Agrostis capillaris</i>	4
<i>Anthoxanthum odoratum</i>	4
<i>Centaurea nigra</i>	4
<i>Cerastium fontanum</i>	1
<i>Cynosurus cristatus</i>	5
<i>Dactylis glomerata</i>	4
<i>Euphrasia</i> sp.	3
<i>Hypochaeris radicata</i>	4
<i>Lathyrus pratensis</i>	2
<i>Leontodon autumnalis</i>	1
<i>Leontodon hispidus</i>	3
<i>Pimpinella major</i>	3
<i>Plantago lanceolata</i>	4
<i>Prunella vulgaris</i>	1
<i>Pteridium aquilinum</i>	1
<i>Ranunculus repens</i>	3
<i>Senecio jacobaea</i>	1
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	2

<b>Relevé code</b>	<b>EC25 R7</b>
Date	06/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS3
Habitat - Annex I	-
Species	Domin
<i>Achillea millefolium</i>	3
<i>Agrostis capillaris</i>	6
<i>Centaurea nigra</i>	2
<i>Cynosurus cristatus</i>	4
<i>Euphrasia</i> sp.	3
<i>Festuca rubra</i>	4
<i>Hypochaeris radicata</i>	4
<i>Leontodon autumnalis</i>	4
<i>Lolium perenne</i>	6
<i>Lotus corniculatus</i>	2
<i>Nardus stricta</i>	1
<i>Polygala serpyllifolia</i>	2
<i>Potentilla erecta</i>	3
<i>Prunella vulgaris</i>	3
<i>Ranunculus repens</i>	4
<i>Senecio jacobaea</i>	2
<i>Trifolium pratense</i>	4

<b>Relevé code</b>	<b>EC25 R8</b>
Date	13/10/2017
Recorder	CC

Relevé code	EC25 R8
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	-
Species	Domin
<i>Calluna vulgaris</i>	1
<i>Carex panicea</i>	4
<i>Erica tetralix</i>	1
<i>Juncus acutiflorus</i>	6
<i>Lythrum salicaria</i>	1
<i>Molinia caerulea</i>	8
<i>Potentilla erecta</i>	4
<i>Sphagnum fallax</i>	5
<i>Succisa pratensis</i>	3

Relevé code	EC29 R1
Date	22/07/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Achillea millefolium</i>	1
<i>Anthoxanthum odoratum</i>	4
<i>Briza media</i>	2
<i>Calliergonella cuspidata</i>	5
<i>Carex hirta</i>	3
<i>Centaurea nigra</i>	6
<i>Cerastium fontanum</i>	1
<i>Cynosurus cristatus</i>	4
<i>Dactylis glomerata</i>	4
<i>Equisetum arvense</i>	3
<i>Euphrasia</i> sp.	4
<i>Festuca rubra</i>	4
<i>Filipendula ulmaria</i>	2
<i>Holcus lanatus</i>	4
<i>Lathyrus pratensis</i>	2
<i>Leontodon hispidus</i>	2
<i>Lolium perenne</i>	3
<i>Lotus corniculatus</i>	3
<i>Plagiomnium undulatum</i>	3
<i>Plantago lanceolata</i>	3
<i>Potentilla anserina</i>	4
<i>Potentilla reptans</i>	3
<i>Ranunculus acris</i>	3
<i>Rhinanthus minor</i>	3
<i>Rhytidadelphus squarrosus</i>	4
<i>Trifolium pratense</i>	5
<i>Trifolium repens</i>	4
<i>Vicia cracca</i>	3

Relevé code	EC31 R1
Date	09/07/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1/GS2
Habitat - Annex I	6510
Species	Domin
<i>Agrostis capillaris</i>	4
<i>Anthoxanthum odoratum</i>	4
<i>Briza media</i>	2
<i>Carex flacca</i>	2
<i>Centaurea nigra</i>	5
<i>Crataegus monogyna</i>	1
<i>Cynosurus cristatus</i>	2
<i>Dactylis glomerata</i>	2
<i>Festuca rubra</i>	3
<i>Helictotrichon pubescens</i>	4
<i>Lathyrus pratensis</i>	2
<i>Leontodon hispidus</i>	4
<i>Linum catharticum</i>	1
<i>Listera ovata</i>	1
<i>Lotus corniculatus</i>	4
<i>Plantago lanceolata</i>	3
<i>Potentilla erecta</i>	2
<i>Pseudoscleropodium purum</i>	2
<i>Ranunculus acris</i>	3
<i>Taraxacum officinale</i> agg.	1
<i>Trifolium pratense</i>	4

Relevé code	EC31 R2
Date	09/07/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1/GS3
Habitat - Annex I	*6210
Species	Domin
<i>Agrostis capillaris</i>	2
<i>Anthoxanthum odoratum</i>	3
<i>Blackstonia perfoliata</i>	2
<i>Briza media</i>	5
<i>Calliergonella cuspidata</i>	3
<i>Dactylis glomerata</i>	3
<i>Festuca rubra</i>	4
<i>Helictotrichon pubescens</i>	3
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	4
<i>Taraxacum officinale</i> agg.	2
<i>Trifolium pratense</i>	4

Relevé code	EC31 R3
Date	26/09/2014
Recorder	JF

Relevé code	EC31 R3
Species recording scale	Domin
Habitat - Fossitt	WN6
Habitat - Annex I	*91E0
Species	Domin
<i>Agrostis stolonifera</i>	6
<i>Alnus glutinosa</i>	8
<i>Atrichum undulatum</i>	1
<i>Centaurea nigra</i>	3
<i>Cirsium palustre</i>	2
<i>Crataegus monogyna</i>	1
<i>Dactylis glomerata</i>	3
<i>Fraxinus excelsior</i>	3
<i>Hedera helix</i>	3
<i>Kindbergia praelonga</i>	5
<i>Prunella vulgaris</i>	2
<i>Ranunculus repens</i>	3
<i>Rubus fruticosus</i> agg.	4
<i>Salix cinerea</i>	3

Relevé code	EC33 R1
Date	07/07/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS2
Habitat - Annex I	6510
Species	Domin
<i>Agrostis capillaris</i>	4
<i>Anthoxanthum odoratum</i>	5
<i>Brachythecium rutabulum</i>	3
<i>Centaurea nigra</i>	3
<i>Cerastium fontanum</i>	2
<i>Cirsium arvense</i>	1
<i>Cynosurus cristatus</i>	4
<i>Festuca rubra</i>	3
<i>Galium verum</i>	1
<i>Helictotrichon pubescens</i>	3
<i>Holcus lanatus</i>	4
<i>Leucanthemum vulgare</i>	4
<i>Lotus corniculatus</i>	1
<i>Plantago lanceolata</i>	3
<i>Prunella vulgaris</i>	3
<i>Pseudoscleropodium purum</i>	3
<i>Ranunculus repens</i>	1
<i>Rumex acetosa</i>	3
<i>Sonchus oleraceus</i>	4
<i>Trifolium dubium</i>	2
<i>Trifolium pratense</i>	3
<i>Trifolium repens</i>	1
<i>Trisetum flavescens</i>	2
<i>Veronica chamaedrys</i>	3

Relevé code	EC33 R2
Date	08/07/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Briza media</i>	5
<i>Carex flacca</i>	3
<i>Carex pulicaris</i>	2
<i>Crataegus monogyna</i>	2
<i>Cynosurus cristatus</i>	4
<i>Dactylis glomerata</i>	2
<i>Dactylorhiza fuchsii</i>	3
<i>Festuca rubra</i>	4
<i>Filipendula ulmaria</i>	1
<i>Galium boreale</i>	3
<i>Hedera helix</i>	3
<i>Hypochaeris radicata</i>	2
<i>Leucanthemum vulgare</i>	3
<i>Lotus corniculatus</i>	3
<i>Molinia caerulea</i>	3
<i>Parnassia palustris</i>	1
<i>Plantago lanceolata</i>	3
<i>Polypodium vulgare</i>	2
<i>Potentilla reptans</i>	1
<i>Prunus spinosa</i>	2
<i>Pseudoscleropodium purum</i>	3
<i>Schoenus nigricans</i>	5
<i>Succisa pratensis</i>	4
<i>Thymus polytrichus</i>	1
<i>Trifolium pratense</i>	3
<i>Valeriana officinalis</i>	1
<i>Vicia cracca</i>	1

Relevé code	EC33 R3
Date	30/09/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	1
<i>Ctenidium molluscum</i>	2
<i>Hedera helix</i>	4
<i>Ilex aquifolium</i>	1
<i>Prunus spinosa</i>	3
<i>Pteridium aquilinum</i>	4
<i>Rosa spinosissima</i>	4
<i>Rubus fruticosus</i> agg.	3
<i>Senecio jacobaea</i>	1
<i>Sesleria caerulea</i>	4
<i>Teucrium scorodonia</i>	1



Relevé code	EC33 R3
Species	Domin
<i>Tortella</i> sp.	1

Relevé code	EC33 R4
Date	01/10/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Anthoxanthum odoratum</i>	4
<i>Ctenidium molluscum</i>	1
<i>Festuca rubra</i>	4
<i>Hypericum tetrapterum</i>	2
<i>Lotus corniculatus</i>	2
<i>Pilosella officinarum</i>	1
<i>Potentilla erecta</i>	1
<i>Prunus spinosa</i>	1
<i>Pteridium aquilinum</i>	3
<i>Rosa spinosissima</i>	1
<i>Rubus fruticosus</i> agg.	2
<i>Sesleria albicans</i>	8
<i>Succisa pratensis</i>	2
<i>Teucrium scorodonia</i>	1
<i>Viola</i> sp.	1

Relevé code	EC36 R1
Date	15/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Species	Domin
<i>Anthoxanthum odoratum</i>	1
<i>Anthyllis vulneraria</i>	3
<i>Briza media</i>	4
<i>Calliargonella cuspidata</i>	1
<i>Carex flacca</i>	4
<i>Carlina vulgaris</i>	1
<i>Ctenidium molluscum</i>	2
<i>Dactylis glomerata</i>	2
<i>Euphrasia officinalis</i> agg.	2
<i>Festuca rubra</i>	4
<i>Gymnadenia conopsea</i>	2
<i>Hedera helix</i>	2
<i>Hypnum lacunosum</i>	4
<i>Leontodon hispidus</i>	4
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	4
<i>Pilosella officinarum</i>	3

Relevé code	EC36 R1
Species	Domin
<i>Plantago lanceolata</i>	3
<i>Plantago maritima</i>	4
<i>Potentilla erecta</i>	3
<i>Pseudoscleropodium purum</i>	2
<i>Rhytidadelphus squarrosus</i>	2
<i>Succisa pratensis</i>	4
<i>Thuidium delicatulum</i>	1
<i>Thymus polytrichus</i>	4
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	1

Relevé code	EC36 R2
Date	16/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Circaea lutetiana</i>	2
<i>Corylus avellana</i>	9
<i>Dryopteris filix-mas</i>	4
<i>Eurhynchium striatum</i>	7
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	3
<i>Ilex aquifolium</i>	2
<i>Isoetes macrospora</i>	2
<i>Melica uniflora</i>	3
<i>Neckera complanata</i>	2
<i>Neckera crispa</i>	2
<i>Phyllitis scolopendrium</i>	1
<i>Potentilla sterilis</i>	1
<i>Primula vulgaris</i>	1
<i>Thamnobryum alopecurum</i>	7
<i>Veronica chamaedrys</i>	1

Relevé code	EC36 R3
Date	16/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Agrostis capillaris</i>	3
<i>Agrostis stolonifera</i>	2
<i>Anthoxanthum odoratum</i>	4
<i>Arrhenatherum elatius</i>	2
<i>Briza media</i>	3
<i>Calliergonella cuspidata</i>	4
<i>Carex flacca</i>	2
<i>Centaurea nigra</i>	5

Relevé code	EC36 R3
Species	Domin
<i>Cerastium fontanum</i>	2
<i>Crepis capillaris</i>	1
<i>Cynosurus cristatus</i>	1
<i>Dactylis glomerata</i>	3
<i>Dactylorhiza fuchsii</i>	2
<i>Daucus carota</i>	3
<i>Dicranella varia</i>	1
<i>Euphrasia officinalis</i> agg.	3
<i>Festuca rubra</i>	4
<i>Helictotrichon pubescens</i>	2
<i>Homalothecium lutescens</i>	1
<i>Leontodon hispidus</i>	5
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	4
<i>Plagiomnium affine</i>	1
<i>Plagiomnium undulatum</i>	2
<i>Plantago lanceolata</i>	3
<i>Potentilla anserina</i>	1
<i>Primula veris</i>	1
<i>Prunella vulgaris</i>	2
<i>Ranunculus acris</i>	1
<i>Rhinanthus minor</i>	3
<i>Rubus fruticosus</i> agg.	2
<i>Succisa pratensis</i>	1
<i>Trifolium pratense</i>	5
<i>Trifolium repens</i>	3
<i>Vicia cracca</i>	4

Relevé code	EC36 R4
Date	24/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	FS2
Habitat - Annex I	6430
Species	Domin
<i>Agrostis stolonifera</i>	2
<i>Calliergonella cuspidata</i>	8
<i>Carex rostrata</i>	4
<i>Epilobium hirsutum</i>	5
<i>Epilobium palustre</i>	7
<i>Equisetum fluviatile</i>	4
<i>Galium palustre</i>	3
<i>Holcus lanatus</i>	1
<i>Iris pseudacorus</i>	3
<i>Menyanthes trifoliata</i>	6
<i>Phragmites australis</i>	5
<i>Prunus spinosa</i>	1
<i>Ranunculus acris</i>	2
<i>Valeriana dioica</i>	4

Relevé code	EC36 R5
Date	24/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	FL6
Habitat - Annex I	*3180
Species	Domin
<i>Agrostis stolonifera</i>	3
<i>Apium inundatum</i>	8
<i>Caltha palustris</i>	3
<i>Eleocharis palustris</i>	5
<i>Glyceria fluitans</i>	3
<i>Myosoton aquaticum</i>	5
<i>Veronica catenata</i>	1

Relevé code	EC36 R6
Date	25/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	2
<i>Carex flacca</i>	3
<i>Centranthus ruber</i>	3
<i>Ceterach officinarum</i>	2
<i>Crataegus monogyna</i>	1
<i>Euphrasia officinalis</i> agg.	3
<i>Festuca ovina</i>	1
<i>Fissidens dubius</i>	1
<i>Fraxinus excelsior</i>	2
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	3
<i>Hypericum</i> sp.	2
<i>Ilex aquifolium</i>	1
<i>Linum catharticum</i>	1
<i>Lotus corniculatus</i>	3
<i>Mycelis muralis</i>	2
<i>Potentilla erecta</i>	2
<i>Prunus spinosa</i>	4
<i>Rosa spinosissima</i>	3
<i>Senecio jacobaea</i>	2
<i>Sesleria albicans</i>	4
<i>Succisa pratensis</i>	3
<i>Teucrium scorodonia</i>	3
<i>Thalictrum minus</i>	3
<i>Thymus praecox</i>	2
<i>Tortella tortuosa</i>	2
<i>Trichostomum crispulum</i>	2

Relevé code	EC36 R7
Date	30/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS2
Habitat - Annex I	6510
Species	Domin
<i>Achillea millefolium</i>	2
<i>Agrimonia eupatoria</i>	2
<i>Agrostis capillaris</i>	2
<i>Anthoxanthum odoratum</i>	2
<i>Briza media</i>	3
<i>Centaurea nigra</i>	4
<i>Cerastium fontanum</i>	2
<i>Cynosurus cristatus</i>	5
<i>Dactylis glomerata</i>	3
<i>Daucus carota</i>	2
<i>Festuca rubra</i>	6
<i>Filipendula ulmaria</i>	3
<i>Holcus lanatus</i>	4
<i>Hypochaeris radicata</i>	3
<i>Lotus corniculatus</i>	3
<i>Phleum pratense</i>	3
<i>Plantago lanceolata</i>	3
<i>Prunella vulgaris</i>	3
<i>Prunus spinosa</i>	3
<i>Ranunculus acris</i>	2
<i>Rhinanthus minor</i>	3
<i>Trifolium pratense</i>	5

Relevé code	EC37 R1
Date	02/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Anthoxanthum odoratum</i>	2
<i>Blackstonia perfoliata</i>	1
<i>Briza media</i>	3
<i>Campylium protensum</i>	1
<i>Carex flacca</i>	5
<i>Carlina vulgaris</i>	1
<i>Centaurea nigra</i>	2
<i>Centranthus ruber</i>	3
<i>Ctenidium molluscum</i>	3
<i>Euphrasia officinalis</i> agg.	2
<i>Festuca ovina</i>	3
<i>Fissidens dubius</i>	1
<i>Geranium robertianum</i>	1
<i>Hypochaeris radicata</i>	2
<i>Linum catharticum</i>	2
<i>Lotus corniculatus</i>	5

Relevé code	EC37 R1
Species	Domin
<i>Pilosella officinarum</i>	2
<i>Plantago maritima</i>	2
<i>Polygala vulgaris</i>	3
<i>Sesleria caerulea</i>	7
<i>Taraxacum officinale</i> agg.	1
<i>Teucrium scorodonia</i>	3
<i>Thymus polytrichus</i>	3
<i>Tortella tortuosa</i>	2
<i>Trifolium pratense</i>	1

Relevé code	EC37 R2
Date	02/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	WN2/ER2
Habitat - Annex I	*8240
Species	Domin
<i>Brachypodium sylvaticum</i>	2
<i>Brachythecium rutabulum</i>	2
<i>Corylus avellana</i>	9
<i>Eurhynchium striatum</i>	3
<i>Fragaria vesca</i>	2
<i>Hedera helix</i>	8
<i>Ilex aquifolium</i>	3
<i>Neckera complanata</i>	1
<i>Neckera crispa</i>	4
<i>Prunus spinosa</i>	4
<i>Pteridium aquilinum</i>	2
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	4
<i>Thuidium tamariscinum</i>	3

Relevé code	EC37 R3
Date	02/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	2
<i>Corylus avellana</i>	4
<i>Crataegus monogyna</i>	1
<i>Ctenidium molluscum</i>	2
<i>Fissidens dubius</i>	3
<i>Fragaria vesca</i>	1
<i>Fraxinus excelsior</i>	1
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	2
<i>Mycelis muralis</i>	2
<i>Phyllitis scolopendrium</i>	2

Relevé code	EC37 R3
Species	Domin
<i>Prunus spinosa</i>	3
<i>Senecio jacobaea</i>	2
<i>Sesleria caerulea</i>	2
<i>Sonchus oleraceus</i>	2
<i>Teucrium scorodonia</i>	4
<i>Tortella tortuosa</i>	2
<i>Tussilago farfara</i>	2

Relevé code	EC37 R4
Date	02/07/2014
Recorder	JD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Species	Domin
<i>Anacamptis pyramidalis</i>	2
<i>Briza media</i>	3
<i>Carex flacca</i>	3
<i>Centaurea nigra</i>	7
<i>Cerastium fontanum</i>	2
<i>Cynosurus cristatus</i>	3
<i>Dactylis glomerata</i>	2
<i>Dactylorhiza fuchsii</i>	1
<i>Daucus carota</i>	2
<i>Festuca ovina</i>	3
<i>Festuca rubra</i>	6
<i>Holcus lanatus</i>	4
<i>Kindbergia praelonga</i>	2
<i>Lathyrus pratensis</i>	3
<i>Lotus corniculatus</i>	5
<i>Plantago lanceolata</i>	4
<i>Ranunculus acris</i>	3
<i>Rubus fruticosus</i> agg.	2
<i>Sesleria caerulea</i>	2
<i>Taraxacum officinale</i> agg.	1
<i>Trifolium dubium</i>	2
<i>Trifolium pratense</i>	3
<i>Trifolium repens</i>	4

Relevé code	EC37 R5
Date	30/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Achillea millefolium</i>	2
<i>Agrostis stolonifera</i>	4
<i>Centaurea nigra</i>	5
<i>Cerastium fontanum</i>	2

<b>Relevé code</b>	<b>EC37 R5</b>
Species	Domin
<i>Cirsium arvense</i>	4
<i>Dactylis glomerata</i>	5
<i>Daucus carota</i>	3
<i>Festuca rubra</i>	2
<i>Heracleum sphondylium</i>	3
<i>Holcus lanatus</i>	4
<i>Lathyrus pratensis</i>	4
<i>Lolium perenne</i>	2
<i>Odontites vernus</i>	2
<i>Phleum pratense</i>	3
<i>Plantago lanceolata</i>	3
<i>Poa pratensis</i>	2
<i>Potentilla anserina</i>	4
<i>Ranunculus acris</i>	3
<i>Trifolium pratense</i>	6
<i>Vicia cracca</i>	2

<b>Relevé code</b>	<b>EC37 R6</b>
Date	30/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Asplenium ruta-muraria</i>	2
<i>Briza media</i>	2
<i>Carlina acaulis</i>	3
<i>Cladonia portentosa</i>	3
<i>Ctenidium molluscum</i>	4
<i>Euphrasia sp.</i>	2
<i>Festuca rubra</i>	4
<i>Fragaria vesca</i>	3
<i>Hypnum cupressiforme</i>	5
<i>Lotus corniculatus</i>	4
<i>Neckera crispa</i>	2
<i>Pilosella officinarum</i>	3
<i>Plantago lanceolata</i>	2
<i>Polygala vulgaris</i>	2
<i>Sesleria albicans</i>	7
<i>Teucrium scorodonia</i>	4
<i>Thymus polytrichus</i>	2
<i>Tortella tortuosa</i>	2



Relevé code	EC37 R7
Date	30/09/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Carex flacca</i>	3
<i>Cotoneaster species</i>	3
<i>Crataegus monogyna</i>	3
<i>Euonymus europaeus</i>	4
<i>Festuca rubra</i>	4
<i>Fragaria vesca</i>	3
<i>Leycesteria formosa</i>	2
<i>Lonicera periclymenum</i>	3
<i>Prunus spinosa</i>	3
<i>Pseudoscleropodium purum</i>	3
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	7
<i>Sesleria albicans</i>	5
<i>Teucrium scorodonia</i>	5
<i>Thuidium tamariscinum</i>	3
<i>Viola</i> sp.	3

Relevé code	EC37 R8
Date	03/09/2015
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Achillea millefolium</i>	3
<i>Agrostis canina</i>	5
<i>Anagallis arvensis</i>	1
<i>Centaurea nigra</i>	4
<i>Cerastium fontanum</i>	1
<i>Corylus avellana</i>	1
<i>Dactylis glomerata</i>	4
<i>Festuca</i> sp.	4
<i>Koeleria macrantha</i>	4
<i>Leontodon hispidus</i>	1
<i>Lolium perenne</i>	4
<i>Odontites vernus</i>	3
<i>Pimpinella major</i>	4
<i>Plantago lanceolata</i>	4
<i>Potentilla anserina</i>	1
<i>Trifolium pratense</i>	4

<b>Relevé code</b>	<b>EC39 R1</b>
Date	07/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	ER2/WN2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	2
<i>Corylus avellana</i>	4
<i>Ctenidium molluscum</i>	2
<i>Epipactis helleborine</i>	1
<i>Fissidens dubius</i>	2
<i>Geranium robertianum</i>	1
<i>Hedera helix</i>	5
<i>Mycelis muralis</i>	1
<i>Prunus spinosa</i>	3
<i>Scapania aspera</i>	2
<i>Sesleria albicans</i>	2
<i>Teucrium scorodonia</i>	2
<i>Tortella tortuosa</i>	1

<b>Relevé code</b>	<b>EC39 R2</b>
Date	07/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	ER2/WN2
Habitat - Annex I	*8240
Species	Domin
<i>Corylus avellana</i>	9
<i>Crataegus monogyna</i>	3
<i>Ctenidium molluscum</i>	2
<i>Epipactis helleborine</i>	1
<i>Eurhynchium striatum</i>	8
<i>Hedera helix</i>	5
<i>Hypericum androsaemum</i>	1
<i>Lonicera periclymenum</i>	2
<i>Neckera complanata</i>	2
<i>Phyllitis scolopendrium</i>	3
<i>Prunus spinosa</i>	3
<i>Rhytidiadelphus triquetrus</i>	2
<i>Rubus fruticosus</i> agg.	3
<i>Thuidium tamariscinum</i>	4
<i>Viola</i> sp.	2

Relevé code	EC39 R3
Date	07/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	PF1
Habitat - Annex I	-
Species	Domin
<i>Agrostis stolonifera</i>	2
<i>Carex elata</i>	4
<i>Carex nigra</i>	6
<i>Carex panicea</i>	7
<i>Epilobium palustre</i>	1
<i>Filipendula ulmaria</i>	2
<i>Hydrocotyle vulgaris</i>	4
<i>Juncus acutiflorus/articulatus</i>	2
<i>Menyanthes trifoliata</i>	3
<i>Molinia caerulea</i>	4
<i>Myrica gale</i>	3
<i>Phragmites australis</i>	3
<i>Ranunculus flammula</i>	1
<i>Schoenus nigricans</i>	5

Relevé code	EC39 R4
Date	22/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	ER2/GS1
Habitat - Annex I	-
Species	Domin
<i>Blackstonia perfoliata</i>	2
<i>Brachythecium rutabulum</i>	2
<i>Carex flacca</i>	6
<i>Centaureum erythraea</i>	2
<i>Cerastium fontanum</i>	2
<i>Cirsium arvense</i>	2
<i>Crepis capillaris</i>	2
<i>Daucus carota</i>	3
<i>Didymodon</i> sp.	5
<i>Holcus lanatus</i>	3
<i>Hypochaeris radicata</i>	2
<i>Leucanthemum vulgare</i>	4
<i>Odontites vernus</i>	4
<i>Prunella vulgaris</i>	4
<i>Pteridium aquilinum</i>	2
<i>Ranunculus acris</i>	4
<i>Rubus fruticosus</i> agg.	4
<i>Sagina nodosa</i>	1
<i>Senecio jacobaea</i>	2
<i>Torilis</i> sp.	1
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	4

<b>Relevé code</b>	<b>EC39 R5</b>
Date	22/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Agrostis capillaris</i>	4
<i>Anthoxanthum odoratum</i>	3
<i>Blackstonia perfoliata</i>	2
<i>Briza media</i>	5
<i>Carex flacca</i>	4
<i>Centaurea nigra</i>	2
<i>Centaureum erythraea</i>	1
<i>Dactylis glomerata</i>	1
<i>Daucus carota</i>	2
<i>Festuca rubra</i>	5
<i>Galium verum</i>	3
<i>Hypochaeris radicata</i>	2
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	3
<i>Medicago lupulina</i>	3
<i>Odontites vernus</i>	2
<i>Plantago lanceolata</i>	5
<i>Polygala vulgaris</i>	1
<i>Potentilla erecta</i>	2
<i>Prunella vulgaris</i>	1
<i>Senecio jacobaea</i>	2
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	3
<i>Vicia cracca</i>	1

<b>Relevé code</b>	<b>EC39 R6</b>
Date	24/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Carex disticha</i>	2
<i>Carex flacca</i>	2
<i>Carex ovalis</i>	2
<i>Carex panicea</i>	2
<i>Epilobium palustre</i>	1
<i>Filipendula ulmaria</i>	4
<i>Juncus conglomeratus</i>	4
<i>Molinia caerulea</i>	9
<i>Poa trivialis</i>	3
<i>Potentilla erecta</i>	4
<i>Vicia cracca</i>	2

<b>Relevé code</b>	<b>EC39 R7</b>
Species	Domin
Date	24/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
<i>Agrostis stolonifera</i>	4
<i>Calliergonella cuspidata</i>	8
<i>Carex nigra</i>	7
<i>Carex panicea</i>	4
<i>Carex viridula</i>	2
<i>Galium palustre</i>	2
<i>Hydrocotyle vulgaris</i>	5
<i>Juncus articulatus</i>	6
<i>Mentha aquatica</i>	4
<i>Potentilla anserina</i>	3
<i>Prunella vulgaris</i>	3
<i>Ranunculus flammula</i>	3

<b>Relevé code</b>	<b>EC39 R8</b>
Date	24/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Calliergonella cuspidata</i>	6
<i>Carex nigra</i>	4
<i>Carex panicea</i>	4
<i>Cirsium dissectum</i>	3
<i>Filipendula ulmaria</i>	4
<i>Hydrocotyle vulgaris</i>	5
<i>Juncus articulatus</i>	5
<i>Mentha aquatica</i>	2
<i>Molinia caerulea</i>	7
<i>Myrica gale</i>	6
<i>Potentilla anserina</i>	3
<i>Potentilla erecta</i>	3
<i>Ranunculus flammula</i>	2

<b>Relevé code</b>	<b>EC39 R9</b>
Date	24/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	PF1
Habitat - Annex I	7230
<b>Species</b>	<b>Domin</b>
<i>Carex lepidocarpa</i>	4
<i>Cirsium dissectum</i>	2
<i>Cladium mariscus</i>	4
<i>Hydrocotyle vulgaris</i>	4
<i>Juncus articulatus</i>	4
<i>Juncus subnodulosus</i>	4
<i>Molinia caerulea</i>	7
<i>Myrica gale</i>	5
<i>Parnassia palustris</i>	1
<i>Schoenus nigricans</i>	6

<b>Relevé code</b>	<b>EC39 R10</b>
Date	22/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
<b>Species</b>	<b>Domin</b>
<i>Agrostis stolonifera</i>	3
<i>Calliergonella cuspidata</i>	6
<i>Carex nigra</i>	8
<i>Carex panicea</i>	3
<i>Epilobium palustre</i>	1
<i>Filipendula ulmaria</i>	3
<i>Galium palustre</i>	3
<i>Hydrocotyle vulgaris</i>	3
<i>Juncus acutiflorus/articulatus</i>	4
<i>Mentha aquatica</i>	2
<i>Molinia caerulea</i>	6
<i>Potentilla anserina</i>	4
<i>Ranunculus flammula</i>	3
<i>Triglochin palustre</i>	2

<b>Relevé code</b>	<b>EC39 R11</b>
Date	27/08/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Anthoxanthum odoratum</i>	4
<i>Anthyllis vulneraria</i>	6
<i>Briza media</i>	3
<i>Calliergonella cuspidata</i>	5
<i>Carex flacca</i>	4
<i>Carex panicea</i>	3
<i>Centaurea nigra</i>	5
<i>Ctenidium molluscum</i>	5
<i>Daucus carota</i>	2
<i>Festuca rubra</i>	5
<i>Helictotrichon pubescens</i>	2
<i>Homalothecium lutescens</i>	2
<i>Hypericum perforatum</i>	?
<i>Hypochaeris radicata</i>	4
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	4
<i>Medicago lupulina</i>	2
<i>Pilosella officinarum</i>	3
<i>Pseudoscleropodium purum</i>	3
<i>Pteridium aquilinum</i>	3
<i>Rhytidiadelphus triquetrus</i>	3
<i>Rubus fruticosus agg.</i>	3
<i>Rumex acetosa</i>	3
<i>Senecio jacobaea</i>	2
<i>Sesleria albicans</i>	2
<i>Teucrium scorodonia</i>	3
<i>Trifolium pratense</i>	2
<i>Trifolium repens</i>	3
<i>Trisetum flavescens</i>	3

<b>Relevé code</b>	<b>EC39 R12</b>
Date	24/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	FS1/FS2
Habitat - Annex I	-
Species	Domin
<i>Agrostis stolonifera</i>	3
<i>Alisma plantago-aquatica</i>	4
<i>Carex lasiocarpa</i>	2
<i>Carex pseudocyperus</i>	2
<i>Carex rostrata</i>	2
<i>Drepanocladus aduncus</i>	9
<i>Eleocharis palustris</i>	6
<i>Equisetum fluviatile</i>	5
<i>Galium palustre</i>	3
<i>Hippuris vulgaris</i>	3
<i>Menyanthes trifoliata</i>	8

<b>Relevé code</b>	<b>EC39 R13</b>
Date	14/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS2
Habitat - Annex I	6510
Species	Domin
<i>Calliergonella cuspidata</i>	5
<i>Carex flacca</i>	2
<i>Cerastium fontanum</i>	1
<i>Daucus carota</i>	1
<i>Festuca rubra</i>	5
<i>Helictotrichon pubescens</i>	5
<i>Holcus lanatus</i>	4
<i>Leontodon hispidus</i>	3
<i>Leucanthemum vulgare</i>	3
<i>Odontites vernus</i>	3
<i>Pimpinella major</i>	2
<i>Plantago lanceolata</i>	4
<i>Potentilla anglica</i>	5
<i>Trifolium dubium</i>	3
<i>Trifolium pratense</i>	6
<i>Trisetum flavescens</i>	3



<b>Relevé code</b>	<b>EC39 R14</b>
Date	03/11/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Agrostis stolonifera</i>	3
<i>Carex panicea</i>	4
<i>Carex</i> sp.	5
<i>Filipendula ulmaria</i>	4
<i>Hydrocotyle vulgaris</i>	3
<i>Juncus articulatus</i>	5
<i>Molinia caerulea</i>	7
<i>Nasturtium officinale</i>	2
<i>Potentilla erecta</i>	1
<i>Ranunculus flammula</i>	1
<i>Schoenus nigricans</i>	5
<i>Succisa pratensis</i>	4

<b>Relevé code</b>	<b>EC39 R15</b>
Date	03/11/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	6410
Species	Domin
<i>Carex lasiocarpa</i>	4
<i>Carex panicea</i>	5
<i>Eriophorum</i> sp.	4
<i>Filipendula ulmaria</i>	3
<i>Hydrocotyle vulgaris</i>	1
<i>Juncus articulatus</i>	4
<i>Juncus bufonius</i>	3
<i>Juncus conglomeratus</i>	4
<i>Molinia caerulea</i>	5
<i>Nasturtium officinale</i>	1
<i>Potentilla erecta</i>	2
<i>Rumex acetosa</i>	2

<b>Relevé code</b>	<b>EC42 R1</b>
Date	25/06/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	ER2/WN2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	2
<i>Carlina vulgaris</i>	1
<i>Ceterach officinarum</i>	2

Relevé code	EC42 R1
Species	Domin
<i>Fragaria vesca</i>	2
<i>Fraxinus excelsior</i>	1
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	1
<i>Hieracium</i> sp.	2
<i>Linum catharticum</i>	1
<i>Lotus corniculatus</i>	1
<i>Prunus spinosa</i>	3
<i>Senecio jacobaea</i>	1
<i>Sesleria caerulea</i>	4
<i>Succisa pratensis</i>	1
<i>Teucrium scorodonia</i>	3
<i>Thymus polytrichus</i>	1

Relevé code	EC42 R2
Date	25/06/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	ER2/WN2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	1
<i>Carex flacca</i>	2
<i>Corylus avellana</i>	6
<i>Geranium robertianum</i>	3
<i>Neckera crispa</i>	3
<i>Polystichum aculeatum</i>	1
<i>Prunus spinosa</i>	3
<i>Pteridium aquilinum</i>	4
<i>Rosa spinosissima</i>	3
<i>Rubia peregrina</i>	1
<i>Rubus fruticosus</i> agg.	3
<i>Sesleria caerulea</i>	3
<i>Teucrium scorodonia</i>	2
<i>Tortella</i> sp.	2

Relevé code	EC42 R3
Date	25/06/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	WN2/ER2/WS1
Habitat - Annex I	*8240
Species	Domin
<i>Brachypodium sylvaticum</i>	4
<i>Brachythecium rutabulum</i>	5
<i>Corylus avellana</i>	8
<i>Crataegus monogyna</i>	2
<i>Fragaria vesca</i>	3
<i>Fraxinus excelsior</i>	5
<i>Geum urbanum</i>	2

Relevé code	EC42 R3
Species	Domin
<i>Hedera helix</i>	5
<i>Ilex aquifolium</i>	3
<i>Lonicera periclymenum</i>	3
<i>Neckera crispa</i>	3
<i>Phyllitis scolopendrium</i>	2
<i>Potentilla sterilis</i>	3
<i>Primula vulgaris</i>	3
<i>Prunus spinosa</i>	2
<i>Rubus fruticosus</i> agg.	3
<i>Taraxacum officinale</i> agg.	2
<i>Thamnobryum alopecurum</i>	7
<i>Thuidium tamariscinum</i>	3
<i>Viola</i> sp.	1

Relevé code	EC42 R4
Date	02/09/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	GS1/WS1
Habitat - Annex I	6210
Species	Domin
<i>Briza media</i>	5
<i>Calliergonella cuspidata</i>	5
<i>Carex flacca</i>	5
<i>Centaurea nigra</i>	4
<i>Ctenidium molluscum</i>	3
<i>Danthonia decumbens</i>	2
<i>Daucus carota</i>	1
<i>Euphrasia</i> sp.	4
<i>Festuca rubra</i>	3
<i>Galium verum</i>	1
<i>Holcus lanatus</i>	3
<i>Hypericum maculatum</i>	3
<i>Hypericum pulchrum</i>	2
<i>Hypochaeris radicata</i>	3
<i>Lathyrus pratensis</i>	4
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	2
<i>Lotus corniculatus</i>	5
<i>Molinia caerulea</i>	5
<i>Potentilla erecta</i>	3
<i>Potentilla sterilis</i>	2
<i>Prunella vulgaris</i>	1
<i>Scleropodium purum</i>	4
<i>Thuidium tamariscinum</i>	3
<i>Thymus polytrichus</i>	2
<i>Trifolium pratense</i>	3
<i>Veronica chamaedrys</i>	1

Relevé code	EC42 R5
Date	02/09/2014
Recorder	JC
Species recording scale	Domin
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Domin
<i>Asplenium ruta-muraria</i>	3
<i>Bryum</i> sp.	3
<i>Carlina vulgaris</i>	3
<i>Ceterach officinarum</i>	1
<i>Corylus avellana</i>	5
<i>Epipactis</i> sp.	1
<i>Festuca rubra</i>	2
<i>Galium verum</i>	1
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	4
<i>Hieracium</i> sp.	2
<i>Leucanthemum vulgare</i>	2
<i>Linum catharticum</i>	1
<i>Lonicera periclymenum</i>	3
<i>Lotus corniculatus</i>	1
<i>Molinia caerulea</i>	1
<i>Mycelis muralis</i>	1
<i>Phyllitis scolopendrium</i>	1
<i>Plantago lanceolata</i>	2
<i>Prunus spinosa</i>	3
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	1
<i>Salix cinerea</i>	1
<i>Senecio jacobaea</i>	1
<i>Sesleria albicans</i>	4
<i>Teucrium scorodonia</i>	3
<i>Thymus polytrichus</i>	3

Relevé code	EC44 R1
Date	16/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Brachypodium sylvaticum</i>	1
<i>Corylus avellana</i>	10
<i>Crataegus monogyna</i>	2
<i>Dryopteris filix-mas</i>	?
<i>Eurhynchium striatum</i>	9
<i>Fragaria vesca</i>	3
<i>Hedera helix</i>	7
<i>Ilex aquifolium</i>	2
<i>Lonicera periclymenum</i>	4
<i>Neckera complanata</i>	3
<i>Prunus spinosa</i>	1

Relevé code	EC44 R1
Species	Domin
<i>Rhytiadelphus triquetrus</i>	1
<i>Rubus fruticosus</i> agg.	3
<i>Sorbus aucuparia</i>	2
<i>Thuidium tamariscinum</i>	2
<i>Viola</i> sp.	2

Relevé code	EC44 R2
Date	16/07/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Brachypodium sylvaticum</i>	3
<i>Calliergonella cuspidata</i>	2
<i>Carex sylvatica</i>	4
<i>Circaea lutetiana</i>	4
<i>Corylus avellana</i>	10
<i>Crataegus monogyna</i>	3
<i>Dryopteris filix-mas</i>	2
<i>Euonymus europaeus</i>	2
<i>Eurhynchium striatum</i>	4
<i>Filipendula ulmaria</i>	2
<i>Fissidens</i> sp.	2
<i>Fragaria vesca</i>	3
<i>Geum urbanum</i>	4
<i>Hedera helix</i>	4
<i>Hypericum androsaemum</i>	2
<i>Ilex aquifolium</i>	3
<i>Kindbergia praelonga</i>	2
<i>Neckera complanata</i>	2
<i>Pellia</i> sp.	1
<i>Plagiomnium undulatum</i>	3
<i>Prunella vulgaris</i>	4
<i>Prunus spinosa</i>	3
<i>Rubus fruticosus</i> agg.	3
<i>Rumex sanguineus</i>	3
<i>Thamnobryum alopecurum</i>	4
<i>Thuidium tamariscinum</i>	4
<i>Veronica chamaedrys</i>	3
<i>Viola</i> sp.	3

Relevé code	EC44 R4
Date	19/08/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Anthoxanthum odoratum</i>	3
<i>Blackstonia perfoliata</i>	1
<i>Briza media</i>	4
<i>Calliergonella cuspidata</i>	?
<i>Carex flacca</i>	5
<i>Centaurea nigra</i>	4
<i>Dactylis glomerata</i>	2
<i>Daucus carota</i>	3
<i>Euphrasia</i> sp.	4
<i>Festuca rubra</i>	4
<i>Holcus lanatus</i>	4
<i>Hypericum androsaemum</i>	1
<i>Leucanthemum vulgare</i>	2
<i>Linum catharticum</i>	3
<i>Lotus corniculatus</i>	5
<i>Potentilla reptans</i>	2
<i>Rubus fruticosus</i> agg.	3

Relevé code	EC47 R1
Date	26/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Corylus avellana</i>	6
<i>Crataegus monogyna</i>	5
<i>Dryopteris filix-mas</i>	2
<i>Eurhynchium striatum</i>	7
<i>Fragaria vesca</i>	3
<i>Fraxinus excelsior</i>	8
<i>Hedera helix</i>	6
<i>Phyllitis scolopendrium</i>	4
<i>Primula veris</i>	2
<i>Rhytidiadelphus triquetrus</i>	5
<i>Rubus fruticosus</i> agg.	4
<i>Scapania</i> sp.	3
<i>Thamnobryum alopecurum</i>	7
<i>Thuidium tamariscinum</i>	5
<i>Veronica chamaedrys</i>	2
<i>Viola</i> sp.	2

<b>Relevé code</b>	<b>EC56 R1</b>
Date	13/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Anagallis tenella</i>	2
<i>Antennaria dioica</i>	3
<i>Blackstonia perfoliata</i>	3
<i>Briza media</i>	3
<i>Calliergonella cuspidata</i>	4
<i>Carex demissa</i>	3
<i>Carex flacca</i>	?
<i>Carex panicea</i>	3
<i>Carex pulicaris</i>	1
<i>Centaurea nigra</i>	3
<i>Cirsium dissectum</i>	1
<i>Ctenidium molluscum</i>	3
<i>Cynosurus cristatus</i>	1
<i>Danthonia decumbens</i>	2
<i>Didymodon ferrugineus</i>	1
<i>Euphrasia sp.</i>	2
<i>Festuca ovina</i>	4
<i>Fissidens adianthoides</i>	2
<i>Gentiana verna</i>	1
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	1
<i>Lotus corniculatus</i>	4
<i>Molinia caerulea</i>	3
<i>Pinguicula vulgaris</i>	3
<i>Plantago maritima</i>	4
<i>Polygala vulgaris</i>	3
<i>Potentilla erecta</i>	3
<i>Prunella vulgaris</i>	1
<i>Rhinanthus minor</i>	2
<i>Schoenus nigricans</i>	3
<i>Selaginella selaginoides</i>	3
<i>Sesleria albicans</i>	2
<i>Succisa pratensis</i>	5
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	3

<b>Relevé code</b>	<b>EC56 R2</b>
Date	13/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	ER2/WS1
Habitat - Annex I	*8240
Species	Domin
<i>Achillea millefolium</i>	1
<i>Agrostis capillaris</i>	1
<i>Antennaria dioica</i>	1
<i>Anthoxanthum odoratum</i>	2
<i>Asplenium ruta-muraria</i>	2
<i>Briza media</i>	3
<i>Carex caryophylla</i>	1
<i>Carex flacca</i>	2
<i>Carlina vulgaris</i>	1
<i>Centaurea nigra</i>	2
<i>Corylus avellana</i>	5
<i>Ctenidium molluscum</i>	3
<i>Dactylorhiza fuchsii</i>	1
<i>Festuca rubra</i>	3
<i>Fissidens dubius</i>	1
<i>Gentiana verna</i>	1
<i>Geranium robertianum</i>	1
<i>Hedera helix</i>	4
<i>Hypericum pulchrum</i>	1
<i>Hypochaeris radicata</i>	1
<i>Juniperus communis</i>	3
<i>Koeleria macrantha</i>	2
<i>Leucanthemum vulgare</i>	2
<i>Lotus corniculatus</i>	2
<i>Mycelis muralis</i>	1
<i>Neckera crispa</i>	2
<i>Pilosella officinarum</i>	3
<i>Plantago lanceolata</i>	3
<i>Plantago major</i>	2
<i>Polygala vulgaris</i>	2
<i>Prunella vulgaris</i>	1
<i>Pteridium aquilinum</i>	2
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	3
<i>Sesleria albicans</i>	3
<i>Succisa pratensis</i>	1
<i>Taraxacum officinale</i> agg.	3
<i>Teucrium scorodonia</i>	2
<i>Thymus praecox</i>	1
<i>Tortella tortuosa</i>	3
<i>Trifolium pratense</i>	3
<i>Ulex europaeus</i>	4
<i>Viola</i> sp.	2



<b>Relevé code</b>	<b>EC56 R3</b>
Date	13/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Agrostis capillaris</i>	1
<i>Anthoxanthum odoratum</i>	3
<i>Breutelia chrysocoma</i>	2
<i>Briza media</i>	3
<i>Calluna vulgaris</i>	2
<i>Campylium protensum</i>	2
<i>Carex caryophylla</i>	3
<i>Carex flacca</i>	3
<i>Carex panicea</i>	3
<i>Centaurea nigra</i>	4
<i>Ctenidium molluscum</i>	4
<i>Cynosurus cristatus</i>	2
<i>Dactylorhiza fuchsii</i>	1
<i>Dicranum scoparium</i>	1
<i>Festuca ovina</i>	4
<i>Fissidens adianthoides</i>	1
<i>Galium verum</i>	3
<i>Helictotrichon pratense</i>	2
<i>Helictotrichon pubescens</i>	2
<i>Hypericum pulchrum</i>	1
<i>Hypochaeris radicata</i>	2
<i>Koeleria macrantha</i>	3
<i>Lathyrus linifolius</i>	1
<i>Linum catharticum</i>	2
<i>Lotus corniculatus</i>	3
<i>Pilosella officinarum</i>	3
<i>Plantago lanceolata</i>	4
<i>Plantago maritima</i>	3
<i>Polygala vulgaris</i>	2
<i>Potentilla erecta</i>	2
<i>Prunella vulgaris</i>	2
<i>Pseudoscleropodium purum</i>	1
<i>Pteridium aquilinum</i>	3
<i>Rhinanthus minor</i>	2
<i>Rosa spinosissima</i>	4
<i>Senecio jacobaea</i>	1
<i>Sesleria caerulea</i>	3
<i>Succisa pratensis</i>	5
<i>Taraxacum officinale</i> agg.	3
<i>Thuidium delicatulum</i>	1
<i>Thymus praecox</i>	3
<i>Tortella tortuosa</i>	3
<i>Trifolium repens</i>	2
<i>Viola</i> sp.	2

<b>Relevé code</b>	<b>EC56 R4</b>
Date	17/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Achillea millefolium</i>	3
<i>Anthoxanthum odoratum</i>	4
<i>Bellis perennis</i>	3
<i>Briza media</i>	4
<i>Carex caryophylla</i>	4
<i>Centaurea nigra</i>	5
<i>Cynosurus cristatus</i>	3
<i>Dactylorhiza fuchsii</i>	1
<i>Danthonia decumbens</i>	2
<i>Euphrasia</i> sp.	4
<i>Festuca rubra</i>	3
<i>Helictotrichon pubescens</i>	2
<i>Holcus lanatus</i>	3
<i>Hypochaeris radicata</i>	3
<i>Lathyrus pratensis</i>	?
<i>Lotus corniculatus</i>	2
<i>Luzula campestris</i>	2
<i>Pilosella officinarum</i>	3
<i>Plantago lanceolata</i>	3
<i>Potentilla anserina</i>	1
<i>Pteridium aquilinum</i>	5
<i>Ranunculus bulbosus</i>	4
<i>Rubus fruticosus</i> agg.	1
<i>Senecio jacobaea</i>	2
<i>Trifolium pratense</i>	3

Relevé code	EC56 R5
Date	17/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Domin
<i>Anthoxanthum odoratum</i>	2
<i>Corylus avellana</i>	1
<i>Crepis capillaris</i>	2
<i>Cynosurus cristatus</i>	2
<i>Festuca rubra</i>	2
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	5
<i>Plantago lanceolata</i>	2
<i>Prunus spinosa</i>	4
<i>Rosa spinosissima</i>	2
<i>Rubus fruticosus</i> agg.	3
<i>Sonchus asper</i>	1
<i>Teucrium scorodonia</i>	3
<i>Tortella tortuosa</i>	2
<i>Trifolium dubium</i>	2
<i>Trisetum flavescens</i>	2

Relevé code	EC56 R6
Date	17/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WN2/ER2
Habitat - Annex I	*8240
Species	Domin
<i>Arum maculatum</i>	2
<i>Corylus avellana</i>	9
<i>Eurhynchium striatum</i>	5
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	4
<i>Lonicera periclymenum</i>	3
<i>Neckera crispa</i>	3
<i>Phyllitis scolopendrium</i>	3
<i>Prunus spinosa</i>	2
<i>Rubus fruticosus</i> agg.	3
<i>Thamnobryum alopecurum</i>	7
<i>Thuidium tamariscinum</i>	5
<i>Veronica montana</i>	2

<b>Relevé code</b>	<b>EC56 R7</b>
Date	28/08/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Achillea millefolium</i>	3
<i>Campanula rotundifolia</i>	2
<i>Carex flacca</i>	3
<i>Centaurea nigra</i>	5
<i>Centaurium erythraea</i>	2
<i>Dactylorhiza</i> sp.	1
<i>Daucus carota</i>	3
<i>Euphrasia</i> sp.	3
<i>Festuca rubra</i>	4
<i>Galium verum</i>	2
<i>Holcus lanatus</i>	4
<i>Hypochaeris radicata</i>	3
<i>Lathyrus pratensis</i>	3
<i>Linum catharticum</i>	2
<i>Lotus corniculatus</i>	2
<i>Odontites vernus</i>	2
<i>Pilosella officinarum</i>	2
<i>Plantago lanceolata</i>	6
<i>Potentilla anserina</i>	3
<i>Prunella vulgaris</i>	4
<i>Rosa spinosissima</i>	2
<i>Rubus fruticosus</i> agg.	3
<i>Solidago virgaurea</i>	2
<i>Succisa pratensis</i>	2
<i>Thymus praecox</i>	2
<i>Trifolium pratense</i>	4
<i>Trifolium repens</i>	4

<b>Relevé code</b>	<b>EC56 R8</b>
Date	27/07/2016
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1/GS2
Habitat - Annex I	-
Species	Domin
<i>Anthoxanthum odoratum</i>	4
<i>Centaurea nigra</i>	2
<i>Cynosurus cristatus</i>	5
<i>Festuca rubra</i>	4
<i>Holcus lanatus</i>	5
<i>Homalothecium lutescens</i>	?
<i>Lathyrus pratensis</i>	3
<i>Plantago lanceolata</i>	4
<i>Rubus fruticosus</i> agg.	4
<i>Rumex acetosa</i>	1
<i>Senecio jacobaea</i>	1
<i>Succisa pratensis</i>	3
<i>Taraxacum officinale</i> agg.	3
<i>Trifolium pratense</i>	3
<i>Trifolium repens</i>	1
<i>Tussilago farfara</i>	2

<b>Relevé code</b>	<b>EC56 R9</b>
Date	12/10/2017
Recorder	KMOC
Species recording scale	DAFOR
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	DAFOR
<i>Achillea millefolium</i>	O
<i>Agrostis stolonifera</i>	A
<i>Cynosurus cristatus</i>	F
<i>Dactylis glomerata</i>	O
<i>Festuca rubra</i>	D
<i>Knautia arvensis</i>	F
<i>Leontodon</i> sp.	O
<i>Plantago lanceolata</i>	A-D
<i>Potentilla anserina</i>	O
<i>Ranunculus repens</i>	F
<i>Rumex</i> sp.	O
<i>Trifolium pratense</i>	F

<b>Relevé code</b>	<b>EC57 R1</b>
Date	17/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Antennaria dioica</i>	3
<i>Anthoxanthum odoratum</i>	2
<i>Anthyllis vulneraria</i>	5
<i>Briza media</i>	4
<i>Carex caryophylla</i>	4
<i>Carex flacca</i>	4
<i>Carex panicea</i>	3
<i>Carlina vulgaris</i>	2
<i>Centaurea nigra</i>	3
<i>Dactylorhiza fuchsii</i>	2
<i>Danthonia decumbens</i>	2
<i>Euphrasia sp.</i>	3
<i>Festuca rubra</i>	3
<i>Galium verum</i>	2
<i>Koeleria macrantha</i>	2
<i>Leontodon hispidus</i>	4
<i>Linum catharticum</i>	4
<i>Lotus corniculatus</i>	3
<i>Pilosella officinarum</i>	3
<i>Plantago maritima</i>	2
<i>Polygala vulgaris</i>	3
<i>Potentilla erecta</i>	3
<i>Rosa spinosissima</i>	2
<i>Sesleria albicans</i>	2
<i>Succisa pratensis</i>	2

Relevé code	EC57 R2
Date	18/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Anthoxanthum odoratum</i>	4
<i>Briza media</i>	4
<i>Carex caryophylla</i>	4
<i>Carex flacca</i>	3
<i>Carex panicea</i>	6
<i>Carex pulicaris</i>	2
<i>Centaurea nigra</i>	3
<i>Conopodium majus</i>	2
<i>Cynosurus cristatus</i>	4
<i>Dactylis glomerata</i>	2
<i>Galium verum</i>	2
<i>Holcus lanatus</i>	3
<i>Juncus bulbosus</i>	3
<i>Lotus corniculatus</i>	2
<i>Luzula campestris</i>	2
<i>Pilosella officinarum</i>	1
<i>Potentilla erecta</i>	4
<i>Pteridium aquilinum</i>	4
<i>Ranunculus acris</i>	2
<i>Succisa pratensis</i>	6
<i>Thymus praecox</i>	1

Relevé code	EC57 R3
Date	18/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Arum maculatum</i>	2
<i>Brachypodium sylvaticum</i>	3
<i>Circaea lutetiana</i>	2
<i>Corylus avellana</i>	9
<i>Crataegus monogyna</i>	3
<i>Dryopteris filix-mas</i>	2
<i>Fragaria vesca</i>	2
<i>Fraxinus excelsior</i>	1
<i>Geum urbanum</i>	1
<i>Hedera helix</i>	3
<i>Lonicera periclymenum</i>	3
<i>Phyllitis scolopendrium</i>	1
<i>Prunus spinosa</i>	1
<i>Rubus fruticosus</i> agg.	3
<i>Taraxacum officinale</i> agg.	2

<i>Viola</i> sp.	2
------------------	---

<b>Relevé code</b>	<b>EC57 R4</b>
Date	19/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Species	Domin
<i>Agrostis capillaris</i>	2
<i>Antennaria dioica</i>	2
<i>Anthoxanthum odoratum</i>	3
<i>Briza media</i>	4
<i>Calluna vulgaris</i>	2
<i>Carex caryophylla</i>	3
<i>Carex flacca</i>	3
<i>Carex panicea</i>	3
<i>Carex pulicaris</i>	3
<i>Carlina vulgaris</i>	1
<i>Centaurea nigra</i>	4
<i>Dactylorhiza fuchsii</i>	2
<i>Danthonia decumbens</i>	1
<i>Festuca rubra</i>	3
<i>Galium verum</i>	3
<i>Hypericum pulchrum</i>	2
<i>Koeleria macrantha</i>	3
<i>Lathyrus linifolius</i>	2
<i>Linum catharticum</i>	3
<i>Listera ovata</i>	2
<i>Lotus corniculatus</i>	3
<i>Molinia caerulea</i>	4
<i>Pedicularis palustris</i>	2
<i>Pilosella officinarum</i>	2
<i>Plantago lanceolata</i>	3
<i>Plantago maritima</i>	2
<i>Platanthera bifolia</i>	2
<i>Polygala vulgaris</i>	2
<i>Potentilla erecta</i>	3
<i>Pteridium aquilinum</i>	4
<i>Sesleria albicans</i>	2
<i>Succisa pratensis</i>	7
<i>Thymus praecox</i>	3



<b>Relevé code</b>	<b>EC57 R5</b>
Date	19/06/2014
Recorder	KD
Species recording scale	Domin
Habitat - Fossitt	WS1/ER2
Habitat - Annex I	*8240
Species	Domin
<i>Brachypodium sylvaticum</i>	2
<i>Corylus avellana</i>	4
<i>Crataegus monogyna</i>	4
<i>Ctenidium molluscum</i>	3
<i>Fragaria vesca</i>	2
<i>Fraxinus excelsior</i>	2
<i>Hedera helix</i>	3
<i>Lathyrus linifolius</i>	1
<i>Lonicera periclymenum</i>	2
<i>Molinia caerulea</i>	2
<i>Prunus spinosa</i>	7
<i>Pteridium aquilinum</i>	4
<i>Rosa spinosissima</i>	4
<i>Rubus fruticosus</i> agg.	5
<i>Scleropodium purum</i>	1
<i>Sesleria albicans</i>	3
<i>Solidago virgaurea</i>	2
<i>Teucrium scorodonia</i>	2
<i>Thuidium tamariscinum</i>	2
<i>Thymus praecox</i>	2
<i>Ulex europaeus</i>	3

<b>Relevé code</b>	<b>LQ-N84 R1</b>
Date	27/07/2016
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Achillea millefolium</i>	3
<i>Anthoxanthum odoratum</i>	4
<i>Anthyllis vulneraria</i>	4
<i>Blackstonia perfoliata</i>	3
<i>Briza media</i>	5
<i>Calliergonella cuspidata</i>	5
<i>Carex flacca</i>	1
<i>Carex panicea</i>	4
<i>Centaurea nigra</i>	5
<i>Centaureum erythraea</i>	2
<i>Festuca rubra</i>	5
<i>Hypericum pulchrum</i>	1
<i>Leontodon hispidus</i>	1
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	1
<i>Lotus corniculatus</i>	3
<i>Pimpinella major</i>	4
<i>Plantago lanceolata</i>	4
<i>Potentilla anserina</i>	1
<i>Prunella vulgaris</i>	1
<i>Rubus fruticosus</i> agg.	3
<i>Taraxacum officinale</i> agg.	1
<i>Trifolium pratense</i>	1
<i>Tussilago farfara</i>	2

<b>Relevé code</b>	<b>LQ-N84 R2</b>
Date	14/08/2014
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Domin
<i>Agrostis capillaris</i>	5
<i>Alchemilla vulgaris</i>	1
<i>Anthoxanthum odoratum</i>	6
<i>Calliergonella cuspidata</i>	4
<i>Calystegia sepium</i>	1
<i>Carex flacca</i>	5
<i>Cirsium dissectum</i>	1
<i>Cotoneaster</i> sp.	3
<i>Ctenidium molluscum</i>	3
<i>Daucus carota</i>	3
<i>Helictotrichon pubescens</i>	3
<i>Hypericum pulchrum</i>	3
<i>Knautia arvensis</i>	4
<i>Lathyrus pratensis</i>	3
<i>Leucanthemum vulgare</i>	4
<i>Lotus corniculatus</i>	2
<i>Pimpinella major</i>	4
<i>Plantago lanceolata</i>	3
<i>Potentilla anglica</i>	3
<i>Potentilla anserina</i>	3
<i>Ranunculus repens</i>	3
<i>Rubus fruticosus</i> agg.	2
<i>Solidago virgaurea</i>	3
<i>Teucrium scorodonia</i>	1

<b>Relevé code</b>	<b>RC-LQ R1</b>
Date	01/06/2016
Recorder	JF
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Agrostis capillaris</i>	5
<i>Anthoxanthum odoratum</i>	5
<i>Bellis perennis</i>	1
<i>Calliergonella cuspidata</i>	3
<i>Carex flacca</i>	0.5
<i>Cerastium fontanum</i>	1
<i>Cirsium arvense</i>	3
<i>Cynosurus cristatus</i>	5
<i>Festuca rubra</i>	20
<i>Holcus lanatus</i>	10
<i>Lolium perenne</i>	5
<i>Luzula campestris</i>	1
<i>Medicago lupulina</i>	1
<i>Plantago lanceolata</i>	10
<i>Prunella vulgaris</i>	1
<i>Ranunculus acris</i>	7
<i>Ranunculus repens</i>	3
<i>Rhynchospora squarrosa</i>	15
<i>Rumex obtusifolius</i>	3
<i>Taraxacum officinale</i> agg.	1
<i>Tortella tortuosa</i>	1
<i>Trifolium pratense</i>	7
<i>Trifolium repens</i>	1

<b>Relevé code</b>	<b>RC-LQ R3</b>
Date	09/10/2017
Recorder	KMOC
Species recording scale	Domin
Habitat - Fossitt	GS4
Habitat - Annex I	-
Species	Domin
<i>Arrhenatherum elatius</i>	5
<i>Filipendula ulmaria</i>	6
<i>Holcus lanatus</i>	5
<i>Juncus effusus</i>	7
<i>Lychnis flos-cuculi</i>	5
<i>Lythrum salicaria</i>	5
<i>Molinia caerulea</i>	8
<i>Plantago lanceolata</i>	5
<i>Potentilla anserina</i>	5
<i>Ranunculus acris</i>	3
<i>Ranunculus repens</i>	4
<i>Trifolium repens</i>	5

<b>Relevé code</b>	<b>RC-LQ R4</b>
Date	03/11/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Domin
<i>Carex flacca</i>	2
<i>Corylus avellana</i>	4
<i>Crataegus monogyna</i>	3
<i>Fragaria vesca</i>	5
<i>Fraxinus excelsior</i>	6
<i>Geranium robertianum</i>	4
<i>Hedera helix</i>	6
<i>Ilex aquifolium</i>	5
<i>Phyllitis scolopendrium</i>	3
<i>Prunus spinosa</i>	4
<i>Rubus fruticosus</i> agg.	5

<b>Relevé code</b>	<b>RC-LQ R5</b>
Date	02/11/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	*8240
Species	Domin
<i>Briza media</i>	5
<i>Carex flacca</i>	6
<i>Centaurea nigra</i>	2
<i>Crepis capillaris</i>	2
<i>Fragaria vesca</i>	4
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	4
<i>Hypericum pulchrum</i>	2
<i>Lotus corniculatus</i>	5
<i>Plantago lanceolata</i>	3
<i>Rubus fruticosus</i> agg.	3
<i>Sesleria caerulea</i>	6
<i>Thymus polytrichus</i>	3

<b>Relevé code</b>	<b>R336-NFMR R1</b>
Date	07/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
<b>Species</b>	<b>Domin</b>
<i>Agrostis stolonifera</i>	4
<i>Calluna vulgaris</i>	4
<i>Campylopus introflexus</i>	1
<i>Daboecia cantabrica</i>	3
<i>Deschampsia flexuosa</i>	1
<i>Erica cinerea</i>	6
<i>Holcus lanatus</i>	2
<i>Hylocomium splendens</i>	3
<i>Hypnum jutlandicum</i>	5
<i>Pteridium aquilinum</i>	4
<i>Ulex gallii</i>	8

<b>Relevé code</b>	<b>R336-NFMR R2</b>
Date	03/09/2015
Recorder	EOS
Species recording scale	Domin
Habitat - Fossitt	HH1
Habitat - Annex I	4030
<b>Species</b>	<b>Domin</b>
<i>Calluna vulgaris</i>	5
<i>Cladonia</i> sp.	1
<i>Daboecia cantabrica</i>	4
<i>Danthonia decumbens</i>	1
<i>Deschampsia flexuosa</i>	1
<i>Erica cinerea</i>	5
<i>Hypnum jutlandicum</i>	1
<i>Molinia caerulea</i>	4
<i>Pteridium aquilinum</i>	1
<i>Rubus fruticosus</i> agg.	3
<i>Scleropodium purum</i>	1
<i>Ulex gallii</i>	7

<b>Relevé code</b>	<b>UD-NUIG R1</b>
Date	29/07/2016
Recorder	JF
Species recording scale	Domin
Habitat - Fossitt	GS1
Habitat - Annex I	-
Species	Domin
<i>Anthoxanthum odoratum</i>	6
<i>Briza media</i>	4
<i>Calliergonella cuspidata</i>	5
<i>Carex flacca</i>	2
<i>Cerastium fontanum</i>	1
<i>Euphrasia sp.</i>	1
<i>Festuca ovina</i>	4
<i>Holcus lanatus</i>	5
<i>Homalothecium lutescens</i>	3
<i>Lathyrus pratensis</i>	3
<i>Linum catharticum</i>	4
<i>Lolium perenne</i>	3
<i>Plantago lanceolata</i>	1
<i>Prunella vulgaris</i>	3
<i>Ranunculus acris</i>	1
<i>Ranunculus repens</i>	3
<i>Trifolium pratense</i>	4

<b>Relevé code</b>	<b>UD-NUIG R3</b>
Date	13/10/2017
Recorder	KMOC
Species recording scale	Domin
Habitat - Fossitt	WN2
Habitat - Annex I	-
Species	Domin
<i>Acer pseudoplatanus</i>	7
<i>Crataegus monogyna</i>	6
<i>Fraxinus excelsior</i>	8
<i>Hedera helix</i>	9
<i>Ilex aquifolium</i>	6
<i>Phyllitis scolopendrium</i>	6
<i>Rubus fruticosus agg.</i>	8
<i>Thuidium sp.</i>	5

<b>Relevé code</b>	<b>UD-NUIG R5</b>
Date	02/11/2017
Recorder	CC
Species recording scale	Domin
Habitat - Fossitt	WN6
Habitat - Annex I	*91E0-
Species	Domin
<i>Alnus glutinosa</i>	5
<i>Circaea lutetiana</i>	3
<i>Crataegus monogyna</i>	4
<i>Filipendula ulmaria</i>	4
<i>Geranium robertianum</i>	3
<i>Hedera helix</i>	6
<i>Heracleum sphondylium</i>	2
<i>Holcus lanatus</i>	1
<i>Phyllitis scolopendrium</i>	2
<i>Ranunculus repens</i>	7
<i>Salix cinerea</i>	6
<i>Scrophularia auriculata</i>	6
<i>Taraxacum officinale</i> agg.	2
<i>Urtica dioica</i>	5



<b>Relevé code</b>	<b>BEC 7</b>
Date	25/07/2013
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	6210
Species	Percentage
<i>Barbula unguiculata</i>	0.1
<i>Briza media</i>	1
<i>Campyliadelphus chrysophyllus</i>	0.1
<i>Carex flacca</i>	0.5
<i>Carlina vulgaris</i>	2
<i>Cladonia pocillum</i>	0.3
<i>Cladonia rangiformis</i>	3
<i>Ctenidium molluscum</i>	7
<i>Ditrichum gracile</i>	0.1
<i>Euphrasia salisburgensis</i>	0.1
<i>Festuca ovina</i>	0.3
<i>Fissidens dubius</i>	0.5
<i>Fragaria vesca</i>	0.3
<i>Galium verum</i>	0.5
<i>Hedera helix</i>	0.7
<i>Hypericum pulchrum</i>	1
<i>Hypochaeris radicata</i>	0.5
<i>Lotus corniculatus</i>	5
<i>Neckera crispa</i>	0.3
<i>Pilosella officinarum</i>	1
<i>Plantago lanceolata</i>	0.1
<i>Potentilla erecta</i>	0.1
<i>Rosa spinosissima</i>	1
<i>Rubus fruticosus</i> agg.	2
<i>Scapania aspera</i>	0.5
<i>Scleropodium purum</i>	1
<i>Sesleria caerulea</i>	45
<i>Teucrium scorodonia</i>	2
<i>Thymus polytrichus</i>	5
<i>Tortella tortuosa</i>	5

<b>Relevé code</b>	<b>BEC 8</b>
Date	25/07/2013
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	WS1
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	1
<i>Carlina vulgaris</i>	0.3
<i>Corylus avellana</i>	30
<i>Ctenidium molluscum</i>	20
<i>Encalypta streptocarpa</i>	0.1
<i>Euphrasia salisburgensis</i>	0.1
<i>Fissidens dubius</i>	1
<i>Fragaria vesca</i>	0.1
<i>Fraxinus excelsior</i>	0.1
<i>Geranium robertianum</i>	0.3
<i>Hedera helix</i>	25
<i>Ilex aquifolium</i>	0.1
<i>Lonicera periclymenum</i>	3
<i>Mycelis muralis</i>	0.1
<i>Neckera complanata</i>	0.1
<i>Neckera crispa</i>	7
<i>Potentilla sterilis</i>	0.1
<i>Rubus fruticosus</i> agg.	10
<i>Senecio jacobaea</i>	0.3
<i>Sesleria caerulea</i>	10
<i>Sorbus aria</i>	5
<i>Taraxacum officinale</i> agg.	0.1
<i>Teucrium scorodonia</i>	0.3
<i>Tortella tortuosa</i>	5
<i>Trichostomum crispulum</i>	1

<b>Relevé code</b>	<b>BEC 9</b>
Date	25/07/2013
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.1
<i>Asplenium trichomanes</i>	0.1
<i>Brachythecium species</i>	0.1
<i>Calliergonella cuspidata</i>	0.1
<i>Corylus avellana</i>	50
<i>Ctenidium molluscum</i>	15
<i>Euonymus europaeus</i>	5
<i>Eurhynchium striatum</i>	35
<i>Fissidens dubius</i>	0.1
<i>Fragaria vesca</i>	0.5
<i>Fraxinus excelsior</i>	5
<i>Geranium robertianum</i>	0.1
<i>Hedera helix</i>	70
<i>Ilex aquifolium</i>	3
<i>Kindbergia praelonga</i>	0.1
<i>Lonicera periclymenum</i>	0.5
<i>Metzgeria furcata</i>	0.1
<i>Neckera crispa</i>	15
<i>Phyllitis scolopendrium</i>	1
<i>Pteridium aquilinum</i>	5
<i>Rosa canina</i>	0.5
<i>Rosa spinosissima</i>	0.3
<i>Rubus fruticosus agg.</i>	7
<i>Scapania aspera</i>	0.1
<i>Scleropodium purum</i>	0.1
<i>Sesleria caerulea</i>	7
<i>Sorbus aria</i>	10
<i>Sorbus aucuparia</i>	10
<i>Thamnobryum alopecurum</i>	5
<i>Thuidium tamariscinum</i>	0.1

<b>Relevé code</b>	<b>BEC 15</b>
Date	14/08/2013
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Species	Percentage
<i>Antennaria dioica</i>	0.3
<i>Anthyllis vulneraria</i>	1
<i>Asperula cynanchica</i>	0.3
<i>Breutelia chrysocoma</i>	0.5
<i>Briza media</i>	6
<i>Campanula rotundifolia</i>	0.3
<i>Carex caryophylla</i>	0.3
<i>Carex flacca</i>	15
<i>Carlina vulgaris</i>	2
<i>Centaurea nigra</i>	1
<i>Ctenidium molluscum</i>	2
<i>Daucus carota</i>	1
<i>Euphrasia officinalis</i> agg.	1
<i>Festuca ovina</i>	6
<i>Fissidens dubius</i>	0.1
<i>Frullania tamarisci</i>	0.5
<i>Galium verum</i>	0.5
<i>Gymnadenia conopsea</i>	0.1
<i>Hypericum pulchrum</i>	0.5
<i>Koeleria macrantha</i>	0.1
<i>Leontodon hispidus</i>	15
<i>Leucanthemum vulgare</i>	3
<i>Linum catharticum</i>	0.5
<i>Lotus corniculatus</i>	30
<i>Pilosella officinarum</i>	0.5
<i>Plantago lanceolata</i>	0.3
<i>Plantago maritima</i>	35
<i>Polygala vulgaris</i>	0.5
<i>Potentilla erecta</i>	5
<i>Prunella vulgaris</i>	1
<i>Rhinanthus minor</i>	0.1
<i>Rosa spinosissima</i>	1
<i>Scapania aspera</i>	0.3
<i>Scleropodium purum</i>	0.1
<i>Scorpidium scorpioides</i>	0.1
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	8
<i>Solidago virgaurea</i>	5
<i>Succisa pratensis</i>	6
<i>Thymus polytrichus</i>	4
<i>Tortella tortuosa</i>	2
<i>Trifolium pratense</i>	1

<b>Relevé code</b>	<b>BEC 17</b>
Date	14/08/2013
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Achillea millefolium	0.5
Species	Percentage
<i>Agrostis capillaris</i>	8
<i>Anthoxanthum odoratum</i>	4
<i>Anthyllis vulneraria</i>	0.1
<i>Asperula cynanchica</i>	0.5
<i>Briza media</i>	2
<i>Calliergonella cuspidata</i>	0.3
<i>Campanula rotundifolia</i>	0.5
<i>Carex caryophylla</i>	0.3
<i>Carex flacca</i>	45
<i>Carex panicea</i>	0.5
<i>Centaurea nigra</i>	20
<i>Ctenidium molluscum</i>	2
<i>Cynosurus cristatus</i>	2
<i>Dactylis glomerata</i>	0.1
<i>Danthonia decumbens</i>	0.1
<i>Daucus carota</i>	2
<i>Euphrasia officinalis</i> agg.	0.5
<i>Festuca ovina</i>	5
<i>Fissidens adianthoides</i>	0.1
<i>Galium verum</i>	0.5
<i>Gymnadenia conopsea</i>	0.1
<i>Helictotrichon pubescens</i>	0.1
<i>Leontodon hispidus</i>	30
<i>Leucanthemum vulgare</i>	0.5
<i>Linum catharticum</i>	0.3
<i>Lotus corniculatus</i>	25
<i>Phleum pratense</i>	0.1
<i>Pilosella officinarum</i>	0.1
<i>Plagiomnium undulatum</i>	0.1
<i>Plantago lanceolata</i>	3
<i>Plantago maritima</i>	3
<i>Potentilla erecta</i>	10
<i>Prunella vulgaris</i>	1
<i>Ranunculus bulbosus</i>	0.1
<i>Rhinanthus minor</i>	0.3
<i>Rosa spinosissima</i>	1
<i>Scleropodium purum</i>	0.1
<i>Sesleria caerulea</i>	5
<i>Succisa pratensis</i>	2
<i>Thymus polytrichus</i>	1
<i>Trifolium pratense</i>	0.5
<i>Trifolium repens</i>	3
<i>Viola riviniana</i>	0.3

<b>Relevé code</b>	<b>BEC 18</b>
Date	14/08/2013
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WS1
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.3
<i>Briza media</i>	0.3
<i>Campylium stellatum</i> v. <i>protensum</i>	0.1
<i>Corylus avellana</i>	35
<i>Crataegus monogyna</i>	40
<i>Ctenidium molluscum</i>	40
<i>Eurhynchium striatum</i>	5
<i>Festuca ovina</i>	6
<i>Fragaria vesca</i>	1
<i>Geranium robertianum</i>	1
<i>Hedera helix</i>	55
<i>Hypericum pulchrum</i>	0.5
<i>Ilex aquifolium</i>	1
<i>Kindbergia praelonga</i>	0.5
<i>Lathyrus linifolius</i>	0.5
<i>Lonicera periclymenum</i>	10
<i>Lotus corniculatus</i>	0.5
<i>Melica uniflora</i>	0.3
<i>Neckera complanata</i>	0.3
<i>Neckera crispa</i>	1
<i>Plantago lanceolata</i>	0.1
<i>Potentilla erecta</i>	3
<i>Potentilla sterilis</i>	0.3
<i>Prunus spinosa</i>	10
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	40
<i>Scleropodium purum</i>	0.3
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	20
<i>Solidago virgaurea</i>	0.3
<i>Succisa pratensis</i>	0.5
<i>Teucrium scorodonia</i>	0.5
<i>Thuidium tamariscinum</i>	5
<i>Tortella tortuosa</i>	0.3
<i>Veronica chamaedrys</i>	0.1

<b>Relevé code</b>	<b>BEC 35</b>
Date	29/08/2013
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	WS1
Habitat - Annex I	*8240
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.3
<i>Asplenium ruta-muraria</i>	0.1
<i>Breutelia chrysocoma</i>	0.1
<i>Bryum capillare</i>	0.1
<i>Carex flacca</i>	3
<i>Ceterach officinarum</i>	0.1
<i>Corylus avellana</i>	70
<i>Crataegus monogyna</i>	3
<i>Ctenidium molluscum</i>	25
<i>Dicranum scoparium</i>	0.3
<i>Festuca ovina</i>	3
<i>Fissidens dubius</i>	0.1
<i>Fragaria vesca</i>	10
<i>Fraxinus excelsior</i>	1
<i>Hedera helix</i>	50
<i>Hypericum pulchrum</i>	0.1
<i>Ilex aquifolium</i>	0.1
<i>Isoetes macrospora</i>	1
<i>Neckera complanata</i>	0.3
<i>Neckera crispa</i>	10
<i>Potentilla erecta</i>	0.1
<i>Prunus spinosa</i>	0.3
<i>Pteridium aquilinum</i>	15
<i>Rosa spinosissima</i>	5
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	25
<i>Solidago virgaurea</i>	0.1
<i>Thuidium tamariscinum</i>	0.3
<i>Thymus polytrichus</i>	0.1
<i>Tortella tortuosa</i>	0.3

<b>Relevé code</b>	<b>BEC 42</b>
Date	18/09/2013
Recorder	FD/JM
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.3
<i>Asplenium trichomanes</i>	0.1
<i>Brachypodium sylvaticum</i>	0.3
<i>Corylus avellana</i>	90
<i>Crataegus monogyna</i>	0.5
<i>Ctenidium molluscum</i>	0.3
<i>Dryopteris affinis</i>	0.3
<i>Eurhynchium striatum</i>	50
<i>Fissidens dubius</i>	0.3
<i>Fragaria vesca</i>	0.3
<i>Fraxinus excelsior</i>	25
<i>Hedera helix</i>	40
<i>Hypnum cupressiforme s.s.</i>	0.1
<i>Ilex aquifolium</i>	60
<i>Kindbergia praelonga</i>	1
<i>Lathyrus linifolius</i>	0.3
<i>Lonicera periclymenum</i>	0.5
<i>Metzgeria furcata</i>	3
<i>Neckera crispa</i>	1
<i>Phyllitis scolopendrium</i>	0.5
<i>Plagiomnium undulatum</i>	0.3
<i>Polypodium vulgare</i>	0.1
<i>Prunus spinosa</i>	0.5
<i>Pteridium aquilinum</i>	1
<i>Rhytidiadelphus triquetrus</i>	0.5
<i>Rubus fruticosus agg.</i>	5
<i>Thamnobryum alopecurum</i>	1
<i>Thuidium tamariscinum</i>	1
<i>Tortella tortuosa</i>	0.3
<i>Viburnum opulus</i>	0.3
<i>Viola sp.</i>	0.7



<b>Relevé code</b>	<b>BEC 102</b>
Date	30/06/2014
Recorder	SB/FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	*6210
Species	Percentage
<i>Aira caryophylla</i>	0.1
<i>Anthoxanthum odoratum</i>	3
<i>Anthyllis vulneraria</i>	0.5
<i>Briza media</i>	5
<i>Calliergonella cuspidata</i>	1
<i>Campanula rotundifolia</i>	0.3
<i>Carex caryophylla</i>	20
<i>Carex flacca</i>	20
<i>Centaurea nigra</i>	5
<i>Ctenidium molluscum</i>	5
<i>Cynosurus cristatus</i>	1
<i>Dactylis glomerata</i>	0.1
<i>Daucus carota</i>	0.5
<i>Euphrasia officinalis</i> agg.	0.5
<i>Festuca rubra</i>	20
<i>Galium verum</i>	0.7
<i>Geranium sanguineum</i>	5
<i>Gymnadenia conopsea</i>	0.7
<i>Koeleria macrantha</i>	0.5
<i>Leontodon hispidus</i>	10
<i>Leucanthemum vulgare</i>	0.5
<i>Linum catharticum</i>	1
<i>Lotus corniculatus</i>	5
<i>Pilosella officinarum</i>	0.5
<i>Plantago lanceolata</i>	1
<i>Polygala vulgaris</i>	0.5
<i>Potentilla erecta</i>	10
<i>Rhinanthus minor</i>	3
<i>Rosa spinosissima</i>	20
<i>Scleropodium purum</i>	2
<i>Sesleria caerulea</i>	1
<i>Solidago virgaurea</i>	5
<i>Succisa pratensis</i>	15
<i>Trifolium repens</i>	0.5

<b>Relevé code</b>	<b>BEC 115</b>
Date	18/07/2014
Recorder	JM
Species recording scale	Percentage
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.3
<i>Briza media</i>	0.1
<i>Carex flacca</i>	1
<i>Carlina vulgaris</i>	0.1
<i>Crataegus monogyna</i>	7
<i>Ctenidium molluscum</i>	0.1
<i>Fissidens dubius</i>	0.3
<i>Hedera helix</i>	20
<i>Hypochaeris radicata</i>	0.1
<i>Ilex aquifolium</i>	0.1
<i>Lotus corniculatus</i>	5
<i>Mycelis muralis</i>	0.3
<i>Pilosella officinarum</i>	0.7
<i>Polygala vulgaris</i>	0.1
<i>Prunus spinosa</i>	5
<i>Rosa spinosissima</i>	7
<i>Rubus fruticosus</i> agg.	5
<i>Senecio jacobaea</i>	0.3
<i>Sesleria caerulea</i>	15
<i>Sonchus asper</i>	0.1
<i>Succisa pratensis</i>	3
<i>Taraxacum officinale</i> agg.	0.1
<i>Teucrium scorodonia</i>	0.3
<i>Thymus polytrichus</i>	0.5
<i>Tortella densa</i>	0.1

<b>Relevé code</b>	<b>BEC 118</b>
Date	18/07/2014
Recorder	JM
Species recording scale	Percentage
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.3
<i>Carex flacca</i>	1
<i>Carlina vulgaris</i>	0.1
<i>Ceterach officinarum</i>	0.1
<i>Crataegus monogyna</i>	0.7
<i>Ctenidium molluscum</i>	0.1
<i>Fissidens dubius</i>	0.3
<i>Geranium robertianum</i>	0.7
<i>Hedera helix</i>	1
<i>Hypochaeris radicata</i>	0.1
<i>Lotus corniculatus</i>	3
<i>Mycelis muralis</i>	0.1
<i>Pilosella officinarum</i>	0.1
<i>Polygala vulgaris</i>	0.1
<i>Prunus spinosa</i>	5
<i>Rubus fruticosus</i> agg.	5
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	7
<i>Sonchus asper</i>	0.1
<i>Teucrium scorodonia</i>	1
<i>Thymus polytrichus</i>	1
<i>Tortella nitida</i>	0.1
<i>Tortella tortuosa</i>	0.1

<b>Relevé code</b>	<b>BEC 120</b>
Date	18/07/2014
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Brachypodium sylvaticum</i>	0.7
<i>Calliergonella cuspidata</i>	0.1
<i>Carex flacca</i>	0.1
<i>Corylus avellana</i>	30
<i>Crataegus monogyna</i>	20
<i>Ctenidium molluscum</i>	3
<i>Euonymus europaeus</i>	3
<i>Eurhynchium striatum</i>	35
<i>Festuca rubra</i>	3
<i>Fragaria vesca</i>	0.1
<i>Fraxinus excelsior</i>	7
<i>Hedera helix</i>	35
<i>Hypericum pulchrum</i>	0.3

<b>Relevé code</b>	<b>BEC 120</b>
<i>Ilex aquifolium</i>	10
<i>Lonicera periclymenum</i>	0.1
<i>Neckera crispa</i>	0.5
<i>Phyllitis scolopendrium</i>	0.1
<i>Potentilla erecta</i>	0.5
<i>Potentilla sterilis</i>	0.1
<i>Prunus spinosa</i>	3
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	15
<i>Scleropodium purum</i>	0.1
<i>Sesleria caerulea</i>	25
<i>Solidago virgaurea</i>	0.1
<i>Succisa pratensis</i>	0.3
<i>Thamnobryum alopecurum</i>	0.7
<i>Thuidium tamariscinum</i>	3
<i>Veronica chamaedrys</i>	0.1
<i>Vicia sepium</i>	0.1
<i>Viola</i> sp.	0.1

<b>Relevé code</b>	<b>BEC 121</b>
Date	18/07/2014
Recorder	JM/FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Cololejeunea minutissima</i>	0.1
<i>Corylus avellana</i>	85
<i>Crataegus monogyna</i>	30
<i>Eurhynchium striatum</i>	40
<i>Fragaria vesca</i>	0.3
<i>Fraxinus excelsior</i>	0.3
<i>Frullania dilatata</i>	0.1
<i>Geum urbanum</i>	0.1
<i>Hedera helix</i>	40
<i>Ilex aquifolium</i>	7
<i>Lonicera periclymenum</i>	3
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	5
<i>Potentilla sterilis</i>	0.3
<i>Prunus spinosa</i>	1
<i>Radula complanata</i>	0.1
<i>Rubus fruticosus</i> agg.	1
<i>Thamnobryum alopecurum</i>	3
<i>Thuidium tamariscinum</i>	3
<i>Ulotia crispa</i>	0.1
<i>Viola</i> sp.	0.3
<i>Zygodon conoideus</i>	0.1

<b>Relevé code</b>	<b>BEC 139</b>
Date	26/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Corylus avellana</i>	80
<i>Crataegus monogyna</i>	7
<i>Eurhynchium striatum</i>	5
<i>Fraxinus excelsior</i>	25
<i>Hedera helix</i>	60
<i>Ilex aquifolium</i>	7
<i>Isothecium alopecuroides</i>	0.1
<i>Lonicera periclymenum</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	1
<i>Plagiomnium undulatum</i>	0.3
<i>Potentilla sterilis</i>	0.1
<i>Rhytidiadelphus triquetrus</i>	5
<i>Rubus fruticosus</i> agg.	0.1
<i>Sorbus aria</i>	20
<i>Thamnobryum alopecurum</i>	1
<i>Thuidium tamariscinum</i>	5
<i>Viola</i> sp.	0.3

<b>Relevé code</b>	<b>BEC 144</b>
Date	28/07/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Brachypodium sylvaticum</i>	0.1
<i>Calluna vulgaris</i>	0.1
<i>Carex flacca</i>	1
<i>Corylus avellana</i>	65
<i>Crataegus monogyna</i>	35
<i>Ctenidium molluscum</i>	3
<i>Eurhynchium striatum</i>	40
<i>Fissidens dubius</i>	0.3
<i>Fragaria vesca</i>	0.3
<i>Geum urbanum</i>	0.1
<i>Hedera helix</i>	30
<i>Hypericum pulchrum</i>	0.1
<i>Ilex aquifolium</i>	15
<i>Isothecium alopecuroides</i>	0.1
<i>Kindbergia praelonga</i>	0.3
<i>Lonicera periclymenum</i>	5
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	3

<b>Relevé code</b>	<b>BEC 144</b>
<i>Neckera crispa</i>	0.5
<i>Phyllitis scolopendrium</i>	0.1
<i>Plagiomnium undulatum</i>	0.1
<i>Potentilla sterilis</i>	3
<i>Prunus spinosa</i>	15
<i>Rosa spinosissima</i>	1
<i>Rubus fruticosus</i> agg.	10
<i>Scleropodium purum</i>	1
<i>Sesleria caerulea</i>	5
<i>Thamnobryum alopecurum</i>	10
<i>Thuidium tamariscinum</i>	15
<i>Tortella tortuosa</i>	0.1
<i>Viola</i> sp.	3
<i>Zygodon conoideus</i>	0.1

<b>Relevé code</b>	<b>BEC 145</b>
Date	29/07/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Brachypodium sylvaticum</i>	3
<i>Carex flacca</i>	0.1
<i>Corylus avellana</i>	65
<i>Crataegus monogyna</i>	15
<i>Ctenidium molluscum</i>	0.1
<i>Epipactis helleborine</i>	0.1
<i>Eurhynchium striatum</i>	0.5
<i>Fragaria vesca</i>	0.3
<i>Frullania dilatata</i>	0.1
<i>Geranium robertianum</i>	0.1
<i>Hedera helix</i>	70
<i>Ilex aquifolium</i>	10
<i>Isoetecium alopecuroides</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	0.3
<i>Neckera crispa</i>	0.1
<i>Plagiochila porelloides</i>	0.1
<i>Potentilla sterilis</i>	3
<i>Prunus spinosa</i>	20
<i>Rubus fruticosus</i> agg.	5
<i>Sesleria caerulea</i>	0.5
<i>Thamnobryum alopecurum</i>	0.3
<i>Thuidium tamariscinum</i>	45
<i>Ulota bruchii</i>	0.1
<i>Veronica chamaedrys</i>	0.1
<i>Viola</i> sp.	1

<b>Relevé code</b>	<b>BEC 148</b>
Date	29/07/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WS1
Habitat - Annex I	*8240
Species	Percentage
<i>Brachypodium sylvaticum</i>	0.7
<i>Briza media</i>	0.3
<i>Calluna vulgaris</i>	15
<i>Carex flacca</i>	1
<i>Corylus avellana</i>	30
<i>Crataegus monogyna</i>	7
<i>Ctenidium molluscum</i>	7
<i>Dactylis glomerata</i>	0.3
<i>Dicranum scoparium</i>	0.1
<i>Euonymus europaeus</i>	5
<i>Eurhynchium striatum</i>	1
<i>Festuca ovina</i>	3
<i>Fragaria vesca</i>	5
<i>Fraxinus excelsior</i>	1
<i>Galium verum</i>	0.1
<i>Hedera helix</i>	35
<i>Homalothecium lutescens</i>	0.1
<i>Hypericum pulchrum</i>	0.1
<i>Hypnum lacunosum</i>	0.1
<i>Ilex aquifolium</i>	0.5
<i>Koeleria macrantha</i>	0.3
<i>Lathyrus linifolius</i>	0.3
<i>Lotus corniculatus</i>	3
<i>Neckera crispa</i>	0.5
<i>Plantago lanceolata</i>	0.1
<i>Potentilla erecta</i>	3
<i>Potentilla sterilis</i>	0.5
<i>Prunus spinosa</i>	0.5
<i>Rosa spinosissima</i>	3
<i>Rubus fruticosus</i> agg.	20
<i>Scleropodium purum</i>	0.5
<i>Sesleria caerulea</i>	30
<i>Solidago virgaurea</i>	0.1
<i>Succisa pratensis</i>	0.7
<i>Teucrium scorodonia</i>	0.3
<i>Thuidium tamariscinum</i>	10
<i>Tortella tortuosa</i>	0.1
<i>Trichostomum brachydontium</i>	0.1
<i>Viola</i> sp.	0.5

<b>Relevé code</b>	<b>BEC 174</b>
Date	19/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WD1
Habitat - Annex I	None
Species	Percentage
<i>Arum maculatum</i>	0.1
<i>Fagus sylvatica</i>	80
<i>Fraxinus excelsior</i>	10
<i>Hedera helix</i>	0.1
<i>Ilex aquifolium</i>	50
<i>Isoetes macrospora</i>	0.3
<i>Kindbergia praelonga</i>	0.1
<i>Lejeunea cavifolia</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	0.1
<i>Radula complanata</i>	0.1
<i>Rhynchostegiella tenella</i>	0.1
<i>Thamnobryum alopecurum</i>	15
<i>Tortella tortuosa</i>	0.1

<b>Relevé code</b>	<b>BEC 175</b>
Date	19/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	None
Species	Percentage
<i>Agrostis stolonifera</i>	1
<i>Calliergonella cuspidata</i>	0.5
<i>Carex disticha</i>	0.3
<i>Carex hirta</i>	3
<i>Centaurea nigra</i>	1
<i>Cerastium fontanum</i>	0.1
<i>Cynosurus cristatus</i>	10
<i>Dactylis glomerata</i>	1
<i>Elytrigia repens</i>	0.1
<i>Festuca rubra</i>	15
<i>Holcus lanatus</i>	1
<i>Lolium perenne</i>	50
<i>Plantago lanceolata</i>	5
<i>Ranunculus acris</i>	5
<i>Ranunculus repens</i>	40
<i>Rumex acetosa</i>	1
<i>Rumex crispus</i>	3
<i>Taraxacum officinale</i> agg.	3
<i>Trifolium pratense</i>	20
<i>Trifolium repens</i>	30



<b>Relevé code</b>	<b>BEC 240</b>
Date	29/08/2014
Recorder	ML
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Brachypodium sylvaticum</i>	0.5
<i>Brachythecium rutabulum</i>	0.1
<i>Carex flacca</i>	1
<i>Corylus avellana</i>	60
<i>Crataegus monogyna</i>	15
<i>Ctenidium molluscum</i>	1
<i>Dactylis glomerata</i>	0.1
<i>Eurhynchium striatum</i>	30
<i>Filipendula ulmaria</i>	0.1
<i>Fissidens taxifolius</i>	0.3
<i>Fraxinus excelsior</i>	40
<i>Geranium robertianum</i>	0.3
<i>Geum urbanum</i>	0.3
<i>Hedera helix</i>	50
<i>Ilex aquifolium</i>	40
<i>Kindbergia praelonga</i>	0.3
<i>Lonicera periclymenum</i>	0.1
<i>Metzgeria furcata</i>	0.3
<i>Neckera complanata</i>	3
<i>Neckera crispa</i>	3
<i>Phyllitis scolopendrium</i>	0.1
<i>Plagiomnium undulatum</i>	0.1
<i>Potentilla sterilis</i>	0.5
<i>Prunus spinosa</i>	10
<i>Pteridium aquilinum</i>	0.1
<i>Rubus fruticosus</i> agg.	15
<i>Scapania aspera</i>	0.5
<i>Solidago virgaurea</i>	0.3
<i>Thuidium tamariscinum</i>	10
<i>Ulotia crispa</i>	0.1
<i>Veronica chamaedrys</i>	1
<i>Viola</i> sp.	1

<b>Relevé code</b>	<b>BEC 246</b>
Date	29/08/2014
Recorder	ML
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	*8240
Species	Percentage
<i>Arum maculatum</i>	0.1
<i>Cololejeunea minutissima</i>	0.1
<i>Crataegus monogyna</i>	75
<i>Eurhynchium striatum</i>	30
<i>Fraxinus excelsior</i>	75
<i>Frullania dilatata</i>	0.1
<i>Hedera helix</i>	95
<i>Hypnum resupinatum</i>	0.1
<i>Ilex aquifolium</i>	20
<i>Kindbergia praelonga</i>	1
<i>Lonicera periclymenum</i>	0.1
<i>Neckera complanata</i>	0.7
<i>Phyllitis scolopendrium</i>	5
<i>Plagiomnium undulatum</i>	0.1
<i>Primula vulgaris</i>	0.5
<i>Prunus spinosa</i>	15
<i>Pteridium aquilinum</i>	5
<i>Radula complanata</i>	0.1
<i>Rubus fruticosus</i> agg.	20
<i>Thamnobryum alopecurum</i>	10
<i>Thuidium tamariscinum</i>	5
<i>Zygodon viridissimus</i>	0.1

<b>Relevé code</b>	<b>BEC 242</b>
Date	26/08/2014
Recorder	ML
Species recording scale	Percentage
Habitat - Fossitt	WS1
Habitat - Annex I	*8240
Species	Percentage
<i>Agrostis stolonifera</i>	0.3
<i>Brachypodium sylvaticum</i>	3
<i>Bryum capillare</i>	0.1
<i>Carex flacca</i>	30
<i>Corylus avellana</i>	55
<i>Crataegus monogyna</i>	10
<i>Ctenidium molluscum</i>	15
<i>Eurhynchium striatum</i>	20
<i>Fissidens taxifolius</i>	0.1
<i>Fragaria vesca</i>	20
<i>Fraxinus excelsior</i>	5
<i>Hedera helix</i>	20
<i>Hypericum pulchrum</i>	0.3
<i>Ilex aquifolium</i>	15
<i>Lonicera periclymenum</i>	5
<i>Lotus corniculatus</i>	0.5
<i>Neckera complanata</i>	1
<i>Neckera crispa</i>	15
<i>Plantago lanceolata</i>	0.1
<i>Populus tremula</i>	10
<i>Potentilla erecta</i>	1
<i>Prunus spinosa</i>	5
<i>Rosa spinosissima</i>	1
<i>Rubus fruticosus</i> agg.	20
<i>Scleropodium purum</i>	0.5
<i>Sesleria caerulea</i>	3
<i>Teucrium scorodonia</i>	0.1
<i>Thuidium tamariscinum</i>	10
<i>Tortella tortuosa</i>	1
<i>Viola</i> sp.	0.3

<b>Relevé code</b>	<b>BEC 258</b>
Date	29/08/2014
Recorder	ML
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	None
Species	Percentage
<i>Agrostis stolonifera</i>	20
<i>Anthoxanthum odoratum</i>	3
<i>Carex flacca</i>	20
<i>Centaurea nigra</i>	10
<i>Cirsium vulgare</i>	5
<i>Epilobium montanum</i>	0.3
<i>Eurhynchium hians</i>	5
<i>Eurhynchium striatum</i>	3
<i>Festuca rubra</i>	1
<i>Fissidens taxifolius</i>	0.5
<i>Fraxinus excelsior</i>	0.1
<i>Lathyrus pratensis</i>	7
<i>Leucanthemum vulgare</i>	1
<i>Lolium perenne</i>	10
<i>Lotus corniculatus</i>	10
<i>Plagiomnium undulatum</i>	0.5
<i>Plantago lanceolata</i>	0.5
<i>Plantago major</i>	3
<i>Potentilla erecta</i>	1
<i>Potentilla sterilis</i>	1
<i>Prunella vulgaris</i>	0.3
<i>Pteridium aquilinum</i>	5
<i>Ranunculus repens</i>	20
<i>Rubus fruticosus</i> agg.	10
<i>Rumex conglomeratus</i>	0.1
<i>Scleropodium purum</i>	3
<i>Senecio jacobaea</i>	0.3
<i>Taraxacum officinale</i> agg.	0.5
<i>Thuidium tamariscinum</i>	1
<i>Trifolium pratense</i>	1
<i>Trifolium repens</i>	5
<i>Veronica chamaedrys</i>	0.3
<i>Viola</i> sp.	0.1

<b>Relevé code</b>	<b>BEC 333</b>
Date	25/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Acer pseudoplatanus</i>	20
<i>Corylus avellana</i>	75
<i>Crataegus monogyna</i>	0.5
<i>Eurhynchium striatum</i>	1
<i>Hedera helix</i>	35
<i>Homalothecium sericeum</i>	0.1
<i>Ilex aquifolium</i>	30
<i>Lonicera periclymenum</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	1
<i>Radula complanata</i>	0.1
<i>Rubus fruticosus</i> agg.	0.5
<i>Thamnobryum alopecurum</i>	70
<i>Thuidium tamariscinum</i>	0.1

<b>Relevé code</b>	<b>BEC 335</b>
Date	26/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Acer pseudoplatanus</i>	0.1
<i>Corylus avellana</i>	90
<i>Crataegus monogyna</i>	5
<i>Ctenidium molluscum</i>	0.1
<i>Euonymus europaeus</i>	0.1
<i>Eurhynchium striatum</i>	3
<i>Fragaria vesca</i>	0.1
<i>Fraxinus excelsior</i>	5
<i>Hedera helix</i>	70
<i>Ilex aquifolium</i>	5
<i>Lonicera periclymenum</i>	5
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	1
<i>Neckera crispa</i>	0.3
<i>Plagiomnium undulatum</i>	0.1
<i>Polypodium sp.</i>	0.1
<i>Quercus robur</i>	0.1
<i>Rhytidiadelphus triquetrus</i>	3
<i>Rubus fruticosus agg.</i>	3
<i>Rubus saxatilis</i>	0.3
<i>Sorbus aucuparia</i>	0.5
<i>Thamnobryum alopecurum</i>	65
<i>Thuidium tamariscinum</i>	1
<i>Tortella tortuosa</i>	0.1
<i>Viola sp.</i>	0.1

<b>Relevé code</b>	<b>BEC 336</b>
Date	26/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Arum maculatum</i>	0.1
<i>Brachypodium sylvaticum</i>	0.3
<i>Carex sylvatica</i>	1
<i>Corylus avellana</i>	65
<i>Crataegus monogyna</i>	25
<i>Eurhynchium striatum</i>	7
<i>Fraxinus excelsior</i>	15
<i>Hedera helix</i>	45
<i>Ilex aquifolium</i>	20
<i>Isoetes macrospora</i>	0.1
<i>Isoetes macrospora</i>	0.1
<i>Kindbergia praelonga</i>	0.1
<i>Lonicera periclymenum</i>	0.3
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	1
<i>Plagiomnium undulatum</i>	0.5
<i>Potentilla sterilis</i>	0.3
<i>Prunus spinosa</i>	0.1
<i>Rhytidadelphus triquetrus</i>	7
<i>Rubus fruticosus</i> agg.	1
<i>Thamnobryum alopecurum</i>	25
<i>Thuidium tamariscinum</i>	7
<i>Viola</i> sp.	1

<b>Relevé code</b>	<b>BEC 340</b>
Date	26/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Acer pseudoplatanus</i>	0.1
<i>Brachypodium sylvaticum</i>	0.3
<i>Carex flacca</i>	0.1
<i>Corylus avellana</i>	80
<i>Crataegus monogyna</i>	20
<i>Eurhynchium striatum</i>	55
<i>Fragaria vesca</i>	0.3
<i>Fraxinus excelsior</i>	7
<i>Frullania dilatata</i>	0.1
<i>Hedera helix</i>	70
<i>Hypnum resupinatum</i>	0.1
<i>Ilex aquifolium</i>	20
<i>Isoetecium myosuroides</i>	0.3
<i>Kindbergia praelonga</i>	0.1
<i>Lonicera periclymenum</i>	0.3
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	0.5
<i>Neckera crispa</i>	1
<i>Potentilla sterilis</i>	0.1
<i>Prunus spinosa</i>	0.3
<i>Quercus robur</i>	0.3
<i>Radula complanata</i>	0.1
<i>Rhytidiadelphus triquetrus</i>	7
<i>Rubus fruticosus</i> agg.	7
<i>Thuidium tamariscinum</i>	3
<i>Ulota bruchii</i>	0.1
<i>Viola</i> sp.	0.3
<i>Zygodon conoideus</i>	0.1



<b>Relevé code</b>	<b>BEC 343</b>
Date	11/09/2014
Recorder	KMN
Species recording scale	Percentage
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Percentage
<i>Anthoxanthum odoratum</i>	0.3
<i>Asplenium ruta-muraria</i>	0.3
<i>Brachythecium rutabulum</i>	0.3
<i>Briza media</i>	3
<i>Calliergonella cuspidata</i>	0.1
<i>Carex flacca</i>	1
<i>Carlina vulgaris</i>	0.3
<i>Corylus avellana</i>	10
<i>Crataegus monogyna</i>	0.3
<i>Ctenidium molluscum</i>	3
<i>Dicranum scoparium</i>	0.1
<i>Epilobium montanum</i>	0.1
<i>Epilobium parviflorum</i>	0.1
<i>Euphrasia salisburgensis</i>	0.1
<i>Festuca ovina</i>	1
<i>Festuca rubra</i>	3
<i>Fissidens dubius</i>	0.1
<i>Fragaria vesca</i>	1
<i>Fraxinus excelsior</i>	0.1
<i>Geranium robertianum</i>	0.3
<i>Hedera helix</i>	3
<i>Homalothecium sericeum</i>	0.1
<i>Hypericum pulchrum</i>	0.1
<i>Hypochaeris radicata</i>	0.1
<i>Koeleria macrantha</i>	0.3
<i>Lapsana communis</i>	0.3
<i>Leucanthemum vulgare</i>	0.3
<i>Linum catharticum</i>	0.1
<i>Lotus corniculatus</i>	1
<i>Neckera complanata</i>	0.1
<i>Neckera crispa</i>	0.1
<i>Pilosella officinarum</i>	0.3
<i>Plantago lanceolata</i>	1
<i>Polygala vulgaris</i>	0.1
<i>Prunus spinosa</i>	0.3
<i>Ranunculus acris</i>	0.1
<i>Rosa spinosissima</i>	1
<i>Rubus idaeus</i>	1
<i>Scleropodium purum</i>	0.3
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	20
<i>Solidago virgaurea</i>	0.5
<i>Sorbus aria</i>	10
<i>Succisa pratensis</i>	0.3
<i>Taraxacum officinale agg.</i>	0.1
<i>Teucrium scorodonia</i>	0.5

Relevé code	BEC 343
Species	Percentage
<i>Thymus polytrichus</i>	1
<i>Tortella tortuosa</i>	1
<i>Trifolium pratense</i>	0.3
<i>Veronica chamaedrys</i>	0.1

Relevé code	BEC 345
Date	04/09/2014
Recorder	JM
Species recording scale	Percentage
Habitat - Fossitt	ER2
Habitat - Annex I	*8240
Species	Percentage
<i>Asplenium ruta-muraria</i>	0.1
<i>Briza media</i>	0.5
<i>Bryum capillare</i>	0.1
<i>Carex flacca</i>	1
<i>Carlina vulgaris</i>	0.5
<i>Centaurea nigra</i>	0.1
<i>Ceterach officinarum</i>	0.1
<i>Crataegus monogyna</i>	0.5
<i>Ctenidium molluscum</i>	15
<i>Dicranum scoparium</i>	0.3
<i>Euphrasia officinalis</i> agg.	0.3
<i>Festuca ovina</i>	1
<i>Festuca rubra</i>	3
<i>Fissidens dubius</i>	0.1
<i>Fraxinus excelsior</i>	0.1
<i>Galium verum</i>	0.3
<i>Grimmia pulvinata</i>	0.1
<i>Hedera helix</i>	1
<i>Homalothecium lutescens</i>	0.1
<i>Hypochaeris radicata</i>	0.3
<i>Koeleria macrantha</i>	1
<i>Leucanthemum vulgare</i>	0.5
<i>Linum catharticum</i>	0.3
<i>Lotus corniculatus</i>	15
<i>Pilosella officinarum</i>	0.5
<i>Plantago lanceolata</i>	0.3
<i>Plantago maritima</i>	0.5
<i>Prunus spinosa</i>	0.5
<i>Rubus fruticosus</i> agg.	0.1
<i>Senecio jacobaea</i>	0.1
<i>Sesleria caerulea</i>	25
<i>Solidago virgaurea</i>	0.3
<i>Sorbus aria</i> agg.	1
<i>Succisa pratensis</i>	5
<i>Teucrium scorodonia</i>	0.3
<i>Thymus polytrichus</i>	0.3
<i>Tortella nitida</i>	0.1

Relevé code	BEC 363
Date	27/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	GS1
Habitat - Annex I	None
Species	Percentage
<i>Agrostis capillaris</i>	15
<i>Agrostis stolonifera</i>	1
<i>Centaurea nigra</i>	0.5
<i>Cerastium fontanum</i>	0.3
<i>Cynosurus cristatus</i>	0.1
<i>Holcus lanatus</i>	85
<i>Lolium perenne</i>	1
<i>Plantago lanceolata</i>	10
<i>Ranunculus acris</i>	0.5
<i>Ranunculus repens</i>	10
<i>Rumex acetosa</i>	0.3
<i>Taraxacum officinale</i> agg.	0.1
<i>Trifolium pratense</i>	0.3
<i>Trifolium repens</i>	5

Relevé code	BEC 366
Date	27/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Brachypodium sylvaticum</i>	0.1
<i>Brachythecium rutabulum</i>	0.1
<i>Corylus avellana</i>	95
<i>Crataegus monogyna</i>	7
<i>Dryopteris affinis</i>	0.5
<i>Eurhynchium striatum</i>	30
<i>Fraxinus excelsior</i>	10
<i>Hedera helix</i>	65
<i>Ilex aquifolium</i>	15
<i>Lonicera periclymenum</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Neckera complanata</i>	1
<i>Potentilla sterilis</i>	0.1
<i>Primula vulgaris</i>	1
<i>Rhytidiadelphus triquetrus</i>	0.5
<i>Rubus fruticosus</i> agg.	0.1
<i>Thamnobryum alopecurum</i>	5
<i>Thuidium tamariscinum</i>	5
<i>Viola</i> sp.	0.3

<b>Relevé code</b>	<b>BEC 404</b>
Date	25/08/2014
Recorder	FON
Species recording scale	Percentage
Habitat - Fossitt	WN2
Habitat - Annex I	None
Species	Percentage
<i>Acer pseudoplatanus</i>	0.1
<i>Carex sylvatica</i>	0.1
<i>Cirriphyllum piliferum</i>	0.1
<i>Corylus avellana</i>	95
<i>Dryopteris dilatata</i>	0.3
<i>Eurhynchium hians</i>	0.1
<i>Eurhynchium striatum</i>	0.5
<i>Fissidens taxifolius</i>	0.1
<i>Fraxinus excelsior</i>	3
<i>Hedera helix</i>	20
<i>Kindbergia praelonga</i>	0.5
<i>Lonicera periclymenum</i>	0.1
<i>Lophocolea bidentata</i>	0.1
<i>Metzgeria furcata</i>	0.1
<i>Phyllitis scolopendrium</i>	0.1
<i>Plagiomnium elatum</i>	0.1
<i>Plagiomnium undulatum</i>	0.1
<i>Polystichum setiferum</i>	1
<i>Potentilla sterilis</i>	0.1
<i>Primula vulgaris</i>	1
<i>Prunus spinosa</i>	1
<i>Pteridium aquilinum</i>	0.1
<i>Quercus robur</i>	1
<i>Rhytidiadelphus triquetrus</i>	7
<i>Rubus fruticosus</i> agg.	0.3
<i>Thamnobryum alopecurum</i>	0.3
<i>Thuidium tamariscinum</i>	0.3
<i>Veronica chamaedrys</i>	0.1
<i>Viola</i> sp.	0.1


## A.8.19.5

---


## Figures

---

Sheet 1:




Sheet 2:





Sheet 3:



Sheet 4:

