

Isotype of Gagea chanae Grossh. [family LILIACEAE]

http://plants.jstor.org/specimen/bm000838144

http://www.jstor.org

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <a href="http://www.jstor.org/page/info/about/policies/terms.jsp">http://www.jstor.org/page/info/about/policies/terms.jsp</a>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the contributing partner regarding any further use of this work. Partner contact information may be obtained at http://plants.jstor.org/page/about/plants/PlantsProject.jsp.


Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact plants@jstor.org.

## Isotype of Gagea chanae Grossh. [family LILIACEAE]

Herbarium	Natural History Museum (BM), BM000838144
Collection	Type specimens
Resource Type	Specimens
Collector	A. Grossheim, No# 55
Collection date	1923-03-12
Locality	Transcaucasia, Tiflis.
Country	Georgia (Georgia)
Identifications	Isotype of Gagea chanae Grossh. [family LILIACEAE] (stored under name); Verified by Zarrei, M., 2004
Data last modified	2009-06-03

http://plants.jstor.org/specimen/bm000838144



## 55. Gagea Chanae Grossheim sp. nova.

4-10 cm, alta. Folia glauca, Folium radicale anguste-lineare, 1-3 mm. latum, subtus carioatum. Folia Iloralia late-lanceolata, inferius basi 3-6 mm. latum, omnia cum pedicellis p. m. ciliato-villosa. Flores 1-4. Perigonii phylla olongo-linearia, obtusa, lutea, 8-13 mm. longa. Pedicoll fructiferi erecti.

Differt a G. pnoilla R. et Sch. foliis glaucis nec vi-ridibus, lattoribus, inflorescentia p. m. villosa, statura rebustiore.

Transcaucasia, Tiflia, in fruticetis, 12-III-23, Leg. A. Grossheim. Ar. geogr. Iberia caucasica,