

POLSKA AKADEMIA NAUK
INSTYTUT ZOOLOGICZNY

A N N A L E S Z O O L O G I C I

Tom XXVI

Warszawa, 15 XI 1968

Nr 3

Maciej MROCKOWSKI

Distribution of the *Dermestidae (Coleoptera)* of the World with a Catalogue of all known Species

Rozmieszczenie *Dermestidae (Coleoptera)* na świecie wraz z katalogiem wszystkich znanych gatunków

Распространение *Dermestidae (Coleoptera)* в мире и каталог
всех известных видов

With 12 maps

A N N A L E S — Z O O L O G I C I

Tom XXVI

Warszawa, 15 XI 1968

Nr 3

Maciej MROCKOWSKI

Distribution of the *Dermestidae (Coleoptera)* of the World with a Catalogue of all known Species

Rozmieszczenie *Dermestidae (Coleoptera)* na świecie wraz z katalogiem wszystkich znanych gatunków

Распространение *Dermestidae (Coleoptera)* в мире и каталог
всех известных видов

[With 12 maps]

CONTENTS

Introduction	16
Remarks on the Arrangement of the Catalogue	17
I. Distribution of the <i>Dermestidae</i> of the World	17
The Neotropical Region	19
The Australian Region	22
The Oriental Region	23
The Ethiopian Region	25
The Palearctic Region	28
The Nearctic Region	31
Some Remarks on the Centers of Origin	32
II. Distribution of the Particular Genera of <i>Dermestidae</i> in the World	35
III. A short Catalogue of all known Species	51
IV. Bibliography	160
V. Alphabetical Index of Names of Subfamilies, Genera, Subgenera and their Synonymical Names	187
Streszczenie	189
Резюме	190

Introduction

The present paper consists of three main parts. In the first part — the most general one — a survey of the *Dermestidae* of the particular zoogeographical regions is given using all available distributional records. The second part contains an analysis of the distribution of the particular genera based on faunistic records of corresponding species. The third part of the paper is the most voluminous one, comprising a taxonomical and nomenclatorial catalogue of all species described so far and an appended list of references. Such a catalogue has never before been published and is filling a gap in the world literature. The only existing catalogue of the *Dermestidae* is that by DALLA TORRE issued in 1911. Unfortunately, it has not been complete and its taxonomical arrangement, as well as its nomenclatorial side, were inaccurate, besides many erroneous literature quotations found in the text. The present catalogue comprises 882 species, 25 subspecies, 127 varieties and 1342 synonymous names. The corresponding figures in DALLA TORRE'S catalogue are 534 species, 2 subspecies, 67 varieties and barely 152 synonymous names.

The preparation of this world catalogue of the *Dermestidae* begun in 1949. During 18 years of intensive work in the Coleopterological Laboratory of the Institute of Zoology of the Polish Academy of Sciences in Warsaw, a card-index with complete literature references as well as available collection data and notes on current investigations by the author have been carefully assembled. They contain information on taxonomy, morphology, anatomy, faunistics, biology, ecology, etc. of the *Dermestidae*. This card-index is now complete with respect to taxonomy and nomenclature and almost complete with regard to distributional data. It formed the basis for the present catalogue. The author is fully aware that the latter is not complete in all its sections, but as far as taxonomy and nomenclature of the species are concerned it is hoped it is as thorough as it could possibly be. The author also tried to present as full an account as possible of distribution records and collecting data with respect to the particular species. Only quotations on those works which contribute substantially to nomenclature (descriptions of new taxa, establishment of new name combinations, new name status and new synonyms) have been included in the catalogue. Results of the author's own studies on taxonomy and zoogeography have also been taken into account.

The manuscript of the catalogue has been sent for criticism to eminent specialists on the *Dermestidae*: to Dr. R. S. BEAL, Jr., Graduate Dean of Northern Arizona University in Flagstaff, to Dr. R. ZHANTIEV of the Lomonosov University in Moscow and to Mr. V. KALIK in Pardubice (Czechoslovakia). They sent their notes and remarks and proposed some corrections — some of them being results of their own unpublished studies. These corrections and suggestions have been fully taken into account in the catalogue and the author wishes to express his best thanks to the persons mentioned for all their kind help and advice.

The 18 years of investigations on the taxonomy, distribution, and nomenclature of the *Dermestidae* are not being closed with the publication of the present catalogue. As mentioned above, these studies and the supplements to the card-index will be continued in the future with the purpose of issuing a more complete catalogue of the *Dermestidae* based on the whole world literature, and enclosing a broader scope of problems.

Remarks on the Arrangement of the Catalogue

All the names, years of publications, page numbers and other data included in the catalogue have been carefully checked with the original literature.¹

Subfamilies and genera within them are arranged in accordance with phyletic relationship, whereas species are placed in their respective genera in alphabetic order. The names of junior synomyms are given twice — first under the valid name (in general the name of the senior synonym) and the second time in the alphabetic list of invalid names which have been used in a given genus.

The names of subfamilies and genera are given as titles; following them are type-genera (for subfamilies) and type species (for genera), as well as synonyms (also with their type genera and type species).

In the sections devoted to the particular genera, a concise alphabetic list of the known species supplemented by their distribution at the margin is given; geographical names in boldface type designate the areas with localities indicated in the original description. Other geographical names have been taken from later publications, not quoted in the present catalogue, mostly faunistic papers, or are based on the author's own studies.

The valid specific name is followed by the name of its author, the year of publication and the corresponding page on which it appeared. These data, in the case when the combination of the specific and generic name is not the original one, are given in parentheses and are followed by the name of the author who first established the combination actually used, with the year of its publication and the page number. In case of synonymous specific names, the name of the author who first published them as synonyms, the year of publication and the number of page are given after a dash. Similarly, when the applied status is not the original one, the name of the author, the year of publication and the number of page referring to the new status of the name are given after a dash.


An alphabetic list of synonymous names, of names of species transferred to other genera or even to other families, of nomina nuda and of all invalid names used in the combination with a given generic name or its synonym, is given within each genus after the list of the known species. Such a system of arranging the text allowed a restriction of the index to generic and family-group names and permitted placing it at the end of the catalogue after the bibliography.

Quotations of papers consist of the author's name and the year of publication of the paper in question. When more than one quoted paper has been published in the same year by a given author, the second paper is marked by the addition of the letter b, the third by the letter c, etc.; the first paper remains without a letter.

I. Distribution of the *Dermestidae* of the World

As a basis for a zoogeographical division of continents, a world map compiled by the specialists of the Institute of Geography and the Institute of Zoology of

¹ Quotations from all but eight works (633) have been taken from their original editions. The eight exceptions are: DUNSTAN, 1923; GRIMMER, 1841; HAUSMANN, 1799; HOPE, 1834; LEACH, 1817; MEGERLE, 1805; SCRIBA, 1790 and 1791.



Map 1. Zoogeographical division of continents.

Regions: I — Neotropical, II — Australian, III — Oriental, IV — Ethiopian, V — Palearctic, VI — Nearctic.

Subregions: I,1 — Antillean, I,2 — Guyano-Brazilian, I,3 — Patagono-Andean, II,1 — Australian, II,2 — New Zealandian, II,3 — Papuan, II,4 — Polynesian, II,5 — Hawaiian, III,1 — Indo-Indochinese, III,2 — Malayan, IV,1 — East-African, IV,2 — West-African, IV,3 — South-African, IV,4 — Madagascar, V,1 — Arctic, V,2 — Euro-Siberian, V,3 — Mediterranean, V,4 — Central-Asiatic, V,5 — Chino-Himalayan, VI,1 — Arctic, VI,2 — Canadian, VI,3 — West-American, VI,4 — East-American.

Provinces: I,2,a — Sonoran, I,2,b — Central-American, I,2,c — other provinces of Guyano-Brazilian Subregion, III,1,a — Indian, III,1,b — Indo-chinese, III,2,a — Sundean, III,2,b — Philippinian, V,2,a — European, V,2,b — Euro-Siberian Taiga, V,3,a — Saharo-Arabian, V,3,b — Mediterranean Proper, V,3,c — Irano-Turanian, V,5,a — Chino-Korean, V,5,b — Japano-Central Chinese, V,5,c — Himalayo-Ammanian.


the Academy of Sciences of the U.S.S.R. has been used. This map is published on pls. 70-71 of the geographical atlas „Fiziko—Geographitsheskiy Atlas Mira” (issued in 1964 in Moscow jointly by the Academy of Sciences of the U.S.S.R. and the Department of Geodesy and Cartography of the Geological State Committee of the U.S.S.R.). An analysis of the distribution of *Dermestidae* revealed that the assumed zoogeographical boundaries do not always correspond to the real distribution of species of that family and should be appropriately altered or corrected. The main changes now proposed are: 1. The Holarctic Region should be divided into two, closely related, regions: the Palearctic and the Nearctic. 2. The Sonoran Province should be placed in the Neotropical rather, than in the Nearctic Region. 3. The Caroline Islands should not be placed in the Australian Region, but transferred to the Philippinian Province of the Oriental Region. The introduced changes and several smaller corrections are indicated on map 1, which shows a zoogeographical division of the continents based on the distribution records of the *Dermestidae*.

Below, the distribution of the *Dermestidae* of the particular zoogeographical regions is discussed. It is worth mentioning that a couple of dozen of Dermestid species are synanthropic and are known as either pests of animal products in storerooms, or as scavengers feeding on various rests and food scraps in human dwellings or in their vicinity. Some of them follow man everywhere and may be regarded as fully cosmopolitan species. Here belong species of the genera *Trogoderma* DEJ. (two species), *Anthrenus* SCHAEFF. (two species), *Attagenus* LATR. (three species) and *Dermestes* L. (five species). It is, of course, difficult to speculate about the original distribution of these 12 cosmopolitan species, as distributional records refer mainly to inhabited areas. We may only surmise that such species of *Trogoderma* DEJ. are Neotropical by origin, while those of *Anthrenus* SCHAEFF. are Palearctic, and the species of *Attagenus* LATR. originated either in the Palearctic or Ethiopian Region; some of the cosmopolitan species of *Dermestes* L. may have originated in the Neotropical Region, some other in the Ethiopian, and still other in the Palearctic Region. Because of the difficulty of establishing the original distribution of those cosmopolitan species, they have been omitted from the descriptions of zoogeographical regions in the following. Species however, which are feebly synanthropic, or are not cosmopolitan in their distribution despite their association as pests with storerooms or their presence in dwellings are, taken into account.

THE NEOTROPICAL REGION

This Region is well delimited and characterised by a great number of endemic genera such as *Hemirhopalum* SHARP — occurring in almost the whole Region (except in the Antillean Subregion, map 2), *Caccoleptus* SHARP — an endemite of the Central-American Province, the genera *Decamerus* SOL. and *Diontolobus* SOL. — known only from Chile, or the genera *Falsocryptorhopalum* PIC and *Pseudothaumaglossa* PIC occurring only in Brazil. A further peculiarity of the Neotropical Region is the complete lack of species of *Anthrenus* SCHAEFF. and *Attagenus* LATR., as well as a small number of species of *Dermestes* L.

The genera *Apsectus* LEC. and *Cryptorhopalum* GUÉR. — although penetrating the southern part of the Nearctic Region — should also be classified as originally Neotropical groups. Their range is given on map 2. The genus *Apsectus* LEC. is mainly distributed in the Central-American and Sonoran Provinces — the latter being wrongly placed in the Nearctic Region. Only one species reaches northward beyond the Sonoran Province. *Cryptorhopalum* GUÉR. is a large genus including about two thirds of the number of species of Dermestidae known from the Neotropical Region. This genus is represented by a few species in the Patagono-Andean Subregion while most species occur mainly in the Guyano-Brazilian Subregion, and are most common in the Central-American Province. Numerous




Map 2. Distribution of the genera *Cryptorhopalum* GUÉR. (1), *Hemirhopalum* SHARP (2) and *Apsectus* LEC. (3).

species of the genus in question are also found in the Sonoran Province, and this clearly indicates its relationship with the Central-American Province and its Neotropical character. Only a few species occur in North America outside the Sonoran Province, and only in areas adjoining it. Other evidence proving the Neotropical character of the Sonoran Province is given below, in the section devoted to the Nearctic Region.

One species of *Apsectus* LEC. and several of *Cryptorhopalum* GUÉR., which occur in the Nearctic Region outside the Sonoran Province, can be regarded as immigrants from the Neotropical Region. Further evidence on certain links between the two regions is the occurrence in the Nearctic of some species of *Trogoderma* DEJ. — a genus which is represented by numerous species in the Neotropical Region, as well as the occurrence of a few species of *Dermestes* L., which is mainly distributed in the Palearctic, but is found in the Nearctic and Neotropical Regions (in the latter four species occur) as well. The links existing between the last two regions are clearly feeble.

There is, however, a distinct affinity between the Neotropical and Australian Regions. Both are well inhabited by species of *Trogoderma* DEJ. (map 3). In the Neotropical Region, 30 species of this genus are found, distributed more or less regularly in all Subregions. In the Australian Region 60 species occur, mostly in the Australian and New Zealandian Subregions, but two species have been also recorded from the Polynesian Subregion.




Map 3. Main distribution areas of the genus *Trogoderma* DEJ.

The fauna of the Ethiopian and Oriental Regions does not show any similarity with that of the Neotropical Region, and the connection between the latter and the Palearctic seems to be extremely feeble; several species of *Trogoderma* DEJ., as mentioned before, occur in the Palearctic, and several species of *Dermestes* L. — mostly a Palearctic genus — are found in the Neotropical Region.

THE AUSTRALIAN REGION

This Region is well delimited. It is characterized by a large number of endemic genera such as *Anthrenocerus* ARR., *Myrmeanthrenus* ARMSTR., *Neoanthrenus* ARMSTR. (endemites of the Australian Subregion), *Hexanodes* BLAIR (endemite of the New Zealand Subregion), *Argocerus* SHARP, *Eocerus* SHARP, *Labrocerus* SHARP (endemites of the Hawaiian Subregion). It is characterized as well by the absence of species of genera widely distributed in other zoogeographical regions, e. g. *Anthrenus* SCHAEFF., *Dermestes* L. and *Attagenus* LATR. The distribution of larger endemic genera is shown on map 4.



Map 4. Distribution of the genera *Anthrenocerus* ARR. (1), *Neoanthrenus* ARMSTR. (2), and *Labrocerus* SHARP (3).


Although being a distinct zoogeographical entity, the Australian Region shows a certain affinity with the Neotropical Region. Both regions are inhabited by many species of *Trogoderma* DEJ. — in the Australian Region 60 species of this genus are found, and this represents a half of the total number of species of *Dermestidae* recorded from this region. They occur mainly in the Australian and New Zealand Subregions. Certain links may be also found between the Australian Region and the Oriental and Ethiopian Regions. An example may be the distribution of the species of *Orphinus* MOTSCH. (map 5) of which 16 species occur in the Australian Region (11 in the Australian Subregion, three in the Polynesian and two in the Papuan Subregion), 25 species in the Oriental Region, 10 species in the Ethiopian Region and three species in Japan (one of these also in Korea). Another example is the presence of one species of *Thaumaglossa* REDT. (map 6) in the Australian Subregion. The remaining species of this genus are distributed in the Oriental (10 species), Ethiopian (10 species) and Nearctic (two species) Regions.

The Carolines, placed in the Polynesian Subregion, do not show any connection with the Australian, but with the Oriental Region and should be regarded as part of the Philippines Province of the latter; this is discussed by the author more in detail in the section devoted to the Oriental Region.

Leaving aside the species introduced by man, the Australian Region exhibits practically no links with the Palearctic and Nearctic Regions.

THE ORIENTAL REGION

Many genera of *Dermestidae* inhabiting the Oriental Region are common to other regions as well, and only one genus — *Evorinea* BEAL — can be regarded as truly endemic to the region in question. A complete absence of species of *Trogoderma* DEJ. is a further



Map 5. Distribution of the genus *Orphinus* MOTSCH.


peculiarity of the Oriental Region (the genus is well represented in the Australian Region).

Faunistic links between the Oriental and Australian Regions are feeble — each is inhabited by only one species of the genus *Adelaidia* BLACKB. Of 54 species of *Orphinus* MOTSCH. (map 5), 25 species occur in the Oriental Region (the majority in the Malayan Subregion), 16 species in the Australian, 10 in the Ethiopian Region and three in Japan. The occurrence in the Carolines of *Evorinea* BEAL, which is endemic to the Oriental Region, indicates that these islands should be included in the Philippines Province; this may be further supported by the absence in the Caroline Islands of genera occurring exclusively in the Australian Region, as well as by the absence of species of *Trogoderma* DEJ.

The distribution of the genus *Orphinus* MOTSCH. indicates, on the other side, a certain affinity between the fauna of the Oriental and Ethiopian Region. This affinity is further shown by the distribution of several other genera such as *Thorictodes* REITT., *Trinodes* DEJ., *Thaumaglossa* REDT. or the subgenus *Aethriostoma* MOTSCH., belonging to the genus *Attagenus* LATR. (map 6). All these genera occur in both regions mentioned.

There are many common elements in the Oriental and Palearctic Regions. Apart from the genus *Trinodes* DEJ. (map 7), already mentioned, which is represented by an equal number of species (seven) in each of them, a further example may be provided by the genera *Anthrenus* SCHAEFF., *Attagenus* LATR. or *Dermestes* L. The Oriental species of these three genera, which occur mainly in the Palearctic, suggest certain links between the Palearctic and the Oriental Region. A great majority of these species are distributed in areas adjacent to the Palearctic Region.

There are no elements common to the Oriental and Neotropical Region, and links between the Oriental and Nearctic Regions are indirect, certain species of *Anthrenus* SCHAEFF., *Dermestes* L. and *Attagenus* LATR. penetrating both regions from the Palearctic. The distribution of *Thaumaglossa* REDT., of which 10 species occur in the Oriental, 10




Map 6. Distribution of the genus *Thaumaglossa* REDT. (1) and the subgenus *Aethriostoma* MOTSCH. of the genus *Attagenus* LATR. (2).

species in the Ethiopian, two species in the Nearctic and one species in the Australian Region (map 6) may provide another example of a certain affinity of the regions mentioned.

THE ETHIOPIAN REGION

Like the Oriental, the Ethiopian Region does not abound in endemics. Only two monospecific genera — *Paratrogoderma* SCOTT from the Seychelles and *Phradonomorphus* PIC from Senegal — may be regarded as endemic to the Ethiopian Region, and the absence of species of *Trogoderma* DEJ. is a further peculiarity of this region.

The links with the Palearctic seem to be the strongest. The large genus *Attagenus* LATR. (which apart from the species of the subgenus *Aethriostoma* MOTSCH., does not occur in the Madagascar Subregion) and *Phradonoma* JACQ. DU VAL are represented almost by an equal number of species in the two regions mentioned: in the Ethiopian Region 64 species of *Attagenus* LATR. and seven species of *Phradonoma* JACQ. DU VAL occur, while 77 species of *Attagenus* LATR. and 11 species of *Phradonoma* JACQ. DU VAL (map 8) are found in the Palearctic Region. Although *Ctesias* STEPH., *Anthrenus* SCHAEFF.




Map 7. Distribution of the genus *Trinodes* DEJ.

and *Dermestes* L. are mainly Palearctic in their distribution, a number of species of these genera has also been recorded from the Ethiopian Region; 14 species of *Anthrenus* SCHAEFF. occur in the East-African and South-African Subregions, six species of *Dermestes* L. are found in all subregions except in the South-African, and one species of *Ctesias* STEPH. is known from Rhodesia and the Congo (Kinshasa). Four species of *Trinodes* DEJ. have been recorded from the East-African and West-African Subregions, which not only accounts for a certain affinity with the Palearctic (where seven species are found) but also with the Oriental Region from which seven species of the said genus (map 7) have been recorded as well.


Further evidence of a certain faunistic affinity between the Oriental and Ethiopian Regions may be provided by the distribution of *Orphinus* MOTSCH. (which in the Ethiopian Region is absent only from the South-African Subregion) and *Thaumaglossa* REDT. (map 6). There is also a certain penetration by Palearctic elements — species of *Anthrenus* SCHAEFF., *Attagenus* LATR. and *Dermestes* L., — into the Ethiopian Region; these are found in the Oriental Region as well.

The Australian and the Ethiopian Regions show hardly anything in common; only the range of *Orphinus* MOTSCH. (map 5) comprises them both. No common elements exist for the Ethiopian and Neotropical Regions.

Links between the Nearctic and Ethiopian Regions are at most indirect; there is a number of species of *Ctesias* STEPH., *Anthrenus* SCHAEFF., *Attagenus* LATR. and *Dermestes* L. which are found in both regions mentioned, the majority of species, however, occur in the Palearctic from where they penetrated into the Ethiopian and Nearctic Regions.



Map 8. Distribution of the genera *Phradonoma* JACQ. DU VAL (1) and *Attagenus* LATR. (without the subgenus *Aethriostoma* MOTSCH.) (2).




Map 9. Distribution of the genera *Anthrenus* SCHAEFF. (1) and *Dermestes* L. (2).

Another example of a link between the two latter may be the distribution of *Thaumaglossa* REDT. (map 6) of which 10 species are spread over the whole of the Ethiopian Region, two species occur in the Nearctic, 10 species in the Oriental and one species in the Australian Region.

THE PALEARCTIC REGION

A seemingly great morphological diversity of the Palearctic *Dermestidae* and the relatively highest number of genera so far recorded from this region are certainly, at least partly, due to a better knowledge of its fauna, thanks to more extensive and more thorough investigations carried out in this region than in other areas. Although a great number of endemites have been found in the Palearctic, they occur mostly in the Mediterranean and Central-Asiatic Subregions; *Ditorines* NORM., *Sefrania* PIC, *Hirtomegatoma* PIC, *Mariouta* PIC, *Egidyella* REITT., *Montandonia* JACQUET and *Globicornis* LATR. (map 10) should be mentioned here. Only the last of these genera has a fairly wide distribution and several of its 21 species occur also in Europe. A great number of Palearctic species belong to genera which may be regarded as in principle Palearctic groups. These are: *Dermestes* L. (with 73 species of which 44 Palearctic), *Anthrenus* SCHAEFF. (with 83 species of which 55 Palearctic) (map 9) and *Ctesias* STEPH. with its six species of which four are Palearctic. Closest links exist between the fauna of the Palearctic and the Ethiopian Region and they are typified by an equal number of species of the large genera — *Attagenus* LATR. and *Phradonoma* JACQ. DU VAL, occurring in each of these regions.

There is hardly a link between the region discussed and the Australian and Neotropical Regions; several species of the Australian and Neotropical genus *Trogoderma* DEJ.,



Map 10. Distribution of the genera *Globicornis* LATR. (1), *Ditorines* NORM. (2), *Egidyella* REITT. (3), *Hirtomegatoma* PIC (4), *Mariouta* PIC (5), *Montandonia* JACQUET (6) and *Sefrania* PIC (7).

which also occurs in the Palearctic Region, and several species of the Palearctic *Dermestes* L. found in the Neotropical Region may be the only examples here.

There seem to be few links between the Palearctic and Oriental Regions. Several species of *Dermestes* L., *Attagenus* LATR. and *Anthrenus* SCHAEFF. are found in the Oriental Region. The distribution of the genus *Trinodes* DEJ., though, may witness a closer affinity of the fauna of the two regions in question: there is an equal number of species of this genus in each region (seven) and four species of *Trinodes* DEJ. also occur in the Ethiopian Region.

The fauna of the Ethiopian and Nearctic Regions indicates the closest relationship with that of the Palearctic. The connection between the Ethiopian and Palearctic Regions seems to be the most striking. Two large genera — *Attagenus* LATR. and *Phradonoma* JACQ. DU VAL are represented by a large number of species in these two regions: 77 species of *Attagenus* LATR. occur in the Palearctic, while 64 species are found in the Ethiopian Region; the corresponding figures for *Phradonoma* JACQ. DU VAL are 11 species in the Palearctic and seven species in the Ethiopian Region (map 8). The absence of species of both these genera in the Eastern Asiatic part of the Palearctic — east of the Lake Baykal is symptomatic. Only two cosmopolitan and synanthropic species of *Attagenus* LATR. are known so far from that area. The Palearctic species of these two genera concentrate around their probable origin centers: the Mediterranean and the Central-Asiatic.

A fair number of Palearctic species also penetrate the Ethiopian Region. They belong to the genera *Dermestes* L. (six species), *Anthrenus* SCHAEFF. (14 species) and *Ctesias* STEPH. (one species); four species of the Palearctic and Oriental genus *Trinodes* DEJ. have also reached the Ethiopian Region.

Links between the Nearctic and Palearctic Regions are less conspicuous. Only two genera, not particularly rich in species, — *Dearthrus* LEC. (four species) and *Megatoma* HERBST (21 species) are more or less similarly represented in both these regions (map 12). One species only of the Nearctic *Orphilus* ER. is found in the Palearctic Region, and a certain number of species of the Palearctic genera — *Anthrenus* SCHAEFF., *Dermestes* L. and *Ctesias* STEPH., as well as the Ethiopian-Palearctic *Attagenus* LATR. penetrate the Nearctic Region. Differences in the fauna of the two regions in question are, on the other hand, considerable. There is not a single species of *Phradonoma* JACQ. DU VAL, *Trinodes* DEJ. and *Globicornis* LATR. in the Nearctic, and numerous genera endemic to the Palearctic are equally absent there. Numerous genera represented in the Nearctic Region, such as *Novelsis* CAS., *Reesa* BEAL, *Apsectus* LEC. or *Cryptorhopalum* GUÉR. — which may be regarded as a Neotropical element in the Nearctic Region — are lacking in the Palearctic.

The eastern part of the Palearctic — more or less east of the Lake Baykal — differs considerably from the rest of the Region. As mentioned before, the eastern part of the Palearctic lacks species of some genera which are abundantly represented in the western Palearctic, such as *Attagenus* LATR. and *Phradonoma* JACQ. DU VAL. Neither species of *Ctesias* STEPH., nor genera endemic to the Palearctic — *Globicornis* LATR., *Montandonia* JACQUET, *Egidyella* REITT., *Mariouta* PIC, *Hirtomegatoma* PIC, *Sefrania* PIC and *Ditorines* NORM. are found in the eastern part of the Palearctic; moreover, rich in species subgenera of *Anthrenus* SCHAEFF.: *Anthrenops* REITT. and *Nathrenus* CAS. are also missing in the area in question.

An analysis of the distribution of the particular species of *Dermestes* L. in the Palearctic also supports this division of the Palearctic into a western and an eastern part: the latter is inhabited by a number of endemic species — *D. tessellatocollis* MOTSCH., *D. vorax* MOTSCH., *D. coarctatus* HAR., *D. murinus auriceps* REITT. as well as *D. nidum* ARR. — which do not range further westward, but, which on the other hand, are known from the Nearctic Region. There are, however, very few species in the eastern part of the Palearctic, which occur in the western Palearctic as well. Some endemic species of *Anthrenus* SCHAEFF. (*A. sinensis* ARR. of the subgenus *Solskinus* MROCKZ. and *A. coreanus* MROCKZ. of the subgenus *Florilinus* MULS. and REY) are distributed in the eastern Asiatic part of the Palearctic, similarly as some species in the genus *Dermestes* L., while an almost complete absence (apart from synanthropic species) of species widely distributed and common in the western Palearctic, is characteristic to that area. Of the seven Palearctic species of *Trinodes* DEJ. three occur in the East: two in Japan and one in northern China; the remaining are distributed in the Mediterranean Subregion, mostly in North Africa.



Map 11. Distribution of the genera *Novelsis* CAS. (1) and *Reesa* BEAL (2).


THE NEARCTIC REGION

Few endemic genera but many common elements with the Palearctic characterize the region in question. Certainly, the fauna of the Nearctic is closely related to that of the Palearctic. The affinities, however, are not so deep as to allow to treat both regions as one — a “Holarctic” — which, unfortunately, is done in many zoogeographical publications.

Endemic are *Novelsis* CAS., occurring only in the West-American Subregion, and the broadly distributed *Reesa* BEAL (map 11).

There are few links between the Nearctic and Neotropical Regions. Few species of the Neotropical *Apsectus* LEC. and *Cryptorhopalum* GUÉR., as well as several species of the Australo-Neotropical *Trogoderma* DEJ., penetrate the Nearctic. The Sonoran Province should be excluded from the Nearctic Region and incorporated into the Neotropical, as there is ample evidence (see Neotropical Region) that it is more closely related with the latter than with the former. The absence in the Sonoran Province of species of *Dearthrus* LEC., *Orphilus* ER., *Reesa* BEAL or *Megatoma* HERBST — typical for the Nearctic — may additionally justify this arrangement.

No direct connections exist between the Nearctic and the Australian Regions. Certain links may be suggested by the presence in the Nearctic of several species of *Trogoderma* DEJ. — a principally Australian (and Neotropical) genus.



Map 12. Distribution of the genera *Megatoma* HERBST (1), *Dearthrus* LEC. (2), and *Ctesias* STEPH. (3)

The Oriental and Ethiopian Regions also show merely indirect links with the Nearctic. The occurrence of Palearctic elements — species of *Anthrenus* SCHAEFF., *Attagenus* LATR. and *Dermestes* L. in the Nearctic, Ethiopian and Oriental Regions may be here an example. The distribution of *Thaumaglossa* REDT. — a genus not occurring in the Palearctic, but represented by two species in the Nearctic, 10 species in the Ethiopian and 11 species in the Oriental Region, is a different case of connections between the three regions.

As mentioned before, the Nearctic and Palearctic are strongly related. The colonisation of the Nearctic by Palearctic elements has been particularly strong. Of the mainly Palearctic genera — *Anthrenus* SCHAEFF., *Attagenus* LATR., *Ctesias* STEPH. and *Dermestes* L. (the genus *Attagenus* LATR. is Palearctic as well as Ethiopian) a certain number of species occur only in the Nearctic: four species of *Anthrenus* SCHAEFF. (54 Palearctic species), nine species of *Attagenus* LATR. (77 in the Palearctic and 64 species known from the Ethiopian Region), one species of *Ctesias* STEPH. (four in the Palearctic), 15 species of *Dermestes* L. (44 Palearctic species) occur exclusively in the said region. The penetration of the Palearctic Region by Nearctic faunal elements seems to be much feebler — the distribution of the genus *Orphilus* ER. with four Nearctic and one Palearctic species seems to be the only example here. One can, thus, apart from certain faunistic affinities consisting of the penetration of certain species of one region into another, hardly speak, of a unity of the two regions; there are, though, two genera — *Megatoma* HERBST and *Dearthrus* LEC. (map 12) evenly distributed in both regions, but they include few species, and they alone can not support the contention that the Palearctic and Nearctic should be treated as an entity. When the entire family is considered, the division of the Holarctic is plainly evident. Some other examples in favour of this division are given in the section devoted to the Palearctic.

Some Remarks on the Centers of Origin

It is very difficult, in view of an almost complete lack of fossils, to hypothesize about the place of origin of the *Dermestidae*. A particular concentration of species in certain areas does not exclude the possibility that their ancestors were of foreign origin and that the speciation process was here simply more intensive than in other areas. The dispersion of distribution centers seems to bear out this contention. Both zoogeographical and morpho-anatomical data reflect only the latest stages of evolution which have been reached by species and particular groups, and do not provide any insight about the past, except by indirect conclusions. These data may clear out certain phylogenetic relationships within a relatively short period of time as they enable us to infer about the origin of genera, the relations between them, or to built up "phylogenetic trees" within them. This has been done, in some monographs of particular genera, for instance in the recent works by BEAL (1967) and the author's own paper (1967) on the genus *Megatoma* HERBST.

However, to get certain hints about the origin and ancestors of the family as a whole, zoogeographical and morphological data must be supplemented by data gained from paleontological material. When fossils are missing altogether one must give up trying to "decipher" the past and limit himself to provide and to explain merely the present pattern of distribution.

Regions

Neotropical Australian Oriental Ethiopian Palearctic Nearctic

Marioutinae WINKL.*Mariouta* PIC**Egidyellinae** SEM.*Egidyella* REITT.**Dermestinae** LATR.*Dermestes* L.*Montandonia* JACQUET**Attageninae** CAST.*Sefrania* PIC*Attagenus* LATR.*Novelsis* CASEY*Decamerus* SOL.**Megatominae** LEACH*Trogoderma* DEJ.*Paratrogoderma* SCOTT*Reesa* BEAL*Phradonomorphus* PIC*Phradonoma* JACQ. DU VAL*Globicornis* LATR.*Hirtomegatoma* PIC*Megatoma* HERBST*Labrocerus* SHARP*Eocerus* SHARP*Argocerus* SHARP*Dearthus* LEC.*Ditorines* NORM.*Caccoleptus* SHARP*Diontolobus* SOL.*Anthrenocerus* ARROW

X = introduced species.

? = species most probably belonging to other genera.

R e g i o n s

Neotropical Australian Oriental Ethiopian Palearctic Nearctic

Myrmeanthrenus ARMSTR.
Adelaidia BLACKB.

Orphinus MOTSCH.
Curtophinus PIC
Ctesias STEPH.
Tiresiomorpha PIC
Thaumaglossa REDT.
Pseudothaumaglossa PIC
Hemirhopalum SHARP
Falsocryptorhopalum PIC

Cryptorhopalum GUÉR.

Anthreninae LEC.

Anthrenus SCHAEFF.

Neoanthrenus ARMSTR.

Orphilinae LEC.

Orphilus ER.

Trinodinae CASEY

Apsectus LEC.

Trinodes DEJ.

Evorinea BEAL


Hexanodes BLAIR

Thylodriadinae MROCK.

Thylodrias MOTSCH.

Thaumaphrastinae ANDERS.

Thorictodes REITT.



x = introduced species.

Actually we see several distribution centers of the *Dermestidae*, differing well from one another. Two such centers exist in the Palearctic Region: one in the Mediterranean Proper Province, the other in the Central-Asiatic Subregion. Other distribution centers are found in the Sonoran Province and Central-American Province of the Guyano-Brazilian Subregion of the Neotropical Region. A distinct center also can be distinguished in the eastern part of the Australian Subregion.

All these centers are characterised by: 1) Occurrence of groups comprising a large number of closely related species, morphologically similar. Such groups may belong in a given center to one or more genera. 2) Occurrence of a considerable number of species producing morphological varieties (subspecies, races or aberrations). 3) Occurrence of a large number of species showing considerable individual variation. 4) Expansion of particular forms into adjacent areas. All these features may suggest a more intensive speciation process in the centers than outside them.

II. Distribution of the Particular Genera of *Dermestidae* in the World

Subfamilies and genera within them are arranged according to their phyletic relationships. An appended diagram (pages 33 and 34) shows number of the species of the particular genera in every zoogeographical region. The height of particular columns corresponds to a number of species in a given region, every species being "represented" by 0.2 mm of the column height.

This system enables an easy comparison of the number of species of every genus in the particular regions. A more detailed analysis of the distribution of each genus is given below — it is not, however, based on diagram figures — which merely serve to give a quick insight into the "distribution pattern" of the family — but resulted from a detailed study of the distribution of the particular species included in part III — the catalogue of the species.

Marioutinae WINKLER

Mariouta PIC (map 10)

The genus comprises two eremic species with disjunctive distribution: one has been recorded from Turkmenia, the other from Egypt and the Maghreb Countries.

Egidyellinae SEMENOV-TIAN-SHANSKIJ

Egidyella REITT. (map 10)

Monospecific eremic genus, endemic to Central Asia: it is known only from Turkmenia and Uzbekistan.

Dermestinae LATREILLE*Dermestes* L. (map 9)

The genus includes 73 species, closely related. The majority of them inhabit the Palearctic and Nearctic. Apart from five cosmopolitan species, known as pests in storerooms, which have been introduced everywhere by man, there is a complete absence of species belonging to this genus in the Australian Region. In the Oriental Region, too, the genus is poorly represented — only three species have been so far recorded — namely from the Indian Province — all were found in northern areas adjacent to the Palearctic; these are: *D. fulvicollis* REITT. in the Himalayas, *D. kurseongensis* LEP. in Sikkim, and *D. leechi* KAL. in India (known from Turkmenia as well). Moreover, a fossil species is known from the amber in Burma. There are also only a few species of *Dermestes* L. in the Ethiopian Region. Only six of them are known to be exclusively Ethiopic, of which two have been taken in Madagascar, two in Congo (Kinshasa), one in Mozambique and one in Kenya and Tanganyika. There is a complete lack of species of the genus in question in the South-African Subregion.

Merely four species are known from the Neotropical Region, one of them — *D. impressicollis* REITT., has been described from South America, unfortunately, without any details about the collecting place; apart from the description, there is nothing known about this species. Still another species has been recorded from Brazil, one from Ecuador, and one is known from Chile, Peru, Argentina, Bolivia, Mexico, the United States and Europe — it has been introduced to the U.S.A. and Europe. In addition, four Nearctic species penetrate the Antillean Subregion.

The great majority of the species of *Dermestes* L. — 59 of them — are known from both the Palearctic and Nearctic, which may indicate a certain affinity of these regions. In particular, the distribution of *D. nidum* ARR. — which occurs almost everywhere in the U.S.A. — as well as in the Chino-Korean Province of the Palearctic Region, and the distribution of *D. undulatus* BRAHM in the whole Palearctic and Nearctic, suggest close links existing between the faunas of the regions in question.

There are only 15 exclusively Nearctic species. In the main they do not range beyond Canada and the U.S.A., where they are fairly uniformly distributed. Only four species are known to occur in the Antilles and two in Mexico, as well.

About half of the number of the species of *Dermestes* L. so far described have been recorded from the Palearctic Region. Three species are found everywhere in the Palearctic, while two are confined to the Asiatic part of the Region; the remaining Palearctic species are distributed over relatively small areas. Three East-Palearctic species: *D. coarctatus* HAR., *D. tessellatocollis* MOTSCH. and *D. vorax* MOTSCH. occur in Japan, Korea, North China and the Soviet territories of the Far East. A similar distribution is also shown by *D. murinus auriceps* REITT. which is a subspecies of a species distributed over the entire Palearctic Region. The distribution of all these forms seems to support the division of the Palearctic into an eastern and a western part. The distribution already discussed of

D. nidum ARR. — which is restricted in the Palearctic to the Far East — may be a further argument in this regard.

The distribution areas of all remaining species overlap and it is difficult, and sometimes impossible, to draw boundaries between them. These combined areas cover North Africa, Europe, Asia Minor and Central Asia as well as Siberia. Only one species penetrates into Sinkiang. There are but few species which have not been taken in Europe. Three of them occur in Central Asia (one of them reaches India), one in Iran, one in Egypt and finally one in the Maghreb Countries. The majority of species have been found at least in some parts of Europe. Thus three species range from Central Europe, over the Caucasus to Central Asia and Siberia. Two species were found to be distributed from South-Eastern Europe to Central Asia, one even reaching Sinkiang. Four species are spread from Central and South Europe to Asia Minor, the Caucasus, Syria and Iraq. One species occurs in South Europe and in the Caucasus. Three species are endemic to Europe; one is found in Sweden and Finland, and two in South Europe. One species is distributed over the whole Mediterranean Proper Province. The last group consists of species found in the Maghreb Countries, in the Pyrenean Peninsula, in France and in Sardinia.

Montandonia JACQUET (map 10)

A genus of only one species, erected originally for *Montandonia latissima* (BIEZL) which occurs in the East Carpathians (Romania).

Attageninae CASTELNAU

Sefrania PIC (map 10)

Monospecific genus, occurring exclusively in the Maghreb Countries.

Attagenus LATR. (maps 6 and 8)

The largest genus of the *Dermestidae*, including 170 species. Eight species differ considerably from the remaining ones in their morphology and are grouped in the subgenus *Aethriostoma* MOTSCH. — which is regarded by some authors as a separate genus. Of these eight species four occur in the Madagascar Subregion, one is very widely distributed — over almost the whole of the Oriental Region, one species is known from South China and the Himalayan Mts., one from Sumatra and one from New Guinea.

The remaining species of the genus in question form a distinctly separated, well characterized group. Here belong the subgenera *Attagenus* s. str., *Lanorus* MULS. et REY,

Ranolus BLAIR and *Telopes* REDT., but, in the author's opinion there are no sufficient grounds for such a classification, at least as far as the first three of the subgenera mentioned are concerned. Only the species of *Telopes* REDT. differ more substantially from the others and placing them in a separate subgenus may be more justified; this, however, would also require treating the subgenus *Aethriostoma* MOTSCH. as another separate genus. For convenience, the distribution of the subgenera *Telopes* REDT., *Ranolus* BLAIR, *Lanorus* MULS. and REY and *Attagenus* s. str. is discussed jointly by the author, as has been done above for the distribution of the subgenus *Aethriostoma* MOTSCH.

The complete absence of species of *Attagenus* LATR. in the Australian, the Neotropical, and in the eastern part of the Palearctic Region (approximately east of the Lake Baykal), as well as the occurrence of only a small number of species in the Oriental and Nearctic Region, and, on the other hand, their abundance in the Ethiopian and western Palearctic Region (particularly in the south of the latter), seem to be most conspicuous features of the distribution of the said genus.

From the Neotropical Region three species are known, but one of them — *A. diversepunctata* PIC — was described from one single specimen found in Dresden (Germany) and considered an import from Costa Rica, what, however, seems very improbable. The second species — *A. adspersus* (BLANCH.) — described from Bolivia as *Megatoma adspersa* BLANCH. has been, probably unjustly, transferred by LACORDAIRE to the genus *Attagenus* LATR. Finally, the last species — *A. brasiliensis* PIC — was described from Brazil, but the description is so superficial that it does not reveal whether the species was an "*Attagenus*" or not. The possibility that this species belongs to some other genus can not be excluded and a thorough examination of the type material may yet solve this question.

Three species are cosmopolitan and one is distributed in the whole Holarctic. They are storage pests and this accounts for their distribution. The author omits them from further consideration.

Nine recent species and two Miocene forms — discovered in Colorado — are known from the Nearctic Region. Of the recent species one occurs in Canada and eight in the United States, of these three have been collected in the southern States only. Furthermore, one Palearctic species which occurs in the Maghreb Countries and in Central Asia — *A. cyphonoides* REITT. — was recently found in the south of the United States and in the Californian Peninsula (R. S. BEAL Jr. in litteris, 1967). It is not improbable that at least some of the Nearctic species will turn up as synonyms of the Palearctic forms following a future revision of the Nearctic species of *Attagenus* LATR. Such a study is currently being undertaken by R. S. BEAL, Jr.

Eleven species are known from the Oriental Region. Four of them have been recorded from the Indian Province, seven from the Indochinese Province — five from South China (of which three from Yunnan only), one from Burma and one from Viet-Nam. Two species occur in the Malayan Subregion — on the Malayan Peninsula — of these one was taken in New Caledonia and Zanzibar as well, where it has been most probably introduced.

Apart from three cosmopolitan species, 61 species have been recorded from the Ethiopian Region. Three of them occur in the Palearctic as well, and their distribution is being discussed in detail below. Of the remaining species one occurs in Yemen, nine in Ethiopia, 26 in Equatorial Africa and 22 in the South African Subregion. They are absent, however, in the Madagascar Subregion. A more detailed zoogeographical analysis is unfortunately not possible due to the scarcity of literature data (generally there are no records aside from the „locus typicus” indicated in the original description) and a lack of preserved material. The above mentioned three Palearctic species which are also found in the Ethiopian Region, have all been recorded from the Maghreb Countries. Moreover, *A. heydeni* (REITT.) is known from Egypt, and the former French West Africa; *A. posticalis* FAIRM. is known from Egypt, Syria, Yemen, the former French West Africa and from the Pyrenean Peninsula; finally *A. cinnamomeus* ROTH was taken in Egypt, Syria, Yemen, Ethiopia and Senegal.

In the Palearctic more species of *Attagenus* LATR. occur than in the Ethiopian Region — 77 species altogether — of which three are cosmopolitan, one is Holarctic, and three enter the Ethiopian Region. The absence of members of this genus in the eastern Palearctic east of the Lake Baykal is striking. Only two cosmopolitan species are known from that area, one of which forms an endemic eastern Palearctic subspecies — *A. unicolor japonicus* REITT., which extends westward to Mongolia. *A. amurensis* PIC, described from Siberia without any indication as to locality appears to be a synonym of *A. unicolor japonicus* REITT., but the description itself is very short and inaccurate and until results of a reexamination of the type material have been known it is difficult to say with certainty.

The Palearctic species mainly concentrate in the southern areas of this region — in the Mediterranean Subregion. From the Maghreb Countries 35 species are known, of which 17 occur there exclusively. Three species occur also in South-Western Europe, four in Egypt and in the Near East, and eight in the whole Mediterranean Proper Province. Of the latter, four species range through Asia Minor to the Caucasus — two of them extending eastward to Central Asia (one even to Mongolia).

Nine species are restricted to Egypt and the Near East (Syria, Israel, Jordan). From Italy, Sicily and Malta three species have been described. Two species are fairly widely distributed in Europe and in the Caucasus, and one in South Europe, Asia Minor and the Caucasus.

The remaining 21 species are distributed in the southern part of the Asiatic Palearctic. Three of them occur in Asia Minor, two also in the Caucasus and one of the latter in Central Asia and Mongolia as well. Two species extend from the Caucasus through Iran to Central Asia and Mongolia. Five species occur only in Transcaucasia, northern Iran and Turkmenia. Similarly, five species have been recorded only from Central Asia, and six species range from Central Asia, through Sinkiang to Mongolia, and even have been found not far from the Baykal Lake. One species is known from Tibet.

Aside of the Maghreb Countries, Central Asia may be considered a second distribution centre of *Attagenus* LATR. Altogether 22 species are known from the area in question, of which one species is common to the Maghreb Countries as well. The remaining Palearctic species are distributed between the two mentioned centers.

Novelsis CASEY (map 11)

A Nearctic genus comprising nine species distributed in the Western American Sub-region, between the Pacific and the Rocky Mts. One species occurs in British Columbia, two in Washington, one in Oregon, two in Idaho, one in Utah, two in Nevada, seven in California, six in Arizona, three in New Mexico, four in Texas; two are known from the Californian Peninsula and three species from the other parts of Mexico.

Decamerus SOLIER

A monospecific Chilean endemite.

Megatominae LEACH*Trogoderma* DEJ. (map 3)

One of the largest genera — comprising 120 species, of which the majority inhabit the Australian (60 species) and Neotropical (30 species) Regions. *T. granarium* Ev. — a synanthropic species — is cosmopolitan in distribution.

In the Australian Region the greatest number of species — 48 of them — are found in the Australian Subregion. The New Zealand Subregion is inhabited by 10 species, and from the Polynesian Subregion two species have been recorded (both from New Caledonia).

The genus is almost completely missing in the Oriental Region. Apart from the above mentioned cosmopolitan species — *T. granarium* Ev. — only two species — *T. koningsbergeri* PIC from Java and *T. unifasciatum* PIC from India are known from this region. Both are described very inaccurately, and there is no information on the distribution of these species apart from certain data from the original descriptions. An examination of the types may yet recommend a transferring of these species to other genera. Similarly, the distribution of the genus *Trogoderma* DEJ. in the Ethiopian Region, from where two species are known — *T. seminigrum* PIC from Madagascar and *T. rufopictum* ARR. from Natal — presents certain difficulties. It is possible that a revision of the type specimens will change the generic position of these species. Thus the opinion that the species of *Trogoderma* DEJ. are foreign elements in the Oriental and Ethiopian Regions seems to have serious grounds.

The second region where the species of the genus in question abound, apart from the Australian, is undoubtedly the Neotropical Region. From the Patagono-Andean Subregion nine species and from the Guyano-Brazilian Subregion 15 species have been described. Three species occur in the Antillean Subregion.

In North America, too, a fair number of species can be listed: six have been recorded from Mexico, and 20 from the United States (one of them reaches Alaska — *T. sinistrum* FALL). Some of these species have even been introduced into South America, e. g. *T. angustum* (SOL.), one is cosmopolitan and one occurs all over the Holarctic.

Not many species of *Trogoderma* DEJ. are found in the Palearctic. Several had been introduced from the Neotropical Region, such as *T. angustum* (SOL.) or *T. megatomoides* REITT., and one — *T. granarium* EV. — is cosmopolitan. Two species show a very wide distribution — *T. glabrum* (HERBST) occurs in the whole Holarctic and *T. versicolor* (CREUTZ.) all over the Palearctic. Fairly widely distributed seems to be *T. variabile* BALL., which occurs from the Caucasus through Iran and Afghanistan to Central Asia and Sinkiang. From China five species, besides the mentioned one, are known and one species has been recorded from Japan. Apart from the above mentioned species, five species, limited in their distribution and very little known, occur in the south-western part of the Palearctic Region. They are *T. syriacum* DALLA TORRE, known only from Syria, *T. quinquefasciatum* JACQ. DU VAL, known from France and Germany (species most probably introduced from South America), *T. albonotatum* MULS. et REY, known only from the Pyrenees, *T. irroratum* REITT., recorded from Egypt, and *T. octaedron* PEYER., collected in Sahara. These species have hardly been studied in their morphology, and their descriptions indicate considerable similarity. An examination of new material, and of the types as well, will certainly shed some light on the systematic position of these species and may alter in a radical manner the picture of the distribution of the genus in South-Western Palearctic.

Paratrogoderma SCOTT

Monospecific genus, known only from the Seychelles in the Madagascar Subregion.

Reesa BEAL (map 11)

Monospecific genus widely distributed in the Nearctic Region.

Phradonomorphus PIC

Monospecific genus known only from Senegal.

Phradonoma JACQ. DU VAL (map 8)

A genus comprising 18 species distributed in the Ethiopian and Palearctic Regions. Two species occurring in Egypt and Syria have been placed in a separate subgenus — *Orbeola* MULS. et REY. They show a more primitive morphology than the remaining ones belonging to the subgenus *Phradonoma* s. str.

Seven species occur in the Ethiopian Region: two in the South-African Subregion, one in Congo (Kinshasa) and Rhodesia, one in Kenya and Tanganyika, one in Zanzibar, one in Yemen and one in the Madagascar Subregion.

In the Palearctic Region, aside from the mentioned species from the subgenus *Orbeola* MULS. et REY, nine species occur; they are widely distributed from the Maghreb

Countries in the west to Sinkiang and Mongolia in the east, two of them have a very wide distribution: *Ph. nobile* (REITT.) occurs in the whole Mediterranean Subregion; *Ph. villosulum* (DUFT.) in Central and South Europe and in Central Asia. Of the remaining species one occurs in the Maghreb Countries, two in Egypt, Syria and Libya, one in Turkmenia, one in Kirghizia, one in Sinkiang and one in Mongolia.

Globicornis LATR. (map 10)

The genus comprises 24 species of which five have been described from the Neotropical Region (three from Argentina, one from Bolivia and one from Brazil). It is very probable that an examination of the types would alter the generic position of these five species. Some characters in their descriptions as well as a very crowded distribution of the other species seem to suggest this. Unfortunately, apart from the short and superficial original descriptions (based on single specimens) there are no other data and more recent material is not available for study.

The remaining 19 species are distributed chiefly in the Mediterranean Subregion — in South Europe, Asia Minor, the Caucasus and in Central Asia, mainly in mountainous regions. Only a few species cross the mentioned boundaries. In the Palearctic part of Africa one species — *G. vaulogeri* PIC — has been recorded from Tunisia and another from both Tunisia and Algeria — the latter also from Sicily and France. Two species have been found in Spain, and eight in France. Corsica, Sardinia and Sicily are inhabited by six species and one species occurs in Crete. One species — *G. fasciata* (FAIRM.) — reaches beyond Sardinia and France to Belgium, Germany and Switzerland. Only three species are distributed in the whole of Europe, one of them has also been recorded from Kazakhstan, and two other from the Caucasus; of the latter one reaches western Siberia. Five species are known to occur in Italy and six in the Balkan Peninsula, but of these only two in Greece. One species has been taken in Asia Minor, and three in Syria (none occurs elsewhere). In the Caucasus five species are found of which two are endemic to that area.

Hirtomegatoma PIC (map 10)

Monospecific genus, probably eremic, known only from the Jordan.

Megatoma HERBST (map 12)

Holarctic genus comprising 20 species. Two species from the Australian Region — *M. tenuefasciata* REITT. recorded from Tasmania and *M. foveolata* LEPESME from Australia — probably do not belong, according to ARMSTRONG (1942:328; 1949:108), to the genus *Megatoma* HERBST, but to *Trogoderma* DEJ.

Three Palearctic species occurring in the whole of Europe and Siberia have been placed in the subgenus *Megatoma* s. str. Some of the eight Nearctic species grouped in a separate subgenus *Perimegatoma* HORN probably also belong to this subgenus; they

occur mainly in the West-American Subregion and north of it (British Columbia, Alberta, Alaska). Only one species — *M. cylindrica* (KIRBY) — crosses eastward the Rocky Mts. It is widely distributed over the whole of the United States. None of the said species has been recorded from the Sonoran Province — one more argument to regard it as a zoogeographically distinct area.

Three Palearctic and five Nearctic species belong to the subgenus *Pseudohadrotoma* KALIK. The Nearctic species occur in areas along the Pacific coast (Alaska, Yukon, British Columbia, Alberta, Oregon, Idaho, Montana, Nevada, California) and none crosses eastward the Rocky Mts. or reaches southwards the Sonoran Province. The ranges of the Palearctic species cover the territory of the U.S.S.R. from Central Asia to the Far East and north-eastern China and Mongolia.

The last subgenus *Caucasotoma* MROCK. comprises only one species inhabiting the Caucasus.

Labrocerus SHARP (map 4)

A fairly large genus (17 species). Endemite to the Hawaiian Subregion.

Eocerus SHARP

Monospecific endemite to the Hawaiian Subregion.

Argocerus SHARP

Genus of two species. Endemite of the Hawaiian Subregion. Considering the fairly small morphological differences between *Labrocerus* SHARP, *Eocerus* SHARP and *Argocerus* SHARP it seems justified to treat them all as subgenera of a single genus.

Dearthrus LEC. (map 12)

Holarctic genus with a disjunctive distribution, comprising two Nearctic and two Palearctic species. One Nearctic species occurs in the West-American Subregion (in Oregon and California), the other in the East-American Subregion (Georgia and Pennsylvania) also reaching the Alleghenies (Illinois and Indiana). The Palearctic species inhabit the eastern part of the Mediterranean Subregion.

Ditorines NORM. (map 10)

Monospecific genus, probably eremic, known only from Tunisia.

Caccoleptus SHARP

A small genus (until now only two species have been described) distributed in the Central-American Province of the Neotropical Region — in Panama and British Honduras.

Diontolobus SOL.

Chilean endemite, comprising two species.

Anthrenocerus ARR. (map 4)

A fairly large genus comprising 19 species, endemic to the Australian Region. The majority of them are distributed in Australia, one — *A. riguus* (ER.), is known only from Tasmania and one — *A. variabilis* (REITT.) — occurs in Australia and in New Caledonia as well. *A. australis* (HOPE) has the widest distribution having been recorded from the whole of Australia, Tasmania and New Zealand and also introduced to Europe (Great Britain — KLOET et HINCKS, 1945:184; Netherlands — MEEUSE, 1950:161).

Myrmeanthrenus ARMSTR.

Monospecific endemite confined to the Australian Subregion.

Adelaidia BLACKB.

Genus of two species. *A. rigua* BLACKB. occurs in Australia and *A. unicolor* MROCK. in the Philippine Islands. The distribution of these species may suggest a certain affinity between the Oriental and Australian Regions.

Orphinus MOTSCH. (map 5)

Large genus (54 species) occurring mainly in the Australian, Oriental and Ethiopian Regions. Beyond these areas only one species — *O. globulicornis* (REITT.) — has been recorded, namely from Cairo; another species — *O. fulvipes* GUÉR. — has a very wide distribution area covering Australia, the Pacific Islands, Java, Madagascar, South and Central America, Florida and Europe. The latter species, partly synanthropic, has been certainly introduced by man to Europe and North and South America. Three species known from Japan belong to the Oriental elements in that country; one of them was recently recorded from Korea.

Eleven species occur in the Australian, two in the Papuan and three in the Polynesian Subregion of which one is found in the Hawaii Is. as well. From Tasmania and New Zealand no species has been recorded.

The majority of the species in the Oriental Region are concentrated in the Malayan Subregion. In Java, Sumatra, Borneo and Celebes nine species occur, while five are distributed in the Philippines and Taiwan. Eight species have been recorded from the Indochinese Province of the Indo-Indochinese Subregion. Ten species are found in the Indian Province.

Ten species inhabit the Ethiopian Region, of which only three the Madagascar Sub-region. The remaining species are distributed in the East-African and West-African Sub-regions, from Nigeria to Kenya in the north and from Angola to Mozambique in the south. There are no representatives of *Orphinus* MORSCH. in the South-African Subregion.

The distribution areas of the particular species of the genus in question are not particularly extensive. The presence of *O. defectus* (WALK.) in Ceylon and Borneo may be of interest. Even more interesting is the distribution of *O. aethiops* ARR. recorded so far from Angola, South Nigeria, Mauritius, Sumatra and Taiwan. It is probable that it has been introduced there as a pest of stored products.

Curtophinus PIC

Monospecific genus recorded from China without any detailed place of collecting.

Ctesias STEPH. (map 12)

A small genus comprising six species, with very wide distribution covering three zoogeographical regions. The distribution areas of the particular species are, however, rather small, and not adjacent to one another. In the Palearctic Region four species occur — one in the whole of Europe and the remaining ones in Syria, Turkmenia and Afghanistan. From the Ethiopian Region only one species has been reported — *C. variegata* ARR. — described from Rhodesia and recorded later from the Congo (Kinshasa). The sixth species — *C. dusmae* BEAL — was recently described from California.

Tiresiomorpha PIC

Monospecific genus, known only from China (no detailed locality given).

Thaumaglossa REDT. (map 6)

A fairly large genus (24 species) inhabiting principally the Oriental and Ethiopian Regions. Only one species — *Th. nigricans* (MACL.) — is known from Australia and two from the United States: one, *Th. americana* (JAYNE), has been taken in Texas and the other, *Th. libochoras* BEAL, in Arizona.

Of the remaining species only one, partly synanthropic — *Th. rufocapillata* REDT. — has a very wide distribution covering the whole Oriental and part of the Palearctic Region: Japan and Korea. The species is also known from Zanzibar and from Europe where it has been introduced to Germany. Apart from the species mentioned, nine more are found in the Oriental Region. All concentrate in the eastern part of this region. Two of them are Far-Eastern: *Th. ovivora* (MATS. et YOK.) is found in Taiwan and Japan, and *Th. hilleri* REITT. in the Philippines and Japan. The remaining species inhabit the Indo-chinese and Philippine Provinces, and one even reaches Sumatra.

In the Ethiopian Region, aside of *Th. rufocapillata* REDT. recorded from Zanzibar, ten species occur. One occurs in South Nigeria, three in the Cameroons, two in Congo (Kinshasa), one in Zanzibar, two in Rhodesia, one in the South-African and one in the Madagascar Subregions.

Pseudothaumaglossa PIC

Monospecific Brazilian endemite.

Hemirhopalum SHARP (map 2)

Neotropical genus comprising 14 species. The majority occur in South America: one species is found in Chile, four in Peru, one in Bolivia and four species have been recorded from Brazil. One species has also been found in Nicaragua, two in Guatemala and two in Mexico.

Falsocryptorhopalum PIC

Monospecific Brazilian endemite.

Cryptorhopalum GUÉR. (map 2)

One of the largest genera of *Dermestidae* (150 species), distributed mainly in the Neotropical Region, but reaching northward to the Nearctic. Few distributional data concerning the particular species in Central and South America, and a lack of a monographic study of the genus, do not allow to speculate about its distribution on the American continent. The available morphological descriptions are also very incomplete and a thorough revision of the genus based on a reexamination of the type specimens may yet drastically reduce the number of species. Any serious study of the distribution of the particular species is of course impossible since corresponding data are very sparse and usually confined to one single locality (*locus typicus*). New material will certainly be welcome.

The genus *Cryptorhopalum* GUÉR. does not extend far southwards. From the Patagono-Andean Subregion — Chile and Uruguay no species of this genus has ever been recorded, and only two species are known from Argentina and two more from Paraguay. Numerous species have been described from the Guyano-Brazilian Subregion: 27 from Brazil, seven from Bolivia, 13 from Peru, one from Ecuador and — north of the equator — two from French Guiana and 15 species from Colombia. A most numerous group of species occurs in the Central-American Province: from Panama five species have been recorded, four from Costa Rica, five from Nicaragua, five from Honduras, twelve from Guatemala and 29 from Mexico.

From the United States 31 species are known. They occur mainly in the south — in the Sonoran Province. This may justify the separation of this Province from the rest of the Nearctic Region and its placing together with the Neotropical Region. Two of the mentioned species are common to Mexico, the remaining are known only from the

United States. Twelve species are known from Arizona, whereas ten have been recorded from California, eight from Texas, five from New Mexico and five from Utah. In the adjacent States (Oregon, Nevada, Colorado, Kansas, Missouri) at most one or two species occur. Two species have been reported from the southern States along the Atlantic coast. In the northern States no member of the genus has ever been taken.

Anthreninae LECONTE

Anthrenus SCHAEFF. (map 9)

One of the largest genera, comprising 83 species. The majority — 56 of them — occur in the Palearctic Region; two are cosmopolitan in their distribution. There, is however, a marked absence of members of the genus in the Neotropical Region (apart from introduced cosmopolitan species). In the Australian Region (aside from imported species) three are known from the Polynesian Subregion: two are Tahitian endemites and one — *A. oceanicus* FAUV., an Oriental species, — is found, in addition to its Oriental stations, in New Caledonia as well. All three belong to the subgenus *Anthrenus* s. str.

From the Nearctic Region ten species are known, but only four can be regarded as autochthonic elements, the remaining species having been most probably introduced from the Palearctic. Similarly, only few species have been listed from the Oriental Region — nine species altogether — of which two are cosmopolitan and one extends through the Palearctic to India. Eighteen species are known from the Ethiopian Region, of which 14 occur exclusively in that region.

The species of *Anthrenus* SCHAEFF. are grouped in eight subgenera. Unfortunately five species could not so far be given a subgeneric position.

In the Nearctic Region, leaving aside the introduced species, only two species of the subgenus *Anthrenus* s. str. are known. One — *A. acomanus* CAS. — occurs in New Mexico, the other — *A. parvus* CAS. — in Texas. Of the subgenus *Florilinus* MULS. et REY, two species occur also in the region in question, one of which — *A. caseyi* DALLA TORRE — was described from the State of New York, and the other — *A. castaneae* MELSH. — is broadly distributed over nearly the whole of the United States. As yet, however, an elaboration of the genus *Anthrenus* SCHAEFF. of the Nearctic is still wanting. Such a revision is now in preparation by the author, and it would be of importance to note that our knowledge concerning the number of species known from the Nearctic as well as their pattern of distribution may change substantially in the future.

Three species of the subgenus *Anthrenops* REITT. have been recorded from the Indian Province of the Oriental Region. One — *A. rufescens* PIC — occurs only in India, the second — *A. subclaviger* REITT. — is known from Yemen and Aden as well, and the third species — *A. coloratus* REITT. — has a very wide distribution in the Palearctic reaching even India. Of the subgenus *Anthrenodes* CHOB. only one species — *A. maculifer* REITT. — occurs in the region in question; it is widely distributed in the Indo-Indochinese Subregion. Two species of the subgenus *Nathrenus* CAS. are known from the Oriental Region — *A. albomaculatus* PIC occurs in Borneo and *A. subsetosus* ARR. in Burma. Finally of the

subgenus *Anthrenus* s. str. only one species — *A. oceanicus* FAUV. — has been recorded from the Oriental Region; it occurs in the Indo-Indochinese Subregion and in New Caledonia.

About a dozen of Ethiopian species are confined principally to the East-African Subregion. They belong to three subgenera. Of the subgenus *Anthrenops* REITT. three species occur in the area mentioned. The majority of the Ethiopian species of *Nathrenus* CAS. (four), however, occur in the South-African Subregion. A fifth species has been recorded from Tanganyika. Finally, three species of *Anthrenus* s. str. are found in the East-African Subregion. There are no members of *Anthrenus* SCHAEFF. known from the West-African and Madagascar Subregions.

The majority of species of *Anthrenus* SCHAEFF., as it has been already mentioned before, are Palearctic in distribution. They are members of all the subgenera.

The subgenus *Ranthenus* MROCKZK. comprises only one species known from Central Asia (eastern Kazakhstan). Of the subgenus *Helocerus* MULS. et REY three species are known, distributed almost over the whole of Europe; none goes beyond Europe. Six species belong to the subgenus *Solskinus* MROCKZK., of which one occurs in Morocco and the remaining ones in mountainous regions of Asia: *A. armeniacus* ZAJC. is known from Armenia, three species were taken in the Central Asiatic Soviet Republics and the last species — *A. sinensis* ARR. — occurs in North China. The three above mentioned subgenera have no members in other zoogeographical regions.

The subgenus *Florilinus* MULS. et REY, comprises two Nearctic and seven Palearctic species. Of the latter, one — *A. coreanus* MROCKZK. — recorded only from Korea, is morphologically related to the Nearctic *A. castaneae* MELSH. and to the eastern Palearctic *A. flavidus* SOLS., which may indicate certain links between the two regions. *A. flavidus* SOLS. occurs from the Caucasus through Central Asia to North China. The remaining species are distributed as follows: *A. museorum* (L.) in the whole Holarctic, *A. caucasicus* REITT. in the Caucasus and in Turkmenia, *A. sordidulus* REITT. in the Dodecanese Islands, Syria and Spain, *A. olgae* KAL. in Central Europe and, finally, *A. oberthueri* REITT. in the Pyrenean Peninsula, Corsica and Algeria.

The subgenus *Anthrenops* REITT. comprises eleven species. They do not cross in the Palearctic the limits of the Mediterranean Subregion. *A. coloratus* REITT. has the widest area of distribution, covering Sudan, the African coast of the Mediterranean, Greece, the southern zone of the European part of the U.S.S.R., Central Asia (Turkmenia, Afghanistan) and India. As mentioned before, three species occur in the east of the Ethiopian Region — in Kenya, Tanganyika, Ethiopia and Socotra. They do not reach Aden and Yemen which link their range with that inhabited by West-Oriental species. In the Maghreb Countries two species occur, of which one — *A. minor* WOLL. — is also known from the Canary Islands and Greece. The remaining species are known from Transcaucasia, Iran, the Turkmen S.S.R. and the Tadzhik S.S.R.

The subgenus *Anthrenodes* CHOB. includes, aside from one Oriental species, two more, of which one occurs in Algeria and the other — *A. sarnicus* MROCKZK. is known only from the Island of Guernsey and has been, most probably, introduced there.

The greatest number of species — 45 — are grouped in two subgenera: *Nathrenus* CAS. and *Anthrenus* s. str. The former comprises two Oriental, five Ethiopian and eleven

Palaearctic species of which one has a cosmopolitan distribution and is known as a storage pest. The remaining Palaearctic species occur in the Mediterranean Subregion, chiefly in North Africa and on Mediterranean Islands (six species) and in the Balkan Peninsula (two species). One species — *A. molitor* AUBÉ — occurs everywhere in the Mediterranean Proper Province. The last species of the subgenus in question occurs in the Irano-Turanian Province.

The subgenus *Anthrenus* s. str. comprises two species already mentioned occurring in the southern part of the Nearctic Region, one Oriental, three Ethiopian and 18 Palaearctic species, of which one is cosmopolitan, two occur not only in the whole Palaearctic but in the whole Nearctic Region as well, where probably they have been introduced; of the last species one is also widely distributed in the Oriental Region. The remaining 15 species are known only from the Palaearctic Region, and, similarly as the species of the subgenus *Nathrenus* CAS., they occur mainly in the Mediterranean Proper Province of the Mediterranean Subregion. Beyond this province three species are known: *A. lindbergi* MROCZK. from Afghanistan, *A. rotundulus* REITT. from Transcaucasia and Turkmenia and the broadly distributed *A. picturatus* SOLS., occurring from the Caucasus through Transcaucasia, Iran, Afghanistan to Central Asia (all Asiatic Soviet Republics) and North China. Of the species distributed in the Mediterranean Proper Province only two are found over the whole of this area. Four species are known only from its eastern part (Balkans, Syria, Israel), five only from the coasts of Africa of which again three only from Maghreb Countries. One species occurs in the western part of the province in question (Pyrenean Peninsula, France, Balearic Islands, Maghreb Countries).

Neoanthrenus ARMSTR. (map 4)

The genus comprising six species is distributed in the Australian Subregion: five species have been recorded from Australia and one, *N. frater* (ARR.), from Tasmania.

Orphilinae LECONTE

Orphilus ER.

A small genus with a disjunctive distribution, comprising five recent and one Miocene species. Three species occur in the West-American Subregion of the Nearctic (British Columbia, Oregon, Idaho, California, Nevada, Arizona, New Mexico, Colorado) where also a fossil species is known (Colorado). The fourth species occurs in the East-American Subregion. The fifth — *O. niger* (Rossi) — inhabits the Mediterranean Subregion of the Palaearctic, reaching northwards to the European Province.

Trinodinae CASEY

Apsectus LECONTE (map 2)

A small genus (eight species) distributed in the Central-American Province of the Neotropical Region. Four species are known from Mexico, one from Panama, one from Guatemala. *A. araneorum* BEAL is known from the south of the United States: Arizona.

The last species — *A. hispidus* (MELSH.) — occurs in Texas, Pennsylvania, Maryland and has been generally recorded from the Central and South States; it constitutes a Neotropical element in the Nearctic fauna.

Trinodes DEJ. (map 7)

A broadly distributed genus, comprising 18 species, of which four occur in the Ethiopian Region, seven in the Palearctic and seven in the Oriental Region. In the Ethiopian Region *T. senegalensis* PIC occurs in Senegal, *T. rufotestaceus* PIC in Congo (Kinshasa), *T. albohirsutus* KAL. in Tanganyika and *T. villosus* BOH. in Natal. In the Palearctic Region two species only are known from Egypt, one from Algeria and one has a wide distribution covering the whole of Europe, the Maghreb Countries, the Caucasus, Asia Minor and Turkmenia. Further three species occur in the eastern part of the Palearctic — in the Japano-Central Chinese Province. In the Oriental Region one species has been described from Taiwan, one from the Philippine Islands, one from Vietnam, one from Borneo and three from the Indian Province: one from India and two from Ceylon.

Evorinea BEAL

Small Oriental genus comprising four species recorded from Ceylon, India (Bombay), Vietnam, Sumatra, the Philippines, the Mariana and the Caroline Islands.

Hexanodes BLAIR

Monospecific endemite of the New Zealand Subregion of the Australian Region.

Thylocardiadinae MROCKZK.

Thylocardia MOTSCH.

Monospecific Holarctic genus recorded from Canada, the United States, the Hawaiian Islands, Egypt, Germany (Hamburg), North Russia (Leningrad), Finland, Trieste, Astrakhan, Transcaucasia, the Turkmen S.S.R., the Uzbek S.S.R. *Th. contractus* MOTSCH. is synanthropic, which explains its scattered "island" distribution. Its area of origin seems to be Central Asia.

Thaumaphrastinae ANDERSON

Thorictodes REITT.

A small genus (four species) with an interesting pattern of distribution. *Th. bennetti* JOHN is known from former British Guiana, *Th. dartevillei* JOHN from Congo (Kinshasa), *Th. erraticus* CHAMP. from India. The last species, *Th. heydeni* REITT., partly synanthropic, is very widely distributed — it occurs in the western part of the Mediterranean, Egypt, Sudan, India, Java, Mariane Is., Guadeloupe, Mexico and the United States.

III. A short Catalogue of all known Species

DERMESTIDAE LATREILLE, 1807 : 3

Type genus: *Dermestes* LINNAEUS, 1758 : 342.

Syn.: *Dermestini* LATREILLE, 1807:3.

Dermestidae: GYLLENHALL, 1808:145.

Dermestides: LEACH, 1815:94.

Dermestidae: LEACH, 1817:

Dermestites: NEWMAN, 1834:416.

Dermestins: CASTELNAU, 1840:32.

Dermestida: HEER, 1841 b:434.

Dermestae: REDTENBACHER, 1845 : 52.

Dermestoidae: AGASSIZ, 1846:120.

Dermestinae: LECONTE, 1883:141.

***Marioutinae* WINKLER, 1926 : 682**

Type genus: *Mariouta* PIC, 1898b : 73.

Genus: *Mariouta* PIC, 1898b : 73

Type species: *Mariouta letourneuxi* PIC, 1898b : 73.

List of species:

Mariouta letourneuxi PIC, 1898b : 73.

Egypt, Algeria,
Morocco

Syn.: *Mariouta letourneuri* ESCALERA, 1914:135 (nomen nudum).

Mariouta strangei REITTER, 1910 : 239.

Turkmen SSR

Mariouta letourneuri ESCALERA, 1914:135 (nomen nudum) — vide *Mariouta letourneuxi* PIC, 1898b: 73

***Egidyellinae* SEMENOV-TIAN-SHANSKIJ, 1916:414 stat. nov.**

Type genus: *Egidyella* REITTER, 1899c : 284.

Syn.: *Epidyellini* SEMENOV-TIAN-SHANSKIJ, 1914:15.

Type genus: *Epidyella* SEMENOV-TIAN-SHANSKIJ, 1914:15.

Egidyellini SEMENOV-TIAN-SHANSKIJ, 1916:414.

Genus: *Egidyella* REITTER, 1899c : 284

Type species: *Egidyella prophetea* REITTER, 1899c : 285.

Syn.: *Trichopleurias* SAHLBERG, 1903:35. — SEMENOV-TIAN-SHANSKIJ, 1914:15.

Type species: *Trichopleurias deserti* SAHLBERG, 1903:36.

Epidyella SEMENOV-TIAN-SHANSKIJ, 1914:15.

Type species: *Egidyella prophetea* REITTER, 1899c:285.

List of species:

Egidyella prophetea REITTER, 1899c : 285.

Syn.: *Trichopleurias deserti* SAHLBERG, 1903:36. — SEMENOV-TIAN-SHANSKIJ, 1914:15.

Uzbek SSR,
Turkmen SSR

Trichopleurias deserti SAHLBERG, 1903:36 — vide *Egidyella prophetea* REITTER, 1899c:285.

Dermestinae LATREILLE, 1807 : 3

Type genus: *Dermestes* LINNAEUS, 1758 : 342.

Syn.: *Dermestini* LATREILLE, 1807:3.

Dermestida: LEACH, 1815:94.

Dermestites: CASTELNAU, 1840:33.

Dermestinae: DALLA TORRE, 1911:39.

Genus: *Dermestes* LINNAEUS, 1758 : 342

Type species: *Dermestes lardarius* LINNAEUS, 1758 : 354.

Subgenus: *Dermestinus* ZHANTIEV, 1967:1352

Type species: *Dermestes murinus* LINNAEUS, 1758 : 356.

List of species:

Dermestes aurichalceus KÜSTER, 1846 : 50.

Syn.: *Dermestes holosericeus* DEJEAN, 1837:139, nomen nudum.

Dermestes mustelinus PERRIS, 1853:638. — REITTER, 1883:89.

Italy, Sardinia,
France,
Tunisia,
Morocco

Dermestes aurichalceus var. *holosericeus* TOURNIER, 1860 : 318. — PIC, 1951c : 8.

Syn.: *Dermestes holosericeus* TOURNIER, 1860:318.

Dermestes caninus GERMAR, 1824 : 84.

Syn.: *Dermestes tessellatus* HERBST, 1792:122 nec FABRICIUS, 1775:56. — GEMMINGER et HAROLD, 1868:913.

Dermestes nubilus SAY, 1832:3. — HENSHAW, 1885:54.

Dermestes murinus LECONTE, 1854:108. — HENSHAW, 1885:54.

Dermestes murinus var. *caninus*: JAYNE, 1882:350.

Piedmont

Kentucky,
Central and
North America

<i>Dermestes caninus</i> var. <i>compactus</i> CASEY, 1900 : 141.	Texas
<i>Dermestes caninus</i> var. <i>mannerheimi</i> LECONTE, 1854 : 107. — CASEY, 1900 : 141.	California, Texas, Mexico
Syn.: <i>Dermestes mannerheimii</i> LECONTE, 1854:107. <i>Dermestes marmoratus</i> MANNERHEIM, 1843:258 nec SAY, 1823:197. — LECONTE, 1854:107. <i>Dermestes marmoratus</i> var. <i>mannerheimi</i> : JAYNE, 1882:350.	
<i>Dermestes caninus</i> var. <i>nubipennis</i> CASEY, 1900 : 141.	Iowa to Florida, whole USA
Syn.: <i>Dermestes nubilus</i> LECONTE, 1854:107 nec SAY, 1832:3. — CASEY, 1900:141.	
<i>Dermestes carnivorus</i> FABRICIUS, 1775 : 55 (<i>carniforous</i> emend. by GOEZE, 1777 : 141).	Australia, New Zealand, nearly cosmopolitan
Syn.: <i>Dermestes versicolor</i> CASTELNAU, 1840:33. — ERICHSON, 1846:429. <i>Dermestes mucoreus</i> LECONTE, 1854:108. — REITTER, 1881:28. <i>Dermestes sobrinus</i> LECONTE, 1854:108. — BLACKWELDER, 1945:395. <i>Dermestes muscoreus</i> REITTER, 1881:28.	
<i>Dermestes carnivorus</i> var. <i>doemmlingi</i> MEIER, 1899 : 99.	Hamburg, Amsterdam
<i>Dermestes coarctatus</i> HAROLD, 1877 : 347.	Japan, Korea, East Siberia, North China
<i>Dermestes coronatus</i> STEVEN in SCHÖNHERR, 1808 : 85. Syn.: <i>Dermestes signaticollis</i> STURM, 1843:93, nomen nudum. — KÜSTER, 1850:15.	Kislari and Georgiefru (?), south-eastern part of Europa, Caucasus, Middle Asia, Afghanistan, Sinkiang
<i>Dermestes cylindricus</i> CASEY, 1916 : 181.	Massachusetts
<i>Dermestes diadema</i> MOTSCHULSKY, 1849 : 89. Syn.: <i>Dermestes normandi</i> KALIK, 1951:154. — syn. nov. ¹	Spain, Algeria, Tunisia

¹ I have examined the two syntypes of *Dermestes diadema* MOTSCHULSKY, 1849:89, labelled „Hisp. mer.”, preserved in the MOTSCHULSKY’s collection in the Museum of Zoology of the Lomonosov Uni-

<i>Dermestes dimidiatus</i> STEVEN in SCHÖNHERR, 1808 : 89.	Caucasus, Asiatic part of USSR, Mongolia, Northern China
Syn.: <i>Dermestes orientalis</i> KUSNEZOVA, 1928:20. — MROCKOWSKI, 1962:236.	
<i>Dermestes dimidiatus orientalis</i> : LEPESME, 1950:50. — MROCKOWSKI, 1962:236.	
<i>Dermestes dimidiatus</i> ab. <i>rosea</i> KUSNEZOVA, 1928 : 22. — MROCKOWSKI, 1962 : 236.	Kazakh SSR, Siberia, Transbaicalia, Mongolia
Syn.: <i>Dermestes orientalis</i> ab. <i>rosea</i> KUSNEZOVA, 1928:22.	
<i>Dermestes elegans</i> GEBLER, 1830 : 99.	Kazakh SSR, Uzbek SSR, Kirghiz SSR, Tadzhik SSR, Turkmen SSR
Syn.: <i>Dermestes elegans</i> SOLSKIJ, 1876:269. — HEYDEN, 1881:98.	
<i>Dermestes elegans</i> var. <i>nadezhdae</i> PLAVILSTSHIKOV, 1924 : 231.	Pamir
Syn.: <i>Dermestes elegans</i> ab. <i>nadeghdae</i> WINKLER, 1926:675.	
<i>Dermestes erichsoni</i> GANGLBauer, 1904b : 16.	Germany, Middle and South Europe, Caucasus, Siberia
Syn.: <i>Dermestes tessellatus</i> ERICHSON, 1846:432 nec FABRICIUS, 1775: 56. — GANGLBauer, 1904 b:16.	
<i>Dermestes fasciatus</i> LECONTE, 1854 : 107.	New Mexico, Colorado, Wyoming
<i>Dermestes fasciventris</i> REITTER, 1881 : 28.	“East Siberia” , Southeast part of European part of RSFSR, Kazakh SSR
<i>Dermestes frischii</i> KUGELANN, 1792 : 478.	“Prussia” : probably Poland, entire Holarctic, Ethiopian and Neotropical Regions
Syn.: <i>Dermestes vulpinus</i> HERBST, 1792:124 nec FABRICIUS, 1781:64. — ERICHSON, 1846:428.	
<i>Dermestes pollinctus</i> HOPE, 1834b:54. — ERICHSON, 1846:429.	
<i>Dermestes frischii</i> heyrovsckyi OBENBERGER, 1917 : 32.	Albania
<i>Dermestes frischii</i> var. <i>rufimembris</i> PIC, 1951c : 7.	France
<i>Dermestes frischii</i> var. <i>sternimaculatus</i> MARCU, 1957 : 531.	Romania

versity at Moscow as well as the Holotype and one Paratype of *Dermestes normandi* KALIK, 1951:154, both labelled „Algeria, Batna”, preserved in the Zoological Museum at Budapest. All examined specimens belong to one species.

Dermestes frischii ab. *uniformis* REY, 1889 : 59.

France,
Europe,
Caucasus,
Turkey

Dermestes fulvicollis REITTER, 1881 : 29.

Himalaya Mts.

Dermestes gyllenhali CASTELNAU, 1840 : 34.

Sweden,
Europe,
Caucasus,
Transcaucasia,
Iran,
Kazakh SSR,
Siberia

Syn.: *Dermestes tessellatus* GYLLENHAL, 1808:150 nec FABRICIUS, 1775:
56. — CASTELNAU, 1840:34.

Dermestes atomarius CRISTOFORI et JAN, 1832:29, nomen nudum.

Dermestes atomarius ERICHSON, 1846:431. — REITTER, 1891:168.

Poland

Dermestes gyllenhali helmi REITTER, 1889c : 279.

South America

Syn.: *Dermestes helmi* REITTER, 1889c:279.

Czechoslovakia,
Hungary,
Romania,
Yugoslavia,
Bulgaria,
Greece, Turkey,
Syria, Iraq

Dermestes atomarius var. *helmi*: KNIEPHOF, 1935:125.

Dermestes atomarius helmi: KALIK, 1951b:45.

Dermestes impressicollis REITTER, 1881 : 29.

Yugoslavia,
Crimea, Cze-
choslovakia,
Hungary,
Romania,
Bulgaria,
Greece, Tur-
key, Caucasus

Dermestes intermedius KALIK, 1951b : 41.

Sikkim

Dermestes kaszabi KALIK, 1950 : 64.

Germany,
entire
Palearctic
Region

Dermestes kuseongensis LEPESME, 1939 : 192. — LEPESME,
1950 : 48.

Syn.: *Dermestes maculatus* var. *kuseongensis* LEPESME, 1939:192.

Dermestes laniarius ILLIGER, 1801 : 85.

Syn.: *Dermestes murinus* HERBST, 1792:120 nec LINNAEUS, 1758:356. —
ERICHSON, 1846:434.

Dermestes macellarium HERBST, 1792:126 nec FABRICIUS, 1781:63. —
ILLIGER, 1801:85.

Dermestes affinis GYLLENHAL, 1808:148. — ERICHSON, 1846:434.

Dermestes catta DUFTSCHMID, 1825:34. — ERICHSON, 1846:434.

Dermestes liniarius STEPHENS, 1829:8.

Hungary,
Uzbek SSR,
Germany,
Bulgaria

Dermestes laniarius ab. *rulei* KALIK, 1951 : 155.

Dermestes laniarius ab. *subfulvicollis* KALIK, 1951 : 155.

South-east part of the European part of RSFSR, Kazakh SSR, Crimea, France

Dermestes leopardinus MULSANT et GODART, 1855 : 273.

Syn.: *Dermestes dimidiatus* KUSNEZOV, 1928:17 nec STEVEN in SCHÖN-HERR, 1808:89. — MROCKOWSKI, 1962:238.

Dermestes dimidiatus dimidiatus: LEPESME, 1950:50. — MROCKOWSKI, 1962:238.

Crimea, Yugoslavia, Bulgaria, Ukrainian SSR, south-east part of European part of RSFSR, Caucasus

Dermestes maculatus DEGEER, 1774 : 223.

Syn.: *Dermestes vulpinus* FABRICIUS, 1781:64. — GYLLENHAL, 1808:147.

Dermestes marginatus THUNBERG, 1781:7. — THUNBERG, 1787:38.

Dermestes australis DEJEAN, 1821:46, nomen nudum. — BLACK-BURN, 1903:160.

Dermestes elongatus HOPE, 1834 b:55. — ARROW, 1915:426.

Dermestes truncatus CASEY, 1916:180. — BEAL, 1961:116.

Dermestes maculatus vulpinus: KALIK, 1955:93.

Surinam, cosmopolitan

Dermestes maculatus var. *lupinus* MANNERHEIM, 1843 : 257.

Syn.: *Dermestes lateralis* STURM, 1826:130, nomen nudum. — STURM, 1843:93.

Dermestes lupinus CRISTOFORI et JAN, 1832:89 — nomen nudum.

Dermestes senegalensis CRISTOFORI et JAN, 1832:30, nomen nudum. — DEJEAN, 1837:139.

Dermestes lupinus MANNERHEIM, 1843:257.

Dermestes vulpinus var. *lupinus*: LECONTE, 1854:109.

Dermestes maculatus var. *cyprius* PIC, 1951c:7, syn. nov. — KALIK in litt.

California, cosmopolitan

Dermestes maculatus ab. *nigropubescens* KALIK, 1955 : 94.

Syn.: *Dermestes maculatus vulpinus* ab. *nigropubescens*: KALIK, 1955b:295.

Cameroons, Congo (Kinshasa)

Dermestes maculatus pakistanius HAVELK, 1951b : 143.

Czechoslovakia (introduced from Pakistan)

Dermestes maculatus ab. *rattulus* MULSANT et REY, 1868 : 30. — KOCHER, 1953 : 38.

Syn.: *Dermestes vulpinus* var. *rattulus* MULSANT et REY, 1868:30.

South parts of Europe, Corsica, Sardinia, Spain, Morocco

<i>Dermestes maculatus</i> ab. <i>senex</i> GERMAR, 1824 : 84. — KALIK, 1948 : 5.	Yugoslavia, whole Palearctic and Ethiopian Regions
Syn.: <i>Dermestes senex</i> GERMAR, 1824:84. <i>Dermestes cinereus</i> REDTENBACHER, 1867:43, syn. nov. — KALIK in litt. <i>Dermestes vulpinus</i> var. <i>senex</i> : REITTER, 1906:377.	
<i>Dermestes maculatus</i> var. <i>sudanicus</i> GREDLER, 1878 : 506.	Sudan
Syn.: <i>Dermestes vulpinus</i> var. <i>sudanicus</i> GREDLER, 1878:506.	
<i>Dermestes madagascariensis</i> LEPESME, 1939 : 192.	Madagascar
<i>Dermestes marmoratus</i> SAY, 1823 : 197.	Missouri, Arkansas, USA, Cuba, Mexico, Canada
<i>Dermestes medialis</i> CASEY, 1900 : 141.	California
<i>Dermestes murinus</i> LINNAEUS, 1758 : 356.	Europe, entire Palearctic Region
Syn.: <i>Dermestes nebulosus</i> DEGEER, 1774:197. — FABRICIUS, 1781:64. <i>Dermestes marinus</i> FABRICIUS, 1787:35. — GMELIN, 1790:1595. <i>Dermestes catta</i> PANZER, 1789:10. — HERBST, 1792:123. <i>Dermestes roseiventris</i> DEJEAN, 1821:46, nomen nudum. — DEJEAN, 1837:139. <i>Dermestes roseiventris</i> CASTELNAU, 1840:34. — ERICHSON, 1846: 429. <i>Dermestes murinus</i> var. <i>roseiventris</i> : HEER, 1841:25.	
<i>Dermestes murinus auriceps</i> REITTER, 1905 : 244. — MROCKOWSKI, 1966 d : 438.	East Siberia, Korea, Manchuria, Primorskiy Kray, env. of Vladivostok
Syn.: <i>Dermestes murinus</i> subvar. <i>auriceps</i> REITTER, 1905:244.	
<i>Dermestes mustelinus</i> ERICHSON, 1846 : 433.	Germany, France, Spain
Syn.: <i>Dermestes tessellatus</i> OLIVIER, 1790:No. 9 p. 9 nec FABRICIUS, 1775:56. — ERICHSON, 1846:433. <i>Dermestes murinus</i> DEJEAN, 1821:46, nomen nudum. — ERICHSON, 1846:433.	
<i>Dermestes mustelinus</i> ab. <i>albomaculatus</i> LEPESME, 1950 : 58.	without terra typica
<i>Dermestes pardalis</i> BILLBERG in SCHÖNHERR, 1808 : 90.	Europe, Portugal, Spain, France, Morocco, Tunisia, Algeria
Syn.: <i>Dermestes thoracicus</i> DEJEAN, 1821:46, nomen nudum. — GENÉ, 1836:22. <i>Dermestes thoracicus</i> GENÉ, 1836:22. — GENÉ, 1836:22.	

Dermestes planus MROCKOWSKI, 1960b : 210. — ZHANTIEV,
1963b : 1053.

Syn.: *Dermestes frischii planus* MROCKOWSKI, 1960b:210.

Dermestes rattus LECONTE, 1854 : 108.

Syn.: *Dermestes murinus* var. *rattus*: JAYNE, 1882:351.

Turkmen SSR

Dermestes roei HOPE, 1834b : 55.

California,
Middle and
Western States
of USA

Dermestes sardous KÜSTER, 1846 : 49.

Egypt

Sardinia, whole
Mediterranea

Dermestes sardous var. *fulvofasciatus* GANGLBAUER, 1904b :
18.

Egypt

Syn.: *Dermestes sardous* var. *striatellus* REITTER, 1908:45, syn.
nov. — KALIK in litt.

Dermestes sibiricus ERICHSON, 1846 : 427.

Siberia, south-
eastern border
of European
part of RSFSR,
Turkey,
Middle Asia,
North China,
Mongolia,
Primorskiy
Kray

Syn.: *Dermestes frischii* var. *sibiricus*: REITTER, 1906:377.
Dermestes frischii sibiricus: KUZNECOVA, 1933:237.
Dermestes sibiricus ZHANTIEV, 1960:1629.

Dermestes solskyi DALLA TORRE, 1911 : 49.

Iran

Syn.: *Dermestes ratus* SOLSKII, 1876:269 nec *D. rattus* LECONTE, 1854:
108. — DALLA TORRE, 1911:49.

Dermestes szekessyi KALIK, 1950 : 61.

Hungary,
Yugoslavia,
Poland,
Germany,
France,
Austria,
Denmark,
Sweden,
Norway, Byelo-
russian SSR,
Ukrainian SSR,
Bulgaria,
Novosibirsk
Oblast,
Kazakh SSR

Dermestes talpinus MANNERHEIM, 1843 : 257.

Syn.: *Dermestes catta* var. *talpinus* DEJEAN, 1837:139, nomen nudum. — MANNERHEIM, 1843:257.

Dermestes undulatus talpinus: LEPESME, 1950:46.

Dermestes tessellatocollis MOTSCHULSKY, 1860 : 124.

Syn.: *Dermestes tessellaticollis* MOTSCHULSKY, 1859:491, nomen nudum.

Dermestes tessellatocollis var. *kaliki* HAVELKA, 1957 : 106.

Dermestes tessellatocollis pici MROCKOWSKI, 1963 : 377.

Syn.: *Dermestes tessellatocollis* ab. *pici* LEPESME, 1950:53.¹

Dermestes tristis FALL, 1897 : 237.

California,
nearly whole
USA, Canada

East Siberia,
Korea, Japan,
China

China

India

Dermestes undulatus BRAHM, 1790 : 114.

Syn.: *Dermestes murinus* OLIVIER, 1790:No. 9 p. 9 nec LINNAEUS, 1758:356. — ILLIGER, 1801:84.

Dermestes vulpecula HERBST, 1792:123. — ILLIGER, 1801:84.

Dermestes tessellatus ILLIGER, 1801:84 nec FABRICIUS, 1775:56. — ILLIGER, 1801:84.

Dermestes tessellatus var. *vulpecula*: DEJEAN, 1821:46.

Dermestes variegatus BRULLÉ, 1832:163. — REITTER, 1881b:73.

Dermestes dissector KIRBY, 1837:115. — ARROW, 1915:426.

California,
Santa Rosa I.,
British
Columbia,
Washington,
Idaho,
Oregon

Germany,
entire
Holarctic
Region

Dermestes undulatus ab. *nigroclava* KALIK, 1946b : 134.

Dermestes undulatus ab. *roseipennis* LEPESME, 1950 : 46.

Dermestes unicolor LEPESME, 1950 : 44.

Czechoslovakia

France

Antilles,
Baja California

Subgenus: *Dermestes* s. str.

List of species:

Dermestes ater DEGEER, 1774 : 223.

Surinam,
cosmopolitan

Syn.: *Dermestes cadaverinus* FABRICIUS, 1775:55. — LEPESME, 1939:192.

Dermestes piceus THUNBERG, 1781:8. — SCHÖNHERR, 1808:90.

Dermestes felinus FABRICIUS, 1787:34. — SCHÖNHERR, 1808:90.

Dermestes domesticus GERMAR, 1824:83. — DOHRN, 1858:42.

Dermestes cadaverinus ab. *domesticus*: ERICHSON, 1846:430.

Dermestes hispidulus MONTROUZIER, 1860:264. — FAUVEL, 1903:
334.

¹ See Art. 10b of International Code of Zoological Nomenclature.

<i>Dermestes chinensis</i> MOTSCHULSKY, 1866:168. — REITTER, 1891: 169.	
<i>Dermestes subcostatus</i> MURRAY, 1867:22. — ARROW, 1915:426.	
<i>Dermestes noxius</i> MULSANT et REY, 1868:55. — HEYDEN, 1879: 336.	
<i>Dermestes favarcqui</i> GODARD, 1883:383. — REITTER, 1891:169.	
<i>Dermestes ater</i> var. <i>domestica</i> : BLACKWELDER, 1945:395.	
<i>Dermestes rufoapicalis</i> PIC, 1951c:7, syn. nov. — KALIK in litt.	
<i>Dermestes ater</i> var. <i>subsulcatus</i> BALLION, 1878 : 277	Sinkiang, Tadzhik SSR, Kazakh SSR
Syn.: <i>Dermestes domesticus</i> var. <i>subsulcatus</i> BALLION, 1878:277.	
<i>Dermestes cadaverinus</i> var. <i>subsulcatus</i> : HEYDEN, 1881:98.	
<i>Dermestes bicolor</i> FABRICIUS, 1781 : 64.	Germany, entire Palearctic Region
Syn.: <i>Dermestes striatus</i> KOLENATI, 1846:58. — ERICHSON, 1847b:108.	
<i>Dermestes depressus</i> GEBLER, 1830 : 99.	South Siberia, Middle Siberia, Altai
Syn.: <i>Montandonia depressa</i> (GEBLER, 1830:99) HEYDEN, 1893:64.	
<i>Dermestes elongatissimus</i> PIC, 1916c : 15.	Ecuador
<i>Dermestes fuliginosus</i> ROSSI, 1792 : 17.	Italy, South and southern part of Middle Europa, Caucasus
Syn.: <i>Dermestes ater</i> DUFTSCHMID, 1825:32 nec DEGEER, 1774:223. — ERICHSON, 1846:436.	
<i>Dermestes ater</i> var. <i>fuliginosus</i> : REITTER, 1887b:45.	
<i>Dermestes gerstaeckeri</i> DALLA TORRE, 1911 : 44.	Tanganyika, Kenya
Syn.: <i>Dermestes subcostatus</i> GERSTAECKER, 1884:45 nec MURRAY, 1867: 22. — DALLA TORRE, 1911:44.	
<i>Dermestes haemorrhoidalis</i> KÜSTER, 1852 : 64.	France, Middle and South Europe
Syn.: <i>Dermestes gulo</i> MULSANT et GODART, 1855:275. — REITTER, 1881b:74.	
<i>Dermestes hirticollis</i> FABRICIUS, 1792 : 229.	North Africa, Spain, Portugal, France, Sardinia
<i>Dermestes hispanicus</i> KALIK, 1952 : 75.	Spain, Algeria
<i>Dermestes impressus</i> PIC, 1898 : 2.	Algeria
<i>Dermestes lardarius</i> LINNAEUS, 1758 : 354.	Europe, cosmopolitan
Syn.: <i>Dermestes luganensis</i> STIERLIN, 1902:425. — REITTER, 1906:377.	
<i>Dermestes lardarius</i> var. <i>atrobasisalis</i> PIC, 1951c : 7.	Sunzel (?)
<i>Dermestes lardarius</i> var. <i>conjunctus</i> REITTER, 1887c : 173.	Hudson Bay

<i>Dermestes lardarius</i> var. <i>proximus</i> REITTER, 1887c : 173.	East Siberia
<i>Dermestes leechi</i> KALIK, 1952b : 45.	India, Turkmen SSR
<i>Dermestes maximus</i> PIC, 1915b : 2.	Congo (Kinshasa)
<i>Dermestes mormonalis</i> CASEY, 1924 : 185.	Utah
<i>Dermestes nidum</i> ARROW, 1915 : 426.	Georgia, south- eastern USA, Korea, Manchuria, Primorskiy Kray
Syn.: <i>Dermestes elongatus</i> LECONTE, 1854:109 nec LINNAEUS, 1781: 141. — ARROW, 1915:426.	
<i>Dermestes olivieri</i> LEPESME, 1939 : 192.	France, South and Middle Europe, Caucasus, Turkey, Turkmen SSR
Syn.: <i>Dermestes ater</i> OLIVIER, 1790: no. 9 p. 9 nec DEGEER, 1774:223. — LEPESME, 1939:192.	
<i>Dermestes palmi</i> SJÖBERG, 1950 : 162.	Sweden, Finland
<i>Dermestes peruvianus</i> CASTELNAU, 1840 : 33.	Peru, Argentina, Bolivia, Chile, Mexico, USA, Europe (introduced)
Syn.: <i>Dermestes peruanus</i> ERICHSON, 1847:95. <i>Dermestes oblongus</i> SOLIER, 1849:365. — LEPESME, 1939:192. <i>Dermestes rufofuscus</i> SOLIER, 1849:365. — LEPESME, 1939:192. <i>Dermestes angustus</i> CASEY, 1900:143. — BLACKWELDER, 1945:396. <i>Dermestes angustatus</i> SCHAEFFER, 1931:174. — SCHAEFFER, 1931:174.	
<i>Dermestes pulcher</i> LECONTE, 1854 : 108.	Philadelphia, USA
<i>Dermestes reductus</i> KALIK, 1952b : 43.	Canada
Syn.: <i>Dermestes lardarius</i> ab. <i>reductus</i> LEPESME, 1950:64.	
<i>Dermestes roubali</i> (KALIK, 1951 : 152) ZHANTIEV, 1967 : 1350.	Altai
Syn.: <i>Montandonia roubali</i> KALIK, 1951:152.	
<i>Dermestes semistriatus</i> BOHEMAN, 1851 : 583.	Mozambique
<i>Dermestes signatus</i> LECONTE, 1874 : 50.	Northern and Western America
Syn.: <i>Dermestes lardarius</i> var. <i>signatus</i> : JAYNE, 1882:353.	
<i>Dermestes subaenescens</i> PIC, 1943 : 15.	Brazil
<i>Dermestes vorax</i> MOTSCHULSKY, 1860 : 123.	East Siberia, North China, Sakhalin, Korea, Japan
Syn.: <i>Dermestes vorax</i> MOTSCHULSKY, 1859:491, nomen nudum. <i>Dermestes lardarius</i> var. <i>vorax</i> : REITTER, 1879:212. <i>Dermestes lardarius</i> <i>vorax</i> : KUZNECOVA, 1933:238.	

<i>Dermestes vorax</i> var. <i>albofasciatus</i> MATSUMURA et YOKO-YAMA, 1928b : 51.	Korea
<i>Dermestes wittei</i> KALIK, 1955 : 95.	Congo (Kinshasa)
Species incertae sedis	
<i>Dermestes impressipennis</i> PIC, 1942b : 9.	Madagascar
† <i>Dermestes larvalis</i> COCKERELL, 1917 : 43.	Burmese amber
† <i>Dermestes pauper</i> HEER, 1847 : 43.	Yugoslavia (Tertiary period)
† <i>Dermestes tertarius</i> WICKHAM, 1912 : 14.	Colorado (Miocene)

- Dermestes abietis* PAYKULL, 1798:291 — non Dermestidae, recte Cryptophagidae: *Micrambe abietis* (PAYKULL, 1798).
- Dermestes absinthii* MARSHAM, 1802:77 — non Dermestidae, recte Coccinellidae, nomen dubium.
- Dermestes acuminatus* (KUGELLAN, 1794) PAYKULL, 1798:299 — non Dermestidae, recte Lathridiidae: *Lathridius lardarius* (DEGEER, 1775).
- Dermestes adstrictor* HERBST, 1792:140 — non Dermestidae, recte Throscidae: *Throscus dermestoides* (LINNAEUS, 1766).
- Dermestes aeneus* (FABRICIUS, 1792) MARSHAM, 1802:79 — non Dermestidae, recte Phalacridae: *Olibrus aeneus* (FABRICIUS, 1792).
- Dermestes aeneus* VILLERS, 1789:61 — non Dermestidae, familia?
- Dermestes aestivus* THUNBERG, 1787:39 — non Dermestidae, familia?
- Dermestes affinis* GYLLENHAL, 1808:148 — vide *Dermestes laniarius* ILLIGER, 1801:85.
- Dermestes affinis* STURM, 1843:93 — nomen nudum.
- Dermestes agaricinus* (LINNAEUS, 1758) SCOPOLI, 1763:16 — non Dermestidae, recte Scaphidiidae: *Scaphosoma agaricinum* (LINNAEUS, 1758).
- Dermestes angustatus* SCHAEFFER, 1931:174 — vide *Dermestes peruvianus* CASTELNAU, 1840:33.
- Dermestes angustus* CASEY, 1900:143 — vide *Dermestes peruvianus* CASTELNAU, 1840:33.
- Dermestes antennulatus* GOEZE, 1777:149 — non Dermestidae, familia?
- Dermestes apriarius* (LINNAEUS, 1758) SCHRANK, 1781:21 — non Dermestidae, recte Cleridae: *Trichodes apriarius* (LINNAEUS, 1758).
- Dermestes aquaticus* FOURCROY, 1785:21 — non Dermestidae, recte Hydrophilidae: *Helophorus aquaticus* (LINNAEUS, 1758).
- Dermestes armadellus* VILLERS, 1789b:216 — non Dermestidae, recte Clambidae: *Clambus armadillo* (DEGEER, 1774).
- Dermestes armadillo* PAYKULL, 1798:296 — non Dermestidae, recte Ptiliidae: *Acrotrichis atomaria* (DEGEER, 1774).
- Dermestes armadillus* DEGEER, 1774:218 — non Dermestidae, recte Clambidae, *Clambus armadillo* (DEGEER, 1774).
- Dermestes armedillus* MOTSCHULSKY, 1850:196 — non Dermestidae, recte Clambidae: *Clambus armadillo* (DEGEER, 1774).
- Dermestes armidilus* VILLERS, 1789:57 — non Dermestidae, recte Clambidae: *Clambus armadillo* (DEGEER, 1774).

- Dermestes assimilis* PILLER et MITTERPACHER, 1783:67 — non *Dermestidae*, familia?
- Dermestes ater* DUFTSCHMID, 1825:32 nec DEGEER, 1774:223 — vide *Dermestes fuliginosus* ROSSI, 1792:17.
- Dermestes ater* MELSHIMER, 1806:7 — nomen nudum.
- Dermestes ater* (HERBST, 1792:95) PANZER, 1795:96 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Dermestes ater* LEPECHIN, 1772:331 — non *Dermestidae*, familia?
- Dermestes ater* OLIVIER, 1790: no. 9 p. 9 nec DEGEER, 1774:223 — vide *Dermestes olivieri* LEPESME, 1939:192.
- Dermestes ater* PANZER, 1794:9 nec DEGEER, 1774:223 — non *Dermestidae*, recte *Cryptophagidae*: *Atomaria fuscata* (SCHÖNHERR, 1808).
- Dermestes ater* PAYKULL, 1798:298 — non *Dermestidae*, recte *Pythidae*: *Sphaeriestes ater* (PAYKULL, 1798).
- Dermestes ater* THUNBERG, 1784:4 nec DEGEER, 1774:223 — non *Dermestidae*, recte *Nitidulidae*: *Cylloides ater* (HERBST, 1792).
- Dermestes ater* var. *domestica*: BLACKWELDER, 1945:395 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes ater* var. *fuliginosus*: REITTER, 1887b:45 — vide *Dermestes fuliginosus* ROSSI, 1792:17.
- Dermestes atomarius* CRISTOFORI et JAN, 1832:29, nomen nudum — vide *Dermestes gyllenhali* CASTELNAU, 1840:34.
- Dermestes atomarius* DEGEER, 1774:219 — non *Dermestidae*, recte *Ptiliidae*: *Acrotrichis atomaria* (DEGEER, 1774).
- Dermestes atomarius* ERICHSON, 1846:431 — vide *Dermestes gyllenhali* CASTELNAU, 1840:34.
- Dermestes atomarius* (FABRICIUS, 1792) THUNBERG, 1794:78 — non *Dermestidae*, recte *Mycetophagidae*: *Mycetophagus atomarius* FABRICIUS, 1792.
- Dermestes atomarius helmi*: KALIK, 1951b:45 — vide *Dermestes gyllenhali helmi* REITTER, 1889c:279.
- Dermestes atomarius* var. *helmi*: KNIEPHOF, 1935:125 — vide *Dermestes gyllenhali helmi* REITTER, 1889c:279.
- Dermestes atricapillus* MARSHAM, 1802:72 — non *Dermestidae*, recte *Hydrophilidae*: *Cercyon nigriceps* (MARSHAM, 1802).
- Dermestes auriculatus* FOURCROY, 1785:20 — non *Dermestidae*, recte *Dryopidae*: *Dryops auriculatus* (FOURCROY, 1785).
- Dermestes auritus* GOEZE, 1777:148 — non *Dermestidae*, familia?
- Dermestes australis* DEJEAN, 1821:46, nomen nudum — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes avidus* CRISTOFORI et JAN, 1832:29 — nomen nudum.
- Dermestes bactridis* LINNAEUS, 1763:392 — non *Dermestidae*, recte *Bruchidae*: *Pachymerus bactris* (LINNAEUS, 1767).
- Dermestes bicinctus* THUNBERG, 1794:79 — non *Dermestidae*, recte *Mycetophagidae*: *Litargus connexus* (FOURCROY, 1785).
- Dermestes bidentatus* FABRICIUS, 1792:233 — non *Dermestidae*, recte *Cucujidae*: *Silvanus bidentatus* (FABRICIUS, 1792).
- Dermestes bidentatus* (HERBST, 1783) SCHNEIDER, 1785:5 — non *Dermestidae*, recte *Scolytidae*: *Pityogenes bidentatus* (HERBST, 1783).
- Dermestes bifasciatus* OLIVIER, 1790: no. 9 p. 13 — vide *Attagenus bifasciatus* (OLIVIER, 1790: no. 9 p. 13).
- Dermestes bifasciatus* ROSSI, 1794:79 nec OLIVIER, 1790: no. 9 p. 13 — vide *Attagenus bifasciatus* var. *rossi* GANGLBauer, 1904b:26.
- Dermestes bifasciatus* GOEZE, 1777:146 — non *Dermestidae*, familia?
- Dermestes bifasciatus* THUNBERG, 1781:6 — non *Dermestidae*, familia?
- Dermestes bimaculatus* PANZER, 1798:7 — non *Dermestidae*, recte *Cryptophagidae*: *Micrambe bimaculata* (PANZER, 1798).
- Dermestes bipunctatus* DEGEER, 1774:197 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Dermestes bipunctatus* FABRICIUS, 1792:231 — non *Dermestidae*, recte *Cucujidae*: *Psammoecus bipunctatus* (FABRICIUS, 1792).
- Dermestes bipunctatus* FOURCROY, 1785:18 — non *Dermestidae*, recte *Nitidulidae*: *Nitidula bipunctata* (LINNAEUS, 1758).
- Dermestes bipunctatus* SCHRANK, 1776:61 — non *Dermestidae*, recte *Anobiidae*: *Coelostethus pertinax* (LINNAEUS, 1758).

- Dermestes bipunctatus* VILLERS, 1789:55 — non Dermestidae, familia?
- Dermestes bipustulatus* FABRICIUS, 1798:72 — non Dermestidae, recte Nitidulidae: *Cateretes bipustulatus* (PAYKULL, 1798).
- Dermestes bipustulatus* PAYKULL, 1798:286 — non Dermestidae, recte Nitidulidae, *Cateretes bipustulatus* (PAYKULL, 1798).
- Dermestes bipustulatus* THUNBERG, 1781:6 — non Dermestidae, recte Erotylidae: *Dacne bipustulata* (THUNBERG, 1781).
- Dermestes bipustulatus* THUNBERG, 1784:4 — non Dermestidae, recte Erotylidae: *Dacne bipustulata* (THUNBERG, 1781) apud GYLLENHAL, 1808:203.
- Dermestes bipustulatus* (FABRICIUS, 1781) VILLERS, 1789:54 — non Dermestidae, recte Hydrophilidae: *Sphaeridium bipustulatum* FABRICIUS, 1781.
- Dermestes boleti* SCOPOLI, 1763:17 — non Dermestidae, recte Cioidae: *Cis boleti* (SCOPOLI, 1763).
- Dermestes boleti* STRÖM, 1783:52 — non Dermestidae, familia?
- Dermestes boleti versicoloris* GOEZE, 1777:147 — non Dermestidae, recte Cioidae, *Cis boleti* (SCOPOLI, 1763).
- Dermestes boletophagus* MARSHAM, 1802:72 — non Dermestidae, recte Hydrophilidae: *Megasternum boletophagum* (MARSHAM, 1802).
- Dermestes brachypterus* FABRICIUS, 1792:235 — non Dermestidae, recte Staphylinidae: *Proteinus brachypterus* (FABRICIUS, 1792).
- Dermestes brachypterus* PANZER, 1793:10 nec FABRICIUS, 1792:235 — non Dermestidae, recte Staphylinidae: *Proteinus macropterus* (GYLLENHAL, 1810).
- Dermestes brassicae* SCOPOLI, 1763:17 — non Dermestidae, familia?
- Dermestes brevicornis* (HERBST, 1792:95) PANZER, 1795:96 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Dermestes brunneus* MARSHAM, 1802:78 — non Dermestidae, recte Ptiliidae: *Nossidium pilosellum* (MARSHAM, 1802).
- Dermestes cadaverinus* FABRICIUS, 1775:55 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes cadaverinus* ab. *domesticus*: ERICHSON, 1846:430 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes cadaverinus* var. *subsulcatus*: HEYDEN, 1881:98 — vide *Dermestes ater* var. *subsulcatus* BALLION, 1878:277.
- Dermestes caerulescens* SCOPOLI, 1763:18 — non Dermestidae, familia?
- Dermestes caeruleus* (THUNBERG, 1781) GMELIN, 1790:1594 — non Dermestidae, recte Cleridae: *Notostenus coeruleus* (THUNBERG, 1781).
- Dermestes cafer* GMELIN, 1790:1598 — non Dermestidae, familia?
- Dermestes calcographus* LINNAEUS, 1761:143 — non Dermestidae, recte Scolytidae: *Pityogenes chalcographus* (LINNAEUS, 1761) (nom. emend.).
- Dermestes calthae* SCOPOLI, 1763:18 — non Dermestidae, familia?
- Dermestes canus* STURM, 1843:93 — nomen nudum.
- Dermestes capensis* (THUNBERG, 1781) GMELIN, 1790:1598 — non Dermestidae, familia?
- Dermestes capucinus* GOEZE, 1777:146 — non Dermestidae, familia?
- Dermestes capucinus* LINNAEUS, 1758:355 — non Dermestidae, recte Bostrichidae: *Bostrychus capucinus* (LINNAEUS, 1758).
- Dermestes carbonarius* SCOPOLI, 1763:15 — non Dermestidae, familia?
- Dermestes cassiodoides* MARSHAM, 1802:77 — non Dermestidae, familia?
- Dermestes castaneus* MARSHAM, 1802:78 — non Dermestidae, recte Leiodidae: *Leiodes marshami* STEPHENS, 1829.
- Dermestes castaneus* (HERBST, 1793) PANZER, 1795:104 — non Dermestidae, familia?
- Dermestes castaneus* THUNBERG, 1794b:134 — nomen nudum.
- Dermestes cathae* VILLERS, 1789:55 — non Dermestidae, familia?
- Dermestes catta* DUFTSCHMID, 1825:34 — vide *Dermestes laniarius* ILLIGER, 1801:85.
- Dermestes catta* PANZER, 1789:10 — vide *Dermestes murinus* LINNAEUS, 1758:356.

- Dermestes catta* var. *talpinus* DEJEAN, 1837:139, nomen nudum — vide *Dermestes talpinus* MANNERHEIM, 1843:257.
- Dermestes cellaris* FABRICIUS, 1792:234 — non Dermestidae, recte Cryptophagidae: *Cryptophagus scanicus* (LINNAEUS, 1758).
- Dermestes cellaris* SCOPOLI, 1763:16 — non Dermestidae, recte Cryptophagidae: *Cryptophagus cellaris* (SCOPOLI, 1763).
- Dermestes chalcographus* GOEZE, 1777:131 — non Dermestidae, recte Scolytidae: *Pityogenes chalcographus* (LINNAEUS, 1761).
- Dermestes chinensis* FABRICIUS, 1775:58 — non Dermestidae, recte Anobiidae: genus *Xyletinus* LATR. apud MOTSCHULSKY, 1858b:146.
- Dermestes chinensis* MOTSCHULSKY, 1866:168 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes chrysomeloides* SCHRANK, 1781:31 — non Dermestidae, familia?
- Dermestes ciliatus* SCOPOLI, 1763:17 — non Dermestidae, familia?
- Dermestes cylindricornis* MELSHEIMER, 1806:7, nomen nudum — vide *Attagenus cylindricornis* SAY, 1825:185.
- Dermestes cinereus* REDTENBACHER, 1867:43 — vide *Dermestes maculatus* ab. *senex* GERMAR, 1824:84.
- Dermestes clavicornis* LINNAEUS, 1758:355 — non Dermestidae, recte Colydiidae: *Orthocerus clavicornis* (LINNAEUS, 1758).
- Dermestes coadunatus* MARSHAM, 1802:76 — non Dermestidae, recte Coccinellidae: *Rhizobius litura* (FABRICIUS, 1787).
- Dermestes coerulescens* VILLERS, 1789:60 — non Dermestidae, familia?
- Dermestes collaris* QUENSEL, 1790:8 — non Dermestidae, recte Serropalpidae: *Tetratoma fungorum* FABRICIUS, 1790.
- Dermestes colon* LINNAEUS, 1758:357 — non Dermestidae, recte Nitidulidae: *Cychramus variegatus* (HERBST, 1792).
- Dermestes complanatus* SCHRANK, 1798:427 — non Dermestidae, familia?
- Dermestes concinnus* MARSHAM, 1802:74 — non Dermestidae, recte Hydrophilidae: *Cercyon nigriceps* (MARSHAM, 1802).
- Dermestes consimilis* MARSHAM, 1802:75 — non Dermestidae, recte Phalacridae: *Stilbus apicalis* (MELSHIMER, 1846).
- Dermestes contractus* FOURCROY, 1785:19 — non Dermestidae, familia?
- Dermestes convexus* MARSHAM, 1802:73 — non Dermestidae, recte Clambidae: *Clambus armadillo* (DEGEER, 1774).
- Dermestes cordatus* THUNBERG, 1794:81 — non Dermestidae, familia?
- Dermestes costatus* GOEZE, 1777:148 — non Dermestidae, familia?
- Dermestes crenatus* FABRICIUS, 1798:72 — non Dermestidae, recte Cryptophagidae: *Pteryngium crenatum* (FABRICIUS, 1798).
- Dermestes crenatus* PANZER, 1795:102 — non Dermestidae, familia?
- Dermestes crenatus* VILLERS, 1789b:220 — non Dermestidae, familia?
- Dermestes cristatus* GOEZE, 1777:148 — non Dermestidae, familia?
- Dermestes cylindricornis* SCHRANK, 1785:315 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Dermestes dentatus* (HERBST, 1793) PANZER, 1795:104 — non Dermestidae, recte Cryptophagidae: *Cryptophagus dentatus* (HERBST, 1793).
- Dermestes dentatus* ROSSI, 1790:34 — non Dermestidae, recte Cleridae: *Enoplium serraticorne* (OLIVIER, 1790).
- Dermestes denticulatus* ROSSI, 1794:80 — non Dermestidae, familia?
- Dermestes destructor* CRISTOFORI et JAN, 1832:29 — nomen nudum.
- Dermestes dimidiatus* BOEBER, 1802:333 — nomen nudum.
- Dermestes dimidiatus* FOURCROY, 1785:25 — non Dermestidae, familia?
- Dermestes dimidiatus* KUSNEZOVA, 1928:17 nec STEVEN in SCHÖNHERR, 1808:89 — vide *Dermestes leopardinus* MULSANT et GODART, 1855:273.

- Dermestes dimidiatus dimidiatus*: LEPESME, 1950:50 — vide *Dermestes leopardinus* MULSANT et GODART, 1855:273.
- Dermestes dimidiatus orientalis*: LEPESME, 1950:50 — vide *Dermestes dimidiatus* STEVEN in SCHÖNHERR, 1808:89.
- Dermestes dissector* KIRBY, 1837:115 — vide *Dermestes undulatus* BRAHM, 1790:114.
- Dermestes domesticus* BRAHM, 1790:17 — non Dermestidae, familia?
- Dermestes domesticus* GERMAR, 1824:83 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes domesticus* LINNAEUS, 1758:356 — non Dermestidae, recte Scolytidae: *Xyloterus domesticus* (LINNAEUS, 1758).
- Dermestes domesticus* SCHRANK, 1781:30 — non Dermestidae, familia?
- Dermestes domesticus* var. *subsulcatus* BALLION, 1878:277 — vide *Dermestes ater* var. *subsulcatus* BALLION, 1878:277.
- Dermestes dubius* ROSSI, 1792:17 — non Dermestidae, recte Bostrychidae: *Psoa dubia* (ROSSI, 1792).
- Dermestes elegans* SOLSKIJ, 1876:269 — vide *Dermestes elegans* GEBLER, 1830:99.
- Dermestes elegans* ab. *nadezhdae* WINKLER, 1926:675 — vide *Dermestes elegans* var. *nadezhdae* PLAVIL-STSHIKOV, 1924:231.
- Dermestes elongatus* DEJEAN, 1837:139 — nomen nudum.
- Dermestes elongatus* HOPE, 1834b:55 — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes elongatus* LECONTE, 1854:109 nec LINNAEUS, 1761:141 — vide *Dermestes nidum* ARROW, 1915:426.
- Dermestes elongatus* LINNAEUS, 1761:141 — non Dermestidae, recte Temnochilidae: *Nemosoma elongatum* (LINNAEUS, 1761).
- Dermestes elongatus* (FABRICIUS, 1787) VILLERS, 1789b:220 — non Dermestidae, familia?
- Dermestes emarginatus* GYLLENHAL, 1808:153 — vide *Globicornis marginata* (PAYKULL, 1798:280).
- Dermestes erythrocephalus* SCHRANK, 1798:426 — non Dermestidae, familia?
- Dermestes eustadius* LINNAEUS, 1758:357 — non Dermestidae, familia?
- Dermestes exilis* GMELIN, 1790:1600 — non Dermestidae, recte Scolytidae: *Xyloterus domesticus* (LINNAEUS, 1758).
- Dermestes faber* (THUNBERG, 1784) SCHÖNHERR, 1808:101 — non Dermestidae, recte Anobiidae: *Xestobium rufovillosum* (DEGEER, 1774).
- Dermestes fagineus* SCHRANK, 1798:424 — non Dermestidae, familia?
- Dermestes fasciatus* FORSTER, 1771:8 — non Dermestidae, familia?
- Dermestes fasciatus* FOURCROY, 1785:23 — non Dermestidae, recte Anobiidae: *Ochina ptinoides* (MARSHAM, 1802).
- Dermestes fasciatus* O. F. MÜLLER, 1764:4 — non Dermestidae, familia?
- Dermestes fasciatus* SCOPOLI, 1772:88 — non Dermestidae, familia?
- Dermestes fasciatus* THUNBERG, 1784:5 — non Dermestidae, recte Mycetophagidae: *Litargus connexus* (FOURCROY, 1785).
- Dermestes fasciatus* (THUNBERG, 1795:105) SCHÖNHERR, 1808:88 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
- Dermestes fascicularis* (HERBST, 1793) PANZER, 1795:101 — non Dermestidae, recte Ptilidae: *Acrotrichis fascicularis* (HERBST, 1793).
- Dermestes favarcqui* GODARD, 1883:383 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes felinus* FABRICIUS, 1787:34 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes fenestralis* LINNAEUS, 1758:356 — non Dermestidae, recte Lathridiidae: *Corticaria fenestralis* (LINNAEUS, 1758).
- Dermestes fenestralis* PAYKULL, 1798:297 — non Dermestidae, recte Lathridiidae: *Corticaria pubescens* (GYLLENHAL, 1827) apud GYLLENHAL, 1827:124.
- Dermestes fenestratus* FABRICIUS, 1801:318 — non Dermestidae, recte Lathridiidae: *Corticaria fenestralis* (LINNAEUS, 1758).
- Dermestes fenestratus* THUNBERG, 1784:3 — non Dermestidae, recte Heteroceridae: *Heterocerus fenestratus* (THUNBERG, 1784).

- Dermestes ferrugineus* (SCOPOLI, 1763) BRAHM, 1790:118 — non Dermestidae, recte Nitidulidae: *Soronia grisea* (LINNAEUS, 1758).
- Dermestes ferrugineus* LINNAEUS, 1761:145 — non Dermestidae, recte Nitidulidae: *Pityophagus ferrugineus* (LINNAEUS, 1761).
- Dermestes ferrugineus* MARSHAM, 1802:74 — non Dermestidae, recte Hydrophilidae: *Megasternum boleto-phagum* (MARSHAM, 1802).
- Dermestes ferrugineus* O. F. MÜLLER, 1764:3 — non Dermestidae, recte Nitidulidae: *Pityophagus ferrugineus* (LINNAEUS, 1761).
- Dermestes fimetarii* PANZER, 1795:103 — non Dermestidae, recte Cryptophagidae: *Atomaria fuscata* (SCHÖNHERR, 1808) apud SCHÖNHERR, 1808:100.
- Dermestes fimetarii* PAYKULL, 1798:293 — non Dermestidae, recte Cryptophagidae: *Atomaria atra* (HERBST, 1793).
- Dermestes fimetarii* FABRICIUS, 1792:231 — non Dermestidae, recte Cryptophagidae: *Atomaria fimetarii* (HERBST, 1793).
- Dermestes fimetarius* HERBST, 1783:21 — non Dermestidae, familia?
- Dermestes fimetarius* (FABRICIUS, 1775) VILLERS, 1789:55 — non Dermestidae, recte Phalacridae: *Phalaerus fimetarius* (FABRICIUS, 1775).
- Dermestes flavescens* FOURCROY, 1785:19 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
- Dermestes flavescens* SCHRANK, 1782:140 — non Dermestidae, familia?
- Dermestes flavigornis* SCHNEIDER, 1785:5 — nomen nudum.
- Dermestes flavus* MARSHAM, 1802:71 — non Dermestidae, recte Hydrophilidae: *Cercyon quisquilius* (LINNAEUS, 1761).
- Dermestes floricola* MELSHEIMER, 1806:7, nomen nudum — vide *Attagenus cylindricornis* var. *floricola* SAY, 1825:185.
- Dermestes formicarius* (LINNAEUS, 1758) SCHRANK, 1781:21 — non Dermestidae, recte Cleridae: *Thanasimus formicarius* (LINNAEUS, 1758).
- Dermestes formicaroides* (SCHRANK, 1776) SCHRANK, 1781:20 — non Dermestidae, recte Cleridae: *Clerus mutillarius* FABRICIUS, 1775.
- Dermestes forniciatus* DEGEER, 1774:216 — non Dermestidae, recte Catopidae: *Catops nigricans* (SPENCE, 1815) apud GYLLENHAL, 1827:314.
- Dermestes frischii planus* MROCKOWSKI, 1960b:210 — vide *Dermestes planus* MROCKOWSKI, 1960b:210.
- Dermestes frischii sibiricus*: KUZNCOVA, 1933:237 — vide *Dermestes sibiricus* ERICHSON, 1846:427.
- Dermestes frischii* var. *sibiricus*: REITTER, 1906:377 — vide *Dermestes sibiricus* ERICHSON, 1846:427.
- Dermestes frontalis* FABRICIUS, 1798:72 — non Dermestidae, familia?
- Dermestes fulvipes* FOURCROY, 1785:22 — non Dermestidae, recte Nitidulidae: *Nitidula rufipes* (LINNAEUS, 1767).
- Dermestes fulvipes* GOEZE, 1777:149 — non Dermestidae, familia?
- Dermestes fumatus* FABRICIUS, 1775:57 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
- Dermestes fumatus* LAICHARTING, 1781:63 — non Dermestidae, recte Byturidae: *Byturus tomentosus* (FABRICIUS, 1775).
- Dermestes fumatus* LINNAEUS, 1767:564 — non Dermestidae, recte Mycetophagidae: *Typhaea stercorea* (LINNAEUS, 1758).
- Dermestes fungorum* PANZER, 1797:H. 39 p. 14 — non Dermestidae, recte Cryptophagidae: *Cryptophagus lycoperdi* (SCOPOLI, 1763).
- Dermestes fuscipes* (HERBST, 1792) PANZER, 1795:99 — non Dermestidae, familia?
- Dermestes fuscus* FABRICIUS, 1787:35 — non Dermestidae, recte Cioidae: *Cis boleti* (SCOPOLI, 1763) apud SCHAUM, 1847:55.
- Dermestes gagates* GOEZE, 1777:149 — non Dermestidae, familia?
- Dermestes gagatinus* FOURCROY, 1785:21 — non Dermestidae, familia?

- Dermestes gibbosus* (HERBST, 1783) PANZER, 1795:100 — non Dermestidae, recte Lathiriidae: *Melanophthalma gibbosa* (HERBST, 1783).
- Dermestes gigantinus* SCHÖNHERR, 1806:101 — non Dermestidae, familia?
- Dermestes glaber* O. F. MÜLLER, 1776:56 — non Dermestidae, familia?
- Dermestes glaber* (HERBST, 1793) PANZER, 1795:101 — non Dermestidae, familia?
- Dermestes glaber* SAHLBERG, 1819:50 — vide *Megatoma undata* (LINNAEUS, 1758:355).
- Dermestes glaberrimus* PAYKULL, 1798:287 — non Dermestidae, recte Nitidulidae: *Cateretes pedicularius* (LINNAEUS, 1758).
- Dermestes gleditsiae* LINNAEUS, 1763:392 — non Dermestidae, recte Bruchidae: *Pachymerus gleditsiae* (LINNAEUS, 1763).
- Dermestes globulus* PAYKULL, 1798:295 — non Dermestidae, recte Cryptophagidae: *Ephistemus globulus* (PAYKULL, 1798).
- Dermestes graphicus* SCHRANK, 1796:160 — non Dermestidae, recte Nitidulidae: *Cryptarcha strigata* (FABRICIUS, 1787).
- Dermestes gravidus* (ILLIGER, 1798) PAYKULL, 1800:448 — non Dermestidae, recte Nitidulidae: *Brachypteron pulicarius* (LINNAEUS, 1758).
- Dermestes gulo* MULSANT et GODART, 1855:275 — vide *Dermestes haemorrhoidalis* KÜSTER, 1852:64.
- Dermestes gyrioides* MARSHAM, 1802:77 — non Dermestidae, recte Cryptophagidae: *Ephistemus globulus* (PAYKULL, 1798).
- Dermestes haemorrhous* SCHRANK, 1781:26 — non Dermestidae, recte Hydrophilidae: *Sphaeridium bipustulatum* FABRICIUS, 1781.
- Dermestes hamatus* (FABRICIUS, 1787) VILLERS, 1789b:218 — non Dermestidae, recte Bostrichidae: *Amphicerus hamatus* (FABRICIUS, 1787).
- Dermestes helmi* REITTER, 1889c:279 — vide *Dermestes gyllenhalii helmi* REITTER, 1889c:279.
- Dermestes hemipterus* LINNAEUS, 1758:358 — non Dermestidae, recte Nitidulidae: *Carpophilus hemipterus* (LINNAEUS, 1758).
- Dermestes hemipterus* PANZER, 1789:11 — non Dermestidae, familia?
- Dermestes hirtus* LINNAEUS, 1767:563 — non Dermestidae, recte Dasytidae: *Enicopus pilosus* (SCOPOLI, 1763).
- Dermestes hispidulus* MONTROUZIER, 1860:264 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes holosericeus* DEJEAN, 1837:139, nomen nudum — vide *Dermestes aurichalceus* KÜSTER, 1846: 50
- Dermestes holosericeus* SAHLBERG, 1819:52 — vide *Attagenus obtusus* (GYLLENHAL in SCHÖNHERR, 1808:88).
- Dermestes holosericeus* TOURNIER, 1860:318 — vide *Dermestes aurichalceus* KÜSTER, 1846:50.
- Dermestes hortensis* FOURCROY, 1785:21 — non Dermestidae, recte Nitidulidae: *Glischrochilus hortensis* (FOURCROY, 1785).
- Dermestes humeralis* MARSHAM, 1802:67 — non Dermestidae, recte Erotylidae: *Tritoma bipustulata* FABRICIUS, 1775.
- Dermestes hypomelanus* MARSHAM, 1802:77 — non Dermestidae, recte Coccinellidae: *Rhizobius litura* (FABRICIUS, 1787).
- Dermestes indicus* GMELIN, 1790:1600 — non Dermestidae, recte Bostrichidae: *Xylopsocus capucinus* (FABRICIUS, 1781).
- Dermestes insulanus* SCHNEIDER, 1785:5 — nomen nudum.
- Dermestes interruptus* THUNBERG, 1781:6 — vide *Trogoderma interrupta* (THUNBERG, 1781:6).
- Dermestes inustus* MARSHAM, 1802:76 — non Dermestidae, familia?
- Dermestes jesuita* (FABRICIUS, 1775) SCHNEIDER, 1785:4 — non Dermestidae, recte Bostrichidae: *Bostrychopsis jesuita* (FABRICIUS, 1775).
- Dermestes jesuita* O. F. MÜLLER, 1776:57 — non Dermestidae, familia?
- Dermestes laevis* MARSHAM, 1802:73 — non Dermestidae, recte Hydrophilidae: *Cercyon nigriceps* (MARSHAM, 1802:72).
- Dermestes lardarius* ab. *reductus* LEPESME, 1950:64 — vide *Dermestes reductus* KALIK, 1952b:43.
- Dermestes lardarius* var. *signatus*: JAYNE, 1882:353 — vide *Dermestes signatus* LeCONTE, 1874:50.

- Dermestes lardarius* var. *vorax*: REITTER, 1879:212 — vide *Dermestes vorax* MOTSCHULSKY, 1860:123.
- Dermestes lardarius* *vorax*: KUZNCOVA, 1933:238 — vide *Dermestes vorax* MOTSCHULSKY, 1860: 123.
- Dermestes lateralis* MARSHAM, 1802:69 — non Dermestidae, recte Hydrophilidae: *Cercyon lateralis* (MARSHAM, 1802).
- Dermestes lateralis* STURM, 1826: 130, nomen nudum — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843:257.
- Dermestes latissimus* BIELZ, 1850:180 — vide *Montandonia latissima* (BIELZ, 1850:180).
- Dermestes lectularius* GOEZE, 1777:147 — nomen dubium.
- Dermestes lepturoides* SCOPOLI, 1763:17 — non Dermestidae, recte Nitidulidae: *Brachypterus pulicarius* (LINNAEUS, 1758).
- Dermestes lepturoides* SCOPOLI, 1772:87 nec SCOPOLI, 1763:17 — non Dermestidae, familia?
- Dermestes lepturoides rufus* GOEZE, 1777:145 — non Dermestidae, familia?
- Dermestes leucophilus* SCHRANK, 1781:28 — non Dermestidae, recte Nitidulidae: *Nitidula rufipes* (LINNAEUS, 1767).
- Dermestes ligniperda* (FABRICIUS, 1787) VILLERS, 1789b:221 — non Dermestidae, recte Scolytidae: *Hyllus ligniperda* (FABRICIUS, 1787).
- Dermestes lili* MOLL, 1784:175 — non Dermestidae, familia?
- Dermestes limbatus* FABRICIUS, 1781:66 — non Dermestidae, recte Colydiidae apud MOTSCHULSKY, 1858b: 146.
- Dermestes limbatus* (HERBST, 1782) SCHNEIDER, 1785:5 — non Dermestidae, recte Scolytidae: *Xyloterus domesticus* (LINNAEUS, 1758).
- Dermestes linearis* GOEZE, 1777:148 — non Dermestidae, familia?
- Dermestes linearis* PAYKULL, 1798:302 — non Dermestidae, recte Lathridiidae: *Corticaria linearis* (PAYKULL, 1798).
- Dermestes linearis* ROSSI, 1788:129 — nomen nudum.
- Dermestes linearis* THUNBERG, 1784:4 — non Dermestidae, recte Bostrichidae: *Lyctus linearis* (GOEZE, 1777).
- Dermestes liniarius* STEPHENS, 1829:8 — vide *Dermestes laniarius* ILLIGER, 1801:85.
- Dermestes listerianus* FOURCROY, 1785:17 — non Dermestidae, recte Silphidae: *Nicrophorus germanicus* (LINNAEUS, 1758).
- Dermestes longicornis* FOURCROY, 1785:22 — non Dermestidae, familia?
- Dermestes longicornis* HERBST, 1783:23 — non Dermestidae, recte Lathridiidae: *Corticaria pubescens* (GYLLENHAL, 1827).
- Dermestes longicornis* (HERBST, 1793) PANZER, 1794:H. 23 p. 10 — non Dermestidae, recte Lathridiidae: *Corticaria longicornis* (HERBST, 1793).
- Dermestes longicornis* THUNBERG, 1794:81 — non Dermestidae, familia?
- Dermestes longipes* MOLL, 1784:176 — non Dermestidae, familia?
- Dermestes luganensis* STIERLIN, 1902:425 — vide *Dermestes lardarius* LINNAEUS, 1758:354.
- Dermestes lugubris* FOURCROY, 1785:22 — non Dermestidae, recte Hydrophilidae: *Cercyon lugubris* (FOURCROY, 1785).
- Dermestes lunata* FABRICIUS, 1787:378 — non Dermestidae, recte Erotylidae: *Diphyllus lunatus* (FABRICIUS, 1787).
- Dermestes lunatus* MARSHAM, 1802:66 — non Dermestidae, recte Hydrophilidae: *Sphaeridium lunatum* FABRICIUS, 1792.
- Dermestes lupinus* CRISTOFORI et JAN, 1832:89, nomen nudum — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843:257.
- Dermestes lupinus* MANNERHEIM, 1843:257 — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843: 257.
- Dermestes luteus* (FABRICIUS, 1787) MARSHAM, 1802:73 — non Dermestidae, recte Nitidulidae: *Cychramus luteus* (FABRICIUS, 1787).

- Dermestes lycoperdi* SCOPOLI, 1763:18 — non Dermestidae, recte Cryptophagidae: *Cryptophagus lycoperdi* (SCOPOLI, 1763).
- Dermestes lycopordi* FABRICIUS, 1792:231 — non Dermestidae, recte Cryptophagidae: *Cryptophagus lycopordi* (SCOPOLI, 1763).
- Dermestes macellarius* DUFTSCHMID, 1825:39 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Dermestes macellarius* FABRICIUS, 1781:63 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Dermestes macellarius* HERBST, 1792:126 nec FABRICIUS, 1781:63 — vide *Dermestes laniarius* ILLIGER, 1801:85.
- Dermestes maculatus* THUNBERG, 1781:7 — non Dermestidae, recte Mycetophagidae: *Mycetophagus* sp. apud KALIK in litt.
- Dermestes maculatus* var. *cyprius* PIC, 1951c:7 — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843:257.
- Dermestes maculatus* var. *kurseongensis* LEPESME, 1939:192 — vide *Dermestes kurseongensis* LEPESME, 1939:192.
- Dermestes maculatus* vulpinus: KALIK, 1955:93 — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes maculatus* vulpinus ab. *nigropubescens*: KALIK, 1955b:295 — vide *Dermestes maculatus* ab. *nigropubescens* KALIK, 1955:94.
- Dermestes mannerheimii* LECONTE, 1854:107 — vide *Dermestes caninus* var. *mannerheimi* LECONTE, 1854:107.
- Dermestes marginatus* MARSHAM, 1802:66 — non Dermestidae, recte Hydrophilidae: *Sphaeridium bipustulatum* FABRICIUS, 1781.
- Dermestes marginatus* PAYKULL, 1798:280 — vide *Globicornis marginata* (PAYKULL, 1798:280).
- Dermestes marginatus* PAYKULL, 1798:300 — non Dermestidae, recte Lathridiidae: *Enicmus minutusa* (LINNAEUS, 1767).
- Dermestes marginatus* THUNBERG, 1781:7 — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes marginatus* (FABRICIUS, 1787) VILLERS, 1789 b:218 — non Dermestidae, familia?
- Dermestes marginellus* QUENSEL, 1790:9 — non Dermestidae, recte Coccinellidae: *Hyperaspis reppensis* (HERBST, 1783).
- Dermestes marinus* FABRICIUS, 1787:35 — vide *Dermestes murinus* LINNAEUS, 1758:356.
- Dermestes marmoratus* MANNERHEIM, 1843:258 nec SAY, 1823:197 — vide *Dermestes caninus* var. *mannerheimi* LECONTE, 1854:107.
- Dermestes marmoratus* MELSHEIMER, 1806:6 — nomen nudum.
- Dermestes marmoratus* var. *mannerheimi*: JAYNE, 1882:350 — vide *Dermestes caninus* var. *mannerheimi* LECONTE, 1854:107.
- Dermestes megatoma* FABRICIUS, 1798:71 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Dermestes melanocephalus* LINNAEUS, 1758:356 — non Dermestidae, recte Hydrophilidae: *Cercyon melanocephalus* (LINNAEUS, 1758).
- Dermestes melanocephalus* (HERBST, 1793) PANZER, 1795:102 — non Dermestidae, recte Cryptophagidae: *Paramecosoma melanocephalum* (HERBST, 1793).
- Dermestes melas* MARSHAM, 1802:78 — non Dermestidae, recte Ptiliidae, *Ptenidium pusillum* (GYLLENHAL, 1808).
- Dermestes mesomelus* HERBST, 1792:143 — non Dermestidae, recte Cryptophagidae: *Atomaria mesomela* (HERBST, 1792).
- Dermestes micographus* LINNAEUS, 1761:143 — non Dermestidae, recte Scolytidae: *Pityophthorus micrographus* (LINNAEUS, 1758).
- Dermestes micrographus* LINNAEUS 1758:355 — non Dermestidae, recte Scolytidae: *Pityophthorus micrographus* (LINNAEUS, 1758).
- Dermestes minimus* MARSHAM, 1802:80 — non Dermestidae, recte Ptiliidae: *Acrotrichis atomaria* (DEGEER, 1774).
- Dermestes minimus* (HERBST, 1793) PANZER, 1795:102 — non Dermestidae, recte Ptiliidae: *Oligella foveolata* (ALLIBERT, 1844).
- Dermestes minimus* SCHRANK, 1798:429 — non Dermestidae, familia?

- Dermestes minutissimus* MOTSCHULSKY, 1850:196 — non *Dermestidae*, recte *Clambidae*: *Clambus armadillo* (DEGEER, 1774).
- Dermestes minutus* FABRICIUS, 1792:235 — non *Dermestidae*, recte *Lathridiidae*: *Melanophthalma gibbosa* (HERBST, 1793).
- Dermestes minutus* (FABRICIUS, 1775) SCHNEIDER, 1785:4 — non *Dermestidae*, recte *Bostrychidae*: *Dinoderus minutus* (FABRICIUS, 1775).
- Dermestes minutus* (FABRICIUS, 1776) SCHNEIDER, 1785:5 — non *Dermestidae*, familia?
- Dermestes minutus* PAYKULL, 1798:294 — non *Dermestidae*, recte *Cryptophagidae*: *Atomaria fimetaria* (HERBST, 1793).
- Dermestes minutus-laevis* VILLERS, 1789:55 — non *Dermestidae*, familia?
- Dermestes minutus-striatus* VILLERS, 1789:55 — non *Dermestidae*, recte *Hydrophilidae*: *Cryptopleurum minutum* (FABRICIUS, 1775).
- Dermestes mitutus* GYLLENHAL, 1827:132 — non *Dermestidae*, recte *Lathridiidae*: *Melanophthalma gibbosa* (HERBST, 1793).
- Dermestes mollis* LINNAEUS, 1758:355 — non *Dermestidae*, recte *Anobiidae*: *Ernobius mollis* (LINNAEUS, 1758).
- Dermestes mollis* (LINNAEUS, 1758) SCHRANK, 1781:22 — non *Dermestidae*, recte *Cleridae*: *Opilo mollis* (LINNAEUS, 1758).
- Dermestes monachus* (FABRICIUS, 1775) SCHNEIDER, 1785:4 — non *Dermestidae*, recte *Bostrychidae*: *Apate monachus* FABRICIUS, 1775.
- Dermestes mucoreus* LECONTE, 1854:108 — vide *Dermestes carnivorus* FABRICIUS, 1775:55.
- Dermestes multiguttatus* SCHRANK, 1798:427 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
- Dermestes multipunctatus* (FABRICIUS, 1792) THUNBERG, 1794:79 — non *Dermestidae*, recte *Mycetophagidae*: *Mycetophagus multipunctatus* FABRICIUS, 1792.
- Dermestes muricatus* LINNAEUS, 1767:562 — non *Dermestidae*, recte *Bostrychidae*: *Sinoxylon sexdentatum* (OLIVIER, 1790).
- Dermestes murinus* DEJEAN, 1821:46, nomen nudum — vide *Dermestes mustelinus* ERICHSON, 1846:433.
- Dermestes murinus* HERBST, 1792:120 nec LINNAEUS, 1758:356 — vide *Dermestes laniarius* ILLIGER, 1801:85.
- Dermestes murinus* LECONTE, 1854:108 — vide *Dermestes caninus* GERMAR, 1824:84.
- Dermestes murinus* OLIVIER, 1790: no. 9 p. 9 nec LINNAEUS, 1758: 356 — vide *Dermestes undulatus* BRAHM, 1790:114.
- Dermestes murinus* STRÖM, 1783:50 nec LINNAEUS, 1758:356 — non *Dermestidae*, recte *Byrrhidae*: *Porcinolus murinus* (FABRICIUS, 1794).
- Dermestes murinus* subvar. *auriceps* REITTER, 1905:244 — vide *Dermestes murinus auriceps* REITTER, 1905: 244.
- Dermestes murinus* var. *caninus*: JAYNE, 1882:350 — vide *Dermestes caninus* GERMAR, 1824:84.
- Dermestes murinus* var. *rattus*: JAYNE, 1882:351 — vide *Dermestes rattus* LECONTE, 1854:108.
- Dermestes murinus* var. *roseiventris*: HEER, 1841:25 — vide *Dermestes murinus* LINNAEUS, 1758:356.
- Dermestes musaeorum* STRÖM, 1783:52 — non *Dermestidae*, familia?
- Dermestes muscoreus* REITTER, 1881:28 — vide *Dermestes carnivorus* FABRICIUS, 1775:55.
- Dermestes museorum* LINNAEUS, 1761:145 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).
- Dermestes mustelinus* PERRIS, 1853:638 — vide *Dermestes aurichalceus* KÜSTER, 1846:50.
- Dermestes navalis* FABRICIUS, 1775:56 — non *Dermestidae*, familia?
- Dermestes navalis* HERBST, 1792:38 — non *Dermestidae*, recte *Tenebrionidae*: *Tribolium ferrugineum* (FABRICIUS, 1787).
- Dermestes nebulosus* DEGEER, 1774:197 — vide *Dermestes murinus* LINNAEUS, 1758:356.
- Dermestes nebulosus* MELSHEIMER, 1806:7 — nomen nudum.
- Dermestes niger* LINNAEUS, 1761:146 — non *Dermestidae*, recte *Dasytidae*: *Dasytes niger* (LINNAEUS, 1761).
- Dermestes niger* MELSHEIMER, 1806:7 — nomen nudum.
- Dermestes nigriceps* MARSHAM, 1802:72 — non *Dermestidae*, recte *Hydrophilidae*: *Cercyon nigriceps* (MARSHAM, 1802).

- Dermestes nigricornis* (FABRICIUS, 1787) THUNBERG, 1794:80 — non Dermestidae, recte Cryptophagidae:
Antherophagus nigricornis (FABRICIUS, 1787).
- Dermestes nigrinus* MARSHAM, 1802:77 — non Dermestidae, recte Hydrophilidae: *Limnebius truncatellus* (THUNBERG, 1794).
- Dermestes nigripennis* (KUGELANN, 1792) PAYKULL, 1798:292 — non Dermestidae, recte Cryptophagidae:
Atomaria nigripennis (KUGELANN, 1792).
- Dermestes nigripes* FABRICIUS, 1792:233 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Dermestes nigripes* OLIVIER, 1790: no. 9 p. 15 — non Dermestidae?
- Dermestes nitens* MARSHAM, 1802:79 — non Dermestidae, familia?
- Dermestes nitidulus* MARSHAM, 1802:79 — non Dermestidae, recte Ptiliidae: *Nossidium pilosellum* (MARSHAM, 1802).
- Dermestes nitidus* MARSHAM, 1802:75 — non Dermestidae, familia?
- Dermestes normandi* KALIK, 1951:154 — vide *Dermestes diadema* MOTSCHULSKY, 1849:89.
- Dermestes notatus* CEDERHIELM, 1798:41 — non Dermestidae, familia?
- Dermestes noxius* MULSANT et REY, 1868:55 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes nubilus* LECONTE, 1854:107 nec SAY, 1832:3 — vide *Dermestes caninus* var. *nubipennis* CASEY, 1900:141.
- Dermestes nubilus* SAY, 1832:3 — vide *Dermestes caninus* GERMAR, 1824:84.
- Dermestes oblongus* FOURCROY, 1785:19 — non Dermestidae, recte Bostrichidae: *Lyctus linearis* (GOEZE, 1777).
- Dermestes oblongus* SOLIER, 1849:365 — vide *Dermestes peruvianus* CASTELNAU, 1840:33.
- Dermestes oblongus* STURM, 1843:93 — nomen nudum.
- Dermestes obscurus* FABRICIUS, 1792:233 — non Dermestidae, recte Cryptophagidae: *Telmatophilus caricis* (OLIVIER, 1790).
- Dermestes obscurus* MARSHAM, 1802:72 — non Dermestidae, recte Hydrophilidae: *Megasternum boleto-phagum* (MARSHAM, 1802).
- Dermestes obscurus* MELSHEIMER, 1806:7, nomen nudum — vide *Attagenus cylindricornis* var. *obscurus* SAY, 1825:185.
- Dermestes obscurus* STURM, 1843:93 — nomen nudum.
- Dermestes obtusus* GYLLENHAL in SCHÖNHERR, 1808:88 — vide *Attagenus obtusus* (GYLLENHAL in SCHÖNHERR, 1808:88).
- Dermestes ochraceus* SCRIBA, 1791:193 — non Dermestidae, recte Byturidae: *Byturus tomentosus* var. *ochraceus* (SCRIBA, 1791).
- Dermestes ochropus* MARSHAM, 1802:64 — non Dermestidae, familia?
- Dermestes orientalis* KUSNEZOVA, 1928:20 — vide *Dermestes dimidiatus* STEVEN in SCHÖNHERR, 1808:89.
- Dermestes orientalis* ab. *rosea* KUSNEZOVA, 1928:22 — vide *Dermestes dimidiatus* ab. *rosea* KUSNEZOVA, 1928:22.
- Dermestes ovalis* MARSHAM, 1802:73 — non Dermestidae, familia?
- Dermestes ovatus* MARSHAM, 1802:76 — non Dermestidae, recte Phalacridae: *Olibrus aeneus* (FABRICIUS, 1792).
- Dermestes pallidus* FOURCROY, 1785:18 — non Dermestidae, familia?
- Dermestes pallidus* MARSHAM, 1802:79 — non Dermestidae, recte Coccinellidae: *Rhizobius litura* (FABRICIUS, 1787).
- Dermestes pallipes* FABRICIUS, 1787:378 — non Dermestidae, familia?
- Dermestes paniceus* LINNÆUS, 1758:357 — non Dermestidae, recte Anobiidae: *Stegobium paniceum* (LINNÆUS, 1758).
- Dermestes pantherinus* AHRENS, 1814:11 — vide *Attagenus pantherinus* (AHRENS, 1814:11).
- Dermestes pectinatus* (LAICHARTING, 1781) SCHNEIDER, 1785:5 — non Dermestidae, recte Anobiidae: *Ptilinus pectinicornis* (LINNÆUS, 1758).
- Dermestes pectinicornis* LINNÆUS, 1758:355 — non Dermestidae, recte Anobiidae: *Ptilinus pectinicornis* (LINNÆUS, 1758).

- Dermestes pedicularius* LINNAEUS, 1758:357 — non Dermestidae, recte Nitidulidae: *Cateretes pedicularius* (LINNAEUS, 1758).
- Dermestes pellio* LINNAEUS, 1758:355 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Dermestes pellucidus* FOURCROY, 1785:22 — non Dermestidae, familia?
- Dermestes pellucidus* GOEZE, 1777:149 — non Dermestidae, familia?
- Dermestes perpusillus* MARSHAM, 1802:80 — non Dermestidae, recte Ptiliidae: *Ptenidium pusillum* (GYLLENHAL, 1808).
- Dermestes persimilis* CROTCH, 1873:41 — nomen nudum.
- Dermestes pertinax* LINNAEUS, 1758:355 — non Dermestidae, recte Anobiidae: *Coelostethus pertinax* (LINNAEUS, 1758).
- Dermestes peruanus* ERICHSON, 1847:95 — vide *Dermestes peruvianus* CASTELNAU, 1840:33.
- Dermestes picatus* MARSHAM, 1802:80 — non Dermestidae, familia?
- Dermestes piceorhaeus* MARSHAM, 1802:78 — non Dermestidae, familia?
- Dermestes piceus* MARSHAM, 1802:69 nec THUNBERG, 1781:8 — non Dermestidae, recte Hydrophilidae: *Cercyon impressus* (STURM, 1807).
- Dermestes piceus* OLIVIER, 1790:no. 9 p. 10 nec THUNBERG, 1781:8 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Dermestes piceus* PANZER, 1795:97 — non Dermestidae, recte Serropalpidae: *Orchesia micans* (PANZER, 1794).
- Dermestes piceus* THUNBERG, 1781:8 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes picinus* MARSHAM, 1802:69 — non Dermestidae, recte Hydrophilidae: *Cercyon haemorrhoidalis* (FABRICIUS, 1775).
- Dermestes picipes* FABRICIUS, 1787:35 — non Dermestidae, recte Cioidae: *Cis boleti* (SCOPOLI, 1763).
- Dermestes pilosellus* MARSHAM, 1802:78 — non Dermestidae, recte Ptiliidae: *Nossidium pilosellum* (MARSHAM, 1802).
- Dermestes pilula* DEGEER, 1774:213 — non Dermestidae, recte Byrrhidae: *Cytilus sericeus* (FORSTER, 1771).
- Dermestes pilula* LINNAEUS, 1758:356 — non Dermestidae, recte Byrrhidae: *Byrrhus pilula* (LINNAEUS, 1758).
- Dermestes piniperda* LINNAEUS, 1758:355 — non Dermestidae, recte Scolytidae: *Myelophilus piniperda* (LINNAEUS, 1758).
- Dermestes pisorum* LINNAEUS, 1758:356 — non Dermestidae, recte Bruchidae: *Bruchus pisorum* (LINNAEUS, 1758).
- Dermestes plantaris* CURTIS, 1838:682 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Dermestes plicatus* SCOPOLI, 1763:19 — non Dermestidae, familia?
- Dermestes poligraphus* LINNAEUS, 1758:355 — non Dermestidae, recte Scolytidae: *Polygraphus poligraphus* (LINNAEUS, 1758).
- Dermestes politus* MARSHAM, 1802:75 — non Dermestidae, familia?
- Dermestes pollinctus* HOPE, 1834b:55 — vide *Dermestes frischii* KUGELANN, 1792:478.
- Dermestes polygraphus* GOEZE, 1777:131 — non Dermestidae, recte Scolytidae: *Polygraphus poligraphus* (LINNAEUS, 1758).
- Dermestes porcatus* (HERBST, 1793) PANZER, 1794:H. 23 p. 8 — non Dermestidae, recte Lathridiidae: *Enicmus minutus* (LINNAEUS, 1767).
- Dermestes psyllius* HERBST, 1783:21 — non Dermestidae, recte Nitidulidae: *Meligethes aeneus* (FABRICIUS, 1775).
- Dermestes psyllius* LINNAEUS, 1758:357 — non Dermestidae, recte Nitidulidae apud MOTSCHULSKY, 1858b: 145.
- Dermestes pubescens* HAUSMANN, 1799:34 — non Dermestidae, familia?
- Dermestes pubescens* ZETTERSTEDT, 1828:136 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Dermestes pulchellus* MARSHAM, 1802:72 — non Dermestidae, recte Hydrophilidae: *Cercyon pygmaeus* (ILLIGER, 1801).

- Dermestes pulicarius* LINNAEUS, 1758:357 — non Dermestidae, recte Nitidulidae: *Brachypterolus pulicarius* (LINNAEUS, 1758).
- Dermestes pulvinatus* SCOPOLI, 1772:87 — non Dermestidae, familia?
- Dermestes punctato-striatus* GOEZE, 1777:148 — non Dermestidae, familia?
- Dermestes punctatus* SCOPOLI, 1772:87 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
- Dermestes punctatus* (FABRICIUS, 1792) THUNBERG, 1794:79 — non Dermestidae, recte Mycetophagidae: *Triphylus bicolor* (FABRICIUS, 1792).
- Dermestes punctum* MARSHAM, 1802:80 — non Dermestidae, familia?
- Dermestes pusillus* PAYKULL, 1798:295 — non Dermestidae, recte Cryptophagidae: *Atomaria pusilla* (PAYKULL, 1798).
- Dermestes pygmaeus* FOURCROY, 1785:24 — non Dermestidae, familia?
- Dermestes pygmaeus* (FABRICIUS, 1787) VILLERS, 1789b:221 — non Dermestidae, recte Scolytidae: *Eccoptogaster pygmaeus* (FABRICIUS, 1787).
- Dermestes quadratus* (HERBST, 1793) PANZER, 1795:101 — non Dermestidae, recte Lathridiidae: *Lathridius lardarius* (DEGEER, 1775).
- Dermestes quadrimaculatus* MARSHAM, 1802:65 — non Dermestidae, recte Hydropilidae: *Sphaeridium bipustulatum* FABRICIUS, 1781.
- Dermestes quadrimaculatus* SCHRANK, 1781:25 — non Dermestidae, recte Hydropilidae: *Sphaeridium scarabaeoides* (LINNAEUS, 1758).
- Dermestes quadrimaculatus* (FABRICIUS, 1787) THUNBERG, 1794:79 — non Dermestidae, recte Mycetophagidae: *Mycetophagus quadrimaculatus* (FABRICIUS, 1787).
- Dermestes quadripunctata* FOURCROY, 1785:21 — non Dermestidae, familia?
- Dermestes quadripunctatus* FUSSLIN, 1775:4 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
- Dermestes quadripustulatus* PANZER, 1793:H. 6 p. 6 — non Dermestidae, recte Erotylidae: *Combocerus glaber* (SCHALLER, 1783).
- Dermestes quadripustulatus* PAYKULL, 1798:297 — non Dermestidae, familia?
- Dermestes quercus* SCHRANK, 1781:34 — non Dermestidae, familia?
- Dermestes quisquilius* (LINNAEUS, 1761) MARSHAM, 1802:71 — non Dermestidae, recte Hydrophilidae: *Cercyon quisquilius* (LINNAEUS, 1761).
- Dermestes rapax* CRISTOFORI et JAN, 1832:30 — nomen nudum.
- Dermestes ratus* SOLSKI, 1876:269 nec *D. rattus* LECONTE, 1854:108 — vide *Dermestes solskyi* DALLA TORRE, 1911:49.
- Dermestes rosae* SCOPOLI, 1763:15 — non Dermestidae, familia?
- Dermestes roseiventris* CASTELNAU, 1840:34 — vide *Dermestes murinus* LINNAEUS, 1758:356.
- Dermestes roseiventris* DEJEAN, 1821:46, nomen nudum — vide *Dermestes murinus* LINNAEUS, 1758:356.
- Dermestes ruber* THUNBERG, 1784:3 — non Dermestidae, recte Leiodidae: *Leiodes dubia* (KUGELANN, 1794).
- Dermestes ruficollis* FABRICIUS, 1775:57 — non Dermestidae, recte Cleridae, *Necrobia ruficollis* (FABRICIUS, 1775).
- Dermestes ruficollis* (OLIVIER, 1792) MARSHAM, 1802:68 — non Dermestidae, recte Leiodidae: *Amphioclytis globus* (FABRICIUS, 1792).
- Dermestes ruficornis* GMELIN, 1790:1600 — non Dermestidae, familia?
- Dermestes ruficornis* STURM, 1843:93 — nomen nudum.
- Dermestes rufipes* FABRICIUS, 1781:65 — non Dermestidae, recte Cleridae: *Necrobia rufipes* (DEGEER, 1775).
- Dermestes rufipes* MARSHAM, 1802:62 — non Dermestidae, recte Anobiidae: *Xyletinus laticollis* DUFTSCHMID, 1825 apud GYLLENHAL, 1827:330.
- Dermestes rufipes* SCOPOLI, 1772:88 — non Dermestidae, familia?
- Dermestes rufitarsis* CREUTZER in PANZER, 1796b:H. 35, t. 6 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Dermestes rufoapicalis* PIC, 1951c:7 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes rufofuscus* SOLIER, 1849:365 — vide *Dermestes peruvianus* CASTELNAU, 1840:33.

- Dermestes rufus* HERBST, 1783:22 — non Dermestidae, recte Coccinellidae: *Coccidula rufa* (HERBST, 1783).
Dermestes rugosus (HERBST, 1793) PANZER, 1795:101 — non Dermestidae, recte Lathridiidae: *Enicmus rugosus* (HERBST, 1793).
Dermestes salicinus O. F. MÜLLER, 1776:57 — non Dermestidae, familia?
Dermestes sambucci SCOPOLI, 1763:16 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
Dermestes sanguinicollis FABRICIUS, 1787:35 — non Dermestidae, recte Cleridae: *Orthopleura sanguinicollis* (FABRICIUS, 1787).
Dermestes sardous var. *striatellus* REITTER, 1908:45 — vide *Dermestes sardous* var. *fulvofasciatus* GANGLBauer, 1904b:18.
Dermestes scaber FABRICIUS, 1775:57 — non Dermestidae, recte Colydiidae: *Ulonotus scaber* (FABRICIUS, 1775).
Dermestes scanicus FABRICIUS, 1775:58 — non Dermestidae, recte Erotylidae: *Dacne bipustulata* (THUNBERG, 1781).
Dermestes scanicus LINNAEUS, 1758:357 — non Dermestidae, recte Cryptophagidae: *Cryptophagus scanicus* (LINNAEUS, 1758).
Dermestes scapularis FOURCROY, 1785:23 — non Dermestidae, familia?
Dermestes scarabaeoides LINNAEUS, 1758:356 — non Dermestidae, recte Hydrophilidae: *Sphaeridium scarabaeoides* (LINNAEUS, 1758).
Dermestes scaraboides SCOPOLI, 1763:15 — non Dermestidae, recte Nitidulidae: *Nitidula bipunctata* (LINNAEUS, 1758).
Dermestes schaefferi (HERBST, 1792:93) PANZER, 1795:95 — vide *Attagenus schaefferi* (HERBST, 1792:93).
Dermestes scolythus SULZER, 1776:I p. 21¹ — non Dermestidae, recte Scolytidae: *Scolytus scolythus* (FABRICIUS, 1775).
Dermestes scolythus (FABRICIUS, 1775) GOEZE, 1777:143 — non Dermestidae, recte Scolytidae: *Scolytus scolythus* (FABRICIUS, 1775).
Dermestes scolythus geoffroi GOEZE, 1777:143¹ — non Dermestidae, recte Scolytidae: *Scolytus scolythus* (FABRICIUS, 1775).
Dermestes scrophulariae LINNAEUS, 1758:356 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
Dermestes semi-coleopterus FOURCROY, 1785:25 — non Dermestidae, familia?
Dermestes semicoleoptratus PANZER, 1794:H. 24 p. 6 — non Dermestidae, recte Staphylinidae: *Anthobium abdominale* (GRAVENHORST, 1806).
Dermestes senegalensis CRISTOFORI et JAN, 1832:30, nomen nudum — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843:257.
Dermestes senex GERMAR, 1824:84 — vide *Dermestes maculatus* ab. *senex* GERMAR, 1824:84.
Dermestes serra FABRICIUS, 1792:234 — vide *Ctesias serra* (FABRICIUS, 1792:234).
Dermestes serra PANZER, 1795:108 — non Dermestidae, recte Anobiidae: *Dorcatoma serra* (PANZER, 1795).
Dermestes serraticornis THUNBERG, 1787:39 — nomen nudum.
Dermestes serratus PAYKULL, 1798:300 — non Dermestidae, recte Lathridiidae: *Corticaria serrata* (PAYKULL, 1798).
Dermestes sexdentatus BÖRNER, 1776:78 — non Dermestidae, recte Scolytidae: *Ips sexdentatus* (BÖRNER, 1776).
Dermestes sexdentatus FABRICIUS, 1792:232 — non Dermestidae, recte Silvanidae: *Oryzaephilus surinamensis* (LINNAEUS, 1758).
Dermestes sibericus ZHANTIEV, 1960:1629 — vide *Dermestes sibiricus* ERICHSON, 1846:427.
Dermestes signaticollis STURM, 1843:93, nomen nudum — vide *Dermestes coronatus* STEVEN in SCHÖNERR, 1808:85.
Dermestes silphoides SCOPOLI, 1772:89 — non Dermestidae, familia?

¹ Rejected and invalid name — see Bull. Zool. Nomencl., London, 20:416 (Opinion No. 683).

- Dermestes similis* MARSHAM, 1802:68 — non Dermestidae, recte Hydrophilidae: *Cercyon haemorrhoidalis* (FABRICIUS, 1775).
- Dermestes sobrinus* LECONTE, 1854:108 — vide *Dermestes carnivorus* FABRICIUS, 1775:55.
- Dermestes sordidus* MARSHAM, 1802:69 — non Dermestidae, recte Hydrophilidae: *Cryptopleurum minutum* (FABRICIUS, 1775).
- Dermestes spadiceus* MELSHEIMER, 1806:7 — nomen nudum.
- Dermestes spadiceus* PANZER, 1795:104 — non Dermestidae, familia?
- Dermestes spinipennis* CRISTOFORI et JAN, 1832:30 — nomen nudum.
- Dermestes spinipes* SCOPOLI, 1772:87 — non Dermestidae, familia?
- Dermestes stercorarius* MARSHAM, 1802:76 — non Dermestidae, recte Hydrophilidae: *Megasternum boleto-phagum* (MARSHAM, 1802).
- Dermestes stercoreus* LINNAEUS, 1758:357 — non Dermestidae, recte Mycetophagidae: *Typhaea stercorea* (LINNAEUS, 1758).
- Dermestes striatus* FOURCROY, 1785:20 — non Dermestidae, recte Hydrophilidae: *Helophorus nubilus* (FABRICIUS, 1776).
- Dermestes striatus* KOLENATI, 1846:58 — vide *Dermestes bicolor* FABRICIUS, 1781:64.
- Dermestes sturmii* CRISTOFORI et JAN, 1832:89 — nomen nudum.
- Dermestes subcostatus* GERSTAECKER, 1884:45 nec MURRAY, 1867:22 — vide *Dermestes gerstaeckeri* DALLA TORRE, 1911:44.
- Dermestes subcostatus* MURRAY, 1867:22 — vide *Dermestes ater* DEGEER, 1774:223.
- Dermestes subfasciatus* GYLLENHAL, 1808:155 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Dermestes subterraneus* FABRICIUS, 1801:318 — non Dermestidae, recte Endomychidae: *Mycetaea hirta* (MARSHAM, 1802).
- Dermestes sulcatus* ROSSI, 1794:80 — non Dermestidae, familia?
- Dermestes sulcatus* THUNBERG, 1784:2 — non Dermestidae, recte Tenebrionidae: *Eledona agricola* (HERBST, 1783).
- Dermestes surinamensis* LINNAEUS, 1758:357 — non Dermestidae, recte Silvanidae: *Oryzaephilus surinamensis* (LINNAEUS, 1758).
- Dermestes suturalis* FABRICIUS, 1801:317 — non Dermestidae, recte Mycetophagidae: *Pseudotriphyllus suturalis* (FABRICIUS, 1801).
- Dermestes suturatus* FOURCROY, 1785:23 — non Dermestidae, familia?
- Dermestes teretiusculus* SCHNEIDER, 1785:5 — nomen nudum.
- Dermestes terminatus* MARSHAM, 1802:70 — non Dermestidae, recte Hydrophilidae: *Cercyon terminatus* (MARSHAM, 1802).
- Dermestes tessellatus* FOURCROY, 1785:25 — non Dermestidae, familia?
- Dermestes tessellaticollis* MOTSCHULSKY, 1859:491, nomen nudum — vide *Dermestes tessellatocollis* MOTSCHULSKY, 1860:124.
- Dermestes tessellatocollis* ab. *pici* LEPESME, 1950:53 — vide *Dermestes tessellatocollis pici* MROCKOWSKI, 1963:377.
- Dermestes tessellatus* ERICHSON, 1846:432 nec FABRICIUS, 1775:56 — vide *Dermestes erichsoni* GANGLBauer, 1904b:16.
- Dermestes tessellatus* FABRICIUS, 1775:56 — non Dermestidae, recte Anobiidae: *Xestobium rufovillosum* (DEGEER, 1774).
- Dermestes tessellatus* GYLLENHAL, 1808:150 nec FABRICIUS, 1775:56 — vide *Dermestes gyllenhali* CASTELNAU, 1840:34.
- Dermestes tessellatus* HERBST, 1792:122 nec FABRICIUS, 1775:56 — vide *Dermestes caninus* GERMAR, 1824:84.
- Dermestes tessellatus* ILLIGER, 1801:84 nec FABRICIUS, 1775:56 — vide *Dermestes undulatus* BRAHM, 1790:114.
- Dermestes tessellatus* OLIVIER, 1790:No. 9 p. 9 nec FABRICIUS, 1775:56 — vide *Dermestes mustelinus* ERICHSON, 1846:433.
- Dermestes tessellatus* var. *vulpecula*: DEJEAN, 1821:46 — vide *Dermestes undulatus* BRAHM, 1790:114.

- Dermestes testaceus* FABRICIUS, 1775:57 — non Dermestidae, familia?
- Dermestes testaceus* FABRICIUS, 1792:233 — non Dermestidae, recte Mycetophagidae: *Typhaea stercorea* (LINNAEUS, 1758).
- Dermestes testaceus* (LINNAEUS, 1767) SCHRANK, 1781:27 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
- Dermestes testaceus* VILLERS, 1789b:214 — non Dermestidae, familia?
- Dermestes testaceus* (FABRICIUS, 1787) VILLERS, 1789b:221 — non Dermestidae, recte Scolytidae: *Myelophilus piniperda* (LINNAEUS, 1758).
- Dermestes testudinarium* FOURCROY, 1785:24 — non Dermestidae, recte Hydrophilidae: *Sphaeridium bipustulatum* FABRICIUS, 1781.
- Dermestes thorace marginato* SCHRANK, 1776:49 — non Dermestidae, recte Cioidae: *Cis boleti* (SCOPOLI, 1763).
- Dermestes thoracicus* DEJEAN, 1821:46, nomen nudum — vide *Dermestes pardalis* BILLBERG in SCHÖNHERR, 1808:90.
- Dermestes thoracicus* FOURCROY, 1785:24 — non Dermestidae, familia?
- Dermestes thoracicus* GENÉ, 1836:22 — vide *Dermestes pardalis* BILLBERG in SCHÖNHERR, 1808:90.
- Dermestes tigrinus* FABRICIUS, 1792:229 — vide *Attagenus bifasciatus* var. *tigrinus* (FABRICIUS, 1792:229).
- Dermestes tomentosus* DEGEER, 1774:199 — non Dermestidae, familia?
- Dermestes tomentosus* FABRICIUS, 1775:57 — non Dermestidae, recte Byturidae: *Byturus tomentosus* (FABRICIUS, 1775).
- Dermestes tomentosus* PANZER, 1797:H. 40 p. 12 — non Dermestidae, recte Dasytidae: *Danacea nigritarsis* (KÜSTER, 1850).
- Dermestes tomentosus* PAYKULL, 1798:282 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
- Dermestes tomentosus* SCHRANK, 1781:30 — non Dermestidae, familia?
- Dermestes tomentosus* var. *ochraceus* ILLIGER, 1798:321 — non Dermestidae, recte Byturidae: *Byturus aestivus* (LINNAEUS, 1758).
- Dermestes tricolor* FOURCROY, 1785:24 — non Dermestidae, recte Hydrophilidae: *Sphaeridium scarabaeoides* (LINNAEUS, 1758).
- Dermestes trifasciatus* FABRICIUS, 1787:34 — vide *Attagenus trifasciatus* (FABRICIUS, 1787:34).
- Dermestes trisulcus* FOURCROY, 1785:23 — non Dermestidae, familia?
- Dermestes truncatus* CASEY, 1916:180 — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes typographus* LINNAEUS, 1758:355 — non Dermestidae, recte Scolytidae: *Ips typographus* (LINNAEUS, 1758).
- Dermestes undatus* LINNAEUS, 1758:355 — vide *Megatoma undata* (LINNAEUS, 1758:355).
- Dermestes undulatus* LINNAEUS, 1761:141 — vide *Megatoma undata* (LINNAEUS, 1758:355).
- Dermestes undulatus talpinus*: LEPESME, 1950:46 — vide *Dermestes talpinus* MANNERHEIM, 1843:257.
- Dermestes unicolor* BRAHM, 1791:144 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Dermestes unidentatus* FABRICIUS, 1792:232 — non Dermestidae, recte Silvanidae: *Silvanus unidentatus* (FABRICIUS, 1792).
- Dermestes unipunctatus* (LINNAEUS, 1758) MARSHAM, 1802:70 — non Dermestidae, recte Hydrophilidae: *Cercyon unipunctatus* (LINNAEUS, 1758).
- Dermestes urticae* FABRICIUS, 1792:235 — non Dermestidae, recte Nitidulidae: *Brachypterus urticae* (FABRICIUS, 1792).
- Dermestes variabilis* FABRICIUS, 1792:234 — non Dermestidae, recte Cioidae: *Cis boleti* (SCOPOLI, 1763).
- Dermestes variabilis* HERBST, 1792:141 — non Dermestidae, recte Mycetophagidae: *Typhaea stercorea* (LINNAEUS, 1758).
- Dermestes variegatus* BRULLÉ, 1832:163 — vide *Dermestes undulatus* BRAHM, 1790:114.
- Dermestes variegatus* FOURCROY, 1785:20 — non Dermestidae, recte Nitidulidae: *Soronia grisea* (LINNAEUS, 1758).

- Dermestes variegatus* SCOPOLI, 1763:16 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Dermestes varius* (FABRICIUS, 1775:60) SCHNEIDER, 1785:5 — vide *Anthrenus verbasci* (LINNAEUS, 1767: 568).
- Dermestes vastator* KUGELANN, 1792:479 — non Dermestidae, recte Throscidae: *Throscus dermestoides* (LINNAEUS, 1766).
- Dermestes vernalis* O. F. MÜLLER, 1776:57 — non Dermestidae, familia?
- Dermestes versicolor* CASTELNAU, 1840:33 — vide *Dermestes carnivorus* FABRICIUS, 1775:55.
- Dermestes versicolor* ILLIGER, 1801:86 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Dermestes vespillo* (LINNAEUS, 1758) SCOPOLI, 1763:13 — non Dermestidae, recte Silphidae: *Nicrophorus vespillo* (LINNAEUS, 1758).
- Dermestes vigintiguttatus* FABRICIUS, 1775:56 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
- Dermestes vigintipunctatus* MEGERLE, 1805:8 — ?
- Dermestes villosus* GMELIN, 1790:1600 — non Dermestidae, recte Heteroceridae: *Heterocerus marginatus* (FABRICIUS, 1787).
- Dermestes vini* LAICHARTING, 1781:65 — non Dermestidae, familia?
- Dermestes vini* PANZER, 1797:H. 40, p. 14 — non Dermestidae, recte Cryptophagidae: *Micrambe vini* (PANZER, 1797).
- Dermestes violaceus* LINNAEUS, 1758:356 — non Dermestidae, recte Cleridae: *Necrobia violacea* (LINNAEUS, 1758).
- Dermestes viridanus* MELSHEIMER, 1806:7 — nomen nudum.
- Dermestes viridis* (THUNBERG, 1781) GMELIN, 1790:1598 — non Dermestidae, recte Cleridae: *Notostenus viridis* (THUNBERG, 1781).
- Dermestes viridis* SCHNEIDER, 1785:5 — nomen nudum.
- Dermestes vittatus* (FABRICIUS, 1787) VILLERS, 1789b:221 — non Dermestidae, recte Scolytidae: *Hylesinus vittatus* (FABRICIUS, 1787).
- Dermestes vorax* MOTSCHULSKY, 1859:491, nomen nudum — vide *Dermestes vorax* MOTSCHULSKY, 1860:123.
- Dermestes vulpecula* HERBST, 1792:123 — vide *Dermestes undulatus* BRAHM, 1790:114.
- Dermestes vulpinus* FABRICIUS, 1781:64 — vide *Dermestes maculatus* DEGEER, 1774:223.
- Dermestes vulpinus* HERBST, 1792:124 nec FABRICIUS, 1781:64 — vide *Dermestes frischii* KUGELANN, 1792: 478.
- Dermestes vulpinus* var. *lupinus*: LECONTE, 1854:109 — vide *Dermestes maculatus* var. *lupinus* MANNERHEIM, 1843:257.
- Dermestes vulpinus* var. *rattulus* MULSANT et REY, 1868:30 — vide *Dermestes maculatus* ab. *rattulus* MULSANT et REY, 1868:30.
- Dermestes vulpinus* var. *senex*: REITTER, 1906:377 — vide *Dermestes maculatus* ab. *senex* GERMAR, 1824:84.
- Dermestes vulpinus* var. *sudanicus* GREDLER, 1878:506 — vide *Dermestes maculatus* var. *sudanicus* GREDLER, 1878:506.
- Dermestes xantocephalus* QUENSEL, 1790:10 — non Dermestidae, recte Coccinellidae: *Hyperaspis reppensis* (HERBST, 1783).

Genus: *Montandonia* JACQUET, 1886b : 320

Type species: *Montandonia catopoides* JACQUET, 1886b : 319.

List of species:

Montandonia latissima (BIELZ, 1850 : 180) REITTER, 1891 : 169.

Romania

Syn.: *Dermestes latissimus* BIELZ, 1850:180.

Montandonia catopoides JACQUET, 1886b:319. — REITTER,
1891:169. —

Montandonia catopoides JACQUET, 1886b:319 — vide *Montandonia latissima* (BIELZ, 1850:180).
Montandonia depressa (GEBLER, 1830:99) HEYDEN, 1893:64 — vide *Dermestes depressus* GEBLER, 1830:99.
Montandonia roubali KALIK, 1951:152 — vide *Dermestes roubali* (KALIK, 1951:152).

Attageninae CASTELNAU, 1840 : 35

Type genus: *Attagenus* LATREILLE, 1802 : 121.

Syn.: *Attagenites* CASTELNAU, 1840:35.

Attageni: LECONTE, 1861:107.

Attagenini: CASEY, 1900:139.

Attageninae: DALLA TORRE, 1911:50.

Genus: *Sefrania* PIC, 1899 : 28

Type species: *Sefrania bleusei* PIC, 1899 : 29.

List of species:

Sefrania bleusei PIC, 1899 : 29.

Algeria,
Tunisia

Genus: *Attagenus* LATREILLE, 1802 : 121

Type species: *Dermestes pellio* LINNAEUS, 1758 : 355.

Syn.: *Megatoma* KUGELANN, 1792: 479 nec HERBST, 1792:92. Type species: *Megatoma schaefferi* HERBST, 1792:93.

Subgenera:

Telopes REDTENBACHER, 1843 : 976.

Type species: *Telopes dispar* REDTENBACHER, 1843 : 976.

Syn.: *Telopus* MULSANT et REY, 1868:101.

Aethriostoma MOTSCHULSKY, 1858 : 47.

Type species: *Aethriostoma undulata* MOTSCHULSKY, 1858 : 47.

Syn.: *Brachysphyrus* BLACKBURN, 1903:159. — ARROW, 1915:426. Type species: *Brachysphyrus irroratus* BLACKBURN, 1903:160.

Pseudotelopes PIC, 1916b:2. — BEAL, 1961:120. Type species: *Pseudotelopes simoni* PIC, 1916b:2.

Lanorus MULSANT et REY, 1868 : 68.

Type species: *Dermestes vigintiguttatus* FABRICIUS, 1775 : 56.

Ranolus BLAIR, 1929 : 382.

Type species: *Attagenus cavernicola* BLAIR, 1929 : 382.

List of species:

Attagenus abbreviatus HEER, 1856 : 140.

Canary Islands

†*Attagenus aboriginalis* WICKHAM, 1913 : 361.

Colorado
(Miocene)

Attagenus adspersus (BLANCHARD, 1843 : 61) LACORDAIRE,
1854 : 466.

Bolivia

Syn.: *Megatoma adspersa* BLANCHARD, 1843:61.

<i>Attagenus aeneus</i> ROTH, 1851 : 122.	Ethiopia
<i>Attagenus africanus</i> MROCKOWSKI, 1958 : 558.	East Africa
Syn.: <i>Attagenus diversepubescens</i> PIC, 1953:112 nec PIC, 1937d:195.	
<i>Attagenus albonotatus</i> PIC, 1927e : 2.	South Africa
<i>Attagenus amurensis</i> PIC, 1942 : 7.	Siberia
<i>Attagenus antennatus</i> CASTELNAU, 1840 : 36.	Spain
<i>Attagenus anthrenoides</i> (WOLLASTON, 1864 : 159) DALLA TORRE, 1911 : 51.	Canary Islands
Syn.: <i>Telopes anthrenoides</i> WOLLASTON, 1864:159.	
<i>Attagenus apicalis</i> PIC, 1942 : 7.	Eritrea
<i>Attagenus apicebrunneus</i> KALIK, 1955b : 316.	Tanganyika
<i>Attagenus arcuatefasciatus</i> PIC, 1951 : 5.	South Viet-Nam
<i>Attagenus aristidis</i> (PIC, 1894d : 119) DALLA TORRE, 1911 : 51.	Egypt, Algeria
Syn.: <i>Telopes aristidis</i> PIC, 1894d: 119.	
<i>Attagenus arrowi</i> KALIK, 1954 : 367.	Yunnan
<i>Attagenus asmaranus</i> PIC, 1942 : 8.	Erythrea
<i>Attagenus assuanensis</i> (PIC, 1899c : 254) comb. nov.	South Egypt
Syn.: <i>Telopes assuanensis</i> PIC, 1899c:254.	
<i>Attagenus astacurus</i> PEYERIMHOFF, 1931 : 64.	Central Sahara
<i>Attagenus ater</i> (PIC, 1937c : 6) comb. nov.	New Guinea
Syn.: <i>Aethriostoma atra</i> PIC, 1937c:6.	
<i>Attagenus atricolor</i> PIC, 1931b : 4.	Equator Africa, Congo (Kinshasa)
<i>Attagenus atripennis</i> PIC, 1938b : 9.	Yemen
<i>Attagenus atrolucens</i> CASEY, 1916 : 183.	Indiana
<i>Attagenus attenuatus</i> (PIC, 1894h : 210) DALLA TORRE, 1911 : 51.	Algeria
Syn.: <i>Telopes attenuatus</i> PIC, 1894h:210.	
<i>Attagenus augustatus</i> BALLION, 1871 : 330.	Tadzhik SSR, China, Mongolia
Syn.: <i>Attagenus angustatus</i> SOLSKIJ, 1876:271. — MROCKOWSKI, 1963: 378.	
<i>Hadrotoma angustata</i> (SOLSKIJ, 1876:271) REITTER, 1889f:163.	
<i>Globicornis angustata</i> (SOLSKIJ, 1876:271) HEYDEN, 1893:64.	
<i>Attagenus augustatus gobicola</i> FRIVALDSZKY, 1892 : 122. — ZHANTIEV, 1963c : 416.	China, Mongolia, Tian-Shan, Alai, Env. of Baykal
Syn.: <i>Attagenus gobicola</i> FRIVALDSZKY, 1892:122.	
<i>Attagenus globicola</i> DALLA TORRE, 1911:53.	
<i>Attagenus angustatus gobicola</i> : ZHANTIEV, 1963c:416.	

<i>Attagenus aurantiacus</i> REITTER, 1900c : 274.	Syria
<i>Attagenus aurantiacus erevanicus</i> ZHANTIEV, 1963c : 417.	Armenia
<i>Attagenus auratofasciatus</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus australis</i> (MONTROUZIER, 1860 : 264) GEMMINGER et HAROLD, 1868 : 915.	New Caledonia, Singapore, Zanzibar
Syn.: <i>Trinodes australis</i> MONTROUZIER, 1860:264.	
<i>Attagenus basalis</i> PIC, 1928c : 257.	East Africa
<i>Attagenus basimaculatus</i> PIC, 1952b : 30.	Ethiopia
<i>Attagenus bifasciatus</i> (OLIVIER, 1790 : no. 9 p. 13) STURM, 1843 : 319.	Sicily, south Europe, North Africa, Asia Minor, Caucasus, Iraq, Iran, Turkmenia, Turkestan, Afghanistan
Syn.: <i>Dermestes bifasciatus</i> OLIVIER, 1790:no. 9 p. 13. <i>Attagenus pulcher</i> FALDERMANN, 1835:226. — KOLENATI, 1846:58. <i>Attagenus wachanrui</i> MULSANT et REY, 1868:91. — REITTER, 1906:378.	
<i>Megatoma tigrina</i> var. <i>pulcher</i> : REITTER, 1887b:49. <i>Megatoma tigrina</i> var. <i>wachanrui</i> : REITTER, 1887b:50. <i>Attagenus tigrinus</i> var. <i>vanchurui</i> DESBROCHERS DES LOGES, 1898: 73.	
<i>Attagenus bifasciatus</i> var. <i>chakouri</i> PIC, 1907 : 153.	Egypt
<i>Attagenus bifasciatus</i> ab. <i>hybridus</i> RAGUSA, 1926 : 74.	Sicily
<i>Attagenus bifasciatus</i> var. <i>persicus</i> REITTER, 1881b : 78.	North Iran, Greece
Syn.: <i>Attagenus tigrinus</i> var. <i>persicus</i> REITTER, 1881b:78. <i>Megatoma tigrina</i> var. <i>persica</i> : REITTER, 1887b:49.	
<i>Attagenus bifasciatus</i> var. <i>rossi</i> GANGLBAUER, 1904b : 26.	Italy, South Europe
Syn.: <i>Dermestes bifasciatus</i> ROSSI, 1794:79 nec OLIVIER, 1790:no. 9 p. 13.	
<i>Attagenus bifasciatus</i> (ROSSI, 1794:79) SCHMIDT, 1844:31. <i>Megatoma bifasciata</i> (ROSSI, 1794:79) REITTER, 1887b:48.	
<i>Attagenus bifasciatus</i> var. <i>sexmaculatus</i> SAHLBERG, 1913c : 107.	Israel, Lebanon
<i>Attagenus bifasciatus</i> var. <i>simonis</i> REITTER, 1881 : 34. — REITTER, 1906 : 378.	Syria, Israel, Egypt
Syn.: <i>Attagenus simonis</i> REITTER, 1881:34. <i>Megatoma simonis</i> (REITTER, 1881:34) REITTER, 1887b:49.	
<i>Attagenus bifasciatus</i> var. <i>tigrinus</i> (FABRICIUS, 1792 : 229) SCHAUM, 1852 : 38. — GANGLBAUER, 1904b : 26.	Italy, Whole Mediterranea, Asia Minor
Syn.: <i>Dermestes tigrinus</i> FABRICIUS, 1792:229.	
<i>Attagenus poecilus</i> GERMAR, 1839:9. — REITTER, 1883:89. <i>Attagenus tigrinus</i> (FABRICIUS, 1792:229) SCHAUM, 1852:38. <i>Megatoma tigrina</i> (FABRICIUS, 1792:229) REITTER, 1887b:49.	

<i>Attagenus bifasciatus</i> var. <i>wegeneri</i> PIC, 1899c : 252.	Israel, Lebanon, Tunisia
Syn.: <i>Attagenus tigrinus</i> v. <i>wegeneri</i> PIC, 1899c:252.	
<i>Attagenus bifasciatus</i> ab. <i>wegeneri</i> NORMAND, 1936:297.	
<i>Attagenus birmanicus</i> ARROW, 1915 : 428.	Burma
<i>Attagenus biskrensis</i> PIC, 1904 : 2.	Algeria
<i>Attagenus brasiliensis</i> PIC, 1923 : 3.	Brasil
<i>Attagenus breviusculus</i> (REITTER, 1881 : 35) DALLA TORRE, 1911 : 52.	South Africa
Syn.: <i>Telopes breviusculus</i> REITTER, 1881:35.	
<i>Attagenus brittoni</i> KALIK, 1954 : 368.	Iran
<i>Attagenus brunneopunctatus</i> (PIC, 1893 : 122) WINKLER, 1926 : 678.	Israel
Syn.: <i>Telopes brunneopunctatus</i> PIC, 1893:122.	
<i>Attagenus brunneopunctatus</i> var. <i>impunctatus</i> (PIC, 1894c : 116) WINKLER, 1926 : 678.	Syria
Syn.: <i>Telopes brunneopunctatus</i> v. <i>impunctatus</i> PIC, 1894c:116.	
<i>Attagenus brunnescens</i> (PIC, 1904d : 73) DALLA TORRE, 1911 : 52.	Turkey
Syn.: <i>Telopes brunnescens</i> PIC, 1904d:73.	
<i>Attagenus brunneus</i> FALDERMANN, 1835 : 227.	Armenia, Mediterranea, Caucasus
Syn.: <i>Attagenus dalmatinus</i> DEJEAN, 1821:46, nomen nudum. — KÜSTER, 1847:44.	
<i>Attagenus sordidus</i> HEER, 1841b:438. — FUENTE, 1930:48.	
<i>Attagenus dalmatinus</i> KÜSTER, 1847:44. — syn. nov.	
<i>Attagenus marginicollis</i> KÜSTER, 1847b:76. — REITTER, 1880b: 510.	
<i>Attagenus fulvipes</i> MULSANT et REY, 1868:71. — TOURNIER, 1869:11.	
<i>Attagenus piceus</i> var. <i>sordidus</i> : REITTER, 1881b:76.	
<i>Attagenus piceus</i> var. <i>dalmatinus</i> : REITTER, 1881b:76.	
<i>Attagenus piceus</i> var. <i>brunneus</i> : REITTER, 1883:89.	
<i>Megatoma picea</i> var. <i>dalmatina</i> : REITTER, 1887b:46.	
<i>Megatoma picea</i> var. <i>sordida</i> : REITTER, 1887b:47.	
<i>Megatoma marginicollis</i> (KÜSTER, 1847b:76) REITTER, 1887b:47.	
<i>Attagenus piceus</i> var. <i>obliquus</i> REY, 1889:59.	
<i>Attagenus longicornis</i> PIC, 1894b:66. — syn. nov.	
<i>Attagenus fulvipes</i> var. <i>obliquus</i> : PIC, 1895b:116.	
<i>Attagenus megatoma dalmatinus</i> : MROCKOWSKI, 1962b:547.	
<i>Attagenus calabricus</i> REITTER, 1881b : 77.	Italy, Sicily
Syn.: <i>Megatoma calabrica</i> (REITTER, 1881b:77) REITTER, 1887b:48.	
<i>Attagenus canadensis</i> CASEY, 1916 : 183.	Canada

<i>Attagenus capensis</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus caracal</i> ZHANTIEV, 1963 : 456.	China, Uzbek SSR
<i>Attagenus cavernicola</i> BLAIR, 1929 : 382.	Selangor
<i>Attagenus cinereus</i> (THUNBERG, 1815 : 151) comb. nov. ¹	South Africa
Syn.: <i>Anthrenus cinereus</i> THUNBERG, 1815:151.	
<i>Attagenus cinnamomeus</i> ROTH, 1851 : 122.	Ethiopia, Senegal, Algeria, Yemen, Egypt, Syria
Syn.: <i>Attagenus unifasciatus</i> FAIRMAIRE et COQUEREL, 1860:168. — REITTER, 1881:33.	Algeria, Tunisia, Morocco
<i>Megatoma cinnamomea</i> (ROTH, 1851:122) REITTER, 1887b:50.	
<i>Attagenus civetta</i> (MULSANT et REY, 1868 : 104) REITTER, 1881b : 80.	Oran, Algeria, Tunisia
Syn.: <i>Telopus civetta</i> MULSANT et REY, 1868:104.	
<i>Telopes civetta</i> (MULSANT et REY, 1868:104) REITTER, 1887b:54.	
<i>Attagenus coquereli</i> (MULSANT et REY, 1868 : 102) DALLA TORRE, 1911 : 52.	Egypt
Syn.: <i>Telopus coquereli</i> MULSANT et REY, 1868:102.	USA
<i>Attagenus curvicornis</i> SAHLBERG, 1913 : 26.	Pennsylvania
<i>Attagenus cylindricornis</i> SAY, 1825 : 185.	Pennsylvania
Syn.: <i>Dermestes cylindricornis</i> MELSHEIMER, 1806:7, nomen nudum — SAY, 1825:185.	
<i>Attagenus cylindricollis</i> MELSHEIMER, 1853:48.	
<i>Attagenus cylindricornis</i> var. <i>floricola</i> SAY, 1825 : 185.	
Syn.: <i>Dermestes floricola</i> MELSHEIMER, 1806:7, nomen nudum — SAY, 1825:185.	
<i>Attagenus cylindricornis</i> var. <i>obscurus</i> SAY, 1825 : 185.	
Syn.: <i>Dermestes obscurus</i> MELSHEIMER, 1806:7, nomen nudum — SAY, 1825:185.	
<i>Attagenus cyphonoides</i> REITTER, 1881 : 34.	Egypt, Tunisia, Morocco, England (introduced), Turkmen SSR, Uzbek SSR, Tadzhik SSR, India, USA, Alabama, Arizona, Kansas, New Mexico, Mexico: Lower California
Syn.: <i>Attagenus alfierii</i> PIC, 1910:17. — ZHANTIEV, 1963c:416.	
<i>Trogoderma cyphonoides</i> (REITTER, 1881:34) ARROW, 1915:427.	

¹ The type-specimen was examined (preserved in Uppsala).

<i>Attagenus decoloratus</i> MULSANT et REY, 1868 : 96.	Mediterranea
<i>Attagenus deficiens</i> CASEY, 1900 : 146.	New York, District of Columbia, Virginia
<i>Attagenus dichrous</i> ROTH, 1851 : 122.	Ethiopia
<i>Attagenus diversepubescens</i> PIC, 1937d : 195.	Dresden (Germany) introduced probably from Costa Rica
<i>Attagenus diversesignatus</i> PIC, 1942b : 10	South Africa
<i>Attagenus diversus</i> REITTER, 1881 : 30.	South Africa, Zanzibar
<i>Attagenus donckieri</i> PIC, 1916c : 15.	Angola, Congo (Kinshasa)
<i>Attagenus donckieri</i> var. <i>latepubescens</i> PIC, 1952b : 30.	Ethiopia
<i>Attagenus duplex</i> (REITTER in HEYDEN, 1890b : 359) HEYDEN, 1893 : 64.	Uzbek SSR, Turkmen SSR, Tadzhik SSR, Kazakh SSR
Syn.: <i>Telopes duplex</i> REITTER in HEYDEN, 1890b:359.	
<i>Attagenus elongatulus</i> CASEY, 1900 : 147.	Nebraska to Utah, New York
Syn.: <i>Attagenus elongatus</i> CASEY, 1916:183.	Cape Verde Is. Sahara, Tunisia Morocco
<i>Attagenus ensicornis</i> WOLLASTON, 1867 : 80.	
<i>Attagenus eremivagus</i> PEYERIMHOFF, 1943 : 11.	
<i>Attagenus escalerai</i> MADER, 1954 : 53.	
Syn.: <i>Attagenus antennatus</i> ESCALERA, 1914:137 nec CASTELNAU, 1840: 36.	
<i>Attagenus extricatus</i> CASEY, 1900 : 146.	Pennsylvania
<i>Attagenus fairmairei</i> MROCKOWSKI, 1958 : 558.	Madagascar
Syn.: <i>Aethriostoma fasciolata</i> FAIRMAIRE, 1897:97.	
<i>Attagenus fasciolatus</i> (FAIRMAIRE, 1897:97) MROCKOWSKI, 1958: 558 nec SOLSKI, 1876:275.	
<i>Attagenus fallax</i> GENÉ, 1839 : 19.	Sardinia, Mediterranea
Syn.: <i>Attagenus sexguttatus</i> STURM, 1843:94. — KÜSTER, 1850b:2. <i>Megatoma fallax</i> (GENÉ, 1839:19) REITTER, 1887b:48.	
<i>Attagenus fallax</i> var. <i>addendus</i> SAHLBERG, 1903e : 34.	Tunisia
<i>Attagenus fallax</i> var. <i>distinctus</i> MULSANT et REY, 1868 : 85. — REITTER, 1881b : 77.	Algeria, Greece, Italy, Tunisia
Syn.: <i>Attagenus distinctus</i> MULSANT et REY, 1868:85. <i>Megatoma fallax</i> var. <i>distincta</i> : REITTER, 1887b:48.	

<i>Attagenus fallax</i> var. <i>octomaculatus</i> PIC, 1894d : 119	Algeria
<i>Attagenus fasciatopunctatus</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus fasciatus</i> (THUNBERG, 1795 : 105) DEJEAN, 1837 : 139.	Sweden, nearly cosmopolitan
Syn.: <i>Anthrenus fasciatus</i> THUNBERG, 1795:105. <i>Anthrenus gloriosae</i> FABRICIUS, 1798:76. — MROCKOWSKI, 1964: 179.	
<i>Dermestes fasciatus</i> (THUNBERG, 1795:105) SCHÖNHERR, 1808:88.	
<i>Attagenus annulatus</i> DEJEAN, 1821:46, nomen nudum. — DEJEAN, 1837:139.	
<i>Attagenus annulifer</i> CASTELNAU, 1840:36. — FAUVEL, 1903:335.	
<i>Attagenus gloriosae</i> (FABRICIUS, 1798:76) LACORDAIRE, 1854:464.	
<i>Aethriostoma gloriosae</i> (FABRICIUS, 1798:76) MOTSCHULSKY, 1858b: 146.	
<i>Trogoderma subfasciatum</i> CHEVROLAT, 1863:617. — FAUVEL, 1903:335.	
<i>Attagenus subfasciatus</i> (CHEVROLAT, 1863:617) REITTER, 1881b:78.	
<i>Attagenus plebeius</i> SHARP in BLACKBURN et SHARP, 1885:147. — FAUVEL, 1903:335.	
<i>Attagenus gossypiatus</i> FAUVEL, 1903:335. — MROCKOWSKI, 1964: 180.	
<i>Attagenus fasciolatus</i> SOLSKIJ, 1876 : 275.	Caucasus, Turkmen SSR, Tadzhik SSR, Mongolia, Uzbek SSR, Kazakh SSR
Syn.: <i>Telopes fasciolatus</i> (SOLSKIJ, 1876:275) REITTER, 1889d:557.	
<i>Attagenus flexicollis</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus floralis</i> PÉRINGUEY, 1885 : 89.	South Africa
<i>Attagenus fruteti</i> GRIMMER, 1841 : 39.	Austria
<i>Attagenus fulvicollis</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus grandjeani</i> PIC, 1942b : 9.	Mozambique
<i>Attagenus grisezensis</i> PIC, 1937c : 5.	India: Bombay
<i>Attagenus hargreavesi</i> PIC, 1935c : 1. — KALIK, 1955b : 297.	Uganda, Congo (Kinshasa), Ethiopia?
Syn.: <i>Attagenus rufimembris</i> var. <i>hargreavesi</i> PIC, 1935c:1.	
<i>Attagenus hargreavesi</i> var. <i>angustegriseus</i> KALIK, 1955b : 303.	Congo (Kinshasa)
<i>Attagenus heydeni</i> (REITTER, 1881 : 35) DESBROCHERS DES LO- GES, 1898 : 73.	Morocco, Tanger, French West Africa
Syn.: <i>Telopes heydeni</i> REITTER, 1881:35.	
<i>Attagenus heydeni</i> var. <i>opacus</i> (PIC, 1946 : 6).	Egypt
Syn.: <i>Telopes heydeni</i> var. <i>opacus</i> PIC, 1946:6.	

<i>Attagenus heydeni</i> var. <i>rotroui</i> (PIC, 1927b : 5).	Oran
Syn.: <i>Telopes heydeni</i> var. <i>rotroui</i> PIC, 1927b:5.	
<i>Attagenus hirtulus</i> ROSENHAUER, 1856 : 109.	Spain, Portugal, Morocco, Tunisia
Syn.: <i>Telopes hirtulus</i> (ROSENHAUER, 1856:109) GEMMINGER et HAROLD, 1868:917.	
<i>Telopus hirtulus</i> (ROSENHAUER, 1856:109) MULSANT et REY, 1868: 104.	
<i>Attagenus hirtulus</i> var. <i>confusus</i> (PIC, 1951 : 5).	Spain
Syn.: <i>Telopes hirtulus</i> var. <i>confusus</i> PIC, 1951:5.	
<i>Attagenus hottentotus</i> (GUÉRIN MÉNEVILLE, 1838b : 65) ERICHSON, 1847b : 108.	South Africa
Syn.: <i>Megatoma hottentotum</i> GUÉRIN MÉNEVILLE, 1838b:65.	
<i>Attagenus laetus</i> PÉRINGUEY, 1885:88. — ARROW, 1915:427.	
<i>Attagenus inapicalis</i> PIC, 1951b : 13.	Tanganyika
<i>Attagenus incertus</i> MULSANT et REY, 1868 : 96.	Mediterranea
<i>Attagenus indicus</i> KALIK, 1954 : 368.	India
<i>Attagenus insignatus</i> PIC, 1942 : 7.	Ethiopia
<i>Attagenus irroratus</i> (THUNBERG, 1815 : 151) comb. nov. ¹	South Africa
Syn.: <i>Anthrenus irroratus</i> THUNBERG, 1815:151.	
<i>Attagenus jacobsoni</i> ZHANTIEV, 1963c : 417.	Tadzhik SSR
<i>Attagenus jucundus</i> PÉRINGUEY, 1885 : 88.	South Africa
<i>Attagenus kratochvili</i> KALIK, 1955b : 305.	Gold Coast, French West Africa
<i>Attagenus lategriseus</i> KALIK, 1955b : 304.	Congo (Kinshasa)
<i>Attagenus leopardinus</i> REITTER, 1881 : 30.	South Africa
<i>Attagenus leprieuri</i> (REITTER, 1887b : 53) DALLA TORRE, 1911 : 53.	Egypt
Syn.: <i>Telopes leprieuri</i> REITTER, 1887b:53.	
<i>Attagenus leprieuri</i> var. <i>aegyptiacus</i> (PIC, 1893 : 122).	Egypt
Syn.: <i>Telopes leprieuri</i> var. <i>aegyptiacus</i> PIC, 1893:122.	
<i>Attagenus lineatus</i> (PIC, 1894g : 71) DALLA TORRE, 1911 : 53.	Algeria
Syn.: <i>Telopes lineatus</i> PIC, 1894g:71.	

¹ The type-specimen was examined (preserved in Uppsala).

Attagenus lobatus ROSENHAUER, 1856 : 108.

Syn.: *Attagenus byturoides* SOLSKI, 1876:272. — ZHANTIEV, 1963c:415.

Attagenus sericeus: REITTER, 1881b:79 nec (GUÉRIN MÉNEVILLE, 1838 b:65).

Spain, Tunisia, Egypt, Greece, Bulgaria, Arabia, Iraq, Afghanistan, Uzbek SSR, Tadzhik SSR, Turkmen SSR, Mongolia

Attagenus longipennis PIC, 1904b : 33.

Syn.: *Attagenus longissimus* SUMAKOV, 1912:430, nomen nudum. — syn. nov.

Sinkiang,
Tadzhik SSR

Attagenus luteithorax PIC, 1931b : 3.

British East Africa

Attagenus luteofasciatus PIC, 1937b : 4.

East Africa

Attagenus lynx (MULSANT et REY, 1868 : 105) REITTER, 1881b : 81.

Israel, Syria, Caucasus, Iran, Turkmen SSR, Uzbek SSR, Kazakh SSR, Tadzhik SSR, Mongolia

Syn.: *Telopex lynx* MULSANT et REY, 1868:105.

Attagenus sareptanus REITTER, 1878b:178. — REITTER, 1881b:81.
Telopex lynx (MULSANT et REY, 1868:105) HEYDEN, 1881:99.

Madagascar

Attagenus madecassus (PIC, 1916b : 2) comb. nov.

Senegal

Syn.: *Pseudotelopex madecassus* PIC, 1916b:2.

Malta

Attagenus madoni PIC, 1942 : 7.

Sardinia, Sicily, Spain, Algeria, Tunisia

Attagenus maltensis PIC, 1894b : 66.

Ethiopia

Syn.: *Attagenus melitensis* WINKLER, 1926:677.

Mongolia, Kazakh SSR

Attagenus maritimus GENÉ, 1839 : 20.

Canary Islands

Syn.: *Telopex maritimus* (GENÉ, 1839:20) GEMMINGER et HAROLD, 1868: 917.

Telopex maritimus (GENÉ, 1839:20) MULSANT et REY, 1868:102.

Attagenus metallicus REITTER, 1881 : 33.

India or.

Attagenus molitor REITTER, 1889d : 557.

China

Attagenus multifasciatus (WOLLASTON, 1863 : 218) DALLA TORRE, 1911 : 54.

Natal

Syn.: *Telopex multifasciatus* WOLLASTON, 1863:218.

Gold Coast, Congo (Kinshasa)

Attagenus multinotatus PIC, 1916c : 15.

Attagenus nankineus PIC, 1916c : 15.

Attagenus natalensis PIC, 1927e : 2.

Attagenus nigripennis ARROW, 1915 : 428.

Attagenus nigripes CASEY, 1916 : 184.

Attagenus nigroapicalis PIC, 1931b : 3.

Attagenus nigroluteus KALIK, 1955b : 308.

Attagenus obtusus (GYLLENHAL in SCHÖNHERR, 1808 : 88) DE-
JEAN, 1821 : 46.

Syn.: *Dermestes obtusus* GYLLENHAL in SCHÖNHERR, 1808 : 88.

Dermestes holosericeus SAHLBERG, 1819:52. — GEMMINGER et HA-
ROLD, 1868:917.

Trogoderma holosericea (SAHLBERG, 1819:52) SCHAUM, 1852:38.

Telopes obtusus (GYLLENHAL in SCHÖNHERR, 1808:88) WOLLA-
STON, 1864:157.

Telopus obtusus (GYLLENHAL in SCHÖNHERR, 1808:88) MULSANT
et REY, 1868:104.

Attagenus obtusus var. *dispar* (REDTENBACHER, 1843 :
976). — REITTER, 1891 : 169.

Syn.: *Telopes dispar* REDTENBACHER, 1843:976.

Attagenus dispar (REDTENBACHER, 1843:976) LACORDAIRE,
1854:464.

Attagenus redtenbacheri PEYRON, 1857:720.

Telopes redtenbacheri (PEYRON, 1857:720) GEMMINGER et HA-
ROLD, 1868:917.

Telopus redtenbacheri (PEYRON, 1857:720) MULSANT et REY,
1868:101.

Telopus dispar (REDTENBACHER, 1843:976) MULSANT et REY,
1868:103.

Telopes obtusus var. *dispar* (REDTENBACHER, 1843:976)
SAHLBERG, 1903d:19.

Attagenus orientalis REITTER, 1878b : 178.

Syn.: *Telopes orientalis* (REITTER, 1878b:178) REITTER, 1887b:52.

Attagenus pantherinus (AHRENS, 1814 : 11) ERICHSON, 1846 :
442.

Syn.: *Dermestes pantherinus* AHRENS, 1814:11.

Trogoderma pantherinum (AHRENS, 1814:11) SCHMIDT, 1844:31.

Megatoma pantherina (AHRENS, 1814:11) REITTER, 1887b:48.

Attagenus pardus ARROW, 1915 : 427.

Attagenus pellio (LINNAEUS, 1758 : 355) LEACH, 1815 : 94.

Syn.: *Dermestes pellio* LINNAEUS, 1758:355.

Dermestes bipunctatus DEGEER, 1774:197. — DEGEER, 1774:198.

Dermestes macellarius FABRICIUS, 1781:63. — ERICHSON, 1846:
440.

Dermestes cylindricornis SCHRANK, 1785:315. — ILLIGER, 1798:
316.

California

British East
Africa, Uganda

Congo
(Kinshasa)

Portugal,
entire
Mediterranea,
Canary
Islands,
Caucasus,
Iran, Sweden

Syria, Iran,
Greece

Caucasus,
Syria, Iraq,
Tunisia

Germany,
Middle
Europa, Spain,
Caucasus

North Rhodesia
Europe,
Cosmopolitan

<i>Megatoma atra</i> HERBST, 1792:95. — ILLIGER, 1798:316.	
<i>Megatoma schrankii</i> KUGELANN, 1792:480. — ILLIGER, 1798:316.	
<i>Dermestes ater</i> (HERBST, 1792:95) PANZER, 1795:96.	
<i>Attagenus macellarius</i> (FABRICIUS, 1781:63) LATREILLE, 1802:121.	
<i>Megatoma pellio</i> (LINNAEUS, 1758:355) CRISTOFORI et JAN, 1832: 30.	
<i>Attagenus bipunctatus</i> (DEGEER, 1774:197) WRADATSCH, 1914:152.	
<i>Attagenus pellio</i> var. <i>pilosissimus</i> ROUBAL, 1932 : 66.	Monte Gargano
<i>Attagenus pici</i> MROCKOWSKI, 1958 : 558.	Madagascar
Syn.: <i>Aethriostoma albonotata</i> PIC, 1927e:3.	
<i>Attagenus albonotatus</i> (PIC, 1927e:3) nec <i>Attagenus albonotatus</i> PIC, 1927e:2.	
<i>Attagenus pici</i> var. <i>atripennis</i> (PIC, 1927e : 3).	Madagascar
Syn.: <i>Aethriostoma albonotata</i> var. <i>atripennis</i> PIC, 1927e:3.	
<i>Attagenus picipennis</i> PIC, 1894b : 66.	Sicily
<i>Attagenus pictus</i> BALLION, 1871 : 330.	Tadzhik SSR, Uzbek SSR, Turkmen SSR, Kazakh SSR, Afghanistan, North Iran
<i>Attagenus placitus</i> NORMAND, 1950 : 72.	Tunisia
<i>Attagenus postfasciatus</i> (PIC, 1951 : 5) KALIK, 1955b : 315.	Senegal
Syn.: <i>Telopes robustior</i> var. <i>postfasciatus</i> PIC, 1951:5.	
<i>Attagenus posticalis</i> FAIRMAIRE, 1879 : 169.	Algeria, Spain, Yemen, Tunisia, Egypt, Syria, French West Africa
Syn.: <i>Telopes posticalis</i> (FAIRMAIRE, 1879:169) REITTER, 1887b:52.	
<i>Attagenus posticalis</i> var. <i>brunneonotatus</i> (PIC, 1894f : 60) DALLA TORRE, 1911 : 56.	Algeria
Syn.: <i>Telopes posticalis</i> v. <i>brunneonotatus</i> PIC, 1894f:60.	
<i>Attagenus bifasciatus</i> var. <i>brunneonotatus</i> : NORMAND, 1936: 297.	
<i>Attagenus posticalis</i> var. <i>subelongatus</i> (PIC, 1929 : 9).	Egypt
Syn.: <i>Telopes brunneonotatus</i> var. <i>subelongatus</i> PIC, 1929:9.	
<i>Attagenus prescutellaris</i> PIC, 1927e : 2.	South Africa
<i>Attagenus pseudomolitor</i> ZHANTIEV, 1963 : 454.	Uzbek SSR
<i>Attagenus pubescens</i> (PIC, 1894b : 66) DALLA TORRE, 1911 : 56.	Egypt
Syn.: <i>Telopes pubescens</i> PIC, 1894b:66.	

Attagenus punctatus (SCOPOLI, 1772 : 87) REITTER, 1906 : 378.

Syn.: *Dermestes punctatus* SCOPOLI, 1772:87.

Dermestes vigintiguttatus FABRICIUS, 1775:56. — SCHÖNHERR, 1808:89.

Dermestes quadripunctatus FUESSLIN, 1775:4. — FABRICIUS, 1781: 64.

Dermestes multiguttatus SCHRANK, 1798:427. — SCHRANK, 1798: 427.

Attagenus vigintiguttatus (FABRICIUS, 1775:56) DEJEAN, 1821:46.

Megatoma vigintipunctatum CRISTOFORI et JAN, 1832:30, nomen nudum.

Megatoma vigintiguttata (FABRICIUS, 1775:56) REITTER, 1887b:47.

Attagenus vigintipunctatus HABERFELNER, 1889:390, syn. nov.

Lanorus vigintiguttatus (FABRICIUS, 1775:56) COUCKE, 1892:73.

Attagenus purpureus (PIC, 1915b : 3) comb. nov.

Syn.: *Aethriostoma purpurea* PIC, 1915b:3.

Attagenus pustulatus (THUNBERG, 1815 : 151) SCOTT, 1926 : 62

(and KALIK in litt., who has examined the type specimens).

Syn.: *Anthrenus pustulatus* THUNBERG, 1815:151.

Attagenus quadricolor (SUMAKOV, 1907 : 15) MROCKOWSKI, 1960 b : 212.

Syn.: *Trogoderma quadricolor* SUMAKOV, 1907:15.

Phradonoma quadricolor (SUMAKOV, 1907:15) DALLA TORRE, 1911: 66.

Attagenus soederbomi PIC, 1935d:2. — MROCKOWSKI, 1960b: 212.

Attagenus quadrimaculatus KRAATZ, 1858 : 145.

Syn.: *Attagenus bivittatus* MULSANT et REY, 1868:97. — REITTER, 1881b:78.

Megatoma quadrimaculata (KRAATZ, 1858:145) REITTER, 1887b: 50.

Attagenus quadrinotatus PIC, 1938b : 10.

Attagenus quadritinctus REITTER, 1889c : 280.

Attagenus reitteri nom. nov.

Syn.: *Telopes antennatus* REITTER, 1887b:54.

Attagenus antennatus (REITTER, 1887b:54) DALLA TORRE, 1911: 51, nec CASTELNAU, 1840:36.

Attagenus reitteri var. *tessellatus* (REITTER, 1887b : 54)
comb. nov.

Syn.: *Telopes antennatus* var. *tessellatus* REITTER, 1887b:54.

Telopes tessellatus REITTER, 1887b:54. — SAHLBERG, 1903c:12.

Carniola,
Middle and
North Europa,
Caucasus

Madagascar

South Africa

Turkmen SSR,
Uzbek SSR,
Tadzhik SSR,
Mongolia

Greece,
Yugoslavia,
Caucasus,
Turkey,
France

Yunnan

Nakhitshevan ASSR,
Armenian
SSR, Iran

Algeria,
Tunisia,
Morocco

Syria, Israel,
Egypt

<i>Attagenus rhodesianus</i> PIC, 1927e : 2.	Rhodesia, Tanganyika
<i>Attagenus robustior</i> (PIC, 1951 : 5) KALIK, 1955 : 97. Syn.: <i>Telopes robustior</i> PIC, 1951:5.	Senegal, Congo (Kinshasa), Tanganyika
<i>Attagenus robustus</i> PIC, 1899c : 253.	Jordan
<i>Attagenus ruficolor</i> PIC, 1918b : 1.	Yunnan
<i>Attagenus rufimembris</i> PIC, 1927c : 246.	Zanzibar, Uganda
<i>Attagenus rufipennis</i> LECONTE, 1859 : 71.	California to Utah
<i>Attagenus rufiventris</i> PIC, 1927c : 246.	South Africa, Uganda
<i>Attagenus rufiventris</i> var. <i>conradsi</i> PIC, 1951b : 13.	Tanganyika
<i>Attagenus rufomaculatus</i> (PIC, 1907c : 274) DALLA TORRE, 1911 : 57. Syn.: <i>Telopes rufomaculatus</i> PIC, 1907c:274.	Algeria
<i>Attagenus scalaris</i> (PIC, 1894f : 60) DALLA TORRE, 1911 : 57. Syn.: <i>Telopes scalaris</i> PIC, 1894f:60.	Egypt
<i>Attagenus schaefferi</i> (HERBST, 1792 : 93) DEJEAN, 1821 : 46. Syn.: <i>Megatoma schaefferi</i> HERBST, 1792:93. <i>Dermestes schaefferi</i> (HERBST, 1792:93) PANZER, 1795:95. <i>Attagenus dichrous</i> LECONTE, 1854:110. — GEMMINGER et HAROLD, 1868:915. <i>Attagenus spucus</i> LECONTE, 1854:109. — CROTCH, 1873:41. <i>Attagenus bicolor</i> HAROLD, 1868:104. — REITTER, 1881:34.	Germany, nearly entire Holarctic Region
<i>Attagenus seminiger</i> FAIRMAIRE, 1863 : 639. Syn.: <i>Telopes seminiger</i> (FAIRMAIRE, 1863:639) REITTER, 1887b:52.	Algeria, Tunisia
<i>Attagenus senegalensis</i> (PIC, 1915b : 2) comb. nov. Syn.: <i>Telopes senegalensis</i> PIC, 1915b:2.	Senegal
<i>Attagenus seniculus</i> (SOLSKIJ, 1876 : 276) REITTER, 1881 : 79. Syn.: <i>Telopes seniculus</i> SOLSKIJ, 1876:276.	Turkmen SSR, Iran, Afghanistan, Tadzhik SSR, Caucasus, Mongolia
<i>Attagenus sericeus</i> (GUÉRIN MÉNEVILLE, 1838b : 65) ERICHSON, 1847b : 108. Syn.: <i>Megatoma sericeum</i> GUÉRIN MÉNEVILLE, 1838b:65.	Egypt, Algeria

<i>Attagenus sexnotatus</i> PIC, 1927e : 3.	West Africa
<i>Attagenus sieversi</i> REITTER, 1896 : 286.	Turkmen SSR, Iran
<i>Attagenus sikkimensis</i> PIC, 1942 : 7.	India
<i>Attagenus similaris</i> MULSANT et REY, 1868 : 85.	Algeria
Syn.: <i>Attagenus trifasciatus</i> var. <i>similaris</i> : SAHLBERG, 1903e:34.	
<i>Attagenus simplex</i> REITTER, 1881b : 79.	Italy, Sicily, Libya, Tunisia, Morocco, Israel
Syn.: <i>Attagenus rufipennis</i> MULSANT et REY, 1868:100 nec LECONTE, 1859:71.	
<i>Megatoma simplex</i> (REITTER, 1881b:79) REITTER, 1887b:51.	
<i>Attagenus simulans</i> SOLSKIJ, 1876 : 272.	Uzbek SSR, Turkmen SSR, Tadzhik SSR, Sinkiang
<i>Attagenus sinensis</i> PIC, 1927e : 3.	China
† <i>Attagenus sopitus</i> SCUDDER, 1900 : 85.	Colorado (Miocene)
<i>Attagenus sparsus</i> CASEY, 1916 : 184.	New Mexico
<i>Attagenus sparsutus</i> (REITTER, 1881 : 36) comb. nov.	China,
Syn.: <i>Aethriostoma sparsuta</i> REITTER, 1881:36.	Himalaya
<i>Attagenus sparsutus</i> var. <i>rufescens</i> (PIC, 1933 : 71) comb. nov.	China
Syn.: <i>Aethriostoma sparsuta</i> var. <i>rufescens</i> PIC, 1933:71.	
<i>Attagenus stachi</i> MROCKOWSKI, 1958 : 557.	Sumatra
Syn.: <i>Aethriostoma multifasciata</i> PIC, 1927e:3.	
<i>Attagenus multifasciatus</i> (PIC, 1927e:3) nec (WOLLASTON, 1864: 159).	
<i>Attagenus steinbergi</i> ZHANTIEV, 1963c : 418.	Turkmen SSR, Iran
<i>Attagenus subundulatus</i> PIC, 1927e : 2.	West Africa
<i>Attagenus suspiciosus</i> SOLSKIJ, 1876 : 274.	Uzbek SSR, Tadzhik SSR, Kazakh SSR, Turkmen SSR, Iran, Afghanistan, Kirghiz SSR, Sinkiang
<i>Attagenus thunbergi</i> KALIK in litt., nom. nov.	South Africa
Syn.: <i>Anthrenus bifasciatus</i> THUNBERG, 1815:151.	
<i>Attagenus bifasciatus</i> (THUNBERG, 1815:151) comb. nov. KALIK in litt., nec (OLIVIER, 1790:no. 9 p. 13).	

Attagenus trifasciatus (FABRICIUS, 1787 : 34) LATREILLE, 1802 : 121.

Syn.: *Dermestes trifasciatus* FABRICIUS, 1787:34.

Attagenus repandus DEJEAN, 1821:46, nomen nudum. — ROSENHAUER, 1856:110.

Megatoma trifasciatum (FABRICIUS, 1787:34) CRISTOFORI et JAN, 1832:30.

Attagenus verbasci JACQUELIN DU VAL, 1859:254. — REITTER, 1891:169.

Attagenus verbasci var. *repandus* DEJEAN, 1821:46. — GEMMINGER et HAROLD, 1868:916.

Lanorus trifasciatus (FABRICIUS, 1787:34) COUCKE, 1892:74.

South Europe,
entire
Mediterranea

Attagenus trifasciatus var. *barbieri* PIC, 1946 : 6.

Egypt

Attagenus trifasciatus var. *subtrifasciatus* FUENTE, 1930 : 49, nomen nudum.

Spain

Attagenus turcomanus ZHANTIEV, 1963 : 455.

Turkmen SSR

Attagenus undulatus (MOTSCHULSKY, 1858 : 47) ARROW, 1915 : 426.

East Indies,
Madagascar,
Mauritius,
India, Ceylon,
Viet-Nam,
Philippine Is.,
Sumatra,
North
Queensland,
S. Mariana Is.,
Hawaiian Is.

Syn.: *Aethriostoma undulata* MOTSCHULSKY, 1858:47.

Attagenus rufipes WALKER, 1859:53. — ARROW, 1915:426.

Aethriostoma rufipes (WALKER, 1859:53) MOTSCHULSKY, 1863 : 433.

Brachysphyrus irroratus BLACKBURN, 1903:160. — ARROW, 1915 : 426.

Pseudotelopessimoni PIC, 1916b:2. — ARROW, 1937:112.

Pseudotelopessetaceipes PIC, 1916b:2. — BEAL, 1961:121.

Telopes undulata (MOTSCHULSKY, 1858:47) FULLAWAY, 1922:77.

Attagenus unicolor (BRAHM, 1791 : 144) comb. nov.

Germany,
cosmopolitan

Syn.: *Dermestes piceus* OLIVIER, 1790:no. 9 p. 10 nec THUNBERG, 1781:8.

Dermestes unicolor BRAHM, 1791:144.

Megatoma brevicornis HERBST, 1792:95. — SCHÖNHERR, 1808:86.

Dermestes brevicornis (HERBST, 1792:95) PANZER, 1795:96.

Dermestes megatoma FABRICIUS, 1798:71. — SCHÖNHERR, 1808 : 85.

Nitidula cylindricornis SCHRANK, 1798:446. — SCHRANK, 1798 : 446.

Attagenus piceus (OLIVIER, 1790:no. 9 p. 10) LATREILLE, 1804:244.

Attagenus megatoma (FABRICIUS, 1798:71) LATREILLE, 1804:244.

Dermestes macellarius DUFTSCHMID, 1825:39. — ERICHSON, 1846 : 441.

Attagenus megadoma STURM, 1826:98.

Attagenus stygialis MULSANT et REY, 1868:73. — FAUVEL, 1903 : 336.

Megatoma picea (OLIVIER, 1790:no. 9 p. 10) REITTER, 1887b:46.

Attagenus piceus var. *megatoma*: REITTER, 1906:378.

Attagenus unicolor japonicus REITTER, 1877 : 375.

Syn.: *Attagenus japonicus* REITTER, 1877:375.

Attagenus piceus japonicus REITTER, 1877:375. — ARROW, 1915:427.

Attagenus megatoma japonicus REITTER, 1877:375. — MROCKOWSKI, 1965b:104.

Japan, Korea,
East Asia,
Mongolia

Attagenus unifasciatus PIC, 1942 : 7.

Attagenus uniformis FAIRMAIRE in FAIRMAIRE et COQUEREL, 1860 : 169.

South Africa
Algeria,
Morocco,
Tunisia, Sicily

Syn.: *Telopes uniformis* (FAIRMAIRE in FAIRMAIRE et COQUEREL, 1860: 169) REITTER, 1887b:52.

Telopus uniformis (FAIRMAIRE in FAIRMAIRE et COQUEREL, 1860: 169) MULSANT et REY, 1868:101.

Attagenus vagepictus FAIRMAIRE, 1889 : 13.

Tibet

Attagenus vestitus KLUG, 1855 : 649.

Mozambique

Attagenus wollastoni MROCKOWSKI, 1964 : 180.

Canary Is.

Syn.: *Telopes fasciatus* WOLLASTON, 1863:218.

Attagenus fasciatus (WOLLASTON, 1863:218) DALLA TORRE, 1911:
53 nec (THUNBERG, 1795:105).

Attagenus zavattarii PIC, 1952b : 29.

Ethiopia

Aethriostoma albonotata PIC, 1927e:3 — vide *Attagenus pici* MROCKOWSKI, 1958:558.

Aethriostoma albonotata var. *atripennis* PIC, 1927e:3 — vide *Attagenus pici* var. *atripennis* (PIC, 1927e:3).

Attagenus alfieri PIC, 1910:17 — vide *Attagenus cyphonoides* REITTER, 1881:34.

Attagenus angularis MANNERHEIM, 1853:216 — vide *Megatoma cylindrica* var. *angularis* (MANNERHEIM, 1853:216).

Attagenus angustatus SOLSKIJ, 1876:271 — vide *Attagenus augustatus* BALLION, 1871:330.

Attagenus annulatus DEJEAN, 1821:46, nomen nudum — vide *Attagenus fasciatus* (THUNBERG, 1795:105).

Attagenus annulifer CASTELNAU, 1840:36 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).

Attagenus antennatus ESCALERA, 1914:137 nec CASTELNAU, 1840:36 — vide *Attagenus escalerai* MADER, 1954:53.

Telopes antennatus REITTER, 1887b:54 — vide *Attagenus reitteri* nom. nov.

Attagenus antennatus var. *tessellatus* REITTER, 1887b:54 — vide *Attagenus reitteri* var. *tessellatus* (REITTER, 1887b:54).

Aethriostoma atricolor PIC, 1915b:2 — vide *Thaumaglossa rufocapillata* REDTENBACHER, 1867:44.

Attagenus bicolor HAROLD, 1868:104 — vide *Attagenus schaefferi* (HERBST, 1792:93).

Attagenus bifasciatus: KORGE, 1961:25 — vide *Trogoderma angustum* (SOLIER, 1849:374).

Attagenus bifasciatus (ROSSI, 1794:79) SCHMIDT, 1844:31 — vide *Attagenus bifasciatus* var. *rossi* GANGLBauer, 1904b:26.

Attagenus bifasciatus (THUNBERG, 1815:151) comb. nov. — vide *Attagenus thunbergi* KALIK in litt., nom. nov.

Attagenus bifasciatus var. *brunneonotatus*: NORMAND, 1936:297 — vide *Attagenus posticalis* var. *brunneonotatus* PIC, 1894f:60.

Attagenus bifasciatus ab. *wegneri* NORMAND, 1936:297 — vide *Attagenus bifasciatus* var. *wegneri* PIC, 1899c:252.

- Attagenus bipunctatus* (DEGEER, 1774:197) WRADATSCH, 1914:152 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Attagenus bivittatus* MULSANT et REY, 1868:97 — vide *Attagenus quadrimaculatus* KRAATZ, 1858:145.
- Telopes brunneonotatus* var. *subelongatus* PIC, 1929:9 — vide *Attagenus posticalis* var. *subelongatus* (PIC, 1929:9).
- Attagenus byturoides* CROTH, 1873:41, nomen nudum — vide *Novelsis uteana* CASEY, 1900:148.
- Attagenus byturoides* SOLSKIJ, 1876:272 — vide *Attagenus lobatus* ROSENHAUER, 1856:108.
- Attagenus capensis* DEJEAN, 1837:139 — nomen nudum.
- Attagenus cayennensis* DEJEAN, 1821:46 — nomen nudum.
- Attagenus cylindricollis* MELSHEIMER, 1853:48 — vide *Attagenus cylindricornis* SAY, 1825:185.
- Attagenus cylindricus* KIRBY, 1837:113 — vide *Megatoma cylindrica* (KIRBY, 1837:113).
- Attagenus dalmatinus* DEJEAN, 1821:46, nomen nudum — vide *Attagenus brunneus* FALDERMANN, 1835: 227.
- Attagenus dalmatinus* KÜSTER, 1847:44 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus? defectus* WALKER, 1858:207 — vide *Orphinus defectus* (WALKER, 1858:207).
- Attagenus dichrous* LECONTE, 1854:110 — vide *Attagenus schaefferi* (HERBST, 1792:93).
- Telopes dispar* REDTENBACHER, 1843:976 — vide *Attagenus obtusus* var. *dispar* (REDTENBACHER, 1843:976).
- Attagenus distinctus* MULSANT et REY, 1868:85 — vide *Attagenus fallax* var. *distinctus* MULSANT et REY, 1868:85.
- Attagenus diversepubescens* PIC, 1953:112 — vide *Attagenus africanus* MROCZKOWSKI, 1958:558.
- Attagenus elongatus* CASEY, 1916:183 — vide *Attagenus elongatus* CASEY, 1900:147.
- Attagenus emarginatus* (GYLLENHAL, 1808:153) STURM, 1826:98 — vide *Globicornis marginata* (PAY-KULL, 1798:280).
- Attagenus exilis* (MULSANT et REY, 1868:158) DALLA TORRE, 1911:53 — vide *Anthrenus exilis* MULSANT et REY, 1868:158.
- Telopes fasciatus* WOLLASTON, 1863:218 — vide *Attagenus wollastoni* MROCZKOWSKI, 1964:180.
- Aethriostoma fasciolata* FAIRMAIRE, 1897:97 — vide *Attagenus fairmairei* MROCZKOWSKI, 1958:558.
- Attagenus flavigornis* DEJEAN, 1821:46 — nomen nudum.
- Attagenus fulvipes* MULSANT et REY, 1868:71 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus fulvipes* var. *obliquus*: PIC, 1895b:116 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus fuscus* DEJEAN, 1837:139, nomen nudum — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Attagenus fuscus* GEBLER, 1841:585 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Attagenus globicola* DALLA TORRE, 1911:53 — vide *Attagenus augustatus globicola* FRIVALDSZKY, 1892:122.
- Attagenus gloriosae* (FABRICIUS, 1798:76) LACORDAIRE, 1854:464 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
- Attagenus gobicola* FRIVALDSZKY, 1892:122 — vide *Attagenus augustatus gobicola* FRIVALDSZKY, 1892:122.
- Attagenus gossypiatus* FAUVEL, 1903:335 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
- Attagenus granarius* DEJEAN, 1837:139 — nomen nudum.
- Attagenus hirtus* STURM, 1826:98 — nomen nudum.
- Attagenus hornii* CROTCH, 1873:41, nomen nudum — vide *Novelsis horni* (JAYNE, 1882:345).
- Attagenus hornii* JAYNE, 1882:345 — vide *Novelsis horni* (JAYNE, 1882:345).
- Brachysphyrus irroratus* BLACKBURN, 1903:160 — vide *Attagenus undulatus* (MOTSCHULSKY, 1858:47).
- Attagenus isabellinus* (KÜSTER, 1848:38) FAIRMAIRE et COQUEREL, 1860:170 — vide *Anthrenus pimpinellae* var. *isabellinus* KÜSTER, 1848:38.
- Attagenus japonicus* REITTER, 1877:375 — vide *Attagenus unicolor japonicus* REITTER, 1877:375.
- Attagenus laetus* PÉRINGERUEY, 1885:88 — vide *Attagenus hottentotus* (GUÉRIN MÉNEVILLE, 1838b:65).
- Attagenus liliputianus* (MULSANT et REY, 1868:159) DALLA TORRE, 1911:53 — vide *Anthrenus liliputianus* MULSANT et REY, 1868:159.
- Attagenus longicornis* PIC, 1894b:66 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus longissimus* SUMAKOV, 1912:430, nomen nudum — vide *Attagenus longipennis* PIC, 1904b:33.
- Attagenus longulus* (LECONTE, 1863:73) CROTCH, 1873:41 — vide *Dearthrus longulus* LECONTE, 1863:73.
- Attagenus luctuosus* DEJEAN, 1837:139 — nomen nudum.

- Attagenus macellarius* (FABRICIUS, 1781:63) LATREILLE, 1802:121 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Attagenus marginatus* (PAYKULL, 1798:280) DEJEAN, 1821:46 — vide *Globicornis marginata* (PAYKULL, 1798:280).
- Attagenus marginicollis* KÜSTER, 1847b:76 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus megadoma* STURM, 1826:98 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Attagenus megaloma* (FABRICIUS, 1798:71) LATREILLE, 1804:244 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Attagenus megaloma dalmatinus*: MROCKOWSKI, 1962b:547 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus melanocerus* DEJEAN, 1837:139 — nomen nudum.
- Attagenus melitensis* WINKLER, 1926:677 — vide *Attagenus maltensis* PIC, 1894b:66.
- Aethriostoma multifasciata* PIC, 1927e:3 — vide *Attagenus stachi* MROCKOWSKI, 1958:557.
- Attagenus nigripes* (FABRICIUS, 1792:233) DEJEAN, 1821:46 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Attagenus niseteoi* DEJEAN, 1821:46 — nomen nudum.
- Attagenus ornatus* (SAY, 1825:185) MELSHEIMER, 1853:48 — vide *Trogoderma ornatum* (SAY, 1825:185).
- Attagenus perplexus* JAYNE, 1882:345 — vide *Novelsis perplexa* (JAYNE, 1882:345).
- Attagenus piceus* (OLIVIER, 1790:no. 9 p. 10) LATREILLE, 1804:244 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Attagenus piceus* var. *brunneus*: REITTER, 1883:89 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus piceus* var. *dalmatinus*: REITTER, 1881b:76 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus piceus* var. *megatoma*: REITTER, 1906:378 — vide *Attagenus unicolor* (BRAHM, 1791:144).
- Attagenus piceus* var. *obliquus* REY, 1889:59 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus piceus* var. *sordidus*: REITTER, 1881b:76 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Attagenus plebeius* SHARP in BLACKBURN et SHARP, 1885:147 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
- Attagenus poecilus* GERMAR, 1839:9 — vide *Attagenus bifasciatus* var. *tigrinus* (FABRICIUS, 1792:229).
- Attagenus pulcher* FALDERMANN, 1835:226 — vide *Attagenus bifasciatus* (OLIVIER, 1790:no. 9 p. 13).
- Attagenus redtenbacheri* PEYRON, 1857:720 — vide *Attagenus obtusus* var. *dispar* (REDTENBACHER, 1843:984).
- Attagenus repandus* DEJEAN, 1821:46, nomen nudum — vide *Attagenus trifasciatus* (FABRICIUS, 1787:34).
- Attagenus reyi* DESBROCHERS DES LOGES, 1898:73 — nomen nudum.
- Telopes robustior* var. *postfasciatus* PIC, 1951:5 — vide *Attagenus postfasciatus* (PIC, 1951:5).
- Attagenus ruficollis* CRISTOFORI et JAN, 1832:30 — nomen nudum.
- Attagenus rufimembris* var. *hargreavesi* PIC, 1935c:1 — vide *Attagenus hargreavesi* PIC, 1935c:1.
- Attagenus rufipennis* MULSANT et REY, 1868:100 — vide *Attagenus simplex* REITTER, 1881b:79.
- Attagenus rufipes* STURM, 1843:93 — nomen nudum.
- Attagenus rufipes* WALKER, 1859:53 — vide *Attagenus undulatus* (MOTSCHELSKY, 1858:47).
- Aethriostoma rufofasciata* PIC, 1915:2 — vide *Phradonomorphus rufofasciatus* (PIC, 1915:2).
- Aethriostoma rufofasciata* var. *postlimbatus* PIC, 1948:14 — vide *Phradonomorphus rufofasciatus* var. *postlimbatus* (PIC, 1948:14).
- Aethriostoma rufomaculata* PIC, 1938c:12 — vide *Thaumaglossa laeta* ARROW, 1915:435.
- Attagenus sareptanus* REITTER, 1878b:178 — vide *Attagenus lynx* (MULSANT et REY, 1868:105).
- Attagenus sericeus*: REITTER, 1881b:79 — vide *Attagenus lobatus* ROSENHAUER, 1856:108.
- Attagenus serra* (FABRICIUS, 1792:234) LATREILLE, 1804:244 — vide *Ctesias serra* (FABRICIUS, 1792:234).
- Attagenus sexguttatus* STURM, 1843:94 — vide *Attagenus fallax* GENÉ, 1839:19.
- Attagenus siculus* REITTER, 1881b:76 — vide *Globicornis variegata* (KÜSTER, 1851:40).
- Attagenus signatus* DEJEAN, 1837:139 — nomen nudum.
- Pseudotelopes simoni* PIC, 1916b:2 — vide *Attagenus undulatus* (MOTSCHELSKY, 1858:47).
- Attagenus simonis* REITTER, 1881:34 — vide *Attagenus bifasciatus* var. *simonis* REITTER, 1881:34.
- Attagenus soederbomi* PIC, 1935d:2 — vide *Attagenus quadricolor* (SUMAKOV, 1907:15).
- Attagenus sordidus* HEER, 1841b:438 — vide *Attagenus brunneus* FALDERMANN, 1835:227.

- Attagenus spurcus* LECONTE, 1854:109 — vide *Attagenus schaefferi* (HERBST, 1792:93).
Attagenus stygialis MULSANT et REY, 1868:73 — vide *Attagenus unicolor* (BRAHM, 1791:144).
Attagenus subfasciatus (CHEVROLAT, 1863:617) REITTER, 1881b:78 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
Attagenus syriacus DEJEAN, 1837:139 — nomen nudum.
Attagenus taeniatus DEJEAN, 1837:139 — nomen nudum.
Telopes tessellatus REITTER, 1887b:54 — vide *Attagenus reitteri* var. *tessellatus* (REITTER, 1887b:54).
Pseudotelopes testaceipes PIC, 1916b:2 — vide *Attagenus undulatus* (MOTSCHULSKY, 1858:47).
Attagenus tigrinus (FABRICIUS, 1792:229) SCHAUM, 1852:38 — vide *Attagenus bifasciatus* var. *tigrinus* (FABRICIUS, 1792:229).
Attagenus tigrinus var. *persicus* REITTER, 1881b:78 — vide *Attagenus bifasciatus* var. *persicus* REITTER, 1881 b:78.
Attagenus tigrinus var. *vancharui* DESBROCHERS DES LOGES, 1898:73 — vide *Attagenus bifasciatus* (OLIVIER, 1790:no. 9 p. 13).
Attagenus tigrinus var. *wegeneri* PIC, 1899c:252 — vide *Attagenus bifasciatus* var. *wegeneri* PIC, 1899c: 252.
Attagenus tomentosus STURM, 1843:94 — nomen nudum.
Attagenus trifasciatus var. *similaris*: SAHLBERG, 1903e:34 — vide *Attagenus similaris* MULSANT et REY, 1868:85.
Attagenus undatus (LINNAEUS, 1758:355) LATREILLE, 1804:243 — vide *Megatoma undata* (LINNAEUS, 1758:355).
Attagenus undulatus (LINNAEUS, 1761:141) STEPHENS, 1829:8 — vide *Megatoma undata* (LINNAEUS, 1758: 355).
Attagenus unifasciatus FAIRMAIRE et COQUEREL, 1860:168 — vide *Attagenus cinnamomeus* ROTH, 1851:122.
Attagenus unifasciatus STURM, 1843:94 — nomen nudum.
Telopes uninotatus PIC, 1942:6 — vide *Phradonoma uninotata* (PIC, 1942:6).
Attagenus varicolor CROTCH, 1873:41, nomen nudum — vide *Novelsis varicolor* (JAYNE, 1882:345).
Attagenus varicolor JAYNE, 1882:345 — vide *Novelsis varicolor* (JAYNE, 1882:345).
Attagenus verbasci JACQUELIN DU VAL, 1859:254 — vide *Attagenus trifasciatus* (FABRICIUS, 1787:34).
Attagenus verbasci var. *repandus* DEJEAN, 1821:46 — vide *Attagenus trifasciatus* (FABRICIUS, 1787:34).
Attagenus vestitus DEJEAN, 1837:139 — nomen nudum.
Attagenus vestitus STURM, 1843:94 — nomen nudum.
Attagenus vigintiguttatus (FABRICIUS, 1775:56) DEJEAN, 1821:46 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
Attagenus vigintipunctatus HABERFELNER, 1889:390 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
Attagenus wachanrui MULSANT et REY, 1868:91 — vide *Attagenus bifasciatus* (OLIVIER, 1790:no. 9 p. 13).
Attagenus xanthocerus DEJEAN, 1837:139 — nomen nudum.

Genus: *Novelsis* CASEY, 1900 : 144

Type species: *Attagenus hornii* JAYNE, 1882 : 345.

Syn.: *Genattus* SHARP, 1902:646. — BEAL, 1954b:75. Type species: *Genattus aequalis* SHARP, 1902:646.

Subgenus: *Novelsis* s. str.

List of species:

Novelsis andersoni BEAL, 1954b : 76.

California,
Arizona,
Lower
California

Novelsis athlophora BEAL, 1954b : 75.

Idaho,
Washington

Novelsis horni (JAYNE, 1882 : 345) CASEY, 1900 : 148.

Pacific States
but only Texas,
New Mexico,
Arizona,
California.
Mexico:
Chihuahua,
Sonora

Syn.: *Attagenus hornii* CROTCH, 1873:41, nomen nudum.

Attagenus hornii JAYNE, 1882:345.

Novelsis picta CASEY, 1900 : 149.

Arizona,
California

Novelsis timia BEAL, 1954b : 76.

California,
Arizona

Novelsis uteana CASEY, 1900 : 148.

Utah, Arizona,
California,
Nevada,
New Mexico,
Texas, Lower
California

Syn.: *Attagenus byturoides* CROTCH, 1873:41, nomen nudum.

Novelsis byturoides (CROTCH, 1873:41) CASEY, 1900:148. — BEAL,
1954b:82.

Subgenus: *Paranovelsis* CASEY, 1900 : 148

Type species: *Attagenus varicolor* JAYNE, 1882 : 345.

List of species:

Novelsis aequalis (SHARP, 1902 : 646) BEAL, 1954b : 75.

Mexico, Texas

Syn.: *Genattus aequalis* SHARP, 1902:646.

Novelsis perplexa (JAYNE, 1882 : 345) CASEY, 1900 : 149.

Syn.: *Attagenus perplexus* JAYNE, 1882:345.

Nevada, British
Columbia,
California,
Washington,
Oregon, Idaho

Novelsis varicolor (JAYNE, 1882 : 345) CASEY, 1900 : 149.

Pacific States,
but only

Syn.: *Trogoderma varicolor* STURM, 1843:94, nomen nudum.

California,

Attagenus varicolor CROTCH, 1873:41, nomen nudum.

Arizona,

Attagenus varicolor JAYNE, 1882:345.

Texas,

New Mexico,

Mexico:

Sonora

Novelsis byturoides (CROTCH, 1873:41) CASEY, 1900:148 — vide *Novelsis uteana* CASEY, 1900:148.

Genus: *Decamerus* SOLIER, 1849 : 369

Type species: *Decamerus haemorrhoidalis* SOLIER, 1849 : 371.

List of species:

Decamerus haemorrhoidalis SOLIER, 1849 : 371.

Chile

***Megatominae* LEACH, 1815 : 94**

Type genus: *Megatoma* HERBST, 1792 : 92.

Syn.: *Megatomida* LEACH, 1815:94.

Megatomini: GANGLBAUER, 1904b:8.

Megatominae: DALLA TORRE, 1911:60.

Genus: *Trogoderma* DEJEAN, 1821 : 46

Type species: *Anthrenus elongatus* FABRICIUS, 1801 : 106.

Syn.: *Macropriion* HOPE, 1840:68 — ERICHSON, 1846:447. Type species: *Anthrenus serraticornis* FABRICIUS, 1792:265.

Eurhopalus SOLIER, 1849:372 — LACORDAIRE, 1854:459. Type species: *Eurhopalus variegatus* SOLIER, 1849:373.

Ocelliger PHILIPPI et PHILIPPI, 1864:283 — GERMAIN, 1892:254. Type species: *Ocelliger ater* PHILIPPI et PHILIPPI, 1864:283.

Psacus PASCOE, 1866:446 — HINTON, 1945b:375. Type species: *Psacus attagenoides* PASCOE, 1866:446.

Asidora MULSANT et REY, 1868: 122 — DALLA TORRE, 1911:67. Type species: *Trogoderma fuscicornis* MULSANT et REY, 1868:122.

Acolpus JAYNE, 1882:347 — BEAL, 1954c:50. Type species: *Acolpus primus* JAYNE, 1882:361. *Eucnocerus* SHARP, 1902:648 — HINTON, 1945b:375. Type species: *Eucnocerus dispar* SHARP, 1902:649.

Entomotrogus GANGLBAUER, 1904b:27 — HINTON, 1945b:375. Type species: *Trogoderma megatomoides* REITTER, 1881b:85.

Pseudomegatoma PIC, 1915b:3 — MROCKOWSKI, 1960d:95. Type species: *Pseudomegatoma boliviensis* PIC, 1915b:4.

List of species:

Trogoderma adelaide BLACKBURN, 1891 : 125.

South Australia

Trogoderma albonotatum MULSANT et REY, 1868 : 125.

Pyrénées

Trogoderma alpicolum BLACKBURN, 1891 : 124.

Victoria, North Queensland

Trogoderma angustum (SOLIER, 1849 : 374) LACORDAIRE, 1854 : 468.

Chile, USA:

New York,
Maine;
Germany
(introduced),
Sweden
(introduced)

Syn.: *Eurhopalus angustus* SOLIER, 1849:374.

Trogoderma bifasciatum REDTENBACHER, 1867:44, syn. nov. — KALIK in litt.

Pseudomegatoma boliviensis PIC, 1915b:4. — MROCKOWSKI, 1960d:95.

Globicornis picta: KLEINE, 1940:17 nec (KÜSTER, 1851:39). — MROCKOWSKI, 1956:29.

Attagenus bifasciatus: KORGE, 1961:25. — KORGE, 1962:73.

- Trogoderma antennale* BROUN, 1893 : 1106. **New Zealand**
- Trogoderma anthrenoides* (SHARP, 1902 : 649) BEAL, 1961 : 112. **Mexico,**
Syn.: *Eucnocerus anthrenoides* SHARP, 1902:649. **Panama,**
West Indies,
Colombia,
Texas,
Hawaiian Is.,
S. Mariana Is.
- Trogoderma antipodum* BLACKBURN, 1891 : 128. **S. Australia**
- Trogoderma apicale* (MACLEAY, 1871 : 170) ARMSTRONG, 1942 : 321. **Queensland,**
Syn.: *Megatoma apicalis* MACLEAY, 1871:170. **New South**
Trogoderma pectinifer ARROW, 1915:429. — ARMSTRONG, 1942: **Wales**
321.
- Trogoderma apicipenne* REITTER, 1881c : 232. **S. Australia,**
Syn.: *Trogoderma baldiense* BLACKBURN, 1891:127. — ARMSTRONG, **Victoria,**
1942:322. **Queensland**
- Trogoderma asperatum* FAUVEL, 1903 : 337. **New Caledonia**
- Trogoderma atrum* (PHILIPPI et PHILIPPI, 1864 : 283) comb. nov. **Chile**
- Syn.: *Ocelliger ater* PHILIPPI et PHILIPPI, 1864:283.
- Trogoderma attagenoides* (PASCOE, 1866 : 466) ARMSTRONG, 1949 : 108. **South Australia**
- Syn.: *Psacus attagenoides* PASCOE, 1866:466.
- Trogoderma baeri* (PIC, 1915b : 5) comb. nov. **Argentina**
- Syn.: *Eucnocerus baeri* PIC, 1915b:5.
- Trogoderma ballfinchae* BEAL, 1954 c : 52. **Arizona**
- Trogoderma beali* nom. nov. **Costa Rica**
- Syn.: *Trogoderma atrum* (PIC, 1937d:193) nec (PHILIPPI et PHILIPPI, 1864:283).
Eucnocerus ater PIC, 1937d:193.
- Trogoderma bicinctum* REITTER, 1881 : 38. **West Indies**
- Trogoderma blackburni* LEA, 1908 : 155. **King I.**
(Victoria)
- Trogoderma boganense* ARMSTRONG, 1942 : 322. **New South**
Wales
- Trogoderma breve* CASEY, 1900 : 154. **California**
- Trogoderma callubriense* (ARMSTRONG, 1945 : 48) ARMSTRONG, 1949 : 108. **New South**
Wales
- Syn.: *Psacus callubriensis* ARMSTRONG, 1945:48.

<i>Trogoderma carteri</i> ARMSTRONG, 1942 : 322	New South Wales
<i>Trogoderma celatum</i> SHARP, 1902 : 647.	Mexico
<i>Trogoderma consors</i> ARROW, 1915 : 429.	N. W. Australia
<i>Trogoderma debilius</i> BLACKBURN, 1903 : 163.	West Australia
<i>Trogoderma difficile</i> BLACKBURN, 1891 : 126.	South Australia
<i>Trogoderma ellipticum</i> ARMSTRONG, 1942 : 321.	South Australia
<i>Trogoderma explanaticolle</i> ARMSTRONG, 1942 : 321.	South Australia
<i>Trogoderma exsul</i> BLACKBURN, 1903 : 163.	Central Australia
<i>Trogoderma eyrense</i> BLACKBURN, 1891 : 124.	South Australia
<i>Trogoderma fasciferum</i> BLATCHLEY, 1914 : 66.	Florida
Syn.: <i>Trogoderma flabellata</i> BLATCHLEY, 1922:28. — BEAL, 1954c:64.	
<i>Trogoderma frater</i> ARROW, 1915 : 430.	New South Wales, Victoria
<i>Trogoderma froggatti</i> BLACKBURN, 1892 : 34.	New South Wales, King I. (Victoria)
<i>Trogoderma glabrum</i> (HERBST, 1783 : 26) DALLA TORRE, 1879 : 100.	Berlin, widely distributed throughout Holarctic Region
Syn.: <i>Anthrenus glaber</i> HERBST, 1783:26.	
<i>Byrrhus glaber</i> (HERBST, 1783:26) SCHNEIDER, 1785:5.	
<i>Anthrenus niger</i> HERBST, 1797:338. — HEYDEN, REITTER et WEISE, 1884:209.	
<i>Anthrenus elongatulus</i> FABRICIUS, 1801:106. — LATREILLE, 1807: 40.	
<i>Dermestes versicolor</i> ILLIGER, 1801:86 nec (CREUTZER, 1799: 117). — ERICHSON, 1846:448.	
<i>Anthrenus elongatus</i> SCHÖNHERR, 1806:114. — SCHÖNHERR, 1806: 114.	
<i>Anthrenus ruficornis</i> LATREILLE, 1807:39. — DEJEAN, 1821:46.	
<i>Dermestes subfasciatus</i> GYLLENHAL, 1808:155. — GYLLENHAL, 1827:283.	
<i>Trogoderma elongatula</i> (FABRICIUS, 1801:106) DEJEAN, 1821:46.	
<i>Trogoderma elongata</i> (SCHÖNHERR, 1806:114) LATREILLE, 1829: 511.	
<i>Trogoderma ruficornis</i> (LATREILLE, 1807:39) LATREILLE, 1829:511.	
<i>Trogoderma subfasciata</i> (GYLLENHAL, 1808:155) LATREILLE, 1829: 511.	
<i>Trogoderma villosum</i> DEJEAN, 1837:139, nomen nudum. — DEJEAN, 1837:139.	
<i>Trogoderma nigrum</i> (HERBST, 1797:338) SCHMIDT, 1844:31.	
<i>Trogoderma fuscicornis</i> MULSANT et REY, 1868:122. — REITTER, 1883:89.	

<i>Trogoderma fusicornis</i> MULSANT et REY, 1868:122. — REITTER, 1891:170.	
<i>Trogoderma nigrescans</i> : HICKS, 1953:13 nec CASEY, 1916:188. — BEAL, 1954c:80.	
<i>Trogoderma boron</i> BEAL, 1954c:53. — BEAL, 1956:561.	
<i>Trogoderma glabrum limbatum</i> PIC, 1934 : 26.	France
Syn.: <i>Trogoderma nigrum limbatum</i> PIC, 1934:26.	
<i>Trogoderma gounellei</i> PIC, 1915b : 4.	Brazil
<i>Trogoderma granarium</i> EVERTS, 1898 : 610.	Rotterdam, very widely distributed, nearly cosmopolitan
Syn.: <i>Trogoderma quinquefasciata</i> LEESBERG, 1906:79 nec JACQUELIN DU VAL, 1859:257. — LEESBERG, 1906:79.	
<i>Trogoderma khapra</i> ARROW, 1917:481. — EVERTS, 1923:169.	
<i>Trogoderma afrum</i> PRIESNER, 1951:133. — HOWE et BURGES, 1956:773.	
<i>Trogoderma granulatum</i> BROUN, 1886 : 900.	New Zealand
<i>Trogoderma grassmani</i> BEAL, 1954c : 54.	Arizona, California, Mexico
<i>Trogoderma hobartense</i> ARMSTRONG, 1942 : 321.	Tasmania
<i>Trogoderma inclusum</i> LECONTE, 1854 : 110.	New York, nearly whole USA, England
Syn.: <i>Trogoderma tarsale</i> : RILEY, 1894:226. — BEAL, 1956:562.	
<i>Trogoderma advena</i> CASEY, 1900:154. — BEAL, 1954c:84.	
<i>Trogoderma obsolescens</i> CASEY, 1900:154. — BEAL, 1954c:84.	
<i>Trogoderma frosti</i> CASEY, 1916:188. — BEAL, 1954c:84.	
<i>Trogoderma nigrescans</i> CASEY, 1916:188. — BEAL, 1954c:84.	
<i>Trogoderma brunnescens</i> CASEY, 1916:189. — BEAL, 1954c:84.	
<i>Trogoderma versicolor</i> : MUTHLER et WEISS, 1927:12 nec (CREUTZER, 1799:117). — BEAL, 1956:563.	
<i>Trogoderma inconspicuum</i> ARMSTRONG, 1942 : 322.	West Australia
<i>Trogoderma insulare</i> CHEVROLAT, 1863 : 617.	Cuba, Puerto Rico
<i>Trogoderma interruptum</i> (THUNBERG, 1781 : 6) comb. nov., KALIK in litt.	Without terra typica
Syn.: <i>Dermestes interruptus</i> THUNBERG, 1781:6.	
<i>Trogoderma irroratum</i> REITTER, 1881 : 41.	Egypt
<i>Trogoderma koenigi</i> PIC, 1954 : 55.	China
<i>Trogoderma koningsbergeri</i> PIC, 1933 : 71.	Java
<i>Trogoderma laevipenne</i> ARMSTRONG, 1942 : 324.	Queensland
Syn.: <i>Trogoderma laevipennae</i> ARMSTRONG, 1942:321 (error).	
<i>Trogoderma laticorne</i> CHAO et LEE, 1966 : 247.	China
Syn.: <i>Trogoderma latecorne</i> CHAO et LEE, 1966:245 (error).	
<i>Trogoderma laticicorne</i> CHAO et LEE, 1966:251 (error).	

<i>Trogoderma leai</i> ARMSTRONG, 1942 : 322.	West Australia
<i>Trogoderma lindense</i> BLACKBURN, 1891 : 125.	South Australia
<i>Trogoderma longisetosum</i> CHAO et LEE, 1966 : 245.	China
<i>Trogoderma longius</i> BLACKBURN, 1903 : 163.	Victoria, Tasmania, New South Wales
<i>Trogoderma macleayi</i> BLACKBURN, 1891 : 126.	South Australia, Victoria
<i>Trogoderma maestum</i> BROUN, 1880 : 241.	New Zealand
Syn.: <i>Trogoderma moestum</i> DALLA TORRE, 1911:68.	
<i>Trogoderma marginicolle</i> ARMSTRONG, 1942 : 321.	South Australia
<i>Trogoderma maurulum</i> BLACKBURN, 1903 : 163.	Victoria
<i>Trogoderma megatomoides</i> REITTER, 1881b : 85.	Czechoslova- kia, Hungary, Germany, Austria, Sweden, Holland, France, England, Mexico, Central America. (Introduced to Europe).
Syn.: <i>Trogoderma filipes</i> SHARP, 1902:647. — HINTON, 1945b:385. <i>Trogoderma gravidum</i> SHARP, 1902:648. — HINTON, 1945b:385. <i>Entomotrogus megatomoides</i> (REITTER, 1881b:85) GANGLBAUER, 1904 b:33.	Czechoslovakia
<i>Trogoderma megatomoides</i> var. <i>nigripubens</i> (REITTER, 1911b : 291).	Mexico
Syn.: <i>Entomotrogus megatomoides</i> var. <i>nigripubens</i> REITTER, 1911b:291.	West Australia
<i>Trogoderma mexicanum</i> REITTER, 1881 : 38.	Tasmania, King I., New South Wales, Victoria, West Australia
<i>Trogoderma meyricki</i> BLACKBURN, 1891 : 128.	Queensland
<i>Trogoderma morio</i> (ERICHSOHN, 1842 : 152) BLACKBURN, 1903 : 163.	North Queensland
Syn.: <i>Megatoma morio</i> ERICHSON, 1842:152.	Brazil
<i>Trogoderma nigrobrunneum</i> ARMSTRONG, 1942 : 322.	West Australia
<i>Trogoderma nigronitidum</i> ARMSTRONG, 1945 : 49.	Sahara
<i>Trogoderma nitens</i> ARROW, 1915 : 432.	
<i>Trogoderma occidentale</i> BLACKBURN, 1891 : 127.	
<i>Trogoderma octaedron</i> PEYERIMHOFF, 1943 : 12.	

<i>Trogoderma okumurai</i> BEAL, 1964 : 81.	Arizona, Utah
<i>Trogoderma oothecophilum</i> CHAO et LEE, 1966 : 245.	China
<i>Trogoderma ornatum</i> (SAY, 1825 : 185) LECONTE, 1854 : 110.	USA
Syn.: <i>Megatoma ornata</i> SAY, 1825:185.	
<i>Trogoderma tarsale</i> MELSHIMER, 1844:116. — HINTON, 1945b: 399.	
<i>Trogoderma pallipes</i> ZIEGLER, 1845:269. — HENSHAW, 1885:54.	
<i>Attagenus ornatus</i> (SAY, 1825:185) MELSHIMER, 1853:48.	
<i>Trogoderma pusillum</i> LECONTE, 1854:111. — JAYNE, 1882:375.	
<i>Trogoderma serriger</i> CASEY, 1900:153. — BEAL, 1954c:66.	
<i>Eucnocerus dispar</i> SHARP, 1902:649. — HINTON, 1945b:376.	
<i>Trogoderma serriferum</i> DALLA TORRE, 1911:70. — BEAL, 1954c: 66.	
<i>Trogoderma procera</i> CASEY, 1916:185. — BEAL, 1954 c:66.	
<i>Trogoderma parvula</i> CASEY, 1916:186. — BEAL, 1954c:67.	
<i>Trogoderma paralia</i> BEAL, 1945c : 53.	California
<i>Trogoderma parvum</i> ARMSTRONG, 1942 : 321.	West Australia
<i>Trogoderma pectinicornis</i> REITTER, 1881 : 39.	Brazil, Argentina
<i>Trogoderma picinum</i> ARMSTRONG, 1945 : 49.	New South Wales
<i>Trogoderma pictulum</i> BROUN, 1911 : 100.	Chatham Isl.
<i>Trogoderma primum</i> (JAYNE, 1882 : 361) BEAL, 1954c : 56.	Texas, Oklahoma
Syn.: <i>Acolpus primus</i> JAYNE, 1882:361.	
<i>Trogoderma punctatum</i> BROUN, 1886 : 953.	New Zealand
<i>Trogoderma puncticolle</i> BROUN, 1914 : 187.	New Zealand
<i>Trogoderma quadrifasciatum</i> BROUN, 1893 : 1105.	New Zealand
<i>Trogoderma quinquefasciatum</i> JACQUELIN DU VAL, 1859 : 257.	France, Germany
<i>Trogoderma reitteri</i> BLACKBURN, 1892b : 207.	New South Wales
<i>Trogoderma rubiginosum</i> (SOLIER, 1849 : 373) LACORDAIRE, 1854 : 468.	Chile
Syn.: <i>Eurhopalus rubiginosus</i> SOLIER, 1849:373.	
<i>Trogoderma ruficeps</i> PIC, 1937 : 410.	former British Guiana
<i>Trogoderma ruficollis</i> REITTER, 1881 : 39.	Brazil
<i>Trogoderma rufipenne</i> ARMSTRONG, 1942 : 322.	New South Wales
<i>Trogoderma rufonotatum</i> PIC, 1942 : 6.	Peru
<i>Trogoderma rufopictum</i> ARROW, 1915 : 432.	Natal

<i>Trogoderma schmorli</i> REITTER, 1881 : 39.	Brazil
<i>Trogoderma seminigrum</i> PIC, 1915b : 5.	Madagascar
<i>Trogoderma serraticorne</i> (FABRICIUS, 1792 : 265) LACORDAIRE, 1854 : 468.	St. Croix (Antilles)
Syn.: <i>Anthrenus serraticornis</i> FABRICIUS, 1792:265. <i>Anthrenus denticornis</i> FABRICIUS, 1792:265. — MOTSCHULSKY, 1858b:147.	
<i>Trogoderma denticornis</i> (FABRICIUS, 1792:265) LACORDAIRE 1854:468.	
<i>Macroprion serraticornis</i> (FABRICIUS, 1792:265) HOPE, 1840:68.	
<i>Trogoderma serrigerum</i> SHARP, 1877 : 270.	New Zealand
<i>Trogoderma setulosum</i> ARMSTRONG, 1942 : 322.	Queensland
<i>Trogoderma signatum</i> SHARP, 1877 : 271.	New Zealand
<i>Trogoderma silvicolum</i> ARMSTRONG, 1949 : 107.	New South Wales
<i>Trogoderma simplex</i> JAYNE, 1882 : 362.	USA: Western States
Syn.: <i>Trogoderma pollens</i> CASEY, 1900:153. — BEAL, 1954c:57. <i>Trogoderma ajax</i> CASEY, 1924:186. — BEAL, 1954c:57.	
<i>Trogoderma sinense</i> PIC, 1927e : 6.	China
<i>Trogoderma singulare</i> BLACKBURN, 1891 : 128.	South Australia, West Australia, New South Wales
<i>Trogoderma sinistrum</i> FALL, 1926 : 193.	Yukon Territory, Alaska, Alberta, British Columbia, Colorado, Illinois, Minnesota, Ontario, Quebec, Utah, Wyoming
<i>Trogoderma socium</i> (LEA, 1895 : 228) BLACKBURN, 1903 : 172.	Sydney
Syn.: <i>Anthrenus socius</i> LEA, 1895:228.	
<i>Trogoderma stachi</i> MROCKOWSKI, 1958 : 559.	Brazil
Syn.: <i>Trogoderma apicale</i> PIC, 1936:1 nec (MACLEAY, 1871:170).	

Trogoderma sternale JAYNE, 1882 : 362.

Syn.: *Trogoderma simulans* CASEY, 1900:153. — BEAL, 1954c:71.
Trogoderma variipes CASEY, 1900:153. — BEAL, 1954c:71.
Trogoderma caseyi ARROW, 1915:429. — BEAL, 1954c:71.

California,
Oregon,
Washington,
British
Columbia

Trogoderma sternale aspericolle CASEY, 1900 : 154. — BEAL,
1954c : 74.

Arizona

Syn.: *Trogoderma aspericollis* CASEY, 1900:154.

Trogoderma sternale complex CASEY, 1900 : 153. — BEAL,
1954c : 74.

Texas, Virginia

Syn.: *Trogoderma complex* CASEY, 1900:153.

Trogoderma virginica CASEY, 1900:154. — BEAL, 1954c:74.
Trogoderma oblongula CASEY, 1900:154. — BEAL, 1954c:74.

Trogoderma sternale deserti BEAL, 1954c : 73.

California,
Arizona

Trogoderma sternale maderae BEAL, 1954c : 71.

California,
Arizona

Trogoderma sternale plagifer CASEY, 1916 : 187. — BEAL,
1954c : 76.

Colorado,
Utah,
New Mexico

Syn.: *Trogoderma plagifera* CASEY, 1916:187.

Trogoderma cylindrella CASEY, 1916:187. — BEAL, 1954c:76.

Trogoderma submarginatum PIC, 1942 : 6.

Argentina

Trogoderma subrotundatum REITTER, 1881 : 39.

Brazil

Trogoderma subtile REITTER, 1881 : 39.

Chile

Trogoderma suffusum BROUN, 1886 : 953.

New Zealand

Trogoderma syriacum DALLA TORRE, 1911 : 70.

Syria

Syn.: *Trogoderma maculifasciatum* REITTER, 1899c:283 nec REITTER,
1887:205.

Trogoderma tasmanicum ARMSTRONG, 1942 : 322.

Tasmania

Trogoderma teukton BEAL, 1956 : 560.

Minnesota,
Iowa, North
Dakota

Syn.: *Trogoderma glabrum*: BEAL, 1954c:53 nec (HERBST, 1783:26). —
BEAL, 1956:565.

Trogoderma thoracicum REITTER, 1881 : 39.

Brazil

Trogoderma thoracicum var. *bicoloripes* PIC, 1936 : 1.

Brazil

Trogoderma thoracicum var. *obscurum* PIC, 1936 : 1.

Brazil

Trogoderma tolarnense BLACKBURN, 1903 : 162.

New South
Wales

Trogoderma unifasciatum PIC, 1942 : 6.

India

Trogoderma variabile BALLION, 1878 : 277.

Syn.: *Phradonoma variabile* (BALLION, 1878:277) DALLA TORRE, 1911 : 66.

Trogoderma persica PIC, 1914:10. — ZHANTIEV, in litt.

Trogoderma scabripennis CASEY, 1916:189. — BEAL, in litt.

Trogoderma parabile BEAL, 1954c:53. — ZHANTIEV, in litt.

Kuldsha

(Sinkiang),
Causasus, Iran,
Afghanistan,
Uzbek SSR,
Kazakh SSR,
Tadzhik SSR,
Turkmen SSR,
USA

Trogoderma variegatum (SOLIER, 1849 : 373) LACORDAIRE, 1854 : 468.

Syn.: *Eurhopalus variegatus* SOLIER, 1849:373.

Trogoderma varipes BLACKBURN, 1892b : 208.

Syn.: *Trogoderma variipes* ARROW, 1915:429.

Trogoderma varium (MATSUMURA et YOKOYAMA, 1928b : 51) HINTON, 1945b : 381.

Syn.: *Megatoma varius* MATSUMURA et YOKOYAMA, 1928b:51.

Trogoderma versicolor (CREUTZER, 1799 : 117) DEJEAN, 1821 : 46.

Syn.: *Anthrenus versicolor* CREUTZER, 1799:117.

Anthrenus grandis THUNBERG, 1815:151. — REITTER, 1891:170.

Anthrenus elongatus DUFTSCHMID, 1825:26. — ERICHSON, 1846 : 448.

Trogoderma costae MULSANT et REY, 1868:127. — REITTER, 1881b:85.

Trogoderma versicolor ab. *major* NORMAND, 1936 : 300, nomen nudum.

Trogoderma versicolor var. *reductum* PIC, 1919 : 17.

Trogoderma versicolor var. *turkestanicum* PIC, 1914 : 10.

Trogoderma versicolor meridionale KRAATZ, 1858 : 146. — MROCKOWSKI, 1962b : 548.

Syn.: *Trogoderma meridionalis* KRAATZ, 1858:146.

Trogoderma flexuosa THOMSON, 1862:207. — REITTER, 1881b : 85.

Trogoderma testaceicorne PERRIS, 1862:196. — REITTER, 1881b:85.

Trogoderma hieroglyphica ABEILLE, 1872:XLII. — REITTER, 1881b:85.

Trogoderma vicinum (SOLIER, 1849 : 374) LACORDAIRE, 1854 : 468.

Syn.: *Eurhopalus vicinus* SOLIER, 1849:374.

Trogoderma vulneratum FAUVEL, 1903 : 337.

Chile

South Australia

Japan

Vienna
(Austria),
nearly whole
Palearctic
Region

Tunis

Egypt

Turkestan

Greece, Sicily,
widely
distributed on
south parts of
areal of species

Chile

New Caledonia

Trogoderma whitei ARMSTRONG, 1942 : 322.

Central
Australia

Trogoderma yorkense BLACKBURN, 1891 : 127.

South Australia

Trogoderma advena CASEY, 1900:154 — vide *Trogoderma inclusum* LECONTE, 1854:110.

Trogoderma aegyptiaca PIC, 1901:33 — vide *Phradonoma aegyptiacum* (PIC, 1901:33).

Trogoderma afrum PRIESNER, 1951:133 — vide *Trogoderma granarium* EVERTS, 1898:610.

Trogoderma ajax CASEY, 1924:186 — vide *Trogoderma simplex* JAYNE, 1882:362.

Trogoderma americanum DEJEAN, 1837:139 — nomen nudum.

Trogoderma amoenula REITTER, 1889d:557 — vide *Phradonoma amoenum* (REITTER, 1889d:557).

Trogoderma apicale PIC, 1936:1 nec (MACLEAY, 1871:170) — vide *Trogoderma stachi* MROCKOWSKI, 1958:559.

Trogoderma aspericollis CASEY, 1900:154 — vide *Trogoderma sternale aspericolle* CASEY, 1900:154.

Trogoderma atrum (PIC, 1937d:193) nec (PHILIPPI et PHILIPPI, 1864:283) — vide *Trogoderma beali* nom. nov.

Trogoderma babaulti PIC, 1921b:12 — vide *Phradonoma babaulti* (PIC, 1921b:12).

Trogoderma baldiense BLACKBURN, 1891:127 — vide *Trogoderma apicipenne* REITTER, 1881c:232.

Trogoderma belfragei LECONTE, 1874:50 — vide *Megatoma belfragei* (LECONTE, 1874:50).

Trogoderma bifasciatum REDTENBACHER, 1867:44 — vide *Trogoderma angustum* (SOLIER, 1849:374).

Pseudomegatoma boliviensis PIC, 1915b:4 — vide *Trogoderma angustum* (SOLIER, 1849:374).

Trogoderma boron BEAL, 1954c:53 — vide *Trogoderma glabrum* (HERBST, 1783:26).

Trogoderma brasiliensis REITTER, 1881:37 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:138).

Trogoderma brunneascens CASEY, 1916:189 — vide *Trogoderma inclusum* LECONTE, 1854:110.

Trogoderma calliesi PEYERIMHOFF, 1928:304 — vide *Phradonoma calliesi* (PEYERIMHOFF, 1928:304).

Trogoderma caseyi ARROW, 1915:429 — vide *Trogoderma sternale* JAYNE, 1882:362.

Trogoderma cercyonoides REITTER, 1887b:62 — vide *Phradonoma cercyonoides* (REITTER, 1887b:62).

Trogoderma complex CASEY, 1900:153 — vide *Trogoderma sternale complex* CASEY, 1900:153.

Trododerma (sic!) *congoana* PIC, 1950:356 — vide *Orphinus congoanus* (PIC, 1950:356).

Trogoderma costae MULSANT et REY, 1868:127 — vide *Trogoderma versicolor* (CREUTZER, 1799:117).

Trogoderma cylindrella CASEY, 1916:187 — vide *Trogoderma sternale plagiifer* CASEY, 1916:187.

Trogoderma cyphonoides (REITTER, 1881:34) ARROW, 1915:427 — vide *Attagenus cyphonoides* REITTER, 1881:34.

Trogoderma defectum (WALKER, 1858:207) OLLIFF, 1883:185 — vide *Orphinus defectus* (WALKER, 1858:207).

Trogoderma denticornis (FABRICIUS, 1792:265) LACORDAIRE, 1854:468 — vide *Trogoderma serraticorne* (FABRICIUS, 1792:265).

Trogoderma dichroa REITTER, 1900b:219 — vide *Phradonoma dichroum* (REITTER, 1900b:219).

Eucocerus dispar SHARP, 1902:649 — vide *Trogoderma ornatum* (SAY, 1825:185).

Trogoderma elongata (SCHÖNHERR, 1806:114) LATREILLE, 1829:511 — vide *Trogoderma glabrum* (HERBST, 1783:26).

Trogoderma elongatula (FABRICIUS, 1801:106) DEJEAN, 1821:46 — vide *Trogoderma glabrum* (HERBST, 1783:26).

Trogoderma eximium ARROW, 1915:431 — vide *Phradonoma eximium* (ARROW, 1915:431).

Trogoderma filipes SHARP, 1902:647 — vide *Trogoderma megatomoides* REITTER, 1881b:85.

Trogoderma flabellata BLATCHLEY, 1922:28 — vide *Trogoderma fasciferum* BLATCHLEY, 1914:66.

Trogoderma flexuosa THOMSON, 1862:207 — vide *Trogoderma versicolor meridionale* KRAATZ, 1858:146.

Trogoderma frosti CASEY, 1916:188 — vide *Trogoderma inclusum* LECONTE, 1854:110.

Trogoderma fulvipes (GUÉRIN-MÉNEVILLE, 1838c:138) LACORDAIRE, 1854:467 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:138).

Trogoderma funestum REITTER, 1881:41 — vide *Phradonoma funestum* (REITTER, 1881:41).

- Trogoderma fuscicornis* MULSANT et REY, 1868:122 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Trogoderma fuscicornis MULSANT et REY, 1868:122 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Trogoderma glabrum: BEAL, 1954c:53 nec (HERBST, 1783:26) — vide *Trogoderma teukton* BEAL, 1956:560.
Trogoderma gravidum SHARP, 1902:648 — vide *Trogoderma megatomoides* REITTER, 1881b:85.
Trogoderma haemorrhoa GERSTAECKER, 1871:45 — vide *Phradonoma haemorrhoum* (GERSTAECKER, 1871: 45).
Trogoderma hieroglyphica ABEILLE, 1872:XLII — vide *Trogoderma versicolor meridionale* KRAATZ, 1858: 146.
Trogoderma hirsutula MULSANT et REY, 1868:134 — vide *Phradonoma hirsutulum* (MULSANT et REY, 1868:134).
Trogoderma holosericea (SAHLBERG, 1819:52) SCHAUM, 1852:38 — vide *Attagenus obtusus* (GYLLENHAL in SCHÖNHERR, 1808:88).
Trogoderma humerale REITTER, 1881:41 — vide *Phradonoma humerale* (REITTER, 1881:41).
Trogoderma khapra ARROW, 1917:481 — vide *Trogoderma granarium* EVERETT, 1898:610.
Trogoderma laevipennae ARMSTRONG, 1942:321 — vide *Trogoderma laevipenne* ARMSTRONG, 1942:324.
Trogoderma latecorne CHAO et LEE, 1966:245 — vide *Trogoderma laticorne* CHAO et LEE, 1966:247.
Trogoderma laticorne CHAO et LEE, 1966:251 — vide *Trogoderma laticorne* CHAO et LEE, 1966:247.
Trogoderma maculifasciata REITTER, 1887:205 — vide *Phradonoma maculifasciatum* (REITTER, 1887:205).
Trogoderma maculifasciatum REITTER, 1899c:283 — vide *Trogoderma syriacum* DALLA TORRE, 1911:70.
Psacus mastersii MACLEAY, 1872:313 — non *Dermestidae* apud ARMSTRONG, 1945:48.
Trogoderma melanarium STURM, 1843:94 — nomen nudum.
Trogoderma meridionalis KRAATZ, 1858:146 — vide *Trogoderma versicolor meridionale* KRAATZ, 1858:146.
Trogoderma moestum DALLA TORRE, 1911:68 — vide *Trogoderma maestum* BROUN, 1880:241.
Trogoderma nigrescans CASEY, 1916:188 — vide *Trogoderma inclusum* LECONTE, 1854:110.
Trogoderma nigrescens: HICKS, 1953:13 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Trogoderma nigrum (HERBST, 1797:338) SCHMIDT, 1844:31 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Trogoderma nigrum limbatum PIC, 1934:26 — vide *Trogoderma glabrum limbatum* PIC, 1934:26.
Trogoderma nobile REITTER, 1881b:85 — vide *Phradonoma nobile* (REITTER, 1881b:85).
Trogoderma nobile paulonotata PIC, 1908b:41 — vide *Phradonoma nobile paulonotatum* (PIC, 1908b:41).
Trogoderma oblongula CASEY, 1900:154 — vide *Trogoderma sternale complex* CASEY, 1900:153.
Trogoderma obsoletes CASEY, 1900:154 — vide *Trogoderma inclusum* LECONTE, 1854:110.
Trogoderma ornata SOLSKIJ, 1876:278 nec (SAY, 1825:185) — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).
Trogoderma pallipes ZIEGLER, 1845:269 — vide *Trogoderma ornatum* (SAY, 1825:186).
Trogoderma pantherinum (AHRENS, 1814:11) SCHMIDT, 1844:31 — vide *Attagenus pantherinus* (AHRENS, 1814:11).
Trogoderma parabile BEAL, 1854c:53 — vide *Trogoderma variabile* BALLION, 1878:277.
Trogoderma parvula CASEY, 1916:186 — vide *Trogoderma ornatum* (SAY, 1825:186).
Trogoderma pectinifer ARROW, 1915:429 — vide *Trogoderma apicale* (MACLEAY, 1871:170).
Trogoderma persica PIC, 1914:10 — vide *Trogoderma variabile* BALLION, 1878:277.
Trogoderma peruviana PIC, 1954c:178 — nomen nudum.
Trogoderma picta KÜSTER, 1851:39 — vide *Globicornis picta* (KÜSTER, 1851:39).
Trogoderma plagifera CASEY, 1916:187 — vide *Trogoderma sternale plagifer* CASEY, 1916:187.
Trogoderma pollens CASEY, 1900:153 — vide *Trogoderma simplex* JAYNE, 1882:362.
Trogoderma procera CASEY, 1916:185 — vide *Trogoderma ornatum* (SAY, 1825:186).
Trogoderma pusillum LECONTE, 1854:111 — vide *Trogoderma ornatum* (SAY, 1825:186).
Trogoderma quadricolor SUMAKOV, 1907:15 — vide *Attagenus quadricolor* (SUMAKOV, 1907:15).
Trogoderma quinquefasciata LEESBERG, 1906:79 nec JACQUELIN DU VAL, 1859:257 — vide *Trogoderma granarium* EVERETT, 1898:610.
Trogoderma riguum ERICHSON, 1842:152 — vide *Anthrenocerus riguum* (ERICHSON, 1842:152).
Trogoderma rubromaculatum SAHLBERG, 1913:29 — vide *Phradonoma rubromaculatum* (SAHLBERG, 1913: 29).

- Trogoderma rubromarginatum* SAHLBERG, 1913c:109 — vide *Phradonoma rubromaculatum* (SAHLBERG, 1913:29).
- Trogoderma ruficornis* (LATREILLE, 1807:39) LATREILLE, 1829:511 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Trogoderma scabripennis* CASEY, 1916:189. — vide *Trogoderma variable* BALLION 1878:277.
- Trogoderma serratum* DEJEAN, 1837:139 — nomen nudum.
- Trogoderma serriferum* DALLA TORRE, 1911:70 — vide *Trogoderma ornatum* (SAY, 1825:186).
- Trogoderma serriger* CASEY, 1900:153 — vide *Trogoderma ornatum* (SAY, 1825:186).
- Trogoderma simulans* CASEY, 1900:153 — vide *Trogoderma sternale* JAYNE, 1882:362.
- Trogoderma solskyi* REITTER, 1881:38 — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).
- Trogoderma subfasciatum* CHEVROLAT, 1863:617 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
- Trogoderma subfasciata* (GYLLENHAL, 1808:155) LATREILLE, 1829:511 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Trogoderma tamaricis* PIC, 1894h:180 — vide *Phradonoma nobile* var. *trizonatum* FAIRMAIRE, 1883:CLVI.
- Trogoderma tarsale* MELSHEIMER, 1844:116 — vide *Trogoderma ornatum* (SAY, 1825:186).
- Trogoderma tarsale*: RILEY, 1894:226 — vide *Trogoderma inclusum* LECONTE, 1854:110.
- Trogoderma testaceicorne* PERRIS, 1862: 196 — vide *Trogoderma versicolor meridionale* KRAATZ, 1858:146.
- Trogoderma tricolor* ARROW, 1915:431 — vide *Phradonoma tricolor* (ARROW, 1915:431).
- Trogoderma trizonatum* FAIRMAIRE, 1883:CLVI — vide *Phradonoma nobile* var. *trizonatum* FAIRMAIRE, 1883:CLVI.
- Trogoderma trizonatum* var. *major* PIC, 1895:79 — vide *Phradonoma nobile* var. *major* (PIC.) 8195:79.
- Trogoderma unicolor* KOLBE, 1910:24 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:133).
- Trogoderma varicolor* STURM, 1843:94, nomen nudum — vide *Novelsis varicolor* (JAYNE, 1882:845).
- Trogoderma variegata* KÜSTER, 1851:40 — vide *Globicornis variegata* (KÜSTER, 1851:40).
- Trogoderma variipes* CASEY, 1900:153 — vide *Trogoderma sternale* JAYNE, 1882:362.
- Trogoderma variipes* ARROW, 1915:429 — vide *Trogoderma variipes* BLACKBURN, 1892b:208.
- Trogoderma versicolor*: MUTCHLER et WEISS, 1927:12 nec (CREUTZER, 1799:117) — vide *Trogoderma inclusum* LECONTE, 1854:110.
- Trogoderma vicinum* DEJEAN, 1837:139 — nomen nudum.
- Trogoderma villosulum* (DUFTSCHMID, 1825:28) STURM, 1843:94 — vide *Phradonoma villosulum* (DUFTSCHMID, 1825:28).
- Trogoderma villosum* DEJEAN, 1837:139, nomen nudum — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Trogoderma virginica* CASEY, 1900:154 — vide *Trogoderma sternale complex* CASEY, 1900:153.

Genus: *Paratrogoderma* SCOTT, 1926 : 63

Type species: *Paratrogoderma mahense* SCOTT, 1926 : 67.

List of species:

Paratrogoderma mahense SCOTT, 1926 : 67.

Seychelles

Genus: *Reesa* BEAL, 1967 : 310

Type species: *Perimegatoma vespulae* MILLIRON, 1939 : 570.

List of species:

Reesa vespulae (MILLIRON, 1939 : 570) BEAL, 1967 : 310.

Minnesota,

Quebec,

British

Columbia,

Michigan,

North Dakota,

California

Syn.: *Perimegatoma vespulae* MILLIRON, 1939:570.

Megatoma vespulae (MILLIRON, 1939:570) SPENCER, 1948:7.

Genus: *Phradonomorphus* PIC, 1916b : 3

Type species: *Aethriostoma rufofasciata* PIC, 1915 : 2.

List of species:

Phradonomorphus rufofasciatus (PIC, 1915 : 2) PIC, 1916b : 3.

Senegal

Syn.: *Aethriostoma rufofasciata* PIC, 1915:2.

Phradonomorphus rufofasciatus var. *postlimbatus* (PIC, 1948 : 14).

Neghelli

Syn.: *Aethriostoma rufofasciata* var. *postlimbata* PIC, 1948:14.

Genus: *Phradonoma* JACQUELIN DU VAL, 1859 : 257

Type species: *Anthrenus villosulus* DUFTSCHMID, 1825 : 28.

Subgenus: *Orbeola* MULSANT et REY, 1868 : 134

Type species: *Trogoderma hirsutula* MULSANT et REY, 1868 : 134.

List of species:

Phradonoma hirsutulum (MULSANT et REY, 1868 : 134) WINKLER, 1926 : 679.

Syria

Syn.: *Trogoderma hirsutula* MULSANT et REY, 1868:134.

Phradonoma rubromaculatum (SAHLBERG, 1913 : 29) comb. nov.

Egypt, Syria

Syn.: *Trogoderma rubromaculatum* SAHLBERG, 1913:29.

Trogoderma rubromarginatum SAHLBERG, 1913c:109. — syn. nov.

Subgenus: *Phradonoma* s. str.

List of species:

Phradonoma aegyptiacum (PIC, 1901 : 33) WINKLER, 1926 : 679.

Egypt

Syn.: *Trogoderma aegyptiaca* PIC, 1901:33.

Phradonoma amoenulum (REITTER, 1889d : 557) DALLA TORRE, 1911 : 65.

Mongolia centr.

Syn.: *Trogoderma amoenula* REITTER, 1889d:557.

Phradonoma babaulti (PIC, 1921b : 12) KALIK, 1965 : 217.

Kenya,
Tanganyika

Syn.: *Trogoderma babaulti* PIC, 1921b:12.

Phradonoma calliesi (PEYERIMHOFF, 1928 : 304) PEYERIMHOFF, 1931 : 65.

Algeria
(Sahara),
Tunisia

Syn.: *Trogoderma calliesi* PEYERIMHOFF, 1928:304.

Phradonoma cercyonoides (REITTER, 1887b : 62) DALLA TORRE, 1911 : 65.

Egypt, Syria,
Lybia

Syn.: *Trogoderma cercyonoides* REITTER, 1887b:62.

<i>Phradonoma cercyonoides</i> var. <i>demaizoni</i> (REITTER, 1889e : 127).	Egypt
Syn.: <i>Trogoderma cercyonoides demaisonii</i> REITTER, 1889e:127.	
<i>Phradonoma dichroum</i> (REITTER, 1900b : 219) DALLA TORRE, 1911 : 65.	Kirghiz SSR
Syn.: <i>Trogoderma dichroa</i> REITTER, 1900b:219.	
<i>Phradonoma eximium</i> (ARROW, 1915 : 431) HINTON, 1945b : 375.	Rhodesia, Congo (Kinshasa)
Syn.: <i>Trogoderma eximium</i> ARROW, 1915:431.	
<i>Phradonoma funestum</i> (REITTER, 1881 : 41) comb. nov.	South Africa
Syn.: <i>Trogoderma funestum</i> REITTER, 1881:41.	
<i>Phradonoma haemorrhoum</i> (GERSTAECKER, 1871 : 45) KALIK, 1965 : 219.	Zanzibar
Syn.: <i>Trogoderma haemorrhoa</i> GERSTAECKER, 1871:45.	
<i>Phradonoma humerale</i> (REITTER, 1881 : 41) comb. nov.	South Africa
Syn.: <i>Trogoderma humerale</i> REITTER, 1881:41.	
<i>Phradonoma impressiceps</i> PIC, 1915b : 6.	Madagascar
<i>Phradonoma maculifasciatum</i> (REITTER, 1887 : 205) DALLA TORRE, 1911 : 66.	Sinkiang
Syn.: <i>Trogoderma maculifasciata</i> REITTER, 1887:205.	
<i>Phradonoma nobile</i> (REITTER, 1881b : 85) DALLA TORRE, 1911 : 66.	Cyprus, whole Mediterranean Region, Portugal, Asia Minor, Caucasus, Turkmen SSR, Afghanistan
Syn.: <i>Trogoderma nobile</i> REITTER, 1881b:85.	
<i>Phradonoma nobile</i> var. <i>distincta</i> KALIK, 1954 : 369.	S. W. Africa, Rhodesia
<i>Phradonoma nobile</i> var. <i>major</i> (PIC, 1895 : 79).	Algeria
Syn.: <i>Trogoderma trizonatum</i> var. <i>major</i> PIC, 1895:79.	
<i>Phradonoma nobile</i> var. <i>majus</i> DALLA TORRE, 1911:66, syn. nov.	
<i>Phradonoma nobile</i> var. <i>trizonatum</i> FAIRMAIRE, 1883 : CLVI. — PIC, 1895b : 118.	Algeria
Syn.: <i>Trogoderma trizonatum</i> FAIRMAIRE, 1883:CLVI.	
<i>Trogoderma tamaricis</i> PIC, 1894h:180. — PIC, 1895b:118.	

<i>Phradonoma nobile paulonotatum</i> (PIC, 1908b : 41) DALLA TORRE, 1911 : 66.	Egypt
Syn.: <i>Trogoderma nobile paulonotata</i> PIC, 1908b:41.	
<i>Phradonoma tricolor</i> (ARROW, 1915 : 431) HINTON, 1945b : 373.	Yemen
Syn.: <i>Trogoderma tricolor</i> ARROW, 1915:431.	
<i>Phradonoma turcomanicum</i> MROCZKOWSKI, 1960b : 214.	Turkmen SSR
<i>Phradonoma uninotata</i> (PIC, 1942 : 6) comb. nov., KALIK in litt.	East Africa
Syn.: <i>Telopes uninotatus</i> PIC, 1942:6.	
<i>Phradonoma villosulum</i> (DUFTSCHMID, 1825 : 28) REITTER, 1906 : 379.	Austria, south and middle Europa, middle Asia
Syn.: <i>Anthrenus villosulus</i> DUFTSCHMID, 1825:28.	
<i>Trogoderma villosulum</i> (DUFTSCHMID, 1825:28) STURM, 1843:94.	

Phradonoma nobile var. *majus* DALLA TORRE, 1911:66, syn. nov. — vide *Phradonoma nobile* var. *major* (PIC, 1895:79).

Phradonoma ornatum (SOLSKII, 1876:278) DALLA TORRE, 1911:66 — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).

Phradonoma quadricolor (SUMAKOV, 1907:15) DALLA TORRE, 1911:66 — vide *Attagenus quadricolor* (SUMAKOV, 1907:15).

Phradonoma variabilis (BALLION, 1878:277) DALLA TORRE, 1911:66 — vide *Trogoderma variable* BALLION, 1878:277.

Genus: *Globicornis* LATREILLE, 1829 : 511

Type species: *Dermestes rufitarsis* CREUTZER in PANZER, 1796b : H. 35, t. 6.

Subgenus: *Elania* MULSANT et REY, 1868 : 111

Type species: *Hadrotoma depressa* MULSANT et REY, 1868 : 111.

List of species:

Globicornis depressa (MULSANT et REY, 1868 : 111) REITTER, 1891 : 169; —

Syn.: *Hadrotoma depressa* MULSANT et REY, 1868:111.

Globicornis vaulogeri PIC, 1900 : 53. —

Globicornis vaulogeri var. *brunneithorax* PIC, 1900 : 53. —

Subgenus: *Globicornis* s. str.

List of species:

Globicornis alpina PIC, 1912 : 75. —

Globicornis bicruciata SAHLBERG, 1913 : 28. —

<i>Globicornis bifasciata</i> (PERRIS, 1866 : 186) REITTER, 1906 : 379.	France, Sicily, Algeria, Tunisia
Syn.: <i>Hadrotoma bifasciata</i> PERRIS, 1866:186.	
<i>Globicornis breviclavis</i> (REITTER, 1878b : 179) REITTER, 1891 : 169.	Caucasus
Syn.: <i>Hadrotoma breviclavis</i> REITTER, 1878b:179.	
<i>Globicornis fasciata</i> (FAIRMAIRE, 1859 : 45) REITTER, 1891 : 169.	France, Belgium, Germany, Switzerland, Sardinia
Syn.: <i>Hadrotoma fasciata</i> FAIRMAIRE, 1859:45.	
<i>Globicornis nigripes</i> (FABRICIUS, 1792 : 233) REITTER, 1891 : 169.	Germany, whole Europe, Caucasus
Syn.: <i>Dermestes nigripes</i> FABRICIUS, 1792:233 nec OLIVIER, 1790 no. 9 p. 15 — name entered on the Official List of Specific Names in Zoology.	
<i>Dermestes rufitarsis</i> CREUTZER in PANZER, 1796b:H. 35, t. 6. — LATREILLE, 1807:35.	
<i>Megatoma rufitarse</i> (CREUTZER in PANZER, 1796b:H. 35, t. 6) LATREILLE, 1807:35.	
<i>Attagenus nigripes</i> (FABRICIUS, 1792:233) DEJEAN, 1821:46.	
<i>Dermestes plantaris</i> CURTIS, 1838:682. — STEPHENS, 1839:143.	
<i>Globicornis rufitarsis</i> (CREUTZER in PANZER, 1796b:H. 35, t. 6) LATREILLE, 1829:511.	
<i>Tiresias plantaris</i> (CURTIS, 1838:682) STEPHENS, 1839:143.	
<i>Hadrotoma nigripes</i> (FABRICIUS, 1792:233) ERICHSON, 1846:446.	
<i>Globicornis picta</i> (KÜSTER, 1851 : 39) REITTER, 1891 : 169.	Dalmatia, whole Balkan Peninsula, Cyprus, Italy, Sicily, Asia Minor, Caucasus
Syn.: <i>Trogoderma picta</i> KÜSTER, 1851:39. <i>Hadrotoma picta</i> (KÜSTER, 1851:39) SCHAUM, 1859:45.	
<i>Globicornis tristis</i> (REITTER, 1881b:84) REITTER, 1891:169.	Greece
Syn.: <i>Hadrotoma tristis</i> REITTER, 1881b:84.	
<i>Globicornis variegata</i> (KÜSTER, 1851 : 40) REITTER, 1891 : 169.	Sardinia, Sicily, Corsica, France, Italy, Crete
Syn.: <i>Trogoderma variegata</i> KÜSTER, 1851:40. <i>Hadrotoma variegata</i> (KÜSTER, 1851:40) KRAATZ, 1858:146. <i>Attagenus siculus</i> REITTER, 1881b:76. — REITTER, 1891:169.	

Subgenus: *Hadrotoma* ERICHSON, 1846 : 424Type species: *Dermestes marginatus* PAYKULL, 1798 : 280.Syn.: *Mesalia* MULSANT et REY, 1868:106. — Type species: *Mesalia guillebelli* MULSANT et REY, 1868:109.

List of species:

<i>Globicornis antoniae</i> (REITTER, 1884 : 7) REITTER, 1891 : 169.	Caucasus
Syn.: <i>Hadrotoma antoniae</i> REITTER, 1883:89 — nomen nudum. <i>Hadrotoma antoniae</i> REITTER, 1884:7.	
<i>Globicornis argentina</i> (PIC, 1906 : 233) DALLA TORRE, 1911 : 63.	Argentina
Syn.: <i>Hadrotoma argentina</i> PIC, 1906:233.	
<i>Globicornis bitaeniata</i> (STEINHEIL, 1869 : 258) DALLA TORRE, 1911 : 63.	Argentina
Syn.: <i>Hadrotoma bitaeniata</i> STEINHEIL, 1869:258.	
<i>Globicornis boliviiana</i> PIC, 1937d : 196.	Bolivia
<i>Globicornis clavata</i> (REITTER, 1881 : 37) DALLA TORRE, 1911 : 63.	Brazil
Syn.: <i>Hadrotoma clavata</i> REITTER, 1881:37.	
<i>Globicornis corticalis</i> (EICHHOFF, 1863 : 437) REITTER, 1891 : 169.	Germany, Europe, Kazakh SSR
Syn.: <i>Hadrotoma corticalis</i> EICHHOFF, 1863:437.	
<i>Globicornis corticalis</i> var. <i>semilimbata</i> (PIC, 1906b : 9).	Greece
Syn.: <i>Hadrotoma corticalis</i> var. <i>semilimbata</i> PIC, 1906b:9.	
<i>Globicornis hispanica</i> PIC, 1908d : 81.	Spain
<i>Globicornis latenotata</i> PIC, 1915b : 4.	Argentina
<i>Globicornis marginata</i> (PAYKULL, 1798 : 280) REITTER, 1891 : 169.	Sweden, Europe, Caucasus, West Siberia
Syn.: <i>Dermestes marginatus</i> PAYKULL, 1798:280. <i>Dermestes emarginatus</i> GYLLENHAL, 1808:153. — SCHÖNHERR, 1808:94.	
<i>Attagenus marginatus</i> (PAYKULL, 1798:280) DEJEAN, 1821:46.	
<i>Attagenus emarginatus</i> (GYLLENHAL, 1808:153) STURM, 1826:98.	
<i>Megatoma marginata</i> (PAYKULL, 1798:280) CRISTOFORI et JAN, 1832:30.	
<i>Hadrotoma marginata</i> (PAYKULL, 1798:280) ERICHSON, 1846:445.	
<i>Mesalia guillebelli</i> MULSANT et REY, 1868:109. — REITTER, 1881b: 82.	
<i>Globicornis rufoguttata</i> PIC, 1899c : 254.	Syria
<i>Globicornis rufoguttata</i> var. <i>signatipennis</i> PIC, 1899 c: 255.	Syria
<i>Globicornis sulcata</i> (BRISOUT, 1866 : 371) REITTER, 1891 : 169.	Spain, France
Syn.: <i>Hadrotoma sulcata</i> BRISOUT, 1866:371.	
<hr/>	
<i>Globicornis angustata</i> (SOLSKI, 1876:271) HEYDEN, 1893:64 — vide <i>Attagenus augustatus</i> BALLION, 1871: 330.	
<i>Hadrotoma bodemeyeri</i> GANGLBauer in BODEMEYER, 1900:148 — vide <i>Dearthrus bodemeyeri</i> (GANGLBauer in BODEMEYER, 1900:148).	
<i>Globicornis cervinus</i> STURM, 1843:319 — nomen nudum.	

- Globicornis fulvipes* GUÉRIN-MÉNEVILLE, 1838c:138 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c: 138).
- Hadrotoma glabra* (SAHLBERG, 1819:50) SCHAUM, 1852:38 — vide *Megatoma undata* (LINNAEUS, 1758:355).
- Mesalia guillebelli* MULSANT et REY, 1868:109 — vide *Globicornis marginata* (PAYKULL, 1798:280).
- Globicornis kulti* KALIK, 1951c:1 — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).
- Hadrotoma longulus* (LECONTE, 1863:73) ZIMMERMANN, 1869:258 — vide *Dearthrus longulus* LECONTE, 1863:73.
- Globicornis obenbergeri* KALIK, 1951c:2 — vide *Megatoma obenbergeri* (KALIK, 1951c:2).
- Globicornis picta*: KLEINE, 1940: 17 nec (KÜSTER, 1851:39) — vide *Trogoderma angustum* (SOLIER, 1849: 374).
- Hadrotoma pubescens*: LEDER, 1881:515 nec (ZETTERSTEDT, 1828:136) — vide *Megatoma riedeli* MROCKOWSKI, 1967:11.
- Hadrotoma pubescens* (ZETTERSTEDT, 1828:136) REITTER, 1880b:510 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Hadrotoma quadriguttata* REITTER, 1878:95 — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).
- Globicornis quadriguttata subfasciata* DALLA TORRE, 1911:64 — vide *Dearthrus quadriguttatus unifasciatus* (PIC, 1908b:41).
- Globicornis quadriguttata unifasciata* PIC, 1908b:41 — vide *Dearthrus quadriguttatus unifasciatus* (PIC, 1908b:41).
- Globicornis quadrinacra* DALLA TORRE, 1911:64 — vide *Dearthrus quadrinaevus* (REITTER, 1908b:213).
- Globicornis quadrinaeva* REITTER, 1908b:213 — vide *Dearthrus quadrinaevus* (REITTER, 1908b:213).
- Globicornis ruftarsis* (CREUTZER in PANZER, 1796b:H. 35, t. 6) LATREILLE, 1829:511 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Globicornis sunica* JABLOKOV-CHNZORJAN, 1961:49, nomen nudum — vide *Dearthrus quadriguttatus* (REITTER, 1878:95).

Genus: *Hirtomegatoma* PIC, 1931 : 5

Type species: *Hirtomegatoma infasciata* PIC, 1931 : 5.

List of species:

Hirtomegatoma infasciata PIC, 1931 : 5.

Jordan

Genus: *Megatoma* HERBST, 1792 : 92

Type species: *Dermestes undatus* LINNAEUS, 1758 : 355.

Syn.: *Asprogramme* GOZIS, 1886:10. — Type species: *Dermestes undatus* LINNAEUS, 1758:355.
Aprogramme REITTER, 1887b:40. — Type species: *Dermestes undatus* LINNAEUS, 1758:355.
Approgramme REITTER, 1887b:55. — Type species: *Dermestes undatus* LINNAEUS, 1758:355.

Subgenus: *Caucasotoma* MROCKOWSKI, 1967 : 5

Type species: *Megatoma riedeli* MROCKOWSKI, 1967 : 11.

List of species:

Megatoma riedeli MROCKOWSKI, 1967 : 11.

Caucasus

Syn.: *Hadrotoma pubescens*: LEDER, 1881:515 nec (ZETTERSTEDT, 1828: 136). — MROCKOWSKI, 1967:12.

Megatoma pubescens: ZAJCEV, 1919:170 nec (ZETTERSTEDT, 1828:136). — MROCKOWSKI, 1867:12.

Megatoma undata: EICHLER, 1930:234 nec (LINNAEUS, 1758: 355). — MROCKOWSKI, 1967:12.

Subgenus: *Megatoma* s. str.

List of species:

Megatoma pubescens (ZETTERSTEDT, 1828 : 136) THOMSON,
1862 : 204.

Syn.: *Dermestes pubescens* ZETTERSTEDT, 1828:136.

Attagenus fuscus DEJEAN, 1837:139, nomen nudum — GEBLER,
1847:446.

Attagenus fuscus GEBLER, 1841:585. — MROCKOWSKI, 1967:9.

Megatoma fusca (GEBLER, 1841:585) GEMMINGER et HAROLD,
1868:917.

Hadrotoma pubescens (ZETTERSTEDT, 1828:136) REITTER, 1880b:
510.

Approgramme pubescens (ZETTERSTEDT, 1828:136) REITTER, 1887b:
55.

Megatoma innotata PIC, 1902b:48. — JANSSON, 1943:23.

Megatoma prolixa BEAL, 1967:299. — syn. nov.

Lapland,
Northern part
of Palearctic
to Amur river,
British
Columbia,
California

Megatoma ruficornis AUBÉ, 1866 : 161.

Syn.: *Megatoma rufovittata* ABEILLE, 1870:XLVI. — MROCKOWSKI,
1967:7.

Approgramme ruficornis (AUBÉ, 1866:161) REITTER, 1887b:55.

Megatoma pici KALIK, 1952:76. — MROCKOWSKI, 1967:7.

France, Italy⁴
Yugoslavia,
Hungary,
Greece, Turkey

Megatoma undata (LINNAEUS, 1758 : 355) LATREILLE, 1807 : 34.

Syn.: *Dermestes undatus* LINNAEUS, 1758:355.

Dermestes undulatus LINNAEUS, 1761:141. — LINNAEUS, 1767:
562.

Megatoma undulata HERBST, 1792:96. — HERBST, 1792:131.

Attagenus undatus (LINNAEUS, 1758:355) LATREILLE, 1804:243.

Dermestes glaber SAHLBERG, 1819:50. — GEMMINGER et HAROLD,
1868:917.

Attagenus undulatus (LINNAEUS, 1761:141) STEPHENS, 1829:8.

Hadrotoma glabra (SAHLBERG, 1819:50) SCHAUM, 1852:38.

Approgramme undata (LINNAEUS, 1758:355) REITTER, 1887b:55.

Europe, Siberia

Megatoma undata var. *unifasciata* GANGLBAUER, 1904b : 28.

?

Megatoma undata ussuriensis MROCKOWSKI, 1967 : 7.

Ussuri

Subgenus: *Pseudohadrotoma* KALIK, 1951c : 2

Type species: *Globicornis obenbergeri* KALIK, 1951c : 2.

List of species:

Megatoma conspersa SOLSKIJ, 1876 : 275.

Syn.: *Approgramme maculosa* REITTER, 1887b:55. — WINKLER, 1926:
678.

Megatoma maculosa (REITTER, 1887b:55) REITTER, 1891:169.

Megatoma m-notata PIC, 1938:1. — MROCKOWSKI, 1967:12.

Uzbek SSR,
Kirghiz SSR,
Kazakh SSR,
China:
Tyan-shan

Megatoma falsa (HORN, 1875 : 135) MROCKOWSKI, 1967 : 14.

Syn.: *Perimegatoma falsum* HORN, 1875:135.

Megatoma cylindrica: SPENCER, 1956:32. — BEAL, 1967:297.

Pseudohadrotoma falsa (HORN, 1875:135) BEAL, 1967:293.

California,
Idaho, Oregon

Megatoma graeseri (REITTER, 1887b : 56) HEYDEN, 1893 : 64.

Syn.: *Approgramme graeseri* REITTER, 1887b:56.

East Asia:
from Kirghiz
SSR over
Transbaicalia
and Mongolia
up to Amur

Megatoma graeseri var. *latenotata* PIC, 1902b : 48.

Megatoma kaliki (BEAL, 1967 : 293) MROCKOWSKI, 1967 : 14.

Syn.: *Pseudohadrotoma kaliki* BEAL, 1967:293.

Megatoma obenbergeri (KALIK, 1951c : 2) MROCKOWSKI,
1967 : 13.

Syn.: *Globicornis obenbergeri* KALIK, 1951c:2.

Pseudohadrotoma obenbergeri (KALIK, 1951c:2) BEAL, 1967:292.

British
Columbia

Megatoma perversa (FALL, 1926 : 194) MROCKOWSKI, 1967 :
14.

Syn.: *Perimegatoma perversa* FALL, 1926:194.

Pseudohadrotoma perversa (FALL, 1926:194) BEAL, 1967:293.

Yukon
Territory,
Alaska,
Alberta, British
Columbia,
California,
Montana

Megatoma trichorhopalum (BEAL, 1967 : 293) MROCKOWSKI,
1967 : 14.

Syn.: *Pseudohadrotoma trichorhopalum* BEAL, 1967:293.

California

Megatoma trogodermoides (BEAL, 1967 : 293) MROCKOWSKI,
1967 : 14.

Syn.: *Pseudohadrotoma trogodermoides* BEAL, 1967:293.

California,
Nevada

Subgenus: *Perimegatoma* HORN, 1875 : 135

Type species: *Perimegatoma variegata* HORN, 1875 : 135.

List of species:

Megatoma ampla (CASEY, 1900 : 150) BEAL, 1967 : 299.

Syn.: *Perimegatoma ampla* CASEY, 1900:150.

California,
Washington,
British
Columbia

Megatoma belfragei (LECONTE, 1874 : 50) BEAL, 1957 : 299.

Syn.: *Trogoderma belfragei* LECONTE, 1874:50.

Perimegatoma belfragei (LECONTE, 1874:50) HORN, 1875:135.

Texas

Megatoma cylindrica (KIRBY, 1837 : 113) CROTCH, 1873 : 41.

Syn.: *Attagenus cylindricus* KIRBY, 1837:113.

Perimegatoma cylindricum (KIRBY, 1837:113) HORN, 1875:135.

Perimegatoma impressa CASEY, 1900:150. — HOPPING, 1932:173.

Perimegatoma monticola CASEY, 1900:150. — HOPPING, 1932: 173.

Rocky Mts.;
Western USA,
Western
Canada

Megatoma cylindrica var. *angularis* (MANNERHEIM, 1853 : 216). — JAYNE, 1882 : 359.

Alaska

Syn.: *Attagenus angularis* MANNERHEIM, 1853:216.

Perimegatoma angularis (MANNERHEIM, 1853:216) CASEY, 1900:151.

Megatoma dolia BEAL, 1967 : 300.

California

Megatoma giffardi (BLAISDELL, 1927 : 163) BEAL, 1967 : 300.

California,

Syn.: *Perimegatoma giffardi* BLAISDELL, 1927:163.

Oregon,
Montana,
Washington,
Alaska

Megatoma leucochlidon BEAL, 1967 : 300.

California,
Oregon, British
Columbia

Megatoma polia BEAL, 1967 : 299.

California

Megatoma variegata (HORN, 1875 : 135) HINTON, 1945b : 366.

Oregon,

Syn.: *Perimegatoma variegatum* HORN, 1875:135.

California,

Perimegatoma jaynei CASEY, 1900:150. — BEAL, 1967:300.

Guadalupe,

Perimegatoma guadalupensis CASEY, 1900:150. — BEAL, 1967: 300.

Nevada, Idaho,
Utah, British
Columbia

Perimegatoma nevadica CASEY, 1900:150. — BEAL, 1967:300.

Species incertae sedis

Megatoma foveolata LEPESME, 1942 : 142.

Australia

Megatoma tenuefasciata REITTER, 1881 : 36.

Tasmania

Megatoma adspersa BLANCHARD, 1843:61 — vide *Attagenus adspersus* (BLANCHARD, 1843:61).

Perimegatoma angularis (MANNERHEIM, 1853:216) CASEY, 1900:151 — vide *Megatoma cylindrica* var. *angularis* (MANNERHEIM, 1853:216).

Megatoma apicalis MACLEAY, 1871:170 — vide *Trogoderma apicalis* (MACLEAY, 1871:170).

Megatoma atra HERBST, 1792:95 — vide *Attagenus pellio* (LINNAEUS, 1758:355).

Megatoma bifasciata (ROSSI, 1794:79) REITTER, 1887b:48 — vide *Attagenus bifasciatus* var. *rossi* GANGLBauer, 1904b:26.

Megatoma bifasciata (THUNBERG, 1781:9) THUNBERG, 1820:372 — non Dermestidae, familia?

Megatoma brevicornis HERBST, 1792:95 — vide *Attagenus unicolor* (BRAHM, 1791:144).

Megatoma calabrica (REITTER, 1881b:77) REITTER, 1887b:48 — vide *Attagenus calabricus* REITTER, 1881b: 77.

- Megatoma cinnamomea* (ROTH, 1851:122) REITTER, 1887b:50 — vide *Attagenus cinnamomeus* ROTH, 1851:122.
- Megatoma cylindrica*: SPENCER, 1956:32 — vide *Megatoma falsa* (HORN, 1875:135).
- Megatoma fallax* (GENÉ, 1839:19) REITTER, 1887b:48 — vide *Attagenus fallax* GENÉ, 1839:19.
- Megatoma fallax* var. *distincta*: REITTER, 1887b:48 — vide *Attagenus fallax* var. *distinctus* MULSANT et REY, 1868:85.
- Megatoma fusca* (GEBLER, 1841:585) GEMMINGER et HAROLD, 1868:917 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Megatoma hottentotum* GUÉRIN-MÉNEVILLE, 1838b:65 — vide *Attagenus hottentotus* (GUÉRIN-MÉNEVILLE, 1838b:65).
- Perimegatoma impressa* CASEY, 1900:150 — vide *Megatoma cylindrica* (KIRBY, 1837:113).
- Megatoma innotata* PIC, 1902b:48 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Megatoma maculosa* (REITTER, 1887b:55) REITTER, 1891:169 — vide *Megatoma conspersa* SOLSKI, 1876: 275.
- Megatoma marginata* (PAYKULL, 1798:280) CRISTOFORI et JAN, 1832:30 — vide *Globicornis marginata* (PAYKULL, 1798:280).
- Megatoma marginicollis* (KÜSTER, 1847b:76) REITTER, 1887b:47 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Megatoma m-notata* PIC, 1938:1 — vide *Megatoma conspersa* SOLSKI, 1876:275.
- Perimegatoma monticola* CASEY, 1900:150 — vide *Megatoma cylindrica* (KIRBY, 1837:113).
- Megatoma morio* ERICHSON, 1842:152 — vide *Trogoderma morio* (ERICHSON, 1842:152).
- Megatoma ornata* SAY, 1825:185 — vide *Trogoderma ornatum* (SAY, 1825:185).
- Megatoma pantherina* (AHRENS, 1814:11) REITTER, 1887b:48 — vide *Attagenus pantherinus* (AHRENS, 1814:11).
- Megatoma pellio* (LINNAEUS, 1758:355) CRISTOFORI et JAN, 1832:30 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Megatoma picea* HERBST, 1792:97 — non *Dermestidae*, recte *Serropalpidae*: *Orchesia micans* (PANZER, 1794).
- Megatoma picea* (OLIVIER, 1790: no. 9 p. 10) REITTER, 1887b:46 — vide *Attagenus unicolor* (BRAHM, 1791: 144).
- Megatoma picea* var. *dalmatina*: REITTER, 1887b:46 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Megatoma picea* var. *sordida*: REITTER, 1887b:47 — vide *Attagenus brunneus* FALDERMANN, 1835:227.
- Megatoma pici* KALIK, 1952:76 — vide *Megatoma ruficornis* AUBÉ, 1866:161.
- Megatoma prolixa* BEAL, 1967:299 — vide *Megatoma pubescens* (ZETTERSTEDT, 1828:136).
- Megatoma pubescens*: ZAJCEV, 1919:170 nec (ZETTERSTEDT, 1828:136) — vide *Megatoma riedeli* MROCKOWSKI, 1967:11.
- Megatoma quadrimaculata* (KRAATZ, 1858:145) REITTER, 1887b:50 — vide *Attagenus quadrimaculatus* KRAATZ, 1858:145.
- Megatoma rufipes* CRISTOFORI et JAN, 1832:30 — nomen nudum.
- Megatoma rufitarse* (CREUTZER in PANZER, 1796b:H. 35, p. 6) LATREILLE, 1807:35 — vide *Globicornis nigripes* (FABRICIUS, 1792:233).
- Megatoma rufovittata* ABEILLE, 1870:XLVI — vide *Megatoma ruficornis* AUBÉ, 1866:161.
- Megatoma schaefferi* HERBST, 1792:93 — vide *Attagenus schaefferi* (HERBST, 1792:93).
- Megatoma schrankii* KUGELANN, 1792:480 — vide *Attagenus pellio* (LINNAEUS, 1758:355).
- Megatoma sericeum* GUÉRIN-MÉNEVILLE, 1838b:65 — vide *Attagenus sericeus* (GUÉRIN-MÉNEVILLE, 1838 b: 65).
- Megatoma serra* (FABRICIUS, 1792:234) LATREILLE, 1807:35 — vide *Ctesias serra* (FABRICIUS, 1792:234).
- Megatoma simonis* (REITTER, 1881:34) REITTER, 1887b:49 — vide *Attagenus bifasciatus* var. *simonis* REITTER, 1881:34.
- Megatoma simplex* (REITTER, 1881b:79) REITTER, 1887b:51 — vide *Attagenus simplex* REITTER, 1881 b: 79.

- Megatoma tigrina* (FABRICIUS, 1792:229) REITTER, 1887b:49 — vide *Attagenus bifasciatus* var. *tigrinus* (FABRICIUS, 1792:229).
- Megatoma tigrina* var. *persica*: REITTER, 1887b:49 — vide *Attagenus bifasciatus* var. *persicus* REITTER, 1881b:78.
- Megatoma tigrina* var. *pulcher*: REITTER, 1887b:49 — vide *Attagenus bifasciatus* (OLIVIER, 1790:no. 9 p. 13).
- Megatoma tigrina* var. *wachanrui*: REITTER, 1887b:50 — vide *Attagenus bifasciatus* (OLIVIER, 1790:no. 9 p. 13).
- Megatoma trifasciatum* (FABRICIUS, 1787:34) CRISTOFORI et JAN, 1832:30 — vide *Attagenus trifasciatus* (FABRICIUS, 1787:34).
- Megatoma undata*: EICHLER, 1930:234 nec (LINNAEUS, 1758:355) — vide *Megatoma riedeli* MROCKOWSKI, 1967:11.
- Megatoma undulata* HERBST, 1792:96 — vide *Megatoma undata* (LINNAEUS, 1758:355).
- Megatoma varius* MATSUMURA et YOKOYAMA, 1928b:51 — vide *Trogoderma varium* (MATSUMURA et YOKOYAMA, 1928b:51).
- Perimegatoma vespulae* MILLIRON, 1939:570 — vide *Reesa vespulae* (MILLIRON, 1939:570).
- Megatoma vigintiguttata* (FABRICIUS, 1775:56) REITTER, 1887b:47 — vide *Attagenus punctatus* (SCOPOLI, 1772:87).
- Megatoma vigintipunctatum* CRISTOFORI et JAN, 1832:30, nomen nudum — vide *Attagenus punctatus* (SCOPOLI, 1772:87).

Genus: *Labrocerus* SHARP in BLACKBURN et SHARP, 1885 : 148

Type species: *Labrocerus jaynei* SHARP in BLACKBURN et SHARP, 1885 : 148.

List of species:

<i>Labrocerus affinis</i> SHARP, 1908 : 410.	Hawaii
<i>Labrocerus concolor</i> SHARP in BLACKBURN et SHARP, 1885 : 149.	Hawaii
<i>Labrocerus curticornis</i> SHARP, 1908 : 407.	Hawaii
<i>Labrocerus dasytoides</i> SHARP, 1908 : 409.	Hawaii
<i>Labrocerus flavigornis</i> SHARP, 1908 : 410.	Hawaii
<i>Labrocerus gravidus</i> SHARP, 1908 : 407.	Hawaii
<i>Labrocerus jaynei</i> SHARP in BLACKBURN et SHARP, 1885 : 148.	Hawaii
<i>Labrocerus laticornis</i> SHARP, 1908 : 408.	Hawaii
<i>Labrocerus moerens</i> SHARP, 1908 : 406.	Hawaii
<i>Labrocerus obscurus</i> BLACKBURN in BLACKBURN et SHARP, 1885 : 149.	Hawaii
<i>Labrocerus obsoletus</i> SHARP, 1908 : 409.	Hawaii
<i>Labrocerus pallipes</i> SHARP, 1908 : 410.	Hawaii
<i>Labrocerus quadrisignatus</i> SHARP, 1908 : 411.	Hawaii
<i>Labrocerus setosus</i> SHARP, 1908 : 410.	Hawaii
<i>Labrocerus simplex</i> SHARP, 1908 : 407.	Hawaii
<i>Labrocerus suffusus</i> SHARP, 1908 : 411.	Hawaii
<i>Labrocerus vestitus</i> SHARP, 1908 : 409.	Hawaii

Genus: *Eocerus* SHARP, 1908 : 412

Type species: *Eocerus depressus* SHARP, 1908 : 412.

List of species:

Eocerus depressus SHARP, 1908 : 412.

Hawaii

Genus: *Argocerus* SHARP, 1908 : 411

Type species: *Argocerus similaris* SHARP, 1908 : 411.

List of species:

Argocerus similaris SHARP, 1908 : 411.

Hawaii

Argocerus subguttatus SHARP, 1908 : 412.

Hawaii

Genus: *Dearthrus* LECONTE, 1861 : 108

Type species: *Dearthrus longulus* LECONTE, 1863 : 73.

Syn.: *Dearthus* JAYNE, 1882:346. — Type species: *Dearthus longulus* LECONTE, 1863:73.

Pseudomesalia GANGLBAUER in BODEMEYER, 1900:149. — BEAL, 1967:290. — Type species: *Hadrotoma bodemeyeri* GANGLBAUER in BODEMEYER, 1900:148.

List of species:

Dearthrus bodemeyeri (GANGLBAUER in BODEMEYER, 1900 : 148)
BEAL, 1967 : 291.

Turkey, Syria

Syn.: *Hadrotoma bodemeyeri* GANGLBAUER in BODEMEYER, 1900:148.
Globicornis bodemeyeri (GANGLBAUER in BODEMEYER, 1900:148)
DALLA TORRE, 1911:63.

Dearthrus longulus LECONTE, 1863 : 73.

Georgia,
Pennsylvania,
Illinois,
Indiana,
Michigan,
Ohio

Syn.: *Hadrotoma longulus* (LECONTE, 1863:73) ZIMMERMANN, 1869:258.
Attagenus longulus (LECONTE, 1863:73) CROTCH, 1873:41.

Caucasus,
Transcaucasia,
Asia Minor,
Turkmen SSR,
Afghanistan,
Uzbek SSR,
Tadzhik SSR

Dearthrus quadriguttatus (REITTER, 1878 : 95) BEAL, 1967 : 291.

Syn.: *Trogoderma ornata* SOLSKIJ, 1876:278 nec (SAY, 1825:186). — MROCKOWSKI, 1960b:212.

Hadrotoma quadriguttata REITTER, 1878:95.

Trogoderma solskyi REITTER, 1881:38. — MROCKOWSKI, 1960b: 212.

Globicornis quadriguttata (REITTER, 1878:95) REITTER, 1891:169.

Phradonoma ornatum (SOLSKIJ, 1876:278) DALLA TORRE, 1911:66.

Globicornis kulti KALIK, 1951c:1. — MROCKOWSKI, 1960b:212.

Globicornis sunica JABLOKOV-CHNZORJAN, 1961:49, nomen nudum — ZHANTIEV in litt.

<i>Dearthrus quadriguttatus unifasciatus</i> (PIC, 1908b : 41) comb. nov.	Syria
Syn.: <i>Globicornis quadriguttata unifasciata</i> PIC, 1908b:41. <i>Globicornis quadriguttata subfasciata</i> DALLA TORRE, 1911: 64. — syn. nov.	
<i>Dearthrus quadrinaevus</i> (REITTER, 1908b : 213) comb. nov.	Kazakh SSR
Syn.: <i>Globicornis quadrinaeva</i> REITTER, 1908b:213. <i>Globicornis quadrinacra</i> DALLA TORRE, 1911:64. — MROCKOWSKI, 1962:247.	
<i>Dearthrus stebbinsi</i> BEAL, 1954 : 232.	California, Oregon

Genus: *Ditorines* NORMAND, 1936 : 298

Type species: *Ditorines denticornis* NORMAND, 1936 : 299.
List of species:

Ditorines denticornis NORMAND, 1936 : 299.

Tunisia

Genus: *Caccoleptus* SHARP, 1902 : 649

Type species: *Caccoleptus rotundus* SHARP, 1902 : 650.
List of species:

Caccoleptus anisotomoides SHARP, 1902 : 650.

British
Honduras
Panama

Caccoleptus rotundus SHARP, 1902 : 650.

Genus: *Diontolobus* SOLIER, 1849 : 367

Type species: *Diontolobus punctipennis* SOLIER, 1849 : 368.
Syn.: *Distontobus* SOLIER, 1854:t. 8, f. 4. — Type species: *Diontolobus punctipennis* SOLIER, 1849:368.
Diodontolobus LACORDAIRE, 1854:458. — Type species: *Diontolobus punctipennis* SOLIER, 1849:368.
Didontolobus FAIRMAIRE, 1884:488. — Type species: *Diontolobus punctipennis* SOLIER, 1849:368.

List of species:

Diontolobus lateritius (FAIRMAIRE, 1884 : 488) comb. nov.

Chile

Syn.: *Didontolobus lateritius* FAIRMAIRE, 1884:488.

Diodontolobus lateritius (FAIRMAIRE, 1884:488) DALLA TORRE,
1911:88.

Diontolobus punctipennis SOLIER, 1849 : 368.

Chile

Syn.: *Distontobus punctipennis* (SOLIER, 1849:368) SOLIER, 1854:t. 8, f. 4.

Diodontolobus punctipennis (SOLIER, 1849:368) LACORDAIRE, 1854:
461.

Didontolobus punctipennis (SOLIER, 1849:368) BLACKWELDER,
1945:398.

Genus: *Anthrenocerus* ARROW, 1915 : 443

Type species: *Anthrenus australis* HOPE, 1843 : 319.

List of species:

Anthrenocerus arrowi ARMSTRONG, 1949 : 109.

Syn.: *Anthrenocerus bicolor* ARMSTRONG, 1943:61 nec ARROW, 1915: 444.

Anthrenocerus australis (HOPE, 1843 : 319) ARROW, 1915 : 444.

Syn.: *Anthrenus australis* HOPE, 1843:319.

Anthremis australis HOPE, 1845:105. — ARMSTRONG, 1943:60.

Cryptorhopalum erichsoni REITTER, 1881:55. — ARMSTRONG, 1943: 58.

Cryptorhopalum australe (HOPE, 1843:319) BLACKBURN, 1903: 167.

Anthrenocerus bicolor ARROW, 1915 : 444.

Anthrenocerus blackburni ARMSTRONG, 1943 : 58.

Anthrenocerus chalceous ARMSTRONG, 1943 : 57.

Anthrenocerus concolorous ARMSTRONG, 1943 : 57.

Anthrenocerus condensus ARMSTRONG, 1943 : 57.

Anthrenocerus confertus (REITTER, 1881 : 55) ARROW, 1915 : 444.

Syn.: *Cryptorhopalum confertum* REITTER, 1881:55.

Anthrenus flinders BLACKBURN, 1891:132. — BLACKBURN, 1903: 168.

Cryptorhopalum flindersi (BLACKBURN, 1891:132) BLACKBURN, 1903:168.

Anthrenocerus convexus ARMSTRONG, 1943 : 58.

Anthrenocerus maculosus ARMSTRONG, 1943 : 57.

Anthrenocerus niger ARMSTRONG, 1943 : 57.

Anthrenocerus pulchellus ARROW, 1915 : 444.

Anthrenocerus quadrifasciatus (BLACKBURN, 1903 : 169) ARROW, 1915 : 444.

Syn.: *Cryptorhopalum quadrifasciatum* BLACKBURN, 1903:169.

New South
Wales,
Queensland,
West Australia

Adelaide, whole
Australia,
New Zealand,
introduced
to Europe
(Britain,
Holland)

N. W. Australia

Victoria, New
South Wales

New South
Wales

New South
Wales

New South
Wales

S. Australia

Cent. Australia

Northern
Territory

New South
Wales

N.W. Australia,
Northern
Territory

Central
Australia

<i>Anthrenocerus riguus</i> (ERICHSON, 1842 : 152) ARROW, 1915 :	Tasmania
444.	
Syn.: <i>Trogoderma riguum</i> ERICHSON, 1842:152.	
<i>Anthrenocerus signatus</i> ARMSTRONG, 1943 : 57.	North Queensland
<i>Anthrenocerus stigmacrophilus</i> ARMSTRONG, 1949 : 107.	New South Wales
<i>Anthrenocerus terzonatus</i> (BLACKBURN, 1903 : 169) ARROW, 1915 : 444.	Queensland
Syn.: <i>Cryptorhopalum terzonatum</i> BLACKBURN, 1903:169.	
<i>Anthrenocerus trimaculatus</i> ARMSTRONG, 1943 : 57.	New South Wales
<i>Anthrenocerus variabilis</i> (REITTER, 1881 : 55) ARROW, 1915 : 444.	South Australia, New Caledonia
Syn.: <i>Cryptorhopalum variable</i> REITTER, 1881:55.	

Anthrenocerus bicolor ARMSTRONG, 1943:61 nec ARROW, 1915:444 — vide *Anthrenocerus arrowi* ARMSTRONG, 1949:109.

Genus: *Myrmeanthrenus* ARMSTRONG, 1945 : 47

Type species: *Myrmeanthrenus frontalis* ARMSTRONG, 1945 : 49.

List of species:

Myrmeanthrenus frontalis ARMSTRONG, 1945 : 49. Victoria

Genus: *Adelaidia* BLACKBURN, 1891 : 130

Type species: *Adelaidia rigua* BLACKBURN, 1891 : 130.

Syn.: *Adelaidea* BLACKBURN, 1903:159.

List of species:

Adelaidia rigua BLACKBURN, 1891 : 130.

South
Australia, New
South Wales

Adelaidia unicolor MROCKOWSKI, 1966 b : 326.

Philippines

Genus: *Orphinus* MOTSCHULSKY, 1858 : 48

Type species: *Orphinus haemorrhoidalis* MOTSCHULSKY, 1858 : 48.

Syn.: *Aethriosia* REITTER, 1908:45 — ARROW, 1915:437. Type species: *Aethriosia globulicornis* REITTER, 1908:46.

Subgenus: *Falsoorphinus* PIC, 1931b : 4Type species: *Falsoorphinus notatipennis* PIC, 1931b : 4.

List of species:

Orphinus kenyensis KALIK, 1965 : 220.

Kenya

Orphinus notatipennis (PIC, 1931b : 4) KALIK, 1965 : 221.

Japan

Syn.: *Falsoorphinus notatipennis* PIC, 1931b:4.*Orphinus ovalis* (ARROW, 1915 : 436) comb. nov. — KALIK in litt.

China

Syn.: *Thaumaglossa ovalis* ARROW, 1915:436.*Orphinus sexmaculatus* (ARROW, 1915 : 435) comb. nov. — KALIK in litt.

Rhodesia

Syn.: *Thaumaglossa sexmaculata* ARROW, 1915:436.Subgenus: *Orphinus* s. str.

List of species:

Orphinus aethiops ARROW, 1915 : 438.Angola,
Mauritius,
Southern
Nigeria,
Taiwan,
SumatraSyn.: *Orphinus formosanus* PIC, 1918b:2. — FLUITER, 1952:X.

Celebes

Orphinus affinis (REITTER, 1881 : 54) ARROW, 1915 : 438.Syn.: *Cryptorhopalum affine* REITTER, 1881:54.*Orphinus africanus* PIC, 1927e : 6.Haut-Niger,
Congo
(Kinshasa)*Orphinus apicalis* PIC, 1918b : 2.

Java, Sumatra

Orphinus apicalis var. *obliteratus* PIC, 1927 : 66.

Sumatra

Orphinus atrofasciatus PIC, 1916c : 16.

India: Madura

Orphinus atrous ARMSTRONG, 1943 : 62.

Queensland

Orphinus australicus (BLACKBURN, 1891 : 130) ARROW, 1915 : 438.

S. Australia

Syn.: *Cryptorhopalum australicum* BLACKBURN, 1891:130.*Orphinus biflexus* (REITTER, 1881 : 55) ARROW, 1915 : 438.

East India

Syn.: *Cryptorhopalum biflexum* REITTER, 1881:55.*Orphinus binotatus* PIC, 1937c : 6.

Madagascar

Orphinus bipunctatus (sic!) (MATSUMURA et YOKOYAMA, 1928b : 53) comb. nov.

Japan

Syn.: *Orphilus bipunctatus* MATSUMURA et YOKOYAMA, 1928b:53.

<i>Orphinus biroi</i> PIC, 1956 : 76.	New Guinea
<i>Orphinus biroi</i> var. <i>seminotatus</i> PIC, 1956 : 76.	New Guinea
<i>Orphinus casuarinae</i> (BLACKBURN, 1903 : 169) ARMSTRONG, 1943 : 57.	Australia
Syn.: <i>Cryptorhopalum casuarinæ</i> BLACKBURN, 1903:169.	
<i>Orphinus ceciliense</i> (BLACKBURN, 1903 : 170) ARMSTRONG, 1943 : 57.	Australia
Syn.: <i>Cryptorhopalum ceciliense</i> BLACKBURN, 1903:170.	
<i>Orphinus congoanus</i> (PIC, 1950 : 356) KALIK, 1955b : 319.	Congo (Kinshasa)
Syn.: <i>Trododerma</i> (sic!) <i>congoana</i> PIC, 1950:356.	
<i>Orphinus conjunctus</i> PIC, 1937c : 6.	Zanzibar
<i>Orphinus convexus</i> PIC, 1956 : 76.	Singapore
<i>Orphinus defectus</i> (WALKER, 1858 : 207) ARROW, 1915 : 437.	Ceylon, Borneo
Syn.: <i>Attagenus?</i> <i>defectus</i> WALKER, 1858:207.	
Trogoderma <i>defectum</i> (WALKER, 1858:207) OLLIFF, 1883:185.	
<i>Orphinus diversus</i> (PIC, 1928b : 2) comb. nov.	Sumatra
Syn.: <i>Cryptorhopalum diversum</i> PIC, 1928b:2.	
<i>Orphinus fasciatus</i> (MATSUMURA et YOKOYAMA, 1928b : 54) comb. nov.	Taiwan
Syn.: <i>Orphilus fasciatus</i> MATSUMURA et YOKOYAMA, 1928b:54.	
<i>Orphinus fouqueti</i> PIC, 1937c : 6.	Senegal
<i>Orphinus fulvipes</i> (GUÉRIN-MÉNEVILLE, 1838c : 138) HINTON, 1945: 368.	Brazil, Antilles, Northern Australia, Pacific Islands, Java, Madagascar, Bolivia, Centr. America, Mexico, Florida, introduced to Germany and France
Syn.: <i>Globicornis fulvipes</i> GUÉRIN-MÉNEVILLE, 1838c:138.	
Trogoderma <i>fulvipes</i> (GUÉRIN-MÉNEVILLE, 1838c:138) LACOR- DAIRE, 1854:467.	
<i>Hadrotoma fulvipes</i> (GUÉRIN-MÉNEVILLE, 1838c:138) REITTER, 1881:37.	
Trogoderma <i>brasiliensis</i> REITTER, 1881:37. — REITTER, 1881:37.	
<i>Cryptorhopalum brevicorne</i> SHARP in BLACKBURN et SHARP, 1885:150. — HINTON, 1945:368.	
<i>Cryptorhopalum fulvipes</i> (GUÉRIN-MÉNEVILLE, 1838c:138) FLEU- TIAUX et SALLÉ, 1890:392.	
Trogoderma <i>unicolor</i> KOLBE, 1910:24. — SCOTT, 1926:61.	
<i>Orphinus brevicornis</i> (SHARP in BLACKBURN et SHARP, 1885:150) ARROW, 1915:438.	
<i>Orphinus funestus</i> ARROW, 1915 : 441.	Ceylon
<i>Orphinus globulicornis</i> (REITTER, 1908 : 46) ARROW, 1915 : 438.	Egypt
Syn.: <i>Aethriosis globulicornis</i> REITTER, 1908:46.	

<i>Orphinus globulicornis</i> var. <i>boehmi</i> (REITTER, 1908 : 46).	Egypt
Syn.: <i>Aethriosia globulicornis</i> var. <i>boehmi</i> REITTER, 1908:46.	
<i>Orphinus globulicornis</i> var. <i>ferrantei</i> (REITTER, 1908 : 46).	Egypt
Syn.: <i>Aethriosia globulicornis</i> var. <i>ferrantei</i> REITTER, 1908:46.	
<i>Orphinus guernei</i> PIC, 1916c : 16.	Ceylon
<i>Orphinus haemorhoidalis</i> MOTSCHULSKY, 1858 : 48.	Burma
Syn.: <i>Cryptorhopalum motschulskyi</i> REITTER, 1881:55.	
<i>Orphinus horni</i> PIC, 1927c : 247.	Ceylon
<i>Orphinus infasciatus</i> PIC, 1926 : 5.	Philippine Isl.
<i>Orphinus interioris</i> (BLACKBURN, 1891 : 131) ARMSTRONG, 1943 : 57.	S. Australia
Syn.: <i>Cryptorhopalum interioris</i> BLACKBURN, 1891:131.	
<i>Orphinus japonicus</i> ARROW, 1915 : 438.	Japan, Korea
<i>Orphinus jucundus</i> ARROW, 1915 : 439.	India: Bombay
<i>Orphinus madecassus</i> PIC, 1924 : 3.	Madagascar
<i>Orphinus malabarensis</i> PIC, 1916c : 16.	India
<i>Orphinus mediojunctus</i> PIC, 1938d : 353.	Thailand
<i>Orphinus minimus</i> ARROW, 1915 : 441.	N. S. Australia
<i>Orphinus minor</i> ARROW, 1915 : 440.	India: Bombay
<i>Orphinus nealense</i> (BLACKBURN, 1903 : 170) ARMSTRONG, 1943 : 57.	New South Wales
Syn.: <i>Cryptorhopalum nealense</i> BLACKBURN, 1903:170.	
<i>Orphinus nesioticus</i> BEAL, 1961 : 113.	S. Mariana Is.
<i>Orphinus nilgirensis</i> ARROW, 1915 : 439.	S. India
<i>Orphinus notaticollis</i> PIC, 1916c : 16.	E. India
<i>Orphinus occidentalis</i> ARMSTRONG, 1943 : 62.	W. Australia
<i>Orphinus oscitans</i> (OLLIFF, 1883 : 185) comb. nov.	N. E. Borneo
Syn.: <i>Orphilus oscitans</i> OLLIFF, 1883:185.	
<i>Orphinus pedestris</i> MOTSCHULSKY, 1858 : 48.	Burma
<i>Orphinus quornensis</i> (BLACKBURN, 1895 : 93) ARROW, 1915 : 438.	Australia
Syn.: <i>Cryptorhopalum quornense</i> BLACKBURN, 1895:93.	
<i>Orphinus rufofasciatus</i> PIC, 1924 : 3.	Philippine Is.
<i>Orphinus subfasciatus</i> PIC, 1927e : 6.	Philippine Is.
<i>Orphinus sulcatopygus</i> PIC, 1927e : 6.	Borneo
<i>Orphinus tabitha</i> ARROW, 1915 : 440.	Ceylon

Orphinus terminalis (SHARP in BLACKBURN et SHARP, 1885 : 150) ARROW, 1915 : 438.

Syn.: *Cryptorhopalum terminale* SHARP in BLACKBURN et SHARP, 1885 : 150.

Orphinus testaceipes (PIC, 1915b : 6) PIC, 1916b : 3.

Syn.: *Thaumaglossa testaceipes* PIC, 1915b:6.

Orphinus tonkineus PIC, 1922 : 6.

Orphinus woodvillensis (BLACKBURN, 1891 : 130) ARROW, 1915 : 438.

Syn.: *Cryptorhopalum woodvillense* BLACKBURN, 1891:130.

Cryptorhopalum eucalypti BLACKBURN, 1903:171. — ARROW, 1915:438.

Hawaiian Is.,
Philippine Is.,
Samoa,
S. Mariana Is.,
Marshall Is.,
Gilbert Is.

New Guinea

Viet-Nam

S. Australia,
entire
Australia

Orphinus brevicornis (SHARP in BLACKBURN et SHARP, 1885:150) ARROW, 1915:438 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:135).

Orphinus formosanus PIC, 1918b:2 — vide *Orphinus aethiops* ARROW, 1915:438.

Orphinus luteovestitus PIC, 1937:410 — vide *Anthrenus luteovestitus* (PIC, 1937:410).

Orphinus rufopygus PIC, 1933:71 — vide *Thaumaglossa nigricans* (MACLEAY, 1871:171).

Orphinus uninotatus PIC, 1954:56 — vide *Thaumaglossa uninotata* (PIC, 1954:56).

Genus: *Curtophinus* PIC, 1954b : 11

Type species: *Curtophinus bicolor* PIC, 1954b : 11.

List of species:

Curtophinus bicolor PIC, 1954b : 11.

China

Genus: *Ctesias* STEPHENS, 1830 : 124

Type species: *Dermestes serra* FABRICIUS, 1792 : 234.

Syn.: *Megatoma* DEJEAN, 1821:46, nec HERBST, 1792:92. Type species: *Dermestes serra* FABRICIUS, 1792:234.

Attagenus LATREILLE, 1829:510 nec LATREILLE, 1802:121. Type species: *Dermestes serra* FABRICIUS, 1792:234.

Tiresias STEPHENS, 1832:410. — Type species: *Dermestes serra* FABRICIUS, 1792:234.

Ctenias REITTER, 1911:151. Type species: *Dermestes serra* FABRICIUS, 1792:234.

List of species:

Ctesias dusmae BEAL, 1960 : 95.

California

Ctesias intermedia MROCKOWSKI, 1961c : 195.

Turkmen SSR

Syn.: *Ctesias syriaca* MROCKOWSKI, 1960b:212 nec GANGLBAUER, 1904b:37.

<i>Ctesias kaliki</i> MROCKOWSKI, 1961c : 193.	Afghanistan
Syn.: <i>Ctesias syriaca</i> MROCKOWSKI, 1960:53 nec GANGLBauer, 1904b: 37.	
<i>Ctesias serra</i> (FABRICIUS, 1792 : 234) STEPHENS, 1830 : 124.	Sweden, all Europe
Syn.: <i>Dermestes serra</i> FABRICIUS, 1792:234.	
<i>Anthrenus viennensis</i> HERBST, 1797:336. — SCHÖNHERR, 1806:292.	
<i>Attagenus serra</i> (FABRICIUS, 1792:234) LATREILLE, 1804:244.	
<i>Megatoma serra</i> (FABRICIUS, 1792:234) LATREILLE, 1807:35.	
<i>Tiresias serra</i> (FABRICIUS, 1792:234) STEPHENS, 1833:34.	
<i>Ctenias serra</i> (FABRICIUS, 1792:234) REITTER, 1911:154.	
<i>Ctesias syriaca</i> GANGLBauer, 1904b : 37.	Syria
<i>Ctesias variegata</i> ARROW, 1915 : 433.	Rhodesia, Congo (Kinshasa)

Ctesias syriaca MROCKOWSKI, 1960:53 nec GANGLBauer, 1904b:37 — vide *Ctesias kaliki* MROCKOWSKI, 1961c:193.

Ctesias syriaca MROCKOWSKI, 1960b:212 nec GANGLBauer, 1904b:37 — vide *Ctesias intermedia* MROCKOWSKI, 1961c:195.

Tiresias plantaris (CURTIS, 1838:682) STEPHENS, 1839:143 — vide *Globicornis nigripes* (FABRICIUS, 1792: 233).

Genus: *Tiresiomorpha* PIC, 1954b : 11

Type species: *Tiresiomorpha klapperichi* PIC, 1954b : 11.

List of species:

Tiresiomorpha klapperichi PIC, 1954b : 11. China

Genus: *Thaumaglossa* REDTENBACHER, 1867 : 43

Type species: *Thaumaglossa rufocapillata* REDTENBACHER, 1867 : 44.

Syn.: *Axinocerus* JAYNE, 1882:347. — CASEY, 1900:145. Type species: *Axinocerus americanus* JAYNE, 1882:368.

Thaumatoglossa CASEY, 1900:145. Type species: *Axinocerus americanus* JAYNE, 1882:368.
Orphilooides MATSUMURA et YOKOYAMA, 1928:130, syn. nov. KALIK in litt. Type species:
Orphilooides ovivorus MATSUMURA et YOKOYAMA, 1928:131.

List of species:

Thaumaglossa africana KALIK, 1955b : 321. Zanzibar

Thaumaglossa americana (JAYNE, 1882 : 368) DALLA TORRE, 1911 : 65. Texas

Syn.: *Axinocerus americanus* JAYNE, 1882:368.

Thaumaglossa basilewskyi KALIK, 1955b : 321. Congo (Kinshasa)

<i>Thaumaglossa bimaculata</i> ARROW, 1916 : cxii.	Transvaal
Syn.: <i>Thaumaglossa bimaculata</i> POOL, 1915:710, nomen nudum.	
<i>Thaumaglossa concavifrons</i> REITTER, 1881 : 42.	Tasmania?
<i>Thaumaglossa conradti</i> PIC, 1927c : 247.	Cameroons, Congo (Kinshasa)
<i>Thaumaglossa hilleri</i> REITTER, 1881 : 42.	Japan, Philippine Is.
<i>Thaumaglossa laeta</i> ARROW, 1915 : 435.	Hongkong, Taiwan, Malaya, Philippine Is., China
Syn.: <i>Orphilus bimaculatus</i> MATSUMURA et YOKOYAMA, 1928b:53. — KALIK, 1952b:48. <i>Aethriostoma rufomaculata</i> PIC, 1938c:12. — KALIK, 1952b:47.	
<i>Thaumaglossa libochoras</i> BEAL, 1952 : 171.	Arizona
<i>Thaumaglossa nigricans</i> (MACLEAY, 1871 : 171) BLACKBURN, 1903 : 166.	Queensland, New South Wales
Syn.: <i>Anthrenus nigricans</i> MACLEAY, 1871:171. <i>Orphinus rufopygus</i> PIC, 1933:71. — ARMSTRONG, 1943:62.	
<i>Thaumaglossa nitidula</i> ARROW, 1915 : 436.	Malaya, Thailand
<i>Thaumaglossa oothecobia</i> ARROW, 1915 : 434.	S. Nigeria
<i>Thaumaglossa ovivora</i> (MATSUMURA et YOKOYAMA, 1928 : 131) comb. nov. KALIK in litt.	Japan, Taiwan
Syn.: <i>Orphiloides ovivorus</i> MATSUMURA et YOKOYAMA, 1928:131.	
<i>Thaumaglossa pauliani</i> PAULIAN, 1953 : 12.	Madagascar
<i>Thaumaglossa pygidialis</i> PIC, 1916c : 16.	Tonkin
<i>Thaumaglossa pygidialis</i> var. <i>sinensis</i> PIC, 1951 : 5.	China
<i>Thaumaglossa pygidialis</i> var. <i>sumatrana</i> PIC, 1951 : 5.	Sumatra
<i>Thaumaglossa rufescens</i> PIC, 1927e : 3.	Cameroons
<i>Thaumaglossa rufiventris</i> PIC, 1927c : 246.	Cameroons
<i>Thaumaglossa rufoapicalis</i> PIC, 1924 : 2.	Tonkin
<i>Thaumaglossa rufocapillata</i> REDTENBACHER, 1867 : 44.	Hongkong, China, Japan, Korea, Malayan pen., Sumatra, Borneo, Zanzibar, Germany (introduced)
Syn.: <i>Anthrenus ovalis</i> FLEUTIAUX, 1887:62. — ARROW, 1915:434. <i>Aethriostoma atricolor</i> PIC, 1915c:2. — ARROW, 1937:112. <i>Thaumaglossa atricolor</i> (PIC, 1915b:2) PIC, 1916c:16. <i>Thaumaglossa capillata</i> RÜSCHKAMP, 1929:154. — RÜSCHKAMP, 1930:11.	
<i>Thaumaglossa rufocincta</i> ARROW, 1915 : 434.	Rhodesia

- Thaumaglossa rufula* PIC, 1931b : 4. **Philippine Is.**
Thaumaglossa tonkinea PIC, 1916c : 16. **Tonkin**
Thaumaglossa uninotata (PIC, 1954 : 56) comb. nov. — KALIK **China: Fukien**
 in litt.
 Syn.: *Orphinus uninotatus* PIC, 1954:56.
-

- Thaumaglossa atricolor* (PIC, 1915b:2) PIC, 1916c:16 — vide *Thaumaglossa rufocapillata* REDTENBACHER, 1867:44.
Thaumaglossa bimaculata POOL, 1915:710, nomen nudum — vide *Thaumaglossa bimaculata* ARROW, 1916:cxii.
Thaumaglossa capillata RÜSKAMP, 1929:154 — vide *Thaumaglossa rufocapillata* REDTENBACHER, 1867: 44.
Thaumaglossa ovalis ARROW, 1915:436 — vide *Orphinus ovalis* (ARROW, 1915:436).
Thaumaglossa sexmaculata ARROW, 1915:435. — vide *Orphinus sexmaculatus* (ARROW, 1915:435).
Thaumaglossa testaceipes PIC, 1915b:6 — vide *Orphinus testaceipes* (PIC, 1915b:6).

Genus: *Pseudothaumaglossa* PIC, 1918b : 1

Type species: *Pseudothaumaglossa anthrenoides* PIC, 1918b : 1.

List of species:

- Pseudothaumaglossa anthrenoides* PIC, 1918b : 1. **Brazil**

Genus: *Hemirhopalum* SHARP, 1902 : 651

Type species: *Hemirhopalum hadrotomoide* SHARP, 1902 : 652.

List of species:

- Hemirhopalum apicale* PIC, 1954c : 177. **Peru**
Hemirhopalum bicolor SHARP, 1902 : 651. **Nicaragua**
Hemirhopalum buprestoide SHARP, 1902 : 652. **Guatemala**
 Syn.: *Hemirhopalum buprestoides* DALLA TORRE, 1911:77.
Hemirhopalum clythraeforme SHARP, 1902 : 651. **Guatemala**
Hemirhopalum curtum PIC, 1942 : 8. **Bolivia**
Hemirhopalum cyaneum PIC, 1927d : 186. **Peru**
Hemirhopalum elongatum PIC, 1942 : 8. **Brazil**
Hemirhopalum hadrotomoide SHARP, 1902 : 652. **Mexico**

Syn.: *Hemirhopalum hadrotomoides* DALLA TORRE, 1911:77.

- Hemirhopalum laticolle* PIC, 1927d : 186. **Peru**
Hemirhopalum longipenne PIC, 1916c : 17. **Peru, Brazil**
Hemirhopalum plaumanni MROCKOWSKI, 1958b : 56. **Brazil**

<i>Hemirhopalum rufipenne</i> PIC, 1916c : 17.	Mexico
<i>Hemirhopalum suturale</i> PIC, 1937c : 5.	Chile
<i>Hemirhopalum testaceipes</i> PIC, 1936 : 2.	Brazil

Hemirhopalum buprestoides DALLA TORRE, 1911:77 — vide *Hemirhopalum buprestoide* SHARP, 1902:652.
Hemirhopalum hadrotomoides DALLA TORRE, 1911:77 — vide *Hemirhopalum hadrotomoide* SHARP, 1902: 652.

Genus: *Falsocryptorhopalum* PIC, 1936 : 1

Type species: *Falsocryptorhopalum impressum* PIC, 1936 : 2.

List of species:

<i>Falsocryptorhopalum impressum</i> PIC, 1936 : 2.	Brazil
---	--------

Genus: *Cryptorhopalum* GUÉRIN-MÉNEVILLE, 1838b : 67

Type species: *Cryptorhopalum quadripunctatum* GUÉRIN-MÉNEVILLE, 1838b : 67.

List of species:

<i>Cryptorhopalum amulae</i> SHARP, 1902 : 666.	Mexico
<i>Cryptorhopalum anthrax</i> CASEY, 1900 : 157.	Arizona
<i>Cryptorhopalum anthrenoides</i> CASEY, 1916 : 194.	Arizona
<i>Cryptorhopalum apicale</i> (MANNERHEIM, 1843 : 258) LECONTE, 1854 : 111.	California, Oregon, Nevada
Syn.: <i>Anthrenus apicalis</i> MANNERHEIM, 1843:258.	

Cryptorhopalum appendiculatum SHARP, 1902 : 658.

Mexico

Cryptorhopalum argentinum PIC, 1916d : 3.

Argentina

Cryptorhopalum argutum SHARP, 1902 : 656.

Mexico,
Guatemala

Cryptorhopalum argutum var. *cordovanum* SHARP, 1902 : 656.

Mexico

Cryptorhopalum aridum CASEY, 1916 : 196.

Utah,
California

Cryptorhopalum atomarium REITTER, 1881 : 47.

Brazil

Cryptorhopalum atripes REITTER, 1881 : 46.

Colombia

Cryptorhopalum atropubescens REITTER, 1881 : 46.

Brazil,
Colombia

Cryptorhopalum atrum PIC, 1937d : 193.

Costa Rica

Cryptorhopalum baeri PIC, 1916d : 3.

Peru

Cryptorhopalum bakeri CASEY, 1916 : 197.

California

Cryptorhopalum balteatum LECONTE, 1854 : 111.

California,
Pacific States,
Arizona

<i>Cryptorhopalum bicolor</i> SHARP, 1902 : 656.	British Honduras
<i>Cryptorhopalum bilimeki</i> REITTER, 1881 : 44.	Mexico
<i>Cryptorhopalum bilimeki</i> var. <i>brumeipenne</i> REITTER, 1881 : 46.	Mexico
Syn.: <i>Cryptorhopalum bilimeki</i> var. <i>brunneipenne</i> DALLA TORRE, 1911:73.	
<i>Cryptorhopalum bimaculatum</i> REITTER, 1881 : 49.	Brazil
<i>Cryptorhopalum biolleyi</i> PIC, 1916d : 3.	Costa Rica
<i>Cryptorhopalum bisignatum</i> SHARP, 1902 : 653.	Panama
<i>Cryptorhopalum bolivianum</i> PIC, 1916d : 3.	Colombia
<i>Cryptorhopalum brevicolle</i> SHARP, 1902 : 663.	Panama
<i>Cryptorhopalum brunnescente</i> PIC, 1916c : 17.	Paraguay
<i>Cryptorhopalum callanganum</i> PIC, 1927e : 5.	Peru
<i>Cryptorhopalum celatum</i> SHARP, 1902 : 659.	Mexico
<i>Cryptorhopalum centromaculatum</i> REITTER, 1881 : 53.	Brazil
<i>Cryptorhopalum centromaculatum</i> var. <i>notatum</i> PIC, 1927e : 5.	Brazil
<i>Cryptorhopalum chontalense</i> SHARP, 1902 : 663.	Nicaragua
<i>Cryptorhopalum claveri</i> PIC, 1916d : 2.	Colombia
<i>Cryptorhopalum cleryi</i> GUÉRIN-MÉNEVILLE, 1838b : 68.	Peru
Syn.: <i>Cryptorhopalum clerii</i> ERICHSON, 1847:95.	
<i>Cryptorhopalum cleryi</i> var. <i>guerini</i> PIC, 1942 : 9.	Peru
<i>Cryptorhopalum coloradense</i> CASEY, 1916 : 192.	Colorado
<i>Cryptorhopalum convexum</i> PIC, 1927d : 186.	Colombia
<i>Cryptorhopalum corumbanum</i> PIC, 1923 : 3.	Brazil
<i>Cryptorhopalum cribiceps</i> SHARP, 1902 : 659.	Nicaragua
<i>Cryptorhopalum cribripenne</i> REITTER, 1881 : 44.	Mexico
<i>Cryptorhopalum cyphonoide</i> SHARP, 1902 : 664.	Mexico
Syn.: <i>Cryptorhopalum cyphonoides</i> DALLA TORRE, 1911:74.	
<i>Cryptorhopalum diehli</i> PIC, 1954c : 177.	Peru
<i>Cryptorhopalum difficile</i> REITTER, 1881 : 47.	Brazil
<i>Cryptorhopalum discedens</i> SHARP, 1902 : 666.	Mexico
<i>Cryptorhopalum divisum</i> SHARP, 1902 : 657.	Nicaragua, Panama
<i>Cryptorhopalum dorcatomoides</i> CASEY, 1900 : 157.	Arizona
<i>Cryptorhopalum dubium</i> SHARP, 1902 : 654.	Mexico, British Honduras
<i>Cryptorhopalum ducale</i> SHARP, 1902 : 665.	Guatemala
<i>Cryptorhopalum equisoleae</i> SHARP, 1902 : 665.	Mexico, British Honduras

<i>Cryptorhopalum eximium</i> ARROW, 1915 : 442.	Brazil
<i>Cryptorhopalum felis</i> ARROW, 1915 : 442.	Brazil
<i>Cryptorhopalum festivum</i> CASEY, 1900 : 156.	Texas
<i>Cryptorhopalum filitarse</i> CASEY, 1900 : 156.	California
<i>Cryptorhopalum flammulatum</i> SHARP, 1902 : 655.	Guatemala
<i>Cryptorhopalum floridanum</i> CASEY, 1916 : 198.	Florida
<i>Cryptorhopalum fontinale</i> CASEY, 1916 : 194.	New Mexico
<i>Cryptorhopalum fraternum</i> SHARP, 1902 : 667.	Mexico
<i>Cryptorhopalum funestum</i> SHARP, 1902 : 661.	Mexico
<i>Cryptorhopalum fusciclave</i> CASEY, 1900 : 158.	Texas, Utah
Syn.: <i>Cryptorhopalum fuscoclave</i> DALLA TORRE, 1911:74.	
<i>Cryptorhopalum fusculum</i> LECONTE, 1854 : 111.	California, Arizona, Mexico
<i>Cryptorhopalum geniculatum</i> PIC, 1916d : 2.	Brazil
<i>Cryptorhopalum germanum</i> SHARP, 1902 : 655.	Mexico, Guatemala
<i>Cryptorhopalum globuloide</i> SHARP, 1902 : 664.	Nicaragua
Syn.: <i>Cryptorhopalum globuloides</i> DALLA TORRE, 1911:74.	
<i>Cryptorhopalum globulum</i> REITTER, 1881 : 44.	Mexico
<i>Cryptorhopalum gounellei</i> PIC, 1916d : 3.	Brazil
<i>Cryptorhopalum grandjeani</i> PIC, 1942 : 9.	Peru
<i>Cryptorhopalum granum</i> CASEY, 1900 : 157.	Arizona
<i>Cryptorhopalum gravidum</i> SHARP, 1902 : 661.	Mexico
<i>Cryptorhopalum grisescens</i> CASEY, 1916 : 195.	Arizona
<i>Cryptorhopalum guatemalenum</i> SHARP, 1902 : 658.	Guatemala
<i>Cryptorhopalum haemorrhoidale</i> (LECONTE, 1824:170) LECONTE, 1854 : 111.	Georgia, Southern, Middle and Western States of USA
Syn.: <i>Anthrenus haemorrhoidalis</i> LECONTE, 1824:170.	
<i>Cryptorhopalum heydeni</i> KIRSCH, 1873 : 152.	Peru
<i>Cryptorhopalum hirsutum</i> PIC, 1927e : 4.	Bolivia
<i>Cryptorhopalum imperiale</i> REITTER, 1881 : 48.	Brazil
<i>Cryptorhopalum incanum</i> REITTER, 1881 : 45.	Mexico
<i>Cryptorhopalum infasciatum</i> PIC, 1942 : 10.	Peru
<i>Cryptorhopalum insigne</i> CASEY, 1916 : 191.	New Mexico
<i>Cryptorhopalum instabile</i> SHARP, 1902 : 654.	Guatemala, Nicaragua, Panama
<i>Cryptorhopalum laterale</i> SHARP, 1902 : 659.	Mexico
<i>Cryptorhopalum laticolle</i> PIC, 1927e : 4.	Brazil

<i>Cryptorhopalum maculatum</i> (FABRICIUS, 1798 : 76) MOTSCHULSKY, 1858b : 148.	America
Syn.: <i>Anthrenus maculatus</i> FABRICIUS, 1798:76.	
<i>Cryptorhopalum major</i> PIC, 1927e : 4.	Bolivia
<i>Cryptorhopalum mexicanum</i> PIC, 1916d : 2.	Mexico
<i>Cryptorhopalum minor</i> PIC, 1942 : 9.	Colombia
<i>Cryptorhopalum misellum</i> SHARP, 1902 : 662.	Guatemala
<i>Cryptorhopalum modestum</i> CASEY, 1900 : 158.	Texas
<i>Cryptorhopalum mordelloide</i> SHARP, 1902 : 664.	Guatemala
Syn.: <i>Cryptorhopalum mordelloides</i> DALLA TORRE, 1911:75.	
<i>Cryptorhopalum nephianum</i> CASEY, 1916 : 196.	Utah
<i>Cryptorhopalum nevermanni</i> PIC, 1937d : 194.	Costa Rica
<i>Cryptorhopalum nitidissimum</i> PIC, 1916d : 3.	Ecuador
<i>Cryptorhopalum oberthuri</i> REITTER, 1881 : 47.	Colombia, Brazil
<i>Cryptorhopalum obesulum</i> CASEY, 1900 : 157.	Texas
<i>Cryptorhopalum obscuriceps</i> PIC, 1916d : 3.	Brazil
<i>Cryptorhopalum obscuriceps</i> var. <i>bruchi</i> PIC, 1928 : 49.	Argentina
<i>Cryptorhopalum ochraceum</i> SHARP, 1902 : 663.	British Honduras, Guatemala
<i>Cryptorhopalum olivaceum</i> PIC, 1927e : 4.	Brazil
<i>Cryptorhopalum orbiculosum</i> REITTER, 1881 : 48.	Colombia
Syn.: <i>Cryptorhopalum orbiculum</i> DALLA TORRE, 1911:75.	
<i>Cryptorhopalum pallens</i> CASEY, 1916 : 193.	Texas
<i>Cryptorhopalum pedestre</i> SHARP, 1902 : 667.	Guatemala
<i>Cryptorhopalum perforatum</i> PIC, 1927d : 185.	Brazil
<i>Cryptorhopalum peruvianum</i> PIC, 1916d : 2.	Peru
<i>Cryptorhopalum piceum</i> CASEY, 1916 : 198.	California
<i>Cryptorhopalum picicorne</i> LECONTE, 1854 : 111.	Pennsylvania to Georgia, Massachusetts, New York
<i>Cryptorhopalum pilosum</i> KIRSCH, 1871 : 355.	Colombia
<i>Cryptorhopalum posticum</i> SHARP, 1902 : 653.	Panama
<i>Cryptorhopalum postnotatum</i> PIC, 1927e : 5.	Brazil
<i>Cryptorhopalum postnotatum</i> var. <i>mediofasciatum</i> PIC, 1927e : 5.	Brazil
<i>Cryptorhopalum presuturale</i> PIC, 1929b : 2.	Peru
<i>Cryptorhopalum pruddeni</i> CASEY, 1900 : 156.	Arizona
<i>Cryptorhopalum puberulum</i> REITTER, 1881 : 44.	Mexico

<i>Cryptorhopalum pumilum</i> CASEY, 1900 : 158.	Arizona
<i>Cryptorhopalum punctatissimum</i> REITTER, 1881 : 46.	Colombia
<i>Cryptorhopalum quadripunctatum</i> GUÉRIN-MÉNEVILLE, 1838b : 67.	Peru, Brazil, British Honduras, Guatemala
Syn.: <i>Cryptorhopalum flavopictum</i> LACORDAIRE, 1855:9. — syn. nov. <i>Cryptorhopalum septemsignatum</i> SHARP, 1902:653. — syn. nov.	
<i>Cryptorhopalum quadrisignatum</i> PIC, 1942 : 9.	Peru
<i>Cryptorhopalum quinquepunctatum</i> REITTER, 1881 : 47.	Bolivia
<i>Cryptorhopalum reversum</i> CASEY, 1900 : 156.	New Mexico
<i>Cryptorhopalum robustum</i> SHARP, 1902 : 660.	Mexico
<i>Cryptorhopalum ruficorne</i> LECONTE, 1854 : 111.	Southern States of USA, New York, Mexico
Syn.: <i>Cryptorhopalum rufipes</i> SHARP, 1902:660. — DALLA TORRE, 1911:76.	
<i>Cryptorhopalum rufipenne</i> PIC, 1916d : 4.	Colombia
<i>Cryptorhopalum rufipes</i> REITTER, 1881 : 44.	Mexico
<i>Cryptorhopalum rufobrunneum</i> PIC, 1942 : 8.	Bolivia
<i>Cryptorhopalum rufofasciatum</i> REITTER, 1881 : 49.	Colombia
<i>Cryptorhopalum rufonotatum</i> PIC, 1916c : 17.	Bolivia
<i>Cryptorhopalum rufum</i> PIC, 1942 : 8.	Peru
<i>Cryptorhopalum rugosipenne</i> PIC, 1916d : 2.	Paraguay
<i>Cryptorhopalum rugulosum</i> PIC, 1923 : 3.	Brazil
<i>Cryptorhopalum rusticolle</i> PIC, 1927e : 5.	Peru
<i>Cryptorhopalum sahlbergi</i> REITTER, 1881 : 46.	Brazil
<i>Cryptorhopalum scutellare</i> ARROW, 1915 : 443.	West Indies
<i>Cryptorhopalum semihirsutum</i> PIC, 1942 : 9.	Colombia
<i>Cryptorhopalum sexpunctatum</i> REITTER, 1881 : 48.	Brazil
<i>Cryptorhopalum sexsignatum</i> REITTER, 1881 : 47.	Brazil
<i>Cryptorhopalum sordidum</i> SHARP, 1902 : 662.	Mexico
<i>Cryptorhopalum splendidum</i> REITTER, 1881 : 47.	Colombia
<i>Cryptorhopalum stachi</i> MROCKOWSKI, 1958 : 559.	French Guiana
Syn.: <i>Cryptorhopalum atrum</i> PIC, 1942:8, nec PIC, 1937d:193.	
<i>Cryptorhopalum subcyanum</i> PIC, 1927e : 5.	Bolivia
<i>Cryptorhopalum subfasciatum</i> SHARP, 1902 : 657.	Mexico
<i>Cryptorhopalum submetallicum</i> PIC, 1937d : 194.	Costa Rica
<i>Cryptorhopalum subtile</i> SHARP, 1902 : 662.	Guatemala
<i>Cryptorhopalum subtrifasciatum</i> REITTER, 1881 : 48.	Brazil
<i>Cryptorhopalum teapense</i> SHARP, 1902 : 658.	Mexico
<i>Cryptorhopalum teffensis</i> REITTER, 1881 : 48.	Brazil
<i>Cryptorhopalum trifasciatum</i> PIC, 1927e : 5.	Bolivia
<i>Cryptorhopalum trisignatum</i> PIC, 1942 : 9.	Peru

Cryptorhopalum triste LECONTE, 1854 : 111.

Syn.: *Cryptorhopalum nigricorne* LECONTE, 1862b:344. — JAYNE, 1882: 367.

Cryptorhopalum affine CASEY, 1900:157 nec REITTER, 1881:54. — BEAL in litt.

Cryptorhopalum caseyi DALLA TORRE, 1911:73.

Cryptorhopalum trogodermoides REITTER, 1881 : 48.

Cryptorhopalum truncatum KIRSCH, 1871 : 355.

Cryptorhopalum tuckeri CASEY, 1916 : 192.

Cryptorhopalum unifasciatum PIC, 1954c : 178.

Cryptorhopalum uniforme PIC, 1927e : 4.

Cryptorhopalum uteanum CASEY, 1916 : 195.

Cryptorhopalum vagesignatum PIC, 1942 : 8.

Cryptorhopalum ventanense SHARP, 1902 : 661.

Cryptorhopalum vestitum SHARP, 1902 : 666.

Cryptorhopalum vianai PIC, 1940 : 12.

Cryptorhopalum vicinum SHARP, 1902 : 667.

Cryptorhopalum villosum REITTER, 1881 : 45.

Cryptorhopalum viridipubens PIC, 1923 : 3.

California,
Arizona,
Atlantic States,
Mexico:
North Sonora

Brazil
Colombia
Arizona
Peru
Brazil
Utah
Brazil
Mexico
Guatemala
Argentina
Mexico
Mexico
French Guiana

Cryptorhopalum affine CASEY, 1900:157 nec REITTER, 1881:54 — vide *Cryptorhopalum triste* LECONTE, 1854:111.

Cryptorhopalum affine REITTER, 1881:54 — vide *Orphinus affinis* (REITTER, 1881:54).

Cryptorhopalum atrum PIC, 1942:8 nec PIC, 1937d:193 — vide *Cryptorhopalum stachi* MROCKOWSKI, 1958:559.

Cryptorhopalum australe (HOPE, 1843:319) BLACKBURN, 1903:167 — vide *Anthrenocerus australis* (HOPE, 1843:319).

Cryptorhopalum australicum BLACKBURN, 1891:130 — vide *Orphinus australicus* (BLACKBURN, 1891:130).

Cryptorhopalum biflexum REITTER, 1881:55 — vide *Orphinus biflexus* (REITTER, 1881:55).

Cryptorhopalum bilimeki var. *brunneipenne* DALLA TORRE, 1911:73 — vide *Cryptorhopalum bilimeki* var. *brunneipenne* REITTER, 1881:46.

Cryptorhopalum brevicorne SHARP in BLACKBURN et SHARP, 1885:150 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:138).

Cryptorhopalum caseyi DALLA TORRE, 1911:73 — vide *Cryptorhopalum triste* LECONTE, 1854:111.

Cryptorhopalum casuarinae BLACKBURN, 1903:169 — vide *Orphinus casuarinae* (BLACKBURN, 1903:169).

Cryptorhopalum ceciliense BLACKBURN, 1903:170 — vide *Orphinus ceciliense* (BLACKBURN, 1903:170).

Cryptorhopalum clerii ERICHSON, 1847:95 — vide *Cryptorhopalum cleryi* GUÉRIN-MÉNEVILLE, 1838b:68.

Cryptorhopalum confertum REITTER, 1881:55 — vide *Anthrenocerus confertum* (REITTER, 1881:55).

Cryptorhopalum cyphonoides DALLA TORRE, 1911:74 — vide *Cryptorhopalum cyphonoide* SHARP, 1902: 664.

Cryptorhopalum diversum PIC, 1928b:2 — vide *Orphinus diversus* (PIC, 1928b:2).

Cryptorhopalum erichsoni REITTER, 1881:55 — vide *Anthrenocerus australis* (HOPE, 1843:319).

Cryptorhopalum eucalypti BLACKBURN, 1903:171 — vide *Orphinus woodvillensis* (BLACKBURN, 1891:130).

Cryptorhopalum flavopictum LACORDAIRE, 1855:9 — vide *Cryptorhopalum quadripunctatum* GUÉRIN-MÉNEVILLE, 1838b:67.

- Cryptorhopalum flindersi* (BLACKBURN, 1891:132) BLACKBURN, 1903:168 — vide *Anthrenocerus confertus* (REITTER, 1881:55).
- Cryptorhopalum fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:138) FLEUTIAUX et SALLÉ, 1890:392 — vide *Orphinus fulvipes* (GUÉRIN-MÉNEVILLE, 1838c:138).
- Cryptorhopalum fuscoclave* DALLA TORRE, 1911:74 — vide *Cryptorhopalum fuscoclave* CASEY, 1900:158.
- Cryptorhopalum globuloides* DALLA TORRE, 1911:74 — vide *Cryptorhopalum globuloide* SHARP, 1902: 664.
- Cryptorhopalum interioris* BLACKBURN, 1891:131 — vide *Orphinus interioris* (BLACKBURN, 1891:131).
- Cryptorhopalum mordelloides* DALLA TORRE, 1911:75 — vide *Cryptorhopalum mordelloide* SHARP, 1902: 664.
- Cryptorhopalum motschulskyi* REITTER, 1881:55 — vide *Orphinus haemorrhoidalis* MOTSCHULSKY, 1858:48.
- Cryptorhopalum nealense* BLACKBURN, 1903:170 — vide *Orphinus nealense* (BLACKBURN, 1903:170).
- Cryptorhopalum nigricorne* LECONTE, 1862b:344 — vide *Cryptorhopalum triste* LECONTE, 1854: 111.
- Cryptorhopalum obscurum* MACLEAY, 1871:171 — non Dermestidae, recte Anobiidae apud ARMSTRONG, 1943:57.
- Cryptorhopalum orbiculum* DALLA TORRE, 1911:75 — vide *Cryptorhopalum orbiculosum* REITTER, 1881: 48.
- Cryptorhopalum quadrifasciatum* BLACKBURN, 1903:169 — vide *Anthrenocerus quadrifasciatus* (BLACKBURN, 1903:169).
- Cryptorhopalum quornense* BLACKBURN, 1895:93 — vide *Orphinus quornensis* (BLACKBURN, 1895:93).
- Cryptorhopalum rufipes* SHARP, 1902:660 — vide *Cryptorhopalum ruficorne* LECONTE, 1854:111.
- Cryptorhopalum septemsignatum* SHARP, 1902:653 — vide *Cryptorhopalum quadripunctatum* GUÉRIN-MÉNEVILLE, 1838b:67.
- Cryptorhopalum terminale* SHARP in BLACKBURN et SHARP, 1885:150 — vide *Orphinus terminalis* (SHARP, 1885:150).
- Cryptorhopalum terzonatum* BLACKBURN, 1903:169 — vide *Anthrenocerus terzonatus* (BLACKBURN, 1903: 169).
- Cryptorhopalum variabile* REITTER, 1881:55 — vide *Anthrenocerus variabilis* (REITTER, 1881:55).
- Cryptorhopalum woodvillense* BLACKBURN, 1891:130 — vide *Orphinus woodvillensis* (BLACKBURN, 1891: 130).

Anthreninae LECONTE, 1861 : 107

Type genus: *Anthrenus* SCHAEFFER, 1766 : [26].

- Syn.: *Anthreni* LECONTE, 1861:107.
Anthrenini: CASEY, 1900:139.
Anthreninae: DALLA TORRE, 1911:77.

Genus: *Anthrenus* SCHAEFFER, 1776 : [26]

Type species: *Dermestes scrophulariae* LINNAEUS, 1758 : 356.

- Syn.: *Byrrhus* SULZER, 1776:[I], 25 nec LINNAEUS, 1767:568. Type species: *Dermestes scrophulaiae* LINNAEUS, 1758:356.
Hypoceuthes GERSTAECKER, 1871:46, syn. nov.¹ Type species: *Hypoceuthes aterrimus* GERSTAECKER, 1871:46.

¹ Two syntypes of the type species, *Hypoceuthes aterrimus* GERSTAECKER, 1871:46, preserved at the Zoological Museum of the Humboldt University in Berlin, were examined. The species belongs to the subgenus *Anthrenus* s. str.

Subgenus: *Anthrenus* s. str.

List of species:

Anthrenus acomanus CASEY, 1916 : 200.*Anthrenus armstrongi* MROCKOWSKI, 1952 : 27.

New Mexico

Yugoslavia,

Greece,

Bulgaria

Tanganyika

Anthrenus aterrimus (GERSTAECKER, 1871 : 46) comb. nov.Syn.: *Hypoceuthes aterrimus* GERSTAECKER, 1871:46.*Anthrenus crustaceus* REITTER, 1881 : 58.Syn.: *Anthrenus cretaceus* KNEUCKER, 1922:22.*Anthrenus festivus* ERICHSON, 1846 : 458.Syn.: *Anthrenus museorum* OLIVIER, 1790:No. 14, p. 8 nec (LINNAEUS, 1761:145). — ERICHSON, 1846 : 458.*Anthrenus museorum* var. *festivus* DEJEAN, 1837:140, nomen nudum.*Anthrenus delicatus* JACQUELIN DU VAL, 1859:t. 62, f. 308. — JACQUELIN DU VAL, 1859:259.*Anthrenus funestus* MULSANT et REY, 1868:140. — REITTER, 1881b:89.*Anthrenus varius* var. *festivus*: BAUDI A SELVE, 1870:63.*Anthrenus flavipes* LECONTE, 1854 : 112.Syn.: *Anthrenus fasciatus* REITTER, 1881b:87 et auctorum, non THUNBERG, 1795:105, nec HERBST, 1797:337. — BARBER, 1951:45.*Anthrenus scrophulariae* var. *flavipes*: JAYNE, 1882:370.*Anthrenus vorax* WATERHOUSE, 1883:61. — BARBER, 1951:45.*Anthrenus importatus* PIC, 1952c:1. — MROCKOWSKI, 1964:181.France,
Portugal,
Spain, Balearic
Is., Morocco,
Tunisia,
Cyprus?New York,
nearly
cosmopolitan*Anthrenus flavipes* var. *alboimpletus* SAHLBERG, 1903e : 34.Syn.: *Anthrenus fasciatus* var. *alboimpletus* SAHLBERG, 1903e:34.*Anthrenus flavipes albopunctatus* PIC, 1894e : 142. — MROCKOWSKI, 1964 : 181.

Algeria

Syria

Syn.: *Anthrenus pimpinellae* var. *albopunctatus* PIC, 1894e:142.
Anthrenus fasciatus var. *isabellae* REITTER, 1899:216. — MROCKOWSKI, 1964:181.*Anthrenus flavipes isabellae*: MROCKOWSKI, 1960b:216.*Anthrenus flavipes* var. *cinnamomeus* GREDLER, 1878 : 507.Sudan,
Balearic Is.Syn.: *Anthrenus pimpinellae* var. *cinnamomeus* GREDLER, 1878:507.
Anthrenus fasciatus var. *cinnamomeus*: REITTER, 1906:380.
Anthrenus vorax var. *cinnamomeus*: MROCKOWSKI, 1952:28.*Anthrenus flavipes* var. *seminiveus* CASEY, 1916 : 199.District of
ColumbiaSyn.: *Anthrenus seminiveus* CASEY, 1916:199.*Anthrenus vorax seminiveus*: BACK et COTTON, 1937:194.*Anthrenus vorax* var. *seminiveus*: HINTON, 1945:8.

Anthrenus goliath MULSANT et REY, 1868 : 140.

Syn.: *Anthrenus pimpinellae* var. *goliath*: REITTER, 1881b:87.

Anthrenus jordanicus PIC, 1934b : 29.

Anthrenus kaliki PIC, 1952c : 1.

Anthrenus lindbergi MROCKOWSKI, 1959 : 100.

Anthrenus miniatulus REITTER, 1899 : 216.

Anthrenus miniopictus BEDEL, 1884 : XXI.

Anthrenus munroi HINTON, 1943 : 14.

Anthrenus oceanicus FAUVEL, 1903 : 338.

Syn.: *Anthrenus fasciatus* HERBST, 1797:337, non THUNBERG, 1795: 105, nec REITTER, 1881b:87 et auctorum. — MROCKOWSKI, 1964:180.

Anthrenus fasciatus var. *latebasalis* PIC, 1927:66. — MROCKOWSKI, 1964:181.

Anthrenus pimpinellae var. *basifasciatus* PIC, 1935:13. — MROCKOWSKI, 1964:181.

Anthrenus vorax var. *latebasalis*: HINTON, 1945:8.

Anthrenus pacificus FAIRMAIRE, 1850 : 183.

Anthrenus parvus CASEY, 1900 : 161.

Anthrenus picturatus SOLSKIJ, 1876 : 280.

Egypt, Algeria,
France, whole
Mediterranea

Jordan

Algeria

Afghanistan

Syria

Algeria,
Morocco,
Tunisia

France,
Corsica,
Algeria, Israel,
Syria, Bulgaria,
Crimea

New Caledonia,
Malaya,
Ceylon

Tahiti

Texas

Uzbek SSR,
Tadzhik SSR,
Kirghiz SSR,
Kazakh SSR,
Turkmen SSR,
Afghanistan,
Iran,
Transcaucasia,
Caucasus

China

Poland

Uzbek SSR,
Tadzhik SSR,
Kirghiz SSR,
Kazakh SSR,
Afghanistan

Anthrenus picturatus hintoni MROCKOWSKI, 1952 : 29.

Anthrenus picturatus makolskii MROCKOWSKI, 1950 : 187.
— MROCKOWSKI, 1952 : 29.

Syn.: *Anthrenus makolskii* MROCKOWSKI, 1950:187.

Anthrenus picturatus melanoleucus SOLSKIJ, 1876 : 279. —
MROCKOWSKI, 1960 : 53.

Syn.: *Anthrenus melanoleucus* SOLSKIJ, 1876:279.

Anthrenus pimpinellae FABRICIUS, 1775 : 61.

Syn.: *Anthrenus scrophulariae* FOURCROY, 1785:27 nec (LINNAEUS, 1758: 356). — OLIVIER, 1790:No. 14, p. 7.

Byrrhus pimpinellae (FABRICIUS, 1775:61) GOEZE, 1777:171.

Anthrenus pimpinellae var. *angustefasciatus* GANGLBAUER, 1904b : 42.

Anthrenus pimpinellae var. *delicatus* KIESENWETTER, 1851 : 579. — REITTER, 1881b : 87.

Syn.: *Anthrenus delicatus* KIESENWETTER, 1851:579.

Anthrenus isabellinae MULSANT et REY, 1868:154. — REITTER, 1906:380.

Anthrenus var. *delicatulus* XAMBEU, 1892:45.

Anthrenus pimpinellae var. *delicatulus*: TENENBAUM, 1915:81.

Anthrenus pimpinellae var. *dimidiatus* HAUSER, 1894 : 23 — nomen nudum.

Anthrenus pimpinellae var. *dorsatus* MULSANT et REY, 1868 : 154.

Anthrenus pimpinellae var. *flavidulus* REITTER, 1889 : 23.

Anthrenus pimpinellae var. *incertus* MULSANT et REY, 1868 : 154.

Anthrenus pimpinellae var. *isabellinus* KÜSTER, 1848 : 38. — SCHAUM, 1862 : 47.

Syn.: *Anthrenus isabellinus* KÜSTER, 1848 : 38.

Attagenus isabellinus (KÜSTER, 1848:38) FAIRMAIRE et COQUEREL, 1860:170.

Anthrenus pimpinellae var. *niveus* REITTER, 1881b:88. — HINTON, 1945b:349.

Anthrenus fasciatus var. *isabellinus*: ESCALERA, 1914:138.

Anthrenus pimpinellae *latefasciatus* REITTER, 1892 : 134. — MROCKOWSKI, 1961c : 192.

Syn.: *Anthrenus pimpinellae* var. *latefasciatus* REITTER, 1892:134.

Anthrenus pimpinellae var. *lepidus* LECONTE, 1854 : 112. — ARROW 1915 : 445.

Syn.: *Anthrenus lepidus* LECONTE, 1854:112.

Anthrenus lepidus var. *conspersus* CASEY, 1900:160.

France,
very widely
distributed,
nearly
cosmopolitan

Dalmatia,
Spain

Pyrenees,
entire
Mediterranea,
Caucasus

Uzbek SSR

Algeria

Caucasus
Istanbul

Spain, France,
Morocco,
Tunisia

Uzbek SSR,
Tadzhik SSR,
Kirghiz SSR,
Kazakh SSR,
Turkmen SSR,
Afghanistan,
Iran,
Caucasus,
Syria, North
China,
Manchuria,
Korea

California

<i>Anthrenus lepidus</i> var. <i>obtectus</i> CASEY, 1900:160.	
<i>Anthrenus lepidus</i> var. <i>suffusus</i> CASEY, 1900:160. — CASEY, 1916:199.	
<i>Anthrenus pimpinellae mroczkowskii</i> KALIK, 1954 : 369. — MROCKOWSKI, 1958c : 4.	Greece, Bulgaria
Syn.: <i>Anthrenus pimpinellae</i> var. <i>mroczkowskii</i> KALIK, 1954:369.	
<i>Anthrenus pimpinellae</i> var. <i>niveipennis</i> SAHLBERG, 1903e : 34.	Algeria
<i>Anthrenus pimpinellae</i> var. <i>occidens</i> CASEY, 1900 : 160. — ARROW, 1915 : 445.	California, Nevada, British Columbia
Syn.: <i>Anthrenus occidens</i> CASEY, 1900:160.	
<i>Anthrenus occidens</i> var. <i>nevadicus</i> CASEY, 1900:160. — CASEY, 1916:199.	
<i>Anthrenus occidens</i> var. <i>pictus</i> CASEY, 1900:160.	
<i>Anthrenus pimpinellae</i> var. <i>pfefferi</i> KALIK, 1954 : 370.	Greece
<i>Anthrenus rauterbergi</i> REITTER, 1908 : 46.	Egypt
Syn.: <i>Anthrenus rautenbergi</i> WINKLER, 1926:680.	
<i>Anthrenus rotundulus</i> REITTER, 1889 : 23.	Transcaucasia, Turkmen SSR
<i>Anthrenus rotundulus</i> var. <i>tolerandus</i> PIC, 1908b : 41. — DALLA TORRE, 1911 : 82.	Egypt
Syn.: <i>Anthrenus rotundatus</i> var. <i>toleranda</i> PIC, 1908b:41.	
<i>Anthrenus scrophulariae</i> (LINNAEUS, 1758 : 356) FABRICIUS, 1775 : 61.	Europe, nearly cosmopolitan
Syn.: <i>Dermestes scrophulariae</i> LINNAEUS, 1758:356.	
<i>Dermestes variegatus</i> SCOPOLI, 1763:16. — SCHRANK, 1781:40.	
<i>Byrrhus scrophulariae</i> (LINNAEUS, 1758:356) LINNAEUS, 1767: 568.	
<i>Anthrenus histrio</i> FABRICIUS, 1792:264. — SCHÖNHERR, 1806:115.	
<i>Anthrenus verbasci</i> HERBST, 1797:328. — ILLIGER, 1801:91.	
<i>Anthrenus scrophulariae</i> ab. <i>histrio</i> : HEER, 1841:27.	
<i>Anthrenus scrophulariae</i> var. <i>verbasci</i> : GEMMINGER et HAROLD, 1868:921.	
<i>Anthrenus scrophulariae</i> var. <i>albida</i> DALLA TORRE, 1879:100. — SCHILSKY, 1889b:350.	
<i>Anthrenus scrophulariae</i> var. <i>flavida</i> DALLA TORRE, 1879:100. — SCHILSKY, 1889b:350.	
<i>Anthrenus scrophulariae</i> var. <i>albidus</i> BRULLÉ, 1832 : 164. — REITTER, 1881b : 90.	Greece, Dalmatia, Italy, Turkey, Caucasus
Syn.: <i>Anthrenus albidus</i> DEJEAN, 1821:47. — nomen nudum.	
<i>Anthrenus albidus</i> BRULLÉ, 1832:164.	
<i>Anthrenus proteus</i> var. <i>albidus</i> : SCHAUM, 1859:45.	
<i>Anthrenus scrophulariae</i> var. <i>signatus</i> REITTER, 1881b:90. — REITTER, 1906:380.	

Anthrenus scrophulariae var. *gravidus* KÜSTER, 1848 : 37. —
KRAATZ, 1864 : 133.

Syn.: *Anthrenus gravidus* KÜSTER, 1848:37.

Anthrenus proteus KRAATZ, 1858:147. — REITTER, 1906:380.
Anthrenus scrophulariae var. *proteus*: KRAATZ, 1864:133.

Dalmatia,
Sicily, Greece,
Crete, Turkey,
Caucasus,
Uzbek SSR,
Bulgaria,
Hungary

Anthrenus scrophulariae var. *hilarei* PIC, 1952c : 1.
Anthrenus scrophulariae ab. *insulicola* OBENBERGER, 1917 :
32. — MROCKOWSKI, 1964 : 186.

Syn.: *Anthrenus insulicola* OBENBERGER, 1917:32.

Pyrenees
Greece

Anthrenus scrophulariae var. *rubricollis* SAHLBERG, 1913c :
109.

Syria

Anthrenus scrophulariae var. *senex* KRAATZ, 1858 : 147. —
REITTER, 1881b : 90.

Greece,
Bulgaria
Turkey

Syn.: *Anthrenus senex* KRAATZ, 1858:147.

Anthrenus proteus var. *senex*: SCHAUM, 1859:45.

Anthrenus albidus var. *senex*: SCHAUM, 1862:47.

Anthrenus scrophulariae var. *suecicus* PALM, 1940 : 1.

Sweden,
Lithuanian
SSR, Germany,
Switzerland,
Italy, Hungary,
Russian FSSR,
Ukrainian SSR

Anthrenus scrophulariae var. *thoracicus* MELSHEIMER, 1844 :
117. — JAYNE, 1882 : 370.

Pennsylvania,
Middle and
Southern States
of USA

Syn.: *Anthrenus thoracicus* MELSHEIMER, 1806:8, nomen nudum —
MELSHEIMER, 1844:117.

Anthrenus thoracicus MELSHEIMER, 1844:117.

Anthrenus seminulum ARROW, 1915 : 446.

South Rhodesia
Senegal

Anthrenus senegalensis PIC, 1927e : 7.

Syria, Tunisia
Tahiti

Anthrenus simoni REITTER, 1881 : 59.

Algeria,
Tunisia, Egypt,
Corsica

Anthrenus sparsutus FAIRMAIRE, 1850 : 184.

Anthrenus x-signum REITTER, 1881b : 88.

Syn.: *Anthrenus x-nigrum* DESBROCHERS DES LOGES, 1898:73.

Subgenus: *Nathrenus* CASEY, 1900 : 161

Type species: *Byrrhus verbasci* LINNAEUS, 1767 : 568.

List of species:

Anthrenus afer PÉRINGUEY, 1888 : 90.

Anthrenus albomaculatus PIC, 1927e : 6.

South Africa
Borneo

Anthrenus albostictus REITTER, 1881 : 57.

Anthrenus basilewskyi KALIK, 1965 : 223.

Anthrenus biskrensis REITTER, 1887b : 69.

Syn.: *Anthrenus biscrensis* REITTER, 1894:116.

South Africa

Tanganyika

**Algeria, Malta,
Sicily, Tunisia,
Morocco**

Tunisia

Algeria, Egypt

Anthrenus biskrensis ab. *pallidus* NORMAND, 1936 : 301.

Anthrenus exilis MULSANT et REY, 1868 : 158.

Syn.: *Attagenus exilis* (MULSANT et REY, 1868:158) DALLA TORRE,
1911:53.

Anthrenus funebris REITTER, 1889b : 256.

Anthrenus liliputanus MULSANT et REY, 1868 : 159.

Syn.: *Attagenus liliputanus* (MULSANT et REY, 1868:159) DALLA TORRE,
1911:53.

Anthrenus luteovestitus (PIC, 1937 : 410) comb. nov. — KALIK
in litt.

Syn.: *Orphinus luteovestitus* PIC, 1937:410.

Anthrenus molitor AUBÉ, 1850 : 335.

Syn.: *Anthrenus ochraceus* MULSANT et REY, 1868:161. — REITTER,
1891:170.

**Dodecanese Is.
Egypt**

Kenya

**Crete, Greece,
Cyprus,
Turkey,
Algeria,
Tunisia,
Morocco,
Spain,
Portugal, Syria**

France, Cyprus

Anthrenus molitor var. *albidus* CASTELNAU, 1840 : 37. —
SCHAUM, 1862 : 47.

Syn.: *Anthrenus albidus* CASTELNAU, 1840:37.

Anthrenus nocivus MULSANT, 1870 : 240.

Anthrenus obscurus THUNBERG, 1815 : 151.

Anthrenus pilosus PIC, 1923 : 3.

Anthrenus pubifer REITTER, 1899 : 217.

Algeria
South Africa

Tibet

Syria

Syria
Austria,
Hungary,
Yugoslavia,
Bulgaria

Bulgaria

Croatia

Anthrenus signatus ab. *bielawskii* MROCKOWSKI, 1958c : 6.
Anthrenus signatus var. *incanus* FRIVALDSZKY, 1882 : 28. —
GANGLBAUER, 1904b : 43.

Syn.: *Anthrenus incanus* FRIVALDSZKY, 1882:28.

Anthrenus subsetosus ARROW, 1915 : 446.

Anthrenus transcaspicus MROCKOWSKI, 1960b : 217.

Anthrenus undatus REITTER, 1881 : 57.

Burma

**Iran,
Turkmen SSR**

South Africa

<i>Anthrenus verbasci</i> (LINNAEUS, 1767 : 568) GMELIN, 1790 : 1614.	France, cosmopolitan
Syn.: <i>Byrrhus verbasci</i> LINNAEUS, 1767:568.	
<i>Bostrichus varius</i> FABRICIUS, 1775:60. — SCHÖNHERR, 1806:115.	
<i>Anthrenus florilegus</i> FOURCROY, 1785:27. — DALLA TORRE, 1911: 84.	
<i>Dermestes varius</i> (FABRICIUS, 1775:60) SCHNEIDER, 1785:5.	
<i>Anthrenus varius</i> (FABRICIUS, 1775:60) FABRICIUS, 1792:264.	
<i>Anthrenus adspersus</i> HERBST, 1797:332. — MROCKOWSKI, 1964: 182.	
<i>Anthrenus tricolor</i> HERBST, 1797:333. — GYLLENHAL, 1808:163.	
<i>Anthrenus pictus</i> GERMAR, 1813:121. — ERICHSON, 1846:455.	
<i>Anthrenus tomentosus</i> THUNBERG, 1815:151. — REITTER, 1891:170.	
<i>Anthrenus verbasci</i> ab. <i>bifasciata</i> HÄNEL, 1935 : 76.	Germany
<i>Anthrenus verbasci</i> var. <i>caseyi</i> HINTON, 1945b : 328.	Indiana
Syn: <i>Anthrenus verbasci</i> var. <i>vorax</i> CASEY, 1900:161. — HINTON, 1945b:328.	
<i>Anthrenus verbasci</i> var. <i>confusus</i> REITTER, 1887b : 69.	Without terra typica! Europe
<i>Anthrenus verbasci</i> var. <i>destructor</i> MELSHEIMER, 1844 : 116. — LACORDAIRE, 1856 : 575.	Pennsylvania
Syn.: <i>Anthrenus destructor</i> MELSHEIMER, 1844:116.	
<i>Anthrenus varius</i> var. <i>destructor</i> : LACORDAIRE, 1856:575.	
<i>Anthrenus verbasci</i> var. <i>nebulosus</i> REITTER, 1887b : 69.	Without terra typica! Mediterranea, Caucasus
Syn.: <i>Anthrenus verbasci</i> var. <i>maculosus</i> GERHARDT, 1910:556, nomen nudum.	
<i>Anthrenus verbasci</i> var. <i>nitidulus</i> KÜSTER, 1847 : 47. — REITTER, 1881b : 91.	Yugoslavia, Bulgaria, Syria, Spain, Tunisia, Balearic Is.
Syn.: <i>Anthrenus nitidulus</i> KÜSTER, 1847:47.	
<i>Anthrenus verbasci</i> var. <i>pistor</i> CASEY, 1900 : 161.	Virginia
<i>Anthrenus verbasci</i> var. <i>substriatus</i> CASEY, 1900 : 161.	Central America
<i>Anthrenus versicolor</i> REITTER, 1887b : 70.	Greece, Sicily
Subgenus: <i>Anthrenodes</i> CHOBAUT, 1898 : 76	
Type species: <i>Anthrenus umbellatarum</i> CHOBAUT, 1898 : 76.	
List of species:	
<i>Anthrenus maculifer</i> REITTER, 1881 : 58.	East India, India, Taiwan, Viet-Nam, Burma, China
Syn.: <i>Anthrenus globiger</i> ARROW, 1915:447. — MROCKOWSKI, 1964: 184.	
<i>Anthrenus multimaculatus</i> PIC, 1918b:2. — MROCKOWSKI, 1964: 183.	
<i>Anthrenus albonotatus</i> PIC, 1922:6. — MROCKOWSKI, 1964:183.	

<i>Anthrenus sarnicus</i> MROCKOWSKI, 1963b : 697.	Guernsey
<i>Anthrenus umbellatarum</i> CHOBAUT, 1898 : 76.	Algeria
<i>Anthrenus umbellatarum</i> var. <i>albidipennis</i> CHOBAUT, 1898:76.	Algeria
Subgenus: <i>Anthrenops</i> REITTER, 1881b : 86	
Type species: <i>Anthrenus coloratus</i> REITTER, 1881b : 91.	
List of species:	
<i>Anthrenus amoenulus</i> REITTER, 1896 : 286.	Caspian Lowlands, Iran, Turkmen SSR
<i>Anthrenus bellulus</i> CHOBAUT, 1897 : 277.	Algeria
Syn.: <i>Anthrenus bullulus</i> DALLA TORRE, 1911:78.	
<i>Anthrenus bellulus ornatus</i> KOCHER, 1955 : 40. — MROCKOWSKI, 1964 : 185.	Morocco
Syn.: <i>Anthrenus ornatus</i> KOCHER, 1955:40.	
<i>Anthrenus coloratus</i> REITTER, 1881b : 91.	Greece, southeast Europe, North Africa, England (introduced), Austria?, Turkmen SSR, Afghanistan, India, Sudan
<i>Anthrenus hissaricus</i> MROCKOWSKI, 1961 : 591.	Tadzhik SSR
<i>Anthrenus longus</i> ARROW, 1915 : 448.	Socotra, Tanganyika
Syn.: <i>Anthrenus longipennis</i> PIC, 1937c:6. — MROCKOWSKI, 1964:186.	
<i>Anthrenus megalops</i> ARROW, 1915 : 447.	Ethiopia
<i>Anthrenus minor</i> WOLLASTON, 1865 : 163.	Canary Is., Algeria, Morocco, Tunisia, Greece
Syn.: <i>Anthrenus claviger</i> WOLLASTON, 1864:161 nec ERICHSON, 1846: 458. — WOLLASTON, 1865:163.	
<i>Anthrenus albidoflavus</i> REITTER, 1881b:91. — REITTER, 1891:170.	
<i>Anthrenops minutus</i> SAHLBERG, 1903c:36, nomen nudum. — SAHLBERG, 1903e:35.	
<i>Anthrenus minor</i> ab. <i>unicolor</i> UYTENBOOGAART, 1937 : 7.	Canary Is.
<i>Anthrenus oculatus</i> ARROW, 1937 : 1111.	Kenya

¹ The examination of six cotypes (labelled as paratypes) preserved in the Institute of Zoology of the Polish Academy of Sciences in Warsaw, proved that the antennae of the species in question are 9-joint ed. It should be, thus, placed not in the subgenus *Florilinus* MULS. et REY, but in the subgenus *Anthrenops* REITT. This has already been stated "in litteris" by Mr. V. KALIK of Pardubice who has had other cotypes at his disposal.

<i>Anthrenus rufescens</i> PIC, 1923 : 3.	India
<i>Anthrenus subclaviger</i> REITTER, 1881 : 59.	Aden, Yemen, India
<i>Anthrenus zebra</i> REITTER, 1889 c : 280.	Azerbaijan SSR
<i>Anthrenus zebra</i> var. <i>tigrinus</i> REITTER, 1889 c : 281.	Azerbaijan SSR
Subgenus: <i>Florilinus</i> MULSANT et REY, 1868 : 141	
Type species: <i>Dermestes museorum</i> LINNAEUS, 1761 : 145.	
List of species:	
<i>Anthrenus caseyi</i> DALLA TORRE, 1911 : 78.	New York
Syn.: <i>Anthrenus rotundulus</i> CASEY, 1900:162 nec REITTER, 1889:23. — DALLA TORRE, 1911:78.	
<i>Anthrenus castaneae</i> MELSHEIMER, 1844 : 116.	USA
Syn.: <i>Anthrenus castaneae pumilae</i> MELSHEIMER, 1806:8, nomen nudum. — MELSHEIMER, 1844:116.	
<i>Anthrenus castaneae</i> var. <i>carolinae</i> CASEY, 1900:162. — CASEY, 1916:199.	
<i>Anthrenus castaneae</i> var. <i>angustulus</i> CASEY, 1900 : 162.	Texas
<i>Anthrenus caucasicus</i> REITTER, 1881 b : 92.	Caucasus, Turkmen SSR, Vienna (introduced)
<i>Anthrenus coreanus</i> MROCKOWSKI, 1966 d : 440.	Korea
<i>Anthrenus flavidus</i> SOLSKII, 1876 : 281.	Uzbek SSR, Tadzhik SSR, Kazakh SSR, Kirghiz SSR, North China, Afghanistan, Iran, Turkmen SSR, Caucasus, Warsaw (introduced)
Syn.: <i>Anthrenus rufulus</i> SOLSKII, 1876:281. — MROCKOWSKI, 1954:1.	
<i>Anthrenus museorum</i> (LINNAEUS, 1761 : 145) FABRICIUS, 1775 : 61.	Sweden, entire Holarctic Region
Syn.: <i>Dermestes museorum</i> LINNAEUS, 1761:145.	
<i>Byrrhus museorum</i> (LINNAEUS, 1761:145) LINNAEUS, 1767:568.	
<i>Anthrenus verbasci</i> FABRICIUS, 1775:61. — ERICHSON, 1846:457.	
<i>Byrrhus muscorum</i> FUSSLIN, 1775:5.	
<i>Byrrhus verbasci</i> (FABRICIUS, 1775:61) GOEZE, 1777:170.	
<i>Anthrenus muscorum</i> FABRICIUS, 1787:39.	
<i>Anthrenus vagus</i> GMELIN, 1790:1615. — MOTSCHULSKY, 1858 b: 148.	

<i>Anthrenus pellio</i> THUNBERG, 1815:151. — syn. nov. ¹	
<i>Anthrenus varius</i> STEPHENS, 1830:130. — ERICHSON, 1846:457.	
<i>Florilinus muscorum</i> (FABRICIUS, 1787:39) COUCKE, 1895:466.	
<i>Anthrenus oberthueri</i> REITTER, 1881b : 91.	Pyrenees, Corsica, Algeria
<i>Anthrenus olgae</i> KALIK, 1946b : 134.	Czechoslo- vakia, Poland, Hungary, Austria, Germany
<i>Anthrenus sordidulus</i> REITTER, 1889b : 256.	Dodecanese Is., Syria, Spain

Subgenus: *Solskinus* MROCKOWSKI, 1962 : 252

Type species: *Anthrenus leucogrammus* SOLSKIJ, 1876 : 282.

Syn.: *Solskinus* MROCKOWSKI, 1951:258 (not valid description — type species not designated).

List of species:

<i>Anthrenus armeniacus</i> ZAJCEV, 1919 : 177.	Armenian SSR
<i>Anthrenus dsungaricus</i> MROCKOWSKI, 1962 : 251.	Kazakh SSR
<i>Anthrenus heptamerus</i> PEYERIMHOFF, 1924 : 62.	Morocco
<i>Anthrenus leucogrammus</i> SOLSKIJ, 1876 : 282.	Tadzhik SSR, Uzbek SSR
<i>Anthrenus sinensis</i> ARROW, 1915 : 445.	North China
<i>Anthrenus tadzhicus</i> MROCKOWSKI, 1961 : 592.	Tadzhik SSR

Subgenus: *Helocerus* MULSANT et REY, 1868 : 141

Type species: *Anthrenus fuscus* OLIVIER, 1789 : 152.

List of species:

<i>Anthrenus fuscus</i> OLIVIER, 1789 : 152.	France, middle Europe
Syn.: <i>Anthrenus obscurus</i> SCHÖNHERR, 1806:117. — SCHÖNHERR, 1806: 117.	
<i>Anthrenus claviger</i> ERICHSON, 1846:458. — ERICHSON, 1846:458.	
<i>Helocerus fuscus</i> (OLIVIER, 1789:152) COUCKE, 1892:75.	
<i>Anthrenus minutus</i> ERICHSON, 1846 : 459.	Sardinia, Portugal, Corsica, Spain
Syn.: <i>Anthrenus verbasci</i> var. <i>minutus</i> : REITTER, 1883:90.	
<i>Anthrenus minutus</i> var. <i>flavosquamatus</i> PIC, 1903b : 121.	Spain

¹ The type-specimen, preserved in coll. THUNBERG at Uppsala University, was examined.

Anthrenus polonicus MROCKOWSKI, 1951 : 253.

Poland,
Romania,
Ukrainian
SSR, Hungary,
Bulgaria,
Yugoslavia,
Russian FSSR

Subgenus: *Ranthenus* MROCKOWSKI, 1962 : 255

Type species: *Anthrenus alatauensis* MROCKOWSKI, 1962 : 254.

List of species:

Anthrenus alatauensis MROCKOWSKI, 1962 : 254.

Kazakh SSR

Species incertae sedis

Anthrenus aegyptiacus PIC, 1899c : 255.

Egypt

Anthrenus bleusei PIC, 1899b : 85.

Algeria

Anthrenus capensis GUÉRIN-MÉNEVILLE, 1838b : 66.

South Africa

Anthrenus palaeoaegyptiacus GRÜSS, 1930 : 181.

Egypt

Anthrenus pulchellus GESTRO, 1889 : 37.

Aden

Anthrenus adspersus HERBST, 1797:332 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).

Anthrenus albidoflavus REITTER, 1881b:91 — vide *Anthrenus minor* WOLLASTON, 1865:163.

Anthrenus albidus BRULLÉ, 1832:164 — vide *Anthrenus scrophulariae* var. *albidus* BRULLÉ, 1832:164.

Anthrenus albidus CASTELNAU, 1840:37 — vide *Anthrenus molitor* var. *albidus* CASTELNAU, 1840:37.

Anthrenus albidus DEJEAN, 1821:47, nomen nudum — vide *Anthrenus scrophulariae* var. *albidus* BRULLÉ, 1832:164.

Anthrenus albidus var. *senex*: SCHAUM, 1862:47 — vide *Anthrenus scrophulariae* var. *senex* KRAATZ, 1858:147.

Anthrenus albonotatus PIC, 1922:6 — vide *Anthrenus maculifer* REITTER, 1881:58.

Anthrenus apicalis KÜSTER, 1847:48 — vide *Anthrenus signatus* ERICHSON, 1846:456.

Anthrenus apicalis MANNERHEIM, 1843:258 — vide *Cryptorhopalum apicale* (MANNERHEIM, 1843:258).

Anthrenus australis HOPE, 1843:319 — vide *Anthrenocerus australis* (HOPE, 1843:319).

Anthrenus basalis DEJEAN, 1837:140. — nomen nudum.

Anthrenus bifasciatus THUNBERG, 1815:151 — vide *Attagenus thunbergi* KALIK in litt., nom. nov.

Anthrenus bisciensis REITTER, 1894:116 — vide *Anthrenus bisciensis* REITTER, 1887b:69.

Anthrenus bullulus DALLA TORRE, 1911:78 — vide *Anthrenus bellulus* CHOBAUT, 1897:277.

Anthrenus castaneae var. *carolinae* CASEY, 1900:162 — vide *Anthrenus castaneae* MELSHEIMER, 1844:116.

Anthrenus castaneae pumilae MELSHEIMER, 1806:8, nomen nudum — vide *Anthrenus castaneae* MELSHEIMER, 1844:116.

Anthrenus cinereus THUNBERG, 1815:151 — vide *Attagenus cinereus* (THUNBERG, 1815:151) comb. nov.

Anthrenus claviger ERICHSON, 1846:458 — vide *Anthrenus fuscus* OLIVIER, 1789:152.

Anthrenus claviger WOLLASTON, 1864:161 nec ERICHSON, 1846:458 — vide *Anthrenus minor* WOLLASTON, 1865:163.

Anthrenus cretaceus KNEUCKER, 1922:22 — vide *Anthrenus crustaceus* REITTER, 1881:58.

Anthrenus var. *delicatulus* XAMBEU, 1892:45 — vide *Anthrenus pimpinellae* var. *delicatus* KIESENWETTER, 1851:579.

- Anthrenus delicatus* JACQUELIN DU VAL, 1859:t. 62, f. 308 — vide *Anthrenus festivus* ERICHSON, 1846:458.
Anthrenus delicatus KIESENWETTER, 1851:579 — vide *Anthrenus pimpinellae* var. *delicatus* KIESENWETTER, 1851:579.
Anthrenus denticornis FABRICIUS, 1792:265 — vide *Trogoderma serraticorne* (FABRICIUS, 1792:265).
Anthrenus destructor MELSHEIMER, 1844:116 — vide *Anthrenus verbasci* var. *destructor* MELSHEIMER, 1844:116.
Anthrenus elongatulus DUFTSCHMID, 1825:26 — vide *Trogoderma versicolor* (CREUTZER, 1799:117).
Anthrenus elongatulus FABRICIUS, 1801:106 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Anthrenus elongatus SCHÖNHERR, 1806:114 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Anthrenus erythrocerus DEJEAN, 1837:140 — nomen nudum.
Anthrenus fasciatus REITTER, 1881b:87 et auctorum, non THUNBERG, 1795:105, nec HERBST, 1797:337 — vide *Anthrenus flavipes* LECONTE, 1854:112.
Anthrenus fasciatus HERBST, 1797:337, non THUNBERG, 1795:105, nec REITTER, 1881b:87 et auctorum — vide *Anthrenus oceanicus* FAUVEL, 1903:338.
Anthrenus fasciatus THUNBERG, 1795:105 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
Anthrenus fasciatus var. *alboimpletus* SAHLBERG, 1903e:34 — vide *Anthrenus flavipes* var. *alboimpletus* SAHLBERG, 1903e:34.
Anthrenus fasciatus var. *cinnamomeus*: REITTER, 1906:380 — vide *Anthrenus flavipes* var. *cinnamomeus* GREDLER, 1878:507.
Anthrenus fasciatus var. *isabellae* REITTER, 1899:216 — vide *Anthrenus flavipes albopunctatus* PIC, 1894e: 142.
Anthrenus fasciatus var. *isabellinus*: ESCALERA, 1914:138 — vide *Anthrenus pimpinellae* var. *isabellinus* KÜSTER, 1848:38.
Anthrenus fasciatus var. *latebasalis* PIC, 1927:66 — vide *Anthrenus oceanicus* FAUVEL, 1903:338.
Anthrenus flavipes isabellae: MROCKOWSKI, 1960b:216 — vide *Anthrenus flavipes albopunctatus* PIC, 1894e:142.
Anthrenus flindersi BLACKBURN, 1891:132 — vide *Anthrenocerus confertus* (REITTER, 1881:55).
Anthrenus florilegus FOURCROY, 1785:27 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).
Anthrenus frater ARROW, 1915:448 — vide *Neoanthrenus frater* (ARROW, 1915:448).
Anthrenus fulvicornis DEJEAN, 1821:46 — nomen nudum.
Anthrenus fulvipes DEJEAN, 1821:47 — nomen nudum.
Anthrenus funestus MULSANT et REY, 1868:140 — vide *Anthrenus festivus* ERICHSON, 1846:458.
Anthrenus fuscus GMELIN, 1790:1615 nec OLIVIER, 1789:152 — non Dermestidae, familia?
Anthrenus glaber CREUTZER in PANZER, 1796:11 — vide *Orphilus niger* (ROSSI, 1790:39).
Anthrenus glaber HERBST, 1783:26 — vide *Trogoderma glabrum* (HERBST, 1783:26).
Anthrenus glabratus FABRICIUS, 1801:109 — vide *Orphilus niger* (ROSSI, 1790:39).
Anthrenus globiger ARROW, 1915:447 — vide *Anthrenus maculifer* REITTER, 1881:58.
Anthrenus gloriosae FABRICIUS, 1798:76 — vide *Attagenus fasciatus* (THUNBERG, 1795:105).
Anthrenus grandis THUNBERG, 1815:151 — vide *Trogoderma versicolor* (CREUTZER, 1799:117).
Anthrenus gravidus KÜSTER, 1848:37 — vide *Anthrenus scrophulariae* var. *gravidus* KÜSTER, 1848:37.
Anthrenus haemorrhoidalis LECONTE, 1824:170 — vide *Cryptorhopalum haemorrhoidale* (LECONTE, 1824: 170).
Anthrenus hirtellus DEJEAN, 1837:140 — nomen nudum.
Anthrenus hirtus (FABRICIUS, 1781:93) FABRICIUS, 1792:265 — vide *Trinodes hirtus* (FABRICIUS, 1781:93).
Anthrenus histrio FABRICIUS, 1792:264 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
Anthrenus importatus PIC, 1952c:1 — vide *Anthrenus flavipes* LECONTE, 1854:112.
Anthrenus incanus FRIVALDSZKY, 1882:28 — vide *Anthrenus signatus* var. *incanus* FRIVALDSZKY, 1882:28.
Anthrenus insulicola OBENBERGER, 1917:32 — vide *Anthrenus scrophulariae* ab. *insulicola* OBENBERGER, 1917:32.
Anthrenus irroratus THUNBERG, 1815:151 — vide *Attagenus irroratus* (THUNBERG, 1815:151) comb. nov.
Anthrenus isabellinae MULSANT et REY, 1868:154 — vide *Anthrenus pimpinellae* var. *delicatus* KIESENWETTER, 1851:579.

- Anthrenus isabellinus* KÜSTER, 1848:38 — vide *Anthrenus pimpinellae* var. *isabellinus* KÜSTER, 1848:38.
- Anthrenus lepidus* LECONTE, 1854:112 — vide *Anthrenus pimpinellae* var. *lepidus* LECONTE, 1854:112.
- Anthrenus lepidus* var. *conspersus* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *lepidus* LECONTE, 1854:112.
- Anthrenus lepidus* var. *obtectus* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *lepidus* LECONTE, 1854:112.
- Anthrenus lepidus* var. *suffusus* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *lepidus* LECONTE, 1854:112.
- Anthrenus longipennis* PIC, 1937c:6 — vide *Anthrenus longus* ARROW, 1915:448.
- Anthrenus maculatus* FABRICIUS, 1798:76 — vide *Cryptorhopalum maculatum* (FABRICIUS, 1798:76).
- Anthrenus makolskii* MROCKOWSKI, 1950:187 — vide *Anthrenus picturatus makolskii* MROCKOWSKI, 1950:187.
- Anthrenus melanoleucus* SOLSKIJ, 1876:279 — vide *Anthrenus picturatus melanoleucus* SOLSKIJ, 1876:279.
- Anthrenus minimus* SCHRANK, 1798:471 — non Dermestidae, familia?
- Anthrenops minutus* SAHLBERG, 1903c:36, nomen nudum — vide *Anthrenus minor* WOLLASTON, 1865:163.
- Anthrenus multimaculatus* PIC, 1918b:2 — vide *Anthrenus maculifer* REITTER, 1881:58.
- Anthrenus murinus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus muscorum* FABRICIUS, 1787:39 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).
- Anthrenus museorum* OLIVIER, 1790:No. 14, p. 8 nec (LINNAEUS, 1761:145) — vide *Anthrenus festivus* ERICHSON, 1846:458.
- Anthrenus museorum* var. *festivus* DEJEAN, 1837:140, nomen nudum — vide *Anthrenus festivus* ERICHSON, 1846:458.
- Anthrenus niger* HERBST, 1797:338 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Anthrenus niger* MELSHEIMER, 1806:8 — nomen nudum.
- Anthrenus nigricans* MACLEAY, 1871:171 — vide *Thaumaglossa nigricans* (MACLEAY, 1871:171).
- Anthrenus nigritus* STURM, 1843:94 — nomen nudum.
- Anthrenus nitidulus* KÜSTER, 1847:47 — vide *Anthrenus verbasci* var. *nitidulus* KÜSTER, 1847:47.
- Anthrenus obscurus* MELSHEIMER, 1806:8 — nomen nudum.
- Anthrenus obscurus* SCHÖNHERR, 1806:117 — vide *Anthrenus fuscus* OLIVIER, 1789:152.
- Anthrenus occidens* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *occidens* CASEY, 1900:160.
- Anthrenus occidens* var. *nevadicus* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *occidens* CASEY, 1900:160.
- Anthrenus occidens* var. *pictus* CASEY, 1900:160 — vide *Anthrenus pimpinellae* var. *occidens* CASEY, 1900:160.
- Anthrenus ocellifer* BLACKBURN, 1891:132 — vide *Neoanthrenus ocellifer* (BLACKBURN, 1891:132).
- Anthrenus ochraceus* MULSANT et REY, 1868:161 — vide *Anthrenus molitor* AUBÉ, 1850:335.
- Anthrenus ornatus* KOCHER, 1955:40 — vide *Anthrenus bellulus* *ornatus* KOCHER, 1955:40.
- Anthrenus ovalis* FLEUTIAUX, 1887:62 — vide *Thaumaglossa rufocapillata* REDTENBACHER, 1867:44.
- Anthrenus pardalinus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus pellio* THUNBERG, 1815:151 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).
- Anthrenus pictus* GERMAR, 1813:121 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).
- Anthrenus pimpinellae* var. *albopunctatus* PIC, 1894e:142 — vide *Anthrenus flavipes* *albopunctatus* PIC, 1894e:142.
- Anthrenus pimpinellae* var. *basifasciatus* PIC, 1935:13 — vide *Anthrenus oceanicus* FAUVEL, 1903:338.
- Anthrenus pimpinellae* var. *cinnamomeus* GREDLER, 1878:507 — vide *Anthrenus flavipes* var. *cinnamomeus* GREDLER, 1878:507.
- Anthrenus pimpinellae* var. *delicatus*: TENENBAUM, 1915:81 — vide *Anthrenus pimpinellae* var. *delicatus* KIESENWETTER, 1851:579.
- Anthrenus pimpinellae* var. *goliath*: REITTER, 1881b:87 — vide *Anthrenus goliath* MULSANT et REY, 1868:140.
- Anthrenus pimpinellae* var. *niveus* REITTER, 1881b:88 — vide *Anthrenus pimpinellae* var. *isabellinus* KÜSTER, 1848:38.
- Anthrenus proteus* KRAATZ, 1858:147 — vide *Anthrenus scrophulariae* var. *gravidus* KÜSTER, 1848:37.

- Anthrenus proteus* var. *albidus*: SCHAUM, 1859:45 — vide *Anthrenus scrophulariae* var. *albidus* BRULLÉ, 1832:164.
- Anthrenus proteus* var. *senex*: SCHAUM, 1859:45 — vide *Anthrenus scrophulariae* var. *senex* KRAATZ, 1858:147.
- Anthrenus pubescens* (FABRICIUS, 1787) FABRICIUS, 1792:265 — non Dermestidae, recte Helodidae, genus *Cyphon* PAYK. apud SCHAUM, 1847:54.
- Anthrenus pulicarius* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus punctatus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus pustulatus* THUNBERG, 1815:151 — vide *Attagenus pustulatus* (THUNBERG, 1815:151).
- Anthrenus rautenbergi* WINKLER, 1926:680 — vide *Anthrenus rauterbergi* REITTER, 1908:46.
- Anthrenus pygmaeus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus rossii* (PONZA, 1805) GEMMINGER et HAROLD, 1868:921 — non Dermestidae, recte Byrrhidae: *Pedilophorus nitidus* (SCHALLER, 1783) apud CAPRA, 1951:44.
- Anthrenus rotundatus* var. *toleranda* PIC, 1908b:41 — vide *Anthrenus rotundulus* var. *tolerandus* PIC, 1908b:41.
- Anthrenus rotundulus* CASEY, 1900:162 nec REITTER, 1889:23 — vide *Anthrenus caseyi* DALLA TORRE, 1911:78.
- Anthrenus ruficornis* LATREILLE, 1807:39 — vide *Trogoderma glabrum* (HERBST, 1783:26).
- Anthrenus rufipes* STURM, 1826:94 — nomen nudum.
- Anthrenus rufulus* SOLSKIJ, 1876:281 — vide *Anthrenus flavidus* SOLSKIJ, 1876:281.
- Anthrenus scrophulariae* FOURCROY, 1785:27 nec (LINNAEUS, 1758:356) — vide *Anthrenus pimpinellae* FABRICIUS, 1775:61.
- Anthrenus scrophulariae* var. *albida* DALLA TORRE, 1879:100 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Anthrenus scrophulariae* var. *flavida* DALLA TORRE, 1879:100 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Anthrenus scrophulariae* var. *flavipes*: JAYNE, 1882:370 — vide *Anthrenus flavipes* LECONTE, 1854:112.
- Anthrenus scrophulariae* ab. *histrio*: HEER, 1841:27 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Anthrenus scrophulariae* var. *proteus*: KRAATZ, 1864:133 — vide *Anthrenus scrophulariae* var. *gravidus* KÜSTER, 1848:37.
- Anthrenus scrophulariae* var. *signatus* REITTER, 1881b:90 — vide *Anthrenus scrophulariae* var. *albidus* BRULLÉ, 1832:164.
- Anthrenus scrophulariae* var. *verbasci*: GEMMINGER et HAROLD, 1868:921 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Anthrenus seminiveus* CASEY, 1916:199 — vide *Anthrenus flavipes* var. *seminiveus* CASEY, 1916:199.
- Anthrenus senex* KRAATZ, 1858:147 — vide *Anthrenus scrophulariae* var. *senex* KRAATZ, 1858:147.
- Anthrenus serraticornis* FABRICIUS, 1792:265 — vide *Trogoderma serraticorne* (FABRICIUS, 1792:265).
- Anthrenus sertarius* STURM, 1843:94 — nomen nudum.
- Anthrenus signatus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus socius* LEA, 1895:228 — vide *Trogoderma socium* (LEA, 1895:228).
- Anthrenus squamosus* DEJEAN, 1837:140 — nomen nudum.
- Anthrenus thoracicus* MELSHEIMER, 1806:8, nomen nudum — vide *Anthrenus scrophulariae* var. *thoracicus* MELSHEIMER, 1844:117.
- Anthrenus thoracicus* MELSHEIMER, 1844:117 — vide *Anthrenus scrophulariae* var. *thoracicus* MELSHEIMER, 1844:117.
- Anthrenus tomentosus* THUNBERG, 1815:151 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).
- Anthrenus tricolor* HERBST, 1797:333 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).
- Anthrenus unifasciatus* LATREILLE, 1825:362 — nomen nudum.
- Anthrenus vagus* GMELIN, 1790:1615 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).
- Anthrenus varius* (FABRICIUS, 1775:60) FABRICIUS, 1792:264 — vide *Anthrenus verbasci* (LINNAEUS, 1767:568).
- Anthrenus varius* STEPHENS, 1830:130 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).

- Anthrenus varius* var. *destructor*: LACORDAIRE, 1856:575 — vide *Anthrenus verbasci* var. *destructor* MEL-SHEIMER, 1844:116.
- Anthrenus varius* var. *festivus*: BAUDI a SELVE, 1870:63 — vide *Anthrenus festivus* ERICHSON, 1846:458.
- Anthrenus verbasci* FABRICIUS, 1775:61 — vide *Anthrenus museorum* (LINNAEUS, 1761:145).
- Anthrenus verbasci* HERBST, 1797:328 — vide *Anthrenus scrophulariae* (LINNAEUS, 1758:356).
- Anthrenus verbasci* var. *maculosus* GERHARDT, 1910:556, nomen nudum — vide *Anthrenus verbasci* var. *nebulosus* REITTER, 1887b:69.
- Anthrenus verbasci* var. *minutus*: REITTER, 1883:90 — vide *Anthrenus minutus* ERICHSON, 1846:459.
- Anthrenus verbasci* var. *vorax* CASEY, 1900:161 — vide *Anthrenus verbasci* var. *caseyi* HINTON, 1945b:328.
- Anthrenus versicolor* CREUTZER, 1799:117 — vide *Trogoderma versicolor* (CREUTZER, 1799:117).
- Anthrenus versicolor* STURM, 1843:94 — nomen nudum.
- Anthrenus viennensis* HERBST, 1797:336 — vide *Ctesias serra* (FABRICIUS, 1791:234).
- Anthrenus villosulus* DUFTSCHMID, 1825:28 — vide *Phradonoma villosulum* (DUFTSCHMID, 1825:28).
- Anthrenus vorax* WATERHOUSE, 1883:61 — vide *Anthrenus flavipes* LECONTE, 1854:112.
- Anthrenus vorax* var. *cinnamomeus*: MROCKOWSKI, 1952:28 — vide *Anthrenus flavipes* var. *cinnamomeus* GREDLER, 1878:507.
- Anthrenus vorax* var. *latebasalis*: HINTON, 1945:8 — vide *Anthrenus oceanicus* FAUVEL, 1903:338.
- Anthrenus vorax* *seminiveus*: BACK et COTTON, 1937:194 — vide *Anthrenus flavipes* var. *seminiveus* CASEY, 1916:199.
- Anthrenus vorax* var. *seminiveus*: HINTON, 1945:8 — vide *Anthrenus flavipes* var. *seminiveus* CASEY, 1916:199.
- Anthrenus x-nigrum* DESBROCHERS des LOGES, 1898:73 — vide *Anthrenus x-signum* REITTER, 1881b:88

Genus: *Neoanthrenus* ARMSTRONG, 1941 : 388

Type species: *Anthrenus ocellifer* BLACKBURN, 1891 : 132.

List of species:

<i>Neoanthrenus armstrongi</i> KALIK, 1957 : 22.	Sydney
<i>Neoanthrenus frater</i> (ARROW, 1915 : 448) ARMSTRONG, 1941 : 389.	Tasmania
Syn.: <i>Anthrenus frater</i> ARROW, 1915:448.	
<i>Neoanthrenus macqueeni</i> ARMSTRONG, 1949 : 111.	Queensland
<i>Neoanthrenus niveosparsa</i> ARMSTRONG, 1941 : 389.	New South Wales
<i>Neoanthrenus ocellifer</i> (BLACKBURN, 1891 : 132) ARMSTRONG, 1941 : 389.	South Australia
Syn.: <i>Anthrenus ocellifer</i> BLACKBURN, 1891:132.	
<i>Neoanthrenus parallelus</i> ARMSTRONG, 1941 : 389.	New South Wales, Queensland

Orphilinae LECONTE, 1861 : 107

Type genus: *Orphilus* ERICHSON, 1846 : 424.

Syn.: *Orphili* LECONTE, 1861:107.

Orphilini: CASEY, 1900:140.

Orphilinae: DALLA TORRE, 1911:87.

Genus: *Orphilus* ERICHSON, 1846 : 424

Type species: *Byrrhus niger* ROSSI, 1790 : 39.

List of species:

Orphilus aequalis CASEY, 1900 : 164.

Orphilus ater ERICHSON, 1846 : 463.

Orphilus chalybeus CASEY, 1900 : 164.

†*Orphilus dubius* WICKHAM, 1912 : 15.

Orphilus niger (ROSSI, 1790 : 39) REITTER, 1891 : 170.

Syn.: *Byrrhus niger* ROSSI, 1790:39.

Anthrenus glaber CREUTZER in PANZER, 1796:11. — SCHÖNHERR,
1806:117.

Anthrenus glabratus FABRICIUS, 1801:109. — SCHÖNHERR, 1806:
117.

Orphilus glabratus (FABRICIUS, 1801:109) ERICHSON, 1846:462.

Orphilus subnitidus LECONTE, 1861 : 109.

Arizona,
Oregon,
British
Columbia

North America

Idaho

Colorado
(Miocene)

Mediterranea,
Caucasus,
Asia Minor,
Middle Asia

California,
Oregon,
Nevada,
Colorado, New
Mexico, British
Columbia

Orphilus bimaculatus MATSUMURA et YOKOYAMA, 1928b:53 — vide *Thaumaglossa laeta* ARROW, 1915:435.

Orphilus bipunctatus (sic!) MATSUMURA et YOKOYAMA, 1928b:53 — vide *Orphinus bipunctatus* (MATSUMURA et YOKOYAMA, 1928b:53).

Orphilus fasciatus MATSUMURA et YOKOYAMA, 1928b:54 — vide *Orphinus fasciatus* (MATSUMURA et YOKOYAMA, 1928b:54).

Orphilus glabratus (FABRICIUS, 1801:109) ERICHSON, 1846:462 — vide *Orphilus niger* (ROSSI, 1790:39).

Orphilus oscitans OLLIFF, 1883:185 — vide *Orphinus oscitans* (OLLIFF, 1883:185).

***Trinodinae* CASEY, 1900 : 140**

Type genus: *Trinodes* DEJEAN, 1821 : 47.

Syn.: *Trinodini* CASEY, 1900:140.

Trinodinae: DALLA TORRE, 1911:85.

***Genus: Apsectus* LECONTE, 1854 : 107**

Type species: *Syncalypta hispidus* MELSHEIMER, 1844 : 117.

List of species:

Apsectus araneorum BEAL, 1959 b : 135.

Apsectus centralis SHARP, 1902 : 669.

Arizona, Utah
Guatemala

<i>Apsectus dichromus</i> BEAL, 1959b : 136.	Panama
<i>Apsectus hispidus</i> (MELSHEIMER, 1844 : 117) LECONTE, 1854 : 113.	United States, but only in the Atlantic and Gulf Coast States
Syn.: <i>Syncalypta hispidus</i> MELSHEIMER, 1844:117.	
<i>Trinodes hispidus</i> (MELSHEIMER, 1844:117) MELSHEIMER, 1853:48.	
<i>Apsectus hystrix</i> SHARP, 1902 : 668.	Mexico
<i>Apsectus mexicanus</i> (REITTER, 1881 : 60) ARROW, 1915 : 449.	Mexico
Syn.: <i>Trinodes mexicanus</i> REITTER, 1881:60.	
<i>Apsectus minutus</i> SHARP, 1902 : 669.	Mexico
<i>Apsectus obscurus</i> SHARP, 1902 : 669.	Mexico

Apsectus hirtellus (WALKER, 1858:207) ARROW, 1915:449 — vide *Evorinea hirtella* (WALKER, 1858:207).
Apsectus indicus ARROW, 1915:450 — vide *Evorinea indica* (ARROW, 1915:450).
Apsectus iota ARROW, 1915:449 — vide *Evorinea iota* (ARROW, 1915:449).
Aspectus (sic!) *minor* (PIC, 1922:6) KALIK, 1954:369 — vide *Evorinea minor* (PIC, 1922:6).

Genus: *Trinodes* DEJEAN, 1821 : 47

Type species: *Nitidula hirta* FABRICIUS, 1781 : 93.

List of species:

<i>Trinodes albohirsutus</i> KALIK, 1965 : 222.	Tanganyika
<i>Trinodes bicoloratus</i> PIC, 1939 : 144.	Egypt
<i>Trinodes carinatus</i> PIC, 1916c : 17.	India
<i>Trinodes cinereohirtus</i> MOTSCHULSKY, 1863 : 433.	Ceylon
<i>Trinodes curtus</i> PIC, 1894d : 119.	Algeria
<i>Trinodes emarginatus</i> ARROW, May 1915 : 451.	Ceylon
Syn.: <i>Trinodes guernei</i> PIC, July 1915b:7. — PIC, 1916:3.	
<i>Trinodes flavus</i> MOTSCHULSKY, 1863 : 433.	Egypt
<i>Trinodes hirtus</i> (FABRICIUS, 1781 : 93) DEJEAN, 1821 : 47.	Europe, Algeria, Caucasus, Turkmen SSR
Syn.: <i>Nitidula hirta</i> FABRICIUS, 1781:93.	
<i>Nitidula pilosa</i> GMELIN, 1790:1631 (<i>pilosa</i> HERBST, 1783:39 not available as published without generic name). — SCHÖNHERR, 1806:116.	
<i>Anthrenus hirtus</i> (FABRICIUS, 1781:93) FABRICIUS, 1792:265.	
<i>Trinodes hirtus</i> v. <i>ater</i> DALLA TORRE, 1879:100. — SCHILSKY, 1889b:350.	
<i>Trinodes hirtus</i> v. <i>castanea</i> DALLA TORRE, 1879:100. — SCHILSKY, 1889b:350.	
<i>Trinodes hirtus</i> v. <i>testacea</i> DALLA TORRE, 1879:100. — SCHILSKY, 1889b:350.	
<i>Trinodes minutus</i> PIC, March 1915 : 2.	Borneo
Syn.: <i>Trinodes rotundus</i> ARROW, May 1915:451. — PIC, 1916:3.	

<i>Trinodes niger</i> MATSUMURA et YOKOYAMA, 1928b : 53.	Taiwan
<i>Trinodes quadrimaculatus</i> MATSUMURA et YOKOYAMA, 1928b : 52.	Japan
<i>Trinodes rufescens</i> REITTER, 1877 : 376.	Japan
<i>Trinodes rufithorax</i> PIC, 1926 : 4.	Luzon
<i>Trinodes rufotestaceus</i> PIC, 1952 : 9.	Congo (Kinshasa)
<i>Trinodes senegalensis</i> PIC, 1915 : 2.	Senegal
<i>Trinodes sinensis</i> FAIRMAIRE, 1886 : 319.	China
<i>Trinodes tonkineus</i> PIC, 1922 : 6.	Tonkin
<i>Trinodes villosus</i> BOHEMAN, 1851 : 584.	Natal

<i>Trinodes australis</i> MONTROUZIER, 1860:264 — vide <i>Attagenus australis</i> (MONTROUZIER, 1860:264).
<i>Trinodes globosus</i> MACLEAY, 1871:171 — non Dermestidae, recte Byrrhidae: <i>Byrrhinus globosus</i> (MACLEAY, 1871) apud ARROW, 1915:450.
<i>Trinodes guernei</i> PIC, July 1915b:7 — vide <i>Trinodes emarginatus</i> ARROW, May 1915:451.
<i>Trinodes hirtellus</i> WALKER, 1858:207 — vide <i>Evorinea hirtella</i> (WALKER, 1858:207).
<i>Trinodes hirtus</i> v. <i>ater</i> DALLA TORRE, 1879:100 — vide <i>Trinodes hirtus</i> (FABRICIUS, 1781:93).
<i>Trinodes hirtus</i> v. <i>castanea</i> DALLA TORRE, 1879:100 — vide <i>Trinodes hirtus</i> (FABRICIUS, 1781:93).
<i>Trinodes hirtus</i> v. <i>testacea</i> DALLA TORRE, 1879:100 — vide <i>Trinodes hirtus</i> (FABRICIUS, 1781:93).
<i>Trinodes hispidus</i> (MELSHEIMER, 1844:117) MELSHEIMER, 1853:48 — vide <i>Apsectus hispidus</i> (MELSHEIMER, 1844:117).
<i>Trinodes mexicanus</i> REITTER, 1881:60 — vide <i>Apsectus mexicanus</i> (REITTER, 1881:60).
<i>Trinodes minor</i> PIC, 1922:6 — vide <i>Evorinea minor</i> (PIC, 1922:6).
<i>Trinodes punctipennis</i> MACLEAY, 1871:171 — non Dermestidae, recte Byrrhidae: <i>Byrrhinus punctipennis</i> (MACLEAY, 1871) apud ARROW, 1915:450.
<i>Trinodes rotundus</i> ARROW, May 1915:451 — vide <i>Trinodes minutus</i> PIC, March 1915:2.

Genus: *Evorinea* BEAL, 1961 : 118

Type species: *Apsectus iota* ARROW, 1915 : 449.

Syn.: *Evorinea* BEAL, 1959b:132, 133 (not valid description — type species not designated).

List of species:

<i>Evorinea hirtella</i> (WALKER, 1858 : 207) BEAL, 1961 : 118.	Ceylon
Syn.: <i>Trinodes hirtellus</i> WALKER, 1858:207.	
<i>Apsectus hirtellus</i> (WALKER, 1858:207) ARROW, 1915:449.	
<i>Evorinea indica</i> (ARROW, 1915 : 450) BEAL, 1961 : 118.	Bombay
Syn.: <i>Apsectus indicus</i> ARROW, 1915:450.	
<i>Evorinea iota</i> (ARROW, 1915 : 449) BEAL, 1961 : 112.	Tanimbar Is., Philippine Is., Caroline Is., S. Mariana Is.
Syn.: <i>Apsectus iota</i> ARROW, 1915:449.	
<i>Evorinea minor</i> (PIC, 1922 : 6) comb. nov.	Tonkin, Sumatra, Mariana Is. (?)
Syn.: <i>Trinodes minor</i> PIC, 1922:6.	
<i>Aspectus</i> (sic!) <i>minor</i> (PIC, 1922:6) KALIK, 1954:369.	

Genus: *Hexanodes* BLAIR, 1941 : 15

Type species: *Hexanodes hudsoni* BLAIR, 1941 : 15.

List of species:

Hexanodes hudsoni BLAIR, 1941 : 15.

New Zealand

***Thylodriadiinae* MROCKOWSKI, 1954c : 10**

Type genus: *Thylodrias* MOTSCHULSKY, 1839 : 75.

Syn.: *Thelydriini* SEMENOV-TIAN-SHANSKII, 1912:498. Type genus: *Thelydrias* AGASSIZ, 1846:368.

Thylodriinae WINKLER, 1926:675. Type genus: *Thylodrias* MOTSCHULSKY, 1839:75.

Thylodriini: BEAL, 1959:99.

Genus: *Thylodrias* MOTSCHULSKY, 1839 : 75

Type species: *Thylodrias contractus* MOTSCHULSKY, 1839 : 76.

Syn.: *Thelodrias* AGASSIZ et ERICHSON, 1846:161. Type species: *Thylodrias contractus* MOTSCHULSKY, 1839:76.

Thelydrias AGASSIZ, 1846:368. Type species: *Thylodrias contractus* MOTSCHULSKY, 1839:76.

Ignodus SLOSSON, 1903:187. — ZAJCEV, 1909:37. Type species: *Ignodus aenigmaticus* SLOSSON, 1903:187.

Hospitopterus PIC, 1921:54. — BARBER, 1947:344. Type species: *Hospitopterus efflatouni* PIC, 1921:54.

List of species:

Thylodrias contractus MOTSCHULSKY, 1839 : 76.

Caucasus,
many places in
the Holarctic
Region

Syn.: *Thelydrias contractus* (MOTSCHULSKY, 1839:76) GEMMINGER et HAROLD, 1869:1704.

Ignodus aenigmaticus SLOSSON, 1903:187. — ZAJCEV, 1909:37.

Hospitopterus efflatouni PIC, 1921:54. — BARBER, 1947:344.

Ignodus aenigmaticus SLOSSON, 1903:187 — vide *Thylodrias contractus* MOTSCHULSKY, 1839:76.

Hospitopterus efflatouni PIC, 1921:54 — vide *Thylodrias contractus* MOTSCHULSKY, 1839:76.

***Thaumaphrastinae* ANDERSON, 1949 : 127**

Type genus: *Thaumaphrastus* BLAISDELL, 1927b : 123.

Syn.: *Thorictini* BEAL, 1959:100. Type genus: *Thorictodes* REITTER, 1875c:45 (nec *Thorictidae* WOLLASTON, 1854). Type genus: *Thorictus* GERMAR, 1834).

Genus: *Thorictodes* REITTER, 1875c : 45

Type species: *Thorictodes heydeni* REITTER, 1875c : 46.

Syn.: *Thaumaphrastus* BLAISDELL, 1927b:123. — BEAL, 1959:99. Type species: *Thaumaphrastus karanisensis* BLAISDELL, 1927b:123.

List of species:

<i>Thorictodes bennetti</i> JOHN, 1961 : 43.	former British Guiana
<i>Thorictodes dartevellei</i> JOHN, 1961 : 41.	Congo (Kinshasa)
<i>Thorictodes erraticus</i> CHAMPION, 1922 : 73.	India
<i>Thorictodes heydeni</i> REITTER, 1875c : 46.	Egypt, Algeria, France, Spain, Sudan, India, Java, Guadeloupe, S. Mariana Is., USA, Mexico
Syn.: <i>Thaumaphrastus karanisensis</i> BLAISDELL, 1927b:123. — BEAL, 1959:99.	

Thaumaphrastus karanisensis BLAISDELL, 1927b:123 — vide *Thorictodes heydeni* REITTER, 1875c:46.

Genera incertae sedis

Genus: *Reeveana* DUNSTAN, 1923 : 48

Type species: *Reeveana minor* DUNSTAN, 1923 : 50.

List of species:

† <i>Reeveana intermedia</i> DUNSTAN, 1923 : 49.	Queensland (Mesozoic)
† <i>Reeveana major</i> DUNSTAN, 1923 : 48.	Queensland (Mesozoic)
† <i>Reeveana minor</i> DUNSTAN, 1923 : 50.	Queensland (Mesozoic)

Genus: *Tryoniopsis* DUNSTAN, 1923 : 51

Type species: *Tryoniopsis punctata* DUNSTAN, 1923 : 51.

List of species:

† <i>Tryoniopsis granulata</i> DUNSTAN, 1923 : 52.	Queensland (Mesozoic)
† <i>Tryoniopsis punctata</i> DUNSTAN, 1923 : 51.	Queensland (Mesozoic)

IV. Bibliography

- ABEILLE DE PERRIN [E.] 1870. [*Megatoma rufovittata* sp. nov.]. Ann. Soc. Ent. France, Paris, ser 4, **10**: XLVI.
- ABEILLE DE PERRIN E. 1872. Études sur les Coléoptères cavernicoles, suivies de la description de Coléoptères nouveaux propres au midi de la France. Ann. Soc. Ent. France, Paris, ser. 5, **2**: XLI-XLIV.
- AGASSIZ L. 1846. Nomenclatoris zoologici index universalis, continens nomina systematica classium, ordinum, familiarum et generum animalium omnium, tam viventium quam fossilium, secundum ordinem alphabeticum unicum disposita, adjectis homonymiis plantarum, nec non variis adnotacionibus et emendationibus. Soloduri, **VIII** + 393 pp.
- AGASSIZ L. et ERICHSON G. F. [1846]. Nomina systelamtica generum Coleopterorum, tam viventium quam fossilium, secundum ordinem alphabeticum disposita, adjectis auctoribus, libris in quibus reperiuntur, anno editionis, etymologia et familiis ad quas pertinent. In: AGASSIZ L. Nomenclator zoologicus, etc. Soloduri, **XII** + 170 pp.
- AHRENS A. 1814. Fauna Insectorum Europae. Fasciculus 2. Halae, 25 pp., 25 pls.
- ANDERSON W. H. 1949. Larval Description and transfer of *Thaumaphrastus karanensis* from *Colydiidae* to a new subfamily of *Dermestidae* (*Coleoptera*). Bull. Brooklyn Ent. Soc., Lancaster, n. s. **44** : 121-127, pls. X-XI.
- ARMSTRONG J. W. T. 1941. On Australian *Dermestidae*. Part I. Descriptions of a new genus and two new species; also a note on the genus *Anthrenus*. Proc. Linn. Soc. N. S. Wales, Sydney, **66** : 388-390, 3 ff.
- ARMSTRONG J. W. T. 1942. On Australian *Dermestidae*. Part II. The genus *Trogoderma* BERTHOLD. Proc. Linn. Soc. N. S. Wales, Sydney, **67** : 321-330.
- ARMSTRONG J. W. T. 1943. On Australian *Dermestidae*. Part III. The genera *Anthrenocerus* ARROW and *Orphinus* MOTSCH.; also a note on the genus *Cryptorhopalum* GUÉR. Proc. Linn. Soc. N. S. Wales, Sydney, **68** : 57-63, 5 ff.
- ARMSTRONG J. W. T. 1945. On Australian *Dermestidae*. Part IV. Notes and the description of a new genus and four new species. Proc. Linn. Soc. N. S. Wales, Sydney, **70** : 47-52, 3 ff.
- ARMSTRONG J. W. T. 1949. On Australian *Dermestidae*. Part V. Notes and the description of four new species. Proc. Linn. Soc. N. S. Wales, Sydney, **74** : 107-111, 1 f.
- ARROW G. J. 1915. Notes on the Coleopterous Family *Dermestidae*, and Descriptions of some new Forms in the British Museum. Ann. Mag. Nat. Hist., London, ser. 8, **15** : 425-451.
- ARROW [G. J.]. 1916. A new Species of *Thaumaglossa* bred from the Egg-clusters of *Mantidae*. Trans. Ent. Soc. London, London, **1915** : cxii-cxiii.
- ARROW G. J. 1917. The Khapra Beetle (*Trogoderma khapra*, sp. n.), an Indian Grain-pest. Ann. Mag. Nat. Hist., London, ser. 8, **19** : 481-482.
- ARROW G. J. 1937. Notes on some Clavicorn *Coleoptera* and Descriptions of a few new Species and Genera. Ann. Mag. Nat. Hist., London, ser. 10, **20** : 101-113.
- AUBÉ Ch. 1850. Description de quelques Insectes Coléoptères appartenant à l'Europe et à l'Algérie. Ann. Soc. Ent. France, Paris, ser. 2, **8** : 299-346.
- AUBÉ Ch. 1866. Description de Nouvelles espèces de Coléoptères de France. Ann. Soc. Ent. France, Paris, ser. 4, **6** : 161-164.
- BACK E. A. et COTTON R. T. 1937. The furniture carpet beetle (*Anthrenus vorax* WATERHOUSE), a pest of increasing importance in the United States. Proc. Ent. Soc. Wash., Washington, **38** : 189-198, 2 ff.
- BALLION E. 1871. Eine centurie neuer Käfer aus der Fauna des russischen Reiches. Bull. Soc. Imp. Nat. Moscou, Moscou, **43** II : 320-353.
- BALLION E. 1878. Verzeichniss der im Kreise von Kuldsha gesammelten Käfer. Bull. Soc. Imp. Nat. Moscou, Moscou, **53** I : 253-389.

- BARBER H. S. 1942. Some synonymy in *Dermestes* (*Coleoptera*). Bull. Brooklyn Ent. Soc., Lancaster, n. s. **37** : 174-176.
- BARBER H. S. 1947. On the Odd, or Tissue-paper, beetle supposed to be *Thylodrias contractus* (*Coleoptera: Dermestidae*). Ann. Ent. Soc. America, **40** : 344-349, 4 ff.
- BARBER H. S. 1951. Another name for the furniture carpet beetle. Coleopt. Bull., Rochester, **5** : 44-45.
- BAUDI a SELVE F. 1870. Coleopterorum messis in insula Cypro et Asia minore ab Eugenio TRUQUI congregatae recensitio: de Europaeis notis quibusdam additis. Pars tertia. Berlin. Ent. Z., Berlin, **14** : 49-90.
- BEAL R. S. 1952. Description of a new Arizona *Thaumaglossa* (*Coleoptera: Dermestidae*). Pan-Pacif. Entomol., San Francisco, **28** : 171-172, 1 f.
- BEAL R. S. 1954. Classification of the Dermestid genus *Dearthrus* with description of a new western species (*Coleoptera*). Pan-Pacif. Entomol., San Francisco, **30** : 231-234, 2 ff.
- BEAL R. S. 1954b. A revision of the species included in the genus *Novelsis* (*Coleoptera: Dermestidae*). Trans. Amer. Ent. Soc., Philadelphia, **80** : 73-90, 3 ff.
- BEAL R. S. 1954c. Biology and Taxonomy of the Nearctic Species of *Trogoderma* (*Coleoptera: Dermestidae*). Univ. Calif. Publ. Ent., Berkeley - Los Angeles, **10** : 35-101, 18 ff.
- BEAL R. S. 1956. Synopsis of the economic species of *Trogoderma* occurring in the United States with description of a new species (*Coleoptera: Dermestidae*). Ann. Ent. Soc. America, Columbia, **49** : 559-566, 9 ff.
- BEAL R. S. 1959. A key to the Nearctic genera of *Dermestidae*. Coleopt. Bull., Rochester, **13** : 99-101.
- BEAL R. S. 1959b. Notes on the biology and systematics of the Dermestid beetle genus *Aspectus* with descriptions of two new species. Ann. Ent. Soc. America, Washington, **52** : 132-137, 2 ff.
- BEAL R. S. 1960. Description of a California species of *Ctesias* (*Dermestidae*). Coleopt. Bull., Washington, **14** : 95-96, 1 f.
- BEAL R. S. 1961. *Coleoptera: Dermestidae*. Insects of Micronesia, Honolulu, **16** : 109-131, 5 ff.
- BEAL R. S. 1964. A new superficially cryptic species of *Trogoderma* from the southwestern United States (*Coleoptera: Dermestidae*). Proc. Ent. Soc. Wash., Washington, **66** : 79-84, 5 ff.
- BEAL R. S. 1967. A Revisionary Study of the North American Dermestid Beetles Formerly Included in the Genus *Perimegatoma* (*Coleoptera*). Misc. Publ. Ent. Soc. Amer., College Park, Maryland, **5** : 281-312, 8 ff.
- BEDEL L. 1884. [*Anthrenus miniopictus* n. sp.]. Ann. Soc. Ent. France, Paris, ser. 6, **4** : XXI.
- BIELZ E. A. 1850. Entomologische Notizen. Verh. Mitt. siebenbürg. Ver. Naturw., Hermannstadt, **1** : 179-181.
- BLACKBURN T. 1891. Further Notes on Australian *Coleoptera*, with Descriptions of New Genera and Species. IX. Trans. R. Soc. S. Austral., Adelaide, **14** : 65-153.
- BLACKBURN T. 1892. Further Notes on Australian *Coleoptera*, with Descriptions of New Genera and Species. XI. Trans. R. Soc. S. Austral., Adelaide, **15** : 20-73.
- BLACKBURN T. 1892b. Further Notes on Australian *Coleoptera* with Descriptions of New Genera and Species. XII. Trans. R. Soc. S. Austral., Adelaide, **15** : 207-261.
- BLACKBURN T. 1895. Notes on Australian *Coleoptera* with descriptions of new species. Part XV. Proc. Linn. Soc. N. S. Wales, Sydney, ser. 2, **9** : 85-108.
- BLACKBURN T. 1903. Further Notes on Australian *Coleoptera*, with Descriptions of New Genera and Species. XXXII. Trans. R. Soc. S. Austral., Adelaide, **27** : 91-182.
- BLACKBURN T. et SHARP D. 1885. Memoirs on the *Coleoptera* of the Hawaiian Islands. Sci. Trans. Roy. Dublin Soc., Dublin, ser. 2, **3**, 6 : 119-290, pls. IV-V.
- BLACKWELDER R. E. 1945. Checklist of the Coleopterous insects of Mexico, Central America the West Indies, and South America. Part 3. Bull. U. S. Nat. Mus., Washington, **185**, 3 : III-IV + 343-550.

- BLAIR K. G. 1929. Fauna of the Batu Caves, Selangor. xvii. *Coleoptera*. J. Feder. Malay Mus., Singapore, **14** : 381–387, 2 ff.
- BLAIR K. G. 1941. A new genus of *Dermestidae* (*Col.*) from New Zealand. Ent. Monthly Mag., London, **77** : 15–16.
- BLAISDELL F. E. 1927. Miscellaneous studies in the *Coleoptera* No. 2. Pan-Pacif. Entomol., San Francisco, **3** : 163–168.
- BLAISDELL F. E. 1927b. A blind beetle excavated from an Egyptian city's ruins dating between 11 and 235 A. D. Proc. Ent. Soc. Wash., Washington, **29** : 121–125, pl. 6.
- BLANCHARD É. 1843. Insectes de l'Amérique Méridionale, recueillis par Alcide d'ORBIGNY. [Without Carabiques, Hydrocanthares and Palpicornes]. In: A. ORBIGNY, Voyage dans l'Amérique Méridionale. **6**, 2. Paris, pp. 60–222, pls. 6–32.
- BLATCHLEY W. S. 1914. Notes on the winter and early spring *Coleoptera* of Florida, with descriptions of new species. Canad. Entomol., London, Ont., **46** : 61–66.
- BLATCHLEY W. S. 1922. Some new and rare *Coleoptera* from Southwestern Florida. Canad. Entomol., Orilla, **54** : 27–33.
- BODEMEYER E. v. 1900. Quer durch Klein-Asien in den Bulghar-Dagh. Eine naturwissenschaftliche Studien-Reise. Emmendingen, V + 169 pp.
- BOEBER [J.]. 1802. Nachträge für die Klasse der im Russischen Reiche vorhandenen Insecten. In: GEORGII J. G. Nachträge für dessen Geographisch-physikalische und Naturhistorische Beschreibung des Russischen Reichs. Königsberg, pp. 331–344.
- BÖRNER I. K. H. 1776. Beschreibung eines neuen Insekts, des *Dermestes 6-dentatus*. Oek. Nachr. Ges. Schlesien, Breslau, **4** : 78–80.
- BOHEMAN C. H. 1851. *Insecta Caffrariae annis 1838–1845 a J. A. WAHLBERG collecta. Coleoptera pars I, fasc. II.* Holmiae, pp. 299–625, 2 pls.
- BRAHM N. J. 1790. Insektenkalender für Sammler und Oekonomen. Erster Theil, [part. I]. Mainz, LXLII + 128 pp.
- BRAHM N. J. 1791. Insektenkalender für Sammler und Oekonomen. Handbuch der ökonomischen Insekten-geschichte in Form eines Kalenders bearbeitet. Erster Theil, [part. II]. Mainz, pp. 129–248.
- BRISOUT DE BARNEVILLE Ch. 1866. Coléoptères nouveaux trouvés en Espagne pendant l'Excursion de la Société en 1865. Ann. Soc. Ent. France, Paris, ser. 4, **6** : 355–426.
- BROUN T. 1880. Manual of the New Zealand *Coleoptera*. Wellington, xx + 651 pp.
- BROUN T. 1886. Manual of the New Zealand *Coleoptera*. Parts III. and IV. Wellington, xvii + 745 – 973 pp.
- BROUN T. 1893. Manual of the New Zealand *Coleoptera*. Parts V., VI., VII. Wellington, xvii + 975–1504 pp.
- BROUN T. 1911. Additions to the Coleopterous Fauna of the Chatham Islands. Trans. N. Z. Inst., Wellington, **43** : 92–115.
- BROUN T. 1914. Descriptions of New Genera and Species of *Coleoptera* (Part III). Bull. N. Z. Inst., Wellington, **1** : 143–266.
- BRULLÉ M. 1832. Expédition scientifique de Morée. Section des sciences physiques. Tome III. 1re Partie. Zoologie. Deuxième Section – Des animaux articulés. Paris, 395 pp., 22 pls.
- CAPRA F. 1951. Che cosa e l' „*Anthrenus*” Rossi PONZA, 1805? (*Coleopt. Dermestidae et Byrrhidae*). Boll. Soc. Ent. Ital., Genova, **81** : 44–45.
- CASEY T. L. 1900. Review of the American *Corylophidae*, *Cryptophagidae*, *Tritomidae*, and *Dermestidae*, with other studies. J. N. Y. Ent. Soc., New York, **8** : 51–172, 21 ff.
- CASEY T. L. 1916. Memoirs on the *Coleoptera*. VII. Lancaster, Pa., [IV] + 300 pp.
- CASEY T. L. 1924. Memoirs on the *Coleoptera*. XI. Lancaster, Pa., 347 pp.

- CASTELNAU [LAPORTE DE, F. L.] 1840. Histoire Naturelle des Insectes Coléoptères. II. Paris, 563 pp., pls. 20–36.
- CEDERHIELM I. 1798. Faunae Ingriæ Prodromus exhibens methodicam descriptionem Insectorum Agri Petropolenensis praemissa Mammalium, Avium, Amphibiorum et Piscium enumeratione. Lipsiae, XVIII + 348 pp., 3 pls.
- CHAMPION G. C. 1922. Some Indian Coleoptera (7). Ent. Monthly Mag., London, 58 : 31–34, 68–76, 1 f.
- CHAO Y. et LEE H. 1966. A study of Chinese *Trogoderma* BERTHOLD (Coleoptera, Dermestidae). Acta Zootax. Sin., Peking, 3 : 245–252, 2 ff., 2 pls.
- CHEVROLAT A. 1863. Coléoptères de l'île de Cuba. (Suite) (1). Notes, synonymies et descriptions d'espèces nouvelles. Quatrième mémoire. Famille des Histériens, Phalacrides, Nitidulaires, Trogositaires, Colydiens, Rhyzodides, Cucujipes, Mycetophagides, Dermestins, Byrrhiens et Chélonariides. Ann. Soc. Ent. France, Paris, ser. 4, 3 : 589–620.
- CHOBAUT A. 1897. Description de deux Coléoptères nouveaux au sud de l'Algérie. Bull Soc. Ent. France, Paris, 1897 : 277–279.
- CHOBAUT A. 1898. Description de quelques espèces et variétés nouvelles de Coléoptères Algériens. Rev. Ent., Caen, 17 : 74–88.
- COCKERELL T. D. A. 1917. Arthropods in Burmese Amber. Psyche, Boston, 24 : 40–45.
- COUCKE E. 1892. Les Brachymères de Belgique et des Régions Voisines. Ann. Soc. Ent. Belg., Bruxelles, 36 : 69–79.
- COUCKE E. 1895. Les *Dermestes bicolor* FABR. trouvés à fleurs par Mr. BIVORT. Changement à apporter à ma liste des Brachymères de Belgique et des Régions Voisines. Ann. Soc. Ent. Belg., Bruxelles, 39 : 466–467.
- CREUTZER Ch. 1799. Entomologische Versuche. Wien, 144 pp., 3 pls.
- CRISTOFORI J. et JAN G. 1832. Catalogus in IV. sectiones divisus rerum naturalium in museo extantium. Sectio III.a Entomologia, pars I.a Conspectus Methodicus Insectorum, fasc. I.us Coleoptera. Mediolani, [4] + 16 + 111 pp.
- CROTCH G. R. 1873. Check list of the Coleoptera of America, North of Mexico. Salem, Mass., 136 pp.
- CURTIS J. 1838. British Entomology; being illustrations and descriptions of the Genera of Insects found in Great Britain and Ireland: containing Coloured Figures from Nature of the most rare and beautiful species, and in many instances of the plants upon which they are found. XV. London, pp. 674–721, pls. 674–721.
- DALLA TORRE K. W. v. 1879. Die Käferfauna von Oberösterreich. Systematisches Verzeichnis der in Oberösterreich bisher beobachteten Käfer. Jber. Ver. Naturk. Linz, Linz, 10 : 1–125.
- DALLA TORRE K. W. v. 1911. Dermestidae. In: Coleopterorum Catalogus, 14, pars 33, Berlin, pp. 39–96.
- DEGEER Ch. 1774. Mémoires pour servir à l'Histoire des Insectes. IV. Stockholm, XII + 457 pp., 19 pls.
- DEJEAN [P. F.J. 1821. Catalogue de la collection de Coléoptères. Paris, VIII + 136 pp.
- DEJEAN [P. F.J. 1837. Catalogue des Coléoptères de la collection. Troisième édition, revue, corrigée et augmentée. Paris, XIV + 503 pp.
- DESBROCHERS DES LOGES J. 1898. Quelques Matériaux pour la Faune des Coléoptères de Barbarie. Frelon, Chateauroux, 6 : 65–82.
- [DOHRN C. A.] 1858. Catalogus Coleopterorum Europæ. Siebente Auflage. Stettin, XV + 104 pp.
- DUFTSCHMID C. 1825. Fauna Austriae. Oder Beschreibung der österreichischen Insecten, für angehende Freunde der Entomologie. III. Linz, 289 pp.
- DUNSTAN B. 1923. Mesozoic Insects of Queensland. Part I. Introduction and Coleoptera. Publ. Queensl. Geol. Surv., Brisbane, 273 : 1–88, 7 pls.
- EICHHOFF W. 1863. Sammelberichte aus Hilchenbach bei Siegen. Berlin. Ent. Z., Berlin, 7 : 436–437.

- EICHLER W. 1930. Chrząszcze okolic Tyflisu. Pol. Pismo Ent., Lwów, **9** : 213–258.
- ERICHSON W. F. 1842. Beitrag zur Insecten-Fauna von Vandiemensland, mit besonderer Berücksichtigung der geographischen Verbreitung der Insecten. Arch. Naturges., Berlin, **8**, 1 : 83–287, pls. IV–V.
- ERICHSON W. F. 1843. Beitrag zur Insecten-Fauna von Angola, in besonderer Beziehung zur geographischen Verbreitung der Insecten in Afrika. Arch. Naturges., Berlin, **9**, 1 : 199–267.
- ERICHSON W. F. 1846. Naturgeschichte der Insecten Deutschlands. Erste Abtheilung. *Coleoptera*. Dritter Band, III Lieferung. Berlin, pp. 321–480.
- ERICHSON W. F. 1847. Conspectus Insectorum Coleopterorum, quae in Republica Peruana observata sunt. Arch. Naturges., Berlin, **13**, 1 : 67–185.
- ERICHSON W. F. 1847 b. Bericht über die wissenschaftlichen Leistungen in der Naturgeschichte der Insecten, Arachniden, Crustaceen und Entomostraceen während des Jahres 1846. Arch. Naturges., Berlin, **13**, 2 : 65–208.
- ESCALERA M. M. 1914. Los Coleópteros de Marruecos. Trab. Mus. Cienc. Nat., Ser. Zool., Madrid, **11** : 1–553.
- EVERTS E. 1898. *Coleoptera Nederlandica*. De schildvleugelige insecten van Nederland en het aangrenzend gebied. I. 'sGravenhage, VIII + 677 pp., 62 ff.
- EVERTS E. 1923. *Trogoderma granarium* EVERTS. (syn.: *khapra* ARROW). De Khaprakever. Ent. Ber. Nederl. Ent. Ver., Enkhuizen, **6** : 168–170.
- FABRICIUS I. Ch. 1775. Systema Entomologie, sistens Insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus. Flensburgi et Lipsiae, 32 + 832 pp.
- FABRICIUS I. Ch. 1781. Species Insectorum exhibentes eorum differentias specificas, synomyma auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus. I. Hamburgi et Kilonii, VIII + 552 pp.
- FABRICIUS I. Ch. 1787. Mantissa Insectorum sistens eorum species nuper detectas adiectis characteribus genericis, differentiis specificis, emendationibus, observationibus. I-II. Hafniae, I:XX + 348 pp., II: 382 pp.
- FABRICIUS J. Ch. 1792. Entomologia Systematica emendata et aucta. Secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus. I. 1–2. Hafniae, 1:XX + 330 pp., 2:538 pp.
- FABRICIUS J. Ch. 1798. Supplementum Entomologiae Systematicae. Hafniae, II + 572 pp.
- FABRICIUS I. Ch. 1801. Systema Eleutherotorum secundum ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus. I. Kiliae, XXIV + 506 pp.
- FAIRMAIRE L. 1850. Essai sur les Coléoptères de la Polynésie. (Suite et fin). Rev. Mag. Zool., Paris, ser. 2, **2** : 181–185.
- FAIRMAIRE L. 1859. Miscellanea entomologica. Troisième partie. Ann. Soc. Ent. France, Paris, ser. 3, **7** : 21–64.
- FAIRMAIRE L. 1863. Coléoptères d'Algérie rapportés par M. OGIER DE BAULNY. Ann. Soc. Ent. France, Paris, ser. 4, **3** : 637–648.
- FAIRMAIRE L. 1879. Descriptions de Coléoptères nouveaux du nord de l'Afrique. 1re partie. Ann. Soc. Ent. France, Paris, ser. 5, **9** : 155–172.
- FAIRMAIRE L. 1883. Descriptions de Coléoptères recueillis par le baron BONNAIRE en Algérie. Ann. Soc. Ent. Belg., Bruxelles, **27**, C.—R. : CLVI—CLIX.
- FAIRMAIRE L. 1884. Note sur quelques Coléoptères de Magellan et de Santa-Cruz. Ann. Soc. Ent. France, Paris, ser. 6, **3** : 483–506.
- FAIRMAIRE L. 1886. Descriptions de Coléoptères de l'intérieur de la Chine. Ann. Soc. Ent. France, Paris, ser. 6, **6** : 303–356.
- FAIRMAIRE L. 1889. Coléoptères de l'intérieur de la Chine. Ann. Soc. Ent. France, Paris, ser. 6, **9** : 5–81.

- FAIRMAIRE L. 1897. Matériaux pour la faune Coléoptérique de la région Malgache. 3e note. Ann. Soc. Ent. Belg., Bruxelles, **41** : 92–119.
- FAIRMAIRE L. et COQUEREL C. 1860. Essai sur les Coléoptères de Barbarie. Deuxième partie (1). Ann. Soc. Ent. France, Paris, ser. 3, **8** : 145–176.
- FALDERMANN F. 1835. Additamenta entomologica ad Faunam Rossicam in itineribus Jussu Imperatoris Augustissimi annis 1827–1831 a Cl. MÉNÉTRIÉS et SZOVITZ susceptis collecta, in lucem adita. *Coleoptera Persico-Armeniaca*. Nouv. Mém. Soc. Natural. Moscou, **4** : 1–4 + 1–310, pls. 1–10.
- FALL H. C. 1897. A list of the *Coleoptera* of the Southern California Islands, with notes and descriptions of new species. Canad. Entomol., Orillia, **29** : 233–244.
- FALL H. C. 1926. A list of the *Coleoptera* taken in Alaska and adjacent parts of the Yukon territory in the summer of 1924 (Continued from page 154). Pan-Pacif. Entomol., San Francisco, **2** : 191–208.
- FAUVEL A. 1903. Faune Analytique des Coléoptères de la Nouvelle Calédonie. Rev. Ent., Caen, **22** : 203–378.
- FLEUTIAUX E. 1887. Descriptions de Coléoptères nouveaux de l'Annam. Rapportés par M. le capitaine DELAUNEY. Ann. Soc. Ent. France, Paris, ser. 6, **7** : 59–68, 4 pls.
- FLEUTIAUX E. et SALLÉ A. 1890. Liste des Coléoptères de la Guadeloupe et descriptions d'espèces nouvelles. Ann. Soc. Ent. France, Paris, ser. 6, **9** : 351–484, 1 map, 8 pls.
- FLUITER [H. J.] 1952. [Discussie]. Tijdschr. Ent., Amsterdam, **95** : IX–X.
- FORSTER J. R. 1771. Novae species Insectorum. Centuria I. Londini, VIII + 100 pp.
- FOURCROY A. F. 1785. Entomologia Parisiensis; sive Catalogus Insectorum quae in Agro Parisiensi reperiuntur; Secundum methodum Geoffraeanam in sectiones, genera et species distributus: Cui addita sunt nomina trivalia et fere trecentae novae Species. Pars Prima. Parisii, VIII + 231 pp.
- FRIVALDSZKY J. 1882. *Coleoptera Europaea nova*. Természetr. Füz., Budapest, **5** : 26–29.
- FRIVALDSZKY J. 1892. *Coleoptera in expeditione D. Comitis Belae SzÉCHENYI in China, praecipue boreali, a Dominis Gustavo KREITNER et Ludovico LÓCZY anno 1879 collecta*. (Pars secunda). Természetr. Füz., Budapest, **15** : 114–125.
- FUENTE J. M. 1930. Catálogo sistemático-geográfico de los Coleópteros observados en la península ibérica, Pirineos propiamente dichos y Baleares. Tomo III, pp. (1)–(16). Bol. Soc. Ent. EspaÑ., Zaragoza, **12** : 43–58.
- FUESSLIN J. C. 1775. Verzeichnis der ihm bekannten Schweizerischen Insekten mit einer ausgemahlten Kupferplatte: nebst der Ankündigung eines neuen Insecten Werks. Zürich und Winterthur, XII + 62 pp., 1 pl.
- FULLAWAY D. T. 1922. Notes on Immigrant *Coleoptera*. Proc. Hawaii. Ent. Soc., Honolulu, **5** : 75–82.
- GANGLBAUER L. 1904. Verzeichnis der auf der dalmatinischen Insel Meleda vorkommenden Koleopteren nach den Sammelergebnissen des Herrn Forstrates Alois GOBANZ. Verh. Zool.-Bot. Ges. Wien, Wien, **54** : 645–660.
- GANGLBAUER L. 1904b. Die Käfer von Mitteleuropa. Die Käfer der österreichisch-ungarischen Monarchie, Deutschlands, der Schweiz, sowie des französischen und italienischen Alpengebietes. Vierter Band, erste Hälfte. *Dermestidae, Byrrhidae, Nosodendridae, Georyssidae, Dryopidae, Heteroceridae, Hydrophilidae*. Wien, 286 pp., 12 ff.
- GBLER F. 1830. Bemerkungen über die Insecten Sibiriens, vorzüglich des Altai. In: Carl Friedrich von LEDEBOUR's Reise durch das Altai-Gebirge und die soongorische Kirgis-Steppe. Zweiter Theil, III. Berlin, 228 pp.
- GBLER F. 1841. Notae et additamenta ad Catalogum Coleopterorum Sibiriae occidentalis et confinis Tatariae operis: (C. F. von LEDEBOUR's Reise in das Altaigebirge und die soongarische Kirgissteppe. Zweyter Theil. Berlin. 1830). Fasciculus secundus. Bull. Soc. Imp. Nat. Moscou, Moscou, **1841**, II : 577–625.

- GEBLER F. 1847. Verzeichniss der im Kolywano-Woskresenskischen Hüttenbezirke Süd-West Sibiriens beobachteten Käfer mit Bemerkungen und Beschreibungen. Bull. Soc. Imp. Nat. Moscou, Moscou, 20, II : 391–512.
- GEMMINGER [M.] et HAROLD [E.]. 1868. Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Tom III. *Histeridae ... Lucanidae*. Monachii, pp. 753–978 + [5] pp.
- GEMMINGER [M.] et HAROLD [E.]. 1869. Catalogus Coleopterorum hucusque descriptorum synonymicus et systematicus. Tom VI. *Rhipidoceridae ... Cioidae*. Monachii, pp. 1609–1800 + [5] pp.
- GENÉ J. 1836. De quibusdam Insectis Sardiniae novis aut minus cognitis. Fasciculus I. Taurini, 39 pp., 1 pl.
- GENÉ J. 1839. De quibusdam Insectis Sardiniae novis aut minus cognitis. Fasciculus II. Taurini, 44 pp., 2 pls.
- GERHARDT J. 1910. Neuheiten der schlesischen Käferfauna aus dem Jahre 1909 (Col.). Dtsch. Ent. Z., Berlin, 1910 : 554–557.
- GERMAIN Ph. 1892. Notes sur les Coléoptères du Chili. Renseignements et observations; descriptions d'espèces nouvelles; rectifications; indications de synonymie. Act. Soc. Sci. Chile, Santiago, 2 : 241–261, 4 ff.
- GERMAR E. F. 1813. Neue Insekten. Mag. Ent. Germar, Halle, 1, 1 : 114–133.
- GERMAR E. F. 1824. Insectorum Species novae aut minus cognitae, descriptionibus illustratae. I — Coleoptera. Halae, XXIV + 624 pp., 2 pls.
- GERMAR E. F. 1839. Fauna Insectorum Europae. Fasciculus 21. Halae, 25 pp., 25 pls.
- GERSTAECKER A. 1871. Beitrag zur Insektenfauna von Zanzibar. III. Coleoptera. Arch. Naturges., Berlin, 37, 1 : 42–86.
- GERSTAECKER A. 1884. Bestimmung der von Herrn Dr. G. A. FISCHER während seiner Reise nach dem Massai-Land gesammelten Coleopteren. Jahresbuch Wiss. Anst. Hamburg, Hamburg, 1883 : 43–63.
- GESTRO R. 1889. Viaggio ad Assab nel Mar Rosso dei signori G. DORIA ed O. BECCARI con il R. Avviso "Esploratore" dal 16 novembre 1879 al 26 febbraio 1880. IV. Coleotteri. Ann. Mus. Stor. Nat. Genova, Genova, 27 : 5–72.
- GMELIN J. F. [1790]. Caroli a LINNÉ Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima tertia, aucta, reformata. I, 4. [LIPSIAE], pp. 1517–2224.
- GODARD A. 1883. Description d'une espèce nouvelle de Dermestide. Ann. Soc. Linn. Lyon, Lyon, 29 : 383–384.
- GOEZE J. A. E. 1777. Entomologische Beyträge zu des Ritter LINNÉ zwölften Ausgabe des Natursystems. Erster Theil. Leipzig, XVI + 736 pp.
- GOZIS M. 1886. Recherche de l'espèce typique de quelques anciens genres. Rectifications synonymiques et notes diverses. Montluçon, 36 pp.
- GREDLER V. P. 1878. Zur Käfer-Fauna Central-Afrikas. Verh. Zool.-Bot. Ges. Wien, Wien, 27 : 501–522.
- GRIMMER C. 1841. Steiermarks Coleopteren mit einhundertsechs neu beschriebenen Species. Graz, 50 + 4 pp.
- GRÜSS J. 1930. Ein altägyptischer blinder Biertrinker vor 5000 Jahren. Wochenschr. f. Brauerei, 47 : 181–182, 2 ff.
- GUÉRIN-MÉNEVILLE F. E. 1838. Insectes du voyage de la Favorite. Mag. Zool., Paris, 8 : 1–80, pls. 225–238.
- GUÉRIN-MÉNEVILLE F. E. 1838b. Iconographie du Règne Animal de G. CUVIER, ou représentation d'après nature de l'une des espèces les plus remarquables, et souvent non encore figurées, de chaque genre d'animaux. Avec un texte descriptif mis au courant de la science. Ouvrage pouvant servir d'atlas à tous les traités de zoologie. Insectes. Paris, 576 pp. [Pages 65–72 were edited in 1838].

- GUÉRIN-MÉNEVILLE [F. E.]. 1838c. Note sur le genre de Coléoptères clavicornes, nommé par LATREILLE *Globicornis*, et description d'une espèce nouvelle de ce genre. Rev. Zool., Paris, **1** : 135–139.
- GYLLENHAL L. 1808. Insecta Suecica. Classis I. Coleoptera sive Eleuterata. Tomus I. Scaris, XII + 572 pp.
- GYLLENHAL L. 1813. Insecta Suecica. Classis I. Coleoptera sive Eleuterata. Tomi I, Pars III. Scaris, 734 pp.
- GYLLENHAL L. 1827. Insecta Suecica. Classis I. Coleoptera sive Eleuterata. Tom I, Pars IV. cum appendice ad partes priores. Lipsiae, X + 762 pp.
- HABERFELNER J. 1889. Ueber entomologische Zuchthäuser. Dtsch. Ent. Z., Berlin, **1889** : 389–394.
- HÄNEL K. 1935. *Anthrenus verbasci* a. nov. *bifasciata*. Ent. Bl., Krefeld, **31** : 76–77.
- HAROLD [E.]. 1868. Abänderungen vergebener Namen. Coleopt. Hefte, München, **4** : 104.
- HAROLD E. 1877. Beiträge zur Käferfauna von Japan. (Zweites Stück). Dtsch. Ent. Z., Berlin, **21** : 337–367.
- HAUSER F. 1894. Beitrag zur Coleopteren-Fauna von Transcaspien und Turkestan. Zusammengestellt unter Mitwirkung der Herren Dr. EPPESHEIM, ESCHERICH, FAUST, Dr. von HEYDEN, KUVERT, REITTER und WEISE. Dtsch. Ent. Z., Berlin, **1894** : 17–74, 4 ff., pl. I, f. 1.
- [HAUSMANN J. F. L.J. 1799. Entomologische Bemerkungen. Braunschweig u. Helmstedt, 4 + 64 pp.
- HAVELKA J. 1951. *Dermestes haemorrhoidalis* KÜST., collected in Prague (Coleoptera, Dermest.). Folia Ent., Brno, **14** : 135–143.
- HAVELKA J. 1951b. *Dermestes maculatus pakistanius* ssp. n. introduced from Pakistan to Č. S. R. (3rd contribution to knowledge of *Dermestes* L.) (Col., Dermestidae). Folia Ent., Brno, **14** : 143–144.
- HAVELKA J. 1957. Contribution to the knowledge of *Dermestes* in China. *Dermestes tessellatocollis* MOTSCH. (Col. Dermestidae) (2nd communication on *Dermestes* L.). Fol. Ent. Hung., Budapest, s. n., **10** : 105–109, 4 ff.
- HEER O. 1841. Die Käfer der Schweiz, mit besonderer Berücksichtigung ihrer geographischen Verbreitung. Erster Theil. Dritte Lieferung. Aus dem fünften Bande der Neuen Denkschriften der allgemeinen schweizerischen Gesellschaft für die gesammten Naturwissenschaften besonders abgedruckt. Neuchatel, 79 pp.
- HEER O. 1841b. Fauna Coleopterorum Helvetica. Pars I. Turici, XII + 652 pp.
- HEER O. 1847. Die Insektenfauna der Tertiärgebilde von Oeningen und von Radoboj in Croatiens. Erste Abtheilung: Käfer. Leipzig, 229 pp., 8 pls.
- HEER O. 1856. Verzeichniss der Insecten. A. Von Lanzarote. B. Von Fuertaventura. In: G. HARTUNG. Die geologischen Verhältnisse der Inseln Lanzarote und Fuertaventura. Neue Denkschr. Allg. Schweiz. Ges. Naturw., Zürich, **1856** : 140–142.
- HENSHAW S. 1885. List of the Coleoptera of America, North of Mexico. Philadelphia, [4] + 161 pp.
- HERBST J. F. W. 1783. Kritisches Verzeichniss meiner Insekten-Sammlung. Arch. Insect. Gesch., Zürich, **4** : 1–72, pls XIX–XXIII.
- HERBST J. F. W. 1792. Natursystem aller bekannten in- und ausländischen Insekten, als eine Fortsetzung der von Büffonschen Naturgeschichte. Der Käfer vierter Theil. Berlin, VIII + 197 pp., 12 pls.
- HERBST J. F. W. 1797. Natursystem aller bekannten in- und ausländischen Insekten, als eine Fortsetzung der von Büffonschen Naturgeschichte. Der Käfer siebenten Theil. Berlin, 346 pp., 26 pls.
- HEYDEN L. 1879. Die coleopterologische Ausbeute des Prof. Dr. REIN in Japan 1874–1875. Dtsch. Ent. Z., Berlin, **23** : 321–365.
- HEYDEN L. 1881. Catalog der Coleopteren von Sibirien mit Einschluss derjenigen der Turanischen Länder, Turkestans und der chinesischen Grenzgebiete. Mit specialler Angabe der einzelnen Fundorte in Sibirien und genauer Citirung der darauf bezüglichen einzelnen Arbeiten nach eigenem Vergleich, sowie mit besonderer Rücksicht auf die geographische Verbreitung der einzelnen Arten über die Grenzländer, namentlich Europa und Deutschland. Berlin, XXIV + 224 pp.

- HEYDEN L. 1890. Aufzählung von Käfer-Arten aus Tunis und Tripolis aus Loosen von M. QUEDENFELDT Dtsch. Ent. Z., Berlin, **1890** : 65–78.
- HEYDEN L. 1890b. XIII. Beitrag zur Coleopteren-Fauna von Turkestan. Unter Mitwirkung der Herrn REITTER und WEISE. Dtsch. Ent. Z., Berlin, **1890** : 353–367.
- HEYDEN L. 1893. Catalog der Coleopteren von Sibirien, mit Einschluss derjenigen des östlichen Caspi-Gebietes, von Turkmenien, Turkestan, Nord-Thibet und des Amur-Gebietes. Mit specieller Angabe der einzelnen Fundorte und genauer Citirung der darauf bezüglichen Literatur. Nachtrag I. Berlin, 217 pp.
- HEYDEN L., REITTER E. et WEISE J. 1884. Berichtigungen und Zusätze zum „Catalogus Coleopterorum Europae et Caucasi“. Wien. Ent. Ztg., Wien, **3** : 177–184, 207–213.
- HICKS E. A. 1953. Observations on the insect fauna of birds' nests. J. Kansas Ent. Soc., Manhattan, Kans., **26** : 11–18.
- HINTON H. E. 1943. Description and figure of a new *Anthrenus* (Col., Dermestidae). Ent. Monthly Mag., London, **79** : 14–16, 3 ff.
- HINTON H. E. 1945. The species of *Anthrenus* that have been found in Britain, with a description of a recently introduced species (Coleoptera, Dermestidae). Entomologist, Wimbledon, **78** : 6–9.
- HINTON H. E. 1945b. A Monograph of the Beetles associated with stored products. Vol. I. London, VIII + 443 pp., 505 ff.
- HOPE F. W. 1834. [Some insects which had been extracted by Mr. PETTIGREW from the skull of a mummy]. Ent. Mag., London, **2** : 186–187.
- HOPE F. W. 1834b. In: PETTIGREW T. J., History of Egyptian Mummies, ... London.
- HOPE F. W. 1834c. Notice of several Species of Insects found in the heads of Egyptian Mummies. Trans. Ent. Soc. London, London, **1** : xi–xiii.
- HOPE F. W. 1840. The Coleopterist's Manual, part the third, containing various families, genera, and species, of beetles, recorded by LINNAEUS and FABRICIUS. Also, descriptions of newly discovered and unpublished Insects. London, [6] + 191 pp., 3 pls.
- HOPE F. W. 1843. Descriptions of new species of insects collected at Adelaide in South-Western Australia by Mr. FORTNUM. Ann. Mag. Nat. Hist., London, **11** : 317–319.
- HOPE F. W. 1845. Descriptions of some new Species of Coleoptera from Adelaide in New Holland. Trans. Ent. Soc. London, London, **4** : 100–113, 2 pls.
- HOPPING R. 1932. A taxonomic note. Canad. Entomol., Ottawa, **64** : 173.
- HORN G. H. 1875. Synonymical Notes and Description of New Species of North American Coleoptera. Trans. Amer. Ent. Soc., Philadelphia, **5** : 126–156, 8 ff.
- HOWE R. W. et BURGES H. D. 1956. *Trogoderma afrum* PRIESNER, a new synonym of *T. granarium* EVERTS and a comparison with *T. versicolor* (CREUTZ.). Bull. Ent. Res., London, **46** : 773–780, 7 ff.
- ILLIGER J. K. W. 1798. Verzeichniss der Käfer Preussens. Halle, XLII + 510 pp.
- ILLIGER K. 1801. Nachtrag und Berichtigungen zum Verzeichnisse der Käfer Preussens. Mag. Insk. k., Braunschweig, **1**, 1 : VI + II + 1–94.
- JABLOKOV-CHNZORJAN S. M. 1961. Opyt vosstanovleniya genezisa fauny zhestkokrylych Armenii. Erevan, 266 pp., 1 map.
- JACQUELIN DU VAL [P. N. C.] 1859. Genera des Coléoptères d'Europe comprenant leur classification en familles naturelles, la description de tous les genres, des Tableaux synoptiques destinés à faciliter l'étude, le Catalogue de toutes les espèces de nombreux dessins au trait de caractères. II. Paris, 288 pp., 67 pls.
- J[ACQUET] E. 1886. Notes entomologiques sur la faune coléoptérologique de Sinaïa (Valachie) et de la Dobroudja (Roumanie) (suite). Échange, Lyon, **2**, 16 : 1–2.
- JACQUET E. 1886b. Note sur le *Montandonia catopoides* Nouveau genre appartenant à la famille des Dermestidae. Ann. Soc. Linn. Lyon, n. s., **32** : 319–321, 3 ff.

- JANSSON A. 1943. Coleopterologiska bidrag. 28. Vad är *Megatoma innotata* Pic? Opusc. Ent., Lund, **8** : 23–25.
- JAYNE H. F. 1882. Revision of the *Dermestidae* of the United States. Proc. Amer. Phil. Soc., Philadelphia, **20** : 343–377, pls. 1–4.
- JOHN H. 1961. Zwei neue Spezies der Gattung *Thorictodes* RTT. (*Col. Dermestidae*). Rev. Zool. Bot. Afr., Bruxelles, **64** : 41–44, 3 ff.
- KALÍK V. 1946. Příspěvek k poznání československých kožojedů (*Dermestidae*). Folia Ent., Brno, **9** : 73–77.
- KALÍK V. 1946b. Noví *Dermestidae* palearktické fauny. Folia Ent., Brno, **9** : 133–138, 4 ff.
- KALÍK V. 1948. Příspěvek k poznání Československých kožojedů (*Dermestidae*). Sborn. Přir. Klubu v Pardubicích, Pardubice, **1948**, 13 pp., 3 pls.
- KALIK V. 1950. Neue Dermestiden aus der paläarktischen Fauna. (3. Beitrag). Fol. Ent. Hung., Budapest, s. n., **3** : 61–66, 3 ff.
- KALIK V. 1951. Neue Dermestiden aus der Palaearktischen Fauna. (4. Beitrag). Ann. Hist.-Nat. Mus. Hung., Budapest, **1** : 152–156, 3 ff.
- KALIK V. 1951b. Notes sur quelques espèces et formes du genre *Dermestes* L. Folia Ent., Brno, **14** : 39–46, 12 ff., 1 map.
- KALIK V. 1951c. New *Dermestidae* of the Palaearctic fauna. (2nd Contribution). Sborn. Entom. Odd. Nář. Mus. Praha, Praha, **26**, nr 362, 4 pp., 4 ff.
- KALIK V. 1952. Neue Dermestiden aus der paläarktischen Fauna (*Col.*) (5. Beitrag). Ann. Hist.-Nat. Mus. Hung., Budapest, s. n., **2** : 75–77, 3 ff.
- KALIK V. 1952b. New and interesting species of *Dermestidae* (*Coleoptera*). Pan-Pacif. Entomol., San Francisco, **28** : 43–49, 3 ff.
- KALIK V. 1954. New and Interesting *Dermestidae* (*Coleoptera*). Ann. Mag. Nat. Hist., London, ser. 12, **7** : 367–370.
- KALIK V. 1955. *Dermestidae*. In: Parc National de L'Upemba. I. Mission G. F. de WITTE en collaboration avec W. ADAM, A. JANSSENS, L. van MEEL et R. VERHEYEN (1946–1949). Fascicule 38 (6). Bruxelles, pp. 93–99, 3 ff.
- KALIK V. 1955b. *Dermestidae* der aethiopischen Region aus den Sammlungen des Königl. Museums von Belgisch Kongo in Tervuren (Mit Bemerkungen über verwandte Arten). Rev. Zool. Bot. Afr., Bruxelles, **52** : 294–322.
- KALIK V. 1957. A new dermestid from Australia (*Col.*). Čas. Čsl. Spol. Ent., Praha, **54** : 22–23, 3 ff.
- KALIK V. 1965. Coleoptera Dermestidae. Mission zoologique de l'I. R. S. A. C. en Afrique orientale. (P. BASILEWSKY et N. LELEUP, 1957). XCII. Ann. Mus. Roy. Afr. Centr., Tervuren, Zool., **138** : 215–224, 15 ff.
- KIESENWETTER H. 1851. Énumération des Coléoptères trouvés dans le midi de la France et en Catalogne. (Ile Partie (1)). Ann. Soc. Ent. France, Paris, ser. 2, **9** : 577–656.
- KIRBY W. 1837. The Insects. In: RICHARDSON J. Fauna Boreali-Americanæ; or the Northern parts of British America: containing descriptions of the objects of Natural History collected on the late Northern Land Expeditions, under command of Captain Sir John FRANKLIN, R. N. Part the Fourth and Last. Norwich, xxxix + 325 pp., 8 pls.
- KIRSCH Th. 1871. Beiträge zur Käferfauna von Bogotà. (Sechstes Stück.). Berlin. Ent. Z., Berlin, **14** : 337–378.
- KIRSCH Th. 1873. Beiträge zur Kenntniss der Peruanischen Käferfauna auf Dr. ABENDROTH's Sammlungen basirt. Berlin. Ent. Z., Berlin, **17** : 121–152.
- KLEINE R. 1940. Übersicht über die in Pommern gefundenen Käfer, die im Verzeichnis von Albert LÜLLWITZ nicht enthalten sind. Nebst einigen Bemerkungen über schon genannte Arten. Dohrniana, Stettin, **19** : 3–28.

- KLOET G. S. et HINCKS W. D. 1945. A check list of British Insects. Stockport, lix + 483 pp.
- KLUG [J. Ch. F.] 1855. Fortsetzung der Diagnosen der neuen (und bereits seit mehreren Monaten vollständig gedruckten) Coleopteren, welche die Insectensendungen des Hrn. Dr. PETERS von Mossambique enthalten hatten, von der Familie der *Staphylinii* an bis zu den *Lamellicornia*, diese mit eingeschlossen. Ber. Bekanntm. Verh. Königl. Akad. Wiss., Berlin, 1855 : 643–660.
- KNEUCKER A. 1922. Zoologische Ergebnisse zweier in den Jahren 1902 und 1904 durch die Sinaihalbinsel unternommener botanischer Studienreisen. II. Teil. Ent. Bl., Berlin, 18 : 20–28.
- KNIEPHOF J. 1935. *Dermestes Helmi* RTT. Ent. Bl., Krefeld, 31 : 125.
- KOCHER L. 1953. Localisations nouvelles ou intéressantes de Coléoptères Marocains. Trav. Inst. Sc. Chérifien, Tanger, 7, 142 pp.
- KOCHER L. 1955. Description d'une nouvelle espèce d'anthrène (*Col. Dermest.*). C. R. Soc. Sci. Nat. Phys. Maroc, Rabat, 1955 : 40–41, 2 ff.
- KOLBE H. 1910. Die Coleopterenfauna der Seychellen. Nebst Betrachtungen über die Tiergeographie dieser Inselgruppe. Mitt. Zool. Mus. Berlin, Berlin, 5 : 1–49.
- KOLENATI F. A. 1846. Meletemata Entomologica. Fasc. V. *Insecta Caucasi. Coleoptera, Dermaptera, Lepidoptera, Neuroptera, Mecoptera, Aphaniptera, Anoplura*. Petropoli, 8 + 170 pp., pls. XVII–XIX.
- KORGE H. 1961. Beiträge zur Koleopterenfauna der Mark Brandenburg (Teil XXV). Mitt. Dtsch. Ent. Ges., Berlin, 20 : 21–27, 5 ff.
- KORGE H. 1962. Beiträge zur Koleopterenfauna der Mark Brandenburg (Teil XXVI). Mitt. Dtsch. Ent. Ges., Berlin, 21 : 73–83, 23 ff.
- KRAATZ G. 1858. Beitrag zur Käferfauna Griechenlands. Drittes Stück: *Staphylinidae* (Schluss), *Trichopterygia, Histeridae, Phalacridae, Nitidulariae, Trogositarii, Colydiidae, Cucujidae, Cryptophagidae, Thorictidae, Mycetophagidae, Dermestini, Byrrhii*. Berlin. Ent. Z., Berlin, 2 : 123–148.
- KRAATZ G. 1864. Zur kritischen Kenntniss europäischer Käferarten. Berlin. Ent. Z., Berlin, 8 : 131–142.
- KÜSTER H. C. 1846. Die Käfer Europa's. Nach der Natur beschrieben. VI Heft. Nürnberg, 100 sheets, 2 pls.
- KÜSTER H. C. 1847. Die Käfer Europa's. Nach der Natur beschrieben. X Heft. Nürnberg, 100 sheets, 3 pls.
- KÜSTER H. C. 1847b. Die Käfer Europa's. Nach der Natur beschrieben. XII Heft. Nürnberg, 1 + 100 sheets, 2 pls.
- KÜSTER H. C. 1848. Die Käfer Europa's. Nach der Natur beschrieben. XIII Heft. Nürnberg, 100 sheets, 3 pls.
- KÜSTER H. C. 1850. Die Käfer Europa's. Nach der Natur beschrieben. XX Heft. Nürnberg, 100 sheets, 2 pls.
- KÜSTER H. C. 1850b. Die Käfer Europa's. Nach der Natur beschrieben. XXI Heft. Nürnberg, 3 + 4 + 100 sheets, 2 pls.
- KÜSTER H. C. 1851. Die Käfer Europa's. Nach der Natur beschrieben. XXII Heft. Nürnberg, 2 + 100 sheets, 3 pls.
- KÜSTER H. C. 1852. Die Käfer Europa's. Nach der Natur beschrieben. XXV Heft. Nürnberg, 2 + 100 sheets, 2 pls.
- KUGELANN J. G. 1792. Verzeichniss der in einigen Gegenden Preussens bis jetzt entdeckten Käfer-Arten, nebst kurzen Nachrichten von denselben. Fortsetzung. Neuestes Mag. für die Liebhaber der Ent., herausgegeben von D. H. SCHNEIDER. Stralsund, 1, 4 : 477–512.
- KUZNECOVA E. 1828. Ueber eine neue *Dermestes*-Art (*Coleoptera, Dermestidae*). Ent. Nachr. Bl., Troppau, 2 : 17–24.
- KUZNECOVA E. 1933. Materialy k faune kozheev (Dermestidae, Coleoptera) Jakutii. Trudy Zool. Inst. Akad. Nauk, Leningrad, 1 : 229–241.
- LACORDAIRE Th. 1854. Histoire naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. II. Paris, 548 pp.

- LACORDAIRE Th. [1855]. Histoire Naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique des tous les genres proposés jusqu'ici dans cet ordre d'insectes. Atlas. Paris, 28 pp., 80 pls.
- LACORDAIRE Th. 1856. Histoire naturelle des Insectes. Genera des Coléoptères ou exposé méthodique et critique de tous les genres proposés jusqu'ici dans cet ordre d'insectes. III. Paris, 594 pp.
- LAICHARTING J. N. E. 1781. Verzeichniss und Beschreibung der Tyrolier-Insecten. I Theil. Käferartige Insecten. I. Band. Zürich, XII + 248 pp.
- LATREILLE P. A. [1802]. Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes. III. Paris, I-XII + 13-468 pp.
- LATREILLE P. A. [1804]. Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes. IX. Paris, 416 pp., pls. 74-80.
- LATREILLE P. A. 1807. Genera Crustaceorum et Insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata. II. Parisiis et Argentorati, 280 pp.
- LATREILLE P. A. 1825. Familles naturelles du Règne Animal, exposées succinctement et dans un ordre analytique, avec l'indication de leurs genres. Paris, VI + 570 pp.
- LATREILLE [P. A.]. 1829. Crustacés, Arachnides et Partie des Insectes. In: CUVIER, Le Règne Animal distribué d'après son organisation, pour servir de la base à l'Histoire Naturelle des Animaux et d'introduction à l'anatomie comparée. Avec figures dessinées d'après nature. Nouvelle édition, revue et augmentée. IV. Paris, XXVII + 584 pp., 20 pls.
- LEA A. M. 1895. Descriptions of new species of australian Coleoptera. Proc. Linn. Soc. N. S. Wales, Sydney, ser. 2, **10** : 224-319.
- LEA A. M. 1908. The Coleoptera of King Island, Bass Strait. Proc. Roy. Soc. Victoria, Melbourne, n. s., **20** : 143-207.
- LEACH W. E. 1815. Entomology. In: BREWSTER, Edinburgh Encyclopaedia. IX, 1. Edinburgh, pp. 57-172.
- LEACH W. E. 1817. In: Encycl. Brit. [fide AGASSIZ].
- LECONTE J. [E.]. 1824. Description of some new Species of North American Insects. Ann. Lyc. Nat. Hist. N. Y., New York, **1**, 1 : 169-173, pl. 11.
- LECONTE J. L. 1854. Synopsis of the *Dermestidae* of the United States. Proc. Acad. Nat. Sci. Philad., Philadelphia, **7** : 106-113.
- LECONTE J. L. 1859. Catalogue of the Coleoptera of Fort Tejon, California. Proc. Acad. Nat. Sci. Philad., Philadelphia, **1859** : 69-90.
- LECONTE J. L. 1861. Classification of the Coleoptera of North America. Part I. Smithson. Misc. Coll., Washington, xxiv + 214 pp., 6 ff.
- LECONTE J. L. 1862. Notes on the Coleopterous Fauna of Lower California. Proc. Acad. Nat. Sci. Philad., Philadelphia, **1861** : 335-338.
- LECONTE J. L. 1862b. New species of Coleoptera inhabiting the Pacific district of the United States. Proc. Acad. Nat. Sci. Philad., Philadelphia, **1861** : 338-359.
- LECONTE J. L. 1863. New species of North American Coleoptera. Part I. [1]. Smithson. Misc. Coll., Washington, **6**, 167 : 1-86.
- LECONTE J. L. 1866. New species of North American Coleoptera. Part I. [2]. Smithson. Misc. Coll., Washington, **6**, 167 : 87-177.
- LECONTE J. L. 1874. Descriptions of New Coleoptera chiefly from the Pacific Slope of North America. Trans. Amer. Ent. Soc., Philadelphia, **5** : 43-72.
- LECONTE J. L. 1883. Classification of the Coleoptera of North America. Smithson. Misc. Coll., Washington, **26**, 507, xxxviii + 567 pp., 5 ff.
- LEDER H. 1881. Beitrag zur kaukasischen Käfer-Fauna. Unter Mitwirkung von Dr. EPPELSHÉIM in Grünstadt und Edmund REITTER in Wien. III. Stück. Verh. Zool.-Bot. Ges. Wien, **30** : 501-518.

- LEESBERG A. F. A. 1906. Insecten uit Arachidennoten en Sesamzaad, in 1905 gevonden. Ent. Ber. Nederl. Ver., 's-Gravenhage, **2** : 78–81.
- LEPECHIN I. 1772. Prodolzhenye dnevnih zapisok puteshestvija akademika i mediciny doktora Ivana LEPECHINA po raznym provincijam rossijskago gosudarstva v 1770 godu. Sanktpeterburg. Pribavlenie. Opisanie pticam, rybam, nasekomym i presmykajushchimsja. Pp. 293–338, 11 pls.
- LEPESME P. 1939. Note synonymique sur les *Dermestes* (*Col.*) et description d'une espèce et d'une variété nouvelles. Bull. Soc. Ent. France, Paris, **44** : 190–193, 6 ff.
- LEPESME P. 1942. Un nouveau *Megatoma* d'Australie (*Col. Dermestidae*). Bull. Soc. Ent. France, Paris, **46** : 142.
- LEPESME P. 1950. Révision des *Dermestes* (*Col. Dermestidae*). Ann. Soc. Ent. France, Paris, **65** : 37–68, 42 ff.
- LINNAEUS C. 1758. Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima, reformata. I. Holmiae, 824 pp.
- LINNAEUS C. 1761. Fauna Suecica sistens Animalia Sueciae Regni: *Mammalia, Aves, Amphibia, Pisces, Insecta, Vermes*. Distributa per classes et ordines, genera et species, cum differentiis specierum, synonymis auctorum, nominibus incolarum, locis natalium, descriptionibus Insectorum. Editio altera, auctior. Stockholmiae, [48] + 578 pp., 2 pls.
- LINNAEUS C. 1763. Centuria Insectorum quam praeside D. D. Carl von LINNÉ. Proposuit Boas JOHANSSON, Calmariensis. Amoen. Acad., Holmiae, **6** : 384–415.
- LINNAEUS C. 1767. Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Editio decima tertia, ad Editionem duodecimam reformatam Holmiensem. I, 2. Vindobonae, pp. 533–1327 + [36 pp.].
- MACLEAY W. 1871. Notes on a collection of Insects from Gayndah. Trans. Ent. Soc. N. S. Wales, Sydney, **2** : 79–205.
- MACLEAY W. 1872. Notes on a collection of Insects from Gayndah. Second Paper. Trans. Ent. Soc. N. S. Wales, Sydney, **2** : 239–318.
- MADEL W. 1941. Zur Kenntnis der Biologie einiger *Dermestes*-Arten. 2. Beitrag. Beobachtungen über die Entwicklungszeiten der Eier, Larven und Puppen von *Dermestes peruvianus* CAST. Ent. Bl., Krefeld, **37** : 17–21, ff. 5–7.
- MADEL W. 1941b. Zur Kenntnis der Biologie einiger *Dermestes*-Arten. 3. Beitrag. Vergleichende Beschreibung von *Dermestes peruvianus* CAST. und dem wieder neu ein zuführenden *Dermestes haemorrhoidalis* KÜSTER. Ent. Bl., Krefeld, **37** : 107–111, ff. 13–14, pl. II.
- MADER L. 1954. Bemerkungen und Richtigstellungen zu einigen paläarktischen Coleopteren. Ent. Nachr.-Bl., Wien, N. F., **1** : 48–53.
- MANNERHEIM C. G. 1843. Beitrag zur Kaefer-Fauna der Aleutischen Inseln, der Insel Sitkha und Neu-Californiens. Bull. Soc. Imp. Nat. Moscou, Moscou, **16** : 175–314.
- MANNERHEIM C. G. 1853. Dritter Nachtrag zur Kaefer-Fauna der Nord-Amerikanischen Laender des Russischen Reiches. Bull. Soc. Imp. Nat. Moscou, Moscou, **26**, II : 95–273.
- MARCU O. 1957. Contribuții la cunoașterea faunei Coleopterelor Transilvaniei. Bul. Univ. Cluj, St. Nat., Cluj, **2** : 527–544.
- MARSHAM Th. 1802. Entomologia Britannica, sistens *Insecta* Britanniae indigena, secundum methodum Linnaeanam disposita. Tomus I. Coleoptera. Londini, xxxi + 548 pp.
- MATSUMURA S. et YOKOYAMA K. 1928. An interesting Dermested beetle from Japan. Ins. Mats., Sapporo, **2** : 130–132.
- MATSUMURA S. et YOKOYAMA K. 1928b. New and hitherto unrecorded species of *Dermestidae* from Japan. Ins. Mats., Sapporo, **3** : 51–54.
- MEEUSE A. D. J. 1950. The Australian Carpet Beetle, *Anthrenocerus australis* (HOPE) (*Coleoptera, Dermestidae—Anthrenini*) in the Netherlands. Ent. Ber. Ned. Ent. Soc., Amsterdam, **13** : 161–162, 3 fot.

- MEGERLE J. K. 1805. Abhandlung über die dem Getreide und Weinstocke im Erzherzogthum Oesterreich vorzüglich nachtheiligen Thiere. Wien. 71 pp., 1 pl.
- MEIER W. 1899. Über Abänderungen einiger Coleopteren-Arten, welche bei Hamburg gefunden wurden. Ent. Nachr., Berlin, **25** : 97–102.
- MELSHEIMER F. V. 1806. Catalogue of Insects of Pennsylvania. Part I. Hanover, York County, VI + 60 pp.
- MELSHEIMER F. E. 1844. Descriptions of New Species of *Coleoptera* of the United States. Proc. Acad. Nat. Sci. Philad., Philadelphia, **2** : 98–118.
- MELSHEIMER F. E. 1853. Catalogue of the described *Coleoptera* of the United States. Smithson. Misc. Coll., Washington, xvi + 174 pp.
- MILLIRON H. E. 1939. A parthenogenetic new species of the genus *Perimegatoma* HORN (*Coleoptera: Dermestidae*). Ann. Ent. Soc. America, Washington, **32** : 570–574, 3 ff.
- MOLL K. E. 1784. Verzeichniss der Salzburger Insekten. IIIte Lieferung. Neues Mag. Ent. Füssli, Winterthur, **2** : 169–197.
- MONTROUZIER P. 1860. Essai sur la Faune Entomologique de la Nouvelle-Calédonie (Balade) et Des îles des Pins, Art, Lifu, etc. Ann. Soc. Ent. France, Paris, ser. 3, **8** : 229–308, 1 pl.
- [MOTSCHULSKY V. =] Victor, 1839. Coléoptères du Caucase et des provinces Transcaucasien. (Continuation). Bull. Soc. Imp. Nat. Moscou, Moscou, **1839** : 68–93, pls. 5–6.
- MOTSCHULSKY V. 1849. Coléoptères reçus d'un Voyage de M. HANDSCHUH dans le Midi de l'Espagne, énumérés et suivis de notes. Bull. Soc. Imp. Nat. Moscou, Moscou, **22**, II : 52–153.
- MOTSCHULSKY V. 1850. Kritische Beurtheilung von Dr. ERICHSON's Naturgeschichte der Insekten Deutschlands. (Fortsetzung). Bull. Soc. Imp. Nat. Moscou, Moscou, **23**, I : 195–257.
- MOTSCHULSKY V. 1858. Insectes des Indes orientales. Étud. Ent., Helsingfors, **7** : 20–122.
- MOTSCHULSKY V. 1858b. Sur les Collections Coléoptérologiques de LINNÉ et de FABRICIUS. (Continuation). Étud. Ent., Helsingfors, **7** : 123–152.
- MOTSCHULSKY V. 1859. Catalogue des insectes rapportés des environs du fl. Amour, depuis la Schilka jusqu'à Nikolaïevsk, examinés et énumérés. Bull. Soc. Imp. Nat. Moscou, Moscou, **32**, II : 487–507.
- MOTSCHULSKY V. 1860. Coléoptères rapportés de la Sibérie orientale et notamment des pays situés sur les bords du fleuve Amour par MM. SCHRENCK, MAACK, DITMAR, VOZNESSENSKI etc. In: SCHRENK L. Reisen und Forschungen im Amur-Lande in den Jahren 1854–1856 im Auftrage der Keiserl. Akademie der Wissenschaften zu St. Petersburg ausgeführt und in Verbindung mit mehreren Gelehrten herausgegeben. Band II. Zweite Lieferung. Coleopteren. St. Petersburg, 258 pp., pls. VI–XI.
- MOTSCHULSKY V. 1863. Essai d'un catalogue des insectes de l'île Ceylan. (Suite). Bull. Soc. Imp. Nat. Moscou, Moscou, **36**, I : 421–532.
- MOTSCHULSKY V. 1866. Catalogue des insectes reçus du Japon. Bull. Soc. Imp. Nat. Moscou, Moscou, **39**–I : 163–200.
- MROCKOWSKI M. 1950. A new species of *Anthrenus* GEOFFR. from Poland (*Coleoptera, Dermestidae*). Ann. Mus. Zool. Pol., Warszawa, **14** : 187–192, pl. XVII.
- MROCKOWSKI M. 1951. *Anthrenus (Helocerus) polonicus* sp. n. from Central and Eastern Europe with a description of a new subgenus of *Anthrenus* GEOFFR. Ann. Mus. Zool. Pol., Warszawa, **14** : 253–260, 2 ff., pl. XXXV.
- MROCKOWSKI M. 1952. Contribution to the knowledge of the *Dermestidae* with description of a new species and a new subspecies (*Coleoptera*). Ann. Mus. Zool. Pol., Warszawa, **15** : 25–32, pl. IV.
- MROCKOWSKI M. 1954. Contribution to the knowledge of *Dermestidae* (*Coleoptera*). *Anthrenus flavidus* SOLSKI. Ann. Zool., Warszawa, **16** : 1–8, 17 ff.
- MROCKOWSKI M. 1954b. *Dermestidae* okolic Warszawy (*Coleoptera*). Fragm. Faun., Warszawa, **7** : 187–197.

- MROCKOWSKI M. 1954c. Skórnikи—*Dermestidae*. In: "Klucze do oznaczania owadów Polski", XIX, 52. Warszawa, 47 pp., 81 ff.
- MROCKOWSKI M. 1956. *Trogoderma angustum* (SOL.) w Europie (*Coleoptera, Dermestidae*). Pol. Pismo Ent., Wrocław, 24, suppl. 1 : 29–31.
- MROCKOWSKI M. 1958. New names in the family *Dermestidae* (*Coleoptera*). Acta Zool. Cracov., Kraków, 2 : 557–561.
- MROCKOWSKI M. 1958b. Notes on the genus *Hemirhopalum* SHARP, with a description of a new species from Brazil (*Coleoptera, Dermestidae*). Ann. Zool., Warszawa, 17 : 49–64, 16 ff.
- MROCKOWSKI M. 1958c. *Dermestidae* (*Coleoptera*) collected in Bulgaria by members of the staff of the Zoological Institute of the Polish Academy of Sciences in the years 1950 and 1956. Fragm. Faun., Warszawa, 8 : 1–9, 2 ff.
- MROCKOWSKI M. 1959. Zweiter Beitrag zur Kenntnis der Dermestiden von Afghanistan, nebst Beschreibung einer neuen Art (*Coleoptera*). Föhr. Fysiogr. Sällsk. Lund, Lund, 29 : 99–101.
- MROCKOWSKI M. 1960. Dritter Beitrag zur Kenntnis der Dermestiden von Afghanistan (*Coleoptera*). Föhr. Fysiogr. Sällsk. Lund, Lund, 30 : 51–55, 5 ff.
- MROCKOWSKI M. 1960b. Kozheedy (*Coleoptera, Dermestidae*) Turkmenii. Trudy Zool. Inst. Akad. Nauk, Moskwa-Leningrad, 27 : 208–219, 19 ff.
- MROCKOWSKI M. 1960c. Data on the distribution of *Dermestidae* (*Coleoptera*) with special reference to the Hungarian fauna. Ann. Hist.-Nat. Mus. Hung., Zool., Budapest, 52 : 247–253.
- MROCKOWSKI M. 1960d. *Trogoderma angustum* SOL. in Deutschland (Col., *Dermestidae*). Mitt. Dtsch. Ent. Ges., Berlin, 19 : 95–96.
- MROCKOWSKI M. 1961. Two New Species of the Genus *Anthrenus* GEOFFR. from the Tadzhik S. S. R. (*Coleoptera, Dermestidae*). Bull. Acad. Pol. Sci., Cl. II, Warszawa, 8 : 591–593, 1 pl.
- MROCKOWSKI M. 1961b. Ergebnisse der Deutschen Afghanistan-Expedition 1956 der Landessammlungen für Naturkunde Karlsruhe. *Dermestidae* (*Coleoptera*). Beitr. Naturk. Forsch. SW-Deutschl., Karlsruhe, 19 : 223–226, 4 ff.
- MROCKOWSKI M. 1961c. Vierter Beitrag zur Kenntnis der Dermestiden von Afghanistan, nebst Beschreibung von zwei neuen Arten der Gattung *Ctesias* STEPH. (*Coleoptera*). Ent. Tidskr., Stockholm, 82 : 191–196, 9 ff.
- MROCKOWSKI M. 1962. *Dermestidae* (*Coleoptera*) from the Kazakh SSR. Ann. Zool., Warszawa, 20 : 229–259, 18 ff., 1 map.
- MROCKOWSKI M. 1962b. *Anthrenus minutus* ER. and Faunistic Notes on other Sardinian *Dermestidae* (*Coleoptera*). Studi Sassarensi, Sez. III, Sassari, 9 : 544–549, 1 map.
- MROCKOWSKI M. 1963. Some remarks on the *Dermestidae* (*Coleoptera*) from the Zoological Museum in Hamburg. Ent. Mitt. Zool. Mus. Hamburg, Hamburg, 2 : 377–380, 3 ff.
- MROCKOWSKI M. 1963b. *Anthrenus sarnicus* sp. n. (*Coleoptera, Dermestidae*) from the Island of Guernsey. Ann. Mag. Nat. Hist., London, ser. 13, 5 : 697–700, 9 ff.
- MROCKOWSKI M. 1964. Systematic and Synonymic Notes Upon Certain Species of *Dermestidae* (*Coleoptera*). Ann. Zool., Warszawa, 22 : 179–187, 26 ff.
- MROCKOWSKI M. 1965. Ergebnisse der Zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei. 11. *Silphidae* partim, *Dermestidae* (*Coleoptera*). Fol. Ent. Hung., Budapest, s. n., 17 : 183–185.
- MROCKOWSKI M. 1965b. 55. *Silpha* LIN. und *Dermestidae* II. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, Dresden, 7 : 103–105.
- MROCKOWSKI M. 1966. *Silphidae* and *Dermestidae* (*Coleoptera*) collected in Mongolia by Polish Zoolologists in the years 1959–1964. Fragm. Faun., Warszawa, 12 : 333–338.
- MROCKOWSKI M. 1966b. *Silphidae*, *Catopidae* and *Dermestidae* of the Noona Dan Expedition to the Philippine and Bismarck Islands. Ent. Medd., København, 34 : 325–328, 5 ff.

- MROCKOWSKI M. 1966c. 79. *Silpha* und *Dermestidae* der III. Expedition. Ergebnisse der zoologischen Forschungen von Dr. Z. KASZAB in der Mongolei (*Coleoptera*). Reichenbachia, Dresden, 7 : 267-269.
- MROCKOWSKI M. 1966 d. Contribution to the Knowledge of *Silphidae* and *Dermestidae* of Korea (*Coleoptera*). Ann. Zool., Warszawa, 23 : 433-443, 11 ff.
- MROCKOWSKI M. 1967. The Palearctic Species of *Megatoma* HERBST (*Coleoptera, Dermestidae*). Pol. Pismo Ent., Wroclaw, 37 : 3-24, 28 ff.
- MÜLLER O. F. 1764. Fauna insectorum Fridrichsdalina, sive methodica descriptio insectorum agra Fridrichsdalensis, cum characteribus genericis et specificis, nominibus trivialibus, locis natalibus, iconibus allegatis, novisque pluribus speciebus additis. Hafniae et Lipsiae, XXIV + 96 pp.
- MÜLLER O. F. 1776. Zoologiae Danicae prodromus, seu Animalium Daniae et Norvegiae indigenarum characteres, nomina, et synonyma imprimis popularium. Havniae, XXXII + 274 pp.
- MULSANT E. 1870. Opuscules Entomologiques. 14 cahier. Paris, 241 pp.
- MULSANT E. et GODART [A.]. 1855. Description de quelques espèces de Coléoptères nouveaux ou peu connus. Ann. Soc. Linn. Lyon, n. s., 2 : 261-283.
- MULSANT [E.] et REY [C.]. 1868. Tribu des Scuticolles. Ann. Soc. Linn. Lyon, n. s., 15 : 1-188, 3 pls.
- MURRAY A. 1867. List of *Coleoptera* received from Old Calabar, on the West Coast of Africa. Ann. Mag. Nat. Hist., London, ser. 3, 20 : 20-23.
- MUTCHLER J. et WEISS H. B. 1927. The dermestid beetles of New Jersey including the carpet beetles and those which feed on animal products. Circ. N. J. Dept. Agric., Trenton, N. J., 108, 31 pp., 4 pls.
- NEWMAN E. 1834. Attempted Division of British Insects into Natural Orders. Ent. Mag., London, 2 : 379-431.
- NORMAND H. 1936. Contribution au Catalogue des Coléoptères de la Tunisie (7e fascicule). Bull. Soc. Hist. Nat. Alger, Alger, 26 : 286-304, 3 ff.
- NORMAND H. 1950. Contribution au catalogue des Coléoptères de la Tunisie (Troisième supplément-fascicule 3). Bull. Soc. Sci. Nat. Tunis, Tunis, 2 : 65-104, pls. VI-VII.
- OBENBERGER J. 1917. II. Beitrag zur Kenntnis der palaearktischen Käfer-fauna. Arch. Naturges., Berlin, 82A, 4 : 9-45.
- OLIVIER [A. G.]. 1789. Insectes. In: Encyclopédie méthodique. Histoire Naturelle. Tome IV. Paris et Liège, CCCLXXIII + 331 pp.
- OLIVIER [A. G.]. 1790. Entomologie, ou histoire naturelle des Insectes, Avec leurs caractères génériques et spécifiques, leur description, leur synonymie, et leur figure enluminée. Coléoptères. Tome Second. No. 9. Dermeste. 16 pp., No. 14. Anthrène. 10 pp. Paris.
- OLLIFF A. S. 1883. Remarks on a small collection of Clavicorn *Coleoptera* from Borneo, with descriptions of new species. Trans. Ent. Soc. London, London, 1883 : 173-186.
- PALM Th. 1940. Systematische Bemerkungen zu den schwedischen *Anthrenus*-Arten (*Col. Dermestidae*). Ent. Tidskr., Stockholm, 61 : 1-3, 3 ff.
- PANZER G. W. F. 1789. Einige seltene Insecten. Naturf., Halle, 24 : 1-35, 1 pl.
- PANZER G. W. F. 1793. Faunae Insectorum Germanicae initia oder Deutschland Insecten. Erster Iahrgang. I-XII. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].
- PANZER G. W. F. 1794. Faunae Insectorum Germanicae initia oder Deutschland Insecten. Zweyter Iahrgang. XIII-XXIV. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].
- PANZER G. W. F. [1795]. Entomologia Germanica exhibens Insecta per Germaniam indigena secundum classes, ordines, genera, species Adiectis synonymis, locis, observationibus. I. Eleuterata. Norimberga, 36 + 372 pp., 12 pls.
- PANZER G. W. F. 1796. Faunae Insectorum Germanicae initia oder Deutschland Insecten. Zweyte Auflage. Erster Iahrgang. I-XII. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].

- PANZER G. W. F. 1796b. Faunae Insectorum Germanicae initia oder Deutschlands Insecten. Dritter Jahrgang. XXV–XXXVI. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].
- PANZER G. W. F. 1797. Faunae Insectorum Germanicae initia oder Deutschland Insecten. Vierter Jahrgang. XXXVII–XLVIII. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].
- PANZER G. W. F. 1798. Faunae Insectorum Germanicae initia oder Deutschlands Insecten. Fünfter Jahrgang. XLIX–LX. Heft. Nürnberg. [24 sheets and 24 pls. in each one volume].
- PASCOE F. P. 1866. Notices of new or little-known genera and species of *Coleoptera*. J. Ent., London, 2 : 443–493, pls. XVIII–XIX.
- PAULIAN R. 1953. Recherches sur les insectes d'importance biologique à Madagascar XII à XX. Mém. Inst. Sci. Madagascar, Paris, sér. E, 3 : 1–27, 30 ff.
- PAYKULL G. [1798]. Fauna Suecica. *Insecta*. Tomus I. Upsaliae, 10 + 358 + 2 pp.
- PAYKULL G. 1800. Fauna Suecica. *Insecta*. Tomus III. Upsaliae, 459 pp.
- PÉRINQUEY L. 1885. First contribution to the South-African Coleopterous Fauna. Trans. S. Afr. Phil. Soc., Cape Town, 3 : 75–150, 4 pls.
- PÉRINQUEY L. 1888. Second contribution to the South-African Coleopterous fauna. Trans. S. Afr. Phil. Soc., Cape Town, 4 : 67–194, 4 pls.
- PERRIS É. 1853. Histoire des Insectes du Pin Maritime. Suite. Ann. Soc. Ent. France, Paris, ser. 3, 1 : 555–644, pls. 17–19.
- PERRIS É. 1862. Histoire des insectes du Pin Maritime. Supplément aux Coléoptères et rectifications. Ann. Soc. Ent. France, Paris, ser. 4, 2 : 173–243, pls. 5–6.
- PERRIS É. 1866. Descriptions de quelques Insectes nouveaux. Ann. Soc. Ent. France, Paris, ser. 4, 6 : 181–196.
- PEYERIMHOFF P. 1924. Nouveaux Coléoptères du Nord-Africain. Quarante sixième note: Récoltes de M. C. DUMONT dans la région d'El-Goléa. Bull. Soc. Ent. France, Paris, 1924 : 62–67.
- PEYERIMHOFF P. 1928. Nouveaux Coléoptères du Nord-Africain. Soixantième note: Récoltes de M. le lieutenant CALLIES dans la région du Mzab. Bull. Soc. Ent. France, Paris, 1927 : 304–306, 1 f.
- PEYERIMHOFF P. 1931. Mission Scientifique du Hoggar envoyée de Février à Mai 1928 par M. Pierre BORDES Gouverneur Général de l'Algérie. Coléoptères. Mém. Soc. Hist. Nat. Afr. N., Alger, 2, 172 pp., 3 pls., 18 ff., 2 maps.
- PEYERIMHOFF P. 1943. Matériaux pour un Catalogue des Coléoptères sahariens. II. Descriptions d'espèces nouvelles. Bull. Soc. Hist. Nat. Afr. N., Alger, 34 : 7–35, 9 ff.
- PEYRON E. 1857. Description de quelques Coléoptères nouveaux et observations diverses. Ann. Soc. Ent. France, Paris, ser. 3, 5 : 715–723.
- PHILIPPI R. A. et PHILIPPI F. 1864. Beschreibung einiger neuen Chilenischen Käfer. Ent. Ztg., Stettin, 25 : 266–284.
- PIC M. 1893. Notes sur quelques Coléoptères, avec descriptions. Échange, Lyon, 9 : 122.
- PIC M. 1894. Excursion entomologique à Tougourt (Fin Avril 1893). Échange, Lyon, 10 : 14–21.
- PIC M. 1894b. Descriptions et notes entomologiques (Coléoptères). Échange, Lyon, 10 : 66.
- PIC M. 1894c. Notes entomologiques. Échange, Lyon, 10 : 116.
- PIC M. 1894d. Description de trois Coléoptères nouveaux du groupe des Dermestides. Échange, Lyon, 10 : 119.
- PIC M. 1894e. Descriptions d'espèces et variétés de Coléoptères asiatiques. Échange, Lyon, 10 : 142–144.
- PIC M. 1894f. Description de Coléoptères de la faune circa-européenne. Feuille Jeun. Natural., Paris, 24 : 60–61.
- PIC M. 1894g. Descriptions de Coléoptères du nord de l'Afrique. Le Naturaliste, Paris, 16 : 71.
- PIC M. 1894h. Contribution à la faune Coléoptérologique d'Algérie. Le Naturaliste, Paris, 16 : 180, 210–211.

- PIC M. 1895. Descriptions de Coléoptères d'Algérie. Échange, Lyon, **11** : 78–82.
- PIC M. 1895b. Notes complémentaires ou observations diverses à propos d'espèces et variétés omises ou cataloguées. Échange, Lyon, **11** : 110–120.
- PIC M. 1898. Variétés et nouvelles espèces de Coléoptères. Misc. Ent., Narbonne, **6** : 2–4.
- PIC M. 1898b. Notes descriptives sur plusieurs Coléoptères et sur un *Ichneumon* (Hyménoptère). Misc. Ent., Narbonne, **6** : 73–75.
- PIC M. 1899. Description d'un genre nouveau de Coléoptère d'Algérie. Bull. Soc. Ent. France, Paris, **1899** : 28–29.
- PIC M. 1899b. Diagnoses de Coléoptères communiqués par M. L. BLEUSE. Bull. Soc. Ent. France, Paris, **1899** : 85–86.
- PIC M. 1899c. Diagnoses de Coléoptères d'Orient. Deuxième Article. Bull. Soc. Hist. Nat. Autun, Autun et Paris, **12**, 2 : 252–261.
- PIC M. 1900. Descriptions de Coléoptères Algériens et Tunisiens. Échange, Moulins, **16** : 53–56.
- PIC M. 1901. Notes diverses et diagnoses (5e article). Échange, Moulins, **17** : 33–35.
- PIC M. 1902. Notes de chasses. Échange, Moulins, **18**, No. 209, third page of cover.
- PIC M. 1902b. Descriptions et notes diverses (3e article). Échange, Moulins, **18** : 47–49.
- PIC M. 1903. Descriptions et habitats de divers Coléoptères du Nord de l'Afrique. Échange, Moulins, **19** : 113–115.
- PIC M. 1903b. Notes entomologiques et descriptions. Échange, Moulins, **19** : 121–125.
- PIC M. 1904. Notes et Descriptions. Échange, Moulins, **20** : 2–5.
- PIC M. 1904b. Diagnoses de Coléoptères paléarctiques et exotiques. Échange, Moulins, **20** : 33–36.
- PIC M. 1904c. Quelques chasses faites avant et après le coucher du soleil. Échange, Moulins, **20** : 60–61.
- PIC M. 1904d. Coléoptères nouveaux de la Turquie d'Asie. Échange, Moulins, **20** : 73–74.
- PIC M. 1906. Descriptions de Coléoptères nouveaux de la République Argentine. Rev. Mus. La Plata, Buenos Aires, **12** : 233–235.
- PIC M. 1906b. Nouveaux Coléoptères d'Europe, Asie, Afrique et Amérique (Suite). Échange, Moulins, **22** : 9–12.
- PIC M. 1907. Notes entomologiques diverses. Échange, Moulins, **23** : 153–154.
- PIC M. 1907b. Notes de Chasses. Échange, Moulins, **23**, No. 275, third page of cover.
- PIC M. 1907c. Nouveaux Coléoptères recueillis en Algérie par M. P. de PEYERIMHOFF. Bull. Soc. Ent. France, Paris, **1907** : 274–275.
- PIC M. 1908. Correspondance. Bull. Soc. Ent. Égypte, Le Caire, **1** : 75–76.
- PIC M. 1908b. Descriptions ou diagnoses et notes diverses. Suite. Échange, Moulins, **24** : 41–46.
- PIC M. 1908c. Notes de chasse. Échange, Moulins, **24**, No. 286, third page of cover.
- PIC M. 1908d. Descriptions ou diagnoses et notes diverses. Suite, Échange, Moulins, **24** : 81–83.
- PIC M. 1910. Descriptions ou diagnoses et notes diverses. Suite. Échange, Moulins, **26** : 17–19.
- PIC M. 1912. Captures de quelques Coléoptères dans les Alpes françaises. Échange, Moulins, **28** : 74–75.
- PIC M. 1912b. Notes sur quelques insectes nouveaux ou peu connus, mutations et synonymies. Mélang. Exot.-Ent., Saint-Amand, **2** : 14–16.
- PIC M. 1914. Notes diverses, descriptions et diagnoses. Suite. Échange, Moulins, **30** : 10–13.
- PIC M. 1915. Nouveautés de diverses familles. Mélang. Exot.-Ent., Moulins, **13** : 2–13.
- PIC M. 1915b. Nouvelles espèces de diverses familles. Mélang. Exot.-Ent., Moulins, **15** : 2–14.
- PIC M. 1916. Coléoptères exotiques en partie nouveaux (Suite). Échange, Moulins, **32** : 3–4.
- PIC M. 1916b. Notes et descriptions abrégées diverses. Mélang. Exot.-Ent., Moulins, **17** : 2–8.
- PIC M. 1916c. Diagnoses Spécifiques. Mélang. Exot.-Ent., Moulins, **17** : 8–20.
- PIC M. 1916d. Diagnoses génériques et spécifiques. Mélang. Exot.-Ent., Moulins, **18** : 2–20.

- PIC M. 1918. Cabinet d'élevage, notes biologiques et captures. Échange, Moulins, **34**, Hors-Texte: 1-4.
- PIC M. 1918b. Courtes descriptions diverses. Mélang. Exot.-Ent., Moulins, **27**: 1-24.
- PIC M. 1919. Notes diverses, descriptions et diagnoses (Suite). Échange, Moulins, **35**: 17-20.
- PIC M. 1921. Nouveaux Coléoptères. Bull. Soc. Ent. Égypte, Le Caire, **6**: 54-55.
- PIC M. 1921b. Rhipidocerides, Helodides, Malacodermes divers, Dermestides, Bruchides, Monommides, Hétéromères (ex parte), Hilophilides et Anthicides. In: Voyage de M. Guy BABAULT en Afrique Orientale (1912-1913). Résultats scientifiques. Insectes Coléoptères. Paris, 32 pp., 2 fot.
- PIC M. 1922. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **36**: 1-32.
- PIC M. 1923. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **40**: 3-32.
- PIC M. 1924. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **41**: 1-32.
- PIC M. 1926. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **45**: 1-32.
- PIC M. 1927. Fauna sumatrensis. (Beitrag Nr. 34). *Byrrhidae, Dermestidae, Anobiidae (Col.)*. Suppl. Ent. Berlin-Dahlem, **15**: 65-66.
- PIC M. 1927b. Notes diverses, descriptions et diagnoses (Suite). Échange, Moulins, **43**: 5-7.
- PIC M. 1927c. Nouveaux Coléoptères exotiques. Ent. Mitt., Berlin-Dahlem, **16**: 246-255.
- PIC M. 1927d. Nouveaux Coléoptères du globe (IV). Bull. Soc. Zool. France, Paris, **52**: 185-189.
- PIC M. 1927e. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **48**: 1-32.
- PIC M. 1928. Nouveaux Coléoptères de la République Argentine. Rev. Soc. Ent. Argentina, Buenos Aires, **2**: 49-52.
- PIC M. 1928b. Notes et Descriptions. Mélang. Exot.-Ent., Moulins, **51**: 1-36.
- PIC M. 1928c. Nouveaux Coléoptères exotiques. Bull. Mus. Hist. Nat., Paris, **34**: 257-258.
- PIC M. 1929. Notes diverses, nouveautés. Échange, Moulins, **45**: 9-10.
- PIC M. 1929b. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **54**: 1-36.
- PIC M. 1931. Notes diverses, nouveautés. Échange, Moulins, **47**: 5-6.
- PIC M. 1931b. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **57**: 1-36.
- PIC M. 1933. Neue Coleopteren-Clavicornia. Ent. Nachr.Bl., Troppau, **7**: 71-72.
- PIC M. 1934. Notes diverses, nouveautés (Suite). Échange, Moulins, **50**: 25-27.
- PIC M. 1934b. Notes diverses, nouveautés (Suite). Échange, Moulins, **50**: 29-31.
- PIC M. 1935. Nouveaux Coléoptères de l'Île Maurice. Bull. Soc. Ent. France, Paris, **40**: 12-14.
- PIC M. 1935b. Excursions dans le massif de la Chartreuse (suite et fin). Échange, Moulins, **51**, Hors-texte: 13-16.
- PIC M. 1935c. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **66**: 1-36.
- PIC M. 1935d. Schwedisch-chinesische wissenschaftliche Expedition nach den nordwestlichen Provinzen Chinas, unter Leitung von Dr. Sven HEDIN und Prof. Sü PING-CHENG. Insecten gesammelt von schwedischen Arzt der Expedition Dr. David HUMMEL 1927-1930. 16. Coleoptera. 2. Helmidae, Dermestidae, Anobiidae, Cleridae Malacodermata, Dascillidae, Heteromera (ex p.), Bruchidae, Cerambycidae, Phytophaga (ex p.). Ark. Zool., Stockholm, **27A**, No. 2, 14 pp.
- PIC M. 1936. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **67**: 1-36.
- PIC M. 1937. Trois nouveaux Coléoptères exotiques. Ann. Mag. Nat. Hist., London, ser. 10, **20**: 409-411.
- PIC M. 1937b. Coléoptères exotiques en partie nouveaux (Suite). Échange, Moulins, **53**: 4.
- PIC M. 1937c. Nouveautés diverses. Mélang. Exot.-Ent., Moulins, **69**: 1-36.
- PIC M. 1937d. Nouveaux Coléoptères du Musée zoologique de Hamburg. Ent. Nachr.Bl., Troppau, **10**: 193-196.
- PIC M. 1938. Notes diverses, nouveautés (Suite). Échange, Moulins, **54**: 1-3.
- PIC M. 1938b. Notes diverses, nouveautés (Suite). Échange, Moulins, **54**: 9-11.

- PIC M. 1938c. Coléoptères exotiques en partie nouveaux (Suite). Échange, Moulins, **54** : 12.
- PIC M. 1938d. Nouveaux Coléoptères du Siam. Bull. Soc. Zool. France, Paris, **63** : 352–356.
- PIC M. 1939. Coléoptères nouveaux d'Egypte et du Sinaï. Bull. Soc. Fouad I Ent., Le Caire, **23** : 143–149.
- PIC M. 1940. Diagnoses de Coléoptères exotiques (Suite). Échange, Moulins, **56** : 10–12.
- PIC M. 1942. Opuscula martialia VII. Échange, Moulins, Numéro spécial, 16 pp.
- PIC M. 1942b. Opuscula martialia VIII. Échange, Moulins, Numéro spécial, 16 pp.
- PIC M. 1943. Coléoptères du globe (suite). Échange, Moulins, **59** : 13–16.
- PIC M. 1946. Coléoptères du globe (suite). Échange, Moulins, **62** : 5–8.
- PIC M. 1948. Notes et nouveautés diverses. Divers. Ent., Moulins, **4** : 12–16.
- PIC M. 1950. Coléoptères nouveaux du Congo de diverses familles. Rev. Zool. Bot. Afr., Tervuren, **43** : 356–362.
- PIC M. 1951. Coléoptères du globe (suite). Échange, Moulins, **67** : 5–8.
- PIC M. 1951b. Coléoptères du globe (suite). Échange, Moulins, **67** : 13–16.
- PIC M. 1951c. Descriptions diverses et notes. Divers. Ent., Moulins, **8** : 7–16.
- PIC M. 1952. Coléoptères nouveaux du Congo Belge. Ann. Mus. Congo Belg. (ser. 8^o), Sci. Zool., Tervuren, **12** : 1–47.
- PIC M. 1952b. Missione Biologica Sagan-Omo diretta dal prof. Edoardo ZAVATTARI. Coléoptères nouveaux. Riv. Biol. Colon., Roma, **11** : 25–36.
- PIC M. 1952c. Coléoptères du globe (suite). Échange, Moulins, **68** : 1–4.
- PIC M. 1953. Coléoptères nouveaux de l'Afrique orientale. Atti Mus. Stor. Nat. Trieste, Bolognà, **19** : 110–114.
- PIC M. 1954. Coléoptères nouveaux de Chine. Bull. Soc. Ent. Mulhouse, Mulhouse, **1954** : 53–59.
- PIC M. 1954b. Coléoptères du globe (suite). Échange, Moulins, **70** : 9–12.
- PIC M. 1954c. *Ptinidae, Anobiidae, Dermestidae, Byrrhidae* (Col.). Beitr. zur Fauna Perus, Jena, **4** : 176–178.
- PIC M. 1956. Nouveaux Coléoptères de diverses familles. Ann. Hist.-Nat. Mus. Hung., Budapest, s. n., **7** : 71–92.
- PILLER M. et MITTERPACHER L. 1783. Iter per Poseganam Sclavoniae provinciam mensibus Junio, Julio Anno MDCCCLXXXII susceptum. Budae, 147 pp., 16 pls.
- PLAVILSTSHIKOV N. N. 1924. Analecta Coleopterologica. Ann. Mag. Nat. Hist., London, ser. 9, **13** : 230–233.
- POOL C. J. C. 1915. Notes from the Caird Insect House. Proc. Zool. Soc. London, London, **1915** : 709–712.
- PRIESNER H. 1951. Description of the Common Egyptian *Trogoderma* (*Coleoptera: Dermestidae*). Bull. Soc. Fouad I Ent., Le Caire, **35** : 133–135, 3 ff.
- QUENSEL C. 1790. Dissertatio historico-naturalis ignotas Insectorum species continens. Respondens Carolus P. LUNDGÅRD. Lundae, 20 pp.
- RAGUSA E. 1926. Coleotteri nuovi o poco conosciuti della Sicilia. Naturalista Sicil., Palermo, **24** : 69–83.
- REDTENBACHER L. 1843. Bemerkungen über die in Syrien von Theodor KOTSCHY gesammelten Käfer [and] Coleopterorum Syriae genera et species novae. In: RUSSEGGER J. Reisen in Europa, Asien und Afrika, mit besonderer Rücksicht auf die naturwissenschaftlichen Verhältnisse der betreffenden Länder, unternommen in den Jahren 1835 bis 1841. Erster Band. Reise in Griechenland, Unteregypten, im nördlichen Syrien und südöstlichen Kleinasiens, mit besonderer Rücksicht auf die naturwissenschaftlichen Verhältnisse der betreffenden Länder, unternommen in dem Jahre 1836. Zweiter Theil. Stuttgart, pp. 973–990.

- REDTENBACHER L. 1845. Die Gattungen der deutschen Kaefer-Fauna nach der analytischen Methode bearbeitet, nebst einem kurz gefassten Leitfaden, zum Studium dieses Zweiges der Entomologie. Wien, 14 + 178 pp., 2 pls.
- REDTENBACHER L. 1867. Coleopteren in Reise der Österreichischen fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den befehlen des commodore B. von WÜLLERSTORF-URBAIR. Zoologischer Theil. Zweiter Band. Wien, IV + 249 pp., 5 pls.
- REITTER E. 1875. Revision der europäischen *Lathridiidae*. Stettin. Ent. Ztg., Stettin, **36** : 297–340.
- REITTER E. 1875b. Revision der europäischen *Lathridiidae*. Stettin. Ent. Ztg., Stettin, **36** : 410–445.
- REITTER E. 1875c. Ueber eine neue Gattung der *Thorictidae*. Coleopt. Hefte, München, **15** : 45–46.
- REITTER E. 1877. Beiträge zur Käferfauna von Japan. (Drittes Stück.). Dtsch. Ent. Z., Berlin, **21** : 369–383.
- REITTER E. 1878. *Hadrotoma quadriguttata* n. sp. Dtsch. Ent. Z., Berlin, **22** : 95.
- REITTER E. 1878b. *Dermestidae*. In: SCHNEIDER O. et LEDER H., Beiträge zur Kenntniss der kaukasischen Käferfauna. Verh. Nat. Ver. Brünn, Brünn, **16** : 3–258.
- REITTER E. 1879. Verzeichniss der von H. CHRISTOPH in Ost-Sibirien gesammelten Clavicornier etc. Dtsch. Ent. Z., Berlin, **23** : 209–226.
- REITTER E. 1880. Coleopterologische Ergebnisse einer Reise nach Croatiens und Slavonien. Verh. Zool.-Bot. Ges. Wien, Wien, **29** : 35–56.
- REITTER E. 1880b. Beitrag zur Synonymie der Coleopteren. Verh. Zool.-Bot. Ges. Wien, Wien, **29** : 507–512.
- REITTER E. 1881. Die aussereuropäischen Dermestiden meiner Sammlung. Mit 70 Diagnosen neuer Arten. Verh. Nat. Ver. Brünn, Brünn, **19** : 27–60.
- REITTER E. 1881b. Bestimmungs-Tabellen der europäischen Coleopteren. III. Enthaltend die Familien: *Scaphidiidae*, *Lathridiidae* und *Dermestidae*. Verh. Zool.-Bot. Ges. Wien, Wien, **30** : 41–94.
- REITTER [E.]. 1881c. Eine neue *Trogoderma* aus Neuholland. Dtsch. Ent. Z., Berlin, **25** : 232.
- REITTER E. 1883. *Dermestidae*. In: Catalogus Coleopterorum Europae et Caucasi. Auctoribus Dr. L. v. HEYDEN, E. REITTER et J. WEISE. Editio tertia. Londini, Berolini, Parisiis, II + 228 pp.
- REITTER E. 1884. Diagnosen neuer Coleopteren aus Lenkoran. Verh. Nat. Ver. Brünn, Brünn, **22** : 3–10.
- REITTER E. 1887. Insecta in itinere Cl. N. PRZEWALSKII in Asia Centrali novissime lecta. VI. *Clavicornia*. *Lamellicornia* et *Serricornia*. Horae Soc. Ent. Ross., Petropoli, **21** : 201–234.
- REITTER E. 1887b. Bestimmungs-Tabellen der europäischen Coleopteren. III. Heft. Enthaltend die Familien: *Scaphidiidae*, *Lathridiidae* und *Dermestidae*. Zweite vermehrte und verbesserte Auflage. Mödling, 75 pp.
- REITTER E. 1887c. Coleopterologische Notizen. XXIII. Wien. Ent. Ztg., Wien, **6** : 171–173.
- REITTER E. 1889. Neue Coleopteren aus Europa, den angrenzenden Ländern und Sibirien, mit Bemerkungen über bekannte Arten. Sechster Theil. Dtsch. Ent. Z., Berlin, **1889** : 17–44.
- REITTER E. 1889b. Berichte über die von E. v. OERTZEN im Jahre 1887 in Griechenland u. Klein-Asien gesammelten Coleopteren. IX. Neue Arten aus verschiedenen Familien. Dtsch. Ent. Z., Berlin, **1889** : 251–259.
- REITTER E. 1889c. Neue Coleopteren aus Europa, den angrenzenden Ländern und Sibirien, mit Bemerkungen über bekannte Arten. Siebenter Theil. Dtsch. Ent. Z., Berlin, **1889** : 273–288.
- REITTER E. 1889d. Insecta, a Cl. G. N. POTANIN in China et in Mongolia novissime lecta. VIII. *Clavicornia*. *Hydrophilidae*. *Bruchidae*. Horae Soc. Ent. Ross., S.-Peterburg, **23** : 555–559.
- REITTER E. 1889e. Coleopterologische Notizen XXXI. Wien. Ent. Ztg., Wien, **8** : 125–128.
- REITTER E. 1889f. Coleopterologische Notizen XXXII. Wien. Ent. Ztg., Wien, **8** : 163.
- REITTER E. 1891. *Dermestidae*. In: Catalogus Coleopterorum Europae, Caucasi et Armeniae rossicae. Auctoribus Dr. L. v. HEYDEN, E. REITTER et J. WEISE cum aliis sociis coleopterologicis. Berlin, Mödling, Caen, VIII + 420 pp.

- REITTER E. 1892. Sechster Beitrag zur Coleopteren-Fauna des Russischen Reiches. Wien. Ent. Ztg., Wien, **11** : 133–136.
- REITTER E. 1894. Coleopterologische Notizen. LI. Wien. Ent. Ztg., Wien, **13** : 115–117.
- REITTER E. 1896. Vierzehnter Beitrag zur Coleopteren-Fauna des russischen Reiches. Wien. Ent. Ztg., Wien, **15** : 285–291.
- REITTER E. 1899. Ueber einige Coleopteren aus der palaearctischen Fauna und aus Japan. Ent. Nachr., Berlin, **25** : 216–220.
- REITTER E. 1899b. EVERTS Chr. Ed. *Coleoptera Nederlandica*. Wien. Ent. Ztg., Wien, **18** : 196.
- REITTER E. 1899c. Abbildungen und Beschreibungen neuer oder wenig gekannter Coleopteren aus der palaearctischen Fauna. Wien. Ent. Ztg., Wien, **18** : 282–287, pl. IV.
- REITTER E. 1900. *Coleoptera*, gesammelt im Jahre 1898 in Chin. Central-Asien von Dr. HOLDERER in Lahr. Wien. Ent. Ztg., Wien, **19** : 153–166, pl. I.
- REITTER E. 1900b. Neue, von Herrn Dr. John SAHLBERG auf seinen Reisen in Corfu, Palästina und Central-Asien gesammelte Coleopteren. Wien. Ent. Ztg., Wien, **19** : 217–220.
- REITTER E. 1900c. Neue Coleopteren aus Europa, der angrenzenden Ländern, russisch Asien und der Mongolei. Dtsch. Ent. Z., Berlin, **1899** : 273–281.
- REITTER E. 1905. Sechzehn neue Coleopteren aus der palaearktischen Fauna. Wien. Ent. Ztg., Wien, **24** : 241–251.
- REITTER E. 1906. *Dermestidae*. In: Catalogus Coleopterorum Europae, Caucasi et Armeniae Rossicae. Auctoribus L. v. HEYDEN, E. REITTER, J. WEISE cum aliis sociis coleopterologicis. Editio secunda. Berlin, Paskau, Caen, VI + 775 pp.
- REITTER E. 1908. Beschreibung einiger neuen Käferarten von Egypten. Bull. Soc. Ent. Égypte, Le Caire, **1** : 39–56.
- REITTER E. 1908b. Sieben neue Coleopteren aus Europa und den angrenzenden Ländern. Wien. Ent. Ztg., Wien, **27** : 213–216.
- REITTER E. 1910. Ueber eine wenig bekannte Gattung der Dermestiden. Ent. Z., Stuttgart, **23** : 239.
- REITTER E. 1911. Fauna Germanica. Die Käfer des Deutschen Reiches. Nach der analytischen Methode bearbeitet. III. Stuttgart, 436 pp., 147 ff., 48 pls.
- REITTER E. 1911b. Zwei neue Coleopteren-Varietäten. Wien. Ent. Ztg., Wien, **30** : 291.
- REY C. 1889. Remarques en passant. Échange, Lyon, **5** : 58–59.
- RILEY C. V. 1894. The insects occurring in the foreign exhibits of the world's Columbian exposition. Ins. Life, Washington, **6** : 213–227.
- ROSENHAUER W. G. 1856. Die Thiere Andalusiens nach dem Resultate einer Reise zusammengestellt, nebst den Beschreibungen von 249 neuen oder bis jetzt noch unbeschriebenen Gattungen und Arten. Erlangen, VIII + 429 pp., 3 pls.
- ROSSI P. 1788. Osservazioni insettologiche. Mem. Mat. Fis. Soc. Ital., Verona, **4** : 122–149, 1 pl.
- ROSSI P. 1790. Fauna Etrusca, sistens *Insecta*, quae in provinciis Florentina et Pisana praesertim collegit. I. Liburni, XXIII + 272 pp.
- ROSSI P. 1792. Mantissa Insectorum exhibens species nuper in Etruria collectas, adiectis faunae Etruscae illustrationibus, ac emendationibus. Pisis, 148 pp.
- ROSSI P. 1794. Mantissa Insectorum exhibens species nuper in Etruria collectas, adiectis faunae Etruscae illustrationibus, ac emendationibus. Tom II. Pisis, 154 pp., 8 pls.
- ROTH J. R. 1851. Diagnosen neuer *Coleoptera* aus Abyssinien. Arch. Naturges., Berlin, **17** : 115–133.
- ROUBAL J. 1932. Addition à la liste des Coléoptères de Monte Gargano. Boll. Soc. Ent. Ital., Genova, **64** : 66.
- RÜSKAMP F. 1929. Zur rheinischen Käferfauna V. Ent. Bl., Berlin, **25** : 150–157.
- RÜSKAMP F. 1930. Zur rheinischen Käferfauna VII. Ent. Bl., Berlin, **26** : 7–13.

- SAHLBERG C. R. 1819. *Dissertatio entomologica Insecta Fennica enumerans.* T. I., pars IV. Resp. J. HIRN. Aboe, pp. 41–56.
- SAHLBERG J. 1903. *Coleoptera mediterranea et rosso-asiatica nova et minus cognita, maxima ex parte itineribus annis 1895–1896 et 1898–1899 collecta.* II. Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **45**, No. 10, 40 pp.
- SAHLBERG J. 1903b. *Messis hiemalis Coleopterorum Corcyraeorum. Enumeratio Coleopterorum mensibus Novembri-Februario 1895–1896 et 1898–1899 nec non primo vere 1896 in insula Corcyra collectorum.* Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **45**, No. 11, 85 pp.
- SAHLBERG J. 1903c. *Entomologiska forskningsresor i Medelhafstrakerna och Centralasien företagna aren 1895–1896 samt 1898–1899. Resenberättelse.* II. Resor i Palestina, Egypten, Tunisien och Algeriet. Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **45**, No. 17, 39 pp.
- SAHLBERG J. 1903d. *Coleoptera Levantina mensibus Februario et Martio 1896 in Palaestina et Aegypto inferiore collecta.* Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **45**, No. 18, 36 pp.
- SAHLBERG J. 1903e. *Coleoptera Numido-Punica mensibus Martio, Aprili et Majo 1899 in Tunisia et Algeria orientali collecta.* Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **45**, No. 19, 70 pp.
- SAHLBERG J. 1913. *Coleoptera mediterranea et rosso-asiatica nova et minus cognita maxima ex parte itineribus annis 1895–1896, 1898–1899 et 1903–1904 collecta.* IV. Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **55A**, No. 8, 88 pp.
- SAHLBERG J. 1913b. *Coleoptera balcanica quae mensibus Octobri et Decembri 1903 atque Martis et Aprili 1906 in peninsula balcanica collegerunt John SAHLBERG et Unio SAALAS.* Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **55A**, No. 15, 108 pp.
- SAHLBERG J. 1913c. *Coleoptera mediterranea orientalia, quae in Aegypto, Palestina, Syria, Caramania atque in Anatolia occidentali anno 1904 collegerunt John SAHLBERG et Unio SAALAS.* Öfvers. Finska Vetensk.-Soc. Förhandl., Helsingfors, **55A**, No. 19, 281 pp.
- SAY Th. 1823. Descriptions of Coleopterous Insects collected in the late Expedition to the Rocky Mountains, performed by order of Mr. CALHOUN, Secretary of War, under the command of Major LONG. Journ. Acad. Nat. Sci. Philad., Philadelphia, **3** : 139–216.
- SAY Th. 1825. Descriptions of new species of Coleopterous Insects inhabiting the United States. J. Acad. Nat. Sci. Philad., Philadelphia, **5** : 160–204.
- SAY Th. 1832. New species of North American Insects, found by Joseph BARABINO, chiefly in Louisiana. New Harmony, 16 pp.
- SCHAFFER Ch. 1931. On a few new and known *Coleoptera*. Bull. Brooklyn Ent. Soc., Lancaster, Pa., **26** : 174–176.
- SCHAFFER I. Ch. 1766. *Elementa Entomologica.* Ratisbonae, 168 pp., CXXXIII pls.
- SCHAUM H. 1847. Bemerkungen über Fabricische Käfer. Stett. Ent. Ztg., Stettin, **8** : 39–57.
- [SCHAUM H.]. 1852. Catalogus Coleopterorum Europae. Vierte Auflage. Berlin, V + 97 pp.
- SCHAUM H. 1859. Catalogus Coleopterorum Europae. In Verbindung mit Dr. G. KRAATZ und H. v. KIESENWETTER. Berlin, IV + 121 pp.
- SCHAUM H. 1862. Catalogus Coleopterorum Europae. Editio secunda aucta et emendata. Berolini, 130 pp.
- SCHILSKY J. 1889. Berichtungen und Ergänzungen zum Verzeichniss der deutschen Käfer. Dtsch. Ent. Z., Berlin, **1889** : 337–344.
- SCHILSKY J. 1889b. Synonymische und andere Bemerkungen zu Dr. Carl W. v. DALLA TORRE'S „Synopsis der Insecten Oberösterreichs“ und „Die Käferfauna von Oberösterreich“. (Fortsetzung und Schluss.). Dtsch. Ent. Z., Berlin, **1889** : 345–356.
- [SCHMIDT]. 1844. Catalogus Coleopterorum Europae. Zusammengestellt auf Veranlassung des Entomologischen Vereins zu Stettin. Stettin, 76 pp.

- [SCHNEIDER D. H.]. 1785. Nomenclator entomologicus oder systematisches Nahmen-Verzeichniss der bis jetzt bekannt gewordenen Insekten. Stralsund, 68 pp.
- SCHÖNHERR C. J. 1806. Synonymia Insectorum, oder: Versuch einer Synonymie Aller bisher bekannten Insecten: nach Fabricii Systema Eleutheratorum geordnet, mit Berichtigungen und Anmerkungen wie auch Beschreibungen neuer Arten und illuminirten Kupfern. Erster Band. *Eleutherata* oder Käfer. Erster Theil. *Lethrus* ... *Scolytes*. Stockholm, XXII + 289 pp., 3 pls.
- SCHÖNHERR C. J. 1808. Synonymia Insectorum, oder: Versuch einer Synonymie Aller bisher bekannten Insecten, nach Fabricii Systema Eleutheratorum &c. geordnet, mit Berichtigungen und Anmerkungen, wie auch Beschreibungen neuer Arten und illuminirten Kupfern. Erster Band. *Eleutherata* oder Käfer. Zweiter Theil. *Spercheus* ... *Cryptocephalus*. Stockholm, IX + 423 pp., pl. IV.
- SCHRANK F. v. P. 1776. Beyträge zur Naturgeschichte. Augsburg, [8] + 140 pp., 7 pls.
- SCHRANK F. v. P. 1781. Enumeratio Insectorum Austriae indigenorum. Augustae Vindelicorum, [24] + 548 pp., 4 pls.
- SCHRANK F. v. P. 1782. Kritische Revision des österreichischen Insectenverzeichnisses. Neues Mag. Ent. Füessli, Winterthur, 1 : 135–168.
- SCHRANK F. v. P. 1785. Verzeichniss beobachteter Insecten im Fürstenthume Berchtesgaden. Neues Mag. Ent. Füessli, Winterthur, 2, 4 : 313–345.
- SCHRANK F. v. P. 1789. Entomologische Beobachtungen. Naturf., Haale, 24 : 60–90.
- SCHRANK F. v. P. 1796. Naturhistorische Beobachtungen um Pöttmes, Neuburg und Weihering. In: Sammlung naturhistorischer und physikalischer Aufsätze. Nürnberg, pp. 97–226.
- SCHRANK F. v. P. 1798. Fauna Boica. Durchgedachte Geschichte der in Baiern einheimischen und zahmen Thiere. Erster Band zweyte Abtheilung. Nürnberg, pp. 293–720.
- SCOPOLI I. A. 1763. *Entomologia Carniolica exhibens Insecta Carnioliae indigena et distributa in ordines, genera, species, varietates. Methodo Linnaeana. Vindobonae*, [32] + 420 + [1] pp.
- SCOPOLI I. A. 1772. Annus V. Historico-Naturalis. Lipsiae, 128 pp.
- SCOTT H. 1926. Coleoptera from the Seychelles and adjacent Islands: *Carabidae* (Supplement), *Cryptophagidae* (Supplement), *Dermestidae*, *Lymexylonidae*, *Rhipiceridae*, *Sphindidae*, *Throscidae*, *Brenthidae*. Ann. Mag. Nat. Hist., London, ser. 9, 18 : 50–76.
- SCRIBA L. G. 1790. Journal für die Liebhaber der Entomologie. Heft 1–2. Frankfurt, 192 pp.
- SCRIBA L. G. 1791. Journal für die Liebhaber der Entomologie. Heft 3. Frankfurt, pp. 193–296.
- SCUDDER S. H. 1900. Adephagous and Clavicorn Coleoptera from the Tertiary deposits at Florissant, Colorado, with descriptions of a few other forms and a systematic list of the non-rhynchophorous Tertiary Coleoptera of North America. Monogr. U. S. Geol. Survey, Washington, 40 : 3–148, 11 pls.
- SEMONOV-TIAN-SHANSKIJ A. 1912. Analecta coleopterologica. XVII. Rus. Ent. Obozr., St.-Pétersbourg, 12 : 497–502.
- SEMONOV-TIAN-SHANSKIJ A. 1914. Analecta coleopterologica. Rus. Ent. Obozr., Petrograd, 14 : 14–22.
- SEMONOV-TIAN-SHANSKIJ A. 1916. Popravka (Corrigendum). Rus. Ent. Obozr., Petrograd, 16 : 414.
- SHARP D. 1877. Descriptions of some new species, and indications of new genera of Coleoptera from New Zealand. Ent. Monthly Mag., London, 13 : 265–272.
- SHARP D. 1902. *Dermestidae*. In: Biologia Centrali-Americanica. Insecta. Coleoptera. Vol. II. Part. 1 [London], pp. 642–669, pl. 19.
- SHARP D. 1908. *Dermestidae*. In: D. SHARP, Fauna Hawaïensis or the zoology of the Sandwich (Hawaiian) Isles: Being Results of the Explorations instituted by the Joint Committee appointed by the Royal Society of London for promoting natural knowledge and the British Association for the advancement of science and carried on with the assistance of those Bodies and of the Trustees of the Bernice Pauahi Bishop Museum at Honolulu. Vol. III, part V. Cambridge, pp. 367–579, pls. XIII–XVI.

- SJÖBERG O. 1950. Drei neue Käfer aus der borealen Region Schwedens. Ent. Tidskr., Stockholm, **71** : 159–164.
- SLOSSON A. T. 1903. A coleopterous conundrum. Canad. Entomol., London, Ont., **35** : 183–187.
- SOLIER [A. J.] 1849. Orden III. Coleopteros. In: GAY C., Historia fisica y politica de Chile segun documentos adquiridos en esta republica durante doce años de residencia en ella y publicada bajo los auspicios del supremo gobierno. Zoologia, tomo cuarto. Paris, Chile, pp. 105–508.
- SOLIER [A. J.] 1854. Coleopteros. In: GAY C., Atlas de la Historia Fisica y Politica de Chile. Tomo segundo. Paris.
- SOLSKIY S. M. 1876. Zhestkokrylyya (*Coleoptera*) tetrad vtoraya. In: FEDTSHENKO A. P., Puteshestvie v Turkestan, t. 2, tsh. 5, tetr. 2. Izv. Imp. Obshtsh. Lyub. Yest. Antrop. Entogr., S.-Peterburg, Moskva, **21**, 1 : 223–398 + I–XXIV, pls. I, IV.
- SPENCER G. J. 1948. Notes on some *Dermestidae* of British Columbia (*Coleoptera*). Proc. Ent. Soc. Brit. Columbia, Vernon, **44** : 6–9.
- SPENCER G. J. 1956. Some unusual records of beetles in Vancouver. Proc. Ent. Soc. Brit. Columbia, Vernon, **52** : 31–32.
- STEINHEIL E. 1869. Symbolae ad historiam Coleopterorum Argentiniae meridionalis, ossia elenco dei coleotteri raccolti dal professore Pellegrino STROBEL, durante il suo soggiorno in Buenos Aires e nei viaggi di là intrapresi a Mendoza e nel Chili, indi a Bahia Blanca ed al Carmen de los Patagones; e descrizione delle specie nuove. Atti Soc. Ital. Sci. Nat., Milano, **12** : 238–260.
- STEPHENS J. F. 1829. The nomenclature of British Insects; being a compendious list of such species as are contained in the systematic catalogue of British Insects, and forming a guide to their classification, &c. &c. London, 4 + 68 pp.
- STEPHENS J. F. 1830. Illustrations of British Entomology; or, a Synopsis of Indigenous Insects: containing their generic and specific distinctions: with an account of their metamorphoses, times of appearance, localities, food, and economy, as far as practicable. *Mandibulata*. III. London, 380 pp., pls. XVI–XIX.
- STEPHENS J. F. 1832. Illustrations of British Entomology; or a Synopsis of Indigenous Insects: containing their generic and specific distinctions; with an account of their metamorphoses, times of appearance localities, food, and economy, as far as practicable. *Mandibulata*. Vol. V. London, 448 pp., pls. 24–27.
- STEPHENS J. F. 1833. The nomenclature of British Insects; together with their synomyms: being a compendious list of such species as are contained in the systematic catalogue of British Insects, and of those discovered subsequently to its publication; forming a guide to their classification, &c. &c. Second Edition. London, iv + 136 pp.
- STEPHENS J. F. 1839. A Manual of British *Coleoptera*, or beetles; containing a brief description of all the species of beetles hitherto ascertained to inhabit Great Britain and Ireland; together with a notice of their chief localities, times and places of appearances, etc. London, XII + 443 pp.
- STIERLIN G. 1902. Beschreibung einiger neuen Species von Coleopteren. Mitt. Schweiz. Ent. Ges., Schaffhausen, **10** : 425–427.
- STRÖM H. 1783. Norske Insecters Beskrivelse med Anmaerkninger. Nye Saml. Danske Viedensk. Selsk. Skrift., Kiobenhavn, **2** : 49–93, 2 pls.
- STURM J. 1826. Catalog meiner Insecten-Sammlung. Erster Theil. Käfer. Nürnberg, VIII + 207 pp., 4 pls.
- STURM J. 1843. Catalog der Kaefer-Sammlung. Nürnberg, XII + 386 pp., 6 pls.
- SULZER [J. H.] 1776. Abgekürzte Geschichte der Insecten Nach dem Linaeischen System. [I] – [II]. Winterthur, XXVIII + 274 + 72 pp., 32 pls.
- SUMAKOV G. G. 1907. Novyy vid iz roda *Trogoderma* LATR. (*Coleoptera, Dermestidae*) i obzor vidov etogo roda. Rus. Ent. Obozr., St. Petersbourg, **7** : 15–19.
- SUMAKOV G. G. 1912. Materialy k poznaniyu fauny zhestkokrylych Sredney Azii. Rus. Ent. Obozr., St.-Petersbourg, **12** : 423–433.

- TENENBAUM Sz. 1915. Fauna koleopterologiczna wysp Balearskich. Warszawa, 150 pp.
- THOMSON C. G. 1862. Skandinaviens *Coleoptera*, synoptiskt bearbetade. IV. Lund, 269 pp.
- THUNBERG C. P. 1781. Dissertatio entomologica novas Insectorum species, sistens. Pars Prima. Upsaliae, 28 pp., 1 pl.
- THUNBERG C. P. 1784. Novae Insectorum species descriptae. Nova Acta Reg. Soc. Sci. Upsaliensis, Upsaliae, 4 : 1–28, pl. I, ff. 1–6.
- THUNBERG C. P. 1787. Museum Naturalium Academiae Upsaliensis. Cujus Partem Tertiam. Donation. Thunbergiana. Upsaliae, pp. 33–42.
- THUNBERG C. P. 1794. Dissertatio Entomologica, sistens *Insecta Suecica*. Pars 5–6. Upsaliae, pp. 63–82.
- THUNBERG C. P. 1794b. Museum Naturalium Academiae Upsaliensis. Appendix III. Donation. Thunbergian. Upsaliae, pp. 131–144.
- THUNBERG C. P. 1795. Dissertatio Entomologica, sistens *Insecta Suecica*. Pars 9. Upsaliae, pp. 105–113.
- THUNBERG C. P. 1815. Anthreni monographia. Nova Acta Reg. Soc. Sci. Upsaliensis, Upsaliae, 7 : 150–156.
- THUNBERG C. P. 1820. *Coleoptera Capensia*, antennarum clava solida et perfoliata, collecta, recensita et descripta. Mém. Acad. Imp. Sci. St.-Pétersb., St. Pétersbourg, 7 : 362–372.
- TOURNIER H. 1860. Drei neue europäische Käferarten. Berlin. Ent. Z., Berlin, 4 : 317–319.
- TOURNIER H. 1869. Rectifications sur les quelques espèces suivantes de coléoptères. Petites Nouv. Ent., Paris, 1 : 11.
- UYTTENBOOGAART D. L. 1937. Report on Canarian *Coleoptera* collected by R. FREY and R. STORÅ in 1931 for the Museum Zoologicum Universitatis Helsingfors. (Contributions to the Knowledge of the Fauna of the Canary-Islands. XVI). Comm. Biol. Soc. Sc. Fenn., Helsingfors, 6, 2, 17 pp., 3 ff.
- VILLERS C. 1789. Caroli LINNAEI Entomologia, Faunae Suecicae Descriptionibus Aucta; DD. SCOPOLI, GEOFFROY, DE GEER, FABRICII, SCHRANK, Etc. speciebus vel in Systemate non enumeratis, vel nuper prime detectis, vel speciebus Galliae Australis locupletata, generum specierumque rariorum iconibus ornata. Tomus Primus. Lugduni, XXIV + 765 pp., 3 pls.
- VILLERS C. 1789b. Caroli LINNAEI Entomologia, Faunae Sueciae descriptionibus aucta; DD. SCOPOLI, GEOFFROY, de GEER, FABRICII, SCHRANK, etc. speciebus vel in Systemate non enumeratis, vel nuper prime detectis, vel speciebus Galliae Australis locupletata, generum specierumque rariorum iconibus ornata. Tomus quartus. Lugduni, cxxiii : 556 pp., pl. XI.
- WALKER F. 1858. Characters of some apparently undescribed Ceylon Insects. Ann. Mag. Nat. Hist., London, ser. 3, 2 : 202–209.
- WALKER F. 1859. Characters of some apparently undescribed Ceylon Insects. (Continued from vol. ii. pp. 286). Ann. Mag. Nat. Hist., London, ser. 2, 3 : 50–56.
- WATERHOUSE Ch. O. 1883. Description of a new Species of *Anthrenus* from India (*Coleoptera, Dermestidae*). Ann. Mag. Nat. Hist., London, ser. 5, 11 : 61.
- WICKHAM H. F. 1912. A report on some recent collections of fossil *Coleoptera* from the Miocene shales of Florissant. Bull. Lab. Nat. Hist. Univ. Iowa, Iowa, 6, 3 : 3–38, 8 pls.
- WICKHAM H. F. 1913. The Princeton collection of fossil Beetles from Florissant. Ann. Ent. Soc. America, Columbus, 6 : 359–366, pls. XXXVIII–XLI.
- WINKLER A. 1926. *Dermestidae*. In: Catalogus Coleopterorum Regionis Palaearcticae. I, 6. Wien, pp. 675–682.
- WOLLASTON T. V. 1863. Diagnostic Notices of New Canarian *Coleoptera*. Ann. Mag. Nat. Hist., London, ser. 3, 11 : 214–221.
- WOLLASTON T. V. 1864. Catalogue of the Coleopterous Insects of the Canaries in the collection of the British Museum. London, xiii + 648 pp.
- WOLLASTON T. V. 1865. *Coleoptera Atlantidum*, being an enumeration of the Coleopterous Insects of the Madeiras, Salvages, and Canaries. London, xlvii + 526 pp. + Appendix 140 pp.

- WOLLASTON T. V. 1867. *Coleoptera Hesperidum*, being an enumeration of the Coleopterous Insects of the Cape Verde Archipelago. London, xl + 285 pp., 1 map.
- WRADATSCH G. 1914. Die Käferausbeute von 1913 und die angewendete Fangtechnik. (Schluss.). Ent. Bl., Berlin, **10** : 150–154.
- XAMBEU [V.]. 1892. Contributions aux faunes locales des régions de l'Est et du Sud-Est. Échange, Lyon, **8** : 45–46.
- ZAJCEV F. A. 1909. A remark on the *Ignotus aenigmaticus*. Canad. Entomol., Guelph, **41** : 37.
- ZAJCEV F. A. 1919. Materialy k faune zhestkokrylych Kavkazskavo kraja. VIII. *Dermestidae et Bostrychidae*. Izv. Kavkaz. Muz., Tiflis, **12** : 161–186.
- ZETTERSTEDT J. W. 1828. Fauna Insectorum Lapponica. Pars I. Hammone, XXI + 563 pp.
- ZHANTIEV R. D. 1960. Materialy po ekologii *Dermestidae* Centralnogo Kazakhstana. Zool. Zhurn., Moskva, **39** : 1628–1636.
- ZHANTIEV R. D. 1963. Novye vidy kozheedov roda *Attagenus* LATR. (*Coleoptera, Dermestidae*) iz Sredney Azii. Zool. Zhurn., Moskva, **42** : 454–457, 12 ff.
- ZHANTIEV R. D. 1963b. Ob ekologii kozheedov (*Coleoptera, Dermestidae*) Sredney Azii. Zool. Zhurn.. Moskva, **42** : 1052–1063.
- ZHANTIEV R. D. 1963c. Kozheedy roda *Attagenus* LATR. (*Coleoptera, Dermestidae*) fauny Sovetskogo Soyuza. Ent. Obozr., Leningrad, **42** : 415–424, 7 ff.
- ZHANTIEV R. D. 1967. Opyt taksonomicheskogo analiza roda *Dermestes* L. (*Coleoptera, Dermestidae*). Zool. Zhurn., Moskva, **46** : 1350–1356.
- ZIEGLER D. 1845. Descriptions of New North American *Coleoptera*. Proc. Ent. Soc. Philad., Philadelphia, **2** : 226–272.
- ZIMMERMANN C. 1869. Synonymical notes on *Coleoptera* of the United States, with descriptions of new species, from the MSS. of the late Dr. C. ZIMMERMANN. Edited by John L. LECONTE, M. D. Trans. Amer. Ent. Soc., Philadelphia, **2** : 243–259.

V. Alphabetical Index of Names of Subfamilies, Genera, Subgenera and their Synonymical Names

- Acolpus* JAYNE, 1882 : 347 (synonym of *Trogoderma* DEJ.) — 99
Adelaidaea BLACKBURN, 1903 : 159 (synonym of *Adelaidia* BLACKB.) — 125
Adelaidia BLACKBURN, 1891 : 130 — 24, 34, 44, 125
Aethriostoma REITTER, 1908 : 45 (synonym of *Orphinus* MOTSCH.) — 125
Aethriostoma MOTSCHULSKY, 1858 : 47 (subgenus of *Attagenus* LATR.) — 24, 25, 26, 37, 38, 79
Anthrenocerus ARROW, 1915 : 443 — 22, 33, 44, 79, 124
Anthrenodes CHOBAUT, 1898 : 76 (subgenus of *Anthrenus* SCHAEFF.) — 47, 48, 146
Anthrenops REITTER, 1881b : 86 (subgenus of *Anthrenus* SCHAEFF.) — 29, 47, 48, 147
Anthrenus SCHAEFFER, 1766 : [26] — 19, 22, 24, 25, 26, 27, 28, 29, 30, 32, 34, 47, 48, 139
Anthrenus s. str. (subgenus of *Anthrenus* SCHAEFF.) — 47, 48, 49, 140
Anthreninae LECONTE, 1861 : 107 — 34, 47, 139
Approgramme REITTER, 1887b : 55 (synonym of *Megatoma* HERBST) — 116
Aprogramme REITTER, 1887b : 40 (synonym of *Megatoma* HERBST) — 116
Apsectus LECONTE, 1854 : 107 — 20, 21, 29, 31, 34, 49, 155
Argocerus SHARP, 1908 : 411 — 22, 33, 43, 122
Asidora MULSANT et REY, 1868 : 122 (synonym of *Trogoderma* DEJ.) — 99
Asprogramme GOZIS, 1886 : 10 (synonym of *Megatoma* HERBST) — 116
Attageninae CASTELNAU, 1840 : 35 — 33, 37, 79
Attagenus LATREILLE, 1802 : 121 — 19, 22, 24, 25, 26, 28, 29, 32, 33, 37, 38, 39, 79
Attagenus s. str. (subgenus of *Attagenus* LATR.) — 37
Attagenus LATREILLE, 1829 : 510 (synonym of *Ctesias* STEPH.) — 129
Axinocerus JAYNE, 1882 : 347 (synonym of *Thaumaglossa* REDT.) — 130
Brachysphyrus BLACKBURN, 1903 : 159 (synonym of subgenus *Aethriostoma* MOTSCH., genus *Attagenus* LATR.) — 79
Byrrhus SULZER, 1776 : [I], 25, nec LINNAEUS, 1767 : 568 (synonym of *Anthrenus* SCHAEFF.) — 139
Caccoleptus SHARP, 1902 : 649 — 19, 33, 43, 123
Caucasotoma MROCKOWSKI, 1967 : 5 (subgenus of *Megatoma* HERBST) — 43, 116
Cryptorhopalum GUÉRIN-MÉNEVILLE, 1838b : 67 — 20, 21, 29, 31, 34, 46, 133
Ctenias REITTER, 1911 : 151 (synonym of *Ctesias* STEPH.) — 129
Ctesias STEPHENS, 1830 : 124 — 25, 26, 28, 29, 31, 32, 34, 45, 129
Curtophinus PIC, 1954b : 11 — 34, 45, 129
Dearthus JAYNE, 1882 : 346 (synonym of *Dearthrus* LE.C.) — 122
Dearthrus LECONTE, 1861 : 108 — 29, 31, 32, 33, 43, 122
Decamerus SOLIER, 1849 : 369 — 19, 33, 40, 99
Dermestes LINNAEUS, 1758 : 342 — 19, 21, 22, 24, 26, 27, 28, 29, 30, 32, 33, 36, 51, 52
Dermestes s. str. (subgenus of *Dermestes* L.) — 59
Dermestinae LATREILLE, 1807 : 3 — 33, 51, 52
Dermestinus ZHANTIEV, 1967 : 1352 (subgenus of *Dermestes* L.) — 52
Didontolobus FAIRMAIRE, 1884 : 488 (synonym of *Diontolobus* SOLIER) — 123
Diodontolobus LACORDAIRE, 1854 : 458 (synonym of *Diontolobus* SOLIER) — 123
Diontolobus SOLIER, 1849 : 367 — 19, 33, 44, 123
Distontobus SOLIER, 1854 : t. 8, f. 4 (synonym of *Diontolobus* SOLIER) — 123
Ditorines NORMAND, 1936 : 298 — 26, 28, 29, 33, 43, 123
Egidyella REITTER, 1899c : 284 — 28, 29, 33, 35, 51
Egidyellinae SEMENOV-TIAN-SHANSKIJ, 1916 : 414 — 33, 35, 51
Elania MULSANT et REY, 1868 : 111 (subgenus of *Globicornis* LATR.) — 111
Entomotrogus GANGLBAUER, 1904b : 27 (synonym of *Trogoderma* DEJ.) — 99
Eocerus SHARP, 1908 : 412 — 22, 33, 43, 122

- Epidyella* SEMENOV-TIAN-SHANSKIJ, 1914 : 15 (synonym of *Egidyella* REITT.) — 52
Epidyellini SEMENOV-TIAN-SHANSKIJ, 1914 : 15 (synonym of *Egidyellinae* WINKL.) — 51
Eucnocerus SHARP, 1902 : 648 (synonym of *Trogoderma* DEJ.) — 99
Eurhopalus SOLIER, 1849 : 372 (synonym of *Trogoderma* DEJ.) — 99
Evorinea BEAL, 1961 : 118 — 23, 24, 34, 50, 157
Falsocrryptorhopalum PIC, 1936 : 1 — 19, 34, 46, 133
Falsoorphinus PIC, 1931b : 4 (subgenus of *Orphinus* MOTSCH.) — 126
Florilinus MULSANT et REY, 1868 : 141 (subgenus of *Anthrenus* SCHAEFF.) — 30, 47, 48, 148
Genattus SHARP, 1902 : 646 (synonym of *Novelsis* CASEY) — 97
Globicornis LATREILLE, 1829 : 511 — 28, 29, 33, 42, 113
Globicornis s. str. (subgenus of *Globicornis* LATR.) — 113
Hadrotoma ERICHSON, 1846 : 424 (subgenus of *Globicornis* LATR.) — 114
Helocerus MULSANT et REY, 1868 : 141 (subgenus of *Anthrenus* SCHAEFF.) — 48, 149
Hemirhopalum SHARP, 1902 : 651 — 19, 20, 34, 46, 132
Hexanodes BLAIR, 1941 : 15 — 22, 34, 50, 158
Hirtomegatoma PIC, 1931 : 5 — 28, 29, 33, 42, 116
Hospitopterus PIC, 1921 : 54 (synonym of *Thylodrias* MOTSCH.) — 158
Hypoceuthes GERSTAECKER, 1871 : 46 (synonym of *Anthrenus* SCHAEFF.) — 139
Ignotus SLOSSON, 1903 : 187 (synonym of *Thylodrias* MOTSCH.) — 158
Labrocerus SHARP, 1885 : 148 — 22, 33, 43, 121
Lanorus MULSANT et REY, 1868 : 68 (subgenus of *Attagenus* LATR.) — 37, 79
Macropriion HOPE, 1840 : 68 (synonym of *Trogoderma* DEJ.) — 99
Mariouta PIC, 1898b : 73 — 28, 29, 33, 35, 51
Mariooutinae WINKLER, 1926 : 682 — 33, 35, 51
Megatoma DEJEAN, 1821 : 46 (synonym of *Ctesias* STEPH.) — 129
Megatoma HERBST, 1792 : 92 — 29, 31, 32, 33, 38, 42, 99, 116
Megatoma s. str. (subgenus of *Megatoma* HERBST) — 42, 117
Megatoma KUGELANN, 1792 : 479 nec HERBST, 1792 : 92 (synonym of *Attagenus* LATR.) — 79
Megatominae LEACH, 1815 : 94 — 33, 40, 99
Mesalia MULSANT et REY, 1868 : 106 (synonym of subgenus *Hadrotoma* ER. of genus *Globicornis* LATR.) — 114
Montandonia JACQUET, 1886b : 320 — 28, 29, 33, 37, 78
Myrmeanthrenus ARMSTRONG, 1945 : 47 — 22, 34, 44, 125
Nathrenus CASEY, 1900 : 161 (subgenus of *Anthrenus* SCHAEFF.) — 29, 47, 48, 49, 130
Neoanthrenus ARMSTRONG, 1941 : 388 — 22, 34, 49, 154
Novelsis CASEY, 1900 : 144 — 29, 30, 31, 33, 40, 97
Novelsis s. str. (subgenus of *Novelsis* CASEY) — 97
Ocelliger PHILIPPI et PHILIPPI, 1864 : 283 (synonym of *Trogoderma* DEJ.) — 99
Orbeola MULSANT et REY, 1868 : 134 (subgenus of *Phradonoma* JACQ. DU VAL) — 41, 111
Orphilinae LECONTE, 1861 : 107 — 34, 49, 154
Orphiloidea MATSUMURA et YOKOYAMA, 1928 : 130 (synonym of *Thaumaglossa* REDT.) — 130
Orphilus ERICHSON, 1846 : 424 — 31, 32, 34, 49, 154, 155
Orphinus MOTSCHULSKY, 1858 : 48 — 23, 24, 26, 29, 34, 44, 45, 125
Orphinus s. str. (subgenus of *Orphinus* MOTSCH.) — 126
Paranovelsis CASEY, 1900 : 148 (subgenus of *Novelsis* CASEY) — 98
Paratrogoderma SCOTT, 1926 : 63 — 25, 33, 41, 110
Perimegatoma HORN, 1875 : 135 (subgenus of *Megatoma* HERBST) — 42, 118
Phradonoma JACQUELIN DU VAL, 1859 : 257 — 25, 26, 28, 29, 33, 41, 111
Phradonoma s. str. (subgenus of *Phradonoma* JACQ. DU VAL) — 33, 111
Phradonomorphus PIC, 1916b : 3 — 25, 33, 41, 111
Psacus PASCOE, 1866 : 446 (synonym of *Trogoderma* DEJ.) — 99
Pseudohadrotoma KALIK, 1951c : 2 (subgenus of *Megatoma* HERBST) — 43, 117

- Pseudomegatoma* PIC, 1915b : 3 (synonym of *Trogoderma* DEJ.) — 99
Pseudomesalia GANGLBAUER, 1900 : 149 (synonym of *Dearthus* LEC.) — 122
Pseudotelopes PIC, 1916b : 2 (synonym of subgenus *Aethriostoma* MOTSCH. of genus *Attagenus* LATR.) — 79
Pseudothaumaglossa PIC, 1918b:1 — 19, 34, 46, 132
Ranolus BLAIR, 1929 : 382 (subgenus of *Attagenus* LATR.) — 38, 79
Ranthenus MROCKOWSKI, 1962 : 255 (subgenus of *Anthrenus* SCHAEFF.) — 48, 150
Reesa BEAL, 1967 : 310 — 29, 30, 31, 33, 41, 110
Reeveana DUNSTAN, 1923 : 48 — 159
Sefrania PIC, 1899 : 28 — 28, 29, 33, 37, 79
Solskinus MROCKOWSKI, 1962 : 252 (subgenus of *Anthrenus* SCHAEFF.) — 30, 48, 149
Telopes REDTENBACHER, 1843 : 976 (subgenus of *Attagenus* LATR.) — 38, 79
Telopus MULSANT et REY, 1868 : 101 (synonym of subgenus *Telopes* REDT. of genus *Attagenus* LATR.) — 79
Taumaglossa REDTENBACHER, 1867 : 43 — 23, 24, 26, 28, 32, 34, 45, 130
Thaumaphrastinae ANDERSON, 1949 : 127 — 34, 50, 158
Thaumaphrastus BLAISDELL, 1927b : 123 (synonym of *Thorictodes* REITT.) — 158
Thaumatoglossa CASEY, 1900 : 145 (synonym of *Thaumaglossa* REDT.) — 130
Thelodrias AGASSIZ et ERICHSON, 1846 : 161 (synonym of *Thylodrias* MOTSCH.) — 158
Thelydrias AGASSIZ, 1846 : 368 (synonym of *Thylodrias* MOTSCH.) — 158
Thelydriini SEMENOV-TIAN-SHANSKIJ, 1912 : 498 (synonym of *Thylodriadinae* MROCKOWSKI, 1954c : 10) — 158
Thorictini BEAL, 1959 : 100 (synonym of *Thaumaphrastinae* ANDERSON, 1949 : 127) — 158
Thorictodes REITTER, 1875c : 45 — 24, 34, 50, 158
Thylodriadinae MROCKOWSKI, 1954c : 10 — 34, 50, 158
Thylodriinae WINKLER, 1926 : 675 (synonym of *Thylodriadinae* MROCKOWSKI, 1954c : 10) — 158
Thylodrias MOTSCHULSKY, 1839 : 75 — 34, 50, 158
Tiresias STEPHENS, 1832 : 410 (synonym of *Ctesias* STEPH.) — 129
Tiresiomorpha PIC, 1954b : 11 — 34, 45, 130
Trichopleurias SAHLBERG, 1903 : 35 (synonym of *Egidyella* REITT.) — 51
Trinodes DEJEAN, 1821 : 47 — 24, 25, 26, 29, 30, 34, 50, 155, 156
Trinodinae CASEY, 1900 : 140 — 34, 49, 155
Trogoderma DEJEAN, 1821 : 46 — 19, 21, 22, 23, 24, 25, 28, 31, 33, 40, 41, 42, 99,
Tryoniopsis DUNSTAN, 1923 : 51 — 159

STRESZCZENIE

Praca niniejsza składa się z trzech części. W pierwszej autor omówił rozmieszczenie *Dermestidae* w poszczególnych obszarach zoogeograficznych, korygując granice tam, gdzie to wynikało z charakteru zasięgów poszczególnych rodzajów. Tak więc autor przyjął rozdzielenie Obszaru Holarktycznego na dwa Obszary: Palearktyczny i Nearktyczny. Prowincja Sonorska, zaliczana do Obszaru Nearktycznego, powinna być, zdaniem autora, wyłączona z tego obszaru i włączona do Obszaru Neotropikalnego. Karoliny powinny być zaliczone do Prowincji Filipińskiej Obszaru Orientalnego, a nie wchodzić w skład Obszaru Australijskiego.

Każdy z obszarów zoogeograficznych został odpowiednio scharakteryzowany materiałem z rodziny *Dermestidae*, przy czym zasięgi ważniejszych rodzajów zilustrowane zostały na mapkach.

W drugiej części autor podał zoogeograficzne charakterystyki wszystkich znanych rodzajów *Dermestidae* na podstawie analizy rozmieszczenia gatunków. Część ta była podstawą do wniosków podanych w części pierwszej.

Trzecią część pracy stanowi taksonomiczno-nomenklatoryczny katalog wszystkich znanych gatunków *Dermestidae*, uzupełniony ich rozmieszczeniem, podanym w możliwie zwięzlej formie. Zawiera on dane o 46 rodzinach, 882 gatunkach, 25 podgatunkach, 127 odmianach oraz 1342 nazwach synonimycznych. Prócz zebrania całości danych z piśmiennictwa, w katalogu uwzględnione są też i nie publikowane dotąd wyniki badań autora z zakresu zoogeografii i taksonomii. W części tej ustalono 26 nowych połączeń nazw gatunkowych z rodzajowymi, 15 nowych synonimów nazw gatunków, dwa nowe synonimy nazw rodzajowych oraz wprowadzono trzy nowe nazwy dla gatunków mających nazwy preokupowane: *Attagenus reitteri* nom. nov. dla *Attagenus antennatus* (REITTER, 1887b : 54) nec CASTELNAU, 1840 : 36, *Attagenus thunbergi* nom. nov. dla *Attagenus bifasciatus* (THUNBERG, 1815 : 151) nec (OLIVIER, 1790 : no. 9 p. 13) oraz *Trogoderma beali* nom. nov. dla *Trogoderma atrum* (PIC, 1937d : 193) nec (PHILIPPI et PHILIPPI, 1864 : 283).

Prace kończy wykaz 640 pozycji piśmiennictwa cytowanych w katalogu oraz skorowidz nazw podrodzin, rodzajów, podrodzajów i ich synonimów.

РЕЗЮМЕ

Настоящая работа состоит из трех частей. В первой автор рассматривает распространение *Dermestidae* в отдельных зоогеографических областях исправляя их границы, где это следует по распространению отдельных родов. И так автор принял разделение Голарктической области на две части: Палеарктическую и Неарктическую. Сонорская провинция должна быть, по мнению автора, выделена из Неарктической области и причислена к Неотропической области. Каролинские о-ва должны быть зачислены к Филиппинской провинции Ориентальной области, а не принадлежат к Австралийской области.

Каждая из зоогеографических областей охарактеризована материалами из семейства *Dermestidae*, при чем ареалы более важных родов представлены на картах.

Во второй части автор дает зоогеографические характеристики всех известных родов *Dermestidae* на основании анализа распространения видов. Эта часть была основой к выводам приведенным в первой части.

Третья часть работы представляет таксономически-номенклатурный каталог всех известных видов *Dermestidae* дополненный краткими данными о их распространении. Он содержит данные о 46 родах, 882 видах, 25 подвидах и 127 разно-

видностях и 1342 синонимических названий. Кроме собранных в полном виде данных по литературе, в каталоге учтены тоже не публикованные до сих пор результаты исследований автора по зоогеографии и таксономии *Dermestidae*. В этой части установлены 26 новых комбинаций видовых названий с родовыми, 15 новых синонимов названий видов, два новые синонимы родовых названий и введены три новые названия для видов имеющих преоккупированные названия: *Attagenus reitteri* nom. nov. для *Attagenus antennatus* (REITTER, 1887b : 54) nec CASTELNAU, 1840 : 36, *Attagenus thunbergi* nom. nov. для *Attagenus bifasciatus* (THUNBERG, 1815 : 151) nec (OLIVIER, 1790 : no. 9 p. 13) и *Trogoderma beali* nom. nov. для *Trogoderma atrum* (PIC, 1937d : 193) nec (PHILIPPI et PHILIPPI, 1864 : 283).

Работа оканчивается списком литературы содержащим 640 заглавий цитированных в каталоге, и указателем названий подсемейств, родов, подродов и их синонимов.

Redaktor pracy — prof. dr T. Jaczewski

Państwowe Wydawnictwo Naukowe — Warszawa 1968
Nakład 1 390+90 egz. Ark. wyd. 15, druk. 11¹/₂. Papier druk. sat. kl. III, 80 g, B. Cena 45 zł
Nr zam. 543/68 — Wrocławska Drukarnia Naukowa