

VOLUME 2

JANUARY, 1935

NUMBER 1

PROCEEDINGS
of The
Helminthological Society
of Washington

EDITORIAL COMMITTEE

JESSE R. CHRISTIE, *Editor*
U. S. Bureau of Plant Industry

EMMETT W. PRICE
U. S. Bureau of Animal Industry

GILBERT F. OTTO
Johns Hopkins University

HENRY E. EWING
U. S. Bureau of Entomology

G. DIKMANS
U. S. Bureau of Animal Industry

PUBLISHED BY THE
HELMINTHOLOGICAL SOCIETY OF WASHINGTON

Subscription \$1.00 a volume, foreign, \$1.25

PROCEEDINGS OF THE HELMINTHOLOGICAL SOCIETY OF WASHINGTON

The Proceedings of the Helminthological Society of Washington is a medium for the publication of notes and papers in helminthology and related subjects. Each volume consists of 2 numbers issued in January and July. Volume 1, number 1, was issued in April, 1934. The Proceedings are intended primarily for the publication of contributions by members of the Society but papers by persons who are not members will be accepted provided the author will contribute toward the cost of publication.

Manuscripts may be sent to any member of the editorial committee. Manuscripts must be typewritten (double spaced) and submitted in finished form for transmission to the printer. Authors should not confine themselves to merely a statement of conclusions but should present a clear indication of the methods and procedures by which the conclusions were derived. Except in the case of manuscripts specifically designated as *preliminary papers* to be published *in extenso* later, a manuscript is accepted with the understanding that it is not to be published, with essentially the same material, elsewhere.

Reprints will be furnished at cost in accordance with the schedule of prices printed below. An author's order for reprints should be attached to the first page of his manuscript. Failure to do so will be regarded as an indication that reprints are not desired.

Copies	1-2 pp.	3-4 pp.	5-8 pp.	9-12 pp.
50	\$1.33	\$2.50	\$4.00	\$6.00
100	2.00	3.50	5.50	7.50
150	2.66	4.50	7.00	9.00
200	3.33	5.50	8.00	10.00
250	4.00	6.00	9.00	11.00

Parcel Post charges extra.

Proceedings of previous meetings.—Previous to independent publication, which began in 1934, the proceedings of the 1st to 15th meeting of the Society were published in *Science*, those of the 16th to 156th meeting were published in the *Journal of Parasitology*. A limited number of sets of these Proceedings, complete except as noted, are available at \$5.00 a set. Price of individual reprints, previous to 51st meeting, 35 cents each; subsequent to 96th meeting, 25 cents each. Reprints contain the proceedings of from one to several meetings as follows: 1-12 (supply exhausted), 13, 14, 15, 16-20, 21, 22-24, 25-26, 27, 28-29, 30-38, 39-44, 45-50, 51-96 (supply exhausted), 97-103, 104-107, 108-110, 111-115, 116-120, 121-126, 127-130, 131-134, 135-139 (supply exhausted), 140-143, 144-148 (supply exhausted), 149-151, 152-154, 155-156.

Remittances should be made payable to Edna M. Buhner, Treasurer.

Correspondence may be addressed to the corresponding secretary, Edna M. Buhner, U. S. Bureau of Plant Industry, Washington, D. C., or to the editor, Jesse R. Christie, U. S. Bureau of Plant Industry, Washington, D. C.

OFFICERS OF THE SOCIETY FOR 1935:

President: LLOYD A. SPINDLER
Vice President: EVERETT E. WEHR
Recording Secretary: G. DIKMANS
Corresponding Secretary-Treasurer: EDNA M. BUHRER

Date of Issue, February 4, 1935

PROCEEDINGS OF THE HELMINTHOLOGICAL SOCIETY OF WASHINGTON

VOLUME 2

WASHINGTON, D. C., JANUARY, 1935

NUMBER 1

A KEY TO THE GENERA OF FREE-LIVING NEMAS

By N. A. COBB

Division of Nematology, U. S. Bureau of Plant Industry

Checked, revised and prepared for the press by Margaret V. Cobb and Corinne Cooper

PREFACE

This key, which was built up and used by N. A. Cobb as a card catalog during forty years of work in nematology, had been reorganized in rough manuscript form during the last two years of his life. As Miss Cooper and I had previously worked with him on this draft, it seemed best for us to carry it to completion. In essentials and in general form it is his key, but we are responsible both for correctness of detail (an appreciable amount of the detailed work was incomplete, in need of revision, or in need of change because of addition or omission of genera), and for such decisions as have to be made in getting such a work printed. Our aim has been to follow his ideas wherever they were known to us, or where we could infer them, and to make as few changes as possible in his outline. Nearly a hundred entries have been omitted, chiefly his own new genera which it has not been possible to publish in advance of publication of the key. In some cases genera have been dropped as not being free-living. Index, glossary and list of abbreviations have been added. The bibliography has been prepared by Mrs. Rowena R. LeHew of the Division of Nematology. The definitions in the glossary are not intended to settle the meaning of terms for other workers in the field, but merely to indicate the terminology used in this key.

A few details in the form of the key stand in need of special explanation. Parentheses have been used around generic names in the key in two ways; one, to indicate that the genus is better placed elsewhere in the key, and two, with an equality sign, to indicate a synonym for the accepted name of a genus. An example of this second use is given by the first genus entry in the key. The symbols ♀ for female, ♂ for male, and J for young or immature specimens are used throughout the key as a means of saving space; "♂ only," for instance, indicates that only the male form is known. The % sign is used to mean per-

cent of body length, or, distant from anterior end, measured in percent of body length.

In general the intention has been to include all synonyms, together with the corresponding accepted generic names. The case of *Di-chromadora* and *Trichromadora* Kreis, 1929, is an exception; these designations seemed to cut across the accepted classification in such a way as to make their inclusion impracticable.

Through the much appreciated courtesy of the United States Bureau of Fisheries, a part of the work on the key had been carried out at its Laboratories at Woods Hole, Mass. Needless to say, the work could not have been completed or completion even attempted by us without the resources and the expert advice of the Division of Nematology, of the Bureau of Plant Industry, U. S. Department of Agriculture, within which the work has been done. Dr. G. Steiner and Dr. J. R. Christie have given never-failing aid. Dr. Maurice Hall and Dr. B. Chitwood of the Division of Zoölogy have also given generously of their time, their knowledge and their experience.

MARGARET V. COBB.

ABBREVIATIONS USED IN KEY

abs, absent	lab, labial	0 absent, none
am'g, among	lat, lateral	+ or more, plus
alt, alternative	long, longitudinal	± more or less, somewhat, about
amph, amphid	mand, mandibular	% per cent of body length
ant, anterior	med, median	♂ male
card, cardiac	no, number	♂ only only the male known
caud, caudal	nr, near	♀ female
ceph, cephalic	oes, oesophagus	'♀ ovaries 2, reflexed
cerv, cervical	oesoph, oesophagus	-♀ ovaries 2, outstretched
circ, circular	onch, onchium	'♀ ovary 1, anterior, reflexed
conspic, conspicuous	opp, opposite	'♀ ovary 1, posterior, reflexed
constrict, constriction	papill, papillae, papilloid	-♀ ovary 1, anterior, outstretched
cutic, cuticular	phar, pharynx, pharyngeal	♀ ovary 1, posterior, outstretched
cyland, cylindroid	pharyng, pharyngeal	
dents, denticles	post, posterior	Bast, 'Bastian
devel, developed	pres, present	Bütsch, Bütschli
diam, diameter	prob, probably	Clap, Claparède
exc, except	prs, pairs	Dies, Diesing
excret, excretory	reg, region	Ditlev, Ditlevsen
ext, external	set, setae	Duf, Dufour
gub, gubernaculum	stri, striae	Duj, Dujardin
gubernac, gubernaculum	subceph, subcephalic	Ehrenb, Ehrenberg
inconspic, inconspicuous	sublat, sublateral	Fil, Filipjev
indef, indefinite	submed, submedian	Leuck, Leuckart
intest, intestine	term, terminal	Linst, von Linstow
irreg, irregular	undev, undeveloped	Metsch, Metschnikoff
J, young, juvenile		Micol, Micoletzky
junction, junction		Strass, zur Strassen

GLOSSARY

adhesion tubes, hollow tube-like setae by means of which some nemas "walk" along a surface
 ambulatory setae, setae, sometimes hollow and tube-like, by means of which some nemas "walk" along a surface
 amphid, a paired lateral sense organ the opening of which is often conspicuous, near the anterior end
 amphidial opening, the pore (or larger aperture) through the cuticle at which amphid opens exteriorly
 amphidial pocket, a chamber, outline often cup-shaped, sometimes present just behind external amphid
 ampulla, widening in canal, forming a reservoir
 annulated, having annules
 annules, definite transverse rings of the cuticle
 apophysate, having an apophysis
 appendicule, a large single ventral pre-anal supplementary organ which is extensible
 bursa, flap-like extensions of the lateral cuticle of the tail end of the male
 calvarium, subcuticular cephalic framework
 cardiac bulb, swelling of oesophageal wall, glandular or muscular, just anterior to beginning of intestine
 cardiac valve, thickening or complication of oesophageal lining in cardiac bulb, functioning as a valve
 caudal glands, 3 cells in or near tail, emptying by separate ducts into a common ampulla at spinneret
 cephalated, having the head or anterior end set off in some way
 cephalic setae, setae of the second or "outer" cirlet around the mouth
 cephaloboid, resembling (that of) the genus *Cephalobus*
 cervical, on the neck
 chromadoroid, resembling (that of) the genus *Chromadora*
 cirri, elaborate cephalic appendages in front of the cephalic setae
 cuirasse, a helmet-like appearance of cuticle of head
 cyatholaimoid, resembling (that of) the genus *Cyatholaimus*
 demanian system, gonenteric system of vessels connecting intestine and uteri with each other, and posteriorly with the exterior
 dentate, bearing teeth
 denticles, minute teeth or "prickles," usually numerous
 denticulate, bearing denticles
 diplogastroid, resembling (that of) the genus *Diplogaster*
 dorylaimoid, resembling (that of) the genus *Dorylaimus*
 onchellid, male form such as has been classified in the genus *Enchelidium*, male of *Symplocostoma* group
 eurystomoid, resembling (that of) the genus *Eurystomina*
 excretory duct, canal leading from renette to ampulla near excretory pore
 excretory pore, opening through cuticle (usually cervical or cephalic), at which the renette empties
 external amphid, the cuticular manifestation of the amphid
 glottoid apparatus, valvular structure at base of pharynx
 gubernaculum, grooved cuticularized piece, sometimes paired, through which the spicula slide
 head, portion of nema anterior to base of mouth cavity
 jaws, cuticular framework around the mouth, for grasping and holding; fundamentally 3-parted
 labial setae, setae of the "inner" cirlet, on the lips or close to the mouth
 male supplements, male organs, usually pre- or post-anal, usually a single ventral row, or paired, subventral
 mandibles, hard, strong, biting or grasping organs around the mouth
 mandibular, of the mandibles, resembling mandibles
 median bulb, swelling of the oesophageal wall at or near the middle of its length
 monospiral, spiral of one wind or not much more
 monospire, spiral, or a spiral, of one wind or not much more
 mucron, a knob-like ending, shaped like that on the end of a lemon
 multispiral, spiral of two or more winds
 multispire, spiral, or a spiral, of two or more winds
 neck, portion of nema anterior to base of oesophagus
 odontium (odontia), labial tooth (teeth)
 oesophagus, portion of alimentary tract between pharynx and intestine, sometimes surrounding pharynx
 onchium (onchia), pharyngeal tooth (teeth)
 "palps," special labial appendages, apparently for touching, grasping, or getting material to mouth
 percent, percent of body length, or distant from anterior end measured in percent of body length
 pharyngeal bulb, muscular swelling of the oesophageal wall around the pharynx
 pharynx, the mouth cavity and its walls
 phasmids, a pair of cuticular pores on the tail
 plectoid, resembling (that of) the genus *Plectus*
 posterior bulb, cardiac bulb, q.v.
 probolae, prominent and elaborate specialized appendages encircling the mouth
 pseudonchs, structures in pharynx which from some points of view resemble onchia
 renette, a cell or group of cells emptying through the excretory duct(s)
 rhabditoid, resembling (that of) the genus *Rhabditis*
 sensory papilla, a structure in the amphidial area in which apparently nerves terminate
 setae, hair-like structures on the cuticle
 sole, the region of attachment of the special ambulatory setae or adhesion tubes
 somatic setae, setae general on the surface of the body (opposed to cephalic, caudal, etc. setae)
 spear, a long, rather slender onchium, in adult usually axial
 spicula, male intromittent organs, often paired each an elongate cuticularized framework, extrusible through anus
 spinneret, outlet, usually terminal, of the caudal glands, enabling nema to attach itself
 striae, fine transverse lines in the cuticle
 striate(d), showing striae
 stylet, a long slender spear
 subcephalic setae, setae on the head but behind the cephalic cirlet
 supplementary organs, see male supplements
 tooth, element of buccal armature,—onchium, odontium, spear etc.
 tylenchoid, resembling (that of) the genus *Tylenchus*
 uvette, rosette or group of cells between uterine afferent duct and efferent duct of demanian system
 vestibule, entrance to the mouth cavity, sometimes a distinct chamber
 wings, longitudinal structures in cuticle, or projecting from it, usually lateral, but sometimes numerous and evenly spaced around the nema

KEY

PHARYNX ABSENT, or so obscure as easily to escape notice (for alternative see page 8)

Oesophagus with median or posterior bulb or swelling, or both (for alternative see next page)

Amphid not known or obscure

Female known; wings 0

Ovary 1, anterior; ceph. set. 0; oesoph. with med. bulb; spinneret 0; bursa pres.; in soil. *Iotonchium* Cobb 1920
(= *Hemicycliophora* de Man 1921)

Ovaries 2, reflexed

Spinneret none; amphid a pore at lips; cepn. set. 2, minute; in beetle larvae... (*Neoaplectana* Steiner 1920)

Spinneret present; oesoph. without median bulb; bursa none; marine

Cephalic setae 0; cuticle not annulated; spicula long, 18%, flexible... *Molgolaimus* Ditlev. 1921

Cephalic setae numerous, in circlets of 4; cuticle annulated, exc. on head... (*Stilbonema* Cobb 1920)

Female not known

Hemispherical cephalic cap suddenly wider than neck, edge set with dots... *Mitrephorus* Linst. 1877

Hemispherical cephalic cap absent

Spinneret absent; cephalic setae 0; wings present; in soil; J... *Litonema* Cobb 1920

Spinneret present (Nuadella ?); ceph. set. present; wings 0, striae uninterrupted; marine

Cuticle not annulated, striae none; cephalic setae small

Neck 8%; cardiac swelling slight, clavate... *Nuadella* Allgen 1927

Neck under 4%; card. bulb broad; ceph. set. in 3 circlets of 4; amph. small, near lips... (*Catanema* Cobb 1920)

Cuticle annulated, exc. on head; amphid near lips

Setae on head, labial 6, ceph. 10; contour of nema ± serrate; pores lateral... (*Leptonemella* Cobb 1920)

Setae on head in circlets of 4; contour of nema crenate; pores 0; nema 10 mm... (*Lazonema* Cobb 1920)

Amphid known

External amphid not spiral, circular nor elliptical

Head a hemispherical cap, suddenly wider than neck, edge set with dots... *Mitrephorus* Linst. 1877

Head not suddenly wider nor cap-like

Habitat soil about roots

Spinneret present; lips 6, long, revolute, flower-like; amph. semicircular; wings 2; '♀' *Anthonema* Cobb 1906

Spinneret absent; lips not long nor revolute; cephalic setae 0

Wings pres. oesoph. ± cephaloboid; striae ± coarse; amph. transverse, ½ head-width; '♀' *Iotalaimus* Cobb 1920

Wings absent, striae not interrupted

Ceph. papill. 6, conspic. amph. crescentic, head-width back; tail 3%, rounded; ♂ only *Bolbinium* Cobb 1920

Ceph. papill. tiny; amph. inconspic. 2 head-widths back; tail 6-12%, slender, conoid *Alaimus* de Man 1880

Habitat marine; cephalic setae and spinneret present

Ambulatory & body setae absent; nema not crooked

Wings pres.; amph. shepherd's crook; annules under 300, of tile-like elements; '♀' *Ceramonema* Cobb 1920

(cf. *Pselionema* Cobb 1933)

Wings 0; amph. a transverse slit, at lips; pores lateral; nema ± 2 mm.; ♂ only... *Leptonemella* Cobb 1920

Ambulatory & body setae pres.; amph. crook-shape to spiral; nema ± S-shaped; '♀'

Oesoph. region hardly swollen; card. bulb faint; cerv. striae alike... *Notochaetosoma* Irwin-Smith 1918

Oesoph. region swollen, ovoid; oesophageal bulbs 1 or 2; band of cervical

striae accentuated, exc. *Tristicochaeta falcatum*

Ventral ambulatory adhesion tubes or setae in 3 or 4 rows... *Tristicochaeta* Panceri 1878

(= *Draconema* Cobb 1913)

Ventral ambulatory adhesion tubes or setae in 2 rows... *Drepanonema* Cobb 1933

(= *Chaetosoma* Clap. 1863)

External amphid spiral, circular or elliptical

Female not known; J only, exc. *Bolbinium*

Annules ± 90, prominent; subdorsal setae 9 pres.; amph. saccate; nema 10%+ wide *Eudemoscolex* Steiner 1916

Annules 0 or not prominent; subdorsal setae 0; amphid not saccate; nema under 5% wide

Spinneret absent; cephalic setae 0; in soil

Cephalic papillae 6, conspicuous, each on circular base... *Bolbinium* Cobb 1920

Cephalic papillae 0 or inconspicuous; wings double; oesophagus ± cephaloboid... *Litonema* Cobb 1920

Spinneret present

Wings single, projecting; amphid almost neck width, barely longer than broad... *Antomicron* Cobb 1920

Wings 0, striae not interrupted laterally

Pharynx with basal cuticularized ring; amphid monospire... (*Cricolaimus* Southern 1914)

Pharynx without basal cuticularized ring; cephalic setae in circlets of 4

Amphid spiral, of 2 winds... (*Alaimonema* Cobb 1920)

Amphid circular or elliptical

Head set off by constriction, spherical; cephalic setae 4; amphid on neck... *Cinctonema* Cobb 1920

Head set off by lack of annules, if at all; ceph. set. many; amph. small, at lips; neck 2%

Cuticle without annules; head not set off; pharynx definite; nema 3-4 mm... (*Catanema* Cobb 1920)

Cuticle with annules exc. on head; pharynx obscure; nema 10 mm... *Lazonema* Cobb 1920

Female known

Ovary 1; spinneret none; wings 0; cephalic setae 0 or papilloid

Gonad ♀ posterior, reflexed; amphid ± obscure; cardiac swelling slight; in soil... (*Alaimus* de Man 1880)

Gonad ♀ anterior, outstretched

Habitat marine; anterior oesophagus fusiform, cardiac bulb massive... (*Solenolaimus* Cobb 1894)

Habitat soil; oesophagus with median bulb; bursa present... *Iotonchium* Cobb 1920

(= *Hemicycliophora* de Man 1921)

Ovaries 2

Gonade ♀ outstretched; cephalic setae present; marine (for alternative see next page)

Spinneret doubtful or absent; striae very fine or seen with difficulty

Cephalic setae 10, subcephalic 4; amphid multispire... (*Laimella* Cobb 1920)

Cephalic setae 4 or 8; amphid circular

Pharynx none; cardiac bulb broad, valvate; tail setaceous... *Terschellingia* de Man 1888

Pharynx obconoid; cardiac bulb pyriform, sometimes valvate... (*Cryptolaimus* Cobb 1933)

Spinneret present

Length of oval monospiral amphid twice its width, equalling head width... *Disconema* Fil. 1918

Length of amphid not much greater than its width

Gland ducts cuticularized, pores ventrad, cerv. & pre-anal; striae coarse *Halaphanolaimus* Southern 1914

Gland ducts and pores 0; striae fine, usually plain

- Oesophagus narrowed from middle; cephalic setae 4; amphid large; head narrowed *Cyartonema* Cobb 1920
 Oesophagus not narrowed; cephalic setae present; wings 0
 Setae on head 10, 4 short specialized at tip; cardiac swelling slight.....(*Linhomoella* Cobb 1920)
 Setae on head 4-8, less than 1 head-width long, tip not specialized; amphid circular
 Pharynx 0; nema small, tapered; card. bulb broad, valvate; tail setaceous.. *Terschellingia* de Man 1888
 Pharynx present though obscure
 Ceph. set. 8-10; card. swelling none or clavate, non-valvate; nema not tapered *Linhomoetus* Bast. 1865
 Ceph. set. 4; cardiac bulb pyriform, sometimes valvate; pharynx obconoid (*Cryptolaimus* Cobb 1933)
 Gonads ♀ reflexed; amphid circular, oval or monospiral (for alternative see preceding page)
 Spinneret none; amphid a pore at lips; ceph. set. 2, minute; in beetle larvae... (*Neocaplectana* Steiner 1929)
 Spinneret present; marine exc. *Haliplectus*
 Ambulatory setae 0; nema not crooked; calvarium absent, or not truncate-conoid
 Med. oes. bulb pres.; ceph. set. 0; pores in submed. rows; water & brackish soil.. (*Haliplectus* Cobb 1913)
 Med. oes. bulb absent; marine
 Gland ducts cuticularized, pores cerv. & pre-anal; striae coarse... (*Halaphanolaimus* Southern 1914)
 Gland ducts and pores none or not cuticularized
 Head expanded opposite monospiral amphid; cephalic setae 4..... *Bolbonema* Cobb 1920
 Head not expanded
 Cephalic setae 0; wings 0; head narrowed from amphid forward..... *Aegialolaimus* de Man 1907
 Cephalic setae present; often in coastal sand
 Amphid oval, 1 head-width long; annules fine; wing single; ceph. set. 4.. *Eutelolaimus* de Man 1922
 Amphid ±equidiametral, less than ½ corresponding neck width
 Neck 21%; striae duplex; wings 2; ceph. setae 4; pharynx narrow, inconspicuous *Apletus* Cobb 1914
 Neck 7% or less; striae not duplex; wings 0
 Cuticle annulated; amph. small, at lips; set. on head many, in circles of 4.. (*Stilbonema* Cobb 1920)
 Cuticle finely striated; amphid ½ to ¾ head-width, often monospiral
 Pharynx straight, closed, with pharynx swelling; nema 4-7 mm.; width 0.3-0.8% *Lazus* Cobb 1894
 Pharynx conoid, often with small onchium; nema 2-4 mm.; width 1.5-3.5% *Spirina* Fil. 1918
 (= *Spira* Bast. 1865)
 Ambulatory set. forming a ventral sole, midbody or pre-anal; nema crooked; calvarium unstriated
 Nema narrowest in cardiac region, ±S-shaped; ambulatory setae hollow, pre-anal; annules finer
 Oesoph. region hardly swollen; card. bulb faint; cervical striae alike... *Notochaetosoma* Irwin-Smith 1918
 Oesoph. region swollen, ovoid; oesophageal bulbs 1 or 2; band of cervical
 striae accentuated, exc. *Tristicochaeta* falcatum
 Ventral ambulatory adhesion tubes or setae in 3 or 4 rows..... *Tristicochaeta* Panceri 1878
 (= *Draconema* Cobb 1913)
 Ventral ambulatory adhesion tubes or setae in 2 rows..... *Drepanonema* Cobb 1933
 (= *Chaetosoma* Clap. 1863)
 Nema narrowest midway, epsilon-shaped; ambulatory setae not open at end; annules coarser
 Sole apparently absent; amphid crescentic; annules 98..... *Epsilonella* Steiner 1930
 Sole present; amphid circular or monospiral
 Oesophagus with median and cardiac bulbs..... *Melepsilonema* Steiner 1927
 Oesophagus with cardiac bulb only
 Annules 80-112; ambulatory setae slender, falcate..... *Bathyepsilonema* Steiner 1927
 Annules 144-192; ambulatory setae straight or slightly arcuate
 Body cylindroid, if crooked faintly so..... *Archepsilonema* Steiner 1927
 Body epsilon-like, i.e. body arches well developed
 Cuticular annules homogeneous in structure..... *Prochaetosoma* Baylis & Daubney 1926
 (= *Rhabdogaster* Metsch. 1867)
 Cuticular annules not homogeneous in structure
 "Axial" part of the cuticular annules hollow ("frame rings")..... *Epsilonema* Steiner 1927
 "Axial" part of the cuticular annules vacuolated, etc..... *Epsilonoides* Steiner 1931
- Oesophagus plain, i.e. without median or posterior bulb or swelling (for alternative see preceding page)**
Amphid not known or obscure (for alternative see next page)
 Female not known
 Habitat fresh water marshes; setae 0; spinneret none; neck 15-25%; wings double *Macroposthonia* de Man 1880
 Habitat marine
 Neck 8%, cylindroid; cephalic setae 4(?); tail cylindroid..... *Nuadella* Allgen 1927
 Neck 16-43%, ±tapering; tail conoid
 Spinneret absent; eyes 0; neck ±40%; cephalic setae 4 or 8..... (*Halalaimoides* Cobb 1933)
 Spinneret present; eyes with lenses; neck 16-25%
 Cephalic setae 4; head set off by cessation of very fine striation..... *Nemella* Cobb 1920
 Cephalic setae 10; striae none; amphid and setae very large; adult ♂ only..... *Enchelidium* Ehrenb. 1836
 (= *Lastomitus* Marion 1870
 = *Parasymplocostoma* Schulz 1932)
- Female known**
 Ovary 1, anterior exc. Antopus & Thalassolaimus (for alternative see next page)
 Gonad ♀ outstretched; setae 4, papilloid; amphid minute, open caudad; marine algae... (*Litotes* Cobb 1920)
 Gonad ♀ reflexed; spinneret none, exc. Antopus & Thalassolaimus
 Cephalic setae 8; posterior ovary longer than anterior; marine
 Pores in dorso-lateral rows..... *Antopus* Cobb 1933
 Pores none..... *Thalassolaimus* de Man 1893
 Cephalic setae 0; vulva near anus; spicula 2; in and around insects
 Spear none or vestigial
 Uterus not evaginated
 Nema serpentine; oesophagus and intestine persisting; in beetles and weevils... *Bradynema* Strass. 1892
 Nema saccate; organs degenerating exc. gonads..... *Allantonema* Leuck. 1884
 (= *Tylenchomorphus* Fuchs 1914)
 Uterus finally evaginated; oesophagus and intestine degenerating early
 Evaginated uterus many times as large as nema..... *Sphaerularia* Duf. 1837
 Evaginated uterus not much larger than nema..... *Aconema* Leuck. 1886
 (= *Atractonema* Leuck. 1887)
- Spear present, at least in larva or ♀; organs degenerating in adult
 Base of spear ±bulbed; oes. fusiform, glands abs. in ♂; bursa pre- and post-anal.. *Seatonema* Bovien 1932
 Base of spear straight; spicula nearly straight
 Vulva degenerating; ♀ free living; bursa pre-anal; in Passalus.. *Chondronema* Christie & Chitwood 1931
 (= *Uracanthus* Dies. 1861)

- Vulva remaining functional; adult ♀ parasitic; gubernaculum small
 Anus none or vestigial; in cucumber beetle, *Diabrotica*.....*Howardula* Cobb 1921
 Anus persisting; in frit-fly, *Oscinella*.....(*Tylenchinema* Goodey 1930)
Ovaries 2; spinneret present exc. Micoletzkyia (for alternative see preceding page)
 Gonads ♀ parallel, posterior; wings 10-12, segmenting the annules.....*Porocoma* Cobb 1920
 Gonads ♀ opposed
 The ovaries outstretched; marine
 Annules few, very coarse; amphid saccate; cephalic setae 4; nema wide
 Number of annules ± 17 , secreted annules prominent, separated; head small...*Desmoscolex* Clap. 1863
 Number of annules 32-91, secreted annules absent; head concave-quadrate pyramidal, *Tricoma* Cobb 1894
 (= *Quadricoma* Fil. 1922)
 Annules and striae 0; amphid not saccate; cuticle thick.....*Phanodermella* Kreis 1928
 The ovaries reflexed
 Habitat freshwater; lips 3, thick; pharynx narrow, deep; onchium minute, deepset (*Tripyla* Bast. 1865)
 (= *Promononchus* Micol. 1923)
Habitat marine; pharynx and onchium absent
 Annules ± 17 , secreted annules prominent; amphid saccate; cephalic setae 4; nema wide *Desmoscolex* Clap. 1863
 Annules not few nor coarse; amphid not saccate; nema \pm slender
 Odontia 6; lip region discoid; cephalic setae 4; gland ducts oft piercing cuticle. *Stephanolaimus* Ditlev. 1918
 Odontia 0; lip region not set off
 Head set off by constrict.; cephalic setae 8; spicula long, slender; gub. complex *Micoletzkyia* Ditlev. 1926
 Head not set off; cephalic setae 4 or 6
 Setae on head 4; eyes with lenses; amphid transverse-oval, at lips; among algae...(*Ionema* Cobb 1920)
 Setae on head 6; eyes none
 Spicula short, 1 anal body-diam.; pores 0; longest cerv. set. $\frac{1}{2}$ neck-width. *Parozystomina* Micol. 1924
 Spicula long, several anal body-diameters
 Cervical setae in 6 longitudinal rows; pores 0; ovaries equally long.....(*Stenolaimus* Marion 1870)
 Cervical setae 0; pores in dorso-sublat. rows; ovary post., anterior a rudiment. *Antopus* Cobb 1933
***Amphid known* (for alternative see preceding page)**
 External amphid not spiral, circular nor elliptical (for alternative see next page)
 Female not known; marine
 Neck $\pm 40\%$; amph. narrow, several head-widths long; tail setaceous; spinneret oft 0 *Halalaimoides* Cobb 1933
 Neck not over 25%; amphid not linear; spinneret present
 Wings absent
 Odontia 6; lip papill. 6; cephalic setae 10(?); subceph. 4, jointed; amph. oval, flat caudad *Apodontium* Cobb 1920
 Odontia 0; mouth & phar. vestigial; eyes, amph. & set. very large; adult ♂ only *Enchelidivum* Ehrenb. 1836
 (= *Lasiomitus* Marion 1870
 = *Parasymphocostoma* Schulz 1932)
Wings conspicuous
 Head mitreform, set off by constriction; striae 0; cephalic setae small, 4 or 6.....*Xennella* Cobb 1920
 Head not mitreform, set off by lack of annulation; annules of "tiles"; amph. crook-shape
 Annules over 700; cephalic setae conspicuous, 8, in 2 circlets; onchia 0; nema 3 mm. + *Pristionema* Cobb 1933
 Annules under 300; cephalic setae 4; onchium present (?); nema under 1 mm.....*Pselionema* Cobb 1933
 (= *Steinera* Fil. 1922)
 (cf. *Ceramonema* Cobb 1920)
- Female known**
Ovary 1
 Gonad ♀ outstretched, anterior; cephalic papill. 4; amph. minute, open caudad; marine algae *Litotes* Cobb 1920
 Gonad ♀ reflexed
 Spinneret present; ovary posterior, anterior branch short; wings 0; marine *Thalassolaimus* de Man 1893
 Spinneret absent; ovary anterior; in soil about roots
 Wings present, striae \pm coarse; cephalic papillae 4; amphid 1 head-width back...*Totalaimus* Cobb 1920
 Wings 0, striae 0; cephalic papillae minute; amphid 2 head-widths back.....*Alaimus* de Man 1880
Ovaries 2
 Gonads ♀ outstretched; spinneret present; marine
 Annules ± 17 , secreted annules prominent; amphid saccate; cephalic setae 4; nema wide *Desmoscolex* Clap. 1863
 Annules not few nor coarse; amphid not saccate; nema \pm slender
 Mouth ventrad; amphid huge, bent, dorsal limb the longer.....*Campylaimus* Cobb 1920
 Mouth axial; amphid not huge
 Amphid narrow, oft many head-widths long; eyes 0; cephalic setae 6, subceph. 4 *Halalaimus* de Man 1888
 Amphid "folded" not over 1 head-width long; eyes 2, sometimes absent
 Oesophagus and lumen widened between eyes and nerve ring; pharynx obvious (*Coinonema* Cobb 1920)
 Oesophagus and lumen not widened; pharynx like oesoph. lining...*Araeolaimoides* (de Man) Fil. 1918
 Gonads ♀ reflexed
 Spinneret 0; tail 3%; cephalic setae 10, cervical long, 1 group at amphid; marine...*Platycomopsis* Ditlev. 1926
 (= *Dactylonema* Fil. 1927)
Spinneret present
 Habitat freshwater; lips 3, thick; pharynx narrow, deep; onchium minute, deepset (*Tripyla* Bast. 1865)
 (= *Promononchus* Micol. 1923)
Habitat marine; onchia absent exc. Actinonema
 Amphid narrow, several head-widths long; head set off, with colorless cuticle
 Cephalic & subceph. setae ± 1 head-width apart; cuticle thin (to $\frac{1}{2}$ radius)...*Halalaimus* de Man 1888
 Cephalic & subceph. setae near together exc. *Nuada* isaitshikovi; cuticle thick, $\frac{1}{2}$ to $\frac{1}{3}$ radius
 The cuticle $\frac{1}{2}$ to $\frac{1}{3}$ radius in thickness.....*Nuada* Southern 1914
 The cuticle $\frac{1}{2}$ to $\frac{1}{3}$ radius in thickness.....*Tychnodora* Cobb 1920
 Amphid not narrow nor very long
 Shape of amphid a shepherd's crook; each annule of tile-like elements.....*Ceramonema* Cobb 1920
 (cf. *Pselionema* Cobb 1933)
 Shape of amphid not a shepherd's crook; annules not of tile-like elements
 Annules ± 17 , secreted annules prominent; amph. saccate; cephalic setae 4; nema wide *Desmoscolex* Clap. 1863
 Annules not few nor coarse; amphid not saccate; nema \pm slender
 Cuticle $\frac{1}{2}$ radius; striae coarse; amphid transverse, $\frac{1}{2}$ head-width; setae 0...(*Actinonema* Cobb 1920)
 Cuticle not thick; amphid not a transverse slit; eyes with lenses; cephalic setae 4
 Amphidial gland large, conspicuous; cephalic setae 4, $\frac{1}{2}$ head-width long...*Ionema* Cobb 1920
 Amphidial gland not evident; external amph. \pm semicircular, internal elongate *Nemella* Cobb 1920

- External amphid spiral, circular or elliptical (for alternative see preceding page)
- Female not known; spinneret present
- Habitat brackish earth; wings faint; ♂ supplements on neck, ventral, papiloid *Deontolaimus* de Man 1880
- Habitat marine
- Annules ± 90 , prominent; subdorsal setae 0 prs.; amph. saccate; nema 10% + wide *Eudesmoscolex* Steiner 1916
- Annules none or not prominent; subdorsal setae 0; amphid not saccate; nema under 5% wide
- Wing small, distinct; amphid almost neck width, joined dorsad..... (*Antomicron* Cobb 1920)
- Wings absent
- Mouth and digestive system vestigial; setae strongly developed
- Eyes absent..... *Barbonema* Fil. 1927
- Eyes and amphid strongly developed; adult ♂ only..... (*Enchelidium* Ehrenb. 1836)
(= *Lasiomitus* Marion 1870
= *Parasymphlocostoma* Schulz 1932)
- Mouth and digestive system functional
- Odontia 6, minute; cephalic setae 10(?), subcephalic 4, jointed..... *Apodontium* Cobb 1920
- Odontia 0; cephalic setae 0 or rather small
- Annules rather coarse; setae minute; amphid transverse-oval..... (*Zygonemella* Cobb 1920)
- Annules and striae none; amphid with internal pocket
- Cephalic setae 0; amphid long-oval, internal pocket conspicuous..... *Schistodera* Cobb 1920
- Cephalic setae small, 10; cervical setae lateral; amphid small, round..... (*Leptosomatides* Fil. 1918)
- Female known
- Ovary 1; amphid not multispire
- Gonad ♀ outstretched
- The ovary posterior; spinneret present
- Amphid large, open caudad; spinneret sometimes doubtful; marine..... (*Alaimella* Cobb 1920)
- Amphid round; anterior ovary rudimentary; ceph. set. 12, in 2 circllets; in salt marsh *Litinium* Cobb 1920
- The ovary anterior; marine
- Pharynx absent; setae 4, papiloid; amph. small, open caudad; spinneret 0; among algae *Litotes* Cobb 1920
- Pharynx present, often minute; setae not papiloid; amphid larger, circular; spinneret sometimes 0
- Onchia 0; nema small..... *Monhystera* Bast. 1865
(= *Tachyhodites* Bast. 1865)
Linhomoetus Bast. 1865
(= *L. obtusicaudatus* de Man 1907)
- Onchium dorsal; nema often several millimetres long.....
- Gonad ♀ reflexed
- Habitat soil or fresh water, or in insects; amphid circular
- The ovary posterior; spinneret absent; in marshy soil..... *Alaimus* de Man 1880
- The ovary anterior
- Spinneret present; in fresh water..... *Helalaimus* de Cillis 1917
- Spinneret absent; in beetle, *Passalus*..... *Chondronema* Christie & Chitwood 1931
(= *Uracanthus* Dies. 1861)
- Habitat marine; ovary posterior; spinneret present (*Alaimella* ?)
- Cephalic setae 0
- Amphid monospire; in sand about algae..... *Nemanema* Cobb 1920
- Amphid long-oval, with posterior cuticularized pocket..... *Schistodera* Cobb 1920
(cf. *Oxystomina* Fil. 1921)
- Cephalic setae present
- Setae on head 4, over 1 head-width long; striae coarse; amphid large, open caudad. *Alaimella* Cobb 1920
- Setae 10 or 16, 4 being subcephalic; striae 0 or fine
- External amphid circular; cephalic setae 12, as long as head is wide..... *Litinium* Cobb 1920
- External amphid oval, pocket cuticularized; cephalic setae 6 (4?)
- Amphidial opening small transverse-oval, or circular..... *Nemanemella* Fil. 1927
- Amphidial opening large, long-oval..... *Oxystomina* Fil. 1921
(= *Oxystoma* Bütsch. 1874)
- Ovaries 2
- Gonads ♀ outstretched; marine
- Spinneret absent
- Pharyngeal bulb definite; ceph. setae 6, stout, jointed; amphid circular; in sand (*Cytolaimium* Cobb 1920)
- Pharyngeal bulb absent
- Tail 3%, conoid; ceph. set. 10; cervical long, bunched; amphid transverse-oval.... *Platycoma* Cobb 1894
- Tail over 12%, conoid then cylindrical; ceph. set. minute, 18, in 3 circllets..... (*Anticyathus* Cobb 1920)
- Spinneret present
- Gland ducts cuticularized, pores cervical & pre-anal; striae coarse..... (*Halaphanolaimus* Southern 1914)
- Gland ducts and pores absent
- Annules few, very coarse; amphid saccate; cephalic setae 4; nema wide
- Number of annules ± 17 , secreted annules prominent, separated; head small... *Desmoscolex* Clap. 1863
- Number of annules 32-91, secreted annules none; head concave-quadrangle pyramidal *Tricoma* Cobb 1894
(= *Quadricoma* Fil. 1922)
- Annules if present not coarse; amphid not saccate
- Amphid multispire
- Cephalic setae 10, shorter 4 specialized at tips; multispire faint; neck 5%.... *Linhomoella* Cobb 1920
- Cephalic setae without specialized tips; multispire distinct; neck 9-16%
- Male ventral pre-anal supplementary organs papiloid..... *Parasabatieria* de Man 1907
- Male ventral pre-anal supplementary organs absent..... *Sabatieria* de Rouville 1903
- Amphid circular (Southernia slightly irregular)
- Oesoph. & lumen with ellipsoid enlargement behind eyes; eyes sometimes 0 *Araeolaimus* de Man 1888
- Oesoph. without median enlargement; eyes 0
- Cephalic setae 4; amphid slightly irregular..... *Southernia* Allgen 1929
- Cephalic setae 6-10
- Lips distinct; pharyngeal bulb pres.; ceph. set. 6, stout, jointed; papillae 6. *Cytolaimium* Cobb 1920
- Lips 0 or confluent
- Setae on head 10, shorter 4 specialized at tips; multispire faint..... (*Linhomoella* Cobb 1920)
- Setae on head without specialized tips; amphid circular..... *Linhomoetus* Bast. 1865
- Gonads ♀ reflexed
- Spinneret absent; amphid circular, elliptical or monospiral; wings 0 (for alternative see next page)
- Habitat soil; internal amphid \pm conspicuous..... (*Bastiania* de Man 1876)
- Habitat marine
- Tail 40%; vulva 33%; ceph. set. 6, subceph. 6; internal amphid \pm conspicuous.... *Trefusia* de Man 1893

- Tail under 10%; vulva $\pm 60\%$; ceph. set. ± 1 head-width long; internal amphid not conspicuous
 Ceph. set 10, cervical long, bunched, in σ^2 2 flat ones at amphid; tail 3%, conoid. *Platycoma* Cobb 1894
 Ceph. set. 4 (8?), other setae 0; nema cephalated by contraction; tail slender, 9%. *Acoma* Steiner 1916
 Spinneret present (for alternative see preceding page)
 Habitat not marine (*Aphanolaimus* ?); contour often crenate
 Amphid obscure, small; pharynx narrow; onch. deep set; ceph. set. 6-10, oft papilloid. (*Tripylla* Bast. 1865)
 (= *Promononchus* Micol. 1923)
 Amphid distinct, circular or spiral; pharynx none or closed; ceph. set. 6, rarely 4 or 0
 Position of amphid $\frac{1}{2}$ -2 head-widths back. *Aphanolaimus* de Man 1880
 Position of amphid 3-4 head-widths back. *Bastiania* de Man 1876
 Habitat marine
 Ambulatory tubes hollow; body setose, annulated
 The amphid not saccate; ambulatory tubes ventral; setae not dense. *Notochaetosoma* Irwin-Smith 1918
 The amphid saccate; annules coarse, 17-91; nema broad
 Body setae not dense; secondary annules secreted, prominent. (*Desmoscolex* Clap. 1863)
 Body setae dense, set along annules; junction oesoph. & intestine indefinite. (*Greefiella* Cobb 1922)
 (= *Trichoderma* Greeff 1869)
 Ambulatory tubes absent; body not densely setose; not annulated exc. Stephanolaimus
 Wings ± 50 ; nema broad, $\pm 16\%$; neck "collared"; amphid spiral; pharynx small. *Richteria* Steiner 1916
 Wings if present not numerous; nema slender, under 5%
 Odontia 6; ceph. set. 4, long; lip reg. discoid; gland ducts oft projecting. *Stephanolaimus* Ditlev. 1918
 Odontia 0; cephalic setae not longer than head is wide
 Gland ducts cuticularized, pores cervical & pre-anal; striae coarse. *Halaphanolaimus* Southern 1914
 Gland ducts and pores none, or not cuticularized
 The amphid multispire; cephalic setae 16, ± 1 head-width long; in sand. *Nannolaimus* Cobb 1920
 The amphid not multispire
 Amphid at lips, transverse-oval, with large gland; eyes with lenses. *Ionema* Cobb 1920
 Amphid not at lips; eyes none exc. *Leptosomatium*
 Anterior part of oesoph. the wider; cervical setae 0; amphid \pm irregular. *Southernia* Allgen 1929
 Anterior part of oesoph. not wider; amphid with internal pocket
 Oesophagus crenate posteriorly
 Calvarium none; labial tubercles none. *Stenolaimus* Marion 1870
 Calvarium in front of ceph. set., margin irregular; labial tubercles 3. *Klugea* Fil. 1927
 (= *Phanodermopsis* Ditlev. 1926, in part)
 Oesophagus not crenate posteriorly
 Cephalic setae 0; pores on anterior half of neck; tail rounded. *Leptosomatium* Bast. 1865
 Cephalic setae present; pores on neck absent
 Setae on neck none; cephalic setae 10, ± 1 head-width long. *Leptosomella* Fil. 1927
 Setae on neck in lateral series; amphid small
 Gubernaculum present; σ^2 pre-anal ventral supplement tubular. *Anticoma* Bast. 1865
 Gubernaculum none; σ^2 pre-anal ventral supplement papilloid. (*Anticomopsis* Micol. 1930)

PHARYNX PRESENT (for alternative see page 4)

Wall of the Pharynx armed (for alternative see page 20)

- Armature spear-like* or apparently so (for alternative see page 11)
 Spear bulbed; cephalic setae usually 0; amph. oft obscure; spinneret 0; not marine exc. *Siphonolaimus* (pg. 10)
 Oesophagus plain, i.e. without median or posterior bulb or swelling; cephalic setae 0
 Female not known; spear with retrorse points midway, base 3-lobed; bursa lobed. *Ecphyadophora* de Man 1921
 Female known; ovary 1
 Gonad \varnothing reflexed
 Habit free-living; spear (dorso-ventral view) arrow-headed; ovary posterior. *Pharetrolaimus* de Man 1921
 Habit parasitic in beetles, etc.; ovary anterior
 Nema saccate; mouth none, organs degenerated; vulva terminal. (*Allantonema* Leuck. 1884)
 (= *Tylenchomorphus* Fuchs 1914)
 Nema not saccate; anus none or vestigial
 Spear none; parasitic in beetles, weevils, etc. (*Bradynema* Strass. 1892)
 Spear present in young \varnothing only; in cucumber beetle, *Diabrotica*. *Howardula* Cobb 1921
 Gonad \varnothing outstretched; mostly parasitic
 Anus none or vestigial; spear \pm vestigial; in cucumber beetle, *Diabrotica*. (*Howardula* Cobb 1921)
 Anus present; spear present, in *Aphelenchulus* reduced; traces median bulb sometimes present
 Junction oesoph. & intest. definite; anterior oes. fusiform, isthmus at nerve ring; in or about plants
 Habitat crop plants; head of 8 sectors; terminus of tail straight. *Neotylenchus* Steiner 1931
 Habitat bladders of *Fucus*, or among marine algae; head of 6 sectors; tail hooked. *Halenchus* Cobb 1933
 Junction oesophagus and intestine indefinite
 Spear with 6 basal lobes; oesoph. isthmus at nerve ring; in decaying vegetation. *Hexatylus* Goodey 1926
 Spear with tripartite base; renette \pm body length; parasitic in insects. *Aphelenchulus* Cobb 1920
 Oesophagus with median or posterior bulb or swelling, or both
 Female not known; spear tripartite at base
 Cephalic setae 4, ± 1 head-width long; oesoph. swellings med. & card. ; bursa striate. *Eutylenchus* Cobb 1913
 Cephalic setae 0 or papilloid
 Spear with retrorse points midway, 3-lobed at base; bursa small, lobed. *Ecphyadophora* de Man 1921
 Spear without retrorse points midway; oesophagus without median bulb
 Pharyngeal bulb 0, cardiac $\frac{1}{2}$ neck length; ceph. set. 0; nema 5% wide; bursa 0. *Tylencholaimellus* Cobb 1915
 Pharyngeal bulb pres.; ceph. set. 6, papilloid; nema 9% wide; tail short, rounded. *Brachynemella* Cobb 1933
 (= *Brachynema* Cobb 1893)
 Female known
 Ovaries 2; cephalic setae 0; spear with tripartite base (for alternative see next page)
 Gonads \varnothing outstretched (for alternative see next page)
 Median oesophageal bulb absent; spear 3-pronged at base; in sandy soil. (*Tyolaimophorus* de Man 1880)
 Median oesophageal bulb present; spear 3-lobed at base
 Junction oesoph. & intestine indefinite; annules plain; tail rounded, with bursa. *Hoplolaimus* Daday 1905
 Junction oesoph. & intestine definite, exc. sometimes *Tylenchus*
 Spear-guide wide as head, forming cuirasse or helmet; bursa none; testes 2. *Nemonchus* Cobb 1913

- Spear-guide smaller, slighter, not forming cuirasse; bursa present
 Nema cephalated by constriction, head \pm discoid; spear long; bursa lobate.... *Dolichodoros* Cobb 1914
 Nema cephalated by contraction or not at all; bursa not lobate
 Anterior end extensible, narrow & beak-like, inner framework cuticular... *Tylenchorhynchus* Cobb 1913
 Anterior end never narrowed & beak-like, without cuticular framework..... *Tylenchus* Bast. 1865
- Gonads ♀ reflexed (for alternative see preceding page)
 Structure of pharynx obscure; oesophagus with cardiac swelling only
 "Spear" 3 rods, ant. hinged to 2 U-shaped pieces, post. 2 stalked swellings. (*Tylopharynx* de Man 1876)
 "Spear" rods 2-3, base heavy, irreg.; setae 0 or papill.; ovary obscure... *Diphtherophora* de Man 1880
 (= *Chaolaimus* Cobb 1893
 = *Archionchus* Cobb 1913)
- Structure of pharynx obvious
 Oesophagus without median bulb, cardiac bulb present
 Cardiac swelling cylindroid (i.e. oesophagus dorylaimoid)
 Oral spear large, over 5%, base 3-bulbed but not trifurcate..... *Xiphinema* Cobb 1913
 Oral spear smaller, under 5%, base trifurcate..... (*Tylencholaimus* de Man 1876)
 Cardiac swelling pyriform to clavate
 Spear 3-bulbed at base, \pm stout; amphid protrusile, tube cuticularized.... *Triplonchium* Cobb 1920
 Spear 3-pronged; amph. $\frac{1}{2}$ head-width, transverse-oval; labial papillae 6 *Tyololaimophorus* de Man 1880
 Oesophagus with median bulb, often with cardiac swelling also
 Body \pm thick, not serpentine; ♀ and larval forms..... *Heterodera* Schmidt 1871
 (= *Meloidogyne* Göldi 1887
 = *Caconema* Cobb 1924)
- Body \pm serpentine
 Junction oesoph. & intestine indef.; annules plain; tail rounded, with bursa. (*Hoplolaimus* Daday 1905)
 Junction oesoph. & intestine definite; bursa none; ♂ and larval forms
 Spear really the apophyses of 3 movable onchia at base anterior pharynx... *Tylenchodon* Fuchs 1930
 Spear tylenchoid..... *Heterodera* Schmidt 1871
 (= *Meloidogyne* Göldi 1887
 = *Caconema* Cobb 1924)
- Ovary 1 (for alternative see preceding page)
 Gonad ♀ reflexed; setae 0
 Spear trifurcate half its length, base swollen; oesoph. with cardiac bulb only... *Doryllium* Cobb 1920
 Spear furcate at base only; bulbs usually 3
 Body thick, inert; parasitic ♀ forms
 Nema body swollen, head and neck narrow; vulva 90%; in citrus roots..... *Tylenchulus* Cobb 1913
 Nema sausage-shaped; organs degenerated exc. gonads; vulva term.; in insects *Allantonema* Leuck. 1884
 (= *Tylenchomorphus* Fuchs 1914)
- Body \pm slender or serpentine
 Oral spear \pm vestigial; median bulb \pm reduced; ♂ and larval forms
 Habitat in citrus roots and soil..... *Tylenchulus* Cobb 1913
 Habitat in and around wood-boring beetles..... *Allantonema* Leuck. 1884
 (= *Tylenchomorphus* Fuchs 1914)
- Oral spear well developed
 Median oesophageal bulb elongate; vulva over 70%, ovary anterior; bursa none
 Spear 3-4%; annules 0; median bulb \pm reduced; anus not functional..... *Tylenchulus* Cobb 1913
 Spear 8-23%; annules under 150, retrorse; median bulb oft valvate; vulva 70-95%. *Ogma* Southern 1914
 (= *Iota* Cobb 1913)
- Median oesophageal bulb absent
 "Spear" rods 2-3, base heavy, irreg.; setae 0 or papilloid; vulva 50%... *Diphtherophora* de Man 1880
 (= *Chaolaimus* Cobb 1893
 = *Archionchus* Cobb 1913)
- "Spear" of usual structure, shaft distinct, base tripartite
 Tip of spear tapering, posterior prongs little swollen; vulva 33-72%... *Tylencholaimus* de Man 1876
 Tip of spear (dorso-ventral view) arrow-headed; vulva 25%, ovary post. *Pharetrolaimus* de Man 1921
- Gonad ♀ outstretched
 Habitat marine; "spear" an evertible pharynx lining; amph. round; neck 4% *Siphonolaimus* de Man 1893
 Habitat not marine; median oesophageal bulb present (*Neotylenchus* ?)
 Cephalic setae 4, $1\frac{1}{2}$ head-widths long; longitudinal striae 10; in cranberry bog... *Atylenchus* Cobb 1913
 Cephalic setae 0
 Dorsal oes. gland emptying at med. bulb; junct. oes. & intest. indef.; bursa 0 *Aphelenchoides* Fischer 1894
 (= *Seinura* Fuchs 1931
 = *Parasitaphelenchus* Fuchs 1930
 = *Pathoaphelenchus* (Cobb) Steiner 1931)
- Dorsal oes. gland emptying at base of spear; genera tylenchoid
 Spear under 5%; junction oesoph. & intest. usually definite; nema under 5% wide; bursa present
 Anterior oesoph. with median bulb; head of 6 sectors; tail without mucron... *Tylenchus* Bast. 1865
 Anterior oesoph. funisiform to isthmus; median bulb not definite; in or about plants
 Habitat crop plants; head of 8 sectors; terminus of tail straight..... *Neotylenchus* Steiner 1931
 Habitat bladders of Fucus, & am'g marine algae; head of 6 sectors; tail hooked *Halenchus* Cobb 1933
 Spear 8-24%; oesoph. lining flexible, looping; nema 5%+ wide; bursa 0; vulva 72-94%
 Base of spear bulbed, not fluked; annules over 100, not retrorse; junct. oes. & intest. definite
 Oral area not raised; annulation fine..... *Paratylenchus* Micol. 1922
 Oral area raised, surrounded by "fossa"; annulation rather coarse..... *Procriconema* Micol. 1925
 Base of spear anchor-shaped (fluked); annules under 150, retrorse; junct. oes. & intest. indef.
 Scales, prickles and fringes absent from annules..... *Criconema* Hoffmänner & Menzel 1914
 Scales, prickles or fringes ornamenting the annules..... *Ogma* Southern 1914
 (= *Iota* Cobb 1913)

- Spear plain, i.e. not bulbed at base (for alternative see page 8)
 Oesophagus plain, i.e. without median or posterior bulb or swelling
- Amphid known**
 External amph. reniform; dorsal onch. small, at spear base; ovary ant.; not marine *Enoplocheilus* Kreis 1932
 External amphid circular, elliptical or monospiral
 Spinneret absent; anus reduced; parasitic in beetles
 Pharyng. bulb strong, head swollen; spear 1.5%; junct. oesoph. & intest. indef.; -♀ (*Rhabdonchus* Cobb 1933)
 Pharyng. bulb none; spear vestigial; in beetle, Passalus; '♀'.... *Chondronema* Christie & Chitwood 1931
 (= *Uracanthus* Dies. 1861)
- Spinneret present; marine
 Annules coarse exc. on head; onch. spear-like, axial; body set. in long. rows; -♀ - *Aculeonchus* Kreis 1928
 Annules none or fine; head not expanded; cephalic setae 4
 Striae of dots, larger laterally; amph. multispire; spicula 2-jointed; -♀ - *Dorylaimopsis* Ditlev. 1918
 (= *Xinema* Cobb 1920)
 Striae 0; amph. monospire; spicula entire; junct. oes. & intest. obscured by glands; eyes present
 Spear 0.3%, a thickening of dorsal pharyngeal wall; ovaries 1 or 2, reflexed.... (*Onchium* Cobb 1920)
 Spear more obvious, 1.5%; '♀'..... *Onchulella* Cobb 1920
- Amphid unknown or obscure**
 Female not known; spear a small prod at lips; marine..... *Rhinonema* Allgen 1927
- Female known**
 Spinneret present; ovaries 2; marine
 Calvarium subcuticular; spear long, slender; set. on head long; nema 6 mm. *Thoracostomopsis* Ditlev. 1918
 Calvarium none; head not suddenly narrowed; spear short; ovaries reflexed.... (*Onchulella* Cobb 1920)
 Spinneret absent; ovary 1; not marine
 The female saccate, sausage-shaped; vulva terminal, ovary reflexed; in beetles *Allantonema* Leuck. 1884
 (= *Tylenchomorpha* Fuchs 1914)
- The female ±elongate, usually not slender; vulva posterior, ovary anterior
 Ovary outstretched; anus present; spear a mere prod, reduced in ♂; wings 0... *Iotonchium* Cobb 1920
 Ovary reflexed exc. *Rhabdonchus*; anus none or reduced; parasitic in insects
 Spear long, slender, in muscular bulb, head swollen; junct. oes. & intest. indef. *Rhabdonchus* Cobb 1933
 Spear vestigial, minute or none; pharyngeal bulb absent
 Mouth and spear absent; in beetles, weevils, etc..... (*Bradynema* Strass. 1892)
 Mouth and spear present in young ♀ only; in beetles, frit fly, etc.
 Vulva degenerating; ♀ free-living; bursa pre-anal; in Passalus *Chondronema* Christie & Chitwood 1931
 (= *Uracanthus* Dies. 1861)
- Vulva remaining functional; adult ♀ parasitic; gubernaculum small
 Anus none or vestigial; in cucumber beetle, *Diabrotica*..... *Howardula* Cobb 1921
 Anus persisting; in frit-fly, *Oscinella*..... (*Tylenchinema* Goodey 1930)
- Oesophagus with median or posterior bulb or swelling, or both**
 Amphid not known or obscure
 Spinneret present; cardiac swelling slight; spear a small prod at lips; J..... (*Rhinonema* Allgen 1927)
 Spinneret absent; cephalic setae 0
 Nema parasitic, saccate female; vulva terminal; in beetles, etc.; '♀'..... (*Allantonema* Leuck. 1884)
 (= *Tylenchomorpha* Fuchs 1914)
- Nema free-living**
 Armature a minute labial prod; expanding head suddenly truncate; vulva 88%; -♀ *Iotonchium* Cobb 1920
 Armature an obvious spear
 Spear closed, bent dorsad at tip when extruded; '♀'..... (*Campydora* Cobb 1920)
 Spear open, a hollow passage; not bent at tip
 Lips petaloid, lip reg. expanded, discoid; oesoph. dorylaimoid; spear small; '♀' *Antholaimus* Cobb 1913
 Lips 0; median bulb valvate, with outlets of 3 glands; junction oesoph. & intestine indefinite; -♀
 Gubernaculum and bursa present..... *Aphelenchus* Bast. 1865
 (= *Isonchus* Cobb 1913)
 Gubernaculum and bursa absent..... *Aphelenchoides* Fischer 1894
 (= *Seinura* Fuchs 1931
 = *Parastaphelenchus* Fuchs 1930
 = *Pathoaphelenchus* (Cobb) Steiner 1931)
- Amphid known**
 External amphid spiral, circular or elliptical (for alternative see next page)
 Spinneret present; amphid labial, ± spiral; "spear" filling posterior pharynx; marine
 Width of nema 3.5%; pharyngeal bulb definite; '♀'..... *Onyx* Cobb 1891
 Width of nema 7%; pharyngeal bulb faint; J..... *Oistolaimus* Ditlev. 1921
 (cf. *Onyx* Cobb 1891)
- Spinneret absent; amphid circular or elliptical
 Anus none; setae 0; pharyng. bulb strong; junct. oes. & intest. indef.; in beetles; -♀ *Rhabdonchus* Cobb 1933
 Anus present; nema free-living
 Cephalic setae 10, *Siphonolaimus* sometimes 0; amphid circular
 Habitat marine; "spear" evertible phar. lining; nema 5-10 mm.; neck 4%; -♀ *Siphonolaimus* de Man 1893
 Habitat moist soil; spear tooth-like; nema 0.85 mm.; neck 16%..... *Odontolaimus* de Man 1880
 (= *Neonchus* Cobb 1893)
- Cephalic setae 0**
 Spear bent dorsad at tip when extruded; amphid small, oval, at lips; '♀'..... *Campydora* Cobb 1920
 Spear not bent at tip
 Med. oes. bulb 0; nema ± dorylaimoid; lab. papillae 6, ceph. 6; ovaries 2 (*Tylolaimophorus* de Man 1880)
 Med. oes. bulb valvate; nema ± tylenchoid; papillae 0; amphid oval, near lips
 Junction of oesophagus and intestine ± definite
 Gubernaculum and bursa absent; tail often with mucron; -♀..... *Paraphelenchus* Micol. 1925
 Gubernaculum small; bursa without ribs; tail slender, long-conoid; -♀ - *Psilenchus* de Man 1921
 Junction oesoph. & intestine indefinite; oes. gland outlets at med. bulb; tail short; -♀
 Bursa and gubernaculum present..... *Aphelenchus* Bast. 1865
 (= *Isonchus* Cobb 1913)
 Bursa and gubernaculum absent..... *Aphelenchoides* Fischer 1894
 (= *Seinura* Fuchs 1931
 = *Parastaphelenchus* Fuchs 1930
 = *Pathoaphelenchus* (Cobb) Steiner 1931)

- External amphid a transverse slit, oft at lip reg.; ceph. set. 0; fresh water or soil (from preceding page)
- Ovary 1, reflexed; spinneret 0 (*Oionchus* ?)
- Spear a closed cutting or pricking organ, \pm tooth-like; vulva 55-60%, ovary anterior
- Spinneret (?) present; spear straight..... *Oionchus* Cobb 1913
- Spinneret absent; tip of spear bent dorsad when extruded; amphid elliptical.... *Campydora* Cobb 1920
- Spear open at end, a hollow passage, tip not bent; oesophagus dorylaimoid
- Anterior $\frac{1}{2}$ of oesophagus narrow; cardiac swelling set off by constriction..... (*Doryllium* Cobb 1920)
- Anterior $\frac{1}{3}$ of oesophagus narrow; both parts cylindroid
- Narrow part of oesophagus set off behind by constriction..... *Azonchium* Cobb 1920
- Narrow part of oesophagus \pm confluent with wider posterior part; ♀'..... (*Dorylaimus* Duj. 1845)
- Ovaries 2, reflexed; oesophagus \pm dorylaimoid
- Spear a closed cutting or pricking organ, \pm tooth-like
- The spear long, slender, flexible; cardiac swelling slight, basal, \pm clavate
- Guide rings for spear absent; spear 30-40% of neck length..... *Trichodorius* Cobb 1913
- Guide rings for spear present; spear \pm 20% of neck length..... *Leptonchus* Cobb 1920
- The spear shorter, less slender, less flexible; cardiac swelling cylindroid, $\frac{1}{3}$ - $\frac{1}{2}$ oesophagus
- Tip of ventral spear oblique, ventrad when extruded; nema 6-10 mm..... *Sectonema* Thorne 1930
- Tip of submedian spear straight; nema under 4.5 mm..... *Nyggolaimus* Cobb 1913
- Spear open at end, a hollow passage
- Pharynx with cup-shaped anterior portion
- Cup-shaped portion with 6 ribs; spear with guiding ring..... *Actinolaimus* Cobb 1913
- Cup-shaped portion without ribs; spear without guide ring; lip region discoid. *Antholaimus* Cobb 1913
- Pharynx without cup-shaped anterior portion
- Oesophagus with fusiform "bulb" behind spear..... *Dorylaimellus* Cobb 1913
- Oesophagus without swelling behind spear
- Lip region discoid, much expanded..... *Discolaimus* Cobb 1913
- Lip region usually not discoid, expanded little or not at all
- Labial papillae 0, cephalic 6; circumoral ridge present; vulva 34%..... *Chrysonema* Thorne 1929
- Labial papillae 6, cephalic 6; circumoral ridge absent..... *Dorylaimus* Duj. 1845
- Armature of one or more teeth (odontia or onchia) (for alternative see page 8)**
- Tooth (onchium) 1 (for alternative see page 15)
- Oesophagus plain, i.e. without median or posterior bulb or swelling; amphid occasionally obscure (pg. 13)
- External amphid not spiral, circular nor elliptical (for alternative see next page)
- Female not known
- Habitat soil; spinneret 0; phar. $\frac{1}{2}$ neck-length; onch. outward-acting; oes. dorylaimoid. *Nanonema* Cobb 1905
(= *Cephalonema* Cobb 1893)
- Habitat marine
- Annules coarse, elements tile-like; amphid crook-shape; ceph. set. 4; nema 0.7 mm. *Pselionema* Cobb 1933
(= *Steinera* Fil. 1922)
- Annules not coarse; amphid not crook-shape
- Onchium small. a labial prod; cephalic setae 6..... *Rhinonema* Allgen 1927
- Onchium pharyngeal
- Labial set. 6, cephalic 10; onchium dorsal; nema 0.5% wide; amphid transverse *Trileptium* Cobb 1933
(= *Trilepta* Cobb 1920)
- Labial set. 0, cephalic 6; onchium ventral; nema over 4% wide..... *Doryonchus* Kreis 1932
(cf. *Symplocostoma* Bast. 1866)
- Female known
- Ovary 1; habitat not marine
- Median onchium spear-like, dorsal onchium minute, basal..... *Enoplocheilus* Kreis 1932
- Median spear-like onchium absent; ovary reflexed; Enopliidae
- Pharynx tubular, depth 10 times width; tooth basal; cephalic setae 0..... *Cryptonchus* Cobb 1913
(= *Dillevenia* Micol. 1925)
- Pharynx conoid when open, depth not over 3 times width
- Tooth minute, basal, deep-set; denticles 0; spinneret terminal..... *Trischistoma* Cobb 1913
- Tooth large, subventral; denticles present; other onchia 2; spinneret ventrad (*Mononchulus* Cobb 1918)
- Ovaries 2
- Gonads ♀ outstretched; spinneret present; striae plain; marine..... *Mononcholaimus* Kreis 1924
- Gonads ♀ reflexed
- Spinneret absent; amphidial opening oval or a transverse slit
- Base of tooth without "flukes"; cephalic setae 6; onchium large, acute; in soil. *Onchulus* Cobb 1920
- Base of tooth anchor-shaped, "flukes" dorsad and ventrad..... *Diphtherophora* de Man 1880
(= *Chaolaimus* Cobb 1893
= *Archionchus* Cobb 1913)
- Spinneret present, exc. sometimes Eurystominae
- Habitat not marine; Trilobinae
- Pharynx closed; lips 3; onchium minute, basal, deepset..... *Tripyla* Bast. 1865
(= *Promononchus* Micol. 1923)
- Pharynx open, domed, larger; lips 6; onchia often more than 1..... *Trilobus* Bast. 1865
- Habitat marine
- Striae conspicuous, resolvable into dots or rod-like elements; Chromadoridae
- Amphid at lips, lenticular, oft obscure; cuticle of "basket-work" on neck *Euchromadora* de Man 1886
(= *Graphonema* Cobb 1898)
- Amphid not at lips, of \pm even width, conspicuous, reaching nearly across head
- Cephalic setae 0; annules of obscure rod-like elements; spicula entire..... *Actinonema* Cobb 1920
- Cephalic setae 10; annules on neck of "basket-work"
- Spicula entire; amphid a wide transverse oval..... *Pareuchromadora* Stekhoven & Adams 1931
- Spicula of 2 sections; amphid a narrow slit of even width..... *Rhips* Cobb 1920
- Striae 0, or plain and very fine
- External amph. bent or crook-shaped, plate oval, 1 head-width; "tooth" minute *Diplopeltis* Cobb 1906
(= *Dipeltis* Cobb 1891
= *Discophora* Villot 1875)
- External amphid not crook-shaped, plate absent; Enopliidae
- Head with elaborate, ornate calvarium..... *Deontostoma* Fil. 1916
- Head without calvarium
- Pharynx of single chamber; cephalic setae 10

- Mouth cavity \pm quadrate; onchium minute, close against wall; eyes 0 *Anoncholaimus* Cobb 1920
 Mouth cavity twice as deep as wide; dorsal onch. bent in over pharynx *Asymmetrella* Cobb 1920
 Pharynx of more than 1 chamber; large onchium subventral
 Onchium conoid to base of short stylet; phar. \pm quadrate; σ supplements eurystomid
 Bulbs of oesophagus 5-8, serial, contiguous.....*Bobella* Cobb 1920
 Bulbs of oesophagus none
 Cervical setae long, about 30; σ supplements weakly developed.....*Ledontia* Fil. 1927
 Cervical setae 0 or minute; σ supplements well developed.....*Eurystomina* Fil. 1921
 (= *Eurystoma* Marion 1870
 = *Marionella* Cobb 1922)
- Onchium a long stylet; phar. much deeper than broad; σ an enchelid
 Anterior oesophagus (pharynx ?) narrower, lumen wide.....*Calyptronema* Marion 1870
 (= *Catalaimus* Cobb 1920)
- Anterior oesophagus not differentiated
 Vestibule set off by row of "comma" markings; pharyng. rings 3 *Symplocoostomella* Micol. 1930
 Vestibule not set off by a ring or rows of denticles or markings.....*Symplocostoma* Bast. 1865
- External amphid spiral, circular or elliptical (for alternative see preceding page)
- Female not known; marine
 Amphid multispire, big; phar. ribbed, complex; onch 3; σ supplements chromadoroid *Pomponema* Cobb 1917
 Amphid circular or oval (Rhinonema, Doryonchus?)
 Annules of dots & rods, altered laterally; amph. transverse-oval *Pareuchromadora* Stekhoven & Adams 1931
 Annules none or not ornamented; cephalic cuticle not thickened
 Cephalic setae 6
 Onchium a minute labial prod.....*Rhinonema* Allgen 1927
 Onchium ventral, pharyngeal, slender, lance-like.....*Doryonchus* Kreis 1932
 (cf. *Symplocostoma* Bast. 1865)
- Cephalic setae 8 or 10
 Oesophagus surrounding phar.; onch. opposed by denticles; amph. long-oval *Anazonchium* Cobb 1920
 Oesophagus censing at base of pharynx; denticles 0; amphid smaller, circular; Epnolidae
 Pharynx of 1 chamber; onchium digitate.....*Cophonchus* Cobb 1920
 Pharynx of 2 chambers; onchium a slender, subventral stylet.....*Isonemella* Cobb 1920
- Female known
 Ovary 1, anterior; onchium dorsal
 Striae of dots; pharynx cyatholaimoid; 'Y'.....*Dentatonema* Kreis 1928
 Striae none or plain; pharynx not cyatholaimoid
 Pharynx large, open; onchium large, apex far forward; amphid with pocket *Pseudodilaimus* Kreis 1928
 Pharynx small or closed, onchium basal or deepset, small
 Gonad σ reflexed; striae 0; spinneret small; vulva 80%; in soil.....*Trischistoma* Cobb 1913
 Gonad σ outstretched; onchium dorsal; amphid circular
 The pharynx very small; oesophagus strongly developed.....*Prosphaerolaimus* Fil. 1918
 The pharynx $\frac{1}{2}$ head-width, as deep as wide; onch. central; in warm salt springs.....*Anticyclus* Cobb 1920
- Ovaries 2
 Gonads σ outstretched; spinneret present; marine
 Amphid not known; striae plain.....*Mononcholaimus* Kreis 1924
 Amphid multispire; striae of dots
 Head setose; striae not altered laterally; spicula long; apophysis absent.....*Comesoma* Bast. 1865
 Head less setose; spicula short; gubernaculum with apophysis
 Male ventral pre-anal supplementary organs small, papilloid.....*Parasabatieria* de Man 1907
 Male supplementary organs absent.....*Sabatieria* de Rouville 1903
- Gonads σ reflexed
 Habitat soil or fresh water; cuticle without pores; wings 0
 Amphid multispire; pharynx conoid; spinneret present; σ supplements none *Nannonchus* Cobb 1913
 Amphid elliptical, with internal pocket; pharynx not conoid
 Cephalic setae 0; labial papillae in 2 circlets; spinneret usually present.....*Mononchus* Bast. 1865
 Cephalic set. 6, jointed; labial papillae 1 circlet; spinneret 0; about roots.....*Cyathonchus* Cobb 1933
- Habitat marine and brackish waters
 Nema broad, 16%; wings \pm 50; neck region "collared"; pharynx narrow.....(*Richtersia* Steiner 1916)
 Nema slender, not over 5%; wings 0 exc. *Rhinema* 12
 Amphid multispire; striae of dots; cuticular pores pres.; vestibule with 12 (6 double ?) ribs
 Large σ supplement tubular, anterior to small supplements; nema viviparous *Acanthonchus* Cobb 1920
 (= *Seuratiella* Ditlev. 1918
 = *Seuratiella* Ditlev. 1922)
- Large σ supplement absent, others if tubular small; nema usually oviparous
 Phar. tubular behind onch.; gubernacula not serrate; σ papill. setose *Paracyatholaimus* Micol. 1922
 Phar. cyathiform; gubernacula serrate distally; σ setose papillae none
 Dorsal onchium large, acute, projecting; striae altered laterally.....*Paracanthonchus* Micol. 1924
 Dorsal onchium 0 or not projecting; pharyngeal ribs extending to base
 Tail setaceous; onchium small; σ papillae present or not.....*Longicyatholaimus* Micol. 1924
 Tail not setaceous
 Dots larger laterally; σ supplements tubular; gubernac. oft joined *Paraacanthonchus* Micol. 1924
 Dots not larger laterally; σ tubular supplements 0; gubernacula joined *Cyatholaimus* Bast. 1865
 (= *Necticonema* Marion 1870)
- Amphid circular, elliptical or monospiral
 Wings 12, markings V-shaped; pharyng. bulb stronger dorsad; amphid circular.....*Rhinema* Cobb 1920
 Wings 0; striae plain
 Dorsal pharyng. wall thick, tooth-like; amph. monospire at lips; ceph. set. 4; renette far back
 Oesoph. glands obscuring junct. intest.; dorsal phar. element \pm uniform, slighter *Onchium* Cobb 1920
 Oesoph. glands not prominent; dorsal pharyng. element not uniformly thick, but heavy
 Tooth-like thickening parallel to axis.....*Camacolaimus* de Man 1889
 (= *Digitonchus* Cobb 1920
 = *Acontiolaimus* Fil. 1918)
- Tooth-like thickening bent outward from beginning of vestibule
 Point of "onchium" single.....*Acmaeolaimus* Fil. 1918
 Point of Y-shaped "onchium" duplex.....*Ypsilon* Cobb 1920

- Dorsal pharyngeal wall not thickened; onchium a distinct projection (from preceding page)
 Cephalic setae 20-30; cardiac glands conspicuous
 Male supplementary organs complex; copulatory muscles conspicuous . . . *Xanthodora* Cobb 1920
 Male supplements simple, papilloid; copulatory muscles not conspic. *Acanthopharynx* Marion 1870
 Cephalic setae not over 10
 Head with elaborately ornate calvarium; striae 0; spinneret present *Deontostoma* Fil. 1916
 Head without calvarium; cuticular pores absent
 Plate round, with "bent" amph.; phar. small; ceph. set. 4, cerv. oft long, many *Diplopettis* Cobb 1905
 (= *Dipeltis* Cobb 1891
 = *Discophora* Villot 1875)
- Plate absent; pharynx large, complex, of more than 1 chamber; onchium large, sharp
 Onchium conoid to base of short stylet; pharynx \pm quadrate; σ^7 supplements eurystomoid
 Long cervical setae \pm 30; σ^7 supplements weakly developed. *Ledovitia* Fil. 1927
 Long cervical setae 0
 Tail rounded; spinneret oft present; σ^7 supplements well developed. . . *Eurystomina* Fil. 1921
 (= *Eurystoma* Marion 1870
 = *Marionella* Cobb 1922)
- Tail spiculate; spinneret 0; σ^7 supplements weakly developed. *Pareurystomina* Micol. 1930
 Onchium a long stylet; phar. much deeper than wide; σ^7 an enchelid
 Oesophageal bulbs 5-8, serial, contiguous. *Polygastrophora* de Man 1922
 Oesophageal bulbs 0
 Anterior oesophagus (pharynx ?) narrower, lumen wide. *Calyptronema* Marion 1870
 (= *Catalaimus* Cobb 1920)
- Anterior oesophagus not differentiated
 Vestibule set off by row of "commas," pharyng. rings 3; eyes 0 *Symplocostomella* Micol. 1930
 Vestibule set off by rows of denticles or markings, or by a ring
 Pharyngeal chambers behind vestibule more than 2. *Symplocostoma* Bast. 1865
 Pharyngeal chambers behind vestibule 2; eyes 2. *Dilaimus* Fil. 1927
 (= *Amphistenus* Marion 1870)
- Oesophagus with median or posterior bulb or swelling, or both (for alternative see page 11)
 Amphid not known or obscure (for alternative see next page)
 Female not known
 Habitat soil; phar. tubular; $\frac{1}{2}$ neck; onch. outward-acting; oes. dorylaimoid; spinneret 0 *Nanonema* Cobb 1905
 (= *Cephalonema* Cobb 1893)
- Habitat marine; pharynx not over $\frac{1}{2}$ neck-length; spinneret present (Rhinionema?)
 Pharyngeal bulb stronger dorsad; onchium pharyngeal; cephalic setae small or 0. . . . *Iotadorus* Cobb 1920
 Pharyngeal bulb absent; onchium a mere prod at lips; cephalic setae 6. *Rhinionema* Allgen 1927
- Female known
 Ovary 1
 Gonad $\frac{1}{2}$ σ^7 outstretched; spinneret pres.; wings 10, marks V-shaped; ceph. set. 4; marine (*Nudora* Cobb 1920)
 Gonad $\frac{1}{2}$ reflexed; spinneret absent; not marine
 Onchium (spear) bent dorsad at tip; amphid small, oval, at lips; in soil (*Campydora* Cobb 1920)
 Onchium not bent obliquely at tip
 Cardiac bulb not valvate; median bulb strong
 Tooth recurved outward; long. striae present; σ^7 supplements papilloid. . . *Diplogaster* M. Schultze 1857
 Tooth forward-pointing, deepset; long. striae none; σ^7 supplements setose . . . *Acrostichus* Rahm 1928
 Cardiac bulb valvate; median bulb absent, oesophagus cephaloboid
 Posterior pharynx muscular, lumen closed; vulva \pm 75%; tail \pm blunt. *Plectonchus* Fuchs 1930
 Posterior pharynx \pm open; vulva 53-65%; tail conoid, usually slender
 Vagina directed inward, not heavily muscular; not viviparous *Macrolaimus* Maupas 1900
 Vagina directed forward, heavily muscular; viviparous or ovoviviparous. . . . *Turbatrix* Peters 1927
 (= *Anguillula* auctores)
- Ovaries 2
 Gonads $\frac{1}{2}$ outstretched; pharyngeal bulb \pm globular. (*Bolbolaimus* Cobb 1920)
 (= *Bulbopharyngiella* Allgen 1929)
- Gonads $\frac{1}{2}$ reflexed, exc. perhaps Demaniella
 Spinneret absent; oesophagus with muscular median bulb, usually valvate
 Cardiac bulb valvate
 Lip region not set off; odontia absent. *Poikilolaimus* Fuchs 1930
 Lip region set off by constriction; curved odontia 2 or 4 *Rhabditis* Duj. 1845
 (= *Diploscapteroides* Rahm 1928)
- Cardiac swelling not valvate, oesophagus diplogastroid
 Onchium small, anterior; median bulb wide as long, wider than oesophagus *Neodiplogaster* Cobb 1924
 Onchia larger, basal; "prod" at lips; median bulb not wider than oesophagus *Demaniella* Steiner 1914
 (= *Demania* Steiner 1914)
- Spinneret present; oesophagus without median bulb; wings often present; marine exc. *Punctodora*
 Lip reg. protrusile; ceph. set. 4; renette long; excretory pore opp. pharynx. . *Ptycholaimellus* Cobb 1920
 Lip region not protrusile
 Pharyngeal swelling stronger dorsad; σ^7 supplements chromadoroid exc. *Odontonema*
 Striae altered laterally; onchium large, dorsal, apex ventral. *Hypodontolaimus* de Man 1886
 Striae not altered laterally
 Onchium solid, inward-pointing. *Prochromadora* Fil. 1922
 Onchium with internal cavity, forward-pointing
 Tooth small. *Chromadorita* Fil. 1922
 Tooth very large, apex ventral; σ^7 supplements papilloid. *Odontonema* Fil. 1930
- Pharyngeal swelling not stronger dorsad
 Cardiac bulb long, 2- or 3-parted; striae altered laterally; tooth large, forward-pointing
 The bulb with dilated lumen. *Spilophorella* Fil. 1918
 The bulb with lumen not dilated; σ^7 supplements chromadoroid. *Chromadorissa* Fil. 1917
 Cardiac bulb short, not divided
 Striae altered laterally, several rows of dots larger. *Chromadora* Bast. 1865
 Striae not altered laterally
 Onchium solid, inward-pointing; σ^7 supplements chromadoroid. *Prochromadora* Fil. 1922
 Onchium with internal cavity, forward-pointing
 Tooth small, sharp; σ^7 supplements chromadoroid. *Chromadorita* Fil. 1922
 Tooth large, irregularly rectangular; cardiac bulb very large, simple *Punctodora* Fil. 1930

- Amphid known (for alternative see preceding page)
- External amphid not spiral, circular nor elliptical, but a transverse slit
- Female not known; amphid large
- Pharyngeal bulb not stronger dorsad; ceph. set. 10; cutic. "basket-work" on neck *Nygmatochus* Cobb 1933
- Pharyngeal bulb stronger dorsad; striae of dots
- Denticles in pharynx numerous; cephalic setae 4; in humus near sea coast. *Denticulella* Cobb 1933
- Denticles 0; cephalic setae if present almost invisible; marine. *Iotadorus* Cobb 1920
- Female known
- Ovary 1, reflexed; habitat soil about roots
- Tooth short, closed, not bent at tip; ovary anterior, vulva 57%; spinneret(?) present *Oionchus* Cobb 1913
- Tooth (spear) bent obliquely dorsad at tip when extruded; spinneret absent. *Campydora* Cobb 1920
- Ovaries 2, reflexed; spinneret present
- Oesophageal bulbs 5-8, contiguous; amphid with pocket; ♂ supplements eurytomoid *Bolbella* Cobb 1920
- Oesophageal bulbs not more than 2, not contiguous
- Tooth small, near lips; setae 0; pharynx long, tubular; in fresh water. *Rhabdolaimus* de Man 1880
- Tooth not small; pharynx ± 1 head-width deep; amphid usually at lips
- Onchium curved, solid, apparently mobile radially. *Prochromadora* Fil. 1922
- Onchium forward-pointing or \pm rectangular, motion apparently \pm parallel to axis
- Lip reg. protrusile; ceph. set. 4; renette long; excretory pore opp. pharynx. *Ptycholaimellus* Cobb 1920
- Lip region not protrusile
- Neck with cuticular "basket-work"; 4 sublat. scales opp. phar.; spicula 2 sections *Rhrips* Cobb 1920
- Neck without "basket-work," or laterally only; scales 0; spicula entire
- Striae of rod-like elements, exc. towards extremities, changing to large dots laterally
- Cardiac bulb broad, well set off, valvate. *Punctodora* Fil. 1930
- Cardiac bulb not very broad, confluent, non-valvate. *Neochromadora* Micol. 1924
- Striae of dots, altered laterally
- Pharyngeal bulb stronger dorsad; onch. large; card. bulb 2-3-zoned. *Hypodontolaimus* de Man 1886
- Pharyngeal bulb if present not stronger dorsad; cardiac bulb not zoned. *Chromadora* Bast. 1865
- External amphid spiral, circular or elliptical
- Female not known; marine or salt marsh (exc. *Amphispira* ?); wings none
- Habitat soil (possibly marine); amphid spiral; cephalic setae 0; spinneret present *Amphispira* Cobb 1920
- Habitat marine or salt marsh
- Amphid multispire, large, on front curvature of head
- Striae altered laterally. *Neonchus* Cobb 1933
- Striae not altered laterally. *Amphispira* Cobb 1920
- Amphid not multispiral
- Onchium spear-like, free at tip. *Oistolaimus* Ditlev. 1921
(cf. *Onyz* Cobb 1891)
- Onchium not spear-like
- Cephalic setae minute; pharyng. bulb stronger dorsad; amphid narrow, transverse *Iotadorus* Cobb 1920
- Cephalic setae 0; amphid circular or monospiral
- Tooth ventral. *Crassolaimus* Kreis 1929
- Tooth dorsal; nema cephalated by expansion and by cessation of striae. *Micromicron* Cobb 1920
- Female known
- Ovary 1
- Gonad ♀ outstretched; amphid circular; cephalic setae 4; spinneret present; marine
- Wings 10, of V-shaped marks; phar. bulb long, set off by constrict.; lab. papill. setose *Nudora* Cobb 1920
- Wings 0; phar. bulb very slight; labial papillae 0; renette large, of ± 6 cells. *Synonema* Cobb 1920
- Gonad ♀ reflexed
- Habitat marine; amphid round; spinneret pres.; wings 6-20, marks V-shaped *Monoposthia* de Man 1889
- Habitat not marine; amphid round or oval; spinneret none exc. *Udonchus*
- Median oesophageal bulb present, usually valvate; tail conoid, then setaceous; bursa none
- Pharynx narrow, post. part 6 times width; ceph. set. 0; onch. small, basal *Diplogasteroides* de Man 1912
- Pharynx not more than twice as deep as wide; cephalic setae usually present
- Onchium anterior, recurved, outward-acting. *Diplogaster* M. Schultze 1857
- Onchium deepest, forward-pointing. *Acrostichus* Rahm 1928
- Median oesophageal bulb absent; striae fine, plain
- Pharynx occupying $\frac{1}{2}$ neck-length; habitat decaying bulbs. (*Odontopharynx* de Man 1912)
- Pharynx occupying $\frac{1}{4}$ neck-length or less
- Tooth (spear) bent obliquely dorsad at lips when extruded; soil near roots. (*Campydora* Cobb 1920)
- Tooth not bent
- Spinneret present; external amphid oval; pharyng. bulb absent; in fresh water *Udonchus* Cobb 1913
- Spinneret absent; amphid circular
- Pharyngeal swelling absent; cardiac bulb valvate. *Macrolaimus* Maupas 1900
- Pharyngeal swelling clavate; in moist soil. *Odontolaimus* de Man 1880
(= *Neonchus* Cobb 1893)
- Ovaries 2
- Gonads ♀ outstretched; spinneret present; habitat marine or brackish soil
- Amphid not multispire; onchia several, often minute
- Pharyngeal bulb none, or slight, confluent; ceph. setae 4; labial papillae 6. *Microlaimus* de Man 1880
- Pharyngeal bulb almost globular; cephalic setae 6; labial setae 6. (*Bolbolaimus* Cobb 1920)
- (= *Bulbopharyngiella* Allgen 1929)
- Amphid multispire; striae finely dotted; pharyngeal bulb absent
- Head setose; striae not altered laterally; spicula long; apophysis none. *Comesoma* Bast. 1865
- Head less setose; spicula short; gubernaculum with apophysis
- Male ventral pre-anal supplementary organs small, papilloid. (*Parasabatieria* de Man 1907)
- Male supplementary organs absent. (*Sabatieria* de Rouville 1903)
- Gonads ♀ reflexed
- Spinneret absent (for alternative see next page)
- Habitat marine; cephalic setae 4; pharyngeal bulb absent, pharynx small. *Spirina* Fil. 1918
(= *Spira* Bast. 1865)
- Habitat not marine; tail \pm setaceous exc. *Neodiplogaster*
- Cardiac bulb valvate. *Poikilolaimus* Fuchs 1930
- Cardiac bulb not valvate
- Pharyngeal bulb clavate, $\frac{1}{2}$ - $\frac{1}{3}$ neck-length; median bulb none; ceph. set. 10 *Odontolaimus* de Man 1880
- Pharyngeal bulb none or short, not tapering; oesoph. diplogastroid; onchium a projection

- Median oesophageal bulb not swollen; ceph. set. 0; striae plain... (*Odontopharynx* de Man 1912)
 Median oesophageal bulb swollen; cardiac swelling non-valvate
 Pharynx $\frac{1}{2}$ wide as deep; ceph. set. present; tail setaceous; bursa 0... *Diplogaster* M. Schultze 1857
 Pharynx narrow, posterior tubular portion not over $\frac{1}{2}$ as wide as deep; ceph. set. 0
 Onchium basal; tail conoid then setaceous; bursa none... *Diplogasteroides* de Man 1912
 Onchium anterior; striae of dots; tail conoid; bursa rhabditoid... *Neodiplogaster* Cobb 1924
 Spinneret present, exc. sometimes *Spirina* (for alternative see preceding page)
 Oesophageal bulbs 5-8, serial, contiguous; ♂ an enchelid; marine... *Polygastrophora* de Man 1922
 Oesophageal bulb 1
 Anterior oesophagus narrow, lumen wide; marine... *Calyptronema* Marion 1870
 (= *Catalaimus* Cobb 1920)
 Anterior oesophagus not differentiated
 Wings present
 Pharyngeal bulb set off by constriction; wings 12, bristles retrorse; marine... *Rhinema* Cobb 1920
 Pharyngeal bulb none or confluent
 Nema cephalated by cuticular change & \pm by expansion; cardiac bulb short, not divided
 Head "punctate"; wings 12-18; ceph. set. 6; denticles pres.; amph. spiral *Desmodorella* Cobb 1933
 Head plain; wing 1; ceph. set. 0; denticles 0; amph. round; caud. glands pre-anal *Xenonema* Cobb 1920
 Nema not cephalated; cardiac bulb long, 2- or 3-zoned; amphid spiral
 Striae of dots; wings scalariform; ♂ supplements 0... *Spilophorium* Cobb 1933
 (= *Spilophora* Bast. 1865)
 Striae plain
 Cephalic setae many, jointed; ♂ pre-anal cuticle thickened... *Neonyx* Cobb 1933
 Cephalic setae 4, short; wing 1; ♂ supplements post-anal... *Metachromadora* Fil. 1918
 (= *Chromadoropsis* Fil. 1918)
 Wings 0; striae not altered laterally, exc. sometimes *Desmodora*
 Habitat freshwater; minute subordinate onchia usually present
 Pharynx behind onchia \pm parallel-sided; pharyngeal bulb definite... *Ethmolaimus* de Man 1880
 Pharynx behind onchia \pm conoid; pharyngeal bulb absent
 Amphid spiral; cephalic setae usually 10; pharynx open, irregular... *Achromadora* Cobb 1913
 Amphid circular; cephalic setae 4; pharynx often obscure... *Prodesmodora* Micol. 1923
 Habitat marine
 Dorsal pharyngeal wall thickened, onchium-like; amphid monospire
 at lips; cephalic setae 4; renette far back
 Oesoph. glands obscuring junction of intest.; dorsal phar. element \pm uniform *Onchium* Cobb 1920
 Oesoph. glands not prominent; dorsal pharyng. element not uniformly thick, but heavy
 Tooth-like thickening parallel to axis... *Camacolaimus* de Man 1889
 (= *Digstonchus* Cobb 1920
 = *Acontiolaimus* Fil. 1918)
 Tooth-like thickening bent outward from beginning of vestibule
 Point of "onchium" single... *Acmaeolaimus* Fil. 1918
 Point of Y-shaped "onchium" duplex... *Ypsilon* Cobb 1920
 Dorsal pharyngeal wall not thickened; onchium a distinct projection
 Cephalic setae 12-26; cardiac bulb elongate; amphid spiral
 Striae of dots, on head also; labial papillae setose; ceph. set. 12... *Bradylaimus* Stekhoven 1931
 Striae plain, absent on head; cephalic setae 24-26; amphid monospire
 Pharynx denticulate; onch. sigmoid ventrally; ♂ supplements sigmoid *Sigmophora* Cobb 1933
 Pharynx not denticulate; onch. not sigmoid; ♂ supplements papilloid *Xanthodora* Cobb 1920
 Cephalic setae 4
 Striae of dots
 Dorsal onchium large, basal; pharyngeal bulb stronger dorsad... *Odontonema* Fil. 1930
 Dorsal onchium anterior; pharyngeal bulb \pm symmetrical... *Chromadorina* Fil. 1918
 Striae plain
 Head set off by lack of striae; pharynx & onchium inconspic.; marine... *Desmodora* de Man 1889
 Head not set off; striae extending to lips
 Pharyngeal bulb 0; cardiac bulb short; onchium minute; neck 6-7%... *Spirina* Fil. 1918
 (= *Spira* Bast. 1865)
 Pharyngeal bulb present; cardiac bulb long, massive, 2- or 3-zoned; neck over 12%
 Onchium spear-like; head rounded... *Onyx* Cobb 1891
 Onchium not spear-like; head truncate... *Metachromadora* Fil. 1918
 (= *Chromadoropsis* Fil. 1918)
 Teeth (onchia or odontia) 2 or more (for alternative see page 11)
 Oesophagus with median or posterior bulb or swelling, or both (for alternative see page 17)
 The teeth \pm equal in size, usually mobile radially (for alternative see next page)
 Stroke of teeth outward; amphid often obscure (for alternative see next page)
 Spinneret absent; ♀ or '♀'
 Cardiac bulb valvate; cuticle without longitudinal striae... *Diploscapter* Cobb 1913
 Cardiac bulb not valvate, median bulb valvate; cuticle with longitudinal striae
 Posterior pharyngeal chamber behind onchia; lips with radiate framework... *Mononchoides* Rahm 1928
 Posterior pharyngeal chamber absent; lips without radiate framework... *Diplogaster* M. Schultze 1857
 (= *Pristionchus* Kreis 1932)
 Spinneret present; median oesophageal bulb absent
 Pharynx tubular, 4-8% of body length
 Amphid long, reaching base of pharynx or mid-body; odontia 6, minute; ♀-♀... *Pseudolella* Cobb 1920
 Amphid small, at lips, with pocket; odontia 3, massive; tail oft striate; ♀'... *Syringolaimus* de Man 1888
 Pharynx conoid, not over 2% of body length
 Striae plain
 Odontia 0; onch. 3, at base shallow phar.; amph. round; card. swelling obvious; ♀-♀ *Aponchium* Cobb 1920
 Odontia 6 or 12; onch. 0; amph. bent or crook-shape; card. bulb 0 or weak; ♀-♀ *Odontophora* Bütsch. 1874
 (= *Trigonolaimus* Ditlev. 1918
 = *Conolaimus* Fil. 1918)
 Striae of dots; cardiac swelling obvious
 Amphid spiral; onch. 3, curved; ♂ supplements 0 or papilloid; gubernac. 0; ♂ only... *Statenia* Allgen 1930
 Amphid a transverse slit, near lips; ♀'
 Lateral fields marked by longitudinal rows of larger dots... *Chromadorella* Fil. 1918
 Lateral fields without longitudinal rows of larger dots... *Prochromadorella* Micol. 1924

- Stroke of teeth inward; amphid occasionally obscure (for alternative see preceding page)
- Spinneret 0; phasmids present; odontia 6; oesophagus cephaloboid; ovaries reflexed
- Nema usually cephalated; cirri 0; amphid round; soil & fresh water; ♀ (*Teratocephalus* de Man 1876)
- Nema not cephalated; cirri 6; amphid oval; tail hooked; in bark, etc.; ♀ (*Chamberstella* Cobb 1920)
- (= *Diatolaimus* Rahm 1928)
- Spinneret present**
- Mandibular jaws 2 or 3; pharyngeal and cardiac bulbs strong; marine
- Jaws 3; amphid not known; ♂ only *Fusonema* Kreis 1928
- Jaws 2, lateral, dentate; amphid multispire *Cheironchus* Cobb 1917
- (= *Dignathonema* Fil. 1918)
- Mandibular jaws absent**
- Amphid multispire; lips flat, confluent; pharynx rather narrow; marine
- Striae very fine; wings 0; labial papillae not seen; ♂ only (*Alaimonema* Cobb 1920)
- Striae ± coarse, of dots; wings pres.; labial papillae 12; ♀ -, or reflexed at tip (*Mesonchium* Cobb 1920)
- (= *Pepsonema* Cobb 1920)
- Amphid not multispire**
- Habitat soil; amph. monospire to crook-shape; phar. 6%; nema small, 0.2 mm.; J (*Pynolaimus* Cobb 1920)
- Habitat marine; pharynx under 2%, not tubular; ovaries not reflexed
- Vulva 15%; odontia 12, minute; amphid "folded"; ♀ - *Synodontium* Cobb 1920
- Vulva 81%; odontia 0; onchia 3; pharyngeal swelling slight; ♀ - (*Aponchium* Cobb 1920)
- The teeth ± unequal in size, usually not mobile radially; amph. usually not obscure (from preceding page)
- External amphid not spiral, circular nor elliptical; ovaries 2; spinneret present
- Gonads ♀ outstretched; amph. long, reaching sometimes to mid-body; odontia 6; ♀ - *Pseudolella* Cobb 1920
- Gonads ♀ reflexed; amphid a transverse slit
- Striae fine, plain; onchia dorsal and ventral *Deltanema* Kreis 1929
- (cf. *Metalinhomoeus* de Man 1907)
- Striae resolvable into elements; dorsal onchium usually the largest
- Spicula of 2 joints; striae forming "basket-work" on neck; amphid opp. base pharynx *Rhyps* Cobb 1920
- Spicula entire; striae of dots; amphid near lips *Chromadora* Bast. 1865
- External amphid spiral, circular or elliptical**
- Female not known; spinneret present; marine
- Amphid multispire; head conoid; onch. minute; ceph. & subceph. setae in sets of 4 *Alaimonema* Cobb 1920
- Amphid circular, elliptical or monospiral
- Pharynx surrounded at base by cuticularized ring *Cricolaimus* Southern 1914
- Pharynx not surrounded by cuticularized ring
- Striae of dots, interrupted laterally; pharynx cyatholaimoid, onchium large . . . *Endolaimus* Fil. 1922
- Striae unaltered laterally; pharynx small; ♂ supplements tubular, post-anal *Catanema* Cobb 1920
- Female known**
- Ovary 1**
- Gonad ♀ outstretched; spinneret pres.; ♂ supplements numerous, complex; marine *Aponchium* Cobb 1920
- Gonad ♀ reflexed; amphid small, circular, obscure; spinneret 0; not marine
- Cardiac bulb valvate, median bulb absent; bursa small, ribs stout *Myolaimus* Cobb 1920
- Cardiac bulb not valvate; oesophagus ± diplogastroid
- Oesophagus without (or with non-valvate, clavate) median swelling *Odontopharynx* de Man 1912
- Oesophagus with median valvate bulb *Diplogaster* M. Schultze 1857
- (= *Pristionchus* Kreis 1932)
- Ovaries 2**
- Gonads ♀ outstretched, Mesonchium reflexed at tips; marine
- Amphid spiral; dots larger laterally; onchia ± equal; spinneret present (*Mesonchium* Cobb 1920)
- (= *Pepsonema* Cobb 1920)
- Amphid circular; striae when present not altered laterally; onchia unequal
- Striae desmodoroid, coarse, plain; amph. behind pharynx, open caudad *Paracotholaimus* Schulz 1932
- Striae none, or not desmodoroid
- Pharyngeal bulb ± globular, set off by constriction; spinneret present *Bolbolaimus* Cobb 1920
- (= *Bulbopharyngiella* Allgen 1929)
- Pharyngeal swelling slight, confluent; base of pharynx denticulate *Linhomoeus* Bast. 1865
- (i.e. subg. *Eulinhomoeus* de Man 1907)
- Gonads ♀ reflexed (Mesonchium at tips only)**
- Habitat marine; spinneret present; amphid spiral**
- Striae of dots; amphid opposite pharynx, of few winds
- Subventral onchium small, dorsal acting out; dots unaltered laterally . . . *Chromadorina* Fil. 1918
- Subventral & dorsal onchia ± equal, apices axial; dots larger laterally *Mesonchium* Cobb 1920
- (= *Pepsonema* Cobb 1920)
- Striae none or plain**
- Head set off by cessation of coarse annulation; amphid usually multispire . . . *Desmodora* de Man 1889
- Head not set off; striae none or very fine
- Male ventral pre-anal supplements tubular, sigmoid, in 2 rows *Polysigma* Cobb 1920
- Male ventral pre-anal supplements simple, in 1 row *Chromaspirina* Fil. 1918
- (= *Mesodorus* Cobb 1920)
- Habitat not marine**
- Spinneret present; amphid spiral, behind pharynx; in fresh water *Achromadora* Cobb 1913
- Spinneret absent; phasmids present; amphid circular; usually in soil
- Cardiac bulb valvate, oesophagus cephaloboid; onchia 4 *Seleneella* Rahm 1932
- Cardiac bulb not valvate; oesophagus diplogastroid
- Anterior oesophagus (corpus) without (or with clavate, non-valvate) swelling
- Cephalic setae 6 or 10; dorsal onch. opposed by ventral onchium or ridge . . . *Butlerius* Goodey 1929
- Cephalic setae 6; dorsal onchium opposed by denticles *Odontopharynx* de Man 1912
- Anterior oesophagus (corpus) ending in ± valvate median bulb
- Lips with radiate framework; post. pharyngeal chamber behind onchia . . . *Mononchoides* Rahm 1928
- Lips without radiate framework; post. pharyngeal chamber none *Diplogaster* M. Schultze 1857
- (= *Pristionchus* Kreis 1932)

- Oesophagus plain, i.e. without median or posterior bulb or swelling (for alternative see page 15)
 The teeth \pm equal in size, usually mobile radially (for alternative see next page)
 Stroke of teeth outward; amphid usually not obscure
 External amphid spiral, circular or elliptical; spinneret present (Eleutherolaimus ?)
 Armature of 2 teeth; ceph. set. 4; amph. monospire, in front of setae; σ^7 only *Diodontolaimus* Southern 1914
 Armature of 6 or 12 odontia
 Odontia 3-jointed, heavy; phar. cylind.; onch. dorsal; amph. round-multispire; - φ *Scaptrella* Cobb 1917
 Odontia not jointed
 Amphid "folded" or shepherd's crook; - φ *Odontophora* Bütsch. 1874
 (= *Trigonolaimus* Ditlev. 1918
 = *Conolaimus* Fil. 1918)
- Amphid circular or elliptical, sometimes obscure; cephalic setae 4
 Pharynx obscure; odontia small; lab. set. 6, setae long; striae coarse; ' φ ' *Stephanolaimus* Ditlev. 1918
 Pharynx obvious; odontia larger; labial and subcephalic setae 0; σ^7 only
 Striae of dots; amphid transverse, open caudad..... *Apodontium* Cobb 1920
 Striae plain; amphid circular..... *Eleutherolaimus* Fil. 1922
- External amphid not spiral, circular nor elliptical; spinneret pres. exc. *Ironus* and *Parironus*
 Cardiac bulb slight; striae of dots; amphid a transverse slit near lips; ' φ '
 Dots of striae larger laterally, in longitudinal rows..... (*Chromadorella* Fil. 1918)
 Dots of striae not altered laterally..... (*Prochromadorella* Micol. 1924)
- Cardiac bulb none; striae none, or plain and very fine
 Odontia 6, sometimes dentate; amphid \pm "folded" or crook-shaped; - φ -
 Amphid opposite basal portion of pharynx; odontia less developed..... *Azonolaimus* de Man 1889
 Amphid opposite anterior portion of pharynx; odontia more developed..... *Odontophora* Bütsch. 1874
 (= *Trigonolaimus* Ditlev. 1918
 = *Conolaimus* Fil. 1918)
- Odontia or onchia 3; amphid transverse, with internal pocket; ovaries reflexed
 Armature 3 awl-shaped onchia, protrusile forward; pharynx 0.5%; ' φ '..... (*Fimbrilla* Cobb 1905)
 (= *Fimbria* Cobb 1894)
- Armature 3 heavy odontia; pharynx tubular, 1-4%; *Ironinae*
 Lips expanded, denticulate, 6; tail 2%; spinneret ventrad; φ '..... *Trissonchulus* Cobb 1920
 Lips not expanded; tail 5% or more; spinneret not ventrad; ' φ '
 Spinneret none; pharynx 2-4%; dorsal tooth duplex; in fresh water and soil..... *Ironus* Bast. 1865
 Spinneret present (in *Parironus* rudimentary or none); marine
 Cephalic setae 0 or papilloid; odontia not duplex; pharyngeal swelling slight
 Pharyngeal wall thin; pharynx 3%; nema 3 mm..... *Dolicholaimus* de Man 1888
 Pharyngeal wall thick; pharynx 1.1%; nema 2-7 mm..... *Thalassironus* de Man 1889
 Cephalic setae present
 Odontia duplex; phar. 2.7%, bulb set off by constriction; ceph. set. 6, subceph. 4. *Ironella* Cobb 1920
 Odontia not duplex; pharynx 1-1.6%, swelling slight; cephalic setae 10..... *Parironus* Micol. 1930
- Stroke of teeth inward; amphid known, in *Enoplidae* with internal pocket and often obscure
 External amphid not spiral, circular nor elliptical; ovaries reflexed
 Ovary 1, post.; phar. bulb strong; "palps" labial; onch. 3, at summit of apophyses *Gammanema* Cobb 1920
 Ovaries 2; wings absent; amphid with internal pocket; *Enoplidae*
 Calvarium a band with undulating margins; glands lateral; odontia large..... *Fiacra* Southern 1914
 Calvarium absent
 Onchia 3, awl-like, protrusile forward, like bristles around open mouth..... *Fimbrilla* Cobb 1905
 (= *Fimbria* Cobb 1894)
- Onchia not awl-like
 Lips confluent, bearing 6 (or 3 duplex) odontia
 Pharynx narrow; onchia 0; odontia probably 6..... *Tubolaimella* Cobb 1933
 Pharynx conoid; onch. 3, midway, alternating paired odontia..... *Rhabdodemania* Baylis & Daubney 1926
 (= *Demania* Southern 1914)
- Lips distinct, 3; armature heavy
 The lips large, conoid, as high as broad; jaws 3
 Mandibles 2-clawed; lips radially striate; spicula long, striate..... *Enoploides* Saveljev 1912
 Mandibles absent; lips not striate; tooth midway on onchial framework..... *Enoplolaimus* de Man 1893
 The lips not conoid, nor as high as broad
 Armature 3 short flattish acute onchia, doming the \pm small pharynx..... *Tridontolaimus* de Man 1893
 Armature of 3 jaws within the pharynx, anteriorly 2-pointed..... *Enoplus* Duj. 1845
 (= *Enoplostoma* Marion 1870)
- External amphid spiral, circular or elliptical
 Female not known
 Amphid monospire, transverse; lips 4; cephalic setae 4; σ^7 supplements 8-12, tubular
 Striae very fine; pharynx conoid; onchia 3; labial papillae 2..... *Dagda* Southern 1914
 Striae coarse; phar. cylind.; onch. 2, lateral or subventral; lab. papill. 4 *Diodontolaimus* Southern 1914
 Amphid multispire; σ^7 supplements \pm chromadoroid
 Pharynx unarmed; lips 6, grasping organs, with longitudinal cuticular ribs..... *Dispira* Cobb 1933
 Pharynx armed, bulb strong; often nemativororous; σ^7 supplements chromadoroid
 Mandibles 3, split exc. at tips, distal hooks 3, subordinates 2; phar. 1 chamber..... *Synonchium* Cobb 1920
 Mandibles 0; phar. 2 chambers; onchia 3, midway, with backward apophyses..... *Trogolaimus* Cobb 1920
- Female known
 Ovary 1; amphid circular or monospiral (for alternative see next page)
 Gonad φ post.; onch. 3, on apophyses; phar. bulb strong; spinneret pres.; marine *Gammanema* Cobb 1920
 Gonad φ anterior
 Spinneret none; wings none; labial "palps" 6, small; marine; - φ
 Amphid large, raised centrally; tail with thorn-like setae..... (*Zanema* Cobb 1920)
 Amphid small, not projecting; tail without setae..... (*Crystallinema* Cobb 1920)
 Spinneret present
 Cephalic setae usually 0 or papilloid; ovoviviparous or viviparous; in gills of land crabs
 Pharynx small, of 2 chambers; cephalic setae 10, often papilloid..... (*Monhystrium* Cobb 1920)
 Pharynx large, open, of 1 chamber; cephalic setae 0; wings none..... *Tripplium* Cobb 1920
 Cephalic setae present; ovary outstretched; marine
 Lips thick, with 3 projecting finger-like "odontia"; wings up to 32 on head..... (*Xyala* Cobb 1920)
 Lips 6, very thin, each with 3 parallel-sided cuticularized ribs..... (*Daptonema* Cobb 1920)
 (cf. *Theristus* Bast. 1865)

- Ovaries 2 (for alternative see preceding page)
 Gonads ♀ outstretched; spinneret present; marine
 Amphid "folded" or crook-shaped; pharynx conoid; striae plain.....(*Axonolaimus* de Man 1889)
 Amphid multispire; striae of dots, altered on lateral fields
 Spicula 2-jointed; nema 2% wide.....*Dorylaimopsis* Ditlev. 1918
 (= *Xinema* Cobb 1920)
 (= *Mesonchium* Cobb 1920)
 (= *Pepsonema* Cobb 1920)
 Spicula entire; nema 4% wide.....
 Gonads ♀ reflexed; spinneret present
 Amphid large, multispire; striae resolvable into elements; often nemativorovous
 Pharynx 2 chambers; mid-pharynx denticulate, apophyses forward & backward; striae of dots
 Circular muscles about anterior pharynx; phar. small; denticles irregular. *Cobbionema* Fil. 1922
 Circular muscles none; pharynx capacious; denticles regularly placed *Halichoanolaimus* de Man 1886
 (= *Smalsundia* Allgen 1929)
 Pharynx not of 2 chambers; denticles absent (*Demonema* ?)
 Mandibles none
 Dots of striae not larger laterally; jaws with many horny teeth anteriorly... *Demonema* Cobb 1894
 Dots of striae larger laterally; onchia equal, apices axial..... *Mesonchium* Cobb 1920
 (= *Pepsonema* Cobb 1920)
 Mandibles 2 or 3; Selachineminae
 Dorsal mandible a mere rod, developed mandibles 2..... *Selachinema* Cobb 1915
 Dorsal mandible developed, making 3, ±split; subordinate hooks pres.; cutic. pores pres.
 Central mandibular prong 1, no. hooks odd; striae of dots; ♂ supplements 2 *Synonchium* Cobb 1920
 Central mand. prongs 2, no. hooks even; striae of dots, dashes; ♂ supplements 20 *Synonchiella* Cobb 1933
 Amphid small, opening circular, obscure; lips 3, large; Enoplinae
 Lips not conoid, nor high as broad; labial papillae seldom setose; jaws 2-pointed. *Enoplus* Duj. 1845
 (= *Enoplostoma* Marion 1870)
 Lips large, conoid, often as high as broad; labial papillae 6, setose
 Mandibles 2-clawed; lips radially striate; spicula long, striate..... *Enoploides* Saveljev 1912
 Mandibles absent; lips not striate
 Basal onchia large, reaching lips
 Framework of jaws absent..... *Saveljevia* Fil. 1927
 Framework of jaws present..... *Filipjevia* Kreis 1928
 Basal onchia absent; "jaws" 3, without hooks
 Jaw framework bearing 3 equal mid-pharyngeal teeth..... *Enoploalaimus* de Man 1893
 Jaw framework without teeth..... *Paronoplus* Fil. 1927
 The teeth ±unequal in size, usually not mobile radially; amph., in Enoplidae, oft obscure (from pg. 17)
 Female not known
 Cephalic setae 0; spiculum 1, gubernaculum absent..... *Dioncholaimus* Kreis 1932
 Cephalic setae pres.; spicula 2, gubernac. pres. (*Hyptiolaimus*, *Cacolaimus*, *Convexolaimus*, J only)
 Amphid multispire, ±head width; striae of dots, wings 2; labial "palps" 12, jointed *Pomponema* Cobb 1917
 Amphid circular, elliptical or monospiral
 Pharynx with denticles, or many subordinate onchia; labial papill. 6, setose; amph. round or oval
 Labial "palps" 12, tips large; onch. dorsal, denticles not basal; wing scalariform *Anazonchium* Cobb 1920
 Labial "palps" 0; base of pharynx filled with subordinate onchia..... *Polydontus* Schulz 1932
 Pharynx without denticles or numerous subordinate onchia
 Bulb around pharynx set off by sudden constriction..... *Hyptiolaimus* Cobb 1930
 Bulb around pharynx none or not set off by constriction
 Striae coarse; onchia 2, lateral or subventral; labial papillae 4..... *Diofantolaimus* Southern 1914
 Striae none or very fine; onchia 3
 Lips 4, 2 of them bearing papillae; cephalic setae 4; ♂ supplements tubular... *Daqda* Southern 1914
 Lips 6, papillae 6, setose; cephalic setae 10, short; J
 Head set off by thicker cuticle; wall of pharynx thin..... *Cacolaimus* Kreis 1932
 Head not set off, its cuticle thin; wall of pharynx distinct..... *Convexolaimus* Kreis 1928
 Female known
 Ovary 1, anterior (*Pseudoncholaimus* ?)
 Median onchium spear-like, dorsal onchium minute, basal..... *Enoplocheilus* Kreis 1932
 Median spear-like onchium absent
 Gonad ♀ outstretched; amphid circular or monospiral
 Onchia 2, like vertical plates, subventral; amphid monospire, longer than wide. *Gonionchus* Cobb 1920
 Onchia not plate-like, basal, outlets of oesoph. glands; amph. round; tail ±setaceous *Cobbia* de Man 1907
 Gonad ♀ reflexed
 Spinneret absent; phasmids present
 Cardiac bulb ±glandular, not valvate; median swelling often present (*Odontopharynx* de Man 1912)
 Card. bulb valvate, oes. cephaloboid; vagina directed forward, muscles heavy *Turbatrix* Peters 1927
 (= *Anguillula* auctores)
 Spinneret present; phasmids absent; amphid with internal pocket, usually small
 Cephalic setae 0; papillae in 2 circllets; in soil and fresh water
 Pharynx large, open, ±parallel-sided; spinneret terminal..... *Mononchus* Bast. 1865
 Pharynx narrow, ±conoid; spinneret ventrad..... *Mononchulus* Cobb 1918
 Cephalic setae present; papillae 1 circllet; marine or in brackish waters; Oncholaiminae
 Demanian system absent
 Ovary posterior; vulva 73-77%; gubernaculum none..... *Pseudoncholaimus* Kreis 1932
 Ovary ant.; vulva 70-75%; gubernaculum pres.; spicula ±tail length. *Probncholaimus* Micol. 1924
 Demanian system present
 Uvette and exit pores present..... *Metoncholaimus* Fil. 1918
 Uvette ampulla-like, exit pores absent
 Male with pre-anal appendicule..... *Oncholaimium* Cobb 1930
 Male without pre-anal appendicule..... *Oncholaimus* Duj. 1845
 Ovaries 2
 Gonads ♀ outstretched; striae of rods, exc. laterally; amph. a slit... *Dicriconema* Steiner & Hoeppli 1926
 Gonads ♀ reflexed
 Habitat soil and fresh water (for alternative see next page)
 Spinneret 0; anterior oesoph. (corpus) with or without non-valvate med. bulb; amph. round or oval
 "Spear" none; dorsal onch. opposed by ventral tooth or ridge; ceph. set. 6-10... *Butlerius* Goodey 1929

- "Spear" a minute prod between lips; cephalic setae 0..... *Demaniella* Steiner 1914
(= *Demania* Steiner 1914)
- Spinneret present, exc. sometimes in *Mononchus*
- Amphid spiral; pharynx \pm conoid..... *Nannonchus* Cobb 1913
- Amphid circular or oval, often small, with internal pocket
- Pharynx irregularly conoid; onchia small..... *Trilobus* Bast. 1865
- Pharynx not conoid; one or more onchia large
- Lips thick; papillae 12, in 2 circles; renette not known..... *Mononchus* Bast. 1865
- Lips thin; papillae 6, ceph. set. 10; renette pres., ampulla conspicuous..... *Adoncholaimus* Fil. 1908
- Habitat marine; spinneret usually present (for alternative see preceding page)
- Cuticle annulated
- Annules not resolvable into dots or rod-like elements
- Head truncate; annules interrupted; ceph. set. 4; phar. ribs long., serrate. *Rhabdotoderma* Marion 1870
- Head rounded; annules plain; setae many, short, \pm irregular on calvarium.... *Croconema* Cobb 1920
- Annules of rod-like marks; pharynx cyatholaimoid, vestibule ribbed; amphid a transverse slit
- Ceph. set. 4, labial papillae setose; annules plain laterally.... *Dicriconema* Steiner & Hoeppli 1926
- Ceph. set. 10; annules with V-shaped marks laterally, "basket-work" on neck.... *Rhyps* Cobb 1920
- Cuticle not annulated; striae if present very fine
- Head with subcuticular calvarium; amphid with internal pocket; Enopliidae
- Calvarium complex, ornate, long-conoid..... *Thoracostoma* Marion 1870
- Calvarium relatively simple, a band with undulating margins; glands lateral
- Pharynx wide; lips with denticles; odontia present(?); nema 20-34 mm.... *Cylicolaimus* de Man 1889
- Pharynx narrow, \pm filled by onchia, dorsal large; cervical setae many; nema 8-17 mm.
- Odontia present; cephalic setae long..... *Fiacra* Southern 1914
- Odontia absent; cephalic setae short or medium..... *Synonchus* Cobb 1894
(= *Jagerskiöldia* Fil. 1916)
- Head without calvarium
- Amphid spiral; pharynx small, irregularly conoid, vestibule ribbed..... *Chromaspirina* Fil. 1918
(= *Mesodorus* Cobb 1920)
- Amphid with internal pocket; pharynx large, open, squarish, ribbed vestibule absent
- Pharynx not divided, walls \pm thick; amph. small; excretory pore oft not at lips; Oncholaiminae
- Demanian system present, with exit pores and developed uvette..... *Adoncholaimus* Fil. 1918
- Demanian system absent, or gonenteric duct only
- Subventral onchia equal in size; amphidial opening longer than wide
- Tail 33%, filiform, spinneret none; gubernaculum none..... *Filoncholaimus* Fil. 1927
(= *Pseudoparonycholaimus* Kreis 1932)
- Tail short, rounded; gubernaculum present..... *Pontonema* Leidy 1855
(= *Paronycholaimus* Fil. 1916)
- Subventral onchia unequal in size; pharynx depth twice its width or more
- Cuticle not viscous; phar. bulb clavate; lip reg. set. off by constriction.... *Steineriella* Allgen 1932
(= *Steineria* Ditlev. 1928)
- Cuticle viscous; pharyngeal bulb none; nema under 3 mm.; tail 6-10%
- Pharynx very deep, \pm filled by subventral onch.; bursa present *Oncholaimellus* de Man 1890
- Pharynx not over twice as deep as wide, not filled by onchium; bursa none
- Dorsal wall of pharynx well developed..... *Viscosia* (de Man) Fil. 1918
- Dorsal wall of pharynx rudimentary, an extension of dorsal tooth.... *Meroviscosia* Kreis 1932
- Pharynx 2-5 chambers, wall thin; amph. large; excretory pore oft at lips; large onch. subventral
- Oesophageal bulbs 5-8, serial, contiguous
- Amphidial opening transverse, slit-like; σ^7 supplements eury stomoid.... *Bolbella* Cobb 1920
- Amphidial opening \pm circular; σ^7 an enchelid, supplements simple.... *Polygastrophora* de Man 1922
- Oesophageal bulbs none
- External amphid transverse, slit-like; pharynx usually \pm square; caudal glands before anus
- Denticles forming a broad mid-pharyngeal band; σ^7 supplements simple.... *Thoönchus* Cobb 1920
- Denticles in 1-3 rows; σ^7 supplements complex, "fuked"..... *Eury stomoides* Fil. 1921
(= *Eury stomoides* Marion 1870
= *Marionella* Cobb 1922)
- External amphid round; pharynx deep; onch. spear-like, exc. *Ditlevseniella*; σ^7 oft an enchelid
- Anterior oesophagus narrower, lumen wide..... *Calyptronema* Marion 1870
(= *Catalaimus* Cobb 1920)
- Anterior oesophagus not differentiated
- Large onchia 3..... *Fenestrolaimus* Fil. 1927
- Large onchium 1, other onchia 2
- The large onchium conoid; nema 4-6 mm..... *Ditlevseniella* Fil. 1927
- The large onchium long, slender, spear-like
- Pharyngeal chambers behind vestibule 2; eyes 2..... *Dilaimus* Fil. 1927
(= *Amphistenus* Marion 1870)
- Pharyngeal chambers behind vestibule more than 2
- Chambers separated by 2-4 rings or by dots or denticles..... *Symplocostoma* Bast. 1865
- Chambers (first 2) separated by "comraas"; rings 3; eyes 0.... *Symplocostomella* Micol. 1930

Wall of the Pharynx unarmed (for alternative see page 8)*Cavity not cylindrical nor prismoid*, i.e. cavity conoid or irregular in long. section (for alt. see page 24)

Oesophagus with median or posterior bulb or swelling, or both (for alternative see next page)

Amphid not known or obscure

Female not known; spinneret present; cephalic setae present; nema 1% wide; marine

Cuticle with distinct, anastomosing annules; amphid a transverse slit.....(*Leptonemella* Cobb 1920)Cuticle without annules; amphid minute, at lips.....(*Catanema* Cobb 1920)Female known; spinneret none, exc. *Dasyilaimus*

Ovaries 2, reflexed; wings present

Spinneret pres. phar. denticulate; oes. without med. bulb; post. ovary undev.; marine *Dasyilaimus* Cobb 1933Spinneret 0; post. pharynx concave-cylind.; oes. with med. bulb; in sheep dung *Rhabditoides* Goodey 1929

Ovary 1

Gonad ♀ outstretched, anterior; cephalic setae 4.....*Walcherenia* de Man 1921

Gonad ♀ reflexed; cephalic setae 0 or minute

Spinneret present; pharynx denticulate; striae of dots; post. ovary undev.; marine *Dasyilaimus* Cobb 1933

Spinneret 0; pharynx not denticulate; oesophagus cephaloboid; not marine

Probolae present; amphid at lips; in soil

Cephalic appendages in 2 circlets.....*Acrobeles* Linst. 1877Cephalic appendages in 1 circlet.....(*Acrobeloides* (Cobb) Steiner & Bührer 1933)

Probolae absent; cephalic appendages none or in 1 circlet

Pharyngeal wall a long. series of elements; onch. 0; vagina directed inward. *Panagrolaimus* Fuchs 1930

Pharyngeal wall not of serial elements; onch. oft pres.; vagina directed ±forward; Turbatricinae

Posterior pharynx closed; vulva ±75%; vagina less muscular; tail ±blunt.....*Plectonchus* Fuchs 1930Posterior pharynx ±open; vulva ±65%; vagina muscular; tail slender.....*Turbatrix* Peters 1927(=*Anguillula* auctores)**Amphid known**

External amphid not spiral, circular nor elliptical

Habitat not marine

Spinneret present; cephalic setae 4; in fresh water; ♀.....*Chronogaster* Cobb 1913

Spinneret none; ovaries 2

Tail setaceous, 19%; amph. semicirc. oesoph. with med. bulb; vulva 21%; ♀ (*Aulolaimoides* Micol. 1915)

Tail not setaceous, 9-15%; amphid inconspicuous, small or slit-like

Med. oesoph. bulb confluent in front; post. phar. closed; in decaying matter; ♀.....(*Cheilobus* Cobb 1924)(=*Rhabditophanes* Fuchs 1930)Med. oesoph. bulb none; pharynx ovoid; wings 0; amphid a slit; in soil; ♀.....*Monhystera* Bast. 1865(=*M. dintheriana* de Man 1885)

Habitat marine; cephalic setae present; spinneret present

Ambulatory tubes forming ventral pre-anal sole; nema ±S-shape, narrowed in card. reg.; marine; ♀

Oesoph. region hardly swollen; card. bulb faint; cerv. striae all alike. *Notochaetosoma* Irwin-Smith 1918

Oesoph. region swollen, ovoid; oesophageal bulbs 1 or 2; band of cervical

striae accentuated, exc. *Tristicochaeta falcatum*Ventral ambulatory adhesion tubes or setae in 2 rows.....*Drepanonema* Cobb 1933(=*Chaetosoma* Clap. 1863)Ventral ambulatory adhesion tubes or setae in 3 or 4 rows.....*Tristicochaeta* Panceri 1878(=*Draconema* Cobb 1913)

Ambulatory tubes absent; nema not crooked, not narrowed in cardiac region

Amphid transverse; pharynx minute; annules plain; ceph. set. 10, jointed; ♂ only *Leptonemella* Cobb 1920

Amphid large, folded or shepherd's crook; pharynx obvious; ovaries outstretched

Pharynx 4-5%, heavy, tubular; amphid 4-50% body; ceph. set. 4, subceph. 12-♀ (*Pseudolella* Cobb 1920)

Pharynx under 2%, conoid, not heavy; cephalic setae large, long; odontia 6 or 12

Ovary 1, posterior; cephalic setae 6; odontia 12; vulva 15%.....(*Synodontium* Cobb 1920)

Ovaries 2; cephalic setae 4; odontia 6; vulva 45-67%

Tail clavate; nema shorter, stouter; amphid larger.....*Axonolaimus* de Man 1889Tail conoid; nema long, slender; amphid smaller.....*Ascolaimus* Ditlev. 1919(=*Bathylaimus* Ditlev. 1918)

External amphid spiral, circular or elliptical

Female not known

Habitat soil (possibly marine); spinneret present; amphid spiral; ceph. set. 0....(*Amphispira* Cobb 1920)

Habitat marine

Amphid multispire; wings 0; annules very fine

Pharyngeal bulb none; lip region not set off; head truncate-conoid; onch. 3, small *Alaimonema* Cobb 1920

Pharyngeal bulb present

Head cylindrical; lip region discoid, set off by constriction.....(*Choniolaimus* Ditlev. 1918)Head hemispherical; lip region not discoid.....(*Amphispira* Cobb 1920)

Amphid circular or monospire; spinneret present

Nema cephalated, striae ceasing at constriction

Pharynx none; head ±spherical; amphid behind head; nema slender, 1.4%....*Cinotonema* Cobb 1920Pharynx definite; head ±hemispherical; amphid on head; nema wider, 3.8% (*Micromicron* Cobb 1920)

Nema not cephalated

Annules strong, each of 8 tile-like elements; oesophagus compound.....*Dasynemella* Cobb 1933(=*Dasyneema* Cobb 1920)

Annules none; oesophagus not compound

Cephalic setae 8 or more; onchium minute; amphid minute, at lips.....*Catanema* Cobb 1920

Cephalic setae 4; onchia 0; amphid large, not at lips

Wing single; amphids on head, large, double, circular, joined dorsally.....*Antomicron* Cobb 1920

Wings 0; amphids behind head, not joined

Base pharynx faintly cuticularized, without ring; amphid circular.....*Monhysteriella* Kreis 1929Base pharynx with cuticular ring, prongs 2, vertical; amphid monospire. *Cricolaimus* Southern 1914

Female known

Ovary 1 (for alternative see next page)

Gonad ♀ outstretched, anterior; amphid circular; striae 0, or fine and plain; wings 0

Pharynx ±quadrate; neck under 5%; crystalline bodies lateral; marine.....(*Crystallonema* Cobb 1920)Pharynx long-conoid; neck 17%; spinneret present; in fresh water.....(*Monhystera* Bast. 1865)(i.e. subg. *Monhysterella* Cobb 1918)

Gonad ♀ reflexed; cephalic setae 0 or minute

- Spinneret present; striae of dots; phar. denticulate; post. ovary undev.; marine *Dasyllaimus* Cobb 1933
 Spinneret absent; pharynx not denticulate; not marine
 Head with 6 biting odontia; amphid round; nema cephalated exc. *Teratocephalus* de Man 1876
 Head with entire or 3-parted lip region; amphid small, at lips; nema not cephalated, exc. *Acrobeles insubricus*
 Probolae absent; cephalic appendages 0 or in 1 circllet. *Panagrolaimus* Fuchs 1930
 Probolae present; in soil
 Cephalic appendages in 2 circllets. *Acrobeles* Linst. 1877
 Cephalic appendages in 1 circllet. *Cephalobus* Bast. 1865
 (= *Acrobeloides* (Cobb) Steiner & Bührer 1933)
- Ovaries 2 (for alternative see preceding page)
 Gonads ♀ outstretched
 Amphid multispire; striae fine; pharynx small, cup-shaped
 Male ventral pre-anal supplementary organs small. (*Parasabatieria* de Man 1907)
 Male supplementary organs none. (*Sabatieria* de Rouville 1903)
- Amphid circular
 Pharynx oboconoid, cutic. rings 1 or more; spinneret pres.; brackish soil. (*Desmolaimus* de Man 1880)
 Pharynx without transverse cuticularized rings; spinneret present or not
 Habitat not marine; cephalic setae 0
 Spinneret present; head with 4 circular "warts"; in fresh water. *Pseudochromadora* Daday 1901
 Spinneret absent; head without "warts"; bursa present; in moss. *Himatidiophila* Rahm 1925
 Habitat marine; striae fine; cephalic setae present
 The pharynx oboconoid, with cylindroid vestibule
 Base pharynx cuticularized; head ± cylindroid; cardiac bulb valvate. *Metalinhomoeus* de Man 1907
 (cf. *Deltanema* Kreis 1929)
 Base pharynx not cuticularized; head ± conoid; lips if everted knob-like *Cryptolaimus* Cobb 1933
 The pharynx minute or none
 Cardiac swelling not strongly developed; ceph. set. 8-10; nema not tapering *Linhomoeus* Bast. 1865
 Cardiac bulb broad, valvate; ceph. set. 4; nema tapering, tail setaceous *Terschellingia* de Man 1888
- Gonads ♀ reflexed
 Habit parasitic, in insects; oesophagus without median bulb. *Neoaplectana* Steiner 1929
 Habit free-living
 Spinneret absent; cephalic setae 0; in soil and fresh water
 Nema cephalated, exc. *T. palustris*; lip reg. lobed, with 6 biting odontia. *Teratocephalus* de Man 1876
 Nema not cephalated; lip region not lobate; bursa present. *Himatidiophila* Rahm 1925
- Spinneret present
 Cephalic setae 0; head with 4 circular "warts"; in fresh water. *Pseudochromadora* Daday 1901
 Cephalic setae present; head without "warts"
 Habitat fresh water or soil; amphid circular to monospiral. *Plectus* Bast. 1865
 Habitat marine
 Lips 2, mouth a dorso-ventral slit; "pseudonchs" dorso-ventral ridges. (*Pseudonchus* Cobb 1920)
 Lips more than 2, mouth not a slit; "pseudonchs" absent
 Amphid multispire
 Pharynx irregular, with tooth-like projections. *Zalonema* Cobb 1920
 Pharynx simply conoid, regular, without tooth-like projections
 Head set off by absence of striae. *Heterodesmodora* Micol. 1924
 Head not set off; striae extending to lips; neck 6-7%; onchium minute. *Spirina* Fil. 1918
 (= *Spira* Bast. 1865)
- Amphid circular or monospiral
 The pharynx denticulate; post. ovary undev.; striae of dots; wings pres. (*Dasyllaimus* Cobb 1933)
 The pharynx not denticulate; both ovaries developed
 Dorsal pharyngeal wall not thickened; amphid not at lips
 Neck 14-21%; wings present; amphid long-oval; head set off by groove at 4 cephalic setae
 Length of neck 14-16%; amphid large, $\frac{1}{2}$ head-width back. *Eutelolaimus* de Man 1922
 Length of neck 21%; amphid 2 head-widths back. *Aplectus* Cobb 1914
 Neck under 7%; wings 0; amphid spiral
 Nema 4-7 mm.; width 0.3-1%; phar. straight, closed, with pharyng. swelling *Laxus* Cobb 1894
 Nema 2-4 mm.; width 1.5-4%; pharynx conoid, oft with small onchium. *Spirina* Fil. 1918
 (= *Spira* Bast. 1865)
 Dorsal pharyng. wall thick, tooth-like; amph. monospire at lips; ceph. set. 4; renette far back
 Oesoph. glands obscuring junct. of intest.; dorsal phar. element ± uniform *Onchium* Cobb 1920
 Oesoph. glands not prominent; dorsal pharyng. element not uniformly thick, but heavy
 Tooth-like thickening parallel to axis. *Camacolaimus* de Man 1889
 (= *Digitonchus* Cobb 1920
 = *Acontiolaimus* Fil. 1918)
- Tooth-like thickening bent outward from beginning of vestibule
 Point of "onchium" single. *Acmaeolaimus* Fil. 1918
 Point of Y-shaped "onchium" duplex. *Ypsilon* Cobb 1920
- Oesophagus plain, i.e. without median or posterior bulb or swelling (for alternative see preceding page)
 Amphid not known or obscure (for alternative see next page)
 Female not known
 Habitat marine; lip region discoid; odontia 6; ceph. set. 4; amphid small. (*Stephanolaimus* Ditlev. 1918)
 Habitat fresh water, marsh or sand
 Hemispherical "cap" wider than neck, edge setose; phar. conoid; spinneret pres. *Diplolaimus* Linst. 1876
 Hemispherical cap none; setae 0; spinneret 0; wings double; bursa striate. *Macropontonia* de Man 1880
- Female known; spinneret present
 Ovary 1; marine
 Gonad ♀ reflexed; ceph. set. 0, pharynx denticulate; post. ovary undev.; wings pres. *Dasyllaimus* Cobb 1933
 Gonad ♀ outstretched; cephalic setae 4
 The ovary anterior; nema 3 mm.; neck 12%; vulva 75%. *Krampia* Ditlev. 1922
 The ovary posterior; nema 1 mm.; neck 19%; vulva 35%. *Asymmetrica* Kreis 1929
- Ovaries 2, reflexed
 Habitat fresh water (for alternative see next page)
 Lips 3, rounded; pharynx small, closed; onchium minute, deepset. (*Trippyla* Bast. 1865)
 (= *Promononchus* Micol. 1923)

- Lips 6, thin; pharynx larger, domed; onchia in posterior chamber (*Trilobus* Bast. 1865)
 Habitat marine (for alternative see preceding page)
 Wings conspic.; striae of dots; cephalic set. 0; pharynx denticulate; post. ovary undeveloped... *Dasytlimus* Cobb 1933
 Wings 0, striae plain, not altered laterally; cephalic setae present; Enoplidae
 Calvarium elaborate, ornate; onchia often present..... *Deontostoma* Fil. 1916
 Calvarium not elaborate, sometimes absent
 Oesophagus crenate towards base; calvarium simple or reduced
 Eyes 0; neck \pm tapering; calvarium reduced; cervical setae in several rows... *Stenolaimus* Marion 1870
 Eyes with lenses
 Pharynx simple; calvarium reduced; neck tapering..... *Nasinema* Fil. 1927
 Pharynx with tooth-like processes; calvarium simple; neck \pm cylindrical... *Phanoderma* Bast. 1865
 (= *Heterocephalus* Marion 1870)
 Oesophagus not crenate; calvarium absent
 Jaws 3, without mandibles or onchia..... (*Paronoplus* Fil. 1927)
 Jaws absent
 The pharynx tapering into oesophagus; cephalic setae 6, cervical small. *Parozystomina* Micol. 1924
 The pharynx broad, thin-walled, base truncate with mound; amph. with pocket *Palagonema* Cobb 1894
 (cf. *Vasculonema* Kreis 1928)
- Amphid known (for alternative see preceding page)
 External amphid not spiral, circular nor elliptical; female known
 Ovary 1; amphid not linear
 Gonad ♀ outstretched; lips 6, thin, each with 3 parallel-sided cutic. ribs; marine *Daytonema* Cobb 1920
 (cf. *Theristus* Bast. 1865)
 Gonad ♀ reflexed
 The ovary posterior; spinneret absent; marine..... (*Rhabdocoma* Cobb 1920)
 The ovary anterior, vulva 80%; spinneret present; soil about sugar cane..... *Trischistoma* Cobb 1913
- Ovaries 2
 Gonads ♀ outstretched; marine exc. Monhystera
 Habitat soil among roots; amphid a transverse slit; pharynx ovoid..... *Monhystera* Bast. 1865
 (= *M. dintheriana* de Man 1885)
- Habitat marine
 Amphid large, long-oval, on triangular plate, \pm as wide as head; spinneret none.... *Didelta* Cobb 1920
 Amphid not on special plate
 Mouth ventrad of axis; amphid large, long, "folded"..... *Campylaimus* Cobb 1920
 Mouth axial; amphid shorter
 Odontia absent; amphid flattened, more or less reniform..... *Margonema* Cobb 1920
 Odontia, if present, 6 or 12; amphid hook-shaped or "folded"
 Tail clavate; body less slender; amphid larger..... *Axonolaimus* de Man 1889
 Tail conoid; body slender; amphid smaller..... *Ascolaimus* Ditlev. 1919
 (= *Bathylaimus* Ditlev. 1918)
- Gonads ♀ reflexed
 Habitat freshwater
 Pharynx larger, domed; lips 6, thin; onchia small, in posterior pharynx..... (*Trilobus* Bast. 1865)
 Pharynx small, closed; lips 3, rounded; onchium minute, deepset..... (*Tripylla* Bast. 1865)
 (= *Promononchus* Micol. 1923)
- Habitat marine
 Wings present; amphid a transverse slit almost as wide as head..... *Actinonema* Cobb 1920
 Wings absent; amphid with internal pocket (*Rhabdocoma* ?); Enoplidae (*Rhabdocoma* ?)
 Oesophagus crenate towards base
 Pharynx without tooth-like processes; cervical setae in longitudinal rows... *Stenolaimus* Marion 1870
 Pharynx with 2-3 forward-pointing tooth-like processes; calvarium pres.... (*Phanoderma* Bast. 1865)
 (= *Heterocephalus* Marion 1870)
- Oesophagus not crenate
 Calvarium present; onchia often 1-3
 The calvarium elaborate, ornate, relatively short..... *Deontostoma* Fil. 1916
 The calvarium a mere band, margin wavy; external amphid transverse-oval *Cycolaimus* de Man 1889
 Calvarium absent
 Lateral series of setae on neck absent; spinneret absent..... (*Rhabdocoma* Cobb 1920)
 Lateral series of setae on neck of few members
 Excretory pore on projection; pharynx definite, \pm quadrate; testis 1.... *Paranticoma* Micol. 1930
 Excretory pore not raised; pharynx \pm conoid, often obscure; testes 2
 Gubernaculum present; ♂ pre-anal ventral supplementary organ tubular *Anticomma* Bast. 1865
 Gubernaculum none; ♂ ventral pre-anal supplementary organ papilloid *Anticomopsis* Micol. 1930
- External amphid spiral, circular or elliptical
 Female not known; marine exc. *Dintheria* (for alternative see next page)
 Spinneret none; cephalic set. 4, long; amph. transverse-oval, $\frac{1}{3}$ head-width, open caudad *Dintheria* de Man 1921
 Spinneret present (*Neurella*, *Bognenia* ?)
 Nema broad, tapering from head to tail; wings 12-24; amphid spiral..... *Pteronium* Cobb 1933
 Nema slender, not widest at head; wings none or few
 Amphid multispire
 Wing single; cephalic setae 4; amphid broad, of few winds; spinneret(?)..... *Newrella* Cobb 1920
 Wings absent; striae not altered laterally
 Onchia 3 at tops apophyses; cephalic set. papilloid; pharynx complex, bulb strong *Trogolaimus* Cobb 1920
 Onchia 0; cephalic setae longer; lips flap-like, at least distally
 Lip supports thickened; phar. cyathiform; ♂ supplements many, chromadoroid *Dispira* Cobb 1933
 Lip supports none; pharynx spheroid, on front of head; ♂ supplements 0. *Dispirella* Cobb 1933
- Amphid circular, elliptical or monospiral
 Wings strong; amphid circular or monospiral
 Oesoph. compound, wide anteriorly, card. bulb faint; amph. spiral; annules wide *Dasynemella* Cobb 1933
 (= *Dasynema* Cobb 1920)
 Oesoph. not compound; amphid circular
 The wings 12, of spine-like markings..... *Xenolaimus* Cobb 1920
 The wing single, with subordinate lines; amphids joined dorsally..... (*Antomicron* Cobb 1920)
 Wings 0; striae not altered laterally
 Onchium \pm spear-like; cephalic setae 10; cephalic cuticle thickened; eyes 2 (*Cophonchus* Cobb 1920)
 Onchium none (*Nudolaimus* ?)

- Nema 9-10 mm.; striae 0; pharyngeal walls heavily cuticularized; amphid small
 Amphid labial, spiral; phar. large, processes basal; ceph. set. 4; eyes 0... (*Nudolaimus* Allgen 1929)
 Amphid 1 head-width back, oval; phar. small; ceph. set. 10; eyes 2... (*Leptosomatides* Fil. 1918)
 Nema under 3 mm.; cephalic setae 4-6
 Cephalic setae at least 2 head-widths long
 Lip region discoid, set off; odontia 6; amphid small, oval... (*Stephanolaimus* Ditlev. 1918)
 Lip region not discoid; odontia none; amphid large, circular... (*Cenolaimus* Cobb 1933)
 Cephalic setae not over 1 head-width long; lip region not discoid; odontia 0
 Pharynx a wide tube; head strongly conoid; width of nema 8%... (*Tubolaimus* Allgen 1929)
 Pharynx shallow; head not strongly conoid
 Cuticle annulated; amphid oval, near lips; lips 3, massive, low... (*Zygonemella* Cobb 1920)
 Cuticle with fine striae or none; amph. circular, ± 1 head-width back; lips \pm confluent
 Ribs of pharynx 6, cuticularized, parallel to axis... (*Rhadinema* Cobb 1920)
 Ribs of pharynx none; pharynx conoid... (*Bognemia* Allgen 1932)
 Female known (for alternative see preceding page)
 Ovary 1; amphid not multispire
 Gonad ♀ reflexed; amphid circular or elliptical
 The ovary posterior; marine
 Spinneret none; cephalic setae 4; tail usually about 40%... (*Rhabdocoma* Cobb 1920)
 Spinneret present; cephalic setae 0; tail about 3%... (*Nemanema* Cobb 1920)
 The ovary anterior; cephalic setae present (in *Monhystrium* sometimes papilloid)
 Habitat not marine; spinneret present
 Pharynx deep; onch. small, basal; lips massive, with flaps; vulva 80%; soil (*Trischistoma* Cobb 1913)
 Pharynx less deep, vestibule definite; onchia 0; in gills of land crabs... (*Monhystrium* Cobb 1920)
 Habitat marine
 Spinneret 0; ceph. set. 10; caudal setae thorn-like; amphid large, center raised (*Zanema* Cobb 1920)
 Spinneret pres.; ceph. set. 4, small; thorn-like setae 0; phar. \pm cylind... (*Conolaimella* Allgen 1930)
 Gonad ♀ outstretched, anterior; usually marine
 Spinneret none; wings 0; amphid circular; marine
 Amphid large, with central mound; caudal setae thorn-like; nema 0.9% wide... (*Zanema* Cobb 1920)
 Amphid not raised centrally; caudal setae if present not thorn-like
 Labial palps 6; pharynx \pm quadrate; tail conoid, 4%... (*Crystallonema* Cobb 1920)
 Labial palps 0; pharynx conoid then spheroid; tail long, filiform... (*Diplolaimella* Allgen 1929)
 Spinneret present; marine exc. *Monhystrium*, and sometimes *Monhystera* and *Theristus*
 Lips 3, each with finger-like odontium; striae coarse; wings 32, fewer towards tail *Xyala* Cobb 1920
 Lips without finger-like processes; wings 0 or few
 Mouth cavity large, complex, of several chambers, the first of radial elements; carnivorous
 Base pharynx unsymmetrical, meeting oesoph. middle chamber denticulate *Sphaerolaimus* Bast. 1865
 Base pharynx symmetrical, oesoph. surrounding 2 chambers; dents. 0 *Parasphaerolaimus* Ditlev. 1918
 Mouth cavity smaller, of 1 or 2 chambers, walls relatively simple
 Lips 6, conoid, twice high as broad, with hook-shaped setae; amph. monospire *Dactylaimus* Cobb 1920
 Lips not as high as broad, without hook-shaped setae; mostly *Monhysterinae*
 Caudal glands pre-anal, reaching vulva; excret. pore nr. phar.; ceph. set. 4 *Conolaimella* Allgen 1930
 Caudal glands post-anal; excret. pore nr. nerve ring, when seen; ceph. set. of many
 Labial ribs 18, cuticularized, parallel-sided; testes 2... (*Daptonema* Cobb 1920
 (cf. *Theristus* Bast. 1865))
 Labial ribs absent or (*Monhystrium*) reduced; genera resembling *Monhystera*, exc. *Sphaerocephalum*
 Margin of transverse-oval amphidial area not definite, sensory papilla at posterior margin
 Intestine narrow; 1 body-width; gubernaculum tubular... (*Leptogastrella* Cobb 1920)
 Intestine wider(?); gubernaculum not tubular(?)... (*Paramonhystera* (Steiner) Fil. 1918)
 Margin of amphidial area definite, sensory papilla not obvious
 Habitat gills of land crabs; cephalic setae 10, sometimes papilloid... (*Monhystrium* Cobb 1920)
 Habitat aquatic; marine exc. sometimes *Monhystera* and *Theristus*
 Pharynx conoid, then spheroid; striae fine or 0; eyes oft present... (*Diplolaimella* Allgen 1929)
 Pharynx without spheroid chamber; amphid circular exc. *Sphaerocephalum*
 Amphid monospire; neck 5-9%; spinneret wide, plug striated... (*Sphaerocephalum* Fil. 1918)
 Amphid circular; neck 10-27%; spinneret without striated plug
 Oesophagus wide in post. half; spicula hamate distally; gubernaculum 0 *Austronema* Cobb 1914
 Oesophagus not wider posteriorly; gubernaculum present
 Setae on head 8-fold; somatic set. oft long... (*Steinera* (Micol.) Stekhoven & de Coninck 1933)
 Setae on head not 8-fold; long paired somatic setae absent
 Gubernaculum tubular, barbed, apophysis none... (*Penzancia* (de Man) Fil. 1918)
 Gubernaculum not tubular
 Spicula not much bent; gubernac. apophysate; mostly freshwater... (*Monhystera* Bast. 1865
 (= *Tachyhodites* Bast. 1865))
 Spicula much bent or sickle-shape; gubernac. plate-like, dorsad... (*Theristus* Bast. 1865
 (= *Allomonhystera* Micol. 1923)
 (cf. *Daptonema* Cobb 1920))
 Ovaries 2
 Gonads ♀ outstretched; marine (for alternative see next page)
 Nema 12-18% wide; neck "collared"; amphid spiral; wings many; spinneret present
 Wings over 60; pharynx narrow; spicula 2, equal... (*Richtersia* Steiner 1916)
 Wings 16-20, in pairs; pharynx capacious, open; spicula unequal, or but 1... (*Richtersiella* Kreis 1929)
 Nema under 3% wide; neck not "collared"; wings if present not numerous
 Spinneret absent; amphid circular exc. *Didelta*
 Setae none; amphid \pm half as wide as head... (*Metalaimus* Kreis 1928)
 Setae present; amphid small exc. *Didelta*
 Amphid very large, long-oval, on triangular plate; cephalic setae 4... (*Didelta* Cobb 1920)
 Amphid small, not on plate; cephalic setae 6 or more
 Pharyngeal bulb distinct; cephalic setae 6, stout, jointed... (*Cytolaimium* Cobb 1920)
 Pharyngeal bulb none; setae on head 18, small; amph. with faint multispire... (*Anticgathus* Cobb 1920)
 Spinneret present; cephalic setae present exc. *Margonema*
 External amphid multispire
 Male ventral pre-anal supplementary organs papilloid... (*Parasabatieria* de Man 1907)
 Male supplementary organs none... (*Sabatieria* de Rouville 1903)
 External amphid not multispire (*Linhomoella* faintly so)

- Amphid wider than long, \pm reniform; lips 3, double, with papillae.....*Margonema* Cobb 1920
 Amphid circular or monospiral
 Lips distinct, with distal flaps; ceph. set. 6, stout, jointed; papillae 6....*Cytolaimium* Cobb 1920
 Lips absent or confluent
 Oesoph. & lumen with oval enlargement behind eyes; eyes sometimes none *Araeolaimus* de Man 1888
 Oesoph. without median enlargement; eyes 0; cephalic setae 6-10
 Setae on head 10, shorter 4 specialized at tip; amph. faint multispire. (*Linhomoea* Cobb 1920)
 Setae on head 6-8, without specialized tips; amphid circular.....*Linhomoeus* Bast. 1865
 Gonads ♀ reflexed (for alternative see preceding page)
 Habitat not marine exc. sometimes *Aphanolaimus*
 Spinneret absent; amphid multispire; setae 0; striae of dots; in sand dunes. *Choanolaimus* de Man 1880
 Spinneret present
 Wings present; contour crenate; glands lateral, serial; amphid monospire. *Aphanolaimus* de Man 1880
 Wings 0, striae not altered laterally; contour not crenate; amphid circular or oval
 Lips, papillae & setae 0; amphid round; pharynx small, conoid, unarmed *Diplohystra* de Cillis 1917
 Lips, papillae & small setae pres.; amph. small, oval; phar. closed; onch. basal *Trippyla* Bast. 1865
 (= *Promononchus* Micol. 1923)
 Habitat marine
 Amphid spiral; spinneret present
 Pharynx with apophyses forward and backward from mid-region
 Mid-pharyngeal region not denticulate; wings numerous, markings V-shaped....*Nunema* Cobb 1933
 Mid-pharyngeal region denticulate.....*Halichoanolaimus* de Man 1886
 (= *Smalsundia* Allgen 1929)
 Pharynx without forward apophyses; mid-pharyngeal region not denticulate
 Wings numerous; neck region "collared"; nema 12-16% wide
 Spicula 2; wings over 50; pharynx narrow.....*Richtersia* Steiner 1916
 Spicula unequal, or 1 only; wings 16-32; pharynx capacious, open.....(*Richtersiella* Kreis 1929)
 Wings absent, striae if present not altered laterally
 Jaws 3, with backward apophyses and numerous horny teeth anteriorly.....*Demonema* Cobb 1894
 Jaws absent, pharynx without apophyses
 Pharynx 2 chambers; ceph. set. 10, oft jointed, tips specialized; striae fine *Trippylodes* de Man 1886
 Pharynx 1 chamber; tips setae unspecialized; striae less fine; eyes present
 Cuticular pores none; pharynx with definite bulb.....*Nannolaimus* Cobb 1920
 Cuticular pores present; pharynx without definite bulb.....*Cyatholaimus* Bast. 1865
 (= *Necticonema* Marion 1870)
 Amphid circular, elliptical or monospiral
 Spinneret absent; cephalic setae 4, stout; tail usually \pm 40%.....*Rhabdocoma* Cobb 1920
 Spinneret present
 Oesophagus crenate towards base; calvarium reduced
 Cervical setae in long. rows; pharynx \pm none; ext. amph. \pm semi-circular (*Stenolaimus* Marion 1870)
 Cervical setae 0.....*Phanodermopsis* Ditlev. 1926
 (= *Galeonema* Fil. 1927)
 Oesophagus not crenate
 Calvarium present; onchia often 1-3
 The calvarium elaborate, ornate.....*Deontostoma* Fil. 1916
 The calvarium a wavy band; often denticles at lips; amphid transverse. *Cytilolaimus* de Man 1889
 Calvarium absent
 Jaws 3, with simple supporting framework.....*Paronoplus* Fil. 1927
 Jaws absent
 Pharynx of 2 chambers
 Chambers of pharynx \pm equal, small; cephalic setae 0.....*Pseudopelagonema* Kreis 1932
 Chambers of pharynx unequal, anterior large; setae 10, often jointed, tips specialized
 Both chambers \pm conoid or irregular in shape.....*Trippylodes* de Man 1886
 Both chambers \pm parallel-sided.....*Bathylaimus* Cobb 1894
 (= *Macrolaimus* Ditlev. 1918
 = *Cothonolaimus* Ditlev. 1919)
 Pharynx of 1 chamber; setae without specialized tips
 Dorsal pharynx. wall thick, tooth-like; amph. monospire at lips; ceph. set. 4; renette far back
 Oesoph. glands obscuring junct. of intest.; dorsal phar. element \pm uniform *Onchium* Cobb 1920
 Oesoph. glands not prominent; dorsal pharyng. element not uniformly thick, but heavy
 Tooth-like thickening parallel to axis.....*Camacolaimus* de Man 1889
 (= *Digitonchus* Cobb 1920
 = *Acontiolaimus* Fil. 1918)
 Tooth-like thickening bent outward from beginning of vestibule
 Point of "onchium" single.....*Acmaeolaimus* Fil. 1918
 Point of Y-shaped "onchium" duplex.....*Ypsilon* Cobb 1920
 Dorsal pharyngeal wall not thickened; amphid not at lips; Enoplidae, exc. *Aphanolaimus*
 The pharynx open, capacious, walls usually heavily cuticularized; amphid with pocket
 External amphid circular; wall of pharynx irregularly thick.....(*Pandolaimus* Allgen 1929)
 External amphid small, transverse; pharyng. wall thin. base with mounds *Pelagonema* Cobb 1894
 (cf. *Vasculonema* Kreis 1928)
 The pharynx small or very narrow
 Cervical setae absent; cephalic setae 4, in front of amphid.....*Aphanolaimus* de Man 1880
 Cervical setae lateral, in short longitudinal series; amphid with pocket
 Gubernaculum present; ♂ supplementary organ tubular.....*Anticoma* Bast. 1865
 Gubernaculum none; ♂ supplementary organ papilloid.....*Anticomopsis* Micol. 1930
 Cavity more or less cylindrical or prismoid, i.e. parallel-sided in long. section (for alt. see page 20)
 Oesophagus with median or posterior bulb or swelling, or both (for alternative see page 27)
 Amphid not known or obscure (for alternative see next page)
 Female not known
 Spinneret none; ceph. set. 0; phar. closed; oesoph. cephaloid; neck over 20%; in swamp *Litonema* Cobb 1920
 Spinneret present; setae on head numerous; neck under 5%; amphid small, near lips; marine
 Pharyngeal bulb present; striae none; length 3-4 mm.; onchia(?) 6, minute.....(*Catanema* Cobb 1920)
 Pharyngeal bulb none; striae annulating the cuticle; length 10 mm.....(*Lazonema* Cobb 1920)
 Female known; spinneret none exc. *Rhabdolaimus*; not marine; phasids usually present

- Ovary 1, anterior; cephalic setae 0 or minute
 Gonad ♀ outstretched
 Pharynx long, tubular, base halfway to nerve ring; spear none; median bulb 0. *Gymnolaimus* Cobb 1913
 Pharynx base $\pm \frac{1}{2}$ distance to nerve ring; spear reduced; oes. bulb strong (*Aphelenchoides* Fischer 1894)
 (= *Seinura* Fuchs 1931
 = *Parasitaphelenchus* Fuchs 1930
 = *Pathoaphelenchus* (Cobb) Steiner 1931)
- Gonad ♀ reflexed
 Median oesophageal bulb present
 Pharynx wide as long; cardiac bulb not valvate, median valvate; bursa none. *Lycolaimus* Rahm 1928
 Pharynx much longer than wide; cardiac bulb valvate; bursa present. *Rhabditis* Duj. 1845
 (= *Diploscapteroides* Rahm 1928
 = *Pseudorhabditis* Kreis 1929)
- Median oesophageal bulb absent; in soil and fresh water
 Cardiac swelling long, non-valvate; pharyng. rods heavily cuticular. *Diphtherophora* de Man 1880
 (= *Chaolaimus* Cobb 1893
 = *Archionchus* Cobb 1913)
- Cardiac bulb valvate, preceded by isthmus, i.e. oesoph. cephaloboid; pharyng. rods none
 Head end lobed by 6 large biting lips or odontia; carnivorous. (*Teratocephalus* de Man 1876)
 Head end not lobed, exc. in *Acrobeles* insubricus
 Pharynx wall simple, forming 2 chambers; onchium oft present; ceph. set. 0-4 *Macrolaimus* Maupas 1900
 Pharynx wall of a longitudinal series of elements, i.e. cephaloboid
 Probolae on head none. *Panagrolaimus* Fuchs 1930
 Probolae on head in 1 or 2 circlets
 Circlets of probolae 2, elements often elaborate. *Acrobeles* Linst. 1877
 (= *Cephalobus* Bast. 1865
 Circlet of probolae single. (= *Acroboloides* (Cobb) Steiner & Bührer 1933)
- Ovaries 2; cephalic setae 0, exc. in *Teratocephalus* minute
 Gonads ♀ outstretched; ceph. set. 0; phar. tubular, reaching oesoph. bulb; in soil *Aulolaimus* de Man 1880
 Gonads ♀ reflexed
 Spinneret present; wings absent; in fresh water and moist soil. *Rhabdolaimus* de Man 1880
 Spinneret absent
 Cuticle with prominent tubercles; lip reg. set off by constrict.; ceph. set. with saccate bases
 Tubercles encrusting dorsal surface; wings broad, thin; in moss. *Craspedonema* Richters 1908
 Tubercles in longitudinal row(s), right lateral; oesophagus rhabditoid. *Bunonema* Jägerskiöld 1905
 (= *Rhodolaimus* Fuchs 1930)
- Cuticle without tubercles
 Pharynx a long straight tube, $\frac{1}{2}$ neck length or more
 Oesoph. short, a muscular bulb only; ♂ supplements few, ventral, pre-anal *Aulolaimus* de Man 1880
 Oesoph. long, of 2 parts; ♂ supplements paired, post-anal, i.e. diplogastroid
 Male papillae 9 pairs; gubernaculum sigmoid; labial papillae 18. *Mycolaimus* Cobb 1920
 Male papillae 10 pairs; gubernaculum straight; bursal membrane narrow. *Cylindrogaster* Goodey 1927
 Pharynx not a simple tube, much shorter
 Oesophagus without median bulb
 Cuticularized rods supporting pharynx; shed cuticle surrounding nema *Diphtherophora* de Man 1880
 (= *Chaolaimus* Cobb 1893
 = *Archionchus* Cobb 1913)
- Cuticularized pharyngeal rods none
 Head end lobed by 6 large biting lips or odontia; carnivorous. (*Teratocephalus* de Man 1876)
 Head end not lobed; bursa present. *Rhabditis* Duj. 1845
- Oesophagus with median and cardiac bulbs
 Cardiac bulb valvate; median bulb well set off; glottoid apparatus present
 Wall of pharynx cylindrical; bursa present; ♂ tail rhabditoid. *Rhabditis* Duj. 1845
 Wall of pharynx convex inward; bursa much reduced; ♂ tail long, spicate *Rhabditoides* Goodey 1929
 (= *Rhabditella* (Cobb) Chitwood 1933)
- Cardiac bulb without valve; median bulb valvate; ♂ tail setaceous, diplogastroid
 Pharyng. rods 3, basal bulbs 2, stalked; tail 25-41%; long. striae fine. *Tylopharynx* de Man 1876
 Pharyng. bulbed supporting-rods none; Diplogasterinae
 Nema with evident longitudinal striation. *Diplogaster* M. Schultze 1857
 Nema without longitudinal striation
 Pharynx wide as deep, wall of several elements; ♂ papillae 0(?) *Peronilaimus* Rahm 1928
 Pharynx tubular, wall entire; ♂ papillae usually 5 pairs. *Rhabditoides* Rahm 1928
- Amphid known (for alternative see preceding page)
 External amphid not spiral, circular nor elliptical (for alternative see next page)
 Habitat marine; cephalic setae and spinneret present; amphid crook-shaped, sometimes \pm spiral
 Ambulatory & somatic set. 0; nema not crooked, not narrowed in card. reg.; ventral arm amph. prolonged
 Phar. wide, heavy, 4-5%; amphid long, 4-50% body; ceph. setae 4, subceph. 12; ♀ ... *Pseudolella* Cobb 1920
 Phar. narrow, inconspic.; amph. under 2%; ceph. set. 6, subceph. 4; annules tiled; ♀ *Ceramonema* Cobb 1920
 (cf. *Pselionema* Cobb 1933)
- Ambulatory set. forming ventral pre-anal sole; nema \pm S-shape, narrowed in card. reg.; marine; ♀
 Oesoph. region hardly swollen; card. bulb faint; cerv. striae all alike. *Notochaetosoma* Irwin-Smith 1918
 Oesoph. region swollen, ovoid; oesophageal bulbs 1 or 2; band of cervical striae
 accentuated, exc. *Tristicochaeta* falcatum
 Ventral ambulatory adhesion tubes or setae in 2 rows. *Drepanonema* Cobb 1933
 (= *Chaetosoma* Clap. 1863)
 Ventral ambulatory adhesion tubes or setae in 3 or 4 rows. *Tristicochaeta* Panceri 1878
 (= *Draconema* Cobb 1913)
- Habitat not marine
 Spinneret present; ♀, exc. *Chronogaster*
 Lips 6, distinct, revolute; pharynx 9%; amphid semicircular; soil about cane roots *Anthonema* Cobb 1906
 Lips not revolute; in fresh water
 Cephalic setae 4; pharynx wider, under 1.5%; ♀ ... *Chronogaster* Cobb 1913
 Cephalic setae 0; pharynx long-tubular; tail 25-30%. *Rhabdolaimus* de Man 1880
 Spinneret absent; cephalic setae 0 or papilloid
 Tail setaceous, 19%; oesoph. with 2 bulbs; amph. semicircular; vulva 21%; ♀ *Aulolaimoides* Micol. 1915
 Tail not setaceous, under 15%; oesophagus without median bulb, exc. *Cheilobus*

- Amphid transverse, curved, wide as head; phar. 7.6%; tail 2%; fresh water; ♂ only *Colpurella* Cobb 1920
 Amphid inconspicuous, small or slit-like; tail over 9%; in soil or decaying material
 Median oesoph. swelling before isthmus; cardiac bulb valvate; wings single; ♀'. *Cheilobus* Cobb 1924
 (= *Rhabditophanes* Fuchs 1930)
- Median oesoph. bulb absent
 Pharynx closed; amphid a pore, with internal cuticularized tube; wings 2; J... *Litonema* Cobb 1920
 Pharynx open, or with anterior open chamber; wings 0; ♂ only
 Tail 15%, terminus setaceous; pharynx without cuticularized rods... (*Monhystera* Bast. 1865)
 (= *M. ditheriana* de Man 1885)
 Tail 8%, conoid; cuticularized rods in pharynx; shed cuticle retained... *Diphtherophora* de Man 1880
 (= *Chaolaimus* Cobb 1893
 = *Archionchus* Cobb 1913)
- External amphid spiral, circular or elliptical (for alternative see preceding page)**
- Female not known; spinneret present
 Habitat soil; ceph. set. 0; ceph. cuticle thick, flaps doming vestibule; amph. spiral *Pycnolaimus* Cobb 1920
 Habitat marine; cephalic setae present
 Amphid multispire; annules not coarse; wings absent
 Head cylindrical; lip reg. set off by constriction; pharyng. bulb present... (*Choniolaimus* Ditlev. 1918)
 Head truncate-conoid; lip reg. not set off; pharyngeal bulb none; onch. 3, minute *Alaimonema* Cobb 1920
 Amphid circular, elliptical or monospiral
 Wings 8; annules coarse; oesoph. to isthmus wide, muscular; card. swelling slight *Dasynemella* Cobb 1933
 (= *Dasynema* Cobb 1920)
- Wings 0**
 Cuticularized pharyng. ring basal; ceph. set. 4; ♂ supplements tubular, 14... *Cricolaimus* Southern 1914
 Cuticularized pharyngeal ring none
 Spiculum 1, gubernaculum absent; cephalic setae 10; amphid large, oval... *Pelagonemella* Kreis 1932
 Spicula 2, gubernaculum present (*Chloronemella* ?); cephalic setae in circlets of 4
 Striae coarse; amphid minute; length 10 mm.; width 0.8%... *Lazonema* Cobb 1920
 Striae fine; length under 4 mm.
 Pharyngeal bulb present; amphid at lips; onchia 6, minute... *Catanema* Cobb 1920
 Pharyngeal bulb none; amphid behind pharynx; onchia 0... *Chloronemella* Allgen 1929
- Female known**
- Ovary 1**
 Gonad ♀ outstretched, anterior; amphid circular; wings absent
 Spinneret present; pharynx long-conoid; neck 17%; in fresh water... *Monhystera* Bast. 1865
 (i.e. subg. *Monhystrella* Cobb 1918)
 Spinneret absent; neck not over 5%; marine
 Pharynx wide as deep; cardiac swelling faint, elongate; intestine not dark... *Crystallonema* Cobb 1920
 Pharynx tubular; posterior $\frac{1}{2}$ oesoph. a massive bulb; intestine dark colored (*Solenolaimus* ?)
 Cephalic setae minute; neck 1.4%; anterior oesoph. fusiform; tail rounded... *Solenolaimus* Cobb 1894
 Cephalic setae often as long as mitreiform head is wide; neck 2-5%
 Anterior oesoph. not fusiform; pharynx not evertible; ♂ supplements 0... *Chromagaster* Cobb 1894
 Anterior oesoph. fusiform; phar "spear" evertible; ♂ supplements pres... *Siphonolaimus* de Man 1893
- Gonad ♀ reflexed; wings often present; not marine exc. rarely Rhabditis**
 The ovary posterior, vulva 2%; med. bulb not valvate; amph. semicirc. imud. *Aulolaimoides* Micol. 1915
 The ovary anterior
 Spinneret pres.; ceph. set. 4; card. bulb 0 or faint; amph. monospire; fresh water (*Anonchus* Cobb 1913)
 Spinneret none; setae none, cephalic papillae sometimes setose
 Oesophageal bulbs median and cardiac; pharynx simple; setae 0
 Cardiac bulb valvate; amphid at lips, obscure; wings & bursa pres.; ubiquitous *Rhabditis* Duj. 1845
 Cardiac bulb not valvate, median valvate; amph. transverse-oval, not at lips; phar. cylindrical
 Dorsal onchium none; bursa present; ectoparasitic on beetle borer... *Rhabditolaimus* Fuchs 1915
 Dorsal onch. small, deepset; bursa 0; in exudate chestnut blight... (*Diplogasteroides* de Man 1912)
 Oesophageal bulb cardiac only, preceded by isthmus; phar. of serial plates; bursa 0 or reduced
 Head with 6 large odontia; amphid not obscure, often large
 Nema mostly cephalated; cirri 0; amphid circular; soil & fresh water... *Teratocephalus* de Man 1876
 Nema not cephalated; cirri 6; amph. transverse-oval; tail hooked; in bark *Chambersiella* Cobb 1920
 (= *Diastolaimus* Rahm 1928)
 Head entire or 3-parted at lips; nema not cephalated, exc. *Acrobelus* insubricus; amph. small
 Probolae absent; onchium 0, or 1 inward-pointing... *Macrolaimus* Maupas 1900
 Probolae on head in 1 or 2 circlets
 Circlets of probolae 2, labial and cephalic; amphid a pore near lips... (*Acrobelus* Linst. 1877)
 Circlet of probolae single; amphid often obscure... *Cephalobus* Bast. 1865
 (= *Acrobeloides* (Cobb) Steiner & Bührer 1933)
- Ovaries 2**
 Gonads ♀ outstretched; cephalic setae present; marine or in brackish soil
 Amphid multispire; cephalic setae 10, unequal
 Cardiac bulb ellipsoid; 6 of ceph. set. short; subcephalic setae 4; tail 30%, fine... *Laimella* Cobb 1920
 Cardiac bulb slight; 4 of ceph. set. short, tips specialized; tail 7%... *Linhomoella* Cobb 1920
 Amphid circular or monospiral; tail not over 18%, not setaceous
 Gland ducts cuticularized, piercing cuticle, ventrad, cerv. & pre-anal... *Halaphanolaimus* Southern 1914
 Gland ducts and pores absent; pharynx minute or none
 Pharynx ± none; ceph. set. 10, 4 short, tips specialized; tail 7%... (*Linhomoella* Cobb 1920)
 Pharynx obconoid, when open $\frac{1}{2}$ as wide as head; spinneret sometimes present
 Cuticularized rings in pharynx 1-3; neck 8-14%; marine & brackish soil (*Desmolaimus* de Man 1880)
 Cuticularized pharyngeal rings absent; ventriculus present; marine
 Base pharynx not cuticularized; cardiac bulb oft valvate; head ± conoid *Cryptolaimus* Cobb 1933
 Base pharynx cuticularized; cardiac bulb valvate; head ± cylindrical *Metainhomoeus* de Man 1907
 (cf. *Dellanema* Kreis 1929)
- Gonads ♀ reflexed**
 Habitat not marine, exc. rarely *Rhabditis* and *Haliplectus* (for alternative see next page)
 Oesophageal bulbs median and cardiac; cephalic setae 0; pharynx ± tubular (for alt. see next page)
 Vulva 21%; anus 81%; amph. semicirc., $\frac{1}{2}$ head-width; wings & bursa none (*Aulolaimoides* Micol. 1915)
 Vulva 40-94%; in soil etc., in fresh water, and marine
 Spinneret pres. amph. monospire, protruding; wings & bursa 0; pores lateral... *Haliplectus* Cobb 1913

- Spinneret none; cardiac bulb valvate; amphid obscure; wings and bursa present *Rhabditis* Duj. 1845
 Oesophageal bulb cardiac only (for alternative see preceding page)
 Spinneret absent; amphid circular, elliptical or monospiral
 Head without odontia; nema not cephalated; setae 0; spear minute; in soil (*Triplonchium* Cobb 1920)
 Head with 6 large odontia; bursa none or much reduced
 Nema mostly cephalated; cirri 0; amph. circular; in soil & fresh water. . . *Teratocephalus* de Man 1876
 Nema not cephalated; cirri 6; amphid transverse-oval; tail hooked; in bark *Chambersiella* Cobb 1920
 (= *Diastolaimus* Rahm 1928)
- Spinneret present
 Pharynx simple; oesophagus not plectoid; neck 15-20%
 Cephalic setae 0; wings 0; pharynx long; amphid with pocket; tail 27% (*Rhabdolaimus* de Man 1880)
 Cephalic setae 4; pharynx short; tail 15%
 Wings present; amphid monospire *Paraphanolaimus* Micol. 1923
 Wings absent; amphid circular; onchium minute, not at lips. *Pradesmodora* Micol. 1923
 Pharynx compound-elongate; oes. off with isthmus to card. bulb (plectoid); wings pres.
 Labial region not expanded; cephalic setae usually 4, sometimes 0 or 6; neck 12-30%
 Cardiac bulb not valvate; pharynx long; neck 14-15%; in fresh water & mud *Paraplectus* Fil. 1930
 Cardiac bulb valvate; amphid circular or transverse-oval *Plectus* Bast. 1865
 Labial region expanded, with complicated appendages; pharynx 4-9%; neck 27-31%
 Lips 6, distinct, revolute; striae fine; amphid semicircular; pharynx 9% (*Anthonema* Cobb 1906)
 Lips not revolute; head with dorso-ventral bladders; striae coarse; amph. round; phar. 3-6%
 Cephalic appendages elaborately palmate *Wilsonema* Cobb 1913
 Cephalic appendages forming a closed, striated dome or cupola. *Bitholinema* de Coninck 1931
- Habitat marine or brackish (for alternative see preceding page)
 Ambulatory set. forming a ventral sole, midbody to pre-anal; nema crooked; calvarium unstriated
 Nema narrowest in card. reg., \pm S-shaped; ambulatory set. hollow, pre-anal; annules finer
 Oesoph. region hardly swollen; card. bulb faint; cerv. striae alike *Notochaetosoma* Irwin-Smith 1918
 Oesoph. region swollen, ovoid; oesophageal bulbs 1 or 2; band of cervical striae accentuated, exc. *Tristichocheata falcatum*
 Ventral ambulatory adhesion tubes or setae in 2 rows. *Drepanonema* Cobb 1933
 (= *Chaetosoma* Clap. 1863)
 Ventral ambulatory adhesion tubes or setae in 3 or 4 rows. *Tristichocheata* Panceri 1878
 (= *Draconema* Cobb 1913)
- Nema narrowest midway, epsilon-shaped; ambulatory setae not open at end; annules coarser
 Sole apparently absent; amphid crescentic; annules 98. *Epsilononema* Steiner 1930
 Sole present; amphid circular or monospiral
 Oesophagus with median and cardiac bulbs. *Metepsilonema* Steiner 1927
 Oesophagus with cardiac bulb only
 Annules 80-112; ambulatory setae slender, falcate *Bathypsilonema* Steiner 1927
 Annules 144-192; ambulatory setae straight or slightly arcuate
 Body cylindrical, if crooked faintly so. *Archepsilonema* Steiner 1927
 Body epsilon-like, i.e. body arches well developed
 Cuticular annules homogeneous in structure. *Prochaetosoma* Baylis & Daubney 1926
 (= *Rhabdogaster* Metsch. 1867)
- Cuticular annules not homogeneous in structure
 "Axial" part of the cuticular annules hollow ("frame rings"). *Epsilononema* Steiner 1927
 "Axial" part of the cuticular annules vacuolated, etc. *Epsilononoides* Steiner 1931
- Ambulatory setae 0; nema not crooked; calvarium absent, or not truncate-conoid
 Lips 2, mouth a slit; amph. faint multispire; "pseudonchs" dorso-ventral ridges *Pseudonchus* Cobb 1920
 Lips not 2, mouth not a slit; "pseudonchs" absent
 Amphid multispire; neck 9-13%
 Pharynx deeper than head width, wall irregular, with projections; striae plain *Zalonema* Cobb 1920
 Pharynx tubular, less deep. wall straight; striae of dots *Kreisia* Allgen 1929
 Amphid circular, elliptical or monospiral; neck 14-25%, exc. *Bolbonema* 8%
 Wings present; anterior end set off by groove at 4 ceph. set. & usually by lack of annulation
 Neck 20-25%
 Species marine; striae duplex; wings 2; phar. tubular, inconspic., prob. double *Aplectus* Cobb 1914
 Species in brackish soil & marsh; lip reg. discoid; ceph. set. papilloid. *Leptolaimus* de Man 1876
 Neck 14-16%; marine
 Gland ducts cuticularized, piercing cuticle, cerv. and pre-anal. *Halaphanolaimus* Southern 1914
 Gland ducts & pores 0; pharynx of several long chambers; amphid round *Polyglaimium* Cobb 1920
- Wings absent
 Median bulb present; ceph. set. 0; amph. protruding; neck 12%; pores lateral *Halipectus* Cobb 1913
 Median bulb absent; pores none; marine
 Nema cephalated by expansion opposite large monospiral amphid; neck 8% *Bolbonema* Cobb 1920
 Nema not cephalated by expansion
 Setae on head 6-10, long, flexible; neck under 5%; pharyngeal bulb present. (*Lazus* Cobb 1894)
 Setae on head 0-4; neck 6% or over; pharyngeal bulb absent
 Head constricted at cephalic setae, not at amphid; amphid 2 head-widths back
 Chambers of pharynx several, long; neck 10-11%; cephalic setae 0(?) *Polyglaimium* Cobb 1920
 Chamber of pharynx single; neck 19%; head conoid; cephalic set. 4 *Dermatolaimus* Steiner 1916
 Head not constricted at setae; amphid near lips or not over 1 head-width back
 Ceph. set. 0; phar. a tube reaching card. bulb; amphid conspicuous. *Aegtaolaimus* de Man 1907
 Ceph. set 4; pharynx shorter
 Amphid small, monospiral, at lips; neck 19%; dorsal phar. wall thickened (*Onchium* Cobb 1920)
 Amphid conspicuous, round, not at lips; neck 6%; phar. wall not thick. *Linolaimus* Cobb 1933
- Oesophagus plain, i.e. without median or posterior bulb or swelling (for alternative see page 24)
 Amphid not known or obscure (for alternative see next page)
 Female not known; marine (for alternative see next page)
 Neck over 40%; amphid a long slit; ceph. setae 4 or 8, fine; spinneret oft none. (*Halalaimoides* Cobb 1933)
 Neck 10-30%; amphid not longitudinal; spinneret present (*Trileptium*, *Fimbriella* ?)
 Cephalic setae 0; onchia 3, awl-shaped, almost parallel to axis. *Fimbriella* Cobb 1905
 (= *Fimbria* Cobb 1894)
- Cephalic setae 8 or more
 Onchium dorsal, anterior, inward-pointing; neck 12%; nema 0.5% wide; amph. a slit (*Trileptium* Cobb 1923)
 (= *Trilepta* Cobb 1920)
 Onchium none; neck 22-30%; nema 2-3% wide; cephalic setae usually 8, in 2 circlets

- Calvarium present; setae 8, near base of pharynx; eyes present *Gullmarnia* Allgen 1929
 Calvarium absent; setae minute, around mouth; eyes 0; pharynx small, tubular. *Fimbrilla* Allgen 1929
- Female known (for alternative see preceding page)
- Ovary 1, anterior
- Ceph. set. 4; pharynx large, open; spinneret present; spicula 2-jointed(?); marine. *Filipjeva* Ditlev. 1928
 Ceph. set. 0 or papilloid; pharynx none or narrow; spinneret usually none; -?
 Pharynx a tube reaching halfway to nerve ring; in soil. *Gymnolaimus* Cobb 1913
 Pharynx closed, or marked by reduced spear, not over $\frac{1}{2}$ distance to nerve ring
 Nema free-living, marine; spinneret doubtful; pharynx none or closed. *Litotes* Cobb 1920
 Nema parasitic in insects; spinneret 0; adult spear reduced, base 3-parted. (*Aphelenchulus* Cobb 1920)
- Ovaries 2
- Gonads ♀ outstretched; spinneret present; marine
 The ovaries extending backward; wings 10-12; vulva 45%. *Porocoma* Cobb 1920
 The ovaries opposed; wings 0
 Annules none or not coarse; amphid not saccate, far back; nema not wide. *Anoplostoma* Bütsch. 1874
 Annules coarse; amphid saccate; nema wide; cephalic setae 4
 Number annules ± 17 , secreted annules prominent, separated; head small. *Desmoscolex* Clap. 1863
 Number annules 32-91, secreted annules 0; head concave-quadrate pyramidal. *Tricoma* Cobb 1894
 (= *Quadrucoma* Fil. 1922)
- Gonads ♀ reflexed
 Spinneret none; wings 0; pharynx 2.5-3%, tubular; tail rounded, under 2%; in soil. *Isolaimium* Cobb 1920
 Spinneret present
 Habitat freshwater; lips thick; pharynx closed; amphid small; onch. small, deepset (*Triplya* Bast. 1865)
 (= *Promononchus* Micol. 1923)
- Habitat marine
 Annules ± 17 , secreted annules prominent; amph. saccate; nema wide; cephalic setae 4. *Desmoscolex* Clap. 1863
 Annules none; amphid not saccate; nema less than 5% wide
 Dorsal pharyngeal wall bent inward at lips, onchium-like; cephalic setae 10. (*Asymmetrella* Cobb 1920)
 Dorsal pharyngeal wall not onchium-like; cephalic setae 0. *Angustinema* Cobb 1933
- Amphid known (for alternative see preceding page)
- External amphid not spiral, circular nor elliptical (for alternative see next page)
- Female not known; Enoplidae (*Trileptium*?)
- Head mitreiform, set off by constriction; wings 6-8; spinneret doubtful; marine. *Xennella* Cobb 1920
 Head not mitreiform; spinneret often absent
 Ceph. set. 0; ext. amph. wide, curved; phar. tubular, halfway to nerve ring; freshwater *Colpurella* Cobb 1920
 Ceph. set. present; amphid small, or long and narrow; marine
 Amphid narrow, many head-widths long; neck $\pm 40\%$; cephalic setae 4-8; tail setaceous *Halalaimoides* Cobb 1933
 Amphid small, transverse-oval; neck 25% or less
 Neck 25%; onchium none; cephalic setae 6, small, inconspicuous; spinneret present. *Vasculonema* Kreis 1928
 (cf. *Pelagonema* Cobb 1894)
 Neck 12%; onchium dorsal; cephalic setae 10, conspicuous; labial setae 6. *Trileptium* Cobb 1933
 (= *Trilepta* Cobb 1920)
- Female known
- Ovary 1, anterior
- Gonad ♀ outstretched; pharynx closed; cephalic setae papilloid; spinneret doubtful; marine. *Litotes* Cobb 1920
 Gonad ♀ reflexed; spinneret present; external amphid a transverse slit, not at lips; not marine
 Onchia 0; amphid far behind the \pm quadrate pharynx; in fresh water. *Prismatolaimus* de Man 1880
 Onchium minute, deepset; pharynx opposite pharynx; in soil about roots
 Pharynx conoid when open; amphid opposite base of pharynx. (*Trischistoma* Cobb 1913)
 Pharynx a long tube, amphid opposite its anterior end, with pocket. *Cryptonchus* Cobb 1913
 (= *Dilleversenia* Micol. 1925)
- Ovaries 2; marine exc. Monhystera
- Gonads ♀ outstretched
- Spinneret absent
 Habitat marine; amphid long-oval, on large triangular plate. *Didelta* Cobb 1920
 Habitat not marine; amphid a transverse slit; in soil about roots. *Monhystera* Bast. 1865
 (= *M. dintheriana* de Man 1885)
- Spinneret present
 Mouth opening \pm ventrad; wing duplex; amph. big, bent, dorsal arm the longer (*Campylaimus* Cobb 1920)
 Mouth opening axial, terminal; wings none exc. sometimes *Pseudolella*
 Annules coarse; amphid saccate; nema wide; cephalic setae 4
 Number of annules ± 17 , secreted annules prominent, separated; head small. *Desmoscolex* Clap. 1863
 Number of annules 32-91, secreted annules 0; head concave-quadrate pyramidal. *Tricoma* Cobb 1894
 (= *Quadrucoma* Fil. 1922)
- Annules none, striae fine; amphid not saccate; nema slender
 Amphid small, opening transverse with internal pocket, far back. (*Anoplostoma* Bütsch. 1874)
 Amphid shepherd's crook, dorsal arm the longer
 Length of amphid extreme, sometimes exceeding neck; eyes 0. (*Pseudolella* Cobb 1920)
 Length of amphid not over 1 head-width, loop-shaped; eyes 2, sometimes none
 Oesophagus & lumen widened between eyes and nerve ring; pharynx obvious *Coinonema* Cobb 1920
 Oesophagus & lumen not widened; pharynx like oesophagus. *Araeolaimoides* (de Man) Fil. 1918
- Gonads ♀ reflexed
- Spinneret absent; amphid with internal pocket
 Cephalic setae 0; pharynx 2-3%, tubular; tail rounded, under 2%; in soil. *Isolaimium* Cobb 1920
 Cephalic setae 10, subcephalic many; marine. *Platycomopsis* Ditlev. 1926
 (= *Dactylonema* Fil. 1927)
- Spinneret present
 Habitat freshwater; amphid usually small or inconspicuous
 Pharynx wide, domed by thin lips; onch. 0; amph. transverse slit, far back. *Prismatolaimus* de Man 1880
 Pharynx closed, lips thick, onch. minute, deepset; external amphid round or oval. (*Triplya* Bast. 1865)
 (= *Promononchus* Micol. 1923)
- Habitat marine
 Cephalic setae none or papilloid; external amphid narrow, transverse (for alt. see next page)
 Amphid $\frac{1}{2}$ head-width; cuticle $\frac{1}{2}$ radius, annules coarse, markings rod-like. (*Actinonema* Cobb 1920)
 Amphid $\frac{1}{2}$ head-width or less, with internal pocket; cuticle less thick; phar. with lumen; Enoplidae
 Eyes 2; protrusile pharyngeal ribs none. *Illium* Cobb 1920

- Eyes 0; protrusile pharyngeal ribs 3, straight..... *Fimbrilla* Cobb 1905
 (= *Fimbria* Cobb 1894)
- Cephalic setae present, not papilloid (for alternative see preceding page)
- Amphid shepherd's crook
 Annules not of tile-like elements..... *Parachromagaster* Allgen 1929
 Annules coarse, each of 8 tile-like elements..... *Ceramonema* Cobb 1920
 (cf. *Pselionema* Cobb 1933)
- Amphid not shepherd's crook
 Nema wide; annules ± 17 , secreted annules obvious; amph. saccate; ceph. set. 4. *Desmoscolex* Clap. 1863
 Nema slender, under 5%; amphid not saccate; annules 0, striae fine or 0; Enopliidae
- Onchium dorsal; labial setae 0; cephalic setae 10, near lips
 Tooth vestigial, on wall; external amphid a small slit; eyes 0..... (*Anoncholaimus* Cobb 1920)
 Tooth a bending inward of dorsal pharyng. wall; amphid larger, faint. (*Asymmetrella* Cobb 1920)
 Onchia none; labial setae 6, obvious; cephalic setae 10 or more, length 1-5 times head width
 Pharynx narrow, closed; odontia (?) minute; ceph. set. 3 times head width (*Tubolaimella* Cobb 1933)
 Pharynx wide, open; odontia none
 Lips 3, deep-cleft, with framework; ceph. set. 2-5 times head width..... *Chaetonema* Fil. 1927
 Lips without framework; ceph. set. 1 head-width; excret. pore raised. (*Paranticoma* Micol. 1930)
- External amphid spiral, circular or elliptical (for alternative see preceding page)
- Female not known; spinneret present; marine
 Nema wide, 14-32%, widest at lip region, tapering to terminus; wings numerous. *Pteronium* Cobb 1933
 Nema slender, not over 5% wide, widest near middle
 Amphid multispire; wings 0; cephalic & labial setae 18; labial framework present.... *Dispira* Cobb 1933
- Amphid circular, elliptical or monospiral
 Odontia 2, large, lateral or subventral..... (*Diodontolaimus* Southern 1914)
 Odontia absent (in *Stephanolaimus* 6 minute?)
 Onchium ventral; cephalic setae 10; amph. $\frac{1}{2}$ head-width, with internal pocket. (*Isonemella* Cobb 1920)
 Onchium absent (*Nudolaimus*?)
 Calvarium reduced to submed. plates; phar. large, wall strong, "onchia" basal. *Nudolaimus* Allgen 1929
 Calvarium absent; cephalic setae present
 Wings present; cephalic setae 6 or 10
 Ceph. set. 6; wings 8; annules coarse, elements tile-like; oesoph. compound. *Dasynemella* Cobb 1933
 (= *Dasyne* Cobb 1920)
 Ceph. set. 10; wings 2; oesophagus simple; pharynx \pm quadrate..... *Pulchranemella* Cobb 1933
- Wings absent; cephalic setae 4-8
 Lip region discoid, set off by constriction; odontia (?) 6..... *Stephanolaimus* Ditlev. 1918
 Lip region not discoid; cephalic setae 4 or 6
 Pharyng. vestibule with rod-like supports; head \pm cylindrical; striae 0; ceph. set. 4
 Rods 6, straight, parallel to body axis; cephalic setae 4, jointed..... *Rhadinema* Cobb 1920
 Rods more than 6, curved, meeting axially..... *Eleutherolaimus* Fil. 1922
 Pharyng. vestibule without supports; striae pres. exc. *Vasculonema* & *Bathylaimella*
- Amphid transverse-oval, near lips; head slightly conoid; setae small
 Setae labial, cephalic and subcephalic, minute..... *Zygonemella* Cobb 1920
 Setae cephalic only, 6, small, inconspicuous..... *Vasculonema* Kreis 1928
 (cf. *Pelagonema* Cobb 1894)
- Amphid circular, near base of wide-tubular pharynx
 Head cylindrical; setae on head 16, longest 4 times head width..... *Cenolaimus* Cobb 1933
 Head conoid, narrowing 50% from base; setae on head 4, short
 Cuticle striated; width (compressed?) 6%..... *Tubolaimus* Allgen 1929
 Cuticle not striated; width 3%..... *Bathylaimella* Allgen 1930
- Female known
- Ovary 1 (for alternative see next page)
- Gonad \varnothing reflexed
 Amphid multispire; ovary posterior; pharynx linear; spinneret doubtful; marine *Alaimella* Cobb 1920
 Amphid not multispire; ovary anterior
 Spinneret none; amph. round, with central mound; caudal setae thorn-like, 2 rows (*Zanema* Cobb 1920)
 Spinneret present; tail without thorn-like setae
 Habitat marine; pharynx tubular; cephalic setae 4; nema $\pm 2\%$ wide..... (*Conolaimella* Allgen 1930)
 Habitat not marine; pharynx open, not small nor narrow; nema 2-5% wide
 Cephalic setae 4; pharynx with supporting pieces; amph. monospire; fresh-water *Anonchus* Cobb 1913
 Cephalic setae 6 or 10, papilloid in *Tripylum*
 Amphid monospire; in gills of land crabs..... (*Tripylum* Cobb 1920)
 Amphid small, circular, transverse-oval or slit-like
 Onchium 0; pharynx \pm quadrate; amphid far back; in fresh water.... *Prismatolaimus* de Man 1880
 Onchium minute, deepset; pharynx conoid; amphid opp. pharynx; in soil (*Trischistoma* Cobb 1920)
- Gonad \varnothing outstretched
 Habitat not marine
 Spear minute; spinneret none; bursa present; parasitic in beetle..... (*Aphelenchulus* Cobb 1920)
 Spear absent; bursa absent; in fresh water..... *Cylindrolaimus* de Man 1886
- Habitat marine
- The ovary posterior
 External amphid multispire; pharynx linear; tail $\pm 10\%$; spinneret doubtful (*Alaimella* Cobb 1920)
 External amphid circular, pocket elongate; tail 34%; spinneret none..... *Halanonchus* Cobb 1920
- The ovary anterior
 Cephalic setae 4, papilloid; pharynx obscure, linear; amphid faint, open caudad.... *Litotes* Cobb 1920
 Cephalic setae not papilloid
 Spinneret absent; labial appendages ("palps") present; amphid circular
 Tail with 2 rows of thorn-like setae; amphid with distinct central mound..... *Zanema* Cobb 1920
 Tail without thorn-like setae; amphid without mound; cardiac bulb faint *Crystallonema* Cobb 1920
 Spinneret present; amphid circular or monospire
 Pharynx 7%, tubular, \pm halfway to nerve ring; head narrow; amphid far back *Rhynconema* Cobb 1920
 Pharynx less than 2%, not a long tube
 Depth of pharynx much greater than width, pharynx small
 Oesophagus almost surrounding conoid pharynx..... *Conolaimella* Allgen 1930
 Oesophagus meeting base of obconoid pharynx..... *Sphaerocephalum* Fil. 1918
 Depth of pharynx not much greater than width, pharynx large, open

- Amphid ± 1 head-width; vulva 50%; caudal set. stiff, 8 prs.; post. ovary small *Halinema* Cobb 1920
 Amphid $\pm \frac{1}{2}$ head-width; vulva 80-85%; caudal set. 0; lips 6, conoid, separate
 Labial setae 6, hook-shaped, on tips of lips; amphid spiral; in mud..... (*Dactylaimus* Cobb 1920)
 Labial setae 6, not hook-shaped nor at tips; amphid round; ceph. set. 18. *Omicronema* Cobb 1920
- Ovaries 2 (for alternative see preceding page)
 Gonads ♀ outstretched; usually marine
 Spinneret absent; marine exc. *Pseudoaulolaimus*
 Habitat mud, rice fields; wings 16-32, fusing to 4 on 4-pronged tail.... *Pseudoaulolaimus* Imamura 1931
 Habitat marine
 Amphid long oval, on wide triangular plate, almost as long as head is wide..... *Didella* Cobb 1920
 Amphid smaller, not on triangular plate
 Cervical setae bunched; 2 flat setae in front of transverse-oval amphid.... (*Platycoma* Cobb 1894)
 Cervical setae if present not bunched; amphid \pm circular
 Pharynx long, simple, tubular; cephalic setae 6..... *Southerniella* Allgen 1932
 Pharynx minute; ceph. set. 18, minute, in 2 circlets; amphid faint multispire *Anticyathus* Cobb 1920
- Spinneret present; marine exc. *Cylindrolaimus*
 Habitat freshwater; ceph. set. 4; ovaries usually unequal; pharynx tubular *Cylindrolaimus* de Man 1880
 Habitat marine
 Annules few, coarse; amphid saccate; cephalic setae 4; nema wide
 Number annules ± 17 , secreted annules prominent, separated; head small... *Desmoscolex* Clap. 1863
 Number annules 32-91, secreted annules none; head concave-quadrate pyramidal *Tricoma* Cobb 1894
 (= *Quadriricoma* Fil. 1922)
- Annules if present not few nor coarse; amphid not saccate
 Gland ducts cuticularized, pores cervical & pre-anal; striae coarse.... *Halaphanolaimus* Southern 1914
 Gland ducts and pores absent
 Cephalic setae 10 or more
 Pharynx with transverse cuticularized ring
 Cavity of pharynx simple, capacious, \pm cylindrical, $\frac{1}{2}$ head-width.... *Eumorpholaimus* Schulz 1932
 Cavity of pharynx small, cup-shaped, rings sometimes 2 or more..... (*Linhomoeus* Bast. 1865)
 (i.e. subg. *Paralinhomoeus* de Man 1907
 = *Paradesmolaimus* Schulz 1932)
- Pharynx without transverse cuticularized rings
 Longest ceph. set. 4, total 16, tips simple; phar. quadrate; amph. \pm neck-width *Halinema* Cobb 1920
 Longest ceph. set. 6, 4 short specialized at tip; phar. 0(?); multispire faint *Linhomoeus* Cobb 1920
- Cephalic setae 4
 Nema wide, 10-16%; wings numerous; neck region "collared"; amphid spiral; Richtersiinae
 Wings ± 50 ; pharynx narrow; spicula 2, equal..... *Richtersia* Steiner 1916
 Wings 16-32; pharynx capacious, open; spicula unequal, or 1 only..... *Richtersiella* Kreis 1929
 Nema slender, under 5%; wings if present not numerous
 Eyes absent; amphid circular; pharynx truncate at base..... *Bathylaimella* Allgen 1930
 Eyes often present; setae behind head usually long, numerous, conspicuous
 Oesophagus lumen not enlarged; amphid loop-shaped..... (*Araolaimoides* (de Man) Fil. 1918)
 Oesophagus and lumen widened between eyes and nerve ring; pharynx obvious
 Shape of amphid loop-like ("folded")..... (*Covonema* Cobb 1920)
 Shape of amphid circular or monospiral; eyes sometimes absent(?)... *Araolaimus* de Man 1888
- Gonads ♀ reflexed
 Spinneret absent; amphid circular, elliptical or monospiral
 Habitat not marine; pharynx long, tubular, \pm half as long as oesophagus
 Wings 16-32, conspic., fusing to 4 on 4-pronged tail; in rice fields.... *Pseudoaulolaimus* Imamura 1931
 Wings 0; end of tail not 4-parted; in soil..... *Isolaimium* Cobb 1920
 Habitat marine; pharynx short
 Pharynx narrow; amphid \pm flattened, preceded by 2 flat setae..... *Platycoma* Cobb 1894
 Pharynx 0 or linear; amphid transverse-oval, with pocket, 4 set. posterior... *Platycomopsis* Ditlev. 1926
 (= *Dactylonema* Fil. 1927)
- Spinneret present
 Habitat freshwater
 Mouth cavity absent or obscure, closed
 Amphid monospire; gland cells lateral; ♂ supplements protrusile tubes... *Aphanolaimus* de Man 1880
 Amphid circular or elliptical; ♂ supplements not protrusile..... *Tripyla* Bast. 1865
 (= *Promononchus* Micol. 1923)
- Mouth cavity definite, open, obvious
 Pharynx narrow, compound; amphid monospire, $\frac{1}{2}$ head-width; cephalic setae 4 *Paraplectus* Fil. 1930
 Pharynx simple, i.e. of 1 chamber
 The amphid small, transverse-oval or a slit; cephalic setae 6 or 10... *Prismatolaimus* de Man 1880
 The amphid monospire, opp. pharynx; ceph. setae 4; gland cells lateral *Paraphanolaimus* Micol. 1923
- Habitat marine
 Nema wide, over 10%
 Annules few, coarse; amphid saccate
 Body setae not dense; secreted annules prominent, separate; head small... *Desmoscolex* Clap. 1863
 Body setae dense, set along annules; junct. oesoph. & intestine indefinite... *Greeffiella* Cobb 1922
 (= *Trichoderma* Greeff 1869)
- Annules if present many; longitudinal ridges or wings 16-50; neck "collared"; Richtersiinae
 Wings ± 50 ; pharynx narrow; spicula 2, equal..... *Richtersia* Steiner 1916
 Wings 16-32; pharynx wide, capacious; spicula unequal, or 1 only..... *Richtersiella* Kreis 1929
 Nema slender, not over 5%
 Calvarium present as a band or plates
 Oesophagus not crenate; calvarium of plates; gland cells lateral..... *Cylicolaimus* de Man 1889
 Oesophagus crenate; calvarium a band..... (*Phanodermopsis* Ditlev. 1926)
 (= *Galeonema* Fil. 1927)
- Calvarium absent
 Gland ducts cuticularized, pores cervical & pre-anal; striae coarse... (*Halaphanolaimus* Southern 1914)
 Gland ducts and pores none, or not cuticularized
 Denticles mid-phar., apophyses forward & back; amph. multispire (*Halichoanolaimus* de Man 1888)
 (= *Smalsundia* Allgen 1929)
- Denticles none: amphid not multispire

- Pharynx open, wide, \pm capacious
 Amphid round; anterior pharyng. chamber large, posterior small; setae 10 *Bathylaimus* Cobb 1894
 (= *Macrolaimus* Ditlev. 1918
 = *Colthonolaimus* Ditlev. 1919)
- Amphid small, with internal pocket (*Pandolaimus* ?); pharynx of 1 chamber
 Depth pharynx \pm equalling width; dorsal onch. small, ant., appressed *Anoncholaimus* Cobb 1920
 Depth pharynx nearly twice width
 Wall of pharynx irregularly thick; amphid apparently circular..... *Pandolaimus* Allgen 1929
 Wall of pharynx thin; amphid small, transverse-oval, with pocket.... *Pelagonema* Cobb 1894
 (cf. *Vasculonema* Kreis 1928)
- Pharynx narrow, often tubular or with closed lumen
 The amphid behind level of cephalic setae
 Cephalic setae 0; pharynx $\frac{1}{2}$ head-width, with 3 refractive elements... *Angustinema* Cobb 1933
 Cephalic setae present
 Nema over 3 mm. long..... *Cynura* Cobb 1920
 Nema 0.5 mm. long..... *Aplectus* Cobb 1914
- The amphid in front of level of cephalic setae; cephalic setae 4
 Dorsal pharyng. wall not thicker; amphidial glands large, conspic.; eyes 2... *Ionema* Cobb 1920
 Dorsal pharyng. wall thick, tooth-like; amph. monospire at lips; renette far back
 Oesoph. glands obscuring junct. of intest.; dorsal phar. element uniform... *Onchium* Cobb 1920
 Oesoph. glands not prominent; dorsal pharyng. element not uniformly thick, but heavy
 Tooth-like thickening parallel to axis..... *Camacolaimus* de Man 1889
 (= *Digitonchus* Cobb 1920
 = *Acontiolaimus* Fil. 1918)
- Tooth-like thickening bent outward from beginning of vestibule
 Point of "onchium" single..... *Acmaeolaimus* Fil. 1918
 Point of Y-shaped "onchium" duplex..... *Ypsilon* Cobb 1920

LITERATURE CITED

- ALLGÉN, C. *Freilebende marine Nematoden von den Campbell-und Staten-Inseln.* Nyt. Mag. Naturvidenskaberne. 66: 305-306. 1927.
- *Südschwedische marine Nematoden.* Göteborgs K. Vetensk. O. Vetterhets = Samhälles. Handl. Ser. B 1: 29. 1929.
- *Neue freilebende marine Nematoden von der Westküste Schwedens.* Zool. Jahrb. Abt. System. Ökol. u. Geogr. Tiere. 57: 433-434, 436-437, 445, 454, 461, 466, 467, 481, 491, 492, 493. 1929.
- *Conolaimella, ein neues Genus mariner Nematoden von der Westküste Schwedens.* Zool. Anz. 88: 138. 1930.
- *Freilebende marine Nematoden von der Stateninsel (Feuerland-Archipel.) I.* Zool. Anz. 89: 257. 1930.
- *Freilebende marine Nematoden von der Stateninsel (Feuerland-Archipel.) II.* Zool. Anz. 90: 28. 1930.
- *Weitere Beiträge zur Kenntnis der marinen Nematodenfauna der Campbellinsel.* Nyt. Mag. Naturv. 70: 186. 1932.
- *Ueber einige freilebende marine Nematoden aus der Umgebung der Biologischen Station auf der Insel Herdla (Norwegen) mit Anhang; zur Richtigestellung älterer und neuerer mariner Nematodengenera I.* Arch. Naturgesch. 1: 424, 432. 1932.
- BASTIAN, H. C. *Monograph on the Anguillulidae, or free nematodes, marine, land, and fresh water; with descriptions of 100 new species.* Trans. Linn. Soc. London 25: 97, 100-101, 103-104, 115, 118, 121-122, 124, 125-126, 132, 141, 142-143, 144, 145, 154-155, 156, 157, 158, 159, 162-163, 165, 167, 178. 1865.
- BAYLIS, H. A. AND DAUBNEY, R. *A synopsis of the families and genera of Nematoda.* London. Pp. 107, 112. 1926.
- BOVIEN, P. *On a new nematode, Scatonema wülkeri gen. et sp. n. parasitic in the body-cavity of Scatopse fuscipes Meig. (Diptera nematocera).* Vidensk. Meddel. Dansk Naturhist. For. Kjøbenhavn. 94: 15-31. 1932.
- BÜTSCHLI, O. *Zur Kenntniss der freilebenden Nematoden.* Abhandl. Senckenb. Naturf. Gesell. Frankfurt a. M. 9: 270-273, 285. 1874.
- CARUS, J. V. *Icones zootomicae. Erste Hälfte: Die wirbellosen Thiere.* Leipzig. Pl. 8, Fig. 1. 1857.
- CHITWOOD, B. G. *Helminthological Society Proceedings.* The Journal of Parasitology. 19: 243. 1933.
- CHRISTIE, J. R. AND CHITWOOD, B. G. *Chondronema passali (Leidy, 1852) n.g. (Nematoda), with notes on its life history.* Jour. Wash. Acad. Sci. 21: 357. 1931.
- CLAPARÈDE, E. *Beobachtungen über Anatomie und Entwicklungsgeschichte wirbelloser Thiere an der Küste von Normandie angestellt.* Pp. 88-90. 1863.
- COBB, M. V. *Some fresh-water nematodes of the Douglas Lake Region of Michigan, U. S. A.* Amer. Micros. Soc. Trans. 34: 28-29. 1915.
- (edited by) N. A. Cobb: *New nemtic genera and species, with taxonomic notes.* Jour. Parasitol. 20: 81-94. 1933.
- COBB, N. A. *Onyx and Dipeltis: New nematode genera, with a note on Dorylaimus.* Linn. Soc. N. S. Wales, Proc. 6: 146-153, 155. 1891.
- *Plant diseases and their remedies. (III. Nematode worms found attacking sugarcane.)* Agric. Gaz. N. S. Wales. Pp. 811, 819-820, 821, 825. 1893.
- *Tricoma and other new nematode genera.* Linn. Soc. N. S. Wales, Proc. (1893) (s. 2). 8: 389-390, 391, 392-393, 399-401, 409-410, 411-413, 413-414, 416, 419, 420-421. 1894.
- *Australian free-living marine nematodes.* Linn. Soc. N. S. Wales, Proc. (s. 1). 23: 406-407. 1898.

- COBB, N. A. 1905. See Stiles and Hassall, 1905.
- *Fungus maladies of the sugar cane, with notes on associated insects and nematodes.* Bull. (5), Hawaii Sugar Planters' Assoc. Expt. Sta., Div. Path. & Physiol., 2nd ed., pp. 187-189. 1906.
- *Draconema: A remarkable genus of marine free-living nematodes.* Jour. Wash. Acad. Sci. 3: 146-149. 1913.
- *Notes on Mononchus and Tylenchulus.* Jour. Wash. Acad. Sci. 3: 287. 1913.
- *New nematode genera found inhabiting fresh water and non-brackish soils.* Jour. Wash. Acad. Sci. 3: 436-444. 1913.
- *Antarctic marine free-living nematodes of the Shackleton Expedition.* Contrib. Sci. Nematol. I, pp. 12, 16. 1914.
- *North American free-living fresh-water nematodes.* Amer. Micros. Soc. Trans. 33: 92-94. 1914.
- *Selachinema a new nematode genus with remarkable mandibles.* Contrib. Sci. Nematol. IV, p. 113. 1915.
- *Notes on nemas.* Contrib. Sci. Nematol. V, pp. 118, 119. 1917.
- *Filter-bed nemas: Nematodes of the slow sand filter beds of American cities (including new genera and species) with notes on hermaphroditism and parthenogenesis.* Contrib. Sci. Nematol. VII, pp. 195-196, 203. 1918.
- *One hundred new nemas (type species of 100 new genera).* Contrib. Sci. Nematol. IX, pp. 230-340. 1920.
- *Howardula benigna, a nema parasite of the cucumber-beetle (Diabrotica).* Contrib. Sci. Nematol. X, pp. 1-2. 1921.
- *Marionella (Eurystoma Marion, 1870); an emendation, with notes on a new birefringent substance, marionellin, found in the intestinal cells.* Jour. Wash. Acad. Sci. 11: 504-509. 1922.
- *Greeffiella.* Jour. Wash. Acad. Sci. 12: 299-303. 1922.
- *The Helminthological Society of Washington.* Jour. Parasitol. 11: 105, 116, 120. 1924.
- *Marine free-living nemas.* Australas. Antarc. Exped. 1911-14. Scient. Rep., Sydney, Ser. C. Zool. & Bot. 6(7): 22-23. 1930.
- *The demanian vessels in nemas of the genus Oncholaimus; with notes on four new oncholaims.* Jour. Wash. Acad. Sci. 20: 227. 1930.
- DADAY, E. VON. *Mikroskopische Süßwasserthiere aus Deutsch-Neu-Guinea.* Termész. Füzetek. 24: 7. 1901.
- *Untersuchungen über die Süßwasser-Mikrofauna Paraguays. Mit einem Anhang von W. Michaelsen.* Zool. 18(44): p. 62. 1905.
- DE CONINCK, L. *Sur trois espèces nouvelles de nématodes libres trouvés en Belgique.* Bull. Mus. Roy. Hist. Nat. Belg. 7(11): 2-5. 1931.
- DIESING, K. M. *Revision der Nematoden.* Sitzber. Akad. Wiss. Wien, Math. Naturw. Kl. (1860) 42: 728. 1861.
- DITLEVSEN, H. *Marine free-living nematodes from Danish waters.* Vidensk. Meddel. Dansk Naturhist. For. Kjøbenhavn. 70: 162-163, 168, 174-177, 177-178, 181-182, 183-184, 185, 188-189, 197, 299. Plate 7, figs. 3, 4 & 6. Plate 8, figs. 2, 3 & 8. 1918.
- *Marine free-living nematodes from the Auckland and Campbell Islands.* Vidensk. Meddel. Dansk Naturhist. For. Kjøbenhavn. 73: 2, 4. 1922. Advance separate apparently issued in 1921.
- *Nematological notes.* Vidensk. Meddel. Dansk Naturhist. For. Kjøbenhavn. 74: 57-58, 60. 1922.
- *Free-living nematodes.* The Danish Ingolf-Expedition (1895-1896) Copenhagen. 4(6): 8, 12, 30-31. 1926.

- DITLEVSEN, H. *Free-living marine nematodes from Greenland waters*. Meddel. Grønland 23 (supplement): 227, 230. 1928.
- DUFOUR, L. *Recherches sur quelques entozoaires et larves parasites des insectes Orthoptères et Hyménoptères*. Ann. Sci. Nat., Zool. 7(2): 9. 1837.
- DUJARDIN, F. *Histoire naturelle des helminthes ou vers intestinaux*. Paris. Pp. 230, 233, 235-236, 239-240. 1845.
- EHRENBERG, HRN. *Über die Akalephen des Rothen Meeres und den Organismus der Medusen der Ostsee*. Abhandl. Akad. Wiss. Berlin (1835). P. 219. 1837.
- FILIPJEV, I. N. *Les nématodes libres contenus dans les collections du Musée Zoologique de l'Académie Impériale des Sciences de Petrograd*. Ann. Mus. Zool. Acad. Imp. Sci. 21: 65, 73-75, 83, 105-106. 1916.
- *Novaia svobodnaia nematoda iz Kaspuskago Moria Chromadorissa gen. nov.* (Chromadoridae, Chromadorina) (also French résumé). Russ. K. Zool. Zhurn. Moskoa. 2: 24-30. 1917.
- *Free-living marine nematodes of the vicinity of Sebastopol. Part I*. Trudy Osob. Zool. Lab. Sebastopol. Biol. Stantsii Russ. Akad. Nauk Petrograd. s. 2(4), pp. 50-51, 110, 122-123, 139-141, 186, 205, 206, 211, 218-219, 226-227, 229-230, 232-233, 256-257, 259, 284, 303, 305, 323-324, 326, 328, 345. 1918.
- *Free-living marine nematodes of the vicinity of Sebastopol. Part II*. Spec. Zool. Lab. and Sebastopol Biol. Sta. Russ. Acad. Sci. s. 2(4). pp. 565, 568. 1921.
- *Encore sur les nématodes libres de la mer noire*. Acta. Inst. Agron. Staupropolitani. (Zool.) 1: 114-115, 121-122, 133-134, 137, 138-139, 150, 173-174. 1922.
- *Les nématodes libres des mers septentrionales appartenant à la famille des Eno-plidae*. Arch. Naturgesch. (1925) Abt. A (6). 91: 65, 66, 79, 81-82, 105-106, 123, 130, 155-156, 160, 176-177, 180-182, 183, 188, 199, 203. 1927.
- *Les nématodes libres de la baie de la Neva et de l'extrémité orientale du Golfe de Finlande. Deuxième partie*. Arch. Hydrobiol. 21: 12-13, 29, 35-36. 1930.
- FISCHER, M. *Ueber eine Clematis-krankheit*. Ber. Physiol. Lab. u. Versuchsanst. Landw. Inst. Halle. 3(11): 1-11. 1894.
- FUCHS, G. *Über Parasiten und andere biologisch an die Borkenkäfer gebundene Nematoden*. Verhandl. Gesell. Deut. Naturf. u. Ärzte. 2nd Part, 1 Hälfte. P. 690. 1914.
- *Die Naturgeschichte der Nematoden und einiger anderer Parasiten*. Zool. Jahrb. Abt. System., Geogr. u. Biol. Tiere. 38: 158-160. 1915.
- *Neue an Borken- und Rüsselkäfer gebundene Nematoden, halbparasitische und Wohnungseinmieter*. Zool. Jahrb., Abt. System., Ökol. u. Geogr. Tiere. 59: 540-544, 560-561, 565-566, 574-576, 586-588, 609-612, 616-618. 1930.
- *Seinura gen. nov.* Zool. Anz. 94: 226-227. 1931.
- GÖLDI, E. A. *Relacao dos estudos tendentes a elucidar a molestia do cafeeiro na provincia do Rio—Resultados dos tres primeiros mezes de estudos*. (34 pp.) 1886.
- *Relatorio sobre a molestia do cafeeiro na provincia do Rio de Janeiro (1887)*. Archivos do Museu Nacional Rio de Janeiro. 8: 68. 1892.
- GOODEY, T. *Hexatyclus viviparus gen. et sp. nov., a nematode found in a diseased potato tuber*. Jour. Helminthol. 4: 28, 30. 1926.
- *On some new and little-known free-living nematodes*. Jour. Helminthol. 7: 27-30, 41-45. 1929.
- *Cylindrogaster coprophaga gen. et sp. nov. a nematode found in a culture of faeces from a wild brown rat*. Jour. Helminthol. 5: 26-31. 1927.
- *On a remarkable new nematode, Tylenchinema oscinellae gen. et sp. n., parasitic in the frit-fly, Oscinella frit L., attacking oats*. Roy. Soc. Proc., Ser. B. (London). 218: 338-339. 1930.

- GREEFF, R. *Untersuchungen über einige merkwürdige Formen des Arthropoden- und Wurm-typus.* Arch. Naturgesch. 35. J., 1: 115. 1869.
- HOFMÄNNER, B. AND MENZEL, R. *Neue Arten freilebender Nematoden aus der Schweiz.* Zool. Anz. 44: 88. 1914.
- IMAMURA, S. *Nematodes in the paddy field, with notes on their population before and after irrigation.* Jour. Col. Agr., Imp. Univ., Tokyo. 11: 210-211. 1931.
- IRWIN-SMITH, V. A. *On the Chaetosomatidae, with descriptions of new species, and a new genus from the coast of New South Wales.* Linn. Soc. N. S. Wales, Proc. (1917) (168). 42: 798. 1918.
- JÄGERSKIÖLD, L. A. *Bunonema richtersi n. g. n. sp. Ein eigentümlicher neuer Land-nematode aus dem Schwarzwald, von Kerguelen und Possession-Inseln (Crozet-Inseln).* Zool. Anz. 28: 557-561. 1905.
- KREIS, H. A. *Zur Kenntnis der freilebenden marinen Nematoden.* Schr. Süßwasser u. Meereskunde. 2(6): 167-168. 1924.
- *Weiterer Beitrag zur Kenntnis der freilebenden marinen Nematoden.* Arch. Naturgesch. (1926). Abt. A(8). 92: 2-4, 10. 1928.
- *Die freilebenden marinen Nematoden der Spitzbergen-Expedition von F. Roemer und F. Schaudinn im Jahr 1898.* Mitt. Zool. Mus. Berlin. 14: 150-151, 158-159, 163-164, 171-172, 172-173, 177, 181-182. 1928.
- *Freilebende terrestrische Nematoden aus der Umgebung von Peking (China).* Zool. Anz. 84: 288. 1929.
- *Freilebende marine Nematoden von der Nordwest-Küste Frankreichs (Trèbeurden: Côtes du Nord).* Capita Zool. 2(7): 26, 44, 51, 54, 62, 67, 69. 1929.
- *Fresh-water nematoda from the Paraguayan Chaco.* Jour. Linn. Soc., Zool. 38(no. 257): 72. 1932.
- *Beiträge zur Kenntnis pflanzenparasitischer Nematoden.* Ztschr. Parasitenk. 5: 185-189. 1932.
- *Freilebende marine Nematoden von den Sunda-Inseln. II. Oncholaiminae.* Vidensk. Meddel. Dansk Naturhist. For. Kjøbenhavn. 93: 28-29, 30, 31, 38-39, 42, 52, 57, 67. 1932-1933.
- LEIDY, J. *Contributions towards a knowledge of the marine invertebrate fauna of the coasts of Rhode Island and New Jersey.* Jour. Acad. Nat. Sci. Phila. 3(s.2): 144. 1855.
- LEUCKART, R. *Ueber einen neuen heterogenen Nematoden.* Tagebl. d. 57. Versamml. Deutsch. Naturf. u. Aertze, Magdeb. (5). P. 320. 1884.
- *Ein Sphaerularia-artiger neuer Nematode.* Zool. Anz. 9: 743-746. 1886.
- *Neue Beiträge zur Kenntniss des Baues und der Lebensgeschichte der Nematoden.* Abhandl. Math.-Phys. Cl. k. Sächs. Gesell. Wiss. 13: 678-703. 1887.
- LINSTOW, O. VON. *Helminthologische beobachtungen.* Arch. Naturgesch. 42.J., 1: 16-17. 1876.
- *Helminthologica.* Arch. Naturgesch. 43.J., 1: 2-3. 1877.
- MAN, J. G., DE. *Onderzoekingen over vrij in de aarde levende nematoden.* Tijdschr. Nederl. Dierk. Vereen. 2: 116-117, 119, 120, 137-138, 168-169, 172. 1876.
- *Die einheimischen, frei in der reinen Erde und im süßsen Wasser lebenden Nematoden.* Tijdschr. Nederl. Dierk. Vereen. 5: 2, 3-4, 5, 14, 15, 21, 28, 30-31, 34, 58-64. 1880.
- *Helminthologische beitraege.* Tijdschr. Nederl. Dierk. Vereen. s.2, 1: 2-3. 1885.
- *Anatomische Untersuchungen über freilebende Nordsee-nematoden.* Leipzig. Pp. 60-76. 1886.
- *Sur quelques nématodes libres de la mer du Nord nouveaux ou peu connus.* Mém. Soc. Zool. France. 1: 2-3, 11, 14-16, 31-32, 34-35. 1888.

- MAN, J. G., DE. *Espèces et genres nouveaux de nématodes libres de la mer du Nord et de la manche.* Mém. Soc. Zool. France. 2: 1, 3, 4, 8, 9-10. 1889.
- *Quatrième note sur les nématodes libres de la mer du Nord et de la manche.* Mém. Soc. Zool. France. 3: 189-190. 1890.
- *Cinquième note sur les nématodes libres de la mer du Nord et de la manche.* Mém. Soc. Zool. France. 6: 81, 84-85, 99-100, 114-115, 118-119. 1893.
- *Sur quelques espèces nouvelles ou peu connues de nématodes libres vivant sur les côtes de la Zélande.* Tijdschr. Nederl. Dierk. Vereen. 10(2): 228, 232-233, 237, 239-240, 241-242. 1907.
- *Helminthologische Beiträge.* Zool. Jahrb. (Supplement 15). 1: 439-440. 1912.
- *Odontopharynx longicaudata n. g., n. sp. Eine neue Form von Anguilluliden.* Zool. Jahrb. Abt. System., Geogr. u. Biol. Tiere. 33: 637-638. 1912.
- *Nouvelles recherches sur les nématodes libres terricoles.* Capita Zool. 1: 9, 14, 33, 35, 36, 42-43. 1921.
- *Neue freilebende Nematoden aus der Zuidersee.* Tijdschr. Nederl. Dierk. Vereen. 18: 126, 131. 1922.
- MARION, A. F. *Recherches zoologiques et anatomiques sur des nématodes non parasites, marins.* Ann. Sci. Nat. Zool. (5) Art. 14. 13: 11-12, 14, 16, 18, 19, 25, 31, 32, 34. 1870.
- MAUPAS, E. *Modes et formes de reproduction des nématodes.* Arch. Zool. Expt. e. Gen. 8 (3): 578-582. 1900.
- METSCHNIKOFF, E. *Beiträge zur Naturgeschichte der Würmer.* Ztschr. Wiss. Zool. 17: 542. 1867.
- MICOLETZKY, H. *Neue Süßwasser-Nematoden aus der Bukowina.* Mitt. Naturw. Ver. Steiermark. (1914) 51: 447. 1915.
- *Die freilebenden Erd-Nematoden.* Arch. Naturgesch. (1921) 87: 119, 374-377, 606-607. 1922.
- *Freilebende Nematoden der Wolga.* Arb. Biol. Wolga-Sta. 7: 13, 14, 17-18, 21-24. 1923.
- *Weitere Beiträge zur Kenntnis freilebender Nematoden aus Suez.* Sitzber. Akad. Wiss. Wien, Math. Naturw. Kl. Abt. 1, 132: 228, 235, 242-244. 1924.
- *Letzter Bericht über freilebende Nematoden aus Suez.* Sitzber. Akad. Wiss. Wien, Math. Naturw. Kl. Abt. 1, 133: 138-139, 140, 157-158, 164-165. 1924.
- *Die freilebenden Süßwasser- und Moornematoden Dänemarks. Nebst. Anhang: Über Amöbosporeidien und andere Parasiten bei freilebenden Nematoden.* K. Dansk. Vidensk. Selsk. Skr. Naturv. Math. Afd. 10: 153, 248-249, 257-258. 1925.
- *Freilebende marine Nematoden von den Sunda-Inseln. 1. Enoplidae.* Vidensk. Meddel. Dansk. Naturh. For. Kjøbenhavn. 87: 261, 266-267, 293, 312, 330-331. 1930.
- ONORATO DE CILLIS, M. I. *Nuovi generi e nuove specie di nematodi liberi d'acqua dolce.* Monit. Zool. Ital. 28: 58-59. 1917.
- PANCERI, P. *Osservazioni intorno a nuove forme di vermi nematodi marini.* Atti. r. Accad. Sci. Fis. e Mat., Napoli. 7(10): 7-9. (1876) 1878.
- PETERS, B. G. *On the nomenclature of the vinegar eelworm.* Jour. Helminth. 5: 141-142. 1927.
- RAHM, G. *Beitrag zur Kenntnis der Moostierwelt der preussischen Rheinlande. I. Systematisch beschreibender Teil.* Arch. Naturgesch. (1924) Abt. A, 90: 176. 1925.
- *Alguns nematodes parasitas e semi-parasitas das plantas culturaes do Brasil.* Arch. Inst. Biol. I: 240, 241, 243-244, 246, 248, 250. 1928.
- *Freilebende Nematoden Rotatorien und Tardigraden aus Südamerika (besonders aus Chile).* Zool. Anz. 98: 96, 99, 106, 107. 1932.

- RICHTERS, F. *Moosfauna-Studien*. Ber. Senckenb. Naturf. Gesell. Frankfurt a. M. (Part 2-Wiss. Mitt.) pp. 23-24. 1908.
- ROUVILLE, M. E., DE. *Énumération des nématodes libres du canal des Bourdigues (Cette)*. Compt. Rend. Soc. Biol. (Paris). 55: 1529. 1903.
- SAVELJEV, S. N. *Zur Kenntnis der freilebenden Nematoden des Kolafjords und des Relictensee Mogilnoje*. Trav. Soc. Nat. St. Petersb. 43: 113-114. 1912.
- SCHMIDT, A. *Ueber den Rüben-Nematoden (Heterodera schachtii A. S.)*. Ztschr. Ver. Rübenz-Indus. im Zollverein. 21: (n.F. v. 8) 1-18. 1871.
- SCHULTZE, M. See Carus, J. V. 1857.
- SCHULZ, E. *Beiträge zur Kenntnis mariner Nematoden aus der Kieler Bucht*. Zool. Jahrb. Abt. Syst. Ökol. u. Geograph. Tiere. 62: 355, 358, 366, 407. 1932.
- SOUTHERN, R. *Clare Island survey part 54 Nematelmia, Kinorhyncha, and Chaetognatha*. Roy. Irish Acad. Proc. 31: 9-10, 11, 20, 29, 31, 34-35, 41-42, 66-67. 1914.
- STEINER, G. *Freilebende Nematoden aus der Schweiz*. Arch. Hydrobiol. u. Planktonkunde. 9: 426, 427. 1914.
- *Freilebende Nematoden aus der Barentssee*. Zool. Jahrb. Abt. System., Geogr. u. Biol. Tiere. 39: 579-580, 583, 603-604, 652. 1916.
- *Helminthological Society of Washington Proceedings*. Jour. Parasitol. 14: 66. 1927.
- *Neoplectana glaseri, n. g., n. sp. (Oxyuridae), a new nemic parasite of the Japanese beetle (Popillia japonica Newm.)*. Jour. Wash. Acad. Sci. 19: 437-438. 1929.
- *Die Nematoden der Deutschen Südpolar-Expedition 1901-1903. I. Teil*. Deut. Südpolar Expedition. Zool. 20: 207. 1930.
- *Die Nematoden der Deutschen Südpolar-Expedition 1901-1903. II. Teil*. Deut. Südpolar Expedition 1901-1903. Zool. 20: 356. 1931.
- *Neotylenchus abulbosus n. g., n. sp. (Tylenchidae, Nematoda) the causal agent of a new nematosis of various crop plants*. Jour. Wash. Acad. Sci. 21: 538. 1931.
- *On the status of the nemic genera Aphelenchus Bastian (sic), Pathoaphelenchus Cobb, Paraphelenchus Micoletzky, Parasitaphelenchus Fuchs, Isonchus Cobb, and Seinura Fuchs*. Jour. Wash. Acad. Sci. 21: 468-475. 1931.
- AND BUHRER, E. M. *The nematode Tylenchus similis Cobb, as a parasite of the tea plant (Thea sinensis, Linn.), its sexual dimorphism, and its nemic associates in the same host*. Ztschr. Parasitenk. 5: 419. 1933.
- AND HOEPLI, R. *Studies on the exoskeleton of some Japanese marine nemas*. Arch. Schiffs-u. Tropenhyg. 30: 573. 1926.
- STEKHOVEN, J. H. SCHUURMANS, JR. *Ökologische und morphologische notizen über Zuiderseenematoden. I. Die westliche Hälfte der Zuidersee*. Ztschr. Morph. u. Ökol. Tiere. Abt. A. 20: 648-649. 1931.
- AND ADAM, W. *The freelifving marine nemas of the Belgian coast*. Mém. (49) Mus. Roy. Hist. Nat. Belg., pp. 42-44. 1931.
- AND CONINCK, L. DE. *Diagnoses of new Belgian marine nemas*. Bul. Mus. Roy. Hist. Nat. Belg. 9: 9-10. 1933.
- STILES, C. W. AND HASSALL, A. *The determination of generic types and a list of round-worm genera with their original and type species*. U. S. Dept. Agr., Bur. Anim. Indus., Bull. No. 79, pp. 100, 101, 107, 122. 1905.
- THORNE, G. *Nematodes from the summit of Long's Peak, Colorado*. Amer. Micros. Soc. Trans. 48: 189-190. 1929.
- *Predacious nemas of the genus Nygolaimus and a new genus, Sectonema*. Jour. Agr. Research (U. S.). 41: 464. 1930.
- VILLOT, A. *Recherches sur les helminthes libres ou parasites des côtes de la Bretagne*. Arch. Zool. Expt. et Gén. 4: 463. 1875.
- ZUR STRASSEN, O. *Bradynema rigidum v. Sieb.* Ztschr. Wiss. Zool. 54: 655-742. 1892.

- | | | | | | | |
|--------------------------------|----------------------------|-------------|-------------------------------------|-------------------------------|---------------------------------------|-------------------------------------|
| Eurystomina..... | 12, 13, 19 | <i>page</i> | <i>Macrolaimus</i> , Ditlevsen..... | 24, 31 | <i>Paradesmolaimus</i> | 30 |
| Eutelolaimus..... | 5, 21 | | <i>Macrolaimus</i> Maupas..... | 13, 14, 25, 26 | <i>Paralinhomoeus</i> (subgenus)..... | 30 |
| Eutylenchus..... | 8 | | <i>Macroposthonia</i> | 5, 21 | <i>Paramonhystera</i> | 23 |
| F | | | <i>Margonema</i> | 22, 24 | <i>Paranticoma</i> | 22, 29 |
| <i>Fenestrolaimus</i> | 19 | | <i>Marionella</i> | 12, 13, 19 | <i>Paraphanolaimus</i> | 27, 30 |
| <i>Fiadra</i> | 17, 19 | | <i>Meloidogyne</i> | 9 | <i>Paraphelenchus</i> | 10 |
| <i>Filipjeva</i> | 28 | | <i>Meroviscosia</i> | 19 | <i>Paraplectus</i> | 27, 30 |
| <i>Filipjevia</i> | 18 | | <i>Mesodorius</i> | 16, 18, 19 | <i>Parasabotia</i> | 7, 12, 14, 21, 23 |
| <i>Filoncholaimus</i> | 19 | | <i>Mesochium</i> | 16, 16, 18, 18 | <i>Parasitaphelenchus</i> | 9, 10, 10, 25 |
| <i>Fimbria</i> | 17, 17, 27, 29 | | <i>Metachromadora</i> | 15, 15 | <i>Parasphaerolaimus</i> | 23 |
| <i>Fimbriella</i> | 28 | | <i>Metalaemus</i> | 23 | <i>Parasymphlocostoma</i> | 5, 6, 7 |
| <i>Fimbrilla</i> | 17, 17, 27, 29 | | <i>Metalinhomoeus</i> | 16, 21, 26 | <i>Paratylenchus</i> | 9 |
| <i>Fusonema</i> | 16 | | <i>Meteppsilonma</i> | 5, 27 | <i>Paratylenchus</i> | 18, 22, 24 |
| G | | | <i>Metoncholaimus</i> | 18 | <i>Pareuchromadora</i> | 11, 12 |
| <i>Galeonema</i> | 24, 30 | | <i>Micoletzkyia</i> | 6 | <i>Pareurystomina</i> | 13 |
| <i>Gammanema</i> | 17, 17 | | <i>Microlaimus</i> | 14 | <i>Parironus</i> | 17 |
| <i>Gonionchus</i> | 18 | | <i>Micromicron</i> | 14, 20 | <i>Paroncholaimus</i> | 19 |
| <i>Graptonema</i> | 11 | | <i>Mitrephephorus</i> | 4, 4 | <i>Paroxystomina</i> | 6, 22 |
| <i>Greiffella</i> | 8, 30 | | <i>Molgolaimus</i> | 4 | <i>Pathoaphelenchus</i> | 9, 10, 10, 25 |
| <i>Gullarnia</i> | 28 | | <i>Monhystera</i> | 7, 20, 20, 22, 23, 26, 26, 28 | <i>Pelagonema</i> | 22, 24, 28, 29, 31 |
| <i>Gymnolaimus</i> | 25, 28 | | <i>M. dinteriana</i> | 20, 22, 26, 28 | <i>Pelagonemella</i> | 26 |
| H | | | <i>Monhysteriella</i> | 20 | <i>Penzancia</i> | 23 |
| <i>Halalaimoides</i> | 5, 6, 27, 28 | | <i>Monhystrella</i> (subgenus)..... | 20, 26 | <i>Pepsinema</i> | 16, 16, 16, 18, 18 |
| <i>Halalaimus</i> | 6, 6 | | <i>Monhystrium</i> | 17, 23, 23 | <i>Peronilaimus</i> | 25 |
| <i>Halanonchus</i> | 29 | | <i>Mononchoioides</i> | 15, 16 | <i>Phanoderma</i> | 22, 22 |
| <i>Halaphanolaimus</i> | 4, 5, 7, 8, 26, 27, 30, 30 | | <i>Mononcholaimus</i> | 11, 12 | <i>Phanodermella</i> | 6 |
| <i>Halenchus</i> | 8, 9 | | <i>Mononchulus</i> | 11, 18 | <i>Phanodermopsis</i> | 8, 24, 30 |
| <i>Halichoanolaimus</i> | 18, 24, 30 | | <i>Mononchus</i> | 12, 18, 19 | <i>Pharetrolaimus</i> | 8, 9 |
| <i>Halinema</i> | 30, 30 | | <i>Monoposthia</i> | 14 | <i>Platycoma</i> | 7, 8, 30, 30 |
| <i>Haliplectus</i> | 5, 26, 27 | | <i>Mytolaimus</i> | 25 | <i>Platycomopsis</i> | 6, 28, 30 |
| <i>Halalaimus</i> | 7 | | <i>Mylolaimus</i> | 16 | <i>Plectonchus</i> | 13, 20 |
| <i>Hemicyclophora</i> | 4, 4 | | <i>Mylolaimus</i> | 25 | <i>Plectus</i> | 21, 27 |
| <i>Heterocephalus</i> | 22, 22 | | N | | <i>Poikilolaimus</i> | 13, 14 |
| <i>Heterodera</i> | 9, 9 | | <i>Nannolaimus</i> | 8, 24 | <i>Polyodontus</i> | 18 |
| <i>Heterodesmodora</i> | 8 | | <i>Nannonchus</i> | 12, 19 | <i>Polygastrophora</i> | 13, 15, 19 |
| <i>Hexatylus</i> | 8 | | <i>Nanonema</i> | 11, 13 | <i>Polyalaimus</i> | 27, 27 |
| <i>Himatidiophila</i> | 21, 21 | | <i>Nasinema</i> | 22 | <i>Polysigma</i> | 16 |
| <i>Hoplolaimus</i> | 8, 9 | | <i>Necticonema</i> | 12, 24 | <i>Pomponema</i> | 12, 18 |
| <i>Howardula</i> | 6, 8, 10, 14 | | <i>Nemanema</i> | 7, 23 | <i>Pontonema</i> | 19 |
| <i>Hypodontolaimus</i> | 13, 14 | | <i>Nemanemella</i> | 7 | <i>Porocoma</i> | 6, 28 |
| <i>Hyptiolaimus</i> | 18 | | <i>Nemella</i> | 5, 6 | <i>Praeaecanthonchus</i> | 12 |
| I | | | <i>Nemonechus</i> | 8 | <i>Prismatolaimus</i> | 28, 28, 29, 30 |
| <i>Ilium</i> | 28 | | <i>Neoaplectana</i> | 4, 5, 21 | <i>Pristionchus</i> | 15, 16, 16 |
| <i>Ionema</i> | 6, 6, 8, 31 | | <i>Neochromadora</i> | 14 | <i>Pristionema</i> | 6 |
| <i>Iota</i> | 9, 9 | | <i>Neodiplogaster</i> | 13, 15 | <i>Prochaetosoma</i> | 5, 27 |
| <i>Iotadorus</i> | 13, 14, 14 | | <i>Neonchus</i> | 10, 14 | <i>Prochromadora</i> | 13, 13, 14 |
| <i>Iotalaimus</i> | 4, 6 | | <i>Neonyx</i> | 15 | <i>Prochromadorella</i> | 15, 17 |
| <i>Iotonchium</i> | 4, 4, 10, 10 | | <i>Neotonchus</i> | 14 | <i>Procciconema</i> | 9 |
| <i>Ironella</i> | 17 | | <i>Neotylenchus</i> | 8, 9 | <i>Prodesmodora</i> | 15, 27 |
| <i>Ironus</i> | 17 | | <i>Neurella</i> | 22 | <i>Promononchus</i> | 6, 6, 8, 11, 21, 22, 24, 28, 28, 30 |
| <i>Isolaimium</i> | 28, 28, 30 | | <i>Notochaetosoma</i> | 4, 5, 8, 20, 25, 27 | <i>Proönccholaimus</i> | 18 |
| <i>Isonchus</i> | 10, 10 | | <i>Nuada</i> | 6 | <i>Prosphaerolaimus</i> | 12 |
| <i>Isonemella</i> | 12, 29 | | <i>Nuadella</i> | 4, 5 | <i>Psellionema</i> | 4, 6, 6, 11, 25, 29 |
| J | | | <i>Nudolaimus</i> | 23, 29 | <i>Pseudoalolaimus</i> | 30, 30 |
| <i>Jägerskiöldia</i> | 19 | | <i>Nudora</i> | 13, 14 | <i>Pseudochromadora</i> | 21, 21 |
| K | | | <i>Nunema</i> | 24 | <i>Pseudodilaimus</i> | 12 |
| <i>Klugia</i> | 8 | | <i>Nygmatochus</i> | 14 | <i>Pseudodella</i> | 15, 16, 20, 25, 28 |
| <i>Krampia</i> | 21 | | <i>Nygolaimus</i> | 11 | <i>Pseudoncholaimus</i> | 18 |
| <i>Kreisia</i> | 27 | | O | | <i>Pseudonchus</i> | 21, 27 |
| L | | | <i>Odontolaimus</i> | 10, 14, 14 | <i>Pseudoparonycholaimus</i> | 19 |
| <i>Laimella</i> | 4, 26 | | <i>Odontonema</i> | 13, 15 | <i>Pseudopelagonema</i> | 24 |
| <i>Lasiomitus</i> | 5, 6, 7 | | <i>Odontopharynx</i> | 14, 15, 16, 16, 18 | <i>Pseudorhabditis</i> | 15 |
| <i>Laxonema</i> | 4, 4, 24, 26 | | <i>Odontophora</i> | 15, 17, 17 | <i>Psilenchus</i> | 10 |
| <i>Laxus</i> | 5, 21, 27 | | <i>Ogma</i> | 9, 9 | <i>Pteronium</i> | 22, 29 |
| <i>Ledovitia</i> | 12, 13 | | <i>Oionchus</i> | 11, 14 | <i>Ptycholaimellus</i> | 13, 14 |
| <i>Leptogastrella</i> | 23 | | <i>Oistolaimus</i> | 10, 14 | <i>Pulchranemella</i> | 29 |
| <i>Leptolaimus</i> | 27 | | <i>Omicronema</i> | 30 | <i>Punctodora</i> | 13, 14 |
| <i>Leptonchus</i> | 11 | | <i>Onchium</i> | 10, 12, 15, 21, 24, 27, 31 | <i>Pycnelaimus</i> | 16, 26 |
| <i>Leptonemella</i> | 4, 4, 20, 20 | | <i>Oncholaimellus</i> | 19 | Q | |
| <i>Leptosomatides</i> | 7, 23 | | <i>Oncholaimium</i> | 18 | <i>Quadricoma</i> | 6, 7, 28, 28, 30 |
| <i>Leptosomatium</i> | 8 | | <i>Oncholaimus</i> | 18 | R | |
| <i>Leptosomella</i> | 8 | | <i>Onchuelia</i> | 10, 10 | <i>Rhabditella</i> | 25 |
| <i>Linhomoea</i> | 5, 7, 7, 24, 26, 26, 30 | | <i>Onchulus</i> | 11 | <i>Rhabditidoides</i> | 25 |
| <i>Linhomoeus</i> | 5, 7, 7, 16, 21, 24, 30 | | <i>Onyx</i> | 10, 10, 14, 15 | <i>Rhabditis</i> | 13, 25, 25, 25, 26, 27 |
| <i>L. obtusicaudatus</i> | 7 | | <i>Oxystoma</i> | 7 | <i>Rhabditoides</i> | 20, 25 |
| <i>Linolaimus</i> | 27 | | <i>Oxystomina</i> | 7, 7 | <i>Rhabditolaimus</i> | 26 |
| <i>Litinium</i> | 7, 29 | | P | | <i>Rhabditophanes</i> | 20, 26 |
| <i>Litonema</i> | 4, 4, 24, 26 | | <i>Panagrolaimus</i> | 20, 21, 25 | <i>Rhabdocoma</i> | 22, 22, 23, 24 |
| <i>Litotes</i> | 5, 6, 7, 28, 28, 29 | | <i>Pandolaimus</i> | 24, 31 | <i>Rhabdodemia</i> | 17 |
| <i>Longicyatholaimus</i> | 12 | | <i>Paracanthonchus</i> | 12 | <i>Rhabdogaster</i> | 5, 27 |
| <i>Lycolaimus</i> | 25 | | <i>Parachromagaster</i> | 29 | <i>Rhabdolaimus</i> | 14, 25, 25, 27 |
| | | | <i>Paraecthonolaimus</i> | 16 | <i>Rhabdonchus</i> | 10, 10, 10 |
| | | | <i>Paracyatholaimus</i> | 12 | <i>Rhabdotoderma</i> | 19 |

	<i>page</i>		<i>page</i>		<i>page</i>
Rhadinema.....	23, 29	Stenolaimus.....	6, 8, 22, 22, 24	Tristricochaeta.....	4, 5, 20, 25, 27
Rhinema.....	12, 15	Stephanolaimus.....	6, 8, 17, 21, 23,	Trogolaimus.....	17, 22
Rhinonema.....	10, 10, 11, 12, 13		29	Tubolaimella.....	17, 29
Rhrips.....	11, 14, 16, 19	Stilbonema.....	4, 5	Tubolaimus.....	23, 29
Rhodolaimus.....	25	Symplocostoma.....	11, 12, 12, 13, 19	Turbatrix.....	13, 18, 20
Rhynconema.....	29	Symplocostomella.....	12, 13, 19	Tycnodora.....	6
Richtersiella.....	8, 12, 23, 24, 30, 30	Syndontium.....	16, 20	Tylenchinema.....	6, 10
Richtersiella.....	23, 24, 30, 30	Synonchiella.....	18	Tylenchodn.....	9
Sabatieria.....	7, 12, 14, 21, 23	Synonchium.....	17, 18	Tylencholaimellus.....	8
Saveljevia.....	18	Synonchus.....	19	Tylencholaimus.....	9, 9
Scaptrella.....	17	Synonema.....	14	Tylenchomorphus.....	5, 8, 9, 9, 10,
Scatonema.....	5	Syringolaimus.....	15		10
Schistodera.....	7, 7	Tachyhodites.....	7, 23	Tylenchorhynchus.....	9
Sectonema.....	11	Teratocephalus.....	16, 21, 21,	Tylenchulus.....	9, 9, 9
Seinura.....	9, 10, 10, 25		25, 25, 26, 27	Tylenchus.....	9, 9
Selachinema.....	18	Terschellingia.....	4, 5, 21	Tyololaimophorus.....	8, 9, 10
Seleneella.....	16	Thalassironus.....	17	Tylopharynx.....	9, 25
Seuratia.....	12	Thalassolaimus.....	5, 6	Udonchus.....	14
Seuratiella.....	12	Theristus.....	17, 22, 23, 23	Uracanthus.....	5, 10
Segmophora.....	15	Thoñchus.....	19	Vasculonema.....	22, 24, 28, 29, 31
Siphonolaimus.....	9, 10, 26	Thoracostoma.....	19	Viscosia.....	19
Smalsundia.....	18, 24, 30	Thoracostomopsis.....	10	Walcherenia.....	20
Solenolaimus.....	4, 26	Trefusia.....	7	Wilsonema.....	27
Southernia.....	7, 8	Trichoderma.....	8, 30	Xanthodora.....	13, 15
Southerniella.....	30	Trichodorus.....	11	Xennella.....	6, 28
Sphaerocephalum.....	23, 29	Tricoma.....	6, 7, 28, 28, 30	Xenolaimus.....	22
Sphaerularia.....	5	Trigonolaimus.....	15, 17, 17	Xenonema.....	15
Spilophora.....	15	Trilepta.....	11, 27, 28	Xinema.....	10, 18
Spilophorella.....	13	Trileptium.....	11, 27, 28	Xiphinema.....	9
Spilophorium.....	15	Trilobus.....	11, 19, 22, 22	Xyala.....	17, 23
Spira.....	5, 14, 15, 21, 21	Triodontolaimus.....	17		
Spirina.....	5, 14, 15, 21, 21	Triplonchium.....	9, 27	Ypsilon.....	12, 15, 21, 24, 31
Statena.....	15	Tripyla.....	6, 6, 8, 11, 21, 22, 24, 23,	Zalonema.....	21, 27
Statena Filipjev.....	6, 11		28, 30	Zanema.....	17, 23, 23, 29, 29
Steinera Ditlevsen.....	19	Tripylum.....	17, 29	Zygonemella.....	7, 23, 29
Steinera Stekhoven and de Coninck.....	23	Tripylodes.....	24, 24		
Steineriella.....	19	Trischistoma.....	11, 12, 22, 23, 28,		
			29		
		Trissonchulus.....	17		

Opuscula miscellanea nematologica, I. G. STEINER, U. S. Bureau of Plant Industry.

There will be published under this heading a series of observations on various nematodes and nematological problems resulting mostly from service work or observations of a more casual nature.




FIG. 1. *Anguillulina gallica*, n. sp.

A—Anterior end; about X 920. B—Vulva, vagina and portion of uterus; about X 1030. C—Posterior end of female with truncate terminus; about X 920. D—Face view of vulva, with dilatatores vaginae; about X 920. E—Tail end of second-stage larva; about X 1030. F—Tail end of male; about X 1030. G—Anterior end of second-stage larva; about X 1030.

- (1) ANGUILLULINA GALLICA, N. SP. (FIG. 1) LIVING IN BURLS OF AN ELM (ULMUS SP.) FROM FRANCE.

Remarks.—This new species of *Anguillulina* is very close to *Anguillulina major* (Fuchs, 1915) amend. The elm burls in which the present species was observed exhibited no mines or other traces of insect activities. The nemas were found throughout the wood, which also harbored a fungus. Larval specimens were most numerous and would immediately swarm into the water when a piece of wood was immersed. This was not true of the adults, which had to be dissected out of the wood. It is thought that this species feeds on fungi.

Anguillulina major, according to Fuchs, is an associate of the bark beetle, *Ips typographus* L., in the mines of which it was observed. Larval specimens are carried under the wing covers of this beetle and apparently distributed to new trees. It is thought that in the present case a bark or woodboring beetle may also be involved as carrier, although apparently the nema migrates by itself through a tree to which it initially may have been transported by a beetle. It is not yet known if it merely follows the fungus in this migration through a tree, or if it is an agent in the spread of the fungus.

Description.—*Larva:* The difference in shape between the larval and adult tails, as observed by Fuchs in *A. major*, is also seen in the present species, where the larva has a pointed (fig. 1, E), the female an obtuse, and the male a bursate, tail end. In addition, the larva has an oesophagus of much greater length in relation to the total body length than the adult. The excretory pore of the larva lies in front of the middle oesophageal bulb, whereas in the adult it is located ventrad of the terminal oesophageal bulb. The posterior portion of the oesophagus of these larvae is of special interest because of its extraordinary length and form (fig. 1, G). The portion following the nerve ring is somewhat swollen and set off posteriorly from the end portion by a faint constriction. It is in this end portion that a large nucleus may be seen, which seemingly belongs to the dorsal oesophageal gland. *Adults:* Cuticle annulated. Two lateral membranes border a lateral field about 1/3 as wide as body. No cervical papillae nor phasmids seen. Tail of female short-conical, with broad-obtuse or even truncate terminus (fig. 1, C). Male tail short-conical; terminus pointed, curved ventrad, surrounded by bursa (fig. 1, F). Head differing from that of *A. major* by being distinctly set off, about half as long as wide, not annulated; lips distinct. Buccal stylet not very strong, with distinct yet very small basal knobs. Middle oesophageal bulb very prominent, with short valves and rather strong muscle fibers. Terminus of oesophagus indistinct, hardly set off from intestine; wall of latter consisting of double series of large cells. Vulva (fig. 1, B,C,D) a long transverse slit, occupying almost 3/4 of body diameter, with 4 dilatator muscles, as shown in figure 1, D. Vagina leads inward and forward; its walls thick. Uterus without a postvulvar branch. Eggs slightly oblong, almost spherical. Male bursa without ribs, surrounding tail end; spicula longer than tail, slightly curved, proximal end not capitate; gubernaculum lineate, slightly curved, about 1/3 as long as spicula.

Measurements.—♀: total length = 1. to 1.1 mm; α = 34.8; β = 14.2; γ = 43.2; ν = 91.7%; ♂: total length, 0.63 to 0.97 mm; α = 39.41; β = 11.8-12.2; γ = 37.2-38.8.

Diagnosis.—*Anguillulina* resembling *A. major*, but differing in the following respects: Head distinctly set off; double lateral membrane; tail of female obtusely rounded or truncate, only 1/3 to 1/2 as long as distance from vulva to anus; uterus without postvulvar branch; spicula longer than male tail.

(2) *ACROBELES GLAPHYRUS*, N. SP. (FIG. 2).

Remarks.—A single adult female of this apparently new species was found in a diseased tuber of *Polyanthes tuberosa*, the tuberose, which originated in Hermosillo, Sonora, Mexico, and was intercepted by Foreign Plant Quarantine officials at Nogales, Arizona (November 1934). The species resembles very closely *Acrobeles punctatus* Thorne, 1925. The intestine was filled from beginning to end with a brownish mass in which no particles could be distinguished, the whole rather suggesting jellied plant material as food of this species.

Description.—Body almost cylindrical, tapering less anteriorly, more rapidly posteriorly to a short concave conoid, sharply pointed tail. Annulation of cuticle very distinct, interrupted on the sides by three lateral membranes separating two non-annulated longitudinal fields, except on a portion of anterior end and on extremity of tail where annulation encompasses the body. Width of annules about 2μ . Width of entire lateral field about $2\frac{1}{2}$ to 3 times width of an annule. Figure 2, A shows mode of ending of lateral field anteriorly and figure 2, D mode of ending posteriorly.

In the region of the lateral fields especially on the tail, but also here and there on the annules, small points are seen in a subcuticular layer, irregularly arranged, not in 4 transverse series. Cervical papillae and phasmids not seen. Labial probolae with 2 outward curved prongs; cephalic probolae separated by deep incision, with short basal prong, flanked on each side by a larger inward curved prong. Cephalic papillae and amphids apparently on these cephalic probolae anterior to the annules, the papillae (only submedial ones were distinctly seen) forming low, rounded elevations, the amphidial opening apparently a curved transverse slit. Buccal armature as follows: cheilorhabdion short, thin; prorhabdions rather long, thin; only dorsal mesorhabdion present, thin, hardly separated from prorhabdion; metarhabdion short, thick, the dorsal with sharply pointed denticle; telorhabdion short, thick. Corpus and middle bulb of oesophagus hardly differentiated, thin, slender; terminal bulb relatively large, almost spherical, with well developed valvular apparatus. Excretory pore situated well cephalad, in the region of the corpus, far in front of nerve ring. Lips of vulva small. No details about female apparatus available.




FIG. 2. *Acrobeles glaphyrus*, n. sp.

A—Anterior end; mode of formation and place of beginning of lateral membranes and lateral field are shown; about X 450. B—Front view of head (ventral side up); about X 1800. C—Head end, lateral view; about X 1800. D—Tail end of female; about X 450.

	5.1	23.	29.	14	32	
Measurements.—♀				59	94.6	.71 mm
	4.7	6.7	7.2	7.2	3.6	

$\alpha = 13.8$; $\beta = 3.4$; $\gamma = 18.5$.

Diagnosis.—*Acrobeles* closely related to *A. punctatus* but different because of absence of dots on surface of annules; annules much smaller, in the region of the buccal cavity 9 instead of 5 as in *A. punctatus*; labial probolae with outward arching prongs; unannulated lateral fields with a central and 2 bordering membranes; excretory pore far in front of nerve ring.

Type locality.—Mexico (Sonora).

Type association.—Diseased tuberose bulb.

(3) CEPHALOBUS MAXIMUS (THORNE, 1925) AMEND. (FIG. 3).




FIG. 3. *Cephalobus maximus* (Thorne) amend.

A—Anterior end; about X 640. B—Vulvar region; about X 640. C—Tail end of female; *dil msc*, dilator muscle of rectum and anus; *phas*, phasid; about X 640. D—Head end; *cph pap*, cephalic papilla; *chlorhd*, cheilorhabdion; *prorhd*, prorhabdion; *metarhd*, metarhabdion; *telorhd*, telorhabdion; about X 824. E—Male tail; *phas*, phasid; about X 824.

Remarks.—Only the female of this large *Cephalobus* species has been known. Specimens (including a male) of what appeared to be this form were found in a bulb of *Iris xiphioides* from San Diego, California (April 1933). Associated with them were *Anguillulina dipsaci* (Kühn, 1857) Gerv. and v. Ben. 1859, *Aphelenchoides parietinus* (Bastian, 1865) Steiner, 1932, and *Acrobeles complexus* Thorne, 1915. Thorne's description of the female of *C. maximus* fits the present specimens fairly well. Most outstanding is the apparent absence of a pronounced middle bulbous swelling of the oesophagus. However, it is not thought that this is enough difference to warrant systematic separation of the present specimens from *C. maximus*. It is not our intention to furnish here a new description of this species, but a few additional characters will be mentioned. There are seemingly 2 submedial and 1 lateral cephalic papillae each. Amphids of the type characteristic of this genus were also seen. Of the buccal armature cheilo-, pro-, meta- and mesorhabdions are present (fig. 3, D). It is interesting to note that the excretory apparatus is composed of 4 cells, 2 larger and 2 smaller ones (fig. 3, A). The dilatator ani is divided into a smaller anterior muscle located about halfway on the rectum and a larger one placed just at the anal opening (fig. 3, C). The uterus of the female still has a posterior branch (fig. 3, B, C).

Description of the male.—Tail similar to that of female, but ventrally bent (fig. 3, E). Copulatory papillae as follows: One ventrosubmedial, a short distance in front of the spicula; 1 ventrosubmedial in latitude of anus; a pair in front of phasmid less than halfway down the tail (1 lateral and 1 ventrosubmedial); 1 dorsosubmedial at about the beginning of the last fourth of the tail, a lateral one and a ventral one close to terminus of tail. Copulatory muscles well developed, extending forward from anus almost $2\frac{1}{2}$ body widths. Spicula slightly curved, distally pointed, widening proximad, but inner end again narrow, not capitate. Gubernaculum about $\frac{3}{4}$ as long as spicula; distal half slightly swollen, proximal half linear. Testis reflexed.

Measurements.— δ : total length = 0.9 mm; $\alpha = 27$; $\beta = 7.2$; $\gamma = 18$.

(4) ON THE SYNONYMY OF ANGUILLULINA COSTATA (DE MAN, 1921) GOODEY, 1932, AND ANGUILLULINA CANCELLATA (COBB, 1925) GOODEY, 1932.

Goodey (1932, J. Helminth. 10: 156) expressed the opinion that these 2 nemie species mentioned above were probably synonyms. The writer recently obtained a female specimen of what he considered to be *A. costata*. It was found on diseased rutabagas (*Brassica campestris*) and was associated with numerous specimens of *Anguillulina pratensis* (de Man, 1880) Goffart, 1929. The host plants came from Arlington Farm, Va., where the specimens originally described by Cobb as *A. cancellata* also were obtained. A careful study of our specimen and a detailed comparison of the description of *A. costata* as well as *A. cancellata* convince us that *A. cancellata* is the synonym of *A. costata*. It may be well to record that the length of the oesophagus as given in Cobb's measurement formula, i. e. 30% of the total length, is a misprint. As may be seen from the figure, this length is actually only 20% of the total length. Our specimen has the following dimensions:

	2.6	12.	18.	^{36-64.}	84.	
♀	<hr/>					.47 mm or $\alpha = 23.2$; $\beta = 5.4$; $\gamma = 6.4$.
	2.5	3.7	4.2	4.3	2.	

Our specimen did not exhibit striae in the head cuticle. The number of longitudinal ridges on the cuticle was 18; they were all equidistant.

The nematode *Neotylechus abulbosus* Steiner (Anguillulidae) as a parasite of sugar-beets. GERALD THORNE and CHARLES PRICE, U. S. Bureau of Plant Industry (Salt Lake City, Utah).

Sugar-beets collected at Chino, California, July 16, 1934, were infested with *Neotylechus abulbosus* Steiner, 1931. Injury consisted of variable-sized lesions covering considerable portions of the beets (fig. 4). In some instances the lesions extended practically through the root; at a depth of 3 inches the root of one beet had been completely severed. The nemas were found invading the living tissues while the dead portions were rapidly decaying and contained large numbers of the saprophytic nema, *Aphelenchus avenae* Bastian, 1865. The infested area covered several square rods and in it the beets were almost completely destroyed (fig. 5). Judging from the severity of the injury in this instance the nema may become an exceedingly destructive parasite if widely disseminated.



FIG. 4
Sugar-beet infested with *Neotylechus abulbosus*. Cortex peeled off to give the lesions greater contrast.



FIG. 5
Sugar-beet field infested with *Neotylechus abulbosus*. (Photograph by Charles Price, April 14, 1934.)

Notes on free-living and plant-parasitic nematodes, I. GERALD THORNE, U. S. Bureau of Plant Industry (Salt Lake City, Utah).

(1) *The occurrence of Diphtherophora perplexans Cobb, syn. Archionchus perplexans Cobb.*—In August 1923, a single specimen of *Diphtherophora perplexans* Cobb was collected near Provo, Utah. During the following 10 years dozens of soil samples from Utah fields were examined without the appearance

of another specimen. However in 1934 large numbers of this species appeared in several collections from widely separated points. In the fields at Salem and Lewiston, where mononch studies have been carried on for 10 years, many specimens of *D. perplezans* were present, the first seen from these fields. There is no apparent reason for this sudden increase in numbers when the population of many nemie species has been greatly reduced by a series of 6 years deficient in rainfall, of which 1934 was the driest. One other collection was made in 1930 in soil from a date garden, Indio, California.

(2) *The effect on nematodes of copper sulphate used in snail control.*—East Canyon Creek near Park City, Wasatch County, Utah, was treated with copper sulphate by Dr. L. E. Swanson, U. S. Bureau of Animal Industry. A concentration of 1:500,000 for 24 hours was used. Bank and bottom material was collected before and after treatment at 2 points about 150 feet and 2 miles respectively below the mixing station. At neither point was there any apparent effect on the nemie population. Species present were: *Dorylaimus stagnalis*, *D. consobrinus*, *Monhystera paludicola* (?), *Aphanolaimus aquaticus*, *Plectus cirratus*, *Trilobus pellucidus* and *Mononchus macrostoma*.

(3) *The taxonomic status of the nematode Tylenchorhynchus cylindricus Cobb, 1913.*—A reexamination of Cobb's original balsam mounts of *Tylenchorhynchus cylindricus* reveals the fact that this species actually is *Anguillulina dubia* (Bütschli, 1873) Goodey, 1932. Cobb's specimens differ only slightly from those described by Bütschli and deMan as *A. dubia*. Apparently such distinguishing characters as are mentioned in Cobb's description and figures were due to fixation distortion. For our present information *Tylenchorhynchus* is best considered a synonym of *Anguillulina* since it exhibits no separating characters.

A review of the fossil nematodes. A. L. TAYLOR.

Three short papers about fossil nematodes have been found in the literature. Two of these papers, both dealing with nematodes in amber, seem to have escaped the attention of most workers in helminthology.

The best known fossil nematode is *Mermis antiqua* reported by v. Heydon (1862, *Paleontographica* 10:62-82, pl. 10). It was found projecting from the anus of a beetle (*Hesthesis immortua* Heydon) from the Rhine lignite. The single specimen was about 25 mm long and is shown in v. Heydon's drawing in a much coiled position (fig. 6, A). Later writers have agreed that this is a nematode and not a gordiid, but it is impossible to be sure that it belongs to the genus *Mermis*. In view of this a new collective fossil genus *Heydonius* is proposed for it and other fossil mermithids. The name of this species becomes *Heydonius antiquus* (v. Heydon, 1862). The Rhine lignite was formed in the Eocene, the lowest division of the Tertiary.

Specimens in a better state of preservation have been reported from amber. As one report comes from Königsberg and the other from Danzig, both cities near the Baltic Sea, it is probable that the amber is from this vicinity and dates from the Lower Oligocene, the next to lowest division of the Tertiary.

H. v. Duisburg (1862, *Schr. Königlichen Phys.-Okonom. Gesell.* 3:31-36, pl. 1) reported 3 specimens. One of these, *Anguillula succini*, was found in a small piece of amber which also contained pollen grains similar to those of *Pinus sylvestris* L. Its description is as follows: Length 855 μ , width 16 μ at widest part (about 1/5 of length from head). Body width diminishes gradually from this point to the knob-like head and pointed tail. Body annulated, annules 2 μ wide. Tail ventrally curved. A strand 1/3 to 1/4 the body width runs from head to a point shortly in front of tail—presumably remains of oesophagus and intestine. From the body shape, and other characters, it seems probable that this species belongs to the Plectidae, and the new collective fossil genus *Oligoplectus* is proposed to contain it and other fossil Plectidae. The name becomes *Oligoplectus succini* (v. Duisburg, 1862) (fig. 6, F).




FIG. 6. Fossil nematodes.

A—*Heydonius antiquus* (v. Heydon, 1862). B-C—*Heydonius matutinus* (Menge, 1866). D—*Vetus pristinus* (Menge, 1866). E—*Vetus capillaceus* (Menge, 1866). F—*Oligo-plectus succini* (v. Duisburg, 1862). G-H—*Vetus duisburgi*, n.g., n.sp. (A, redrawn from v. Heydon; B, C, D and E, redrawn from Menge; F, G and H, redrawn from v. Duisburg.)

Two other specimens were also included by v. Duisburg under the name *Anguillula succini*. They were found in a small piece of amber which contained the cast off cuticle of an unidentifiable insect but there is no evidence to show that this association is anything but accidental. As they are not congeneric with the previous specimen, the genus *Vetus* is proposed as a new collective genus to contain these and other fossil free living species which cannot be identified to family. This species becomes *Vetus duisburgi* n.g., n. sp. The two specimens are described as follows: Lengths 850 μ and 650 μ . Body widths 18 μ and 17 μ respectively. Body fusiform, head rounded, tail bluntly pointed. Anterior portion of the smaller specimen evidently crushed and measures 23 μ at widest part. Annules about 1 μ wide on both specimens (figs. 6, G, H).

A. Menge (1866, Schr. Naturf. Gesell. Danzig v. 1, pt. 3-4) reported a number of specimens, also in amber from the Baltic. His first species, *Mermis matutina* Menge, 1866, is represented by 3 specimens associated with a chironomid (Diptera) (fig. 6, B). One of these is reported to have its head end between 2 segments of the abdomen of the insect, while the other 2 were coiled around and near it. Description: Length 3.5 mm, width 0.1 mm. Body cylindrical, bluntly pointed at head and tail. Color yellowish white, in life probably white or reddish white. Annules about 300. The only sign of internal organs is a cylindrical canal which does not run to the end of the body. The specimen which had its "head end" between the segments of the insect also had a protuberance near the other end from which 2 "spicules" projected. It seems probable that the worms were just leaving the host. If this assumption is correct, the "spicules" noted by Menge were on the wrong end of the worm, for mermithids leave the host head first. If we assume that the objects really were spicules and not some accidental bit of trash, the specimens would be excluded from the genus *Mermis* as we know it today, for *Mermis* has only 1 spicule. The annules observed by Menge were explained as postmortem changes due to the contraction of the longitudinal muscles. It seems quite possible that this could be the case.

As these specimens are probably mermithids, but not members of the genus *Mermis*, it is proposed that they be placed in the genus *Heydonius* to be known as *H. matutinus* (Menge, 1866).

Under the name *Anguillula pristina* Menge includes over 30 specimens of small, evidently free-living nematodes, contained in another piece of amber. The length is given as 139μ and the width as 3μ , but his drawings (fig. 6, D) show the specimens to be much thicker than these dimensions would indicate. Body fusiform, annulated and widest in front of the middle. Head rounded, tail pointed. Some specimens show remains of intestine. Some variation in size and annulation between individuals. Not enough description is given to determine the systematic position. It is proposed that these be included in the genus *Vetus* as *V. pristinus* (Menge, 1866).

The same author also describes about 50 nematodes in a third piece of amber under the name *Anguillula capillacea*. These vary in length from 0.7 to 2.0 mm and are 0.1 mm wide. Body cylindrical, head rounded, tail pointed. (Head and tail seem to be confused in the original description of both this and the other free-living species.) There is no trace of annulation. Here, also, not enough description is given to place the specimens in their proper systematic position, so it is proposed that these specimens be placed in the genus *Vetus* to be known as *V. capillaceus* (Menge, 1866) (fig. 6, E).

Thanks are due to Dr. R. S. Bassler, of the U. S. National Museum, for geological information and to Dr. B. G. Chitwood for help and the use of the drawing of *Heydonius antiquus*.

A new nematode, *Camacolaimus prytherchi*, n. sp. (Camacolaimidae).

B. G. CHITWOOD, U. S. Bureau of Animal Industry.

During the past summer the writer made numerous collections of marine nematodes while at the Bureau of Fisheries Laboratory, Beaufort, N. C. Representatives of most of the groups of marine nematodes were collected and are being used for anatomic study. Some of these appear to be new and are being described. The histology of the species described here will be published later.

Camacolaimus prytherchi, n. sp.

Description.—Oral opening subtriangular. Cephalic papillae consisting of 6 small papillae of internal circle and 4 setae of external circle (?dorsodorsals and ventroventrals). Dorsal tooth short, bluntly truncate. Cuticle striated; lateral alae present; sublateral somatic setae present. Excretory cell situated about $2/3$ to 1 esophageal length from anterior end of body. Male 1.31 mm long; $a = 43$, $\beta = 5.9$, $\gamma = 13.1$. Testis extending to about 40 percent of body length from anterior extremity. Female 1.69 mm long; $a = 52$, $\beta = 8.45$, $\gamma = 19$. Vulva dividing body in proportions 52:48. Anterior ovary extending 280μ anteriorad, reflexed; posterior ovary extending 180μ posteriorad, reflexed. Mature eggs (in 1- to 2-cell stage 87μ long by 25μ wide; 1 to 2 present.

Habitat.—Beach sand, just below low tide mark.

Locality.—United States (Beaufort, N. C.).

Specimens.—U. S. N. M. Helm. Coll. No. 40501 (cotypes).




FIG. 7. *Camacolaimus prytherchi*.

A—Head, en face view. B—Head, lateral view. C—Esophageal region, lateral view. D—Tail of male, lateral view. E—Excretory cell.

Camacolaimus prytherchi appears to be more closely related to *C. zostericola* (Filipjev, 1918) than to any of the other species of this genus, and differs from it in the dorsal tooth which is conical in *C. zostericola* and bluntly truncate in *C. prytherchi*.

Two new nematodes of the genus *Heth* Cobb, 1898 (Atractidae). M. B. CHITWOOD.

Two species of the genus *Heth* were found in a collection of nematodes from preserved specimens of millepeds in the U. S. National Museum. These appear to be new species, one being very similar to *Heth juli* Cobb, 1898 (Agric. Gaz. N. S. Wales 9:249, figs. 10, I-IV), the other similar to *Heth spinosum* Artigas, 1929 (? 1930, Systematica dos nematoideos dos arthropodes, These, São Paulo, pp. 81-83, pl. 32, fig. 1; pl. 33, fig. 1). Sexual dimorphism is so extreme in this group that it is difficult to be certain which males and females belong to the same species. Fortunately the males were found in the same collections as the females on several occasions and only one species of each in a collection. The new species are separated from previously described species by means of a key following the descriptions.

Heth dimorphum, n. sp.

Description.—**Male** 2.03 to 2.04 mm long by 90 μ wide. Oral opening surrounded by an internal circle of 6 papillae and an external circle of 4 papillae; 3 feather-like structures projecting from stoma. Cuticular ornamentations absent. Esophagus 610 to 660 μ long. Excretory pore not observed. Anus 260 μ from posterior extremity; tail attenuated. Genital papillae consisting of at least 1 pair of preanal and at least 4 pairs of postanal papillae. Spicules 140 to 150 μ long; nearly completely fused. Preanal organ vesiculate in lateral view, sucker-like in oblique view (fig. 8, D).

Female 2.28 to 4.22 mm long by 110 to 263 μ wide. Head in form of 2 lateral lobes, each bearing 2 papillae of external circle and an amphid. Subcephalic collar laterally shield-like; 2 pairs (in tandem) of double postcephalic lateral spines. Esophagus 400 to 454 μ long. Excretory pore not observed. Anus 45




FIG. 8.

Heth dimorphum, n. sp. A—Head of female. B—Head of male. C—Tail of male. D—Preanal organ, slightly oblique view. E—Esophageal region of male. F—Head of female, en face view. *Heth spinosum*, n. sp. G—Head of female. H—Esophageal region of female. I—Esophageal region of male. J—Tail of male. (All lateral views except as noted.)

to 67 μ from posterior extremity; tail attenuated. Vulva 18 to 22 μ anterior to anus; vagina directed anteriorly; uteri parallel. Eggs 130 to 170 μ long by 53 to 72 μ wide.

Host.—*Spirostreptus* sp. (milleped).

Location.—Intestine.

Localities.—Sumatra (Kepahiang) (type locality); Philippine Islands (Mt. Makiling, Luzon).

Specimens.—U. S. N. M. Helm. Coll. Nos. 31636 (type) and 31640 (paratypes).

Heth hexaspinosum, n. sp.

Description.—*Male* 1.45 to 1.52 mm long by 82 to 91 μ wide. Oral opening surrounded by internal circle of 6 papillae and an external circle of 4 papillae. Labial structures apparently absent, ornamentations absent. Esophagus 410 to 490 μ long. Excretory pore 210 μ from anterior extremity. Anus 298 to 306 μ from posterior extremity; tail attenuated. Genital papillae consisting of 1 pair of adanal and 6 pairs of postanal papillae; phasmids lateral. Spicules 68 to 70 μ long, nearly completely fused. Preanal organ vesiculate (fig. 8, J).

Female 1.82 to 2.07 mm long by 91 to 109 μ wide. Head in form of 2 lateral lobes (fig. 8, G), each bearing 2 papillae of external circle and an amphid. Subcephalic collar not shield-like; 2 median and 4 submedian postcephalic spines. Esophagus 327 to 363 μ long. Excretory pore 160 μ from anterior extremity. Anus 500 to 545 μ from posterior extremity; tail attenuated. Vulva 89 to 109 μ anterior to anus; vagina directed anteriorly, uteri parallel. Eggs 94 to 100 μ long by 36 to 40 μ wide.

Host.—*Spirobolus* sp. (milleped).

Location.—Intestine.

Type locality.—Panama (Upper Pequena River).

Specimens.—U. S. N. M. Helm. Coll. Nos. 31631 (type) and 31630 (paratypes).

Key to the species of the genus Heth

1. Subcephalic collar of female shield-like..... 2
Subcephalic collar of female not shield-like..... 3
2. Female bearing 2 pairs (tandem) of lateral spines.....*H. dimorphum*, n. sp.
Female bearing 2 pairs (tandem) of lateral, distally bifid scales
H. juli Cobb, 1898
3. Female bearing 1 group of postcephalic spines (2 median, 4 submedian)
H. hexaspinosum, n. sp.
Female bearing 3 groups (tandem) of postcephalic spines (2 in each group) 4
4. Postcephalic spines lateral.....*H. duvidosum* Artigas, 1929
Postcephalic spines median.....*H. spinosum* Artigas, 1929

Nomenclatorial notes, I. B. G. CHITWOOD, U. S. Bureau of Animal Industry.

In the process of revising the classification of nematodes several cases have been found in which nomenclatorial corrections appear to be necessary. These corrections fall in 4 groups, as follows: (1) A case of priority involving the status of *Anguina* Scopoli, 1777, *Anguillulina* Gervais & Beneden, 1859, and *Tylenchus* Bastian, 1865; (2) renaming of homonyms; (3) emendation of supergeneric names; and (4) designation of type species. These subjects will be covered under separate headings.

(1) THE STATUS OF ANGUINA SCOPOLI, 1777

It appears that *Anguina* invalidates *Anguillulina* and *Tylenchus* as generic names for the wheat eelworm. The oldest generic name for this important nematode appears to be *Anguina* Scopoli, 1777, a name which Baylis and Daubney (1926, Synopsis fam. & gen. Nematoda, p. 65) have placed as a synonym of *Anguillulina* Gervais & Beneden, 1859 (= *Tylenchus* Bastian, 1865). Goodey (1932, J. Helminth. 10:76) does not appear to regard this worm as congeneric with that described by Steinbuch (1799, Der Naturforscher 28:256) as *Vibrio triticii* (= *Anguillulina tritici*, = *Tylenchus tritici*).

Study of Scopoli's paper (1777, *Introductio ad Historiam naturalem sistens genera Lapidum, Plantarum, et Animalium*, Prague, pp. 373, 374) shows that he clearly stated that he was making a new genus; he gave host, location, an attempted description, and referred to Linnaeus.

(Quoted from Scopoli, p. 373)

Sequor hic doctissimum Virum *Othonem Fredericum* MÜLLERIVM, qui mira industria in hac Tribu ea præstitit, quæ nemo alius. Nova tamen & a Mullerianis diversa Genera ANGVINAE, & ROTARIAE, nec non FVRIAM Ill. LINN. adiungere visum est, de cuius tamen existentia dubitant etiamnum multi.

(Quoted from Scopoli, p. 374)

*5 ANGVINA. SCOP. Corpus filiforme.

Habitat in Galla inter glumas & folliculos ce-realium, directionem medullæ ita sæpe mutans, ut plura germina prodeant.

Mas. brevior, pene porrecto.

Fem. longior, vivipara & ovipara.

Neutrum mediocre, pene & vulva destitutum, FONTANA.

Tritici grana abbreviata & rotundata LINN. in *not.* ad CHAOS.

The Linnean reference (1767, *Systema Naturae*, p. 1326) also gave host, location and an attempted description.

(Quoted from Linnaeus, p. 1326)

1326 VERMES. ZOOPHYTA. Chaos.

354. CHAOS. *Corpus* liberum, uniforme, redivivum: Artubus sensusque organ-
nis externis nullis.

Ustilago. 4. C. fructificationis vegetabilis, pulveraceum. *Munch-
hausv.* I. p. 149.

Habitat in destructis granis Hordei, Tritici, Graminumque aliorum, inque *Flosculis* Tragopogonis, Scorzonæræ, forma nigri pulveris. Hic pulvis aliquot dies in aqua tepente maceratus transit in Animalcula oblonga, hyalina, piscium instar ludentia, armato oculo videnda. *Munch. confer. Dissert. nost. de Mundo invisibili.*

TRITICI Grana abbreviata illa & rotundata, exsiccata etiam post annos, in aqua tepidiuscula intra horulam egerminant in ascaridiformem quasi vermiculum; animatum vix dixero.

Scopoli and Linnaeus undoubtedly referred to the same species. There is no doubt as to what species this is because the form now known as *Tylenchus tritici* = *Anguillulina tritici* is the only species found in grains of wheat causing the formation of galls (rounded) instead of grains (oblong).

The wheat eelworm was given a specific name, *Vibrio tritici*, by Steinbuch (1799, loc. cit.), basing his species on forms described by Roffredi (1775, *Obs. Mem. Phys. Nat.* 5 (1):1-19).

The international rules of zoological nomenclature do not invalidate old genera which have been described without a specific name being mentioned. The genera *Anguillulina* Gervais & Beneden, 1859 (type *A. tritici* (Steinbuch, 1799) Gervais & Beneden, 1859), and *Tylenchus* Bastian, 1865 (type *T. davainii* Bastian, 1865) are both congeneric with the form described under *Anguina*. The former has an identical species as type, hence *Anguina* is monotypic even though no specific name was mentioned. The type can only be *Tylenchus tritici* (Steinbuch, 1799). Since *Anguillulina* Gervais & Beneden, 1859, and *Tylenchus* Bastian, 1865, are clearly antedated by *Anguina* Scopoli, 1777, according to the rules of priority they must be considered synonyms of *Anguina*.

It is difficult to understand why Baylis & Daubney (1926, loc. cit.) and Goodey (1932, loc. cit.) failed to recognize priority in this case. Either *Anguina* should be used on the basis of priority or the rules of priority should be suspended and *Tylenchus* recognized, since the latter name is best known and most widely used. A request that *Tylenchus* be placed on the official list of genera has been sent to Dr. C. W. Stiles, Secretary of the International Commission on Zoological Nomenclature, and it is hoped that an agreement can be reached to this effect.

(2) RENAMING OF HOMONYMS

Asconema Leuckart, 1886 (type *A. gibbosum* Leuckart, 1886), is a homonym of *Asconema* Schmidt, 1880, sponge, and *Atractonema* Leuckart, 1887 = *Asconema* renamed (type *A. gibbosum* (Leuckart, 1886) Leuckart, 1887) is a homonym of *Atractonema* Stein, 1878, protozoan; the new generic name *Tripius* is proposed for *Asconema* (type *Tripius gibbosus* (Leuckart, 1886) n. comb.).

Leptodera Dujardin, 1845 (type *L. flexilis* Dujardin, 1845), is a homonym of *Leptodera* Audinet-Serville, 1839, orthopteran, and *Leptoderes* Dujardin, 1845 (= *Leptodera* renamed), is a homonym of *Leptoderes* Audinet-Serville, 1839, orthopteran; the new generic name *Agfa* is proposed for *Leptodera* (type *Agfa flexilis* (Dujardin, 1845) n. comb.).

(3) EMENDATION OF SUPERGENERIC NAMES

The original name is given first, the emended form second.

Bunoneminae Mic., 1922	Bunonematinae Mic., 1922, emend.
Angiostominae Wülker, 1923	Angiostomatinae Wülker, 1923, emend.
Ransomneminae Travassos, 1929	Ransomnematinæ, Trav., 1929, emend.
Rhigonemidae Artigas, 1930	Rhigonematodæ Artigas, 1930, emend.
Rhigoneminae Artigas, 1930	Rhigonematinae Artigas, 1930, emend.
Steinerneminae Filipjev, 1934	Steinernematinae Filip., 1934, emend.
Drilonemidae Pierantoni, 1916	Drilonematidae, Pieran., 1916, emend.
Allantoneminae Pereira, 1932	Allantonematinae Periera, 1932, emend.
Avioserpensinae Wehr & Chit., 1934	Avioserpentinae, W. & C., 1934, emend.
Oxystominae Micoletzky, 1934	Oxystomininae, Mic., 1924, emend.
Eurystominae ?Filipjev, 1934	Eurystomininae ?Filipjev, 1934, emend.

(4) DESIGNATION OF TYPE SPECIES

The type designated is enclosed in parenthesis.

- Poikilolaimus* Fuchs, 1930. (*P. micoletzkyi* Fuchs, 1930).
Rhodolaimus Fuchs, 1930. (*R. pusillus* Fuchs, 1930).
Panagrolaimus Fuchs, 1930. (*P. detritophagus* Fuchs, 1930).
Rhabdoltolaimus Fuchs, 1931. (*R. corinthiacus* Fuchs, 1931).
Rhabditophanes Fuchs, 1930. (*R. brassicae* Fuchs, 1930).
Mesonema Pierantoni, 1916. (*M. acuminatum* Pierantoni, 1916).
Opistonema Pierantoni, 1916. (*O. minutum* Pierantoni, 1916).
Parasitaphelenchus Fuchs, 1930. (*P. conjunctus* Fuchs, 1930).
Chitinoaphelenchus Micoletzky, 1922. (*Aphelenchus* (C.) *ormerodis* (Ritzema Bos, 1891) Micoletzky, 1922 = *Aphelenchoides parietinus* (Bastian, 1865)).
 Subgenus, s. o. *Aphelenchoides*. Originally included species type of older genus.

- Parasitylenchus* Micoletzky, 1922. (*P. dispar* (Fuchs, 1914) Mic., 1922.)
Bathymermis Daday, 1911. (*B. fuhrmanni* Daday, 1911).
Limnomermis Daday, 1911. (*L. bathybia* Daday, 1911).
Mesomermis Daday, 1911. (*M. zschokkei* Daday, 1911).
Dichromadora Kreis, 1929. (*D. cephalata* (Steiner, 1916) Kreis, 1929).
Trichromadora Kreis, 1929. (*T. longicaudata* Kreis, 1929).
Paraeuchromadora Stekhoven & Adams, 1931. (*P. amphidiscata* Stekhoven & Adams, 1931).
Epsilonema Steiner, 1927. (*E. steineri* n.n.=*Rhabdogaster cygnoides* Metschn. of Steiner, 1916 and 1927; type not *E. cyrtum* Steiner, 1931 (not an originally included species).
Brachydesmodora, Allgen, 1932. (*Desmodora* (B.) *similis* Allgen, 1932) [subgenus].

Survival of horse strongyle eggs under anerobic conditions. JOHN T. LUCKER,
 U. S. Bureau of Animal Industry.

The object of the experiments reported in this paper was to determine the survival time of horse strongyle eggs under the anerobic conditions existing in a dilute mixture of feces and water. The data supplied by these experiments have some practical application because the conditions under which the experiments were performed are comparable to those obtaining in tanks employed for the storage of stable manure mixed with water. Peterson (1927, Hoard's Dairyman 72:888) reports that stable manure is frequently stored in liquid form by European farmers. He states that the usual period of storage is from a month to 6 weeks.

In the writer's experiments, a liquid suspension of horse feces was prepared by macerating fresh stable manure in small amounts of water. The mixture was screened to remove coarse particles, further diluted with water, and finally placed in one of the receptacles described later, where it remained unagitated. Portions of the sludge containing strongyle eggs were removed at certain intervals through a valve-controlled outlet at the bottom of the container. Eggs were isolated from a part of the sludge sample for immediate examination and for culturing in water, and the remaining sludge was mixed with charcoal and cultured in Petri dishes.

In the first experiment, the depth of liquid in the small metal tank which was used as a container for the mixture was about 24 inches. The tank was 17½ inches in diameter and the volume of the liquid mixture was 5,760 cubic inches. Sludge was withdrawn 7, 12, 15, 19, and 41 days after the tank was filled and, following each withdrawal of sludge, cultures were prepared and the eggs were immediately examined. In no case was there any progress in the development of viable eggs, and attempts to isolate larvae from freshly drawn sludge gave negative results. In all cases some eggs appeared to have been killed and failed to develop subsequently in water cultures, the proportion of non-viable eggs increasing with the duration of the anerobic exposure. However, despite submergence in the liquid mixture for 41 days, the majority of the eggs cultured in water hatched in the course of a day or two. Many infective larvae were recovered from a sludge and charcoal culture prepared 19 days after the container was filled. Infective larvae were recovered also from the culture prepared from sludge removed after storage for 41 days. However, the number of larvae recovered from this culture was considerably lower than that recovered from the earlier culture.

In the second experiment, the depth of the liquid in a glazed tile receptacle was 105 inches and the volume of the mixture was 8,245 cubic inches at the beginning of the experiment; the depth was reduced to about 65 inches and the volume to 5,000 cubic inches at the conclusion of the experiment, due to the withdrawal of sludge from time to time. Sludge was withdrawn after 18, 31, 45, 53 and 63 days. The results obtained in cultures are given below in tabular form:

TABLE 1.—*Effect of exposure to anerobic conditions on horse strongyle eggs in experiment 2.*

Duration of exposure <i>Days</i>	Results in cultures of eggs in water	Results in sludge and charcoal cultures
18	About 85 percent of the eggs hatched	Many infective larvae recovered
31	About 75 percent of the eggs hatched	Several infective larvae recovered
45	About 2.5 percent of the eggs hatched	7 infective larvae recovered
53	Of several hundred eggs present, 4 hatched	5 infective larvae recovered
63	Of 24,600 eggs present, 4 hatched	No infective larvae recovered

An examination of the data given in table 1 shows that horse strongyle eggs submerged for 63 days in a dilute mixture of feces and water at depths varying from about 5½ to 8 feet failed to yield infective larvae when cultured in a medium of charcoal and feces, although 0.016 percent of the eggs isolated from another portion of the same sludge sample hatched subsequently in water culture. Two of the four eggs which hatched in the latter medium yielded infective larvae. A few infective larvae were found in solid cultures prepared from eggs submerged at these depths for 45 and 53 days, respectively. Following submergence for 41 days at a depth of 24 inches, the proportion of eggs which hatched in a water culture and the number of larvae found in a solid culture medium were significantly greater than in the case involving submergence of the eggs for 45 days at greater depths. After submergence for 1 month (experiment 2) a majority of eggs were still viable and considerable numbers of larvae were recovered in a solid culture medium.

Both experiments reported in this paper and the somewhat similar experiments reported by Gackstatter (1932, *Ztschr. Veterinärk.* 44: 334-339) indicate that horse strongyle eggs are completely arrested in their development when they are submerged in a dilute mixture of feces and water. However, the progressive destruction of the vitality of submerged eggs, shown particularly by the writer's second experiment, is not in agreement with Gackstatter's statement to the effect that all of the originally viable strongyle eggs in partially or completely submerged mule feces retained the capability of further development despite maintenance under these conditions for from 4 to 5 weeks. The greater depths to which eggs were submerged in the writer's experiments may account for this difference in results. It is interesting to note that Zavadovskii (1916, *Compt. rend. Soc. Biol.* 79: 595-598) found that eggs of *Ascaris equorum* were unaffected by maintenance under anerobic conditions for from 4 to 5 months.

In the experiments described in this paper, eggs were subjected to anerobic conditions, to the influences of products of the putrefaction and decomposition of the medium, and to other influences possibly unfavorable to survival and development. While no attempt was made to determine the individual effects of these various factors on the eggs, the writer assumes that the lack of oxygen in the dilute fecal mixture was largely responsible for the failure of the eggs to develop while in this medium and for the ultimate destruction of their vitality. This assumption is supported by data published by Looss in 1896 (*Centbl. Bakt. [etc.]*, I Abt. 20: 865-870) and 1911 (*Rec. School Med., Min. Ed., Egypt* 4: 163-613), by Lambinet in 1903 (*Bull. Acad. Roy. Méd. Belg.* 4. s. 17: 534-544), by Bruns in 1904 (*Klin. Jahrb.* 12: 1-28), by Zavadovskii in 1916 (*Compt. rend. Soc. Biol.* 79: 798-802) and 1927 (*Arch. Entwicklmechl. Organ.* 109: 14-23), by Zavadovskii and Orloff in 1927 (*Arch. Entwicklmechl. Organ.* 109: 750-769), by Minagawa in 1920 (*Mitt. Med. Fakult. Kais. Univ. Tokyo* 23: 149-170; *ibidem.* 23: 171-187), by Kosmin in 1928 (*Tr. Lab. Exp. Biol. Zoopark Moscow* 4: 207-218), by McCoy in 1930 (*Am. J. Hyg.* 11: 413-448), and by Nolf in 1932 (*Am. J. Hyg.* 16: 288-322).

Observations on the life history of *Toxascaris leonina* (Nematoda: Ascaridae). WILLARD H. WRIGHT, U. S. Bureau of Animal Industry.

The prevailing conception that the life history of *Toxascaris leonina* is identical with that of *Ascaris lumbricoides* and *Toxocara canis* is apparently based on the observations of Fülleborn. Actually, Fülleborn (1922, Arch. Schiffsu. Tropen-Hyg. 26: 59-60) expressed considerable doubt concerning this point and stated that in his opinion it was entirely possible that *T. leonina* had another mode of development. Feeding experiments carried out by the writer with albino rats, mice, guinea-pigs and dogs indicate that the larvae of *T. leonina* do not regularly migrate throughout the body of the host.

There are 4 larval stages in the life history of this parasite. Ova of *T. leonina*, maintained at a temperature of 30° C. in an atmosphere saturated with moisture, contain vermiform embryos, or 1st-stage larvae, on the 2nd day of incubation, and ensheathed, infective embryos, or 2nd-stage larvae, on the 3rd day of incubation. Unlike the ova of *T. canis*, ova of *T. leonina* are able to develop to infective, 2nd-stage larvae, at a temperature of 37° C., in an atmosphere saturated with moisture.

Evidence indicates that when ova containing 2nd-stage larvae are ingested by the host, the ova usually hatch in the duodenum. The first ecdysis occurs at the time of hatching or shortly thereafter. The liberated larvae then penetrate into the wall of the small intestine, particularly that of the lower part of the duodenum, and come to rest in the crypts of Lieberkühn, in the submucosa and in the circular muscle of the wall. The larvae undergo considerable growth while in the intestinal wall and begin to emerge about the 9th or 10th day after infection. It is probable that the 2nd ecdysis occurs on emergence or shortly before emergence of the larvae, as we have recovered exsheathed 3rd-stage larvae from the mucosa of the duodenum on the 10th day after infection. The genital primordium of 3rd-stage larvae recovered from the lumen of the intestine of a dog 10 days after infection contained 8 cells, indicating that the larvae were in the early 3rd-stage. Larvae at this stage were 0.669 to 0.795 mm long. Later 3rd-stage larvae were recovered from the lumen of the small intestine of a dog killed 14 days after infection. These larvae were 0.794 to 0.890 mm in length, and the genital primordium contained 16 cells. Following their reentrance into the lumen of the small intestine, the larvae grow rapidly and by the 18th day after infection some of them have undergone the 3rd ecdysis. The 4th-stage larva later molts to become an adult.

Occasionally, particularly in very heavy infestations, some larvae will penetrate the wall of the small intestine and develop in the abdominal cavity, or reach such organs as the mesenteric lymph nodes, pancreas, liver and lungs. Larvae in these locations are capable of developing into 3rd-stage larvae, as larvae recovered from these organs have been in the same stage of development as those recovered from the lumen of the small intestine in the same length of time after infection. Larvae which reach the lungs are capable of finding their way into the trachea, where they are carried upward, are swallowed and again reach the small intestine, and such larvae probably develop to maturity.

A note on the identity of *Nematodirus tarandi* Hadwen, 1922, and *Nematodirus skrjabini* Mitzkewitsch, 1929 (Nematoda: Trichostrongylidae). G. DIKMANS, U. S. Bureau of Animal Industry.

On comparing the published figures of *Nematodirus skrjabini* described as a new species by Mitzkewitsch in 1929 from the reindeer with specimens of *Nematodirus tarandi* Hadwen, 1922, in the National Museum Helminthological Collection, I find that these 2 nematodes are identical. *Nematodirus skrjabini* Mitzkewitsch, 1929, therefore, becomes a synonym of *Nematodirus tarandi* Hadwen, 1922.

An unidentified nematode from the eye of the moose, *Alces americana*.

G. DIKMANS and E. E. WEHR, U. S. Bureau of Animal Industry.

A short time ago Dr. C. P. Fitch, Chief of the Veterinary Division, College of Agriculture, University of Minnesota, sent to the U. S. Bureau of Animal Industry 2 nematodes, collected by Dr. Fenstermacher from the eyes of moose in northern Minnesota. The worms submitted for examination were immature females, about 40 mm long and threadlike. The head is without buccal capsule or circumoral elevation, and the genital opening is in close proximity to the anal opening in the posterior part of the body.

As far as we are able to ascertain only 2 genera of nematodes have been reported as parasites of the eye of ruminants, viz., *Setaria* and *Thelazia*. The worms from the eye of the moose do not belong in either of these genera because in *Setaria* the mouth is surrounded by a chitinous peribuccal ring, and in *Thelazia* there is a small buccal capsule, and in both *Setaria* and *Thelazia* the vulva is located in the esophageal region. The structure of the head of the nematode collected from the eye of the moose resembles that of members of the *Metastrongylidae*, and the appearance of the tail end is similar to that of members of the *Protostrongylinae*. There is one genus in this subfamily, viz., *Elaphostrongylus* which appears to be a tissue parasite. This genus has now 2 species, *Elaphostrongylus cervi*, reported by Cameron from the connective tissues between the latissimus dorsi and the external intercostal muscles of the Scottish red deer, *Cervus elaphus*; and *Elaphostrongylus odocoilei* reported by the Hobmaiers from the connective tissues of the black-tailed deer, *Odocoileus columbianus*. Since these worms are tissue parasites they probably do considerable wandering before reaching their final destination. It is possible that the worms recovered from the eye of the moose belong in this genus. An interesting point in connection with *Elaphostrongylus odocoilei* is that this nematode is described as a lungworm, the authors claiming that eggs and larvae were found in nodules in the lungs, and adults were embedded in the connective tissues of the vessels beneath the spine and their nearby branches, in the muscles surrounding the body cavity and along the vessels of the upper part of the hind legs. The authors further state that ordinarily the worms were coiled up in the lymphatic spaces in the connective tissues and that some specimens were found penetrating the walls of larger veins of the hind legs. Accumulations of eggs were found in these places and in lymphatic vessels.

Nematode parasites of the Saiga antelope, *Saiga tatarica*. G. DIKMANS, U. S. Bureau of Animal Industry.

The following nematodes have been collected from the viscera of a Saiga antelope submitted to the U. S. Bureau of Animal Industry for examination on November 20, 1934: *Cameloststrongylus mentulatus* (*Ostertagia mentulata*), *Haemonchus contortus*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Nematodirus* sp., and *Trichuris* sp. The host catalogue of the Zoological Division contains no record of any nematodes having been collected from this animal. According to authorities of the National Museum, the Saiga antelope is becoming a rare animal. It formerly ranged the eastern part of Russia and the western part of Siberia; its present range appears to be much smaller. The animal from which the above mentioned nematodes were obtained was 1 of a shipment of 3 imported from Russia for the National Zoological Park. According to a letter received from the importers, these antelopes had been caught in the Kalmuck Steppe in South Russia, shipped to the Zoological Park in Moscow, and then transferred to a Zoological Park in Hannover, Germany. They remained in Hannover 3 months. The nematode infection may, therefore, have been acquired, at least in part, since their capture in Russia.

A note on *Nematodirella* (Nematoda: Trichostrongylidae) from three different hosts. G. DIKMANS, U. S. Bureau of Animal Industry.

The genus *Nematodirella* Yorke and Maplestone, 1926, was first reported from the reindeer, *Rangifer tarandus*, by Romanovitch (1915) under the name of *Microcephalus longissime spiculatus*. It has since been collected from the moose, *Alces americana*, and from the antelope, *Antilocapra americana*. It is a trichostrongylid characterized by the presence of very long, filiform spicules in the male, and by the atrophy of the anterior portion of the genital apparatus in the female. The figure on page 144 of Mitzkewitsch's paper on the trichostrongylid fauna of the reindeer, purporting to illustrate the ovejectors of *Nematodirella longispiculata*, is undoubtedly an error. This figure shows anterior and posterior ovejectors both functioning normally, whereas in *Nematodirella* the anterior ovarian complex is always sterile. It is probable that the ovejectors of *Nematodirus tarandi* Hadwen, 1922 (Syn. *Nematodirus skrjabini* Mitzkewitsch, 1929), were unintentionally substituted for those of *Nematodirella longispiculata*.

Some time ago Dr. W. E. Swales of the Institute of Parasitology, MacDonald College, McGill University, Quebec, Canada, called my attention to the difference in length of the spicules in specimens of *Nematodirella* collected from the moose and from the prong-horned antelope. I have, therefore, compared specimens of *Nematodirella* collected from 3 different hosts, viz., reindeer, moose and prong-horned antelope, as to the length of spicules, length of body of male, length of bursa and length of posterior ovejector in female with the following results:

TABLE 1.—Size variations in *Nematodirella longispiculata* from different hosts.

Host	Length of male	Length of spicules	Length of bursa	Length of posterior ovejector
<i>Rangifer tarandus</i>	13.5-18 mm	8.5-10.5 mm	200-225 μ	6-8 mm
<i>Alces americana</i>	10-16.5 mm	6.25-8 mm	140-150 μ	4-6 mm
<i>Antilocapra americana</i>	13-16 mm	4-5 mm	140-150 μ	3-4 mm

While the measurements here given, with the exception of those of the spicules, may not be entirely accurate, because not all the material was fixed in the same manner, and the material available for examination from both the moose and the antelope was in a very poor state of preservation, except for a few specimens collected from the moose by Dr. Swales, yet I believe them sufficiently accurate for the purpose of comparison.

On the basis of spicule length, specimens of *Nematodirella* from these 3 different hosts appear to fall in 3 groups, with the maximum length of the spicules of those from antelope approaching the minimum length of spicules of those from moose, and the maximum length of spicules of those from moose approaching the minimum length of spicules of those from reindeer. The bursa of the specimens collected from




FIG. 9. *Nematodirella longispiculata* longispiculata, Posterior ovejector.

the reindeer is somewhat larger than the bursa of the specimens collected from the moose and antelope, which are alike. As far as the length of the posterior, functional ovejector is concerned, we have a situation similar to that obtaining for the spicules. Morphologically there appears to be no difference in the structures mentioned except for size. I am, therefore, not inclined to give these nematodes different specific names but I do believe that the differences obtaining among them should be recognized by the addition of subspecific names as follows: For the form from the reindeer, *Nematodirella longispiculata longispiculata*; for the form from the moose, *Nematodirella longispiculata alcidis*; for the form from the antelope, *Nematodirella longispiculata antilocaprae*.

Lungworms collected from deer, *Odocoileus virginianus*, in Michigan. G. DIKMANS, U. S. Bureau of Animal Industry.

A little over a year ago the U. S. Bureau of Animal Industry began to receive correspondence from the Conservation Department of Michigan concerning the prevalence and identity of lungworms infesting deer in the upper peninsula of Michigan. At first only material containing eggs and larvae was received for examination. The larvae resembled the larvae of members of the Protostrongylinae, but it was, of course, impossible to make a specific determination on the basis of these larvae. Later, the posterior portion of a male and an entire female were received for study. These were identified as a species of *Protostrongylus*. Shortly thereafter additional specimens were received and identified as *Dictyocaulus filaria*. Next 3 sets of lungs from deer were received and examined. From these lungs was collected a nematode which has been placed in the genus *Pneumostrongylus*. Later the Zoological Division received from the University of Michigan through the Biological Survey some nematodes collected from the lungs of deer and these were identified as *Dictyocaulus viviparus*. Therefore, at the present time, there have been collected from the lungs of deer in Michigan 4 nematodes representing 3 genera, viz., *Protostrongylus*, *Dictyocaulus* and *Pneumostrongylus*.

New avian and insect hosts for *Gongylonema ingluvicola* (Nematoda: Spiruridae). ELOISE B. CRAM, U. S. Bureau of Animal Industry.

In the undilated esophagus of a captive mountain quail (*Oreortyx picta*), which died soon after it had been shipped from Oregon to Frederick, Md., there were found 2 female and 1 male specimens of *Gongylonema ingluvicola*. Embryonated eggs from the uteri of the worms were fed to 2 laboratory-reared cockroaches (*Blattella germanica*); 34 days later the cockroaches were killed and contained larvae tightly coiled in cysts in the body cavity. The cysts were slightly oval, measuring about 670 by 545 μ , opaque, thick-walled; the coil of the body showed no reversal of direction, during its course, as with some other avian spirurids. The 3rd-stage larvae of *G. ingluvicola* recovered from the cysts measured 1.67 mm long by 75 μ in greatest width; 2 very flat lateral lips gave appearance of an annular thickening around mouth, followed by shallow vestibule about 6 μ deep. Total length of pharynx and esophagus, 1.17 mm, thus somewhat over two-thirds total body length; excretory pore, 175 μ from head end. Tail short and blunt, anus 88 μ from end; 2 or 3 inconspicuous conical papillae at tip of tail. These measurements show only moderate variations from those of *G. scutatatum*, after development of the larvae for 7 weeks in the cockroach, as given by Ransom and Hall, as follows: Cysts 0.5 mm in diameter; larvae 1.9 mm long by 60 μ wide; total length of pharynx and esophagus 1.235 mm; excretory pore 210 μ from head end; anus 90 μ from tail end. Five larvae of *G. ingluvicola* were fed to a chicken but without subsequent development, no worms being found when the chicken was killed 79 days later.

Additional notes on two bat parasites, *Dicrocoelium lasiuri* McIntosh, 1933 (Trematoda: Dicrocoeliidae) and *Litomosa americana* McIntosh, 1932 (Nematoda: Filariidae). ALLEN MCINTOSH, U. S. Bureau of Animal Industry and GERTRUDE E. MCINTOSH.

Two parasites previously reported only in abstract form (McIntosh, 1933, J. Parasitol. 20: 137-138; 1932, J. Parasitol. 19: 170), are described here in greater detail.

Dicrocoelium lasiuri McIntosh, 1933

Description.—Body lancet-shaped, 2.47 mm long by 650μ wide, anterior sucker 160μ and acetabulum 170μ in diameter, respectively, the latter about 570μ from anterior end. Pharynx 62 by 60μ , apparently preceded by short prepharynx and followed by slightly longer esophagus. Ceca slightly undulating, extending about $2/3$ length of body; lemon yellow cuboidal crystals of some substance noted in ceca of living specimens, which did not dissolve in processes of fixation and mounting. Excretory bladder Y-shaped, with a very long stem extending from level of cecal tips to excretory pore. Testes 140 to 190μ in diameter, subtriangular, lobed or irregular in outline, situated somewhat obliquely. Vasa efferentia uniting near level of anterior margin of acetabulum, forming short vas deferens and passing immediately into cirrus sac; each vas efferens slightly expanded near its middle and filled with spermatozoa. Cirrus sac 210 by 125μ , between acetabulum and intestinal fork, containing coiled seminal vesicle. Genital pore at fork of intestine, 340μ from anterior end of body. Ovary 200 to 210μ in diameter, somewhat irregular in outline, immediately caudal to posterior testis. Laurer's canal and Mehlis' gland present. Vitellaria in zone occupied by postovarian portion of ceca, arranged in 2 lateral groups of approximately 20 follicles each. Uterus filling posterior portion of body and extending anteriorly in median field to genital pore. Metraterm to right of cirrus sac. Eggs 38 by 22μ .

Type specimen.—U.S.N.M. Helm. Coll. No. 40241; paratypes No. 40242.

Habitat.—Gall bladder and biliary tubules of liver of *Lasiurus borealis*, Washington, D. C.

Remarks.—This parasite was described from 15 specimens collected September 15, 1933, from a bat found at the base of the Washington Monument. Only one bat of several specimens, representing three species, was infected. *Dicrocoelium lasiuri* differs from *D. rileyi* Macy, 1931, a closely related form, taken from specimens of a western bat captured in Oklahoma and Kansas, in that the ovary of *D. lasiuri* is as large as, or larger than, the testes, while the ovary of *D. rileyi* is much smaller than the testes; *D. lasiuri* has twice as many vitelline follicles as *D. rileyi*. In Macy's key (1931, J. Parasitol. 18: 28-33), *D. lasiuri* will key out with *D. panduriforme* Raillet, 1900. The latter, however, is a bird parasite having a huge acetabulum. In addition to comparing our material with Macy's type specimen, several individuals of *D. rileyi*, collected in Texas and in Dr. P. D. Harwood's collection, were also examined. The differences between *D. rileyi* and *D. lasiuri* noted above were fairly constant in all specimens examined.

In addition to *Dicrocoelium rileyi* and *D. lasiuri*, two other species belong-



FIG. 10

Dicrocoelium lasiuri McIntosh, 1933, ventral aspect.

ing to the family Dicrocoeliidae have been reported from chiropterans, *Platynosomum philippinorum* Tubangui, 1928 (Philippine J. Sci. 36: 367-368), from a Philippine bat, and *Dicrocoelium hospes* Looss, 1907, of Northup, 1928 (J. Burma Research Soc. 18: 80-97), from two species of bats in Burma. From the few words of description given by Northup it is evident that the species is not *D. hospes*, but her description is too inadequate to distinguish it from the other species described from bats.

Litomosa americana McIntosh, 1932

Description.—Male 12 mm long by 110μ wide, attenuated posteriorly. Head bluntly rounded; oral opening oval, laterally elongated, bounded by an oral membrane; buccal cavity 28μ long by 20μ in diameter, conical, with apex directed posteriorly. Cephalic papillae of external circle consisting of medium-sized dome-shaped laterodorsals and lateroventrals, and of small dorsodorsals and ventroventrals; internal circle represented by 2 small internolaterals. Amphids lateral, slightly more posterior than laterodorsals and lateroventrals. Esophagus cylindrical, 300 to 350μ long by 40 to 50μ wide; nerve ring and excretory pore in region of 2nd fourth of esophagus. Spicules unequal; left 260μ long by 8μ wide, ending distally in a long lash, right 90μ long by 7μ wide, terminating in short bipartite process. Posterior part of body with 2 to 3 spiral coils; anus 95μ from tip of tail; tail tapering gradually, rounded at tip.

Female head, buccal cavity, and tail similar to those of male. Vulva near level of nerve ring, 150μ from anterior extremity; ovejector long. Sheathed microfilariae in blood, 80 to 90μ long by 5 to 7.5μ wide.

Type specimen.—U. S. N. M. Helm. Coll. No. 34160 (male); paratype No. 40243 (2 males and 1 female).

Habitat.—Abdominal cavity of *Nycticeius humeralis*, Washington, D. C.

Remarks.—The above filariid was described from 4 males and 1 female collected by the writers from 2 specimens of bats, *Nycticeius humeralis*, taken September 28, 1932, at the base of the Washington Monument. The female




FIG. 11. *Litomosa americana* McIntosh, 1932.

A—Male, esophageal region. B—Head, en face view showing oral membrane, cephalic papillae and amphids. C—Female tail. D—Male, complete worm. E—Microfilaria, sheathed, from blood of bat. F—Microfilaria, from uterus of female worm. G—Female, esophageal region showing position of vulva and ovejector. H—Male, tail showing spicules.

and a male were somewhat mutilated on opening the abdominal cavity of the host, and because of the damage to the female it was impossible to make as thorough a study of this sex as might be desired. The host is apparently a rare visitor to the District of Columbia, since in more than 2 years of collecting no additional specimens of *Nycticeius humeralis* have been captured. Mr. Gerrit S. Miller, Jr., of the U. S. National Museum, who kindly identified the hosts for the 2 parasites reported in this paper, stated at the time of identification that very few specimens of *Nycticeius humeralis* had been received from the District of Columbia. Two other species of bats, *Lasiurus borealis* and *Eptesicus fuscus*, taken from the same location did not harbor filariids.

Genus *Litomosa* Yorke and Maplestone, 1926

The type species of this genus is *Litomosa filaria* (v. Beneden, 1873) Yorke and Maplestone, 1926, which was described from a European species of bat. A second species, *L. chiropterorum* Ortlepp, 1932, was described from South African bats captured in caves in Pretoria. *L. americana* McIntosh, 1932, was the third species to be placed in the genus *Litomosa*. A fourth species, *L. witei* Krepkogorskaja, 1933, emend. (= *L. wite* Krepk., 1933) was described from a rodent from Kasakstan.

The four species may be separated with the aid of the following key:

1. Buccal capsule flask-shaped; male tail without terminal spikes; female tail with only 2 terminal processes.....*L. chiropterorum*
Not with above combination of characters..... 2
2. Buccal cavity conical with apex directed posteriorly; tails of male and female without terminal processes.....*L. americana*
Buccal cavity infundibuliform with apex directed anteriorly; tail with terminal spikes or processes..... 3
3. Vulva at level of posterior end of esophagus; female tail with 1 pair of small divergent terminal processes between which are 2 minute spines; male tail with subterminal spike.....*L. filaria*
Vulva 140 to 180 μ posterior to level of posterior end of esophagus; female tail terminating in 2 lateral processes between which is a somewhat larger dorsal process, the intervening space between the processes covered with a number of fine tubercules; male unknown.....*L. witei*

Tropical rat mite attacking man in St. Louis area. ALLEN MCINTOSH, U. S. Bureau of Animal Industry.

Several mites were received December 10, 1934, from Dr. C. F. Payne, inspector in the Meat Inspection Division of the U. S. Bureau of Animal Industry, with the following note:

"These parasites were obtained from employees in the office of the _____ Commission Company, National Stock Yards, Illinois, from clothing and from arms and legs. Employees of this firm have been complaining for the past month of a condition somewhat resembling harvest or chigger mite bites. The employees' arms and legs appear as if they had been subjected to severe chigger or mosquito bites. Similar conditions are being reported from dressing rooms in the public stock yards and around scale houses. One case was reported from the _____ Packing Company, where a watchman was complaining of a similar condition to the employees in the _____ Commission Company's office.

"The employees in the _____ Commission Company state that they find parasites in the desk drawers, on their clothing, arms and legs, and in linen closet. The ones forwarded in the bottle were obtained from the linen closet and it appears as if they have been engorged with blood from a host other than human beings, possibly rats.

"As the season of the year appears past for the prevalence of the chigger mite, and with the possibility that these parasites may be the carrier of some diseased condition, it is thought that the existence of these parasites here might be of interest to the public health service."

The mites were identified by the writer as *Liponyssus bacoti* Hirst, 1914. This mite has recently been shown experimentally by Dove and Shel mire (1932,

J. Parasitol. 18: 159-168) to transmit endemic typhus. In the above reference Dove and Shelmire pointed out that from their reports this mite probably occurs in St. Louis, Missouri. This note is given here to establish definitely the fact that the tropical rat mite does occur in the St. Louis area.

A new trematode, *Lyperosomum monenteron*, n. sp. (Dicrocoeliidae), from a robin. E. W. PRICE and ALLEN MCINTOSH, U. S. Bureau of Animal Industry.

About three years ago, Mr. Alex. D. Baker, MacDonald College, Quebec, forwarded to the U. S. Bureau of Animal Industry a number of trematodes for identification. Among these were several specimens from a robin, *Turdus migratorius*, which appeared to represent a new species belonging to the genus *Lyperosomum* Looss. Subsequently 3 additional lots of specimens have been secured, one from Mr. L. M. Dickerson collected from a robin at Charlottesville, Va., another collected by one of us (A. M.) from the same host at Washington, D. C., and a third from a bluebird found dead at Falls Church, Va.

Lyperosomum monenteron, n. sp.

Description.—Body lanceate, 1.9 to 5.2 mm long by 630 to 670 μ wide at level of acetabulum, narrowing more or less abruptly in preacetabular region, and with short, lip-like projection dorsal to oral sucker; cuticula smooth. Oral sucker subterminal, 120 to 170 μ long by 123 to 150 μ wide; acetabulum 170 to 320 μ long by 200 to 320 μ wide, about 1/5 of body length from anterior end. Pharynx subglobular, 42 to 60 μ long by 50 to 70 μ wide, its anterior end in contact with oral sucker; esophagus slender; intestine single, extending between testes and between posterior testis and ovary, then continuing in more or less wide undulations (undulations wider, as a rule, than shown in illustration) and terminating blindly about 1/3 of body length from posterior end. Excretory aperture terminal; excretory vesicle tubular, extending anterior to about level of anterior limits of vitellaria, then bifurcating and forming short lateral limbs; remainder of excretory system not observed. Genital aperture median, about midway between anterior margin of acetabulum and tip of body, its position varying somewhat, depending on amount of extension or contraction of preacetabular portion of body. Cirrus pouch elongate piriform to pestle-shaped, 160 to 320 μ long by 65 to 100 μ wide, containing a relatively long, folded seminal vesicle, a small pars prostatica, and cirrus. Testes transversely oval, 140 to 260 μ long by 250 to 460 μ wide, one diagonally in front of other in posterior part of anterior third of body. Ovary transversely oval, 90 to 125 μ long by 100 to 220 μ wide, posttesticular and to right of median line. Seminal receptacle globular or piriform, median, at level of posterior margin of ovary; Mehlis' gland moderately developed, postovarial; Laurer's canal long and slender, opening dorsal to ovary. Vitellaria in equatorial zone, consisting of relatively large follicles tending to meet anteriorly in median line. Uterus with greatly convoluted descending and ascending limbs, filling greater part of postovarial portion of body. Eggs 32 μ long by 16 μ wide.

Host.—*Turdus migratorius* and *Sialia sialis*.

Location.—Gall bladder and bile ducts.

Distribution.—United States and Canada.

Specimens.—U. S. N. M. Helm. Coll. Nos. 40237 (type), 40238 (paratypes), 40239, 40240 and 34337.

This species differs from other species of the genus in having a single intestinal cecum. It is not improbable that some of the other species that have been assigned to the




FIG. 12. *Lyperosomum monenteron*, n. sp., dorsal view.

genus *Lyperosomum* by various workers may show this feature, but it appears not to have been mentioned. Some of the descriptions and illustrations of species indicate that the specimens studied were in such poor condition that it was possible to make out only the more obvious characters. Such descriptions and figures make it virtually impossible to determine the validity of many of the species.

***Mesocoelium danforthi*, n. sp. (Dicrocoeliidae), from a lizard, *Celestus pleii*, in Puerto Rico.** W. A. HOFFMAN, School of Tropical Medicine, San Juan, P. R.

This work was done in the laboratory of the Zoological Division of the U. S. Bureau of Animal Industry, Washington, D. C. The writer is indebted to Dr. E. W. Price for aid and counsel rendered during the course of this study.

Mesocoelium danforthi, n. sp.

Description.—Body elongate oval, 1.31 to 1.65 mm long by 0.54 to 0.58 mm wide, posterior end slightly more rounded than anterior; cuticula apparently without spines. Oral sucker subterminal, slightly longer than wide, 0.296 to 0.300 mm wide by 0.328 to 0.336 mm long, covered by a membrane with a longitudinal median slit. Acetabulum nearly circular, 0.148 to 0.172 mm in diameter; longitudinal diameter about half that of oral sucker, its anterior margin situated near junction of anterior and median thirds of body length. Prepharynx present, indistinct; pharynx semiglobular, about 0.110 by 0.088 mm. In all specimens examined the esophagus is so short as to appear nonexistent. Intestinal caeca extending horizontally, curving at an angle of almost 90 degrees, and then running parallel to lateral margins, barely attaining posterior third of body. Only posterior portion of excretory vesicle visible. Genital pore at level of base of pharynx; cirrus pouch narrowly piriform, about 0.146 mm long, its base beneath anterior margin of acetabulum. Testes broadly ovoid, 0.104 to 0.108 by 0.072 to 0.076 mm, situated obliquely on either side of acetabulum, one slightly anterior and the other somewhat posterior to it. Ovary subspherical, 0.100 to 0.136 mm by 0.08 to 0.096 mm, immediately behind and sometimes contiguous with right testis. Vitellaria extending from level of oral sucker into proximal portion of posterior third of body, terminating just beyond caeca. Seminal receptacle, yolk reservoir and transverse vitelline ducts not discernible, obscured by the numerous ova filling posterior half of body.

Habitat.—Intestine of a lizard, *Celestus pleii*, collected by Dr. S. T. Danforth at El Yunque, a mountain in the Luquillo National Forest, Puerto Rico, March 29, 1934; captured at altitude of 2000 ft.

Specimen.—U. S. N. M. Helm. Coll. Nos. 39570 (type) and 39571 (paratypes).

Mesocoelium danforthi is closely allied to *M. sociale* (Lühe, 1901), *microon* Nicoll, 1914, *americanum* Harwood, 1932, *meggitti* Bhalero, 1927, and *monodi* Dollfus, 1929. From all these, *danforthi* differs in having the longitudinal diameter of the oral sucker approximately twice that of the acetabulum, in the small size of its eggs, and with possible exception of *americanum*, in its ovary being larger than either testis. The cirrus pouch of *americanum* does not extend to the acetabulum, as it does in the case of *danforthi*.




FIG. 13. *Mesocoelium danforthi*, n. sp., ventral view.

A note on the life history of *Telorchis robustus* Goldb. (Trematoda: Telorchhiidae). WENDELL H. KRULL, U. S. Bureau of Animal Industry.

Experiments have been conducted to determine the life history of *Telorchis robustus*, a trematode parasite of the common land turtle, *Terrapene carolina*. The turtle from which the material used in these experiments was obtained was collected and the flukes determined by Mr. Allen McIntosh of the Zoological Division.

Eggs of the parasite were obtained from mature specimens of the fluke and fed to laboratory-raised snails, *Pseudosuccinea columella*. The snails became infected and cercariae were shed by the first snail 28 days after exposure to infection, and all snails were shedding cercariae 4 days later.

The cercaria, a xiphidiocercaria, is spined anteriorly, and has 4 penetration glands on one side and 5 on the other. The excretory bladder is Y-shaped, the main excretory tubes entering the cornua near their posterior ends. A pocket or vestibule is present at the posterior end of the body, which contains the excretory pore and in which the simple tail is fastened. The cuticula of the 2 dorsolateral portions of the cavity is greatly thickened, and each thickened area is provided with about 50 long spines which project into the cavity.

Laboratory-raised *Pseudosuccinea columella*, *Helisoma trivolvis* and *Lymnaea traskii* were exposed to the cercariae and became infested, the cercariae penetrating the exposed portions of the snails. Encysted metacercariae were subsequently found in almost any soft part of the snail; they were never abundant, even though the snails had been exposed repeatedly to thousands of cercariae.

The experiments demonstrated that *Pseudosuccinea columella* may serve as first and second intermediate hosts, and *Helisoma trivolvis* and *Lymnaea traskii* as second intermediate hosts for this fluke. Further details of the life history and complete descriptions of the larval stages will be given later.

Minutes.

One hundred sixty-fifth to one hundred sixty-seventh meetings.

The 165th meeting was held on October 20, 1934, at Beltsville, Md., on the occasion of the dedication of the new Zoological Division laboratory. Officers were elected as follows: President, L. A. Spindler; Vice President, E. E. Wehr; Recording Secretary, G. Dikmans; Corresponding Secretary-Treasurer, Miss Edna M. Buhner. No program was presented at this meeting.

The 166th meeting was held at the School of Hygiene, Baltimore, Md., on November 17, 1934. A paper was presented by Dr. Cram. Informal notes were presented by Miss Wolfson and by Messrs. Alicata, Bartsch, Andrews, Atchley, Otto, and Spindler. Dr. Cram's paper appears in this issue.

The 167th meeting was held on December 15, 1934. Papers were presented by Mrs. Chitwood and Mrs. Cobb and by Messrs. Chitwood, Dikmans, Krull, Luckner, McIntosh, Price, Steiner, and Taylor. (See this issue.) The financial report was presented by Mr. E. E. Wehr. Dr. Christie called attention to the fact that "A key to the genera of free-living nemas" which appears in this issue, is the 27th and concluding number of "Contributions to a science of nematology" by N. A. Cobb. Any one wishing to purchase a complete set of this series should communicate with Miss Margaret V. Cobb, Tilton, N. H.

G. DIKMANS, *Secretary.*

CONTENTS

	PAGE
CHITWOOD, B. G. A new nematode, <i>Camacolaimus prytherchi</i> , n. sp. (Camacolaimidae)	49
—————. Nomenclatorial notes, I.	51
CHITWOOD, M. B. Two new nematodes of the genus <i>Heth</i> Cobb, 1898 (Atractidae)	50
COBB, N. A. A key to the genera of free-living nemas.....	1
CRAM, ELOISE B. New avian and insect hosts for <i>Gongylonema ingluvicola</i> (Nematoda: Spiruridae)	59
DIKMANS, G. A note on the identity of <i>Nematodirus tarandi</i> Hadwen, 1922, and <i>Nematodirus skrjabini</i> Mitzkewitsch, 1929 (Nematoda: Trichostrongylidae)	56
—————. Nematode parasites of the Saiga antelope, <i>Saiga tatarica</i> ...	57
—————. A note on <i>Nematodirella</i> (Nematoda: Trichostrongylidae) from three different hosts	58
—————. Lungworms collected from deer, <i>Odocoileus virginianus</i> , in Michigan	59
————— and WEHR, E. E. An unidentified nematode from the eye of the moose, <i>Alces americana</i>	57
HOFFMAN, W. A. <i>Mesocoelium danforthi</i> , n. sp. (Dicrocoeliidae), from a lizard, <i>Celestus pleii</i> , in Puerto Rico.....	64
KRULL, WENDELL H. A note on the life history of <i>Telorchis robustus</i> Goldb. (Trematoda: Telorchidae)	65
LUCKER, JOHN T. Survival of horse strongyle eggs under anerobic conditions	54
MCINTOSH, ALLEN. Tropical rat mite attacking man in St. Louis Area.....	62
————— and MCINTOSH, GERTRUDE E. Additional notes on two bat parasites, <i>Dicrocoelium lasiuri</i> McIntosh, 1933 (Trematoda: Dicrocoeliidae) and <i>Litomosa americana</i> McIntosh, 1932 (Nematoda: Filariidae)	60
PRICE, E. W., and MCINTOSH, ALLEN. A new trematode, <i>Lyperosomum monenteron</i> , n. sp. (Dicrocoeliidae), from a robin.....	63
STEINER, G. Opuscula miscellanea nematologica, I.....	41
TAYLOR, A. W. A review of the fossil nematodes.....	47
THORNE, GERALD. Notes on free-living and plant-parasitic nematodes, I.....	46
————— and PRICE, CHARLES. The nematode <i>Neotylenchus abulbosus</i> Steiner (Anguilluliniidae) as a parasite of sugar-beets.....	46
WRIGHT, WILLARD H. Observations on the life history of <i>Toxascaris leonina</i> (Nematoda: Ascaridae)	56