

Widespread infection of the Eastern red-spotted newt (*Notophthalmus viridescens*) by a new species of *Amphibiocystidium*, a genus of fungus-like mesomycetozoan parasites not previously reported in North America

T. R. RAFFEL^{1,2*}, T. BOMMARITO³, D. S. BARRY⁴, S. M. WITIAK⁵
and L. A. SHACKELTON¹

¹ Center for Infectious Disease Dynamics, Biology Department, Penn State University, University Park, PA 16802, USA

² Department of Biology, University of South Florida, Tampa, FL 33620, USA

³ Cooperative Wildlife Research Lab, Department of Zoology, Southern Illinois University, Carbondale, IL 62901, USA

⁴ Department of Biological Sciences, Marshall University, Huntington, WV 25755, USA

⁵ Department of Plant Pathology, Penn State University, University Park, PA 16802, USA

(Received 21 March 2007; revised 17 August 2007; accepted 20 August 2007; first published online 12 October 2007)

SUMMARY

Given the worldwide decline of amphibian populations due to emerging infectious diseases, it is imperative that we identify and address the causative agents. Many of the pathogens recently implicated in amphibian mortality and morbidity have been fungal or members of a poorly understood group of fungus-like protists, the mesomycetozoans. One mesomycetozoan, *Amphibiocystidium ranae*, is known to infect several European amphibian species and was associated with a recent decline of frogs in Italy. Here we present the first report of an *Amphibiocystidium* sp. in a North American amphibian, the Eastern red-spotted newt (*Notophthalmus viridescens*), and characterize it as the new species *A. viridescens* in the order Dermocystida based on morphological, geographical and phylogenetic evidence. We also describe the widespread and seasonal distribution of this parasite in red-spotted newt populations and provide evidence of mortality due to infection.

Key words: Dermocystida, *Dermocystidium*, *Amphibiothecum*, amphibian decline, salamander, fungal infection.

INTRODUCTION

Emerging diseases are of increasing concern for both humans and wildlife, and determining the identity of their causative agents and potential impacts on their hosts will be important for developing control measures (Daszak *et al.* 2001). This is especially true for amphibians, which are declining precipitously worldwide due, in large part, to emerging diseases. The pathogens and parasites most commonly implicated in amphibian declines and mortality events include the chytrid fungus (*Batrachochytrium dendrobatidis*), ranaviruses, *Saprolegnia* spp. fungi, and *Ribeiroia* spp. trematodes, which cause limb deformities (Green *et al.* 2002; Jancovich *et al.* 2005; Johnson and Lunde, 2005; Lips *et al.* 2006). However, mortality and morbidity events in North America have also been attributed to *Ichthyophonus* sp., *Amphibiothecum* (*Dermosporidium*) spp., and a *Perkinsus*-like organism, all of which are fungus-like mesomycetozoan organisms (Jay and Pohley, 1981;

Green and Sherman, 2001; Green *et al.* 2002, 2003; Feldman *et al.* 2005; Raffel *et al.* 2007).

Outbreaks of another mesomycetozoan, *Amphibiocystidium ranae*, have been associated with declining populations of *Rana lessonae* in Italy, although the role of the parasite in these declines is still undetermined (Pascolini *et al.* 2003). *Amphibiocystidium* spp. have been reported for over a century in frogs and salamanders within Europe, where the parasite has been found in adults of 3 types of anurans (*Rana temporaria*, *Rana esculenta*, and *Alytes obstetricians*) and 2 newts (*Triturus marmoratus* and *Triturus cristatus*) (Pascolini *et al.* 2003). *Amphibiocystidium* spp. infections have not previously been reported in New World amphibians (Pascolini *et al.* 2003), although Carini (1940) described a similar infection, *Dermosporidium hyalarum*, in Brazilian frogs. Green *et al.* (2002) reported a *Dermocystidium*-like organism in North American ranid tadpole livers, but this was later shown to be more closely related to *Perkinsus* spp. parasites of marine shellfish (Green *et al.* 2003). Another similar parasite, *Amphibiothecum penneri*, was described in American toads but is now considered to be outside the genus *Amphibiocystidium* based on morphological and phylogenetic evidence

* Corresponding author: 4202 E. Fowler Avenue, SCA 110, Tampa, Florida 33620, USA. Tel: +813 974 6210. Fax: +813 974 3263. E-mail: traffel@cas.usf.edu

(Feldman *et al.* 2005). The lack of any reports of *Amphibiocystidium* spp. in North America is striking given that these parasites produce clearly visible macroscopic subcutaneous cysts (Pascolini *et al.* 2003).

Little is known about the extent or pathogenicity of *Amphibiocystidium* spp. in natural populations, aside from sparse prevalence data from a few sites and sampling dates (Poisson, 1937; Broz and Privora, 1951; Pascolini *et al.* 2003). Several authors have reported general malaise and mortality of infected individuals – which they attributed to the infection (Moral, 1913; Gambier, 1924) – but other authors have not observed any noticeable effect and, to date, no study has compared mortality rates of infected amphibians to uninfected controls (Pérez, 1907; Guyenot and Naville, 1922; Pascolini *et al.* 2003). In contrast, the closely related and better characterized *Dermocystidium* spp., which infect salmonid fishes, are known to cause high mortality and have been associated with fish kills in the U.S. Pacific Northwest (Olson *et al.* 1991).

Here we present evidence of widespread *Amphibiocystidium* sp. infection in adults of a North American amphibian species, the eastern red-spotted newt (*Notophthalmus viridescens*), first reported as a *Candida*-like cyst-producing organism by Raffel (2006). We describe the unique pathology of this organism in newts, provide morphological and phylogenetic evidence for its relationship to other mesomycetozoans, and describe the seasonal distribution of this infection in multiple newt populations across the northeastern United States. We also present preliminary evidence of mortality and morbidity due to infection and the potential importance of secondary infections as a source of mortality.

MATERIALS AND METHODS

Observations and collections of newts

Newts were sampled or observed for visible signs of infection from 19 wetland locations in central Pennsylvania between 2002 and 2006, from 3 locations in the MeadWestvaco Wildlife and Ecosystem Research Forest (Cassity, West Virginia; N38°48'45", W80°3'45") in April and May 2004, and from a population in Hampshire County, Massachusetts in December 2006 (Table 1). The Pennsylvania and West Virginia newts were sampled with a combination of dip-nets and minnow traps. The Massachusetts newts were obtained from a biological supply company. For the Pennsylvania populations, a subsample of up to 10 newts was collected for dissection on each sampling date for each of the ponds included in a 2004 spatial survey or a 2003–2005 seasonal survey (Table 1). Newts examined for a 2005–2006 mark-recapture study were not collected; any additional newts were collected or observed for

unpublished experiments and surveys (Table 1). The numbers and locations of subcutaneous cysts were recorded for all Pennsylvanian newts.

Newts collected for surveys in Pennsylvania were euthanized by decapitation within 3 h of collection and dissected for internal parasite examination. Blood smears were produced and examined as described by Raffel *et al.* (2006, 2007), and 25 mg of the liver was ground up and cultured for bacteria on TSA (trypticase soy) blood agar, as described by Raffel (2006). Bacterial isolates were identified using the Biolog[®] system (Biolog Inc., Hayward, California). Culturing of cyst contents was attempted on 6 occasions, variously using TSA (trypticase soy) blood agar, corn meal agar or YPD (yeast peptide dextrose) agar. See Raffel (2006) for further details concerning collection and dissection methods for the Pennsylvanian populations. Following dissection, newts were fixed and stored in 70% ethanol. Eleven of the Massachusetts newts, including 3 containing visible subcutaneous cysts, were preserved in 70% ethanol within 24 h of death and dissected. Subcutaneous and liver cysts were counted for all dissected newts.

Mortality data collection

Upon arrival at Southern Illinois University on 18 December 2006, the Massachusetts newts were divided into groups of approximately 35 individuals and transferred to a set of 20 gallon aquariums each containing 10 gallons of deionized water treated using ASTM (1988) methods (0.03 g/l calcium sulfate, 0.03 g/l magnesium sulfate, 0.048 g/l sodium bicarbonate, and 0.002 g/l potassium chloride) and aged under aeration for 24 h prior to addition to aquaria. Aquaria were maintained at 23.8 °C with a photoperiod of 16L:8D and water was changed every other day. Newts were fed approximately 0.25 ml of captive-raised blood worms (*Chironomus tentans*) per newt following each water change, and tanks were checked for food consumption at 3 and 24 h following each feeding. Newts with visible signs of infection were observed on the day of arrival and immediately placed in a separate aquarium. A subset of 120 uninfected newts was removed for use in a separate experiment on 28 December 2006. Mortality data were recorded for all other newts until 18 January 2007. Eleven newts were preserved in 70% ethanol within 24 h of death and necropsied for confirmation of the causative agent, including 3 newts with visible subcutaneous cysts (Table 2).

Histology

Of the 15 infected newts which were collected for dissection during surveys or experiments, cysts from 12 were further examined histologically. Tissue

Table 1. Observations of red-spotted newts (*Notophthalmus viridescens*) infected with *Amphibiocystidium ranae*, compared to the total numbers of newts observed in different seasons and locations

(Observations are divided into winter (Dec, Jan, Feb), early spring (Mar, Apr), late spring (May, Jun), summer (Jul, Aug) and fall (Sep, Oct, Nov). Numbers indicate the total number of newts observed. Ponds in which the infection was found are highlighted in bold type, and bolded entries each have an additional number to indicate how many of infected newts were observed in each season and pond (# infected/ total # observed).)

Location	State	Winter	Early Spring	Late Spring	Summer	Fall	Latitude (N)	Longitude (W)	Wetland Type
Beaver 1*	PA	0	50	463	176	176	40° 45' 52.6"	78° 0' 43.6"	Permanent pond
Clearcut Pond*	PA	0	0	13	0	0	40° 46' 27.7"	77° 57' 0.0"	Ephemeral pond
Catty Ninetails*	PA	0	0	17	0	0	40° 47' 45.5"	77° 57' 15.5"	Ephemeral pond
Colyer Lake*	PA	0	0	42	0	0	40° 46' 41.8"	77° 41' 9.2"	Human impoundment
Cranberry Lake*	PA	0	0	88	0	0	40° 46' 2.6"	78° 0' 15.5"	Permanent pond
Deep Woods*	PA	0	0	38	0	0	40° 52' 9.0"	78° 4' 54.6"	Beaver wetland
False Beaver*‡	PA	55	2667	1989	360	967	40° 42' 38.3"	77° 52' 54.3"	Human impoundment
Greenbriar 1*	PA	0	0	31	0	0	40° 46' 41.3"	78° 0' 27.4"	Ephemeral pond
Irrigation Pond*	PA	0	0	81	0	0	40° 42' 18.4"	77° 56' 48.2"	Permanent pond
Little Acre*†	PA	11	10/89	77	41	50	40° 48' 5.8"	77° 56' 36.5"	Permanent pond
Mothersbaugh*†‡	PA	1/488	3/2388	5419	2798	1/1659	40° 39' 12.2"	77° 54' 9.6"	Beaver wetland
Mystery Newt*†	PA	0	109	278	190	14	40° 45' 53.0"	78° 0' 49.2"	Ephemeral pond
Muskrat Pond*	PA	0	0	37	0	0	40° 53' 8.4"	78° 4' 3.8"	Beaver wetland
Parking Lot Pond	PA	0	0	337	38	0	40° 45' 51.4"	78° 0' 58.6"	Permanent pond
Penn Roosevelt*	PA	0	0	2/ > 200	0	0	40° 43' 36.8"	77° 42' 8.3"	Human impoundment
Rock Springs	PA	0	0	0	0	3/11	40° 42' 40.0"	77° 56' 29.4"	Plastic wading pools
Ten Acre Pond	PA	0	0	459	137	218	40° 48' 4.2"	77° 56' 36.5"	Human impoundment
Turtle Shell*†‡	PA	7	566	1314	292	246	40° 52' 26.1"	78° 4' 35.6"	Beaver wetland
Twin Pond*†	PA	0	74	117	12	0	40° 46' 49.1"	78° 0' 13.9"	Ephemeral pond
Massachusetts	MA	13/180	0	0	0	0	—	—	Unknown (supply co.)
Permanent Pond	WV	0	11/34	40	0	0	—	—	Permanent pond
Marsh 2	WV	0	2/6	0	0	0	—	—	Ephemeral wetland
Ephemeral pond	WV	0	3/4	0	0	0	—	—	Ephemeral pond

* 2004 late spring survey; † 2003–2005 seasonal survey; ‡ 2005–2006 mark-recapture study.

Table 2. Results for all dissected newts, including the location, date of euthanasia/mortality, newt sex, numbers of subcutaneous and liver cysts, types of cysts observed. Bolded type indicates the cyst type from which spore measurements were taken, mean spore diameter \pm S.D. (μm), mean size of the large spore inclusion \pm S.D. (μm), and approximate numbers of granules observed within spores (min-max)

Location	Date	Sex	Cysts		Types	Spore Size	Inclusion Size	Granules
			skin	liver				
Little Acre	3/30/2004	F	163†	0	A,B	5.2 \pm 0.3	3.4 \pm 0.4	5–15
Little Acre	3/30/2004	M	10	0	A,B	7.2 \pm 0.5	4.9 \pm 0.5	5–15*
Little Acre	3/14/2006	M	3	0	A,B	7.3 \pm 0.7	5.2 \pm 0.8	5–15*
Rock Springs	11/27/2006	F	9	0	A	5.4 \pm 0.5	4.1 \pm 0.6	0–5
Rock Springs	11/27/2006	M	2	0	A	5.1 \pm 0.3	3.2 \pm 0.4	0–5
Rock Springs	11/27/2006	M	6	0	A	5.0 \pm 0.4	3.6 \pm 0.4	0–5
Penn Roosevelt	5/9/2005	M	7	1	A,B	7.3 \pm 0.5	6.0 \pm 0.5	20–40
Penn Roosevelt	5/9/2005	F	9	1	A,B	8.8 \pm 0.1	6.9 \pm 1.2	0–40*
Mothersbaugh	10/13/2004	—	5	0	A,B	4.7 \pm 0.5	2.6 \pm 0.3	0–15*
Massachusetts	1/1/2007	F	20	13†	A,B,C	6.3 \pm 0.5	4.9 \pm 0.5	20–40
Massachusetts	1/7/2007	F	2	6†	A,B,C	6.9 \pm 0.4	5.1 \pm 0.6	20–40
Massachusetts	1/19/2007	M	1	0	—	7.4 \pm 0.6	6.2 \pm 0.6	0–40*

* Granules difficult to count due to poor sample preservation; † sampled for genetic analysis.

containing 1 or more cysts was fixed in 10% buffered formalin for 24 h prior to paraffin embedding and sectioning. Cysts were sectioned at 10 μm thickness with a Shandon Finesse[®] Paraffin microtome (Thermo Electron Corporation, Waltham, Massachusetts) and stained with haematoxylin and eosin.

Diameters of spores and their inclusions from all cysts examined histologically were measured using light micrographs (1000 \times magnification). Spores were selected for measurements by zooming into a 25 μm^2 region of the cyst and measuring all spores that were in focus (6–22 spores depending on the sample) using Image-J image analysis software (Wayne Rasband, National Institutes of Health, USA). Length and diameter of 54 cysts from 3 newts, including all 3 morphologies, were measured using a dissecting microscope. Length was measured as the maximum linear dimension of the cyst and diameter was measured halfway between the cyst ends and perpendicular to the longest dimension (not necessarily the maximum diameter of the cyst due to the dumbbell shape of type C cysts, described below). Photomicrographs were taken using a Nikon Coolpix[®] 4500 camera (Melville, New York).

Genetic analysis

Cysts from 1 Little Acre newt and 2 Massachusetts newts were microdissected for DNA analysis. Samples were frozen and ground in liquid nitrogen. DNA was extracted with the Qiagen DNeasy[®] Blood and Tissue kit, using the spin-column protocol for purification of total DNA from animal tissues. DNA was eluted from spin columns with 100 μl of elution buffer. Undiluted DNA was used as template

for polymerase chain reactions (PCR). PCR was performed in 20 μl reactions using GoTaq (Promega) per manufacturer's instructions with 1.5 mM MgCl₂ and 2 μM of each primer. Touchdown PCR was performed using the following parameters: an initial 2 min denaturation at 94 °C, followed by cycles of denaturation at 94 °C for 2 min, annealing at 60 °C for 30 s (decreasing 1 °C each cycle for 10 cycles to reach 50 °C), and extension at 72 °C for 1.5 min. These cycles were followed by 30 cycles with an annealing temperature of 50 °C and a final 5 min extension at 72 °C. The PCR products were electrophoresed on 2% agarose gels to ensure that a single product was produced. For DNA sequencing, a 5 μl aliquot of each PCR reaction was incubated for 37 °C for 30 min with 5 U ExoI and 1 U shrimp alkaline phosphatase (USB) in an 8 μl reaction vol. The reaction was stopped by heating the reactions to 80 °C for 15 min. DNA sequencing was performed with the same primers used for PCR in 10 μl reactions using the ABI Prism (ABI/Perkin Elmer) Dye Terminator Cycle Sequencing Ready Reaction Kit according to the manufacturer's instructions.

Primer design

We used a modification of the universal non-metazoan primers (Bower *et al.* 2004), 18s-EUK581-F (5'-GTGCCAGCAGCCGCG-3') and 18s-EUK1134-R (5'-TTT**AARK**TTTCAGCCTT-GSG-3'), designed to amplify a 544-base pair fragment of 18s rDNA from protists without amplifying animal DNA (modifications in bold type). Specificity is provided by the final base of the reverse primer (Bower *et al.* 2004). To amplify a longer region of the 18s rDNA gene, we designed an

additional forward primer, 18s-EUK581L-F (5'-ATCAACTTTCGRTGGTAAGGTA-3'), based on an alignment of 16 mesomycetozoon 18s rDNA sequences (GenBank Accession numbers AY550245, AY692319, AY772001, AY772000, AF533941, Y19155, AF070445, AJ130859, Y16260, AY267346, AF436886, AF192386, U25637, AF232303, U21337, U21336). We verified the specificity of the reverse primer by alignment with an amphibian (toad) sequence (M59386).

Phylogenetic analysis

Mesomycetozoon 18s rRNA sequences were assembled and aligned with the program MUSCLE (Edgar, 2004). After trimming hanging ends, the resulting alignment was comprised of sequences ranging from 778 to 1780 nt in length. Accession numbers: *Dermocystidium* sp. CM-2002, AF533950; *Rhinosporidium* sp. ex. *Canis familiaris*, AY372365; *Rhinosporidium seeberi* 1, AF158369; *Rhinosporidium seeberi* 2, AF118851; *Dermocystidium salmonis*, U21337; *Rhinosporidium cygnus* from Florida swans, AF399715; *Amphibiocystidium ranae* strain 2-04, AY692319; *Dermocystidium* sp., U21336; *Amphibiocystidium ranae*, AY550245; *Sphaerothecum destruens* isolate BML, AY267345; *Choanoflagellate*-like sp., L29455; *Sphaerothecum destruens* isolate WA, AY267344; *Sphaerothecum destruens* isolate SK, AY267346; *Dermocystidium percae*, AF533941; *Amphibiothecum penneri* 1, AY772001; *Amphibiothecum penneri* 2, AY772000; *Amoebidium parasiticum* 1, Y19155; *Paramoebidium* sp. CMJ-2003 isolate KS61W6, AY336708; *Sphaeroforma arctica*, Y16260; *Amoebidium parasiticum* 2, AF274051; *Pseudoperkinsus tapetis*, AF192386; *Ichthyophonus irregularis*, AF232303; *Ichthyophonus hoferi*, U25637. The program MODELTEST (Posada and Crandall, 1998) was employed to select the optimal model of nucleotide substitution for the alignment. Both a hierarchical likelihood ratio test and Akaike information criterion found the TrN+I+ Γ model of substitution to best fit the data. A maximum likelihood (ML) tree, based on this model, was then inferred using the PAUP* (version 4.0) package (Swofford, 2003). The support for each node was determined with bootstrap resampling analysis based on 1000 pseudo-replicates of neighbour-joining trees estimated under the ML substitution model. The phylogeny was midpoint rooted, consistent with specifying the members of the order Ichthyophonida as an outgroup.

Alternative phylogenetic topologies, based off the ML tree, were constructed with the program TreeView version 1.5.3 (Page, 1996). The likelihoods of each of these topologies were compared with the Shimodaira-Hasegawa (1999) topology test (RELL distribution with 1000 bootstrap replicates), also implemented in PAUP*.

Statistical analyses

Differences in prevalence among seasons were analysed for the Little Acre, Mothersbaugh and West Virginia Permanent Pond populations by binomial regression analysis using the data presented in Table 1. To determine the degree of aggregation of infection intensity in Little Acre, the distribution of cyst counts from newts caught in early spring was compared to expected values for the Poisson and negative binomial distributions using chi-square goodness of fit tests (Parasite load categories 1–9 and 10–166 were binned to ensure expected values of ≥ 5), and the variance to mean ratio was calculated for comparison with published values for other parasites. The locations on the body (head, back, throat, stomach, forelimbs, hindlimbs or tail) of 262 cysts were recorded for 20 of the infected newts observed or collected in Pennsylvania, and the total number of cysts observed on each body part was compared to expected numbers (the proportion surface area of the body part times the total number of cysts observed) using a chi-square goodness of fit test as described by Raffel *et al.* (2007). We calculated the proportion of the total skin surface area accounted for by different body parts by analysis of newt photos using Image-J[®]. The effect of cyst type on cyst length and diameter was analysed with one-way analysis of variance using available cyst measurements, and pairwise differences between cyst types were assessed by Tukey's test for honestly significant differences (family-wise error rate of 0.05).

Mortality due to infection in the Massachusetts population was estimated using a survival analysis regression on the time to death of infected versus uninfected newts, with censoring for the 120 newts which were removed from the experiment and for those still alive at the end of the experiment. A model with logistic errors fitted significantly better than one with exponential errors ($AIC_{\text{logistic}} = 205.0$, $AIC_{\text{exponential}} = 305.6$). Statistical analyses were conducted using the 'R' statistical software system (www.cran.r-project.org).

RESULTS

Visible subcutaneous cysts were observed in newts from 4 Pennsylvania populations, 3 populations in West Virginia and in a shipment of newts collected in Massachusetts (Fig. 1A). Infections were most commonly observed in early spring but were also observed in fall and winter, with infections being found in 4 of the 9 populations sampled in early spring (Table 1). There was a significant effect of season on cyst prevalence in Little Acre ($\chi^2 = 25.2$, D.F. = 4, $P < 0.001$), Mothersbaugh ($\chi^2 = 11.2$, D.F. = 4, $P < 0.001$) and the Permanent Pond in West Virginia ($\chi^2 = 20.6$, D.F. = 4, $P < 0.001$), with early spring having the highest prevalence in all 3 locations (Table 1). Infections were seldom observed in late




Fig. 1. Macroscopic appearance of the three cyst types, shown (A) under the skin (ventral view of throat) and (B) in the liver. Representative cysts of each type are indicated by lowercase letters (a = Type A; b = Type B; c = Type C).




Fig. 2. Distributions of cyst counts. (A) Aggregated distribution of visible cyst counts for newts caught in early spring from Little Acre pond. Open diamonds indicate the expected values for the negative binomial distribution in each cyst count category. (B) Distribution of subcutaneous cysts on the newt body. Grey bars indicate the total number of cysts observed on each body part (sums of counts from 20 newts). White bars indicate cyst numbers when the single most heavily infected newt was excluded from the analysis. Black lines indicate the expected cyst counts based on the relative proportion of surface area for each body part.

spring and summer for any population despite these being the most extensively sampled seasons (Table 1).

Cyst counts were highly aggregated in the Little Acre population, with a variance to mean ratio of 133.0 (mean = 2.58, variance = 300.3, Fig. 2A). The observed distribution differed significantly from a Poisson distribution ($\chi^2 = 689$, D.F. = 5, $P < 0.001$), but not from a negative binomial distribution ($\chi^2 = 0.2$, D.F. = 2, $P > 0.8$, Fig. 2A). Subcutaneous cysts were observed on every body part, but with significantly more cysts than expected on the stomach ($\chi^2 = 174.7$, D.F. = 6, $P < 0.001$, Fig. 2B), even when the single newt with 163 cysts was removed from the analysis ($\chi^2 = 17.0$, D.F. = 6, $P = 0.009$, Fig. 2B). Of 95 newts randomly selected for dissection from Little Acre pond during the seasonal survey, 2 had visible subcutaneous cysts (Table 2) and 3 apparently

uninfected newts each had a single cyst in the liver (1 in 10 dissected newts on 5 October 2004, 1 in 8 on 28 January 2005 and 1 in 10 on 8 June 2005), providing an estimated false negative rate for presumptive diagnosis of 3.2% due to internal infections. Liver cysts were also found in 2 of the 11 dissected Massachusetts newts and 3 newts from Penn Roosevelt, all of which also had multiple subcutaneous cysts (Table 2, Fig. 1B). Cyst samples from all 12 preserved newts were confirmed by histology to be *Amphibiocystidium* sp. (the Little Acre newt liver samples were not available for histology). Syntype specimens were submitted to the U. S. National Parasite Collection (Accession numbers: 99608-99619).

Subcutaneous cysts appeared as raised bumps under the skin, which appeared white under the




Fig. 3. Differences in size between the three cyst types. Grey bars indicate cyst length (straight-line = length of the longest dimension) and white bars indicate cyst diameter (measured halfway between the cyst ends and perpendicular to the longest dimension). Different lowercase letters indicate significant differences among cyst types. Error bars represent 95% confidence intervals.

translucent skin of the throat and abdomen. Cyst shape was variable, including unbent elongated egg- to rice-shaped cysts (type A, Fig. 1A), curved C-shaped cysts (type B, Fig. 1A), and bent dumb-bell-shaped cysts with incompletely restricted globules at the ends (type C, Fig. 1C). These resembled descriptions of cyst stages by Broz and Privora (1951), except that types A and B were larger than they described, and type C cysts were not as acutely bent. Cysts of types A and B were observed on newts from all infected populations except for the Rock Springs population (Table 2, Fig. 1A). Type C cysts were only observed in the Massachusetts newts (Table 2, Fig. 1B). Cyst length and diameter both varied significantly among the cyst types (respectively: $F = 44.9$, D.F. = [2, 47], $P < 0.001$; $F = 5.3$, D.F. = [2, 47], $P = 0.008$), with type A and type C cysts having the smallest (~ 1 mm) and largest (~ 2 mm) lengths, respectively, and type B cysts having significantly larger diameters (~ 0.5 mm) than type A or type C cysts (Fig. 3). When dissected, cysts were found to contain white pus composed of many spherical hyaline spores, each containing a nucleus and a large inclusion body comprising most of the cell (Fig. 4). Each cyst was bounded by a cyst wall attached to the host connective tissue (Fig. 4A, E). Spores frequently contained additional smaller ($\sim 1 \mu\text{m}$ diameter) inclusions in the cytoplasm, hereafter referred to as 'granules', ranging from 0–5 granules in many of the type A cysts to > 20 in the type C cysts (Table 2, Fig. 3C, D). Spore diameter was generally larger in type C cysts than in types A or B, but was consistent within

individual cysts (Table 2, Fig. 3C, D). The diameter of the single large inclusion tended to be greater in cysts containing larger spores (Table 2). Although most cysts lacked membrane-delimited chambers surrounding spores (Fig. 3A), we observed that the margin of one cyst consisted of individual spores similar to the immature spores described by Poisson (1937) in that they were contained within small chambers, which appeared to disintegrate toward the centre of the cyst (Fig. 3E).

Visibly infected newts from the Massachusetts shipment had a significantly lower survival rate than uninfected newts (Coef. = -9.4 , $\chi^2 = 21.1$, D.F. = 1, $P < 0.001$), with a life expectancy of only 19.9 days after arrival compared to 29.3 days for the (visibly) uninfected newts.

The most heavily infected newt observed in the Pennsylvania populations (163 cysts) also had a severe *Pseudomonas aeruginosa* infection (> 2500 colonies from 12.5 mg liver), whereas all 9 other newts (including the other infected newt with 10 subcutaneous cysts) caught at Little Acre at the same time-point had little to no evidence of bacterial infection (3 or fewer colonies). The most heavily infected newt also had elevated numbers of lymphocytes and neutrophils (361 lymphocytes and 1105 neutrophils per 5000 erythrocytes) relative to the other 9 newts (51–161 lymphocytes and 4–27 neutrophils per 5000 erythrocytes). The 2 infected newts also had significantly less food in their stomachs than the other 8 newts, as shown using a one-tailed *t*-test assuming unequal variances (2.0 ± 2.0 mg vs 21.3 ± 8.2 mg respectively [\pm s.e.]; $t = 2.29$, D.F. = 8, $P = 0.026$).

Amphibiocystidium viridescens n. sp.

Phylogenetic analysis of Mesomycetozoean 18S rRNA regions showed that the subcutaneous and liver cysts observed in widely separated populations of red-spotted newts were all caused by phylogenetically similar parasites (Fig. 5). Although many clades within the Mesomycetozoeans cannot be sufficiently separated or their relative positions clearly resolved, due to limited sequence availability and a mix of long and short branches, it is apparent that the 3 isolates sequenced in this report (Genbank Accession numbers: EF493028–EF493030) belong to the order Dermocystida. The most likely solution is that these 3 isolates form a clade distinct from other members of the order, though additional sequence data would be needed to verify this conclusion due to low bootstrap values (Fig. 5). Indeed, the low bootstrap values make it clear that current sequence data are insufficient to strongly support or refute any of the genus designations within the order Dermocystida.

The red-spotted newt isolates resembled *A. ranae* morphologically, with subcutaneous cysts varying in




Fig. 4. Light micrographs of cysts and spores stained with haematoxylin and eosin, including (A) a typical cross-section of a subcutaneous cyst in the dermis and (B) higher magnification of the same cyst showing an aggregation of eosinophils. Typical spores from (C) a type A cyst and (D) a type C cyst, illustrating the larger size and greater number of granules for spores of the type C cyst. The lower panels depict (E) developing spores in the cyst periphery and (F) phagocytes attacking spores of a ruptured cyst. Specific features are indicated with lowercase letters (**c** = cyst; **e** = eosinophil; **g** = skin gland; **i** = large inclusion; **is** = immature spore; **j** = small granular inclusion; **m** = muscle; **n** = nucleus; **p** = phagocyte; **s** = spore; **w** = cyst wall).

shape from elongated ellipsoids to curved 'U'-shaped cysts with globular ends and containing spores with large inclusions and granules (Guyenot and Naville, 1922; Broz and Privora, 1951). Although there was no available phylogenetic evidence to determine its relationship with *A. pusula*, which infects European newts (Pérez, 1913), the presence of curved cysts distinguishes this parasite from all other closely-related subgroups except

A. ranae. The elongate cyst shape and absence of chambers containing multiple spores distinguishes this parasite from *Amphibiothecum penneri*, the only other member of this clade to infect North American amphibians (Jay and Pohley, 1981; Feldman *et al.* 2005). The presence of cysts in the liver represents an apparently unique pathology for members of this group that infect fish and amphibians (Pascolini *et al.* 2003). This, together with the fact that this




Fig. 5. ML phylogenetic tree of partial 18s rRNA sequences from the Mesomycetozoa. The sequences characterized in this study are highlighted. The tree is midpoint rooted, consistent with an Ichthyophonida outgroup and branch lengths are drawn to scale. Only highly-supported or relevant nodes within the Dermocystida are depicted with bootstrap values.

parasite represents both a new host record and a new geographical range for *Amphibiocystidium*, seems to warrant the designation of a new species.

Pascolini *et al.* (2003) postulated that *Dermocystidium*, *Dermomycoides* and *Dermosporidium* spp. infecting amphibians are monophyletic and placed them into the new genus *Amphibiocystidium*, a grouping supported by Pereira *et al.* (2005). Feldman *et al.* (2005) cast doubt upon this hypothesis by providing phylogenetic evidence for placement of *Dermosporidium penneri* (now *Amphibiothecum penneri*) basal to the *Amphibiocystidium*, *Dermocystidium*, and *Rhinosporidium* clade. We tested this hypothesis by constructing 2 alternative trees, based on our ML tree, which repositioned only the *Amphibiothecum* subgroup either immediately basal to the *Amphibiocystidium*, *Dermocystidium*, and *Rhinosporidium* subgroups or immediately basal to *Amphibiocystidium*, *Dermocystidium*, *Rhinosporidium*, and *D. percae*. The likelihoods of these two alternative topologies were not significantly better than that of the original ML tree ($P=0.109, 0.087$), indicating that, with the current data, the branching order of subgroups cannot be stated with certainty. Alternative trees with the red-spotted newt isolates forming a sister clade to *A. ranae* were not significantly different from our ML tree ($P=0.159, 0.206$). Thus, we

follow the recommendation of Pascolini *et al.* (2003) and tentatively name the new species *Amphibiocystidium viridescens* based on its morphological resemblance to *A. ranae* and its use of an amphibian host.

Presumptive identification based on the presence of visible cysts appears to be accurate provided that cysts are examined by a trained observer. All 12 randomly selected infected newts were confirmed to be infected with *A. viridescens* upon histological examination, but this only provides sufficient power to conclude that the false positive rate for presumptive diagnosis is less than 22% (based on a binomial test with $\alpha=0.05$). Nevertheless, the disease signs are distinguishable from other parasites which infect red-spotted newts. The most likely parasites to confuse with *A. viridescens* are *Clinostomum* sp. trematodes, which encyst under the skin and grow to a similar size, and an unidentified metacercarial trematode that commonly encysts in the newt liver (Raffel, 2006). Both trematodes form spherical cysts distinct from *A. viridescens*, but identifications by untrained observers should be confirmed by histology.

DISCUSSION

To our knowledge this is the first report of *Amphibiocystidium* sp. infection in a North American

amphibian species, though a separate genus designation for *Amphibiotheicum penneri*, a parasite of the American toad (*Bufo americanus*), might be unwarranted unless additional sequence data are found to support the separation. The lack of previously published accounts of this parasite in red-spotted newts is striking given the visible signs of *A. viridescens* infection, its widespread distribution in red-spotted newt populations, the early description of *Amphibiocystidium* spp. in European amphibians (Pérez, 1907), and the large number of studies which have been devoted to red-spotted newt ecology and physiology (e.g. Holl, 1932; Russell, 1951; Gill, 1979; Rohr *et al.* 2002; Sever, 2002; Muzzall *et al.* 2003). Other common parasites of newts with visible signs, such as *Clinostomum* sp. and *Ichthyophonus* sp., have been known in North American amphibians for many decades (Hopkins, 1933; Goodchild, 1953). The seasonal distribution of *A. viridescens* and its low prevalence in most populations might limit the ability of researchers to detect this parasite. However, our detection of this parasite in 3 different states over a 5-year period suggests that it should have been noted by previous researchers if its historical incidence was comparable to current levels.

Although *A. viridescens* might be a newly acquired parasite of red-spotted newts, it is unclear where the parasite would have originated, given the geographical separation of *A. viridescens* from other members of the genus *Amphibiocystidium*. Spillover of *Dermocystidium* sp. infection from fish seems unlikely since most of the infected newts were observed in fishless ponds. In addition, we are aware of an unpublished observation of heavily infected newts collected by Thomas Pauley on 28 February 1990 in Grandview State Park (Raleigh Co., W.V.), which we confirmed based on photos of infected newts and histological preparations (James Joy, personal communication). We suggest that a simpler explanation for the emergence of *A. viridescens* in red-spotted newts is a recent increase in incidence of a previously rare endemic parasite. Examination of museum specimens would be necessary to address whether the parasite was present in red-spotted newts prior to 1990.

The winter and early spring peaks in prevalence that we observed for *A. viridescens* are consistent with published accounts of *Amphibiocystidium* spp. in Europe, where infections have generally been reported in winter and early spring. Over 3 years of sampling *Triton* spp. in France, Pérez (1907, 1913) consistently found *A. pusula* infections in February and March but not at any other time of year. Over 2 years of sampling, Poisson (1937) found a consistently higher (25–30%) prevalence of *A. armoriacus* infecting *T. palmatus* in March and April than in the rest of the field season. Guyenot and Naville (1922) reported *A. ranae* infection in 12 of 200 *R. temporaria* caught in November 1921 and *A. pusula* infection of

several *T. cristatus* over the winter of 1918–1919; and Remy (1931) reported an infected *R. esculenta* individual in February 1924. Broz and Privora (1951) reported early-stage infections of *A. ranae* infecting *R. temporaria* in all seasons, but only found late-stage cysts in April. This consistent seasonal pattern differs from the seasonality found for the majority of red-spotted newt parasites, which mostly have peak infection rates in the late spring and summer (Raffel, 2006). However, winter peaks in prevalence are not uncommon amongst fungal pathogens of fish and amphibians such as *Saprolegnia* spp. and *Batrachochytrium dendrobatidis*, for which high winter infection rates have been attributed to cold-induced immune suppression coupled with the low optimal growth temperatures of the pathogens (Bly *et al.* 1993; Berger *et al.* 2004).

The considerable variability in cyst size and shape, spore size and numbers of granules in spores from different isolates of *A. viridescens* supports the argument of Pascolini *et al.* (2003) that these are poor diagnostic characteristics for distinguishing *Amphibiocystidium* species. The type C cysts of *A. viridescens* resemble the mature cysts of *A. ranae* (Guyenot and Naville, 1922; Pascolini *et al.* 2003), and type A cysts might be difficult to distinguish from the ‘spherical’ and ‘egg-shaped’ cysts of *A. pusula* and *A. hyalarum*, respectively (Pérez, 1913; Carini, 1940). Broz and Privora (1951) described a range of cyst sizes and types similar to our 3 cyst types in *A. ranae*, attributing the differences to different developmental stages. The steady increase in cyst size from type A through to type C cysts observed in this study support their conclusion, as do the larger and more granular spores we observed in type C cysts.

This developmental progression is not unlike the development of closely related *Dermocystidium salmonis*, which transmits by tiny flagellated zoospores that develop within spores from granules similar to those in *Amphibiocystidium* spp. spores (Olson *et al.* 1991). *D. salmonis* spores each possess a large inclusion similar to those seen in *A. viridescens*, which become progressively smaller as the zoospores mature (Olson *et al.* 1991). Zoospores have not been observed in *Amphibiocystidium* spp. infections (Pascolini *et al.* 2003), but this is not surprising given the 2-week incubation period before release of zoospores in *D. salmonis* (Olson *et al.* 1991). Several authors have described progressive weakening of the skin covering cysts of *Amphibiocystidium* spp., leading to cyst rupture and the release of spores onto the skin surface (Pérez, 1913; Broz and Privora, 1951). Perhaps zoospore development follows cyst rupture in *Amphibiocystidium* spp., so that zoospores are released from spores resting in the benthos. This mechanism would help explain the high numbers of cysts on the ventral surface of newts observed in this study, since the source of infection would be underneath the newt.

This is, to our knowledge, the first report of *Amphibiocystidium* sp. infection in the liver. Since many authors have examined this parasite solely through external examination and biopsies (e.g. Pascolini *et al.* 2003) and the actual number of specimens dissected is unclear in other studies (Moral, 1913; Pérez, 1913; Remy, 1931; Broz, 1944), it seems possible that liver infections may have simply been missed by other researchers. The related parasite *Rhinosporidium seeberi*, associated with human disease, has been reported from the liver and other soft tissues in very rare disseminated infections (Branscomb, 2002). However, spores or other identifiable stages have not been observed outside cysts for any *Amphibiocystidium* species (Pascolini *et al.* 2003), and an analysis of the parasite distribution in the Little Acre population provided no support for the hypothesis that *A. viridescens* replicates within the host to produce additional cysts. This hypothesis predicts that the distribution of cyst counts in the host population should be more highly aggregated than in macroparasitic infections. The distribution of cysts in the Little Acre population was aggregated but fit a negative binomial distribution, as do most macroparasitic helminth infections (Shaw *et al.* 1998). The observed log-variance of 2.5 in this study was higher than the expected value of 1.9 for macroparasites with the same log-mean intensity, but within the range observed in a large sample of helminth and arthropod parasites (Shaw and Dobson, 1995). Although we cannot rule out disseminated infection based on this analysis, the result is consistent with the hypothesis that each cyst represents a discrete infection event, despite the high numbers of cysts observed in some individuals. Perhaps infectious stages of *A. viridescens* access the liver through the bile duct following ingestion of mature spores by foraging newts, as an alternative to the hypothesis of disseminated infection.

Previous studies have asserted that *Amphibiocystidium* spp. are relatively benign parasites of amphibians (Pérez, 1913; Guyenot and Naville, 1922; Broz and Privora, 1951), and although *A. ranae* was associated with population declines of *Rana lessonae*, the lack of systemic post-mortem examinations has made it difficult to determine whether the parasite contributed to declines (Pascolini *et al.* 2003). Our finding of significantly higher mortality in the visibly infected newts from Massachusetts suggests that *A. viridescens* can reduce red-spotted newt survival, though we cannot rule out the possibility of an additional mortality factor in the infected newts' tank or that the stress of shipping newts and placing them in a new environment exacerbated the disease. Newts in this shipment also had substantially suppressed appetites compared to newts in previous shipments (Bommarito, T., personal observation), consistent with the low appetite in an infected *Triton cristatus* observed by Moral (1913). Preliminary

evidence suggests that infected newts also experience decreased appetite in natural ponds, since the most heavily infected newts in Little Acre pond had less food in their stomachs than uninfected newts. In addition, the single most heavily infected newt in the seasonal survey was co-infected with *Pseudomonas auruginosa*, suggesting that the proximate cause of mortality due to *A. viridescens* infection might be secondary infections.

The incidence of *A. viridescens* in red-spotted newt populations might be higher than suggested by the low prevalence recorded in this study, since parasites with short infectious periods are less likely than persistent parasitic infections to be detected in horizontal surveys. While the length of the infectious period in natural ponds remains unknown, red-spotted newts which survive the infection appear to lose their cysts in a relatively short amount of time. The one infected newt, still alive at the end of the mortality study apparently cleared its infection within 25 days of arrival (Bommarito, T., personal observation). In addition, an infected newt collected in 2004 from a Pennsylvania site also lost its single subcutaneous cyst after being held in the lab for approximately 2 weeks, although the site of infection still appeared inflamed (Raffel, T., personal observation). The one internally ruptured cyst observed in this study was infiltrated by a large number of eosinophils and other phagocytic cells. These observations are consistent with published accounts of *A. pusula* and *A. ranae* cysts, which are infiltrated by neutrophils shortly following cyst rupture, and healed in a matter of days (Pérez, 1913; Broz, 1944). The timing of cyst rupture appears to be controlled more by the parasite than by the host, since the host response to intact cysts is usually minimal (Pascolini *et al.* 2003), although we observed an eosinophilic response to one intact cyst – consistent with that described by Broz (1944) for *A. ranae*.

Although the significance of *A. viridescens* to red-spotted newt populations remains an open question, its apparently recent emergence in North America is troubling. Other pathogens and parasites also seem to be increasing in incidence and geographical range amongst North American amphibians, including deformity-inducing trematodes (Johnson *et al.* 2003), ranaviruses (Chinchar, 2002), and the chytrid fungus associated with amphibian declines (Green *et al.* 2002; Weldon *et al.* 2004), which has recently been found for the first time in red-spotted newts (Padgett-Flohr *et al.* 2007). Reasons for the emergence of these parasites remain unclear, but human-induced environmental changes and transportation of infected carriers of the diseases appear to be important (Johnson and Chase, 2004; Jancovich *et al.* 2005). Red-spotted newts and other North American amphibians play crucial roles in North American ecosystems (Kurzava and Morin, 1998), and determining the potential causes and severity of

emerging infections will be important for ensuring the future conservation of these species.

We would like to thank P. Hudson and J. Rohr of Penn State University, T. Pauley of the WV Herpetological Museum, J. Joy and R. Gain of Marshall University, and the Cooperative Wildlife Research Lab of Southern Illinois University-Carbondale for their time, consultation and supplies. Histological sectioning was performed by R. Haldeman of the Penn State EM Facility. C. Barry, Z. Loughman, J. Sias, J. Falkenbach, J. Dillard, R. LeGros, and R. Huang provided significant assistance with field observations and dissections. M. Lopez performed the bacterial identification with the help of F. Zambito from the PSU Animal Diagnostic Lab. W. Marshall and S. Feldman provided advice about primer selection, and K. Vandegrift and S. Lass provided comments on early drafts of the manuscript. This work was supported in part by a National Science Foundation (NSF) Fellowship to T. Raffel, NSF Dissertation Improvement Grant #0508847, and NSF Grant #0516227. Funding for the Massachusetts newts was provided by the Barton Springs Salamander Conservation Fund and the City of Austin, Texas. Funding for West Virginia sampling was provided by the MeadWestvaco Corporation.

REFERENCES

- ASTM.** (1988). *Standard Practice for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates, and Amphibians*. ASTM, West Conshohocken, PA, USA.
- Berger, L., Speare, R., Hines, H. B., Marantelli, G., Hyatt, A. D., McDonald, K. R., Skerratt, L. F., Olsen, V., Clarke, J. M., Gillespie, G., Mahony, M., Sheppard, N., Williams, C. and Tyler, M. J.** (2004). Effect of season and temperature on mortality in amphibians due to chytridiomycosis. *Australian Veterinary Journal* **82**, 434–439.
- Bly, J. E., Lawson, L. A., Szalai, A. J. and Clem, L. W.** (1993). Environmental factors affecting outbreaks of winter saprolegniosis in Channel catfish, *Ictalurus punctatus* (Rafinesque). *Journal of Fish Diseases* **16**, 541–549.
- Bower, S. M., Carnegie, R. B., Goh, B., Jones, S. R. M., Lowe, G. J. and Mak, M. W. S.** (2004). Preferential PCR amplification of parasitic protistan small subunit rDNA from metazoan tissues. *Journal of Eukaryotic Microbiology* **51**, 325–332.
- Branscomb, R.** (2002). Rhinosporidiosis update. *Laboratory Medicine* **33**, 631–633.
- Broz, O.** (1944). Die Herkunft der Zystenmembran von *Dermocystidium ranae*. *Vestník Československé Společnosti Zoologické* **9**, 16–25.
- Broz, O. and Privora, M.** (1951). Two skin parasites of *Rana temporaria*: *Dermocystidium ranae* Guyenot & Naville and *Dermosporidium granuloso* n. sp. *Parasitology* **42**, 65–69.
- Carini, A.** (1940). Sobre um parasito semelhante ao “*Rhinosporidium*” encontrado em quistos da pele de uma “*Hyla*”. *Arquivos do Instituto Biológico* **11**, 93–98.
- Chinchar, V. G.** (2002). Ranaviruses (family Iridoviridae): emerging cold-blooded killers. *Archives of Virology* **147**, 447–470.
- Daszak, P., Cunningham, A. A. and Hyatt, A. D.** (2001). Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Tropica* **78**, 103–116.
- Edgar, R. C.** (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**, 1792–1797.
- Feldman, S. H., Wimsatt, J. H. and Green, D. E.** (2005). Phylogenetic classification of the frog pathogen *Amphibiotheicum* (*Dermosporidium*) *penneri* based on small ribosomal subunit sequencing. *Journal of Wildlife Diseases* **41**, 701–706.
- Gambier, H.** (1924). Sur un Protiste parasite et pathogene des Tritons: *Hepatosphera molgarum* n. g., n. sp. *Comptes Rendus des Séances de la Société de Biologie et de ses Filiales* **90**, 439–441.
- Gill, D. E.** (1979). Density dependence and homing behavior in adult red-spotted newts, *Notophthalmus viridescens* (Rafinesque). *Ecology* **60**, 800–813.
- Goodchild, C. G.** (1953). A subcutaneous, cyst-parasite of Bullfrogs: *Histocystidium ranae*, n. g., n. sp. *Journal of Parasitology* **39**, 395–405.
- Green, D. E., Converse, K. A. and Schrader, A. K.** (2002). Epizootiology of sixty-four amphibian morbidity and mortality events in the USA, 1996–2001. *Annals of the New York Academy of Sciences* **969**, 323–339.
- Green, D. E., Feldman, S. H. and Wimsatt, J. H.** (2003). Emergence of a *Perkinsus*-like agent in anuran liver during die-offs of local populations: PCR detection. *American Association of Zoo Veterinarians Conference Proceedings, Milwaukee, Wisconsin*, 120–121.
- Green, D. E. and Sherman, C. K.** (2001). Diagnostic histological findings in Yosemite toads (*Bufo canorus*) from a die-off in the 1970s. *Journal of Herpetology* **35**, 92–103.
- Guyenot, E. and Naville, A.** (1922). Un nouveau protiste du genre *Dermocystidium* parasite de la Grenouille *Dermocystidium ranae* nov. spec. *Revue Suisse de Zoologie* **29**, 133–145.
- Holl, F. J.** (1932). The ecology of certain fishes and amphibians with special reference to their helminth and linguatulid parasites. *Ecological Monographs* **2**, 83–107.
- Hopkins, S. H.** (1933). Note on the life history of *Clinostomum marginatum* (Trematoda). *Transactions of the American Microscopical Society* **52**, 147–149.
- Jancovich, J. K., Davidson, E. W., Parameswaran, N., Mao, J., Chinchar, V. G., Collins, J. P., Jacobs, B. L. and Storfer, A.** (2005). Evidence for emergence of an amphibian iridoviral disease because of human-enhanced spread. *Molecular Ecology* **14**, 213–224.
- Jay, J. M. and Pohley, W. J.** (1981). *Dermosporidium penneri* sp n from the skin of the American toad, *Bufo americanus* (Amphibia, Bufonidae). *Journal of Parasitology* **67**, 108–110.
- Johnson, P. T. and Lunde, K. B.** (2005). Parasite infection and limb malformations: A growing problem in amphibian conservation. In *Amphibian Declines: the Conservation Status of United States Species* (ed. Lannoo, M. J.). pp. 124–138. University of California Press, Berkeley, USA.
- Johnson, P. T. J. and Chase, J. M.** (2004). Parasites in the food web: linking amphibian malformations and aquatic eutrophication. *Ecology Letters* **7**, 521–526.

- Johnson, P. T. J., Lunde, K. B., Zelmer, D. A. and Werner, J. K.** (2003). Limb deformities as an emerging parasitic disease in amphibians: Evidence from museum specimens and resurvey data. *Conservation Biology* **17**, 1724–1737.
- Kurzava, L. M. and Morin, P. J.** (1998). Tests of functional equivalence: Complementary roles of salamanders and fish in community organization. *Ecology* **79**, 477–489.
- Lips, K. R., Brem, F., Brenes, R., Reeve, J. D., Alford, R. A., Voyles, J., Carey, C., Livo, L., Pessier, A. P. and Collins, J. P.** (2006). Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proceedings of the National Academy of Sciences, USA* **103**, 3165–3170.
- Moral, H.** (1913). Über das Auftreten von *Dermocystidium pusula* (Pérez), einem einzelligen Parasiten der Haut des Molches bei *Triton cristatus*. *Archiv für mikroskopische Anatomie* **81**, 381–393.
- Muzzall, P. M., Peterson, J. D. and Gilliland, M. G.** (2003). Helminths of *Notophthalmus viridescens* (Caudata: salamandridae) from 118th Pond, Michigan, USA. *Comparative Parasitology* **70**, 214–217.
- Olson, R. E., Dungan, C. F. and Holt, R. A.** (1991). Water borne transmission of *Dermocystidium salmonis* in the laboratory. *Diseases of Aquatic Organisms* **12**, 41–48.
- Padgett-Flohr, G. E., Bommarito, T. and Sparling, D.** (2007). Amphibian chytridiomycosis: Implications regarding amphibian research. *Herpetological Review* (in the Press).
- Page, R. D. M.** (1996). TREEVIEW: An application to display phylogenetic trees on personal computers. *Computer Applications in the Biosciences* **12**, 357–358.
- Pascolini, R., Daszak, P., Cunningham, A. A., Tei, S., Vagnetti, D., Bucci, S., Fagotti, A. and Di Rosa, I.** (2003). Parasitism by *Dermocystidium ranae* in a population of *Rana esculenta* complex in central Italy and description of *Amphibiocystidium* n. gen. *Diseases of Aquatic Organisms* **56**, 65–74.
- Pereira, C. N., Di Rosa, I., Fagotti, A., Simoncelli, F., Pascolini, R. and Mendoza, L.** (2005). The pathogen of frogs *Amphibiocystidium ranae* is a member of the order Dermocystida in the class Mesomycetozoa. *Journal of Clinical Microbiology* **43**, 192–198.
- Pérez, C.** (1907). *Dermocystis pusula* organisme nouveau parasite de la peau des Tritons. *Comptes Rendus de la Société de Biologie* **63**, 445–447.
- Pérez, C.** (1913). *Dermocystidium pusula*: Parasite de la peau des Tritons. *Archives de Zoologie Expérimentale et Générale* **52**, 343–357.
- Poisson, C.** (1937). Sur une nouvelle espèce du genre *Dermomycoïdes* Granata 1919: *Dermomycoïdes armoriacus* Poisson 1936 parasite cutané de *Triturus palmatus* (Schneider). Genèse et structure de la zoospore. *Bulletin Biologique de la France et de la Belgique* **71**, 91–116.
- Posada, D. and Crandall, K. A.** (1998). Modeltest: testing the model of DNA substitution. *Bioinformatics* **14**, 817–818.
- Raffel, T. R.** (2006). Drivers of seasonal infection dynamics in the parasite community of red-spotted newts (*Notophthalmus viridescens*). Ph. D. thesis. The Pennsylvania State University, University Park, USA.
- Raffel, T. R., Dillard, J. R. and Hudson, P. J.** (2007). Field evidence for leech-borne transmission of amphibian *Ichthyophonus*. *Journal of Parasitology* **92**, 1256–1264.
- Raffel, T. R., Rohr, J. R., Kiesecker, J. M. and Hudson, P. J.** (2006). Negative effects of changing temperature on amphibian immunity under field conditions. *Functional Ecology* **20**, 819–828.
- Remy, P.** (1931). Presence de *Dermocystidium ranae* (Guyenot et Naville) chez une *Rana esculenta* L. de Lorraine. *Annales de Parasitologie Humaine et Comparée* **9**, 1–3.
- Rohr, J. R., Madison, D. M. and Sullivan, A. M.** (2002). Sex differences and seasonal trade-offs in response to injured and non-injured conspecifics in red-spotted newts, *Notophthalmus viridescens*. *Behavioral Ecology and Sociobiology* **52**, 385–393.
- Russell, C. M.** (1951). Survey of the intestinal helminths of *Triturus v. viridescens* in the vicinity of Charlottesville, Virginia. *Virginia Journal of Science* **2**, 215–219.
- Sever, D. M.** (2002). Female sperm storage in amphibians. *Journal of Experimental Zoology* **292**, 165–179.
- Shaw, D. J. and Dobson, A. P.** (1995). Patterns of macroparasite abundance and aggregation in wildlife populations: A quantitative review. *Parasitology* **111** (Suppl.), S111–S133.
- Shaw, D. J., Grenfell, B. T. and Dobson, A. P.** (1998). Patterns of macroparasite aggregation in wildlife host populations. *Parasitology* **117**, 597–610.
- Shimodaira, H. and Hasegawa, M.** (1999). Multiple comparisons of log-likelihoods with applications to phylogenetic inference. *Molecular Biology and Evolution* **16**, 1114–1116.
- Swofford, D. L.** (2003). *PAUP*: Phylogenetic Analysis using Parsimony (and other Methods). Version 4.0*. Sinauer Associates, Sunderland, MA, USA.
- Weldon, C., du Preez, L. H., Hyatt, A. D., Muller, R. and Speare, R.** (2004). Origin of the amphibian chytrid fungus. *Emerging Infectious Diseases* **10**, 2100–2105.