

SHORT COMMUNICATION

Heteropneustes microps, a junior synonym of *H. fossilis* (Osteichthyes: Heteropneustidae)

Rohan Pethiyagoda* and Mohamed M. Bahir*

Heteropneustes microps (Günther, 1864) (type locality: Dambuwa, Sri Lanka) is distinguished from its nominal congeners only by its caudal and anal fins being confluent (vs. separate). The other species of this genus from South Asia, *H. singio* (Hamilton, 1822) (Ganges Basin) and *H. microcephalus* (Günther, 1864) (Sri Lanka) were treated by Day (1875-78), Hora (1936), Jayaram (1981) and Talwar & Jhingran (1991) as synonyms of *H. fossilis* (Bloch, 1794) (Tranquebar, south-east India).

Pethiyagoda (1991: 165) observed that in Sri Lanka, *H. microps* is found to occur only rarely and even then, only together with *H. fossilis*, representing about two percent of the catch of *H. fossilis*, where there is a fishery. At the same time, Pethiyagoda (*op. cit.*; 1994) also expressed concern for the continued survival of this species.

Confluence of unpaired fins has been observed occasionally in *Clarias brachysoma* (Günther, 1864) in Sri Lanka (Pethiyagoda, 1994: 161), but these have been thought to be aberrant specimens.

We tested the possibility that the confluent anal and caudal fins of *H. microps* could be the result of fusion following an injury. We made a small incision into the lower hypural area—between the caudal and anal fins—of five living specimens of *H. fossilis* and found that in the course of healing, the anal and caudal fins became completely fused (in the case of four of the five specimens tested, the caudal fin was entirely lost, but when regeneration occurred it was in confluence with the anal fin). No visible external evidence of injury remained.

An examination of the syntypes of *H. microps* and their radiographs at the BMNH (3 ex., BMNH 1859.5.31.9-11) and recently-collected material from Sri Lanka confirmed that these specimens all had damaged or malformed hypurals. Figure 1 shows the caudal skeleton of an "*H. microps*" from Navinna, (Galle: Southern Province) and Fig. 2, that of an *H. fossilis* from Attidiya-Bellanwila (Western Province, Sri Lanka).

It would appear therefore that *H. microps* is nothing more than the result of anomalous fin regeneration in *H. fossilis*, injury being at least one of the possible causes. We therefore consider *H. microps* a junior synonym of *H. fossilis*.

The occurrence of '*H. microps*' elsewhere in the range of *H. fossilis* should therefore occasion no surprise. Datta Munshi & Srivastava (1988: 298) reported a single specimen from Bihar, India. Talwar & Jhingran's (1991: 691) state-

* Wildlife Heritage Trust of Sri Lanka, 95 Cotta Road, Colombo 8, Sri Lanka.



Figure 1. Radiograph of tail portion of *Heteropneustes microps* from Navinna (Galle: Southern Province), Sri Lanka, WHT 1323, 86.1 mm SL.




Figure 2. Radiograph of tail portion of *Heteropneustes fossilis* from Attidiya-Bellanwila (Western Province) Sri Lanka, 117.4 mm SL, WHT 1324.

ment that *H. microps* (which they diagnosed correctly) '... is esteemed as food fish for its invigorating qualities. In Bihar tanks are frequently stocked with this fish during the rainy season', however, is mystifying.

Literature cited

- Bloch, M.E. 1794. Naturgeschichte der ausländischen Fische. Berlin. 8: i-iv, 1-174; pls. 361-396.
- Datta Munshi, J.S. & M.P. Srivastava. 1988. Natural history of fishes and systematics of freshwater fishes of India. Narendra Publishing House, Delhi, 403 pp.
- Day, F. 1875-78. The Fishes of India; being a natural history of the fishes known to inhabit the seas and fresh waters of India, Burma and Ceylon. Bernard Quaritch, London. 1: (text) 778 pp.; 2: (atlas) 198 pl.
- Günther, A. 1864. Catalogue of the fishes in the British Museum, 5: Catalogue of the Physostomi containing the families Siluridae, Characinidae, Haplochitonidae, Sternoptychidae, Scopelidae, Stomiatidae. British Mus., London, xxii + 455 pp.
- Hamilton, F. 1822. An account of the fishes found in the River Ganges and its branches. Archibald Constable (Edinburgh) & Hurst Robinson (London), vii + 405 pp, 39 pl.
- Hora, S.L. 1936. Siluroid fishes of India, Burma and Ceylon, 5: Fishes of the genus *Heteropneustes* Müller. Rec. Indian Mus., 38(2): 208-209.
- Jayaram, K.C. 1981. The freshwater fishes of India, Pakistan, Bangladesh, Burma and Sri Lanka — a handbook. Zool. Surv. India, 475 pp., 13 pl.
- Pethiyagoda, R. 1991. Freshwater fishes of Sri Lanka. Wildlife Heritage Trust, Colombo, xiv+362 pp.
- Pethiyagoda, R. 1994. Threats to the indigenous freshwater fishes of Sri Lanka and remarks on their conservation. Hydrobiologia, 285:189-201.
- Talwar, P.K. & A.G. Jhingran. 1991. Inland fishes of India and adjacent countries. Oxford & IBH, New Delhi, 1: liv+1-542 pp; 2: 543-1158 pp.