

Dietary Habits and the Predators of the Bengal Monitor *Varanus bengalensis* in Sri Lanka

SURANJAN KARUNARATHNA¹, THILINA SURASINGHE^{2,*}, DUMINDA DISSANAYAKE³,
MADHAVA BOTEJUE^{4,5}, DINESH GABADAGE^{4,6} & MAJINTHA MADAWALA⁷

¹*Nature Explorations and Education Team*
No: B-1 / G-6, De Soysapura, Moratuwa 10400, SL
E-mail: dmsameera@gmail.com

²*Department of Biological Sciences, Bridgewater State University*
Bridgewater, MA 02325, USA
E-mail: tsurasinghe@bridgew.edu
*Corresponding author

³*Department of Biological Sciences, Faculty of Applied Sciences, Rajarata University*
Mihintale 50300, SL
E-mail: duminda.rusl@gmail.com

⁴*Biodiversity Conservation Society*
150/6 Stanley Thilakarathne Mawatha, Nugegoda 10250, SL

⁵E-mail: madhavabotejue@gmail.com

⁶E-mail: degabadage@gmail.com

⁷*South Australian Herpetology Group, South Australian Museum*
North Terrace, Adelaide, SA 5000, Australia
E-mail: majintham@yahoo.com

Abstract - The Bengal monitor (*Varanus bengalensis*) is the second largest lizard species in Sri Lanka, and is well adapted to living in a variety of terrestrial habitats. *Varanus bengalensis* is a diurnal generalist predator, but sometimes function as a scavenger. Given the ecological plasticity and generalist foraging strategy of *V. bengalensis*, its feeding ecology and role in trophic networks could vary among different environments. Thus, in this study we documented the prey and natural predators of *V. bengalensis* across different landscapes along the urban-rural gradient and geo-climatic zones of Sri Lanka through field observations, literature surveys, and personal communications with other biologists. We documented 82 species of vertebrate prey in the diet of *V. bengalensis*, which included 20 mammals, 21 birds, 20 reptiles, 14 amphibians, and seven fishes. Although numerous invertebrates were recorded, their identification to species level was challenging. *Varanus bengalensis* also fed on road-killed animals and organic waste, such as rotting fruits and vegetables. Although *V. bengalensis* is largely terrestrial, our dietary analyses revealed that it also preys on aquatic prey; their predation on introduced species as well as threatened species was also remarkable. Our study confirmed that *V. bengalensis* has a greater dietary selection than *V. salvator* in Sri Lanka, which remained consistent across a diverse array of habitats. Our study also noted that *V. bengalensis* is predated by 24 species of vertebrates, including nine mammals, 10 birds, and five reptiles, which largely targeted juveniles. Neither the predator nor prey communities of *V. bengalensis* varied across the urban-rural gradient or across different geo-climatic zones of Sri Lanka.

Introduction

Varanid lizards, or monitors, are a highly diverse reptilian group that is broadly distributed throughout Africa, South and Southeast Asia, and Australasia (Pianka *et al.*, 2004). In general, varanids have a highly conservative body plan, but vary greatly in size, and have successfully colonized a wide range of habitats in various geo-climatic zones (Pianka & King, 2004). Their ecological success has frequently been attributed to elevated seasonal and daily activity levels, high stamina and metabolic rates, cardiovascular efficiency, respiratory efficiency due to gular pumping, and greater chemosensory perception (Pianka *et al.*, 2004).

With a few exceptions (*e.g.*, *Varanus bitatawa*, *V. olivaceus*, *V. indicus*), most varanids are highly-adaptive habitat generalists and opportunistic carnivores (Shine, 1986; Losos & Greene, 1988; Molnar, 2004). Certain species, such as, the Bengal monitor, *V. bengalensis*, have successfully established themselves in highly altered anthropogenic habitats, and also occur in a variety of natural landscapes including dense forests, sparse woodlands, grasslands, and thorny scrublands (Pianka, 2004). The geographic distribution of *V. bengalensis* extends from southeastern Iran through much of southern Asia, into the Malay Archipelago (Pianka, 2004). As in many wide-ranging varanids, the ecology and natural history of *V. bengalensis* vary remarkably throughout its range in response to the different habitats and geo-climatic zones they inhabit (Losos & Greene, 1988; Auffenberg, 1994; Pianka, 1995). In this study, we focused on the trophic ecology of *V. bengalensis* in Sri Lanka, an Indian oceanic island where varanid ecology still remains largely unexplored.

Two species of monitor lizard (genus *Varanus*) occur on Sri Lanka - *V. bengalensis* and *V. salvator* (de Silva, 2006). *Varanus bengalensis* has been recorded throughout Sri Lanka up to 500 m in elevation and across all geo-climatic zones (Das & de Silva, 2005). *Varanus bengalensis* is diurnal, and in Sri Lanka, due to year-round uniformly high temperatures, is active throughout the year. Daily activity is mostly limited to mid-day (1000 to 1400 h), and *V. bengalensis* is rarely observed during early mornings or late evenings (Deraniyagala, 1953; Wikramanayake & Dryden, 1993; Karunarathna *et al.*, 2012). Being a moderately large (average total length [TL] 100 cm) varanid, *V. bengalensis* is the second largest lizard species in Sri Lanka (Auffenberg, 1994), with the longest individual recorded measuring 174 cm TL at Hambegamuwa, a rural village in southern Sri Lanka (S. Karunarathna, pers. obs., 2013). *Varanus*

bengalensis is well adapted to occupy a variety of terrestrial habitats, and can also climb vertical surfaces and trees if necessary (Auffenberg, 1994). Although the species is found in floodplain woodlands, it tends to avoid aquatic habitats and is inept at swimming or diving (Pianka, 2004). In Sri Lanka, the thermoregulatory behavior, microhabitat use, metabolism (with respect to osmotic balance and energy), and behavioral aspects (particularly conspecific agnostic interactions) of *V. bengalensis* have been investigated (Wikramanayake & Green, 1989; Dryden *et al.*, 1992; Wikramanayake and Dryden, 1993); however, their trophic interactions remain largely unexplored. *Varanus bengalensis* are active generalist predators, feeding mostly on small invertebrate prey (Deraniyagala, 1953). They are also human commensals, and thrive in anthropogenic habitats such as home gardens and homesteads, and can even occupy households and built-up urban environments (Koch *et al.*, 2013).

Resource and microhabitat availability and environmental complexity differs remarkably across urban-rural gradients as well as geo-climatic gradients (Faeth *et al.*, 2005). Given the ecological plasticity and generalist foraging strategy of *V. bengalensis*, its feeding ecology and role in trophic networks could vary in different environments. Thus, documenting the diet and natural predators of *V. bengalensis* across different landscapes and geo-climatic zones are salient elements of ecological theory. In this paper, we studied the prey and predators of *V. bengalensis* in Sri Lanka.

Methods

Study Area

Sri Lanka is a tropical island (area: 65,610 km²) located in the Indian Ocean off the southern tip of peninsular India (5° 55' 0.12" – 9° 50' 60" N; 79° 40' 59.88" E – 81° 54' 0" E). The island consists of three elevation-based geographic zones (lowlands: < 300 m, uplands: 300-900 m, and highlands: > 900 m) and three major climatic zones based on average annual precipitation (dry zone: < 1900 mm, wet zone: > 2500 mm, and intermediate zone: 1900-2500 mm) (Survey Department of Sri Lanka, 2012).

Data collection

We collected data on the prey and predators of *V. bengalensis* using various methods. Primarily, we made extensive field observations through opportunistic field

excursions conducted over a six-year (2010-2016) period in various regions of Sri Lanka representing different geo-climatic zones as well as habitats along the urban-rural gradient. Our field observations were made at distances of 2 to 20 m from the animals using 8×40 binoculars. Our field surveys were limited to periods of activity for *V. bengalensis*, between 0600 and 1800 h (Wikramanayake & Green, 1989; Wikramanayake & Dryden, 1993). We dissected ~12 fresh road-killed *V. bengalensis* from various parts of Sri Lanka and identified their stomach contents. Our direct feeding observations also included *V. bengalensis* feeding on road kills. During our field excursions, we also interviewed residents of local communities (~200 local residents) regarding their random observations on the trophic roles of *V. bengalensis*.

Additionally, we consulted veteran herpetologists and field biologists (a total of ~25) who have conducted field research in Sri Lanka in recent decades (1990-2016) on their experiences with *V. bengalensis* through in-person communication and social media. Whenever possible, we requested photographic evidence to validate their observations.

Finally, we conducted a comprehensive literature review utilizing online databases and scholarly search engines (Google Scholar, MEDLINE, Science Direct, Biological Sciences Collection, Academic Search Premier, EBSCOhost, JSTOR, and PubMed). We used a combination of the following keywords in the literature review: “*Varanus bengalensis*”, “Bengal monitor”, “Clouded monitor”, “Common Indian monitor”, “feeding”, “foraging”, “prey”, “predator”, “trophic”, “diet”, “Sri Lanka”, and “Ceylon”. We categorized the prey and predators based on their IUCN conservation status (Vulnerable, Endangered, and Critically Endangered) and residential status (endemic, residential, and non-native).

Results

Prey species

Our field-based observations, interviews with professional biologists, and review of the literature revealed a diverse array of prey consumed by *V. bengalensis* which included vertebrates, invertebrates, as well as dead organic matter including road-killed animals and household garbage and urban trash (Table 1; Fig. 1). We noted that the vertebrate prey of *V. bengalensis* was remarkably high in diversity: 20 mammals, 21 birds, 20 reptiles, 14 amphibians, and

7 freshwater fishes (a total of 82 species; Table 1). Among invertebrate prey, species-level identification was limited. Most of the invertebrate taxa preyed by *V. bengalensis* were arthropods. Live prey included different life-history stages (eggs, larvae, juveniles, and adults). *Varanus bengalensis* appeared to consume animal carcasses of different stages of decomposition ranging from fresh carcasses (such as road-kills) to carrion (rotting animal flesh). Among live prey, these monitors consumed both endemic species (Sri Lankan toque monkey [*Macaca sinica*] and Sri Lankan yellow-fronted barbet [*Megalaima flavifrons*]) and non-native species including invasive species (1 species) as well as domesticated species (2 species). Moreover, 10 endemics and a total of 5 IUCN Red Listed species (2 critically endangered, 2 endangered, and 1 vulnerable) were also documented in our survey.

Predators

Our study revealed that *V. bengalensis* has several natural predators, including 9 mammals, 10 birds, and 5 reptiles (Table 2; Fig. 2). Among these predators, we found one endemic species (Sri Lankan grey hornbill [*Ocyrceros gingalensis*]) and one non-native species (domestic cat [*Felis catus*]). Of the 24 total records of predation we documented, juveniles were the victims in 14 (58.3%) instances, suggesting that juveniles are the most susceptible life-history stage (Table 2). In five instances, consumers of *V. bengalensis* were carrion feeders scavenging on road-killed individuals (jackal [*Canis aureus*], wild boar [*Sus scrofa*], and jungle crow [*Corvus leuallantii*]). Live, adult *V. bengalensis* suffered direct predation in a handful of instances (by five species). Most *V. bengalensis* predators were visually-oriented, active foragers (domestic cats, domestic dogs [*Canis familiaris*], leopards [*Panthera pardus*], mongoose [*Herpestes brachyurus*], birds of prey [e.g., *Spilornis cheela*], Indian cobra [*Naja naja*], and rat snakes [*Ptyas mucosa*]), while a few were sit-and-wait foragers (mugger crocodile [*Crocodylus palustris*], stork-billed kingfisher [*Pelargopsis capensis*], and lesser adjutant [*Leptoptilos javanicus*]). Moreover, one predator is a threatened species listed in the IUCN Red List (leopard, *Panthera pardus*) while another (Sri Lankan grey hornbill, *Ocyrceros gingalensis*) is an endemic species. Humans are also considered a key predator of *V. bengalensis*, and harvest these monitors from the wild for their flesh, hide, and fat. Neither the natural predators nor prey of *V. bengalensis* were restricted or specific to a particular geo-climatic zone of




Fig. 1. *Varanus bengalensis* preying on an (a) Indian tiger centipede (*Scolopendra hardwickei*), (b) juvenile black-napped hare (*Lepus nigricollis*), (c) adult common garden lizard (*Calotes versicolor*), (d) adult variegated kukri snake (*Oligodon taeniolata*), (e) flying termites, (f) household trash, and (g) adult Asian common toad (*Duttaphrynus melanostictus*). Photographs by Craig Moore, Mevan Piyasena, Senehas Karunaratna, Nilupul Rangana, and Bushana Kalhara.

Table 1. Currently known prey items of *Varanus bengalensis* in Sri Lanka and their current status in Sri Lanka in terms of IUCN national conservation status, endemism, and residential (native vs introduced) status. Abbreviations used: E = Endemic; NA = not applicable; NV = native; NN = non-native; VU = Vulnerable; EN = Endangered; CR = Critically Endangered. Non-native species include domesticated species (domestic cats and dogs), naturalized alien species, and invasive species.

Prey Class	Prey taxon	Common name	Species Status	Description of Prey	Geo-climatic Zone	Location (Habitat)	References
Mammalia	<i>Bandicota bengalensis</i>	Mole rat	NV	Freshly killed by home owner	Lowland wet zone	Ratmalana (Urban)	This study
	<i>Bandicota indica</i>	Malabar bandicoot	NV	Road kill	Lowland wet zone	Kelaniya (Residential)	This study
	<i>Bubolbus bubalis</i>	Buffalo	NV	Scavenged from slaughter houses and carcasses	Lowland wet zone	Aluthgama (Residential)	M. Wickramasinghe, pers. comm.
	<i>Canis familiaris</i>	Domestic dog	NN	Road kill	Lowland wet zone, Lowland arid zone	Beruwala (Urban), Hambantota (Forest)	M. Wickramasinghe, pers. comm.
	<i>Crocidura sp.</i>	Shrew sp.	NV	Live adult	Lowland wet zone	Kudawa (Residential)	This study
	<i>Felis catus</i>	Domestic cat	NN	Live juveniles, road-killed carcass	Lowland wet zone	Kalutara (Forest)	M. Wickramasinghe, pers. comm.
	<i>Funambulus palmarum</i>	Palm squirrel	NV	Juveniles hunted from the nest, road kill	Lowland intermediate zone, Lowland dry zone, Upland wet zone	Hambegamuwa (Residential), Yala (Forest), Thammimalai (Residential)	S. Kumara, M. Karunarathna & T. Kusuminda, pers. comm.
	<i>Kerivoula picta</i>	Painted bat	NV	Road kill	Lowland dry zone	Polonnaruwa (Residential)	This study
	<i>Lepus nigricollis</i>	Black-napped hare	NV	Live subadult	Lowland dry zone	Yala (Forest)	M. Pyasena, pers. comm.
	<i>Macaaca sinica</i>	Sri Lankan toque monkey	E	Juvenile carcass	Lowland wet zone	Dombagaskanda (Forest)	This study
Aves	<i>Manis crassicaudata</i>	Pangolin	NV	Adult carcass	Lowland dry zone	Gaioya (Forest)	D. Kulathunga, pers. comm.
	<i>Mus booduga</i>	Field mouse	NV	Freshly killed by homeowner	Lowland intermediate zone	Tanamalwila (Residential)	S. Kumara, pers. comm.
	<i>Mus musculus</i>	Indian house mouse	NV	Freshly killed by homeowner	Lowland wet zone	Ambalangoda (Urban)	This study
	<i>Pteropus giganteus</i>	Flying fox	NV	Electrocuted carcass	Lowland wet zone	Piliyandala (Urban)	This study
	<i>Rattus rattus</i>	Black rat	NN	Freshly killed by homeowner	Lowland wet zone	Ratmalana (Urban)	This study
	<i>Rusa unicolor</i>	Sambar deer	NV	Adult carcass	Lowland wet zone	Sinharaja (Forest)	This study
	<i>Semnopithecus priam</i>	Grey langur	NV	Adult carcass	Lowland dry zone	Wilpattu (Forest)	D. Ramasinghe, pers. comm.
	<i>Semnopithecus vetulus</i>	Sri Lankan Purple-faced langur	E/EN	Electrocuted carcass	Lowland wet zone	Awissawella (Urban)	S. Kolambage, pers. comm.
	<i>Suncus murinus</i>	Common musk shrew	NV	Adult carcass	Lowland wet zone	Ganemulla (Urban)	This study
	<i>Tandeleuria oleracea</i>	Long-tailed tree mouse	VU	Live adult	Lowland dry zone	Karawalagaswewa (Forest)	This study
	<i>Acridotheres tristis</i>	Common myna	NV	Nestlings	Lowland wet zone	Ganemulla (Urban)	This study
	<i>Amaurornis phoeniceus</i>	White-breasted waterhen	NV	Road kill	Lowland dry zone	Tissamaharamaya (Residential)	This study
	<i>Chalcophaps indica</i>	Emerald dove	NV	Adult carcass	Lowland wet zone	Aluthgama (Residential)	This study
	<i>Columba livia</i>	Rock pigeon	NV	Adult carcass	Lowland wet zone	Ratmalana (Urban)	This study
	<i>Copsychus saularis</i>	Oriental magpie robin	NV	Eggs	Lowland intermediate zone	Bibila (Residential)	This study
	<i>Dinopium benghalense</i>	Black-rumped flameback	NV	Hatchlings from the nest	Lowland dry zone	Polonnaruwa (Residential)	L. Dayaratne, pers. comm.
	<i>Eudynamis scolopaceus</i>	Asian koel	NV	Juvenile carcass	Lowland wet zone	Kadawatha (Urban)	R. Saranga, pers. comm.
	<i>Francolinus pictus</i>	Painted francolin	CR	Leftover meat discarded by hunters	Lowland intermediate zone	Nilgala (Forest)	D. Kulathunga, pers. comm.
<i>Halcyon smyrnensis</i>	White-throated kingfisher	NV	Hatchlings from the nest	Lowland wet zone	Galle (Residential)	M. Dahanayake, pers. comm.	
<i>Megalaima flavifrons</i>	Sri Lankan yellow-fronted barbet	E	Live chicks from the nest	Lowland wet zone	Dombagaskanda (Forest)	This study	
<i>Megalaima zeylanica</i>	Brown-headed barbet	NV	Live chicks from the nest	Lowland dry zone	Polonnaruwa (Forest)	L. Dayaratne, pers. comm.	
<i>Perdix asiatica</i>	Jungle bush-quail	CR	Trash body parts discarded by hunters	Lowland intermediate zone	Mahiyanganaya (Forest)	D. Kulathunga, pers. comm.	
<i>Psittacula calthropae</i>	Sri Lankan emerald collared parakeet	E	Dislodged live hatchlings and eggs	Lowland wet zone	Deraniyagala (Forest)	M. Wickramasinghe, pers. comm.	
<i>Psittacula krameri</i>	Rose-ringed parakeet	NV	Dislodged live hatchlings and eggs	Lowland wet zone	Mideripitiya (Forest)	M. Wickramasinghe, pers. comm.	
<i>Pycnonotus cafer</i>	Red vented bulbul	NV	Dislodged hatchlings	Lowland wet zone	Matara (Residential)	N. Rangana, pers. comm.	
<i>Pycnonotus luteolus</i>	White browed bulbul	NV	Road kill carcass	Lowland wet zone	Ingiriya (Residential)	This study	
<i>Saxicoloides fulvicaus</i>	Indian robin	NV	Eggs and hatchlings	Lowland dry zone	Rajanganaya (Forest)	D. Ramasinghe, pers. comm.	
<i>Stigmatopelia chinensis</i>	Spotted dove	NV	Adult carcass	Lowland dry zone	Chilaw (Urban)	A. Kumasinghe, pers. comm.	
<i>Turdoides affinis</i>	Yellow billed babbler	NV	Injured adult	Lowland wet zone	Panadura (Urban)	This study	
<i>Turdix suscitator</i>	Barred buttonquail	NV	Trash body parts discarded by hunters	Lowland dry zone	Giradurukotte (Forest)	D. Kulathunga, pers. comm.	
<i>Vanellus indicus</i>	Red-wattled lapwing	NV	Live fledglings and eggs from the nest	Lowland wet zone	Seenigama (Residential)	S. de Soya, pers. comm.	

Table 1 (continued).

Prey Class	Prey taxon	Common name	Species Status	Description of Prey	Geo-climatic Zone	Location (Habitat)	References	
Reptilia	<i>Amphisma stollatum</i>	Buff striped keelback	NV	Live adult, Road kill	Lowland intermediate zone, Lowland dry intermediate zone, Lowland wet zone	Thanamalwila (Residential), Kumana (Forest), Waga (Residential)	This study, G. Jayantha & M. Wickramasinghe, pers. comm.	
		Green garden lizard	NV	Live adults	Lowland dry zone, Lowland wet zone	Anuradapura (Residential), Maharagama (Urban)	This study	
	<i>Calotes calotes</i>	Common garden lizard	NV	Live adult, road-killed specimen	Lowland dry zone, Lowland arid zone	Mihintale (Forest), Yala (Forest)	This study, S. Karunarathna, pers. comm.	
		Sri Lankan chameleon	EN	Live adults	Lowland dry zone	Vanatavilluwa (Residential)	Karunarathna et al., 2009	
		Mugger crocodile	NV	Carcass	Lowland dry zone	Kanadara (Residential)	This study	
		Russell's viper	NV	Live juvenile	Lowland intermediate zone	Udawalawe (Forest)	S. Kumara, pers. comm.	
		Common skink	NV	Live adults	Lowland intermediate zone	Hambugamuwa (Residential)	This study	
		Star tortoise	NV	Live juvenile	Lowland dry zone	Eluwankulam (Residential)	D. Ramasinghe, pers. comm.	
		Common house-gecko	NV	Live adults	Lowland wet zone	Ganemulla (Residential)	This study, M. Wickramasinghe, pers. comm.	
		<i>Hemidactylus parvimaaculatus</i>	Spotted housegecko	NV	Live adults	Lowland wet zone	Ganemulla (Residential)	This study
			Merrem's hump nose viper	NV	Killed by villagers	Lowland wet zone	Nitambuwu (Residential)	This study
			Blind snake species	NV	Live adults	Lowland wet zone	Ganemulla (Residential)	This study
	Wolf snake		NV	Live adults	Lowland dry zone	Gampaha (Urban)	This study	
	Black turtle		NV	Live juvenile	Lowland wet zone	Maharagama (Urban)	This study	
	Common kukri snake		NV	Road kill	Lowland dry zone	Padaviya (Forest)	This study	
	Variagated kukri snake		NV	Live adults	Lowland dry zone	Yala (Forest)	This study	
	Black spotted kangaroo lizard		E	Live adults	Lowland intermediate zone	Niigala (Forest)	This study	
	<i>Ptyas mucosa</i>	Rat snake	NV	Live juvenile	Lowland wet zone	Ragama (Urban)	This study	
		Devaka's fanthroat lizard	E	Live adults	Lowland dry zone	Vanatavilluwa (Residential)	This study	
		Checked keelback	NV	Live adults	Lowland wet zone	Dediya (Forest)	M. Wickramasinghe, pers. comm.	
			NV	Live adults	Lowland wet zone	Matarra (Residential)	This study	
	Amphibia	<i>Duttaphrynus atukoralei</i>	Aukoral's toad	E	Live adults	Lowland wet zone	Kalutara (Residential), Polgasovita (Urban)	This study, B. Kalhara & T. Ramasinghe, pers. comm.
			Common house toad	NV	Live adult, road kill	Lowland wet zone	Polgasovita (Urban)	This study
<i>Duttaphrynus scaber</i>		Schneider's toad	NV	Live adults	Lowland dry zone	Jaffna (Residential)	This study	
		Indian skipper frog	NV	Live adults	Lowland intermediate zone	Hambugamuwa (Forest)	This study	
		Indian green frog	NV	Live adults	Lowland intermediate zone	Hambugamuwa (Residential)	This study	
		Common paddy field frog	NV	Live adults	Lowland intermediate zone	Hettipola (Residential)	Jolley & Meek, 2006	
		Jurdon's bullfrog	NV	Live adults	Lowland dry zone	Yala (Forest)	S. Roelofs, pers. comm.	
		Sri Lankan bullfrog	NV	Live adult, road kill	Lowland dry zone, Lowland wet zone	Puttalam (Urban), Dambulla (Residential), Matugama (Forest)	This study, M. Wickramasinghe, pers. comm.	
<i>Polypedates maculatus</i>		Spotted tree frog	NV	Live adults, road kill	Lowland intermediate zone, Lowland arid zone	Monaragala (Forest), Hambantota (Residential)	This study, M. Wickramasinghe, pers. comm.	
		Common shrub frog	E	Live adults	Lowland wet zone	Ganemulla (Urban)	This study	
<i>Pseudophilautus popularis</i>		Variagated ramanella	NV	Live adults	Lowland arid zone	Hambantota (Residential)	M. Wickramasinghe, pers. comm.	
		Short-headed burrowing frog	NV	Live adults	Lowland dry zone	Yala (Forest), Puttalam (Urban), Hambantota (Residential)	This study, M. Wickramasinghe, pers. comm.	
<i>Sphaerotheca breviceps</i>	Roland's burrowing frog	NV	Live adults	Lowland dry zone, Lowland arid zone	Karawalagasweva (Forest), Mannar (Urban)	This study, M. Wickramasinghe, pers. comm.		
	Marbled balloon frog	NV	Live adults	Lowland dry zone	Yala (Forest), Anuradapura (Residential), Wilpattu (Forest)	This study, M. Wickramasinghe, pers. comm.		
Pisces	<i>Davkinsia singhala</i>	Sri Lankan filament Barb	E	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
		Giant danio	NV	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
	<i>Devario malabaricus</i>	Sri Lankan flying barb	E	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
		Green chromide	NV	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
	<i>Epiplatys suratisensis</i>	Tilapia	NN	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
		Long-snouted barb	NV	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
	<i>Oreochromis mossambicus</i>	Broad line Strip rasbora	NV	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	
			NV	Live specimens trapped in small pools	Lowland dry zone	Udawalawe (Forest)	S. Atapattu, pers. comm.	

Table 1 (continued).

Prey Class	Prey taxon	Common name	Species Status	Description of Prey	Geo-climatic Zone	Location (Habitat)	References
Arachnida	<i>Buthoscorpio sarasinorum</i>	Scorpion	NV	Live adults	Lowland dry zone	Eluwankulam (Residential)	A. Kumara, pers. comm.
	<i>Heierometrus gravimanus</i>	Scorpion	NV	Live adults	Lowland wet zone	Amalangoda (Residential), Akuressa (Residential)	This study, S. Karunaratna, pers. comm.
Chilopoda	<i>Heterometrus serratus</i>	Scorpion	NV	Live adults and juveniles	Lowland intermediate zone	Nilgala (Forest)	This study
	<i>Reddyanus jayarathnei</i>	Scorpion	NV	Live adults	Lowland wet zone	Kanneliya (Forest)	S. Akmeemana, pers. comm.
	<i>Reddyanus loebli</i>	Scorpion	NV	Live adults	Lowland dry zone	Sangamankande (Residential)	This study
	<i>Rhyssida</i> sp.	Common centipede	NV	Live adults	Lowland dry zone	Bibila (Urban)	This study
	<i>Scelopendra harawicki</i>	Indian tiger centipede	NV	Live adults	Lowland dry zone	Yala (Forest)	C. Moore, pers. comm.
Gastropoda	<i>Cryptozona biserialis</i>	Common translucent snail	NV	Live adults	Lowland wet zone	Ganemulla (Urban)	This study
	<i>Lissachatina fulica</i>	Giant African snail	NN	Live adults	Lowland wet zone	Ganemulla (Urban), Gampaha (Urban)	This study, T. Ramasinghe, pers. comm.
Malacostraca	<i>Oziohelphusa</i> sp.	Freshwater crab	E	Live adults	Lowland dry zone	Mihintale (Forest)	This study
	<i>Oziohelphusa hippocastanum</i>	Freshwater crab	E/EN	Live adults, road kill	Lowland dry zone	Jaffna (Residential)	This study
	<i>Perbrinckia</i> sp.	Freshwater crab	E	Live adults	Lowland intermediate zone	Nilgala (Forest), Pokunntenna (Forest)	This study, M. Wickramasinghe, pers. comm.
Insecta	Suborder: Coelifera	Grasshoppers	NV	Live adults	Lowland intermediate zone, Upland wet zone	Sigiriya (Residential), Peradeniya (Urban)	This study
	Family: Scarabaeidae	Dung beetles	NV	Live adults	Lowland dry zone	Mihintale (Forest), Udawalawe (Forest)	This study, M. Wickramasinghe, pers. comm.
	Infraorder: Isoptera	Termites	NV	Live adults, flying and larvae	Lowland dry zone, Lowland dry zone	Mihintale (Forest), Suriyawewa (Residential), Sigiriya (Residential), Dambulla (Forest)	N. Rangana & M. Rathnayake, pers. comm.
	Subfamily: Formicidae Order: Coleoptera	Black ants Beetle species	NV NV	Live adults	Lowland dry zone Lowland wet zone, Lowland dry zone	Mihintale (Forest) Moratuwa (Urban), Yala (Forest)	This study A. Godahewa & M. Wickramasinghe, pers. comm.
Suborder: Oniscidea Order: Blattodea	Family: Gryllidae	Crickets species	NV	Live adults and juveniles	Lowland wet zone, Lowland dry zone	Ganemulla (Urban), Hambegamuwa (Residential)	This study
	Suborder: Oniscidea	Doodlebug	NV	Live adults and larvae	Lowland dry zone, Lowland wet zone	Habarana (Residential), Galle (Residential)	This study, T. Ramasinghe, pers. comm.
	Order: Blattodea	Wild cockroaches	NV	Live specimens	Lowland wet zone	Aluthigama (Urban), Nittambuwa (Residential)	This study
	<i>Heteropoda venatoria</i> <i>Gryllotalpa</i> sp.	Huntsman spider Mole cricket	NV NV	Live specimens Live specimens	Lowland wet zone Lowland intermediate zone, Lowland dry zone	Elpitiya (Forest) Nilgala (Forest), Ampara (Urban)	This study D. Kulathunga, pers. comm.
Oligochaeta	Suborder Megadriacea	Earthworm	Live specimens	Lowland wet zone	Homagama (Urban), Ratmalana (Urban)	This study	
Garbage	n/a	Cooked anchovy	NA	Household trash	Lowland dry zone	Higurakgoda (Residential)	This study
	n/a	Cooked chicken	NA	Household trash	Lowland wet zone	Kuruwita (Forest)	This study
	n/a	Cooked shrimps	NA	Household trash	Lowland dry zone	Yala (Forest)	C. Amarasinghe, pers. comm.
	n/a	Fruits	NA	Road-killed specimen (from inside the stomach)	Lowland dry zone	Trincomalee (Urban)	M. Rathnayake, pers. comm.

Table 2. Currently known predators of *V. bengalensis*, in Sri Lanka. Abbreviations used: E = Endemic; NV = native; NN = non-native; EN = non-native; EN = Endangered.

Predatory Class	Predator Taxon	Common name	Species Status	Prey Description	Location	References
Mammalia	<i>Homo sapiens</i>	Human	NV	Hunting on juveniles subadult, and adult stages, cats tongue and heart, skin use for drums, fat to make an oil	Lowland wet zone: Galle; Lowland intermediate zone: Kurunegala, Nilgala; Lowland dry zone: Jaffna	This study; Deraniyagala (1953), de Silva (1996)
	<i>Panthera pardus</i>	Leopard	EN	Live subadult	Lowland dry zone: Yala, Wilpattu	D. Ramasinghe & M. Piyasena, pers. comm.
	<i>Canis familiaris</i>	Domestic dog	NN	Live adult	Lowland wet zone: Raddolugama, Ambalangoda	This study; S. Atapattu pers. com.
	<i>Felis catus</i>	Domestic cat	NN	Live juvenile	Lowland wet zone: Rambukkana	This study
	<i>Canis aureus</i>	Jackal	NV	Carcass	Lowland intermediate zone: Illukkumbura; Lowland wet zone: Mirrigama	This study
	<i>Herpestes brachyurus</i>	Brown mongoose	NV	Live juvenile	Lowland wet zone: Galle	M. Dahanayake, pers. comm.
	<i>Herpestes edwardsii</i>	Grey mongoose	NV	Live juveniles	Lowland wet zone: Ganemulla	This study
	<i>Herpestes smithii</i>	Ruddy mongoose	NV	Live juveniles	Lowland dry zone: Yala	M. Piyasena & S. Karunaratna, pers. comm.
	<i>Sus scrofa</i>	Wild boar	NV	Road kill	Lowland dry zone: Wilpattu, Galoya	This study; D. Kulathunga, pers. comm.
Aves	<i>Ocyerous gingalensis</i>	Sri Lankan grey hornbill	E	Live juveniles	Lowland dry zone: Mahiyangamaya	D. Kulathunga, pers. comm.
	<i>Pelargopsis capensis</i>	Stork-billed kingfisher	NV	Live juveniles	Lowland dry zone: Chilaw	A. Kumarasinghe, pers. comm.
	<i>Centropus sinensis</i>	Greater coucal	NV	Live juveniles	Lowland dry zone: Pannala	This study
	<i>Spilornis cheela</i>	Crested serpent-eagle	NV	Live juveniles	Lowland dry zone: Yala, Giritale; Lowland intermediate zone: Udawala we	This study; C. Amarasinghe, pers. comm.
	<i>Accipiter badius</i>	Shikra	NV	Live juveniles	Lowland wet zone: Panadura, Kaluara	S. Dayananda, pers. comm.
	<i>Spizaetus cirrhatus</i>	Changeable hawk-eagle	NV	Live adult	Lowland dry zone: Yala	S. Karunaratna, pers. comm.
	<i>Mycteria leucocephala</i>	Painted stork	NV	Live juvenile	Lowland wet zone: Maharagama	K. Buddika pers. comm.
	<i>Leptoptilos javanicus</i>	Lesser adjutant	NV	Live juvenile	Lowland dry zone: Pokunutenna	N. Perera, pers. comm.
	<i>Corvus splendens</i>	House Crow	NV	Road kill	Lowland intermediate zone: Nilgala, Thanamalwila;	This study
	<i>Corvus levaillantii</i>	Jungle crow	NV	Road kill	Lowland dry zone: Eluwankulam	This study
	<i>Crocodylus palustris</i>	Mugger crocodile	NV	Live adult	Lowland intermediate zone: Thanamalwila; Lowland dry zone: Nilgala	This study
Reptilia	<i>Ptyas mucosa</i>	Rat snake	NV	Live juveniles	Lowland dry zone: Wilpattu; Lowland dry zone: Yala	D. Ramasinghe & N. Kamalgoda, pers. com.
	<i>Python molurus</i>	Indian python	NV	Live adult	Lowland intermediate zone: Nilgala, Hambegamuwa	This study, S. Kumara, pers. comm.
	<i>Naja naja</i>	Indian cobra	NV	Live juveniles	zone: Yala, Wilpattu; Upland wet zone: Kundsale	D. Jayantha, N. Kamalgoda & S. Karunaratna, pers. comm.
	<i>Varanus salvator</i>	Water monitor	NV	Road kill	Lowland intermediate zone: Bulupitiya; Lowland dry zone: Polonnaruwa	This study; D. Kulathunga, pers. comm.
					Lowland wet zone: Kandana	This study




Fig. 2. Predators of *V. bengalensis*: (a) Sri Lankan leopard (*Panthera pardus*), (b) domestic dog attack (*Canis familiaris*), (c) changeable hawk-eagle (*Spizaetus cirrhatus*), (d) crested serpent-eagle (*Spilornis cheela*), (e & f) grey mongoose (*Herpestes edwardsii*), (g) local drum using the skin of *V. bengalensis*. Photographs by Mevan Piyasena, Sanjaya Atapattu, Mendis Wickramasinghe, and Dinesh Silva.

Sri Lanka or habitat type.

Discussion

Our study confirmed that *V. bengalensis* is a widely-foraging generalist predator that can also function as an opportunistic scavenger (Auffenberg, 1994; Pianka, 2004). Broad dietary niches have been reported for several other varanids throughout the Indo-Malayan and Australasian realms (Sutherland, 2011). In Sri Lanka, *V. bengalensis* seems to forage mostly as an active predator, and most of its prey are vertebrates. In

contrast, studies conducted elsewhere have suggested that invertebrates (flying insects, annelids, arachnids, mollusks, and crustaceans) comprised the primary diet, while vertebrates were an alternative prey depending on their availability (Auffenberg, 1994; Pianka, 2004). The dietary bias we noted in our study can be an artifact of biased sampling and detectability since predation on vertebrates is more noticeable to an opportunistic observer. *Varanus bengalensis* is also considered an egg predator of both reptiles and birds (Deraniyagala, 1953; Somaweera & Somaweera, 2009); however, our study only confirmed *V. bengalensis* predation of bird

eggs in Sri Lanka. Dietary studies on varanids have suggested a strong positive correlation between prey size and the monitor's body size, with large-bodied monitors foraging predominantly on vertebrate prey such as mammals while smaller-sized monitors practice insectivory (Arbuckle, 2009). Gut content analyses of *V. bengalensis* have supported this notion, where juvenile monitors feed nearly exclusively on insects such as orthopterans and coleopterans (Auffenberg & Ipe, 1983). However, vertebrate prey may account for a greater caloric (energetic) intake by monitors irrespective of the relative proportion of vertebrates in monitor diets (Arbuckle, 2009).

Plasticity in food selection and the catholic dietary habits of *V. bengalensis* have previously been documented from southeastern and southern Asian mainland, Indo-Malayan island, and Indian oceanic island populations (Jolley & Meek, 2006). Although predominantly terrestrial, a minor proportion of the diet of *V. bengalensis* consists of both aquatic and semi-aquatic prey (freshwater fish, amphibians, waterfowl, crocodiles, and freshwater crabs) in aquatic and riparian environments (S. Karunarathna, pers. obs.). Therefore, *V. bengalensis*, like many other varanids, may even shift its prey choice in response to seasonal and spatial variation in food availability, ontogeny, and intensity of competitive interactions (Shine, 1986; Losos & Greene, 1988; Sutherland, 2011). In Sri Lanka, the diversity of large lizards (only two species of *Varanus*) is lower than other regions within the natural range of varanids such as the Indo-Malayan region; thus, inter-specific competition is lower, which may have enabled *V. bengalensis* to retain a generalist diet. The other extant Sri Lankan monitor species, *V. salvator*, is predominantly aquatic, which further minimizes niche overlap with *V. bengalensis* (Karunarathna *et al.*, 2012).

Although *V. bengalensis* is mostly terrestrial, we have documented that the species can forage in arboreal habitats, where individuals feed on birds (common myna [*Acridotheres tristis*]; Jolley & Meek, 2006). Sometimes, *V. bengalensis* forages in vegetation alongside river banks or at the land-water interphase of rivers, lakes, and wetlands. When feeding on larger vertebrate prey, these monitors are known to kill their prey by violently shaking it or slamming it against hard substrates, then swallow the prey head first by pushing the prey against the ground and then using inertial movements until ingestion is complete (Loop, 1974; Jolley & Meek, 2006; Rahman *et al.*, 2015). When handling or capturing larger prey, these monitors can dismember their prey by holding them with forelimbs;

they are also known to dig in search of prey with their forelimbs and jaws (Auffenberg & Ipe, 1983; Rahman *et al.*, 2015). Moreover, when feeding on invertebrates with hard exoskeletons, the monitors masticate the shells first prior to ingestion (Jolley & Meek, 2006). We noted that *V. bengalensis* forages in multiple habitat types, including forests and woodlands with trees of variable trunk densities, home gardens and homesteads, urban and suburban environments, lotic and lentic aquatic habitats of both natural and anthropogenic origins, and brackish water habitats.

Our study indicates that *V. bengalensis* is capable of feeding on live prey as well as dead organic matter, including both native and introduced species. Carrion feeding has been observed throughout its range (Karunarathna *et al.*, 2012). Populations inhabiting homestead environments are known to spend nearly 50% of their activity budget on scavenging among household garbage, particularly kitchen trash where they feed on cooked and uncooked vegetables, grains, and fruits (Rahman *et al.*, 2015). The ability of several varanid species to scavenge human-generated trash is a remarkable feature which has enabled them to successfully exploit built-up environments and as a human commensal (Uyeda, 2009). Human-assisted food subsidies for varanids could have profound ecological effects on wild populations, such as increased abundance, altered community and intraspecific interactions, modified social hierarchies, and aggravated human-wildlife conflicts. Dramatic changes in wildlife behavior and community structure have been recognized among many other human commensals such as coyotes, crows, raccoons, mongoose, and gulls upon "supplementary feeding" (O'Connor, 2013).

Our study also established the fact that *V. bengalensis* may not be a top predator in Sri Lanka's ecosystems, although we only documented a few predators of *V. bengalensis*. Throughout its biogeographic range, a number of predators including pythons and other large snakes, eagles, mongooses, wild and domesticated dogs, feral cats, humans, and other varanids are known to predate on *V. bengalensis* (Auffenberg, 1994; Pianka, 2004). Similar to our findings in Sri Lanka, predation mostly occurs in early life-history stages (eggs, hatchlings, and juveniles), with only a small portion of predation involving fully-grown adults (Auffenberg, 1994; Pianka, 2004). Although cannibalism has been cited elsewhere, we do not have any evidence for active intraspecific predation of *V. bengalensis* in Sri Lanka (Auffenberg, 1994; Pianka, 2004).

Humans' role as a predator of *V. bengalensis* in

Sri Lanka is noteworthy. The human exploitation of *V. bengalensis* for food in Sri Lanka has historically been recorded as far back as 800 BC - 200 AD, and continues today to a greater extent and includes both consumption and trade (Abayaratna & Mahaulpatha, 2006; de Silva, 2006). Human exploitation of *V. bengalensis* has also been reported from South and Southeast Asia (Koch *et al.*, 2013). The hunting pressure on *V. bengalensis* in Sri Lanka as well as in some other parts of its range have led to localized population declines and reductions in their former ranges and habitats (Amarasinghe *et al.*, 2009; Koch *et al.*, 2013). In some rural parts of Sri Lanka such as Hambegamuwa (southeastern Sri Lanka), local inhabitants use the hides of *V. bengalensis* for making drums (Fig. 2).

Given its broad dietary preferences, which include many insects and rats, *V. bengalensis* can be considered a biological pest control agent (Karunarathna *et al.*, 2008; Karunarathna *et al.*, 2012). Moreover, as scavengers, these monitors contribute to organic matter cycling and the removal of carrion (Somaweera & Somaweera, 2009; Karunarathna *et al.*, 2012). Their role as a mesopredator provisions many ecosystem services which involves regulating populations in lower trophic levels while also serving as prey for top predators. Behavioral studies have suggested that *V. bengalensis* allocate a greater portion of their daily activity budgets to foraging, where they roam extensively over great distances exploring profitable foraging grounds (Losos & Greene, 1988; Sweet & Pianka, 2007). Thus, their contribution to energy flow and nutrient cycling is salient for proper ecosystem functioning. Quantifying their role in food webs based on gut content surveys and carbon isotope analyses using network theory (carbon and energy flow across different trophic nodes) may provide considerable insight on their ecological role. Moreover, our study has indicated that *V. bengalensis* feeds on non-native species; therefore, these monitors could have some impacts on controlling invasive species. This potential requires further investigation.

Acknowledgements - We would like to thank Tharindu Ranasinghe, Avishka Godahewa, Chamara Amarasinghe, Nilupul Rangana, Craig Moore, Bushana Kalhara, Mevan Piyasena, Stuart Roelofs, Senehas Karunarathna, Sameera de Soyza, Dammika Ramasinghe, Dissanayake Kulathunga, Sampath Kolombage, Samantha Kumara, Lele Dayarathne, Malaka Dahanayake, Namal Kamalgoda, Rasika Saranga, Anushka Kumarasinghe, Dinesh Silva, Gamini Jayantha, Sameera Akmeemana, Sanjaya Atapattu,

Sanjaya Karunarathna, Tharaka Kusuminda, Mendis Wickramasinghe, Manoj Rathnayake, Mathesha Karunarathna, and Dinesh Silva for valuable discussions, information and photographs. Finally, we thank an anonymous reviewer for constructive comments on the manuscript.

References

- Abayaratna, M.G.T.H. & W.A.D. Mahaulpatha. 2006. Activity budgets and habitat preference of land monitor, thalagoya *Varanus bengalensis* in a residential area. *Vidyodaya Journal of Science* 13: 127–138.
- Amarasinghe, A.A.T., G. Chathuranga & D.M.S.S. Karunarathna. 2009. *Varanus salvator* (Laurenti, 1768) in Rathgama lagoon in Galle District, Sri Lanka. *Biawak* 3(3): 81–84.
- Arbuckle, K. 2009. Ecological function of venom in *Varanus*, with a compilation of dietary records from the literature. *Biawak* 3(2): 46–56.
- Auffenberg, W. 1994. *The Bengal monitor*. University Press of Florida, Gainesville. 590 pp.
- Auffenberg, W. & I. Ipe. 1983. The food and feeding of juvenile Bengal monitor lizards (*Varanus bengalensis*). *Journal of the Bombay Natural History Society* 80(1): 119–124.
- Das, I. & A. de Silva 2005. *A Photographic Guide to Snakes and Other Reptiles of Sri Lanka*. New Holland Publishers, London. 144 pp.
- de Silva, A. 2006. Current status of the reptiles of Sri Lanka. Pp. 134–163 *In: Bambaradeniya, C.N.B. (ed.), Fauna of Sri Lanka: Status of Taxonomy, Research and Conservation*. The World Conservation Union and Government of Sri Lanka, Colombo, Sri Lanka.
- Deraniyagala, P. 1953. *A Coloured Atlas of Some Vertebrates from Ceylon. Tetrapod Reptilia, Volume 2*, National Museums of Sri Lanka. The Ceylon Government Press, Colombo. 102 pp.
- Dryden, G., B. Green, E.D. Wikramanayake & K. Dryden. 1992. Energy and water turnover in two tropical varanid lizards, *Varanus bengalensis* and *V. salvator*. *Copeia* 1992: 102–107.
- Faeth, S.H., P.S. Warren, E. Shochat & W.A. Marussich. 2005. Trophic dynamics in urban communities. *Bioscience* 55(5): 399–407.
- Jolley, E. & R. Meek. 2006. *Varanus bengalensis* (Bengal monitor): Unusual behaviour and feeding. *Herpetological Bulletin* 95: 31–32.
- Karunarathna, D.M.S.S, A.A.T. Amarasinghe &

- A. De Vos. 2008. Preliminary notes on the monitor lizards (Family: Varanidae) within the National Zoological Gardens (NZG) Dehiwala, Colombo District, Sri Lanka. *Biawak* 2(3): 109–118.
- Karunaratna, D.M.S.S., A.A.T. Amarasinghe, B.M. Majintha & H.K.D. Kandambi. 2012. Population status of two *Varanus* species (Reptilia: Sauria: Varanidae) in Sri Lanka's Puttalam lagoon system, with notes on their diet and conservation status. *Biawak* 6(1): 22–33.
- Karunaratna, D.M.S.S., T.D. Surasinghe, M.C. De Silva, M.B. Madawala, D.E. Gabadage & W.M.S. Botejue, 2015. Dietary habits of *Varanus salvator salvator* in Sri Lanka with a new record of predation on an introduced clown knifefish, *Chitala ornata*. *Herpetological Bulletin* 133: 23–28.
- Koch, A., T. Ziegler, W. Boehme, E. Arida & M. Auliya. 2013. Pressing problems: Distribution, threats, and conservation status of the monitor lizards (Varanidae: *Varanus* spp.) of Southeast Asia and the Indo-Australian Archipelago. *Herpetological Conservation and Biology* 8(3): 1–62.
- Loop, M.S. 1974. The effect of relative prey size on the ingestion behavior of the Bengal monitor, *Varanus bengalensis* (Sauria: Varanidae). *Herpetologica* 30(2): 123–127.
- Losos, J.B. and H.W. Greene. 1988. Ecological and evolutionary implications of diet in monitor lizards. *Biological Journal of the Linnean Society* 35(4): 379–407.
- Molnar, R.E. 2004. The long and honorable history of monitors and their kin. Pp. 10–67. *In*: Pianka, E.R., D.R. King & R.A. King (eds.), *Varanoid Lizards of the World*. Indiana University Press, Bloomington.
- O'Connor, T. 2013. *Animals as Neighbors: The Past and Present of Commensal Animals*. Michigan State University Press, Lansing. 184 pp.
- Pianka, E.R. 1995. Evolution of body size: Varanid lizards as a model system. *The American Naturalist* 146(3): 398–414.
- Pianka, E.R. 2004. *Varanus bengalensis*. Pp. 157–160. *In*: Pianka, E.R., D.R. King & R.A. King (eds.), *Varanoid Lizards of the World*. Indiana University Press, Bloomington.
- Pianka, E.R., D. King & R.A. King. 2004. *Varanoid Lizards of the World*. Indiana University Press, Bloomington. 608 pp.
- Pianka, E.R. and D.R. King. 2004. Introduction. Pp. 3–9. *In*: Pianka, E.R., D.R. King & R.A. King (eds.), *Varanoid Lizards of the World*. Indiana University Press, Bloomington.
- Rahman, K.M., M. Khan & I. Rakhimov. 2015. Scavenging behavior of the bengal monitor (*Varanus bengalensis*) in Jahangirnagar University campus, Bangladesh. *Journal of Scientific Research and Reports* 7(7): 539–550.
- Shine, R. 1986. Food habits, habitats and reproductive biology of four sympatric species of varanid lizards in tropical Australia. *Herpetologica* 42(3): 346–360.
- Survey Department of Sri Lanka. 2012. *The National Atlas of Sri Lanka*. 2nd Edition. Survey Department of Sri Lanka, Colombo, Sri Lanka.
- Sutherland, D.R. 2011. Dietary niche overlap and size partitioning in sympatric varanid lizards. *Herpetologica* 67(2): 146–153.
- Sweet, S.S. & E.R. Pianka. 2007. Monitors, mammals, and wallace's line. Pp. 79–99. *In*: Horn, H.-G., W. Böhme & U. Krebs (eds.), *Advances in Monitor Research III*, Mertensiella 16. Deutsche Gesellschaft für Herpetologie und Terrarienkunde, e.V., Rheinbach.
- Uyeda, L. 2009. Garbage appeal: Relative abundance of water monitor lizards (*Varanus salvator*) correlates with presence of human food leftovers on Tinjil Island, Indonesia. *Biawak* 3(1): 9–17.
- Wikramanayake, E.D. & G.L. Dryden. 1993. Thermal ecology of habitat and microhabitat use by sympatric *Varanus bengalensis* and *V. Salvator* in Sri Lanka. *Copeia* 1993: 709–714.
- Wikramanayake, E.D. & B. Green. 1989. Thermoregulatory influences on the ecology of two sympatric varanids in Sri Lanka. *Biotropica* 21(1): 74–79.