



Site Name: Killarney National Park, Macgillycuddy's Reeks and Caragh River Catchment SAC

Site Code: 000365

This very large site encompasses the mountains, rivers and lakes of the Iveragh Peninsula, and the Paps Mountains which stretch eastward from Killarney towards Millstreet. The majority of the site is in Co. Kerry, with a small portion in Co. Cork. This is the most mountainous region in Ireland and includes Carrauntoohil, the highest peak in the country at 1,039 m. The underlying geology is almost entirely Old Red Sandstone, although Carboniferous limestone occurs on the eastern shores of Lough Leane, and rhyolitic lavas occur above Lough Guitane. The dramatic sandstone ridges and valleys have been shaped by glacial processes and many of the lakes are impounded by glacial moraines. Located close to the Atlantic in the south-west of Ireland, the site is subject to strong oceanic influences. Generally, Lusitanian flora and fauna is well-represented, while the high peaks and cliffs support arctic-alpine relicts.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes).

- [3110] Oligotrophic Waters containing very few minerals
- [3130] Oligotrophic to Mesotrophic Standing Waters
- [3260] Floating River Vegetation
- [4010] Wet Heath
- [4030] Dry Heath
- [4060] Alpine and Subalpine Heaths
- [5130] Juniper Scrub
- [6130] Calaminarian Grassland
- [6410] *Molinia* Meadows
- [7130] Blanket Bogs (Active)*
- [7150] Rhynchosporion Vegetation
- [91A0] Old Oak Woodlands
- [91E0] Alluvial Forests*
- [91J0] Yew Woodlands*

- [1024] Kerry Slug (*Geomalacus maculosus*)
- [1029] Freshwater Pearl Mussel (*Margaritifera margaritifera*)
- [1065] Marsh Fritillary (*Euphydryas aurinia*)
- [1095] Sea Lamprey (*Petromyzon marinus*)

- [1096] Brook Lamprey (*Lampetra planeri*)
- [1099] River Lamprey (*Lampetra fluviatilis*)
- [1103] Twaite Shad (*Alosa fallax*)
- [1106] Atlantic Salmon (*Salmo salar*)
- [1303] Lesser Horseshoe Bat (*Rhinolophus hipposideros*)
- [1355] Otter (*Lutra lutra*)
- [1421] Killarney Fern (*Trichomanes speciosum*)
- [1833] Slender Naiad (*Najas flexilis*)

The Oak woodlands, occurring mostly around the Killarney lakes, are the habitat for which the area is perhaps best known. They form the most extensive area of native woodland remaining in Ireland and include Derrycunihy Wood, described as perhaps the most natural Sessile Oak (*Quercus petraea*) wood in the country. The woods are typically dominated by Sessile Oak, with an understorey of Holly (*Ilex aquifolium*). The Strawberry-tree (*Arbutus unedo*) is a notable component of the woods and there are scattered areas of Yew (*Taxus baccata*). The herb layer is not particularly species-rich, but the woods support perhaps the best developed Atlantic bryophyte community in Europe. Several rare species are present including *Lejeunea flava*, *Cyclodictyon laetivirens*, *Daltonia splachnoides*, *Sematophyllum demissum* and *Radula carringtonii*.

The only sizeable Yew woodland in Ireland is found on the limestone of the Muckross peninsula. Here, some of the trees are up to 200 years old. The dense shade beneath the tree results in few herbs in the ground flora, but the bryophyte layer is well-developed and almost continuous.

Wet woodland, or carr, occurring on the low-lying limestone areas within the floodplain of Lough Leane, forms one of the most extensive areas of this woodland type in Ireland. The dominant canopy species are Alder (*Alnus glutinosa*), willows (*Salix* spp.), Ash (*Fraxinus excelsior*) and Downy Birch (*Betula pubescens*), while the field layer is dominated by Remote Sedge (*Carex remota*) and Creeping Bent (*Agrostis stolonifera*).

Adding to the diversity of the woodland component of this site are a number of mixed woodlands, including those of Ross Island which support one of the richest herb layers of the Killarney woods.

The most common habitat types within the overall site are blanket bog, heath and upland grassland. The heath and grassland generally occur on areas with shallow peat and on the mineral soils of the steep mountain sides, while the blanket bog occurs on the more gentle slopes, plateaux and other level ground. Often the habitats occur in a mosaic, with exposed rock frequently occurring.

A variety of blanket bog types are represented from lowland valley to mountain blanket bog. Some of the best include: Cumeragh River Bog Nature Reserve, a

domed bog which is perhaps the most southern intact blanket bog in the country; Ballygisheen, which contains one of the most extensive areas of intact lowland blanket bog in Co. Kerry; Coomacheo/Caherbarnagh, which combine to form the largest mountain blanket bog in the south-west; Eirk Bog Nature Reserve, a classic example of a bog intermediate between a raised and blanket bog; Mangerton Bog, an upland bog which grades into an unusual lichen heath seen at no other site; and Oolagh East, a quaking basin mire. Generally, the bogs have a characteristic flora. The Lusitanian species, Large-flowered Butterwort (*Pinguicula grandiflora*), is common. The bogs also support a number of unusual species, including mosses (*Sphagnum pulchrum*, *S. fuscum*, *S. platyphyllum*, *S. strictum*, *S. contortum* and *Calliergon stramineum*), liverworts (*Cladopodiella francisci* and *Calypogeia azurea*) and lichens (*Cladonia mediterranea*, *C. macilenta*, *C. rangiferina*, *C. arbuscula* and *Cetraria islandica*).

Rhynchosporion vegetation is confined to wet areas within the lowland blanket bogs, with one of the best areas for the habitat being to the north-east of the Ballygisheen Pass. On a portion of this bog there is an extensive area of quaking flats and pools dominated by the bog mosses *Sphagnum cuspidatum* and *S. auriculatum*. These areas have a typically species-poor flora which includes Bogbean (*Menyanthes trifoliata*), White Beak-sedge (*Rhynchospora alba*), Bog Asphodel (*Narthecium ossifragum*), Common Cottongrass (*Eriophorum angustifolium*) and Great Sundew (*Drosera anglica*). Brown Beak-sedge (*R. fusca*), a locally rare plant of wet bog pools, is occasional within the site. Although the habitat is best developed in very wet areas of intact bog, it may also occur in wet areas of regenerating cutover blanket bog.

Wet heath often occurs in association with blanket bog and features Cross-leaved Heath (*Erica tetralix*). Dry heath is more frequent in this site, and is dominated by Heather (*Calluna vulgaris*), Bell Heather (*Erica cinerea*) and Western Gorse (*Ulex gallii*), with occasional Bilberry (*Vaccinium myrtillus*). This habitat is well-developed on the Paps Mountains. Elsewhere it is often over-grazed, with upland grassland becoming more frequent. Some of the highest ridges support alpine heath (referable to the *Lycopodium alpinum* - *Racomitrium lanuginosum* association). Widespread plant species of the alpine heath include Bog-myrtle (*Vaccinium myrtillus*), Crowberry (*Empetrum nigrum*) and Fir Clubmoss (*Huperzia selago*), while species such as Juniper (*Juniperus communis* subsp. *nana*) and Dwarf Willow (*Salix herbacea*) have a much more restricted distribution.

The site contains many lakes, but these can be broadly divided into two types: small upland corrie lakes and larger lowland lakes. Examples of the first type are Lough Murtagh and Lough Gortavehy in the Paps Mountains. They are oligotrophic and typically species-poor, with Quillwort (*Isoetes lacustris*), Water Lobelia (*Lobelia dortmanna*) and Shoreweed (*Littorella uniflora*) occurring most commonly. The lowland lakes are mostly oligotrophic, although Lough Leane, the largest freshwater body in the region, has become somewhat mesotrophic as a result of pollution from Killarney town. These lowland lakes tend to be more species-rich than those at higher altitudes, with additional species such as Awlwort (*Subularia aquatica*), Six-stamened Waterwort (*Elatine hexandra*) and Alternate Water-milfoil (*Myriophyllum*

alterniflorum). Good examples include Lough Caragh, Upper Lake and Muckcross Lake.

The rivers associated with these lakes are also of importance. The Caragh is relatively unpolluted from headwater to estuary, a rare phenomenon in Europe. The Flesk runs over Old Red Sandstone in its upper reaches and limestone as it nears Lough Leane. Both rivers support floating and submerged vegetation and rare invertebrates. Rocks around the smaller mountain streams often support a lush vegetation of ferns and bryophytes, most notably at Torc Waterfall.

Other habitats of note include: Juniper scrub found on islands in the Upper Lake and on dry ridges in nearby Newfoundland Bog; damp meadows, with Purple Moor-grass (*Molinia caerulea*), supporting scarce species such as Whorled Caraway (*Carum verticillatum*) and Ivy-leaved Bellflower (*Wahlenbergia hederacea*); and Calaminarian grasslands, associated with the old copper mines on Ross Island, with species such as Sea Campion (*Silene vulgaris* subsp. *maritima*) and Thrift (*Armeria maritima*).

A large number of plant and animal species of interest occur within the site. For example, two plant species listed on Annex II of the E.U. Habitats Directive occur. Slender Naiad (*Najas flexilis*) is found in some of the lakes at the site. The Killarney Fern (*Trichomanes speciosum*) is another listed and well-known rarity. An additional twenty-two Red Data Book plant species have been recorded, but only twelve of these have been seen recently. These are Pillwort (*Pilularia globulifera*), Kerry Lily (*Simethis planifolia*), Irish Lady's-tresses (*Spiranthes romanzoffiana*), Slender Cottongrass (*Eriophorum gracile*), Small Cudweed (*Logfia minima*), Betony (*Stachys officinalis*), Heath Cudweed (*Omalotheca sylvatica*), Alder Buckthorn (*Frangula alnus*), Alpine Saw-wort (*Saussurea alpina*), Hoary Whitlowgrass (*Draba incana*), Smooth Brome (*Bromus racemosus*) and Holly Fern (*Polystichum lonchitis*). The first seven of these species are legally protected under the Flora (Protection) Order, 1999, as are Slender Naiad and Killarney Fern.

Additional plant species of interest include a fern (*Dryopteris affinis* subsp. *stilluppensis*) and a Whitebeam (*Sorbus anglica*), both at their only Irish locations.

The site is very important for oceanic bryophytes, particularly the woodland species. It also contains good representative examples of the Northern Atlantic Hepatic Mat community and other oceanic montane communities. Killarney Oak woods and mountains have been nominated as a site of international importance for bryophytes.

The Killarney Woods are notable for the number of rare species of Myxomycete fungus that have been recorded, namely *Collaria arcyriomena*, *Craterium muscorum*, *Cribraria microcarpa* (only known Irish site), *C. rufa*, *C. violacea*, *Diderma chondrioderma*, *D. lucidum*, *D. ochraceum*, *Fuligo muscorum* and *Licea marginata*.

The site has six bird species which are listed on Annex I of the E.U. Birds Directive. A small flock of Greenland White-fronted Goose, which winters on the boglands within the National Park, is now the only regular flock in the south-west. The site has one of

the highest concentrations of breeding Peregrines in the country, as well as some breeding Merlin. Chough is found both in the coastal and inland areas of the site, with possibly up to 30 pairs breeding. Kingfisher is a species associated with the lakes and rivers, especially in the National Park and probably breeds. Finally, a few pairs of Common Tern breed within the site.

The woodlands provide habitat for a variety of breeding birds, most notably Garden Warbler, Blackcap, and probably a few pairs each of the rare Redstart and Wood Warbler. Lough Leane is a site for wintering wildfowl with the following average counts for the two winters 1995/96 and 1996/97: Teal (208), Mallard (350), Pochard (81), Tufted Duck (323) and Coot (169).

The site supports most of the Irish mammal species. Of particular note is the occurrence of two E.U. Habitats Directive Annex II species: Lesser Horseshoe Bat, with a total population of about 300 individuals distributed at several locations, including both nursery and hibernation sites, and Otter. Perhaps the best known mammals of the Killarney National Park are the Red Deer, which form the only remaining native herd in Ireland, comprised of around 600 animals. Sika Deer also occur. Pine Marten is another notable species.

The site is valuable for its rare fish species, five of which are listed on Annex II of the E.U. Habitats Directive: Brook Lamprey (*Lampetra planeri*), River Lamprey (*Lampetra fluviatilis*), Sea Lamprey (*Petromyzon marinus*), Atlantic Salmon (*Salmo salar*) and Killarney Shad (*Alosa fallax killarnensis*). The Killarney Shad is a unique land-locked subspecies confined to the Killarney lakes. Also of note is the glacial relict, Arctic Char (*Salvelinus alpinus*), a Red Data Book species, a unique form of which is found in Lough Coomasaharn.

There are numerous rare invertebrates within the site. These include three E.U. Habitats Directive Annex II species: Kerry Slug (*Geomalacus maculosus*), the Freshwater Pearl Mussel (*Margaritifera margaritifera*) and the Marsh Fritillary (*Euphydryas aurinia*). The Kerry Slug and Pearl Mussel populations are of particular importance in a national context. Other species of note include: three chironomids of international importance found in the River Flesk; a wood ant (*Formica lugubris*) at one of only four Irish sites; a snail (*Limnaea involuta*), in Lough Crincaum, at its only known location; two dragonflies (*Cordulea aenea* and *Somatochlora arctica*), the former at one of only two known sites in Ireland and the latter at its only known Irish location; and several other aquatic and woodland species at their only known Irish locations.

The main land use within the site is grazing by sheep. In and around the National Park deer grazing is also common. The extensive grazing has caused damage to many of the terrestrial habitats, resulting in degradation of heath and blanket bogs and prevention of woodland regeneration. In the upland habitats the erosion caused by grazing is exacerbated by the exposed nature of the terrain. Apart from grazing, the woodlands are particularly threatened by Rhododendron (*Rhododendron ponticum*) invasion: approximately two thirds of the Oak woodlands are affected,

although a Rhododendron removal programme is underway in the National Park. The Yew wood has been adversely affected by heavy grazing for many years, but it is intended to control this in the near future by erection of a deer fence. The bogs are sensitive to grazing and are also threatened by turbary, burning and afforestation. Most of the lakes are very acid-sensitive and therefore vulnerable to afforestation within the catchment areas. Lough Leane has been subject to some eutrophication, although water quality appears to have improved since phosphates were removed from the sewage in 1985.

A management plan was drawn up for the Killarney National Park in 1991. The park is managed primarily for conservation purposes although recreation is also provided for.

Overall, the site is of high ecological value because of the diversity, quality and extensiveness of many of the habitats, and impressive list of rare species of flora and fauna. In recognition of its importance the Killarney National Park has been designated a World Biosphere Reserve.

*For inspection purposes only.
Consent of copyright owner required for any other use.*