

# Aquatic fauna refuges in Margaret River and the Cape to Cape region of Australia's Mediterranean-climatic Southwestern Province

Mark G. Allen<sup>1,2</sup>, Stephen J. Beatty<sup>1</sup> and David L. Morgan<sup>1\*</sup>

1. Freshwater Fish Group & Fish Health Unit, Centre for Fish & Fisheries Research, School of Veterinary and Life Sciences, Murdoch University, South St, Murdoch 6150, Western Australia
2. Current address: Department of Aquatic Zoology, Western Australian Museum, Locked Bag 49, Welshpool DC, Perth, Western Australia, 6986, Australia

\* correspondence to D.Morgan@murdoch.edu.au

## SUMMARY

Margaret River and the Cape to Cape region in the extreme south-western tip of Australia are located between Cape Naturaliste in the north and Cape Leeuwin in the south and encompass all intervening catchments that drain westward to the Indian Ocean. The region has a Mediterranean climate and houses 13 native, obligate freshwater macrofauna species (i.e. fishes, decapod crustaceans and a bivalve mollusc), four of which are listed as threatened under State and/or Commonwealth legislation. The most imperiled species are the Margaret River Burrowing Crayfish (*Engaewa pseudoreducta*) and Hairy Marron (*Cherax tenuimanus*), both of which are endemic to the Margaret River catchment and listed as critically endangered (also by the IUCN), and Balston's Pygmy Perch (*Nannatherina balstoni*) which is vulnerable. The region also houses several fishes that may represent new, endemic taxa based on preliminary molecular evidence. Freshwater ecosystems in the region face numerous threats including global climate change, a growing human population, introduced species, destructive land uses, riparian degradation, water abstraction, declining environmental flows, instream barriers, and fire. Here we review the current knowledge of the considerable aquatic biodiversity values of the region to provide a contemporary checklist, and to highlight actions that may be considered to protect these values in the face of both current and future conservation threats.

**Keywords:** Threatened species; teleosts; decapod crustaceans; agnathans; mussels

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

**Citation:** Allen MG, Beatty SJ, Morgan DL (2017) Aquatic fauna refuges in Margaret River and the Cape to Cape region of Australia's Mediterranean-climatic Southwestern Province. *Fishes in Mediterranean Environments* 2017.002: 26p.  
<https://doi.org/10.29094/FiSHMED.2017.002>

## INTRODUCTION

The south-west of Western Australia is a global hotspot of species endemism that is influenced by a Mediterranean climate (Myers *et al.*, 2000; Olson & Dinerstein, 2002; Morgan *et al.*, 2014a). The aquatic fauna of the region is species impoverished, yet the region has the highest rate of species endemism of any major drainage division in Australia, with >90% of the native aquatic macrofauna (i.e. fishes, decapod crustaceans, and bivalve molluscs) being endemic to this region (Morgan *et al.*, 2011, 2014a). Some of these endemic species have suffered dramatic range declines during the past century, mainly due to impacts associated with deforestation and agricultural development, whilst others are naturally restricted to narrow ranges (Morgan *et al.*, 1998, 2011; Unmack, 2001; Allen *et al.*, 2002; Klunzinger *et al.*, 2015). While land clearing is now less prevalent, the entire regional aquatic fauna is becoming increasingly jeopardised by a number of anthropogenic stressors and environmental threats, not the least of which is human-induced global climate change (Suppiah *et al.*, 2007; Morrongiello *et al.*, 2011; Beatty *et al.*, 2014; IPCC, 2014; Morgan *et al.*, 2014a). Consequently, several of the region's endemic aquatic species are now imperiled and officially recognised as threatened under State and/or Commonwealth legislation (Morgan *et al.*, 2011, 2014a).

Situated within south-western Australia, the Cape to Cape region (CCR) (Figure 1) is globally recognised as a premier food and wine producing region, as well as a popular tourist destination. The population of the CCR has expanded in recent decades and land use patterns have shifted accordingly (CCG, 2005a, 2005b, 2006; DoW, 2009, 2010). For instance, agriculture has intensified (particularly viticulture), resulting in a

growing demand for surface and subsurface water resources (CCG, 2005a, 2005b, 2006). In light of this, the Department of Water (Government of Western Australia) recently proclaimed several unregulated water resources in the CCR for allocation under license (DoW, 2009) and addressed the need for water management reform in the South West Regional Water Plan 2010 – 2030 (DoW, 2010). Balancing human water demands with ecological water requirements in a drying climatic region, where water demand is predicted to be at the limit of supply within 20 years, looms as a key management challenge (DoW, 2010). Moreover, the projected continuation of the current drying trend beyond this timeframe places further uncertainty on the long-term sustainability of some of the region's aquatic ecosystems (Suppiah *et al.*, 2007; CSIRO, 2009a, b; Morrongiello *et al.*, 2011).

The obligate freshwater macrofauna of the CCR comprises 13 native species (Morgan *et al.*, 2011), four of which are listed as threatened under the Wildlife Conservation Act 1950 and/or Environment Protection and Biodiversity Conservation (EPBC) Act 1999. Two additional species are listed as 'priority taxa' by the Department of Parks and Wildlife, Government of Western Australia (Morgan *et al.*, 2011, 2014a). The most imperiled are the Hairy Marron *Cherax tenuimanus* (Smith 1912) and the Margaret River Burrowing Crayfish *Engaewa pseudoreducta* Horwitz & Adams 2000, two range-restricted endemics that are listed as critically endangered under the EPBC Act 1999 (and also by the IUCN), while Balston's Pygmy Perch *Nannatherina balstoni* Regan 1906, and Carter's Freshwater Mussel *Westralunio carteri* (Iredale 1934) are either vulnerable (*N. balstoni*) or qualify as vulnerable (*W. carteri*) under the EPBC Act 1999 (Morgan *et al.*, 2014b, Klunzinger *et al.*, 2015). Notwithstanding the threat posed by climate change, these

rare species also face a number of further specific threats (see Burnham *et al.*, 2012; Duffy *et al.*, 2014; Klunzinger *et al.*, 2015).

With almost half of the CCR's native freshwater macrofauna recognised as threatened or requiring conservation management, freshwater ecosystems in the region have justifiably received considerable attention from natural resource managers, scientists and conservation organisations in recent years. Increasing resources have been channelled into conservation and land-care initiatives in the region, such as the construction of two rock ramp fishways on the Margaret River (Beatty *et al.*, 2007), comprehensive foreshore assessments (e.g. CCG, 2005a, 2006, 2008), and research, monitoring and recovery work on the region's critically endangered species (e.g. Bunn, 2004; Bunn *et al.*, 2008; Burnham *et al.*, 2012; Duffy *et al.*, 2014; Morgan *et al.*, 2014b; Klunzinger *et al.*, 2015). Given the burgeoning amount of unpublished literature pertaining to the catchments and resident aquatic macrofauna of the CCR, the aim of this review is to collate and interpret the available information and data, thus providing a framework for environmental managers to develop a conservation strategy that may be used to protect and enhance the region's aquatic biodiversity values.

## STUDY AREA

This review focuses on the area between Cape Naturaliste (33.53° S, 115.00° E) and Cape Leeuwin (34.38° S, 115.14° E) and encompasses all intervening catchments draining westward to the Indian Ocean (Figure 1). The geology of the region features a coastal strip of sandy soil overlying limestone with occasional outcroppings of the underlying granitic and gneissic bedrock (Leeuwin Block), whilst further inland the topography is undulating and low-relief (20-100 m above sea level) with a mixture of gravelly and sandy soils in upland areas and alluvial sandy loams on drainage lines and floodplains (Tille & Lantzke, 1990; CCG, 2003).

The CCR has a Mediterranean climate characterised by hot, dry summers and mild, wet winters. Long-term mean annual precipitation at Margaret River is 1131 mm, and is similar at other weather stations in the vicinity, but considerably higher than areas to the region's north (BoM 2015). Precipitation decreases to around 900 mm further inland near the headwaters of the Margaret River catchment (BoM, 2015). On average, >75% of precipitation falls in the five months from May to September (BoM, 2015). Evaporation rates in the region are around 1.5 times higher than precipitation rates (Taylor & Tinley, 1999; BoM, 2015) resulting in highly seasonal flows, with most drainage systems becoming a series of isolated pools during the dry summer-autumn period.

## CATCHMENT CONDITION IN THE CAPE TO CAPE REGION

Although small on a global scale, the Margaret River is the largest river system in the CCR, draining an area of 477 km<sup>2</sup> and features extensive permanent pool habitats throughout the middle and upper reaches. The river is in relatively good condition and has not become impacted by secondary salinisation, unlike many other rivers in south-western Australia (Morgan *et al.*, 2003; Beatty *et al.*, 2011). Almost the entire upper Margaret River catchment and significant portions of the lower catchment are forested (Green *et al.*, 2010) and only 21% of the catchment has been cleared of native vegetation (CCG, 2003; DoW, 2008). In contrast, other CCR catchments are much smaller and most have been cleared more extensively (Table 1). Agriculture (predominantly livestock and viticulture) is the dominant land use across the CCR, with the proportion of cleared land typically exceeding 60% (Table 1). Wilyabrup Brook is the most highly modified catchment in the CCR (Table 1) with unrestricted access of livestock to riparian habitats prevalent (CCG, 2006).

Threatened fauna of Australia's Cape to Cape region  
DOI: 10.29094/FiSHMED.2017.002




FIGURE 1. Major catchments and towns of the Mediterranean climatic Cape to Cape region of Western Australia.

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

**TABLE 1.** Attributes of major catchments of the Cape to Cape region, south-western Australia. The categorisation of foreshore condition (A – pristine; B – weedy; C – erosion prone/eroding; D - ditch) was derived from the application of methods developed by Pen and Scott (1995).

Catchment	Area (km <sup>2</sup> )	Stream length (km)	Foreshore condition	Dominant land use types	Key environmental issues	References
<b>Yallingup Brook</b>	<10	5.5	Not available	Native vegetation (64%); agriculture (36%)	Erosion & sedimentation; degradation & destruction of riparian vegetation	Taylor & Tinley, 1999
<b>Gunyulgup Brook</b>	47	24.3	A <1%; B 29%; C 58%; D 6.5%	Agriculture (65%); native vegetation (35%)	Erosion & sedimentation; excessive livestock access to riparian zone; water abstraction / barriers (110 dams)	Hunt <i>et al.</i> , 2002; CCG, 2005b
<b>Wilyabrup Brook</b>	89	~100	A 6%; B 17%; C 22%; D 44%; Dams 11%	Agriculture (mainly livestock & viticulture) (84%); native vegetation (12%); residential (4%)	Excessive livestock access to riparian zone; intensification of land use; reduced flows / water abstraction / barriers (~100 dams)	Hunt <i>et al.</i> , 2002; CCG, 2006
<b>Cowaramup Brook</b>	24	30	A 15%; B 32%; C 25%; D 28%	Agriculture (mainly viticulture) (64%); native vegetation (36%)	Excessive livestock access to riparian zone; reduced flows /water abstraction / barriers (19 dams); sedimentation; salinisation of springs; limited refuge habitat	Hunt <i>et al.</i> , 2002; CCG, 2008
<b>Ellen Brook</b>	29	40	A 5%; B 29%; C 27%; D 37%; unassessed 2%	Agriculture (70%); native vegetation (30%)	Excessive livestock access to riparian zone; expanding human population; water abstraction / barriers (32 dams + 1 weir); algal blooms	Hunt <i>et al.</i> , 2002; CCG, 2005a
<b>Margaret River</b>	477	~250	Main channel: A 50%; B 45%; C 5%; D 0%. Lower tributaries: A 11%; B 46%; C 20%; D 23%. Bramley Brook: A 28%; B 31%; C 6%; D 35%	Forested (mostly native, some plantations) (79%); cleared (agriculture, residential) (21%)	Degradation & destruction of riparian vegetation; water abstraction / barrier (670 dams, 43 > 8 ML capacity); alien species introductions; pollution; livestock access to riparian zone	Pen, 1999; CCG, 2003, 2009a, 2011; DoW, 2008; Green <i>et al.</i> , 2010
<b>Boodjidup Brook</b>	60	55	A 15%; B 38%; C 30%; D 17%	Agriculture (49%); native vegetation (36%); rural residential (10%)	Excessive livestock access to riparian zone; reduced flows /water abstraction / barriers (90 dams); erosion / sedimentation	CCG, 2009b
<b>Qunninup Brook</b>	19	20.4	A 2%; B 41%; C 14%; D 43%	Agriculture (livestock & viticulture) (72%); native vegetation (27%)	Water abstraction / flow reductions/ barriers; degradation & destruction of riparian vegetation	Hunt <i>et al.</i> , 2002; CCG, 2013



The proliferation of on-stream dams is a key issue impacting catchments across the CCR (Table 1). Wilyabrup Brook, for instance, has ca 100 on-stream dams (CCG, 2006), and the density of dams is even higher in the Gunyulgup Brook and Boodjidup Brook catchments (Green *et al.*, 2010; Table 1). The exception to this trend is the Margaret River main channel, which has only three low-level weirs, although there are 670 dams on its numerous tributaries including 43 of commercial size (i.e. >8 ML storage capacity) (Green *et al.*, 2010; Table 1). Water is also pumped directly from the river in the middle and lower reaches for anthropogenic uses (Green *et al.*, 2010). A report on the Ecological Water Requirements (EWR) in two representative reaches of the lower and middle section of Margaret River suggested that current levels of water abstraction were within ecologically sustainable limits but ongoing monitoring was required to ensure these limits would not be exceeded with future allocation of water licenses and climate change (Green *et al.*, 2010). In Cowaramup Brook, however, a similar study found that water abstraction was already close to the ecologically sustainable yield, with only a limited capacity for the system to accommodate increased consumption (Donohue *et al.*, 2010).


## FRESHWATER MACROFAUNA OF THE CAPE TO CAPE REGION

The CCR houses a total of six native fish species (Table 2; Figure 2) including one agnathan, the Pouched Lamprey *Geotria australis* Gray 1851, and five teleosts, i.e. Western Minnow *Galaxias occidentalis* Ogilby 1899, Western Mud Minnow *Galaxiella munda* McDowall at 1978, Balston's Pygmy Perch *N. balstoni*, Western Pygmy Perch *Nannoperca cf. vittata* and Nightfish

*Bostockia cf. porosa*. Preliminary genetic analyses have revealed that the latter two taxa represent cryptic species that are as yet undescribed (Unmack, 2013; Morgan *et al.*, 2014a). Additionally, five introduced teleost species have been recorded in fresh waters of the CCR, i.e. Rainbow Trout *Oncorhynchus mykiss* (Walbaum 1792), Goldfish *Carassius auratus* Linnaeus 1758, Common Carp *Cyprinus carpio* Linnaeus 1758, Eastern Gambusia *Gambusia holbrooki* (Girard 1859) and Redfin Perch *Perca fluviatilis* Linnaeus 1758 (Table 2; Figure 3); although some of these species may no longer be present in the region (see below).


The CCR also houses nine decapod crustacean species (Table 3), seven of which are native (Figure 4), i.e. Hairy Marron *Cherax tenuimanus*, Gilgie *Cherax quinquecarinatus* Gray 1845, Restricted Gilgie *Cherax crassimanus* Riek 1967, Koonac *Cherax preissii* (Erichson 1846), Glossy Koonac *Cherax glaber* Riek 1967, Margaret River Burrowing Crayfish *Engaewa pseudoreducta*, Augusta Burrowing Crayfish *Engaewa similis* Riek 1967, and an additional two that have been introduced from elsewhere in Australia (Figure 3), i.e. Smooth Marron *Cherax cainii* Austin 2002 and Yabby *Cherax destructor* Clark 1936. The freshwater bivalve mollusc Carter's Freshwater Mussel (*Westralunio carteri*) also occurs in the CCR (Table 3, Figure 4).

A number of euryhaline species (e.g. *Leptatherina wallacei* (Prince, Ivantsoff & Potter 1982), *Pseudogobius olorum* (Sauvage 1880), *Afurcagobius suppositus* (Sauvage 1880)) have occasionally been reported from fresh waters of the CCR; however, these species generally only use freshwater habitats opportunistically and will not be covered further in this review.



**FIGURE 2.** Native freshwater fishes of the Cape to Cape region: a) maturing Pouched Lamprey (*Geotria australis*) (image: S. Beatty); b) suckling disc of sub-adult Pouched Lamprey (image: S. Beatty); c) metamorphosed juvenile or downstream migrant Pouched Lamprey (image: D. Morgan); d) larva (ammocoete) Pouched Lamprey (image: D. Morgan); e) Western Minnow (*Galaxias occidentalis*) (image: M. Allen); f) Western Mud Minnow (*Galaxiella munda*) (image: G. R. Allen); g) Nightfish (*Bostockia porosa*) (image: S. Beatty); h) Western Pygmy Perch (*Nannoperca vittata*) (image: M. Allen); i) Balston's Pygmy Perch (*Nannatherina balstoni*) (image: D. Morgan).





**FIGURE 3.** Introduced freshwater macrofauna species of the Cape to Cape region: a) Rainbow Trout (*Oncorhynchus mykiss*) (image: R. Kuitert); b) Goldfish (*Carassius auratus*) (image: S. Beatty); c) Common Carp (*Cyprinus carpio*) (image: S. Beatty); d) Redfin Perch (*Perca fluviatilis*) (image: R. Kuitert); e) female Eastern Gambusia (*Gambusia holbrooki*) (image: D. Morgan); f) male Eastern Gambusia (image: M. Allen); g) Smooth Marron (*Cherax cainii*) (image: D. Morgan); h) Yabby (*Cherax destructor*) (image: D. Morgan).



Threatened fauna of Australia's Cape to Cape region  
DOI: 10.29094/FiSHMED.2017.002



**FIGURE 4.** Native freshwater decapod crustaceans and bivalve molluscs in the Cape to Cape region: a) Margaret River Burrowing Crayfish (*Engaewa pseudoreducta*) (image: Q. Burnham); b) Augusta Burrowing Crayfish (*Engaewa similis*) (image: Q. Burnham); c) Hairy Marron (*Cherax tenuimanus*) (image: S. Visser); d) Gilgie (*Cherax quinquecarinatus*) (image: D. Morgan); e) Restricted Koonac (*Cherax crassimanus*) (image: S. Beatty); f) Koonac (*Cherax preissii*) (image: D. Morgan); g) Glossy Koonac (*Cherax glaber*) (image: R. McCormack); h) Carter's Freshwater Mussel (*Westralunio carteri*) (image: D. Morgan).

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

**TABLE 2.** Freshwater fish species present in the Cape to Cape region, south-western Australia. NB -\* denotes an introduced species; ^ denotes a species that is presumed to have been locally extirpated from a catchment.

Family/Species	Common name	Catchments	References
<b>Geotriidae</b>			
<i>Geotria australis</i>	Pouched Lamprey	Margaret	Morgan <i>et al.</i> , 1998; Morgan & Beatty, 2003, 2004a, 2007a; Beatty & Morgan, 2008
<b>Galaxiidae</b>			
<i>Galaxias occidentalis</i>	Western Minnow	Margaret, Wilyabrup, Boodjidup, Turner	Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003, 2004a, 2005, 2007a, 2008; Beatty <i>et al.</i> , 2006, 2007, 2008; Beatty & Morgan, 2008; Beatty & Allen, 2008; Morgan <i>et al.</i> , 1998; Morgan & Beatty, 2003, 2004a, 2008; Beatty & Allen, 2008; Beatty <i>et al.</i> , 2008
<i>Galaxiella munda</i>	Western Mud Minnow	Margaret, Wilyabrup, Boodjidup	Morgan <i>et al.</i> , 1998; Morgan & Beatty, 2003, 2004a, 2008; Beatty & Allen, 2008; Beatty <i>et al.</i> , 2008
<b>Percichthyidae</b>			
<i>Bostockia porosa</i>	Nightfish	Margaret, Wilyabrup, Ellen, Boodjidup, Turner^	Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003, 2004a, 2005, 2008; Beatty & Allen, 2008; Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003, 2004a, 2005; Beatty <i>et al.</i> , 2006, 2008; Beatty & Allen, 2008
<i>Nannoperca vittata</i>	Western Pygmy Perch	Margaret, Wilyabrup, Ellen, Turner^	Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003, 2004a, 2005; Beatty <i>et al.</i> , 2006, 2008; Beatty & Allen, 2008
<i>Nannatherina balstoni</i>	Balston's Pygmy Perch	Margaret., Turner^	Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003
<b>Salmonidae</b>			
* <i>Oncorhynchus mykiss</i>	*Rainbow Trout	Margaret, private farm dams throughout CCR	Morgan <i>et al.</i> , 1998; CCG, 2011
<b>Cyprinidae</b>			
* <i>Carassius auratus</i>	*Goldfish	Margaret^, Wilyabrup	Beatty <i>et al.</i> , 2008; Allen <i>et al.</i> , 2013
* <i>Cyprinus carpio</i>	*Common Carp	Margaret^, Wilyabrup^	R. Paice, pers. comm.; D. McKenzie, pers. comm.
<b>Poeciliidae</b>			
* <i>Gambusia holbrooki</i>	*Eastern Gambusia	Margaret, Cowaramup, Ellen, Boodjidup, Turner	Morgan <i>et al.</i> , 1998, 2013; Morgan & Beatty, 2003, 2004a, 2005, 2008; Beatty & Morgan, 2008; Beatty <i>et al.</i> , 2008
<b>Percidae</b>			
* <i>Perca fluviatilis</i>	*Redfin Perch	Margaret^	Morgan <i>et al.</i> , 1998



**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

**TABLE 3.** Freshwater decapod crustaceans (Parastacidae) and bivalve mollusc (Hyriidae) species present in the Cape to Cape region, south-western Australia. NB -\* denotes an introduced species; ^ denotes a species that is presumed to have been locally extirpated from a catchment; (?) denotes an unconfirmed record.

Family/Species	Common name	Catchments	References
<b>Parastacidae</b>			
<i>Engaewa pseudoreducta</i>	Margaret River Burrowing Crayfish	Margaret	Burnham <i>et al.</i> , 2012; DEC 2008
<i>Engaewa similis</i>	Augusta Burrowing Crayfish	Margaret, Boodjidup, Turner^	Horwitz & Adams 2000; Morgan <i>et al.</i> 2013; Q. Burnham pers. comm.
<i>Cherax tenuimanus</i>	Hairy Marron	Margaret	Austin & Ryan 2002; Bunn 2004; Molony <i>et al.</i> 2004; de Graaf <i>et al.</i> 2009
<i>Cherax quinquecarinatus</i>	Gilgie	Margaret, Wilyabrup, Ellen, Cowaramup, Gunyulgup, Boodjidup, Calgardup	Morgan & Beatty 2003, 2004a, 2005, 2008; Beatty <i>et al.</i> 2006, 2008; Beatty & Allen 2008;
<i>Cherax crassimanus</i>	Restricted Koonac	Margaret, Ellen, Turner^	Morgan & Beatty 2003, 2004a, 2005, 2008; Beatty <i>et al.</i> 2006, 2008; Beatty & Allen 2008;
<i>Cherax preissii</i>	Koonac	Margaret, Ellen, Calgardup, Turner	Morgan & Beatty 2005; Morgan <i>et al.</i> 2013
<i>Cherax glaber</i>	Glossy Koonac	Calgardup (?)	Austin & Knott 1996
* <i>Cherax cainii</i>	*Smooth Marron	Margaret, Wilyabrup, Cowaramup, Ellen, Gunyulgup, Boodjidup	Morgan & Beatty 2005, 2008; Beatty <i>et al.</i> 2006; de Graaf <i>et al.</i> 2009
* <i>Cherax destructor</i>	*Yabby	Margaret, Wilyabrup, Gunyulgup, Boodjidup, Turner	Morgan & Beatty 2005, 2008; Beatty <i>et al.</i> 2006, 2008; Beatty & Allen 2008; Allen <i>et al.</i> 2013; Morgan <i>et al.</i> 2013
<b>Hyriidae</b>			
<i>Westralunio carteri</i>	Carter's Freshwater Mussel	Margaret, Wilyabrup, Ellen, Boodjidup	Klunzinger <i>et al.</i> 2012



## THREATENING PROCESSES IN THE CAPE TO CAPE REGION

### *Climate change*

South-western Australia has experienced a 10-15% reduction in rainfall since the mid-1970s with some parts of the region experiencing a 50% reduction in surface runoff during this time (Silberstein *et al.*, 2012). Global Climatic Models unanimously project further reductions in rainfall (ca 8%) and runoff (ca 24%), and an increase in the number of no-flow days (by up to 4 months) by 2030 (Suppiah *et al.*, 2007; CSIRO, 2009a, b; Barron *et al.*, 2012; Silberstein *et al.*, 2012). Compared with baseline period between 1986-2005, median estimates of average annual rainfall decrease for the region by 2090 are 12% under intermediate emissions scenario (RCP 4.5), and 18% under high emissions scenario (RCP 8.5); the latter could result in ca 65% reductions in annual runoff (Hope *et al.*, 2015). Drought periods and days of extreme heat are also projected to increase, which will cause higher evaporation rates (CSIRO, 2009a, b, Hope *et al.*, 2015). These changes are likely to have a profound impact on freshwater ecosystems in the region, in particular on the quantity and quality of available baseflow refuge habitat. The anticipated decline in biological carrying capacity of freshwater ecosystems due to climate change may lead to localised extirpations or even extinctions of the rarest and most vulnerable species (IPCC, 2013, 2014; DEC, 2008).

A recent survey of key baseflow refuge habitats in the Margaret River has revealed dramatic declines in the abundance of both Pouched Lamprey and Western Mud Minnow. Pouched Lamprey larvae (i.e. ammocoetes) were not found at 75% of previous known sites and not a single individual Western Mud Minnow was captured despite thorough sampling of sites where the species was previously common (see Morgan & Beatty, 2003). Land use and water quality in the Margaret River catchment have not changed markedly over this time period; indeed, the historical Western Mud Minnow sites were all located within a conservation

reserve that was established many years prior to the initial fish survey. Whilst the spread of invasive Eastern Gambusia may be a factor in the decline of Western Mud Minnow (discussed in the Introduced species section below), a more likely explanation for the observed fish declines in the Margaret River is the sharp drop in annual stream discharge in recent years. Five of the lowest discharge years on record (since 1970 when record keeping commenced in the catchment) have occurred since 2006. Western Mud Minnow has been shown to spawn in temporarily flooded riparian zones and its recruitment success is likely to be linked to stream flow as has been demonstrated for other south-western Australian fishes including lampreys (Morgan *et al.*, 1998; Beatty *et al.*, 2007, 2014). Moreover, Western Mud Minnow typically has a short lifespan with most individuals spawning at age one and perishing shortly afterwards (Pen *et al.*, 1991), therefore population numbers are probably susceptible to dramatic inter-annual fluctuations depending on the suitability of environmental conditions from year to year. The prolonged drought conditions in recent years may account for the severe population decline of these species in Margaret River, as well as providing a bellwether of the potential for climate change to impact freshwater species in this system and throughout the south-west region.

The burrowing crayfishes (*Engaewa* spp.) are likely to be particularly vulnerable to climate change as they are restricted to ephemeral habitats that may cease to function effectively in the future due to rainfall reductions and declining water tables (Commander, 2000; DEC, 2008). It is unknown if these species will have the capacity to burrow to sufficient depths in order to withstand predicted water table declines in the region and interventionary actions may be required to maintain populations in their natural habitats. The Margaret River Burrowing Crayfish (*E. pseudoreducta*) is already critically endangered due to its small geographic range, limited dispersal capability, specialised ecological requirements and small population size (DEC, 2008). These

attributes combine to make it the most imperiled aquatic species in the CCR (and arguably in the entire south-western Australian region), and its conservation warrants top priority. Captive breeding programs may need to be considered in the future to avoid extinction of this species (DEC, 2008).

### **Water abstraction**

Groundwater abstraction can reduce flow from natural springs (CSIRO, 2009b), which has been shown to be vital in maintaining important aquatic refuges in some catchments in south-western Australia (Beatty *et al.*, 2014). It will be critical to ensure that abstraction from major aquifers, particularly the south-west Yarragadee and Leederville, does not exacerbate the effects of climate change in reducing the amount and quality of baseflow refuge pools; particularly those in the upper Margaret River. Other water abstraction issues include the illegal pumping of river water, non-compliance by private land holders in the installation and use of flow-bypass valves in dams, and the impact of timber plantations intercepting run-off and causing draw-down of water tables (DoW, 2009). In the Cowaramup Brook catchment, water abstraction is already very close to the maximum sustainable level necessary to maintain aquatic ecological functions (Donohue *et al.*, 2010). Further EWR studies would be beneficial in order to determine if water use is encroaching on ecologically sustainable limits in other catchments of the region. The State Government's water authorities have embarked on a massive public educational campaign to raise awareness of the precarious state of potable water supplies in south-western Australia and to encourage strategies for limiting water wastage. Clearly, authorities will need to closely monitor and, where required, impose restrictions on human water use in the CCR and across the broader region in order to balance human demands with ecological requirements as the region continues its transition to drier and warmer climatic conditions.

### **Introduced species**

Over the past 40 years, the rate of species introductions has soared (75% increase since 1970) in fresh waters of south-western Australia (Beatty & Morgan, 2013). The earliest introductions were government sanctioned for the purpose of creating enhanced recreational angling opportunities (e.g. Redfin Perch, Rainbow Trout) and for the biological control of mosquitos (i.e. Eastern Gambusia). More recently, popular aquarium and aquaculture species (e.g. Goldfish, Carp, Yabby, Smooth Marron) have become established in natural waterways via the dumping of unwanted animals or via their escape from ponds and dams during flooding (Morgan & Beatty, 2004b, 2007b; Morgan *et al.*, 2004, 2011; Beatty & Morgan, 2013). Invasive aquatic species have been shown to have adverse impacts on Australian native species through predation, competition for food and space resources, agonistic behaviour, habitat degradation, and as vectors of diseases and parasites (e.g. Fletcher *et al.*, 1985; Horwitz, 1990; Gill *et al.*, 1999; Morgan *et al.*, 2002, 2004; Morgan & Beatty, 2007b; Tay *et al.*, 2007; Lymbery *et al.*, 2010; Beatty & Morgan, 2013).


Six introduced aquatic macrofauna species have been recorded in fresh waters of the CCR (Tables 2; 3). Additionally, despite being a south-western Australian endemic, translocation of Smooth Marron into the Margaret River catchment has pushed the Hairy Marron to the brink of extinction (Bunn, 2004; de Graaf *et al.*, 2009; Duffy *et al.*, 2014). In response, the Department of Fisheries, Government of Western Australia, has established a captive breeding program of genetically validated pure-strain Hairy Marron for supplementation of wild stocks (de Graaf *et al.*, 2009; Duffy *et al.*, 2014), and has overseen the removal of large numbers of Smooth Marron and hybrid specimens from the upper Margaret River since 2004, which has slowed but not reversed the overall decline of Hairy Marron in this system (de Graaf *et al.*, 2009).

Generally, once established, introduced species prove impossible to eradicate

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

from the wild (Horwitz, 1990; Rowe *et al.*, 2008). On the one hand, Smooth Marron and Eastern Gambusia have become well established and widespread in the CCR (Figures 5 and 6), with the latter species recently being discovered for the first time in Canebrake Pool, one of the most important baseflow refuge habitats in the Margaret River (Allen *et al.*, 2015). This noxious pest species has rapidly become one of the most abundant fishes at this site, underlining its highly invasive traits. Its impact on the native aquatic fauna has been significant, with >90% of native percichthyids


captured from this site showing signs of caudal fin damage due to fin-nipping by Eastern Gambusia (Figure 7). Its establishment in Canebrake Pool may have also contributed to the decline of Western Mud Minnow at this site. However, as the abundance of Western Mud Minnow has also dropped to undetectable levels in refuge pools further upstream that are not yet colonised by Eastern Gambusia, the decline is more likely attributable to climate change driven flow reductions.




**FIGURE 5.** Distribution of introduced fishes in the Cape to Cape region of Western Australia.



Threatened fauna of Australia's Cape to Cape region  
 DOI: 10.29094/FiSHMED.2017.002



**FIGURE 6.** Distribution of introduced crayfishes in the Cape to Cape region of Western Australia.



**FIGURE 7.** Caudal fin damage on percichthyids captured from Canebrake Pool, Margaret River (March-April 2015) caused by agonistic behaviour of Eastern Gambusia; a) Balston's Pygmy Perch (damaged tail above, undamaged tail below); b) Nightfish; c) Western Pygmy Perch (images: M. Allen).

The Yabby is also in the early stages of establishing itself in the region (Figure 6). Unfortunately, this species is very adaptable to local conditions and eradication is unlikely via conventional control methods. The Yabby is of particular concern as it occupies an equivalent ecological niche to endemic *Cherax* spp. and directly competes with them for food and habitat resources (Beatty, 2006; Lynas *et al.*, 2007). It is also capable of breeding multiple times per year, allowing it to rapidly colonise habitats and potentially outnumber native species (Beatty *et al.*, 2005).

The remaining alien species have either failed to establish self-sustaining populations (e.g. Rainbow Trout and Redfin Perch), or their spread in the region has, thus far, been effectively contained through targeted fishing (e.g. Goldfish and Common Carp). Goldfish have been eradicated from Darch Brook (a tributary of Margaret River) via a combination of habitat dewatering, netting and chemical ichthyocide (rotenone) application (Allen *et al.*, 2013). As far as the authors are aware, this represents the first successful eradication of the species from an aquatic system in Western Australia. The only other record of Goldfish in the CCR is from an artificial pond in the Wilyabrup Brook catchment near Cowaramup, where >1,000 individuals were removed in 2011 (Freshwater Fish Group, Murdoch University, unpubl. data). However the species remains at liberty in the catchment.

The strategy of targeted fishing effort has had mixed success in mitigating the environmental risks of introduced freshwater species in the CCR, but undoubtedly a more cost-effective and practical strategy would be to prevent further introductions from occurring in the first instance. This is probably best achieved through educational programs designed to raise awareness and appreciation of native biodiversity whilst also emphasising the negative impacts that introduced species can have on native wildlife. The biosecurity branch of the Department of Fisheries (Government of Western Australia) has recently launched an educa-

tional campaign to address this growing threat.

### ***Instream barriers***

Artificial barriers on waterways (e.g. dams, weirs, road crossings) can impact on aquatic species in a number of ways. Most crucially, they reduce longitudinal connectivity in lotic systems, which blocks or restricts the access of migratory species to vital habitats such as breeding grounds, seasonal feeding areas, or nurseries (Lucas & Baras, 2001; Morgan & Beatty, 2003; Koehn & Crook, 2013). For example, a number of native freshwater fishes (e.g. Western Minnow, Nightfish) move upstream to spawn in lower order tributaries with the onset of winter flows (Beatty *et al.*, 2014). Not only do instream barriers restrict the migration of such species through physical blockage, they also delay the onset of seasonal flows that provide spawning cues, thus shortening the duration of the breeding season. In the CCR, hundreds of instream barriers (mainly gully dams) have been installed in recent decades (e.g. CCG, 2005a, 2006), yet only three fish passage structures have been built during this time (Beatty *et al.*, 2007; Beatty & Allen, 2008). Furthermore, these fishways have been shown to effectively allow passage of only a subset of the resident fish fauna (i.e. Western Minnow, Pouched Lamprey and Western Pygmy Perch) while blocking upstream passage of other fishes (Morgan & Beatty, 2004; Beatty & Allen, 2008; Beatty & Morgan, 2008). Thus, there is a need for further research into modifications or alternative designs that will allow the full suite of migratory native aquatic species to bypass instream barriers in the CCR and more broadly throughout south-western Australia.

A simpler and more effective means of mitigating the impacts of instream barriers on migratory species may be the decommissioning and removal of artificial barriers (see Beatty *et al.*, 2013). However, these structures are rarely redundant and may even provide ecological benefits such as the provision of permanent refuge habitats in their associated impoundments (an eco-

logical function that will become increasingly valuable in a drying climatic region), or by limiting the dispersal of introduced species. In Boodjidup Brook for instance, the absence of Eastern Gambusia from the majority of the upper catchment is presumably due to the presence of an instream barrier, which may, in part, explain why the Western Mud Minnow persists in reasonable numbers in this system (Morgan & Beatty, 2008).

### ***Loss and degradation of riparian vegetation***

Riparian vegetation provides multiple benefits for both terrestrial and aquatic fauna and plays a vital role in maintaining aquatic ecosystem function and health (Pen, 1999). In the past, riparian vegetation was commonly destroyed during the development of agricultural land. The infamous case of the extirpation of a population of critically endangered Margaret River Burrowing Crayfish at the type locality in 1985 following the construction of a dam and establishment of a timber plantation (DEC, 2008) highlights the severity of the impact that such practices can have on aquatic macrofauna. Although the rate of clearing of riparian vegetation has declined in recent years, degradation of remnant vegetation is an ongoing issue across the region, mainly due to unfettered access of livestock to stream lines and this threatens aquatic ecosystems through processes of erosion, sedimentation and increased nutrient inputs.

Mature overstorey vegetation is vital in moderating water temperature and evaporation rates through shading (Pen, 1999; Davies *et al.*, 2007), and provides instream habitat for native aquatic species such as Nightfish and Marron in the form of woody debris (Pen, 1999). Additionally, the leafy material that falls into the water is a vital source of carbon that drives aquatic food webs in a region where primary productivity in streams is naturally low (Pen, 1999). Deep rooted riparian vegetation also keeps salt-laden groundwater at a sufficient depth to prevent secondary salinisation of streams, which has led to the extirpation of

native species in some of the major rivers of south-western Australia (Pen, 1999; Morgan *et al.*, 1998, 2003).

Much riparian rehabilitation and fencing work has already been undertaken in the region (D. McKenzie, Cape to Cape Catchments Group pers. comm.) and our review revealed that further work is needed. From the standpoint of conserving aquatic fauna, future works should focus on sites that house threatened or rare aquatic species, particularly Burrowing Crayfishes (*Engaewa* spp.), Balston's Pygmy Perch, and Western Mud Minnow.

### ***Fire***

Native vegetation has long been burnt on a multi-annual rotational schedule as a land management tool for reducing fuel loads and associated risk of destructive wildfires, and to stimulate native plant growth and reproduction (Christensen & Kimber, 1975). Given the prevalence of fire in the Australian landscape (and particularly in the south-west of Australia); it is surprising that its impact on aquatic ecosystems has not yet been studied in detail. Concerns have been raised about the impact of fire on *Engaewa* spp., especially the highly restricted *E. pseudoreducta* (DEC, 2008). The restricted range of this species leaves it extremely vulnerable to habitat disturbance that may occur during fire management activities such as the clearing of firebreaks and containment lines, or in the use of fire retardant chemicals (DEC, 2008). In the *Engaewa* Recovery Plan it was stated that there was an "urgent need to improve the understanding of the effects of fire on *Engaewa* spp." (DEC, 2008) but little has been done to address this knowledge gap, and we contend that the same applies to other native aquatic species, and aquatic ecosystem more generally, as well. The threat of wild fires is likely to escalate as the region's climate becomes hotter and drier; therefore managing associated risks is likely to become more prevalent in the future.



## CONCLUSIONS

The Cape to Cape Region is an iconic part of Mediterranean climatic Australia. It has an expanding population and is also one of the premier tourist destinations in Western Australia due to the climate and natural beauty. This review has highlighted that the aquatic macrofauna of the region, whilst depauperate, has exceptional rates of endemism and contains a number of threatened species. We recommend that sites within the CCR known to house these species (Figures 8 and 9) be monitored on a regular basis in order to maintain an up-to-date knowledge of the conservation status of threatened populations and their habitats so that immediate threats can be identified and addressed (e.g. riparian zone degradation, water quality issues, novel species introductions). The importance of such monitoring was highlighted during the recent survey of baseflow refuge pools in Margaret River, where a severe decline in abundance of both Western Mud Minnow and Pouched Lamprey was detected, as was the spread of invasive Eastern Gambusia into the upper Margaret River (Allen *et al.*, 2015).

One of the major environmental challenges faced in the region is finding a balance between human water demands and those of the environment. Many of the aquatic ecosystems in the region are degraded, while the areas that remain near pristine face a number of serious threats. The work on improving the management of aquatic ecosystems in the CCR to date has

been commendable, but our review reveals a clear need for ongoing and adaptive management, as well as further monitoring and onground works that will contribute to the protection of the aquatic fauna values of the CCR.


## ACKNOWLEDGEMENTS

We wish to thank the Department of Parks and Wildlife (DpaW, formerly DEC), Government of Western Australia, for providing the funding to undertake this project. We thank Drew McKenzie from the Cape to Cape Catchments Group for assisting with the project, Dr Quinton Burnham (Griffith University) for providing images and valuable feedback on Engaewa, Dr Rodney Duffy (Department of Fisheries) for information on Hairy Marron conservation initiatives, Mike Klunzinger (Murdoch University) for information on freshwater bivalves, and Sue Morrison and Andrew Hosie (Western Australian Museum) for contributing historical collection records. We thank Dr Gerald R. Allen (Western Australian Museum), Rudie Kuitert, Robert McCormack, and Simon Visser for allowing us to use their images. Lastly, we also acknowledge the original environmental custodians and traditional owners of the CCR - the Wardandi people.

## AUTHOR CONTRIBUTIONS


Each of the authors contributed to the collection of data, and writing of the manuscript

Threatened fauna of Australia's Cape to Cape region  
 DOI: 10.29094/FiSHMED.2017.002



**FIGURE 8.** Distribution of threatened fish and crayfish species in the Cape to Cape region, Western Australia. Black diamonds indicate known sites where the species is likely to remain extant; pink diamonds indicate historical sites where species is presumed to have been lost.

Threatened fauna of Australia's Cape to Cape region  
DOI: 10.29094/FiSHMED.2017.002



**FIGURE 9.** Conservation priority areas in the Cape to Cape region housing threatened, endangered and priority species (i.e. Hairy Marron, Margaret River Burrowing Crayfish, Balston's Pygmy Perch, Western Mud Minnow, and Pouched Lamprey)



## CITED REFERENCES

- Allen GR, Midgley SH, Allen M (2002) Field Guide to the Freshwater Fishes of Australia. Western Australian Museum, Perth, 394 pp.
- Allen MG, Beatty SJ, Morgan DL (2013) Fish survey of Brookfield Estate, Darch Brook and Margaret River: evidence of eradication of goldfish from a Western Australian river. Freshwater Fish Group and Fish Health Unit (Murdoch University), Report to the Cape to Cape Catchments Group, August 2013.
- Allen MG, Beatty SJ, Morgan DL (2015) Baseline survey of key fish refuge habitats in the Margaret River. Freshwater Fish Group & Fish Health Unit, Centre for Fish & Fisheries Research, Murdoch University, Report to the Cape to Cape Catchments Group and South West Catchments Council, Perth, Western Australia.
- Austin CM, Knott B (1996) Systematics of the freshwater crayfish genus *Cherax* Erichson (Decapoda: Parastacidae) in south-western Australia: electrophoretic, morphological and habitat variation. Australian Journal of Zoology 44:223–258.
- Austin CM, Ryan SG (2002) Allozyme evidence for a new species of freshwater crayfish of the genus *Cherax* Erichson (Decapoda: Parastacidae) from the south-west of Western Australia. Invertebrate Systematics 16:357–367.
- Barron O, Silberstein R, Ali R, Donohue R, McFarlane DJ, Davies P, Hodgson G, Smart N, Donn M (2012) Climate change effects on water-dependent ecosystems in south-western Australia. Journal of Hydrology 434–435:95–109.
- Beatty SJ (2006) The diet and trophic positions of translocated, sympatric populations of *Cherax destructor* and *Cherax cainii* in the Hutt River, Western Australia: evidence of resource overlap. Marine and Freshwater Research 57:825–835.
- Beatty S, Allen M (2008) Preliminary assessment of the functioning of the bypass fishway on Wilyabrup Brook. Report to Cape to Cape Catchments Group. Centre for Fish and Fisheries Research, Murdoch University, WA.
- Beatty SJ, Morgan DL (2008) Monitoring the Margaret River fishways - 2007. Report to the Cape to Cape Catchments Group and Department of Water. Centre for Fish and Fisheries Research, Murdoch University, WA.
- Beatty SJ, Morgan DL (2013) Introduced freshwater fishes in a global endemic hotspot and implications of habitat and climatic change. BioInvasions Records 2:1–9.
- Beatty S, Allen M, Lymbery A, Storer T, White G, Morgan D, Ryan T (2013) Evaluating small barrier removal to improve refuge connectivity: a global review of barrier decommissioning and a process for southern Australia in a drying climate. In: Novel methods for managing freshwater refuges against climate change in southern Australia (eds B J Robson, E T Chester, M Allen, S Beatty, P Close, B Cook, P Davies, R Lester, A Lymbery, T Matthews, D Morgan). Final Report. NCCARF, Brisbane. 81 pp.
- Beatty SJ, Morgan DL, Rashnavadi M, Lymbery AJ (2011) Salinity tolerances of endemic freshwater fishes of south-western Australia: implications for conservation in a biodiversity hotspot. Marine and Freshwater Research 62:91–100.
- Beatty S, Morgan D, Allen M (2008) Freshwater fish and crayfish communities of the tributaries of the Margaret River. Technical report to Cape to Cape Catchments Group. Cen-

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

- tre for Fish and Fisheries Research, Murdoch University, WA, 25 pp.
- Beatty SJ, Morgan DL, Gill HS (2005) Role of life history strategy in the colonisation of Western Australian aquatic systems by the introduced crayfish *Cherax destructor* Clark, 1936. *Hydrobiologia* 549:219–237.
- Beatty S, Morgan D, Jury C, Mitchell J (2006) Fish and freshwater crayfish in streams in the Cape Naturalist region and Wilyabrup Brook. Report prepared for the Cape to Cape Catchments Group and GeoCatch, 28 pp.
- Beatty SJ, Morgan DL, Lymbery AJ (2014) Implications of climate change for potamodromous fishes. *Global Change Biology* 20:1794–1807.
- Beatty SJ, Morgan DL, Torre A (2007) Restoring ecological connectivity in the Margaret River: Western Australia's first rock-ramp fishways. *Ecological Management and Restoration* 8:224–229.
- Bunn JJS (2004) Investigation of the replacement of Margaret River hairy marron *Cherax tenuimanus* (Smith) by smooth marron *C. cainii* Austin: 41–74. M.Sc. Thesis, Edith Cowan University, Perth.
- Bunn JJS, Koenders A, Austin CM, Horwitz P (2008) Identification of hairy, smooth and hybrid marron (Decapoda: Parastacidae) in the Margaret River: morphology and allozymes. *Freshwater Crayfish* 16:113–121.
- Bureau of Meteorology (2015) (webpage) viewed July 28 2015, <http://www.bom.gov.au/climate/>
- Burnham QF, Koenders A, Horwitz P (2012) The status of the critically endangered freshwater crayfish *Engaewa pseudoreducta* (Crustacea: Parastacidae) in south-western Australia. *Records of the Western Australian Museum* 27:45–54.
- Cape to Cape Catchments Group (2003) Margaret River Action Plan. Water and Rivers Commission, 65 pp.
- Cape to Cape Catchments Group (2005a) Ellen Brook, River Action Plan. Department of Environment, Perth, 68 pp.
- Cape to Cape Catchments Group (2005b) Gunyulgup Brook, River Action Plan. Department of Environment, Perth, 84 pp.
- Cape to Cape Catchments Group (2006) Wilyabrup Brook, River Action Plan. Department of Environment, Perth, 95 pp.
- Cape to Cape Catchments Group (2008) Cowaramup Creeks Action Plan, 78 pp.
- Cape to Cape Catchments Group (2009a) Tributaries of the lower Margaret River Action Plan, 74 pp.
- Cape to Cape Catchments Group (2009b) Boodjidup Brook Action Plan, 77 pp.
- Cape to Cape Catchments Group (2011) Bramley Brook Action Plan, 73 pp.
- Cape to Cape Catchments Group (2013) Quinninup Brook Action Plan, 68 pp.
- Christensen PE, Kimber PC (1975) Effect of prescribed burning on the flora and fauna of south west Australian forests. *Proceedings of the Ecological Society of Australia* Vol. 1975.
- Commander DP (2000) Potential effects of climate change on groundwater resources in Western Australia. *Hydro 2000*, Perth, Western Australia, 20–23 November, Institution of Engineers, Australia, pp. 234–239.
- CSIRO (2009a) Surface water yields in south-west Western Australia. A report to the Australian Government from the CSIRO South-West Western Australia Sustainable Yields Project. CSIRO Wa-

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

- ter for a Healthy Country Flagship, Australia, 171 pp.
- CSIRO (2009b) Groundwater yields in south-west Western Australia. A report to the Australian Government from the CSIRO South-West Western Australia Sustainable Yields Project. CSIRO Water for a Healthy Country Flagship, Australia, 330 pp.
- Davies PM, Bunn S, Mosisch T, Cook BA, Walshe TV (2007) Temperature and light. *In*: Principles for riparian lands management. Land and Water Australia, Canberra, ACT.
- de Graaf M, Lawrence C, Vercoe P (2009) Rapid replacement of the critically endangered Hairy Marron by the introduced Smooth Marron (Decapoda, Parastacidae) in the Margaret River (Western Australia). *Crustaceana* 82:1469–1476.
- Department of Environment and Conservation (2008) Dunsborough Burrowing Crayfish (*Engaewa reducta*), Margaret River Burrowing Crayfish (*Engaewa pseudoreducta*) and Walpole Burrowing Crayfish (*Engaewa walpolea*). Recovery Plan 2007-2016. Species and Communities Branch. Department of Environment and Conservation, Bentley, Western Australia, 33 pp.
- Department of Water (2008) Margaret River Hydrology Summary. Surface water hydrology series, Report No. 27, Department of Water, Perth, Western Australia, 30 pp.
- Department of Water (2009) Whicher area surface water allocation plan. Water resource allocation planning series, Report No. 19, Department of Water, Perth, Western Australia, 74 pp.
- Department of Water (2010) South West regional water plan: Supporting detail. Department of Water, Perth, Western Australia, 135 pp.
- Donohue R, Green A, Pauli N, Storey A, Lynas J, Bennett K (2010) Ecological Water Requirements of Cowaramup Brook. Department of Water, Government of Western Australia, Environmental Water Report No. 10, 56 pp.
- Duffy R, Ledger J, Dias J, Snow M (2014) The critically endangered hairy marron, *Cherax tenuimanus* Smith, 1912: A review of current knowledge and actions required to prevent extinction of a species. *Journal of the Royal Society of Western Australia* 97:297–306.
- Fletcher AR, Morison AK, Hume DJ (1985) Effects of carp (*Cyprinus carpio* L.) on aquatic vegetation and turbidity of waterbodies in the lower Goulburn River Basin. *Australian Journal of Marine and Freshwater Research* 36:311–327.
- Gill HS, Hambleton SJ, Morgan DL (1999) Is *Gambusia holbrooki* a major threat to the native freshwater fishes of south-western Australia? *In*: Proceedings 5th Indo-Pacific Fish Conference (Noumea, 3-8 November 1997) (eds B Seret, J -Y Sire) pp. 79–87. Paris: Societe Francaise d'Ichtyologie and Institut de Recherche pour le Development.
- Green A, Donohue R, Storey A, Lynas J, Pauli N (2010) Ecological water requirements of the Margaret River. Environmental water report series, Report No. 11, Department of Water, Western Australia, 72 pp.
- Hope P, Abbs D, Bhend J, Chiew F, Church J, Ekström M, Kirono D, Lenton A, Lucas C, McInnes K, Moise A, Monselesan D, Mpelasoka F, Timbal B, Webb L, Whetton P (2015) Southern and South-Western Flatlands Cluster Report, Climate Change in Australia Projections for Australia's Natural Resource Management Regions: Cluster Reports, eds. Ekström, M., Whetton, P., Gerbing, C., Grose, M., Webb, L., Risbey, J. CSIRO and Bureau of Meteorology, Australia.



## Threatened fauna of Australia's Cape to Cape region

DOI: 10.29094/FiSHMED.2017.002

- Horwitz P (1990) The translocation of freshwater crayfish in Australia: potential impact, the need for control and global relevance. *Biological Conservation* 54:291–305.
- Horwitz P, Adams M (2000) The systematics, biogeography and conservation status of species in the freshwater crayfish genus *Engaewa* Riek (Decapoda: Parastacidae) from south-western Australia. *Invertebrate Taxonomy* 14:655–680.
- Hunt K, Oldham C, Sivapalan M, Smettem K (2002) Stream condition in the Cape to Cape subregion, southwest Western Australia. Centre for Water Research, University of Western Australia.
- IPCC (2013) *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1535 pp.
- IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* [Barros, V.R., C.B. Field, D.J. Dokken, M.D. Mastrandrea, K.J. Mach, T.E. Bilir, M. Chatterjee, K.L. Ebi, Y.O. Estrada, R.C. Genova, B. Girma, E.S. Kissel, A.N. Levy, S. MacCracken, P.R. Mastrandrea, and L.L. White (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 688.
- Klunzinger MW, Beatty SJ, Morgan DL, Lymbery AJ, Pinder AM, Cale DJ (2012) Distribution of *Westralunio carteri* Iredale, 1934 (Bivalvia: Unionoida: Hyriidae) on the south coast of south-western Australia, including new records of the species. *Journal of the Royal Society of Western Australia* 95:77–81.
- Klunzinger MW, Haag WR, Beatty SJ, Morgan DL, Lymbery AJ (2015) Range decline and conservation status of *Westralunio carteri* Iredale, 1934 (Bivalvia: Hyriidae) from south-western Australia. *Australian Journal of Zoology* 63:127–135.
- Koehn JD, Crook DA (2013) Movements and migration. In: *Ecology of Australian Freshwater Fishes* (eds P Humphries and K Walker), pp. 105–129. CSIRO Publishing, Victoria, Australia.
- Lucas MC, Baras E (2001) *Migration of Freshwater Fishes.* Oxford: Blackwell Science, 420 pp.
- Lymbery AJ, Hassan M, Morgan DL, Beatty SJ, Doupe RG (2010). Parasites of native and exotic freshwater fishes in south-western Australia. *Journal of Fish Biology* 76:1770–1785.
- Lynas J, Storey A, Knott B (2007) Introduction and spread of crayfish (Parastacidae) in Western Australia and their potential to displace indigenous species. In: *Biological invaders in inland waters: Profiles, distribution, and threats.* Vol. 2 (ed F Gherardi), Springer, Netherlands, pp. 577–596.
- Molony BW, Wilkinson IS, Montes B (2004) Draft interim Recovery Plan for *Cherax tenuimanus* Smith. Report to the Western Australian Department of Conservation and Land Management.
- Morgan D, Beatty S (2003) Fish fauna of Margaret River Western Australia. Centre for Fish and Fisheries Research, Murdoch University Report to the Margaret River Regional Environment Centre, 15 pp.
- Morgan D, Beatty S (2004a) Margaret River Fishway. Report to the Margaret River Regional Environment Centre, 23 pp.

## Threatened fauna of Australia's Cape to Cape region

DOI: 10.29094/FiSHMED.2017.002

- Morgan D, Beatty S (2004b) Fish fauna of the Vasse River and the colonisation by feral goldfish (*Carassius auratus*). Centre for Fish & Fisheries Research, Murdoch University Report to the Department of Environment, Government of Western Australia.
- Morgan D, Beatty S (2005) Fish and Crayfish Fauna of Ellen Brook, Cowaramup Brook and Gunyulgup Brook in the Cape to Cape Region of Western Australia. Report to Ribbons of Blue/Waterwatch WA. Centre for Fish and Fisheries Research, Murdoch University, WA, 38 pp.
- Morgan DL, Beatty SJ (2007a) Fish migrations patterns on the Margaret River fishways: 2006. Centre for Fish and Fisheries Research (Murdoch University) report to Department of Water and Cape to Cape Catchments Group.
- Morgan DL, Beatty SJ (2007b) Feral Goldfish (*Carassius auratus*) in Western Australia: a case study from the Vasse River. *Journal of the Royal Society of Western Australia* 90:51–156.
- Morgan DL, Beatty SJ (2008) Fish and freshwater crayfish of Boodjidup Brook, south-western Australia. Centre for Fish and Fisheries Research, Murdoch University Report to Cape to Cape Catchments Group, 16 pp.
- Morgan DL, Beatty SJ, Allen MG (2013) Fishes and crayfishes of Turner Brook: Past and present. Freshwater Fish Group & Fish Health Unit (Murdoch University). Report to the Cape to Cape Catchments Group.
- Morgan DL, Beatty SJ, Allen MG, Keleher J, Moore G (2014b) Long live the King River Perchlet (*Nannatherina balstoni*). *Journal of the Royal Society of Western Australia* 97: 307–312.
- Morgan DL, Beatty SJ, Klunzinger MW, Allen MG, Burnham QF (2011) A field guide to freshwater fishes, crayfishes and mussels of south-western Australia. South East Regional Centre for Urban Landcare (SERCUL), Perth, Western Australia, 72 pp.
- Morgan DL, Gill HS, Maddern MG, Beatty SJ (2004) Distribution and impacts of introduced freshwater fishes in Western Australia. *New Zealand Journal of Marine and Freshwater Research* 38:511–523.
- Morgan DL, Gill HS, Potter IC (1998) Distribution, identification and biology of freshwater fishes in south-western Australia. *Records of the Western Australian Museum Supplement* 56:1–97.
- Morgan DL, Hambleton SJ, Gill HS, Beatty SJ (2002) Distribution, biology and likely impacts of the introduced redfin perch (*Perca fluviatilis*) (Percidae) in Western Australia. *Marine and Freshwater Research* 53:1211–1221.
- Morgan DL, Thorburn DC, Gill HS (2003) Salinization of south-western Western Australian rivers and the implications for the inland fish fauna – the Blackwood River, a case study. *Pacific Conservation Biology* 9:161–171.
- Morgan DL, Unmack PJ, Beatty SJ, Ebner BC, Allen MG, Keleher JJ, Donaldson JA, Murphy J (2014a) An overview of the 'freshwater fishes' of Western Australia. *Journal of the Royal Society of Western Australia* 97:263–278.
- Morrongiello JR, Beatty SJ, Bennett JC, Crook DA, Ikedife DNEN, Kennard MJ, Kerezszy A, Lintermans M, McNeil DJ, Pusey BJ, Rayner T (2011) Climate change and its implications for Australia's freshwater fish. *Marine and Freshwater Research* 62:1082–1098.
- Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. *Nature* 403:853–858.
- Olson DM, Dinerstein E (2002) The global 200: priority ecoregions for global con-

**Threatened fauna of Australia's Cape to Cape region**  
**DOI: 10.29094/FiSHMED.2017.002**

- ervation. *Annals of the Missouri Botanical Garden* 89:199–224.
- Pen L (1999) *Managing our rivers: A guide to the nature and management of the streams of south-west Western Australia*. Water and Rivers Commission, Perth.
- Pen LJ, Potter IC, Hilliard RW (1991) Biology of *Galaxiella munda* McDowall (Teleostei: Galaxiidae), including a comparison of the reproductive strategies of this and three other local species. *Journal of Fish Biology* 39:717-731.
- Rowe DK, Moore A, Giorgetti A, Maclean C, Grace P, Wadhwa S, Cooke J (2008) Review of the impacts of gambusia, redfin perch, tench, roach, yellowfin goby and streaked goby in Australia. Prepared for the Australian Government Department of the Environment, Water, Heritage and the Arts.
- Silberstein RP, Aryal SK, Durrant J, Pearcey M, Braccia M, Charles SP, Boniecka L, Hodgson GA, Bari MA, Viney NR, McFarlane DJ (2012) Climate change and runoff in south-western Australia. *Journal of Hydrology* 475:441–455.
- Suppiah R, Hennessy KJ, Whetton PH, McInnes K, Macadam I, Bathols J, Ricketts J, Page CM (2007) Australian climate change projections derived from simulations performed for the IPCC 4<sup>th</sup> assessment report. *Australian Meteorological Magazine* 131:131–152.
- Tay MY, Lymbery AJ, Beatty SJ, Morgan DL (2007) Predation by Rainbow Trout (*Oncorhynchus mykiss*) on a Western Australian icon: Marron (*Cherax cainii*). *New Zealand Journal of Marine and Freshwater Research* 41:197–204.
- Taylor S, Tinley K (1999) Yallingup Brook Action Plan. Report for the Geographie Catchment Council – GeoCatch and the Yallingup LCDC.
- Tille P, Lantzke N (1990) Busselton - Margaret River - Augusta land capability study: methodology and results. Department of Agriculture and Food, Western Australia, Perth
- Unmack PJ (2001) Biogeography of Australian freshwater fishes. *Journal of Biogeography* 28:1053–1089.
- Unmack PJ (2013) Biogeography. In: *Ecology of Australian Freshwater Fishes* (eds P Humphries and K Walker). CSIRO Publishing, Collingwood, pp. 25–48.