

# TAXONOMIC REVIEW OF SELECTED INVERTEBRATE GROUPS COLLECTED DURING THE CAMPAIGNS OF THE PRINCE ALBERT I OF MONACO IN THE AZOREAN WATERS

CRISTINA GARCÍA-DIEZ, FILIPE M. PORTEIRO, ANA MEIRINHO, FREDERICO CARDIGOS & FERNANDO TEMPERA

GARCÍA-DIEZ, C., F.M. PORTEIRO, A. MEIRINHO, F. CARDIGOS & F. TEMPERA 2005. Taxonomic review of selected invertebrate groups collected during the Campaigns of the Prince Albert I of Monaco in the Azorean waters. *Arquipélago. Life and Marine Sciences* 22A: 35-59.

In the late 19th and early 20th centuries Prince Albert I of Monaco promoted 13 cruises that sampled the Azorean waters. During those cruises a total of 2624 nominal marine species were reported in the area. This work assembles the biological data provided by these expeditions to the Azores in a geo-referenced database. Faunal lists for seven invertebrate groups (Echiura, Sipuncula, Cephalopoda, Annelida, Brachiopoda, Chaetognatha and Echinodermata) are compiled. The checklist includes 331 nominal species, of which 310 are valid names: 1 echiurid; 11 sipunculids; 32 cephalopods; 130 annelids; 4 brachiopods; 14 chaetognaths; and 118 echinoderms. Eighteen percent are synonyms, 29% of the species were allocated to a different genus, 2.8% were misspellings and corrections due to gender or concordance rules, 0.8% were specific epithets allocated to sub-specific level or vice-versa, the rest were validated directly (without any modification in their nomenclature).

Cristina García-Diez (e-mail: [cristigar10@yahoo.es](mailto:cristigar10@yahoo.es)), Centre of Biodiversity and Environmental Management, Faculty of Marine Sciences, University of Las Palmas de Gran Canaria, ES-35017 Las Palmas de G.C., Spain; Filipe M. Porteiro, A. Meirinho, F. Cardigos & F. Tempera Department of Oceanography and Fisheries, University of the Azores, PT-9901-862 Horta, Portugal.

## INTRODUCTION

From 1885 to 1914 HRH Prince Albert I of Monaco (PAM) promoted 13 cruises which sampled the Azorean waters. During those cruises many biological samples were collected. About 50 types of collecting gear were used to prospect the marine life from coastal waters to the open ocean, as well as from the surface down to 5000 m depth. Eight hundred forty-three sampling locations were completed and 2624 nominal marine species reported across this region.

The results of these activities were published by an international bureau of more than 70 scientists in 110 volumes (*Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ier Prince Souverain du Monaco*) between 1889 and 1950.

The scientific heritage from these works

comprises the most comprehensive inventories of benthic and pelagic fauna living in the Azorean waters. The investigations covered from bacteria and phytoplankton to marine mammals, including the most important groups of invertebrates. However, only a few of the high level taxonomic groups reported in the Prince's expeditions have been subsequently reviewed. The records of fish caught during those campaigns have been critically studied, synonymised and assembled in a checklist of the fishes from the Azores (SANTOS et al. 1997). Available information for some other groups have also been updated (e.g. gastropods, ÁVILA 2000; octopods, GONÇALVES 1991; barnacles, SOUTHWARD 1998, and YOUNG 1998, 2001). However, the majority of the recent inventories of Azorean marine fauna deal with specific collections and normally do not provide reviews of historical records. This is the case of

works such as CHAPMAN & DALES (1954) and BELLAN (1978), who listed 27 and 156 species of annelid worms from the coastal waters of the Azores, respectively; and CHAPMAN (1955), who collected three sipunculids in the Azores. Beyond that we only find other references in general field guides (i.e. SALDANHA 1995; WIRTZ 1995; WIRTZ & DEBELIUS 2003). A checklist of echinoderms (41 species) was compiled by PEREIRA (1997) but it covers only the littoral zone. In groups such as chaetognaths, echiurids and brachiopods the situation is even worst as there are no lists of the species inhabiting the region.

The compilation of species records of different origins and times poses several challenges. Revisiting of the samples kept in different museums is not always possible in the short-term and requires expensive and time-consuming work by experts. An alternative based on the use of literature, web resources and e-mail consultation of specialists was investigated. This paper resumes a tentative assessment of the work involved to try establishing the synonymy of historical records by using the database of the results of Prince Albert I of Monaco expeditions as an example. Works like this are expected to become more and more required as individual datasets need to be integrated into broader consolidated databases such as the ones informing assessments of long-term, large-scale patterns of the distribution of marine organisms. This has already been initiated in European waters through project Biomare and is currently under the scope of the MARBEF network (more info: <http://www.pml.ac.uk/biomare>; <http://www.marbef.org/>). A series of Azorean marine areas were selected as reference sites for the network. An *All Taxon Biodiversity Inventory* (ATBI) is expected for sites in which catalogues are already available for a large number of components of the biota.

In order to compile everything that is known about the biodiversity of these sites (e.g., the channel between the Azorean islands of Faial and Pico), some data-mining still needs to be done regarding the many species that have been recorded since they started to be visited by historical scientific expeditions. The major achievement so far consisted of building a database with all the species recorded in Azorean waters by the Prince Albert of Monaco

expeditions and published along the range of 110 volumes. The PAM database includes species occurrences, meta-information about the location, the sampling strategy and some basic characterization of the place where the collections were done.

The extensive collections performed by these campaigns in the waters surrounding the archipelago are still a valuable source of information for many taxonomic groups. However, the fact that this information was only in paper format has made it quite cryptic to modern scientific search engines and prevented an overall assessment of what is known for the region and for particular locations.

We present faunal lists for seven invertebrate groups: Echiura, Sipuncula, Cephalopoda, Annelida, Brachiopoda, Chaetognatha and Echinodermata. These particular groups were selected because they are poorly known in the area, especially their deep-sea forms.

In the absence of historical sample revisiting by taxonomy experts, interim synonymy is assumed as a temporary solution preventing overestimations of biodiversity.

## MATERIAL AND METHODS

A database was created to assemble and store the information collected by the PAM cruises in the Azorean waters, by including biological data (species occurrences), meta-data on the stations (e.g. cruise, vessel, date, latitude, longitude, depth, gear operated) and bibliographic reference. The Azores region was defined as the area confined by a rectangle involving the 200 miles Economic Exclusive Zone ( $33^{\circ} 46'N$  -  $42^{\circ} 57'N$ ,  $035^{\circ} 45'W$  -  $021^{\circ} 5'W$ ; Fig.1). The data used in this contribution were retrieved from the database. The sampling locations that produce the material treated in this checklist are plotted in Figure 1 and summarised in Annex 1.

Although more than 90% of the sampling locations were defined by their geographical position (RICHARD 1934), some are referred only by local names (e.g. Baía de Porto Pim, Faial; Santa Cruz, Flores). In those cases the probable latitude and longitude of the sample sites have been estimated from official nautical charts

[Instituto Hidrográfico: 113 (1987), 46401 (2002), 46403 (1999), 46405 (2001), 46406 (2001); 46407 (2000)] in a GIS environment.




Fig 1. Area selected: rectangle involving the 200 miles Economic Exclusive Zone (ZEE). Sampling locations around the Azores region.

The species occurrences were compiled from the summary tables (by sampling location) presented at the end of the species accounts in the *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ier Prince Souverain du Monaco*. All the 110 volumes have been reviewed. However, the data compiled in this paper was originally treated in 15 volumes of the above mentioned publications (Sipuncula and Echiura [SLUITER 1900, 1912], Mollusca-Cephalopods [JOUBIN 1895, 1900, 1920, 1924], Annelida [FAUVEL 1914, 1916, 1932], Brachiopoda [FISCHER & OEHLMER 1892], Chaetognatha [GERMAIN & JOUBIN 1916] and Echinodermata [HÈROUARD 1902; KOEHLER 1898, 1909; PERRIER 1896]).

The checklist is organised by phylum, and then by infra-phylum taxa to species (e.g. superclass, class, subclass, superorder, order, suborder, family and genus). The nomenclature adopted follows the NEAT initiative (North East Atlantic Taxa; [www.tmbi.gu.se/libdb/taxon/taxa.html](http://www.tmbi.gu.se/libdb/taxon/taxa.html); HANSSON 1997, 1998). To highlight the taxa level the following code was adopted: **PHYLUM**, **SUPERCLASS**, **Class**, Subclass, **SUPERORDER**, **Order**, **SUBORDER**, FAMILY, Genus & species.

Under each species entry there is the reference number(s) of the PAM sampling **station(s)** **location(s)** that produced the record(s). Those numbers linked to Annex 1, where sampling location data are available.

The following method was used to validate or synonymised a nominal name presented in the PAM faunal lists. If a species name is considered valid in most of the inventories used for reference (see above), then it was accepted as valid species for the area.

If a nominal species name listed in the database is not considered valid at present, a bibliographic and a web based search was made to find the valid name; then the scientific names used in PAM were relegated to synonyms. Synonyms were found in CUTLER (1994) and HANSSON (1997) for Sipuncula; CLARKE (1996), GONÇALVES (1991) and [www.cephbase.dal.ca](http://www.cephbase.dal.ca) for Cephalopoda; COSTELLO et al. (2001), HANSSON (1998), BELLAN (2001) and KATO & PLEIJEL (2002) for Annelida; LOGAN (1998) and HOWSON (2001) for Brachiopoda; VAN DER LAND & KAPP (2001) and HANSSON (1997) for Chaetognatha; and HANSSON (1998, 2001), MÜLLER (1999) and PEREIRA (1997) for Echinodermata. Those contributions which are part of the European Register of Marine Species (ERMS, COSTELLO et al. 2001) were also searched on the world wide web. Other web faunal inventories and resources were also consulted such as: *La Fauna Ibérica* for Annelida and Brachiopoda (ALVAREZ 2004); *The Echinoid Directory* for Echinodermata (SMITH 2003); and the website of the Muséum National d'Histoire Naturelle, Paris for Echinodermata (AMEZIANE 1999).

Synonyms and misspelling of nominal names are preceded by = and are compiled under each species entry.

Biological material that was identified to genus or species in PAM reports, which could not be found in the taxonomic resources used, are listed in a section at the end of each **group** as "uncertain species".

The following checklist includes 310 current valid names: 1 echiurid; 11 sipunculids; 32 cephalopods; 130 annelids; 4 brachiopods; 14 chaetognaths; and 118 echinoderms.

## CHECKLIST

### ECHIURA

#### Echiuroidea

##### Bonelliida

###### BONELLIIDAE

*Bonellia viridis* Rolando, 1821

Station: 866

Station: 2210

*Nephasoma diaphanes diaphanes* (Gerould, 1913) = *Phascolosoma minutum* Théel, 1911

Station: 663

*Thysanocardia catharinae* (Grube, 1868) = *Phascolosoma catharinae* Grube, 1868

Station: 226

### SIPUNCULA

#### Phascolosomatidea

##### Aspidosiphonida

###### ASPIDOSIPHONIDAE

*Aspidosiphon muelleri* Diesing, 1851

Stations: 196, 198, 216, 594, 882, 2214

#### PHASCOLIONIDAE

*Phascolion strombus strombus* (Montagu, 1804) = *Phascolion strombi* Montagu, 1804

Station: 858

*Phascolion tuberculatum* Théel, 1875 = *Phascolion hirondellei* Sluiter, 1900

Stations: 112, 184, 203, 211, 233, 244, 575, 624, 683, 684, 703, 719, 743, 832, 1311, 1334, 2214

### MOLLUSCA

#### Cephalopoda

##### Coleoidea

##### Spirulida

###### SPIRULIDAE

*Spirula spirula* (Linnaeus, 1758) = *Spirula peroni* Lamarck, 1822

Station: 603

#### Sepiolida

###### SEPIOIDAE

*Rossia macrosoma* (Chiaie, 1830) = *Rossia* sp.

Station: 211

*Heteroteuthis dispar* (Rüppell, 1844)

Stations: 554, 3256

#### Teuthida

###### ANCISTROCHEIRIDAE

*Ancistrocheirus lesueuri* (Orbigny, 1842)

Station: 588

###### ARCHITEUTHIDAE

*Architeuthis* sp. Steenstrup, 1857 = *Dubiotheuthis physeteris* (Joubin, 1900)

Stations: 244, 588

#### Sipunculidea

##### Sipunculida

###### SIPUNCULIDAE

*Sipunculus norvegicus* Danielssen, 1868

Station: 553

= *Sipunculus nitidus* Sluiter, 1900

Station: 650

#### Golfingida

###### GOLFINGIIDAE

*Golfingia vulgaris vulgaris* (Blainville, 1827) = *Phascolosoma vulgare* Blainville, 1827

Station: 198

*Golfingia (Golfingiella) innoxia* Sluiter (in Stephen & Edmonds, 1972), 1912 = *Phascolosoma innoxium* Sluiter, 1900

BATHYTEUTHIDAE	MASTIGOTEUTHIDAE
<i>Bathyteuthis abyssicola</i> Hoyle, 1885 =	<i>Mastigoteuthis grimaldii</i> (Joubin, 1895) =
<i>Benthoteuthis megalops</i> Verrill, 1885	<i>Chiroteuthis grimaldii</i> Joubin, 1895
Station: 3131	Stations: 220, 3131, 3526
BRACHIOTEUTHIDAE	OCTOPOTEUTHIDAE
<i>Brachiotheuthis beani</i> Verrill, 1881	<i>Octopoteuthis sicula</i> Rüppell, 1844 =
Station: 3147	<i>Octopodoteuthis sicula</i> Krohn, 1845
CHIROTEUTHIDAE	Station: 2172
<i>Grimalditeuthis bonplandi</i> (Verany, 1839) =	<i>Tanigia danae</i> Joubin, 1931 = <i>Cucioteuthis</i>
<i>Grimalditeuthis richardi</i> Joubin, 1898	<i>unguiculata</i> Joubin, 1898
Station: 750	Station: 588
CRANCHIIDAE	OMMASTREPHIDAE
<i>Cranchia scabra</i> Leach, 1817	<i>Todarodes sagittatus</i> (Lamarck, 1798) =
Station: 2153	<i>Ommatostrephes sagittatus</i> Pfeffer, 1908
<i>Liocranchia reinhardtii</i> (Steenstrup, 1856) =	Station: 3279
<i>Fusocranchia alpha</i> Joubin, 1920	<i>Ommastrephes bartrami</i> (Lesueur, 1821) =
Stations: 1856, 1872	<i>Stenoteuthis bartrami</i> Lesueur, 1821
<i>Galiteuthis armata</i> Joubin, 1898	Station: 2184
Station: 3147	ONYCHOTEUTHIDAE
<i>Teuthowenia megalops</i> (Prosche, 1847)	<i>Onychoteuthis banksii</i> (Leach, 1817)
Stations: 1849, 1874, 2187	Stations: 576, 1874
ENOPLOTEUTHIDAE	PYROTEUTHIDAE
<i>Abraliopsis lineata</i> (Goodrich, 1896) =	<i>Pyroteuthis margaritifera</i> (Rüppell, 1844)
<i>Abralia morisii</i> (Verany, 1839)	Stations: 3147, 3150, 3279
Station: 3131	<b>Octopoda</b>
HISTIOTEUTHIDAE	OPISTHOTEUTHIDAE
<i>Histioteuthis bonnellii</i> (Férussac, 1834) =	<i>Opistoteuthis grimaldii</i> (Joubin, 1903) =
<i>Histioteuthis bonelliana</i> (Férussac, 1834) =	<i>Cirroteuthis grimaldii</i> Joubin, 1903
<i>Histioteuthis ruppelli</i> Vérany, 1835	Station: 1339
Stations: 588, 3279	BOLITAENIDAE
<i>Histioteuthis meleagroteuthis</i> (Chun, 1910) =	<i>Japetella diaphana</i> Hoyle, 1885 = <i>Eledonella</i>
<i>Meleagroteuthis hoylei</i> (Goodrich, 1896)	<i>diaphana</i> Verrill, 1884 = <i>Bolitaena</i>
Station: 3285	<i>diaphana</i> (Verrill, 1884)
<i>Histioteuthis reversa</i> (Verrill, 1880) =	Station: 2168, 3131
<i>Calliteuthis meneghini</i> Pfeffer, 1912	OCTOPODIDAE
Station: 3279	<i>Octopus vulgaris</i> Cuvier, 1797 = <i>Octopus</i>
LEPIDOTEUTHIDAE	<i>granulatus</i> Lamarck, 1799
<i>Lepidoteuthis grimaldii</i> Joubin, 1895	Station: 882
Stations: 588, 849	<i>Octopus macropus</i> Risso, 1826
	Station: 225

- Pteroctopus tetricirrus* (Delle Chiaie, 1830)  
   = *Scaergus tetricirrus* Tiberi, 1830  
 Station: 866
- Benthocotopus levis* (Hoyle, 1885) = *Octopus levis* Houye, 1885  
 Station: 719
- VITRELEDONELLIDAE**  
*Vitreledonella richardi* Joubin, 1918 =  
   *Vitreledonella alberti* Joubin, 1924  
 Station: 3131
- Vampyromorphida**
- VAMPYROTEUTHIDAE**  
*Vampyroteuthis infernalis* Chun, 1903 =  
   *Melanoteuthis lucens*  
 Station: 2168
- Uncertain species (cephalopods)**
- Abraliopsis* sp.  
 Stations: 1834, 3526
- Architeutis* sp.  
 Stations: 244, 588
- Mastigoteuthis* sp.  
 Stations: 1334, 1408, 2148
- ANNELIDA**
- POLYCHAETA**
- Palpata**  
 Aciculata
- Phyllodocida**
- PHYLLODOCIDAE**  
*Nereiphylla rubiginosa* (Saint-Joseph, 1888) =  
   *Phyllodoce rubiginosa* Saint-Joseph  
 Station: 226
- Notophyllum foliosum* (Sars, 1835)  
 Station: 1373
- Phyllodoce madeirensis* Langerhans, 1880  
 Stations: 234, 584, 594, 866, 2214
- Eulalia viridis* (Linnaeus, 1767)  
 Station: 216
- ALCIOPIDAE**  
*Vanadis formosa* Claparède, 1870
- Station: 2194
- Rhynchonereella angelini* (Kinberg, 1866) =  
   *Callizona angelini* Kinberg  
 Stations: 1851, 1874, 2153, 2185, 2187, 2194,  
   2200, 2212, 3526
- Rhynchonereella petersii* (Langerhans, 1880)  
   = *Callizona setosa* Apstein, 1900  
 Stations: 1872, 3132
- TOMOPTERIDAE**  
*Tomopteris elegans* Chun, 1888  
 Stations: 1849, 1851, 2187, 2194, 2212
- Tomopteris (Johnstonella) apsteini* Rosa,  
 1908  
 Station: 1844
- Tomopteris (Tomopteris) ligulata* Rosa, 1908  
 Stations: 1849, 2168, 2194
- Tomopteris (Tomopteris) nisseni* Rosa, 1908  
 Station: 1849
- Tomopteris (Tomopteris) planktonis* Apstein,  
 1900  
 Stations: 2185, 2187, 2212
- Tomopteris (Tomopteris) septentrionalis*  
 Quatrefages, 1866  
 Station: 1805
- LOPADORHYNCHIDAE**  
*Lopadorhynchus appendiculatus* Southern,  
 1909  
 Station: 2159
- Lopadorhynchus uncinatus* Fauvel, 1915  
 Station: 1856
- Pelagobia longicirrata* Greeff, 1879  
 Stations: 17, 1849, 2185, 2244
- Pelagobia serrata* Southern, 1909  
 Station: 2212
- Pedinosoma curvum* Reibich, 1895  
 Station: 1839

IOSPILIDAE

*Phalacrophorus pictus pictus* Greeff, 1879 =  
*Phalacrophorus pictus* Greeff

Station: 1805

*Phalacrophorus uniformis* Reibich, 1875  
Station: 2218

TYPHLOSCOLEDIDAE

*Typhloscolex phylloides* Reibich, 1895 =  
*Typhloscolex phyllodes* Reibich, 1895  
Stations: 2212, 2244

*Sagitella kowalewskii* Wagner, 1872  
Stations: 1805, 2168, 2187, 2194, 2200, 2212

*Travisiopsis lanceolata* Southern, 1910  
Stations: 1851, 1874, 2168, 2187, 2200, 3526,  
3608

*Travisiopsis levinseni* Southern, 1910  
Stations: 2185, 2244

GLYCERIDAE

*Glycera lapidum* Quatrefages, 1866  
Stations: 112, 832, 874

*Glycera tesselata* Grube, 1863  
Stations: 226, 234, 616, 702, 866, 1344, 1349,  
1373, 2210, 2214

*Glycerella magellanica* (McIntosh, 1885)  
Stations: 229, 234, 584, 2211

HESIONIDAE

*Leocrates atlanticus* (McIntosh, 1885)  
Station: 198, 578, 587, 597, 616, 837, 1344,  
2214

*Dalhousiella carpenteri* McIntosh, 1901 =  
*Dalhousasiella carpenteri* McIntosh, 1901  
Station: 838

*Nereimyra punctata* (Müller, 1776) = *Castalia*  
*punctata* Müller, 1776  
Stations: 184, 213, 832, 1311, 2214

SYLLIDAE

*Eusyllis monilicornis* Malmgren, 1867  
Station: 226

*Myrianida brachycephala* (Marenzeller, 1874)  
= *Autolytus brachycephala* Marenzeller,  
1874  
Station: 227

*Opisthosyllis brunnea* Langerhans, 1879  
Station: 216

*Pionosyllis weissmanni* Langerhans, 1879 =  
*Pionosyllis weissmanni* Langerhans, 1879  
Station: 226

*Pseudosyllis brevipennis* (Grube, 1863) =  
*Syllis brevipennis* Grube, 1863  
Station: 226

*Typosyllis armillaris* (Müller, 1776) = *Syllis*  
*alternosetosa* Saint-Joseph, 1887  
Station: 226

*Syllis gracilis* Grube, 1840  
Station: 226

*Syllis cornuta* Rathke, 1843  
Stations: 184, 238, 244, 743, 2214

*Syllis krohnii* Ehlers, 1864  
Station: 191

*Syllis prolifera* Krohn, 1852  
Station: 1412

*Syllis variegata* Grube, 1860  
Station: 226

*Syllis vittata* (Grube, 1840)  
Stations: 195, 202, 622

*Trypanosyllis coeliaca* Claparède, 1868  
Station: 226

*Trypanosyllis gigantea* McIntosh, 1885  
Station: 584

- NEREIDIDAE**
- Neanthes kerguelensis* (McIntosh, 1885) =  
*Nereis kerguelensis* McIntosh, 1885  
 Stations: 226, 234, 597, 749
- Neanthes irrorata* (Malmgren, 1867) = *Nereis irrorata* Malmgren, 1867  
 Stations: 193, 842
- Nereis pelagica* Linnaeus, 1758  
 Station: 226
- Nereis rava* Ehlers, 1868  
 Stations: 226, 234, 866, 1373, 3293
- Perinereis cultrifera* (Grube, 1840)  
 Station: 216
- Platynereis coccinea* (Delle Chiaje, 1841)  
 Station: 216
- Platynereis dumerilli* (Audouin & H. Milne-Edwards, 1834)  
 Stations: 226, 1311, 1373
- NEPTYIDAE**
- Nephtys hystricis* McIntosh, 1900 = *Nephthys hystricis* McIntosh, 1900  
 Station: 673
- APHRODITIDAE**
- Aphrodisia aculeata* Linnaeus, 1758 =  
*Aphrodite aculeata* Linnaeus, 1758  
 Stations: 3150, 576, 866
- Pontogenia sericoma* Ehlers, 1867  
 Station: 594
- POLYNOIDAE**
- Polynoe antillicola* (Augener, 1906)  
 Station: 738
- Acanthicolepis asperrima* (Sars, 1861)  
 Stations: 616, 618, 702, 838, 869, 1349
- Adyte assimilis* (McIntosh, 1876) =  
*Scalisetosus assimilis* McIntosh, 1876  
 Station: 2214
- Malmgrenia andreapolis* McIntosh, 1874 =  
*Harmothoe lunulata* Delle Chiaje, 1841  
 Stations: 193, 196, 218
- Malmgrenia glabra* (Malmgren, 1865) =  
*Nereis longisetis* McIntosh, 1885  
 Station: 683
- Subadyte pellucida* (Ehlers, 1864) =  
*Scalisetosus pellucidus* Ehlers, 1864  
 Station: 614
- Harmothoe echinopustulata* Fauvel, 1913  
 Station: 889
- Harmothoe imbricata* (Linnaeus, 1767)  
 Station: 594
- Harmothoe impar* (Johnston, 1839)  
 Stations: 226, 578, 584, 702, 866, 2210, 2214
- Harmothoe johnstoni* (McIntosh, 1876)  
 Stations: 213, 837, 874, 1412, 1849, 1851
- Harmothoe spinifera* (Ehlers, 1864) = *Scione spinifera* Ehlers, 1864  
 Station: 1344, 1349
- Lagisca talismani* (Roule, 1898)  
 Stations: 105, 244, 584, 587, 597, 837, 838, 874, 1344
- Lepidasthenia maculata* Potts, 1910  
 Station: 616, 873
- Lepidonotus clava* (Montagu, 1808)  
 Station: 225
- Macellicephalia incerta* Fauvel, 1914  
 Station: 2185
- Macellicephalia mirabilis* (McIntosh, 1885)  
 Station: 1344
- Nectochaeta grimaldii* Marenzeller, 1892  
 Station: 2194
- Robertianella synophthalma* McIntosh, 1885  
 Stations: 198, 1344, 2214

PHOLOIDAE

*Pholoe synopthalmica* Claparède, 1868  
Station: 838

*Pholoides dorsipapillata* (Marenzeller, 1893)  
= *Pholoe dorsipapillata* Marenzeller, 1893  
Stations: 226, 1349

SIGALIONIDAE

*Neoleanira tetragona* (Örsted, 1845) =  
*Leanira tetragona* Örsted, 1845  
Station: 581

*Eunicida*

AMPHINOMINA

AMPHINOMIDAE

*Amphinome rostrata* (Pallas, 1776) =  
*Amphinome pallasii* Quatrefages, 1865  
Stations: 699, 2215

*Hermodice carunculata* (Pallas, 1766)  
Stations: 103, 225, 238, 243, 1355

*Hipponoe gaudichaudi* Audouin & Milne-  
Edwards, 1830  
Stations: 181, 182, 536, 545, 699

EUPHROSINIDAE

*Euphrosine foliosa* Audouin & Milne-  
Edwards, 1834 = *Euphrosyne foliosa*  
Audouin & Milne-Edwards, 1834  
Station: 594

*Palmyreuphrosyne paradoxa* Fauvel, 1914  
Station: 2210

EUNICINA

ONUPHIDAE

*Rhamphobrachium agassizii* Ehlers, 1887  
Stations: 203, 553

*Paradiopatra quadricuspis* (Sars, 1872) =  
*Onuphis quadricuspis* Sars, 1872  
Stations: 553, 703

*Hyalinoecia tubicola* (Müller, 1776)  
Stations: 226, 889

*Nothria conchylega* (Sars, 1835) = *Onuphis*

*conchylega* Sars, 1835

Stations: 198, 234, 527, 536, 553, 575, 602,  
614, 616, 618, 673, 719, 738, 863, 866,  
1334, 1349

EUNICIDAE

*Eunice norvegica* (Linnaeus, 1767) = *Eunice*  
*floridana* (Pourtales, 1867)  
Stations: 227, 584, 837, 838, 1349

*Eunice oerstedi* Stimpson, 1854  
Stations: 584, 587, 838, 2210

*Eunice pennata* (Müller, 1776)

Stations: 112, 198, 213, 226, 227, 616, 663,  
673, 683, 684, 698, 702, 719, 738, 743, 866,  
873, 889, 1331, 1334, 1344, 1349, 3293

*Eunice vittata* (Delle Chiaje, 1829)

Stations: 112, 244, 553, 702, 1349

*Lysidice ninetta* Audouin & Milne-Edwards,  
1833

Stations: 198, 226, 234, 738, 882

LUMBRINERIDAE

*Lumbrineriopsis paradoxa* (Saint-Joseph,  
1888) = *Lumbriconereis paradoxa* Saint-  
Joseph, 1888  
Station: 226

*Lumbrineris coccinea* (Renier, 1804) =  
*Lumbriconereis coccinea* Renier, 1804  
Station: 236

*Lumbrineris funchalensis* (Kinberg, 1865) =  
*Lumbriconereis funchalensis* Kinberg, 1865  
Station: 594

*Scoletoma fragilis* (Müller, 1776) =  
*Lumbriconereis fragilis* Müller, 1776  
Station: 553

OENONIDAE

*Arabella iricolor* (Montagu, 1804) = *Maclovia*  
*iricolor* Montagu, 1804  
Station: 594

- Canalipalpata
- Spionida***
- SPIONIDAE
- Prionospio steenstrupi* Malmgren, 1867  
Stations: 1834, 1872
- POECILOCHAETIDAE
- Poecilochaetus serpens* Allen, 1904  
Stations: 1834, 1860
- CHAETOPTERIDAE
- Spiochaetopterus typicus* Sars, 1856  
Stations: 226, 882
- Magelonidae
- Magelona mirabilis* (Johnston, 1865) =  
*Magelona papillicornis* Müller, 1858  
Stations: 244, 553
- Terebellida***
- PECTINARIIDAE
- Petta pusilla* Malmgren, 1866  
Station: 198
- TEREBELLIDAE
- Amphitrite alcicornis* Fauvel, 1909  
Stations: 112, 616, 873, 874, 1349
- Amphitrite cirrata* Müller, 1771  
Stations: 112, 190, 612, 1344
- Thelepus cincinnatus* (Fabricius, 1780)  
Stations: 226, 587
- Sabellida***
- SABELLARIIDAE
- Phalacrostemma cidariophilum* Marenzeller, 1895  
Stations: 232, 703, 838, 1311, 3144
- SABELLIDAE
- Amphiglena mediterranea* (Leydig, 1851)  
Station: 226
- Jasmineira candela* (Grube, 1863)  
Station: 837
- Megalomma vesiculosum* (Montagu, 1815) =  
*Branchiomma vesiculosum* Montagu, 1815  
Station: 226
- Potamis spathiferus* Ehlers, 1887  
Stations: 702, 874, 2210
- SERPULIDAE
- Hyalopomatus marenzelleri* (Langerhans, 1884) = *Hyalopomatopsis marenzelleri* Langerhans, 1884  
Stations: 184, 584, 703, 1311, 1349, 3144
- Filograna implexa* Berkeley, in M. Sars, 1851  
Station: 226
- Serpula vermicularis* Linnaeus, 1767  
Stations: 703, 882
- Serpula concharum* Langerhans, 1880  
Station: 226
- Spirodiscus grimaldii* Fauvel, 1909  
Stations: 698, 1334
- Hydroides norvegica* Gunnerus, 1768  
Stations: 226, 584
- Protis arctica* (Hansen, 1878)  
Station: 213
- Filogranula calyculata* (Costa, 1861) =  
*Omphalopoma aculeata* Fauvel, 1909  
Station: 866
- Placostegus tridentatus* (Fabricius, 1779)  
Stations: 105, 234, 2214
- Pomatostegus polytrema* (Philippi, 1844)  
Stations: 112, 226, 703, 861, 882, 1349
- Ditrupa arietina* (Müller, 1776)  
Stations: 112, 743, 1334, 1349
- Protula alberti* Fauvel, 1909  
Stations: 198, 587, 866, 2214
- Protula tubularia* (Montagu, 1803)  
Station: 234
- Metavermilia multicristata* (Philippi, 1844) =  
*Vermiliopsis multicristata* Philippi, 1844  
Station: 226

*Vermiliopsis striaticeps* (Grube, 1862) =  
*Vermiliopsis langerhansi* Fauvel, 1903

Station: 527

*Vermiliopsis infundibulum* (Linnaeus, 1788)  
Station: 600

*Circeis spirillum* (Linnaeus, 1758) = *Spirorbis spirillum* Linnaeus, 1758  
Station: 2211

#### SPIORBIDAE

*Dexiospira corrugata* (Montagu, 1803) =  
*Spirorbis corrugatus* Montagu, 1803  
Station: 236

#### Scolecida

##### CAPITELLIDAE

*Notomastus latericeus* Sars, 1850  
Station: 553

##### MALDANIDAE

*Maldane sarsi* Malmgren, 1865  
Station: 2199

##### SCALIBREGMATIDAE

*Asclerocheilus intermedius* (Saint-Joseph, 1894) = *Lipobranchius intermedius* Saint-Joseph, 1894  
Station: 553

#### Uncertain species (Annelida)

*Harmothoe benthophila bimucronata* Fauvel, 1914 = *Harmothoe benthophila* Ehlers

Station: 1856, 2244

Note: the validity of the species needs further evaluation.

*Tharyx* sp.

Station: 232

*Onuphis* sp.

Note: Probably *Nothria conchylega* (Fauvel, 1914).

Station: 527

*Nereis* sp.

Station: 3150

*Spirorbis cornuarietis* Philippi

Station: 236

Note: The species could not be found in recent references

*Chone acustica* Claparède, 1870

Station: 236

Note: The validity of the species needs further clarification.

## BRACHIOPODA

#### Rhynchonellata

##### Terebratulida

##### TEREBRATULIDAE

*Stenosarina sphenoidea* (Jeffreys, 1878) =  
*Terebratula sphenoidea* Philippi, 1844

Station: 233, 242

Note: the status of the species needs further investigation; LOGAN (1998) described *Stenosarina davidsoni* from the "species misidentified by Jeffreys (1878) as *Terebratula vitrea* var. *sphenoidea* [non Philippi, 1844]".

##### DYSCOLIIDAE

*Dyscolia wyvillei* (Davidson, 1878)  
Stations: 203, 213, 227

##### PLATIDIIDAE

*Platidia davidsoni* (Deslongchamp, 1885)  
Station: 234

##### DALLINIDAE

*Dallina septigeria* (Lovén, 1846) =  
*Magellania septigera* Lovén, 1846  
Station: 242

## CHAETOGNATHA

#### Sagittoidea

##### Phragmophora

##### SPAPELLIDAE

*Spadella cephaloptera* (Busch, 1851)  
Station: 185

##### EUKROHNIIIDAE

*Eukrohnia hamata* (Mobius, 1875)  
Stations: 1834, 1844, 1849, 1851, 1856, 1876, 2153, 2168, 2185, 2187, 2194, 2200, 2244

#### Aphragmophora

##### PTEROSAGITTIDAE

*Pterosagitta draco* (Krohn, 1853)  
Stations: 1851, 2149

**SAGITTIDAE**

- Sagitta bipunctata* Quoy & Gaimard, 1828  
Stations: 15, 17, 21, 208, 1844, 1847, 1849, 1851, 1856, 1872, 2149, 2150, 2151, 2153, 2159, 2162, 2168, 2170, 2185, 2187, 2191, 2194, 2198, 2200, 2212, 2241, 2244, 2245, 2249, 2252, 2263, 2267
- Sagitta elegans* (Verrill, 1873)  
Stations: 1322, 2159, 2168, 2185, 2194, 2200, 2244
- Sagitta hexaptera* (d'Orbigny, 1836)  
Stations: 215, 246, 1333, 1844, 1849, 1851, 1856, 1874, 2149, 2153, 2159, 2162, 2168, 2185, 2187, 2194, 2200, 2212, 2244, 2264  
= *Sagitta hexaptera magna* d'Orbigny, 1836  
Stations: 2149, 2185, 2187, 2194, 2200, 2244
- Sagitta enflata* (Grassi, 1881) = *Sagitta inflata* Grassi, 1881  
Stations: 21, 246, 2149
- Sagitta lyra* (Krohn, 1853)  
Stations: 16, 115, 1358, 1844, 1849, 1851, 1856, 1874, 2149, 2153, 2159, 2168, 2185, 2187, 2194, 2200, 2212, 2244, 2264
- Sagitta serratodentata* Krohn, 1853 = *Sagitta serratodenta* Krohn, 1853  
Stations: 16, 1844, 1851, 1860, 2168, 2185, 2187, 2191, 2194, 2195, 2200, 2212, 2242, 2244, 2266
- Sagitta macrocephala* (Fowler, 1904)  
Stations: 1849, 1851, 1856, 1874, 2159, 2168, 2185, 2187, 2200, 2212  
= *Spadella macrocephala* Fowler, 1904  
Station: 2244
- Sagitta minima* (Grassi, 1881)  
Station: 2187
- Sagitta planctonis* (Steinhaus, 1896)  
Stations: 740, 2153, 2185, 2187, 2194, 2200, 2212
- Sagitta tenuis* Conant, 1896  
Station: 2172

**Flabellodontina**

- KROHNITTIDAE  
*Krohnitta subtilis* (Grassi, 1881)  
Stations: 1872, 2187, 2200, 2244

**Uncertain species (Chaetognatha)**

- Sagitta* sp.  
Stations: 18, 115, 116, 125, 1851, 1860, 2143, 2151, 2161, 2171, 2196, 2204, 2266

**ECHINODERMATA****Crinoidea**

- Bourgueticrinida*  
BOURGUETICRINIDAE  
*Democrinus rawsonii* (Pourtalès, 1874) =  
*Rhizocrinus rawsoni* Pourtalès, 1874  
Stations: 683, 738

**BATHYCRINIDAE**

- Comatulida*  
PENTAMETROCRINIDAE  
*Pentametrocrinus atlanticus* (Perrier, 1883) =  
*Eudioecrinus atlanticus* Perrier, 1883  
Station: 578

**Asteroidea**

- Paxillosida*  
ASTROPECTINIDAE  
*Dytaster grandis grandis* (Verrill, 1884) =  
*Dytaster agassizi* Perrier, 1894  
Station: 527

- Dytaster intermedius* Perrier, 1891  
Station: 248

- Persephonaster patagiatus* (Sladen, 1889) =  
*Psilasteropsis patagiatus* Sladen, 1889  
Stations: 575, 684, 719, 858, 1311, 1331, 1334, 1344, 1349, 553

- Persephonaster sphenoplax* (Bell, 1892) =  
*Psilasteropsis humilis* Koehler, 1907  
Stations: 582, 584  
= *Astropecten sphenoplax* Bell, 1892  
Station: 1311

- Plutonaster agassizi notatus* Sladen, 1889 =  
*Plutonaster granulosus* Perrier, 1891

- Stations: 203, 213  
 = *Plutonaster inermis* Perrier, 1885  
 Station: 213  
 = *Plutonaster notatus* Sladen, 1889  
 Stations: 244, 536, 624, 698, 703, 719, 743, 858, 863, 1331, 1334  
 = *Plutonaster rigidus* Sladen, 1889  
 Stations: 575, 602, 654, 683, 684, 832, 858, 1311, 1331, 1334, 1349  
*Psilaster andromeda andromeda* (J. Müller & Troschel, 1842) = *Psilaster andromeda* J. Müller & Troschel, 1842  
 Stations: 211, 213, 233
- Luidiidae**  
*Luidia sarsi sarsi* Düben & Koren, in Düben, 1845 = *Luidia sarsi* Düben & Koren, 1845  
 Station: 899
- Ctenodiscidae**  
*Ctenodiscus crispatus* (Retzius, 1805) = *Asterias polaris* Sabine, 1824  
 Station: 104
- Porcellanasteridae**  
*Styrcaster armatus* Sladen, 1883 = *Styrcaster spinosus* Perrier, 1885  
 Stations: 527, 745
- Styrcaster elongatus* Koehler, 1907  
 Station: 527
- Notomyotida**  
**BENTHOPECTINIDAE**  
*Cheiraster sepitus* (Verrill, 1885) = *Pontaster venustus* Sladen, 1889  
 Stations: 112, 203, 213, 233, 244, 575, 663, 683, 684, 703, 719, 724, 743, 833, 858, 874, 1311, 1331, 1348
- VALVATIDA**  
**GONIASTERIDAE**  
*Ceramaster granularis granularis* (Müller, 1776) = *Pentagonaster granularis* Müller, 1776  
 Station: 213
- Ceramaster grenadensis grenadiensis* (Perrier, 1881) = *Pentagonaster gosselini* Perrier, 1884  
 Stations: 203, 213, 244, 575, 684, 703, 1344
- Paragonaster subtilis* (Perrier, 1881)  
 Stations: 527, 652
- Plinthaster dentatus* (Perrier, 1884) = *Pentagonaster perrieri* Sladen, 1889  
 Stations: 663, 683, 684, 703, 1311, 1344
- Pseudarchaster gracilis gracilis* (Sladen, 1889) = *Astrogonium necator* Perrier, 1894  
 Station: 1334  
 = *Astrogonium marginatum* Koehler, 1909  
 Station: 1331  
 = *Astrogonium aequabile* Koehler, 1907  
 Station: 1334  
 = *Astrogonium eminens* Koehler, 1907  
 Stations: 698, 863, 1344
- Pseudarchaster parelii* (Düben & Koren, 1846) = *Astrogonium fallax* Perrier, 1885  
 Stations: 553, 575  
 = *Astrogonium annectens* Perrier, 1894  
 Stations: 213, 683, 743, 1331, 1334
- OPHIDIASTERIDAE**  
*Hazelia attenuata* Gray, 1840 = *Hacelia attenuata* Gray, 1840  
 Station: 882
- Ophidiaster ophidianus* (Lamarck, 1816)  
 Station: 550
- Spinulosida**  
**ECHINASTERIDAE**  
*Henricia cylindrella* (Sladen, 1883) = *Cribrella abyssalis* Perrier, 1894  
 Stations: 683, 858, 1331
- Henricia oculata* (Pennant, 1777) = *Cribrella oculata* (Linck) Forbes, 1841  
 Stations: 203, 213, 244
- Velatida**  
**PTERASTERIDAE**  
*Hymenaster giboryi* Perrier, 1894  
 Stations: 527, 652, 738
- Hymenaster pellucidus* Thompson, 1873  
 Station: 248
- Hymenaster roseus* Koehler, 1907  
 Stations: 624, 698, 1334

*Pteraster personatus* Sladen, 1891 = *Pteraster reductus* Koehler, 1907  
Stations: 184, 248, 698, 738, 863

#### ***Brisingida***

##### **BRISINGIDAE**

*Brisingella coronata* (Sars, 1871) = *Brisinga coronata* Sars, 1871  
Stations: 248, 575, 578, 624, 673, 719, 858, 861

##### **FREYELLIDAE**

*Freyastera sexradiata* (Perrier, 1885) =  
*Freyella sexradiata* Perrier, 1885  
Station: 527

#### ***Forcipulatida***

##### **ASTERIIDAE**

*Marthasterias glacialis* (Linnaeus, 1758) =  
*Stolasterias madeirensis* (Stimpson, 1862)  
Station: 216  
= *Asterias glacialis* Linnaeus, 1758  
Station: 550

##### **PEDICELLASTERIDAE**

*Hydrasterias sexradiata* (Perrier, in Milne-Edwards, 1882) = *Pedicellaster sexradiatus* Perrier in Milne Edwards, 1882  
Stations: 248, 536, 624, 663, 673, 698, 724, 738, 745, 847, 861, 863, 866, 1318, 1331, 1334, 1407

##### **NEOMORPHASTERIDAE**

*Neomorphaster margaritaceus* (Perrier, 1882) =  
*Calycaster monoecus* Perrier, 1891  
Station: 203  
= *Neomorphaster talismani* (Perrier, 1885)  
Stations: 213, 244, 575, 624, 663, 683, 684, 698, 702, 719, 724, 738, 858, 1331, 1334

#### ***Ophiuroidea***

##### ***Phrynomphiurida***

##### **OPHIOMYXINA**

##### **OPHIOMYXIDAE**

*Astrogeron supinus* (Lyman, 1883)  
Station: 1349

*Ophiomyxia serpentaria* Lyman, 1883 =  
*Ophiodera serpentina* Lyman, 1883  
Stations: 866, 1344

##### **EURYALINA**

##### **ASTEROSCHEMATIDAE**

*Asteroschema inornatum* Koehler, 1906 =  
*Astrochisma inornatum* Koehler, 1906  
Station: 2248

#### ***Ophiurida***

##### ***GNATHOPHURINA***

##### **OPHIOTRICHIDAE**

*Ophiothrix fragilis* (Abildgaard, in Müller 1789)  
Stations: 226, 882

*Ophiothrix luetkeni* Wyville Thomson, 1873 =  
*Ophiothrix lutkeni* Wyville Thomson, 1873  
Station: 594

##### **OPHIACTIDAE**

*Histampica duplicata* (Lyman, 1875) =  
*Ophiacis duplicata* (Lyman, 1875)  
Stations: 536, 684, 719, 832, 833, 1334, 1344, 1349  
= *Amphiura duplicata* Lyman, 1875  
Station: 184, 233

*Ophiacis abyssicola* (Sars, 1861)

Stations: 233, 602  
= *Ophiacis corallicola* Koehler, 1896  
Stations: 112, 203, 213, 233, 242, 552, 553, 575, 584, 602, 616, 698, 719, 832, 837, 838, 861, 1311, 1344, 1349, 1412, 1420, 2183  
= *Ophiacis echinata* Koehler, 1898  
Station: 198

*Ophiacis profundi* Lütken & Mortensen, 1829  
Station: 2214

##### **AMPHIURIDAE**

*Amphipholis squamata* (Delle Chiaje, 1829)  
= *Amphiura squamata* Delle Chiaje, 1829  
Station: 226  
= *Amphiura tenuispina* Ljungman, 1872  
Stations: 198, 2210

*Amphiura grandisquama* Lyman, 1869

Stations: 838, 866  
= *Amphiura longispina* Koehler, 1???

Station: 242

- Amphiura otteri* Ljungman, 1872 = *Amphiura grandis* Koehler, 1896  
 Station: 1334
- Amphiura richardi* Koehler, 1906  
 Stations: 578, 602, 743
- LAEMOPHIURINA*  
*OPHIACANTHIDAE*  
*Ophiacantha abyssicola* Sars, 1871  
 Station: 837
- Ophiacantha aristata* Koehler, 1896  
 Stations: 233, 874, 1344, 1349
- Ophiacantha bidentata* (Retzius, 1805)  
 Station: 112
- Ophiacantha composita* Koehler, 1907  
 Stations: 738, 1318
- Ophiacantha crassidens* Verrill, 1885  
 Stations: 703, 1344
- Ophiacantha cuspidata* Lyman, 1882  
 Station: 1420
- Ophiacantha enopla veterna* (Koehler, 1907)  
 = *Ophiacantha veterna* Koehler, 1907  
 Stations: 1412, 1420
- Ophiacantha lineata* Koehler, 1909  
 Station: 578
- Ophiacantha valenciennesi* Lyman, 1879  
 Station: 584
- Ophiochondrus armatus* (Koehler, 1909) =  
*Ophioplus armatus* Koehler, 1909  
 Station: 1349
- Ophiomitrella clavigera* (Ljungman, 1865) =  
*Ophiomitrella cordifera* Koehler, 1909  
 Station: 618
- Ophiotoma barletti* (Lyman, 1883) =  
*Ophiopora bartletti* Lyman, 1883  
 Station: 745
- Ophiotrema alberti* Koehler, 1896  
 Station: 527
- CHILOPHIURINA*  
*OPHIURIDAE*  
*Amphiophiura bullata convexa* (Lyman, 1878)  
 = *Ophioglypha convexa* Lyman, 1878  
 Station: 652
- Aspidophiura minuta* (Lyman, 1878) =  
*Ophioglypha minuta* Lyman, 1878  
 Station: 527
- Homophiura tessellata* (Verrill, 1894) =  
*Ophioglypha tessellata* Verrill, 1894  
 Stations: 248, 738
- Ophiocten hastatum* Lyman, 1878  
 Stations: 184, 738, 863, 1334
- Ophiothycis mirabilis* Koehler, 1901  
 Station: 578
- Ophiosphalma planum* (Lyman, 1878) =  
*Ophiomusium planum* Lyman, 1878  
 Stations: 527, 652, 749
- Ophiora carnea* (Lutken, 1858) =  
*Ophioglypha carnea* Lutken, 1858  
 Station: 866
- Ophiura ljunghmani* (Lyman, 1878) =  
*Ophioglypha thouleti* Koehler, 1896  
 Stations: 663, 719, 723, 858  
 = *Ophioglypha ljunghmanni* Lyman, 1878  
 Station: 1334
- Ophiura mundata* Koehler, 1906 =  
*Ophioglypha mundata* Koehler, 1906  
 Station: 738
- Ophiomusium lymani* Wyville Thomson, 1873  
 Stations: 184, 213, 623, 624, 663, 673, 698,  
 719, 724, 738, 847, 858, 1331, 1334
- OPHIOLEUCIDAE*  
*Ophiernus vallincola* Lyman, 1878 =  
*Ophiernus abyssalis* Koehler, 1896  
 Stations: 663, 738, 1334

OPHIONEREIDIDAE  
*Ophionereis reticulata* (Say, 1825)  
Station: 882

OPHIODERMATIDAE  
*Ophioconis forbesi* (Heller, 1863)  
Stations: 226, 698, 882

**Echinoidea**  
Perischoechinoidea

**Cidaroida**

CIDARIDAE

*Cidaris cidaris* (Linnaeus, 1758) =  
*Dorocidaris papillata* (Leske, 1778)  
Stations: 112, 198, 227, 233, 234, 242, 553,  
578, 584, 587, 602, 616, 618, 838, 866,  
1311, 1344, 1367, 2210, 2214

*Cidaris tribuloides* Lamarck, 1816  
Station: 226

Euechinoidea

**DIADEMATOIDEA**

**Echinothuroidea**

ECHINOTHURIDAE

*Calveriosoma hystrix* (Wyville Thomson,  
1872) = *Asthenosoma hystrix* (Wyville  
Thomson, 1872)  
Stations: 112, 198, 242, 244  
= *Araeosoma hystrix* (Wyville Thomson,  
1872)  
Stations: 1311, 1338, 1344, 1349, 1384

*Hygrosoma petersii* (Agassiz, 1880)  
Stations: 575, 578, 1334  
= *Phormosoma uranusp* Wyville Thomson,  
1898  
Stations: 184, 203, 213, 233, 248

*Sperosoma grimaldii* Koehler, 1897  
Stations: 184, 224, 698, 863, 1331

**Pedinoida**

PEDINIDAE

*Caenopedina cubensis* Agassiz, 1869 =  
*Hemipedina cubensis* Agassiz, 1869  
Station: 1311

**ECHINOIDEA**

**Salenoida**

SALENIIDAE

*Salenocidaris profundi profundi* (Duncan,  
1877) = *Salenia hastigera* Agassiz, 1869  
Stations: 105, 203, 233, 244, 575, 578, 584,  
587, 602, 616, 624, 654, 663, 683, 698, 702,  
703, 738, 743, 833, 838, 858, 863, 874,  
1334, 1344, 1349

**Arbacioida**

ARBACIIDAE

*Arbacia lixula* (Linnaeus, 1758) = *Arbacia*  
*pustulosa* Leske, 1778  
Stations: 104, 195, 202

**Temnopleuroidea**

TEMNOPLEURIDAE

*Genocidaris maculata* Agassiz, 1869  
Station: 882  
= *Temnechinus maculatus* Agassiz, 1869  
Station: 226

*Trigonocidaris albida* Agassiz, 1869  
Stations: 234, 570, 597, 600

TOXOPNEUSTIDAE

*Sphaerechinus granularis* (Lamarck, 1816)  
Station: 236

**Echinoida**

ECHINIDAE

*Echinus acutus* Lamarck, 1816  
Stations: 184, 211

*Echinus affinis* Mortensen, 1903

Stations: 536, 738, 853, 858, 861, 863, 1331,  
1334

*Echinus alexandri* Danielssen & Koren, 1883  
Stations: 575, 743, 858, 863, 1331

*Echinus melo* Lamarck, 1816

Station: 899

*Paracentrotus lividus* (Lamarck, 1816) =  
*Strongylocentrotus lividus* (Lamarck, 1816)  
Station: 236

***GNATHOSTOMATA******Clypeasteroida***

## FIBULARIIDAE

*Echinocyamus grandiporus* Mortensen, 1907  
Stations: 553, 584, 587, 597, 614, 616, 618,  
866, 1349, 2210, 2214

*Echinocyamus macrostomus* Mortensen, 1907  
Stations: 536, 581, 719

*Echinocyamus pusillus* (Müller, 1776)  
Stations: 198, 233, 234, 1349, 2214

***ATELESTOMATA******Spatangoida***

## BRISSIDAE

*Brissopsis lyrifera* (Forbes, 1841)  
Station: 226

## AEROPSIDAE

*Acaste bellidifera* Wyville Thomson, 1877  
Station: 703

## PALAEOPNEUSTIDAE

*Palaeotropus josephinae* Lovén, 1898 =  
*Palaeotropus hirondellei* Koehler, 1???  
Stations: 105, 553, 575, 578, 584, 602, 616,  
703, 719, 743, 833, 838, 1344, 1349

*Peripatagus cinctus* Koehler, 1895  
Stations: 616, 743, 838

***Holothuroidea******Dendrochirotida***

## CUCUMARIIDAE

*Abyssocucumis abyssorum* (Théel, 1886) =  
*Cucumaria abyssorum* Théel, 1886  
Station: 248

*Pawsonia saxicola* (Brady & Robertson,  
1871) = *Cucumaria montagui* Fleming,  
1828  
Station: 226

*Thyone inermis* Heller, 1868  
Stations: 226, 553, 575

***Aspidochirotida***

## SYNALLACTIDAE

*Mesothuria lactea* (Théel, 1886)  
Stations: 575, 624, 663, 698, 703, 719, 863,  
1334, 3150

*Mesothuria murrayi* Théel, 1886

Note: The species *M. murrayi* was described by Théel, 1886; *M. murrayi* var. *grandipes* is referred to Hérouard (see also HANSSON 2001)

Stations: 536, 719, 3150

*Mesothuria verrilli* (Théel, 1886)

Stations: 1311, 1318, 1334, 3150, 3293  
= *Allantis intestinalis verrilli* Théel, 1886  
Stations: 553, 575, 683, 684, 743, 858, 874  
= *Holothuria verrilli* Théel, 1886  
Stations: 233, 244

*Paelopatides atlantica* Hérouard, 1902 =  
*Poelopatides atlantica* Hérouard, 1902

Station: 527

*Pseudostichopus lapidus* Hérouard, 1902

Station: 527

*Pseudostichopus occultatus* Marenzeller, 1893

Station: 527

*Pseudostichopus villosus* Théel, 1886

Station: 652

## HOLOTHURIIDAE

*Holothuria lentiginosa* Marenzeller, 1892  
Station: 226

*Holothuria mexicana* Ludwig, 1875

Station: 882

***Elasipodida***

## ELPIDIIDAE

*Amperima furcata* (Hérouard, 1899) = *Kolga furcata* Hérouard, 1899  
Station: 698

*Ellipinion delagei* (Hérouard, 1896)

Station: 3293

= *Scotoplanes delagei* Hérouard

Station: 553, 578, 743

*Peniagone azorica* Marenzeller, 1893

Stations: 248, 527

## LAETMOGONIDAE

*Benthogone rosea* Koehler, 1896  
Station: 1334

*Laetmogone violacea* Théel, 1879 =  
*Laetmogone wyville-thomsoni* Théel  
Station: 683

*Pannychia glutinosa* Hérourard, 1902  
Station: 624

#### PSYCHROPOTIDAE

*Benthodytes janthina* Marenzeller, 1893  
Stations: 248, 673

*Benthodytes typica* Théel, 1882  
Station: 248

*Psychropotes depressa* (Théel, 1882) =  
*Euphronides talismani* Perrier, 1896  
Station: 673

*Psychropotes grimaldii* Hérourard, 1896  
Station: 527

#### *Apodida*

##### SYNAPTIDAE

*Synaptula hydriformis* (Lesueur, 1824)  
Station: 1349

##### CHIRIDOTIDAE

*Chiridota abyssicola* Marenzeller, 1893  
Station: 248

#### Uncertain species (echinoderms)

*Pannychia* sp. Théel, 1882

Note: according to Hansson (1998; European Echinodermata Checklist) the record is a synonym of *Laetmophasma* sp.  
Station: 624

*Prognaster grimaldii* Perrier, 1896

Note: It was impossible to determine the validity of this species name, even considering its description by Perrier (1896). The author included it in the family Zoroasteridae.  
Station: 248

*Ophiacantha pentagona* var *armata* Koehler

Note: Possibly a synonym of *Ophiochondrus armatus* (Koehler, 1909) (Hansson, 2001). Both genera *Ophiacantha* and *Ophiochondrus* belong to the family Ophiacanthidae.  
Station: 248

#### *Ophiactis hirta* Lyman

Note: the validity of this nominal species could not be verified. The genus exists (family Ophiactidae) but the species could not be found.  
Station: 1344

#### *Zoroaster trispinosus* Koehler

Note: Included in the family Zoroasteridae. Might be *Zoroaster fulgens* Thompson, 1873 (Hansson, 1998, 2001).  
Station: 745

#### *Stellosphaera mirabilis* Koehler & Vaney

Note: As Koehler (1909) refers this is a larva of asteroid, probably belonging to an abyssal form. The author included the larvae in Forcipulida.

Stations: 1874, 2159, 2168, 2194, 2242, 2264

## CONCLUSIONS

The main objective in compiling this checklist was to find the currently valid names of the species reported by Prince Albert's campaigns, as the systematics of most groups evolved since his pioneering programme.

The 331 nominal species associated to the seven phyla selected represent about 13% of the total nominal species (2624) assembled in the PAM Azores database. Since 15 species have 2 or more synonyms (all echinoderms and 1 cephalopod) a list of 310 species names was considered to relate to valid species. About 50% of those were validated directly (e.g. without any modification in their nomenclature), but the systematics and/or nomenclature of the remaining species names were modified. Those major changes are summarised in the table 1. Annelids, chaetognaths and brachiopods were the phyla that showed fewer modifications.

Since it lacks a critical observation of the original specimens sampled, the method adopted to review and validate the original species names is obviously prone to potential flaws which cannot be easily assessed with the available knowledge. Working directly with the specimens would potentially provide additional information and accuracy about the actual species systematics. However, and at least for several groups of invertebrates, this is impracticable because the

original collections are in very poor condition and will require systematics expertise in many disparate groups and consequently the involvement of a vast group of taxonomists. In the absence of the sustained and co-ordinated

effort required to evaluate the original collections, the present review (which took approximately 500 hours work), based on the published bibliography and electronic documentation, is justified.

Table 1

Systematics modifications made in the original lists of species reported in the results of the Prince Albert I of Monaco for the Azores waters (by phylum). Nom. sp: species included in the original lists; Val. sp: species considered valid, included those synonymised; Gen %: species that were allocated to different genus; Syn %: nominal species relegated to synonyms; Missp %: misspellings and corrections due to gender or concordance rules; Sub-sp: specific epithets allocated to sub-specific level or vice-versa.

Phylum	Nom. sp	Val. sp	Gen %	Syn %	Missp %	Sub-sp %
<b>Echiura</b>	1	1	0	0	0	0
<b>Sipuncula</b>	11	11	54,5	27,3	0	0
<b>Mollusca (Cephalopoda)</b>	37	32	45,9	64,8	0	0
<b>Annelida</b>	130	130	20,0	7,7	3,1	0
<b>Brachiopoda</b>	4	4	50,0	0	0	0
<b>Chaetognatha</b>	14	14	7,1	0	14,3	0
<b>Echinodermata</b>	134	118	25,4	31,4	2,5	5,9
<b>Total</b>	331	310	28,9	18,7	2,8	0,8

The historical aspect of this biodiversity assessment is also particularly relevant in the view of the study of global changes in biodiversity patterns and species distributions. These data will provide a baseline for future inventories of the Azorean fauna. Data processing is an important scientific activity because it gives the baseline for future research. This contribution provides historical data, which were not readily available, in a useful format that can be used in the present by the scientific community in general. Moreover, the data may contribute to some international initiatives which aims to improve their databases to include e.g. geographical information (e.g. ERMS, COSTELLO et al. 2001; NEAT, HANSSON 1997, 1998).

At the regional level, the knowledge obtained by the critical analysis of the PAM results is crucial for an appropriate assessment of the biodiversity of the Azores region. The inventories of biodiversity for the region are limited and the several faunistic lists published do not include systematic reviews of historical data. As the data presented here are geo-referenced, distribution maps can be generated to support efforts of biodiversity conservation.

The availability of an on-line database assembling sampling locations and biological

data produced by the immense work done under the auspicious of the Prince Albert I of Monaco is expected to contribute to improve the dissemination of the knowledge about the marine fauna of the Northeast Atlantic.

#### ACKNOWLEDGEMENTS

The European programme Leonardo da Vinci supported this work in the Department of Oceanography and Fisheries of the University of the Azores (DOP/UAç). Thanks are due to the head of DOP/UAç Ricardo Serrão Santos for his support and all the people who did the previous work to create the PAM Azores database: Ricardo Medeiros, Fátima Mendes and Rogério Ferraz. Edward Cutler is acknowledged for his help in solving some issues related to the systematics of sipunculids.

#### REFERENCES

- ALVAREZ F. Brachiopoda of the Iberian Peninsula and Balearic Islands. Internet resources. Available from <http://www.fauna-iberica.mncn.csic.es/htmlfauna/faunibe/zoolist/brachiopoda.html>. Last accessed March 2004.

- AMEZIANE, N. 1999. Muséum National d'Histoire Naturelle. Echinoderms. Internet resources. Available from <http://www.mnhn.fr/mnhn/bimm/protection/fr/Espesies/Fiches/Paracentrotuslividus.htm>. Last accessed February 2004.
- ÁVILA, S.P. 2000. Shallow-water marine mollusks if the Azores: biogeographical relationships. *Arquipélago. Life and Marine Sciences. Supplement 2A*: 99-131.
- BELLAN, G. 1978. Une petite collection d'annelides polychetes recoltées dans l'île de São Miguel (Archipel des Açores). *Boletim da Sociedade Portuguesa de Ciências Naturais*. 18: 57-67.
- BELLAN, G. 2001. Annelida. Pp. 214-232 in: COSTELLO, M.J., C.S. EMBLOW & R. WHITE (Eds). *European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification*. Patrimoines naturels, 50: 463 pp.
- CHAPMAN, G. & R. DALES 1954. Aspects of the fauna and flora of the Azores. II. Polychaeta. *Annales and Magazine Natural History Series* 12 (7) 678-683.
- CHAPMAN, G. 1955. Aspects of the fauna and flora of the Azores. III. Gephyrea. *Annales and Magazine Natural History Series* 12 (8) 351-352.
- CLARKE, M.R. 1996. A review of the systematics and Ecology of Oceanic Squids. *Advances in Marine Biology* 4: 91-300.
- COSTELLO, M.J., C.S. EMBLOW & R. WHITE (Eds) 2001. European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification. *Patrimoines naturels* 50: 463 pp.
- CUTLER, E.B. 1994. *The Sipuncula, their systematics, biology, and evolution*. Cornell University Press. 480 pp.
- FAUVEL, P. 1914. Annélides Polychètes non pélagiques provenant des campagnes de l' Hirondelle et de la Princesse-Alice (1885-1910). *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 46, 432 pp.
- FAUVEL, P. 1916. Annélides Polychètes pélagiques provenant des campagnes des yachts Hirondelle et Princesse-Alice (1885-1910). *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 48, 152 pp.
- FAUVEL, P. 1932. Annélides Polychètes provenant des campagnes de l' Hirondelle II (1911-1915). *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 85, 50 pp.
- FISCHER, P. & D.-P. OEHLENT 1892. Brachiopodes de l'Atlantique Nord. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 3, 30 pp.
- GERMAIN, L. & L. JOUBIN 1916. Chétognathes provenant des campagnes des yachts Hirondelle et Princesse-Alice. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 49, 118 pp.
- GONÇALVES, J.M. 1991. The Octopoda (Mollusca: Cephalopoda) of the Azores. *Arquipélago. Ciências da Natureza* 9:75-81.
- HANSSON, H.G. (Comp.), NEAT (North East Atlantic Taxa): South Scandinavian marine Annelida Check-List. Internet resources pdf Ed., Aug. 1998. [<http://www.tmbi.gu.se>]. Last accessed April 2004.
- HANSSON, H.G. (Comp.), NEAT (North East Atlantic Taxa): South Scandinavian marine Echinodermata Check-List. Internet pdf Ed., Aug. 1998. [<http://www.tmbi.gu.se>]. Last accessed May 2004.
- HANSSON, H.G. (Comp.) 1997. NEAT (North East Atlantic Taxa): South Scandinavian marine "Aschelminth" (excl. Nematoda) & other species poor phyla Check-List. Internet pdf Ed., May. 1997. [<http://www.tmbi.gu.se>]. Last accessed April 2004.
- HANSSON, H.G. 2001. Echinodermata. Pp. 336-351 in: COSTELLO, M.J., C.S. EMBLOW & R. WHITE (Eds). *European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification*. Patrimoines naturels, 50: 463 pp.
- HÈROUARD, E. 1902. Holothuries provenant des campagnes de la Princesse-Alice (1892-1897). *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 21, 61 pp.
- HOWSON, C.M. 2001. Brachiopoda. Pp. xx-xx in: COSTELLO, M.J., C.S. EMBLOW & R. WHITE (Eds). *European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification*. Patrimoines naturels, 50: 463 pp.
- JOUBIN, L. 1895. Contribution à l'étude des céphalopodes de l'Atlantique Nord. *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ière Prince Souverain de Monaco*. IX:63 pp.
- JOUBIN, L. 1900. Céphalopodes provenant des campagnes de la Princesse-Alice (1891-1897). *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ière Prince Souverain de Monaco*. XVII:135 pp.
- JOUBIN, L. 1920. Céphalopods provenant des campagnes de la Princesse-Alice (1898-1910) (3ème Série). *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ière Prince Souverain de Monaco*. LIV:135 pp.
- JOUBIN, L. 1924. Contribution à l'étude des céphalopodes de l'Atlantique Nord (4ème Série). *Résultats des Campagnes Scientifiques Accomplies sur son Yacht par Albert Ière Prince Souverain de Monaco*. LXVII:113 pp.

- KATO, T., & F. PLEIJEL 2002. A revision of *Notophyllum* Orsted, 1834 (Phyllodocidae, Polychaeta). *Journal of Natural History* 36 (10): 1135-1178.
- KOEHLER, R. 1898. Echinides et Ophiures provenant des campagnes du yacht l'Hirondelle. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 12: 78 pp.
- KOEHLER, R. 1909. Échinoderms provenant des campagnes du yacht Princesse-Alice. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 34: 317pp.
- LOGAN, A. 1998. Recent Brachiopoda from the oceanographic expedition SEAMOUNT 2 to the north-eastern Atlantic in 1993. *Zoosystema* 20(4): 549-562.
- MÜLLER, S. 1999. Internet resources. Available from Web - Echinodermata Sammlung Marburg. Last accessed February 2004.
- PEREIRA, R.M.O. 1997. Checklist of the littoral Echinoderms of the Azores. *Açoreana* 8(3): 331-337.
- PERRIER, E. 1896. Contribution à l'étude des Stellérides de l'Atlantique Nord. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 11: 57 pp.
- RICHARD, J. 1934. Liste général des stations des campagnes scientifique du Prince Albert of Monaco. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 89: 472 pp + viii plates.
- SANTOS, R.S., F.M. PORTEIRO & J.P. BARREIROS 1997. Marine fishes of the Azores: An annotated checklist and bibliography. *Arquipélago*. Life and Marine Sciences. Supplement 1: xxii + 231pp.
- SALDANHA, L. 1995. *Fauna Submarina Atlântica*. Publicações Europa-América. 364 pp.
- SLUITER, G.-P. 1900. Géphyriens (Sipunculides et Echiurides) provenant des campagnes de l'Hirondelle et de la Princesse-Alice. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 15, 29 pp.
- SLUITER, G.-P. 1912. Géphyriens (Sipunculides et Echiurides) provenant des campagnes de la Princesse-Alice. *Résultats des Campagnes Scientifiques du Monaco*, Fascicule 36, 36 pp.
- SMITH, A.B. (Ed.) 2003. *The Echinoid Directory*. Internet resources. Available from World Wide Web electronic publication <http://www.nhm.ac.uk/palaeontology/echinoids>. Last accessed February 2004.
- SOUTHWARD, A.J. 1998. New observations on barnacles (Crustacea: Cirripedia) of the Azores region. *Arquipélago*. Life and Marine Sciences 16A: 11-27
- VAN DER LAND, J. & H. KAPP. 2001. Chaetognatha. P. 335 in: COSTELLO, M.J., C.S. EMBLOW & R. WHITE (Eds). *European Register of Marine Species. A check-list of the marine species in Europe and a bibliography of guides to their identification*. Patrimoines Naturels, 50: 463 pp.
- YOUNG, P.S. 1998. Cirripedia (Crustacea) from the "Campagne Biaçores" in the Azores region, including a generic revision of Verrucidae. *Zoosystema* 20(1): 31-92.
- YOUNG, P.S. 2001. Deep-sea Cirripedia Thoracica (Crustacea) from the northeast Atlantic collected by French expeditions. *Zoosystema* 23(4): 705-756.
- WIRTZ, P. 1995. *Unterwasserführer Madeira, Kanaren Azoren*. Ed. Naglschmid. 247 pp.
- WIRTZ, P. & H. DEBELIUS 2003. *Mediterranean and Atlantic Invertebrate Guide*. ConchBooks. 305 pp.

Accepted 12 September 2005

## APPENDIX 1

Sampling location data from the campaigns of Prince Albert I of Monaco (PAM) in the Azores waters (1885-1914). Only sampling locations with positive catches of species of the phyla Echiura, Sipuncula, Mollusca, Annelida, Brachiopoda, Chaetognatha and Echinodermata are presented. Data from Richard (1934). The sampling location numbers follows the PAM registration; \* denotes locations which latitude and longitude were estimated (see material and methods).

PAM Stat	Lat N	Long W	Depth	Date	Vessel	Gear
15	38° 15' 20"	28° 22' 45"	Surface	02-08-1885	Hirondelle	Plankton net
16	38° 6' 0"	26° 57' 45"	Surface	03-08-1885	Hirondelle	Plankton net
17	37° 31' 0"	26° 44' 45"	Surface	05-08-1885	Hirondelle	Plankton net
18	37° 40' 0"	25° 29' 45"	Surface	20-08-1885	Hirondelle	Plankton net
21	42° 55' 0"	22° 34' 45"	Surface	23-08-1885	Hirondelle	Plankton net
103	38° 32' 5"	28° 37' 30"	15	21-06-1887	Hirondelle	Trap

104*	38° 32' 10"	28° 36' 52"	Intertidal	22-06-1887	Hirondelle	
105	38° 23' 45"	28° 31' 15"	927	25-06-1887	Hirondelle	Bottom trawl
112	38° 34' 30"	28° 06' 15"	1287	01-07-1887	Hirondelle	Bottom trawl
115	38° 38' 45"	28° 23' 0"	0-1	04-07-1887	Hirondelle	Plankton high speed net
116	38° 35' 55"	28° 10' 35"	0-1	04-07-1887	Hirondelle	Plankton high speed net
125	38° 23' 0"	27° 06' 15"		13-07-1887	Hirondelle	
182	41° 48' 22"	22° 28' 45"	Surface	11-07-1888	Hirondelle	Floating wreck
184	40° 5' 0"	27° 27' 45"	1850	14-07-1888	Hirondelle	Bottom trawl
185	39° 27' 0"	27° 55' 45"	Surface	15-07-1888	Hirondelle	Plankton net
190	38° 46' 30"	28° 20' 43"	696	19-07-1888	Hirondelle	Bottom trawl
191*	38° 32' 10"	28° 36' 52"		20-07-1888	Hirondelle	
193*	38° 32' 10"	28° 36' 52"	20	22-07-1888	Hirondelle	Dredge
195*	38° 31' 28"	28° 37' 43"	4	23-07-1888	Hirondelle	Dredge
196*	38° 32' 10"	28° 36' 52"	5-6	23-07-1888	Hirondelle	Dredge
198	38° 26' 25"	28° 38' 55"	800	25-07-1888	Hirondelle	Bottom trawl
202*	39° 28' 10"	31° 7' 44"	Intertidal	30-07-1888	Hirondelle	
203	39° 27' 5"	30° 55' 5"	1557	30-07-1888	Hirondelle	Bottom trawl
208	39° 22' 15"	31° 12' 0"	Surface	31-07-1888	Hirondelle	Plankton net
211	39° 18' 5"	31° 12' 0"	1372	01-08-1888	Hirondelle	Bottom trawl
213	39° 22' 48"	31° 25' 15"	1384	02-08-1888	Hirondelle	Bottom trawl
215	39° 34' 0"	31° 14' 15"	Surface	03-08-1888	Hirondelle	Midwater trawl
216	39° 26' 30"	31° 9' 0"	Intertidal	03-08-1888	Hirondelle	
218*	39° 28' 10"	31° 7' 44"	40	04-08-1888	Hirondelle	Dredge
220	39° 42' 59"	31° 1' 24"	1445	05-08-1888	Hirondelle	Bottom trawl
224	39° 44' 20"	34° 5' 7"	1213	07-08-1888	Hirondelle	Bottom trawl
225	38° 31' 0"	28° 35' 35"	129	13-08-1888	Hirondelle	Trap
226	38° 31' 19"	28° 34' 30"	130	14-08-1888	Hirondelle	Bottom trawl
227	38° 23' 0"	28° 26' 37"	1135	15-08-1888	Hirondelle	Bottom trawl
229	38° 22' 0"	28° 14' 24"	736	16-08-1888	Hirondelle	Bottom trawl
232	38° 33' 21"	28° 8' 39"	1300	17-08-1888	Hirondelle	Trap
233	38° 33' 21"	28° 8' 39"	1300	18-08-1888	Hirondelle	Bottom trawl
234	39° 1' 40"	27° 55' 25"	454	19-08-1888	Hirondelle	Bottom trawl
236	38° 3' 25"	27° 57' 45"	Intertidal	20-08-1888	Hirondelle	
238	39° 3' 0"	27° 56' 45"	95	21-08-1888	Hirondelle	Bottom long-line
242	38° 48' 30"	27° 58' 45"	861	22-08-1888	Hirondelle	Bottom trawl
243	38° 31' 55"	28° 35' 45"	120	25-08-1888	Hirondelle	Trap
244	38° 33' 57"	28° 19' 15"	1266	27-08-1888	Hirondelle	Bottom trawl
246	38° 27' 45"	28° 8' 0"	Surface	29-08-1888	Hirondelle	Midwater trawl
248	41° 40' 41"	26° 44' 8"	2870	02-09-1888	Hirondelle	Bottom trawl
527	38° 8' 0"	23° 18' 45"	4020	25-06-1895	Princesse-Alice	Bottom trawl
536	37° 54' 0"	24° 43' 15"	2178	27-06-1895	Princesse-Alice	Bottom trawl
545	37° 16' 0"	24° 44' 45"	Surface	01-07-1895	Princesse-Alice	By hand
550*	37° 16' 16"	24° 47' 17"	Intertidal	02-07-1895	Princesse-Alice	By hand
552	37° 42' 40"	25° 5' 15"	1385	03-07-1895	Princesse-Alice	Sounder
553	37° 42' 40"	25° 5' 15"	1385	03-07-1895	Princesse-Alice	Bottom trawl
554	37° 43' 00"	25° 5' 45"	1385	03-07-1895	Princesse-Alice	Trap
570	37° 54' 0"	25° 38' 15"	550	12-07-1895	Princesse-Alice	Swab bar
575	38° 27' 0"	26° 30' 15"	1165	13-07-1895	Princesse-Alice	Bottom trawl
576	38° 26' 0"	26° 32' 45"	Surface	13-07-1895	Princesse-Alice	Dip net
578	38° 26' 0"	26° 30' 45"	1165	14-07-1895	Princesse-Alice	Bottom trawl
581	38° 26' 0"	26° 38' 15"	2139	15-07-1895	Princesse-Alice	Bottom trawl
582	38° 31' 0"	26° 49' 15"	845	15-07-1895	Princesse-Alice	Buchanan bottle
584*	38° 31' 0"	26° 49' 15"	845	16-07-1895	Princesse-Alice	Swab bar
587	38° 36' 40"	27° 17' 15"	793	18-07-1895	Princesse-Alice	Swab bar
588	38° 34' 45"	27° 16' 45"	Surface	18-07-1895	Princesse-Alice	Dip net
594*	38° 38' 31"	27° 12' 28"	54	22-07-1895	Princesse-Alice	
597	38° 27' 0"	28° 3' 25"	523	23-07-1895	Princesse-Alice	Swab bar
600	38° 30' 35"	28° 16' 20"	349	24-07-1895	Princesse-Alice	Swab bar
602	38° 38' 30"	28° 13' 5"	1230	24-07-1895	Princesse-Alice	Bottom trawl

603	38° 31' 28"	28° 37' 43"	Intertidal	26-07-1895	Princesse-Alice	
612	38° 26' 40"	28° 40' 5"	778	29-07-1895	Princesse-Alice	Buchanan bottle
614	38° 27' 12"	28° 39' 15"	778	31-07-1895	Princesse-Alice	Swab bar
616	38° 47' 40"	28° 17' 5"	1022	01-08-1895	Princesse-Alice	Swab bar
618	38° 52' 45"	28° 6' 0"	1143	01-08-1895	Princesse-Alice	Bottom trawl
622*	39° 3' 10"	27° 57' 37"		05-08-1895	Princesse-Alice	Gill net
623	38° 59' 0"	28° 18' 5"	2102	04-08-1895	Princesse-Alice	Buchanan bottle
624	38° 59' 0"	28° 18' 5"	2102	04-08-1895	Princesse-Alice	Bottom trawl
652	36° 55' 0"	22° 22' 45"	4261	23-06-1896	Princesse-Alice	Bottom trawl
654	36° 58' 30"	24° 58' 15"	1495	24-06-1896	Princesse-Alice	Swab bar
663	37° 28' 30"	25° 31' 45"	1732	27-06-1896	Princesse-Alice	Bottom trawl
664	37° 37' 40"	25° 37' 45"	Surface	27-06-1896	Princesse-Alice	Dip net
673	37° 51' 0"	26° 53' 45"	2252	05-07-1896	Princesse-Alice	Bottom trawl
682	38° 20' 0"	28° 5' 45"	Surface	07-07-1896	Princesse-Alice	Harpoon
683	38° 20' 0"	28° 4' 45"	1550	07-07-1896	Princesse-Alice	Bottom trawl
684	38° 20' 0"	28° 4' 45"	1550	08-07-1896	Princesse-Alice	Bottom trawl
698	39° 11' 0"	30° 44' 40"	1846	18-07-1896	Princesse-Alice	Bottom trawl
699	39° 12' 0"	30° 48' 45"	Surface	18-07-1896	Princesse-Alice	
702	39° 21' 20"	31° 5' 53"	1360	19-07-1896	Princesse-Alice	Gill net
703	39° 21' 20"	31° 5' 45"	1360	19-07-1896	Princesse-Alice	Bottom trawl
709	39° 13' 40"	30° 43' 45"		22-07-1896	Princesse-Alice	By hand
719	39° 11' 0"	30° 24' 15"	1600	27-07-1896	Princesse-Alice	Bottom trawl
723	37° 17' 40"	28° 15' 10"	1692	31-07-1896	Princesse-Alice	Trap
724	38° 18' 0"	28° 14' 45"	1692	31-07-1896	Princesse-Alice	Gill net
726	38° 18' 0"	28° 14' 45"	Surface	01-08-1896	Princesse-Alice	Harpoon
738	37° 40' 0"	26° 26' 15"	1919	07-08-1896	Princesse-Alice	Bottom trawl
740	37° 39' 0"	26° 26' 45"	1000	07-08-1896	Princesse-Alice	Modified Giesbrecht net
743	37° 35' 45"	25° 17' 15"	1494	11-08-1896	Princesse-Alice	Bottom trawl
745	38° 5' 0"	23° 50' 15"	3465	15-08-1896	Princesse-Alice	Bottom trawl
750	38° 55' 00"	21° 18' 45"	Surface	17-08-1896	Princesse-Alice	Dip net
832	37° 39' 0"	25° 17' 45"	1230	21-07-1897	Princesse-Alice	Gill net
833	37° 39' 0"	25° 17' 45"	1230	21-07-1897	Princesse-Alice	Swab bar
837	37° 55' 0"	25° 24' 15"	880	22-07-1897	Princesse-Alice	Trap
838	37° 55' 0"	25° 23' 45"	880	22-07-1897	Princesse-Alice	Swab bar
842*	37° 53' 35"	25° 29' 41"	Surface	23-07-1897	Princesse-Alice	Dip net
844	37° 55' 0"	25° 24' 15"	Surface	24-07-1897	Princesse-Alice	Dip net
847	38° 1' 30"	25° 25' 45"	1638	24-07-1897	Princesse-Alice	Gill net
848	38° 4' 0"	25° 22' 45"	Surface	25-07-1897	Princesse-Alice	Dip net
849	38° 4' 0"	25° 42' 45"	Surface	25-07-1897	Princesse-Alice	Harpoon
853	38° 15' 0"	26° 44' 15"	1531	29-07-1897	Princesse-Alice	Gill net
858	38° 45' 0"	26° 35' 45"	1482	31-07-1897	Princesse-Alice	Midwater trawl
861	38° 53' 0"	26° 40' 45"	1935	31-07-1897	Princesse-Alice	Midwater trawl
863	39° 22' 0"	26° 55' 45"	1940	01-08-1897	Princesse-Alice	Bottom trawl
866	38° 52' 50"	27° 23' 5"	599	02-08-1897	Princesse-Alice	Bottom trawl
869	39° 3' 0"	27° 42' 45"	1240	03-08-1897	Princesse-Alice	Bottom trawl
873	38° 37' 45"	28° 14' 20"	1260	04-08-1897	Princesse-Alice	Trap
874	38° 37' 45"	28° 14' 20"	1260	04-08-1897	Princesse-Alice	Gill net
882	38° 30' 40"	28° 34' 45"	98	07-08-1897	Princesse-Alice	Bottom trawl
889	37° 57' 30"	29° 15' 10"	208	10-08-1897	Princesse-Alice	3 sinkers de sonda suiffée
896	38° 1' 0"	29° 22' 15"	1260	11-08-1897	Princesse-Alice	Trap
899	37° 57' 0"	29° 14' 45"	200	12-08-1897	Princesse-Alice	Bottom trawl
1311	37° 37' 0"	25° 20' 45"	1187	31-07-1902	Princesse-Alice II	Bottom trawl
1318	38° 6' 0"	26° 13' 45"	3018	05-08-1902	Princesse-Alice II	Bottom trawl
1322	38° 39' 0"	25° 6' 15"	3020	06-08-1902	Princesse-Alice II	Towed double trap
1331	38° 40' 0"	26° 0' 45"	1805	09-08-1902	Princesse-Alice II	Bottom trawl
1333	39° 30' 0"	29° 2' 45"	1900	13-08-1902	Princesse-Alice II	Towed double trap
1334	39° 30' 0"	29° 2' 15"	1900	13-08-1902	Princesse-Alice II	Bottom trawl
1338	38° 41' 30"	28° 45' 15"	950	14-08-1902	Princesse-Alice II	Bottom trawl
1339	38° 41' 30"	28° 45' 15"	Surface	14-08-1902	Princesse-Alice	Dip net

1344	38° 45' 30"	28° 7' 45"	1095	18-08-1902	Princesse-Alice II	Bottom trawl
1348	38° 35' 40"	28° 8' 15"	1250	19-08-1902	Princesse-Alice II	Gill net
1349	38° 35' 30"	28° 5' 45"	1250	19-08-1902	Princesse-Alice II	Bottom trawl
1355	37° 58' 30"	29° 17' 45"	78	23-08-1902	Princesse-Alice II	Hand line
1358	37° 42' 0"	29° 2' 45"	403	24-08-1902	Princesse-Alice II	3 sinkers de sonda suiffée
1367	37° 34' 0"	28° 56' 45"	563	25-08-1902	Princesse-Alice II	Bottom trawl
1373	37° 34' 30"	29° 7' 45"	1685	25-08-1902	Princesse-Alice II	3 sinkers de sonda suiffée
1384	37° 45' 0"	28° 46' 45"	1528	26-08-1902	Princesse-Alice II	Bottom long-line
1407	38° 1' 30"	26° 16' 45"	2755	03-09-1902	Princesse-Alice II	Bottom long-line
1408	38° 01' 30"	26° 16' 45"	?	03-09-1902	Princesse-Alice	From the line de sonda
1412	42° 44' 0"	28° 22' 45"	2200	05-09-1902	Princesse-Alice II	Bottom long-line
1420	42° 53' 0"	28° 30' 45"	2460	06-09-1902	Princesse-Alice II	Bottom trawl
1805	34° 0' 0"	25° 30' 0"	Surface	27-08-1904	Princesse-Alice II	Plankton high-speed net
1834	37° 28' 0"	25° 53' 30"	0-1000	06-09-1904	Princesse-Alice II	Large Richard trawl
1839*	37° 43' 56"	25° 39' 40"	Surface	06-09-1904	Princesse-Alice II	Plankton high speed net
1844	37° 8' 0"	28° 28' 30"	0-1500	07-09-1904	Princesse-Alice II	Large Richard trawl
1847	37° 3' 0"	28° 52' 0"	Surface	07-09-1904	Princesse-Alice II	Plankton high speed net
1849	36° 14' 0"	28° 53' 0"	0-3000	08-09-1904	Princesse-Alice II	Large Richard trawl
1851	36° 17' 0"	28° 53' 0"	0-3000	08-09-1904	Princesse-Alice II	Large Richard trawl
1855	36° 46' 0"	26° 41' 0"	3620	09-09-1904	Princesse-Alice II	3 sinkers de sonda suiffée and bait
1856	36° 46' 0"	26° 41' 0"	0-3250	09-09-1904	Princesse-Alice II	Large Richard trawl
1860	37° 0' 0"	26° 20' 0"	Surface	09-09-1904	Princesse-Alice II	Plankton high speed net
1872	37° 35' 0"	24° 40' 0"	Surface	11-09-1904	Princesse-Alice II	Plankton high speed net
1874	37° 20' 0"	21° 40' 0"	0-2000	12-09-1904	Princesse-Alice II	Large Richard trawl
1875	37° 20' 0"	21° 40' 0"	Surface	12-09-1904	Princesse-Alice II	Dip net
2143	34° 0' 0"	34° 20' 0"	Surface	20-08-1905	Princesse-Alice II	Plankton high speed net
2148	33° 51' 0"	34° 3' 0"	?	20-08-1905	Princesse-Alice	By hand from the cable de sonda
2149	33° 51' 0"	34° 3' 0"	0-2000	20-08-1905	Princesse-Alice II	Large Richard trawl
2150	34° 10' 0"	33° 50' 0"	Surface	20-08-1905	Princesse-Alice II	Plankton high speed net
2151	34° 50' 0"	32° 30' 0"	Surface	21-08-1905	Princesse-Alice II	Plankton high speed net
2153	35° 4' 0"	32° 11' 0"	0-2000	21-08-1905	Princesse-Alice II	Large Richard trawl
2159	36° 24' 0"	30° 0' 0"	0-2500	22-08-1905	Princesse-Alice II	Large Richard trawl
2161	36° 30' 0"	29° 50' 0"	Surface	22-08-1905	Princesse-Alice II	Plankton high speed net
2162	36° 40' 0"	29° 0' 0"	Surface	22-08-1905	Princesse-Alice II	Plankton high speed net
2168	36° 35' 0"	27° 12' 0"	0-2000	23-08-1905	Princesse-Alice II	Large Richard trawl
2170	36° 38' 0"	27° 6' 0"	Surface	23-08-1905	Princesse-Alice II	Plankton high speed net
2171	36° 50' 0"	26° 50' 0"	Surface	23-08-1905	Princesse-Alice II	Dip net
2172	36° 50' 0"	26° 5' 50"	Surface	23-08-1905	Princesse-Alice II	Plankton high speed net
2183	38° 4' 45"	25° 54' 0"	1998	28-08-1905	Princesse-Alice II	Bottom long-line
2184	38° 04' 45"	25° 54' 0"	Surface	28-08-1905	Princesse-Alice	Dip net from the fanal électrique
2185	38° 4' 0"	26° 7' 30"	0-3000	29-08-1905	Princesse-Alice II	Large Richard trawl
2187	38° 4' 0"	26° 7' 30"	0-2500	29-08-1905	Princesse-Alice II	Large Richard trawl
2191	39° 9' 0"	26° 13' 0"	Surface	30-08-1905	Princesse-Alice II	Plankton high speed net
2194	39° 36' 0"	26° 5' 0"	0-2500	30-08-1905	Princesse-Alice II	Large Richard trawl
2195	39° 36' 0"	26° 5' 0"	Surface	30-08-1905	Princesse-Alice II	Dip net
2196	39° 38' 0"	26° 40' 0"	Surface	30-08-1905	Princesse-Alice II	Plankton high speed net
2198	39° 44' 0"	28° 25' 0"	Surface	31-08-1905	Princesse-Alice II	Plankton high speed net
2199	39° 44' 0"	28° 25' 0"	1943	31-08-1905	Princesse-Alice II	3 sinkers de sonda suiffée and bait
2200	39° 44' 0"	28° 25' 0"	0-1500	31-08-1905	Princesse-Alice II	Large Richard trawl
2204	39° 44' 0"	29° 29' 0"	Surface	31-08-1905	Princesse-Alice II	Plankton high speed net
2210	39° 25' 0"	31° 22' 30"	1229	01-09-1905	Princesse-Alice II	Bottom trawl
2211	39° 27' 0"	31° 22' 30"	1229	01-09-1905	Princesse-Alice II	Bottom long-line
2212	39° 26' 0"	31° 23' 30"	0-1200	02-09-1905	Princesse-Alice II	Large Richard trawl
2214	39° 26' 10"	31° 21' 30"	914-650	02-09-1905	Princesse-Alice II	Bottom trawl
2215	39° 26' 10"	31° 21' 30"	Surface	02-09-1905	Princesse-Alice II	Several
2218	39° 20' 0"	31° 0' 0"	Surface	02-09-1905	Princesse-Alice II	Plankton high speed net
2241	37° 45' 0"	29° 0' 0"	Surface	05-09-1905	Princesse-Alice II	Plankton high speed net
2242	37° 6' 0"	28° 10' 0"	Surface	06-09-1905	Princesse-Alice	Plankton high speed net
2244	37° 4' 0"	28° 1' 0"	0-3000	06-09-1905	Princesse-Alice II	Large Richard trawl

2245	37° 3' 0"	27° 46' 0"	Surface	06-09-1905	Princesse-Alice II	Plankton high speed net
2248	37° 2' 30"	27° 35' 0"	1478	06-09-1905	Princesse-Alice II	3 sinkers de sonda
2249	37° 2' 30"	27° 35' 0"	Surface	06-09-1905	Princesse-Alice II	Plankton high speed net
2252	38° 0' 0"	26° 10' 0"	Surface	07-09-1905	Princesse-Alice II	Plankton high speed net
2263	37° 33' 0"	23° 30' 0"	Surface	13-09-1905	Princesse-Alice II	Plankton high speed net
2264	37° 30' 0"	22° 39' 0"	0-3000	13-09-1905	Princesse-Alice II	Large Richard trawl
2266	37° 30' 0"	22° 35' 0"	Surface	13-09-1905	Princesse-Alice II	Plankton high speed net
2267	37° 28' 0"	21° 30' 0"	Surface	13-09-1905	Princesse-Alice II	Plankton high speed net
3131	35° 09' 00"	21° 21' 0"	0-3500	16-08-1911	Hirondelle II	High speed Bourée trawl
3132	35° 30' 00"	22° 2' 0"	Surface	16-08-1911	Hirondelle II	Plankton high speed net
3137	37° 0' 0"	25° 0' 0"	1330	17-08-1911	Hirondelle II	Swab bar
3144	37° 40' 30"	25° 58' 0"	919	25-08-1911	Hirondelle II	Swab bar
3147	38° 12' 00"	25° 49' 0"	0-550	26-08-1911	Hirondelle II	High speed Bourée line
3150	38° 1' 0"	25° 21' 0"	1740	27-08-1911	Hirondelle II	Midwater trawl
3231	35° 35' 0"	25° 40' 0"	0-100	11-08-1912	Hirondelle II	Plankton pump
3234	36° 39' 20"	25° 56' 0"	0-100	12-08-1912	Hirondelle II	Plankton pump
3256	38° 16' 40"	28° 23' 0"	0-1200	19-08-1912	Hirondelle II	High speed Bourée trawl
3257	38° 16' 40"	28° 23' 0"	0-100	19-08-1912	Hirondelle II	Plankton pump
3260	38° 30' 0"	28° 59' 0"	0-100	19-08-1912	Hirondelle II	Plankton pump
3279	38° 55' 00"	34° 07' 30"	0-3000	23-08-1912	Hirondelle II	High speed Bourée trawl
3282	39° 19' 0"	35° 27' 30"	0-100	24-08-1912	Hirondelle II	Plankton pump
3285	39° 23' 0"	35° 18' 30"	0-1000	24-08-1912	Hirondelle II	High speed Bourée trawl
3293	38° 47' 0"	30° 16' 0"	1331	26-08-1912	Hirondelle II	Bottom trawl
3526	38° 30' 00"	34° 40' 0"	0-2000	29-09-1913	Hirondelle II	Large Richard trawl
3608	38° 35' 0"	22° 57' 30"	0-2600	29-07-1914	Hirondelle II	High speed Bourée trawl