

Polychaete assemblages in the Argentinean Biogeographical Province, between 34° and 38°S

CLAUDIA BREMEC and DIEGO GIBERTO

Consejo Nacional de Investigaciones Científicas y Técnicas. Instituto Nacional de Investigación y Desarrollo Pesquero.
V. Ocampo 1, 7600 Mar del Plata, Argentina. e-mail: cbremec@inidep.edu.ar

SUMMARY: Polychaete assemblages were studied from samples taken in the Southwestern Atlantic, between 34°S and 38°S (Argentinean Biogeographical Province) during 1993, 2001 and 2002 from a series of oceanographic cruises in mixohaline to oceanic waters. Polychaetes represented almost 25% (number of taxa) of the macrobenthic assemblages; 57 taxa corresponding to 29 families were identified. The MDS plot shows an ordination of samples according to the following areas: the central estuary and mouth of Río de la Plata (group 1), the coastal shelves of Uruguay and Argentina and the transect off the mouth of Río de la Plata (groups 2 and 3) and other coastal stations from both countries (groups 4 and 5). The abiotic variable that best explained the distribution pattern (BIO-ENV analysis) was the bottom salinity, while the best combination included 2 variables: bottom salinity and type of bottom. Trophic diversity increased with salinity. The distributional range of four species: *Sabellaria bellis* Hansen, 1882; *Travisia amadoi* Elías *et al.*, 2003a; *Terebellides lanai* Solis-Weiss *et al.*, 1991 and *Aglaophamus uruguayi* Hartman, 1953, was extended.

Keywords: benthos, polychaetes, assemblages, SW Atlantic.

RESUMEN: AGRUPACIONES DE POLIQUETOS EN LA PROVINCIA BIOGEOGRÁFICA ARGENTINA, ENTRE 34° Y 38°S. – Se estudiaron las agrupaciones de poliquetos a partir de muestras recogidas en el Atlántico Sudoccidental, entre 34°S y 38°S (Provincia Biogeográfica Argentina) durante los años 1993, 2001 y 2002, en el marco de campañas oceanográficas llevadas a cabo desde aguas mixohalinas hasta marinas. Los poliquetos representan casi el 25% (número de taxones) de la agrupación macrobentónica; fueron identificados 57 taxones pertenecientes a 29 familias. El ordenamiento multidimensional (MDS) de estaciones distingue diferentes áreas: estuario central y boca del Río de la Plata (grupo 1), plataformas costeras de Uruguay y Argentina, y transecta fuera del Río de la Plata (grupos 2 y 3) y otras estaciones costeras de ambos países (grupos 4 y 5). La variable abiótica que mejor explicó el patrón de distribución (análisis BIO-ENV) fue la salinidad de fondo; la mejor combinación incluyó dos variables: salinidad de fondo y tipo de fondo. La diversidad trófica incrementó con la salinidad. Se extiende el rango de distribución de cuatro especies: *Sabellaria bellis* Hansen, 1882, *Travisia amadoi* Elías *et al.*, 2003a, *Terebellides lanai* Solis-Weiss *et al.*, 1991 y *Aglaophamus uruguayi* Hartman, 1953.

Palabras clave: bentos, poliquetos, asociaciones, Atlántico Sudoccidental.

INTRODUCTION

Few studies on benthic polychaetes have been undertaken in the Southwestern Atlantic, especially given the wide geographical area involved. As a result, the group has been excluded from previous analyses of the structure and spatial distribution of benthic communities in the Argentine Sea (i.e. Bastida *et al.*, 1992). Currently, the presence of

polychaetes in southern regions is mainly monitored as part of the benthic assemblages in areas subjected to different anthropogenic impacts: trawling fisheries (e.g. Bremec and Lasta, 2002; Roux *et al.*, 2002, 2005) or pollution (e.g. Elías and Bremec, 1994; Elías *et al.*, 2004).

Our study area comprises the Uruguayan and northern Argentinean shelves, both belonging to the Argentinean Biogeographical Province (Boschi,

2000a). Polychaetes from these areas were studied in the past decades (e.g. Orensanz, 1972; 1976; Orensanz and Gianuca, 1974; Rullier and Amoureux, 1979). More recently, Salazar-Vallejo and Orensanz (1991), Lana and Bremec (1994), Bremec and Elías (1999), Elías and Bremec (1997, 2003), Elías *et al.* (2000, 2003b) and Palacios *et al.* (2005) provided taxonomical information on many other families, both for the present area as well as for a wider range from Patagonia to southern Brazil.

In this paper we analyse macrobenthic samples taken between 34°S and 38°S, in 1993, 2001 and 2002 during oceanographic surveys in mixohaline to oceanic waters. This contribution provides a preliminary characterisation of local polychaete assemblages and extends biogeographic knowledge of the polychaete fauna in the Argentinean Biogeographical Province.

MATERIALS AND METHODS

The material studied was collected at 49 stations, between 34 and 38°S, in a depth range of 8-270 m (Fig. 1). Samples were obtained during 3 surveys:

cruise EH-09-1993 (epibenthic sledge, 200 × 50 cm mouth opening, soft bottoms); cruise EH-09-2001 (Picard dredge, 65 × 20 cm mouth opening, in sandy bottoms and van Veen grab, 0.1 m², in muddy bottoms); cruise CC-02-2002 (rectangular dredge, 2.5 × 0.5 m mouth opening, in gravel-sandy bottoms of a mussel bed and a lemon-fish feeding ground). As different gears were used to collect the samples from different types of bottoms, we present the results in terms of the relative abundance of the taxa found.

Benthic samples were sieved (0.5 mm mesh screen) and specimens were fixed in 4% formalin, preserved in 70% alcohol and identified to the lowest taxonomic level possible. In order to indicate the relative proportion of polychaetes within the benthic assemblages the total number of taxa of the major invertebrate groups was summed and the percentage of taxa of each group was estimated for the 3 surveys.

Cluster analysis (Bray-Curtis similarity index, UPGMA, Q mode) (Clarke and Warwick, 1993) and non-metric multidimensional scaling (MDS) (Clarke, 1993) were applied to the 4th root transformed relative abundance (each species count divided by the total abundance per sample) of poly-




FIG. 1. – Location of the 49 sampling stations between 34°S and 38°S.

TABLE 1. – Depth, bottom temperature, bottom salinity and bottom type of the sampling stations between 34°S and 38°S.

Station	Depth (m)	Bottom temperature (°C)	Bottom salinity(psu)	Bottom type
1	11,5	16.782	31.114	sandy
2	20	14.973	32.088	sandy
3	33	14.736	32.407	sandy
4	44	12.8865	32.9025	sandy
5	54	11.037	33.398	sandy
6	74	10.262	33.589	sandy
7	79	8.145	33.609	sandy
8	270	4.705	34.094	sandy
9	125	7.007	33.757	sandy
10	60	9.257	33.603	sandy
11	46	12.019	33.240	sandy
12	24	14.575	32.270	sandy
13	10	19.348	18.971	muddy
14	8	19.507	17.100	muddy
15	15	12.949	20.285	sandy
16	26	13.095	32.363	sandy
17	39	11.558	33.468	sandy
18	23	12.607	31.526	sandy
19	37	11.139	33.442	sandy
20	22	10.098	29.991	sandy
21	21	11.480	33.532	sandy
22	31	10.297	32.247	sandy
23	36	12.098	29.629	sandy
24	25	11.435	33.571	sandy
25	23	10.290	30.788	sandy
26	27	11.833	33.759	sandy
27	20	10.239	32.108	sandy
28	22	10.542	30.226	muddy
29	8	11.281	15.056	muddy
30	14	11.399	13.715	muddy
31	11	11.054	13.101	muddy
32	9	10.350	14.655	muddy
33	11	9.726	20.235	muddy
34	14	9.274	21.633	muddy
35	15	9.025	33.750	muddy
36	11	9.850	32.988	muddy
37	17	9.750	33.568	muddy
38	18	9.780	33.246	sandy
39	21	9.500	33.705	sandy
40	27	19.940	32.690	sandy-gravel
41	25	19.590	32.680	sandy-gravel
42	38	19.780	33.350	sandy-gravel
43	38	19.780	33.400	sandy-gravel
44	38	19.780	33.350	sandy-gravel
45	24	19.793	32.690	sandy-gravel
46	24	19.780	33.350	sandy-gravel
47	36	19.780	33.350	sandy-gravel
48	22	10.484	33.736	sandy
49	25	19.850	32.700	sandy-gravel

chaetes. The SIMPER analysis (Clarke, 1993) was used to identify which species contributed most to the similarity between samples. Stations 48 and 49, where only one polychaete taxon was collected, were excluded from the data matrix.

The classification of taxa into trophic categories was based on Fauchald and Jumars (1979), following the general trophic assumptions adopted in similar investigations worldwide (Muniz and Pires, 1999; 2000; Hutchings and Peyrot-Clausade, 2002; Arasaki *et al.*, 2004; Hughes and Gage, 2004). Cluster analysis (Bray-Curtis similarity index, UPGMA, Q mode) was applied to the presence-

absence data of trophic categories (carnivore, deposit-feeder, omnivore, filter/suspension feeder) .

Environmental data (Table 1) were taken from Giberto *et al.* (2004, EH-09-1993) and Giberto and Bremec (2003a, b, EH-09-2001 and CC-02-02 respectively). The BIO-ENV analysis was used to determine which set of variables (bottom salinity, depth, bottom temperature, type of bottom, similarity calculated with the Euclidean distance coefficient) best explains biological variation patterns (relative abundance data using Bray-Curtis similarity measure) (Clarke and Ainsworth, 1993; Clarke and Warwick, 2001).

TABLE 2. – List of taxa and their feeding types found between 35°S-38°S during the cruises in 1993, 2001 and 2002. (DF, deposit feeder; FF/SS, filter/suspension feeder; CC, carnivore; OM, omnivore).

Taxa	Feeding type	Taxa	Feeding type
ORBINIIDAE		ONUPHIDAE	
Orbiniidae unid.1	DF	<i>Diopatra viridis</i> Kinberg, 1865	OM
SPIONIDAE		<i>Diopatra</i> sp.	OM
Spionidae unid.1	DF	<i>Onuphis aff. eremita</i> Audouin and Milne Edwards, 1834	OM
MAGELONIDAE		<i>Onuphis tenuis</i> Hansen, 1881	OM
<i>Magelona riojai</i> Jones, 1963	DF	<i>Onuphis</i> sp.	OM
CHAETOPTERIDAE		Onuphidae unid.1	OM
<i>Phyllochaetopterus socialis</i> Claparède, 1870	FF/SS	EUNICIDAE	
Chaetopteriidae unid.1	FF/SS	<i>Eunice argentinensis</i> (Treadwell, 1929)	CC
CIRRATULIDAE		<i>Eunice magellanica</i> McIntosh, 1885	CC
Cirratulidae unid.1	DF	Eunicidae unid.1	CC
CAPITELLIDAE		LUMBRINERIDAE	
<i>Capitella</i> sp.	DF	<i>Augeneria aff. tentaculata</i> Monro, 1930	CC
MALDANIDAE		<i>Lumbrineriopsis mucronata</i> (Ehlers, 1908)	CC
Maldanidae unid.1	DF	<i>Lumbrineris</i> sp.	CC
OPHELIIDAE		<i>Ninoe brasiliensis</i> Kinberg, 1865	CC
<i>Armandia loboii</i> Elías and Bremec, 2003	DF	<i>Ninoe falklandica</i> Monro, 1936	CC
<i>Travisia amadoi</i> Elías <i>et al.</i> , 2003	DF	<i>Ninoe</i> sp.	CC
PHYLLODOCIDAE		Lumbrineridae unid.1	CC
Phyllodocidae unid.1	CC	Lumbrineridae unid.2	CC
APHRODITIDAE		OENONIDAE	
<i>Aphrodita longicornis</i> Kinberg, 1855	CC	<i>Drilonereis</i> sp.	CC
<i>Aphroditella alta</i> (Kinberg, 1855)	CC	Oeononidae unid.1	CC
POLYNOIDAE		<i>Lysarete brasiliensis</i> Kinberg, 1865	CC
<i>Lagisca</i> sp.	CC	OWENIIDAE	
Polynoidae unid.1	CC	<i>Owenia</i> sp.	DF/SS
SIGALIONIDAE		SABELLARIIDAE	
<i>Sigalion cirriferum</i> Orensanz and Gianuca, 1974	CC	<i>Idanthyrsus armatus</i> Kinberg, 1867	FF/SS
SYLLIDAE		<i>Sabellaria bellis</i> Hansen, 1882	FF/SS
Syllidae unid.1	CC	PECTINARIIDAE	
NEREIDIDAE		<i>Cistenides ehlersi</i> (Hessle, 1917)	DF
<i>Neanthes succinea</i> (Frey and Leuckart, 1847)	OM	Pectinariidae unid.1	DF
Nereididae unid.1	OM	AMPHARETIDAE	
GLYCERIDAE		<i>Ampharete</i> sp.	DF
<i>Glycera americana</i> Leidy, 1855	CC	Ampharetidae unid.1	DF
Glyceridae unid.1	CC	TEREBELLIDAE	
GONIADIDAE		Terebellidae unid.1	DF
Goniadidae unid.1	CC	TRICHOBRANCHIDAE	
NEPHTYIDAE		<i>Terebellides lanai</i> Solis-Weiss <i>et al.</i> , 1991	DF
<i>Nephtys</i> sp.	CC	Trichobranchiidae unid.1	DF
<i>Aglaophamus uruguayi</i> Hartman, 1953	CC	SABELLIDAE	
Nephtyidae unid.1	CC	Sabellidae unid.1	FF/SS
		SERPULIDAE	
		Serpulidae unid.1	FF/SS

RESULTS

Faunistic analysis

A total of 57 polychaete taxa distributed among 29 families was found (Table 2). The percentage of polychaetes which represent the fraction of the macrofauna larger than 0.5mm in the bottom samples was rather constant (20-24%) in the different habitats studied between 34°S and 38°S. Lumbrineridae and Onuphidae were the most diverse, with 8 and 6 taxa respectively. The highest frequency of occurrence was 44% (21 stations) for Onuphidae and 24% (13 stations) for Lumbrineridae. The other families were collected in 1-12 sampling stations.

The number of invertebrate species collected in soft bottoms was 128 and 141, during 1993 and 2001 respectively, and 118 in mussel and lemon-fish grounds during 2002. The faunistic composition of the benthic assemblages sampled in the 3 cruises showed a similar relative percentage of polychaete species. Molluscs (25-28%), Crustaceans (22-37%) and Polychaetes (20-24%) were the most diverse groups.

Polychaete assemblages

The cluster analysis produced 5 main groupings of stations (15% similarity level) and the corresponding MDS plot (Fig. 2) shows an ordination of samples according to the following areas: -the central




FIG. 2. – MDS ordination (Bray-Curtis similarity index) of the sampling stations based on relative abundance of polychaetes within them. Groups 1 (\diamond), 2 (Δ), 3 (\square), 4 (O) and 5 (V).

estuary and mouth of Río de la Plata from 8 to 27 m depth, reaching the Uruguayan coast at 39 m depth (group 1), - the coastal shelves of Uruguay (15-39 m) and Argentina (20-74 m) and the transect off the mouth of Río de la Plata, between 24 and 270 m depth (groups 2 and 3). Other coastal stations from both countries are clustered in groups 4 and 5. The results of the SIMPER test show that group 1 clusters stations (average similarity 21.40) where Polynoidae, Onuphidae and Glyceridae are the most representative taxa. Group 2 (average similarity 29.95) represents areas with *Ampharete* sp. and Serpulidae and group 3 (average similarity 15.21) with *Lagisca* sp., *Eunice argentinensis*, *Lumbrineris* sp., *Diopatra viridis*, *Phyllochaetopterus socialis*, and Maldanidae and Cirratulidae, as the most rep-

TABLE 3. – Combinations of n variables giving the largest rank Spearman correlations between biotic (4th root transformed relative abundance data, Bray-Curtis similarity) and environmental (Euclidean distance) similarity matrices (BIOENV analysis). Bold type indicates overall optimum.

n	Spearman's Correlation Coefficient (ρ_w)	Variables
1	0.205	Bottom salinity
	0.160	Bottom type
2	0.011	Bottom temperature
	0.237	Bottom salinity, bottom type
	0.133	Bottom temperature, bottom salinity
3	0.011	Bottom temperature, bottom type
	0.159	Bottom temperature, bottom salinity, bottom type
	-0.126	Depth, bottom salinity, bottom type
	-0.085	Depth, bottom temperature, bottom salinity
	-0.046	Depth, bottom temperature, bottom type

representative taxa. The species that contribute most to the average similarity of group 4 (35.00) are *Nephtys* sp. and *Glycera americana*, and *Onuphis tenuis* to the average similarity of group 5 (51.41). Stations 32 and 47 are characterized by Nephtyidae.

Environmental analysis

The matching of the environmental and faunal matrices (BIO-ENV analysis) is outlined in Table 3. No single variable provided the best correlation with biotic patterns. The abiotic variable that best explained the distribution pattern was the bottom salinity, while the best combination included 2 variables: bottom salinity and type of bottom.




FIG. 3. – Cluster analysis of stations (Bray-Curtis similarity index) according to presence-absence of polychaete feeding types in the study area (34°S-38°S). (cluster 1, deposit and filter feeders; cluster 2, omnivores; cluster 3, carnivores; cluster 4, mainly carnivores and deposit-feeders with the addition of omnivores and suspension feeders).

Feeding types

The cluster analysis performed with general feeding types (carnivore, omnivore, deposit-feeder, filter/suspension-feeder) as variables shows 4 main groups at 40% similarity level (Fig. 3). Group 1 clusters uruguayan and argentinian coastal stations where deposit-feeders (Ampharetidae) and filter-feeders (Serpulidae) are prevalent. Group 2 clusters stations with omnivores (Onuphidae) from the estuarine area. Group 3 clusters other coastal stations where carnivores (Polynoidae, Glyceridae, Lumbrineridae, Nephtyidae) are conspicuously present. The sampling stations deeper than 23 m are clustered in group 4, with carnivores (Eunicidae, Lumbrineridae, Polynoidae) and deposit-feeders (Ampharetidae, Maldanidae, Cirratulidae) as the main feeding types, together with omnivores and suspension feeders.

DISCUSSION

As the sampled area included a variety of substrata ranging from consolidated sandy-gravel to soft sediments, it was not feasible to use the same sampling device. Consequently, due to this methodological constraint, the information presented here is based on the relative abundance and distribution of taxa. This contribution intends to compile information for a large area and future studies will attempt to obtain comparative quantitative samples from the different habitats, throughout the geographical range under study.

The dominance of polychaetes in a variety of macrofaunal assemblages and geographical regions is well documented, even though comparing data sets is difficult due to differences in sampling techniques, mesh size used and number of replicates (i.e. Hutchings, 1998 and references). In previous studies in the region, in northern temperate latitudes (32°S - 34°S), the number of polychaete species reaches 36% of the total number of species collected in a large variety of bottoms between the inner shelf and the upper slope, and the number of specimens represents 48% of the total number of macroinvertebrates (Capitoli, 2002). In southern areas of the southwest Atlantic, these estimates reach 9% in sandy bottoms on the Patagonian shelf and 35% in diverse substrata in the Magellan Strait (Gambi and Mariani, 1999; Bremec *et al.*, 2000).

The multivariate analysis produced 5 different clusters of stations characterized by different suites of polychaetes in the study area. The spatial distribution of these sampling stations and the results of the SIMPER analysis clearly show that, in general, the polychaete fauna from the central estuary and mouth of the river (group 1) at almost 30m depth, is different from that collected at other coastal stations (groups 2, 4 and 5). Secondly, the deeper stations are clustered in group 3 and characterized by another suite of polychaetes. The BIO-ENV analysis indicates that species are distributed mainly according to a salinity gradient and bottom type. These variables serve to explain the patterns of faunal distribution in this study area that comprises an estuarine muddy sector and sandy or gravel-sandy bottoms of the adjacent marine waters, and a few stations on the open shelf. The granulometric type is the main abiotic variable that determines the spatial distribution of polychaete assemblages in other South American systems influenced by continental waters, like Lagoa dos Patos on the south Brazilian shelf (Capitoli *et al.*, 2004) and Bahía Blanca estuary on the Argentinean shelf (Elías and Bremec, 1994). The close relationship between benthic fauna and sediment type has previously been recognised in classical papers regarding spatial distribution of benthos (e.g. Thorson, 1957; Sanders, 1958; 1968; Lie and Kelley, 1970; Day *et al.*, 1971) and sediment was considered as an indicator of the availability of feeding resources (Sanders *et al.*, 1962; Rhoads and Young, 1970; Young and Rhoads, 1971). Recent investigations continue to show that the sediment characteristics are primary factors determining the spatial patterns of macrobenthos and polychaete assemblages (e.g. Muniz and Pires, 2000; Probert *et al.*, 2001; Hernández-Arana *et al.*, 2003; Rodríguez-Villanueva *et al.*, 2003; Dauvin *et al.*, 2004; Díaz-Castañeda and Harris, 2004).

Mainly omnivores (Onuphidae) and carnivores (Polynoidae, Glyceridae, Lumbrineridae, Nephtyidae) characterize the different stations in the estuarine muddy areas. A variety of feeding types were found in the rest of the sandy coastal stations and in areas deeper than 23m. Studies on spatial distribution of macrobenthos in the study area show a higher biomass of deposit-feeders and carnivorous molluscs in the estuarine area and various feeding types widely distributed in the marine and open waters (Giberto *et al.*, 2004). Polychaetes, as a group, reflect the trend observed for the benthic

community as a whole. It was found that generally trophic diversity increased with salinity, meaning that a more even distribution of trophic structure is found at higher salinities (Gaston *et al.*, 1998), which suggests differences in the availability of resources and in the interactions of food webs (Brown *et al.*, 2000). The area studied comprises the Río de la Plata estuary and the adjacent marine coastal and open waters and little is known about trophic webs. The large estuarine zone (38,000 km²) has both bottom and surface salinity fronts, important in fish reproductive processes and where high concentrations of zooplankton occur, and a turbidity front in the inner estuary (Mianzan *et al.*, 2001a), with high biomass of deposit-feeding bivalves (Giberto *et al.*, 2004). The whole study area is disturbed by trawling fisheries, many of the catches of which are dependent on benthic food resources. Thus future studies need to be conducted to assess the functional role of these benthic communities and in particular of polychaetes.

Our present records extend the distributional range of *Sabellaria bellis* from the mouth of the Río de la Plata estuary at 35°S (Bremec and Giberto, 2004) to 38°S (Station 7, 79 m depth), and the bathymetric range of *Travisia amadoi* from intertidal and shallow sandy bottoms (Elías *et al.*, 2003a) to depths of 125 and 270 m (Stations 9 and 8 respectively). Additionally, the southern distribution range of two more species is extended: *Terebellides lanai*, originally described from southern Brazil (Bremec and Elías, 1999), was collected at 35°S (Station 22) and *Aglaophamus uruguayi*, described for muddy bottoms at 33°S 51°W (Orensanz and Gianuca, 1974), was collected between 35°S and 37°S, at 24 m depth (Stations 45 and 12) and in deeper sandy substrata at 125-270 m depth (Stations 9 and 8). Our study material was collected in warm temperate waters of the Argentinean Biogeographical Province (Boschi, 2000a). It includes coastal waters from latitude 43-44°S (Patagonia) reaching 23°S (southern Brazil). The Río de la Plata (35°S) is the major hydrographical feature within this province (Boschi, 2000b), and considered a significant zoogeographic barrier. However, this barrier was found to be variable for plankton and fish species, according to climatic variation and discharge patterns of the river (Mianzan *et al.*, 2001b). In the case of the decapod *Ebalia rotundata* (A. Milne Edwards, 1880), found in marine waters off Argentina and Uruguay, it was considered that both thermal and salinity tolerance could con-

strain the dispersal of the species from south to north and within estuarine waters respectively (Giberto and Bremec, 2003c). Local patterns of larval transport could allow the colonization of new areas from Brazil to the south and the role of the Río de la Plata regime as a biological barrier to the distribution of coastal polychaete fauna between Uruguay and Argentina still has to be assessed. More frequent sampling and detailed examination of the polychaetes will almost certainly lead to new records and provide more detailed distributional patterns.

ACKNOWLEDGEMENTS

We thank the reviewers for valuable comments, which greatly improved the manuscript. This study was partially supported by Antorchas Foundation 13817-5 and 13900-13, ANPCyT PICT n° 8424 and 15080, PNUD/GEF/RLA/99/G31/Biología N°022 and PEI 6026 (CONICET). This is INIDEP Contribution N° 1369.

REFERENCES

- Arasaki, E., P. Muniz and A.M.S. Pires-Vanin. – 2004. A functional analysis of the benthic macrofauna of the São Sebastião Channel (Southeastern Brazil). *P.S.Z.N.: Mar. Ecol.*, 25(4): 249-263.
- Bastida, R., A. Roux and D. Martínez. – 1992. Benthic communities of the Argentine continental shelf. *Oceanol. Acta*, 15(6): 687-698.
- Boschi, E.E. – 2000a. Species of Decapod Crustaceans and their distribution in the American marine zoogeographic provinces. *Rev. Invest. Des. Pesq.*, 13: 7-136.
- Boschi, E.E. – 2000b. Biodiversity of marine decapod brachyurans of the Americas. *J. Crust. Biol.*, 20: 337-342.
- Bremec, C. and R. Elías. – 1999. Species of *Terebellides* from South Atlantic waters off Argentina and Brazil (Polychaeta: Trichobranchidae). *Ophelia*, 51(3): 177-186.
- Bremec, C. and D. Giberto. – 2004. New records of two species of *Sabellaria* (Polychaeta, Sabellariidae) from the Argentinean Biogeographic Province. *Rev. Biol. Mar. Oceanogr.*, 39(2): 101-105.
- Bremec, C. and M. Lasta. – 2002. Epibenthic assemblage associated with scallop (*Zygochlamys patagonica*) beds in the Argentinian shelf. *Bull. Mar. Sci.*, 70(1): 89-105.
- Bremec, C., R. Elías and M.C. Gambi. – 2000. Comparison of polychaete community composition from the patagonian shelf and Strait of Magellan, preliminary results from cruises "Shinkai Maru" IV, V, X and XI (1978-1979) and 2nd. P.N.R.A. Oceanographic Cruise (1991). *Bull. Mar. Sci.*, 67(1): 189-197.
- Brown, S., G. Gaston, C. Rakocinski and R. Heard. – 2000. Effects of sediment contaminants and environmental gradients on macrobenthic community trophic structure in Gulf of Mexico estuaries. *Estuaries*, 23(3), 411-424.
- Capitoli, R. – 2002. *Distribuição e abundância dos macroinvertebrados bentônicos da plataforma continental e talude superior no extremo sul do Brasil*. Ph. D. thesis, Univ. Rio Grande.
- Capitoli, R., C. Bremec, R. Elías and D. Giberto. – 2004. Polychaetes assemblages from the Southbrazilian Shelf. *Res. VIII International Polychaete Conference*, 5-9 jul., Madrid, p. 45.
- Clarke, K.R. 1993. – Non-parametric multivariate analysis of

- changes in community structure. *Austr. J. Ecol.*, 18: 117-143.
- Clarke, K.R. and M. Ainsworth. – 1993. A method of linking multivariate community structure to environmental variables. *Mar. Ecol. Progr. Ser.*, 92: 205-219
- Clarke, K.R. and R. Warwick. – 1993. Environmental effects on benthic communities. Training Workshop on "Multivariate Analysis of Benthic Community Data". Lecture Notes for SEAS/EPOSII Workshop, Plymouth Marine Laboratory
- Clarke, K.R. and R. Warwick. – 2001. *Change in marine communities: an approach to statistical analysis and interpretation*, 2nd ed. PRIMER-E, Plymouth.
- Dauvin, J.C., E. Thiébaud, J.L., J.L. Gomes Gesterira, K. Ghertsois, F. Gentil, M. Ropert, B. Sylvand. – 2004. Spatial structure of a subtidal macrobenthic community in the Bay of Veys (western Bay of Seine, English Channel). *J. Exp. Mar. Biol. Ecol.*, 307: 217-235.
- Day, J.M., J. Field and M. Montgomery. – 1971. The use of numerical methods to determine the distribution of the benthic fauna across the continental shelf of North Carolina. *J. Anim. Ecol.*, 40: 93-125.
- Díaz-Castañeda, V. and L.H. Harris. – 2004. Biodiversity and structure of the polychaete fauna from soft bottoms of Bahía Todos Santos, Baja California, Mexico. *Deep-Sea Res. II*, 51: 827-847.
- Elías, R. and C.S. Bremec. – 1994. Biomonitoring of water quality using benthic communities in Blanca Bay (Argentina). *Sc. Total Envir.*, 158(1-3): 45-49.
- Elías, R. and C.S. Bremec. – 1997. First record of Magelonidae, *Magelona riojai*, Jones 1963, (Polychaete) in coastal waters of Argentina. *Nerítica*, 11(1/2): 111-117.
- Elías, R. and C.S. Bremec. – 2003. First record of the genus *Armandia* (Polychaeta, Opheliidae) in coastal waters of Argentina, with the description of a new species *Armandia loboi*. *Bull. Mar. Sci.*, 72(1): 181-186.
- Elías, R., C.S. Bremec, P.C. Lana and J.M. Orenzan. – 2003a. Opheliidae (Polychaeta) from the southwestern Atlantic ocean, with the description of *Travisia amadoi* n. sp., *Ophelina gaucha* n. sp. and *Ophelina alata* n. sp. *Hydrobiologia*, 496(1/3): 75-85.
- Elías, R., C. Bremec and E.A. Vallarino. – 2000. *Protoariciella uncinata* Hartmann-Schröder, 1962 (Polychaeta, Orbiniidae): a new record for intertidal mussel beds of the Southwestern Atlantic shore. *Rev. Biol. Mar. Oceanogr.*, 35(2): 181-184.
- Elías, R., M.S. Rivero and E.A. Vallarino. – 2003. Sewage impact assessment based on the composition and distribution of Polychaetes associated to intertidal mussel beds of the Southwestern Atlantic shore. *Iheringia*, 93(3): 309-318.
- Elías, R., E.A. Vallarino, M. Scagliola and F.I. Isla. – 2004. Macrobenthic distribution pattern at a sewage disposal site in the inner shelf off Mar del Plata (SW Atlantic). *J. Coast. Res.*, 20(4): 1176-1182.
- Fauchald, K. and P. Jumars. – 1979. The diet of worms: a study of polychaete feeding guilds. *Oceanogr. Mar. Biol. Ann. Rev.*, 17: 193-284.
- Gambi, M.C. and S. Mariani. – 1999. Polychaetes of the soft bottoms of the Straits of Magellan collected during the Italian oceanographic cruise in February-March 1991. *Sci. Mar.*, 63 (Supl. 1): 233-242.
- Gaston, G., C. Rakocinski, S. Brown and C. Cleveland. – 1998. Trophic function in estuaries: response of macrobenthos to natural and contaminant gradients. *Mar. Freshw. Res.*, 49: 833-846.
- Giberto, D. and C.S. Bremec – 2003a. Benthic diversity of the Río de la Plata estuary and adjacent marine waters. *Proyecto de las Naciones Unidas para el Desarrollo. PNUD Project/Gef RLA/99/G31*, 48 pp.
- Giberto, D. and C. Bremec. – 2003b. Diversidad bentónica de fondos duros de plataforma intermedia: bancos del Pez Limón y Mejillón (35-37° S). *Res. V Jornadas Nacionales de Ciencias del Mar*, 8-12 Dec, Mar del Plata., p. 113.
- Giberto, D. and C. Bremec. – 2003c. *Ebalia rotundata* A. Milne-Edwards, 1880 (Brachyuran, Leucosiidae) in marine waters off Uruguay and Argentina. *Crustaceana*, 76: 307-312.
- Giberto, D.A., C.S. Bremec, E.M. Acha and H. Mianzan. – 2004. Large-scale spatial patterns of benthic assemblages in the SW Atlantic: the Río de la Plata estuary and adjacent shelf waters. *Estuar. Coast. Shelf Sci.*, 61(1): 1-13.
- Hansen, A. – 1882. Recherches sur le Annélides recueillies par M. Le professeur Edouard Van Beneden pendant son voyage au Brésil et à la Plata. *Mém. Cour. Acad. Roy. Sci. Belgique*, Bruxelles, 44: 1-29.
- Hartman, O. – 1953. Non-pelagic polychaetes of the Swedish Antarctic Expedition. 1901-1903. *Forth. Zool. Swed. Antarct. Exped.*, 4(11): 1-83.
- Hernández-Arana, H.A., A.A. Rowden, M. Attrill, R. Warwick and G. Gold-Bouchot. – 2003. Large-scale environmental influences on the benthic macroinfauna of the southern Gulf of Mexico. *Estuar. Coast. Shelf Sci.*, 58: 825-841.
- Hughes, D.J. and J.D. Gage. – 2004. Benthic metazoan biomass, community structure and bioturbation at three contrasting deep-water sites on the northwest European continental margin. *Progr. Oceanogr.*, 63: 29-55.
- Hutchings, P. – 1998. Biodiversity and functioning of polychaetes in benthic sediments. *Biodiv. Conserv.*, 7: 1133-1145.
- Hutchings, P. and M. Peyrot-Clausade. – 2002. The distribution and abundance of boring species of polychaetes and sipunculans in coral substrates in French Polynesia. *J. Exp. Mar. Biol. Ecol.*, 269: 101-121.
- Lana, P.C. and C. Bremec. – 1994. Sabellariidae (Annelida, Polychaeta) from South America. *Mém. Mus. nat. Hist. nat.*, 162: 209-221.
- Lie, U. and J. Kelley. – 1970. Benthic infauna communities off the coast of Washington and in Puget Sound. Identification and distribution of the communities. *J. Fish. Res. Bd. Can.*, 27: 621-651.
- Mianzan, H., C. Lasta, M. Acha, R. Guerrero, G. Macchi and C. Bremec. – 2001a. Río de la Plata estuary: ecological profile of a large scale salt wedge. In: U. Seeliger and B. Kjerfve (eds.), *Coastal Marine Ecosystems of latin America*, pp. 185-204. Springer Verlag, Berlin.
- Mianzan, H.W., E.M. Acha, R. Guerrero, F. Ramírez, D. Sorraín, C. Simionato and J. Borus. – 2001b. South Brazilian marine fauna in the Río de la Plata estuary: discussing the barrier hypothesis. *Resúmenes Ampliados IXº Congreso Latinoamericano de Ciencias del Mar (COLACMAR)*, San Andrés, Colombia, (CD-ROM).
- Muniz, P. and A.M.S. Pires. – 1999. Trophic structure of polychaetes in the São Sebastião Channel (southeastern Brazil). *Mar. Biol.*, 134: 517-528.
- Muniz, P. and A.M.S. Pires. – 2000. Polychaete associations in a subtropical environment (São Sebastião Channel, Brazil): a structural analysis. *P.S.Z.N.: Mar. Ecol.*, 21(2): 145-160.
- Orensan, J.M. – 1972. Los anélidos poliquetos de la Provincia Biogeográfica Argentina. I. Palmyridae (=Chrysopetalidae), Amphinomidae y Euphrosinidae. *Physis*, 31(83): 485-501.
- Orensan, J.M. – 1976. Los anélidos poliquetos de la Provincia Biogeográfica Argentina. IX. Poecilochaetidae y Cossuridae. *Comunicaciones Zoológicas Mus. Hist. Nat. Montevideo*, 10(140): 1-11.
- Orensan, J.M. and N. Gianuca. – 1974. Contribuição ao conhecimento dos anelídeos poliquetos do Rio Grande do Sul, Brasil. I. Lista sistemática preliminar e descrição de tres novas espécies. *Comun. Mus. Ci. PUCRS*, P. Alegre (4): 1-37.
- Palacios, J.R., C. Bremec, M.S. Rivero and R. Elías. – 2005. First records of *Parandalia tricuspis* (Müller 1858) and *Sigambra cf. tentaculata* (Treadwell 1941) (Pilargidae: Polychaeta) in Argentina. *Rev. Biol. Mar. Oceanogr.*, 40(1): 1-6.
- Probert, P.K., G.B. Read, S.L. Grove and A.A. Rowden. – 2001. Macrobenthic polychaete assemblages of the continental shelf and upper slope off the west coast of the South Island, New Zealand. *N. Z. J. Mar. Freshwat. Res.*, 35(5): 971-984.
- Rhoads, D. and D. Young. – 1970. The influence of deposit feeding organisms on sediment stability and community trophic structure. *J. Mar. Res.*, 28: 151-178.
- Rodriguez-Villanueva, V., R. Martinez-Lara and V. Macías Zamora. – 2003. Polychaete community structure of the northwestern coast of Mexico: patterns of abundance and distribution. *Hydrobiologia*, 496(1-3): 385-399.
- Roux, A., J. Calcagno and C. Bremec. – 2002. Macrobenthic assemblages associated with demersal fisheries around South Georgias islands. R/V Eduardo Holmberg Survey, February-March, 1994. *Arch. Fish. Mar. Res.*, 49(3): 231-241.
- Roux, A., C. Bremec, L. Schejter and D. Giberto. – 2005. Benthic invertebrates by-catch of demersal fisheries: a comparison between Subantarctic and Antarctic shelf waters (45°S-57°S). *Ber. Polarforsch. Meeresforsch.*, 507: 179-181.
- Rullier, F. and L. Amoureux. – 1979. Annélides polychètes. Campagne de la Calypso au large des côtes atlantiques de

- l'Amérique du Sud (1961-1962). *Ann. Inst. Océanogr.*, 55, fasc. suppl.: 10-206.
- Salazar-Vallejo, S. and J. M. Orensanz. – 1991. Pilárgidos (Annelida: Polychaeta) de Uruguay y Argentina. *Cah. Biol. Mar.*, 32: 267-279.
- Sanders, H. – 1958. Benthic studies in Buzzards bay. I. Animal-sediment relationships. *Limnol. Oceanogr.*, 3: 245-258.
- Sanders, H. – 1968. Marine benthic diversity: a comparative study. *Am. Nat.*, 102: 243-282.
- Sanders, H., E. Gondsmith, E. Mills and G. Hampson. – 1962. A study of the intertidal fauna of Barnstable Harbour, Massachusetts. *Limnol. Oceanogr.*, 7: 63-79.
- Solis-Weiss, V., K. Fauchald and A. Blankenstejn. – 1991. Trichobranchidae (Polychaeta) from shallow warm water areas in the western Atlantic Ocean. *Proc. Biol. Soc. Wash.*, 104: 147-158.
- Thorson, G. – 1957. Bottom communities (sublittoral or shallow shelf). In: J.W. Hedgpeth (ed.), *Treat. Mar. Ecol Palaeoecol.*, pp. 461-534. Geol. Soc. Amer. Mem.
- Young, D. and D. Rhoads. – 1971. Animal-sediment relations in Cape Cod Bay, Massachusetts. I. A transect study. *Mar. Biol.*, 11: 242-254.

Received July 9, 2004. Accepted September 12, 2005.