

Lower Badenian fish otoliths of the Styrian and Lavanttal basins, with a revision of WEINFURTER's type material

By Dirk NOLF¹ & Rostislav BRZOBOHATÝ²

(With 2 figures, 8 plates and 2 tables)

Manuscript submitted on March 31st 2008,
the revised manuscript on September 24th 2008

Abstract

Study of the Middle Miocene (Badenian) otolith collections in the Landesmuseum Joanneum in Graz, the Landesmuseum Kärnten in Klagenfurt, and the Krahuletz Museum in Eggenburg allowed for the reconstruction of a teleost fauna of 57 taxa, among which *Lethrinus styriacus* is new. The most diversified associations occur in the St. Florian Beds; the Mühldorf Formation provided 18 taxa. Five of those were only recorded from the last named formation. “genus Gobiidarum” *noricus* is reported only from the non-marine younger sediments in the Lavanttal Basin. Skeleton based taxa from the Eastern Styrian Basin (SCHULTZ 2000, 2004, 2006; SCHULTZ & BELLWOOD 2004) were not documented by otoliths from similar taxa in the Western Styrian Basin or the Lavanttal Basin. The associations reflect tropical to subtropical nearshore marine faunas that are quantitatively dominated by gobiids. Although the reconstructed fauna shows most affinities to present-day Atlantic and Mediterranean forms, the number of Indo-Pacific taxa (6) is also relatively high. Eight taxa (*Thrissa floriani*, *Arius nucleus*, *Ambassis* sp., *Gazza hilberi*, *Gerres* sp. 2, “genus Haemulidarum” *lucidus*, *Lethrinus styriacus*, and *Xyrichtys* sp.) have a Badenian record in the Styrian Basin only, and “genus Atherinidarum” *bavayi* occurs in the Lavanttal Basin only.

Keywords: Fish otoliths, palaeoenvironment, Styrian Basin, Lavanttal Basin, Central Paratethys, Austria, Badenian, Middle Miocene.

Zusammenfassung

Die mittel-miozänen (badenischen) Fischfaunen des Weststeirischen Beckens und des Lavanttal-Beckens wurden anhand der Otolithen in den Sammlungen des Landesmuseums Joanneum in Graz, des Landesmuseums Kärnten in Klagenfurt und des Krahuletz Museums in Eggenburg einer systematischen Revision unterzogen. Insgesamt konnten 57 Taxa nachgewiesen werden, darunter 1 neue Art: *Lethrinus styriacus*. Die artenreichste Assoziation der Otolithen kommt in den St. Florianer Schichten vor, während die Mühldorf-Formation nur 18 Arten beinhaltet. “genus Gobiidarum” *noricus* wurde nur in jüngeren ausgesäusten Ablagerungen des Lavanttales aufgefunden. Ein Nachweis der Formen die im Oststeirischen Becken auf Grund von osteologischem Material bekannt sind (SCHULTZ 2000, 2004, 2006; SCHULTZ & BELLWOOD 2004) war im Weststeirischen Becken und im Lavanttal anhand von Otolithen nicht möglich.

Die nachgewiesenen Otolithen spiegeln eine tropische bis subtropische Seichtwasserfauna mit einer Dominanz von Gobiiden wider. Auf Grund der Anwesenheit der tropischen Fische können die studierten Fundorte mit dem unteren Teil der Weissenegg-Formation im Oststeirischen Becken korreliert werden.

¹ Institut royal des Sciences naturelles de Belgique, 29 rue Vautier, 1000 Bruxelles, Belgique; e-mail: Dirk.Nolf@natuurwetenschappen.be

² Department of Geological Sciences, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic; e-mail: rosta@sci.muni.cz




Fig. 1. Position of the Styrian and Lavanttal basins (modified after BECHTEL et al. 2007).

Abgesehen davon, dass die behandelte Otolithenfauna Ähnlichkeiten mit den rezenten Atlantischen und Mittelmeeranen Formen zeigt, ist die Zahl von heute auf den Indo-Pazifik beschränkten Taxa (6) verhältnismäßig hoch. Acht Taxa (*Thrissa floriana*, *Arius nucleus*, *Ambassis* sp., *Gazza hilberi*, *Gerres* sp. 2, "genus *Haemulidarum*" *lucidus*, *Lethrinus styriacus* und *Xyrichtys* sp.) kommen im Badenium nur im Steirischen Becken und "genus *Atherinidarum*" *bavayi* nur im Lavanttal vor.

Schlüsselwörter: Pisces, Otolithen, Paläökologie, Steirisches Becken, Lavanttal-Becken, Zentrale Paratethys, Österreich, Badenium, Mittel-Miozän.

Introduction

The Badenian bony fishes of the Styrian Basin and the Lavanttal Basin are represented by otoliths, and rarely by isolated teeth and skeletal remnants. Results of primary studies on otoliths were published by WEINFURTER (1949, 1952b, c – Mühldorf Formation and overlaying brackish and freshwater beds, Lavanttal, Carinthia; 1952d – different localities, St. Florian Beds, Western Styrian Basin). Otoliths from the claypit Gasselsdorf (Eibiswald Beds, the southern bay of the Western Styrian Basin) were both taxonomically and stratigraphically evaluated by HIDEN & STINGL (1998). An additional species, "*Gobius* (n.) sp." was recently cited from the lower part ("Fish Shales") of the Mühldorf Formation by REISCHENBACHER et al. (2007).

The teeth and tooth plates of sparids and labrids were mentioned by SCHULTZ (in BRZOBOHATY & SCHULTZ 1978) from different localities in the Styrian Basin. Teeth and skeletal remnants of the species *Epinephelus casottii* (COSTA, 1858), *Trigonodon jugleri* (MÜNSTER, 1846) and *Balistes muensteri* SCHULTZ, 2004 were recently mentioned in SCHULTZ (2000 – locality Retznei; 2004 – localities Wiesfleck and Retznei) and in SCHULTZ & BELLWOOD (2004 – localities Wiesfleck, Weissenegg and Retznei). SCHULTZ (2006) described a fragmentary jaw as *Oligodiodon* sp. (porcupinefish) from the Retznei outcrop.

The present paper provides a revision of all otolith material from the Styrian and the Lavanttal basins based on the type material of the various papers of WEINFURTER and of additional unstudied material in the collections of the Joanneum Museum in Graz and the Kärntner Landesmuseum in Klagenfurt.




Fig. 2. Simplified stratigraphic chart of the Badenian fill of the Styrian and Lavanttal basins (modified after GROSS et al. 2007; BECHTEL et al. 2007; REISCHENBACHER et al. 2007; RÖGL et al. 2007).

Geological context

The Styrian Basin, palaeogeographically a western part of the Central Paratethys, is situated in the foothills of the Eastern Alps and represents the western part of the Pannonian Basin System belonging to the back-arc basin domain within the Alpine-Carpathian arc (fig. 1). By the Middle Styrian Swell, it is subdivided into the Western and the Eastern Styrian Sub basins. It is filled with Karpatian (Early Miocene) and Badenian (Middle Miocene) mostly marine sediments separated by the “Styrian Tectonic Phase” unconformity (RÖGL et al. 2002; POLESNY 2003; KOVÁČ et al. 2007) and various older and younger non-marine (or reduced marine) deposits.

Sedimentation of the Early Badenian in the Eastern Styrian Basin (Kreuzberg and Weissenegg formations) generally started with silts, fine sands, or conglomerates with *Praeorbulina sicana* and *P. transitoria* in the late NN4 nannoplankton Zone (fig. 2). Stratigraphically higher the corallinacean limestones and the transgressive coral facies (e.g., directly on top of the Karpatian Kreuzkrumpel Formation) with *P. circularis* and nannoplankton of the NN5 Zone were deposited along the shoreline or around basement highs, while deepwater marly and pelitic sediments prevailed in the basin and graben structures (Weissenegg Fm.). During the early Badenian marine sediments reached their

largest extent, whereas during the middle and late Badenian depocenters shifted eastwards (SPEZZAFERRI et al. 2004; RÖGL et al. 2007; GROSS et al. 2007). Tuffs in the clastic sediments overlaying corallinean limestones in the Retznei section (northeastern part of the Styrian Basin) show a radiometric age of 14.2 ± 0.1 Ma (BOJAR et al. 2004). Many features of sediments (sands, deltaic and limnic complexes) indicate a strong regression at the end of the Badenian time and erosion at the Badenian/Sarmatian boundary (EBNER & SACHSENHOFER 1995).

In the Western Styrian Subbasin, despite the presence of different coarse clastics at the western border, the Early Badenian is represented by calcareous clays and clayey sands and silts of the St. Florian Beds (an old German name is the “Florianer Tegel”) filling the Bay of St. Florian. These beds are palaeogeographically interpreted as lagoonal deposits with rich mollusks, sharks, foraminifera, and otoliths and stratigraphically equivalent to the Weissenegg Formation. To the south, they pass in the upper part of the Eibiswald Beds (sands and clayey silts) interpreted as shallow deltaic deposits of the Eibiswald Bay (HIDEN & STINGL 1998; PILLER et al. 2004; GROSS et al. 2007). Volcaniclastic rocks intercalated within the St. Florian Beds show a radiometric age of 15.75 ± 0.17 Ma (Lower Badenian, HANDLER et al. 2006).

The Lavanttal Basin represents a small pull-apart basin west of the Styrian Basin. It is downfaulted along the Pöls-Lavanttal Fault between the crystalline of the Saualpe and Koralpe and filled with Lower to Middle Miocene sediments (e.g., POLESNY 2003). The marine ingressions started in the Lower Badenian and the marine connection to the Central Paratethys ceased during the Middle and Late Badenian (KOVÁČ et al. 2007). The Lower Badenian Mühldorf Formation is separated by an unconformity and mainly consists of grey or brown mudstones and shelly marls with silty and sandy layers and with a lacustrine-brackish fauna at the base and higher with diverse marine molusc, bryozoa, foraminifera, and otolith faunas. In the marine upper part benthic foraminifera assemblages are similar with those of the Lower Lagenidae Zone in the Vienna Basin (RÖGL et al. 2002). Rich nannoplankton assemblages in combination with *Praeorbulina* and ostracods indicate the lower part of the nannoplankton Zone NN5 (REISCHENBACHER et al. 2007). Overlaying younger Badenian and Sarmatian beds are brackish to freshwater (BECHTEL et al. 2007).

Material & Methods

A list of all otolith-based taxa from the Badenian of the Western Styrian Basin and the Lavanttal Basin is given in Table 1. Table 2 provides an alphabetical list of all taxa cited in WEINFURTER's (1952b, c, d) papers, and their present taxonomical interpretation. Most of the material (from sites in the proximity of Wetzelsdorf and St. Florian, including the types of taxa described by WEINFURTER 1952d) belongs to the collections of the Landesmuseum Joanneum in Graz. The material from Mühldorf in Kärnten (including the type material described by WEINFURTER 1952c) is from the Kärnter Landesmuseum in Klagenfurt, and finally, a small collection of the Lavanttal is from the Krahuletz Museum, Eggenburg. For general matter about otoliths and their morphological nomenclature, the reader is referred to NOLF (1985 and 1995).

Abbreviations

The abbreviation “aff.” in the binomina was used in the cases of well-preserved material where specific identity could not be unequivocally determined. The abbreviation “cf.” was inserted in cases where the poor preservation of otoliths did not allow an adequate specific identification. Twelve species appear in open nomenclature due to limited or poorly preserved material.

GBW	Geologische Bundesanstalt Wien.
IRSNB	Institut royal des Sciences naturelles de Belgique, Bruxelles.
KME	Krahuletz Museum in Eggenburg.
LMJG	Landesmuseum Joanneum in Graz.
LMKK	Landesmuseum Kärnten in Klagenfurt.

Localities studied

All studied material is from the above cited museum collections. After the study of this material, D. NOLF did a short exploration trip in the area, which is not very appropriate for rapid exploration, due to the exuberant vegetation and the scattered but relatively dense countryside habitations. He sampled a rather indurated silty marl at $x = 675.700$, $y = 196.400$, in a steep ravine that runs off northward from the way that runs eastward from the road from St. Josef to Wetzelsdorf, at 500 m from the junction with this road, south of the place marked Bramberg on the topographic map. This sample (about 20 kg) only provided a single *Pomadasys* aff. *incisus* otolith (Coll. IRSNB). For the material in the rather old (mainly before 1950) museum collections, there are no data on the size of the samples or the method of collecting (surface-picked or screenwashed). Therefore, we did not provide data on the number of collected specimens. Based on the small size of many specimens and the great abundance of small gobiid otoliths, it is probable that most specimens came from screenwashed samples. Locality data for the Wetzelsdorf area are from HOLLER (1900), who provides a map and some descriptive data.

Gross Sankt Florian, Kogelbauer.

Österreichische Karte 1/50 000, sheet 189 (Deutschlandsberg),

$x = 672.950$, $y = 186.200$

Coll. Landesmuseum Joanneum.

Gross Sankt Florian, Mühlbauer.

Exact location unknown. Coll. Landesmuseum Joanneum, Graz.

Lavanttal, Mühlendorf.

Exact location unknown. Coll. Landesmuseum Kärnten in Klagenfurt.

Lavanttal, Weinzettel bei Ober-Aigen.

Exact location unknown. Coll. Landesmuseum Kärnten in Klagenfurt.

Wetzelsdorf, Kreuzschaller.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), $x = 675.450$, $y = 196.250$

Coll. Landesmuseum Joanneum, Graz.

Wetzeldorf, Rinngraben.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), x = 675.900, y = 194.250
 Coll. Landesmuseum Joanneum, Graz.

Wetzeldorf, Simihansl.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), x = 676.300, y = 193.800
 Coll. Landesmuseum Joanneum, Graz.

Wetzeldorf, Thomihiaslgraben.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), x = 675.400, y = 195.800
 Coll. Landesmuseum Joanneum, Graz.

Wetzeldorf, Wenzelsteffi.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), x = 675. 550, y = 196.150
 Coll. Landesmuseum Joanneum, Graz.

Wetzeldorf, Winkeltoni.

Österreichische Karte 1/50 000, sheet 190 (Leibnitz), x = 676.000, y = 195.250
 Coll. Landesmuseum Joanneum, Graz.

Remarks on taxa requiring comments

***Thrissa floriani* (WEINFURTER, 1952) (1)**

Otoliths of this species, originally described as a percoid by WEINFURTER (1952d), clearly belong to the engraulid genus *Thrissa*; see RIVATON & BOURRET (1999: pl. 2, fig. 15-16) for otoliths of the Recent *Thrissa baelama* (FORSSKAL, 1775) and SMALE et al. (1995: pl. 7, fig. D1-D3 and E1-E3) for respectively *T. setirostris* (BROUSSONET, 1782) and *T. vitrirostris* (GILCHRIST & THOMPSON, 1908).

***Arius nucleus* (WEINFURTER, 1952) (2)**

Otoliths of this species, originally mentioned as generic incertae sedis, are utricular otoliths of an ariid.

“genus Atherinidarum” *bavayi* STEURBAUT, 1984 (3)

This otolith, originally referred to *Dentex* aff. *latior* SCHUBERT by WEINFURTER (1952c) represents the first Paratethys record of “genus Atherinidarum” *bavayi*, an atherinid known from the Aquitaine Basin.

***Tylosurus wetzeldorfensis* (WEINFURTER, 1952) (4)**

WEINFURTER’s holotype, originally, described as a generic incertae sedis in WEINFURTER (1952d), shows most affinities with the belonid genus *Tylosurus*, see NOLF & GIRONE (2006: pl. 5, figs 1-4) for otoliths of the Recent *Tylosurus acus* (LACEPÈDE, 1803). The fossil species has higher otoliths and a more blunt anterior portion than the Recent form.

***Gazza hilberi* (WEINFURTER, 1952) (5)**

This otolith, referred to a triglid by WEINFURTER (1952d), shows most affinities with those of the Recent leiognathid genus *Gazza*, see RIVATON & BOURRET (1999: pl. 35, figs. 8-9) and SMALE et al. (1995: pl. 75, figs. 1-3) for otoliths of the Recent *Gazza minuta* (BLOCH, 1797).

Sciaenidae indet. (6)

Under the name *Ot. (Sciaenidarum) styriacus*, WEINFURTER (1952d) published a strongly eroded sciaenid otolith, non-diagnostic at species or genus level. This specimen certainly does not belong to the American sciaenid genus *Pogonias*, as guessed by SCHWARZHANS (1993: 86) from WEINFURTER's drawing only. This specimen is refigured here (pl. 6, fig. 2), together with another non-diagnostic sciaenid otolith from the Badenian of the Vienna Basin, the holotype of *Ot. (Sciaenidarum) kittli* SCHUBERT, 1902. Those two otoliths show some similarity and may be related to each other, but neither is sufficiently well preserved to define a species or even a genus.

"genus Gobiidarum" sp. 2 (7)

The otolith figured here apparently belongs to the same species as the three otoliths figured by BRZOBOHATÝ et al. (2007: pl. 8, figs 9-11) under the name *Priolepis* sp. This identification was based on the comparison with a picture of a *Priolepis cincta* (REGAN, 1908) otolith published by SMALE et al. (1995: pl. 129, fig. H 1). For a more detailed look at Recent *Priolepis* otoliths, see, e.g., *P. hipoliti* (METZELAAR, 1922) (see pl. 7, fig. 4), which shows that our fossils are too different from *Priolepis* to be assigned to the same genus. Moreover, fishes of the genus *Priolepis* usually do not reach such a large size as those that provided our fossil otoliths. As already stated by BRZOBOHATÝ et al. (2007: 173), otoliths described as *Gobius* sp. 6 from the Badenian at Gainfarn (BRZOBOHATÝ, 1994: pl. 6, fig. 17, not figs 18-20) and *Gobius praetiosus* PROCHAZKA, 1893 from the Lower Badenian of the Transylvanian Basin (WEILER, 1950: pl. 4, figs 27, ? 25; not PROCHAZKA, 1893) are conspecific with the treated taxon.

***Syacium syacioides* (WEINFURTER, 1952) (8)**

As already stated by SCHWARZHANS (1999), the otoliths described as *Ot. (Pleuronectidarum) syacioides* by WEINFURTER (1952d: 480) belong to the genus *Syacium*. They are here compared to those of the Recent *Syacium papillosum* (LINNAEUS, 1758) (pl. 8, fig. 1-2).

***Arnoglossus taureri* (WEINFURTER, 1952) (9)**

The otoliths described as *Solea taureri* by WEINFURTER (1952c: 166) and *Arnoglossus hollerii* by WEINFURTER (1952d: 479) are all conspecific, and the generic name *Arnoglossus* is the correct one, see CHAINE (1936: pl. 3) for comparative material of the Recent species *Arnoglossus thori* KYLE, 1913 and *A. laterna* (WALBAUM, 1792).

Description of new species

Order Perciformes BLEEKER, 1859

Family Lethrinidae REGAN, 1913

Lethrinus styriacus nov. spec. (pl. 7, figs 7-9)

Type material: Holotype: a right otolith (pl. 7, fig. 7); from Gross Sankt Florian, Mühlbauer (Landesmuseum Joanneum in Graz 62 082/41), one paratype from Wetzelsdorf, Winkeltoni (LMJG, nr. 55 863) (pl. 7, fig. 8), and one paratype from Gross Sankt Florian, Mühlbauer (LMJG 62 082/41) (pl. 7, fig. 9).

Dimensions of the holotype: Length: 4.8 mm, height: 3.0 mm, thickness: 0.9 mm.

Type locality: Gross Sankt Florian, Mühlbauer, St. Florian Beds.

Typeymology: *Styriacus*, a, um (Latin) = from Styria, alludes to the type area where the species occurs.

Diagnosis: This species is characterised by globally elliptical, biaculeate otoliths with crenulated margins. The outer face is concave in all directions, with the strongest concavity in the antero-posterior direction. The inner face has a very smooth surface and is strongly convex in all directions. The sulcus is constituted by a rather wide ostium, especially in its dorsal part, and a narrower cauda that is about twice as long as the ostium and has a general antero-dorsal/postero-ventral orientation. There is a marked depression in the dorsal area, just above the caudal crista superior and accentuating this crest. The holotype shows no ventral furrow, but in the two paratypes some traces of a ventral furrow are visible. Although not perfectly preserved in any of the available specimens, there is a well marked rostrum, a clear excisura, and a small but rather strong antirostrum.

Remarks: Lethrinids are reef-associated fishes represented by many species in the present-day Indo-West Pacific realm. An abundant iconography of the otoliths of Recent lethrinids can be found in NOLF (1993: p. 232, fig. 7), SMALE et al. (1995: pls 92-93) and RIVATON & BOURRET (1999: pls 46-52). Otoliths of our fossil species differ from most of them by their rather regular elliptical outline with acculate ends.

Conclusions

Systematics

Fifty-seven taxa based on otoliths of bony fishes were identified in the Lower Badenian sediments of the Western Styrian Basin and the Lavanttal Basin. Among them, the genus *Xyrichtys* is documented for the first time from deposits older than the Pliocene. The most diversified associations occur in the St. Florian Beds; the Mühlendorf Formation pro-

vided 18 taxa. Five of those were only recorded from the last named formation. “Genus *Gobiidarum*” *noricus* is reported only from the non-marine younger sediments in the Lavanttal Basin (tab. 2). Skeleton-based taxa from the Eastern Styrian Basin (SCHULTZ 2000, 2004, 2006; SCHULTZ & BELLWOOD 2004) were not documented by otoliths from similar taxa in the Western Styrian Basin or the Lavanttal Basin.

Palaeoenvironment

Based on an actualistic analysis, the otolith associations are indicative of exclusively marine shallow-water fish faunas with dominant demersal and benthopelagic taxa (33 taxa). We did not attach much importance to the quantitative composition of each of the associations because little is known of how these old museum collections were collected (and about eventual lost material). Gobiid otoliths nevertheless seem to constitute a dominant portion in each of the associations that are documented by significant numbers of specimens, e. g., for the Wetzelsdorf sites, gobiid otoliths are represented by 91 % at Ringraben, by 81 % at Winkeltoni, 56 % at Kreuzsholler, 29 % at Thomihiaslgraben and 18 % at Wenzelsteffi.

In the marine upper part of the Mühldorf Formation, this high portion of bottom fishes contrasts with the interpretations of prevailing suboxic condition in the bottom waters during the deposition (REISCHENBACHER et al. 2007). Reef-associated groups are represented by a relatively high number of taxa (*Apogon imberbis*, *Lethrinus*, *Pristigenys*, and approximately *Tylosurus*, *Hemiramphus*, *Serranus*, *Syacium*, *Xyrichtyhs*). This is in agreement with the presence of small reefs and coral carpets along the Styrian Basin margins in the Early Badenian. Fishes living in somewhat deeper water are individually very scarce (*Phycis blennioides*, *Trigla lyra*, *Hoplobrotula*), and bathypelagic fishes are completely absent. Pelagic (*Bregmaceros*, *Thrissa*, *Tylosurus*) and a poorly diversified amount of mesopelagic fishes (Myctophidae) are rare. This indicates an environment that was only poorly exposed to the deeper waters of the open sea. The myctophids however, are more diversified (3 genera) than the ones of shallow-water assemblages of Middle and Upper Badenian age in the Vienna Basin (e.g. Kienberg – BRZOBOHATÝ et al. 2007; Devínská Nová Ves, Rohožník – HOLEC 1973, 1975). This can be interpreted as resulting from a deep connection of the Central Paratethys with the Mediterranean during the Early Badenian (e.g. BRZOBOHATÝ 1995).

Climate

Climatic requirements of fishes of the St. Florian Beds show clearly tropic-subtropic climate (3 genera, *Thrissa*, *Lethrimus*, *Syacium*, are tropic and others tropic or subtropic). The mesopelagic myctophids exhibit a wider climatic range. This is also supported by data on molluscs (HARZHAUSER et al. 2003), echinoderms (KROH 2007), and sharks (HIDEN 1995) in the Early Badenian of the area under study. The extant *Phycis blennioides* lives mainly in warm temperate waters, including the Mediterranean. Other gadoids requiring cooler waters were not found; this feature was also noted by WEINFURTER (1952d). In the southern basins of the Central Paratethys, the first occurrence of diversified gadoids begins stratigraphically higher – in the Upper Lagenidae Zone (BRZOBOHATÝ 1978; BRZOBOHATÝ et al. 2007). The presence of tropic fishes proves that the studied localities in the St. Florian Beds and the Mühldorf Formation stratigraphi-

Families	Taxa	Iconography
Heterenchelyidae	<i>Panturichthys subglaber</i> (SCHUBERT, 1906)	BNK, 2007, pl. 1, figs 6-8
Congridae	<i>Ariosoma balearicum</i> (DELAROCHE, 1809)	pl. 1, figs 1-3
	<i>Rhynchoconger pantanellii</i> (BASSOLI & SCHUBERT, 1906)	pl. 1, fig. 4
Clupeidae	Clupeidae	-
Engraulidae	<i>Thrissa floriana</i> (WEINFURTER, 1952) (1)	* pl. 1, figs 6-7
Ariidae	<i>Arius nucleus</i> (WEINFURTER, 1952) (2)	* pl. 1, fig. 5
Synodontidae	<i>Saurida germanica</i> (WEILER, 1942)	BNK, 2007, pl. 1, figs 12-13
Myctophidae	<i>Hygophum murbani</i> (WEINFURTER, 1952)	pl. 2, figs 1-2
	<i>Diaphus regani</i> TAANING, 1932	pl. 2, figs 7-9
	<i>Diaphus</i> sp.	-
Bregmacerotidae	<i>Notoscopelus cf. mediterraneus</i> (KOKEN, 1891)	-
Carapidae	<i>Bregmaceros albyi</i> (SAUVAGE, 1880)	pl. 2, figs 5-6
Ophidiidae	<i>Carapus aff. acus</i> (BRÜNNICH, 1768)	pl. 2, figs 3-4
Phycidae	<i>Hoplobrotula acutangula</i> (KOKEN, 1884)	pl. 3, fig. 2
Atherinidae	<i>Phycis blennioides</i> (BRÜNNICH, 1768)	pl. 3, fig. 1
Belonidae	"genus Atherinidarum" bavayi STEURBAUT, 1984 (3)	* pl. 3, fig. 5
Hemiramphidae	<i>Tylosurus wetzelsdorffensis</i> (WEINFURTER, 1952) (4)	pl. 3, fig. 4
Triglidae	<i>Hemiramphus miocenicus</i> (WEINFURTER, 1952)	pl. 3, fig. 7
Ambassidae	<i>Trigla</i> aff. <i>lyra</i> LINNAEUS, 1758	pl. 4, fig. 10
Serranidae	<i>Ambassis</i> sp.	* pl. 3, fig. 6
Priacanthidae	<i>Serranus integer</i> (SCHUBERT, 1906)	pl. 4, figs 6-7
Apogonidae	<i>Pristigenys rhombicus</i> (SCHUBERT, 1906)	BNK, 2007, pl. 4, fig. 13
Sillaginidae	<i>Apogon aff. imberbis</i> LINNAEUS, 1758	pl. 3, fig. 3
Carangidae	<i>Sillago schwarzhansi</i> STEURBAUT, 1984	pl. 4, figs 8-9
Leiognathidae	<i>Trachurus</i> aff. <i>picturatus</i> (BOWDICH, 1825)	BNK, 2007, pl. 7, figs 13-15
Gerreidae	<i>Gazza hilberi</i> (WEINFURTER, 1952) (5)	* pl. 5, fig. 1
Haemulidae	<i>Gerres</i> sp. 1	* pl. 5, fig. 5
	<i>Gerres</i> sp. 2	* pl. 5, figs 2-3
	<i>Brachydeuterus speronatus</i> (BASSOLI, 1906)	BNK, 2007, pl. 5, figs 1-10
Lethrinidae	<i>Pomadasys</i> aff. <i>incisus</i> (BOWDICH, 1825)	BNK, 2007, pl. 5, fig. 11
Sparidae	"genus Haemulidarum" <i>lucidus</i> (BASSOLI, 1906)	* pl. 5, fig. 6
	<i>Lethrinus styriacus</i> n. sp.	* pl. 7, figs 7-9
	<i>Boops boops</i> (LINNAEUS, 1758)	pl. 5, fig. 8
Sciaenidae	<i>Dentex</i> aff. <i>maroccanus</i> VALENCIENNES, 1830	BNK, 2007, pl. 5, figs 12-14
	<i>Diplodus</i> cf. <i>puntazzo</i> (CETTI, 1777)	BNK, 2007, pl. 6, fig. 1
	<i>Lithognathus mormyrus</i> (LINNAEUS, 1758)	pl. 5, fig. 7
	<i>Spondylisoma cantharus</i> (LINNAEUS, 1758)	BNK, 2007, pl. 6, figs 10-12
	<i>Pagellus erythrinus</i> (LINNAEUS, 1758)	BNK, 2007, pl. 6, figs 8-9
	<i>Sciaena</i> sp.	pl. 6, fig. 4
Polynemidae	[<i>Sciaenidae</i> ind.] (6)	pl. 6, figs 1-2
Cepolidae	"g. Polynemidarum" <i>huyghebaertae</i> STEURBAUT, 1984	pl. 5, fig. 4
Labridae	<i>Cepola rubescens</i> LINNAEUS, 1766	pl. 6, fig. 3
Trachinidae	<i>Xyrichtys</i> sp.	* pl. 6, fig. 6
Gobiidae	<i>Trachinus</i> <i>vipera</i> CUVIER, 1829	pl. 6, fig. 5
	<i>Deltentosteus telleri</i> (SCHUBERT, 1906)	BNK, 2007, pl. 8, figs 2-5
	<i>Gobius</i> aff. <i>multipinnatus</i> (VON MEYER, 1852)	BNK, 2007, pl. 8, figs 6-8
	<i>Lesueurigobius</i> aff. <i>vicinalis</i> (KOKEN, 1891)	BNK, 2007, pl. 8, figs 12-14
	? <i>Oxyurichtys</i> sp.	pl. 7, figs 1-3
	"genus Gobiidarum" noricus WEINFURTER, 1952	L pl. 8, fig. 12
	"genus Gobiidarum" sp. 1	pl. 7, fig. 5-6
	"genus Gobiidarum" sp. 2 (7)	pl. 7, fig. 10
	[<i>Gobiidae</i> ind.]	-
Citharidae	<i>Citharus linguatula</i> (LINNAEUS, 1758)	pl. 8, fig. 3
Paralichthyidae	<i>Syacium syacioides</i> (WEINFURTER, 1952) (8)	pl. 8, fig. 4
Bothidae	<i>Arnoglossus taureri</i> (WEINFURTER, 1952) (9)	pl. 8, figs 8-10
Soleidae	<i>Buglossidium frequens</i> STEURBAUT, 1984	pl. 8, figs 5-6
	<i>Dicologlossa hexophthalma</i> (BENNETT, 1831)	pl. 8, fig. 11
	<i>Cynoglossus leuchsii</i> WEINFURTER, 1952	pl. 8, fig. 7

cally correspond to a lower part of the Weissenegg Formation in the Eastern Styrian Basin. In the upper part of this formation, a deposition of red algal and corallinaceous limestones terminated in connection with a cooling, as indicated by isotopic studies (BOJAR et al. 2004).

Palaeogeography

Taxa inhabiting the present-day Atlantic or the Eastern Atlantic–Mediterranean region dominate, but the number of taxa with the present-day Indo-Pacific distribution is also relatively high (*Thrissa*, *Diaphus regani*, *Hoplobrotula*, *Ambassis*, *Gazza*, *Sillago*, *Lethrinus*, *Oxyurichthys*). However, five of those taxa are already known in the European region before the beginning of the Middle Miocene and can be considered as descendants from earlier established residents.

The non neglectable Indo-Pacific component, the tropical to subtropical affinities of the association, and the presence of nine taxa (*Thrissa floriana*, *Arius nucleus*, “g. Atherinidarum” *bavayi*, *Ambassis* sp., *Gazza hilberi*, *Gerres* sp. 2, “genus Haemulidarum” *lucidus*, *Lethrinus styriacus*, and *Xyrichtys* sp.) representing the Badenian record from the Styrian Basin only (tab. 1), support a specific nature of the West Styrian otolith association among known Central Paratethys ones. However, a comparison with contemporaneous faunas from different areas is not evident. Otolith associations from the Transylvanian Basin and the Carpathian Foredeep in Moravia represent bathymetrical deeper fish assemblages. In the Vienna Basin, only a small fauna is reported from the Lower Lagenidae Zone (Niederleis, SCHUBERT 1906). More than 20 species (43 %) of the Styrian taxa occur contemporaneously in the Korytnica Clay (Carpathian Foredeep, southern Poland, RADWAŃSKA 1992). In contrast to the assemblages under study, the Korytnica otoliths have a markedly higher number of non-termophylllic gadoids, which agrees with a marginal northern position of the Korytnica Bay in the Central Paratethys (HARZHAUSER et al. 2003; BRZOBOHATÝ et al. 2007; KROH 2007; HARZHAUSER & PILLER 2007).

Outside the Paratethys, in the Mediterranean, contemporaneous neritic otolith associations are scarce. In fact, they are only well documented from Burdigalian/Langhian boundary sediments in the Vallés-Penedés area, Catalunya, Spain, (HOEDEMAKERS & BATTLORI 2005) where quite similar faunas have been found. Thirteen (28 %) of the nominal species from the Styrian Basin also occur in this last region.

Acknowledgments

This study is entirely based on existing collections conserved in the collections of the Landesmuseum Joanneum in Graz and the Landesmuseum Kärnten in Klagenfurt. We are very obliged to the respective curators, Martin GROSS and Cornelia BOCKRATH for their kind assistance. We visited the Graz museum on

◀ Table 1. List of teleost species represented by otoliths in the Badenian of the Styrian and Lavanttal basins. Numbers in brackets behind taxon names indicate problematic taxa discussed in the text. Bold taxa are refer to taxa recorded from Mühldorf only, taxa occurring in the Lavanttal only are marked by a bold L. Asterisks mark taxa that are recorded from the Badenian of the Styrian Basin only. BNK, 2007 = BRZOBOHATÝ, NOLF & KROUPA, 2007.

WEINFURTER (1952b, 1952c, 1952d) (1)

Apogon imberboides WEINFURTER, 1952d
Arnoglossus hollerii WEINFURTER, 1952d
? *Box insignis* (PROCHAZKA) (in WEINFURTER, 1952 d)
Chrysophris doderleini SCHUBERT & BASSOLI (in W., 1952 d)
Centropricus integer SCHUBERT (in WEINFURTER, 1952 d)
Cepola praerubescens BASSOLI & SCHUBERT (in W., 1952c, 1952d)
Chirodorus miocenicus WEINFURTER, 1952d
Clupea aff. testis (KOKEN) (in WEINFURTER, 1952c)
Clupea sp. (in WEINFURTER, 1952d)
Congermuraena moravica (SULC, 1932) (in WEINFURTER, 1952c)
? *Cottus germanicus* WEILER (in WEINFURTER, 1952c)
Cynoglossus leuchsi WEINFURTER, 1952d
Dentex latior SCHUBERT (in WEINFURTER, 1952d)
Dentex aff. latior SCHUBERT (in WEINFURTER, 1952c)
Dentex nobilis KOKEN (in WEINFURTER, 1952d)
Eucitharus rheanensis (KOKEN) (in WEINFURTER, 1952c)
Fierasfer nuntius KOKEN (in WEINFURTER, 1952c, 1952d)
Gobius noricus WEINFURTER, 1952 b
Gobius carinifrons WEINFURTER, 1952 b
Gobius aff. pretiosus PROCHAZKA (in WEINFURTER, 1952b, 1952d)
Gobius telleri SCHUBERT (in WEINFURTER, 1952c, 1952d)
Gobius pretiosus PROCHAZKA (in WEINFURTER, 1952c)
Gobius vicinalis KOKEN (in WEINFURTER, 1952c, 1952d)
Mugil sp. (in WEINFURTER, 1952d)
Ot. (*Congridarum*) *pantanellii* (BASSOLI & SCHUBERT) (in W., 1952c)
Ot. (inc. sed.) *hassovicus* KOKEN (in WEINFURTER, 1952 d)
Ot. (inc. sed.) *nucleus* WEINFURTER, 1952d
Ot. (inc. sed.) *wetzelesdorffensis* WEINFURTER, 1952d
Ot. (*Ophidiidarum*) *joachimicus* (KOKEN) (in WEINFURTER, 1952c)
Ot. ?(*Pagellus*) *gregarius* (KOKEN) (in WEINFURTER, 1952c)
Ot. (*Percidarum*) *floriani* WEINFURTER, 1952d
Ot. (*Pleuronectidarum*) *syacioides* WEINFURTER, 1952d
Ot. (*Sciaenidarum*) *styriacus* (WEINFURTER, 1952d)
Ot. (*Smaris*) *elegans* (PROCHAZKA) (in WEINFURTER, 1952 d)
Ot. (*Spardarum*) *gregarius* KOKEN (in WEINFURTER, 1952 d)
 (errenous reference to iconography in WEINFURTER, 1952 d)
Ot. (*Spardarum*) *giganteus* WEINFURTER, 1952 d
Ot. (?*Xenodermichthys*) *catus* SCHUBERT (in W., 1952c, 1952d)
Phycis simplex (KOKEN) (in WEINFURTER, 1952d)
Pristipoma arcuata (BASSOLI & SCHUBERT) (in WEINFURTER, 1952d)
Scopelus debilis austriacus (KOKEN) (in WEINFURTER, 1952c, 1952d)
Scopelus mediterraneus (KOKEN) (in WEINFURTER, 1952d)
Scopelus pulcher (PROCHAZKA) (in WEINFURTER, 1952c, 1952d)
Scopelus tenuis murbani WEINFURTER, 1952d
Serranus noetlingi KOKEN (in WEINFURTER, 1952d)
Serranus sp. (in WEINFURTER, 1952c)
 (= htp. of *Serranus noetlingi muhldorfensis* WEINFURTER, 1949)
Solea kokeni BASSOLI & SCHUBERT (in WEINFURTER, 1952d)
Solea latior SCHUBERT (in WEINFURTER, 1952d)
Solea subglaber SCHUBERT (in WEINFURTER, 1952c, 1952d)
Solea subvulgaris SCHUBERT (in WEINFURTER, 1952d)
Solea taureri WEINFURTER, 1952c
Sphyraena cf. hansfuchi SCHUBERT (in WEINFURTER, 1952d)
Trachinus biscutis KOKEN (in WEINFURTER, 1952d)
Trigla asperoides SCHUBERT (in WEINFURTER, 1952c)
Trigla cf. asperoides SCHUBERT (in WEINFURTER, 1952d)
Trigla (?) hilberti WEINFURTER, 1952d
Trigla rhombica Schubert (in WEINFURTER, 1952d)
Umbrina gibberula (KOKEN) (in WEINFURTER, 1952d)

(1) evaluation of the species described previously to WEINFURTER (1952 b, c and d) concern the Styrian material cited by WEINFURTER only, not necessary the primary type material of the concerned species. In the first column, spelling is the original one used by WEINFURTER.

This paper

Apogon aff. imberbis LINNAEUS, 1758
Arnoglossus taureri (WEINFURTER, 1952)
indeterminata (Percoidei)
indeterminata (Percoidei)
Serranus integer (SCHUBERT, 1906)
Cepola rubescens LINNAEUS, 1766
Hemiramphus miocenicus (WEINFURTER, 1952)
Clupeidae indet.
Clupeidae indet.
Ariosoma balearicum (DELAROCHE, 1809)
Saurida germanica (WEILER, 1942)
Cynoglossus leuchsi WEINFURTER, 1952
Brachydeuterus speronatus (BASSOLI, 1906)
"genus *Atherinidarum*" *bavayi* STEURBAUT, 1984
indeterminata (Percoidei)
Citharus linguatula (LINNAEUS, 1758)
Carapus acus (BRÜNNICH, 1768)
"genus *Gobiadarum*" *noricus* (WEINF., 1952)
Gobiidae indet.
Gobiidae indet.
Deltentosteus telleri (SCHUBERT, 1906)
Gobiidae indet.
Lesueurigobius aff. *vicinalis* (KOKEN, 1891)
indeterminata (Perciformes)
Rhynchoconger pantanellii (B. & SCH., 1891)
Sillago schwarzianus STEURBAUT, 1984
Arius nucleus (WEINFURTER, 1952)
Tylosurus wetzelsdorffensis (WEINFURTER, 1952)
Hoplobrotula acutangula (KOKEN, 1884)
Dentex aff. maroccanus VALENCIENNES, 1830
Thrissa floriana (WEINFURTER, 1952)
Syacium syacioides (WEINFURTER, 1952)
Sciaenidae indet.
indeterminata (Percoidei)
Dentex aff. maroccanus VALENCIENNES, 1830
indeterminata (Percoidei)
Bregmaceros albyi (SAUVAGE, 1880)
Phycis blennioides (BRÜNNICH, 1768)
Pomadasys aff. *incisus* (BOWDICH, 1825)
Diaphus sp. indet.
Notoscopelus cf. *mediterraneus* (KOKEN, 1891)
Diaphus sp. indet.
Hygophum murbani (WEINFURTER, 1952)
indeterminata (Percoidei)
Lithognathus mormyrus (LINNAEUS, 1758)
Arnoglossus taureri (WEINFURTER, 1952)
indeterminata (Soleidae)
Panturichthys subglaber (SCHUBERT, 1906)
indeterminata (Soleidae)
Arnoglossus taureri (WEINFURTER, 1952)
Saurida germanica (WEILER, 1942)
Trachinus viperus CUVIER, 1829
Trigla aff. lyra LINNAEUS, 1758
indeterminata (Triglidae)
Gazza hilberti (WEINFURTER, 1952)
Pristigenys rhombicus (SCHUBERT, 1906)
Sciaena sp.

three different occasions, and enjoyed the generous hospitality of the town of Graz at their Schloss Eggenberg guest room. Irene ZORN assisted us in examination of the otolith collections (mainly type material of the otoliths described by SCHUBERT) in the Geologische Bundesanstalt Wien, and Bettina REICHENBACHER (Universität München) showed us some WEINFURTER type specimens that she had on loan from the the Krahuletz Museum, Eggendorf. The material described by WEINFURTER (1952c) and conserved in the Klagenfurt Museum was loaned in totality to us, which greatly facilitated our investigations. It is also a pleasure for us to express our sincere gratitude to Anne WOUTERS and Jean Pierre DE BLAUWE for help in the preparation of the iconography and the tables. The research of R. BRZOBHATÝ was supported by the MSM Project 0021622412 (Czech Republic).

References

- For all references concerning only author names of Recent species, we refer to ESCHMEYER (1998).
- BECHTEL, A., REISCHENBACHER, D., SACHSENHOFER, R.S., GRATZER, R., LÜCKE, A. & PÜTTMANN, W. (2007): Relations of petrographical and geochemical parameters in the middle Miocene Lavanttal lignite (Austria). – International Journal of Coal Geology, **70**/4: 325-349.
- BOJAR, A.-V., HIDEN, H., FENNINGER, A. & NEUBAUER, F. (2004): Middle Miocene seasonal temperature changes in the Styrian Basin, Austria, as recorded by the isotopic composition of pectinid and brachiopod shells. – Palaeogeography, Palaeoclimatology, Palaeoecology, **203**: 95-105.
- BRZOBHATÝ, R. (1978): Die Fisch-Otolithen aus dem Badenien von Baden-Sooß, NÖ. – Annalen des Naturhistorischen Museums in Wien, **81**: 163-171.
- (1994): Die Fischotolithen des Badenien von Gainfarn, Niederösterreich (Mittelmiozän, Wiener Becken). – Annalen des Naturhistorischen Museums in Wien, Serie A, **96**: 67-93.
- (1995): Macrouridae (Teleostei, Otolithen) im Oligozän und Miozän der Zentralen Paratethys und ihre paläogeographische Bedeutung. – Jahrbuch der Geologischen Bundesanstalt, **138**/4: 615-637.
- , NOLF, D. & KROUPA, O. (2007): Fisch Otoliths from the Middle Miocene of Kienberg at Mikulov, Czech Republic, Vienna Basin: Their paleoenvironmental and paleogeographic significance. – Bulletin de l’Institut royal des sciences naturelles de Belgique, Sciences de la Terre, **77**: 167-196.
- & SCHULTZ, O. (1978): Die Fischfauna des Badenien. – In: PAPP, A., CICHA, I., SENEŠ, J. & STEININGER, F.: M4, Badenien. – Chronostratigraphie und Neostratotypen, **6**: 441-465. Veda, Bratislava.
- CHAINE, J. (1936): Recherches sur les otolithes de poissons. Étude descriptive et comparative de la sagitta des Téléosteens. – Actes de la Société Linnéenne de Bordeaux, **89**: 1-252.
- EBNER, F. & SACHSENHOFER, R.F. (1995): Palaeogeography, subsidence and thermal history of the Neogene Styrian Basin (Pannonian basin system, Austria). – Tectonophysics, **242**: 133-150.
- ESCHMEYER, W.N. (1998): Catalog of Fishes. – Special Publication of the Center for Biodiversity Research and Information, California Academy of Sciences, **1**: 1- 2950.

► Table 2: Alphabetic list of the taxa cited by WEINFURTER (1952b, c, d) from the Styrian and Lavanttal basins.

- GROSS, M., FRITZ, I., PILLER, W.E., SOLIMAN, A., HARZHAUSER, M., HUBMANN, B., MOSER, B., SCHOLGER, R., SUTTNER, T.J. & BOJAR, H.P. (2007): The Neogene of the Styrian Basin – Guide to excursions. – *Joannea, Geologie und Paläontologie*, **9**: 117-193.
- HANDLER, R., EBNER, F., NEUBAUER, F., BOJAR, A.-V. & HERMAN, S. (2006): $^{40}\text{Ar}/^{39}\text{Ar}$ dating of Miocene tuffs from the Styrian part of the Pannonian Basin: an attempt to refine the basin stratigraphy. – *Geologica Carpathica*, **57/6**: 483-494.
- HARZHAUSER, M., MANDIĆ, O. & ZUSCHIN, M. (2003): Changes in Paratethyan marine molluscs at the Early/Middle Miocene transition: diversity, palaeogeography and palaeoclimate. – *Acta Geologica Polonica*, **53/4**: 323-339.
- & PILLER, W.E. (2007): Benchmark data of a changing sea – Palaeogeography, Palaeobiogeography and events in the Central Paratethys during the Miocene. – In: BRUCH, A.A., UHL, D. & MOSBRUGGER, V. (eds): Miocene Climate in Europe – Patterns and Evolution. First Synthesis of Neclime. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, **253**: 8-31.
- HIDEN, H. (1995): Elasmobranchier (Pisces, Chondrichthyes) aus dem Badenium (Mittleres Miozän) des Steirischen Beckens (Österreich). – *Mitteilungen der Abteilung für Geologie und Paläontologie, Landesmuseum Joanneum*, **52/53**: 41-110.
- & STINGL, K. (1998): Neue Ergebnisse zur Stratigraphie und Paläogeographie der Eibiswalder Schichten (Miozän, Weststeirisches Becken, Österreich): Die Otolithenfauna der Tongrube Gasseldorf. – *Geologisch-Paläontologische Mitteilungen Innsbruck*, **23**: 77-75.
- HOEDEMAKERS, K. & BATLLORI, J. (2005): Fish otoliths from the Early and Middle Miocene of the Penedès (Catalunya, Spain). – *Batalleria*, **12/2004-5**: 105-134.
- HOLEC, P. (1973): Fisch-Otolithen aus dem oberen Baden (Miozän) des nordöstlichen Teiles des Wiener Beckens (Gebiet von Rohožník). – *Geologický Zborník, Geologica Carpathica*, **24/2**: 393-414.
- (1975): Fisch-Otolithen aus dem Baden (Miozän) des nördlichen Teiles des Wiener Beckens und der Donau-Tiefebene. – *Geologický Zborník, Geologica Carpathica*, **26/2**: 253-266.
- HOLLER, A. (1900): Über die Fauna der Meeressbildungen von Wetzelsdorf bei Preding in Steiermark. – *Mittheilungen des Naturwissenschaftlichen Vereins für Steiermark*, **1899**: 47-71.
- KOVÁČ, M., ANDREYeva-GRIGOROVICH, A., BAJRAKTAREVIĆ, Z., BRZOBOHATÝ, R., FILIPESCU, S., FODOR, L., HARZHAUSER, M., OSZCZYPKO, N., NAGYMAROSY, A., PAVELIĆ, D., RÖGL, F., SAFTIĆ, B., SLIVA, L. & STUDENCKA, B. (2007): Badenian evolution of the Central Paratethys Sea: Paleogeography, climate and eustatic sea level changes. – *Geologica Carpathica*, **58/6**: 579-606.
- KROH, A. (2007): Climate changes in the Early to Middle Miocene of the Central Paratethys and the origin of its echinoderm fauna. – In: BRUCH, A.A., UHL, D. & MOSBRUGGER, V. (eds): Miocene Climate in Europe – Patterns and Evolution. First Synthesis of Neclime. – *Palaeogeography, Palaeoclimatology, Palaeoecology*, **253**: 185-223.
- NOLF, D. (1985): Otolithi Piscium. In: SCHULTZE, H.P. (ed.). *Handbook of Paleoichthyology*, **10**. – pp. 1-145, Stuttgart et New York (Fischer).
- (1993): A survey of perciform otoliths and their interest for phylogenetic analysis, with an iconographical synopsis of the Percoidei. – *Bulletin of Marine Science*, **52/1**: 220-239.

- (1995): Studies on fossil otoliths. – The state of the art. – In: SECOR, D.H., DEAN, J.M. & CAMPANA, S. E. (eds): Recent developments in fish otolith research. – pp. 513-544. University of South Carolina press.
- & GIRONE, A. (2006): Otolithes de poissons du Pliocène inférieur (Zancléen) des environs d'Alba (Piemont) et de la côte ligure. – Rivista Piemontese di Storia Naturale, **27**: 77-114.
- PILLER, W.E., EGGER, H., ERHART, C.W., GROSS, M., HARZHAUSER, M., HUBMANN, B., VAN HUSEN, D., KRENMayR, H.-G., KRYSYn, L., LEIN, R., LUKENEDER, A., MANDL, G.W., RÖGL, F., ROETZEL, R., RUPP, C., SCHNABEL, W., SCHÖNLAUB, H.P., SUMMESBERGER, H., WAGREICH, M. & WESSELY, G. (2004): Die stratigraphische Tabelle von Österreich 2004 (sedimentäre Schichtenfolgen). – Kommission für paläontologische und stratigraphische Erforschung Österreichs der Österreichischen Akademie der Wissenschaften und Österreichischen Stratigraphischen Kommission, 1. Auflage.. Gerin, Wolkersdorf.
- POLESNY, H. (2003): The Karpatian in the Styrian Basin. – In: BRZOBHATÝ, R., CICHA, I., KOVÁČ, M. & RÖGL, F. (eds): The Karpatian – a Lower Miocene Stage of the Central Paratethys. – pp. 127-132, Brno (Masaryk University).
- RADWAŃSKA, U. (1992). Fish otoliths in the Middle Miocene (Badenian) deposits of southern Poland. – Acta Geologica Polonica, **42**/3-4: 141-328.
- REISCHENBACHER, D., RIFEJL, H., SACHSENHOFER, R.F., JELEN, B., ČORIĆ, S., GROSS, M. & REICHENBACHER, B. (2007): Early Badenian paleoenvironment in the Lavanttal Basin (Mühldorf Formation; Austria): Evidence from geochemistry and paleontology. – Austrian Journal of Earth Science, **100**: 202-229.
- RIVATON, J. & BOURRET, P. (1999): Les otolithes des poissons de l'Indo-Pacifique. – Documents scientifiques et techniques. Institut de recherche pour le développement, Nouméa, 2/2: 1-378.
- RÖGL, F., ČORIĆ, S., HOHENEGGER, J., PERVESLER, P., ROETZEL, R., SCHOLGER, R., SPEZAFFERI, S. & STINGEL, K. (2007): Cyclostratigraphy and Transgressions at the Early/Middle Miocene (Karpatian/Badenian) Boundary in the Austrian Neogene Basins (Central Paratethys). – Scripta Facultatis scientiarum naturalium Universitatis Masarykianae Brunensis, Geology, **36**: 7-13.
- RÖGL, F., SPEZZAFERRI, S. & ČORIĆ, S. (2002): Micropaleontology and biostratigraphy of the Karpatian-Badenian transition (Early-Middle Miocene boundary) in Austria (Central Paratethys). – Courier Forschungsanstalt Senckenberg, **237**: 47-67.
- SCHUBERT, R.J. (1906): Die Fischotolithen des österr.-ungar. Tertiärs. III. – Jahrbuch der Kaiserlich-Königlichen Geologischen Reichsanstalt, **56**: 623-706.
- SCHULTZ, O. (2000): Ein Zackenbarsch (*Epinephelus*, Serranidae, Pisces) aus dem Mittelmiozän von Retznei, Steiermark. – Joannea, Geologie und Paläontologie, **2**: 5-56.
- (2004): A Triggerfish (Osteichthyes: Balistidae: *Balistes*) from the Badenian (Middle Miocene) of the Vienna Basin and the Styrian Basin (Central Paratethys). – Annalen des Naturhistorischen Museum in Wien, Serie A, **106**: 345-369.
- (2006): *Oligodiodon*, ein Igelfisch aus dem Mittel-Miozän (Badenium) der Steiermark, Österreich (Diodontidae, Osteichthyes). – Joannea, Geologie und Paläontologie, **8**: 25-46.
- & BELLWOOD, D.R. (2004): *Trigonodon oweni* and *Asima jugleri* are different parts of the same species *Trigonodon jugleri*, a Chiseltooth Wrasse from the Lower and Middle

- Miocene in Central Europe (Osteichthyes, Labridae, Trigonodontinae). – *Annalen des Naturhistorischen Museum in Wien, Serie A*, **105**: 287-305.
- SCHWARZHANS, W. (1993): A comparative morphological treatise of recent and fossil otoliths of the family Sciaenidae (Perciformes). – In: PFEIL, H.F. (ed.): *Piscium catalogus, Part Otolithi Piscium 1*. – pp. 1-245, München (Verlag Dr. Friedrich Pfeil).
- (1999): A comparative morphological treatise of recent and fossil otoliths of the order Pleuronectiformes. – In: PFEIL, H.F. (ed.): *Piscium Catalogus, Part Otolithi Piscium 2*. – pp. 1-391, München (Verlag Dr. Friedrich Pfeil).
- SMALE, M. J., WATSON, G. & HECHT, T. (1995): Otolith atlas of South African marine fishes. – *Ichthyological monographs*, **1**: 1-253.
- SPEZZAFERRI, S., RÖGL, F., ČORIĆ, S. & HOHENEGGER, J. (2004): Paleoenvironmental changes and agglutinated foraminifera across the Karpatian/Badenian (Early/Middle Miocene) boundary in the Styrian Basin (Austria, Central Paratethys). – In: BUBÍK, M. & KAMINSKI, M.A. (eds): Proceedings of the six International Workshop on agglutinated foraminifera. – Grzybowski Foundation. Special Publication, **8**: 423-459.
- WEILER, W. (1950): Die Otolithen aus dem Jungtertiär Süd-Rumäniens. 2. Mittel-Miozän, Torton, Buglow und Sarmat. – *Senckenbergiana Lethaea*, **31**: 209-258.
- WEINFURTER, E. (1949): Eine neue Otolithenfauna aus dem Miozän von Mühldorf in Kärnten. – *Anzeiger der mathematisch-naturwissenschaftlichen Klasse der Österreichischen Akademie der Wissenschaften*, **1949/7**: 1-3.
- (1952b): Otolithen aus miozänen Brack- und Süßwasserschichten des Lavanttales in Kärnten. – *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, Mathematisch-naturwissenschaftliche Klasse I*, **161/2-3**: 141-148.
- (1952c): Die Otolithen aus dem Torton (Miozän) von Mühldorf in Kärnten. – *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse I*, **161/2-3**: 149-172.
- (1952d): Die Otolithen der Wetzelsdorfer Schichten und des Florianer Tegels (Miozän, Steiermark). – *Sitzungsberichte der Österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse I*, **161/7**: 455-498.

Explanation of the Plates

All figured fossil specimens are deposited in the collections of the Landesmuseum Joanneum, Graz (LMJG), the Kärnter Landesmuseum, Klagenfurt (LMKK), and the Krahuletz Museum, Eggenburg (KME). Some additional comparative specimens from the Institut royal des Sciences naturelles de Belgique (IRSNB) and from the Geologische Bundesanstalt Wien (GBW) are also figured. The Recent otoliths are part of the reference collection of Recent otoliths of the IRSNB. The latter collection is arranged in systematic order without numbering. Therefore, such specimens, when figured, bear only the notation "coll. IRSNB". The annotations F and R in the upper right corner of each compartment of the plates indicate if the figures in that compartment are fossils (F) or Recent (R). Recent species occurring as fossils are labeled with [(F)]. The annotations a, b and c in each compartment of the plates are used to indicate respectively ventral, inner (= mesial) and posterior views. Figures with only numbers and no letter show inner views. The scale bar in each compartment equals 1 mm. In the captions, L stands for left otolith and R for right otolith.

Plate 1

Figs 1-3. *Ariosoma balearicum* (DELAROCHE, 1809), L, Wetzeldorf, Winkeltoni (LMJG 2362, coll. 76970).

Fig. 4. *Rhynchoconger pantanellii* (BASSOLI & SCHUBERT, 1906), L, Lavanttal, Mühldorf (coll. LMKK).

Fig. 5. *Arius nucleus* (WEINFURTER, 1952), R, Gross Sankt Florian, Mühlbauer, Holotype (LMJG 62 082/8).

Figs 6-7. *Thrissa floriani* (WEINFURTER, 1952), Gross Sankt Florian, Mühlbauer, 6 = L, holotype (LMJG 62 082/10), 7 = R, paratype (LMJG 62 082/11).




Plate 2

Figs 1-2. *Hygophum murbani* (WEINFURTER, 1952), 1 = L, Brno – Královo Pole, Czech Republic (IRSNB P 6600), 2 = R, Wetzelsdorf, Thomihiaslgraben, Holotype (LMJG 62 082/14).

Figs 3-4. *Carapus aff. acus* (BRÜNNICH, 1768), R, Wetzelsdorf, Winkeltoni (LMJG no 2362, coll. 76 970).

Figs 5-6. *Bregmaceros albyi* (SAUVAGE, 1880), 5 = R, 6 = L, Lavanttal, Mühldorf (coll. LMKK).

Figs 7-9. *Diaphus regani* TAANING, 1932, L, Wetzelsdorf, Winkeltoni (LMJG 76971 or 55863).




Plate 3

- Fig. 1. *Phycis blennioides* (BRÜNNICH, 1768), R, Wetzelsdorf, Wenzelsteffi (LMJG 62082/28).
- Fig. 2. *Hoplobrotula acutangula* (KOKEN, 1884), R, Lavanttal, Mühldorf (coll. LMKK).
- Fig. 3. *Apogon aff. imberbis* LINNAEUS, 1758, L, Gross Sankt Florian, Mühlbauer, holotype of *Apogon imberboides* WEINFURTER, 1952 (LMJG inv. nr. 62 082/1).
- Fig. 4. *Tylosurus wetzelsdorffensis* (WEINFURTER, 1952), L, Wetzelsdorf, Wenzelsteffi, holotype (LMJG 62082/7).
- Fig. 5. "genus Atherinidarum" *bavayi* STEURBAUT, 1984, R, Lavanttal, Mühldorf (coll. LMKK).
- Fig. 6. *Ambassis* sp., L, Gross Sankt Florian, Mühlbauer, (LMJG 62082/39).
- Fig. 7. *Hemiramphus miocenicus* (WEINFURTER, 1952), R, Gross Sankt Florian, Kogelbauer, holotype (LMJG 62082/13).




Plate 4

- Figs 1-5. *Trigla lyra* LINNAEUS, 1758, L. Recent, 1-2 = eastern Atlantic, off Casablanca, Morocco, 3 = Mallorca, 4-5 = Mediterranean. Côte d'Azur, France (coll. IRSNB).
- Figs 6-7. *Serranus integer* (SCHUBERT, 1906). 6 = R, Wetzelsdorf, Winkeltoni (LMJG coll. nr. 55863), 7 = L, Steinabrunn, holotype (GBW 1906/01/13).
- Figs 8-9. *Sillago schwarzhansii* STEURBAUT, 1984, L, 8 = Wetzelsdorf, Thomihieslgraben (LMJG coll. nr. 55861), 9 = Wetzelsdorf, Winkeltoni (LMJG coll. nr. 55863).
- Fig. 10. *Trigla aff. lyra* LINNAEUS, 1758, L, Lavanttal, Mühldorf (coll. LMKK).




Plate 5

- Fig. 1. *Gazza hilberi* (WEINFURTER, 1952), L, Gross Sankt Florian, Mühlbauer, holotype (LMJG 62 082).
- Figs 2-3. *Gerres* sp. 2, L, Gross Sankt Florian, Mühlbauer (LMJG 62082).
- Fig. 4. "genus Polynemidarum" *huyghebaertae* STEURBAUT, 1984, L, Wetzelsdorf, Wenzelsteffi (LMJG nr 55865).
- Fig. 5. *Gerres* sp. 1, R, Wetzelsdorf, Wenzelsteffi (LMJG 55863).
- Fig. 6. "genus Haemulidarum" *lucidus* (BASSOLI, 1906), R, Wetzelsdorf, Wenzelsteffi (LMJG 62 082/29).
- Fig. 7. *Lithognathus mormyrus* (LINNAEUS, 1758), R, Wetzelsdorf, Kreuzschaller (LMJG 5331).
- Fig. 8. *Boops boops* (LINNAEUS, 1758), R, Wetzelsdorf, Winkeltoni (LMJG coll. nr. 55 863).




Plate 6

Figs 1-2. Sciaenidae ind., 1 = L, holotype of *Ot. (Sciaenidarum) kittli* SCHUBERT, 1902 (GBW 1860/1/50/2), Kienberg, Czech Republic. 2 = R, holotype of *Ot. (Sciaenidarum) styriacus* (WEINFURTER, 1952), Wetzelsdorf, Kreuzschaller (LMJG 62 082/4).

- Fig. 3. *Cepola rubescens* LINNAEUS, 1758, R, Lavanttal, Mühldorf (coll. LMKK).
Fig. 4. *Sciaena* sp., R, Wetzelsdorf, Wenzelsteffi (LMJG 62 082/19).
Fig. 5. *Trachinus viperus* CUVIER, 1829, R, Wetzelsdorf, Wenzelsteffi (LMJG 62 082/18).
Fig. 6. *Xyrichtys* sp., R, Wetzelsdorf, Winkeltoni (LMJG nr. 2362, coll. 76970).




Plate 7

Figs 1-3. ? *Oxyurichthys* sp., 1-2 = L, 3 = R, Wetzelsdorf, Winkeltoni (LMJG nr. 2362, coll. 76970).

Fig. 4. *Priolepis hipoliti* (METZELAAR, 1922), L, Recent, Caribbean, off Tobago (coll. IRSNB).

Figs 5-6. "genus Gobiidarum" sp. 1, L, Wetzelsdorf, Winkeltoni (coll. LMJG).

Figs 7-9. *Lethrinus styriacus* n. sp., R, 7 and 9 = Gross Sankt Florian, Mühlbauer, 7 = holotype (LMJG 62 082/41), 9 = paratype (LMJG 62 082/41), 8 = Wetzelsdorf, Winkeltoni, paratype (LMJG, nr. 55 863).

Fig. 10 "genus Gobiidarum" sp. 2, R, Wetzelsdorf, Thomihiaslgraben (LMJG, nr. 55 861).




Plate 8

- Figs. 1-2. *Syacium papillosum* (LINNAEUS, 1758), 1 = L, 2 = R, Recent, Atlantic, USA (coll. IRSNB).
- Fig. 3. *Citharus linguatula* (LINNAEUS, 1758), R, Lavanttal, Mühldorf (coll. LMKK).
- Fig. 4. *Syacium syacioides* (WEINFURTER, 1952), R, Gross Sankt Florian, Mühlbauer, holotype (LMJG 62082/2).
- Figs 5-6. *Buglossidium frequens* STEURBAUT, 1984, 5 = L, 6 = R, Wetzelsdorf, Winkeltoni (LMJG nr. 2362, coll. 76970).
- Figs 7. *Cynoglossus leuchsi* WEINFURTER, 1952, R, Wetzelsdorf, Winkeltoni, holotype (LMJG 62082/5).
- Figs 8-10. *Arnoglossus taureri* (WEINFURTER, 1952), R, 8 and 10 = Lavanttal, Mühldorf, 8 = holotype, 10 = paratype (coll. LMKK), 9 = Wetzelsdorf, Kreuzschaller, holotype of *Arnoglossus holleri* WEINFURTER, 1952 (LMJG 62 082).
- Fig. 11. *Dicologlossa hexophthalma* (BENNETT, 1831), R, Gross Sankt Florian, Mühlbauer, (LMJG 62 082/84).
- Fig. 12. “genus Gobiidarum” *noricus* WEINFURTER, 1952, R, Lavanttal, Weinzel bei Ober-Aigen, holotype (coll. KME).

