

WORLD CATALOGUE AND BIBLIOGRAPHY OF THE RECENT PYCNOGONIDA

(date of publication: 26 Oct. 1993)

compiled and distributed by

Hans-Georg Müller

Wissenschaftlicher Verlag,
Laboratory for Tropical Ecosystems
Research & Information Service
P.O. Box 2268
D-35532 Wetzlar
Germany

Sea spiders or pycnogonids are exclusively marine invertebrates of worldwide distribution, which rarely occur in large numbers. They are often cryptic and have usually very little economic significance. Most species are epibenthic ectoparasites of invertebrates, few are interstitial or bathypelagic. Revisions for major taxa of pycnogonids do not exist, except for STOCK's (1957) contribution on the Austrodecidae. Moreover, the most part of information on pycnogonids is scattered in many short papers. Descriptions, especially in the older literature, are often too poor to allow a reliable determination.

The purpose of the present work is therefore to partially address the lack of comprehensive works through publication of this overview, which is intended to have worldwide coverage and to include the valid taxa. Catalogues, checklists or reviews are a valuable tool for anyone needing an up-to-date reference to a currently accepted name and frequently to other pertinent information such as bibliographic and distributional data. This is so because most information is filed in publications under a species' scientific name, which then becomes the key to retrieval information from the literature. The taxonomic literature is an often confusing subject, because it is constantly changing to reflect current work, and the majority of species are known by several names. Thus a comprehensive listing of names, including synonyms, is an important starting point for locating information, whether as the basis for applied or basic research.

The information included in such an overview is usually arranged in a logical and organized format that allows for its convenient and rapid conveyance a quick and easy storage and retrieval system. The format and amount of information presented varies greatly. My use of the term catalogue herein is intended to convey a comprehensive treatment, including information on all valid names and most synonymies. Remarks on distribution, depth range and habitat preference of the included species are also given. The bibliographic section includes all references reporting on pycnogonids that came to my knowledge.

There are certain arguments that speak for publishing such a catalogue on computer disk, instead of distributing it regularly as a printed book. From the financial point of view, the former way is considerably cheaper and allows new editions in short intervals. If you prefer to have parts of this catalogue printed, you can do it yourself easily. The layout is prepared, genus and species names written in italics. Using the catalogue in a computer, you can locate certain species automatically with the search function of "Word Perfect 5.1". This allows convenient access to the respective species in the file PYCNO.CAT on the first floppy disk, even when only older names/synonyms are available to you.

Users should keep in mind, that we have a considerable steady increase of information within a short time. Thus, to avoid unnecessary inflation of the volume of this catalogue, I included only the more substantial information, which is (surely) subjective to some extend. For example, when a species is mentioned only in a key in a certain paper, this is often not noted in the reference-list under the respective species. When you are searching for keys to genera or species, you should check the information given for the respective families, subfamilies and genera. The inclusion of recent literature has been given priority, because these papers usually summarize information of older publications.

A complete bibliography of publications refering on Pycnogonida is available in the file PYCNO.LIT on the second floppy disc. I included all references that came to my knowledge.

Helpful suggestions for improvement of this catalogue are always welcome.

© 1993, H.-G. Müller, for personal use only. You are allowed to take one copy of this file for security purposes and to print yourself one copy of the files PYCNO.CAT and PYCNO.LIT. The price of 69 US-\$ for the floppy disk version payable in advance includes packing and registered airmail worldwide for individuals and institutions. With the acquisition of these files you have the possibility to obtain the update-versions for a reduced fee, which are compiled from time to time. You are informed free of charge by air mail, as soon as an update will be available. However, a printed version of this catalogue is available on request as paper-bound photocopy for 119 US-\$, or 139 US-\$ as cloth-bound volume with gold tooling. The prices of the printed versions include the floppy disks and registered surface-mailing. Add 25 US-\$, if you prefer registered air-mailing for the printed version. The contents of computer- and printed versions are identical, except for the lack of an index in the former. ISBN 3-930311-04-6, 26 Oct. 1993.

Helpful suggestions for improvement of this catalogue are always welcome.

How to use this catalogue in your Computer

To look for a certain species with the search function of "Word Perfect", you should consider the following hints:

1. Typewrite the complete name of the species you are looking for without the name of the author and date of the year. Ignore subgenera, subspecies and varieties. With this method you should be able to locate a certain species, even when only an older name is available to you. Unfortunately one is sometimes confronted with erroneous assignments of species. To avoid such problems I suggest to use for determinations only the most recent literature.

2. To locate certain countries use only the current name, i.e. Sri Lanka instead of Ceylon, Taiwan instead of Formosa. I found it to costly in this catalogue to summarize in each case the species known from a certain country, because many species are widely distributed in the Indo-Pacific and cannot be assigned to a certain country, particularly when they have been caught offshore.

SOME GENERAL INFORMATION ON PYCNOGONIDA

- 1949 HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 234-240 [zoogeographical remarks on Japanese species].
- 1963 CLARK, Rec. Austr. Mus., **26** (1): 1-5 [general remarks of Australian fauna, checklist of the up to then known species from Australia].
- 1963 HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1315-1324 [remarks on North American Arctic species].
- 1969 FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 1-139 [Antarctic fauna].
- 1974 KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 13 [key to British families].
- 1983 NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 4-5 [key to Japanese families].
- 1987 ARNAUD, Mesogee, **47**: 37-58 [review of the Mediterranean species].
- 1987 ARNAUD & BAMBER, Adv. Mar. Biol., **24**: 1-96 [the most up-to-date summary on the biology of pycnogonids].
- 1991 ARNAUD & BRANCH, S. Afr. J. Antarct. Res., **21** (1): 65-71 [illustrated key to species of Marion and Prince Edward Islands].
- 1933 MUNILLA, - J. mar. biol. Ass. U.K., **73**: 551 [checklist of species recorded from southern Spain].

AMMOTHEIDAE

- 1948 Ammotheidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 240-241 [diagnosis, remarks, key to American genera].
- 1954 Ammotheidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 132 [diagnosis, key to South African genera].
- 1969 Ammotheidae, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 66-67 [remarks, list of genera].
- 1974 Ammotheidae, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 23 [diagnosis, key to British species].
- 1983 Ammotheidae, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 5-6 [key to Japanese genera].
- 1986 Ammotheidae, - STOCK, Bull. mar. Sci., **38** (3): 406-407 [remarks, diagnostic

features].

- 1992 Ammoothidae, - CHILD, Mem. Hourglass Cruises, **9** (1): 7-9 [key of species from Gulf of Mexico], tab. 3 [species known or expected in the Gulf of Mexico].

Achelia HODGE 1864

- 1932 *Achelia*, - GORDON, Discovery Rep., **6**: 110 [remarks].
 1948 *Achelia*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 241-242 [diagnosis, remarks, key to NE-American species].
 1954 *Achelia*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 137-138 [diagnosis, remarks, key to South African species].
 1969 *Achelia*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 99-103 [remarks, variation, distribution, key to treated in the report], 103 [diagnosis].
 1969 *Aduncorostris*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 113 [diagnosis].
 1983 *Achelia*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 6 [key to Japanese species].
 1986 *Achelia*, - KRAPP, Bonn. Zool. Beitr., **37** (4): 302 [tab. 1, comparison with *Ammothea*].
 1986 *Achelia*, - STOCK, Bull. mar. Sci., **38** (3): 415.
 1992 *Achelia*, - CHILD, Mem. Hourglass Cruises, **9** (1): 11 [diagnosis, comparison with *Tanystylum*].

Achelia adelpha CHILD 1970

- 1970 *Achelia adelpha* CHILD, Proc. biol. Soc. Wash., **83** (27): 289-292, fig. 1.

Distribution: Society Islands (Tahiti, Huahine).

Depth range: 0-11m.

Habitat: Anchor chain; sandy shoal and wooden post.

Achelia alaskensis (COLE 1904)

- 1904 *Ammothea alaskensis* COLE, Harriman Alaska Exped., **10**: 266-268, pl. 12, fig. 4, pl. 17, figs. 4-12.
 1913 *Ammothea nudiuscula* HALL, Univ. Calif. Publ. Zool., **11** (6): 135-137, pl. 3, figs. 1-8.
 1929 *Ammothea (Achelia) alaskensis*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 151-156, figs. 42, 45.
 1933 *Ammothea alaskensis*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 60, fig. 10.
 1933 *Ammothea (Achelia) alaskensis*, - OHSHIMA, Bull. Biogeogr. Soc. Japan, **4**: 144-146, fig. 1.
 1934 *Ammothea alaskensis*, - SCHMITT, J. Wash. Acad. Sci., **24** (1): 70.
 1936 *Ammothea alaskensis*, - EXLINE, Proc. U.S. natn. Mus., **83** (2991): 421.
 1936 *Ammothea (Achelia) alaskensis*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 866.
 1939 *Ammothea nudiuscula*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 32.
 1940 *Ammothea alaskensis*, - OKUDA, J. Fac. Sci., Hokkaido Imp. Univ., (Zool. 6) **7** (2): 73-86, figs. 1-10.
 1941 *Ammothea nudiuscula*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26):

256 [key].

- 1942 *Ammothea nudiuscula*, - HILTON, J. Ent. Zool. Pomona College, **34** (4): 93.
 1949 *Achelia alaskensis*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 289.
 1954 *Achelia alaskensis*, - UTINOMI, Publ. Akkeshi Mar. Biol. Stat., **3**: 14, figs. 6-7.
 1955 *Ammothea alaskensis*, - LOSINA-LOSINSKY, in: Atlas of Invertebrates of the Far-Eastern Seas of the U.S.S.R., Izd. Akad. Nauk, S.S.S.R.: 160, pl. 42, fig. 6.
 1961 *Achelia alaskensis*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 91.
 1961 *Achelia gurjanovii* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 93-95, fig. 18.
 1964 *Achelia nudiuscula*, - HEDGPETH, in: R.I. Smith et al., eds., Intertidal Invertebrates of the Central California Coast, Berkeley; Univ. Calif. Press: 208 [key], fig. 94b.
 1971 *Achelia alaskensis*, - UTINOMI, Publ. Seto Mar. Biol. Lab, **18** (5): 329.
 1986 *Achelia alaskensis*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 44-46, fig. 7.
 1987 *Achelia alaskensis*, - CHILD, Proc. biol. Soc. Wash., **100** (3): 552-553.

Distribution: Connecting a line between mid-Californian, Alaskan, Russian and Korean coasts to central Japan.

Depth range: Littoral-180m.

Habitat: Larvae found on hydromedusa *Polyorchis karaftuoensis*. Adults on subtidal hydroids and algae.

***Achelia anomala* ARNAUD 1974**

- 1974 *Achelia anomala* ARNAUD, Bull. Zool. Mus. Univ. Amsterdam, **3** (21): 174-175, figs. 15-20.

Distribution: Acores.

Depth range: intertidal.

Habitat: rocky shore.

***Achelia armata* (BOUVIER 1916)**

- 1916 *Ammothea (Achelia) armata* BOUVIER, Bull. Mus. natn. Hist. nat., Paris, **22**: 81-83.
 1937 *Achelia armata*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 38-42, figs. 11-13.
 1990 *Achelia armata*, - STOCK, Zool. Meded., **63** (16): 211-213, fig. 3.

Distribution: North-Atlantic.

Depth range: 307-405m.

***Achelia aspera* (LOMAN 1923)**

- 1923 *Ammothea (Achelia) aspera* LOMAN, Ark. Zool., **15** (10): 1-4, figs. 1-5.
 1935 *Ammothea (Achelia) aspera*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1969 *Achelia (Pigropycnon) aspera*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 103-104, figs. 152, 153, 155, 157-159.

Distribution: southern oceans.
Depth range: 0-99m.

Achelia assimilis (HASWELL 1875)

- 1875 *Ammothea assimilis* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1026-1027, pl. LIV, figs. 5-9.
- 1887 *Ammothea Wilsoni* SCHIMKEWITSCH, Zool. Anz., **10** (251): 271.
- 1907 *Ammothea Wilsoni*, - HODGSON, Rep. natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 10-13, fig. 3.
- 1913 *Achelia assimilis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 140.
- 1920 *Achelia assimilis*, - FLYNN, Pap. & Proc. Roy. Soc. Tasm., 1919: 87-90, Pl. 87-90, Pl. XXII, figs. 22-26.
- 1923 *Ammothea (Achelia) wilsoni*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 25.
- 1954 *Achelia assimilis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 97-100, figs. 45-46 [small and large form].
- 1954 *Achelia variabilis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 100-105, figs. 47-48.
- 1957 *Achelia wilsoni*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 100.
- 1961 *Achelia assimilis*, - HEDGPETH, Lunds Univ. Arsskr., N.F., (2) **57** (3): 7-11, figs. 3-7.
- 1963 *Achelia assimilis*, - CLARK, Rec. Austr. Mus., **26** (1): 59.
- 1963 *Achelia variabilis*, - CLARK, Rec. Austr. Mus., **26** (1): 59-60.
- 1968 *Achelia variabilis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 16, fig. 4c.
- 1969 *Achelia (Ignavogriphus) assimilis* Fry & Hedgpeth, - N.Z. Oceanogr. Inst. Mem., **49**: 106, figs. 152, 153, 156.
- 1973 *Achelia assimilis*, - STOCK, Beaufortia, **20** (266): 106.
- 1973 *Achelia assimilis*, - STOCK, Beaufortia, **21** (279): 92.
- 1975 *Achelia assimilis*, - CHILD, Smiths. Contr. Zool., **190**: 24.
- 1988 *Achelia assimilis*, - CHILD, Smiths. Contr. Zool., **468**: 2.
- 1989 *Achelia assimilis*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 124, figs. 1-2.
- 1990 *Achelia assimilis*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 312.
- 1990 *Achelia assimilis*, - GONZALEZ & EDDING, Proc. biol. Soc. Wash., **103** (1): 151.
- 1991 *Achelia assimilis*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 138.
- 1992 *Achelia assimilis*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 155.

Distribution: Widely distributed in tropical and temperate waters of the southwest Pacific, with some scattered locations known from Indonesia, Malaysia, French Polynesia and Chile.

Depth range: 0.5-28m.

Habitat: hard bottom; stones; dead corals; red algae.

Achelia australiensis STOCK 1954

- 1884 *Achelia laevis* var. *australiensis* MIERS, Rep. Zool. Coll. in the Indo-Pac. during Voyage of HMS Alert, 1881-82: 323-324, Pl. XXXV, fig. A.
- 1891 *Ammothea laevis*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 124-127, pl. XIII, figs. 2a-m.

- 1908 *Achelia laevis* var. *australiensis*, - LOMAN, Siboga Exped. Monogr., **40**: 62.
 1929 *Achelia laevis* var. *australiensis*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 139.
 1935 *Ammothea australiensis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1954 *Achelia australiensis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 105-107, fig. 49.
 1956 *Achelia australiensis*, - STOCK, Mitt. Zool. Mus. Hamburg, **54**: 43.
 1961 *Achelia laevis*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 91.
 1969 *Achelia (Ignavogriphus) australiensis*, - FRY & HEDGEPTH, N.Z. Oceanogr. Inst. Mem., **40**: 106, figs. 152, 153, 156.

Distribution: Australia.

Depth range: 5-9m.

Habitat: In mud.

***Achelia barnardi* STOCK 1959**

- 1959 *Achelia barnardi* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 552-553, fig. 2.

Distribution: South Africa.

Depth range: 9-20m.

Habitat: sand, shells and rocks.

***Achelia bituberculata* HEDGPETH 1949**

- 1949 *Achelia bituberculata* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 287-289, fig. 49a-g.
 1951 *Achelia ohshimai* UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (5): 163, fig. 2.
 1954 *Achelia bituberculata*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 94-95, fig. 44.
 1962 *Achelia bituberculata*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 98, fig. 4.
 1965 *Achelia ohshimai*, - UTINOMI, in: New Illustr. Encyclopaedia Fauna Japan, **2**: 336, fig. 20.
 1971 *Achelia bituberculata*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 329.
 1971 *Achelia ohshimai*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 330.
 1983 *Achelia bituberculata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 6-7.
 1984 *Achelia bituberculata*, - KIM, J. Kangreung Natn. Univ., **7**: 537, fig. 6a-i.
 1986 *Achelia bituberculata*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 46.
 1987 *Achelia bituberculata*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 141.
 1987 *Achelia bituberculata*, - NAKAMURA, Biol. Lab. Imperial Household: 18-19, pl. 16.
 1991 *Achelia bituberculata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 3.

Distribution: Japan and Korea.

Depth range: Intertidal-75m.

Habitat: coralline algae; under stones

***Achelia borealis* (SCHIMKEWITSCH 1895)**

- 1895 *Ammothea borealis* SCHIMKEWITSCH, Trudy St. Petersburg Obshch. Iestestvoispytateli, **25**: 36-40, pl. 2, figs. a-b
- 1908 *Ammothea borealis*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 225.
- 1923 *Ammothea borealis japonica*, - LOSINA-LOSINSKY, Izvest. Nauchn. Inst. imeni P.F. Lesgafta, **7**: 45-47, figs. 3a-i.
- 1933 *Ammothea borealis* var. *japonica* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 57-59, 78, fig. 9.
- 1947 *Achelia borealis*, - HEDGPETH, Smiths. Misc. Coll., **106** (18): 24, 27, fig. 13b.
- 1949 *Achelia borealis*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 286-287, fig. 41h-m.
- 1961 *Achelia borealis*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 52 [tab.].
- 1963 *Achelia borealis*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1338.
- 1986 *Achelia borealis*, - KRAPPA, Bonn. zool. Beitr., **37** (4): 301-310 [lit.].

Distribution: White Sea, Kara-Sea, fjords of Novaja Zemlja and Barents-Sea.

Depth range: 130m.

Habitat: muddy bottom.

***Achelia boschi* STOCK 1992**

- 1992 *Achelia boschi* STOCK, Bijdr. Dierk., **62** (2): 86-89, figs. 4-5.

Distribution: Oman.

Depth range: intertidal.

Habitat: "algae"; "under rocks and stones"; "muddy tidal flat with stones and algae" (STOCK).

***Achelia brevicauda* (LOMAN 1904)**

- 1904 *Ammothea brevicauda* LOMAN; Zool. Jb. Syst., XX: 376, pl. 14, figs. 1-4.
- 1954 *Achelia brevicauda*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 140.
- 1959 *Achelia brevicauda*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 552.

Distribution: South Africa.

Depth range: 11-27.5m.

Habitat: on rocks.

***Achelia brevirostris* LOSINA-LOSINSKY 1961**

- 1961 *Achelia brevirostris* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 95-97, fig. 19.
- 1991 *Achelia brevirostris*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 3-5, fig. 1a-f.

Distribution: Sea of Okhotsk (Sakhalin Island); Japan.

Depth range: 48-479m.

***Achelia chelata* (HILTON 1939)**

- 1939 *Ammothea chelata* Hilton, J. Ent. Zool. Pomona College, **31** (2): 30-31.
 1978 *Achelia chelata*, - ARNAUD, Zool. J. Linn. Soc., **63**: 104 [text].

Distribution: California (Vera Cruz).

Habitat: parasitic on the mollusc *Mytilus californicus*; known to be also free-living.

***Achelia columnaris* STOCK 1992**

- 1992 *Achelia columnaris* STOCK, Tijdschr. Ent., **135**: 116-118, 139, figs. 9-18.

Distribution: southern Brazil.

Depth range: 34m.

***Achelia communis* (BOUVIER 1906)**

- 1906 *Ammothea communis* BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, 142: 20, 44-50, figs. 23-32.
 1907 *Ammothea affinis* BOUVIER, L'Expéd. Antarct. Française, 1903-1905: 50-53, figs. 33-36.
 1908 *Ammothea communis*, - HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1): 177-179, pl. II, fig. 1-1a.
 1911 *Ammothea communis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 1140.
 1915 *Achelia communis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 57 [key], 61, fig. 16A-C.
 1915 *Achelia brucei* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 61, fig. 16.
 1923 *Ammothea (Achelia) hoekii* LOMAN, Ark. Zool., **15** (9): 24-25 [part.].
 1927 *Achelia communis*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool., 2): 344.
 1932 *Achelia hoekii*, - GORDON, Discovery Rep., **6**: 110-112, figs. 59a-b, 60a-c.
 1932 *Achelia brucei*, - GORDON, Discovery Rep., **6**: 113.
 1935 *Ammothea communis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1938 *Achelia brucei*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C) (Zool. & Bot.), **2** (8): 22 [key].
 1938 *Achelia communis*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C) (Zool. & Bot.), **2** (8): 22 [key].
 1940 *Achelia brucei*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 79, 81 [key].
 1944 *Achelia brucei*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Exped., (B) **5** (1): 52.
 1944 *Achelia communis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B) **5** (1): 53.
 1950 *Achelia communis*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 153.
 1950 *Achelia* sp. (?*brucei*), - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 154.
 1951 *Achelia brucei*, - SAWAYA, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), **16**: 274 [key].
 1952 *Achelia brucei*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (2): 185.
 1952 *Achelia brucei*, - Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 263.
 1955 *Achelia communis*, - STOCK, Res. Norw. Sci. Exped. Tristan da Cunha, 1937-1938, **33**: 3.

- 1969 *Achelia (Ignavogriphus) communis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 110-112, figs. 152-154, 157, 171-172.
 1969 *Achelia (Ignavogriphus) hoekii*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 106-107, figs. 152-154, 157, 164-165

Distribution: antarctic and subantarctic waters; Tristan da Cunha.

Depth range: 180-300m.

***Achelia crurispinifera* KIM & KIM 1985**

- 1985 *Achelia crurispinifera* KIM & KIM, Kor. J. Zool., **28** (2): 120-124, figs. 1-2.
 1991 *Achelia crurispinifera*, - Smiths. Contr. Zool., **512**: 5.

Distribution: Japan and Korea.

Depth range: 59-150m.

***Achelia deodata* MÜLLER 1990**

- 1990 *Achelia deodata* MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (10): 103-105, figs. 1-4.

Distribution: Kenya.

Habitat: coral reef.

***Achelia discoidea* (EXLINE 1936)**

- 1936 *Ammothea discoidea* EXLINE, Proc. U.S. natn. Mus., **83** (2991): 418-421, fig. 33e-f.

Distribution: temperate north-eastern Pacific (Puget Sound).

Depth range: 40-70m.

Habitat: among hydroids.

***Achelia dohrni* (THOMSON 1884)**

- 1884 *Ammothea dohrni* THOMSON, Trans & Proc. N.Z. Inst., **16** (1883): 243-245, Pl. XIV, figs. 5-9.
 1904 *Ammothea dohrni*, - HUTTON, Index Faunae N.Z.: 247.
 1935 *Ammothea dohrni*, - Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1954 *Achelia dohrni*, - STOCK, Videns. Medd. dansk naturhist. Foren., **116**: 107-109, figs. 50-51.
 1956 *Achelia dohrni*, - STOCK, Mitt. Zool. Mus. Hamburg, **54**: 42.
 1968 *Achelia dohrni*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 17.
 1969 *Achelia (Ignavogriphus) dohrni*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 101-103, 106, fig. 156.
 1971 *Achelia dohrni*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 334.
 1972 *Achelia dohrni*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (3 & 4) (1971): 441-448, figs. 6A-H, 7A-J, 8A-I, 9A-H, 10A-R [variability].

Distribution: New Zealand; subantarctic Islands.

Depth range: Intertidal-192m.

Habitat: rocky shore; coralline algae; muddy bottom and mud with shells and stones; *Cystophora* and *Obelia* hydrozoans; bryozoans.

Achelia echinata HODGE 1864

- 1881 *Ammothea fibulifera* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 141-145, 15, 49, 51, 81, pl. IV, figs. 1-22.
- 1890 ?*Achelia echinata* var. *japonica* ORTMANN, Zool. Jb. (Syst.), **5** (1): 164.
- 1891 *Ammothea echinata*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 120-124, pl. XIII, figs. 1a-m.
- 1899 ?*Achelia echinata* var. *japonica*, - SHISHIDA, Dôzatsu, **11** (128): 199.
- 1919 ?*Achelia echinata* var. *japonica*, - FUKUI, Rigakukai[?], **16** (2): 98.
- 1923 *Ammothea (Achelia) echinata*, - BOUVIER, Faune de France, **7**: 55.
- 1929 *Ammothea echinata*, - GILTAY, Bull. & Ann. Roy. Soc. Belg., **69**: 175.
- 1933 *Ammothea echinata* subsp. *orientalis* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 55-57, 77-78, fig. 7.
- 1936 *Ammothea (Achelia) echinata* var. *sinensis*, LOU, Contr. Inst. Zool., Natn. Acad. Peiping, **3** (1): 19, figs. 7-9, pls. 2-4.
- 1936 ?*Achelia echinata* var. *japonica*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 865.
- 1952 *Achelia echinata*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (6): 530.
- 1938 *Achelia echinata*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 161.
- 1940 *Achelia echinata* ssp. *nasuta* MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 84, 128.
- 1945 *Ammothea echinata*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 163.
- 1949 *Achelia echinata orientalis*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 318.
- 1954 *Achelia echinata*, - BOURDILLON, Notes Stat. Océanogr. Salammbô, **35**: 3.
- 1955 *Achelia echinata*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 213, 245.
- 1956 *Achelia echinata* ssp. *nasuta*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 98-99, fig. 16a.
- 1958 *Achelia echinata*, - STOCK, Bull. Res. Council Israel, (B. Zool.), **7B** (3-4): 138.
- 1958 *Achelia echinata*, - Bull. Sea Fish. Res. Stat. Haifa, **16**: 3.
- 1959 *Achelia echinata*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 201.
- 1962 *Achelia echinata*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 97.
- 1966 *Achelia echinata*, - FAGE & STOCK, Ann. Inst. Océanogr. Monaco, **44**: 317.
- 1966 *Achelia echinata*, - STOCK, Atlantide Rep., **9**: 45.
- 1967 *Achelia echinata sinensis*, - NESIS, Issöedov. Fauny Morei, **5** (13): 249.
- 1971 *Achelia echinata sinensis*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 328-329.
- 1972 *Achelia echinata*, - KING, J. nat. Hist., **6**: 623.
- 1973 *Achelia echinata*, - ARNAUD, Tethys, **5** (1): 151.
- 1973 *Achelia echinata*, - KRAPPA, Beaufortia, **21** (277): 58.
- 1973 *Achelia echinata tuberculata* LOSINA-LOSINSKY, Zool. Zh., **52** (7): 1084.
- 1974 *Achelia echinata*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 23 [key], 28-30, figs. 11-12, 14C.
- 1975 *Achelia echinata*, - KRAPPA-SCHICKEL & KRAPPA, Vie Milieu, (B) **25** (1): 14 fig. 4, 17, 18-20 (tab. II), 21-23 (ecology).
- 1976 *Achelia echinata*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.

- 1976 *Achelia echinata*, - WOLFF, Netherlands J. Sea Res., **10** (4): 473-474, fig. 2b (distribution in estuary).
- 1978 *Achelia echinata*, - CHIMENTZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 98.
- 1980 *Achelia echinata*, - MUNILLA LEON, Cah. Biol. Mar., **21**: 115-121, figs. 1-3, tab. I (annual development and reproduction).
- 1981 *Achelia echinata*, - MUNILLA & de HARO, - Bijdr. Dierk., **51** (2): 191-197, fig. 3, 11, tab. I (electrophoretical and immunological study).
- 1983 *Achelia echinata*, - KRAPF, Bonn. zool. Beitr., **34** (1-3): 408.
- 1983 *Achelia echinata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 7-8.
- 1984 *Achelia echinata sinensis*, - KIM, J. Kangreung Natn. Univ., **7**: 537, fig. 7a-i.
- 1984 *Achelia echinata*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 531-535, fig. 1, tab I-IV.
- 1986 *Achelia echinata sinensis*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 2 (2): 46.
- 1986 *Achelia echinata*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
- 1987 *Achelia echinata*, - ARNAUD, Mesogee, **47**: 38.
- 1987 *Achelia echinata*, - BITAR, Bull. Inst. Sci., Rabat, **11**: 163.
- 1987 *Achelia echinata sinensis*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 141-142.
- 1987 *Achelia echinata sinensis*, - NAKAMURA, Biol. Lab., Imperial Household: 19-21, pl. 17.
- 1987 *Achelia echinata*, - SANCHEZ & MUNILLA, Vieraea, **17**: 182.
- 1987 *Achelia echinata*, - STOCK, Cah. Biol. Mar., **28**: 506.
- 1989 *Achelia echinata*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 287 [tab. 1, 2], 291, 293 (phenology).
- 1990 *Achelia echinata*, - STOCK, Zool. Meded., **63** (16): 217.
- 1991 *Achelia echinata sinensis*, - Smiths. Contr. Zool., **512**: 5.
- 1992 *Achelia echinata*, - PÉREZ-RUZAFA, A. & T. MUNILLA, Sci. Mar., **56** (1): 23.
- 1993 *Achelia echinata*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: in boreal waters of the N-Atlantic, Mediterranean and NW-Pacific.

Depth range: 0-336m.

Habitat: In a variety of substrata, often very abundant; *Cystoseira adriatica*, *Halopteris*; *Laurencia obtusa*; *Dictyobteris membranacea*; *Peyssonnelia squamaria*; *Sargassum linifolium* (algae); *Posidonia* (seagrass); feeds on *Flustra foliacea*, *Dynamene pumila* (hydroids) and on detritus in *Bugula* colonies; ectocommensal on *Eunicella*; on *Suberites* spp. and *Diazona violacea*; amongst hydroids on top of boulder; amongst mud; amongst red weed; amongst *Serturella*, *Lepralia* and commensals on top of boulder; tunicates on gravel and sand; amongst sponges, polyzoa, tunicates and *Serturella*; amongst *Luidia*, *Cellaria* and debris; amongst epiphytes on *Pecten*; on *Dynamena pumila* and *Flustra foliacea* and feeds on the latter polyzoan by thrusting its proboscis into the operculum when it opens.

***Achelia euchelata* (HEDGPETH 1940)**

- 1940 *Ammothaea euchelata* HEDGPETH, J. Wash. Acad. Sci., **30** (2): 84-86, fig. 1.

Distribution: California.
Habitat: Bugula on reef.

Achelia fernandeziana (LOMAN 1920)

- 1920 *Ammothea fernandeziana* LOMAN, in: Nat. Hist. Juan Fernandez, **3**: 142-144, fig. 2.
- 1934 *Achelia fernandeziana*, - GILTAY, Bull. Mus. Roy. Hist. Nat. Belg., **10** (18): 5.
- 1935 *Ammothea fernandeziana*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
- 1940 *Achelia fernandeziana*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 81 [key].
- 1951 *Achelia fernandeziana*, - SAWAYA, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), **16**: 274 [key].
- 1969 ***Achelia fernandeziana***, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 104, figs. 152, 153, 155, 160.
- 1992 *Achelia fernandeziana*, - CHILD, Smiths. Contr. Zool., **526**: 3-4.

Distribution: Juan Fernandez Island, SE-Pacific (off Chile); Chile.

Depth range: 10-180m.

Habitat: coralline algae.

Achelia gracilipes (COLE 1904)

- 1904 *Ammothea gracilipes* COLE, Harriman Alaska Exped., **10**: 269-270, pl. XII, fig. 5, pl. XVIII, figs. 1-6.
- 1933 *Ammothea gracilipes* var. *borealis* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 78.
- 1951 *Achelia gracilipes*, - HEDGPETH, Wasmann J. Biol., **9** (1): 106.

Distribution: California, E-U.S.S.R.

Depth range: ?intertidal-4m.

Habitat: hydroids.

Achelia gracilis VERRILL 1900

- 1900 *Achelia ?gracilis* VERRILL, Trans. Conn. Acad. Sci., **10** (15): 582, pl. 70, fig. 10.
- 1904 *Ammothea gracilis*, - COLE, Proc. Boston Soc. Nat. Hist., **31** (8): 317-323, pl. 21, fig. 4-14.
- 1934 *Ammothea (Achelia) gracilis*, - GILTAY, Bull. Mus. Roy. Hist. Nat. Belg., **10** (18): 5.
- 1935 *Ammothea gracilis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
- 1940 *Achelia gracilis*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 79.
- 1948 *Achelia gracilis*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 244.
- 1954 *Achelia gracilis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 117.
- 1955 *Achelia gracilis*, - BOUDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 597.

- 1969 *Achelia (Pigrolavatus) gracilis*, - FRY & HEDGPETH, N.Z. oceanogr. Inst. Mem., **40**: 104, figs. 152, 153, 155.
- 1975 *Achelia gracilis*, - STOCK, Bull. mar. Sci., **24** (4): 983.
- 1979 *Achelia gracilis*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 10.
- 1986 *Achelia gracilis*, - STOCK, Bull. mar. Sci. **38** (3): 400 tab. 1, 416.
- 1992 *Achelia gracilis*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 43, fig. 1.
- 1992 *Achelia gracilis*, - STOCK, Tijdschr. Ent., **135**: 118, 139.

Distribution: Tropical and subtropical W-Atlantic, Bermuda to Brazil.

Depth range: 0-157 m. Records from deep-water probably from fouling on ship's hulls, when deep-water samples have been taken.

Habitat: coral rubble; fouling on pilings; sponges; hydroids, octocorals, bryozoans; hydroids, sponges and algae on shipwreck.

***Achelia hoekii* (PFEFFER 1889)**

- 1889 *Ammothea hoekii* PFEFFER, Jb. Hamburg. Wiss. Anst., **6** (2): 46.
- 1923 *Ammothea (Achelia) hoekii*, - LOMAN, Ark. Zool., **15** (9): 4.
- 1923 *Ammothea (Achelia) hoekii*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 24.
- 1932 *Achelia hoekii*, - GORDON, Discovery Rep., **6**: 110-112, figs. 59a, b, 60a, c.
- 1935 *Ammothea hoekii*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
- 1938 *Achelia hoekii*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 22 [key].
- 1951 *Achelia hoekii*, - SAWAYA, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), **16**: [key].
- 1969 *Achelia (Ignavogriphus) hoekii*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 106.

Distribution: Antarctic.

Depth range: 5-60m.

***Achelia kamtschatica* LOSINA-LOSINSKY 1961**

- 1961 *Achelia kamtschatika* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 92-93, fig. 17.

Distribution: NW-Pacific.

Depth range: 30-50m.

***Achelia kiiensis* UTINOMI 1951**

- 1951 *Achelia kiiensis* UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 161-163, fig. 1.
- 1971 *Achelia kiiensis*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 330.

Distribution: Japan.

Depth range: 5-20m.

Achelia kurilensis LOSINA-LOSINSKY 1961

1961 *Achelia kurilensis* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, 7: 98-100, fig. 21.

Distribution: NW-Pacific.
Depth range:

Achelia lagenaria STOCK 1992

1992 *Achelia lagenaria* STOCK, Bijdr. Dierk., 62 (2): 89-92, figs. 7-8.

Distribution: Oman.
Habitat: on algae.

Achelia langi (DOHRN 1881)

- 1881 *Ammothea langi* DOHRN, Monogr. Fauna Flora Golf. Neapel, 3: 146-147, pl. 5, figs. 1-8.
 1901 *Ammothea Langii*, - d'ARCY THOMPSON, in: A catalogue of Crustacea and Pycnogonida contained in the Museum of University College, Dundee: 55.
 1923 *Ammothea (Achelia) Langi*, - BOUVIER, Faune de France, 7: 55.
 1925 *Ammothea (Achelia) Langi*, - LOMAN, Bull. Soc. Sci. Nat. Maroc, 5 (3): 50.
 1942 *Achelia Langi*, - FAGE, Arch. Zool. Ge., 82 (Notes & Rev., 2): 87-88.
 1967 *Achelia langi*, - NOGUEIRA, Arqu. Mus. Bocage, (2) 1 (15): 324-325.
 1968 *Achelia langi*, - STOCK, Vie Milieu, 19 (1A): 16, fig. 9.
 1972 *Achelia langi*, - LEUNG, Téthys, 3 (4): 821.
 1979 *Achelia cf. langi*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., 59 (184): 10-11.
 1986 *Achelia langi?*, - STOCK, Bull. mar. Sci., 38 (3): 400 tab. 1.
 1987 *Achelia langi*, - ARNAUD, Mesogee, 47: 39.
 1990 *Achelia langi*, - Stock, Zool. Meded., 63 (16): 209-211, fig. 2.

Distribution: Mediterranean and temperate E-Atlantic;
?Caribbean (Anguilla).

Depth range: 1-100m.
Habitat: algae and seagrass.

Achelia latifrons (COLE 1904)

- 1904 *Ammothea latifrons* COLE, Harriman Alaska Exped., 10: 263-266, pl. XI, fig. 3, pl. XVI, figs. 1-9, pl. XVII, figs. 1-3.
 1934 *Ammothea latifrons*, - SCHMITT, J. Wash. Acad. Sci., 24 (1): 68-69.
 1941 *Ammothea latifrons*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., 9 (26): 256 [key].
 1942 *Ammothea latifrons*, - HILTON, Rep. Allan Hancock Pac. Exped., 5 (9): 295-296, pl. 41.
 1942 *Ammothea latifrons*, - HILTON, J. Ent. Zool. Pomona College, 34 (4): 93.
 1954 *Achelia latifrons*, - Vidensk. Medd. dansk naturhist. Foren., 116: 96.

Distribution: Bering Sea, Alaska, California, and Japan.
Depth range: ?110-183m.
Habitat: Washed from *Sertularella*.

Achelia lavrentii (LOSINA-LOSINSKY 1933)

1933 *Ammothea lavrentii* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 49-50, 75-76, fig. 3.

Distribution: NW-Pacific (U.S.S.R.).
Depth range: shallow water.

Achelia litke (LOSINA-LOSINSKY 1933)

1933 *Ammothea litke* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 52-53, 76-77, fig. 5.

1933 *Ammothea litke* var. *intermedia* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 53-54, 77, fig. 6.

Distribution: NW-Pacific (U.S.S.R.).
Depth range: 10m.

Achelia megacephala HODGSON 1915

1927 *Achelia megacephala*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 345-346, fig. 12.

Distribution: Kerguelen (Observatory Bay).

Achelia nana (LOMAN 1908)

1908 *Ammothea nana* LOMAN, Siboga Exped. Monogr., **40**: 60-61, pl. I, figs. 1-13.
 1953 *Achelia nana*, - STOCK, Temminckia, **9**: 300-301, fig. 14-15, fig. 14.
 1954 *Achelia nana*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 97.
 1965 *Achelia nana*, - STOCK, Beaufortia, **13** (151): 14-15, figs. 1-3.
 1968 *Achelia nana*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 16.
 1971 *Achelia nana*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 329-330.
 1974 *Achelia nana*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 13-14.
 1977 *Achelia nana*, - CHILD, Proc. biol. Soc. Wash., **90** (2): xxx.
 1983 *Achelia nana*, - CHILD, Proc. biol. Soc. Wash., **96** (4): 699.
 1988 *Achelia nana*, - CHILD, Bull. biol. Soc. Wash., **8**: 50.
 1991 *Achelia nana*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 161.

Distribution: Tropical to temperate Indo-West Pacific. Several localities from Madagascar to Japan, New Caledonia and Micronesia.

Depth range: 0-435m.

Habitat: coral reefs; dead corals, rubble; algae; sponges; octocorals.

Achelia neotenica KRAPP 1986

- 1907 *Ammothea (Achelia) borealis*, - SCHIMKEWITSCH, Rés. Sci. Expéd. polaire Russe 1900-1903, ... (8) **18** (6): 5-8, figs. 1-6.
 1929 *Ammothea borealis*, - LOSINA-LOSINSKY, Ber. Wiss. Meeresinst. Moskau, **4** (3): 73, figs. 3 (6a+b), 4 (A, A1, B, B1, C, C1, D, D1).
 1930 *Ammothea (Achelia) borealis*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 139-144, pl 2, fig. 2; figs. 34-37.
 1948 *Ammothea borealis*, - BRONSTEIN, Opredel. Fauni Flori Severnich Morei SSSR: 350, pl. XCIV: fig. 4.
 1986 *Achelia neotenica* KRAPP, Bonn. zool. Beitr., **37** (4): 303-309, figs. 1-9.

Distribution: Arctica. - White Sea, Kara-Sea, fjords of Novaja Zemlja and Barents-Sea.

Achelia orientalis SCHIMKEWITSCH 1913

- 1913 *Ammothea orientalis* SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St. Petersburg, 18: 241-242, pl. III, figs. 1-7.

Distribution: NW-Pacific (Kurile Strait).

Achelia orpax NAKAMURA & CHILD 1983

- 1983 *Achelia orpax* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 8-10, fig. 2.
 1986 *Achelia latifrons*, KIM & HONG, Kor. J. Syst. Zool., **2** (2): 46-48, fig. 8.
 1991 *Achelia orpax*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 5-6, fig. 1g-h.

Distribution: Japan, Korea.

Depth range: 7-30m.

Achelia paradisiaca LOMAN 1923

- 1923 *Ammothea (Ammothella) paradisiaca* LOMAN, Ark. Zool., **15** (9): 4-6, fig. B1-3.

Distribution: subantarctic area (Valparaiso).

Depth range: shallow water.

Habitat: "from roots of sea-weed" (LOMAN).

Achelia parvula (LOMAN 1923)

- 1923 *Ammothea (Achelia) parvula* LOMAN, Ark. Zool., **15** (9): 2-4, fig. A1-6.
 1932 *Achelia parvula*, - GORDON, Discovery Rep., **6**: 113-114, fig. 62.
 1935 *Ammothea (Achelia) parvula*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1938 *Achelia parvula*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 22 [key].
 1940 *Achelia parvula*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 80 [key].

- 1950 *Achelia parvula*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 153.
 1951 *Achelia parvula*, - SAWAYA, Bol. Fac. Fil., Cién. Letr. Univ. S. Paulo,
 (Zool.), **16**: 274 [key].
 1961 *Achelia parvula*, - HEDGPETH, Lunds Univ. Arsskr., N.F., (2) **57** (3): 12.
 1969 *Achelia (Pigrolavatus) parvulus*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst.
 Mem., **49**: 103-104, figs. 157-159.
 1992 *Achelia parvula*, - CHILD, Smiths. Contr. Zool., **526**: 4.

Distribution: Ecuador; Straits of Magellan; Falkland Islands.

Depth range: 0-20m.

***Achelia pribilofensis* (COLE 1904)**

- 1904 *Ammothea pribilofensis* COLE, Harriman Alaska Exped., **10**: 270-273, pl. 12,
 fig. 6, pl. 18, figs. 7-8, pl. 19, figs. 1-8.
 1930 *Ammothea (Achelia) pribilovensis*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo
 Akad. Nauk SSSR, **2**: 156-160, figs. 46-49.
 1949 *Achelia pribilofensis*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 287.

Distribution: Alaska, Kuriles, Japan.

***Achelia quadridentata* (HODGSON 1910)**

- 1910 *Ammothea quadridentata* HODGSON, Denkschr. Med.-Nat. Ges. Jena, **16**: 223-225,
 figs. 2-3.
 1954 *Achelia quadridentata*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 138-139, fig.
 24.
 1957 *Achelia quadridentata*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 84.
 1959 *Achelia quadridentata*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 552.
 1962 *Achelia quadridentata*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 273.
 1974 *Achelia quadridentata*, - ARNAUD, Tethys, **5** (4): 787-789, figs. 20-25.
 1988 *Achelia quadridentata*, - MUNILLA, Monogr. Zool. Mar., **3**: 195-196, fig. 11.

Distribution: South Africa; Namibia; St. Paul and Amsterdam islands.

Depth range: 0-21m.

Habitat: on rocks and in kelp vegetation.

***Achelia salebrosa* LOSINA-LOSINSKY 1961**

- 1961 *Achelia salebrosa* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR,
 Leningrad, **7**: 97-98, fig. 20.

Distribution: NW-Pacific.

Depth range: 414m.

***Achelia sawayai* MARCUS 1940**

- 1940 *Achelia sawayai* MARCUS, Bolm. Fac. Fil. Cién. Letr. Univ. S. Paulo, ser. **19**
 (Zool.), **4**: 81-86, figs. 10a-f, 17.

- 1948 *Achelia sawayai*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 244.
- 1949 *Achelia Sawayai*, - FAGE, Arch. Zool. Exp. Gen., **86** (Notes & Rev., 1): 28.
- 1951 *Achelia besnardi*, - SAWAYA, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), **16**: 271-279, figs. 1-2.
- 1954 *Achelia sawayai*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 117.
- 1955 *Achelia sawayai*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 245.
- 1966 *Achelia sawayai*, - STOCK, Atlantide Rep., **9**: 46.
- 1966 *Achelia sawayai*, - FAGE & STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 317.
- 1969 *Achelia besnardi*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 105, figs. 153, 155.
- 1969 *Achelia (Pigrolavatus) sawayai*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 104, figs. 152, 153, 155.
- 1974 *Achelia sawayai*, - CHILD, Proc. biol. Soc. Wash., **87** (43): 497.
- 1974 *Achelia sawayai*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 14.
- 1975 *Achelia sawayai*, - STOCK, Bull. mar. Sci., **24** (4): 982-983.
- 1975 *Achelia besnardi*, - STOCK, Ent. Ber., **35** (5): 70-71, figs. 1-9.
- 1976 *Achelia sawayai*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 342-343, fig. 3.
- 1979 *Achelia sawayai*, - CHILD, Smiths. Contr. Zool., **293**: 7-8.
- 1979 *Achelia sawayai*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 9-10.
- 1982 *Achelia sawayai*, - CHILD, Smiths. Contr. mar. Sci., **12**: 356-357.
- 1986 *Achelia sawayai* f. *typica*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
- 1986 *Achelia sawayai* f. *besnardi*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
- 1989 *Achelia sawayai*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 124, fig. 40.
- 1989 *Achelia sawayai*, - STOCK, Bijdr. Dierk., **59** (2): 87-89.
- 1990 *Achelia sawayai*, - MÜLLER, Senckenbergiana biol., **70** (1989) (1/3): 186-188, figs. 1-10.
- 1990 *Achelia sawayai*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (10): 105-106.
- 1990 *Achelia sawayai*, - MÜLLER, Bonn. zool. Beitr., **41** (3/4): 277.
- 1990 *Achelia sawayai*, - STOCK, Zool. Meded., **63** (16): 205.
- 1992 *Achelia sawayai*, - CHILD, Mem. Hourglass Cruises, **9** (1): 7 [key], 9 [tab. 3], 11-12, fig. 3, 76 [tab. 12].
- 1992 *Achelia sawayai*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 43.
- 1992 *Achelia besnardi*, - STOCK, Tijdschr. Ent., **135**: 116, 139.
- 1992 *Achelia sawayai*, - STOCK, Tijdschr. Ent., **135**: 118-119, 139.

Distribution: pantropical, most records from the W-Atlantic.

Depth range: 0-157m. Records from deeper water probably from fouling of ship's hulls, when deep-water samples have been taken.

Habitat: Lives in a variety of substrata, more often found associated with seagrass (*Thalassia*).

***Achelia scabra* WILSON 1880**

- 1880 *Achelia scabra* WILSON, Rep. U.S. Fish. Comm., **6**: 475-476.
- 1942 *Achelia scabra*, - GILTAY, J. Fish. Res. Board Ca., **5** (5): 460.
- 1943 *Ammothea scabra*, - NEEDLER, Ca. Atl. Fauna, **10**: 16, fig. 21a-c.
- 1948 *Achelia scabra*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 244.

Distribution: NW-Atlantic, Canada.

Depth range: 40-82m.

Achelia segmentata UTINOMI 1954

- 1954 *Achelia segmentata* UTINOMI, Publ. Akkeshi Mar. Biol. Star., 3: 20, figs.
[not seen].
1954 *Achelia segmentata* UTINOMI, Publ. Seto Mar. Biol. Lab., 18 (5): 330.

Distribution: Japan.

Habitat: on red algae and *Strongylocentrotus nudus*.

Achelia serratipalpis (BOUVIER 1911)

- 1911 *Ammothea serratipalpis* BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, 152:
1140.
1913 *Achelia serratipalpis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-
1910), 6: 140-144, figs. 90-95.
1920 *Ammothea (Achelia) serratipalpis*, - LOMAN, in: The Nat. Hist. of Juan
Fernandez and Easter Isl., 3: 141.
1923 *Ammothea (Achelia) serratipalpis*, - LOMAN, Further Zool. Res. Swed. Antarct.
Exped., 1 (2): 25.
1932 *Achelia serratipalpis*, - GORDON, Discovery Rep., 6: 113-114.
1935 *Ammothea serratipalpis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs,
5 (4) (2): 285.
1938 *Achelia serratipalpis*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C),
(Zool. & Bot.), 2 (8): 24.
1969 *Achelia (Ignavogriphus) serratipalpis*, - FRY & HEDGPETH, N.Z. Oceanogr.
Inst. Mem., 49: 107-109, figs. 152-154, 156, 166-167.

Distribution: Antarctic and subantarctic area.

Depth range: 64-391m.

Achelia setulosa (LOMAN 1912)

- 1912 *Ammothea setulosa* LOMAN, Bull. Inst. Océanogr. Monaco, 238: 13-14.
1917 *Ammothea (Achelia) setulosa*, - BOUVIER, Res. Campagnes Sci. accomplies sur
son Yacht par Albert Ier Prince Souverain de Monaco, 51: 41-42, pl. IV,
figs. 9-10..
1966 *Achelia setulosa*, - FAGE & STOCK, Ann. Inst. Oceanogr. Monaco, 44: 317-318,
fig. 1.
1990 *Achelia setulosa*, - STOCK, Zool. Meded., 63 (16): 218.

Distribution: Cape Verde Islands.

Depth range: 20-120m.

Habitat: sand; coralline algae; bryozoans.

Achelia sheperdi STOCK 1973

- 1973 *Achelia sheperdi* STOCK, Beaufortia, 21 (279): 92-95, fig. 1.
1975 *Achelia sheperdi*, - CHILD, Smiths. Contr. Zool., 190: 22-24, fig. 10.

Distribution: Australia, incl. Tasmania.
Depth range: ?intertidal-28m.
Habitat: rocks and algae.

Achelia simplex GILTAY 1934

- 1934 *Achelia simplex* GILTAY, Bull. Mus. roy. Hist. nat. Belg., **X** (18): 1-4, figs. 1-3.
 1952 *Achelia simplex*, - STOCK, Boll. Soc. Veneziana Stor. Nat., **6** (2): 182.
 1971 *Achelia simplex*, - WYER, KING & JARVIS, Ir. Nat., **17**: 92-94.
 1974 *Achelia simplex*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 23 [key], 32-33, fig. 13, 14D.
 1975 *Achelia simplex*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 14 fig. 4, 17, 18-20 (tab. II) 21-23 (ecology).
 1987 *Achelia simplex*, - ARNAUD, Mesogee, **47**: 39.
 1990 *Achelia simplex*, - STOCK, Cah. Biol. Mar., **31**: 516.

Distribution: Mediterranean and temperate NE-Atlantic.

Depth range: 0-8m.

Habitat: *Corallina mediterranea*; *Halimeda tuna*; *Halopteris scoparia*; *Peyssonnelia squamaria*; *Cutleria* sp.; *Cystoseira abrotanifolia*, *adriatica*, *crinita* and *discors*; *Padina pavonia*; *Sargassum linifolium*.

Achelia simplissima (HILTON 1939)

- 1939 *Ammothaea simplissima* HILTON, J. Ent. Zool. Pomona College, **31** (2): 31-32.
 1951 *Achelia simplissima*, - HEDGPETH, Wasmann J. Biol., **9** (1): 106-108.

Distribution: California.

Habitat: bryozoans.

Achelia socors (LOMAN 1908)

- 1908 *Ammothaea socors* LOMAN, Siboga Exped. Monogr., **40**: 61-62, pl. I, figs. 14-19.
 1985 *Achelia socors*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **10** (18): 157.
 1991 *Achelia socors*, - STOCK, Mém. Mus. natn. Hist. nat (A), **151**: 220.

Distribution: Indonesia (between Ceram, and Buru and Strait of Makassar), Philippines.

Depth range: 24-97m.

Achelia spatula NAKAMURA & CHILD 1983

- 1983 *Achelia spatula* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 10-12, fig. 3.

Distribution: Japan.

Depth range: 7-15m.

Achelia spicata (HODGSON 1915)

- 1915 *Austrothea spicata* HODGSON, Ann. Mag. Nat. Hist., (8) **15** (85): 147.
 1915 *Achelia intermedia* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3**: 57 [key], 60, fig. 15.
 1915 *Achelia spicata*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 57-60, figs. 13-14.
 1927 *Achelia intermedia*, - HODGSON, Dt. Südpolar Exped., **19** (Zool. II): 345 [key].
 1932 *Achelia intermedia*, - GORDON, Discovery Rep., **6**: 112, fig. 61.
 1935 *Ammothea spicata*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1938 *Achelia spicata*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot), **2** (8): 22 [key], 23.
 1940 *Achelia intermedia*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 80, 81 [key].
 1944 *Achelia intermedia*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Exped., (B), **5** (1): 52.
 1944 *Achelia spicata*, - GORDON, Rep. Brit. Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 52 [key].
 1947 *Achelia intermedia*, - STEPHENSEN, Sci. Res. Norw. Antarct. Exped., 1927-1928, **27**: 85.
 1950 *Achelia intermedia*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 153, 159.
 1950 *Achelia spicata*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 153, 159.
 1951 *Achelia intermedia*, - SAWAYA, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), **16**: 274 [key].
 1952 *Achelia intermedia*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (2): 181.
 1952 *Achelia intermedia*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 263.
 1952 *Achelia spicata*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 263.
 1969 *Achelia (Ignavogriphus) spicata*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 103 [key], 109-110, figs. 168-170, 152-154, 157.
 1980 *Achelia spicata*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 2.

Distribution: Arctic and subantarctic area.

Depth range: 30-470m.

***Achelia spinosa* WILSON 1878**

- 1853 *Zetes spinosa* STIMPSON, Smiths. Contr. Knowl., **6** (5): 37.
 1878 *Achelia spinosa* WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 7-8, pl. 2, fig. 1a-h.
 1878 *Ammothea achelioides* WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 16-17, pl. 5, fig. 1a-e.
 1880 *Achelia spinosa*, - WILSON, Rep. U.S. Fish. Comm., **6**: 473-476, pl. 1, fig. 1, pl. 2, fig. 8.
 1880 *Ammothea achelioides*, - WILSON, Rep. U.S. Fish. Comm., **6**: 484-485, pl. 4, figs. 19-20.
 1901 *Achelia spinosa*, - WHITEAVES, Geol. Surv. Can., **772**: 262.
 1901 *Ammothea achelioides*, - WHITEAVES, Geol. Surv. Can., **772**: 203.
 1908 *Ammothea echinata*, - J. Linn. Soc. London (Zool.), **30**: 224.
 1930 *Ammothea (Achelia) echinata?* var. *spinosa*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 133-136.
 1943 *Ammothea spinosa*, - NEEDLER, Can. Atl. Fauna, **10**: 16, fig. 20a-d.

- 1948 *Achelia spinosa*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 212-214, fig. 38a-b.
 1954 *Achelia spinosa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 96.
 1963 *Achelia spinosa*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1337-1338.

Distribution: E-coast of N-America; Japan; Bering Sea; Point Barrow.

Depth range: ?-732m.

***Achelia spinoseta* (HILTON 1939)**

- 1939 *Ammothea spinoseta* HILTON, J. Ent. Zool. Pomona College, **31** (2): 31.
 1939 *Achelia spinoseta*, - HEDGPETH, Wasmann J. Biol., **9** (1): 106.

Distribution: California.

Depth range: intertidal.

Habitat: rocky shore.

***Achelia sufflata* GORDON 1944**

- 1944 *Achelia sufflata* GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B), **5** (1): 53-54, fig. 18a-e.
 1969 *Achelia (Pigrolavatus) sufflata*, - FRY & HEDGPETH, N.Z. Oceanogr. Onst. Mem., **49**: 105-106, figs. 153, 155, 156, 161, 162.

Distribution: Arctic.

Depth range: 180-300m.

***Achelia superba* (LOMAN 1911)**

- 1911 *Ammothea superba* LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 11-12, pls. 1-2, figs. 14-24.
 1936 *Ammothea superba*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 866.
 1947 *Ammothea superba*, - OHSHIMA & KISHIDA, in: Illustr. Encyclopaedia Fauna Japan: 1008, fig. 2860.
 1949 *Achelia superba*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 288.
 1951 *Achelia superba*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 160-161.
 1954 *Achelia superba*, - STOCK, Vidensk. Medd. dansk naturhist. Foren, **116**: 96.
 1955 *Achelia superba*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 23-24, fig. 13.
 1959 *Achelia superba*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 202.
 1962 *Achelia superba*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 97.
 1971 *Achelia superba*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 328.
 1983 *Achelia superba*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 12.
 1987 *Achelia superba*, - NAKAMURA, Biol. Lab., Imperial Household: 21-23, pls. 18, 36.
 1990 *Achelia superba*, - TURPAYEVA, Zool. Zh., **69** (9): 19-20, fig. 2.
 1991 *Achelia superba*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 6.

Distribution: NW-Pacific, from the Sea of Okhotsk off Sakhalin south to Kii Strait.

Depth range: 7-300m.

Habitat: sand.

Achelia tenuipes STOCK 1990

1990 *Achelia tenuipes* STOCK, Zool. Meded., **63** (16): 215-217, fig. 5.

Distribution: Cape Verde Islands.
Depth range: 1200m.

Achelia transfuga STOCK 1954

1954 *Achelia transfuga* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 109-112, figs. 52-53.
 1969 *Aduncorostris transfuga*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 113.

Distribution: Antarctica; Lyttelton harbour.
Depth range: 2-10m.
Habitat: muddy bottom.

Achelia transfugoides STOCK 1973

1973 *Achelia transfugoides* STOCK, Beaufortia, **20** (266): 104-106, fig. 2.
 1973 *Achelia transfugoides*, - STOCK, Beaufortia, **21** (279): 92.

Distribution: Australia (West Island, Toad Head).
Depth range: 5-28m.
Habitat: *Phacelocarpus* (algae).

Achelia turba STOCK 1990

1990 *Achelia turba* STOCK, Zool. Meded., **63** (16): 213-215, fig. 4.

Distribution: Canary Islands.
Depth range: 0-1m.
Habitat: rocky shore; sublittoral sand; under stones with actinaria.

Achelia uschakovi (LOSINA-LOSINSKY 1933)

1933 *Ammothea uschakovi* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 50-52, 76, fig. 4.

Distribution: Bering Sea (Lawrentji Bay).
Depth range: 8-10m.

Achelia vulgaris (COSTA 1861)

1861 *Alcinous vulgaris* COSTA, Microdoride mediterranea, o descrizione de' poco ben conosciuti ignoti viventi e microscopici del Mediterraneo, etc., **1**: 11.

- 1881 *Ammothea fransciscana* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 135-141, pl. III.
- 1923 *Achelia vulgaris*, - BOUVIER, Faune de France, **7**: 55.
- 1954 *Achelia vulgaris*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 100.
- 1958 *Achelia vulgaris*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 3.
- 1959 *Achelia vulgaris*, - GOTTLIEB, Thesis; Hebrew Univ. Jerusalem: 114.
- 1962 *Achelia vulgaris*, - STOCK, Ent. Ber., Amsterdam, **22** (11): 219.
- 1966 *Achelia vulgaris*, - de HARO, Invest. Pesqu., **30**: 665.
- 1966 *Achelia vulgaris*, - LEDOYER, Rec. Trav. Stat. Mar. d'Endoume, **41** (57): 162.
- 1966 *Achelia vulgaris*, - STOCK, Atlantide Rep., **9**: 45-46.
- 1966 *Achelia vulgaris*, - FAGE & STOCK, Ann. Inst. Océanogr. Monaco., **44**: 316.
- 1968 *Achelia vulgaris*, - LEDOYER, Rec. Trav. Stat. Mar. d'Endoume, **44** (60): 170.
- 1973 *Achelia vulgaris*, - ARNAUD, Tethys, **5** (1): 151.
- 1973 *Achelia vulgaris*, - KRAPP, Beaufortia, **21** (277): 58.
- 1975 *Achelia vulgaris*, - KRAPP, Bonn. zool. Beitr., **26** (1-3): 284.
- 1976 *Achelia vulgaris*, - ARNAUD, Acta Ecol. Iranica, **1**: 71.
- 1978 *Achelia vulgaris*, - MUNILLA, Publ. Univ. Autonom. Barcelona Ballaterra, Fac. Cien. (Tesis Doc.): 8.
- 1981 *Achelia vulgaris*, - MUNILLA, Bull. Inst. Catalana Hist. Nat., **47** (Sec. Zool., 4): 80.
- 1983 *Achelia vulgaris*, - KRAPP, Bonn. zool. Beitr., **34** (1-3): 408.
- 1984 *Achelia vulgaris*, - MUNILLA & de HARO, in: Els sistemes de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 531, 533-535, tab. III-IV.
- 1987 *Achelia vulgaris*, - ARNAUD, Mesogee, **47**: 39.
- 1987 *Achelia vulgaris*, - SANCHEZ & MUNILLA, Vieraea, **17**: 182.
- 1990 *Achelia vulgaris*, - STOCK, Zool. Meded., **63** (16): 218.

Distribution: Mediterranean, Atlantic coast of Morocco, Canary and Cape Verde Islands.

Depth range: 0-300m.

Habitat: found in a variety of substrata.

***Acheliana* ARNAUD 1971**

- 1971 *Acheliana* ARNAUD, Beaufortia, **18** (241): 199 [diagnosis], 201 [remarks].

***Acheliana tropicalis* ARNAUD 1971**

- 1971 *Acheliana tropicalis* ARNAUD, Beaufortia, **18** (241): 201-204, figs. 1-9.

Distribution: Madagascar.

Depth range: 0.5m.

Habitat: on reef, "platier externe algal" (ARNAUD).

***Ammothea* LEACH 1814**

- 1932 *Ammothea*, - GORDON, Discovery Rep., **6**: 94 [remarks], 95-96 [key to antarctic and subantarctic species].
- 1956 *Ammothea*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **54**: 45 [diagnosis].
- 1969 *Ammothea*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 67-63 [general remarks], 71 [key to species], 73 [diagnosis].
- 1969 *Magnammothea* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 90 [diagnosis].

- 1969 *Thavmastopycnon* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 91 [diagnosis].
 1969 *Athernopycnon* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 94 [diagnosis].
 1969 *Anamnothea* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 97 [diagnosis].
 1977 *Ammothea*, - CLARK, J. Roy. Soc. N.Z., **7** (2): 171-176 [remarks, checklist and key to species].
 1983 *Ammothea*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 12 [remarks], 13 [key to Japanese species].
 1986 *Ammothea*, - KRAPF, Bonn. Zool. Beitr., **37** (4): 202 [tab. 1, comparison with *Achelia*].

Additional synonymy: *Leionymphon*.

Ammothea allopodes FRY & HEDGPETH 1969

- 1944 *Ammothea spinosa* var., - GORDON, Rep. Brit., Austr., and N.Z. Antarct. Res. Exped., (B), **5** (1): 50-51, figs. 16a-e, 17.
 1969 *Ammothea (Mathoma) allopodes* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 85-87, figs. 104, 105, 106-129.

Distribution: Antarctic and Subantarctic area.

Depth range: 474-540m.

Ammothea antipodensis CLARK 1972

- 1972 *Ammothea antipodensis* CLARK, N.Z. J. Mar. Freshw. Res., **5** (3-4): 435-438, fig. 4A-K.
 1977 *Ammothea antipodensis*, - CLARK, J. Roy. Soc. N.Z., **7** (2): 180-181, figs. 15-24.

Distribution: New Zealand.

Depth range: 0-24m.

Habitat: on red and brown algae, bryozoans.

Ammothea australiensis FLYNN 1919

- 1919 *Ammothea australiensis* FLYNN, Pap. Proc. Roy. Soc. Tasm., 1918: 95-99, pl. XIV, figs. 4-6.
 1940 *Achelia flynni*, - MARCUS, Bolm. Fac. Fil., Ciênc. Letr. Univ., S. Paulo, **19** (Zool. 4): 84, 124.
 1963 *Ammothea australiensis*, - CLARK, Rec. Austr. Mus., **26** (1): 57-59, fig. 29A-G, 59 [comparison with *A. magniceps*].
 1973 *Ammothea australiensis*, - STOCK, Beaufortia, **20** (266): 102.
 1975 *Ammothea australiensis*, - CHILD, Smiths. Contr. Zool., **190**: 24.
 1977 *Ammothea australiensis*, - CLARK, J. Roy. Soc. N.Z., **7** (2): 176-180, figs. 1, 3, 5, 7A, 9, 10-11, 13.

Distribution: Australia and New Zealand.

Depth range: 0-124m.

Habitat: red algae (*Gelidium*), under rocks, in tide pools.

Ammothea calmani GORDON 1932

- 1915 *Ammothea striata?* CALMAN, - Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 55-56.
 1932 *Ammothea calmani* GORDON, Discovery Rep., **6**: 104-106, figs. 54-55.
 1935 *Ammothea calmani*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 285.
 1969 *Ammothea (Mathoma) calmani*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 83-85, figs. 104, 105, 123-125.

Distribution: Antarctica.

Depth range: 99-549m.

Ammothea carolinensis LEACH 1814

- 1814 *Ammothea carolinensis* LEACH, Zool. Misc., **1**: 34, pl. 13.
 1840 *Nymphon carolinensis*, - H. MILNE-EDWARDS, Hist. Nat. Crust., etc. ..., 3. Ordre des Araneiformes ou Pycnogonides: ?126.
 1889 *Ammothea grandis* PFEFFER, Jb. Hamburg. Wiss. Anst., **6** (2): 43.
 1902 *Colossendeis gibbosa* MÖBIUS, Wiss. Ergeb. dt. Tiefsee-Exped. "Valdivia", 1898-1899, 3: 192, pl. XXX, figs. 1-5.
 1905 *Colossendeis (?) charcoti*, - BOUVIER, Bull. Mus. natn. Hist. nat., Paris, **11**: 296.
 1906 *Ammothea curculio* BOUVIER, - C.R. Séanc. Hebd. Acad. Sci., Paris, **142**: 19.
 1906 *Ammothea curculio*, - BOUVIER, L'Expéd. Antarct. Française, 1903-1905: 40.
 1906 *Leionymphon grande*, - BOUVIER, L'Expéd. Antarct. Française, 1903-1905: 40-48, 60-65, pl. III, fig. 6.
 1907 *Leionymphon gibbosum*, - HODGSON, Rep. natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 40.
 1907 *Leionymphon grande*, - HODGSON, Rep. Natn. Antarct. Expéd. 1901-1904, Nat. Hist., **3**: 41-43, pl. VI, fig. 1.
 1908 *Leionymphon grande*, - HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1) (6): 169-170.
 1908 *Ammothea carolinensis*, - LOMAN, Siboga Exped. Monogr., **40**: 10.
 1911 *Leionymphon grande*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 1140.
 1913 *Ammothea carolinensis*, - BOUVIER, Deuxième Exped. Antarct. Française (1908-1910), **6**: 127-131, figs. 78-82.
 1913 *Ammothea grandis*, - BOUVIER, Deuxième Exped. Antarct. Française (1908-1910), **6**: 126-127.
 1915 *Ammothea carolinensis*, - CALMAN, Ann. Mag. Nat. Hist., (9) **15**: 310-315, figs. 1-3.
 1915 *Ammothea carolinensis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool. **3** (1): 51.
 1915 *Ammothea carolinensis*, - HODGSON, Ann. Mag. Nat. Hist., (8) **15** (85): 146.
 1927 *Ammothea carolinensis*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 341-342 [key].
 1927 *Ammothea curculio*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 341.
 1927 *Ammothea gibbosa*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 341.
 1927 *Ammothea grandis*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 341.
 1932 *Ammothea carolinensis*, - GORDON, Discovery Rep., **6**: 95, 109.
 1935 *Leionymphon carolinensis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ord. Tierreichs, **5** (4) (2): 284.
 1938 *Ammothea gibbosa*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 20-21.

- 1944 *Ammothea gibbosa*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 52.
 1950 *Ammothea carolinensis*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 152.
 1950 *Ammothea gibbosa*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 152.
 1957 *Ammothea carolinensis*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., 55: 100.
 1969 *Ammothea (Ammothea) carolinensis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 73-75, figs. 104-108.

Distribution: W-Antarctic, from South Georgia southward.
Depth range: 10-640m.

Ammothea clausi PFEFFER 1889

- 1889 *Ammothea clausi* PFEFFER, Jb. Hamburg. Wiss. Anst., 6 (2): 45.
 1905 *Ammothea antarctica* BOUVIER, Bull. Mus. natn. Hist. nat., Paris, 11: 296.
 1906 *Ammothea antarctica*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, 142: 19.
 1907 *Leionymphon australe* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 46-49, pl. VII, fig. 1.
 1907 *Leionymphon clausi*, - HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 40.
 1908 *Leionymphon clausi*, - HODGSON, Trans. Roy. Soc. Edinburgh, 46 (1) (6): 160.
 1911 *Leionymphon clausi*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, 152: 1140.
 1913 *Ammothea australis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), 6: 123 [key].
 1913 *Ammothea clausi*, - BOUVIER, Deuxième Exped. Antarct. Française (1908-1910), 6: 135-138, figs. 88-89.
 1915 *Ammothea australis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool. 3 (1): 53.
 1923 *Ammothea australis*, - LOMAN, Further Zool. Res. Swed. Anatarct. Exped., 1 (2): 23.
 1927 *Ammothea australis*, - HODGSON, Dt. Südpolar-Exped., 19 (Zool. II): 341, 342 [key].
 1927 *Ammothea clausi*, - HODGSON, Dt. Südpolar-Expéd., 19 (Zool. II): 342, 343.
 1932 *Ammothea clausi*, - GORDON, Discovery Rep., 6: 109-110.
 1935 *Leionymphon clausi*, - HELFER & SCHLÖTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 284.
 1938 *Ammothea australis*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot), 2 (8): 20.
 1950 *Ammothea clausi*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 152.
 1969 *Ammothea (Theammoa) clausi*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 77.

Distribution: Antarctic (apparently common in the South Georgia-Bellingshausen Sea area).

Depth range: 3-860m.

Ammothea depolaris STOCK 1966

- 1966 *Ammothea depolaris* STOCK, Ann. Inst. Océanogr. Monaco, 44: 386-388, fig. 1.

Distribution: South of the Rio de la Plata, Argentina.

Depth range: 220–270m.

Ammothea dorsiplicata (Hilton, 1942)

1992 *Ammothea dorsiplicata*, - CHILD, Smiths. Contr. Zool., **526**: 4–6, fig. 1.

Distribution: Chile (NW of Valdivia).

Depth range: 1238m.

Synonymy: *Leionymphon dorsiplicata*.

Ammothea gigantea GORDON 1932

1932 *Ammothea gigantea* GORDON, Discovery Rep., **6**: 97–99, figs. 46, 47a.

1935 *Ammothea gigantea*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5**, Abt. IV, 2 (1): 285.

1969 *Magnammothea gigantea*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 90–91, figs. 104, 105, 132, 137, 138.

Distribution: Antarctica.

Depth range: 99–1115m.

Ammothea glacialis (HODGSON 1907)

1907 *Leionymphon glaciale* HODGSON, Rep. Natn. Antarct. Exped. 1901–1904, Nat. Hist., **3**: 50, pl. 7, fig. 3.

1913 *Ammothea glacialis*, - BOUVIER, Deuxième Exped. Antarct. Française (1908–1910), **6**: 123 [key].

1915 *Ammothea gibbosa*, - CALMAN, Ann. Mag. nat. Hist., **8** (15): 51–52 [part.].

1915 *Ammothea glacialis*, - CALMAN, Ann. Mag. nat. Hist., **8** (15): 50–51.

1927 *Ammothea glacialis*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 342–343.

1932 *Ammothea glacialis*, - GORDON, Discovery Rep., **6**: 96 [key].

1935 *Leionymphon glaciale*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 234.

1938 *Ammothea glacialis*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 20–21.

1944 *Ammothea glacialis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 51–52.

1950 *Ammothea gibbosa*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 152.

1950 *Ammothea glacialis*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 152.

1952 *Ammothea glacialis*, - FAGE, Bull. Mus. natn. Hist. Paris, (2) **24** (3): 271.

1969 *Ammothea (Ammothea) glacialis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 75–77, figs. 104, 105, 109–111.

Distribution: Antarctica.

Depth range: 163–640m.

Ammothea hedgpethi (UTINOMI 1959)

1949 *Lecythorhynchus* sp., - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296, figs. 44e–f.

- 1959 *Lecytorhynchus hedgpethi* UTINOMI, Publ. Seto. Mar. Biol. Lab., **7** (2): 212-216, figs. 7-9.
- 1962 *Lecytorhynchus hedgpethi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 103.
- 1971 *Lecytorhynchus hedgpethi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 336.
- 1977 *Ammothea hedgpethi*, - CLARK, J. Roy. Soc. N.Z., **7** (2): 173, 176 [key].
- 1983 *Ammothea hedgpethi*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 13 [key].
- 1987 *Ammothea hedgpethi*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 142-143, fig. 2.
- 1987 *Ammothea hedgpethi*, - NAKAMURA, Biol. Lab., Imperial Household: 23-24, pls. 19-20.
- 1991 *Ammothea hedgpethi*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 6.

Distribution: Japan.

Depth range: 8-95 m.

***Ammothea hilgendorfi* (BÖHM 1879)**

- 1879 *Corniger Hilgendorfi* BÖHM, Monatsber. Kgl. Preuss. Akad. Wiss. Berlin, 1879: 187-189, pl. 2 fig. 3-3d.
- 1879 *Lecytorhynchus Hilgendorfi*, - BÖHM, Sitz.-Ber. naturforsch. Fr. Berlin, 1879 (9): 140.
- 1881 *Lecytorhynchus hilgendorfi*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 27.
- 1899 *Lecytorhynchus Hilgendorfi*, - SHIGIDO, Dôzatsu, **11** (128): 199.
- 1904 *Lecytorhynchus marginatus* COLE, Harriman Alaska Exped., **10**: 260, pl. 11, figs. 1-2, pl. 15, figs. 1-8.
- 1911 *Lecytorhynchus Hilgendorfi*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-naturwiss. Kl.), Suppl. **2** (4): 8, pl. 2, figs. 28-29.
- 1913 *Lecytorhynchus marginatus*, - SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **18**: 240.
- 1915 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **7** (1): 68.
- 1951 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **7** (3): 204.
- 1919 *Lecytorhynchus Hilgendorfi*, - FUKUI, Rigakukai[?], **16** (2): 99.
- 1920 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **12** (4): 93.
- 1927 *Lecytorhynchus hilgendorfi*, - OHSHIMA, Proc. Imp. Acad., Tokyo, **3** (9): 610, figs. 1-3.
- 1929 *Lecytorhynchus hilgendorfi*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 54, figs. 10-11.
- 1929 *Lecytorhynchus marginatus*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 50, figs. 7-9.
- 1933 *Lecytorhynchus marginatus*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 61-62.
- 1934 *Lecytorhynchus marginatus*, - SCHMITT, J. Wash. Acad. Sci., **24** (1): 69.
- 1936 *Lecytorhynchus hilgendorfi*, - LOU, Contr. Inst. Zool., Natn. Acad. Peiping, **3** (5): 133, figs. 1-4, 5A, pls. 11-13.
- 1938 *Lecytorhynchus marginatus*, - HELFER, Sitz.-Ber. Ges. naturforsch. Fr. Berlin, 1937 (2): 184.
- 1938 *Lecytorhynchus hilgendorfi*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 867.
- 1939 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 34.

- 1940 *Lecytorhynchus marginatus*, - HEDGPETH, J. Wash. Acad. Sci., **30** (2): 86.
 1941 *Lecytorhynchus marginatus*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 255 [key].
 1942 *Lecytorhynchus ovatus* HILTON, Occ. Pap. Bernice P. Bishop Mus., **17** (3): 54, fig. 10a-b.
 1943 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **35** (1): 4.
 1947 *Lecytorhynchus hilgendorfi*, - OHSHIMA & KISHIDA, Illustr. Encyclopaedia Fauna Japan: 1009, fig. 2864.
 1949 *Lecytorhynchus hilgendorfi*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296, fig. 44a-b.
 1949 *Lecytorhynchus marginatus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296.
 1949 *Ammothea ovata*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296.
 1949 *Lecytorhynchus ovatus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296, fig. 44a-b.
 1951 *Lecytorhynchus marginatus*, - HEDGPETH, Wasmann J. Biol., **9** (1): 110.
 1951 *Lecytorhynchus hilgendorfi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 166.
 1954 *Lecytorhynchus marginatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 139.
 1956 *Lecytorhynchus marginatum*, - STOCK, Mitt. Zor): 124-127, fig. 1.

Distribution: South Africa, Agulhas Bank to Transkei coast.
Depth range: 90-900m.

***Ammothella spinifera* COLE 1904**

- 1904 *Ammothella spinifera* COLE, - Harriman Alaska Exped., **10**: 275-277, pl. 12. fig. 8, pl. 20, figs. 7-9, pl. 21, figs. 1-6.
 1913 *Ammothella spinifera*, - HALL, Univ. Calif. Publ. Zool., **11** (6): 132. [key].
 1941 *Ammothella spinifera*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 256 [key].
 1942 *Ammothella spinifera*, - HILTON, J. Ent. Zool. Pomona College, **34** (4): 97.
 1954 *Ammothella spinifera*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., 116, 119 [key].
 1955 *Ammothella spinifera*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 253.
 1979 *Ammothella spinifera*, - CHILD, Smiths. Contr. Zool., **293**: 11; Literatur.
 1986 *Ammothella spinifera*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
 1990 *Ammothella spinifera*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 278.
 1992 *Ammothella spinifera*, - CHILD, Smiths. Contr. Zool., **526**: 9-11.
 1992 *Ammothella spinifera*, - Müller, Stud. Nat. Hist. Caribb. Reg., 71: 43-44, figs. 2-3.
 1992 *Ammothella spinifera*, - STOCK, Tijdschr. Ent., **135**: 116, 139.

Distribution: California to Pacific Coast of Panamá; Ecuador; Caribbean (Martinique; Caribbean Coast of Panamá); Brazil.

Depth range: 0-13m.

Habitat: in a variety of substrates.

***Ammothella stauromata* CHILD 1982**

- 1982 *Ammothella stauromata* CHILD, Proc. biol. Soc. Wash., **95** (2): 271-273, fig. 1.
- 1988 *Ammothella stauromata*, - CHILD, Smiths. Contr. Zool., **468**: 5-6.
- 1988 *Ammothella stauromata*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 809.
- 1990 *Ammothella stauromata*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 316.
- 1989 *Ammothella stauromata*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 125.
- 1990 *Ammothella stauromata*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 66.
- 1990 *Ammothella stauromata*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierkunde Dresden, **45** (10): 106.

Distribution: Marshall Islands; Samoa; Philippines; Society Islands (Moorea); Fiji Islands.

Depth range: 0-3m.

Habitat: on pilings; dead corals on reef; rubble.

***Ammothella stocki* CLARK 1963**

- 1963 *Ammothella stocki* CLARK, Rec. Austr. Mus., **26** (1): 66-68, fig. 32A-J.

Distribution: Australia (Port Jackson).

Depth range: shallow water.

Habitat: "amongst marine growth on oregon test piece of timber" (CLARK).

***Ammothella symbius* CHILD 1979**

- 1979 *Ammothella symbius*, - Smiths. Contr. Zool., **293**: 12-15, fig. 4.

Distribution: Pacific Coast of Costa Rica and Mexico.

Depth range: shallow water to littoral depths.

Habitat: *Abietinaria*, *Pocillopora*.

***Ammothella thetidis* CLARK 1963**

- 1963 *Ammothella thetidis* CLARK, Rec. Austr. Mus., **26** (1): 68-70, figs. 33A-G, 34A-F.

Distribution: Australia.

Depth range: 40-115m.

***Ammothella tippula* CHILD 1983**

- 1983 *Ammothella tippula* CHILD, Proc. biol. Soc. Wash., **96** (4): 701-705, fig. 2.
- 1988 *Ammothella tippula*, - CHILD, Smiths. Contr. Zool., **468**: 7.
- 1988 *Ammothella tippula*, - CHILD, Bull. biol. Soc. Wash., **8**: 51-52.
- 1990 *Ammothella tippula*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 67.
- 1990 *Ammothella tippula*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (9): 97.
- 1990 *Ammothella tippula*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45**

(10): 106.

1991 *Ammothella tippula*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 138-139.

Distribution: Guam; Philippines; Seychelles (Aldabra Atoll); La Réunion; Sri Lanka.

Depth range: 0-11m.

Habitat: coral reefs. - rubble; dead corals; *Halimeda*; algal mats; coralline algae; dead barnacles covered with algae.

***Ammothella tuberculata* COLE 1904**

1904 *Ammothella tuberculata* COLE, Harriman Alaska Exped., **10**: 273-275, pl. XII, fig. 7, pl. XX, figs. 1-6.

1951 *Ammothella tuberculata*, - HEDGPETH, Wasmann J. Biol., **9** (1): 108.

Distribution: California.

***Ammothella tubicen* STOCK 1978**

1978 *Ammothella tubicen* STOCK, Cah. Biol. Mar., **19**: 194-197, fig. 3.

Distribution: N-Atlantic (Bay of Biscay).

Depth range: 1980-1995m.

***Ammothella uniunguiculata* (DOHRN 1881)**

1968 *Ammothella uniunguiculata*, - STOCK, Vie Milieu, **19** (1A): 13, fig. 5.

1973 *Ammothella uniunguiculata*, - KRAPP, Beaufortia, **21** (277): 59, fig. 1.

1976 *Ammothella uniunguiculata*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.

1979 *Ammothella uniunguiculata*, - CHIMENEZ-GUSSO, BRIGNOLI & BASCIANO, Cah. Biol. Mar., **20**: 484.

1980 *Ammothella uniunguiculata*, - MUNILLA-LEON, Cah. Biol. Mar., **21**: 2.

1981 *Ammothella uniunguiculata*, - MUNILLA & de HARO, Bijdr. Dierk., **51** (2): 191-197, figs. 4, 11, tab. I (electrophoretical and immunological study).

1984 *Ammothella uniunguiculata*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: tab. III-IV.

1987 *Ammothella uniunguiculata*, - ARNAUD, Mesogee, **47**: 41.

Distribution: E-Atlantic; Mediterranean.

Depth range: ?-11-16m.

Habitat: *Halopteris scoparia*; *Cystoseira*; *Posidonia*; *Corallina*; *Ulva*.

***Ammothella vanninii* STOCK 1982**

1982 *Ammothella vanninii* STOCK, Monit. zool. Ital., **17** (7): 183-187, figs. 1-7.

Distribution: Red Sea (Egypt; Israel); Somalia; Tanzania.

Depth range: shallow water

Habitat: on *Pocillopora* corals.

***Ascorhynchus* SARS 1877**

- 1948 *Ascorhynchus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 253-255 [diagnosis, key to NW-Atlantic species].
 1953 *Ascorhynchus*, - STOCK, Temminckia, **9**: 304-305 [key to species].
 1969 *Ascorhynchus*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 123 [diagnosis].
 1979 *Ascorhynchus*, - CHILD, Smiths. Contr. Zool., **293**: 15 [key to Middle American species].
 1983 *Ascorhynchus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 19 [key to Japanese species].
 1986 *Ascorhynchus*, - STOCK, Bull. mar. Sci., **38** (3): 412.
 1987 *Ascorhynchus*, - CHILD, Proc. biol. Soc. Wash., **100** (4): 906-907 [key to the giant species].
 1992 *Ascorhynchus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 15 [diagnosis].

***Ascorhynchus abyssi* SARS 1877**

- 1877 *Ascorhynchus abyssi* SARS, Arch. Math. Naturvidensk., Kristiania, **2**: 367.
 1888 *Ascorhynchus abyssi*, - SARS, Arch. Math. Naturvidensk., Oslo: **12**: 256.
 1891 *Ascorhynchus abyssi*, - SARS, Norw. North-Atl. Exp., 1876-1878, **6** (Zool. 20): 133, pl. XIV, fig. 2a-t.
 1898 *Ascorhynchus tridens* MEINERT, Danske Ingolf-Exped., **3** (1): 52-53, pl. V, figs. 7-18.
 1899 *Ascorhynchus tridens*, - MEINERT, Danish Ingolf-Exped., **3** (1): 55-56, pl. V, figs. 7-18.
 1901 *Ascorhynchus tridens*, - A catalogue of Crustacea and of Pycnogonida contained in the Museum of University College, Dundee: 55.
 1901 *Ascorhynchus abyssi*, - MÖBIUS, Fauna Arctica, **2** (1): 54.
 1901 *Ascorhynchus tridens*, - MÖBIUS, Fauna Arctica, **2** (1): 54.
 1903 *Ascorhynchus abyssi*, - LÖHNBERG, Kungl. Svenska Vetensk. Akad. Handl., **10**: 358.
 1908 *Ascorhynchus abyssi*, - NORMAN, J. Linn. Soc. Lond. (Zool.), **30**: 227.
 1908 *Ascorhynchus tridens*, - NORMAN, J. Linn. Soc. Lond. (Zool.), **30**: 227.
 1909 *Ascorhynchus abyssi*, - d'ARCY THOMPSON, Cambr. Nat. Hist., 274-275, fig. 266A-B.
 1912 *Ascorhynchus abyssi*, - STEPHENSEN, Danmark Exp. Grönl. Nordostkyst., 1906-1908, **5** (11): 564-570.
 1912 *Ascorhynchus castelli*, - LOMAN, Bull. Inst. Océanogr., Monaco, **238**: 8.
 1917 *Ascorhynchus abyssi* var. *apicalis* BOUVIER, Res. Campagnes Sci. accomplies sur son yacht par Albert Ier Prince Souverain de Monaco, **51**: 37-38, pl. IV, fig. 8.
 1923 *Ascorhynchus abyssi*, - BOUVIER, Faune de France, **7**: 49, fig. 45.
 1930 *Ascorhynchus abyssi*, - SCHIMKEWITSCH, FAUNA SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 100, 102, 106.
 1956 *Ascorhynchus abyssi*, - NOGUEIRA, Arq. Mus. Bocage, **27** (1): 92-93.
 1964 *Ascorhynchus abyssi*, - LOSINA-LOSINSKY, Trudy Arch. Antarkt. Nauchno-Issledov. Inst., Leningrad, **259**: 334-336, fig. 1.
 1984 *Ascorhynchus castelli*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 534-535, tab. III-IV.
 1984 *Ascorhynchus abyssi*, - STOCK, J. mar. biol. Ass. U.K.: 745.

Distribution: NE-Atlantic, Arctica.

Depth range: 906-2396m.

Habitat: clay; mud.

Ascorhynchus agassizi SCHIMKEWITSCH 1893

- 1893 *Ascorhynchus agassizii* SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard **25** (2): 36-39, pl. I, figs. 4-6, pl. II, figs. 12-13, 16-17.
 1909 *Ascorhynchus agassizi*, - COLE. Bull. Mus. Comp. Zool., **52** (11): 185.

Distribution: E-Pacific (Panamá).

Depth range: 1018-1416m.

Habitat: green sand, large globules.

Ascorhynchus antipodus CHILD 1987

- 1987 *Ascorhynchus antipodus* CHILD, Proc. biol. Soc. Wash., **100** (4): 902-904, fig. 1.

Distribution: SW-Pacific, off Antipodes Islands.

Depth range: 5340m.

Ascorhynchus arenicola (DOHRN 1881)

- 1881 *Barana arenicola* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 129-133, pl. II, figs. 2-8.
 1986 *Ascorhynchus arenicola*, - CHIMENTZ & COTTARELLI, Oebalia, 13, N.S.: 138.
 1987 *Ascorhynchus arenicola*, - ARNAUD, Mesogee, **47**: 41.

Distribution: Mediterranean Sea.

Depth range: 5-20m.

Habitat: "charactéristique de fonds détritiques sableux" (ARNAUD); is found on sandy and sandy-muddy bottom.

Ascorhynchus armatus (WILSON 1881)

- 1881 *Scaeorrhynchus armatus* WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 248-249, pl. 2, figs. 3-4, pl. 5, figs. 26-31.
 1937 *Ascorhynchus agassizi*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10) **20** (1): 38.
 1943 *Ascorhynchus armatus*, - HEDGPETH, Proc. New England Zool. Club, **22**: 49-50.
 1943 *Ascorhynchus armatus*, - NEEDLER, Can. Atl. Fauna, **10**: 15, fig. 19a-e.
 1948 *Ascorhynchus armatus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 255, figs. 42a, 43.
 1975 *Ascorhynchus armatus*, - STOCK, Bull. mar. Sci., **24** (4): 965-966, fig. 3.
 1990 *Ascorhynchus armatus*, - STOCK, Zool. Meded., **63** (16): 209.

Distribution: NW-Atlantic, including Caribbean; Sénégal.

Depth range: 1600-2300m.

Ascorhynchus athernum CHILD 1982

1982 *Ascorhynchus athernum* CHILD, Smiths. Contr. Zool., **349**: 4-6, fig. 1.

Distribution: North American Basin.

Depth range: 2862m.

***Ascorhynchus auchenicum* (SLATER 1879)**

- 1879 *Parazetes auchenicus* SLATER, Ann. Mag. Nat. Hist., (5) **3** (16): 281-283.
 1881 *Parazetes auchenicus*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 26.
 1881 *Ascorhynchus minutus* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 55-57, pl. VI, figs. 10-16.
 1890 *Ascorhynchus bicornis* ORTMANN, Zool. Jb. (Syst.), **5** (1): 162.
 1890 *Ascorhynchus glabroides* ORTMANN, Zool. Jb. Syst., **5** (1): 160-161, pl. 24 figs. 3a-b.
 1899 *Parazetes auchenicus*, - SHISHIDO, Dôzatsu, **11** (128): 199.
 1911 *Ascorhynchus glabroides*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturw. Klasse), Suppl. **2** (4): 7
 1922 *Ascorhynchus auchenicus*, - CALMAN, Ann. Mag. Nat. Hist., **9** (9): 199-203, figs. 1-4.
 1929 *Ascorhynchus minutus*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 111, figs. 26-27.
 1936 *Ascorhynchus auchenicus*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 864.
 1949 *Ascorhynchus auchenicus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 291.
 1949 *Ascorhynchus glabroides*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 291-292.
 1951 *Ascorhynchus glabroides*, - UTINOMI, Publ. Seto. Mar. Biol. Lab., **1** (4): 166.
 1953 *Ascorhynchus auchenicum*, - STOCK, Temminckia, **9**: 304 [key].
 1954 *Ascorhynchus auchenicus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 124-126, figs. 59-60.
 1955 *Ascorhynchus glabroides*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 25.
 1959 *Ascorhynchus auchenicum*, - UTINOMI, - Publ. Seto Mar. Biol. Lab., **7** (2): 206.
 1965 *Ascorhynchus auchenicus*, - UTINOMI, New Illustrated Encyclopaedia of the Fauna of Japan, 2: 337, fig. 27.
 1971 *Ascorhynchus auchenicum*, - Publ. Seto Mar. Biol. Lab., **18** (5): 332.
 1971 *Ascorhynchus glabroides*, - Publ. Seto Mar. Biol. Lab., **18** (5): 333.
 1983 *Ascorhynchus auchenicum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 20.
 1987 *Ascorhynchus auchenicum*, - NAKAMURA, Biol. Lab. Imperial Household: 27-28, pls. 24, 37.

Distribution: apparently endemic to Japan.

Depth range: 18-550m.

Habitat: hard bottom; gravel; sand.

***Ascorhynchus bacescui* STOCK 1975**

- 1975 *Ascorhynchus bacescui* STOCK, Trav. Mus. Hist. nat. "Grigore Antipa", **16**: 128-131, figs. 1-9.

Distribution: E-Africa.

Ascorhynchus birsteini Turpaeva, 1971

- 1971 *Ascorhynchus birsteini* TURPAEVA, Byull. Moskovsk. obsh. Ispyt. Prirod., Otd. Biol., **76** (3): 107-109, fig. 2.
 1992 *Ascorhynchus birsteini*, - CHILD, Smiths. Contr. Zool., **526**: 11.

Distribution: northern Parts of the Peru-Chile Trench.
Depth range: 5986-6260m.

Ascorhynchus breviscapus STOCK 1968

- 1968 *Ascorhynchus breviscapus* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 9-10, fig. 1.
 1991 *Ascorhynchus breviscapus*, - STOCK, Mém. Mus. natn. Hist. nat., (A), **151**: 127-128, fig. 1.

Distribution: South Africa, New Caledonia.

Depth range: 825-1361m.

Habitat: hard bottom, grab.

Ascorhynchus bucerus TURPAEVA 1971

- 1971 *Ascorhynchus bucerus* TURPAEVA, Byull. Moskovsk. obsh. Ispyt. Prirod., Otd. Biol., **76** (3): 105-107, fig. 1.
 1991 *Ascorhynchus bucerus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 129 [text].

Distribution: Pacific Ocean (Kurile-Kamtchatka Trench).
Depth range: 5035-5210 m.

Ascorhynchus cactoides STOCK 1954

- 1954 *Ascorhynchus cactoides* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 132-135, figs. 63d-f, 66.
 1991 *Ascorhynchus cactoides*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 128-129, fig. 2bd-d.
 1991 *Ascorhynchus cactoides*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 128-129, fig. 2b-d.

Distribution: Kei Islands (Indonesia), New Caledonia, Philippines.

Depth range: 90-515.

Habitat: sand.

Ascorhynchus castelli (DOHRN 1881)

- 1881 *Barana castelli* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 123-129, pl. I, figs. 1-16, pl. II, fig. 1.
 1891 *Barana castelnaudi* ORTMANN, Zool. Jb. (Syst.), **5** (1): 159.
 1908 *Barana castelli*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 227.
 1909 *Barana castelli*, - d'ARCY THOMPSON, in: The Cambridge Nat. Hist.: 533.

- 1912 *Ascorhynchus castelli*, - LOMAN, Bull. Inst. Océanogr. Monaco, **238**: 8.
 1917 *Ascorhynchus castelli*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 5, 6.
 1923 *Ascorhynchus castelli*, - BOUVIER, Faune de France, **7**: 49, fig. 46.
 1929 *Ascorhynchus castelli*, - GILTAY, Bull. & Ann. Soc. Ent. Belg., **69**: 174-175.
 1966 *Ascorhynchus castelli*, - de HARO, Invest. Pesqu., **30**: 665.
 1966 *Ascorhynchus castelli*, - de HARO, Bol. Real. Soc. Espan. Hist. Nat., Madrid (Secc. Biol.), **64**: 8, 10.
 1966 *Ascorhynchus castelli*, - SOYER, Doriane, **47** (174): 2.
 1968 *Ascorhynchus castelli*, - LEDOYER, Rec. Trav. Stat. Mar. d'Endoume, **44** (60): 170.
 1966 *Ascorhynchus castelli*, - STOCK, Vie Milieu, **17** (1B): 407.
 1968 *Ascorhynchus castelli*, - STOCK, Vie Milieu, **19** (1A): 10-11, fig. 2.
 1975 *Ascorhynchus castelli*, - STOCK, Bull. mar. Sci., **24** (4): 966-968, fig. 4.
 1976 *Ascorhynchus castelli*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 340-342
 1978 *Ascorhynchus castelli*, - CHIMENTZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 98.
 1987 *Ascorhynchus castelli*, - ARNAUD, Mesogee, **47**: 41.

Distribution: Mediterranean and temperate-tropical N-Atlantic.

Depth range: upper sublittoral-130m.

Habitat: on a variety of substrata.

***Ascorhynchus castelliooides* STOCK 1957**

- 1957 *Ascorhynchus castelliooides* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 82-84, fig. 2.
 1970 *Ascorhynchus castelliooides*, - BAYER, VOSS & ROBINS, in: Rep. Mar. Fauna and Benthic Shelf-slope communities of the Isthmian Region; Univ. Miami: A48, A117.
 1975 *Ascorhynchus castelliooides*, - STOCK, Bull. mar. Sci., **24** (4): 968.
 1976 *Ascorhynchus castelliooides*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 133.
 1979 *Ascorhynchus castelliooides*, - CHILD, Smiths. Contr. Zool., **293**: 15.
 1986 *Ascorhynchus castelliooides*, - STOCK, Bull. mar. Sci., **38** (3): 400.
 1992 *Ascorhynchus castelliooides*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 45.

Distribution: W-Atlantik, Florida south to Venezuela.

Depth range: 0-30m.

Habitat: in a variety of substrata, often found in fouling of pilings; "Feeds probably on sponges" (STOCK 1975); red algae, sponges, and hydroids on wreck.

***Ascorhynchus colei* HEDGPETH 1943**

- 1943 *Ascorhynchus colei* HEDGPETH, Proc. New England Zool. Club, **22**: 50.
 1948 *Ascorhynchus colei*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 257-258, fig. 44g-k.
 1986 *Ascorhynchus colei*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1, 412.

Distribution: W-Atlantic (Florida, Bahamas, Dominican Republic).

Depth range: 0-293 m.

Ascorhynchus comatum CHILD 1992

1992 *Ascorhynchus comatum* CHILD, Smiths. Contr. Zool., **526**: 11-13, fig. 4.

Distribution: Ecuador.

Depth range: 1363-1369m.

Ascorhynchus compactum CLARK 1963

1963 *Ascorhynchus compactum* CLARK, Rec. Austr. Mus., **26** (1): 61-63, fig. 30A-H.

Distribution: Australia (Port Jackson).

Depth range: intertidal.

Habitat: on rocks.

Ascorhynchus cooki CHILD 1987

1987 *Ascorhynchus cooki* CHILD, Proc. biol. Soc. Wash., **100** (4): 904-906, fig. 2.

Distribution: SW-Pacific, SW of Macquarie Island; New Zealand, N of Chatham Rise, SE of Cook Strait.

Depth range: 2610-2992m.

Ascorhynchus corderoi du BOIS-REYMOND MARCUS 1952

1952 *Ascorhynchus corderoi* du BOIS-REYMOND MARCUS, An. Acad. Bras. Cien., **24** (1): 23-30, figs. 1-9.

1953 *Ascorhynchus corderoi*, - STOCK, Temminckia, **9**: 304.

1965 *Ascorhynchus corderoi*, - STOCK, Beaufortia, **13** (151): 14.

1988 *Ascorhynchus corderoi*, - CHILD, Bull. biol. Soc. Wash., **8**: 52.

1992 *Ascorhynchus corderoi*, - STOCK, Bijdr. Dierk., **62** (2): 82.

Distribution: Brazil (off Sao Paulo); Mauritius; Seychelles (Aldabra Atoll); Oman.

Depth range: shallow water.

Habitat: coral reefs. - in coral.

Ascorhynchus crenatum CHILD 1992

1992 *Ascorhynchus crenatum* CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 15-17, fig. 5, 19 [text], 76 [tab. 12], 78 [tab. 14].

Distribution: off central west Florida.

Depth range: 55-76 m.

Ascorhynchus cryptopygium ORTMANN 1890

- 1890 *Ascorhynchus cryptopygium* ORTMANN, Zool. Jb. (Syst.), **5** (1): 159-160, pl. XXIV fig. 2.
- 1899 *Ascorhynchus cryptopygium*, - SHISHIDO, Dôzatsu, **11** (128): 199.
- 1911 *Ascorhynchus cryptopygium*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss (Math.-Naturwiss. Kl.) Suppl. **2** (4): 7.
- 1918 *Ascorhynchus cryptopygium*, - FUKUI, Rigakukai(?), **16** (2): 98.
- 1936 *Ascorhynchus cryptopygium*, - OHSHIMA, Dôbutsgaku Zasshi, **48** (8-10): 865.
- 1953 *Ascorhynchus cryptopygium*, - STOCK, Temminckia, **9**: 304 [key].
- 1954 *Ascorhynchus cryptopygium*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 121.
- 1955 *Ascorhynchus cryptopygium*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 24-25, fig. 14.
- 1971 *Ascorhynchus cryptopygius*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 333.
- 1983 *Ascorhynchus cryptopygium*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 19 [key], 20.

Distribution: Japan.

Depth range: 75-274 m.

Habitat: hard and sand bottom.

***Ascorhynchus cuculum* FRY & HEDGPETH 1969**

- 1969 *Ascorhynchus cuculus* FRY & HEDGPETH, Fanua of the Ross Sea, 7. Mem. N.Z. Oceanogr. Inst., **49**: 123-124, figs. 203-204.
- 1974 *Ascorhynchus hedgpethi* TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **98**: 285-287, fig. 4.
- 1982 *Ascorhynchus cuculum*, - CHILD, Smiths. Contr. Zool., **349**: 6, fig. 2a-b.

Distribution: Scotia Sea Basin; Argentine Basin.

Depth range: 3475-3910m.

***Ascorhynchus discrepans* STOCK 1955**

- 1955 *Ascorhynchus discrepans* STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 257-260, figs. 22-23.
- 1986 *Ascorhynchus discrepans*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.

Distribution: off western coast of Thatch Island (near St. John); north of Congo Cay (northwest of St. John).

Depth range: 26-29m.

***Ascorhynchus endoparasiticus* ARNAUD 1978**

- 1978 *Ascorhynchus endoparasiticus* ARNAUD, Zool. J. Linn. Soc., **63**: 99-104, figs. 1-2.

Distribution: Azores.

Depth range: 1590-1655m.

Habitat: parasitic inside the pallial cavity of the opisthobranch mollusc (*Scaphander punctostriatus*).

***Ascorhynchus foresti* STOCK 1991**

1991 *Ascorhynchus foresti* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 214-216, figs. 1-2.

Distribution: Philippines.
Depth range: 92-97m.

***Ascorhynchus fragilis* STOCK 1991**

1991 *Ascorhynchus fragilis* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 129-131, figs. 3-4.

Distribution: New Caledonia.
Depth range: 2365-3700m.

***Ascorhynchus fusticulum* NAKAMURA & CHILD 1983**

1983 *Ascorhynchus fusticulum* NAKAMURA & CHILD, Siths. Contr. Zool., **386**: 20-24, fig. 6.

Distribution: Japan.
Depth range: 75-85m.

***Ascorhynchus glaber* HOEK 1881**

1881 *Ascorhynchus glaber* HOEK, Challenger Rep., London, **3** (10): 53-55, pl. VI, figs. 5-9, pl. XV, fig. 16

1956 *Ascorhynchus glaber*, - FAGE, Galathea Repths., **2**: 175.

1991 *Ascorhynchus glaber*, - TURPAEVA, Zool. Zh., **70** (12): 37.

Distribution: off Madagascar.
Depth range: 2500-2500m.

***Ascorhynchus glaberrimum* SCHIMKEWITSCH 1913**

1913 *Ascorhynchus glaberrimus* SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **18**: 242, pl. 3a figs. 8-14.

1930 *Ascorhynchus glaberrimus*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 107.

1949 *Ascorhynchus glaberrimus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 293.

1955 *Ascorhynchus glaberrimum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 26, fig. 15.

1959 *Ascorhynchus glaberrimum*, - UTINOMI, Publ. Seto. Mar. Biol. Lab., **7** (2): 208.

1962 *Ascorhynchus glaberrimum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 100.

1971 *Ascorhynchus glaberrimus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 334.

1983 *Ascorhynchus glaberrimum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 23-

24, fig. 7.

- 1986 *Ascorhynchus glaberrimum*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 48-50.
 1987 *Ascorhynchus glaberrimum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 146.
 1987 *Ascorhynchus glaberrimum*, - NAKAMURA, Biol. Lab., Imperial Household: 29-30,
 pls. 25-26.
 1991 *Ascorhynchus glaberrimum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 7-
 8.

Distribution: Japan and Korea.

Depth range: 0-230 m.

***Ascorhynchus horologium* CHILD 1992**

- 1992 *Ascorhynchus horologium* CHILD, Mem. Hourglass Cruises, **9** (1): 9 [key, tab.
 3], 17-20, fig. 6, 76 [tab. 12], 78 [tab. 14].

Distribution: off central west Florida.

Depth range: 73-90 m.

***Ascorhynchus inflatum* STOCK 1963**

- 1963 *Ascorhynchus inflatum* STOCK, Ann. S. Afr. Mus., **46** (12): 323-326, figs. 2-3.
 1971 *Ascorhynchus inflatum*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR,
92: 284-286, fig. 6.
 1992 *Ascorhynchus inflatum*, - CHILD, Smiths. Contr. Zool., **526**: 13.

Distribution: Cape Basin off South Africa; Kurile-Kamtchatka Trench; Scotia Sea; Peru-Chile Trench.

Depth range: 2743-6070m.

***Ascorhynchus insularum* CLARK 1971**

- 1971 *Ascorhynchus insularum* CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 335-336,
 fig. 2A-L.
 1977 *Ascorhynchus insularum*, - CLARK, N.Z. H. mar. Freshw. Res., **11** (1): 176.

Distribution: New Zealand (Snares Islands).

Depth range: lower littoral.

Habitat: on stalked barnacles; "on limpets incrusted with polychaete tubes and bryozoa, in mid-littoral crevices" (CLARK).

***Ascorhynchus japonicum* IVES 1891**

- 1891 *Ascorhynchus japonicus* IVES, Proc. Acad. Nat. Sci. Philadelphia, 1892-1893:
 219-221, pl. XII, figs. 4-10.
 1911 *Ascorhynchus japonicus*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-
 Naturwiss. Kl.), Suppl. **2** (4): 5.
 1919 *Ascorhynchus japonicus*, - FUKUI, Rigakukai(?), **16** (2): 98.
 1936 *Ascorhynchus japonicus*, - OHSIMA, Dōbutsugaku Zasshi, **48** (8-10): 865.
 1942 *Ascorhynchus agassizi*, - HILTON, J. Ent. Zool. Pomona College, **34**: 98.

- 1947 *Ascorhynchus japonicus*, - OHSIMA & KISHIDA, Illustrated Encyclopaedia of the Fauna of Japan: 1008, fig. 2859.
- 1949 *Ascorhynchus japonicus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 292-293.
- 1951 *Ascorhynchus japonicus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 166.
- 1953 *Ascorhynchus japonicum*, STOCK, Temminckia, **9**: 305 [key].
- 1954 *Ascorhynchus japonicum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 126-128, figs. 61-62.
- 1955 *Ascorhynchus japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 24.
- 1959 *Ascorhynchus japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 205.
- 1961 *Ascorhynchus japonicus*, - LOSINA-LOSINSKY, Issl. Dalnyevost. Morei SSSR, Leningrad, **7**: 107-108, fig. 25.
- 1971 *Ascorhynchus japonicus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 333.
- 1973 *Ascorhynchus japonicus*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **91**: 181.
- 1983 *Ascorhynchus japonicum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 19 [key].
- 1987 *Ascorhynchus japonicum*, - NAKAMURA, Biol. Lab. Imperial Household: 30-31, pls. 27, 38.

Distribution: NW-Pacific and ?Oregon coast of America.

Depth range: 53-1700m.

Habitat: sand bottom.

***Ascorhynchus laterospinum* HILTON 1942**

- 1942 *Ascorhynchus laterospinus* HILTON, J. Ent. Zool. Pomona College, **34** (4): 98.
- 1953 *Ascorhynchus laterospinum*, - STOCK, Temminckia, **9**: 304.
- 1971 *Ascorhynchus laterospinum*, - CHILD & HEDGPETH, J. nat. Hist., **5**: 628-631, fig. 8.

Distribution: Galápagos Archipelago.

Depth range: 1159m.

***Ascorhynchus latipes* (COLE 1906)**

- 1906 *Barana latipes* COLE, Am. Nat., **40** (471): 217-222, pls. 1-2.
- 1940 *Ascorhynchus latipes*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 93.
- 1948 *Ascorhynchus latipes*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 256-257, fig. 42b.
- 1952 *Ascorhynchus latipes*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (6): 530.
- 1953 *Ascorhynchus latipes*, - STOCK, Temminckia, **9**: 304 [key].
- 1954 *Ascorhynchus latipes*, - HEDGPETH, Fish. Bull. Fish & Wildlife Service U.S., **89** (55): 427.
- 1954 *Ascorhynchus latipes*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 116.
- 1975 *Ascorhynchus latipes*, - STOCK, Bull. mar. Sci., **24** (4): 969.
- 1976 *Ascorhynchus latipes*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 133.
- 1979 *Ascorhynchus latipes*, - CHILD, Smiths. Contr. Zool., **293**: 15; Literatur.
- 1979 *Ascorhynchus latipes*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 969.
- 1982 *Ascorhynchus latipes*, - CHILD, Smiths. Contr. mar. Sci., **12**: 359.

- 1986 *Ascorhynchus latipes*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
 1992 *Ascorhynchus latipes*, - CHILD, Mem. Hourglass Cruises, **9** (1): 9 [key, tab. 3], 21, fig. 7.

Distribution: amphi-atlantic; W-Atlantic from the Bahamas to the northern coast of South America; E-Atlantic from Senegal.

Depth range: 0-2m.

Habitat: in a variety of substrata.

***Ascorhynchus latum* CALMAN 1923**

- 1923 *Ascorhynchus latus* CALMAN, Rec. Indian Mus., **24** (3): 270-272, figs. 2-3.
 1954 *Ascorhynchus latum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 128, fig. 63a-c.
 1990 *Ascorhynchus latum*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (1): 106.

Distribution: Gulf of Manaar; Gulf of Siam; Thailand.

Depth range: 3-17m.

Habitat: brownish sand.

***Ascorhynchus levissimum* LOMAN 1908**

- 1908 *Ascorhynchus levissimus* LOMAN, Siboga Exp. Monogr., **40**: 33-34, pl. IV, figs. 46-51
 1953 *Ascorhynchus levissimus*, - STOCK, Temminckia, **9**: 304 [key].
 1990 *Ascorhynchus levissimus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **103** (2): 305, fig. 1A-D.

Distribution: Flores Sea.

Depth range: 630-794m.

***Ascorhynchus longicolle* (HASWELL 1884)**

- 1884 *Ascorhynchus longicollis* HASWELL, Proc. Linn. Soc. N.S.W. **9**: 1028-1029, pl. 56, figs. 1-4.
 1889 *Ascorhynchus longicollis*, - WHITELEGGE, J. Proc. Roy. Soc. N.S.W., **23** (2): 223.
 1908 *Ascorhynchus longicollis*, - LOMAN, Siboga Exp. Monogr., **40**: 32.
 1909 *Eurydyce longicollis*, - d' ARCY THOMPSON, Cambr. Nat. Hist., 533.
 1920 *Ascorhynchus longicollis*, - FLYNN, Pap. Proc. Roy. Soc. Tasmania, 1910: 81-83, pl. XXI, figs. 16-17.
 1963 *Ascorhynchus longicollis*, - CLARK, Rec. Austr. Mus., **26** (1): 60-61, fig. 29H-M.

Distribution: Australia.

Habitat: among seaweeds.

***Ascorhynchus losinalosinskii* TURPAEVA 1971**

1971 *Ascorhynchus losinalosinskii* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **92**: 280-281, fig. 5.

Distribution: Kurile-Kamtchatka Trench.
Depth range: 3145-3250m.

***Ascorhynchus mariae* TURPAEVA 1971**

1971 *Ascorhynchus mariae* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **92**: 280-281, fig. 4.

Distribution: Kurile-Kamtchatka Trench.
Depth range: 3145-3250m.

***Ascorhynchus melwardi* FLYNN 1929**

1929 *Ascorhynchus melwardi* FLYNN, Mem. Qd. Mus., **9** (3): 252-256, figs. 1-3.

1954 *Ascorhynchus melwardi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 128-132, figs. 64-65.

1968 *Ascorhynchus melwardi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 8.

Distribution: Kei Islands; Singapore.
Depth range: 0-30m.
Habitat: sand; shells.

***Ascorhynchus meteor* MÜLLER 1989**

1989 *Ascorhynchus meteor* MÜLLER, Senckenbergiana marit., **20** (5/6): 278-279, figs. 1-7.

Distribution: Gulf of Aden.
Depth range: 35-45m.

***Ascorhynchus minutum* HOEK 1881**

1881 *Ascorhynchus minutus* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 55-57, pl. VI, figs. 10-16.

1908 *Ascorhynchus minutus*, - LOMAN, Siboga Exp. Monogr., **40**: 33.

1922 *Ascorhynchus auchenicus*, - CALMAN, Ann. Mag. Nat. Hist., **9** (9): 199-203 [part.].

1953 *Ascorhynchus minutum*, - STOCK, Temminckia, **9**: 305.

1954 *Ascorhynchus minutum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 121-124, fig. 57d-h.

1963 *Ascorhynchus minutum*, - CLARK, Rec. Austr. Mus., **26** (1): 61.

1975 *Ascorhynchus minutum*, - CHILD, Smiths. Contr. Zool., **190**: 26.

Distribution: Indonesia; Australia (Queensland, New South Wales).

Depth range: 30-100m.

Habitat: sand, mud, silt.

Ascorhynchus okai NAKAMURA & CHILD 1983

1983 *Ascorhynchus okai* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 24-26, fig. 8.

Distribution: Japan.
Depth range: 30-85m.

Ascorhynchus ornatum (HELFER 1938)

1938 *Ainigma ornatum* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 181-183, fig. 10a-f.
 1953 *Ascorhynchus ornatum*, - STOCK, Beaufortia, **35** (4): 41-44, fig. 6a-g.
 1953 *Ascorhynchus ornatum*, - STOCK, Temminckia, **9**: 304 [key].
 1954 *Ainigma ornatum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 144, fig. 28B
 1988 *Ascorhynchus ornatum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 127.
 1989 *Ascorhynchus ornatum*, - MÜLLER, Senckenbergiana marit., **20** (5/6): 280-281, figs. 8-14.
 1991 *Ascorhynchus ornatus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 131, fig. 2a.

Distribution: South Africa, New Caledonia.
Depth range: 90-708m.

Ascorhynchus orthorhynchus HOEK 1881

1881 *Ascorhynchus orthorhynchus* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 57-59, pl. V, figs. 11-13, pl. Vi, figs. 1-4, pl. XV, figs. 14-15.
 1908 *Ascorhynchus orthorhynchus*, - LOMAN, Siboga Exp. Monogr., **40**: 31.
 1953 *Ascorhynchus orthorhynchus*, - STOCK, Temminckia, **9**: 301.

Distribution: East Indies.
Depth range: 439m.

Ascorhynchus ovicoxa STOCK 1975

1975 *Ascorhynchus ovicoxa* STOCK, Bull. mar. Sci., **24** (4): 969-972, figs. 5-6.
 1982 *Ascorhynchus ovicoxa*, - CHILD, Smiths. Contr. Zool., **349**: 6.
 1986 *Ascorhynchus ovicoxa*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1, 412.

Distribution: W-Atlantic (Florida; North American Basin, north of Bermuda).
Depth range: 333-2223m.

Ascorhynchus pararmatus STOCK 1975

1943 *Ascorhynchus pararmatus*, - HEDGPETH, Proc. New England Zool. Club., **22**: 49-50.
 1975 *Ascorhynchus pararmatus* STOCK, Bull. mar. Sci., **24** (4): 960-966, figs. 1-2.

1986 *Ascorhynchus pararmatus*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1, 412.

Distribution: Bahamas to Florida.

Depth range: 101-2195m.

Ascorhynchus parvituberculatum STOCK 1953

1953 *Ascorhynchus parvituberculatum* STOCK, Temminckia, **9**: 301-304, fig. 15.

Distribution: Indonesia (near Timor).

Depth range: 10-15m.

Ascorhynchus paxillum CHILD 1992

1992 *Ascorhynchus paxillum* CHILD, Smiths. Contr. Zool., **526**: 13-15, fig. 5.

Distribution: Costa Rica (Cocos Ridge).

Depth range: 1892m.

Ascorhynchus pennai (MELLO-LEITAO 1946)

1946 *Acantammothella pennai* MELLO-LEITAO, An. Acad. Bras. Cien., **18** (4): 292-296, figs. 1-4.

Distribution: Brazil (Baia de Guanabara).

Ascorhynchus pilipes STOCK 1991

1991 *Ascorhynchus pilipes* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 131-134, figs. 5-6.

Distribution: New Caledonia.

Depth range: 1600-1640m.

Ascorhynchus prosum NAKAMURA & CHILD 1983

1983 *Ascorhynchus prosum* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 26-29, fig. 9.

Distribution: Japan.

Depth range: 30m.

Ascorhynchus pudicum STOCK 1970

1970 *Ascorhynchus pudicum* STOCK, "Meteor"-Forsch.-Ergebn., D, 5: 6-7, figs. 1-7.

1991 *Ascorhynchus pudicus*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 137.

Distribution: NW-Africa (Josephine Bank); NE-Atlantic.

Depth range: 205-310m.

Ascorhynchus pyriginosum McCLOSKEY 1967

- 1967 *Ascorhynchus pyriginosum* McCLOSKEY, J. nat. Hist., **1**: 125-128, figs. 12-17.
 1967 *Nanphon grasslei* McCLOSKEY, J. nat. Hist., **1**: 123-125, figs. 7-11.
 1973 *Ascorhynchus pyriginosum*, - KRAEUTER, J. nat. Hist., **7**: 495.
 1976 *Ascorhynchus pyriginosum*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 342.
 1982 *Ascorhynchus pyriginosum*, - CHILD, Smiths. Contr. Zool., **349**: 6-8.
 1992 *Ascorhynchus pyriginosum*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 21-24, fig. 8.

Distribution: Gulf of Alabama and Florida, off Georgia and North Carolina, New England continental slope.

Depth range: 32-811m.

Ascorhynchus ramipes (BÖHM 1879)

- 1879 *Gnaptorhynchus ramipes* BÖHM, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1879 (4): 56-59, fig. 1.
 1881 *Ascorhynchus ramipes*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 25.
 1890 *Ascorhynchus ramipes*, - ORTMANN, Zool. Jb. (Syst.), **5** (1): 161-162, pl. 24.
 1899 *Ascorhynchus ramipes*, - SHISHIDO, Dôzatsu, **11** (128): 199.
 1911 *Ascorhynchus ramipes*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 6 [part.].
 1923 *Ascorhynchus latus*, - CALMAN, Rec. Indian Mus., **25** (3), 270.
 1936 *Ascorhynchus ramipes* var. *tsingtaensis*, - LOU, Contr. Inst. Zool., Natn. Acad. Peiping, **3** (5): 2.
 1936 *Ascorhynchus ramipes*, OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 865.
 1936 *Ascorhynchus ramipes* var. *tsingtaensis*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 865.
 1949 *Ascorhynchus ramipes*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 292.
 1953 *Ascorhynchus latum*, - STOCK, Temminckia, **9**: 304.
 1954 *Ascorhynchus latum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 128.
 1959 *Ascorhynchus ramipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 207, fig. 4B.
 1962 *Ascorhynchus ramipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 99.
 1971 *Ascorhynchus ramipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 332-333.
 1983 *Ascorhynchus ramipes*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 29.
 1983 *Ascorhynchus ramipes*, - NAKAMURA & SEKIGUCHI, Bull. Biogeogr. Soc. Japan, **38** (6): 39.
 1986 *Ascorhynchus ramipes*, - KIM, Proc. Nat. Sci. Res. Inst. KANU, **2** (1): 6.
 1986 *Ascorhynchus ramipes*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 50.
 1987 *Ascorhynchus ramipes*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 146.
 1987 *Ascorhynchus ramipes*, - NAKAMURA, Biol. Lab., Imperial Household: 31-32, pl. 28.
 1991 *Ascorhynchus ramipes*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 8.

Distribution: Japan to Gulf of Thailand and Gulf of Manaar,

India.

Depth range: 10-200m.

Ascorhynchus serratum HEDGPETH 1948

- 1948 *Ascorhynchus serratum* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 259-260, fig. 44a-f.
 1975 *Ascorhynchus serratum*, - STOCK, Bull. mar. Sci., **24** (4): 969.
 1982 *Ascorhynchus* sp. cf. *serratum*, - CHILD, Smiths. Contr. mar. Sci., **12**: 359.
 1986 *Ascorhynchus serratus*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.
 1992 *Ascorhynchus serratum?*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 24, fig. 976 [tab. 12], 78 [tab. 14].

Distribution: Yucatan Channel of Mexico, off Belize, Northwest Providence Channel in the Bahamas.

Depth range: 27-711m.

Habitat: ?in coral and algae on outer reef slope.

Ascorhynchus seticauda STOCK 1991

- 1991 *Ascorhynchus seticauda* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 134-137, figs. 7-8.

Distribution: New Caledonia (Chesterfield Islands).

Depth range: 600m.

Ascorhynchus simile FAGE 1942

- 1942 *Ascorhynchus similis* FAGE, - Arch. Zool. Exp. Gén., **82**: 85-86, fig. 7.
 1957 *Ascorhynchus simile*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 81.
 1965 *Ascorhynchus simile*, - STOCK & SOYER, Vie et Milieu, **16** (1B): 416-417.
 1968 *Ascorhynchus simile*, - STOCK, Vie et Milieu, **19** (1A), Suppl.: 11, fig. 3.
 1987 *Ascorhynchus simile*, - ARNAUD, Mesogee, **47**: 42.

Distribution: W-Africa; Mediterranean.

Depth range: 5-100m.

Habitat: *Cymodocea truncata*; *Halarachnion spatulatum* (red alga).

Ascorhynchus simplex NAKAMURA & CHILD 1991

- 1991 *Ascorhynchus simplex* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 8-10, fig. 2.

Distribution: Japan.

Depth range: 110-122m.

Ascorhynchus stocki HONG & KIM 1987

- 1987 *Ascorhynchus stocki* HONG & KIM, Kor. J. Syst. Zool., **3** (2): 146-149, figs.

6-7.

Distribution: Korea.

Habitat: mud bottom.

Ascorhynchus tenuirostris CARPENTER 1892

- 1892 *Ascorhynchus tenuirostris* CARPENTER, Sci. Proc. Roy. Dublin Soc., **7** (5) (40): 555-557, pl. XXII, figs. 7-14.

Distribution: Torres Straits.

Ascorhynchus tuberosum UTINOMI 1962

- 1962 *Ascorhynchus tuberosum* UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 100, figs. 5-6.
- 1971 *Ascorhynchus tuberosus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 334.
- 1983 *Ascorhynchus tuberosum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 19 [key].
- 1987 *Ascorhynchus tuberosum*, - NAKAMURA, Biol. Lab., Imperial Household: 32-33 pl. 29.
- 1991 *Ascorhynchus tuberosum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 10-12, fig. 3.

Distribution: Japan.

Depth range: 123-450m.

Ascorhynchus turritus STOCK 1978

- 1978 *Ascorhynchus turritus* STOCK, Cah. Biol. Mar., **19**: 192-194, fig. 2.

- 1982 *Ascorhynchus turritus*, - CHILD, Smiths. Contr. Zool., **349**: 8.

Distribution: North American Basin, N of Bermuda; Bay of Biscay.

Depth range: 1900-4400m.

Ascorhynchus utinomii NAKAMURA & CHILD 1982

- 1982 *Ascorhynchus utinomii* NAKAMURA & CHILD, Proc. biol. Soc. Wash., **95** (2): 283-285, fig. 1.
- 1983 *Ascorhynchus utinomii*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 29-30.
- 1991 *Ascorhynchus utinomii*, - NAKAMURA & CHILD, - Smiths. Contr. Zool., **512**: 12.

Distribution: Japan.

Depth range: subtidal-41m.

Ascorhynchus verrucosus STOCK 1991

- 1991 *Ascorhynchus verrucosus* STOCK, Mem. Mus. natn. Hist. nat. (A), **151**: 216-219, figs. 3-4.

Distribution: Philippines.
Depth range: 92-97 m.

Austroraptus HODGSON 1907

- 1932 *Austroraptus*, - GORDON, Discovery Rep., **6**: 114 [key to species].
 1954 *Austroraptus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 146 [diagnosis, remarks].
 1956 *Austroraptus*, - Stock, Vidensk. Medd. Dansk. naturh. Foren., **118**: 92-93 [comparison with *Tanystylum*, diagnostic features].
 1969 *Austroraptus*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 113-116 [diagnosis, remarks, key to species].

Austroraptus calcaratus GORDON 1944

- 1944 *Austroraptus calcaratus* GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B) **5** (1): 58-62, fig. 21a-e.
 1969 *Austroraptus calcaratus*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 116, 120-121, figs. 174, 176-183, 194-196.

Distribution: Antarctica.
Depth range: 219m.

Austroraptus juvenilis CALMAN 1915.

- 1915 *Austroraptus juvenilis* CALMAN, Br. Antarct. Terra Nova Exped. 1910, Zool. **3** (1): 63-65, fig. 18.
 1932 *Austroraptus juvenilis*, - GORDON, Discovery Rep., **6**: 114-115.
 1935 *Austroraptus juvenilis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5**, Abt. IV, 2 (1): 291.
 1938 *Austroraptus juvenilis*, - GORDON, Sci. Rep. Australas.. AAntarct. Exped., ser. C, (Zool. & Bot.), **2** (8): 24.
 1952 *Austroraptus juvenilis*, - FAGE, Bull. Mus. natn. Hist. nat. Paris, ser. 2, **24** (3): 263, 272.
 1969 *Austroraptus juvenilis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem. **40**: 117-119, figs. 174-183, 187-189, 196.

Distribution: Antarctic.
Depth range: 40-472m.

Austroraptus polaris HODGSON 1907

- 1907 *Austroraptus polaris* HODGSON, Natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 54, pl. VIII, fig. 2.
 1915 *Austroraptus polaris*, - CALMAN, Br. Antarct. Terra Nova Exped. 1910, Zool. **3** (1): 62-63, fig. 17A-c.
 1915 *Austroraptus polaris*, - HODGSON, Ann. Mag. nat. Hist., ser. 8, **15**: 147.
 1923 *Austroraptus polaris*, - LOMAN, Zool. Res. Swed. Antarct. Exped., **1** (2): 30-31.
 1927 *Austroraptus polaris*, - HODGSON, Dt. Südpolar-Exped., **19**: 350.

- 1932 *Austroraptus polaris*, - GORDON, Discovery Rep., **6**: 114.
 1935 *Austroraptus polaris*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5**, Abt. 4, 2 (1): 291, fig. 207.
 1938 *Austroraptus polaris*, - GORDON, Sci. Rep. Australas. Antarct. Exped., ser. C., (Zool. & Bot.) **2** (8): 24.
 1950 *Austroraptus polaris*, - HEDGPETH, Proc. U.S. natn. Mus., **100** ((3260)): 154.
 1969 *Austroraptus polaris*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 116-117, figs. 174-186.

Distribution: Antarctica.

Depth range: 50-439m.

***Austroraptus praecox* CALMAN 1915**

- 1915 *Austroraptus praecox* CALMAN, Br. Antarct. Terra Nova Exped. 1901-1904, Nat. Hist., **3**: 63-64, fig. 18A-E.
 1932 *Austroraptus praecox*, - GORDON, Discovery Rep., **6**: 114.
 1935 *Austroraptus praecox*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5**, Abt. 4, 2 (1): 291.
 1938 *Austroraptus praecox*, - GORDON, Sci. Rep. Australas. Antarct. Exped., ser. C., (Zool. & Bot.) **2** (8): 24.
 1944 *Austroraptus praecox*, - GORDON, Rep. B.A.N.Z. Antarct. Res. Exped., ser. B, **5** (1): 57-58, fig. 20a-b.
 1969 *Austroraptus praecox*, - FRY & HEDGPETH, - N.Z. Oceanogr. Inst. Mem., **49**: 119-120, figs. 174-183, 190-193.

Distribution: Antarctica.

Depth range: 50-209m.

***Austroraptus sicarius* FRY & HEDGPETH 1969**

- 1969 *Austroraptus sicarius* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 121-123, figs. 174, 176-182, 196-200.

Distribution: Antarctica.

Depth range: 380m.

***Bathyzetes* STOCK 1955**

- 1955 *Bathyzetes* Stock, Vidensk. Medd. dansk naturhist. Foren., **117**: 261 [diagnosis, remarks].

***Bathyzetes setiger* (LOMAN 1908)**

- 1908 *Euryzyde setiger* LOMAN, Siboga Exped. Monogr., **40**: 29-30, pl. V, figs. 52-58.
 1955 *Bathyzetes setiger*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 261, 262, [key], fig. 24a-c.
 1990 *Bathyzetes setiger*, - NAKAMURA & CHILD, - Proc. biol. Soc. Wash., **103** (2): 305-307, fig. 1E-G.
 1991 *Bathyzetes setiger*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 152.

Distribution: Indonesia (near Menado, Flores Sea), New Caledonia.

Depth range: 550-1260 m.

***Bathyzetes virago* (LOMAN 1908)**

- 1908 *Eurycyde virago* LOMAN, Siboga Exped. Monogr., **40**: 30-31, pl. V, figs. 59-65.
 1955 *Bathyzetes virago* STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 262,
 fig. 24d-f.

Distribution: SE-Asia (E of Ceram).

Depth range: 567m.

Habitat: mud.

***Boehmia* HOEK 1881**

- 1954 *Boehmia*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 132-133 [diagnosis, key to South African species].
 1988 *Boehmia*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 129 [key to species].

***Boehmia chelata* (MÖBIUS 1879)**

- 1879 *Pycnogonum*(?) *chelatum* BÖHM, Monatsber. Königl. Preuss. Akad. Wiss. Berlin, 1879: 192, pl. 2, figs. 5-5d.
 1902 *Boehmia chelata*, - MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 189, pl. 28, fig. 15.
 1909 *Boehmia chelata*, - SCHIMKEWITSCH, Zool. Anz., **34** (1): 3, fig. 1A, C.
 1928 *Boehmia chelata*, - FLYNN, Repts. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Repts, I): 30, fig. 17.
 1954 *Boehmia chelata*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 133-136, fig. 22.
 1956 *Boehmia chelata*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 89.

Distribution: South Africa.

Depth range: 28-79m.

***Boehmia dubia* HEDGPETH 1950**

- 1950 (?) *Boehmia dubia* HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 155-157, fig. 18.

Distribution: Antarctica (off Cape Royds, Ross Island).
Depth range: 106m.

***Boehmia longirostris* STOCK 1957**

- 1957 *Boehmia longirostris* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 95-96, fig. 13.

Distribution: South Africa.

Boehmia tuberosa MÖBIUS 1902

- 1902 *Boehmia tuberosa*, MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 189-190, pl. 28 (figs. 13-14).
 1902 *Boehmia spinosa*, - MÖBIUS, - Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 190 [lapsus calami].
 1909 *Boehmia tuberosa*, - SCHIMKEWITSCH, Zool. Anz., **34** (1): 3-4, fig. 1B, D.
 1954 *Boehmia tuberosa*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 136-137, fig. 23.
 1957 *Boehmia tuberosa*, - STOCK, Mitt. Zool. Mus. Hamburg, **55**: 96.
 1962 *Boehmia tuberosa*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 278.
 1988 *Boehmia tuberosa*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 128-129, fig. 3H.

Distribution: South Africa.

Depth range: 11-550m.

Habitat: rocks; shells.

Calypsopycnon HEDGPETH 1948

- 1948 *Calypsopycnon* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 263-264 [diagnosis, remarks].

Calypsopycnon georgiae HEDGPETH 1948

- 1948 *Calypsopycnon georgiae* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 264-266, fig. 48.

Distribution: NW-Atlantic.

Chonothaea NAKAMURA & CHILD 1983

- 1983 *Chonothaea* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 30 [diagnosis, remarks].
 1991 *Chonothaea*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 13.

Chonothaea hians NAKAMURA & CHILD 1983

- 1983 *Chonothaea hians* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 30-32, fig. 10.

Distribution: Japan.

Depth range: 160-174m.

Chonothaea verrucosa NAKAMURA & CHILD 1991

- 1991 *Chonothaea verrucosa* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 14-16, fig. 6.

Distribution: Japan.

Depth range: 355-381m.

Cilunculus LOMAN 1908

- 1908 *Cilunculus* LOMAN, Siboga Exped. Monogr., **40**: 54 [diagnosis, remarks].
 1949 *Cilunculus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 293-294 [remarks].
 1955 *Cilunculus*, - Stock, Vidensk. Medd. Dansk naturhist. Foren., **117**: 255-256 [key to species].
 1969 *Cilunculus*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 124 [diagnosis, remarks].
 1978 *Cilunculus*, - STOCK, Cah. Biol. Mar., **19**: 197-198 [remarks, key to species].
 1983 *Cilunculus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 32-33 [remarks, key to Japanese species].
 1990 *Cilunculus*, - TURPAEVA, Zool. Zh., **69** (9): 20-21.

Cilunculus acanthus FRY & HEDGPETH 1969

- 1969 *Cilunculus acanthus* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 126-127, figs. 207-209.
 1978 *Cilunculus acanthus*, - STOCK, Cah. Biol. Mar., **19**: 197.
 1982 *Cilunculus acanthus*, - CHILD, Smiths. Contr. Zool., **349**: 9.

Distribution: Antarctic; Argentine Basin.

Depth range: 2440-2700m.

Cilunculus achelioides STOCK 1991

- 1991 *Cilunculus achelioides* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 143-146, figs. 14-15.

Distribution: New Caledonia.

Depth range: 505-515 m.

Cilunculus alcicornis STOCK 1978

- 1978 *Cilunculus alcicornis* STOCK, Cah. Biol. Mar., **19**: 201-204, fig. 6.
 1983 *Cilunculus alcicornis*, - BAMBER, Zool. J. Linn. Soc., **77**: 69.
 1991 *Cilunculus alcicornis*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 137.

Distribution: NE-Atlantic.

Depth range: 650-1140m.

Habitat: sandy mud, sometimes clayish, nullipores.

Cilunculus antillensis STOCK 1955

- 1955 *Cilunculus antillensis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 253-257, figs. 19-20.
 1986 *Cilunculus antillensis*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.

Distribution: west of St. Croix, Virgin Islands.
Depth range: about 500 fathoms.

***Cilunculus armatus* (BÖHM 1879)**

- 1879 *Lecythorhynchus armatus* BÖHM, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1879 (9): 141-142.
- 1891 *Parazetes pubescens*, - ORTMANN, Zool. Jb. Syst., 5 (1): 163-164, pl. 24, fig. 5a-d.
- 1909 *Ammothea armata*, - SCHIMKEWITSCH, Zool. Anz., 34 (1): 4, fig. 2.
- 1911 *Cilunculus armatus*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. 2 (4): 9, pl. I, figs. 1-8.
- 1918 *Cilunculus armatus*, - FUKUI, Rigakukai(?), 16 (2): 98.
- 1929 *Cilungulus armatus*, - SCHIMKEWITSCH, Fauna SSSR, Izdatljstvo Akad. Nauk SSSR, 1: 20, fig. 5A-C, 81, figs. 18, 19A-C, 20.
- 1936 *Cilunculus armatus*, - OHSHIMA, Dôbutsugaku Zasshi, 48 (8-10): 866.
- 1949 *Cilunculus armatus*, - HEDGPETH, Proc. U.S. natn. Mus., 98 (3231): 294-295, fig. 43.
- 1955 *Cilunculus armatus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., 5 (1): 27-28, fig. 16.
- 1959 *Cilunculus armatus*, - UTINOMI, - Publ. Seto Mar. Biol. Lab., 7 (2): 205.
- 1961 *Cilunculus armatus*, - LOSINA-LOSINSKY, Issled. Dalnyev. Morei SSSR, Leningrad, 7: 108.
- 1965 *Cilunculus armatus*, - UTINOMI, in: New Illustr. Encyclopaedia Fauna Japan, 2: 337, fig. 25.
- 1971 *Cilunculus armatus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., 18 (5): 334-335.
- 1983 *Cilunculus armatus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., 386: 33.
- 1987 *Cilunculus armatus*, - NAKAMURA, Biol. Lab., Imperial Household: 33-34, pls. 30, 39.

Distribution: Japan and Korea.
Depth range: 27-618 m.

***Cilunculus australiensis* CLARK 1963**

- 1963 *Cilunculus australiensis* CLARK, Rec. Austr. Mus., 26 (1): 71-73, fig. 35A-K.
- 1991 *Cilunculus australiensis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), 151: 146.

Distribution: Australia (New South Wales), New Caledonia.
Depth range: 40-435 m.

***Cilunculus cactoides* FRY & HEDGPETH 1969**

- 1969 *Cilunculus cactoides* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 124-126, figs. 205-206.

Distribution: Antarctica (off South Shetlands).
Depth range: 72-433m.

***Cilunculus compactus* STOCK 1991**

- 1991 *Cilunculus compactus* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 146-149,
figs. 16-17.
1991 *Cilunculus compactus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 219.

Distribution: New Caledonia, Philippines.

Depth range: 92-610m.

***Cilunculus crinitus* STOCK 1991**

- 1991 *Cilunculus crinitus* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 149-151,
figs. 18-19.

Distribution: New Caledonia.

Depth range: 1395 m.

***Cilunculus europaeus* STOCK 1978**

- 1978 *Cilunculus europaeus* STOCK, Cah. Biol. Mar., **19**: 198-200, figs. 4, 5a-c.

Distribution: NE-Atlantic; Mediterranean.

Depth range: 660-1200m.

Habitat: calcareous bottom with large sandstone cobbles and fine granite.

***Cilunculus frontosus* LOMAN 1908**

- 1908 *Cilunculus frontosus* LOMAN, Siboga Exped., **40**: 54-55, pl. VII, figs. 83-95.
1991 *Cilunculus frontosus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 149-151.

Distribution: New Caledonia; near Menado.

Depth range: 515-1260 m.

***Cilunculus galeritus* NAKAMURA & CHILD 1991**

- 1991 *Cilunculus galeritus* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 16-18.
fig. 7, 20-21 [text].

Distribution: Japan.

Depth range: 353-492 m.

***Cilunculus gracilis* NAKAMURA & CHILD 1991**

- 1991 *Cilunculus gracilis* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 18-20, fig.
8.

Distribution: Japan.

Depth range: 700-1440m.

***Cilunculus haradai* NAKAMURA & CHILD 1983**

1983 *Cilunculus haradai* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 33-35, fig. 11.

Distribution: Japan.

Depth range: 30-416m.

Cilunculus hirsutus CLARK 1963

1963 *Cilunculus hirsutus* CLARK, Rec. Austr. Mus., **26** (1): 73-75, fig. 36A-L.

1968 *Cilunculus hirsutus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 13.

1991 *Cilunculus hirsutus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 219.

Distribution: Australia; Philippines.

Depth range: 55-107m.

Habitat: gravel.

Cilunculus kravcovi PUSHKIN 1973

1973 *Cilunculus kravcovi* PUSHKIN, Zool. Zh., **52** (3): 432-434, figs. 1-12.

Distribution: off Croset Archipelago.

Depth range: 255-309m.

Cilunculus pedatus STOCK 1991

1991 *Cilunculus pedatus* STOCK, Mém. Mus. natn. Hist. nat (A), **151**: 152-154, figs. 20-21.

Distribution: New Caledonia.

Depth range: 825m.

Cilunculus perspicax LOMAN 1908

1908 *Cilunculus perspicax* LOMAN, Siboga Exped. Monogr., **40**: 55-56, pl. VII, figs. 96-101.

Distribution: SE-Asia (near Menado, north of Celebes).

Depth range: 1165-1260m.

Habitat: stony bottom.

Cilunculus profundus (HEDGPETH 1949)

1949 *Ammothella profunda* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 289-291, fig. 42.

1991 *Cilunculus profundus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 20 [text].

Distribution: Japan.

Depth range: 1142m.

Cilunculus sekiguchi NAKAMURA & CHILD 1983

1983 *Cilunculus sekiguchi* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 35-38,
fig. 12.

Distribution: Japan.
Depth range: 30-492m.

Cilunculus sewelli CALMAN 1938

1938 *Cilunculus sewelli* CALMAN, Sci. Rep. John Murray Exped., **5** (6): 161-163,
figs. 8, 9.
1956 *Ammothella gigas*, - FAGE, Galathea Rep., **2**: 173-175, figs. 9-14.
1963 *Cilunculus sewelli*, - CLARK, Rec. Austr. Mus., **26** (1): 75.
1968 *Cilunculus sewelli*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**:
13.
1988 *Cilunculus sewelli*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 129-130.
1990 *Cilunculus sewelli*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45**
(10): 106.
1990 *Cilunculus sewelli*, - TURPAYEVA, Zool. Zh., **69** (9): 21-22, fig. 3.
1991 *Cilunculus sewelli*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 21.

Distribution: Known from scattered places in the Indo-West Pacific: off Zanzibar, Natal (South Africa), Kenya, Mozambique Channel, New Zealand and Japan.

Depth range: 183-1789m.

Cilunculus spinicristus CHILD 1987

1987 *Cilunculus spinicristus* CHILD, Proc. biol. Soc. Wash., **100** (4): 907-909,
fig. 3.

Distribution: New Zealand (Antipodes Islands).
Depth range: 476-540m.

Cilunculus tubicinis CHILD 1982

1982 *Cilunculus tubicinis* CHILD, Smiths. Contr. Zool., **349**: 9-11, fig. 3.

Distribution: West European Basin, SW of Ireland.
Depth range: 2868-2890m.

Dromedopycnon CHILD 1982

1982 *Dromedopycnon* Child, Smiths. Contr. Zool., **349**: 15-16 [diagnosis, remarks].

Dromedopycnon acanthus CHILD 1982

1982 *Dromedopycnon acanthus* CHILD, Smiths. Contr. Zool., **349**: 16-18, fig. 5.

Distribution: southeast of Cape Horn, and from Tierra del Fuego-Burdwood Bank areas and the Brazil Slope off Recife.

Depth range: 124-903m.

Elassorhis CHILD 1982

1982 *Elassorhis* Child, Smiths. Contr. Zool., **349**: 11-12 [diagnosis, remarks].

Elassorhis laterospina CHILD 1982

1982 *Elassorhis laterospina* CHILD, Smiths. Contr. Zool., **349**: 12-14, fig. 4.

Distribution: Brazil Slope, off Recife.

Depth range: 943-1007m.

Ephyrogymna HEDGPETH 1943

1948 *Ephyrogymna*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 261 [diagnosis].

Ephyrogymna circularis HEDGPETH 1943

1943 *Ephyrogymna circularis* HEDGPETH, Proc. New England Zool. Club, **22**: 51-52, pl. 9 figs. a-g.

1948 *Ephyrogymna circularis*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 261-262, fig. 46.

1986 *Ephyrogymna circularis*, - STOCK, Bull. mar. Sci., **38** (3): 400 tab. 1.

Distribution: off Martinique, Caribbean Sea.

Depth range: 961-1034m.

Eurycyde SCHIÖDTE 1857

1948 *Eurycyde*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 260 [diagnosis, remarks].

1979 *Eurycyde*, - STOCK, Smiths. Contr. Zool., 293: 16 [key to Middle American species].

1986 *Eurycyde*, - STOCK, Bull. mar. Sci., **38** (3): 407.

1988 *Eurycyde*, - Child, Smiths. Contr. Zool., **468**: 7-8 [remarks, key to species].

1992 *Eurycyde*, - CHILD, Mem. Hourglass Cruises, **9** (1): 25 [diagnosis].

Eurycyde acanthopus STOCK 1979

1979 *Eurycyde acanthopus* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 3-8, figs. 2, 3g-h.

Distribution: off Venezuela.

Depth range: 69.5-155.5m.

Eurycyde antarctica CHILD 1987

1987 *Eurycyde antarctica* CHILD, Proc. biol. Soc. Wash., **100** (4): 909-912, fig. 4.

Distribution: Antarctic.

Depth range: 523-714m.

Eurycyde clitellaria STOCK 1955

- 1955 *Eurycyde clitellaria* STOCK, Vidensk. Medd. Dansk naturhist. Foren., **117**: 263-266, figs. 25-26.
 1967 *Eurycyde clitellaria*, - McCLOSKEY, J. nat. Hist., **1**: 128-131, figs. 18-21.
 1979 *Eurycyde clitellaria*, - CHILD, Smiths. Contr. Zool., **293**: 16.
 1986 *Eurycyde clitellaria*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.
 1988 *Eurycyde clitellaria*, - CHILD, Smiths. Contr. Zool., **468**: 7.
 1992 *Eurycyde clitellaria*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 25-26, fig. 10, 28 [text], 76 [tab. 12], 78 [tab. 14].

Distribution: Gulf of Mexico; Virgin Islands; off North Carolina and Pacific Coast of Mexico.

Depth range: intertidal-100 m.

Habitat: coral rubble.

Eurycyde curvata CHILD 1979

- 1979 *Eurycyde curvata* CHILD, Smiths. Contr. Zool., **293**: 16-19, fig. 5a-h.
 1986 *Eurycyde curvata*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1, 406 [text].
 1988 *Eurycyde curvata*, - CHILD, Smiths. Contr. Zool. **468**: 7.

Distribution: Caribbean Coast of Colombia.

Depth range: 24m.

Eurycyde diacantha STOCK 1990

1990 *Eurycyde diacantha* STOCK, Zool. Meded., **63** (16): 206-208, fig. 1.

Distribution: Cape Verde Islands.

Depth range: 140-160m.

Habitat: corse calcareous sand, gravel and stones, Foraminifera, grab.

Eurycyde extenuata CALMAN 1938

1938 *Eurycyde extenuata* CALMAN, Sci. Rep. John Murray Exped., **5** (6): 149-151, figs. 1-2.

Distribution: Zanzibar area.

Depth range: 925-2926m.

Eurycyde gorda CHILD 1979

1979 *Eurycyde gorda* CHILD, Smiths. Contr. Zool., **293**: 19-21, fig. 6.

Distribution: Panamá Caribbean.

Depth range: intertidal

Habitat: reef flats.

Eurycyde hispida (KRØYER 1844)

1891 *Eurycyde hispida* SARS, Norw. North-Atl. Exped., **6** (Zool., 20): 128-132, pl. XIV, fig. 1a-g.

1943 *Eurycyde hispida*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 88.

1963 *Eurycyde hispida*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1339.

Distribution: W-Atlantic; eastern fringe of arctic Canada to Kara Sea, Baffin Land and Greenland, as far south as Kristiansund in Norway.

Depth range: littoral-68m.

Eurycyde longioculata MÜLLER 1990

1990 *Eurycyde longioculata* MÜLLER, Senckenbergiana biol., **70** (1989) (1/3): 188-191, figs. 11-18.

Distribution: Society Islands (Bora Bora).

Depth range: 0.5-1m.

Habitat: dead corals covered with sponges and algae.

Eurycyde longisetosa HILTON 1942

1942 *Eurycyde longisetosa* HILTON, Re. Allan Hancock Exped., **5** (9): 305-307, pl. 46.

1971 *Eurycyde encantada* CHILD & HEDGPETH, J. nat. Hist., **5**: 617-619, fig. 4.

1986 *Eurycyde longisetosa*, - STOCK, Bull. mar. Sci., **38** (3): 409 [text].

1992 *Eurycyde longisetosa*, - CHILD, Smiths. Contr. Zool., **526**: 15-17, fig. 6.

Distribution: Colombia; Galapagos Archipelago; Ecuador.

Depth range: 0-37m.

Habitat: algae; corals; rocks.

Eurycyde platyspina STOCK 1992

1992 *Eurycyde platyspina* STOCK, Tijdschr. Ent., **135**: 113-114, figs. 1-8, 139.

Distribution: southern Brazil.

Depth range: 34m.

Eurycyde raphiaster LOMAN 1912

- 1912 *Eurycyde raphiaster* LOMAN, Bull. Inst. Oceanogr., **238**: 13.
 1917 *Eurycyde raphiaster*, - BOUVIER, Res. Campagnes Sci. accomplies sur son yacht par Albert Ier Prince Souverain de Monaco, **51**: 33-35, pl. 4, figs. 2-7.
 1948 *Eurycyde raphiaster* - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 260-261, fig. 45.
 1952 *Eurycyde raphiaster*, - FAGE, Bull. Mus. natn. Hist. nat., (2) **24** (6): 530.
 1954 *Eurycyde raphiaster*, - HEDGPETH, Fish Bull. Fish. Wildlife Service U.S., **89** (55): 427.
 1954 *Eurycyde raphiaster*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 116.
 1966 *Eurycyde raphiaster*, - FAGE & STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 316.
 1975 *Eurycyde raphiaster*, - STOCK, Bull. mar. Sci., **24** (4): 979; Literatur.
 1976 *Eurycyde raphiaster*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 133.
 1979 *Eurycyde raphiaster*, - CHILD, Smiths. Contr. Zool., **293**: 21.
 1979 *Eurycyde raphiaster*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 979.
 1982 *Eurycyde raphiaster*, - CHILD, Smiths. Contr. mar. Sci., **12**: 360.
 1988 *Eurycyde raphiaster*, - CHILD, Smiths. Contr. Zool., **468**: 7.
 1990 *Eurycyde raphiaster*, - STOCK, Zool. Meded., **63** (16): 209.
 1992 *Eurycyde raphiaster*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 27-28, fig. 11.
 1992 *Eurycyde raphiaster*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 45.

Distribution: amphi-atlantic; most records from West-Indian localities; in the E-Atlantic known from the Cape Verde Islands and Sénégal.

Depth range: 0-90m.

Habitat: in a variety of substrata, as dead corals, algae and seagrass beds.

Eurycyde sertula CHILD 1991

- 1991 *Eurycyde sertula* CHILD, Proc. biol. Soc. Wash., **104** (1): 139-142, fig. 1.

Distribution: W-Pacific. - Ypan.

Depth range: 1m.

Habitat: coral reefs; branching coralline algae on algal ridge.

Eurycyde setosa CHILD 1988

- 1988 *Eurycyde setosa* CHILD, Smiths. Contr. Zool., **468**: 8-10, fig. 3.
 1990 *Eurycyde setosa*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 316-317.
 1992 *Eurycyde setosa*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 156, figs. 1-5.

Distribution: Malaysia, Philippines, Australian barrier reef.

Depth range: 0-1 m.

Habitat: dead corals.

Eurycyde spinosa HILTON 1916

1916 *Eurycyde spinosa* HILTON, J. Ent. Zool. Pomona College, **8** (1): 19-24, figs. 1-6.

1992 *Eurycyde spinosa*, - CHILD, Smiths. Contr. Zool., **526**: 17.

Distribution: California; Galapagos Archipelago.

Depth range: intertidal.

Habitat: under rock.

Eurycyde unispina STOCK 1986

1986 *Eurycyde unispina* STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1, 407-409, fig. 1.

Distribution: Florida.

Depth range: 183m.

Hedgpethius CHILD 1974

1974 *Hedgpethius* Child, Proc. biol. Soc. Wash., **87** (43): 493-494 [diagnosis, remarks].

1982 *Hedgpethius*, - Child, Smiths. Contr. mar. Sci., **12**: 360 [emended diagnosis].

Hedgpethius interstitialis STOCK 1989

1989 *Hedgpethius interstitialis* STOCK, Bijdr. Dierk., **59** (2): 89-91, figs. 1-10.

Distribution: Curaçao.

Depth range: shallow water.

Habitat: strongly exposed intertidal rockpool, in coarse sand; sand and coral rubble.

Hedgpethius mamillatus CHILD 1982

1982 *Hedgpethius mamillatus* CHILD, Smiths. Contr. mar. Sci., **12**: 360-362, fig. 163a-f.

1986 *Hedgpethius mamillatus*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.

Distribution: Belize (Carrie Bow Cay).

Depth range: intertidal.

Habitat: broken rock and rubble.

Hedgpethius tridentatus CHILD 1974

1974 *Hedgpethius tridentatus* CHILD, Proc. biol. Soc. Wash., **87** (43): 494-497, fig. 1.

1982 *Hedgpethius tridentatus*, - CHILD, Smiths. Contr. mar. Sci., **87** (43): 362-363, fig. 163g-k.

1986 *Hedgpethius tridentatus*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.

Distribution: Florida (Key West); Belize (Carrie Bow Cay).

Depth range: littoral.

Habitat: among intertidal and subtidal rocks; mat of *Caulerpa verticillata* and mangrove rootlets beneath *Rhizophora*.

***Hemicheila* STOCK 1954**

1954 *Hemicheila* Stock, Vidensk. Medd. Dansk naturh. Foren., **116**: 90-94, figs. 42-43 [diagnosis].

***Hemicheila longiunguis* STAPLES 1982**

1982 *Hemicheila longiunguis* STAPLES, Mem. Qd. Mus., **20** (3): 463-465, fig. 4A-L.

Distribution: Australia (Queensland).

Depth range: shallow water.

***Hemicheila micrasterias* STOCK 1954**

1954 *Hemicheila micrasterias* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 90-94, figs. 42-43.

1978 *Hemicheila micrasterias*, - FRY, Zool. J. Linn. Soc., **63**: 44.

1985 *Hemicheila micrasterias*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **10** (18): 153-155, figs. 1-11.

1988 *Hemicheila micrasterias*, - CHILD, Smiths. Contr. Zool., **468**: 10.

1990 *Hemicheila micrasterias*, - NAKAMURA & CHILD, - Proc. biol. Soc. Wash., **103** (2): 307-308.

1991 *Hemicheila micrasterias*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 21.

Distribution: Indonesia (flores Sea), Philippines, Japan.

Depth range: 20-84 m.

Habitat: sandy mud with shells.

***Heterofragilia* HEDGPETH 1943**

1948 *Heterofragilia*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 262 [diagnosis].

***Heterofragilia amica* STOCK 1954**

1954 *Heterofragilia amica* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 136-139, figs. 67-68.

1955 *Heterofragilia amica*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 28-31, figs. 17-18.

1971 *Heterofragilia amica*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 335.

Distribution: Japan.

Depth range: 165-203m.

Habitat: hard bottom.

Heterofragilia brevicauda STOCK 1991

1991 *Heterofragilia brevicauda* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 141-143, figs. 12-13.

Distribution: New Caledonia.

Depth range: 650-680 m.

Heterofragilia fimbriata HEDGPETH 1943

1943 *Heterofragilia fimbriata* HEDGPETH, Proc. New England Zool. Club, **22**: 53.

1948 *Heterofragilia fimbriata*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 262-263, fig. 47.

Distribution: off Martinique, Caribbean Sea.

Depth range: 871m.

Heterofragilia hamisetosa LOMAN 1908

1908 *Fragilia hamisetosa* LOMAN, Siboga Exped. Monogr., **40**: 56-58, pl. VIII, figs. 112-121.

Distribution: SE-Asia, Philippines (Sulu Archipelago; near Menado and Ceram).

Depth range: 450-1260m.

Heterofragilia hirsuta NAKAMURA & CHILD 1991

1991 *Heterofragilia hirsuta* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 21-24, fig. 9.

Distribution: Japan.

Depth range: 454-495 m.

Heterofragilia major STOCK 1986

1986 *Heterofragilia major* STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1, 409-412, fig. 2.

Distribution: Caribbean (east of St. Vincent).

Depth range: 1053-1540 m.

Lecythorhynchus BÖHM 1879

1879 *Lecythorhynchus* BÖHM, Sitz.-Ber. Ges. naturforsch. Fr. Berlin, 1879, **9**: 140-141 [diagnosis].

Lecytorhynchus marginatum COLE 1904

- 1904 *Lecytorhynchus marginatus* COLE, Harriman Alaska Exped., **10**: 260-262, pl. XI, figs. 1-2, pl. XV, figs. 1-8.
- 1913 *Lecytorhynchus marginatus*, - SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **18**: 240.
- 1927 ?*Lecytorhynchus hilgendorfi*, - OHSHIMA, Proc. Imp. Acad., Tokyo, **3** (9): 610-611, figs. 1-3.
- 1929 *Lecytorhynchus marginatus*, - SCHIMKEWITSCH, Fauna SSSR, Izdatljstvo Akad. Nauk. SSSR, **1**: 50-53, figs. 7-9.
- 1933 *Lecytorhynchus marginatus*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 61-62.
- 1934 *Lecytorhynchus marginatus*, - SCHMITT, J. Wash. Acad. Sci., **24** (1): 69.
- 1938 *Lecytorhynchus marginatus*, - HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 184-185.
- 1940 *Lecytorhynchus marginatus*, - HEDGPETH, J. Wash. Acad. Sci., **30** (2): 86-87.
- 1941 *Lecytorhynchus marginatus*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 255.
- 1943 *Lecytorhynchus marginatus*, - HILTON, J. Ent. Zool. Pomona College, **35** (1): 4.
- 1949 *Lecytorhynchus marginatus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 296, fig. 44c-d.
- 1951 *Lecytorhynchus marginatus*, - HEDGPETH, Wasmann J. Biol., **9** (1): 111.
- 1954 *Lecytorhynchus marginatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 139-141, fig. 69.

Distribution: California; Sea of Okhotsk; Japan.

Habitat: Corallina.

Megarhetus CHILD 1982

- 1982 *Megarhetus* Child, Smiths. Contr. Zool., **349**: 14 [diagnosis, remarks].

Megarhetus brevicheliferus (HEDGPETH 1948)

- 1948 *Achelia brevichelifera* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 245-246, fig. 38h-l.
- 1968 *Scipiolus brevicheliferus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 15.
- 1982 *Megarhetus brevicheliferus*, - CHILD, Smiths. Contr. Zool., **349**: 14-15, fig. 2f-i.

Distribution: North American Slope and Basin.

Depth range: 750-2500m.

Nymphonella OHSHIMA 1927

- 1927 *Nymphonella* OHSHIMA, Annot. Zool. Jap., **11** (3): 262 [diagnosis].

Nymphonella lambertensis STOCK 1959

- 1959 *Nymphonella lambertensis* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 549-551, fig. 1.
 1988 *Nymphonella lambertensis*, - MUNILLA, Monogr. Zool. Mar., **3**: 192-194, fig. 10.

Distribution: South Africa.

Depth range: 0-15m.

Habitat: sand, rocks.

***Nymphonella tapetis* OHSHIMA 1927**

- 1927 *Nymphonella tapetis* OHSHIMA, Annot. Zool. Jap., **11** (3): 262.
 1927 *Nymphonella tapetis*, - OHSHIMA, Bulteno Sci. Fak. Terkultura, Kyushu Imp. Univ., **2** (5): 367.
 1933 *Nymphonella tapetis*, - OHSHIMA, Annot. Zool. Jap., **14** (1): 53.
 1933 *Nymphonella tapetis*, - OHSHIMA, Dôbutsugaku Zasshi, **45**: 94.
 1935 *Nymphonella tapetis*, - OHSHIMA, Annot. Zool. Jap., **15** (1): 95.
 1936 *Nymphonella tapetis*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 846.
 1936 *Nymphonella tapetis*, - ARITA, Annot. Zool. Jap., **15** (4): 469.
 1937 *Nymphonella tapetis*, - ARITA, J. Dep. Agric. Kyushu Imp. Univ., **5** (6): 271.
 1937 *Nymphonella tapetis*, - OHSHIMA, C.R. XII'eme Congr. Int. Zool., 1935: 1616.
 1942 *Nymphonella tapetis*, - Proc. Imp. Acad., Tokyo, **18** (8): 520.
 1943 *Nymphonella tapetis*, - OHSHIMA, Bulteno Sci. Fak. Terkultura, Kyushu Imp. Univ., Tokyo, **10** (4): 374.
 1950 *Nymphonella tapetis*, - CALVEZ, Arch. Zool. Exp. Gen., **86** (Notes & Rev.): 114.
 1968 *Nymphonella lecalvezi* GUILLE & SOYER, Vie Milieu (A) **18** (2A): 346.
 1971 *Nymphonella tapetis*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 331.
 1978 *Nymphonella tapetis*, - de HARO, Zool. J. Linn. Soc., **63** (1-2):
 1987 *Nymphonella tapetis*, - ARNAUD, Mesogee, **47**: 42.
 1987 *Nymphonella tapetis*, - ARNAUD & BAMBER, Adv. Mar. Biol., **24**: 50.
 1988 *Nymphonella lecalvezi*, - MUNILLA, Monogr. Zool. Mar., **3**: 194-195.

Distribution: Japan; Namibia; Mediterranean.

Depth range: 0-15m.

Habitat: rocky and sandy bottoms.

***Nymphopsis* HASWELL 1884**

- 1908 *Nymphopsis*, - LOMAN, Siboga Exped. Monogr., **40**: 49-50 [diagnosis, remarks].
 1948 *Nymphopsis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 249-250 [diagnosis, remarks, key to species].
 1954 *Nymphopsis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 141-142 [diagnosis, remarks].
 1967 *Nymphopsis*, - McCLOSKEY, J. nat. Hist., **1**: 122.
 1992 *Nymphopsis*, - CHILD, Mem. Hourglass Cruises, **9** (1): 28 [diagnosis].

***Nymphopsis abstrusus* LOMAN 1923**

- 1923 *Nymphopsis abstrusus* LOMAN, Ark. Zool., **15** (9): 7-10, fig. D1-5.

Distribution: subantarctic area.

Nymphopsis acinacispinus WILLIAMS 1933

- 1933 *Nymphopsis acinacispinus* WILLIAMS, Ann. Mag. Nat. Hist., **10** (12): 173.
 1975 *Nymphopsis acinacispinus*, - CHILD, Smiths. Contr. Zool., **190**: 26-27, fig. 11c-f.
 1980 *Nymphopsis acinacispinus*, - HOOPER, Aust. Zool., **20** (3): 475.

Distribution: Australia.

Depth range: 0-2.5m.

Habitat: Sargassum.

Nymphopsis anarthra LOMAN 1928

- 1928 *Nymphopsis anarthrus* LOMAN, Tijdschr. Nederl. Dierk. Ver., ser. 3, **1**: 39-42, fig. 1.
 1948 *Nymphopsis anarthra*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 250.
 1986 *Nymphopsis anarthra*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.
 1992 *Nymphopsis anarthra*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 28-29, fig. 12, 76 [tab. 12], 78 [tab. 14].

Distribution: central West Florida Shelf; off Tortuga Island, Venezuela.

Depth range: 45-73 m.

Nymphopsis armatus HASWELL 1884

- 1884 *Nymphopsis armatus* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1025.
 1908 *Nymphopsis armatus*, - LOMAN, Siboga Exped. Monogr., **40**: 49.
 1919 *Nymphopsis armatus*, - FLYNN, Pap. Proc. Roy. Soc. Tasm., 1919: 84.
 1929 *Nymphopsis armatus*, - FLYNN, Mem. Qd. Mus., **9** (3): 256-257, figs. 4-5.

Distribution: Australia.

Depth range: 17m.

Nymphopsis bathursti Williams, 1940

- 1963 *Nymphopsis acinacispinus bathursti*, - CLARK, Rec. Austr. Mus., **26** (1): 5.
 1992 *Nymphopsis bathursti*, - STOCK, Bijdr. Dierk., **62** (2): 82-83, fig. 1.

Distribution: Western Australia; Oman.

Depth range: 0-15m.

Habitat: algae; muddy tidal flats with stones and algae.

Nymphopsis curtiscapus STOCK 1974

- 1974 *Nymphopsis curtiscapus* STOCK, Bull. Zool. Univ. Amsterdam, **4** (3): 14-15, fig. 2.
 1982 *Nymphopsis curtiscapus*, - STOCK, Monit. zool. ital., Suppl. **17** (7): 187.

1990 *Nymphopsis curtiscapus*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 67.

Distribution: Madagascar; Somalia; Kenya.

Depth range: 0-0.5m.

Habitat: dead corals covered with algae.

***Nymphopsis cuspidata* (HODGSON 1910)**

- 1910 *Ammothaea cuspidata* HODGSON, in: Schultze, Zool. und anthropol. Ergebni. einer Forschungsreise im westl. und zentr. Südafrika 1903-1905..., **4** (1): 221-223, fig. 1.
- 1923 *Nymphopsis cuspidata*, - LOMAN, Medd. Göteborgs Mus., Zool. Avd., **22** (6): 7, fig. D.
- 1954 *Nymphopsis cuspidata*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 142-144, figs. 25-27.
- 1959 *Nymphopsis cuspidata*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 551-552.

Distribution: South Africa.

Depth range: 11-27.5m.

Habitat: sand; rocks; algae; shells.

***Nymphopsis denticulata* GORDON 1932**

- 1932 *Nymphopsis denticulata* GORDON, Discovery Rep., **6**: 123-125, figs. 69-70.

Distribution: Lively Islands, East Falkland Islands.

Depth range: 79m.

***Nymphopsis dromedaria* McCLOSKEY 1967**

- 1967 *Nymphopsis dromedaria* McCLOSKEY, J. nat. Hist., **1**: 119-123, figs. 1-6.

Distribution: off North Carolina.

Depth range: 100m.

***Nymphopsis duodorsospinosa* HILTON 1942**

- 1942 *Nymphopsis duodorsospinosa* HILTON, Allan Hancock Pac. Exped., **5** (9): 303-305, pl. 45.
- 1948 *Nymphopsis duiodorsospinosa*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 250-252, fig. 40.
- 1971 *Nymphopsis duodorsospinosa*, - CHILD & HEDGPETH, J. nat. Hist., **5**: 609 [list].
- 1973 *Nymphopsis duodorsospinosa*, - KRAEUTER, J. nat. Hist., **7**: 496.
- 1975 *Nymphopsis duodorsospinosa*, - STOCK, Bull. mar. Sci., **24** (4): 978.
- 1976 *Nymphopsis duodorsospinosa*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 134, 157.
- 1976 *Nymphopsis duodorsospinosa*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27** (3/4): 342.
- 1979 *Nymphopsis duodorsospinosa*, - CHILD, Smiths. Contr. Zool., **293**: 21; Literatur.

- 1982 *Nymphopsis duodorsospinosum*, - CHILD, Smiths. Contr. mar. Sci., **12**: 363.
 1986 *Nymphopsis duodorsospinosa*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.
 1992 *Nymphopsis duodorsospinosa*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key],
 9 [tab. 3], 30-31, fig. 13.

Distribution: E-Pacific. - Galápagos-archipelago, Gulf of California south to Panamá; W-Atlantic. - South Carolina (U.S.A.) to the northern coast of South America.

Depth range: 0-60m.

Habitat: in a variety of substrata.

***Nymphopsis korotnewi* SCHIMKEWITSCH 1887**

- 1887 *Nymphopsis korotnewi* SCHIMKEWITSCH, Zool. Jb., **3**: 127-134, figs. 1-12.
 1908 *Nymphopsis korotnewi*, - LOMAN, Siboga Exped. Monogr., **40**: 50-52, pl. XIII,
 figs. 179-181.
 1975 *Nymphopsis korotnewi*, - CHILD, Smiths. Contr. Zool., **190**: 27.
 1991 *Nymphopsis korotnewi*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 219-220.

Distribution: Sunda Islands (Indonesia), Western Australia, Philippines.

Depth range: 34-144m.

***Nymphopsis melidae* SAWAYA 1947**

- 1947 *Nymphopsis melidae* SAWAYA, Bol. Fac. Fil., Cien. Letr. Univ. S. Paulo (Zool.), **12**: 83-89, pl. I, figs. 1-2, pl. II, figs. 3-6.
 1992 *Nymphopsis melidae*, - STOCK, Tijdschr. Ent., **135**: 115, 139.

Distribution: Brazil.

Depth range: upper littoral-41m.

***Nymphopsis muscosa* LOMAN 1908**

- 1908 *Nymphopsis muscosus* LOMAN, Siboga Exped. Monogr., **40**: 52-53, pl. 13, figs. 175-188.
 1911 *Nymphopsis muscosus*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 11.
 1918 *Nymphopsis muscosus*, - FUKUI, Rigakukai[?], **16** (2): 99.
 1936 *Nymphopsis muscosus*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 864.
 1953 *Nymphopsis muscosa*, - STOCK, Temminckia, **9**: 307.
 1954 *Nymphopsis muscosa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 121.
 1959 *Nymphopsis muscosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 205.
 1971 *Nymphopsis muscosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 335.
 1987 *Nymphopsis muscosa*, - NAKAMURA, Biol. Lab., Imperial Household: 34-36, pls. 32, 32, 40.
 1991 *Nymphopsis muscosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 24.

Distribution: Malaysia, Philippines and Japan.

Depth range: 12-300m.

***Nymphopsis spinosissima* (HALL 1912)**

- 1912 *Ammothella spinosissima* HALL, Ann. Rep. Laguna Mar. Lab., 1: 95-99, figs. 51, 52B, D, H.
- 1913 *Ammothella spinosissima*, - HALL, Univ. Calif. Publ. Zool., 11 (6): 130, 132 [key].
- 1915 *Ammothella spinosissima*, - HILTON, J. Ent. Zool. Pomona College, 7 (1): 68.
- 1915 *Ammothella spinosissima*, - HILTON, J. Ent. Zool. Pomona College, 7 (3): 204.
- 1920 *Ammothella spinosissima*, - HILTON, J. Ent. Zool. Pomona College, 12 (4): 93.
- 1934 *Ammothella spinosissima*, - SCHMITT, J. Wash. Acad. Sci., 24 (1): 69.
- 1939 *Nymphopsis spinosissimus*, - HEDGPETH, Am. Midl. Nat., 22 (2): 461-463, fig. 2m-r.
- 1939 *Ammothella spinosissima*, - HILTON, J. Ent. Zool. Pomona College, 31 (2): 32.
- 1939 *Nymphopsis spinosissimus*, - HILTON, J. Ent. Zool. Pomona College, 31 (4): 72.
- 1941 *Nymphopsis spinosissimus*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., 9 (26): 256 [key], pl. 11.
- 1942 *Nymphopsis spinosissima*, - HILTON, Re. Allan Hancock Exped., 5 (9): 278, 300-302, pl. 44.
- 1943 *Nymphopsis spinosissima*, - J. Ent. Zool. Pomona College, 34 (4): 98.
- 1951 *Nymphopsis spinosissima*, - HEDGPETH, Wasmann J. Biol., 9 (1): 110.
- 1955 *Nymphopsis spinosissima*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., 117: 253.
- 1964 *Nymphopsis spinosissima*, - HEDGPETH, in: R.I. Smith et al., eds, Intertidal Invertebrates of the Central Calif. Coast, Berkeley: Univ. Calif. Press: 206.
- 1987 *Nymphopsis spinosissima*, - CHILD, Proc. biol. Soc. Wash., 100 (3): 554.

Distribution: California.

Depth range: 0-31m.

Habitat: on weeds.

***Nymphopsis varipes* STOCK 1962**

- 1962 *Nymphopsis varipes* STOCK, Trans. Roy. Soc. S. Afr., 36 (4): 273-276, figs. 1-2.

Distribution: South Africa.

Depth range: 29m.

Habitat: on rocks.

***Paranymphon* CAULLERY 1896**

- 1948 *Paranymphon*, - HEDGPETH, Proc. U.S. natn. Mus., 97 (3216): 253 [diagnosis].
- 1986 *Paranymphon*, - STOCK, Bull. mar. Sci., 38 (3): 412.
- 1992 *Paranymphon*, - CHILD, Mem. Hourglass Cruises, 9 (1): 31 [diagnosis].

***Paranymphon filarium* STOCK 1986**

- 1986 *Paranymphon filarium* STOCK, Bull. mar. Sci., 38 (3): 401 tab. 1, 413-415, fig. 3.

Distribution: W- Atlantic (Santaren Channel, north of Cuba; Straits of Florida).

Depth range: 333-576m.

***Paranymphon magnidigitatum* HONG & KIM 1987**

- 1987 *Paranymphon magnidigitatum* HONG & KIM, Kor. J. Syst. Zool., **3** (2): 150-152, figs. 10-11.
 1991 *Paranymphon magnidigitatum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 24.

Distribution: Japan and Korea.

Depth range: 49-80m.

***Paranymphon spinosum* CAULLERY 1896**

- 1896 *Paranymphon spinosum* CAULLERY, Ann. Univ. Lyon, **26**: 361, pl. 12 figs. 1-6.
 1899 *Paranymphon spinosum*, - MEINERT, Danish Ingolf Exped., **3** (1): 46-47, pl. 4, figs. 20-28.
 1908 *Paranymphon spinosum*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 222-224, pl. 30, figs. 10-14.
 1917 *Paranymphon spinosum*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 17, pl. 3, figs. 3-6.
 1933 *Paranymphon spinosum*, - STEPHENSEN, Medd. Grönland, **79** (6): 6.
 1948 *Paranymphon spinosum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 253, fig. 41.
 1965 *Paranymphon spinosum*, - STOCK & SOYER, Vie Milieu, **16** (1B): 415-416.
 1966 *Paranymphon spinosum*, - STOCK, Vie Milieu, **17** (1B): 408-409, fig. 1.
 1973 *Paranymphon spinosum*, - ARNAUD, Tethys, **5** (1): 151.
 1978 *Paranymphon spinosum*, - STOCK, Cah. Biol. Mar., **19**: 204-205, fig. 5d-g.
 1982 *Paranymphon spinosum*, - CHILD, Smiths. Contr. Zool., **349**: 18-19.
 1983 *Paranymphon spinosum*, - BAMBER, Zool. J. Linn. Soc., **77**: 69.
 1983 *Paranymphon spinosum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 38.
 1986 *Paranymphon spinosum*, - Chimenz & Cottarelli, Oebalia, **13**, N.S.: 138-139, figs. 2-13.
 1987 *Paranymphon spinosum*, - ARNAUD, Mesogee, **47**: 43.
 1987 *Paranymphon spinosum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 149-150, figs. 8-9.
 1987 *Paranymphon spinosum*, - STOCK, Cah. Biol. Mar., **28**: 506-507.
 1991 *Paranymphon spinosum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 24.
 1992 *Paranymphon spinosum*, CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 31-33, fig. 14.

Distribution: N-Atlantic including Caribbean; Japan and Korea.

Depth range: 7-2000m.

Habitat: sandy bottom with *Cymodocea nodosa*; mud; coarse bottom, with stones and sponges.

***Proboehmia* STOCK 1991**

- 1991 *Proboehmia* STOCK, Mém. Mus. natn. Hist. nat (A), **151**: 155 [diagnosis,

remarks].

Proboehmia tubirostris STOCK 1991

1991 *Proboehmia tubirostris* STOCK, - Mém. Mus. natn. Hist. nat. (A), **151**: 155-158, figs. 22-23.

Distribution: New Caledonia.
Depth range: 530m.

Prototrygaeus STOCK 1975

1975 *Prototrygaeus* Stock, Bull. mar. Sci., **24** (4): 979 [diagnosis, remarks].
 1992 *Prototrygaeus*, - Child, Smiths. Contr. Zool., **526**: 17 [emended diagnosis].

Prototrygaeus ammotelloides STOCK 1975

1975 *Prototrygaeus ammotelloides* STOCK, Bull. mar. Sci., **24** (4): 979-981, fig. 9.
 1986 *Prototrygaeus ammotelloides*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.

Distribution: Guyana Shelf.
Depth range: 23-27.5m.
Habitat: muddy sand; sandy mud.

Prototrygaeus contrarius Child, 1992

1992 *Prototrygaeus contrarius* CHILD, Smiths. Contr. Zool., **526**: 17-19, fig. 8.

Distribution: Ecuador (outer Gulf of Guayaquil).
Depth range: 80m.

Prototrygaeus jordanae CHILD 1990

1990 *Prototrygaeus jordanae* CHILD, Proc. biol. Soc. Wash., **103** (1): 157-160, fig. 1.

Distribution: California.
Depth range: 50-250 m.
Habitat: soft sediment with hydroids on molluscs and wood.

Scipiolus LOMAN 1908

1908 *Scipiolus* LOMAN, Siboga Exped. Monogr., **40**: 58 [diagnosis].

Scipiolus bifidus STOCK 1968

1968 *Scipiolus bifidus* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 14-15,

fig. 3.

Distribution: South Africa (off False Bay).

Depth range: 1361m.

Habitat: hard bottom.

***Scipiolus plumosus* LOMAN 1908**

1908 *Scipiolus plumosus* LOMAN, Siboga Exped. Monogr., **40**: 60-61, pl. I, figs. 1-13.

1991 *Scipiolus plumosus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 160-161, fig. 26.

Distribution: Kei Islands (Indonesia), Loyalty Islands, New Caledonia.

Depth range: 90-300m.

***Scipiolus spinosus* UTINOMI 1955**

1955 *Scipiolus spinosus* UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 31-33, fig. 19.

1971 *Scipiolus spinosus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 335.

Distribution: Japan.

Depth range: 71m.

***Scipiolus thermophilus* TURPAEVA 1988**

1988 *Scipiolus thermophilus* TURPAEVA, Zool. Zh., **LXVII** (6): 950-952, figs. 1-2.

Distribution: Juan de Fuca Ridge.

Depth range: 1800m.

Habitat: hydrothermal vents.

***Scipiolus validus* STOCK 1957**

1957 *Scipiolus validus* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 91-94, figs. 8-9, 10b, d.

1971 *Scipiolus validus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 335.

Distribution: Japan.

Depth range: 38m.

***Sericosura* FRY & HEDGPETH 1969**

1969 *Sericosura* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 112 [diagnosis].

***Sericosura bifurcata* STOCK 1991**

1991 *Sericosura bifurcata* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 158-160, figs. 24-25.

Distribution: New Caledonia.

Depth range: 3680-3700 m.

***Sericosura cochleifovea* CHILD 1989**

1989 *Sericosura cochleifovea* CHILD, Proc. biol. Soc. Wash., **102** (3): 732-737, fig. 1.

Distribution: Marianas Back-Arc Basin.

Depth range: 3660m.

Habitat: hydrothermal Snail Pits Vent.

***Sericosura mitrata* (GORDON 1944)**

1944 *Achelia mitrata* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., ser. B., **5** (1): 54-57, figs. 19a-e, 22b.

1969 *Sericosura mitrata*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 112, 113, 152, 153, 173.

1982 *Sericosura mitrata*, - CHILD, Smiths. Contr. Zool., **349**: 19-21, fig. 6.

Distribution: Antarctica; of SW Africa (Walvis Ridge).

Depth range: 219-2154m.

***Sericosura venticola* CHILD 1987**

1987 *Sericosura venticola* CHILD, Proc. biol. Soc. Wash., **100** (4): 896-899, fig. 2.

Distribution: NE-Pacific (Endeavour Segment).

Depth range: 2208-2216m.

Habitat: hydrothermal vents, associated with vestimentiferans.

***Tanystylum* Miers 1879**

1932 *Tanystylum*, - GORDON, Discovery Rep., **6**: 117-118 [remarks, key to Antarctic and Subantarctic species].

1948 *Tanystylidae*, *Tanystylum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 266 [diagnosis, remarks, key to NE-American species].

1954 *Tanystylum*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 141-142 [remarks].

1956 *Tanystylum*, - STOCK, Vidensk. Medd. Dansk. naturh. Foren., **118**: 92-93 [comparison with *Austroraptus*, diagnostic features].

1977 *Tanystylum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 313-318 [remarks on and synopsis of species from southern Oceans; diagnosis, key to species south of 30°S].

1979 *Tanystylum*, - CHILD, Smiths. Contr. Zool., **293**: 21-22 [remarks on Caribbean

- species, key to Middle American species].
 1983 *Tanystylum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 38 [key to Japanese species].
 1992 *Tanystylum*, - CHILD, Mem. Hourglass Cruises, **9** (1): 33 [diagnosis].

***Tanystylum acuminatum* STOCK 1954**

- 1954 *Tanystylum acuminatum* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 125-127, fig. 29.
 1979 *Tanystylum acuminatum*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 11.
 1986 *Tanystylum acuminatum*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.
 1990 *Tanystylum acuminatum*, - MÜLLER, Bonn. zool. Beitr. **41** (3-4): 278.
 1992 *Tanystylum acuminatum*, - STOCK, Tijdschr. Ent., **135**: 119, 139, figs. 19-21.

Distribution: E-Caribbean Islands; Brazil.

Depth range: 0-15m.

Habitat: rocky shore; dead corals and algae; *Dyctiota*.

***Tanystylum anthomasti* HEDGPETH 1949**

- 1949 *Tanystylum anthomasti* HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 297-298, fig. 45.
 1954 *Tanystylum anthomasthi*, - UTINOMI, Publ. Akkeshi Mar. Biol. Stat., **3**: 23-26, figs. 10-11, pl. 1, fig. 2.
 1963 *Tanystylum anthomasti*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1339-1340, figs. 7-10.
 1965 *Tanystylum anthomasthi*, - UTINOMI, New Illustr. Encyclopaedia Fauna Japan, **2**: 338.
 1971 *Tanystylum anthomasti*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 317-347.

Distribution: Japan; Alaska; Oregon.

Depth range: 24-66m.

Habitat: on *Metalcyonium pacificum* (YAMADA) and on the nephteid octocorall *Gersemia rubiformis* (EHRENBERG).

***Tanystylum antipodum* CLARK 1977**

- 1972 *Tanystylum cavidorum*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 334-335.
 1972 *Tanystylum cavidorum*, - CLARK, N.Z. J. mar. Freshw. Res., **5** (3-4) (1971): 440-441, fig. 5G-J.
 1977 *Tanystylum antipodum* CLARK, J. Roy. Soc. N.Z., **7** (3): 318-321, figs. 2-9, 18, 19.

Distribution: New Zealand (Campbell, Snares, Antipodes, Stewart Islands).

Depth range: shallow water.

Habitat: under *Durvillea holdfast*; intertidal rocks; on *Xiphophora hydroids*; among algae and hydroids in pool; associated with *Margarella hinemoa* and *Exosphaeroma gigas*.

Tanystylum beuroisi ARNAUD 1974

1971 *Tanystylum brevipes*, - ARNAUD, Tethys, **3**: 160 [part.], figs. 3-4.

1974 *Tanystylum beuroisi* ARNAUD, Tethys, **5**: 784-787, figs. 10-17.

1977 *Tanystylum beuroisi*, - CLARK, - J. Roy. Soc. N.Z., **7** (3): 321.

Distribution: St. Paul and Amsterdam Islands.

Depth range: 80-100m.

Habitat: *Laminaria pallida*.

Tanystylum bigibbosum FAGE & STOCK 1966

1966 *Tanystylum bigibbosum* FAGE & STOCK, Ann. Inst. Océanogr. Monaco, **44**: 321-323, fig. 3.

Distribution: Cape Verde Islands.

Depth range: 17m.

Habitat: "fond rocheux" (FAGE & STOCK).

Tanystylum birkelandi CHILD 1979

1979 *Tanystylum birkelandi* CHILD, Smiths. Contr. Zool., **293**: 23, fig. 7.

1982 *Tanystylum birkelandi*, - CHILD, Smiths. Contr. mar. Sci., **12**: 363.

1986 *Tanystylum birkelandi*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.

1990 *Tanystylum birkelandi*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 278-279, figs. 1-5.

Distribution: Panamá Caribbean; Belize (Carrie Bow Cay); Martinique.

Depth range: 0-2m.

Habitat: reef flats, rubble and calcareous algae; dead corals in exposed location.

Tanystylum bredini CHILD 1970

1970 *Tanystylum bredini* CHILD, Proc. biol. Soc. Wash., **83** (27): 296-299, fig. 3.

1977 *Tanystylum bredini*, - CHILD, Proc. biol. Soc. Wash., **90** (2): 441.

1988 *Tanystylum bredini*, - CHILD, Bull. biol. Soc. Wash., **8**: 52.

1989 *Tanystylum bredini*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 125, figs. 11-21.

1990 *Tanystylum bredini*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 67, figs. 7-15.

1992 *Tanystylum bredini*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 156-159, figs. 6-11.

1992 *Tanystylum bredini*, - STOCK, Bijdr. Dierk., **62** (2): 92-93.

Distribution: French Polynesia, Seychelles, Kenya, Sri Lanka, Malaysia; Oman.

Depth range: 0-2 m.

Habitat: coral reefs and sandstone reefs; dead corals; stones in tidal pools; under stones.

Tanystylum brevicaudatum FAGE & STOCK 1966

- 1966 *Tanystylum brevicaudatum* FAGE & STOCK, Ann. Inst. Océanogr. Monaco, **44**: 319-321, figs. 2a-e.
 1971 *Tanystylum brevicaudatum*, - ARNAUD, Tethys, **3** (1): 162-163, figs. 5-9.
 1977 *Tanystylum brevicaudatum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 321.
 1990 *Tanystylum brevicaudatum*, - STOCK, Zool. Meded., **63** (16): 219.

Distribution: St. Paul and Amsterdam Islands; Cape Verde Islands.

Depth range: 0-15m.

Habitat: sandy bottom, rock ledges; among bryozoans.

Tanystylum brevipes (HOEK 1881)

- 1881 *Discoarachne brevipes* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 74-76, 105, pl. 7, figs. 8-12.
 1901 *Discoarachne brevipes*, - COLE, Zool. Jb. (Syst., Geogr. Biol.), **15** (2): 243-248, pl. 13.
 1904 *Discoarachne brevipes*, - LOMAN, Zool. Jb., **20** (4): 379, pl. 14, figs. 7-11.
 1910 *Discoarachne brevipes*, HODGSON, in: Denkschr. Med.-Naturwiss. Ges. Jena, **16**: 227.
 1914 *Tanystylum paulovense*, - HODGSON, Zool. Anz., **45** (4): 164.
 1923 *Discoarachne brevipes*, - LOMAN, Ark. Zool., **15** (9): 6-7, fig. C1-2.
 1927 *Tanystylum paulovense*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. II): 352-353.
 1928 *Discoarachne brevipes*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 35.
 1929 *Tanystylum paulovensis*, - SCHIMKEWITSCH, Fauna SSSR, Izdatljstvo Akad. Nauk SSSR, **1**: 64.
 1951 *Discoarachne brevipes*, - STOCK, Mem. Inst. Roy. Sci. Nat. Belg., ser. **2** (43): 3.
 1954 *Discoarachne brevipes*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 149-151, fig. 31a-b.
 1956 *Discoarachne brevipes*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 93.
 1959 *Discoarachne brevipes*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 555-556.
 1962 *Tanystylum brevipes*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 276.
 1971 *Tanystylum brevipes*, - ARNAUD, Tethys, **3** (1): 160.
 1974 *Tanystylum brevipes*, - ARNAUD, Tethys, **5** (4): 787, figs. 18-19.
 1977 *Tanystylum brevipes*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 322.
 1988 *Tanystylum brevipes*, - MUNILLA, Monogr. Zool. Mar., **3**: 189-192, figs. 8-9 [juvenile stages].

Distribution: South Africa; St. Paul and Amsterdam Islands.

Depth range: 45-100m.

Habitat: Sand; rocks; algae; mussels; kelp; in pools under granite boulders.

Tanystylum calcirostrum SCHIMKEWITSCH 1889

- 1889 *Tanystylum calcirostre* SCHIMKEWITSCH, Atti Accad. Lincei 1889, Mem. Cl. Sci., Fisiche, Matematiche e Naturel, ser. **4** (6): 331-333, figs. 5-7.

- 1940 *Tanystylum calcirostre*, - MARCUS, Bolm. Fac. Fil., Ciênc. Letr., Univ. S. Paulo, **19** (Zool.), 4: 97.
 1948 *Tanystylum calcirostre*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 268, fig. 49b-d.
 1979 *Tanystylum calcirostrum*, - CHILD, Smiths. Contr. Zool., **293**: 23-26, fig. 8.
 1986 *Tanystylum calcirostrum*, - STOCK, Bull. mar. Sci., **38** (3): 401 tab. 1.

Distribution: Bermuda.

Tanystylum californicum HILTON 1939

- 1939 *Tanystylum californicum* HILTON, J. Ent. Zool. Pomona College, **31** (2): 32-33.
 1939 *Tanystylum californicum* HILTON, J. Ent. Zool. Pomona College, **31** (4): 72-73, pl. I, fig. 6.
 1951 *Tanystylum californicum*, - HEDGPETH, Wasmann J. Biol., **9** (1): 115.

Distribution: California.

Habitat: on hydroid Aglaophenia; bryozoans.

Tanystylum cavidorsum STOCK 1957

- 1954 *Tanystylum neorhetum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 149-151, figs. 73-74.
 1957 *Tanystylum cavidorsum* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 98-100, fig. 15.
 1961 *Tanystylum cavidorsum steatopygidium* HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 12-14, fig. 9.
 1970 *Tanystylum cavidorsum*, - ARNAUD, Bull. Mus. Natn. Hist. Nat., Paris, (2) **41** (6): 1424-1426, figs. 1-2, 6.
 1977 *Tanystylum cavidorsum*, - CLARK, N.Z. J. mar. Freshw. Res., **11** (1): 176.
 1977 *Tanystylum cavidorsum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 323.

Distribution: New Zealand; Chile; South Georgia; Possession Is., Crozet Is.

Depth range: 0-26m.

Habitat: holdfast fauna; hydroids and algae in pool.

Tanystylum cinctum CHILD 1992

- 1992 *Tanystylum cinctum* CHILD, Smiths. Contr. Zool., **526**: 21-23, fig. 9.

Distribution: Ecuador.

Depth range: 3m.

Tanystylum conirostre (DOHRN 1881)

- 1881 *Clotenia conirostris* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 161.
 1953 *Tanystylum conirostre*, - BACESCU, Bul. Stiint. Acad. Rep. Pop. Rom., Sect. Stiint. Biol., Agr., Geol. si Geogr., **5** (2): 266.
 1973 *Tanystylum conirostrum*, - KRAPP, Beaufortia, **21** (277): 60-61, fig. 3.
 1975 *Tanystylum conirostre*, - KRAPP-SCHICKEL & KRAPP, - Vie Milieu, (B) **25** (1):

- 14 figs. 4, 17, 18-20 (tab. II), 21-23 (ecology).
- 1976 *Tanystylum conirostre*, - ARNAUD, Acta Ecol. Iranica, **1**: 69-70.
- 1978 *Tanystylum conirostre*, - CHIMENZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 98-100, fig. 4.
- 1979 *Tanystylum conirostre*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **41** (3): 141.
- 1986 *Tanystylum conirostrum*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.
- 1987 *Tanystylum conirostre*, - ARNAUD, Mesogee, **47**: 43.
- 1989 *Tanystylum conirostre*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 293.
- 1992 *Tanystylum conirostre*, - PÉREZ-RUZAFA, A. & T. MUNILLA, Sci. Mar., **56** (1): 25, 27-28, fig. 3 [biology and ecology].

Distribution: Mediterranean Sea, probably Amphi-Atlantic.

Depth range: 0-3m.

Habitat: in a variety of substrata, more often found on algae.

***Tanystylum distinctum* CHILD & HEDGPETH 1971**

- 1971 *Tanystylum distinctum* CHILD & HEDGPETH, J. nat. Hist., **5**: 623-625, fig. 6.
- 1992 *Tanystylum distinctum*, - CHILD, Smiths. Contr. Zool., **526**: 21.

Distribution: Ecuador; Galápagos Archipelago; Chile.

Depth range: 0-75m.

Habitat: algae in wave-exposed locations; *Sargassum*; rubble in intertidal zone.

***Tanystylum dowi* CHILD 1979**

- 1979 *Tanystylum dowi* CHILD, Smiths. Contr. Zool., **293**: 2626-28, fig. 9a-g.

Distribution: Panamá Pacific.

Depth range: intertidal.

Habitat: Abietinaria.

***Tanystylum duospinum* HILTON 1939**

- 1939 *Tanystylum duospinum* HILTON, J. Ent. Zool. Pomona College, **31** (2): 33.
- 1990 *Tanystylum duospinum*, - RUSSEL & HEDGPETH, Bijdr. Dierk., **60** (3/4): 215-224, figs. 1-2 (biology).

Distribution: California.

Habitat: on hydroids; ectoparasitic larvae.

***Tanystylum evelinae* MARCUS 1940**

- 1940 *Tanystylum evelinae* MARCUS, Bolm. Fac. Fil. Cienc. Letr., Univ. S. Paulo, **4**: 99-102, fig. 13.
- 1992 *Tanystylum evelinae*, - STOCK, Tijdschr. Ent., **135**: 119, 139.

Distribution: Brazil.

Depth range: infralittoral.
Habitat: seaweeds on rocky shore.

Tanystylum excratum STOCK 1954

- 1954 *Tanystylum excratum* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 146-149, figs. 71-72.
 1977 *Tanystylum excratum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 323-327, figs. 14-17.

Distribution: New Zealand.
Depth range: 1 fathom-47m.
Habitat: in mud.

Tanystylum geminum STOCK 1954

- 1954 *Tanystylum geminum* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 120-122, fig. 26.
 1955 *Tanystylum geminum*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 600-601, pl. 3, figs. 5-7.
 1975 *Tanystylum geminum*, - STOCK, Bull. mar. Sci., **24** (4): 983-984, fig. 10.
 1979 *Tanystylum geminum*, - CHILD, Smiths. Contr. Zool., **293**: 28.
 1979 *Tanystylum geminum*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 13, fig. 9h-i.
 1982 *Tanystylum geminum*, - STOCK, Monit. zool. Ital., Suppl. **17** (7): 187-188.
 1986 *Tanystylum geminum*, - STOCK, Bull. mar. Sci. **38** (3): 402 tab. 1.
 1990 *Tanystylum geminum*, - MÜLLER, Bonn. zool. Beitr. **41** (3-4): 279-280.
 1992 *Tanystylum geminum*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 45.

Distribution: tropical W-Atlantic, Caribbean Sea and Gulf of Mexico.

Depth range: 0-4m.

Habitat: in a variety of substrata, often in algae and seagrass (*Thalassia*).

Tanystylum haswelli CHILD 1990

- 1990 *Tanystylum haswelli* CHILD, Proc. biol. Soc. Wash., **103** (2): 317-319, fig. 2.

Distribution: Australia (Lizard Island).

Depth range: 2m.

Habitat: coral reefs; rubble bank on barrier reef.

Tanystylum hooperi CLARK 1977

- 1977 *Tanystylum hooperi* CLARK, J. Roy. Soc. N.Z., **7** (3): 326-327, figs. 20-30.
 1980 *Tanystylum hooperi*, - HOOPER, Aust. Zool., **20** (3): 474.

Distribution: Australia.

Depth range: 0-2.5m.

Habitat: *Sargassum*.

Tanystylum hummeli STOCK 1954

- 1954 *Tanystylum hummeli* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., 5 (24): 122-125, figs. 27-28.
 1986 *Tanystylum hummeli*, - STOCK, Bull. mar. Sci., 38 (3): 402 tab. 1.

Distribution: Venezuela; Caribbean Sea of Colombia.

Depth range: 1-2m.

Habitat: sandy debris.

Tanystylum intermedium COLE 1904

- 1904 *Tanystylum intermedium* COLE, -Harriman Alaska Exped., 10: 278-280, pl. 21, figs. 7-9, pl. 22, figs. 1-7, pl. 23, figs. 1-3.
 1913 *Tanystylum intermedium*, - HALL, Univ. Calif. Publ. Zool., 11 (6): 132.
 1934 *Tanystylum intermedium*, - SCHMITT, J. Wash. Acad. Sci., 24 (1): 69.
 1941 *Tanystylum intermedium*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., 9 (26): 255.
 1942 *Tanystylum panamum* HILTON, J. Ent. Zool. Pomona College, 34 (3): 69-70.
 1955 *Tanystylum intermedium*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., 117: 249.
 1961 *Tanystylum intermedioides* HEDGPETH, Lunds Univ. Arsskr., N.F., 2 (57): 14-16, fig. 10.
 1975 *Tanystylum intermedium*, - STOCK, Ent. Ber., 35: 71-74, figs. 10-15.
 1976 *Tanystylum intermedium*, - BIRKELAND, REIMER & YOUNG, Rep. Mar. Fauna and Benthic Shelf-slope communities of the Isthmian Region (Univ. Miami): 157.
 1992 *Tanystylum intermedium*, - CHILD, Smiths. Contr. Zool., 526: 21.

Distribution: California; Mexico Pacific; Panamá Pacific; Ecuador; Galapagos Archipelago; Peru; Chile.

Depth range: 0-18m.

Habitat: on weeds from rocky coast; *Abietinaria*; from piers in harbours and bays; on rocks and among coral rubble in intertidal zone.

Tanystylum isabellae MARCUS 1940

- 1940 *Tanystylum isabellae* MARCUS, Bolm. Fac. Fil. Cién. Letr. Univ. S. Paulo, 10 (Zool.) 4: 102.
 1943 *Tanystylum isabellae*, - HEDGPETH, Proc. New England Zool. Club, 22: 55.
 1949 *Tanystylum isabellae*, - PAGE, Arch. Zool. Exp. Gen., 86 (Notes & Rev.), 2: 29.
 1950 *Tanystylum isabellae*, - SAWAYA, Mem. Inst. Oswaldo Cruz, 47 (1-2): 73.
 1955 *Tanystylum tubirostre*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., 19 (4): 600.
 1955 *Tanystylum isabellae*, - STOCK, Vidensk. Medd. Dansk naturhist. Foren., 117: 247.
 1986 *Tanystylum isabellae*, - STOCK, Bull. mar. Sci., 38 (3): 402 tab. 1.
 1992 *Tanystylum isabellae*, - STOCK, Tijdschr. Ent., 135: 119-120, 139, figs. 22-26.

Distribution: amphi-atlantic; Eastern Caribbean Islands south to Brazil; in the E-Atlantic known from Kongo.

Depth range: 0-37m

Habitat: on seaweeds on rocky shore.

***Tanystylum isthmiacum* STOCK 1955**

- 1955 *Tanystylum isthmiacum* STOCK, Vidensk. Medd. Dansk naturhist. Foren., **117**: 247-249, fig. 17.
 1966 *Tanystylum isthmiacum difficile*, - FAGE & STOCK, Ann. Inst. Oceanogr. Monaco **44**: 318.
 1966 *Tanystylum isthmiacum difficile*, - STOCK, Ann. Inst. Oceanogr. Monaco., **44**: 389-391, figs. 2a-h.
 1975 *Tanystylum isthmiacum isthmiacum*, - STOCK, Ent. Ber., **35** (5): 74, figs. 16-21.
 1979 *Tanystylum isthmiacum difficile*, - CHILD, Smiths. Contr. Zool., **293**: 29.
 1979 *Tanystylum isthmiacum isthmiacum*, - CHILD, Smiths. Contr. Zool., **293**: 29.
 1979 *Tanystylum isthmiacum difficile*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 11-13, fig. 3b-f.
 1986 *Tanystylum isthmiacum difficile*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.
 1990 *Tanystylum isthmiacum*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 280.

Distribution: In the E-Pacific recorded from Mexico, Costa Rica, Panamá and Ecuador; in the W-Atlantic known from Panamá, Curaçao, Martinique and Brazil; one E-Atlantic record from the Cape Verde Islands.

Depth range: 0-51m.

Habitat: in a variety of substrata.

***Tanystylum malpelensis* CHILD 1979**

- 1979 *Tanystylum malpelensis* CHILD, Smith. Contr. Zool., **293**: 30-32, fig. 10.
 1992 *Tanystylum malpelensis*, - CHILD, Smiths. Contr. Zool., **526**: 21.

Distribution: Colombia Pacific (Isla Malpelo); Galapagos Archipelago.

Depth range: 0-36m.

Habitat: rocks in intertidal zone.

***Tanystylum mexicanum* CHILD 1979**

- 1979 *Tanystylum mexicanum* CHILD, Smiths. Contr. Zool., **293**: 32-34, fig. 11a-h.

Distribution: Mexico Pacific (Gulf of California).

Depth range: 18-64m.

***Tanystylum neorhetum* MARCUS 1940**

- 1889 *Clotenia dohrnii* PFEFFER, Jb. Hamburg. Wiss. Anst., **6** (2): 48.
 1913 *Tanystylum pfefferi* BOUVIER, Deuxième Exped. Antarct. Française (1908-1910),

- 6:** 5, 45, 122.
- 1932 *Tanystylum pfefferi*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 26-28, figs. D1-4.
- 1932 *Tanystylum pfefferi*, - GORDON, Ann. Mag. Nat. Hist., (10) **10**: 90-93, figs. 3-4.
- 1932 *Tanystylum pfefferi*, - GORDON, Discovery Rep., **6**: 119-122, figs. 65a, 66-68.
- 1938 *Tanystylum pfefferi*, - GORDON, Sci. Rep. Australas. Antarct. Exp. (C) (Zool. & Bot.), **2** (8): 27.
- 1940 *Tanystylum neorhetum* MARCUS, Bol. Fac. Fil, Ciênc. Letr., Univ. S. Paulo, **19** (Zool. 4): 95-96, 129.
- 1944 *Tanystylum neorhetum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 67.
- 1955 *Tanystylum neorhetum*, - STOCK, Res. Norw. Sci. Exped. Tristan da Cunha, 1937-1938, **33**: 3.
- 1957 *Tanystylum neorhetum*, - STOCK, Mitt. Zool. Mus. Hamburg., **55**: 98.
- 1961 *Tanystylum neorhetum*, - HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 12.
- 1962 *Tanystylum neorhetum*, - KENNY & HAYSON, Pac. Sci., **16** (3): 260.
- 1969 *Tanystylum neorhetum*, - ARNAUD, Les Terres Austr. Françaises, **47**: 37.
- 1971 *Tanystylum neorhetum*, - STOCK, Com. natn. français rech. Antarct., **30**: 129.
- 1972 *Tanystylum neorhetum*, - ARNAUD, Bull. Mus. natn. Hist. nat., Paris, (3) **51** (65): 809-811.
- 1977 *Tanystylum neorhetum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 327-330, figs. 31-39.

Distribution: New Zealand; South Georgia; E-Patagonian Bank; Kerguelen Islands, Macquarie Island. Tristan da Cunha.

Depth range: 0-46m.

Habitat: under stones; on *Melobesia*.

***Tanystylum nesiotes* CHILD 1970**

- 1970 *Tanystylum nesiotes* CHILD, Proc. biol. Soc. Wash., **83** (27): 299-302, fig. 4.
- 1989 *Tanystylum nesiotes*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 126.

Distribution: Society Islands (Tahiti, Moorea, Bora Bora).

Depth range: 0-11m.

Habitat: coral reefs; algae and dead corals.

***Tanystylum occidentalis* (COLE 1904)**

- 1904 *Clotenia occidentalis* COLE, Harriman Alaska Exped., **10**: 281-282, pl. XIII, fig. 9, pl. XXIII, figs. 4-8.

Distribution: Pacific Grove, California.

***Tanystylum oculospinosum* HILTON 1942**

- 1942 *Tanystylum oculospinosum* HILTON, Allan Hancock Pac. Exped., **5** (9): 70.
- 1955 *Tanystylum oculospinosum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 248-249.
- 1966 *Tanystylum oculospinosum*, - STOCK, Ann. Inst. Océanogr., Monaco, **44**: 390.
- 1979 *Tanystylum oculospinosum*, - CHILD, Smiths. Contr. Zool., **293**: 34.

- 1992 *Tanystylum oculospinosum*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 35-37, fig. 16.
 1992 *Tanystylum oculospinosum*, - CHILD, Smiths. Contr. Zool., **526**: 23-24.

Distribution: In the West-Atlantic from Bermuda to the Caribbean. In the E-Pacific known from Panamá, Peru, Ecuador, Baja California and the Galapagos Archipelago.

Depth range: intertidal-10 m.

Habitat: shallow shore localities and *Sargassum*.

***Tanystylum oedinotum* LOMAN 1923**

- 1923 *Tanystylum oedinotum* LOMAN, Further Zool. Res. Swed. Antarct. Exped. (1901-1903), **1** (2): 29-30, fig. F1-5.
 1932 *Tanystylum oedinotum*, - GORDON, Discovery Rep., **6**: 119-122, figs. 65a, 66-68.
 1938 *Tanystylum pfefferi*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (c), (Zool. & Bot.), **2** (8): 27 [part.].
 1972 *Tanystylum oedinotum*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 811-812, figs. 2-6.
 1977 *Tanystylum oedinotum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 330-332, figs. 44-50.

Distribution: Macquarie Island.

Depth range: 0-60m.

Habitat: sand; hydroids and *Mytilus* clumps; debris on stones.

***Tanystylum orbiculare* WILSON 1878**

- 1844 ?*Pasithoe umbonata* GOULD, Proc. Soc. Nat. Hist., Boston[?], **1**: 92-93.
 1878 *Tanystylum orbiculare* WILSON, Trans. Connecticut Acad. Arts Sci., **5**: 5-7, fig. 2a-f.
 1880 *Tanystylum orbiculare*, - WILSON, Rep. U.S. Commissioner Fish., 1878 (6): 471-473, pl. 3, fig. 11.
 1891 *Tanystylum orbiculare*, - MORGAN, Stud. Biol. Lab. Johns Hopkins Univ., Baltimore, **5** (1): 37-49 [embryology].
 1913 *Tanystylum orbiculare*, - SUMNER, OSBURN & COLE, Bull. Bureau Fish., Washington, **31** (1, 2): 141-142.
 1925 *Tanystylum orbiculare*, - FISH, Bull. Bureau Fish., Washington, **41**: 161.
 1929 *Clotenia orbiculare*, - GILTAY, Bull. & Ann. Soc. Roy. Ent. Belg., **69**: 175.
 1940 *Tanystylum orbiculare*, - MARCUS, Bolm. Fac. Fil., Ciênc. Letr., Univ. S. Paulo, **19** (Zool. 4): 105-108, pl. XV, fig. 15a-h.
 1943 *Tanystylum orbiculare*, - HEDGPETH, Proc. New England Zool. Club, **22**: 54-55.
 1948 *Tanystylum orbiculare*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 266-268, figs. 8, 49a.
 1949 *Tanystylum orbiculare*, - FAGE, Arch. Zool. Exp. Gen., 86 (Notes & Rev.), **2**: 29-30, fig. 6.
 1950 *Tanystylum orbiculare*, - SAWAYA, Mem. Inst. Oswaldo Cruz, **47** (1-2): 73.
 1951 *Tanystylum orbiculare*, - STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **2** (43): 18-20.
 1952 *Tanystylum orbiculare*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (6): 530.

- 1954 *Tanystylum orbiculare*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 117.
- 1954 *Tanystylum orbiculare*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 145.
- 1955 *Tanystylum orbiculare*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 599.
- 1958 *Tanystylum orbiculare*, - Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
- 1958 *Tanystylum orbiculare*, - STOCK, Bull. Res. Council Israel, (B. Zool.), **7B** (3-4): 138.
- 1962 *Tanystylum orbiculare*, - STOCK, Ent. Ber., **22** (11): 219.
- 1963 *Tanystylum orbiculare*, - CLARK, Rec. Austr. Mus., **26** (1):
- 1966 *Tanystylum orbiculare*, - FAGE & STOCK, Ann. Inst. Océanogr. Monaco, **44**: 319.
- 1966 *Tanystylum orbiculare*, - STOCK, Atlantide Rep., **9**: 46.
- 1973 *Tanystylum orbiculare*, - KRAEUTER, J. nat. Hist., **7** (5): 496.
- 1973 *Tanystylum orbiculare*, - KRAPP, Beaufortia, **21** (277): 60, 64, fig. 4.
- 1975 *Tanystylum orbiculare*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 17, 18-20 (tab. II) (ecology).
- 1975 *Tanystylum orbiculare*, - STOCK, Bull. mar. Sci., **24** (4): 985.
- 1976 *Tanystylum orbiculare*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.
- 1976 *Tanystylum orbiculare*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27** (3/4): 343.
- 1977 *Tanystylum orbiculare*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 332.
- 1981 *Tanystylum orbiculare*, - MUNILLA & de HARO, Bijdr. Dierk., **51** (2): 191-197, figs. 5, 11, tab. I (electrophoretical and immunological study).
- 1983 *Tanystylum orbiculare*, - KRAPP, Bonn. zool. Beitr., **34** (1-3): 409, 412.
- 1984 *Tanystylum orbiculare*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 531, 533-535, tab. III-IV.
- 1986 *Tanystylum orbiculare*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.
- 1987 *Tanystylum orbiculare*, - ARNAUD, Mesogee, **47**: 43-44.
- 1992 *Tanystylum orbiculare*, - CHILD, Mem. Hourglass Cruises, **9** (1): 8 [key], 9 [tab. 3], 33-35, fig. 15.
- 1992 *Tanystylum orbiculare*, - STOCK, Tijdschr. Ent., **135**: 121, 139.

Distribution: Massachusetts to Brazil, Indo-Pacific; ?Mediterranean.

Depth range: intertidal-?7m.

Habitat: rocky shore; *Peyssonnelia squamaria*, *Sargassum*.

Tanystylum ornatum FLYNN 1928

- 1928 *Tanystylum ornatum* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., I): 33-34, figs. 20-21.
- 1954 *Tanystylum ornatum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 148-149.
- 1977 *Tanystylum ornatum*, - CLARK, J. Roy. Soc. N.Z., **7** (3): 332.

Distribution: South Africa.

Depth range: 46-55m.

Tanystylum philippinensis CHILD 1988

- 1988 *Tanystylum philippinensis* CHILD, Smiths. Contr. Zool., **468**: 10-12, fig. 4.

Distribution: Philippines.
Depth range: 15-20m.

Tanystylum rehderi CHILD 1970

- 1970 *Tanystylum rehderi* CHILD, Proc. biol. Soc. Wash., **83** (27): 302-306, fig. 5.
 1983 *Tanystylum rehderi*, - CHILD, Proc. biol. Soc. Wash., **96** (4): 705.
 1988 *Tanystylum rehderi*, - CHILD, Bull. biol. Soc. Wash., **8**: 53-54.
 1989 *Tanystylum rehderi*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 126, figs. 22-39.
 1991 *Tanystylum rehderi*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 142.

Distribution: Society Islands (Bora Bora, Moorea);
 Seychelles (Aldabra Atoll); Koror; Guam.

Depth range: 0-12m.

Habitat: coral reefs; dead corals.

Tanystylum scrutator STOCK 1954

- 1954 *Tanystylum scrutator* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 142-145, fig. 70.
 1971 *Tanystylum scrutator*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 337.
 1983 *Tanystylum scrutator*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 41.
 1986 *Tanystylum scrutator*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 50.
 1991 *Tanystylum scrutator*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 24-25.

Distribution: Japan and Korea.

Depth range: ?0-40 m.

Habitat: Corallina.

Tanystylum styligerum (MIERS 1875)

- 1875 *Nymphon styligerum* MIERS, Ann. Mag. Nat. Hist., **16** (4): 76.
 1879 *Tanystylum styligerum*, - MIERS, Phil. Trans. Roy. Soc. London, **168**: 213, pl. XI, figs. 9-9d.
 1907 *Tanystylum longicaudatum* HODGSON, Ergebn. Hamburg. Magalhaens. Sammelreise 1892-93, **2**: 13.
 1915 *Tanystylum styligerum*, - HODGSON, Ann. Mag. Nat. Hist., (8) **15** (85): 148.
 1923 *Tanystylum kentrodes* LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 26-28, fig. E1-4.
 1927 *Tanystylum styligerum*, - Dt. Südpolar Exped., **19** (Zool. II): 117.
 1932 *Tanystylum styligerum*, - GORDON, Ann. Mag. Nat. Hist., **10** (10): 88-90, figs. 1-2
 1932 *Tanystylum styligerum*, - GORDON, Discovery Rep., **6**: 118-119, fig. 65b.
 1938 *Tanystylum styligerum*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 26-27.
 1940 *Tanystylum styligerum*, - MARCUS, Bolm. Fac. Fil., Ciênc. Letr., Univ. S. Paulo, **19** (Zool. 4): 96.
 1944 *Tanystylum styligerum*, - GORDON, Re. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 67.
 1961 *Tanystylum styligerum*, - HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 12.
 1962 *Tanystylum styligerum*, - KENNY & HAYSON, Pac. Sci., **16** (3): 260.
 1972 *Tanystylum styligerum*, - ARNAUD, Bull. Mus. natn. Hist. nat., Paris, (3) **51**

(65): 809.

- 1977 *Tanystylum styligerum*, - CLARK, J. Roy. Soc. N.Z., 7 (3): 333-336, figs. 51-61, figs. 51-61.

Distribution: Straits of Magellan; Kerguelen Islands; Macquarie Island; Falkland Islands.

Depth range: 0-91m.

Habitat: *Macrocystis pyrifera*.

***Tanystylum thermophilum* (BARNARD 1946)**

- 1946 *Austroraptus thermophilus* BARNARD, Ann. Magh. Nat. Hist., 13 (11): 62.
 1954 *Austroraptus thermophilus*, - BARNARD, Ann. S. Afr. Mus., 41 (3): 146-148, figs. 29-30.
 1956 *Tanystylum thermophilum* STOCK, Vidensk. Medd. dansk naturhist. Foren., 118: 92-93.
 1957 *Tanystylum thermophilum*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., 55: 94-95, fig. 12a-c.
 1962 *Tanystylum thermophilum*, - STOCK, Trans. Roy. Soc. S. Afr., 36 (4): 277-278.
 1977 *Tanystylum thermophilum*, - CLARK, J. Roy. Soc. N.Z., 7 (3): 336.
 1988 *Tanystylum thermophilum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., 98 (6): 130.

Distribution: South Africa; Macquarie Island.

Depth range: 62-71m-?>100m.

Habitat: rocks.

***Tanystylum tubirostrum* STOCK 1954**

- 1954 *Tanystylum tubirostre* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., 5: 117-120, figs. 24-25.
 1955 non *Tanystylum tubirostre* (= *isabellae*), - BOURDILLON, Rev. Trav. Inst. Pêches Marit., 19 (4): 600, pl. 3, figs. 2-4.
 1975 *Tanystylum tubirostrum*, - STOCK, Bull. mar. Sci., 24 (4): 984.
 1979 *Tanystylum tubirostrum*, - CHILD, Smiths. Contr. Zool., 293: 34-35.
 1982 *Tanystylum tubirostrum*, - CHILD, Smiths. Contr. mar. Sci., 12: 363.
 1986 *Tanystylum tubirostrum*, - STOCK, Bull. mar. Sci., 38 (3): 402 tab. 1.
 1990 *Tanystylum tubirostrum*, - MÜLLER, Bonn. zool. Beitr., 41 (1): 67-68, figs. 16-20.

Distribution: tropical W-Atlantic and Kenya.

Depth range: 0-1m.

Habitat: in a variety of substrata.

***Tanystylum ulreungum* KIM 1983**

- 1983 *Tanystylum ulreungum* KIM, J. Kangreung Natn. Univ., 5 (2): 467, figs. 1-2.
 1983 *Tanystylum nabetensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., 386: 39-41, fig. 13.
 1986 *Tanystylum ulreungum*, - KIM & HONG, Kor. J. Syst. Zool., 2 (2): 50.
 1987 *Tanystylum ulreungum*, - HONG & KIM, Kor. J. Syst. Zool., 3 (2): 152.

Distribution: Japan; Korea.

Depth range: 7-15m.

***Trygaeus* DOHRN 1881**

1881 *Trygaeus* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 164 [diagnosis].

***Trygaeus communis* DOHRN 1881**

- 1881 *Trygaeus communis* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 164-168, pl. IX, figs. 6-14, pl. X, figs. 1-5.
 1909 *Trygaeus communis*, - d'ARCY THOMPSON, The Cambridge Nat. Hist.: 535.
 1923 *Trygaeus communis*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10) **6** (3, 4): 122.
 1923 *Trygaeus communis*, - BOUVIER, Faune de France, **7**: 58.
 1929 *Trygaeus communis*, - GILTAY, Bull. & Ann. Soc. Ent. Belg., **69**: 176.
 1940 *Trygaeus communis*, - FARAGGIANA, Boll. Mus. Zool. Anat. Comp. Reale Univ. Torino, (3) **48** (112): 12.
 1954 *Trygaeus communis*, - BOURDILLON, Rec. Trav. Stat. Mar. d'Endoume, **12** (7): 155.
 1968 *Trygaeus communis*, - STOCK, Vie Milieu, Suppl. **19** (1A): 19-20, fig. 13.
 1973 *Trygaeus communis*, - KRAPP, Bonn. zool. Beitr., **21** (277): 64.
 1975 *Trygaeus communis*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 14 fig. 4, 17, 18-20 (tab. II), 21-23 (ecology).
 1979 *Trygaeus communis*, - CHIMENEZ, BRIGNOLI & BASCIANO, Cah. Biol. Mar., **20** (4): 489.
 1984 *Trygaeus communis*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 534-535, tabs. III-IV.
 1987 *Trygaeus communis*, - ARNAUD, Mesogee, **47**: 44.
 1989 *Trygaeus communis*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 287 [tab. 1], 288 (phenology).

Distribution: Mediterranean.

Depth range: 0.5-3m.

Habitat: algae. - (*Cutleria*; *Cystoseira*; *Halopteris scoparia*; *Padina pavonia*; *Peyssonelia squammaria*; *Sargassum*) and on hydroids (*Aglaophenia*).

AUSTRODECIDAE

- 1954 Austrodecidae STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 151 [diagnosis, remarks].
 1957 Austrodecidae, - Stock, Beaufortia, **68** (6): 1-81 [revision of family].

***Austrodecus* HODGSON 1907**

- 1954 *Austrodecus*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 151 [remarks, key to species].
 1991 *Austrodecus*, - Stock, Mem. Mus. natn. Hist. nat., (A), **151**: 170 [diagnosis of subgenus *Tubidecus*].

Austrodecus aconae (HEDGPETH & McCAIN 1971)

1971 *Pantopipetta aconae* HEDGPETH & McCAIN, in: LLANO & WALLEN (eds): Biology of the Antarctic Seas, Antarct. Res. Ser., **17**: 227-228, figs 1, 6, tab. 4.

Distribution: NE-Pacific.
Depth range: 1400-1420m.

Austrodecus bathyale STOCK 1991

1991 *Austrodecus (Tubidecus) bathyale* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 174-175, fig. 34.

Distribution: Between New Caledonia and Loyality Islands.
Depth range: 1590-1665 m.

Austrodecus breviceps GORDON 1938

1938 *Austrodecus breviceps* GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 25-26, figs. 7-8.
 1950 *Austrodecus breviceps*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 154.
 1957 *Austrodecus breviceps*, - STOCK, Beaufortia, **6** (68): 59-61, figs. 30, 31b, 32c-d.
 1971 *Austrodecus breviceps*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (3-4): 449-450, fig. 11A-E.
 1972 *Austrodecus breviceps*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 806.

Distribution: New Zealand (Antipodes Islands); Kerguelen Islands.

Depth range: 0-140m.
Habitat: pools.

Austrodecus calcaricauda STOCK 1957

1957 *Austrodecus calcaricauda* STOCK, Beaufortia, **6** (68): 34-35, fig. 13.

Distribution: South Georgia (Cumberland Bay).
Depth range: 230-250m.

Austrodecus calvum STOCK 1991

1991 *Austrodecus calvum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 168-170, fig. 31.

Distribution: New Caledonia.
Depth range: 1490-1620m.

Austrodecus confusum STOCK 1957

1957 *Austrodecus confusum* STOCK, Beaufortia, **6** (68): 69-71, fig. 38.

Distribution: Three Kings, north of New Zealand.

Depth range: 119m..

Habitat: hard bottom.

***Austrodecus conifer* STOCK 1991**

1991 *Austrodecus conifer* STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 139-141, figs. 1-7.

Distribution: NE-Atlantic (banc Galice).

Depth range: 675-685m.

***Austrodecus curtipes* STOCK 1957**

1957 *Austrodecus curtipes* STOCK, Beaufortia, **6** (68): 36-39, fig. 14.

1961 *Austrodecus curtipes*, - HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 16.

1969 *Austrodecus curtipes*, - HEDGPETH, Antarct. Map Folio Ser., **11**: pl. 14.

1972 *Austrodecus curtipes*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 806-807.

Distribution: Tierra del Fuego; Falkland Islands; Kerguelen Islands.

Depth range: 0-130m.

Habitat: *Macrocystis pyrifera*.

***Austrodecus elegans* STOCK 1957**

1957 *Austrodecus elegans* STOCK, Beaufortia, **6** (68): 68-69, figs. 36, 37a, c.

Distribution: just north of Marion Island (Prince Edward group).

Depth range: 99-113m.

***Austrodecus enzoi* CLARK 1971**

1971 *Austrodecus enzoi* CLARK, N.Z. J. Mar. Fresh. Res., **5** (2): 338-340, fig. 3H-M.

1991 *Austrodecus enzoi*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 170.

Distribution: New Zealand (Snares Islands).

Depth range: 81m.

Habitat: coarse shelly sand.

***Austrodecus excelsum* STOCK 1991**

1991 *Austrodecus (Tubidecus) excelsum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 170-172, fig. 32.

Distribution: New Caledonia.
Depth range: 440m.

Austrodecus fagei STOCK 1957

1957 *Austrodecus fagei* STOCK, Beaufortia, **6** (68): 3942, figs. 15-16.

Distribution: Eastern part of Antarctic continent.
Depth range: 26-393m.

Austrodecus frigorifugum STOCK 1954

1954 *Austrodecus frigorifugum* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 156-159, fig. 77.
 1957 *Austrodecus frigorifugum*, - STOCK, Beaufortia, **6** (68): 72-73, fig. 40.
 1968 *Austrodecus frigorifugum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 28-29, fig. 10a.

Distribution: New Zealand (Three Kings Island); Raoul Island; Kermadec Islands.

Depth range: 60-119m.

Habitat: hard bottom; stones.

Austrodecus glabrum STOCK 1957

1957 *Austrodecus glabrum* STOCK, Beaufortia, **6** (68): 42-45, figs. 17-18.

Distribution: Shelf around South Georgia.
Depth range: 18-250m.

Austrodecus glaciale HODGSON 1907

1907 *Austrodecus glaciale* HODGSON, Rep. natn. Antarct. Exp., 1901-1904, Nat. Hist., **3**: 53, pl. VIII, fig. 1.
 1927 *Austrodecus glaciale*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 347-349.
 1932 *Austrodecus glaciale*, - GORDON, Discovery Rep., **6**: 115-117, figs. 63-64.
 1944 *Austrodecus glaciale*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B) **5** (1): 62-68, figs. 24-25.
 1950 *Austrodecus glaciale*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 154.
 1952 *Austrodecus glaciale*, - Bull. Mus. natn. Hist. nat., Paris, (2) **24** (2): 185.
 1952 *Austrodecus glaciale*, - Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 272.
 1954 *Austrodecus glaciale*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 14, 159.
 1957 *Austrodecus glaciale*, - STOCK, Beaufortia, **6** (68): 45-50, figs. 19-21.

Distribution: Antarctic continent, probably circumpolar.
Depth range: 0-640m.
Habitat: mud.

Austrodecus gordonae STOCK 1954

- 1954 *Austrodecus gordonae* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 153-156, figs. 75-76.
 1957 *Austrodecus gordonae* STOCK, Beaufortia, **6** (68): 73-75, figs. 41-42.

Distribution: New Zealand (Three Kings Island).
Depth range: 119m.
Habitat: hard bottom.

***Austrodecus goughense* STOCK 1957**

- 1957 *Austrodecus goughense* STOCK, Beaufortia, **6** (68): 63-65, fig. 34.
 1974 *Austrodecus goughense*, - ARNAUD, Tethys, **5** (4): 781.

Distribution: St. Paul and Amsterdam Islands; Gough Island.
Depth range: 50-120m.
Habitat: coralline algae; bryozoans.

***Austrodecus kelpi* PUSHKIN 1977**

- 1977 *Austrodecus kelpi* PUSHKIN, Expl. Fauna Seas, **21** (24): 56-58, fig. 3.
 1991 *Austrodecus kelpi*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 170 [text].

Distribution: Antarctica (off South Georgia).
Depth range: shallow water.
Habitat: among algae (kelps).

***Austrodecus latum* STOCK 1991**

- 1991 *Austrodecus (Tubidecus) latum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 174-177, fig. 35.

Distribution: Between New Caledonia and Loyality Islands.
Depth range: 570-595 m.

***Austrodecus longispinum* STOCK 1957**

- 1957 *Austrodecus longispinum* STOCK, Beaufortia, **6** (68): 50-52, figs. 22-23.

Distribution: Shelf of Kerguelen Island.
Depth range: 91-150m.

***Austrodecus minutum* CLARK 1971**

- 1971 *Austrodecus minutum* CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 336-338, fig. 3A-G.

Distribution: New Zealand (Snares Islands).
Depth range: 132m.
Habitat: sand with shells.

Austrodecus oblongus PUSHKIN 1977

1977 *Austrodecus oblongus* PUSHKIN, Expl. Fauna Seas, **21** (24): 59, fig. 4.
 1991 *Austrodecus oblongus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 170
 [text].

Distribution: Antarctica.
Depth range: 100-140m.

Austrodecus palauense CHILD 1983

1983 *Austrodecus palauense* CHILD, Proc. biol. Soc. Wash., **96** (4): 711-713, fig. 5.

Distribution: Palau, SE Koror Island.
Depth range: 0.1m.
Habitat: sea reef inside cove next to cave, from sponges.

Austrodecus pentamerum STOCK 1968

1968 *Austrodecus pentamerum* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 26-28, fig. 9.

Distribution: Raoul Islan;, Kermadec Islands.
Depth range: 60m.
Habitat: stones.

Austrodecus profundum STOCK 1957

1957 *Austrodecus profundum* STOCK, Beaufortia, **6** (68): 52-54, fig. 24.

Distribution: Graham Region of the Antarctic zone.
Depth range: 920m.
Habitat: mud and stones.

Austrodecus simulans STOCK 1957

1957 *Austrodecus simulans* STOCK, Beaufortia, **6** (68): 54-56, figs. 25, 26, 37b, d.

Distribution: Shelf around Kerguelen.
Depth range: 91-150m.

Austrodecus sinuatum STOCK 1957

1957 *Austrodecus sinuatum* STOCK, Beaufortia, **6** (68): 56-57, fig. 27.

Distribution: Auckland Islands (Masked Island, Carnley Harbour).

Habitat: rocky coast, from Melobesia.

Austrodecus staplesi STOCK 1990

1990 *Austrodecus staplesi* STOCK, Tijdschr. Ent., **133**: 269-272, figs. 1-9.

Distribution: Australia (New South Wales, Split Solitary Island).

Depth range: 12-14m.

Habitat: "rocky area with small caves, some corals, algae and some sand" (STOCK).

Austrodecus stocki CHILD 1988

1988 *Austrodecus stocki* CHILD, Bull. biol. Soc. Wash., **8**: 54-55, fig. 1.

Distribution: Seychelles (Aldabra Atoll).

Depth range: 20m.

Habitat: "reef slope, coral rubble in hollows between coral heads" (CHILD).

Austrodecus tristanense STOCK 1955

1955 *Austrodecus tristanense* STOCK, Res. Norw. Sci. Exped. Tristan da Cunha 1937-1938, **33**: 5-7, fig. 2.

1957 *Austrodecus tristanense* STOCK, Beaufortia, **6** (68): 66-67, fig. 35.

Distribution: Tristan da Cunha.

Depth range: 0-55m.

Habitat: from *Lithothamnion* in surf zone; algae and sand; intertidal, under stones; from rocks covered with *Lithothamnion* and *Balanus*; hard bottom with algae.

Austrodecus tuberculatum STOCK 1991

1991 *Austrodecus (Tubidecus) tuberculatum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 172-174, fig. 33.

Distribution: New Caledonia.

Depth range: 440-610 m.

Austrodecus tubiferum STOCK 1957

1954 *Austrodecus gordonae* (part), - STOCK, Vidensk. Medd. dansk naturhist. Foren, **116**: 153.

1957 *Austrodecus tubiferum* STOCK, Beaufortia, **6** (68): 75-77, fig. 43.

1971 *Austrodecus tubiferum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 337.

1988 *Austrodecus tubiferum*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 668.

1991 *Austrodecus tubiferum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 35.

Distribution: Japan (Okinose, Sagami Bay).
Depth range: 37-187m.
Habitat: hard bottom.

Austrodecus valdiviensis TURPAEVA 1990

1990 *Austrodecus valdiviensis* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **126**: 116-118, fig. 5

Distribution: southern Atlantic.
Depth range: 620m.

Pantopipetta STOCK 1963

1963 *Pantopipetta* STOCK, Ann. S. Afr. Mus., **46** (12): 184 [diagnosis, remarks], 185 [key to species].
 1971 *Pantopipetta*, - HEDGPETH & McCAIN, in: G. Llano & Wallen: Biology of the Anarct. Seas, 4. Antarct. Res. Ser., **17**: 217-222 [diagnosis, remarks], 222-223 [key to species].
 1981 *Pantopipetta*, - STOCK, Cah. Biol. Mar., **22**: 465-466 [remarks on longituberculata-complex].
 1988 *Pantopipetta*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 131-132 [key to species].

Pantopipetta angusta STOCK 1981

1981 *Pantopipetta angusta* STOCK, Cah. Biol. Mar., **22**: 468-470, fig. 8.
 1990 *Pantopipetta angusta* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **126**: 119-123.

Distribution: southern Atlantic; Walvis Basin.
Depth range: 4560-5218m.

Pantopipetta armata ARNAUD & CHILD 1988

1988 *Pantopipetta armata* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 132-134, fig. 2.

Distribution: South Africa.
Depth range: 680-1300m.

Pantopipetta armoricana STOCK 1978

1978 *Pantopipetta armoricana* STOCK, Cah. Biol. Mar., **19**: 206-208, fig. 7.
 1991 *Pantopipetta armoricana*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 138.

Distribution: NE Atlantic.
Depth range: 180-1210m.

Pantopipetta australis (HODGSON 1914)

- 1914 *Pipetta australis* HODGSON, Zool. Anz., **45** (4): 159.
 1915 *Pipetta australis*, - HODGSON, Ann. Mag. Nat. Hist., (8) **15** (85): 141-142.
 1927 *Pipetta australis*, - HODGSON, Dt. Südpolar-Exped., **19** (Zool. 2): 314-315,
 fig. 2.
 1955 *Pipetta australis*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad.
 Nauk SSSR, **12**: 327.
 1971 *Pantopipetta australis*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**.
 Antarct. Res. Ser., 17: 223, figs. 1-2.

Distribution: Antarctica.

Depth range: 2450-3725m.

Pantopipetta auxiliata STOCK 1968

- 1968 *Pantopipetta auxiliata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**:
 24-26, fig. 8.
 1971 *Pantopipetta auxiliata*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**.
 Antarct. Res. Ser., 17: 219-221, 223 [key].
 1988 *Pantopipetta auxiliata*, - CHILD, Bull. biol. Soc. Wash., **8**: 55-56.

Distribution: Seychelles (Aldabra Atoll); South Africa (off
 Natal).

Depth range: 20-550m.

Habitat: coral reef slope. - among coral rubble in hollows
 between coral heads.

Pantopipetta bilobata ARNAUD & CHILD 1988

- 1988 *Pantopipetta bilobata* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 135-137,
 fig. 3A-G.

Distribution: South Africa.

Depth range: 90-550m.

Pantopipetta brevicauda STOCK 1963

- 1963 *Pantopipetta brevicauda* STOCK, Ann. S. Afr. Mus., **46** (12): 336-388, figs. 9,
 10a.
 1971 *Pantopipetta brevicauda*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**.
 Antarct. Res. Ser., 17: 223-226, figs. 1, 3-4.
 1978 *Pantopipetta brevicauda*, - STOCK, Cah. Biol. Mar., **19**: 205.
 1990 *Pantopipetta brevicauda*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR,
 126: 119.

Distribution: NE Atlantic Basin and southern Atlantic,
 including Antarctica; SW-Africa; Chile.

Depth range: 2580-5024m.

Pantopipetta brevipilata TURPAEVA 1990

1990 *Pantopipetta brevipilata* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **126**: 123-125, figs. 8-9.

Distribution: southern Atlantic.
Depth range: 6050-6150m.

Pantopipetta capensis (BARNARD 1946)

1946 *Pipetta capensis* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 60.

1954 *Pipetta capensis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 86-88, fig. 1.

1963 *Pantopipetta capensis*, - STOCK, Ann. S. Afr. Mus., **46** (12): 336, fig. 8a-e.

1971 *Pantopipetta capensis*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**. Antarct. Res. Ser., **17**: 226-227, figs. 1, 5.

Distribution: South Africa.
Depth range: 841-2890m.

Pantopipetta lata STOCK 1981

1981 *Pantopipetta lata* STOCK, Cah. Biol. Mar., **22**: 466-468, fig. 7.

Distribution: Walvis Basin.
Depth range: 942-973m.

Pantopipetta longituberculata (TURPAEVA 1955)

1955 *Pipetta longituberculata* TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **12**: 324-326, fig. 2.

1963 *Pantopipetta brevicauda* STOCK, Ann. S. Afr. Mus., **46** (12): 336.

1963 *Pantopipetta longituberculata*, - STOCK, Ann. S. Afr. Mus., **46** (12): 336.

1971 *Pantopipetta longituberculata*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**. Antarct. Res. Ser., **17**: 219, 221, 223.

1971 *Pantopipetta brevicauda*, - HEDGPETH & McCAIN, in: Biol. Antarct. Seas, **4**. Antarct. Res. Ser., **17**: 219, 220, 222, 223.

1971 *Pantopipetta longituberculata*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **92**: 290.

1974 *Pantopipetta longituberculata*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **98**: 292.

1975 *Pantopipetta brevicauda*, - STOCK, Bull. mar. Sci., **24** (4): 993.

1975 *Pantopipetta longituberculata*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **103**: 242.

1978 *Pantopipetta brevicaudata*, - STOCK, Cah. Biol. Mar., **19**: 205-206.

1982 *Pantopipetta longituberculata*, - CHILD, Smiths. Contr. Zool., **349**: 49.

1986 *Pantopipetta brevicauda*, - STOCK, Bull. mar. Sci., **38** (3): 405 tab. 1.

Distribution: Guiana Basin; Brazil Slope and Basin, off Recife; Argentine Basin, off Rio de la Plata; North American Basin; Scotia Sea; Walvis Ridge; Pacific trenches.

Depth range: 2456-6700m.

***Pantopipetta oculata* STOCK 1968**

1968 *Pantopipetta oculata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 22-24, fig. 7.

Distribution: Andaman Islands.

Depth range: 66m.

CALLIPALLENIDAE

- 1948 Pallenidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 199-202 [diagnosis, remarks, key to genera from American waters].
 1949 Pallenidae, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 275 [remarks].
 1954 Pallenidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 105 [diagnosis, key to South African genera].
 1974 Pallenidae, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 59.
 1978 Callipallenidae, - CHILD, Smiths. Contr. Zool., **293**: 41 [key to Middle American species].
 1982 Callipallenidae, - CHILD, Smiths. Contr. Zool., **349**: 24-25 [remarks, key to genera].
 1983 Callipallenidae, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 57 [key to Japanese genera].
 1986 Callipallenidae, - STOCK, Bull. mar. Sci., **38** (3): 421 [remarks].
 1992 Callipallenidae, - CHILD, Mem. Hourglass Cruises, **9** (1): 58, tab. 9.

***Anoropallene* STOCK 1956**

1956 *Anoropallene* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **54**: 46 [diagnosis].

***Anoropallene laysani* CHILD 1972**

- 1972 *Anoropallene laysani* CHILD, Proc. biol. Soc. Wash., **85** (11): 147-150, fig. 1.
 1983 *Anoropallene laysani*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 57.
 1987 *Anoropallene laysani*, - NAKAMURA, Biol. Lab., Imperial Household: 8-9, pl. 6, figs. 4-10.

Distribution: Japan.

Depth range: 30-80m.

***Anoropallene palpida* (HILTON 1939)**

- 1939 *Palene palpida* HILTON, J. Ent. Zool. Pomona College, **31** (2): 30.
 1942 *Oropallene palpida*, - HILTON, J. Ent. Zool. Pomona College, **34** (2): 38.
 1942 *Oropallene heterodonta* HILTON, J. Ent. Zool. Pomona College, **34** (2): 38.
 1954 *Oropallene* (?) spec., - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 29.

- 1956 *Anoropallene palpida*, - STOCK, Mitt. Zool. Mus. Hamburg, **1**: 46.
 1956 *Anoropallene heterodonta*, - STOCK, Mitt. Zool. Mus. Hamburg, **1**: 46.
 1956 *Anoropallene crenispina* STOCK, Mitt. Zool. Mus. Hamburg, **1**: 46-48, fig. 7.
 1979 *Anoropallene palpida*, - CHILD, Smiths. Contr. Zool., **293**: 40-41, fig. 3d-e.
 1992 *Anoropallene palpida*, - CHILD, Smiths. Contr. Zool., **526**: 24-25.

Distribution: California; Mexico and Panamá Pacific;
 Ecuador; Peru.

Depth range: 8-30m.

Habitat: often on muddy bottoms.

***Anoropallene valida* (HASWELL 1884)**

- 1884 *Nymphon validum* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1024-1025, pl. 54,
 figs. 6-9.
 1908 *Parapallene valida*, - LOMAN, Siboga Exped. Monogr., **40**: 48-49.
 1920 *Pallene (?) valida*, - FLYNN, Pap. & Proc. Roy. Soc. Tasm., 1919: 75-77, pl.
 XIX, figs. 7-8.
 1940 *Oropallene valida*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19**
 (Zool. 4): 33.
 1954 *Oropallene valida*, - STOCK, Vidensk. Medd. dansk naturhist. Foren, **116**: 29.
 1954 *Pallene valida*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 31.
 1956 *Anoropallene (?) valida*, - STOCK, Mitt. Zool. Mus. Hamburg, **54**: 46.
 1963 *Anoropallene valida*, - CLARK, Rec. Austr. Mus., **26** (1): 16-18, fig. 9A-J.

Distribution: Australia (Port Jackson).

***Austropallene* HODGSON 1907**

- 1927 *Austropallene*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 329 [diagnosis,
 remarks].
 1944 *Austropallene*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped.,
 (B), **5** (1): 36-37 [remarks, key to antarctic and sub-antarctic species].

***Austropallene brachyura* (BOUVIER 1913)**

- 1915 *Austropallene brachyura*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910,
 Zool., **3** (1): 39.
 1927 *Austropallene brachyura*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 332-
 334, fig. 8.
 1932 *Austropallene brachyura*, - GORDON, Discovery Rep., **6**: 86.
 1944 *Austropallene brachyura* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res.
 Exped., (B), **5** (1): 38-39, fig. 10b-d, 14b.
 1972 *Austropallene brachyura*, - ARNAUD, Tethys, Suppl. **4**: 146-147.

Distribution: Antarctica, probably circumpolar.

Depth range: 180-640m.

***Austropallene calmani* GORDON 1944**

- 1944 *Austropallene calmani* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res.

Exped., (B), 5 (1): 42-45, fig. 12a, figs. 13a-c, 14a.

Distribution: off Antarctic continent, in Gauss and Victoria Quadrants.

Depth range: 163-640m.

***Austropallene cornigera* (MÖBIUS 1902)**

- 1902 *Pseudopallene cornigera* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. Valdivia, 1898-1899, 3: 186, pl. XXVII, figs. 14-20.
 1915 *Austropallene cornigera*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 38.
 1923 *Austropallene cornigera*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., 1 (2): 22.
 1927 *Austropallene cornigera*, - HODGSON, Dt. Südpolar-Exped., 19 (Zool. 11): 329-330, fig. 7.
 1932 *Austropallene cornigera*, - GORDON, Discovery Rep., 6: 85-86, figs. 42-43.
 1938 *Austropallene cornigera*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), 2 (8): 15.
 1944 *Austropallene cornigera*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 36 [key], 37.
 1950 *Austropallene cornigera*, - HEDGPETH, Proc. U.S. Natn. Mus., 100 (3260): 151.
 1965 *Austropallene cornigera*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., 41 (31): 2.
 1972 *Austropallene cornigera*, - ARNAUD, Téthys, suppl. 4: 147.
 1980 *Austropallene cornigera*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., 52 (18): 4.

Distribution: Antarctica.

Depth range: 120-520m.

***Austropallene cristata* (BOUVIER 1911)**

- 1913 *Pseudopallene cristata*, - BOUVIER, 2ème Expéd. Antarct. Française 1908-1910: 102-105, figs. 55-58.
 1927 *Austropallene cristata*, - HODGSON, Dt. Südpolar-Exped., 19, Zool. 11: 331-332.
 1932 *Austropallene cristata*, - GORDON, Discovery Rep., 6: 86.
 1938 *Austropallene cristata*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), 2 (8): 15.
 1944 *Austropallene cristata*, - GORDON: Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 37.
 1980 *Austropallene cristata*, - KRAPP, Bull. Bull. Inst. Roy. Sci. Nat. Belg., 52 (18): 6.

Distribution: Antarctica, probably circumpolar.

Depth range: 350m.

***Austropallene gracilipes* GORDON 1944**

- 1944 *Austropallene gracilipes* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 39-41, fig. 10a, 11a, c, 14c, d.

Distribution: off the Antarctic Continent, in Gauss and Victoria Quadrants.

Depth range: 219-640m.

Austropallene tibicina CALMAN 1915

1915 *Austropallene tibicina*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 39-41, figs. 7-8.

1944 *Austropallene tibicina*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 41-42, fig. 12b-d, 14e.

1950 *Austropallene tibicina*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 151.

Distribution: Antarctica (Ross Sea to South Georgia).

Depth range: 180-209m.

Bradypallene KIM & HONG 1987

1987 *Bradypallene* KIM & HONG, Kor. J. Zool., **30** (3): 272 [diagnosis]

Bradypallene espina KIM & HONG 1987

1987 *Bradypallene espina* KIM & HONG, Kor. J. Zool., **30** (3): 2772-273, figs. 1-2.

Distribution: Korea.

Depth range: 5-7m.

Habitat: rocky bottom.

Callipallene FLYNN 1929

1948 *Callipallene*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 202 [diagnosis, remarks, key to NE-American species].

1954 *Callipallene*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 105-106 [diagnosis].

1954 *Callipallene*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 37 [diagnosis].

1979 *Callipallene*, - CHILD, Smiths. Contr. Zool., 293: 41 [key to Middle American species].

1983 *Callipallene*, - NNAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 57-58 [key to Japanese species].

1992 *Callipallene*, - CHILD, Mem. Hourglass Cruises, 9 (1): 60 [diagnosis], 59 [tab. 9].

Callipallene acribica KRAPP 1975

1975 *Callipallene acribica* KRAPP, Bonn. zool. Beitr., **26** (1-3): 284-288, fig. 1a-f.

1987 *Callipallene acribica*, - ARNAUD, Mesogee, **47**: 48.

Distribution: Mediterranean.

Depth range: 40-85m.

Habitat: "sur les fonds détriques côtiers" (ARNAUD).

***Callipallene acus* (MEINERT 1898)**

- 1898 *Pallene acus* MEINERT, Dansk. Ingolf-Exped., **3** (1): 45-46, pl. IV, figs. 8-13.
 1898 *Pallene hasata* MEINERT, Dansk. Ingolf Exped., **3** (1): 46-47, pl. IV, figs. 14-19.
 1899 *Pallene acus*, - MEINERT, Danish Ingolf-Exped., **3** (1): 48-49, pl. IV, figs. 8-13.
 1899 *Pallene hasata*, - MEINERT, Danish Ingolf-Exped., **3** (1): 49, pl. 4, figs. 14-19.
 1901 *Pallene acus*, - MÖBIUS, Fauna Arctica, **2** (1): 50.
 1901 *Pallene hasata*, - MÖBIUS, Fauna Arctica, **2** (1): 50.
 1926 *Pallene acus*, - BOUVIER, Res. Campagnes Sci. acc. Yacht Albert Ier Prince Souverain de Monaco, **51**: 26.
 1933 *Pallene acus*, - STEPHENSEN, Medd. om Grönland, **79** (6): 20.
 1948 *Callipallene acus*, - CORRÉA, Avuls. Dept. Zool., S. Paulo, **9** (1): 5 [key].
 1948 *Callipallene acus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 204-205, fig. 18c-e.
 1964 *Callipallene acus*, - STOCK, Beaufortia, **135** (11): 46, fig. 1

Distribution: N-Atlantic Basin; between Iceland and Greenland.

Depth range: 900-3800m.

Habitat: mud.

***Callipallene africana* ARNAUD & CHILD 1988**

- 1988 *Callipallene africana* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 138-140, fig. 4.

Distribution: South Africa (Transkei area).

Depth range: 150-775m.

***Callipallene amaxana* (OHSHIMA 1938)**

- 1933 ?*Pallene producta*, - OHSHIMA, Annot. Zool. Jap., **14** (2): 96.
 1938 *Pallene amaxana* OHSHIMA, Annot. Zool. Jap., **14** (2): 216.
 1936 *Pallene amaxana*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 863.
 1942 *Callipallene brevirostris*, - OHSHIMA, Proc. Imp. Acad., Tokyo, **18** (5): 257.
 1943 *Callipallene brevirostris*, - OHSHIMA, Bul. Sci. Fac. Terkult., Kyūsyū Imp. Univ., Tokyo, **10** (4): 371.
 1948 *Callipallene brevirostris*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 202-203, fig. 18a.
 1954 *Callipallene amaxana*, - STOCK, Vidensk. Medd. Dansk naturhist. Foren., **116**: 37.
 1962 *Callipallene phantoma*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 95.
 1968 *Callipallene phantoma amaxana*, - STOCK, Vidensk. Medd. Dansk naturhist. Foren., **131**: 37-38, fig. 14a-d.
 1971 *Callipallene phantoma amaxana*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 322.

- 1983 *Callipallene amaxana*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 58.
 1986 *Callipallene amaxana*, - KIM, Proc. Nat. Sci. Res. Inst. KANU, **2** (1): 1-3,
 fig. 1a-g.
 1986 *Callipallene amaxana*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 38.
 1987 *Callipallene amaxana*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 153.
 1991 *Callipallene amaxana*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 35-36.

Distribution: Japan; Korea; south to Straits of Malacca.
Depth range: 4-160m.

***Callipallene brevirostris* (JOHNSTON 1837)**

- 1837 *Pallene brevirostris* JOHNSTON, Mag. Zool. Bot., **1**: 380, pl. 12, figs. 7-8.
 1840 *Pallene brevirostris*, - MILNE-EDWARDS, Hist. Nat. des Crustacés..., 3. Ordre
 des Araneiformes ou Pycnogonides: 534-535.
 1845 ?*Phoxichilus spinosus* QUATREFAGES, Ann. Sci. Nat., Paris, **3** (4): 41, pl. I,
 fig. 2a, pl. 2, fig. 1.
 1859 *Pallene brevirostris*, - FORBES, in: Nat. Hist. of European Seas: page?
 1869 *Pallene brevirostris*, - GRUBE, Abh. Schles. Ges. Vaterländische Cultur,
 Breslau, 1868/1869: 118-119, 125, pl. I, fig. 5a-c.
 1873 *Pallene sp.*, - VERRILL, Rep. U.S. Comm. Fish., **1**: 415.
 1877 *Pallene brevirostris*, - HOEK, Niederl. Arch. Zool., **3**: 237-240, pl. XV,
 figs. 4-7, pl. XVI, figs. 14, 21-22.
 1878 *Pallene empusa* WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 9, pl. 3, fig. 2a-
 g.
 1880 *Pallene empusa*, - WILSON, Rep. U.S. Fish. Comm., **6**: 476-477, pl. 2, figs. 5-
 7.
 1881 *Pallene brevirostris*, - HOEK, Arch. Zool. Exp. Gén., **9**: 511-512, pl. XXVI,
 fig. 17, pl. XXIX, fig. 36.
 1881 *Pallene empusa*, - RATHBUN, Proc. U.S. Natn. Mus., **3**: 118.
 1884 *Pallene brevirostris*, - HANSEN, in: Zoologia Danica Afbildn. danske Dyr med
 populaer Text, **4**: pl. VII, fig. 20.
 1884 *Pallene brevirostris*, - HANSEN, Naturhist. Tidsskr., Kobenhavn, **14** (3): 649.
 1885 *Pallene brevirostris*, - GOSSE, in: Manual of Mar. Zool. Brit. Isles: 120,
 fig. 192.
 1886 *Pallene brevirostris*, - HALHED, Ann. Rep. Fauna Liverpool Bay and the
 Neighbouring Seas, **1** (1): 230.
 1888 *Pallene brevirostris*, - SARS, Arch. Math. Naturvidensk., Oslo, **12**: ?page.
 1890 *Pallene brevirostris*, - TOPSENT, Bull. Soc. Linéenne Normandie, Caen, **2** (4):
 62.
 1891 *Pallene empusa*, - MORGAN, Stud. Biol. Lab. John Hopkins Univ., Baltimore, **5**
 (1): 8-22.
 1891 *Pallene brevirostris*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool.
 20): 32.
 1893 *Pallene brevirostris*, - MÖBIUS, Sitz.-Ber. Preuss. Akad. Wiss. Berlin, 1893:
 86.
 1896 *Pallene brevirostris*, - HERDMAN, Rep. 66th Meeting Brit. Ass. Adv. Sci.:
 442.
 1901 *Pallene brevirostris*, - COLE, Biol. Bull. Woods Hole Mar. Biol. Lab., **2** (5):
 195-207.
 1901 *Pallene brevirostris*, - MÖBIUS, Fauna Arctica, **2** (1): 50.
 1901 *Pallene brevirostris*, - d'ARCY THOMPSON, in: The Cambr. Nat. Hist.: 54.
 1904 *Pallene brevirostris*, - HALLEZ, Arch. Zool. Exp. Gén., **3** (4) (Notes & Rev.,
 3): p. L.
 1908 *Pallene brevirostris*, - NORMAN, J. Linn. Soc. Zool., London (Zool.), **30**:

- 204-205 [part].
- 1908 *Pallene brevirostris*, - SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **13** (1908): 430.
- 1909 *Pallene brevirostris*, - d'ARCY THOMPSON, in: The Cambr. Nat. Hist.: 541, 542, fig. 275A, 285 8part.].
- 1910 *Pallene brevirostris*, - TESCH, Jaarbork Rijksinst. Onderzoek der Zee, 1909 (4): 51, 55.
- 1912 *Pallene brevirostris*, - LOMAN, Bull. Inst. Océanogr. Monaco, **238**: 8 [part].
- 1913 *Pallene brevirostris*, - SUMMER, OSBURNE & COLE, Bull. Bureau Fish., Washington, **31** (1): 677.
- 1918 *Pallene brevirostris*, - WIRÉN, Zoologisk Bidrag Uppsala, **6**: 42.
- 1923 *Pallene brevirostris*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10) **6** (3,4): 119 [part].
- 1923 *Pallene brevirostris*, - BOUVIER, Faune de France, **7**: 34-36, fig. 28.
- 1925 *Pallene brevirostris*, - FISH, Bull. Bureau Fish., Washington, **41**: 161.
- 1925 *Pallene brevirostris*, - MEISENHEIMER, Die Tierwelt der Nord- und Ostsee, **11**: 2, 5.
- 1927 *Pallene brevirostris*, - MEISENHEIMER, Die Tierwelt der Nord- und Ostsee (Nachtrag), **11a**: 14.
- 1928 *Pallene brevirostris*, - GILTAY, Bull. & Ann. Soc. Roy. Ent. Belg., **68** (9-10): 211-213, fig. 8.
- 1928 *Pallene brevirostris*, - LOMAN, Die Tierwelt Deutschlands, **8** (3): 79-80, fig. 7.
- 1930 *Pallene brevirostris*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 245-253, figs. 58-59 [part].
- 1932 *Pallene brevirostris*, - SCHLOTTKE, Wiss. Meeresunters. Kommiss. Wiss. Unters. der Deutschen Meere, Abt. Helgoland, (N.F.), **18** (9): 1-10.
- 1933 *Pallene brevirostris*, - STEPHENSEN, Danmarks Fauna, **38**: 33-35, fig. 10.
- 1935 *Pallene brevirostris*, - HELFER, Der Naturforscher: 268, fig. 130B, 178.
- 1936 *Pallene brevirostris*, - HELFER, Ministry of Commerce & Industry, Egypt, Notes & Mem., **16**: 2 [map].
- 1936 *Pallene brevirostris*, - STEPHENSEN, Bergens Mus. Arbok, 1935, Naturvidensk. Rekke, **7**: 24.
- 1936 *Pallene brevirostris*, - STEPHENSEN, Medd. Göteborgs Mus. Zool. Avd., **69**: 34-36, fig. 7.
- 1940 *Pallene brevirostris*, - FARAGGIANA, Boll. Mus. Zool. Anat. Comp. Real. Univ. Torino, (3) **48** (112): 2-4 [part].
- 1945 *Pallene brevirostris*, - LEBOUR, J. bar. biol. Ass. U.K., **26**: 144.
- 1948 *Callipallene brevirostris*, - CORREA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 6.
- 1948 *Callipallene brevirostris*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 202-203, fig. 18a.
- 1949 *Callipallene brevirostris*, - STOCK, Strand-Werk-Gemeenschap Tabellenser., **7**: 3, fig. 4.
- 1954 *Callipallene brevirostris brevirostris*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 115.
- 1954 *Callipallene brevirostris*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 43-44, fig. 18.
- 1955 *Callipallene brevirostris*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 589.
- 1959 *Callipallene brevirostris*, - BACESCU, Rev. Biol. Acad. Rep. Pop. Roumanie, **4** (1): 127.
- 1970 *Callipallene brevirostris brevirostris*, - STOCK, "Meteor" Forschungs-Ergebnisse, Reihe D, No. **5**: 9.
- 1972 *Callipallene brevirostris*, - KING, J. nat. Hist., **6**: 621-622.
- 1974 *Callipallene brevirostris*, - KING, A Synopsis of the British Sea Spiders.

- Synopses of the British Fauna (New Ser.), **5**: 59 [key], 60-61, fig. 27.
- 1975 *Callipallene brevirostrum*, - STOCK, Bull. mar. Sci., **24** (4): 1010.
- 1976 *Callipallene brevirostris*, - ARNAUD, Acta Ecol. Iranica, **1**: 70.
- 1976 *Callipallene brevirostris*, - WOLFF, Netherlands J. Sea Res., **10** (4): 476-477, Fig. 4b (distribution in estuary).
- 1978 *Callipallene brevirostris*, - CHIMENZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 94-95, fig. 2.
- 1979 *Callipallene brevirostris*, - STOCK, Stud. Fauna Curaçao and other Caribb. Ils., **59** (184): 14.
- 1982 *Callipallene brevirostris*, - CHILD, Smiths. Contr. Zool., **349**: 26.
- 1986 *Callipallene brevirostris*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1, 424.
- 1987 *Callipallene brevirostris*, - ARNAUD, Mesogee, **47**: 48-49
- 1987 *Callipallene brevirostris*, - STOCK, Cah. Biol. Mar., **28**: 512.
- 1992 *Callipallene brevirostris*, - CHILD, Mem. Hourglass Cruises, **9** (1): 59 [key], 60-61, fig. 27.

Distribution: N-Atlantic inclusive of Caribbean, Mediterranean; Black Sea; West Africa.

Depth range: 0-316m.

Habitat: in a variety of substrata.

***Callipallene bullata* NAKAMURA & CHILD 1991**

- 1991 *Callipallene bullata* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 36, fig. 13.

Distribution: Japan.

Depth range: 37-41m.

***Callipallene californiensis* (HALL 1913)**

- 1913 *Pallene californiensis* HALL, Univ. Calif. Publ. Zool., **11** (6): 131, 133-135, pl. 4, figs. 9-13.
- 1915 *Pallene californiensis*, - HILTON, J. Ent. Zool. Pomona College, **7** (1): 67.
- 1915 *Palene californiensis*, - HILTON, J. Ent. Zool. Pomona College, **7** (3): 204.
- 1916 *Pallene californiensis*, - HILTON, J. Comp. Neur., **26** (5): 465.
- 1920 *Pallene californiensis*, - HILTON, J. Ent. Zool. Pomona College, **12** (4): 93.
- 1939 *Palene californiensis*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 29.
- 1941 *Callipallene californiensis*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 257 [key], pl. 11.
- 1942 *Callipallene californiensis*, - HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 277, 281.
- 1942 *Callipallene californiensis*, - HILTON, J. Ent. Zool. Pomona College, **34** (2): 38.
- 1948 *Callipallene californiensis*, - CORRÉA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 6 [key].
- 1979 *Callipallene solicitatus* CHILD, Smiths. Contr. Zool., **293**: 44-46, fig. 15.
- 1987 *Callipallene californiensis*, - CHILD, Proc. biol. Soc. Wash., **100** (3): 555.
- 1992 *Callipallene californiensis*, - CHILD, Smiths. Contr. Zool., **526**: 25.

Distribution: California; Pacific Coast of Mexico and Panamá; Galapagos Archipelago; Chile (Islas Juan Fernandez).

Depth range: shallow subtidal to 10m.
Habitat: under stones in channel.

Callipallene conirostris STOCK 1954

- 1954 *Callipallene conirostris* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 39-41, figs. 16, 20f-g.
 1971 *Callipallene conirostris*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 323.

Distribution: Japan.
Depth range: 146-220m.

Callipallene cuspidata STOCK 1954

- 1954 *Callipallene cuspidata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 37-39, fig. 15.
 1991 *Callipallene cuspidata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 36-38.

Distribution: Philippines; Japan.
Depth range: 300-732m.

Callipallene dubiosa HEDGPETH 1949

- 1949 *Callipallene dubiosa* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 275-277, fig. 35.
 1954 *Callipallene dubiosa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 41.
 1957 *Callipallene dubiosa*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 88.
 1965 *Callipallene dubiosa*, - UTINOMI, New Illustr. Fauna Japan, **2**: 335.
 1971 *Callipallene dubiosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 322.
 1983 *Callipallene dubiosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 58.
 1984 *Callipallene dubiosa*, - KIM, J. Kangreung Natn. Univ., **7**: 535.
 1986 *Callipallene dubiosa*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 38.
 1987 *Callipallene dubiosa*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 153.
 1990 *Callipallene dubiosa*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 71.
 1991 *Callipallene dubiosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 38.
 1992 *Callipallene dubiosa*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159.

Distribution: recorded from East Africa (Kenya, Zanzibar) to Japan and Korea.

Depth range: 1-46m.
Habitat: dead corals; rocky bottom.

Callipallene emarginata (DOHRN 1881)

- 1881 *Pallene emarginata* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 193-196, pl. 14, figs. 10-21.
 1885 *Pallene emarginata*, - CARUS, Prodromus faunae mediterraneae sive descriptio animalium maris Mediterranei incolarum, **2**: 287.

- 1908 *Pallene brevirostris*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 204-205 [part].
- 1909 *Pallene brevirostris*, - d'ARCY THOMPSON, The Cambridge Nat. Hist.: 541 [part.].
- 1912 *Pallene brevirostris*, - LOMAN, Bull. Inst. Océanogr., **238**: 8 [part.].
- 1923 *Pallene emacinata*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10), **6**, (3,4): 119 [part.].
- 1923 *Pallene emacinata*, - BOUVIER, Faune de France, **7**: 34, 36.
- 1929 *Pallene emacinata*, - GILTAY, Bull. & Ann. Soc. Roy. Ent. Belg., **69**: 172-173, fig. 1 [part.].
- 1930 *Pallene brevirostris*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 245-253 [part.].
- 1940 *Pallene brevirostris*, - FARAGGIANA, Boll. Mus. Zool. Anat. Comp. Real. Univ. Torino, (3) **48** (112): 2-4 [part.].
- 1948 *Callipallene emacinata*, - CORRÉA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 5, 7, 8.
- 1948 *Callipallene emacinata*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 204, fig. 18f-g.
- 1952 *Callipallene emacinata*, - STOCK, Beaufortia, **1** (13): 8-11, figs. 1-4, 7-8, 15-19.
- 1954 *Callipallene emacinata* ssp. *micracantha* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 44-46, figs. 19a-g, 20a-b.
- 1954 *Callipallene emacinata* subsp., - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 46-48, figs. 19h-i, 20c-e.
- 1958 *Callipallene emacinata emacinata*, - STOCK, Bull. Res. Counc. Israel., **7B**: 137.
- 1963 *Callipallene emacinata micracanta*, - CLARK, Rec. Austr. Mus., **26** (1): 21-23, fig. 11A-I.
- 1974 *Callipallene emacinata*, - ARNAUD, Bull. Zool. Mus. Univ. Amsterdam, **3** (21): 175.
- 1974 *Callipallene emacinata*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 59 [key], 64.
- 1975 *Callipallene emacinata micracantha*, - CHILD, Smiths. Contr. Zool., **190**: 10.
- 1975 *Callipallene emacinata*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 16-17, 18-20 (tab. II), 23 (ecology).
- 1975 *Callipallene emacinata*, - STOCK, Bull. mar. Sci., **24** (4): 1011.
- 1976 *Callipallene emacinata*, - ARNAUD, Acta Ecol. Iranica, **1**: 70.
- 1979 *Callipallene emacinata*, - CHILD, Smiths. Contr. Zool., **293**: 41-42.
- 1981 *Callipallene emacinata emacinata*, - MUNILLA & de HARO, Bijdr. Dierk., **51** (2): 191-197, figs. 8, 11, tab. I (electrophoretical and immunological study).
- 1982 *Callipallene emacinata*, - CHILD, Smiths. Contr. mar. Sci., **12**: 365-366.
- 1984 *Callipallene emacinata emacinata*, - MUNILLA & de HARO, in: Els sistemes naturals de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 534-535, tab. III-IV.
- 1986 *Callipallene emacinata*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1.
- 1987 *Callipallene emacinata*, - ARNAUD, Mesogee, **47**: 49.
- 1987 *Callipallene emacinata*, - STOCK, Cah. Biol. Mar., **28**: 513.
- 1990 *Callipallene emacinata*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 280.

Distribution: N-Atlantic. - Mediterranean Sea; Caribbean Sea.

Depth range: shallow water.

Habitat: in a variety of substrata, often on algaee.

Callipallene evelinae MARCUS 1940

- 1940 *Callipallene evelinae* MARCUS, Bolm. Fac. Fil., Cienc. Letr., Univ. S. Paulo, **4**: 31-33, figs. 2 A-D.
 1992 *Callipallene evelinae*, - STOCK, Tijdschr. Ent., **135**: 128, 139.

Distribution: Brazil.

Depth range: 0-130m.

Habitat: rocky shore, on *Dyctiota*.

Callipallene gabriellae CORRÉA 1948

- 1948 *Callipallene gabriellae* CORRÉA, Pap. Avuls., Dept. Zool., S. Paulo, **9** (1): 1-4, figs. 1-6.
 1992 *Callipallene gabriellae*, - STOCK, Tijdschr. Ent., **135**: 128, 139.

Distribution: Brazil.

Depth range: infralittoral.

Habitat: seaweeds growing on rocks.

Callipallene hastata MEINERT 1899

- 1899 *Pallene hastata* MEINERT, Danish Ingolf-Exped., **3** (1): 49, pl. IV, figs. 14-19.

Distribution: southern part of Davis Strait.

Depth range: 2626m.

Callipallene kenyensis MÜLLER 1990

- 1990 *Callipallene kenyensis* MÜLLER, Bonn. zool. Beitr., **41** (1): 71-74, figs. 21-26.
 1992 *Callipallene gabriellae*, - STOCK, Bijdr. Dierk., **62** (2): 94.

Distribution: Kenya (Watamu area); Oman.

Depth range: 0-3m.

Habitat: dead corals covered with algae; also "under rocks and stones"; "on algae"; "tidal pools, on stones"; "from stones in tidal pools"; "muddy tidal flat with stones and algae" (STOCK).

Remarks: Species was erroneously considered conspecific with *C. gabriellae* from Brazil by STOCK (1992).

Callipallene longicoxa STOCK 1955

- 1955 *Callipallene brevirostris longicoxa* STOCK, Vidensk. Medd. Dansk naturhist. Foren., **117**: 223-226, figs. 4-5.
 1982 *Callipallene belizae* CHILD, - Smiths. Contr. mar. Sci., **12**: 363-365, fig. 164.
 1986 *Callipallene longicoxa*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1, 424.

Distribution: Lesser Antilles, Belize and Florida.

Depth range: 18-549m.

Habitat: *Halimeda* and rubble, algae.

***Callipallene margarita* (GORDON 1932)**

- 1932 *Pallene margarita* GORDON, Discovery Rep., **6**: 82-85, figs. 40-41.
 1948 *Callipallene margarita*, - CORRÉA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 6.
 1961 ?*Callipallene margarita*, - HEDGPETH, Lunds Univ. Arsskr., N.F., **2** (57): 3, 4, fig. 1.
 1982 *Callipallene margarita*, - CHILD, Smiths. Contr. Zool., **349**: 26.
 1992 *Callipallene margarita*, - STOCK, Tijdschr. Ent., **135**: 128, 139.

Distribution: along Argentine Shelf and Slope, Burdwood Bank, and South Georgia area; Golfo San Jorge, Argentina; Brazil.

Depth range: ?<120-2323m.

***Callipallene novaezealandiae* (THOMSON 1884)**

- 1884 *Pallene novae-zealandiae* THOMSON, Trans. & Proc. N.Z. Inst., **16** (1883): 246-247, pl. 14, figs. 1-4.
 1904 *Pallene novae-zealandiae*, - HUTTON, Index Faunae N.Z.: 247.
 1954 *Callipallene brevirostris* ssp. *novae-zealandiae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **386**: 48-50, fig. 21a-h.
 1972 *Callipallene brevirostris* cf. ssp. *novaezealandiae*, - ARNAUD, Tethys, Suppl. **3**: 162-163.
 1975 *Callipallene brevirostris novaezealandiae*, - CHILD, Smiths. Contr. Zool., **190**: 10.
 1976 *Callipallene brevirostrum* ssp. *novaezealandiae*, - STOCK, Trav. Mus. Hist. Nat. "Gr. Antipa", **16**: 132.
 1982 *Callipallene* sp. cf. *C. novaezealandiae*, - CHILD, Proc. biol. Soc. Wash., **95** (2): 277.
 1983 *Callipallene novaezealandiae*, - CHILD, Proc. biol. Soc. Wash., **96** (4): 708.
 1983 *Callipallene brevirostris novaezealandiae*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 58-59.
 1988 *Callipallene novaezealandiae*, - CHILD, Smiths. Contr. Zool., **468**: 21.
 1988 *Callipallene novaezealandiae*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 664.
 1990 *Callipallene novaezealandiae*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk., Dresden, **45** (10): 106.
 1991 *Callipallene novaezealandiae*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 145.
 1991 *Callipallene novaezealandiae*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 38.

Distribution: from E-Africa (Tanzania) to the Philippines; Palau Islands; Ryukyus; Japan (Sagami Bay).

Depth range: 0.3-300m.

Habitat: various coral reef locations, often among seaweeds (*Thalassodendron ciliatum*, *Syringodium* et *Cymodocea serrulata*).

Callipallene pacifica (HEDGPETH 1939)

1939 *Pallene pacifica* HEDGPETH, Am. Midland Nat., **22** (2): 460-461, pl. 1, figs. f-i, pl. 2, figs. j-l.

Distribution: off S-California.

Depth range: 103m.

Habitat: "rock (?) " (HEDGPETH).

Callipallene panamensis CHILD 1979

1979 *Callipallene panamensis* CHILD, Smiths. Contr. Zool., **293**: 42-44, fig. 14.

1983 *Callipallene panamensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 59.

1991 *Callipallene panamensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 38.

Distribution: Panamá Pacific and Japan.

Depth range: littoral-39m.

Callipallene pectinata (CALMAN 1923)

1923 *Pallene pectinata* CALMAN, Rec. Indian Mus., **24** (3): 275-276, fig. 5.

1938 *Pallene pectinata*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 154.

1948 *Callipallene pectinata*, - CORRÉA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 5 [key].

1964 *Callipallene pectinata*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **35**: 28-31, figs. 2-4.

1972 *Callipallene pectinata*, - ARNAUD, Tethys, Suppl. **3**: 161.

1968 *Callipallene pectinata*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 38-39.

Distribution: Red Sea; Arabian Coast; Andaman Islands; Madagascar.

Depth range: 0.8-66m.

Habitat: from brown algae on dead *Acropora* corals.

Callipallene phantoma (DOHRN 1881)

1881 *Pallene phantoma* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 196, pl. 14, figs. 1-9.

1885 *Pallene phantoma*, - CARUS, Prodromus faunae mediterraneae sive descriptio animalium maris Mediterranei incolarum, **2**: 287.

1908 *Pallene phantoma*, - NORMAN, J. Linn. Soc. London, (Zool.), **30**: 205.

1912 *Pallene phantoma*, - LOMAN, Bull. Inst. Océanogr. Monaco, **238**: 9.

1923 *Pallene phantoma*, - BOUVIER, Faune de France, **7**: 34, 37.

1934 *Pallene phantoma*, - GILTAY, Bull. Mus. Roy. d'Hist. Nat. Belg., **10** (35): 1-5.

1942 *Callipallene brevirostris*, - OHSHIMA, Proc. Imp. Acad., Tokyo, **18** (5): 257.

1948 *Callipallene phantoma*, - CORRÉA, Pap. Avuls. Dept. Zool., S. Paulo, **9** (1): 6, 8.

1948 *Callipallene phantoma*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 204, fig. 18b.

1952 *Callipallene phantoma*, - STOCK, Beaufortia, **1** (13): 3-4, figs. 12-14, 20-27

- [ssp. *phantoma*, *crinita*].
- 1953 *Callipallene phantoma*, - BACESCU, Bul. Stint., Acad. Rep. Pop. Rom., Sect. de Stint. Biol., Agr. Geol. si Geogr., **5** (2): 264.
- 1954 *Callipallene phantoma phantoma*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **24**: 116.
- 1962 *Callipallene phantoma*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 95.
- 1968 *Callipallene phantoma phantoma*, - STOCK, Vie Milieu, **19** (**1A**): 23, fig. 16.
- 1973 *Callipallene phantoma phantoma*, - KRAPP, Beaufortia, **21** (277): 73.
- 1974 *Callipallene phantoma*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 59 [key], 62-63, fig. 28.
- 1975 *Callipallene phantoma*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 17, 18-20 (tab. II), 22 (ecology).
- 1976 *Callipallene phantoma*, - ARNAUD, Acta Ecol. Iranica, **1**: 70.
- 1982 *Callipallene phantoma*, - CHILD, Smiths. Contr. Zool., **349**: 26.
- 1984 *Callipallene phantoma phantoma*, - MUNILLA & de HARO, in: Els sistemes de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 532, tab. I, 535, tab. IV.
- 1986 *Callipallene phantoma*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1.
- 1987 *Callipallene phantoma*, - ARNAUD, Mesogee, **47**: 49.
- 1987 *Callipallene phantoma*, - STOCK, Cah. Biol. Mar., **28**: 513.
- 1988 *Callipallene phantoma*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 137-138.
- 1989 *Callipallene phantoma phantoma*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 287 [tab. 1], 289 [tab. 3], 292, fig. 4.

Distribution: N-Atlantic; Black Sea; Mediterranean to New England; Florida; Bahamas; Japan; North American Slope, S of Cape Cod; South Africa.

Depth range: 0-850m.

Habitat: in a variety of substrata, often on algae.

***Callipallene producta* (SARS 1888)**

- 1952 *Callipallene brevirostris producta*, - STOCK, Beaufortia, **13**: 6-7, figs. 9-11.
- 1978 *Callipallene producta*, - STOCK, Cah. Biol. Mar., **19**: 215.
- 1984 *Callipallene brevirostris producta*, - MUNILLA & de HARO, in: Els sistemes de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 533, 535, tab. II, IV.
- 1987 *Callipallene producta*, - STOCK, Cah. Biol. Mar., **28**: 513.
- 1990 *Callipallene producta*, - STOCK, Zool. Meded., **63** (16): 227.
- 1992 *Callipallene producta*, - STOCK, Tijdschr. Ent., **135**: 128, 139.

Distribution: NE-Atlantic, from Norway to Cape Verde Islands; Brazil.

Depth range: 70-1550m.

***Callipallene sagamiensis* NAKAMURA & CHILD 1983**

- 1983 *Callipallene sagamiensis* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 59-61, fig. 20.
- 1986 *Callipallene sagamiensis*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 38, fig. 3a-h.
- 1987 *Callipallene sagamiensis*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 153.

- 1987 *Callipallene sagamiensis*, - NAKAMURA, Biol. Lab., Imperial Household: 9-10, pl. 7.
 1991 *Callipallene sagamiensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 38-39.

Distribution: Japan and Korea.

Depth range: 100-160m.

***Callipallene seychellensis* CHILD 1988**

- 1988 *Callipallene seychellensis* CHILD, Bull. biol. Soc. Wash., **8**: 61-65, fig. 4.

Distribution: Seychelles (Aldabra Atoll, Mahé Island).

Depth range: 0-6m.

Habitat: "dense Halimeda on dead coral"; "offshore, corallines"; "dead coral rubble" (CHILD).

***Callipallene spectrum* (DOHRN 1881)**

- 1975 *Callipallene spectrum*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 16-17, 18-20 (tab. II) (ecology).
 1984 *Callipallene emacinata spectrum*, - MUNILLA & de HARO, in: Els sistemes de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 532-533, tab. II, 535, tab. IV.
 1987 *Callipallene spectrum*, - ARNAUD, Mesogee, **47**: 49-50
 1989 *Callipallene emacinata spectrum*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 287 [tab. 1], 289 [tab. 4], 291-294, fig. 3 (phenology).

Distribution: NE-Atlantic; Mediterranean Sea; ? Black Sea.

Depth range: 0-95m.

Habitat: often found on algae.

***Callipallene tiberi* (DOHRN 1881)**

- 1952 *Callipallene emacinata tiberi*, - STOCK, Beaufortia, **1** (13): 8-11, figs. 3, 4, 17, 18.
 1972 *Callipallene emacinata tiberi*, - KING, J. nat. Hist., **6**: 622.
 1975 *Callipallene tiberi*, - KRAPP-SCHICKEL & KRAPP, Vie Milieu, (B) **25** (1): 17, 18-20 (tab. II), 21-22 (ecology).
 1987 *Callipallene tiberi*, - ARNAUD, Mesogee, **47**: 50.
 1987 *Callipallene tiberi*, - STOCK, Cah. Biol. Mar., **28**: 513-514.
 1989 *Callipallene emacinata tiberi*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 287 [tab. 1], 289 [tab. 5] (phenology).
 1990 *Callipallene tiberi*, - STOCK, Zool. Meded., **63** (16): 227-228.

Distribution: Scilly Islands; Mediterranean; Canary Islands; Portugal; Atlantic coasts of Europe.

Depth range: 0-523m.

Habitat: In a variety of substrata, often in algal vegetation; amongst *Alcyonidium* and tunics on horizontal rock surface over sandy bottom; shell gravel; red seaweeds; amongst *Leparalia* and *Nemertesia* on boulders; above sandy

substrate and shelly sand; amongst sponges and tunicates.

Callipallene tridens NAKAMURA & CHILD 1988

1988 *Callipallene tridens* NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 664-667, fig. 1.

Distribution: Japan (Ryukyu Islands).
Depth range: 3m.

Callipallene vexator STOCK 1956

1956 *Callipallene vexator* STOCK, Vidensk. Medd. dansk. naturhist. Foren., **118**: 79-81, fig. 4.

Distribution: South Africa.
Depth range: 64m.
Habitat: sandy mud.

Cheilopallene STOCK 1955

1955 *Cheilopallene* STOCK, Vidensk. Medd. Dansk naturh. Foren., **117**: 230 [diagnosis, remarks].

1992 *Cheilopallene*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [diagnosis, remarks, key to species].

Cheilopallene brevichelata CLARK 1961

1961 *Cheilopallene brevichelata* CLARK, Ann. Mag. Nat. Hist., **13** (3): 293-296, figs. 8-14.

1992 *Cheilopallene brevichelata*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key], 163 [text].

Distribution: Maldives.
Depth range: 30m.
Habitat: "from Vermetes colony in outer reef" (CLARK).

Cheilopallene clavigera STOCK 1955

1955 *Cheilopallene clavigera* STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 231-233, figs. 8-9.

1986 *Cheilopallene clavigera*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1.

1992 *Cheilopallene clavigera*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key].

Distribution: Caribbean (Virgin Islands).
Depth range: 26-28m.

Cheilopallene coralliophila MÜLLER 1992

1992 *Cheilopallene coralliophila* MÜLLER, Bonn. zool. Beitr., **43** (1): 159-163, figs. 12-17.

Distribution: Malaysia (Pulau Tioman).

Depth range: 1-3m.

Habitat: dead corals.

***Cheilopallene gigantea* CHILD 1987**

1987 *Cheilopallene gigantea* CHILD, Proc. biol. Soc. Wash., **100** (4): 912-914, fig. 5.

1992 *Cheilopallene gigantea*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key].

Distribution: Antarctic (Palmer Peninsula, Weddell Sea; off Larsen Ice Shelf).

Depth range: 581-610m.

***Cheilopallene hirta* CHILD 1988**

1988 *Cheilopallene hirta* CHILD, Bull. biol. Soc. Wash., **8**: 65-66, fig. 5.

1988 *Cheilopallene hirta*, - CHILD, Proc. biol. Soc. Wash., **101** (3): 667.

1992 *Cheilopallene hirta*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key].

Distribution: Japan (Ryukyu Islands), Seychelles (Aldabra Atoll).

Depth range: 0-21m.

Habitat: "channel edge, root mass and dead algae"; "rubble in hollows between coral heads"; "Reef crest"; "Seaward of reef crest, seagrass bed"; "coralline algae on beach rock"; "Lagoon..., algae on limestone pavement" (CHILD, 1988).

***Cheilopallene nodulosa* HONG & KIM 1987**

1987 *Cheilopallene nodulosa* HONG & KIM, Kor. J. Syst. Zool., **3** (2): 153-155, fig. 12.

1990 *Cheilopallene nodulosa*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 321.

1991 *Cheilopallene nodulosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 38.

1992 *Cheilopallene nodulosa*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key].

Distribution: Japan, Korea, Australian barrier reef (Lizard Island).

Depth range: 5-42m.

Habitat: rubble on the bottom of a cliff.

***Cheilopallene trappa* CLARK 1971**

1972 *Cheilopallene trappa* CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 331-332, fig. 1A-J.

1977 *Cheilopallene trappa*, - CLARK, N.Z. J. Mar. Fresw. Res., **11** (1): 174.

1992 *Cheilopallene trappa*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 159 [key].

Distribution: New Zealand (Snares Islands).

Depth range: 16-81m.

Habitat: coarse shally sand; on *Caulerpa brownii*.

***Decachela* HILTON 1939**

1939 *Decachela* HILTON, J. Ent. Zool. Pomona College, **31** (2): 34 [diagnosis].

***Decachela discata* HILTON 1939**

1939 *Decachela discata* HILTON, J. Ent. Zool. Pomona College, **31** (2): 34.

1942 *Decachela discata*, - HILTON, J. Ent. Zool. Pomona College, **34** (3): 70.

1949 *Decachela discata*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 280.

1971 *Decachela discata*, - UTINOMI, Publ. Seto Mart. Biol. Lab., **18** (5): 325.

1991 *Decachela discata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 39.

Distribution: California; Japan; Korea Strait.

Depth range: shallow water-180m.

***Decachela dogieli* LOSINA-LOSINSKY 1961**

1961 *Decachela dogieli* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 88.

1987 *Decachela dogieli*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 155.

Distribution: Korea

Habitat: on starfish *Pteraster*; "from the ambulacrinal grooves of starfishes" (HONG & KIM).

***Hannonia* HOEK 1881**

1954 *Hannonia*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 123-124 [diagnosis, remarks].

1993 *Hannonia*, - MUNILLA, J. mar. biol. Ass. U.K., **73**: 549 [key to species].

***Hannonia echinata* STOCK 1990**

1990 *Hannonia echinata* STOCK, Cah. Biol. Mar., **31**: 513-516, figs. 1-7.

Distribution: NE-Atlantic.

Depth range: 51m.

Habitat: "fond dur, riche en éponges, hydriaires et tuniciers" (STOCK).

***Hannonia spinipes* STOCK 1956**

1956 *Hannonia spinipes* STOCK, Vidensk. Medd. dansk. naturhist. Foren., **118**: 89-92, fig. 12.

Distribution: South Africa.

Depth range: 64m.

Habitat: bottom with sponges and corals.

***Hannonia stocki* MUNILLA 1993**

1993 *Hannonia stocki* MUNILLA, J. mar. biol. Ass. U.K., **73**: 544-548, fig. 2.

Distribution: Gibraltar area.

Depth range: 76-80m.

Habitat: gravel and rocks.

***Hannonia typica* HOEK 1881**

- 1881 *Hannonia typica* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 92, pl. 14, figs. 8-11.
 1904 *Hannonia typica*, - LOMAN, Zool. Jb., **20** (4): 383, pl. 14, figs. 12-15.
 1910 *Hannonia typica*, - HODGSON, Denkschr. Med.-Nat. Ges. Jena, **16**: 227.
 1923 *Hannonia typica*, - LOMAN, Medd. Göteborgs Mus., Zool. Avd., **22** (6): 7.
 1927 *Hannonia typica*, - CALMAN, Trans. Zool. Soc. London, **22** (3): 410.
 1954 *Hannonia typica*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 124-125, fig. 18.
 1959 *Hannonia typica*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 561.
 1962 *Hannonia typica*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 285.
 1970 *Hannonia typica*, - PENRITH & KENSLEY, Cimbebasia, (A), **1** (9): 232.

Distribution: Somalia; South Africa.

Depth range: 0-40m.

Habitat: on *Pocillopora* corals; sand banks at low tide; *Zostera* flats; sand and shells; Khaki mud; on rocky shores.

***Metapallene* SCHIMKEWITSCH 1909**

- 1954 *Metapallene*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 113-114 [diagnosis, remarks].

***Metapallene dubitans* (HODGSON 1910)**

- 1910 *Heteropallene dubitans* HODGSON, Denkschr. Med.-Nat. Ges. Jena, **16**: 226.
 1938 *Procidella gibber* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 172-174, fig. 5.
 1953 *Procidella gibber*, - STOCK, Beaufortia, **4** (35): 38-39, fig. 3a-e.
 1954 *Metapallene dubitans*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 114-115, fig. 14a.

Distribution: South Africa.

***Mimipallene* CHILD 1982**

- 1982 *Mimipallene* Child, Smiths. Contr. Zool., **349**: 27 [diagnosis, remarks].

Mimipallene atlantis CHILD 1982

1982 *Mimipallene atlantis* CHILD, Smiths. Contr. Zool., **349**: 27-29, fig. 8.

Distribution: Argentine Basin and Slope, off Rio de la Plata.

Depth range: 993-2323m.

Neopallene DOHRN 1881

1954 *Neopallene*, - Stock, Vidensk. Medd. Dansk naturh. Foren., **116**: 34 [diagnosis].

Neopallene antipoda STOCK 1954

1954 *Neopallene antipoda* STOCK, Vidensk. Medd. dansk. naturhist. Foren., **116**: 34-36, figs. 13-14.

Distribution: New Zealand (N.W. of Cape Maria van Diemen).

Depth range: 55m.

Habitat: hard bottom.

Neopallene azorensis ARNAUD 1974

1974 *Neopallene azorensis* ARNAUD, Bull. Zool. Mus. Univ. Amsterdam, **3** (21): 173-174, figs. 8-14.

Distribution: Azores.

Depth range: 1240-2100m.

Habitat: "fond de vase sableuse" (ARNAUD).

Neopallene campanellae DOHRN 1881

1881 *Neopallene campanellae* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 200-202, pl. XV, figs. 11-15.

1923 *Neopallene campanellae*, - BOUVIER, Faune de France, **7**: 32.

1987 *Neopallene campanellae*, - ARNAUD, Mesogee, **47**: 50.

Distribution: Mediterranean.

Depth range: 70-90m.

Habitat: "dans des algues Mélobésiéées mortes et roulées ("Melobesien-Geröll")" (ARNAUD).

Oropallene SCHIMKEWITSCH 1930

1954 *Oropallene*, - Stock, Vidensk. Medd. Dansk naturh. Foren., **116**: 28-29 [diagnosis].

Oropallene dimorpha (HOEK 1898)

- 1898 *Pallene dimorpha* HOEK, Tijdschr. Nederl. Dierk. Ver., (2) **5** (2-4): 290-293, pl. II, figs. 1-6.
 1930 *Oropallene dimorpha*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 245, 291-292.
 1944 *Oropallene dimorpha*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 36.
 1954 *Oropallene dimorpha*, - STOCK, Vidensk. Medd. dansk. naturhist. Foren., **116**: 29.
 1987 *Oropallene dimorpha*, - CHILD, Proc. biol. Soc. Wash., **100** (4): 915.

Distribution: SW-Pacific, NE of Macquarie Island.**Depth range:** 93-124m.***Oropallene minor*** CLARK 1963

- 1963 *Oropallene minor* CLARK, Rec. Austr. Mus., **26** (1): 18-21, fig. 10A-L.

Distribution: Australia.**Depth range:** 39-144m.**Habitat:** on mud and from conglomerate boulder.***Oropallene ovigerosetosus*** HILTON 1942

- 1942 *Oropallene ovigerosetosus* HILTON, Occ. Pap. Bernice P. Bishop Mus., **17** (3): 43-44, fig. 1.

Distribution: Hawaii.***Oropallene polaris*** HEDGPETH 1963

- 1963 *Oropallene polaris* HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1334-2336, fig. 8.

Distribution: Arctica (Pt. Barrow).**Depth range:** 66m.**Habitat:** rocks, stones.***Pallenoides*** STOCK 1951

- 1954 *Pallenoides*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 115 [diagnosis].
 1955 *Pallenoides*, - Stock, Vidensk. Medd. Dansk naturh. Foren., **117**: 226-227 [remarks].
 1975 *Pallenoides*, - Stock, Bull. mar. Sci., **24** (4): 1015 [key to species].

Pallenoides amazonica STOCK 1975

- 1975 *Pallenoides amazonica* STOCK, Bull. mar. Sci., **24** (4): 1012-1015, figs. 23-24.

1992 *Pallenoides amazonica*, - STOCK, Tijdschr. Ent., **135**: 129, 139.

Distribution: Brazil (near mouth of Amazonas Rivier; other records along the north coast of Brazil, between Amazonas and Fortaleza).

Depth range: 24-72m.

***Pallenoides magnicollis* STOCK 1951**

1951 *Pallenoides magnicollis* STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **2** (43): 11-12, figs. 7-13.

1954 *Pallenoides magnicollis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 115-116, fig. 14b-c.

Distribution: South Africa.

Depth range: 15m.

Habitat: among algae.

***Pallenoides opuntia* STOCK 1965**

1965 *Pallenoides opuntia* STOCK, Beaufortia, **13** (151): 24-26, figs. 28-37.

Distribution: Madagascar (region of Fort Dauphin).

Depth range: 50m.

Habitat: shelly sand.

***Pallenoides proboscideum* BARNARD 1955**

1955 *Pallenoides proboscideum* BARNARD, Ann. S. Afr. Mus., **43** (1): 105-106, fig. 53d-e.

Distribution: South Africa (Algoa Bay).

Depth range: 5-7m.

***Pallenoides spinulosa* STOCK 1955**

1955 *Pallenoides* (?) *spinulosa* STOCK, Vidensk. Medd. Dansk naturhist. Foren., **117**: 227-230, figs. 6-7.

1970 *Pallenoides spinulosa*, - CAPRILES, Carrib. J. Sci., **10** (1-2): 105.

1975 *Pallenoides spinulosa*, STOCK, Bull. mar. Sci., **24** (4): 1012.

1986 *Pallenoides spinulosa*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1.

Distribution: Caribbean.

Depth range: 28-586m.

Habitat: *Sargassum* on pilings.

***Pallenoides stylirostrum* STOCK 1973**

1973 *Pallenoides stylirostrum* STOCK, Beaufortia, **20** (266): 113-115, fig. 6.

Distribution: SE-Australia (West Island, Toad Head).

Depth range: 5m.

Habitat: on *Acrocarpia paniculata* (Phaeophyta).

***Parapallene* CARPENTER 1892**

- 1954 *Parapallene*, - BARNARD, Ann. S. Afr. Mus., **108** [diagnosis, key to South African species].
 1954 *Parapallene*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 50 [diagnosis, remarks].
 1963 *Parapallene*, - CLARK, Rec. Austr. Mus., **26** (1): 31 [key to Australian species].
 1986 *Parapallene*, - STOCK, Bull. mar. Sci., **38** (3): 421 [remarks].
 1991 *Parapallene*, - STOCK, Mém. Mus. natn. Hist. nat., (A) **151**: 191-194 [key to species].
 1992 *Parapallene*, - CHILD, Mem. Hourglass Cruises, **9** (1): 61 [diagnosis].

***Parapallene algoae* BARNARD 1946**

- 1946 *Parapallene algoae* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 61.
 1954 *Parapallene algoae* BARNARD, Ann. S. Afr. Mus., **41** (3): 112-113, fig. 12.
 1956 *Parapallene algoae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 81-82, fig. 5.
 1962 *Parapallene algoae*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 284.
 1988 *Parapallene algoae*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 143.
 1991 *Parapallene algoae*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: Southern Africa.

Depth range: 49-550m.

Habitat:

***Parapallene arnaudae* STOCK 1991**

- 1991 *Parapallene arnaudae* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 191, figs. 45-46, 193 [key].

Distribution: New Caledonia.

Depth range: 57m.

***Parapallene australiensis* (HOEK 1881)**

- 1881 *Pallene australiensis* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 76-78, pl. 11, figs. 1-7 [part.].
 1884 *Pallene australiensis*, - HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1022.
 1892 *Parapallene australiensis*, - CARPENTER, Sci. Proc. Roy. Dublin Soc., **7** (5): 553.
 1908 *Parapallene australiensis*, - LOMAN, Siboga Exped. Monogr., **40**: 48.
 1937 *Parapallene australiensis*, - CALMAN, Ann. Mag. Nat. Hist., **10** (20): 531-532.
 1954 *Parapallene australiensis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 50, fig. 24d-e.
 1963 *Parapallene australiensis*, - CLARK, Rec. Austr. Mus., **26** (1): 25-26.

- 1973 *Parapallene australiensis*, - STOCK, Beaufortia, **20** (266): 119.
 1975 *Parapallene australiensis*, - CHILD, Smiths. Contr. Zool., **190**: 12.
 1982 *Parapallene australiensis*, - STAPLES, Mem. Qd. Mus., **20** (3): 455.
 1991 *Parapallene australiensis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 190, 194 [key].

Distribution: Australia, New Caledonia.

Depth range: 44-137m.

Habitat: sand, sponges and bryozoa; also found on the hydroid *Halopteris buski*.

***Parapallene avida* STOCK 1973**

- 1973 *Parapallene avida* STOCK, Beaufortia, **20** (266): 119-121, fig. 9.
 1973 *Parapallene avida*, - STOCK, Beaufortia, **21** (279): 92, 95-96, fig. 3a-c.
 1980 *Parapallene avida*, - HOOPER, Aust. Zool. **20** (3): 473-483 (reproductive biology).
 1991 *Parapallene avida*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].

Distribution: Australia.

Depth range: 0-28m.

Habitat: red algae, from *Sargassum*, and in association with hydroids (*Aglaophenia plumosa*).

***Parapallene bermudensis* LEBOUR 1949**

- 1949 *Parapallene bermudensis* LEBOUR, Proc. Zool. Soc. London, **118** (4): 930.
 1982 *Parapallene bermudensis*, - STOCK, Smiths. Contr. mar. Sci., **12**: 366-367, fig. 165.
 1991 *Parapallene bermudensis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].
 1992 *Parapallene bermudensis?*, - CHILD, Mem. Hourglass Cruises, **9** (1): 59 [key], 61-62, fig. 28.

Distribution: Bermuda to Belize.

Depth range: 27-55m.

Habitat: coral, sand and rubble substrate from behind outer coral reef ridge.

***Parapallene calmani* FLYNN 1928**

- 1928 *Parapallene calmani* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 16, figs. 7-8.
 1954 *Parapallene calmani*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 110-111, fig. 11.
 1962 *Parapallene calmani*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 284.
 1991 *Parapallene calmani*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].

Distribution: Southern Africa.

Depth range: 37-86m.

Habitat: sand and rocks; sand and shells.

Parapallene capillata STOCK 1954

- 1954 *Parapallene capillata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 52-57, figs. 22-23, 24a-c.
 1991 *Parapallene capillata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: Singapore.**Depth range:** low tide.***Parapallene challengereri*** CALMAN 1937

- 1937 *Parapallene challengereri* CALMAN, Ann. Mag. Nat. Hist., **10** (20): 532-534, figs. 1-6.
 1991 *Parapallene challengereri*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].

Distribution: SE-Australia.**Depth range:** 73 or 220m.***Parapallene exigua*** STOCK 1954

- 1954 *Parapallene exigua* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 57-59, fig. 25.
 1991 *Parapallene exigua*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: New Zealand (Three Kings).**Depth range:** 119m.**Habitat:** hard bottom.***Parapallene famelica*** FLYNN 1929

- 1929 *Parapallene famelica* FLYNN, Mem. Qd. Mus., **9** (3): 258-260, figs. 6-9.
 1963 *Parapallene famelica* CLARK, Rec. Austr. Mus., **26** (1): 28, fig. 14A-G.
 1991 *Parapallene famelica*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: E-Australia (off Port Philip; near Lindeman Island, Whitsunday Passage).**Depth range:** 17m.***Parapallene haddoni*** CARPENTER 1892

- 1892 *Parapallene haddoni* CARPENTER, Sci. Proc. Roy. Dublin Soc., **7** (5) (40): 553-555, pl. 22, figs. 1-6.
 1908 *Parapallene haddoni*, - LOMAN, Siboga Exped. Monogr., **40**: 44.
 1963 *Parapallene haddoni*, - CLARK, Rec. Austr. Mus., **26** (1): 26-18, fig. 13A-J.
 1975 *Parapallene haddoni*, - CHILD, Smiths. Contr. Zool., **190**: 12.
 1991 *Parapallene haddoni*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: Australia (Torres Straits; Holothuria Bank; Bluff Point; Rottnest Island).

Depth range: 27-139m.

Parapallene hodgsoni BARNARD 1946

- 1946 *Parapallene hodgsoni* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 61.
 1954 *Parapallene hodgsoni*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 112-113, fig. 13.
 1965 *Parapallene hodgsoni*, - STOCK, Beaufortia, **13** (151): 26-28, figs. 38-44.
 1968 *Parapallene hodgsoni*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 39.
 1991 *Parapallene hodgsoni*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: Southern Africa; Madagascar (Fort Dauphin region); off Lorenzo Marquez.

Depth range: 22-73m.

Habitat: shelly sand; rocks.

Parapallene hospitalis LOMAN 1908

- 1908 *Parapallene hospitalis* LOMAN, Siboga Exped. Monogr., **40**: 45-46, pl. VIII, figs. 102-111.
 1923 *Parapallene hospitalis*, - CALMAN, Rec. Indian Mus., **24** (3): 278-279.

Distribution: SE-Asia.

Depth range: 82-88m.

Parapallene invertichelata ARNAUD & CHILD 1988

- 1988 *Parapallene invertichelata* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 143-146, fig. 5.
 1991 *Parapallene invertichelata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: South Africa (Zululand area).

Depth range: 550m.

Parapallene longipes CALMAN 1938

- 1938 *Parapallene longipes* CALMAN, Sci. Rep. John Murray Exped., **5** (6): 156-159, figs. 5-6.
 1990 *Parapallene longipes*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (10): 106.
 1991 *Parapallene longipes*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].
 1991 *Parapallene longipes*, - TURPAEVA, Zool. Zh., **70** (12): 39-41, fig. 3.

Distribution: shelf and slope of south-eastern Africa,

Zanzibar.

Depth range: 183-194m.

Parapallene nierstraszi LOMAN 1908

- 1908 *Parapallene nierstraszi* LOMAN, Siboga Exped. Monogr., **40**: 44-45, pl. IX, figs. 122-127.
 1928 *Parapallene nierstraszi*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep. 1): 18.
 1938 *Parapallene nierstraszi*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 158-159, fig. 7.
 1954 *Parapallene nierstraszi*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 113.
 1954 *Parapallene nierstraszi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 52, fig. 24f.
 1955 *Parapallene nierstraszi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 18-19, fig. 10.
 1971 *Parapallene nierstraszi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 323.
 1991 *Parapallene nierstraszi*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 41.
 1991 *Parapallene nierstraszi*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 223.

Distribution: Japan, Philippines, Indonesia, Western Australia, South Africa.

Depth range: 73-84m.

Habitat: Lithothamnion bottom.

Parapallene obtusirostris CLARK 1963

- 1963 *Parapallene obtusirostris* CLARK, Rec. Austr. Mus., **26** (1): 29-31, fig. 15A-I.
 1991 *Parapallene obtusirostris*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194 [key].

Distribution: Australia (Port Philip, Victoria).

Parapallene parviunguicularis STOCK 1986

- 1986 *Parapallene parviunguicularis* STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1, 421-424, fig. 7.
 1991 *Parapallene parviunguicularis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].

Distribution: Florida.

Depth range: 851-933m.

Parapallene spinosa (MÖBIUS 1902)

- 1902 *Anoplodactylus spinosus* MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. Valdivia, 1898-1899, **3**: 188-189, pl. 28, figs. 8-12.
 1954 *Parapallene spinosa*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 108-110, fig. 10a-e.
 1956 *Parapallene spinosa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren, **118**:

82, fig. 6.

- 1962 *Parapallene spinosa*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 284.
 1991 *Parapallene spinosa*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 193 [key].

Distribution: Southern Africa.

Depth range: 10-75m.

Habitat: sand.

***Pigrogromitus* CALMAN 1927**

- 1992 *Pigrogromitus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 62-64 [diagnosis].

***Pigrogromitus timsanus* CALMAN 1927**

- 1927 *Pigrogromitus timsanus* CALMAN, Trans. Zool. Soc. London, **22** (3): 408-410, fig. 104a-f.
 1942 *Clotenopsa prima* HILTON, Occ. Pap. Bernice P. Bishop Mus., **17** (3): 52-53, fig. 8.
 1948 *Pigrogromitus timsanus*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 214-216, fig. 23.
 1968 *Pigrogromitus timsanus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 46.
 1971 *Pigrogromitus timsanus*, - LIPKIN & SAFRIEL, - J. Ecol., **59** (1): 9.
 1972 *Pigrogromitus timsanus*, - ARNAUD, Tethys, Suppl., **3**: 159-160.
 1975 *Pigrogromitus timsanus*, - STOCK, Bull. mar. Sci., **24** (4): 1015-1016.
 1976 *Pigrogromitus timsanus*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 134, 158.
 1976 *Pigrogromitus timsanus*, - REIMER, Mar. Biol., **35**: 229.
 1979 *Pigrogromitus timsanus*, - CHILD, - Smiths. Contr. Zool., **293**: 46.
 1982 *Pigrogromitus timsanus*, - CHILD, Smiths. Contr. mar. Sci., **12**: 367-368.
 1982 *Pigrogromitus timsanus*, - STAPLES, Mem. Qd. Mus., **20** (3): 457.
 1986 *Pigrogromitus timsanus*, - STOCK, Bull. mar. Sci. **38** (3): 403 tab. 1.
 1987 *Pigrogromitus timsanus*, - ARNAUD, Mesogeee, **47**: 80.
 1988 *Pigrogromitus timsanus*, - CHILD, Smiths. Contr. Zool., **468**: 21.
 1988 *Pigrogromitus timsanus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 667.
 1989 *Pigrogromitus timsanus*, - STOCK, Bijdr. Dierk., **59** (2): 95.
 1990 *Pigrogromitus timsanus*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (9): 97-98.
 1990 *Pigrogromitus timsanus*, - MÜLLER, Bonn. zool. Beitr. **41** (1): 74.
 1990 *Pigrogromitus timsanus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **103** (2): 304-310.
 1991 *Pigrogromitus timsanus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 194-195.
 1992 *Pigrogromitus timsanus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 59 [key], 64-65, fig. 29.
 1992 *Pigrogromitus timsanus*, - STOCK, Bijdr. Dierk., **62** (2): 93.

Distribution: primarily circumtropical, some records from subtropical areas.

Depth range: 0-108 (2050)m.

Habitat: in a variety of substrata, often among algae in

exposed locations; feeds on small anthozoans.

Propallene SCHIMKEWITSCH 1909

- 1954 *Propallene*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 31 [diagnosis].
 1975 *Propallene*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (11): 89-90 [remarks], 93 [diagnosis], 94 [key to species].
 1983 *Propallene*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 61 [key to Japanese species].

Propallene ardua STOCK 1975

- 1975 *Propallene ardua* STOCK, Trav. Mus. Hist. Nat. "Gr. Antipa", **16**: 131-132.
 1975 *Propallene ardua*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (11): 92-93, figs. 27-41.

Distribution: Tanzania.

Depth range: 0-24m.

Habitat: in *Syringodium* grass; sandy bottom, and washings of brown algae.

Propallene crassimanus STOCK 1959

- 1959 *Propallene crassimanus* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 559-561, fig. 5.
 1968 *Propallene crassimanus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 37.
 1992 *Propallene crassimanus*, - STOCK, Bijdr. Dierk., **62** (2): 93-94, fig. 10.

Distribution: South Africa; off Lorenzo Marquez; Oman.

Depth range: 1-22m.

Habitat: sand, rocks, and algae.

Propallene crinipes STOCK 1968

- 1968 *Propallene crinipes* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 35-37, fig. 13.

Distribution: Strait of Malacca.

Propallene curtipalpus CHILD 1988

- 1988 *Propallene curtipalpus* CHILD, Smiths. Contr. Zool., **468**: 21-23, fig. 9.
 1988 *Propallene curtipalpus*, - CHILD, Proc. biol. Soc. Wash., **101** (3): 667-668.

Distribution: Japan (Ryukyu Islands); Philippines (Siquijor Island).

Depth range: 0-20m.

Habitat: coral rubble, sand, on mud flats.

Propallene cyathus STAPLES 1979

1979 *Propallene cyathus* STAPLES, Trans. Roy. Soc. S. Austr., **103** (3-4): 85-90, figs. 1A-P, 2A-B.

Distribution: Australia (Bass Strait).

Depth range: 13m.

Propallene kempfi (CALMAN 1923)

- 1923 *Parapallene kempfi* CALMAN, Rec. Indian Mus., **25** (3): 277-278, fig. 6.
 1937 *Propallene kempfi*, - CALMAN, Ann. Mag. Nat. Hist., **20** (10): 534.
 1950 *Propallene kempfi*, - GNANAMUTHU, Proc. Zool. Soc. Bengal, **3** (1): 39-47, figs. 1-4.
 1954 *Propallene kempfi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 33-34, fig. 12c-f.
 1972 *Propallene kempfi*, - ARNAUD, Tethys, Suppl. **3**: 161-162, figs. 3-6.
 1975 *Propallene kempfi*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (11): 3.

Distribution: Gulf of Manaar, Waltair (Madras); Madagascar; Singapore.

Depth range: 0-4m.

Habitat: among weeds (*Diplanthera uninervis*; *Syringodium isoetifolium*; *Cymodocea serrulata*; *Halophila cf. minor*).

Propallene longiceps (BÖHM 1879)

- 1879 *Pallene longiceps* BÖHM, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1879 (4): 59-60.
 1881 *Pallene longiceps*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 30.
 1890 *Pallene longiceps*, - ORTMANN, Zool. Jb. (Syst.), **5** (1): 165-166, pl. 24, fig. 7.
 1899 *Pallene longiceps*, - SHISHIDO, Dôzatsu, **11** (128): 199.
 1919 *Pallene longiceps*, - FUKUI, Rigakukai[?], **16** (2): 99.
 1930 *Pallene longiceps*, - OHSHIMA, Dôbutsgaku Zasshi, **42** (495): 34-35.
 1930 *Propallene longiceps*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 282-287.
 1933 *Pallene longiceps*, - OHSHIMA, Annot. Zool. Jap., **14** (2): 212-216, figs. 1-7.
 1933 *Pallene longiceps*, - OHSHIMA, Dôbutsgaku Zasshi, **45** (532-533): 96.
 1936 *Propallene longiceps*, - OHSHIMA, Dôbutsgaku Zasshi, **48** (8-10): 863.
 1954 *Propallene longiceps*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 31.
 1959 *Propallene longiceps*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 199.
 1962 *Propallene longiceps*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 96.
 1965 *Propallene longiceps*, - UTINOMI, New Illustr. Encyclopaedia Fauna Japan, **2**: 334.
 1971 *Propallene longiceps*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.
 1975 *Propallene longiceps*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (11): 90-92, 94, figs. 1-20.

- 1980 *Propallene longiceps*, - NAKAMURA & SEKIGUCHI, Mar. Ecol., Progress Ser., **2**: 163-167, figs. 1-5 [mating behavior and oviposition].
1981 *Propallene longiceps*, - NAKAMURA, J. nat. Hist., **15**: 49-61, figs. 1-17 [post embryonic development].
1983 *Propallene longiceps*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 61-62.
1983 *Propallene longiceps*, - NAKAMURA & SEKIGUCHI, Bull. Biogeogr. Soc. Japan, **38** (6): 39.
1986 *Propallene longiceps*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 41, fig. 4a-1.
1987 *Propallene longiceps*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 158.
1987 *Propallene longiceps*, - NAKAMURA, Biol. Lab., Imperial Household: 14.
1988 *Propallene longiceps*, - NAKAMURA & CHILD, - Proc. biol. Soc. Wash., **101** (3): 668.
1991 *Propallene longiceps*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 41-42.

Distribution: Japan and Korea.

Depth range: 6-103m.

Habitat: gravel.

***Propallene saengeri* STAPLES 1979**

- 1979 *Propallene saengeri* STAPLES, Trans. Roy. Soc. S. Austr., **103** (3-4): 90-93, figs. 2D, 4A-L.
1982 *Propallene saengeri*, - STAPLES, Mem. Qd. Mus., **20** (3): 456.
1983 *Propallene saengeri*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 62.

Distribution: Australia and Japan.

Depth range: 4.8-30m.

Habitat: soft mud.

***Propallene similis* BARNARD 1955**

- 1955 *Propallene similis* BARNARD, Ann. S. Afr. Mus., **43** (1): 105.
1959 *Propallene similis*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 558-559, fig. 4.

Distribution: South Africa.

Depth range: 9-10m.

Habitat: rock; sand, broken shell, and rocky patches.

***Propallene stocki* FAGE 1956**

- 1956 *Propallene stocki* FAGE, Bull. Mus. natn. Hist. nat., Paris, **2** (28) (3): 291-295, figs. 2-6.

Distribution: Sierra Leone.

***Propallene vagus* STAPLES 1979**

- 1979 *Propallene vagus* STAPLES, Trans. Roy. Soc. S. Austr., **103** (3-4): 88-90, figs. 2C, 3A-N.

Distribution: Australia (Port Phillip).

Pseudopallene WILSON 1878

- 1943 *Pseudopallene*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 88.
 1948 *Pseudopallene*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 205 [diagnosis, remarks, key to species].
 1948 *Cordylochele*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 206-207 [diagnosis, remarks, key to W-Atlantic species].
 1953 *Pseudopallene*, - Stock, Temminckia, **9**: 294-297 [remarks].
 1954 *Pseudopallene*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 107 [diagnosis, remarks].
 1954 *Pseudopallene*, - Stock, Vidensk. Medd. Dansk naturh. Foren., **116**: 61 [diagnosis].

Pseudopallene ambigua STOCK 1956

- 1956 *Pseudopallene ambigua* STOCK, Mitt. Zool. Mus. Hamburg, **54**: 40-42, fig. 5.
 1963 *Pseudopallene ambigua*, - CLARK, Rec. Austr. Mus., **26** (1): 31-33, fig. 16A-F.
 1973 *Pseudopallene ambigua*, - STOCK, Beaufortia, **20** (266): 115-117, fig. 7.
 1973 *Pseudopallene ambigua*, - STOCK, Beaufortia, **21** (279): 92, 95.

Distribution: Australia.

Depth range: 3-126m.

Habitat: red algae, coralline algae, bryozoa.

Pseudopallene brevicollis (SARS 1891)

- 1891 *Cordylochele brevicollis*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 51-54, pl. IV, fig. 3a-g.
 1933 *Cordylochele brevicollis*, - STEPHENSEN, Medd. Grönland, **79** (6): 25.

Distribution: Arctica. - Greenland; Iceland; Kara Sea; Barents Sea; Spitsbergen; Siberian Polar Sea.

Depth range: littoral-860m.

Pseudopallene centrotus Pushkin, 1990

- 1990 *Pseudopallene centrotus* Pushkin, Zool., Zh., **69** (2): 155-157, pl. 1, figs. 1-7.

Distribution: Antarctica.

Depth range: 250m.

Pseudopallene circularis (GOODSIR 1842)

- 1842 *Pallene circularis* GOODSIR, Edinburgh New Phil. J., **33**: 136, pl. 3.
 1853 *Pallene hispida*, STIMPSON, Smiths. Contr. Knowl., **6** (5): 37.
 1878 *Pseudopallene hispida*, - WILSON, Am. J. Sci., **15**: 200.

- 1878 *Pseudopallene hispida*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 10, pl. 3, fig. 1a-e.
- 1878 *Pseudopallene discoidea*, WILSON, Trans. Conn. Acad. & Sci., **5**: 12, pl. 3, fig. 3a-c.
- 1880 *Pseudopallene hispida*, - WILSON, Rep. U.S. Fish. Comm., **6**: 479-480, pl. 2, fig. 10.
- 1880 *Pseudopallene discoidea*, - WILSON, Rep. U.S. Fish. Comm., **6**: 479.
- 1891 *Pseudopallene circularis*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 38-42, pl. III, fig. 3a-h.
- 1899 *Pseudopallene circularis*, - MEINERT, Danish Ingolf-Exped., **3** (1): 50-51.
- 1901 *Pseudopallene hispida*, - WHITEAVES, Ann. Mag. Nat. Hist., **4** (10): 263.
- 1908 *Phoxichilus circularis*, - NORMAN, J. Linn. Soc. Lond. (Zool.), **30**: 207.
- 1933 *Pseudopallene circularis*, - STEPHENSEN, Medd. om Grönland, **79** (6): 20-21.
- 1938 *Spicularia discoidea*, - HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 177-179, fig. 8.
- 1940 *Halosoma discoideum*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 46.
- 1943 *Pseudopallene circularis*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 88.
- 1943 *Pseudopalene circularis*, - NEEDLER, Can. Atl. Fauna, **10**: 12, fig. 15a-d.
- 1948 *Pseudopallene circularis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 206, fig. 19.
- 1963 *Pseudopallene circularis*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1333-1334.

Distribution: Boreal-Arctic, from Okhotsk Sea to NE-America, perhaps circumpolar.

Depth range: 10-135m.

Habitat: shells, hydroids.

Pseudopallene dubia CLARK 1963

- 1963 *Pseudopallene dubia* CLARK, Rec. Austr. Mus., **26** (1): 33-35, figs. 17A-H, 18A-B.

Distribution: Tasmania.

Depth range: shallow water.

Habitat: "in kelp weed" (CLARK).

Pseudopallene gilchristi FLYNN 1928

- 1928 *Pseudopallene gilchristi* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 23, fig. 13.
- 1954 *Pseudopallene gilchristi*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 107.
- 1968 *Pseudopallene gilchristi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 39, fig. 14e-g.
- 1988 *Pseudopallene gilchristi*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 146.

Distribution: apparently restricted to the Indian Ocean from Transkei and Port Natal to Lourenço Marques.

Depth range: 73-775m.

Pseudopallene glutus PUSHKIN 1975

1975 *Pseudopallene glutus* PUSHKIN, Zool. Zh., **54** (9): 1402-1404, fig. 1.

Distribution: subantarctic.

Depth range: 320m.

***Pseudopallene longicollis* (SARS 1888)**

- 1888 *Cordylochele longicollis* SARS, Arch. Math. Naturvidensk., Oslo, **12**: 344.
 1891 *Cordylochele longicollis*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 49-51, pl. IV, fig. 2a-g.
 1933 *Cordylochele longicollis*, - STEPHENSEN, Medd. Grönland, **79** (6): 25-26.
 1948 *Cordylochele longicollis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 207-209, fig. 20b.

Distribution: NE-Atlantic, south of the Wyville Thomson Ridge and in Davis Strait.

Depth range: 494-1326m.

***Pseudopallene malleolata* (SARS 1879)**

- 1879 *Pallene malleolata* SARS, Arch. Math. Naturvidensk., Kristiana, **4**: 48.
 1883 *Pallene malleolata*, - HOEK, Trans. Roy. Soc. Edinburgh, **32**: 6-7, pl. I, fig. 7.
 1891 *Cordylochele malleolata*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 45-49, pl. IV, fig. 1a-k.
 1933 *Cordylochele malleolata*, - STEPHENSEN, Medd. Grönland, **79** (6): 25.
 1936 *Cordylochele malleolata*, - STEPHENSEN, Bergens Mus. Arbok Naturv. Rek., **2**: 26-27.
 1948 *Cordylochele malleolata*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 207.

Distribution: around Iceland, Spitsbergen, Faroes, Kara Sea; westward occurring in Denmark and Davis Straits to latitude 66°35'N.

Depth range: 348-1512m.

***Pseudopallene pachycheira* HASWELL 1884**

- 1884 *Pallene pachycheira* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1030-1031, pl. LVII, figs. 6-9.
 1908 *Parapallene pachycheira*, - LOMAN, Siboga Exped. Monogr., **40**: 47-48.
 1918 *Pseudopallene pachycheira*, - FLYNN, Pap. Proc. Roy. Soc. Tasm., 1919: 77-79, pl. XIX, fig. 9, pl. XX, figs. 10-11.

Distribution: Australia (Port Jackson).

Depth range: shallow water.

***Pseudopallene pacifica* LOSINA-LOSINSKY 1961**

- 1961 *Pseudopallene pacifica* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 77-78, fig. 10.

Distribution: NW-Pacific.
Depth range: 138m.

Pseudopallene spinipes (FABRICIUS 1780)

- 1891 *Pseudopallene spinipes* SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 42-44, pl. III, fig. 4a-g.
 1933 *Pseudopallene spinipes*, - STEPHENSEN, Medd. Grønland, **79** (6): 21.
 1936 *Pseudopallene spinipes*, - STEPHENSEN, Bergens. Mus. Arbok Naturv. Rek., **2**: 25.
 1943 *Pseudopallene spinipes*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 88.
 1963 *Pseudopallene spinipes*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1334.

Distribution: Arctica, known from western Norway, Kara Sea, Franz Josef Land and West Greenland.

Depth range: sublittoral-380m.

Habitat: gravel; seaweed.

Pseudopallene zamboangae STOCK 1953

- 1953 *Pseudopallene zamboangae* STOCK, Temminckia, **9**: 297-300, figs. 12-13.
 1954 *Pseudopallene zamboangae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren, **116**: 61-63, fig. 27d-f.
 1971 *Pseudopallene zamboangae*, - STOCK, Publ. Seto Mar. Biol. Lab., **18** (5): 323.
 1991 *Pseudopallene zamboangae*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 223.

Distribution: Kei Islands (Indonesia), Philippines.

Depth range: 17-84m.

Habitat: sand, corals.

Pycnothea LOMAN 1921

- 1921 *Pycnothea* LOMAN, Nat. Hist. Juan Fernandez, **3**: 137-138 [diagnosis].

Pycnothea flynni WILLIAMS 1940

- 1940 *Pycnothea flynni* WILLIAMS, J. Roy. Soc. W.-Australia, **25**: 202-204, figs. 6-9.
 1963 *Pycnothea flynni*, - CLARK, Rec. Austr. Mus., **26** (1): 46-48, fig. 23.
 1972 *Pycnothea flynni*, - ARNAUD, Tethys, Suppl. 3: 160-161, fig. 2.
 1973 *Pycnothea flynni*, - CLARK, Pac. Sci., **27** (1): 30.
 1973 *Pycnothea flynni*, - STOCK, Beaufortia, **266** (20): 121.
 1975 *Pycnothea flynni*, - CHILD, Smiths. Contr. Zool., **190**: 16.
 1980 *Pycnothea flynni*, - HOOPER, Aust. Zool., **20** (3): 475.

Distribution: Australia; New Britain; Madagascar.

Depth range: 0-2.5m.

Habitat: among algae (*Sargassum*; *Gelidium*); under rocks and stones; boulders with detritus.

Pycnothea selkirkii LOMAN 1921

1921 *Pycnothea selkirkii* LOMAN, Nat. Hist. Juan Fernandez, **3**: 138-142, fig. 1A-F.
 1992 *Pycnothea selkirkii*, - CHILD, Smiths. Contr. Zool., **526**: 30-31, fig. 12.

Distribution: Chile (Isla Juan Fernandez; Isla San Felix).
Depth range: 0-40m.
Habitat: on coralline algae.

Queubus BARNARD 1946

1954 *Queubus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 156 [diagnosis, remarks].

Queubus jamesanus BARNARD 1946

1946 *Queubus jamesanus* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 63.
 1954 *Queubus jamesanus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 157-158, fig. 34.
 1955 *Queubus jamesanus*, - BARNARD, Ann. S. Afr. Mus., **43** (1): 107.
 1959 *Queubus jamesanus*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 562-563.

Distribution: South Africa.
Depth range: shallow water, low tide.
Habitat: in pools under granite boulders.

Safropallene ARNAUD & CHILD 1988

1988 *Safropallene* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 147 [diagnosis, remarks].

Safropallene longimana ARNAUD & CHILD

1988 *Safropallene longimana* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 147-150,
 fig. 6.

Distribution: South Africa (Transkei and Durban areas)
Depth range: 710-850m.

Seguapallene PUSHKIN 1975

1975 *Seguapallene* PUSHKIN, Zool. Zh., **54** (9): 1404 [diagnosis].
 1983 *Seguapallene*, - Child, Proc. biol. Soc. Wash., **96** (4): 709 [diagnosis, remarks].

Seguapallene aculeata (STOCK 1954)

1954 *Parapallene(?) aculeata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 59-61, figs. 26, 27a-c.

Distribution: W-Atlantic.

Seguapallene crassa CHILD 1990

1990 *Seguapallene crassa* CHILD, Proc. biol. Soc. Wash., **103** (2): 321-324, fig. 4.

Distribution: Australia (Great Barrier Reef, Lizard Island).

Depth range: 2m.

Habitat: rubble bank; small rubble in sand; coral with *Halimeda* clumps and red algae.

Seguapallene echinata (CALMAN 1938)

1938 *Parapallene echinata* CALMAN, Sci. Rep. John Murray Exped., **5** (6): 154-156, fig. 4A-E.

Distribution: Indian Ocean. - Arabian Coast.

Depth range: 40m.

Seguapallene insignata PUSHKIN 1975

1975 *Seguapallene insignatus* PUSHKIN, Zool. Zh., **54** (9): 1404-1405, fig. 2.

1991 *Seguapallene insignata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 190 [text].

Distribution: Kerguelen Islands.

Depth range: 3-30m.

Seguapallene micronesica CHILD 1983

1983 *Seguapallene micronesica* CHILD, Proc. biol. Soc. Wash., **96** (4): 709-711, fig. 4.

1991 *Seguapallene micronesica*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 145.

Distribution: Palau, SE Koror Island; Guam.

Depth range: 0.1-11 m.

Habitat: sea reef inside cove next to cave, from sponges; on dead branching coral.

Seguapallene tricuspidata STOCK 1991

1991 *Seguapallene tricuspidata* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 188-190, fig. 44.

Distribution: New Caledonia.

Depth range: 570-610 m.

Spasmopallene STOCK 1968

1968 *Spasmopallene* STOCK, Vidensk. Medd. Dansk naturh. Foren., **131**: 39-40 [diagnosis, remarks].

***Spasmopallene clarki* STOCK 1968**

1968 *Spasmopallene clarki* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 42-44, fig. 16.

Distribution: Great Australian Bight.

Depth range: 72m.

Habitat: corals.

***Spasmopallene reflexa* STOCK 1968**

1968 *Spasmopallene reflexa* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 40-42, fig. 15.

Distribution: Great Australian Bight.

Depth range: 72m.

Habitat: corals.

***Stylopallene* CLARK 1963**

1963 *Stylopallene* CLARK, Rec. Austr. Mus., **26** (1): 35-36 [diagnosis], 42 [key to species].

***Stylopallene cheilorhynchus* CLARK 1963**

1963 *Stylopallene cheilorhynchus* CLARK, Rec. Austr. Mus., **26** (1): 36-38, fig. 19A-I.

1973 *Stylopallene cheilorhynchus*, - STOCK, Beaufortia, **20** (266): 117.

1973 *Stylopallene cheilorhynchus*, - STOCK, Beaufortia, **21** (279).

Distribution: Australia (Pearson Island; West Island; Tasmania).

Depth range: 12-90m.

Habitat: on bryozoans; in caves; on hydroids; among kelp.

***Stylopallene dorsospinum* CLARK 1963**

1963 *Stylopallene dorsospinum* CLARK, Rec. Austr. Mus., **26** (1): 38-40, fig. 20A-I.

Distribution: Australia (Twofold Bay; near Botany Bay).

Depth range: 90-93m.

***Stylopallene longicauda* STOCK 1973**

1973 *Stylopallene longicauda* STOCK, Beaufortia, **20** (266): 117-118, fig. 8.

Distribution: SE-Australia (Western Port).

Depth range: 10-20m.

Habitat: associated with *Amathea* sp. (Bryozoa).

***Stylopallene tubirostrum* CLARK 1963**

- 1963 *Siphopallene tubirostris* CLARK, Rec. Austr. Mus., **26** (1): 40-42, fig. 21.
 1968 *Siphopallene tubirostris*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 45-46, fig. 17a-b.
 1973 *Siphopallene tubirostrum*, - STOCK, Beaufortia, **21** (279): 92, 96 fig. 3d.
 1975 *Stylopallene tubirostris*, - CHILD, Smiths. Contr. Zool., **190**: 15.

Distribution: Australia.

Depth range: 24m.

Habitat: on bryozoa; sand.

COLOSSENDEIDAE

- 1948 Colossendeidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 268 [diagnosis, remarks, key to NW-Atlantic genera].
 1954 Colossendeidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 84 [diagnosis, key to South African genera].
 1969 Colossendeidae, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 23-24 [diagnosis, remarks, key to genera].
 1986 Colossendeidae, - STOCK, Bull. mar. Sci., **38** (3): 416.

***Colossendeis* JARZYNSKY 1870**

- 1927 *Colossendeis*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 310-312 [remarks, key to Antarctic species].
 1932 *Colossendeis*, - GORDON, Discovery Rep., **6**: 11-12 [remarks, key to species of longitarsus-group].
 1948 *Colossendeis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 268-269 [diagnosis, key to W-Atlantic species].
 1954 *Colossendeis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 84 [diagnosis, key to South African species].
 1969 *Colossendeis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 24-29 [diagnosis, general remarks, distribution], 28-29 [key to species occurring south of 40°S remarks on megalonyx-group].
 1986 *Colossendeis*, - STOCK, Bull. mar. Sci., **38** (3): 416-417 [remarks].

***Colossendeis angusta* SARS 1877**

- 1877 *Colossendeis angusta* SARS, Arch. Math. Naturvidensk., Kristiana, **2**: 268-269.
 1881 *Colossendeis angusta*, - WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 243-244, pl. 3, figs. 8, 13.
 1885 *Colossendeis angusta*, - SMITH (in VERRILL), Rep. U.S. Comm. Fish., 1883: 560.
 1891 *Colossendeis angusta*, - SARS; Norw. North-Atl. Exped., 1876-1878, **6** (Zool.

- 20): 140-143, pl. 15, fig. 2a-f.
- 1891 *Colossendeis angusta*, - TOPSENT, Bull. Soc. Zool. France, **16**: 178.
- 1899 *Colossendeis angusta*, - MEINERT, Danish Ingolf-Exped., **3** (1): 59.
- 1902 *Colossendeis angusta*, - MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 191.
- 1908 *Colossendeis angusta*, - LOMAN, Siboga Exped. Monogr., **40**: 22.
- 1908 *Colossendeis angusta*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 228.
- 1913 *Colossendeis angusta*, - OLSEN, Rep. Sci. Res. M. Sars North Atl. Deep Sea Exped., 1910, **3** (1) (Zool.): 3.
- 1917 *Colossendeis angusta*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Alber Ier Prince Souverain de Monaco, **51**: 8-9.
- 1930 *Colossendeis angusta*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 39.
- 1932 *Colossendeis angusta*, - GORDON, Discovery Rep., **6**: 12 [key].
- 1933 *Colossendeis angusta*, - STEPHENSEN, Medd. Grönland, **79** (6): 28-30, figs. 6-7.
- 1935 *Colossendeis angusta*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
- 1935 *Colossendeis angusta*, - STEPHENSEN, Bergens Mus. Arbok, 1935, Naturvidensk. Rekke, **7**: 33.
- 1937 *Colossendeis angusta*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 25-26.
- 1938 *Colossendeis angusta*, - CALMAN, Sci. Rep. John Murray Exped., 1933-34, **5** (6): 148-149.
- 1938 *Colossendeis angusta*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 8 [key].
- 1940 *Colossendeis angusta*, - MARCUS, Bol Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 110.
- 1943 *Colossendeis angusta*, - HILTON, J. Ent. Zool. Pomona College, **35** (1): 3.
- 1943 *Colossendeis angusta*, - NEEDLER, Can. Atl. Fauna, **10**: 5, fig. 2a-d.
- 1943 *Colossendeis angusta*, - STEPHENSEN, Medd. Grönland, **121** (8): 1.
- 1948 *Colossendeis angusta*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 269-271, fig. 50a.
- 1949 *Colossendeis angusta*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 298-299.
- 1956 *Colossendeis angusta*, - FAGE, Galathea Rep., **2**: 175-176, fig. 15.
- 1963 *Colossendeis angusta*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1340-1341.
- 1964 *Colossendeis angusta*, - LOSINA-LOSINSKY, Trudy Arktichesk. i Antarktichesk. Nauchno-Issledov. Inst., Leningrad, **259**: 336.
- 1969 *Colossendeis angusta*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 52-53, figs. 7-8.
- 1980 *Colossendeis angusta*, - DANIEL & SEN, J. mar. biol. Ass. India, **17** (2) ("1975"): 165.
- 1982 *Colossendeis angusta*, - CHILD, Smiths. Contr. Zool., **349**: 50.
- 1983 *Colossendeis angusta*, - BAMBER, Zool. J. Linn. Soc., **77**: 70.
- 1987 *Colossendeis angusta*, - STOCK, Cah. Biol. Mar., **28**: 507.
- 1991 *Colossendeis angusta*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 62.
- 1992 *Colossendeis angusta*, - CHILD, Smiths. Contr. Zool., **526**: 41.

Distribution: apparently cosmopolitan in deep cold waters.

Depth range: 77-5480m.

Habitat: foraminifera and pteropod mud.

- 1885 *Colossendeis arcuata* MILNE-EDWARDS, in FILHOL, La vie au fond des mers. Les explor. sous-marines et les voyages du Travailleur et du Talisman: 151.
- 1886 *Colossendeis Titan* PERRIER, in: Les Expl. Sous-Marines, **17**: 302.
- 1913 *Colossendeis michaelsarsi* OHLSEN, Rep. Sci. Res. M. Sars North Atl. Deep Sea Exped., 1910, **3** (1) (Zool.): 4-5, figs. 1-4, pl. 1, fig. A.
- 1937 *Colossendeis arcuata* BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 26-30, figs. 2-8.
- 1948 *Colossendeis michaelsarsi*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 274-275, fig. 50f.
- 1956 *Colossendeis michaelsarsi*, - FAGE, Galathea Rep., **2**: 177.
- 1978 *Colossendeis arcuata*, - STOCK, Cah. Biol. Mar., **19**: 403-405, fig. 1g-j.
- 1983 *Colossendeis arcuata*, - BAMBER, Zool. J. Linn. Soc., **77**: 71.
- 1984 *Colossendeis arcuata*, - STOCK, J. mar. biol. Ass. U.K., **64**: 745.
- 1986 *Colossendeis arcuata*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 417-418.
- 1987 *Colossendeis arcuata*, - STOCK, Cah. Biol. Mar., **28**: 508.
- 1992 *Colossendeis angusta*, - CHILD, Smiths. Contr. Zool., **526**: 41.

Distribution: N- and W-Atlantic deep Basin waters; Chile.

Depth range: 730-2177m.

Habitat: shells, foraminifera, pteropods; foraminifera and pteropod mud.

***Colossendeis australis* HODGSON 1907**

- 1907 *Colossendeis australis* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 59, pl. IX, fig. 1, pl. X, figs. 1-2.
- 1913 *Colossendeis australis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 63-65, figs. 20-21.
- 1915 *Colossendeis australis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 10 [key], 14-15.
- 1932 *Colossendeis australis*, - GORDON, Discovery Rep., **6**: 12, 15-16, figs. 2a, 3a, c, 4b.
- 1935 *Colossendeis australis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **3**: 293.
- 1938 *Colossendeis australis*, - GORDON. Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 8 [key].
- 1944 *Colossendeis australis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 10.
- 1969 *Colossendeis australis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 49-50, figs. 7, 8, 10, 65-71.

Distribution: Arctic and subantarctic area.

Depth range: 91-640m.

***Colossendeis avidus* PUSHKIN 1970**

- 1970 *Colossendeis avidus* PUSHKIN, Zool. Zh., **49** (10): 1488-1490, fig. 1.

Distribution: Antarctica.

Depth range: 270m.

***Colossendeis bicincta* SCHIMKEWITSCH 1893**

- 1893 *Colossendeis bicincta* SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 27-29, pl. I, figs. 1-3.
 1978 *Colossendeis bicincta*, - STOCK, Cah. Biol. Mar., **19**: 409-411, figs. 1a-f, 3.
 1984 *Colossendeis bicincta*, - STOCK, J. mar. biol. Ass. U.K.: 745-746.
 1991 *Colossendeis bicincta*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 163.

Distribution: New Caledonia (Valufa Ridge, Lau Basin), W-coast of Panamá, NE-Atlantic.

Depth range: 1920-3058m.

***Colossendeis brevitarsis* LOSINA-LOSINSKY 1958**

- 1961 *Colossendeis brevitarsis*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 109.

Distribution: NW-Pacific (U.S.S.R.).

***Colossendeis bruuni* FAGE 1956**

- 1956 *Colossendeis bruuni* FAGE, Galathea Rep., **2**: 176-177, figs. 16-17.

Distribution: Kermadec Trench.

Depth range: 4390m.

***Colossendeis californica* HEDGPETH 1939**

- 1939 *Colossendeis californica* HEDGPETH, Am. Midl. Nat., **22** (2): 459-460, pl. 1, figs. a-e.

Distribution: off California.

Depth range: 330m.

Habitat: green mud.

***Colossendeis clavata* MEINERT 1899**

- 1899 *Colossendeis clavata* MEINERT, Danish Ingolf-Exped., **3** (1): 57-58, pl. 5, figs. 19-20.
 1917 *Colossendeis clavata*, - BOUVIER, Res. Camp. Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 9-10.
 1937 *Colossendeis clavata*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), **20** (10) (1): 26, fig. 1.
 1948 *Colossendeis clavata*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 273-274, fig. 50e.
 1978 *Colossendeis clavata*, - STOCK, J. mar. biol. Ass. U.K.: 746.
 1978 *Colossendeis clavata*, - STOCK, Cah. Biol. Mar., **19**: 399.
 1990 *Colossendeis clavata*, - STOCK, Zool. Meded., **63** (16): 219.

Distribution: N-Atlantic between Iceland and Cape Ortegal (Spain); Canary Islands; NW-Atlantic (U.S.A.).

Depth range: 994-3100m.

Habitat: deep-sea clay.

***Colossendeis colossea* WILSON 1881**

- 1881 *Colossendeis colossea* WILSON, Bull. Mus. Comp. Zool. Harv., **8** (12): 244-246, pl. 1, fig. 1, pl. 3, figs. 5-7.
- 1881 *Colossendeis colossea*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 61-64, 147, pl. VIII, figs. 1-2, pl. X, figs. 1-5.
- 1881 *Colossendeis gigas* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 61-64, pl. VIII, figs. 1-2, pl. X, figs. 1-5, 147.
- 1885 *Colossendeis colossea*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 560, fig. 169.
- 1891 *Colossendeis colossea*, - TOPSENT, Bull. Soc. Zool. France, **16**: 177.
- 1893 *Colossendeis colossea*, - SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 29.
- 1899 *Colossendeis colossea*, - MEINERT, Danish Ingolf-Exped., **3** (1): 58.
- 1908 *Colossendeis colossea*, - LOMAN, Siboga Exped. Monogr., **40**: 21.
- 1908 *Colossendeis colossea*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 229.
- 1909 *Colossendeis colossea*, - COLE, Bull. Mus. Comp. Zool., Harvard, **52** (11): 187.
- 1911 *Colossendeis colossea*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 4.
- 1913 *Colossendeis colossea*, - OLSEN, Rep. Sci. Res. M. Sars North Atl. Deep Sea Exped., 1910, **3** (1) (Zool.): 4.
- 1913 *Colossendeis colossea*, - STEPHENSEN, Medd. Grönland, **52**: 409.
- 1917 *Colossendeis colossea*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 13-16, pl. I, fig. 2, pl. II, fig. 1.
- 1923 *Colossendeis colossea*, - BOUVIER, Faune de France, **7**: 25.
- 1923 *Colossendeis colossea*, - CALMAN, Rec. Indian Mus., **24** (3): 266.
- 1928 *Colossendeis colossea*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 7.
- 1935 *Colossendeis colossea*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
- 1936 *Colossendeis colossea*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 866-867.
- 1937 *Colossendeis colossea*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 31-32.
- 1938 *Colossendeis colossea*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 12.
- 1940 *Colossendeis colossea*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 109, 112 [key].
- 1943 *Colossendeis colossea*, - HEDGPETH, Proc. New England Zool. Club., **22**: 55-56.
- 1943 *Colossendeis colossea*, - NEEDLER, Can. Atl. Fauna, **10**: 4, fig. 1a-c.
- 1944 *Colossendeis colossea*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 11.
- 1947 *Colossendeis gigas*, - OHSHIMA & KISHIDA, in: Illustr. Encyclopaedia Fauna Japan: 1009, fig. 2863.
- 1948 *Colossendeis colossea*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 271-272, fig. 50b.
- 1949 *Colossendeis colossea*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 299.
- 1953 *Colossendeis colossea*, - STOCK, Temminckia, **9**: 311.
- 1954 *Colossendeis colossea*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 85.
- 1956 *Colossendeis colossea*, - PAGE, Galathea Rep., **2**: 178, fig. 21A.
- 1969 *Colossendeis colossea*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 53-54, fig. 8

- 1975 *Colossendeis colossea*, - STOCK, Bull. mar. Sci., **24** (4): 987.
 1983 *Colossendeis colossea*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **5**
 (sec. A) (1): 300.
 1986 *Colossendeis colossea*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 418.
 1987 *Colossendeis colossea*, - STOCK, Cah. Biol. Mar., **28**: 508.
 1990 *Colossendeis colossea*, - STOCK, Zool. Meded., **63** (16): 219.
 1991 *Colossendeis colossea*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 62.
 1991 *Colossendeis colossea*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 163.
 1991 *Colossendeis colossea*, - TURPAEVA, Zool. Zh., **70** (12): 41.
 1991 *Colossendeis colossea*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**,
 sec. A, nos. 1-2: 137-138.
 1992 *Colossendeis colossea*, - CHILD, Smiths. Contr. Zool., **526**: 41-42.

Distribution: almost world-wide in the oceanic deep-sea basins.

Depth range: 800-4140m.

Habitat: shells, foraminifera, pteropods.

***Colossendeis cucurbita* COLE 1909**

- 1881 *Colossendeis gigas-leptorhynchus* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 65-66.
 1909 *Colossendeis cucurbita* COLE, Bull. Mus. Comp. Zool. Harvard College, **52** (11): 188-191, pl. 2, figs. 3-4, pl. 3, figs. 8-12.
 1937 *Collosendeis gigas-leptorhynchus*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 32.
 1956 *Colossendeis cucurbita*, - FAGE, Galathea Rep., **2**: 178-179, fig. 21B.
 1971 *Colossendeis macerrima*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **92**: 289.
 1975 *Colossendeis macerrima leptorhynchus*, - TURPAEVA, Trudy Inst. Okeanol P.P. Shirshov, Akad. Nauk SSSR, **103**: 234.
 1978 *Colossendeis cucurbita*, - STOCK, Cah. Biol. Mar., **19**: 406.
 1985 *Colossendeis cucurbita*, - BAMBER, Zool. J. Linn. Soc., **83**: 305.

Distribution: N-Atlantic and -Pacific.

Depth range: 3294-4400m.

***Colossendeis curtirostris* STOCK 1963**

- 1963 *Colossendeis curtirostris* STOCK, Ann. S. Afr. Mus., **46** (12): 330-332, fig. 6b-h.

Distribution: South Africa.

Depth range: 2708m.

***Colossendeis drakei* CALMAN 1915**

- 1915 *Colossendeis drakei* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 11, 22-23, fig. 3.
 1932 *Colossendeis drakei*, - GORDON, Discovery Rep., **6**: 22-24, figs. 4c, 7e-f, 8b.
 1935 *Colossendeis drakei*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 294.

- 1938 *Colossendeis drakei*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 8 [key].
 1944 *Colossendeis drakei*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 10.
 1969 *Colossendeis drakei*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 39-40, figs. 7, 8, 10, 34-39.

Distribution: Antarctic.
Depth range: 3-549m.

Colossendeis enigmatica TURPAEVA 1974

- 1974 *Colossendeis enigmatica* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **98**: 301, fig. 12 (1-8).

Distribution: Antarctica.
Depth range: 315-335m.

Colossendeis gardineri CARPENTER 1908

- 1908 *Colossendeis gardineri* CARPENTER, Trans. Linn. Soc. London, **12** (2): 98-99, pl. 13, figs. 20-24.
 1923 *Colossendeis gardineri*, - CALMAN, Rec. Indian Mus., **25** (3): 267-268.
 1984 *Colossendeis gardineri*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **6** (sec. A) (3): 705, figs. 7-9.
 1986 *Colossendeis gardineri*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 417, fig. 5a-c.
 1991 *Colossendeis gardineri*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 164.

Distribution: Indian Ocean (Saya de Malha Bank); Caribbean (off St. Vincent).
Depth range: 576-824m.

Colossendeis geoffroyi MANE-GARZON 1944

- 1944 *Colossendeis geoffroyi* MANE-GARZON, Comun. Zool. Mus. Hist. Nat. Montevideo, **1** (15): 1-6, figs. 1-5.
 1966 *Colossendeis geoffroyi*, - STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 396.
 1992 *Colossendeis geoffroyi*, - STOCK, Tijdschr. Ent., **135**: 134-135, 139.

Distribution: SW-Atlantic.
Depth range: 65-130m.

Colossendeis gibbosa MÖBIUS 1902

- 1902 *Colossendeis gibbosa* MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 192-193, pl. XXX, figs. 1-5.

Distribution: Bouvet-Island.
Depth range: 439-567m.

Colossendeis gracilis HOEK 1881

- 1881 *Colossendeis gracilis* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 69-70, pl. IX, figs. 6-8, pl. X, figs. 6-7.
 1963 *Colossendeis gracilis*, - STOCK, Ann. S. Afr. Mus., **46** (12): 330.
 1968 *Colossendeis gracilis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 17.
 1975 *Colossendeis gracilis*, - STOCK, Bull. mar. Sci., **24** (4): 990.
 1990 *Colossendeis gracilis*, - STOCK, Zool. Meded., **63** (16): 220.
 1990 *Colossendeis gracilis*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **126**: 114-115, fig. 4.

Distribution: worldwide in tropical areas and in the southern hemisphere.

Depth range: 1500-3200m.

Colossendeis hoeki GORDON 1944

- 1944 *Colossendeis hoeki* GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B), **5** (1): 8 [key], 11-13, fig. 2a-b.
 1969 *Colossendeis hoeki*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 51, figs. 7-8.

Distribution: Antarctica.

Depth range: 3112m.

Colossendeis korotkevitschi PUSHKIN 1984

- 1984 *Colossendeis korotkevitschi* PUSHKIN, Zool. Zh., **63** (8): 1259-1261, figs. 1-11.

Distribution: off Kerguelen Islands and Crozet Islands, in Weddell Sea and near D'Urville Island.

Depth range: 132-660m.

Colossendeis leptorhynchus HOEK 1881

- 1881 *Colossendeis leptorhynchus* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 64-65, pl. VIII, figs. 3-7.
 1970 *Colossendeis pennata* PUSHKIN, Zool. Zh., **49** (10): 1490-1492, fig. 2.
 1978 *Colossendeis leptorhynchus*, - STOCK, Cah. Biol. Mar., **19**: 402.
 1981 *Colossendeis leptorhynchus*, - STOCK, Cah. Biol. Mar., **22**: 454-455
 1983 *Colossendeis leptorhynchus*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **5** (sec. A) (1): 299-300.
 1986 *Colossendeis leptorhynchus*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 417.
 1990 *Colossendeis leptorhynchus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **103** (2): 308.
 1992 *Colossendeis leptorhynchus*, - CHILD, Smiths. Contr. Zool., **526**: 42.

Distribution: predominantly in the Southern Hemisphere;

probably cosmopolitan.

Depth range: 531-3675m.

***Colossendeis lilliei* CALMAN 1915**

- 1915 *Colossendeis lilliei* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 25-27, fig. 4.
 1935 *Colossendeis lilliei*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 294.
 1838 *Colossendeis lilliei*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 11 [key].
 1944 *Colossendeis lilliei*, - Rep. Brit., Austr. and N.Z. Antarct. Exped., (B), **5** (1): 11.
 1969 *Colossendeis lilliei*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 43-45, figs. 7, 8, 10, 47-52.

Distribution: Antarctica.

Depth range: 110-1119m.

***Colossendeis longirostris* GORDON 1938**

- 1938 *Colossendeis longirostris* GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 8 [key], 9-10, fig. 1.
 1959 *Colossendeis longirostris*, - UTINOMI, Special Publ. Seto Mar. Biol. Lab. (= Contr. Seto Mar. Biol. Lab., **341**): 10-12, fig. 6.
 1969 *Colossendeis longirostris*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 42-43, figs. 7, 8, 45, 46.

Distribution: Antarctica.

Depth range: 2.5-2400m.

***Colossendeis macerrima* WILSON 1881**

- 1881 *Colossendeis macerrima* WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 246-247, pl. 1, fig. 2, pl. 4, figs. 9-12, pl. 5, fig. 32.
 1881 *Colossendeis macerrima*, - MILNE-EDWARDS, C.R. Hebd. Séanc. Acad. Sci., Paris, **93**: 933.
 1885 *Colossendeis macerrima*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 560, fig. 170.
 1893 *Colossendeis macerrima*, - SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 30.
 1896 *Colossendeis leptorhynchus* var. *septentrionalis*, - CAULLERY, Ann. Univ. Lyon, **26**: 362.
 1898 *Colossendeis japonica* HOEK, Tijdschr. Nederl. Dierk. Ver., ser. 2, vol. **1**: 295-296, pl. III, figs. 11-13.
 1908 *Colossendeis macerrima*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 229.
 1915 *Colossendeis macerrima*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 11.
 1917 *Colossendeis macerrima*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 10-13, pl. I, fig. 1, pl. III, figs. 1-2.
 1923 *Colossendeis macerrima*, - BOUVIER, Faune de France, **7**: 25.

- 1923 *Colossendeis macerrima*, - CALMAN, Rec. Indian Mus., **25** (3): 267-268.
- 1928 *Colossendeis macerrima*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 7.
- 1935 *Colossendeis macerrima*, - Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
- 1937 *Colossendeis gigas-leptorhynchus*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) **20** (1): 32.
- 1938 *Colossendeis macerrima*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 12.
- 1942 *Colossendeis macerrima*, - FAGE, Arch. Zool. Gen., **82** (Notes & Rev., 2): 75-76.
- 1948 *Colossendeis macerrima*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 273, fig. 50d.
- 1949 *Colossendeis japonica*, - HEDGEPETH, Proc. U.S. Natn. Mus., **98** (3231): 299-300, fig. 46a-d.
- 1949 *Colossendeis macerrima*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 299.
- 1953 *Colossendeis macerrima*, - STOCK, Temmickia, **9**: 307, fig. 17e-h.
- 1954 *Colossendeis macerrima*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 85.
- 1956 *Colossendeis japonica*, - FAGE, Galathea Rep., **2**: 176.
- 1956 *Colossendeis macerrima*, - FAGE, Galathea Rep., **2**: 180.
- 1963 *Colossendeis macerrima*, - CLARK, Rec. Austr. Mus., **26** (1): 75-76.
- 1963 *Colossendeis macerrima*, - STOCK, Ann. S. Afr. Mus., **46** (12): 326.
- 1969 *Colossendeis macerrima*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 53, figs. 7-8.
- 1975 *Colossendeis macerrima*, - STOCK, Bull. mar. Sci., **24** (4): 985-987, fig. 11a-b.
- 1978 *Colossendeis macerrima*, - STOCK, Cah. Biol. Mar., **19**: 400-402, fig. 2m.
- 1980 *Colossendeis macerrima*, - DANIEL & SEN, J. mar. biol. Ass. India, **17** (2) ("1975"): 165.
- 1981 *Colossendeis macerrima*, - STOCK, Cah. Biol. Mar., **22**: 455.
- 1982 *Colossendeis macerrima*, - CHILD, Smiths. Contr. Zool., **349**: 51.
- 1983 *Colossendeis macerrima*, - BAMBER, - Zool. J. Linn. Soc., **77**: 71.
- 1984 *Colossendeis macerrima*, - STOCK, J. mar. biol. Ass. U.K., **64**: 746.
- 1986 *Colossendeis macerrima*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 417, fig. 5e.
- 1987 *Colossendeis macerrima*, - STOCK, Cah. Biol. Mar., **28**: 508.
- 1988 *Colossendeis macerrima*, - MUNILLA, Monogr. Zool. Mar., **3**: 185.
- 1990 *Colossendeis macerrima*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **103** (2): 309.
- 1990 *Colossendeis macerrima*, - STOCK, Zool. Meded., **63** (16): 220.
- 1991 *Colossendeis macerrima*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 62.
- 1991 *Colossendeis macerrima*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 163, 164.
- 1991 *Colossendeis macerrima*, - TURPAEVA, Zool. Zh., **70** (12): 41-42.
- 1991 *Colossendeis macerrima*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 138.
- 1992 *Colossendeis macerrima*, - CHILD, Smiths. Contr. Zool., **526**: 42.

Distribution: cosmopolitan.

Depth range: 121-4000m.

Habitat: mud, clay, shells, foraminifera, pteropods; foraminifera and pteropod mud.

Additional synonymy: *Colossendeis spei*.

- 1881 *Colossendeis media* HOEK, Challenger Rep. (Zool., 3): 71, pl. X, figs. 10-11.
 1881 *Colossendeis media* HOEK, Challenger Rep. (Zool., 3): 72-73, pl. X, figs. 8-9
 1974 *Colossendeis media*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, 98: 297, fig. 11.

Distribution: Antarctica.

Depth range: 4072-4850m.

Habitat: mud.

Colossendeis megalonyx HOEK 1881

- 1881 *Colossendeis megalonyx* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, 3 (10): 67-69, pl. IX, figs. 1-3.
 1902 *Colossendeis frigida* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 63.
 1907 *Colossendeis rugosa* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 64, pl. IX, fig. 3, pl. X, figs. 5-6.
 1908 *Colossendeis orcadense*, - HODGSON, Trans. Roy. Soc. Edinburgh, 46 (1) (6): 184.
 1915 *Colossendeis megalonyx*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 15-16.
 1915 *Colossendeis rugosa*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 16-17.
 1915 *Colossendeis frigida*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool. 3 (1): 17-18.
 1923 *Colossendeis frigida*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., 1 (2): 7.
 1932 *Colossendeis rugosa*, - GORDON, Discovery Rep., 6: 12, 17.
 1932 *Colossendeis frigida*, - GORDON, Discovery Rep., 6: 16-18, figs. 5a-b, 6a, d, 7c-d.
 1932 *Colossendeis scoresbii* GORDON, Discovery Rep., 6: 18-21, figs. 5c, 6b-c, 7a-b.
 1935 *Colossendeis frigida*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 293.
 1935 *Colossendeis megalonyx*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 293.
 1935 *Colossendeis rugosa*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 293.
 1935 *Colossendeis orcadense*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 294.
 1944 *Colossendeis frigida*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 10.
 1956 *Colossendeis megalonyx*, - FAGE, Galathea Rep., 2: 177-178, figs. 18-20.
 1963 *Colossendeis orcadensis*, - STOCK, Ann. S. Afr. Mus., 46 (12): 334.
 1969 *Colossendeis megalonyx*, - ARNAUD, Les Terres Antarct. Françaises, 47: 36.
 1969 *Colossendeis megalonyx*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 30-32, figs. 7-8, 11-16, 23.
 1969 *Colossendeis megalonyx megalonyx*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 30.
 1969 *Colossendeis megalonyx scoresbii*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 32.
 1969 *Colossendeis megalonyx arundorostris*, - FRY & HEDGPETH, - N.Z. Oceanogr. Inst. Mem., 49: 33.
 1969 *Colossendeis megalonyx orcadense*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 34.

- 1969 *Colossendeis megalonyx megalonyx*, - HEDGPETH, Antarct. Map Folio Ser. Am. Geogr. Soc., Folio **11**: 27.
- 1972 *Colossendeis megalonyx megalonyx*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 805-806.
- 1972 *Colossendeis megalonyx*, - ARNAUD, Téthys, suppl. **4**: 145.
- 1980 *Colossendeis m. megalonyx*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 7.
- 1991 *Colossendeis megalonyx orcadense*, - TURPAEVA, Zool. Zh., **70** (12): 42.

Distribution: Antarctic, subantarctic area; extending to the southern coast of South Africa and America.

Depth range: 128-3410m.

***Colossendeis melancholicus* STOCK 1975**

- 1975 *Colossendeis melancholicus* STOCK, Bull. mar. Sci., **24** (4): 988-990, figs. 11c-f, 12.
- 1986 *Colossendeis melancholicus*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.

Distribution: Grand Bahama.

Depth range: 779m.

***Colossendeis mica* PUSHKIN 1970**

- 1970 *Colossendeis mica* PUSHKIN, Zool. Zh., **49** (10): 1492-1494, fig. 3.
- 1991 *Colossendeis mica*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 164.

Distribution: Antarctica.

Depth range: 1400m.

***Colossendeis minor* SCHIMKEWITSCH 1893**

- 1893 *Colossendeis macerrima minor* SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 30-32, pl. I, figs. 7-10, pl. II, figs. 14-15
- 1917 *Colossendeis macerrima*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 10-13
- 1971 *Colossendeis macerrima minor*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **92**: 289-290, pl. 7, 6-7.
- 1975 *Colossendeis macerrima*, - STOCK, Bull. mar. Sci., **24** (4): 985-987 (Pillsbury stat. 526 only).
- 1984 *Colossendeis minor*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **6** (sec. A) (3): 702-704, figs. 1-6.
- 1991 *Colossendeis minor*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 163-164.
- 1991 *Colossendeis minor*, - TURPAEVA, Zool. Zh., **70** (12): 42.

Distribution: eastern tropical Pacific off Central and South America; Mid Indian Basin; south-eastern Madagascar.

Depth range: 1035-3344m.

***Colossendeis minuta* HOEK 1881**

- 1881 *Colossendeis minuta* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 73-74, pl. X, figs. 12-14.
- 1893 *Colossendeis minuta*, - SCHIMKEWITSCH. Bull. Mus. Comp. Zool., Harvard, **25** (2): 34.
- 1898 *Colossendeis japonica* HOEK, Tijdschr. Nederl. Dierk. Ver., (2) **5** (2-4): 295.
- 1944 *Colossendeis minuta*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 13.
- 1948 *Colossendeis minuta*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 272, fig. 50c.
- 1963 *Colossendeis minuta*, - STOCK, Ann. S. Afr. Mus., **46** (12): 326-328, fig. 4.
- 1975 *Colossendeis minuta*, - STOCK, Bull. mar. Sci., **24** (4): 987.
- 1975 *Colossendeis minuta*, - TUIRPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **103**: 241.
- 1990 *Colossendeis minuta*, - STOCK, Zool. Meded., **63** (16): 221.

Distribution: Atlantic.

Depth range: 2000-3200m.

***Colossendeis nasuta* HEDGPETH 1949**

- 1949 *Colossendeis nasuta* HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 302-303, fig. 46e-h.
- 1955 *Colossendeis nasuta*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 35-36, fig. 21.

Distribution: Japan.

Depth range: 519-716m.

***Colossendeis oculifera* STOCK 1963**

- 1963 *Colossendeis oculifera* STOCK, Ann. S. Afr. Mus., **46** (12): 328-330, fig. 5.

Distribution: South Africa.

***Colossendeis pipetta* STOCK 1991**

- 1991 *Colossendeis pipetta* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 164-166, fig. 28.

Distribution: New Caledonia, Chesterfield Islands, Loyalty Islands.

Depth range: 270-1475m.

***Colossendeis proboscidea* (SABINE 1824)**

- 1824 *Phoxichilus proboscideus* SABINE, Suppl. to Append. of Capt. Parry's Voyage in the years 1819-20: p. CCXXVI.
- 1881 *Anomorhynchus Smithii* MIERS, Ann. Mag. Nat. Hist., **5** (7): 50-51, pl. VII, figs. 6-8.
- 1899 *Colossendeis proboscidea*, - MEINERT, Danish Ingolf-Exped., **3** (1): 57.
- 1933 *Colossendeis proboscidea*, - STEPHENSEN, Medd. Gronl., **79** (6): 28, fig. 6.

- 1936 *Colossendeis proboscidea*, - STEPHENSEN, Bergens Mus. Arbok Naturv. Rek., **2**: 32-33.
 1943 *Colossendeis proboscidea*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 88-89.
 1963 *Colossendeis proboscidea*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1340.

Distribution: high Arctic area, possibly circumpolar.
Depth range: shallow water to about 915m.

***Colossendeis robusta* HOEK 1881**

- 1881 *Colossendeis robusta* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 66-67, pl. IX, figs. 4-5.
 1902 *Colossendeis robusta*, - MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 190, pl. XXIX, figs. 1-5.
 1907 *Colossendeis glacialis* HODGSON, Rep. Natn. Antarct. Exped., 1901-1904, Nat. Hist., **3**: 61, pl. IX, fig. 2, pl. X, figs. 3-4.
 1911 *Colossendeis gracilipes* BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 1137.
 1913 *Colossendeis gracilipes*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 58-63, figs. 12-19.
 1913 *Colossendeis robusta*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 54-58, figs. 8-11.
 1915 *Colossendeis glacialis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 20-22.
 1915 *Colossendeis robusta*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 24.
 1932 *Colossendeis glacialis*, - GORDON, Discovery Rep., **6**: 21-22, figs. 4d, 7g, 8a.
 1932 *Colossendeis robusta*, - GORDON, Discovery Rep., **6**: 12 [key].
 1935 *Colossendeis glacialis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
 1935 *Colossendeis gracilipes*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
 1935 *Colossendeis robusta*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 293.
 1938 *Colossendeis glacialis*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 8 [key].
 1938 *Colossendeis robusta*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 8 [key].
 1944 *Colossendeis glacialis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 10.
 1969 *Colossendeis robusta*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 35-38, figs. 7, 8, 10, 24-31.

Distribution: Antarctic and subantarctic area.
Depth range: 0.3-3610m.

***Colossendeis scotti* CALMAN 1915**

- 1915 *Colossendeis scotti* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 10 [key], 11-13, fig. 1.
 1932 *Colossendeis scotti*, - GORDON, Discovery Rep., **6**: 11 [key], 12.
 1935 *Colossendeis scotti*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5**

- (4) (2): 294.
- 1938 *Colossendeis scotti*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 8 [key], 12.
- 1969 *Colossendeis scotti*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 45-47, figs. 7, 8, 10, 53-58.
- 1974 *Colossendeis scotti*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **98**: 299, fig. 12 (9-11).

Distribution: Antarctica.

Depth range: 36-250m.

***Colossendeis smirnovi* PUSHKIN 1988**

- 1988 *Colossendeis smirnovi* PUSHKIN, Zool. Zh., **67** (6): 953-955, figs. 1-8.

Distribution: southern Oceans.

Depth range: 750-1070m.

***Colossendeis stramenti* FRY & HEDGPETH 1969**

- 1969 *Colossendeis stramenti* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 38-40, figs. 32-33.

Distribution: Magellanic region.

Depth range: 645-845m.

***Colossendeis subminuta* SCHIMKEWITSCH 1893**

- 1893 *Colossendeis subminuta* SCHIMKEWITSCH, Bull. Mus. Comp. Zool., **25** (2): 33-36, pl. II, figs. 26-27.

Distribution: W-coast of America, south of Cocos-Island.

Depth range: 2150m.

***Colossendeis tethya* TURPAEVA 1974**

- 1974 *Colossendeis tethya* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **98**: 302-303, fig. 13.

Distribution: Antarctica.

***Colossendeis tortipalpis* GORDON 1932**

- 1932 *Colossendeis tortipalpis* GORDON, Discovery Rep., **6**: 12-15, figs. 2b-e, 3b, d, 4a.
- 1935 *Colossendeis tortipalpis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 294.
- 1944 *Colossendeis tortipalpis*, - GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B), **5** (1): 9-10.
- 1969 *Colossendeis tortipalpis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**:

41-43, figs. 7, 8, 10, 40-45.

Distribution: Antarctica.

Depth range: 219-1400m.

***Colossendeis wilsoni* CALMAN 1915**

- 1915 *Colossendeis wilsoni* CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 11 [key], 18-20, fig. 2.
 1932 *Colossendeis wilsoni*, - GORDON, Discovery Rep., **6**: 12 [key].
 1935 *Colossendeis wilsoni*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 294.
 1938 *Colossendeis wilsoni*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), Zool. & Bot.), **2** (8): 8 [key], 12.
 1969 *Colossendeis wilsoni*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 47-49, figs. 7, 8, 10, 59-64.

Distribution: Antarctica.

Depth range: 110-314m.

***Decolopoda* EIGHTS 1835**

- 1969 *Decolopoda*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 54 [diagnosis].

***Decolopoda australis* EIGHTS 1835**

- 1835 *Decolopoda australis* EIGHTS, Boston J. Nat. Hist., **1** (2): 203-206, pl. VII.
 1902 *Decolopoda australis*, - STEEBING, Knowledge, **25**: 185-189.
 1905 *Decolopoda antarctica* BOUVIER, Bull. Mus. natn. Hist. nat., Paris, **11**: 295.
 1905 *Decolopoda australis*, - COLE, Ann. Mag. Nat. Hist., **15** (7): 405.
 1905 *Decolopoda australis*, - BOUVIER, Proc. Roy. Phys. Soc. Edinburgh, **16** (1): 36-42, pl. III.
 1905 *Decolopoda australis*, - HODGSON, Zool. Anz., **29** (8): 254-256.
 1905 *Decolopoda australis*, - LOMAN, Zool. Anz., **28**: 722-723.
 1906 *Decolopoda australis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., **142**: 16-18.
 1906 *Decolopoda australis*, - BOUVIER, L'Expéd. Antarct. Française, 1903-1905: 21-29, pl. II, figs. 6-9.
 1908 *Decolopoda australis*, - HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1): 181-184, pl. III.
 1909 *Decolopoda australis*, - CALMAN, Sci. Progr., London, 1909: 687-693.
 1909 *Decolopoda australis*, - d'ARCY THOMPSON, The Cambr. Nat. Hist., **4**: 531.
 1910 *Decolopoda australis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **151**: 27.
 1911 *Decolopoda australis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 491.
 1911 *Decolopoda australis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 1136.
 1913 *Decolopoda antarctica*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910): 48.
 1913 *Decolopoda australis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910): 48-52, figs. 1-7.

- 1920 *Decolopoda antarctica*, - CALMAN, Ann. Mag. Nat. Hist., **6** (9): 244.
 1932 *Decolopoda antarctica*, - GORDON, Discovery Rep., **6**: 9-11, figs. I, 73B, 74B.
 1932 *Decolopoda australis*, - GORDON, Discovery Rep., **6**: 8-9, fig. 73A.
 1935 *Decolopoda antarctica*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 280.
 1935 *Decolopoda australis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 280.
 1944 *Decolopoda* sp.?, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 8.
 1969 *Decolopoda australis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 54-56, figs. 7, 8, 10, 75, 76, 78-82.
 1969 *Decolopoda australis*, - ARNAUD, Les Terres Austr. Antarct. Françaises, **47**: 36.
 1969 *Decolopoda australis*, - HEDGPETH, Antarct. Map Folio Ser. Am. Geogr. Soc., Folio **11**: 27.
 1972 *Decolopoda australis*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 804-805.
 1982 *Decolopoda australis*, - PUSHKIN, Zool. Zh., **61** (11): 1657-1659, figs. A-D.
 1982 *Decolopoda antarctica*, - PUSHKIN, Zool. Zh., **61** (11): 1659-1661, figs. E-J.

Distribution: Antarctic and subantarctic area.

Depth range: 0-1119m.

***Dodecolopoda* CALMAN & GORDON 1933**

- 1969 *Dodecolopoda* , - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 56 [diagnosis].

***Dodecolopoda mawsoni* CALMAN & GORDON 1933**

- 1933 *Dodecolopoda mawsoni* CALMAN & GORDON, Proc. Roy. Soc. London, (B) **113**: 107-115, fig. 1.
 1935 *Dodecolopoda mawsoni*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 280.
 1944 *Dodecolopoda mawsoni*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 7-9, fig. 1a-c.
 1947 *Dodecolopoda mawsoni*, - HEDGPETH, Smiths. Misc. Coll., **106** (18): 17, figs. 8-10.
 1969 *Dodecolopoda mawsoni*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 56-57, figs. 7-10, 77-82.
 1974 *Dodecolopoda mawsoni*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **98**: 294-295, fig. 10.

Distribution: Antarctica.

Depth range: 534-549m.

***Hedgpethia* TURPAEVA 1973**

- 1973 *Hedgpethia* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **91**: 183 [diagnosis, remarks], 184 [key to species].

Hedgpethia articulata (LOMAN 1908)

- 1908 *Colossendeis articulata* LOMAN, Siboga Exped., **XL**: 22-23, pl. VI, figs. 66-73.
 1973 *Hedgpethia articulata* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **91**: 185-189, fig. 2.
 1991 *Hedgpethia articulata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166 [text].

Distribution: Indo-West Pacific (type locality: Gulf of Boni).

Depth range: 1944-3610m.

Habitat: mud.

Hedgpethia atlantica (STOCK 1970)

- 1970 *Rhopalorhynchus atlanticum* STOCK, Meteor-Forsch.-Ergebn., (D), **5**: 7-9, figs. 8-13.
 1979 *Rhopalorhynchus atlanticum*, - RACK, Mitt. Hamburg. Zool. Mus. Inst., **67**: 113.
 1987 *Rhopalorhynchus atlanticum*, - STOCK, Cah. Biol. Mar., **28**: 508-509, fig. 12.
 1991 *Hedgpethia atlantica*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166 [text].
 1991 *Hedgpethia atlantica*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 138.

Distribution: NE-Atlantic.

Depth range: 135-1125m.

Habitat: shelly sand

Hedgpethia bicornis (TURPAEVA 1953)

- 1958 *Colossendeis bicornis*, - LOSINA-LOSINSKY, Biull. Mosov. Obsh. Ispyt. Prirod., **63** (1): 27-29, pl. 3, figs. 1-7.
 1961 *Colossendeis bicornis*, - LOSINA-LOSINSKY, Issledov. Dalnevost. Morei SSSR, Leningrad, **7**: 109-110.
 1991 *Hedgpethia bicornis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166 [text].

Distribution: NW-Pacific (U.S.S.R.).

Depth range: 64-515m.

Hedgpethia brevitarsis (LOSINA-LOSINSKY & TURPAEVA 1958)

- 1958 *Colossendeis brevitarsis* LOSINA-LOSINSKY & TURPAEVA, Bull. Moskovskovo Obschest. Ispytatelei Prirody, **63** (1): 26-27, fig. 2.
 1970 *Rhopalorhynchus brevitarsis*, - STOCK, Meteor Forsch.-Ergebn., (D), **5**: 9 [key].
 1983 *Hedgpethia brevitarsis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 63.
 1991 *Hedgpethia brevitarsis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 62.

Distribution: Japan (Sagami Bay) to Gulf of Alaska.

Depth range: 290-563m.

Hedgpethia californica (HEDGPETH 1939)

- 1939 *Colossendeis californica* HEDGPETH, Am. Midl. Nat., **22** (2): 459-460, pl. 1, figs. a-e
 1973 *Hedgpethia californica* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **91**: 186-189.
 1991 *Hedgpethia californica*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166 [text].

Distribution: California.

Depth range: 330m.

Habitat: green mud.

Hedgpethia chitinosa (HILTON 1943)

- 1943 *Colossendeis chitinosa* HILTON, J. Ent. Zool. Pomona College, **35** (1): 4.
 1949 *Colossendeis chitinosa*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 301, fig. 47e-h.
 1954 *Colossendeis chitinosa*, - STOCK, Viensk. Medd. dansk naturhist. Foren., **116**: 161.
 1955 *Colossendeis chitinosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 34-35, fig. 20.
 1958 *Colossendeis chitinosa*, - LOSINA-LOSINSKY, Biull. Moskov. Obsh. Ispyt. Prirod., **63** (1): 23-26, pl. 1, figs. 1-6.
 1961 *Colossendeis chitinosa*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 109.
 1962 *Colossendeis chitinosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 103.
 1970 *Rhopalorhynchus chitinosum*, - STOCK, Meteor Forsch-Ergebn., (D), **5**: 9 [key].
 1971 *Colossendeis chitinosa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 337.
 1973 *Hedgpethia californica chitinosa*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **91**: 186, tab. 3.
 1983 *Hedgpethia chitinosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 63.
 1987 *Hedgpethia chitinosa*, - NAKAMURA, Biol. Lab., Imperial Household: 36.
 1991 *Hedgpethia chitinosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 63.

Distribution: Common in the NW-Pacific (Aleutian Islands to west of Kyushu); Korea Straits.

Depth range: 20-950m

Habitat: sand, shells.

Hedgpethia dampieri (CHILD 1975)

- 1975 *Rhopalorhynchus dampieri* CHILD, Smiths. Contr. Zool., **190**: 8-10, fig. 4.

Distribution: Australia.

Depth range: 113-122m.

Habitat: "beam trawl with bryozoans" (CHILD).

Hedgpethia dofleini (LOMAN 1911)

- 1911 *Colossendeis dofleini* LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 4-5, pl. I, figs. 6-13.
 1919 *Colossendeis dofleini*, - FUKUI, Rigakukai[?], **16** (2): 98.
 1936 *Colossendeis dofleini*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 867.
 1949 *Colossendeis dofleini*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 300, fig. 47a-d.
 1951 *Colossendeis dofleini*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 167.
 1955 *Colossendeis dofleini*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 33.
 1958 *Rhopalorhynchus dofleini*, - STOCK, Tijdschr. Ent., **101** (2): 114.
 1971 *Colossendeis dofleini*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 337.
 1991 *Hedgpethia dofleini*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 63.

Distribution: from the Bering Sea to south of Japan (Kyushu).

Depth range: 20-923m.

Hedgpethia magnirostris ARNAUD & CHILD 1988

- 1988 *Hedgpethia magnirostris* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 153-155, fig. 7.
 1991 *Hedgpethia magnirostris*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166.

Distribution: South Africa.

Depth range: 680m.

Hedgpethia tibialis STOCK 1991

- 1991 *Hedgpethia tibialis* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 166-168, figs. 29-30.

Distribution: New Caledonia.

Depth range: 520-550m.

Pentacolossendeis HEDGPETH 1943

- 1948 *Pentacolossendeis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 275 [diagnosis, remarks].

Pentacolossendeis reticulata HEDGPETH 1943

- 1943 *Pentacolossendeis reticulata* HEDGPETH, Proc. New England Zool. Club, **22**: 56-57.
 1948 *Pentacolossendeis reticulata*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 275-278, fig. 51a-e.
 1986 *Pentacolossendeis reticulata*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 416.

Distribution: Bahamas and Florida, south to Guyana.

Depth range: 135-1298m.

Rhopalorhynchus WOOD-MASON 1873

- 1873 *Rhopalorhynchus* WOOD-MASON, J. Asiat. Soc. Bengal, **42** (2): 172 [diagnosis].
 1954 *Rhopalorhynchus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 88 [diagnosis].
 1970 *Rhopalorhynchus*, - STOCK, Meteor-Forsch.-Ergebn., (D), **5**: 9 [key to species of *articulatum*-group].

Rhopalorhynchus claudus STOCK 1975

- 1975 *Rhopalorhynchus claudus* STOCK, Bull. mar. Sci., **24** (4): 991-993, fig. 13.
 1986 *Rhopalorhynchus claudus*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.

Distribution: Barbados.

Depth range: 100m.

Habitat: muddy sand, shell debris; sponge bottom.

Rhopalorhynchus clavipes CARPENTER 1893

- 1893 *Rhopalorhynchus clavipes* CARPENTER, Sci. Proc. Roy. Dublin Soc., **8** (1) (4): 24-26, pl. II, figs. 1-10.
 1923 *Rhopalorhynchus kroeyeri* CALMAN, Rec. Indian Mus., **25** (3): 268 [part].
 1958 *Rhopalorhynchus clavipes*, - STOCK, Tijdschr. Ent., **101** (2): 127.
 1968 *Rhopalorhynchus clavipes*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 17.
 1970 *Rhopalorhynchus clavipes*, - CLARK, Trans. Roy. Soc. N.Z., **12** (3): 17.

Distribution: NE-Australia.

Depth range: 8-86m.

Habitat: found among corals.

Rhopalorhynchus filipes STOCK 1991

- 1991 *Rhopalorhynchus filipes* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 161-163, fig. 27.

Distribution: Chesterfield Islands; between Loyality Islands and New Caledonia.

Depth range: 200-320m.

Rhopalorhynchus gracillimus CARPENTER 1908

- 1908 *Rhopalorhynchus gracillimus* CARPENTER, Trans. Linn. Soc. London, **12** (2): 99-100, pl. 13, figs. 25-32 [except Saya de Malha record].
 1908 *Rhopalorhynchus kröyeri*, - LOMAN, Siboga Exped. Monogr., **40**: 26-27 [only Siboga sta. 310 specimen].
 1923 *Rhopalorhynchus kröyeri*, - CALMAN, Rec. Indian Mus., **25** (3): 268-270 [part].
 1954 *Rhopalorhynchus kröyeri*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 88-89, fig. 2.
 1958 *Rhopalorhynchus gracillimum*, - STOCK, Tijdschr. Ent., **101** (2): 128-132,

figs. 39-56.

- 1988 *Rhopalorhynchus gracillimus*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 156.

Distribution: East Indies and Maldives Islands to the east-coast of South Africa.

Depth range: 0-156m.

***Rhopalorhynchus kroeyeri* WOOD-MASON 1873**

- 1873 *Rhopalorhynchus kröyeri* WOOD-MASON, J. Asiatic Soc. Bengal, **42** (2): 172-175, pl. XIII, figs. 1-5.
 1954 *Rhopalorhynchus kröyeri*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 88-89, fig. 2.
 1954 *Rhopalorhynchus kröyeri*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 161 [only Kei-Island record].
 1957 *Rhopalorhynchus kroeyeri*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 97.
 1958 *Rhopalorhynchus kroeyeri*, - STOCK, Tijdschr. Ent., **101** (2): 118.

Distribution: East Indian Archipelago (Andaman Islands; Philippines, Kei Islands).

Depth range: 35-46m.

***Rhopalorhynchus lomani* STOCK 1958**

- 1958 *Rhopalorhynchus lomani* STOCK, Tijdschr. Ent., **101** (2): 119-124, figs. 7-21.
 1965 *Rhopalorhynchus lomani*, - STOCK, Beaufortia, **13** (151): 16.
 1973 *Rhopalorhynchus lomani*, - ARNAUD, Tethys, **4** (4): 954.

Distribution: Tropical W-Indian Ocean.

Depth range: 4-86m.

Habitat: Halophila; hydroids and Zoantharia.

***Rhopalorhynchus mortenseni* STOCK 1958**

- 1954 *Rhopalorhynchus kröyeri*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 161-162 [only Jolo record].
 1958 *Rhopalorhynchus mortenseni* STOCK, Tijdschr. Ent., **101** (2): 132-133, figs. 57-61.
 1968 *Rhopalorhynchus mortenseni*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 18.

Distribution: Philippines and Australia (Coral Sea).

Depth range: 46-55m.

Habitat: gravel; sand and corals.

***Rhopalorhynchus pedunculatum* STOCK 1957**

- 1957 *Rhopalorhynchus pedunculatum* STOCK, Bull. Sea Fish. Res. Stat. Haifa, **13**: 13-14, fig. 1.
 1958 *Rhopalorhynchus pedunculatum*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**:

3.

- 1958 *Rhopalorhynchus pedunculatum*, - STOCK, Tijdschr. Ent., **101** (2): 125.
 1970 *Rhopalorhynchus pedunculatum*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **2** (1): 1.

Distribution: probably endemic in the Red Sea area.
Depth range: littoral-6m.

***Rhopalorhynchus sibogae* STOCK 1958**

- 1908 *Rhopalorhynchus kröyeri*, - LOMAN, Siboga Exped. Monogr., **40**: 217 [part], pl. XV, fig. 217.
 1958 *Rhopalorhynchus sibogae* STOCK, Tijdschr. Ent., **101** (2): 124-125, figs. 22-27.
 1975 *Rhopalorhynchus sibogae*, - CHILD, Smiths. Contr. Zool., **190**: 7-8.

Distribution: Indonesia (Flores) to Australia (near Perth).

Depth range: 27-139m.

Habitat: in beam trawls with bryozoans, corals and sponges; mud and sand.

***Rhopalorhynchus tenuissimum* (HASWELL 1884)**

- 1884 *Colossendeis tenuissima* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1029-1030, pl. LVI, figs. 5-8.
 1919 *Rhopalorhynchus tenuissimum*, - FLYNN, Pap. Proc. Roy. Soc. Tasm., 1918: 71.
 1958 *Rhopalorhynchus tenuissimum*, - STOCK, Tijdschr. Ent., **101** (2): 125.
 1982 *Rhopalorhynchus tenuissimum*, - STAPLES, Mem. Qd. Mus., **20** (3): 465.

Distribution: Australia (Queensland area).

Depth range: shallow water.

NYMPHONIDAE

- 1891 Nymphonidae, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 54 [diagnosis, remarks].
 1948 Nymphonidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 179-180 [diagnosis, remarks, key to genera].
 1954 Nymphonidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 89 [diagnosis].
 1955 Nymphonidae, - STOCK, Vidensk. Medd. Dansk. Naturhist. Foren., **117**: 219 [key to genera].
 1974 Nymphonidae, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 14.
 1979 Nymphonidae, - Child, Smiths. Contr. Zool., **293**: 35 [key to Middle American species].
 1986 Nymphonidae, - STOCK, Bull. mar. Sci., **38** (3): 418 [remarks].
 01992 Nymphonidae, - CHILD, Mem. Hourglass Cruises, **9** (1): 69-70, tab. 11.

***Boreonymphon* SARS 1888**

1972 *Boreonymphon*, - JUST, Sarsia, **49**: 3-4 [diagnosis, key to species].

***Boreonymphon abyssorum* (NORMAN 1873)**

1972 *Boreonymphon abyssorum*, - JUST, Sarsia, **49**: 6-8, figs. 3, 7D.

1983 *Boreonymphon abyssorum*, - BAMBER, Zool. J. Linn. Soc., **77**: 68.

Distribution: NE-Atlantic.

Depth range: 0-1040m.

Habitat: clay with stones, red clay, clay with stones and gravel, mud, hard clay, hard brown clay, clay with gravel and large stones, clay and *Fucus*.

***Boreonymphon compactum* JUST 1972**

1972 *Boreonymphon compactum* JUST, Sarsia, **49**: 11-14, figs. 6, 7B.

Distribution: N-Atlantic (Bylot Sund).

Depth range: 110-260m.

Habitat: clay and large stones.

***Boreonymphon ossiansarsi* KNABEN 1972**

1972 *Boreonymphon ossiansarsi* KNABEN, in JUST, Sarsia, **49**: 8-11, figs. 4, 5, 7A.

Distribution: N-Atlantic.

Depth range: 0-1098m.

Habitat: reddish brown clay, dark sand and clay, white viscuous clay with stones.

***Boreonymphon robustum* (BELL 1855)**

1891 *Boreonymphon robustum*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 115-119, pl. XII, fig. 3a-d.

1933 *Boreonymphon robustum*, - STEPHENSEN, Medd. Grönland, **79** (6): 4-5, fig. 1 (map).

1972 *Boreonymphon robustum*, - JUST, Sarsia, **49**: 4-6, figs. 2, 7C.

Distribution: Arctica.

Depth range: 97-640m.

Habitat: mud, clay, gravel, and stones.

***Heteronymphon* GORDON 1932**

1932 *Heteronymphon* GORDON, Discovery Rep., **6**: 79-80 [diagnosis, remarks].

1985 *Heteronymphon*, - NAKAMURA, Bull. Biogeogr. Soc. Japan, **40** (4): 33 [key to species].

***Heteronymphon abyssale* (STOCK 1968)**

- 1968 *Nymphon abyssale* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 33-35, fig. 12.
 1970 *Heteronymphon abyssale*, - TURPAEVA, Zool. Zh., 49 (11): 1723-1724, fig. 1 (5), 2 (5).

Distribution: Pacific Ocean off Costa Rica.

Depth range: 3570m.

Habitat: dark muddy clay.

***Heteronymphon bioculatum* TURPAEVA 1956**

- 1956 *Heteronymphon bioculatum* TURPAEVA, Byull. Moskov. Obsh. Ispyt. Prir., Otd. Biol., **61** (2): 69-71, fig. 2.
 1970 *Heteronymphon bioculatum*, - TURPAEVA, Zool. Zh., 49 (11): 1723-1724, fig. 1 (3), 2 (4).
 1971 *Heteronymphon bioculatum*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **92**: 280.

Distribution: NW-Pacific.

Depth range: 2770-3940m.

***Heteronymphon birsteini* TURPAEVA 1954**

- 1956 *Heteronymphon birsteini*, - TURPAEVA, Byull. Moskov. Obsh. Ispyt. Prir., Otd. Biol., **61** (2): 71-72, fig. 3.
 1970 *Heteronymphon birsteini*, - TURPAEVA, Zool. Zh., 49 (11): 1723-1724, fig. 1 (2), 2 (2).

Distribution: NW-Pacific.

Depth range: 1060m.

***Heteronymphon exiguum* (HODGSON 1927)**

- 1932 *Heteronymphon kempfi* GORDON, Discovery Rep., **6**: 80-82, figs. 38-39.
 1938 *Heteronymphon kempfi* var. *dimorpha* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 171-172, fig. 4a-c.
 1944 *Heteronymphon exiguum*, - GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B), **5** (1): 35.
 1953 *Heteronymphon exiguum*, - STOCK, Beaufortia, **35** (4): 37-38, fig. 2a-e.
 1970 *Heteronymphon exiguum*, - TURPAEVA, Zool. Zh., 49 (11): 1723-1724, fig. 1 (1), fig. 2 (1).
 1972 *Heteronymphon exiguum*, - ARNAUD, Tethys, Suppl., **4**: 149-150.

Distribution: Antarctica.

Depth range: 18-393m.

***Heteronymphon horikoshii* NAKAMURA 1985**

- 1985 *Heteronymphon horikoshii* NAKAMURA, Bull. Biogeogr. Soc. Japan, **40** (4): 31-33, fig. 2.

1991 *Heteronymphon horikoshii*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 42.

Distribution: Japan
Depth range: 700-2000m.

Heteronymphon ponsitor CHILD 1982

1982 *Heteronymphon ponsitor* CHILD, Smiths. Contr. Zool., **349**: 31-33, fig. 9.

Distribution: North American Basin; West European Basin.
Depth range: 2022-2862m

Heteronymphon profundum TURPAEVA 1956

- 1956 *Heteronymphon profundum* TURPAEVA, Byull. Moskov. Obsh. Ispyt. Prir., Otd. Biol., **61** (2): 67-69, fig. 1.
 1970 *Heteronymphon profundum*, - TURPAEVA, Zool. Zh., **49** (11): 1723-1724, fig. 1 (4), 2 (3).
 1971 *Heteronymphon profundum*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **92**: 280.

Distribution: NW-Pacific.
Depth range: 6150-6800m.

Neonymphon STOCK 1955

- 1955 *Neonymphon* STOCK 1955, Vidensk. Medd. dansk naturhist. Foren., **117**: 219 [diagnosis].
 1986 *Neonymphon*, - STOCK, Bull. mar. Sci., **38** (3): 420-421 [remarks].

Neonymphon caecum STOCK 1955

- 1955 *Neonymphon caecum* STOCK 1955, Vidensk. Medd. dansk naturhist. Foren., **117**: 219-223, figs. 2-3.
 1986 *Neonymphon caecum*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 421, fig. 6a.

Distribution: tropical W-Atlantic (west of St. Croix, U.S. Virgin Islands; off Surinam).
Depth range: 914-1271m.

Nymphon FABRICIUS 1794

- 1932 *Nymphon*, - GORDON, Discovery Rep., **6**: 26-31 [remarks], 31-36 [key to adult males of Antarctic and subantarctic waters].
 1944 *Nymphon*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 17-22 [remarks, key to antarctic and subantarctic species].
 1948 *Nymphon*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 180-182 [diagnosis, remarks, key to W-Atlantic species south of Newfoundland].
 1949 *Nymphon*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 243-247 [remarks, key

- to species from N-Pacific].
- 1954 *Nymphon*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 89 [diagnosis], 90 [key to South African species].
- 1963 *Nymphon*, - CLARK, Rec. Austr. Mus., **26** (1): 15-16 [key to Australian species].
- 1971 *Nymphon*, - STOCK, Zool. Meded., **47**: 259.
- 1975 *Nymphon*, STOCK, Bull. mar. Sci., **24** (4): 1006-1007 [key to *albatrossi*-group].
- 1986 *Nymphon*, - STOCK, Bull. mar. Sci., **38** (3): 418 [remarks].
- 1988 *Nymphon*, - CHILD, Bull. biol. Soc. Wash., **8**: 67-68 [remarks, key to *Nymphon aequidigitatum*-group].
- 1992 *Nymphon*, - CHILD, Mem. Hourglass Cruises, **9** (1): 70 [diagnosis].

***Nymphon adami* GILTAY 1937**

- 1937 *Nymphon adami* GILTAY, Mem. Mus. Roy. Hist. Nat. Belg., **2** (9): 84-87, figs. 2-5.
- 1956 *Nymphon adami*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, **3** (28) (3): 290-291, fig. 1.
- 1959 *Nymphon adami*, - FAGE, Res. Sci. Campagne Calypso, **4**: 236, fig. 1a.
- 1967 *Nymphon adami*, - STOCK, Atlantide Rep., **9**: 47.
- 1975 *Nymphon adami*, - STOCK, Bull. mar. Sci., **24** (4): 1001-1004, figs. 18, 19a-d.
- 1990 *Nymphon adami*, - STOCK, Cah. Biol. Mar., **63** (16): 221.

Distribution: W-Africa, Mauritania south to Nigeria.

Depth range: 5-22m.

Habitat: ascidians, shells; brown and red algae; muddy sand with shell gravel.

***Nymphon adareanum* HODGSON 1907**

- 1907 *Nymphon adareanum* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 23-25, pl. 3, fig. 3.
- 1913 *Nymphon adareanum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 73.
- 1915 *Nymphon adareanum*, - CALMAN, Brit. Antarct (Terra Nova) Exped., 1910, Zool., **3** (1): 33-34.
- 1923 *Nymphon adareanum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14.
- 1932 *Nymphon adareanum*, - GORDON, Ann. Mag. Nat. Hist., **10** (9): 98-101, figs. 1-2.
- 1950 *Nymphon adareanum*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 149.
- 1952 *Nymphon adareanum*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (2): 181.

Distribution: Antarctica.

Depth range: 31-92m.

***Nymphon adenense* MÜLLER 1989**

- 1989 *Nymphon adenense* MÜLLER, - Senckenbergiana marit., **20** (5/6): 283-284, figs. 21-27.

Distribution: Gulf of Aden.
Depth range: 76m.

Nymphon adenopus STOCK 1991

- 1991 *Nymphon adenopus* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 177-179, figs. 36-37, 185 [text].

Distribution: New Caledonia.
Depth range: 440-450m.

Nymphon aemulum STOCK 1975

- 1948 *Nymphon floridanum*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 196 (only short-necked form).
 1955 *Nymphon floridanum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 215 (only short-necked form).
 1975 *Nymphon aemulum* STOCK, Bull. mar. Sci., **24** (4): 998-1001, fig. 16-17.
 1986 *Nymphon aemulum*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.
 1992 *Nymphon aemulum*, - CHILD, Mem. Hourglass Cruises, **9** (1): 70, fig. 32.

Distribution: tropical W-Atlantic, Florida to Surinam.
Depth range: 0.5-37m.
Habitat: in a variety of substrata.

Nymphon aequidigitatum HASWELL 1885

- 1885 *Nymphon aequidigitatum* HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1022-1024, pl. 54, figs. 1-5.
 1889 *Nymphon aequidigitatum*, - WHITELEGGE, J. & Proc. Roy. Soc. N.S.W., **23** (2): 233.
 1908 *Nymphon aequidigitatum*, - LOMAN, Siboga Exped. Monogr., **40**: 38.
 1919 *Nymphon aequidigitatum*, - FLYNN, Pap. Proc. Roy. Soc. Tasm., 1919: 72-75, pl. XVIII, figs. 4-5, pl. XIX, fig. 6.
 1963 *Nymphon aequidigitatum*, - CLARK, Rec. Austr. Mus., **26** (1): 5-7, fig. 3A-G.
 1973 *Nymphon aequidigitatum*, - STOCK, Beaufortia, **20** (266): 107-108, fig. 3a-d.

Distribution: Australia (Western Port, Crawfish Rock).
Depth range: 10m.
Habitat: on *Amathea* sp. (Bryozoa) and *Aglaophenia* (Hydrozoa).

Nymphon affinis STOCK 1951

- 1951 *Nymphon affinis* STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **2** (43): 5-7, figs. 1-6.
 1954 *Nymphon affinis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 94.

Distribution: South Africa.
Depth range: 15m.

Habitat: rocky shores.

Nymphon akane NAKAMURA & CHILD 1983

- 1983 *Nymphon akane* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 54-56, fig. 19.
1986 *Nymphon akane*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 37-38, fig. 2a-h.
1987 *Nymphon akane*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 158.
1991 *Nymphon akane*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 42.

Distribution: Japan; Korea.

Depth range: 7-30m.

Nymphon albatrossi HEDGPETH 1949

- 1949 *Nymphon albatrossi* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 263-266, fig. 30.

Distribution: Japan.

Depth range: 209-781m.

Nymphon aldabraense CHILD 1988

- 1988 *Nymphon aldabraense* CHILD, Bull. biol. Soc. Wash., **8**: 68-70, fig. 6.

Distribution: Seychelles (Aldabra Atoll).

Depth range: 20m.

Habitat: "Reef slope, coral rubble in hollows between coral heads" (CHILD).

Nymphon altioculatum MÖBIUS 1902

- 1902 *Nymphon altioculatum* MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. Valdivia, 1898-1899, **3**: 181-182, pl. XXVI, figs. 1-6.

Distribution: Bouvet Island.

Depth range: 450m.

Nymphon andamanense CALMAN 1923

- 1923 *Nymphon andamanense* CALMAN, Rec. Indian Mus., **24** (3): 273-275, fig. 4a-f.
1938 *Nymphon andamanense*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 151-152.

Distribution: Arabian Coast; Andaman Islands.

Depth range: 40m.

Nymphon angolense GORDON 1932

- 1932 *Nymphon angolense* GORDON, Discovery Rep., **6**: 77-80, figs. 36-37.
1937 *Nymphon angolense*, - GILTAY, Mem. Mus. Roy. Hist. Nat. Belg., **9** (2): 88.

1951 *Nymphon angolense*, - STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **2** (43): 7-8.
 1954 *Nymphon angolense*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 102-103.

Distribution: South Africa; Angola.

Depth range: 73-110m.

Habitat: muddy bottom.

***Nymphon apheles* CHILD 1979**

1979 *Nymphon apheles* CHILD, Smiths. Contr. Zool., **293**: 35-37, fig. 12.

Distribution: Panamá Pacific.

Depth range: intertidal.

Habitat: in tide pool.

***Nymphon apicatum* STOCK 1991**

1991 *Nymphon apicatum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 180-182, figs. 38-39.

Distribution: New Caledonia.

Depth range: 440-610m.

***Nymphon arabicum* CALMAN 1938**

1938 *Nymphon arabicum* CALMAN, Sci. Rep. John Murray Exped., **5** (6): 152-153, fig. 3A-C.

1965 *Nymphon arabicum*, - STOCK, Beaufortia, **13** (151): 16-18, figs. 4-12.

Distribution: Arabian Coast; Madagascar.

Depth range: 35-83m.

Habitat: muddy sand.

***Nymphon aritai* NAKAMURA & CHILD 1991**

1991 *Nymphon aritai* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 42-44, fig. 14.

Distribution: Japan.

Depth range: 400-479m.

***Nymphon articulare* HODGSON 1908**

1932 *Nymphon articulare*, - GORDON, Discovery Rep., **6**: 65-66, fig. 23a, 28, 29a-b.

Distribution: Antarctica.

Depth range: 24-36m.

***Nymphon australe* HODGSON 1902**

- 1902 *Nymphon australe* HODGSON, Rep. Coll. Nat. Hist. in the Antarct. Regions during the voyage of the "Southern Cross": 257, pl. XL.
- 1902 *Chaetonymphon altioculum*, - MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, 3: 181, pl. XXVI, figs. 1-6.
- 1913 *Nymphon stylops*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), 6: 73-80, figs. 25-31.
- 1915 *Nymphon australe*, - CALMAN, Brit. Antarct (Terra Nova) Exped., 1910, Zool., 3 (1): 36-37.
- 1928 *Nymphon australe*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., 6 (Special Rep., 1): 16.
- 1932 *Nymphon australe*, - GORDON, Discovery Rep., 6: 59-63, figs. 25d, 26b.
- 1938 *Nymphon australe*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), 2 (8): 13.
- 1944 *Nymphon australe*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 24.
- 1944 *Nymphon australe* var. *caecum* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 25-27, fig. 5a-d.
- 1950 *Nymphon australe*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 148.
- 1954 *Nymphon australe*, - BARNARD, Ann. S. Afr. Mus., 41 (3): 90-91.
- 1965 *Nymphon australe*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., 41 (31): 1.
- 1972 *Nymphon australe*, - ARNAUD, Téthys, suppl. 4: 150.
- 1974 *Nymphon australe* var. *caecum*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, 98: 280.
- 1980 *Nymphon australe*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., 52 (18): 2.

Distribution: Antarctica.

Depth range: 8-640m.

***Nymphon banzare* GORDON 1944**

- 1944 *Nymphon banzare* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B) 5 (1): 29-30, fig. 6c-e.
- 1965 *Nymphon banzare*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., 41 (31): 1.
- 1980 *Nymphon banzare* KRAPP, - Bull. Inst. Roy. Sci. Nat. Belg., 52 (18): 3.

Distribution: Antarctica.

Depth range: 130-400m.

***Nymphon barnardi* ARNAUD & CHILD 1988**

- 1988 *Nymphon barnardi* ARNAUD & CHILD, Ann. S. Afr. Mus., 98 (6): 156-159, fig. 8.

Distribution: South Africa.

Depth range: 150-200m.

***Nymphon basispinosum* HILTON 1942**

- 1942 *Nymphon basispinosum* HILTON, J. Ent. Zool. Pomona College, 34 (1): 5.
- 1949 *Nymphon basispinosum*, - HEDGPETH, Proc. U.S. natn. Mus., 98 (3231): 273, fig. 34e.

Distribution: Japan.

Depth range: 104m.

Nymphon benthos HEDGPETH 1949

- 1949 *Nymphon benthos* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 256-257, fig. 25.
 1991 *Nymphon benthos*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 43-44.

Distribution: Japan.

Depth range: 923-1249m.

Nymphon bergi LOSINA-LOSINSKY 1961

- 1961 *Nymphon bergi* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 72-74, fig. 7.

Distribution: NW-Pacific.

Depth range: 291m.

Nymphon biarticulatum (HODGSON 1907)

- 1932 *Nymphon biarticulatum*, - GORDON, Discovery Rep., **6**: 71-73, figs. 27d, 31a, c, 32a.
 1944 *Nymphon biarticulatum*, - GORDON, Rep. Brit. Austr. and N.Z. Antarct. Res. Exped., (B) **5** (1): 22 [key], 27.
 1965 *Nymphon biarticulatum*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., **41** (31): 1-2.
 1969 *Nymphon biarticulatum*, - ARNAUD, Les Terres Austr. Antarct. Françaises, **47**: 36.
 1972 *Nymphon biarticulatum*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 807-808.

Distribution: Antarctica.

Depth range: 211-300m.

Nymphon bicornum ARNAUD & CHILD 1988

- 1988 *Nymphon bicornum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 159-161, fig. 9.

Distribution: South Africa (Zululand and Durban area).

Depth range: 800-850m.

Nymphon biformidens STOCK 1974

- 1974 *Nymphon biformidens* STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 15-16, fig. 4.

Distribution: Madagascar (near Nossi-Be).

Depth range: 1m.

Habitat: dead corals and sand.

Nymphon bigibbulare LOSINA-LOSINSKY, 1961

1961 *Nymphon bigibbulare* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, 7: 59-60, fig. 1.

Distribution: NW-Pacific.
Depth range: 12-137m.

Nymphon birsteini TURPAEVA 1955

1955 *Nymphon birsteini* TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, 12: 322-324, fig. 1.

Distribution: Kurile-Kamtchatka Trench.
Depth range: 1060m.

Nymphon bisseratum LOSINA-LOSINSKY, 1961

1961 *Nymphon bisseratum* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, 7: 66-67, fig. 4.

Distribution: NW-Pacific.
Depth range: 101m.

Nymphon boraborae MÜLLER 1990

1990 *Nymphon boraborae* MÜLLER, Senckenbergiana biol., 70 (1989) (1/3): 192-194, figs. 19-27.

Distribution: Society Islands (Bora Bora).
Depth range: 0.5-1m.
Habitat: fringing reef, dead corals covered with sponges and algae.

Nymphon bouvieri GORDON 1932

1932 *Nymphon bouvieri* GORDON, Discovery Rep., 6: 35 [key], 62 [tab.], 73-75, figs. 25a, 27b, 32c, 33.

1947 *Nymphon bouvieri*, - SPEHENSEN, Sci. Res. Norw. Antarct. Exped., 1927-1928 et seqq., 27: 83.

1974 *Nymphon bouvieri*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, 98: 281.

1980 *Nymphon bouvieri*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., 52 (18): 3.

Distribution: Antarctica.
Depth range: 245-583m.

Nymphon brachyrhynchum HOEK 1881

- 1881 *Nymphon brachyrhynchum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 47-48, pl. IV, figs. 2-7.
- 1913 *Nymphon brachyrhynchum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 73 [key].
- 1923 *Nymphon brachyrhynchum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].
- 1932 *Nymphon brachyrhynchum*, - GORDON, Ann. Mag. Nat. Hist., **10**, (9): 101-104, figs. 3, 4a-b.

Distribution: Kerguelen (Christmas Harbour).

Depth range: 82-220m.

***Nymphon braschnikowi* SCHIMKEWITSCH 1906**

- 1906 *Nymphon braschnikowi* SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **11** (1906): 248-251, figs. 1-7.
- 1930 *Nymphon braschnikowi*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 507-512, figs. 154-160, pl. 9, figs. 1-2.
- 1936 *Nymphon braschnikowi*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 863.
- 1949 *Nymphon braschnikowi*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 250, fig. 21a-c.
- 1955 *Nymphon braschnikowi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 6-7, fig. 2.
- 1961 *Nymphon braschnikowi*, - LOSINA-LOSINSKY, Islledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 62-63.
- 1971 *Nymphon braschnikowi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 319.

Distribution: NW-Pacific; Japan.

Depth range: 95-494m.

***Nymphon brevicaudatum* MIERS 1875**

- 1875 *Nymphon brevicaudatum* MIERS, Ann. Mag. Nat. Hist., (4) **16** (92): 117-118.
- 1879 *Nymphon horridum* BÖHM, Monatsber. Kgl. Preuss. Akad. Wiss. Berlin, 1879: 175, pl. I, fig. 3-3f.
- 1881 *Nymphon brevicaudatum*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (1): 49-52, pl. IV, figs. 12-13, pl. V, figs. 1-5.
- 1889 *Nymphon brevicaudatum*, - PFEFFER, Jb. Hamburg. Wiss. Anst., **6** (2): 41.
- 1902 *Chaetonymphon brevicaudatum*, - MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 181, pl. XXV, fig. 9.
- 1913 *Nymphon brevicaudatum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 95.
- 1932 *Nymphon brevicaudatum*, - GORDON, Discovery Rep., **6**: 69-71, figs. 25b, 26c, 31b, d.
- 1944 *Nymphon brevicaudatum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 27.
- 1972 *Nymphon brevicaudatum*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 808.

Distribution: probably circumpolar in Antarctica. - Kerguelen; ?South Georgia.

Depth range: 27-1080m.

Nymphon brevirostre HODGE 1863

- 1863 *Nymphon brevirostre* HODGE, Ann. Mag. Nat. Hist., **11** (3): 464.
- 1865 *Nymphon rubrum* HODGE, Nat. Hist. Trans. Northumberland and Durham, **1** (1): 41, pl. 10, fig. 1.
- 1877 *Nymphon gracile*, - HOEK, Niederl. Arch. Zool., **3**: 241-243, pl. XV, figs. 11-13, pl. XVI, fig. 18.
- 1881 *Nymphon gracile*, - HOEK, Arch. Zool. Exp. Gén., **9**: 498-501, pl. XXIII, figs. 1-5.
- 1888 *Nymphon gracile*, - SARS, Arch. Math. Naturvidensk., Oslo, **12**: 345.
- 1891 *Nymphon gracile*, - SARS, Norw. Nort-Atl. Exped., 1876-1878, **6** (Zool. 20): 55-58, pl. 5, fig. a-h.
- 1891 *Nymphon rubrum*, - SARS, Norw. Atl.-Exped., 1876-1878, **6** (Zool. 20): 58-61, pl. V, fig. 2a-k.
- 1896 *Nymphon rubrum*, - HERDMAN, Rep. 66th Meeting Brit. Ass. Advancement Sci.: 442.
- 1897 *Nymphon rubrum*, - HURST, in: W.A. Herdman, 10th Ann. Rep. Liverpool Mar. Biol. Committee: 22, 27.
- 1901 *Nymphon grossipes*, - MÖBIUS, Fauna Arctica, **2** (1): 41-44.
- 1908 *Nymphon rubrum*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 208-209, pl. 29, figs. 4-7.
- 1909 *Nymphon grossipes* var. *balthica* HELFER, Inaugural-Diss. Kgl. Christian-Albrechts-Univ. Kiel: 8-10, fig. 2.
- 1909 *Nymphon rubrum*, SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **13** (1908): 432-433.
- 1909 *Nymphon rubrum*, - d'ARCY THOMPSON, in: The Cambr. Nat. Hist.: 541.
- 1910 *Nymphon rubrum*, - TESCH, Jaarb. Rijksinst. Onderzoek der Zee, 1909 (4): 52, 56-57.
- 1928 *Nymphon rubrum*, - GILTAY, Bull. & Ann. Soc. Roy. Ent. Belg., **68** (9-10): 201-205, figs. 2-4.
- 1928 *Nymphon rubrum*, - LOMAN, Die Tierwelt Deutschlands, **8** (3): 78, fig. 4.
- 1930 *Nymphon rubrum*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 381-389, figs. 93-97.
- 1932 *Nymphon rubrum*, - SCHLÖTTKE, Wiss. Meeresunters. Kommiss. Wiss. Unters. Dt. Meere, Abt. Helgoland, (N.F.), **18** (9): 5.
- 1935 *Nymphon brevirostre*, - DERJUGIN, Arktichesti Inst. SSSR, **4**: 11-57.
- 1936 *Nymphon rubrum*, - STEPHENSEN, Bergens Mus. Arbok, 1935, Naturvidensk. Rekke, **7**: 9-10.
- 1936 *Nymphon brevirostre*, - STEPHENSEN, Bergens Mus. Arbok, 1935, Naturvidensk. Rekke, **7**: 10.
- 1942 *Nymphon microcollis* HILTON, J. Ent. Zool. Pomona College, **34** (1): 5.
- 1942 *Nymphon gracile*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 7.
- 1943 *Nymphon rubrum*, - NEEDLER, Can. Atl. Fauna: 11, fig. 13.
- 1945 *Nymphon brevirostre*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 144.
- 1945 *Nymphon rubrum*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 143-144.
- 1948 *Nymphon rubrum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 192, fig. 14.
- 1949 *Nymphon brevirostre*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 248-249.
- 1949 *Nymphon rubrum*, - STOCK, Strand-Werk-Gemeenschap Tabellenser., **7**: 4-5, figs. 1, 8.
- 1955 *Nymphon rubrum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 214, 217-218.
- 1961 *Nymphon brevirostre*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 67-68.
- 1972 *Nymphon brevirostre*, - KING, J. nat. Hist., **6**: 621.
- 1974 *Nymphon brevirostre*, - KING, A Synopsis of the British Sea Spiders. Synopses

- of the British Fauna (New Ser.), **5**: 14 [key], fig. 7.
- 1974 *Nymphon rubrum*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 14 [key], 18-19, fig. 6.
- 1976 *Nymphon rubrum*, - WOLFF, Netherlands J. Sea Res., **10** (4): 473, 474 fig. 2a (distribution in estuary).
- 1978 *Nymphon rubrum*, - STOCK, Zool. J. Linn. Soc., **63**: 59-74 (chemical sense, food preference).
- 1982 *Nymphon brevirostre*, - BAMBER, Biol. J. Linn. Soc., **17**: 275-288.
- 1982 *Nymphon rubrum*, - BAMBER, Biol. J. Linn. Soc., **17**: 275-288, fig. 1.
- 1993 *Nymphon rubrum*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: Norway and Sweden to south of Belgium; Japan; Nova Scotia, Str. of Georgia; Russian Arctic.

Depth range: littoral to 60m.

Habitat: usually amongst hydroids growing on algae; feeds on *Dynamena*, *Tubularia* and *Laomedea* hydroids, also on *Actinia* and *Metridium* anthozoans; associated with tubularians *Halichondria panicea* and *Antennularia* spp.

Remarks: Breeding in spring (incl. March) and summer.

Nymphon brevis NAKAMURA & CHILD 1991

- 1991 *Nymphon brevis* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 44-45, fig. 15.

Distribution: Japan.

Depth range: 102m.

Nymphon brevitarse KRØYER 1838

- 1891 *Nymphon brevitarse*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 61-63, pl. V, fig. 3a-g.
- 1933 *Nymphon brevitarse*, - STEPHENSEN, Medd. Grönland, **79** (6): 10-11.
- 1943 *Nymphon brevitarse*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 85.
- 1963 *Nymphon brevitarse*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1328-1329.

Distribution: Arctica, from Spitzbergen to NW-Greenland and Fox Basin.

Depth range: 10-100m.

Habitat: rocky bottom; from floating seaweed.

Nymphon bullatum STOCK 1992

- 1992 *Nymphon bullatum* STOCK, Tijdschr. Ent., **135**: 121-123, 139, figs. 27-34, 47-51.

Distribution: southern Brazil.

Depth range: 65-250m.

Nymphon bunyipi CLARK 1963

- 1963 *Nymphon bunyipi* CLARK, Rec. Austr. Mus., **26** (1): 9-10, fig. 5A-K.

Distribution: Australia.

Depth range: 81-144m.

Habitat: "found intermingled with hydrozoa and polyzoa"
(CLARK) .

Nymphon caementarum STOCK 1975

1975 *Nymphon caementarum* STOCK, Bull. mar. Sci., **24** (4): 1004-1006, figs. 19e-g,
20.

1990 *Nymphon caementarum*, - STOCK, Zool. Meded., **63** (16): 221-222, fig. 7.

Distribution: Ivory Coast; Liberia; off Mauritania.

Depth range: 37-82m.

Habitat: muddy sand; sandy bottom.

Nymphon caldarium STOCK 1987

1987 *Nymphon caldarium* STOCK, Cah. Biol. Mar., **28**: 509-512, figs. 1-11.

1992 *Nymphon caldarium*, - STOCK, Tijdschr. Ent., **135**: 123 [text].

Distribution: NE-Atlantic.

Depth range: 340-580m.

Habitat: shelly sand; rock, corals.

Nymphon calypso FAGE 1959

1959 *Nymphon calypso* FAGE, Rés. Sci. Campagne Calypso, **4**: 236-239, figs. 2-3.

1966 *Nymphon calypso*, - STOCK, Atlantide Rep., **9**: 49, fig. 1c.

Distribution: W-Africa (off Gambia).

Depth range: 18-79m.

Habitat: mud and sand.

Nymphon chaetochir UTINOMI 1971

1971 *Nymphon chaetochir* UTINOMI, J. Roy. Soc. N.Z., **1** (3/4): 197-200, figs. 1-2.

Distribution: New Zealand (Chatham Rise).

Depth range: 995-1110m.

Nymphon chainae CHILD 1982

1982 *Nymphon chainae* CHILD, Smiths. Contr. Zool., **349**: 33-36, fig. 10.

Distribution: West European Basin, SW of Ireland; Brazil Slope, off Recife.

Depth range: 587-1550m.

Nymphon citerium NAKAMURA & CHILD 1991

1991 *Nymphon citerium* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 45-47, fig. 16.

Distribution: Japan.

Depth range: 1225-1970m.

Nymphon clarencei GORDON 1932

1932 *Nymphon clarencei* GORDON, Discovery Rep., **6**: 54-56, figs. 21-22, 23b.

Distribution: Antarctica.

Depth range: 106-342m.

Nymphon cognatum LOMAN 1928

1928 *Nymphon cognatum* LOMAN. Bull. Soc. Sci. nat. Phys. Maroc., **8** (1-3): 65-67, fig. A.

1937 *Nymphon cognatum*, - GILTAY, Mem. Mus. Roy. Hist. Nat. Belg., **9** (2): 87.

Distribution: NW-Africa (Cap Bojador, Rio de Oro).

Depth range: 64m.

Nymphon comes FLYNN 1928

1928 *Nymphon comes* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 14-16, figs. 4-6.

1951 *Nymphon affinis* STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **43** (2); 5.

1954 *Nymphon affinis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 92-93, figs. 3, 5C.

1954 *Nymphon comes*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 92-93, figs. 3, 5c.

1956 *Nymphon comes*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 78-79, fig. 3.

1962 *Nymphon comes*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 282.

1965 *Nymphon affinis*, - STOCK, Beaufortia, **13** (151): 21-23 [list, text].

1988 *Nymphon comes*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 161.

Distribution: South Africa.

Depth range: 15-580m.

Habitat: sand, shells, corals.

Nymphon compactum HOEK 1881

1881 *Nymphon compactum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 41-43, pl. II, figs. 6-8, pl. XV, fig. 10.

1908 *Nymphon compactum*, - HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1) (6): 167, pl. I, figs. 5, 5a.

1913 *Nymphon compactum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 72 [key].

1923 *Nymphon compactum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].

- 1932 *Nymphon compactum*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 115-117, fig. 10a-c.
 1974 *Nymphon compactum*, - TURPAEVA, Trudy Inst. Okeanol. Akad. Nauk SSSR, **98**: 278-280, pl. 1, figs. 1-7.

Distribution: New Zealand (east of Auckland), ?sub-Antarctic.

Depth range: 2013-3248m.

***Nymphon compactum* LOSINA-LOSINSKY, 1961**

- 1961 *Nymphon compactum* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 75-76, fig. 9.

Distribution: NW-Pacific.

Depth range: 2013-3248m

Remarks: homonym of *N. compactum* HOEK 1881.

***Nymphon conirostrum* STOCK 1973**

- 1973 *Nymphon conirostrum* STOCK, Beaufortia, **20** (266): 111-113, fig. 5.

Distribution: SE-Australia (West Island, Oedipus Point).

Depth range: 13m.

Habitat: "on cave floor and side; on hydroids" (STOCK).

***Nymphon crenatiunguis* BARNARD 1946**

- 1946 *Nymphon crenatiunguis* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 60.
 1954 *Nymphon crenatiunguis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 101-102, fig. 7.
 1988 *Nymphon crenatiunguis*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 162.

Distribution: South Africa.

Depth range: 80-200m.

***Nymphon crosnieri* STOCK 1965**

- 1965 *Nymphon crosnieri* STOCK, Beaufortia, **13** (151): 19-23, figs. 18-27.

Distribution: Madagascar.

Depth range: 35m.

Habitat: muddy sand.

***Nymphon curvidens* STOCK 1990**

- 1990 *Nymphon curvidens* STOCK, Zool. Meded., **63** (16): 223-226, figs. 8-9.

Distribution: Mauritania.

Depth range: 32m.

Habitat: shell gravel and bivalve shells, on *Pyura*.

Nymphon diabolus CHILD 1988

- 1988 *Nymphon diabolus* CHILD, Smiths. Contr. Zool., **468**: 23-25, fig. 10.
 1988 *Nymphon diabolus*, - CHILD, Bull. biol. Soc. Wash., **8**: 75.
 1988 *Nymphon diabolus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 668-669.
 1991 *Nymphon diabolus*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 145-146.

Distribution: Japan (Ryukyu Islands); Guam; Philippines (Negros, Siquijor and Apo Islands); Seychelles (Aldabra Atoll).

Depth range: 1-21m.

Habitat: coral reefs. - on reef front, found on dead branching coral.

Nymphon discorsicoxae CHILD 1982

- 1982 *Nymphon discorsicoxae* CHILD, Smiths. Contr. Zool., **349**: 36-38, fig. 11.
 1991 *Nymphon discorsicoxae*, - STOCK, Mém. Mus. natn. Hist. (A), **151**: 177 [text].

Distribution: Brazil slope, off Recife.

Depth range: 587-1493m.

Nymphon dissimilis HEDGPETH 1949

- 1949 *Nymphon dissimilis* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 262-263, fig. 28.

Distribution: Japan.

Depth range: 1914m.

Nymphon distensum MÖBIUS 1902

- 1902 *Nymphon distensum* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 179-180, pl. XXV, figs. 1-6.
 1923 *Nymphon distensum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].
 1928 *Nymphon distensum*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 10-11.
 1932 *Nymphon distensum*, - GORDON, Ann. Mag. Nat. Hist., **41** (3): 105-106, fig. 6a-b.
 1932 *Nymphon distensum*, - GORDON, Discovery Rep., **6**: 28-29, 35 [key].
 1944 *Nymphon distensum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 20 [key].
 1954 *Nymphon distensum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 98-100, fig. 5b.
 1988 *Nymphon distensum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 163.
 1988 *Nymphon distensum*, - MUNILLA, Monogr. Zool. Mar., **3**: 187-189, figs. 6-7.

Distribution: South Africa (Cape Point to Zululand region;

Namibia).

Depth range: 80-560m.

Nymphon draconis CHILD 1990

1990 *Nymphon draconis* CHILD, Proc. biol. Soc. Wash., **103** (2): 324-327, fig. 5.

1992 *Nymphon draconis*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 163.

Distribution: Australian Barrier Reef; Malaysia (Pulau Tioman).

Depth range: 1-3m.

Habitat: dead corals, rubble.

Nymphon dubitabile STOCK 1973

1973 *Nymphon dubitabile* STOCK, Beaufortia, **20** (266): 109-111, fig. 4.

Distribution: SE-Australia (Western Port, Crawfish Rock).

Depth range: 10m.

Habitat: on *Amathea* sp. (Bryozoa) and *Aglaophenia plumosa* (Hydroida)

Nymphon duospinum (HILTON 1942)

1942 *Chaetonymphon duospinum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 6.

1949 *Nymphon duospinum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 274, fig. 331.

Distribution: Alaska (Kiska Harbor).

Depth range: 18m.

Nymphon elegans HANSEN 1886

1891 *Nymphon elegans*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 86-89, pl. IX, fig. 1a-g.

1933 *Nymphon elegans*, - STEPHENSEN, Medd. Grönland, **79** (6): 17.

1943 *Nymphon elegans*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 87.

1963 *Nymphon elegans*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1332.

Distribution: European Arctic across Russia to about 80° E.

Depth range: eurybathic (100-1358m).

Nymphon elongatum HILTON 1942

1942 *Nymphon elongatum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 5.

1949 *Nymphon elongatum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 251-252, fig. 22, 273, fig. 34f.

1971 *Nymphon elongatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 320.

1987 *Nymphon elongatum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 158, fig. 15A-

H.

Distribution: Japan; Korea.
Depth range: 110-132m.

Nymphon falcatum UTINOMI 1955

- 1955 *Nymphon falcatum* UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 15-17, fig. 9.
 1956 *Nymphon falcatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 94.
 1965 *Nymphon falcatum*, - STOCK, Beaufortia, **13** (151): 21.
 1971 *Nymphon falcatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.
 1991 *Nymphon falcatum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 47.

Distribution: Japan.
Depth range: 120-421m.

Nymphon femorale FAGE 1956

- 1956 *Nymphon femorale* FAGE, Galathea Rep., **2**: 161-163, figs. 5-7.
 1966 *Nymphon femorale*, - BELYAEV, Trudy Inst. Okeanol. Akad. Nauk SSSR, **591**: 77.
 1968 *Nymphon femorale*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 32-33, fig. 10i-j.
 1969 *Nymphon femorale*, - TURPAEVA, In: V.G. Kort, ed., Biol. Tikhogo okeana, **2**, Glubokavod naia donna fauna pleuston; Akad. Nauk SSR, Inst. Okeanol: tab. 14.
 1979 *Nymphon femorale*, - MARSHALL, Developments in Deep Sea Biol.: 204, fig. 87.
 1984 *Nymphon femorale*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4), **6**, (3): 701-702.

Distribution: Indian Ocean (between Seychelles and Sri Lanka; Carpenter Ridge).

Depth range: 4040-6650m.

Nymphon flindersi CHILD 1975

- 1975 *Nymphon flindersi* CHILD, Smiths. Contr. Zool., **190**: 3-5, fig. 2.

Distribution: Australia (West of Garden Island).
Depth range: 17-33m.

Nymphon floridanum HEDGPETH 1948

- 1910 *Nymphon* sp., - COLE, Biol. Woods Hole Mar. Biol. Lab., **18** (4): 196.
 1948 *Nymphon floridanum* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 196-199, fig. 17 (only long-necked form).
 1955 *Nymphon floridanum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 215, fig. 1a (only long-necked form).
 1973 *Nymphon floridanum*, - KRAEUTER, J. nat. Hist., **7**: 494.
 1975 *Nymphon floridanum*, - STOCK, Bull. mar. Sci., **24** (4): 994-998, figs. 14-15.
 1976 *Nymphon floridanum*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27** (3-4): 336-337.

- 1979 *Nymphon floridanum*, - CHILD, Smiths. Contr. Zool., **293**: 37.
1982 *Nymphon floridanum*, - CHILD, Smiths. Contr. mar. Sci., **12**: 374.
1986 *Nymphon floridanum*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1.
1992 *Nymphon floridanum*, - CHILD, Mem. Hourglass Cruises, **9** (1): 70 [key], 71-72,
fig. 33, 76 [tab. 12], 78 [tab. 14].

Distribution: U.S.A. (Georgia) to north-eastern coast of South America (French Guiana).

Depth range: 1-402m.

Habitat: in a variety of substrata.

Nymphon forceps NAKAMURA & CHILD 1991

- 1991 *Nymphon forceps* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 47-49, fig. 17.

Distribution: Japan.

Depth range: 410-440m.

Nymphon foresti FAGE 1953

- 1953 *Nymphon foresti* FAGE, Bull. Mus. natn. Hist. nat., Paris, (2), **25** (4): 378-380, figs. 3-4.

Distribution: E-Atlantic (Guibn  e fran  aise).

Depth range: 30m.

Habitat: in samples with shells, sponges and hydroids.

Nymphon foxi CALMAN 1927

- 1927 *Nymphon foxi* CALMAN, Trans. Zool. Soc. London, **22** (3): 403-405, fig. 102.
1938 *Nymphon foxi*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 152.
1957 *Nymphon foxi*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 86-87, fig. 4.
1968 *Nymphon foxi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 31-32.
1989 *Nymphon foxi*, - M  LLER, Senckenbergiana marit., **20** (5/6): 284-287, figs. 28-34.

Distribution: Suez Canal; Arabian Coast; Gulf of Aden; SE Australia.

Depth range: 40-1315m.

Habitat: gravel.

Nymphon frigidum HODGSON 1907

- 1907 *Nymphon frigidum* HODGSON, Rep. Natn. Antarct. Exped., 1901-1904, Nat. Hist., **3**: 25-26, pl. III, fig. 4, pl. X, fig. 10.
1938 *Nymphon longitarse* var. *antarctica*, - HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 167-168, fig. 2a, c, e.
1944 *Nymphon frigidum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 21 [key].
1953 *Nymphon frigidum*, - STOCK, Beaufortia, **35** (4): 36.

Distribution: Antarctica.

Nymphon fuscum HOEK 1881

1881 *Nymphon fuscum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 48-49, pl. IV, figs. 8-11.

Distribution: off Kerguelen Island.

Depth range: 46m.

Nymphon galatheaæ FAGE 1956

1956 *Nymphon galatheaæ* FAGE, Galathea Rep., **2**: 159-161, figs. 1-4.

Distribution: Southern Oceans.

Depth range: 5340-5850m.

Nymphon gerlachei GILTAY 1935

1935 *Nymphon gerlachei* GILTAY, Res. Voy. Belgica, Zool.: 7-10, figs. 5-9.

Distribution: Antarctica.

Depth range: 460m.

Nymphon giltayi HEDGPETH 1948

1948 *Nymphon giltayi* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 195-196, fig. 16.

Distribution: NW-Atlantic.

Nymphon giraffa LOMAN 1908

1908 *Nymphon giraffa* LOMAN, Siboga Exped. Monogr., **40**: 39-40, pl. VI, figs. 74-82.

1962 *Nymphon giraffa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 92, figs. 1-2.

1971 *Nymphon giraffa*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.

1987 *Nymphon giraffa*, - NAKAMURA, Biol. Lab., Imperial Household: 3-4, pl. 1.

Distribution: Makassar Strait, Indonesia; Sagami Bay, Japan.

Depth range: 16-70m.

Nymphon gracile LEACH 1814

1923 *Nymphon gracile*, - BOUVIER, Faune de France, **7**: 30.

1932 *Nymphon gracile*, - FAGE, Arch. Zool. Exp. Gén., **74** (14): 251-261 [occurrence in plankton].

- 1945 *Nymphon gracile*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 143.
 1972 *Nymphon gracile*, - KING, J. nat. Hist., **6**: 621.
 1973 *Nymphon gracile* ARNAUD, Tethys, **5** (1): 152.
 1973 *Nymphon gracile*, - JARVIS & KING, Mar. Behav. Physiol., **2**: 1-13 [ultrastructure of photoreceptors].
 1974 *Nymphon gracile*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 14 [key], 16-17, fig. 5.
 1976 *Nymphon gracile*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.
 1978 *Nymphon gracile*, - STOCK, Zool. J. Linn. Soc., **63**: 59-74 (chemical sense, food preference).
 1979 *Nymphon gracile*, - JARVIS & KING, Mar. Behav. Physiol., **6**: 307-317 [response to light stimuli].
 1987 *Nymphon gracile*, - Arnaud, Mesogee, **47**: 53.
 1990 *Nymphon gracile*, - STOCK, Zool. Meded., **63** (16): 226.
 1993 *Nymphon gracile*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: NE-Atlantic, from Denmark to the Mediterranean.

Depth range: 0-52m.

Habitat: feeds on *Laomedea* and *Dynamena pumila* hydroids; also on *Actina equina* *Bowerbankia* and feeding on *Nucella* eggs; further on drifting weed, on rocks among sponges, polyzoans, tunicates and *Serturella*.

***Nymphon gracilipes* MIERS 1875**

- 1891 *Nymphon gracilipes*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 83-86, pl. VIII, fig. 3a-g.
 1902 *Nymphon gracilipes*, - MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 180, pl. XXV, figs. 7-8.
 1915 *Nymphon gracilipes*, - CALMAN, Ann. Mag. Nat. Hist., **15**: 584-588, figs. 1-4.
 1944 *Nymphon gracilipes*, - GORDON, Rep. Brit., Austr., and N.Z. Antarct. Res. Exped., (B), **5** (1): 22-23.
 1972 *Nymphon gracilipes*, - ARNAUD, Bull. Mus. natn. Hist. nat., (3) **51** (65): 808.

Distribution: Antarctica (off Gauss and Weddell Quadrants); Kerguelen.

Depth range: 10-3055m.

***Nymphon gracillimum* CALMAN 1915**

- 1915 *Nymphon gracillimum* CALMAN, Rep. Brit. Antarct. "Terra Nova" Exped., Zool. **3**: 30-32, fig. 5.
 1932 *Nymphon gracillimum*, - GORDON, Discovery Rep., **6**: 42-43, figs. 12b, 13a.
 1992 *Nymphon gracillimum*, - STOCK, Tijdschr. Ent., **135**: 124, 126 [text].

Distribution: cold to polar waters in Southern Hemisphere; Antarctica.

Depth range: 200-234m.

***Nymphon granulatum* ARNAUD & CHILD 1988**

1988 *Nymphon granulatum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 163-166, fig. 10.

Distribution: South Africa.
Depth range: 620-850m.

***Nymphon groenlandicum* MEINERT 1899**

1899 *Nymphon groenlandicum* MEINERT, Danish Ingolf-Exped., **3** (1): 41-42, pl. III, figs. 14-22.

1933 *Nymphon groenlandicum*, - STEPHENSEN, Medd. Grönland, **79** (6): 15.

Distribution: Arctica (Greenland).

Depth range: 339fms-740m.

Habitat: soft, gray clay, numerous pebbles, mostly granite.

***Nymphon grossipes* KRØYER 1845**

1780 ?*Pycnogonum grossipes* FABRICIUS, Fauna Groenlandica: 229.

1945 *Nymphon grossipes* KRØYER, Naturhistorisk Tidskr., Kobenhav, **1** (2): 108.

1853 *Nymphon grossipes*, - STIMPSON, Smiths. Contr. Knowl., **6** (5): 38.

1874 *Nymphon grossipes*, - VERRILL, Am. J. Sci. & Arts, **3** (7): 411.

1874 *Nymphon grossipes*, - VERRILL, Am. J. Sci. & Arts, **3** (7): 502.

1878 *Nymphon grossipes*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 20-22, pl. 17, fig. 1a-q.

1880 *Nymphon grossipes*, - WILSON, Rep. U.S. Fish. Comm., **6**: 491-494, pl. 6, figs. 32-37, pl. 7, fig. 42

1881 *Nymphon grossipes*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 44-45, pl. III, figs. 9-12, pl. IV, fig. 1.

1881 *Nymphon grossipes*, - Wilson, Bull. Mus. Comp. Zool., Harvard, **8** (12): 253.

1885 *Nymphon grossipes*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 561.

1891 *Nymphon glaciale*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 63-65, pl. VI, fig. 1a-q

1891 *Nymphon grossipes*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 65-68, pl. 6, fig. 2a-i

1891 *Nymphon mixtum*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 68-71, pl. 6, fig. 3a-i

1901 *Nymphon grossipes*, - WHITEAVES, Ann. Mag. Nat. Hist. **10** (4): 264.

1913 *Nymphon grossipes*, - SUMNER, OSBURN & COLE, Bull. Bur. Fish., Washington, **31** (1, 2): 677.

1916 *Nymphon grossipes*, - APPELLÖF, Kungl. Svensk. Vetensk. Handl., **54** (5): 13-16, figs. 5-6.

1930 *Nymphon grossipes* var. *mixtum* SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 416-421, figs. 107-109.

1930 *Nymphon grossipes* var. *glaciale* SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 421-425, figs. 110-112.

1933 *Nymphon grossipes*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 70-71.

1933 *Nymphon grossipes* subsp. *glaciale* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 69-70.

1933 *Nymphon grossipes*, - STEPHENSEN, Medd. Grönland, **79** (6): 11-12.

1934 *Nymphon grossipes mixtum*, - GILTAY, Can. Field Naturalist, **48**: 50.

- 1936 *Nymphon turritum* EXLINE, Proc. U.S. natn. Mus., **83** (2991): 416-418, figs. g-k.
- 1936 *Nymphon mixtum*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 862.
- 1942 *Nymphon glaciale*, - GILTAY, J. Fish. Res. Board Canada, **5** (5): 459.
- 1942 *Nymphon mixtum*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 5.
- 1942 *Nymphon oculospinum* HILTON, - J. Ent. Zool. Pomona College, **34** (1): 4-5.
- 1942 *Nymphon nigrognathum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 6.
- 1942 *Nymphon turritum*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 6.
- 1942 *Nymphon grossipes*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 7.
- 1943 *Nymphon grossipes*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 85-86.
- 1943 *Nymphon glaciale*, - NEEDLER, Can. Atl. Fauna, **10**: 8, fig. 7a-e.
- 1943 *Nymphon grossipes*, - NEEDLER, Can. Atl. Fauna, **10**: 5-7, fig. 5a-e.
- 1943 *Nymphon mixtum*, - NEEDLER, Can. Atl. Fauna, **10**: 7, fig. 6a-e.
- 1943 *Nymphon grossipes*, - STEPHENSEN, Medd. Grönland, **121** (8): 18-20, fig. 6.
- 1948 *Nymphon grossipes*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 187-189, fig. 13a.
- 1949 *Nymphon grossipes*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 247.
- 1949 *Nymphon nigrognathum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 274, fig. 34g.
- 1955 *Nymphon grossipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 4-5, fig. 1.
- 1959 *Nymphon grossipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 198.
- 1961 *Nymphon grossipes*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 69-70.
- 1961 *Nymphon grossipes* n. var. *longidactylus* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 70, fig. 5.
- 1963 *Nymphon grossipes*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1329-1330.
- 1971 *Nymphon grossipes*, - TURPAEVA, Trudy Inst. Okeanol. Akad. Nauk SSSR, **92**: 275, pl. 1, figs. 1-5.
- 1971 *Nymphon grossipes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 318.
- 1991 *Nymphon grossipes*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 49.

Distribution: circum boreal, more often taken in N-Atlantic and Alaskan-Russian Arctic localities.

Depth range: 11-1025m.

Habitat: mud and stones.

***Nymphon grus* STOCK 1991**

- 1991 *Nymphon grus* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 220-223, figs. 5-6.

Distribution: Philippines.

Depth range: 266m.

***Nymphon gruveli* BOUVIER 1910**

- 1910 *Nymphon gruveli* BOUVIER, Act. Soc. Linn., Bordeaux, **64**: 331-335, figs. 1-6.
- 1928 *Nymphon gruveli*, - LOMAN, Bull. Soc. Sci. Nat. Phys. Maroc, **8** (1-3): 63-64.
- 1929 *Nymphon gruveli*, - LOMAN, Bull. Soc. Sci. Nat. Maroc, **9** (1-6): 71.
- 1937 *Nymphon gruveli*, - GILTAY, Mém. Mus. Roy. Hist. nat. Belg., **9** (2): 83-84, fig. 1.
- 1942 *Nymphon gruveli*, - FAGE, Arch. Zool. Gen., 82 (Notes & Rev., 2): 76-77.
- 1951 *Nymphon gruveli*, - STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **43** (2): 8.

- 1959 *Nymphon gruveli*, - FAGE, Res. Sci. Campagne Calypso, **4**: 236, fig. 1b.
 1966 *Nymphon gruveli*, - STOCK, Atlantide Rep., **9**: 46-47.
 1990 *Nymphon gruveli*, - STOCK, Zool. Meded., **63** (16): 226-227, fig. 10.

Distribution: NW-Africa, from Morocco to Senegal.

Depth range: 18-150m.

Habitat: sand, shells, bryozoans, muddy sand, hard bottom, sponges, brown algae, worm tubes, calcareous gravel.

***Nymphon gunteri* HEDGPETH 1949**

- 1949 *Nymphon gunteri* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 257-259, fig. 26.
 1987 *Nymphon gunteri*, - NAKAMURA, Biol. Lab., Imperial Household: 4-5, pl. 2.
 1991 *Nymphon gunteri*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 49.

Distribution: Japan.

Depth range: 100-2000m.

***Nymphon hadale* CHILD 1982**

- 1982 *Nymphon hadale* CHILD, Smiths. Contr. Zool., **349**: 38-40, fig. 12.

Distribution: Argentine Basin.

Depth range: 4435-5781m.

***Nymphon hamatum* HOEK 1881**

- 1881 *Nymphon hamatum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 36, pl. I, figs. 1-9.
 1913 *Nymphon hamatum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 72 [key].
 1923 *Nymphon hamatum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].
 1932 *Nymphon hamatum*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 109-112, fig. 9b.

Distribution: southern Oceans.

Depth range: 2516-2928m.

***Nymphon hampsoni* CHILD 1982**

- 1982 *Nymphon hampsoni* CHILD, Smiths. Contr. Zool., **349**: 40-42, fig. 13.

Distribution: Argentine Basin, off the Rio de la Plata.

Depth range: 993-2323m.

***Nymphon hedgpethi* STOCK 1953**

- 1953 *Nymphon hedgpethi* STOCK, Temminckia, **9**: 285-288, fig. 7.

Distribution: East Indian Archipelago.
Depth range: 686m.

Nymphon heterodenticulatum HEDGPETH 1941

1941 *Nymphon heterodenticulatum* HEDGPETH, Am. Midl. Nat., **25** (2): 447-449, pl. 1 (figs. a-h).

Distribution: southern California.
Depth range: 146m.
Habitat: dead and bleached corallines.

Nymphon heterodentum TURPAEVA 1991

1991 *Nymphon heterodentum* TURPAEVA, Zool. Zh., **70** (12): 34-37, fig. 1.

Distribution: shelf and slope of south-eastern Africa.
Depth range: 355-529m.

Nymphon heterospinum HEDGPETH 1949

1949 *Nymphon heterospinum* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 259-260, fig. 27.

1991 *Nymphon heterospinum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 49.

Distribution: Japan.
Depth range: 102-563m.

Nymphon hiemale HODGSON 1907

1932 *Nymphon hiemale*, - GORDON, Discovery Rep., **6**: 39-42, figs. 12a, 13c.
 1974 *Nymphon hiemale*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **98**: 278.
 1992 *Nymphon hiemale*, - STOCK, Tijdschr. Ent., **135**: 124, 126 [text].

Distribution: cold to polar waters in Southern Hemisphere;
 Antarctica.

Depth range: 110-342-?1613m.

Nymphon hirtipes BELL 1853

1853 *Nymphon hirtipes* BELL, In: The Last of the Arct. Voyages, being a Narrative
 of the Exped. of HMS Assistance, under the command of Captain Sir Edward
 Belcher, **2**: 403, pl. 35, fig. 3.
 1878 *Nymphon hirtipes*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 22-23, pl. 5,
 figs. 2-3, pl. 6, fig. 2a-k.
 1880 *Nymphon hirtum* WILSON, - Rep. U.S. Fish. Comm., **6**: 495-497, pl. 7, figs. 38-
 41.
 1891 *Chaetonymphon hirtipes*, SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool.
 20): 103-107, pl. XI, fig. 2a-k.
 1901 *Nymphon hirtum*, - WHITEAVES, Ann. Mag. Nat. Hist., **10** (4): 264.

- 1933 *Chaetonymphon hirtipes*, - STEPHENSEN, Medd. Grönland, **79** (6): 8-9, figs. 2 (map), 10.
- 1943 *Nymphon hirtipes*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 83, 85.
- 1943 *Chaetonymphon hirtipes*, - NEEDLER, Can. Atl. Fauna, **10**: 11-12, fig. 14a-e.
- 1943 *Chaetonymphon hirtipes*, - STEPHENSEN, Medd. Grönland, **121** (8): 9-14, figs. 2-3.
- 1948 *Nymphon hirtipes*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 183-185, figs. 10b, 11b.
- 1963 *Nymphon hirtipes*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1326-1327.

Distribution: Arctic and boreal-Arctic, distributed in the N-Atlantic and from Kara Sea to NW-Greenland; also known from eastern U.S.A.

Depth range: 2-399m.

Habitat: gravel; mud and pebbles; mud and stones.

***Nymphon hirtum* KRØYER 1838**

- 1891 *Chaetonymphon hirtum*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 101-103, pl. XI, fig. 1a-g.
- 1898 *Chaetonymphon hirtum*, - HOEK, Tijdschr. Nederl. Dierk. Ver., ser. 2, vol. **1**: 297-300, pl. III, figs. 14-19.
- 1963 *Nymphon hirtum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1327.
- 1974 *Nymphon hirtum*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 14 [key], 15, fig. 4.

Distribution: Iceland to Murmansk Sea, along the coast of Norway, south to Belgium.

Depth range: 79-below 200m.

***Nymphon hodgsoni* SCHIMKEWITSCH 1913**

- 1913 *Nymphon hodgsoni* SCHIMKEWITSCH, Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, **18**: 244-248, pl. 3, figs. 15-25.
- 1930 *Nymphon hodgsoni*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nau SSSR. **2**: 512-517, figs. 161-166, pl. 10.
- 1933 *Nymphon hodgsoni*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 71.
- 1936 *Nymphon hodgsoni*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 863.
- 1949 *Nymphon hodgsoni*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 250, fig. 21d-g.
- 1961 *Nymphon hodgsoni*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 63.
- 1971 *Nymphon hodgsoni*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **92**: 275.
- 1971 *Nymphon hodgsoni*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 319.

Distribution: Japan Sea; Tatary Straits; Okhotsk Sea; southern Kuriles.

Depth range: 72-414m.

***Nymphon hoekii* MEINERT 1899**

1899 *Nymphon hoekii* MEINERT, Danish Ingolf Exped., 3 (1): 39-41, pl. III, fig. 7-13.

Distribution: Arctica, between Greenland and Iceland (Denmark Strait).

Depth range: 1376m.

***Nymphon immane* STOCK 1954**

1954 *Nymphon immane* STOCK, Vidensk. Medd. dansk naturhist. Foren., 116: 25-28, figs. 9-10.

1963 *Nymphon immane*, - CLARK, Rec. Austr. Mus., 26 (1): 7, fig. 4A-B.

Distribution: Australia (off Eden, New South Wales).

Depth range: 37-540m.

Habitat: soft bottom.

***Nymphon improcerum* NAKAMURA & CHILD 1991**

1991 *Nymphon improcerum* NAKAMURA & CHILD, Smiths. Contr. Zool., 512: 49-51, fig. 18.

Distribution: Japan (Sagami Bay).

Depth range: 33-41m.

***Nymphon inaequipes* STOCK 1992**

1992 *Nymphon inaequipes* STOCK, Tijdschr. Ent., 135: 124-126, 128 [text], 139, figs. 35-46.

Distribution: southern Brazil.

Depth range: 26-500m.

***Nymphon inerme* FAGE 1956**

1956 *Nymphon inerme* FAGE, Galathea Rep., 2: 163-164, figs. 8-10.

1969 *Nymphon inerme*, - TURPAEVA, in: V.G. Kort, ed., Biol. Tikhogo okeana, 2, Glubokavod naia donna fauna pleuston, Akad. Nauk, SSR, Inst. Okeanol: pl. 14.

1971 *Nymphon inerme*, - STOCK, Bull. Zoöl. Mus., Univ. Amsterdam, 2 (4): 25-26, figs. 2-9.

1974 *Nymphon inerme*, - ARNAUD, Bull. Zoöl. Mus., Univ. Amsterdam, 3 (21): 170-171.

1981 *Nymphon inerme* STOCK, Cah. Biol. Mar., 22: 455.

1982 *Nymphon inerme*, - CHILD, Smiths. Contr. Zool., 349: 42.

Distribution: Kermadec Trench in the Pacific; between Greenland and Labrador Sea; Azores; Argentine Basin; Walvis Basin.

Depth range: 3550-4410m.

Nymphon infundibulum NAKAMURA & CHILD 1991

1991 *Nymphon infundibulum* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 51-53, fig. 19.

Distribution: Japan (Kii Strait, central Honshu).

Depth range: 353-492m.

Nymphon japonicum ORTMANN 1890

- 1891 *Nymphon japonicum* ORTMANN, Zool. Jb. (Syst.), **5** (1): 158-159, pl. 24, fig. 1.
 1899 *Nymphon japonicum*, - SHISHIDO, Dôzatsu, **11** (128): 198.
 1911 *Nymphon japonicum*, - LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Nat. Kl.), Suppl. **2** (4): 8.
 1919 *Nymphon japonicum*, - FUKUI, Rigakukai[?], **16** (2): 99.
 1936 *Nymphon japonicum*, - OHSHIMA, Dôbutusugaku Zasshi, **48** (8-10): 8.
 1949 *Nymphon japonicum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 249-250, fig. 20a-e.
 1951 *Nymphon japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 159-160.
 1954 *Nymphon japonicum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 18-20, fig. 6a-c.
 1955 *Nymphon japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 5-6.
 1959 *Nymphon japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 199.
 1962 *Nymphon japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 92.
 1971 *Nymphon japonicum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 320.
 1983 *Nymphon japonicum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 56.
 1987 *Nymphon japonicum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 158-160, fig. 16A-H.
 1987 *Nymphon japonicum*, - NAKAMURA, Biol. Lab., Imperial Household: 5.

Distribution: Japan.

Depth range: 20-5488m.

Habitat: hard bottom.

Nymphon kensleyi CHILD 1988

- 1988 *Nymphon kensleyi* CHILD, Bull. biol. Soc. Wash., **8**: 70-72, fig. 7.

Distribution: Belize (Carrie Bow Cay).

Depth range: 15m.

Habitat: "rubble from outer reef ridge" (CHILD).

Nymphon kodanii HEDGPETH 1949

- 1949 *Nymphon kodanii* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 252-254, fig. 23a-j.
 1954 *Nymphon kodanii*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 21.
 1955 *Nymphon kodanii*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 7-8, fig. 3.
 1962 *Nymphon kodanii*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 92.
 1971 *Nymphon kodanii*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 320.
 1971 *Nymphon kodanii*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk

SSSR, **92**: 275-276, pl. 1, figs. 6-9.
 1991 *Nymphon kodanii*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 53.

Distribution: Sea of Japan.

Depth range: 75-1970m.

Habitat: sand bottom.

Nymphon kurilense LOSINA-LOSINSKY, 1961

1961 *Nymphon kurilense* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 74-75, fig. 8.

Distribution: NW-Pacific.

Depth range: 3-52m.

Nymphon kurilokamchaticum TURPAEVA 1971

1971 *Nymphon kurilokamchaticum* TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **92**: 277-279, pl. 3, figs. 1-8.

Distribution: Kurile-Kamtchatka.

Depth range: 2770-2820m.

Nymphon lanare HODGSON 1907

1907 *Nymphon lanare* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., **3**: 22, pl. III fig. 2, pl. X fig. 9.
 1913 *Nymphon charcoti*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 81-86, figs. 32-34.
 1915 *Nymphon charcoti*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 29.
 1923 *Nymphon lanare*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 16.
 1932 *Nymphon charcoti*, - GORDON, Discovery Rep., **6**: 28-29 [tab.], 34 [key], 51-54, fig. 10b.
 1944 *Nymphon lanare*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 18 [tab. + key].
 1965 *Nymphon lanare*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., **41** (31): 2.
 1974 *Nymphon charcoti*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **98**: 280.
 1980 *Nymphon lanare*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 3.

Distribution: Antarctica.

Depth range: 60-391m.

Nymphon laterospinum STOCK 1963

1963 *Nymphon laterospinum* STOCK, Ann. S. Afr. Mus., **46** (12): 322-323, fig. 1.
 1978 *Nymphon laterospinum*, - STOCK, Cah. Biol. Mar., **19**: 211-212, fig. 9a-d.
 1982 *Nymphon laterospinum*, - CHILD, Smiths. Contr., **349**: 42.
 1984 *Nymphon laterospinum*, - STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **6**

(sec. A) (3): 702.

1984 *Nymphon laterospinum*, - STOCK, J. mar. biol. Ass. U.K., **64**: 746.

1991 *Nymphon laterospinum*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 180.

Distribution: NE-Atlantic; off Saldanha Bay; off Rio de la Plata; Newfoundland Basin; off Durban, Indian Ocean.

Depth range: 1894-4715.

***Nymphon leptochelus* SARS 1888**

1891 *Nymphon leptochelus*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 78-80, pl. VIII, fig. 1a-i.

1933 *Nymphon leptochelus*, - STEPHENSEN, Medd. Grönland, **79** (6): 15-16.

1983 *Nymphon leptochelus* BAMBER, Zool. J. Linn. Soc., **77**: 66-68.

Distribution: N-Atlantic, inclusive Arctica.

Depth range: 600-1300m.

***Nymphon lituus* CHILD 1979**

1979 *Nymphon lituus* CHILD, Smiths. Contr. Zool., **293**: 38-40, fig. 13.

Distribution: Mexico Pacific (Gulf of California); Panamá Pacific.

Depth range: 0-9m.

Habitat: associated with floating *Sargassum*; among hydroids and bryozoans on rocky shore.

***Nymphon lobatum* STOCK 1962**

1962 *Nymphon lobatum* STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 278-281, figs. 3-4.

1966 *Nymphon lobatum*, - STOCK, Atlantide Rep., **9**: 47-48, fig. 1a-b.

Distribution: E-Atlantic; South Africa.

Depth range: 40-93m.

Habitat: rocks.

***Nymphon lomani* GORDON 1944**

1944 *Nymphon lomani* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 24-24, fig. 4a-c.

Distribution: Antarctica

Depth range: 180-220m.

***Nymphon longicaudatum* CARPENTER 1904**

1904 *Nymphon longicaudatum* CARPENTER, Ceylon Pearl Oyster Fish., 1904, Suppl. Rep., **13**: 183, figs. 8-14.

1968 *Nymphon longicaudatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**:

31, fig. 10b-d.

Distribution: India (Pamban Pass, Gulf of Manaar).
Depth range: 5-7m.

***Nymphon longicollum* HOEK 1881**

- 1881 *Nymphon longicollum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 40-41, pl. II, figs. 1-5, pl. XV, figs. 8-9.
 1913 *Nymphon longicollum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 72 [key].
 1923 *Nymphon longicollum*, - BOUVIER, Faune de France, **7**: 14 [key].
 1932 *Nymphon longicollum*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 106-109, figs. 7a-c, 8a.

Distribution: off the coast of Chile; New Zealand (E of Auckland).

Depth range: 2013-4072m.

***Nymphon longicoxa* HOEK 1881**

- 1881 *Nymphon longicoxa* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 38, 39, pl. 2, figs. 1-5, pl. XV, figs. 8-9.
 1908 *Nymphon longicoxa*, - HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1) (6): 165, pl. I, figs. 3, 3a.
 1913 *Nymphon longicoxa*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 72 [key].
 1923 *Nymphon longicoxa*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].
 1932 *Nymphon longicoxa*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 106-109, figs. 7a-c, 8a.
 1932 *Nymphon longicoxa*, - GORDON, Discovery Rep., **6**: 28, 32, 34.
 1935 *Nymphon longicoxa*, - GILTAY, Res. Voy Belgica 1897-99, Zool.: 5-7, figs. 1-4.
 1956 *Nymphon longicoxa*, - FAGE, Galathea Rep., **2**: 164.
 1956 *Nymphon longicoxa*, - FAGE, Galathea Rep., **2**: 168.
 1958 *Nymphon longicoxa*, - CLARK, Zool. Publ. Univ. Wellington, **23**: 1-2.
 1974 *Nymphon longicoxa*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **98**: 281-282.
 1976 *Nymphon longicoxa*, - ZHARKOVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **99**: 241-243, fig. 1.
 1982 *Nymphon longicoxa*, - CHILD, Smiths. Contr. Zool., **349**: 43.

Distribution: southeast of New Zealand; off Antarctica; Scotia Sea; Argentine Basin, off Rio de la Plata.

Depth range: 318-2578m.

Habitat: grey mud.

***Nymphon longimanum* G.O. SARS 1888**

- 1891 *Nymphon longimanum*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 93-95, pl. X, fig. 1a-f.

1956 *Nymphon longimanum*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **54**: 33-48, fig. 1.

Distribution: Arctica.
Depth range: 110m.

***Nymphon longisetosum* (HODGSON 1915)**

1927 *Chaetonymphon longisetosum*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 326-327, fig. 5.

Distribution: Antarctica.
Depth range: 385m.

***Nymphon longispinum* NAKAMURA & CHILD 1991**

1991 *Nymphon longispinum* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 53-55, fig. 20.

Distribution: Japan (northern Honshu).
Depth range: 700-950m.

***Nymphon longitarse* KRØYER 1844**

- 1844 *Nymphon longitarse* KRØYER, Naturhistorisk Tidskr., Kobenhavn, **1** (2): 112.
 1878 *Nymphon longitarse*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 19, pl. 7, fig. 2a-h.
 1880 *Nymphon longitarse*, - WILSON, Rep. U.S. Fish. Comm., **6**: 489-491, pl. 6, fig. 30.
 1891 *Nymphon longitarse*, - SARS, Norw. North-Atl. Exped., 1878-1878, **6**: 75-78, pl. VII, fig. 3a-h.
 1901 *Nymphon longitarse*, - WHITEAVES, Ann. Mag. Nat. Hist., **10** (4): 264.
 1908 *Nymphon longitarse*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 212-213.
 1916 *Nymphon longitarse*, - APPELLÖF, Kungl. Svensk. Vetensk. Handl., **54** (5): 4-5.
 1933 *Nymphon longitarse*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 67-68.
 1933 *Nymphon longitarse*, - STEPHENSEN, Medd. Grönland, **79** (6): 13-14, fig. 3 (map).
 1936 *Nymphon longitarse*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 862.
 1943 *Nymphon longitarse*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 86.
 1943 *Nymphon longitarse*, - NEEDLER, Can. Atl. Fauna, **10**: 9, fig. 9a-e.
 1943 *Nymphon longitarse*, - STEPHENSEN, Medd. Grönland, **121** (8): 21-22.
 1947 *Nymphon longitarse*, - OHSHIMA & KISHIDA, Illustr. Encyclopaedia Fauna Japan: 1006, fig. 2855.
 1948 *Nymphon longitarse*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 190, fig. 13b.
 1949 *Nymphon longitarse*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 247-248.
 1955 *Nymphon longitarse*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 214, 217.
 1961 *Nymphon longitarse*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 65.
 1963 *Nymphon longitarse*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1330-1331.

- 1971 *Nymphon longitarse*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 318.
1982 *Nymphon longitarse*, - CHILD, Smiths. Contr. Zool., **349**: 43.

Distribution: circum polar in boreal-arctic waters.

Depth range: 16-137m.

Habitat: mud; stones.

***Nymphon longituberculatum* OHLSEN 1913**

- 1913 *Nymphon longituberculatus* OHLSEN, Rep. Sci. Res. M. Sars N-Atl. Deep-Sea Exped. 1910, vol. 3 (pt. 1, Zool.): 5-6, figs. 5-9, pl. 1, fig. B.

Distribution: N-Atlantic.

Depth range: 83m.

***Nymphon macabou* MÜLLER 1990**

- 1990 *Nymphon macabou* MÜLLER, Bonn. zool. Beitr., **41** (3-4): 280-283, figs. 6-13.

Distribution: Martinique.

Depth range: 0-2m.

Habitat: dead corals on nearshore patch reef; exposed reef-flat and reef-margin.

***Nymphon macilentum* STOCK 1981**

- 1981 *Nymphon macilentum*, STOCK, Rés. Camp. MUSORSTOM, 1, Mém. ORSTOM, **91**: 314-315, figs. 1-11.

Distribution: Philippines.

Depth range: 188-200m.

***Nymphon macronyx* SARS 1877**

- 1881 *Nymphon macronyx* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 95-77, pl. XV, figs. 1-7.
1891 *Chaetonymphon macronyx*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 111-114, pl. XII, figs. 2a-k.
1902 *Nymphon macronyx*, - MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 181.
1933 *Chaetonymphon macronyx*, - STEPHENSEN, Medd. Grönland, **79** (6): 9-10.

Distribution: boreal Atlantic and Arctica.

Depth range: 300-1326m.

***Nymphon macrum* WILSON 1880**

- 1880 *Nymphon macrum* WILSON, Rep. U.S. Fish. Comm., **6**: 487-489, pl. 4, figs. 21-23.
1881 *Nymphon brevicollum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3**

- (10): 45-47, pl. III, figs. 13-15.
- 1891 *Nymphon macrum*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 89-91, pl. IX, fig. 2a-g.
- 1901 *Nymphon brevicollum*, - WHITEAVES, Ann. Mag. Nat. Hist., **10** (4): 263.
- 1901 *Nymphon macrum*, - WHITEAVES, Ann. Mag. Nat. Hist., **10** (4): 263.
- 1913 *Nymphon brevicollum*, - OLSEN, Rep. Sci. Res. M. Sars North Atl. Deep-Sea Exped., 1910, **3** (1) (Zool.): 5-6.
- 1933 *Nymphon macrum*, - STEPHENSEN, Medd. Grönland, **79** (6): 17-18.
- 1935 *Nymphon macrum*, - STEPHENSEN, Bergens Mus. Arbok, 1935, Naturvidensk. Rekke, **7**: 21-22.
- 1936 *Nymphon macrum*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 862.
- 1942 *Nymphon macrum*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 3.
- 1943 *Nymphon macrum*, - NEEDLER, Can. Atl. Fauna, **10**: 10, fig. 11a-e.
- 1948 *Nymphon macrum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 193-195, figs. 13d, 15.
- 1949 *Nymphon macrum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 248.
- 1963 *Nymphon macrum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1332.
- 1982 *Nymphon macrum*, - CHILD, Smiths. Contr. Zool., **349**: 43-44.
- 1986 *Nymphon macrum*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 419.

Distribution: NW-Atlantic, with southernmost records from Florida and Cuba.

Depth range: <100-1700m.

***Nymphon maculatum* CARPENTER 1910**

- 1910 *Nymphon maculatum* CARPENTER, J. Linn. Soc. London (Zool.), **31** (207): 256-257, pl. 26, figs. 1-9.
- 1964 *Nymphon maculatum*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **35**: 28.

Distribution: Red Sea (Aabak, Dahlak Archipelago).

Depth range: 0.5m.

Habitat: soft corals on reef.

***Nymphon maldivensis* CLARK 1961**

- 1961 *Nymphon maldivensis* CLARK, Ann. Mag. Nat. Hist., **3** (13): 291-293, figs. 1-7.

Distribution: Maldives Wadewaru Island, Fadiffolu Atoll).

Depth range: shallow water.

Habitat: from *Lithotamnion* encrusted dead *Acropora* coral in lagoon reef.

***Nymphon maoriana* CLARK 1958**

- 1958 *Nymphon maoriana* CLARK, Zool. Publ. Univ. Wellington, **23**: 2-3, figs. 1-8.

Distribution: New Zealand (Cook Strait).

Depth range: 37-73m.

Habitat: from kelp holdfasts; in shell with *Eupagurus novaezelandiae*.

Nymphon maruyamai NAKAMURA & CHILD 1991

1991 *Nymphon maruyamai* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 55-56, fig. 21.

Distribution: Japan.
Depth range: 410-440m.

Nymphon mauritanicum FAGE 1942

1942 *Nymphon mauritanicum* FAGE, Arch. Zool. Gen., **82** (Notes & Rev., 2): 77-80, figs. 1-3.

Distribution: W-Africa (Cap Blanc).
Depth range: 90-100m.

Nymphon megacheles CHILD 1988

1988 *Nymphon megacheles* CHILD, Smiths. Contr. Zool., **468**: 25-27, fig. 11.

Distribution: Philippines (Batan Island).
Depth range: 0.4m.
Habitat: "sea cliff and shallow caves" (CHILD).

Nymphon megalops G.O. SARS 1891

1891 *Nymphon megalops* SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 98-100, pl. X, fig. 3a-g.

1933 *Nymphon megalops*, - STEPHENSEN, Medd. Grönland, **79** (6): 19.

1943 *Nymphon megalops*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 87-88, fig. 2.

1963 *Nymphon megalops*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1333.

Distribution: Arctica, from Western Norway to Fox Basin, south to about 61°30'N in the Faroe Channel.

Depth range: 46-1444m.

Nymphon mendosum (HODGSON 1907)

1944 *Nymphon mendosum* GORDON, Rep. Brit. and N.Z. Antarct. Res. Exped., (B), **5** (1): 27-28.

1952 *Nymphon mendosum*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 263 [list], 265-267, fig. 1.

1972 *Nymphon mendosum*, - ARNAUD, Téthys, suppl. **4**: 150.

1980 *Nymphon mendosum*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 3-4.

Distribution: Antarctica.
Depth range: 180-219m.

Nymphon meridionale HOEK 1881

1881 *Nymphon meridionale* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 43-44, pl. III, figs. 4-8.

Distribution: Antarctica.

Depth range: 3065m.

Nymphon microcollis HILTON 1942

1942 *Nymphon microcollis* HILTON, J. Ent. Zool. Pomona College, **34** (1): 5.

1949 *Nymphon microcollis*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 273, fig. 34d.

Distribution: Alaska.

Nymphon microctenatus BARNARD 1946

1946 *Nymphon microctenatus* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 60.

1954 *Nymphon microctenatus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 95-96, fig. 4.

1965 *Nymphon microctenatus*, - STOCK, Beaufortia, **13** (151): 22-23 [list, key].

1988 *Nymphon microctenatus*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 166-167.

Distribution: South Africa.

Depth range: 357-650m.

Nymphon micronesicum CHILD 1982

1982 *Nymphon micronesicum* CHILD, Proc. biol. Soc. Wash., **95** (2): 277-280, fig. 3.

Distribution: Bikini Atoll (Namu islan).

Depth range: shallow water.

Habitat: coral reefs.

Nymphon micronyx SARS 1888

1891 *Nymphon micronyx* SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 90-93, pl. IX, fig. 3a-g.

1933 *Nymphon micronyx*, - STEPHENSEN, Medd. Grönland, **79** (6): 18.

1961 *Nymphon micronyx* ssp. n. *orientalis* LOSINA LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 65.

1963 *Nymphon micronyx*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1332.

Distribution: Kamtchatka; Greenland; American Arctic.

Depth range: 4-35m.

Nymphon micropedes HEDGPETH 1949

1949 *Nymphon micropedes* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 254-256, fig. 24.

1955 *Nymphon micropedes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 9-10, fig.

4.

- 1971 *Nymphon micropedes*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.
1983 *Nymphon micropedes*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 56.

Distribution: Japan.

Depth range: 924m.

***Nymphon microrhynchum* SARS 1888**

- 1891 *Nymphon microrhynchum*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 71-73, pl. VII, fig. 1a-g.
1902 *Nymphon microrhynchum*, - MÖBIUS, Fauna Arctica, **2**: 44-45.
1933 *Nymphon microrhynchum*, - STEPHENSEN, Medd. Grönland, **79** (6): 13.
1956 *Nymphon microrhynchum*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., 54: 33-48.
1963 *Nymphon microrhynchum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1330.

Distribution: Arctica.

Depth range: 30-200m.

***Nymphon microsetum* HILTON 1942**

- 1942 *Nymphon microsetum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 6.
1949 *Nymphon microsetum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 274, fig. 34h.

Distribution: Japan.

Depth range: 79-95m.

***Nymphon modestum* STOCK 1959**

- 1959 *Nymphon modestum* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 556-558, fig. 3.
1988 *Nymphon modestum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 167-168.

Distribution: South Africa.

Depth range: 23-550m.

Habitat: rock and mussels.

***Nymphon molleri* CLARK 1963**

- 1963 *Nymphon molleri* CLARK, Rec. Austr. Mus., **26** (1): 10-12, fig. 10A-H.
1968 *Nymphon cf. molleri*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 32, fig. 10h.
1973 *Nymphon molleri*, - STOCK, Beaufortia, **20** (266): 99.
1973 *Nymphon molleri*, - STOCK, Beaufortia, **21** (279): 91.

Distribution: Australia (Great Australian Bight; Western Port, Crawfish Rock; Port Jackson area).

Depth range: 10-112m.

Habitat: red algae; on *Amathaea* sp. (Bryozoa) and *Aglaophenia plumosa* (Hydroida); bryozoan gravel.

Nymphon molum HILTON 1942

- 1942 *Nymphon molum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 4.
 1949 *Nymphon molum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 271, fig. 34a.

Distribution: Japan.**Depth range:** 75m.***Nymphon multidens*** GORDON 1932

- 1932 *Nymphon multidens* GORDON, Discovery Rep., **6**: 75-76, figs. 34-35.
 1950 *Nymphon multidens*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 149-150.
 1990 *Nymphon multidens*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **126**:
 110-111, pl. 2, figs. 1-7.

Distribution: Antarctica (Bouvet Island, Palmer Peninsula).

Depth range: 40-45m.***Nymphon multituberculatum*** GORDON 1944

- 1944 *Nymphon multituberculatum* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B) **5** (1): 32-35, figs. 8a, 9a-d.
 1965 *Nymphon multituberculatum*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., **41** (31): 4.
 1980 *Nymphon multituberculatum*, - KRAPP, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 4.

Distribution: Antarctica.**Depth range:** 180-640m.***Nymphon natalense*** FLYNN 1928

- 1928 *Nymphon natalense* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., I): 11, fig. 3.
 1954 *Nymphon natalense*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 94.

Distribution: South Africa.***Nymphon neumayri*** GORDON 1932

- 1932 *Nymphon neumayri* GORDON, Discovery Rep., **6**: 66-69, figs. 27e, 29c-e, 30.
 1990 *Nymphon neumayri*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **126**:
 109-110, pl. 1, figs. 2-4.

Distribution: Antarctica (Palmer Archipelago).**Depth range:** 259-270m.***Nymphon nipponense*** HEDGPETH 1949

1949 *Nymphon nipponeense* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 267-269, fig. 32.

Distribution: Japan.

Depth range: 928m.

***Nymphon novaecaledoniae* STOCK 1991**

1991 *Nymphon novaecaledoniae* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 182-185, figs. 40-41.

Distribution: New Caledonia.

Depth range: 440-450m.

***Nymphon novaehollandiae* CLARK 1963**

1963 *Nymphon novaehollandiae* CLARK, Rec. Austr. Mus., **26** (1): 12-15, figs. 7A-I, 8A-C.

1968 *Nymphon novaehollandiae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 32.

Distribution: Australia (Sydney area; Great Australian Bight).

Depth range: 55-100m.

Habitat: gravel; bryozoan gravel.

***Nymphon nugax* STOCK 1966**

1966 *Nymphon nugax* STOCK, Atlantide Rep., **9**: 49-52, fig. 2.

Distribution: W-Africa (off Senegal).

Depth range: 41-55m.

Habitat: "brown sand, shells" (STOCK).

***Nymphon obesum* ARNAUD & CHILD 1988**

1988 *Nymphon obesum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 168-170, fig. 11.

Distribution: South Africa (off East London).

Depth range: 80m.

***Nymphon ochoticum* LOSINA-LOSINSKY, 1961**

1961 *Nymphon ochoticum* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 70-72, fig. 6.

Distribution: NW-Pacific.

Depth range: 515m.

Nymphon ohshimae HEDGPETH 1949

1949 *Nymphon ohshimae* HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 266-267, fig. 31.

Distribution: Japan.
Depth range: 997-1303m.

Nymphon okudai NAKAMURA & CHILD 1991

1991 *Nymphon okudai* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 56-58, fig. 22.

Distribution: Japan.
Depth range: 216-328m.

Nymphon orcadense (HODGSON 1908)

1908 *Chaetonymphon orcadense* HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1):
 1932 *Nymphon orcadense*, - GORDON, Discovery Rep., **6**: 63-65, fig. 26a.
 1944 *Nymphon orcadense*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 27.

Distribution: Antarctica and sub-antarctic areas (South Orkneys; ? South Georgia; Magellan District).
Depth range: 18-163m.

Nymphon ortmanni HELFER 1938

1938 *Nymphon ortmanni* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 164-167, fig. 1a-g.
 1953 *Nymphon ortmanni*, - STOCK, Beaufortia, **35** (4): 34-36, fig. 1.
 1954 *Nymphon ortmanni*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 20-21, fig. 6d-e.
 1955 *Nymphon ortmanni*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 10, fig. 5.
 1962 *Nymphon ortmanni*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 92.
 1971 *Nymphon ortmanni*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 320.
 1983 *Nymphon ortmanni*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 56.
 1987 *Nymphon ortmanni*, - NAKAMURA, Biol. Lab., Imperial Household: 6-7, pls. 4, 35.
 1991 *Nymphon ortmanni*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 58.

Distribution: Japan (Tsugaru Strait to Suruga Bay).
Depth range: 75-450m.
Habitat: sand bottom.

Nymphon paralobatum ARNAUD & CHILD 1988

1988 *Nymphon paralobatum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 170-173, fig. 12.

Distribution: South Africa (off Transkei coast).
Depth range: 710-775m.

Nymphon parasiticum MERTON 1906

- 1906 *Nymphon parasiticum* MERTON, Mitt. Zool. Stat. Neapel, **18** (1): 136-141, pl. 7, figs. 1-6.
1987 *Nymphon parasiticum*, - ARNAUD, Mesogee, **47**: 53.
1987 *Nymphon parasiticum*, - ARNAUD & BAMBER, Adv. in Mar. Biol., **24**: 52.

Distribution: Mediterranean.

Habitat: "ectoparasite du pied bouclier céphalique d'un gastropode Nudibranche, *Tethys leporina*" (ARNAUD).

Nymphon parum STOCK 1991

- 1991 *Nymphon parum* STOCK, in: A. CROSNIER (ed.), Rés. Camp. MUSORSTOM, 8. Mém. Mus. natn. Hist. nat., (A), **151**: 185, figs. 42-43.

Distribution: New Caledonia.

Depth range: 440-530m.

Nymphon paucidens GORDON 1932

- 1932 *Nymphon paucidens* GORDON, Discovery Rep., **6**: 48-51, figs. 17a, e, 19-20.

Distribution: Antarctica.

Depth range: 22-235m.

Nymphon paucituberculatum GORDON 1944

- 1944 *Nymphon paucituberculatum* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 30-32, fig. 7a-d, 8b.

Distribution: Antarctica (off Gauss Quadrant).

Depth range: 180-219m.

Nymphon pedunculatum ARNAUD & CHILD 1988

- 1988 *Nymphon pedunculatum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 173-176, fig. 13.

Distribution: South Africa (off East London).

Depth range: 90m.

Nymphon perlucidum HOEK 1881

- 1881 *Nymphon perlucidum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 52-53, pl. V, figs. 6-10.

Distribution: Indo-West Pacific (between Celebes and Halmahera).

Depth range: 1510m.

Habitat: rocks.

***Nymphon pfefferi* LOMAN 1923**

1923 *Nymphon pfefferi* LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 17-18, fig. B.

1932 *Nymphon pfefferi*, - GORDON, Discovery Rep., **6**: 46-48, figs. 16, 17b-d, 18a,-b, d.

Distribution: Antarctica.

Depth range: 17-250m.

***Nymphon phasmatodes* BÖHM 1879**

1879 *Nymphon phasmatodes* BÖHM, Monatsber. Kgl. Preuss. Akad. Wiss. Berlin, 1879: 173-174, pl. I, fig. 2.

1908 *Nymphon capense* HODGSON, Trans. Roy. Soc. Edinburgh, **46** (1) (6): 169-170, pl. I, figs. 2, 2a.

1913 *Nymphon capense*, - BOUVIER, Deuxième Expédition Antarctique (1908-1910), **6**: 72.

1923 *Nymphon capense*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14.

1928 *Nymphon capense*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 4.

1928 *Nymphon phasmatodes*, - FLYNN, Rep. Fish & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 4.

1928 *Nymphon phasmatodes*, - LOMAN, Bull. Soc. Sci. Nat. Phys. Maroc, **8** (1-3): 67, figs. 2, 6, 7.

1932 *Nymphon capense*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 117-120, figs. 11, 12a-c.

1954 *Nymphon capense*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 91-92.

1956 *Nymphon phasmatodes*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 76-78, fig. 2.

1959 *Nymphon phasmatodes*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 558.

1962 *Nymphon phasmatodes*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 283-284.

1965 *Nymphon phasmatodes*, - STOCK, Beaufortia, **13** (151): 22-23 [key].

1988 *Nymphon phasmatodes*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 176.

Distribution: South Africa.

Depth range: 13-90m.

Habitat: sand, broken shells and *Phyllochaetopterus*; shell and sand; fine khaki sand; shelly khaki sand; shell, coarse sand.

***Nymphon pilosum* MÖBIUS 1902**

1902 *Nymphon pilosum* MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 179, pl. XXIV, figs. 8-12.

- 1928 *Nymphon bipunctatum* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 8-10, figs. 1-2.
- 1954 *Nymphon pilosum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 100-101, fig. 6.
- 1956 *Nymphon pilosum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 76.
- 1962 *Nymphon pilosum*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 278.
- 1968 *Nymphon pilosum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 29.
- 1988 *Nymphon pilosum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 177.

Distribution: off South Africa.

Depth range: 20-850m.

Habitat: rocks, shells.

***Nymphon pixellae* SCOTT 1912**

- 1912 *Nymphon pixellae* SCOTT, Ann. Mag. Nat. Hist., **10** (8): 206-209, pl. 7, figs. 1-11.
- 1936 *Nymphon solitarium* EXLINE, Proc. U.S. natn. Mus., **83** (2991): 414-416, fig. 33a-d.
- 1939 *Nymphon solitarium*, - HEDGPETH, Am. Midl. Nat., **22** (2): 458-459.
- 1941 *Nymphon pixellae*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 255 [key], pls. 9-10.
- 1942 *Nymphon pixellae*, - HILTON, J. Ent. Zool. Pomona College, **34** (1): 4.
- 1942 *Nymphon variatum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 4.
- 1942 *Nymphon pixellae*, - HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 279-280, pl. 35.
- 1949 *Nymphon pixellae*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 246 [key].
- 1949 *Nymphon variatum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 271, fig. 34b.
- 1953 *Nymphon pixellae*, - HENRY, Microentomology, **18** (1): 17-18, figs. 1-5.
- 1955 *Nymphon pixellae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 216-217, fig. 1b-d.
- 1973 *Nymphon pixellae*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **91**: 178-179.
- 1979 *Nymphon pixellae*, - CHILD, Smiths. Contr. Zool., **293**: 40.
- 1984 *Nymphon pixellae*, - WICKSTEN, Calif. Fish and Game, **70** (3): 136.

Distribution: Mexico Pacific (Gulf of California); California; Japan.

Depth range: 28-128m.

Habitat: sandy mud; mud and sponges; mud and stones.

***Nymphon pleodon* STOCK 1962**

- 1962 *Nymphon pleodon* STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 281-282, figs. 5-6.

Distribution: South Africa.

Depth range: 27m.

Habitat: rocks.

***Nymphon polare* (HODGSON 1915)**

- 1915 *Chaetonymphon polare* HODGSON, Ann. Mag. Nat. Hist., ser. 8, **15**: 143.

1927 *Chaetonymphon polare*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 324-326.

Distribution: Antarctica.

Depth range: 350m.

***Nymphon primacoxa* STOCK 1968**

1968 *Nymphon primacoxa* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 29-31, fig. 11.

Distribution: Mozambique Channel.

Depth range: 1600-1628m.

***Nymphon proceroides* BOUVIER 1913**

1913 *Nymphon proceroides* BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 90-94, figs. 42-48.

1932 *Nymphon proceroides*, - GORDON, Discovery Rep., **6**: 36-37, figs. 9, 10a, 75.

1944 *Nymphon proceroides*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 22.

1990 *Nymphon proceroides*, - TURPAEVA, Trudy Inst. Okeanol. Akad. Nauk SSSR, **126**: 111-113, pl. 3, figs. 1-9.

Distribution: Antarctica (Palmer Archipelago, South Shetlands).

Depth range: 160-420m.

***Nymphon procerum* HOEK 1881**

1881 *Nymphon procerum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 39, pl. II, figs. 9-12.

1913 *Nymphon procerum*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 72 [key].

1923 *Nymphon procerum*, - LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 14 [key].

1932 *Nymphon procerum*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 112-113, fig. 9a.

1932 *Nymphon procerum*, - GORDON, Discovery Rep., **6**: 34 [key].

1944 *Nymphon procerum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 19.

1951 *Nymphon procerum*, - FAGE, Rep. Swed. Deep-Sea Exped., **2** (7): 95-97, figs. 1-2.

1965 *Nymphon procerum*, - STOCK, Beaufortia, **13** (151): 22 [list].

1971 *Nymphon procerum*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **92**: 276-277, fig. 2.

1973 *Nymphon procerum*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **91**: 178.

1974 *Nymphon procerum*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **98**: 281.

1981 *Nymphon procerum*, - STOCK, Cah. Biol. Mar., **22**: 460, fig. 3.

1982 *Nymphon procerum*, - CHILD, Smiths. Contr. Zool., **349**: 44.

1990 *Nymphon procerum*, - TURPAEVA, Trudy Inst. Okeanol. Akad. Nauk SSSR, **126**: 108-109.

Distribution: North and South Pacific; North and South Atlantic.

Depth range: 2450-6135m.

Nymphon profundum HILTON 1942

- 1942 *Nymphon profundum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 3.
1942 *Nymphon noctum* HILTON, J. Ent. Zool. Pomona College, **34** (1): 3.
1949 *Nymphon noctum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 270-271, fig. 33a-f.
1949 *Nymphon profundum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 270.
1965 *Nymphon profundum*, - STOCK, Beaufortia, **13** (151): 22.
1991 *Nymphon profundum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 58.

Distribution: Japan.

Depth range: 3232-4220m.

Nymphon prolatum FAGE 1942

- 1942 *Nymphon prolatum* FAGE, Arch. Zool. Gen., **82** (Notes & Rev., 2): 80-82, figs. 4-5.
1966 *Nymphon prolatum*, - STOCK, Atlantide Rep., **9**: 48-49.

Distribution: W-Africa (off Senegal; off Dakar).

Depth range: 65-89m.

Nymphon proximum CALMAN 1915

- 1915 *Nymphon proximum* CALMAN, Rep. Brit. Antarct. "Terra Nova" Exped., Zool. **3**: 34-36, fig. 6.
1944 *Nymphon proximum* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 28-29, fig. 6a-b.

Distribution: Antarctica (Ross Sea area).

Depth range: 300-603m.

Nymphon puellula KRAPP 1973

- 1973 *Nymphon puellula* KRAPP, Beaufortia, **21** (277): 64, 68.
1987 *Nymphon puellula*, - ARNAUD, Mesogee, **47**: 53.

Distribution: Mediterranean.

Depth range: 35m.

Nymphon pumillum NAKAMURA & CHILD 1991

- 1991 *Nymphon pumillum* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 58-60, fig. 23.

Distribution: Japan (SE of Akkeshio Bay, Hokkaido).
Depth range: 102m.

Nymphon quadriclavus NAKAMURA & CHILD 1986

- 1986 *Nymphon quadriclavus* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 893-895,
 figs. 1-2.
 1986 *Nymphon quadriclavus*, - NAKAMURA, Biol. Lab., Imp. Household: 7-8, pls. 5-6,
 figs. 1-3

Distribution: Japan (Sagami Bay).
Depth range: 450m.

Nymphon quadrispinum (HILTON 1942)

- 1949 *Nymphon quadrispinum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 274,
 fig. 33m.

Distribution: Alaska (Kiska Harbor).
Depth range: 35m.

Nymphon residuum STOCK 1971

- 1971 *Nymphon residuum* STOCK, Zool. Meded., **47**: 257-259, figs. 1-8.

Distribution: off Kenya.
Depth range: 1510m.
Habitat: pure globigerina-ooze.

Nymphon robustum BELL 1853

- 1881 *Nymphon robustum*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3**
 (10): 97-98, fig.
 1899 *Nymphon robustum*, - MEINERT, Danish Ingolf Exped., **3** (1): 45-46, pl. III,
 figs. 23-24.
 1943 *Boreonymphon robustum*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 83.
 1963 *Nymphon robustum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1325.

Distribution: Arctica, perhaps circumpolar.
Depth range: 2-2000m.
Habitat: may live on *Umbellula encrinus* and other
 octocorals.

Nymphon rottnesti CHILD 1975

- 1975 *Nymphon rottnesti* CHILD, Smiths. Contr. Zool., **190**: 5-7, fig. 3.

Distribution: W-Australia (West of Rottnest Island).
Depth range: 137-143m.

Nymphon sandersi CHILD 1982

1982 *Nymphon sandersi* CHILD, Smiths. Contr. Zool., **349**: 44-46, fig. 14.

Distribution: Argentine Basin, off Rio de la Plata.
Depth range: 3815-4435m.

Nymphon sarsi MEINERT 1899

1899 *Nymphon sarsi* MEINERT, Danish Ingolf Exped., 3 (1): 38-39, pl. III, figs. 1-6.

Distribution: subarctic area.
Depth range: 480m.
Habitat: clay and gravel.

Nymphon schimkewitschi LOSINA-LOSINSKY 1929

1956 *Nymphon schimkewitschi*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **54**: 33-48, fig. 2.

Distribution: Russian part of Arctica.

Nymphon schmidti LOSINA-LOSINSKY 1961

1961 *Nymphon schmidti* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 63-65, fig. 3.

Distribution: NW-Pacific.
Depth range: 130m.

Nymphon serratidentatum ARNAUD & CHILD 1988

1988 *Nymphon serratidentatum* ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 177-180, fig. 14.

Distribution: South Africa (off Transkei coast).
Depth range: 710-775m.

Nymphon serratum G.O. SARS 1879

1891 *Nymphon serratum*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3): 95-97, pl. X, fig. 2a-h.
1933 *Nymphon serratum*, - STEPHENSEN, Medd. Grönland, **79** (6): 18-19.
1942 *Nymphon serratum*, - GILTAY, J. Fish. Res. Board Canada, **5**: 459.
1943 *Nymphon serratum*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 87.
1963 *Nymphon serratum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1333.

Distribution: An arctic species, known in the N-Atlantic

from Siberia to Greenland and the Canadian Atlantic Coast (Hudson bay).

Depth range: 45-1444m.

Habitat: rocky bottom.

***Nymphon setimanus* BARNARD 1946**

- 1946 *Nymphon setimanus* BARNARD, Ann. Mag. Nat. Hist., **13** (11): 61.
 1954 *Nymphon setimanus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 103-1205, fig. 8.
 1962 *Nymphon setimanus*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 278.
 1965 *Nymphon setimanus*, - STOCK, Beaufortia, **13** (151): 18-19, figs. 13-17.
 1989 *Nymphon setimanus*, - MÜLLER, Senckenbergiana marit., **20** (5/6): 287-288, figs. 35-41.

Distribution: Gulf of Aden; Madagascar; South Africa.

Depth range: 0-751m.

Habitat: shelly sand; on algaee.

***Nymphon setipedes* CHILD 1988**

- 1988 *Nymphon setipedes*, Bull. biol. Soc. Wash., **8**: 72-75, fig. 8.

Distribution: Belize (Carrie Bow Cay).

Depth range: 24.4m.

Habitat: "fore-reef slope" (CHILD).

***Nymphon signatum* MÖBIUS 1902**

- 1902 *Nymphon signatum* MÖBIUS, Wiss. Ergebn. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 178-179, pl. XXIV, figs. 1-7.
 1932 *Nymphon signatum*, - GORDON, Ann. Mag. Nat. Hist., **9** (10): 104-105, fig. 5a-b.
 1932 *Nymphon signatum*, - GORDON, Discovery Rep., **6**: 28-29, 35 [key].
 1954 *Nymphon signatum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 96-98, fig. 5a.

Distribution: South Africa (Agulhas Bank, off Cape St. Blaize, off Gt. Fish Point, off Glendower Beacon).

Depth range: 100m.

***Nymphon similis* Child, 1992**

- 1992 *Nymphon similis* CHILD, Smiths. Contr. Zool., **526**: 31-33, fig. 13.

Distribution: Ecuador.

Depth range: 25m.

***Nymphon simulatum* NAKAMURA & CHILD 1991**

- 1991 *Nymphon simulatum* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 60-61, fig. 24.

Distribution: Japan (eastern Honshu).
Depth range: 1950-2000m.

Nymphon singulare STOCK 1954

- 1954 *Nymphon singulare* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 21-25,
figs. 7-8.
1963 *Nymphon singulare*, - CLARK, Rec. Austr. Mus., **26** (1): 7-9, 15.

Distribution: Australia (disaster Bay, New South Wales).
Depth range: 55-92m.
Habitat: sand, mud.

Nymphon sluiteri HOEK 1877

- 1891 *Nymphon sluiteri*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3):
73-75, pl. VII, fig. 2a-g.
1916 *Nymphon sluiteri*, - APPELLÖF, Kungl. Svensk. Vetensk. Handl., **54** (5): 5-6.
1933 *Nymphon sluiteri*, - STEPHENSEN, Medd. Grönland, **79** (6): 14-15, fig. 4 (map).
1943 *Nymphon sluiteri*, - HEDGPETH, J. Wash. Acad. Sci., **33**: 86-87.
1963 *Nymphon sluiteri*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1331.

Distribution: circumpolar arctic.
Depth range: 2-1000m.
Habitat: mud; stones.

Nymphon soyoi UTINOMI 1955

- 1955 *Nymphon soyoi* UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 13-15, fig. 8.
1971 *Nymphon soyoi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.

Distribution: Japan.
Depth range: 188m.

Nymphon spicatum CHILD 1982

- 1982 *Nymphon spicatum* CHILD, Smiths. Contr. Zool., **349**: 46-48, fig. 15.

Distribution: Argentine Basin, off Rio de la Plata.
Depth range: 3815-3822m.

Nymphon spiniventris STOCK 1953

- 1953 *Nymphon spiniventris* STOCK, Temminckia, **9**: 282-285, figs. 5-6.

Distribution: East Indian Archipelago.
Depth range: 17m.

Nymphon spinosissimum (NORMAN 1894)

- 1891 *Chaetonymphon spinosum*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 107-109, pl. XI, fig. 3a-i.
- 1894 *Chaetonymphon spinosissimum* NORMAN, Ann. Mag. Nat. Hist., **13** (6): 154.
- 1908 *Chaetonymphon spinosissimum*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 219-220.
- 1930 *Chaetonymphon spinosissimum*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nau SSSR, **2**: 335-336.
- 1933 *Chaetonymphon spinosissimum*, - STEPHENSEN, Medd. Grönland, **79** (6): 6-8, fig. 2 (map).
- 1943 *Chaetonymphon spinosissimum*, - STEPHENSEN, Medd. Grönland, **121** (8): 14-15, fig. 4.
- 1948 *Nymphon spinosissimum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 183, figs. 10a, 11a.
- 1963 *Nymphon spinosissimum*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1327.
- 1982 *Nymphon spinosissimum*, - CHILD, Smiths. Contr. Zool., **349**: 48.

Distribution: Norway to western Greenland; Labrador Sea; rare in Arctica.

Depth range: 62-862m.

Habitat: sand, clay.

Nymphon stipulum CHILD 1990

- 1990 *Nymphon stipulum* CHILD, Beaufortia, **41** (6): 40-43, fig. 1.

Distribution: southern California.

Depth range: 395m.

Nymphon stocki UTINOMI 1955

- 1955 *Nymphon stocki* UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 10-13, figs. 6-7.
- 1971 *Nymphon stocki*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 321.

Distribution: Japan.

Depth range: 324m.

Nymphon stocki TURPAEVA 1974

- 1974 *Nymphon stocki* TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **98**: 282-283, pl. 3, figs. 1-8.

Distribution: Scotia Sea.

Remarks: homonym of *N. stocki* UTINOMI 1955.

Nymphon striatum LOSINA-LOSINSKY 1929

- 1929 *Nymphon striatum* LOSINA-LOSINSKY, Zool. Jb. (Syst.), **57**: 538-539, fig. 1.
- 1933 *Nymphon striatum*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 64-66, fig. 13.

- 1954 *Nymphon striatum*, - UTINOMI, Publ. Akkeshi Mar. Biol. Stat., **3**: 2.
 1967 *Nymphon striatum*, - NESIS, Issledov. Fauny Morei, **5** (13): 248.
 1971 *Nymphon striatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 319.
 1987 *Nymphon striatum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 160.
 1991 *Nymphon striatum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 61.

Distribution: Sea of Japan; Gulf of Tataray.

Depth range: littoral-212m.

***Nymphon stromi* KRØYER 1844**

- 1844 *Nymphon strömi* KRØYER, Naturhistorisk Tidskr., Kobenhavn, **1** (2): 111.
 1872 *Nymphon giganteum* WHITEAVES, Ann. Mag. nat. Hist., **10** (4): 349.
 1874 *Nymphon giganteum*, - VERRILL, Am. J. Sci. & Arts, **3** (7): 411.
 1878 *Nymphon Strömi*, - WILSON, Trans. Conn. Acad. Arts Sci., **5**: 17-18, pl. 6,
 fig. 1a-h.
 1880 *Nymphon Strömi*, - WILSON, Rep. U.S. Fish. Comm., **6**: 483-487, pl. 5, pl. 6,
 fig. 29.
 1881 *Nymphon Strömi*, - WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 253.
 1885 *Nymphon Strömi*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 561.
 1891 *Nymphon strömii*, - SARS, Norw. North-Atl. Exped., 1876-1878, Zool., **6** (3):
 80-83, pl. VIII, fig. 2a-k.
 1916 *Nymphon strömi*, - APPELLÖF, Kungl. Svensk. Vetensk. Handl., **54** (5): 8-13,
 figs. 1-4.
 1933 *Nymphon strömi*, - STEPHENSEN, Medd. Grönland, **79** (6): 16-17.
 1943 *Nymphon stroemii*, - NEEDLER, Can. Atl. Fauna, **10**: 9, fig. 10a-d.
 1948 *Nymphon strömi*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 190-192, fig.
 13c.
 1963 *Nymphon stromi*, - HEDGPETH, J. Fish. Res. Board Can., **20** (5): 1331-1332.
 1975 *Nymphon stromi*, - STOCK, Bull. mar. Sci., **24** (4): 994.
 1982 *Nymphon stromi*, - CHILD, Smiths. Contr. Zool., **349**: 48.
 1983 *Nymphon stroemii*, - BAMBER, Zool. J. Linn. Soc., **77**: 68.
 1986 *Nymphon stromi*, - STOCK, Bull. mar. Sci., **38** (3): 403 tab. 1.

Distribution: widely distributed in the boreal arctic area.

Depth range: 15->1000m.

Habitat: mud stones.

***Nymphon subtile* LOMAN 1923**

- 1923 *Nymphon subtile* LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 19-
 21, fig. C.
 1932 *Nymphon subtile*, - GORDON, Discovery Rep., **6**: 43-45, figs. 14, 15, 24b.

Distribution: Falkland Islands.

Depth range: 22m.

***Nymphon surinamense* STOCK 1975**

- 1975 *Nymphon surinamense* STOCK, Bull. mar. Sci., **24** (4): 1007-1010, figs. 21-22.
 1986 *Nymphon surinamense*, - STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 420.

Distribution: continental shelf of northern South America, from Venezuela (Isla Margarita) to the Guyanas.

Depth range: 26-91m.

***Nymphon tanypalpes* CHILD 1988**

1988 *Nymphon tanypalpes* CHILD, Bull. biol. Soc. Wash., **8**: 75-77, fig. 9.

Distribution: Seychelles (Aldabra Atoll).

Depth range: 0.5-21m.

Habitat: "Reef slope coral rubble in hollows between coral heads"; "mangrove roots in lagoon" (CHILD).

***Nymphon tenellum* (SARS 1888)**

1881 *Nymphon pallenoides*, - WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 254.

1885 *Nymphon pallenoides*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 561.

1888 *Chaetonymphon tenellum* SARS, Arch. Math. Naturvidensk., **4**: 353.

1891 *Chaetonymphon tenellum*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 109-111, pl. XII, fig. 1a-h.

1933 *Chaetonymphon tenellum*, - STEPHENSEN, Medd. Grönland, **79** (6): 10.

1948 *Nymphon tenellum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 185, fig. 10c.

1982 *Nymphon tenellum*, - CHILD, Smiths. Contr. Zool., **349**: 49.

Distribution: both sides of N-Atlantic.

Depth range: 1501-2452m.

***Nymphon tenuipes* BOUVIER 1911**

1913 *Nymphon tenuipes*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 86-90, figs. 35-41.

1932 *Nymphon tenuipes*, - GORDON, Discovery Rep., **6**: 37-39, figs. 10c, 11.

1944 *Nymphon tenuipes*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 22.

Distribution: Antarctica (South Shetlands; Palmer Archipelago).

Depth range: 122-220m.

***Nymphon tuberculare* LOSINA-LOSINSKY, 1961**

1961 *Nymphon tuberculare* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 60-62, fig. 2.

Distribution: NW-Pacific.

Depth range: 130m.

***Nymphon tubiferum* STOCK 1978**

1978 *Nymphon tubiferum* STOCK, Cah. Biol. Mar., **19**: 208-211, fig. 8.

Distribution: NE-Atlantic Basin.

Depth range: 740m.

Habitat: soft mud, sandy and granular, broken shells.

***Nymphon typhlops* (HODGSON 1915)**

1927 *Chaetonymphon typhlops*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 327-329, fig. 6.

Distribution: Antarctica.

Depth range: 2450m.

***Nymphon unguiculatum* HODGSON 1927**

1927 *Nymphon unguiculatum* HODGSON, Dt. Südpolar-Exped., **XIX**, Zool. XI: 318-319.

1974 *Nymphon unguiculatum*, - TURPAEVA, Trudy Inst. Okeanol. Akad. Nauk SSSR, **98**: 280-281.

Distribution: Scotia Sea.

Depth range: 330-450m.

***Nymphon uniunguiculatum* LOSINA-LOSINSKY 1933**

1933 *Nymphon uniunguiculatum* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., **17**: 62-64, 78-79, fig. 12.

1949 *Nymphon uniunguiculatum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 263, fig. 29.

1961 *Nymphon uniunguiculatum*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 72.

1987 *Nymphon uniunguiculatum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.

Distribution: NW-Pacific.

***Nymphon vilosum* (HODGSON 1915)**

1927 *Chaetonymphon villosum*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 323-324.

1952 *Nymphon villosum*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (3): 267-269, fig. 2.

1980 *Nymphon villosum*, - KRAPP; Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 4 [text].

Distribution: Antarctica.

Depth range: 385m.

***Nymphon vulcanellum* STOCK 1992**

1992 *Nymphon vulcanellum* STOCK, Tijdschr. Ent., **135**: 124 [text], 126-128, 139, figs. 52-61.

Distribution: southern Brazil.
Depth range: 21-250m.

Nymphon vulsum STOCK 1986

1986 *Nymphon vulsum* STOCK, Bull. mar. Sci., **38** (3): 402 tab. 1, 406 [text], 419-420, fig. 6b-j.

Distribution: Venezuela (Isla Margarita).
Depth range: 26-27 m.

Nymphon walvisense STOCK 1981

1981 *Nymphon walvisense* STOCK, Cah. Biol. Mar., **22**: 455-460, figs. 1-2.

Distribution: Walvis Basin.

Pentanymphon HODGSON 1904

1904 *Pentanymphon antarcticum* HODGSON, Ann. Mag. nat. Hist., **7** (14): 459 [diagnosis].

Pentanymphon antarcticum HODGSON 1904

- 1904 *Pentanymphon antarcticum* HODGSON, Ann. Mag. nat. Hist., **14** (7): 459, pl. XIV.
 1932 *Pentanymphon antarcticum*, - GORDON, Discovery Rep., **6**: 24-26.
 1938 *Pentanymphon antarcticum* var. *gaussi* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 169-170, fig. 3a-c.
 1944 *Pentanymphon antarcticum*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 14.
 1950 *Pentanymphon antarcticum*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 150 [lit., syn.].
 1952 *Pentanymphon antarcticum*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (3): 263 [list], 264.
 1953 *Pentanymphon antarcticum* var. *gaussi*, - STOCK, Beaufortia, **35** (4): 37.
 1965 *Pentanymphon antarcticum*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., **41** (31): 2.
 1972 *Pentanymphon antarcticum*, - ARNAUD, Téthys, suppl. **4**: 149, figs. 1-2.
 1980 *Pentanymphon antarcticum*, - KRAPPA, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 4.

Distribution: Antarctica.
Depth range: 163-640m.

Pentanymphon minutum GORDON 1944

1944 *Pentanymphon minutum* GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res.

- Exped., (B), **5** (1): 14-16, fig. 3a-f.
 1952 *Pentanymphon minutum*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (3):
 264-265.

Distribution: Antarctica.

Depth range: 163-180m.

***Sexanymphon* HEDGPETH & FRY 1964**

- 1964 *Sexanymphon* HEDGPETH & FRY, Ann. Mag. Nat. Hist., **13** (7): 161 [diagnosis].

***Sexanymphon mirabilis* HEDGPETH & FRY 1964**

- 1964 *Sexanymphon mirabilis* HEDGPETH & FRY, Ann. Mag. Nat. Hist., **13** (7): 161-169,
 figs. 1-4.
 1974 *Sexanymphon mirabilis*, - TURPAEVA, Trudy Inst. Okeanol, Akad. Nauk SSSR, **98**:
 284.

Distribution: Antarctica; Scotia Sea.

Depth range: 2760-2810m.

PHOXICHILIDIIDAE

- 1948 Phoxichilidiidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 216
 [diagnosis, remarks, key to NW-Atlantic genera].
 1945 Phoxichilidae, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 144-146.
 1954 Phoxichilidiidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 125 [diagnosis, key
 to South African genera].
 1954 Phoxichilidiidae, - STOCK, Vidensk. Medd. Dansk naturh. Foren., 116: 68-71
 [remarks].
 1974 Endeidae, - KING, A Synopsis of the British Sea Spiders. Synopses of the
 British Fauna (New Ser.), **5**: 48.
 1992 Phoxichilidiidae, - CHILD, Mem. Hourglass Cruises, **9** (1): 37-38, tab. 5.
 1992 Endeididae, - CHILD, Mem. Hourglass Cruises, **9** (1): 56, tab. 8.

***Anoplodactylus* WILSON 1878**

- 1945 *Anoplodactylus*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 154-155.
 1948 *Anoplodactylus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 218-222
 [diagnosis, remarks, key to NW-Atlantic species].
 1954 *Anoplodactylus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 126 [diagnosis,
 remarks, key to South African species].
 1963 *Anoplodactylus*, - CLARK, Rec. Austr. Mus., **26** (1): 55-56 [key to Australian
 species].
 1975 *Anoplodactylus*, - STOCK, Bull. mar. Sci., **24** (4): 1075-1076 [remarks on
pygmaeus-complex].
 1979 *Anoplodactylus*, - CHILD, Smiths. Contr. Zool., 293: 47-48 [key to Middle
 American species].
 1979 *Anoplodactylus*, - STOCK, Bijdr. Dierk., **48** (2): 158 [list of valid
Anoplodactylus species], 158-159 [list of species inquirenda, homonyms,
 synonyms and alternative names].

- 1983 *Anoplodactylus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 41-42 [key to Japanese species].
 1992 *Anoplodactylus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 39 [diagnosis].

***Anoplodactylus aculeatus* MÖBIUS 1902**

- 1902 *Anoplodactylus aculeatus* MÖBIUS, Wiss. Ergebn. dtsch. Tiefsee-Exped. 1898-1899, 3: 188, pl. XXVIII, figs. 1-7.
 1954 *Anoplodactylus aculeatus*, - BARNARD, Ann. S. Afr. Mus., **XLI** (3): 127-128.

Distribution: South Africa.

Depth range: 126m.

***Anoplodactylus allotrius* CHILD 1979**

- 1979 *Anoplodactylus allotrius* CHILD, Smiths. Contr. Zool., **293**: 48-50, fig. 16.
 1989 *Anoplodactylus allotrius*, - STOCK, Bijdr. Dierk., **59** (2): 94.

Distribution: Panamá Caribbean; Curaçao.

Depth range: 3-4m.

Habitat: Acanthophora; coarse coral sand.

***Anoplodactylus anarthrus* LOMAN 1908**

- 1908 *Anoplodactylus anarthrus* LOMAN, Siboga Exped. Monogr., **40**: 76-77, pl. IV, figs. 40-45.
 1990 *Anoplodactylus anarthrus*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 74.
 1990 *Anoplodactylus anarthrus*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk., **45** (10): 106-109, figs. 7-13.

Distribution: Kenya; Indonesia (Timor); Fiji Islands.

Depth range: 0-0.5m.

Habitat: dead corals covered with algae; dead corals on reef flat.

***Anoplodactylus angulatus* (DOHRN 1881)**

- 1881 *Phoxichilidium angulatum* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: 184-188, pl. XII, figs- 1-12.
 1929 *Anoplodactylus angulatus*, - GILTAY, Bull. & Ann. Roy. Soc. Ent. Belg., **69**: 173.
 1945 *Anoplodactylus angulatus*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 155-157.
 1958 *Anoplodactylus angulatus*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
 1968 *Anoplodactylus angulatus*, - STOCK, Vie et Milieu, **19** (1A): 28, fig. 22.
 1972 *Anoplodactylus angulatus*, - KING, J. nat. Hist., **6**: 622.
 1973 *Anoplodactylus angulatus*, - KRAPP, Beaufortia, **21** (277): 72, fig. 6.
 1974 *Anoplodactylus angulatus*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 42-43, fig. 17.
 1978 *Anoplodactylus angulatus*, - CHIMENTZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRIONZATO, Mem. Biol. Mar. Ocean., **8** (4): 96.
 1979 *Anoplodactylus angulatus*, - CHIMENTZ, BRIGNOLI & BASCIANO, Cah. Biol. Mar.,

20: 479.

- 1987 *Anoplodactylus angulatus*, - ARNAUD, Mesogee, **47**: 44-45.
 1990 *Anoplodactylus angulatus*, - STOCK, Zool. Meded., **63** (16): 229.
 1992 *Anoplodactylus angulatus*, - STOCK, Bijdr. Dierk., **62** (2): 94.
 1993 *Anoplodactylus angulatus*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24,
 tab. 3.

Distribution: Mediterranean; England, Ireland and Wales;
 North Sea; Portugal; Azores; Canary Islands.

Depth range: 0-25m.

Habitat: shallow water algae in sheltered locations (*Halopteris scoparia*, *Dictyopteris* et *Jania rubens*); *Eunicella cavolini*; *Corallina mediterranea*; *Cystoseira stricta*; in *Posidonia oceanica* seagrass; rocky and sandy coasts; on intertidal stones; muddy tidal flats with stones and algae.

***Anoplodactylus aragaoi* SAWAYA 1950**

- 1950 *Anoplodactylus aragaoi* SAWAYA, Mem. Inst. Oswaldo Cruz, **47** (1-2): 63-67, 77-
 80, pls. I-II.
 1992 *Anoplodactylus aragaoi*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: Brazil.

Depth range: 35-38m.

Habitat: among Hydrozoa, Bryozoa and algae.

***Anoplodactylus arcuatus* CHILD 1977**

- 1977 *Anoplodactylus arcuatus* CHILD, Proc. biol. Soc. Wash., **90** (3): 584-587, fig.
 1.
 1992 *Anoplodactylus arcuatus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab.
 5], 39 [key], 41, fig. 17.
 1992 *Anoplodactylus arcuatus*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 45-47,
 figs. 4-14.

Distribution: Barbados, Curaçao, Florida (Dry Tortugas),
 Texas.

Depth range: 2.5-55m.

Habitat: various coral reef locations.

***Anoplodactylus arescus* Du BOIS-REYMOND MARCUS 1959**

- 1959 *Anoplodactylus arescus* du BOIS-REYMOND MARCUS 1959, Kieler Meeresforsch. **15**
 (1): 105-107, pl. 1.
 1968 *Anoplodactylus arescus*, - STOCK, Vidensk. Medd. Dansk Naturhist. Foren, **131**:
 53.
 1973 *Anoplodactylus arescus*, - ARNAUD, Tethys, **4** (4): 954.
 1975 *Anoplodactylus arescus*, - Trav. Mus. Hist. Nat. "Grigore Antipa", **16**: 133,
 figs. 10-12.
 1988 *Anoplodactylus arescus*, - CHILD, Smiths. Contr. Zool., **468**: 12.
 1988 *Anoplodactylus arescus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4):
 810.

Distribution: Madaascar; Tanzania; Red Sea; Philippines.
Depth range: 0-5m.
Habitat: algae; in sand.

Anoplodactylus arnaudae STOCK 1978

- 1978 *Anoplodactylus arnaudae* STOCK, Cah. Biol. Mar., **19**: 217-218, fig. 10.
 1984 *Anoplodactylus arnaudae*, - STOCK, J. mar. biol. Ass. U.K.: 749.
 1987 *Anoplodactylus arnaudae*, - STOCK, Cah. Biol. Mar., **28**: 514.
 1991 *Anoplodactylus arnaudi*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 139.

Distribution: NE-Atlantic.
Depth range: 330-1350m.

Anoplodactylus australis (HODGSON 1914)

- 1914 *Phoxichilidium australe* HODGSON, Zool. Anz., **45** (4): 161.
 1915 *Phoxichilidium australe*, - CALMAN, Brit. Antarctic Exp., 1910, Zool., **3** (1): 46-48, fig. 10A-D.
 1915 *Phoxichilidium australe*, - HODGSON, Ann. Mag. Nat. Hist., (8) **15** (85): 145.
 1927 *Phoxichilidium australe*, - HODGSON, Dtsch. Südpolar-Exp., **19** (Zool. II): 340.
 1932 *Phoxichilidium australe*, - GORDON, Discovery Rep., **6**: 92.
 1938 *Phoxichilidium australe*, - Sci. Rep. Australasian Antarctic Exp. (C), (Zool. & Bot.), **2** (8): 19.
 1944 *Phoxichilidium australe*, - Rep. British, Australian and New Zealand Antarct. Res. Exp., (B) **5** (1): 48.
 1952 *Phoxichilidium australe*, - FAGE, Bull. Mus. Natn. Hist. Nat., Paris, (2) **24** (2): 184.
 1952 *Phoxichilidium australe*, - FAGE, Bull. Mus. Natn. Hist. Nat., Paris, (2) **24** (3): 271.
 1955 *Anoplodactylus australis*, - STOCK, Res. Norw. Sci. Exped. Tristan da Cunha 1937-1938, **33**: 4.

Distribution: circumpolar in the Antarctic, northernmost records from Tasmania and Tristan da Cunha.

Depth range: 0-549m.

Anoplodactylus baldarus CHILD 1988

- 1988 *Anoplodactylus baldarus* CHILD, Bull. Biol. Soc. Wash., **8**: 56-59, fig. 2.

Distribution: Seychelles (Aldabra Atoll).

Depth range: 0-55m.

Habitat: "Reef crest"; "Seaward of reef crest, sand in coral hollows" (CHILD); also among coralline algae.

Anoplodactylus bahamensis CHILD 1977

- 1977 *Anoplodactylus bahamensis* CHILD, Proc. biol. Soc. Wash., **90** (3): 587-589, fig. 2.
 1982 *Anoplodactylus bahamensis*, - CHILD, Smiths. Contr. mar. Sci., **12**: 368.
 1986 *Anoplodactylus bahamensis*, - STOCK, Bull. mar. Sci., **38** (3): 403.

Distribution: Caribbean Sea. - Bahamas and Belize.

Depth range: 0.5-12m.

Habitat: tidal flats of coral reefs.

***Anoplodactylus batangensis* (HELFER 1938)**

- 1938 *Pycnosoma batangense* HELFER, Sitz.-Ber. naturforsch. Freunde Berlin, 1937: 174-176, fig. 6a-c.
 1940 *Pycnosoma batangense*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 47-48.
 1948 *Anoplodactylus stylirostris* HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 232-234, fig. 33.
 1949 *Anoplodactylus tenuirostris* LEBOUR, Proc. Zool. Soc. London, **118** (4): 929-930, fig. 1.
 1953 *Anoplodactylus batangense*, - STOCK, Beaufortia, **35** (4): 39-41, fig. 4a-e.
 1954 *Anoplodactylus batangense*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 127-128.
 1955 *Anoplodactylus batangense*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 595-598, pl. II, figs. 4-8.
 1968 *Anoplodactylus batangensis*, - STOCK, Medd. dansk. naturhist. Foren., **131**: 54.
 1973 *Anoplodactylus batangensis*, - ARNAUD, Tethys, **4** (4) (1972): 957, figs. 3-4.
 1974 *Anoplodactylus batangensis*, - STOCK, Bull Zoöl. Mus. Univ. Amsterdam, **4** (3): 17.
 1975 *Anoplodactylus batangensis*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 191.
 1975 *Anoplodactylus batangensis*, - STOCK, Trav. Mus. Hist. Nat. "Grigore Antipa", **16**: 133.
 1975 *Anoplodactylus batangensis*, - STOCK, Bull. mar. Sci., **24** (4): 1082-1083, fig. 43c-d.
 1977 *Anoplodactylus batangensis*, - CHILD, Proc. biol. Soc. Wash., **90** (2): 444.
 1979 *Anoplodactylus batangensis*, - CHILD, Smiths. Contr. Zool., **293**: 50.
 1979 *Anoplodactylus batangensis*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 27.
 1982 *Anoplodactylus batangensis*, - CHILD, Smiths. Contr. mar. Sci., **12**: 368.
 1988 *Anoplodactylus batangensis*, - CHILD, Smiths. Contr. Zool., **468**: 14.
 1989 *Anoplodactylus batangensis*, - STOCK, Bijdr. Dierk., **59** (2): 95.
 1990 *Anoplodactylus batangensis*, - CHILD, Proc. biol. Soc. Wash., **103** (82): 311-335.
 1990 *Anoplodactylus batangensis*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 283-284.
 1992 *Anoplodactylus batangensis*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 41-42, fig. 18.
 1992 *Anoplodactylus batangensis*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 47.

Distribution: circumtropical.

Depth range: 0-40 m.

Habitat: in a variety of substrata, often in seagrass (*Thalassia*) and coral rubble.

Anoplodactylus bourboni MÜLLER 1990

1990 *Anoplodactylus bourboni* MÜLLER, Zool. Abh. Staat. Mus. Tierk., **45** (9): 98-100, figs. 1-6.

Distribution: La Réunion.

Depth range: 0.5-1m.

Habitat: crest and lagoon of fringing reef, from dead corals.

Anoplodactylus bova CHILD 1979

1979 *Anoplodactylus bova* CHILD, Smiths. Contr. Zool., **293**: 50-52, fig. 17.

Distribution: Panamá Pacific.

Depth range: low intertidal.

Anoplodactylus brasiliensis HEDGPETH 1948

1940 *Anoplodactylus pygmaeus*, - MARCUS, Bolm. Fac. Fil. Cienc. Letr., Univ. S. Paulo, 4: 63-65, Pl. VI fig. A-D.

1992 *Anoplodactylus brasiliensis*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: Brazil.

Depth range: infralittoral.

Habitat: seaweeds on rocky shore.

Anoplodactylus brevicollis LOMAN 1908

1908 *Anoplodactylus brevicollis* LOMAN, Siboga Exped. Monogr., **40**: 74-75, pl. III, figs. 29-32.

Distribution: SE-Asia.

Depth range: 13m.

Habitat: sand and shells.

Anoplodactylus brevirostris CHILD 1988

1988 *Anoplodactylus brevirostris* CHILD, Smiths. Contr. Zool., **468**: 14-16, fig. 6.

1992 *Anoplodactylus brevirostris*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 163-164.

Distribution: Philippines (Batan Island); Malaysia (Pulau Tioman).

Depth range: 1-3 m.

Habitat: dead corals.

Anoplodactylus brucei CHILD 1990

1990 *Anoplodactylus brucei* CHILD, Proc. biol. Soc. Wash., **103** (2): 327-330, fig.

6.

1992 *Anoplodactylus brucei*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 166 [text].

Distribution: Australian barrier reef.

Depth range: 6.2m.

Habitat: coral reefs. - rubble.

***Anoplodactylus bruuni* Child, 1992**

1992 *Anoplodactylus bruuni* CHILD, Smiths. Contr. Zool., **526**: 35-37, fig. 15.

Distribution: Chile (Islas Juan Fernandez).

Depth range: 0-180m.

Habitat: "rocks and sand" (CHILD).

***Anoplodactylus californicus* HALL 1912**

1849 ?*Nymphon dubium*, - NICOLET, in: GAY, Hist. fis. polit. Chile, 3. Atlas zool., Crustacéos: 307.

1912 *Anoplodactylus californicus* HALL, First Ann. Rep. Laguna Mar. Lab., **1**: 91.

1913 *Anoplodactylus californicus*, - HALL, Univ. Calif. Publ. Zool., **11** (6): 129.

1915 *Anoplodactylus californicus*, - HILTON, J. Ent. Zool. Pomona College, **7** (1): 69.

1915 *Anoplodactylus californicus*, - HILTON, J. Ent. Zool. Pomona College, **7** (3): 201, 205.

1920 *Anoplodactylus californicus*, - HILTON, J. Ent. Zool. Pomona College, **12** (4): 93.

1927 *Anoplodactylus portus* CALMAN, Trans. Zool. Soc. London, **22** (3): 405-408, fig. 103.

1939 *Anoplodactylus californicus*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 29.

1939 *Anoplodactylus robustus*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 28-29.

1940 *Anoplodactylus californicus*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 40 [key].

1941 *Anoplodactylus californiensis*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 257 [key], pl. 11.

1942 *Anoplodactylus californicus*, - HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 277 [list], 288-291, pl. 39.

1942 *Anoplodactylus californicus*, - HILTON, J. Ent. Zool. Pomona College, **34** (3): 72.

1942 *Anoplodactylus projectus* HILTON, Occ. Pap. Bernice P. Bishop Mus., **17** (3): 45-47, fig. 3a-b.

1943 *Anoplodactylus carvalhoi*, - HEDGPETH, Proc. New England Zool. Club, **22**: 46.

1948 *Anoplodactylus carvalhoi*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 230-232, fig. 30e-g.

1950 *Anoplodactylus portus*, - SAWAYA, Mem. Inst. Oswaldo Cruz, **47** (1-2): 70 [key].

1954 *Anoplodactylus portus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 128.

1955 *Anoplodactylus carvalhoi*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 592.

1955 *Anoplodactylus portus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 238-239.

- 1958 *Anoplodactylus portus*, - STOCK, Bull. Res. Council Israel, **7B**: 140-141, fig. 2.
- 1958 *Anoplodactylus portus*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
- 1962 *Anoplodactylus portus*, - STOCK, Ent. Ber., **22**: 218.
- 1971 *Anoplodactylus portus*, - LIPKIN & SAFRIEL, J. Ecol., **59** (1): 9.
- 1975 *Anoplodactylus portus*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 201.
- 1975 *Anoplodactylus portus*, - STOCK, Bull. mar. Sci., **24** (4): 1052-1053.
- 1976 *Anoplodactylus portus*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser.: 158.
- 1978 *Anoplodactylus portus*, - CHILD, Zool. J. Linn. Soc., **63** (1-2): 133-144, figs. 1-4 [gynandromorphism].
- 1979 *Anoplodactylus portus*, - CHILD, Smiths. Contr. Zool., **293**: 58.
- 1979 *Anoplodactylus portus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 15.
- 1982 *Anoplodactylus portus*, - CHILD, Smiths. Contr. mar. Sci., **12**: 373.
- 1987 *Anoplodactylus portus*, - ARNAUD, Mesogee, **47**: 46.
- 1987 *Anoplodactylus californicus*, - CHILD, Proc. biol. Soc. Wash., **100** (3): 544-555.
- 1990 *Anoplodactylus californicus*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 284.
- 1992 *Anoplodactylus californicus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 43, fig. 19.
- 1992 *Anoplodactylus californicus*, - CHILD, Smiths. Contr. Zool., **526**: 37.

Distribution: tropical and temperate N-Atlantic, NE-Pacific south to Chile; Mediterranean.

Depth range: 0-<100m.

Habitat: in a variety of substrata, often in algal vegetation.

***Anoplodactylus calliopus* STAPLES 1982**

- 1982 *Anoplodactylus calliopus* STAPLES, Mem. Qd. Mus., **20** (3): 459-461, fig. 3A-K.

Distribution: Australia (Queensland).

Depth range: 1-5.6m.

***Anoplodactylus carnatus* NAKAMURA & CHILD 1983**

- 1983 *Anoplodactylus carnatus* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 42-44, fig. 14.
- 1991 *Anoplodactylus carnatus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 25.

Distribution: Japan.

Depth range: 5-15 m.

***Anoplodactylus chamorus* CHILD 1983**

- 1983 *Anoplodactylus chamorus* CHILD, Proc. biol. Soc. Wash., **96** (4): 705-707, fig. 3.
- 1988 *Anoplodactylus chamorus*, - Smiths. Contr. Zool., **468**: 16.
- 1990 *Anoplodactylus chamorus*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 330-331.
- 1991 *Anoplodactylus chamorus*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 143.
- 1992 *Anoplodactylus chamorus*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 164.

Distribution: Philippines; Guam; Australian barrier reef; Malaysia (Pulau Babi Besar, Tioman Archipelago).

Depth range: 0-12 m.

Habitat: dead corals; coralline and other algae.

***Anoplodactylus compactus* (HILTON 1939)**

- 1939 *Phoxichilidium compactum* HILTON, J. Ent. Zool. Pomona College, **31** (2): 27-28.
 1939 *Phoxichilidium compactum*, - HILTON, J. Ent. Zool. Pomona College, **31** (4): 72.
 1940 *Halosoma compactum*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 45-46 [list].
 1941 *Phoxichilidium compactum*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 257.
 1942 *Halosoma compactum*, - HILTON, J. Ent. Zool. Pomona College, **34** (3): 74.
 1954 *Halosoma compactum*, - HEDGPETH, in: R.I. SMITH et al. (Eds.), Intertidal Invertebrates of the Central California coast: 205.
 1975 *Anoplodactylus compactus*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 191-193, fig. 1.

Distribution: California (Laguna Beach, Monterey Bay).

Depth range: intertidal.

***Anoplodactylus compositus* CHIMENZ, COTTARELLI & TOSTI 1991**

- 1991 *Anoplodactylus compositus* CHIMENZ, COTTARELLI & TOSTI, Fragm. Ent., **23** (1): 7-12, figs. 1-9.

Distribution: Mediterranean Sea (Apulian coast).

Depth range: 21m.

Habitat: "Coralligenous Biocenosis" (CHIMENZ, COTTARELLI & TOSTI).

***Anoplodactylus coxalis* STOCK 1968**

- 1968 *Anoplodactylus coxalis* STOCK, Vidensk. Meddr. dansk naturh. Foren., **131**: 54-56, fig. 20.

Distribution: Kermadec Islands (Raoul Island).

Depth range: 75-85m.

Habitat: gravel, stones.

***Anoplodactylus crassus* NAKAMURA & CHILD 1988**

- 1986 *Anoplodactylus viridintestinalis*, - KIM, Proc. Nat. Sci. Res. Inst. KANU, **2** (1): 3-5.
 1986 *Anoplodactylus viridintestinalis*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 44.
 1987 *Anoplodactylus viridintestinalis*, - HONG & KIM, Jor. J. Syst. Zool., **3** (2):

- 161.
- 1988 *Anoplodactylus crassus* NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 810-813, fig. 1.
- 1988 *Anoplodactylus viridintestinalis*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 812-813 [text], fig. 2a-f.
- 1991 *Anoplodactylus crassus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 25.

Distribution: American Samoa, Korea and Japan.
Depth range: 3m.

Anoplodactylus cribellatus CALMAN 1923

- 1923 *Anoplodactylus cribellatus* CALMAN, Rec. Ind. Mus., **24** (3): 285-287, fig. 12a-e.

Distribution: Andam Islands.
Depth range: 4-17m.

Anoplodactylus dentimanus STOCK 1979

- 1923 *Anoplodactylus virescens*, - BOUVIER, Faune de France, **7**: 41, fig. 37.
- 1945 *Anoplodactylus virescens*, - LEBOUR, J. Mar. Biol. Assoc. U.K., **26**: 150.
- 1974 *Anoplodactylus virescens*, - ARNAUD, Tethys **5** (4): 784, figs. 6-9.
- 1979 *Anoplodactylus dentimanus* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **29**: 24-26, fig. 8.

Distribution: Indian Ocean (St. Paul and Amsterdam Islands).

Anoplodactylus derjugini (LOSINA-LOSINSKY 1929)

- 1933 *Halosoma derjugini*, - LOSINA-LOSINSKY, Isledov. Morei S.S.S.R. Leningrad, **17**: 47-49, pl. 2, figs. a-e.

Distribution: Pacific U.S.S.R.

Anoplodactylus digitatus (BÖHM 1879)

- 1879 *Phoxichilidium (Anoplodactylus) digitatum* BÖHM, Monatsber. Königl. Preuss. Akad. Wiss. Berlin, 1879: 184-185, pl. II, figs. 2a-b.
- 1908 *Anoplodactylus digitatus*, - LOMAN, Siboga Exp. Monogr., **40**: 74-76, pl. II, figs. 25-28.
- 1923 *Anoplodactylus saxatilis* CALMAN, Rec. Indian Mus., **24** (3): 287-288, fig. 13.
- 1923 *Anoplodactylus investigatoris* CALMAN, Rec. Indian Mus., **24** (3): 288-289, fig. 14.
- 1927 *Anoplodactylus saxatilis*, - CALMAN, Trans. Zool. Soc. London, **22** (3): 405.
- 1954 *Anoplodactylus investigatoris*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 85-86, figs. 37a-b, 38b.
- 1955 *Anoplodactylus investigatoris*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 593-594, pl. II, figs. 1-3.
- 1958 *Anoplodactylus saxatilis*, - STOCK, Bull. Res. Coun. Israel, **7B**: 139-140,

fig. 1a-c.

- 1958 *Anoplodactylus investigatoris*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
- 1965 *Anoplodactylus digitatus*, - STOCK, Beaufortia, **13** (151): 28-29.
- 1968 *Anoplodactylus digitatus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 49.
- 1974 *Anoplodactylus cf. digitatus*, - ARNAUD, Bull. Zool. Mus. Univ. Amsterdam., **3** (21): 174.
- 1987 *Anoplodactylus digitatus*, - ARNAUD, Mesogee, **47**: 45.
- 1992 *Anoplodactylus digitatus*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 164-166, figs. 18-26.
- 1992 *Anoplodactylus digitatus*, - STOCK, Bijdr. Dierk., **62** (2): 94.

Distribution: almost worldwide in tropical and subtropical waters. - tropical Indo-Pacific (Malaysia (Tioman Archipelago, Pulau Babi Besar; Oman; Singapore), Caribbean and Mediterranean).

Depth range: 0-2 m.

Habitat: dead corals and under stones.

Anoplodactylus erectus COLE 1904

- 1904 *Anoplodactylus erectus* COLE, Harriman Alaska Exped., **10**: 289-291, pl. 14 fig. 12, pl. 26 figs. 1-9.
- 1913 *Anoplodactylus erectus*, - HALL, Univ. Calif. Publ. Zool., **11** (6): 133.
- 1916 *Anoplodactylus erectus*, - HILTON, J. Ent. Zool. Pomona College, **8** (1): 25-34, figs. 1-6.
- 1934 *Anoplodactylus erectus*, - SCHMITT, J. Wash. Acad. Sci., **24** (1): 69.
- 1939 *Anoplodactylus erectus*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 28.
- 1939 *Anoplodactylus erectus*, - HILTON, J. Ent. Zool. Pomona College, **31** (4): 72, pl. II, fig. 9.
- 1941 *Anoplodactylus erectus*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 257 [key].
- 1942 *Anoplodactylus erectus*, - HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 283-286, pl. 37.
- 1942 *Anoplodactylus erectus*, - HILTON, J. Ent. Zool. Pomona College, **34** (3): 72.
- 1955 *Anoplodactylus erectus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren, **117**: 239-243, figs. 13-14.
- 1970 *Anoplodactylus erectus*, - CHILD, Proc. biol. Soc. Wash., **83** (27): 288.
- 1975 *Anoplodactylus erectus*, - STOCK, Ent. Ber., **35**: 74-76, figs. 22-23.
- 1976 *Anoplodactylus erectus*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser., EPA-600/3-76-028: 158.
- 1979 *Anoplodactylus erectus*, - CHILD, Smiths. Contr. Zool., **293**: 52-53.
- 1986 *Anoplodactylus erectus*, - HONG & KIM, Kor. J. Syst. Zool., **2** (2): 41-42, fig. 5.
- 1987 *Anoplodactylus erectus*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.
- 1988 *Anoplodactylus erectus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 813.
- 1989 *Anoplodactylus erectus*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 126, figs. 41-44.
- 1990 *Anoplodactylus erectus*, - CHILD, Proc. biol. Soc. Wash., **103** (1): 157.
- 1990 *Anoplodactylus erectus*, - MÜLLER, Senckenbergiana biol., **70** (1989) (1/3): 194-195, figs. 28-34.
- 1991 *Anoplodactylus erectus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 25-26.

1992 *Anoplodactylus erectus*, - CHILD, Smiths. Contr. Zool., **526**: 37.

Distribution: Pacific coast of North-America (Canada to Panamá), French Polynesia (Society Archipelago; Tuamotu Archipelago), Samoa, Korea and Japan; Chile (Isla San Felix); Ecuador.

Depth range: 0-40m.

Habitat: coral reef locations; sand.

***Anoplodactylus eroticus* STOCK 1968**

1968 *Anoplodactylus eroticus* STOCK, Vidensk. Medd. dansk naturhist. Foren, **131**: 49-51, fig. 18.

Distribution: India (Pamban Pass, Gulf of Manaar); Hawaii (Honolulu).

***Anoplodactylus evansi* CLARK 1963**

1963 *Anoplodactylus evansi* CLARK, Rec. Austr. Mus., **26** (1): 51-53, fig. 26A-G.

1973 *Anoplodactylus evansi*, - STOCK, Beaufortia, **20**: 121.

Distribution: Australia (Tasmania; West Island).

Depth range: 0-13m.

Habitat: on *Acrocarpia* (Phaeophyta); on seaweed attached to rock; on hydroids.

***Anoplodactylus evelinae* MARCUS 1940**

1940 *Anoplodactylus evelinae* MARCUS, Bolm. Fac. Fil. Cién, Letr. Univ. S. Paulo, ser. **19** (Zool.), 4: 55-58, fig. 4.

1948 *Anoplodactylus evelinae*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 232, fig. 31.

1950 *Anoplodactylus evelinae*, - SAWAYA, Mem. Inst. Oswaldo Cruz, **47** (1-2): 72.

1954 *Anoplodactylus (Labidodactylus) evelinae*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **24**: 128.

1975 *Anoplodactylus (Labidodactylus) evelinae*, - STOCK, Bull. mar. Sci., **24** (4): 1083.

1979 *Anoplodactylus evelinae*, - CHILD, Smiths. Contr. Zool., **293**: 53.

1979 *Anoplodactylus evelinae*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 27.

1982 *Anoplodactylus evelinae*, - CHILD, Smiths. Contr. mar. Sci., **12**: 368.

1986 *Anoplodactylus evelinae*, - STOCK, Bull. mar. Sci., **38** (3): 404.

1992 *Anoplodactylus evelinae*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: in W-Atlantic from Florida to Brazil; Kongo and Pacific Coast of Panamá.

Depth range: 0-30m.

Habitat: in a variety of substrata.

Anoplodactylus excelsus NAKAMURA & CHILD 1983

- 1983 *Anoplodactylus excelsus* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 44-46,
fig. 15.
1991 *Anoplodactylus excelsus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 26,
27 [key].

Distribution: Japan (Sagami Bay).
Depth range: 37-416m.

Anoplodactylus falciclavus CHILD 1988

- 1988 *Anoplodactylus falciclavus* CHILD, Smiths. Contr. Zool., **468**: 16-18, fig. 7.

Distribution: Philippines (Negros Island).
Depth range: littoral.

Anoplodactylus galetensis CHILD 1979

- 1979 *Anoplodactylus galetensis* CHILD, Smiths. Contr. Zool., **293**: 54-55, fig. 18.

Distribution: Panamá Caribbean.
Depth range: subtidal-6m.

Anoplodactylus gestiens (ORTMANN 1890)

- 1891 *Phoxichilidium gestiens* ORTMANN, Zool. Jb. (Syst.), **5** (1): 166, pl. XXIV,
fig. 8.
1911 *Anoplodactylus gestiens*, - LOMAN, Abh. Kayserl. Bayr. Akad. Wiss. (Math. -
Naturwiss. Kl.), Suppl. **2** (4): 13.
1919 *Anoplodactylus gestiens*, FUKUI, Rigakukai(?), **16** (2): 98.
1936 *Anoplodactylus gestiens*, - OHSHIMA, Dōbutsugaku Zasshi, **48** (8-10): 864.
1949 *Anoplodactylus gestiens*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 284,
fig. 40a-d.
1954 *Anoplodactylus gestiens*, - STOCK, Vidensk. Medd. dansk. naturhist. Foren.,
116: 71-72, fig. 31a-b.
1962 *Anoplodactylus gestiens*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 96.
1965 *Anoplodactylus gestiens*, - UTINOMI, New Illustr. Encycl. Fauna Japan, **2**:
335.
1971 *Anoplodactylus gestiens*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 325-
326.
1982 *Anoplodactylus gestiens*, - CHILD & NAKAMURA, Proc. biol. Soc. Wash.,
95 (2): 292-295, fig. 1 [gynandromorphism].
1983 *Anoplodactylus gestiens*, - NAKAMURA & CHILD 1983, Smiths. Contr. Zool., **386**:
46.
1987 *Anoplodactylus gestiens*, - NAKAMURA, Biol. Lab. Imperial Household: 15-16,
pl. 12.
1990 *Anoplodactylus gestiens*, - TURPAYEVA, Zool. Zh., **69** (9): 18-19, fig. 1 (4-
5).

Distribution: Japan.
Depth range: 0-479m.

Habitat: *Zostera*-belt; hard bottom.

Anoplodactylus gibbifemoris TURPAEVA 1991

1991 *Anoplodactylus gibbifemoris* TURPAEVA, - Zool. Zh., **70** (3): 151-154, figs. 1-7.

Distribution: SE-Pacific (Naska ridge, underwater mountain).

Anoplodactylus glandulifer STOCK 1954

- 1954 *Anoplodactylus glandulifer* STOCK, Vidensk. Medd. Dansk naturhist. Foren., **116**: 80-84, fig. 36.
 1958 *Anoplodactylus glandulifer*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 3.
 1968 *Anoplodactylus glandulifer*, - STOCK, Vidensk. Meddr. dansk naturhist. Foren. **131**: 49.
 1973 *Anoplodactylus glandulifer*, - ARNAUD, Tethys, **4** (4): 955, figs. 1-2.
 1974 *Anoplodactylus glandulifer*, - STOCK, Bull. Zoöl. Mus. Univ. Amsterdam, **4** (3): 16.
 1977 *Anoplodactylus multiclavus* CHILD, Proc. biol. Soc. Wash., **90** (3): 593, fig. 4.
 1979 *Anoplodactylus multiclavus*, - CHILD, Smiths. Contr. Zool., **293**: 58.
 1982 *Anoplodactylus glandulifer*, - CHILD, Proc. biol. Soc. Wash., **95** (2): 273-274.
 1982 *Anoplodactylus multiclavus*, - CHILD, Smiths. Contr. mar. Sci., **12**: 372.
 1986 *Anoplodactylus multiclavus*, - STOCK, Bull. mar. Sci., **38** (3): 404, 439.
 1988 *Anoplodactylus glandulifer*, - CHILD, Bull. biol. Soc. Wash., **8**: 59.
 1988 *Anoplodactylus glandulifer*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 813.
 1990 *Anoplodactylus glandulifer*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 331.
 1990 *Anoplodactylus glandulifer*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 74.
 1991 *Anoplodactylus glandulifer*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 143-144.
 1992 *Anoplodactylus dauphinus* CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab 5, key], 43-46, fig. 20.
 1992 *Anoplodactylus glandulifer*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 166, figs. 27-30.
 1992 *Anoplodactylus glandulifer*, - STOCK, Bijdr. Dierk., **62** (2): 94-95.

Distribution: tropical Indo-Pacific and Caribbean Sea.

Depth range: 0-78 m.

Habitat: in a variety of substrata.

Anoplodactylus haswelli (FLYNN 1918)

- 1918 *Halosoma haswelli* FLYNN, Pap. Proc. Roy. Soc. Tasmania, 1918: 11-15, pl. I, figs. 1-6.
 1954 *Anoplodactylus haswelli*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 86-88, fig. 40a-b.
 1963 *Anoplodactylus haswelli*, - CLARK, Rec. Austr. Mus., **26** (1): 48-49, fig.

24A-D.

1982 *Anoplodactylus haswelli*, - STAPLES, Mem. Qd. Mus., **20** (3): 461.

Distribution: Australia; New Zealand.

Depth range: intertidal.

Habitat: amongst mussels; from weeds on rocky coast; rocks.

***Anoplodactylus hwanghaensis* KIM & HONG 1986**

1986 *Anoplodactylus hwanghaensis* KIM & HONG, Kor. J. Syst. Zool., **2** (2): 41-44, fig. 6.

1987 *Anoplodactylus hwanghaensis*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.

1991 *Anoplodactylus hwanghaensis*, - Smiths. Contr. Zool., **512**: 27.

Distribution: Japan and Korea.

Depth range: intertidal.

***Anoplodactylus imperialis* NAKAMURA & CHILD 1986**

1983 *Anoplodactylus* sp. NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 54, fig. 18.

1986 *Anoplodactylus imperialis* NAKAMURA & CHILD, Zool. Sci., **3**: 895-897, fig. 3.

1987 *Anoplodactylus imperialis*, - NAKAMURA, Biol. Lab. Imperial Household: 16, pl. 13.

1991 *Anoplodactylus imperialis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 26 [key], 27.

Distribution: Japan (Sagami Bay).

Depth range: 38-50 m.

***Anoplodactylus imswe* CHILD 1982**

1982 *Anoplodactylus imswe* CHILD, Smiths. Contr. mar. Sci., **12**: 369-371, fig. 166.

Distribution: Belize (Tobacco Reef; Carrie Bow Cay).

Depth range: 0.5m.

Habitat: tidal flats.

***Anoplodactylus inermis* LOSINA-LOSINSKY, 1961**

1961 *Anoplodactylus inermis* LOSINA-LOSINSKY, Issledov. Dalnevost. Morei SSSR, Leningrad, **7**: 80-82, fig. 12.

Distribution: NW-Pacific.

Depth range: 414m.

***Anoplodactylus insigniformis* STOCK 1975**

1975 *Anoplodactylus insigniformis* STOCK, Bull. mar. Sci., **24** (4): 1058-1063, figs. 45-47.

- 1979 *Anoplodactylus insigniformis*, - CHILD, Smiths. Contr. Zool., **293**: 54.
 1979 *Anoplodactylus insigniformis*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 15.
 1986 *Anoplodactylus insigniformis*, - STOCK, Bull. mar. Sci., **38** (3): 404.

Distribution: Caribbean Coast of Panamá, Jamaica, Aruba, Curaçao, Venezuela.

Depth range: 0-7m.

Habitat: in a variety of substrata, often among fouling on pilings.

***Anoplodactylus insignis* (HOEK 1881)**

- 1881 *Phoxichilidium insigne* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 82-84, pl. 14, figs. 5-7.
 1904 *Anoplodactylus insignis bermudensis* COLE, Boston Soc. nat. Hist., **31** (8): 325-327, pl. 20, figs. 1-3.
 1940 *Anoplodactylus insignis bermudensis*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 40.
 1940 *Anoplodactylus insignis*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 58-60.
 1943 *Anoplodactylus insignis*, - HEDGPETH, Proc. New England Zool. Club, **22**: 45.
 1948 *Anoplodactylus insignis*, - HEDGPETH, Proc. U.S. natn. Mus., **3216**: 226-228, fig. 28d-g
 1954 *Anoplodactylus insignis*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 127.
 1955 *Anoplodactylus insignis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 235.
 1973 *Anoplodactylus insignis*, - KRAEUTER, J. nat. Hist., **7**: 494
 1975 *Anoplodactylus insignis*, - STOCK, Bull. mar. Sci., **24** (4): 1056-1058, fig. 44
 1976 *Anoplodactylus insignis*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27** (3/4) 337.
 1986 *Anoplodactylus insignis*, - STOCK, Bull. mar. Sci., **38** (3): 437.
 1986 *Anoplodactylus insignis* var. *calcaratus*, - STOCK, Bull. mar. Sci., **38** (3): 437-438, fig. 15a-b [n. var.].
 1992 *Anoplodactylus insignis*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 46-49, fig. 21, 76 [tab. 12], 78 [tab. 14].

Distribution: Virginia to Brazil and Bermuda.

Depth range: 0-90m.

Habitat: Intertidal estuaries.

***Anoplodactylus iuleus* STOCK 1975**

- 1975 *Anoplodactylus iuleus* STOCK, Bull. mar. Sci., **24** (4): 1069-1072, figs. 51-52.
 1976 *Anoplodactylus iuleus*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27** (3/4): 337.
 1986 *Anoplodactylus iuleus*, - STOCK, Bull. mar. Sci., **38** (3): 438-439.

Distribution: Florida and Cuba.

Depth range: 218-476m.

***Anoplodactylus jonesi* CHILD 1974**

- 1974 *Anoplodactylus jonesi* CHILD, Proc. biol. Soc. Wash., **87** (43): 497-500, fig. 2.
- 1975 *Anoplodactylus(?) antillanus* STOCK, Bull. mar. Sci., **24** (4): 1081-1082, fig. 57.
- 1979 *Anoplodactylus jonesi*, - CHILD, Smiths. Contr. Zool., **293**: 56, fig. 19a-b.
- 1982 *Anoplodactylus jonesi*, - CHILD, Smiths. Contr. mar. Sci., **12**: 371.
- 1986 *Anoplodactylus jonesi*, - Bull. mar. Sci., **38** (3): 404.

Distribution: Florida, Bonaire, Caribbean Coast of Panamá and Belize.

Depth range: 0-100m.

Habitat: in a variety of substrata; *Halimeda* on coral reefs.

***Anoplodactylus jungersi* FAGE 1949**

- 1949 *Anoplodactylus jungersi* FAGE, Arch. Zool. Exp. Gen., **86** (Notes & Rev.), 2: 25-27, fig. 2.

Distribution: Congo Belge.

***Anoplodactylus justi* MÜLLER 1992**

- 1992 *Anoplodactylus justi* Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 48-50, figs. 15-19.

Distribution: Barbados.

Depth range: 54m.

Habitat: "sand and fine coral rubble, low sponges" (MÜLLER).

***Anoplodactylus krappi* MÜLLER 1990**

- 1990 *Anoplodactylus krappi* MÜLLER, Bonn. zool. Beitr., **41** (1): 74-77, figs. 27-33.

Distribution: Kenya (Watamu area).

Depth range: 0-0.5m.

Habitat: dead corals covered with algae.

***Anoplodactylus lagenus* NAKAMURA & CHILD 1983**

- 1983 *Anoplodactylus lagenus* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 48-49, fig. 16.
- 1987 *Anoplodactylus lagenus*, - NAKAMURA, Biol. Lab. Imperial Household: 17, pl. 14.
- 1991 *Anoplodactylus lagenus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 27.

Distribution: Japan (Sagami Bay).
Depth range: 37-40m.

Anoplodactylus laminifer ARNAUD 1974

1974 *Anoplodactylus laminifer* ARNAUD, Tethys, 5 (4): 782-784, figs. 1-5.

Distribution: Southern Indian Ocean (St. Paul and Amsterdam Islands).

Depth range: 80-100m.

Anoplodactylus lentus WILSON 1878

- 1821 *Anaphia pallida* SAY 1821, J. Acad. Nat. Sci., 2 (1): 59-61, pl. 5, fig. 7.
 1873 *Phoxichilidium maxillare* VERRILL, Rep. U.S. Comm. Fish., 1: 250, pl. 7, fig. 35.
 1878 *Anoplodactylus lentus* WILSON, Am. J. Sci. Arts, 3 (15): 200.
 1878 *Anoplodactylus lentus*, - WILSON, Trans. Conn. Acad. Arts & Sci., 5: 14-15, pl. 4, fig. 3a-c.
 1880 *Anoplodactylus lentus*, - WILSON, Rep. U.S. Fish. Comm., 6: 482-483, pl. 3, figs. 16-18.
 1891 *Phoxichilidium maxillare*, - MORGAN, Stud. Biol. Lab. Johns Hopkins Univ., Baltimore, 5 (1): (embryology).
 1901 *Anoplodactylus lentus*, - COLE, Biol. Bull. Woods Hole Mar. Biol. Lab., 2 (5): 195-207 (habits).
 1908 *Anaphia lenta*, - NORMAN, J. Linn. Soc. London (Zool.), 30: 204.
 1913 *Anoplodactylus lentus*, - SUMNER, OSBURN & COLE, Bull. Bureau Fish., Washington, 31 (1, 2): 142, 677, chart 121 (distribution map).
 1925 *Anoplodactylus lentus*, - FISH, Bull. Bureau Fish., Washington, 41: 161.
 1934 *Anoplodactylus lentus*, - DAWSON, Biol. Bull. Woods Hole Mar. Biol. Lab., 66: 62-68, pl. 1, figs. 1-17 (character of blood).
 1943 *Anoplodactylus lentus*, - HEDGPETH, Proc. New England Zool. Club, 22: 45.
 1943 *Anoplodactylus lentus*, - NEEDLER, Can. Atl. Fauna, 10: 14, fig. 17a-d.
 1948 *Anoplodactylus lentus*, - HEDGPETH, Proc. U.S. Natn. Mus., 97 (3216): 225--226, fig. 28a-c
 1973 *Anoplodactylus lentus*, - KRAEUTER, J. nat. Hist., 7: 494.
 1975 *Anoplodactylus lentus*, - STOCK, Bull. mar. Sci., 24 (4): 1055-1056, figs. 42f-i, 43a-b
 1976 *Anoplodactylus lentus*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., 27: 338, fig. 1.
 1986 *Anoplodactylus lentus*, - STOCK, Bull. mar. Sci., 38 (3): 439.
 1992 *Anoplodactylus lentus*, - CHILD, Mem. Hourglass Cruises, 9 (1): 38 [tab. 5], 39 [key], 49-52, fig. 22, 67 [text], 76 [tab. 12], 78 [tab. 14].

Distribution: tropical and temperate W-Atlantic, from Massachusetts (U.S.A.) to northern coast of South America.

Depth range: 20-600 m.

Habitat: sand and soft bottom

Anoplodactylus lineatus NAKAMURA & CHILD 1991

1991 *Anoplodactylus laminifer* NAKAMURA & CHILD, Smiths. Contr. Zool., 512: 27

[key], 27-29, fig. 10.

Distribution: Japan.
Depth range: 79-80 m.

Anoplodactylus longiceps STOCK 1951

- 1941 *Anoplodactylus longicollis* WILLIAMS, Mem. Qd. Mus., **12** (1): 36-38, figs. 2-5.
 1951 *Anoplodactylus longiceps* STOCK, Rés. Sci. Crois. Navire-École Belge "Mercator", **5**. Mem. Inst. Roy. Sci. Nat. Belg., sér. 2, fasc. 43: 16.
 1954 *Anoplodactylus longiceps*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 83.
 1956 *Anoplodactylus longiceps*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 97-98, figs. 14c-d.
 1963 *Anoplodactylus longiceps*, - Rec. Austr. Mus., **26** (1): 56 [key].
 1975 *Anoplodactylus longiceps*, - CHILD, Smiths. Contr. Zool., **190**: 20, fig. 9f.
 1990 *Anoplodactylus longiceps*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 331.
 1991 *Anoplodactylus longiceps*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 144.

Distribution: Australia (Great Barrier Reef; NW of Point Cloates); Guam.

Depth range: 2-21m.

Habitat: dead coral rubble; dredged with crustaceans and shells.

Anoplodactylus longiformis CHILD 1977

- 1977 *Anoplodactylus longiformis* CHILD, Proc. biol. Soc. Wash., **90** (2): 441-444, fig. 1.

Distribution: New Caledonia (Ile des Pins, Kuto).
Habitat: algae.

Anoplodactylus mamillous STOCK 1954

- 1954 *Anoplodactylus mamillous* STOCK, Vidensk. Medd. dansk. naturhist. Foren., **116**: 72-74, figs. 31c-e, 32.
 1971 *Anoplodactylus mamillous*, - UTINOMI, Publ. Seto. Mar. Biol. Lab., **18** (5): 327.
 1991 *Anoplodactylus mamillous*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 29.

Distribution: Japan.
Depth range: 96-732 m.
Habitat: hard bottom.

Anoplodactylus maritimus HODGSON 1914

- 1914 *Anoplodactylus maritimus* HODGSON, Zool. Anz., **45** (4): 164.
 1915 *Anoplodactylus maritimus*, - HODGSON, Ann. Mag. nat. Hist., (8) **15** (85): 148.

- 1927 *Anoplodactylus maritimus*, - HODGSON, Dtsch. Südpolar-Exp., **19** (Zool. II): 357.
- 1934 *Anoplodactylus parvus* GILTAY, Bull. Mus. roy. Hist. nat. Belg., **10** (42): 1-3, figs. 1-5.
- 1940 *Anoplodactylus maritimus*, - MARCUS, Bol. Fac. Filos., Cién. Letr., Univ. S. Paulo, **19** (Zool. 4): 60.
- 1948 *Anoplodactylus parvus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 223-224, fig. 27e-f.
- 1951 *Anoplodactylus parvus*, - STOCK, Mem. Inst. roy. Sci. nat. Belg., sér. **2**, fasc. 43: 13, figs. 14-16.
- 1954 *Anoplodactylus parvus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **24**: 127.
- 1955 *Anoplodactylus parvus*, - BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 590-591, fig. 1, pl. 1.
- 1957 *Anoplodactylus parvus*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 85.
- 1966 *Anoplodactylus parvus*, - FAGE & STOCK, Ann. Inst. Océanogr. Monaco., **44**: 326.
- 1973 *Anoplodactylus parvus*, - KRAEUTER, J. nat. Hist., **7** (5): 494-495.
- 1975 *Anoplodactylus parvus*, - STOCK, Bull. mar. Sci., **24** (4): 1069-1074, fig. 54.
- 1982 *Anoplodactylus maritimus*, - CHILD, Smiths. Contr. Zool., **349**: 21.
- 1990 *Anoplodactylus maritimus*, - STOCK, Zool. Meded., **63** (16): 229.
- 1992 *Anoplodactylus maritimus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 52-53, fig. 23.
- 1992 *Anoplodactylus maritimus*, - STOCK, Tijdschr. Ent., **135**: 131, 139.

Distribution: amphi-Atlantic; known from the Azores and Cape Verde Islands, as well as from Chesapeake Bay south to Brazil.

Depth range: 0-100 m.

Habitat: *Thalassia*; often associated with floating Sargassum.

***Anoplodactylus marshallensis* CHILD 1982**

- 1982 *Anoplodactylus marshallensis* CHILD, Proc. Biol. Soc. Wash., **95** (2): 274-275, fig. 2a-f.

Distribution: Marshall Islands (Enewetak Atoll).

Depth range: littoral.

Habitat: on pier pilings.

***Anoplodactylus massiliensis* BOUVIER 1916**

- 1916 *Anoplodactylus massiliensis* BOUVIER, Bull. Mus. natn. Hist. nat., Paris (1), 22: 14, fig. 34.
- 1923 *Anoplodactylus massiliensis*, - BOUVIER, Faune de France, **7**: 40.
- 1967 *Anoplodactylus massiliensis*, - STOCK, Atlantide Rep., **9**: 52-53, fig. 3a-b.
- 1970 *Anoplodactylus massiliensis*, - STOCK, "Meteor"-Forsch. - Ergebn., D, 5: 9-10, figs. 14-17.
- 1987 *Anoplodactylus massiliensis*, - ARNAUD, Mesogee, **47**: 45.

Distribution: W-Africa; off Canary Islands; Mediterranean.

Depth range: 300m.

Habitat: hard bottom.

Anoplodactylus massiliformis STOCK 1975

- 1975 *Anoplodactylus massiliformis* STOCK, Bull. mar. Sci., **24** (4): 1063-1066, figs. 48-49.
1979 *Anoplodactylus massiliformis*, - CHILD, Smiths. Contr. Zool., **293**: 56.
1986 *Anoplodactylus massiliformis*, - STOCK, Bull. mar. Sci., **38** (3): 439.

Distribution: Brazil; northern border of South America to Panamá; Barbados.

Depth range: 0-100m.

Habitat: muddy sand, shell debris, sponge bottom; gravel; calcareous sand.

Anoplodactylus minusculus CLARK 1970

- 1970 *Anoplodactylus minusculus* CLARK, Trans. R. Soc. N.Z., Biol. Sci., **12** (3): 15-17, figs. 9-16.

Distribution: Australia (Queensland).

Depth range: littoral.

Anoplodactylus minutissimus STOCK 1954

- 1954 *Anoplodactylus minutissimus* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 78-80, fig. 34.

Distribution: Singapore.

Depth range: intertidal.

Anoplodactylus micros BOURDILLON 1955

- 1955 *Anoplodactylus micros* BOURDILLON, Rev. Trav. Inst. Pêches Marit., **19** (4): 591-592, pl. I, figs. 3-8.
1986 *Anoplodactylus micros*, - STOCK, Bull. mar. Sci., **38** (3): 404.

Distribution: with certainty known only from Martinique, French Antilles; other records doubtful.

Depth range: 2-4m.

Habitat: *Sargassum*.

Anoplodactylus monotrema STOCK 1979

- 1979 *Anoplodactylus monotrema* STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 15-18, figs. 4-5.
1979 *Anoplodactylus monotrema*, - CHILD, Smiths. Contr. Zool., **293**: 56-58, fig. 19c.
1982 *Anoplodactylus monotrema*, - CHILD, Smiths. Contr. mar. Sci., **12**: 372.
1986 *Anoplodactylus monotrema*, - STOCK, Bull. mar. Sci., **38** (3): 404.

- 1990 *Anoplodactylus monotrema*, - MÜLLER, Bonn. zool. Beitr., **41** (3-4): 284.
 1992 *Anoplodactylus monotrema*, - CHILD, Smiths. Contr. Zool., **526**: 37-38.
 1992 *Anoplodactylus monotrema*, - STOCK, Tijdschr. Ent., **135**: 131, 139.

Distribution: W-Atlantic, Florida to Brazil; Galapagos Archipelago.

Depth range: 0-18m.

Habitat: in a variety of substrata, often in algal vegetation.

***Anoplodactylus neglectus* HOEK 1898**

- 1898 *Anoplodactylus neglecta* HOEK, Tijdschr. Nederl. Dierk. Ver., ser. 2, vol. 1: 293-295, pls. I-II, figs. 7-10.

Distribution: between Prince Edwards and Crozet Islands.

Depth range: 2928m.

***Anoplodactylus nodosus* HILTON 1939**

- 1939 *Anoplodactylus nodosus* HILTON, J. Ent. Zool. Pomona College, **31** (2): 29.
 1940 *Anoplodactylus nodosus*, - MARCUS, Bol. Fac. Filos., Ciênc. Letr., Univ. S. Paulo, **19** (Zool. 4): 41 [list].
 1941 *Anoplodactylus nodosus*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 257 [key].
 1942 *Anoplodactylus nodosus*, - HILTON, J. Ent. Zool. Pomona College, **34** (3): 72.
 1975 *Anoplodactylus nodosus*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 193-196, fig. 2.

Distribution: California (Catalina Island).

***Anoplodactylus oculatus* CARPENTER 1905**

- 1905 *Anoplodactylus oculatus* CARPENTER, Sci. Invest. Fish. Branch Ireland, **4**: 4-5, pl. II, figs. 7-11.
 1978 *Anoplodactylus oculatus*, - STOCK, Cah. Biol. Mar., **19**: 216.

Distribution: NE-Atlantic.

Depth range: 607-850m.

***Anoplodactylus oculospinosus* HILTON 1942**

- 1942 *Anoplodactylus oculospinosus* HILTON, J. Ent. Zool. Pomona College, **34** (3): 72-73.
 1975 *Anoplodactylus oculospinosus*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 196-198, fig. 3.

Distribution: California (Monterey Bay).

Depth range: 91m.

Habitat: from rocks.

Anoplodactylus ophiurophilus STOCK 1979

1979 *Anoplodactylus ophiurophilus* STOCK, Bijdr. Dierk., **48** (2): 156-160, figs. 1-9.

Distribution: Seychelles (Aldabra Atoll).

Depth range: littoral and shallow subtidal.

Habitat: associated with ophiuroid *Ophiocoma doederleini*.

Anoplodactylus pacificus HILTON 1942

1942 *Anoplodactylus pacificus* HILTON, J. Ent. Zool. Pomona College, **34** (3): 73.

1955 *Anoplodactylus pacificus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 242-243.

1975 *Anoplodactylus pacificus*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 198-201, fig. 4.

Distribution: California (Santa Catalina Island).

Depth range: 86m.

Anoplodactylus paradigitatus CHILD 1988

1988 *Anoplodactylus paradigitatus* CHILD, Smiths. Contr. Zool., **468**: 18-20, fig. 8.

Distribution: Philippines (Negros and Batan Island).

Depth range: 0.5-2m

Anoplodactylus pectinus HEDGPETH 1948

1948 *Anoplodactylus pectinus*, HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 234-236, fig. 34.

1954 *Anoplodactylus pectinus*, - HEDGPETH, Fish. Bull. Fish & Wildlife Service U.S., **89** (55): 427.

1955 *Anoplodactylus pectinus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 235, fig. 11.

1973 *Anoplodactylus pectinus*, - ARNAUD, Tethys, **4** (4): 955-957.

1974 *Anoplodactylus pectinus*, - CHILD, Proc. biol. Soc. Wash., **87** (43): 500.

1974 *Anoplodactylus pectinus*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 17.

1975 *Anoplodactylus pectinus*, - STOCK, Bull. mar. Sci., **24** (4): 1050-1052, fig. 41a.

1979 *Anoplodactylus pectinus*, - CHILD, Smiths. Contr. Zool., **293**: 58.

1979 *Anoplodactylus pectinus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **59** (184): 15.

1982 *Anoplodactylus pectinus*, - CHILD, Smiths. Contr. mar. Sci., **12**: 372-373.

1986 *Anoplodactylus pectinus*, - STOCK, Bull. mar. Sci., **38** (3): 404.

1988 *Anoplodactylus pectinus*, - CHILD, Smiths. Contr. Zool., **468**: 20.

1988 *Anoplodactylus pectinus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 662.

Distribution: probably cosmopolitan, most often recorded from the Caribbean.

Depth range: 0-34m.

Habitat: in a variety of substrata.

***Anoplodactylus pelagicus* FLYNN 1928**

- 1928 *Anoplodactylus pelagicus* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., I): 25-27, fig. 14.
 1954 *Anoplodactylus pelagicus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 128, fig. 19.
 1963 *Anoplodactylus pelagicus*, - STOCK, Ann. S. Afr. Mus., **46** (12): 340.
 1981 *Anoplodactylus pelagicus*, - STOCK, Cah. Biol. Mar., **22**: 463-464, fig. 6.

Distribution: North and South Atlantic, extending into the Indian Ocean in the sub-Antarctic zone of Prince Edward and Crozet islands; also taken in the Gulf of Alaska.

Depth range: 549-3600m.

***Anoplodactylus perforatus* NAKAMURA & CHILD 1982**

- 1982 *Anoplodactylus perforatus* NAKAMURA & CHILD, Proc. biol. Soc. Wash., **95** (2): 289-291, fig. 3.
 1983 *Anoplodactylus perforatus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 49.
 1988 *Anoplodactylus perforatus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 663.
 1988 *Anoplodactylus perforatus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 813-815.
 1991 *Anoplodactylus perforatus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 29-30.

Distribution: Samoa, Japan (incl. Ryukyu Islands).

Depth range: 7-113m.

***Anoplodactylus petiolatus* (KRØYER 1844)**

- 1844 *Phoxichilidium petiolatum* KRØYER, Naturhist. Tidskr., **2** (4): 123.
 1891 *Anoplodactylus petiolatus*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 25-29, pl. II, fig. 2a-i.
 1916 *Anaphia petiolata*, - LEBOUR, J. mar. biol. Ass. U.K., **11** (1): 51-56, figs. 1-3 (life history).
 1932 *Anoplodactylus petiolatus*, - TIMMERMANN, - Z. Morph. Oek. Tiere, **25**: 327, fig. 3.
 1940 *Anoplodactylus petiolatus*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 61-62, fig. 5a-b.
 1945 *Anoplodactylus petiolatus*, - LEBOUR, J. mar. biol. Ass. U.K., **26**: 157-159, fig. 6a-h.
 1948 *Anoplodactylus petiolatus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 222, fig. 27a-d.
 1950 *Anoplodactylus petiolatus*, - SAWAYA, Mem. Inst. Oswaldo Cruz, **47** (1-2): 73.
 1951 *Anoplodactylus petiolatus*, - Mem. Inst. Roy. Sci. Nat. Belg., **2** (43): 16.

- 1957 *Anoplodactylus petiolatus*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 85.
 1959 *Anoplodactylus petiolatus*, - BACESCU, Rev. Biol. Acad. Rep. Pop. Roum., **4** (1): 117.
 1958 *Anoplodactylus petiolatus*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
 1962 *Anoplodactylus petiolatus*, - STOCK, Ent. Ber., **12**: 218.
 1966 *Anoplodactylus petiolatus*, - STOCK, Atlantide Rept., **9**: 52.
 1972 *Anoplodactylus petiolatus*, - KING, J. nat. Hist., **6**: 622.
 1973 *Anoplodactylus petiolatus*, - ARNAUD, Tehys, **5** (1): 151-152.
 1973 *Anoplodactylus petiolatus*, - KRAEUTER, J. nat. Hist., **7**: 495.
 1973 *Anoplodactylus petiolatus*, - KRAPP, Beaufortia, **21** (277): 72.
 1974 *Anoplodactylus petiolatus*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 44-45, fig. 18.
 1975 *Anoplodactylus petiolatus*, - STOCK, Bull. mar. Sci., **24** (4): 1072-1075, fig. 53.
 1976 *Anoplodactylus petiolatus*, - ARNAUD, Acta Ecol. Iranica **1**: 69.
 1976 *Anoplodactylus petiolatus*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 338-340, fig. 2.
 1977 *Anoplodactylus guyanensis* CHILD, Proc. biol. Soc. Wash., **90** (3): 591-593, fig. 3.
 1978 *Anoplodactylus petiolatus*, - STOCK, Cah. Biol. Mar., **19**: 216.
 1982 *Anoplodactylus petiolatus*, - CHILD, Smiths. Contr. Zool., **349**: 21.
 1986 *Anoplodactylus petiolatus*, - Chimenz & Cottarelli, Oebalia, **13**, N.S.: 138.
 1986 *Anoplodactylus guyanensis*, - STOCK, Bull. mar. Sci., **38** (3): 404.
 1987 *Anoplodactylus petiolatus*, - ARNAUD, Mesogee, **47**: 45-46.
 1987 *Anoplodactylus petiolatus*, - BITAR, Bull. Inst. Sci., Rabat, **11**: 163.
 1987 *Anoplodactylus petiolatus*, - STOCK, Cah. Biol. Mar., **28**: 514.
 1989 *Anoplodactylus petiolatus*, - SCHÜLLER, Bonn. zool. Beitr., **40** (3/4): 290.
 1992 *Anoplodactylus petiolatus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 53-55, fig. 24.
 1993 *Anoplodactylus petiolatus*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: tropical and temperate Atlantic; Mediterranean, Black Sea.

Depth range: 0-1200m.

Habitat: in a variety of substrata.

Anoplodactylus pharus STOCK 1975

- 1975 *Anoplodactylus pharus* STOCK, Bull. mar. Sci., **24** (4): 1053-1055, fig. 42a-e.

Distribution: Florida.

Depth range: 256m.

Anoplodactylus polignaci BOUVIER 1914

- 1914 *Anoplodactylus polignaci* BOUVIER, Bull. Soc. ent. France, 1914: 224-226, figs. 1-3.
 1942 *Anoplodactylus polignaci*, - FAGE, Arch Zool. Exp. Gén., 82 (Notes & Rev., 2): 82-83.
 1943 *Anoplodactylus polignaci*, - HEDGPETH, Proc. New England Zool. Club, **22**: 45-46.
 1948 *Anoplodactylus polignaci*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 230,

fig. 30a-d.

1966 *Anoplodactylus polignaci*, - STOCK, Atlantide Rep., **9**: 53-54, fig. 3c.

Distribution: Florida; off Guinea, W-Africa.

Depth range: 32-50m.

Anoplodactylus pseudotarsalis MÜLLER 1992

1992 *Anoplodactylus pseudotarsalis* MÜLLER, Bonn. zool. Beitr., **43** (1): 167-170, figs. 31-36, 173 [text].

Distribution: Malaysia (Tioman Archipelago, Pulau Babi Besar).

Depth range: 1-2m.

Habitat: dead corals.

Anoplodactylus pycnosoma (HELFER 1938)

1938 *Peritrichia pycnosoma* HELFER, Sitz.-Ber. Ges. naturforsch. Freunde Berlin, 1937 (2): 176-177, fig. 7.

1940 *Halosoma pycnosoma*, - MARCUS, Bol. Fac. Filos., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 45-46.

1953 *Anoplodactylus pycnosoma*, - STOCK, Beaufortia, **35** (4): 41, fig. 5.

1954 *Anoplodactylus pycnosoma*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 75-77, fig. 33.

1971 *Anoplodactylus pycnosoma*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 326.

1974 *Anoplodactylus pycnosoma*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 16.

1975 *Anoplodactylus pycnosoma*, - CHILD, Smiths. Contr. Zool., **190**: 20.

1975 *Anoplodactylus pycnosoma*, - STOCK, Trav. Mus. Hist. nat. "Grigore Antipa", **16**: 132.

1983 *Anoplodactylus pycnosoma*, - CHILD, Proc. biol. Soc. Wash., **96** (4): 708.

1983 *Anoplodactylus pycnosoma*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 50.

1986 *Anoplodactylus pycnosoma*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 44.

1987 *Anoplodactylus pycnosoma*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.

1988 *Anoplodactylus pycnosoma*, - CHILD, Bull. biol. Soc. Wash., **8**: 59.

1988 *Anoplodactylus pycnosoma*, - CHILD, Smiths. Contr. Zool., **468**: 20.

1990 *Anoplodactylus pycnosoma*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, **45** (9): 100-101.

1990 *Anoplodactylus pycnosoma*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 77.

1991 *Anoplodactylus pycnosoma*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 144.

1991 *Anoplodactylus pycnosoma*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 30.

1992 *Anoplodactylus pycnosoma*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 173, figs. 43-48.

Distribution: known from scattered places in the W-Indian Ocean to Japan.

Depth range: 1.5-122m.

Habitat: dead corals.

Anoplodactylus pygmaeus (HODGE 1864)

- 1864 *Pallene pygmaea* HODGE, Ann. Mag. nat. Hist., **3** (13): 116, pl. XIII, figs. 16-17.
- 1881 *Phoxichilidium exiguum* DOHRN, Monogr. Fauna Flora Golf. Neapel, **3**: pp. 34, 36, 56, 76, 99, 181-184, pl. XII, figs. 19-22.
- 1889 *Phoxichilidium pygmaeum*, - TOPSENT, Bull. Soc. Linéenne Normandie, Caen, **4** (2) (1888-1889): 62.
- 1905 *Anoplodactylus pygmaeus*, - CARPENTER, Sci. Invest. Fish. Branch Ireland, **4**: 6.
- 1908 *Anoplodactylus petiolata*, - NORMAN, J. Linn. Soc. London (Zool.), **30**: 202.
- 1912 *Halosoma exiguum*, - LOMAN, Bull. Inst. Océanogr. Monaco, **238**: 6.
- 1913 *Anoplodactylus pygmaeus*, - DOGIEL, Zeitschr. Wiss. Zool., **107** (4): 665-685, pl. XXII, figs. 1-8.
- 1918 *Anoplodactylus exiguum*, - WIRÉN, Zoologisk. Bidr. Uppsala, **6**: 42.
- 1923 *Anoplodactylus exiguum*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10) **6** (3,4): 120.
- 1923 *Anoplodactylus exiguum*, - BOUVIER, Faune de France, **7**: 43, fig. 39.
- 1928 *Anoplodactylus pygmaeus*, - GILTAY, Bull. & Ann. Soc. Roy. Ent. Belg., **68** (9-10): 217.
- 1932 *Anoplodactylus pygmaeus*, - SCHLOTTKE, Wiss. Meeresunters. Kommiss. Wiss. Unters. Dtsch. Meere, Abt. Helgoland, (N.F.), **18** (9): 7, fig. 6.
- 1936 *Anoplodactylus exiguum*, - STEPHENSEN, Göteborgs Kungl. Vetensk. Vitterhets. Samh. Handl., (5) (B) **4** (14): 44-45, fig. 10.
- 1945 *Anoplodactylus pygmaeus*, - LEBOUR, J. mar. biol. Assoc. U.K., **26**: 159-162, fig. 7a-1.
- 1948 *Anoplodactylus pygmaeus*, - HEDGPETH, Proc. U.S. natn. Mus., **3216**: 224-225, fig. 27.
- 1954 *Anoplodactylus pygmaeus*, - STOCK, Vidensk. Medd. dansk. naturhist. Foren., **116**: 77.
- 1958 *Anoplodactylus pygmaeus*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **16**: 4.
- 1962 *Anoplodactylus pygmaeus*, - STOCK, Ent. Ber., **22**: 218.
- 1970 *Anoplodactylus pygmaeus*, - STOCK, "Meteor"-Forsch.-Ergebn., D, **5**: 10.
- 1970 *Anoplodactylus pygmaeus*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **2** (1): 1.
- 1972 *Anoplodactylus pygmaeus*, - KING, J. nat. Hist., **6**: 622.
- 1973 *Anoplodactylus pygmaeus*, - ARNAUD, Tethys, **5** (1): 152.
- 1973 *Anoplodactylus pygmaeus*, - KRAPP, Beaufortia, **21** (277): 72.
- 1974 *Anoplodactylus pygmaeus*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 46-47, fig. 19.
- 1976 *Anoplodactylus pygmaeus*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.
- 1978 *Anoplodactylus pygmaeus*, - CHIMENTZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 94-96.
- 1981 *Anoplodactylus pygmaeus*, - STOCK, Cah. Biol. Mar., **22**: 464-465.
- 1983 *Anoplodactylus pygmaeus*, - KRAPP, Bonn. zool. Beitr. **34** (1-3): 412.
- 1987 *Anoplodactylus pygmaeus*, - ARNAUD, Mesogee, **47**: 46.
- 1987 *Anoplodactylus pygmaeus*, - BITAR, Bull. Inst. Sci., Rabat, **11**: 163.
- 1990 *Anoplodactylus pygmaeus*, - STOCK, Zool. Meded., **63** (16): 230.
- 1992 *Anoplodactylus pygmaeus*, - CHILD, Mem. Hourglass Cruises, **9** (1): 38 [tab. 5], 39 [key], 55-56, fig. 25.
- 1992 *Anoplodactylus pygmaeus*, - PÉREZ-RUZAFA, A. & T. MUNILLA, Sci. Mar., **56** (1): 25.
- 1993 *Anoplodactylus pygmaeus*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: Sweden to West Africa and Azores; Mediterranean Sea; Red Sea; Virginia to Florida; Gulf of

Mexico; Caribbean Sea; ?Argentina.

Depth range: 0-587-?4558m

Habitat: in a variety of substrata, also on drifting weed; from the polyps of *Obelia* sp. and *Podocoryne carneae* and from galls in *Coryne muscoides*; adults frequently collected on *Bowerbankia*.

Anoplodactylus quadratispinosus HEDGPETH 1943

- 1943 *Anoplodactylus quadratispinosus* HEDGPETH, Proc. New England Zool. Club, **22**: 47-48, pl. 8, figs. a-g.
 1948 *Anoplodactylus quadratispinosus*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 232.
 1986 *Anoplodactylus quadratispinosus*, - STOCK, Bull. mar. Sci., **38** (3): 404.

Distribution: Florida (near Key West).

Anoplodactylus reimerae CHILD 1979

- 1979 *Anoplodactylus reimerae* CHILD, Smiths. Contr. Zool., **293**: 59-61, fig. 20.
 1992 *Anoplodactylus reimerae*, - CHILD, Smiths. Contr. Zool., **526**: 38.

Distribution: Panamá Pacific; Ecuador.

Depth range: 0-9m.

Anoplodactylus robustus (DOHRN 1881)

- 1945 *Phoxichilidium robustum*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 153.
 1979 *Anoplodactylus robustus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **184**: 18-22, fig. 6.
 1987 *Anoplodactylus robustus*, - ARNAUD, Mesogee, **47**: 46-47.

Distribution: Mediterranean; Congo Belge.

Depth range: 4-6m.

Habitat: "sur algues calcaires Mélobésiéées" (ARNAUD).

Anoplodactylus shimodaensis NAKAMURA & CHILD 1982

- 1982 *Anoplodactylus shimodaensis* NAKAMURA & CHILD, Proc. biol. Soc. Wash., **95** (2): 285-287, fig. 2.
 1983 *Anoplodactylus shimodaensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 50.
 1991 *Anoplodactylus shimodaensis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 30.

Distribution: Japan.

Depth range: 2-40m.

Anoplodactylus simplex CLARK 1963

1963 *Anoplodactylus simplex* CLARK, Mem. Qd. Mus., **20** (3): 50-51, fig. 25A-F.
1982 *Anoplodactylus simplex*, - STAPLES, Mem. Qd. Mus., **20** (3): 461.

Distribution: Australia.

Depth range: 5m.

Habitat: sand and weeds.

***Anoplodactylus simulator* STOCK 1975**

1975 *Anoplodactylus simulator* STOCK Bull. mar. Sci., **24** (4): 1066-1068, fig. 50.
1986 *Anoplodactylus simulator*, - STOCK, Bull. mar. Sci., **38** (3): 440, fig. 15c.

Distribution: Straits of Florida.

Depth range: 176-183m.

***Anoplodactylus spinirostrum* STOCK 1973**

1973 *Anoplodactylus spinirostrum* STOCK, Beaufortia, **20**: 122-123, fig. 11.

Distribution: SE-Australia.

Depth range: 10m.

Habitat: on *Amathea* sp. (Bryozoa) and *Aglaophenia plumosa* (Hydrozoa).

***Anoplodactylus spinosus* MÖBIUS 1902**

1902 *Anoplodactylus spinosus* MÖBIUS, Wiss. Ergebni. Dt. Tiefsee-Exped., 1898-1899, 3: 188-189, pl. XXVIII, figs. 8-12.

Distribution: Agulhas Bank.

Depth range: 10m or less.

***Anoplodactylus spurius* STOCK 1992**

1992 *Anoplodactylus spurius* STOCK, Tijdschr. Ent., **135**: 132-134, 139, figs. 65-75.

Distribution: southern Brazil.

Depth range: 19-220m.

***Anoplodactylus squalidus* CLARK 1973**

1973 *Anoplodactylus squalida* CLARK, Pac. Sci., **27** (1): 30-33, fig. 2A-G.
1988 *Anoplodactylus rimulus* CHILD, Proc. biol. Soc. Wash., **101** (4): 59-61, fig. 3.
1988 *Anoplodactylus rimulus*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (4): 815.
1989 *Anoplodactylus squalidus*, - MÜLLER, Bonn. zool. Beitr., **40** (2): 126-127, figs. 45-56.

Distribution: New Britain; Seychelles (Aldabra atoll); Society Islands (Moorea).

Depth range: 1-6m.

Habitat: coral reefs. - *Halimeda* on dead coral; dead corals.

Anoplodactylus stellatus NAKAMURA & CHILD 1983

1983 *Anoplodactylus stellatus* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 50, fig. 17.

Distribution: Japan.

Depth range: 7-15m.

Anoplodactylus stictus MARCUS 1940

1940 *Anoplodactylus stictus* MARCUS, Bol. Fac. Filos., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 65-68, pl. VI, fig. 7E, pl. VII, fig. 7A-D.

1957 *Anoplodactylus stictus*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 84.

1992 *Anoplodactylus stictus*, - STOCK, Tijdschr. Ent., **135**: 132-134 [text], 134, 139, fig. 62.

Distribution: Brazil to northern Argentina.

Depth range: 0-250m.

Habitat: seaweeds on rocky shore.

Anoplodactylus stocki BACESCU 1959

1959 *Anoplodactylus stocki* BACESCU, Rev. Biol. Acad. Rep. Pop. Roum., **4** (1): 123-127, fig. 3A-F.

1987 *Anoplodactylus stocki*, - ARNAUD, Mesogee, **47**: 47.

Distribution: Black Sea.

Depth range: 71m.

Anoplodactylus stri CHILD 1979

1979 *Anoplodactylus stri* CHILD, Smiths. Contr. Zool., **293**: 61-63, fig. 21.

Distribution: Panamá Caribbean.

Depth range: intertidal.

Anoplodactylus tanseii NAKAMURA & CHILD 1991

1991 *Anoplodactylus tanseii* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 27 [key], 30-32, fig. 11.

Distribution: Japan.

Depth range: 262-563m.

Anoplodactylus tarsalis STOCK 1968

- 1968 *Anoplodactylus tarsalis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 52-54, fig. 19.
 1973 *Anoplodactylus tarsalis*, - ARNAUD, Tethys **4** (4): 955.
 1988 *Anoplodactylus tarsalis*, - CHILD, Smiths. Contr. Zool., **468**: 20.
 1990 *Anoplodactylus tarsalis*, - MÜLLER, Bonn. zool. Beitr., **41** (1): 78.
 1992 *Anoplodactylus tarsalis*, - MÜLLER, Bonn. zool. Beitr., **43** (1): 170 [text], 171-173, figs. 37-42.
 1992 *Anoplodactylus tarsalis*, - STOCK, Bijdr. Dierk., **62** (2): 95.

Distribution: Philippines; Madagascar; Kenya; Oman.

Depth range: 0-0.5m.

Habitat: dead corals; algae.

Anoplodactylus tenuicorpus CHILD 1991

- 1988 *Anoplodactylus attenuatus* CHILD, Smiths. Contr. Zool., **468**: 12-14, fig. 5.
 1988 *Anoplodactylus attenuatus*, - CHILD, Bull. biol. Soc. Wash., **8**: 56.
 1991 *Anoplodactylus tenuicorpus* CHILD, Proc. biol. Soc. Wash., **104** (1): 142-143.

Distribution: Guam; Seychelles (Aldabra Atoll).

Depth range: 6-10m.

Habitat: coral rubble.

Anoplodactylus torus CHILD & HEDGPETH 1971

- 1971 *Anoplodactylus torus* CHILD & HEDGPETH, J. nat. Hist., **5**: 610-612, fig. 1a-f.

Distribution: Galápagos Archipelago.

Depth range: shallow to 6.5m.

Habitat: from coral and coralline heads; *Padina*.

Anoplodactylus trispinosus STOCK 1951

- 1951 *Anoplodactylus trispinosus* STOCK, Mém. Inst. Roy. Sci. Nat. Belg., **2** (43): 14-16, figs. 17-22.
 1954 *Anoplodactylus trispinosus*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 127.
 1954 *Anoplodactylus trispinosus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 35.
 1964 *Anoplodactylus trispinosus*, - STOCK, Bull. Sea Fish. Res. Stat. Haifa, **35**: 31, figs. 8-9 (Curaçao record only).
 1975 *Anoplodactylus trispinosus*, - STOCK, Bull. mar. Sci., **24** (4): 1076-1079, figs. 55, 56a.
 1979 *Anoplodactylus trispinosus*, - CHILD, Smiths. Contr. Zool., **293**: 63.
 1986 *Anoplodactylus trispinosus*, - STOCK, Bull. mar. Sci., **38** (3): 404.

Distribution: amphiatlantic.

Depth range: 0-10m.

Habitat: in a variety of substrata.

Anoplodactylus tubiferus (HASWELL 1884)

- 1884 *Phoxichilidium tubiferum* HASWELL, Proc. linn. Soc. N.S. Wales, **9**: 1032, pl. 57 figs. 1-5.
- 1904 *Anoplodactylus tubiferus*, - COLE, Harriman Alaska Exped., **10**: 288.
- 1908 *Anoplodactylus stylops* LOMAN, Siboga Exped. Monogr., **40**: 71, pl. 11 figs. 20-24.
- 1908 *Anoplodactylus tubiferus*, - LOMAN, Siboga Exp. Monogr., **40**: 72-73.
- 1907 *Anoplodactylus pulcher* CARPENTER, Trans. Linn. Soc. London, **2** (12): 97-98, pl. 12 figs. 13-19.
- 1919 *Anoplodactylus tubiferus*, - FLYNN, Pap. Proc. Roy. Soc. Tasm., 1918: 79.
- 1941 *Anoplodactylus tubiferus*, - WILLIAMS, Mem. Qd. Mus., **12** (1): 35.
- 1954 *Anoplodactylus pulcher*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 84-85, figs. 38a, 39.
- 1963 *Anoplodactylus tubiferus*, - CLARK, Rec. Austr. Mus., **26** (1): 49.
- 1965 *Anoplodactylus pulcher*, - STOCK, Beaufortia, **13** (151): 29, fig. 46.
- 1968 *Anoplodactylus pulcher*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 49.
- 1973 *Anoplodactylus pulcher*, - ARNAUD, Tethys, **4** (4): 957.
- 1973 *Anoplodactylus pulcher*, - STOCK, Beaufortia, **21** (279): 97.
- 1982 *Anoplodactylus tubiferus*, - STAPLES, Mem. Qd. Mus., **20** (3): 457-459, fig. 2c-f.
- 1983 *Anoplodactylus tubiferus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 49.
- 1988 *Anoplodactylus tubiferus*, - CHILD, Bull. Biol. Soc. Wash., **8**: 61.
- 1989 *Anoplodactylus tubiferus*, - MÜLLER, Senckenbergiana marit., **20** (5/6): 281-282, figs. 15-20.
- 1991 *Anoplodactylus tubiferus*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 32.
- 1991 *Anoplodactylus tubiferus*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 224.

Distribution: widely distributed in the Indo-West Pacific.

Depth range: subtidal-135m.

Habitat: coralline algae; muddy sand; shelly sand.

Anoplodactylus turbidus STOCK 1975

- 1964 *Anoplodactylus trispinosus*, - STOCK, Bull. Sea Fish. Res. Sta. Haifa, **35**: 31.
- 1974 *Anoplodactylus turbidus*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, **4** (3): 16.
- 1975 *Anoplodactylus turbidus* STOCK, Bull. mar. Sci., **24** (4): 1080, fig. 56b-g.

Distribution: Red Sea (Dahlak Archipelago); Madagascar.

Depth range: shallow water.

Habitat: on reefs.

Anoplodactylus typhloides STOCK 1991

- 1991 *Anoplodactylus typhloides* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 204-206, figs. 53-54.

Distribution: New Caledonia.

Depth range: 825-1475m.

***Anoplodactylus typhlops* SARS 1888**

- 1888 *Anoplodactylus typhlops* SARS, Arch. Math. Naturvidensk., Oslo, **12**: 29-31, pl. 2, fig. 3a-c.
- 1891 *Anoplodactylus typhlops*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 29-31, pl. II, fig. 3a-e.
- 1905 *Anoplodactylus typhlops*, - CARPENTER, Fish. Ireland Sci. Invest., 1904: 5, pl. III, figs. 12-19.
- 1908 *Anaphia typhlops*, - NORMAN, J. Linn. Soc. London, **30**: 203-
- 1933 *Anoplodactylus typhlops*, - STEPHENSEN, Medd. Grönland, **79** (6): 44-45, fig. 12.
- 1935 *Anoplodactylus typhlops*, - STEPHENSEN, Bergens Mus. Arbok. Naturv. Rek., 1935 (2): 29-30.
- 1948 *Anoplodactylus typhlops*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 228-229, fig. 29a-c.
- 1955 *Anoplodactylus typhlops*, - STOCK, Viddensk. Medd. dansk naturhist. Foren., **117**: 235-236, fig. 12a.
- 1973 *Anoplodactylus typhlops*, - TURPAEVA, Trudy Inst. Okeanol. P..P. Shirshov, Akad. Nauk. SSSR, **91**: 181-183, pl. 1, figs. 1-8.
- 1974 *Anoplodactylus typhlops*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **98**: 288.
- 1982 *Anoplodactylus typhlops*, - CHILD, Smiths. Contr. Zool., **349**: 21.
- 1987 *Anoplodactylus typhlops*, - STOCK, Cah. Biol. Mar., **28**: 514-515.
- 1988 *Anoplodactylus typhlops*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 130-131.
- 1991 *Anoplodactylus typhlops*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 207, fig. 55.
- 1992 *Anoplodactylus typhlops*, - CHILD, Smiths. Contr. Zool., **526**: 38.

Distribution: North and South Atlantic; Gulf of Alaska; Costa Rica (Cocos Ridge).

Depth range: 915-3620m.

***Anoplodactylus unilobus* STOCK 1959**

- 1959 *Anoplodactylus unilobus* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 564-565, fig. 8.

Distribution: South Africa.

Depth range: 11m.

Habitat: sand.

***Anoplodactylus velamellus* NAKAMURA & CHILD 1991**

- 1991 *Anoplodactylus velamellus* NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 32-34, fig. 12.

Distribution: Japan.

Depth range: 50-60m.

Anoplodactylus vema CHILD 1982

1982 *Anoplodactylus vema* CHILD Smiths. Contr. Zool., **349**: 21-24, fig. 7a-g.

Distribution: Argentine Slope, E of Bahia Samborombon.
Depth range: 676m.

Anoplodactylus versluysi LOMAN 1908

- 1908 *Anoplodactylus versluysi* LOMAN, Siboga Exp. Monogr., **40**: 73-74, pl. III, figs. 33-39.
 1954 *Anoplodactylus versluysi*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 84-85, figs. 38a, 39.
 1959 *Anoplodactylus versluysi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 200.
 1965 *Anoplodactylus versluysi*, - STOCK, Beaufortia, **13** (151): 1: 29-30, fig. 46.
 1971 *Anoplodactylus versluysi*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 326.
 1983 *Anoplodactylus versluysi*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 52-54.
 1983 *Anoplodactylus versluysi*, - Bull. Mus. natn. Hist. nat., Paris, 4e sér., **5**, sect. A., no. 1: 304.
 1987 *Anoplodactylus versluysi*, - NAKAMURA, Biol., Lab., Imperial Household: 17-18, pl. 15.
 1991 *Anoplodactylus versluysi*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 224.

Distribution: Singapore, Indonesia, Philippines, Madagascar.

Depth range: 0-400m.

Habitat: coral bottom; corals and *Lithothamnion*; shelly sand; on *Dendronephtha*.

Anoplodactylus virescens (HODGE 1864)

- 1945 *Phoxichilidium virescens*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 150-152, fig. 4a-f.
 1973 *Anoplodactylus virescens*, - KRAPP, Beaufortia, **21** (277): 70-71, fig. 7.
 1974 *Phoxichilidium virescens*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 40.
 1976 *Anoplodactylus virescens*, - ARNAUD, Acta Ecol. Iranica, **1**: 69.
 1979 *Anoplodactylus virescens*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **184**: 22-24, fig. 7.
 1987 *Anoplodactylus virescens*, - ARNAUD, Mesogee, **47**: 47.

Distribution: Mediterranean; NW-Atlantic.

Depth range: 0-16m.

Habitat: *Corallina mediterranea*; *Posidonia* seagrass; on *Syncoryne eximia* and *Coryne muscoides*.

Anoplodactylus viridintestinalis COLE 1904

1904 *Halosoma viridintestinale* COLE, Harriman Alaska Exped., **10**: 286-288, pl. 14,

- fig. 11, pl. 24, figs. 6-8, pl. 25, figs. 1-4.
- 1951 *Halosoma viridintestinale*, - HEDGPETH, Wasmann J. Biol., **9** (1): 106.
- 1955 *Anoplodactylus viridintestinalis*, - STOCK, Vidensk. Medd. dansk naturist. Foren., **117**: 239.
- 1960 *Anoplodactylus viridintestinalis*, - ZIEGLER, The Veliger, **3** (1): 19.
- 1975 *Anoplodactylus viridintestinalis*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 202.
- 1976 *Anoplodactylus viridintestinalis*, - BIRKELAND, REIMER & YOUNG, Ecol. Res. Ser., EPA-600/3-76-028: 158.
- 1976 *Anoplodactylus viridintestinalis*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 340.
- 1976 *Anoplodactylus viridintestinalis*, - REIMER, Mar. Biol., **35** (3): 299.
- 1979 *Anoplodactylus viridintestinalis*, - CHILD, Smiths. Contr. Zool., **293**: 63.

Distribution: California south to Panama.

Depth range: littoral.

Habitat: on bryozoans; washed from weeds.

***Anoplodactylus vulcanus* Child, 1992**

- 1992 *Anoplodactylus vulcanus* CHILD, Smiths. Contr. Zool., **526**: 38-40, fig. 16.

Distribution: Ecuador.

***Anoplodactylus xenus* STOCK 1980**

- 1980 *Anoplodactylus xenus* STOCK, Steenstrupia, **6** (1): 1-4, figs. 1-6.

Distribution: New Zealand (Cape Maria van Diemen).

Depth range: intertidal

***Endeis* Philippi 1843**

- 1923 *Endeis*, - CALMAN, Rec. Indian Mus., **24** (3): 290-291 [remarks].
- 1948 *Endeis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 238 [diagnosis, remarks].
- 1954 Endeidae, *Endeis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 129 [diagnosis, key to South African species].
- 1968 *Endeis*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **131**: 59 [key to species].
- 1969 Endeidae, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 63 [diagnosis].
- 1969 *Endeis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 63-64 [diagnosis, remarks], 65-66 [key to species].
- 1979 *Endeis*, - Child, Smiths. Contr. Zool., **293**: 66 [key to Middle American species].
- 1983 *Endeis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 41 [key to Japanese species].
- 1992 *Endeis*, - CHILD, Mem. Hourglass Cruises, **9** (1): 37.

***Endeis australis* (HODGSON 1907)**

- 1907 *Phoxichilus australis* HODGSON, Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 5-7, pl. I, fig. 1.
- 1911 *Phoxichilus australis*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, 152: 1139.
- 1913 *Phoxichilus australis*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), 6: 118-120, fig. 74.
- 1915 *Endeis australis*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 49, fig. 11.
- 1932 *Endeis australis*, - GORDON, Discovery Rep., 6: 93-94.
- 1935 *Endeis australis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 297.
- 1935 *Phoxichilus australis*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, 5 (4) (2): 279.
- 1938 *Endeis australis*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), 2 (8): 20.
- 1944 *Endeis australis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 48.
- 1952 *Endeis australis*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) 24 (2): 184.
- 1969 *Endeis australis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 64.
- 1976 *Endeis australis*, - PUSHKIN, Zool. Zh., 55 (6): 924, figs. k-m [text].

Distribution: Antarctic-subantarctic areas.

Depth range: 5-1000m.

***Endeis biseriata* STOCK 1968**

- 1908 *Phoxichilus meridionalis*, - LOMAN, Siboga Exped. Monogr., 40: 78-79, pl. XI, figs. 153-155 [part.].
- 1968 *Endeis biseriata* STOCK, Vidensk. Medd. dansk naturhist. Foren., 131: 57-60, fig. 21.
- 1970 *Endeis biseriata*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, 2 (1): 1.
- 1974 *Endeis biseriata*, - STOCK, Bull. Zool. Mus. Univ. Amsterdam, 4 (3): 17.
- 1979 *Endeis biseriata*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., 59 (184): 28-30, fig. 9.
- 1988 *Endeis biseriata*, - CHILD, Smiths. Contr. Zool., 468: 20.
- 1990 *Endeis biseriata*, - CHILD, Proc. biol. Soc. Wash., 103 (2): 332-333.
- 1992 *Endeis biseriata*, - STOCK, Tijdschr. Ent., 135: 134, 139.

Distribution: Brazil, Red Sea, Indian Ocean, Indonesia, Australia, Philippines and Hawaii.

Depth range: 0-37m.

Habitat: stones; algae; reef and sand flat rubble; seaweeds on rocks; amongst ascidians, bryozoans and sponges growing on boulders.

***Endeis charybdea* (DOHRN 1881)**

- 1881 *Phoxichilus charabdaeus* DOHRN, Monogr. Fauna Flora Golf. Neapel, 3: 174, pl. X, figs. 7-13, pl. Xa, figs. 14-15, 21-22, pl. XI, figs. 11, 14-15.
- 1923 *Chilophoxus charabdaeus*, - BOUVIER, Faune de France, 7: 46, fig. 43.
- 1937 *Chilophoxus charybdaeus* var. *bispinatus*, - BOUVIER, Ann. Sci. Nat., Paris, (Zool.), (10) 20 (1): 36, fig. 10.

- 1942 *Chilophoxus charibdaeus*, - FAGE, Arch. Zool. Exp. Gen., **82** (Notes & Rev., 2): 84, fig. 6.
 1965 *Endeis charybdaea*, - STOCK & SOYER, Vie Milieu, **16** (1B): 417-418, fig. 1A-B.
 1970 *Endeis charybdaea*, - STOCK, "Meteor" Forsch.-Ergebn., D (5): 10.
 1975 *Endeis charybdaea*, - KRAPP, Sarsia, **59**: 85-94, fig. 1.
 1987 *Endeis charybdaea*, - ARNAUD, Mesogee, **47**: 52.
 1990 *Endeis charybdaea*, - STOCK, Zool. Meded., **63** (16): 230-231.

Distribution: Mediterranean; Cape Verde Islands and temperate NE-Atlantic.

Depth range: 10-800m.

Habitat: in a variety of substrata.

***Endeis clipeata* (MÖBIUS 1902)**

- 1902 *Phoxichilus clipeatus* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 196, pl. 30, figs. 6-10.
 1928 *Endeis clipeatus*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 29.
 1954 *Endeis clipeatus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 129-130, fig. 20.
 1956 *Endeis clipeatus*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 89.
 1959 *Endeis clipeatus*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 563.
 1962 *Endeis clipeatus*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 285.
 1965 *Endeis clipeata*, - STOCK, Beaufortia, **13** (151): 31.

Distribution: South Africa; Madagascar.

Depth range: 0-80m.

Habitat: rocks; shelly sand, gravel, carpets of *Phyllochaetopterus socialis*.

***Endeis flaccida* CALMAN 1923**

- 1908 ?*Phoxichilus* sp., - LOMAN, Siboga Exped. Monogr., **40**: 79, pl. 13, fig. 189.
 1923 *Endeis flaccidus* CALMAN, Rec. Indian Mus., **24** (3): 295-297, fig. 17.
 1953 *Endeis flaccida*, - STOCK, Temminckia, **9**: 300.
 1968 *Endeis flaccida*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 56, 59 [key].
 1970 *Endeis flaccida*, - STOCK, Ent. Ber., **30**: 3-4.
 1975 *Endeis flaccida*, - STOCK, Bull. mar. Sci., **24** (4): 1085, fig. 58a-b.
 1979 *Endeis flaccida*, - CHILD, Smiths. Contr. Zool., **293**: 66.
 1986 *Endeis flaccida*, - STOCK, Bull. mar. Sci., **38** (3): 440.

Distribution: probably circumtropical.

Depth range: 0-69m.

Habitat: fouling on pilings, floating oil-pipes and buoys; seagrass (*Thalassia*).

***Endeis laevis* (GRUBE 1871)**

- 1974 *Endeis laevis*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 48 [key], 52-54, figs. 20B, D, 21B, D, 22B, 23B, D, E.

Distribution: NE-Atlantic (Great Britain).

Depth range: sublittorally to about 15m.

Habitat: associated with hydroids (*Obelia dichotoma* and *Tubularia larynx*).

***Endeis meridionalis* (BÖHM 1879)**

- 1879 *Phoxichilus meridionalis* BÖHM, Monatsber. Kgl. Preuss. Akad. Wiss. Berlin, 1879: 189-191, pl. 2, fig. 4.
- 1890 *Phoxichilus meridionalis*, - SCHIMKEWITSCH, Izvest. Imperatorskavo Obshch. Lubitelei Estestvozn. Anthrop. Ethn., Moskva, 67: 20.
- 1891 *Phoxichilus meridionalis*, - SCHIMKEWITSCH, Arch. Zool. Exp. Gén., 9 (2): 509-510.
- 1908 *Phoxichilus meridionalis*, - LOMAN, Siboga Exped. Monogr., 40: 78.
- 1923 *Endeis meridionalis*, - CALMAN, Rec. Indian Mus., 24 (3): 291-293, fig. 15.
- 1929 *Phoxichilus meridionalis*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, 1: 186-188.
- 1965 *Endeis meridionalis*, - STOCK, Beaufortia, 151 (13): 30-31.
- 1968 *Endeis meridionalis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., 131: 59 [key].
- 1969 *Endeis meridionalis*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., 49: 64, 66 [key].
- 1971 *Endeis meridionalis*, - UTINOMI, Publ. Seto Mar. Biol. Lab., 18 (5): 327-328.
- 1973 *Endeis meridionalis*, - ARNAUD, Tethys, 4 (4): 959.
- 1973 *Endeis meridionalis*, - CLARK, Pac. Sci., 27 (1): 33, fig. 3A-D.
- 1979 *Endeis meridionalis*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., 59 (184): 27-28.
- 1980 *Endeis meridionalis*, - DANIEL & SEN, J. mar. biol. Ass. India, 17 (2) ("1975"): 165.
- 1982 *Endeis meridionalis*, - STOCK, Monit. Zool. ital., Suppl. 7, (7): 189.
- 1989 *Endeis meridionalis*, - MÜLLER, Bonn. zool. Beitr., 40 (2): 127.
- 1990 *Endeis meridionalis*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, 45 (9): 101.
- 1990 *Endeis meridionalis*, - MÜLLER, Zool. Abh. Staatl. Mus. Tierk. Dresden, 45 (10): 109.
- 1990 *Endeis meridionalis*, - STOCK, Zool. Meded., 63 (16): 231.

Distribution: circumtropical.

Depth range: 0.5-90m.

Habitat: on a variety of substrata.

***Endeis mollis* (CARPENTER 1904)**

- 1887 ?*Phoxichilus charybdeus*, - SCHIMKEWITSCH, Zool. Anz., 10 (251): 271.
- 1904 *Phoxichilus molle* CARPENTER, Rep. Governm. Ceylon Pearl Oyster Fish. Gulf Manaar, Suppl. Rep. 13: 182-183, figs. 1-7.
- 1907 *Phoxichilus mollis*, - CARPENTER, Trans. Linn. Soc. London, 12 (2): 98.
- 1908 *Phoxichilus mollis*, - LOMAN, Siboga Exped. Monogr., 40: 77-78.
- 1923 *Endeis mollis*, - CALMAN, Rec. Indian Mus., 25 (3): 293-294, fig. 16.
- 1927 *Endeis mollis*, - CALMAN, Trans. Zool. Soc. London, 22 (3): 408.
- 1938 *Endeis mollis*, - CALMAN, Sci. Rep. John Murray Exped., 1933-34, 5 (6): 160.
- 1942 ?*Chilophoxus charibdeus*, - PAGE, Arch. Zool. Exp. Gén., 82 (Notes & Rev.,

- 2): 84, fig. 6.
- 1951 *Endeis mollis*, - STOCK, Mem. Inst. Roy. Sci. Nat. Belg., **43** (2): 17-18, figs. 23-24.
- 1954 *Endeis mollis*, - BOURDILLON, Notes Stat. Océanogr. Salammbô, **35**: 4-8, figs. 1-3.
- 1954 *Endeis mollis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 130-132, fig. 21.
- 1957 *Endeis mollis*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 85-86.
- 1965 *Endeis mollis*, - STOCK, Beaufortia, **151**, 13: 31.
- 1966 *Endeis mollis*, - STOCK, Atlantide Rept., **9**: 54.
- 1971 *Endeis mollis*, - UTINOMI, Publ. Seto. Mar. Biol. Lab., **18** (5): 327; Literatur.
- 1975 *Endeis mollis*, - STOCK, Bull. mar. Sci., **24** (4): 1083-1085.
- 1975 *Endeis mollis*, - STOCK, Ent. Ber., **35**: 76.
- 1977 *Endeis mollis*, - CHILD, Proc. biol. Soc. Wash., **90** (2): 441.
- 1979 *Endeis mollis*, - CHILD, Smiths. Contr. Zool., **293**: 66.
- 1983 *Endeis mollis*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 41.
- 1986 *Endeis mollis*, - STOCK, Bull. mar. Sci., **38** (3): 440.
- 1987 *Endeis mollis*, - ARNAUD, Mesogee, **47**: 52.
- 1988 *Endeis mollis*, - CHILD, Smiths. Contr. Zool., **468**: 20-21.
- 1988 *Endeis mollis*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 664.
- 1991 *Endeis mollis*, - CHILD, Proc. biol. Soc. Wash., **104** (1): 144-145.

Distribution: pantropical.

Depth range: 0-60m.

Habitat: in a variety of substrata, often in coral reef locations.

***Endeis nodosa* HILTON 1942**

- 1942 *Endeis (Phoxichilus) nodosa* HILTON, Occ. Pap. Bernice P. Bishop Mus., **17** (3): 47-48, fig. 4.
- 1968 *Endeis nodosa*, - Vidensk. Medd. dansk naturhist. Foren., **131**: 59 [key].
- 1982 *Endeis nodosa*, - CHILD, Proc. biol. Soc. Wash., **95** (2): 275-277, fig. 2g-i.
- 1988 *Endeis nodosa*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., **101** (3): 663-664.
- 1991 *Endeis nodosa*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 35.
- 1992 *Endeis nodosa*, - Müller, Stud. Nat. Hist. Caribb. Reg., **71**: 50-51, figs. 20-22.

Distribution: Hawaii; Marshall Islands; Japan (Ryukyu Islands, Honshu); Caribbean Sea (Barbados).

Depth range: 0-45m.

Habitat: rocks and pilings; algae on buoys; hydroids growing on anchor chain.

***Endeis pauciporosa* STOCK 1970**

- 1970 *Endeis pauciporosa* STOCK, Bull. Zoöl. Mus. Univ. Amsterdam, **2** (1): 2-3, fig. 1.

Distribution: Red Sea; Caribbean Sea of Colombia.

Depth range: 5m.

Habitat: "harbour pier" (STOCK).

Endeis procera (LOMAN 1908)

- 1908 *Phoxichilus procerus* LOMAN, Siboga Exped. Monogr., **40**: 79-80, pl. XI, figs. 146-152.
 1954 *Endeis procera*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 88-89, fig. 41.

Distribution: SE-Asia.

Depth range: 50m.

Habitat: sand, shells.

Endeis spinosa (MONTAGU 1808)

- 1808 *Phallangium spinosum* MONTAGU, Trans. Linn. Soc. London, **9**: 100, pl. 5, fig. 7.
 1881 *Phoxichilus vulgaris* DOHRN, Fauna Flora Golf. Neapel, **3**: 169-174, pls. 10-11.
 1891 *Phoxichilus spinosus*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 15-20, pl. 1, fig. 3a-g.
 1908 *Endeis spinosus*, - NORMAN, J. Linn. Soc. London, (Zool.), **30**: 233.
 1910 *Endeis spinosus*, - COLE, Biol. Bull. Woods Hole mar. biol. Lab., **18** (4): 193-203, figs. 1-2.
 1913 *Endeis spinosus*, - SUMNER, OSBURN & COLE, Bull. Bureau Fish., Wash., **31** (1-2): 143.
 1917 *Phoxichilus spinosus*, - BOUVIER, Res. Campagnes Sci. accomplies sur son Yacht par Albert Ier Prince Souverain de Monaco, **51**: 30-31, pl. 2, fig. 2.
 1923 *Chilophoxus spinosus*, - BOUVIER, Faune de France, **7**: 45-46, figs. 42-43.
 1932 *Chilophoxus spinosus*, - TIMMERMANN, Z. Morph. Oek. Tiere, **25**: 327, 333.
 1935 *Phoxichilus spinosus*, - STEPHENSEN, Bergens Mus. Arbok, 1935 (2): 30.
 1940 *Endeis spinosa*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 73-75, fig. 9a-e.
 1942 *Chilophoxus spinosus*, - FAGE, Arch. Zool. Gen., **82** (Notes & Rev., 2): 83.
 1943 *Endeis spinosa*, - HEDGPETH, Proc. New England Zool. Club, **22**: 48.
 1945 *Endeis spinosa*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 163.
 1948 *Endeis spinosa*, - Proc. U.S. Natn. Mus., **97** (3216): 238-240, fig. 37.
 1952 *Endeis spinosa*, - STOCK, Boll. Soc. Venez. Stor. Nat., **6** (2): 185-186.
 1954 *Endeis spinosa*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 128.
 1957 *Endeis spinosa*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 85.
 1962 *Endeis spinosa*, - STOCK, Ent. Ber., **22**: 218.
 1965 *Endeis spinosa*, - STOCK & SOYER, Vie Milieu, **16** (1B): 418.
 1966 *Endeis spinosa*, - de HARO, Bol. Real Soc. Espan. Hist. Nat., Madrid (Sec. Biol.), **64**: 9.
 1966 *Endeis spinosa*, - SOYER, Doriania, **47** (174): 3.
 1966 *Endeis spinosa*, - STOCK, Atlantide Rep., **9**: 54.
 1967 *Endeis spinosa*, - de HARO, Publ. Inst. Biol. appl., Barcelona, **43**: 109, 112-113, fig. 5.
 1970 *Endeis spinosa*, - BAYER, VOSS & ROBINS, in: Bioenv. and Radiolog. Safety Feasibility Stud. Atl.-Pac. Interoc. Canal, Univ. Miami: 48.
 1972 *Endeis spinosus*, - KING, J. nat. Hist., **6**: 622.
 1973 *Endeis spinosa*, - KRAPP, Beaufortia, **21** (277): 72.
 1974 *Endeis spinosa*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 48 [key], 50-51, figs. 20A, C, 21A, C, 22A, 23A, C.

- 1979 *Endeis spinosa*, - CHILD, Smiths. Contr. Zool., **293**: 66.
 1979 *Endeis spinosa*, - CHIMENEZ, BRIGNOLI & BASCIANO, Cah. Biol. Mar., **20**: 480.
 1982 *Endeis spinosa*, - CHILD, Smiths. Contr. mar. Sci., **12**: 373-374.
 1982 *Endeis spinosa*, - CHILD, Smiths. Contr. Zool., **349**: 24.
 1987 *Endeis spinosa*, - ARNAUD, Mesogee, **47**: 52.
 1992 *Endeis spinosa*, - CHILD, Mem. Hourglass Cruises, **9** (1): 57 [tab. 8], 57-58,
 fig. 26.

Distribution: from Norway to Argentina, including the Mediterranean; ?Gulf of Mexico.

Depth range: 0.2-537m.

Habitat: in a variety of substrata, often found in floating *Sargassum*; in littoral zone on *Asparagopsis*; on top of a rock; ectocommensal on *Thecocarpus*; ectocommensal on *Eunicella*; amongst hydroids and sponges; on *Laminaria* holdfast; amongst hydroids and polyzoa; on *Pecten* valves; on sand and gravel; amongst *Nemertesia*; amongst *Antennularia*.

***Endeis straughani* CLARK 1970**

- 1884 ?*Phoxichilus charybdaeus*, - HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1033.
 1970 *Endeis straughani* CLARK, Trans. Roy. Soc. N.Z., **12** (3): 13-15, figs. 1-8.
 1979 *Endeis picta* BAMBER, Zool. J. Linn. Soc., **65**: 251-254, fig. 1.
 1982 *Endeis straughani*, - STAPLES, Mem. Qd. Mus., **20** (3): 461-462, fig. 5K-M, pl.
 1, fig. A-B.
 1990 *Endeis straughani*, - STOCK, Zool. Meded., **63** (16): 231.

Distribution: Australia (Queensland); Ghana; Azores.

Depth range: 10-20m.

***Endeis viridis* PUSHKIN 1976**

- 1976 *Endeis viridis* PUSHKIN, Zool. Zh., **55** (6): 923-925, figs. a-u.

Distribution: Crozet Archipelago, Subantarctica.

Depth range: 150-377m.

***Pallenopsis* WILSON 1881**

- 1948 *Pallenopsis*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 209 [diagnosis,
 remarks, key to NW-Atlantic species].
 1954 *Pallenopsis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 116 [diagnosis, key to
 South African species].
 1954 *Pallenopsis*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **116**: 63
 [diagnosis].
 1956 *Pallenopsis*, - STOCK, Vidensk. Medd. Dansk naturhist. Foren., **118**: 87
 [diagnosis of subgen. nov. *Pallenopsodon*].
 1957 *Pallenopsis*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 100-101
 [remarks], 102 [key to South American species of *fluminensis*-group].
 1975 *Pallenopsis*, - STOCK, Bull. mar. Sci., **24** (4): 1016-1017 [remarks], 1017-
 1018 [key to W-Atlantic, South American species and diagnosis of subgen.
 Pallenopsis], 1032-1036 [diagnosis of subgen. *Bathypallenopsis*, species

- groups], 1039-1040 [key to species of *mollissima*-group].
 1986 *Pallenopsis*, - STOCK, Bull. mar. Sci., **38** (3): 425 [remarks].
 1992 *Pallenopsis*, - CHILD, Mem. Hourglass Cruises, **9** (1): 37-38, 64-65
 [diagnosis].

Pallenopsis alcocki CALMAN 1923

- 1923 *Pallenopsis (Rigona) alcocki* CALMAN, Rec. Indian Mus., **24** (3): 282-284,
 figs. 9-10.

Distribution: Andaman Sea.

Depth range: 97m.

Pallenopsis angusta STOCK 1991

- 1991 *Pallenopsis (Pallenopsis) angusta* STOCK, Mém. Mus. natn. Hist. nat. (A),
151: 195-197, figs. 47-48.

Distribution: New Caledonia, Loyality Islands.

Depth range: 150-570m.

Pallenopsis annandalei CALMAN 1923

- 1923 *Pallenopsis annandalei* CALMAN, Rec. Indian Mus., **24** (3): 279-281, figs. 7-8.
 1991 *Pallenopsis (Bathypallenopsis) annandalei*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 198 [text].

Distribution: Laccadive Archipelago.

Depth range: 1164m.

Pallenopsis antipoda CLARK 1971

- 1971 *Pallenopsis (Pallenopsodon) antipoda* CLAR, N.Z. J. Mar. Freshw. Res., **5** (3-4): 428-430, fig. 1A-G.

Distribution: Antipodes Islands.

Depth range: 1280m.

Habitat: sand, rocks and pebbles.

Pallenopsis arctica STOCK 1956

- 1856 *Pallenopsis arctica* STOCK, Mitt. Hamburg. Zool. Mus. Inst., **54**: 37-40, figs. 3-4.

Distribution: Arctica (W-Spitzbergen).

Pallenopsis bicuspisdata STOCK 1968

- 1968 *Pallenopsis bicuspisdata* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**:

46-47, fig. 17c-h.

Distribution: off Lorenzo Marquez, Indian Ocean.

Depth range: 1628-1600m.

***Pallenopsis boehmi* SCHIMKEWITSCH 1930**

1973 *Pallenopsis boehmi*, - STOCK, Nederl. J. Zool., **23** (3): 347-352, figs. 1-10.
 1992 *Pallenopsis (Pallenopsis) boehmi*, - STOCK, Tijdschr. Ent., **135**: 129, 139.

Distribution: southern South America, Magellan Strait to southern Brazil.

Depth range: 35-100m.

***Pallenopsis brevidigitata* MÖBIUS 1902**

- 1902 *Pallenopsis brevidigitata* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 185-186, pl. 27, figs. 7-13.
 1928 *Pallenopsis brevidigitata*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 19.
 1938 *Pallenopsis brevidigitata*, - CALMAN, Sci. Rep. John Murray Exped., **5** (6): 160.
 1954 *Pallenopsis brevidigitata*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 120, fig. 15b-c.
 1956 *Pallenopsis (Pallenopsodon) brevidigitata* STOCK, - Vidensk. Medd. dansk naturhist. Foren., **118**: 87-88, fig. 10.
 1991 *Pallenopsis (Pallenopsis) brevidigitata*, - TURPAEVA, Zool. Zh., **70** (12): 37.

Distribution: Indo-West Pacific (E-Africa off Durban, off Zanzibar; SE-Asia).

Depth range: 57-500m.

Habitat: sand; sponges.

***Pallenopsis bulbifera* MUNILLA & STOCK 1984**

- 1984 *Pallenopsis bulbifera* MUNILLA & STOCK, Res. Exp. Cien., **12**: 33-37, figs. 1-14.
 1988 *Pallenopsis bulbifera*, - MUNILLA, Monogr. Zool. Mar., **3**: 183-185, figs. 1a, 2a, 5, tab. 1.

Distribution: Nambia.

Depth range: 183-329m.

Habitat: muddy-sandy, sandy, rocky and coraligenous bottom.

***Pallenopsis calcanea* STEPHENSEN 1933**

- 1932 ?*Pallenopsis* sp., - GORDON, Discovery Rep., **6**: 91-92, fig. 45a-c, 74c.
 +1933 *Pallenopsis calcanea* STEPHENSEN, Medd. Grönland, **79** (6): 21-24, fig. 5.
 1948 *Pallenopsis calcanea*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 211, fig. 21e-g.

- 1962 *Pallenopsis calcanea*, - HEDGPETH, Deep-Sea Res., **9**: 487-490, figs. 1-2.
 1968 *Pallenopsis calcanea*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 48.
 1970 *Pallenopsis calcanea*, - NESIS, Trans. P.P. Shirshov Inst. Oceanol., **88**: 168.
 1971 *Pallenopsis calcanea*, - TURPAEVA, Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **92**: 286.
 1978 *Pallenopsis aff. calcanea*, - STOCK, Cah. Biol. Mar., **19**: 216.
 1984 *Pallenopsis calcanea*, - MAUCHLINE, - J. nat. Hist., **18**: 316.
 1986 *Pallenopsis (Bathypallenopsis) calcanea*, - STOCK, Bull. mar. Sci., **38** (3): 437.
 1991 *Pallenopsis (Bathypallenopsis) calcanea*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 40.

Distribution: worldwide, scattered records.

Depth range: 660-2000m

Habitat: bathypelagic.

***Pallenopsis californica* SCHIMKEWITSCH 1893**

- 1893 *Pallenopsis californica* SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 39-41, pl. I, fig. 11, pl. II, figs. 18-23.
 1942 *Pallenopsis californica*, - HILTON, J. Ent. Zool. Pomona College, **34** (2): 40.
 1954 *Pallenopsis oscitans*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 122, fig. 17.
 1975 *Pallenopsis (Bathypallenopsis) californica* SCHIMKEWITSCH, Bull. mar. Sci., **24** (4): 1036.
 1991 *Pallenopsis (Bathypallenopsis) californica*, - TURPAEVA, Zool. Zh., **70** (12): 39.

Distribution: Gulf of California, Mozambique Strait.

Depth range: 1700-3000m.

***Pallenopsis candidoi* MELLO-LEITAO 1949**

- 1949 *Pallenopsis candidoi* MELLO-LEITAO, Arqu. Mus. Paran., **7** (7): 299-307, pls. IX-X.
 1957 *Pallenopsis candidoi*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 105, figs. 19-20.
 1966 *Pallenopsis candidoi*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 393.
 1975 *Pallenopsis (Pallenopsis) candidoi*, - STOCK, Bull. mar. Sci., **24** (4): 1018 [key], 1030.
 1976 *Pallenopsis candidoi*, - KRAPP & KRAEUTER, Bonn. zool. Beitr., **27**: 337.
 1992 *Pallenopsis (Pallenopsis) candidoi*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: Georgia (U.S.A.) south to southern Brazil.

Depth range: 15-102m.

***Pallenopsis capensis* BARNARD 1946**

- 1954 *Pallenopsis capensis* BARNARD, Ann. S. Afr. Mus., **41** (3): 118-119.
 1956 *Pallenopsis capensis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 83-84, fig. 7a.
 1962 *Pallenopsis capensis*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 285.

- 1988 *Pallenopsis capensis*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 141-142.
1988 *Pallenopsis capensis*, - MUNILLA, Monogr. Zool. Mar., **3**: 183-184, 186, figs. 1b, 2b, 3, 4, 5, tab. 1.

Distribution: South Africa.

Depth range: 51-342m.

Habitat: rocks; coarse sand, corals and shell.

***Pallenopsis childi* STOCK 1986**

- 1986 *Pallenopsis (Pallenopsis) childi*, - Bull. mar. Sci., **38** (3): 429-431, figs. 9-10.

Distribution: Florida to south of Hispaniola.

Depth range: 18-230m.

***Pallenopsis cidaribatus* CHILD 1975**

- 1975 *Pallenopsis cidaribatus* CHILD, Smiths. Contr. Zool., **190**: 16-18, fig. 8.

Distribution: Australia.

Depth range: 100-141m.

***Pallenopsis comosa* STOCK 1975**

- 1975 *Pallenopsis (Bathypallenopsis) comosa* STOCK, Bull. mar. Sci., **24** (4): 1043-1045, fig. 37.
1992 *Pallenopsis (Bathypallenopsis) comosa*, - CHILD, Smiths. Contr. Zool., **526**: 30.

Distribution: Gulf of Panamá; Peru-Chile Trench.

Depth range: 2945-3200m.

***Pallenopsis conirostris* TURPAEVA 1991**

- 1991 *Pallenopsis (Pallenopsis) conirostris* TURPAEVA, Zool. Zh., **70** (12): 37-39, fig. 2.

Distribution: south of Madagascar.

Depth range: 415-460m.

***Pallenopsis crosslandi* CARPENTER 1910**

- 1910 *Pallenopsis crosslandi* CARPENTER, J. Linn. Soc. London (Zool.), **31** (207): 257.
1972 *Pallenopsis crosslandi*, - ARNAUD, Tethys, Suppl. **3**: 158-159.

Distribution: E-Africa (Wasin Channel); Madagascar.

Depth range: 25-31m.

Habitat: reef slope; "substrat de dalle corallienne avec

nombreux Hydroïdes et Zoanthaires" (ARNAUD).

Pallenopsis denticulata HEDGPETH 1944

- 1944 *Pallenopsis denticulata* HEDGPETH, Proc. New England Zool. Club, **23**: 55-58,
pls. XI-XII.
1963 *Pallenopsis denticulata*, - CLARK, Rec. Austr. Mus., **26** (1): 4 [list].
1975 *Pallenopsis denticulata*, - CHILD, Smiths. Contr. Zool., **190**: 18-19.

Distribution: Australia.

Depth range: 9-15m.

Habitat: sand; rubble; sponges; corals.

Pallenopsis dentifera STOCK 1983

- 1983 *Pallenopsis dentifera* STOCK, Bull. Mus. natn. Hist. nat., (4) **5** (1): 300-304, figs. 1-10.
1988 *Pallenopsis dentifera*, - CHILD, Smiths. Contr. Zool., **468**: 23.

Distribution: Philippines.

Depth range: 166-172m.

Pallenopsis fluminensis (KRØYER 1844)

- 1844 *Phoxichilidium fluminensis* KRØYER, Naturhist. Tidskr., Kobenhavn, **1** (2): 104, 124, pl. I, fig. 1a-f.
1898 *Pallenopsis fluminensis*, - MEINERT, Dansk. Ingolf-Exped., **3** (1): 49, pl. V, figs. 1-6.
1899 *Pallenopsis fluminensis*, - MEINERT, Danish Ingolf-Exped., **3** (1): 52, pl. V, figs. 1-6.
1935 *Pallenopsis fluminensis*, - HELFER & SCHLÖTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 265, fig. 174.
1940 *Pallenopsis fluminensis*, - MARCUS, Revta. Ent., Rio de Janeiro, **11** (1-2): 182, figs. 1-6.
1940 *Pallenopsis fluminensis*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 24, fig. 1A-B.
1975 *Pallenopsis (Pallenopsis) fluminensis*, - STOCK, Bull. mar. Sci., **24** (4): 1020.
1992 *Pallenopsis (Pallenopsis) fluminensis*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: Brazil, from the State of Sao Paulo to north of Pernambuco.

Depth range: 0-49m.

Pallenopsis forficifera WILSON 1881

- 1881 *Pallenopsis forficifer* WILSON, Bull. Mus. comp. Zool., **8** (12): 250-252, pl. 4, figs. 15-18, pl. 5, fig. 23.
1943 *Pallenopsis forficifer*, - HEDGPETH, Proc. New England Zool. Club, **22**: 43.
1948 *Pallenopsis forficifer*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 209-

- 210, fig. 21c-d.
1975 *Pallenopsis (Pallenopsis) forcifer*, - STOCK, Bull. mar. Sci., **24** (4): 1030-1032, fig. 30a-b.
1986 *Pallenopsis (Pallenopsis) forficifera*, - STOCK, Bull. mar. Sci., **38** (3): 425.

Distribution: tropical and temperate northern W-Atlantic.
Depth range: 128-1000m.

***Pallenopsis gippslandiae* STOCK 1954**

- 1954 *Pallenopsis gippslandiae* STOCK 1954, Vidensk. Medd. dansk naturhist. Foren., **116**: 65-68, figs. 28-29.
1963 *Pallenopsis gippslandiae*, - CLARK, Rec. Austr. Mus., **26** (1): 43-44, fig. 22I.

Distribution: Australia.
Depth range: 61-236m.

***Pallenopsis glabra* MÖBIUS 1902**

- 1902 *Pallenopsis glabra* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 184-185, pl. XXVII, figs. 1-6.
1913 *Pallenopsis glabra*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 109-111, figs. 62-65.
1915 *Pallenopsis glabra*, - CALMAN, Rep. Brit. Antarct. "Terra Nova" Exped., **3**: 41-42.

Distribution: Antarctica.
Depth range: 250-450m.

***Pallenopsis guineensis* STOCK 1975**

- 1975 *Pallenopsis (Bathypallenopsis) guineensis* STOCK, Bull. mar. Sci., **24** (4): 1049-1050, fig. 40.

Distribution: Gulf of Guinea.
Depth range: 1949-1986m.

***Pallenopsis hiemalis* HODGSON 1907**

- 1952 *Pallenopsis hiemalis*, - FAGE, Bull. Mus. natn. Hist. nat., ser. 2, **24** (3): 270-271.
1975 *Pallenopsis hiemalis*, - PUSHKIN, Byull. Sov. Antarkt. Exped., **8** (90): 686-687, fig. 3 a-k, tab. 1.

Distribution: Antarctica (Ross, Mawson and Davis Seas, Sea of Cooperation and Sea of the Cosmonauts).
Depth range: 3-540m.

Pallenopsis hodgsoni GORDON 1938

- 1938 *Pallenopsis hodgsoni* GORDON Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 16-17, figs. 3a, 4d, 5d.
 1944 *Pallenopsis hodgsoni*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 45.
 1950 *Pallenopsis hodgsoni*, - HEDGPETH, Proc. U.S. natn. Mus., **100** (3260): 151.

Distribution: Antarctica, probably circumpolar.

Depth range: less than 641m.

Pallenopsis hoekii MIERS 1884

- 1884 *Pallenopsis hoekii* MIERS, Rep. Zool. Coll. made in the Indo-Pac. during the Voyage of HMS Alert 1881-82: 324-326, pl. XXXV, fig. B.
 1893 *Pallenopsis hoekii*, - CARPENTER, Sci. Rep. Roy. Dublin Soc., **8** (1) (4): 23, pl. II, fig. 11.
 1884 ?*Phoxichilidium hoekii*, - HASWELL, Proc. Linn. Soc. N.S.W., **9**: 1022 [list].
 1908 ?*Phoxichilidium hoekii*, - LOMAN, Siboga Exped. Monogr., **40**: 70-71.
 1908 *Pallenopsis (Rigona) rigens* LOMAN, Siboga Exped. Monogr., **40**: 68, pl. IX, figs. 128-133.
 1929 *Pallenopsis (Rigona) hoekii*, - FLYNN, Mem. Qd. Mus., **9** (3): 257-258.
 1953 *Pallenopsis* cf. *hoekii*, - STOCK, Temminckia, **9**: 288.
 1963 *Pallenopsis hoekii*, - CLARK, Rec. Austr. Mus., **26** (1): 42, fig. 24E.
 1971 *Pallenopsis hoekii*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 324.
 1975 *Pallenopsis hoekii*, - CHILD, Smiths. Contr. Zool., **190**: 19.

Distribution: Philippines, Indonesia, Australia (off Cape York, Dirk Hartogs Island).

Depth range: 7-134m.

Pallenopsis hoekiana SCHIMKEWITSCH 1930

- 1881 *Phoxichilidium fluminense*, - HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 81-82, pl. XIV, figs. 1-4.
 1930 *Pallenopsis hoekiana* SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 237.
 1940 *Pallenopsis hoekiana*, - MARCUS, Revta. Ent., Rio de Janeiro, **11** (1-2): 187-188.
 1975 *Pallenopsis (Pallenopsis) hoekiana*, - STOCK, Bull. mar. Sci., **24** (4): 1021.

Distribution: W-Atlantic.

Depth range: 94m.

Pallenopsis intermedia FLYNN 1928

- 1928 *Pallenopsis intermedia* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 20-23, figs. 10-12.
 1954 *Pallenopsis intermedia*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 117-118, figs. 15a, 16a-d.
 1956 *Pallenopsis intermedia*, - STOCK, Vidensk. Medd. dansk naturhist. Foren.,

118: 82-83, fig. 7b.

1959 *Pallenopsis intermedia*, - STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 561.

1962 *Pallenopsis intermedia*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 284.

1988 *Pallenopsis intermedia*, - MUNILLA, Monogr. Zool. Mar., **3**: 185.

Distribution: South Africa; Namibia.

Depth range: 29-233m.

Habitat: on a variety of substrata.

***Pallenopsis juttingae* STOCK 1964**

1964 *Pallenopsis (Pallenopsodon) juttingae* STOCK, Beaufortia, **11** (135): 46-51,
fig. 2a-g.

1973 *Pallenopsis (Pallenopsodon) juttingae*, - ARNAUD, Tethys, **5** (1): 148-149,
figs. 3-4.

Distribution: N-Atlantic (off Bay of Biscay, Golfe de Gascogne).

Depth range: 400m.

Habitat: found in plankton.

***Pallenopsis kempfi* STOCK 1975**

1975 *Pallenopsis (Pallenopsis) kempfi* STOCK, Bull. mar. Sci., **24** (4): 1025-1028,
figs. 28-29.

1986 *Pallenopsis (Pallenopsis) kempfi*, - STOCK, Bull. mar. Sci., **38** (3): 427-429,
fig. 8a-f.

1992 *Pallenopsis (Pallenopsis) kempfi*, - STOCK, Tijdschr. Ent., **135**: 139.

Distribution: Florida south to southern Brazil.

Depth range: 11-600m.

Habitat: sandy bottom.

***Pallenopsis knipovichii* TURPAEVA 1974**

1974 *Pallenopsis knipovichii* TURPAEVA, Trudy Inst. Okeanol., **98**: 291-293, pl. 7,
figs. 1-6.

Distribution: Scotia Sea.

Depth range: 583-540m.

***Pallenopsis kupei* CLARK 1971**

1971 *Pallenopsis kupei* CLARK, N.Z. J. Mar. Freshw. Res., **5** (3 & 4): 430-433, fig.
2A-J.

Distribution: New Zealand.

Depth range: 146-1530m.

Habitat: sand and shells.

Pallenopsis longicoxa STOCK 1966

1966 *Pallenopsis longicoxa* STOCK, Ann. Inst. Océanogr. Monaco, **44**: 394-396, fig. 4.

Distribution: Atlantic coast of South America, type locality uncertain.

Depth range: ?63m.

Pallenopsis longimana STOCK 1991

1991 *Pallenopsis (Bathypallenopsis) longimana* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 198-200, figs. 49-50.

Distribution: New Caledonia.

Depth range: 505-550m.

Pallenopsis longirostris WILSON 1881

1881 *Pallenopsis longirostris* WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 252-253, pl. 4, figs. 19-22, pl. 5, figs. 24-25.

1885 *Pallenopsis longirostris*, - VERRILL, Rep. U.S. Comm. Fish., 1883: 561.

1899 *Pallenopsis plumipes* MEINERT, Danish Ingolf-Exped., **3** (1): 51-52, pl. 4, figs. 1-7.

1942 *Pallenopsis longirostris*, - GILTAY, J. Fish. Res. Board Ca., **5** (5): 459.

1943 *Pallenopsis longirostris*, - NEEDLER, Can. Atl. Fauna, **10**: 13, fig. 16.

1973 *Pallenopsis longirostris*, - ARNAUD, Tethys, **5** (1): 150.

1981 *Pallenopsis longirostris*, - STOCK, Cah. Biol. Mar., **22**: 461-463, fig. 5a-f.

1986 *Pallenopsis (Bathypallenopsis) longirostris*, - STOCK, Bull. mar. Sci., **38** (3): 434.

1988 *Pallenopsis longirostris*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 142.

Distribution: N-Atlantic (Canadian Atlantic coast (off New Jersey) south to Bahamas), Indian Ocean (Zululand area) and South Africa (off Cape Point).

Depth range: 135-3065m.

Pallenopsis longiseta TURPAEVA 1958

1893 *Pallenopsis mollissima*, - SCHIMKEWITSCH, Bull. Mus. Comp. Zool., Harvard, **25** (2): 41-42.

1958 *Pallenopsis longiseta* TURPAEVA Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **27**: 359-361, pl. 2, figs. 1-6.

1975 *Pallenopsis (Bathypallenopsis) longiseta*, - STOCK, Bull. mar. Sci., **24** (4): 1042-1043, fig. 36.

Distribution: U.S.S.R. (Bering Sea, Sea of Okhotsk); Gulf of Panamá.

Depth range: 3060m.

Pallenopsis macneilli CLARK 1963

- 1963 *Pallenopsis macneilli* CLARK, Rec. Austr. Mus., **26** (1): 45-46, fig. 22A-H.
 1968 *Pallenopsis macneilli*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 48.
 1973 *Pallenopsis macneilli*, - STOCK, Beaufortia, **20** (266): 121, fig. 10A.
 1975 *Pallenopsis macneilli*, - CHILD, Smiths. Contr. Zool., **190**: 19.

Distribution: Australia.

Depth range: sublittoral-6m.

Habitat: growths in rock pool; *Ecklonia*.

Pallenopsis macronyx BOUVIER 1911

- 1913 *Pallenopsis macronyx*, - BOUVIER, Deuxième Expédition Antarctique Française (1908-1910), **6**: 112-117, figs. 66-72.

Distribution: Antarctica.

Depth range: 420m.

Pallenopsis mariae LARRAMENDY & de CASTELLANOS 1978

- 1978 *Pallenopsis mariae* LARRAMENDY & de CASTELLANOS, Neotropica, La Plata, **24** (72): 91-95, figs. A-E.

Distribution: Argentina.

Pallenopsis mauii CLARK 1958

- 1958 *Pallenopsis mauii* CLARK, Zool. Publ. Univ. Wellington, **23**: 4-6, figs. 11, 14, 16, 18, 19.
 1971 *Pallenopsis mauii*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (3-4): 433-435, fig. 3A-G.

Distribution: New Zealand.

Depth range: 695-1098m.

Habitat: mud, stones, rocks.

Pallenopsis meinerti SCHIMKEWITSCH 1930

- 1887 *Pallenopsis fluminensis*, - SCHIMKEWITSCH, Zool. Anz., **10** (251): 271.
 1930 *Pallenopsis meinerti* SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 237.
 1940 *Pallenopsis meinerti*, - MARCUS, Revta. Ent., Rio de Janeiro, **11** (1-2): 22.
 1975 *Pallenopsis (Pallenopsis) meinerti*, - STOCK, Bull. mar. Sci., **24** (4): 1018 [key], 1021-1024, figs. 26-27.
 1982 *Pallenopsis (Pallenopsis) meinerti*, - Smiths. Contr. Zool., **349**: 29-30.

Distribution: SW-Atlantic.

Depth range: 270-1011m.

Pallenopsis meridionalis HODGSON 1915

- 1927 *Pallenopsis meridionalis*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 337-338.
 1975 *Pallenopsis meridionalis*, - PUSHKIN, Byull. Sov. Antarkt. Exped., **8** (90): 685-686, fig. 2 a-k, tab. 1.

Distribution: Antarctica (Davis Sea).
Depth range: 3-355m.

Pallenopsis mixta STOCK 1986

- 1986 *Pallenopsis (Pallenopsis) mixta* STOCK, Bull. mar. Sci., **38** (3): 433-434, fig. 12.

Distribution: Venezuela (Isla Margarita).
Depth range: 91m.

Pallenopsis mollissima (HOEK 1881)

- 1881 *Phoxichilidium mollissimum* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 87-88, pl. XIII, figs. 6-9.
 1893 *Pallenopsis mollissima*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 41-42, pl. II, fig. 24.
 1949 *Pallenopsis mollissima*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 277, fig. 36f.
 1953 *Pallenopsis mollissima*, - STOCK, Temminckia, **9**: 288.
 1956 *Pallenopsis mollissima*, - FAGE, Galathea Rep., **2**: 172-173, figs. 5-8.
 1975 *Pallenopsis (Bathypallenopsis) mollissima*, - STOCK, Bull. mar. Sci., **24** (4): 1040-1042, fig. 35.
 1985 *Pallenopsis mollissima*, - BAMBER, Zool. J. Linn. Soc., **83**: 302.
 1986 *Pallenopsis (Bathypallenopsis) mollissima atlantica* STOCK, Bull. mar. Sci., **38** (3): 434-437, figs. 13-14.

Distribution: probably cosmopolitan, scattered records.
Depth range: 1235-3500m.

Pallenopsis notiosa (Child, 1992)

- 1992 *Pallenopsis (Pallenopsis) notiosa* CHILD, Smiths. Contr. Zool., **526**: 25-27, fig. 10.

Distribution: Chile (W of Bahia Las Canas).
Depth range: 290-450m.

Pallenopsis obliqua (THOMSON 1884)

- 1884 *Phoxichilidium obliquum* THOMSON, Trans. & Proc. N.Z. Inst., **16** (1883): 247-248, pl. XV, fig. 6, pl. XVI, figs. 1-2.
 1904 *Phoxichilidium obliquum*, - HUTTON, Index Fauna N.Z.: 247.
 1954 *Pallenopsis obliqua*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**:

- 64-65, fig. 30b-f.
- 1956 *Pallenopsis obliqua*, - STOCK, Medd. dansk naturhist. Foren., **118**: 85-86, fig. 9C.
- 1958 *Pallenopsis obliqua*, - CLARK, Zool. Publ. Univ. Wellington, **23**: 3-4.
- 1971 *Pallenopsis obliqua*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (2): 330.
- 1971 *Pallenopsis obliqua*, - CLARK, N.Z. J. Mar. Freshw. Res., **5** (3 & 4): 435.
- 1977 *Pallenopsis obliqua*, - CLARK, N.Z. J. Mar. Freshw. Res., **11** (1): 173-174, fig. 1A-C.

Distribution: New Zealand (incl. subantarctic islands).

Depth range: 0-24m.

Habitat: rocks; rock pool in *Durvilla* zone; algae.

***Pallenopsis oscitans* (HOEK 1881)**

- 1881 *Phoxichilidium oscitans* HOEK, Rep. Sci. Res. Expl. Voyage HMS Challenger, **3** (10): 89-90, pl. XIII, figs. 1-5.
- 1981 *Pallenopsis oscitans*, - STOCK, Cah. Biol. Mar., **22**: 461.
- 1984 *Pallenopsis (Bathypallenopsis) oscitans*, - STOCK, J. mar. biol. Ass. U.K., **64**: 746-747.

Distribution: Walvis Basin in southern Atlantic; Azores; entrance to English Channel.

Depth range: 1970-3550m.

***Pallenopsis ovalis* LOMAN 1908**

- 1908 *Pallenopsis ovalis* LOMAN, Siboga Exped. Monogr., **40**: 68-69, pl. X, figs. 136-137.
- 1916 *Pallenopsis ovalis*, - LOMAN, Zool. Meded., **2** (1): 20-21.
- 1923 *Pallenopsis (Rigona) ovalis*, - CALMAN, Rec. Indian Mus., **25** (3): 284-285, fig. 11.
- 1928 *Pallenopsis ovalis*, - FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 23.
- 1954 *Pallenopsis (Rigona) ovalis*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 120-121, fig. 15d.
- 1954 *Pallenopsis ovalis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 64.
- 1968 *Pallenopsis ovalis*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 48.

Distribution: Indian Ocean (Sri Lanka to Kei Islands).

Depth range: 4-86m.

Habitat: sand; muddy bottom.

***Pallenopsis paramollis* STOCK 1975**

- 1975 *Pallenopsis (Bathypallenopsis) paramollis* STOCK, Bull. mar. Sci., **24** (4): 1045-1048, figs. 38-39.

Distribution: NE-Pacific; Japan.

Depth range: 924-1303m.

***Pallenopsis patagonica* (HOEK, 1881)**

- 1932 *Pallenopsis patagonica*, - GORDON, Discovery Rep., **6**: 88-90, fig. 44.
 1938 *Pallenopsis patagonica*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 15.
 1940 *Pallenopsis patagonica*, - MARCUS, Revta. Ent., Rio de Janeiro, **11** (1-2): 191.
 1944 *Pallenopsis patagonica*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 46-48, fig. 15b, d, e.
 1950 *Pallenopsis patagonica*, - HEDGPETH, Proc. U.S. Natn. Mus., **100** (3260): 150-151.
 1957 *Pallenopsis patagonica*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 102-103, fig. 17.
 1961 *Pallenopsis patagonica*, - HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 5.
 1974 *Pallenopsis patagonica*, - TURPAEVA, Trudy Inst. Okeanol., Akad. Nauk SSSR, **98**: 288-290, pl. 5, figs. 1-10.
 1975 *Pallenopsis patagonica*, - PUSHKIN, Byull. Sov. Antarkt. Exped., **8** (90): 682-685, figs. 1 a-k, tab. 1.

Distribution: Antarctic to atlantic coast of South America.

Depth range: 36-344m

Habitat: broken shells and sand.

***Pallenopsis persimilis* STOCK 1956**

- 1956 *Pallenopsis persimilis* STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 84-87, figs. 8, 9a.
 1962 *Pallenopsis persimilis*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 285.

Distribution: South Africa.

Depth range: 43-92m.

Habitat: rocks; gravel with worm tubes.

***Pallenopsis pilosa* (HOEK 1881)**

- 1881 *Phoxichilidium pilosum* HOEK, Challenger Rep. (Zool.), **X**: 90-92, pl. XIII, figs. 10-13.
 1913 *Pallenopsis pilosa*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 107-109, figs. 60-61.
 1915 *Pallenopsis pilosa*, - CALMAN, Rep. Brit. Antarct. "Terra Nova" Exped., **3**: 42-43.
 1927 *Pallenopsis pilosa*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 335-336.
 1932 *Pallenopsis pilosa*, - GORDON, Discovery Rep., **6**: 87-88.
 1944 *Pallenopsis pilosa*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 45.
 1974 *Pallenopsis pilosa*, - TURPAEVA, Trudy Inst. Okeanol., **98**: 290-291, pl. 6, figs. 1-9.

Distribution: Antarctica; Scotia Sea.

Depth range: 25-3650m.

Pallenopsis safari STOCK 1984

1984 *Pallenopsis (Bathypallenopsis) safari* STOCK, Bull. Mus. natn. Hist. nat., Paris, (4) **6** (sec. A) (3): 707-709, figs. 10-21.

Distribution: west of Sri Lanka.

Depth range: 1035m.

Pallenopsis schmitti HEDGPETH 1943

1943 *Pallenopsis schmitti*, HEDGPETH, Proc. New England Zool. Club, **22**: 44.

1948 *Pallenopsis schmitti*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 212-214, fig. 22.

1955 *Pallenopsis schmitti*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **117**: 233-234, fig. 10.

1975 *Pallenopsis schmitti*, - STOCK, Bull. mar. Sci., **24** (4): 1028-1030, fig. 30c-d.

1979 *Pallenopsis schmitti*, - CHILD, Smiths. Contr. Zool., **293**: 46.

1986 *Pallenopsis (Pallenopsis) schmitti*, - STOCK, Bull. mar. Sci., **38** (3): 425-427, fig. 8g-i.

1992 *Pallenopsis (Pallenopsis) schmitti*, - CHILD, Mem. Hourglass Cruises, **9** (1): 65-67, fig. 30, 76 [tab. 12], 78 [tab. 14].

Distribution: tropical W-Atlantic. - Florida to Brazil; E-Pacific. - Panama, south of Balboa in the Canal zone.

Depth range: 27-600m.

Habitat: rock and broken shell; coralline bottom; bryozoans; sand; mud.

Pallenopsis scoparia FAGE 1956

1956 *Pallenopsis scoparia* FAGE, Galathea Rep., **2**: 171-172, figs. 1-4.

1973 *Pallenopsis scoparia*, - ARNAUD, Tethys, **5** (1): 149-150, figs. 1-2.

1975 *Pallenopsis (Bathypallenopsis) scoparia*, - STOCK, Bull. mar. Sci., **24** (4): 1032 [text].

1984 *Pallenopsis scoparia*, - MAUCHLINE, J. nat. Hist., **18**: 318.

1986 *Pallenopsis (Bathypallenopsis) scoparia*, - CHILD & HARBISON, J. mar. biol. Ass. U.K., **66**: 113-116, figs. 1-2.

1987 *Pallenopsis (Bathypallenopsis) scoparia*, - STOCK, Cah. Biol. Mar., **28**: 516, figs. 13-15.

1991 *Pallenopsis (Bathypallenopsis) scoparia*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 40.

1991 *Pallenopsis (Bathypallenopsis) scoparia*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 198.

1992 *Pallenopsis (Bathypallenopsis) scoparia*, - CHILD, Mem. Hourglass Cruises, **9** (1): 67-69, fig. 31.

Distribution: cosmopolitan (except for Polar seas).

Depth range: 400-1520m.

Habitat: bathypelagic; predator of the scyphomedusa *Periphylla periphylla*.

Pallenopsis sibogae LOMAN 1911

- 1908 *Pallenopsis plumipes*, - LOMAN, Siboga Exped. Monogr., **40**: 66-67, pl. XII, figs. 160-164.
- 1911 *Pallenopsis sibogae* LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-Naturwiss. Kl.), Suppl. **2** (4): 14.
- 1911 *Pallenopsis sibogae*, - LOMAN, Codicil to Siboga Monogr., **40**: 1.
- 1954 *Pallenopsis sibogae*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 63-64, fig. 30h-i.
- 1955 *Pallenopsis sibogae*, LOMAN, Publ. Seto Mar. Biol. Lab., **5** (1): 19.
- 1959 *Pallenopsis sibogae*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 199.
- 1962 *Pallenopsis sibogae*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **10** (1): 62.
- 1971 *Pallenopsis sibogae*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 324.
- 1987 *Pallenopsis sibogae*, - NAKAMURA, Biol. Lab., Imperial Household: 10-11, pl. 8.
- 1991 *Pallenopsis (Pallenopsis) sibogae*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 41.

Distribution: Indonesia to Japan.

Depth range: 37-160m.

Habitat: sand; corals.

Pallenopsis spicata HODGSON 1914

- 1915 *Pallenopsis spicata*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 44-46, fig. 9.
- 1927 *Pallenopsis spicata*, - HODGSON, Dt. Südpolar-Exped., **19**, Zool. 11: 338-339, fig. 10.
- 1932 *Pallenopsis spicata*, - GORDON, Discovery Rep., **6**: 90-91.
- 1938 *Pallenopsis spicata*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 19.
- 1944 *Pallenopsis spicata*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 48.

Distribution: Antarctica, probably circumpolar.

Depth range: 180-549m.

Pallenopsis spinipes CARPENTER 1907

- 1907 *Pallenopsis spinipes* CARPENTER, Trans. Linn. Soc. London, **12** (2): 96, pl. 12, figs. 1-12.
- 1916 *Pallenopsis spinipes*, - LOMAN, Zool. Meded., **2** (1): 20.
- 1957 *Pallenopsis spinipes*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 88, figs. 5-7.
- 1987 *Pallenopsis spinipes*, - NAKAMURA, Biol. Lab., Imperial Household: 11-12, pl. 9.
- 1988 *Pallenopsis (Pallenopsis) spinipes*, - CHILD, Bull. biol. Soc. Wash., **8**: 66.

Distribution: Maldives; Amirantes Islands; Seychelles (Aldabra Atoll); Fiji Islands; Japan (Sagami Bay).

Depth range: 0-71m.

***Pallenopsis stschapovae* TURPAEVA 1958**

1958 *Pallenopsis stschapovae* TURPAEVA Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, **27**: 356-358, pl. 1, figs. 1-8.

Distribution: NW-Pacific.
Depth range: ?0-5500m.

***Pallenopsis styliostre* HEDGPETH 1949**

1949 *Pallenopsis styliostre* HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 278-280, fig. 36a-e.

1991 *Pallenopsis (Bathypallenopsis) styliostris*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 40.

Distribution: Japan.
Depth range: 1000-1300m.

***Pallenopsis temperans* STOCK 1953**

1953 *Pallenopsis temperans* STOCK, Temminckia, **9**: 291-293, fig. 10.

1954 *Pallenopsis temperans*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 63, fig. 30g.

1968 *Pallenopsis temperans*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 47.

1971 *Pallenopsis temperans*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 324.

Distribution: Philippines, Japan, Korea Straits, Central Pacific (Fanning and Line Island).

Depth range: 0.5-284m.

Habitat: sand; shells; corals.

***Pallenopsis tongaensis* CLARK 1973**

1973 *Pallenopsis tongaensis* CLARK, Pac. Sci., **27** (1): 28-30, fig. 1A-J.

Distribution: Tonga.

***Pallenopsis triregia* CLARK 1962**

1962 *Pallenopsis triregia* CLARK, N.Z. J. Sci., **5** (4): 517-519, fig. 1A-H.

Distribution: New Zealand.
Depth range: 82-137m.

***Pallenopsis tritonis* HOEK 1883**

1883 *Pallenopsis tritonis* HOEK, Trans. Roy. Soc. Edinburgh, **32** (1): 7-10, pl. 1,

figs. 1-6.

- 1905 *Pallenopsis holti* CARPENTER, Sci. Invest. Fish. Branch Ireland, **4**: 4, pl. I, figs. 1-6.
- 1909 *Pallenopsis tritonis*, - d'ARCY THOMPSON, in: The Cambr. Nat. Hist.: 542.
- 1982 *Pallenopsis (Bathypallenopsis) tritonis*, - CHILD, Smiths. Contr. Zool., **349**: 40.
- 1983 *Pallenopsis tritonis*, - BAMBER, Zool. J. Linn. Soc., **77**: 72-73, fig. 6.
- 1984 *Pallenopsis tritonis*, - MAUCHLINE, J. nat. Hist., **18**: 317.
- 1984 *Pallenopsis (Bathypallenopsis) tritonis*, - STOCK, J. mar. biol. Ass. U.K. (1984): 747-748.
- 1991 *Pallenopsis (Bathypallenopsis) tritonis*, - STOCK, Bull. Mus. natn. Hist. nat., 4e sér., **13**, sec. A, nos. 1-2: 138-139.

Distribution: NE-Atlantic (Faroe Channel; NW-Ireland; entrance to English Channel; Newfoundland; New England).

Depth range: ?344-2884m.

***Pallenopsis truncatula* CHILD 1992**

- 1992 *Pallenopsis (Pallenopsis) truncatula* CHILD, Smiths. Contr. Zool., **526**: 27-29, fig. 11.

Distribution: Peru; Chile.

Depth range: 290-735m.

***Pallenopsis tumidula* LOMAN 1923**

- 1923 *Pallenopsis tumidula* LOMAN, Further Zool. Res. Swed. Antarct. Exped., **1** (2): 32-34, fig. G1-8.
- 1940 *Pallenopsis tumidula*, - MARCUS, Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 23-24 [key].
- 1957 *Pallenopsis tumidula*, - STOCK, Mitt. Hamburg. Zool. Mus. Inst., **55**: 103-105, fig. 18.
- 1966 *Pallenopsis tumidula*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 393, fig. 3e.
- 1992 *Pallenopsis (Pallenopsis) tumidula*, - STOCK, Tijdschr. Ent., **135**: 130, 139.

Distribution: off Rio de la Plata, Uruguay, to southern Brazil.

Depth range: 42-270m

***Pallenopsis tydemani* LOMAN 1908**

- 1908 *Pallenopsis tydemani* LOMAN, Siboga Exped. Monogr., **40**: 65-66, pl. X, figs. 139-145.
- 1949 *Pallenopsis tydemani*, - HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 277, fig. 36i-j.
- 1951 *Pallenopsis tydemani*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 160.
- 1971 *Pallenopsis tydemani*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 323.
- 1975 *Pallenopsis (Bathypallenopsis) tydemani caraibica*, - STOCK, Bull. mar. Sci., **24** (4): 1033-1036, figs. 31d, 32.
- 1982 *Pallenopsis (Bathypallenopsis) tydemani*, - CHILD, Smiths. Contr. Zool., **349**:

- 30-31.
- 1984 *Pallenopsis (Bathypallenopsis) tydemani caraibica*, - STOCK, J. mar. biol. Ass. U.K.: 748.
- 1986 *Pallenopsis (Bathypallenopsis) tydemani caraibica*, - STOCK, Bull. mar. Sci., 38 (3): 437.
- 1990 *Pallenopsis (Bathypallenopsis) tydemani*, - NAKAMURA & CHILD, Proc. biol. Soc. Wash., 103 (2): 308.
- 1991 *Pallenopsis (Bathypallenopsis) tydemani*, - NAKAMURA & CHILD, Smiths. Contr. Zool., 512: 40.
- 1991 *Pallenopsis (Bathypallenopsis) tydemani*, - STOCK, Mém. Mus. natn. Hist. nat. (A), 151: 224.

Distribution: Atlantic and Indo-Pacific except for Polar Seas.

Depth range: 100-2862m.

***Pallenopsis vanhoffeni* HODGSON 1915**

- 1915 *Pallenopsis vanhoffeni*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 43.
- 1927 *Pallenopsis vanhoffeni*, - HODGSON, Dt. Südpolar-Exped., 19, Zool. 11: 336-337.
- 1938 *Pallenopsis vanhoffeni*, - Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), 2 (8): 17-19, figs. 3b, 4a, 5c, 6c-d
- 1944 *Pallenopsis vanhoffeni*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 45.
- 1950 *Pallenopsis vanhoffeni*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 151.
- 1952 *Pallenopsis vanhoffeni*, - FAGE, Bull. Mus. natn. Hist. nat., ser. 2, 24 (2): 184.
- 1952 *Pallenopsis vanhoffeni*, - FAGE, Bull. Mus. natn. Hist. nat., ser. 2, 24 (3): 269-270.

Distribution: Antarctica (Ross Sea; off Victoria and Gauss Quadrants; Magellan District).

Depth range: 40-603m.

***Pallenopsis varioculata* STOCK 1986**

- 1986 *Pallenopsis (Pallenopsis) varioculata* STOCK, Bull. mar. Sci., 38 (3): 431-433, fig. 11.

Distribution: W-Atlantic (SE of Trinidad, off Surinam).
Depth range: 48-59m.

***Pallenopsis verrucosa* STOCK 1953**

- 1953 *Pallenopsis verrucosa* STOCK, Temminckia, 9: 288-291, figs. 8-9.
- 1968 *Pallenopsis verrucosa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., 131: 48.

Distribution: Philippines and Coral Sea off E-Australia.
Depth range: 55-80m.

Habitat: gravel.

Pallenopsis villosa HODGSON, 1907

- 1938 *Pallenopsis villosa*, - GORDON, Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 16, figs. 3c, 4b, 5a.
 1944 *Pallenopsis villosa*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), **5** (1): 45-46.

Distribution: Antarctica (Ross Sea; off Victoria and Gauss Quadrants).

Depth range: 180-540m.

Pallenopsis virgata LOMAN 1908

- 1908 *Pallenopsis virgatus* LOMAN, Siboga Exped. Monogr., **40**: 69-70, pl. 9, fig. 134, pl. 10, figs. 135-136.
 1949 *Pallenopsis virgatus*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 277-278, fig. 36g-h.
 1951 *Pallenopsis virgatus*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **1** (4): 160.
 1959 *Pallenopsis virgata*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 200.
 1965 *Pallenopsis virgata*, - UTINOMI, in: New Illustr. Encyclopaedia Fauna Japan, **2**: 335, fig. 17.
 1971 *Pallenopsis virgata*, - LOMAN, Publ. Seto Mar. Biol. Lab., **18** (5): 324.
 1983 *Pallenopsis (Pallenopsis) virgata*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 61.
 1987 *Pallenopsis virgata*, - NAKAMURA, Biol. Lab., Imperial Household: 13, pl. 10.
 1991 *Pallenopsis (Pallenopsis) virgata*, - CHILD, Smiths. Contr. Zool., **512**: 41.
 1991 *Pallenopsis (Pallenopsis) virgata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 197.
 1991 *Pallenopsis (Pallenopsis) virgata*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 223-224.

Distribution: Indonesia, Japan, Philippines, New Caledonia, Chesterfield Islands.

Depth range: 30-82m.

Philyra LOSINA-LOSINSKY 1961

- 1961 *Philyra* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 82 [diagnosis].

Philyra quadriarticulata LOSINA-LOSINSKY 1961

- 1961 *Philyra quadriarticulata* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 84-87, fig. 14.

Distribution: NW-Pacific.

Depth range: 40-50m.

Philyra robusta LOSINA-LOSINSKY 1961

1961 *Philyra robusta* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 82-84, fig. 13.

Distribution: NW-Pacific.

Depth range: 24m.

Phoxichilidium MILNE-EDWARDS 1840

- 1948 *Phoxichilidium*, - HEDPETH, Proc. U.S. natn. Mus., **97** (3216): 216 [diagnosis, remarks].
 1954 *Phoxichilidium*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 125 [diagnosis].
 1949 *Phoxichilidium*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 280-281 [remarks].

Phoxichilidium australe HODGSON 1914

- 1915 *Phoxichilidium australe*, - CALMAN, Brit. Antarct. (Terra Nova) Exped., 1910, Zool., **3** (1): 46-48, fig. 10.
 1927 *Phoxichilidium australe*, - BOUVIER, Dt. Südpolar-Exped., **19**, Zool. 11: 340-341.
 1932 *Phoxichilidium australe*, - GORDON, Discovery Rep., **6**: 92-93.
 1938 *Pallenopsis australe*, - Sci. Rep. Australas. Antarct. Exped., (C), (Zool. & Bot.), **2** (8): 19.
 1952 *Phoxichilidium australe*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (2): 184.
 1952 *Phoxichilidium australe*, - FAGE, Bull. Mus. natn. Hist. nat., 2e sér., **24** (3): 271.
 1991 *Phoxichilidium australe*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 202 [text].

Distribution: subantarctic area, probably circumpolar;
Tasmania.

Depth range: 15-400m.

Habitat: amongst scallops; Bryozoa.

Phoxichilidium capense FLYNN 1928

- 1928 *Phoxichilidium capense* FLYNN, Rep. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Rep., 1): 27, figs. 15-16.
 1954 *Phoxichilidium capense*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 126.

Distribution: South Africa.

Phoxichilidium femoratum (RATHKE 1799)

- 1799 *Nymphon femoratum* RATHKE, Skrift. Naturhist., Kobenhavn, **5** (1) (4): 201.
 1853 *Phoxichilidium maxillare* STIMPSON, Smiths. Contr. Knowl., **6** (5): 37.
 1874 *Phoxichilidium femoratum* VERRILL, Am. J. Sci. & Arts, **3** (7): 45.
 1878 *Phoxichilidium maxillare*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 12-

- 13, pl. 4, fig. 1a-e.
- 1878 *Phoxichilidium minor* WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 13-14, pl. 4, fig. 2a-f.
- 1880 *Phoxichilidium maxillare*, WILSON, Rep. U.S. Fish. Comm., **6**: 480-481, pl. 3, figs. 12-15.
- 1891 *Phoxichilidium femoratum*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 21-25, pl. II, fig. 1a-g.
- 1901 *Phoxichilidium maxillare*, - WHITEAVES, Geol. Surv. Canada, **772**: 263.
- 1914 *Phoxichilidium maxillare*, - PEARSE, Bull. Wisconsin Nat. Hist. Soc., N.S., **12**: 77.
- 1933 *Phoxichilidium femoratum*, - LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 47.
- 1933 *Phoxichilidium femoratum*, - STEPHENSEN, Medd. Grönloand, **79** (6): 26-27.
- 1936 *Phoxichilidium femoratum*, - EXLINE, Proc. U.S. Natn. Mus., **83** (2991): 421.
- 1943 *Phoxichilidium femoratum*, - NEEDLER, Can. Atl. Fauna, **10**: 14, fig. 18a-f.
- 1945 *Phoxichilidium femoratum*, - LEBOUR, J. mar. biol. Ass. U.K., **26**: 146, figs. 1 a-c, 2a.
- 1948 *Phoxichilidium femoratum*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 216-217, fig. 24a.
- 1961 *Phoxichilidium femoratum*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 79.
- 1974 *Phoxichilidium femoratum*, KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 36-37, fig. 15.
- 1975 *Phoxichilidium femoratum*, - STOCK, Ent. Ber., **35**: 74.
- 1993 *Phoxichilidium femoratum*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: N-Atlantic (Greenland and Norway to France and Long Island Sound); Pacific Coast of N-America.

Depth range: shallow water-101m.

Habitat: panels; in cysts on *Syncoryne eximia*; *Aglaophenia latirostris*.

***Phoxichilidium forfex* STOCK 1991**

- 1991 *Phoxichilidium forfex* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 202-204, fig. 52.

Distribution: New Caledonia.

Depth range: 530m.

***Phoxichilidium horribilis* HEDGPETH 1949**

- 1949 *Phoxichilidium horribilis* HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 283-284, fig. 39.
- 1991 *Phoxichilidium horribile*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 202 [text].

Distribution: Sea of Okhotsk north of Kurile chain.

Depth range: 419m.

***Phoxichilidium micropalpidum* HILTON 1942**

- 1942 *Phoxichilidium micropalpidum* HILTON, J. Ent. Zool. Pomona College, 34 (3): 72.
 1975 *Phoxichilidium micropalpidum*, - CHILD, Proc. biol. Soc. Wash., 88 (19): 202-205, fig. 5.

Distribution: Russian Arctic (Bering Island; Bering Sea).
Depth range: 132m.

Phoxichilidium parvum HILTON 1939

- 1939 *Phoxichilidium parvum* HILTON, J. Ent. Zool. Pomona College, 31 (2): 28.
 1941 *Phoxichilidium parvum*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., 9 (26): 257 [key].
 1942 *Phoxichilidium parvum*, - HILTON, J. Ent. Zool. Pomona College, 34 (3): 71-72.
 1954 *Phoxichilidium hokkaidoense* UTINOMI, Publ. Akkeshi Mar. Biol. Stat., 3: 4-7, figs. 2-3, pl. I fig. 1.
 1965 *Phoxichilidium hokkaidoensis*, - UTINOMI, in: New Illustr. Encyclopaedia Fauna Japan, 2: 335.
 1971 *Anoplodactylus hokkaidoensis*, - UTINOMI, Publ. Seto Mar. Biol. Lab., 18 (5): 326.
 1975 *Phoxichilidium parvum*, - CHILD, Proc. biol. Soc. Wash., 88 (19): 205, fig. 6a.
 1991 *Phoxichilidium parvum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., 512: 34.

Distribution: California; Japan.
Depth range: ?-381m.
Habitat: shallow-water hydroids.

Phoxichilidium quadradentatum HILTON 1942

- 1942 *Phoxichilidium quadradentatum* HILTON, J. Ent. Zool. Pomona College, 34 (3): 71.
 1954 *Phoxichilidium quadridentatum*, - HEDGPETH, in: Intertidal Inv. Central Calif coast, Univ. Calif. Press: 204.
 1955 *Phoxichilidium quadridentatum*, - MacGINITIE, Smiths. Misc. Coll., 128 (9): 171.
 1963 *Phoxichilidium quadradentatum*, - HEDGPETH, J. Fish. Res. Board Can., 20 (5): 1336-1337, fig. 9.
 1975 *Phoxichilidium quadradentatum*, - CHILD, Proc. biol. Soc. Wash., 88 (19): 205-207.

Distribution: NE-Pacific.
Habitat: stony mud, among hydroids and bryozoans.

Phoxichilidium spinosum LOSINA-LOSINSKY 1961

- 1961 *Phoxichilidium spinosum* LOSINA-LOSINSKY, Issledov. Dalnevost. Morei SSSR, Leningrad, 7: 79-80, fig. 11.
 1991 *Phoxichilidium spinosum*, - STOCK, Mém. Mus. natn. Hist. nat. (A), 151: 202 [text].

Distribution: NW-Pacific.
Depth range: 151-414m.

Phoxichilidium tuberculatum STOCK 1991

1991 *Phoxichilidium tuberculatum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 200-202, fig. 51.

Distribution: New Caledonia.
Depth range: 435-610m.

Phoxichilidium tubulariae LEBOUR 1945

- 1945 *Phoxichilidium tubulariae* LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 147-150, figs. 1d-g, 2b, 3a-c.
 1974 *Phoxichilidium tubulariae*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 35 [key], 38-39, fig. 16.
 1993 *Phoxichilidium tubulariae*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: NE-Atlantic.
Depth range: shallow water.
Habitat: on *Tubularia larynx*.

Phoxichilidium ungellatum HEDGPETH 1949

- 1949 *Phoxichilidium ungellatum* HEDGPETH, Proc. U.S. Natn. Mus., **98** (3231): 281-283, fig. 38.
 1954 *Phoxichilidium ungellatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 71.
 1955 *Phoxichilidium ungellatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 22-23, fig. 12.
 1971 *Phoxichilidium ungellatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 325.
 1990 *Phoxichilidium ungellatum*, - TURPAEVA, Zool. Zh., **69** (9): 17, fig. 1 (1-3).
 1991 *Phoxichilidium ungellatum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 34-35.

Distribution: Japan.
Depth range: 0-479m.
Habitat: hard bottom and surface plancton (0-5m).

Pycnosomia LOSINA-LOSINSKY 1961

1961 *Pycnosomia* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 86.

Pycnosomia asterophila STOCK 1981

- 1981 *Pycnosomia asterophila* STOCK, Res. Camp. MUSORSTOM, 1, Mém. ORSTOM, **91** (13): 309-312, figs. 1-8.
 1991 *Pycnosomia asterophila*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 225 [text].

Distribution: Philippines.

Depth range: 379-407m.

Habitat: On starfish *Calliaster corynetes*.

***Pycnosomia coxata* STOCK 1991**

- 1991 *Pycnosomia coxata* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 225-226, fig. 7.

Distribution: Philippines.

Depth range: 119-204m.

***Pycnosomia strongylocentroti* LOSINA-LOSINSKY 1933**

- 1933 *Pycnosoma strongylocentroti* LOSINA-LOSINSKY, Issledov. Morei S.S.S.R., Leningrad, **17**: 43.
 1942 *Pigrogromitus robustus* HILTON, J. Ent. Zool. Pomona College, **34** (2): 40.
 1947 *Pycnosoma strongylocentroti*, - HEDGPETH, Smiths. Misc. Coll., **106** (18): 7 [footnote].
 1961 *Pycnosomia strongylocentroti*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 86.
 1975 *Pycnosoma strongylocentroti*, - CHILD, Proc. biol. Soc. Wash., **88** (19): 207.

Distribution: Alaska and Russian Siberia.

Depth range: 16-75m.

***Pycnosomia tuberculata* LOSINA-LOSINSKY, 1961**

- 1961 *Pycnosomia tuberculata* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 86-87, fig. 15.

Distribution: NW-Pacific.

Depth range: 591m.

PYCNOGONIDAE

- 1948 Pycnogonidae, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 276 [diagnosis, key to genera].
 1954 Pycnogonidae, - BARNARD, Ann. S. Afr. Mus., **41** (3): 151 [diagnosis, remarks].
 1969 Pycnogonidae, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: [diagnosis, key to genera].
 1979 Pycnogonidae, - CHILD, Smiths. Contr. Zool., **293**: 72 [key to Middle American species].

Pentapycnon BOUVIER 1910

- 1948 *Pentapycnon*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 281 [diagnosis, remarks].
 1966 *Pentapycnon*, - STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 400 [key to species].

Pentapycnon charcoti BOUVIER 1910

- 1910 *Pentapycnon charcoti* BOUVIER, C. r. hebd. Séanc. Acad. Sci., Paris, **151**: 29.
 1911 *Pentapycnon charcoti*, - BOUVIER, C. r. hebd. Séanc. Acad. Sci., Paris, **152**: 492.
 1911 *Pentapycnon charcoti*, - BOUVIER, C. r. hebd. Séanc. Acad. Sci., Paris, **152**: 1137.
 1913 *Pentapycnon charcoti*, - BOUVIER, Deuxième Exped. Antarct. fr.: 98, 151-156, figs. 98-100.
 1935 *Pentapycnon charcoti*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordin. Tierreichs, **5**, Abt. IV, 2 (1): 298.
 1969 *Pentapycnon charcoti*, - FRY & HEDGPETH, Mem. N.Z. Oceanogr. Inst., **49**: 62-63, figs. 88, 92-95.

Distribution: Antarctica.

Depth range: 240-1420m.

Pentapycnon geayi BOUVIER 1911

- 1911 *Pentapycnon geayi* BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 491-494.
 1911 *Pentapycnon geayi*, - BOUVIER, C.R. Séanc. Hebd. Acad. Sci., Paris, **152**: 1140.
 1913 *Pentapycnon geayi*, - BOUVIER, Deuxième Expéd. Antarct. Française (1908-1910), **6**: 161-166, figs. 105-109.
 1948 *Pentapycnon geayi*, - HEDGPETH, Proc. U.S. Natn. Mus., **97** (3216): 281, fig. 53.
 1975 *Pentapycnon geayi*, - STOCK, Bull. mar. Sci., **24** (4): 1088.
 1986 *Pentapycnon geayi*, - STOCK, Bull. mar. Sci., **38** (3): 405 tab. 1.
 1992 *Pentapycnon geayi*, - STOCK, Tijdschr. Ent., **135**: 137, 139.

Distribution: French Guiana; Surinam; north of Puerto Rico; Brazil.

Depth range: 8-70m.

Pycnogonum BRÜNNICH 1764

- 1948 *Pycnogonum*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 277 [remarks, key to NW-Atlantic species].
 1954 *Pycnogonum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 151-152 [diagnosis, remarks, key to South African species].
 1963 *Pycnogonum*, - CLARK, Rec. Austr. Mus., **26** (1): 79 [key to Australian species].
 1966 *Pycnogonum*, - STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 400-402 [key to species].
 1968 *Pycnogonum*, - STOCK, Vidensk. Medd. Dansk naturh. Foren., **131**: 60-61

[remarks].

- 1969 *Pycnogonum*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 58 [remarks].
1986 *Pycnogonum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 64 [key to Japanese species].

***Pycnogonum africanum* CALMAN 1938**

- 1938 *Pycnogonum africanum* CALMAN, Sci. Rep. John Murray Exp., **5** (6): 163-165, fig. 10A-C.
1968 *Pycnogonum (Nulloviger) africanum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 61, fig. 22a-b.
1988 *Pycnogonum (Nulloviger) africanum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 150-151.
1991 *Pycnogonum (Nulloviger) africanum*, - TURPAEVA, Zool. Zh., **70** (12): 42.

Distribution: shelf and slope of south-eastern Africa.

Depth range: 183-1228m.

***Pycnogonum angulirostrum* STOCK 1959**

- 1959 *Pycnogonum angulirostrum* STOCK, Trans. Roy. Soc. S. Afr., **35** (5): 565-567, fig. 9.

Distribution: South Africa.

Depth range: littoral.

Habitat: under stone in pool.

***Pycnogonum anovigerum* CLARK 1956**

- 1956 *Pycnogonum anovigerum* CLARK, Rec. Canterbury Mus., **7** (2): 171-173, figs. 1-5.

Distribution: New Zealand.

Depth range: littoral.

Habitat: from underside of boulder in rockpool.

***Pycnogonum arbustum* STOCK 1966**

- 1966 *Pycnogonum arbustum* STOCK, Atlantide Rep., **9**: 54-56, fig. 4.

Distribution: W-Africa (off Guinea).

Depth range: 45m.

Habitat: shells, Foraminifera.

***Pycnogonum asiaticum* MÜLLER 1992**

- 1992 *Pycnogonum asiaticum* MÜLLER, Bonn. zool. Beitr., **43** (1): 173-177, figs. 50-55.

Distribution: Malaysia (Tioman Archipelago, Pulau Babi

Besar)

Depth range: 1-2 m.

Habitat: dead corals on fringing reef.

Pycnogonum aurilineatum FLYNN 1919

- 1919 *Pycnogonum aurilineatum* FLYNN, Pap. Proc. Roy. Soc. Tasm., 1918: 92-95, pl. XIII, figs. 1-2, pl. XIV, fig. 3.
 1973 *Pycnogonum aurilineatum*, - STOCK, Beaufortia, 20 (266): 125.

Distribution: Australia.

Depth range: 10-15m.

Habitat: in red algae.

Pycnogonum benokianum OHSIMA 1935

- 1935 *Pycnogonum benokianum* OHSIMA, Dōbutsugaku Zasshi, 47 (557): 137-139.
 1947 *Pycnogonum benokianum*, - OHSIMA & KISHIDA, Illustr. Encyclopaedia Fauna Japan: 1010, fig. 2866.
 1949 *Pycnogonum benokianum*, - HEDGPETH, Proc. U.S. natn. Mus., 98 (3231): 304-307, fig. 49.
 1971 *Pycnogonum benokianum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., 18 (5): 339.

Distribution: Japan (Benoki, Okinawa Islands).

Habitat: infesting sea anemone *Phellia decora*.

Pycnogonum buticulosum HEDGPETH 1949

- 1949 *Pycnogonum buticulosum* HEDGPETH, Proc. U.S. natn. Mus., 98 (3231): 308-309, fig. 48a, 50a-b.

Distribution: Japan.

Pycnogonum callosum LOSINA-LOSINSKY 1961

- 1961 *Pycnogonum callosum* LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, 7: 110-111, fig. 26.

Distribution: NW-Pacific.

Depth range: 200-400m.

Pycnogonum cataphractum MÖBIUS 1902

- 1902 *Pycnogonum cataphractum* MÖBIUS, Wiss. Ergebni. dtsch. Tiefsee-Exp. "Valdivia", 1898-1899, 3: 194, pl. XXX, fig. 11.
 1954 *Pycnogonum cataphractum*, - BARNARD, Ann. S. Afr. Mus., 41 (3): 152.

Distribution: South Africa (St. Francis Bay).

Depth range: shallow water.

Pycnogonum cessaci BOUVIER 1911

- 1911 *Pycnogonum cessaci* BOUVIER, Compt. Rend. Acad. Sci., **152**: 493.
 1940 *Pycnogonum pamphorum* MARCUS, - Bol. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, **19** (Zool. 4): 115-117, fig. 16.
 1945 *Pycnogonum leticiae*, - MELLO-LEITAO, Bol. Mus. Nac., Rio de Janeiro, N.S. (Zool.), **42**: 1-4, figs. 1-5.
 1952 *Pycnogonum cessaci*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **24** (6): 531-533, fig. 1.
 1954 *Pycnogonum cessaci*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **5** (24): 129.
 1966 *Pycnogonum cessaci*, - STOCK, Ann. Inst. Océanogr. Monaco, **440**: 400 [key].
 1967 *Pycnogonum cessaci*, - McCLOSKEY, J. nat. Hist., **1**: 131-133, figs. 25-27.
 1975 *Pycnogonum cessaci*, - STOCK, Bull. mar. Sci., **24** (4): 1085-1086, fig. 58d.
 1979 *Pycnogonum cessaci*, - CHILD, Smiths. Contr. Zool., **293**: 72.
 1986 *Pycnogonum cessaci*, - STOCK, Bull. mar. Sci., **38** (3): 405 tab. 1.
 1990 *Pycnogonum cessaci*, - STOCK, Zool. Meded., **63** (16): 231.
 1992 *Pycnogonum (Retroviger) cessaci*, - STOCK, Tijdschr. Ent., **135**: 136, 139, 136-137 [text].
 1992 *Pycnogonum leticiae*, - STOCK, Tijdschr. Ent., **135**: 137 [text].

Distribution: amphi-atlantic, in tropical and temperate waters; Pacific-Coast of Panamá.

Depth range: 0-10m.

Habitat: on scleractinian coral *Oculina arbuscula*; rock debris and sand; rocky and sandy coast; seaweeds on rocky shore.

Pycnogonum crassirostre SARS 1888

- 1888 *Pycnogonum crassirostre* SARS, Arch. Math. Naturvidensk., Oslo, **12**: 340.
 1891 *Pycnogonum crassirostre*, - SARS, Norw. North-Atl. Exp., 1876-1878 (Zool. **20**): 12-14, pl. 1, fig. 1A-H.
 1933 *Pycnogonum crassirostre*, - STEPHENSEN, Danmarks Fauna, **38**: 30, fig. 8 (map).
 1936 *Pycnogonum crassirostre*, - STEPHENSEN, Bergens Mus. Arbok Naturv. Rek., **2**: 35.
 1948 *Pycnogonum crassirostre*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 279, fig. 52B.
 1966 *Pycnogonum crassirostre*, - STOCK, Ann. Inst. Océanogr., Monaco, **44**: 402 [key].
 1975 *Pycnogonum crassirostrum*, - STOCK, Bull. mar. Sci., **24** (4): 1086.
 1988 *Pycnogonum crassirostrum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 151.

Distribution: off Iceland, Greenland, Norway, eastern N-America, South Africa.

Depth range: 100-620m.

Habitat: dead corals; *Lophohelia*.

Pycnogonum crosnieri STOCK 1991

1991 *Pycnogonum crosnieri* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 206-208, fig. 56.

Distribution: New Caledonia.
Depth range: 440-450m.

Pycnogonum elephas STOCK 1966

1966 *Pycnogonum elephas* STOCK, Ann. Inst. Oceanogr. Monaco, **44**: 397-400, fig. 5.

Distribution: W-Atlantic (between Rio de Janeiro and Rio de la Plata).
Depth range: 30-740m.

Pycnogonum eltanin FRY & HEDGPETH 1969

1969 *Pycnogonum eltanin* FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **40**: 61-62, figs. 85, 90, 91.

Distribution: Antarctica.
Depth range: 300m.

Pycnogonum forte FLYNN 1928

1928 *Pycnogonum forte* FLYNN, Repts. Fish. & Mar. Biol. Surv. Union S. Afr., **6** (Special Repts. 1): 31-33, figs. 18-19.

1954 *Pycnogonum forte*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 152-153, fig. 31C-D.

1998 *Pycnogonum forte*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 151-152.

Distribution: South Africa.
Depth range: 90-775m

Pycnogonum gaini BOUVIER 1910

1910 *Pycnogonum gaini* BOUVIER, C. r. hebd. Séanc. Acad. Sci., Paris, **151**: 30.

1913 *Pycnogonum gaini*, - BOUVIER, Deuxième Exped. Antarct. fr.: 156-160, figs. 101-104.

1915 *Pycnogonum gaini*, - CALMAN, Br. Antarct. Terra Nova Exped., 1910, Zool. **3** (1): 68, fig. 22.

1927 *Pycnogonum gaini*, - HODGSON, Dt. Südpolar-Expe., **19**: 354-356.

1935 *Pycnogonum gaini*, - HELFER & SCHLÖTTKE, Bronn's Kl. Ord. Tierreichs, **5**, Abt. IV, 2, (1): 298.

1938 *Pycnogonum gaini*, - GORDON, Sci. Rep. Australas. Antarct. Exped., ser. C., (Zool. and Bot.) **2** (8): 28.

1944 *Pycnogonum gaini*, - GORDON, Rep. B.A.N.Z. Antarct. Res. Exped., ser. B, **5** (1): 68-69, fig. 26a-d, 27a.

1969 *Pycnogonum gaini*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **40**: 58-59, figs. 83-84, map. 85.

1972 *Pycnogonum gaini*, - ARNAUD, Téthys, suppl. **4**: 145.

1980 *Pycnogonum gaini*, - KRAPF, Bull. Inst. Roy. Sci. Nat. Belg., **52** (18): 7.

Distribution: Antarctica.

Depth range: upper sublittoral-2495m.

***Pycnogonum gibberum* MARCUS & du BOIS REYMOND-MARCUS 1963**

- 1963 *Pycnogonum gibberum* MARCUS & du BOIS REYMOND-MARCUS, Bol. Inst. Oceanogr. S Paulo, **12** (3): 3-6, figs. 1-5.
1992 *Pycnogonum (Retroviger) gibberum*, STOCK, Tijdschr. Ent., **135**: 136, 139.

Distribution: Brazil.

Depth range: 32-56m.

***Pycnogonum gordonaee* PUSHKIN 1984**

- 1984 *Pycnogonum gordonaee* PUSHKIN, Zool. Zh., **63** (7): 1102-1104, figs. 1-15.

Distribution: Antarctic.

Depth range: 260-400m.

***Pycnogonum guyanae* STOCK 1975**

- 1975 *Pycnogonum guyanae* STOCK, Bull. mar. Sci., **24** (4): 1086-1088, fig. 59a-e.
1986 *Pycnogonum guyanae*, - STOCK, Bull. mar. Sci., **38** (3): 405.

Distribution: French Guiana.

Depth range: 104-130m.

Habitat: sandy calcarenite; from gorgonarians.

***Pycnogonum hancocki* SCHMITT 1934**

- 1934 *Pycnogonum hancocki* SCHMITT, J. Wash. Acad. Sci., **24** (1): 65-67, fig. 2.
1942 *Pycnogonum hancocki*, - HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 308-310, pl. 47.
1943 *Pycnogonum hancocki*, - HILTON, J. Ent. Zool. Pomona College, **35** (2): 19.
1966 *Pycnogonum hancocki*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 402 [key].
1971 *Pycnogonum hancocki*, - CHILD & HEDGPETH, J. nat. Hist., **5**: 631-633.

Distribution: Galápagos Archipelago.

Depth range: intertidal.

Habitat: under rocks.

***Pycnogonum indicum* SUNDARA RAJ 1930**

- 1930 *Pycnogonum indicum* SUNDARA RAJ, Bull. Madras. Governm. Mus. N.H., **1**: 73-75, figs. IA-B, II, figs. A-B.

Distribution: India (Shingle Island, Gulf of Manaar).

Depth range: intertidal.

Pycnogonum koreanum KIM & STOCK 1984

- 1984 *Pycnogonum koreanum* KIM & STOCK, J. nat. Hist., **18**: 685-688, figs. 1-6.
 1987 *Pycnogonum koreanum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.

Distribution: Korea.

Depth range: intertidal.

Habitat: under stones and in crevices of sandy rocks.

***Pycnogonum littorale* (STRÖM 1762)**

- 1762 *Pycnogonum littorale* STRÖM, Phys. Oecon. Beskr. Sondmor, beliggende Bergens Stift i Norge, **1**: 209, pl. 1, fig. 17.
 1853 *Pycnogonum pelagicum*, - STIMPSON, Smiths. Contr. Knowledge, **6** (5): 37.
 1872 *Pycnogonum pelagicum*, - WHITEAVES, Ann. Mag. nat. Hist, ser. **4**, 10: 347.
 1873 *Pycnogonum littorale*, - VERRILL, Rep. U.S. Commissioner Fish., **1**: 415.
 1874 *Pycnogonum littorale*, - SMITH & HARGER, Trans. Conn. Acad. Arts & Sci., **3**: 10.
 1874 *Pycnogonum pelagicum*, - VERRILL, Am. J. Sci. & Arts, **3** (7): 502.
 1875 *Pycnogonum littorale*, - VERRILL, Am. J. Sci. & Arts, **3** (10): 38.
 1878 *Pycnogonum littorale*, - WILSON, Trans. Conn. Acad. Arts & Sci., **5**: 4-5, pl. 1, fig. 1a-b, pl. 2, fig. 3a-b.
 1880 *Pycnogonum littorale*, - WILSON, Rep. U.S. Commissioner Fish., 1878 (6): 469-471, pl. 1, figs. 1-3.
 1881 *Pycnogonum littorale*, - WILSON, Bull. Mus. Comp. Zool., Harvard, **8** (12): 242.
 1885 *Pycnogonum littorale*, - VERRILL, Rep. U.S. Commission Fish., 1883: 560.
 1891 *Pycnogonum littorale*, - SARS, Norw. North-Atl. Exped., 1876-1878, **6** (Zool. 20): 7-12, pl. I, figs. 1a-i.
 1896 *Pycnogonum littorale*, - CAULLERY, Ann. Univ. Lyon, **26**: 363.
 1902 *Pycnogonum littorale*, - MÖBIUS, Wiss. Ergeb. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 193.
 1913 *Pycnogonum littorale*, - SUMNER, OSBURN & COLE, Bull. Bureau Fish., Washington, **31** (1,2): 677.
 1914 *Pycnogonum littorale*, - BOUVIER, J. Mar. Biol. Ass., Plymouth, **10**: 207-210.
 1923 *Pycnogonum littorale*, - BOUVIER, Faune de France, **7**: 61.
 1930 *Pycnogonum littorale*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **2**: 7-15.
 1933 *Pycnogonum littorale*, - STEPHENSEN, Godthaab Exp. 1928, Medd. Gronland, **79** (6): 30, fig. 8 (map).
 1936 *Pycnogonum littorale*, - STEPHENSEN, Bergens Mus. Arbok Naturv. Rek., **2**: 33-34.
 1943 *Pycnogonum littorale*, - NEEDLER, Can. Atl. Fauna, **10**: 5, fig. 3a-d.
 1945 *Pycnogonum littorale*, - LEBOUR, J. Mar. Biol. Ass. U.K., **26**: 163-164.
 1948 *Pycnogonum littorale*, - HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 277.
 1972 *Pycnogonum littorale*, - KING, J. nat. Hist., **6**: 622.
 1973 *Pycnogonum littorale*, - ARNAUD, Tethys, **5** (1): 152.
 1973 *Pycnogonum littorale*, - JARVIS & KING, Mar. Behav. Physiol., **2**: 1-13 [ultrastructure of photoreceptors].
 1974 *Pycnogonum littorale*, - KING, A Synopsis of the British Sea Spiders. Synopses of the British Fauna (New Ser.), **5**: 56-58, figs. 24-25.
 1976 *Pycnogonum littorale*, - WOLFF, Netherlands J. Sea Res., **10** (4): 476-477, fig. 4a (distribution in estuary).
 1978 *Pycnogonum littorale*, - KING & EL-HAWAII, Acta zool., **59**: 97-103 [spermiogenesis].

- 1979 *Pycnogonum littorale*, - JARVIS & KING, Mar. Behav. Physiol., **6**: 307-317 [response to light stimuli].
 1980 *Pycnogonum littorale*, - HOOPER, Aust. Zool., **20** (3): 477-478, 480 [text].
 1984 *Pycnogonum littorale*, - BEHRENS, Zoomorphology, **104**: 266-279 (larval-development, feeding).
 1985 *Pycnogonum littorale*, - BAMBER, Zool. J. Linn. Soc., **83**: 306.
 1993 *Pycnogonum littorale*, - HARMS, Helgoländer Meeresuntersuch., **47**: 24, tab. 3.

Distribution: N-Atlantic (Iceland and Greenland to west coast of France, Mediterranean and E-coast of U.S.A.)

Depth range: 0-800m.

Habitat: on hydroids (*Clava multicornis*) and Anthozoa (*Actinia equina*, *Metridium senile*, *Tealia felina*); associated with *Tealia*.

Remarks: Breeding season from late spring to early autumn.

***Pycnogonum lobipes* STOCK 1991**

- 1991 *Pycnogonum (Nulloviger) lobipes* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 208-210, fig. 57.

Distribution: New Caledonia.

Depth range: 680-700m.

***Pycnogonum madagascariensis* BOUVIER 1911**

- 1971 *Pycnogonum madagascariensis*, - ARNAUD, Tethys, Suppl. **1**: 161-163, figs. 1-6.

Distribution: Madagascar.

***Pycnogonum magellanicum* HOEK 1898**

- 1898 *Pycnogonum magellanicum* HOEK, Tijdschr. Nederl. Dierk. Ver., (2) **5** (2-4): 296, pl. III, figs. 20-21.
 1902 *Pycnogonum magellanicum*, - MÖBIUS, Fauna Arctica, **2**: 60-61.
 1932 *Pycnogonum magellanicum*, - GORDON, Discovery Rep., **6**: 126-128, fig. 72.
 1966 *Pycnogonum magellanicum*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 397, 402.

Distribution: off Falkland Islands and Gough Island.

Depth range: 102-236m.

***Pycnogonum magnirostre* MÖBIUS 1902**

- 1902 *Pycnogonum magnirostre* MÖBIUS, Wiss. Ergebni. dt. Tiefsee-Exped. "Valdivia", 1898-1899, **3**: 194-195, pl. XXX, figs. 12-14.

Distribution: Kerguelen Islands.

Depth range:

***Pycnogonum microps* LOMAN 1904**

- 1904 *Pycnogonum microps* LOMAN, Zool. Jb. Syst., **20** (4): 378, pl. 14, figs. 5-6.
 1954 *Pycnogonum microps*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 154-155, fig. 33.
 1956 *Pycnogonum microps*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **118**: 93.
 1962 *Pycnogonum microps*, - STOCK, Trans. Roy. Soc. S. Afr., **36** (4): 285.

Distribution: South Africa.

Depth range: 15m.

Habitat: shelly khaki sand, rocks.

Pycnogonum minutum LOSINA-LOSINSKY & KOPANEVA 1973

- 1973 *Pycnogonum minutum* LOSINA-LOSINSKY & KOPANEVA, Zool. Zh., **52** (7): 1083-1084, figs. 1-2.

Distribution: Guinean Republic.

Habitat: offshore.

Pycnogonum moolenbecki STOCK 1992

- 1992 *Pycnogonum moolenbecki* STOCK, Bijdr. Dierk., **62** (2): 95-97, fig. 11.

Distribution: Oman.

Depth range: low tide.

Habitat: "stones and algae" (STOCK).

Pycnogonum moniliferum STOCK 1991

- 1991 *Pycnogonum (Nulloviger) moniliferum* STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 210-211, fig. 58.

Distribution: New Caledonia.

Depth range: 570-825m.

Pycnogonum nodulosum DOHRN 1881

- 1881 *Pycnogonum nodulosum* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 203-207, pl. 16, figs. 1-3.
 1928 *Pycnogonum nodulosum*, - LOMAN, Bull. Soc. Sci. nat. Phys. Maroc., **8** (1-3): 62, 64.
 1953 *Pycnogonum nodulosum*, - FAGE, Bull. Mus. natn. Hist. nat., Paris, (2) **25** (4): 381 [list].
 1954 *Pycnogonum nodulosum*, - BOURDILLON, Rec. Trav. Stat. Mar. d'Endoume, **12** (7): 155.
 1955 *Pycnogonum nodulosum*, - BARNARD, Ann. S. Afr. Mus., **43** (1): 106.
 1958 *Pycnogonum nodulosum*, - STOCK, Bull. Sea Fish. Res. Sta. Haifa, **16**: 5.
 1966 *Pycnogonum nodulosum*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 401 [key].
 1968 *Pycnogonum nodulosum*, - STOCK, Vie Milieu, Suppl. **19** (1A): 34-35, fig. 28.
 1978 *Pycnogonum nodulosum*, - CHIMENZ-GUSSO, FRESI, CINELLI, MAZZELLA, PANSINI & PRONZATO, Mem. Biol. Mar. Ocean., **8** (4): 100.

- 1981 *Pycnogonum nodulosum*, - MUNILLA & de HARO, Bijdr. Dierk., **51** (2): 191-197, figs. 9, 11, tab. I (electrophoretical and immunological study).
- 1984 *Pycnogonum nodulosum*, - MUNILLA & de HARO, in: Els sistemes de les illes Medes, ROS, OLIVELLA & GILI eds., Barcelona: 532, tab. I, 534-535, tab. III-IV.
- 1988 *Pycnogonum nodulosum*, - ARNAUD & CHILD, Ann. S. Afr. Mus., **98** (6): 152-153.

Distribution: South Africa; W-Africa; Mediterranean.

Depth range: 9-49m.

Habitat: in *Posidonia oceanica* seagrass; *Lithothamnion racemus*, *Pessonnellia polymorpha*.

***Pycnogonum occa* LOMAN 1908**

- 1908 *Pycnogonum occa* LOMAN, Siboga Exped. Monogr., **40**: 35-36, pl. 12, figs. 171-174.
- 1922 *Pycnogonum occa*, - BOUVIER, Ann. Sci. nat., Paris (Zool.), (10) **5** (1-2): 116.
- 1966 *Pycnogonum occa*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 61-62, fig. 22c-e.
- 1988 *Pycnogonum occa*, - CHILD, Smiths. Contr. Zool., **468**: 27.

Distribution: ?Kermadec Trench; Indonesia; Philippines.

Depth range: 567-2470m.

Habitat: grey clay.

***Pycnogonum ornans* STOCK 1992**

- 1992 *Pycnogonum ornans* STOCK, Tijdschr. Ent., **135**: 136-137, 139, figs. 76-80.

Distribution: southern Brazil.

Depth range: 38-47m.

***Pycnogonum panamum* HILTON 1942**

- 1849 ?*Pycnogonum littorale*, - NICOLET, Hist. fisica y politica Chile, **3**. Atlas zool., Crust.: 308.
- 1942 *Pycnogonum panamum* HILTON, Rep. Allan Hancock Pac. Exped., **5** (9): 310-311, pl. 47.
- 1961 *Pycnogonum panamum*, - HEDGPETH, Lunds Univ. Arsskr., N.F. (2) **57** (3): 17, fig. 11.
- 1966 *Pycnogonum panamum*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 402 [key].

Distribution: Panamá Pacific; ?Chile.

Depth range: shallow water.

***Pycnogonum paragaini* MUNILLA 1989**

- 1989 *Pycnogonum paragaini* MUNILLA, Actas Tercer. Symp. Espan. Estud. Antarct.: 76-78, fig. 2.

Distribution: Antarctica.
Depth range: 205-230m.

Pycnogonum planum STOCK 1954

1954 *Pycnogonum planum* STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 165-168, fig. 81.

Distribution: New Zealand.
Depth range: littoral
Habitat: shore, sandstone.

Pycnogonum platylophum LOMAN 1923

- 1923 *Pycnogonum platylophum* LOMAN, Ark. Zool., **15** (10): 10-12, fig. E1-5.
 1932 *Pycnogonum platylophum*, - GORDON, Discovery Rep., **6**: 126, fig. 71a-b.
 1935 *Pycnogonum platylophum*, - HELFER & SCHLOTTKE, Bronn's Kl. Ord. Tierreichs, **5**, Abt. IV, 2 (1): 298.
 1938 *Pycnogonum platylophum*, - GORDON, Sci. Rep. Australas. Antarct. Exped. (C), (Zool. & Bot.), **2** (8): 28.
 1944 *Pycnogonum platylophum*, - GORDON, Rep. B.A.N.Z. Antarct. Res. Exped., ser. B, **5** (1): 69-70, fig. 27b-d.
 1947 *Pycnogonum (magellanicum HOEK ?)*, - STEPHENSEN, - Sci. Res. Norw. Antarct. Exped., 1927-1928 et sqq., **27**: 86.
 1969 *Pycnogonum platylophum*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 61, figs. 88-89.
 1970 *Pycnogonum platylophum*, - ARNAUD, Bull. Mus. natn. Hist. nat., 2e sér., **41** (6): 1426.

Distribution: (sub)-Antarctic.
Depth range: 18-540m.

Pycnogonum plumipes STOCK 1960

- 1960 *Pycnogonum plumipes* STOCK, Vie Milieu, **11** (1): 124-126, fig. 1a-c.
 1968 *Pycnogonum plumipes*, - STOCK, Vie Milieu, **19** (1A): 35, fig. 29.
 1987 *Pycnogonum plumipes*, - ARNAUD, Mesogee, **47**: 50-51.

Distribution: Mediterranean.
Depth range: 2-40m.
Habitat: *Paramuricea*, *Corallium*, *Leptopsammia*, *Halopteris scoparia*, hydroids.

Pycnogonum portus BARNARD 1946

- 1946 *Pycnogonum portus* BARNARD, Ann. Mag. Nat. Hist., **11** (13): 62.
 1954 *Pycnogonum portus*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 153-154, fig. 32.

Distribution: South Africa.
Depth range: littoral.

***Pycnogonum pusillum* DOHRN 1881**

- 1881 *Pycnogonum pusillum* DOHRN, Monogr. Fauna Flora Golf Neapel, **3**: 207, pl. 16, figs. 4-8.
- 1885 *Pycnogonum pusillum*, - CARUS, Prodr. faunae medit. sive descr. maris Medit. incalarum, **2**: 288.
- 1909 *Pycnogonum pusillum*, d'ARCY THOMPSON, The Cambr. Nat. Hist., 540.
- 1922 *Pycnogonum pusillum*, - BOUVIER, Ann. Sci. Nat., Paris (Zool.), (10) **5** (1-2): 115.
- 1923 *Pycnogonum pusillum*, - BOUVIER, Faune de France, **7**: 61.
- 1929 *Pycnogonum pusillum*, - SCHIUMKEWITSCH, Fauna SSSR, Izdatljstvo Akad. Nauk SSSR, **1**: 6, 7, 18.
- 1935 *Pycnogonum pusillum*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 298.
- 1952 *Pycnogonum pusillum*, - STOCK, Bull. Inst. Roy. Sci. Nat. Belg., **38** (14): 1-6, fig. 1a-b.
- 1954 *Pycnogonum pusillum*, - BARNARD, Ann. S. Afr. Mus., **41** (3): 155-156, fig. 33.
- 1954 *Pycnogonum pusillum*, - BOURDILLON, Rec. Trav. Sta. Mar. d'Endoume, **12** (7): 155.
- 1955 *Pycnogonum pusillum*, - HOENIGMAN & STOCK, Ent. Ber., **15**: 534.
- 1962 *Pycnogonum pusillum*, - MONNIOT, Vie Milieu, **13** (2): 273.
- 1964 *Pycnogonum pusillum*, - STOCK, Bull. Sea Fish. Res. Sta. Haifa, **35**: 32.
- 1968 *Pycnogonum pusillum*, - STOCK, Vie Milieu, **19** (1A), Suppl.: 36, fig. 30.
- 1968 *Pycnogonum littorale*, - ZAVODNIK, Thalasia Jugosl., **4**: 46, 49.
- 1983 *Pycnogonum pusillum*, - KRAPP, Bonn. zool. Beitr., **34** (1-3): 412.
- 1987 *Pycnogonum pusillum*, - ARNAUD, Mesogee, **47**: 51.
- 1990 *Pycnogonum (Retroviger) pusillum*, - STOCK, Zool. Meded., **63** (16): 232.

Distribution: Western Mediterranean; Adriatic Sea; Morocco; Angola; South Africa (False Bay); Cape Verde Islands.

Depth range: 0.5-1200m.

Habitat: *Dictyota dichotoma*; *Cystoseira stricta*, *abrotanifolia* and *barbata*; *Halopteris scaparia*.

***Pycnogonum reticulatum* HEDGPETH 1948**

- 1947 *Pycnogonum* sp., HEDGPETH, Smiths. Misc. Coll., 106 (18): 13, fig. 5c.
- 1948 *Pycnogonum reticulatum* HEDGPETH, Proc. U.S. natn. Mus., **97** (3216): 279-281, fig. 52c-f.
- 1954 *Pycnogonum reticulatum*, - HEDGPETH, Fish Bull. Fish & Wildlife Service U.S., **89** (55): 427.
- 1954 *Pycnogonum reticulatum*, - STOCK, Stud. Fauna Curaçao and other Caribb. Isl., **24**: 129.
- 1968 *Pycnogonum reticulatum*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 62.
- 1975 *Pycnogonum reticulatum*, - STOCK, Bull. mar. Sci., **24** (4): 1085.
- 1979 *Pycnogonum reticulatum*, - CHILD, Smiths. Contr. Zool., **293**: 73.
- 1986 *Pycnogonum reticulatum*, - STOCK, Bull. mar. Sci., **38** (3): 405.

Distribution: Florida; Caribbean coast of Mexico and Panamá; Venezuela; El Salvador and Panamá Pacific.

Depth range: 1.5m.

Habitat: on sponge; on coral.

Pycnogonum rhinoceros LOMAN 1923

- 1923 *Pycnogonum rhinoceros* LOMAN, Ark. Zool., **15** (9): 7-9, fig. A.
 1932 *Pycnogonum rhinoceros*, - GORDON, Discovery Rep., **6**: 125, fig. 71c.
 1935 *Pycnogonum rhinoceros*, - HELFER & SCHLOTTKE, Bronn's Kl. Ordn. Tierreichs, **5** (4) (2): 298.
 1969 *Pycnogonum rhinoceros*, - FRY & HEDGPETH, N.Z. Oceanogr. Inst. Mem., **49**: 59-61, figs. 86-88.

Distribution: Antarctic (South Georgia, South Shetlands and Weddell Sea).

Depth range: 154-1115m.

Pycnogonum rickettsi SCHMITT 1934

- 1934 *Pycnogonum rickettsi* SCHMITT, J. Wash. Akad. Sci., **24** (1): 62-65, fig. 1a-d.
 1951 *Pycnogonum rickettsi*, - HEDGPETH, Wasmann J. Biol., **9** (1): 116.
 1980 *Pycnogonum rickettsi*, - HOOPER, Aust. Zool., **20** (3): 475.

Distribution: Australia.

Depth range: 0-2.5m.

Habitat: *Sargassum*.

Pycnogonum sivertseni STOCK 1955

- 1955 *Pycnogonum sivertseni* STOCK, Res. Norw. Sci. Exped. Tristan da Cunha 1937-1938, **33**: 7-10, figs. 3-5.

Distribution: Tristan da Cunha.

Depth range: 102-141m.

Pycnogonum stearnsi IVES 1892

- 1892 *Pycnogonum stearnsi* IVES, Proc. Acad. Nat. Sci. Philadelphia 1892-1893: 142-144, pl. 10.
 1904 *Pycnogonum stearnsi*, - COLE, Harriman Alaska Exped., **10**: 251, 259, 292-294, pl. 14, figs. 13-15.
 1913 *Pycnogonum stearnsi*, - HALL, Univ. Calif. Publ. Zool., **11** (6): 133 [key].
 1915 *Pycnogonum stearnsi*, - HILTON, J. Ent. Zool. Pomona College, **7** (1): 69.
 1915 *Pycnogonum stearnsi*, - HILTON, J. Ent. Zool. Pomona College, **7** (3): 202-203, 205-206.
 1920 *Pycnogonum stearnsi*, - HILTON, J. Ent. Zool. Pomona College, **12** (4): 93.
 1933 *Pycnogonum stearnsi*, - OHSHIMA, Bull. Biogeogr. Soc. Japan, **4**: 147-149, fig. 2.
 1934 *Pycnogonum stearnsi*, - SCHMITT, J. Wash. Acad. Sci., **24** (1): 69.
 1939 *Pycnogonum stearnsi*, - HILTON, J. Ent. Zool. Pomona College, **31** (2): 34.
 1940 *Pycnogonum stearnsi*, - HEDGPETH, J. Wash. Acad. Sci., **30** (2): 86-87.
 1941 *Pycnogonum stearnsi*, - HEDGPETH, Trans. San Diego Soc. Nat. Hist., **9** (26): 254 [key], pl. 10 [part.].
 1943 *Pycnogonum stearnsi*, - HILTON, J. Ent. Zool. Pomona College, **35** (2): 19.

- 1949 *Pycnogonum stearnsi*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 307 [text].
 1951 *Pycnogonum stearnsi*, - HEDGPETH, Wasm. J. Biol., **9** (1): 115-116.
 1960 *Pycnogonum stearnsi*, - ZIEGLER, The Veliger, **3** (1): 21.
 1964 *Pycnogonum stearnsi*, - HEDGPETH, in: Intertidal Invertebrates of the Central Calif. Coast: 209 [key].
 1966 *Pycnogonum stearnsi*, - STOCK, Ann. Inst. Océanogr., **44**: 402 [key].
 1979 *Pycnogonum stearnsi*, - CHILD, Smiths. Contr. Zool., **293**: 73.

Distribution: Mexico Pacific; California; Kurile Islands.

Pycnogonum tenue KISHIDA 1927

- 1879 *Pycnogonum littorale* var. *tenue* SLATER, Ann. Mag. Nat. Hist., (5) **3** (16): 281-283.
 1890 *Pycnogonum littorale* var. *tenue*, - ORTMANN, Zool. Jb. Syt., **5** (1): 167.
 1899 *Pycnogonum littorale* var. *tenue*, - SHISHIDO, Dôzatsu, **11** (128): 200.
 1919 *Pycnogonum littorale* var. *tenue*, - FUKUI, Rigakukai(?), **16** (2): 99.
 1927 *Pycnogonum tenue* KISHIDA, in: Nikon Dôbutsu tsaran: 989, fig. 1905.
 1929 *Pycnogonum littorale* var. *tenue*, - SCHIMKEWITSCH, Fauna SSSR, Izdateljstvo Akad. Nauk SSSR, **1**: 15.
 1936 *Pycnogonum tenue*, - OHSHIMA, Dôbutsugaku Zasshi, **48** (8-10): 867.
 1947 *Pycnogonum tenue*, - OHSHIMA & KISHIDA, - in: Illustr. Encyclopaedia Fauna Japan: 1010, fig. 2865.
 1949 *Pycnogonum tenue*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 303-304, fig. 48b, 50c-d.
 1954 *Pycnogonum tenue*, - STOCK, Vidensk. Medd. dansk naturhist. Foren., **116**: 162.
 1955 *Pycnogonum tenue*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **5** (1): 36-38, figs. 22-23.
 1959 *Pycnogonum tenue*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **7** (2): 216.
 1965 *Pycnogonum tenue*, - UTINOMI, in: New illustr. Encyclopaedia Fauna Japan, **2**: 338.
 1966 *Pycnogonum tenue*, - STOCK, Ann. Inst. Océanogr. Monaco, **44**: 401.
 1971 *Pycnogonum tenue*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 338.
 1983 *Pycnogonum tenue*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 64.
 1987 *Pycnogonum tenue*, - NAKAMURA, Biol. Lab., Imperial Household: 37.
 1991 *Pycnogonum tenue*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **512**: 63.

Distribution: Japan; Korea.

Depth range: 27-300m.

Habitat: sand.

Pycnogonum tessellatum STOCK 1968

- 1968 *Pycnogonum tessellatum* STOCK, Vidensk. Medd. dansk naturhist. Foren., **131**: 63-64, fig. 22f-i.

Distribution: Pakistan (Karachi harbour).

Pycnogonum torresi CLARK 1963

- 1963 *Pycnogonum torresi* CLARK, Rec. Austr. Mus., **26** (1): 76-77, fig. 37A-D.

Distribution: Torres Strait (Murray Island).
Depth range: 10-15m.

Pycnogonum tuberculatum CLARK 1963

1963 *Pycnogonum tuberculatum* CLARK, Rec. Austr. Mus., **26** (1): 77-79, fig. 38A-D.

Distribution: Australia.
Depth range: 119-148m.

Pycnogonum uedai NAKAMURA & CHILD 1983

1983 *Pycnogonum uedai* NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 64-66, fig. 21.

1984 *Pycnogonum koreanum* KIM & STOCK, J. nat. Hist., **18**: 685.

1986 *Pycnogonum koreanum*, - KIM & HONG, Kor. J. Syst. Zool., **2** (2): 50.

1987 *Pycnogonum koreanum*, - HONG & KIM, Kor. J. Syst. Zool., **3** (2): 161.

Distribution: Japan.
Depth range: 7-49m.

Pycnogonum ungellatum LOMAN 1911

1911 *Pycnogonum ungellatum* LOMAN, Abh. Kaiserl. Bayr. Akad. Wiss. (Math.-naturwiss. Kl.), Suppl. **2** (4): 7-8, pl. 2, figs. 25-27.

1949 *Pycnogonum ungellatum*, - HEDGPETH, Proc. U.S. natn. Mus., **98** (3231): 304, figs. 48c, 50e-g.

1961 *Pycnogonum ungellatum*, - LOSINA-LOSINSKY, Issledov. Dalnyevost. Morei SSSR, Leningrad, **7**: 111, fig. 27.

1971 *Pycnogonum ungellatum*, - UTINOMI, Publ. Seto Mar. Biol. Lab., **18** (5): 339.

1983 *Pycnogonum ungellatum*, - NAKAMURA & CHILD, Smiths. Contr. Zool., **386**: 66.

Distribution: NW-Pacific; Japan.
Depth range: 7-419m.

RHYNCHOTHORACIDAE

Rhynchothorax COSTA 1861

1966 *Rhynchothorax*, - STOCK, Vie Milieu, **17** (1B): 415 [key to species].

1976 *Rhynchothorax*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 294-295 [key to species].

1979 *Rhynchothorax*, - CHILD, Smiths. Contr. Zool., **293**: 67 [diagnostic features, remarks, key to Middle American species].

Rhynchothorax alcicornis KRAPP 1973

1973 *Rhynchothorax alcicornis* KRAPP, Bull. Zoöl. Mus. Univ. Amsterdam, **3** (17): 120, fig. 1.

- 1976 *Rhynchothorax alcicornis*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 294.
 1979 *Rhynchothorax alcicornis*, - CHILD, Smiths. Contr. Zool., **293**: 67.
 1987 *Rhynchothorax alcicornis*, - ARNAUD, Mesogee, **47**: 47, 54.
 1987 *Rhynchothorax alcicornis*, - ARNAUD & BAMBER, Adv. Mar. Biol., **24**: 32.
 1990 *Rhynchothorax alcicornis*, - ARNAUD & KRAPP, Beaufortia, **41** (1): 1-4, figs. 1-2.

Distribution: Mediterranean.

Depth range: 1-12m.

Habitat: calcareous algae; *Posidonia oceanica* seagrass; sponges and hydroids with sand.

***Rhynchothorax architectus* CHILD 1979**

- 1979 *Rhynchothorax architectus* CHILD, Smiths. Contr. Zool., **293**: 68-72, figs. 23, 24a-g, 25a-e.
 1982 *Rhynchothorax architectus*, - CHILD, Smiths. Contr. mar. Sci., **12**: 374.
 1986 *Rhynchothorax architectus*, - STOCK, Bull. mar. Sci., **38** (3): 405 tab. 1.
 1992 *Rhynchothorax architectus*, - CHILD, Smiths. Contr. Zool., **526**: 33-34.

Distribution: Belize and both sides of the Isthmus of Panamá; Caribbean Sea of Colombia; Galapagos Archipelago.

Depth range: intertidal-3m.

Habitat: between broken rocks and rubble.

***Rhynchothorax arenicolus* STOCK 1989**

- 1989 *Rhynchothorax arenicolus* STOCK, Bijdr. Dierk., **59** (2): 92-95, figs. 19-30.

Distribution: Curaçao.

Habitat: coarse sand of rockpool.

***Rhynchothorax articulatus* STOCK 1968**

- 1968 *Rhynchothorax articulatus* STOCK, Vidensk. Meddr. dansk naturhist. Foren, **131**: 18-20, fig. 5a-f.
 1970 *Rhynchothorax articulatus*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., **21**: 3.
 1972 *Rhynchothorax articulatus*, - ARNAUD, Tethys, **3** (2): 408.
 1973 *Rhynchothorax articulatus*, - KRAPP, Bull. Zoöl. Mus. Univ. Amsterdam, **3** (17): 121, 123.
 1976 *Rhynchothorax articulatus*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 287-292, figs. 2-9, 10-11.

Distribution: New Zealand.

Depth range: 146-500m.

Habitat: clay and sand.

***Rhynchothorax australis* HODGSON 1907**

- 1907 *Rhynchothorax australis* HODGSON, Rep. National Antarctic Exp. 1901-1904,

- Nat. Hist., 3: ..., Pl. 8.
- 1915 *Rhynchothorax australis*, - CALMAN, Rep. Brit. Antarct. "Terra Nova" Exped., Zool., 3: 67-68, fig. 21.
- 1938 *Rhynchothorax australis*, - HELFER, Sitz.-Ber. naturf. Freunde Berlin: 182.
- 1944 *Rhynchothorax australis*, - GORDON, Rep. Brit., Austr. and N.Z. Antarct. Res. Exped., (B), 5 (1): 67-68.
- 1950 *Rhynchothorax australis*, - HEDGPETH, Proc. U.S. natn. Mus., 100 (3260): 155.
- 1966 *Rhynchothorax australis*, - STOCK, Vie et Milieu, 17 (1B): 415.
- 1970 *Rhynchothorax australis*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., 21: 3.
- 1972 *Rhynchothorax australis*, - ARNAUD, Tethys, 3 (2): 408.
- 1973 *Rhynchothorax australis*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, 3 (17): 121.
- 1976 *Rhynchothorax australis*, - CLARK, J. Roy. Soc. N.Z., 6 (3): 295.

Distribution: Kerguelen Islands; Antarctica (off Wilkes Land to Ross Sea).

Depth range: 275m.

***Rhynchothorax barnardi* CHILD & HEDGPETH 1971**

- 1971 *Rhynchothorax barnardi* CHILD & HEDGPETH, J. nat. Hist., 5: 626-628, fig. 7.
- 1973 *Rhynchothorax barnardi*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, 3 (17): 121.
- 1976 *Rhynchothorax barnardi*, - CLARK, J. Roy. Soc. N.Z., 6 (3): 295.
- 1992 *Rhynchothorax barnardi*, - CHILD, Smiths. Contr. Zool., 526: 34-35, fig. 14.

Distribution: Galápagos Archipelago.

Depth range: intertidal-9m.

Habitat: rocks.

***Rhynchothorax crenatus* CHILD 1982**

- 1982 *Rhynchothorax crenatus* CHILD, Smiths. Contr. mar. Sci., 12: 374-376, fig. 167.
- 1986 *Rhynchothorax crenatus*, - STOCK, Bull. mar. Sci., 38 (3): 405 tab. 1.

Distribution: Belize (Carrie Bow Cay).

Depth range: 27m.

Habitat: coral sand and rubble in sand.

***Rhynchothorax malaccensis* STOCK 1968**

- 1968 *Rhynchothorax malaccensis* STOCK, Vidensk. Meddr. dansk. naturh. Foren., 131: 20-22, fig. 6.
- 1970 *Rhynchothorax malaccensis*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., 21: 3.
- 1972 *Rhynchothorax malaccensis*, - ARNAUD, Tethys, 3 (2): 408.
- 1973 *Rhynchothorax malaccensis*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, 3 (17): 121.
- 1976 *Rhynchothorax malaccensis*, - CLARK, J. Roy. Soc. N.Z., 6 (3): 295.

Distribution: Malacca Straits.
Depth range: 77m.

Rhynchothorax mediterraneus COSTA 1861

- 1861 *Rhynchothorax mediterraneus* COSTA, Microdoride mediterranea, o descr. de' poco ben conosc. od affato ignoti viventi, etc.: 8-9, pl. 1.
- 1881 *Rhynchothorax mediterraneus*, - DOHRN, Monogr. Fauna Flora Golf Neapel, 3: 211-215, pl. 17, figs. 1-9.
- 1909 *Rhynchothorax mediterraneus*, - d'ARCY THOMPSON, in: HARMER & SHIPLEY (Eds.) The Cambridge Nat. Hist.: 535, fig. 284.
- 1951 *Rhynchothorax mediterraneus*, - HEDGPETH, Wasman J. Biol., 9 (1): 114.
- 1963 *Rhynchothorax mediterraneus*, - ZILBERBERG, Bolm. Inst. Oceanogr. Univ. S. Paulo, 13 (2): 21.
- 1966 *Rhynchothorax mediterraneus*, - FAGE & STOCK, Ann. Inst. Océanogr. Monaco, 44: 325.
- 1966 *Rhynchothorax mediterraneus*, - STOCK, Vie et Milieu, 17 (1B): 415.
- 1970 *Rhynchothorax mediterraneus*, - ZAGO, Contr. Avuls. Inst. Océanogr. Univ. S. Paulo, 21: 1.
- 1972 *Rhynchothorax mediterraneus*, - ARNAUD, Téthys, 3 (2): 121 [key].
- 1973 *Rhynchothorax mediterraneus*, - KRAPP, Bull. Zoöl. Mus. Univ. Amsterdam, 3 (17): 121.
- 1974 *Rhynchothorax mediterraneus*, - STOCK, Bull. Zoöl. Mus. Univ. Amsterdam, 4 (3): 15-16, fig. 4.
- 1976 *Rhynchothorax mediterraneus*, - ARNAUD, Acta Ecol. Iranica, I: 70.
- 1976 *Rhynchothorax mediterraneus*, - CLARK, J. Roy. Soc. N.Z., 6 (3): 293, 295.
- 1987 *Rhynchothorax mediterraneus*, - ARNAUD, Mesogee, 47: 47-48.
- 1987 *Rhynchothorax mediterraneus*, - STOCK, Cah. Biol. Mar., 28: 507.
- 1988 *Rhynchothorax mediterraneus*, - CHILD, Bull. Biol. Soc. Wash., 8: 56.
- 1990 *Rhynchothorax mediterraneus*, - ARNAUD & KRAPP, Beaufortia, 41 (1): 4.
- 1992 *Rhynchothorax mediterraneus*, - STOCK, Tijdschr. Ent., 135: 135-136, 139.

Distribution: Mediterranean; Straits of Gibraltar; southern Brazil; Madagascar.

Depth range: 1-?1100m.

Habitat: "une espèce des fonds détriques et des fonds concrétionnement coralligène" (ARNAUD, 1987).

Rhynchothorax monnioti ARNAUD 1974

- 1974 *Rhynchothorax monnioti* ARNAUD, Bull. Zoöl. Mus. Univ. Amsterdam, 3 (21): 171-172, figs. 1-5.
- 1976 *Rhynchothorax monnioti*, - CLARK, J. Roy. Soc. N.Z., 6 (3): 295.
- 1987 *Rhynchothorax monnioti*, - ARNAUD, Mesogee, 47: 48.
- 1990 *Rhynchothorax monnioti*, - ARNAUD & KRAPP, Beaufortia, 41 (1): 4-5.

Distribution: Mediterranean; Canary Islands.

Depth range: 15-62m.

Habitat: coralline algae.

Rhynchothorax orientalis CHILD 1988

- 1988 *Rhyncothorax orientalis* CHILD, Smiths. Contr. Zool., **468**: 28-29, fig. 12.
 1991 *Rhyncothorax orientalis*, - STOCK, Mém. Mus. natn. Hist. nat. (A), **151**: 227.

Distribution: Philippines.

Depth range: 1-97m.

***Rhyncothorax percivali* CLARK 1976**

- 1976 *Rhyncothorax percivali* CLARK, J. Roy. Soc. N.Z., **6** (3): 292-294, figs. 1, 12-21.
 1988 *Rhyncothorax percivali*, - CHILD, Smiths. Contr. Zool., **468**: 28.

Distribution: New Zealand.

Depth range: littoral.

Habitat: rockpools.

***Rhyncothorax philopsammum* HEDGPETH 1951**

- 1951 *Rhyncothorax philopsammum* HEDGPETH, Wasmann J. Biol., **9** (1): 111-115, pl. 3.
 1966 *Rhyncothorax philopsammum*, - STOCK, Viet et Milieu, **17** (1B): 415 [key].
 1970 *Rhyncothorax philopsammum*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., 21: 3 [key].
 1972 *Rhyncothorax anophthalmus* ARNAUD, Tethys, **3** (2): 405-409, figs. 1-7.
 1973 *Rhyncothorax anophthalmus*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, **3** (17): 121 [key].
 1972 *Rhyncothorax philopsammum*, - KRAPP, Tethys, **3** (2): 406.
 1973 *Rhyncothorax philopsammum*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, **3** (17): 121.
 1974 *Rhyncothorax anophthalmus*, - ARNAUD, Bull. Zoöl. Mus. Univ. Amsterdam, **3** (21): 172-173, figs. 6-7.
 1976 *Rhyncothorax anophthalmus*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 295 [key].
 1976 *Rhyncothorax philopsammum*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 295.
 1979 *Rhyncothorax philopsammum*, - CHILD, Smiths. Contr. Zool., **293**: 67, 72, fig. 24h.
 1987 *Rhyncothorax anophthalmus*, - ARNAUD, Mesogee, **47**: 47.
 1987 *Rhyncothorax anophthalmus*, - ARNAUD & BAMBER. Adv. Mar. Biol., **24**: 32.
 1990 *Rhyncothorax philopsammum*, - CHILD, Proc. biol. Soc. Wash., **103** (2): 311.
 1990 *Rhyncothorax philopsammum*, - ARNAUD & KRAPP, Beaufortia, **41** (1): 5.
 1990 *Rhyncothorax philopsammum*, - MÜLLER, Senckenbergiana biol., **70** (1989) (1/3): 197.

Distribution: California; Pacific Coast of Mexico; Bora Bora, Society Islands; Caribbean Sea of Colombia.

Depth range: 0-77m.

Habitat: sand and gravel; dead coral substratum.

***Rhyncothorax tiahurensis* MÜLLER 1889**

- 1989 *Rhyncothorax tiahurensis* MÜLLER, Bonn. zool. Beitr., **40** (2): 127-128, figs. 58-62.

Distribution: Society Islands (Moorea).

Depth range: 0.5-1m.

Habitat: dead corals on barrier reef.

***Rhyncothorax unicornis* FAGE & STOCK 1966**

- 1966 *Rhyncothorax unicornis* FAGE & STOCK, Ann. Inst. Océanogr., **44**: 324-326, fig. 4.
1966 *Rhyncothorax unicornis*, - STOCK, Viet et Milieu, **17** (1B): 415.
1970 *Rhyncothorax unicornis*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., **21**: 3.
1972 *Rhyncothorax unicornis*, - ARNAUD, Tethys, **3** (2): 408.
1973 *Rhyncothorax unicornis*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, **3** (17): 121.
1976 *Rhyncothorax unicornis*, - CLARK, J. Roy. Soc. N.Z., **6** (3): 295.

Distribution: Cape Verde Islands.

Depth range: 40m.

Habitat: on rocks.

***Rhyncothorax vallatus* CHILD 1990**

- 1990 *Rhyncothorax vallatus* CHILD, Proc. biol. Soc. Wash., **103** (2): 333-334, fig. 7.

Distribution: Australia (Lizard Island).

Depth range: 2m.

Habitat: coral reef rubble.

***Rhyncothorax voxorinum* STOCK 1966**

- 1966 *Rhyncothorax voxorinum* STOCK, Vie Milieu, **17** (1B): 412-415, fig. 3a-e.
1970 *Rhyncothorax voxorinum*, - ZAGO, Contr. Avuls. Inst. oceanogr. Univ. S. Paulo, sér. Ocean. Biol., **21**: 3.
1972 *Rhyncothorax voxorinus*, - ARNAUD, Tethys, **3** (2): 406.
1973 *Rhyncothorax voxorinus*, - KRAPP, Bull. Zool. Mus. Univ. Amsterdam, **3** (17): 121.
1987 *Rhyncothorax voxorinum*, - ARNAUD, Mesogee, **47**: 48.
1990 *Rhyncothorax voxorinum*, - STOCK, Beaufortia, **41** (1): 5.
1990 *Rhyncothorax voxorinum*, - ARNAUD & KRAPP, Beaufortia, **41** (1): 5.

Distribution: Mediterranean.

Depth range: 10-18m.

Habitat: in crevices or endopsammic.

BIBLIOGRAPHY OF THE RECENT PYCNOGONIDA

This bibliography lists in alphabetic order the publications on pycnogonids.

For your convenience to find articles reporting on different subjects I included several keywords for the most important publications, which can be used in the search-function of Word Perfect:

alimentary system
anatomy - for papers reporting on organs of Pycnogonida, functional morphology, or histology.
bivalve host
bryozoan host
circulatory system
cytology
development
echinoderm host
ecology - different kind of ecological aspects.
embryology
feeding - the food of pycnogonids.
gastropod host
genital system
gland
gynandromorph
hormones
hydroid host
mollusca host
musculature
nervous system
optical sense
physiology
pollution - runs to all different kind of pollution.
polychaete host
reproduction
respiration

When you are searching for a certain genus or species in PYCNO.LIT typewrite the name without author and year. Be sure to run with the search function through the complete bibliography and check the synonyms in the systematic index PYCNO.CAT, because older names may exist in the references.

BIBLIOGRAPHY

Abildgaard, P.C. (1789): In O.F. Müller, *Zoologica Danica seu animalium Daniae et Norvegiae rariorum ac minus notorum et Historia*, 3: 67-68; CXIX; Havniae.

Acloque, A. (1899): Classe de Pycnogonides, ordre des Pantopodes, in: Faune de France, pp. 219-220; Paris, J.B. Baillièvre & Fils.

Adams, J. (1800): Descriptions of some marine animals found off the coast of Wales. - Trans. Linn. Soc. Lond, 1800 (4), 5: 7-12; II.

Adlerz, G. (1888): Bidrag till Pantopodernas morfologi och utvecklings-historia. - Kungl. Svensk. Vetensk.-Akad. Handl., Stockholm, Bihang, (4) 13 (11): 1-25; I-II.

Allman, G.J. (1859): On parasitic larval forms in hydroids. - Rep. Brit. Assoc. Adv. Sci., 1859. [hydroid host]

Allman, G.J. (1862): On a remarkable form of parasitism among the Pycnogonidae. - Ann. Mag. Nat. Hist., (3) 9: 36-43.

Allman, G.J. (1872): Pycnogonidae, in: A monograph of the gymnoblastic or tubularian hydroids, part II: 200-201, 314; 75; London.

Anadón, R. (1975): Aportación al conocimiento de la fauna bentónica de la ría de Vigo (NW de España). 1. Pycnogónidos y crustáceos de Panjón. - Inv. Pesq., 39 (1): 199-218.

André, M. & E. Lamy (1938): Pycnogonides parasites de Mollusques. - J. Conch., 82: 326-331. [mollusca host]

Andrews, H.J. (1977): Observations on the Fauna of the Metropolitan River Thames during the Drought in 1976. - London Nat., 56: 44-56.

Angel, H. (1977): Sea Feature. - Sea Frontiers, 23 (5): 321.

Anonymous (1948): Cover photo, in: Science, 107 (2778).

Anonymous (1978): Water Quality Investigation of Estuaries in Georgia, 1977. Brunswick Junior College (Georgia), Ann. Rep., 1978: 1-166.

Appellöf, A. (1910): Pycnogoniden. The Norwegian Arctic Expedition of the Fram, 1898-1902, 2 (26).

Appellöf, A. (1912): Invertebrate bottom fauna of the Norwegian Sea and North Atlantic. Ch. 8, in: J. Murray & J. Hjort. The depths of the sea: 821 pp., London; Macmillan.

Appellöf, A. (1912): Die Pycnogoniden des Eisfjorden. Zoologische Ergebnisse der schwedischen Expedition nach Spitzbergen 1908, II (5). Kungl. Svensk. Vetensk.-Akad. Handl., Stockholm, 54 (5): 1-29.

- Ardizzone, G.D., C. Chimenz & E. Taramelli (1976): Valore dell'indice di diversità nello studio delle comunità fouling del Porto di Civitavecchia. - Boll. Pesca, Piscicolt. Idrobiol., Roma, 31 (1-2): 313-318.
- Arita, K. (1936): Ein überzähliges Bein bei einer Pantopoden-Art (*Nymphonella tapetis* Ohshima). - Annot. Zool. Japon., 15 (4): 469-477, 1 pl.
- Arita, K. (1937): Beiträge zur Biologie der Pantopoden. - J. Dep. Agric. Kyushu Imp. Univ., 5 (6): 271-288. [biology]
- Arnaud, F. (1969): Les Pycnogonides: un aspect de la biologie marine dans les Terres Australes et Antarctiques Françaises. Les Terres Australes Antarctiques Françaises, 47: 35-41; La Documentation français, Paris.
- Arnaud, F. (1970): Pycnogonides subantarctiques des îles Crozet. - Bull. Mus. natn. Hist. nat., Paris, (2) 41 (6): 1423-1428.
- Arnaud, F. (1971): *Acheliana tropicalis* n. gen., n. sp., Pycnogonide des récifs coralliens du sud-ouest de Madagascar. - Beaufortia, 18 (241): 199-204.
- Arnaud, F. (1971): Pycnogonides des récifs coralliens de Madagascar. 2. Redéscription de *Pycnogonum madagascariensis* Bouvier, 1911, Suppl. 1: 161-163.
- Arnaud, F. (1971): Pycnogonides littoraux des îles Saint-Paul et Amsterdam (Océan Indien). - Téthys, 3 (1): 159-166.
- Arnaud, F. (1972): Un nouveau Pycnogonide de Méditerranée nord-occidentale: *Rhynchothorax anophthalmus* n. sp. et redécouvert de *Rhynchothorax mediterraneus* Costa, 1861. - Téthys, 3 (2): 405-409.
- Arnaud, F. (1972): 9. Pycnogonides. Invertébrés marins. XIIeme et XVeme expéditions antarctiques françaises en Terre Adélie. - Téthys, Suppl. 4: 135-156.
- Arnaud, F. (1972): Pycnogonides des îles Kerguelen (Sud Océan Indien); matériel nouveau et révision des spécimens du Muséum national d'Histoire naturelle de Paris. - Bull. Mus. natn. Hist. nat., Paris, (3) 51 (65): 801-815.
- Arnaud, F. (1973): Pycnogonides des récifs coralliens de Madagascar. 4. Colossendeidae, Phoxichilidiidae et Endeidae. - Tethys, 4 (4): 953-960.
- Arnaud, F. (1973): Les Pycnogonides du golfe de Gascogne (Atlantique Nord-Est). - Téthys, 5 (1): 147-154.

- Arnaud, F. (1974): Pycnogonides recoltes aux Açores par les campagnes 1969 et Biacores 1971. - Bull. Zool. Mus. Univ. Amsterdam, 3 (21): 169-186.
- Arnaud, F. (1974): Nouveaux Pycnogonides des îles Saint Paul et Amsterdam (Océan Indien). - Téthys, 5 (4): 779-790.
- Arnaud, F. (1976): Sur quelques Pycnogonides de Turquie et de la mer Egée (Méditerranée orientale). - Acta Ecol. Iranica, 1: 68-72.
- Arnaud, F., P.M. Arnaud, A. Intes & P. le Loeuff (1976): Transport dinvertébrés benthiques entre l'Afrique du Sud et Sainte Hélène par les Laminaires (Phaeophyceae). - Bull. Mus. natn. Hist. nat., Paris, (3), 384: 49-55.
- Arnaud, F. (1978): A new species of *Ascorhynchus* (Pycnogonida) found parasitic on an opisthobranchiate mollusc. - Zool. J. Linn. Soc., 63 (1-2): 99-104. [gastropod host]
- Arnaud, F. (1988): Les Pycnogonides (Chelicerata) de Méditerranée: Distribution écologique, bathymétrique et biogéographie. - Mésogée - Bull. Mus. Hist. Nat. Marseille, 1987, 47: 37-58.
- Arnaud, F. & R.N. Bamber (1987): The Biology of Pycnogonida. - Adv. Mar. Biol., 24: 1-96.
- Arnaud, F. & C.A. Child (1988): The South African Museum's Meiring Naude Cruises, Part 17: Pycnogonida. - Ann. S. Afr. Mus., 98 (6): 121-187.
- Arnaud, F. & F. Krapp (1990): The genus *Rhynchothorax* (Pycnogonida) in the Mediterranean Sea. - Beaufortia, 41 (1): 1-7; Amsterdam.
- Arnaud, P.M. (1970): Frequency and ecological significance of necrophagy among the benthic species of Antarctic coastal waters. In: M.W. Holdgate (ed.). Antarctic Ecology, 2: 259-267. Scientific Committee for Antarctic Research. London & New York; Academic Press.
- Arnaud, P.M. (1973): Contributions à la bionomie marine benthique des régions antarctique et subantarctique. Thèse, Doct. Sci. nat., Univ. Aix-Marseille, 7819.
- Arnaud, P.M. (1974): Contribution à la bionomie marine benthique des régions antarctiques et subantarctiques. - Téthys, 6 (3): 467-653.

Arnaud, P.M., K. Jazdzewski, P. Dresler & J. Sicinski (1986) : Preliminary survey of benthic invertebrates collected by Polish Antarctic Expeditions in Admiralty Bay (King George Island, South Shetland Islands, Antarctica). - Pol. Polar Res., 7 (1-2) : 7-24.

Arndt, W. (1913) : Zoologische Ergebnisse der ersten Lehrexpedition der Dr. P. Schlotländerschen Jubiläums-Stiftung. - Jb. Schles. Ges. Vaterl. Kult., 90 (1912) (1) (2)b (Zool. Bot.) : 110-136.

Arndt, W. & K. Viets (1938) : Die biologischen (parasitologischen) Beziehungen zwischen Arachnoideen und Spongien. - Z. Parasitenk., 10 (1) : 67-93.

Bacescu, M. (1953) : Contributii la cunoasterea picnogonidelor marii negre. - Bull. Stiint., Acad. Rep. Pop. Române, Sec. Stiint. Biol., Agr., Geol., 5 (2) : 263-270.

Bacescu, M. (1958) : Pycnogonide noi pentru fauna Maarii Negre: *Anoplodactylus petiolatus* (Kr.), *A. stocki* n.sp. si *Callipallene brevirostris* (John.). - Stud. Cercetari Biol., ser. "Biol. Anim.", 10 (4) : 321-334.

Bacescu, M. (1959) : Pycnogonides nouveaux pour la faune de la Mer Noire: *Anoplodactylus petiolatus* (Kr.), *A. stocki* n.sp. et *Callipallene brevirostris* (John.). - Rev. Biol., Acad. Rep. Pop. Roum., 4 (1) : 117-128.

Bacescu, M. & C. Margineanu (1959) : Eléments méditerranéens nouveaux dans la faune de la Mer Noire, rencontrés dans les eaux de Roumérie (Nord-Ouest-Bosphore). - Arch. Oceanogr. Limnol., 11, Suppl. 63-74.

Bacescu, M. (1968) : Pycnogonida. In: Guide to the Identification of the Black Sea and Azov Sea Fauna. Naukova Dumko, Kiev, 1968. [in Russian]

Bacescu, M. (1972) : Tom tretj: Swobodnosiwutcie bespozvonocnye: Clenistonosie (krome rakoobraznych), mollujski, islokoscie, teminkoceljustnye, chordo ye. In: F.D. Morduchai-Boltowskoi, Opredelitel fauny Cernogo i Azowskogo Morei (v trech tomach). Naukova Dumko (Kiev), 1972: 340. [in Russian]

Bacescu, M. (1972) : Rapport sur les travaux parus, entre 1968 et 1970, sur le benthos de Méditerranée, ses mers annexes y comprises et la mer Rouge, I: Rapport sur les travaux concernant le zoobenthos. - Rapp. P.-v. Comm. Int. Expl. Sci. Mer Médit., 21 (9) : 549-576.

Bacescu, M. (1972) : Klass Pantopody, ili morskiye pauki-Pantopoda Gerstaeker, 1777. In: Opredel. Fauny Chornovo Azovsk. Morei, 3: 32-37; Kiev.

- Bain, B. (1988): A preliminary look at the systematics of the Pycnogonida. - Am. Zool., 28 (4): 9A. [Abstract]
- Bain, B. (1991): Some observations on biology and feeding behavior in two southern California pycnogonids. - Bidr. Dierk., 61 (1): 63-64.
- Balfour, F.M. (1880): Pycnogonida. In: Treatise of comparative embryology, 1: 448; Jena, von Vetter.
- Balfour, F.M. (1880): Notes on the development of the Araneida. - Quart. J. Microsc. Sci., (N.S.), 20: 167-189.
- Bamber, R.N. (1978): The morphological taxonomy of *Nymphon rubrum* (Pycnogonida). - Porcupine News., 1 (5): 79.
- Bamber, R.N. (1979): A new species of *Endeis* (Pycnogonida) from West Africa. - Zool. J. Linn. Soc., 65: 251-254.
- Bamber, R.N. (1980): British Sea Spiders by P.E. King [review]. - Irish. Nat. J., 20 (3): 127.
- Bamber, R.N. (1982): Variation in *Nymphon brevirostre* (Hodge), and the status of *N. rubrum* (Hodge) (Arthropoda: Pycnogonida). - Biol. J. Linn. Soc., 17: 275-288.
- Bamber, R.N. & M.H. Davis (1982): Feeding of *Achelia echinata* Hodge (Pycnogonida) on marine algae. - J. Exp. mar. Biol. Ecol., 60: 181-187.
- Bamber, R.N. (1983): The marine fauna of the Cullercoats District, no. 12: Pycnogonida. - Rep. Dove Mar. Lab., 3rd ser., 25: 1-24.
- Bamber, R.N. (1983): Some deep water Pycnogonids from the North-East Atlantic. - Zool. J. Linn. Soc., 77: 65-74.
- Bamber, R.N. & P.A. Henderson (1983): Epifaunal arthropods from the tide pools at Rhosneigr. - Porcupine News., 2 (8): 196.
- Bamber, R.N. (1985): A second *Pallenopsis mollissima* (hoeck), with other deep water pycnogonids from the Glasgow Museum. - Zool. J. Linn. Soc., 83: 301-306.
- Bamber, R.N. (1985): The intinerant sea-spider *Ammothaea hilgendorfi* (Böhm) in British Waters. - Proc. Hampsh. Field Club Archaeol. Soc., 41: 269-270.
- Bamber, R.N. (1985): Why do Pycnogonids prefer inaccessible anemones?. - Porcupine News., 3: 67-71.

Bamber, R.N. (1986): Epifaunal collections from the Channel Islands, September 1986. - Porcupine News., 3 (9): 235-239.

Barnard, K.H. (1946): Diagnoses of a new species and a new genus of Pycnogonida in the South African Museum. - Ann. Mag. Nat. Hist., (11) 13: 60-63.

Barnard, K.H. (1954): South African Pycnogonida. - Ann. S. Afr. Mus., 41 (3): 81-158.

Barnard, K.H. (1955): Additions to the fauna-list of South African Crustacea and Pycnogonida. - Ann. S. Afr. Mus., 43 (1): 1-107.

Barnes, R.D. (1968): Class Pycnogonida or Pantopoda. In: Invertebrate Zoology: 423-426; New York: Saunders.

Barrett, J. & C.M. Yonge (1964): Pycnogonida: Sea spiders. In: Collins pocket guide to the sea shore: 122-123; London: Collins.

Baster, J. (1765): In: Natuurkundige Uitspanningen, behelzende waaremingen over sommige Zee-planten en Zee-Insecten, benevens derzelver zaadhuisjes en eijernesten, 2: 155; Haarlem: J. Bosch.

Baster, J. (1765): In: Opuscula subseciva, observationes miscellaneas de animalculis et plantis quibusdam marinis, eorumque ovariis et seminibus continentia, 2: 139-142; Haarlem.

Bastida, R.M., M. Trivi de Mandri, L. de Bastida & M. Stupak (1980): Ecological aspects of marine fouling at the Port of Mar del Plata (Argentina). V. Congr. Int. Corros. Mar. e Incrust., pp. 299-320.

Bayer, F.M., G.L. Voss & C.R. Robbins (1970): Report on the marine fauna and benthic shelf-slope communities of the Isthmian Region. - In: Bioenvironmental and Radiological Safety Feasability Studies Atlantic-Pacific Interceanic Canal, pp. 1-99, appendix, pp. 1-311. [processed Rep., Univ. Miami, Florida]

Beaumont, W.I. (1900): The fauna and flora of Valencia Harbour on the west coast of Ireland, II. The benthos (dredging and shore collecting), VIII... - Proc. Roy. Ir. Acad., Dublin, (3) 5: 754-798.

Behrens, W. (1984): Larvenentwicklung und Metamorphose von *Pycnogonum litorale* (Chelicerata, Pantopoda). - Zoomorphol., 104: 266-279. [development]

Behrens, W. & D. Bückmann (1981): Der Nachweis von Häutungshormonen bei einem Pantopoden, *Pycnogonum litorale* Ström. - Verh. Dtsch. Zool. Ges., 74: 195. [physiology, hormones]

Behrens, W. & D. Bückmann (1983): Ecdysteroids in the Pycnogonid *Pycnogonum littorale* (Strom) (Arthropoda, Pantopoda). - Gen. Comp. Endocrinol., 51: 8-14. [physiology, hormones]

Beklemishev, V.N. (1964): Pantopodi. In: Ossnovi Sravnityelnoi Anatomii Bespozvonochnikh, 1: 207, 210, 298, 299-301, 305, 312; 149, 2: 100, 105, 158, 205, 357, 358, 360; Moscow, Akad. Nauk SSSR.

Bell, T. (1853): Account of the Crustacea. In: The last of the Arctic Voyages, being a narrative of the expedition of HMS Assitance, under the command of Captain Sir Edward Belcher, 2: 408-409; London.

Bellan-Santini, D. (1967): Contribution à l'étude des peuplements infralittoraux sur substrat rocheux... Mémoire présentée le 20 Décembre 1967 pour l'obtention du Grade de Docteur dès Sciences naturelles: 1-294.

Bellisio, N.B. (1966): Fauna marina Arctica. Secretaria de Marina, Servico de Hydrografia naval, Rep. Argentina, Buenos Aires, H 907: 1-91.

Belloc, G. (1962): Catalogue des types de Pycnogonides du Musée océanographique de Monaco. - Bull. Inst. Océanogr., Monaco, 1215: 3 pp.

Belyaev, G.M. (1966): Pycnogonids (Sea spiders) - Pantopoda. In: Hadal (Ultraabyssal) Fauna of the World Ocean: 102-103; Trudy Inst. Okeanol. Akad. Nauk SSSR, 591: 1-248. [in Russian]

Béanard, F. (1960): La faunule associée au *Lithophyllum incrustans* Phil. dans les cuvettes de la région de Roscoff. - Cah. Biol. Mar., 1 (1): 89-102.

Benson, P.H. & D.C. Chivers (1960): A Pycnogonid infestation of *Mytilus californicus*. - The Veliger, 3 (1): 16-18. [Bivalvia host]

Berry, M.B. (1980): The embryological development of *Achelia sawayai* (Ammotheidae, Pycnogonida), with notes on some phases of the behavior and anatomy of the adult, and on the phylogenetic position of the Pycnogonida. - Diss. Abstr. Int., (B) 41 (3): 849. [embryology]

Berry, M.B. (1982): Gastrulation by delamination in *Achelia sawayai*. - Am. Zool., 22 (4): 876. [embryology]

- Bertrand, H. (1941): Les Pycnogonides de la région dinardaise. - Bull. Lab. Mar. Dinard, 23: 48-52.
- Bertrand, H. (1943): Récoltes de Crustacés Malacostracés et de Pycnogonides dans la rivière de Trieux. - Bull. Lab. marit. Dinard, 25: 6-18.
- Beyerley, I. (1855): Fauna of Liverpool. - Proc. Literary & Phil. Soc. Liverpool, 8 (1853/4), Appendix: 125.
- Bitar, G. (1987): Sur une collection de Crustaces et de Pycnogonides du Maroc. - Bull. Inst. Sci., Univ. Mohammed V, Rabat, 1987: 155-166.
- Björck, W. (1913): Biologisch-faunistische Untersuchungen aus dem Öresund. 1. Pantopoda, Mysidacea und Decapoda. - Lunds Univ. Arsskr., N.F. (2) 9 (17): 39pp., 1 pl.
- Björck, W. (1915): Biologisk-faunistika Undersökningar av Öressund. 2. Crustacea Malacostraca och Pantopoda. - Lunds Univ. Arsskr., N.F. (2) 11 (7): 1-98.
- Blake, C.H. (1933): Class Pycnogonida: Sea Spiders. In: W. Procter, Biological Survey of the Mount Desert Region, Part V.: Marine Fauna. The Wistar Inst. of Anat. Biol., Philadelphia, 1933: 266-267.
- Boaden, P.J.S. & R.J. O'Connor & R. Seed (1975): Pycnogonida. In: The composition of the *Fucus serratus* community in Strangford Lough, Co. Down. - J. Exp. Mar. Biol. Ecol., 17 (2): 111-136.
- Böhm, R. (1879): Ueber die Pycnogoniden des Kgl. Zoolog. Museums zu Berlin, insbesondere über die von S.M.S. Gazelle mitgebrachten Arten. - Monatsber. Königl. Preuss. Akad. Wiss. Berlin, 1879: 170-195, 2 pls.
- Böhm, R. (1879): Ueber zwei neue von Herrn Dr. Hilgendorf in Japan gesammelte Pycnogoniden. - Sitz.-Ber. Ges. Naturforsch. Fr. Berlin, 1879 (4): 53-60.
- Böhm, R. (1879): Ueber Pycnogoniden. - Sitz.-Ber. Ges. Naturforsch. Fr. Berlin, 1879 (9): 140-142.
- Bogdanos, C. & J. Satsmadijs (1983): The macrozoobenthos of an Aegean embayment. - Thalassogr., 6: 77-105.
- Boisseau, J.P. (1952): Notes sur la faune d'Arcachon (Première Sér.). - Bull. Soc. zool. France, 77: 148-151.
- Börner, C. (1903): Die Beingliederung der Arthropoden. 2. - Sitz.-Ber. Ges. Naturforsch. Fr. Berlin, 1903: 292-335.

Borradaile, L.A., F.A. Potts, L.E.S. Eastham & J.T. Saunders (1951): Class Pantopoda (Pycnogonida). In: The Invertebrata: A manual for the use of students: 538-539; 370A; Cambridge Univ. Press.

Bosc, L.A.G. (1802): Histoire Naturelle des Crustacés, 2: 202-207; Paris.

Boudreaux, H.B. (1979): Significance of intersegmental Tendon Systems in arthropod phylogeny and a monophyletic classification of arthropoda. In: A.P. Gupta (ed.). Arthropod Phylogeny, pp. 551-586. Van Nostrand Reinhold Co.

Boudreaux, H.B. (1979): Phylogeny of the arthropod classes, in: Arthropod phylogeny with special reference to insects, pp. 82-88, 115-122. New York; John Wiley & Sons.

Bourdillon, A. (1952): Note préliminaire sur diverses réactions parasitaires d'un Hydroïde aux larves de Pycnogonides. - Rec. Trav. Stat. Mar. d'Endoume, 6: 19-26. [hydroid host]

Bourdillon, A. (1953): Note sur *Pallene brevirostris* (Pycnogonida). - Rec. Trav. Stat. Mar. d'Endoume, 8: 37-41.

Bourdillon, A. (1954): Les Pycnogonides de Marseille et ses environs. - Rec. Trav. Stat. Mar. d'Endoume, 12 (7): 145-158.

Bourdillon, A. (1954): Contribution à l'étude des Pycnogonides de Tunisie. - Notes Stat. Océanogr. Salammbô, 35: 1-8, 1 pl.

Bourdillon, A. (1955): Les Pycnogonides de la croisière 1951 du Président Théodore Tissier. - Rev. Trav. Inst. Pêches Marit., 19 (4): 581-609.

Bourne, G.C. (1890): Report of a trawling cruise in H.M.S. "Research" off the south-west coast of Ireland. - J. mar. Biol. Ass. Plymouth, 1: 306-323.

Bousfield, E.L. & A.H. Leim (1958): The fauna of Minas Basin and Minas Channel. - Bull. natn. Mus. Canada, 166: 1-30.

Bouvier, E.L. (1905): Observations préliminaires sur les Pycnogonides recueillis dans la région antarctique par la mission du "Français". - Bull. Mus. natn. Hist. nat., Paris, 11: 294-298.

Bouvier, E.L. (1906): Nouvelles observations sur les Pycnogonides recueillis dans les régions antarctiques au cours de la campagne dirigée par M. Jean Charcot. - C.R. Séanc. Hebd. Acad. Sci., Paris, 142: 15-22.

Bouvier, E.L. (1906): Pycnogonides du "Français". L'expédition Antarctique Française, 1903-1905: 1-69.

Bouvier, E.L. (1910): Les Pycnogonides à cinq paires de pattes recueillis par la Mission antarctique Jean Charcot à bord du "Pourquoi Pas?". - C.R. Séanc. Hebd. Acad. Sci., Paris, 151: 26-32.

Bouvier, E.L. (1910): Note sur les Arthropodes marins recueillis par M. Rallier du Batys aux îles Kerguelen. - Bull. Mus. natn. Hist. nat., Paris, 16: 178-179.

Bouvier, E.L. (1910): Mission en Mauritanie occidentale. III. Partie zoologique. Crustacés et Pycnogonides. - Act. Soc. Linnéenne, Bordeaux, 64: 221-226, 1pl.

Bouvier, E.L. (1911): Observations sur les Pycnogonomorphes et principalement sur le *Pentapycnon Geayi*, espèce tropicale à dix pattes. - C.R. Séanc. Hebd. Acad. Sci., Paris, 152: 491-494.

Bouvier, E.L. (1911): Les Pycnogonides du "Pourquoi Pas?". - C.R. Séanc. Hebd. Acad. Sci., Paris, 152: 1136-1141.

Bouvier, E.L. (1911): Pycnogonides. Algunos animales marinos de las Islas Sandwich. - An. Mus. Nac. Hist. Nat. Buenos Aires, (3) 15: 413-414.

Bouvier, E.L. (1911): Les Pycnogonides décapodes et la classification des Pycnogonides. - Proc. Int. Congr. Ent., Bruxelles, 1910: 345-355.

Bouvier, E.L. (1913): Pycnogonides du "Pourquoi Pas?". - Deuxième Expéd. Antarct. Française (1908-1910), 6: 1-169.

Bouvier, E.L. (1914): Quelques mots sur la variabilité du *Pycnogonum littorale* Ström. - J. mar. Biol. Ass., Plymouth, 10: 207-210.

Bouvier, E.L. (1914): Les Crustacés de profondeur et les Pycnogonides recueillis par le "Pourquoi Pas?..." 1913. - Bull. Mus. natn. Hist. nat., Paris, 20: 215-221.

Bouvier, E.L. (1916): Sur un Nymphonomorphe nouveau capturé par le "Travailleur", dans les mers européennes au cours de sa campagne de 1881. - Bull. Mus. natn. Hist. nat., Paris, (1) 22: 14-16.

Bouvier, E.L. (1916): Un nouveau Pycnogonide, *Ammothea (Achelia) armata*, trouvé par le "Talisman". - Bull. Mus. natn. Hist. nat., Paris, 22: 81-83.

Bouvier, E.L. (1917): Sur un Pycnogonide (Arach.) nouveau recueilli en Guinée portugaise par le comte de Polignac, au cours de sa campagne de 1913 sur le yacht "Sylvana". - Bull. Soc. ent. France, Paris, 10: 223-226.

Bouvier, E.L. (1917): Pycnogonides provenant des campagnes scientifiques de S.A.S. le Prince de Monaco (1885-1913). Rés. Camp. Sci. accomplies sur son Yacht par Albert I^{er} Prince Souverain de Monaco, 51: 1-56.

Bouvier, E.L. (1920): On a collection of Pycnogonida from the South Orkney Islands. - Ann. Mag. Nat. Hist., (9) 6: 244-247.

Bouvier, E.L. (1922): Classification et caractères distinctifs essentiels des Pycnogonides appartenants à la série des Pycnogonomorphes. - Ann. Sci. Nat., Paris (Zool.), (10) 5 (1,2): 113-117.

Bouvier, E.L. (1923): Quelques notes sur les Pycnogonides des côtes de France. - Ann. Sci. Nat., Paris (Zool.), (10) 6 (3,4): 117-124.

Bouvier, E.L. (1923): Pycnogonides. Faune de France, 7: 1-69.

Bouvier, E.L. (1937): Étude sur les Pycnogonides du "Travailleur" et du "Talisman" précédée d'observations systématiques sur les Articulées de ce groupe. - Ann. Sci. Nat., Paris, (Zool.), (10) 20 (1): 1-42.

Brady, G.S. (1903): Report on the dredging and other marine research off the N.E. coast of England in 1901. - Nat. Hist. Trans. Northumberland, Durham and Newcastle-upon-Tyne, 14: 87-101.

Bremec, C.S., D.E. Martinez & E. Elias (1986): Pycnogónidos de los alrededores de Bahía Blanca (Argentina). - Spheniscus, 3: 31-49.

British Museum (Natural History) (1927): Guide to the Crustacea exhibited in the Department of Zoology, British Museum (Natural History): 1-81; London.

Broch, H. (1961): Benthonic problems in Antarctic and Arctic waters. Sci. Ress. Norw. Antarct. Exped. 1927-1928 et sqq., 3 (38): 1-32; Oslo, Norske Vidensk.-Akad.

Bromley, J.E.C. (1979): A preliminary checklist of marine fauna of Minas Basin and Minas Channel. - Proc. Nova Scotian Inst. Sci., Halifax, 29 (4): 517-541.

Bronstein, Z.S. (1948): Check list of fauna and flora of the northern seas of the USSR. In: N.S. Gajevskoi (ed.), Opredel. Fauni Flori Severnich Morei SSSR: 344-351. [in Russian]

Bruce, J.R., J.S. Colman & N.S. Jones (1963): Pycnogonida pp. 180-182. In: marine fauna of the Isle of Man and its surrounding seas: i-xi, 1-307; Liverpool Univ. Press.

Brunel, P. (1970): Catalogue d'Invertébrés benthiques du Golfe Saint-Laurent recueillis de 1951 à 1966 par la Station de Biologie marine de Grand-Rivière. - Trav. Pêch. Quebec, 32: 1-55.

Brünnich, M.T. (1764): Entomologia sistens Insectorum Tabulas Systematicas, cum Introductione et Iconibus, etc.: 87 pp., 1 pl.

Buchholz, R. (1874): Crustaceen der zweiten Deutschen Nordpolarfahrt. Anhang Pycnogonida. Die 2. deutsche Nordpolarfahrt in den Jahren 1869 und 1870, 2 (7): 396-397.

Bückmann, D. (1979): Induction and control of metamorphosis in arthropods, a review under comparative aspects. - Fortschr. Zool. Syst. Evol.-Forsch., 1: 166-171. [physiology, hormones]

Bückmann, D., G. Starnecker, K.H. Tomaschko, E. Wilhelm, R. Lafont & J.P. Girault (1986): Isolation and identification of major ecdysteroids from the pycnogonid *Pycnogonum litorale* Strøm (Arthropoda: Pantopoda). - J. Comp. Physiol., (B. Biochemistry, systematics, environment, physiology), 156 (6): 759-766. [physiology, hormones]

Bullivant, J.S. (1959): Photographs of the bottom fauna in the Ross Sea. - N.Z. J. Sci., 2 (4): 485-497.

Bullivant, J.S. (1961): Photographs of the Antarctic bottom fauna. - Polar Record, Cambr., 10 (68): 505-508, 5 pls.

Calman, W.T. (1901): A catalogue of Crustacea and of Pycnogonida contained in the Museum of the University College, Dundee. Dundee, Univ. St. Andrews, v + 56pp.

Calman, W.T. (1909): The problem of the Pycnogonids. - Sci. Progr., London, 1909: 687-693.

Calman, W.T. (1910): Antarctic Pycnogons. - Nature, Lond., 84: 104.

Calman, W.T. (1915): The holotype of *Ammothea carolinensis* Leach (Pycnogonida). - Ann. Mag. Nat. Hist., (9), 15: 310-314.

- Calman, W.T. (1915): The holotype of *Nymphon gracilipes* Miers (Pycnogonida). - Ann. Mag. Nat. Hist., 15: 584-588.
- Calman, W.T. (1915): Pycnogonida. Brit. Antarct. (Terra Nova) Exped., 1910, Zool., 3 (1): 1-74.
- Calman, W.T. (1920): On a collection of Pycnogonida from the South Orkney Isles. - Ann. Mag. Nat. Hist., (9) 6: 244-247.
- Calman, W.T. (1922): The holotype of *Parazetes auchenicus* Slater (Pycnogonida). - Ann. Mag. Nat. Hist., (9) 9: 199-203.
- Calman, W.T. (1923): Pycnogonida of the Indian Museum. - Rec. Indian Mus., 25 (3): 265-299.
- Calman, W.T. (1927): 28. Report on the Pycnogonida. Zoological results of the Cambridge Expedition to the Suez Canal, 1924. - Trans. Zool. Soc. Lond., 22 (3): 403-410.
- Calman, W.T. (1929): The Pycnogonida. - J. Quekett Microsc. Club, London, (2)16(96): 95-106.
- Calman, W.T. (1933): A dodecapodous Pycnogonid. - Nature, London, 131 (3309): 242.
- Calman, W.T. & I. Gordon (1933): A dodecapodous Pycnogonid. - Proc. Roy. Soc. Lond., (B.), 113: 107-115.
- Calman, W.T. (1937): James Eights, a pioneer Antarctic naturalist. - Proc. Linn. Soc. Lond., 1936-1937, 149(4): 171-184.
- Calman, W.T. (1937): The type-specimens of *Pallene australiensis* Hoek (Pycnogonida). - Ann. Mag. Nat. Hist., (10) 20: 530-534.
- Calman, W.T. (1938): Pycnogonida. Sci. Rep. John Murray Exped., 1933-34, 5 (6): 147-166.
- Calvez, J.L. (1950). Un Pycnogonide nouveau pour la Méditerranée: *Nymphonella tapetis* Ohshima. - Arch. Zool. Exp. Gén., 86 (Notes & Rev., 3): 114-117.
- Camp, D.K., N.H. Whiting & R.E. Martin (1977): Pycnogonida. In: Nearshore marine ecology at Hutchinson Island, Florida, 1971-1974, V: Arthropoda. - Fla. Mar. Res. Publ., St. Petersburg, 25; p.10.
- Camp, J. & J. Ros (1980): Comunidades de sustrato duro del litoral NE Espanol, 8: Sistemática de los grupos menores. - Invest. Pesqu., 44 (1): 199-209.

- Campbell, A.C. & J.A. Nicholls (1976): Pycnogonida, p. 230. In: The Hamlyn Guide to the Seashore and Shallow Seas of Britain and Europe: 320 pp., London, etc.; Hamlyn.
- Candela, A., R. Sconfietti & A.R. Tornelli (1983): Ricerche sperimentali sulla dinamica stagionale delle zoocenosi intermareali della laguna di Venezia, 1. - Boll. Mus. Civ. Stor. Nat. Venezia, 33 (1982): 28-72.
- Candela, A., R. Sconfietti & A.R. Tornelli (1983): Ricerche sperimentali sulla dinamica stagionale delle zoocenosi intermareali della laguna di Venezia, 2. - Boll. Mus. Civ. Stor. Nat. Venezia, 34 (1983): 7-28.
- Capriles, V.A. (1970): Note on the occurrence of the pycnogonid *Pallenoides* (?) *spinulosa* Stock in Puerto Rican waters. - Caribb. J. Sci., 10 (1-2): 105.
- Carpenter, G.H. (1892): A new *Nymphon* and *Phoxichilus spinosus* on the Irish Coast. - Ir. Nat., 1: 42.
- Carpenter, G.H. (1892): Pycnogonida from Killala Bay. - Ir. Nat., 1: 168.
- Carpenter, G.H. (1892): Pycnogonida. Reports on the zoological collections made in Torres Straits by Prof. A.C. Haddon, 1888-89. - Sci. Proc. Roy. Dublin Soc., 7 (5) (40): 552-558, 1pl.
- Carpenter, G.H. (1893): Pycnogonids. Reports on the zoological collections made in Torres Straits by Professor A.C. Haddon, 1888-89. - Sci. Proc. Roy. Dublin Soc., 8 (1) (4): 21-27.
- Carpenter, G.H. (1893): On some Pycnogonida from the Irish coasts. - Sci. Proc. Roy. Dublin Soc., (NS) 8: 195-205, 1pl.
- Carpenter, G.H. (1895): A new British Pantopod (*Tanystylum conirostre* Dohrn). - Ir. Nat., 4: 297-302.
- Carpenter, G.H. (1898): On Pantopoda collected by Mr. W.S. Bruce in the neighbourhood of Franz-Jose Land, 1896-97. - J. Linn. Soc. Lond. (Zool.), 24: 626-634.
- Carpenter, G.H. (1900): Pycnogonida from the Arctic Seas, dredged by Mr. W. Bruce, 1897-98. - Sci. Proc. Roy. Dublin Soc., 9 (3): 279-282.
- Carpenter, G.H. (1904): Report on the Pantopoda collected by Prof. Herdman at Ceylon in 1902. Rep. Governm. Ceylon Pearl Oyster Fish. Gulf Manaar, 2, Suppl. Rep. XIII: 181-184, 1pl.

- Carpenter, G.H. (1905): The marine fauna of the coast of Ireland. Part VI. Pycnogonida. - Sci. Inv. Fish. Branch Ireland, 4: 171-178.
- Carpenter, G.H. (1905): Notes on the segmentation and phylogeny of the Arthropoda. - Quart. J. Microsc. Sci., 40: 469-491.
- Carpenter, G.H. (1907): Pycnogonida collected by Mr. J. Stanley Gardiner. - Proc. Linn. Soc. Lond., 1907: 11.
- Carpenter, G.H. (1908): Pycnogonida. The Percy Sladen Trust Expedition to the Indian Ocean. - Trans. Linn. Soc. Lond., (2) 12: 95-101.
- Carpenter, G.H. (1910): XVI. Pycnogonida from the Red Sea and Indian Ocean collected by Mr. Cyril Crossland. Reports on the Marine Biology of the Sudan and the Red Sea. - J. Linn. Soc. London (Zool.), 31 (207): 256-258.
- Carpenter, G.H. (1912): Pycnogonida. Clare Island Survey. - Proc. Roy. Irish Acad., Dublin, 31 (34): 1-4.
- Carus, J.V. (1863): Arthropoden. In: A. Gerstaecker (ed.), Handbuch der Zoologie, 2: 348-350.
- Carus, J.V. (1885): Pantopoda. In: Prodromus fauna mediterraneae sive descriptio animallum maris Mediterranei incolarum, 2: 283-288; Stuttgart.
- Caspers, H. (1950): Pantopoda, p. 138. In: Die Lebensgemeinschaft der Helgoländer Austernbank. - Helgol. Wiss. Meeresunters., 3: 119-169.
- Caspers, H. (1980): Fauna adapted to inhabiting the sponge *Damiriana hawaiiana* in Kanehoe Bay, Hawaii. (Abstr.). Int. Sympos. Biol. Management of Mangroves and Tropical Shallow Water Communities, 2: 20.
- Caullery, M. (1896): Pycnogonides. Résultats scientifiques de la campagne du "Caudan" dans le golfe de Gascogne, août-sept. 1895. - Ann. Univ. Lyon, 26: 361-364.
- Cavanna, G. (1876): Riassunto di una memoria sui Pignogonidi. - Bull. Soc. Ent. Ital., VIII: 292-297.
- Cavanna, G. (1876): Studi e ricerche sui Picnogonidi. - Pubbl. Real. Inst. Stud. Superiori Pratici Perfez. Firenze, 1: 249-264, 19pls.
- Cavanna, G. (1878): Alcune osservazioni intorno ad un recante lavoro del dott. A. Dohrn, sui Picnogonidi. - Bull. Soc. Ent. Ital., Anno X, trim. 4: 244-247.

- Cerame-Vivas, M.J. & I.E. Gray (1966): The distributional pattern of benthic invertebrates of the continental shelf off North Carolina. - *Ecology*, 47: 260-270.
- Chaudonneret, J. (1980): Table ronde sur la métamérie, 6: Autres Chélicérates. - *Bull. Soc. Zool. France*, 104 (3): 287-290.
- Child, C.A. (1970): Pycnogonida of the Smithsonian-Bredin Pacific-Expedition, 1957. - *Proc. biol. Soc. Wash.*, 83 (27): 287-308.
- Child, C.A. & J.W. Hedgpeth (1971): Pycnogonids of the Galápagos Islands. - *J. nat. Hist.*, 5: 609-634.
- Child, C.A. (1972): A new species of *Anoropallene* (Pycnogonida) from the Hawaiian Islands. - *Proc. biol. Soc. Wash.*, 85 (11): 147-150.
- Child, C.A. (1974): *Hedgpethius tridentatus*, a new genus and species, and other Pycnogonida from Key West, Florida. - *Proc. biol. Soc. Wash.*, 87 (43): 493-500.
- Child, C.A. (1975): Pycnogonida of Western Australia. - *Smiths. Contr. Zool.*, 190: 1-29.
- Child, C.A. (1975): The Pycnogonida types of William A. Hilton. I. Phoxichilidiidae. - *Proc. biol. Soc. Wash.*, 88 (19): 189-210.
- Child, C.A. (1977): On some Pycnogonida of French Oceania. - *Proc. biol. Soc. Wash.*, 90 (2): 440-446.
- Child, C.A. (1977): Four new species of *Anoplodactylus* (Pycnogonida) from the Western North Atlantic. - *Proc. biol. Soc. Wash.*, 90 (3): 584-596.
- Child, C.A. (1978): Gynandromorphs of the pycnogonid *Anoplodactylus portus*. - *Zool. J. Linn. Soc. Lond.*, 63 (1-2): 133-144. [genital system]
- Child, C.A. (1979): Shallow-water Pycnogonida of the Isthmus of Panamá and the coasts of Middle America. - *Smiths. Contr. Zool.*, 293: 1-86; Washington.
- Child, C.A. (1982): Pycnogonida from Carrie Bow Cay, Belize. - *Smiths. Contr. mar. Sci.*, 12: 355-380; Washington.
- Child, C.A. (1982): Pycnogonida of the Western Pacific Islands. I. The Marshall Islands. - *Proc. biol. Soc. Wash.*, 95 (2): 270-281; Washington.
- Child, C.A. (1982): Deep-sea Pycnogonida from the North and South Atlantic Basins. - *Smiths. Contr. Zool.*, 349: 1-54.

- Child, C.A. & K. Nakamura (1982): A gynandromorph of the Japanese Pycnogonid *Anoplodactylus gestiens* (Ortmann). - Proc. biol. Soc. Wash., 95 (2): 292-296. [genital system]
- Child, C.A. (1983): Pycnogonida of the western Pacific Islands, II. Guam and the Palau Islands. - Proc. biol. Soc. Wash., 96 (4): 698-714.
- Child, C.A. (1986): Spotlight on a sea-going "spider". - Anim. Kingdom, 89 (1): 36-38.
- Child, C.A. & G.R. Harbison (1986): A parasitic association between a Pycnogonid and a Scyphomedusa in Midwater. - J. Mar. biol. Ass. U.K., 66: 113-117.
- Child, C.A. (1987): The Pycnogonida types of H.V.M. HALL. - Proc. biol. Soc. Wash., 100 (3): 552-558; Washington.
- Child, C.A. (1987): Pycnogonida of Enewetak Atoll. In: D.M. Devaney et al. (eds.). The natural history of Enewetak Atoll, Vol. II, Biogeography and Systematics, chapter 16, pp. 179-180. Dept. Energy, Off. Sci. Tech. Inf., Oak Ridge, Tennessee.
- Child, C.A. (1987): *Ammothea verenae* and *Sericosura venticola*, two new Hydrothermal Vent-associated Pycnogonids from the Northeast Pacific. - Proc. Biol. Soc. Wash., 100 (4): 892-901.
- Child, C.A. (1987): New and little known Pycnogonida from Antarctic and Subantarctic Waters. - Proc. biol. Soc. Wash., 100 (4): 902-916.
- Child, C.A. (1988): Pycnogonida of the Western Pacific Islands, III: recent Smithsonian-Philippine expeditions. - Smiths. Contr. Zool., 468: 1-32; Washington.
- Child, C.A. (1988): Pycnogonida from Aldabra Atoll. - Bull. Biol. Soc. Wash., 8: 45-78.
- Child, C.A. (1988): 45: Pycnogonida. In: R.P. Higgins & H. Thiel (eds.). - Introduction to the study of meiofauna, 1988, pp. 423-424; Washington.
- Child, C.A. (1989): Pycnogonida of the Western Pacific Islands. 6. *Sericosura cochleifovea*, a new hydrothermal vent species from the Marianas Back-Arc Basin. - Proc. biol. Soc. Wash., 102 (3): 732-737.
- Child, C.A. (1990): *Prototrygaeus jordanae*, a new species of pycnogonid from Monterey Bay, California. - Proc. biol. Soc. Wash., 103 (1): 157-160.

- Child, C.A. (1990): Pycnogonida of the Western Pacific Islands, VIII. Recent collections from islands of the Great Barrier Reef, Australia. - Proc. biol. Soc. Wash., 103 (2): 311-335; Washington.
- Child, C.A. (1990): *Nymphon stipulum*, a new pycnogonid species from southern California. - Beaufortia, 41 (6): 39-43.
- Child, C.A. (1991): Pycnogonida of the Western Pacific Islands, IX. A shallow-water Guam survey, 1984. - Proc. biol. Soc. Wash., 104 (1): 138-146.
- Chimenz, C., P.M. Brignoli & G. Basciano (1979): Pantopodi del porto di Civitavecchia e Dintorni (Italia Centrale). - Cah. Biol. Mar., 20: 471-497.
- Chimenz, C. & V. Cottarelli (1986): Soft bottom Pycnogonida from the Gulf of Salerno (Italy). - Oebalia, 13: 137-146.
- Chimenz, C., V. Cottarelli & M. Tosti (1991): Researches on the Mediterranean Pycnogonida. 1. *Anoplodactylus compositus* n.sp. (Arthropoda, Chelicerata). - Fragm. Ent., 23 (1): 7-13.
- Chimenz-Gusso, C. & E.T. Rivosecchi (1972): Nuovi richercche sul fouling del Porto di Civitavecchia. 2. Rendiconti Accad. Naz. dei XL, (4) 20/21: 199-216.
- Chimenz-Gusso, C., E. Fresi, F. Civelli, L. Mazella, M. Pansini & R. Pronzato (1978): Distribution of hard bottom communities along a light gradient in a superficial marine cave, II; Pantopoda. - Mem. Biol. Mar. Oceanogr., new ser., 8 (4): 91-103.
- Citarella, G. (1982): Le zooplancton de la baie de Shediac (Nouveau-Brunswick). - J. Plankton Res., 4 (4): 791-812.
- Claparède, R.E. (1863): Beobachtungen über Anatomie und Entwicklungsgeschichte wirbelloser Thiere, an der Küste der Normandie angestellt; pp. 102-105.
- Clark, W.C. (1956): A new species of *Pycnogonum* from Banks Peninsula, New Zealand. - Rec. Canterbury Mus., 7 (2): 171-173.
- Clark, W.C. (1958): Some Pycnogonida from Cook Strait, New Zealand, with descriptions of two new species. - Zool. Publ. Victoria Univ. Wellington, 23: 1-7.
- Clark, W.C. (1961): Two new pycnogonids from the Maldives Islands. - Ann. Mag. Nat. Hist., 13 (3): 291-296.

- Clark, W.C. (1962): *Pallenopsis triregia*, a new species of Pycnogonida from near Three Kings Islands, New Zealand. - N.Z. J. Sci., Wellington, 5 (4): 517-520.
- Clark, W.C. (1963): Australian Pycnogonida. - Rec. Austr. Mus., 26 (1): 1-81.
- Clark, W.C. (1970): New Pycnogonida from Queensland. - Trans. Roy. Soc. N.Z. (Biol. Sci.), 12 (3): 13-20.
- Clark, W.C. (1972): Pycnogonida of the Snares Islands. - N.Z. J. Mar. Freshw. Res., 5 (2): 329-341.
- Clark, W.C. (1972): Pycnogonida of the Antipodes Islands. - N.Z. J. Mar. Freshw. Res., 5 (3/4): 427-452.
- Clark, W.C. (1973): New species of Pycnogonida from New Britain and Tonga. - Pac. Sci., 27 (1): 28-33.
- Clark, W.C. (1976): The genus *Rhynchothorax* Costa (Pycnogonida) in New Zealand waters. - J. Roy. Soc. N.Z. (Biol. Sci.), 6 (3): 287-296.
- Clark, W.C. (1977): The genus *Ammothea* Leach (Pycnogonida) in New Zealand waters: New species and a review. - J. Roy. Soc. N.Z., 7 (2): 171-187.
- Clark, W.C. (1977): The genus *Tanystylum* Miers, 1879 (Pycnogonida) in the Southern Oceans. - J. Roy. Soc. N.Z., 7 (3): 313-338.
- Clark, W.C. & A. Carpenter (1977): Swimming behavior in a pycnogonid. - N.Z. J. Mar. Freshw. Res., 11: 613-615.
- Clark, W.C. (1983): Platyhelmenthes, Nematoda and Pycnogonida. In: P.J. Brownsey & A.N. Baker (eds.). The New Zealand Biota: What do we know after 200 years? - Natn. Mus. N.Z., Misc. Ser., 7: 60-61.
- Clarke, J.M. (1916): The reincarnation of James Eights, Antarctic explorer. - Illustr. Sci. Monthly ?, 2 (2): 189-202.
- Clarke, K.U. (1973): The Pycnogonida. In: The Biology of the Arthropoda: 185, 187, 188; 9.5 London: Edward Arnold.
- Clememçon, H. (1961): Asselspinnen-Bewohner der Meeresküsten. - Mikrokosmos, 50: 262-270.
- Cloudsley-Thompson, J.L. (1973): Some littoral and terrestrial arthropoda from Cumbrae, Scotland. - Ent. Monthly Mag., 109 (1304-1306): 59.

- Codreanu, R. & M. Bacescu (1951): Clasificarea regnului animal; Crustacea; Pycnogonida. In: Academia Republ. Pop. Rom. Indrumtdor, 1: 7-12, 147-169, 182-184.
- Coelho, P.A. & G. Calabria (1987): Litoral de Alagoas e Sergipe: Pantopodas. - An. Soc. Nordest., Zool., 1 (1): 55.
- Cole, L.J. (1901): Notes on the habits of pycnogonids. - Biol. Bull. Woods Hole Mar. Biol. Lab., 2 (5): 195-207.
- Cole, L.J. (1901): On *Discoarachne brevipes* Hoek, a pycnogonid from South Africa. - Zool. Jb. (Syst., Geogr., Biol.), 15 (2): 243-248.
- Cole, L.J. (1904): Pycnognida of the West Coast of North America. - Harriman Alaska Exped., 10: 249-330.
- Cole, L.J. (1904): Pycnognida collected at Bermuda in the summer of 1903. - Proc. Soc. Nat. Hist., 31 (8): 315-328.
- Cole, L.J. (1905): Ten-legged pycnogonids, with remarks on the classification of the Pycnogonida. - Ann. Mag. Nat. Hist., (7) 15: 405-415.
- Cole, L.J. (1906): Feeding habits of the pycnogonid *Anoplodactylus latus*. - Zool. Anz., 29 (24): 740-741.
- Cole, L.J. (1906): A new Pycnogonid from the Bahamas. - Am. Nat., 40 (471): 217-222.
- Cole, L.J. (1909): XIX. Pycnogonida. Reports on the scientific results of the expedition to the Eastern Tropical Pacific in charge of Alexander Agassiz, by the U.S. Fish Commission Steamer Albatross from October 1904 to March 1905... - Bull. Mus. Comp. Zool. Harvard Coll., 52 (11): 185-192.
- Cole, L.J. (1910): Peculiar habitat of a pycnogonid (*Endeis spinosus*) new to North America, with observations on the heart and circulation. - Biol. Bul. Woods Hole Mar. Biol. Lab., 18 (4): 193-203. [circulatory system]
- Cole, L.J. (1921) Report on the Pycnogonida collected by the Canadian Arctic Expeditions, 1913-1918. Crustacea, F. Rep. Can. Arct. Exped., 7: 1-6.
- Collingbourne, H. (1976): The seaspider is a pycnogonid. - Aquarist and Pondkeeper (London), 41 (1): 23-24.
- Corrêa, D.D. (1948): *Callipallene gabriellae*, novo Pantópodo de Santos. - Pap. Avuls. Dept. Zool., São Paulo, 9 (1): 1-12.

Costa, O.G. (1836): Aracnidi. Famiglia II. Picnogonidi. Fauna del Regno di Napoli: 5-10, 2 pls.

Costa, O.G. (1861): Microdoride mediterranea, o descrizione de' poco ben conosciuti od affatto ignoti viventi minuti e microscopici del Mediterraneo, etc., 1: i-xviii, 1-80.

Costa, S. (1960): Recherches sur les fonds à *Halarachnion spatulatum* de la Baie de Marseille. - Vie Milieu, 11 (1): 1-68.

Coustalin, J.-B. (1972): Méthodologie expérimentale en vue de la création de récifs artificiels. - Téthys, 3 (4): 827-840.

Crapp, G.B. (1967): The ecology and identification of Anglesey pycnogonids. M.Sc. Thesis, Univ. Wales.

Crawshay, L.R. (1912): On the fauna of the English Channel. - J. Mar. Biol. Ass. Plymouth, (N.S.) 9: 292.

Crawshay, L.R. (1912): On the fauna of the outer western area of the English Channel. - J. Mar. Biol. Ass. U.K., (N.S.), 9: 360.

Crome, W. (1961): Pantopoda-Asselsspinnen. In: E. Stresemann (ed.): Exkursionsfauna von Deutschland, Wirbellose I, pp. 363-369, 2nd. edition. Berlin: VEB Volk und Wissen.

Crothers, J.H. (1966): Dale Fort Marine Fauna, 2nd ed. - Field Stud., 2 (Suppl.): i-xxiv, 1-169.

Crowder, W. (1928): Pycnogonida. In: A naturalist at the seashore: 31-37, 33, 48. New York, Century Publ. Co.

Crowder, W. (1931): Order Pycnogonida, in: Between the tides: 334-339; 319-326; New York, Dodd Mead.

Cubit, J. & S. Williams (1983): The invertebrates of Galeta Reef (Caribbean, Panama): A species list and bibliography. - Atoll Res. Bull., 269: 1-45.

Cuénot, L. (1891): Etudes sur la sang et les glandes lymphatiques dans la série animale. 2ème partie. Invertébrés. Pantopoda. - Arch. Zool. Exp. Gén., (2) 9: 406-407.

Cuénot, L. (1921): Contributions à la faune du Bassin d'Arcachon. VIII. Pycnogonida. - Arch. Zool. Exp. Gén., 60 (Notes & Rev., 2): 21-32.

Czeczuga, B. (1981): Presence of carotenoids in some invertebrates of the Antarctic Coast. - Comp. Biochem. Physiol., London (B: Comp. Biochem.), 69 (3): 611-616.

- Daguzan, J. (1969): Contribution à l'étude de la faune intertidale de la pointe rocheuse de Penvins (Morbihan). Liste faunistique à l'exclusion des Mollusques. - Bull. Soc. Sci. Bretagne, 44: 249-255.
- Dahl, E., L. Laubier, M. Sibuet & J.O. Strömberg (1976): Some quantitative results on benthic communities of the deep Norwegian Sea. - Astarte, 9: 61-74.
- Daiber, M. (1921): Die Pantopoden (Pycnogoniden). In: Lang's Handbuch der Morphologie der Wirbellosen Tiere: 635-643; Jena, G. Fischer.
- Dakin, W.J., I. Bennett & E. Pope (1960): Pycnogonida. In: Australian Seashores: pl. 49.4; Sydney, London, Melbourne; Angus & Robertson.
- Dall, W.H. (1876): On the marine faunal regions of the North Pacific; an introductory note to the report on Alaskan Hydrozoans, by Mr. Clark. - Proc. Acad. Nat. Sci. Philadelphia 1876, 8: 205-208.
- Dana, J.D. (1849): *Astridium orientae* n.gen. n.sp. - Proc. Am. Acad. Arts Sci., Boston, 2: 61.
- Dana, J.D. (1852): *Pycnogonum orientale* (Dana). U.S. Expl. Exped. 1838-42, under the command of Charles Wilkes, 13: 1382-1391.
- Daniel, A. & J.K. Sen (1980): Studies on the pycnogonids from the collections of the zoological survey of India, Calcutta, together with notes on their distribution in the Indian Ocean. - J. Mar. biol. Ass. India, 17 (2) (1975): 160-167.
- D'Arcy Thompson, W. (1901): A catalogue of Crustacea and of Pycnogonida contained in the Museum of University College, Dundee. Dundee, Univ. St. Andrews, v + 56 pp.
- D'Arcy Thompson, W. (1909): Pycnogonida. In: S.F. Harmer & B.E. Shipley (eds.). The Cambridge Natural History: 504-542; London: Macmillan.
- Daro, M.H. (1969): Étude écologique d'un brise-lames de la côte belge. 1. Description et zonation des organismes. - Ann. Soc. roy. Zool. Belg., 99 (3-4): 111-152.
- Daro, M.H. (1971): Étude écologique d'un brise-lames de la côte belge 2. Biologie et développement saisonnier des espèces. - Ann. Soc. roy. Zool. Belg., 110 (3): 159-189.

Davenport, J.N., D.A. Blackstock, D.A. Davies & M. Yarrington (1987): Observations on the physiology and integumentary structure of the Antarctic Pycnogonid *Decolopoda australis*. - *J. Zool.*, 211 (3): 451-465.

Dawson, A.B. (1934): The coloured corpuscles of the blood of the purple sea-spider, *Anoplodactylus latus* Wilson. - *Biol. Bull. Woods Hole Mar. Biol. Lab.*, 66: 62-68, 1pl. [circulatory system]

Dawson, E.W. (1979): Pycnogonida. In: Catalogue of type and figured specimens in the New Zealand Oceanographic Institute. - *N.Z. Oceanogr. Inst. Mem.*, 76: 58-60.

Dawydooff, C. (1928): *Traité d'Embryologie Comparée des Invertébrés*: 1-930. Paris: Masson. [embryology]

Day, J.H. (1969): Class Arachnida: Pycnogonida (Sea Spiders). In: *A guide to marine life on South African Shores*: 120-122; Cape Town, A.A. Balkema.

Dayton, P.K., G.A. Robilliard & R.T. Paine (1970): Benthic faunal zonation as a result of anchor ice at McMurdo Sound, Antarctica. In: M.W. Holdgate, *Antarctic Ecology*: 244-258. Sci. Committ. Antarct. Res.; London & New York, Academic Press.

Dearborn, J.H. (1967): Food and reproduction of *Glyptonotus antarcticus* (Crustacea: Isopoda) at McMurdo Sound, Antarctica. - *Trans. Roy. Soc. N.Z. (Zool.)*, 8 (15): 163-168.

Dearborn, J.H. (1968): Benthic invertebrates. *Austr. Nat. Hist.*, Sydney, 1968 (Dec.): 134-139.

de Bruyne, R.H. & A. Oosterbaan (1987): Strandwerkkamp 1986. - *Terschelling Zeepaerd*, 47 (4): 94-101.

de Castellanos, Z.J.A. (1965): Adiciones a la Pantopodofauna Argentina. - *Physis*, 25 (69): 13-16.

de Haro, A. (1965): Comunidad de Picnogónidos en el alga parda *Halopteriss scoparia* (L.). - *Bol. Real. Soc. Esp. Hist. Nat.*, Madrid (Secc. Biol.), 63: 213-218.

de Haro, A. (1965): Picnogónidos de la fauna española. Picnogónidos de las islas Medas (Gerona). - *Publ. Inst. Biol. Apl.*, Barcelona, 39: 137-145, pls. 1-5.

- de Haro, A. (1966): Distribución ecológica de los pícnogónidos entre algas y posidonia mediterráneas. - Inv. Pesq., 30: 661-667.
- de Haro, A. (1966): Pícnogónidos del alga parda *Halopteris scoparia* (L.) de las islas Medas (Gerona). - Bol. Real Soc. Esp. Hist. Nat., Madrid (Secc. Biol.), 64: 5-14.
- de Haro, A. (1967): Relaciones entre Pícnogónidos e Hidroideos en el medio posidónico. - Bol. Real Soc. Esp. Hist. Nat., Madrid (Secc. Biol.), 65: 301-303.
- de Haro, A. (1967): Pícnogónidos de la fauna española. Pícnogónidos de las posidonia de Blanes (Gerona). - Publ. Inst. Biol. Apl., Barcelona, 43: 103-123.
- de Haro, A. (1967): Los Pícnogónidos y su posición sistemática dentro de los Artrópodos. - Bol. Real. Soc. Esp. Hist. Nat., Madrid (Secc. Biol.), 65: 367-375.
- de Haro, A. (1969): Relaciones filogenéticas de los Pícnogónidos. I. Simposio internacional de Zoofilogenia, 13-17 octubre de 1969: 293-304; Univ. Salamanca; Fac. Cienc.
- de Haro, A. (1978): Ecological distribution of pycnogonids on the Catalan coast. In: Sea spiders (Pycnogonida). - Zool. J. Linn. Soc. Lond., 63 (1-2): 181-196.
- de la Fuente, J.A. (1978): Consideraciones sobre filogenia y taxonomía de los artrópodos. - Boln. Asoc. Esp. Ent., 2: 33-40.
- de Lamarck, J.B. (1818): Les Pycnogonides. In: Histoire naturelle des animaux sans vertèbres, 5: 72-77; Paris.
- Dell, R.K. (1972): Antarctic benthos. - Adv. Mar. Biol., 10: 1-216. [Pycnogonids: 60-64]
- Demir, M. (1952): Pycnogonida. Les Invertébrés benthoniques des Détroits et des îles Marmara. - Hidrobiol., 2 (A): 1-654. [in Turkish].
- de Quatrefages, A. (1844): Recherches sur les Pycnogonides. Extr. P.-v. Séanc. Soc. Phil. Paris, 1844: 86.
- de Quatrefages, A. (1844): Observations générales sur le phlébentérisme: anatomie des Pycnogonides. - C.R. Acad. Sci., Paris, 19: 1150-1157.
- de Quatrefages, A. (1845): Études sur les types inférieures de l'embranchement des Annelés. Mémoire sur l'organisation des Pycnogonides. - Ann. Sci. Nat., Paris, (4) 3: 69-83.

- Derjugin, K.M. (1935): Pantopoda of the Polar Seas within the U.S.S.R. - Arkticheski Inst. SSSR, (Material for the Study of the Arctic), 4: 1-40. [in Russian]
- de Villiers, A. (1976): Littoral ecology of Marion and Prince Edward Islands (Southern Ocean). - S. Afr. J. Antarct. Res., suppl., 1: 1-40.
- Diaz, M.Z., M.I. Garcia, O. Rivero B. & T. Eligio, C. (1985): Lista de especies y bibliografia de los picnogónidos (Arthropoda) del Mediterraneo Americano. - Revta. Invest. Mar., 6 (1): 29-38.
- Dittmer, J.D. (1981): The distribution of subtidal macrobenthos in the estuaries of the Rivers Ems and Weser. In: N. Dankers, H. Kühl & J. Wolff (eds.). Invertebrates of the Wadden Sea, pp. 188-206. Leiden: Wadden Sea Working Group, Rep. 4.
- Dörjes, J. (1971): Der Golf von Gaeta (Tyrrhenisches Meer). IV. Das Makrobenthos und seine küstenparallele Zonierung. - Senckenbergiana marit., 3: 302-346.
- Dogiel, V.A. (1911): A short account of work on Pycnogonida done during June, 1922, at Cullercoats. - Rep. Sci. Invest., Northumberland Sea Fish. Comm., 1910-11: 26-27.
- Dogiel, V.A. (1911): Ein interessanter Fall von atavistischer Missbildung bei einer Pantopodenlarve. - Zool. Anz., 38 (13): 321-323.
- Dogiel, V.A. (1911): Studien über die Entwicklungsgeschichte der Pantopoden. - Nervensystem und Drüsen der Pantopodenlarven. - Z. Wiss. Zool., 98: 110-146. [anatomy, nervous system, gland]
- Dogiel, V.A. (1911): Note on pycnogonid researches. In: Ann. Rep. Mar. Biol. Ass. W.-Scotland, 1911.
- Dogiel, V.A. (1913): Materials for the natural history of the Pantopoda. Doc. Diss., St. Petersburg: 1-229 [in Russian]
- Dogiel, V.A. (1951): Class Pycnogonida (Pantopoda). In: Rukovodstvo po zoologii, 3: 45-106; Sovietskaya Nauk [in Russian]
- Dogiel, V.A. (1954): Oligomersation of homologous organs: 1-368; Izdateljstvo Leningrad Gosudarstvo Univ. [in Russian]
- Dohrn, A. (1869): Ueber Entwicklung und Bau der Pycnogoniden. - Jenaische Z. Med. Naturwiss., Jena, 5: 138-157. [anatomy]
- Dohrn, A. (1879): Untersuchungen über Pycnogoniden. - Mitt. Zool. Stat. Neapel, 1: 28-39.

- Dohrn, A. (1881): Die Pantopoden des Golfes von Neapel und der angrenzenden Meeresabschnitte. - Monogr. Fauna Flora Golf. Neapel, 3: 1-252.
- Dons, C. (1913): Et og andet fra mine ekskursioner (Foreløbig meddelelse). - Tromsø Mus. Aarsberetning, 1913: 35-60.
- Dons, C. (1930): Zoologiske Notiser VI. Om enkelte Pantopoder. - Kongel. Norske Vidensk. Selsk. Forhandl., 5: 194-197.
- Dons, C. (1930): Zoologiske Notiser VIII. Endel pantopoder fra Nord-Norge. - Kongel. Norske Vidensk. Selsk. Forhandl., 2 (46): 163-165.
- Dons, C. (1933): Norges Strandfauna. 1. Pantopoder. - Kongel. Norske Vidensk. Selsk. Forhandl., 5 (50): 194-197.
- Douglas, E.L., J.W. Hedgpeth & E. Hemmingsen (1969): Oxygen consumption of some Antarctic pycnogonids. - Antarct. J. U.S., 4: 109. [respiration]
- Drago, N. & G. Albertelli (1978): Etude faunistique et bionomique du littoral de Cogoleto (Golfe de Genes). - Tethys, 8 (2): 203-212.
- Drago, N., G. Albertelli & H. Cattaneo (1978): Osservazioni faunistiche sul bentos dell'Isole di Capraia. - Ann. Mus. Civ. Stor. Nat. "Fiacomo Doria", 82: 72-77.
- Dresco-Derouet, L. (1978): Métabolisme respiratoire de *Nymphon gracile* Leach et d'*Endeis spinosa* (Montagu) (Pycnogonida): Comparison avec quelques espèces de crustacés intercotidaux. - Cah. Biol. Mar., 19: 309-315. [physiology, respiration]
- Dujardin, F. (1849): Résumé d'un mémoire sur les Pycnogonides. - C.R. Hebd. Séanc. Acad. Sci., Paris, 29: 28-29.
- Duris, Z. (1985): Interesting factss on arthropods of the White Sea. - Czech-Ziva, 33 (1): 21-23. [in Czechoslovakian]
- Duris, Z. & P.E. King (1983): Spermatogenesis in a Pycnogonid *Achelia echinata* Hodge. - Acta Zool., Stockholm, 64 (4): 227-233. [cytology, genital system]
- Durrell, L.G. (1977): Pycnogonida. In: The Black Bock: 238; London, Faber & Faber.
- Eales, N.B. (1950): The littoral fauna of Great Britain, 2nd ed.: i-xvii, 1-305; Cambr. Univ. Press.

Eights, J. (1835): Description of a new animal belonging to the Arachnides of Latreille; discovered in the sea along the shores of the New South Shetland Islands. - Boston J. Nat. Hist., 1 (2): 203-206.

Eldredge, L.G. (?1968): Catalog of invertebrate type specimens. Pac. Sci. Inf. Center, Honolulu: 22pp.

Ekman, S. (1953): Zoogeography of the Sea, 417. Textbooks of Marine Biology and Ecology, 33: 213-221.

El-Hawawi, A.S.N. & P.E. King (1978): Salinity and temperature tolerance by *Nymphon gracile* (LEACH) and *Achelia echinata* (HODGE) (Pycnogonida). - J. exp. mar. Biol., 33: 213-221;

El-Hawawi, A.S.N. & P.E. KING (1978): Spermatogenesis in a Pycnogonid *Nymphon gracile* (Leach). - J. Submicrosc. Cytol., 10 (3): 345-356. [cytology, genital system]

El-Hawawi, A.S. (1980): Aspects of sperm axoneme patterns in pycnogonids. - Proc. Conf. Biol. Aspects Saudi Arabia, 4: 54. [cytology, genital system]

Elliot, P., P.E. King, C.I. Morgan, P.J.A. Bugh, A. Smith & S.L.A. Wheeler (1990): Chelicerata, Uniramia, and Tardigrada; pp. 553-627, in: Hayward, P.J. & J.S. Ryland (eds.). The marine fauna of the British Isles and north-west Europe. Vol. 1. Introduction and protozoans to arthropods. Oxford Univ. Press: i-xvi, 1-627.

Ellis, D.V. (1960): Marine infaunal benthos in arctic North America. - Tech. Pap. Arct. Inst. North America, 5: 1-53.

El Maghraby, A.M. & E.J. Perkins (1956): Additions to the marine fauna of Whitstable. - Ann. Mag. Nat. Hist., (12), 9: 481-496.

Elmhirst, R. (1910): List of Pycnogonida collected in the Clyde area. - Trans. Nat. Hist. Soc. Glasgow, (N.S.), 3 (2): 146-149.

Entrop, B. (1951): *Pycnogonum littorale* en de Paardenanemmon. - De Kor, 1951 (3): 16-17.

Esmark (?) (1861): Esmark foreviste en *Phoxichilus*, som i sin gigantiske størrelse, i sin tykke kraftfulle Bygning legner meget *Ph. proboscideus*, som Sabine har opdaget ved Nord-Georgiens Øer i Baffinsbugten, men fra hvilken den i enkelte Dele synes at adskille sig. In: Expl. ble funnet flytende i havoverflat mellom Spitsbergen og Bjørnøya og innsendt til Universitetet av stortingsman Rostad i Hammerfest. - Forhandl. i Vidensk.-Selsk. Christiania, 1860: 64.

Ewing, H.E. (1928): The legs and the leg-bearing segments of some primitive arthropod groups, with notes on leg segmentation in the Arachnida. - Smiths. Misc. Coll., 80 (11): 1-41, 12pls.

Exline, H. (1936): Pycnogonids from Puget Sound. - Proc. U.S. Natn. Mus., 83 (2991): 414-422.

Fabricius, J.C. (1787): Mantissa Insectorum. Vol. 2, p. 368; Hafniae.

Fabricius J.C. (1794): Entomologia Systematica Emendata et aucta, 4: 416-417; Hafniae.

Fabricius, O. (1780): Fauna Groenlandica; xvi + 452 pp., 1 pl.

Fage, L. (1932): Pêches planctoniques à la lumiere effectuès à Banyuls-sur-Mer et à Concarneau. II. Pycnogonides. - Arch. Zool. Exp. Gén., 74 (14): 249-261.

Fage, L. (1933): Migrations verticales des animaux benthiques litoraux. - Rapp. P.-v. Cons. Perm. Expl. Mer, 85: 60-69.

Fage, L. (1942): Pycnogonides de la côte occidentale d'Afrique. - Arch. Zool. Exp. Gén., 82 (Notes & Rev., 2): 75-90.

Fage, L. (1949): Pycnogonides du Congo Belge. - Arch. Zool. Exp. Gén., 86 (Notes & Rev., 1): 20-31.

Fage, L. (1949): A propos de quelques Pycnogonides du Congo belge. - Bull. Inst. Roy. Colon. Belge, 20 (2): 568-574.

Fage, L. (1949): Classe des Pycnogonides. In: P.-P. Grasse (ed.), Traité de Zoologie: Anatomie, Systématique, Biologie, 6: 906-941; Paris: Masson.

Fage, L. (1951): Sur un Pycnogonide de l'expédition suédoise des grands fonds, 1947-48. - Rep. Swed. Deep-Sea Exped., 2 (7): 93-97.

Fage, L. (1952): 2. Pycnogonides. Missions du bâtiment polaire Commandant Charcot, récoltes faites en Terre Adélie (1950). - Bull. Mus. natn. Hist. nat., Paris, (2)24(2): 180-186.

Fage, L. (1952): Pycnogonides de la Terre Adélie. - Bull. Mus. natn. Hist. nat., Paris, (2)24(3): 263-273.

Fage, L. (1952): Sur quelques Pycnogonides de Dakar. - Bull. Mus. natn. Hist. nat., Paris, (2)24(6): 530-533.

Fage, L. (1953): Deux Pycnogonides nouveaux de la côte occidentale d'Afrique. - Bull. Mus. Natn. Hist. Nat., Paris, (2)25(4): 376-382.

- Fage, L. (1954): Remarques sur les Pycnogonides abyssaux. - Un. Int. Sci. Biol., (B), 16: 49-56.
- Fage, L. (1956): Sur deux espèces de Pycnogonides de Sierra Leone. - Bull. Mus. natn. Hist. nat., Paris (2)28(3): 290-295.
- Fage, L. (1956): Les Pycnogonides du genre *Nymphon*. - Galathea Rep., 2: 159-165.
- Fage, L. (1956): Les Pycnogonides (excl. le genre *Nymphon*). - Galathea Rep., 2: 167-182.
- Fage, L. (1959): Pycnogonides. - Res. Sci. Camp. Calypso, 4: 235-239.
- Fage, L. & J.H. Stock (1966): Pycnogonides. Résultats Scientifiques du Campagne de la Calypso aux îles de Cap Vert (1959). - Ann. Inst. Océanogr. Monaco, 44: 315-327.
- Faraggiana, R. (1940): Pantopodi del Mare Ligure. - Boll. Mus. Zool. Anat. Comp. Real. Univ. Torino, (3)48(112): 145-158.
- Fauvel, P. (1905): Histoire naturelle de la Presqu'île du Cotentin. III. La faune. In: Cherbourg et le Cotentin (Congr. A.F.A.S. Cherbourg, 3-10 août 1905). Cherbourg. Le Maout.
- Faxon, W. (1882): Bibliography to accompany selections from embryological monographs. Harvard Mus. Coll., 9: 247 ?.
- Filhol, H. (1884): Explorations sous-marines, voyage du "Talisman". - La Nature (Paris), 12 (1): 198-202.
- Filhol, H. (1885): La vie au fonds des mers. - La Nature (Paris), 13 (2): 227-230.
- Filhol, H. (1885): Pycnogonides. In: La vie au fonds des mers. Les explorations sous-marines et les voyages du Travailleur et du Talisman: 150-154.
- Fine, M.L. (1970): Faunal variation on pelagic *Sargassum*. - Mar. Biol., 7 (2): 112-122.
- Firstman, B. (1971): The relationship of the cheliceral arterial system to the evolution of the endosternite. Ph.D. Thesis, Stanford Univ. [anatomy, circulatory system]
- Firstman, B. (1973): The relationship of the Chelicerate arterial system to the evolution of the endosternite. - J. Arachnol., 1 (1): 1-54. [anatomy, circulatory system]

- Fish, C.J. (1925): Seasonal distribution of the plankton of the Woods Hole region. - Bull. Bur. Fish., Wash., 41: 91-179.
- Flynn, T.T. (1918): On a pycnogonid of the genus *Halosoma*, from New South Wales. - Pap. Proc. Roy. Soc. Tasmania, 1918: 11-15.
- Flynn, T.T. (1919): Two new Australian Pycnogonida. - Pap. Proc. Roy. Soc. Tasmania, 1918: 91-100.
- Flynn, T.T. (1919): A re-examination of Professor Haswell's types of Australian Pycnogonida. - Pap. Proc. Roy. Soc. Tasmania, 1919: 70-92.
- Flynn, T.T. (1928): The Pycnogonida of the marine survey of South Africa. - Rep. Fish. & Mar. Biol. Surv. Union S. Afr., 6 (Spec. Rep., I): 3-36
- Flynn, T.T. (1929): Pycnogonida from the Queensland coast. - Mem. Qd. Mus., 9 (3): 252-260.
- Forbes, E. (1859): In: R. Godwin-Austen (ed.). Natural History of the European Seas.
- Forsman, B. (1972): Evertebrater vid svenska Östersjö-kusten. - Zoologisk Revy, 34: 32-56.
- Fourtner, C.R. & R.G. Sherman (1973): Chelicerate skeletal nouromuscular systems. - Am. Zool., 13 (2): 271-289. [anatomy, musculature, nervous system]
- Franc, A. (1951): Le zooplancton de la région de Dinard-Saint Malo. - Bull. Lab. Mar. Dinard, 34: 25-40.
- Franc, A. (1952): Le répartition du zooplancton dans les eaux de Dinard. - Bull. Lab. Mar. Dinard, 37: 37-47.
- Franganillo, P. (1918): Aracnidos nuevos a hallados por primera vez en España. - Bol. Soc. Ent. España, Zaragoza, 1: 123.
- Fredj, G. (1964): Contribution à l'étude bionomique de la Méditerranée occidentale...2. La région de Saint-Tropez... - Bull. Inst. Océanogr. Monaco, 63 (1311): 1-55.
- Frey, H. & R. Leuckart (1847): Pycnogonida: 164-165. Verzeichnis der zur Fauna Helgolands gehörenden wirbellosen Seethiere. In: Beiträge zur Kenntniss wirbelloser Thiere: 136-168; Braunschweig.
- Frith, D.W. (1976): Animals associated with sponges at North Hayling, Hampshire. - Zool. J. Linn. Soc. Lond, 58: 353-362.

- Fry, W.G. (1962): Relative growth of Antarctic Pycnogonida. M.Sc. Thesis. University of the Pacific.
- Fry, W.G. (1964): The Pycnogonida and the coding of biological information for numerical taxonomy. - *Syst. Zool.*, 13 (1): 32-41.
- Fry, W.G. (1964): The pycnogonid fauna of the Antarctic continental shelf. In: *Biologie Antarctique. Actualités Scientifiques et Industrielles*, 1312: 263-269; Paris: Herman.
- Fry, W.G. (1965): The feeding mechanisms and preferred foods of three species of Pycnogonida. - *Bull. Brit. Mus. (Nat. Hist.) (Zool.)*, 12 (6): 195-233, 5 pls.
- Fry, W.G. & J.W. Hedgpeth (1969): Pycnogonida, 1. Colossendeidae, Pycnogonidae, Endeidae, Ammotheidae. Fauna of the Ross Sea, 7. - *Mem. N.Z. Oceanogr. Inst.*, 49: 1-139.
- Fry, W.G. (1975): Review of British Sea Spiders. - *J. nat. Hist.*, 9 (6): 715.
- Fry, W.G. (1975): Pycnogonida: 539. In: Raft fouling in the Menai Strait, 1963-1971. - *Hydrobiologia*, 47 (3-4): 527-558.
- Fry, W.G. (1978): Sea spiders (Pycnogonida). - *Zool. J. Linn. Soc. Lond.*, 63 (1-2): i-x, 1-238.
- Fry, W.G. (1978): A classification within the pycnogonids. In: Sea Spiders (Pycnogonida). - *Zool. J. Linn. Soc. Lond.*, 63 (1-2): 35-58.
- Fry, W.G. & J.H. Stock (1978): A pycnogonid bibliography. In: Sea spiders (Pycnogonida). - *Zool. J. Linn. Soc. Lond.*, 63 (1-2): 197-238.
- Fry, W.G. (1980): Pycnogonids in NE Atlantic food webs: The uneaten sea spider? - *NERC Newsj.*, Nat. Environm. Res. Council, 2 (9): 9-10.
- Fukui, T. (1919): The structure of the Pycnogonida and their species hitherto known from Japan. - *Rigakukai ?, 16 (2)*: 95-99. [in Japanese]
- Fuse, S., T. Nishikawa, N. abe, Y. Fukui, T. Yamamoto & R. Yamanishi: Report on a benthic survey of the inter- and subtidal rocky zone off Nada, in the Middle of Wakayama Prefecture, Japan, 3. - *Nanki Seibutsu*, 22 (2): 80-87.
- Gadeau de Kerville, H. (1885): Pycnogonides. In: G. Lennier: *L'Estuaire de la Saeine*, 2: 185.

- Gadeau de Kerville, H. (1897): Recherches sur les faunes marine et maritime de la Normandie, 2ème voyage... - Bull. Soc. Amis Sci. Nat., Rouen, 1897: 311-447.
- Gadeau de Kerville, H. (1898): Pycnogonides, in: Recherches sur les faunes marines et maritimes de la Normandie: 353-354; Paris, J.-B. Baillièvre & Fils.
- Gadeau de Kerville, H. (1898): Supplément au compte-rendu de mon voyage dans la région de Granville et aux îles Chausey (Manche): 435-443. In: Gadeau de Kerville (1898).
- Gadeau de Kerville, H. (1901): Pycnogonides. In: Recherches sur les faunes marines et maritimes de la Normandie, 3ème voyage: 190; Paris, J.B. Baillièvre & Fils.
- Garneri, A.G. (1911): Contribuzione alle Fauna Sarda. Aracnidi. - Boll. Soc. Zool. Ital., Roma (2) 3 (4-6): 57-103.
- Gaubert, P. (1892): Autotomie chez les Pycnogonides. - Bull. Soc. Zool. France, 17: 224.
- Gegenbauer, C. (1854): Zur Lehre vom Generationswechsel und der Fortpflanzung bei Medusen und Polypen: 1-64, 2 pls.; Würzburg.
- Gegenbauer, C. (1870): Pantopoda: In Grundzüge der vergleichenden Anatomie, 2nd ed., p. 349.
- Gerstaecker, C.E.A. (1863): Pantopoda. In: Carus & Gerstaecker: Handbuch der Zoologie, 2: 248-350.
- Giltay, L. (1928): Note sur les Pycnogonides de la Belgique. - Bull. & Ann. Soc. Roy. Ent. Belg., 68 (9-10): 193-229.
- Giltay, L. (1929): Quelques Pycnogonides des environs de Banyuls (France). - Bull. & Ann. Soc. Ent. Belg., 69: 172-176.
- Giltay, L. (1934): Pycnogonida from the coast of British Columbia. - Can. Field Nat., 48: 49-50.
- Giltay, L. (1934): Remarques sur le genre *Ammothea* Leach et description d'une espèce nouvelle, de la mer d'Irlande. - Bull. Mus. Roy. Hist. Nat. Belg., 10 (18): 1-6.
- Giltay, L. (1934): Pycnogonides. Résultats de voyage de la Belgica en 1897-99. Rapp. Sci. Rés. Voyage Belgica 1897-99, Zool.: 1-16.
- Giltay, L. (1934): Notes sur quelques Pycnogonides de Villefranche-sur-Mer (Alpes-Maritimes). - Bull. Mus. Roy. Hist. Nat. Belg., 10 (35): 1-5.

Giltay, L. (1934): A new Pycnogonid from Bermuda. - Bull. Mus. Roy. Hist. nat. Belg., 10 (42): 1-3.

Giltay, L. (1937): V. Pycnogonida. Résultats scientifiques des croisières du navire-école Belge "Mercator", 1. - Mém. Mus. Roy. Hist. Nat. Belg., (2) 9: 83-89.

Giltay, L. (1942): New records of Pycnogonida from the Canadian Atlantic coast. - J. Fish. Res. Board Can., 5 (5): 459-460.

Glynne-Williams, J. & J. Hobart (1952): Studies on the crevice fauna of a selected shore in Anglesey. - Proc. Zool. Soc. Lond, 122 (3): 797-824.

Gnanamuthu, C.P. (1950): Notes on the morphology and development of a pycnogonid, *Propallene kempfi* (Calman). - Proc. Zool. Soc. Bengal, 3 (1): 39-47.

Gonzalez, S.A. & M.E. Edding (1990): *Achelia assimilis* (Haskell, 1884) in the *Heterozostera* bed of Puerto Aldea, Coquimbo: first record from the northern chilean coast (Pycnogonida: Ammotheidae). - Proc. biol. Soc. Wash., 103 (1): 151-156.

Goodsir, H. (1842): Descriptions of some new species of Pycnogonidae. - Edinb. New Phil. J., 33: 136-139, 1 pl.

Goodsir, H. (1842): Descriptions of some new crustaceous animals found in the Firth of Forth. - Edinb. New Philos. J., 33: 363-368, 1 pl.

Goodsir, H. (1844): On the specific and generic characters of the araneiform Crustacea. - Ann. Mag. Nat. Hist., 14: 1-3, 1 pl.

Goodsir, H. (1844): Description of a new species of *Nymphon*. - Ann. Mag. Nat. Hist., 15: 293.

Gorbunov, G. (1946): Bottom life of the New Siberian shoalwaters and the central part of the Arctic Ocean. Rep. Drift. Exped. Icebreaker Sedov, 1937-40, 3: 30-138. [in Russian]

Gordon, D.P. & W.J. Ballantine (1976): Cape Rodnes to Okokirt Point Marine Reserve: Review of knowledge and bibliography to December, 1976. - Tane, suppl., 22: 1-146.

Gordon, I. (1932): Re-description of some type-specimens of Pycnogonida of the genus *Nymphon*. - Ann. Mag. Nat. Hist., (10), 9: 97-120.

- Gordon, I. (1932): Redescription of two species of Pycnogonida of the genus *Tanystylum*. - Ann. Mag. Nat. Hist., (10), 10: 87-93.
- Gordon, I. (1932): Pycnogonida. - Discovery Rep., 6: 1-138.
- Gordon, I. (1933): A dodecapodous Pycnogonid. - Proc. Linn. Soc. Lond., 1932-33 (2): 91-93.
- Gordon, I. (1938): Pycnogonida. Sci. Rep. Australas. Antarct. Exped. (C) (Zool. & Bot.), 2 (8): 1-40.
- Gordon, I. (1944): Pycnogonida. Rep. Brit., Austral., N.Z. Antarct. Res. Exped., (B) 5(1): 1-72.
- Gosse, P.H. (1855): Pycnogonida. In: A manual of Marine Zoology for the British Isles: pp. 118-120; 189-193; London.
- Gottlieb, E. (1959): Studies of the benthos in Haifa Bay - Ecology and Zoogeography of Invertebrates. Thesis. Hebrew Univ. Jerusalem: i-iii, 1-131.
- Gould, A.A. (1844): Description of a new species of crustacean of the genus *Pasithoe* (Goodsir). - Proc. Soc. Nat. Hist. Boston ?, 1: 92-93.
- Gourret, P. (1896): Documents zoologiques sur l'Etang de Thau. - Trav. Inst. Zool. Univ. Montpellier et Stat. mar. Cette, (N.S.), 5: 1-55.
- Grassle, J.F., et al. (1975): Pattern and zonation: a study of the bathyal megafauna using the research submersible Alvin. - Deep-Sea Res., 22 (7): 457-481.
- Green, C.R. (1981): A variation of the smooth septate junction in sea spiders *Ammothea clausi* (Pycnogonida). - Tissue Cell, 13: 189-195. [cytology]
- Grieg, J.A. (1888): Undersøgelser over dyrelivet i de vestlandske fjorde, II. - Bergens Mus. Aarnsberetning, 1888: 1-11.
- Grieg, J.A. (1907): Invertébrés du fond. In: Duc d'Orleans (ed.): Croisière océanographique dans la Mer du Grönland en 1905, 2: 502-568; Bruxelles, Impr. Sci.
- Grieg, J.A. (1913): Bidrag til kundkaperen om Hardanger-fjordens fauna. - Bergens Mus. Arbok, 1913 (1): 1-147.
- Grieg, J.A. (1914): Evertebratenfaunaen paa havdybet uenfor "Tampen". - Bergens Mus. Aarbok 1914, 3: 1-26.

Grieg, J.A. (1924): Evertebrater fra bankerne ved Spitsbergen
indsamlet av M.K. "Blaafjeld" og M.K. "Tovik" somrene 1923
og 1924. Bergens Mus. Aarbook 1923-1923, 9: 1-33

Grieg, J.A. (1924): Evertebrater fra bankerne ved Spitsbergen
indsamlet av "Torvik" og "Armauer Hansen" somrene 1924 og
1926. - Bergens Mus. Aarbook 1926, 5: 1-28.

Grindley, J.R. & B.F. Kensley (1966): Benthonic Marine Fauna
obtained off the Orange River Mouth by the Diamond Dredge
Emerson K.- Cimbebasia, 16: 3-14.

Grube, E. (1869): Ueber einige vom Zoologischen Museum der
Universität erworbene Pantopoden (Pycnogoniden). -
Jahresber. Schles. Ges. Vaterl. Cult., Breslau, 1868: 54-
55.

Grube, E. (1869): Beschreibungen einiger Pycnogoniden und
Crustaceen. - Abh. Schles. Ges. Vaterl. Cult., Breslau,
1869: 115-119.

Grube, E. (1869): Mittheilungen über St. Vaast-la-Houge und
seine Meeres-, besonders Annelidenfauna. - Abh. Schles.
Ges. Vaterl. Cult., Breslau, 1868/1869: 91-129.

Grube, E. (1871): Ueber zwei neue Heteronereisformen und
Pycnogoniden. - Jb. Schles. Ges. Vaterl. Cult., Breslau,
1870 (48): 84-86.

Grube, E. (1872): Mittheilungen über St. Malo und Roscoff und
die dortige Meeres, und besonders Annelidenfauna. - Abh.
Schles. Ges. Vaterl. Cult., Breslau, 1869-1872: 75-176.

Gruet, Y. (1970): Faune associée des "recifs" édifiés par
l'Annélide *Sabellaria alveolata* (Linné) en baie du Mont
Saint Michel: banc des Hermelles. - Mém. Soc. Natn. Sci.
Nat. Math., Cherbourg, 54: 73-93.

Gruet, Y. (1972): Morphologie, croissance et faune associé des
récifs de *Sabellaria alveolata* (Linné) de la Bernerie-en-
Retz (Liure Atlantique). - Téthys, 3 (2): 321-380.

Gruet, Y. (1972): Fauna associée de "récifs" d'Hermelles... -
Bull. Soc. sci. Bretagne, 47: 69-80.

Grygier, M.J. (1984): Ascothoracida (Crustacea: Maxillopoda)
parasitic on *Chrysogorgia* (Gorgonacea) in the Pacific and
Western Atlantic. - Bull. Mar. Sci., 34 (1): 141-169.

Guerin-Méneville, F.E. (1844): Arachnides. Iconographie du
Régne Animal de G. Cuvier, 3: 12, 2 pls.

Guille, A. (1964): Contribution à l'étude de la systématique et d'écologie d'*Ophiothrix quinquemaculata* d.Ch. - Vie Milieu, 15 (2): 243-308.

Guille, A. & J. Soyer (1968): Nouvelle signalisation du genre *Nymphonella* Ohshima à Banyuls-sur-Mer: *Nymphonella lecalvezii* n.sp. - Vie Milieu (A), 18(2A): 345-353.

Guille, A. (1973): Bionomie benthique du plateau continental de la côte catalane française. VI. Données autécologiques. - Vie Milieu (B), 22(3B): 496-527.

Gunther, K. (1956): Systematik und Stammesgeschichte der Tiere 1939-1953. - Fortschr. Zool., new ser., 10: 32-278. [Pantopoda, p. 146]

Gutierrez, M., J.C. Iturroundobeitia, J.C. Arresti & A. Rallo (1989): Crustaceos y picnogonidos en el abra de Bilbao (Golfo de Vizcaya, Atlantico Nororiental). - Cuad. Inv. Biol., 14: 135-165.

Haddad, M.A., K.N. Suzuki & J.M. Durigan (1987): Ocorrencia e distribucao de Pycnogonida (Arthropoda) em costões rochosos do litoral do Paraná (Abstract). In: VI Mini-Simp. Biol. Mar., 30 Oct.-2 Nov. 1987. Brazil: SMO Sebastiao.

Haeckel, E. (1866): Pantopoda. In: Generelle Morphologie der Organismen, 2: XIV.

Haeckel, E. (1896): Systematische Phylogenie der wirbellosen Thiere, 2: 680pp.

Hagerman, L. (1966): The macro- and microfauna associated with *Fucus serratus* L., with some ecological remarks. - Ophelia, 3: 1-43.

Hagerman, L. (1969): Pycnogonida. In: Marina djur 1. (Fätf fauna 1). In: Natur og Kultur: 27-38; V. Pettersons, Bokindustri AB.

Haig, J. (1978): A catalog of arthropod types in the Allan Hancock Foundation, part I: Non-Crustacean Arthropoda. - Tech. Rep. Allan Hancock Found., 1: 1-4.

Halhed, W.B. (1886): Report on the Pycnogonida of the L.M.B.C. District. - Ann. Rep. Fauna Liverpool Bay and Neighbouring Seas, 1 (1): 227-231.

Halhed, W.B. (1896): List of the Pycnogonida. Report of the Marine Zoology, Botany and Geology of the Irish Sea. - Rep. Brit. Asss. Adv. Sci., 1896: 442.

Hall, H.V.M. (1912): Studies on Pycnogonida. 1. Ann. Rep. Laguna Mar. Lab., 1: 91-99.

Hall, H.V.M. (1913): Pycnogonida from the coast of California with descriptions of two new species. - Univ. Calif. Publ. Zool., 11 (6): 127-142.

Hallez, P. (1899): Le laboratoire de zoologie du Portel et les campagnes de dragages du Berœ. Ouvrage offert par la Ville de Boulogne-sur-Mer aux membres du XXVIIIe Congr. Ass. française pour l'Avancement des Sciences, tenu en cette ville en 1899: 1-28.

Hallez, P. (1905): Notes faunistiques. - Arch. Zool. Exp. Gén., (4) 3 (Notes & Rev., 3): xlvii-lii.

Hallez, P. (1905): Observations sur le parasitisme des larves de *Phoxichilidium* chez *Bougainvillea*. - Arch. Zool. Exp. Gén., (4) 3: 133-144. [hydrroid host]

Hamond, R. (1963): A preliminary report on the marine fauna of the North Norfolk coast. - Trans. Norfolk & Norwich Naturalists Soc., 20 (1): 1-31.

Hamond, R. (1973): The Chelicerate marine arthropodss (Acarina and Pycnogonida) of Norfolk. - Trans. Norfolk Norwich Naturalists' Soc., 22 (6): 424-429.

Hansen, H.J. (1884): Fortegnelse over hidtil i danske Have fundne Pycnogonider eller Søspindler. - Naturhist. Tidsskr., Kjøbenhavn, (3), 14: 647-652. ?

Hansen, H.J. (1884): Spindeldyr, Søspindler (Pantopoder eller Pycnogonidae). In: Zoologia Danica Afbildninger af danske Dyr med pop. Text, 4: 117-131; Copenhagen.

Hansen, H.J. (1886): Vorläufige Mitteilung über Pycnogoniden und Crustaceen aus dem nördlichen Eismeer, von der Dijmphna-Expedition mitgebracht. - Zool. Anz., 9: 638-643.

Hansen, H.J. (1887): Kara Havets Pycnogonider. In: Dijmphna-Togets zoologisk-botaniske Udbytte: 157-181; Kjøbenhavn, Bianco Lunos.

Hansen, H.J. (1889): Liste over Crustacea og Pycnogonida. In: T Holm, Beretning om de paa "Fylla" s Togt i 1884 foretagne zoolog. Undersøgelser i Grønland. - Medd. Grønland, 8: 166.

Hansen, H.J. (1895): Pycnogonider og Malakostrake Krebsdyr. - Medd. Grønland, 19: 123-125.

Hansen, H.J. & W. Sørensen (1898): The order Palpigradi Thor. and its relationship to the other Arachnida. - Ent. Tidsskr., 18 (3-4): 223-240.

Hansen, H.J. (1930): Studies on Arthropoda, 3: 333-335; Copenhagen.

Hanström, B. (1919): Zur Kenntnis des zentralen Nervensystems der Arachnoiden und Pantopoden nebst Schlussfolgerungen betreffs der Phylogenie der genannten Gruppen. - Diss., Univ. Lund: 1-191. [nervous system]

Hanström, B. (1926): Eine genetische Studie über die Augen und Sehzentren von Turbellarien, Anneliden und Arthropoden. - Kung. Svensk. Vetensk. Akad. Handl., Stockholm, (3) 4. [genetics]

Hanström, B. (1927): Neue Beobachtungen über Augen und Sehzentren von Entomostraken, Schizopoden und Pantopoden. - Zool. Anz., 70: 236-251. [optical sense]

Hanström, B. (1965): Indications of neurosecretion and the structure of Sokolow's organ in pycnogonids. - Sarsia, 18: 24-36. [gland, nervous system]

Harada, E. (1991): Inventory of zoological type specimens in the museum of the Seto Marine Biological Laboratory. - Publ. Seto Mar. Biol. Lab., 35 (1-3): 171-233.

Harms, J. (1993): Checklist of species (algae, invertebrates and vertebrates found in the vicinity of the island Helgoland (North Sea, German Bight) - a review of recent records. - Helgoländer Meeresunters., 47: 1-34.

Harrison, F.W., D.R. Calder, B.C. Coull & F.C. James (1979): Status Report: Lower invertebrates and miscellaneous phyla. Proc. 12st S-Carolina endangered species Symp 1976: 34-40.

Hartlaub, C. (1900): Zoologische Ergebnisse einer Untersuchungsfahrt des deutschen Seefischerei-Vereins nach der Bäreninsel und Westspitzbergen, ausgeführt im Sommer 1898 auf S.M.S. "Olga". Kommission zur wissenschaftlichen Untersuchung der deutschen Meere in Kiel. - Wiss. Meeresunters., new ser., 4: 171-193.

Hartley, J.P. (1986): The macrobenthos of the Beatrice Oilfield, Moray Firth, Scotland. - Proc. Zool. Soc. Edinburgh, 8 (91): 221-245.

Haswell, W.A. (1885): Pycnogonida of the Australian coast with description of new species. - Proc. Linn. Soc. N.S. Wales, 9: 1021-1033.

Hedgpeth, J.W. (1939): Some pycnogonids found off the coast of Southern California. - Am. Midl. Nat., 22 (2): 458-465, 2 pls.

Hedgpeth, J.W. (1940): A new pycnogonid from Pescadero, Calif., and distributional notes on other species. - J. Wash. Acad. Sci., 30 (2): 84-87.

Hedgpeth, J.W. (1941): On a species of *Nymphon* from the waters of Southern California. - Am. Midl. Nat., 25 (2): 447-449, 1 pl.

Hedgpeth, J.W. (1941): A key to the Pycnogonida of the Pacific coast of North America. - Trans. San Diego Soc. Nat. Hist., 9 (26): 253-264.

Hedgpeth, J.W. (1941): A key to the Pycnogonids of the Pacific coast of North America (especially of central California). In: S.F. Light (ed.), Laboratory and Field Text in Invertebrate Zoology: 119-124; Berkeley & Los Angeles, Univ. Calif. Press.

Hedgpeth, J.W. (1942): Sea spiders. - Nature Mag.: 413-414.

Hedgpeth, J.W. (1943): Pycnogonids of the Bartlett collections. - J. Wash. Acad. Sci., 33 (3): 83-90.

Hedgpeth, J.W. (1943): Pycnogonida from the West Indies and South America collected by the Atlantis and earlier expeditions. - Proc. New England Zool. Club, 22: 41-58.

Hedgpeth, J.W. (1943): On a species of pycnogonid from the North Pacific. - J. Wash. Acad. Sci., 33 (7): 223-224.

Hedgpeth, J.W. (1944): On a new species of *Pallenopsis* (Pycnogonida) from Western Australia. - Proc. New England Zool. Club, 23: 55-58.

Hedgpeth, J.W. (1947): Pycnogonida. Encyclopedia Britannica: 2pp.

Hedgpeth, J.W. (1947): On the evolutionary significance of the Pycnogonida. - Smiths. Misc. Coll., 106 (18): 1-54.

Hedgpeth, J.W. (1948): The Pycnogonida of the Western North Atlantic and the Caribbean. - Proc. U.S. natn. Mus., 97 (3216): 157-342.

Hedgpeth, J.W. (1949): Report on the Pycnogonida collected by the Albatross in Japanese waters in 1900 and 1906. - Proc. U.S. natn. Mus. 98 (3231): 233-321.

Hedgpeth, J.W. (1950): Pycnogonida of the United States Navy Expedition, 1947-48. - Proc. U.S. Natn. Mus., 100 (3260): 147-160.

Hedgpeth, J.W. (1951): Pycnogonids from Dillon Beach and vicinity, California, with descriptions of two new species. - Wasman J. Biol., 9 (1): 105-117, 3 pls.

Hedgpeth, J.W. (1952): Class (or Subphylum) Pycnogonida. Sea Spiders. In: E.F. Ricketts & J. Calvin (eds.). Between Pacific Tides, 3rd ed.: 430; Stanford Univ. Press.

Hedgpeth, J.W. (1953): Class Pycnogonida. In: R.I. Smith et al. (eds.). Intertidal invertebrates of the Central California coast: 201-210; Univ. Calif. Press.

Hedgpeth, J.W. (1954): Pycnogonida. In: P.S. Galtsoff (ed.). The Gulf of Mexico, its origin, waters and marine life. - Fish. Bull. Fish & Wildl. Serv. U.S., 89 (55): 425-427.

Hedgpeth, J.W. (1954): On the phylogeny of the Pycnogonida. - Acta Zool., Stockholm, 35: 193-213.

Hedgpeth, J.W. (1954): 1. The Pycnogonida. Reports on the dredging results of the Scripps Institution of Oceanography Trans-Pacific Expedition. July-December, 1953. - Syst. Zool., 3 (4): 147.

Hedgpeth, J.W. (1956): On the phylogeny of the Pycnogonida. - Proc. XIVth Int. Congr. Zool., Copenhagen, 1953, sec. XV: 506-507.

Hedgpeth, J.W. (1957): Miscellaneous arthropods - annotated bibliography. In: J.W. Hedgpeth (ed.). Treatise on marine ecology and paleoecology. 1. Ecology: 1175-1176; Geol. Soc. Am., Mem., 67; Boulder, Geol. Soc. Am.

Hedgpeth, J.W. (1959): Pycnogonida. In: Encyclopaedia Britannica: 2pp.

Hedgpeth, J.W. (1961): Pycnogonida. In: Encyclopedia of Science and Technology: 105-106; New York: McGraw Hill.

Hedgpeth, J.W. (1961): Pycnogonida. In: The Encyclopedia of the Biological Sciences: 851-852. New York: Van Nostrand Reinhold.

Hedgpeth, J.W. (1961): Pycnogonida. Rep. Lund Univ. Chile Exped. 1948-49. - Lunds Univ. Arsskr., N.F. (2): 1-18.

Hedgpeth, J.W. (1962): A bathypelagic pycnogonid. - Deep-Sea Res., 9: 487-491.

- Hedgpeth, J.W. (1962): Taxonomy: Man's oldest profession. 11th Ann. Univ. Pac. Fac. Res. Lect., May 22, 1961: i, 18pp.
- Hedgpeth, J.W. (1962): Pycnogonida. In: Introduction to seashore life of the San Francisco Bay Region and the coast of Northern California: 74-75; Univ. Calif. Press.
- Hedgpeth, J.W. (1963): Pycnogonida of the North American Arctic. - J. Fish. Res. Board Can., 20 (5): 1315-1348.
- Hedgpeth, J.W. (1963): A comparative study of Arctic and Antarctic Pycnogonida (Abstract). Proc. XVI Int. Congr. Zool., Washington, 1: 107.
- Hedgpeth, J.W. (1963): Arctic and Antarctic Pycnogonida: A comparison (Abstract.). - Bull. Ecol. Soc. Amer., 44 (2): 40.
- Hedgpeth, J.W. (1964): Notes on the peculiar egg laying habit of an antarctic prosobranch (Mollusca: Gastropoda). - The Veliger, 7 (1): 45-46.
- Hedgpeth, J.W. & W.G. Fry (1964): Another dodecopodous pycnogonid. - Ann. Mag. Nat. Hist., (13) 7: 161-169.
- Hedgpeth, J.W. (1967): Review of Basic Arthropodan Stock. - Quart. Review Biol., 42: 69.
- Hedgpeth, J.W. (1968): Pycnogonid studies. - Antarct. J. U.S., 3 (4): 129-130.
- Hedgpeth, J.W. (1968): Pycnogonids. In: E.F. Ricketts & J. Calvin. Between Pacific Tides (4th ed.): 78, 102-104, 157, 170, 186, 188, 202, 316-317, 349, 351-352, 366-368, 477-479; Stanford Univ. Press.
- Hedgpeth, J.W. (1969): Introduction to Antarctic Zoogeography. - Antarct. Map Folio Ser. Am. Geogr. Soc., Folio 11: 1-9.
- Hedgpeth, J.W. (1969): Pycnogonida. Antarct. Map Folio Ser. Am. Geogr. Soc., Folio 11: 26-28, pls. 13-14.
- Hedgpeth, J.W. (1969): Pycnogonida. In: P. Gray (ed.). Encyclopedia of the Biological Sciences; pp. 780-781; New York, Van Nostrand Reinhold.
- Hedgpeth, J.W. (1969): An intertidal reconnaissance of rocky shores of the Galápagos. - Wasman J. Biol., 27 (1): 1-24.
- Hedgpeth, J.W. (1969): Preliminary observations of life between tidemarks at Palmer Station, 64°45'S, 64°05'W. - Antarct. J. U.S., 4 (4): 106-107.

Hedgpeth, J.W. (1970): Marine biogeography of the Antarctic regions. In: M.W. Holdgate (ed.). Antarctic Ecology: 97-104. Sci. Comm. Antarct. Res.; London & New York, Academic Press.

Hedgpeth, J.W. (1971): Subphylum Pycnogonida, Class Pantopoda. Sea Spiders. In: K.L. Gosner. Guide to identification of marine and estuarine invertebrates - Cape Hatteras to the Bay of Fundy: 400-402; New York, Wiley Interscience.

Hedgpeth, J.W. (1971): James Eights of the Antarctic. In: Research in the Antarctic. Am. Ass. Adv. Sci.

Hedgpeth, J.W. (1971): Perspective of benthic ecology in Antarctica. In: Research in the Antarctic: 93-136. Am. Ass. Adv. Sci.

Hedgpeth, J.W. & J.C. McCain (1971): A review of the pycnogonid genus *Pantopipetta* (family Austrodecidae, emend.), with the description of a new species. In: G. Llano & I.E. Wallen (eds.). Biology of the Antarctic Seas. - Antarct. Res. Ser., 17: 217-229.

Hedgpeth, J.W. (1974): One hundred years of Pacific Oceanography. In: The biology of the Oceanic Pacific: 137-155; Oregon State Univ. Press.

Hedgpeth, J.W. (1974): Phylum Arthropoda. Subphylum Chelicerata. Class Pycnogonida. In: E.N. Kozloff (ed.). Keys to the marine invertebrates of the Puget Sound, the San Juan Archipelago, and adjacent regions: 125-128; Seattle, Univ. Wash. Press.

Hedgpeth, J.W. (1974): Review of Pycnogonids. - Science, 184: 150.

Hedgpeth, J.W. (1975): Pycnogonida. In: S.F. Light. Intertidal invertebrates of the Central California Coast; Univ. Calif. Press.

Hedgpeth, J.W. (1975): Review of Pycnogonids and of British Sea Spiders. - Quart. Review Biol., 50 (3): 330-331.

Hedgpeth, J.W. (1977): Pycnogonida. In: McGraw Hill Encyclopedia of Science Technology, 11: 110-112.

Hedgpeth, J.W. & E.C. Haderlie (1978 ?): Pycnogonids. In: R. Morris, D.P. Abbott & E.C. Haderlie. Invertebrates of California Shores (?); Stanford Univ. Press.

Hedgpeth, J.W. (1979 ?): At sea with provinces and plates. In: Biogeography. Biol. Colloq. Oregon State Univ. Press.

- Hedgpeth, J.W. (1982): Pycnogonida. In: S.P. Parker. *Synopsis and classification of living organisms*, 2: 169-173. New York: McGraw-Hill.
- Heezen, B.C. & C.D. Hollister (1971): Pycnogonida. In: *The Face of the Deep*: 8, 28, 92-94, 98-100, 121; New York, Oxford Univ. Press.
- Hegner, R.W. & K.A. Stiles (1959): Subclass Pycnogonida. In: *College Zoology*: 268, 279; New York, Macmillan.
- Helfer, H. (1909): Biologisch-faunistische Beobachtungen an Pantopoden der Nord- und Ostsee. Diss. Kgl. Christian-Albrechts Univ. Kiel: 1-49.
- Helfer, H. (1915): Pantopoda. In: C.A. Apstein (ed.). *Nomina conservanda*. - Sitz.-Ber. Ges. naturforsch. Fr. Berlin, 5: 147.
- Helfer, H. (1932): Pantopoda. In: Kükenthal-Krumbach, *Handbuch der Zoologie*, 3 (2) 4: 1-66.
- Helfer, H. (1932): Das Pantopoden-Schrifttum. - Sitz.-Ber. Ges. naturforsch. Fr. Berlin: 235-254.
- Helfer, H. & E. Schlottke (1935): Pantopoda. Dr. H.G. Bronn's Klassen und Ordnungen des Tierreichs, 5 (4) (2): 1-314.
- Helfer, H. (1936): The fishery grounds near Alexandria (Egypt). VIII. Pantopoda. Ministry of Commerce & Industry, Egypt, Notes & Mem., 16: 1-6.
- Helfer, H. (1936): Pantopoda. Die Tierwelt der Nord- und Ostsee, 31 (XIa3): 1-5.
- Helfer, H. (1938): Einige neue Pantopoden aus der Sammlung des Zoologischen Museums in Berlin. - Sitz.-Ber. Ges. naturforsch. Fr. Berlin, 1937 (2): 162-185.
- Heller, C. (1875): Neue Crustaceen und Pycnogoniden. Gesammelt während der K.K. österr.-ungar. Nordpol-Expedition. Vorläufige Mitteilung. - Sitz.-Ber. math.-naturw. Cl. K.K. Akad. Wiss., Wien, 71 (1): 609-612.
- Heller, C. (1875): Die Crustaceen, Pycnogoniden und Tunicaten der K.K. österr.-ungar. Nordpol-Expedition. - Denkschr. Kaiserl. Akad. Wiss. (math-naturwiss. Cl.), Wien, 35: 25-46.
- Henderson, J.R. (1885): Recent additions to the invertebrate fauna of the Firth of Forth. - Proc. Roy Soc. Edinb., 8: 307-313.

Henry, L.M. (1953): The nervous system of Pycnogonida. - Microent., 18 (1): 16-36.

Herdman, W.A. (1892): Notes on the collections made during the cruise of the S.Y. "Argo" up the West Coast of Norway in July, 1891. - Trans Liverpool Biol. Soc., 6: 70-93, pls. V-VI.

Herdman, W.A. (1896): List of Pycnogonida. In: The marine zoology, botany and geology of the Irish Sea. Fourth and final report of the Committee... Rep. 66th Meeting Brit. Ass. Adv. Sci.: 442.

Herdman, W.A. (1896): Pycnogonida. In: 9th Ann. Rep. Liverpool Mar. Biol. Committee: 19, 21, 24.

Herdman, W.A., Thompson, I.C. & H.C. Chadwick (1900): Fauna distribution lists for the south end of the Isle of Man. 14th Rep. Mar. Biol. Stat. Port Erin: 36.

Herdman, W.A. (1901): Fourteenth annual report of the Liverpool Marine Biological Committee and their Biological Station at Port Erin. - Proc. & Trans. Liverpool Biol. Soc., 15 (1900-1901): 19-84.

Herdman, W.A. (1903): The Marine Biological Station at Port Erin. 17th Ann. Rep. Liverpool Mar. Biol. Committee: 5-32.

Herdman, W.A. (1897): The Tenth Annual Report of the Liverpool Marine Biology Committee, and their biological station at Port Erin. - Trans. Liverpool Biol. Soc., 11: 3-47.

Herring, P.J. (1983): The spectral characteristics of luminous marine organisms. - Proc. Roy. Soc. London, ser. B, 220: 183-217. [bioluminescence]

Hesse, M. (1867): Observations sur des Crustacés rares ou nouveaux des côtes de France. - Ann. Sci. Nat., Paris, (5) 7: 199-216.

Hesse, M. (1974): Mémoire sur les Crustacés rares ou nouveaux. - Ann. Sci. Nat., Paris, (5): 20: 1-18, 1 pl.

Hicks, G.R.F., M.J. Huaki, W.R. Weber & J.C. Yaldwyn (1991): Inventory of cnidarian, pycnogonid and crustacean type specimens in the National Museum of New Zealand. - Natn. Mus. N.Z., misc. ser., 22: 1-23.

Highnam, K.C. (1981): A survey of invertebrate metamorphosis. In: L.I. Gilbert & E. Frieden (eds.). Metamorphosis: A problem in developmental biology, 2nd ed., 2: 43-73. New York & London: Plenum Press.

Hilton, W.A. (1915): The central nervous system of the pycnogonid *Lecythorhynchus*. - J. Ent. Zool. Pomona Coll., 6: 134-136.

Hilton, W.A. (1915): Pycnogonids collected during the summer of 1914 at Laguna Beach. - J. Ent. Zool. Pomona College, 7 (1): 67-70.

Hilton, W.A. (1915): Pycnogonids collected during the summer of 1915 at Laguna Beach. - J. Ent. Zool. Pomona Coll., 7 (3): 201-206.

Hilton, W.A. (1916): A remarkable pycnogonid. - J. Ent. Zool. Pomona Coll., 8 (1): 19-24.

Hilton, W.A. (1916): The life history of *Anoplodactylus erectus* Cole. - J. Ent. Zool. Pomona Coll., 8 (1): 25-34.

Hilton, W.A. (1916): The nervous system of pycnogonids. - J. Comp. Neur., 26 (5): 463-472.

Hilton, W.A. (1920): Notes on Pacific Coast Pycnogonids. - J. Ent. Zool. Pomona Coll., 12 (4): 93.

Hilton, W.A. (1931): Nervous system and sense organs. XXXVI. Pantopoda. - J. Ent. Zool. Pomona Coll., 23 (1): 5-18.

Hilton, W.A. (1934): Notes on parasitic pycnognids. - J. Ent. Zool. Pomona Coll., 26 (4): 57.

Hilton, W.A. (1939): A preliminary list of pycnognids from the shores of California. - J. Ent. Zool. Pomona Coll., 31 (2): 27-35.

Hilton, W.A. (1939): A collection of pycnognids from Santa Cruz Island. - J. Ent. Zool. Pomona Coll., 31 (4): 72-74.

Hilton, W.A. (1942): Pantopoda chiefly from the Pacific. I. Nymphonidae. - J. Ent. Zool. Pomona Coll., 34 (1): 3-7.

Hilton, W.A. (1942): Pycnogonids from the Allan Hancock Expeditions. - Rep. Allan Hancock Exped., 5 (9): 227-339.

Hilton, W.A. (1942): Pantopoda chiefly from the Pacific. II. Callipallenidae. - J. Ent. Zool. Pomona Coll., 34 (2): 38-41.

Hilton, W.A. (1942): Pycnogonids from Hawaii. - Occ. Pap. Bernice P. Bishop Mus., Honolulu, Hawaii, 17 (3): 43-55.

Hilton, W.A. (1942): Pycnogonids from the Pacific. Family Tanystilidae. - J. Ent. Zool. Pomona Coll., 34 (3): 69-70.

- Hilton, W.A. (1942): Pycnogonids from the Pacific. Family Phoxichilididae Sars 1891. - J. Ent. Zool. Pomona Coll., 34 (3): 71-74.
- Hilton, W.A. (1943): Pycnogonids from the Pacific. Family Ammotheidae. - J. Ent. Zool. Pomona Coll., 34 (4): 93-99.
- Hilton, W.A. (1943): Pycnogonids from the Pacific. Family Colossendeidae. - J. Ent. Zool. Pomona Coll., 35 (1): 2-4.
- Hilton, W.A. (1943): Pycnogonids from the Pacific. Family Pycnogonidae. - J. Ent. Zool. Pomona Coll., 35 (2): 19.
- Hodge, G. (1862): Observations on a species of pycnogon (*Phoxichilidium coccineum*). With an attempt to explain the order of its development. - Ann. Mag. Nat. Hist., (3) 9: 33-43.
- Hodge, G. (1863): Descriptions of two new species of Pycnogonidea. - Ann. Mag. Nat. Hist., (3) 11: 463-464.
- Hodge, G. (1863): Report on the Pycnogonidea with descriptions of two new species. In: Report of the Dredging Expedition to the Dogger Bank and the coast of Northumberland. - Trans. Tyneside Naturalists' Field Club, 5: 281.
- Hodge, G. (1864): List of the British Pycnogonidea, with descriptions of several new species. - Ann. Mag. Nat. Hist., (3) 13: 113-117.
- Hodge, G. (1865): Pycnogonidea. Reports of the deep-sea dredging on the coasts of Northumberland and Doggerbank. - Nat. Hist. Trans. Northumberland Durham, 1 (1): 41-42.
- Hodgson, T.V. (1902): Crustacea (Pycnogonida). In: Report on the Collections of Natural History made in the Antarctic Regions during the voyage of the "Southern Cross": 256-258; London.
- Hodgson, T.V. (1904): On a new pycnogonid from the South Polar regions. - Ann. Mag. Nat. Hist., (7) 14: 458-462.
- Hodgson, T.V. (1905): Preliminary report of the biological collections of the Discovery. - Geogr. J., London, 25: 396-400.
- Hodgson, T.V. (1905): Decolopoda and Colossendeis. - Zool. Anz., 29 (8): 254-256.
- Hodgson, T.V. (1905): Scotia collections, VII. On *Decolopoda australis* Eights - an old pycnogonid rediscovered. - Proc. Roy. Phys. Soc. Edinburgh, 16 (1): 35-42.

- Hodgson, T.V. (1907): Pycnogoniden. Ergebnisse der Hamburger Magalhaensischen Sammelreise 1892-93, 2: 1-20.
- Hodgson, T.V. (1907): Pycnogonida. National Antarctic Expedition 1901-1904. Rep. Natn. Antarct. Exped. 1901-1904, Nat. Hist., 3: 1-72, 10 pls.
- Hodgson, T.V. (1908): The Pycnogonida of the Scottish National Antarctic Expedition. - Trans. Roy. Soc. Edinburgh, 46 (1) (6): 159-188.
- Hodgson, T.V. (1910): The Pycnogonida of Devonshire. - Rep. Trans. Devonsh. Ass. Adv. Sci., Lit. & Arts, 42: 425-439.
- Hodgson, T.V. (1910): Pantopoda. In: L. Schultze. Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika ausgeführt in den Jahren 1903-1905 von Dr. Leonhard Schultze, 4 (1): 221-228. - Denkschr. Med.-Naturwiss. Ges. Jena, 16: 219-228.
- Hodgson, T.V. (1914): Preliminary report on the Pycnogonida of the German Southpolar Expedition, 1901-1903. - Zool. Anz., 45 (4): 158-165.
- Hodgson, T.V. (1915): The Pycnogonida collected by the "Gauss" in the Antarctic regions, 1901-3; preliminary report. - Ann. Mag. Nat. Hist., (8) 15 (85): 141-149.
- Hodgson, T.V. (1927): Die Pycnogoniden der Deutschen Südpolar-Expedition 1901-3. - Dtsch. Südpolar-Exped., 19 (Zool. II): 303-358.
- Hoek, P.P.C. (1877): Pycnogoniden en hun plaatss in het system. - Versl. Vergad. Nederl. Dierk. Ver., 1877: 1-4.
- Hoek, P.P.C. (1877): Ueber Pycnogoniden. 1. Die Arten der niederländischen Fauna. - Nederl. Arch. Zool., 3: 235-254.
- Hoek, P.P.C. (1879): Pycnogonida. In: Verslag van de wetenschappelijke vergadering der Nederlandse Dierkundige Vereeniging, etc. - Tijdschr. Nederl. Dierk. Ver., 4: 11-12.
- Hoek, P.P.C. (1881): Report on the Pycnogonida dredged by HMS Challenger 1873-76. - Rep. Sci. Res. Expl. Voyage HMS Challenger, 3 (10): 1-167, 21 pls.
- Hoek, P.P.C. (1881): The pycnogonids, dredged during the cruises of the Willem Barents in the years 1878 and 1879. - Nederl. Arch. Zool., Suppl. 1: 1-28.
- Hoek, P.P.C. (1881): Nouvelles études sur les Pycnogonides. - Arch. Zool. Exp. Gén., 9: 445-542.

- Hoek, P.P.C. (1882): Report on the Pycnogonida. - Proc. Roy. Soc. Edinburgh, 11: 689-690.
- Hoek, P.P.C. (1883): The Pycnogonida dredged in the Faroe Channel during the cruise of HMS Triton in August 1882. - Trans. Roy. Soc. Edinburgh, 32 (1): 1-10, 1 pl.
- Hoek, P.P.C. (1884): Crustacés de l'Escaut de l'Est. - Tijdschr. Dierk. Ver., Suppl. 1: 517-545.
- Hoek, P.P.C. (1898): On four pycnogonids dredged during the cruise of the Challenger (investigated and described after the completion of the report). With an appendix. - Tijdschr. Nederl. Dierk. Ver., (2) 5 (2-4): 290-301.
- Hoenigman, J. & J.H. Stock (1955): Das männliche Geschlecht von *Pycnogonium pusillum* Dohrn, 1881 (Pantopoda). - Ent. Ber., Amsterdam, 15 (24): 534-536.
- Holme, N.A. (1966): The bottom fauna of the English Channel, II. - J. Mar. biol. Ass. U.K., 46: 401-493.
- Hong, J.-S. & I.H. Kim (1987): Korean Pycnogonids chiefly based on the collections of the Korea Ocean Research and Development Institute. - Korean J. Syst. Zool., 3 (2): 137-164.
- Honma, Y. & T. Kitami (1978): Fauna and Flora in the waters adjacent to the Sado Marine Biological Station, Niigata University. - Ann. Rep. Sado Mar. Biol. Stat., Niigata Univ., 8: 7-81.
- Hooper, J.N.A. (1981): Some aspects of the reproductive biology of *Parapallene avida* Stock (Pycnogonida: Callipallenidae) from Northern New South Wales. - Australian Zool., 20 (3): 473-483.
- Hunter, J. & D. Rendall (1986): The sublittoral fauna of the Inverness, Cromarty and Dornoch Firths. - Proc. Zool. Soc. Edinburgh, 8 (91): 263-274.
- Hurst, C.H. (1897): Some specimens of *Nymphon* dredged from the "Turbot hole", near Puffin Island. In: W.A. Herdman, The Tenth Annual Report of the Liverpool Marine Biology Committee and their biological station at Port Erin. - Trans. Biol. Soc. Liverpool, 11: 27.
- Hurst, C.H. (1897): Pycnogonida. In: W.A. Herdman, 10th Ann. Rep. Liverpool Mar. Biol. Committee: 22, 27.
- Hutton, F.W. (1904): In: Index Faunae Novae Zealandiae: 246-247. London: Dulau.

- Huxley, T.H. (1877): Pycnogonida. In: Manual of Anatomy of Invertebrated Animals: 386.
- Ichikawa, S. & M. Yamada (1957): Catalogue of marine invertebrates of the Osyoro Bay area; 10pp. Sapporo. [in Japanese]
- Ihle, J.E.W. (1898): Ueber die Phylogenie und systematische Stellung der Pantopoden. - Biol. Zentralbl., 18: 603-609.
- Inaba, A. (1963): Pycnogonida. In: Fauna and Flora of the Inland Sea of Seto. - Publ. Mukaishima Mar. Biol. Stat.: 205-206. [in Japanese].
- Isaac, M.J. & J.H. Jarvis (1973): Endogenous tidal rhythmicity in the littoral pycnogonid *Nymphon gracile* (Leach). - J. Exp. Mar. Biol. Ecol., 13: 83-90.
- Ives, J.E. (1891): Echinoderms and arthropods from Japan. - Proc. Acad. Nat. Sci. Philadelphia, 1891-1892: 210-221.
- Ives, J.E. (1892): A new species of *Pycnogonum* from California. - Proc. Acad. Nat. Sci. Philadelphia, 1892-1893: 142-144.
- Iwaki, T., K. Hibino & T. Kawahara (1977): Seasonal changes in the initial development of fouling communities in Matoya Bay. - Bull. Fac. Fish., Mie Univ., 4: 11-29.
- Jarvis, J.H. & P.E. King (1972): Reproduction and development in the pycnogonid *Pycnogonum littorale*. - Mar. Biol., 13: 146-155.
- Jarvis, J.H. & P.E. King (1973): Ultrastructure of the photoreceptors in the pycnogonid species *Nymphon gracile* (Leach), and *Pycnogonum littorale* (Ström). - Mar. Behav. Physiol., 2: 1-13. [cytology, optical sense]
- Jarvis, J.H. (1974): Aspects of the reproductive biology of pycnogonids. Ph.D. Tehesis, Univ. Wales. [reproduction]
- Jarvis, J.H. (1975): Egg development and the reproductive cycle in the pycnogonid *Endeis laevis*. - Mar. Biol., 33: 331-339. [reproduction]
- Jarvis, J.H. & P.E. King (1978): Reproductive biology of British pycnogonids (Oogenesis and the reproductive cycle). - Zool. J. Linn. Soc., 63 (1-2): 105-131. [reproduction]
- Jarvis, J.H. & P.E. King (1979): The response of the Pycnogonids *Nymphon gracile* Leach and *Pycnogonum littorale* (Strom) to light stimuli of different wavelength and intensity. - Mar. Behav. Physiol., 6 (4): 307-317. [optical sense]

Jarzynsky, T. (1870): Praemissus catalogus Pycnogonidarum, inventarum in mari Glaciali, ad oras Lapponiae rossicae et in mari Albo, anno 1869 et 1870. - Ann. Soc. Nat. St.-Pétersbourg, 1: 319-320. Die Wirbellosen Thiere des Weissen Meeres: 168-171; Leipzig.

Johnson, M.E. & H.J. Snook (1927): Seashore animals of the Pacific Coast: i-xiv, 1-659.

Johnston, G. (1828): Contributions to the British Fauna. - Zool. J. Zool. Soc. London, 3: 486-491.

Johnston, G. (1833): Illustrations in British Zoology. - Mag. Nat. Hist., 6: 40-43.

Johnston, G. (1837): Miscellanea zoologica, I. An attempt to ascertain the British Pycnogonida. - Mag. Zool. Bot, 1: 371-382; 1 pl.

Jones, D.J. (1971): Ecological studies on macroinvertebrate populations associated with polluted kelp forests in the North Sea. - Helgol. Wiss. Meeresunters., 22: 417-441. [ecology, pollution]

Jones, T.R. (1855): Pycnogonidae. In: General Outline of the Organisation of the Animal Kingdom and Manual of Comparative Anatomy: 437-439; London, Van Voorst.

Jordan, H.E. (1916): The microscopic structure of the leg muscle of the sea-spider, *Anoplodactylus latus*. - Anat. Rec., (10) 7: 493-508. [anatomy, musculature]

Jourdain, S. (1892): Affinités zoologiques des Pantopodes. - C.R. Séanc. Soc. Biol., Paris, (9) 4: 945.

Juan Garcia, J. (1988): Contribucion al estudio de los Picnogónidos de las Baleares. VI Simp. Iberico Estud. Bentos Mar., Palma de Mallorca, 19-22 Sep. 1988: 77.

Just, J. (1970): Marine biological investigations of Jørgen Brønlund Fjord, North Greenland. - Medd. Grønland, 184 (5) ?.

Just, J. (1970): Pycnogonida from Jørgen Brønlund Fjord, North Greenland. - Medd. Grønland, 184 (7): 23-27.

Just, J. (1972): Revision of the genus *Boreonymphon* G.O. Sars (Pycnogonida) with a description of two new species, *B. ossiansarsi* Knaben and *B. compactum* Just. - Sarsia, 49: 1-28.

Kaesler, C.B. (1964): The Mediterranean crevice habitat. - Vie Milieu, 15 (4): 947-977.

- Kaesler, C.B. (1968): Pycnogonida. Sea spiders. In: Invertebrate Zoology, 2: 292-301; New York: John Wiley, Interscience.
- Kawahara, T., T. Iwaki, K. Hibino & Y. Sugimura (1979): Fouling communities in Yokkaichi Harbor. - Publ. Amakusa Mar. Biol. Lab., Kyushu Univ., 5 (1): 19-30, I+II.
- Kawecki, Z. (1978): The contributions of Maria Dohrn b. Baranowska (1856-1918) as a co-founder of the Zoological Station in Naples (on the 60-tieth Anniversary of Her Deadth). - Przeglad Zool., 22 (4): 309-314.
- Keegan, B.F., B.D.S. O'Connor, D. McGarth, G. Konnercker & D.D. Foighil (1987): Littoral and benthic investigations on the south coast of Ireland, II: The macrobenthic fauna off Carnsore Point. - Proc. Roy. Acad., ser. B, 87B (1): 1-14.
- Kenny, R. & N. Haysom (1962): Ecology of rocky shore organisms at Macquarie Island. - Pac. Sci., 16 (3): 245-263.
- Kerneis, A. (1960): Contribution à l' étude faunistique et écologique des herbiers de Posidoniens de la région de Banyuls. - Vie Milieu, 11 (2): 145-187.
- Khayrallah, N. & A.M. Jones (1975): A survey of the benthos of the Tay estuary. - Proc. Roy. Soc. Edinburgh, (B) 75 (7): 113-135.
- Kiaer, H. (1905): Om dyrelivet i Balsfjorden og denne fjords udløb til havet. - Tromsø Mus. Aarsberetning, 28: 13-50.
- Kikuchi, T. (1965): Faunal list of the *Zostera marina* belt in Tomioka Bay, Amakusa, Kyushu. - Publ. Amakusa Mar. Biol. Lab., Kyushu Univ., 1 (2): 163-192.
- Kikuchi, T. Rediscovery of the bivalve-infesting Pycnogonida, *Nymphonella tapetis* Ohshima, 1927 in Amakusa. - Publ. Amakusa Mar. Biol. Lab., Kyushu Univ., 4 (1): 51-56.
- Kikuchi, T. (1985): Check list of the shore fauna in Tsujishima Islet, Amakusa, West Kyushu. - Publ. Amakusa Mar. Biol. Lab., 8 (1): 65-88.
- Kim, H.S. & I.H. Kim (1985): *Achelia crurispinifera*, a new Pycnogonid species from Korean waters. - Kor. J. Zool., 28 (2): 120-124.
- Kim, I.H. (1982): Description of a sea-spider (Pycnogonida), new to Korea. - J. Kangreung Natn. Univ., 4: 275-281. [in Korean with English abstract]

- Kim, I.H. (1983): *Tanystylum ulreungum*, a new pycnogonid species from Korean water. - J. Kangreung Natn. Univ., 5 (2): 467-471.
- Kim, I.H. (1984): Common Pycnogonid species from East Sea and South Sea of Korea. - J. Kangreung Natn. Univ., 7: 531-551.
- Kim, I.I. & J.H. Stock (1984): A new Pycnogonid, *Pycnogonum koreanum*, sp. nov. from the Sea of Japan. - J. nat. Hist., 18: 685-688.
- Kim, I.H. (1986): Four Pycnogonid species new to Korean Fauna. - Proc. Nat. Sci. Res. Inst., Kangreung Natn. Univ., 2 (1): 1-9.
- Kim, I.H. & J.-S. Hong (1986): Korean shallow-water Pycnogonids based on the collections of the Korea Ocean Research and Development Institute. - Kor. J. Syst. Zool., 2 (2): 35-52.
- Kim, I.H. & J.-S. Hong (1987): *Bradypallene espina*, new genus and new species, a Pycnogonid from the East Sea of Korea (Pycnogonida). - Kor. J. Zool., 30 (3): 272-276.
- Kim, I.H. & J.-S. Hong (1989): Korean shallow-water Pycnogonids based on the collections of the Korea Ocean Research and Development Institute. - Korean J. Syst. Zool., 2 (2): 35-52.
- Kindle, E. & E.J. Whittaker (1918): Bathymetric check-list of the marine invertebrates of eastern Canada. - Contr. Can. Biol. Fish.: Prof. Pap., 38A: 229-284.
- King, L.A. (1913): Clyde fauna; supplementary list. - Ann. Rep. Mar. Biol. Ass. W-Scotland, 1912: 60-97.
- King, P.E. (1969): Sea spiders on Gower shores. - Gower, 20: 19-23.
- King, P.E. (1970): Sea spiders in Glamorgan. - Nature in Wales, 12 (1): 24-31.
- King, P.E. & J.H. Jarvis (1970): Egg development in a littoral pycnogonid *Nymphon gracile*. - Mar. Biol., 7 (4): 294-304. [embryology]
- King, P.E. & G.B. Crapp (1971): Littoral pycnogonids of the British Isles. - Field Stud., 3 (3): 455-480.
- King, P.E., D. Wyer & J.H. Jarvis (1971): Littoral pycnogonids of Galway Bay. - Ir. Nat. J., 17 (3): 78-85.

- King, P.E. (1972): The marine flora and fauna of the Isles of Scilly. Pycnogonida. - J. Nat. Hist., 6: 621-624.
- King, P.E. (1972): Pycnogonids: 1-144; London; St. Martin's Press.
- King, P.E. (1974): British Sea Spiders (Arthropoda: Pycnogonida). Keys and notes for the identification of the species. Synopses of the British Fauna (New Ser.), 5: i-vi, 1-68; London & New York, Academic Press.
- King, P.E. & A.S.N. El-Hawwi (1978): Spermatogenesis in the Pycnogonid *Pycnogonum littorale* (Ström). - Acta Zool., 59: 97-103. [cytology, genital system]
- King, P.E. (1982): Pycnogonids of the Channel Islands. - Ann. Bull., Soc. Jersiaise, 23 (2): 230-232.
- King, P.E. (1986): Sea Spiders: A revised key to the adults of littoral Pycnogonida in the British Isles. - Field Stud., 6 (3): 493-516.
- King, P.E. & R.M. Case (1986): Sea spiders (Pycnogonida) in and around Milford Haven (South West Wales). - Field Stud., 6 (3): 517-529.
- King, P.E., J.P. Thorpe & G.P. Wallis (1986): A biochemical, genetic and morphological investigation of the species within the genus *Endeis Philippe* (Pycnogonida: Endeidae). - J. Exp. mar. Biol. Ecol., 98 (1-2): 115-128. [anatomy, biochemistry, genetics]
- Kingsley, J.S. (1894): The classification of the Arthropoda. - Am. Nat., 28: 118-135.
- Kishida, K. (1927): Pantopoda. In: Nikon Dōbutsu tsaran: p. 989; Tokyo Hokoryūkan. [in Japanese]
- Kitching, J.A. & T.T. Macan (1934): A study in sublittoral ecology. - J. Mar. Biol. Ass. U.K., 19 (2): 677-706.
- Kitching, J.A. & V.M. Thain (1983): The ecological impact of the sea urchin *Paracentrotus lividus* (Lamarck) in Lough Ine, Ireland. - Phil. Trans. Roy. Soc., London, (B) 300: 513-552.
- Kito, K. (1975): Pycnogonida. In: Preliminary report on the phytal animals in the *Sargassum confusum* region in Oshoro Bay, Hokkaido. - J. Fac. Sci., Hokkaido Univ., (6) Zoology 20 (1): 141-158.

Klausnitzer, B. & K. Richter (1981): Stammesgeschichte der Gliedertiere. Die neue Brehm-Bücherei, 541: 1-160; A. Ziemsen Verl., Wittenberg-Lutherstadt [Pycnogonida pp. 69-70]

Knight, G.S. (1974): Benthic community structure in Lyttelton Harbour, New Zealand. - N.Z. J. Mar. Freshw. Res., 8 (2): 291-306.

Knight-Jones, E.W. & A. MacFadyen (1959): The metachronism of limb and body movements in annelids and arthropods. - Proc. XVth Int. Congr. Zool., 1959: 969-971.

Knox, G.A. (1970): Antarctic marine ecosystems. In: M.W. Holdgate (ed.). Antarctic Ecology, 2: 69-96; Sci. Committee Antarct. Res.; London & New York; Academic Press.

Kocatas, A. (1970): Contribution à l'étude des peuplements des horizons supérieurs de substrat rocheux du Golfe d'Izmir (Turquie). - Sci. Monogr. Fac. Sci. d'Ege Univ., 12: i-viii, 1-93.

Koch, L.E. (1980): The primary types of Arachnida, Chilopoda, Diplopoda, Insecta, Onychophora and Pycnogonida in the Western Australian Museum. - Rec. W. Austr. Mus., 8 (2): 295-326.

Koehler, R. (1885): Recherches sur la faune marine des îles Anglo-Normandes. - Bull. Soc. Sci., Nancy, Fasc. 17 (2): 51-120.

Kölbl, C. (1886): Crustaceen, Pycnogoniden und Arachniden von Jan Mayen. Die österreichische Polarstation, Jan Mayen... - Die internationale Polarforschung, 1882-1883, 3: 53-55.

Kölliker, A. (1834): Beiträge zur Entwicklungsgeschichte wirbelloser Tiere. 1. Ueber die ersten Vorgänge im befruchteten Ei. - Arch. Anat. Physiol., 1843: 136-137.

Konno, K. & H. Uchida (1972): A preliminary list of marine animals found in the vicinity of the Fukaura Marine Biological Laboratory, part 4. - Rep. Fukaura Mar. Biol. Lab., 4: 2-10.

Kowalewsky, A. (1892): Ein Beitrag zur Kenntnis der Exkretionsorgane der Pantopoden. - Mém. Acad. imp. Sci. St. Pétersbourg, (7) 38 (12): 1-9, 1 pl. [anatomy, excretion]

Kozloff, E.N. (1974): Keys to the marine invertebrates of Puget Sound, the San Juan Archipelago, and adjacent regions. ix + 226 pp. Seattle & London: Univ. Wash. Press.

- Kraeuter, J.N. (1973): Pycnogonida from Georgia, U.S.A. - J. nat. Hist., 7 (5): 493-498.
- Kraeuter, J.N. & D.R. Calder (1978): Class Pycnogonida. In: R.G. Zingmark (ed.). An annotated checklist of the biota of the coastal zone of South Carolina, p. 153. Columbia, S.C.: Univ. S. Carolina Press.
- Kramp, P.L. (1963): Pycnogonida. In: Summary of the zoological results of the Godthaab Expedition, 1928: 71-72. - Medd. Grønland, 81 (7): 1-115.
- Krapp, F. (1973): Pycnogonida from Pantelleria and Catania, Sicily. - Beaufortia, 21 (277): 55-74.
- Krapp, F. (1973): A fourth Mediterranean *Rhynchothorax* and remarks on the genus (Pycnogonida). - Bull. Zoöl. Mus., Univ. Amsterdam, 3 (17): 119-124.
- Krapp, F. (1975): Bathyale und zirkalittoriale Pantopoden (Pycnogonida) aus dem Adriatischen und Ligurischen Meer, mit *Callipallene acribica* n.sp. - Bonn. zool. Beitr., 26 (1-3): 280-290.
- Krapp, F. (1975): New records of *Endeis charybdea* (Dohrn, 1881) (Pycnogonida) in the Atlantic and Mediterranean Seas. - Sarsia, 59: 85-94.
- Krapp, F. & J.N. Kraeuter (1976): Additions to the Pycnogonida of Georgia. - Bonner Zool. Beitr., 27 (3/4): 336-346.
- Krapp, F. (1980): Neue Pantopodenfunde (Pycnogonida, Arthropoda) nahe der belgischen Antarktis-Station. - Bull. Inst. Roy. Sci. Nat. Belg., Bruxelles, Biol., 52 (18): 1-8.
- Krapp, F. (1981): Pantopodi o Picnogonidi. In: A. Mondalori, (ed.). Grande Enciclopedia illustrata degli Animali, Invertebrati, 3: 95-96.
- Krapp, F. (1983): Pantopoden aus Nordwestafrika (Pycnogonida). - Bonner Zool. Beitr., 34 (1-3): 405-415.
- Krapp, F. & R. Sconfietti (1983): *Ammothea hilgendorfi* (Böhm, 1879), an adventitious Pycnogonid new for the Mediterranean Sea. - Mar. Ecol., 4 (2): 123-132.

Krapp, F. (1986): *Achelia borealis* (Schimkewitsch, 1895) und *Achelia neotenica* sp.n. - ein Vergleich (Pantopoda, Pycnogonida). - Bonn. Zool. Beitr., 37 (4): 301-310.

Krapp, F. & J. Nieder (1993): Pycnogonids as prey of blenniid fishes. - Cah. Biol. Mar., 34: 383-386. [*Tanystylum orbiculare*, *Tanystylum conirostre*, *Ammothella cf. longipes*, *Anoplodactylus virescens*, *Achelia* sp., food for fishes, Blenniidae, *Lipophrys pholis*].

Krapp-Schickel, G. & F. Krapp (1975): Quelques traits de l'écologie d'Amphipodes et de Pycnogonides provenant d'un îlot nord-adriatique. - Viet Milieu (B), 25 (1): 1-31.

Krishnan, G. (1955): The nature of the cuticle of Pycnogonida. - Nature, London, 175: 904.

Krohn, A. (1849): Notiz über die Eierstöcke der Pycnogoniden. - Frorieps Notizen, (3) 9 (191): 225-226.

Krohn, A. (1855): Ueber das Herz und den Blutumlauf in den Pycnogoniden. - Arch. Naturgesch., 1: 6-8, 1 pl.

Krøyer, H. (1838): Grønlands Amfipoder beskrevne. - Kongl. Vidensk. Selsk. Skrif., 7: 229-236.

Krøyer, H. (1840): Om Pycnogonidernes Forvandlinger. - Naturhist. Tidskr., Kjøbenhavn, (1) 3: 299-306.

Krøyer, H. (1845): Bidrag til Kundskab om Pycnogoniderne eller Søspindlerne. - Naturhist. Tidskr., Kjøbenhavn, (2), 1: 90-139.

Krøyer, H. (1846): Beitrag zur Kenntnis der Pycnogoniden. - Isis von Oken, 1846 (6) (2): 429-448.

Krøyer, H. (1849): Gaimards Voyage en Scandinavie, en Laponie, au Spitzberg et aux Feroe. Crustacés: XXXV-XXXIX; Paris.

Kühlmorgen-Hille, G. (1965): Qualitative und quantitative Veränderungen der Bodenfauna der Kieler Bucht in den Jahren 1953-1965. - Kieler Meeresforsch., 21 (2): 167-191.

Kühne, C. (1952): Untersuchungen über das Grossplankton in der Deutschen Bucht und im Nordsylter Wattenmeer. - Helgol. Wiss. Meeresunters., 4 (1): 1-54.

Kurian, C.V. (1948): A collection of pycnogonids from the Vizhinjam coast. - Proc. 35th Ind. Sci. Congr., 3: 195.

Kurian, C.V. (1953): A preliminary survey of the bottom fauna and bottom deposits of the Travancore coast within the 15-fathom line. - Proc. Natn. Inst. Sci. India, 19 (6): 746-775.

- Kussakin, O.G. (1975): A list of the macrofauna in the intertidal zone of the Kurile Islands, with remarks on zoogeographical structure of the region. - Publ. Seto Mar. Biol. Lab., 22 (1-4): 42-74.
- Lacourt, A.W. (1974): De mariene fauna van Ameland, een vergeten eiland. - Het Zeepard, 33 (4-5): 67-78.
- Lagardère, F. (1972): Les fonds de pêche de la côte ouest de l'Ile d'Oléron - cartographie bionomique. 3. Les peuplements benthiques. - Téthys, 3 (3): 507-539.
- Lamarck, J.B.P.A. (1838): Arachnidae. In: Histoire naturelle des animaux sans vertèbres: 100-105; 2nd ed., 5. Paris.
- Lankester, E.R. (1905): The structure and classification of the Arachnida. Subclass I: Pantopoda. - Quart. J. Microsc. Sci., (N.S.), 48: 223-225.
- Larramendy, M.L. (1974): Anotaciones sobre *Colossendeis geoffroyi* (Mane-Garzon, 1944). - Neotropica, 20 (63): 149-152.
- Larramendy, M.L. (1975): Un nuevo *Pallenopsis* Wilson, 1881, para el Mar Argentino (Pycnogonida, Pallenidae). - Neotropica, 21 (65): 94-98.
- Larramendy, M.L. & Z.L.A. de Castellanos (1978): Sobre un nuevo picnogónido del Mar Argentino (Pycnogonida, Pallenidae). - Neotropica, 24 (72): 91-97.
- Latreille, P.A. (1804): Histoire Naturelle. Générale et Particulière des Crustacés et des Insectes, 7: 330-333.
- Latreille, P.A. (1804): Nouveau Dictionnaire d'Histoire Naturelle, 24: 137.
- Latreille, P.A. (1806): Genera Crustaceorum et Insectorum, 1: 143-144; Paris.
- Latreille, P.A. (1810): Considérations Générales sur l'ordre natural des animaux composant les classe des Crustacés, des Arachnides et des Insectes: 39, 107, 115; Paris.
- Latreille, P.A. (1817): Les Cruistacés, les Arachnides et les Insectes. Pycnogonides. In: G. Cuvier. La Régne Animal distribu'd'après son Organisation, 3: 109-112.
- Latreille, P.A. (1818): Nouveau dictionnaire d'Histoire Naturelle, nouvelle éd., 26: 14.
- Latreille, P.A. (1825): Pycnogonida. In: Familles naturelles du règne animal: 318-319; Paris.

Latreille, P.A. (1831): Cours d'Entomologie ou de l'Histoire Naturelle des Crustacés, des Arachnides, des Myriapodes et des Insectes: 543-545.

Laubier, L. (1966): Le coralligène des Alberes: Monographie biocenotique. - Ann. Inst. Oceanogr., Monaco, 43 (2): 137-316, 1 pl.

Lauterbach, K.E. (1973): Schlüsselereignisse in der Evolution der Stammgruppe der Euarthropoda. - Zool. Beitr., new ser., 19 (2): 251-299.

Leach, W.E. (1814): The Zoology Miscellany, 1: 33-34, 43-45; London.

Leach, W.E. (1815): A tabular view of the external characters of four classes of animals which Linné arranged under Insecta; with the distribution of the genera composing three of these classes into orders etc. and descriptions of new genera and species. - Trans. Linn. Soc. London, 11: 305-400.

Lebour, M.V. (1916): Notes on the life history of *Anaphia petiolata* (Kröyer). - J. Mar. Biol. Ass. U.K., 11 (1): 51-56.

Lebour, M.V. (1945): Notes on the Pycnogonida of Plymouth. - J. Mar. Biol. Ass. U.K., 26: 139-165.

Lebour, M.V. (1949): Two new pycnogonids from Bermuda. - Proc. Zool. Soc. London, 118 (4): 929-932.

Ledoyer, M. (1966): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome. II. Données analytiques sur les herbiers de phanérogames (1.). - Rec. Trav. Stat. Mar. d'Endoume, 41 (57): 135-164.

Ledoyer, M. (1966): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome. III. Données analytiques sur les biotopes de substrat meuble. - Rec. Trav. Stat. Mar. d'Endoume, 41 (57): 165-186.

Ledoyer, M. (1968): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonomes (region de Marseille principalement), IV: Synthèse de l'étude écologique. - Rec. Trav. Stat. Mar. d'Endoume, 44 (60): 125-295.

Ledoyer, M. (1969): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome. V. Étude des phénomènes nycthéméraux. - Téthys, 1 (2): 291-308.

Ledoyer, M. (1969): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome. VI. Étude des phénomènes nycthéméraux. 2. - *Téthys*, 1 (2): 309-320.

Ledoyer, M. (1969): Écologie de la faune vagile des biotopes méditerranéens accessibles en scaphandre autonome, VII: Étude des phénomènes nycthéméraux: Les variations nycthémérales des gaz dissous au sein des biotopes. - *Téthys*, 1 (2): 321-340.

Legendre, R. (1968): Morphologie et développement des Chélicérates. Embryologie, développement et anatomie des Xiphosures, Scorpions, Pseudoscorpions, Opilions, Palpigrades, Uropyges, Amblipyges et Pycnogonides. - *Fortschr. Zool.*, (N.F.), 19 (1): 1-50.

Lewes, G.H. (1859): Naturstudien am Seestrande (Seaside Studies). In: Deutsche Übersetzung von J. Frese, 194 pp, pl. 5; Berlin.

Leuckart, R. (1848): Ueber die Morphologie und die Verwandtschaftsverhältnisse der wirbellosen Thiere: 99, 120.

Leung Tack Kit, T. (1972): Étude du milieu pollué: le vieux-port de Marseille: Influence des conditions physiques et chimiques sur la physiognomie du peuplement de Quasi. - *Téthys*, 3 (4): 767-825. [pollution]

Leung Tack Kit, T. (1976): Étude de la faune marine de substrat rocheux dans la zone de rejet des eaux résiduaires de la ville de Marseille (France). - *Téthys*, 7 (2-3): 191-212.

Leunis, J. (1886): Pantopoda. In: H. Ludwig (ed.). *Synopsis der Thierkunde*, 2: 622-625.

Levine, R.J.C. & J.L. Woodhead (1987): Structural effects of *limulus* myosin light chain phosphorylation. - *Biol. J.*, 51 (2, part 2): 32A.

Lichti, P.A. (1914): Notes on the eggs of some Laguna Beach Invertebrates. - *J. Ent. Zool. Pomona Coll.*, 6 (4): 215-216.

Light, S.F. et al. (1954): Intertidal invertebrates of the Central California Coast: i-xiv, 1-446; Univ. Calif. Press.

Liljeborg, W. (1851): Bidrag till Norra Rysslands och Norriges fauna, samlade under en vetenskaplig resa is dess länder 1848. - *Kungl. Svensk. Vetensk.-Akad. Handl.*, 1850 (2): 235-341.

Linnaeus, C. (1767): *Systema Naturae per Regna tria naturae*,
13th ed., 2 (2): 1027-1029.

Lipkin, Y. & U. Safriel (1971): Intertidal zonation on rocky
shores at Mikhmoret (Mediterranean, Israel). - *J. Ecol.*,
59 (1): 1-30, pls. 1-3. [ecology]

Littlepage, J. & J.S. Pearse (1962): Biological and
oceanographic observations under an Antarctic ice shelf. -
Science, 137 (3531): 679-681.

Loeb, J. (1896): Ueber die Regeneration des Rumpfes bei
Pantopoden. Zur Theorie der Regenerationserscheinungen. -
Arch. Entwicklungs-Mechanik der Organismen, 2: 250-256.

Lönnberg, E. (1901): Über die, während der schwedischen
zoologischen Polarexpedition von 1900 eingesammelten
Pycnogoniden. - *Zool. Anz.*, 24: 205.

Lönnberg, E. (1902): List of pycnogonids collected by the
Swedish Zoological Expedition to Spitzbergen and East
Greenland in 1900. - *Kungl. Svensk. Vetensk. Akad. Handl.*,
10: 353-359.

Loman, J.C.C. (1903): Eenige mededelingen over de Pycnogoniden
der Siboda-Expeditie. - *Tijdschr. Nederl. Dierk. Ver.*, (2)
8: xxix.

Loman, J.C.C. (1904): *Pipetta weberi* n. g., n. sp. with notes
about the proboscis of the Pycnogonida. - *Tijdschr.
Nederl. Dierk. Vereenig.*, (2) 8 (3): 259-266.

Loman, J.C.C. (1904): Pycnogoniden aus der Capkolonie und
Natal. - Beiträge zur Kenntnis der Fauna von Süd-Afrika,
V. - *Zool. Jb.*, 20 (4): 375-386.

Loman, J.C.C. (1905): *Decolopoda* Eight oder *Colossendeis* Jarz.
- *Zool. Anz.*, 28: 722-723.

Loman, J.C.C. (1906): Mitteilung über den Kommensalismus
zwischen Infusorien und Pantopoden. - *Wetenschap. Vergad.
Nederl. Dierk. Ver.* Amsterdam, 1906: xix-xx.

Loman, J.C.C. (1907): Biologische Beobachtungen an einem
Pantopoden. - *Tijdschr. Nederl. Dierk. Ver.*, (2) 10 (3):
255-284.

Loman, J.C.C. (1908): Die Pantopoden der Siboga-Expedition. -
Siboga Exped., 40: 1-88.

Loman, J.C.C. (1911): Japanische Podosomata: Beiträge zur
Naturgeschichte Ostasiens, herausgegeben von F. Doflein. -
*Abh. Kaiserl. Bayerisch. Akad. Wiss. (Math.-Naturwiss.
Kl.)*, Suppl. 2 (4): 1-18.

- Loman, J.C.C. (1911): Erratum Monographie 40: Loman, Die Pantopoden der Siboga-Expedition, 1908. Codicil to Siboga Monograph, 40.
- Loman, J.C.C. (1912): Note préliminaire sur les "Podosomata" (Pycnogonides) du Musée Océanographique de Monaco. - Bull. Inst. Ocánogr., Monaco, 238: 1-14.
- Loman, J.C.C. (1915): Les Pycnogonides et les règles de la nomenclature zoologique. - Tijdschr. Nederl. Dierk. Ver., (2) 14 (2): 187-223.
- Loman, J.C.C. (1916): *Pallenopsis* and *Rigona*, with description of a new species. - Zool. Meded. uitgeg. vanwege 's Rijks-Mus. Natuurlijke Hist. Leiden, 2 (1): 15-26.
- Loman, J.C.C. (1917): Vitale kleuring bij Pantopoden. - Tijdschr. Nederl. Dierk. Ver., (2) 15 (1): 20-21.
- Loman, J.C.C. (1917): Beiträge zur Anatomie und Biologie der Pantopoden. - Tijdschr. Nederl. Dierk. Ver., (2) 16 (1): 53-102. [anatomy, biology]
- Loman, J.C.C. (1920): Pycnogoniden von Juan Fernandez. In: C. Skottsberg (ed.). The natural history of Juan Fernandez and Easter Island, 3: 137-144.
- Loman, J.C.C. (1923): Subantarctic Pantopoda from the Stockholm Museum. - Ark. Zool., 15 (9): 1-13.
- Loman, J.C.C. (1923): A new pycnogonid from Australia. - Ark. Zool., 15 (10): 1-4.
- Loman, J.C.C. (1923): Pycnogoniden aus Süd- und Westafrika. - Medd. Göteborgs. Mus., Zool. Avd., 22 (6): 1-7.
- Loman, J.C.C. (1923): The Pycnogonida. - Further Zool. Res. Swed. Antarct. Exped., 1 (2): 1-41.
- Loman, J.C.C. (1924): Les affinités des Pycnogonides. - Ann. Sci. Nat., Paris (Sér Bot., Zool.), (10) 7 (3-4): 309-333.
- Loman, J.C.C. (1925): Pycnogonides du Maroc (côte atlantique) et de la Mauritanie. - Bull. Soc. Sci. Nat. Maroc, 5 (3): 50-53.
- Loman, J.C.C. (1928): Ein neuer Pantopode aus Westindien. - Tijdschr. Nederl. Dierk. Ver., (3) 1 (2): 39-42.
- Loman, J.C.C. (1928): Note complémentaire sur les Pycnogonides de la côte atlantique du Maroc. - Bull. Soc. Sci. Nat. Phys. Maroc, 8 (1-3): 61-67.

Loman, J.C.C. (1928): Pantopoda oder Asselspinnen (Meerspinnen). - Tierwelt Deutschlands, 8 (3): 73-83.

Loman, J.C.C. (1930 ?): Addendum à la liste des Pycnogonides recoltés sur la côte atlantique du Maroc par le Dr. J. Louiville et R.-Ph. Dollfus. - Bull. Soc. Sci. Nat. Maroc, 9 (1-6): 71.

Loman, J.C.C. (1938): Note préliminaire sur les "Podosomata" (Pycnogonides) du Musée Océanographique de Monaco. - Rés. Camp. Sci. accomplies par le Prince Albert I, Monaco, 97: 277-286.

Lombas, I. & N. Anadón (1985): Estudio de la fauna de microhabitats esciafilos intermareales en Luanco (Asturias). - Revta. Biol. Univ. Oviedo, 3: 107-120.

Losina-Losinsky, L.K. (1923): White Sea Pantopods. - Izvest. Nauchn. Inst. imeni P.F. Lesgafta, 7: 39-48, 1 pl. [in Russian]

Losina-Losinsky, L.K. (1929): Ueber einige neue Formen der Pantopoda. - Zool. Jb. (Syst.), 57: 537-544.

Losina-Losinsky, L.K. (1929): Die Pantopoden des Kola-Fjords und des Barents-Meeres längs des Meridians 33°30' und 38° östl. Länge, gesammelt durch die Biologische Murman-Station 1924-28. - Raboty Murmanskoi Biologichesk. Stantsii, 3: 1-49. [in Russian, German summary]

Losina-Losinsky, L.K. (1929): Opredelitelj Pantopoda morej SSSR. - Trav. Soc. Nat. Leningrad, 59: 63-82.

Losina-Losinsky, L.K. (1930): Report on the Pantopoda of the seas of the U.S.S.R. - Trudy Leningr. Obshchest. Estestvoiput., 59 (1): 63-82.

Losina-Losinsky, L.K. (1933): Die Pantopoden der östlichen Meere der U.S.S.R. - Issledov. Morei S.S.S.R., Leningrad, 17: 43-80. [in Russian, German summary]

Losina-Losinsky, L.K. (1935): Pantopoda of the Arctic Seas of the U.S.S.R. - Vsesoyuzni Arkticheskii Inst. ?, 4: 1-140. [in Russian, German summary]

Losina-Losinsky, L.K. (1955): Class Pycnogonida - Pantopoda. In: Atlas of Invertebrates of the Far-Eastern Seas of the U.S.S.R.: 159-161; Moscow & Leningrad, Izdatyel Akad. Nauk., S.S.S.R. [in Russian]

Losina-Losinsky, L.K. & E.P. Turpaeva (1958): The genus Colossendeis (Pantopoda) in the northern part of the Pacific Ocean. - Byull. Mosk. Obshchest. Ispyt. Prirody, 63 (1): 23-33. [in Russian, English summary]

Losina-Losinsky, L.K. (1961): Pantopoda of the far-eastern seas of the U.S.S.R.. - Issledov. Dalnyevost. Morei SSSR, Leningrad, 7: 47-117. [in Russian]

Losina-Losinsky, L.K. (1964): Pantopoda from the collections of the expeditions of the "F. Litke" in 1955 and the "Ob" in 1956. In: Scientific results of the oceanographic expeditions to the northern parts of the Greenland Sea and neighbouring regions of the Arctic Basin in the years 1955-1958. - Trudy Arktichesk. Antarktichesk. Nauchno-Issledov. Inst., 259: 330-339. [in Russian]

Losina-Losinsky, L.K. & L.M. Kopanyeva (1973): New Pantopoda of the Guinean Republic coast. - Zool. Zh., 52 (7): 1083-1084 [in Russian, English summary]

Losina-Losinsky, L.K. (1982): Modern concepts of Schimkewitsch's "Periodic System" of Pantopoda and evolution of this group. - Zool. Zh., 61 (4): 485-499.

Lou, Ting-Heng (1936): Sur deux nouvelles variétés de Pycnogonides recueillies à Tsing-Tao, dans la baie de Kiao-Chow, Chine. - Contr. Inst. Zool., Natn. Acad. Peiping, 3 (1): 1-34, 4 pls.

Lou, Ting-Heng (1936): Note sur *Lecythorhynchus hilgendorfi* Böhm (Pycnogonida). - Contr. Inst. Zool., Natn. Acad. Peiping, 3 (5): 133-163.

Lubet, P. (1953): Variations saisonnières du zooplancton du Basin d'Arcachon. - Bull. Soc. Zool. Fr., 78 (2/3): 204-216.

Lütken, C. (1875): List of the Fishes, Tunicata, Polyzoa, Crustacea, Annulata, Entozoa, Echinodermata, Anthozoa, Hydrozoa and Sponges, known from Greenland. Compiled for the use of the British North Polar Expedition. XV. Crustacea of Greenland, Appendix: 163-164.

Lundstrom, A. (1934): Über den Bau des zentralen Nervensystems bei *Pycnogonum littorale*. - Kungl. Fysiogr. Salsk. Lund, Forh., 5 (1): 1-14. [anatomy, nervous system]

MacBride, E.W. (1914): Pantopoda. In: Text-book of embryology, 1: 169, 242-243, 285; London: Macmillan.

MacCloskey, L.R. (1967): New and little known benthic pycnogonids from North Carolina. - J. nat. Hist., 1: 129-134.

MacCloskey, L.R. (1973): Pycnogonida. Marine flora and fauna of the northeastern United States. U.S. Dept. Commerce, NOAA Tech. Rep. NMFS CIRC-386: 1-12.

- MacDonald, J.A. (1981): Temperature compensation in the peripheral nervous system: Antarctic vs. temperate poikilotherms. - *J. Comp. Physiol., A, Sens. Neur. Behav. Physiol.*, 142 (3): 411-418.
- MacFadyen, A. (1954): The invertebrate fauna of Jan Mayen Islands (East Greenland). - *J. Anim. Ecol.*, 23: 261-297.
- MacGintie, G.E. (1955): Distribution and ecology of marine invertebrates at Point Barrow, Alaska. - *Smiths. misc. Coll.*, 128 (): 1-201.
- Mañé-Garzon, F. (1944): Notas sobre pantopodas. 1. *Colossendeis geoffroyi*, nov. sp., de la plataforma continental frente al Rio de la Plata. - *Comm. Zool. Mus. Hist. Nat. Montevideo*, 1 (15): 1-7.
- Mañé-Garzon, F. (1948): Morfologia y filogenia de animales marinos. - *Cién. Invest., Buenos Aires*, 4 (5): ?.
- Manton, S.M. (1973): Arthropod phylogeny - a modern synthesis. - *J. Zool. Soc. London*, 171 (1): 111-130.
- Manton, S.M. (1977): The Arthropoda; Habits, functional morphology and evolution: 540 pp; I-VIII, 1-184; Oxford Univ. Press.
- Manton, S.M. (1978): Habits, functional morphology and the evolution of pycnogonids. - *Zool. J. Linn. Soc. London*, 63 (1-2): 1-21.
- Marcus, E. (1940): *Pallenopsis fluminensis* (Kröyer) e as *Pallenopsis* sud-atlanticas restantes (Arthropoda, Pantopoda). - *Revta. Ent., Rio de Janeiro*, 11 (1-2): 180-199.
- Marcus, E. (1940): Os Pantopoda brasileiros e os demais sul-americanos. *Bol. Fac. Fil., Ciên. Letr. Univ. Sao Paulo*, 19 (Zool. 4): 3-179, 17 pls.
- Marcus, E. (= E. Du-Bois Reymond) (1952): A hermaphrodite pantopod. - *An. Acad. Bras. Ciên.*, 24 (1): 23-30.
- Marcus, E. (= E. Du-Bois Reymond) (1959): Ein neuer Pantopode aus Foraminiferensand. - *Kieler Meeresforsch.*, 15 (1): 105-107.
- Marcus, E. (= E. Du-Bois Reymond) (1963): A new *Pycnogonum* from Brazil. - *Bol. Inst. Oceanogr., Sao Paulo*, 12 (3): 3-10.

- Markham, J. (1986): Class Pycnogonida (= Pantopoda) (Sea Spiders). In: W. Sterrer (ed.). *Marine Fauna and Flora of Bermuda: A systematic Guide to the identification of marine organisms*, pp. 275-277; New York, etc.: John Wiley & Sons.
- Marshall, A.J. & W.D. Williams (1972): Subphylum Pantopoda. In: *Textbook of Zoology, Invertebrates*: 480-481; London: Macmillan.
- Marshall, N.B. (1954): Pycnogonids. In: *Aspects of Deep Sea Biology*: 121, 151, 356; London: Hutchinson.
- Marshall, N.B. (1979): *Developments in deep-sea biology*. i-ix + 566 pp. Poole, Dorset (U.K.); Blandford Press.
- Marshall, W. (1895): Asselsspinnen. In: *Die Deutschen Meere und ihre Bewohner*: 287-289; Leipzig.
- Martin, A.L. (1982): Fauna distribution maps of the sea shore monitoring sites in Long Island and Roaringwater Bays, County Cork, Ireland 1975-1982. Sherkin Isl. Mar. Stat. Shore Surv. Unit, Sherkin Isl., County Cork, XIII: 102.
- Massy, A.L. (1912): Report of a survey of trawling grounds on the coasts of the counties Down, Louth, Meath and Dublin. Part 3, Invertebrate Fauna. - Sci. Inv. Fish. Branch Ireland, 1911 (1): 1-125.
- Matthews, S.W. (1971): Antarctica's nearer side. - Natn. Geogr. Mag., 140 (5): 622-655.
- Mauchline, J. & P.F. Harvey (1983): The Scyphomedusae of the Rockall Trough, Northeastern Atlantic Ocean. - J. Plankt. Res., 5 (6): 881-890.
- Mauchline, J. (1984): Pycnogonids caught in bathypelagic samples from the Rockall Trough, Northeastern Atlantic Ocean. - J. nat. Hist., 18 (2): 315-322.
- Maurer, D. & J.A. Vargas (1984): Diversity of soft bottom benthos in a tropical estuary (Gulf of Nicoya, Costa Rica). - Mar. Biol., Berlin (81): 97-106.
- Meek, A. & B. Storrow (1912): Biological investigations. Rep., Dove mar. Lab., Cullercoats, (N.S.), 1: 61-67.
- Meglitsch, P.A. (1967): Invertebrate Zoology. Oxford Univ. Press.
- Meinert, F. (1898): Pycnogonida. Dansk. Ingolf-Exped., 3 (1): 1-68.

Meinert, F. (1899): Om Ingolf-Expedition op de paa den
indsamlede Pycnogonider. Overs. Kongl. Dansk. Videnskab.
Selsk. Forhandl., 1898: 367-377.

Meinert, F. (1899): Pycnogonida. Danish Ingolf-Exped., 3 (1):
1-71. [translation of Meiner, 1898]

Meisenheimer, J. (1902): Ueber die Pantopoden und ihre
systematische Stellung. - Verh. Dtsch. Zool. Ges., 11-12:
57-64.

Meisenheimer, J. (1902): Beiträge zur Entwicklungsgeschichte
der Pantopoden. I. Die Entwicklung von *Ammothea echinata*
Hodge bis zur Ausbildung der Larvenform. - Z. Wiss. Zool.,
72 (2-3): 191-248. [development, embryology]

Meisenheimer, J. (1912): Pantopoden. Handwörterbuch der
Naturwissenschaften, Jena, 7: 490-495, 2nd. ed.

Meisenheimer, J. (1925): Pantopoda. Die Tierwelt der Nord- und
Ostsee, 11: 1-12.

Meisenheimer, J. (1927): Pantopoda (Nachtrag). Die Tierwelt der
Nord- und Ostsee, 11a: 1-3.

Mello-Leitao, A.C. da G. de (1945): Uma espécie nova do género
Pycnogonum Brünnich, 1764 (Pycnogonidae, Pantopoda). -
Bolm. Mus. Nac., Rio de Janeiro, N.S. (Zool.), 42: 1-4.

Mello-Leitao, A.C. da G. de (1946): Nova gênero de Pantopodes
de Baía de Guanabara. - An. Acad. Bras. Ciên., 18 (4):
291-296.

Mello-Leitao, A.C. da G. de (1949): Uma nova espécie de
Pallenopsis do Atlântico Sul (Pantopoda, Pallenidae). -
Arq. Mus. Paranaense, 7 (7): 299-307.

Mello-Leitao, A.C. da G. de (1949): Nova espécie de *Halosoma*
Cole, 1904 (Pantopoda, Phoxichlidiidae Sars, 1891). - An.
Acad. Bras. Cién., 21 (2): 167-173.

Mello-Leitao, A.C. da G. de (1955): Novo Pantopodo de Baie de
Guanabara (*Melloleitanianus candidoi*, Phoxichilidiidae,
Sars 1891). - Revta. Biol. Mar., 5 (1952): 119-129.

Menzies, R.J., R.Y. George & G.T. Rowe (1973): Pycnogonida:
scattered references. In: Abyssal Environment and Ecology
of the World Oceans: 5-18; New York: John Wiley.

Merton, H. (1906): Eine auf *Tethys leporina* parasitisch lebende
Pantopodenlarve (*Nymphon parasiticum* n.sp.). - Mitt. Zool.
Stat. Neapel, 18 (1): 136-141. [gastropod host]

Meyer, K.-E. (1964): Die Häutung und die neurosekretorischen Zellen im Zentralnervensystem bei dem adulten Pantopoden *Pycnogonum litorale* (Ström). Diss., Math.-Naturwiss. Fak. Georg-August-Univ., Göttingen: 1-67. [nervous system]

Mielch, W. (1914): Eine Untersuchungsfahrt des deutschen Forschungsdampfers Poseidon nach der Barentssee (Weisses Meer) im Juni und Juli 1913. - Mitth. Dtsch. Seefisch.-Ver. ?, 2: 3-30.

Miers, E.J. (1875): Descriptions of new species of Crustacea collected at Kerguelen's Island by the Rev. A.E. Eaton. - Ann. Mag. Nat. Hist., (4) 16: 73-76.

Miers, E.J. (1875): Descriptions of three additional species of Crustacea from Kerguelens Land and Crozet Island with remarks upon the genus *Paramaera*. - Ann. Mag. Nat. Hist., (4) 16 (92): 115-118.

Miers, E.J. (1877): List of the species of Crustacea collected by the Rev. A.E. Eaton at Spitzbergen in the summer of 1873, with their localities and notes. - Ann. Mag. Nat. Hist., (4) 19: 131-140.

Miers, E.J. (1877): Report on the Crustacea collected by the naturalists of the Arctic Expedition in 1875-76. - Ann. Mag. Nat. Hist., (4) 20: 96-110.

Miers, E.J. (1877): Crustacea, Pycnogonida. In: Zoology of Kerguelen's Island: 12-15; London.

Miers, E.J. (1879): Crustacea (and Pycnogonida) of the Transit of Venus Expedition, Kerguelen Island. - Phil. Trans. Roy. Soc. London, 168: 200-214.

Miers, E.J. (1881): On a small collection of Crustacea and Pycnogonida from Franz-Josefs-land, collected by Mr. B. Leigh Smith. - Ann. Mag. Nat. Hist., (5) 7: 45-51.

Miers, E.J. (1881): A further note on *Anomorhynchus* (or *Colossendeis*) *Smithii*. - Ann. Mag. Nat. Hist., 5 (7): 264-265.

Miers, E.J. (1884): Pycnogonida. In: Report on the Zoological Collections made in the Indo-Pacific during the Voyage of HMS Alert 1881-82: 323-326.

Milne-Edwards, A. (1881): Compte rendu sommaire d'une exploration zoologique, faite dans le Méditerranée, à bord du navire de l'Etat le "Travailleur". - C.R. Acad. Sci., Paris, 93: 876-882.

Milne-Edwards, A. (1881): Compte rendu sommaire d'une exploration zoologique, faite dans l'Atlantique, à bord du navire le "Travailleur". - C.R. Acad. Sci., Paris, 93: 931-936.

Milne-Edwards, A. (1882): Raports sur les travaux de la commission... d'étudier la faune sous-marine dans les grandes profondeurs de la Méditerranée et de l'Océan Atlantique. - Arch. Miss. Sci. litt., (3), 9: 1-59.

Milne-Edwards, H. (1840): Histoire Naturelle des Crustacés... 3. Ordre des Araneiformes ou Pycnogonides: 530-537.

Milne-Edwards, H. (1845): Recherches zoologiques faites pendant un voyage sur les côtes de la Sicile: Rapport adressé à M. le ministre de L'instruction publique. - Ann. Sci. Nat., Paris (3, Zool.), 3: 129-145.

Minaard, V.A. & M. Zamponi (1985): Estudios sistemáticos de algunos pantópodos de la región subantártica. - Hist. Nat., 4 (28): 257-279.

Miner, R.W. (1950): Class Pycnogonida (sea spiders). In: Fieldbook of Seashore Life: 538-547, 4 pls.; New York: G.P. Putnam.

Mitchell, R., R.C. Earll & F.A. Dipper (1983): Shallow sublittoral ecosystems in the Inner Hebrides. - Proc. Roy. Soc. Edinburgh, 83B: 161-184.

Miyazaki, K. & T. Makioka (1988): Observations on the ovarian structure and oogenesis of some Pycnogonids. - Proc. Arthropodan Embryol. Soc. Japan (XXIII ann. meeting, May 22-23, 1987, Sugadaira, Nagano): 15-16. [embryology, genital system]

Miyazaki, K. & T. Makioka (1991): Structure of the adult female reproductive system and oogenetic mode in the sea spider, *Endeis nodosa* (Pycnogonida; Endeidae). - J. Morphol., 209 (3): 257-263.

Moazzam, M. (1979): Some observations on the seasonal population changes and breeding biology of *Propallene kempfi* Calman (Pycnogonida). Abstr. Pap. submitted for the 42nd Ann. Meeting, Am. Soc. Limnol. Oceanogr. (18-21 June 1979). New York: State Univ., Stony Brook, N.Y.

Moazzam, M. (1987): Studies on the sea spiders (Pycnogonida) of Pakistan coast. 1. Range extension of *Pigrogromitus timsanus* Calman 1927 into the waters of Karachi. - Pak. J. Zool., 19 (3): 301-302.

Möbius, K. (1873): Die wirbellosen Tiere der Ostsee. Bericht über die Expedition zur Untersuchung der Ostsee im Sommer 1871 auf S.M. Avisodampfer "Pommerania", 1873: 126; Berlin.

Möbius, K. (1873): Die auf der Fahrt nach Arendal gesammelten Tiere. In: Bericht über die Expedition der Untersuchung der Ostsee im Sommer 1871 auf S.M. Avisodampfer Pommeriana: 174-184; Berlin.

Möbius, K. (1893): Pycnogonida. In: Ueber die Tiere der schleswig-holsteinischen Austernbänke und ihre physikalischen und biologischen Lebensverhältnisse. - Sitz.-Ber. Preuss. Akad. Wiss. Berlin, 1893: 67-92.

Möbius, K. (1901): Arktische und subarktische Pantopoden. - Fauna Arctica, 2 (1): 37-64.

Möbius, K. (1902): Die Pantopoden der Deutschen Tiefsee-Expedition, 1898-99. Wiss. Ergebni. dt. Tiefsee-Exped. Dampfer "Valdivia", 1898-1899, 3: 177-196.

Monniot, C. (1962): Recherches sur les graviers à *Amphioxus* de la région de Banyuls-sur-Mer. - Vie Milieu, 13 (2): 231-322.

Monniot, C. (1965): Les "blocs à *Microcosmus*" des fonds chalutables de la région de Banyuls-sur-Mer. - Vie Milieu, 16 (2B): 819-849.

Monod, T. (1971): A propos d'un *Rhopalorhynchus* australien. - Bull. Mus. natn. Hist. nat., (2) 42 (6): 1263-1267.

Montagu, G. (1808): Description of several marine animals found on the south coast of Devonshire. - Trans. Linn. Soc. Lond., 9: 81-113.

Moore, H.B. (1933): New faunistic records for the Manx region. - Rep. Mar. Biol. Stat. Port Erin, 46: 30-34.

Moore, H.B. (1934): New faunistic records for the Manx region. - Rep. Mar. Biol. Stat. Port Erin, 47: 27-28.

Moore, H.B. (1937): Pycnogonida. In: Marine Fauna of the Isle of Man: 142-144. [reprint from Proc. & Trans. Liverpool. Biol. Soc., 50]

Morgan, E., A. Nelson-Smith & E.W. Knight-Jones (1964): Responses of *Nymphon gracile* (Pycnogonida) to pressure cycles of tidal frequency. - J. Exp. Biol., 41: 825-836.

Morgan, E. (1971): The swimming of *Nymphon gracile* (Pycnogonida). The mechanisms of the leg-beat cycle. - J. Exp. Biol., 55 (1): 273-287.

Morgan, E. (1972): The swimming of *Nymphon gracile* (Pycnogonida). The swimming gait. - J. Exp. Biol., 56 (2): 421-432.

Morgan, E. (1972): The pressure sensitivity of marine invertebrates - a resumé after 25 years. - Proc. Roy. Soc. Edinburgh, (B) 73 (29): 287-299.

Morgan, E. (1977): The swimming of *Nymphon gracile* (Pycnogonida). The energetics of swimming at constant depth. - J. Exp. Biol., 71: 205-211.

Morgan, E. & S.V. Hayes (1977): The swimming of *Nymphon gracile* (Pycnogonida): The vertical lift forces generated while swimming at constant depth. - J. Exp. Biol., 71: 187-283.

Morgan, E. (1978): The energy cost of off-shore migration in *Nymphon gracile*. In: Sea Spiders (Pycnogonida). - Zool. J. Linn. Soc. London, 63 (1-2): 171-179. [physiology]

Morgan, T.H. (1890): A preliminary note on the embryology of the Pycnogonida. Circulars Johns Hopkins Univ., Baltimore ?, 9 (80).

Morgan, T.H. (1891): A contribution to the embryology and phylogeny of the pycnogonids. Studies of the pycnogonids. - Stud. Biol. Lab. Johns Hopkins Univ., Baltimore, 5 (1): 1-76.

Morgan, T.H. (1891): The relationships of the sea-spiders. Biological Lectures delivered at the Marine Biological Laboratory, Woods Hole, 1891: 142-167.

Morgan, T.H. (1892): The embryology and phylogeny of pycnogonids. - J. Roy. Microsc. Soc., 1891: 341-342.

Morgans, J.F.C. (1967): Records of breeding activity of marine invertebrates at Kaikoura. - Trans. Roy. Soc. N.Z., Zool., 9 (12): 169-174.

Morley, C. (1940): Arachnida of Suffolk: The remaining Arachnids. - Trans. Suffolk Naturalists' Soc., 4 (3): 165-174.

Morri, C.A., A. Occhipinti Ambrogi & R. Sconfietti (1982): Specie nuovo o critiche del benthos lagunare italiano. C.N.R., Progetto Finalizzato Promozione Qualita dell'Ambiente, Notiziario, 35.

Morris, B.F. & D.D. Mogelberg (1975): Identification manual to the pelagic *Sargassum* Fauna. - Spec. Publ. Bermuda Biol. Stat., Res., 11 (1973): 1-63.

Mortensen, T. (1923): The Danish Expedition to the Kei Islands 1922. - Vidensk. Medd. Dansk Naturhist. Foren. Kjøbenhavn, 76: 55-99, pls. 1-3.

Morton, J. & M. Miller (1968): Pycnogonida. In: The New Zealand Sea Shore: 217-218; London & Auckland: Collins.

Moseley, H.N. (1879): On the structure of the Stylasteridae, a family of hydroid stony corals. - Phil. Trans. Roy. Soc., London, 169 (2): 425-503, pls. 33-44.

Moseley, H.N. (1881): Report on certain Hydroid, Alcyonarian and Madreporarian Corals procured during the voyage of H.M.S. Challenger, in the years 1873-1876. - Rep. Sci. Res. Expl. Voyage HMS Challenger (Zool.), 2: 1-248, pls. I-XIV, I-II, I-XVI.

Motas, C. & Z. Popoivici (1941): Le premirer Pycnogonide recueilli dans les eaux roumaines de la Mer Noire. - Bull. Sec. Sci. Acad. Roum., Bucarest, 23 (9): 446.

Müller, G.J. (1973): Note sur la faune des fonds infralittoraux conchilifères de la partie nord-ouest de la mer Noire. - Rapp. P.-v. Comm. Int. Expl. sci. Mer. Medit., 21 (9): 627-629.

Müller, H.-G. (1989): Shallow-water Pycnogonida from coral reefs at Moorea, Society Islands, with description of *Rhynchothorax tiahurensis* n.sp. - Bonner zoologische Beiträge 40 (2): 123-139.

Müller, H.-G. (1989): Pycnogonida from the Gulf of Aden, northern Indian Ocean. - Senckenbergiana maritima 20 (5/6): 277-290.

Müller, H.-G. (1990): Shallow-water Pycnogonida from Kenya and Sri Lanka, with descriptions of three new species. - Bonner zoologische Beiträge 41 (1): 63-79.

Müller, H.-G. (1990): Flachwasser-Pantopoden von Bora Bora, Gesellschaftsinseln, S-Pazifik, mit zwei Neubeschreibungen (Pantopoda). - Senckenbergiana biologica 70 (1989) (1/3): 185-201.

Müller, H.-G. (1990): Pycnogonida from coral reefs at Réunion Island, southern Indian Ocean. - Zoologische Abhandlungen Staatliches Museum für Tierkunde Dresden 45 (9): 97-102.

Müller, H.-G. (1990): On some Indo-West-Pacific Pycnogonida from the Zoologisk Museum, Copenhagen. - Zoologische Abhandlungen Staatliches Museum für Tierkunde Dresden 45 (10): 103-110.

- Müller, H.-G. (1990): Shallow-water Pycnogonida from Martinique, French Antilles, with description of *Nymphon macabou* n.sp. - Bonner zoologische Beiträge 41 (3-4): 277-285.
- Müller, H.-G. & M. Roth (1990): Asselspinnen. Morphologie, Lebensweise, Präparation. - Mikrokosmos 79 (6): 162-174.
- Müller, H.-G. (1992): Pycnogonida from Malaysian coral reefs, including descriptions of three new species. - Bonner zoologische Beiträge, 43 (1): 155-178.
- Müller, H.-G. (1992): Shallow-water Pycnogonida from Barbados, Lesser Antilles with description of *Anoplodactylus justi* n.sp. - Studies on the Natural History of the Caribbean Region, 71: 42-52.
- Mulicki, Z. & L. Zmudzinski (1969): The stocks of zoobenthos in the southern Baltic. - Prace Morsk. Inst. Ryback. Gdnyi, (A), 15: 77-101. [in Polish]
- Munilla T. (1978): Contribución al estudio de los Picnogónidos: Aspectos morfológicos, biológicos, electroforéticos e immunológicos en relación con la sistemática de los mismos. - Publ. Univ. Auton. Barcelona Ballaterra, Fac. Cien. (Tesis Doctorales): 1-48.
- Munilla T. (1980): Dimorfismo de dos especies de Picnogónidos: *Ammothella longipes* (Hodge) y *Achelia echinata* Hodge. - Graellsia, Revta. Ent. Iber., 1978, 34: 211-221.
- Munilla T. (1980): Ciclo postlarvario anual de *Ammothella longipes* (Hodge, 1864) (Pycnogonida). - Cah. Biol. Mar., 21: 1-9.
- Munilla T. (1980): Desarrollo anual y reproducción de *Achelia echinata* HODGE, 1864 (Pycnogonida). - Cah. Biol. Mar., 21: 115-121; Roscoff.
- Munilla T. (1981): Contribución al coneixement de la distribución ecológica delas Picnogónides alla Costa Brava Catalana. - Bull. Inst. Catal. Hist. Nat., 47 (Sec. Zool., 4): 77-86.
- Munilla T. & A. De Haro (1981): An electrophoretical and immunological study of Pycnogonida, with phylogenetic considerations. - Bijdr. Dierk., 51 (2): 191-198.
- Munilla T. (1982): Pignogonifauna existente en la foce de *Halopteris* de la costa alicantina. - Actas II Simp. Iber. Estud. Bentos Mar., 3: 331-224.
- Munilla T. (1982): Algunas etapas del ciclo anual de *Tanystylum orbiculare* Wilson, 1878. - Misc. Zool., 6 (1980): 77-79.

- Munilla T. (1983): Algunas etapas del ciclo anual de *Ammothella uniunguiculata* (Dohrn, 1881) (Pycnogonida). In: Primer Simposio de Bentos, San Sebastian, 9-11 Abril 1979, (Actas), pp. 405-411.
- Munilla T. (1984): Pycnogónidos costeros de Castellon de la Plana. - Bol. Asoc. Esp. Ent., 8: 13-20.
- Munilla T. & A. De Haro (1984): 28: Pycnogónidos de les Illes Medes. In: J. Ros, I. Olivella & J.M. Gili (eds), Les sistemes naturals de les Illes Medes. In: Arxius Sec. Cien., 73. Barcelona: I.E.C.
- Munilla, T. & J.H. Stock (1984): Nuevo pycnogónido del genero *Pallenopsis* de costas de Namibia (Atlantico sudoriental). - Res. Exo. Cien., 12: 31-37.
- Munilla T. (1985): Pycnogónidos costeros del estrecho de Gibraltar (Lado Iberico). - Bolm. Soc. Portug. Ent., 76: A1.
- Munilla T. (1986): Biometría de una población de *Ammothella longipes* (Hodge, 1864) (Pycnogonida). - Orsis, 2: 103-114.
- Munilla T. (1988): Premiers Pycnogonides côtiers du détroit de Gibraltar (Côté Ibérique). - Bull. Inst. Catalana Hist. Nat., 55 (Sec. Zool., 7): 59-65.
- Munilla T. (1988): A collection of pycnogonids from Namibia/South West Africa. - Monogr. Zool. Marino, 3: 177-204.
- Munilla T. (1988): Pycnogonida, pp. 163-173. In: Barrientos, J.A. (ed.). Bases para un curso practico de entomología. - As. esp. Ent., Salamanca, 1988: i-xii, 1-754.
- Munilla T. & E. Sanchez (1988): Ecología de los primeros Pycnogónidos litorales Canarios. VI. Simp. Iber. Estud. Bentos Mar., Palma de Mallorca, 9-22 Sep. 1988: 61.
- Munilla, T. (1989): Pycnogónidos costeros del norte de España. - Misc. Zool. (Barc.), 11 (1987): 369-373.
- Munilla, T. (1991): Pycnogónidos litorales reconocidos sobre el alga *Mesophyllum lichenoides* en Nerja (Málaga, Mar de Alborán). - ORSIS (Org. Sist.), 6: 95-99.
- Munilla, T. (1993): Pycnogonids from southern Spain: Fauna 1 Project. - J. mar. biol. Ass. U.K., 73: 543-553.

- Munoz-Cuevas, A. (1974): Embryogenèse, organogenèse et rôle des organes ventraux et neuraux de *Pachylus quinamavidensis* Muoz (Arachnides, Opilions, Gonyleptidae): Comparison avec les Annélides et d'autres Arthropodes. - Bull. Mus. Natn. Hist. Nat., (3), (196, Zool.) 128: 1517-1537.
- Munroe, R. (1972): Chelicerate type-specimens in the Queensland Museum. - Mem. Qd. Mus., 16 (2): 291-307.
- Nakamura, K. & K. Sekiguchi (1980): Mating behavior and oviposition in the Pycnogonid *Propallene longiceps*. - Mar. Ecol., Progr. Ser., 2: 163-168.
- Nakamura, K. (1981): Post embryonic development of a Pycnogonid, *Propallene longiceps*. - J. nat. Hist., 15: 49-62.
- Nakamura, K. & C.A. Child (1982): Three new species of Pycnogonida from Sagami Bay, Japan. - Proc. biol. Soc. Wash., 95 (2): 282-291.
- Nakamura, K. & C.A. Child (1983): Shallow-water Pycnogonida from the Izu Peninsula, Japan. - Smiths. Contr. Zool., 386: 1-71.
- Nakamura, K. & K. Sekiguchi (1983): Seasonal occurrence of four species of Pycnogonids in Nabeta Bay, Shimoda, Japan. - Bull. Biogeogr. Soc. Japan, 38 (6): 39-43.
- Nakamura, K. (1985): A new species of Pycnogonid, *Heteronymphon horikoshii*, from the waters adjacent to Honshu, Japan. - Bull. Biogeogr. Soc. Japan, 40 (4): 31-34.
- Nakamura, K. & C.A. Child (1986): New species of Pycnogonida from waters adjacent to Japan. - Zool. Sci., 3: 893-897.
- Nakamura, K. (1987): The sea spiders of Sagami Bay. - Biol. Lab., Imp. Household, 1987: 1-43, pls. 1-40.
- Nakamura, K. & C.A. Child (1988): Pycnogonida of the Western Pacific Islands IV. On some species from the Ryukyu Islands. - Proc. biol. Soc. Wash., 101 (3): 662-670; Washington.
- Nakamura, K. & C.A. Child (1988): Pycnogonida of the Western Pacific Islands V. A collection by the Kakuyo Maru from Samoa. - Proc. biol. Soc. Wash., 101 (4): 809-816; Washington.
- Nakamura, K. & K. Sekiguchi (1988): A: Pycnogonida. In: K. Dan, K. Sekiguchi, H. Ando & H. Watanabe (eds.). Embryology of invertebrates, 2: 12-A, 2-11; Tokyo, Baihu-Kan.

Nakamura, K. & C.A. Child (1990): Pycnogonida of the Western Pacific Islands. 7. On some rare species from the Flores Sea, Indonesia. - Proc. biol. Soc. Wash., 103 (2): 304-310.

Nakamura, K. & C.A. Child (1991): Pycnogonida from Waters Adjacent to Japan. - Smiths. Contr. Zool., 512: 1-74.

Needler, A.B. (1943): Pantopoda. Canadian Atlantic Fauna, 10: 1-16.

Nesis, K.N. (1958): The Faroe-Icelandic sill as a zoogeographic boundary for fauna.- Dokl. Akad. Nauk SSSR, 122 (6): 1011-1013. [in Russian]

Nesis, K.N. (1960): Pantopoda of the shores of eastern Murman. - Trudy Murmansk. Biologichesk. Inst., 2 (6): 137-161.

Nesis, K.N. (1962): Pacific elements in the benthos of the north-western Atlantic. - In: Soviet Fish. Invest. NW Atl.: 83-98; Polar Res. Inst., Moscow.

Nesis, K.N. (1964): Variations in the bottom fauna of the Barents Sea under the influence of fluctuations of the hydrological regime. (On the Section of the Kola meridian). In: Sov. Fish. Invest. N. Europ. Seas: 129-138; Moscow. [in Russian]

Nesis, K.N. (1966): Food, water masses and depth as factors of the distribution of sea bottom animals. (On the example of Pantopods of Labrador and Newfoundland). Abstr. Pap. 2nd Int. Oceanogr. Congr., 30 May - 9 June 1966; Moscow: Nauka.

Nesis, K.N. (1967): Pycnogonida (Pantopoda) of Possjet Bay and South Sakhalin taken by the underwater exploring expedition of the Zoological Institute of the Academy of Science of the USSR in 1962-1963. - Issled. Fauny Morei, 5 (13): 248-251. [in Russian]

Nesis, K.N. (1970): Distribution of Pantopoda in the Labrador and Newfoundland Areas in connection with food, water masses and depth. In: Ecology and distribution of the sea bottom fauna and Flora. - Trans. P. Shirshov. Inst. Oceanol., 88: 149-173; Moscow.

Nesis, K.N. (1976): Class Pantopoda. In: A.N. Golikov & A.V. Zhirmunsky (eds.). Fauna and flora of Peter the Great Gulf, 363 pp., Leningrad. [in Russian, Pycnogonids pp. 76-77].

- Nesis, K.N. & A.F. Pushkin (1986): Pantopoda. In: K.V. Galaktionov (ed.). Life and conditions for its existence in the benthic zone of the Barents Sea, pp. 122-124; Apatity: Akad. Nauk SSSR. [in Russian]
- Newell, G.E. (1954): The marine fauna of Whitstable. - Ann. Mag. Nat. Hist., ser. 12 (7): 322-350.
- Nichols, D., J.A. Cooke & J. Whiteley (1971): Pycnogonids. In: The Oxford Book of Invertebrates: 126, 127, 204; Oxford Univ. Press.
- Nicol, J.A.C. (1967): Pycnogonids. In: The Biology of marine Animals: 123, 244, 257. London: Pitman.
- Nicolet, H. (1849): Araneiformes, I. Picnogonidos. In: C. Gay, Historia fisica y politica de Chile, 3. Atlas zoológico, Crustacéos: 304-308.
- Nogueira, M. (1956): Contribucao para o estudo dos Pantópodos das costas de Portugal. - Arq. Mus. Bocage, (1), 27: 65-105.
- Nogueira, M. (1967): Bases para a determinacao dos Pantópodos das costas portuguesas. - Arq. Mus. Bocage, (2) 1 (15): 283-341.
- Nordgaard, O. (1897): Nogle oplysninger om Puddelfjorden (temperatur, saltgehalt, plankton, etc. 1896-97). - Bergens Mus. Aarbog, 1897 (15): 1-19.
- Nordgaard, O. (1898): 2den beretning om de ved den bioogiske station i Bergen foreagne usklaeknings forsøg med lakserogen i saltvand. - Bergens Mus. Aarbog, 1898 (8): 9.
- Nordgaard, O. (1898): Undersgelser i fjordene ved Bergen 1897-1898. - Bergens Mus. Aarbog, 1898 (10): 20, pls.
- Norman, A.M. (1865): Report of deep-sea dredging off the coast of Northumberland and Durham. - Nat. Hist. Trans. Northumberland, Durham & Newcastle-upon-Tyne, 1: 12.
- Norman, A.M. (1866): Report of the Committee appointed for the purpose of exploring the coast of Hebrides by means of the dredge, part II: On the Crustacea, Echinodermata, Polyzoa, Actinozoa and Hydrozoa. - Rep. Brit. Ass. Adv. Sci., 1866: 193-206.
- Norman, A.M. (1869): Last report on dredging among the Shetland Isles, Part II, Crustacea etc. - Rep. Brit. Ass. Adv. Sci., 1868: 247-346.
- Norman, A.M. (1870): Final Shetland dredging report. - Rep. Brit. Ass. Adv. Sci., 1868-1869.

- Norman, A.M. (1873): Pycnogonida. In: C. Wyville Thompson: The Depths of the Sea: 129; London: Macmillan.
- Norman, A.M. (1894): A month on the Trondhjem Fjord. - Ann. Mag. Nat. Hist., 13 (6): 151-164.
- Norman, A.M. (1903): Notes on the natural history of East Finmark. - Ann. Mag. Nat. Hist., 11 (7): 168.
- Norman, C.A. (1901): Pantopoda (Pycnogonoidea). - Mus. Normanianum, pt. 2, 2nd ed.
- Norman, C.A. (1908): The Podosomata (= Pycnogonida) of the temperate Atlantic and Arctic Ocean. - J. Linn. Soc. Lond. (Zool.), 30: 198-238.
- O'Connell, G.W. (1980): Biology and systematics of the Ammotheidae (Arthropoda: Pycnogonida). - Am. Zool., 20 (4): 786.
- Oersted, A.S. (1844): De regionibus marinis. Elementa topographiae historiconaturalis freti Öresund: 71-73; Havniae.
- Ogawa, K. & K. Matsuzaki (1985): Discovery of bivalve-infesting Pycnogonida, *Nymphonella tapetis*, in a new host, *Hiatella orientalis*. - Zool. Sci., 2 (4): 583-589. [bivalve host]
- Ohshima, H. (1927): *Nymphonella tapetis* n. g., n. sp., a pycnogon parasite in a bivalve. - Annot. Zool. Japon., 11 (3): 257-263. [bivalve host]
- Ohshima, H. (1927): Notes on some pycnogons living semiparasitic on holothurians. - Proc. Imp. Acad., Tokyo, 3 (9): 610-613. [echinoderm host]
- Ohshima, H. (1927): Picnogono parazite vivanta en Bivalvo. - Bulteno Sci. Fak. Terkult., Kyushu Imp. Univ., Tokyo, 2 (5): 366-379. [in Japanese, Esperanto summary] [bivalve host]
- Ohshima, H. (1927): Piknogonoj alkrocigantaj al Holoturioj. - Bulteno Sci. Fak. Terkult., Kyushu Imp. Univ., Tokyo, 2 (5): 380-388. [in Japanese, Esperanto summary]
- Ohshima, H. (1930): Floating Pycnogonids. - Dōbutsugaku Zasshi, 42 (495): 34-35. [in Japanese]
- Ohshima, H. (1933): The adult of the bivalve-infesting pycnogonid, *Nymphonella tapetis* Ohshima. - Annot. Zool. Japon., 14 (1): 53-60. [bivalve host]

- Ohshima, H. (1933): Young pycnogonids found parasitic on nudibranchs. - Annot. Zool. Japon., 14 (1): 61-66. [gastropod host]
- Ohshima, H. (1933): Pycnogonida as plankton-organisms. - Dôbutsugaku Zasshi, 45 (532-533): 96. [in Japanese]
- Ohshima, H. (1933): Pycnogonids taken with a tow-net. - Annot. Zool. Japon., 14 (2): 211-220.
- Ohshima, H. (1933): Pycnogonids of the North Kuriles. - Bull. Biogeogr. Soc. Japan, 4: 143-150. [in Japanese]
- Ohshima, H. (1933): (Four topics on Pantopoda, lecture abstract). - Dôbutsugaku Zasshi, 45 (532-533): 94-97. [in Japanese]
- Ohshima, H. (1935): On a sea spider inhabiting the Okinawa region. - Dôbutsugaku Zasshi, 47 (557): 137-139. [in Japanese]
- Ohshima, H. (1935): A further note on *Nymphonella tapetis*: the egg-carrying mature male. - Annot. Zool. Japon., 15 (1): 95-102.
- Ohshima, H. (1935): A list of Pycnogonida recorded from Japanese and adjacent waters. - Dôbutsugaku Zasshi, 48 (8-10): 861-869. [in Japanese]
- Ohshima, H. (1937): The life-history of *Nymphonella tapetis* (Pantopoda, Eurycydiidae). - C.R. XIIème Congr. Int. Zool., Lisbonne, 1935: 1616-1625.
- Ohshima, H. (1938): Nymphonellidae, a new family of Pantopoda. - Annot. Zool. Japon., 17 (3-4): 229-233.
- Ohshima, H. (1939): La vivistorio de Piknogono (*Nymphonella tapetis*) vivanta parazite en Bivalvo. - Vol. jubil. Prof. Sadao Yoshida: 415-434. [in Japanese]
- Ohshima, H. (1942): Six-legged Pantopod, an extraordinary case of hypomery in arthropods. - Proc. Imp. Acad., Tokyo, 18 (5): 257-262.
- Ohshima, H. (1942): A remarkable case of malformed appendages in a pantopod, *Nymphonella tapetis*. - Proc. Imp. Acad., Tokyo, 18 (8): 520-523.
- Ohshima, H. (1943): A remarkable case of malformed appendages in a pantopod, *Nymphonella tapetis*. - Re. Ann. Meet. Zool. Soc. Japon., 1943: 1 p. [in Japanese]

- Ohshima, H. (1943): Du maloftaj kazoj de nenormaleco en pantopodoj. - Bull. Scu. Fac. Terkult., Kyushu imp. Univ., Tokyo, 10 (4): 371-382. [in Japanese, Esperanto summary]
- Ohshima, H. & K. Kishida (1947): Pantopoda (Encyclopaedia of the Fauna of Japan): 1005-1010; Tokyo: Honkoryukan. [in Japanese]
- Ohta, S. (1983): Photographic census of large-sized benthic organisms in the bathyal zone of Suruga Bay, Central Japan. - Bull. Ocean Res. Inst., Univ. Tokyo, 15: 1-244.
- Okuda, S. (1934): On a tubicolous polychate living in commensal with a pycnogonid. - Annot. Zool. Japon., 14 (4): 437-439. [polychate host]
- Okuda, S. (1940): Metamorphosis of a pycnogind parasitic in a hydromedusa. - J. Fac. Sci., Hokkaido Univ. (Zool. 6) 7 (2): 73-86. [hydroid host, development]
- Olsen, Ø. (1913): Pycnogonida from the "Michael Sars" North Atlantic deep-sea expedition, 1910. Rep. Sci. Res. "Michael Sars" N.-Atl. Deep Sea Exped., 1910, 3 (1) (Zool.): 3-7.
- Ortmann, A.E. (1890): Bericht über die von Herrn Dr. Döderlein in Japan gesammelten Pycnogoniden. - Zool. Jb. (Syst.), 5 (1): 157-168.
- Ortmann, A.E. (1901): Crustacea and Pycnogonida collected during the Princeton expedition to North Greenland. - Proc. Acad. Nat. Sci. Philad., 53: 144-168.
- Ottaway, J.R. (1977): Predators of sea anemones. - Tuatara, 22: 213-221. [Actinaria host].
- Packard, A.S. (1867): Observations on the glacial-phenomena of Labrador and Maine: with a view of the recent invertebrate fauna of Labrador. - Mem. Boston Nat. Hist. Soc., 1: 210-303.
- Pallas, P.S. (1766): 2. Acarus marinus seu Polygonopus. XVI. Descriptiones Insectorum. - Misc. Zool., 1766: 188-189.
- Parker, R.H. (1975): Pycnogonida. In: The study of benthic communities. A model and a review: 116, 139, 206, 266; Amsterdam, Oxford & New York: Elsevier.
- Pascoe, F.P. (1887): Notes on American Pycnogonidae. - Am. Monthly Micscosc. J., New York, 8 (1): 26-27.
- Pearse, A.S. (1914): Observations on the fauna of the rock beaches at Nahant, Mass., Pt. 2. - Bull. Wiscons. Nat. Hist. Soc., N.S., 12: 72-80.

- Peattie, M.E. & R. Hoare (1981): The sublittoral ecology of Menai Strait, II: The sponge *Halichondria panicea* (Pallas) and its associated fauna. - Estuarine, Cstl. Shelf Sci., 13: 621-635.
- Pelt, W.G. (1936): Pantopoda. Flora en Fauna der Zuiderzee, suppl.: 131-133.
- Pennant, T. (1777): Crustaceous Animals. Brit. Zool., 4: 8; London.
- Penrith, M.-L. & B. Kensley (1970): The constitutions of the intertidal fauna of rocky shores of South West Africa. Part. 1. Lüderitzbucht. - Cimbebasia, (A) 1 (9): 191-239.
- Pérez-Ruzafa, A. & T. Munilla (1992): Pycnogonid ecology in the Mar Menor (Murcia, SW Mediterranean). - Sci. Mar., 56 (1): 21-28.
- Perrier, E. (1886): Pycnogonides. In: Les Explorations Sous-Marines, 17: 302-303, 337; Paris: Hachette.
- Pfeffer, G. (1889): Zur Fauna von Süd-Georgien. - Jb. Hamburg. Wiss. Anst., 6 (2): 39-55.
- Pfeffer, G. (1890): Die Fauna der Insel Jeretik, Port Wladimir, an der Murman-Küste, 1. Nach den Sammlungen des Herrn Kapitän Horn. - Jb. Hamburg. Wiss. Anst., 7 (3): 63-96.
- Philippi, A. (1843): Über die neapolitanischen Pycnogoniden. - Arch. Naturgesch., (1) 9: 175-182.
- Picton, B.E., C.M. Howson, D.W. Connor & A. Williams (1983): Sublittoral survey of Scarba, Lunga and the Garvellachs, Firth of Lorne. - Nature Cons. Coun. CST Rep. (Banbury), 436: 1-56.
- Piel, W.H. (1991): Pycnogonid predation on nudibranchs and ceratal anatomy. - Veliger, 34 (4): 366-367. [nudibranch host]
- Pipe, A.R. (1982): Epizoites of marine invertebrates: With particular reference to those associated with the Pycnogonid *Phoxichilidium tubulariae* Lebour, the amphipod *Caprella linearis* (L.) and the Decapod *Corystes cassivelaunus* (Pennant). - Chem. Ecol., 1 (1): 61-74. [epizoans]
- Plymouth Marine Fauna, 1957, 3rd ed.
- Poore, G.C.B. (1968): Succession of a wharf-pile fauna at Lyttelton, New Zealand. - N.Z. J. Mar. Freshw. Res., 2 (4): 577-590.

- Préfontaine, G. (1933): Additions à la liste des espèces animales de l'estuaire de Saint Laurent. - Trans. Roy. Soc. Can., (3) 27 (5): 253-258.
- Préfontaine, G. & P. Brunel (1962): Liste d'invertébrés marins recueillis dans l'estuaire du Saint-Laurent de 1929 à 1934. - Naturaliste Canadien, 79 (8-9): 237-263.
- Prell, H. (1910): Beiträge zur Kenntnis der Lebensweise einiger Pantopoden. - Bergens Mus. Aarbok (N.R.), 1909 (10): 1-30.
- Preston, A. & P.G. Moore (1988): The flora and fauna associated with *Cladophora albida* (Huds.) from rockpools on Great Cumbrae Island, Scotland. - Ophelia, 29 (3): 169-186.
- Propp, M.V. (1971): Ecology of the near shore bottom communities in the Murmansk Shore Region of the Barents Sea. - Izdat. Nauk., Leningrad: 1-125. [in Russian]
- Pross, A. (1977): Diskussionsbeitrag zur Segmentierung des Cheliceraten-Kopfes. - Zoomorphologie, 86: 183-196. [anatomy]
- Pushkin, A.F. (1970): New species of the genus *Colossendeis* (Pantopoda). - Zool. Zh., 49 (10): 1488-1496. [in Russian, English summary]
- Pushkin, A.F. (1973): A new species of the genus *Cilunculus* (Pantopoda). - Zool. Zh., 52 (3): 432-434. [in Russian, English summary]
- Pushkin, A.F. (1974): *Clavigeropallene* gen. n. (Pantopoda) from the Antarctic. - Zool. Zh., 53 (6): 938-940. [in Russian, English summary]
- Pushkin, A.F. (1974): Some ecological and zoogeographic peculiarities of the Pantopoda-fauna in the Davis Sea. In: V.A.V. Golikov (ed.). Hydrobiology and biogeography of the shelf of the temperate and cold waters of the ocean: 43-55; Izdatyel Nauka, Leningrad. [in Russian]
- Pushkin, A.F. (1975): Revision of the pycnogonids (Pantopoda) of the *Pallenopsis patagonica* group from Antarctica and adjacent waters. - Byull. Sovietsk. Antarkt. Eksped., 90: 72-83. [in Russian]
- Pushkin, A.F. (1975): New species of Callipallenidae (Pantopoda) from the subantarctic. - Zool. Zh., 54 (9): 1402-1406. [in Russian, English summary]
- Pushkin, A.F. (1976): New species of Endeidae (Pantopoda) from the subantarctic. - Zool. Zh., 55 (6): 923-925. [in Russian, English summary]

- Pushkin, A.F. (1977): New species of Austrodecidae (Pantopoda) from the Antarctic. In: New species and genera of marine invertebrates. Expl. Fauna Seas, XXI (XXIV): 54-59. [in Russian with English summary]
- Pushkin, A.F. (1982): On the species composition of the genus *Decolopoda* (Pantopoda). - Zool. Zh., 59 (11): 1657-1661. [in Russian with English summary]
- Pushkin, A.F. (1984): A new species of the genus *Pycnogonum* (Pantopoda, Pycnogonidae) from the Antarctic. - Zool. Zh., 63 (7): 1102-1104. [in Russian with English summary]
- Pushkin, A.F. (1984): A new species of the genus *Colossendeis* (Pantopoda) from Antarctic. - Zool. Zh., 63 (8): 1259-1261. [in Russian with English summary]
- Pushkin, A.F. (1988): On revision of the *Colossendeis robusta* (Pantopoda) from the Southern Ocean. - Zool. Zh., 67 (6): 953-956. [in Russian with English summary]
- Pushkin, A.F. (1990): A new species of the genus *Pseudopallene* (Pycnogonida, Callipallenidae) from Antarctica. - Zool. Zh., 69 (2): 155-157. [in Russian, English summary]
- Pyefinch, K.A. (1943): The intertidal ecology of Bardsey Island, North Wales... - J. Anim. Ecol., 12: 82-108.
- Rack, G. (1971): Die entomologischen Sammlungen des Zoologischen Instituts und Zoologischen Museums Hamburg. - Mitt. Zool. Mus. Hamburg, 67: 109-133.
- Rainer, S.F. (1981): Soft-bottom benthic communities in Otago Harbor and Blueskin Bay, New Zealand. - N.Z. Oceanogr. Inst. Mem., 80: 1-38.
- Raj, B.S. (1927): Class Pycnogonida Latr. In: The littoral fauna of Krusadai Island in the Gulf of Madras. - Bull. Madras Governm. Mus., (N.S.) (Nat. Hist.), 1 (1): 157-160.
- Raj, B.S. (1930): Pycnogonida of Krusadai Island (Supplement). - Bull. Madras Gov. Mus., (N.S.), 1 (2) (7): 73-76.
- Rajagopal, A. (1963): On a new species of pycnogonid *Endeis ghaziei* from Ennore. - Proc. Ind. Acad. Sci., 57 (B) (4): 235-238.
- Rakusa-Suszczewski, S. (1981): Hypostenothermic organisms. - Polsk. Badania Polarne, 1 (4) (1980): 231-241.
- Rasmussen, E. (1973): Pycnogonida. In: Systematics and ecology of the Isefjord marine fauna (Denmark). - Ophelia, 11: 224-225.

Rasmussen, E. (1973): Systematics and ecology of the Isefjord marine fauna (Denmark). - *Ophelia*, 11: 1-507. [Pycnogonids pp. 224-225]

Rathbun, R. (1881): The littoral marine fauna of Provincetown, Cape Cod. - *Proc. U.S. Natn. Mus.*, 3: 116-133.

Rathke, J. (1799): Entomologiske Jagttagelser. - *Skr. Naturhist.-Selsk.*, Kjøbenhavn, 5 (1) (4): 191-207.

Redmond, J.R. & C.D. Swanson (1968): Preliminary studies of the physiology of the Pycnogonida. - *Antarct. J. U.S.*, 3 (4): 130-131.

Reimer, A.A. (1976): Description of a *Tetraclita stalactifera panamensis* community on a rocky intertidal Pacific shore of Panama. - *Mar. Biol.*, 35 (3): 225-238.

Ribeiro, M.A.G., P. de Alcantara-Filho, F.M. Fantinato & L.R. Tommasi (1982): Sobre a occorencia de Pantopoda na regiao da Baia de Santos, Sao Paulo (Brasil). - *Dusenia*, Curitiba, Brasil, 13 (1): 37-40.

Richards, P.R. (1976): Aspects of the biology of polar pycnogonids. D.Phil. Thesis, Council Natn. Acad. Awards, Lutton College.

Richards, P.R. & W.G. Fry (1978): Pycnogonid digestion: A study of polar forms. - *Zool. J. Linn. Soc. Lond.*, 63 (1-2): 75-97. [alimentary system]

Richters, R. (1912): Eine Pantopodenlarve von Gabelsflach (Kieler Förde). - *Zool. Jb.*, Suppl. 15 (1): 45-50, 1 pl.

Richter, S. (1982): Zur Ultrastruktur der seitlichen Sinnensorgane am Augenhügel von *Anoplodactylus pygmaeus* (Pycnogonida). - *Helgol. Wiss. Meeresunters.*, 35: 465-478.

Ricketts, E.F. & J. Calvin (1939): Between Pacific Tides. 1st. ed.: i-xxii, 1-320; Stanford Univ. Press.

Riedl, R. (1966): Pantopoda. In: Biologie der Meereshöhlen. Topographie, Faunistik und Ökologie eines unterseeischen Lebensraumes. Eine Monographie: 173; Hamburg & Berlin: P. Parey.

Riedl, R. (1983): Klasse Pantopoda (Asselsspinnen). In: Fauna und Flora des Mittelmeeres: 542-546; Hamburg & Berlin: P. Parey.

Roberts, D. (1981): Pycnogonids from Strangford Lough, Northern Ireland. - *Ir. Nat. J.*, 20 (5): 189-192.

Rodger, A. (1893): Preliminary account of natural history collections made on a voyage to the Gulf of St. Lawrence and Davis Strait. - Proc. Roy. Soc. Edinburgh, 20: 154-163.

Rogers, D.M. (1973): Distribution of meio-benthic organisms in a Texas, USA Bay in relation to season and habitat disturbance. - Am. Zool., 13 (4): 1333.

Runnström, S. (1928): Amphipoda, Isopoda and Pycnogonida from the Siberian Arctic Ocean. - Sci. Res. Norw. N.-Polar Exped. "Maud" (1918-1925), 5 (8): 1-18.

Russel, D.J. & J.W. Hedgpeth (1990): Host utilization during ontogeny by two pycnogonid species (*Tanystylum duospinum* and *Ammothea hilgendorfi*) parasitic on the hydroid *Eucopella everta* (Coelenterata: Campanulariidae). - Bijdr. Dierk., 60 (3-4): 215-224. [hydroid host]

Ryland, J.S. & A. Nelson-Smith (1975): Littoral benthic investigations on the West coast of Ireland, IV. - Proc. Roy. Ir. Acad., (B) 75 (11): 246-266.

Ryland, J.S. (1976): Pycnogonida. In: Physiology and ecology of marine bryozoans. - Adv. Mar. Biol., 14: 417-421. [bryozoan host]

Sabine, E. (1824): Marine Invertebrate Animals. Supplement to the appendix of Captain Parry's voyage for the discovery of a North-West passage in the years 1819-20: 219-240; London.

Sachi, C.F., C. Morri, A. Occhipinti-Ambrogi & R. Sconfietti (1983): Nouveaux éléments pour la zoogeographie lagunaire de la Haute Adriatique. - Rapp. Comm. Int. Mer Médit., 28 (6): 226-228.

Saenger, P., W. Stephenson & J. Moverley (1980): The estuarine macrobenthos of the Calliope River and Auckland Creek, Queensland. - Mem. Qd. Mus., 20 (1): 143-161.

Saenger, P., W. Stephenson & J. Moverley (1982): Macrobenthos of the cooling water discharge canal of the Gladstone Power Station, Queensland. - Austr. J. Mar. Freshw. Res., 33: 1083-1095.

Saint-Hilaire, K.K. (1913): Pantopoda. In: Eine systematische Sammlung der Tiere des Kowda-Fjords, 1908, 1911, 1912 und 1913. Pantopoda v katalog ekskursioni visavki ustraivaimoi krujkom studentov Fiziko-Matematicheskavo Fakulteta Imperatorskavo Yuryevskao Universiteta, Jurieff (Dorpat), 17-21 Okt. 1913: 54 pp. [in Russian]

Saito, M. & H. Suzuki (1974): Fauna in the littoral zone of Niisaki Beach in the western part of Sagami Bay, 1. Sci. Rep. Yokohama Natn. Univ., 2 (21): 31-48.

Salazar-Vallejo, S.I. & J.H. Stock (1987): Apparent parasitism of *Sabellida melanostigma* (Polychaeta) by *Ammothella spinifera* (Pycnogonida) from the Gulf of California. - Revta. Biol. Trop., 35 (2): 269-275. [Annelida host]

Saldanha, L. (1974): Estudo de povamente dos horizontes superiores da rocha litoral da costa da Arrábida (Portugal). - Arq. Mus. Bocage, (2) 5 (1): i-xiv, 1-382.

Sanchez, E.L. & T. Munilla (1987): Pycnogónidos (Pycnogonida) de la comunidad de *Dendrophyllia ramea* en el sureste de Tenerife, 17: 179-188.

Sanchez, S. (1953): Contributions à l'étude du développement des Pycnogonides. - Arch. Zool. Exp. Gén., 90 (N. & R.1): 1-7. [development]

Sanchez, S. (1954): Les glandes neurosécrétrices des Pycnogonides. - Bull. Acad. Sci., Paris ?: 1078-1080.

Sanchez, S. (1958): Cellules neurosécrétrices et organes infracérébraux de *Peripatopsis moseleyi* Wood (Onychophora) et neurosécrétion chez *Nymphon gracile* Leach (Pycnogonides). - Arch. Zool. Exp. Gén., 96 (N. & R.2): 57-62.

Sanchez, S. (1959): Le développement des Pycnogonides et leurs affinités avec les Arachnides. - Arch. Zool. Exp. Gén., 98 (1): 1-101.

Sando, H. (1964): Pycnogonida. In: Faunal list of the *Zostera marina* region at Kugurizaka coastal waters, Aomori Bay. - Bull. Mar. Biol. Stat. Asamushi, 12 (1): 30.

Sars, G.O. (1877): Prodromus descriptionis Crustaceorum et Pycnogonidarum, quase in expeditione norvegica anno 1876 observavit. - Arch. Math. Naturvidensk., Kristiania, 2: 237-271.

Sars, G.O. (1879): Crustacea et Pycnogonida nova in itinere 2do et 3tio expeditionis Norvegicae anno 1877, 1878 collecta. - Arch. Math. Naturv., Christiania, 4: 427-476.

Sars, G.O. (1879): Crustacea et Pycnogonida nova, etc. - Arch. Math. Naturvidensk., Kristiania, 4: 469-472.

Sars, G.O. (1888): Pycnogonida borealia et arctica enumerat (Prodromus descriptionis). - Arch. Math. Naturvidensk., Oslo, 12: 339-356.

- Sars, G.O. (1891): Pycnogonidea. Norw. N.-Atl. Exped., 1876-1878, 6 (Zool. 20): 1-163.
- Sars, G.O. (1891): Pycnogonidea of the Norwegian North Sea Expedition. - J. Roy. Microsc. Soc. London, 1891: 185-186.
- Savigny, L.C. (1816): Théorie des Organes de la Bouche des Crustacés et des Insectes, I, II; Paris.
- Savigny, L.C. (1816): Mémoires sur les animaux sans vertèbres, 1: 54-57.
- Savory, T. (1964): Pycnogonida. In: Arachnida: i-viii, 1-291; London & New York: Academic Press.
- Sawaya, M.P. (1941): Sobre uma larva de *Pycnogonum pamphorum* Marc. - Bolm. Fac. Fil., Ciênc. Letr., Univ. S. Paulo, 22 (Zool. 5): 278-282.
- Sawaya, M.P. (1945): *Anoplodactylus stictus* Marc. (Pantopoda) em Caiobá, Estado do Paraná. - Arqu. Mus. Paran., 4 (10): 231-236.
- Sawaya, M.P. (1947): *Nymphopsis melidae* n.sp. e observações sobre a respiração nos Pantopoda. - Bolm. Fac. Fil., Ciênc. Letr., Univ. S. Paulo, (Zool.), 12: 83-97.
- Sawaya, M.P. (1950): *Anoplodactylus aragaoi*, n.sp. and other pycnogonids collected by the hydrographic research-ship Rio Branco. - Mem. Inst. Oswaldo Cruz, 47 (1-2): 63-86, 2 pls.
- Sawaya, M.P. (1951): *Achelia besnardi*, n.sp. (Pantopoda-Ammotheidae). - Bolm. Fac. Fil., Ciênc. Letr. Univ. S. Paulo, (Zool.), 16: 271-280.
- Say, T. (1821): An account of the Arachnides of the United States. - J. Acad. Nat. Sci., Philad., 2 (1): 59-62.
- Scanland, T.B. (1979): The epibionts of *Arca zebra imbricata*: A community analysis. - Veliger, 21 (4): 475-485.
- Schimkewitsch, W. (1886): Les Arachnides et leurs affinités. - Arch. Slav. Biol., Paris, 1.
- Schimkewitsch, W. (1887): Sur les Pantopodes de l'expédition du Vettor Pisani. (Note préliminaire). - Zool. Anz., 10 (251): 271-272.
- Schimkewitsch, W. (1888): Ueber eine von Dr. Korotnew auf den Sunda-Inseln gefundene Pantopoden-Form. - Zool. Jb. (Syst.), 3: 127-133.

Schimkewitsch, W. (1888): Sur les Pantopodes recueillis par M. le lieutenant G. Chierchia, pendant le voyage de la Corvette Vettor Pisani. - Trudy Imp. S.-Petersburgskavo Obschest. Estest., (Otdel. Zool., Phiziol.), 19 (1): 94-96. [in Russian]

Schimkewitsch, W. (1890): Sur les Pantopodes recueillis par M. le lieutenant G. Chierchia, pendant le voyage de la Corvette Vettor Pisani en 1882-1885. - Atti Acad. Lin. 1889, Mem. Classe Sci., Fis., Math. Nat., (4) 6: 329-347, 1 pl.

Schimkewitsch, W. (1890): Sur la collection de Pantopodes du Musée Zoologique de l'Université de Moscou. - Izvest. Imper. Obshchest. Lub. Estest. Anthropol. Ethnogr. Moskva, 67 (2): 20. [in Russian]

Schimkewitsch, W. (1891): Note sur les genres de Pantopodes *Phoxichilus* (Latr.) et *Tanystylum* (Miers). - Arch. Zool. Exp. Gén., (2) 9: 503-522.

Schimkewitsch, W. (1893): Report on the dredging operations off the West Coast of Central America to the Galapagos, to the West Coast of Mexico, and in the Gulf of California, in charge of Alexander Agassiz, carried on by the U.S. Fish Commission Steamer Albatross, during 1891, Lieut. Commander Z.L. Tanner, U.S.N., commanding. Compte rendu sur les Pantopodes recueillis pendant les Explorations de l'Albatross 1891. - Bull. Mus. Comp. Zool., Harvard, 25 (2): 27-43.

Schimkewitsch, W. (1895): Sur les Pantopodes de l'Océan Glacial et de la Mer Blanche. - Trav. Soc. Imp. Nat. St.-Petersbourg, 25 (1): 1-20. [in Russian]

Schimkewitsch, W. (1895): On some forms of Pantopoda. - Trudy Imp. S.-Petersburgs. Obshchest. Estest., 27 (1): 1-14.

Schimkewitsch, W. (1896): On the Pantopoda of the Whote Sea and the Glacial Ocean. - Trudy Imp. S.-Ptersburgs. Obshchest. Estest., 27 (1): 1-14. [in Russian]

Schimkewitsch, W. (1906): Ueber die Periodizität in dem System der Pantopoda. - Zool. Anz., 30 (1, 2): 1-22.

Schimkewitsch, W. (1906): Über die Entwicklung von *Thelyphonus caudatus* (L.) verglichen mit derjenigen einiger anderen Arachniden. - Z. Wiss. Zool., 81 (1): 1-95.

Schimkewitsch, W. (1907): Übersicht der von P. Schmidt und W. Braschnikow in den Ostasiatischen Ufergewässern gesammelten Pantopoden. - Ann. Mus. Zool. Acad. Imp. Sci., St. Pétersbourg, 11 (1906): 246-252.

Schimkewitsch, W. (1907): Zur Pantopoden-Fauna des Sibirischen Eismeeres. - Résultats Scientifiques de l'Expédition polaire Russe en 1900-1903, sous la direction du Baron E. Toll, 1 (6). In: Mém. Acad. Imp. Sci. St. Pétersbourg, (8) 18 (6): 1-10.

Schimkewitsch, W. (1909): Ueber die Pantopoden von St. Vaast-laHouge und Roscoff. - Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersburg, 13 (1908): 427-436.

Schimkewitsch, W. (1909): Nochmals über die Periodicität in dem System der Pantopoden. - Zool. Anz., 34 (1): 1-13.

Schimkewitsch, W. (1913): Ein Beitrag zur Klassifikation der Pantopoden. - Zool. Anz., 41 (13): 507-615.

Schimkewitsch, W. (1913): Einige neue Pantopoden. - Ann. Mus. Zool. Acad. Imp. Sci. St.-Pétersbourg, 18: 240-248.

Schimkewitsch, W. & V. Dogiel (1913): Ueber Regeneration bei Pantopoden. - Bull. Acad. Imp. Sci. St.-Pétersbourg, 1913: 1147-1156.

Schimkewitsch, W. (1918): Pantopoda of the Russian Fauna. - Zool. Vyest. Petrograd, 3: 222-248. [in Russian]

Schimkewitsch, W. (1929): Pycnogonida (Pantopoda). Fauna SSSR, Izdatel. Akad. Nauk. SSSR, part 1: I-CXV, 1-224. [in Russian and Latin]

Schimkewitsch, W. (1930): Pycnogonida (Pantopoda). Fauna SSSR, Izdatel. Akad. Nauk. SSSR, part 2: 225-555. [in Russian and Latin]

Schioedte, J.C. (1875): Udsigt over Grønlands Land Ferskvands og Strandbreds-Arthropoder ?. In: H. Rink, Grønland geografisk og statistisk beskrevet, 2 (3): 50-74 (Pycnogonida: p. 71); Kjøbenhavn.

Schioedte, J.C. (1875): Naturhistoriske Bidrag til en Beskrivelse af Gronland. Kjøbenhavn: Louis Kleins Bogtrykkeri.

Schlottke, E. (1932): Die Pantopoden der deutschen Küsten. - Wiss. Meeresunters. Dtsch. Meere, Abt. Helgoland, (N.F.), 18 (9): 1-10.

Schlottke, E. (1933): Darm und Verdauung bei Pantopoden. - Z. Mikrosk.-Anat. Forsch., Leipzig, 32 (4): 633-658. [anatomy, alimentary system]

Schmidt, H.W. (1971): Die Beeinflussung der Häutungen von *Pycnogonum litorale* (Ström) durch exogene und endogene Faktoren. - Oecologia, 7 (3): 249-261.

Schmidt, H.W. & D. Bückman (1971): Beobachtungen zur Lebensweise von *Pycnogonum litorale* (Ström) (Pantopoda). - *Oecologia*, 7 (3): 242-248.

Schmitt, W.L. (1934): Notes on certain pycnogonids including descriptions of two new species of *Pycnogonum*. - *J. Wash. Acad. Sci.*, 24 (1): 61-70.

Schmitt, W.L. (1945): Miscellaneous zoological material collected by the U.S. Antarct. Service Exped., 1939-1941. - *Proc. Am. Phil. Soc.*, Philadelphia, 89 (1): 297.

Schneider, G. (1884): Crustacea og Pycnogonider indsamlet i Kvaenangsfjorden (Trømsö): ?, 5 pls.

Schneider, S.J. (1884): Undersøgelser af dyrelivet i de arktiske fjorde, II. Crustacea og Pycnogonida indsamlede i Kvärangsfjorden 1881. - *Aarshefter Trømsö Mus.*, 7: 47-132.

Schram, F.R. & J.W. Hedgpeth (1978): Locomotory mechanisms in antarctic pycnogonids. - *Zool. J. Linn. Soc.*, 63 (1-2): 145-169.

Schüller, S. (1989): Die Pantopodenfauna von Rovinj (Nördliche Adria) und der Jahreszyklus einiger Arten. - *Bonn. zool. Beitr.*, 40 (3-4): 285-295; Bonn.

Scott, F.M. (1912): On a species of *Nymphon* from the North Pacific. - *Ann. Mag. Nat. Hist.*, (8) 10: 206-208.

Sekiguchi, K. (1971): Egg-carrying habit and embryonic development in a pycnogonid, *Propallene longiceps*. - *Dōbutsugaku Zasshi*, Tokyo, 80 (4): 137-139. [embryology]

Semper, C. (1874): Ueber Pycnogoniden und ihre in Hydroiden schmarotzenden Larvenformen. - *Arb. Zool.-Zootom. Inst. Würzburg*, 1: 264-286, 2 pls. [hydroid host]

Sergeev, V.N., S.M. Clarke & S.A. Sheperd (1988): Motile macroepifauna of the seagrasses, *Amphibolis* and *Posidonia*, and unvegetated sandy substrata in Holdfast Bay, South Australia. - *Trans. Roy. Soc. S. Austr.*, 112 (3-4): 97-108.

Sharov, A.G. (1966): Basic Arthropodan Stock with special Reference to Insects: 1-271; London: Pergamon Press.

Shigido, I. (1899): Sea spiders occurring in our country. - *Dōzatsu*, 11 (128): 195-200. [in Japanese]

- Shin, P.K.S., M.E. Conneely & B.F. Keegan (1982): Littoral and benthic investigations on the west coast of Ireland - XV: The macrobenthic communities of North Bay (Galway Bay). - Proc. Roy. Ir. Acad., (B) 82 (8): 133-152.
- Shishikura, F. & K. Nakamura (1985): Two types of hemocytes, granular amebocytes and spherule cells, in a pycnogonid, *Propallene longiceps*. - Zool. Sci., 2: 571-581. [circulatory system, cytology]
- Sims, R.W. (1980): Animal identification: A reference guide, Vol. I: Marine and Brackish Water Animals. i-x + 111 pp. London, etc.: John Wiley & Sons.
- Slater, H.H. (1879): On a new genus of Pycnogon and a variety of *Pycnogonum littorale* from Japan. - Ann. Mag. Nat. Hist., (5) 3 (16): 281-283.
- Sloan, N.A. (1979): A Pycnogonid-Ophiuroid association. - Mar. Biol., 52 (2): 171-176. [echinoderm host]
- Smaldon, G., D. Heppell & K.R. Watt (1976): Type specimens of invertebrates (excluding insects) held at the Royal Scottish Museum, Edinburgh. - Roy. Scott. Mus. Inf. Ser., Nat. Hist., 4: IV + 118 pp. [Pycnogonids pp. 58-59]
- Smith, E., G. Chapman, R.B. Clark, D. Nichols & J.D. Carthy (1971): Pycnogonids. In: The Invertebrate Panorama: 30; London: Weidenfeld & Nicolson.
- Smith, S.J. (1873): Pycnogonida. In: Report on the invertebrate animals of Vineyard Sound....: 415, 544. U.S. Comm. Fish & Fish., Washington, 1873.
- Smith, S.J. & O. Harger (1874): Report on the dredging in the region of the St. George's Banks in 1872. - Trans. Conn. Acad. Arts Sci., 3: 1-55.
- Smith, S.J. (1885): In: A.E. Verrill. Results of the explorations made by the steamer Albatross off the northern coast of the United States in 1883. - Rep. U.S. Comm. Fish., 1883: 503-601, 38pls.
- Snodgrass, R.E. (1938): Evolution of the Annelida, Onychophora and Arthropoda. - Smiths. Mis. Coll., 97 (6): 1-159.
- Snodgrass, R.E. (1951): Comparative studies on the head of the mandibulate arthropods; i-vii, 1-118; Ithaca: Comstock Publ. [anatomy]
- Snodgrass, R.E. (1952): Textbook of arthropod anatomy: i-viii, 1-363; New York Cornell Univ. Press.

- Sobre, J.C. (1982): Observaciones preliminares del suprabenthos en un transecto batimetrico de la plataforma continental aquitana (suroeste de Francia). - *Oecol. Aquat.*, 6: 9-17.
- Sokolow, J. (1911): Ueber den Bau der Pantopodenauge. - *Z. Wiss. Zool.*, 98 (3): 339-380. [anatomy, optical sense]
- Soot-Ryen, T. (1925): The Folden Fjord. - *Troms Mus. Skrift.*, 1 (1): 1-11.
- Soot-Ryen, T. (1927): Pycnogonida (of the Folden Fjord, N. Norway). - *Troms Mus. Skrift.*, 1 (5): 19-20.
- Sowinsky, W. (1904): Introduction à l'étude de la faune du bassin marin Ponto-Aralo-Kaspien sous le point de vue d'une province zoo-géographique indépendante. - *Mém. Soc. Nat. Kieff*, 18: I-XIII, 1-487; Suppl. 1: 1-193; Suppl. 2: 197-216.
- Soyer, J. (1966): Sur quelques Pycnogonides du Golfe de Gênes. - *Doriana*, 47 (174): 1-5.
- Spamer, E.E. & A.E. Bogan (1992): General invertebrates collection of the Academy of Natural Sciences of Philadelphia, parts 1 and 2. - *Tryonia*, 26: i-vi, 1-305.
- Spindler, K.D. (1989): Hormonal role of ecdysteroids in Crustacea, Chelicerata and other arthropods; pp. 290-295, in: Koolman, J. (ed.). *Ecdysone: from chemistry to mode of action*. G. Thieme Verl., Stuttgart & New York, 1989: i-xiii, 1-482. [hormones]
- Staples, D.A. (1977): Pycnogonida...the "Nobodies". - *Australia's Oceans*, J. Oceans Soc. Austr., 1 (3): 122-125.
- Staples, D.A. (1979): Three new species of *Propallene* (Pycnogonida: Callipallenidae) from Australian Waters. - *Trans. Roy. Soc. S. Austr.*, 103 (4): 85-93.
- Staples, D.A. (1982): Pycnogonida of the Calliope River & Auckland Creek, Queensland. - *Mem. Qd. Mus.*, 20 (3): 455-471, pl.1; Brisbane.
- Staples, D.A. (1985): Pycnogonida. In: *Museum of Victoria Bass Strait Survey, 1979-1984*, pp. 17-18. Melbourne, Australia; Mus. Victoria.
- Staples, D.A. & J.. Watson (1987): 20. Associations between pycnogonids and hydroids. In: *Modern trends in the systematics, ecology and evolution of hydroids & Hydromedusa*, pp. 215-226. Oxford Univ. Press. [hydroid host]

Stappers, L. (1909): Notes biologiques. In: Duc d'Orleans (ed.). La revanche de la Banquise, Suppl. 4: 256-272, 3 pls; Paris.

Stebbing, T.R.R. (1902-1903): The Nobodies - a sea-faring family. - Knowledge, 25: 37-40, 73-76, 137-140, 185-189; 26: 14-17, 145-148.

Stebbing, T.R.R. (1915): On some enigmatical names in conchology and pycnogonology. - Ann. Mag. Nat. Hist., (8) 16: 328-331.

Stephensen, K. (1912): Report on the Malacostraca, Pycnogonida and some Entomostraca collected by the Danmark expedition to northeast Greenland 1906-1908. - Medd. Grønland, 45: 501-630, 5 pls.

Stephensen, K. (1913): Account of the Crustacea and the Pycnogonida collected by Dr. V. Nordman in the summer of 1911 from northern Strömfjord and Giesecke Lake in West Greenland. - Medd. Grønland, 51 (2): 55-77.

Stephensen, K. (1913): Grønlands Krebsdyr og Pycnogonider (Conspectus Crustaceorum et Pycnogonidorum Groenlandiae). - Medd. Grønland, 52: 382-409.

Stephensen, K. (1917): Zoogeographical investigation of certain fjords in southern Greenland with special reference to Crustacea, Pycnogonida and Echinodermata including a list of Alcyonaria and Pisces. - Medd. Grønland, 53: 231-378.

Stephensen, K. (1918): Grønlands Krebsdyr or Pycnogonider. - Medd. Grønland, 22: 479 pp.

Stephensen, K. (1929): Pycnogonida. In: A.S. Jensen et al. (eds.). Zoology of the Faroes: 1-8.

Stephensen, K. (1933): The Scoresby Committee's 2nd East Greenland Expedition in 1932 to King Christian IX's Land: Crustacea and Pycnogonida. - Medd. Grønland, 104 (15): 1-12.

Stephensen, K. (1933): Pycnogonida. In: The Godthaab Expedition 1928. - Medd. Grønland, 79 (6): 1-46.

Stephensen, K. (1933): Havedderkopper (Pycnogonida) og Rankefødder (Cirripedia). Danmarks Fauna, 38: 1-158.

Stephensen, K. (1936): Pycnogonida from Norway and adjacent waters. - Bergens Mus. Arbok., 1935; Naturvidensk. Rekke, 7: 1-39.

Stephensen, K. (1936): Sveriges pycnogonider. Göteborgs. Kungl. Vetensk.-Vitterhets-Samhälles Handl., (5) (B) 4 (14): 1-56.

- Stephensen, K. (1937): Pycnogonida. The Zoology of Iceland, 3 (58): 1-13.
- Stephensen, K. (1938): Amphipoda, Tanaidacea and Pycnogonida. - Zool. Ergebn. Reise Dr. Kohl Larsen..., 20 (3-4): 236-264.
- Stephensen, K. (1943): Pycnogonida. The Zoology of East Greenland. - Medd. Grønland, 121 (8): 1-41.
- Stephensen, K. (1947): Tanaidacea, Isopoda, Amphipoda and Pycnogonida. Sci. Res. Norw. Antarct. Exped., 1927-1928 et seq., 27: 1-90 (Pycnogonida: 80-86).
- Stimpson, W. (1853): Synopsis of the marine Invertebrata of Grand Manan, or the region round the Bay of Fundy, New Brunswick. - Smiths. Contr. Knowl., 6 (5): 1-67, 3 pls.
- Stimpson, W. (1864): Descriptions of new species of marine Invertebrata from Puget Sound, collected by the naturalists of the North-West Boundary Commission, A.H. Campbell, Esq. Commissioner. - Proc. Acad. Nat. Sci., Philadelphia, 16: 153-161.
- Stock, J.H. (1949): Zeespinnen. - Strand-Werk-Gem. Tabellenser., 7: 1-8.
- Stock, J.H. (1950): The taxonomic status of "*Pycnogonum*" *claudum* Loman, 1908. - Treubia, 20 (3): 627-629.
- Stock, J.H. (1951): 5. Pantopoda. Résultats Scientifiques des Croisières de la Navire-École Belge Mercator. - Mém. Inst. Roy. Sci. Nat. Belg., (2) 43: 1-2.
- Stock, J.H. (1952): *Pycnogonum pusillum* Dohrn, 1881: retrouvé après 70 années sur la côte africaine. - Bull. Inst. Roy. Sci. Nat. Belg., 38 (14): 1-8.
- Stock, J.H. (1952): Revision of the European representatives of the genus *Callippallene* Flynn. - Beaufortia, 1 (13): 1-15.
- Stock, J.H. (1952): The pycnogonids of the lagoon of Venice. - Boll. Soc. Venez. Stor. Nat., 6 (2): 179-186.
- Stock, J.H. (1953): Re-description of some of Helfer's pycnogonid type-specimens. - Beaufortia, 4 (35): 33-45.
- Stock, J.H. (1953): Contribution to the knowledge of the pycnogonid fauna of the East Indian Archipelago. - Biol. Res. Snellius Exped., XVII. - Temminckia, 9: 276-313.
- Stock, J.H. (1954): Four new *Tanystylum* species, and other Pycnogonida from the West Indies. - Stud. Fauna Curaçao and other Caribb. Isl., 5 (24): 115-129.

- Stock, J.H. (1954): Pycnogonida from Indo-West-Pacific, Australian, and New Zealand Waters. - Vidensk. Medd. Dansk Naturhist. Foren., 116: 1-168.
- Stock, J.H. (1955): Pycnogonida from Tristan da Cunha. - Res. Norw. Sci. Exped. Tristan da Cunha, 1937-1938, 33: 1-13.
- Stock, J.H. (1955): Pycnogonida from the West Indies, Central America and the Pacific Coast of North America. Papers from Dr. Th. Mortensen's Pacific Expedition 1914-1916. - Vidensk. Medd. Dansk Naturhist. Foren., 117: 209-266.
- Stock, J.H. (1956): Pantopoden aus dem Zoologischen Museum Hamburg, 1. - Mitt. Zool. Mus. Hamburg, 54: 33-48.
- Stock, J.H. (1956): Verslag van de werkzaamheden van J.H. Stock gedurende zijn verblijf (10 Juli - 9 September 1954) aan het Station Biologique de Roscoff. Verslag der gewone Vergadering van de Afdeling Natuurkunde, Koninklijke Nederlandse Akademie van Wetenschappen te Amsterdam, 65 (9): 138-140.
- Stock, J.H. (1956): Tropical and subtropical Pycnogonida, chiefly from South Africa. - Vidensk. Medd. Dansk. Naturhist. Foren. Kjøbenhavn, 118: 71-113.
- Stock, J.H. (1957): Pycnogonida from the Gulf of Aqaba. - Bull. Sea Fish. Res. Stat., Haifa, 13: 13-14.
- Stock, J.H. (1957): Pantopoden aus dem Zoologischen Museum Hamburg, 2. - Mitt. Zool. Mus. Hamburg, 55: 81-106.
- Stock, J.H. (1957): The pycnogonid family Austrodecidae. - Beaufortia, 6 (68): 1-81.
- Stock, J.H. (1958): The Pycnogonida of the Erythrean and of the Mediterranean coasts of Israel. Contributions to the knowledge of the Red Sea, 5. - Bull. Sea Fish. Res. Stat., Haifa, 16: 3-5.
- Stock, J.H. (1958): The pycnogonid genus *Rhopalorhynchus* Wood-Mason, 1873. - Tijdschr. Ent., 101 (2): 113-137.
- Stock, J.H. (1958): Pycnogonida from the Mediterranean coast of Israel. - Bull. Res. Council Israel (B. Zool.), 7B (3-4): 137-142.
- Stock, J.H. (1959): On some South African Pycnogonida of the University of Cape Town ecological survey. - Trans. Roy. Soc. S. Afr., 35 (5): 549-567.

- Stock, J.H. (1960): *Pycnogonum plumipes* n.sp., nouveau Pacnogonide de la région de Banyuls. - Vie Milieu, 11 (1): 124-126.
- Stock, J.H. (1961): A new pycnogonid, *Endeis holthuisi* n.sp., from New Guinea. - Ent. Ber., Amsterdam, 21 (2): 28-29.
- Stock, J.H. (1962): Some Turkish pycnogonid records. - Ent. Ber., Amsterdam, 22 (11): 218-219.
- Stock, J.H. (1962): Second list of Pycnogonida of the University of Cape Town ecological survey. - Trans. Roy. Soc. S. Afr., 36 (4): 273-286.
- Stock, J.H. (1963): South African deep-sea Pycnogonida, with descriptions of five new species. - Ann. S. Afr. Mus., 46 (12): 321-340.
- Stock, J.H. (1964): Report on the Pycnogonida of the Israel South Red Sea Expedition. - Bull. Sea Fish. Res. Stat., Haifa, 35: 27-34.
- Stock, J.H. (1964): Deep-sea Pycnogonida collected by the "Cirrus" in the Northern Atlantic. - Beaufortia, 11 (135): 45-52.
- Stock, J.H. (1965): Pycnogonida from the southwestern Indian Ocean. - Beaufortia, 13 (151): 13-33.
- Stock, J.H. (1965): Pycnogonides provenant de la base Antarctique Belge. - Bull. Inst. Roy. Sci. Nat. Belg., 41 (31): 1, 2.
- Stock, J.H. & J. Soyer (1965): Sur quelques Pycnogonides rares de Banyuls-sur-Mer. - Vie Milieu, 16 (1B): 415-421.
- Stock, J.H. (1966): 4. Pycnogonida. Campagne de la Calypso au large des côtes atlantiques de l'Amerique du sud (1961-62). - Rés. Sci. Campagne Calypso, 7. In: Ann. Inst. Océanogr. Monaco, 44: 385-406.
- Stock, J.H. (1966): Sur quelques Pycnogonides de la région de Banyuls. 3ème Note. - Vie Milieu, 17 (1B): 407-417.
- Stock, J.H. (1966): Pycnogonida from West Africa. Scientific results of the Danish expedition to the coasts of tropical West Africa, 1945-46. - Atlantide Rep., 9: 45-57.
- Stock, J.H. (1968): Pycnogonides. Faune marine des Pyrénées-Orientales. - Vie Milieu, 19 (1A), Suppl.: 1-38.
- Stock, J.H. (1968): Pycnogonida collected by the Galathea and Anton Bruun in the Indian and Pacific Oceans. - Vidensk. Medd. Dansk naturhist. Foren., 131: 7-65.

Stock, J.H. (1970): *Endeis flaccida* Calman, 1923, in Florida: a pycnogonid new to the Atlantic Ocean. - Ent. Ber., Amsterdam, 30 (1): 3-4.

Stock, J.H. (1970): The Pycnogonida collected off northwestern Africa during the cruise of the "Meteor". - Meteor Forsch.-Ergebn., (D) 5: 6-10.

Stock, J.H. (1970): A new species of *Endeis* and other pycnogonid records from the Gulf of Aqaba. - Bull. Zoöl. Mus., Univ. Amsterdam, 2 (1): 1-4.

Stock, J.H. (1971): Pycnogonides récoltés durant la campagne "Noratlante" en Atlantique Nord. - Bull. Zoöl. Mus., Univ. Amsterdam, 2 (4): 25-28.

Stock, J.H. (1971): Pycnogonides. In: P. Grua (Ed.). Invertébrés de l'infra-littoral rocheux dans l'archipel de Kerguelen: 29. Com. Natn. française Rech. antarct., 30.

Stock, J.H. (1971): A new deep-sea Nymphon (Pycnogonida) collected by the Galathea off Kenya. - Zool. Meded., Leiden, 47: 257-260.

Stock, J.H. (1973): Pycnogonida from south-eastern Australia. - Beaufortia, 20 (266): 99-127.

Stock, J.H. (1973): *Achelia shepherdii* n.sp. and other Pycnogonida from Australia. - Beaufortia, 21 (279): 91-97.

Stock, J.H. (1973): A re-description of the pycnogonid *Pallenopsis boehmi* Schimkewitsch, 1930 based on the original material from Strait Magellan. - Netherl. J. Zool., 23 (3): 347-352.

Stock, J.H. (1974): Medio- and infralittoral Pycnogonida collected during the I.I.O.E. near the landbase of Nossi-Bé, Madagascar. - Bull. Zoöl. Mus., Univ. Amsterdam, 4 (3): 11-22.

Stock, J.H. (1975): Pycnogonida from the continental shelf, slope, and deep sea of the tropical Atlantic and East Pacific. - Biological results of the University of Miami deep-sea expedition, 108. - Bull. Mar. Sci., 24 (4): 957-1092.

Stock, J.H. (1975): The pycnogonid genus *Propallene* Schimkewitsch, 1909. - Bull. Zoöl. Mus., Univ. Amsterdam, 4 (11): 89-97.

Stock, J.H. (1975): Pycnogonida found on fouling panels from the East and West coast of America. - Ent. Ber., Amsterdam, 35 (5): 70-77.

- Stock, J.H. (1976): Infralittoral Pycnogonida from Tanzania. - Trav. Mus. Hist. Nat. "Gr. Antipa", 16: 127-134.
- Stock, J.H. (1978): Abyssal Pycnogonida from the North-eastern Atlantic Basin, part I. - Cah. Biol. Mar., 19: 189-219.
- Stock, J.H. (1978): Abyssal Pycnogonida from the North-eastern Atlantic Basin, part II. - Cah. Biol. Mar., 19: 397-413.
- Stock, J.H. (1978): Experience on food preference and chemical sense in pycnogonids. In: Sea Spiders (Pycnogonida). - Zool. J. Linn. Soc. Lond., 63 (1-2): 59-74.
- Stock, J.H. (1979): Pycnogonida from the mediolittoral and infralittoral zones in the tropical Western Atlantic. - Stud. Fauna Curaçao and other Caribb. Isl., 59 (184): 1-32; Amsterdam.
- Stock, J.H. (1979): *Anoplodactylus ophiurophilus* n.sp., a sea spider associated with brittle stars in the Seychelles. - Bijdr. Dierk., 48 (2): 156-160. [echinoderm host]
- Stock, J.H. (1979): Pycnogonida from the mediolittoral and infralittoral zones in the tropical Western Atlantic. - Stud. Fauna Curaçao and other Caribb. Isl., 59 (184): 1-32.
- Stock, J.H. (1980): *Anoplodactylus xenus* n.sp. (Pycnogonida), a species from New Zealand confused with *A. haswelli* (Flynn, 1918). - Steenstrupia, 6 (1): 1-4.
- Stock, J.H. (1981): 13: Pycnogonides, I.: *Pycnosoma asterophila*, a sea spider associated with the starfish *Calliaster* from the Philippines. In: Rés. Campagnes MUSORSTOM, Paris, 91: 309-312. [echinoderm host]
- Stock, J.H. (1981): 14: Pycnogonides, II.: Description de *Nymphon macilentum* sp. nov. In: Rés. Campagnes MUSORSTOM, Paris, 91: 313-315.
- Stock, J.H. (1981): Abyssal Pycnogonida from the Walvis Basin, Southern Atlantic. - Cah. Biol. Mar., 22: 453-471.
- Stock, J.H. (1982): Researches on the coast of Somalia. Shallow-water Pycnogonida. - Monit. zool. ital., N.S. Suppl. 17 (7): 183-190;
- Stock, J.H. (1983): Pycnogonides des campagnes CORINDON et MUSORSTOM II (detroit de Makassar et Philippines), avec description de *Pallenopsis dentifera* sp. nov. - Bull. Mus. natn. Hist. nat., Paris, (4) 5 (sec. A) (1): 299-305.

Stock, J.H. (1984): Deep-water Pycnogonida of the INCAL, BIOGAS, GEOMANCHE and SAFARI cruises. - *J. mar. Biol. Ass. U.K.*, 64: 743-749.

Stock, J.H. (1984): The deep-water Pycnogonida of the Safari Cruises to the Indian Ocean. - *Bull. Mus. natn. Hist. nat.*, Paris, (4) 6 (sec.A) (3): 701-709.

Stock, J.H. (1985): Deux pycnogonides rares, récoltés pendant la campagne CORINDON II dans le détroit de Makassar, Indonésie. - *Bull. Zoöl. Mus., Univ. Amsterdam*, 10 (18): 153-158.

Stock, J.H. (1986): Pycnogonida (= Pantopoda). In: *Stygofauna Mundi: A Faunistic, distributional, and ecological synthesis of the World Fauna inhabiting Subterranean Waters (including the Marine Interstitial)*, pp. 698-700. Leiden: E.J. Brill/Dr. W. Backhuys.

Stock, J.H. (1986): Pycnogonida from the Caribbean and the Straits of Florida. - *Bull. mar. Sci.*, 38 (3): 399-441; Miami.

Stock, J.H. (1987): Faunistic transit between the Atlantic and the Mediterranean: the deep-water Pycnogonida. - *Cah. Biol. Mar.*, 28: 505-519; Roscoff.

Stock, J.H. (1989): Pycnogonida collected in interstitia of coral sand and rubble in the Netherlands Antilles. - *Bijdr. Dierk.*, 59 (2): 87-96; Den Haag.

Stock, J.H. (1990): Macaronesian Pycnogonida. CANCAP-project. Contribution no.78. - *Zool. Meded.*, 63 (16): 205-233; Leiden.

Stock, J.H. (1990): A new species of *Austrodecus* (Pycnogonida) from New South Wales, Australia. - *Tijdschr. Ent.*, 133 (2): 269-272.

Stock, J.H. (1991): Pycnogonides de la campagne Seamount 1 au large de la péninsule ibérique et dans le golfe ibéro-marocain. - *Bull. Mus. natn. Hist. nat.*, Paris. 4e sér., 13, sec. A, (1-2): 135-142.

Stock, J.H. (1991): Deep-water Pycnogonida from the surroundings of New Caledonia. In: A. Crosnier (ed.). *Résultats des Campagnes MUSORSTOM*, Vol. 8. - *Mém. Mus. natn. Hist. nat.*, a (151): 125-212.

Stock, J.H. (1991): Pycnogonida of the MUSORSTOM campaigns to the Philippines. In: A. Crosnier (ed.). *Résultats des Campagnes MUSORSTOM*, Vol. 8. - *Mém. Mus. natn. Hist. nat.*, a (151): 213-228.

Stock, J.H. (1992): Pycnogonida from southern Brazil. - Tijdschr. Ent., 135: 113-139.

Størmer, L. (1944): On the relationships and phylogeny of fossil and recent Arachnomorpha. - Skrift. utgitt Norske Vidensk.-Akad. Oslo, Math.-Naturvidensk. Klasse, 1944 (5): 1-158.

Storrow, B. (1912): Material collected during July and August 1911, when on board trawlers fishing on the local prawn ground, together with the catches of edible fish then made. - Rep. Dove Mar. Lab., Cullercoats, (N.S.), I: 68-70.

Straughan, J.H. & R.W. Klink (1980): A taxonomic listin of common marine invertebrate species from southern California, USA. - Allan Hancock Found., Tech. Rep., 1980, (3): I-IV, 1-281.

Ström, H. (1762): Physik og Oeconomisk Beskrevelse over Fogderiet Søndmør, beliggende i Bergens Stift i Norge, 1: 208-209.

Studder, T. (1889): Die Forschungsreise S.M.S. Gazelle in den Jahren 1874 bis 1876...III. Zoologie und Geologie, i-vi, 1-322; Berlin: E.S. Mittler & Sohn.

Stuxberg, A. (1887): Vega-Expeditionens vetenskapliga Jaktagelser, 5.

Sumner, F.P., R. Osburn & L.J. Cole (1913): A biological survey of the waters of Woods Hole and vicinity. - Bull. Bur. Fish., Washington, 31 (1-2): 1-860.

Suzuki, S. (1963): Pycnogonida. In: Fauna of the Coast of Shonai and the adjacent Sea: 12. Publ. by author, Yamagata. [in Japanese]

Taramelli, E., C. Chimenz & L. Blundo (1981): Contributo delle zoocenosi di Torvaldaliga. - Quad. Lab. Tech. Pesc., Ancona, 3 (suppl.), 1: 535-550.

Templeton, R. (1836): Catalogue of Irish Crustacea, Myriapoda and Arachnida, selected from the papers of the late John Templeton, Esq. - Mag. Nat. Hist. & J. Zool. & Bot., 9: 9-14.

Terry, L.A. & D. Sell (1986): Rocky shores in the Moray Firth. - Proc. Roy. Soc., Edinburgh, (B) 91: 169-191.

Tesch, J.J. (1910): V. Pycnogonida, verzameld met de Wodan. Bijdragen tot de fauna der Zuidelijke Noordzee. Jaarb. Rijksinst. Onderzoek Zee, 1909 (4): 43-64.

Thiel, M.E. (1976): Wirbellose Meerestiere als Parasiten, Kommensalen oder Symbionten in oder an Scyphomedusen. - Helgol. Wiss. Meeresunters., 28: 417-446.

Thomassin, A. (1974): Soft bottomcarcinological fauna sensu lato on Tuléar coral reef complexes (S.W. Madagascar) ... - Proc. 2nd. Int. Coral Reef Symp., 1: 297-320; Brisbane: Great Barrier Reef Committee.

Thompson, C.W. (1883): Pycnogonida. In: The Depths of the Sea: 129; London.

Thompson, W. (1844): Report on the Fauna of Ireland. Div. Invertebrata. Proc. 13th Meeting Brit. Ass. Adv. Sci., 1843: 245-290.

Thomson, G.M. (1883): New Zealand Pycnogonida of the "Challenger". - N.Z. J. Sci., 1: 28.

Thomson, G.M. (1884): On the New Zealand Pycnogonida, with descriptions of new species. - Trans & Proc. N.Z. Inst., 16 (1883): 242-248.

Tjønneland, A., H. Kryvi, J.P. Ostnes & S. Okland (1985): The heart ultrastructure of two species of Pycnogonids, and its phylogenetic implications. - Zool. Scr., 14 (3): 215-219. [cytology]

Topsent, E. (1889): Notice concernant les Pantopodes de Luc. - Bull. Soc. Lin. Normandie, Caen, (4) 2 (1888-1889): 60-63.

Topsent, E. (1891): Les Pycnogonides provenant des campagnes du yacht "l'Hirondelle". - Bull. Soc. Zool. France, 16: 176-180.

Topsent, E. (1897): Pycnogonides recueillis par le yacht "Princesse-Alice". - Bull. Soc. Zool. France, 22: 106-107.

Topsent, E. (1938): Les Pycnogonides provenant des campagnes du yacht "l'Hirondelle", 1886-1887-1888 (Golfe de Gascogne, Terre Neuve, Açores). - Rés. Camp. Sci. Monaco, 97: 272-276. [reprint of Topsent, 1891]

Topsent, E. (1938): Pycnogonides recueillis par le yacht "Princesse-Alice". - Res. Camp. Sci. Monaco, 97: 276-277. [reprint of Topsent, 1897]

Totland, G.K. & H. Kryvi (1986): The fine structure of the somatic muscles and their attachment to the cuticle in two species of Pycnogonida. - Zool. Scr., 15: 69-72.

Tucker, M.J. (1988): Temporal distribution and brooding behaviour of selected benthic species from the shallow marine waters off the Vestfold Hills, Antarctica. In: J.M. Ferris, H.R. Burton, G.W. Johnstone & I.A.E. Bayly (eds.). Biology of the Vestfold Hills, Antarctica. - Hydrobiologia, 165: 151-159.

Turpaeva, E.P. (1955): New species of sea spiders (Pantopoda) from the Kurile-Kamchatka trench. - Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, 12: 323-327. [in Russian]

Turpaeva, E.P. (1956): Pantopoda of the genus *Heteronymphon* from the northwest part of the Pacific Ocean. - Byull. Moskv. Obshchest. Ispyt. Prir., Otdel Biol., 61 (2): 67-72.

Turpaeva, E.P. (1957): New species of seaspiders (Pantopoda) of the genus *Pallenopsis* from the north-western Pacific Ocean. - Trudy Inst. Okeanol. P.P. Shirshov Akad. Nauk SSSR, 27: 356-361.

Turpaeva, E.P. (1969): Morskie pauhi (Pycnogonida). In: V.G. Kort (ed.). Biologija Tikhogo okeana (Biology of the Pacific Ocean), part 2, Glubokavod naia donna fauna pleuston (Deep-sea bottom fauna), pp. 63-66. Moscow: Akademiia Nauk, SSR, Institut Okeanologii.

Turpaeva, E. (1970): Belonging of *Nymphon abyssale* Stock to the genus *Heteronymphon*. - Zool. Zh., 49 (11): 1723-1725. [in Russian, English title]

Turpaeva, E. (1971): The deepwater Pantopoda collected in the Kurile-Kamchatka trench. In: Fauna of the Kurile-Kamchatka Trench. - Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, 92: 274-291.

Turpaeva, E. (1971): The genus *Ascorhynchus* (Pantopoda) in the deep-sea fauna of the Pacific Ocean. - Byull. Mosk. Obshchest. Ispyt. Prirody, Otdel Biol., 76 (3): 104-110. [in Russian, English title]

Turpaeva, E. (1973): Pantopoda from the Northeast Pacific Ocean. - Trudy Inst. Okeanol. P.P. Shirshov., Akad. Nauk SSSR, 91: 178-191. [in Russian]

Turpaeva, E. (1974): The Pycnogonids of the Scotia Sea and surrounding waters. - Trudy Inst. Okeanol. P.P. Shirshov, Akad. Nauk SSSR, 98: 277-305. [in Russian]

Turpaeva, E. (1975): On some deep-water pantopods (Pycnogonida) collected in north-western and south-eastern Pacific. - Trudy Inst. Okeanol P.P. Shirshov, Akad. Nauk SSSR, 103: 230-246.

Turpaeva, E.P. (1988): The finding of Pycnogonida in Hydrothermal Fauna. - Zool. Zh., 67 (6): 950-953.

Turpaeva, E.P. (1989): Some morphological features of the deep-sea Pycnogonida. - Trudy Inst. Okeanol., Akad. Nauk SSSR, 124: 127-133.

Turpaeva, E.P. (1990): Pycnogonida from southern Atlantic. - Trudy Inst. Okeanol., Akad. Nauk SSSR, 126: 108-126.

Turpaeva, E.P. (1990): Pycnogonida collected on the Pacific slope of the Hondo Island. - Zool. Zh., 69 (9): 17-26.

Turpaeva, E.P. (1991): Pantopods (Pycnogonida) from the shelf of south-eastern Africa and surrounding waters. - Zool. Zh., 70 (12): 33-43.

Turpaeva, E.P. (1991): *Anoplodactylus gibbefemoris* sp.n. (Pycnogonida, Phoxichilidiidae) from the underwater mountains. - Zool. Zh., 70 (3): 151-154.

Uchida, H., K. Konno & S. Chiba (1970): Pycnogonida. In: A preliminary list of marine animals found in the vicinity of Fukaura Marine Biological Laboratory. II. - Rep. Fukaura Mar. Biol. Lab., 2: 7-29. [in Japanese]

Uhlmann, D. & W. Dunger (1955): Biologische Beobachtungen auf einer Fangreise des Loggers "Anton Saefkow" im Barentsmeer im Juli-August 1952. - Wiss. Z. Karl-Marx Univ., Leipzig, 4 (1954/55) (Math.-naturwiss. Reihe (1-2)): 51-71.

Ushakob, P.V. (1951): On the marine fauna of the region of the Kuril Islands. - Dokl. Akad. Nauk SSSR, 80 (1): 125-128. [in Russian]

Ushakob, P.V. (1953): Pycnogonida. In: Fauna of the Sea of Okhotsk and its natural resources. - Izdat. Zool. Inst. Akad. Nauk SSSR: 1-459. [in Russian]

Utinomi, H. (1950): Pycnogonida from the Kii Province (Records of collections by dredging from off Minabe, Kii Province, I.). - Nanki Seibutu, Wakayama, 1 (3): 127-132. [in Japanese]

Utinomi, H. (1951): On some pycnogonids from the sea around Kii Peninsula. - Publ. Seto Mar. Biol. Lab., 1 (4): 159-168.

Utinomi, H. (1954): The fauna of Akkeshi Bay. XIX. Littoral Pycnogonida. - Publ. Akkeshi Mar. Biol. Stat., 3: 1-28.

Utinomi, H. (1955): Report on the Pycnogonida collected by the Sôyô-Maru Expedition made on the continental shelf bordering Japan during the years 1926-30. - Publ. Seto Mar. Biol. Lab., 5 (1): 1-42.

- Utinomi, H. (1958): Pycnogonida. In: Coloured Illustrations of Sea Shore Animals of Japan: 47-48; Osaka: Hoikusha.
- Utinomi, H. (1959): Pycnogonida of Sagami Bay. - Publ. Seto Mar. Biol. Lab., 7 (2): 197-222.
- Utinomi, H. (1959): Pycnogonida of the Japanese Antarctic Research Expeditions 1956-1958. Biological Results of the Japanese Antarctic Research Expedition. 8. Spec. Publ. Seto Mar. Biol. Lab.: 1-12. [= Contr. Seto Mar. Biol. Lab., 341]
- Utinomi, H. (1962): Pycnogonida of Sagami Bay - Supplement. - Publ. Seto Mar. Biol. Lab., 10 (1): 91-104.
- Utinomi, H. (1965): Pycnogonida. In: New illustrated Encyclopaedia of the Fauna of Japan, 2: 333-338; Tokyo: Hokoryû-kan. [in Japanese]
- Utinomi, H. (1966): Subphylum Pycnogonida. - Dôbutsu Keitôbunruigaku, 7 (B): 1-14; Tokyo: Nakayamashoten. [in Japanese]
- Utinomi, H. (1971): Records of Pycnogonida from shallow waters of Japan. - Publ. Seto Mar. Biol. Lab., 18 (5): 317-347.
- Utinomi, H. (1971): *Nymphon chaetochir* n.sp., a new deepsea pycnogonid from Chatham Rise, east of New Zealand. - J. Roy. Soc. N.Z., 1 (3-4): 197-202.
- Vachon, M. (1963): Pycnogonida. In, Chélicerates. Encyclopédie de la Pléiade, Zoologie, 2: 244-253.
- Vader, W. (1975): The sea anemone *Bolocera tuediae*, and its copepod parasite, *Antheacheres duebeni*, in northern Norway. - Astarte, 8 (1): 37-39.
- van Beneden, P.-J. (1845): Recherches sur la circulation dans quelques animaux inférieurs. - Bull. Acad. Roy. Bruxelles, 12 (5): 1-19.
- van Deurs, B. (1973): Axonemal 12 + 0 pattern in the flagellum of the motile spermatozoan of *Nymphon leptochaetes*. - J. Ultrastruct. Res., 42: 594-598. [cytology, genital system]
- van Deurs, B. (1974): Pycnogonid sperm. An example of inter- and intraspecific axonemal variation. - Cell Tissue Res., 149: 105-111. [cytology, genital system]
- van Deurs, B. (1974): Spermatology of some Pycnogonida (Arthropoda), with special reference to a microtubule-nuclear envelope complex. - Acta Zool., 55: 151-162. [cytology, genital system]

- van der Hammen, L. (1978): The evolution of the Chelicerate life-cycle. - *Acta Biotheoret.*, 27 (1/2): 44-60.
- Verrill, A.E. (1873): Brief contribution to zoology... Results of recent dredging expeditions off the coast of New England. - *Am. J. Sci. & Arts*, (3) 6: 431-435.
- Verrill, A.E. (1873): Report upon the invertebrate animals of Vineyard Sound and the adjacent waters... - *Rep. U.S. Comm. Fish.*, 1: 295-778.
- Verrill, A.E. (1874): Results of recent dredging expeditions on the coast of New England. - *Am. J. Sci. & Arts*, 3 (7): 38-46.
- Verrill, A.E. (1875): Results of dredging expeditions off the New England coast in 1874. - *Am. J. Sci. & Arts*, 3 (10): 36-43.
- Verrill, A.E. (1881): Notice of the remarkable marine fauna occupying the outer banks of the southern coast of New England, no. 2. - *Am. J. Sci. Arts*, ser. 3: 292-303.
- Verrill, A.E. (1881): Recent papers on the marine invertebrates of the Atlantic Coast of North America. - *Am. J. Sci. Arts*, ser. 3, 22: 411-415.
- Verrill, A.E. (1882): Notice of the remarkable marine fauna occupying the outer banks off the southern coast of New England, no. 7, and of some additions of the fauna of Vineyard Ground. - *Am. J. Sci. Arts*, ser. 3, 24: 360-371.
- Verrill, A.E. (1885): Results of the explorations made by the steamer Albatross off the northern coast of the United States in 1883. - *Rep. U.S. Comm. Fish.*, 1883: 503-601.
- Verrill, A.E. (1900): Additions to the Crustacea and Pycnogonida of the Bermudas. - *Trans. Conn. Acad. Arts & Sci.*, 10 (2) (15): 580-582.
- Versluys, J. & R. Demoll (1920): De verwantschap der Merostomata met de Arachnida en met andere klassen der Arthropoden (eerste mededeling). Verslag van de gewone Vergadering der Wis- en Natuurkundige Afdeling der Koninklijke Akademie van Wetenschappen te Amsterdam, 29: 448-460.
- Versluys, J. & R. Demoll (1921): Die Verwandtschaft der Merostomata mit den Arachniden und den anderen Abteilungen der Arthropoda. - *Proc. Koninklijke Akad. Wetenschapen Amsterdam*, 23 (5): 739-765.

- Visweswara, R.V. & K. Hanamatha (1975): Macro- and microfaunal associations of the fouling Dresseinid *Mytilopsis sallei* (Reclus) in the Visakhapatnam Harbor. - Bull Dep. Mar. Sci., Univ. Cochin, 7 (3): 623-629.
- von Dencker, D. (1974): Das Skelettmuskelssystem von *Nymphon rubrum* Hodge, 1862 (Pycnogonida: Nymphonidae). - Zool. Jb. Anat., 93: 272-287. [anatomy]
- von Duben, M.W. (1845): Über die Meeresfauna Norwegens. - Arch. Skand. Beitr. Naturgeschaft, 1: 136-138.
- von Lendenfeld, R. (1833): Die Larvenentwicklung von *Phoxichilidium plumularie* nov. sp. - Z. wiss. Zool., 38: 323-329. [embryology, development]
- von Lotz, G. (1968): Nahrungsaufnahme und Beutefang bei einem Pantopoden, *Anoplodactylus petiolatus* Krøyer. - Oecologia (Berlin), 1: 171-175. [feeding]
- von Lotz, G. & D. Bückmann (1968): Die Häutung und die Exuvie bei *Pycnogonum littorale* (Ström) (Pantopoda). - Zool. Jb. (Anat.), 85: 529-536. [anatomy, development]
- von Salvini-Plawen, L. (1972): Cnidaria as food-sources for marine invertebrates. - Cah. Biol. Mar., 13: 385-400. [feeding]
- Waddington, H.J. (1914): Marine fauna: Pycnogonida. In: Morris, The Natural History of Bournemouth and District: 227.
- Walker, A.O. (1901): Contributions to the malacostracan fauna of the Mediterranean. - J. Linn. Soc. Lond., (Zool.), 28: 290-306.
- Warner, G.F. (1971): On the ecology of a dense bed of the brittle-star *Ophiothrix fragilis*. - J. Mar. Biol. Ass. U.K., 51 (2): 267-282.
- Warren, E. (1908): On a collection of hydroids, mostly from the Natal Coast. - Ann. Natal Mus., 1 (3): 269-355, pls. 45-48.
- Warren, R. (1892): *Nymphon* and *Caprella* on the coast of Sligo. - The Zoologist, (3) 16 (190): 367-368.
- Weygoldt, P. & H.F. Paulus (1979): Untersuchungen zur Morphologie, Taxonomie und Phylogenie der Chelicerata, I: Morphologische Untersuchungen. - Z. Zool. Syst. Evolutionsforsch., 17 (2): 85-116.

- Weygoldt, P. & H.F. Paulus (1979): Untersuchungen zur Morphologie, Taxonomie und Phylogenie der Chelicerata, II: Cladogramme und die Entfaltung der Chelicerata. - Z. Zool. Syst. Evolutionsforsch., 17 (3): 177-200.
- Weygoldt, P. (1990): Arthropoda-Chelicerata: sperm transfer; pp. 77-119, in: Adiyodi, K. & R.G. Adiyodi (eds.). Reproductive biology of invertebrates, vol. 4, part B. Fertilization, development and parental care. J. Wiley & Sons, Chichester, New York, etc.: i-xxiii, 1-527. [mating]
- White, A. (1847): Descriptions of new or little-known Crustacea in the collection of the British Museum. - Proc. Zool. Soc. Lond., 15: 118-126.
- Whiteavers, J.F. (1872): Notes on a deepsea dredging-expedition round the island of Anticosti in the Gulf of St. Lawrence. - Ann. Mag. Nat. Hist., ser. 4, 10: 341-354.
- Whitelegge, T. (1889): List of the marine and fresh-water invertebrate fauna of Port Jackson and the neighborhood. - J. Proc. Roy. Soc. N.S. Wales, 23 (2): 163-325.
- Whitten, H.L., F. Rosene & J.W. Hedgpeth (1950): The invertebrate fauna of Texas coast jetties; a preliminary survey. - Publ. Inst. mar. Sci., Univ. Texas: 53-87.
- Wicksten, M.K. (1980): Crustacea and Pycnogonida. - Tech. Rep. Allan Hancock Fdn., 3: 196-223.
- Wicksten, M.K. (1984): Distributions of some common Decapod Crustaceans and a Pycnogonid from the continental shelf of Northern California. - Calif. Fish Game, 70 (3): 132-139.
- Wieser, W. (1952): Investigations on the microfauna inhabiting seaweeds on rocky coasts. IV. Studies on the vertical distribution of the fauna inhabiting seaweeds below the Plymouth Laboratory. - J. Mar. Biol. Ass. U.K., 31 (1): 145-174.
- Williams, G. (1933): On *Nymphopsis acinacispinatus*, a new Pycnogonid from Queensland. - Ann. Mag. Nat. Hist., ser. 10 (12): 173-180.
- Williams, G. (1940): Contributions to the fauna of Rottnest Island, II. In: Pycnogonida of Western Australia. - J. Roy. Soc. W. Australia, 25: 197-205.
- Williams, G. (1941): A revision of the genus *Anoplodactylus* with a new species from Queensland. - Mem. Qd. Mus., 12: 33-39.
- Williams, G. (1954): Fauna of Strangford Lough and neighbouring coasts. - Proc. Roy. Irish Acad., (B) 56 (3): 29-133.

- Wilson, E.B. (1878): Descriptions of two new genera of Pycnogonida. - Am. J. Sci., 15: 200-203.
- Wilson, E.B. (1878): Synopsis of the Pycnogonida of New England. - Trans. Conn. Acad. Arts & Sci., 5: 1-26.
- Wilson, E.B. (1880): Synopsis of the Pycnogonida of New England. Rep. U.S. Fish. Comm., 6: 465-506. [almost identical with Wilson, 1878]
- Wilson, E.B. (1880): Report on the Pycnogonida of New England and adjacent waters. - Rep. U.S. Comm. Fish., 1878 (6): 463-506, 7 pls.
- Wilson, E.B. (1881): Report on the Pycnogonida. Reports on the results of dredging... "Blake". - Bull. Mus. Comp. Zool., Harvard, 8 (12): 239-256.
- Winter, G. (1980): Beiträge zur Morphologie und Embryologie des vorderen Körperabschnitts (Cephalosoma) der Pantopoda Gerstaecker, 1863. - Z. Zool. Syst. Evolutionsforsch., 18 (1): 27-61.
- Wirén, E. (1918): Zur Morphologie und Phylogenie der Pantopoden. - Zool. Bidr. Uppsala, 6: 41-181.
- Wolff, T. (1958): Notiser om danske havedderkopper. - Flora og Fauna, 64 (1): 1-7.
- Wolff, T. (1960): Strejflys over dybhavets dyreliv. - Naturens Verden, 1960 (June): 161-191.
- Wolff, T. (1961): Animal life from a single abyssal trawling. - Galathea Rep., 5: 129-162.
- Wolff, T. (1970): The concept of the hadal or ultraabyssal fauna. - Deep-Sea Res., 17: 983-1003.
- Wolff, T. (1971): Krebsdyr og Havedderkopper. In: Rosenskilde og Bagger: Danmarks Dyreverden, pp. 97-181.
- Wolff, W.J. (1976): Distribution of Pantopoda in the estuarine area in the southwestern part of the Netherlands. - Netherl. J. Sea Res., 10 (4): 472-478.
- Wolff, W.J. & N. Dankers (1981): Preliminary checklist of the Zoobenthos and nekton species of the Wadden Sea. In: N. Dankers, H. Kuhl & W.J. Wolff (eds.). Invertebrates of the Wadden Sea, pp. 24-60. Leiden: Wadden Sea Working Group.
- Wood-Mason, J. (1873): On *Rhopalorhynchus Kröyeri*, a new genus and species of Pycnogonida. - J. As. Soc. Bengal., 42 (2): 171-175.

Wood-Mason, J. & A. Alcock (1891): On the results of the deep-sea dredging during the season 1890-91. Ser. II., no. 1 of Natural History Notes from H.M. Indian Marine Survey Steamer Investigator. - Ann. Mag. Nat. Hist., (6) 8: 16-17, 269-286, 353-362.

Wright, T.S. (1863): On the development of Pycnogon larvae within the polyps of *Hydractinia echinata*. - Quart. J. Microsc. Sci., (N.S.) 3: 51-52.

Wright, T.S. (1863): On the development of Pycnogon larvae within the polyps of *Hydractinia echinata*. - Proc. Roy. Soc. Edinburgh, 2: 439-442. [hydroid host]

Wyer, D., P.E. King & J.H. Jarvis (1971): *Achelia simplex* (Giltay) a pycnogonid new to the Irish Fauna. - Irish Nat. J., 17 (3): 92-95.

Wyer, D. (1972): Aspects of the nutritional biology of pycnogonids. Ph.D. thesis, Univ. Wales.

Wyer, D. & P.E. King (1973): Relationship between some British littoral and sublittoral bryozoans and pycnogonids. 18. In: G.P. Larwood (ed.). Living and fossil Bryozoa: Recent advances in Research: 199-207.

Wyer, D. & P.E. King (1974): Feeding in British littoral Pycnogonids. - Estuarine cstl. mar. Sci., 2 (2): 177-184.

Wyer, D. & P.E. King (1974): An investigation into the validity of the species *Achelia hispida* Hodge and *Achelia longipes* (Hodge). - Irish Nat. J., 18 (4): 114-120.

Yamashita, H. (1980): Studies on the benthic organisms collected from the East China Sea and the Yellow Sea, IX: Distribution of Sipunculoidea, Annelida, Tentaculata, Arthropoda and Mollusca. - Bull. Seikai Reg. Fish. Res. Lab., 55: 1-11. [in Japanese with English abstract]

Yonge, C.M. (1966): Pycnogonids, sea spiders. In: The sea shore: 25, 82, 105; London: Collins.

Zago, M.S.A. (1970): Sobre o Pantópodo *Rhynchothorax mediterraneus* Costa, 1861. - Contr. Avuls. Inst. Oceanogr., Univ. S. Paulo, 21: 1-5.

Zambrana Diaz, M., M.I. García Palmer, O. Rivero Bustos & T. Eligio Castillo (1985): Lista de especies y bibliografía de los picnogónidos (Arthropoda) del Mediterráneo Americano. - Revta. Invest. Mar., 6 (1): 29-38.

- Zavodnik, D. (1965): A contribution to the knowledge of the community *Cystoseira barbata* in the North Adriatic. - Biol. Vestnik, 13: 87-101. [in Serbo-Croatic]
- Zavodnik, D. (1968): Beitrag zur Kenntnis der Asselspinnen (Pantopoda) der Umgebung von Rovinj (Nördl. Adria). - Thalasia Jugoslavica, 4: 45-53.
- Zavodnik, D. (1971): Contribution to the dynamics of the benthic communities in the region of Rovinj (northern Adriatic). - Thalasia Jugoslavica, 7 (2): 447-514.
- Zenker, W. (1852): Untersuchungen über die Pycnogoniden. - Arch. Anat. Physiol., 1852: 379-391.
- Zenkewitsch, L.A., J.A. Birstein & G.M. Beliaev (1955): The bottom fauna of the Kurile-Kamchatka Trench. - Trudy Inst. Okeanol., 12: 375-381; Akad. Nauk. SSSR.
- Zharkova, I.S. (1976): Structural variability of vision organs of *Nymphon longicoxa* (Pantopoda). - Trudy Inst. Okeanol., Akad. Nauk SSSR, 99: 241-243. [in Russian]
- Ziegelmeier, E. (1978): Macrobenthos investigations in the eastern part of the German Bight from 1950 to 1974. - Rapp. P.-v. Reun. Cons. Int. Expl. Mer, 172: 432-444.
- Ziegler, A.C. (1960): Annotated list of Pycnogonida collected near Bolinas, California. - The Veliger, 3 (1): 19-22.
- Zilberberg, F. (1964): Pantopoda. In: P.E. Vanzolini (ed.). História natural de organismos aquáticos do Brasil - Bibliografia comentada. - Publ. Fund. Amparo Pesq. Estado São Paulo, 1-452.
- Zilberberg, F. (1965): Notes on Pantopoda. - Bolm. Inst. Oceanogr., Univ. São Paulo, 13 (2): 21-32.