

Isolation, Characterization and Crystal Structure of Cytotoxic *ent*-Kaurane Diterpenoids from *Isodon weisiensis* C. Y. Wu

Lan Ding^a, Zhang-Jing Zhang^b, Guo-An Liu^a, Dong-Juan Yang^a, Guo-Cong Guo^b, Han Wang^a, and Kun Sun^a

^a College of Life Science, Northwest Normal University, Lanzhou 730070, P. R. China

^b State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, The Chinese Academy of Sciences, Fuzhou, Fujian 350002, P. R. China

Reprint requests to Prof. Lan Ding. E-mail: dinglan641@sina.com; Fax: 86-931-7971207.

Z. Naturforsch. **60b**, 805 – 810 (2005); received February 14, 2005

The new *ent*-kaurane diterpenoid weisiensin C (**1**) ($1\alpha,7\alpha,14\beta,18,20$ -pentahydroxy-*ent*-kaur-16-en-15-one), together with four known *ent*-kaurene diterpenoids, glaucocalyxin A (**2**), kamebanin (**3**), macrocalyxin D (**4**), and excisanin K (**5**), was isolated from the leaves of *Isodon weisiensis* C. Y. Wu. The structure of the new constituent (**1**) was elucidated by spectroscopic methods and confirmed by single-crystal X-ray diffraction.

Key words: *Isodon Weisiensis* C. Y. Wu, *Ent*-kaurane Diterpenoid, Weisiensin C, Crystal Structure