

Nepetolide: A New Diterpene from *Nepeta suaveis*

Javid Hussain^a, Farman Ullah^a, Hidayat Hussain^b, S. Tasleem Hussain^a, and M. Raza Shah^c

^a Department of Chemistry Kohat University of Science and Technology Kohat, NWFP Pakistan

^b Laboratory of Organic Synthesis, UMR CNRS 6011, Université de Maine, F-72085,
Le Mans Cedex 9, France

^c International Center for Chemical Sciences, H. E. J. Research Institute of Chemistry, University of
Karachi, Karachi-75270, Pakistan

Reprint requests to Dr. Javid Hussain. Fax: (+92)-922-554556. E-mail: javidhej@yahoo.com

Z. Naturforsch. **2008**, *63b*, 591 – 594; received September 10, 2007

A new tricyclic clerodane-type diterpene, nepetolide (**1**), has been isolated from *Nepeta suaveis* along with three known compounds namely β -sitosterol, stigmasterol, and ursolic acid. The structure elucidation of the isolated compounds was based primarily on two-dimensional (2D)-NMR techniques including correlation spectroscopy (COSY), heteronuclear multiple quantum coherence (HMQC), and heteronuclear multiple bond correlation (HMBC) experiments.

Key words: *Nepeta suaveis*, Labiateae, Nepetolide, Clerodane Type Diterpene Lactone