



Database tool

Boechera microsatellite website: an online portal for species identification and determination of hybrid parentage

Fay-Wei Li^{1,2,*†}, Catherine A. Rushworth^{1,2}, James B. Beck^{3,4} and Michael D. Windham^{1,*†}

¹Department of Biology, Duke University, Durham, NC 27708, USA, ²University and Jepson Herbaria and Department of Integrative Biology, University of California, Berkeley, CA 94720, USA, ³Department of Biological Sciences, Wichita State University, Wichita, KS 67260, USA and ⁴Botanical Research Institute of Texas, 1700 University Drive, Fort Worth, TX 76107, USA

*Corresponding author: Email: fayweili@gmail.com, mdw26@duke.edu

Present address: Fay-Wei Li, Boyce Thompson Institute, Ithaca, NY 14850, USA

†These authors contributed equally to this work

Citation details: Li, F.W., Rushworth, C.A., Beck, J.B., *et al.* *Boechera* microsatellite website: an online portal for species identification and determination of hybrid parentage. *Database* (2016) Vol. 2016: article ID baw169; doi:10.1093/database/baw169

Received 3 September 2016; Revised 10 December 2016; Accepted 13 December 2016

Abstract

Boechera (Brassicaceae) has many features to recommend it as a model genus for ecological and evolutionary research, including species richness, ecological diversity, experimental tractability and close phylogenetic proximity to *Arabidopsis*. However, efforts to realize the full potential of this model system have been thwarted by the frequent inability of researchers to identify their samples and place them in a broader evolutionary context. Here we present the *Boechera* Microsatellite Website (BMW), a portal that archives over 55 000 microsatellite allele calls from 4471 specimens (including 133 nomenclatural types). The portal includes analytical tools that utilize data from 15 microsatellite loci as a highly effective DNA barcoding system. The BMW facilitates the accurate identification of *Boechera* samples and the investigation of reticulate evolution among the ± 83 sexual diploid taxa in the genus, thereby greatly enhancing *Boechera*'s potential as a model system.

Database URL: <http://sites.biology.duke.edu/windhamlab/>

Introduction

Among the close relatives of *Arabidopsis* Heynh., the genus *Boechera* Á. Löve & D. Löve (Brassicaceae) is

notable for its species richness, ecological diversity and experimental tractability, features that have contributed to its emergence as a model genetic system for studies of

evolutionary biology (1). However, efforts to fully leverage *Boecheera* as a model system have been hampered by the taxonomic complexity of the genus and resultant misidentifications of materials used in various research studies. The taxonomic uncertainty in *Boecheera* arises from a “perfect storm” of hybridization, polyploidy, and apomixis, all of which are rampant in the genus (2–7). Through these processes, the already substantial diversity of sexual diploid *Boecheera* taxa (± 83) has given rise to hundreds of additional taxa involving diverse combinations of at least 64 of these known sexual genomes (see 8–14).

In groups prone to such hybridization, determining which taxa represent the products of divergent evolution (i.e. sexual diploids) and which are the result of reticulate evolution is a prerequisite to developing a useable taxonomy (15–18). However, sexual diploids can be very difficult to detect morphologically in the apomictic hybrid milieu (14), and traditional molecular tools for species identification and hybrid diagnosis have proven of little use in *Boecheera* due to widespread allele-sharing across species (19–21) and possible plastid capture (21). To address this problem, we have developed a 15-locus nuclear microsatellite ‘barcode’ that distinguishes all known sexual diploids and simultaneously provides a framework for determining the genomic make-up of hybrid individuals. These short genetic markers offer another advantage, in that they can be reliably amplified from herbarium specimens of *Boecheera* collected over the past 130 years. Here we present a database of 56 353 microsatellite allele calls from 4471 specimens including 133 nomenclatural types. We also present novel tools for genotype-based species identification and determination of the genomic constitution of hybrid individuals.

Materials and methods

Microsatellite data

The database consists of microsatellite genotypes and basic locality data for 4471 *Boecheera* specimens representing ca. 95% of all previously named taxa. Each population of every taxon included in the database is represented by voucher specimens deposited in one or more of the 36 herbaria supporting this effort. DNA extraction and multiplex microsatellite genotyping were performed using protocols outlined in Beck *et al.* (12). In addition to the 13-locus microsatellite set employed in that study, we incorporated two additional loci (BF19 and a3; Table 1). These loci are useful for sample identification but are excluded from heterozygosity calculations because these loci occasionally exhibit more than the expected number of alleles in plants of known ploidy. Extra alleles likely indicate amplification of

multiple regions of the genome, and these loci are considered unreliable for heterozygosity estimates. The need for consistency in calling alleles is critical, and we have developed a set of locus-by-locus guidelines accessible through the *Boecheera* Microsatellite Website (BMW) entry page by clicking on ‘About & Contact’.

Inferring reproductive modes

The two primary reproductive pathways in *Boecheera* are sexuality (largely self-compatible and self-pollinating) and apomixis, the latter defined here as asexual seed production through either diplospory or apospory (25). The reproductive pathway of each accession in the database has been inferred based on a series of genetic and morphological assessments summarized in Figure 1. Foremost among these are analyses of meiotic chromosome pairing and segregation (14). Plants exhibiting normal pairing behavior resulting in the formation of spore tetrads are considered sexual; those with little or no pairing that produce malformed spores or functional diplospores in dyads are inferred to be apomictic. Including both published (summarized in 26) and unpublished chromosome studies, we have cytogenetic data for ~5% of all accessions in the database. Inferring the reproductive mode of the remaining accessions requires some additional observations.

The differences in chromosome pairing cited above lead to easily observed disparities in pollen morphology between sexuals and apomicts. Following Windham and Al-Shehbaz (8) and Beck *et al.* (12), accessions with a high percentage of well-formed, narrowly ellipsoid, tricolpate and symmetrical pollen (and no diplosporous pollen) are inferred to be sexual; those with a preponderance of malformed (highly variable) and/or ovoid-spheroid, multicolpate and asymmetric pollen are classified as apomictic. Approximately 60% of *Boecheera* herbarium specimens have flowers with sufficient pollen to ascertain reproductive mode. Assessment of reproductive mode in remaining specimens involves looking for evidence of stabilized hybridity in the available microsatellite data. Specimens exhibiting allelic additivity (involving either distinct taxa or inbred lines) at most loci are inferred to be apomicts, with the exception of a small number of sexual allotetraploids in the dataset. Accessions in which nearly all alleles are associated with a single taxon or inbred line are considered sexual.

Inferring ploidy levels

Many previous studies have reported variation in ploidy between and within species of *Boecheera* (see 23 and references therein). Voucher specimens for most published

Table 1. Information for the 15 microsatellite loci presented in the database

Marker	Primer sequences	Chr. no.	nt repeat	Number of alleles, N _a	Original publication
ICE3	GACTAATCATCACCGACTCAGCCAC ATTCTTCTTCACTTTTCTTGATCCCG	5	CT	62	(22)
ICE14	TCGAGGTGCTTTCTGAGGTT TACCTCACCTTTTGACCCA	2	GAT	17	(22)
a1	GTCTATTCGAGGACGCC AGGTTGGGTAGGTGAAG	2	GAT	15	(23)
a3	AGCTTTGTTTGCATGGAG GTGAGAATAATATTGACC	2	AG, AT	45	(23)
b6	GCAAAAGATCTTCATGGGAC TGCCATTTCTTCCCTAGTG	1	CT, GT	62	(23)
c8	TTCCGGGTATCATTCTAG GTTGTAAGTTCTTTCTCAG	5	CTT	58	(23)
e9	GCGTATCTCGAATCACCTTTG CTCCCCCTGAGTTTTTCAAG	4	CT	56	(23)
BF3	TTTTTAGACAGTAGTGGCTGTGAG ACTTCGTTCCAGGCTCGTC	4	GA	58	(24)
BF9	AAACACATCCCCTCAGCTC TTGATTGAATCCTGCGTTTG	3	GA	55	(24)
BF11	TCCTCCATTGTAGAGCAGAGC CCATTGCTTAAACCCTAAACC	2	GA	27	(24)
BF15	CAGCATCTCCTTTGGGTTTG ACTTGCTCCTTTGCATGACC	5	GA	48	(24)
BF18	AACCTCCCAAGATTCGCTTC TTCGCCATTGTTGTGATTTG	1	CT	30	(24)
BF19	ACCGCATTGGTGTGTGTC ATAACGGACGCGACCAAAG	-	GA	63	(24)
BF20	TTCTCGGGAAAGTAATGAGGAG GCAAAATCTGACCAATGCAAG	2	CT	47	(24)
Bdru266	TTTAATTTGTGCGTTTGATCC CAAAATCGCAGAATGAGAGG	-	AT	54	(24)

Loci a3 and BF19 occasionally exhibit more than the expected number of alleles in plants of known ploidy and were therefore excluded from heterozygosity calculations. Null alleles were treated as missing data and are not included in the total number of alleles.




Figure 1. Summary of the steps involved in inferring reproductive mode in accessions of *Boechea*.

chromosome counts in the genus have been included in the database, as have the vouchers for several hundred unpublished counts (Windham *et al.*, in prep.). These data reveal a very strong association between the chromosome number documented for each individual and the maximum number of alleles observed at 13 of the 15 microsatellite loci analyzed (excluding loci a3 and BF19; Table 1). Cytogenetically known diploids in *Boecheira* range from completely homozygous to having two alleles at 12 of these 13 putatively single-copy loci, whereas documented apomictic triploid and apomictic tetraploid individuals exhibit at least two loci with three or four alleles, respectively. These observations provide a pathway for inferring the ploidy of accessions that have not been studied cytogenetically but have appropriate microsatellite data. These inferences are based on the maximum number of alleles at two or more loci, a conservative threshold adopted to prevent the inflation of inferred ploidy based on a single aberrant locus or allele call. For example, a specimen would be called triploid if two or more of the loci used to assess heterozygosity (Table 1) exhibited three alleles. Plants that are homozygous at all loci are, by default, diploid. Although this approach works well for distinguishing diploids from triploids and apomictic tetraploids, sexual tetraploids are less tractable. The database includes 56 sexual tetraploid accessions representing two taxa: *holmgrenii* × *lemmonii* and *laevigata* × *stricta*. These hybrid species exhibit

disomic inheritance (fixed heterozygosity) and rarely show more than two alleles per locus, as is common in sexual allotetraploids derived from relatively divergent species (26). Thus they are largely indistinguishable from highly heterozygous (apomictic) diploids based on the maximum number of alleles per locus. However, sexual tetraploids are rarely encountered in *Boecheira* (27, 28), and they are readily separated from apomictic diploids by other criteria, including pollen morphology and chromosome number (Fig. 1).

Taxonomic coverage

In the database as well as this article, known or inferred sexual diploid samples are designated by single epithets (e.g. ‘gracilipes’), most validly published in *Boecheira* but with a few still pending. Inferred apomicts (diploid, triploid or tetraploid) and the few sexual tetraploids are identified by the appropriate number and combination of epithets, each separated by the hybrid symbol ‘×’ (e.g. ‘gracilipes × perennans × texana’). After removing 43 accessions that lack data at eight or more loci, the remaining 4428 accessions included 2594 presumed or known sexual individuals and 1834 likely apomicts (Table 2). These represent 485 unique entities (sexual diploid taxa and their hybrid derivatives), a phenomenal amount of diversification given the relatively short evolutionary history of the

Boecheira Microsatellite Website

Microsatellite Database Wizard

- SEARCH DATABASE
 - BY EXTRAC#
 - BY TAXON
 - BY LOCALITY
 - BY ALLELES
 - BATCH
- IDENTIFY PARENTS
 - BY EXTRAC#
 - BY ALLELES
 - BATCH
- TUTORIAL
- ABOUT & CONTACT

PRIUS: Parental Relationship Identification Using Subtraction

DNA extraction ID:

Query: PJA154A Heterozygosity: 0.923076923077 No. empty loci: 0

This appears to be a triploid hybrid

Cumulative score	Putative parent 1	Allele similarity score	Putative parent 2	Allele similarity score	Putative parent 3	Allele similarity score
2.1333	perennans (FW812)	0.8667	texana (JB1454)	0.7333	gracilipes (JB226)	0.5333
2.1333	perennans (FW812)	0.8667	texana (PJA269J)	0.7333	gracilipes (JB907)	0.5333
2.1333	perennans (FW812)	0.8667	texana (PJA269G)	0.7333	gracilipes* (JB603)	0.5333
2.1333	perennans (FW812)	0.8667	texana (JB1448)	0.7333	gracilipes (FW1621)	0.5333
2.1333	perennans (FW812)	0.8667	texana (PJA269H)	0.7333	gracilipes* (JB603)	0.5333
2.1333	perennans (FW812)	0.8667	texana (PJA269J)	0.7333	gracilipes (JB905)	0.5333
2.1333	perennans (FW812)	0.8667	texana (JB1454)	0.7333	gracilipes (FW1628)	0.5333
2.1333	perennans (FW812)	0.8667	texana (JB1454)	0.7333	fendleri (PJA225D)	0.5333
2.1333	perennans (FW812)	0.8667	texana (JB1454)	0.7333	gracilipes* (JB603)	0.5333
2.1333	perennans (FW812)	0.8667	texana (PJA269J)	0.7333	gracilipes (FW1621)	0.5333

Figure 2. Screen shot of entry page for the BMW listing basic functions (left banner) and showing the result of a PRIUS search on Extrac# PJA154a.

Table 2. Summary of inferred ploidy levels and reproductive modes in 4428 *Boecheera* database accessions after removing 43 accessions with >50% missing data

	Diploid	Triploid	Tetraploid	Total
Sexual	2538	0	56	2594
Apomictic	863	959	12	1834
Total	3401	959	68	4428

group (29). A full list of sexual diploid taxa and the current sampling of each can be accessed by clicking on ‘By Taxon’ on the entry page of the website. As of this publication, all known sexual diploid taxa except the critically endangered *B. perstellata* are represented in the database, with an average of 31 samples each. Several previously recognized rare endemics (e.g. *Boecheera hoffmannii*, *Boecheera serotina*, *Boecheera yorkii*) remain poorly sampled, as do many of the newly discovered sexual diploids (e.g. ‘*roguensis*’, ‘*tahoensis*’, ‘*wallowaensis*’). In terms of geography, the region with the greatest taxonomic complexity (i.e. western North America) is generally well sampled. However, the various taxa confined to eastern North America and eastern Asia are currently under-sampled. Our ongoing research efforts are focused on filling these gaps, and the database will be continuously updated as new results become available.

Database structure

In order to facilitate the broadest usage of these data, the database has been made openly accessible through a web portal (<http://sites.biology.duke.edu/windhamlab/>), constructed using custom Python and HTML scripts, and connected by the Common Gateway Interface. The core is a Python library for interacting with the database (scripts available at <https://bitbucket.org/fayweili/boecheradata> base.git). Users can search and retrieve basic metadata using the Extrac# (a unique ID number annotated on every voucher specimen), taxon name, or locality (Fig. 2). Searches based on taxon name can be limited to sexual diploids (i.e. just the epithet entered) or expanded to include all hybrids containing that genome. The output of a taxon query can be used to produce working state and county level distribution lists free of the misidentifications common in other on-line resources. The search by locality function allows investigators to quickly assemble checklists of confirmed *Boecheera* identifications for any state, province, or county. This can improve the efficiency of morphology-based identification by reducing the diversity of taxa and the number of dichotomous key couplets that need to be negotiated. To further facilitate this process, exemplar

specimens of each sexual diploid taxon as well as all documented hybrid combinations are being imaged and will be available via the portal. In addition to the basic search and display functions, we also include two novel algorithms: ‘Taxon Enquiry based on Similarity of Loci and Alleles’ (TESLA) for sample identification, and ‘Parental Relationship Identification Using Subtraction’ (PRIUS) for inferring the genomic constitution of hybrid individuals (see below). Although many microsatellite databases have been developed for plants, the majority of them focus on crop plants with the main purpose to assist breeding and genotyping marker selection (e.g. 30, 31). To our knowledge, the only comparable database for biodiversity research and species identification is the Olive Genetic Diversity Database (32).

Taxon enquiry based on similarity of loci and alleles (TESLA)

Taxonomic identification of *Boecheera* specimens is notoriously challenging, even for specialists (11). With about ± 83 sexual diploid taxa that hybridize whenever they come into contact to produce true-breeding apomictic diploids, triploids and tetraploids, it is impossible to identify many accessions based on gross morphology alone (14). To address this problem, we have developed TESLA, a microsatellite-based identification method. Once a user enters allelic information for their sample, TESLA compares this multi-locus genotype (MLG) to every accession in the database, calculating an Allele Similarity (AS) score between two accessions as follows:

$$AS = \frac{(\text{No. shared alleles}) \times (\text{No. loci} - \text{No. loci without data})}{(\text{Total allele count in query}) \times (\text{No. loci})}$$

The AS score represents the proportion of alleles that are shared relative to the query, scaled by the amount of missing data. The output of a TESLA query consists of the specimens with the 100 highest AS scores listed in descending order. In addition to researching samples already included in the database, user-generated MLGs can be entered individually (using the ‘by alleles’ function) or a query spreadsheet can be uploaded to the database in batch mode (Fig. 2).

Both the rank value and magnitude of the AS scores derived from TESLA are useful in identifying unknown *Boecheera* accessions. For example, in queries involving 100 random sexual diploid samples of known identity, the highest AS scores coincided with the ‘correct’ identification 96% of the time. In an additional 3%, the highest scoring matches belonged to the same major clade as the expected taxon (clade assignments of all sexual diploid taxa can be

accessed through ‘Search Database By Taxon’). These phylogenetically localized mismatches were attributable to limited sampling of the taxon involved, poor differentiation between the target taxon and its close relatives, and/or missing data. In the single instance in which the individual with the highest AS score belonged to an unrelated taxon, the analysis was impeded by a lack of data at the majority of loci. In terms of magnitude, the highest AS scores from TESLA searches resulting in correct identifications ranged from 1.000 to 0.375. Misidentifications that attribute a sample to the wrong clade became common below 0.500 and we therefore recommend this as the threshold for considering a TESLA result a hypothesis worthy of serious consideration.

Parental relationship identification using subtraction (PRIUS)

Boecheira hybrids are ubiquitous, often comprising >50% of plants randomly sampled from natural populations (M. D. Windham, personal observation). They are also hyperdiverse, drawing nearly every known sexual genome into the apomictic milieu (12, 13). Few extant analytical programs can determine the genomic constitution of hybrid individuals based on microsatellite data (33–35), and none of the available implementations are capable of handling the tri-genomic hybrids common in *Boecheira*. To address this need, we have developed a program called ‘PRIUS’, available via the online database portal.

The basic workflow of PRIUS is as follows: (i) identify the most similar sexual diploid specimens in the database based on AS score, (ii) subtract from the hybrid’s MLG the corresponding alleles from each top-scoring specimen and (iii) repeat the previous steps using the subtracted genotype for one or two more iterations (see below). The end result is a list of diploid specimen combinations that best explain the hybrid’s MLG. The corresponding species of these diploid specimens then form a working hypothesis of the putative parental species.

The proper functioning of PRIUS depends on the ability to distinguish between sexual diploids (the pool from which potential parents are drawn) and apomicts (generally the products of hybridization; see 12, 36). Fortunately, this information resides within the microsatellite data themselves. Based on a broad taxonomic sampling of 13 microsatellite loci in 1393 *Boecheira* specimens, Beck *et al.* (12) demonstrated clear bimodality in the number of heterozygous loci, a metric that was strongly associated with inferred breeding system. In this study, the mean heterozygosity of diploid plants with predominantly sexual pollen was 0.232 (3.02/13), whereas the mean for diploid plants with predominantly apomictic pollen was 0.723 (9.4/13).

Only one sexual diploid taxon (*parishii*) included multiple samples in which heterozygosity exceeded 0.5, and only 6 of 82 apomictic diploid taxa exhibited heterozygosity <0.5 (12). Based on this empirical evidence, we set the PRIUS heterozygosity threshold at 0.5, thus creating essentially non-overlapping pools of potential parents (sexual diploids) and apomictic (or sexual tetraploid) hybrids subject to a PRIUS query.

To begin the PRIUS process, the program infers the ploidy of the query accession based on the maximum number of alleles at two or more of the 13 loci used to calculate heterozygosity (see Table 1). If the accession is determined to be triploid, PRIUS identifies the five sexual diploid specimens in the full dataset with the highest AS scores relative to the queried MLG. For each tri-allelic locus in the query accession, PRIUS independently subtracts from the MLG any matching alleles observed in each of the five sexual diploids, generating five new query MLGs. These roughly approximate apomictic diploid progenitors that could have hybridized with the subtracted genome to form the original triploid MLG. Each of the five subtracted MLGs is then queried in a second iteration, where PRIUS identifies the five sexual diploid specimens with the highest AS scores relative to each of the subtracted MLGs. For each di-allelic and tri-allelic locus in the five queried MLGs, the program subtracts any matching alleles from each of the five most similar sexual diploids. This step generates 25 (5^2) new query MLGs that very roughly approximate a third sexual diploid genome that might have combined with the two subtracted genomes to form the original triploid MLG. A third iteration then identifies the five sexual diploid specimens with the highest AS scores relative to each of the 25 new subtracted MLGs. After three iterations, there will be 125 (5^3) different specimen combinations, and PRIUS ranks them by cumulative AS scores. If a query accession is determined to be diploid (i.e. exhibits a maximum of two alleles at 12 of the 13 loci used to assess heterozygosity), PRIUS begins with the second iteration. As with TESLA, a user generated MLG can be entered individually (using the ‘by alleles’ function) or a query spreadsheet can be uploaded to the database in batch mode (Fig. 2).

PRIUS is not designed to directly address the parentage of the few tetraploid apomicts encountered (12/4428 specimens or 0.3% of the dataset) because a fourth round of subtraction produces a high percentage of spurious matches. PRIUS searches are also negatively affected by the presence of undetected null alleles in a queried accession. This is because the AS scores of appropriate sexual diploid accessions with null alleles are lowered, reducing their likelihood of being identified as possible parents. The effect can be significant, resulting in a bias against certain

sexual diploid taxa (e.g. *retrofracta*) that are extensively involved in the formation of apomictic hybrids (11). This problem is not easily addressed, but we have instituted an approach that improves the success rate in both PRIUS and TESLA for many triploid accessions.

In sexual taxa, null alleles are hypothesized to exist in the homozygous state if both of the following conditions were met: (i) the remaining multiplexed loci from that individual amplified, and (ii) individuals in the same genotyping run representing other taxa that lack null alleles amplified. Because most *Boecheira* triploids (even those arising within named species) combine divergent genomes with different alleles, we expect a given triploid sample to be tri-allelic or, at least, di-allelic at the majority of these highly variable loci (Table 1). A broad survey of the database suggests that the occurrence of apparently homozygous loci in such accessions is more often the result of undetected null alleles than the improbable incorporation of three identical alleles from three different parental genomes. Both PRIUS and TESLA therefore add a null allele (as '0') to apparently homozygous loci in triploid queries. This protocol has no effect on the AS scores of accessions supporting the 'three identical alleles' hypothesis but increases the probability that sexual diploids with null alleles will be considered as possible parents.

It should be stressed that although PRIUS is a powerful tool for data exploration and generating hypotheses regarding the genomic constitution of hybrid *Boecheira* individuals, the hypothesized hybrid origin should be evaluated with the widest array of data that can be brought to bear, including morphology, cytology, ecology, geography, genomics and reproductive biology (13).

Results

The following examples demonstrate the functionality of TESLA (Table 3) and PRIUS (Fig. 2) for identifying hybrids. The sample queried (Extrac# PJA154a) was subject to a variety of analyses by Alexander *et al.* (13) and thus provides a good basis for comparing the outputs of these algorithms to other approaches used to identify taxa and propose hybrid parentages. Table 3 shows the result of a TESLA search on PJA154a, demonstrating that all of the most similar accessions in the database (with AS scores ranging from 1.0000 to 0.9143) have been identified as triploid hybrids incorporating genomes derived from *gracilipes*, *perennans* and *texana*. To facilitate presentation of this TESLA query, the full 15-locus dataset has been reduced to the eight loci that show allelic variability within this hybrid cluster. In addition, the standard output of TESLA (i.e. the 100 most similar accessions in the

database) has been condensed by grouping hybrids with identical genotypes and assigning members of individual sexual diploid taxa with identical AS scores to a single row (the number of condensed accessions indicated parenthetically in the Extrac# column).

As indicated earlier, the top hits for a TESLA query on PJA154a are all specimens identified as *gracilipes* × *perennans* × *texana*. These comprise 13 different MLGs representing 30 individual samples (Table 3). In fact, all but two of the samples in the dataset with this hypothesized parentage are clustered at the top of this figure with AS scores ≥ 0.9143 . One exception is Extrac# FW765 (Table 3; row 19), which is identical to PJA222e (row 12) except that it lacks data at two loci (c8 and b6) and has incomplete data at two others (ICE14 and Bdru266). The remaining exception to the tight clustering of samples identified as *gracilipes* × *perennans* × *texana* is Extrac# JB1453 (Table 3; last row), which has an AS score of 0.6111 reflecting divergent allele profiles at seven of the eight illustrated loci. We suspect that Extrac# JB1453 represents an independent hybrid origin of this triploid apomict, a hypothesis discussed in more detail below.

Random sampling of apomictic hybrids in the current dataset indicate that ca. 60% of TESLA searches correctly specify their genomic make-up, placing plants with identical parentage at the top of the results list. In the remaining 40% of cases in which highly similar hybrid genotypes have not yet been sampled, the probable parents typically are discernable among the sexual diploid taxa with high AS scores. Our query involving PJA154a (Table 3) can be used to illustrate this approach as well. Ignoring for the moment the hybrid origin suggested by rows 3–15, the next highest AS score (0.8667) is shared by individuals of *perennans* and *gracilipes*. A third sexual diploid taxon (*texana*) appears at an AS score of 0.7333. Repeated occurrences of these three species [as well as *gracilipes* × *perennans*, a possible apomictic diploid bridge discussed by Alexander *et al.* (13)] suggest the same hybrid parentage reflected in rows 3–15.

PRIUS provides an alternative, taxon-centric approach to assessing hybrid parentage and is especially informative in cases where TESLA fails to provide a clear result. Figure 2 presents a screen shot of the PRIUS query result for Extrac# PJA154a, the same individual used in the preceding TESLA search. The cumulative scores on the left serve the same function as the AS scores in TESLA. In the case of PJA154a, the highest cumulative score is 2.1333, which is represented by 92 rows (different combinations of individual accessions), with 10 presented in Figure 2, for clarity. In every case, putative parent 1 is an accession of *perennans* while putative parent 2 is *texana*. In 82/92 combinations (89.1%), an accession of *gracilipes* appears as

Table 3. Output from a TESLA query on Extrac# PJA154a

AS_score	Extrac#	Taxon ID	State	County	ICE3	ICE3	ICE3	c8	c8	ICE14	BF3	BF3	BF3	BF3	b6	b6	BF19	BF19	BF15	BF15	BF15	Bdru266	Bdru266	
	Query=	gracilipes x	NM	Eddy	71	75	93	244	250	211	223	107	111	129	299	309	311	145	0	93	104	115	133	146
	PJA154a	perennans x texana																						
1	PJA154b (3)	gracilipes x	NM	Eddy	71	75	93	244	250	211	223	107	111	129	299	309	311	145		93	104	115	133	146
		perennans x texana																						
0.9722	PJA163h (6)	gracilipes x	NM	Doña Ana	71	75	93	244	250	211	223	107	111	129	299	309	311	145		93	104	115	133	154
		perennans x texana																						
0.9722	PJA382z (2)	gracilipes x	NM	Torrance	71	75	93	244	250	211	223	107	111	129	299	309	311	145		93	104	115	133	150
		perennans x texana																						
0.9722	PJA380a	gracilipes x	NM	Torrance	71	75	93	244	250	211	223	107	111	129	299	309	311	145		93	104	115	133	152
		perennans x texana																						
0.9722	PJA385f	gracilipes x	NM	Otero	71	75	93	244	250	211	223	107	111	129	299	309	311	145		93	104	115	133	142
		perennans x texana																						
0.9459	PJA380z	gracilipes x	NM	Torrance	71	75	93	244	250	211	223	107	111	129	299	309	311	143	145		93	104	115	133
		perennans x texana																						
0.9444	PJA230a (2)	gracilipes x	NM	Doña Ana	71	75	93	244	250	211	223	107	111	129	299	307	311	145		93	104	115	133	150
		perennans x texana																						
0.9444	PJA230c (2)	gracilipes x	NM	Doña Ana	71	75	93	244	250	211	223	107	111	129	299	307	311	145		93	104	115	133	152
		perennans x texana																						
0.9211	JB1346 (2)	gracilipes x	NM	San Miguel	71	75	93	244	250	211	223	107	111	129	299	309	311	143	145		93	104	115	130
		perennans x texana																						
0.9189	PJA222e (2)	gracilipes x	NM	Doña Ana	71	75	93	244	250	211	223	107	111	127	299	309	311	145		93	104	115	130	150
		perennans x texana																						
0.9189	PJA385z (3)	gracilipes x	NM	Otero	71	75	93	244	248	250	211	223	107	111	129	299	309	311	145		93	104	115	131
		perennans x texana																						
0.9167	JB1345	gracilipes x	NM	Socorro	71	75	93	244	250	211	223	111	127	299	309	311	145		93	104	115	133	136	
		perennans x texana																						
0.9143	PJA395c (4)	gracilipes x	NM	Lincoln	71	75	77	244	250	211	223	107	111	129	299	309	145		93	104	115	131	150	
		perennans x texana																						
0.8667	FW812	perennans	AZ	Maricopa	75			244	258	223		111	155	307			145	153		93			141	
		gracilipes																						
0.8667	FW483	gracilipes	AZ	Gila	71			250	252	223		105	107	317	321		147			104			125	
		perennans																						
0.8	FW799 (4)	perennans	AZ	Gila	75			244		0		111	135	305	309		157	163		93			154	
		gracilipes x																						
0.7407	FW765	gracilipes x	NM	Doña Ana	71	75	93			223		107	111	127			145			93	104	115	130	
		perennans x texana																						
0.7333	PJA151h (14)	perennans	NM	Doña Ana	75			248	254	223		99	129	305	319		149			93			125	
		texana																						
0.7333	JB1454 (8)	gracilipes x	TX	Jeff Davis	73			232		211		97	123	313		0				101	115		146	
		perennans																						
0.7143	JB595	gracilipes x	AZ	Cochise	71	75		248	252	223		105	129	311	313		145	151	155	93	104		141	
		perennans																						
0.6667	FW811 (30)	perennans	AZ	Maricopa	75			264		223		119		305		0				93			133	
		gracilipes																						
0.6667	JB905 (6)	gracilipes	AZ	Coconino	71			258		223		99		321		145	151			104			0	
		spatifolia																						
0.6667	JB183 (2)	texana	CO	Fremont	75			242		223		101		311		143				99			133	
		perennans																						
0.6667	JB1447	gracilipes x	TX	Brewster	73			232	240	211		107	115	313	319		0			101	137		144	
		perennans																						
0.6538	JB648	gracilipes x	AZ	Graham	73	75		252	262	223		111	119	309	315		145			93	104		156	
		perennans																						
0.6111	JB1453	gracilipes x	TX	Culberson	73	75		232	242	248	211	223	107	113	125	299	309	315	147	151	155	93	104	137
		perennans x texana																						

putative parent 3. In another six cases (6.5%), *perennans* is repeated as the third parent and, in four others (4.4%), *fendleri* is identified as putative parent 3. All four of these sexual diploid taxa belong to the same major clade of *Boechea* (13, 21).

In Alexander *et al.* (13) Extrac# PJA154a is labeled PJA195a (the original collector's number) and is identified as *Boechea porphyrea* (the correct binomial name for hybrids of this proposed parentage). In a phylogenetic analysis incorporating data from the nuclear *pistillata* locus, different clone sequences from this sample associated with *gracilipes*, *perennans* and *texana*, respectively (13). STRUCTURE analysis of the microsatellite data available at the time for this taxon revealed admixture proportions averaging 25% *gracilipes*, 37% *perennans* and 23% *texana*, with a lesser contribution attributed to *fendleri*. Thus, the results reported by Alexander *et al.* (13) are nearly identical to those obtained from our TESLA and PRIUS queries on PJA154a. Our new algorithms have two clear advantages, however. First, using TESLA and PRIUS requires no specialized taxonomic knowledge of the group. To conduct their analyses, both Alexander *et al.* (13) and Beck *et al.* (12) had to decide *a priori* which *Boechea* species were likely involved in the hybridization events they were trying to reconstruct. Second, TESLA and PRIUS results are obtained in seconds, as opposed to the time-consuming STRUCTURE algorithm.

Discussion

In addition to providing microsatellite-based tools to facilitate the identification of *Boechea* specimens, the database also can be used to explore large-scale evolutionary patterns within the genus. Insights regarding the relationships among ploidy, heterozygosity and reproductive mode, and processes of hybrid formation and evolution are discussed below.

Relationship between reproductive mode and ploidy

Boechea ploidy is non-randomly associated with reproductive mode (28). The clearest such association involves plants with three full sets of chromosomes ($2n = 21$) that have no chance of pairing normally in meiosis. The current dataset includes 959 triploid accessions (Table 2), all of which are inferred to be apomictic based on the stepwise analyses outlined in Figure 1. Whereas triploids represent 21.7% of the current dataset, tetraploids are rare (1.5%) and higher ploidy levels have not been observed. Based on cytogenetic and pollen data, the majority of tetraploid individuals are sexual; though 12 apomictic tetraploids

(representing seven different genomic combinations) are present as well (Table 3). The two types of tetraploids are readily separated based on the maximum number of alleles per locus (two or rarely three in sexual tetraploids vs. four in apomicts). The remaining accessions (77.8%) are considered diploid based on the allele number criterion discussed earlier. Although ca. 75% of these appear to be sexual, our sampling was biased toward sexual individuals, and apomictic diploids are likely to be more prevalent in nature. This is a critical feature of the *Boechea* model system given the rarity of apomictic diploids among flowering plants (25).

Relationship between reproductive mode and heterozygosity

As suggested by previous studies, apomicts exhibit higher levels of heterozygosity than sexually reproducing *Boechea* accessions (Fig. 3). This is not surprising given that the observed heterozygosity is largely the result of hybridization between different inbred genotypes, which may be a trigger for the development of apomixis (12, 37). Results from a Wilcoxon rank sum test of the current dataset indicate that apomict heterozygosity is significantly higher than that of sexuals (apomict mean 0.78 ± 0.14 SD, median 0.77 vs. sexual mean 0.11 ± 0.18 SD, median 0; Wilcoxon $W = 4709860$, $P < 2.2e^{-16}$; Fig. 3). Although mean heterozygosity of triploid (0.85 ± 0.10 SD) and tetraploid apomicts (0.90 ± 0.07 SD) is higher than that of diploid apomicts (0.70 ± 0.13 SD), this result does not change qualitatively when comparing diploid only sexuals and apomicts.




Figure 3. Histogram of heterozygosity for 4428 *Boechea* accessions included in this study, colored by reproductive mode (sexuals in purple and apomicts in orange) and ploidy level (darker with increasing ploidy).

The data presented in Figure 3 provide strong empirical support for using the 0.5 heterozygosity threshold implemented in PRIUS to separate potential parents (sexual diploids) from the non-segregating hybrids subject to analysis. Just 147 accessions (3.3% of the dataset) exhibit heterozygosity values higher or lower than expectations based on reproductive mode inferred via the pathway outlined in Figure 3. Unexpectedly high values are observed in 56 accessions of the two sexual tetraploid taxa (*holmgrenii* × *lemmonii* and *laevigata* × *stricta*) discussed earlier and 28 individuals of the sexual diploid taxon *parishii*. This species exhibits unusually high allelic diversity and possesses a suite of traits that suggest or enforce outcrossing. Unexpectedly low values are observed in a series of apomictic diploids that include: (i) a variety of hybrids between closely related sexual diploids that are insufficiently divergent to meet the 0.5 threshold when crossed, and (ii) a series of apomictic diploid hybrids that include *retrofracta* or other species with null alleles in their parentage.

Hybrid formation and evolution

Earlier articles hint at the important role hybridization has played in the evolutionary diversification of *Boecheera* (12, 13), but we clearly establish that hybridization is ubiquitous and largely unconstrained by phylogeny. Of the 485 taxa included in the current database, 400 are confirmed or inferred apomictic hybrids. Eighteen of these have arisen through crosses between divergent inbred lines within particular sexual diploid taxa; the remaining 382 are the products of hybridization between distinct taxa. Of the 83 sexual diploid taxa, 64 have been involved in at least one hybridization event and the 19 sexual diploid taxa not currently known to hybridize are those that rarely co-occur with other taxa and/or are currently under-sampled.

Hybridization has also occurred repeatedly among the same sexual diploid taxa, producing hybrid lineages with similar taxonomic constitutions but different parental alleles. The results for a TESLA query for Extrac# PJA154a, provide a good example (Table 3). The queried specimen, an apomictic triploid hybrid incorporating genomes derived from *gracilipes*, *perennans* and *texana* (13), belongs to a cluster of 13 very similar MLGs with AS scores ≥ 0.9143 and one outlier (Extrac# FW765, AS 0.7407) identical to one of these except for missing data. There is, however, a 14th MLG represented by a single sample (Table 3; Extrac# JB1453, row 28) with a much lower AS score of 0.6111. The lower score reflects the fact that there are allelic differences between JB1453 and the other 13 MLGs at nearly every locus analyzed. Given the lack of chromosome pairing and allelic segregation in these apomictic triploids, it is very likely that the apomictic triploid

gracilipes × *perennans* × *texana* had at least two independent origins. A cursory survey of the data included in the BMW indicates that multiple origins of *Boecheera* hybrid apomicts are the rule rather than the exception.

Conclusion

The taxonomic complexity of *Boecheera* has long stymied those investigating the diversity, biogeography, and evolutionary dynamics of the group (8, 11). Here we present a database containing information for 4428 specimens collected throughout the native range in North America, Greenland and Russia and encompassing >95% of all named taxa. A variety of search algorithms facilitate assigning the correct taxon name, ploidy, reproductive mode, and hybrid parentage to any *Boecheera* specimen with appropriate microsatellite data. Two novel identification algorithms, TESLA and PRIUS, facilitate the identification of unknown *Boecheera* specimens using simple, standardized methods. This database, publicly available and regularly updated, will provide valuable new tools for researchers interested in *Boecheera* as a model genus.

Acknowledgements

We thank the curators of the following herbaria (acronyms from 38) for permission to remove pollen and small leaf fragments: ALA, ARIZ, ASC, ASU, BHSU, BRY, CAN, CAS, COLO, CS, DAO, DS, DUKE, GH, JEPS, MO, MONTU, NCU, NDG, NY, ORE, OSC, POM, RM, RSA, SJNM, UBC, UC, UNLV, UNM, USFS, UT, UTC, WILLU, WS, WTU and also Ryan Hardy at Duke Office of Information Technology for helping to setup the website server. Kathleen Pryer, Tom Mitchell-olds, Carl Rothfels, and three anonymous reviewers also provided insightful suggestions.

Funding

This work was funded by National Science Foundation DEB-0816560 (to M.D.W.), Doctoral Dissertation Improvement Grant DEB-1407158 (to F.-W.L.) and DEB-1311269 (to C.A.R.). F.-W.L. and C.A.R. were each supported by a National Science Foundation Graduate Research Fellowship.

Conflict of interest. None declared.

References

1. Rushworth, C.A., Song, B.H., Lee, C.R.R. *et al.* (2011) *Boecheera*, a model system for ecological genomics. *Mol. Ecol.*, 20, 4843–4857.
2. Rollins, R.C. (1941) A monographic study of *Arabis* in western North America. *Rhodora*, 43, 289–325.
3. Böcher, T.W. (1951) Cytological and embryological studies in the amphipomictic *Arabis holboellii* complex. *Biol. Skr. Dan. Vid. Selsk.*, 6, 1–59.
4. Böcher, T.W. (1964) Further studies in the *Arabis holboellii* complex. *Bot. Tidsskr.*, 64, 141–161.

5. Böcher, T.W. (1954) Experimental taxonomic studies in the *Arabis holboellii* complex. *Svensk. Bot. Tidsskr.*, 48, 31–44.
6. Rollins, R.C. *The Cruciferae of Continental North America*; Stanford University Press, Palo Alto, 1993.
7. Roy, B.A., and Rieseberg, L.H. (1989) Evidence for apomixis in *Arabis*. *J. Hered.*, 80, 506–508.
8. Windham, M.D., and Al-Shehbaz, I.A. (2006) New and noteworthy species of *Boechea* (Brassicaceae) I: sexual diploids. *Harvard Pap. Bot.*, 11, 61–88.
9. Windham, M.D., and Al-Shehbaz, I.A. (2007) New and noteworthy species of *Boechea* (Brassicaceae) II: apomictic hybrids. *Harvard Pap. Bot.*, 11, 257–274.
10. Windham, M.D. and Al-Shehbaz, I.A. (2007) New and noteworthy species of *Boechea* (Brassicaceae) III: additional sexual diploids and apomictic hybrids. *Harvard Pap. Botany*, 12, 235–257.
11. Al-Shehbaz, I.A. and Windham, M.D. (2010) *Boechea*. In: Flora of North America Editorial Committee (ed). *Flora of North America North of Mexico*. Oxford University Press, Oxford, pp. 348–412.
12. Beck, J.B., Alexander, P.J., Allphin, L. *et al.* (2012) Does hybridization drive the transition to asexuality in diploid *Boechea*? *Evolution*, 66, 985–995.
13. Alexander, P.J., Windham, M.D., Beck, J.B. *et al.* (2015) Weaving a tangled web: Divergent and reticulate speciation in *Boechea fendleri* sensu lato (Brassicaceae: Boecheae). *Syst. Bot.*, 40, 572–596.
14. Windham, M.D., Beck, J.B., Li, F.W. *et al.* (2015) Searching for diamonds in the apomictic rough: a case study involving *Boechea lignifera* (Brassicaceae). *Syst. Bot.*, 40, 1031–1044.
15. Rieseberg, L.H., Carter, R. and Zona, S. (1990) Molecular tests of the hypothesized hybrid origin of two diploid *Helianthus* species (Asteraceae). *Evolution*, 44, 1498–1511.
16. Wendel, J.F., Schnabel, A. and Seelanan, T. (1995) Bidirectional interlocus concerted evolution following allopolyploid speciation in cotton (*Gossypium*). *Proc. Natl. Acad. Sci. USA.*, 92, 280–284.
17. Doyle, J.J., Doyle, J.L. and Brown, A.H. (1999) Origins, colonization, and lineage recombination in a widespread perennial soybean polyploid complex. *Proc. Natl. Acad. Sci. USA.*, 96, 10741–10745.
18. Beck, J.B., Windham, M.D., Yatskievych, G. *et al.* (2010) A diploids-first approach to species delimitation and interpreting polyploid evolution in the fern genus *Astroblepis* (Pteridaceae). *Syst. Bot.*, 35, 223–234.
19. Kiefer, C., Dobeš, C. and Koch, M.A. (2009) *Boechea* or not? Phylogeny and phylogeography of eastern North American *Boechea* species (Brassicaceae). *Taxon*, 58, 1109–1121.
20. Kiefer, C., and Koch, M.A. (2012) A continental-wide perspective: the genepool of nuclear encoded ribosomal DNA and single-copy gene sequences in North American *Boechea* (Brassicaceae). *PLoS One*, 7, e36491.
21. Alexander, P.J., Windham, M.D., Beck, J.B. *et al.* (2013) Molecular phylogenetics and taxonomy of the genus *Boechea* and related genera (Brassicaceae: Boecheae). *Syst. Bot.*, 38, 192–209.
22. Clauss, M.J., Cobban, H., and Mitchell-Olds, T. (2002) Cross-species microsatellite markers for elucidating population genetic structure in *Arabidopsis* and *Arabis* (Brassicaceae). *Mol. Ecol.*, 11, 591–601.
23. Dobeš, C., Mitchell-Olds, T., and Koch, M.A. (2004) Intraspecific diversification in North American *Boechea stricta* (= *Arabis drummondii*), *Boechea xdivaricarpa*, and *Boechea holboellii* (Brassicaceae) inferred from nuclear and chloroplast molecular markers—an integrative approach. *Am. J. Bot.*, 91, 2087–2101.
24. Song, B.H., Clauss, M.J., Pepper, A. *et al.* (2006) Geographic patterns of microsatellite variation in *Boechea stricta*, a close relative of *Arabidopsis*. *Mol. Ecol.*, 15, 357–369.
25. Asker, S. and Jerling, L. (1992) *Apomixis in Plants*; CRC Press, Boca Raton, Florida.
26. Dobeš, C., Koch, M. and Sharbel, T.F. (2006) Embryology, karyology, and modes of reproduction in the North American genus *Boechea* (Brassicaceae): a compilation of seven decades of research. *Ann. Missouri Bot. Gard.*, 93, 517–534.
27. Doyle, J.J. and Egan, A.N. (2010) Dating the origins of polyploidy events. *New Phytol.*, 186, 73–85.
28. Schranz, M.E., Dobeš, C., Koch, M.A. *et al.* (2005) Sexual reproduction, hybridization, apomixis, and polyploidization in the genus *Boechea* (Brassicaceae). *Am. J. Bot.*, 92, 1797–1810.
29. Song, B.H., Lee, C.R.R., Pais, A. *et al.* Rapid adaptive speciation and divergence population genetics in *Boechea*, a model system in ecology and evolution. in review.
30. Doddamani, D., Katta, M.A.V.S.K., Khan, A.W. *et al.* (2014) CicArMiSatDB: the chickpea microsatellite database. *BMC Bioinformatics*, 15, 212.
31. Krishnan, R.R., Sumathy, R., Bindroo, B.B. *et al.* (2014) MulSatDB: a first online database for mulberry microsatellites. *Trees*, 28, 1793–1799.
32. Ben Ayed, R., Ben Hassen, H., Ennouri, K. *et al.* (2016) OGDD (Olive Genetic Diversity Database): a microsatellite markers' genotypes database of worldwide olive trees for cultivar identification and virgin olive oil traceability. *Database*, 2016, bav090–bav099.
33. Anderson, E.C. and Thompson, E.A. (2002) A model-based method for identifying species hybrids using multilocus genetic data. *Genetics*, 160, 1217–1229.
34. Spielmann, A., Harris, S.A., Boshier, D.H. *et al.* (2015) Orchard: paternity program for autotetraploid species. *Mol Ecol Resour*, 15, 915–920.
35. Walling, C.A., Pemberton, J.M., Hadfield, J.D. *et al.* (2010) Comparing parentage inference software: reanalysis of a red deer pedigree. *Mol. Ecol.*, 19, 1914–1928.
36. Kantama, L., Sharbel, T.F., Schranz, M.E. *et al.* (2007) Diploid apomicts of the *Boechea holboellii* complex display large-scale chromosome substitutions and aberrant chromosomes. *Proc. Natl. Acad. Sci. USA.*, 104, 14026–14031.
37. Carman, J.G. (1997) Asynchronous expression of duplicate genes in angiosperms may cause apomixis, bispority, tetraspority, and polyembryony. *Biol. J. Linn. Soc.*, 61, 51–94.
38. Thiers, B. (2016) Index Herbariorum: A global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> (August 2016, last accessed).