

Milestones in understanding transport, sensing, and signaling of the plant nutrient phosphorus

Shu-Yi Yang ^{1,†} Wei-Yi Lin ^{2,†} Yi-Min Hsiao ^{3,†} and Tzzy-Jen Chiou ^{3,*}

1 Institute of Plant Biology, National Taiwan University, Taipei 106319, Taiwan

2 Department of Agronomy, National Taiwan University, Taipei 106319, Taiwan

3 Agricultural Biotechnology Research Center, Academia Sinica, Taipei 115201, Taiwan

*Author for correspondence: tjchiou@gate.sinica.edu.tw

†These authors contributed equally.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (<https://academic.oup.com/plcell/pages/General-Instructions>) is: Tzzy-Jen Chiou (tjchiou@gate.sinica.edu.tw).

Review

Abstract

As an essential nutrient element, phosphorus (P) is primarily acquired and translocated as inorganic phosphate (Pi) by plant roots. Pi is often sequestered in the soil and becomes limited for plant growth. Plants have developed a sophisticated array of adaptive responses, termed P starvation responses, to cope with P deficiency by improving its external acquisition and internal utilization. Over the past 2 to 3 decades, remarkable progress has been made toward understanding how plants sense and respond to changing environmental P. This review provides an overview of the molecular mechanisms that regulate or coordinate P starvation responses, emphasizing P transport, sensing, and signaling. We present the major players and regulators responsible for Pi uptake and translocation. We then introduce how P is perceived at the root tip, how systemic P signaling is operated, and the mechanisms by which the intracellular P status is sensed and conveyed. Additionally, the recent exciting findings about the influence of P on plant-microbe interactions are highlighted. Finally, the challenges and prospects concerning the interplay between P and other nutrients and strategies to enhance P utilization efficiency are discussed. Insights obtained from this knowledge may guide future research endeavors in sustainable agriculture.

Introduction

Phosphorus (P) is a macronutrient that is indispensable for growth, development, and reproduction because it is an essential component of several important biomolecules, such as DNA, RNA, ATP, NADPH, and phospholipids, and is also involved in respiration and photosynthesis (Raghothama 1999; Jiang *et al.* 2019a). Orthophosphate (Pi, $\text{H}_2\text{PO}_4^-/\text{HPO}_4^{2-}$) is the predominant form of P absorbed by plant roots through loading into the xylem and translocation into shoots. Still, it is quickly fixed by Fe^{3+} and Al^{3+} at low pH (<5 or acidic soils) or by tricalcium at high pH (>7 or alkaline soils), resulting in low availability and mobility in the soil (Holford 1997; Prathap *et al.* 2022). Modern agriculture leans heavily on P fertilizer to cope

with the low available soil Pi. However, due to the low efficiency of applied P absorbed by plants and the finite nature of P fertilizer, which is resourced from natural P rock reserves, increasing the P use efficiency (PUE) of crops has become an important issue to fulfill the demand for food security and agricultural sustainability (Baker *et al.* 2015; van de Wiel *et al.* 2016; Heuer *et al.* 2017). Endeavors to increase PUE will rely on knowledge of how plants sense and respond to environmental P levels.

In response to P deficiency, plants have developed sophisticated mechanisms to enhance the external acquisition and improve the internal utilization of P (Fig. 1), designated as P starvation responses (PSRs). Local PSRs are regulated by the external P concentration in the soil surrounding the root




Figure 1. An overview of adaptive P starvation responses in plants. There are 2 major categories of adaptive PSRs. One is to enhance the acquisition of external P, including root system architecture (RSA) modification; increased Pi release via acid/enzyme secretion; elevated Pi uptake through PHT1 transporters; and facilitated interactions with beneficial microbes such as arbuscular mycorrhizal fungi. The other is to improve utilization of internal P, including increased root-to-shoot growth ratio, PHO1-mediated root-to-shoot Pi reallocation, phospholipid replacement with galacto- and sulfolipids, ATP-conserving metabolic bypasses, and regulation of vacuolar Pi storage/release by PHT5 and VPE. Created with BioRender.com.

and are responsible for enhancing the acquisition of external P. Systemic PSRs depend on the adjustment of internal P concentration and improve the utilization of internal P through P recycling and reallocation via coordinating with external P availability (Bates and Lynch 1996; Thibaud et al. 2010; Lin et al. 2014; Chien et al. 2018). Regarding the enhanced acquisition of external P, the modification of the root system architecture, including an increase in the number of lateral roots and root hairs and a reduction of the primary root growth, allows plants to forage for P in the topsoil and enlarge root-soil surface efficiently (Bates and Lynch 2001; López-Bucio et al. 2003; Lynch 2011). In addition, increased high-affinity Pi transporter (PHT1) activity in the root surface facilitates Pi uptake. The organic acids, nucleases, and phosphatase secreted from the root enable Pi release from insoluble or organic matter (Taylor et al. 1993; Bariola et al. 1994; Raghothama 1999; Plaxton and Tran 2011). Interactions with beneficial microorganisms triggered by P starvation could also promote P acquisition (Oldroyd and Leyser 2020). Regarding the improved utilization of internal P, an increase in the root-to-shoot growth ratio, a reduction in shoot branching, and an increase in the erectness of leaves could allow plants to adjust growth and photosynthetic efficiency to adapt to P deficiency (Ruan et al. 2018; López-Bucio et al. 2003). The increase in the xylem loading activity of Pi facilitates root-to-shoot Pi reallocation (Hamburger et al. 2002). The replacement of phospholipids by galacto- and sulfolipids, the activation of metabolic bypasses to conserve ATP, and

the maintenance of cytosolic Pi homeostasis by modulating vacuolar Pi storage and release also occur (Härtel et al. 2001; Kelly and Dörmann 2002; Gaude et al. 2008; Plaxton and Tran 2011; Wang et al. 2012; Nakamura 2013; Okazaki et al. 2013; Liu et al. 2015, 2016; Wang et al. 2015; Xu et al. 2019).

In this review, we first summarize the historical key discoveries pertaining to the molecular players involved in P sensing, signaling, uptake, and utilization in plants since PHT1 Pi transporters were identified in 1996 (Fig. 2). We then highlight the major players and regulators in Pi uptake and translocation in coordinating PSRs. We illustrate local P sensing at the root tip, systemic P signaling, and intracellular P sensing. Furthermore, the recent exciting findings about P in plant-microbe interactions are discussed. Finally, the remaining challenges and perspectives on crosstalk with other nutrients and how to improve PUE are presented.

Historical overview—the key players in P transport, sensing, and signaling

This section briefly introduces the discoveries of key players in Pi transport, sensing, and signaling chronologically (Fig. 2). The details are elaborated afterward.

Since the identification of the genes encoding Pi transporters, molecular studies on P acquisition and its regulation have been inspired and enthusiastically pursued. Members




Figure 2. Milestones in understanding the genes involved in P transport, sensing, and signaling pathways. The genes with different roles in Pi uptake and translocation; local (root tip), systemic, and intracellular P sensing and signaling; and P-mediated plant-microbe interactions are depicted along the timeline and chronologically arranged according to the original publication dates.

of the plasma membrane-localized Pi transporter PHT1 family were first identified from Arabidopsis by yeast functional complementation in 1996 (Muchhal *et al.* 1996), followed by the second Pi transporter PHT2 in 1999, which is localized in the chloroplast (Daram *et al.* 1999). Later, in 2002, PHOSPHATE 1 (PHO1) was identified as a Pi efflux transporter mediating root-to-shoot Pi translocation (Hamburger *et al.* 2002). Then, in 2017, an additional group of Pi transporters belonging to the family of sulfate transporters (SULTRs), named SULTR-like phosphorus distribution transporter (SPDT), was uncovered (Yamaji *et al.* 2017). More recently, Pi transporters facilitating Pi translocation between the cytoplasm and vacuole have been identified, including Arabidopsis PHT5;1/VPT1 (VACUOLAR PHOSPHATE TRANSPORTER), a Pi influx transporter; and rice VPE1 and VPE2 (VACUOLAR PHOSPHATE EFFLUX), Pi efflux transporters (Liu *et al.* 2015, 2016; Xu *et al.* 2019). After uncovering the molecular identity of these Pi transporters, their regulation at the transcriptional and posttranslational levels was revealed. For example, upregulation of PHT1 transcription by low P is mediated by PHOSPHATE STARVATION RESPONSE (PHR) proteins binding to PHR1 binding sequence (P1BS, GNATATNC) (Rubio *et al.* 2001; Zhou *et al.* 2008), the proper targeting of PHT1 Pi transporters to plasma membranes requires PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR 1 (PHF1) (Gonzalez *et al.* 2005), and the protein abundance of PHT1 and PHO1 in response to Pi availability is controlled by PHOSPHATE 2 (PHO2/UBC24), a ubiquitin E2 conjugase, (Aung *et al.* 2006; Bari *et al.* 2006;

Liu *et al.* 2012; Huang *et al.* 2013), and/or NITROGEN LIMITATION ADAPTATION (NLA), a RING-type E3 ligase (Lin *et al.* 2013). It is worth mentioning that PHO1 and PHO2 were identified initially from genetic screenings through having significantly reduced or increased shoot Pi content, respectively (Poirier *et al.* 1991; Delhaize and Randall 1995).

To coordinate the P acquisition in the roots and P utilization at the whole-plant level for optimizing plant growth and development, sensing and signaling P availability are requisite. P sensing and signaling can be classified into local and systemic. The local signaling to sense external P at the root tips was shown to be mediated by Low Phosphate Root1 (LPR1) and LPR2, 2 multicopper oxidases (Svistoonoff *et al.* 2007). Other components, such as PHOSPHATE DEFICIENCY RESPONSES 2 (PDR2) and ALUMINUM ACTIVATED MALATE TRANSPORTER 1 (ALMT1), were later found to work together with LPR1 to regulate root meristem differentiation (Ticconi *et al.* 2009; Balzergue *et al.* 2017). On the other hand, systemic signaling plays an essential role in the improved utilization of internal P. From 1997 to 2000, non-coding RNAs (*TPS11*, *Mt4*, *At4* and *IPS1*) were discovered in different plant species that are highly induced by P starvation and share a conserved 22-nucleotide sequence (Liu *et al.* 1997; Burleigh and Harrison 1999; Martín *et al.* 2000; Shin *et al.* 2006). Their biological function remained unknown until the discovery of microRNA399 (miR399), the first miRNA identified to be involved in P starvation response (Fujii *et al.* 2005). Upon P starvation, miR399 is upregulated, which

suppresses *PHO2* expression and leads to the activation of Pi uptake and translocation (Aung et al. 2006; Bari et al. 2006; Chiou et al. 2006). *AT4/IPS1* homologs function to modulate the action of miR399 by sequestering miR399 through the 22 conserved nucleotides via a mechanism of target mimicry (Franco-Zorrilla et al. 2007). Interestingly, miR399 was found to serve as a shoot-to-root systemic signal (Lin et al. 2008; Pant et al. 2008).

With regard to intracellular Pi sensing and signaling, SPX proteins were discovered to be involved in regulating PSR genes in 2008 (Duan et al. 2008), and their role as P sensors through SPX-PHR interaction was reported in 2014 (Puga et al. 2014; Wang et al. 2014b). SPX proteins function as repressors to sequester PHRs from transcriptional activation of PSRs. Notably, a structure-function analysis revealed that inositol polyphosphate (InsP) and inositol pyrophosphate (PP-InsP) rather than Pi could be bound by SPX in vitro, but PP-InsPs could potentially serve as the pertinent signaling molecules in vivo (Wild et al. 2016). Later genetic analyses revealed that bis-diphosphoinositol tetrakisphosphate 1,5(PP)₂-InsP₄ (1,5-InsP₈) acts as a bona fide intracellular signaling molecule to regulate PSRs in plants (Dong et al. 2019a; Zhu et al. 2019).

Plants actively modulate the interactions with various microbes to adapt to the Pi availability. Strikingly, PHR was discovered to serve as a negative and positive regulator to mediate immune-related genes and arbuscular mycorrhizal-related genes, respectively (Castrillo et al. 2017; Isidra-Arellano et al. 2021; Shi et al. 2021; Das et al. 2022). The role of SPX in arbuscular mycorrhizal (AM) symbiosis was also revealed (Shi et al. 2021; Wang et al. 2021; Liao et al. 2022), suggesting the central role of SPX-PHR in integrating signals involved in nutritional response and biotic interaction.

Players in Pi uptake and translocation

Pi is initially acquired by the root epidermal and/or cortical cells and subsequently transported radially to the central vascular tissues where Pi is loaded into the xylem for translocation up to the shoot. Pi is then allocated among different tissues and distributed in organelles to sustain growth and development or stored inside vacuoles when excessive. Therefore, Pi transporters in the plasma membrane or organellar membranes are essential to facilitate Pi transport across membranes for uptake, distribution and remobilization (Fig. 3). While vegetative cells generally store Pi inside the vacuole, the primary form of P stored in the vacuole of seeds is inositol hexakisphosphate (InsP₆, phytate) mediated by MULTIDRUG RESISTANCE-ASSOCIATED PROTEIN 5 (MRP5), a vacuolar InsP₆ transporter (Nagy et al. 2009).

Pi transporters comprise different families, such as the phosphate transporter (PHT) family, the SYG1/Pho81/XPR1 (SPX) domain-containing protein family and the SPDT family. Members in the PHT family are classified into 5 types according to sequence identity and subcellular localization (Versaw and Garcia 2017), comprising PHT1 (plasma




Figure 3. Subcellular localization of players in Pi uptake and translocation. Arrows indicate the direction of Pi transport. Plasma membrane-localized PHT1 and SPDT are responsible for Pi uptake. PHT2 (chloroplast), PHT3 (mitochondria), and PHT4 (the Golgi apparatus, chloroplast, and nonphotosynthetic plastid) mediate Pi import or export from the corresponding organelles. Pi influx and efflux through the tonoplast are operated by PHT5 and VPE, respectively. PHO1 protein localizes to the plasma membrane and Golgi bearing the Pi efflux activity for Pi allocation. Pi translocators (PT) localized in the inner envelope membrane of plastids transport Pi in exchange with different substrates, including triose-phosphate/phosphate translocator (TPT), phosphoenolpyruvate/phosphate translocator (PPT), glucose-6-phosphate/phosphate translocator (GPT), and pentose xylulose-5-phosphate/phosphate translocator (XPT). Coupling with proton (H⁺) or sodium (Na⁺) transport is indicated. Created with BioRender.com.

membrane), PHT2 (chloroplast), PHT3 (mitochondria), PHT4 (the Golgi apparatus, chloroplast and nonphotosynthetic plastid) and PHT5 (vacuole). SPX domain-containing proteins are divided into 4 subfamilies: SPX, SPX-EXS (ERD1/XPR1/SYG1), SPX-MFS (Major Facilitator Superfamily), and SPX-RING (Really Interesting New Gene). Among them, SPX-MFS domain- and SPX-EXS domain-proteins have been identified as Pi transporters (e.g. members of the PHT5 family and PHO1, respectively) (Wang et al. 2012, 2015; Liu et al. 2015, 2016).

PHT1 transporters

PHT1 transporters are proton-Pi symporters and play a critical role in Pi acquisition from the soil and in Pi translocation between tissues (Nussaume et al. 2011; Remy et al. 2012; Lopez-Arredondo et al. 2014; Ayadi et al. 2015). There are 13 members in rice (*Oryza sativa*) and 9 in *Arabidopsis thaliana*. Most of them are expressed in roots and upregulated by the PHR transcription factor under P starvation (Karthikeyan et al. 2002; Mudge et al. 2002). In *Arabidopsis*, Pi uptake is mainly mediated by PHT1;1 and PHT1;4 (Shin et al. 2004; Remy et al. 2012; Ayadi et al. 2015), and root-to-shoot mobilization is facilitated by PHT1;8/PHT1;9 (Lapis-Gaza et al. 2014). In rice, OsPT2, a low-affinity Pi transporter exclusively expressed in the stele, facilitates the movement of Pi from

roots to shoots. In contrast, OsPT6 and OsPT8 function as high-affinity Pi transporters responsible for Pi uptake and subsequent translocation (Ai *et al.* 2009; Jia *et al.* 2011). OsPT9 and OsPT10, 2 other high-affinity Pi transporters, are also involved in Pi uptake (Wang *et al.* 2014a). The expression of these 5 rice PHT1 transporters were all induced by Pi starvation in the roots (Ai *et al.* 2009; Jia *et al.* 2011; Wang *et al.* 2014a). The members of *PHT1* are also expressed in the reproductive tissues responsible for anther and embryo development, for example, Arabidopsis *PHT1;6* and *PHT1;7* and rice *OsPht1;7* in the anthers (Mudge *et al.* 2002; Dai *et al.* 2022) and *OsPht1;4* in the embryo (Zhang *et al.* 2015).

Regarding the proper localization of PHT1 proteins, PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR 1 (PHF1), related to yeast SECRETORY 12 (SEC12), facilitates the exit of PHT1 proteins from the endoplasmic reticulum (ER) to plasma membranes in rice and Arabidopsis (Gonzalez *et al.* 2005; Bayle *et al.* 2011; Chen *et al.* 2011b). Additionally, the phosphorylation of rice OsPHT1;8 at Ser-517 by Casein Kinase 2 (OsCK2) interfered with the interaction between OsPHT1;8 and PHF1 and resulted in ER retention of OsPHT1;8 (Chen *et al.* 2015). Intriguingly, PHT1 protein abundance in the ER is controlled by PHO2-mediated degradation according to cellular P status, which determines the PHT1 amount directed to the plasma membrane (Huang *et al.* 2013).

Once at plasma membranes, PHT1 proteins are regulated by NLA-mediated degradation, a RING-type ubiquitin E3 ligase containing the SPX domain that can perceive cellular P levels (as discussed below) (Lin *et al.* 2013; Park *et al.* 2014; Yue *et al.* 2017; Yang *et al.* 2020). It was proposed that the cooperation between NLA and PHO2 controls the final protein level of PHT1 (Lin *et al.* 2013; Park *et al.* 2014). Intriguingly, different from Arabidopsis AtNLA, which is cleaved by P starvation-induced miR827 (Hsieh *et al.* 2009), rice *OsNLA1* is not the target of miR827, but its upstream open reading frame (uORF) is essential for its P-induced transcriptional expression (Lin *et al.* 2018; Yang *et al.* 2020).

PHT2/3/4 transporters

PHT2, PHT3, and PHT4 are localized to organelles, such as chloroplasts/plastids, mitochondria, and the Golgi apparatus. In Arabidopsis, PHT4;1, PHT4;2, and PHT2;1 control the Pi concentration in chloroplasts and regulate ATP synthesis and starch accumulation (Versaw and Harrison 2002; Karlsson *et al.* 2015; Miyaji *et al.* 2015). Similarly, mitochondrial localized PHT3 also plays a role in ATP metabolism (Zhu *et al.* 2012). Unexpectedly, PHT4;4 was shown to transport ascorbate into chloroplasts required to tolerate high light stress (Miyaji *et al.* 2015). In rice, OsPHT2;1 is also characterized as a chloroplast Pi influx transporter. Overexpression and loss of OsPHT2;1 result in altered Pi content in the shoot under low P conditions (Liu *et al.* 2020). How different organelles coordinate with each other to mediate cellular Pi homeostasis between the cytosol and organelles awaits more studies.

Vacuolar Pi transporters—PHT5 and VPE

Inside plant vegetative cells, the vacuole is the largest organelle and stores about 70% to 95% of the intracellular Pi (Yang *et al.* 2017), thus serving as an important reservoir in buffering cytoplasmic Pi concentrations. The Pi influx into the vacuole is operated by Arabidopsis PHT5 or VPT and rice SPX-MFS (Wang *et al.* 2012; Liu *et al.* 2015, 2016). These transporters contain N-terminal SPX and C-terminal MFS domains. OsSPX-MFS1 controls Pi homeostasis in leaves and mediates Pi influx to yeast vacuoles when heterologously expressed (Wang *et al.* 2012; Liu *et al.* 2016). Loss of Arabidopsis PHT5;1/VPT1 reduces total Pi level but results in necrotic leaves during P replenishment after starvation (Liu *et al.* 2015, 2016). Patch clamp analysis revealed a reduced Pi influx current in the vacuoles of *vpt1* mutants compared with wild-type ones (Liu *et al.* 2015). In addition, decreased vacuolar Pi and increased cytosolic Pi levels were observed in *pht5* mutants by ³¹P-NMR analysis or FRET-based Pi sensor (Liu *et al.* 2016; Sahu *et al.* 2020). Notably, the expression of *AtPHT1s* is affected in *pht5* mutants and *PHT5* overexpressor, implying that coordination between the storage capacity and acquisition activity would be achieved to maintain cytosolic Pi concentration (Liu *et al.* 2016). Conversely, rice vacuolar Pi efflux proteins VPE1/2 operate Pi efflux from the vacuole. The *vpe1/2* double mutant and overexpression plants show higher and lower vacuolar Pi levels, respectively (Xu *et al.* 2019). Unlike *VPE1/2*, the expression of *PHT5* is not responsive to P status, suggesting that the activity of PHT5 might be regulated at the protein level (Liu *et al.* 2016; Xu *et al.* 2019). Indeed, a recent study showed that the activity of PHT5 is impaired under Pi-deficient conditions or when the InsP/PP-InsP binding pocket in its SPX domain is mutated, thus linking the cellular P status to modulate the vacuolar Pi storage activity (Luan *et al.* 2022).

PHO1

Arabidopsis PHO1 and its close homolog PHO1;H1 with Pi efflux activity were able to facilitate the unloading of Pi into the xylem apoplastic space and allow for the subsequent translocation of Pi to the shoots (Poirier *et al.* 1991; Hamburger *et al.* 2002; Arpat *et al.* 2012; Liu *et al.* 2012). The significance of PHO1 in the root-to-shoot Pi allocation is supported by the higher Pi accumulation in roots and reduced Pi levels in shoots of *pho1* mutants in Arabidopsis and rice (Poirier *et al.* 1991; Secco *et al.* 2010; Che *et al.* 2020). AtPHO1 also facilitates the transfer of Pi from maternal tissues to developing seeds in the chalazal seed coat (Vogiatzaki *et al.* 2017). It is also expressed in leaf guard cells to play a role in the stomatal response to abscisic acid (ABA) (Zimmerli *et al.* 2012). In rice, OsPHO1;2 are involved in the allocation of Pi during grain filling (Che *et al.* 2020; Ma *et al.* 2021). In *Medicago truncatula*, PHO1 mediates the transfer of Pi from nodule-infected cells to bacteroids (Nguyen *et al.* 2021). Earlier results showed that the subcellular localization of Arabidopsis PHO1 is in endomembrane systems, such as

the Golgi and trans-Golgi networks, different from the plasma membrane localization of OsPHO1;2 (Arpat et al. 2012; Liu et al. 2012; Che et al. 2020). Nevertheless, a recent report showed that AtPHO1 is constitutively internalized from the plasma membrane (Vetal and Poirier 2023).

Arabidopsis PHO1 is negatively regulated at the translational level by its uORF (Reis et al. 2020) or at the transcriptional level by WRKY6 (Chen et al. 2009). In rice, the OsPHO1;2 mRNA translation can be enhanced by interacting with its cis-natural antisense transcript (Jabnourne et al. 2013; Reis et al. 2021). PHO1 has a cytosolic N-terminal SPX domain, 4 transmembrane domains, and a C-terminal EXS domain (Wege et al. 2016). The EXS domain is required for PHO1 membrane localization and Pi efflux activity (Wege et al. 2016). The SPX-spanning region of PHO1 facilitates its interaction with PHO2, which is responsible for PHO1 degradation (Liu et al. 2012). The absence of the SPX domain in PHO1 does not affect its efflux activity in transient expression assay. However, it fails to restore the impaired root-to-shoot Pi translocation activity in *pho1* mutants, indicating that the SPX domain is crucial for PHO1's functionality regulated by InsP/PP-InsP binding (Wege et al. 2016; Wild et al. 2016).

PHO2-mediated degradation of PHO1 occurs in endomembranes and is involved in vacuole proteolysis mediated by multi-vesicular bodies (Liu et al. 2012). Similarly, rice OsPHO2 also interacts with OsPHO1 to direct its degradation via a multi-vesicular body-mediated pathway (Wang et al. 2020), suggesting the conservation of the PHO2-PHO1 regulatory module among plants. The expression of *PHO2* is negatively regulated by miR399 (Bari et al. 2006) and positively regulated by the HD-ZIP III transcription factor PHB, which is inhibited by SHORT-ROOT (SHR) (Xiao et al. 2022). Studies have also revealed that OsCK2, a protein kinase, can enhance the degradation of OsPHO2 through the phosphorylation of OsPHO2 by OsCK2 α 3 (Wang et al. 2020).

SULTR-like phosphorus distribution transporter (SPDT)

SPDT was first shown to play a crucial role in regulating P allocation in rice (Yamaji et al. 2017). SPDT is a plasma-membrane-localized transporter that facilitates Pi transport and is expressed in the xylem region of both enlarged- and diffuse-vascular bundles of the nodes. Loss-of-function of SPDT in rice decreased P levels in grains but increased P levels in leaves. These findings indicate that SPDT functions as a switch in the rice node to allocate Pi to the grains preferentially. Likewise, barley HvSPDT, mainly expressed in the nodes, also plays an essential role in loading P into grains (Gu et al. 2022). In addition, the primary function of vascular cambium-localized Arabidopsis AtSPDT was shown to preferentially distribute Pi to the growing tissues (Ding et al. 2020).

Phosphate translocators

Phosphate translocators (PTs) localized in the inner envelope membrane of plastids play a crucial role in establishing the

metabolic connection between the plastid stroma and cytosol. Different from the Pi transporters mentioned above, these PT facilitate a precise counter-exchange of Pi with phosphorylated intermediates, including triose-phosphate/phosphate translocator, phosphoenolpyruvate/phosphate translocator, glucose-6-phosphate/phosphate translocator, and pentose xylulose-5-phosphate/phosphate translocator. They were reported to be involved in vegetative tissue and gametophyte development, redox signaling in leaves, and acclimation responses of photosynthesis (Młodzińska and Zboińska 2016; Fabiańska et al. 2019).

PHR1 as a central transcriptional regulator of PSRs

Arabidopsis PHR1 or rice PHR2 encoding an MYB coiled-coil transcription factor is a central positive regulator of the PSRs. PHR1 was initially identified from a genetic screen in Arabidopsis harboring a reporter gene driven by the IPS1 promoter highly induced by P starvation (Rubio et al. 2001). *phr1* mutants displayed multiple impairments in PSRs, including altered root-to-shoot Pi allocation, reduced accumulation of anthocyanin and carbohydrate, altered lipid composition and lipid remodeling genes, and impaired induction of several P starvation-induced (PSi) genes under P starvation (Rubio et al. 2001; Nilsson et al. 2007; Pant et al. 2015). Arabidopsis PHR1 transcript and protein accumulation are weakly responsive to P starvation; however, its activity is regulated by interacting with SPX proteins coupled with the cellular P status (Rubio et al. 2001; Puga et al. 2014).

PHR1 forms homodimers or heterodimers with its close homolog PHR1-like 1 (PHL1) and binds to P1BS elements present in the promoter of many PSi genes (Rubio et al. 2001; Bustos et al. 2010; Jiang et al. 2019b). Arabidopsis PHR1 and partially redundant PHL1 regulate the majority of transcriptional responses to P starvation, and P1BS elements are enriched in the promoter of PSi genes but not in P starvation-repressed (PSr) genes (Bustos et al. 2010). In P-starved *phr1phl1* mutants, nearly 70% of PSi genes and 50% of PSr genes showed decreased and increased expression, respectively, suggesting a direct and indirect regulation of transcriptional activation and repression responses (Bustos et al. 2010).

The other 3 Arabidopsis PHR1 homologs, PHL2, PHL3, and PHL4, have also been found to participate in PSRs to varying extents. PHL4 has a redundant function with PHR1/PHL1 but a minimal effect (Wang et al. 2018). PHL2 and PHL3 exhibit a specific interaction, and their expression is induced by P starvation, unlike PHR1 and PHL1. The transcriptomic analysis further suggests that PHR1/PHL1 and PHL2/PHL3 regulate various aspects of PSRs as separate regulatory modules (Wang et al. 2022). PHR1/PHL1 regulates genes involved in the cellular response to P starvation, Pi transport, and galactolipid biosynthetic process, whereas PHL2/PHL3 regulates genes related to the cellular response to P starvation, translation, and ribosome biogenesis. In rice, Arabidopsis PHR1 homolog *OsPHR2* plays a crucial role in Pi homeostasis, and

overexpressing *OsPHR2* increases *PSi* gene expression and results in P toxicity (Zhou *et al.* 2008). *PHR1* homologs participating in cellular Pi homeostasis have also been identified in various plant species (Wang *et al.* 2022), suggesting that *PHR* proteins have a universal and predominant role in regulating *PSRs*.

Despite not being as prevalent as *PHR1*, other transcription factors have been identified as regulators of *PSRs* to different extents. *MYB62*, as a negative regulator of *PSRs*, is involved in the interplay between *PSRs* and gibberellic acid responses (Devaiah *et al.* 2009), and *WRKY75* is a positive regulator of Pi uptake and a subset of *PSi* genes (Devaiah *et al.* 2007a). As mentioned, *WRKY6* is a repressor of *PHO1*. It interacts with 2 *W*-box motifs in the promoter of *PHO1* and is degraded by 26S proteasome to relieve the suppression upon P starvation (Chen *et al.* 2009). Basic Helix-Loop-Helix 32 and zinc finger of *Arabidopsis thaliana* 6 were also reported to have roles in regulating Pi content and root architecture (Chen *et al.* 2007; Devaiah *et al.* 2007b).

Intracellular P sensing and signaling

SPX proteins as intracellular P sensors

Proteins harboring a hydrophilic and poorly conserved *SPX* domain have been identified in many eukaryotes, such as yeasts, plants, and mammals (Secco *et al.* 2012). Many yeast *SPX* domain-containing proteins are involved in Pi homeostasis, including Pi transporters *Pho87*, *Pho90*, and *Pho91*; polyphosphate synthase subunits of vacuole transporter chaperone *VTC2-5*; the cyclin-dependent kinase inhibitor *Pho81*; and the glycerophosphocholine phosphodiesterase 1 (Secco *et al.* 2012; Desfougères *et al.* 2016). Yeast *Pho87* and *Pho90* are plasma membrane-localized low-affinity Pi influx transporters, and their *SPX* domains act as auto-inhibitory domains to regulate Pi import (Hürlimann *et al.* 2009). The *SPX* domain of vacuolar Pi exporter *Pho91* also regulates its activity through binding with PP-InsP (Potapenko *et al.* 2018). Additionally, PP-InsP-dependent binding on the *SPX* domains of yeast *VTC* complex is required for its polyphosphate polymerase activity (Gerasimaite *et al.* 2017).

Arabidopsis SPX1 and *SPX2* are nuclear proteins and function to repress *PHR1* activity through physical interaction in the nucleus to sequester *PHR1* from binding to the *P1BS* element of *PSi* genes (Fig. 4A) (Duan *et al.* 2008; Puga *et al.* 2014). Notably, the interaction between *PHR1* and *SPX1/2* highly depends on the cellular P status, with a strong interaction under P-sufficient conditions. Intriguingly, *SPX1/2* transcription is induced by *PHR1* during P starvation, revealing a negative feedback regulatory loop between *SPX1/2* and *PHR1* (Puga *et al.* 2014). This mechanism is conserved in rice; *OsSPX1/2* interacts with *OsPHR2* to suppress the binding of *OsPHR2* to the *P1BS* element (Zhou *et al.* 2008; Wang *et al.* 2014b). Other than in the nucleus, *Arabidopsis SPX4* and rice *OsSPX4* and *OsSPX6* interact with *PHR* in the cytoplasm to restrain the nuclear translocation of *PHR* when P is replete (Lv *et al.* 2014; Zhong *et al.* 2018). Upon P deficiency, *OsSPX4*


and *OsSPX6* undergo ubiquitin-mediated proteasomal degradation, thus releasing *OsPHR2*.

InsP₈ as a metabolic messenger for intracellular P signaling

Pi was initially considered to be a signaling molecule because of the attenuation of *PSRs* by phosphite (Phi, a nonmetabolized homolog of Pi) (Ticconi *et al.* 2001; Varadarajan *et al.* 2002). However, this concept was revised when the InsP/PP-InsP binding site of the *SPX* domain was revealed after solving the structure (Wild *et al.* 2016). The conserved amino acid residues defined as Pi binding cluster and lysine surface cluster together in the α -helix 2 and 4 of *SPX* domains form a positively charged surface, with higher affinities for InsP₆ and 5-InsP₇ than Pi. Moreover, the binding of InsP₆ and 5-InsP₇ but not Pi to *OsSPX4* can facilitate the interaction with *OsPHR2*, with 5-InsP₇ showing a higher affinity than InsP₆ (Wild *et al.* 2016), supporting the notion that PP-InsP rather than Pi is the signaling molecule representing intracellular P status.

In *Arabidopsis*, the synthesis and hydrolysis of 5-InsP₇ and 1,5-InsP₈ are catalyzed by 2 bifunctional enzymes, inositol 1,3,4-trisphosphate 5-/6-kinase *ITPK1/2* and diphosphoinositol pentakisphosphate kinase *VIH1/2*, respectively (Desai *et al.* 2014; Laha *et al.* 2015, 2019; Zhu *et al.* 2019; Whitfield *et al.* 2020; Riemer *et al.* 2021). The activity of *ITPK1/2* and *VIH1/2* can respond to changes in cellular ATP and Pi levels: high ATP favors kinase activity to synthesize 1,5-InsP₈, as ATP levels decrease in response to P starvation, and ADP phosphotransferase activity of *ITPK1* and phosphatase activity of *VIH1/2* are stimulated to catalyze the hydrolysis of 5-InsP₇ and 1,5-InsP₈ (Laha *et al.* 2019; Zhu *et al.* 2019; Whitfield *et al.* 2020; Riemer *et al.* 2021). Inhibition of *VIH1/2* phosphatase activity by Phi may explain the early observations of the Phi-mediated suppression of *PSRs*. *Arabidopsis vih1vih2* mutants displaying severe growth retardation exhibit an undetectable level of InsP₈ but overaccumulation of Pi and constitutive expression of *PSi* genes when grown under P-sufficient conditions, implying that InsP₈ is critical for plants to sense cellular P status (Dong *et al.* 2019b; Zhu *et al.* 2019). Furthermore, whereas the levels of InsP₆ and InsP₇ were relatively unchanged in response to P starvation, the level of InsP₈ was significantly decreased (Kuo *et al.* 2018; Dong *et al.* 2019a). 1,5-InsP₈ but not 5-InsP₇ can restore the interaction between *SPX1* and *PHR1* in the Pi-depleted cell lysate (Dong *et al.* 2019b). These findings indicate that InsP₈ is a signaling molecule communicating the cellular P level to control cellular Pi homeostasis through the *SPX* domain. A recent yeast study also revealed that 1,5-InsP₈ specifically inactivates *Pho81* through its *SPX* domain and that the reduction of 1,5-InsP₈ upon P starvation triggers the yeast *PHO* pathway (Chabert *et al.* 2023). The authors further proposed 1,5-InsP₈ as a universal signaling molecule in P sensing and signaling pathways among fungi, plants, and mammals (Chabert *et al.* 2023). It is worth noting that defective

A Intracellular and systemic P signaling



B P sensing at the root tip




Figure 4. Intracellular and systemic signaling and local sensing to P availability at root tips. **A)** During P sufficiency, the synthesis of InsP₈ is favored, which enables the interaction between SPX and PHR, suppressing PSi genes. Excess Pi in the cytosol is transported into the vacuole by PHT5. In the seeds, P is stored as InsP₆ (IP6) in the vacuole by MRP5. PHO2 and NLA promote the degradation of Pi transporters PHT1 and PHO1 via ubiquitination (ub). Upon P deficiency, InsP₈ (IP8) level is reduced. PHR is released and binds to the P1BS element as a dimer to activate PSi genes. PHF1 facilitates the targeting of PHT1 Pi transporters to plasma membranes. miR399 and miR827 repress PHO2 and NLA, respectively, to increase the abundance of PHT1 and PHO1. AT4/IPS1 transcripts can sequester miR399 to antagonize the effects of miR399. As indicated, miR399, miR827, sugar, strigolactone (SL), and cytokinin (tZ and iP) are potential systemic signals traveling via vascular tissues. **B)** Under P-sufficient conditions, ARSK1 is induced to phosphorylate RAPTOR1B, stabilizing the TOR1 complex and promoting root growth. Under P-deficient conditions, STOP1-ALMT1 modulates LPR1 activity, which controls Fe distribution in the root apical meristem, blocking SHR cell-to-cell movement by callose deposition. Meanwhile, the reception of CLE14 peptide activates POL/PLL1 phosphatases to inhibit root cell proliferation at transcript levels. Abbreviations: CALS, callose synthase; ROS, reactive oxygen species. Created with BioRender.com.

vacuolar storage of InsP_6 , as seen in *mrp5* mutants, results in elevated InsP_7 and InsP_8 levels (Desai *et al.* 2014; Riemer *et al.* 2021), suggesting InsP_6 compartmentation may also play a role in P signaling.

Recent studies have investigated the structure-functional relationship of the SPX-PP- InsP -PHR complex. InsP_8 promotes the binding between the InsP_8 -SPX1 complex and the coiled-coil domain of PHR1 (Ried *et al.* 2021). In the presence of InsP_6 (used as a substitute for InsP_8), OsSPX1 interacts with the MYB and the coiled-coil domains of OsPHR2 to form a complex with 1:1 stoichiometry, which disrupts the OsPHR2 dimer and further inhibits its binding to the P1BS element (Zhou *et al.* 2021). On the other hand, Guan *et al.* (2022) proposed that upon binding with InsP_6 , OsSPX2 forms a domain-swapped dimer and blocks the Myb and coiled-coil domains of OsPHR2 to abolish OsPHR2 dimerization and DNA binding; the SPX- InsP_6 -PHR complex was described as a 2:2:2 complex. This discrepancy might be due to the difference in conformational changes between different SPX proteins (Guan *et al.* 2022).

P sensing and signaling at the root tips

Root tips are critical in sensing environmental P status and regulating root meristem activities in response to external stimuli. P starvation causes premature differentiation in the root meristem, inhibiting primary root growth (Sanchez-Calderon *et al.* 2005). The increase of cell wall stiffness in the root transition zone can be observed 2 hours after plants transfer to P-deficient conditions. This results in the restriction of root cell extension and inhibition of root growth (Balzergue *et al.* 2017). LPR1 and LPR2, 2 multicopper oxidases with ferroxidase activity, which function with PDR2, a P5-type ATPase, are involved in the control of iron (Fe) distribution in the root apical meristem (RAM) (Reymond *et al.* 2006; Svistoonoff *et al.* 2007; Ticconi *et al.* 2009; Müller *et al.* 2015; Naumann *et al.* 2022). Intriguingly, Fe distribution in P-depleted roots overlays with the callose deposition pattern in the cell wall, restricting the movement of SHR for stem cell specification (Müller *et al.* 2015). Loss of LPR significantly reduced class III peroxidase activity, which might affect ROS-mediated callose deposition, leading to cell wall softening (Müller *et al.* 2015; Balzergue *et al.* 2017). These studies demonstrate the role of the LPR1-PDR2 module in mediating Fe- and peroxidase-dependent P starvation-induced root meristem differentiation (Fig. 4B) (Balzergue *et al.* 2017; Mora-Macias *et al.* 2017). Later studies revealed the involvement of PDR2-mediated ER stress-dependent autophagy in controlling Pi starvation-induced root growth retardation (Naumann *et al.* 2019).

Organic acids such as malate and citric acid in root exudates are needed for solubilizing metal ion-conjugated Pi in soils (Ryan *et al.* 2001). The activation of ALMT1, a malate efflux channel, by SENSITIVE TO PROTON RHIZOTOXICITY (STOP1) is not only critical for aluminum tolerance (Liu *et al.* 2009; Sawaki *et al.* 2009) but also influences malate export in the apoplast of root tips and Fe distribution. The

reduction of LPR1 mRNA in *stop1* and *almt1* mutants and ineffective malate treatment on the long primary root phenotype in *lpr1* indicate that ALMT1-mediated malate-triggered responses promote LPR1-dependent root meristem differentiation (Fig. 4B) (Balzergue *et al.* 2017).

Arabidopsis CLAVATA3/ENDOSPERM SURROUNDING REGION 14 (CLE14) is a low-P-induced peptide signal found specifically in the cortex, endodermis, and stele of RAM, functioning downstream of LPR1/LPR2. Applying CLE14 peptides can trigger RAM differentiation even under P-sufficient conditions but does not affect callose deposition. Instead, CLE14 perceived by CLAVATA 2 (CLV2) and PEP1 RECEPTOR2 (PEPR2) activates 2 phosphatases, POLTERGEIST (POL) and POL-LIKE1, to repress SHR and SCARECROW at mRNA levels (Fig. 4B) (Gutierrez-Alanis *et al.* 2017); however, the underlying mechanism is still missing. Moreover, the crosstalk of these signaling pathways with phytohormone-mediated root growth regulation remains to be studied.

Although Fe distribution in root tips determines the root meristem activities, an Fe-independent regulatory pathway is suggested because the reduction of root growth was observed before changes in Fe accumulation in the short-term responses to P deficiency. Arabidopsis-root-specific kinase 1 (ARSK1), a receptor-like kinase, is expressed explicitly in P-sufficient roots and can phosphorylate regulatory-associated protein of TOR 1B (RAPTOR1B), a scaffold protein of the target of rapamycin complex 1 (TOR1) that integrates environmental cues to modulate cellular growth (Wu *et al.* 2019). Overexpressing ARSK1 or phosphorylated RAPTOR1B inhibited P deficiency-triggered primary root retardation, indicating the crucial role of ARSK1 in maintaining primary root growth under P-sufficient conditions via stabilizing TOR1 complex (Fig. 4B) (Cho *et al.* 2023). Therefore, PDR2-LPR1 and STOP1-ALMT1 modules are activated by low P and collectively regulate RAM activities after gradually reducing ARSK1 levels.

Systemic signaling

In addition to local sensing and signaling, optimizing adaptive responses to P starvation requires communication between roots and shoots via systemic signals traveling long distances in the vasculature. Several molecules are considered long-distance signals that regulate PSRs systemically (Fig. 4A).

miR399 and miR827 are evolutionarily conserved and induced explicitly by P starvation (Fujii *et al.* 2005; Hsieh *et al.* 2009; Lin *et al.* 2018). Their levels are highly increased in phloem sap during P starvation (Pant *et al.* 2008, 2009), suggesting their potential roles as P starvation signals moving from shoot to root to coordinate the shoot Pi demand and root Pi acquisition and translocation activity. miR399 and its target gene *PHO2* are coexpressed in vascular tissues (Fujii *et al.* 2005; Aung *et al.* 2006; Chiou *et al.* 2006). Suppression of *PHO2* by overexpression of miR399 results in overaccumulation of Pi in shoots due to increased uptake and root-to-shoot translocation of Pi (Chiou *et al.* 2006).

The wild-type root grafted with miR399 overexpressing scion shows a high level of mature miR399, suppressing root *PHO2* transcripts and enhancing shoot Pi accumulation (Lin et al. 2008; Pant et al. 2008). Recently, the molecular basis underlying the miR399-mediated long-distance silencing was uncovered, by which miR399f/miR399f* duplex moves as a mobile entity and is unloaded in phloem pore pericycle in the root through plasmodesmata in a dose-dependent manner independent of its biogenesis, sequence context, and length (Chiang et al. 2023). Arabidopsis miR827 has also been reported to move from shoot to root (Huen et al. 2017). It is interesting to note that miR827 shifted its target preference during evolution; it targets *NLA* in Brassicaceae and Cleomaceae but targets *PHT5* homologs in other species (Lin et al. 2018).

Plants accumulate sugars and starch in leaves upon P starvation (Nielsen et al. 1998; Nilsson et al. 2007), and the induction of *PSi* genes increased with increasing concentrations of exogenous sugars (Karthikeyan et al. 2007). Increased sucrose loading and shoot-to-root translocation enhanced the expression of *PSi* genes (Lei et al. 2011; Dasgupta et al. 2014); in contrast, impairment of sucrose translocation via stem girdling of white lupin suppressed the expression of *PSi* genes in Pi-starved roots (Liu et al. 2005). These findings support the systemic role of sugar in P signaling, but the underlying mechanism requires further investigation.

Phytohormones also play a role in P signaling. Strigolactones (SLs), a group of carotenoid-derived terpenoid lactones, inhibit shoot branching in plants (Gomez-Roldan et al. 2008; Umehara et al. 2008). It has been demonstrated that the reduced shoot branching under P starvation could be attributed to the increased levels of SL in xylem sap (Kohlen et al. 2011), indicating a systemic role for SLs in regulating shoot branching during P starvation to coordinate the shoot development with nutrient supply from roots. N^6 -(Δ^2 -isopentenyl) adenine (iP) and *trans*-zeatin (tZ) are primary forms of natural isoprenoid cytokinins in Arabidopsis with strong physiological activities (Sakakibara 2006). Early studies proposed cytokinins to be negative regulators of PSRs because the exogenous application of cytokinin suppresses the expression of *PSi* genes in Arabidopsis (Martín et al. 2000; Wang et al. 2006). iP and tZ are predominantly present in the phloem and xylem, respectively, suggesting cytokinins are able to function as systemic signals (Hirose et al. 2007). Silva-Navas et al. (2019) reported that even though tZ has a greater substantial repression effect on PSR genes, an increase in the *cis*-zeatin (cZ):tZ ratio in P-deprived plants to improve root architecture and sustain necessary cytokinin responses was observed. Nevertheless, the detailed systemic role of cytokinins in response to P starvation remains to be determined.

Involvement of PSR factors in biotic interaction

Plants live with a diverse composition of microorganisms, and their relationship can be affected by plant internal nutrient levels or immune responses. It has been shown that plant

P status significantly affects the interaction with beneficial and pathogenic microbes (Paries and Gutjahr 2023). Moreover, recent evidence reveals the involvement of PSR genes in regulating biotic relationships, providing profound insights into plant-microbe interaction.

Regulation of arbuscular mycorrhizal (AM) symbiosis via SPX-PHR modules

Due to the scarcity of Pi in soil, plants not only modulate Pi uptake efficiency but also form beneficial mutualistic relationships with microbes to enhance Pi acquisition (Fig. 5). Arbuscular mycorrhizal fungi (AMF) are soil-born microbes that can establish endophytic symbiosis with more than 80% of land plant species and provide mineral nutrients via highly branched structures, termed arbuscules, in root cortical cells in exchange for carbon source for survival (Chiu and Paszkowski 2019). Interestingly, P1BS elements are present in the promoter regions of several AM symbiosis-responsive genes, such as symbiosis-induced *PHT1* genes (Chen et al. 2011a; Lota et al. 2013). Rice *PHR2* is activated in arbuscule-containing cells (Shi et al. 2021), and overexpressing *OsPHR2* induces symbiosis-responsive genes even without fungal infection (Shi et al. 2021; Das et al. 2022). Mutation at 3 rice *PHR* proteins severely impairs AMF colonization and arbuscular development. It alters the expression pattern of more than 60% symbiosis-responsive genes, including genes involved in pre-contact signaling, fungal entry, arbuscular development, and nutrient exchange (Shi et al. 2021; Das et al. 2022), demonstrating that *PHR* also functions in the center of AM symbiotic regulatory network (Fig. 5C).

As mentioned, modulation of *PHR* activity via the interaction with *SPX* proteins is widely validated in many plant species. Similarly, *Medicago SPX1* and *SPX3* are induced by low P, and *SPX*-*PHR* modules work in a P-dependent manner. In mycorrhizal roots, the activation of these 2 *SPX* genes is more confined to arbuscule-containing cells. Although the colonization efficiency in *spx1spx3* mutants is reduced, the enrichment of large arbuscules and the repression of arbuscular degeneration-related genes indicate that these 2 genes function redundantly in the control of arbuscular degradation (Wang et al. 2021). In addition, *MtSPX1* and *MtSPX3* control SL biosynthesis genes, the critical phytohormone in modulating shoot branching and AMF hyphae branching. The effects of *MtSPX1* and *MtSPX3* on the regulation of PSR and symbiosis were shown to be partially through the control of SL levels (Wang et al. 2021). In contrast, in rice *spx1/2/3/5* quadruple mutants, AMF colonization efficiency and arbuscular abundance are increased (Shi et al. 2021). In tomatoes, *Sspx1* mutants enhance AM colonization under Pi-replete conditions, whereas overexpression of *SISPX1* inhibits the formation of AM symbiosis (Liao et al. 2022). Further characterization is required to determine whether the distinct effects of *SPX* protein are due to the difference of family members or the interacting partners in different species.




Figure 5. Versatility of SPX-PHR modules in response to P status and plant-microbe interaction. The SPX-PHR module not only regulates adaptive responses to P availability **A**) and **B**), but it also regulates arbuscular mycorrhizal symbiosis **C**), and the responses to pathogen infection **D**), through controlling various sets of genes (see text for details). H in a circle represents a proton. Abbreviations: PHT1s, phosphate transporter 1; PHR, phosphate homeostasis regulator; SPX, SPX domain-containing protein; P1BS, phosphate 1-binding site; PSR, phosphate starvation response; RALFs, root-associated lipopeptides; FLS2, FLAGELLIN SENSING 2; MYC, Myc factor; S1P, site-one protease; SL, strigolactone; SYMRK, Symbiosis Receptor-like Kinase.

Control of rhizobium-mediated nitrogen fixation via P status

Rhizobium-mediated nitrogen (N) fixation is critical for N supply and cycling in agriculture and ecosystems. In nodulating legumes, P supply to rhizobia is essential for maintaining nodule function and mutualistic relationship. Depletion of environmental P impairs nodulation and N fixation (Tang *et al.* 2001; Valverde *et al.* 2002). Soybean (*Glycine max*) PHT1;14, a high-affinity PHT1 family member, is induced by low P treatment, mainly in the junction area between roots and young nodules and in the nodule vascular bundles. It is one of the key players in transporting plant Pi to nodules (Qin *et al.* 2012). In addition to taking up Pi indirectly from host plants, Pi can be acquired directly by nodules from growth environments. Soybean PHT1;1 is upregulated in the plasma membrane of the outer cortex and the fixation

zone of nodules in P-depleted roots. Overexpressing *GmPHT1;1* increases the fresh weight of nodules and the nitrogenase activity both under normal and P-deficient conditions, leading to the enhancement of N and P accumulation in plants and the final yield, supporting the significance of Pi transporters in Pi absorption by nodules and N fixation (Chen *et al.* 2019). *GmPHR1* and *GmPHR4* are expressed in the entire nodules and non-N-fixation region, respectively. Overexpressing *GmPHR1* enhances the expression of *GmPHT1;11* in nodules, leading to an increase in nodule Pi content and the size of the nodule, showing the importance of PHR-PHT1 module in nodules (Lu *et al.* 2020).

SPX proteins also regulate Pi homeostasis in nodules but not always via their association with PHR proteins. Soybean SPX5 and SPX8 are predominantly expressed in nodules and upregulated by P deficiency. Overexpressing either

GmSPX5 or *GmSPX8* increases the number and the fresh weight of nodules, resulting in higher N and P content in nodules than in the wild type (Zhuang et al. 2021; Xing et al. 2022). *GmNF-YC4*, a member of nuclear factor Y transcription factors, physically interacts with *GmSPX5*, and the association with *GmSPX5* enhances the binding capability of *GmNF-YC4* to the promoters of downstream target genes, such as asparagine synthetase-related genes. The phenotype of nodulation in overexpressing *GmNF-YC4* is largely overlaid with overexpressing *GmSPX5*, supporting the involvement of *GmSPX5-GmNF-YC4* in nodule development and functions (Zhuang et al. 2021).

Modulation of plant immunity against pathogens via P status

Excess P supply or Pi overaccumulation reduced the expression of defense-related genes, increasing the susceptibility to fungal pathogen *Magnaporthe oryzae* in rice (Campos-Soriano et al. 2020) and indicating the influence of external P in plant responses to pathogens. In contrast, P deficiency also suppresses plant immune response to promote the association with plant growth-promoting bacteria, which might facilitate Pi uptake efficiency or relieve the pressure of P scarcity (Tang et al. 2022). Thus, Pi uptake is tightly regulated to optimize the cellular P content for recruiting beneficial microbes. Among Arabidopsis PHT1 members, PHT1;4 is the major player in Pi uptake after recovering from P deficiency. Mutation at PHT1;4 reduces the susceptibility to the pathogen and upregulates pathogen elicitor-induced defense-related genes only under low P conditions (Dindas et al. 2022). Coincidentally, overexpressing rice PT8 reduces the pathogen resistance (Dong et al. 2019b). These studies reveal the direct link between PHT1-dependent Pi uptake and immune responses.

The influences of PSR genes on the structure of root microbe communities also point out the connection between Pi signaling and immune responses. Results of transcriptomic and chromatin immunoprecipitation-sequencing analyses disclose that PHR1 regulates not only PSR genes but also a set of salicylic acid- and jasmonic acid-inducible genes, which are under the transcriptional control of PHR1 via the direct binding on the promoter regions (Castrillo et al. 2017). Moreover, the flagellin peptide flg22-responsive defense genes are upregulated in *phr1phl1* mutants. FERONIA (FER), a *Catharanthus roseus* receptor-like kinase 1-like (CrRLK1L) family member, and its peptide ligand, rapid alkalization factor 23 (RALF23), can inhibit flg22-induced immune signaling (Stegmann et al. 2017). Interestingly, RALF23 and several RALF genes are also the direct targets of PHR1 that are upregulated by low P treatment. Studies of overexpressing RALF23 or *fer* mutants support the involvement of the FER-RALF23 module in P starvation-mediated immune inhibition. All these observations together revealed that PHR1 has a negative role in regulating immune responses through the enhancement of the FER-RALK module (Fig. 5D) (Tang et al. 2022).

Challenges and perspectives

Facing the limitation of Pi availability in nature, an increase in P fertilizer supply promotes crop yield, but plants acquire less than 30% of it (Vance et al. 2003). Tremendous progress has been made in establishing the regulatory network of Pi sensing and signaling over years of study, providing opportunities to accelerate high PUE crop development. Yet, many fundamental questions and gaps remain if such information is to be usefully translated to agriculture and ecological systems. For example, the underlying mechanisms of apoplastic responses to low external P to adjust the growth in root tips have been established; however, it is still unclear whether other plant cells can also sense extracellular P availability and how. Concerning intracellular sensing, the perception of InsP₈ by SPX proteins determines the activation of PHR-mediated PSRs (Wild et al. 2016; Ried et al. 2021). Multiple enzymes are required to catalyze PP-InsPs synthesis, but understanding of the coordination of the enzyme activities in response to extra- or intracellular P levels remains limited. In addition, the direct link between external and internal P sensing is still missing.

Many studies have focused on increasing Pi acquisition to improve PUE. These include modifying root architecture to enlarge root surface area for Pi scavenging (Jia et al. 2018; Sun et al. 2018; Wang et al. 2019), enhancing the expression of PHTs to increase Pi acquisition and translocation (Song et al. 2014; Ouyang et al. 2016; Pontigo et al. 2023), modulating the secretion of root exudates involved in soil Pi solubilization (Pang et al. 2018; Robles-Aguilar et al. 2019; Wen et al. 2019) and promoting the interaction of plants with beneficial microbes to extend Pi source (Wen et al. 2019; Yahya et al. 2021).

Natural germplasms are valuable genetic resources for identifying novel alleles to improve the current cultivars' PUEs. Phenotypic and molecular characterization of various germplasm lines in crop species identified many quantitative trait loci associated with low P tolerance. Genome-wide association studies of soybean germplasms pinpointed a variant in the uORF of *GmPHF1* that contributes to P acquisition diversity (Guo et al. 2022b). Rice phosphate-starvation tolerance 1 (PSTOL1) is an enhancer of root growth and Pi uptake of aus-type cultivars grown in poor soils (Wissuwa et al. 2002; Gamuyao et al. 2012). Sequencing PSTOL1 alleles in wild rice (*Oryza rufipogon*) grown in diverse environments found novel alleles associated with high Pi content, which might be used for breeding high PUE rice cultivars (Neelam et al. 2017). Ectopic expression of *OsPSTOL1* in wheat enhanced plant growth and root plasticity and promoted the induction of PHR-mediated P-starvation responses under low P conditions (Kettenburg et al. 2023). However, overexpressing wheat endogenous *TaPSTOL* homolog decreased PUE due to reduced yield but increased P accumulation in grains (Milner et al. 2018). Genome-wide association studies of soybean germplasm identified a variant in the uORF of soybean PHF1 as a key determinant for P acquisition

efficiency in the field (Guo *et al.* 2022b). Nevertheless, most studies were conducted in greenhouses. Reproducing the traits in field conditions and integrating the knowledge on P sensing/uptake/recycling to improve plant PUE will be the main challenges for further application in breeding programs.

Coordinated acquisition and distribution of P and other nutrients are also required for optimizing plant growth and development. Thus, the interplay between P and other nutrients must be considered when modulating PUE via genetic strategies. Perception of nitrate strengthens the interaction of the nitrate sensor, OsNRT1;1B, with OsSPX4, resulting in OsSPX4 degradation, which leads to the activation of OsPHR2-mediated PSRs and NIN-like protein NLP3-mediated nitrate signalings (Hu *et al.* 2019). Meanwhile, Medici *et al.* (2019) identified PHO2 as another integrator of N signals to PSR pathways. These findings give insights into nitrate-regulated PSR responses, but little is known about the effects of ammonium and the role of P signaling in nitrate responses. Additionally, diverse functions of SPX and PHR homologs imply the different SPX-PHR modules present *in vivo* to exert transcriptional regulation. The underlying mechanism for controlling SPX-PHR formation and their roles in the interplay between P and other nutrient signaling requires further investigation.

As mentioned, Fe mediates P sensing in the root tips (Fig. 4B). Reciprocally, transcriptomic analysis also revealed the importance of P level in regulating Fe deficiency- or excess-responsive genes (Sanchez-Calderon *et al.* 2005; Thibaud *et al.* 2010). Rice HRZ, a Fe starvation-induced ubiquitin E3 ligase, mediates the protein degradation of OsPHR2. Conversely, OsPHR2 negatively regulates the HRZ transcript level to modulate Fe-deficient responses (Guo *et al.* 2022a), pointing out the regulatory loop in P-Fe interaction. Zn-dependent regulation of Pi accumulation has also been observed (Khan *et al.* 2014; Kisko *et al.* 2018), but knowledge about the underlying signaling and the coordination of P and Zn homeostasis is still lacking.

In terms of the role of plant P status in plant-microbe interaction, it is crucial to integrate PSR and immune responses for recruiting a specified set of microbes for survival. How to identify friends or foes and balance pathogen resistance and symbiotic interaction are fascinating topics about which little is known. Interestingly, PHR can directly activate genes required for AM symbiosis but repress genes involved in defense, suggesting that PHR appears to be a hub that coordinates immunity and symbiosis. However, the diversity and complexity of soil microbe composition challenge the integration of genetic studies and crop breeding programs. Large-scale studies and field trials are required to elucidate the interaction between P signaling and immune responses.

In nature, plants undergo multiple stresses. The rising temperature and CO₂ levels aggravate the pressure and frequency of such stresses. However, the coordination between Pi signaling and other stress responses is rarely

discussed. Recent research reported that overexpressing a chloroplast-localized Pi importer resulted in the overaccumulation of chloroplast Pi and phytic acids, leading to growth retardation under CO₂-elevated conditions (Bouain *et al.* 2022). This result points out the importance of chloroplast Pi homeostasis in maintaining plant growth when plants face elevated CO₂ levels. Despite being a challenging topic, more studies must be conducted to investigate the interplay between P and other stresses and underlying mechanisms, which need to be scrutinized in the field when applied to crop improvement.

Acknowledgments

We thank the anonymous reviewers for their constructive comments. We apologize to our colleagues whose work could not be cited due to space constraints.

Author contributions

S.-Y.Y., W.-Y.L., Y.-M.H., and T.-J.C. conceptualized the review outline. S.-Y.Y., W.-Y.L. and Y.-M.H. prepared the figures and wrote the original draft, in which S.-Y.Y. contributed to the introduction and P transport, Y.-M.H. to intracellular P sensing and signaling and W.-Y.L. to local sensing, biotic interaction, and perspectives. T.-J.C. compiled, reviewed, and edited the manuscript. All authors contributed to the critical revision and its final approval.

Funding

We thank Academia Sinica (AS-GCS-112-L03) for funding to T.-J.C. laboratory and the National Science and Technology Council, Taiwan for funding to T.-J.C. (MOST 110-2311-B-001 -021 -MY3, NSTC 112-2311-B-001 -040 -MY3), S.-Y.Y. (MOST 111-2311-B-002 -005 -MY3) and W.-Y.L. (MOST 111-2313-B-002 -009 -MY3) laboratories. Y.-M.H. was supported by the Taiwan International Graduate Program (TIGP) scholarship.

Conflict of interest statement. None declared.

References

- Ai P, Sun S, Zhao J, Fan X, Xin W, Guo Q, Yu L, Shen Q, Wu P, Mille A, *et al.* Two rice phosphate transporters, OsPht1;2 and OsPht1;6 have different functions and kinetic properties in uptake and translocation. *Plant J.* 2009;57(5):798–809. <https://doi.org/10.1111/j.1365-313X.2008.03726.x>
- Arpat AB, Magliano P, Wege S, Rouached H, Stefanovic A, Poirier Y. Functional expression of PHO1 to the Golgi and trans-Golgi network and its role in export of inorganic phosphate. *Plant J.* 2012;71(3):479–491. <https://doi.org/10.1111/j.1365-313X.2012.05004.x>
- Aung K, Lin SI, Wu CC, Huang YT, Su CL, Chiou TJ. Pho2, a phosphate overaccumulator, is caused by a nonsense mutation in a microRNA399 target gene. *Plant Physiol.* 2006;141(3):1000–1011. <https://doi.org/10.1104/pp.106.078063>

- Ayadi A, David P, Arrighi JF, Chiarenza S, Thibaud MC, Nussaume L, Marin E.** Reducing the genetic redundancy of Arabidopsis PHOSPHATE TRANSPORTER1 transporters to study phosphate uptake and signaling. *Plant Physiol.* 2015;**167**(4):1511–1526. <https://doi.org/10.1104/pp.114.252338>
- Baker A, Ceasar SA, Palmer AJ, Paterson JB, Qi W, Muench SP, Baldwin SA.** Replace, reuse, recycle: improving the sustainable use of phosphorus by plants. *J Exp Bot.* 2015;**66**(12):3523–3540. <https://doi.org/10.1093/jxb/erv210>
- Balergue C, Dartevelle T, Godon C, Laugier E, Meisrimler C, Teulon JM, Creff A, Bissler M, Brouchoud C, Hagege A, et al.** Low phosphate activates STOP1-ALMT1 to rapidly inhibit root cell elongation. *Nat Commun.* 2017;**8**(1):15300. <https://doi.org/10.1038/ncomms15300>
- Bari R, Datt Pant B, Stitt M, Scheible W-R.** PHO2, microRNA399, and PHR1 define a phosphate-signaling pathway in plants. *Plant Physiol.* 2006;**141**(3):988–999. <https://doi.org/10.1104/pp.106.079707>
- Bariola PA, Howard CJ, Taylor CB, Verburg MT, Jaglan VD, Green PJ.** The Arabidopsis ribonuclease gene RNS1 is tightly controlled in response to phosphate limitation. *Plant J.* 1994;**6**(5):673–685. <https://doi.org/10.1046/j.1365-313x.1994.6050673.x>
- Bates TR, Lynch JP.** Stimulation of root hair elongation in Arabidopsis thaliana by low phosphorus availability. *Plant, Cell Environ.* 1996;**19**(5):529–538. <https://doi.org/10.1111/j.1365-3040.1996.tb00386.x>
- Bates TR, Lynch JP.** Root hairs confer a competitive advantage under low phosphorus availability. *Plant Soil.* 2001;**236**(2):243–250. <https://doi.org/10.1023/A:1012791706800>
- Baile V, Arrighi JF, Creff A, Nespoulous C, Vialaret J, Rossignol M, Gonzalez E, Paz-Ares J, Nussaume L.** Arabidopsis thaliana high-affinity phosphate transporters exhibit multiple levels of posttranslational regulation. *Plant Cell.* 2011;**23**(4):1523–1535. <https://doi.org/10.1105/tpc.110.081067>
- Bouain N, Cho H, Sandhu J, Tuiwong P, Prom-u-thai C, Zheng LQ, Shahzad Z, Rouached H.** Plant growth stimulation by high CO₂ depends on phosphorus homeostasis in chloroplasts. *Curr Biol.* 2022;**32**(20):4493–4500. <https://doi.org/10.1016/j.cub.2022.08.032>
- Burleigh SH, Harrison MJ.** The down-regulation of Mt4-like genes by phosphate fertilization occurs systemically and involves phosphate translocation to the shoots. *Plant Physiol.* 1999;**119**(1):241–248. <https://doi.org/10.1104/pp.119.1.241>
- Bustos R, Castrillo G, Linhares F, Puga MI, Rubio V, Pérez-Pérez J, Solano R, Leyva A, Paz-Ares J.** A central regulatory system largely controls transcriptional activation and repression responses to phosphate starvation in Arabidopsis. *PLoS Genet.* 2010;**6**(9):e1001102. <https://doi.org/10.1371/journal.pgen.1001102>
- Campos-Soriano L, Bundo M, Bach-Pages M, Chiang SF, Chiou TJ, San Segundo B.** Phosphate excess increases susceptibility to pathogen infection in rice. *Mol Plant Pathol.* 2020;**21**(4):555–570. <https://doi.org/10.1111/mpp.12916>
- Castrillo G, Teixeira PJPL, Paredes SH, Law TF, de Lorenzo L, Feltcher ME, Finkel OM, Breakfield NW, Mieczkowski P, Jones CD, et al.** Root microbiota drive direct integration of phosphate stress and immunity. *Nature.* 2017;**543**(7646):513–518. <https://doi.org/10.1038/nature21417>
- Chabert V, Kim G, Qiu D, Liu G, Mayer L, Jamsheer K, Jessen H, Mayer A.** Inositol pyrophosphate dynamics reveals control of the yeast phosphate starvation program through 1,5-IP₈ and the SPX domain of Pho81. *Elife.* 2023;**12**:RP87956. <https://doi.org/10.7554/eLife.87956.2>
- Che J, Yamaji N, Miyaji T, Mitani-Ueno N, Kato Y, Shen R, Ma J.** Node-localized transporters of phosphorus essential for seed development in rice. *Plant Cell Physiol.* 2020;**61**(8):1387–1398. <https://doi.org/10.1093/pcp/pcaa074>
- Chen AQ, Gu MA, Sun SB, Zhu LL, Hong SA, Xu GH.** Identification of two conserved cis-acting elements, MYCS and P1BS, involved in the regulation of mycorrhiza-activated phosphate transporters in eudicot species. *New Phytol.* 2011a;**189**(4):1157–1169. <https://doi.org/10.1111/j.1469-8137.2010.03556.x>
- Chen J, Liu Y, Ni J, Wang Y, Bai Y, Shi J, Gan J, Wu Z, Wu P.** OsPHF1 regulates the plasma membrane localization of low- and high-affinity inorganic phosphate transporters and determines inorganic phosphate uptake and translocation in rice. *Plant Physiol.* 2011b;**157**(1):269–278. <https://doi.org/10.1104/pp.111.181669>
- Chen L, Qin L, Zhou L, Li X, Chen Z, Sun L, Wang W, Lin Z, Zhao J, Yamaji N, et al.** A nodule-localized phosphate transporter GmPT7 plays an important role in enhancing symbiotic N₂ fixation and yield in soybean. *New Phytol.* 2019;**221**(4):2013–2025. <https://doi.org/10.1111/nph.15541>
- Chen J, Wang Y, Wang F, Yang J, Gao M, Li C, Liu Y, Liu Y, Yamaji N, Ma JF, et al.** The rice CK2 kinase regulates trafficking of phosphate transporters in response to phosphate levels. *Plant Cell.* 2015;**27**(3):711–723. <https://doi.org/10.1105/tpc.114.135335>
- Chen YF, Li LQ, Xu Q, Kong YH, Wang H, Wu WH.** The WRKY6 transcription factor modulates PHOSPHATE1 expression in response to low Pi stress in Arabidopsis. *Plant Cell.* 2009;**21**(11):3554–3566. <https://doi.org/10.1105/tpc.108.064980>
- Chen ZH, Nimmo GA, Jenkins GI, Nimmo HG.** BHLH32 modulates several biochemical and morphological processes that respond to Pi starvation in Arabidopsis. *Biochem J.* 2007;**405**(1):191–198. <https://doi.org/10.1042/bj20070102>
- Chiang CP, Li LJ, Chiou TJ.** Dose-dependent long-distance movement of microRNA399 duplex regulates phosphate homeostasis in Arabidopsis. *New Phytol.* 2023;**240**(2):802–814. <https://doi.org/10.1111/nph.19182>
- Chien PS, Chiang CP, Leong SJ, Chiou TJ.** Sensing and signaling of phosphate starvation: from local to long distance. *Plant Cell Physiol.* 2018;**59**(9):1714–1722. <https://doi.org/10.1093/pcp/pcy148>
- Chiou TJ, Aung K, Lin SI, Wu CC, Chiang SF, Su CL.** Regulation of phosphate homeostasis by microRNA in Arabidopsis. *Plant Cell.* 2006;**18**(2):412–421. <https://doi.org/10.1105/tpc.105.038943>
- Chiu CH, Paszkowski U.** Mechanisms and impact of symbiotic phosphate acquisition. *Cold Spring Harb Perspect Biol.* 2019;**11**(6):a034603. <https://doi.org/10.1101/cshperspect.a038281>
- Cho H, Banf M, Shahzad Z, Van Leene J, Bossi F, Ruffel S, Bouain N, Cao P, Krouk G, De Jaeger G, et al.** ARSK1 activates TORC1 signaling to adjust growth to phosphate availability in Arabidopsis. *Curr Biol.* 2023;**33**(9):1778–1786. <https://doi.org/10.1016/j.cub.2023.03.005>
- Dai C, Dai X, Qu H, Men Q, Liu J, Yu L, Gu M, Xu G.** The rice phosphate transporter OsPHT1; 7 plays a dual role in phosphorus redistribution and anther development. *Plant Physiol.* 2022;**188**(4):2272–2288. <https://doi.org/10.1093/plphys/kiac030>
- Daram P, Brunner S, Rausch C, Steiner C, Amrhein N, Bucher M.** Pht2; 1 encodes a low-affinity phosphate transporter from Arabidopsis. *Plant Cell.* 1999;**11**(11):2153–2166. <https://doi.org/10.1105/tpc.11.11.2153>
- Das D, Paries M, Hobecker K, Gigl M, Dawid C, Lam H, Zhang J, Chen M, Gutjahr C.** PHOSPHATE STARVATION RESPONSE transcription factors enable arbuscular mycorrhiza symbiosis. *Nat Commun.* 2022;**13**(1):447. <https://doi.org/10.1038/s41467-022-27976-8>
- Dasgupta K, Khadilkar AS, Sulpice R, Pant B, Scheible W-R, Fisahn J, Stitt M, Ayre BG.** Expression of sucrose transporter cDNAs specifically in companion cells enhances phloem loading and long-distance transport of sucrose but leads to an inhibition of growth and the perception of a phosphate limitation. *Plant Physiol.* 2014;**165**(2):715–731. <https://doi.org/10.1104/pp.114.238410>
- Delhaize ER, Randall PJ.** Characterization of a phosphate-accumulator mutant of Arabidopsis thaliana. *Plant Physiol.* 1995;**107**(1):207–213. <https://doi.org/10.1104/pp.107.1.207>
- Desai M, Rangarajan P, Donahue JL, Williams SP, Land ES, Mandal MK, Phillippy BQ, Perera IY, Raboy V, Gillaspay GE.** Two inositol hexakisphosphate kinases drive inositol pyrophosphate synthesis in plants. *Plant J.* 2014;**80**(4):642–653. <https://doi.org/10.1111/tj.12669>
- Desfougères Y, Gerasimaitė RU, Jessen HJ, Mayer A.** Vtc5, a novel subunit of the vacuolar transporter chaperone complex, regulates

- polyphosphate synthesis and phosphate homeostasis in yeast. *J Biol Chem*. 2016;**291**(42):22262–22275. <https://doi.org/10.1074/jbc.M116.746784>
- Devaiah BN, Karthikeyan AS, Raghothama KG.** WRKY75 transcription factor is a modulator of phosphate acquisition and root development in Arabidopsis. *Plant Physiol*. 2007a;**143**(4):1789–1801. <https://doi.org/10.1104/pp.106.093971>
- Devaiah BN, Madhuvanathi R, Karthikeyan AS, Raghothama KG.** Phosphate starvation responses and gibberellic acid biosynthesis are regulated by the MYB62 transcription factor in Arabidopsis. *Mol Plant*. 2009;**2**(1):43–58. <https://doi.org/10.1093/mp/ssn081>
- Devaiah BN, Nagarajan VK, Raghothama KG.** Phosphate homeostasis and root development in Arabidopsis are synchronized by the zinc finger transcription factor ZAT6. *Plant Physiol*. 2007b;**145**(1):147–159. <https://doi.org/10.1104/pp.107.101691>
- Dindas J, DeFalco TA, Yu G, Zhang L, David P, Bjornson M, Thibaud MC, Custodio V, Castrillo G, Nussaume L, et al.** Direct inhibition of phosphate transport by immune signaling in Arabidopsis. *Curr Biol*. 2022;**32**(2):488–495.e485. <https://doi.org/10.1016/j.cub.2021.11.063>
- Ding G, Lei GJ, Yamaji N, Yokosho K, Mitani-Ueno N, Huang S, Ma JF.** Vascular cambium-localized AtSPDT mediates xylem-to-phloem transfer of phosphorus for its preferential distribution in Arabidopsis. *Mol Plant*. 2020;**13**(1):99–111. <https://doi.org/10.1016/j.molp.2019.10.002>
- Dong J, Ma G, Sui L, Wei M, Satheesh V, Zhang R, Ge S, Li J, Zhang T-E, Wittwer C, et al.** Inositol pyrophosphate InsP8 acts as an intracellular phosphate signal in Arabidopsis. *Mol Plant*. 2019a;**12**(11):1463–1473. <https://doi.org/10.1016/j.molp.2019.08.002>
- Dong Z, Li W, Liu J, Li L, Pan S, Liu S, Gao J, Liu L, Liu X, Wang GL, et al.** The rice phosphate transporter protein OsPT8 regulates disease resistance and plant growth. *Sci Rep*. 2019b;**9**(1):5408. <https://doi.org/10.1038/s41598-019-41718-9>
- Duan K, Yi K, Dang L, Huang H, Wu W, Wu P.** Characterization of a sub-family of Arabidopsis genes with the SPX domain reveals their diverse functions in plant tolerance to phosphorus starvation. *Plant J*. 2008;**54**(6):965–975. <https://doi.org/10.1111/j.1365-313X.2008.03460.x>
- Fabińska I, Bucher M, Häusler R.** Intracellular phosphate homeostasis—a short way from metabolism to signaling. *Plant Sci*. 2019;**286**:57–67. <https://doi.org/10.1016/j.plantsci.2019.05.018>
- Franco-Zorrilla JM, Valli A, Todesco M, Mateos I, Puga MI, Rubio-Somoza I, Leyva A, Weigel D, García JA, Paz-Ares J.** Target mimicry provides a new mechanism for regulation of microRNA activity. *Nat Genet*. 2007;**39**(8):1033–1037. <https://doi.org/10.1038/ng2079>
- Fujii H, Chiou TJ, Lin SI, Aung K, Zhu JK.** A miRNA involved in phosphate-starvation response in Arabidopsis. *Curr Biol*. 2005;**15**(22):2038–2043. <https://doi.org/10.1016/j.cub.2005.10.016>
- Gamuyao R, Chin JH, Pariasca-Tanaka J, Pesaresi P, Catausan S, Dalid C, Slamet-Loedin I, Tecson-Mendoza EM, Wissuwa M, Heuer S.** The protein kinase Pstol1 from traditional rice confers tolerance of phosphorus deficiency. *Nature*. 2012;**488**(7412):535. <https://doi.org/10.1038/nature11346>
- Gaude N, Nakamura Y, Scheible N, Ohta H, Dörmann P.** Phospholipase C5 (NPC5) is involved in galactolipid accumulation during phosphate limitation in leaves of Arabidopsis. *Plant J*. 2008;**56**(1):28–39. <https://doi.org/10.1111/j.1365-313X.2008.03582.x>
- Gerasimaite R, Pavlovic I, Capolicchio S, Hofer A, Schmidt A, Jessen HJ, Mayer A.** Inositol pyrophosphate specificity of the SPX-dependent polyphosphate polymerase VTC. *ACS Chem Biol*. 2017;**12**(3):648–653. <https://doi.org/10.1021/acscchembio.7b00026>
- Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pagès V, Dun EA, Pillot J-P, Letisse F, Matusova R, Danoun S, Portais J-C, et al.** Strigolactone inhibition of shoot branching. *Nature*. 2008;**455**(7210):189–194. <https://doi.org/10.1038/nature07271>
- Gonzalez E, Solano R, Rubio V, Leyva A, Paz-Ares J.** PHOSPHATE TRANSPORTER TRAFFIC FACILITATOR1 is a plant-specific SEC12-related protein that enables the endoplasmic reticulum exit of a high-affinity phosphate transporter in Arabidopsis. *Plant Cell*. 2005;**17**(12):3500–3512. <https://doi.org/10.1105/tpc.105.036640>
- Gu M, Huang H, Hisano H, Ding G, Huang S, Mitani-Ueno N, Yokosho K, Sato K, Yamaji N, Ma J.** A crucial role for a node-localized transporter, HvSPDT, in loading phosphorus into barley grains. *New Phytol*. 2022;**234**(4):1249–1261. <https://doi.org/10.1111/nph.18057>
- Guan Z, Zhang Q, Zhang Z, Zuo J, Chen J, Liu R, Savarin J, Broger L, Cheng P, Wang Q, et al.** Mechanistic insights into the regulation of plant phosphate homeostasis by the rice SPX2—PHR2 complex. *Nat Commun*. 2022;**13**(1):1581. <https://doi.org/10.1038/s41467-022-29275-8>
- Guo MN, Ruan WY, Zhang YB, Zhang YX, Wang XQ, Guo ZH, Wang L, Zhou T, Paz-Ares J, Yi KK.** A reciprocal inhibitory module for Pi and iron signaling. *Mol Plant*. 2022a;**15**(1):138–150. <https://doi.org/10.1016/j.molp.2021.09.011>
- Guo ZL, Cao HR, Zhao J, Bai S, Peng WT, Li J, Sun LL, Chen LY, Lin ZH, Shi C, et al.** A natural uORF variant confers phosphorus acquisition diversity in soybean. *Nat Commun*. 2022b;**13**(1):3796. <https://doi.org/10.1038/s41467-022-31555-2>
- Gutierrez-Alanis D, Yong-Villalobos L, Jimenez-Sandoval P, Alatorre-Cobos F, Oropeza-Aburto A, Mora-Macias J, Sanchez-Rodriguez F, Cruz-Ramirez A, Herrera-Estrella L.** Phosphate starvation-dependent iron mobilization induces CLE14 expression to trigger root meristem differentiation through CLV2/PEPR2 signaling. *Dev Cell*. 2017;**41**(5):555–570.e3. <https://doi.org/10.1016/j.devcel.2017.05.009>
- Hamburger D, Rezzonico E, MacDonald-Comber Petétot J, Somerville C, Poirier Y.** Identification and characterization of the Arabidopsis PHO1 gene involved in phosphate loading to the xylem. *Plant Cell*. 2002;**14**(4):889–902. <https://doi.org/10.1105/tpc.000745>
- Härtel H, Dörmann P, Benning C.** Galactolipids not associated with the photosynthetic apparatus in phosphate-deprived plants. *J Photochem Photobiol B: Biol*. 2001;**61**(1-2):46–51. [https://doi.org/10.1016/S1011-1344\(01\)00144-0](https://doi.org/10.1016/S1011-1344(01)00144-0)
- Heuer S, Gaxiola R, Schilling R, Herrera-Estrella L, López-Arredondo D, Wissuwa M, Delhaize E, Rouached H.** Improving phosphorus use efficiency: a complex trait with emerging opportunities. *Plant J*. 2017;**90**(5):868–885. <https://doi.org/10.1111/tpj.13423>
- Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H.** Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J Exp Bot*. 2007;**59**(1):75–83. <https://doi.org/10.1093/jxb/erm157>
- Holford ICR.** Soil phosphorus: its measurement, and its uptake by plants. *Soil Res*. 1997;**35**(2):227–240. <https://doi.org/10.1071/S96047>
- Hsieh L-C, Lin S-I, Shih AC-C, Chen J-W, Lin W-Y, Tseng C-Y, Li W-H, Chiou T-J.** Uncovering small RNA-mediated responses to phosphate deficiency in Arabidopsis by deep sequencing. *Plant Physiol*. 2009;**151**(4):2120–2132. <https://doi.org/10.1104/pp.109.147280>
- Hu B, Jiang Z, Wang W, Qiu Y, Zhang Z, Liu Y, Li A, Gao X, Liu L, Qian Y, et al.** Nitrate-NRT1.1B-SPX4 cascade integrates nitrogen and phosphorus signalling networks in plants. *Nat Plants*. 2019;**5**(4):401–413. <https://doi.org/10.1038/s41477-019-0384-1>
- Huang TK, Han CL, Lin SI, Chen YJ, Tsai YC, Chen YR, Chen JW, Lin WY, Chen PM, Liu TY, et al.** Identification of downstream components of ubiquitin-conjugating enzyme PHOSPHATE2 by quantitative membrane proteomic in Arabidopsis roots. *Plant Cell*. 2013;**25**(10):4044–4060. <https://doi.org/10.1105/tpc.113.115998>
- Huen AK, Rodriguez-Medina C, Ho AYY, Atkins CA, Smith PMC.** Long-distance movement of phosphate starvation-responsive microRNAs in Arabidopsis. *Plant Biology*. 2017;**19**(4):643–649. <https://doi.org/10.1111/plb.12568>
- Hürlimann HC, Pinson B, Stadler-Waibel M, Zeeman SC, Freimoser FM.** The SPX domain of the yeast low-affinity phosphate transporter Pho90 regulates transport activity. *EMBO Rep*. 2009;**10**(9):1003–1008. <https://doi.org/10.1038/embor.2009.105>

- Isidra-Arellano M, Delaux P, Valdés-López O.** The phosphate starvation response system: its role in the regulation of plant–microbe interactions. *Plant Cell Physiol.* 2021;**62**(3):392–400. <https://doi.org/10.1093/pcp/pcab016>
- Jabnoue M, Secco D, Lecampion C, Robaglia C, Shu Q, Poirier Y.** A rice cis-natural antisense RNA acts as a translational enhancer for its cognate mRNA and contributes to phosphate homeostasis and plant fitness. *Plant Cell.* 2013;**25**(10):4166–4182. <https://doi.org/10.1105/tpc.113.116251>
- Jia H, Ren H, Gu M, Zhao J, Sun S, Zhang X, Chen J, Wu P, Xu G.** The phosphate transporter gene *OsPht1*; 8 is involved in phosphate homeostasis in rice. *Plant Physiol.* 2011;**156**(3):1164–1175. <https://doi.org/10.1104/pp.111.175240>
- Jia XC, Liu P, Lynch JP.** Greater lateral root branching density in maize improves phosphorus acquisition from low phosphorus soil. *J Exp Bot.* 2018;**69**(20):4961–4970. <https://doi.org/10.1093/jxb/ery252>
- Jiang M, Calderaru S, Zaehle S, Ellsworth DS, Medlyn BE.** Towards a more physiological representation of vegetation phosphorus processes in land surface models. *New Phytol.* 2019a;**222**(3):1223–1229. <https://doi.org/10.1111/nph.15688>
- Jiang M, Sun L, Isupov MN, Littlechild JA, Wu X, Wang Q, Wang Q, Yang W, Wu Y.** Structural basis for the target DNA recognition and binding by the MYB domain of phosphate starvation response 1. *FEBS J.* 2019b;**286**(14):2809–2821. <https://doi.org/10.1111/febs.14846>
- Karlsson PM, Herdean A, Adolfsen L, Beebo A, Nziengui H, Irigoyen S, Ünneper R, Zsiros O, Nagy G, Garab G, et al.** The Arabidopsis thylakoid transporter *PHT4*; 1 influences phosphate availability for ATP synthesis and plant growth. *Plant J.* 2015;**84**(1):99–110. <https://doi.org/10.1111/tpj.12962>
- Karthikeyan A, Varadarajan D, Mukatira U, Paino D'Urzo M, Damsz B, Raghothama K.** Regulated expression of Arabidopsis phosphate transporters. *Plant Physiol.* 2002;**130**(1):221–233. <https://doi.org/10.1104/pp.020007>
- Karthikeyan AS, Varadarajan DK, Jain A, Held MA, Carpita NC, Raghothama KG.** Phosphate starvation responses are mediated by sugar signaling in Arabidopsis. *Planta.* 2007;**225**(4):907–918. <https://doi.org/10.1007/s00425-006-0408-8>
- Kelly AA, Dörmann P.** DGD2, an arabidopsis gene encoding a UDP-galactose-dependent digalactosyldiacylglycerol synthase is expressed during growth under phosphate-limiting conditions. *J Biol Chem.* 2002;**266**(2):1166–1173. <https://doi.org/10.1074/jbc.M110066200>
- Kettenburg AT, Lopez MA, Yogendra K, Prior MJ, Rose T, Bimson S, Heuer S, Roy SJ, Bailey-Serres J.** PHOSPHORUS-STARVATION TOLERANCE 1 (*OsPSTOL1*) is prevalent in upland rice and enhances root growth and hastens low phosphate signaling in wheat. *Plant Cell Environ.* 2023;**46**(7):2187–2205. <https://doi.org/10.1111/pce.14588>
- Khan GA, Bouraine S, Wege S, Li YY, de Carbonnel M, Berthomieu P, Poirier Y, Rouached H.** Coordination between zinc and phosphate homeostasis involves the transcription factor *PHR1*, the phosphate exporter *PHO1*, and its homologue *PHO1*; H3 in *Arabidopsis*. *J Exp Bot.* 2014;**65**(3):871–884. <https://doi.org/10.1093/jxb/ert444>
- Kisko M, Bouain N, Safi A, Medici A, Akkers RC, Secco D, Fouret G, Krouk G, Aarts MGM, Busch W, et al.** *LPCAT1* controls phosphate homeostasis in a zinc-dependent manner. *Elife.* 2018;**7**:e32077. <https://doi.org/10.7554/eLife.32077>
- Kohlen W, Charnikhova T, Liu Q, Bours R, Domagalska MA, Beguerie S, Verstappen F, Leyser O, Bouwmeester H, Ruyter-Spira C.** Strigolactones are transported through the xylem and play a key role in shoot architectural response to phosphate deficiency in nonarbuscular mycorrhizal host Arabidopsis. *Plant Physiol.* 2011;**155**(2):974–987. <https://doi.org/10.1104/pp.110.164640>
- Kuo H, Hsu Y, Lin W, Chen K, Munnik T, Brearley C, Chiou T.** Arabidopsis inositol phosphate kinases *IPK1* and *ITPK1* constitute a metabolic pathway in maintaining phosphate homeostasis. *Plant J.* 2018;**95**(4):613–630. <https://doi.org/10.1111/tpj.13974>
- Laha D, Johnen P, Azevedo C, Dynowski M, Weiß M, Capolicchio S, Mao H, Iven T, Steenbergen M, Freyer M, et al.** *VIH2* regulates the synthesis of inositol pyrophosphate *InsP8* and jasmonate-dependent defenses in Arabidopsis. *Plant Cell.* 2015;**27**(4):1082–1097. <https://doi.org/10.1105/tpc.114.135160>
- Laha D, Parvin N, Hofer A, Giehl RFH, Fernandez-Rebollo N, von Wirén N, Saiardi A, Jessen HJ, Schaaf G.** Arabidopsis *ITPK1* and *ITPK2* have an evolutionarily conserved phytic acid kinase activity. *ACS Chem Biol.* 2019;**14**(10):2127–2133. <https://doi.org/10.1021/acscchembio.9b00423>
- Lapis-Gaza H, Jost R, Finnegan P.** Arabidopsis *PHOSPHATE TRANSPORTER1* genes *PHT1*; 8 and *PHT1*; 9 are involved in root-to-shoot translocation of orthophosphate. *BMC Plant Biol.* 2014;**14**(1):334. <https://doi.org/10.1186/s12870-014-0334-z>
- Lei M, Liu Y, Zhang B, Zhao Y, Wang X, Zhou Y, Raghothama KG, Liu D.** Genetic and genomic evidence that sucrose is a global regulator of plant responses to phosphate starvation in Arabidopsis. *Plant Physiol.* 2011;**156**(3):1116–1130. <https://doi.org/10.1104/pp.110.171736>
- Liao D, Sun C, Liang H, Wang Y, Bian X, Dong C, Niu X, Yang M, Xu G, Chen A, et al.** *SISPX1-SIPHR* complexes mediate the suppression of arbuscular mycorrhizal symbiosis by phosphate repletion in tomato. *Plant Cell.* 2022;**34**(10):4045–4065. <https://doi.org/10.1093/plcell/koac212>
- Lin SI, Chiang SF, Lin WY, Chen JW, Tseng CY, Wu PC, Chiou TJ.** Regulatory network of *MicroRNA399* and *PHO2* by systemic signaling. *Plant Physiol.* 2008;**147**(2):732–746. <https://doi.org/10.1104/pp.108.116269>
- Lin W, Lin Y, Chiang S, Syu C, Hsieh L, Chiou T.** Evolution of *microRNA827* targeting in the plant kingdom. *New Phytol.* 2018;**217**(4):1712–1725. <https://doi.org/10.1111/nph.14938>
- Lin WY, Huang TK, Chiou TJ.** *NITROGEN LIMITATION ADAPTATION*, a target of *microRNA827*, mediates degradation of plasma membrane-localized phosphate transporters to maintain phosphate homeostasis in Arabidopsis. *Plant Cell.* 2013;**25**(10):4061–4074. <https://doi.org/10.1105/tpc.113.116012>
- Lin WY, Huang TK, Leong SJ, Chiou TJ.** Long-distance call from phosphate: systemic regulation of phosphate starvation responses. *J Exp Bot.* 2014;**65**(7):1817–1827. <https://doi.org/10.1093/jxb/ert431>
- Liu C, Muchhal U, Raghothama K.** Differential expression of *TPS11*, a phosphate starvation induced gene in tomato. *Plant Mol Biol.* 1997;**33**(5):867–874. <https://doi.org/10.1023/a:1005729309569>
- Liu J, Samac DA, Bucciarelli B, Allan DL, Vance CP.** Signaling of phosphorus deficiency-induced gene expression in white lupin requires sugar and phloem transport. *Plant J.* 2005;**41**(2):257–268. <https://doi.org/10.1111/j.1365-313X.2004.02289.x>
- Liu J, Yang L, Luan M, Wang Y, Zhang C, Zhang B, Shi J, Zhao FG, Lan W, Luan S.** A vacuolar phosphate transporter essential for phosphate homeostasis in Arabidopsis. *Proc Natl Acad Sci U S A.* 2015;**112**(47):E6571–E6578. <https://doi.org/10.1073/pnas.1514598112>
- Liu JP, Magalhaes JV, Shaff J, Kochian LV.** Aluminum-activated citrate and malate transporters from the *MATE* and *ALMT* families function independently to confer Arabidopsis aluminum tolerance. *Plant J.* 2009;**57**(3):389–399. <https://doi.org/10.1111/j.1365-313X.2008.03696.x>
- Liu TY, Huang TK, Tseng CY, Lai YS, Lin SI, Lin WY, Chen JW, Chiou TJ.** *PHO2*-dependent degradation of *PHO1* modulates phosphate homeostasis in Arabidopsis. *Plant Cell.* 2012;**24**(5):2168–2183. <https://doi.org/10.1105/tpc.112.096636>
- Liu TY, Huang TK, Yang SY, Hong YT, Huang SM, Wang FN, Chiang SF, Tsai SY, Lu WC, Chiou TJ.** Identification of plant vacuolar transporters mediating phosphate storage. *Nat Commun.* 2016;**7**(1):11095. <https://doi.org/10.1038/ncomms11095>
- Liu XL, Wang L, Wang XW, Yan Y, Yang XL, Xie MY, Hu Z, Shen X, Ai H, Lin H, et al.** Mutation of the chloroplast-localized phosphate transporter *OsPHT2*; 1 reduces flavonoid accumulation and UV

- tolerance in rice. *Plant J.* 2020;**102**(1):53–67. <https://doi.org/10.1111/tpj.14611>
- Lopez-Arredondo DL, Leyva-Gonzalez MA, Gonzalez-Morales SI, Lopez-Bucio J, Herrera-Estrella L.** Phosphate nutrition: improving low-phosphate tolerance in crops. *Annu Rev Plant Biol.* 2014;**65**(1):95–123. <https://doi.org/10.1146/annurev-arplant-050213-035949>
- López-Bucio J, Cruz-Ramírez A, Herrera-Estrella L.** The role of nutrient availability in regulating root architecture. *Curr Opin Plant Biol.* 2003;**6**(3):280–287. [https://doi.org/10.1016/s1369-5266\(03\)00035-9](https://doi.org/10.1016/s1369-5266(03)00035-9)
- Lota F, Wegmuller S, Buer B, Sato S, Brautigam A, Hanf B, Bucher M.** The cis-acting CTTC-P1BS module is indicative for gene function of LjVTI12, a Qb-SNARE protein gene that is required for arbuscule formation in *Lotus japonicus*. *Plant J.* 2013;**74**(2):280–293. <https://doi.org/10.1111/tpj.12120>
- Lu M, Cheng Z, Zhang XM, Huang P, Fan C, Yu G, Chen F, Xu K, Chen Q, Miao Y, et al.** Spatial divergence of PHR-PHT1 modules maintains phosphorus homeostasis in soybean nodules. *Plant Physiol.* 2020;**184**(1):236–250. <https://doi.org/10.1104/pp.19.01209>
- Luan M, Zhao F, Sun G, Xu M, Fu A, Lan W, Luan S.** A SPX domain vacuolar transporter links phosphate sensing to homeostasis in *Arabidopsis*. *Mol Plant.* 2022;**15**(10):1590–1601. <https://doi.org/10.1016/j.molp.2022.09.005>
- Lv Q, Zhong Y, Wang Y, Wang Z, Zhang L, Shi J, Wu Z, Liu Y, Mao C, Yi K, et al.** SPX4 negatively regulates phosphate signaling and homeostasis through its interaction with PHR2 in rice. *Plant Cell.* 2014;**26**(4):1586–1597. <https://doi.org/10.1105/tpc.114.123208>
- Lynch JP.** Root phenes for enhanced soil exploration and phosphorus acquisition: tools for future crops. *Plant Physiol.* 2011;**156**(3):1041–1049. <https://doi.org/10.1104/pp.111.175414>
- Ma B, Zhang L, Gao Q, Wang J, Li X, Wang H, Liu Y, Lin H, Liu J, Wang X, et al.** A plasma membrane transporter coordinates phosphate reallocation and grain filling in cereals. *Nat Genet.* 2021;**53**(6):906–915. <https://doi.org/10.1038/s41588-021-00855-6>
- Martín AC, Del Pozo JC, Iglesias J, Rubio V, Solano R, De La Peña A, Leyva A, Paz-Ares J.** Influence of cytokinins on the expression of phosphate starvation responsive genes in *Arabidopsis*. *Plant J.* 2000;**24**(5):559–567. <https://doi.org/10.1046/j.1365-313x.2000.00893.x>
- Medici A, Szponarski W, Dangeville P, Safi A, Dissanayake IM, Saenchai C, Emanuel A, Rubio V, Lacombe B, Ruffel S, et al.** Identification of molecular integrators shows that nitrogen actively controls the phosphate starvation response in plants. *Plant Cell.* 2019;**31**(5):1171–1184. <https://doi.org/10.1105/tpc.18.00656>
- Milner MJ, Howells RM, Craze M, Bowden S, Graham N, Wallington EJ.** A PSTOL-like gene, controls a number of agronomically important traits in wheat. *BMC Plant Biol.* 2018;**18**(1):115. <https://doi.org/10.1186/s12870-018-1331-4>
- Miyaji T, Kuromori T, Takeuchi Y, Yamaji N, Yokosho K, Shimazawa A, Sugimoto E, Omote H, Ma J, Shinozaki K, et al.** AtPHT4; 4 is a chloroplast-localized ascorbate transporter in *Arabidopsis*. *Nat Commun.* 2015;**6**(1):11. <https://doi.org/10.1038/ncomms6928>
- Młodzińska E, Zboińska M.** Phosphate uptake and allocation—a closer look at *Arabidopsis thaliana* L. and *Oryza sativa* L. *Front Plant Sci.* 2016;**7**:1198. <https://doi.org/10.3389/fpls.2016.01198>
- Mora-Macias J, Ojeda-Rivera JO, Gutierrez-Alanis D, Yong-Villalobos L, Oropeza-Aburto A, Raya-Gonzalez J, Jimenez-Dominguez G, Chavez-Calvillo G, Rellán-Alvarez R, Herrera-Estrella L.** Malate-dependent Fe accumulation is a critical checkpoint in the root developmental response to low phosphate. *Proc Natl Acad Sci U S A.* 2017;**114**(17):E3563–E3572. <https://doi.org/10.1073/pnas.1701952114>
- Muchhal US, Pardo JM, Raghothama KG.** Phosphate transporters from the higher plant *Arabidopsis thaliana*. *Proc Natl Acad Sci U S A.* 1996;**93**(19):10519–10523. <https://doi.org/10.1073/pnas.93.19.10519>
- Mudge S, Rae A, Diatloff E, Smith F.** Expression analysis suggests novel roles for members of the Pht1 family of phosphate transporters in *Arabidopsis*. *Plant J.* 2002;**31**(3):341–353. <https://doi.org/10.1046/j.1365-313x.2002.01356.x>
- Müller J, Toev T, Heisters M, Teller J, Moore KL, Hause G, Dinesh DC, Bürstenbinder K, Abel S.** Iron-dependent callose deposition adjusts root meristem maintenance to phosphate availability. *Dev Cell.* 2015;**33**(2):216–230. <https://doi.org/10.1016/j.devcel.2015.02.007>
- Nagy R, Grob H, Weder B, Green P, Klein M, Frelet-Barrand A, Schjoerring J, Brearley C, Martinoia E.** The *Arabidopsis* ATP-binding cassette protein AtMRP5/AtABCC5 is a high affinity inositol hexakisphosphate transporter involved in guard cell signaling and phytate storage. *J Biol Chem.* 2009;**284**(48):33614–33622. <https://doi.org/10.1074/jbc.M109.030247>
- Nakamura Y.** Phosphate starvation and membrane lipid remodeling in seed plants. *Prog Lipid Res.* 2013;**52**(1):43–50. <https://doi.org/10.1016/j.plipres.2012.07.002>
- Naumann C, Heisters M, Brandt W, Janitzka P, Alfs C, Tang N, Nienguesso AT, Ziegler J, Imre R, Mechtler K, et al.** Bacterial-type ferroxidase tunes iron-dependent phosphate sensing during *Arabidopsis* root development. *Curr Biol.* 2022;**32**(10):2189–2205. <https://doi.org/10.1016/j.cub.2022.04.005>
- Naumann C, Mueller J, Sakhonwasee S, Wiegand A, Hause G, Heisters M, Buerstenbinder K, Abel S.** The local phosphate deficiency response activates endoplasmic reticulum stress-dependent autophagy. *Plant Physiol.* 2019;**179**(2):460–476. <https://doi.org/10.1104/pp.18.01379>
- Neelam K, Thakur S, Neha, Yadav IS, Kumar K, Dhaliwal SS, Singh K.** Novel alleles of phosphorus-starvation tolerance 1 gene (*PSTOL1*) from *Oryza rufipogon* confers high phosphorus uptake efficiency. *Front Plant Sci.* 2017;**8**:509. <https://doi.org/10.3389/fpls.2017.00509>
- Nguyen N, Clua J, Vetat P, Vuarambon D, De Bellis D, Pervert M, Lepetit M, Udvardi M, Valentine A, Poirier Y.** PHO1 family members transport phosphate from infected nodule cells to bacteroids in *Medicago truncatula*. *Plant Physiol.* 2021;**185**(1):196–209. <https://doi.org/10.1093/plphys/kiaa016>
- Nielsen TH, Krapp A, Röper-Schwarz U, Stitt M.** The sugar-mediated regulation of genes encoding the small subunit of Rubisco and the regulatory subunit of ADP glucose pyrophosphorylase is modified by phosphate and nitrogen. *Plant Cell Environ.* 1998;**21**(5):443–454. <https://doi.org/10.1046/j.1365-3040.1998.00295.x>
- Nilsson L, Müller R, Nielsen TH.** Increased expression of the MYB-related transcription factor, PHR1, leads to enhanced phosphate uptake in *Arabidopsis thaliana*. *Plant Cell Environ.* 2007;**30**(12):1499–1512. <https://doi.org/10.1111/j.1365-3040.2007.01734.x>
- Nussaume L, Kanno S, Javot H, Marin E, Pochon N, Ayadi A, Nakanishi T, Thibaud M.** Phosphate import in plants: focus on the PHT1 transporters. *Front Plant Sci.* 2011;**2**:83. <https://doi.org/10.3389/fpls.2011.00083>
- Okazaki Y, Otsuki H, Narisawa T, Kobayashi M, Sawai S, Kamide Y, Kusano M, Aoki T, Hirai MY, Saito K.** A new class of plant lipid is essential for protection against phosphorus depletion. *Nat Commun.* 2013;**4**(1):1510. <https://doi.org/10.1038/ncomms2512>
- Oldroyd GED, Leyser O.** A plant's diet, surviving in a variable nutrient environment. *Science.* 2020;**368**(6486):eaba0196. <https://doi.org/10.1126/science.aba0196>
- Ouyang X, Hong X, Zhao X, Zhang W, He X, Ma W, Teng W, Tong Y.** Knock out of the PHOSPHATE 2 gene TaPHO2-A1 improves phosphorus uptake and grain yield under low phosphorus conditions in common wheat. *Sci Rep.* 2016;**6**(1):29850. <https://doi.org/10.1038/srep29850>
- Pang JY, Bansal R, Zhao HX, Bohuon E, Lambers H, Ryan MH, Ranathunge K, Siddique KHM.** The carboxylate-releasing phosphorus-mobilizing strategy can be proxied by foliar manganese concentration in a large set of chickpea germplasm under low phosphorus supply. *New Phytol.* 2018;**219**(2):518–529. <https://doi.org/10.1111/nph.15200>
- Pant BD, Buhtz A, Kehr J, Scheible W-R.** MicroRNA399 is a long-distance signal for the regulation of plant phosphate homeostasis. *Plant J.* 2008;**53**(5):731–738. <https://doi.org/10.1111/j.1365-313x.2007.03363.x>

- Pant BD, Burgos A, Pant P, Cuadros-Inostroza A, Willmitzer L, Scheible W-R.** The transcription factor PHR1 regulates lipid remodeling and triacylglycerol accumulation in *Arabidopsis thaliana* during phosphorus starvation. *J Exp Bot.* 2015;**66**(7):1907–1918. <https://doi.org/10.1093/jxb/eru535>
- Pant BD, Musialak-Lange M, Nuc P, May P, Buhtz A, Kehr J, Walther D, Scheible W-Rd.** Identification of nutrient-responsive Arabidopsis and rapeseed MicroRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. *Plant Physiol.* 2009;**150**(3):1541–1555. <https://doi.org/10.1104/pp.109.139139>
- Paries M, Gutjahr C.** The good, the bad, and the phosphate: regulation of beneficial and detrimental plant-microbe interactions by the plant phosphate status. *New Phytol.* 2023;**239**(1):29–46. <https://doi.org/10.1111/nph.18933>
- Park BS, Seo JS, Chua NH.** NITROGEN LIMITATION ADAPTATION recruits PHOSPHATE2 to target the phosphate transporter PT2 for degradation during the regulation of Arabidopsis phosphate homeostasis. *Plant Cell.* 2014;**26**(1):454–464. <https://doi.org/10.1105/tpc.113.120311>
- Plaxton WC, Tran HT.** Metabolic adaptations of phosphate-starved plants. *Plant Physiol.* 2011;**156**(3):1006–1015. <https://doi.org/10.1104/pp.111.175281>
- Poirier Y, Thoma S, Somerville C, Schiefelbein J.** Mutant of Arabidopsis deficient in xylem loading of phosphate. *Plant Physiol.* 1991;**97**(3):1087–1093. <https://doi.org/10.1104/pp.97.3.1087>
- Pontigo S, Parra-Almuna L, Luengo-Escobar A, Poblete-Grant P, Nunes-Nesi A, Mora M, Cartes P.** Biochemical and molecular responses underlying the contrasting phosphorus use efficiency in Ryegrass cultivars. *Plants.* 2023;**12**(6):1224. <https://doi.org/10.3390/plants12061224>
- Potapenko E, Cordeiro CD, Huang G, Storey M, Wittwer C, Dutta AK, Jessen HJ, Starai VJ, Docampo R.** 5-Diphosphoinositol pentakisphosphate (5-IP7) regulates phosphate release from acidocalcisomes and yeast vacuoles. *J Biol Chem.* 2018;**293**(49):19101–19112. <https://doi.org/10.1074/jbc.ra118.005884>
- Prathap V, Kumar A, Maheshwari C, Tyagi A.** Phosphorus homeostasis: acquisition, sensing, and long-distance signaling in plants. *Mol Biol Rep.* 2022;**49**(8):8071–8086. <https://doi.org/10.1007/s11033-022-07354-9>
- Puga MI, Mateos I, Charukesi R, Wang Z, Franco-Zorrilla JM, Lorenzo L, Irigoyen ML, Masiero S, Bustos R, Rodríguez J, et al.** SPX1 is a phosphate-dependent inhibitor of PHOSPHATE STARVATION RESPONSE 1 in Arabidopsis. *Proc Natl Acad Sci U S A.* 2014;**111**(41):14947–14952. <https://doi.org/10.1073/pnas.1404654111>
- Qin L, Zhao J, Tian J, Chen LY, Sun ZA, Guo YX, Lu X, Gu MA, Xu GH, Liao H.** The high-affinity phosphate transporter GmPT5 regulates phosphate transport to nodules and nodulation in soybean. *Plant Physiol.* 2012;**159**(4):1634–1643. <https://doi.org/10.1104/pp.112.199786>
- Raghothama KG.** Phosphate acquisition. *Annu Rev Plant Physiol Plant Mol Biol.* 1999;**50**(1):665–693. <https://doi.org/10.1146/annurev.arplant.50.1.665>
- Reis R, Deforges J, Schmidt R, Schippers J, Poirier Y.** An antisense noncoding RNA enhances translation via localised structural rearrangements of its cognate mRNA. *Plant Cell.* 2021;**33**(4):1381–1397. <https://doi.org/10.1093/plcell/koab010>
- Reis R, Deforges J, Sokoloff T, Poirier Y.** Modulation of shoot phosphate level and growth by PHOSPHATE1 upstream open reading frame. *Plant Physiol.* 2020;**183**(3):1145–1156. <https://doi.org/10.1104/pp.19.01549>
- Remy E, Cabrito TR, Batista RA, Teixeira MC, Sa-Correia I, Duque P.** The Pht1; 9 and Pht1; 8 transporters mediate inorganic phosphate acquisition by the *Arabidopsis thaliana* root during phosphorus starvation. *New Phytol.* 2012;**195**(2):356–371. <https://doi.org/10.1111/j.1469-8137.2012.04167.x>
- Reymond M, Svistoonoff S, Loudet O, Nussaume L, Desnos T.** Identification of QTL controlling root growth response to phosphate starvation in *Arabidopsis thaliana*. *Plant Cell Environ.* 2006;**29**(1):115–125. <https://doi.org/10.1111/j.1365-3040.2005.01405.x>
- Ried MK, Wild R, Zhu JS, Pipercevic J, Sturm K, Broger L, Harmel RK, Abriata LA, Hothorn LA, Fiedler D, et al.** Inositol pyrophosphates promote the interaction of SPX domains with the coiled-coil motif of PHR transcription factors to regulate plant phosphate homeostasis. *Nat Commun.* 2021;**12**(1):384. <https://doi.org/10.1038/s41467-020-20681-4>
- Riemer E, Qiu D, Laha D, Harmel RK, Gaugler P, Gaugler V, Frei M, Hajirezaei M-R, Laha NP, Krusenbaum L, et al.** ITPK1 is an InsP6/ADP phosphotransferase that controls phosphate signaling in *Arabidopsis*. *Mol Plant.* 2021;**14**(11):1864–1880. <https://doi.org/10.1016/j.molp.2021.07.011>
- Robles-Aguilar AA, Pang JY, Postma JA, Schrey SD, Lambers H, Jablonowski ND.** The effect of pH on morphological and physiological root traits of *Lupinus angustifolius* treated with struvite as a recycled phosphorus source. *Plant Soil.* 2019;**434**(1-2):65–78. <https://doi.org/10.1007/s11104-018-3787-2>
- Ruan W, Guo M, Xu L, Wang X, Zhao H, Wang J, Yi K.** An SPX-RLI1 module regulates leaf inclination in response to phosphate availability in rice. *Plant Cell.* 2018;**30**(4):853–870. <https://doi.org/10.1105/tpc.17.00738>
- Rubio Y, Linhares F, Solano R, Martín AC, Iglesias J, Leyva A, Paz-Ares J.** A conserved MYB transcription factor involved in phosphate starvation signaling both in vascular plants and in unicellular algae. *Genes Dev.* 2001;**15**(16):2122–2133. <https://doi.org/10.1101/gad.204401>
- Ryan PR, Delhaize E, Jones DL.** Function and mechanism of organic anion exudation from plant roots. *Annu Rev Plant Physiol Plant Mol Biol.* 2001;**52**(1):527–560. <https://doi.org/10.1146/annurev.arplant.52.1.527>
- Sahu A, Banerjee S, Raju A, Chiou T-J, Garcia L, Versaw W.** Spatial profiles of phosphate in roots indicate developmental control of uptake, recycling, and sequestration. *Plant Physiol.* 2020;**184**(4):2064–2077. <https://doi.org/10.1104/pp.20.01008>
- Sakakibara H.** Cytokinins: activity, biosynthesis, and translocation. *Annu Rev Plant Biol.* 2006;**57**(1):431–449. <https://doi.org/10.1146/annurev.arplant.57.032905.105231>
- Sanchez-Calderon L, Lopez-Bucio J, Chacon-Lopez A, Cruz-Ramirez A, Nieto-Jacobo F, Dubrovsky JG, Herrera-Estrella L.** Phosphate starvation induces a determinate developmental program in the roots of *Arabidopsis thaliana*. *Plant Cell Physiol.* 2005;**46**(1):174–184. <https://doi.org/10.1093/pcp/pci011>
- Sawaki Y, Iuchi S, Kobayashi Y, Kobayashi Y, Ikka T, Sakurai N, Fujita M, Shinozaki K, Shibata D, Kobayashi M, et al.** STOP1 regulates multiple genes that protect Arabidopsis from proton and aluminum toxicities. *Plant Physiol.* 2009;**150**(1):281–294. <https://doi.org/10.1104/pp.108.134700>
- Secco D, Baumann A, Poirier Y.** Characterization of the rice PHO1 gene family reveals a key role for OsPHO1; 2 in phosphate homeostasis and the evolution of a distinct clade in dicotyledons. *Plant Physiol.* 2010;**152**(3):1693–1704. <https://doi.org/10.1104/pp.109.149872>
- Secco D, Wang C, Arpat BA, Wang Z, Poirier Y, Tyerman SD, Wu P, Shou H, Whelan J.** The emerging importance of the SPX domain-containing proteins in phosphate homeostasis. *New Phytol.* 2012;**193**(4):842–851. <https://doi.org/10.1111/j.1469-8137.2011.04002.x>
- Shi J, Zhao B, Zheng S, Zhang X, Wang X, Dong W, Xie Q, Wang G, Xiao Y, Chen F, et al.** A phosphate starvation response-centered network regulates mycorrhizal symbiosis. *Cell.* 2021;**184**(22):5527–5540. <https://doi.org/10.1016/j.cell.2021.09.030>
- Shin H, Shin HS, Chen R, Harrison MJ.** Loss of At4 function impacts phosphate distribution between the roots and the shoots during phosphate starvation. *Plant J.* 2006;**45**(5):712–726. <https://doi.org/10.1111/j.1365-313X.2005.02629.x>
- Shin H, Shin HS, Dewbre GR, Harrison MJ.** Phosphate transport in Arabidopsis: Pht1; 1 and Pht1; 4 play a major role in phosphate acquisition from both low- and high-phosphate environments. *Plant J.*

- 2004;**39**(4):629–642. <https://doi.org/10.1111/j.1365-313X.2004.02161.x>
- Silva-Navas J, Conesa CM, Saez A, Navarro-Neila S, Garcia-Mina JM, Zamarreño AM, Baigorri R, Swarup R, Del Pozo JC.** Role of cis-zeatin in root responses to phosphate starvation. *New Phytol.* 2019;**224**(1):242–257. <https://doi.org/10.1111/nph.16020>
- Song HN, Yin ZT, Chao MN, Ning LH, Zhang D, Yu DY.** Functional properties and expression quantitative trait loci for phosphate transporter GmPT1 in soybean. *Plant Cell Environ.* 2014;**37**(2):462–472. <https://doi.org/10.1111/pce.12170>
- Stegmann M, Monaghan J, Smakowska-Luzan E, Rovenich H, Lehner A, Holton N, Belkhadir Y, Zipfel C.** The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. *Science.* 2017;**355**(6322):287–289. <https://doi.org/10.1126/science.aal2541>
- Sun BR, Gao YZ, Lynch JP.** Large crown root number improves topsoil foraging and phosphorus acquisition. *Plant Physiol.* 2018;**177**(1):90–104. <https://doi.org/10.1104/pp.18.00234>
- Svistonoff S, Creff A, Reymond M, Sigoillot-Claude C, Ricaud L, Blanchet A, Nussaume L, Desnos T.** Root tip contact with low-phosphate media reprograms plant root architecture. *Nat Genet.* 2007;**39**(6):792–796. <https://doi.org/10.1038/ng2041>
- Tang C, Hinsinger P, Drevon JJ, Jaillard B.** Phosphorus deficiency impairs early nodule functioning and enhances proton release in roots of *Medicago truncatula* L. *Ann Bot.* 2001;**88**(1):131–138. <https://doi.org/10.1006/anbo.2001.1440>
- Tang J, Wu D, Li X, Wang L, Xu L, Zhang Y, Xu F, Liu H, Xie Q, Dai S, et al.** Plant immunity suppression via PHR1-RALF-FERONIA shapes the root microbiome to alleviate phosphate starvation. *EMBO J.* 2022;**41**(6):e109102. <https://doi.org/10.15252/embj.2021109102>
- Taylor CB, Bariola PA, delCardayre SB, Raines RT, Green PJ.** RNS2: a senescence-associated RNase of Arabidopsis that diverged from the S-RNases before speciation. *Proc Natl Acad Sci U S A.* 1993;**90**(11):5118–5122. <https://doi.org/10.1073/pnas.90.11.5118>
- Thibaud MC, Arrighi JF, Bayle V, Chiarenza S, Creff A, Bustos R, Paz-Ares J, Poirier Y, Nussaume L.** Dissection of local and systemic transcriptional responses to phosphate starvation in Arabidopsis. *Plant J.* 2010;**64**(5):775–789. <https://doi.org/10.1111/j.1365-313X.2010.04375.x>
- Ticconi CA, Delatorre CA, Abel S.** Attenuation of phosphate starvation responses by phosphite in Arabidopsis. *Plant Physiol.* 2001;**127**(3):963–972. <https://doi.org/10.1104/pp.010396>
- Ticconi CA, Lucero RD, Sakhonwasee S, Adamson AW, Creff A, Nussaume L, Desnos T, Abel S.** ER-resident proteins PDR2 and LPR1 mediate the developmental response of root meristems to phosphate availability. *Proc Natl Acad Sci U S A.* 2009;**106**(33):14174–14179. <https://doi.org/10.1073/pnas.0901778106>
- Umehara M, Hanada A, Yoshida S, Akiyama K, Arite T, Takeda-Kamiya N, Magome H, Kamiya Y, Shirasu K, Yoneyama K, et al.** Inhibition of shoot branching by new terpenoid plant hormones. *Nature.* 2008;**455**(7210):195–200. <https://doi.org/10.1038/nature07272>
- Valverde C, Ferrari A, Wall LG.** Phosphorus and the regulation of nodulation in the actinorhizal symbiosis, between *Discaria trinervis* (Rhamnaceae) and *Frankia* BC110501. *New Phytol.* 2002;**153**(1):43–51. <https://doi.org/10.1046/j.0028-646X.2001.00298.x>
- van de Wiel CCM, van der Linden CG, Scholten OE.** Improving phosphorus use efficiency in agriculture: opportunities for breeding. *Euphytica.* 2016;**207**(1):1–22. <https://doi.org/10.1007/s10681-015-1572-3>
- Vance CP, Uhde-Stone C, Allan DL.** Phosphorus acquisition and use: critical adaptations by plants for securing a nonrenewable resource. *New Phytol.* 2003;**157**(3):423–447. <https://doi.org/10.1046/j.1469-8137.2003.00695.x>
- Varadarajan DK, Karthikeyan AS, Matilda PD, Raghothama KG.** Phosphite, an analog of phosphate, suppresses the coordinated expression of genes under phosphate starvation. *Plant Physiol.* 2002;**129**(3):1232–1240. <https://doi.org/10.1104/pp.010835>
- Versaw WK, Garcia LR.** Intracellular transport and compartmentation of phosphate in plants. *Curr Opin Plant Biol.* 2017;**39**:25–30. <https://doi.org/10.1016/j.pbi.2017.04.015>
- Versaw WK, Harrison MJ.** A chloroplast phosphate transporter, PHT2; 1, influences allocation of phosphate within the plant and phosphate-starvation responses. *Plant Cell.* 2002;**14**(8):1751–1766. <https://doi.org/10.1105/tpc.002220>
- Vetal P, Poirier Y.** The Arabidopsis PHOSPHATE 1 exporter undergoes constitutive internalization via clathrin-mediated endocytosis. *Plant J.* 2023;**116**(5):1477–1491. <https://doi.org/10.1111/tpj.16441>
- Vogiatzaki E, Baroux C, Jung J-Y, Poirier Y.** PHO1 exports phosphate from the chalazal seed coat to the embryo in developing Arabidopsis seeds. *Curr Biol.* 2017;**27**(19):2893–2900. <https://doi.org/10.1016/j.cub.2017.08.026>
- Wang C, Huang W, Ying Y, Li S, Secco D, Tyerman S, Whelan J, Shou H.** Functional characterization of the rice SPX-MFS family reveals a key role of OsSPX-MFS1 in controlling phosphate homeostasis in leaves. *New Phytol.* 2012;**196**(1):139–148. <https://doi.org/10.1111/j.1469-8137.2012.04227.x>
- Wang C, Yue W, Ying Y, Wang S, Secco D, Liu Y, Whelan J, Tyerman SD, Shou H.** Rice SPX-major facility superfamily3, a vacuolar phosphate efflux transporter, is involved in maintaining phosphate homeostasis in rice. *Plant Physiol.* 2015;**169**(4):2822–2831. <https://doi.org/10.1104/pp.15.01005>
- Wang F, Deng M, Chen J, He Q, Jia X, Guo H, Xu J, Liu Y, Zhang S, Shou H, et al.** CASEIN KINASE2-dependent phosphorylation of PHOSPHATE2 fine-tunes phosphate homeostasis in rice. *Plant Physiol.* 2020;**183**(1):250–262. <https://doi.org/10.1104/pp.20.00078>
- Wang P, Snijders R, Kohlen W, Liu J, Bisseling T, Limpens E.** Medicago SPX1 and SPX3 regulate phosphate homeostasis, mycorrhizal colonization, and arbuscule degradation. *Plant Cell.* 2021;**33**(11):3470–3486. <https://doi.org/10.1093/plcell/koab206>
- Wang W, Ding GD, White PJ, Wang XH, Jin KM, Xu FS, Shi L.** Mapping and cloning of quantitative trait loci for phosphorus efficiency in crops: opportunities and challenges. *Plant Soil.* 2019;**439**(1-2):91–112. <https://doi.org/10.1007/s11104-018-3706-6>
- Wang X, Wang Y, Piñeros M, Wang Z, Wang W, Li C, Wu Z, Kochian L, Wu P.** Phosphate transporters OsPHT1; 9 and OsPHT1; 10 are involved in phosphate uptake in rice. *Plant Cell Environ.* 2014a;**37**:1159–1170. <https://doi.org/10.1111/pce.12224>
- Wang X, Yi K, Tao Y, Wang F, Wu Z, Jiang D, Chen X, Zhu L, Wu P.** Cytokinin represses phosphate-starvation response through increasing of intracellular phosphate level. *Plant Cell Environ.* 2006;**29**(10):1924–1935. <https://doi.org/10.1111/j.1365-3040.2006.01568.x>
- Wang Z, Ruan W, Shi J, Zhang L, Xiang D, Yang C, Li C, Wu Z, Liu Y, Yu Y, et al.** Rice SPX1 and SPX2 inhibit phosphate starvation responses through interacting with PHR2 in a phosphate-dependent manner. *Proc Natl Acad Sci U S A.* 2014b;**111**(41):14953–14958. <https://doi.org/10.1073/pnas.1404680111>
- Wang Z, Zheng Z, Song L, Liu D.** Functional characterization of Arabidopsis PHL4 in plant response to phosphate starvation. *Front Plant Sci.* 2018;**9**:1432. <https://doi.org/10.3389/fpls.2018.01432>
- Wang Z, Zheng Z, Zhu Y, Kong S, Liu D.** PHOSPHATE RESPONSE 1 family members act distinctly to regulate transcriptional responses to phosphate starvation. *Plant Physiol.* 2022;**191**(2):1324–1343. <https://doi.org/10.1093/plphys/kiac521>
- Wege S, Khan G, Jung J, Vogiatzaki E, Pradervand S, Aller I, Meyer A, Poirier Y.** The EXS domain of PHO1 participates in the response of shoots to phosphate deficiency via a root-to-shoot signal. *Plant Physiol.* 2016;**170**(1):385–400. <https://doi.org/10.1104/pp.15.00975>
- Wen ZH, Li HB, Shen Q, Tang XM, Xiong CY, Li HG, Pang JY, Ryan MH, Lambers H, Shen JB.** Tradeoffs among root morphology, exudation and mycorrhizal symbioses for phosphorus-acquisition strategies of 16 crop species. *New Phytol.* 2019;**223**(2):882–895. <https://doi.org/10.1111/nph.15833>

- Whitfield H, White G, Sprigg C, Riley AM, Potter BVL, Hemmings AM, Brearley CA.** An ATP-responsive metabolic cassette comprised of inositol tris/tetrakisphosphate kinase 1 (ITPK1) and inositol pentakisphosphate 2-kinase (IPK1) buffers diphosphoinositol phosphate levels. *Biochem J.* 2020;**477**(14):2621–2638. <https://doi.org/10.1042/BCJ20200423>
- Wild R, Gerasimaite R, Jung J-Y, Truffault V, Pavlovic I, Schmidt A, Saiardi A, Jessen HJ, Poirier Y, Hothorn M, et al.** Control of eukaryotic phosphate homeostasis by inositol polyphosphate sensor domains. *Science.* 2016;**352**(6288):986–990. <https://doi.org/10.1126/science.aad9858>
- Wissuwa M, Wegner J, Ae N, Yano M.** Substitution mapping of Pup1: a major QTL increasing phosphorus uptake of rice from a phosphorus-deficient soil. *Theor Appl Genet.* 2002;**105**(6):890–897. <https://doi.org/10.1007/s00122-002-1051-9>
- Wu Y, Shi L, Li L, Fu LW, Liu YL, Xiong Y, Sheen J.** Integration of nutrient, energy, light, and hormone signalling via TOR in plants. *J Exp Bot.* 2019;**70**(8):2227–2238. <https://doi.org/10.1093/jxb/erz028>
- Xiao X, Zhang J, Satheesh V, Meng F, Gao W, Dong J, Zheng Z, An G-Y, Nussaume L, Liu D, et al.** SHORT-ROOT stabilizes PHOSPHATE1 to regulate phosphate allocation in Arabidopsis. *Nat Plants.* 2022;**8**(9):1074–1081. <https://doi.org/10.1038/s41477-022-01231-w>
- Xing X, Du H, Yang Z, Li X, Kong Y, Li W, Zhang C.** GmSPX8, a nodule-localized regulator confers nodule development and nitrogen fixation under phosphorus starvation in soybean. *BMC Plant Biol.* 2022;**22**(1):161. <https://doi.org/10.1186/s12870-022-03556-2>
- Xu L, Zhao H, Wan R, Liu Y, Xu Z, Tian W, Ruan W, Wang F, Deng M, Wang J, et al.** Identification of vacuolar phosphate efflux transporters in land plants. *Nat Plants.* 2019;**5**(1):84–94. <https://doi.org/10.1038/s41477-018-0334-3>
- Yahya M, ul Islam E, Rasul M, Farooq I, Mahreen N, Tawab A, Irfan M, Rajput L, Amin I, Yasmin S.** Differential root exudation and architecture for improved growth of wheat mediated by phosphate solubilizing bacteria. *Front Microbiol.* 2021;**12**:744094. <https://doi.org/10.3389/fmicb.2021.744094>
- Yamaji N, Takemoto Y, Miyaji T, Mitani-Ueno N, Yoshida KT, Ma JF.** Reducing phosphorus accumulation in rice grains with an impaired transporter in the node. *Nature.* 2017;**541**(7635):92–95. <https://doi.org/10.1038/nature20610>
- Yang SY, Huang TK, Kuo HF, Chiou TJ.** Role of vacuoles in phosphorus storage and remobilization. *J Exp Bot.* 2017;**68**(12):3045–3055. <https://doi.org/10.1093/jxb/erw481>
- Yang SY, Lu WC, Ko SS, Sun CM, Hung JC, Chiou TJ.** Upstream open reading frame and phosphate-regulated expression of rice OsNLA1 controls phosphate transport and reproduction. *Plant Physiol.* 2020;**182**(1):393–407. <https://doi.org/10.1104/pp.19.01101>
- Yue W, Ying Y, Wang C, Zhao Y, Dong C, Whelan J, Shou H.** OsNLA1, a RING-type ubiquitin ligase, maintains phosphate homeostasis in *Oryza sativa* via degradation of phosphate transporters. *Plant J.* 2017;**90**(6):1040–1051. <https://doi.org/10.1111/tpj.13516>
- Zhang F, Sun Y, Pei W, Jain A, Sun R, Cao Y, Wu X, Jiang T, Zhang L, Fan X, et al.** Involvement of OsPht1; 4 in phosphate acquisition and mobilization facilitates embryo development in rice. *Plant J.* 2015;**82**(4):556–569. <https://doi.org/10.1111/tpj.12804>
- Zhong Y, Wang Y, Guo J, Zhu X, Shi J, He Q, Liu Y, Wu Y, Zhang L, Lv Q, et al.** Rice SPX6 negatively regulates the phosphate starvation response through suppression of the transcription factor PHR2. *New Phytol.* 2018;**219**(1):135–148. <https://doi.org/10.1111/nph.15155>
- Zhou J, Hu Q, Xiao X, Yao D, Ge S, Ye J, Li H, Cai R, Liu R, Meng F.** Mechanism of phosphate sensing and signaling revealed by rice SPX1-PHR2 complex structure. *Nat Commun.* 2021;**12**(1):7040. <https://doi.org/10.1038/s41467-021-27391-5>
- Zhou J, Jiao F, Wu Z, Li Y, Wang X, He X, Zhong W, Wu P.** OsPHR2 is involved in phosphate-starvation signaling and excessive phosphate accumulation in shoots of plants. *Plant Physiol.* 2008;**146**(4):1673–1686. <https://doi.org/10.1104/pp.107.111443>
- Zhu J, Lau K, Puschmann R, Harmel RK, Zhang Y, Pries V, Gaugler P, Broger L, Dutta AK, Jessen HJ, et al.** Two bifunctional inositol pyrophosphate kinases/phosphatases control plant phosphate homeostasis. *Elife.* 2019;**8**:e43582. <https://doi.org/10.7554/eLife.43582>
- Zhu W, Miao Q, Sun D, Yang G, Wu C, Huang J, Zheng C.** The mitochondrial phosphate transporters modulate plant responses to salt stress via affecting ATP and gibberellin metabolism in *Arabidopsis thaliana*. *PLoS One.* 2012;**7**(8):e43530. <https://doi.org/10.1371/journal.pone.0043530>
- Zhuang Q, Xue Y, Yao Z, Zhu S, Liang C, Liao H, Tian J.** Phosphate starvation responsive GmSPX5 mediates nodule growth through interaction with GmNF-YC4 in soybean (*Glycine max*). *Plant J.* 2021;**108**(5):1422–1438. <https://doi.org/10.1111/tpj.15520>
- Zimmerli C, Ribot C, Vavasseur A, Bauer H, Hedrich R, Poirier Y.** PHO1 expression in guard cells mediates the stomatal response to abscisic acid in Arabidopsis. *Plant J.* 2012;**72**(2):199–211. <https://doi.org/10.1111/j.1365-313X.2012.05058.x>