

yellow alfalfa

Medicago sativa ssp. *falcata* (L.) Arcang.

Synonyms: *Medicago falcata* L., *Medicago falcata* L. var. *romanica* (Prodan) O. Schwarz & Klink.

Other common names: alfalfa

Family: Fabaceae

Invasiveness Rank: 64 The invasiveness rank is calculated based on a species' ecological impacts, biological attributes, distribution, and response to control measures. The ranks are scaled from 0 to 100, with 0 representing a plant that poses no threat to native ecosystems and 100 representing a plant that poses a major threat to native ecosystems.

Description

Yellow alfalfa is a perennial herb that grows up to 91 cm tall. Multiple, erect stems grow from the thick, somewhat woody crown. Leaves are alternate and composed of 3 leaflets. Leaflets are ovate, hairy, and 13 to 25 ½ mm long. Central leaflets are stalked, while lateral leaflets are sessile. Flowers are yellow and 13 mm long. They grow in oblong clusters in leaf axils. Pods are curved (falcate) or nearly straight. Each pod is 6 to 13 mm long and encloses several seeds (Hultén 1968, Hitchcock and Cronquist 1973, Royer and Dickinson 1999).



Medicago sativa ssp. *falcata* (L.) Arcang.

Similar species: Yellow alfalfa readily hybridizes with common alfalfa, *M. sativa* ssp. *sativa*, to produce intermediate forms (Small and Brookes 1984). Common alfalfa can be distinguished from yellow alfalfa because it has purple flowers and spirally-coiled pods. Both alfalfa and yellow alfalfa can be easily confused with other weedy, trifoliate legumes, such as *Melilotus* and *Trifolium* species. Alfalfa can be distinguished from other trifoliate legumes, even when not in flower or fruit, by the presence of longer stalks on the central leaflets than on the lateral leaflets and toothed margins on the leaves.



Leaves of *Medicago sativa* ssp. *falcata* (L.) Arcang.

Ecological Impact

Impact on community composition, structure, and interactions: 27 bird species and 46 mammal species are known to consume yellow alfalfa (Graham 1941). Most big game animals, including moose and mule deer, eat yellow alfalfa (Leach 1956, Kufeld 1973). Many small mammals, including marmots, mice, and ground squirrels, graze on yellow alfalfa. Waterfowl, such as American wigeons and mallards, eat the leaves, flowers, and seeds. Seeds are also consumed by rodents, rabbits, and upland birds. Yellow alfalfa is a source of nectar and pollen for insects (Graham 1941, Stanton 1974), and it is particularly attractive to solitary bees (Carlson pers. obs.). Dabbling ducks (mallards, blue-winged teals, northern pintail, northern shovelers, and American wigeons) nest in alfalfa stands (Klett et al. 1984). Undisturbed alfalfa fields provide food and cover for a variety of birds, including sharp-tailed grouse, American bitterns, marsh hawks, short-eared owls, and passerines (Duebbert et al. 1981).

Impact on ecosystem processes: *Rhizobia* bacteria located in the roots of yellow alfalfa alter soil conditions by fixing atmospheric nitrogen (USDA 2002). By increasing the nutrient contents of the soil, yellow

alfalfa may facilitate the establishment of other exotic or native species.

Biology and Invasive Potential

Reproductive potential: Yellow alfalfa reproduces by seeds only (USDA 2002). The average number of seeds produced by an individual plant has been documented at 5,320 (Stevens 1932).

Role of disturbance in establishment: Yellow alfalfa generally establishes on disturbed ground. However, it establishes easily after seeding on undisturbed ranch lands (Hultén 1968, Hitchcock and Cronquist 1973, Royer and Dickinson 1999, MAFRI 2004).

Potential for long-distance dispersal: Yellow alfalfa seeds are large and do not disperse easily. Herbivores facilitate the spread of seeds (Leach 1956, Kufeld 1973, Duebber et al. 1981).

Potential to be spread by human activity: Yellow alfalfa is cultivated worldwide. It is planted for erosion-control projects and wildlife habitat and rangeland restoration. It is cultivated for hay and as a nectar source for honey bees (McLean et al. 1971). The utility of this species has probably contributed to its spread.

Germination requirements: Yellow alfalfa seeds require scarification, a burial depth of 6 to 13 mm, and moist conditions to germinate efficiently. Sufficient soil moisture is also critical for root establishment (MAFRI 2004).

Growth requirements: Yellow alfalfa is best adapted to medium-textured soils with pH between 6 and 8. It requires a minimum of 26 cm of precipitation annually, at least half of which must be received as rain (as opposed to snow). Alfalfa can withstand temperatures as low as -36°C. It requires 100 frost-free days to grow and reproduce successfully. Alfalfa is highly tolerant of drought and fire, but it cannot tolerate flooding, poor soil drainage, salinity, or shading (USDA 2002).

Congeneric weeds: Alfalfa (*Medicago sativa* ssp. *sativa*), black medick (*M. lupulina*), burclover (*M. polymorpha*), and burr medick (*M. minima*) are known to occur as non-native weeds in North America (USDA 2002). Burclover is considered a noxious weed in Arizona (Invaders 2010).


Legal Listings

- Has not been declared noxious
- Listed noxious in Alaska
- Listed noxious by other states
- Federal noxious weed
- Listed noxious in Canada or other countries

Distribution and Abundance

Yellow alfalfa occasionally establishes in roadsides, waste areas (Hultén 1968, Hitchcock and Cronquist 1973), and active or abandoned agricultural fields (Royer and Dickinson 1999).

Native and current distribution: Yellow alfalfa is native to southwestern Asia. It was first cultivated in Iran and now has a worldwide distribution as an agricultural crop. This species is widely planted in the continental U.S. and Canada, and it has naturalized in many of the areas in which it has been cultivated. Yellow alfalfa is not well adapted for cultivation in Alaska (J. Conn – pers. com.). It has been documented from the Pacific Maritime and Interior-Boreal ecogeographic regions of Alaska (AKEPIC 2010).



Distribution of yellow alfalfa in Alaska

Management

Control measures have not been developed because of the value of yellow alfalfa as an agricultural crop. Yellow alfalfa is susceptible to herbicides (Bowes 1982, Cogliastro et al. 1990).

References:

- AKEPIC database. Alaska Exotic Plant Information Clearinghouse Database. 2010. Available: <http://akweeds.uaa.alaska.edu/>
- Bowes, G.G. 1982. Changes in the yield of forage following the use of herbicides to control aspen poplar. *Journal of Range Management*. 35: 246-248.
- Carlson, M. L., Assistant Research Professor, Alaska Natural Heritage Program, University of Alaska Anchorage, 707 A St, Anchorage, AK. Tel: (907) 257-2790 – Pers. obs.
- Cogliastro, A., D. Gagnon, D. Coderre, P. Bhereur. 1990. Response of seven hardwood tree species to herbicide, rototilling, and legume cover at two southern Quebec plantation sites. *Canadian Journal of Forestry*. 20: 1172-1182.
- Conn, J. Weed Scientist, USDA Agricultural Research Service PO Box 757200 Fairbanks, Alaska 99775 tel: (907) 474-7652; fax (907) 474-6184. – Pers. com.
- Duebber, H.F., E.T. Jacobson, K.F. Hoggins, E.B. Podoll. 1981. Establishment of seeded

- grasslands for wildlife habitat in the prairie pothole region. Special Scientific Report – Wildlife No. 234. Washington, DC: U.S. Department of the Interior, Fish and Wildlife Service. 21 p.
- Graham, E.H. 1941. Legumes for erosion control and wildlife. Misc. Publ. 412. Washington, DC: U.S. Department of Agriculture. 153 p.
- Hitchcock, C. L., A. Cronquist. 1973. Flora of the Pacific Northwest. University of Washington Press, Seattle and London. 730 p.
- Hultén, E. 1968. Flora of Alaska and Neighboring Territories. Stanford University Press, Stanford, CA. 1008 p.
- Invaders Database System. 2010. University of Montana. Missoula, MT.
<http://invader.dbs.umt.edu/>
- ITIS. 2010. Integrated Taxonomic Information System.
<http://www.itis.gov/>
- Klett, A.T., H.F. Duebbert, G.L. Heismeyer. 1984. Use of seeded native grasses as nesting cover by ducks. Wildlife Society Bulletin. 12: 134-138.
- Kufeld, R.C. 1973. Food eaten by the Rocky Mountain elk. Journal of Range Management. 26: 106-113.
- Leach, H.R. 1956. Food habits of the Great Basin deer herds of California. California Fish and Game. 38: 243-308.
- MAFRI - Manitoba Agriculture, Food and Rural Initiatives. 2004. Crops, Alfalfa seed production. Available:
<http://www.gov.mb.ca/agriculture/index.shtml> [October 4, 2005].
- McLean, A., T.M. Lord, A.J. Green. 1971. Utilization of the major plant communities in the Similkameen Valley, British Columbia. Journal of Range Management. 24: 346-142.
- Royer, F., and R. Dickinson. 1999. Weeds of the Northern U.S. and Canada. The University of Alberta press. 434 pp.
- Small, E. and B.S. Brookes. 1984. Taxonomic circumscription and identification in the *Medicago sativa-falcata* (Alfalfa) continuum. Economic Botany. 38: 83-96.
- Stanton, F. 1974. Wildlife guidelines for range fire rehabilitation. Tech. Note 6712. Denver, CO: U.S. Department of the Interior, Bureau of Land Management. 90 p.
- Stevens, O.A. 1932. The number and weight of seeds produced by weeds. American Journal of Botany, 19: 784-794.
- USDA (United States Department of Agriculture), NRCS (Natural Resource Conservation Service). 2002. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.
- Weeds of Alaska Database. 2005. AKEPIC Mapping Project Inventory Field Data. Alaska Natural Heritage Program, University of Alaska – US Forest Service – National Park Service. Available: <http://akweeds.uaa.alaska.edu/>