

Contributions to the knowledge of the diversity of the chewing lice fauna in Turkey

Gökhan EREN^{1*}, Ömral Ünsal ÖZKOÇ², Mustafa AÇICI¹

¹Department of Parasitology, Faculty of Veterinary Medicine, Ondokuz Mayıs University, Samsun, Turkey

²Turkish Nature Research Society, Ankara, Turkey

Received: 06.06.2022

Accepted/Published Online: 10.10.2022

Final Version: 10.11.2022

Abstract: The aim of the study was to identify the chewing lice which were sampled from the birds found dead during ornithological studies in the natural areas of Samsun, and transported to the laboratory of Ondokuz Mayıs University, Veterinary Faculty, Department of Parasitology, between 2018 and 2020. All samples were collected by ectoparasite examination. The collected lice specimens were preserved in tubes with 70% alcohol. The specimens were cleared in 10% KOH for 24 h before being mounted in Canada balsam and identified under the light microscope. A total of 35 species under 3 families were detected in this study: Menoponidae: *Colpocephalum apivorus*, *C. impressum*, *C. nanum*, *C. napiforme*, *C. turbinatum*, *Colpocephalum* sp., *Pseudomenopon dolium*, *Kurodaia subpachygaster*, *Ciconiphilus cygni*, *C. decimfasciatus*, *C. quadripustulatus*, *Menacanthus agilis*, *M. curuccae*, *M. eurysternus*, *M. gonophaeus*, *M. stramineus*, *Menacanthus* sp., *Trinoton anserinum*, *T. querquedulae*; Laemobothriidae: *Laemobothrion maximum*; Philopteriidae: *Anaticola crassicornis*, *A. mergiserrati*, *Anatoecus cygni*, *Anatoecus* sp., *Aquanirmus podiceps*, *Brueelia merulensis*, *Columbicula columbae*, *Cummingsiella ovalis*, *Degeeriella fulva*, *D. fusca*, *D. phlyctopygus*, *Craspedorrhynchus aquilinus*, *Incidifrons fulicae*, *Ornithobius cygni*, *Philopterus atratus*, *Saemundssonina lari*, *Strigiphilus cursitans*, and *Quadriceps punctatus*. The first records for the fauna of Turkey are: *Anatoecus cygni*, *Ciconiphilus cygni*, *Colpocephalum apivorus*, *C. napiforme*, *Kurodaia subpachygaster*, *Ornithobius cygni*, and *Philopterus atratus*. As a result of this study, the number of chewing lice species (Phthiraptera, Amblycera, and Ischnocera) detected on birds in Turkey reached 195 species.

Key words: Amblycera, birds, chewing lice, Ischnocera

1. Introduction

Around 4000 species of lice-infested birds have been identified in studies carried out so far around the world (Price et al. 2003). Species infesting birds are classified in the suborders Amblycera and Ischnocera and feed on the feathers and other epidermal structures of their hosts. Species in these two suborders are also called “chewing lice” because of their types of feeding. Most of the chewing lice species show high host specificity and exhibit one of the most important models of host-parasite evolution (Clayton et al., 2003). Mild infestations are usually subclinical, while heavy infestations may cause clinical symptoms. Furthermore, some lice species are vectors of filarial nematodes (Clayton et al., 2008; Cohen et al., 1991). Studies on bird chewing lice (Amblycera and Ischnocera) in Turkey have increased in recent years. Although the ornithofauna of Turkey is represented by nearly 500 species (Furtun et al., 2021), only quarter of these species have been studied in terms of lice infestation. With this study, 195 species of chewing lice have been detected from 126 different bird species (exotic and domestic and wild

species) in Turkey so far (Table 1). When 43 species of chewing and blood-sucking lice (Merdivenci, 1970; Dik, 2020; Eren et al., 2021; Keskin and Dik, 2022) that were identified in mammals, including bloodsucking species in humans, were also accounted for, the lice fauna of Turkey (Phthiraptera; Amblycera, Anoplura, and Ischnocera) consists of 238 species.

The aim of this study is to contribute to the ornithoparasitological studies that have increased in recent years and the Phthiraptera fauna of Turkey.

2. Materials and methods

This study was carried out using lice samples collected from birds found dead in the natural areas of Samsun, and transported to the laboratory of Ondokuz Mayıs University, Faculty of Veterinary Medicine, Department of Parasitology, between 2018 and 2020. Birds were identified by an ornithologist based on morphological criteria before systemic ectoparasite examination. After the collection of the lice samples, they were preserved in 70% ethanol. After 1 day of clearing in 10% KOH, the specimens were

* Correspondence: gokhaneren54@gmail.com

Table 1. List of identified taxa of chewing lice reported in Turkey so far.

	Family Menoponidae
Suborder Amblycera	<i>Actornithophilus multisetosus</i> , <i>A. patellatus</i> , <i>A. piceus lari</i> , <i>A. pustulosus</i> , <i>A. spinulosus</i> , <i>A. stictus</i> , <i>A. totani</i> , <i>A. umbrinus</i> ; <i>Afrimenopon waar</i> ; <i>Austromenopon alpinum</i> , <i>A. atrofulvum</i> , <i>A. durisetosum</i> , <i>A. lutescens</i> , <i>A. transversum</i> , <i>Austromenopon</i> sp.; <i>Ciconiphilus cygni</i> *, <i>C. decimfasciatus</i> *, <i>C. quadripustulatus</i> ; <i>Colpocephalum apivorus</i> *, <i>C. eucarenum</i> , <i>C. heterosoma</i> , <i>C. impressum</i> , <i>C. milvi</i> , <i>C. nanum</i> , <i>C. napiforme</i> *, <i>C. polonum</i> , <i>C. subzerafae</i> , <i>C. trachelioti</i> , <i>C. turbinatum</i> , <i>C. zebra</i> , <i>Colpocephalum</i> sp.; <i>Comatomenopon elongatum</i> ; <i>Cuculiphilus fasciatus</i> ; <i>Dennyus hirundinis</i> ; <i>Eidmanniella pellucida</i> ; <i>Holomenopon clypeilargum</i> , <i>H. leucoxanthum</i> , <i>H. obscurum</i> , <i>H. tadornae</i> , <i>Holomenopon</i> sp.; <i>Kurodaia fulvofasciata</i> , <i>K. longipes</i> , <i>K. subpachygaster</i> *; <i>Menacanthus abdominalis</i> , <i>M. agilis</i> , <i>M. Alaudae</i> , <i>M. camelinus</i> , <i>M. cornutus</i> , <i>M. curuccae</i> , <i>M. eurysternus</i> , <i>M. fertilis</i> , <i>M. gonophaeus</i> *, <i>M. lyali</i> , <i>M. pusillus</i> , <i>M. stramineus</i> , <i>Menacanthus</i> sp.; <i>Menopon gallinae</i> ; <i>Meromenopon meropis</i> ; <i>Myrsidea cucullaris</i> , <i>M. isostoma</i> , <i>M. picae</i> , <i>M. rustica</i> , <i>Myrsidea</i> sp.; <i>Piagetiella titan</i> ; <i>Pseudomenopon concretum</i> , <i>P. dolium</i> , <i>P. pilosum</i> , <i>P. scopulacorne</i> ; <i>Trinoton anserinum</i> , <i>T. femoratum</i> , <i>T. querquedulae</i> , <i>Trinoton</i> sp.
	Family Laemobothriade
	<i>Laemobothrion atrum</i> , <i>L. maximum</i> , <i>L. tinnunculi</i> , <i>Laemobothrion</i> sp.
	Family Ricinidae
	<i>Ricinus dolicocephalus</i> , <i>R. elongatus</i> , <i>R. frenatus</i> , <i>R. fringillae</i>
	Family Philopteridae
Suborder Ischnocera	<i>Alcedoffula alcedinis</i> ; <i>Anaticola anseris</i> , <i>A. crassicornis</i> , <i>A. magnificus</i> , <i>A. mergiserrati</i> , <i>A. phoenicopteri</i> ; <i>Anatoecus cygni</i> *, <i>A. icterodes</i> , <i>A. dentatus</i> , <i>A. pygaspis</i> , <i>A. regina</i> , <i>Anatoecus</i> sp.; <i>Ardeicola celeris</i> , <i>A. ciconiae</i> , <i>A. ixobrychae</i> , <i>A. plataleae</i> ; <i>Aquanirmus podiceps</i> ; <i>Brueelia apiastri</i> , <i>B. biocellata</i> , <i>B. cruciata</i> , <i>B. domestica</i> , <i>B. iliaci</i> , <i>B. jacobi</i> , <i>B. lais</i> , <i>B. locustellae</i> , <i>B. marginata</i> , <i>B. merulensis</i> , <i>B. munda</i> , <i>B. nebulosa</i> , <i>B. tasniemae</i> , <i>B. turdinulae</i> , <i>Brueelia</i> sp.; <i>Campanulotes bidentatus</i> ; <i>Carduiceps meinertzhageni</i> , <i>C. scalaris</i> , <i>C. zonarius</i> ; <i>Chelopistes meleagridis</i> ; <i>Coloceras chinensea</i> , <i>C. hilli</i> , <i>C. piageti</i> ; <i>Columbicola bacillus</i> , <i>C. columbae</i> ; <i>Cuclotogaster heterographus</i> , <i>C. cinereus</i> , <i>Cuclotogaster</i> sp.; <i>Cuculicola latirostris</i> ; <i>Cuculoecus latifrons</i> ; <i>Cummingsiella ambigua</i> , <i>C. ovalis</i> ; <i>Craspedorrhynchus fraterculus</i> , <i>C. platystomus</i> , <i>Craspedorrhynchus</i> sp.; <i>Degeeriella aquilarum</i> , <i>D. fulva</i> , <i>D. fusca</i> , <i>D. leucopleura</i> , <i>D. nisus</i> , <i>D. phlyctopygus</i> , <i>D. rufa</i> , <i>Degeeriella</i> sp.; <i>Falcolipeurus suturalisa</i> ; <i>Fulicoffula gallinula</i> , <i>F. lurida</i> ; <i>Goniocotes gallinae</i> , <i>G. hologaster</i> , <i>G. megaloccephalus</i> , <i>G. pusillus</i> ; <i>Goniodes astrocephalus</i> , <i>G. colchici</i> , <i>G. dissimilis</i> , <i>G. gigas</i> , <i>G. dispar</i> , <i>G. astrocephalus</i> , <i>G. pavonis</i> ; <i>Ibidococcus plataleaea</i> ; <i>Incidifrons fulica</i> ; <i>Lipeurus caponis</i> ; <i>Lunaceps drosti</i> , <i>L. actophilus</i> , <i>L. holophaeus</i> , <i>L. incoenis</i> , <i>L. numeni numeni</i> , <i>Lunaceps</i> sp.; <i>Meropoecus meropis</i> ; <i>Multicola hypoleucus</i> ; <i>Neophilopterus incompletus</i> , <i>N. tricolor</i> ; <i>Ornithobius cygni</i> *; <i>Pectinopygus brevicornis</i> , <i>P. bifasciatus</i> , <i>P. excornis</i> , <i>P. forficulatus</i> , <i>Pectinopygus</i> sp.; <i>Penenirmus affectator</i> , <i>P. longuliceps</i> , <i>P. pikulai</i> , <i>P. rarus</i> , <i>P. serrilimbus</i> , <i>P. silvicultrix</i> , <i>Penenirmus</i> sp.; <i>Philopterus atratus</i> *, <i>P. desertus</i> , <i>P. eurasiaticus</i> , <i>P. fringillae</i> , <i>P. mirificus</i> , <i>P. rapax</i> , <i>P. reguli</i> , <i>P. sitta</i> ; <i>Rallicola fulica</i> , <i>R. lugens</i> , <i>R. minutus</i> , <i>R. ortyometrae</i> ; <i>Rhynonirmus scolopacis</i> ; <i>Quadriceps anagrapsus</i> , <i>Q. nyctemerus</i> , <i>Q. obscurus</i> , <i>Q. punctatus</i> , <i>Q. punctatus sublingulatus</i> ; <i>Rhynonirmus helvolus</i> , <i>R. scolopacis</i> ; <i>Saemundssonsonia clayae</i> , <i>S. lari</i> , <i>S. lobaticeps</i> ; <i>Strigiphilus barbatus</i> , <i>S. cursitans</i> , <i>S. strigis</i> ; <i>Sturnidoecus pflergi</i> , <i>S. sturni</i> ; <i>Struthiolipeurus struthionisa</i> ; <i>Upupicola upupae</i>
	References

mounted using the Canada balsam technique (Palma, 1978). The specimens were examined under a light microscope (Nikon Eclipse 80i) and comparisons with the literature were performed for species identification (Clay and Hopkins, 1950; Timmermann, 1952; Clay, 1957; Price and Beer, 1963a; Price and Beer, 1963b; Price and Beer, 1965; Nelson and Price, 1965; Edwards, 1965; Price, 1974; Eichler et al., 1981; Clayton and Price, 1984; Soler-Cruz et

al., 1985; Gallego et al., 1987; Price and Hellenthal, 1998; Adams et al., 2005; Arnold, 2005; Adam and Daróczy, 2006; Castro and Cicchino, 1983; Naz et al., 2010). All identified specimens are being preserved in the same laboratory.

3. Results

In this study, 24 species of birds from nine orders (Accipitriformes, Anseriformes, Charadriiformes,

Ciconiiformes, Columbiformes, Falconiformes, Passeriformes, Pelecaniformes, Podicipediformes and Strigiformes) were examined for ectoparasites between 2018 and 2020. In total, 35 species belonging to 3 different families of chewing lice were identified (Table 2): Menoponidae: *Colpocephalum apivorus*, *C. impressum*, *C. nanum*, *C. turbinatum*, *Colpocephalum* sp., *Pseudomenopon dolium*, *Kurodaia subpachygaster*, *Ciconiphilus decimfasciatus*, *C. quadripustulatus*, *C. cygni*, *Menacanthus agilis*, *M. curuccae*, *M. eurysternus*, *M. gonophaeus*, *M. stramineus*, *Menacanthus* sp., *Trinoton anserinum*, *T. querquedulae*; Laemobothriidae: *Laemobothrion maximum*; Philopteridae: *Anaticola crassicornis*, *A. mergiserrati*, *Anatoecus cygni*, *Anatoecus* sp., *Aquanirmus podiceps*, *Brueelia merulensis*, *Columbicula columbae*, *Cummingsiella ovalis*, *Degeeriella fulva*, *D. fusca*, *D. turbinatum*, *Craspedorrhynchus aquilinus*, *Incidifrons fulicae*, *Ornithobius cygni*, *Philopterus atratus*, *Saemundssonina lari*, *Strigiphilus cursitans* and *Quadriceps punctatus*. The first records for the fauna of Turkey are: *Anatoecus cygni*, *Ciconiphilus cygni*, *Colpocephalum apivorus*, *C. napiforme*, *Kurodaia subpachygaster*, *Ornithobius cygni*, and *Philopterus atratus*. As a result of this study, the number of chewing lice species (Phthiraptera: Amblycera, Ischnocera) detected on birds in Turkey reached 195. Infestations of *Colpocephalum* and *Menacanthus* spp. were more intense on the hosts. Photographs of the lice species, the species recorded for the first time by this study, are presented in Figure.

4. Discussion

Chewing lice (Phthiraptera: Amblycera and Ischnocera) can cause many direct or indirect effects on birds. For dermatitis, hyperkeratosis, pruritis, insomnia, hemorrhagic ulcerative stomatitis in *Piagetiella* infestations, excessive feather growth, deterioration in the quality of the feathers, disruption of thermoregulation by making small holes in the feathers and breaking of the feathers, the prolongation of the migration period and the delay of reproduction due to the deterioration of the feather structure, especially in migratory birds, and the decrease in the chance of survival as a result of the decrease in the body condition score are the most important direct effects (Clayton et al., 2008; Moreno-Rueda and Hoi, 2012). The indirect effect is negative sexual selection—birds with a low parasite load have more sexual feathers, and females prefer males with more sexual feathers to mate with, thus decreasing breeding success because of the shortening of the singing duration and reducing the quality of the singing of infested birds (Clayton, 1991; Garamszegi, 2005).

Although studies about chewing lice on birds have increased in recent years in Turkey, the number of birds examined and reported lice species is still not at the desired

level. More than 120 bird species have been examined in studies and 188 lice species—together with this study, a total of 195—species have been reported (Table 1). Some individuals of chewing lice can be overlooked during macroscopic examination due to the rapid movement of chewing lice species (Johnson and Clayton, 2003). In addition, factors such as the size of the host, the density of feathers, and the severity of the infestation make detection difficult. Synthetic pyrethroid insecticides are also used to make the collection of ectoparasite samples easier in ornitho-parasitological studies (Clayton and Walther, 1997). In ectoparasite studies conducted on dead birds, infestation rates may be lower than in other studies since the host is not alive and the ectoparasites may have left the host due to the long time elapsed since death. As chewing lice are host-obligate ectoparasites, they leave the host shortly after its death. The members of Ischnocera feed on the skin for a while, while the members of Amblycera leave the host more rapidly (Johnson and Clayton 2003).

Chewing lice of bird species in the order Columbiformes have been extensively studied throughout the world and in Turkey. As a result of these studies, species belonging to the genus *Auricotes*, *Bonomiella*, *Campanulotes*, *Cavifera*, *Coloceras*, *Colpocephalum*, *Hohorstiella*, *Goniocotes*, *Kodocephalon*, *Menacanthus*, *Menopon*, *Neomenopon*, *Physconelloides*, *Syrrhaptoecus*, *Turturicola*, and *Quateia* have been reported (Price et al., 2003; Adang et al., 2008; Jahantigh et al., 2016). *Columbicola columbae*, *Goniocotes bidentatus* (*Campanulotes compar*), *Goniocotes hologaster* (*Goniocotes gallinae*), *Lipeurus caponis*, *Menopon gallinae*, *Coloceras israelensis* were identified in the studies on domestic pigeons *Columba livia domestica* and rock pigeons (wild type of *Columba livia*) in Turkey. These studies show that *Columbicola columbae* is the dominant species causing infestation on rock dove *Columba livia* both in the world and Turkey. Besides these, infestation of *Columbicola bacillus*, *Coloceras hillia*, and *Coloceras piageti* on Eurasian collared dove *Streptopelia decaocto* (Dik, 2010; 2013; Girişgin et al., 2013); *Columbicola bacillus* on European turtle dove *Streptopelia turtur* (Girişgin et al., 2013; Açııcı et al., 2021b); and *Coloceras chinense*, *Columbicola bacillus*, *C. columbae*, and *Cuclotogaster heterographus* on laughing dove *Spilopelia senegalensis* (Girişgin et al., 2013; 2022) were determined. In this study, *Columbicola columbae* and *Menacanthus stramineus* were determined on rock dove *Columba livia* that has been reported as a host of *M. stramineus* in Turkey for the first time.

Infestations in owls belonging to the order Strigiformes are caused by the genera *Colpocephalum*, *Kurodaia*, and *Strigiphilus* (Price et al., 2003). A limited number of studies have been conducted on the chewing lice of owls in Turkey: *Strigiphilus cursitans* on little owl *Athene noctua* (Girişgin

Table 2. The identified chewing lice species and their hosts (*new record, **new host record).

Family	Genus	Species	Host
Philopteridae	<i>Quadriceps</i>	<i>Q. punctatus</i>	<i>Larus michahellis</i> (n: 3)
	<i>Saemundssonina</i>	<i>S. lari</i>	
Menoponidae	<i>Colpocephalum</i>	<i>C. impressum</i>	<i>Aquila chrysaetos</i> (n: 2)
Philopteridae	<i>Degeeriella</i>	<i>D. fulva</i>	
	<i>Craspedorrhynchus</i>	<i>C. aquilinus</i>	
Menoponidae	<i>Colpocephalum</i>	<i>C. apivorus</i>	<i>Pernis apivorus</i> (n: 1)
Philopteridae	<i>Degeeriella</i>	<i>D. phlyctopygus</i> **	
Menoponidae	<i>Colpocephalum</i>	<i>Colpocephalum</i> sp.	<i>Buteo rufinus</i> (n: 2)
Philopteridae	<i>Craspedorrhynchus</i>	<i>C. platystomus</i>	
	<i>Degeeriella</i>	<i>D. fulva</i>	
Menoponidae	<i>Colpocephalum</i>	<i>C. napiforme</i> *	<i>Pandion haliaetus</i> (n: 1)
Philopteridae	<i>Degeeriella</i>	<i>D. phlyctopygus</i>	<i>Podiceps cristatus</i> (n: 1)
Philopteridae	<i>Aquanirmus</i>	<i>A. podiceps</i>	
Menoponidae	<i>Pseudomenopon</i>	<i>P. dolium</i>	<i>Buteo buteo</i> (n: 5)
Laemobothriidae	<i>Laemobothrion</i>	<i>L. maximum</i>	
	Philopteridae	<i>Degeeriella</i>	
Menoponidae	<i>Colpocephalum</i>	<i>C. nanum</i> <i>C. turbinatum</i>	
Philopteridae	<i>Strigiphilus</i>	<i>S. cursitans</i>	<i>Athene noctua</i> (n: 1)
Menoponidae	<i>Kurodaia</i>	<i>K. subpachygaster</i> *	<i>Tyto alba</i> (n: 1)
Menoponidae	<i>Ciconiphilus</i>	<i>C. decimfasciatus</i>	<i>Ardea cinerea</i> (n: 1)
Menoponidae	<i>Ciconiphilus</i>	<i>C. decimfasciatus</i> **	<i>Egretta garzetta</i> (n: 2)
Menoponidae	<i>Ciconiphilus</i>	<i>C. quadripustulatus</i>	<i>Ciconia ciconia</i> (n: 2)
Philopteridae	<i>Degeeriella</i>	<i>D. fusca</i>	<i>Circus aeruginosus</i> (n: 1)
Menoponidae	<i>Menacanthus</i>	<i>M. gonophaeus</i> **	<i>Corvus frugilegus</i> (n: 1)
Philopteridae	<i>Philopterus</i>	<i>P. atratus</i> *	
Philopteridae	<i>Anatoecus</i>	<i>Anatoecus</i> sp.	<i>Netta rufina</i> (n: 1)
Philopteridae	<i>Cummingsiella</i>	<i>C. ovalis</i>	<i>Numenius arquata</i> (n: 1)
Philopteridae	<i>Anatoecus</i>	<i>A. cygni</i> *	<i>Cygnus cygnus</i> (n: 1)
	<i>Ornithobius</i>	<i>O. cygni</i> *	
	<i>Anaticola</i>	<i>A. crassicornis</i> **	
Menoponidae	<i>Ciconiphilus</i>	<i>C. cygni</i> *	
	<i>Trinoton</i>	<i>T. anserinum</i>	
Philopteridae	<i>Brueelia</i>	<i>B. merulensis</i>	<i>Turdus merula</i> (n: 1)
Menoponidae	<i>Menacanthus</i>	<i>M. eurysternus</i>	<i>Lanius collurio</i> (n: 1)
Menoponidae	<i>Menacanthus</i>	<i>Menacanthus</i> sp..	
Menoponidae	<i>Menacanthus</i>	<i>M. agilis</i>	<i>Phylloscopus collybita</i> (n: 2)
Menoponidae	<i>Menacanthus</i>	<i>M. curuccae</i>	<i>Sylvia atricapilla</i> (n: 2)
Philopteridae	<i>Columbicola</i>	<i>C. columbae</i>	<i>Columba livia</i> (n: 1)
Menoponidae	<i>Menacanthus</i>	<i>M. stramineus</i> **	
Philopteridae	<i>Incidifrons</i>	<i>I. fulicae</i>	<i>Fulica atra</i> (n: 1)
Menoponidae	<i>Trinoton</i>	<i>T. querquedulae</i> **	<i>Aythya ferina</i> (n: 1)
Philopteridae	<i>Anaticola</i>	<i>A. mergiserrati</i>	




Figure. Original photos of chewing lice identified in this study: A, *Anatoecus cygni* (♀); B, *Ciconiphilus cygni* (♀); C, *Ciconiphilus decimfasciatus* (♀); D, *Colpocephalum apivorus* (♂); E, *Colpocephalum apivorus* (♀); F, *Colpocephalum napiforme* (♂); G, *Kurodaia subpachygaster* (♀); H, *Ornithobius cygni* (♀); I, *Philopterus atratus* (♀) (bar: 1 mm).

et al., 2013; 2022); *Strigiphilus barbatus* and *Columbicola columbae* on long-eared owl *Asio otus* (İnci et al., 2010; Dik et al., 2013); *Strigiphilus barbatus*, *Strigiphilus strigis* and *Kurodaia longipes* on Eurasian eagle-owl *Bubo bubo* (Dik and Uslu, 2007; Dik, 2010; Dik et al., 2015); *Colpocephalum* sp. on western barn owl *Tyto alba* (Girişgin et al., 2022) caused infestations. In this study, *S. cursitans* on little owl and *Athene noctua* and *K. subpachygaster* on western barn owl *Tyto alba* were identified. *Kurodaia subpachygaster* is a new record for the Phthiraptera fauna of Turkey.

Studies conducted on species in the order Podicipediformes (Price et al., 2003) have identified the lice of genera *Aquanirmus*, *Laemobothrion*, and *Pseudomenopon*. The studies of the effects of chewing lice on the hosts of this order are extremely limited in Turkey. Only infestations of *Aquanirmus podiceps* and *Pseudomenopon dolium* were detected in this order (Dik et al., 2017). In this study, both species were found on great crested grebe *Podiceps cristatus*.

Studies on the members of the Rallidae (rails) and Gruidae (cranes) in Turkey are extremely insufficient. In species of the order Gruiformes listed in the fauna of Turkey (Furtun et al. 2021), genera *Esthiopterum*, *Fulicoffula*, *Gruimenopon*, *Heleonomus*, *Incidifrons*, *Laemobothrion*, *Pseudomenopon*, *Rallicola*, and *Saemundssonina* can cause infestations (Price et al., 2003). The cases reported for infestations caused by genera *Fulicoffula*, *Incidifrons*, *Laemobothrion*, and *Pseudomenopon* species were as follows: infestations of *Fulicoffula lurida*, *Incidifrons fulicae*, *Laemobothrion atrum*, *Pseudomenopon pilosum*, and *Rallicola fulicae* on Eurasian coot *Fulica atra* (Dik et al., 2011a; 2017); *Fulicoffula gallinula*, *Rallicola minutus*, and *Pseudomenopon pilosum* on common moorhen *Gallinula chloropus* (Dik et al., 2011a; 2017; Girişgin et al., 2022); *Pseudomenopon concretum* and *Rallicola lugens* on Western swamphen *Porphyrio porphyrio* (Açıcı et al., 2021b); *Pseudomenopon scopulacorne* on little crane *Zapornia parva* (Dik et al., 2011c); and *Pseudomenopon scopulacorne* on water rail *Rallus aquaticus* (Dik et al., 2011c) were reported. In this study, an infestation of *Incidifrons fulicae* was reported in the Eurasian coot *Fulica atra*.

As a result of studies on storks (Ciconiiformes), it has been reported that *Anaticola*, *Anatoecus*, *Austromenopon*, *Ardeicola*, *Ciconiphilus*, *Colpocephalum*, *Comatomenopon*, *Laemobothrion*, *Neophilopteris*, and *Trinoton* species cause infestation (Price et al., 2003). In Turkey, infestation of *Ardeicola ciconiae*, *Ciconiphilus quadripustulatus*, *Colpocephalum zebra* and *Neophilopteris incompletus* on white stork *Ciconia ciconia* (İnci et al., 2010; Dik et al., 2011, Girişgin et al., 2013; Dik et al., 2013; 2021; Açıcı et al., 2021b; Girişgin et al., 2022); infestation of *Neophilopteris tricolor* were reported on black stork *Ciconia nigra* (Açıcı

et al., 2021b). In this study, an infestation of *Ciconiphilus quadripustulatus* was reported on white stork *Ciconia ciconia*.

Species of belonging to the genera *Ardeiphagus*, *Ardeiphilus*, *Ciconiphilus*, *Comatomenopon*, *Colpocephalum*, *Eucolpocephalum*, *Ibidoecus*, *Laemobothrion*, *Pectinopygus*, *Piagetiella*, and *Plegadiphilus* cause infestation on pelicans, herons, and ibises in the order Pelecaniformes (Price et al., 2003). Infestation of *Comatomenopon elongatum* on great egret *Ardea alba* (İnci et al., 2010; Girgin et al., 2022); *Ciconiphilus decimfasciatus* on grey heron *Ardea cinerea* (Girişgin et al., 2022); *Ardeicola celeris* and *Ardeicola ixobrychae* on little bittern *Ixobrychus minutus* (Dik et al., 2017b; Girişgin et al., 2022); *Colpocephalum eucarenum*, *Piagetiella titan*, *Pectinopygus forficulatus*, and *Pectinopygus* sp. on great white pelican *Pelecanus onocrotalus* (Dik, 2006b; Dik & Uslu, 2008; Girişgin et al., 2013; Dik et al., 2012; 2017a; 2021); *Piagetiella titan* and *Pectinopygus* sp. on Dalmatian pelican *Pelecanus crispus* (Girişgin et al., 2013); and *Ibidoecus platalaeae* and *Ardeicola platalaeae* on Eurasian spoonbill *Platalea leucorodia* (Girişgin et al., 2013) were reported in the studies on this order conducted in Turkey. *Ciconiphilus decimfasciatus* was reported on grey heron *Ardea cinerea* and little egret *Egretta garzetta*. Along with this study, *Ciconiphilus decimfasciatus* is a new host for *Ardea cinerea* in Turkey.

Waders, gulls, and terns of the order Charadriiformes were listed in the ornithofauna of Turkey (Kirwan et al., 2010; Furtun et al., 2021). In the studies around the world, genera *Actornithophilus*, *Austromenopon*, *Carduiceps*, *Cirrophthirius*, *Cummingsiella*, *Haffneria*, *Lunaceps*, *Rediella*, *Rhynonirmus*, *Saemundssonina*, and *Quadriceps* cause infestation on the members of these groups (Price et al., 2003). The infestation of *Saemundssonina lari* and *Quadriceps punctatus* on yellow-legged gull *Larus michahellis* (Collinson et al., 2008)—this host is probably misidentified because Armenian gull *Larus armenicus* is distributed in the region where the study was carried out, but yellow-legged gull is not found in the same region (Furtun et al., 2021) and it is difficult to identify these two species (van Duivendijk, 2011)—; *Actornithophilus piceus* and *Saemundssonina lari* on Caspian gull *Larus cachinnans* and slender-billed gull *Chroicocephalus genei* (Dik et al., 2011; Girişgin et al., 2022); *Austromenopon transversum*, *Saemundssonina lari* and *Quadriceps punctatus* (Dik et al., 2017); *Actornithophilus umbrinus*, *Austromenopon alpinum*, *Carduiceps meinertzhagani* and *Lunaceps actophilus* on dunlin *Calidris alpina* (Dik et al., 2010; Açıcı et al., 2011; 2021b); *Actornithophilus umbrinus*, *Austromenopon lutescens*, *Carduiceps zonarius* and *Lunaceps drosti* on little stint *Calidris minuta* (Dik et al., 2010); *Lunaceps incoensis* on Temminck's stint *Calidris temminckii* (Dik et al., 2010); *Austromenopon atrofulvum*, *Quadriceps anagrapsus*, and

Saemundssonina lobaticeps on white-winged tern *Chlidonias leucopterus* (Dik et al., 2010;2011a); *Actornithophilus stictus*, *Austromenopon durisetosum* and *Rhynonirmus scolopacis* on *Gallinago gallinago* (Dik et al., 2010; 2011b); *Actornithophilus spinulosus* and *Luniceps* sp. on black-tailed godwit *Limosa limosa* (Dik et al., 2017); *Actornithophilus multisetosus* on jack snipe *Lymnocyptes minimus* (Dik et al., 2011b); *Actornithophilus patellatus*, *Cummingsiella ovalis*, and *Luniceps numenii numenii* on Eurasian curlew *Numenius arquata* (Açııcı et al., 2021b); *Actornithophilus pustulosus*, *Austromenopon lutescens*, *Carduiceps scalaris*, and *Luniceps holophaeus* on ruff *Calidris pugnax* (Dik et al., 2010); *Quadriceps anagrapsus* on little tern *Sternula albifrons* (Açııcı et al., 2011); *Rhynonirmus helvolus*, *Saemundssonina clayae*, and *Saemundssonina* sp. on Eurasian woodcock *Scolopax rusticola* (Dik et al., 2015; Girişgin et al., 2022); *Actornithophilus totani*, *Austromenopon* sp., and *Quadriceps obscurus* on wood sandpiper *Tringa glareola* (Dik et al., 2010;2017) were reported in Turkey. *Saemundssonina lari* and *Quadriceps punctatus* on yellow-legged gull *Larus michahellis* and *Cummingsiella ovalis* on *Numenius arquata* were reported in this study. Along with this study, *Larus michahellis* is a new host for *Saemundssonina lari* and *Quadriceps punctatus* in Turkey.

Species belonging to the genera *Acidoproctus*, *Anaticola*, *Anatoecus*, *Austrogoniodes*, *Ciconiphilus*, *Holomenopon*, *Ornithobius*, *Saemunssonina*, and *Trinoton* cause infestation on the hosts of the order Anseriformes (ducks, geese and swans) (Price et al., 2003). The infestation of *Anaticola crassicornis*, *Anatoecus icterodes*, *Holomenopon clypeilargum*, *Holomenopon* sp., and *Trinoton querquedulae* on Northern Pintail *Anas acuta* (Dik et al., 2011c; Dik et al., 2012;b2017); *Anaticola crassicornis*, *Trinoton querquedulae*, and *Holomenopon* sp. on northern shoveler *Spatula clypeata* (Dik et al., 2012); *Anaticola crassicornis*, *Anatoecus dentatus*, *A. icterodes*, and *Trinoton querquedulae* on Eurasian teal *Anas crecca* (Dik et al., 2012; 2017); *Anaticola crassicornis*, *Anatoecus icterodes*, *Holomenopon clypeilargum*, *Holomenopon leucoxanthum*, *Trinoton querquedulae*, and *Anatoecus* sp. on mallard *Anas platyrhynchos* (Aksin, 2010; Dik et al., 2012; Göz et al., 2015; Açııcı et al., 2021b); *Anaticola* sp. on gadwall *Mareca strepera* (Dik et al., 2017); *Trinoton querquedulae querquedulae* on garganey *Spatula querquedula* (Dik et al., 2012); *Anaticola anseris*, *Trinoton anserinum* and *Holomenopon* sp. on greylag goose *Anser anser* (Göz et al., 2015); *Trinoton anserinum* on whooper swan *Cygnus cygnus* and mute swan *Cygnus olor* (Açııcı et al., 2015; Oğuz et al., 2015; Açııcı et al., 2021b); *Anaticola mergiserrati* and *Anatoecus icterodes* on common pochard *Aythya ferina* (Dik et al., 2017); *Trinoton querquedulae* on tufted duck *Aythya fuligula* (Dik et al., 2017); *Anatoecus icterodes* on red-crested pochard *Netta rufina* (Açııcı et al., 2021b);

Anaticola magnificus, *Anatoecus regina*, *Holomenopon tadornae* and *Trinoton querquedulae* on ruddy shelduck *Tadorna ferruginea* (Dik et al., 2012); *Anaticola crassicornis* and *Trinoton* sp. on common shelduck *Tadorna tadorna* (Dik et al., 2012); *Anatoecus icterodes* on marbled duck *Marmaronetta angustirostris* (Dik, 2010) were reported in Turkey. *Anaticola mergiserrati* and *Trinoton querquedulae* on common pochard *Aythya ferina*; *Anatoecus* sp. on red-crested pochard *Netta rufina*; and *Anaticola crassicornis*, *Anatoecus cygni*, *Ciconiphilus cygni*, *Ornithobius cygni*, and *Trinoton anserinum* on whooper swan *Cygnus cygnus* were reported in this study. *Anatoecus cygni*, *Ciconiphilus cygni*, and *Ornithobius cygni* are reported as the first record for chewing lice fauna of Turkey.

The diurnal predators found in Turkey are classified under the orders Accipitriformes (Accipitridae: hawks, buzzards, kites, harriers, eagles and vultures; Pandionidae: ospreys) and Falconiformes (Falconidae: falcons) (Kirwan et al., 2010; Mindell et al., 2018). The genera *Acutifrons*, *Aegypocetus*, *Colpocephalum*, *Craspedorhynchus*, *Cuculiphilus*, *Degeeriella*, *Falcomenopon*, *Falcolipeurus*, *Falcolius*, *Kurodaia*, *Laemobothrion*, *Neopsittaconirmus*, *Nosopon*, and *Pterophilus* can cause infestation in members of these orders (Price et al., 2003). Many studies have been carried out on the chewing lice of diurnal raptors in Turkey. The infestation of *Colpocephalum polonum* on northern goshawk *Accipiter gentilis* (Dik et al., 2011a); *Colpocephalum nanum*, *C. polonum* and *Degeeriella nisis* on Eurasian sparrowhawk *Accipiter nisis* (Dik et al., 2013; 2017; Girişgin et al., 2022); *Colpocephalum trachelioti*, *Falcolipeurus quadripustulatus*, and *Laemobothrion vulturis* on cinereous vulture *Aegypius monachus* (Dik et al., 2013); *Laemobothrion* sp. on golden eagle *Aquila chrysaetos* (Göz et al., 2015); *Laemobothrion maximum* on golden eagle *Aquila chrysaetos* (Açııcı et al., 2021b); *Laemobothrion maximum* on booted eagle *Hieraaetus pennatus* (Girişgin et al., 2013); *Laemobothrion vulturis* on lesser spotted eagle *Clanga pomarina* (Açııcı et al., 2021b); *Craspedorhynchus* sp., *Colpocephalum nanum*, *C. turbinatum*, *C. platystomus*, *Colpocephalum* sp., *Degeeriella fulva*, *D. nisis*, *Degeeriella* sp., *Falcolipeurus suturalis*, *Kurodaia fulvofasciata*, *Laemobothrion maximum*, and *Laemobothrion* sp. on common buzzard *Buteo buteo* and long-legged buzzard *Buteo rufinus* (Dik, 2006a; Gülanber et al., 2006; Dik et al., 2007; İnci et al., 2010; Dik, 2010; Girişgin et al., 2013; Dik et al., 2015; 2021; Göz et al., 2015; Açııcı et al., 2021b; Girişgin et al., 2022); *Colpocephalum turbinatum*, *Craspedorhynchus rotundatus*, *Degeeriella fusca*, and *Kurodaia fulvofasciata* on *Circus aeruginosus* (Dik et al., 2013; Göz et al., 2015; Dik et al., 2017; Açııcı et al., 2021b); *Degeeriella leucopleura* and *Laemobothrion maximum* on *Circaetus gallicus* (Girişgin et al., 2013; Açııcı et al., 2021b); *Degeeriella rufa* and *Laemobothrion*

tinnunculi on lesser kestrel *Falco naumanni* (Göz et al., 2015); *Colpocephalum subzerafae*, *Degeeriella rufa*, *Degeeriella* sp., and *Laemobothrion tinnunculi* on common kestrel *Falco tinnunculus* (Esatgil et al., 2012; Girişgin et al., 2013; Dik et al., 2013; Girişgin et al., 2022); *Colpocephalum milvi* and *L. maximum* on black kite *Milvus migrans* (İnci et al., 2010); *Degeeriella fulva*, *D. phlyctopygus*, and *Colpocephalum* sp. on European honey buzzard *Pernis apivorus* (İnci et al., 2010; Dik et al., 2013) were reported in Turkey. *Colpocephalum nanum*, *C. turbinatum*, *Degeeriella fulva*, and *Laemobothrion maximum* on common buzzard *Buteo buteo*; *Colpocephalum nanum*, *Craspedorrhynchus platystomus* and *Degeeriella fulva* on long-legged buzzard *Buteo rufinus*; *Colpocephalum apivorus* and *Degeeriella phlyctopygus* on European honey buzzard *Pernis apivorus*; *Degeeriella fusca* on *Circus aeruginosus*; *Colpocephalum impressum*, *Craspedorrhynchus aquilinus*, and *Degeeriella fulva* on golden eagle *Aquila chrysaetos*; *Colpocephalum napiforme* and *Degeeriella phlyctopygus* on western osprey *Pandion haliaetus* were reported in this study. Along with this study, golden *Aquila chrysaetos* is a new host for *Craspedorrhynchus aquilinus* in Turkey

Chewing lice studies on families Acrocephalidae, Aegithalidae, Alaudidae, Cettiidae, Corvidae, Emberizidae, Fringillidae, Hirundinidae, Laniidae, Locustellidae, Motacillidae, Muscipapidae, Oriolidae, Paridae, Passeridae, Phylloscopidae, and Regulicidae belonging to order Passeriformes (Passerines) have been carried out in Turkey. In studies conducted around the world, it has been reported that species that belong to genera *Brueelia*, *Colpocephalum*, *Machaerilaeumus*, *Menacanthus*, *Myrsidea*, *Penenirmus*, *Philoaterus*, *Ricinus*, and *Sturnidoecus* cause infestations (Price et al., 2003). The chewing lice of passerines have been studied in recent years, and many new records have been reported for the Turkish Phthiraptera fauna. The infestations of *Philoaterus mirificus* and *Sturnidoecus pflergi* on marsh warbler *Acrocephalus palustris* (Açııcı et al., 2021b); *Menacanthus curuccae* on Eurasian reed warbler *Acrocephalus scirpaceus* (Açııcı et al., 2011; Dik et al., 2017; Açııcı et al., 2021b); *Menacanthus pusillus* on water pipit *Anthus spinoletta* (Dik et al., 2011c); *Penenirmus longuliceps* on Cetti's warbler *Cettia cetti* (Dik et al., 2015); *Philoaterus eurasiaticus* on hawfinch *Coccothraustes coccothraustes* (Açııcı et al., 2021b); *Brueelia* sp. and *Menacanthus gonophaeus* on northern raven *Corvus corax* (Açııcı et al., 2021b); *Brueelia tasniemae* and *Myrsidea isostoma* on rook *Corvus frugilegus* (Girişgin et al., 2022); *Brueelia* sp. and *Penenirmus* sp. on black-headed bunting *Emberiza melanocephala* (Dik et al., 2015); *Menacanthus chrysophaeus* on common reed bunting *Emberiza schoeniclus* (Dik et al., 2011b); *Menacanthus eurysternus* and *Menacanthus* sp. on European robin *Erithacus rubecula* (Dik et al., 2011c; 2015; 2017); *Brueelia*

sp. on collared flycatcher *Ficedula albicollis* (Dik et al., 2017); *Ricinus fringillae* on common chaffinch *Fringilla coelebs* (Açııcı et al., 2011; 2021b); *Philoaterus rapax* on brambling *Fringilla montifringilla* (Dik et al., 2017); *Brueelia domestica*, *Myrsidea rustica* and *Menacanthus* sp. on barn swallow *Hirundo rustica* (Açııcı et al., 2011; Dik et al., 2011b; 2015; 2017; Açııcı et al., 2021b); *Menacanthus curuccae* on eastern oliveaceous warbler *Iduna pallida* (Dik et al., 2011b); *Brueelia cruciata* and *Menacanthus camelinus* on red-backed shrike *Lanius collurio* (Açııcı et al., 2011; Dik et al., 2011b; 2017; Açııcı et al., 2021b); *Menacanthus camelinus* on lesser grey shrike *Lanius minor* (Açııcı et al., 2021b); *Brueelia locustellae* on Savi's warbler *Locustella luscinioides* (Açııcı et al., 2021b); *Brueelia lais*, *Menacanthus curuccae* on thrush nightingale *Luscinia luscinia* (Dik et al., 2017; Açııcı et al., 2021b); *Brueelia lais* on common nightingale *Luscinia megarhynchos* (Açııcı et al., 2021b); *Menacanthus alaudae* and *Menacanthus pusillus* on *Melanocorypha calandra* (Dik et al., 2011b; 2011c); *Menacanthus pusillus* and *Menacanthus* sp. on western yellow wagtail *Motacilla flava* (Dik et al., 2011b; 2017); *Philoaterus desertus* and *Penenirmus* sp. on spotted flycatcher *Muscicapa striata* (Dik et al., 20137); *Brueelia munda*, *Menacanthus oriolii*, *Ricinus dolicocephalus* and *Maculinirmus mundus* on Eurasian golden oriole *Oriolus oriolus* (Dik et al., 2013b; 2017); *Menacanthus eurysternus* and *Philoaterus fringillae* on house sparrow *Passer domesticus* (Girişgin et al., 2013; Dik et al., 2013b; 2015; 2017); *Penenirmus silvicultrix* on common redstart *Phoenicurus phoenicurus* (Dik et al., 2015); *Menacanthus agilis*, *M. eurysternus*, *M. curuccae*, *Penenirmus rarus*, and *Penenirmus* sp. on common chiffchaff *Phylloscopus collybita* (Dik et al., 2011b; 2011c; 2015; 2017; Açııcı et al., 2021b); *Brueelia* sp., *Menacanthus agilis* and *M. eurysternus* sp. on willow warbler *Phylloscopus trochilus* (Dik et al., 2017; Açııcı et al., 2021b); *Brueelia biocellata*, *Menacanthus eurysternus*, and *Myrsidea picae* on Eurasian magpie *Pica pica* (Dik et al., 2011c; 2013a; Girişgin et al., 2022); *Philoaterus reguli* and *Ricinus frenatus* on *Regulus regulus* (Dik et al., 2017); *Myrsidea* sp. on sand martin *Riparia riparia* (Dik et al., 2017); *Menacanthus curuccae* and *Menacanthus* sp. on Eurasian blackcap *Sylvia atricapilla* (Dik et al., 2015; 2017); *Menacanthus curuccae*, *Penenirmus affectator*, and *Myrsidea* sp. on garden warbler *Sylvia borin* (Açııcı et al., 2011; Dik et al., 2015; 2017; Açııcı et al., 2021b); *Menacanthus curuccae* and *Myrsidea* sp. on common whitethroat *Curruca communis* (Dik et al., 2015; 2017; Açııcı et al., 2021b); *Menacanthus* sp. and *Penenirmus* sp. on Sardinian warbler *Curruca melanocephala* (Açııcı et al., 2011; Dik et al., 2015; 2017); *Penenirmus pikulai*, *Menacanthus curuccae* and *Myrsidea* sp. on barred warbler *Curruca nisoria* (Dik et al., 2015; 2017; Açııcı et al., 2021b); *Brueelia nebulosa*, *Myrsidea cucullaris*, *Sturnidoecus sturni*

and *Brueelia* sp. on common starling *Sturnus vulgaris* (Dik et al., 2009; 2013b); *Brueelia iliacy* on redwing *Turdus iliacus* (Dik et al., 2015); *Brueelia jacobi*, *Brueelia merulensis*, *Menacanthus eurysternus*, *Menacanthus* sp., and *Ricinus elongatus* on common blackbird *Turdus merula* (Açııcı et al., 2011; Dik et al., 2011c; Dik and Dinçer, 2012; Dik et al., 2017; Açııcı et al., 2021b; Girişgin et al., 2022); *Brueelia turdinulae*, *B. marginata*, and *Menacanthus eurysternus* on song thrush *Turdus philomelos* (Açııcı et al., 2011; Dik et al., 2015; 2017; Açııcı et al., 2021b) were reported in Turkey. The infestation of *Menacanthus* sp. on red-backed shrike *Lanius collurio*; *Brueelia merulensis* and *Menacanthus eurysternus* on common blackbird *Turdus merula*; *Menacanthus agilis* on *Phylloscopus collybita*; *Menacanthus currucae* on Eurasian blackcap *Sylvia*

atricapilla; *Menacanthus gonophaeus*, and *Philopterus atratus* on rook *Corvus frugilegus* were reported on this study. *Philopterus atratus* are reported as the first records, and also *Menacanthus gonophaeus* is a new host for *Corvus frugilegus* for chewing lice fauna of Turkey.

Anatoecus cygni, *Colpocephalum apivorus*, *C. napiforme*, *Ciconiphilus cygni*, *Kurodaia subpachygaster*, *Ornithobius cygni*, and *Philopterus atratus* identified in this study are new records for the fauna of Turkey. Although Turkey has a rich ornithofauna, approximately 1/4 of the bird species have been studied for chewing lice. With each ornitho-parasitological study, new records are being reported for the Phthiraptera fauna of Turkey and host-parasite relationships are being revealed.

References

- Açııcı M, Adam C, Gürler A, Erciyas K, Bölükbaş C et al. (2011). Chewing Lice (Phthiraptera: Amblycera, Ischnocera) From Some Wild Birds in the Kızılırmak Delta (Turkey). *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 54 (2): 395-407. doi: 10.2478/v10191-011-0025-z
- Açııcı M, Eren G, Özkoç ÖÜ, Öztürk M, Bölükbaş CS (2021a). Occurrence of *Neophilopterus tricolor* (Burmeister, 1838) on a black stork (*Ciconia nigra*) in the Kızılırmak Delta. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* 68 (1): 177-180. doi: 10.33988/auvfd.720717
- Açııcı M, Erciyas Yavuz K, Gürler AT, Eren G, Tuygun T et al. (2021b). Kızılırmak deltası kuşlarının bitleri üzerine çalışmalar: Türkiye bit faunası için yeni bildirimler. In: *Proceedings of XXII. Parazitoloji Kongresi 2021; Didim, Aydın, Turkey*. p. 295. (in Turkish).
- Adams RJ, Price RD, Clayton DH (2005). Taxonomic revision of Old World members of the feather louse genus *Columbicola* (Phthiraptera: Ischnocera), including descriptions of eight new species. *Journal of Natural History*, 39 (41): 3545-3618. doi: 10.1080/00222930500393368
- Adam C, Daróczy SJ (2006). The chewing lice (Phthiraptera: Amblycera, Ischnocera) collected on some Falconiformes and Strigiformes (Aves) from Romania. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 49: 145-168.
- Adang KL, Oniye SJ, Ezealor AU, Abdu PA, Ajanusi OJ (2008). Ectoparasites of Domestic Pigeon (*Columba livia domestica*, Linnaeus) in Zaria, Nigeria. *Research Journal of Parasitology* 3: 79-84. doi: 10.3923/jp.2008.79.84
- Aksın N (2003). The presence of Mallophaga species on wild partridge in the Elazığ District. *Turkish Journal of Veterinary and Animal Sciences* 27: 559-565. (in Turkish).
- Aksın N (2004). The presence of lice species on wild geese in the Elazığ District. *Turkish Journal of Veterinary and Animal Sciences* 28: 87-90. (in Turkish).
- Aksın N (2011). Chewing lice (Insecta: Phthiraptera) on Mallards (*Anas platyrhynchos*) in Turkey. *Journal of Animal and Veterinary Advances* 10: 1656-1659. doi: 10.3923/javaa.2011.1656.1659
- Aldemir OS (2004). Ectoparasites in chickens in the province of Kars. *Türkiye Parazitoloji Dergisi* 28 (3): 154-157. (in Turkish).
- Arnold DC (2005). Review of the genus *Ornithobius* (Phthiraptera: Ischnocera: Philopteridae), with descriptions of two new species. *Journal of the Kansas Entomological Society* 78 (2): 158-166.
- Castro DDC, Cicchino C (1983). Contributions to the Knowledge of the Argentine Mallophaga XIII. On Two New Species of the Genus *Incidifrons* Ewing, 1929 (Mallophaga Philopteridae) Parasitic on the Genus *Fulica* (Aves-Gruiformes) In Argentina. *Revista de la Sociedad Entomológica Argentina* 42 (1-4): 273-281.
- Clay T, Hopkins GHE (1950). The early literature on Mallophaga. Part I. 1758-62. *Bulletin of the British Museum (Natural History) Entomology*. London, England.
- Clay T (1957). The *Degeeriella* (Insecta: Mallophaga) parasitic on *Pemis* (Aves: Falconiformes). *Proceedings of the Zoological Society of Calcutta Mookerjee Memorial Volume*: 339-347.
- Clayton DH (1991). The influence of parasites on host sexual selection. *Parasitology Today* 7 (12): 329-334.
- Clayton DH, Price RD (1984). Taxonomy of the *Strigiphilus cursitans* group (Ischnocera: Philopteridae), parasites of owls (Strigiformes). *Annals of the Entomological Society of America* 77 (4): 340-363. doi: 10.1093/aesa/77.4.340
- Clayton DH, Al-Tamimi S, Johnson KP (2003). The ecological basis of coevolution history. In: Page RDM (editor). *Tangled Trees: Phylogeny, Cospeciation and Coevolution*. 1st ed. Chicago, USA: University of Chicago Press, pp. 310-341.

- Clayton DH, Adams RJ, Bush SH (2008). Phthiraptera, the Chewing lice. In: Atkinson CT, Thomas NJ, Bruce Hunter D (editors): Parasitic Diseases of Wild Birds. 1st ed. New Iowa, USA: Wiley-Blackwell Publication, pp. 513-526. doi: 10.1002/9780813804620.ch29
- Cohen M, Greenwood T, Fowler JA (1991). The louse *Trinoton anserium* (Amblycera: Phthiraptera), an intermediate host of *Sarconema eurycerca* (Filarioidea: Nematoda), a heartworm of swans. Medical and Veterinary Entomology 5: 101-110. doi: 10.1111/j.1365-2915.1991.tb00527.x.
- Collinson JM, Parkin DT, Knox AG, Sangster G, Svensson L (2008). Species boundaries in the Herring and Lesser Black-backed Gull complex. British Birds 101 (7): 340.
- Değer MS, Biçek K, Özdal N, Yılmaz AB, Denizhan V et al. (2010). Louse infestations of Domestic Pigeons (*Columba livia domestica*) in Van area Yüzüncü Yıl Üniversitesi Veteriner Fakültesi Dergisi 21 (1): 69-70. (in Turkish).
- Dik B, Yaman M, Köse M, Gülbahçe S (1999). Determination of Mallophaga species on poultry in Konya. Türkiye Parazitoloji Dergisi 23 (3): 327-330. (in Turkish).
- Dik B (2006a). Mallophaga species on Long-legged Buzzards (*Buteo rufinus*): new records from Turkey. Türkiye Parazitoloji Dergisi 30 (3): 226-230.
- Dik B (2006b). Erosive stomatitis in a white pelican (*Pelecanus onocrotalus*) caused by *Piagetiella titan* (Mallophaga: Menoponidae). Journal of Veterinary Medicine 53 (3): 153-154.
- Dik B, Uslu U (2006a). *Cuclotogaster heterographus* (Mallophaga: Lipeuridae) infestation on Ring-Necked Pheasants (*Phasianus colchicus*) in Konya. Türkiye Parazitoloji Dergisi 30 (2): 125-127. (in Turkish).
- Dik B, Uslu U (2006b). Mallophaga (Insecta) species occurring on Storks (*Ciconia ciconia* Linnaeus, 1758). Türkiye Parazitoloji Dergisi 30 (3): 220-225. (in Turkish)..
- Dik B, Özkayhan MA (2007). Mallophaga species on Long-legged Buzzards (*Buteo rufinus*) in Turkey. Türkiye Parazitoloji Dergisi 31 (4): 298.
- Dik B, Uslu U (2007). *Strigiphilus strigis* (Mallophaga: Philopteridae) in a Eurasian Eagle Owl (*Bubo bubo interpositus*) in Turkey. Türkiye Parazitoloji Dergisi 31 (1): 69-71. (in Turkish).
- Dik B, Yamaç E (2008). First report of *Colpocephalum trachelioti* (Amblycera: Menoponidae) on a Black Vulture (*Aegypius monachus* L.) in Turkey. Türkiye Parazitoloji Dergisi 32 (2): 149-152. (in Turkish).
- Dik B, Uslu U (2008). Mallophaga species observed in White Pelicans (*Pelecanus onocrotalus*, Linnaeus) in Turkey. Türkiye Parazitoloji Dergisi 32 (1): 71-76. (in Turkish).
- Dik B (2009). The first report of *Mulcticola hypoleucus* (Denny, 1842) (Phthiraptera: Ischnocera) from Nightjars (*Caprimulgus europaeus* L.) in Turkey. Türkiye Parazitoloji Dergisi 33 (3): 212 - 214. (in Turkish).
- Dik B, Uslu U (2009). Chewing-lice (Phthiraptera: Amblycera, Ischnocera) occurring on birds in the Konya Zoo. Türkiye Parazitoloji Dergisi 33 (1): 43-49. (in Turkish).
- Dik B, Uslu U, Derinbay Ekici Ö, Işık N (2009). Chewing lice (Phthiraptera: Ischnocera, Amblycera) of Starlings (*Sturnus vulgaris*, L.) in Turkey. Türkiye Parazitoloji Dergisi 33 (4): 316-320. (in Turkish).
- Dik B (2010a). New records of chewing lice (Phthiraptera) from some bird species in Turkey. Türkiye Parazitoloji Dergisi 34: 168-73. doi: 10.5152/tpd.2010.06.
- Dik B (2010b). Chewing-lice species (Phthiraptera) found on domestic and wild birds in Turkey. Türkiye Parazitoloji Dergisi 34 (1): 55-60.
- Dik B, Şekercioğlu ÇH, Kıprik MA, İnək S, Uslu U (2010). Chewing lice (Phthiraptera) species found on Turkish shorebirds (Charadriiformes). Kafkas Üniversitesi Veteriner Fakültesi Dergisi 16 (5): 867-874. doi:10.9775/kvfd.2010.2971
- Dik B, Yamaç EE, Uslu U (2011a). Chewing lice (Phthiraptera) found on wild birds in Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 17 (5), 787-794.
- Dik B, Kıprik MA, Şekercioğlu Ç, Şaşmaz Y (2011b). Chewing lice (Phthiraptera) found on songbirds (Passeriformes) in Turkey. Türkiye Parazitoloji Dergisi 35 (1): 34-39. doi: 10.5152/tpd.2011.09.
- Dik B, Sekercioğlu CH, Kıprik MA (2011). Chewing lice (Phthiraptera) species found on birds along the Aras River, Iğdır, Eastern Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 17: 567-573.
- Dik B, Dinçer S (2012). Chewing lice species (Phthiraptera: Ischnocera) found on blackbirds (*Turdus merula*): new records from Turkey. Türkiye Parazitoloji Dergisi 36 (1): 23-27. doi: 10.5152/tpd.2012.06
- Dik B, Uslu U (2012). Studies on chewing lice (Phthiraptera) species found on some duck (Anseriformes: Anatidae) species at lake Akşehir, Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 18 (6): 1055-1060. doi: 10.9775/kvfd.2012.7020
- Dik B, Işık N, Ekici ÖD (2013a). Chewing Lice (Phthiraptera) of Magpie (*Pica pica* L.) (Aves: Passeriformes: Corvidae) in Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 19 (3): 439-441. doi: 10.9775/kvfd.2012.7987
- Dik B, Yamaç E, Uslu U (2013b). Studies on chewing lice (Phthiraptera: Amblycera, Ischnocera) species from domestic and wild birds in Turkey. Kafkas Üniversitesi Veteriner Fakültesi Dergisi 19 (4): 553-560. doi: 10.9775/kvfd.2012.8207
- Dik B, Albayrak T, Adanır R, Uslu U (2013c). Chewing lice (Phthiraptera; Ischnocera, Amblycera) species found on some songbirds (Aves: Passeriformes). Kafkas Üniversitesi Veteriner Fakültesi Dergisi 19 (5): 755-760. (in Turkish). doi: 10.9775/kvfd.2013.8740
- Dik B, Per E, Yavuz KE, Yamaç E (2015). Chewing lice (Phthiraptera: Amblycera, Ischnocera) species found on birds in Turkey, with new records and a new host association. Turkish Journal of Zoology 39 (5): 790-798. doi: 10.3906/zoo-1411-45
- Dik B, Hügül F, Ceylan O (2017a). Chewing lice (Phthiraptera: Amblycera, Ischnocera) of some aquatic birds in Konya province, Turkey, new records for Turkish fauna. Ankara Üniversitesi Veteriner Fakültesi Dergisi 64: 307-312. doi: 10.1501/Vetfak_0000002814

- Dik B, Erciyas-Yavuz K, Per E (2017b). Chewing lice (Phthiraptera: Amblycera, Ischnocera) on birds in the Kızılırmak delta, Turkey. *Revue de Médecine Vétérinaire* 167 (1-2): 53-62.
- Dik B (2020). Türkiye'deki Evcil ve Yabani Memelilerde Görülen Bit (Phthiraptera) Türleri. İzmir, Turkey: Türkiye Parazitoloji Derneği pp. 67 (in Turkish).
- Dik B, Kandir EH (2021). Ectoparasites in Some Wild Birds (Aves) in Turkey. *Progress in Nutrition* 23 (2): e2021261. doi: 10.23751/pn.v23iS2.11919
- Döner A, Yaman M (2015). Mallophaga species in the chickens of Mardin Province. *Van Veterinary Journal* 26 (1): 7-12.
- Edwards RL (1965). Revision of the genus *Aquanirmus* (Mallophaga: Philopteridae), parasitic on grebes (Podicipidae). *The Canadian Entomologist* 97 (9): 920-935.
- Eichler W, Mitwirkung U, Vasjukova TT (1981). Die Mallophagengattung *Trinoton*. Mitteilungen aus dem Museum für Naturkunde in Berlin. *Zoologisches Museum und Institut für Spezielle Zoologie (Berlin, Germany)* 57 (1): 23-62. (in German).
- Elmacioğlu S, Yaman M, Zerek A, Akkücüük Ş, Karagöz M et al. (2018). Ectoparasites of domestic pigeons (*Columba livia domestica*) in Antakya region. *Turkish Journal of Veterinary Research* 2 (2): 23-27.
- Eren G, Gürler AT, Topçu EBG, Tuygun T Açıcı M (2021). First recorded *Felicola (Suricatoecus) vulpis* (Phthiraptera: Trichodectidae) infestation in a Red Fox (*Vulpes vulpes* in Turkey).. *Türkiye Parazitoloji Dergisi* 45 (2): 157-159. (in Turkish).
- Esatlıgil MU, Efil İİ, Tüzer E (2012). Two Chewing Lice Species, *Colpocephalum subzerafae* and *Laemobothrion tinnunculi*, on a Kestrel (*Falco tinnunculus*) in Istanbul: Two New Records from Turkey. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 18 (Suppl-A): A241-A244. doi: 10.9775/kvfd.2012.6313
- Furtun ÖL, Erciyas Yavuz K, Karataş A (Editors) (2021). *Trakus Türkiye'nin Kuşları*. 2nd ed. İstanbul, Turkey: Türkiye İşbankası Kültür Yayınları, pp. 414. (in Turkish).
- Garamszegi LZ (2005). Bird song and parasites. *Behavioral Ecology and Sociobiology* 59 (2): 167-180.
- Gallego Berenguer J, Martín Mateo MP, Aguirre Martí JM (1987). Malófagos de rapaces españolas: II. Las especies del género *Craspedorrhynchus* Keler, 1938 parásitas de falconiformes, con descripción de tres especies nuevas. *Eos* 63: 31-66. (in Spanish).
- Gıcık Y (1999). Ectoparasites in wild pigeons of Ankara and vicinity. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 5 (1): 71-74. (in Turkish with an abstract in English).
- Girişgin AO, Dik B, Girişgin O (2013). Chewing lice (Phthiraptera) species of wild birds in northwestern Turkey with a new host record. *International Journal for Parasitology: Parasites and Wildlife* 30 (2): 217-221. doi: 10.1016/j.ijppaw.2013.07.001.
- Girişgin O, Girişgin AO, Çimenlikaya N, Saygin B (2022). A Survey of the Ectoparasites Found on Wild Birds in Northwest Turkey. *Indian Journal of Animal Research* BF-1474: 1-7. doi: 10.18805/IJAR.BF-1474.
- Göz Y, Dik B, Kılınç ÖZ, Yılmaz AB, Aslan L (2015). Chewing lice (Phthiraptera: Amblycera, Ischnocera) on several species of wild birds around the lake Van Basin, Van, eastern Turkey. *Kafkas Üniversitesi Veteriner Fakültesi Dergisi* 21 (3): 333-338. doi : 10.9775/kvfd.2014.12484
- Gülenber A, Tüzer E, Çetinkaya H (2002). A survey on lice infestations of pigeons in Istanbul, Turkey. *İstanbul Üniversitesi Veteriner Fakültesi Dergisi* 28 (1): 231-234.
- Gülenber A, Kaya Ü, Vassen EWAM, Yavuz E (2006). Chewing-lice on long-legged buzzard. *Indian Veterinary Journal* 83: 1238-1239.
- Güralp N, Doğru C (1966). The Distribution of External Parasites of Poultry in Ankara and Its Vicinity, and Their Treatment with Neguvon Ankara Üniversitesi Veteriner Fakültesi Dergisi 13, 299-305. (in Turkish).
- Güralp N, Mayilmayıl A (1971). An outbreak of caecal trichostrongylosis and Mallophaga infestations in pheasants (*Phasianus colchicus*) in the Samsun area of Turkey. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* 18 (2): 271-275. (in Turkish).
- İnci A, Dik B, Kibar M, Yıldırım A, Düzlü Ö (2010). Chewing lice (Phthiraptera) species on wild birds in Cappadocia region, Turkey. *Türkiye Parazitoloji Dergisi* 34 (4): 174-178. doi: 10.5152/tpd.2010.07
- İpek DNS, Şaki CE (2009). Lice (Mallophaga) species on chickens and distribution of these species in Diyarbakır region. *Dicle Üniversitesi Veteriner Fakültesi Dergisi* 2 (5): 64-68. (in Turkish).
- Jahantigh M, Dizaji RE, Teymooori Y (2016). Prevalence of external parasites of pigeon in Zabol, southeast of Iran. *Journal of Parasitic Diseases* 40 (4): 1548-1551. doi: 10.1007/s12639-015-0725-6
- Johnson KP, Clayton DH (2003). The biology, ecology, and evolution of chewing lice. In: Price RD, Henthall RA, Palma RL, Johnson KP, Clayton DH (editors). *The Chewing Lice: World Checklist and Biological Overview*. Champaign, IL, USA: Illinois Natural History Survey Special Publication, pp. 449-476.
- Karataş A, Göçmen B, Karataş A (2008). A new record for Turkish lice fauna: *Dennyus hirundinis* (Linnaeus, 1761) (Mallophaga: Menoponidae). *Türkiye Parazitoloji Dergisi* 32 (1): 77-8.
- Karatepe M, Dik B, Karatepe B (2017). Chewing lice species (Phthiraptera) found on a European Shag (*Phalacrocorax aristotelis*) in Turkey: new records of a genus and two species for the Turkish fauna of Phthiraptera. *Turkish Journal of Zoology* 41 (3): 576-582. doi: 10.3906/zoo-1603-60
- Keskin A, Dik A (2022). First data on the ectoparasites (fleas, ticks, and lice) of the stone marten, *Martes foina* (Erxleben) in Turkey. *Research Square Preprint (Version 1)* doi: 10.21203/rs.3.rs-1680340/v1
- Kılınç ÖO, Biçek K, Özdal N, Bekir O (2013). Chewing lice (Phthiraptera) found on gull (*Larus michahellis*, Naumann 1840) lived around Van Lake. *Yüzüncü Yıl Üniversitesi Veteriner Fakültesi Dergisi* 24 (3): 117-121. (in Turkish).

- Koroğlu E, Şaki CE, Aktaş M, Dumanlı N, Angın M (1999). Distribution of lice in chicken in Elazığ region. *Fırat Üniversitesi Sağlık Bilimleri Dergisi* 13 (2): 57-60. (in Turkish).
- Koroğlu E, Şimşek S (2001). Distribution of ectoparasites of pigeons (*Columba livia*) in Elazığ. *Fırat Üniversitesi Sağlık Bilimleri Dergisi* 15 (1): 195-198. (in Turkish).
- Merdivenci A (1963). Incidence of ectoparasites of pigeons (*Columba livia*) in Istanbul. *Türk Biyoloji Dergisi* 13: 81-86. (in Turkish).
- Merdivenci A (1970). *Türkiye Parazitleri ve Parazitolojik Yayınları*. 1st ed. İstanbul, Turkey: İstanbul Üniversitesi Cerrahpaşa Tıp Fakültesi Yayınları Kurtuluş Matbaası İstanbul, pp. 324. (in Turkish).
- Mimioğlu M (1952). *Türkiye'de tavuklarda Mallophaga'lar (tavuk bitleri) ve en uygun mücadele metotları üzerinde araştırmalar*. 1st ed. Ankara, Turkey. Ankara Üniversitesi Veteriner Fakültesi Yayınları No: 32, Ankara Üniversitesi Basımevi, pp. 60. (in Turkish).
- Mindell DP, Fuchs J, Johnson JA (2018). Phylogeny, taxonomy, and geographic diversity of diurnal raptors: Falconiformes, Accipitriformes, and Cathartiformes. In: Sarasola JH, Grande JM, Negro JJ (editors). *Birds of Prey: Biology and Conservation in the XXI century*. Berlin, Germany: Springer, pp. 3-32.
- Moreno-Rueda G, Hoi H (2012). Female house sparrows prefer big males with a large white wing bar and fewer feather holes caused by chewing lice. *Behavioral Ecology* 23 (2): 271-277. doi: 10.1093/beheco/arr182
- Naz S, Rizvi SA, Sychra O (2010). *Anaticola crassicornis* (Phthiraptera: Ischnocera: Philopteridae) on wild geese from Pakistan. *Zootaxa* 2659 (1): 60-66. doi: 10.11646/zootaxa.2659.1.3
- Nelson RC, Price RD (1965). The *Laemobothrion* (Mallophaga: Laemobothriidae) of the Falconiformes. *Journal of Medical Entomology* 2 (3): 249-257.
- Oğuz B, Kılınç ÖO, Değer MS (2015). First Reports of *Sarconema eurycerca* and *Trinoton anserinum* in The Whooper Swan (*Cygnus cygnus*) in Van, Turkey, Kafkas Üniversitesi Veteriner Fakültesi Dergisi 21 (6): 933-936. doi: 10.9775/kvfd.2015.13682
- Oruç Ö, Biçer K (2009). Determination of parasite fauna of chicken in the Van region. *Türkiye Parazitoloji Dergisi* 33 (2), 162-164. (in Turkish).
- Palma RL (1978). Slide-mounting of lice: a detailed description of the Canada balsam technique. *New Zealand Entomologist* 6 (4): 432-436. doi: 10.1080/00779962.1978.9722313
- Price RD, Beer JR (1963a). The *Kurodaia* (Mallophaga: Menoponidae) parasitic on the Strigiformes, with a key to the species of the genus. *Annals of the Entomological Society of America* 56 (6): 849-857.
- Price RD, Beer JR (1963b). Species of *Colpocephalum* (Mallophaga: Menoponidae) parasitic upon the Falconiformes. *The Canadian Entomologist* 95 (7): 731-763.
- Price RD, Beer JR (1965). A review of *Ciconiphilus bedford* (Mallophaga: Menoponidae). *The Canadian Entomologist* 97 (6): 657-666.
- Price RD (1974). A review of the genus *Pseudomenopon* (Mallophaga: Menoponidae). *Annals of the Entomological Society of America* 67 (1): 73-84.
- Price RD, Hellenthal RA (1998). Taxonomy of *Philopterus* (Phthiraptera: Philopteridae) from the Corvidae (Passeriformes), with descriptions of nine new species. *Annals of the Entomological Society of America* 91 (6): 782-799. doi: 10.1093/aesa/91.6.782
- Price RD, Hellenthal RA, Palma RL, Johnson KP, Clayton DH (2003). *The Chewing Lice: World Checklist and Biological Overview*. Illinois, USA: Illinois Natural History Survey Special Publication, pp. 501. doi: 10.1080/10635150490468521
- Soler-Cruz MD, Guevara-Benítez DC, Alcántara-Ibáñez F (1985). *Brueelia merulensis* (Denny, 1842), *Brueelia marginata* (Burmeister, 1838) y *Menacanthus polonicus* Eichler y Zlotorzycza, 1963 (Mallophaga, Insecta): estudio morfológico y biométrico. *Revista iberica de parasitología* (1941), 45 (2): 107-117 (in Spanish).
- Şenlik B, Güleğen E, Akyol V (2005). Ectoparasites of Domestic Pigeons (*Columba livia domestica*) in Bursa province. *Türkiye Parazitoloji Dergisi* 29 (2): 100-102. (in Turkish).
- Tigin Y (1973). Ectoparasites of Domestic Pigeons (*Columba livia*) in Turkey. *Ankara Üniversitesi Veteriner Fakültesi Dergisi* 20 (2-3): 372-390. (in Turkish).
- Timmermann G (1952). XXIII.—The species of the genus *Quadriceps* (Mallophaga) from the Larinae. With some remarks on the systematics and the phylogeny of the gulls. *Annals and Magazine of Natural History* 5 (51): 209-222.
- Van Duivendijk N (2011). *Advanced Bird ID Handbook*. 2nd ed. New Holland, United Kingdom: Bloomsbury Publishing PLC, pp. 308.