

ADA033130

卷之三

THE LARGEST PUBLISHER

三

**INFORMAL
MANUSCRIPT
REPORT**

(14) NO⁰-IM-0-45-62

TITLE

LITERATURE SURVEY, BIOLOGY IN THE TONGUE OF THE OCEAN AND EXUMA SOUND

(10)
Good

AUTHOR

ANTHONY J. KRAL, LT. USN


DATE

11 JUN 1962
12 49 p.

**COPY AVAILABLE TO DDC BUREAU
PERMIT FULLY LEGIBLE PRODUCTION**

This manuscript has a limited distribution, therefore in citing it in a bibliography, the reference should be followed by the phrase UNPUBLISHED MANUSCRIPT.

Copy available to DDC does not



~~THE~~ MOST Project

PREFACE

→ The purpose of this paper was to attempt to digest all the available information in the literature on the marine biology of the Tongue of the Ocean and Exuma Sound, and to summarize present knowledge of the oceanography and hydrography of these areas. This paper also points out holidays in the biological information available in these areas. This paper should prove helpful in assessing the role of marine biology in the overall oceanographic environment.

A

ACCESSION NO.	
8708	WHITE SAILOR
REC	DATA SHEET
MANUFACTURER	
JUSTIFICATION	
Polaris on 36	
BY	
DISTRIBUTION/AVAILABILITY	
MAIL	AVAIL. MM/W
A	

TABLE OF CONTENTS

	Page
INTRODUCTION	1
AREA DESCRIPTION	1
General	1
Bathymetry	3
Geology	3
Oceanography	4
Description of Life	7
PLANT LIFE	9
Algae	9
Sea Grasses	10
Sponges	10
Coelenterates	10
Annelids and Sipunculoids	11
Echinoderms	11
Mollusks	11
Arthropods	12
Cartilaginous Fishes	12
Bony Fishes	13
Reptiles	13
Mammals	13
SOUND PRODUCERS	14
Sound Attenuators	14
False Targets	15
Arthroods	15
Fishes	16
Fouling	16
DANGEROUS ORGANISMS	18
CONCLUSIONS	19
BIBLIOGRAPHY	20

FIGURES

1. Tongue of the Ocean and Exuma Sound	2
--	---

APPENDIXES

1. Organisms in the Tongue of the Ocean (TOTO)	26
2. Possible Organisms in the Tongue of the Ocean (TOTO) and Exuma Sound	28

LITERATURE SURVEY - BIOLOGY IN THE TONGUE
OF THE OCEAN AND EXUMA SOUND


INTRODUCTION

Recently there has been increased emphasis in the study of the oceanography of the Tongue of the Ocean (TOTO) and Exuma Sound areas of the Bahamas. Emphasis has been placed primarily on the study of the physical and geological oceanography in these areas; however, some interest has also been shown in biological oceanography. Unfortunately, the marine life of the TOTO and Exuma Sound has not been well studied. The bulk of biological information currently available for the Bahamas is primarily the result of studies conducted in the Bimini's. Since environmental similarities exist from region to region in the Bahamas it can be inferred that biological similarities may also exist. The following study was conducted to assist in the present and future evaluation of the biological contribution to the overall oceanic environment of the TOTO and Exuma Sound.

AREA DESCRIPTION

General

The TOTO and Exuma Sound are two deep channels on the Great Bahama Bank (Fig. 1). The TOTO is a long channel with a narrow northerly section and a nearly circular cul-de-sac southern extremity. The TOTO is about 105 miles long with a minimum width of about 20 miles in the northern section and a maximum width of about 40 miles in the southern cul-de-sac. Exuma Sound is a broad channel with its northern boundary terminating



in a shallow flat bank of about 6 fathoms depth. Exuma Sound is about 108 miles long with an average width of about 33 miles. The channel of the TOTO extends NNE with the northern mouth open to the deep sea. The full length of the TOTO and Exuma Sound are separated by a shallow flat bank of 4 to 6 fathoms depth which is about 40 miles wide at its widest point and 20 miles wide at its narrowest point.

Bathymetry

The bathymetry of the TOTO has been studied by Armstrong (1953) and Miami University (1958). In general the walls of the TOTO and Exuma Sound are quite steep. According to Heezen *et al.* (1959) slope gradients are in the order of 1:4 to 1:8. Busby (1962) has speculated as to the geological origin of the TOTO in reference to the present bathymetry in the area.

Geology

A limited study of the geology of the TOTO has made, but little is known about Exuma Sound. Sediment studies in the TOTO have been conducted by Armstrong (1953), Miami University (1958), USNHO (1961), and Busby (personal communication). In general, the sediments examined range in size from silty sand to silty clay. The highest sediment frequency is in the clay silt range. Limited chemical analyses (Busby, 1962) indicate that the dominant sediment is calcium carbonate, with analyzed samples from the TOTO ranging from 89 to 98 percent CaCO_3 .

Oceanography

Oceanographic data from the TOTO have been reported by Armstrong (1953), Miami University (1958), Magnitzky and French (1960), and Ridley (1962). Results of other investigations in the Bahamas have been reported by Smith (1940) at Andros Island and Berry Island and by Turekian (1957) at Bimini.

Temperature in the upper 100 meters of the TOTO varies between 25°C in the winter to above 30°C in the summer. The temperature is about 22°C at 200 meters, about 15°C at 500 meters, and about 5°C at 1,000 meters. A sharp thermocline is found at a depth of about 50 meters. Wind mixing during the winter results in a deep mixed layer which can extend to about 130 meters.

Salinities in the TOTO range from 34.9 to 37.2‰ (parts per thousand) with the salinity maximum at about 100 meters. Surface salinities in the TOTO were found by Miami University (1958) to increase from 34.2 to 34.3‰ proceeding from north to south. Seasonally, marked changes in the salinity in the upper 125 to 150 meters occurs because of the effects of deep wind mixing in the winter and high evaporation rates in the summer. Salinities are approximately 36.5‰ at 250 meters, 36.0‰ at 500 meters, and 35.0‰ between 1,000 meters and the bottom. Isolated cells of high salinity water from the surface to 200 meters have been observed by Miami University (1958). These cells may be the result of the intrusion of highly saline water from the shallow bank areas. These waters are rapidly cooled and consequently high in density. Smith (1940) observed that extremely high saline

waters were formed on the shallow western shore of Andros Island because of the excess of evaporation as compared to precipitation in this geographical region. Extensive seasonal shallow water oceanographic measurements which could corroborate this hypothesis are not available from the TOTO and Exuma Sound.

Oxygen data from the TOTO have been reported by Smith (1958). In the winter, oxygen values ranged from 3.06 to 4.48 ml/l, while in the summer oxygen values ranged from 3.31 to 5.20 ml/l. Oxygen values reach their minimum at about 500 to 600 meters and reach their maximum at about 2,000 meters. In the summer there is a slight increase in oxygen from the surface to 50 meters. This increase is probably due to the increased biological activity by phytoplankton. Below 50 meters the oxygen values decrease to the depths of the oxygen minimum. Although no conclusive evidence has been published, it is quite probable that since there is an intrusion of water from the banks, lenses of high oxygen content water and lenses of low oxygen content water should exist in both the TOTO and Exuma Sound.

Phosphate data in the TOTO have been published by Miami University (1958). The $\text{PO}_4^{3-}\text{-P}$ values ranged from 0.01 to 1.43 ug at/l in the winter and 0.04 to 1.61 ug at/l in the summer. The $\text{PO}_4^{3-}\text{-P}$ minimum was found at a depth of about 50 meters and the maximum at a depth of 800 meters. In the winter, owing to wind mixing, the $\text{PO}_4^{3-}\text{-P}$ content is greater from the surface to the wind-mixed layer. In the summer $\text{PO}_4^{3-}\text{-P}$ content is less from the surface to the wind-mixed layer. In the summer $\text{PO}_4^{3-}\text{-P}$ levels decrease markedly in the region of greatest biological activity.

Some current measurements have been conducted in the TOTC. The results of these studies are reported by Armstrong (1953), Miami University (1958), Magnitzky and French (1960), and USNHO (1961). Currents in the TOTC and Exuma Sound are generated by winds, tides, and by variations in density distribution. On the basis of analysis of density distribution, relative to the 300-meter level, Magnitzky and French (1960) found that geostrophic currents flowed southward in the eastern part of the TOTC and then gradually curved westward and turned northward in the western part of the TOTC. In all probability, these currents are not too significant since maximum velocities observed below 1,000 meters are about 0.2 knot.

A study of the winds in the TOTC and Exuma Sound prepared by DeLeonibus (1961) shows a high percentage of the winds in a northeasterly to easterly direction. Current measurements between the surface and 200 meters indicated the presence of currents ranging from 0.2 to 1.0 knot with an average velocity of 0.5 knot. Recent rogue measurements by USNHO (1961) showed that current direction and velocity did not coincide with tidal changes. In all probability, wind-induced currents are of primary importance in both the TOTC and Exuma Sound.

Accurate tide data from the TOTC and Exuma Sound are lacking. Present approximations, from USCGS (1961), are that astronomic tides in these areas are semidiurnal with a mean tide level of 2.0 feet (MHWL is 3.3 feet and MLWL is 0.7 feet). Tidal generated currents have been estimated by Smith (1958) to be of the order of 0.2 to 0.5 knot.

Meteorological tide and storm surge data are lacking from the TOTO and Exuma Sound. Studies by Redfield and Miller (1955) have shown that meteorological tides and storm surges, which occur in conjunction with hurricanes and tropical storms, can increase water height 9 feet greater than astronomic tides. In view of the incidence of hurricanes and tropical storms in the vicinity of the TOTO and Exuma Sound, the currents generated by meteorological tides and storm surges are probably quite important.

Submarine daylight measurements, conducted in three regions of the TOTO, have been reported by Miami University (1958). These measurements were made in August 1958 using a submarine photometer with a photoelectric cell (selenium rectifier). From these measurements, it was found that the transparency of the waters in the TOTO is extremely high. Incident light was reduced to 50 percent in the upper 5 to 10 meters and was reduced to 1 percent at depths of 80 to 90 meters. Miami University was able to conclude that the transparency in the TOTO is similar to that of the Sargasso Sea, which means that the waters in the TOTO are among the clearest of oceanic waters.

Description of Life

Limited biological data in the TOTO have been published by Armstrong (1953) and Miami University (1958). The USNHO has conducted some biological sampling during the period of August through October 1961. This sampling was limited to short-period (15 to 20 minute) surface horizontal hauls with a #10-mesh half meter net. The results of analysis of these samples, with

respect to the different species collected, are shown in Appendix 1. (Analyses were conducted by James Bruce, Marine Surveys Division, USNHO, Washington 25, D. C.) Since this sampling was limited geographically and seasonally, Appendix 1 is undoubtedly a very incomplete representation of the marine flora and fauna of the TOTO and Exuma Sound.

Since there has been no complete study of the marine biology of the TOTO and Exuma Sound, it was considered advisable to prepare a detailed list, Appendix 2 of some of the marine plants and animals which might be found in these two areas. This tabulation should assist in the taxonomic analysis of future biological samples collected in the TOTO and Exuma Sound. The method employed in preparing Appendix 2 was to conduct a comprehensive survey of the literature available on the marine biology of the Bahamas. The assumption made was that since there are similarities in the physical environment of the Bahamas from north to south, there is probably a similarity in the species distribution of organisms throughout the Bahamas.

Appendix 2 is by no means a complete listing of all the possible biological organisms which might be found in the TOTO and Exuma Sound. The smaller flora and fauna such as bacteria and nanoplankton have been omitted since future sampling in these areas will probably be limited to one-half meter nets, Clarke-Bumpus sampler hauls (#10- to #20-mesh nets), midwater trawls, and dredging operations. Examination of the reference sources for Appendix 2 will indicate that the majority of the data for this appendix are from studies conducted in the northern region of the Bahamas.

PLANT LIFE

Algae

Numerous species of marine algae have been collected in the Bahamas. Howe (1904) provided some of the first information on the algae in this oceanic area. In addition marine algal studies have been carried out in the vicinity of the Bahamas by Almodovar and Blomquist (1959), Diaz-Pferrer and Lopez (1959), and Voss and Voss (1960). No studies pertaining strictly to the marine algae of the TOTO and Exuma Sound have been reported. In view of the physical environmental data and the availability of nutrients from the shallow water areas, marine algae should be abundant throughout the year.

Hanlon (1957-1961) has discussed the presence of the narrow-leaf Sargasso weed (Sargassum natans) and the broad-leaf Sargasso weed (Sargassum vulgare) in the Bahamas. These brown algae can occur in mats from 1 to 1,000 square feet. Woodcock (1950) has published information on biotic communities associated with submerged rafts of Sargasso weed.

Numerous studies have been conducted on the occurrence of the "red tide" phenomenon associated with blooms of the marine alga Gymnodinium brevis. This alga is found in the Bahamas and "red tides" could occur in the TOTO and Exuma Sound. Other marine algae would have considerable importance with respect to military operations in the Bahamas.

Sea Grasses

The presence of sea grasses in the Bahamas has been discussed by Voss and Voss (1960). These grasses are present near shore and provide a habitat for numerous marine invertebrates. Marine grasses present in a widespread area of the Bahamas are Thalassia testudinum and Cymodocea manatorum.

Sponges

The sponges in the Bahamas have been studied by de Laubenfels (1949), Pearse (1950), and Voss and Voss (1960). These organisms should be present in the inshore areas of the TOTC and Exuma Sound in large numbers. The sponges provide niches for numerous marine invertebrates, and certain sponges such as the Tedania ignis (Fire Sponge) can be dangerous to man.

Coelenterates

The most conspicuous of all the coelenterates found in the Bahamas are the corals. The corals in this area have been studied by Vaughn (1915, 1919a, and 1919b), Newell et al (1959), and Squires (1958). Extensive growths of corals are found off the east coast of Andros Island and in the vicinity of the bank separating the TOTC and Exuma Sound. Coral in these areas is of considerable importance with respect to the presence of other plants and animals. Corals are carnivorous and require a rich food supply. Further, voluminous coral growth is an indication of well oxygenated, freely flowing water in the vicinity of the living part of the reef.

In addition to the corals, a number of hydrozoan and scyphozoan coelenterates are found in these waters. These organisms have been discussed by Newell et al (1959), Voss and Voss (1960), and De Palma

'1961). Cassiopeia sp., the lagoon jellyfish, is found in abundance in sheltered water and should be present in the TOTO and Exuma Sound. Conspicuous among the jellyfish in this area, according to Bolst (1962), is Physalia physalis, the Portuguese-man-of-war, which can be extremely dangerous to man.

Annelids and Sipunculoids

Numerous marine worms are found in the Bahamas. These organisms have been extensively studied by Pearse (1950), Andrew and Andrew (1953), Renaud (1956), and Voss and Voss (1960). These benthic organisms should be found in abundance in the littoral zones of the TOTO and Exuma Sound.

Echinoderms

The echinoderms are represented by numerous species in the Bahamas. These animals have been studied by Ives (1891), Clarke (1942), Deichmann (1957), Newell et al (1959), and Voss and Voss (1960). These organisms are generally littoral and should be abundant in the benthic fauna of the TOTO and Exuma Sound. Included in this group are sea urchins such as Diadema antillarum, Black Urchin, which are poisonous to man.

Mollusks

The Bahamas provide the habitat for numerous mollusks. Primary among these are the polycepods (clams and oysters). The study of the mollusks in the Bahamas has been treated by Pearse (1950), Newell et al (1959), Voss (1960), and Voss and

Voss (1960). In all probability boring mollusks are present in the TOTO and Exuma Sound. These boring forms could cause considerable damage to installations constructed below water.

Arthropods

The marine arthropods in the Bahamas have received only superficial study. These studies have been limited primarily to those arthropods found in the littoral zone. Reports of these arthropods have been published by Pearse (1950), Clarke (1955), Chappuis and Deboutteville (1956), Newall *et al* (1959), and Voss and Voss (1960). There has been no detailed study of the marine arthropods in the neritic zone. Examination of Appendix 1 will indicate that in the oceanic surface samples a large variety of arthropods were encountered. Records at the USNHO indicate that there are numerous sonic arthropods (i.e., snapping shrimp) present in these areas. These sonic arthropods, if present in abundance, may result in high ambient noise levels.

Cartilaginous Fishes

Numerous sharks and rays are found in the waters of the Bahamas. It is quite possible that these animals may be found in considerable numbers in the TOTO and Exuma Sound. Reports of rays and sharks in the Bahamas have been published by La Garce (1919), Gudger (1939), and Bigelow and Schroeder (1948). In view of the numerous reports of shark attacks, these animals are a definite hazard to humans.

Bony Fishes

The bony fishes in the Bahamas have been studied by Fish et al (1952), Woods (1952), Fish (1954), Mowbray (1956), Moulton (1958), Tavelga (1958), Fish and Mowbray (1959), and Krumholz (1959). There probably are a large number of fishes in the TOTO and Exuma Sound throughout the year, especially in the shallow water regions. A large number of the bony fishes from the Bahamas are sonic.

Reptiles

The most important marine reptiles in the Bahamas are the marine turtles. These animals have been discussed by Ingle and Smith (1949) and Hanlon (1957-1961). Several species of these large reptiles are hunted commercially. Andros Island serves as a nesting ground for Thallasochelys kempii, Kemp's turtle.

Mammals

Reports by Kellogg (1929), Bullis and Moore (1956), Caldwell et al (1956), and Moore (1958) indicate that there is the possibility of six different species of whales appearing in the TOTO and Exuma Sound. In all probability these animals would be found as transients. Included in this group is the Finback Whale, Balaenoptera musculus, which, according to Harvey (1959), has been known to measure up to 100 feet long.

Ridley (personal communication) has reported the presence of porpoises in the TOTO. The reaction and behavior of these animals to sound has been reported by Kellogg and Kohler (1952).

SOUND PRODUCERS

Marine organisms capable of producing sounds use these sounds for a variety of purposes. Numerous studies have been conducted to show that sound can be used as a means of echo location and orientation [Griffin (1956), McBride (1956), and Kellogg (1953, 1956, and 1959)]. Organisms in the TOTO and Exuma Sound may use their sonic capability for this purpose.

The contribution of marine animal noises to the total ambient noise in deep water in the TOTO and Exuma Sound will probably not be too great. However, in shallow water areas (0 to 30 feet) this noise contribution could be significant. Unfortunately, marine organisms are notoriously curious and underwater devices (i.e., sensitive hydrophones and transducers) could attract sound producers. Further, any underwater device will provide additional surface area for the growth of organisms which might act as the food for some producers. Consequently, the possible incidence of attachment of these organisms could be increased.

Sound Attenuators

Studies by Hansen and Barnum (1962) have shown that large plankton blooms, such as occur during "red tides," can cause considerable extinction of sea life in surface water layers. Further, large populations of algae can cause the introduction of a considerable amount of oxygen into the water in the proximity of these algae. The level of oxygen introduced can

result in saturation in the upper levels of the water column. Carsola and Kelly (1960) and Ramsey (.962) have found that oxygen saturation can have a serious effect on sound transmission in the region of saturation.

False Targets

Since there are numerous large fishes, sea turtles, and possibly whales in the vicinity of the TOTO and Exuma Sound it is quite probable that these organisms will act as false targets during sonar operations. Further, subsurface rafts of Sargasso weed can also act as false targets. With some practical experience, operators of sonar equipments will probably be able to differentiate between the real targets and "false targets" in these areas.

Arthropods

Studies by Johnson et al (1947), off the coast of southern California, led to the conclusion that major contributors of the biological noise encountered were shrimp of the genera Crangon and Synalpheus. These shallow water shrimp should be found in the TOTO and Exuma Sound (USNHO Snapping Shrimps Cards, 1962). In all probability the area in the vicinity of the shrimp beds should be noisy. The strongest components of shrimp noise are in the frequency range from 2 to 20 kc, and the sound levels over snapping shrimp beds may be 30 db higher than sea state 1 noise. A study by Everest et al (1948) lists the spectra of sound emitted by shrimp from numerous areas.

Moulton (1957 and 1958) has studied the sound production of the lobster, Panulirus argus, found in the Bahamas. This organism produces a rasping sound which is a single burst lasting for about 0.1 second. The frequency of the sound is from 0.04 to 9.00 kc with the greatest intensity at 8.00 kc. In addition to the rasping sound this organism can produce rattling noises. These rattling noises are made by lobsters even when not being disturbed.

Fishes

The study of fish noise in the Bahamas has received considerable attention by Fish et al (1952), Fish (1954), Moulton (1958), Tavolga (1958), and Fish and Mowbray (1959). The fishes studied produced sounds in a wide spectrum of frequencies and for varying intervals of time. Fish noise can be significant when fishes are in schools. According to Mowbray (1956) gamefishes in the Bahamas are not found in large schools, but this does not preclude the possibility of seasonal schooling for migration or mating purposes.

Fouling

A deep-sea fouling pretest was conducted in the TOTO by the USNHO from 17 June to 22 August 1961 (De Palma, 1962). Test panels were placed from the surface to about 970 meters. Fouling organisms attached in decreasing amounts to a depth of approximately 170 meters. Severe fouling was noted from the

surface to 12 meters with green algae, diatoms, and foraminifera being the most abundant fouling organisms. Other fouling organisms were various hydroids, parasitic copepods (Lepeophthirus sp.), and barnacles (Balanus sp., Conchoderma virgatum, and Lepas anatifera). De Palma (1962) noted that fouling on piers and pilings at Andros Island and New Providence Island was slight to moderate in the tidal zone. This fouling was dominated by balanoid barnacles.

On the basis of the numbers of encrusting and boring organisms in the shallow waters of the Bahamas it would seem that fouling could become a serious problem. However, fouling in an area is not necessarily limited to shallow water as Murray (1895) has shown in his summary of "Challenger" bottom studies.

A unique biological feature in a highly productive area is the lack of adequate space for the settling of the larvae of benthic organisms. Man-made intrusions into the benthic environment provide additional settling sites. These intrusions, to mention a few, may be pilings, platforms, components of acoustical systems, ships, and mooring cables. This is particularly true if these intrusions occur in a region of sufficient illumination to permit growth of holotrophic organisms. Plant fouling organisms require a sufficient food supply in the surrounding medium to remain alive. In many instances the fouling organisms are able to use the materials which they are attached to as a source of raw materials for food production.

On the basis of the following criteria:

- (1) Clear water in near shore areas and
- (2) Abundant coral growth, indicating a good supply of water borne foodstuffs,

it is estimated that fouling in the shallow water areas of the TOTO and Exuma Sound will be significant.

DANGEROUS ORGANISMS

Numerous biological organisms (sponges, coelenterates, echinoderms, annelids, and fishes) which are present in the Bahamas could be injurious to man. This is particularly true of personnel whose duties will require them to spend any time in the water (i.e., divers and nearshore construction workers). Contact with poisonous marine organisms could result in death, but it is more likely to result in painful inflammation of the site of contact. Owre (1956) and Lane and Dodge (1958) have studied the effects of stings by coelenterates.

Several of the gamefish in the Bahamas are poisonous so that the possibility of poisoning from eating these fish (ictyosarcotoxism) could exist if gamefish are improperly prepared.

Dinoflagellates responsible for using the "red tide" phenomenon can cause skin irritations and irritations to the respiratory track. Chew (1956) has suggested a method for prediction of the occurrence of "red tides" and Collier (1958) has reported on the biochemical aspects of the "red tide."

Since there are numerous sharks present in the TOTC and Exuma Sound the possibility of shark attack exists. There is also the danger of attack by skates (rays) and moray eels.

CONCLUSIONS

At present it is impossible to completely assess the biological contribution to the overall oceanic environment of the TOTC and Exuma Sound. The information presented as to the possible occurrence of various organisms is purely speculative. However, it would seem that the biology in this area may be extremely important with respect to naval operations conducted in these areas. It is considered that a comprehensive biological study conducted in the TOTC and Exuma Sound is necessary. This study should include both qualitative and quantitative examinations of the pelagic, planktonic, and benthic organisms in the neritic and oceanic provinces and in the littoral and deep-sea zones. Productivity studies should be conducted in the area. Ambient noise measurements should be made, with special emphasis on the biological noise in the region of shrimp beds. Fouling studies should be increased with some emphasis placed on the study of fouling in the neritic province. Dangerous marine plants and animals should be thoroughly studied so that proper assessment of possible danger from these organisms can be made. Behavior of specific organisms (fish schools or clusters of benthic organisms) should be observed in order to determine the location of regions of excessive fouling or abnormal animal activity. When a comprehensive study is completed it may be determined that some areas in the TOTC and Exuma Sound are unfit for specific naval operations because of the biological aspects of these regions.

BIBLIOGRAPHY

- Almodovar, L.R. and H. L. Blomquist
1959. The benthic algae of Bahia Fosforecente, Puerto Rico.
Quart. J. Fla. Acad. Sci. 2(3): 163-168.
- Andrew, Warren and Nancy V. Andrew
1953. Some annelid and sipunculoid worms of the Bimini region.
Amer. Mus. Novitates, No. 1617: 1-16
- Armstrong, John C.
1953. Oceanography in the Tongue of the Ocean, Bahamas, BWI.
Am. Mus. Nat. Hist., Dept. Fishes and Aquatic Biology, N.Y., 38 pp.
- Bigelow, H.B. and Schroeder, W. C.
1948. Fishes of the western north Atlantic. Part . Lancelets,
cyclostomes and sharks. Chapt. 3-Sharks. New Haven, Conn.,
Sears Found. for Mar. Res., Yale Univ. Press, pp. 59-546.
- Bullis, H.R. and J.C. Moore
1956. The occurrences of false killer whales and a summary of
American records. Amer. Mus. Novitates, No. 1756: 1-5
- Busby, Roswell F.
1962. Evidence of subaerial exposure on the floor of the Tongue
of the Ocean, Bahamas. Unpublished Manuscript, USNHO, Washington 25,
D.C., 10 pp.
- Caldwell, David K.
1951. Evidence of home range of an Atlantic bottlenose dolphin.
J. Mammal 36(3): 467-470.
- Caldwell, David K., James N. Layne and J.B. Siebenaler
1956. Notes on a killer whale (*Orcinus orca*) from northeastern
Gulf of Mexico. Quart. J. Fla. Acad. Sci. 19(4): 189-196.
- Carsola, A. J. and N. C. Kelly
1960. Acoustic transmission in oxygen-saturated sea water. Naval
Elect. Laboratory, Tech Rept. No. 955.
- Chappuis, P. A. and C. D. Deboutteville
1956. Etudes sur la faune interstitielle des îles Bahamas recoltes
par Madame Renaud-Debyser. I. Copepodes et Isopodes. Vie et Milieu,
Bull. Lab. Arago. Univ. Paris 7(3): 373-396.
- Chew, Frank
1956. A tentative method for prediction of the Florida red tide
outbreaks. Bull. Mar. Sci. Gulf and Carib. 6(4): 292-304.
- Clark, H. L.
1942. The echinoderm fauna of Bermuda. Bull. Mus. Comp. Zool.
82: 367-394.

- Clarke, W. D.
1955. A new species of the genus *Heteromysis* (Crustacea, Mysidae) from the Bahama Islands, commensal with a sea-anemone. Amer. Mus. Novitates, No. 1751: 1-13
- Collier, Albert
1958. Some biochemical aspects of red tide and related oceanographic problems. Limnol and Oceanogr. 3(1): 33-39.
- de Laubenfels, M.W.
1949. Sponges of the western Bahamas. Amer. Mus. Novitates, No. 1431: 1-25.
- de Leonibus, P. S.
1961. Observed and theoretical significant wave heights-Tongue of the Ocean and Exuma Sound. USNHO Inform. Mem. No. 4-61, Unpublished Manuscript, Washington 25, D.C.
- De Palma, John R.
1962. Results of a deep-sea marine fouling and corrosion pre-test in the Tongue of the Ocean. Inform. Mem. No. 13-62, Unpublished Manuscript, Washington 25, D.C., USNHO.
- Deichmann, E.
1957. The littoral holothurians of the Bahama Islands. Amer. Mus. Novitates, No. 1821: 1-20.
- Diaz-Piferrer, Manuel and Hady Lopez
1959. Taxonomia, ecologia y valor nutrimental de algas marinas cubanas. I. Ser. Estudios sobre Trabajos de Investigacion, Inst. Cubano Investigaciones Tech. Rept. No. 6.
- Everest, F.A., R. W. Young and M. W. Johnson
1948. Acoustical characteristics of noise produced by snapping shrimp. J. Acoust. Soc. Amer. 20(2):137-142
- Fish, M.P., A.S. Kelsey Jr. and W. H. Mowbray
1952. Studies on the production of underwater sound by North Atlantic coastal fishes. J. Mar. Res. 11(2): 180-193
- Fish, Marie F.
1954. The character and significance of the sound production among fishes of the western north Atlantic. Bull. Bingham Oceanogr. Coll. 14(3): 1-109.
- Fish, M.P. and W.H. Mowbray
1959. The production of underwater sound by *Opsanus* sp., a new toadfish from Bimini, Bahamas. Zoologica 44(2): 71-76.
- Griffin, D.
1955. Hearing and acoustic orientation in Marine animals. Pap. Mar. Biol., Deep Sea Res., Suppl. to Vol. 3, pp 406-417

- Wade, W.
1939. A general review of the Bahama Islands. Nature 143(3611): 79-80.
- Manlon, R. V.
1957-1961 Bahamian Nat. Hist. Series. Nassau Daily Tribune.
- Hansen, J. and E.G. Barham
1962. Resonant cavity measurements of the effects of "red water" plankton on the attenuation of under water sound. Limnol and Oceanogr. 7(1): 8-13.
- Hardy, Alister
1950. The Open Sea. The Natural History Library II, Fish and Fisheries. Houghton Mifflin Co., Boston, Mass.
- Henderson, C. M. (Editor)
1957. The Biology of the Bahamas. The Naturalistic Survey of the British West Indies, Vol. 1, Part 1.
- Hewitt, G.
1900. Contributions to marine fauna from the Bahamas. J. N.Y. Bot. Garden 8(1): 1-11.
- Ingle, R.M. and W.C. Smith
1940. Game fishes and sport fishing industry of the West Indies, Florida and Gulf of Mexico with annotated bibliography. Spec. Publ., Marine Lab., Univ. Miami and Carib. Research Committee.
- Ives, J. E.
1891. Echinoderms from the Bahama Islands. Proc. Acad. Nat. Sci. Philadelphia pp. 337-341
- Johnson, M. W., F.A. Everest and R. W. Young
1947. Shrimp noise. Biol Bull. 93(2): 127-138.
- Kellogg, Remington
1920. What is known of the migrations of some whalebone whales. Smithsonian Inst. Ann. Rept., 1928, pp. 467-494.
- Kellogg, W. N. and R. Kohler
1952. Reactions of porpoise to ultrasonic frequencies. Science 116 (3010): 250-252
- Kellogg, W.N., R. Kohler and H. W. Morris
1953. Porpoise sounds as sonar signals. Science 117(3036): 239-243.
- Kellogg, W.N.
1959. Size of discrimination by reflected sound in a bottlenose porpoise. J. comp. physiol. Psychol. 52(5): 509-514.
- Kellogg, W.N.
1959. Auditory perception of submerged objects by porpoises. J. acoust. soc. America. 31(1): 1-6.

- Krumholz, Louis A.
1959. Stomach contents and organ weights of some bluefin tuna,
Thunnus thynnus (L), near Bimini Bahamas. Zoologica 44(3): 127-131
- La Garce, John O.
1919. Devil fishing in the gulf stream. Nat. Geo. Mag. 35(6): 476-488.
- Lane, C. E. and Eleanor Dodge
1958. The toxicity of Physalia nematocysts. Biol. Bull. 115(2): 219-226.
- Magnitzky, A. W. and H.V. French
1960. Tongue of the Ocean. ASWEPs Rept. 3, USNHO, TR-94
- McBride, Arthur F.
1956. Evidence of echolocation by cetaceans. Deep Sea Res. 3(2): 153-154.
- Miami University
1958. Oceanographic survey of the Tongue of the Ocean, Vol. 1, University of Miami Tech. Rept.
- Moore, J. C.
1958. A beaked whale from the Bahamas Islands and comments on the distribution of Mesoplodon densirostris. Amer. Mus. Novitates, No. 1897: 1-12.
- Moulton, J.M.
1957. Sound production in the spiny lobster, Panulirus argus (Latrelle). Biol. Bull. 113(3): 286-295.
- Moulton, J. M.
1958. The acoustical behaviour of some fishes of the Bimini area. Biol. Bull., 114(3): 357-374
- Mowbray, L.S.
1956. Game fishes of Bermuda. International Gamefish Committee.
- Murray, John
1895. Summary of the scientific results obtained at sounding, dredging and trawling stations of H.M.S. Challenger, Repts., Summary, Pt.2, pp 797-1608.
- Newell, N. D., John Imbrie, E.G. Purdy and D.L. Thurber
1959. Organism communities and bottom facies, Great Bahama Bank. Bull. Amer. Mus. Nat. Hist. 117(4): 183-228, pls. 58-69.
- Owre, Harding B.
1956. Coelenterate nematocysts as a cause of certain kinds of sea stings. Bull. Mar. Sci. Gulf and Carib. 6(4): 309-314

- Pearse, S.S.
1950. Notes on the inhabitants of certain sponges at Bimini.
Ecology 31(1): 149-151
- Ramsey, W. L.
1962. Bubble growth from dissolved oxygen near the surface.
Limnol and Oceanogr 7(1): 1-7
- Redfield, A. C. and A. R. Miller
1955. Memorandum on water levels accompanying Atlantic coast hurricanes. Woods Hole Ocean Inst. Ref. No. 55-28, 195 pp.
- Renaud, Jeanne
1956. A report on some polychaetous annelida from the Miami-Bimini area. Amer. Mus. Novitates, No. 1812: 1-40.
- Rivas, Luis R.
1954. A preliminary report on the spawning of the western north Atlantic bluefin tuna (*Thunnus thynnus*) in the Straits of Florida. Bull. Mar. Sci. Gulf and Carib. 4(4): 302-322
- Smith, C. L.
1940. The Great Bahama Bank. I. General hydrographical and chemical features. J. Mar. Res. 3(2): 147-170.
- Squires, Donald F.
1958. Stony corals from the vicinity of Bimini, Bahamas, British West Indies. Bull. Amer. Mus. Nat. Hist. 115(4): 215-262
- Tavolga, W.N.
1958. Underwater sounds produced by two species of toadfish, Opsanus tau and Opsanus beta. Bull. Mar. Sci. Gulf and Carib. 8(3): 278-284.
- Turekian, K.K.
1957. Salinity variations in sea water in the vicinity of Bimini, Bahamas, British West Indies. Amer. Mus. Novitates, No. 1822: 1-12
- USNHO
1961. Current drogue measurements and bottom sediment analyses, preliminary report. Tongue of the Ocean. Unpublished Manuscript. 7 pp.
- USNHO
1962. Snapping shrimp cards. Oceanographic Analysis Division.
- U.S. Coast and Geodetic Survey
1961. Tide tables. east coast U.S.A. and South America.

Vaughn, T.W.

1915. Reef corals of the Bahamas and southern Florida. Carnegie Inst. of Wash., Yearbook No. 13, pp. 222-226.

Vaughn, T. W.

1919a. Fossil corals from central America, Cuba and Puerto Rico with an account of the tertiary, pleistocene and recent coral reefs. Bull. U.S. Nat. Mus. 103: 189-524, pls 68-152.

Vaughn, T.W.

1919b. Corals and formation of coral reefs. Ann. Rept. Smithsonian Inst. for 1917: 189-276, pl 1-37.

Voss, Gilbert L.

1960. Bermudan cephalopods. Fieldiana, Zool. 39(40): 419-446.

Voss, G.L. and N.A. Voss

1960. An ecological survey of the marine invertebrates of Bimini, Bahamas with consideration of their zoogeographical relationships. Bull. Mar. Sci. Gulf and Carib. 10(1): 96-116.

Woodcock, A.H.

1950. Subsurface pelagic Sargassum. J. Mar. Res. 9(2): 77-92.

Woods, L.P.,

1952. Fishes attracted to surface light at night in the Gulf of Mexico. Copeia 1: 40-41.

Personal Communications:

Busby, Roswell F., USNHO, Washington 25, D.C.

Bolet, A., LP, USN, USNHO, Washington 25, D.C.

Ridley, E., USNHO, Washington 25, D.C.

APPENDIX I

Organisms in the Tongue of the Ocean (TOTO)

<u>Organism</u>	<u>Scientific Name</u>
<u>ALGAE</u>	
Blue green algae	
Dinoflagellate	<u>Ceratium</u> sp.
Diatom	<u>Chaetoceros</u> sp.
Diatom	<u>Coscinodiscus</u> sp.
Foraminiferan	<u>Globigerina</u> sp.
Radiolarian	<u>Radiolaria</u> sp.
Brown algae	<u>Sargassum</u> sp.
<u>COELENTERATA</u>	
Scyphozoan medusa	
Cyphonautes larva	
<u>ANNELIDA</u>	
Polychaete larva	
<u>MOLLUSKS</u>	
Gastropod	
Lamellibranch	
Scaphopod	
Pteropod	
<u>CHAETOGNATHS</u>	
Arrow worm	
<u>ECHINODERMS</u>	
Starfish larva	
Echinopluteus larva	

APPENDIX I (Continued)

<u>Organism</u>	<u>Scientific Name</u>
<u>ARTHROPODS</u>	
Copepod	<u>Acartia tonsa</u>
Copepod	<u>Acrocalanus longicornis</u>
Copepod	<u>Calanus sp.</u>
Copepod	<u>Calocalanus pavo</u>
Copepod	<u>Calocalanus pavonina</u>
Copepod	<u>Copilia sp.</u>
Copepod	<u>Corycaeus sp.</u>
Copepod	<u>Corycaeus speciosus</u>
Brachiopod	<u>Evadne sp.</u>
Copepod	<u>Farranula sp.</u>
Copepod	<u>Jubstockia sp.</u>
Copepod	<u>Lucicutia longicornis</u>
Copepod	<u>Macrostella sp.</u>
Decapod	<u>Maia sp.</u>
Copepod	<u>Microstella sp.</u>
Copepod	<u>Oithona sp.</u>
Copepod	<u>Oncaea sp.</u>
Decapod	<u>Parcellanid sp.</u>
Copepod	<u>Sapphirina sp.</u>
Copepod	<u>Temora stylifera</u>
Euphausiid	<u>Thysanoessa sp.</u>
Copepod	<u>Undeuchaeta minor</u>
<u>VERTEBRATES</u>	
Salp	<u>Doliolum sp.</u>
Fish eggs	

APPENDIX B

Possible Organisms in the Tongue of the Ocean (TOTO)
and Exuma Sound

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ALGAE</u>		
<u>Acanthophora spicifera</u>	Red algae	Voss and Voss (1960)
<u>Acetularia sp.</u>	Green algae	Voss and Voss (1960)
<u>Batophora Oerstedi</u>	Green algae	Voss and Voss (1960)
<u>Bostrychia tenella</u>	Red algae	Voss and Voss (1960)
<u>Caulpera paspaloides</u>	Green algae	Newell et al (1959)
<u>Cladophora glaucescens</u>	Green hair	Voss and Voss (1960)
<u>Cladophoropsis membranacea</u>	Green algae	Voss and Voss (1960)
<u>Coralina cubensis</u>	Red algae	Voss and Voss (1960)
<u>Dictyosphaerium cavernosum</u>	Brown algae	Newell et al (1959)
<u>Dictyota indica</u>	Brown algae	Voss and Voss (1960)
<u>Entophysalis deusta</u>	Blue-green algae	Newell et al (1959)
<u>Enteromorpha sp.</u>	Green algae	Voss and Voss (1960)
<u>Galaxaura subverticillata</u>	Red algae	Newell et al (1959)
<u>Goniolithon strictum</u>	Red algae	Newell et al (1959)
<u>Halimeda incrassata</u>	Green algae	Voss and Voss (1960)
<u>Halimeda opuntia</u>	Green algae	Voss and Voss (1960)
<u>Halimeda scabra</u>	Green algae	Voss and Voss (1960)
<u>Halimeda tridens</u>	Green algae	Voss and Voss (1960)
<u>Heterociphonia secunda</u>	Red algae	Voss and Voss (1960)
<u>Jania capillacea</u>	Red algae	Newell et al (1959)
<u>Laurencia intricata</u>	Red algae	Newell et al (1959)
<u>Laurencia portei</u>	Red algae	Voss and Voss (1960)

APPENDIX 2 (continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ALGAE (Continued)</u>		
<u>Lithothamnion sp.</u>	Red algae	Voss and Voss (1960)
<u>Penicillus capitatus</u>	Green algae	Newell <u>et al</u> (1959)
<u>Penicillus dumetosus</u>	Green algae	Newell <u>et al</u> (1959)
<u>Penicillus pyriformis</u>	Green algae	Voss and Voss (1960)
<u>Podina Sanctae-Grucis</u>	Brown algae	Voss and Voss (1960)
<u>Porolithon pachydermum</u>	Red algae	Newell <u>et al</u> (1959)
<u>Rhipocephalus phoenix</u>	Green algae	Newell <u>et al</u> (1959)
<u>Rotalea rossa</u>	Foraminiferan	Newell <u>et al</u> (1959)
<u>Sargassum natans</u>	Narrow-leaf Sargassum weed	Hanlon (1957-1961)
<u>Sargassum polyceratum ovatum</u>	Brown algae	Newell <u>et al</u> (1959)
<u>Sargassum Vulgare</u>	Wide-leaf Sargassum weed	Hanlon (1957-1961)
<u>Turbinaria tubrinata</u>	Brown algae	Newell <u>et al</u> (1959)
<u>Zonaria zonalis</u>	Brown algae	Newell <u>et al</u> (1959)
<u>SEA GRASSES</u>		
<u>Thalassia testudinum</u>	Turtle grass	Newell <u>et al</u> (1959)
<u>Cymodocea Manatorum</u>	Sea grass	Newell <u>et al</u> (1959)
<u>PORIFERA</u>		
<u>Agelas sparsus</u>	Sponge	de Laubenfels (1949)
<u>Anthosigmella varians</u>	Dingy Sponge	de Laubenfels (1949)
<u>Aulospongus schoemus</u>	Branching sponge	Pearse (1950)
<u>Axocella spinosa</u>	Thin sponge	de Laubenfels (1949)
<u>Callyspongia fallax</u>	Tube sponge	de Laubenfels (1949)

APPENDIX 2 (continued)

<u>Scientific Name</u>	<u>*Common Name</u>	<u>Reference Source</u>
<u>PORIFERA (Continued)</u>		
<u>Callyspongia vaginalis</u>	Tube sponge	de Laubenfels (1949)
<u>Chondrilla nucula</u>	Chicken-liver sponge	de Laubenfels (1949)
<u>Ginuchyra cavernosa</u>	Sponge	de Laubenfels (1949)
<u>Cliona vastifica</u>	Boring sponge	Voss and Voss (1960)
<u>Cribrochalina infundibula</u>	Funnel sponge	de Laubenfels (1949)
<u>Cryptotheya cripta</u>	Sponge	de Laubenfels (1949)
<u>Dysidea etheria</u>	Heavenly sponge	de Laubenfels (1949)
<u>Fibulia politangere</u>	Bun sponge	Voss and Voss (1960)
<u>Geodia gibberosa</u>	White sponge	de Laubenfels (1949)
<u>Haliclona longloyi</u>	Finger sponge	de Laubenfels (1949)
<u>Haliclona molibita</u>	Sponge	de Laubenfels (1949)
<u>Haliclona rubens</u>	Red sponge	Pearse (1950)
<u>Haliclona viridis</u>	Green sponge	de Laubenfels (1949)
<u>Higginsia coralloides</u>	Coral sponge	de Laubenfels (1949)
<u>Hircina strobilina</u>	Sponge	Pearse (1950)
<u>Ianthella ianthella</u>	Big sponge	de Laubenfels (1949)
<u>Iotrochata brotulata</u>	Brown sponge	Pearse (1950)
<u>Ircinia fasciculata</u>	Stink sponge	de Laubenfels (1949)
<u>Ircinia strobilina</u>	Loggerhead sponge	de Laubenfels (1949)
<u>Oligoceras hemorrhages</u>	Bleeding sponge	de Laubenfels (1949)
<u>Pseudaxinella rosacea</u>	Rosette sponge	de Laubenfels (1949)
<u>Siphonochalina siphona</u>	Yellow-brown sponge	de Laubenfels (1949)
<u>Spheciospongia vesperia</u>	Manjack sponge	Pearse (1950)
<u>Spirastrella coccinea</u>	Sponge	de Laubenfels (1949)

APPENDIX 2 (continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>PORIFERA (Continued)</u>		
<u>Spongia officinalis</u>	Reef sponge	de Laubenfels (1949)
<u>Tedania ignis</u>	Fire sponge	de Laubenfels (1949)
<u>Verongia fistularis</u>	Candle sponge	de Laubenfels (1949)
<u>Xytopsene sigmatum</u>	Lagoon sponge	de Laubenfels (1949)
<u>COELENTERATA - CORALS</u>		
<u>Acropora cervicornis</u>	Staghorn coral	Newell <u>et al</u> (1959)
<u>Acropora palmata</u>	Outlet coral	Squires (1958)
<u>Agarcia agaricites</u> var. <u>crassa</u>	Purple coral	Squires (1958)
<u>Agarcia agaricites</u> var. <u>purpurea</u>	Purple coral	Squires (1958)
<u>Ichocoenia stokesii</u>	Small stone coral	Vaughn (1915)
<u>Aploria clivosa</u>	Mat coral	Squires (1958)
<u>Diploria strigosa</u>	Boulder coral	Squires (1958)
<u>Eusmilia</u> sp.	Coral	Vaughn (1915)
<u>Favia fragum</u>	Caupiliiform coral	Squires (1958)
<u>Isophyllastrea rigida</u>	Coral	Vaughn (1915)
<u>Isohyllia sinuosa</u>	Coral	Vaughn (1915)
<u>Manicina areolata</u>	Rose coral	Newell <u>et al</u> (1959)
<u>Millepora alcicornis</u>	Coral	Vaughn (1915)
<u>Millepora complanata</u>	Coral	Vaughn (1915)
<u>Montastrea annularis</u>	Coral	Vaughn (1915)
<u>Monastrea cavernosa</u>	Coral	Vaughn (1915)
<u>Mussa</u> aff. <u>dispacea</u>	Coral	Vaughn (1915)
<u>Oculina diffusa</u>	Branching coral	Vaughn (1915)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
COELENTERATA - CORALS (Continued)		
<u>Palythoa sp.</u>	Coral	Newell <u>et al</u> (1959)
<u>Paracyanthus confertus</u>	Coral	Vaughn (1915)
<u>Porites asteroides</u>	Star coral	Vaughn (1915)
<u>Porites porites</u> <u>var. clavaria</u>	Finger coral	Squires (1958)
<u>Porites porites</u> <u>var. divaricata</u>	Finger coral	Squires (1958)
<u>Porites porites</u> <u>var. furcata</u>	Finger coral	Squires (1958)
<u>Siderastrea radians</u>	Spheroid coral	Squires (1958)
<u>S. siderea</u>	Starlet coral	Squires (1958)
<u>Stephanocoenia intersepta</u>	Coral	Vaughn (1915)
<u>Stephanocoenia michelini</u>	Coral	Vaughn (1915)
COELENTERATA - OTHER COELENTERATES		
<u>Aiptasia annulata</u>	Sea anenome	Newell <u>et al</u> (1959)
<u>Bartholomea annulata</u>	Sea anenome	Newell <u>et al</u> (1959)
<u>Cassiopeia sp.</u>	Lagoon jellyfish	Voss & Voss (1960)
<u>Cassiopeia xamachina</u>	Lagoon jellyfish	Voss & Voss (1960)
<u>Condylactis gigantica</u>	Sea anenome	Voss & Voss (1960)
<u>Gorgonia fabellum</u>	Gorgonacean	Newell <u>et al</u> (1959)
<u>Phymianthus crucifer</u>	Sea anenome	Newell <u>et al</u> (1959)
<u>Physalia physalis</u>	Portugese-man-of-war	Bolst (personal communication)
<u>Stoichaeatis helianthus</u>	Sea anenome	Newell <u>et al</u> (1959)
<u>Zoanthus sociatus</u>	Sea anenome	Newell <u>et al</u> (1959)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
ANNELIDA & SIPUNCULOIDEA		
<u>Amphione jamaicensis</u>	Bristle worm	Voss & Voss (1960)
<u>Arenicola cristata</u>	Lug worm	Andrew & Andrew (1953)
<u>Arenicola marina</u>	W. Indian lug worm	Andrew & Andrew (1953)
<u>Armandia agilis</u>	Ophelid worm	Renaud (1956)
<u>Armandia maculata</u>	Sea worm	Pearse (1950)
<u>Aspidosiphon cumingii</u>	Sipunculoid worm	Andrew & Andrew (1953)
<u>Autolytus sp.</u>	Transparent bristle worm	Andrew & Andrew (1953)
<u>Chloea viridis</u>	Red bristle worm	Andrew & Andrew (1953)
<u>Cirriformia filigera</u>	Cirratulid worm	Renaud (1956)
<u>Cirriformia tortugaensis</u>	Sea worm	Pearse (1950)
<u>Dasybranchus sp.</u>	Capitellid worm	Andrew & Andrew (1953)
<u>Dasybranchus tumulatus</u>	Sea worm	Pearse (1950)
<u>Eunice denticulata</u>	Eunicid worm	Renaud (1956)
<u>Eunice filamentosa</u>	Sea worm	Pearse (1950)
<u>Eunice longicirrata</u>	Eunicid worm	Renaud (1956)
<u>Eunice rubra</u>	Red sponge worm	Renaud (1956)
<u>Eunice tibania</u>	Eunicid worm	Andrew & Andrew (1953)
<u>Eupolydentes cornishii</u>	Sea worm	Andrew & Andrew (1953)
<u>Eupolymnia crassicornis</u>	Verbellid worm	Andrew & Andrew (1953)
<u>Eurythoe complanata</u>	Bristle worm	Andrew & Andrew (1953)
<u>Glycera americana</u>	Sargassum worm	Andrew & Andrew (1953)
<u>Glycera tesselata</u>	Pink worm	Renaud (1956)
<u>Haplosyllis spongicola</u>	Bristle worm	Renaud (1956)
<u>Hermodice carunculata</u>	Bristle worm	Andrew & Andrew (1953)
<u>Hysicomis midoculi</u>	Purple feather duster worm	Andrew & Andrew (1953)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ANNELIDA & SIPUNCULOIDA (Continued)</u>		
<u>Lepidontotus variabilis</u>	Sea worm	Renaud (1956)
<u>Lumbrenereis latrelli</u>	Sea worm	Pearse (1950)
<u>Lysidice ninetta</u>	Red worm	Andrew & Andrew (1953)
<u>Marpheya sanguinea</u>	Eunicid worm	Andrew & Andrew (1953)
<u>Mystides elongata</u>	Phylloicid worm	Renaud (1956)
<u>Namanereii ovenaryensis</u>	Nereid worm	Andrew & Andrew (1953)
<u>Nicidion kinbergi</u>	Eunicid worm	Renaud (1956)
<u>Notopygus crinita</u>	Small bristle worm	Andrew & Andrew (1953)
<u>Odontosyllis enoola</u>	Bristle worm	Andrew & Andrew (1953)
<u>Perenereis andersoni</u>	Nereid worm	Andrew & Andrew (1953)
<u>Perenereis floridana</u>	Nereid worm	Andrew & Andrew (1953)
<u>Phascolosoma sp.</u>	Sipunculoid worm	Andrew & Andrew (1953)
<u>Physocosoma microdentigerum</u>	Sipunculoid worm	Andrew & Andrew (1953)
<u>Physoccsoma varians</u>	Sipunculoid worm	Andrew & Andrew (1953)
<u>Platynereis dumarillii</u>	Sargassum worm	Andrew & Andrew (1953)
<u>Polymnia nebulosa</u>	Terebellid worm	Voss & Voss (1960)
<u>Sabella melanostigma</u>	Feather duster worm	Voss & Voss (1960)
<u>Sabellastarte magnifica</u>	Feather duster worm	Renaud (1956)
<u>Scyphonroctus sp.</u>	Capitellid worm	Andrew & Andrew (1953)
<u>Sthenelais setosa</u>	Sea worm	Pearse (1950)
<u>Styliariodes plumosa</u>	Sea worm	Pearse (1950)
<u>Terebella magnifica</u>	Sea worm	Pearse (1950)
<u>Trypanopsyllis sp.</u>	Sea worm	Pearse (1950)
<u>Vanadis crystallina</u>	Red-eyed worm	Renaud (1956)
<u>MOLLUSKS</u>		
<u>Acanthopleura granulata</u>	Chiton	Voss & Voss (1960)

A PENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>MOLLUSKS (Continued)</u>		
<u>Aequipecten gibbus</u>	Pecten	Newell <u>et al</u> (1959)
<u>Anadaca mutabilis</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Asaphis deflorata</u>	- - -	Voss & Voss (1960)
<u>Potillaria minima</u>	- - -	Voss & Voss (1960)
<u>Chiton squamosus</u>	Snake chiton	Pearse (1950)
<u>Chiton viridis</u>	Chiton	Newell <u>et al</u> (1959)
<u>Chlamys benedicti</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Cione cancellata</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Codakia orbicularis</u>	- - -	Voss & Voss (1960)
<u>Cumingia coarata</u>	Clam	Pearse (1950)
<u>Cypraea cinerea</u>	Clam	Pearse (1950)
<u>Diodora listeri</u>	Limpet	Voss & Voss (1960)
<u>Divaricella quadrivalvis</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Echininus nodulosus</u>	Snail	Voss & Voss (1960)
<u>Fasciolaria tulipa</u>	- - -	Voss & Voss (1960)
<u>Fissurella barbadensis</u>	Keyhole limpet	Voss & Voss (1960)
<u>Gastrochaena hirs</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Glycimeris undata</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Gouldia cerina</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Irvitia sp.</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Isognomon alata</u>	Bivalve	Voss & Voss (1960)
<u>Isognomon listeri</u>	Clam	Pearse (1950)
<u>Isognomon radiata</u>	Clam	Newell <u>et al</u> (1959)
<u>Laevicardium laevigatum</u>	Pelycepod	Newell <u>et al</u> (1959)
<u>Limnoria sp.</u>	Wood borer	Voss & Voss (1960)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>MOLLUSKS (Continued)</u>		
<u><i>Linia antillensis</i></u>	Clam	Pearse (1950)
<u><i>Lithophaga nigra</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Littorina meleagris</i></u>	Gastropod	Voss & Voss (1960)
<u><i>Littorina mespilium</i></u>	Brown gastropod	Voss & Voss (1960)
<u><i>Littorina ziczac</i></u>	Periwinkle	Voss & Voss (1960)
<u><i>Livona pica</i></u>	W. Indian top shell	Voss & Voss (1960)
<u><i>Lucina pennsylvanica</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Lucopina sawbubii</i></u>	Limpet	Voss & Voss (1960)
<u><i>Macrocalista maculata</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Musculus coralliphagus</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Musculus lateralis</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Nerita peleronta</i></u>	Gastropod	Voss & Voss (1960)
<u><i>Nerita tessellata</i></u>	Gastropod	Voss & Voss (1960)
<u><i>Nerita versicolor</i></u>	Gastropod	Voss & Voss (1960)
<u><i>Octopus macroous</i></u>	Grass octopus	Voss (1960)
<u><i>Octopus vulgaris</i></u>	Rock octopus	Voss (1960)
<u><i>Onychotestis banksi</i></u>	Squid	Voss (1960)
<u><i>Papyridae sevisculatum</i></u>	Rugose clam	Pearse (1950)
<u><i>Pecten antillarum</i></u>	Pecten	Voss & Voss (1960)
<u><i>Penicardia reliata</i></u>	Pen shell	Voss & Voss (1960)
<u><i>Pincilda radiata</i></u>	Clam	Voss & Voss (1960)
<u><i>Pter simpsoni</i></u>	- - -	Newell <u>et al</u> (1959)
<u><i>Plicatulata gibbosa</i></u>	Polycepod	Newell <u>et al</u> (1959)
<u><i>Polinices duplicata</i></u>	Moon-shell	Voss & Voss (1960)
<u><i>Polinices lacteus</i></u>	Snail	Voss & Voss (1960)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>MOLLUSKS (Continued)</u>		
<u>Postularca odonsi</u>	Snail	Pearse (1950)
<u>Siphonaria alternata</u>	Snail	Voss & Voss (1960)
<u>Spiroglyphus irregularis</u>	Snail	Newell et al (1959)
<u>Strigilla mirabilis</u>	Pelycepod	Newell et al (1959)
<u>Strombus sambæ</u>	Conch	Newell et al (1959)
<u>Strombus gigas</u>	Pink conch	Voss & Voss (1960)
<u>Tectarius muricatus</u>	- - -	Voss & Voss (1960)
<u>Tellina interrupta</u>	Pelycepod	Newell et al (1959)
<u>Tellina radiata</u>	Pelycepod	Newell et al (1959)
<u>Teredo sp.</u>	Wood borer	Voss & Voss (1960)
<u>Thais patula</u>	Purple shell	Voss & Voss (1960)
<u>Thais rusticata</u>	Purple shell	Voss & Voss (1960)
<u>Varicorbula operculata</u>	Pelycepod	Newell et al (1959)
<u>Vermicularis spirata</u>	Pelycepod	Voss & Voss (1960)
<u>Verticordia ornata</u>	Pelycepod	Newell et al (1959)
<u>ECHINODERMATA</u>		
<u>Actinopyga agassizii</u>	Sea cucumber	Deichmann (1957)
<u>Amphipoda repens</u>	Brittle star	Clark (1942)
<u>Amphipholis gracillima</u>	Brittle star	Clark (1942?)
<u>Amphipholis squemata</u>	Brittle star	Clark (1942)
<u>Amphiura Steransi</u>	Brittle star	Ives (1891)
<u>Asterina folium</u>	Sea star	Clark (1942)
<u>Asteroporpa annulata</u>	Brittle star	Clark (1942)
<u>Astracea longispina</u>	Sea star	Ives (1891)
<u>Astropecten articulatus</u>	Sea star	Ives (1891)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ECHINODERMATA (Continued)</u>		
<u>Astropecten duplicatus</u>	Sea star	Ives (1891)
<u>Astrophyton costatum</u>	Brittle star	Ives (1891)
<u>Cidaris tribuloides</u>	Feather stars	Ives (1891)
<u>Clypeaster rosaceus</u>	- - -	Newell <i>et al</i> (1959)
<u>Diodena antillarum</u>	Black urchin	Voss & Voss (1960)
<u>Echinanthus rosaceus</u>	Feather stars	Ives (1891)
<u>Echinaster sentus</u>	Starfish	Newell <i>et al</i> (1959)
<u>Echinaster spinosus</u>	Sea star	Ives (1891)
<u>Echinometra lucunter</u>	Boring sea urchin	Voss & Voss (1960)
<u>Echinometra subangularis</u>	Feather star	Ives (1891)
<u>Euapta lappa</u>	Sea cucumber	Deichmann (1957)
<u>Eucardis tribuloides</u>	Sea urchin	Voss & Voss (1960)
<u>Hysponae esculenta</u>	Feather stars	Ives (1891)
<u>Holothuria cubana</u>	Sea cucumber	Clark (1942)
<u>Holothuria glaberrima</u>	Soft-skin sea cucumber	Deichmann (1957)
<u>Holothuria grisea</u>	Sea cucumber	Deichmann (1957)
<u>Holothuria impatiens</u>	Warted sea cucumber	Deichmann (1957)
<u>Holothuria parvula</u>	Sea cucumber	Deichmann (1957)
<u>Holothuria princeps</u>	Sea cigar	Deichmann (1957)
<u>Linckia Guildingii</u>	Sea star	Ives (1891)
<u>Luidia clathrata</u>	Sea star	Ives (1891)
<u>Lytechinus variegatus</u>	Echinoid	Newell <i>et al</i> (1959)
<u>Mellita sexiesperforata</u>	Echinoid	Newell <i>et al</i> (1959)
<u>Myona ventricosa</u>	- - -	Newell <i>et al</i> (1959)
<u>Nemaster iowensis</u>	Feather star	Clark (1942)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ECHINODERMATA (Continued)</u>		
<u>Ophiactis algicola</u>	Brittle star	Clark (1942)
<u>Ophiactis mulleri</u>	Brittle star	Ives (1891)
<u>Ophiactis savignyi</u>	Brittle star	Clark (1942)
<u>Ophiocoma pumila</u>	Brittle star	Pearse (1950)
<u>Ophiocoma riisei</u>	Brittle star	Clark (1942)
<u>Ophiomyxa flaccida</u>	Brittle star	Clark (1942)
<u>Ophiostigma isacanthum</u>	Brittle star	Clark (1942)
<u>Ophiothrix angulata</u>	Brittle star	Clark (1942)
<u>Ophiothrix oerstedi</u>	Brittle star	Ives (1891)
<u>Ophiothrix suensonii</u>	Brittle star	Clark (1942)
<u>Ophiura cincerea</u>	Brittle star	Ives (1891)
<u>Oreaster reticulatus</u>	Sea star	Clark (1942)
<u>Parathyone surinamensis</u>	Button sea cucumber	Deichmann (1957)
<u>Pentaceros reticulatus</u>	Sea star	Ives (1891)
<u>Pentacucumis planci</u>	Sea cucumber	Deichmann (1957)
<u>Stephanasterias gracilis</u>	Sea star	Clark (1942)
<u>Stichopus badinotus</u>	Sea cucumber	Deichmann (1957)
<u>Stylocidaris affinis</u>	Sea urchin	Clark (1942)
<u>Stolasterias tenuispina</u>	Sea star	Clark (1942)
<u>Synaptus sp.</u>	Sea cucumber	Newell <u>et al</u> (1959)
<u>Toxopneustes variegatus</u>	Feather star	Ives (1891)
<u>Tripneustes esculentus</u>	Sea urchin	Voss & Voss (1960)
<u>ARTHROPODA</u>		
<u>Arenopontia longiremis</u>	Copepod	Chappuis & Deboutteville (1956)
<u>Arenostella palpilabra</u>	Copepod	Chappuis & Deboutteville (1956)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ARTHROPODA (Continued)</u>		
<u>Callinectes exasperatus</u>	Brachyuran crab	Voss & Voss (1960)
<u>Chthamalus angustitergum</u>	Barnacle	Newell et al (1959)
<u>Chthamalus stellatus</u>	Barnacle	Voss & Voss (1960)
<u>Clibanarius sp.</u>	Decapod	Pearse (1950)
<u>Crangon cylindricus</u>	Pistol shrimp	Pearse (1950)
<u>Crangon floridonus</u>	Pistol shrimp	Pearse (1950)
<u>Crangon formosus</u>	Pistol shrimp	Pearse (1950)
<u>Crangon normani</u>	Pistol shrimp	Pearse (1950)
<u>Crangon ornatus</u>	Pistol shrimp	Pearse (1950)
<u>Dromidia antillensis</u>	Sponge crab	Voss & Voss (1960)
<u>Galappa flammula</u>	Crab	Newell et al (1959)
<u>Gnathophyllum americanus</u>	Decapod	Pearse (1950)
<u>Gonodactylus oerstedii</u>	Stomatapod	Pearse (1950)
<u>Grapsus grapsus</u>	Sally-light-foot	Voss & Voss (1960)
<u>Heteractae ceratopus</u>	Decapod	Pearse (1950)
<u>Heteromyysis sp.</u>	Mysid	Clarke (1955)
<u>Horisella trisaetosa</u>	Copepod	Chappuis & Deboutteville (1956)
<u>Laphontia arenicola</u>	Copepod	Chappuis & Deboutteville (1956)
<u>Laphontia Renaudi</u>	Copepod	Chappuis & Deboutteville (1956)
<u>Leucothoe commensalis</u>	Stomatapod	Pearse (1950)
<u>Lithotryxa dorsalis</u>	Boring barnacle	Voss & Voss (1960)
<u>Macrocelona trispinosum</u>	Crab	Pearse (1950)
<u>Microcerbus Renaudii</u>	Isopod	Chappuis & Deboutteville (1956)
<u>Microphyne bicornuties</u>	Crab	Voss & Voss (1960)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>ARTHROPODA (Continued)</u>		
<u>Microcerbus mirabilis</u>	Isopod	Chappuis & Deboutteville (1956)
<u>Microcerbus littoralis</u>	Isopod	Chappuis & Deboutteville (1956)
<u>Modulus sp.</u>	Hermit crab	Voss & Voss (1960)
<u>Murex sp.</u>	Hermit crab	Voss & Voss (1960)
<u>Pachygrapsus traversus</u>	Piling crab	Voss & Voss (1960)
<u>Panopeus herbstii</u>	Crab	Pearse (1950)
<u>Panopeus occidentalis</u>	Brachyuran crab	Voss & Voss (1960)
<u>Panulinus argus</u>	W. Indian crawfish	Newell <u>et al</u> (1959)
<u>Periclimenes americanus</u>	Shrimp	Pearse (1950)
<u>Periclimenes sp.</u>	Shrimp	Pearse (1950)
<u>Porcellina sayna</u>	Crab	Pearse (1950)
<u>Processa sp.</u>	Crab	Pearse (1950)
<u>Synalpheus brevicarpus</u>	Alpheid shrimp	Pearse (1950)
<u>Synalpheus brookski</u>	Alpheid shrimp	Pearse (1950)
<u>Synalpheus longicarpus</u>	Alpheid shrimp	Pearse (1950)
<u>Synalpheus minus</u>	Alpheid shrimp	Pearse (1950)
<u>Synalpheus pectinger</u>	Alpheid shrimp	Pearse (1950)
<u>Synalpheus rathbunae</u>	Alpheid shrimp	Pearse (1950)
<u>Tetraclita squamosa stalactifera</u>	Barnacle	Voss & Voss (1960)
<u>Tetraclita sp.</u>	Ribbed barnacle	Voss & Voss (1960)
<u>Tipton tortugae</u>	Crab	Pearse (1950)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>FISH - SHARKS and RAYS</u>		
<u>Actobatis narinari</u>	Spotted eagle ray	Bigelow and Schroeder (1948)
<u>Alopias vulpinus</u>	Thresher shark	Bigelow and Schroeder (1948)
<u>Carcharhinus lionbatus</u>	Ground shark	Bigelow and Schroeder (1948)
<u>Carcharhinus longimanus</u>	White tip shark	Bigelow and Schroeder (1948)
<u>Carcharodon charcharias</u>	Great white shark	Bigelow and Schroeder (1948)
<u>Galeocerdo cuvier</u>	Tiger shark	Bigelow and Schroeder (1948)
<u>Ginglymostoma cirratum</u>	Nurse shark	Bigelow and Schroeder (1948)
<u>Hexanchus griseus</u>	Mud shark	Bigelow and Schroeder (1948)
<u>Isurus oxyrinchus</u>	Mackerel shark	Bigelow and Schroeder (1948)
<u>Manta sp.</u>	Devil fish	La Garce (1919)
<u>Mustelus canis</u>	Smooth dogfish	Bigelow and Schroeder (1948)
<u>Negaprion brevirostris</u>	Lemon shark	Bigelow and Schroeder (1948)
<u>Parmaturus xanthurus</u>	Filetail shark	Bigelow and Schroeder (1948)
<u>Prionace glauca</u>	Blue shark	Bigelow and Schroeder (1948)
<u>Rhincodon typus</u>	Whale shark	Gundger (1939)
<u>Scyliorhinus carriculus</u>	Cat shark	Gundger (1939)
<u>Sphyrna zygaena</u>	Hammerhead shark	Gundger (1939)
<u>Squalus acanthias</u>	Spiny dogfish	Gundger (1939)
<u>FISH - BONY FISH</u>		
<u>Abudeduf exustilis</u>	Seargent major	Woods (1952)
<u>Acanthurus bahianus</u>	Ocean surgeon	Fish (1954)
<u>Acanthurus chirurgus</u>	Doctor fish	Fish (1954)
<u>Acanthurus coeruleus</u>	Blue tang	Moulton (1958)
<u>Albula vulpes</u>	Bone fish	Fish (1954)
<u>Anisotremus virginicus</u>	Pork fish	Fish (1954)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>FISH - BONY FISH (Continued)</u>		
<u><i>Bagre marinus</i></u>	Gaftopsail catfish	Fish (1954)
<u><i>Balistes vetula</i></u>	Queen triggerfish	Moulton (1958)
<u><i>Calamus calamus</i></u>	Saucerey porgy	Fish (1954)
<u><i>Calamus providens</i></u>	Porgy	Moulton (1958)
<u><i>Canthidermis sabaco</i></u>	Ocean triggerfish	Moulton (1958)
<u><i>Caranx hippos</i></u>	Horse-eye jack	Moulton (1958)
<u><i>Chaetodipterus faber</i></u>	Spade fish	Fish (1952)
<u><i>Chilomycterus atinga</i></u>	Spotted burrfish	Fish (1954)
<u><i>Coryphaena sp.</i></u>	Dolphin fish	Fish (1954)
<u><i>Cypsalurus sp.</i></u>	Flying fish	Mowbray (1956)
<u><i>Diodon holacanthus</i></u>	Ballon fish	Fish (1954)
<u><i>Diodon hystrix</i></u>	Porcupine fish	Moulton (1958)
<u><i>Diplectrum formosum</i></u>	Sand perch	Fish 1954)
<u><i>Epinephelus adscensionis</i></u>	Rock hind	Moulton (1958)
<u><i>Epinephelus striatus</i></u>	Nassau grouper	Moulton (1958)
<u><i>Eupomacentrus laucosticus</i></u>	Bean-gregory	Fish (1954)
<u><i>Euthynnus pelamis</i></u>	Skipjack tuna	Mowbray (1956)
<u><i>Galeichthys felis</i></u>	Sea catfish	Mowbray (1956)
<u><i>Gymnothorax funebris</i></u>	Green moray	Mowbray (1956)
<u><i>Haemulon album</i></u>	Margate	Moulton (1958)
<u><i>Haemulon carbonarium</i></u>	Caesar grunt	Fish (1954)
<u><i>Haemulon flavolineatum</i></u>	French grunt	Moulton (1958)
<u><i>Haemulon macrostomum</i></u>	Spanish grunt	Fish (1954)
<u><i>Haemulon melanurum</i></u>	Cotton wick	Fish (1954)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>FISH - BONY FISH (Continued)</u>		
<u>Haemulon parrai</u>	Sailor's choice	Fish (1954)
<u>Haemulon plumieri</u>	White grunt	Fish (1954)
<u>Haemulon sciurus</u>	Yellow grunt	Moulton (1958)
<u>Halichoeres bivittatus</u>	Slippery dick	Fish (1954)
<u>Halichoeres radiatus</u>	Pudding wife	Fish (1954)
<u>Holocentrus ascensionis</u>	Squirrel fish	Moulton (1958)
<u>Kyphosus sectatrix</u>	Bermuda chub	Fish (1954)
<u>Lachnolaimus maximus</u>	Hog fish	Fish (1954)
<u>Lactophrys bicaudalis</u>	Spotted trunkfish	Fish (1954)
<u>Lactophrys tricornis</u>	Cowfish	Moulton (1958)
<u>Lutjanus apodus</u>	Schoolmaster	Fish (1954)
<u>Lutjanus griseus</u>	Gray snapper	Moulton (1958)
<u>Lutjanus synagris</u>	Lane snapper	Fish (1954)
<u>Melacanthus plumieri</u>	- - - - -	Moulton (1958)
<u>Megalops atlantica</u>	Tarpon	Mowbray (1956)
<u>Mycteroperca bonaci</u>	Black grouper	Fish (1954)
<u>Myceteroperca venosa</u>	Yellowfin grouper	Mowbray (1956)
<u>Myrophamprus</u>	Flying fish	Mowbray (1956)
<u>Naucrates ductor</u>	Pilotfish	Mowbray (1956)
<u>Ocyurus chrysurus</u>	Yellowtail snapper	Mowbray (1956)
<u>Oligoplites saurus</u>	Leatherjacket	Mowbray (1956)
<u>Opsanus beta</u>	Oyster toadfish	Tavolga (1958)
<u>Opsanus tau</u>	Gulf toadfish	Tavolga (1958)
<u>Pomacentrus leucostriatus</u>	Demoiselle	Moulton (1958)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>FISH - BONY FISH (Continued)</u>		
<u>Pomacanthus itaira</u>	- - - - -	Moulton (1958)
<u>Pomacanthus arcuatus</u>	Black angelfish	Moulton (1958)
<u>Pomacanthus paru</u>	French angelfish	Moulton (1958)
<u>Scarus coeruleus</u>	Blue parrotfish	Fish (1954)
<u>Scarus guacamia</u>	Rainbow parrotfish	Fish (1954)
<u>Comber comberus</u>	Atlantic mackel	Mowbray (1956)
<u>Scorpaena sp.</u>	Scorpean fish	Moulton (1958)
<u>Seriola dumerili</u>	Amberjack	Fish (1954)
<u>Sphaeroides maculatus</u>	Common puffer	Moulton (1958)
<u>Sphaeroides spengleri</u>	Checkered puffer	Moulton (1958)
<u>Sphyraena barracuda</u>	Barracuda	Mowbray (1956)
<u>Synodus saphidoma</u>	Houndfish	Mowbray (1956)
<u>Synodus longylura sp.</u>	Needlefish	Mowbray (1956)
<u>Thunnus thynnus</u>	Bluefin tuna	Krumholz (1959)
<u>REPTILES</u>		
<u>Caretta caretta</u>	Loggerhead turtle	Ingle & Smith (1949)
<u>Chelonia mydas</u>	Green turtle	Ingle & Smith (1949)
<u>Dermochelys coriacea</u>	Leatherback turtle	Ingle & Smith (1949)
<u>Thallasochelys imbricata</u>	Hawksbill turtle	Ingle & Smith (1949)
<u>Thallasochelys kempii</u>	Kemp's turtle	Hanlon (1957-1961)
<u>MAMMALS</u>		
<u>Balaenoptera physalus</u>	Finback whale	Kellogg (1929)

APPENDIX 2 (Continued)

<u>Scientific Name</u>	<u>Common Name</u>	<u>Reference Source</u>
<u>MAMMALS (Continued)</u>		
<u>Megaptera novaeangliae</u>	Humpback whale	Kellogg (1929)
<u>Mesoplodon densirostris</u>	Beaked whale	Moore (1958)
<u>Orcinus orca</u>	Killer whale	Caldwell <u>et al</u> (1956)
<u>Pseudorca crassidens</u>	False Killer whale	Bullis & Moore (1956)
<u>Sibbaldus musculus</u>	Blue whale	Kellogg (1929)