Australian Tropical Rainforest Plants - Online edition

Phaleria chermsideana (F.M.Bailey) C.T.White

Family:

Thymelaeaceae

White, C.T. (1919) Queensland Department of Agriculture and Stock. Botany Bulletin 21: 19.

Common name:

Scrub Daphne

Stem

Grows into a small tree but probably does not exceed 30 cm dbh. Sometimes flowers and fruits as a shrub.

Leaves

Leaf blades about $7-8 \times 2.5-3.5$ cm, petioles about 0.5-0.6 cm long. Lateral veins forming loops inside the blade margin. Twig bark strong and fibrous, emitting a faint hydrocyanic odour when stripped.

Flowers

Umbels about 4-10-flowered. Peduncle about 2-20 mm long. Flowers +/- sessile. Perianth tube (hypanthium) about 8-17 mm long. Tepals about 3-6 mm long, alternating with five indistinct glands or lobes. Stamens attached near the apex of the perianth tube (hypanthium), filaments about 2 mm long. Pollen yellow. Ovary +/- surrounded by a glandular cupule, about 0.5 mm long. Style about 12-14 mm long. Stigma large, globular to cylindrical.

Fruit

Fruit about 10-15 mm long. Endocarp tough and fibrous. Seeds ovoid, about 7 x 3.5 mm.

Seedlings

Features not available.

Distribution and Ecology

Occurs in NEQ, south-eastern Queensland and north-eastern New South Wales. Altitudinal range in NEQ from 700-1100 m. Grows as an understory plant in mountain rain forest or on the boundary of mountain rain forest and wet sclerophyll forest.

Natural History & Notes

A bushy shrub or small tree with glossy dark green leaves and fragrant white flowers. It is becoming more common in cultivation.

Synonyms

Leucosmia chermsideana F.M.Bailey, *Queensland Agric. J.* 14: 35(1904), Type: Queensland, Killarney, J. Wedd; syn; Beechmont, Oct. 1918, C. England; syn: NSW.


RFK Code

1082

Copyright © CSIRO 2020, all rights reserved.







Buds. © A. Ford



Flowers and buds. © Stanley Breeden



Fruit [not vouchered]. © G. Sankowsky



Scale bar 10mm. © CSIRO















Web edition hosted at https://apps.lucidcentral.org/rainforest