





cm 1 2 3 4 5 6 SciELO 10 11 12 13 14 15



SciELO






DEPARTAMENTO DE ZOOLOGIA
SECRETARIA DA AGRICULTURA DO ESTADO DE SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

DEPARTAMENTO DE ZOOLOGIA
BIBLIOTECA
Secretaria da Agricultura

SÃO PAULO
[1766/67]



595.77
S239c
v.1

CANCELADO

DEDALUS - Acervo - MZ

Catalogue of the diptera of the americas south of united states /



12400006017



A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

1

INTRODUCTION

PREFACE


CURTIS W. SABROSKY

*Entomology Research Division, Agr. Res. Serv.
U. S. Department of Agriculture
Washington, D.C.*

Catalogues of the genera and species of living organisms are vitally essential as guides and reference works to the continually expanding knowledge of the fauna and flora. Nowhere is this more true than in the class Insecta, which includes approximately three-fourths of the known kinds of animals. And in this age of literature explosion, it is urgently necessary that there be adequate catalogues, supplemented or revised at suitable intervals. A complete summary and index to the literature would be desirable, but for many groups their preparation would be an insuperably massive task. In the large orders of insects, even a mere listing of the genera and species is a formidable project that is not lightly undertaken and not easily carried forward to completion.

The order Diptera, comprising the true flies and including those variously called gnats, midges, mosquitoes, etc., includes about one-eighth of the known species of insects in the world as a whole, and approximately one-fifth in the relatively well-studied region of America north of Mexico. The world's tropics are especially rich in species of flies, and the varied fauna of the Neotropical Region may be expected to total much more than that of the Nearctic Region, for which there are 1,971 genera and 16,130 species recognized in the recently completed "A Catalog of the Diptera of America North of Mexico" (Stone *et al.*, 1965).


No comprehensive catalogue of Neotropical Diptera has ever been published. Hunter's "A Catalogue of the Diptera of South America" (two parts, 1900-1901) was incomplete, ending with the Mydaiidae. Townsend's (1892) catalogue of the "Calyprate Muscidae" has covered only a part of the remainder. Catalogues of local areas or countries (e.g., Stuardo, 1946, for Chile) are necessarily limited in scope. The North American catalogues by Osten Sacken (1858, 1878) and Aldrich



(1905) were complete only for the area south to and including Panama, and for the West Indies. For individual families, there are catalogues or monographs of the Neotropical fauna, such as Aezél's catalogue of the "Tylidae" (1949) and John Lane's two-volume monograph of the "Neotropical Culicidae" (1953), but such contributions do not always include all the geographical area specifically covered by the present project. For occasional families or genera — all too few — modern world catalogues cover the Neotropical Region (e.g., "A synoptic catalog of the mosquitoes of the world" by Stone, Knight & Stareke, 1959, and M. R. Wheeler's catalogue of the genus *Drosophila*, 1959). In a few families, world catalogues are to be found in the fascicles of the "Genera Insectorum", but the last on families of Diptera appeared in 1937 (Muscidae) and 1952 (Scatophagidae), and most of the relatively few fascicles on Diptera are now over a half century old. Faunal studies, such as the "Diptera of Patagonia and South Chile" published by the British Museum (Natural History) and the "Biologia Centrali-Americana", may be important contributions for large areas, but they are necessarily limited in scope, and the coverage is often uncertain in extent and depth.

Entomologists will therefore welcome the "Catalogue of the Diptera of the Americas south of the United States" now being initiated by the Departamento de Zoologia, Secretaria da Agricultura, São Paulo. This catalogue, in bringing together for the first time a comprehensive, organized summary of the genera and species of Diptera in the Neotropical Region, will be of inestimable significance and usefulness. Particular appreciation is due to the dipterists of the Departamento for their energy and enthusiasm in planning and carrying out the project, and to the Director of the Departamento, Dr. P. E. Vanzolini, for his interest and support. Authors of individual parts are to be commended for their cooperation in preparing the manuscripts on the separate families.

This catalogue, including as it does the continent south of the U.S.A., the adjacent oceanic islands such as Galapagos, Juan Fernandez and Falklands, and the West Indies, will exactly complement the one for America north of Mexico. The two catalogues necessarily overlap slightly in content because of the arbitrary geographical dividing line. The "Nearctic" one includes Neotropical genera and species that extend their ranges into the southern United States and Bermuda, and the "Neotropical" will include Nearctic taxa that range into Mexico. The formats of the two catalogues are similar, with only minor differences, and concomitant use of the two will be easy and efficient. Together the two catalogues will provide complete modern coverage for the western hemisphere in one of the largest orders of insects and one of outstanding importance in public health, preventive medicine, and agriculture.



FOREWORD

P. E. VANZOLINI

Director

*Departamento de Zoologia, Secretaria da Agricultura,
São Paulo, Brasil*


Museum zoologists have been traditionally concerned with amassing collections, cataloguing faunas, and assembling morphological data, such things for a long time taken as ends in themselves. In the last 25 years, however, the analysis of evolution at work and the interrelationships between ecology, behaviour and systematics have become very much our province, and nowadays even experimental biology has its place among the pinned, stuffed and alcoholic specimens.

This broadened scope, which affords far more intellectual satisfaction and attracts good minds to the field, at times results into subtraction of effort and time from the traditional tasks, which tend to lose prestige more than deserved by their fundamental importance. This ambivalence is especially acute in developing countries, where the need and desire to do advanced research clashes everyday with the tremendous volume of the materials to be collected, catalogued and described. Thus it becomes a matter of museum policy to search for an equilibrium between the two extremes and to acknowledge and enhance their complementarity.

This Catalogue is meant to be both a significant step in the systematic assessment of the Neotropical Diptera and a basis for further revisionary work, leading to an understanding of the history of the fauna and of the mechanisms operative in its evolution.

The size of the task is obviously too large not only for an individual South American museum, but even for the aggregate of all zoological institutions in the continent. We asked for help, and the uniformly warm response from all quarters is perhaps the most satisfying result of the venture. To the contributors to this Catalogue we are deeply indebted, and special mention must be made of Dr. Curtis W. Sabrosky, whose advice and help, asked from the beginning, were always more than generously given.

We have patterned our work after the "A Catalog of the Diptera of America North of Mexico", partly as a matter of convenience for consultation, but mostly in recognition of its excellent features.



SciELO

EXPLANATORY NOTE

NELSON PAPAVERO

*Departamento de Zoologia, Secretaria da Agricultura,
São Paulo, Brasil*

A few words are necessary to explain the format of our Catalogue. Unlike the North American one, this Catalogue will be issued in fascicles, each one treating a single family.

The Neotropical fauna is being actively studied; were we to wait for the preparation of all manuscripts in order to publish the Catalogue in one single volume, many of the treatments would have become obsolete at the time of publication. On the other hand, supplementation and substitution of individual fascicles are much easier.

Fascicles will be numbered according to the systematic arrangement of the families but published as soon as available, regardless of the sequence.

With some minor modifications, the format proper follows that of the North American Catalogue in order to permit concomitant use of both.

The list of intended fascicles is as follows:

- | | |
|--------------------------------|------------------------|
| 1. Introduction ✓ | 21. Hyperoscelididae ✓ |
| X 2. List of collaborators NED | 22. Scatopsidae ✓ |
| 3. Trichoderidae ✓ | 23. Cecidomyiidae ✓ |
| X 4. Tipulidae ✓ | 24. Xylophagidae ✓ |
| 5. Tanyderidae ✓ | 25. Xylomyiidae ✓ |
| 6. Psychodidae A ✓ | 26. Stratiomyidae ✓ |
| 7. Nymphomyiidae ✓ | 27. Pelecorhynchidae ✓ |
| 8. Blephariceridae ✓ | 28. Tabanidae ✓ |
| X 9. Deuterophlebiidae ✓ | 29. Rhagionidae ✓ |
| 10. Dixidae ✓ | 30. Pantoptalmidae ✓ |
| 11. Chaoboridae ✓ | X 31. Therevidae ✓ |
| X 12. Culicidae ✓ | 32. Scenopinidae ✓ |
| 13. Thaumaleidae ✓ | 33. Apiceridae ✓ |
| 14. Ceratopogonidae ✓ | 34. Mydidae ✓ |
| X 15. Chironomidae ✓ | 35. Asilidae A ✓ |
| 16. Simuliidae ✓ | 36. Nemestrinidae ✓ |
| 17. Anisopodidae ✓ | 37. Acroceridae ✓ |
| 18. Bibionidae ✓ | 38. Bombyliidae ✓ |
| 19. Mycetophilidae AE ✓ | 39. Empididae ✓ |
| X 20. Sciaridae ✓ | 40. Dolichopodidae ✓ |

X - now published

v

- | | |
|-----------------------|--|
| 41. Lonchopteridae ✓ | 76. Canaceidae ✓ |
| 42. Sciadoceridae ✓ | 77. Ephydriidae ✓ |
| 43. Phoridae ✓ | 78. Curtonotidae ✓ |
| 44. Platypezidae ✓ | 79. Drosophilidae ✓ |
| 45. Pipunculidae ✓ | 80. Diastatidae ✓ |
| 46. Syrphidae ✓ | 81. Chloropidae ✓ |
| 47. Conopidae ✓ | 82. Odiniidae ✓ |
| 48. Micropezidae ✓ | 83. Agromyzidae ✓ |
| 49. Neriidae A ✓ | 84. Clusiidae JA ✓ |
| 50. Psilidae ✓ | 85. Heleomyzidae JB ✓ |
| 51. Megamerinidae ✓ | 86. Trixoscelididae ✓ |
| 52. Tanypezidae ✓ | 87. Rhinotoridae ✓ |
| 53. Richardiidae ✓ | 88. Anthomyzidae A ✓ |
| 54. Otitidae ✓ | 89. Opomyzidae ✓ |
| 55. Platystomatidae ✓ | 90. Chyromyidae ✓ |
| 56. Pyrgotidae ✓ | 91. Aulacigastridae ✓ |
| 57. Tephritidae ✓ | 92. Asteiidae ✓ |
| 58. Tachiniscidae ✓ | 93. Cryptochetidae ✓ |
| 59. Helcomyzidae ✓ | 94. Somatiidae ✓ |
| 60. Ropalomeridae ✓ | 95. Acalyprate genera of uncertain family position ✓ |
| 61. Coelopidae ✓ | 96. Anthomyiidae AB ✓ |
| 62. Dryomyzidae ✓ | 97. Muscidae ✓ |
| 63. Sepsidae ✓ | 98. Gasterophilidae ✓ |
| 64. Sciomyzidae ✓ | 99. Hippoboscidae ✓ |
| 65. Lauxaniidae A ✓ | 100. Streblidae ✓ |
| 66. Chamaemyiidae ✓ | 101. Nycteribiidae ✓ |
| 67. Periscelididae ✓ | 102. Calliphoridae ✓ |
| 68. Piophilidae ✓ | 103. Sarcophagidae ✓ |
| 69. Thyreophoridae ✓ | 104. Tachinidae ✓ |
| 70. Pallopteridae ✓ | 105. Cuterebridae ✓ |
| 71. Lonchaeidae ✓ | 106. Ocstridae ✓ |
| 72. Sphacrocridae ✓ | 107. Final Remarks ✓ |
| 73. Braulidae ✓ | 108. General bibliography ✓ |
| 74. Tethinidae ✓ | 109. General index ✓ |
| 75. Milichiidae ✓ | 110. List of type-localities ✓ |

GEOGRAPHICAL SCOPE

The area covered by this Catalogue is the American continent south of the United States-Mexico border plus the following islands: (i) all continental islands, including those of the Gulf of California; (ii) on the Atlantic side: Bahamas, Greater and Lesser Antilles, Fernando Noronha, São Pedro, São Paulo, Martin Vaz, South Trinidad, Falklands, South Georgia, Sandwiches and South Orkneys; (iii) on the Pacific side: Galapagos, Juan Fernandez, Clipperton, Cocos, Revillagigedo. That part of Antarctica just opposite the southern end of South America is also included.

For brevity's sake, the area covered by this Catalogue, as well as the taxa occurring in it, will be referred to as "Neotropical", regardless of the real limits of the zoogeographical region.

CLASSIFICATION

The classification here adopted follows mainly that of Hennig (1954, 1958) and modified in the North American Catalogue (Stone *et al.*, 1965):



Suborder Nematocera

Superfamily Tipuloidea

Family Trichoceridae .. ✓

Family Tipulidae .. ✓

Superfamily Psychodoidea

Family Tanyderidae ✓

Family Psychodidae.

Family Nymphomyiidae

Superfamily Culicoidea

Family Blepharictridae ..

Family Deuterophlebiidae

Family Dixidae ✓

Family Chaoboridae ..

Family Culicidae .. ✓

Family Thaumaleidae *

Family Ceratopogonidae ✓

Family Chironomidae ✓

Family Simuliidae. ✓

Superfamily Anisopodoidea

Family Anisopodidae .. ✓

Superfamily Bibionoidea

Family Bibionidae. ✓

Superfamily Mycetophiloidea

Family Mycetophilidae ✓

Family Sciaridae .. ✓

Family Hyperoscelididae

Family Scatopsidae ✓

Family Cecidomyiidae ✓

Suborder Brachycera

Superfamily Tabanoidea

Family Xylophagidae

Family Xylomyidae

Family Stratiomyidae

Family Pelecorhynchidae

Family Tabanidae

Family Rhagionidae

Family Pantophtalmidae

Superfamily Asiloidea

Family Therevidae

Family Scenopinidae

Family Apioceridae

Family Mydidae

Family Asilidae

Family Nemestrinidae

Family Acroceridae

Family Bombyliidae



Superfamily Empidoidea
Family Empididae
Family Dolichopodidae

Suborder Cyclorrhapha

Division Aschiza

Superfamily Lonchopteroidea
Family Lonchopteridae

Superfamily Phoroidea
Family Sciadoceridae
Family Phoridae
Family Platypezidae

Superfamily Syrphoidea
Family Pipunculidae
Family Syrphidae
Family Conopidae

Division Schizophora

Section Acalyptratae

Superfamily Micropezoidea
Family Micropezidae
Family Neriidae

Superfamily Nothyboidea
Family Psilidae
Family Megamerinidae
Family Tanypzidae

Superfamily Tephritoidea
Family Richardiidae
Family Otitidae
Family Platystomatidae
Family Pyrgotidae
Family Tephritidae
Family Tachiniscidae

Superfamily Sciomyzoidea
Family Helcomyzidae
Family Ropalomeridae
Family Coelopidae
Family Dryomyzidae
Family Sepsidae
Family Sciomyzidae

Superfamily Lauxanioidea
Family Lauxaniidae



Family Chamaemyiidae
Family Periseelididae

Superfamily Pallopteroidea

Family Piophilidae
Family Thyracophoridae
Family Pallopteridae
Family Lonchaeidae

Superfamily Milichioidea

Family Sphaeroceeridae
Family Braulidae
Family Tethinidae
Family Milichiidae
Family Canaceidae

Superfamily Drosophiloidea

Family Ephydriidae
Family Curtonotidae
Family Drosophilidae
Family Diastatidae

Superfamily Chloropoidea

Family Chloropidae

Unplaced families of Acalyptratae


Family Odiniidae
Family Agromyzidae
Family Clusiidae
Family Heleomyzidae
Family Trixoseelididae
Family Rhinotoridae
Family Anthemyzidae
Family Opomyzidae
Family Chyromyidae
Family Aulacigastridae
Family Asteiidae
Family Cryptoheidae
Family Somatiidae

Acalyptrate genera of uncertain
family position

Section Calyptratae

Superfamily Museoidea

Family Anthomyiidae
Family Muscidae
Family Gasterophilidae
Family Hippoboscidae
Family Streblidae
Family Nycteribiidae



Superfamily Oestroidea

Family Calliphoridae
Family Sarcophagidae
Family Tachinidae
Family Cuterebridae
Family Oestridae

INFORMATION CONTAINED

Each fascicle treats an individual family. In the introductory paragraphs a brief survey is made of the present state of knowledge of the biology, geographical distribution, morphology, anatomy, etc., of the family treated. Explicit mention is made of basic papers on relevant topics (revisions, catalogues, monographs, bibliographies, etc.). No nomenclatural references are made to the family. The same applies to subfamilies and tribes. When available, reference is made to a taxonomic revision of the whole taxon or of a significant group of genera.

Genera and subgenera — For each generic or subgeneric name, whether valid or a synonym, are given author, date, page, type-species, nature of type fixation and the present valid name of the type-species, if there has been a change. Generic synonyms are arranged chronologically. In the synonymic lists generic names based on exotic species are not included unless the names have come into significant usage in the American literature or unless the author of the fascicle, for some reason, considers it necessary. Generic and subgeneric headings are arranged either alphabetically or phylogenetically, according to the author's choice. Citation for the typical subgenus follows the generic rather than the subgeneric heading. For type-species, the original combination is used and the generic name omitted when it is the same as that of the genus for which it is the type. Authors are requested to follow the 1964 edition of the Code of Zoological Nomenclature for the fixation of type-species. Following the citations are given useful references to the genus or subgenus, especially of taxonomic nature, to the whole taxon or to significant groups of species.

Species and subspecific categories — Each specific or subspecific name is accompanied by the name of the author, date, page, figure, plate, name of the genus in which it was originally described (in parentheses), type-locality, and distribution.

The type-locality is explicitly stated without the use of abbreviations of geographical names (excepting U.S.A. and D.F.) in the following order: country, state (or equivalent administrative unit), locality proper (as complete as possible). The *Index to Map of Hispanic America 1:1,000,000* (Hanson, 1945) is here used as the basic standard reference work for geographical names. In cases of multiple type-localities or of allopatric syntypes, all localities are stated in the same order as they appeared in the original description of the taxon.

Distributional data are as complete as possible; in cases of a few scattered localities, full citation is given, in a north-south order. If the area covered by the species is too large, only outer limits are cited. However, in all of these cases a full list of islands is given.

Specific names are arranged alphabetically, following their respective generic or subgeneric headings. For specific synonyms and varieties, the same information (except distributional data) is furnished as for the species; such names are indented and listed chronologically.



Explicit mention is made of at least one good redescription, figure or useful detail, as an aid to the identification of the species, when this is available. Citation of the location of types is optional.

Errors and emendations are indented after the name misspelled or emended. Parentheses, indicating a shift in the original generic position of the species, are used only for valid specific names.

In cases of misidentifications, one of two procedures are followed (according to the North American Catalogue): (i) if the name of an exotic species has been erroneously applied to only one Neotropical species, it is listed in the synonymy of that species; the citation is thus given: "*punctatus*, authors, nec Fabricius"; a reference to the misuse may be given if it has been used only once but in a widely used publication; (ii) if the name has been misapplied to several valid Neotropical species, or if the identity of the Neotropical species apparently misnamed is unknown, then the name is cited in the proper alphabetical order in italics, without citation, followed by the notation: "not Neotropical".

Names of dubious identity are listed in regular alphabetical order when their generic position is reasonably certain and with the notation "unrecognized"; names of uncertain generic position are placed separately at the end of the respective tribe, subfamily or family; names of uncertain subgeneric position are placed at the end of the genus.


Nomina nuda are placed in synonymy if the application is known; otherwise, they are to be referred to the editors to be included in a special fascicle.

Unnamed species (e.g., *Rhinotora* sp.), accompanied by a description or figure, are also placed in synonymy when the application is known; otherwise, they are to be included in the regular alphabetical order under the respective generic or subgeneric headings. If an unnamed species was not accompanied by a description or figure, when it was originally published, it is included in this Catalogue when it represents the only published citation of the genus in the region covered by this Catalogue. Unpublished records of the existence of a certain taxon in the Neotropical region, or details on its geographical distribution based on collections data, are at times included, but this is explicitly stated by the author.

All other nomenclatural matters are to be solved with reference to the 1964 edition of the International Code of Zoological Nomenclature.

Each fascicle contains its own bibliography and index. Bibliographical references follow the pattern of the Zoological Record, but latitude is given to collaborators to adopt any consistent, easily understandable system. The index contains the name of the taxon, author, date, and, in cases of specific and subspecific names, the genus under which they now stand and (in parentheses) the genus in which they were originally described.

The pagination of each fascicle is independent, but the page number is preceded by the number of the fascicle. Thus, page 72.3 is the third page of fascicle 72. Each fascicle will bear its respective date of issue, for priority purposes.



LITERATURE CITED

ACZÉL, M.

1949. Catálogo de la familia Tylidae (+ Neriidae) de la región neotropical. *Acta zool. Lilloana* 8:309-389.

ALDRICH, J. M.

1905. A catalogue of North American Diptera. *Smiths. Misc. Coll.* 46(2):1-680.

HANSON, E. P. (ed.)

1945. *Index to Map of Hispanic America 1:1,000,000*, 923 pp. American Geographical Society, Publ. n° 5, U.S. Government Printing Office, Washington, D.C.

HENNIG, W.

1954. Flügelgeäder und System der Dipteren, unter Berücksichtigung der aus dem Mesozoikum beschriebenen Fossilien. *Beitr. z. Ent.* 4:245-388, 272 figs.
1958. Die Familien der Diptera Schizophora und ihre phylogenetischen Verwandtschaftsbeziehungen. *Beitr. z. Ent.* 8:505-688, 365 figs.

HUNTER, W. D.

- 1900-1901. A catalogue of the Diptera of South America, Parts I, II. *Trans. Amer. Ent. Soc.* 26:260-298; 27:121-155.

LANE, J.

1953. *Neotropical Culicidae*, 2 vols., 1,112 pp., 1,071 figs. Universidade de São Paulo.

OSTEN SACKEN, C. R.

1858. Catalogue of the described Diptera of North America. *Smiths. Misc. Coll.* 3(1): xx + 92 pp.
1878. Catalogue of the described Diptera of North America (ed. 2). *Smiths. Misc. Coll.* 16(2):1-276.

STONE, A., K. L. KNIGHT & H. STARCKE

1959. A synoptic catalog of the mosquitoes of the world. *Thomas Say Foundation* 6:358 pp., Washington, D.C.

STONE, A., C. W. SABROSKY, W. W. WIRTH, R. H. FOOTE & J. R. COULSON

1965. A catalog of the Diptera of America North of Mexico. *U.S. Department of Agriculture, Agriculture Handbook* 276:1547 pp., Washington, D.C.



STUARDO, C.


1946. *Catálogo de los dípteros de Chile*, 253 pp. Ministerio de Agricultura, Santiago.

TOWNSEND, C. H. T.

1892. Catalogue of the described South American species of Calyptrate Muscidae. *Ann. New York Acad. Sci.* 7:1-44.

WHEELER, M. R.

1959. A nomenclatural study of the genus *Drosophila*. *Univ. Texas Publ.* 5914:181-205.





SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

3

FAMILY TRICHOCERIDAE

(PETAURISTIDAE, MELUSINIDAE)

CHARLES P. ALEXANDER AND MABEL M. ALEXANDER

Amherst, Massachusetts

The family Trichoceridae, including the so-called winter crane-flies, from their occurrence as adults being chiefly in the colder months of the year, has approximately 90 known species that are distributed in 6 genera. These include the major group *Trichocera* Meigen with 49 species in the typical subgenus, and 11 others in the recently proposed subgenus *Metatrachocera* Dahl, 1966, all species being Holarctic in distribution, with a few that have been transported to various parts of the world by commerce. The remaining genera are *Diazosma* Bergroth (2), Holarctic; *Nothotrachocera* Alexander (5), of southeastern Australia and the subantarctic islands of New Zealand (Campbell Island); *Palaeopetaurista* Séguy (1), Kerguelen Island, Indian Ocean; *Parocladura* Brunetti (23), with most of the species occurring in New Zealand and southern South America; and the very distinct and isolated genus *Kawasemyia* Alexander (1), known only from central Japan.

The immature stages of various species of *Trichocera* are known, being found in animal manure, decaying vegetable matter, fungi, and in stored roots and tubers (Karandikar, 1931; Keilin, 1912; Laurence, 1956; Rhynehart, 1925).

In the "Catalog of the Diptera of America North of Mexico" (1965) this family was placed with the Tipulidae in the superfamily Tipuloidea, and this arrangement was followed by Papavero in the *Introduction to the present Catalogue*. In our opinion the Trichoceridae should be placed with the Anisopodidae in the superfamily Anisopodoidea, while the Tipuloidea include only the family Tipulidae. The various stages of the Trichoceridae show little in common with the Tipulidae but have undisputed characters that are found in the Anisopodidae, as discussed by various authors (Anthon, 1943; Edwards, 1928; Keilin, 1912; Rhynehart, 1925).

Genus TRICHOCERA Meigen

Petaurista Meigen, 1800:15 (preocc. Link, 1795). Type-species, *Tipula hiemalis* De Geer (Coquillett, 1910:587). Suppressed by I.C.Z.N., 1963:339.



- Melusina* Meigen, 1800:19. Type-species, *Tipula regelationis* Linnaeus (Hendel, 1908:50). Suppressed by I.C.Z.N., 1963:339.
- TRICHOcera Meigen, 1803:262. Type-species, *Tipula hiemalis* De Geer (mon.).
- Trichocerodes* Brèthes, 1925:169. Type-species, *georgianus* Brèthes (mon.) = *regelationis* (Linnaeus).
- MEXICANA Alexander, 1946:219, fig. (venation). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- REGELATIONIS (Linnaeus), 1758:587 (*Tipula*). Type-locality: "Sweden". Distr. — Europe; eastern North America; Falkland and South Georgia Islands.
- georgianus* Brèthes, 1925:169 (*Trichocerodes*). Type-locality: South Georgia and Falkland Islands (South Atlantic; British).

Genus PARACLADURA Brunetti

- PARACLADURA Brunetti, 1911:286. Type-species, *gracilis* Brunetti (orig. des.).
- CHILENSIS Alexander, 1929:225, fig. C5 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- EDWARDSI Alexander, 1929:223, figs. C1 (venation), C4 (male hypopygium), C8 (ovipositor). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Argentina, Chile.
- PATAGONICA Alexander, 1929:226, figs. C2 (venation), C6 (male hypopygium), C9 (ovipositor). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi. Distr. — Argentina, Chile.
- PIRIONI Alexander, 1929:227, figs. C3 (venation), C7 (male hypopygium). Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.

LITERATURE CITED

ALEXANDER, C. P.

1929. Trichoceridae, in *Diptera of Patagonia and South Chile*, Part I: i-xvi, 1-240, 12 pls. with 259 figs.
1946. Records and descriptions of Mexican crane-flies (Dipt. Tipulidae). *Anales Escuela Nac. Cienc. Biol.* 4:213-253, 28 figs.

ANTHON, HENNING


1943. Der Kopfbau der Larven einiger nematoceren Dipteren-Familien. *Spoelia Zool. Mus. Hauniensis*, Copenhagen 3:1-61, 11 pls. with 62 figs.

BRÈTHES, JEAN

1925. Un Coléoptère et un Diptère nouveaux de la Géorgie du Sud. *Comun. Mus. Nac. Hist. Nat. "Bernardino Rivadavia"*, Buenos Aires 2:169-173, 2 figs.

BRUNETTI, E.

1911. Revision of the Oriental Tipulidae with descriptions of new species. *Rec. Indian Mus.* 6:231-314.



COQUILLET, D. W.

1910. The type-species of the North American genera of Diptera. *Proc. U. S. Nat. Mus.* 37:499-647.

DAHL, CHRISTINE

1966. Notes on the taxonomy and distribution of Swedish Trichoce-
ridae (Dipt. Nemat.). *Opuscula Entomologica* 31:93-118, 72 figs.

EDWARDS, F. W.

1928. Diptera. Fam. Protorhyphidae, Anisopodidae, Pachyneuridae,
Trichoceridae, in P. Wytsman: *Genera Insectorum* 190:40 pp., pl.
1, 14 figs., pl. 2, 18 figs.

HENDEL, F.

1908. Nouvelle classification des mouches à deux ailes (Diptera L.),
d'après un plan tout nouveau par J. G. Meigen, Paris, an VIII
(1800 v.s.). Mit einem Kommentar. *Verhandl. k. k. zool.-bot. Ges.
Wien* 58:43-69.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. Opinion 678. The suppression under the Plenary Powers of
the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.*
20:339-342.

KARANDIKAR, K. R.

1931. The early stages and bionomics of *Trichocera maculipennis*
(Meigen) (Diptera, Tipulidae). *Trans. R. ent. Soc. London*
79:249-262, figs.

KEILIN, DAVID

1912. Recherches sur les Diptères du genre *Trichocera*. *Bull. Scient.
France et Belgique* (2) 46(2):172-191, 4 pls. with 27 figs.

LAURENCE, B. R.

1956. On the life history of *Trichocera saltator* (Harris) (Diptera,
Trichoceridae). *Proc. Zool. Soc. London* 126(2):235-243, 12 figs.

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*. Ed. 10, Vol. 1, 824
pp. Holmiae (= Stockholm).


MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.),
.. d'après un plan tout nouveau*, 40 pp., Paris.

1803. Versuch einer neuen Gattungseintheilung der europäischen
zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2:259-281.


RHYNEHART, J. G.

1925. The larva and pupa of *Trichocera regelationis* L. (Diptera,
Rhyphidac). *Proc. Belfast Nat. Hist. and Phil. Soc.*, 102nd session
(1922-23):14 pp., 3 pls. with 27 figs.



INDEX

<i>chilensis</i> Alexander, 1929, <i>Paracladura</i>	3.2
<i>edwardsi</i> Alexander, 1929, <i>Paracladura</i>	3.2
<i>georgianus</i> (Brèthes), 1925 (<i>Trichocerodes</i>), <i>Trichocera</i>	3.2
<i>Melusina</i> Meigen, 1800	3.2
<i>mexicana</i> Alexander, 1946, <i>Trichocera</i>	3.2
<i>Paracladura</i> Brunetti, 1911	3.2
<i>patagonica</i> Alexander, 1929, <i>Paracladura</i>	3.2
<i>Petaurista</i> Meigen, 1800	3.1
<i>pirioni</i> Alexander, 1929, <i>Paracladura</i>	3.2
<i>regelationis</i> (Linnaeus), 1758 (<i>Tipula</i>), <i>Trichocera</i>	3.2
<i>Trichocera</i> Meigen, 1803	3.1, 3.2
<i>Trichocerodes</i> Brèthes, 1925	3.2



SUBORDER NEMATOCERA






SUPERFAMILY TIPULOIDEA

Family Trichoceridae

Family Tipulidae





SciELO

MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
OF THE UNITED STATES

4

FAMILY TIPULIDAE

CHARLES P. ALEXANDER AND MABEL M. ALEXANDER

Amherst, Massachusetts

The family Tipulidae, comprising the true crane flies, is one of the largest families of Diptera in the Neotropical fauna, at present including slightly more than 3,000 species of the more than 13,000 described members of the family. The present fauna is still poorly known and the number of species here included presumably is not more than one-half of those that actually occur.

A number of the species that were named by the early workers remain in question. The original descriptions of these are brief and unsatisfactory and for several, accurate identification is impossible without a comparison with type material. A lack of exact geographical data for several of these species further limits the chances of identifying such materials. A major proportion of the taxonomic work on this family of flies has been done by the senior author of the present catalogue (see Literature cited. Alexander, 1911-date).

Very little is known regarding the biology and immature stages of Neotropical species in this family, the more important references being by Alexander (for *Sigmatomera*, 1930a), Bruch (1939b), Edwards (1934), and Picado (*Trentepohlia*, as *Mongoma*, 1913). Various genera occurring in North America, where the biology is known, have representatives in the Neotropics and these presumably have similar or comparable life histories. Of particular importance are four papers by the late J. S. Rogers considering *Orimarga* (*Diotrepha*), 1927a; *Atarba*, 1927b; *Polymera*, 1933; and *Megistocera*, 1949. The most important single paper on the biology of Neotropical crane flies is the one by Bruch, cited above. Further references to the immature stages of North American crane flies that will prove helpful to workers on the species found within the geographical limits of the catalogue include those by Alexander, 1920i, including a general summary of crane fly biology; Eyers (*Dolichozeza*, 1961), and Hynes (*Cryptolabis*, 1963, and *Lipsothrix* 1965).

The arrangement of genera in the various subfamilies and tribes is shown, with an indication of the number of species, and a summary of the total number of species included in the major divisions of the family.

TIPULINAE

Number of species

Brachypremna Osten Sacken	35
Ctenophora Meigen	3
<i>Pselliophora</i> Osten Sacken	(3)
Dolichocheza Curtis	8
<i>Megistomastix</i> Alexander	(8)
Elnoretta Alexander	1
Ischnotoma Skuse	31
<i>Icriomastax</i> Enderlein	(10)
<i>Ischnotoma</i> Skuse	(17)
<i>Neotipula</i> Alexander	(4)
Leptotarsus Guérin-Méneville	89
<i>Aldrovandia</i> Enderlein	(1)
<i>Araucomyia</i> Alexander	(3)
<i>Ceoneura</i> Alexander	(1)
<i>Leptotarsus</i> Guérin-Méneville	(1)
<i>Limoniodes</i> Alexander	(1)
<i>Longurio</i> Loew	(35)
<i>Pehlkea</i> Enderlein	(4)
<i>Tanypremna</i> Osten Sacken	(30)
<i>Tanypremnella</i> Alexander	(10)
<i>Tanypremnodes</i> Alexander	(2)
<i>Xenotipula</i> Alexander	(1)
Megistocera Wiedemann	1
Nephrotoma Meigen	21
Ozodicera Macquart	54
<i>Dihexaclonus</i> Enderlein	(21)
<i>Ozodicera</i> Macquart	(33)
Phacelodocera Enderlein	1
Tipula Linnaeus	410
<i>Bellardina</i> Edwards	(9)
<i>Eumicrotipula</i> Alexander	(236)
<i>Lunatipula</i> Edwards	(3)
<i>Microtipula</i> Alexander	(145)
<i>Pectinotipula</i> Alexander	(4)
<i>Pterelachisus</i> Rondani	(1)
<i>Trichotipula</i> Alexander	(6)
<i>Triplicitipula</i> Alexander	(1)
<i>Yamatotipula</i> Matsumura	(5)
Valdiviana Alexander	4
Zelandotipula Alexander	50
Unplaced Tipulinae	10

Total: 718

CYLINDROTOMINAE

Stibadocerina Alexander	1
-------------------------	---

LIMONIINAE

LIMONINI

Antocha Osten Sacken	1
Dicranoptycha Osten Sacken	3

Helius Lepeletier & Serville	39
Limonia Meigen	598
<i>Atypophthalmus</i> Brunetti	(1)
<i>Caenoglochina</i> Alexander	(18)
<i>Caenolimonia</i> Alexander	(15)
<i>Dicranomyia</i> Stephens	(188)
<i>Discobola</i> Osten Sacken	(1)
<i>Geranomyia</i> Haliday	(142)
<i>Idioglochina</i> Alexander	(3)
<i>Neoglochina</i> Alexander	(21)
<i>Neolimnobia</i> Alexander	(11)
<i>Neolimonia</i> Alexander	(33)
<i>Peripheroptera</i> Schiner	(36)
<i>Rhipidia</i> Meigen	(106)
<i>Zahsa</i> Enderlein	(1)
<i>Zelandoglochina</i> Alexander	(19)
Unplaced <i>Limonia</i>	(3)
Orimarga Osten Sacken	45
<i>Diotrepha</i> Osten Sacken	(14)
<i>Orimarga</i> Osten Sacken	(31)
Protohelius Alexander	2
Tonnoiromyia Alexander	1

Total: 689

PEDICIINI

Dicranota Zetterstedt	3
<i>Plectromyia</i> Osten Sacken	(1)
<i>Rhaphidolabis</i> Osten Sacken	(2)
Pedicia Latreille	9
<i>Tricyphona</i> Zetterstedt	(9)
Unplaced Pediciini	1

Total: 13

HEXATOMINI

Atarba Osten Sacken	87
<i>Atarba</i> Osten Sacken	(61)
<i>Ischnothrix</i> Bigot	(26)
Austrolimnophila Alexander	57
<i>Austrolimnophila</i> Alexander	(46)
<i>Limnophilella</i> Alexander	(9)
<i>Mediophragma</i> Alexander	(2)
Chilelimnophila Alexander	1
Ctenolimnophila Alexander	7
<i>Abitagua</i> Alexander	(1)
<i>Campbellomyia</i> Alexander	(3)
<i>Ctenolimnophila</i> Alexander	(3)
Edwardsomyia Alexander	1
Elephantomyia Osten Sacken	31
<i>Elephantomyia</i> Osten Sacken	(29)
<i>Elephantomyia</i> Alexander	(1)
<i>Xenoelephantomyia</i> Alexander	(1)
Epiphragma Osten Sacken	60

Gynoplistia Westwood	18
<i>Dirhipis</i> Enderlein	(3)
<i>Gynoplistia</i> Westwood	(15)
Hexatoma Latreille	146
<i>Cladolipes</i> Loew	(1)
<i>Eriocera</i> Macquart	(139)
Lecteria Osten Sacken	14
<i>Lecteria</i> Osten Sacken	(3)
<i>Psaronius</i> Enderlein	(11)
Limnophila Macquart	40
<i>Araucolimnophila</i> Alexander	(1)
<i>Habrolimnophila</i> Alexander	(1)
<i>Limnophila</i> Macquart	(38)
Mesolimnophila Alexander	2
Paradelphomyia Alexander	7
<i>Oxyrhiza</i> Meijere	(7)
Paralimnophila Alexander	14
Pilaria Sintenis	2
Polymera Wiedemann	52
<i>Polymera</i> Wiedemann	(44)
<i>Polymerodes</i> Alexander	(8)
Pseudolimnophila Alexander	5
Shannonomyia Alexander	74
<i>Roraimomyia</i> Alexander	(1)
<i>Shannonomyia</i> Alexander	(73)
Unplaced Hexatomini	10

Total: 622

ERIOPTERINI

Amphineurus Skuse	12
<i>Amphineurus</i> Skuse	(3)
<i>Rhamphoneurus</i> Alexander	(9)
Aphrophila Edwards	8
Aymaramyia Alexander	1
Baeoura Alexander	1
Cheilotrichia Rossi	25
<i>Empeda</i> Osten Sacken	(25)
Cryptolabis Osten Sacken	41
<i>Cryptolabis</i> Osten Sacken	(39)
<i>Procryptolabis</i> Alexander	(2)
Empedomorpha Alexander	1
Erioptera Meigen	68
<i>Erioptera</i> Meigen	(18)
<i>Eriopterella</i> Alexander	(3)
<i>Mesocyphona</i> Osten Sacken	(42)
<i>Psilocoenopa</i> Zetterstedt	(1)
<i>Symplecta</i> Meigen	(3)
<i>Trimicra</i> Osten Sacken	(1)
Eugnophomyia Alexander	19
Gnophomyia Meigen	91
Gonomyia Meigen	216
<i>Euptilostena</i> Alexander	(2)
<i>Gonomyia</i> Meigen	(55)

4.04



<i>Gonomyia</i> Alexander	(4)
<i>Idiocera</i> Dale	(2)
<i>Lipophleps</i> Bergroth	(83)
<i>Neolipophleps</i> Alexander	(17)
<i>Paralipophleps</i> Alexander	(14)
<i>Progonomyia</i> Alexander	(39)
Gymnastes Brunetti	1
<i>Paragymnastes</i> Alexander	(1)
Jivaromyia Alexander	1
Lipsothrix Loew	1
Maietta Alexander	2
Molophilus Curtis	199
<i>Eumolophilus</i> Alexander	(4)
<i>Molophilus</i> Curtis	(191)
<i>Rhynchomolophilus</i> Alexander	(1)
<i>Trichomolophilus</i> Alexander	(3)
Neognophomyia Alexander	30
Neolimnophila Alexander	1
Neophilippiana Alexander	1
Quechuamyia Alexander	1
Rhabdomastix Skuse	20
<i>Rhabdomastix</i> Skuse	(14)
<i>Sacandaga</i> Alexander	(6)
Sigmatomera Osten Sacken	19
<i>Austrolimnobia</i> Alexander	(5)
<i>Eufurina</i> Alexander	(1)
<i>Sigmatomera</i> Osten Sacken	(13)
Styringomyia Loew	5
Tasiocera Skuse	2
<i>Dasymolophilus</i> Goetghebuer	(2)
Teucholabis Osten Sacken	203
<i>Euparatropesa</i> Alexander	(14)
<i>Euteucholabis</i> Alexander	(2)
<i>Paratropesa</i> Schiner	(11)
<i>Teucholabis</i> Osten Sacken	(176)
Toxorhina Loew	34
<i>Ceratocheilus</i> Wesché	(11)
<i>Toxorhina</i> Loew	(23)
Trentepohlia Bigot	42
<i>Mongoma</i> Westwood	(1)
<i>Neomongoma</i> Alexander	(6)
<i>Paramongoma</i> Brunetti	(34)
<i>Promongoma</i> Alexander	(1)
Total Tipulidae: 3088.	Total: 1045

Subfamily TIPULINAE

Genus BRACHYPREMNA Osten Sacken

BRACHYPREMNA Osten Sacken, 1886b:161. Type-species, *Tipula dispellens* Walker (Coquillett, 1910:516, as *dispellans*, error) (Neotropical).
Ref. — Alexander, 1912g (revision).



- ABITAGUAE Alexander, 1945n:390. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- ANGUSTA Alexander, 1945e:210. Type-locality: Brazil, Rondônia, Pôrto Velho, Rio Madeira. Distr. — Brazil.
- APPENDIGERA Alexander, 1945n:393. Type-locality: Ecuador, Napo-Pastaza, Rio Jatun Yacu. Distr. — Ecuador.
- ARAJUNO Alexander, 1945a:8. Type-locality: Ecuador, Napo-Pastaza, Rio Arajuno, Napo Watershed. Distr. — Ecuador.
- ARCUARIA ARCUARIA Alexander, 1937c:481. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador.
- ssp. TRIANGULARIS Alexander, 1945g:60. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- AUSTRALIS Alexander, 1923b:185. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- BASILICA Alexander, 1921d:92. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- BREVIGENUA Alexander, 1945j:2. Type-locality: Brazil, São Paulo, Ribeirão Pires, Barreira das Camélias. Distr. — Brazil.
- BREVITEREBRA Alexander, 1944m:296, figs. 2, 4 (ovipositor). Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- BREVIVENTRIS (Wiedemann), 1821:43 (*Tipula*). Type-locality: "America." (Later Osten Sacken, 1886b:161, Brazil.) Distr. — Venezuela, Guyana, Surinam, Peru, Brazil, Bolivia. Ref. — Alexander, 1912g:230, fig. d (wing).
- CANDIDA Alexander, 1912g:233, fig. f (wing) (nom. nov. for *albimana* Wiedemann). Type-locality: "Surinam". Distr. — Panama, Surinam, Ecuador, Peru, Brazil.
- albimana* Wiedemann, 1830:615 (*Tipula*; preocc. Fabricius, 1787). Type-locality: "Surinam".
- CLYMENE Alexander, 1945e:214. Type-locality: Brazil, Guanabara, Rio de Janeiro, Alto da Boa Vista. Distr. — Brazil.
- DISPELLENS (Walker), 1861:334 (*Tipula*). Type-locality: "Mexico". Distr. — Guatemala, Costa Rica, Panama, Surinam, Venezuela, Brazil, Trinidad. Ref. — Alexander, 1912g:228, fig. b (wing).
- DIVERSIPES Alexander, 1941a:237. Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- GEJSKESI Alexander, 1945b:49. Type-locality: Surinam, Brownsberg. Distr. — Surinam.
- ILLUDENS Alexander, 1945o:579. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- INTEGRISTIGMA Alexander, 1940f:391, fig. 3 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ITATIAYANA Alexander, 1944m:293, fig. 1 (ovipositor). Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- KARMA Alexander, 1945a:11. Type-locality: Ecuador, Napo-Pastaza, Rio Arajuno, Napo Watershed. Distr. — Ecuador.
- LAETIVENTRIS Alexander, 1945g:61. Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela.
- NIROFEMORATA Alexander, 1936c:298. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.



- PHREXUS Alexander, 1953a:7, fig. 1 (wing). Type-locality: Ecuador, Manabí, Cojimies. Distr. — Ecuador.
- PICTIPES Osten Sacken, 1887b:239. Type-locality: Brazil, Rio Grande do Sul, Caçapava do Sul. Distr. — Brazil. Ref. — Alexander, 1912g:234.
- PICTIVENTRIS Alexander, 1945e:212. Type-locality: Brazil, Rondônia. Pôrto Velho, Rio Madeira. Distr. — Brazil. Ref. — Alexander, 1912g:234.
- QUASIMODO Alexander, 1943c:199. Type-locality: Ecuador, Santiago-Zamora, Río Nanganitza, Magaico. Distr. — Ecuador.
- SAPPHO Alexander, 1943f:292, fig. 1 (wing). Type-locality: Brazil, Espírito Santo, São João Petrópolis. Distr. — Brazil.
- SIMILIS Williston, 1900:229. Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico, Nicaragua, Costa Rica, Panama, Venezuela. Ref. — Alexander, 1912g:232, fig. a (wing).
- SUBEVANESCENS Alexander, 1962c:17, fig. 1 (wing). Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- SUBSIMILIS Alexander, 1921a:76. Type-locality: Argentina, Corrientes, San Roque. Distr. — Brazil, Argentina.
- SUBUNIFORMIS Alexander, 1945g:64, fig. 1 (male hypopygium). Type-locality: Venezuela, Bolívar, Akuriman, Gran Sabana. Distr. — Venezuela.
- THYESTES Alexander, 1953e:599, fig. 1 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- UNICOLOR Osten Sacken, 1887b:239. Type-locality: "Puerto Rico". Distr. — Cuba, Haiti, Dominican Republic, Jamaica, Puerto Rico, Grenada. Refs. — Alexander, 1912g:235, fig. c (wing); 1932:354, fig. 2 (wing); 1964a:14, fig. 2 (wing).
- UNIFORMIS Alexander, 1920a:75. Type-locality: Brazil, Amazonas, Parintins. Distr. — Brazil, Peru.
- VARIANTIBIA Alexander, 1936c:300. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- WILLIAMSONI Alexander, 1912g:231, fig. e (wing). Type-locality: Guyana, Wismar. Distr. — Guyana, Surinam, Peru.

Genus CTENOPHORA Meigen

- CTENOPHORA Meigen, 1803:263. Type-species, *Tipula pectinicornis* Linnaeus (Westwood, 1840:128) (Western Palaearctic). (See note in Alexander, 1965d:19).
- Flabellifera* authors, not Meigen.

Subgenus PSELLIOPHORA Osten Sacken

- PSELLIOPHORA Osten Sacken, 1886b:165. Type-species, *Tipula laeta* Fabricius (Enderlein, 1912:21; Brunetti, 1912:290) (Oriental).
- MARGARITA Alexander, 1965a:401. Type-locality: Mexico, Durango, 48 miles west of El Salto. Distr. — Mexico.
- MESAMERICANA (Alexander), 1944e:92 (*Pseliophora*). Type-locality: Guatemala, Cayuga. Distr. — Guatemala.
- VENEZUELENSIS (Alexander), 1944c:38, fig. 2 (wing) (*Pseliophora*). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.



Genus DOLICHOPEZA Curtis

DOLICHOPEZA Curtis, 1825: vol. 2, pl. 62. Type-species, *sylicola* Curtis (orig. des.) = *albipes* (Ström) (Wertern Palaearctic).

Subgenus MEGISTOMASTIX Alexander

- MEGISTOMASTIX Alexander, 1912a:63 (as genus). Type-species, *portoricensis* Alexander (mon.) (Neotropical).
- ACUTILOBA Alexander, 1937a:179, fig. 6 (male hypopygium). Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- CUBENSIS (Alexander), 1928b:47 (*Megistomastix*). Type-locality: Cuba, Pinar del Río, Sierra Rangel. Distr. — Cuba.
- DARLINGTONI Alexander, 1939d:105, fig. 11 (male hypopygium). Type-locality: Cuba, Pico Turquino, Sierra Maestra. Distr. — Cuba.
- DEVEXA Alexander, 1937e:523, figs. 1 (wing), 9 (male hypopygium). Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- DOMINGENSIS Alexander, 1939d:106, figs. 2 (wing), 12 (male hypopygium). Type-locality: Hispaniola: Dominican Republic, Loma Rucilla. Distr. — Dominican Republic.
- OBTUSILOBA Alexander, 1937a:180, fig. 7 (male hypopygium). Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- PORTORICENSIS (Alexander), 1912a:63, figs. 1 (head), 2 (wing), 3 (male hypopygium) (*Megistomastix*). Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico. Ref. — Alexander, 1932:355, fig. 3 (wing).
- VITTI NERVIS Alexander, 1937a:181, fig. 1 (wing). Type-locality: Cuba, Oriente, Sierra del Cobre, Loma del Gato. Distr. — Cuba.

Genus ELNORETTA Alexander

- ELNORETTA Alexander, 1929a:15. Type-species, *acracanthoides* Alexander (orig. des.) (Neotropical).
- ACRACANTHODES Alexander, 1929a:16, figs. 31 (wing), 111 (antenna). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.

Genus ISCHNOTOMA Skuse

- ISCHNOTOMA Skuse, 1890b:114. Type-species, *Tipula serricornis* Macquart (Alexander, 1920g:53) — *eburnea* Walker (Australasian). Ref. — Vane-Wright, 1967, figs.

Subgenus ICRIMASTAX Enderlein

- ICRIMASTAX Enderlein, 1912:9 (as genus). Type-species, *ocellata* Enderlein (orig. des.) (Neotropical). Ref. — Vane-Wright, 1967.
- ANTINYMPHA (Alexander), 1942i:48, fig. 13 (wing) [*Holorusia* (*Holorusia*)]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- CALLIPE (Alexander), 1945j:8, fig. 2 (male hypopygium) [*Holorusia* (*Holorusia*)]. Type-locality: Brazil, São Paulo, Guarujá. Distr. — Brazil.
- EUTERPE (Alexander), 1945e:230 [*Holorusia* (*Holorusia*)]. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.

- HELIOS (Alexander), 1949b:467, fig. 7 (male hypopygium) (*Holorusia*). Type-locality: Brazil, Paraná, Curitiba. Distr. — Brazil.
- JUJUYENSIS (Alexander), 1920f:220 (*Holorusia*). Type-locality: Argentina, Jujuy. Distr. — Argentina. (May be *nudicornis*, auct. Vane-Wright, 1967:516, fig. 8 (wing); page 529, fig. 43 (male hypopygium); Bruch, 1939a:12-16, pls. 2 (habitat); 3, fig. 7 (head, adult); 7, fig. 26 (wing), figs. 27, 28 (immature stages); 8, figs. 1-16 (immature stages).
- NITRA (Alexander), 1945c:227, fig. 1 (male hypopygium) [*Holorusia* (*Holorusia*)]. Type-locality: Brazil, Rio de Janeiro, Serra dos órgãos. Distr. — Brazil.
- NUDICORNIS (Macquart, 1838a:59(55) (*Tipula*). Type-locality: Argentina. Buenos Aires. Distr. — Argentina.
- OCELLATA (Enderlein), 1912:9 (*Ieriomastax*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil. Ref. — Alexander, 1926c:388.
- PHAETON (Alexander), 1945j:10, fig. 3 (male hypopygium) [*Holorusia* (*Holorusia*)]. Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantarcira. Distr. — Brazil.
- ZIKANI (Alexander), 1936a:14 (*Holorusia*). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.

Subgenus ISCHNOTOMA Skuse

- CONCINNA (Philippi), 1865:605 (*Tipula*). Type-locality: Chile, Valdivia, Port Corral. Distr. — Chile. Ref. — Alexander, 1929a:22.
- DECORATA DECORATA (Philippi), 1865:604 (*Tipula*). Type-locality: Chile, Valdivia, Santiago, Illapel. Distr. — Chile. Ref. — Alexander, 1929a:30, fig. 6 (wing).
- ssp. ARAUCANA (Alexander), 1929a:30 (footnote) (*Holorusia*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile. Ref. — Vane-Wright, 1967:514, fig. (wing), page 526, fig. (male hypopygium).
- DLPONTEI (Alexander), 1929a:27, figs. 4 (wing), 115 (male hypopygium) (*Holorusia*). Type-locality: Chile, Llanquihue, Ensenada. Distr. — Chile, Argentina.
- FAGETORUM FAGETORUM (Alexander), 1929a:29, figs. 5 (wing), 117 (male hypopygium) (*Holorusia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina. Ref. — Alexander, 1962b:337, fig. 8 (male hypopygium).
- ssp. TRUNCULATA (Alexander), 1962b:340, fig. 7 (male hypopygium) (*Holorusia*). Type-locality: Argentina, Neuquén, Lago Lacar. Distr. — Argentina.
- FRAUENFELDI (Schiner), 1868:36 (*Tipula*). Type-locality: "Chile". Distr. — Chile. (May be *decorata*). Ref. — Alexander, 1929a:23.
- FUSCOSTIGMOSA (Alexander), 1929a:32 (*Holorusia*). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- IIOMOCHROA (Alexander), 1926c:391, figs. 1, 2 (male hypopygium) (*Holorusia*). Type-locality: Chile, Malleco, Victoria. Distr. — Chile. Ref. — Alexander, 1929a:23.
- LAROTYPA (Alexander), 1929a:31, fig. 7 (wing) (*Holorusia*). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.



- PENAI (Alexander), 1952b:99, figs. 1 (male hypopygium), 5 (wing) [*Holorusia (Ischnotoma)*]. Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- guzmani* Alexander, 1953b:8 [*Holorusia (Ischnotoma)*] (lapsus for *penai*).
- PORTERI (Alexander), 1929a:24, fig. 33 (wing) (*Holorusia*). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile. Ref. — Vane-Wright, 1967:526, fig. 39 (male hypopygium), page 528, fig. 42 (male hypopygium).
- POSTNOTALIS (Alexander), 1929a:28 (*Holorusia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina. Ref. — Vane-Wright, 1967:526, fig. 39 (male hypopygium), page 528, fig. 42 (male hypopygium).
- PROBLEMATICA (Alexander), 1945f:8, figs. 5 (antenna), 7 (wing) (*Holorusia*). Type-locality: Chile, Ñuble, Guapé, near Chillán. Distr. — Chile.
- RUFPOSTIGMOSA (Macquart), 1838a:60(56) (*Tipula*). Type-locality: "Chile". Distr. — Chile. Ref. — Alexander, 1929a:22. (May be prior name of *decorata*).
- SCHINERIANA (Alexander), 1928f:635 (*Holorusia*). Type-locality: Chile, Valparaíso, Marga-Marga. Distr. — Chile.
- SHANNONIANA (Alexander), 1929a:27, fig. 116 (male hypopygium) (*Holorusia*). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile. Ref. — Vane-Wright, 1967:517, fig. 11 (wing).
- SILVAI (Alexander), 1929a:26, fig. 114 (male hypopygium) (*Holorusia*). Type-locality: Chile, Llanquihue, Puerto Montt. Distr. — Chile.
- VITTIGERA (Philippi), 1865:607 (*Tipula*). Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1929a:23.

Subgenus NEOTIPULA Alexander

- Tipula*, subg. NEOTIPULA Alexander, 1940c:105. Type-species, *Tipula pectinella* Alexander (orig. des.) (Neotropical).
- MAYA (Alexander), 1912f:358, figs. a (wing), t (ovipositor) (*Tipula*). Type-locality: Guatemala, Aguna. Distr. — Guatemala.
- PAPRZYCKII (Alexander), 1941a:238 [*Tipula (Neotipula)*]. Type-locality: Peru, Junín, Satipo. Distr. — Peru. Ref. — Alexander, 1951b:299, fig. 17 (wing).
- PECTINELLA (Alexander), 1940c:106 [*Tipula (Neotipula)*]. Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Panama.
- PENATA (Alexander), 1965b:337 [*Holorusia (Holorusia)*]. Type-locality: Ecuador, Quito. Distr. — Ecuador, Peru.

Genus LEPTOTARSUS Guérin-Ménéville

- LEPTOTARSUS Guérin-Ménéville, 1838:286, pl. 20, fig. 1. Type-species, *macquarti* Guérin-Ménéville (mon.) (Australasian).

Subgenus ALDROVANDIA Enderlein

- ALDROVANDIA Enderlein, 1912:52 (as genus). Type-species, *gesneri* Enderlein (orig. des.) (Neotropical).

GESNERI (Enderlein), 1912:52, fig. F¹ (wing) (*Aldrovandia*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil. Ref. — Alexander, 1936c:303, fig. 1 (wing).

Subgenus ARAUCOMYIA Alexander

Macromastix, subg. ARAUCOMYIA Alexander, 1929a:14. Type-species, *paulseni* (Philippi) (orig. des.) (Neotropical).

BREVIHIRSUTUS (Alexander), 1934d:174 [*Macromastix (Araucomyia)*]. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.

PAULSENI (Philippi), 1865:606 (*Tipula*). Type-locality: Chile, Santiago. Distr. — Chile. Ref. — Alexander, 1929a:15, fig. 30 (wing).

PENITUS (Alexander), 1944i:118 [*Macromastix (Araucomyia)*]. Type-locality: Chile, Arauco, Cerros de Nahuelbuta. Distr. — Chile.

Subgenus CEONEURA Alexander

Tanyremna, subg. CEONEURA Alexander, 1924b:289. Type-species, *idioneura* Alexander (orig. des.) (Neotropical).

IDIONEURUS (Alexander), 1924b:289 [*Tanyremna (Ceoneura)*]. Type-locality: Guyana, Kartabo. Distr. — Guyana. Ref. — Alexander, 1936c:303, fig. 4 (wing).

Subgenus LEPTOTARSUS Guérin-Ménéville

ANNULIPES (Philippi), 1865:605 (*Tipula*). Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1929a:21, fig. 3 (wing).

Subgenus LIMONIODES Alexander

Tanyremna, subg. LIMONIODES Alexander, 1938b:337. Type-species, *sulphurella* (Alexander) (orig. des.) (Neotropical).

SULPHURELLUS (Alexander), 1938b:337, fig. 1 (wing) [*Tanyremna (Limoniodes)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

Subgenus LONGURIO Loew

LONGURIO Loew, 1869:3 (Cent. 8, no. 2) (as genus). Type-species, *testaceus* Loew (mon.) (Nearctic).

ATRIROSTRIS (Alexander), 1945f:5, figs. 1, 2 (male hypopygium) (*Macromastix*). Type-locality: Chile, Nuevo Imperial. Distr. — Chile.

BRASILIAE (Alexander), 1935a:313 [*Macromastix (Macromastix)*]. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.

BROWNI (Alexander), 1940d:194 (*Macromastix*). Type-locality: Ecuador, Napo-Pastaza, Río Upano, Macas. Distr. — Ecuador.

BULLOCKI (Alexander), 1931c:622 [*Macromastix (Araucomyia)*]. Type-locality: Chile, Malleco, Nahuelbuta. Distr. — Chile, Argentina. Ref. — Alexander, 1951a:82, figs. 1 (male hypopygium), 2 (wing).



- CAPARAONUS (Alexander), 1944m:298 [*Macromastix (Macromastix)*]. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- CARRERANUS (Alexander), 1945j:3 (*Macromastix*). Type-locality: Brazil, São Paulo, Guarujá. Distr. — Brazil.
- CHRYSOSTIGMA (Alexander), 1944m:300 [*Macromastix (Macromastix)*]. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- ESPINOZAI (Alexander), 1938i:276 [*Macromastix (Macromastix)*]. Type-locality: Chile, Cautín, Temuco. Distr. — Chile.
- EUCRYPTA (Alexander), 1953b:7, fig. 3 (wing) [*Longurio (Longurio)*]. Type-locality: Chile, Chiloé Is., Aulén. Distr. — Chile.
- EXEMPTUS (Alexander), 1936a:10 (*Habromastix*). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- GOYAZANUS (Alexander), 1940f:386, fig. 11 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- GULMARAESI (Alexander), 1942i:46, figs. 12 (wing), 16 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- GYMNOCERA (Alexander), 1938c:321 (*Macromastix*). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Maromba. Distr. — Brazil.
- HUANUCENSIS (Alexander), 1953e:613, fig. 7 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Peru, Huanuco, Pillao. Distr. — Peru.
- INSIDIOSUS (Alexander), 1949b:463, figs. 3 (wing), 5 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- IXION (Alexander), 1949b:464, figs. 2 (wing), 6 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- LEMNISCATUS (Alexander), 1929b:91 (*Habromastix*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- LUSTRALIS (Alexander), 1936d:750 (*Tipula*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LUTEISTIGMA (Alexander), 1940f:389, fig. 12 (male hypopygium) [*Macromastix (Araucomyia)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NAHUEL BUTAE (Alexander), 1945f:7, figs. 3, 4 (male hypopygium) (*Macromastix*). Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- NOCIVUS (Alexander), 1953b:5, figs. 4 (wing), 5 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Chile, Chiloé Is., Toi-Goi. Distr. — Chile.
- PARAGUAYANUS (Riedel), 1921:126, fig. (wing) (*Tipulodina*). Type-locality: Paraguay, Pikade. Distr. — Paraguay.
- PARAGUAYENSIS (Alexander), 1935a:315 (*Tipula*). Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.

- PERGLABRATUS (Alexander), 1953e:614, fig. 8 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- PHIABEDRA (Alexander), 1949b:461, figs. 1 (wing), 4 (male hypopygium) (*Macromastix*). Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- PYGMAEUS (Alexander), 1914f:351, pl. 15, fig. 1 (wing) (*Macromastix*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- RABELLOI (Alexander), 1953c:615, fig. 9 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- SEROTINELLUS (Alexander), 1928e:245 (as 1927) (*Macromastix*). Type-locality: Chile, Valdivia, Panguipulli. Distr. — Chile. Ref. — Alexander, 1929a:14, fig. 29 (wing).
- SESSORIS (Alexander), 1953e:612, fig. 6 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- STENOSTYLA (Alexander), 1953a:8, figs. 2 (wing), 4 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Ecuador, Tungurahua, Río Blanco near Baños. Distr. — Ecuador.
- STYX (Alexander), 1954a:31, figs. 1 (wing), 3 (male hypopygium) [*Longurio (Longurio)*]. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- TIJUCANUS (Alexander), 1943f:294, figs. 2 (wing), 4 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, Guanabara, Rio de Janeiro, Pico da Tijuca. Distr. — Brazil.
- TRAVASSOSANUS (Alexander), 1942i:45, figs. 11 (wing), 15 (male hypopygium) [*Macromastix (Macromastix)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- ZIKANELLUS (Alexander), 1953e:616 [*Longurio (Longurio)*]. Type-locality: Brazil, Rio de Janeiro, São Gonçalo, Barracão. Distr. — Brazil.

Subgenus PEHLKEA Enderlein

- PEHLKEA Enderlein, 1912:15 (as genus). Type-species, *columbiana* Enderlein (orig. des.) (Neotropical).
- COLUMBIANUS (Enderlein), 1912:15 (*Pehlkea*). Type-locality: "Colombia". Distr. — Colombia. Ref. — Alexander, 1914d:213, fig. 3 (wing).
- PALLITARSIS (Alexander), 1937c:483 [*Tanypremna (Pehlkea)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador.
- REGINA (Alexander), 1914d:214, fig. 4 (wing) (*Tanypremna*). Type-locality: Colombia, Cordillera Central. Distr. — Colombia.
- REGULUS (Alexander), 1943b:104 [*Tanypremna (Pehlkea)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

Subgenus TANYPREMNA Osten Sacken

- TANYPREMNA Osten Sacken, 1886a:19 (as genus). Type-species, *opilio* Osten Sacken (mon.) (Neotropical).
- Stegasmaonotus* Enderlein, 1912:11. Type-species, *longissimus* Enderlein (orig. des.).

- AURANTIOTHORAX (Alexander), 1953e:608 [*Longurio (Tanypremna)*]. Type-locality: Peru, Loreto, Contamana, Cerro Azul. Distr. — Peru.
- BEZZIANUS (Alexander), 1921d:99 (*Tipula*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- BORGMEIERI (Alexander), 1942g:167, figs. 1 (wing), 7 (male hypopygium) (*Tanypremna*). Type-locality: Brazil, Guanabara, Rio de Janeiro, Jardim Botânico. Distr. — Brazil.
- CALLOPE (Alexander), 1945c:243 (*Tanypremna*). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- CARBONIPES (Alexander), 1938b:342, fig. 3 (wing) (*Tanypremna*). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CERRITA (Alexander), 1945e:215 (*Tanypremna*). Type-locality: Brazil, Rondônia, Guajará-Mirim, Vila Murinho. Distr. — Brazil.
- CLOTHO (Alexander), 1944h:370 (*Tanypremna*). Type-locality: Brazil, Rio de Janeiro, Palmeiras. Distr. — Brazil.
- ELEGANTIOR (Alexander), 1945b:51 (*Tanypremna*). Type-locality: "Colombia". Distr. — Colombia.
- FIELDIANUS (Alexander), 1956a:22 [*Longurio (Tanypremna)*]. Type-locality: Panama, Canal Zone. Distr. — Panama.
- FUSCITARSIS (Alexander), 1919a:149, fig. 16 (wing) (*Tanypremna*). Type-locality: Colombia, Magdalena, Mt. Santa Marta. Distr. — Colombia.
- HODGEI (Alexander), 1939d:92, figs. 1 (wing), 10 (male hypopygium) (*Tanypremna*). Type-locality: Lesser Antilles: Dominica, Morne Trois Pitons. Distr. — Dominica.
- HORRIDUS (Alexander), 1941a:235 (*Tanypremna*). Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- INCOMPLETA (Alexander), 1944L:172, fig. 3 (male hypopygium) (*Tanypremna*). Type-locality: Venezuela, Miranda, Naiguata. Distr. — Venezuela.
- INVARIPES (Alexander), 1936d:749 (*Tanypremna*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- KADENI (Alexander), 1941a:232 (*Tanypremna*). Type-locality: "Venezuela". Distr. — Venezuela.
- LONGIPES (Fabricius), 1805:25 (*Tipula*). Type-locality: "South America." Distr. — Colombia, Guyana, Peru, Brazil. Ref. — Alexander, 1914d:208, fig. 1 (wing).
- LONGISSIMUS (Enderlein), 1912:11 (*Stegasmonotus*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil. Ref. — Alexander, 1914d:209.
- MANICATUS (Osten Sacken), 1887b:240 (*Tanypremna*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Alexander, 1914d:211.
- MIRANDA (Alexander), 1944j:146, fig. 4 (male hypopygium) (*Tanypremna*). Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela.
- OPILO (Osten Sacken), 1886a:19, pl. 1, fig. 2 (*Tanypremna*). Type-locality: Guatemala, Verapaz, Purula. Distr. — Panama, Guatemala, Venezuela. Ref. — Alexander, 1914d:211, fig. 2 (wing).

- ORNATIPES (Alexander), 1941a:233 (*Tanypremna*). Type-locality: "Brazil".
Distr. — Brazil.
- PERORNATUS (Alexander), 1929a:90 (*Tanypremna*). Type-locality: Venezuela, Táehira, La Fría. Distr. — Venezuela, Eeuador, Peru.
- PHYLAX (Alexander), 1953e:610 [*Longurio (Tanypremna)*]. Type-locality: Peru, Huanueo, Fundo Sinehono, Cordillera Azul. Distr. — Peru.
- PICTURELLUS (Alexander), 1941d:105 (*Tanypremna*). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Eeuador.
- PORTERIANUS PORTERIANUS (Alexander), 1936e:304, fig. 2 (wing) (*Tanypremna*). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- ssp. LIMAI (Alexander), 1945e:245 (*Tanypremna*). Type-locality: Brazil, São Paulo, São Paulo, Sacomã. Distr. — Brazil.
- PROAVITA (Alexander), 1940f:390, figs. 2 (wing), 13 (male hypopygium) (*Tanypremna*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SALOME (Alexander), 1945j:6, fig. 1 (male hypopygium) (*Tanypremna*). Type-locality: Brazil, São Paulo, Guarujá. Distr. — Brazil.
- SALTATRIX (Alexander), 1944h:369 (*Tanypremna*). Type-locality: Costa Rica, Talamanca, Rivas. Distr. — Costa Rica.
- SANCTAECATHARINAE (Alexander), 1945e:217 (*Tanypremna*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- UNIGUTTATUS (Alexander), 1953e:611 [*Longurio (Tanypremna)*]. Type-locality: Peru, Huanueo, Fundo Sinehono, Cordillera Azul, Divisoria. Distr. — Peru.

Subgenus TANYPREMNELLA Alexander

- Tanypremna*, subg. TANYPREMNELLA Alexander, 1938b:339. Type-species, *crystallina* Alexander (orig. des.). (Neotropical).
- ANTENNIFERUS (Alexander), 1942b:343, fig. 2 (wing) [*Tanypremna (Tanypremnella)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- CRYSTALLINA (Alexander), 1938b:340, fig. 2 (wing) [*Tanypremna (Tanypremnella)*]. Type-locality: Eeuador, Napo-Pastaza, Abitagua. Distr. — Eeuador.
- GENTILIS (Alexander), 1943e:18 [*Tanypremna (Tanypremnella)*]. Type-locality: Venezuela, Bolívar, Akuriman, Gran Sabana. Distr. — Venezuela.
- MALDONADOI (Alexander), 1953d:147 [*Longurio (Tanypremnella)*]. Type-locality: Venezuela, Pacaraima Mts., Camp Benítez near foot of Mt. Marahuaca. Distr. — Venezuela.
- MEDIOCORNIS (Alexander), 1944d:300 [*Tanypremna (Tanypremnella)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- MEGACERA (Alexander), 1943b:105 [*Tanypremna (Tanypremnella)*]. Type-locality: Eeuador, Napo-Pastaza, Abitagua. Distr. — Eeuador.
- MICROCERA (Alexander), 1943b:106 [*Tanypremna (Tanypremnella)*]. Type-locality: Eeuador, Napo-Pastaza, Abitagua. Distr. — Eeuador.
- PERDISTINCTUS (Alexander), 1942b:341, fig. 1 (wing) [*Tanypremna (Tanypremnella)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.



- SEGNIPES (Alexander), 1953d:148 [*Longurio (Tanypremnella)*]. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- TRANSFASCIATUS (Alexander), 1938b:342 [*Tanypremnna (Tanypremnella)*]. Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador.

Subgenus TANYPREMNODES Alexander

- Tanypremnna*, subg. TANYPREMNODES Alexander, 1924b:290. Type-species, *leucoplaca* Alexander (orig. des.) (Neotropical).
- LEUCOPLACA (Alexander), 1924b:291 [*Tanypremnna (Tanypremnodes)*]. Type-locality: Brazil, Pará, Upper Rocana (?). Distr. — Brazil. Ref. — Alexander, 1936c:303, fig. 3 (wing).
- SUBAPICALIS (Alexander), 1941a:236 [*Tanypremnna (Tanypremnodes)*]. Type-locality: Brazil, Santa Catarina, Blumenau. Distr. — Brazil.

Subgenus XENOTIPULA Alexander

- XENOTIPULA Alexander, 1921b:171 (as genus). Type-species, *munroi* Alexander (orig. des.) (Ethiopian).
- CISATLANTICUS (Alexander), 1937c:485 (*Xenotipula*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus MEGISTOCERA Wiedemann

- MEGISTOCERA Wiedemann, 1820:41(1821:41) (as *Maekistocera*, 1828:55). Type-species, *Tipula filipes* Fabricius (Macquart, 1838a:63; 1838:59) (Ethiopian), (See note in Alexander, 1965d:18).
- LONGIPENNIS (Macquart), 1838a:61(57), pl. 5, fig. 1 (whole insect) (*Tipula*). Type-locality: "Cuba". Distr. — Southeastern U. S. A., Guyana, Peru, Brazil, Bolivia, Argentina, Paraguay, Cuba, Jamaica, Puerto Rico, Trinidad. Refs. — Alexander, 1914d:216; 1932:354, fig. 1 (wing); 1964a:13, fig. 1 (wing).
- tenuis* Wulp, 1885:7 (*Tipula*). Type-locality: "Surinam". Distr. — Surinam. Ref. — Neebham, 1908, pl. 16, fig. 2 (wing).

Genus NEPHROTOMA Meigen

- Pales* Meigen, 1800:14. Type-species, *Tipula dorsalis* Fabricius (Hendel, 1908:46). Suppressed by I.C.Z.N., 1963:339.
- NEPHROTOMA Meigen, 1803:262. Type-species, *Tipula dorsalis* Fabricius (mon.) (Western Palaearctic).
- Pachyrhina* Macquart, 1834:88. Type-species, *Tipula crocata* Linnaeus (Westwood, 1840:128).
- Pachyrhina*, *Pachyrina*, *Pachyrhyna*, emends. or errors.
- AFFINIS (Bellardi), 1859:10(1861:210) (*Tipula*). Type-locality: "Mexico". Distr. — Mexico. Identity uncertain.
- ALLENI (Alexander), 1913c:206, pl. 7, fig. 5 (wing) (*Pachyrhina*). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- BOLIVIANA Alexander, 1947a:174. Type-locality: Bolivia, Cochabamba, Chapore. Distr. — Bolivia.

- CACUMINIS Alexander, 1945p:753. Type-locality: Ecuador. El Oro, Morro Morro, Pinas. Distr. — Ecuador.
- CIRCUMSCRIPTA (Loew), 1863:180 (*Pachyrrhina*). Type-locality: "Cuba". Distr. — Cuba, Jamaica. Ref. — Alexander, 1964a:14, fig. 3 (wing).
- CONSULARIS CONSULARIS (Osten Sacken), 1886a:17 (*Pachyrrhina*). Type-locality: Costa Rica, Cache. Distr. — Nicaragua, Costa Rica, Venezuela, Peru, Brazil, Argentina, Paraguay.
- ssp. EMINENS Alexander, 1944j:149. Type-locality: Venezuela, Mérida, San Diego. Distr. — Venezuela.
- DAMPFI Alexander, 1925a:358, fig. 12 (wing). Type-locality: Mexico. Mt. Tlaloc. Distr. — Mexico.
- DURANGENSIS Alexander, 1953d:149. Type-locality: Mexico, Durango, 10 miles east of El Salto. Distr. — Mexico.
- ELEGANS (Fabricius), 1805:26 (*Tipula*). Type-locality: "South America." (Identity not known).
- ELEGANTULA (Williston), 1896a:295, pl. 10, fig. 70 (wing) (*Pachyrrhina*). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.
- FERRUGINEA (Fabricius), 1805:28 (*Tipula*). Type-locality: "North America". Distr. — North America.
- quadrilineata* Macquart, 1838a:54(50) (*Pachyrrhina*). Type-locality: "Mexico" (auct. Osten Sacken). Apparently correct.
- proxima* Bellardi, 1859:9 (1861:209) (*Tipula*). Type-locality: "Mexico" (auct. Osten Sacken). Apparently correct.
- GLABRICRISTATA Alexander, 1939d:102. Type-locality: Cuba, Oriente, Sierra del Cobre, Loma del Gato. Distr. — Cuba, Haiti, Dominica.
- LATEROPOLITA Alexander, 1938h:432. Type-locality: Costa Rica, San Pedro de Montes de Oca. Distr. — Costa Rica.
- MEDIOLIGULA Alexander, 1945g:67, fig. 2 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Colombia, Venezuela.
- MELANOXANTHA Alexander, 1962c:36, fig. 5 (wing). Type-locality: Bolivia, Yungas de Arcpucho, Sihuencos. Distr. — Bolivia.
- MEXICANA (Macquart), 1846:140 (1846:12) (*Pachyrrhina*). Type-locality: "Mexico". Distr. — Mexico.
- NIGROLUTEA (Bellardi), 1859:11 (*Tipula*). Type-locality: "Mexico". Distr. — Mexico, Costa Rica. Supplementary notes, Osten Sacken, 1886a:16.
- ORDINARIA (Osten Sacken), 1886a:16 (*Pachyrrhina*). Type-locality: Mexico, Durango, Durango. Distr. — Mexico.
- PULCHELLA (Rondani), 1850:195 (*Pachyrrhina*). Type-locality: Brazil, São Paulo, Ilha de São Sebastião. Distr. — Brazil.
- PUNCTIFRONS (Macquart), 1838a:53(49) (*Pachyrrhina*). Type-locality: Brazil, "campos gerais, southern part". Distr. — Brazil, Argentina. Identity uncertain.
- USTA (Osten Sacken), 1886a:17 (*Pachyrrhina*). Type-locality: Costa Rica, Mt. Irazu, Río Sucio. Distr. — Costa Rica.

Genus OZODICERA Macquart

- OZODICERA Macquart, 1834:92. Type-species, *ochracea* Macquart (mon.) = *pectinata* (Wiedemann) (Neotropical).
- Hemictaina* Westwood, 1835c:450. Type-species, *gracilis* Westwood (mon.).

Subgenus DIHEXACLONUS Enderlein

- DIHEXACLONUS Enderlein, 1912:28 (as genus). Type-species, *Ozodicera apicalis* Macquart (orig. des.) (Neotropical).
- APICALIS Macquart, 1838a:52(48), pl. 4, fig. 1 (whole insect). Type-locality: Brazil, Western Minas Gerais. Distr. — Brazil.
- BLACULEATA Alexander, 1953e:604. Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- EFFECTA Alexander, 1944f:19. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- FUMIPENNIS Loew, 1851:386, fig. 4 (antenna). Type-locality: "Brazil". Distr. — Brazil.
- GRACILIRAMA Alexander, 1941a:231. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.
- GUIANENSIS Alexander, 1934b:303, figs. 11 (wing), 12, 13 (male hypopygium). Type-locality: Guyana, Bartica District, Kartabo. Distr. — Guyana.
- JESSEANA Alexander, 1945e:219. Type-locality: Brazil, Rondônia, Guajará-Mirim, Vila Murtinho. Distr. — Brazil.
- LANEI Alexander, 1942i:44, figs. 3 (wing), 5 (antenna), 10 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- LONGISECTOR Alexander, 1944c:36, fig. 1 (wing). Type-locality: Venezuela, Bolívar, Surukam, Caroni River. Distr. — Venezuela.
- MACRACANTHA Alexander, 1940f:384, fig. 10 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NEIVAI Alexander, 1940f:383, figs. 1 (wing), 8 (antenna), 9 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PANAMENSIS Alexander, 1922c:63. Type-locality: Panama, Río Chico. Distr. — Panama.
- PERFUGA Alexander, 1944f:20. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- PUMILA Alexander, 1942h:424, fig. 2 (antenna). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- SPILOPHAEA Alexander, 1942b:338, fig. 3 (male hypopygium). Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- SUPERARMATA Alexander, 1942h:422, figs. 1 (antenna), 7, 8 (male hypopygium). Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil.
- TERRIFICA Alexander, 1942i:40, figs. 1 (wing), 4 (antenna), 8 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- TRIGUTTATA Alexander, 1921d:94. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil. Ref. — Alexander, 1934b:303, figs. 14, 15 (male hypopygium).
- TRIPALLENS Alexander, 1942i:42, figs. 2 (wing), 6, 7 (antenna), 9 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- UMBRIFERA Alexander, 1937b:236. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- XANTHOSTOMA Loew, 1851:387, fig. 5 (antenna). Type-locality: "Brazil". Distr. — Brazil.



Subgenus OZODICERA Macquart

- ATTENUATA Alexander, 1920b:8. Type-locality: Brazil, Pará, Prata. Distr. — Brazil.
- BIMACULATA Enderlein, 1912:27. Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- BISPINIFER Alexander, 1921d:96. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- CARRERELLA Alexander, 1956a:23. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- CAUDIFERA Alexander, 1945j:12, fig. 4 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- CINEREIPENNIS Alexander, 1937b:235. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- CORRIENTESANA Alexander, 1962b:336 figs. 1 (wing), 6 (male hypopygium). Type-locality: Argentina, Corrientes, Manantiales. Distr. — Argentina.
- CYGNIFORMIS Alexander, 1953a:10, figs. 3 (wing), 5 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Puyo. Distr. — Ecuador.
- DUIDENSIS Alexander, 1931b:2, figs. 1 (wing), 11, 12 (male hypopygium). Type-locality: Venezuela, Mount Duida. Distr. — Venezuela.
- ELIANA Alexander, 1953e:605. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- EPICOSMA Alexander, 1926d:386. Type-locality: Brazil, Espírito Santo, Pôrto do Cadi. Distr. — Brazil.
- EURYSTYLA Alexander, 1953e:607, fig. 5 (male hypopygium). Type-locality: Peru, Loreto, Cachiyacu, Upper Ucayali River. Distr. — Peru.
- EXTENSA Alexander. 1921d:95. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- GRACILIS (Westwood), 1835c:450, figs. 16, 17 (*Hemictena*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Westwood, 1881:380, pl. 18, fig. 8 (antenna, wing).
- GRISEIPENNIS Loew, 1851:389, figs. 6-8. Type-locality: "Brazil". Distr. — Brazil.
- LONGIMANA (Fabricius), 1805:26 (*Tipula*). Type-locality: "South America". Distr. — South America.
- MULTIERMIS Alexander, 1942d:135. Type-locality: Ecuador, Napo-Pastaza, Rio Jatun Yaeu. Distr. — Ecuador. Ref. — Alexander, 1953a:14, fig. 6 (male hypopygium).
- NIROMARGINATA Alexander, 1938c:319. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- NOCTIVAGANS Alexander, 1914e:253. Type-locality: Guyana, Bartica. Distr. — Guyana.
- PECTINATA (Wiedemann), 1828:47 (*Tipula*). Type-locality: "Guyana". Distr. Guyana.
ochracea Macquart, 1834:92 (unnecessary nom. nov. for *pectinata* Wiedemann). Type-locality: "South America".
- PHALACANTHA Alexander, 1942b:340, fig. 4 (male hypopygium). Type-locality: Peru, Huanuco, Leonpampa. Distr. — Peru.
- PIATRIX Alexander, 1944d:299. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.



- PLACATA Alexander, 1965b:339, fig. 1 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- SCHWARZMAIERANA Alexander, 1942h:426, figs. 4, 5 (antenna). Type-locality: Brazil, São Paulo, Pindamonhangaba. Distr. — Brazil.
- SEPTEMTRIONIS Alexander, 1940b:140. Type-locality: Mexico, Chiapas, Finca San Cristóbal. Distr. — Mexico.
- SIMPLEX (Walker), 1856:446 (*Ptilogyna*). Type-locality: "South America." Distr. — South America.
- STRIATIPENNIS Alexander. 1941b:139. Type-locality: Venezuela, Antimano. Distr. — Venezuela.
- STROHMI Alexander, 1945e:21. Type-locality: Brazil, Rondônia, Guajará-Mirim, Vila Murtinho. Distr. — Brazil.
- SUBVITTATA Alexander, 1938c:320. Type-locality: Brazil, Rio de Janeiro, Campo Belo. Distr. — Brazil.
- THAUMASTA Alexander, 1953e:608. Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- FRISPINIFER Alexander, 1940a:275. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador. Ref. — Alexander, 1953a:14, fig. 7 (male hypopygium).
- WITTEANA Alexander, 1942h:428, figs. 6 (antenna), 9, 10 (male hypopygium). Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil.
- ZIKANIANA Alexander, 1937b:234. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.

Genus PHACELODOCERA Enderlein

- PHACELODOCERA Enderlein, 1912:26. Type-species, *Ptilogyna flabellifera* Loew (orig. des.) (Neotropical).
- FLABELLIFERA (Loew), 1851:391, figs. 1-3 (venation, antenna, wing) (*Ptilogyna*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Alexander, 1953e:600, figs. 2 (antenna), 3 (wing).

Genus TIPULA Linnaeus

- TIPULA Linnaeus, 1758:585. Type-species, *oleracea* Linnaeus (Latreille, 1810:442) (Western Palearctic). Refs. — Alexander, 1945k, 1945m, 1946d, 1951b, 1965c.

Subgenus BELLARDINA Edwards

- Tipula*, subg. BELLARDINA Edwards, 1931:82. Type-species, *craverii* Bellardi (orig. des.; as *cravieri*) (Neotropical).
- CRAVERII Bellardi, 1859:7, pl. 1, fig. 1 (wing); also 1861:207, pl. 1, fig. 1. Type-locality: "Mexico". Distr. — Mexico.
- CYDIPPE Alexander, 1947f:77, fig. 2 (male hypopygium). Type-locality: Venezuela, Aragua, Choroní. Distr. — Guatemala, Venezuela.
- EDWARDSI Bellardi, 1859:8, pl. 1, fig. 2 (wing) (also 1861:208, pl. 1, fig. 2). Type-locality: "Mexico". Distr. — Mexico.
- LARGA Alexander, 1946b:222, fig. 4 (wing and male hypopygium). Type-locality: Mexico, Michoacán, Uruapán. Distr. — Mexico.
- OBLIQUEFASCIATA Macquart, 1846:143(15), pl. 1, fig. 10 (female). Type-locality: Colombia [New Grenada] temperate region. Enderlein, 1912:6,

records it from Ecuador and Venezuela. Identity uncertain, presumed to belong to *Bellardina* and possibly the same as *theobromina*.

- PARRAI Alexander, 1940b:141. Type-locality: Mexico, Chiapas, Porvenir. Distr. — Mexico.
- SCHIZOMERA Alexander, 1940b:142. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Southwestern U. S. A. to Mexico.
- THEOBROMINA Edwards, 1919:159, fig. 21 (ovipositor). Type-locality: Ecuador, Cuenca, 2532 meters. Distr. — Venezuela, Ecuador, Peru, Bolivia.
- WETMOREANA Alexander, 1947a:173. Type-locality: Guatemala, Tecpán, Sierra Santa Elena, 9500 feet. Distr. — Guatemala.

Subgenus EUMICROTIPULA Alexander

- Microtipula*, subg. EUMICROTIPULA Alexander, 1922d:74 Type-species, *macrotrichiata* Alexander (orig. des.) (Neotropical).
- ABORTIVA Alexander, 1914f:358. Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- ABSONA Alexander, 1937c:486. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- ACCIPITRINA Alexander, 1946d:193. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- ACCUMULATRIX Alexander, 1951b:297. Type-locality: Guatemala, Chicacao El Naranjo. Distr. — Guatemala.
- AEDON Alexander, 1947f:80, fig. 3 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- AGLOSSA Alexander, 1962c:40, figs. 25 (wing), 28 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- AGRIPPINA Alexander, 1945o:581. Type-locality: Ecuador, Bolivar, Hacienda Talahua. Distr. — Ecuador.
- ALBIFASCIATA Macquart, 1838a:62 (58). Type-locality: "Chile". Ref. — Alexander, 1929a:35. Identity uncertain.
- AMBLYTHRIX Alexander, 1967b:457, fig. 7 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- AMPHION Alexander, 1952b:101, fig. 2 (male hypopygium) (as 1951). Type-locality: Chile, Malleco, Curacautin, Termas de Río Blanco. Distr. — Chile.
- ANDALGALA Alexander, 1919a:150. Type-locality: Argentina, Catamarca, Cerro de Carocal. Distr. — Argentina.
- ANDINA Brèthes, 1909:88. Type-locality: Argentina, Mendoza, Puente del Inca. Distr. — Argentina.
- ANDROMACHE Alexander, 1950a:36. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- ANGOLENSIS Alexander, 1944i:119, fig. 1 (male hypopygium). Type-locality: Chile, Malleco, Angol. Distr. — Chile.
- ANTARCTICA Alexander, 1920e:19, figs. 12, 16 (male hypopygium). Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:38.
- ANTHONYMPHA Alexander, 1928f:629. Type-locality: Argentina, Río Negro, Bariloche, L. Nahuel Huapí. Distr. — Chile, Argentina.



- APTEROGYNE Philippi, 1865:606. Type-locality: Chile, Valdivia. Distr. — Chile. Refs. — Alexander, 1928i:277, figs. 1, 2 (male hypopygium). 1929a:36.
- ARAGUENSIS Alexander, 1950a:37, fig. 2 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- ARAUCANIA Alexander, 1928i:280, figs. 6 — 8 (male hypopygium). Type-locality: Chile, Santiago, Macul. Distr. — Chile.
- ARECUNA Alexander, 1931b:3, figs. 13, 14 (male hypopygium). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela.
- ARMLLATA Alexander, 1916a:30, pl. 4, fig. 8 (antenna) pl. 5, fig. 8 (male hypopygium). Type-locality: Colombia, Magdalena, La Cumbrec. Distr. — Colombia.
- ASAROTON Alexander, 1962b:345, fig. 14 (male hypopygium). Type-locality: Argentina, Santa Cruz, Cachillo, Pié del Cerro, Lago Argentino. Distr. — Argentina.
- ASTERIA Alexander, 1951d:6, fig. 1 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- ATACAMA Alexander, 1912f:357, figs. d (wing), u (ovipositor). Type-locality: Bolivia, San Antonio. Distr. — Bolivia.
- ATAMELES Alexander, 1965b:345, fig. 5 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- ATROSCAPA Alexander, 1949b:470. Type-locality: Brazil, São Paulo, São Paulo, Ipiranga. Distr. — Brazil.
- ATROVELUTINA (Alexander), 1928f:624 [*Microtipula (Microtipula)*]. Type-locality: Paraguay, Santa Bárbara. Distr. — Paraguay.
- AURICOMATA Alexander, 1942b:366, figs. 14, 15 (male hypopygium). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- AUSTROANDINA Alexander, 1929a:54, pl. 5, figs. f (wing), 142, 143, 144 (male hypopygium). Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- AZTECA Alexander, 1925a:357. Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico.
- BAECKSTROEMI Alexander, 1920d:26. Type-locality: Chile, Juan Fernández, Masatierra. Distr. — Chile. Refs. — Alexander, 1929a:37; 1952c:39, figs. 1 (wing), 2 (male hypopygium).
- BALLOUI Alexander, 1938h:428. Type-locality: Costa Rica, Coliblanco. Distr. — Costa Rica.
- BARRETOI Alexander, 1923c:312. Type-locality: Argentina, Buenos Aires, Monte Veloz. Distr. — Argentina.
- BATHROMECES Alexander, 1962c:41, fig. 27 (antenna). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- BIGOTIANA Alexander, 1920c:15. Type-locality: Chile, Magallanes. Distr. — Chile. Ref. — Alexander, 1929a:38.
- BOGOTANA Alexander, 1938d:327. Type-locality: Colombia, Bogotá. Distr. — Colombia.
- ERETHESIANA Alexander, 1929a:45, figs. 121, 122, 123 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina, Chile.
- BREVICOMA Alexander, 1944i:121. Type-locality: Chile, Malleco, Curacautin, Termas Rio Blanco. Distr. — Chile.

- BROWNIANA Alexander, 1940c:107. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- BRUCHI Alexander, 1920b:10. Type-locality: Argentina, Buenos Aires. Distr. — Argentina.
- CALLISTO Alexander, 1944g:297, fig. 4 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- CALLITHRIX Alexander, 1944g:295, fig. 3 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- CAMPA Alexander, 1914f:358, pl. 15, figs. 5 (wing), 9-12 (male hypopygium), 18 (antenna). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- CAPUCINA Alexander, 1947f:82, fig. 4 (male hypopygium). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- CARIZONA Alexander, 1913c:208, pl. 7, figs. 2 — 4 (male hypopygium), 7 (wing). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia, Ecuador, Peru. Ref. — Alexander, 1946d:195.
- sphaerulifera* Edwards, 1919:156, figs. 15 (ovipositor), 16, 17 (male hypopygium). Type-locality: Ecuador, Carchi, Tulcán. Distr. — Ecuador.
- CHACOPATA CHACOPATA Alexander, 1944L:174, fig. 1 (male hypopygium). Type-locality: Venezuela, Aragua, road between Maracay-Choroni. Distr. — Venezuela.
- ssp. CHANCA Alexander, 1945i:244. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- CHARMOSYNE Alexander, 1967b:459, fig. 8 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- CHICANA Alexander, 1945a:12. Type-locality: Ecuador, Tungurahua, Minza Chica. Distr. — Ecuador. Ref. — Alexander, 1945L:378, fig. 1 (male hypopygium).
- CHILENSIS Alexander, 1920e:13. Type-locality: Chile (unknown station in southern Chile). Distr. — Chile. Ref. — Alexander, 1929a:65, pl. 5, fig. c (wing).
- CHILLANICA Alexander, 1945f:10, fig. 11 (male hypopygium). Type-locality: Chile, Ñuble, Termas de Chillán. Distr. — Chile.
- CHILOTA Alexander, 1929a:46, figs. 125, 126 (male hypopygium). Type-locality: Chile, Chiloé, near Puntra. Distr. — Chile.
- CLARKIANA Alexander, 1929a:55. Type-locality: Chile, Los Andes. Distr. — Chile.
- CLAVARIA Alexander, 1945o:583. Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- CONSONATA Alexander, 1940d:202. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- CONSPICILLATA Alexander, 1945i:234. Type-locality: Peru, Junín, Tarma, Moyobamba. Distr. — Peru, Bolivia. Ref. — Alexander, 1951d:7, fig. 2 (male hypopygium).
- CORONARIA Alexander, 1940d:207, fig. 4 (male hypopygium). Type-locality: Ecuador, Río Blanco near Baños. Distr. — Ecuador.
- COSTARICENSIS (Alexander), 1922a:6 [*Microtipula* (*Microtipula*)]. Type-locality: Costa Rica, Atirro. Distr. — Costa Rica.
- CREPERA Alexander, 1952b:103, fig. 3 (male hypopygium) (as 1951). Type-locality: Chile, Malleco, Termas de Río Blanco. Distr. — Chile.



- CRISTATA* Alexander, 1945g:72, fig. 5 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- CROSSOSPILA* Alexander, 1929a:50, figs. 9 (wing), 133-135 (male hypopygium). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile.
- CURINAO* Alexander, 1941f:361, pl. 15, figs. 3 (wing), 16 (antenna). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- CYCLOMERA* Alexander, 1951b:298, fig. 19 (male hypopygium). Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- DARLINGTONIANA* Alexander, 1939d:101. Type-locality: Hispaniola: Dominican Republic, Valle Nuevo, southeast of Constanza. Distr. — Dominican Republic.
- DELECTATA* Alexander, 1940d:203. Type-locality: Ecuador, Tungurahua, El Tablón near Baños. Distr. — Ecuador.
- DICTYOPHORA* Alexander, 1962c:43, fig. 29 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- DIMORPHIA* Alexander, 1929a:63, fig. 35; pl. 5, figs. h (wing), 159, 161 (male hypopygium). Type-locality: Argentina, Misiones, Río Pichileufu. Distr. — Argentina, Chile.
- DUIDAE* Alexander, 1931b:4, fig. 15 (male hypopygium). Type-locality: Venezuela, Mt. Duida. Distr. — Venezuela.
- DUSENI* Alexander, 1920e:30, pl. 1, fig. 9 (male hypopygium). Type-locality: Chile, Aysén, Río Aysén. Distr. — Chile. Ref. — Alexander, 1929a:39.
- EFFICAX* Alexander, 1945a:14. Type-locality: Ecuador, Tungurahua, Pastaza, Río Blanco. Distr. — Ecuador.
- EMERITA* Alexander, 1944d:304. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador.
- ENDERLEINANA* Alexander, 1929a:49, figs. 130-132 (male hypopygium). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina, Chile.
- EXILIS* Alexander, 1916a:24, pl. 3, fig. 1 (wing); pl. 4, fig. 5 (antenna); pl. 5, figs. 2, 5 (male hypopygium). Type-locality: Peru, Lima, San Cristóbal Hill. Distr. — Peru.
- EXPLETA* Alexander, 1962c:44, fig. 26 (wing). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- FATIDICA* Alexander, 1940a:280. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- FAZI* Alexander, 1928f:630. Type-locality: Chile, Valdivia, L. Panguipulli. Distr. — Chile.
- FLAVIDULA* Alexander, 1940d:205, fig. 3 (male hypopygium). Type-locality: Ecuador, Mt. Tunguragua. Distr. — Ecuador, Peru, Bolivia.
- FLAVOANNULATA* Jacobs, 1900:106. Type-locality: Argentina, Magallanes, Morrochico, Valle Río Gallegas. Distr. — Argentina. Refs. — Jacobs, 1906:68, pl. 3, fig. 5 (entire male, color); Alexander, 1929a:37.
- FOERSTERI* Alexander, 1962c:46, figs. 30 (wing), 31 (male hypopygium). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- FOERSTERIANA* Alexander, 1967b:461, fig. 9 (male hypopygium). Type-locality: Colombia, Manizales. Distr. — Colombia.

- FORTIOR FORTIOR Alexander, 1951b:310, fig. 23 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- ssp. AREPUCHOENSIS Alexander, 1962c:48, fig. 32 (male hypopygium). Type-locality: Bolivia, Yungas de Arepucho, Sihuencas. Distr. — Bolivia.
- ssp. KLAUSMA Alexander, 1962c:50. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- FRAUDULENTA Alexander, 1928f:626. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- FUEGIENSIS Alexander, 1920e:27, pl. 1, figs. 1, 6 (male hypopygium). Type-locality: Chile, Tierra del Fuego, Río Grande. Distr. — Chile. Ref. — Alexander, 1929a:64, figs. 11 (wing), 162-164 (male hypopygium).
- GLAPHYROPTERA Philippi, 1865:604. Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1929a:53, figs. g (wing), 139-141 (male hypopygium).
- GLOSSOPHORA Alexander, 1962c:51, fig. 33 (male hypopygium). Type-locality: Bolivia, Yungas de Arepucho, Sihuencas. Distr. — Bolivia.
- GRAPHICA Schiner, 1868:36. Type-locality: "South America." Distr. — South America. Identity uncertain; apparently close to *monilifera* Loew (q.v.).
- GUARANI Alexander, 1914f:357. Type-locality: Brazil, Rio Grande (?). Distr. — Brazil.
- HEDYMOPA Alexander, 1944d:303. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- HOSTIFICA Alexander, 1937d:600. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- HUANCA Alexander, 1945i:237. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- HYLONYMPIA Alexander, 1929a:57, figs. 147-149 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- IGUAZUENSIS (Alexander), 1928f:625 [*Microtipula (Microtipula)*]. Type-locality: Argentina, Misiones, Iguazu Falls. Distr. — Argentina.
- IMMERENS Alexander, 1944c:40 [*Tipula (Microtipula)*]. Type-locality: Venezuela, Miranda, Río Chicaíto. Distr. — Venezuela.
- IMMORSA Alexander, 1945i:239, fig. 1 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.
- INAEQUIARMATA Alexander, 1962c:52, fig. 34 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- INAEQUIDENS Alexander, 1946d:196, fig. 5 (male hypopygium). Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- INCA Alexander, 1912f:351, figs. c (wing), 1 (male hypopygium). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- INCONDITA Alexander, 1951b:307. Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia. Ref. — Alexander, 1962c:54.
- INFIDELIS Alexander, 1951b:306. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- INFINITA Alexander, 1945g:76, fig. 6 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

- INNUBENS Alexander, 1942b:363, fig. 16 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- INVIGILANS Alexander, 1951b:312. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- ITATIAYENSIS Alexander, 1944m:302. Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- JACOBSIANA Alexander, 1929a:43, figs. 118-120 (male hypopygium); pl. 5, fig. d (wing). Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- JAENNICKEANA Alexander, 1929a:47, pl. 5, fig. a (wing), figs. 127-129 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- JIVARO Alexander, 1916a:25, pl. 3, fig. 2 (wing), pl. 4, fig. 6 (antenna), pl. 5, figs. 1, 3, 6 (male hypopygium). Type-locality: Ecuador, Chimborazo, Alaousi. Distr. — Ecuador.
- JUBILANS Alexander, 1940a:279. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- JUVENTA Alexander, 1937c:488. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- KATHEMA Alexander, 1965b:346, fig. 6 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- KUEHLIORNI Alexander, 1962c:55, fig. 35 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- KUSCHELI Alexander, 1967b:463, fig. 11 (male hypopygium). Type-locality: Chile, Tierra del Fuego, Beagle Canal, Navarino Is., P. Williams. Distr. — Chile.
- LANIGERA LANIGERA Alexander, 1928f:628. Type-locality: Argentina, Rio Negro, Bariloche, L. Nahuel Huapi. Distr. — Argentina.
- ssp. TAU Alexander, 1944i:122. Type-locality: Chile, Cautín, Temuco. Distr. — Chile.
- LATEROSETOSA Alexander, 1931c:623. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- LATIFOLIA Alexander, 1944i:124, fig. 2 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- LEGITIMA (Alexander), 1928f:623 [*Microtipula (Microtipula)*]. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- LIGULATA Alexander, 1929a:61, figs. 155-157 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile, Argentina. Ref. — Alexander, 1962b:347, fig. 13 (male hypopygium).
- LIGULPENICILLATA Alexander, 1946b:229, fig. 12 (male hypopygium). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- LONGIBASIS Alexander, 1946d:179. Type-locality: Mexico, Morcos, Lagunas de Zempole. Distr. — Mexico.
- LONGURIODES Alexander, 1954d:140. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- MACINTYREANA Alexander, 1941d:120, fig. 8 (male hypopygium). Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- MACROTRICHIATA (Alexander), 1922d:74 [*Microtipula (Eumicrotipula)*]. Type-locality: Brazil, Rio de Janeiro, Serra dos Órgãos. Distr. — Brazil.

- MAGELLANICA Alexander, 1920e:21, figs. 10, 11, 14 (male hypopygium). Type-locality: Chile, Magallanes, Punta Arcnas. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:62, fig. 36; pl. 5, figs. c (wing), 158 (male hypopygium).
- magellanicola* Alexander, 1920e:22. Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile. Ref. — Alexander, 1946d:177.
- MARMORIPENNIS Rondani, 1850:194. Type-locality: "Venezuela". Identity uncertain.
- MARTINEROWNI Alexander, 1946d:198, fig. 6 (male hypopygium). Type-locality: Ecuador, Chimborazo, Urbina. Distr. — Ecuador.
- MECOGLOSSA Alexander, 1967b:464, fig. 12 (male hypopygium). Type-locality: Colombia, Cundinamarca, Monterredondo. Distr. — Colombia.
- MEDIODENTATA Alexander, 1944g:293, fig. 2 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- MERIDIANA Edwards, 1919:160, figs. 23, 24 (male hypopygium), 25 (ovipositor). Type-locality: Ecuador, Tio Loma, 4260 meters. Distr. — Ecuador.
- MICROPILOTA Alexander, 1928f:634. Type-locality: Chile, Malleco. Distr. — Chile.
- MIRANHA Alexander, 1913c:210, pl. 7, fig. 1 (wing). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- MITHRODATES Alexander, 1951b:303, figs. 21, 22 (male hypopygium). Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- MITUA Alexander, 1916a:29, pl. 3, fig. 4 (wing), pl. 4, fig. 7 (antenna), pl. 5, figs. 4, 7 (male hypopygium). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- MOCOA Alexander, 1913c:210, pl. 7, fig. 6 (wing). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- MOCTEZUMAE Alexander, 1925a:356, fig. 26 (male hypopygium). Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico.
- MONILIFERA Loew, 1851:404, figs. 26 (male antenna), 27 (wing). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- MONILIFEROIDES Alexander, 1920f:221. Type-locality: Argentina, Buenos Aires, Monte Veloz. Distr. — Argentina, Brazil, Colombia. Ref. — Bruch, 1939a, pl. 3, fig. 6 (head of male), pl. 4, fig. 13 (wing), pl. 5, fig. 18 (entire male).
- MONILIFORMIS Röder, 1886:259. Type-locality: Colombia, Cerro del Altar, 3800 meters. Distr. — Colombia.
- MORDAX Alexander, 1945c:256, fig. 6 (male hypopygium). Type-locality: Brazil, São Paulo, Serra da Cantarreira. Distr. — Brazil.
- MORPHEA Alexander, 1946d:200, fig. 7 (male hypopygium). Type-locality: Ecuador, Mt. Tungurahua. Distr. — Ecuador.
- NAVARINOENSIS Alexander, 1962a:33. Type-locality: Chile, Tierra del Fuego, Beagle Canal, Is. Navarino. Distr. — Chile.
- NEIVAI Alexander, 1940c:895. Type-locality: Venezuela, Mérida, Páramo de Mucuchics. Distr. — Venezuela.
- NETHIS Alexander, 1965b:348. Type-locality: Peru, Cuzco, Quincmil. Distr. — Peru.
- NIGRISCAPA Alexander, 1946d:184. Type-locality: Surinam, Marowijne near Gran Creek. Distr. — Surinam.



- NIMBINERVIS Alexander, 1946b:227, fig. 11 (male hypopygium). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- NOLENS Alexander, 1965b:349, fig. 7 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Libertad. Distr. — Ecuador.
- NORDENSKJOELDI Alexander, 1920e:18, pl. 1, figs. 4, 8 (male hypopygium). Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:39.
- NOTHIOFAGETORUM Alexander, 1929a:59, figs. 150, 151 (male hypopygium). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina.
- NOTORIA Alexander, 1942d:137. Type-locality: Ecuador, Napo-Pastaza, Pundoa. Distr. — Ecuador. Ref. — Alexander, 1951b:312, fig. 24 (male hypopygium).
- NOVALEONENSIS Alexander, 1940b:147. Type-locality: Mexico, Nuevo León, Cerro Potosi. Distr. — Mexico.
- NOVATRIX Alexander, 1942b:355. Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- NUBIFERA Wulp, 1881:151, pl. 15, fig. 6. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Identity uncertain.
- OBIRATA Alexander, 1940a:277. Type-locality: Ecuador, Napo-Pastaza, Río Blanco, Yunguilla. Distr. — Ecuador.
- OBSCURICINCTA Alexander, 1940b:145. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- ODONTOMERA Alexander, 1967b:467, fig. 13 (male hypopygium). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- OLSSONIANA Alexander, 1943c:19. Type-locality: Venezuela, Mérida, Páramo de Mueuchies. Distr. — Venezuela.
- OMNILUTEA Alexander, 1967b:468. Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- ONA Alexander, 1920e:11. Type-locality: Chile, Tierra del Fuego, Río Grande. Distr. — Chile.
- OREONYMPHIA Alexander, 1929a:56, figs. 145, 146 (male hypopygium). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina.
- ORIZABENSIS Alexander, 1941c:323, fig. 1 (wing). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- ORNATICORNIS Wulp, 1891:195, pl. 12, fig. 2 (male hypopygium). Type-locality: "Colombia" [New Grenada]. Distr. — Colombia.
- OSCULATA Alexander, 1944d:301. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PALENA Alexander, 1944L:176, fig. 2 (male hypopygium). Type-locality: Venezuela, Mérida, Tabay. Distr. — Venezuela.
- PALITANS Alexander, 1965b:350, fig. 8 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- PALLIDINEURIS Macquart, 1846:144(16), pl. 2, fig. 1. Type-locality: "Colombia" (New Granada). Distr. — Colombia.
- pallidinervis* Alexander, 1914f:353. error.
- PALLIDISIGNATA PALLIDISIGNATA Alexander, 1928i:284, figs. 10-12 (male hypopygium). Type-locality: Chile, Malleco. Distr. — Chile.
- ssp. SALUTATORIA Alexander, 1945f:12. Type-locality: Chile, Llanquihue, Cayutue. Distr. — Chile.

- PANTHERINA Alexander, 1941a:242. Type-locality: "Venezuela". Distr. — Venezuela.
- PARANENSIS Alexander, 1945d:262. Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil.
- PARVILOBA Alexander, 1929a:60, figs. 152-154 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina, Chile.
- PATAGONICA Alexander, 1920e:25, pl. 1, figs. 3, 5, 7 (male hypopygium). Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:38.
- PEDIFORMIS Alexander, 1962c:57, fig. 36 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- PERJOVIALIS Alexander, 1944g:290. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- PERSTUDIOSA Alexander, 1937d:599. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PETALURA Alexander, 1953b:20, figs. 1, 2 (male hypopygium). Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- PETIOLARIS Alexander, 1940a:282. Type-locality: Ecuador, Napo-Pastaza, Mapotó, Río Pastaza. Distr. — Ecuador.
- PHALANGIODES Alexander, 1945a:16. Type-locality: Ecuador, Tungurahua, Mt. Tungurahua, Minza Chica. Distr. — Ecuador. Ref. — Alexander, 1953a:11, fig. 8 (male hypopygium and antenna).
- PHILIPPANA Alexander, 1920b:12. Type-locality: Argentina, Santa Cruz, Valle Tunel. Distr. — Argentina, Chile. Ref. — Alexander, 1929a:37, fig. 10 (wing).
- PICTIPENNIS Walker, 1837:333. Type-locality: Chile, Magallanes, Port Famine, Straits of Magellan. Distr. — Chile. Ref. — Alexander, 1929a:35.
- FILULIFERA Edwards, 1919:154, figs. 9, 10 (male hypopygium). Type-locality: Ecuador, El Pelado, 4150 meters. Distr. — Ecuador.
- PIRIONI Alexander, 1928i:282, fig. 9 (male hypopygium). Type-locality: Chile, Bío-Bío. Distr. — Chile.
- PIRO Alexander, 1914f:360, figs. 4 (wing), 13 (male hypopygium), 17 (antenna). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- PLATYTERGATA Alexander, 1962c:58, fig. 37 (male hypopygium). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- PROCERICORNIS Edwards, 1919:155, figs. 11, 12 (male hypopygium). Type-locality: Ecuador, Chillacocha, 3900 meters. Distr. — Ecuador. Ref. — Alexander, 1967b:462, fig. 10 (male hypopygium).
- PROFUGA Alexander, 1937f:218. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PRCTRUDENS Alexander, 1952e:84, fig. 3 (male hypopygium). Type-locality: Argentina, Neuquén, Chapelco. Distr. — Argentina.
- PSITTACINA Alexander, 1962c:60, fig. 38 (male hypopygium). Type-locality: Bolivia, Yungas de Forestal. Distr. — Bolivia, Argentina.
- QUADRISSETOSA Alexander, 1951d:10, fig. 5 (male hypopygium). Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.



- QUICHUA Alexander, 1916a:27, pl. 3, fig. 3 (wing). Type-locality: Peru, Lima, Matucana. Distr. — Peru.
- RECIPROCA Alexander, 1946d:181. Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- RBDUNCA Alexander, 1967b:469, fig. 14 (male hypopygium). Type-locality: Colombia, Manizales. Distr. — Colombia.
- REEDI Alexander, 1934d:176. Type-locality: Chile, Llanquihue, Cayutue, Lago Todos los Santos. Distr. — Chile.
- RESPLENDENS Alexander, 1946d:186, fig. 1 (male hypopygium). Type-locality: Ecuador, Tungurahua, Mt. Tungurahua, Minza Chico. Distr. — Ecuador.
- RIVETI RIVETI Edwards, 1919:158, figs. 18 (wing), 19, 20 (male hypopygium). Type-locality: Ecuador, El Pelado, 4151 meters. Distr. — Ecuador.
- ssp. TOLMENSIS Alexander, 1941a:244. Type-locality: Colombia, Cordillera Central, Mt. Tolima. Distr. — Colombia.
- UCANA Alexander, 1946d:189, fig. 2 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- RUFIROSTRIS Bigot, 1888:5, pl. 1, fig. 2 (wing). Type-locality: Chile, Tierra del Fuego. Distr. — Chile. Ref. — Alexander, 1929a:36.
- RUNTUNENSIS Alexander, 1940d:209, fig. 5 (male hypopygium). Type-locality: Ecuador, Tungurahua, Runtun near Baños. Distr. — Ecuador.
- SARIAPAMPAE Alexander, 1967b:471, fig. 15 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- SATRAPA Alexander, 1952b:104, fig. 4 (male hypopygium) (as 1951). Type-locality: Chile, Chiloé Is., Río Coluco. Distr. — Chile.
- SCHACHOVSKOYI Alexander, 1952e:86, fig. 4 (male hypopygium). Type-locality: Argentina, Neuquén, Mt. Chapelco, near San Martín de los Andes. Distr. — Argentina.
- SCRIPTELLA Alexander, 1940c:109. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- SEMIVULPINA Alexander, 1944g:292. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- SERRIOBATA Alexander, 1941a:246. Type-locality: Chile, Malleco, Curacautín, Termas Río Blanco. Distr. — Chile.
- SERVAL Alexander, 1937c:490. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SONGOANA Alexander, 1962c:62. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- SPATULIFERA Alexander, 1928f:632. Type-locality: Chile, Valdivia, L. Panquipulli. Distr. — Chile.
- SPILOTA Wiedemann, 1828:553. Type-locality: "Brazil". Distr. — Brazil. Identity uncertain.
- STEINBACHII Alexander, 1946d:191, fig. 3 (male hypopygium). Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.
- STENOGLOSSA STENOGLOSSA Alexander, 1942b:359, fig. 12 (male hypopygium). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- ssp. PROPITIA Alexander, 1944g:295. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- SUAVISSIMA Alexander, 1951d:8, fig. 3 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.

- SUBANDINA Philippi, 1865:605. Type-locality: Chile, Santiago, foot of Andes. Distr. — Chile. Refs. — Alexander, 1928i:278, figs. 3-5 (male hypopygium), 13 (ovipositor); 1929a:35.
- SUBCANANA Edwards, 1919:155, figs. 13, 14 (male hypopygium). Type-locality: Ecuador, Narihuíña, 3904 meters. Distr. — Ecuador.
- SUBGLABRATA Alexander, 1951b:301, fig. 20 (male hypopygium). Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- SUBLIGULATA Alexander, 1941a:244. Type-locality: Chile, Malleco, Curacautín, Termas Río Blanco. Distr. — Chile.
- TANYMETRA Alexander, 1962e:64, figs. 39, 40 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- TEHUELCHÉ Alexander, 1920e:17, figs. 13, 16, 17, 18 (male hypopygium), 19 (antenna), 20 (ovipositor). Type-locality: Chile, Tierra del Fuego, Río Grande. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:38.
- TEPHRONOTA Alexander, 1942b:368. Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- TERSA Alexander, 1928f:633. Type-locality: Chile, Bio-Bío. Distr. — Chile.
- TERSOIDES Alexander, 1945f:13, fig. 12 (male hypopygium). Type-locality: Chile, Nuble, Termas de Chillán. Distr. — Chile.
- THALIA Alexander, 1945o:586. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- TOVARENSIS Alexander, 1947d:40, fig. 1 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- TRIEMARGINATA Alexander, 1935b:99, figs. 1-3 (male hypopygium). Type-locality: Chile, Metrengo. Distr. — Chile.
- TRISPILOTA Alexander, 1942b:356, fig. 11 (male hypopygium). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- TRISTILLATA Alexander, 1929a:52, figs. 8 (wing), 136-138 (male hypopygium). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina, Chile.
- TUNGURAGUANA Alexander, 1940d:211. Type-locality: Ecuador, Mt. Tunurahua, Chaupí. Distr. — Ecuador.
- TYRANNA Alexander, 1946d:182. Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- UNISTRILIATA Alexander, 1941e:242. Type-locality: "Brazil". Distr. — Brazil.
- URSULA Alexander, 1951d:9, fig. 4 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- VALDIVIANA Philippi, 1865:606. Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1929a:36.
- VARIINEURA (Bigot), 1888:6, pl. 1, fig. 4 (*Nephrotoma*). Type-locality: Chile, Tierra del Fuego. Distr. — Chile. Ref. — Alexander, 1929a:36. Kertész, 1902:325, corrects the name to *variinervis*.
- VIRGULATA Williston, 1900:228. Type-locality: Mexico, Morelos, Amecameca. Distr. — Mexico.
- oreomyzoides* Alexander, 1926a:175. Type-locality: Mexico, Ajusco.
- VOTIVA Alexander, 1944h:374. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- WILLINKI Alexander, 1962b:348, fig. 16 (male hypopygium). Type-locality: Argentina, Santa Cruz, Lago Argentino, Calafate. Distr. — Argentina.

- WITTEI Alexander, 1920b:11. Type-locality: Argentina, Santa Cruz, Valle Tunch. Distr. — Argentina. Refs. — Alexander, 1929a:37, fig. 34 (wing); 1962b:350, fig. 15 (male hypopygium).
- WOYTKOWSKIANA Alexander, 1942b:361, fig. 13 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- YANAMONTEANA Alexander, 1945i:242, fig. 2 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- YUNGASENSIS Alexander, 1962c:65. Type-locality: Bolivia, Yungas de Songo Valley. Distr. — Bolivia.
- ZELTALE Alexander, 1928a:102. Type-locality: Mexico, Chiapas, near Gruta del Elcanto. Distr. — Mexico.
- ZEUGMATA Alexander, 1951d:12, fig. 6 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- ZOTZIL Alexander, 1928a:103. Type-locality: Mexico, Chiapas, Hueitcepec. Distr. — Mexico.

Subgenus LUNATIPULA Edwards

- Tipula*, subg. LUNATIPULA Edwards, 1931:81. Type-species, *lunata* Linnaeus (orig. des.) (Western Palaearctic).
- ABSCISSA Alexander, 1946e:527. Type-locality: Mexico, Morelos, Tepoztlán. Distr. — Mexico.
- DAMPFIANA Alexander, 1946e:526. Type-locality: Mexico, Mt. Telapón, Valle de México. Distr. — Mexico.
- MICHOACANA Alexander, 1946b:230, figs. 7 (male wing), 8 (female wing). Type-locality: Mexico, Michoacán, Pedregal de Tancitaro. Distr. — Mexico.

Subgenus MICROTIPULA Alexander

- MICROTIPULA Alexander, 1912f:360 (as genus). Type-species, *amazonica* Alexander (orig. des.) (Neotropical).
- Tipula*, subg. *Nephrotomodes* Alexander, 1946b:224. Type-species, *smilodon* Alexander (orig. des.). Refs. — Alexander, 1945k:330, 9 figs.; 1945m:415, 25 figs.
- AEQUITORIALIS Alexander, 1940d:213. Type-locality: Ecuador, Napo-Pastaza, Río Upano, Macar. Distr. — Ecuador.
- insolabilis* Alexander, 1944g:287, fig. 1 (male hypopygium) [*Tipula* (*Nephrotomodes*)]. Type-locality: Peru, Huanuco, Afilador. Distr. — Peru.
- AFFABILIS Alexander, 1945c:247, fig. 1 (male hypopygium). Type-locality: Brazil, São Paulo, Serra da Cantareira. Distr. — Brazil.
- AKESTRA Alexander, 1965b:340, fig. 2 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- ALECTRO Alexander, 1945m:422, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- AMARA Alexander, 1951b:288, fig. 11 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- AMAZONICA (Alexander), 1912f:361, figs. i (wing), q (male hypopygium) (*Microtipula*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil, Surinam.

- AMOENICORNIS Alexander, 1922b:210. Type-locality: Argentina, Tucumán, San Pedro de Colalao. Distr. — Argentina. Ref. — Alexander, 1962b:341, fig. 10 (male hypopygium).
- APOLLYON Alexander, 1951b:292, fig. 13 (male hypopygium). Type-locality: Bolivia, Cochabamba, Paractito. Distr. — Bolivia.
- APPENDENS (Enderlein), 1912:14 (*Macromastix*). Type-locality: Ecuador, Santa Inez. Distr. — Ecuador.
- ARIRANIIAE Alexander, 1945k:338, fig. 2 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Paraguay.
- ARMATIPENNIS ARMATIPENNIS Alexander, 1912f:347, figs. g (wing), o (male hypopygium), v (ovipositor). Type-locality: Brazil, Mato Grosso, Chapada [dos Guimarães]. Distr. — Brazil, Argentina.
ssp. NAPOENSIS Alexander, 1921d:98. Type-locality: Peru, Loreto, Napo River. Distr. — Peru.
- AURICULARIS Alexander, 1942d:136 [*Tipula (Nephrotomodes)*]. Type-locality: Ecuador, Manabí, Palmar, Playones. Distr. — Venezuela, Ecuador. Ref. — Alexander, 1945m:421, fig. 6 (male hypopygium).
tecta Alexander, 1926a:166 (preocc. Scudder, 1877). Type-locality: Venezuela, Táchira, Táchira. Ref. — Alexander, 1945m:421, fig. 5 (male hypopygium).
subtecta Alexander, 1964b:282 (unnecessary nom. nov. for *tecta* Alexander). Type-locality: Venezuela, Táchira, Táchira.
- AUSTROVOLENS Alexander, 1951b:270, fig. 1 (male hypopygium). Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- AYMARA Alexander, 1912f:353, figs. b (wing), m (male hypopygium), s (ovipositor). Type-locality: Bolivia, San Antonio. Distr. — Bolivia. Ref. — Alexander, 1962c:67, fig. 41 (male hypopygium).
- BILIMEKI Alexander, 1941c:325, fig. 2 (wing). Type-locality: Mexico, Veraeruz, Orizaba. Distr. — Mexico.
- BIPROLATA Alexander, 1967b:449, fig. 1 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- BITRIBULA Alexander, 1967b:450, fig. 2 (male hypopygium). Type-locality: Peru, Madre de Dios, Avispas, 150 km west of Puerto Maldonado. Distr. — Peru.
- BLASERI Alexander, 1942g:171, figs. 3 (wing), 9 (male hypopygium). Type-locality: Brazil, Minas Gerais. Distr. — Brazil.
- BRASILIENSIS (Wiedemann), 1828:554, pl. VIb, figs. 13 (male antenna), 14 (wing) (*Megistocera*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Osten Sacken, 1886b:184, note on type. Identity uncertain.
- BRUESI Alexander, 1945m:431, fig. 17 (male hypopygium). Type-locality: Lesser Antilles: Grenada, Grand Etang. Distr. — Grenada.
- CEROGAMA Alexander, 1944d:306 [*Tipula (Nephrotomodes)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CITHARIFORMIS Alexander, 1967b:451, fig. 3 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Libertad, Tena. Distr. — Ecuador.
- COLOMBICOLA Alexander, 1929d:768. Type-locality: Colombia, Chiriguana Dist., Lake Sapatoza Region. Distr. — Colombia.



- CONTEMPLATA Alexander, 1951b:287, fig. 10 (male hypopygium). Type-locality: Guatemala, Nahualate. Distr. — Guatemala.
- CRASSISTYLA Alexander, 1965b:342, fig. 3 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- CTENOPYGA Alexander, 1940c:110. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- DECENS Alexander, 1942b:351, figs. 9, 10 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- DECOLORATA (Alexander), 1935a:314 (*Habromastix*). Type-locality: Paraguay, Santa Bárbara. Distr. — Paraguay, Brazil. Ref. — Alexander, 1962b:342, fig. 12 (male hypopygium).
- DETECTA Alexander, 1926a:165. Type-locality: Venezuela, Táchira, Táchira. Distr. — Venezuela.
- DIACANTHOS Alexander, 1921d:97. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- effera* Alexander, 1929b:89. Type-locality: Peru, San Ramón. Distr. — Peru.
- DIADEXIA Alexander, 1967b:453, fig. 4 (male hypopygium). Type-locality: Peru, Huanuco, Cordillera Azul, Divisoria. Distr. — Peru.
- DIDACTYLA Alexander, 1942b:348, fig. 7 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- DIDOLOS Alexander, 1962b:343, fig. 11 (male hypopygium). Type-locality: Argentina, Tucumán, Quebrada Cainzo, Tafi. Distr. — Argentina.
- DIRHABDOPHORA Alexander, 1951b:274, fig. 3 (male hypopygium). Type-locality: Panama, Canal Zone, Summit. Distr. — Panama.
- DISCOPHORA Alexander, 1951b: 281, fig. 7 (male hypopygium). Type-locality: Peru, Loreto, Pucallpa. Distr. — Peru.
- EFFERON Alexander, 1945c:249, fig. 2 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Cidade Jardim. Distr. — Brazil.
- EFFETA Alexander, 1921d:100. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- EFFULTA Alexander, 1941d:109, fig. 2 (male hypopygium). Type-locality: Ecuador, Tungurahua, Río Blanco near Baños. Distr. — Ecuador. Ref. — Alexander, 1945m:425, fig. 10 (male hypopygium).
- EPIONE Alexander, 1944h:376. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- EROSTRATA Alexander, 1945m:424, fig. 8 (male hypopygium). Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia.
- EURYMERA EURYMERA Alexander, 1945c:250, fig. 3 (male hypopygium). Type-locality: Brazil, São Paulo, Serra da Cantareira. Distr. — Brazil.
- ssp. GOYAZICOLA Alexander, 1945c:252. Type-locality: Brazil, Goiás. Distr. — Brazil.
- ssp. PARAGUAYICOLA Alexander, 1945c:252. Type-locality: Paraguay, Santa Bárbara. Distr. — Paraguay.
- FALCIFER Alexander, 1944k:595. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- FELICIANA Alexander, 1945c:253, fig. 4 (male hypopygium). Type-locality: Brazil, São Paulo, Serra da Cantareira. Distr. — Brazil.

- FEROCIA Alexander, 1936e:57. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FIEBRIGI Alexander, 1941a:241. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- FLAVOPOLITA Alexander, 1942g:172, fig. 10 (antenna). Type-locality: Brazil, Rio de Janeiro, Serra do Itatiaia ("Südabhang, Waldreg"). Distr. — Brazil.
- GLADIATOR Alexander, 1914f:356, figs. 6 (wing), 7 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- CUATO Alexander, 1912f:349, figs. e (wing), p (male hypopygium). Type-locality: Brazil, Mato Grosso, Chapada [dos Guimarães]. Distr. — Brazil.
- GUERREROENSIS Alexander, 1940b:149. Type-locality: Mexico. Guerrero, Taxco. Distr. — Mexico.
- GUTTICELLULA Alexander, 1936a:13. Type-locality: Brazil, Rio de Janeiro, Recznde, Campo Belo. Distr. — Brazil.
- HETERODACTYLA Alexander, 1944k:593. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- HEXAMELANIA Alexander, 1951b:276, fig. 4 (male hypopygium). Type-locality: Guatemala, Chieacao, El Naranjo. Distr. — Guatemala.
- HISTRIONICA Alexander, 1945k:350. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.
- HORRIBILIS Alexander, 1945k:352, fig. 8 (male hypopygium). Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- ICASTA Alexander, 1941d:118, fig. 7 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador. Ref. — Alexander, 1945m:433, fig. 18 (male hypopygium).
- IMPATIENS Alexander, 1951b:284, fig. 9 (male hypopygium). Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.
- INAEQUILOBATA Alexander, 1937f:220. Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu, Napo Watershed. Distr. — Ecuador.
- INARMATA Alexander, 1928f:627. Type-locality: Argentina, Misiones, Puerto Bemberg. Distr. — Argentina.
- INFIDA Alexander, 1941d:116, fig. 6 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador. Ref. — Alexander, 1945m:433, fig. 19 (male hypopygium).
- INTEMPERATA Alexander, 1943c:200 [*Tipula* (*Nephrotomodes*)]. Type-locality: Ecuador, Manabi, Palmar, Río Maizito. Distr. — Ecuador.
- JIVARONIS Alexander, 1943c:202 [*Tipula* (*Nephrotomodes*)]. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- JORDANENSIS Alexander, 1949b:469, fig. 8 (male hypopygium). Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- JUQUIELLA Alexander, 1945c:255, fig. 5 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiã. Distr. — Brazil.
- KLAGESI Alexander, 1945k:348, fig. 7 (male hypopygium). Type-locality: Brazil, Pará, Benevides. Distr. — Brazil.
- LACOTIS Alexander, 1942g:169, figs. 2 (wing), 8 (male hypopygium). Type-locality: Brazil, Minas Gerais ?Río Bandeiro [Riacho dos Pan-deiros?]. Distr. — Brazil.

- LANGUIDULA Alexander, 1942b:347, fig. 6 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- LATICOSTATA Alexander, 1937d:598. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LETALIS Alexander, 1936e:55. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LICHYANA Alexander, 1945g:69, figs. 3, 4 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- LUCTIFICA Alexander, 1940d:200. Type-locality: Ecuador, Napo-Pastaza, San Francisco. Distr. — Ecuador.
- LUTEILIMBATA Alexander, 1944g:305. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- LYRIFORMIS Alexander, 1942b:350, fig. 8 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- MACROSTERNA (Alexander), 1912f:344, figs. k (male hypopygium), r (ovipositor) (*Pachyrhina*). Type-locality: Guatemala, Antigua. Distr. — Guatemala, Honduras.
- MANDATOR Alexander, 1945m:416, fig. 2 (male hypopygium). Type-locality: Costa Rica, San Matco, Higuito. Distr. — Costa Rica.
- MANNIANA Alexander, 1945m:419, fig. 4 (male hypopygium). Type-locality: Bolivia, Beni, Ivón. Distr. — Bolivia.
- MEDIOCOPRESSA Alexander, 1944k:589. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador, Peru.
- MEGALYRA Alexander, 1945m:436, fig. 24 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru, Ecuador.
- MONOCERA Alexander, 1967b:455, fig. 5 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- MULFORDI Alexander, 1945d:259. Type-locality: Bolivia, Beni, Cavinás. Distr. — Bolivia. Ref. — Alexander, 1945k:342, fig. 4 (male hypopygium).
- MULTIMODA Alexander, 1938d:328. Type-locality: Colombia, Puerto Boyaca, Magdalena Basin. Distr. — Colombia.
- MYRIATRICHA Alexander, 1942b:344, fig. 5 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- NEOLENTA Alexander, 1945k:334 (nom. nov. for *lenta* Alexander). Type-locality: Ecuador, Los Rios, Playas de Montalvo. Distr. — Ecuador. Ref. — Alexander, 1945m:436, fig. 22 (male hypopygium).
lenta Alexander, 1941d:112, fig. 4 (male hypopygium) (preocc. Harris, 1776). Type-locality: Ecuador, Los Rios, Playas de Montalvo.
- NICOYA Alexander, 1944h:377. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- NIGROABDOMINALIS (Alexander), 1936a:12 (*Habromastix*). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- NIGROVARIEGATA Alexander, 1928f:627. Type-locality: Ecuador, Guayas, Bucay, Ventura. Distr. — Ecuador.
- NIOBE Alexander, 1947f:75, fig. 1 (male hypopygium). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- OPIPARA Alexander, 1951b:294, fig. 14 (male hypopygium). Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.



- ORTONI Alexander, 1945d:261. Type-locality: Bolivia, Beni, Cavinás. Distr. — Bolivia. Ref. — Alexander, 1945m:425, fig. 11 (male hypopygium).
- PALA Alexander, 1939c:429, figs. 1 (wing), 9, 10 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PALAEOGAMA Alexander, 1944d:307 [*Tipula (Nephrotomodes)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1951b:292, fig. 15 (male hypopygium).
- PALOIDES Alexander, 1940f:393, fig. 15 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PARALENTA Alexander, 1950a:35. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- PARARIA Alexander, 1965b:343, fig. 4 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- PARISHI Alexander, 1912f:355, fig. n (male hypopygium). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- PENANA Alexander, 1951b:273, fig. 2 (male hypopygium). Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- PERANGUSTULA Alexander, 1938h:431. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia, Venezuela.
- PERCOMPRESSA Alexander, 1944k:587. Type-locality: Ecuador, Pichincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- PERCOMPTARIA Alexander, 1945m:433, fig. 20 (male hypopygium). Type-locality: Ecuador, Santiago-Zamora, Río Nanganitza, Mayaico. Distr. — Ecuador.
- PERDELECTA Alexander, 1942b:353 [*Tipula (Nephrotomodes)*]. Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- PERLATICOSTA Alexander, 1941a:240. Type-locality: "Brazil". Distr. — Brazil.
- PLAUMANNINA Alexander, 1945d:258. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1945m:428, fig. 13 (male hypopygium).
- PLUMBETHORAX Alexander, 1921d:101. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- PONTIFEX Alexander, 1967b:456, fig. 6 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- PORRECTA Alexander, 1951b:280, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Tarma, Utuyacu. Distr. — Peru.
- PRETIOSA Alexander, 1945m:428, fig. 14 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PRITCHARDI PRITCHARDI Alexander, 1940b:148. Type-locality: Mexico, Michoacán, Pedregal de Tancitaro. Distr. — Mexico.
- ssp. GLABRISTYLA Alexander, 1946b:226 [*Tipula (Nephrotomodes)*]. Type-locality: Mexico, Michoacán, Pedregal de Tancitaro. Distr. — Mexico.
- PROCTOTRICHA Alexander, 1945m:430, fig. 16 (male hypopygium). Type-locality: Ecuador, Santiago-Zamora, Nanganitza, Mayaico. Distr. — Ecuador.
- PROLIXISTERNA Alexander, 1941d:111, fig. 3 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador. Ref. — Alexander, 1945m:436, fig. 23 (male hypopygium).
- QUADRICOLLIS Alexander, 1945k:354, fig. 9 (male hypopygium). Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.



- RECTANGULUS Alexander, 1951b:289, fig. 12 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- REGRESSA Alexander, 1950a:34, fig. 1 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- RETROSA Alexander, 1962c:68, fig. 43 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SCAPHULA Alexander, 1945m:438, fig. 25 (male hypopygium). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- SCELESTA Alexander, 1945m:415, fig. 1 (male hypopygium). Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- SCHILDEANA Alexander, 1945k:345, fig. 6 (male hypopygium). Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.
- SCHWARZMAIERI Alexander, 1940f:392, figs. 4 (wing), 14 (male hypopygium). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- SEPTEMHASTATA Alexander, 1962c:70, figs. 44, 45 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SEXINCINTA Alexander, 1942i:50, fig. 14 (wing). Type-locality: Brazil, São Paulo, Serra da Cantareira. Distr. — Brazil.
- SMILODON Alexander, 1940d:196, fig. 1 (male hypopygium). Type-locality: Ecuador, Los Ríos, Playas de Montalvo. Distr. — Ecuador. Ref. — Alexander, 1945m:419, fig. 3 (male hypopygium).
- SMITHI Alexander, 1912f:350, figs. f (wing), w (ovipositor). Type-locality: Brazil, Mato Grosso, Chapada [dos Guimarães]. Distr. — Brazil.
- SPARSIPHA Alexander, 1962c:71, fig. 46 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SPINICAUDA Alexander, 1919a:152. Type-locality: Panama, Río Chico near mouth of Río Porcona. Distr. — Panama.
- STERNOHIRSUTA Alexander, 1941d:115, fig. 5 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador. Ref. — Alexander, 1945m:436, fig. 21 (male hypopygium).
- SUBARMATA Alexander, 1914a:247. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay. Ref. — Alexander, 1942g:175, fig. 9 (male hypopygium).
- SUBAYMARA Alexander, 1962c:72, figs. 47-49 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SUBEFFETA Alexander, 1951b:296, fig. 16 (male hypopygium). Type-locality: Peru, Tarapoto, San Martín. Distr. — Peru.
- SUBINFUSCATA Williston, 1896a:295, pl. 10, fig. 69 (wing). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.
- TANCITARO Alexander, 1946b:226, figs. 3 (wing), 6 (male hypopygium) [*Tipula (Nephrotomodes)*]. Type-locality: Mexico, Michoacán, Pedregal de Tancitaro. Distr. — Mexico.
- TEMPERATA Alexander, 1940b:146. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- TENUICULA Enderlein, 1912:7. Type-locality: "Colombia". Distr. — Colombia.
- TENULOBATA Alexander, 1940d:198, fig. 2 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Jatun Yacu. Distr. — Ecuador, Peru.

- TERPISCHORE Alexander, 1951b:277, fig. 5 (male hypopygium). Type-locality: Guatemala, Chicacao, El Naranjo. Distr. — Guatemala.
- TERRIBILIS Alexander, 1944k:591. Type-locality: Ecuador, Pinchincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- TIJUCENSIS Alexander, 1943f:296, figs. 3 (wing), 5, 6 (male hypopygium). Type-locality: Brazil, Guanabara, Rio de Janeiro, Pico da Tijuca. Distr. — Brazil.
- TOPOENSIS Alexander, 1951b:283, fig. 8 (male hypopygium). Type-locality: Ecuador, Tungurahua, Río Topo. Distr. — Ecuador, Peru.
- TRICHOPROCTA Alexander, 1945k:335, fig. 1 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- TRIHASTATA Alexander, 1945k:340, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- TRINIDADENSIS (Alexander), 1912f:346, figs. h (wing), j (male hypopygium) (*Pachyrhina*). Type-locality: Trinidad, Port of Spain. Distr. — Trinidad.
- TRINITATIS Alexander, 1941c:327, figs. 3 (wing), 6 (male hypopygium). Type-locality: Trinidad, Fyzabad. Distr. — Trinidad.
- TUCUMANENSIS Alexander, 1945m:425, fig. 12 (male hypopygium). Type-locality: Argentina, Tucumán, Quebrada Famailla. Distr. — Argentina.
- UROPHORA Alexander, 1938h:429. Type-locality: Ecuador, Los Ríos, Playas de Montalvo. Distr. — Colombia, Ecuador. Ref. — Alexander, 1945m:423, fig. 9 (male hypopygium).
- VIRGILIA Alexander, 1951b:291. Type-locality: Peru, Loreto, Aguaitía. Distr. — Peru.
- VOLENS Alexander, 1944g:286 [*Tipula* (*Nephrotomodes*)]. Type-locality: Peru, Huanuco, Afilador. Distr. — Peru.
- ZETEKI Alexander, 1945k:342, fig. 5 (male hypopygium). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- ZONALIS Alexander, 1926a:164. Type-locality: Venezuela, Táchira, Táchira. Distr. — Venezuela.

Subgenus PECTINOTIPULA Alexander

- PECTINOTIPULA Alexander, 1920b: 9 (as genus). Type-species, *Ozodicera argentina* Wulp (orig. des.) (Neotropical).
- ARGENTINA (Wulp), 1881:147, pl. 15, figs. 1, 2 (*Ozodicera*). Type-locality: "Argentina". Distr. — Argentina. Refs. — Alexander, 1962c:38, fig. 21 (male hypopygium); Bruch, 1939a, pl. 3, fig. 8 (head of male).
- BOLIVIENSIS (Alexander), 1946f:297 (*Pectinotipula*). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia. Ref. — Alexander, 1962c:38, fig. 22 (male hypopygium).
- TITICACAE (Alexander), 1944g:284 (*Pectinotipula*). Type-locality: Peru, Puno, Chucuito. Distr. — Peru, Bolivia. Ref. — Alexander, 1962c:38, fig. 24 (male hypopygium, antenna).
- TUCUMANA (Alexander), 1946f:299 (*Pectinotipula*). Type-locality: Argentina, Tucumán, Quebrada Famailla. Distr. — Argentina. Ref. — Alexander, 1962c:38, fig. 23 (male hypopygium).



Subgenus PTERELACHISUS Rondani

- PTERELACHISUS Rondani, 1842:243 (as genus). Type-species, *bertei* Rondani (mon.) (Western Palaearctic).
Oreomyza Pokorny, 1887:50 (as genus). Type-species, *glacialis* Pokorny (Coquillett, 1910:580).
Tipula, subg. *Geotipula* Svtshenko, 1964:15. Type-species, *variipennis* Meigen (orig. des.).
BELLARDIANA Alexander, 1926a:173. Type-locality: Mexico, Chapingo. Distr. — Mexico.

Subgenus TRICHOTIPULA Alexander

- Tipula*, subg. TRICHOTIPULA Alexander, 1915d:468. Type-species, *oropezoides* Johnson (orig. des.) (Nearctic).
APLECTA Alexander, 1946b:220, figs. 2 (wing), 5 (male hypopygium). Type-locality: Mexico, Michoacán, Pedregal de Tancitaro. Distr. — Mexico.
KRAUSSI Alexander, 1946e:522. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
RELIGIOSA Alexander, 1946a:119. Type-locality: Mexico, México, D.F. Distr. — Mexico.
SELANDERI Alexander, 1956a:24. Type-locality: Mexico, Guanajuato, Irapuato. Distr. — Mexico.
UNORIA Alexander, 1946e:525. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
VULTUOSA Alexander, 1946e:523. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.

Subgenus TRIPLICITIPULA Alexander

- Tipula*, subg. TRIPLICITIPULA Alexander, 1965c:354. Type-species, *triplex* Walker (orig. des.) (Nearctic).
HOOGSTRAALI Alexander, 1940b:144. Type-locality: Mexico, Nuevo León, Cerro Potosí. Distr. — Mexico.

Subgenus YAMATOTIPULA Matsumura

- YAMATOTIPULA Matsumura, 1916:461 (as genus). Type-species, *nohirae* Matsumura (orig. des.) = *nova* Walker (Eastern Palaearctic).
JAMAICENSIS Alexander, 1928g:27, fig. 2 (wing) [*Tipula* (*Tipula*)]. Type-locality: Jamaica, Clarendon. Distr. — Jamaica. Ref. — Alexander, 1964a:15, fig. 4 (wing).
LUDOVICIANA Alexander, 1919c:196. Type-locality: U.S.A., Louisiana. Distr. — Southeastern U.S.A., Cuba.
SAYI Alexander, 1911a:194 (nom. nov. for *costalis* Say). Type-locality: "Eastern North America". Distr. — Eastern North America, Bermuda.
costalis Say, 1823:23 (1859:48) (preocc. Swederus, 1787). Type-locality: "Eastern North America". Ref. — Johnson, 1913:443.
SUBVIRGO Alexander, 1951b:268, fig. 18 (male hypopygium). Type-locality: Mexico, 44 km west of Toluca. Distr. — Mexico.
VIRGO Osten Sacken, 1886a:14. Type-locality: Mexico, Durango, Durango City. Distr. — Mexico.

Genus VALDIVIANA Alexander

- VALDIVIANA Alexander, 1929a:17. Type-species, *edwardsina* Alexander (orig. des.) (Neotropical).
- EDWARDSINA Alexander, 1929a:17, figs. 1 (wing), 113 (male hypopygium). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
- NEUQUENENSIS Alexander, 1962b:338, figs. 2 (wing), 9 (male hypopygium). Type-locality: Argentina, Neuquén, Cerro Malo. Distr. — Chile, Argentina.
- SHANNONINA Alexander, 1929a:19, fig. 32 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- SYNEMPORA Alexander, 1929a:20, fig. 2 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina.

Genus ZELANDOTIPULA Alexander

- Holorusia*, subg. ZELANDOTIPULA Alexander, 1922f:522. Type-species, *Tipula novarae* Schiner (orig. des.) (Australasian: New Zealand). Ref. — Vane-Wright, 1967.
- ACUTISTYLA (Alexander), 1945p:749 [*Holorusia (Holorusia)*]. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- ASSOCIANS (Walker), 1861:333 (*Tipula*). Type-locality: "Southern Mexico". Distr. — Mexico, Guatemala. Ref. — Vane-Wright, 1967:517, fig. 12 (wing), p. 523, figs. 31, 32 (male hypopygium).
guatemalensis Alexander, 1926a:161 [*Holorusia (Holorusia)*]. Type-locality: "Guatemala".
- BISATRA (Alexander), 1962c:19, figs. 2 (wing), 6 (antenna) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- CALVICORNIS (Edwards), 1919:161, fig. 26 (male hypopygium) (*Prionocera*). Type-locality: Ecuador, Carchi, La Rinconada, 3200 meters. Distr. — Ecuador. Ref. — Vane-Wright, 1967:517, fig. 13 (wing).
- CALYPSO (Alexander), 1945p:740 (*Holorusia*). Type-locality: Ecuador, Tungurahua, Río Blanco, near Baños. Distr. — Ecuador.
- CATAMARCENSIS (Alexander), 1920f:219 (*Holorusia*). Type-locality: Argentina, Catamarca, Esquina Grande. Distr. — Argentina.
- CELESTISSIMA (Alexander), 1953e:602 [*Holorusia (Holorusia)*]. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- CHIMBORAZO (Alexander), 1945p:736 [*Holorusia (Holorusia)*]. Type-locality: Ecuador, Chimborazo, El Calera, near San Juan. Distr. — Ecuador.
- CRISTALTA (Alexander), 1946f:296 [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia. Ref. — Alexander, 1962c:22, fig. 14 (male hypopygium).
- CRISTIFERA (Alexander), 1945p:751 [*Holorusia (Holorusia)*]. Type-locality: Ecuador, Santiago-Zamora, Río Nangarico, Mayaico. Distr. — Ecuador.
- CRISTOBTUSA (Alexander), 1946f:295 [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia. Ref. — Alexander, 1962e:22, fig. 15 (male hypopygium).



- FASLIANA (Alexander), 1962c:22, figs. 7 (antenna), 16 (male hypopygium) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- FLAVICORNIS (Alexander), 1914f:352, pl. 15, fig. 2 (wing) (*Holorusia*). Type-locality: "Venezuela". Distr. — Venezuela.
- FLAVOGENUALIS (Alexander), 1945p:742 [*Holorusia (Holorusia)*]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- FORSTERIANA (Alexander), 1962c:24, figs. 8 (antenna), 17 (male hypopygium) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- FURCIFERA (Alexander), 1944h:373 [*Holorusia (Holorusia)*]. Type-locality: Costa Rica, Talamanca, Piedregoso. Distr. — Costa Rica.
- GRACILIPES (Walker), 1836:333 (*Tipula*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Vane-Wright, 1967:519, fig. 16 (wing base), p. 523, fig. 35 (male hypopygium).
- HORNI (Alexander), 1926c:389 (*Holorusia*). Type-locality: Brazil, São Paulo, São Paulo, Butantã. Distr. — Brazil.
- INFERNALIS (Alexander), 1947d:38 [*Holorusia (Holorusia)*]. Type-locality: Venezuela, Aragua, road between Maracay and Choroní. Distr. — Venezuela.
- JUTURNA (Alexander), 1945p:738 [*Holorusia (Holorusia)*]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru, Paraguay.
- LAEVIS (Alexander), 1914f:353, pl. 15, figs. 8 (male hypopygium), 15 (antenna) (*Holorusia*). Type-locality: Paraguay, Asunción. Distr. — Brazil, Paraguay. Ref. — Vane-Wright, 1967:514, fig. 3 (wing), p. 520, fig. 18 (antenna).
- LASSULA (Alexander), 1940a:276 (*Holorusia*). Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- LONGITARSIS (Macquart), 1846:145 (17), pl. 2, fig. 2 (*Tipula*). Type-locality: "Venezuela". Distr. — Colombia, Venezuela.
- LUTEIVENA (Alexander), 1946f:293 [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia. Ref. — Alexander, 1962c:26, fig. 18 (male hypopygium).
- MONOSTICTULA (Alexander), 1945p:745 [*Holorusia (Holorusia)*]. Type-locality: Peru, Huanuco. Distr. — Ecuador, Peru.
- NEUROTRICHIA (Alexander), 1962c:27, fig. 9 (antenna) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, La Paz. Distr. — Bolivia.
- NIGROSETOSA (Alexander), 1945p:744 [*Holorusia (Holorusia)*]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- OROPHILA (Alexander), 1914f:354, pl. 15, fig. 14 (antenna) (*Holorusia*). Type-locality: Colombia, San Lorenzo Mountain. Distr. — Colombia.
- FARVICEPS (Speiser), 1909:56 (*Tipula*; nom. nov. for *microcephala* Wulp). Type-locality: Lesser Antilles: Is. Guadalupe. Distr. — Guadalupe Is.
microcephala Wulp, 1881:150, pl. 15, fig. 4 (*Tipula*; preocc. Bigot, 1858). Type-locality: Lesser Antilles, Guadalupe Is.
vanderwulpi Alexander, 1912f:344 (*Tipula*; unnecessary nom. nov. for *microcephala* Wulp). Type-locality: Lesser Antilles, Guadalupe Is.

- PARVIMACULA (Alexander), 1945d:257 [*Holorusia (Holorusia)*]. Type-locality: Brazil, Guanabara, Rio de Janeiro, Manguinhos. Distr. — Brazil.
- PERSTRANGALIA (Alexander), 1962c:29, fig. 3 (wing). [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- PERUVIANA (Alexander), 1914f:355 (*Holorusia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- FLAGIFERA (Alexander), 1943d:125 [*Holorusia (Holorusia)*]. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- RINGENS (Alexander), 1935a:316 (*Holorusia*). Type-locality: Paraguay, Guaira, Villarrica. Distr. — Brazil, Paraguay.
- SCHINERI (Alexander), 1934b:303 (*Holorusia*; nom. nov. for *eluta* Schiner). Type-locality: "Brazil". Distr. — Guyana, Brazil.
eluta Schiner, 1868:351 (*Tipula*; preocc. Loew, 1863). Type-locality: "Brazil".
- SINUOSA (Alexander), 1926a:160 (*Holorusia*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- SONGOENSIS (Alexander), 1962c:30, fig. 4 (wing) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- STRANGALIA (Alexander), 1926a:162 (*Holorusia*). Type-locality: Venezuela, Táchira, La Fría. Distr. — Venezuela.
- SUBFURCIFERA (Alexander), 1945L:375 [*Holorusia (Holorusia)*]. Type-locality: Honduras, Puerto Castilla. Distr. — Honduras.
- SUBLAEVIS (Alexander), 1938c:322 (*Holorusia*). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- SUBTARDA (Alexander), 1935a:318 (*Holorusia*). Type-locality: Paraguay, Caraveni. Distr. — Paraguay.
- TARDA (Alexander), 1935a:317 (*Holorusia*). Type-locality: Paraguay, Guaira, Villarrica. Distr. — Brazil, Paraguay.
- TRIATRA (Alexander), 1962c:31, figs. 11 (antenna), 19 (male hypopygium) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- TRICHONEURA (Alexander), 1962c:33, fig. 10 (antenna) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- TUBERCULIFERA (Alexander), 1953e:603, fig. 4 (male hypopygium) [*Holorusia (Holorusia)*]. Type-locality: Peru, Tarapoto, San Martín. Distr. — Peru.
- UNIATRA (Alexander), 1945p:734 [*Holorusia (Holorusia)*]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- VIVIDA (Alexander), 1941d:107, fig. 1 (male hypopygium) (*Holorusia*). Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- VULPES (Alexander), 1945d:256 [*Holorusia (Holorusia)*]. Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil.
- YUNGASICOLA (Alexander), 1962c:34, fig. 13 (antenna) [*Holorusia (Holorusia)*]. Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia.
- ZAMORAE (Alexander), 1945p:747 [*Holorusia (Holorusia)*]. Type-locality: Ecuador, Santiago-Zamora, Rio Zamora, Zumbi. Distr. — Ecuador.

Unplaced Tipulinae

- ALBOCOSTATA Macquart, 1846:143 (16) (*Tipula*). Type-locality: Colombia, Santa Fe de Bogotá. Kertész (1902) gives this as "Brazil". Type lost; Vanc-Wright (1967) places it in list of species dubiae, and indicates that it possibly might be an *Ischnotoma* or a *Tipula*.
- ELONGATA Macquart, 1838:54 (50) pl. 4, fig. 2 (whole insect), 2a (head) (*Pachyrhina*). Type-locality: unknown (possibly may be neotropical).
- FILIGERA Walker, 1856:443 (*Tipula*). Type-locality: "South America". Identity uncertain.
- FLAVIPENNIS Philippi, 1865:607 (*Tipula*). Type-locality: Chile, Valdivia. (May belong to *Ischnotoma*). Ref. — Alexander, 1929a:36.
- PRAESES Alexander, 1942j:29 (*Tipula*; nom. nov. for *fumipennis* Alexander). Type-locality: Peru, Pichís and Perené Valleys.
fumipennis Alexander, 1912f:359 (preocc. Hudson, 1892). Type-locality: Peru, Pichís and Perené Valleys.
- QUADRIMACULATA Bellardi, 1859:9, pl. 1, fig. 2 (wing) [1861:209, pl. 1, fig. 3 (wing)] (*Tipula*; a primary homonym of *Tipula quadrimaculata* Linnaeus, 1761). Type-locality: Mexico, Córdoba. Identity uncertain.
- SETICORNIS Macquart, 1846:144 (16) pl. 1, fig. 11 (whole insect) (*Tipula*). Type-locality: "Colombia" (as "New Grenada"). Identity uncertain; *Tipula seticornis* Walker, 1856:443, from Colombia, possibly distinct.
- TABIDA Enderlein, 1912:8 (*Tipula*). Type-locality: "Peru". Identity uncertain: Vane-Wright (1967) suggests its position among species dubiae. Possibly a *Tipula* (*Microtipula*) sp.
- TRIMACULATA Macquart, 1838:59 (55), pl. 1, fig. 2 (wing) (*Tipula*). Type-locality: "Brazil or Chile" (very doubtfully from Chile). Identity uncertain; may be a *Zelandotipula*.
- TRINA Wiedemann, 1824:11 (*Tipula*). Type-locality: "Brazil": Identity uncertain; perhaps a *Zelandotipula*.

Subfamily CYLINDROTOMINAE

Genus STIBADOCERINA Alexander

- STIBADOCERINA Alexander, 1929a:66. Type-species, *chilensis* Alexander (orig. des.) (Neotropical).
- CHILENSIS Alexander, 1929a:67, fig. 37 (wing). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile.

Subfamily LIMONIINAE

Tribe LIMONINI

Genus ANTOCHA Osten Sacken

- ANTOCHA Osten Sacken, 1859:219. Type-species, *saxicola* Osten Sacken (mon.). (Nearctic). Refs. — Alexander, 1921f:799 (biology); 1967a.

MONTICOLA Alexander, 1917a:23. Type-locality: U.S.A., Colorado, Platte Canyon. Distr. — Western Nearctic region southward to Mexico (Cuernavaca, Morelos).

Genus DICRANOPTYCHA Osten Sacken

- DICRANOPTYCHA Osten Sacken, 1859:217. Type-species, *germana* Osten Sacken (Coquillett, 1910:533) (Nearctic). Ref. — Alexander, 1919d (biology).
- COSTARICENSIS Alexander, 1939e:622, figs. 1 (wing), 9 (male hypopygium). Type-locality: Costa Rica. Talamanca, San Juan Miramar. Distr. — Costa Rica.
- HARPYIA Alexander, 1946b:232, figs. 9 (wing), 13 (male hypopygium). Type-locality: Mexico, Michoacán, La Majada. Distr. — Mexico.
- LEUCOPODA Alexander, 1953d:156. Type-locality: Guatemala. Chicacao. El Naranjo. Distr. — Guatemala.

Genus HELIUS Lepeletier & Serville

- Megarhina* Lepeletier & Serville in Latreille *et al.*, 1828:585. Type-species, *Limnobia longirostris* Meigen (mon.).
- HELIUS Lepeletier & Serville in Latreille *et al.*, 1828:831 (unjustified nom. nov. for *Megarhina* Lepeletier & Serville; as *Hélius*). Type-species, *Limnobia longirostris* Meigen (aut.) (Western Palaearctic). [See note in Alexander, 1965d:50].
- Rhamphidia* Meigen, 1830:281. Type-species, *Limnobia longirostris* Meigen (Westwood, 1840:129). Ref. — Alexander, 1967a:344.
- ACANTHOSTYLUS Alexander, 1944b:167, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- ALBITARSIS ALBITARSIS (Osten Sacken), 1887b:184 (*Rhamphidia*). Type-locality: "Puerto Rico". Distr. — Colombia, Guyana, Venezuela, Cuba, Jamaica, Puerto Rico. Refs. — Alexander, 1932:366, fig. 5 (wing); 1964a:16, fig. 5 (wing).
- ssp. FUMPENNIS (Alexander), 1919a:137, fig. 5 (wing) (*Rhamphidia*). Type-locality: Panama, Isla de Colón, Bocas del Toro, Silico Creek. Distr. — Panama.
- ssp. FUSCIPES (Alexander), 1919a:137, fig. 4 (wing) (*Rhamphidia*). Type-locality: Panama, Bocas del Toro, San San. Distr. — Costa Rica, Panama.
- ALBOGENICULATUS Alexander, 1936d:751. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- ANGUSTALBUS Alexander, 1953d:154. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru, Bolivia. Ref. — Alexander, 1967a:345, fig. 69 (wing).
- ARAUCARIAE Alexander, 1945f:15, figs. 6 (male hypopygium), 8 (wing). Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- BITERGATUS Alexander, 1950a:45, figs. 12 (wing), 15 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- CAPNIOPTERUS Alexander, 1945e:222. Type-locality: Brazil, Guanabara, Rio de Janeiro, Tinguá. Distr. — Brazil.

- CREPER Alexander, 1926b:225. Type-locality: Jamaica, Troy. Distr. — Jamaica. Ref. — Alexander, 1964a:17, fig. 6 (wing).
- DISTINERVIS Alexander, 1940c:112. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- FRAGOSUS Alexander, 1937b:238. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- HAUSCHBACHII Alexander, 1954a:35. Type-locality: Brazil, Paraná, Paranguá, Matinhos. Distr. — Brazil.
- INEPTUS Alexander, 1938g:433. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- INVAREGATUS Alexander, 1938g:436. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- LAROTYPA Alexander, 1926d:379. Type-locality: Paraguay, St. Catharine, Santa Bárbara. Distr. — Brazil, Paraguay.
- LEUCOPLACA Alexander, 1936d:752. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- LOBULIFERUS Alexander, 1938g:434. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- MELANOPHALLUS Alexander, 1938g:435. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- MICRACANTHUS Alexander, 1945j:16, fig. 7 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- MIRABILIS (Alexander), 1914e:230, pl. 3, fig. 5 (wing) (*Rhamphidia*). Type-locality: Guyana, Bartica. Distr. — Guyana, Peru, Brazil.
- MIRANDA (Alexander), 1921d:54 (*Rhamphidia*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- MULTIVOLUS Alexander, 1945d:276. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- MYRSIELLUS Alexander, 1935a:322. Type-locality: Venezuela, between Mau-rukau (?) and Arabopó. Distr. — Venezuela, Guyana, Brazil.
- PALLIDIPES Alexander, 1926d:379. Type-locality: Paraguay, St. Catharina, Santa Bárbara. Distr. — Brazil, Paraguay.
- PANAMENSIS PANAMENSIS Alexander, 1934a:63. Type-locality: Panama, Chiriqui, Finca Parada. Distr. — Panama, Venezuela.
- ssp. LATERALIS Alexander, 1938g:437. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- PARVIDENS Alexander, 1944b:164. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PERPALLIDUS Alexander, 1942g:177. Type-locality: Brazil, Rio de Janeiro, Serra do Itatiaia, "Südabhang". Distr. — Brazil.
- PERVENUSTUS Alexander, 1953d:153. Type-locality: Costa Rica, La Suiza. Distr. — Costa Rica. Ref. — Alexander, 1967a:346, figs. 70 (wing), 73, 74 (male hypopygium).
- PHASMATIS Alexander, 1945b:56. Type-locality: Hispaniola: Dominican Republic. Distr. — Dominican Republic.
- PLEBEIUS Alexander, 1954o:592. Type-locality: Ecuador, Santiago-Zamora, Zamora. Distr. — Ecuador.
- PRODUCTELLUS Alexander, 1944b:165, fig. 6 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- QUADRIFIDUS Alexander, 1926a:161, fig. 3 (wing). Type-locality: Mexico, Chiapas. Distr. — Mexico. Ref. — Alexander, 1967a:346, fig. 75 (male hypopygium).

- RECTISPINA Alexander, 1947d:43, fig. 2 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- RECTUS Alexander, 1945e:224. Type-locality: Brazil, Guanabara, Rio de Janeiro, Tinguá. Distr. — Brazil.
- REGIUS Alexander, 1944b:162, fig. 5 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- RUBICUNDUS (Alexander), 1922c:37 (*Rhamphidia*). Type-locality: Paraguay, San Bernardino. Distr. — Paraguay, Brazil, Bolivia.
- SANGUINOLENTUS (Alexander), 1921d:55 (*Rhamphidia*). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- SCHILDI Alexander, 1945b:54. Type-locality: Costa Rica, San Mateo, Híguito. Distr. — Costa Rica.
- TETRACRADUS Alexander, 1967a:346, figs. 71 (wing), 76 (male hypopygium). Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- UNIFORMIS (Alexander), 1914e:230, pl. 3, fig. 4 (wing) (*Rhamphidia*). Type-locality: Guyana, Bartica. Distr. — Guyana, Peru, Brazil.

Genus LIMONIA Meigen

- Amphinome* Meigen, 1800:15 (preocc. Bruguière, 1792). Type-species, *Tipula tripunctata* Fabricius (Coquillett, 1910:505). Suppressed by I.C.Z.N., 1963:339.
- LIMONIA Meigen, 1803:262. Type-species, *Tipula tripunctata* Fabricius (Westwood, 1840:129) (Western Palaearctic).
- Limnobia* Meigen, 1818:116 (unjustified nom. nov. for *Limonia* Meigen). Type-species, *Tipula tripunctata* Fabricius (aut.). Ref. — Alexander, 1950b:161, figs.; 1967L:279, figs.

Subgenus ATYPOPTHALMUS Brunetti

- ATYPOPTHALMUS Brunetti, 1911:273 (as genus). Type-species, *holopticus* Brunetti (Brunetti, 1912:408) = *umbrata* (Meijere). (Oriental).
- Limonia*, subg. *Atypophthalmus*; Alexander, 1967a:280.
- UMBRATA (Meijere), 1911b:25, fig. 7 (wing) (*Dicranomyia*). Type-locality: Java, Batavia. Distr. — Mexico, Peru, Cuba; Palaetropical, including Hawaii. Refs. — Meijere, 1916:197 (description of immature stages); Byers, 1966:708, pl. (male hypopygium).
- fissilis* Alexander, 1926a:159. Type-locality: Mexico, Vera Cruz, Tres Valles.

Subgenus CAENOGLOCHINA Alexander

- Limonia*, subg. CAENOGLOCHINA Alexander, 1964a:31, figs. 20 (wing), 29 (male hypopygium). Type-species, *apicata* Alexander (orig. dcs.) (Neotropical). Ref. — Alexander, 1967a:280.
- ACUMINATA (Alexander), 1921d:51, fig. 9 (male hypopygium) (*Dicranomyia*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- APICATA APICATA (Alexander), 1914e:225, pl. 3, fig. 1 (wing) (*Dicranomyia*). Type-locality: Guyana, Bartica. Distr. — Mexico, Panama, Venezuela, Guyana.

- ssp. *DOMINICENSIS* Alexander, 1939d:93, fig. 13 (male hypopygium) [*Limonia*]. Type-locality: Lesser Antilles: Dominica, Layou River, on Hatton Garden Trail. Distr. — Dominica.
- BASISTYLATA* Alexander, 1928g:19 (*Limonia*). Type-locality: Jamaica, Cinchona. Distr. — Jamaica. Ref. — Alexander, 1964a:31, figs. 20 (wing), 29 (male hypopygium).
- CAPTONIUS* Alexander, 1945c:259, figs. 8 (wing), 11 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- EGAE* (Alexander), 1921d:49, fig. 10 (male hypopygium) (*Dicranomyia*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- FIELDI* Alexander, 1967a:281, figs. 1 (wing), 4 (male hypopygium). Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- HOFFMANI* Alexander, 1927c:265 [*Limonia (Limonia)*]. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Cuba, Puerto Rico. Ref. — Alexander, 1932:356, figs. 4 (wing), 6 (male hypopygium).
- MYCTERA* Alexander, 1967a:282, fig. 5 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- NAPOENSIS* (Alexander), 1921d:50, fig. 11 (male hypopygium) (*Dicranomyia*). Type-locality: Peru, Loreto, Napo River. Distr. — Peru.
- PAUCIOBATA* Alexander, 1940c:111 [*Limonia (Limonia)*]. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- PUGNAX* Alexander, 1946f:300 [*Limonia (Limonia)*]. Type-locality: Surinam, 30 miles south of Paramaribo. Distr. — Surinam.
- RAPAX* (Alexander), 1921d:50, fig. 12 (male hypopygium) (*Dicranomyia*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- SCAENALIS* Alexander, 1951a:100 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- SICA* Alexander, 1941g:346 (also 1946h:92) [*Limonia (Limonia)*]. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- SINGULARIS* Alexander, 1944k:598 [*Limonia (Limonia)*]. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- SOMNIFICA* Alexander, 1943e:204 [*Limonia (Limonia)*]. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- SUBACUMINATA* Alexander, 1967a:284, fig. 6 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- VORAX* Alexander, 1941g:345 (also 1946h:91) [*Limonia (Limonia)*]. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.

Subgenus CAENOLIMONIA Alexander

- Limonia*, subg. *CAENOLIMONIA* Alexander, 1967a:285. Type-species, *neorepanda* Alexander (orig. des.) (Neotropical).
- AMARYLLIS* Alexander, 1944j:149, fig. 1 (wing) [*Limonia (Limonia)*]. Type-locality: Venezuela, Mérida, Zea. Distr. — Venezuela.
- BRACHYCANTHA* Alexander, 1945d:264 [*Limonia (Limonia)*]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1945L:378, fig. 2 (male hypopygium).
- COMBOSTENA* Alexander, 1967a:287, figs. 2 (wing), 7 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.

- CONTRADISTINCTA Alexander, 1936d:752 [*Limonia (Limonia)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- GALBIPES Alexander, 1967a:288, fig. 3 (wing). Type-locality: Ecuador, Napo-Pastaza, Río Pastaza, Mapoto. Distr. — Ecuador.
- INTERSTITIALIS Alexander, 1940a:283 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Río Pastaza, Mapoto. Distr. — Ecuador.
- MECONEURA Alexander, 1962c:97, figs. 70 (wing), 73 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- MELAXANTHIA Alexander, 1934a:56. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.
- MERIDENSIS Alexander, 1940e:897, fig. 1 (wing). Type-locality: Venezuela, Mérida. Distr. — Venezuela.
- NEOREPANDA Alexander, 1964b:282 (nom. nov. for *repanda* Alexander). Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador. Refs. — Alexander, 1954c:240, fig. 13 (male hypopygium); 1967a:286, figs. 2, 3, 7.
- repanda* Alexander, 1928f:637 (preocc. Edwards, 1923). Type-locality: Ecuador, Chimborazo, Pallatanga.
- ORTHOAGONIA Alexander, 1945a:18 [*Limonia (Neolimnobia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1954c:244, fig. 17 (male hypopygium).
- OSTERHOUTI (Alexander), 1912b:336, fig. q (wing) (*Furcomyia*). Type-locality: Cuba, Pinar del Río. Distr. — Panama, Cuba.
- PAPRZYCKII Alexander, 1941f:326, figs. 1 (wing), 2 (male hypopygium) [*Limonia (Neolimnobia)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- TRANSLUCIDA TRANSLUCIDA (Alexander), 1912b:335, fig. r (wing) (*Furcomyia*). Type-locality: Panama, Bocas del Toro. Distr. — Panama.
- ssp. NIGROTINCTA Alexander, 1945L:378 [*Limonia (Neolimnobia)*]. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- XANTHOMELA Alexander, 1962c:75, fig. 50 (wing) [*Limonia (Limonia)*]. Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.

Subgenus DICRANOMYIA Stephens

- Furcomyia* Meigen, 1818:133. Type-species, *Limonia lutea* Meigen (Coquillett, 1910:546). Unavailable name, cited in specific synonymy.
- DICRANOMYIA Stephens, 1829:53 (also 1829b:243) (as genus). Type-species, *Limnobia modesta* Meigen (Coquillett, 1910:533) (Western Palaearctic). Ref. — Alexander, 1967a:289, figs. 8-30.
- ACUPRODUCTA Alexander, 1962c:84, fig. 59 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- AFFABILIS Alexander, 1952c:42, figs. 3 (wing), 7 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Quebrada de las Casas. Distr. — Chile.
- ALBOAPICALIS Alexander, 1928h:360. Type-locality: Chile, Valparaíso, Perales de Marga-Marga. Distr. — Brazil, Chile, Argentina.
- ALFAROI (Alexander), 1922a:2 (*Dicranomyia*). Type-locality: Costa Rica, San José. Distr. — Mexico, Costa Rica, Venezuela, Bolivia.



- ALTANDINA Alexander, 1956b:111. Type-locality: Peru, Cuzco, La Raya. Distr. — Peru, Bolivia. Refs. — Alexander, 1962c:85, figs. 57 (wing), 60 (male hypopygium); 1967a:293, figs. 8 (wing), 10 (male hypopygium).
- AMBIGUA Alexander, 1929a:99, figs. 43 (wing), 185 (male hypopygium). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
- AMPHIONIS Alexander, 1952c:44, fig. 6 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Plazoleta del Yunque. Distr. — Chile.
- ANAX Alexander, 1942a:222, fig. 2 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- ANDICOLA (Alexander), 1912b:362, fig. h (wing) (*Furcomyia*). Type-locality: Bolivia, San Antonio. Distr. — Colombia, Bolivia. Ref. — Alexander, 1962c:85, fig. 61 (male hypopygium).
- ANDINALTA Alexander, 1956b:112. Type-locality: Peru, Cuzco, La Raya. Distr. — Peru. Ref. — Alexander, 1967a:203, fig. 11 (male hypopygium).
- APPOSITA Alexander, 1942a:230, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- ARGENTINA (Alexander), 1912b:363, fig. k (wing) (*Furcomyia*). Type-locality: Argentina, Neuquén. Distr. — Argentina.
- ARIADNE Alexander, 1942a:225. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- ATRITARSIS Alexander, 1949b:473 [*Límonia* (*Límonia*)]. Type-locality: Brazil, São Paulo, Pindamonhangaba. Distr. — Brazil.
- ATROSTYLA Alexander, 1942a:235, fig. 9 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- AURANTIOTHORAX Alexander, 1941g:348 (also 1946h:95) Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Panama.
- AXIERASTA Alexander, 1952c:45, figs. 4 (wing), 5 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- BICOMIFERA Alexander, 1943d:128, fig. 1 (male hypopygium). Type-locality: Venezuela, Mérida, Chachopo. Distr. — Venezuela.
- BIGLADIA Alexander, 1942f:760, fig. 1 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- BOLIVIANA Alexander, 1930c:723. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia, Peru.
- BORINQUENIA Alexander, 1968c:210. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- BREVICUBITALIS Alexander, 1947f:87, figs. 7 (wing), 8 (male hypopygium). Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela, Bolivia.
- BREVIPLADIA Alexander, 1962c:87, fig. 62 (male hypopygium). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- BREVIVENA BREVIVENA (Osten Sacken), 1869:66 (*Dicranomyia*). Type-locality: U.S.A.: District of Columbia. Distr. — U.S.A. (New York, District of Columbia), Bolivia.
- ssp. CAPRA Alexander, 1947e:57, figs. 1 (wing), 5 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

- ssp. *TORRIDA* Alexander, 1932:358. Type-locality: Puerto Rico, Puerto Real, Vieques Is. Distr. — Jamaica, Puerto Rico.
- BULLOCKIANA* Alexander, 1938i:277. Type-locality: Chile, Toitén. Distr. — Chile.
- CALLIERGON* *CALLIERGON* Alexander, 1939d:109. Type-locality: Hispaniola: Haiti, La Selle Range, La Visite. Distr. — Haiti.
- ssp. *POLYGRAPHIA* Alexander, 1939d:111. Type-locality: Hispaniola: Dominican Republic, Valle Nuevo, southeast of Constanza. Distr. — Dominican Republic.
- CATAMARCANA* Alexander, 1929d:769 [*Limonia* (*Limonia*)]. Type-locality: Argentina, Catamarca, Quebrada Volcan at Agua Amarillo. Distr. — Argentina.
- CAUTINENSIS* Alexander, 1941a:251. Type-locality: Chile, Malleco, Curacautin. Termas Río Blanco. Distr. — Chile.
- CERBEREANA* Alexander, 1929a:87, figs. 16 (wing), 177 (male hypopygium) [*Limonia* (*Limonia*)]. Type-locality: Argentina, Río Negro, L. Correntoso. Distr. — Argentina.
- CHILENSIS* (Alexander), 1913b:487 (*Furcomomyia*; nom. nov. for *guttata* Philippi). Type-locality: Chile, Valdivia and Aconcagua.
- guttata* Philippi, 1865:613 (*Limnobia*; preocc. Meigen, 1838). Type-locality: Chile, Valdivia and Aconcagua. Ref. — Alexander, 1929a:94, figs. 21 (wing), 183 (male hypopygium).
- CHIMBORAZICOLA* Alexander, 1945n:395. Type-locality: Ecuador, Chimborazo, Mt. Chimborazo. Distr. — Ecuador.
- CHLOROTICA* (Philippi), 1865:614 (*Limnobia*). Type-locality: "Chile". Identity uncertain.
- CLARIBASIS* Alexander, 1945d:267. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1945L:380, fig. 7 (male hypopygium).
- CLAVIGERA* Alexander, 1931a:109. Type-locality: Chile, Malleco, Angol. Distr. — Chile.
- CLAVISTYLA* Alexander, 1956b:113. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru. Ref. — Alexander, 1967a:294, figs. 9 (wing), 12 (male hypopygium).
- COIHERI* Alexander, 1964a:38, figs. 31 (wing), 36 (male hypopygium). Type-locality: Jamaica, Portland, Ecclesdown. Distr. — Jamaica.
- COMMUNA* Alexander, 1967a:294, fig. 14 (male hypopygium). Type-locality: Chile, Osorno, Antillanca. Distr. — Chile.
- COMPLACITA* Alexander, 1942f:768, fig. 5 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- CONTRISTANS* Alexander, 1945p:756. Type-locality: Peru, Ayacucho, Huanta. Distr. — Peru.
- CROCIAPICALIS* Alexander, 1962c:88, fig. 65 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- DAMPEI* (Alexander), 1925a:346, fig. 6 (wing) (*Dicranomyia*). Type-locality: Mexico, México, D.F., Desierto de los Leones. Distr. — Mexico.
- DISSOLUTA* Alexander, 1929a:108, fig. 28 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina, Chile.

- DISTANS (Osten Sacken), 1859:211 (*Dicranomyia*). Type-locality: U.S.A., Florida. Distr. — U.S.A. (Florida), British Honduras, Venezuela, Brazil, Paraguay, Jamaica, Puerto Rico. Ref. — Alexander, 1932:358, fig. 7 (male hypopygium).
- mediatlantica* Freeman, 1962:77, fig. 1 (male hypopygium). Type-locality: Gough Is., South Atlantic. Distr. — Gough Is.
- parishi* Alexander, 1914e:226 (*Dicranomyia*). Type-locality: Guyana. Bartica. Distr. — Guyana.
- tricornis* Alexander, 1925a:344, figs. 3 (wing), 15 (male hypopygium) (*Dicranomyia*). Type-locality: Mexico, México, D.F. Distr. — Mexico.
- DIURA Alexander, 1942a:229, fig. 5 (male hypopygium). Type-locality: Peru, Jurín, Tarma, Huacapistana. Distr. — Peru.
- DIVERSIGLADIA DIVERSIGLADIA Alexander, 1942f:769, fig. 6 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru, Bolivia.
- ssp. *PIABELIS* Alexander, 1944c:42. Type-locality: Venezuela, Mérida, Páramo La Negra. Distr. — Venezuela.
- DIVISA Alexander, 1929f:247 (nom. nov. for *diversa* Osten Sacken). Type-locality: U.S.A., District of Columbia. Distr. — U.S.A., Haiti, Jamaica, Puerto Rico. Ref. — Alexander, 1932:359, fig. 8 (male hypopygium).
- diversa* Osten Sacken, 1859:212 (*Dicranomyia*; preocc. Osten Sacken, 1859:207). Type-locality: U.S.A., District of Columbia.
- DOLEROSA Alexander, 1931b:8, fig. 3 (wing). Type-locality: Brazil. Mt. Roraima. Distr. — Brazil.
- DORSOLORATA Alexander, 1945n:398. Type-locality: Ecuador, Chimborazo, Riobamba. Distr. — Ecuador. Ref. — Alexander, 1967a:296, fig. 13 (male hypopygium).
- ELEGANTULA (Alexander), 1913c:195, pl. 5, fig. 2 (wing) (*Dicranomyia*). Type-locality: Colombia, Cauca, Almaguer. Distr. — Colombia.
- ELNORA Alexander, 1929a:107, fig. 27 (wing). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- ELQUENSIS (Blanchard), 1852:341 (*Limnobia*). Type-locality: Chile, Coquimbo, La Serena. Distr. — Chile. Subgenus incertum.
- EMPELA Alexander, 1962c:89, fig. 58 (wing). Type-locality: Bolivia, Cuticuco, Yungas de Songo Valley. Distr. — Bolivia.
- ERRABUNDA Alexander, 1929a:102, figs. 23 (wing), 189 (male hypopygium). Type-locality: Chile, Santiago. Distr. — Chile.
- EXAETA (Alexander), 1927d:305 (*Dicranomyia*). Type-locality: Mexico, Chiapas, near summit of Zontehuitz. Distr. — Mexico.
- EXERCITA Alexander, 1929a:87, figs. 41 (wing), 178 (male hypopygium) [*Limonia* (*Limonia*)]. Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- EXTRANEA Alexander, 1944f:25, figs. 2 (wing), 5 (male hypopygium). Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- FALCIGUIA Alexander, 1962c:90, fig. 66 (male hypopygium). Type-locality: Bolivia, Cuticuco, Yungas de Songo Valley. Distr. — Bolivia.
- FARRI Alexander, 1964a:40, fig. 32 (wing). Type-locality: Jamaica, St. Andrew, Cinchona. Distr. — Jamaica.

- FILICAUDA (Alexander), 1925a:345, fig. 4 (wing) (*Dicranomyia*). Type-locality: Mexico, México, D.F., Desierto de los Leones. Distr. — Mexico.
- FLAVAPERTA Alexander, 1941a:252. Type-locality: Chile, Malleco. Curacautín, Termas Río Blanco. Distr. — Chile.
- FLAVIDA (Philippi), 1865:612 (*Limnobia*). Type-locality: Chile, Valdivia. Corral. Distr. — Chile. Identity uncertain.
- FLAVOFASCIALIS (Alexander), 1924e:61 (*Dicranomyia*). Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- GALAPAGOENSIS Alexander, 1962d:1, figs. 1 (wing), 2 (male hypopygium). Type-locality: Ecuador, Galápagos Is., Santa Cruz, Finca Castro. Distr. — Ecuador.
- GIBBERA (Alexander), 1916a:6 (*Dicranomyia*). Type-locality: Peru, Lima. Distr. — Peru.
- GLOBULICORNIS (Alexander), 1924a:63 (*Dicranomyia*). Type-locality: Argentina, Jujuy, Tilcara. Distr. — Argentina. Ref. — Bruch, 1939a: pl. 5, fig. 16 (entire male) (as *globulithorax*, error).
- GLORIOSA (Alexander), 1912b:337, fig. j (wing) (*Furcomyia*). Type-locality: Guatemala, Totonicapán. Distr. — Guatemala.
- HALOPHILA Alexander, 1929a:91, fig. 18f (male hypopygium) [*Limonia* (*Limonia*)]. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- HARPAX Alexander, 1952c:46, figs. 8 (male hypopygium), 11 (wing). Type-locality: Chile, Juan Fernández, Masatierra, Quebrada de Villagra. Distr. — Chile.
- HIRSUTISSIMA Alexander, 1962c:92, fig. 63 (wing). Type-locality: Bolivia, Chacaltaya, Cordillera Real. Distr. — Bolivia.
- HOSTICA Alexander, 1938a:24 [*Limonia* (*Limonia*)]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- HUMEROSA Alexander, 1942f:762, fig. 2 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- IGNARA Alexander, 1967a:296, figs. 15 (wing), 18 (male hypopygium). Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- ILLEPIDA Alexander, 1944f:22. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- IMITABILIS Alexander, 1942f:766, fig. 4 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru, Bolivia.
- INDEFENSA Alexander, 1939d:112. Type-locality: Hispaniola: Haiti, La Selle Range, La Visite. Distr. — Haiti.
- INFUMATA (Philippi), 1865:613 (*Limnobia*). Type-locality: Chile, Valdivia, Corral. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:105, fig. 25 (wing).
- INGRATA (Alexander), 1928a:104 (*Dicranomyia*). Type-locality: Mexico, Chiapas, Cerro San Cristóbal. Distr. — Mexico.
- INSIGNIFICA (Alexander), 1912b:363, fig. i (wing) (*Furcomyia*). Type-locality: Peru, Iquico. Distr. — Colombia, Peru. Ref. — Alexander, 1913c:197, pl. VI, fig. 1 (male hypopygium).
- INSOLABILIS Alexander, 1946b:234, fig. 16 (wing). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- INVALIDA (Alexander), 1916a:7, pl. 1, fig. 4 (wing) (*Dicranomyia*). Type-locality: Peru, Lima, Matucana. Distr. — Peru.




- ITATIAYANA Alexander, 1944m:308, fig. 7 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- JORGENSENI (Alexander), 1919a:134, fig. 3 (wing) (*Dicranomyia*). Type-locality: Argentina, Catamarca, Esquina Grande. Distr. — Argentina.
- JUJUYENSIS JIJUYENSIS (Alexander), 1924a:65 (*Dicranomyia*). Type-locality: Argentina, Jujuy, Tilcara. Distr. — Argentina. Ref. — Alexander, 1962c:93, fig. 67 (male hypopygium).
- ssp. OBTUSIROSTRIS Alexander, 1962c:94, figs. 64 (wing) 68 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- KNABI (Alexander), 1912b:340, figs. m (wing), w (male hypopygium) (*Furcomyia*). Type-locality: Guatemala, Totonicapán. Distr. — Mexico, Guatemala. Ref. — Alexander, 1946b:233, fig. 15 (male hypopygium).
- KUSCHELIANA Alexander, 1952c:47, figs. 9 (male hypopygium), 12 (wing). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- LABECULA Alexander, 1942a:234, fig. 8 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- LAISTES Alexander, 1967a:298, figs. 16 (wing), 19 (male hypopygium). Type-locality: Mexico, Guadalajara, Barranca de Oblatos. Distr. — Mexico.
- LAPAZENSIS Alexander, 1962c:95, figs. 69 (wing), 71 (male hypopygium). Type-locality: Bolivia, La Paz. Distr. — Bolivia.
- LATEBRA Alexander, 1967a:299, figs. 17 (wing), 20 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Caramávida. Distr. — Chile.
- LATISPINA Alexander, 1942a:226, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- LEWISI Alexander, 1964a:41, figs. 33 (wing), 37 (male hypopygium). Type-locality: Jamaica, Portland, Hardwar Gap. Distr. — Jamaica.
- LIBERTA (Osten Sacken), 1859:209 (*Dicranomyia*). Type-locality: Eastern U.S.A. Distr. — Eastern U.S.A., Bermuda. Ref. — Johnson, 1913:443.
- LINEICOLLIS (Blanchard), 1852:341 (*Limnobia*). Type-locality: Chile, Coquimbo, La Serena. Distr. — Chile. Subgenus incertum.
- LIVIDA (Say), 1829:150 (1859 (2):349) (*Limnobia*). Type-locality: "Mexico". Distr. — Mexico. Identity uncertain.
- LONGIVENTRIS (Alexander), 1913c:198, pl. VI, fig. 2 (male hypopygium); pl. V, fig. 3 (wing) (*Dicranomyia*). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia, Venezuela.
- LUTEAPICALIS Alexander, 1962c:96, fig. 72 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- LYDIA Alexander, 1950b:185, fig. 16 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- MALIGNA Alexander, 1945d:265. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- MALITIOSA Alexander, 1942a:232, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.

- MASAFUERAE Alexander, 1952c:48, figs. 13 (wing), 19 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Las Chozas. Distr. — Chile.
- MELANOCERA (Alexander), 1925a:346, figs. 5 (wing), 16 (male hypopygium) (*Dicranomyia*). Type-locality: Mexico. Xochimilco. Distr. — Mexico.
- MERIDICOLA Alexander, 1943d:130, fig. 2 (male hypopygium). Type-locality: Venezuela, Mérida, Apartaderos. Distr. — Venezuela.
- MICROSCOLA Alexander, 1967a:300, fig. 21 (male hypopygium). Type-locality: Chile, Osorno, Pucatrihue. Distr. — Chile.
- MICROSOMA Alexander, 1944f:23, figs. 1 (wing), 4 (male hypopygium). Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- MICROSOMODES Alexander, 1954a:33, fig. 4 (male hypopygium). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil.
- MISERANDA Alexander, 1945c:258, fig. 7 (wing). Type-locality: Brazil, São Paulo, Jiquiá. Distr. — Brazil.
- MISTURA Alexander, 1942i:52, figs. 18 (wing), 21 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Cabeceiras do M'Boy Guaçu. Distr. — Brazil.
- MULLERCUOLA Alexander, 1942f:764, fig. 3 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- MULSA (Alexander), 1916a:5, pl. 1, figs. 1, 2 (wing) (*Dicranomyia*). Type-locality: Peru, Lima, Matucana. Distr. — Peru, Bolivia.
- MUTATA Alexander, 1935a:321. Type-locality: Venezuela, near Mt. Roraima, along Arabopó River. Distr. — Brazil.
- NEFASTA Alexander, 1944b:161, fig. 4 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- NOTHOFAGI Alexander, 1929a:93, figs. 20 (wing), 182 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Chile, Llanquihue. Casa Panque. Distr. — Chile, Argentina.
- OBTUSISTYLUS Alexander, 1925a:347, figs. 7 (wing), 17 (male hypopygium) (*Limonia*). Type-locality: Mexico, Mexico, D.F., Desierto de los Leones. Distr. — Mexico.
- OHLINI (Alexander), 1920e:2 (*Dicranomyia*). Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile, Argentina.
- OMISSA (Alexander), 1912b:340, fig. o (wing) (*Furcomyia*). Type-locality: Guatemala, Aguna. Distr. — Guatemala, Costa Rica.
- OMISSIVENA (Alexander), 1922b:207 (*Dicranomyia*). Type-locality: Argentina, Tucumán, Quebrada Famailla. Distr. — Argentina.
- ONEROSA (Alexander), 1928a:104 (*Dicranomyia*). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- ORNATIPENNIS (Blanchard), 1852:342 (*Limnobia*). Type-locality: Chile, Coquimbo, La Serena. Distr. — Chile. Subgenus incertum.
- PALLIDIIFERGA Alexander, 1942i:51, figs. 17 (wing), 22 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Cabeceiras do M'Boy Guaçu. Distr. — Brazil.
- PAMPOECIA Alexander, 1922a:1 [*Limonia (Limonia)*]. Type-locality: Costa Rica, Tiribi. Distr. — Panama, Venezuela, Ecuador, Peru, Brazil. Ref. — Alexander, 1954c:239, fig. 11 (male hypopygium).



- PATRUELIS (Alexander), 1924a:62 (*Dicranomyia*). Type-locality: Argentina. Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- PEDESTRIS Alexander, 1952c:49, fig. 10 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Inocentes Bajos. Distr. — Chile.
- PELATES Alexander, 1962c:99, figs. 74 (wing), 78 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- PENANA Alexander, 1956b:114. Type-locality: Peru, Cuzco, El Cuzco. Distr. — Peru. Ref. — Alexander, 1967a:302, figs. 22 (wing), 25 (male hypopygium).
- PENNIGERA Alexander, 1945d:266. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1945L:380, fig. 9 (male hypopygium).
- PERALTA Alexander, 1945n:397. Type-locality: Ecuador, Chimborazo, Cumbres de Titilac. Distr. — Ecuador. Ref. — Alexander, 1967a:302, fig. 26 (male hypopygium).
- PERDISTALIS Alexander, 1943b:108. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PEREXCELSIOR Alexander, 1962c:100, figs. 75 (wing), 79 (male hypopygium). Type-locality: Bolivia, Illimani, "Westhang." Distr. — Bolivia.
- PERFLAVEOLA Alexander, 1927d:304 (*Limonia*). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico, Panama.
- PERPUNCTICOSTA Alexander, 1950b:186 [*Limonia (Limonia)*]. Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.
- PERRTRACTA Alexander, 1942a:221. Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru, Bolivia. Ref. — Alexander, 1962a:103, fig. 76 (wing).
- PERSERENA Alexander, 1946e:528. Type-locality: Mexico, Morelos, Tepoztlán. Distr. — Mexico.
- PERTRUNCATA Alexander, 1962c:103, figs. 77 (wing), 80 (male hypopygium). Type-locality: Bolivia, Yungas de Unduavi. Distr. — Bolivia.
- PIATTA (Philippi), 1865:614 (*Limnobia*). Type-locality: Chile, Valdivia. Distr. — Chile, Argentina. Ref. — Alexander 1929a:90, figs. 18 (wing), 180 (male hypopygium) [*Limonia (Limonia)*].
- stictica* Blanchard, 1852:342 (*Limnobia*; preocc. Meigen, 1818). Type-locality: Chile, Coquimbo, La Serena.
- blanchardi* Alexander, 1913b:488 (*Furcomyia*; unnecessary nom. nov. for *stictica* Blanchard). Type-locality: Chile, Coquimbo, La Serena.
- PIETATIS Alexander, 1943h:486, fig. 1 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- PINODES Alexander, 1929a:102, figs. 44 (wing), 188 (male hypopygium). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- PLUVIALIS PLUVIALIS Alexander, 1929a:95, figs. 22 (wing), 184 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Chile, Concepción. Distr. — Chile, Argentina.
- ssp. CORRENTOSANA Alexander, 1929a:96 [*Limonia (Limonia)*]. Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- ssp. FUSCOLINEATA Alexander, 1929a:96 [*Limonia (Limonia)*]. Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.



- POLYSTICTA (Philippi), 1865:613 (*Limnobia*). Type-locality: Chile, Valdivia. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:89, figs. 17 (wing), 179 (male hypopygium) [*Limonia (Limonia)*].
- POLYSTONYX Alexander, 1968a:21, fig. 3 (male hypopygium). Type-locality: Chile, Chiloé Is., Chaitén. Distr. — Chile.
- PRAECELLENS Alexander, 1944b:156, fig. 2 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- PRAEPOSTERA (Alexander), 1925a:342, figs. 1 (wing), 13 (male hypopygium) (*Dicranomyia*). Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico.
- PRAEVIA Alexander, 1952b:106 [*Limonia (Limonia)*]. Type-locality: Chile, Malleco, Río Blanco. Distr. — Chile.
- PUNOENSIS Alexander, 1945o:588. Type-locality: Peru, Puno. Distr. — Bolivia, Peru. Ref. — Alexander, 1962c:104, fig. 81 (male hypopygium).
- QUADRIGLADIA Alexander, 1944b:159, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- QUADRITUBERCULATA Alexander, 1937d:601. Type-locality: Colombia, Usme, near Bogotá. Distr. — Colombia.
- RAPIDA Alexander, 1946b:233, figs. 10 (wing), 14 (male hypopygium). Type-locality: Mexico, Michoacán, El Soledad. Distr. — Mexico.
- RAVIDA (Alexander), 1925a:343, figs. 2 (wing), 14 (male hypopygium) (*Dicranomyia*). Type-locality: Mexico, Chapingo. Distr. — Mexico.
- REGIFICA (Alexander), 1916a:5, pl. 2, fig. 1 (wing) (*Dicranomyia*). Type-locality: Peru, Lima, Matucana. Distr. — Peru, Bolivia, Argentina. Refs. — Alexander, 1962c:105, fig. 84 (male hypopygium); Bruch, 1939a: pl. 4, fig. 12 (wing).
- REPENTINA Alexander, 1929a:103, fig. 190 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Gutiérrez. Distr. — Argentina.
- RETICULATA (Alexander), 1912b:334, fig. p (wing) (*Furcomyia*). Type-locality: Cuba, Pinar del Río, Sierra Rangel. Distr. — Mexico, Cuba, Jamaica. Ref. — Alexander, 1964a:42, fig. 38 (male hypopygium).
- SANCTAECRUZAE SANCTAECRUZAE (Alexander), 1920b:3 (*Dicranomyia*). Type-locality: Argentina, Santa Cruz, Valle Tuncl. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:104, fig. 24 (wing).
- ssp. IMMACULOSA Alexander, 1929a:105. Type-locality: Chile, Chiloé Is., Castro. Distr. — Argentina, Chile.
- ssp. VANA Alexander, 1929a:105. Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
- SCINDLERI Alexander, 1962c:105, figs. 82 (wing), 85 (male hypopygium). Type-locality: Bolivia, Illimani, "Westhang". Distr. — Bolivia.
- SCIMITAR Alexander, 1942a:228, fig. 4 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Ecuador, Peru.
- SELKIRKI (Alexander), 1920d:25 (*Dicranomyia*). Type-locality: Chile, Juan Fernández, Masatierra. Distr. — Chile. Ref. — Alexander, 1952c:50, figs. 14 (wing), 20 (male hypopygium).
- SEPOSTA Alexander, 1929a:88, fig. 42 (wing) [*Limonia (Limonia)*]. Type-locality: Chile, Valparaíso, Marga-Marga. Distr. — Chile, Argentina.
- SERRATILoba Alexander, 1950a:41, fig. 8 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

- SIBYLLINA Alexander, 1929a:106, fig. 26 (wing). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina.
- SIMILLIMA (Alexander), 1912b:361, fig. n (wing) (*Furcomyia*). Type-locality: Guatemala, Totonicapán. Distr. — Mexico, Guatemala, Venezuela, Peru.
- SIMULANS CONCINNA (Williston), 1893:60 (*Limnobia*). Type-locality: U.S.A., Washington State. Distr. — Western Nearctic, Mexico.
- SMYTHIANA SMYTHIANA Alexander, 1942a:219, fig. 1 (wing). Type-locality: Peru, Chucuito, Lake Titicaca, Puno. Distr. — Peru.
- ssp. CURTISPINA Alexander, 1962c:107, figs. 83 (wing), 88 (male hypopygium). Type-locality: Bolivia, Illimani, "Westhang". Distr. — Peru, Bolivia.
- SPARSITUBER Alexander, 1962c:108, fig. 90 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- SPLENDIDULA Alexander, 1930c:721 [*Limonia (Limonia)*]. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.
- STUARDOI Alexander, 1952c:52, figs. 15 (wing), 21 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- SUBANDICOLA Alexander, 1947e:60, fig. 6 (male hypopygium). Type-locality: Mexico, México, D.F., El Junquito. Distr. — Mexico, Venezuela, Peru, Bolivia.
- SUBANDINA (Alexander), 1913b:488 (*Furcomyia*; nom. nov. for *stigmatica* Blanchard). Type-locality: Chile, Coquimbo, La Serena. Distr. — Chile. Identity uncertain.
- stigmatica* Blanchard, 1852:341 (*Limnobia*; preocc. Meigen, 1830). Type-locality: Chile, Coquimbo, La Serena.
- SUBDOLA (Alexander), 1913d:404, fig. 1 (wing) (*Dicranomyia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- SUBFLAVIDA Alexander, 1929a:99, fig. 186 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Gutiérrez. Distr. — Chile, Argentina. Ref. — Alexander, 1967a:302, fig. 27 (male hypopygium).
- SUBMUTATA Alexander, 1942a:236, figs. 10 (wing), 11 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- SUBRAVIDA (Alexander), 1928a:105 (*Dicranomyia*). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- SUBRETICULATA Alexander, 1943b:107 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1954c:240, fig. 14 (male hypopygium).
- TEINOTERGA Alexander, 1967a:303, figs. 23 (wing), 28 (male hypopygium). Type-locality: Colombia, Mount Redondo. Distr. — Colombia.
- THAMYRIS Alexander, 1950a:39, fig. 6 (male hypopygium) (*Limonia (Limonia)*). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- THIXIS Alexander, 1967a:304, figs. 24 (wing), 29 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Pichinahuel. Distr. — Chile.
- TITICACANA Alexander, 1945d:268. Type-locality: Peru, Puno, Chucuito, near Lake Titicaca. Distr. — Peru, Bolivia. Refs. — Alexander, 1945L:380, fig. 8 (male hypopygium); 1962c:109, figs. 86 (wing), 87 (male hypopygium).

- TORULOSA Alexander, 1968c:211. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- TRAGICA Alexander, 1946a:120 [*Limonia (Limonia)*]. Type-locality: Mexico, Chiapas, Finca Prussia. Distr. — Mexico. Subgenus incertum may be *Neolimonia*.
- TRIOBIFERA Alexander, 1967a:305, fig. 30 (male hypopygium). Type-locality: Chile, Cautín, La Cabana. Distr. — Chile.
- TRINITATIS Alexander, 1931a: 110. Type-locality: Cuba, Trinidad Mts., Buenos Aires. Distr. — Cuba.
- TRITUBERCULATA TRITUBERCULATA Alexander, 1929a:100, fig. 187 (male hypopygium). Type-locality: Chile, Mechuque Is. Distr. — Chile.
- ssp. INGLORIA Alexander, 1929a:101. Type-locality: Argentina, Río Negro, Bariloche. Distr. — Chile, Argentina.
- TROGLOPHILA Alexander, 1929a:92, fig. 19 (wing) [*Limonia (Limonia)*]. Type-locality: Argentina, Río Negro, Grotto de los Leones, 12 km. east of Bariloche. Distr. — Argentina.
- VENATRIX Alexander, 1952c:53, fig. 16 (wing). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile
- VENERIS Alexander, 1952c:54, figs. 17 (wing), 25 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Quebrada de las Casas. Distr. — Chile.
- VENUSTOR Alexander, 1950b:192 [*Limonia (Limonia)*]. Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru, Bolivia.
- VERNALIS (Philippi), 1865:612 (*Limnobia*). Type-locality: Chile, Aconcagua, Catemu. Distr. — Chile. Identity uncertain.
- VILLARICAE Alexander, 1929d:770 [*Limonia (Limonia)*]. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- VIRILIS (Alexander), 1916a:4, pl. 1, fig. 5 (wing) (*Dicranomyia*). Type-locality: Peru, Lima, Matucana. Distr. — Peru.
- WEISERIANA Alexander, 1929d:771. Type-locality: Argentina, Catamarca, Quebrada Volcán, Agua Amarilla. Distr. — Argentina.
- YUNQUEANA Alexander, 1952c:55, fig. 18 (wing). Type-locality: Chile, Juan Fernández, Masatierra, Summit of El Yunque. Distr. — Chile.

Subgenus DISCOBOLA Osten Sacken

- DISCOBOLA Osten Sacken, 1865:226 (as genus). Type-species, *Limnobia argus* Say (Coquillett, 1910:534) = *annulata* (Linnaeus).
- Trochobola* Osten Sacken, 1869:97 (unjustified nom. nov. for *Discobola* Osten Sacken). Type-species, *Limnobia argus* Say (aut.) = *annulata* (Linnaeus). Ref. — Alexander, 1950b:194.
- GOWDEYI Alexander, 1932:384, fig. 26 (male hypopygium). Type-locality: Jamaica, Cinchona, Blue Mts. Distr. — Cuba, Jamaica. Ref. — Alexander, 1964a:32, figs. 19 (wing), 26 (male hypopygium).

Subgenus GERANOMYIA Haliday

- GERANOMYIA Haliday, 1833:154 (as genus). Type-species, *unicolor* Haliday (mon.) (Western Palaearctic).



- Limnobiorrhynchus* Westwood, 1835b:683. Type-species, *brasiliensis* Westwood (Coquillett, 1910:561).
- Aporosa* Macquart, 1839:100. Type-species, *maculipennis* Macquart (Coquillett, 1910:509) = *canariensis* Bergroth.
- Plettusa* Philippi, 1865:597. Type-species, *virescens* Philippi (Coquillett, 1910:592) = *valida* Loew. Refs. — Alexander, 1929f:242; 1953f:77-95, figs. 1-9; 1954b:51-67, figs. 10-32; 1967a:306-313, figs. 31-39.
- AQUABILIS AQUABILIS (Alexander), 1923c:311 [*Geranomyia* (*Geranomyia*)]. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- ssp. DEPLEXA Alexander, 1941f:331. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- AMBLYTYLOS Alexander, 1964a:42, figs. 34 (wing), 39 (male hypopygium). Type-locality: Jamaica, Portland, Eccledown. Distr. — Jamaica.
- AMOENALIS Alexander, 1945d:270. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1945L:380, fig. 10 (male hypopygium).
- ANDUZEANA Alexander, 1943d:133, fig. 3 (male hypopygium). Type-locality: Venezuela, Mérida, Apartaderos. Distr. — Venezuela.
- ANISACANTHA Alexander, 1964a:43, figs. 35 (wing), 40 (male hypopygium). Type-locality: Jamaica, St. Andrew, Bellevue. Distr. — Jamaica.
- ANTHINA Alexander, 1945d:271. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1945L:380, fig. 11 (male hypopygium).
- ARECUNA Alexander, 1931b:11, figs. 5 (wing), 17 (male hypopygium). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela.
- ARGENTINENSIS (Alexander), 1920b:1 (*Geranomyia*). Type-locality: Argentina, Buenos Aires. Distr. — Brazil, Argentina.
- ASSUETA Alexander, 1943b:111. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador. Ref. — Alexander, 1953f:87, fig. 1 (male hypopygium).
- AUSTROANDINA Alexander, 1929d:778. Type-locality: Argentina, Catamarca, Ciénega, near Agua Amarilla. Distr. — Argentina (preocc. by *austroandina* Alexander, 1929f:248).
- AVARA Alexander, 1944m:304, fig. 5 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- BAHIENSIS Alexander, 1930c:724. Type-locality: Brazil, Bahia (Salvador). Distr. — Brazil.
- BANKSIANA Alexander, 1939d:114, fig. 14 (male hypopygium). Type-locality: Cuba, Soledad, near Cienfuegos. Distr. — Cuba.
- SEATRIX Alexander, 1945a:21. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Refs. — Alexander, 1945L:380, fig. 12 (male hypopygium); 1953f:88, fig. 2 (male hypopygium).
- BIARGENTATA Alexander, 1929c:398. Type-locality: Paraguay, Guaira, Villarica. Distr. — Brazil, Paraguay.
- BICINCTA BICINCTA (Alexander), 1921d:44 (*Geranomyia*). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- ssp. ANGUSTICINCTA (Alexander), 1921d:45 (*Geranomyia*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.

- BRASILIENSIS (Westwood), 1835:683 (*Limnobia rhynchus*; part, only male).
Type-locality: Brazil, Bahia vicinity (Salvador). Distr. — Brazil.
- BREVISPINULA Alexander, 1930b:110. Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico.
- BUSTILLOI Alexander, 1938h:435. Type-locality: El Salvador, San Salvador. Distr. — El Salvador. Ref. — Alexander, 1967a:310, fig. 32 (male hypopygium).
- CALLINOTA Alexander, 1941f:318. Type-locality: Peru, Junín, Tarma, Carapata. Distr. — Venezuela, Peru.
- CARUNCULATA CARUNCULATA Alexander, 1941g:349 (also 1946h:95, as 1938). Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Panama, Ecuador, Peru. Ref. — Alexander, 1953f:88, fig. 3 (male hypopygium).
- ssp. MANABIANA Alexander, 1953f:89. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador, Bolivia.
- CERBERUS (Alexander), 1927c:266 (*Geranomyia*). Type-locality: Guatemala, Escuintla. Distr. — Mexico, Guatemala, Costa Rica, Panama.
- CERTHIA (Alexander), 1916d:492 (*Geranomyia*). Type-locality: Guatemala, Antigua. Distr. — Guatemala, El Salvador, Venezuela, Peru, Brazil.
- CINEREINOTA (Alexander), 1913d:407, fig. 4 (wing) (*Geranomyia*). Type-locality: Bolivia, Yungas de Coroico. Distr. — Venezuela, Guyana, Ecuador, Peru, Brazil, Bolivia, Paraguay, Dominican Republic, Jamaica, Puerto Rico. Refs. — Alexander, 1932:362; 1953f:89, fig. 4 (male hypopygium); 1964a:46, fig. 41 (male hypopygium).
- domingensis* Alexander, 1916d:490 (*Geranomyia*). Type-locality: Hispaniola: Dominican Republic, San Francisco Mts.
- CIVICA Alexander, 1939a:70. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico, Jamaica. Ref. — Alexander, 1964a:46, fig. 42 (male hypopygium).
- CONQUISITA Alexander, 1941f:330, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- CONTORTA Alexander, 1941f:320, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- COSTARICENSIS (Alexander), 1916d:494 (*Geranomyia*). Type-locality: Costa Rica, Cartago. Distr. — Costa Rica.
- CUBANA Alexander, 1929e:396. Type-locality: Cuba, Hanabanilla Falls. Distr. — Cuba.
- DAMICOI Alexander, 1942i:54, figs. 19 (wing), 20 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Cabeceiras do M'Bov Guaçu. Distr. — Brazil.
- DELICIOSA Alexander, 1934a:61. Type-locality: Panama, Chiriquí, Finca Parada. Distr. — Panama, Peru.
- DERICTA Alexander, 1940a:287. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Venezuela, Ecuador, Peru. Ref. — Alexander, 1953f:90, fig. 5 (male hypopygium).
- LEVOTA Alexander, 1929f:247 (nom. nov. for *pulchella* Alexander). Type-locality: Guyana, Bartica. Distr. — Guyana.
- pulchella* Alexander, 1914e:228, pl. 3, fig. 3 (wing) (*Geranomyia*; preocc. Mcigen, 1830). Type-locality: Guyana, Bartica.



- DIABOLICA Alexander, 1944L:181, fig. 4 (male hypopygium). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- DIARGYRIA Alexander, 1941f:321, fig. 4 (male hypopygium). Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- DISPARILIS Alexander, 1946e:529. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico, Peru.
- DOMINICANA Alexander, 1939d:94. Type-locality: Lesser Antilles, Dominica, Sylvania Estate. Distr. — Dominica.
- ENDERLEINI Alexander, 1913b:487 (nom. nov. for *annulata* Enderlein). Type-locality: "Costa Rica". Distr. — Costa Rica.
annulata Enderlein, 1912:72, fig. T1 (wing) (*Geranomyia*; preocc. Linnaeus, 1758). Type-locality: "Costa Rica".
- EURYGRAMMA EURYGRAMMA (Alexander), 1928a:107 (*Geranomyia*). Type-locality: Mexico, Veracruz, Córdoba, Barranca of the Río San Antonio. Distr. — Mexico, Panama, south to Ecuador. Refs. — Alexander, 1953f:90, fig. 6 (male hypopygium); 1964a:50, fig. 48 (male hypopygium).
 ssp. STENOMERA Alexander, 1964a:47, figs. 43 (wing), 47 (male hypopygium). Type-locality: Jamaica, St. Andrew, Second Breakfast Spring. Distr. — Jamaica.
- FLUXA Alexander, 1941g:350 (also 1946h:97). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Mexico, Panama, Venezuela, Peru.
- FORSTERIANA Alexander, 1962c:111, fig. 91 (wing). Type-locality: Bolivia, Illimani, "Westhang". Distr. — Bolivia.
- FUROR Alexander, 1944L:183, fig. 5 (male hypopygium). Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela.
- GAUDENS (Alexander), 1922b:208 [*Geranomyia* (*Geranomyia*)]. Type-locality: Argentina, Tucumán, San Pedro de Colalao. Distr. — Argentina.
- GLAUCA (Alexander), 1916a:12, pl. 2, fig. 7 (wing) (*Geranomyia*). Type-locality: Ecuador, Huigra. Distr. — Peru, Ecuador. Ref. — Alexander, 1953f:91, fig. 7 (male hypopygium).
- GUATEMALENSIS (Alexander), 1916d:487 (*Geranomyia*). Type-locality: Guatemala, Aguna. Distr. — Southwestern U.S.A., Mexico, Guatemala, Costa Rica, Venezuela.
- GUIANENSIS Alexander, 1930c:725. Type-locality: Guyana, Koriabo, Barima River. Distr. — Guyana.
- HETEROXIPHA Alexander, 1942f:787, fig. 7 (male hypopygium). Type-locality: Peru, Ayacucho, Sivia. Distr. — Peru.
- HIRSUTINOTA Alexander, 1943b:114. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Peru. Ref. — Alexander, 1953f:91, fig. 8 (male hypopygium).
- IMMERITA Alexander, 1929e:402. Type-locality: Paraguay, Gualra, Villarrica. Distr. — Brazil, Paraguay.
- INAEQUISPINOSA Alexander, 1940a:285. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador, Bolivia. Refs. — Alexander, 1953f:92, fig. 9 (male hypopygium); 1962c:113, fig. 92 (male hypopygium).
- INAEQUITUBERCULATA Alexander, 1929e:401. Type-locality: Paraguay, Gualra, Villarrica. Distr. — Paraguay.


- INFAMOSA Alexander, 1937e:59. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INQUISITA Alexander, 1942f:785. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Ecuador, Peru, Brazil, Bolivia. Ref. — Alexander, 1954b:51, fig. 10 (male hypopygium).
- INSIGNIS (Loew), 1851:395 (*Aporosa*). Type-locality: "Brazil". Distr. — Guyana, Ecuador, Brazil. Ref. — Alexander, 1916d:547, fig. 9 (wing).
- INTERMEDIA (Walker), 1848:47 (*Limnobia*). Type-locality: "Jamaica". Distr. — U.S.A. (Louisiana), Jamaica, Cuba. Refs. — Alexander, 1916d:489, fig. 3 (wing); 1964a:48, figs. 44 (wing), 50 (male hypopygium).
- remingtoni* Alexander, 1946j:247. Type-locality: USA: Louisiana, Orleans Parish, Chalmette. Ref. — Alexander, 1964a:48, figs. 44 (wing), 50 (male hypopygium).
- KNABIANA (Alexander), 1916d:487 (*Geranomyia*). Type-locality: Panama, Canal Zone. Distr. — Guatemala, Panama.
- LACHRYMALIS (Alexander), 1916a:9, pl. 2, fig. 3 (wing) (*Geranomyia*). Type-locality: Ecuador, Napo-Pastaza, Huigra. Distr. — Mexico, Guatemala, Costa Rica, Panama, Ecuador, Peru, Brazil. Refs. — Alexander, 1916d:492, fig. 6 (wing); 1954b:52, fig. 11 (male hypopygium).
- LACTEITARSIS (Alexander), 1922c:34 (*Geranomyia*). Type-locality: Colombia, Condoto. Distr. — Colombia.
- LATITUDINIS Alexander, 1954a:34. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LAUDANDA Alexander, 1938d:330. Type-locality: Colombia, Usme. Distr. — Colombia, Venezuela. Ref. — Alexander, 1943d:133.
- LEMNISCATA Alexander, 1929c:399. Type-locality: Paraguay, Santa Barbara. Distr. — Paraguay.
- LICHYI Alexander, 1943d:136. Type-locality: Venezuela, Mérida, Chachapo. Distr. — Venezuela.
- LINEATA (Enderlein), 1912:73, fig. U1 (wing) (*Geranomyia*). Type-locality: Colombia, Hacienda Pehlke. Distr. — Colombia.
- LUTEIMANA Alexander, 1938g:425. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador, Peru, Brazil. Ref. — Alexander, 1954b:52, fig. 12 (male hypopygium).
- LYCAON Alexander, 1952f:326 (nom. nov. for *pallidula* Alexander). Type-locality: St. Vincent. Distr. — Guyana, St. Vincent, Trinidad.
- pallida* Williston, 1896a:284, pl. 9, fig. 53 (wing) (*Geranomyia*; preocc. Macquart, 1838). Type-locality: St. Vincent.
- pallidula* Alexander, 1929f:247 (nom. nov. for *pallida* Williston; preocc. Abreu, 1923). Type-locality: St. Vincent.
- MACRAUCHENIA Alexander, 1956b:116. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru. Ref. — Alexander, 1967a:311, fig. 33 (male hypopygium).
- MACTA Alexander, 1945d:274. Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil. Ref. — Alexander, 1967a:311, fig. 34 (male hypopygium).



- MARTHAE Alexander, 1930c:726. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Bolivia, Colombia, Venezuela. Ref. — Alexander, 1967a:311, fig. 35 (male hypopygium).
- MEXICANA (Ecllardi), 1861:1 (*Aporosa*). Type-locality: Mexico, Tuxpango, Orizaba. Distr. — Mexico.
- MICROPHAEA Alexander, 1939d:95, fig. 15 (male hypopygium). Type-locality: Lesser Antilles; Dominica, Layou River, on Hatton Garden Trail. Distr. — Dominica.
- MILITARIS Alexander, 1952f:326 (nom. nov. for *viridula* Alexander). Type-locality: Mexico, Quintana Roo, Payo Obispo (=Ciudad Chetumal). Distr. — Mexico, Jamaica. Ref. — Alexander, 1964a:52, fig. 54 (male hypopygium).
- viridula* Alexander, 1930b:110 (preocc. Alexander, 1922h). Type-locality: Mexico, Quintana Roo, Payo Obispo (= Ciudad Chetumal).
- MYERSIANA Alexander, 1929e:397. Type-locality: Cuba, Hababanilla Falls. Distr. — Cuba, Jamaica, Puerto Rico. Refs. — Alexander, 1932:363; 1964a:49, figs. 45 (wing), 49 (male hypopygium).
- NEANTHINA Alexander, 1954d:142. Type-locality: Bolivia, Cochabamba, Chaparc, Locotal, El Limbo. Distr. — Bolivia.
- NEOGAUDENS Alexander, 1942f:784. Type-locality: Peru, Ayacucho, Huanta, Lake Rasvilca. Distr. — Colombia, Peru.
- NEONUMENIUS Alexander, 1930c:729. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- NEOPARILIS Alexander, 1962c:113, fig. 93 (male hypopygium). Type-locality: Bolivia, Cuticuecho, Yungas de Songo Valley. Distr. — Bolivia.
- NEOPENTHERES Alexander, 1930b:109. Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico.
- NIGRIPLEURA (Alexander), 1919a:136, fig. 1 (wing) (*Geranomyia*). Type-locality: Panama, Bocas del Toro, San San. Distr. — Panama.
- NUGATORIA Alexander, 1943b:112. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:53, fig. 13 (male hypopygium).
- NUMENIUS (Alexander), 1913d:406, fig. 3 (wing) (*Geranomyia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Panama, Ecuador, Peru. Ref. — Alexander, 1954b:53, fig. 14 (male hypopygium).
- ONERIS Alexander, 1956b:117. Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru. Ref. — Alexander, 1967a:311, fig. 37 (male hypopygium).
- OPINATOR Alexander, 1950a:42, fig. 16 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- OPULENS Alexander, 1945n:405. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:54, fig. 15 (male hypopygium).
- ORTHORHABDA Alexander, 1940b:152. Type-locality: Mexico, Nuevo León, Horse-Tail Falls. Distr. — Mexico.
- ITALIDAPEX Alexander, 1945n:404. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:54, fig. 16 (male hypopygium).

- PARILIS Alexander, 1945n:407. Type-locality: Ecuador, Tungurahua, Runtun, above Baños. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:55, fig. 17 (male hypopygium).
- PASTAZINA Alexander, 1944d:312. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:55, fig. 18 (male hypopygium).
- PENTHERES (Alexander), 1928a:106 (*Geranomyia*). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- PERFECTA (Alexander), 1928a:109 (*Geranomyia*). Type-locality: Mexico, Nayarit, near Tepic. Distr. — Mexico.
- PLATENSIS (Alexander), 1923c:310 [*Geranomyia* (*Geranomyia*)]. Type-locality: Argentina, La Plata. Distr. — Argentina. Ref. — Bruch, 1939a, pl. 3, fig. 4 (head); pl. 4, fig. 11 (wing); pl. 5, fig. 15 (entire female).
- PLUMBEICOLOR Alexander, 1938c:325. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- PLUMBEIPLEURA (Alexander), 1916a:10, pl. 2, fig. 4 (wing) (*Geranomyia*). Type-locality: Ecuador, Huigra. Distr. — Panama, Colombia, Venezuela, Ecuador, Peru, Brazil, Dominica, Trinidad. Ref. — Alexander, 1954b:56, fig. 19 (male hypopygium).
- PROPERA Alexander, 1949b:476, figs. 9 (wing), 13 (male hypopygium). Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- PROVOCATOR Alexander, 1941f:327, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- RABULA Alexander, 1940a:284. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:57, fig. 20 (male hypopygium).
- RECISA (Alexander), 1927c:267 (*Geranomyia*). Type-locality: El Salvador, Agronomia, Sonsonate. Distr. — Mexico, El Salvador, Panama, Ecuador, Peru, Brazil.
- RECONDITA (Alexander), 1921d:42 (*Geranomyia*). Type-locality: Peru, Loreto, Iquitos. Distr. — Guyana, Panama, Ecuador, Peru, Brazil. Ref. — Alexander, 1954b:57, fig. 21 (male hypopygium).
- gelatifex* Edwards, 1934:633. Type-locality: "Guyana".
- REFUGA Alexander, 1944d:311. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:58, fig. 22 (male hypopygium).
- RELATA Alexander, 1944m:306, fig. 6 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Alto Itatiaia. Distr. — Brazil.
- ROSTRATA ANTILLARUM Alexander, 1929e:395. Type-locality: Cuba, Soledad. Distr. — Cuba, Jamaica, Puerto Rico. Refs. — Alexander, 1932:362; 1964a:49, figs. 46 (wing), 51 (male hypopygium).
- RUBIGINOSA Alexander, 1931b:11, fig. 6 (wing). Type-locality: Venezuela, Mt. Duida. Distr. — Venezuela, Brazil.
- RUFESCENS (Loew), 1851:396 (*Aporosa*). Type-locality: "Puerto Rico". Distr. — Puerto Rico. Ref. — Alexander, 1932:363.
- SATIPOANA Alexander, 1945d:272. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru. Ref. — Alexander, 1945L:380, fig. 13 (male hypopygium).

- SCOLOPAX (Alexander), 1913d:408 (*Geranomyia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Guatcmala, Venezuela, Ecuador, southward to Peru. Refs. — Alexander, 1916a:12, pl. 2, fig. 5 (wing); 1954b:58, fig. 23 (male hypopygium).
- SEPARATA (Alexander), 1921d:45 (*Geranomyia*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Ecuador, Peru, Brazil, Bolivia. Ref. — Alexander, 1954b:59, fig. 24 (male hypopygium).
- SEROTINA (Alexander), 1923c:309 [*Geranomyia (Geranomyia)*]. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- STENOLEUCA Alexander, 1956b:118. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.
- STENOPHALLUS Alexander, 1944d:310. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Venezuela, Ecuador, Peru, Brazil. Ref. — Alexander, 1954b:59, fig. 25 (male hypopygium).
- STOICA Alexander, 1941f:325, fig. 5 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru, Bolivia.
- STYLOBTUSA Alexander, 1967a:311, figs. 31 (wing), 36 (male hypopygium). Type-locality: Peru, Huanuco, Tingo Maria. Distr. — Peru.
- SUBGAUDENS Alexander, 1941f:324. Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- SUBINSIGNIS (Alexander), 1916d:495 (*Geranomyia*). Type-locality: Guatemala, Aguna. Distr. — Mexico, Guatemala, El Salvador, Panama.
- SUBPARILIS Alexander, 1962c:115, fig. 94 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chaparc, Locotal, El Limbo. Distr. — Bolivia.
- SUBPENTHERES Alexander, 1943b:113. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador, Brazil. Ref. — Alexander, 1954b:60, fig. 26 (male hypopygium).
- SUBRECEISA Alexander, 1932:364. Type-locality: Puerto Rico, Vieques Is., Puerto Real. Distr. — Puerto Rico.
- SUBSEROTINA (Alexander), 1921a:72 (*Geranomyia*). Type-locality: Argentina, Jujuy. Distr. — Argentina.
- SUBVIRESCENS SUBVIRESCENS Alexander, 1930b:112. Type-locality: Cuba, Trinidad Mts. Distr. — Cuba, Venezuela.
- ssp. CLEMENTIS Alexander, 1945o:591. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- ssp. JAMAICAE Alexander, 1939d:115. Type-locality: Jamaica, Port Antonio. Distr. — Jamaica. Ref. — Alexander, 1964a:51, fig. 52 (male hypopygium).
- SUMPTUOSA Alexander, 1942f:782. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- SYLVANIA Alexander, 1939d:97, fig. 16 (male hypopygium). Type-locality: Lesser Antilles: Dominica, Sylvania Estate. Distr. — Dominica.
- TATEI Alexander, 1931b:9, figs. 4 (wing), 16 (male hypopygium). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- TIBIALIS (Loew), 1851:397 (*Aporosa*). Type-locality: "Brazil". Distr. — Mexico, Venezuela, Guyana, Galápagos, Peru, Brazil, Bolivia, Argentina, Paraguay, Dominican Republic, Jamaica, Puerto Rico,



- Dominica, Granada Is. Refs. — Alexander, 1916d:491, fig. 5 (wing); 1932:365; 1954b:60, fig. 27 (male hypopygium); 1962d:3, figs. 3 (wing), 4 (male hypopygium); 1964a:52, fig. 53 (male hypopygium).
- TIMENS** Alexander, 1950a:43, figs. 10 (wing), 17 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- TOWNSENDI** (Alexander), 1916a:11, pl. 2, fig. 6 (wing) (*Geranomomyia*). Type-locality: Peru, Lima, Matucana. Distr. — Peru.
- TRANSITORIA** Alexander, 1941f:322. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- TRICHOMERA** Alexander, 1939a:72. Type-locality: Mexico, Chiapas, Finca Belem. Distr. — Mexico.
- TRISTELLA** Alexander, 1929f:247 (nom. nov. for *tristis* Loew). Type-locality: "Brazil". Distr. — Brazil.
tristis Loew, 1851:398 (*Aporosa*; preocc. Schummel, 1829). Type-locality. "Brazil".
- TULUMAYOENSIS** Alexander, 1944g:303, fig. 5 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- TUMIDIBASIS** Alexander, 1938g:426. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador, Brazil, Bolivia. Ref. — Alexander, 1954b:61, fig. 28 (male hypopygium).
- TURBIDA** (Alexander), 1928a:107 (*Geranomomyia*). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- UBERIS** Alexander, 1939a:71. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- UMBRI-COLOR** Alexander, 1937c:491. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador, Peru, Brazil. Ref. — Alexander, 1954b:61, fig. 29 (male hypopygium).
- UNISPINFERA** Alexander, 1938c:326. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- VALIDA** (Loew), 1851:398 (*Aporosa*). Type-locality: "Chile". Distr. — Chile, Argentina. Refs. — Alexander, 1913d:406, fig. 2 (wing); 1929a:71, fig. 39 (wing).
virescens Philippi, 1865:597 (*Plettusa*; preocc. Loew, 1851). Type-locality: Chile, Valdivia.
testacea Philippi, 1865:598 (*Plettusa*). Type-locality: "Chile".
philippii Alexander, 1913b:487 (*Geranomomyia*; unnecessary nom. nov. for *virescens* Philippi). Type-locality: Chile, Valdivia.
- VARIEGATA** (Walker), 1837:333 (*Geranomomyia*). Type-locality: Brazil, São Paulo, São Paulo. Distr. — Brazil.
- VERSUTA** Alexander, 1930c:728. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Mexico, Colombia. Ref. — Alexander, 1967a:312, fig. 38 (male hypopygium).
- VILLARICENSIS** Alexander, 1929e:400. Type-locality: Paraguay, Guaira, Villarica. Distr. — Paraguay.
- VINDICTA VINDICTA** Alexander, 1943c:21. Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela. Refs. — Alexander, 1944c:43, fig. 5 (male hypopygium); 1954b:62, fig. 31 (male hypopygium).
ssp. **DILUCIDA** Alexander, 1954b:62, fig. 32 (male hypopygium). Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Brazil, Ecuador.



- VIRESCENS (Loew), 1851:398 (*Aporosa*). Type-locality: Lesser Antilles: Virgin Is., St. Thomas. Distr. — U.S.A. (Florida), Guyana, Puerto Rico, Virgin Is. Ref. — Alexander, 1932:365.
- VIRIDELLA Alexander, 1930b:111. Type-locality: British Honduras, Cayo District, Yalbac. Distr. — British Honduras.
- WALKERI Alexander, 1930c:730. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia, Venezuela, Brazil.
- XANTHOPLACA (Alexander), 1921d:43 (*Geranomyia*). Type-locality: Peru. Loreto, Iquitos. Distr. — Ecuador, Peru, Brazil. Ref. — Alexander, 1954b:63, fig. 30 (male hypopygium).
- YUNQUENSIS Alexander, 1956b:119. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico. Ref. — Alexander, 1967a:312, fig. 39 (male hypopygium).

Subgenus IDIOGLOCHINA Alexander

- Dicranomyia*, subg. IDIOGLOCHINA Alexander, 1921f:207. Type-species, *Rhipidia tusttala* Alexander (orig. des.) (Australasian).
- Limonia*, subg. *Idioglochina*; Alexander, 1929f:243. Ref. — Alexander, 1967b:314, figs. 40, 41, 47.
- AMBROSIANA Alexander, 1962a:38. Type-locality: Chile, Isla San Ambrosio. Distr. — Chile. Ref. — Alexander, 1967a:314, figs. 40 (wing), 47 (male hypopygium).
- LATIBASIS Alexander, 1967a:314, fig. 41 (wing). Type-locality: Chile, Tarapacá, Arica. Distr. — Chile.
- PORTERI (Alexander), 1919a:133 (*Dicranomyia*). Type-locality: Chile, Antofagasta. Distr. — Chile.

Subgenus NEOGLOCHINA Alexander

- Limonia*, subg. NEOGLOCHINA Alexander, 1967a:316. Type-species, *felix* Alexander, (orig. des.) (Neotropical).
- AURIGENA Alexander, 1944d:309 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Cunibunda. Distr. — Ecuador. Ref. — Alexander, 1954c:235, fig. 2 (male hypopygium).
- CAPNORA CAPNORA (Alexander), 1921d:47 (*Dicranomyia*). Type-locality: Peru, Loreto, Napo River. Distr. — Peru.
- ssp. METAE Alexander, 1938h:434 [*Limonia (Limonia)*]. Type-locality: Colombia, Meta, Restrepo. Distr. — Colombia.
- CURRANIANA Alexander, 1944h:378 [*Limonia (Limonia)*]. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama. Ref. — Alexander, 1945L:378, figs. 3 (wing), 4 (male hypopygium).
- ESAU Alexander, 1947f:85, figs. 5 (wing), 6 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela, Ecuador.
- FELIX Alexander, 1950b:180 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Tarma, Uteyuacu. Distr. — Peru.
- FUMOSA (Alexander), 1912b:364, fig. t (wing) (*Furcomyia*). Type-locality: Guyana, Amatuik. Distr. — Panama, Venezuela, Guyana.



- CROSSA Alexander, 1938g:422 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador. Ref. — Alexander, 1954c:236, fig. 4 (male hypopygium).
- IMPETURBATA Alexander, 1951a:99 [*Limonia (Limonia)*]. Type-locality: Guatemala, Chicacao. Distr. — Guatemala.
- INFUCATA Alexander, 1927d:303 (*Limonia*). Type-locality: Mexico, Chiapas, San Cristobal. Distr. — Mexico.
- INGENS Alexander, 1945e:232, fig. 2 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Serra dos Orgãos. Distr. — Brazil.
- INSULARIS (Williston), 1896a:287, fig. 58 (wing) (*Limnobia*). Type-locality: Lesser Antilles: St. Vincent. Distr. — Dominica, St. Vincent.
- LEUCOSCELIS Alexander, 1950b:183, figs. 11 (wing), 15 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Peru, Loreto, Cerro Azul, Upper Ucayali River. Distr. — Peru.
- LIMBINERVIS Alexander, 1941a:249 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- LUTZI (Alexander), 1912b:339 (*Furcomya*). Type-locality: Guyana. Tukeit. Distr. — Guyana.
- MESOTRICHA Alexander, 1944h:379 [*Limonia (Limonia)*]. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama. Ref. — Alexander, 1945L:378, figs. 5 (wing), 6 (male hypopygium).
- MONILIGERA Alexander, 1967a:318, figs. 42 (wing), 49 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Libertad, Tena. Distr. — Ecuador.
- MULTISIGNATA Alexander, 1936d:753 [*Limonia (Limonia)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- PERNOBILIS Alexander, 1941a:248 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Satipo. Distr. — Peru.
- PRAECLARA Alexander, 1928f:636 [*Limonia (Limonia)*]. Type-locality: Argentina, Misiones, Iguazu Falls. Distr. — Brazil, Argentina. Ref. Alexander, 1950b:190, fig. 19 (male hypopygium).
- SCIASMA Alexander, 1950b:190, figs. 13 (wing), 20 (male hypopygium). [*Limonia (Limonia)*]. Type-locality: Peru, Loreto, Cerro Azul, Contamana. Distr. — Peru.
- TEIALBOCINCTA Alexander, 1941g:347 (also 1946h:93, as 1938) [*Limonia (Limonia)*]. Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Panama.

Subgenus NEOLIMNOBIA Alexander


- Dicranomyia*, subg. NEOLIMNOBIA Alexander, 1928g:20. Type-species, *Limnobia diva* Schiner (orig. des.) (Neotropical).
- Limonia*, subg. *Neolimnobia*; Alexander, 1929f:244; 1950b:193.
- ANTHRACOPODA Alexander, 1938c:324. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil, Bolivia.
- ARCHANGELICA Alexander, 1942g:173, figs. 4 (wing), 11 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CORALLINA Alexander, 1938g:423. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador. Ref. — Alexander, 1954c:243, fig. 16 (male hypopygium).



- DIVA (Schiner), 1868:46 (*Limnobia*). Type-locality: "Brazil". Distr. — Mexico, Panama, Venezuela, Brazil, Bolivia, Cuba, Jamaica, Puerto Rico. Refs. — Alexander, 1932:357; 1943d:138; 1964a:13, figs. 18 (wing), 25 (male hypopygium).
- EXCELSIOR Alexander, 1944g:300. Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- HYPOCRITA Alexander, 1935a:320. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- IMMACULIPES Alexander, 1935a:319. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- MUSCOSA (Enderlein), 1912:75, fig. W1 (wing) (*Dicranomyia*). Type-locality: Ecuador, Santa Inez. Distr. — Ecuador.
- PUGLIS Alexander, 1943b:110. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Mayorga Playa. Distr. — Ecuador. Ref. — Alexander, 1954c:244, fig. 18 (male hypopygium).
- STYGIORNIS Alexander, 1962c:78, fig. 51 (wing). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- TRICINCTA TRICINCTA (Alexander), 1913d:405 (*Dicranomyia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru, Bolivia.
ssp. *TEXTRINA* Alexander, 1967a:319. Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.

Subgenus NEOLIMONIA Alexander

- Limonia*, subg. NEOLIMONIA Alexander, 1964a:28. Type-species, *Furcomyia eiseni* Alexander, (orig. des.) (Neotropical). Ref. — Alexander, 1967a:320, fig. 48.
- AMAZONICA (Alexander), 1920b:2 (*Dicranomyia*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- AUSTERA Alexander, 1950b:175, fig. 14 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- BIMUCRONATA Alexander, 1938g:419 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador, Bolivia. Ref. — Alexander, 1954c:235, fig. 3 (male hypopygium).
- BORINQUENSIS Alexander, 1950b:177 [*Limonia (Limonia)*]. Type-locality: Puerto Rico, Luquillo National Forest, El Yunque. Distr. — Puerto Rico.
- CARIBAEA Alexander, 1932:383, fig. 25 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Cuba, Trinidad Mts., Santa Clara, San Blas. Distr. — Cuba.
- CORDILLERENSIS (Alexander), 1913c:196, pl. 5, fig. 1 (wing) (*Dicranomyia*). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- CUZCOENSIS Alexander, 1967a:321, figs. 43 (wing), 48 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- DECEPTRIX Alexander, 1945j:14 [*Limonia (Limonia)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- DICAX Alexander, 1941f:313 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.



- DOMBALLAH Alexander, 1939d:108, fig. 3 (wing) [*Limonia (Limonia)*]. Type-locality: Hispaniola: Dominican Republic, Loma Rucilla. Distr. — Dominican Republic.
- EISENI (Alexander), 1912b:338, fig. s (wing) (*Furcomyia*). Type-locality: Guatemala, Aguna. Distr. — Guatemala, Panama, Guyana, Brazil. Ref. — Alexander, 1964a:28, figs. 17 (wing), 24 (male hypopygium).
- EURYLEON Alexander, 1950b:179, fig. 10 (wing) [*Limonia (Limonia)*]. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- HESIONE Alexander, 1950b:182 [*Limonia (Limonia)*]. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.
- HORRENDA Alexander, 1944b:154, fig. 1 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Peru, Ayacucho, La Mar, Tamba. Distr. — Peru.
- HUACAPISTANAE Alexander, 1941f:314 [*Limonia (Limonia)*]. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- HYPERPHALLUS Alexander, 1938g:418 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador. Ref. — Alexander, 1954c:236, fig. 5 (male hypopygium).
- IMMODICA Alexander, 1938a:25 [*Limonia (Limonia)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INDOMITA Alexander, 1938a:23 [*Limonia (Limonia)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- JAMAICENSIS Alexander, 1926b:223. Type-locality: Jamaica, Troy. Distr. — Jamaica. Refs. — Alexander, 1950b:189, fig. 17 (male hypopygium); 1964a:29, figs. 17 (wing), 24 (male hypopygium).
- LACHESIS Alexander, 1942d:139 [*Limonia (Limonia)*]. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador, Peru. Ref. — Alexander, 1954c:237, fig. 6 (male hypopygium).
- LAWLORI Alexander, 1934a:55 [*Limonia (Limonia)*]. Type-locality: Panama, Chiriquí, Lerida. Distr. — Panama, Ecuador. Ref. — Alexander, 1954c:237, fig. 7 (male hypopygium).
- LUDIBUNDA Alexander, 1927d:304. Type-locality: Mexico, Chiapas, Hueitepec. Distr. — Mexico, Panama.
- LUSTRALIS Alexander, 1938g:421 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador.
- MACINTYREI Alexander, 1938b:344 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Peru. Ref. — Alexander, 1954c:238, figs. 1 (wing), 8 (male hypopygium).
- ONOMA Alexander, 1950a:38 [*Limonia (Limonia)*]. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- OPTABILIS Alexander, 1921d:48 [*Limonia (Limonia)*]. Type-locality: Peru, Loreto, Napo River. Distr. — Peru, Ecuador. Ref. — Alexander, 1954c:238, fig. 9 (male hypopygium).
- PASTAZICOLA Alexander, 1938g:416 [*Limonia (Limonia)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador. Ref. — Alexander, 1954c:240, fig. 12 (male hypopygium).
- PORRECTA Alexander, 1950b:188, figs. 12 (wing), 18 (male hypopygium) [*Limonia (Limonia)*]. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.



- RORAIMAE Alexander, 1931b:5, fig. 2 (wing). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- SANCTAEMARTAE Alexander, 1930c:722 [*Limonia* (*Limonia*)]. Type-locality: Colombia, Magdalena, Mt. Santa Marta, La Cumbre. Distr. — Colombia, Peru.
- SCABRISTYLA Alexander, 1968e:212. Type-locality: Mexico, Tenancingo. Distr. — Mexico.
- SUBDOMITA Alexander, 1945e:234 [*Limonia* (*Limonia*)]. Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil.
- VELASTEGUII Alexander, 1938b:345, fig. 4 (wing) [*Limonia* (*Limonia*)]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1954e:241, fig. 15 (male hypopygium).

Subgenus PERIPHEROPTERA Schiner.

- PERIPHEROPTERA Schiner, 1866:933 (as genus) (emended to *Peripheroptera* by Schiner, 1868, and thus adopted by all subsequent authors). Type-species, *nitens* Schiner (orig. des.) (Neotropical).
- Limonia*, subg. *Peripheroptera*; Alexander, 1929f:242. Ref. — Alexander, 1967a:322, figs. 44, 45, 46, 50.
- ABERRANS (Schiner), 1868:47 (*Rhamphidia*). Type-locality: (original description said from Colombia; actually from Venezuela). Distr. — Venezuela. Ref. — Alexander, 1944L:188.
- ANGUSTIFASCIATA (Alexander), 1922e:35 (*Peripheroptera*). Type-locality: "Venezuela". Distr. — Venezuela.
- ARCUATA (Alexander), 1913d:411, fig. 8 (wing) (*Peripheroptera*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- ATROSIGNATA Alexander, 1934a:58. Type-locality: Panama, Chiriqui, Finca Parada. Distr. — Panama.
- AUSTROANDINA Alexander, 1929f:248 (nom. nov. for *subandina* Alexander). Type-locality: Argentina, Catamarca, El Suncho. Distr. — Argentina.
- subandina* Alexander, 1919a:135, fig. 2 (wing) (*Peripheroptera*; preoce. Alexander, 1913b). Type-locality: Argentina, Catamarca, El Suncho.
- COCHABAMBAE Alexander, 1945b:53. Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- CROCEIBASIS Alexander, 1942a:240. Type-locality: Peru, Junin, Tarma, Carpapata. Distr. — Peru.
- CYNARA Alexander, 1942f:788. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Ecuador, Peru.
- DIS Alexander, 1944g:301. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- EUDORAE (Alexander), 1913d:410, fig. 7 (wing) (*Peripheroptera*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- EURYPTERA Alexander, 1953d:150. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru. Ref. — Alexander, 1967a:323, fig. 44 (wing).
- FULVISTIGMA Alexander, 1953d:151. Type-locality: Peru, Tarma, Uteuyacu. Distr. — Peru.



- FUMBASALIS Alexander, 1937e:58. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1939c:432, fig. 3 (wing).
- GLOCHINOIDES (Alexander), 1922c:36 (*Peripheroptera*). Type-locality: Venezuela, from "Novara" Expedition. Distr. — Venezuela.
- INCOMMODA (Osten Sacken), 1887b:176 (*Peripheroptera*). Type-locality: "Brazil". Distr. — Brazil, Bolivia. Ref. — Alexander, 1913d:410, fig. 5 (wing).
- INCOMMOTES Alexander, 1928f:638. Type-locality: Peru, "Tambillo". Distr. — Peru.
- LANKESTERI Alexander, 1946f:303. Type-locality: Costa Rica, Cartago, Hacienda "Las Cóncavas". Distr. — Costa Rica.
- LICHYELLA Alexander, 1947f:90, fig. 9 (wing). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- LISSOMELANIA Alexander, 1967a:323, figs. 45 (wing), 50 (male hypopygium). Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- MACHUPICHUANA Alexander, 1953d:152. Type-locality: Peru, Cuzco, Machu Picchu. Distr. — Peru.
- MORGANA Alexander, 1944k:597. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- NEARCUATA Alexander, 1940d:215. Type-locality: Ecuador, Napo-Pastaza, Mapoto. Distr. — Ecuador, Peru.
- NITENS (Schiner), 1868:47 (*Peripheroptera*). Type-locality: Colombia (type male), Venezuela (type female). Distr. — Venezuela, Peru, Colombia.
- oliviae* Alexander, 1915b:441, fig. 1 (wing) (*Peripheroptera*). Type-locality: Panama, Porto Bello. Ref. — Alexander, 1944L:189.
- ORDINARIA Alexander, 1945L:380. Type-locality: Bolivia, Cochabamba, Chapore. Distr. — Bolivia.
- PERAMOENA Alexander, 1945a:20. Type-locality: Ecuador, Tungurahua, Mt. Tungurahua, Minza Chica. Distr. — Ecuador.
- PERDELECTA Alexander, 1940d:217. Type-locality: Ecuador, Bolívar, Balzapamba. Distr. — Ecuador.
- FRINDLEI Alexander, 1938g:424. Type-locality: Ecuador, Napo-Pastaza, Pundoa. Distr. — Ecuador.
- REDIVIVA Alexander, 1942a:238. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- RHODA Alexander, 1939c:430, fig. 2 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SCHINERI (Osten Sacken), 1887b:177 (*Peripheroptera*). Type-locality: "Brazil". Distr. — Brazil.
- SUBAMOENA Alexander, 1942a:239, fig. 2 (wing). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru, Bolivia.
- TEUCHOLABODES (Alexander), 1913d:410, fig. 6 (wing) (*Peripheroptera*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- THIOPTERA Alexander, 1941a:250. Type-locality: Ecuador, Tungurahua, Mt. Tungurahua, Minza Chica. Distr. — Ecuador. Ref. — Alexander, 1967a:324, fig. 46 (wing).



- TRIMELANIA Alexander, 1942a:242. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- TRINIGRINA Alexander, 1940d:214. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- VIVASBERTHIERI Alexander, 1940e:899, fig. 2 (wing). Type-locality: Venezuela, Mérida, Andes. Distr. — Venezuela.

Subgenus RHIPIDIA Meigen

- RHIPIDIA Meigen, 1818:153 (as genus). Type-species, *maculata* Meigen (mon.) = *duplicata* Doane (Western Palaearctic).
- Rhipidia*, subg. *Arhipidia* Alexander, 1912c:6. Type-species, *domestica* Osten Sacken (Alexander, 1950b:195).
- Rhipidia*, subg. *Monorhipidia* Alexander, 1912c:6. Type-species, *fidelis* Osten Sacken (Alexander, 1950b:195).
- Rhipidia*, subg. *Conorhipidia* Alexander, 1914c:117. Type-species, *conica* Alexander (orig. des.).
- Limonia*, subg. *Rhipidia*; Alexander, 1929f:241. Refs. — Alexander, 1950b: 195-221, figs. 21-42; 1967a:324-343, figs. 51-68.
- AGGLOMERATA (Alexander), 1926a:160 [*Rhipidia* (*Arhipidia*)]. Type-locality: Mexico, Veracruz, Córdoba, Río San Antonio. Distr. — Mexico.
- ANNULICORNIS (Enderlein), 1912:80, fig. Y1 (wing) (*Rhipidia*). Type-locality: Colombia, Hacienda Pehlke. Distr. — Colombia, Venezuela, Guyana, Ecuador, Peru, Brazil, Paraguay.
- ANTROTTRICHIA Alexander, 1967a:325, fig. 54 (male hypopygium). Type-locality: Ecuador, Ambato, Hacienda Pau Carlos. Distr. — Ecuador.
- APIRODITE Alexander, 1942f:771. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- ASPILOTA Alexander, 1967a:328, fig. 55 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Abitagua, Engaño. Distr. — Ecuador.
- EANOSENSIS Alexander, 1945p:757. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- BELLINGERI Alexander, 1964a:33, figs. 21 (wing), 22 (male hypopygium). Type-locality: Jamaica, St. Andrew, Morces Gap. Distr. — Jamaica.
- BIPECTINATA (Williston), 1896a:285, fig. 54 (wing) (*Rhipidia*). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.
- BREVIPETALIA Alexander, 1950a:40, fig. 7 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- BREVIRAMOSA BREVIRAMOSA Alexander, 1936d:754. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ssp. TOVARENSIS Alexander, 1944L:191. Type-locality: Venezuela, El Junquito, Colonia Tovar. Distr. — Venezuela.
- BRUCHIANA Alexander, 1929d:774. Type-locality: Argentina, La Plata. Distr. — Venezuela, Brazil, Argentina, Paraguay. Ref. — Bruch, 1939a: pl. 3, fig. 9 (body of male).
- CALVERTI (Alexander), 1912c:8, figs. d (wing), h (antenna) (*Rhipidia*). Type-locality: Costa Rica, Río Surubres. Distr. — Costa Rica.



- CASSANDRA Alexander, 1945c:261, fig. 12 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- CERMATOLEUCA Alexander, 1967a:329, figs. 51 (wing), 56 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- CHILOEANA Alexander, 1967a:331, figs. 52 (wing), 57 (male hypopygium). Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- COMMELINA Alexander, 1946b:235, figs. 17 (wing), 22 (male hypopygium). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- COMPLEXA Alexander, 1950b:200, fig. 31 (male hypopygium). Type-locality: Peru, Junin, Tarma, Utcuyacu. Distr. — Peru.
- CONICA CONICA (Alexander), 1914e:227, pl. 3, fig. 2 (wing) (*Rhipidia*). Type-locality: Guyana, Bartica. Distr. — Guyana.
- ssp. TURRIFERA Alexander, 1931b:8 [*Limonía (Rhipidia)*]. Type-locality: Venezuela, Mt. Duida. Distr. — Venezuela.
- COSTALOIDES (Alexander), 1920b:4 [*Rhipidia (Rhipidia)*]. Type-locality: Panama, Quebrada Richa, branch of Río Tapaliza, a branch of Río Puerro. Distr. — Panama.
- CRAMPTONI (Alexander), 1912c:10, figs. c (wing), i (antenna) (*Rhipidia*). Type-locality: "North Brazil". Distr. — Brazil.
- CYTHIEREA Alexander, 1942f:773. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- DISTELA Alexander, 1967a:332, figs. 53 (wing), 58 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Libertad, 6 km southeast of Tena. Distr. — Ecuador.
- DOMESTICA DOMESTICA (Osten Sacken), 1859:208 (*Rhipidia*). Type-locality: U.S.A., District of Columbia. Distr. — Nearctic; Guyana, Venezuela, Brazil, Bolivia, Cuba, Jamaica, Puerto Rico, Lesser Antilles: Dominica. Ref. — Alexander, 1932:360.
- ssp. AMAZONENSES (Alexander), 1912c:17, fig. b (wing) (*Rhipidia*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Peru, Brazil.
- ssp. ANGUSTIFRONS (Alexander), 1912c:16, fig. f (wing) (*Rhipidia*). Type-locality: Ecuador, Guayas, Guayaquil. Distr. — Ecuador, Peru.
- DOTALIS Alexander, 1942e:251. Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- ELIANA Alexander, 1950b:202, fig. 32 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- EREMNOPTERA Alexander, 1967a:334, figs. 59 (wing), 63 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru, Ecuador.
- EURYPHALLUS Alexander, 1967a:336, fig. 64 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- FLABELLIFORMIS FLABELLIFORMIS Alexander, 1934a:59. Type-locality: Panama, Chiriquí, Finca Lerida. Distr. — Panama, Venezuela, Peru, Bolivia.
- ssp. BRACHYNEMA Alexander, 1936a:15. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- GRACILIRAMA GRACILIRAMA Alexander, 1940b:150. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- ssp. LASSULA Alexander, 1940b:151. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.



- GUERRERENSIS Alexander, 1946b:237. Type-locality: Mexico, Guerrero, Chilapa. Distr. — Mexico.
- HIRTILOBATA Alexander, 1939a:75. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- ILLUMINATA Alexander, 1967a:337, figs. 60 (wing), 66 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- IMPROPERATA Alexander, 1936d:757. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INAEQUIPECTINATA Alexander, 1929d:772. Type-locality: Argentina, Misiones, Iguazu Falls. Distr. — Argentina. Ref. — Alexander, 1950b:204, figs. 35, 35A (male hypopygium).
- INGENUA Alexander, 1945b:52. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- INVARIPENNIS INVARIPENNIS Alexander, 1941f:335, fig. 8 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
ssp. CARPAPATAE Alexander, 1941f:337. Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.
- JUBILATA Alexander, 1938g:427. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- JUNINENSIS Alexander, 1942a:243. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- LAETITARSIS Alexander, 1937f:221. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LAIIS Alexander, 1950b:204, figs. 22 (antenna), 26 (wing), 33 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.
- LATILUTEA Alexander, 1942h:430, fig. 11 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- LEDA Alexander, 1945d:275. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- LICHNOPHORA Alexander, 1962c:79, figs. 52 (wing), 54 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- LONGISPINA (Alexander), 1922a:4 [*Rhipidia* (*Rhipidia*)]. Type-locality: Costa Rica, San José. Distr. — Mexico, Costa Rica.
- LONGURIO Alexander, 1938a:28. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LUQUILLOENSIS Alexander, 1950b:207, fig. 34 (male hypopygium). Type-locality: Puerto Rico, Luquillo National Forest, El Yunque. Distr. — Jamaica, Puerto Rico. Ref. — Alexander, 1964a:35, fig. 27 (male hypopygium).
- LUXURIOSA Alexander, 1929f:247 (nom. nov. for *vicina* Alexander). Type-locality: Colombia, Magdalena, La Cumbre. Distr. — Colombia.
vicina Alexander, 1916a:8, pl. 2, fig. 2 (wing) [*Rhipidia* (*Arhipidia*); procc. Macquart, 1838]. Type-locality: Colombia, Magdalena, La Cumbre.
- MEGALOPYGA Alexander, 1967a:338, figs. 61 (wing), 65 (male hypopygium). Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- MELANARIA Alexander, 1942f:774. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- MICROSTICTA (Alexander), 1921b:175 (*Rhipidia*). Type-locality: Argentina, Jujuy, Ciudad. Distr. — Argentina. Ref. — Bruch, 1939a, pl. 4, fig. 10 (wing); pl. 5, fig. 17 (entire female).

- MONNULA Alexander, 1962e:80, fig. 53 (wing). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- MONOXANTHA Alexander, 1944j:152, fig. 2 (wing). Type-locality: Venezuela, Mérida, Cano del Tigre. Distr. — Venezuela.
- MORDAX Alexander, 1950b:208, figs. 27 (wing), 36 (male hypopygium). Type-locality: Peru, Junín, Tarma, Utcuyacu. Distr. — Peru.
- MULTIFIDA (Alexander), 1926a:159, figs. 2 (wing), 13 (male hypopygium) [*Rhipidia* (*Rhipidia*)]. Type-locality: Mexico, Veraacruz, Córdoba. Distr. — Mexico.
- MULTIGUTTATA (Alexander), 1912e:14, fig. a (wing) (*Rhipidia*). Type-locality: Guatemala, Tetonicapán. Distr. — Guatemala, Panama.
- MULTIPUNCTIGERA Alexander, 1945o:590. Type-locality: Ecuador, Santiago-Zamora, Río Zamora. Zumbi. Distr. — Ecuador.
- MULTIRAMOSA Alexander, 1950b:210, figs. 24 (antenna), 39 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- MYRIOSTICTA Alexander, 1941f:333. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Ecuador, Peru.
- MYSTICA Alexander, 1931c:627. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- NEGLECTA NEGLECTA Alexander, 1936d:755. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ssp. SUBNEGLECTA Alexander, 1942g:175. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NIGROROSTRATA NIGROROSTRATA Alexander, 1938a:30. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ssp. MUZOENSIS Alexander, 1938h:434. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia.
- NOBILISSIMA Alexander, 1941f:332. Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- NUBILOSA Alexander, 1967a:340, fig. 67 (male hypopygium). Type-locality: Peru, Cuzeo, Quincemil. Distr. — Peru.
- OCELLANA Alexander, 1942h:432, fig. 12 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- PALLATANGAE Alexander, 1929d:775. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- PARAGUAYANA Alexander, 1929d:776. Type-locality: Paraguay, Guaira, Villarriea. Distr. — Paraguay.
- PAULUS Alexander, 1947e:62, figs. 2 (wing), 7 (male hypopygium). Type-locality: Venezuela, Táchira, San Camilo. Distr. — Venezuela.
- PERARMATA (Alexander), 1921d:52 [*Rhipidia* (*Rhipidia*)]. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- PERSIMPLEX Alexander, 1950b:212, figs. 21 (antenna), 28 (wing), 38 (male hypopygium). Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.
- PERSUFFUSA Alexander, 1967a:341, figs. 62 (wing), 68 (male hypopygium). Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- PIAON Alexander, 1950b:214, figs. 23 (antenna), 29 (wing), 37 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.



- POLYCLADA Alexander, 1943b:115. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PRATTI Alexander, 1950b:217, fig. 41 (male hypopygium). Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- PROCTIGERICA Alexander, 1950b:218, fig. 40 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- PROFANA Alexander, 1941f:338, fig. 9 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- PROLIFERATA Alexander, 1940b:151. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- PROSENI Alexander, 1954d:141. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- PUNCTORIA Alexander, 1929f:246 (nom. nov. for *punctipennis* Alexander). Type-locality: Costa Rica, Turrucare, Río Siquiare. Distr. — Costa Rica, Peru.
- punctipennis* Alexander, 1914c:117, figs. 1 (wing), 10 (male hypopygium) [*Rhipidia* (*Conorhipidia*); preocc. Skuse, 1890]. Type-locality: Costa Rica, Turrucare, Río Siquiare.
- SCHADEI Alexander, 1929d:777. Type-locality: Paraguay, Santa Barbara. Distr. — Brazil, Paraguay.
- SCHWARZI (Alexander), 1912c:13, fig. e (wing) [*Rhipidia* (*Arhipidia*)]. Type-locality: Hispaniola: Dominican Republic. Distr. — U.S.A. (Southern Florida), Cuba, Dominican Republic, Jamaica, Venezuela.
- SEJUGATA Alexander, 1939a:73. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- SHANNONI (Alexander), 1914g:581, fig. 23 (wing) [*Rhipidia* (*Arhipidia*)]. Type-locality: Panama, Canal Zone. Distr. — Panama.
- SIMPLICICORNIS Alexander, 1938a:27. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SPRUCEI Alexander, 1943b:117. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- STONEI Alexander, 1938d:328. Type-locality: Colombia, Usme. Distr. — Colombia.
- SUBCOSTALIS (Alexander), 1922a:3 [*Rhipidia* (*Rhipidia*)]. Type-locality: Costa Rica, Tiribi. Distr. — Costa Rica, Jamaica. Refs. — Alexander, 1928g:21; 1964a:35, figs. 23 (antenna), 28 (male hypopygium), 30 (wing).
- SUBPECTINATA SUBPECTINATA (Williston), 1896a:237, figs. 57 (wing), 57a (antenna) (*Rhipidia*). Type-locality: Lesser Antilles: St. Vincent, Distr. — St. Vincent, Grenada.
- ssp. PLEURALIS (Alexander), 1912c:12 (*Rhipidia*). Type-locality: Guatemala, Alta Verapaz, Trece Aguas, Cacao. Distr. — Mexico, Guatemala, Panama, Colombia.
- SUBTERMINALIS (Alexander), 1921b:175 (*Rhipidia*). Type-locality: Argentina, La Plata, Bosque. Distr. — Argentina, Brazil.
- SUBVAFRA Alexander, 1962c:82, fig. 55 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- SUCCENTIVA Alexander, 1941f:340, fig. 10 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.



- SUPERARMATA Alexander, 1942f:776. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- SURINAMICA Alexander, 1946f:301. Type-locality: Surinam, Brownsberg. Distr. — Surinam.
- SYBARITA Alexander, 1945n:400. Type-locality: Ecuador, Bolívar, Hacienda Talahua. Distr. — Ecuador.
- SYCOPHANTA Alexander, 1945n:402. Type-locality: Ecuador, Bolívar, Hacienda Talahua. Distr. — Ecuador.
- TABESCENS (Enderlein), 1912:81, fig. Z¹ (wing) (*Rhipidia*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- TETRALEUCA Alexander, 1937a:182. Type-locality: Puerto Rico, El Yunque. Distr. — Puerto Rico.
- THYSEE Alexander, 1942f:780. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- TIRESIAS Alexander, 1950b:220, figs. 30 (wing), 42 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- TRIDIGITATA Alexander, 1943h:488, fig. 2 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- TRIPLECTINATA Alexander, 1931c:625. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia. Ref. — Alexander, 1950b:221, fig. 25 (antenna).
- TURRITELLA Alexander, 1943c:205. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador, Peru.
- UNIPECTINATA (Williston), 1896a:286, fig. 55 (antenna) (*Rhipidia*). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent. Ref. — Alexander, 1940d:221, fig. 25 (male hypopygium).
- UXOR Alexander, 1942f:778. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru, Bolivia. Ref. — Alexander, 1962c:83, fig. 56 (male hypopygium).
- VAFRA Alexander, 1941f:342, fig. 11 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru, Ecuador.
- VARICOSTA Alexander, 1934a:60. Type-locality: Panama, Chiriquí, Finca Parada. Distr. — Panama.
- WILLISTONIANA Alexander, 1929f:246 (nom. nov. for *costalis* Williston). Type-locality: Lesser Antilles: St. Vincent. Distr. — Costa Rica, Dominica, St. Vincent. Ref. — Alexander, 1914c:118, fig. 2 (wing).
- costalis* Williston, 1896a:286, fig. 56 (antenna) (*Rhipidia*; preocc. Wiedemann, 1824). Type-locality: "St. Vincent".

Subgenus ZALUSA Enderlein

- ZALUSA Enderlein, 1906:71 (as genus). Type-species, *falklandica* Enderlein (Neotropical).
- Limonía*, subg. *Zalusá*; Alexander, 1929a:85; 1967a:343.
- FALKLANDICA (Enderlein), 1906:71, figs. 1 (halter, wing), 2 (male hypopygium) (*Zalusá*). Type-locality: Falkland Is., Port Darwin. Distr. — Falkland Is.; Patagonia, Missier Channel (Alexander, 1929a:85).



Subgenus ZELANDOGLOCHINA Alexander

- Dicranomyia*, subg. ZELANDOGLOCHINA Alexander, 1924f:449. Type-species, *huttoni* Edwards (orig. des.) (Australasian: New Zealand).
- Limonia* (*Zelandoglochina*) Alexander, 1929a:71. Ref. — Alexander 1967a:344.
- ANCELICA Alexander, 1929a:74, fig. 12 (wing). Type-locality: Argentina, Rio Negro, L. Nahuel Huapi. Distr. — Chile, Argentina. Ref. — Alexander, 1953g:10, fig. 4 (male hypopygium).
- APIHANTA Alexander, 1929a:83, fig. 175 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina.
- BIGOTI (Alexander), 1913b:488 (*Furcomyia*; nom. nov. for *longicollis* Bigot). Type-locality: Chile, Tierra del Fuego, Cape Horn. Distr. — Chile, Argentina. Ref. — Alexander, 1929a:72.
- longicollis* Bigot, 1888:8 (preocc. Maequart, 1846). Type-locality: Chile, Tierra del Fuego, Cape Horn.
- FAGETORUM Alexander, 1929a:79, fig. 171 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Argentina.
- FLABELLIFERA Alexander, 1929a:77, figs. 15 (wing), 168 (antenna), 169 (male hypopygium). Type-locality: Chile, Chiloé Is., Aneud. Distr. — Chile.
- FULVITHORAX (Philippi), 1865:598 (*Plettusa*). Type-locality: Chile, Valdivia. Distr. — Chile. Identity uncertain. Ref. — Alexander, 1929a:72.
- MINIATA Alexander, 1929a:84, fig. 176 (male hypopygium). Type-locality: Chile, Chiloé Is., Aneud. Distr. — Chile, Argentina.
- MULTIARMATA Alexander, 1929a:76, figs. 14 (wing), 167 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Chile, Argentina.
- MULTINODOSA Alexander, 1929a:81, fig. 174 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina.
- NODULIFERA Alexander, 1929a:80, figs. 172 (head), 173 (male hypopygium). Type-locality: Chile, Llanquihue, Puerto Montt. Distr. — Chile.
- OFELLA Alexander, 1953g:10, figs. 1 (wing), 5 (male hypopygium). Type-locality: Chile, Chiloé Is., Aulen, Duhatao. Distr. — Chile.
- OMISSISTYLA Alexander, 1929a:78, figs. 40 (wing), 170 (male hypopygium). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile.
- PARVISPINOSA Alexander, 1928h:359. Type-locality: Chile, Concepción. Distr. — Chile. Refs. — Alexander, 1929a:73; 1953g:11, fig. 6 (male hypopygium).
- PERVINETA PERVINCTA Alexander, 1936d:757. Type-locality: Argentina, Rio Negro, L. Correntoso, L. Nahuel Huapi. Distr. — Argentina.
- ssp. PERCELESTIS Alexander, 1953b:10. Type-locality: Chile, Malleco, Curacautin. Distr. — Chile, Argentina.
- PILOSIPENNIS Alexander, 1929a:75, figs. 13 (wing), 166 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- SETULIPENNIS Alexander, 1928h:356. Type-locality: Chile, Concepción. Distr. — Chile. Refs. — Alexander, 1929a:72; 1953g:12, fig. 7 (male hypopygium).



- FEHUELICHE Alexander, 1929a:82. Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina.
- TENUIPALPIS Alexander, 1928h:357. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile. Refs. — Alexander, 1929a:72; 1953g:12, fig. 8 (male hypopygium).
- TORTICORNIS Alexander, 1928h:356. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile. Refs. — Alexander, 1929a:73; 1953g:12, fig. 9 (male hypopygium).

Unplaced *Limonia*


- MULTIPUNCTATA (Fabricius), 1805:31 (*Tipula*). Type-locality: "South America". Identity uncertain, presumed to belong to *Rhipidia*.
- OCCELLATA (Röder), 1886:259 (*Limnobia*). Type-locality: Colombia, Cerro del Altar.
- VALLEIDA (Macquart), 1838a:76(72) (*Limnobia*). Type-locality: "Chile". Ref. — Blanchard, 1852:340. Unrecognized species.

Genus ORIMARGA Osten Sacken

- ORIMARGA Osten Sacken, 1869:120. Type-species, *Limnobia alpina* Zetterstedt (Coquillett, 1910:580) (Western Palearctic).

Subgenus DIOTREPHA Osten Sacken

- DIOTREPHA Osten Sacken, 1878:219 (as genus). Type-species, *mirabilis* Osten Sacken (mon.) (Nearctic).
- Thambeta* Williston, 1896b:32. Type-species, *Diotrepha mirabilis* Osten Sacken (Alexander, 1965d:53).
- ACROLEUCA Alexander, 1964a:22, figs. 9 (wing), 14 (male hypopygium). Type-locality: Jamaica, St. Thomas, Bath. Distr. — Jamaica.
- ARAWAK Alexander, 1964a:23, figs. 10 (wing), 15 (male hypopygium). Type-locality: Jamaica, St. Thomas, Corn Puss Gap. Distr. — Jamaica.
- ATREBASIS (Alexander), 1914e:240, pl. 4, fig. 4 (wing) (*Diotrepha*). Type-locality: Guyana, Bartica. Distr. — Guyana, Surinam, Peru, Brazil.
- CONCINNA (Williston), 1896a:291, fig. 66 (wing) (*Diotrepha*). Type-locality: Lesser Antilles, St. Vincent. Distr. — St. Vincent.
- ELONGATA Alexander, 1943c:23 (as *funicosta* ssp.). Type-locality: Venezuela, Carabobo, Borburata. Distr. — Venezuela, Bolivia.
- FLAVICOSTA (Alexander), 1928g:23 (*Diotrepha*). Type-locality: Jamaica, Cinchona, Logwood. Distr. — Cuba, Jamaica. Ref. — Alexander, 1964a:24, figs. 11 (wing), 16 (male hypopygium).
- FUMICOSTA (Alexander), 1921d:58 (*Diotrepha*). Type-locality: Brazil, Pará, Prata. Distr. — Venezuela, Ecuador, Brazil, Lesser Antilles: Grenada.
- MIRABILIS (Osten Sacken), 1878:220 (*Diotrepha*). Type-locality: U.S.A., Georgia. Distr. — Southern U.S.A., Cuba. Refs. — Alexander, 1932:387, fig. 9 (wing); Rogers, 1927a:23-26 (biology).
- MYERSIANA (Alexander), 1930c:732 (*Diotrepha*). Type-locality: Guyana, Koriabo, Barima River. Distr. — Guyana.
- OMISSINERVIS (Alexander), 1913c:447, pl. 14, fig. 9 (wing) (*Diotrepha*). Type-locality: Bolivia, Cuticcho, Yungas de Songo Valley. Distr. — Bolivia.



- PROFUSA Alexander, 1945o:593. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- SUBCONCINNA Alexander, 1940b:154. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- SUBPROTUSA Alexander, 1968a:22, fig. 4 (male hypopygium). Type-locality: Peru, Chanchamayo. Distr. — Peru.
- SYNDACTYLA Alexander, 1942d:140. Type-locality: Ecuador, Pichincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- TRAVASSOSI Alexander, 1943f:299, figs. 8 (wing), 13 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.

Subgenus ORIMARGA Osten Sacken

- ANDINA Alexander, 1916a:14, pl. 1, fig. 7 (wing). Type-locality: Colombia, Magdalena, Mt. Santa Marta, La Cumbre. Distr. — Colombia.
- ARGENTEOPLEURA Alexander, 1913a:48, fig. f (wing). Type-locality: Guatemala, Alta Verapaz, Trece Aguas, Cacao. Distr. — Mexico, Guatemala.
- BAHIANA Alexander, 1930c:732. Type-locality: Brazil, Bahia. Distr. — Brazil.
- CHIONOMERA Alexander, 1945c:263, fig. 9 (wing). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- CHIONOPUS Alexander, 1943b:119. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CUBENSIS Alexander, 1932:385. Type-locality: Cuba, Pinar del Río, Sierra Rangel. Distr. — Cuba.
- DAMPFI Alexander, 1926a:162, fig. 4 (wing). Type-locality: Mexico, Veracruz. Distr. — Mexico.
- DICHOPTERA Alexander, 1950a:44, fig. 11 (wing). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- EXCESSIVA EXCESSIVA Alexander, 1926d:380. Type-locality: Venezuela, Táchira, Táchira. Distr. — Venezuela. Ref. — Alexander, 1962e: 118, fig. 97 (male hypopygium).
- ssp. CHAPARENSIS Alexander, 1962c:117, fig. 98 (male hypopygium). Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- FARRIANA Alexander, 1964a:18, figs. 7 (wing), 12 (male hypopygium). Type-locality: Jamaica, St. Andrew. Distr. — Jamaica.
- FUNERULA Alexander, 1929b:93. Type-locality: Peru, Tambo Eneñas to Dos de Mayo, Campo del Pichis. Distr. — Peru.
- LANEI Alexander, 1942h:433, figs. 13 (wing), 17 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil.
- MAJUSCULA Alexander, 1940b:154. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico.
- MELAMPODIA Alexander, 1940e:112. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- MULTIPUNCTATA Alexander, 1938g:429. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- NEOGAUDENS Alexander, 1945c:264, fig. 10 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.



- NIGROAPICALIS Alexander, 1941c:328. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Costa Rica, Panama.
- NIVEITARSIS Alexander, 1915b:443, fig. 5 (wing). Type-locality: Panama, Porto Bello. Distr. — Panama.
- PALLIDIBASIS Alexander, 1921d:56. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru. Ref. — Alexander, 1967a:350, as *palliditarsus*, lapsus.
- PARVIPUNCTA Alexander, 1938g:430. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- PERPALLENIS Alexander, 1964a:21, figs. 8 (wing), 13 (male hypopygium). Type-locality: Jamaica, Portland, Spring Hill Post Office. Distr. — Jamaica.
- PUNCTIPENNIS Alexander, 1914e:239, pl. 4, fig. 3 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana.
- SATURNINA Alexander, 1943b:118. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- SCABRISETA Alexander, 1938g:431. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- SPECIOSA Alexander, 1953d:155. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru. Ref. — Alexander, 1967a:351, figs. 72 (wing), 78 (male hypopygium).
- SUBCOSTATA Alexander, 1954d:144. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SUBSPECIOSA Alexander, 1962c:118, fig. 96 (wing). Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- SUBTARTARUS Alexander, 1946e:531. Type-locality: Mexico, Jalisco, Tequila. Distr. — Mexico.
- TARTARUS Alexander, 1946e:530. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- TINGUANA Alexander, 1943f:297, fig. 7 (wing). Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil.
- TRISPINGERA Alexander, 1945a:24. Type-locality: Ecuador, Santiago-Zamora. Distr. — Ecuador.

Genus PROTOHELIUS Alexander

- PROTOHELIUS Alexander, 1928j:466. Type-species, *issikii* Alexander (orig. des.) (Oriental).
- CISATLANTICUS Alexander, 1938b:347, fig. 5 (wing). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- VENEZOLANUS Alexander, 1950a:45, fig. 13 (wing). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

Genus TONNOIROMYIA Alexander

- TONNOIROMYIA Alexander, 1926e:190. Type-species, *tasmaniensis* Alexander (orig. des.) (Australasian).
- PATAGONICA Alexander, 1929a:68, figs. 38 (wing), 165 (male hypopygium). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Chile, Argentina.



Tribe PEDICIINI

Genus DICRANOTA Zetterstedt

DICRANOTA Zetterstedt, 1838:851. Type-species, *guerini* Zetterstedt (mon.).

Subgenus PLECTROMYIA Osten Sacken

PLECTROMYIA Osten Sacken, 1869:282 (as genus). Type-species, *modesta* Osten Sacken (mon.) (Nearctic).

Astrolabis Osten Sacken, 1865:225. *Nomen nudum*.

TERGATA (Alexander), 1926a:172, figs. 11 (wing), 19 (male hypopygium) [*Rhaphidolabis* (*Plectromyia*)]. Type-locality: Mexico, Ajusco. Distr. — Mexico.

Subgenus RHAPHIDOLABIS Osten Sacken

RHAPHIDOLABIS Osten Sacken, 1869:284 (as genus). Type-species, *tenuipes* Osten Sacken (orig. des.) (Nearctic).

MEXICANA Alexander, 1946a:121. Type-locality: Mexico, Mexico State, Dos Ríos. Distr. — Mexico.

ROSTRIFERA Alexander, 1946a:122. Type-locality: Mexico, Mexico State, Río Borja, D.F. Distr. — Mexico.

Genus PEDICIA Latreille

PEDICIA Latreille, 1809:255. Type-species, *Tipula rivosa* Linnaeus (mon.). (Palearctic).

Peditia, error.

Subgenus TRICYPHONA Zetterstedt

TRICYPHONA Zetterstedt, 1837:65 (as genus). Type-species, *Limonia immaculata* Meigen (mon.).

Amalopis Haliday, 1856:xv. Type-species, *Limnobia occulta* Maigen (orig. des.).

ALBICENTRA Alexander, 1962b:352, fig. 3 (wing). Type-locality: Argentina, Neuquén, Lago Nonthue. Distr. — Argentina.

AYSENENSIS Alexander, 1944i:127, figs. 3 (wing), 4 (male hypopygium). Type-locality: Chile, Aysén. Distr. — Chile.

CHILOTA (Alexander), 1929a:110, figs. 46 (wing), 193-196 (male hypopygium) (*Tricyphona*). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile, Argentina.

CRASSIPYGA (Alexander), 1928b:48 (*Tricyphona*). Type-locality: Chile, Valparaíso, Perales de Marga-Marga. Distr. — Chile. Refs. — Alexander, 1929a:110; 1944i:130, fig. 6 (male hypopygium).

GUTTISTIGMA Alexander, 1941a:253. Type-locality: Chile, Angol. Distr. — Chile.

PENAI Alexander, 1953b:11, fig. 6 (male hypopygium). Type-locality: Chile, Malleco, Río Blanco. Distr. — Chile.

PHAEOSTIGMA Alexander, 1962c:353, fig. 4 (wing). Type-locality: Argentina, Neuquén, San Martín de los Andes. Distr. — Argentina.



FLATYPTERA (Alexander), 1929a:109, figs. 45 (wing), 191 (antenna), 192 (male hypopygium) (*Tricyphona*). Type-locality: Chile, Chiloé Is., Aneud. Distr. — Chile.

SERRIMARGA Alexander, 1944i:129, fig. 5 (male hypopygium). Type-locality: Argentina, Rio Negro, Lago Gutiérrez. Distr. — Argentina.

Unplaced Pedicelini

FUSILLA (Bigot), 1888:12 (*Tricyphona*). Type-locality: Argentina, Tierra del Fuego. Distr. — Argentina. Evidently not a *Tricyphona* and may be found to be a species of *Paracladura* Brunetti in the family Trichoeridae.

Tribe HEXATOMINI

Genus ATARBA Osten Sacken

ATARBA Osten Sacken, 1869:127. Type-species, *piticornis* Osten Sacken (mon.) (Nearectic). Refs. — Alexander, 1948c:536, figs. 21-26; Rogers, 1927b:49 (biology).

Subgenus ATARBA Osten Sacken

ALMEDAI Alexander, 1946e:2, fig. 2 (wing). Type-locality: Brazil, São Paulo. Boracéia. Distr. — Brazil.

AMABILIS Alexander, 1928a:113. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.

ANGUSTIPENNIS Alexander, 1928b:57. Type-locality: Cuba, Pinar del Río, Sierra Rangel. Distr. — Mexico, Cuba.

ANTHRAICINA Alexander, 1937d:609. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

APERTA APERTA Alexander, 1926a:170, fig. 9 (wing). Type-locality: Mexico, Córdoba, Pena Blanca. Distr. — Mexico, Brazil.

ssp. SUBAPERTA Alexander, 1945L:383, figs. 14 (wing), 15 (male hypopygium). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.

APICISPINOSA Alexander, 1934a:66. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.

BIFILOSA Alexander, 1948c:546, figs. 26 (wing), 32 (male hypopygium). Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica.

BIFUREULA Alexander, 1931a:115. Type-locality: Brazil, Santa Catarina, Jaguará. Distr. — Brazil.

BIPENDULA Alexander, 1966a:224. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.

BIPRODUETA Alexander, 1966a:224. Type-locality: Ecuador, Napo-Pastaza, Libertad. Distr. — Ecuador.


BOLIVIANA Alexander, 1930c:736. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.

BREVICORNIS Alexander, 1929d:781. Type-locality: Paraguay, Guaira, Villarica. Distr. — Paraguay.

BREVISSIMA Alexander, 1944f:30. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.



- BRUNNEICORNIS Alexander, 1916a:15, pl. 1, fig. 6 (wing); pl. 4, fig. 2 (male hypopygium). Type-locality: Colombia, Magdalena, La Cumbre. Distr. — Colombia.
- BULBIFERA Alexander, 1943i:744, fig. 4 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- CINCTICORNIS CINCTICORNIS Alexander, 1916d:499. Type-locality: Guyana, Mallali, Demerara River. Distr. — Guyana, Ecuador.
- ssp. THAMIA Alexander, 1950a:50, fig. 27 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- CIRCE Alexander, 1945n:414. Type-locality: Ecuador, Santiago-Zamora, Zamora. Distr. — Ecuador.
- COLUMBIANA Alexander, 1913e:199, pl. 5, fig. 4 (wing). Type-locality: Colombia, Cauca Almaguer. Distr. — Colombia.
- EUCELLATA Alexander, 1945n:413. Type-locality: Ecuador, Santiago-Zamora, Zamora. Distr. — Ecuador.
- DASYCERA Alexander, 1948e:548, figs. 24 (wing), 33 (male hypopygium). Type-locality: Peru, Loreto, Previsto, Upper Ucayali River. Distr. — Peru.
- DIACANTHA Alexander, 1945a:26. Type-locality: Ecuador, Napo-Pastaza Abitagua. Distr. — Ecuador.
- DILATISTYLA Alexander, 1966a:225. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- DINEMATOPHORA Alexander, 1943i:740, fig. 2 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- FIEBRIGI Alexander, 1922e:51. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- FORTICORNIS Alexander, 1947f:99, fig. 13 (wing). Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- FUSCOPICALIS Alexander, 1944f:29. Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia. Distr. — Brazil.
- HETERACANTHA Alexandre, 1943a:219. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- HIRTICORNIS Alexander, 1943a:221. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- IDONEA Alexander, 1939b:206. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- INCISURATA Alexander, 1938f:250. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LADDEYANA Alexander, 1944k:600. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- LATEROSPINA Alexander, 1962e:131, figs. 106 (wing), 108 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Loetol. Distr. — Bolivia.
- LONGITERGATA Alexander, 1945j:25, fig. 10 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- LYRIFORMIS Alexander, 1966a:226. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- MACRACANTHA Alexander, 1943a:217. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru, Bolivia.



- MEGAPHALLUS Alexander, 1921d:59. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- MELANOMERA Alexander, 1943g:540, fig. 6 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- MERITA Alexander, 1938f:250. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- MEXICANA Alexander, 1926a:170, fig. 8 (wing). Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico.
- MICROPHALLUS Alexander, 1944k:602. Type-locality: Ecuador, Los Ríos, Playas de Montalvo. Distr. — Ecuador.
- MULTIARMATA MULTIARMATA Alexander, 1943a:224. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- ssp. TARMAE Alexander, 1943a:225. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- NODULOSA Alexander, 1939b:207. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PALLIDAPEX Alexander, 1943a:222. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PERINCISA Alexander, 1948c:549, fig. 34 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- PROCERICORNIS Alexander, 1944d:316. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Cunibunda. Distr. — Ecuador.
- PUNCTISCUTA Alexander, 1922c:52. Type-locality: Paraguay, San Bernardino. Distr. — Brazil, Paraguay.
- QUASIMODO Alexander, 1950a:50, figs. 20 (wing), 24 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- RELIGIOSA Alexander, 1946a:126. Type-locality: Mexico, Mexico, D.F., Desierto de los Leones. Distr. — Mexico.
- RESTRICTA Alexander, 1943g:539, fig. 3 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- SCABROSA Alexander, 1944b:175, fig. 9 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- SCUTATA Alexander, 1939a:80. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- SETLOBATA Alexander, 1964a:66, figs. 65 (wing), 68 (male hypopygium). Type-locality: Jamaica, Portland, Hardwar Gap. Distr. — Jamaica.
- STIGMOSA Alexander, 1930c:737. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.
- SUBDENTATA Alexander, 1952a:121. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- TATEI Alexander, 1929d:782. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- TETRACANTHA Alexander, 1945a:28. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- TUBERCULIFERA TUBERCULIFERA Alexander, 1943i:742, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- ssp. EDAX Alexander, 1948c:551, fig. 35 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.



- TUNGURAHUENSIS Alexander, 1945p:762. Type-locality: Ecuador, Tungurahua, above Baños. Distr. — Ecuador.
- UNILATERALIS Alexander, 1931b:14, figs. 9 (wing), 18 (male hypopygium). Type-locality: Venezuela, Mt. Duida. Distr. — Venezuela.
- VARICORNIS Alexander, 1913e:448, pl. 14, fig. 10 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Guyana, Peru.
- VARIISPINA Alexander, 1938f:252. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Subgenus ISCHNOTHRIX Bigot

- ?*Lachnocera* Philippi, 1865:615. Type-species, *delicatula* Philippi (mon.). [In the interests of stability, the Author rejects this prior apparent generic name].
- ISCHNOTHRIX Bigot, 1888:7 (as genus). Type-species, *aetherea* Bigot (mon.) (Neotropical).
- Oromyia* Alexander, 1913c:203. Type-species, *lloydi* Alexander (orig. des.).
- Orolimnophila* Alexander, 1921b:178 (nom. nov. for *Oromyia* Alexander, on the wrong assumption that it was preocc. by *Orosmyia* Rondani, 1856). Type-species, *Oromyia lloydi* Alexander (aut.). Rcf. — Alexander, 1929a:167.
- AETHEREA (Bigot), 1888:7, pl. 2, fig. 1 (wing) (*Ischnothrix*). Type-locality: Chile, Tierra del Fuego. Distr. — Chile.
- APRICA Alexander, 1962c:129, figs. 105 (wing), 107 (male hypopygium) Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- ARGENTINICOLA (Alexander), 1921b:178 (*Orolimnophila*). Type-locality: Argentina, Tucumán, Quebrada Famailla. Distr. — Argentina.
- BERTHAE Alexander, 1948c:541, figs. 22 (wing), 31 (male hypopygium) Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- BREVIYRA Alexander, 1966a:223. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- BREVISECTOR (Alexander), 1935a:326 (*Ischnothrix*). Type-locality: Venezuela, Silla da Caracas. Distr. — Venezuela.
- CAMPTELLA Alexander, 1945p:759. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- DELICATULA (Philippi), 1865:615 (*Lachnocera*). Identity uncertain. Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1913b:502.
- DIGITIFERA Alexander, 1945p:761. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- FIDELIS (Alexander), 1929a:169, fig. 87 (wing) [*Ischnothrix* (*Orolimnophila*)]. Type-locality: Chile, Llanquihue, Puerto Montt. Distr. — Chile.
- GEMINATA Alexander, 1943g:535, figs. 1 (wing), 4 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- HELENAE Alexander, 1948c:543, fig. 29 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina.
- IGNITHORAX Alexander, 1929a:172, fig. 95 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina.
- INTEGRIOBA Alexander, 1943i:738, fig. 1 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- LLOYDI (Alexander), 1913c:204, pl. 5, fig. 7 (wing); pl. 6, figs. 7-9 (male hypopygium) (*Oromyia*). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.



- MESOCERA Alexander, 1929a:171, fig. 94 (wing). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile, Argentina.
- OBTUSILOBA Alexander, 1944b:172, fig. 8 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- PATENS (Alexander), 1940c:113 [*Ischnothrix (Ischnothrix)*]. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- PICTURATA (Alexander), 1929a:169, fig. 88 (wing) [*Ischnothrix (Orolimnophila)*]. Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Chile.
- RECTANGULARIS Alexander, 1954d:145. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SCUTELLATA (Alexander), 1929a:173 [*Ischnothrix (Ischnothrix)*]. Type-locality: Argentina, Río Negro, Lake Correntoso. Distr. — Chile, Argentina.
- SETICORNIS Alexander, 1946c:1, fig. 1 (wing). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- SPINITUBER Alexander, 1950a:49, figs. 22 (wing), 26 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- SUPPLICATA Alexander, 1943g:537, figs. 2 (wing), 5 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- TENUISSIMA (Alexander), 1929a:171 [*Ischnothrix (Orolimnophila)*]. Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina.
- VORACIS Alexander, 1948c:544, figs. 21 (wing), 30 (male hypopygium). Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.

Genus AUSTROLIMNOPHILA Alexander


Limnophila, subg. AUSTROLIMNOPHILA Alexander, 1920e:4. Type-species, *eutaeniata* Bigot (orig. des.) (Neotropical).

Polymoria Philippi of Alexander, 1924c:367.

Refs. — Alexander, 1929a:115; 1948a:153, figs. 3-4 (venation), 7-17 (male hypopygium) (discussion).

Subgenus AUSTROLIMNOPHILA Alexander

- ACUTERGATA Alexander, 1939c:434, figs. 4 (wing), 11 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- BIFIDARIA Alexander, 1942i:56, figs. 23 (wing), 30 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- BRADLEYI Alexander, 1929b:93. Type-locality: Argentina, Tucumán, Parque Aconquija. Distr. — Argentina. Ref. — Alexander, 1948a:161, fig. 10 (male hypopygium).
- BULBIFERA Alexander, 1948b:137. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil. Ref. — Alexander, 1966c:465, fig. 6 (male hypopygium).
- BYERSIANA Alexander, 1968c:213. Type-locality: Mexico, near Huauchinango. Distr. — Mexico.



- CANDIDITARSIS Alexander, 1937d:603. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CAPARAËNSIS Alexander, 1944f:27. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- CHILOEANA Alexander, 1953b:15, figs. 9 (wing), 13 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- COMANTIS Alexander, 1948a:161. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- DUSENI (Alexander), 1920e:5 (*Limnophila*). Type-locality: Chile, Aysén, Rio Aysén. Distr. — Chile. Ref. — Alexander, 1929a:124, fig. 50 (wing).
- ELNORA Alexander, 1929a:127, figs. 52 (wing), 202 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina.
- EUTAENIATA (Bigot), 1888:9, pl. 2, fig. 3 (wing) (*Limnophila*). Type-locality: Chile, Tierra del Fuego. Distr. — Chile, Argentina. Refs. — Alexander, 1920e:5; 1929a:115; 1948a:163, fig. 11 (male hypopygium).
- FUSCOHALTERATA Alexander, 1929a:123, figs. 64, 65 (wing). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- HAZELAE Alexander, 1929a:125, fig. 51 (wing). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile, Argentina. Ref. — Alexander, 1948a:163.
- INFIDELIS Alexander, 1929a:122, fig. 63 (wing). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile, Argentina.
- IRIS Alexander, 1929a:117, fig. 61 (wing). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile, Argentina.
- JOANA Alexander, 1929a:126, figs. 66 (wing), 201 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile. Ref. — Alexander, 1953b:15, figs. 8 (wing), 12 (male hypopygium).
- MACROPYGA Alexander, 1953b:12, figs. 7 (wing), 11 (male hypopygium). Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- MANNHEIMSI Alexander, 1960:15. Type-locality: Peru, near Taulis, Hacienda Udima. Distr. — Peru. Ref. — Alexander, 1966c:466, figs. 1 (wing), 7 (male hypopygium).
- MARTINEZI Alexander, 1958:197, figs. 1 (wing), 6 (male hypopygium). Type-locality: Bolivia. Cochabamba, Chapare, Loetol, El Limbo. Distr. — Bolivia.
- MERKLEI Alexander, 1928b:50. Type-locality: Argentina, Rio Negro, L. Nahuel Huapí, Bariloche. Distr. — Argentina, Chile. Ref. — Alexander, 1948a:164, fig. 13 (male hypopygium).
- MICHAELSENI Alexander, 1928h:361. Type-locality: Argentina, Tierra del Fuego, Ushuaia "Wald". Distr. — Argentina.
- MICROPILOTA Alexander, 1943f:300, fig. 9 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- MICROSTICTA Alexander, 1929a:119, fig. 49 (wing). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile, Argentina. Ref. — Alexander, 1948a:164, fig. 14 (male hypopygium).
- NAHUELICOLA Alexander, 1958:198, figs. 3 (wing), 7 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Pichinahuel. Distr. — Chile.



- NEUQUENENSIS Alexander, 1952e:90, figs. 5 (wing), 7 (male hypopygium).
Type-locality: Argentina, Neuquém, San Martín de los Andes.
Distr. — Chile, Argentina.
- NYMPHA Alexander, 1943e:206. Type-locality: Ecuador, Santiago-Zamora,
Río Zamora, Zumbi. Distr. — Ecuador. Ref. — Alexander,
1948a:166, fig. 15 (male hypopygium).
- OROENSIS Alexander, 1943h:490, fig. 3 (wing). Type-locality: Ecuador, El
Oro, Morro Morro. Distr. — Ecuador.
- PACIFERA Alexander, 1936e:60. Type-locality: Brazil, Santa Catarina, Nova
Teutônia. Distr. — Brazil.
- PALLIDISTYLA PALLIDISTYLA Alexander, 1942i:57, figs. 24 (wing), 32 (male
hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Serra
da Cantareira. Distr. — Brazil.
- ssp. PERLIMBATA Alexander, 1942i:59. Type-locality: Brazil, São Paulo,
São Paulo, Serra da Cantareira. Distr. — Brazil.
- PERCARA Alexander, 1958:200, figs. 4 (wing), 9 (male hypopygium). Type-
locality: Chile, Osorno, Antillanca. Distr. — Chile.
- PERSESSILIS Alexander, 1939b:200. Type-locality: Ecuador, Napo-Pastaza,
Abitagua. Distr. — Ecuador. Ref. — Alexander, 1948a:166, fig.
16 (male hypopygium).
- PLATENSIS (Alexander), 1923b:184 (*Limnophila*). Type-locality: Argentina,
La Plata, Palo Blanco. Distr. — Brazil, Argentina. Ref. —
Alexander, 1948a:166, fig. 17 (male hypopygium).
- POLYSPILOTA Alexander, 1937b:239. Type-locality: Brazil, Rio de Janeiro,
Rezende, Campo Belo. Distr. — Brazil.
- PUNCTIPENNIS (Philippi), 1865:609 (*Polymoria*). Type-locality: "Chile".
Distr. — Chile. Ref. — Alexander, 1929a:115.
- SPINICAUDATA Alexander, 1937d:602. Type-locality: Brazil, Santa Catarina,
Nova Teutônia. Distr. — Brazil.
- STERNOLOBATA Alexander, 1958:201, figs. 2 (wing), 8 (male hypopygium).
Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo.
Distr. — Bolivia.
- SUBPACIFERA Alexander, 1942i:61, figs. 26 (wing), 33 (male hypopygium).
Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira.
Distr. — Brazil.
- TENULOBATA Alexander, 1942i:59, figs. 25 (wing), 31 (male hypopygium).
Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira.
Distr. — Brazil.
- TERGIFERA Alexander, 1953g:13, figs. 2 (wing), 10 (male hypopygium).
Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- TREMULA Alexander, 1929a:128, fig. 53 (wing). Type-locality: Chile, Llan-
quihue, Peulla. Distr. — Chile. Ref. — Alexander, 1958:202,
fig. 10 (male hypopygium).
- TUNGURAGUENSIS Alexander, 1940d:218. Type-locality: Ecuador, Mt. Tun-
gurahua. Distr. — Ecuador.
- VARITARSIS Alexander, 1929a:118, fig. 48 (wing). Type-locality: Chile,
Chiloé Is., Castro. Distr. — Chile.



- VIVASBERTHIERI Alexander, 1938h:436. Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela. Ref. — Alexander, 1950a:46.
- WYGODZINSKYI Alexander, 1948a:167, figs. 4 (wing), 18 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Serra dos Órgãos, Sítio Bonfim, Nova Friburgo. Distr. — Brazil.
- XANTHOPTERA XANTHOPTERA Alexander, 1929a:121, fig. 62 (wing). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- ssp. CAYUTUENSIS Alexander, 1944i:130. Type-locality: Chile, Llanquihue, Cayutue. Distr. — Chile.

Subgenus LIMNOPHILELLA Alexander

- Limnophila*, subg. LIMNOPHILELLA Alexander, 1919a:146. Type-species, *epiphragmoides* Alexander (orig. des.) (Neotropical).
- DIVERSIPES (Alexander), 1921d:62 (*Limnophila*). Type-locality: Peru, Loreto, Yurimaguas. Distr. — Ecuador, Peru.
- EPIPHRAGMOIDES (Alexander), 1913b:543 (*Limnophila*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- INQUIETA INQUIETA Alexander, 1943e:208. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- ssp. REFRACTOR Alexander, 1943c:209. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- MANTISSA Alexander, 1966a:221. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- MULTIPICTA Alexander, 1939e:433, fig. 3 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1948a:156, fig. 7 (male hypopygium).
- PATAGONICA (Alexander), 1928b:51 [*Limnophila (Limnophilella)*]. Type-locality: Argentina, Río Negro, L. Nahuel Huapi, Bariloche. Distr. — Chile, Argentina. Refs. — Alexander, 1929a:114; 1948a:159, fig. 8 (male hypopygium).
- SCHUNKEANA Alexander, 1948a:158, figs. 3 (wing), 9 (male hypopygium). Type-locality: Peru, Loreto, Pueallpa, Río Ucayali. Distr. — Peru.
- SUBVICTOR Alexander, 1948a:160. Type-locality: Costa Rica, La Suiza. Distr. — Costa Rica.
- VICTOR (Alexander), 1919a:146, fig. 6 (wing) [*Limnophila (Limnophilella)*]. Type-locality: Panama, Darien, Quebrada Richa, a branch of Río Tapaliza, a branch of Río Puero. Distr. — Panama.

Subgenus MEDIOPHRAGMA Alexander

- Austrolimnophila*, subg. MEDIOPHRAGMA Alexander, 1954a:36. Type-species, *delectissima* Alexander (orig. des.) (Neotropical).
- DELECTISSIMA Alexander, 1954a:37. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- PARAGUAYANA Alexander, 1952a:119. Type-locality: Paraguay, Colonia Carlos Pfannel. Distr. — Paraguay.

Genus CHILELIMNOPHILA Alexander

- CHILELIMNOPHILA Alexander, 1968a:23. Type-species, *Limnophila lyra* Alexander (orig. des.).

LYRA (Alexander), 1952b:108, fig. 7 (male hypopygium) (*Limnophila*).
Type-locality: Chile, Malleco, Curacautín. Distr. — Chile. Ref.
— Alexander, 1968a; figs. 2 (wing), 7 (male hypopygium); 1968c:23,
figs. 1 (wing), 8 (male hypopygium).

Genus CTENOLIMNOPHILA Alexander

CTENOLIMNOPHILA Alexander, 1921d:61. Type-species, *bivena* Alexander
(orig. des.) (Neotropical).

Subgenus ABITAGUA Alexander

Ctenolimnophila, subg. ABITAGUA Alexander, 1944d:315. Type-species, *longifusa* Alexander (orig. des.).

LONGIFUSA Alexander, 1944d:315. Type-locality: Ecuador, Napo-Pastaza,
Abitagua. Distr. — Ecuador. Ref. — Alexander, 1948c:513, fig. 4
(wing).

Subgenus CAMPBELLOMYIA Alexander

Gnophomyia, subg. CAMPBELLOMYIA Alexander, 1925b:70. Type-species,
Gnophomyia alpina Alexander (orig. des.) (Australasian: New
Zealand).

NEOLIMNOPHILOIDES Alexander, 1942i:64, fig. 28 (wing). Type-locality:
Brazil, São Paulo, Juquiá. Distr. — Brazil. Ref. — Alexander,
1948c:513, fig. 3 (wing).

PAULISTAE Alexander, 1943f:301, fig. 10 (wing). Type-locality: Brazil, São
Paulo, Batéia. Distr. — Brazil.

SEVERA Alexander, 1943e:209. Type-locality: Ecuador, Santiago-Zamora,
Zamora. Distr. — Ecuador, Bolivia. Ref. — Alexander, 1948c:513,
fig. 2 (wing).

Subgenus CTENOLIMNOPHILA Alexander

BIVENA Alexander, 1921d:61. Type-locality: Peru, Napo River. Distr. —
Peru, Brazil. Ref. — Alexander, 1948c:513, fig. 1 (wing).

DECISA (Alexander), 1914e:245, pl. 3, fig. 8 (wing) (*Gnophomyia*). Type-
locality: Guyana, Mallali. Distr. — Guyana, Brazil.

FUSCOANALIS Alexander, 1946f:304. Type-locality: Brazil, Pará, Prata.
Distr. — Surinam, Brazil.

Genus EDWARDSOMYIA Alexander

EDWARDSOMYIA Alexander, 1929a:112. Type-species, *chiloensis* Alexander
(orig. des.) Ref. — Alexander, 1948a:175.

CHILOENSIS Alexander, 1929a:112, figs. 47 (wing), 197 (head), 198 (male
hypopygium). Type-locality: Chile, near Puntra. Distr. — Chile.

Genus ELEPHANTOMYIA Osten Sacken

ELEPHANTOMYIA Osten Sacken, 1859:200. Type-species, *Limnobiaorhynchus*
canadensis Westwood (mon. misident.) = *westwoodi* Osten



Sacken (Nearctic). Ref. — Alexander, 1948c:552, figs. (discussion).

Subgenus ELEPHANTOMYIA Osten Sacken

- ALTIKOLA Alexander, 1925a:348, fig. 8 (wing). Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico.
- ANGUSTISSIMA Alexander, 1931c:632. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Mexico. Colombia.
- ARCUARIA Alexander, 1945L:384. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- BANKSI Alexander, 1927c:273. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- BOLIVIENSIS Alexander, 1930c:736. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia. Ref. — Alexander, 1962c:132, fig. 111 (male hypopygium).
- BRUNNEIPENNIS Alexander, 1938b:352. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CHINOPODA Alexander, 1943a:226. Type-locality: Peru, Junin, Tarma, Tulumayo Valley. Distr. — Peru.
- CHIRIQUIANA Alexander, 1945L:385. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- CINCTIVENTRIS Alexander, 1967c:375. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- CITILLARIA Alexander, 1929a:173, fig. 96 (wing). Type-locality: Chile, Llanquihue, Ensenada. Distr. — Chile.
- DECINCTA Alexander, 1962c:133, fig. 112 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- DISTINCTIOR Alexander, 1952a:122. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- PUMIPES Alexander, 1928a:114. Type-locality: Mexico, Chiapas. Hueitepec. Distr. — Mexico.
- HUMILIS Alexander, 1931c:631. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- JUQUIENSIS Alexander, 1945j:27, fig. 11 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- LUTEIANNULATA LUTEIANNULATA Alexander, 1939a:81. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- ssp. CHIRIQUIENSIS Alexander, 1941g:351 (also 1946h:98). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- MERIDIONALIS Alexander, 1913b:489 (nom. nov. for *longirostris* Williston). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.
- longirostris* Williston, 1896a:288 (*Toxohrina*; preocc. Loew, 1851). Type-locality: "St. Vincent".
- PICTIVENTRIS Alexander, 1945o:603. Type-locality: Ecuador, Santiago-Zamora, Zamora. Distr. — Ecuador.
- PRIMITIVA Alexander, 1943b:121. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PRIMOGENIA Alexander, 1948c:554, fig. 36 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.



- SERRULIFERA Alexander, 1949c:592. Type-locality: Peru, Junín, Tarma, Utcuyacu. Distr. — Peru.
- SETULISTYLA Alexander, 1938b:351. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Venezuela, Ecuador.
- SORDIDIPES Alexander, 1962c:134, fig. 109 (wing). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SUBHUMILIS Alexander, 1938b:350. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- TARSALBA Alexander, 1928h:363. Type-locality: Surinam, Paramaribo. Distr. — Surinam.
- TENUISSIMA Alexander, 1944b:177, fig. 10 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- TIGRIVENTRIS Alexander, 1942e:253. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador.
- UNICINCTA Alexander, 1962c:135, figs. 110 (wing), 113 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal. Distr. — Bolivia.
- WESTWOODI ANTILLARUM Alexander, 1932:386, fig. 13 (wing). Type-locality: Cuba, Trinidad Mts., Buenos Aires. Distr. — Cuba, Haiti.

Subgenus ELEPHANTOMYINA Alexander

- Elephantomyia*, subg. ELEPHANTOMYINA Alexander, 1938b: 349. Type-species, *supernumeraria* Alexander (orig. des.).
- SUPERNUMERARIA Alexander, 1921d:60. Type-locality: Peru, Loreto, Napo River. Distr. — Peru. Ref. — Alexander, 1938b:349.

Subgenus XENOELEPHANTOMYIA Alexander

- Elephantomyia*, subg. XENOELEPHANTOMYIA Alexander, 1965a:402. Type-species, *penai* Alexander (orig. des.).
- PENAI Alexander, 1965a:403. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.

Genus EPIPHRAGMA Alexander

- Limnophila*, subg. EPIPHRAGMA Osten Sacken, 1859:238. Type-species, *pa-
vonina* Osten Sacken (Coquillett, 1910:539) = *fasciapennis* (Say).
Ref. — Alexander, 1948a:168, fig. (general discussion).
- ADSPERSA (Wiedemann), 1828:550 (*Limnobia*). Type-locality: "Brazil".
Distr. — Guyana, Brazil. Ref. — Alexander, 1913b:551, fig. 34
(wing).
- AMPHILEUCA Alexander, 1941c:329. Type-locality: Panama, Chiriquí, Po-
trerillos. Distr. — Panama.
- ANNULICORNE Alexander, 1921b:176. Type-locality: Argentina, Tucumán,
Quebrada Famaila. Distr. — Peru, Bolivia, Argentina.
- ATROTERMINATA Alexander, 1943g:526. Type-locality: Peru, Ayacucho, La
Mar, Yanamonte. Distr. — Peru.
- AURICOSTA Alexander, 1939d:116. Type-locality: Lesser Antilles: Dominica,
Valle Nuevo, Southeast of Constanza. Distr. — Dominica.
- BINIGROCINCTA Alexander, 1962c:120, figs. 99 (wing), 102 (male hypopygium).
Type-locality: Bolivia, Yungas de Arepuchó, Silhuencas. Distr. —
Bolivia.



- GUSCKI Alexander, 1913b:540, fig. 36 (wing). Type-locality: Hispaniola: Dominican Republic, San Francisco Mts. Distr. — Haiti, Dominican Republic.
- CELATOR Alexander, 1946b:238. Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- CIRCINATA Osten Sacken, 1886a:9. Type-locality: Costa Rica, Mt. Irazu. Distr. — Costa Rica, Panama.
- CLAUDIA Alexander, 1945o:597. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- CORDILLERENSE Alexander, 1913c:202, pl. 5, fig. 8 (wing). Type-locality: Colombia, Cauca, Popayán. Distr. — Colombia, Peru, Bolivia.
- CUBENSE Alexander, 1929e:403. Type-locality: Cuba, Oriente, Los Llanos, Maisi. Distr. — Cuba.
- CYNOTIS Alexander, 1945o:595. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- DELIBERATA Alexander, 1939b:189. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- DELICATULA Osten Sacken, 1887b:208. Type-locality: "Colombia". Distr. — Colombia, Venezuela.
- DIADEMA Alexander, 1939b:193. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- ENIXA Alexander, 1939b:190. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Venezuela, Ecuador.
- ENIXOIDES Alexander, 1962c:121. Type-locality: Bolivia, Yungas de Forestal. Distr. — Bolivia.
- FABRICII Alexander, 1913b:537, fig. 33 (wing) (nom. nov. for *maculata* Fabricius). Type-locality: "South America." Distr. — Peru, Brazil. *maculata* Fabricius, 1805:27 (*Tipula*; procc. Linnaeus, 1758). Type-locality: "South America".
- FARRI Alexander, 1951a:101. Type-locality: Guatemala, Chicacao, El Naranjo. Distr. — Guatemala.
- FELIX Alexander, 1943g:522. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- FILIFORME Alexander, 1940d:219. Type-locality: Ecuador, Napo-Pastaza, Río Pastaza, San Francisco. Distr. — Ecuador. Ref. — Alexander, 1953c:33, figs. 2 (antenna), 3 (male hypopygium).
- GAIGEI Alexander, 1928h:362. Type-locality: Panama, Chiriqui, Boquete. Distr. — Panama.
- GENUALE Alexander, 1934a:64. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.
- GRACILICORNE Alexander, 1916a:21, pl. 1, fig. 11 (wing). Type-locality: Colombia, Magdalena, La Cumbre. Distr. — Colombia.
- HIRTISTYLATA Alexander, 1939b:198. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1953c:33, fig. 4 (male hypopygium).
- HISTRIO Schiner, 1868:41. Type-locality: "Venezuela" (not "Colombia", as stated). Distr. — Venezuela. Refs. — Alexander, 1947d:45; 1948a:171.
- IMITANS Alexander, 1913b:536, fig. 35 (wing). Type-locality: Bolivia, San Antonio. Distr. — Brazil, Bolivia, Argentina. Closely related to *solatrix* (Osten Sacken). Ref. — Bruch, 1939a:16-20; text fig. 3 (pupa). pl. 4, fig. 14 (entire female); pl. 7, figs. 23, 24 (immature stages); pls. 9, 10, figs. 17-43 (details of larva, pupa).

- IMMACULIPES Alexander, 1941c:332. Type-locality: Panama, Chiriqui, Potrerrillos. Distr. — Panama.
- INAEQUICINCTA Alexander, 1941c:331. Type-locality: Panama, Chiriqui, Potrerrillos. Distr. — Panama, Costa Rica, Venezuela.
- INORNATIPES Alexander, 1939d:118. Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- INTERSPERSA Alexander, 1953c:34, figs. 1 (wing), 5 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Bolivia.
- JUQUICOLA Alexander, 1945c:266. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- JURATOR Alexander, 1948a:172. Type-locality: Peru, Junín, Huacapistana. Distr. — Peru.
- MELAXANTHA Alexander, 1943g:524. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- MEPHISTOPHELICA Alexander, 1947f:94, fig. 11 (wing). Type-locality: Venezuela, Aragua, Choroní. Distr. — Venezuela.
- MITHRAS Alexander, 1949d:253. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- MUSCICOLA Alexander, 1943g:528. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- NEBULOSA (Bellardi), 1861:206, pl. 1, fig. 4 (wing) (*Tipula*). Type-locality: "Mexico". Distr. — Mexico.
- NEPHELE Alexander, 1948a:174. Type-locality: Peru, Junín, Huacapistana. Distr. — Peru.
- NIGRIPLEURALIS Alexander, 1945j:17, fig. 8 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NIGROPLAGIATA Alexander, 1939b:187. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Peru. Ref. — Alexander, 1953c:35, fig. 6 (male hypopygium)
- OREONYMPHA Alexander, 1928a:110. Type-locality. Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- OXYPHALLUS Alexander, 1939b:196. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1953c:36, fig. 9 (male hypopygium).
- PARVISETA Alexander, 1941d:122. Type-locality: Ecuador, Mt. Tungurahua. Distr. — Ecuador. Ref. — Alexander, 1953c:36, figs. 7 (antenna), 8 (male hypopygium).
- PERSANCTA Alexander, 1938f:248. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Venezuela, Brazil. Ref. — Alexander, 1948a:153, fig. 5 (wing).
- PETALINA Alexander, 1947f:95. Type-locality: Venezuela, Aragua, Choroní. Distr. — Venezuela.
- PIAEOXANTHA Alexander, 1944d:313. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Cunibunda. Distr. — Ecuador. Ref. — Alexander, 1953c:37.
- FUNCTATISSIMA (Wiedemann), 1828:30 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil.
- PUPILLATA Alexander, 1913b:539, fig. 37 (wing). Type-locality: Brazil, Mato Grosso, Chapada [dos Guimarães]. Distr. — Brazil.
- SACKENI Williston, 1896a:294. Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.



- SAPPHO Alexander, 1943g:530. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- SEPTUOSA Alexander, 1962c:123, fig. 103 (male hypopygium). Type-locality: Bolivia, Yungas de Arcpucho, Silhuencas. Distr. — Bolivia.
- SERRISTYLA Alexander, 1945j:19, fig. 9 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Tercoópolis, Serra dos Órgãos. Distr. — Brazil.
- SOLATRIX Osten Sacken, 1859:238; 1869:195, pl. 2, fig. 8 (wing). Type-locality: U.S.A., District of Columbia. Distr. — Southern U.S.A., to Mexico. Refs. — Alexander, 1919b:915, fig. 159 (wing); 1921f:844, pls. 37, 38 (immature stages); 1939b:186.
- SUBENIXA Alexander, 1939b:192. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- SUBSOLATRIX Alexander, 1939b:186. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Venezuela, Ecuador.
- SYBARITICA Alexander, 1947f:92, fig. 10 (wing). Type-locality: Venezuela, Aragua, Choroní. Distr. — Venezuela.
- VARIA (Wiedemann), 1828:573 (*Limnobia*). Type-locality: "Brazil". Distr. — Panama, Venezuela, Guyana, Ecuador, Peru, Brazil.
- XANTHOMELA Alexander, 1939b:195. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

Genus GYNOPLISTIA Westwood

- GYNOPLISTIA Westwood, 1835a:280. Type-species, *nervosa* Westwood (Brunetti, 1918:332) = *vilis* (Walker) (Australasian).
- Ctedonia* Philippi 1865:602. Type-species, *bicolor* Philippi (Enderlein, 1917:57). Refs. — Alexander, 1929a:148; 1948c:524.
- Scepasma* Enderlein, 1917:60. Type-species, *Ctedonia bipunctata* Philippi (orig. des.).

Subgenus DIRHIPIS Enderlein

- DIRHIPIS Enderlein, 1917:58 (as genus). Type-species, *riedeliana* Enderlein (orig. des.).
- FUSCA (Jaenicke), 1867:322 (*Ctedonia*). Type-locality: "Chile". Distr. — Chile.
- RIEDELIANA (Enderlein), 1917:58, figs. 1 (male antenna), 2 (wing), 3 (palpus) (*Dirhipis*). Type-locality: Chile, Concepción. Distr. — Chile.
- STRIATIPENNIS Alexander, 1928e:246 (for 1927) [*Gynoplistia* (*Ctedonia*)]. Type-locality: Chile, Valdivia, Panguipulli. Distr. — Chile.

Subgenus GYNOPLISTIA Westwood

- BIARMATA Alexander, 1931c:629. Type-locality: Chile, Malleco, Angol. Distr. — Chile.
- BICOLOR BICOLOR (Philippi), 1865:603 (*Ctedonia*). Type-locality: Chile, Llanquihue, Puerto Montt. Distr. — Chile. Refs. — Alexander, 1929a:158, fig. 82 (wing); 1948c:527, fig. 14 (wing).
- ssp. DILLMANI Alexander, 1952b:109. Type-locality: Chile, Arauco, Cerros de Nahuelbuta. Distr. — Chile.
- BIPUNCTATA (Philippi), 1865:603 (*Ctedonia*). Type-locality: Chile, Valdivia, Pantanos. Distr. — Chile. Refs. — Alexander, 1929a:151; Enderlein, 1917:60 (*Scepasma*).



- ELNORAE (Alexander), 1929a:159 (*Ctedonia*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- FLAVIPENNIS (Philippi), 1865:602 (*Ctedonia*). Type-locality: Chile, Valdivia. Distr. — Chile.
- GILVIPPENNIS Alexander, 1928e:248 [*Gynoplusia* (*Ctedonia*)]. Type-locality: Chile, Valdivia, Panguipull. Distr. — Chile, Argentina.
- HYLONYMPHA Alexander, 1929a:163, fig. 84 (wing). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile, Argentina.
- LEUCOPEZA LEUCOPEZA Alexander, 1929a:161, figs. 78 (wing), 213 (male hypopygium). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
ssp. POSTICA Alexander, 1929a:162, fig. 93 (wing). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- MANICATA Alexander, 1929a:160, fig. 77 (wing). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- PENANA Alexander, 1967e:373. Type-locality: Chile, Cordillera Curicó, El Coigual. Distr. — Chile.
- PICTIPPENNIS (Philippi), 1865:603 (*Ctedonia*). Type-locality: Chile Valdivia. Distr. — Chile.
- SCHACHOVSKOYANA Alexander, 1960:17. Type-locality: Argentina, Neuquén, Lago Lacar. Distr. — Argentina.
- TRISTILLATA Alexander, 1929a:165, fig. 85 (wing). Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Chile.
- VARIATA VARIATA Alexander, 1931c:628. Type-locality: Chile, Malleco, Angol. Distr. — Chile.
ssp. FLAVIDULA Alexander, 1952b:109. Type-locality: Chile, Malleco, Rio Blanco. Distr. — Chile.
- VARICALCARATA Alexander, 1929a:165, fig. 86 (wing). Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Chile.
- VARIPEX Alexander, 1929a:162, figs. 83 (wing), 214, 215 (male hypopygium). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.

Genus HEXATOMA Latreille

HEXATOMA Latreille, 1809:260. Type-species, *nigra* Latreille (mon.).

Subgenus CLADOLIPES Loew

- CLADOLIPES Loew, 1865:424 (as genus). Type-species, *simplex* Loew (mon.) (Western Palaearctic).
- CISATLANTICA Alexander, 1937d:608. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1948c:527, fig. 16 (wing).

Subgenus ERIOCERA Macquart

- Caloptera* Guérin-Méneville, 1831: pl. 20, fig. 2 (as genus). Type-species, *fasciata* Guérin-Méneville (orig. des.; as gen. n., sp. n.) [See note in Alexander, 1965d:69].
- ERIOCERA Macquart, 1838a:78(74) (as genus). Type-species, *Limnobia nigra* Wiedemann (mon.) = *wiedemanni* Alexander (Neotropical).
- Arrhenica* Osten Sacken, 1859:242. Type-species, *spinosa* Osten Sacken (Coquillett, 1910:510).



- Pentoptera* Schiner, 1863:220. Type-species, *Tipula chirothecata* Scopoli (orig. des.). Ref. — Alexander, 1948c:529, figs.
- ACUNAI (Alexander), 1928b:56 (*Eriocera*). Type-locality: Cuba, Pinar del Río, Sierra Rangcl, Las Ánimas. Distr. — Cuba.
- ATHIEREA (Alexander), 1916c:346, fig. 5 (wing) (*Eriocera*). Type-locality: Hispaniola: Dominican Republic, San Lorenzo. Distr. — Dominican Republic.
- AGLAIJA Alexander, 1945c:601. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- AMAZONICOLA (Alexander), 1920c:144 (*Eriocera*). Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- ANDICOLA (Alexander), 1923c:312 (*Eriocera*). Type-locality: Argentina, Jujuy, Ciudad. Distr. — Argentina.
- ANTENNATA (Alexander), 1930c:735 (*Eriocera*). Type-locality: Colombia, Magdalena, Seville. Distr. — Colombia.
- ARGENTINA (Alexander), 1919a:148, fig. 7 (wing) (*Eriocera*). Type-locality: Argentina, Catamarca, La Plaza. Distr. — Argentina.
- ATROSIGNATA Alexander, 1941h:259. Type-locality: Peru, Huanuco, Puma-huasi. Distr. — Peru.
- AURANTIONOTA Alexander, 1939a:78. Type-locality: Mexico, Chiapas, Huixtla. Distr. — Mexico.
- BATESI (Alexander), 1921d:67 (*Pentoptera*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- BEEBEANA Alexander, 1950a:49, fig. 23 (wing). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- BEQUARTIANA Alexander, 1938h:437. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia.
- BIFURCATA Alexander, 1947:97. Type-locality: Venezuela, Aragua, Choroni. Distr. — Venezuela.
- BITUBERCULATA (Macquart), 1838a:76(72) (*Limnobia*). Type-locality: Unknown; Osten Sacken, 1869:248 says "Brazil." Identity doubtful.
- BRACONIDES (Enderlein), 1912:47, fig. B1 (wing) (*Eriocera*). Type-locality: "Colombia". Distr. — Colombia.
- BREVIUSCULA (Alexander), 1929b:97 (*Eriocera*). Type-locality: Peru, Putumayo, La Chorerra. Distr. — Peru. Ref. — Alexander, 1948c:527, fig. 19 (wing).
- BRUNERI (Alexander), 1928b:55 (*Eriocera*). Type-locality: Cuba, Oriente, Jarahuca. Distr. — Cuba.
- BRUNNEIPES (Williston), 1900:227, p. 4, fig. 5 (*Eriocera*). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- CABRALENSIS Alexander, 1945j:24, fig. 6 (wing). Type-locality: Brazil, São Paulo, Pôrto Cabral, Alto Paraná River. Distr. — Brazil.
- CAMINARIA (Wiedemann), 1828:79 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Alexander, 1922c:53.
- CANDIDIPES (Alexander), 1923a:20 (*Pentoptera*). Type-locality: Venezuela, Táchira, Táchira. Distr. — Venezuela.
- CAPTIOSA Alexander, 1945o:600. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CARRERAI Alexander, 1942i:65, fig. 29 (wing). Type-locality: Brazil, São Paulo, Guarujá. Distr. — Brazil.
- CHALYBEIVENTRIS Alexander, 1949c:591. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.



- CHRYSOPTERA (Walker), 1856:438 (*Limnobia*). Type-locality: "South America." Distr. — Brazil. Ref. — Alexander, 1914b:214, pl. 9, fig. 13 (wing).
- CHRYSOPTEROIDES (Alexander), 1922c:58 (*Eriocera*). Type-locality: "Brazil". Distr. — Brazil.
- COLUMBIANA (Alexander), 1919a:147, fig. 8 (wing) (*Eriocera*). Type-locality: Colombia, Bolívar. Distr. — Costa Rica, Colombia.
- CONJUNCTA (Alexander), 1914a:44, fig. 2 (wing) (*Penthoptera*). Type-locality: Guatemala, Patulue. Distr. — Guatemala.
- CORNIGERA (Alexander), 1914a:40, fig. 3 (wing) (*Eriocera*). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Bolivia.
- COROICOENSIS Alexander, 1962c:128, fig. 104 (wing). Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia.
- CRAMPTONI (Alexander), 1928g:23 (*Eriocera*). Type-locality: Jamaica, Cinchona. Distr. — Jamaica.
- CUBENSIS (Alexander), 1926b:229 (*Eriocera*). Type-locality: Cuba, Trinidad Mountains. Distr. — Cuba. Refs. — Alexander, 1932:370, fig. 12 (wing); 1948c:527, fig. 18 (wing).
- DOMINGENSIS (Alexander), 1916c:346, fig. 6 (wing) (*Eriocera*). Type-locality: Hispaniola: Dominican Republic, Sánchez. Distr. — Dominican Republic.
- ERYTHRAEA (Osten Sacken), 1886a:11 (*Eriocera*). Type-locality: Guatemala, San Gerónimo. Distr. — Guatemala, Costa Rica.
- ESMERALDA Alexander, 1952a:120. Type-locality: Ecuador, Esmeraldas, Río Salinas. Distr. — Ecuador.
- EXQUISITA (Alexander), 1914c:123, fig. 5 (wing) (*Eriocera*). Type-locality: Costa Rica, Alajuela. Distr. — Costa Rica.
- FARRIANA Alexander, 1964a:63, fig. 62 (wing). Type-locality: Jamaica, Portland, Hardwar Gap. Distr. — Jamaica.
- FASCIATA (Guérin-Méneville), 1830:287, pl. 20, fig. 2 (wing) (*Evanioptera*). Type-locality: Brazil, Santa Catarina. Distr. — Brazil.
- FERAX Alexander, 1937b:242. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- FLAMMEINOTA (Alexander), 1930c:734 (*Eriocera*). Type-locality: Brazil, Minas Gerais. Distr. — Brazil.
- FLAMMEIPENNIS Alexander, 1942g:178, fig. 5 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- FLAVICEPS (Wiedemann), 1828:550 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil.
- FLAVIDA (Williston), 1900:227 (*Eriocera*). Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- FULVITHORAX Alexander, 1966c:473. Type-locality: Argentina, Santa Cruz Lago Argentino, Calafate. Distr. — Argentina.
- GOMESIANA Alexander, 1945e:225. Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia. Distr. — Brazil.
- GOYAZENSIS Alexander, 1940f:396, fig. 6 (wing). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- GRACILIS (Osten Sacken), 1886a:12 (*Eriocera*). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- HAEMORRHOA (Osten Sacken), 1886a:11 (*Eriocera*). Type-locality: Mexico, Mexico, Veracruz, Orizaba. Distr. — Mexico.



- INTERLINEATA Alexander, 1934a:67. Type-locality: Panama, Chiriqui, Finca Parada. Distr. — Mexico, Costa Rica, Panama.
- INTERMEDIA (Alexander), 1927c:271 [*Eriocera* (*Penthoptera*)]. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- JOCULARIS Alexander, 1948c:534. Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- JULIANA Alexander, 1937e:529, fig. 6 (wing). Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- JURATA Alexander, 1936d:759. Type-locality: Brazil, Rio de Janeiro, Itaitaia. Distr. — Brazil.
- KAJETURENSIS (Alexander), 1914a:41, fig. 1 (wing) (*Eriocera*). Type-locality: Guyana, Kaieteur Falls, Potaro River. Distr. — Guyana.
- LADDEYI Alexander, 1942d:142. Type-locality: Ecuador, Pichincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- LATICOSTATA Alexander, 1937d:607. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LATISSIMA (Alexander), 1922c:55 (*Eriocera*). Type-locality: "Venezuela" ["Novara Reise"]. Distr. — Venezuela.
- LESSEPSI (Osten Sacken), 1886a:13 (*Eriocera*). Type-locality: "Panama". Distr. — Panama.
- LONGIPENNIS (Alexander), 1923a:17 (*Eriocera*). Type-locality: Venezuela, Macuto. Distr. — Venezuela.
- LONGISTYLA (Alexander), 1914a:39, fig. 6 (wing) (*Eriocera*; nom. nov. for *erythrocephala* Fabricius). Type-locality: "South America". Distr. — Guyana.
erythrocephala Fabricius, 1805:31 (*Tipula*; procc. De Geer, 1776). Type-locality: "South America".
- LOPESI Alexander, 1942g:178, fig. 6 (wing). Type-locality: Brazil, Goiás, Campinas. Distr. — Brazil.
- LUTEICOLOR Alexander, 1968c:215. Type-locality: Puerto Rico, Maricao State Forest. Distr. — Puerto Rico.
- LUTEITARSIS Alexander, 1964a:64, fig. 63 (wing). Type-locality: Jamaica, St. Thomas, Whitfield Hall. Distr. — Jamaica.
- MACQUARTI (Enderlein), 1912:45 (*Eriocera*; nom. nov. for *nigra* Macquart). Type-locality: "Brazil." Distr. — Brazil.
nigra Wiedemann of Macquart, 1838a:79(75) pl. 10, fig. 2 (*Eriocera*).
- MACROCERA (Alexander), 1914a:40, fig. 4 (wing) (*Eriocera*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- MAGISTRA Alexander, 1942d:144. Type-locality: Ecuador, Napo-Pastaza, Río Huagra Yacu. Distr. — Ecuador.
- MAGNIFICA (Alexander), 1914a:37, fig. 7 (wing) (*Eriocera*). Type-locality: Guatemala, Alta Verapaz, Cacao Trece Aguas. Distr. — Guatemala.
- MALDONADOI Alexander, 1952d:245. Type-locality: Venezuela, Camp Benitez, Marahuaca Region, foot of Pacaraima Mts. Distr. — Venezuela.
- MANABIANA Alexander, 1942d:141. Type-locality: Ecuador, Pichincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- MARGARITAE Alexander, 1954a:38. Type-locality: Brazil, Rio de Janeiro — São Paulo Road. Distr. — Brazil.
- MELANACRA (Wiedemann), 1828:548 (*Eriocera*). Type-locality: "Brazil". Distr. — Brazil.



- MELANOLITHA (Alexander), 1927c:272 [*Eriocera* (*Pentoptera*)]. Type-locality: Guatemala, Antigua, El Salto. Distr. — Mexico, Guatemala, El Salvador.
- MELANONOTA Alexander, 1968c:216. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- MELINA (Alexander), 1922c:63 (*Eriocera*). Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- MESOXANTHA (Osten Sacken), 1886a:10 (*Eriocera*). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- MULTIGUTTULA Alexander, 1939d:124. Type-locality: Hispaniola: Dominican Republic, Valle Nuevo, southeast of Constanza. Distr. — Dominican Republic.
- MYRTEA (Alexander), 1922c:62 (*Eriocera*). Type-locality: "Brazil". Distr. — Brazil.
- NEOSAGA Alexander, 1947d:51, fig. 5 (wing). Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- NGROCHALYBEA (Alexander), 1922c:57 (*Eriocera*). Type-locality: "Brazil". Distr. — Brazil.
- OBSOLETA (Williston), 1900:227 (*Eriocera*). Type-locality: Honduras, Ruan-tan Is. Distr. — Honduras.
- OCELLIFERA (Alexander), 1915c:104 (*Eriocera*). Type-locality: Puerto Rico, Mayaguez. Distr. — Puerto Rico. Ref. — Alexander, 1932:370.
- OGLOBLINI Alexander, 1935a:325. Type-locality: Argentina, Misiones, Lorcto. Distr. — Argentina.
- OHAUSIANA (Enderlein), 1912:45, fig. A1 (*Eriocera*). Type-locality: Ecuador, Sabanilla. Distr. — Colombia, Ecuador, Peru, Bolivia.
- OPULENTA Alexander, 1951a:104. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- ORNATICORNIS Alexander, 1939d:123, fig. 7 (wing). Type-locality: Cuba, Oriente, Sierra del Cobre, Loma del Gato. Distr. — Cuba.
- PALLIDIPES (Alexander), 1926a:172, fig. 10 (wing) (*Eriocera*). Type-locality: Mexico, Veracruz. Distr. — Mexico.
- PATENS Alexander, 1938b:348, fig. 6 (wing). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador. Ref. — Alexander, 1948c:527, fig. 20 (wing).
- PERDECORA (Alexander), 1914b:213, pl. 9, fig. 11 (wing) (*Eriocera*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- PERENENSIS (Alexander), 1923a:18 (*Eriocera*). Type-locality: Peru, Junín, Colonia del Perené. Distr. — Peru.
- PEREXIGUA Alexander, 1941h:258. Type-locality, Peru, Huanuco, Pumahuasi. Distr. — Peru.
- PERFESTIVA Alexander, 1951a:106. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- PERLAETA (Alexander), 1922c:54 (*Eriocera*). Type-locality: "Colombia". Distr. — Brazil, Colombia.
- PERPULCHRA (Alexander), 1914a:38, fig. 8 (wing) (*Eriocera*). Type-locality: "Northern Brazil". Distr. — Brazil.
- FERRARA Alexander, 1936a:15. Type-locality: Brazil, Rio de Janeiro, Rezen-de, Campo Belo. Distr. — Brazil.
- PERUVIANA (Alexander), 1914a:42, fig. 5 (wing) (*Eriocera*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- PERVERSA Alexander, 1937c:495. Type-locality: Ecuador, Napo-Pastaza, Río Blanco, Yunquilla. Distr. — Ecuador.



- PERVIA Alexander, 1966a:227. Type-locality: Ecuador, Napo-Pastaza, Libertad. Distr. — Ecuador.
- PIATRIX Alexander, 1942d:143. Type-locality: Ecuador, Pichincha, Santo Domingo de los Colorados. Distr. — Ecuador.
- PLAUMANNI PLAUMANNI Alexander, 1936e:61. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ssp. LATAURATA Alexander, 1937d:608. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PLUMBEICOLOR Alexander, 1942d:145. Type-locality: Ecuador, Los Ríos, Playas de Montalvo. Distr. — Ecuador.
- PLUMBEINOTA Alexander, 1940b:159. Type-locality: Mexico, Guerrero, Taxco. Distr. — Mexico.
- PRETIOSA (Osten Sacken), 1886a:12 (*Eriocera*). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- PROPINQUUA Alexander, 1936d:760. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- PULCHRIPES (Alexander), 1922c:59 (*Eriocera*). Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia.
- PULLATIPES Alexander, 1951a:105. Type-locality: Peru, Huanuco, Fundo Sinchono, Cordillera Azul. Distr. — Peru.
- REVERENTIA Alexander, 1949b:480, fig. 12 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- RORAIMELLA Alexander, 1935a:325. Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- RUFICORNIS (Macquart), 1838b:176 (*Cylindrotoma*). Type-locality: "Brazil". Distr. — Brazil.
- RUPUNUNI Alexander, 1945L:381. Type-locality: Guyana, Pirara, Rupununi. Distr. — Guyana.
- SAGA Alexander, 1940e:900, fig. 3 (wing). Type-locality: Venezuela, Trujillo, Quebrada de las Avispas. Distr. — Venezuela.
- SANCTAEMARTAE (Alexander), 1919a:148, fig. 9 (wing) (*Pentoptera*). Type-locality: Colombia, Magdalena, Mt. Santa Marta. Distr. — Colombia.
- SCHAUSEANA Alexander, 1966c:474. Type-locality: Guatemala, Cayuga. Distr. — Guatemala.
- SCHILDEANA Alexander, 1966c:475, fig. 5 (wing). Type-locality: Costa Rica, San Matco, Higuito. Distr. — Costa Rica.
- SCHINERI (Alexander), 1922c:60 (*Eriocera*; nom. nov. for *fuliginosa* Schiner). Type-locality: "Colombia". Distr. — Colombia, Venezuela.
- fuliginosa* Schiner, 1868:42 (*Pentoptera*; preocc. Osten Sacken, 1859). Type-locality: "Colombia".
- SCULLENIANA Alexander, 1965a:402. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- SEMIRUFA (Alexander), 1927c:273 (*Eriocera*; nom. nov. for *dimidiata* Alexander). Type-locality: Venezuela, Táchira, Táchira. Distr. — Venezuela.
- dimidiata* Alexander, 1923a:17 (*Eriocera*; preocc. Henriksen, 1922). Type-locality: Venezuela, Táchira, Táchira.
- SPECIOSA (Alexander), 1914e:252, pl. 3, fig. 10 (wing) (*Eriocera*). Type-locality: Guyana, Bartica. Distr. — Guyana.
- STOLIDA Alexander, 1934a:68. Type-locality: Panama, Chiriqui, Finca Parada. Distr. — Panama.



- SUBGRACILIS Alexander, 1939a:79. Type-locality: Mexico, Chiapas, Huixtla. Distr. — Mexico.
- SUBLIMA (Alexander), 1914b:214, pl. 9, fig. 12 (wing) (*Eriocera*). Type-locality: Brazil, Minas Gerais. Distr. — Brazil.
- SUBCELLATA Alexander, 1964a:65, fig. 64 (wing). Type-locality: Jamaica, Portland, Hardwar Gap. Distr. — Jamaica.
- SUBSAGA Alexander, 1941h:261. Type-locality: Peru, Junin, Satipo. Distr. — Peru.
- SUBSTOLIDA Alexander, 1939a:77. Type-locality: Mexico, Chiapas, Finca Belén. Distr. — Mexico.
- TAENIOPTERA (Wiedemann), 1828:28 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Alexander, 1922c:56.
- TENE BROSA (Walker), 1856:439 (*Limnobia*). Type-locality: "South America". Distr. — South America.
- THOLOPA (Alexander), 1931c:630 (*Eriocera*). Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- TOWNSENDI (Alexander), 1914a:42 (*Eriocera*). Type-locality: Mexico, Chihuahua, Sierra Madre. Distr. — Mexico.
- TRANQUILLA (Alexander), 1922c:59 (*Eriocera*). Type-locality: "Brazil" (Blumenau). Distr. — Brazil.
- TRIALROSNATA Alexander, 1941h:262. Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- TRIFASCIATA (Röder), 1885:338 (*Eriocera*). Type-locality: "Puerto Rico". Distr. — Puerto Rico. Ref. — Alexander, 1932:370.
- VARIEGATA Alexander, 1936d:761. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- VENAVITTA Alexander, 1940f:394, fig. 5 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- VIRGULATIVENTRIS (Enderlein), 1912:47, fig. C1 (*Eriocera*). Type-locality: Colombia, Tolima, Natagaima. Distr. — Colombia.
- VITTINERVIS Alexander, 1936a:17. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- WEBERI Alexander, 1951a:103. Type-locality: Guyana, Courantyne River, Waricabra Falls. Distr. — Guyana.
- WIEDEMANNI Alexander, 1933:149 (nom. nov. for *nigra* Wiedemann). Type-locality: "Brazil". Distr. — Brazil.
nigra Wiedemann, 1828:27 (*Limnobia*; preocc. Latreille, 1809). Type-locality: "Brazil".
- WILLIAMSONI (Alexander), 1923a:19 (*Eriocera*). Type-locality: Peru, Colonia del Perené. Distr. — Peru.
- WILLISTONI (Alexander), 1913b:488 (*Eriocera*; nom. nov. for *fasciata* Williston). Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
fasciata Williston, 1900:226 (*Eriocera*; preocc. Guérin-Méncville, 1830). Type-locality: Mexico, Morelos, Cuernavaca.
- ZONATA (Osten Sacken), 1886a:10 (*Eriocera*). Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico

Genus LECTERIA Osten Sacken

- LECTERIA Osten Sacken, 1887a:206. Type-species, *Tipula armillaris* Fabricius (orig. des.). Refs. — Alexander, 1913b:493; 1948a:176.



Subgenus LECTERIA Osten Sacken

- ARMILLARIS (Fabricius), 1805:26 (*Tipula*). Type-locality: "South America".
Distr. — Costa Rica, Panama, Peru, Brazil. Refs. — Alexander, 1913b:497, fig. 42 (wing); 1948a:179, figs. 19-20 (wing).
- CALOPUS (Walker), 1856:439 (*Limnobia*). Type-locality: "South America."
Distr. — South America.
- MATTOGROSSAE Alexander, 1913b:496, fig. 39 (wing). Type-locality: Brazil, Mato Grosso, Corumbá. Distr. — Brazil.

Subgenus PSARONIUS Enderlein

- PSARONIUS Enderlein, 1912:50 (as genus). Type-species, *lituratus* Enderlein (orig. des.) = *obscura* (Fabricius). Ref. — Alexander, 1948a:176, figs.
- ABNORMIS Alexander, 1914b:211, pl. 9, fig. 9 (wing). Type-locality: "Paraguay". Distr. — Bolivia, Argentina, Paraguay. Ref. — Alexander, 1948a:179, fig. 24 (wing).
- BREVISECTOR Alexander, 1936d:761. Type-locality: Brazil, Bahia, Ilhéus. Distr. — Brazil.
- BREVIPTIBIA (Alexander), 1920a:74 (*Psaronius*). Type-locality: Brazil, Pará, Óbidos. Distr. — Brazil.
- FUSCIPENNIS (Alexander), 1914e:249 (*Psaronius*). Type-locality: Guyana, Bartica. Distr. — Guyana, Brazil.
- LEGATA Alexander, 1948a:179. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- MANCA (Alexander), 1921d:64 (*Psaronius*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- OBLITERATA Alexander, 1913b:494, fig. 41 (wing). Type-locality: Guyana, Tukeit. Distr. — Guyana.
- OBSCURA (Fabricius), 1805:27 (*Tipula*). Type-locality: "South America."
Distr. — Guyana, Brazil. Ref. — Alexander, 1913b:495.
liturata Enderlein, 1912L:50, fig. E1 (wing) (*Psaronius*). Type-locality: Guyana, Demerara.
- PALLIPES (Alexander), 1920a:73 (*Psaronius*). Type-locality: Brazil, Pará, Prata. Distr. — Brazil.
- PYGMAEA (Alexander), 1914e:249, pl. 4, fig. 9 (wing) (*Psaronius*). Type-locality: Guyana, Bartica. Distr. — Guyana, Peru, Brazil.
- TRIANGULIFERA (Alexander), 1921d:63 (*Psaronius*). Type-locality: Peru, Loreto, Yuriniaguas. Distr. — Peru.

Genus LIMNOPHILA Macquart

- LIMNOPHILA Macquart, 1834:95. Type-species, *Limnobia pictipennis* Meigen Westwood, 1840:128). Ref. — Alexander, 1948c:513, figs. (general discussion).

Subgenus ARAUCOLIMNOPHILA Alexander

- Limnophila*, subg. ARAUCOLIMNOPHILA Alexander, 1939f:177. Type-species, *wolffhuegeli* Alexander (orig. des.).



WOLFFHUEGELI Alexander, 1939f:177. Type-locality: Chile, Llanquihue, Cayutue, near Lago de Todo los Santos. Distr. — Chile. Ref. — Alexander, 1948c:515, fig. 7 (wing).

Subgenus HABROLIMNOPHILA Alexander

Limnophila, subg. HABROLIMNOPHILA Alexander, 1968a:23. Type-species, *Austrolimnophila celestissima* Alexander (orig. des.).
CELESTISSIMA (Alexander), 1945f:17, fig. 9 (wing) (*Austrolimnophila*). Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile. Ref. — Alexander, 1958:195, fig. 5 (male hypopygium).

Subgenus LIMNOPHILA Macquart

The numerous members of the genus in this faunal area do not belong to the restricted subgenus and at the present time are not assigned to any subgeneric group. The species are recorded in species groups that may be found to represent valid subgenera.

Group I

The species show certain features of *Shannonomyia* and may more properly be placed therein.

ARAUCANIA ARAUCANIA Alexander, 1928b:49. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.

ssp. MACRACANTHA Alexander, 1952b:107. Type-locality: Chile, Malleco, Curacautin. Distr. — Chile.

CANCELATA Alexander, 1962c:126, fig. 101 (wing). Type-locality: Bolivia, Yungas de Corani. Distr. — Bolivia.

DICTYOPTERA Alexander, 1922a:5. Type-locality: Costa Rica, San José. Distr. — Costa Rica, Panama, Venezuela.

EXPRESSA Alexander, 1937d:605. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

GUTTULATISSIMA Alexander, 1913b:546. Type-locality: Guatemala, Totonicapán. Distr. — Guatemala, Costa Rica, Venezuela.

HOFFMANNIANA Alexander, 1937f:223. Type-locality: Brazil, Santa Catarina, Ibirama, Nova Bremen. Distr. — Brazil.

KERTESZI Alexander, 1914b:212, pl. 9, fig. 10 (wing). Type-locality: Brazil, São Paulo. Distr. — Brazil.

LEUCOSTIGMA Alexander, 1937d:606. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

LITIGIOSA Alexander, 1928f:639. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.

LLOYDI Alexander, 1913c:205, pl. 5, fig. 9 (wing). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.

MADIDA Alexander, 1928a:113. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.

NEMORIVAGA Alexander, 1929a:139, figs. 54, 55 (wing), 210 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile. Ref. — Alexander, 1953b:19, fig. 17 (male hypopygium).

PROCELLA Alexander, 1944b:169. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.



- PULLIPES Alexander, 1937f:225. Type-locality: Brazil, Santa Catarina, Ibirama, Nova Bremen. Distr. — Brazil.
- RORAIMA Alexander, 1931a:13, fig. 7 (wing). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- RORAIMICOLA Alexander, 1931a:13, fig. 8 (wing). Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- RUBECULA Alexander, 1944b:168. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- SCHIADEI Alexander, 1926d:385. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- THERESIAE Alexander, 1945e:237. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Serra dos Órgãos. Distr. — Brazil.
- UNDULATA Bellardi, 1861:1. Type-locality: "Mexico". Distr. — Mexico.

Group II

- ABSTRUSA Alexander, 1929a:140, figs. 56 (wing), 211 (male hypopygium). Type-locality: Argentina, Rio Negro, Rio Pichileufu. Distr. — Chile, Argentina.
- ANGUSTILINEATA Alexander, 1926d:386. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- ARMIGERA Alexander, 1945f:20, fig. 10 (wing). Type-locality: Chile, Malleco. Angol. Distr. — Chile.
- COLOPHALLUS Alexander, 1966c:466, figs. 2 (wing), 8 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Caramavida. Distr. — Chile.
- EUTHETA Alexander, 1935b:102, fig. 4 (wing). Type-locality: Chile, Malleco. Angol. Distr. — Chile.
- HUMIDICOLA Alexander, 1929a:138, figs. 72 (wing), 209 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile, Argentina.
- INCULTA Alexander, 1929a:137, figs. 71 (wing), 208 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile, Argentina.
- JORDANICA Alexander, 1949b:477, fig. 10 (wing). Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- KAJETURANA Alexander, 1930c:733. Type-locality: Guyana, Kaieteur Falls. Distr. — Guyana, Venezuela.
- MELICA Alexander, 1929a:141, fig. 57 (wing). Type-locality: Chile, Valparaíso, Marga-Marga. Distr. — Chile, Argentina.
- OPTICAI Alexander, 1948c:517. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- PENANA Alexander, 1966c:470, figs. 4 (wing), 9 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Pillim Pilli. Distr. — Chile.
- SPINULOSA Alexander, 1945n:411. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

Group III

- FILIFORMIS Alexander, 1929a:95. Type-locality: Chile, Chiloé Is., Butaleura. Distr. — Chile.
- NEMATOCERA Alexander, 1939b:201. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PERGRACILIS Alexander, 1943b:120. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Cunibunda. Distr. — Ecuador.



Group IV

CHARON Alexander, 1937d:604. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Group V

CTENONYCHA Alexander, 1929a:136, figs. 70 (wing), 207 (male hypopygium). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile, Argentina. Ref. — Alexander, 1966c:468, fig. 3 (wing).

Genus MESOLIMNOPHILA Alexander

Polymoria Philippi, 1865:608 (preocc. Förster, 1856). Type-species, *lutea* Philippi (Alexander, 1927b:15).

MESOLIMNOPHILA Alexander, 1929a:129 (nom. nov. for *Polymoria* Philippi). Type-species, *Polymoria lutea* Philippi (aut.).

Polypraesidia Miller, 1943:72 (unnecessary nom. nov. for *Polymoria* Philippi). Type-species, *Polymoria lutea* Philippi (aut.). Ref. — Alexander, 1948c:512, fig.

HIRSUTIPES Alexander, 1929a:131, figs. 67 (wing), 204 (male hypopygium). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile.

LUTEA (Philippi), 1865:608, fig. (*Polymoria*). Type-locality: Chile, Valdivia. Distr. — Chile. Refs. — Alexander, 1929a:131; 1948c:517, fig. 5 (wing).

Genus PARADELPHOMYIA Alexander

Adelphomyia, subg. PARADELPHOMYIA Alexander, 1936f:184. Type-species, *crossospila* Alexander (orig. des.).

Subgenus OXYRHIZA Meijere

Oxydiscus Meijere, 1913:350 (preocc. Koken, 1889). Type-species, *nebulosus* Meijere (mon.).

OXYRHIZA Meijere, 1946:68 (as genus; nom. nov. for *Oxydiscus* Meijere). Type-species, *Oxydiscus nebulosus* Meijere (aut.).

AEQUATORIALIS (Alexander), 1944k:605 [*Oxydiscus (Oxydiscus)*]. Type-locality: Ecuador, Santiago-Zamora, Zamora. Distr. — Ecuador.

COSTARICENSIS (Alexander), 1922a:6 (*Adelphomyia*). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.

DESTITUTA (Alexander), 1945b:56 [*Oxydiscus (Oxydiscus)*]. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.

MEXICANA MEXICANA (Alexander), 1940b:158 [*Oxydiscus (Oxydiscus)*]. Type-locality: Mexico, Chiapas, Finca Germania. Distr. — Mexico

ssp. ACUTISSIMA (Alexander), 1940b:159 [*Oxydiscus (Oxydiscus)*]. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.

MORELOSENSIS (Alexander), 1946c:531 [*Oxydiscus (Oxydiscus)*]. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.

OAXACENSIS (Alexander), 1946a:124 [*Oxydiscus (Oxydiscus)*]. Type-locality: Mexico, Oaxaca, Sierra de Juárez, Yotao. Distr. — Mexico. Ref. — Alexander, 1948a:153, fig. 2 (wing).

VENEZUOLANA Alexander, 1950a:46, figs. 14 (wing), 18 (male hypopygium).
Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.

Genus PARALIMNOPHILA Alexander

- Limnophila*, subg. PARALIMNOPHILA Alexander, 1921g:559. Type-species, *Limnophila leucophaeta* Skuse (orig. des.) (Australasian).
- CONSPERSA (Enderlein), 1912:49, fig. D1 (wing) (*Dactylobasis*). Type-locality: Brazil, Santa Catarina. Distr. — Brazil.
- DIFFUSIOR Alexander, 1968b:87. Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- INFESTIVA (Alexander), 1929a:155, fig. 79 (wing) [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Llanquihue, Ensenada. Distr. — Chile.
- IRRORATA (Philippi), 1865:608 (*Polymoria*). Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Alexander, 1929a:157, fig. 81 (wing).
- PACHYSPILA (Alexander), 1928f:639 [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Malleco. Distr. — Chile.
- PALLITARSIS (Alexander), 1929a:153, fig. 76 (wing) [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile, Argentina.
- PERIRRORATA (Alexander), 1928d:218 [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.
- PERREDUCTA (Alexander), 1938i:278 [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- PIRIONI (Alexander), 1928b:53 [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Valparaiso, Perales de Marga-Marga. Distr. — Chile. Ref. — Alexander, 1948c:527, fig. 15 (wing).
- PLATENSIS (Alexander), 1923b:184 (*Limnophila*). Type-locality: Argentina, La Plata, Palo Blanco. Distr. — Argentina.
- RARA (Alexander), 1929a:156, fig. 80 (wing) [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- STYGHES (Alexander), 1929a:154 [*Gynoplistia (Paralimnophila)*]. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- SUBFUSCATA (Alexander), 1921b:177 (*Limnophila*). Type-locality: Argentina, Buenos Aires. Distr. — Argentina.
- TORTILIS Alexander, 1968b:88. Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.

Genus PILARIA Sintenis

- PILARIA Sintenis, 1888:398. Type-species, *Limnobia pilicornis* Zetterstedt (Coquillett, 1910:591). Ref. — Alexander, 1948c:523, fig.
- Limnophila*, subg. *Eulimnophila* Alexander, 1919b:917. Type-species, *Limnobia tenuipes* Say (Ishida, 1959:8).
- RUBELLA Alexander, 1926a:169, fig. 7 (wing). Type-locality: Mexico, Veracruz, Córdoba, Hacienda San Francisco. Distr. — Mexico. Ref. — Alexander, 1948c:524, fig. 10 (wing).
- TENUIPES (Say), 1823:21 (1859:46) (*Limnobia*). Type-locality: U.S.A., Pennsylvania. Distr. — Eastern U.S.A., south to northern Mexico.



Genus POLYMERA Wiedemann

POLYMERA Wiedemann, 1820:40 (1821:40). Type-species, *Chironomus hirticornis* Fabricius (mon.). Refs. — Alexander, 1948a:182, figs.; Rogers, 1933:1-14, 2 pls. (biology).

Subgenus POLYMERA Wiedemann

- ALBIDITARSIS Alexander, 1953b:16, fig. 10 (wing). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- ALBITARSIS ALBITARSIS Williston, 1896a:296, pl. x, figs. 71, 71A (antenna), 71B (wing). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent.
- ssp. DOMINICAE Alexander, 1939d:98. Type-locality: Lesser Antilles: Dominica, Sylvania Estate. Distr. — Dominica.
- ALBOGENICULATA Alexander, 1926d:383. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- ALBOGENUALIS Alexander, 1939b:205. Type-locality: Ecuador, Tungurahua, Río Blanco, near Baños. Distr. — Ecuador.
- ANTICALBA Alexander, 1939c:436, fig. 5 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ARAWAK Alexander, 1964a:53, fig. 55 (wing). Type-locality: Jamaica, St. Andrew, Hermitage Reservoir. Distr. — Jamaica.
- BRACHYNEURA Alexander, 1962c:125, fig. 100 (wing). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- BRUCHI Alexander, 1926d:384. Type-locality Argentina, La Plata, Palo Blanco. Distr. — Argentina.
- CAVERNICOLA Alexander, 1964a:54, fig. 56 (wing). Type-locality: Jamaica, St. Catherine, St. Clair Cave. Distr. — Jamaica.
- CHIRIQUENSIS Alexander, 1941c:334. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- CINEREIPENNIS Alexander, 1926d:382. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- CLAUSA Alexander, 1939b:204. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CRYSTALLOPTERA Alexander, 1921d:67. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- FUSCA Wiedemann, 1828:58. Type-locality: "Brazil". Distr. — Brazil. Ref. — Alexander, 1948a:187, fig. 29 (wing).
- FUSCITARSIS Alexander, 1937b:240. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- GENICULATA GENICULATA Alexander, 1915c:106. Type-locality: Puerto Rico, Carolina. Distr. — Puerto Rico. Ref. — Alexander, 1932:367, fig. 10 (wing).
- ssp. PALLIPES Alexander, 1964a:55, fig. 57 (wing). Type-locality: Jamaica, Trelawny, Windsor Cave. Distr. — Jamaica.
- GRISEA Alexander, 1913b:535, fig. 6 (wing). Type-locality: Panama, Canal Zone, Ancon. Distr. — Panama.
- HIRTICORNIS (Fabricius), 1805:39 (*Chironomus*). Type-locality: "South America." Distr. — Guyana, Brazil. Ref. — Alexander, 1913b:532, fig. 3 (wing).
- HONESTA Alexander, 1940b:156. Type-locality: Mexico, Chiapas. Distr. — Mexico.



- INORNATA Alexander, 1913b:534, fig. 5 (wing). Type-locality: Guyana, Tukeit. Distr. — Guyana.
- LEUCOPEZA Alexander, 1940b:155. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- MELANOSTERNA Alexander, 1945c:236. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Serra dos Órgãos. Distr. — Brazil.
- MICROSTICTULA Alexander, 1929e:404. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Peru, Brazil, Paraguay.
- MINUTOR Alexander, 1942e:253. Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- MONOSTICTA Alexander, 1948a:187. Type-locality: Peru, Huanuco. Distr. — Peru.
- NEOCLAUSA Alexander, 1967c:373. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- NIMBIPENNIS Alexander, 1945n:409. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- NIVEITARSIS Alexander, 1913b:532, fig. 16 (wing). Type-locality: Panama, Bocas del Toro. Distr. — Guatemala, Panama, Venezuela, Guyana, Surinam, Ecuador, Peru, Brazil. Ref. — Alexander, 1948a:185, fig. 31 (male antenna).
- NODULIFERA Alexander, 1940b:157. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- OBSCURA Macquart, 1838a:69(65). Type-locality: "Brazil". Distr. — Mexico, Guyana, Peru, Brazil, Bolivia, Argentina. Refs. — Alexander, 1913b:529, fig. 18 (wing); 1948a:185, fig. 32 (male antenna).
- OMINOSA Alexander, 1938d:333. Type-locality: Colombia, Cundinamarca, Fusagasugá. Distr. — Colombia.
- PLEURALIS Alexander, 1913b:528, fig. 17 (wing). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- PROLIXICORNIS Alexander, 1927c:269. Type-locality: El Salvador, Agronomla, Sonsonata. Distr. — Mexico, El Salvador.
- PULCHRICORNIS Alexander, 1914e:251, pl. 4, fig. 10 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana.
- REGINA Alexander, 1926d:381. Type-locality: Paraguay, Santa Bárbara. Distr. — Brazil, Paraguay.
- SCELEROSA Alexander, 1948a:188. Type-locality: Bolivia, Santa Cruz, Buenavista. Distr. — Bolivia.
- SORDIDIPES Alexander, 1938d:332. Type-locality: Colombia, Cundinamarca, Fusagasugá. Distr. — Colombia.
- STENOPTERA Alexander, 1949c:589. Type-locality: Peru, Loreto, Pucallpa. Distr. — Peru.
- SUBSUPERBA Alexander, 1926d:382. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Venezuela, Bolivia, Paraguay.
- SUPERBA SUPERBA Alexander, 1913b:530, fig. 2 (wing). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Panama, Ecuador, Brazil.
- ssp. DISCALIS Alexander, 1929e:405. Type-locality: Paraguay, Caroveni. Distr. — Brazil, Paraguay.
- THORACICA Alexander, 1913b:533, fig. 4 (wing). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- TIBIALIS Alexander, 1922c:50. Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- UNPUNCTATA Alexander, 1921b:179. Type-locality: Argentina, Jujuy, Jujuy. Distr. — Argentina.

VERTICILLATA Alexander, 1948a:189, 30 (wing). Type-locality. Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil.

Subgenus POLYMERODES Alexander

- Polymera*, subg. POLYMERODES Alexander, 1920c:143. Type-species, *parishi* Alexander (orig. des.). Ref. — Alexander, 1948a:185, fig. 27 (wing).
- CATHARINAE Alexander, 1931a:111. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- CONJUNCTA Alexander, 1913b:529, fig. 7 (wing). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Guyana, Brazil.
- CONJUNCTOIDES Alexander, 1920a:74. Type-locality: Brazil, Amazonas, Itacoatiara. Distr. — Venezuela, Ecuador, Peru, Brazil.
- EVANESCENS Alexander, 1948a:185, fig. 28 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LEUCOSTROPHA Alexander, 1966a:222. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- MINUTISSIMA Alexander, 1945a:25. Type-locality: Ecuador, Napo-Pastaza, Río Arajuno, Napo Watershed. Distr. — Ecuador.
- PARISHI Alexander, 1920c:143. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil, Paraguay.
- TASIOCEROIDES Alexander, 1948a:186. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus PSEUDOLIMNOPHILA Alexander


- Limnophila*, subg. PSEUDOLIMNOPHILA Alexander, 1919b:917 (in key). Type-species, *luteipennis* Osten Sacken (Alexander, 1920i:849). Ref. — Alexander, 1948a:181.
- LUTEIPENNIS (Osten Sacken), 1859:236 (*Limnophila*). Type-locality: U.S.A., Washington, D. C. Distr. — Nearctic region south to Mexico and Central America.
- MEGALOPS Alexander, 1942i:62, figs. 27 (wing), 34 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil. Ref. — Alexander, 1948a:179, fig. 25 (wing).
- PLUTO Alexander, 1939b:202. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PLUTOIDES Alexander, 1948b:138. Type-locality: Peru, Huanuco, Carpiash. Distr. — Peru. Ref. — Alexander, 1948a:179, fig. 26 (wing).
- SUPPLEMENTA Alexander, 1945j:22, fig. 5 (wing). Type-locality: Brazil, São Paulo, Ribeirão Pires, Barreira das Camélias. Distr. — Brazil.

Genus SHANNONOMYIA Alexander

- SHANNONOMYIA Alexander, 1929a:142. Type-species, *Limnophila lenta* Osten Sacken (orig. des.) (Nearctic). Ref. — Alexander, 1948c:518, figs. (general discussion).

Subgenus RORAIMOMYIA Alexander

- Limnophila*, subg. RORAIMOMYIA Alexander, 1935a:323. Type-species, *Limnophila permonstrata* Alexander (orig. des.).



PERMONSTRATA (Alexander), 1935a:323 [*Limnophila (Roraimomyia)*]. Type-locality: Venezuela, Summit of Mount Roraima. Distr. — Venezuela. Ref. — Alexander, 1948c:516.

Subgenus SHANNONOMYIA Alexander

- ABORTIVA Alexander, 1966c:471. Type-locality: Argentina, Santa Cruz, Lago Argentino, Calafate. Distr. — Argentina.
- ABRA Alexander, 1949c:590. Type-locality: Peru, Junín, Chanehamayo. Distr. — Peru.
- ADUMBRATA Alexander, 1946b:239, figs. 18 (wing), 25 (male antenna). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- AENIGMATICA Alexander, 1929a:144, fig. 73 (wing). Type-locality: Chile, Chiloe, Mechuque Is. Distr. — Chile.
- ANTARCTICA (Walker), 1837:332 (*Gonomyia?*). Type-locality: Chile, Magalanes, Port Famine. Distr. — Chile, Argentina, Straits of Magellan. Ref. — Alexander, 1929a:143, fig. 58 (wing).
- ARAGUAE Alexander, 1947d:48, figs. 3 (wing), 4 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- ARGENTICEPS Alexander, 1929d:780. Type-locality: Ecuador, Ventura. Distr. — Ecuador.
- ATROAPICALIS Alexander, 1934a:65. Type-locality: Panama, Chiriqui, Finca Parada. Distr. — Panama.
- AUSTROLATHRAEA Alexander, 1930e:733. Type-locality: Bolivia, Santa Cruz. Distr. — Ecuador, Bolivia.
- BARILOCHENSIS Alexander, 1929a:147, fig. 59 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Chile, Argentina.
- BATESI Alexander, 1939d:119, fig. 4 (wing). Type-locality: Hispaniola: Haiti, La Selle Range, La Visite. Distr. — Haiti.
- BOGOTENSIS Alexander, 1938d:331. Type-locality: Colombia, Usme. Distr. — Colombia.
- BREVICULA Alexander, 1931a:113. Type-locality: Cuba, Buenos Aires, Trinidad Mts., Eiba. Distr. — Cuba.
- BREVINERVIS Alexander, 1929a:146, fig. 74 (wing). Type-locality: Chile, Chiloé Is., Aneud. Distr. — Chile.
- BRUNERIANA BRUNERIANA Alexander, 1937e:528, fig. 4 (wing). Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
ssp. FORTICORNIS Alexander, 1937e:528. Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- BULLOCKIANA (Alexander), 1939f:178 (*Limnophila*). Type-locality: Chile, Malleco, Curacautín, Termas Río Blanco. Distr. — Chile.
- CACOXENA CACOXENA Alexander, 1929a:148, fig. 60 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
ssp. MENDICA Alexander, 1929a:148. Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
- CAESIA Alexander, 1937c:492. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CERBEREANA Alexander, 1945c:598. Type-locality: Ecuador, Tungurahua, Río Blanco near Baños. Distr. — Ecuador.
- CINGARA Alexander, 1948c:520, figs. 8 (wing), 11 (male hypopygium). Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru.



- CRASSICORNIS Alexander, 1964a:57, figs. 58 (wing), 66 (male hypopygium).
Type-locality: Jamaica, St. Andrew, Cinchona. Distr. — Jamaica.
- DAMPMI Alexander, 1939a:76. Type-locality: Mexico, Chiapas, Finca Germania. Distr. — Mexico.
- ERUBESCENS Alexander, 1944m:310. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- FERIATA FERIATA (Alexander), 1929a:135, fig. 69 (wing) (*Limnophila*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- ssp. PAMPHAEA (Alexander), 1929a:136 (*Limnophila*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- FUSCOSTIGMALIS (Alexander), 1966c:469 (*Limnophila*). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- GLOBULICORNIS Alexander, 1968a:25, fig. 5 (antenna), 9 (male hypopygium).
Type-locality: Chile, Las Brizas. Distr. — Chile.
- GRACILIOR Alexander, 1944m:311. Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- HAITENSIS Alexander, 1939d:120, fig. 5 (wing). Type-locality: Hispaniola: Haiti, La Selle Range, La Visite. Distr. — Haiti.
- HALTERATA Alexander, 1966c:472, fig. 10 (male hypopygium). Type-locality: Chile, Tierra del Fuego, Port Williams, Navarino Is., Beagle Canal. Distr. — Chile.
- HOFFMANI Alexander, 1936b:877, fig. 1 (wing). Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- IGNAVA Alexander, 1943g:533. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- JAFFUELI Alexander, 1928d:217. Type-locality: Chile, Valparaiso, Margamarga. Distr. — Chile.
- JUSTA Alexander, 1937c:493. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- KUSCHELI Alexander, 1952c:57, fig. 26 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Inocentes Bajos. Distr. — Chile.
- LATHRAEA (Alexander), 1926d:386 (*Pilaria*). Type-locality: Colombia, San Lorenzo Mt., Cincinnati. Distr. — Colombia, Venezuela.
- LENITATIS Alexander, 1946a:125. Type-locality: Mexico, Chiapas, Sierra Madre. Distr. — Mexico.
- LENTINA Alexander, 1928a:112. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- LENTOIDES (Alexander), 1913b:545, fig. 21 (wing) (*Limnophila*). Type-locality: Guatemala, Totonicapán. Distr. — Guatemala.
- LEONARDI Alexander, 1932:368, fig. 11 (wing). Type-locality: Puerto Rico, Luquillo National Forest, El Yunque. Distr. — Puerto Rico.
- LONGIRADIALIS Alexander, 1929b:96. Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile. Ref. — Alexander, 1948c:520, fig. 12 (male hypopygium).
- MASATIERRAE Alexander, 1952c:58. Type-locality: Chile, Juan Fernández, Miradero de Selkirk. Distr. — Chile.
- MESOPHRAGMA Alexander, 1928b:52. Type-locality: Cuba, Jarahuca. Distr. — Cuba.
- MESOPHRAGMOIDES Alexander, 1937e:527, fig. 3 (wing). Type-locality: Cuba, Oriente, Sierra Macstra, Pico Turquino. Distr. — Cuba.
- MINUTIPENNIS Alexander, 1929a:145. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.



- MOCTEZUMA Alexander, 1928a:111. Type-locality: Mexico, Mexico, D.F., Desierto de los Leoncs. Distr. — Mexico.
- MYERSIANA Alexander, 1931a:112. Type-locality: Jamaica, Blue Mts. Distr. — Jamaica. Ref. — Alexander, 1964a:58, fig. 59 (wing).
- NACREA (Alexander), 1913b:544, fig. 19 (wing) (*Limnophila*). Type-locality: Jamaica, St. Andrew, Cinchona. Distr. — Jamaica. Ref. — Alexander, 1964a:58.
- NUDIPENNIS Alexander, 1964a:59, fig. 60 (wing). Type-locality: Jamaica, St. Andrew, Newcastle. Distr. — Jamaica.
- OLSSONI (Alexander), 1919a:144, fig. 12 (wing) (*Gnophomyia*). Type-locality: Panama, Bocas del Toro, Río Cana. Distr. — Panama.
- OROPHILA (Alexander), 1913c:205 (*Limnophila*). Type-locality: Colombia, Cauca, Almaguer. Distr. — Colombia.
- OVALIFORMIS Alexander, 1946b:240, figs. 19 (wing), 26 (antenna). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- PARAGUAYENSIS Alexander, 1929d:780. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- PARVICELLULA Alexander, 1968c:214. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico.
- PENUMBROSA Alexander, 1939f:180. Type-locality: Chile, Malleco, Curacautín, Termas Río Blanco. Distr. — Chile.
- PHAEOSTIGMOSA Alexander, 1943g:532. Type-locality: Peru, Ayacucho, La Mar, Yanamontc. Distr. — Peru.
- PHRAGMOPHORA Alexander, 1937e:525, fig. 2 (wing). Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- POMERANTZI Alexander, 1964a:61, figs. 61 (wing), 67 (male hypopygium). Type-locality: Jamaica, St. Andrew, Cinchona. Distr. — Jamaica.
- PROTUBERANS Alexander, 1946a:124. Type-locality: Mexico, Morelos, Lagunas de Zempole. Distr. — Mexico.
- PROVIDENS Alexander, 1950a:47, figs. 19 (wing), 25 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- RORAIMENSIS Alexander, 1935a:324. Type-locality: Venezuela, Mount Roraima. Distr. — Venezuela, Brazil.
- SCARAMUZZAI Alexander, 1937a:183, fig. 2 (wing). Type-locality: Cuba, Oriente, Sierra del Cobre, Loma del Gato. Distr. — Cuba.
- SECLUSA (Alexander), 1928f:638 (*Limnophila*). Type-locality: Chile, Malleco, Pailahueque. Distr. — Chile.
- SELKIRKIANA Alexander, 1952c:58, figs. 22 (wing), 27 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Bahía Cumberland. Distr. — Chile.
- SEPTEMPUNCTATA Alexander, 1939d:121, fig. 6 (wing). Type-locality: Hispaniola: Dominican Republic, Valle Nuevo, southeast of Constanza. Distr. — Dominican Republic.
- SOPORA Alexander, 1944b:171. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- SPARSIPUNCTATA Alexander, 1940d:221. Type-locality: Ecuador, Napo-Pastaza, Río Pastaza, San Francisco. Distr. — Ecuador.
- SPARSISSIMA (Alexander), 1929a:134, fig. 68 (wing) (*Limnophila*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- STIGMATICA (Philippi), 1865:610 (*Limnophila*). Type-locality: Chile, Valdivia. Distr. — Chile.

- TRIANGULARIS (Alexander), 1927c:270 (*Pilaria*). Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico. Ref. — Alexander, 1932:369.
- TUBER (Alexander), 1953b:17, figs. 11 (male hypopygium), 14 (wing) (*Limnophila*). Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- UMBRA Alexander, 1948c:522, figs. 9 (wing), 13 (male hypopygium). Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.
- VOCATOR Alexander, 1944k:604. Type-locality: Ecuador, Bolívar, Hacienda Talahua. Distr. — Ecuador.
- ZERNYANA Alexander, 1949b:479, figs. 11 (wing), 14 (male hypopygium). Type-locality: Brazil, São Paulo, Alto da Serra (= Paranapiacaba). Distr. — Brazil.

Unplaced Hexatomini

- CHILENSIS Blanchard, 1852:339 (*Limnophila*). Type-locality: Chile, Coquimbo, La Serena.
- CINERACEA Philippi, 1865:611 (*Limnophila*). Type-locality: Chile, Santiago. Ref. — Alexander, 1913b:546.
- CINEREA Philippi, 1865:609 (*Polymoria*). Type-locality: Chile, Valdivia.
- DECASEILA Wiedemann, 1828:549 (*Limnobia*). Type-locality: "Brazil".
- FLAVICAUDA Bigot, 1888:10, pl. 2, fig. 4 (wing) (*Poecilostola*). Type-locality: Chile, Tierra del Fuego.
- INSULARIS Johnson, 1913:443, fig. 2 (*Limnophila*). Type-locality: "Bermuda". (Perhaps in *Pilaria*, as suggested by Johnson).
- PALLENS Philippi, 1865:610 (*Limnophila*). Type-locality: Chile, Valdivia. Identity uncertain; may belong to the present genus or possibly to *Aphrophila* Edwards.
- TENELLA Philippi, 1865:609 (*Polymoria*). Type-locality: Chile, Valdivia.
- VENOSA Philippi, 1865:611 (*Limnophila*). Type-locality: unknown (Chile).
- VERECUNDA Philippi, 1865:611 (*Limnophila*). Type-locality: unknown (Chile).

Tribe ERIOPTERINI

Genus AMPHINEURUS Skuse


- AMPHINEURUS Skuse, 1890a:800. Type-species, *umbraticus* Skuse (Alexander, 1920g:52) (Australasian). Ref. — Alexander, 1947c:337.

Subgenus AMPHINEURUS Skuse

- CASTROENSIS Alexander, 1929a:190, fig. 105 (wing). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- EXTRAORDINARIUS Alexander, 1938i:280. Type-locality: Chile, Pinares de Lonquimai, Villa Portales. Distr. — Chile. Ref. — Alexander, 1947c:329, fig. 18 (wing).
- LONGIPES (Philippi), 1865:616 (*Erioptera?*). Type-locality: Chile, Valdivia. Distr. — Chile. Identity uncertain.

Subgenus RHAMPHONEURUS Alexander

- Amphineurus*, subg. RHAMPHONEURUS Alexander, 1929a:186. Type-species, *nothofagetorum* Alexander (orig. des.).



- APICULATUS Alexander, 1968a:29, fig. 15 (male hypopygium). Type-locality: Chile, Chaihuin. Distr. — Chile.
- FUSCIFUSUS Alexander, 1929a:190. Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- GLABRISTYLATUS Alexander. 1929a:188, fig. 224 (male hypopygium). Type-locality: Chile, Mechuque Is. Distr. — Chile, Argentina.
- INSANUS Alexander, 1952b:111, fig. 9 (male hypopygium). Type-locality: Chile, Malleco, Río Blanco. Distr. — Chile.
- NONNULUS Alexander, 1966c:491. Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile. Ref. — Alexander, 1968a, fig. 17 (male hypopygium).
- NOTHOFAGETORUM Alexander, 1929a:187, figs. 90 (wing), 223 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina.
- NULLUS Alexander, 1966c:492. Type-locality: Chile, Chiloé Is., Aulendukateo. Distr. — Chile. Ref. — Alexander, 1968a, figs. 11 (wing), 16 (male hypopygium).
- RUTRISTYLUS Alexander, 1968a:30, fig. 18 (male hypopygium). Type-locality: Chile, Valdivia, Enco. Distr. — Chile.
- SANUS Alexander, 1929a:189, fig. 225 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.

Genus APHROPHILA Edwards

- Gnophomyia*, subg. APHROPHILA Edwards, 1923:297. Type-species, *neozelandica* Edwards (orig. des.) (Australasian: New Zealand). Ref. — Alexander, 1947b:99.
- ANTENNATA Alexander, 1953g:16, figs. 3 (wing), 11 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- AURANTIACA Alexander, 1944i:131, fig. 8 (ovipositor). Type-locality: Chile, Aysén. Distr. — Chile.
- BIDENTATA Alexander, 1968a:27, figs. 10 (wing), 14 (male hypopygium). Type-locality: Argentina, Neuquen, Lago Nonthué. Distr. — Argentina.
- CARBONARIA Alexander, 1931a:115. Type-locality: Chile, Malleco, Angol. Distr. — Chile. Ref. — Alexander, 1947b:99, fig. 30 (wing).
- CHILENA Alexander. 1928f:640. Type-locality: Chile, Bio-Bío. Distr. — Chile. Ref. — Alexander, 1944i:135, fig. 11 (ovipositor).
- CORONATA Alexander, 1944i:132, fig. 7 (male hypopygium). Type-locality: Chile, Aysén. Distr. — Chile.
- MULTIDENTATA Alexander, 1931a:116. Type-locality: Chile, Malleco, Angol. Distr. — Chile. Ref. — Alexander, 1944i:134, fig. 10 (ovipositor).
- SUBTERMINALIS Alexander, 1966c:489, figs. 12 (wing), 16 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Chacay. Distr. — Chile.
- VRIDINERVIS Alexander, 1934d:177, fig. 42 (wing). Type-locality: "Chile". Distr. — Chile. Refs. — Alexander, 1944i:134, fig. 9 (ovipositor); 1953g:17, fig. 12 (male hypopygium).

Genus AYMARAMYIA Alexander

- AYMARAMYIA Alexander, 1943a:236. Type-species, *dubia* Alexander (orig. des.). Ref. — Alexander, 1947b:80, fig. 13 (wing).



DUBIA Alexander, 1943a:237, figs. 3 (wing), 4 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru. Ref. — Alexander, 1947b:81, fig. 13 (wing).

Genus BAEOURA Alexander

Erioptera, subg. BAEOURA Alexander, 1924d:67. Type-species, *nigrolatera* Alexander (orig. des.) (Ethiopian). Refs. Alexander, 1929a:191; 1947c:322.

ADVENA (Alexander), 1928c:180 [*Cryptolabis (Baeoura)*]. Type-locality: Chile, Valparaiso, Perales de Marga-Marga. Distr. — Chilc. Ref. — Alexander, 1947c:323, figs. 3 (wing), 7 (male hypopygium).

Genus CHEILOTRICHIA Rossi

CHEILOTRICHIA Rossi. 1848:12. Type-species, *Erioptera imbuta* Meigen (mon.) (Palaeartic).

Subgenus EMPEDA Osten Sacken

EMPEDA Osten Sacken, 1869:153 (as genus). Type-species, *stigmaticea* Osten Sacken (Coquillett, 1910:537) (Nearctic). Ref. — Alexander, 1947c:335.

ABITAGUAI (Alexander), 1941d:131, fig. 15 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Venezuela. Ecuador.

ALTICOLA (Alexander), 1925a:353, fig. 22 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico.

AUSTRONYMPHICA (Alexander), 1943a:241 [*Erioptera (Empeda)*]. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru, Bolivia. Ref. — Alexander, 1947c:329, fig. 17 (wing).

BASALIS (Alexander), 1921e:293 [*Rhodomastix (Sacandaga)*]. Type-locality: Argentina, Catamarca, Punta de Balasto. Distr. — Argentina.

BOLIVIANA (Alexander), 1930c:739 [*Erioptera (Empeda)*]. Type-locality: Bolivia, Santa Cruz. Distr. — Bolivia.

COANGUSTATA (Alexander), 1944k:615 [*Erioptera (Empeda)*]. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.

COMPLICATA (Alexander), 1922b:209 [*Rhodomastix (Sacandaga)*]. Type-locality: Argentina, Catamarca, Masao. Distr. — Argentina.

CURTA (Alexander), 1925a:354, figs. 11 (wing), 23 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Chapingo. Distr. — Mexico.

DELUDENS (Alexander), 1939a:87 [*Erioptera (Empeda)*]. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.

DESTITUTA (Alexander), 1942c:443 [*Erioptera (Empeda)*]. Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.

DIVARICATA (Alexander), 1939a:86 [*Erioptera (Empeda)*]. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.

INSTRENUA (Alexander), 1942c:444, fig. 12 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.

LONGIFURCATA (Alexander), 1940a:293 [*Erioptera (Empeda)*]. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.

- LONGISQUAMA (Alexander), 1937f:226 [*Erioptera (Empeda)*]. Type-locality: Brazil, Santa Catarina, Ibirama, Nova Bremen. Distr. — Brazil.
- MAYANYMPHICA (Alexander), 1946b:243, fig. 24 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Chiapas, Finca Germania. Distr. — Mexico.
- NIGROLINEATA (Enderlein), 1912:56, fig. H1 (venation) (*Empeda*). Type-locality: "Costa Rica". Distr. — Mexico, Guatemala, Costa Rica. Ref. — Alexander, 1913b:520.
- NYMPHICA (Alexander), 1928g:24 [*Erioptera (Empeda)*]. Type-locality: Jamaica, Cinchona. Distr. — Jamaica. Ref. — Alexander, 1964a:81, figs. 89 (wing), 93 (male hypopygium).
- OCHRICAUDA (Alexander), 1925a:352, fig. 21 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Mt. Tlaloc. Distr. — Mexico. Ref. — Alexander, 1947c:329, fig. 16 (wing).
- ORESITROPHA (Alexander), 1927d:310 [*Erioptera (Empeda)*]. Type-locality: Mexico, north of San Cristóbal, summit of the Zontehuitz. Distr. — Mexico.
- PERCUPIDA (Alexander), 1941e:255 [*Erioptera (Empeda)*]. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- PUBESCENS (Alexander), 1913b:520 [*Erioptera (Empeda)*; as *nigrolineata* var.] Type-locality: Guatemala, Totonicapán. Distr. — Guatemala.
- STYGIA (Alexander), 1927d:309 [*Erioptera (Empeda)*]. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico, Costa Rica, Peru.
- SUTRINA (Alexander), 1941g:354 (also 1946h:102) [*Erioptera (Empeda)*]. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- TRIDENTATA (Alexander), 1925a:355, fig. 24 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Mexico, D.F., Desierto de los Leones. Distr. — Mexico.
- UNIDENTATA (Alexander), 1925a:355, fig. 25 (male hypopygium) [*Erioptera (Empeda)*]. Type-locality: Mexico, Mexico, D.F., Desierto de los Leones. Distr. — Mexico.

Genus CRYPTOLABIS Osten Sacken

- CRYPTOLABIS Osten Sacken, 1859:224. Type-species, *paradoxa* Osten Sacken (mon.) (Nearctic). Ref. — Alexander, 1947b:322, figs.

Subgenus CRYPTOLABIS Osten Sacken

- ALTICOLA Alexander, 1944a:56. Type-locality: Ecuador, Bolívar, Hacienda Talahua. Distr. — Ecuador. Ref. — Alexander, 1947c:323, fig. 6 (wing).
- ATMOPHORA Alexander, 1929a:192, figs. 107 (wing), 227 (male hypopygium). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile.
- CHILOTA Alexander, 1929a:194, fig. 228 (male hypopygium). Type-locality: Chile, Mechuque Is. Distr. — Chile.
- CORTESI Alexander, 1952b:110, figs. 6 (wing), 8 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- DIVERSIPES Alexander, 1939e:631, figs. 5 (wing), 14 (male hypopygium). Type-locality: Costa Rica, Talamanca, Rivas. Distr. — Costa Rica.
- ECALCARATA Alexander, 1947c:326, figs. 5 (wing), 8 (male hypopygium). Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.



- FUSCOVENOSA Alexander, 1928a:118. Type-locality: Mexico, Sonora, Río Mayo near Navojoa. Distr. — Mexico.
- HILARIS Alexander, 1945d:278. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru. Ref. — Alexander, 1945L:379, fig. 21 (wing).
- INVARIPIES Alexander, 1939e:634, fig. 7 (wing). Type-locality: Costa Rica, Talamanca, Pedrogoso. Distr. — Costa Rica.
- JOVIALIS Alexander, 1945d:279. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- JUBILATA Alexander, 1943h:497, figs. 12 (wing), 16 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- LADDEYI Alexander, 1943h:500, figs. 14 (wing), 18 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- LATICOSTATA Alexander, 1940a:296. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- LONGIRADIALIS Alexander, 1939a:89. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- LUTEICEPS Alexander, 1927d:315. Type-locality: Mexico, Tonalá. Distr. — Mexico.
- LUTEICOSTA Alexander, 1934a:71. Type-locality: Panama, Chiriquí, Finca Parada. Distr. — Panama.
- LUTEOLA Alexander, 1946a:137. Type-locality: Mexico, Chiapas, Huehuetán. Distr. — Mexico.
- MOLOPHILOIDES Alexander, 1943i:763. Type-locality: U.S.A., Wyoming, Yellowstone National Park. Distr. — U.S.A., south to Mexico.
- MENACANTHA Alexander, 1943h:498, figs. 13 (wing), 17 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- NEBULICINCTA Alexander, 1941b:147. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- PARRAI Alexander, 1946a:138. Type-locality: Mexico, Chiapas, San Antonio Nexapa. Distr. — Mexico.
- PENDULIFERA Alexander, 1949c:613, fig. 38 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- PERDISTANS Alexander, 1967c:376. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- PHALLOSTENA Alexander, 1968b:94. Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- RECURVATA Alexander, 1942e:260. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador.
- ROUNDSI Alexander, 1939e:630, figs. 4 (wing), 13 (male hypopygium). Type-locality: Costa Rica, Talamanca, Rivas. Distr. — Costa Rica.
- SCHADEI Alexander, 1935a:330. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- SEMIFLAVA Alexander, 1949c:614, fig. 39 (male hypopygium). Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.
- SEPULCHRALIS Alexander, 1922c:40. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- SORDIPIES Alexander, 1943h:501, figs. 15 (wing), 19 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.

- SPATULATA Alexander, 1929a:193, fig. 230 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina, Chile.
- TACITURNA Alexander, 1945e:240, fig. 5 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- TENUICINCTA Alexander, 1921d:73. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- TRAVASSOSI Alexander, 1945c:275, fig. 22 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- TRIDENTICULATA Alexander, 1967c:376. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- TROPICALIS Alexander, 1913b:509, fig. 8 (wing). Type-locality: Guatemala, Alta Verapaz, Cacao Trece Aguas. Distr. — Guatemala.
- UMBROSA Alexander, 1938f:258. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- VALLICOLA Alexander, 1947c:327, fig. 9 (male genitalia). Type-locality: Panama, El Valle. Distr. — Panama.
- VARIPIES Alexander, 1939e:632, figs. 6 (wing), 15 (male hypopygium). Type-locality: Costa Rica, Talamanca, Rivas. Distr. — Costa Rica.

Subgenus PROCRYPTOLABIS Alexander

- Cryptolabis*, subg. PROCRYPTOLABIS Alexander, 1923b:183. Type-species, *argentinensis* Alexander (orig. des.).
- ARGENTINENSIS Alexander, 1923b:184. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- BARILOCHENSIS Alexander, 1929a:192, figs. 106 (wing), 226 (male hypopygium). Type-locality: Argentina, Rio Negro, Bariloche, Rio Nirico. Distr. — Argentina. Ref. — Alexander, 1947c:323, fig. 4 (wing).

Genus EMPEDOMORPHA Alexander


- EMPEDOMORPHA Alexander, 1916d:507. Type-species, *Trimicra empedoides* Alexander (orig. des.) (Nearctic).
- APACHEANA (Alexander), 1946a:135 [*Erioptera (Empedomorpha)*]. Type-locality: Mexico, Nuevo León, Granja Rodríguez. Distr. — Mexico. Refs. — Alexander, 1946e:529, fig. 26 (male hypopygium); 1947c:329, fig. 14 (wing).

Genus ERIOPTERA Meigen

- Polymeda* Meigen, 1800:14. Type-species, *Erioptera lutea* Meigen (Coquillett, 1910:593). Suppressed by I.C.Z.N., 1963:339.
- ERIOPTERA Meigen, 1803:262. Type-species, *lutea* Meigen (Coquillett, 1910:540). [See note in Alexander, 1965d:80].

Subgenus ERIOPTERA Meigen

- ANDINA Alexander, 1913c:200, pl. 5, fig. 5 (wing), pl. 6, fig. 6 (male hypopygium). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia. Ref. — Alexander, 1947c:329, fig. 10 (wing).
- ANNULIPES Williston, 1896a:294. Type-locality: Lesser Antilles: St. Vincent. Distr. — Guatemala, Colombia, Peru, Brazil, Paraguay, St. Vincent. Ref. — Alexander, 1913b:513.




- APICIALBA Alexander, 1921d:70. Type-locality: Brazil, Amazonas, Tefé. Distr. — Guyana, Brazil. Ref. — Alexander, 1934b:305.
- BEEBEANA Alexander, 1950a:54, figs. 32 (wing), 37 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grandc. Distr. — Venezuela.
- CELESTIS Alexander, 1940a:294. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Venezuela, Ecuador.
- CLADOPHORA Alexander, 1920c:142. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil, Argentina.
- CLADOPHOROIDES Alexander, 1921c:162. Type-locality: Argentina, La Plata. Distr. — Argentina. Ref. — Bruch, 1929a:20-24, pl. 6, fig. 22 (adult male), pl. 11, figs. 45-55 (immature stages).
- DAMPFI Alexander, 1927d:311. Type-locality: Mexico, near summit of the Zontehuitz. Distr. — Mexico.
- LAETIPLEURA Alexander, 1938e:406. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- LUNIGERA Alexander, 1929a:195, figs. 91 (wing), 231 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- MICROMYIA Alexander, 1920b:8. Type-locality: Brazil, Pará, Prata. Distr. — Peru, Brazil. Ref. — Alexander, 1921d:69, fig. 8 (male hypopygium).
- MULTIANNULATA Alexander, 1937c:499. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Bolivia. Ref. — Alexander, 1962c:154, fig. 135 (wing).
- POLYDONTA Alexander, 1944b:179, fig. 11 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- POLYTRICHA Alexander, 1945a:35. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- QUADRICINCTA Alexander, 1927d:310. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico, Guatemala.
- QUINQUECINCTA Alexander, 1927c:278. Type-locality: Colombia, Caldas. Distr. — Colombia.
- SUSURRA Alexander, 1945i:258. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- URANIA Alexander, 1944k:614. Type-locality: Ecuador, Santiago-Zamora. Zamora. Distr. — Ecuador, Bolivia.

Subgenus ERIOPTERELLA Alexander

- Erioptera*, subg. ERIOPTERELLA Alexander, 1929a:196. Type-species, *jaffueli* Alexander (orig. des.).
- BREVISETA (Alexander), 1968b:89 (*Neophilippiana*). Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile.
- JAFFUELI Alexander, 1929a:196, figs. 92 (wing), 232 (male hypopygium). Type-locality: Chile, Valparaíso, Marga-Marga. Distr. — Chile. Ref. — Alexander, 1947c:329, fig. 15 (wing).
- PILOSIPES (Alexander), 1929a:177, pl. 5, fig. 98 (wing) (?*Philippiana*). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.

Subgenus MESOCYPHONA Osten Sacken

- Erioptera*, subg. MESOCYPHONA Osten Sacken, 1869:152. Type-species, *caloptera* Say (Brunetti, 1912:458, as *caloptera*, error).



- AGLAIA Alexander, 1945j:30, fig. 12 (wing) [*Erioptera (Erioptera)*]. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- APICINIGRA Alexander, 1927d:313. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- BICINCTIPES Alexander, 1913b:519. Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Guyana, Peru, Brazil.
- BIEVEXA Alexander, 1967c:377. Type-locality: Nicaragua, Guapinolar. Distr. — Nicaragua.
- CALIPTERA CALIPTERA (Say), 1823:17 (1859:44). Type-locality: U.S.A., Missouri. Distr. — Antilles and southward, Bolivia, Puerto Rico; Nearctic. Refs. — Alexander, 1919b:908, fig. 77 (wing); 1932:376; 1942k:457, fig. 52 I (male hypopygium).
- ssp. FEMORANIGRA Alexander, 1913b:518. Type-locality: Costa Rica, Cartago, Juan Vinas. Distr. — Costa Rica.
- ssp. SUBEVANESCENS Alexander, 1940a:296. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- CELESTIOR Alexander, 1956b:121. Type-locality: Panama, Chiriqui, Cerro Punta. Distr. — Panama.
- COSTALIS COSTALIS Alexander, 1913b:517, fig. 24 (wing). Type-locality: Guatemala, Aguna. Distr. — Mexico, Guatemala, Cuba.
- ssp. TANTILLA Alexander. 1916d:502. Type-locality: Southeastern U.S.A. Distr. — U.S.A., Jamaica. Ref. — Alexander, 1964a:82, fig. 91 (venation).
- CYNTHIA Alexander, 1945o:608. Type-locality: Ecuador, El Oro, Morro Morro, Pinas. Distr. — Ecuador.
- DIFFUSA Alexander. 1921d:71. Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- EISENI Alexander. 1913b:516, fig. 26 (wing). Type-locality: Guatemala, Aguna. Distr. — Southwestern U.S.A. to Guatemala, Colombia, Peru.
- EUPHIROSYNE Alexander, 1945j:32, fig. 15 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- FACTIOSA Alexander, 1943i:762, fig. 13 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Ecuador, Peru.
- FUSCODISCALIS Alexander, 1938e:407. Type-locality: Mexico, Chiapas, Finca Germania. Distr. — Mexico. Ref. — Alexander, 1947c:329, fig. 2 (wing).
- GULOSA Alexander, 1943i:763, fig. 14 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- HISTRIO Alexander, 1945i:260. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Ecuador, Peru.
- IMMACULATA IMMACULATA Alexander, 1913b:518, fig. 20 (wing). Type-locality: Nicaragua, Bluefield. Distr. — Mexico, Guatemala, Nicaragua, Guyana.
- ssp. FUSCIVENA Alexander, 1923c:311. Type-locality: Argentina, Córdoba, Alta Gracia, La Granja. Distr. — Brazil, Paraguay, Argentina.
- INORNATIPES Alexander, 1925a:351, fig. 10 (wing). Type-locality: Mexico, Peñon Viejo. Distr. — Mexico.
- INTERCEPTA Alexander, 1947a:181. Type-locality: Surinam, Paramaribo. Distr. — Surinam.
- INVARIEGATA Alexander, 1921d:71. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Ecuador, Peru.

- IQUITOSENSIS* Alexander, 1945o:606. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- KNABI* Alexander, 1913b:515, fig. 25 (wing). Type-locality: Mexico, Oaxaca, Salina Cruz. Distr. — U.S.A., south to Mexico.
- LEONENSIS* Alexander, 1946a:136. Type-locality: Mexico, Neuvo León, Montemorelos. Distr. — Mexico.
- LEUCOPASTA* Alexander, 1927d:313. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico.
- MODICA* Alexander, 1927d:314. Type-locality: Mexico, Chiapas, Santa Lucia. Distr. — Mexico.
- PACHYRHAMPHA* Alexander, 1967c:378. Type-locality: Nicaragua, Villa So-moza. Distr. — Nicaragua.
- PARVA PARVA* Osten Sacken, 1859:227 (*Erioptera*). Type-locality: U.S.A., District of Columbia. Distr. — Costa Rica, Guyana, Ecuador; Nearectic.
- ssp. *BRASILIENSIS* Alexander, 1913b:514, fig. 27 (wing). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Peru, Brazil, Bolivia.
- PORTORICENSIS* Alexander, 1932:377, fig. 18 (wing). Type-locality: Puerto Rico, Luquillo National Forest, El Yunque. Distr. — Jamaica, Puerto Rico. Ref. — Alexander, 1964a:83, figs. 90 (wing), 94, 95 (male hypopygium).
- QUADRIFURCATA* Alexander, 1928a:117. Type-locality: Mexico, Chiapas, Rancho Sabino Pérez. Distr. — Mexico, El Salvador, Venezuela, Ecuador. Ref. — Alexander, 1947e:73, fig. 14 (male hypopygium).
- SATURATA* Alexander, 1927d:312. Type-locality: Mexico, Chiapas, San Cris-tóbal. Distr. — Mexico.
- SCABRIFOLIA* Alexander, 1967c:379. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- SPLENDIDA* Alexander, 1913b:514, fig. 28 (wing). Type-locality: Guatemala, Totonicapán. Distr. — Mexico, Guatemala.
- SUBCYNTHIA* Alexander, 1967c:380. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- SUBDULCIS* Alexander, 1937e:533. Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- SUBHISTRIO* Alexander, 1966c:490. Type-locality: Ecuador, Napo-Pastaza, Libertad. Distr. — Ecuador.
- SURINAMENSIS* Alexander, 1947a:180. Type-locality: Surinam, Paramaribo. Distr. — Surinam.
- THALIA* Alexander, 1945a:34, fig. 14 (male hypopygium). Type-locality: São Paulo, Juquiá. Distr. — Brazil.
- TRIANGULARIS* Alexander, 1945a:35, fig. 13 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- TROGLODYTA* Edwards, 1918:425, fig. (male hypopygium). Type-locality: Trinidad, in Guacharo Cave. Distr. — Trinidad.
- TURRIALBAE* Alexander, 1945h:290. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- VENUSTIFES* Alexander, 1926a:162. Type-locality: Mexico, Coscomatepec. Distr. — Mexico.
- WHITEI* Alexander, 1930b:119. Type-locality: Guatemala, Bananera. Distr. — Guatemala.
- WITHYCOMBEI* Alexander, 1929e:406. Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.

Subgenus PSILOCONOPA Zetterstedt

- PSILOCONOPA Zetterstedt, 1838:847 (as genus). Type-species, *meigeni* Zetterstedt (mon.) (Western Palaearctic).
WINTHEMI Alexander, 1922c:38. Type-locality: "Brazil". Distr. — Brazil. (Ex Winthem collection, Vienna Museum. Not re-discovered and the locality presumably in question, all related species being Holarctic.)

Subgenus SYMPLECTA Meigen

- Helobia* Lepeletier & Serville in Latrille *et al.*, 1828:585 (preocc. Stephens, 1827). Type-species, *Limnobia punctipennis* Meigen (mon.) = *hybrida* (Meigen).
SYMPLECTA Meigen, 1830:282 (as genus). Type-species, *Limnobia punctipennis* Meigen (Westwood, 1840:128) = *hybrida* (Meigen).
Idioneura Philippi, 1865:615. Type-species, *macroptera* Philippi (mon.).
CANA (Walker), 1848:48 (*Limnobia*). Type-locality: Canada, Ontario. Distr. — Mexico, Guatemala; Nearctic.
COLOMBIANA COLOMBIANA (Alexander), 1937d:616 (*Helobia*). Type-locality: Colombia, Usme. Distr. — Colombia. Ref. — Alexander, 1947c:329, fig. 12 (wing).
ssp. MICROPTERA Alexander, 1945a:36 [*Erioptera* (*Helobia*)]. Type-locality: Ecuador, Chimborazo, Riobamba. Distr. — Ecuador.
MACROPTERA MACROPTERA (Philippi), 1865:615 (*Idioneura*). Type-locality: unknown. Distr. — Chile, Argentina.
ssp. ARGENTINA Alexander, 1944h:383. Type-locality: Argentina, Jujuy, Hornadita. Distr. — Bolivia, Argentina. Ref. — Bruch, 1939a: pl. 6, fig. 21 (adult in resting position), pl. 7, fig. 25 (adult female).

Subgenus TRIMICRA Osten Sacken

- TRIMICRA Osten Sacken, 1861:290 (as genus). Type-species, *anomala* Osten Sacken (mon.) = *pilipes* (Fabricius). Ref. — Alexander, 1947c:332.
PILIPES (Fabricius), 1787:324 (*Tipula*). Type-locality: "Europe". Distr. — Virtually cosmopolitan. Refs. — Alexander, 1962d:4 (distribution, list of all synonyms), figs. 5 (venation), 6 (male hypopygium); Bruch, 1939a:pl. 6, fig. 20 (entire male).
reciproca Walker, 1848:50 (*Limnobia*). Type-locality: Uruguay, Montevideo.
obscurata Blanchard in Gay, 1852:343 (*Limnobia*). Type-locality: "Chile".
trichopus Philippi, 1865:610 (*Limnophila*). Type-locality: Chile, Santiago.
apocela Philippi, 1865:611 (*Limnophila*). Type-locality: "Chile".
andensis Alexander, 1916a:20, pl. 2, fig. 8 (wing) (*Trimicra*). Type-locality: Ecuador, Alausi.

Genus EUGNOPHOMYIA Alexander

- Gnophomyia*, subg. EUGNOPHOMYIA Alexander, 1947b:72. Type-species, *luc-tuosa* Osten Sacken (orig. des.) (Nearctic).



- AZRAEL (Alexander), 1943g:542. fig. 7 (male hypopygium) [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. -- Peru.
- CURRANIANA (Alexander), 1945h:285 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- DARLINGTONI (Alexander), 1937e:530, figs. 7 (wing), 10 (male hypopygium) [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Cuba, Oriente, Sierra Maestra, Pico Turquino. Distr. — Cuba.
- EXCORDIS (Alexander), 1951a:108 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Guatemala, Chicacao, El Naranjo. Distr. — Guatemala.
- FLAGRANS (Alexander), 1928h:364 (*Gnophomyia*). Type-locality: Costa Rica, Cartago. Distr. — Costa Rica.
- FLAMMEITHORAX (Alexander), 1949a:108 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Peru, Huanuco, Carpish. Distr. — Peru.
- FUNEBRIS (Alexander), 1922c:45 (*Gnophomyia*). Type-locality: Brazil, Bahia. Distr. — Brazil.
- FUNEREA (Alexander), 1938f:255 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FUSCOCOSTATA (Alexander), 1959:446 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.
- GLABRIPENNIS (Alexander), 1948b:145 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- COLBACHI (Alexander), 1962b:354, figs. 17 (wing), 20 (male hypopygium) [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Argentina, Tucumán, Quebrada Cainzo, Tafi. Distr. — Argentina.
- JUNINIANA (Alexander), 1951a:109 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- LEUCOPLACA (Alexander), 1921d:76 (*Gnophomyia*). Type-locality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- LUCTUOSA (Osten Sacken), 1859:224 (*Gnophomyia*). Type-locality: U.S.A., Florida. Distr. — Nearctic; Guatemala, Colombia. (Neotropical records presumably in error).
- nigricola* Walker, 1861:333 (*Limnobia*). Type-locality: "U.S.A."
- MELANCHOLICA (Alexander), 1922c:46 (*Gnophomyia*). Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Paraguay.
- PAMMELAS (Alexander), 1922c:47 (*Gnophomyia*). Type-locality: Paraguay, Trinidad, Asunción. Distr. — Paraguay.
- POSTICATA (Alexander), 1939e:628, figs. 3 (wing), 12 (male hypopygium) [*Gnophomyia* (*Gnophomyia*)]. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- TEMPESTIVA (Alexander), 1943a:239 [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- VIVASBERTHIERI (Alexander), 1944c:48, fig. 8 (male hypopygium) [*Gnophomyia* (*Eugnophomyia*)]. Type-locality: Venezuela, Caracas, El Valle. Distr. — Venezuela.

Genus GNOPHOMYIA Osten Sacken

- GNOPHOMYIA Osten Sacken, 1859:223. Type-species, *tristissima* Osten Sacken (Coquillett, 1910:547) (Nearctic). Ref. — Alexander, 1947b:73, figs. 8-12.



- ACRICULA Alexander, 1931c:637. Type-locality: Colombia, Magdalena, Mt. Santa Marta. Vista Nieve. Distr. — Colombia.
- ADJUSTA Alexander, 1949a:115. Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru. Ref. — Alexander, 1949c:594, fig. 11 (male hypopygium).
- APICULARIS Alexander, 1945b:59. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- ARCUATA Alexander, 1914e:244, pl. 4, fig. 6 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana.
- ARGUTULA Alexander, 1944a:45. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- ASTUTA Alexander, 1965a:404. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- AXILLARIS Alexander, 1921d:75. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- BANKSIANA Alexander, 1945L:388, fig. 18 (male hypopygium). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- BIPECTINATA Alexander, 1965a:404. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- BULBIBASIS Alexander, 1944a:46. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador. Ref. — Alexander, 1949c:594, fig. 12 (male hypopygium).
- CALOPTERA Osten Sacken, 1887b:200. Type-locality: "Brazil" (with a question). Distr. — Brazil.
- CERTA Alexander, 1952a:130. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- CHILOTA Alexander, 1929a:179, figs. 99 (wing), 220 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- CONITALIS Alexander, 1944d:317. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- DELECTABILIS Alexander, 1965a:405. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- DIACANTHA Alexander, 1966c:476. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- DIAZI Alexander, 1937a:184, fig. 3 (wing). Type-locality: Puerto Rico, Luquillo Mountains, Nevis. Distr. — Puerto Rico. Ref. — Alexander, 1947b:75, fig. 12 (male hypopygium).
- DICTENA Alexander, 1942e:257. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador.
- DIGITIFORMIS Alexander, 1941b:144. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- DISTIFURCULA Alexander, 1943g:546, fig. 8 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- DUPLEX Alexander, 1937c:498. Type-locality: Peru, Yunquilla. Distr. — Peru.
- EMARGINATA Alexander, 1945L:389, fig. 19 (male hypopygium). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- FERRUGINEA Williston, 1900:226. Type-locality: Mexico, Tabasco, Tcapa. Distr. — Mexico.
- FESSA Alexander, 1944a:47. Type-locality: Ecuador, Manabí, Palmar, Río Maizito. Distr. — Venezuela, Ecuador.
- FLEBILIS Alexander, 1944d:320. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.



- FUSCOCOSTALIS Alexander, 1947c:69, fig. 4 (wing). Type-locality: Venezuela, Miranda. Río Chacaito. Distr. — Venezuela.
- GLABRITERGATA Alexander, 1965a:406. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- JUSTA Alexander, 1938c:330. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- JUSTOIDES Alexander, 1938f:254. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- KERTESZIANA Alexander, 1929c:168, fig. 3 (male hypopygium). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- LACHRYMOSA Alexander, 1919a:143, fig. 13 (wing). Type-locality: Panama, Porto Bello. Distr. — Panama.
- LATA Alexander, 1943i:757. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- LATICINCTA Alexander, 1919a:142, fig. 14 (wing). Type-locality: Panama, Porto Bello. Distr. — Panama.
- LATILOBATA Alexander, 1949c:594, fig. 13 (male hypopygium). Type-locality: Peru, Loreto, Cerro Azul. Distr. — Peru.
- LATISSIMA Alexander, 1966c:478. Type-locality: Ecuador, Río Frío, Balao Chico. Distr. — Ecuador.
- LONGITEREBRA Alexander, 1943a:240. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- LONGITERGATA Alexander, 1943g:551, fig. 11 (male hypopygium) Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- MAESTITIA Alexander, 1914b:208, pl. 9, fig. 8 (wing). Type-locality: Peru, Vilcanota. Distr. — Peru. Ref. — Alexander, 1929c:168, fig. 4 (male hypopygium).
- MAGICA Alexander, 1945L:391. Type-locality: Guyana, Pacaraima Mountains near the headwaters of the Mazaruni River. Distr. — Guyana.
- MAGNIARCUATA Alexander, 1949a:108. Type-locality: Peru, Huanuco, Huamincha. Distr. — Peru.
- MEDIOTUBERCULATA Alexander, 1949a:118. Type-locality: Peru, Loreto, Contamana, Upper Ucayali River. Distr. — Peru. Ref. — Alexander, 1949c:596, fig. 14 (male hypopygium).
- MOLINAE Alexander, 1929c:164, fig. 1 (male hypopygium). Type-locality: Ecuador, Guayas, Bucay, Ventura. Distr. — Ecuador.
- MONOPHAEA Alexander, 1946a:134. Type-locality: Mexico, Chiapas, Nueva America. Distr. — Mexico.
- NAHUELBTAE Alexander, 1966c:479, figs. 11 (wing), 13 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Pillim Pilli. Distr. — Chile.
- NECTARAEA Alexander, 1944a:49. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- NGRINA (Wiedemann), 1828:37 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil, Peru.
- NIMBIFERA Alexander, 1943g:548, fig. 9 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- OSTENSACKENI Skuse, 1890a:825 (nom. nov. for *fascipennis* Osten Sacken). Type-locality: Brazil, Amazon River. Distr. — Brazil.
- fascipennis* Osten Sacken, 1887b:199 (preocc. Thomson, 1869). Type-locality: Brazil, Amazon River.

- OXYMERA Alexander, 1943g:549, fig. 10 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- PALLIDAPEX Alexander, 1929d:783. Type-locality: Paraguay, Santa Barbara. Distr. — Brazil, Paraguay. Ref. — Alexander, 1947b:75, fig. 8 (wing).
- PAUCISETA Alexander, 1949a:116. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:596, fig. 15 (male hypopygium).
- PERACUTOR Alexander, 1949a:113. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:596, fig. 16 (male hypopygium).
- PERDEBILIS Alexander, 1949a:109. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru, Bolivia. Ref. — Alexander, 1949c:596, fig. 17 (male hypopygium).
- PERLATA Alexander, 1944d:320. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- PERMAGICA Alexander, 1947b:76, fig. 10 (male hypopygium). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- PERSEVERA Alexander, 1949a:117. Type-locality: Peru, Loreto, Cerro Azul. Distr. — Peru. Ref. — Alexander, 1949c:597, fig. 18 (male hypopygium).
- PETENTIS Alexander, 1949a:110. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:597, fig. 19 (male hypopygium).
- PODACANTHA Alexander, 1944h:380. Type-locality: Costa Rica, Talamanca, San Juan Miramar. Distr. — Costa Rica.
- PORTERI Alexander, 1929c:165, fig. 2 (male hypopygium). Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- PROPATULA Alexander, 1947a:175. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- PULVINARIS Alexander, 1945h:286. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- REGNATRIX Alexander, 1943g:544. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- RUBICUNDULA Alexander, 1921d:74. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- SAGITTA Alexander, 1949c:597, fig. 20 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- SAGITTODES Alexander, 1951a:109. Type-locality: Peru, Huanuco, Fundo Sinchono, Cordillera Azul. Distr. — Peru.
- SOCIALIS Alexander, 1949a:112. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru. Ref. — Alexander, 1949c:598, fig. 21 (male hypopygium).
- SPINIBASIS Alexander, 1943i:758, fig. 11 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- STENOCHOREMA Alexander, 1962c:137, figs. 115 (wing), 119 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chaparc, Locotal. Distr. — Bolivia.
- STENOPHALLUS Alexander, 1941b:145. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- STUPENS (Walker), 1861:333 (*Limnobia*). Type-locality: "Mexico". Distr. — Mexico. Identity uncertain.

- STYLACUTA Alexander, 1947b:77, fig. 11 (male hypopygium). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- SUBAPICULARIS Alexander, 1945b:60. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- SUBARCUATA Alexander, 1946e:533. Type-locality: Mexico, Chiapas, Finca Vergel. Distr. — Mexico.
- SUBFLEBILIS Alexander, 1949a:113. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador. Ref. — Alexander, 1949c:598, fig. 22 (male hypopygium).
- SUBHYALINA Alexander, 1913b:523, fig. 23 (wing). Type-locality: Guatemala, Alta Verapaz, Cacao Trece Aguas. Distr. — Guatemala, Costa Rica, Panama, Guyana, Brazil.
- SUBNIMBIFERA Alexander, 1966c:480. Type-locality: Ecuador, Napo-Pastaza, Libertad, 6 km southeast of Tena. Distr. — Ecuador.
- SUBOBLITERATA Alexander, 1946e:532. Type-locality: Mexico, Morelos, Tezotlán. Distr. — Mexico.
- TELENEURA Alexander, 1944d:318. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- TETRAEENA Alexander, 1966c:481. Type-locality: Ecuador, Napo-Pastaza, Libertad, 6 km. southeast of Tena. Distr. — Ecuador.
- TRESIAS Alexander, 1949c:598, fig. 23 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- TOLERANDA Alexander, 1954d:148. Type-locality: Bolivia, Cochabamba, Chapore, Locotal, El Limbo. Distr. — Bolivia.
- TRANSVERSA Alexander, 1949c:600, fig. 24 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- TRIATRATA Alexander, 1968b:90. Type-locality: Brazil, São Paulo, Ribeirão Pires. Distr. — Brazil.
- TRICEPS Alexander, 1947b:78, fig. 9 (male hypopygium). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- TRICEPOIDES Alexander, 1966c:482. Type-locality: Peru, Cuzco, Quíncemil, along the Marcapata river. Distr. — Peru.
- TRICORNIS Alexander, 1949a:114. Type-locality: Peru, Huanuco, Sariapampa. Distr. — Peru. Ref. — Alexander, 1949c:602, fig. 25 (male hypopygium).
- TRISETIGERA Alexander, 1949a:111. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:602, fig. 26 (male hypopygium).
- TUBER Alexander, 1944a:50. Type-locality: Ecuador, Los Ríos, Playas de Montalvo. Distr. — Ecuador.
- TUNGURAHUANA Alexander, 1945n:417. Type-locality: Ecuador, Tungurahua, Mt. Tungurahua. Distr. — Ecuador.
- VILIS Alexander, 1931c:638. Type-locality: Colombia, Orihueca. Distr. — Colombia.
- VITRIPENNIS Alexander, 1942e:258. Type-locality: Ecuador, Manabi, Palmar. Distr. — Ecuador.

Genus GONOMYIA Meigen

- GONOMYIA Meigen, 1818:146. Type-species, *Limnobia tenella* Meigen (mon.). [See note in Alexander, 1965d:73].
- Gonomyia*, emend.
Ref. — Alexander, 1947b:83-99, figs. 17-29.



Subgenus EUPTILOSTENA Alexander

- Gonomyia*, subg. EUPTILOSTENA Alexander, 1938j:126. Type-species, *reticulata* Alexander, (orig. des.) (Eastern Palaearctic)
- DAMPFIANA Alexander, 1938e:395. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico. Ref. — Alexander, 1947b:87, fig. 19 (wing).
- POLINGI Alexander, 1946e:523. Type-locality: U.S.A., Texas, Davis Mts., Old Fort Davis. Distr. — Southwestern U.S.A., Mexico.

Subgenus GONOMYIA Meigen

- AEQUALIS Alexander, 1916b:323. Type-locality: Guatemala, Totonicapán. Distr. — Guatemala. Ref. — Alexander, 1916d:526, fig. 86 (male hypopygium).
- ANDICOLA Alexander, 1913c:199, pl. 5, fig. 6 (wing), pl. 6, fig. 3 (male hypopygium). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- ANDUZEI Alexander, 1940e:902, figs. 4 (wing), 7 (male hypopygium). Type-locality: Venezuela. Naiguata, D.F., Los Canales. Distr. — Venezuela.
- ANIMULA Alexander, 1966b:703. Type-locality: Honduras, Lancetilla. Distr. — Honduras
- ANSERINA Alexander, 1943i:752, fig. 8 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- APPENDICULATA Alexander, 1943i:751, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- ASPERA Alexander, 1943a:231, fig. 1 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- AYMARA Alexander, 1962c:148, figs. 127 (wing), 129 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- BIFURCIFER Alexander, 1939e:626, figs. 2 (wing), 10 (male hypopygium). Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- BIFURCULA Alexander, 1946a:130. Type-locality: Mexico, Chiapas, San Antonio Nexapa. Distr. — Mexico.
- EIRAMA Alexander, 1941e:251, fig. 9 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru. Ref. — Alexander, 1947b:85, fig. 21 (wing).
- BREVICULA Alexander, 1926b:225. Type-locality: Cuba, Hanabanilla Falls near Cumanayagua. Distr. — Cuba.
- BREVISSIMA Alexander, 1926b:227. Type-locality: Cuba, Trinidad Mountains. Distr. — Cuba.
- CATAMARCAE Alexander, 1929d:785. Type-locality: Argentina, Catamarca, Bolsón. Distr. — Argentina. Ref. — Alexander, 1962b:355, fig. 21 (male hypopygium).
- CHIAPASENSIS Alexander, 1927d:306. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- CONNIVENS Alexander, 1946b:242. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- CRINITA Alexander, 1941d:127, fig. 11 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Peru.
- DASYPHALLUS Alexander, 1949c:610, fig. 32 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.



- DEBILIS Alexander, 1928a:116. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico.
- DELICATA Alexander, 1913b:506. Type-locality: Guatemala, Totonicapán. Distr. — Guatemala.
- EFFICIENS Alexander, 1943a:232. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- EXPANSA Alexander, 1938e:403. Type-locality: Mexico, Chiapas, Finca Lubeca. Distr. — Mexico.
- FLAVIBASIS Alexander, 1916b:317. Type-locality: U.S.A., California, Monterey County. Distr. — Western U.S.A., Mexico.
tuberculata Alexander, 1925a:349, figs. 9 (wing), 18 (male hypopygium). Type-locality: Mexico, Penonmarquez.
- FURCULA FURCULA Alexander, 1962c:149, figs. 128 (wing), 130 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
ssp. BREVISPIDA Alexander, 1962c:150, fig. 131 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- GEMULA Alexander, 1946c:4, figs. 4 (wing), 6 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- GUERREROENSIS Alexander, 1940b:161. Type-locality: Mexico, Guerrero, Taxco. Distr. — Mexico.
- ILICIS Alexander, 1921a:74. Type-locality: Argentina, Jujuy, Ciudad. Distr. — Argentina.
- JAMAICANA Alexander, 1964a:76, figs. 81 (wing), 85 (male hypopygium). Type-locality: Jamaica, St. Thomas, Whitfield Hall. Distr. — Jamaica.
- JEJUNA Alexander, 1916a:17, pl. 1, fig. 8 (wing), pl. 4, fig. 1 (male hypopygium). Type-locality: Peru, Lima. Distr. — Peru.
- JUAREZI Alexander, 1946a:131. Type-locality: Mexico, Oaxaca, Sierra de Juárez, Yotao. Distr. — Mexico.
- MEGARHOPALA Alexander, 1946a:129. Type-locality: Mexico, Morcos, Lagunas de Zempole. Distr. — Mexico.
- METHODICA Alexander, 1931c:635. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- MEXICANA Alexander, 1916b:321. Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico. Rcf. — Alexander, 1916d:527, figs. 90, 91 (male hypopygium).
- MICROSERRATA Alexander, 1938e:399. Type-locality: Mexico, Chiapas, Finca Germania. Distr. — Mexico.
- MULTISPICATA Alexander. 1938e:402. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- OSTENTATOR Alexander, 1946c:537. Type-locality: Mexico, Morcos, Cuernavaca. Distr. — Mexico.
- OTIOSA Alexander, 1949c:612, fig. 3 (male hypopygium). Type-locality: Peru, Junin, Chanchamayo. Distr. — Peru.
- PLATYMERINA Alexander, 1944k:612. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- PLATYMEROIDES Alexander, 1952a:132. Type-locality: Peru, Junin, Chanchamayo. Distr. — Peru.
- QUAESITA Alexander, 1938e:400. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- QUERIBUNDA Alexander, 1941c:252, fig. 10 (male hypopygium). Type-locality: Peru, Junin, Tarma, Huacapistana. Distr. — Peru.

- RECTANGULA Alexander, 1966b:704. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- REMIGERA Alexander, 1946e:536. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- REMOTA REMOTA Alexander, 1926a:164, fig. 5 (wing). Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico, El Salvador.
- ssp. OBTUSISTYLA Alexander, 1940c:115. Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Panama.
- SALMANI Alexander, 1927c:274. Type-locality: El Salvador, Agronomía, Sonsonate. Distr. — El Salvador.
- SERPENTINA Alexander, 1938a:34. Type-locality: Brazil. Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SPICULISTYLA Alexander, 1950a:53, figs. 30 (wing), 35 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- STELLATA Alexander, 1946e:535. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- SUBAPPENDICULATA Alexander, 1954a:39, fig. 5 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- SUBBREVICULA Alexander, 1947b:89, fig. 20 (wing). Type-locality: Puerto Rico, El Semil. Distr. — Puerto Rico.
- SUBREMOTA Alexander, 1938c:401. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- SUBUNICOLOR Alexander, 1948b:146. Type-locality: Panama, Cerro Peña, near El Valle. Distr. — Panama.
- TRIACULEATA Alexander, 1938e:398. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- UNICOLOR Alexander, 1913b:507, fig. 15 (wing). Type-locality: Guatemala, Aguna. Distr. — Mexico, Guatemala, El Salvador. Ref. — Alexander, 1916d:527, pl. 30, figs. 88, 89 (male hypopygium).
- WYGODZINSKYI Alexander, 1962b:356, figs. 19 (wing), 23 (male hypopygium). Type-locality: Argentina, Tucumán, La Angostura, Tafi del Valle. Distr. — Argentina.

Subgenus GONOMYINA Alexander

- Gonomyia*, subg. GONOMYINA Alexander, 1946c:5 Type-species, *durabilis* Alexander (orig. des.).
- DURABILIS Alexander, 1946c:6, figs. 3 (wing), 7 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- PARISHI (Alexander), 1913a:46, fig. k (male hypopygium) (*Teucholabis*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- PERSIMILIS (Alexander), 1920b:6 (*Teucholabis*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- RUNA Alexander, 1947b:84, fig. 24 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Subgenus IDIOCERA Dale

- Limnobia*, subg. IDIOCERA Dale, 1842:431, 433. Type-species, *sexguttata* Dale (mon.) (Western Palaearctic).
- ANGUSTISSIMA Alexander, 1928b:57 [*Gonomyia* (*Ptilostena*)]. Type-locality: Cuba, Oriente, Jarahueca. Distr. — Cuba.



HOOGSTRAALI Alexander, 1946b:241, figs. 20 (wing), 23 (male hypopygium).
Type-locality: Mexico, Michoacán, La Majada. Distr. — Mexico.
Ref. — Alexander, 1947b: 87, fig. 18 (wing).

Subgenus LIPOPHLEPS Bergroth

Leiponeura Skuse, 1890a:795. Type-species, *gracilis* Skuse (Alexander, 1913b:503) = *skusei* Alexander. [See note in Alexander, 1965d:74].

LIPOPHLEPS Bergroth, 1915:55 (as genus; unjustified nom. nov. for *Leiponeura* Skuse). Type-species, *Leiponeura gracilis* Skuse (aut.) = *skusei* Alexander.

Ref. — Alexander, 1947b:91.

ADUNCA Alexander, 1921e:294 [*Gonomyia* (*Leiponeura*)]. Type-locality: Argentina, Catamarca, Ampajango. Distr. — Argentina.

ANDUZEANA Alexander, 1941b:141. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela. Ref. — Alexander, 1947e:73, fig. 10 (male hypopygium).

ARAJUNO Alexander, 1945n:416. Type-locality: Ecuador, Napo-Pastaza, Río Arajuno, Napo Watershed. Distr. — Ecuador. Ref. — Alexander, 1945L:386, fig. 16 (male hypopygium).

EASISPINOSA Alexander, 1936e:63. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1947b:89, fig. 27 (male hypopygium).

BATESI Alexander, 1945L:386, fig. 17 (male hypopygium). Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.

BICORNUTA Alexander, 1927c:276. Type-locality: Puerto Rico, Luquillo National Forest. Distr. — Puerto Rico. Ref. — Alexander, 1932:372, fig. 22 (male hypopygium).

BIFILIGERA Alexander, 1932:372, figs. 15 (wing), 21 (male hypopygium). Type-locality: Puerto Rico, Las Cruces. Distr. — Puerto Rico.

BISPINOSA Alexander, 1921d:82, figs. 4,4A (male hypopygium) [*Gonomyia* (*Leiponeura*)]. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.

BORBURATANA Alexander, 1941b:143. Type-locality: Venezuela, Carabobo, Borburata. Distr. — Venezuela. Ref. — Alexander, 1947e:73, fig. 11 (male hypopygium).

BRUCHI Alexander, 1920f:217 [*Gonomyia* (*Leiponeura*)]. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Bruch, 1939a: pl. 6, fig. 19 (entire female).

CALVERTI CALVERTI Alexander, 1914c:122, figs. 7, 8 (male hypopygium) [*Gonomyia* (*Leiponeura*)]. Type-locality: Costa Rica, Alajuela. Distr. — Costa Rica.

ssp. PLEUROTAENIATA Alexander, 1942j:31 (nom. nov. for *pleurostriata* Alexander). Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.

pleurostriata Alexander, 1939e:627, fig. 11 (male hypopygium) (preocc. Alexander, 1936). Type-locality: Costa Rica, Talamanca, Pedregoso.

CANTAREIRAE Alexander, 1945c:270, fig. 17 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.

CARRERAI Alexander, 1943f:304, fig. 14 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.

- CERVARIA Alexander, 1938d:334. Type-locality: Colombia, Cundinamarca, Fusagasugá. Distr. — Colombia.
- CLAVIFERA Alexander, 1943i:755, fig. 10 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- CREPUSCULA Alexander, 1921d:82, figs. 5, 5A (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Peru, Loreto, Napo River. Distr. — Cuba.
- CTENOPHORA Alexander, 1921d:81, fig. 6 (male hypopygium) (*Gonomyia (Leiponeura)*). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- CUBANA Alexander, 1931a:120. Type-locality: Cuba, Tiisdad Mts. Distr. — Cuba.
- CULTRATA Alexander, 1941d:130, fig. 14 (male hypopygium). Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- DIACANTHOPHORA Alexander, 1941a:253, fig. 11 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.
- DOTATA Alexander, 1945i:253, fig. 4 (male hypopygium). Type-locality: Peru, Junín. Satipo, Jauja. Distr. — Peru.
- DUURVOORTI Alexander, 1928h:366. Type-locality: Surinam, Paramaribo. Distr. — Guyana, Surinam.
- EXTENSA Alexander, 1914e:243 [*Gonomyia (Leiponeura)*]. Type-locality: Guyana, Bartica. Distr. — Panama, Guyana, Trinidad.
- FORTIBASIS Alexander, 1966c:486. Type-locality: Ecuador, Esmeralda, San Mateo. Distr. — Ecuador.
- GILLOTTAE Alexander, 1928h:365. Type-locality: Costa Rica, near Cartago. Distr. — Costa Rica, Panama.
- HAPLOA Alexander, 1926a:167, fig. 18 (male hypopygium). Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico.
- HAPLOIDES Alexander, 1938e:397. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico, Costa Rica.
- HOFFMANIANA Alexander, 1947b:93, figs. 22 (wing), 29 (male hypopygium). Type-locality: Puerto Rico, El Semil. Distr. — Puerto Rico.
- IMPEDITA Alexander, 1946a:132. Type-locality: Mexico, Oaxaca, Cacalotepec, Sierra de Juárez. Distr. — Mexico.
- INERMIS Alexander, 1914e:242, pl. 4, fig. 5 (wing) [*Gonomyia (Leiponeura)*]. Type-locality: Guyana, Bartica. Distr. — Guyana.
- JUQUIANA Alexander, 1945c:273, figs. 15 (wing), 20 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- LEONURA Alexander, 1941d:128, fig. 12 (male hypopygium). Type-locality: Ecuador, Bolívar, Balzapamba. Distr. — Ecuador.
- LUSTRALIS Alexander, 1945h:289. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- MACINTYREI Alexander, 1937d:609. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- MACSWAINI Alexander, 1940c:115. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Costa Rica, Panama.
- MAYA Alexander, 1927d:307. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico.
- MECOPHALLUS Alexander, 1966b:702. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- MELANACANTHA Alexander, 1954a:40, fig. 6 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.



- MISERA Alexander, 1921e:164 [*Gonomyia (Leiponeura)*]. Type-locality: Argentina, Tucumán, Arroyo Famailla. Distr. — Argentina.
- MYTHICA Alexander, 1945i:250, fig. 3 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- NESTOR Alexander, 1947b:95, fig. 26 (male hypopygium). Type-locality: Brazil, São Paulo. Distr. — Brazil.
- ORTHOMERA Alexander, 1937a:185, fig. 10 (male hypopygium). Type-locality: Puerto Rico, Tanama River. Distr. — Puerto Rico.
- ORTHOMEROIDES Alexander, 1939a:85. Type-locality: Mexico, Chiapas, Huixtla. Distr. — Mexico.
- PARINERMIS Alexander, 1940e:903, fig. 8 (male hypopygium), Type-locality: Venezuela, Naiguata, D.F., Los Canales. Distr. — Venezuela.
- PEDICA Alexander, 1947b:96, fig. 25 (male hypopygium). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- PETRONIS Alexander, 1942h:434, figs. 14 (wing), 18 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- PHILOMELA Alexander, 1945i:251, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PHOROCENIA Alexander, 1921d:80, fig. 7 (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- PHLOSISPINA Alexander, 1936e:65. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PRODUCTA Alexander, 1919a:139 [*Gonomyia (Leiponeura)*]. Type-locality: Lesser Antilles: Antigua, Blubber Valley. Distr. — Mexico, Ecuador, Puerto Rico, Antigua. Ref. — Alexander, 1932:374, fig. 23 (male hypopygium).
- PROJECTA Alexander, 1941d:129, fig. 13 (male hypopygium). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador, Peru, Bolivia.
- PROLIXISTYLUS Alexander, 1930b:118. Type-locality: British Honduras, Blue Creek. Distr. — Mexico, British Honduras.
- PROLONGATA Alexander, 1940e:904, figs. 5 (wing), 9 (male hypopygium). Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela, Ecuador.
- PUELLA (Williston), 1896a:288, fig. 60 (*Atarba*). Type-locality: Lesser Antilles: St. Vincent. Distr. — St. Vincent. Refs. — Alexander, 1913b:505; 1947b:95, fig. 28 (male hypopygium).
- PUER Alexander, 1913b:506, fig. 14 (wing) [*Gonomyia (Leiponeura)*]. Type-locality: Hispaniola: Dominican Republic. Distr. — Mexico, Guyana, Ecuador, Peru, Jamaica, Dominican Republic. Ref. — Alexander, 1916d:516, figs. 21 (wing), 63 (male hypopygium).
- PYENSONI Alexander, 1939e:441. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- RAMUS Alexander, 1943e:210. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- RASTRIFORMIS Alexander, 1946e:534. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- RECURVATA Alexander, 1941c:121, fig. 6 (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Costa Rica, Alajuela. Distr. — Costa Rica, Ecuador.
- REYESI Alexander, 1946a:133. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.

- RHADINOSTYLA** Alexander, 1964a:78, figs. 82 (wing), 86 (male hypopygium).
 Type-locality: Jamaica, St. Thomas, Bath. Distr. — Jamaica.
- SANA** Alexander, 1936d:763. Type-locality: Paraguay, Guaira, Villarrica.
 Distr. — Paraguay.
- SANDERSI** Alexander, 1931a:121. Type-locality: Cuba, Soledad. Distr. —
 Cuba.
- SCELERATA** Alexander, 1945i:254, fig. 5 (male hypopygium). Type-locality:
 Peru, Junín, Satipo, Jauja. Distr. — Peru.
- SCIMITAR** Alexander, 1914e:243 [*Gonomyia (Leiponeura)*]. Type-locality:
 Guyana, Bartica. Distr. — Guyana.
- SECESPITA** Alexander, 1936e:64. Type-locality: Brazil, Santa Catarina,
 Nova Teutônia. Distr. — Brazil.
- SENARIA** Alexander, 1943a:234, fig. 2 (male hypopygium). Type-locality:
 Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- SPICATA** Alexander, 1921d:83, fig. 3 (male hypopygium) [*Gonomyia (Leipo-*
neura)]. Type-locality: Peru, Loreto, Napo River. Distr. — Peru.
- SPINITERGA** Alexander, 1948b:147. Type-locality: Brazil, São Paulo, Bora-
 céia. Distr. — Brazil. Ref. — Alexander, 1966c:487, fig. 14 (male
 hypopygium).
- SUBINERMIS** Alexander, 1939a:84. Type-locality: Mexico, Chiapas, Escuintla.
 Distr. — Mexico.
- SUBLUSTRALIS** Alexander, 1966c:487. Type-locality: Peru, Santa Margarita,
 Río Perené. Distr. — Peru.
- SUBSCIMITAR** Alexander, 1945c:605. Type-locality: Peru, Junín, Satipo,
 Jauja. Distr. — Peru. Ref. — Alexander, 1949c:613, fig. 34 (male
 hypopygium).
- SUBTERMINALIS** Alexander, 1927c:275. Type-locality: Puerto Rico, Luquillo
 National Forest. Distr. — Cuba, Puerto Rico. Ref. — Alexander,
 1932:374, fig. 24 (male hypopygium).
- SUBTRIBULATOR** Alexander, 1943i:754, fig. 9 (male hypopygium). Type-loca-
 lity: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- SUBTROLUS** Alexander, 1966c:488. Type-locality: Peru, Santa Margarita,
 Río Perené. Distr. — Peru.
- TAFIENSIS** Alexander, 1962b:357, figs. 18 (wing), 22 (male hypopygium).
 Type-locality: Argentina, Tucumán, Quebrada Cainzo, Tafi. Distr.
 — Argentina.
- TERGOFIMBRIATA** Alexander, 1928h:367. Type-locality: Guyana, Bartica. Distr.
 — Guyana.
- TERSA** Alexander, 1949d:263. Type-locality: Peru, Loreto, Pucallpa. Distr.
 — Peru. Ref. — Alexander, 1949c:613, fig. 35 (male hypopygium).
- TRIBULATOR** Alexander, 1942c:439, fig. 11 (male hypopygium). Type-locali-
 ty: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- TRIONYX** Alexander, 1949b:494, fig. 25 (male hypopygium). Type-locality:
 Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. —
 Brazil.
- TROLUS** Alexander, 1949d:262. Type-locality: Peru, Junín, Tarma, Utcuyacu.
 Distr. — Peru. Ref. — Alexander, 1949c:613, fig. 36 (male hypo-
 pygium).
- VEHEMENS** Alexander, 1949d:264. Type-locality: Peru, Junín, Tarma,
 Utcuyacu. Distr. — Peru. Ref. — Alexander, 1949c:613, fig. 37
 (male hypopygium).

VINDEX Alexander, 1941b:142. Type-locality: Venezuela, Carabobo, Borburata. Distr. — Venezuela. Ref. — Alexander, 1947e:73, fig. 12 (male hypopygium).

Subgenus NEOLIPOPHLEPS Alexander

- Gonomyia*, subg. NEOLIPOPHLEPS Alexander, 1947b:98. Type-species, *Dicranomyia cinerea* Doane (orig. des.).
- ACUMINATA Alexander, 1921d:77, fig. 1 (male hypopygium) [*Gonomyia Leiponeura*]. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru, Argentina.
- AEQUIDENS Alexander, 1938e:396 [*Gonomyia (Lipophleps)*]. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico, Costa Rica.
- AEQUISPINOSA Alexander, 1926d:390 [*Gonomyia (Lipophleps)*]. Type-locality: Paraguay, Santa Barbara. Distr. — Paraguay.
- CONDENSA Alexander, 1938a:33 [*Gonomyia (Lipophleps)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- EXTENUATA Alexander, 1945e:271, fig. 18 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- FALCIFERA Alexander, 1921d:79, fig. 2 (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- GLABRISPINA Alexander, 1926d:390 [*Gonomyia (Lipophleps)*]. Type-locality: Argentina, Riacho Pilaga. Distr. — Argentina.
- HELOPHILA Alexander, 1916e:343, figs. 1 (wing), 3 (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Peru, Lima. Distr. — Mexico, Dominica, Central America to Peru. Refs. — Alexander, 1932:373, fig. 20 (male hypopygium); 1937b:190, fig. 9 (male hypopygium).
- MACHAERIA Alexander, 1921d:79 [*Gonomyia (Leiponeura)*]. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- MONACANTHA Alexander, 1937b:184, fig. 8 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Puerto Rico, Vieques Island. Distr. — Puerto Rico.
- NEOFALCIFERA Alexander, 1943c:24 [*Gonomyia (Lipophleps)*]. Type-locality: Venezuela, Bolívar, Akurimán, Gran Sabana. Distr. — Venezuela.
- PLATYNERA Alexander, 1939d:128, fig. 18 (male hypopygium) [*Gonomyia (Lipophleps)*; as *monacantha* ssp.]. Type-locality: Cuba, Soledad, near Cienfuegos. Distr. — Jamaica, Cuba. Ref. — Alexander, 1964a:79, fig. 84 (venation).
- SCHADEANA Alexander, 1935a:328 [*Gonomyia (Lipophleps)*]. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Brazil, Paraguay.
- STRIGILIS Alexander, 1926a:165, fig. 14 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Mexico, between Coscomatepec and Tomatlán. Distr. — Mexico.
- SUBFALCIFERA Alexander, 1922c:44 [*Gonomyia (Leiponeura)*]. Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Paraguay.
- TOPOENSIS Alexander, 1952a:133. Type-locality: Ecuador, Tungurahua, Upper Río Topo. Distr. — Ecuador.
- TRISPINOSA Alexander, 1921a:75 [*Gonomyia (Leiponeura)*]. Type-locality: Argentina, Jujuy, Ciudad. Distr. — Argentina.

Subgenus PARALIPOPHLEPS Alexander

- Gonomyia*, subg. PARALIPOPHLEPS Alexander, 1947b:97. Type-species, *Atarba pleuralis* Williston (orig. des.).
- AMAZONA Alexander, 1921e:418, figs. 1 (wing), 2 (male hypopygium) [*Gonomyia (Leiponeura)*]. Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Ecuador, Peru, Brazil.
- DIPLOCANTHA Alexander, 1966c:485. Type-locality: Peru, Santa Margarita, Río Perené. Distr. — Peru.
- GLADIATOR Alexander, 1919a:138, fig. 11 (wing) [*Gonomyia (Leiponeura)*]. Type-locality: Panama, Real de Santa Maria. Distr. — Panama.
- GUAYAQUILENSIS Alexander, 1938h:438. Type-locality: Ecuador, Guayas, Guayaquil. Distr. — Ecuador.
- HETEROMERA Alexander, 1942c:438, fig. 10 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru, Bolivia.
- INDOTATA Alexander, 1966b:701. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- IATISTYLA Alexander, 1926a:165, fig. 15 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Mexico, Veracruz. Distr. — Mexico, Honduras.
- LEMNISCATA Alexander, 1931c:634 [*Gonomyia (Lipophleps)*]. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia, Venezuela, Brazil.
- naiguatana* Alexander, 1940e:905, fig. 10 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Venezuela, Miranda, Naiguata, D.F., Los Canales.
- MICRACANTHA Alexander, 1926d:391 [*Gonomyia (Lipophleps)*]. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- MICROMERA Alexander, 1945c:274, fig. 19 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- NEOBIFIDA Alexander, 1952f:327 (nom. nov. for *bifida* Alexander). Type-locality: Mexico, Tospán near Córdoba. Distr. — Mexico.
- bifida* Alexander, 1926a:167, fig. 17 (male hypopygium) [*Gonomyia (Lipophleps)*]; preocc. Tonnoir, 1920]. Type-locality: Mexico, Tospán near Córdoba.
- PERACUTA PERACUTA Alexander, 1928a:117 [*Gonomyia (Lipophleps)*]. Type-locality: Mexico, Chiapas, Santa Lucia. Distr. — Mexico. Ref. — Alexander, 1964a:79, figs. 83 (venation), 88 (male hypopygium).
- ssp. CONIFERA Alexander, 1941g:354 (also 1946h:101). Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Costa Rica, Panama.
- PLEURALIS (Williston), 1896a:289 (*Atarba*). Type-locality: Lesser Antilles: St. Vincent. Distr. — Southeastern U.S.A., Bermuda, through Antilles to Peru, Brazil, Bolivia, Guyana. Refs. — Alexander, 1912e:419, figs. 3, 4 (male hypopygium); 1913b:504; 1916d:516, fig. 20 (wing); 1932:374; 1947c:97, fig. 23 (wing); 1964a:79, fig. 87 (male hypopygium); Johnson, 1913:443 (Bermuda).
- SPINISTYLA Alexander, 1926a:166, fig. 16 (male hypopygium) [*Gonomyia (Lipophleps)*]. Type-locality: Mexico, Chiapas. Distr. — Mexico, Brazil.



Subgenus PROGONOMYIA Alexander

- Gonomyia*, subg. *Gonomyella* Alexander, 1917b:152 (preocc. Meunier, 1899).
Type-species, *slossonae* Alexander (mon.)
- Gonomyia*, subg. PROGONOMYIA Alexander, 1920i:938 (nom. nov. for *Gonomyella* Alexander), Type-species, *slossonae* Alexander (aut.)
- ACANTHIAS Alexander, 1941a:247, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- ACRISSIMA Alexander, 1944a:53. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- ADELA Alexander, 1949d:261. Type-locality: Peru, Huanuco, Chinchao. Distr. — Peru.
- ALTIVOLANS Alexander, 1942c:437, fig. 9 (male hypopygium). Type-locality: Peru, Huanuco, Piedra Grande. Distr. — Peru.
- ARGENTINENSIS Alexander, 1920f:216 [*Gonomyia* (*Gonomyella*)]. Type-locality: Argentina, Rioja. Distr. — Argentina.
- ATROAPICATA Alexander, 1938c:394. Type-locality: Mexico, Chiapas, Huixtla. Distr. — Mexico.
- BALZAPAMBAE Alexander, 1941d:124, fig. 9 (male hypopygium). Type-locality: Ecuador, Bolívar. Balzapamba. Distr. — Ecuador.
- BIFASCIOLATA Alexander, 1937e:531, fig. 8 (wing). Type-locality: Panama, Chiriquí, Potrerillos. Distr. — Mexico, Panama, Cuba, Jamaica. Ref. — Alexander, 1964a:75, fig. 79 (venation).
- CATAMARCENSIS Alexander, 1941d:126, fig. 10 (male hypopygium). Type-locality: Argentina, Catamarca, Santa María. Distr. — Argentina.
- COMPACTA Alexander, 1941b:140. Type-locality: Venezuela, Carabobo, Borburata. Distr. — Venezuela. Ref. — Alexander, 1947c:73, fig. 9 (male hypopygium).
- DESTRUCTA Alexander, 1939c:440, figs. 7 (wing), 13 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Alexander, 1947c:85, fig. 17 (wing).
- DOLOROSA Alexander, 1922c:41. Type-locality: Brazil, São Paulo, Iguape. Distr. — Brazil.
- ERIOPTEROIDES Alexander, 1926d:389. Type-locality: Paraguay, Santa Barbara. Distr. — Paraguay.
- FIELDI Alexander, 1966b:700. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- FORCEPS Alexander, 1941e:250, fig. 8 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- HISTRONICA Alexander, 1943i:748, fig. 6 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- HYPERPLATYS Alexander, 1944a:55. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- MAESTA Alexander, 1921a:73 [*Gonomyia* (*Gonomyella*)]. Type-locality: Argentina, Jujuy, Ciudad. Distr. — Argentina.
- MAGISTRATA Alexander, 1949c:606, fig. 30 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- OMINOSA Alexander, 1926d:388. Type-locality: Paraguay, Santa Barbara. Distr. — Paraguay.
- OVALIS Alexander, 1949c:609, fig. 31 (male hypopygium). Type-locality: Peru, Junín, Utcuyacu. Distr. — Peru, Bolivia.
- PARAENSIS Alexander, 1920b:7 [*Gonomyia* (*Gonomyella*)]. Type-locality: Brazil, Pará, Prata. Distr. — Brazil.

- PARAMOENSIS Alexander, 1944c:46, figs. 4 (wing), 7 (male hypopygium). Type-locality: Venezuela, Merida, Páramo La Negra. Distr. — Venezuela.
- PATRUELIS Alexander, 1930b:117. Type-locality: Mexico, Payo Obispo (= Ciudad Chetumal), Quintana Roo. Distr. — Mexico.
- PERUVIANA Alexander, 1922c:42. Type-locality: Peru, Lima, Matucana. Distr. — Peru.
- PLATYMERELLA Alexander, 1947c:84 (nom. nov. for *platymera* Alexander). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
platymera Alexander, 1943i:745, fig. 5 (male hypopygium) (preocc. Alexander, 1939). Type-locality: Peru, Ayacucho, La Mar, Ayna.
- PLEUROLINEATA Alexander, 1929d:784. Type-locality: Argentina,, Catamarca, Quebrada Volcán at Agua Amarilla. Distr. — Argentina.
- QUINQUEPLAGIATA Alexander, 1931a:119. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil, Argentina.
- SATURATA Alexander, 1937b:245. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- SAXICOLA Alexander, 1923b:182. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- SERENA Alexander, 1922c:43. Type-locality: "South America". Distr. — South America.
- SLOSSONAE Alexander, 1914g:588, fig. 26 (wing). Type-locality: Panama, Canal Zone, Paraiso. Distr. — Southeastern U.S.A., Mexico, Panama, Jamaica. Ref. — Alexander, 1964a:76, fig. 80 (wing).
- SUBCOSTATA Alexander, 1919a:137, fig. 10 (wing) [*Gonomyia* (*Gonomyella*)]. Type-locality: Panama, Río Chico. near mouth of Río Porcona. Distr. — Panama.
- SUBSATURATA Alexander, 1942i:67, figs. 36 (wing), 37 (male hypopygium). Type-locality: Brazil, São Paulo, Cabeceiras do M'Boy Guaçu. Distr. — Brazil.
- SYNCHROA SYNCHROA Alexander, 1929a:180, figs. 100 (wing), 221 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Correntoso. Distr. — Argentina.
- ssp. SETOSIVENA Alexander, 1929a:181. Type-locality: Chile, Valparaiso. Marga-Marga. Distr. — Chile.
- TESSELATA Alexander, 1943i:749. Type-locality: Peru, Ayacucho. La Mar. Ayna. Distr. — Peru.
- THIOSEMA Alexander, 1921c:164 [*Gonomyia* (*Gonomyella*)]. Type-locality: Argentina, Tucumán, Famailla. Distr. — Argentina.
- VELUTINA Alexander, 1916a:18, pl. 2, figs. 9, 10 (wing). Type-locality: Peru. Lima, Matucana. Distr. — Peru.
- WEISERI Alexander, 1920f:215 [*Gonomyia* (*Gonomyella*)]. Type-locality: Argentina, Jujuy, Cueva Iturbe. Distr. — Argentina.

Genus GYMNASTES Brunetti

- GYMNASTES Brunetti, 1911:281. Type-species, *violacea* Brunetti (mon.) = *cyaneus* (Edwards).

Subgenus PARAGYMNASTES Alexander

- PARAGYMNASTES Alexander, 1922g:583 (as genus). Type-species, *Limnobia fascipennis* Thomson (orig. des.).



PEREXQUISITA Alexander, 1938f:256. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Refs. — Alexander, 1939c:439, figs. 8 (wing), 14 (male hypopygium); 1947b:81, fig. 14 (wing).

Genus JIVAROMYIA Alexander

JIVAROMYIA Alexander, 1943h:493. Type-species, *problematica* Alexander (orig. des.).

PROBLEMATICA Alexander, 1943h:494, figs. 3 (wing), 10 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador. Ref. — Alexander, 1947b:81, fig. 15 (wing).

Genus LIPSOTHRIX Loew

LIPSOTHRIX Loew, 1873:67, 69. Type-species, *nobilis* Loew (Alexander, 1965d:77) (Western Palaearctic).

NEOTROPICA Alexander, 1940c:114. Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama. Ref. — Alexander, 1947b:99, fig. 31 (wing).

Genus MAIETTA Alexander

MAIETTA Alexander, 1929a:184. Type-species, *squamigera* Alexander (orig. des.). Ref. — Alexander, 1947c:338.

SQUAMIGERA Alexander, 1929a:185, figs. 104 (wing), 233 (male hypopygium), 234 (ovipositor). Type-locality: Chile, Mechuque Is. Distr. — Chile.

TRIMEDIA Alexander, 1966c:493. Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chilc. Ref. — Alexander, 1968a:30, figs. 19 (wing), 20 (male hypopygium).

Genus MOLOPHILUS Curtis

MOLOPHILUS Curtis, 1833: pl. 444. Type-species, *brevipennis* Curtis (mon.) = *ater* (Melgen). [See note in Alexander, 1965d:88].

Archimolophilus Enderlein, 1940:669. Type-species, *selkirkianus* Enderlein (mon.). Ref. — Alexander, 1947c:338, figs.

Subgenus EUMOLOPHILUS Alexander


Molophilus, subg. EUMOLOPHILUS Alexander, 1921d:72. Type-species, *thau-mastopodus* Alexander (orig. des.).

ANGUSTIOR Alexander, 1936a:21. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.

PENNIPES Alexander, 1921d:72. Type-locality: Brazil, ? Santa Felpe. Distr. — Brazil, Peru.

SABETHOIDES Edwards, 1927:45. Type-locality: Brazil, Mato Grosso, Chapada [dos Guimarães]. Distr. — Brazil.

THAUMASTOPODUS Alexander, 1913b:510, fig. 12 (wing). Type-locality: Brazil, Pará. Distr. — Brazil.



Subgenus *MOLOPHILUS* Curtis

- AMPHIACANTHUS* Alexander, 1952c:64, fig. 30 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Alto Francés. Distr. — Chile.
- ANERASTA* Alexander, 1934a:71. Type-locality: Panama, Chiriqui, Finca Palo Alto. Distr. — Panama.
- ANTIMENUS* Alexander, 1952c:65, fig. 31 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Quebrada de las Vacas. Distr. — Chile.
- APICIDENS* Alexander, 1951c:16, fig. 9 (male hypopygium). Type-locality: Chile, Malleco, Curacautín. Distr. — Chile.
- APPRESSOPINUS* Alexander, 1952c:65, fig. 32 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Alto Francés. Distr. — Chile.
- APPRESSUS* Alexander, 1929a:213, fig. 252 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Argentina.
- ARAUCANUS* Alexander, 1929a:219, fig. 259 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- ARCIFERUS* Alexander, 1952c:66, fig. 33 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Quebrada de las Vacas. Distr. — Chile.
- ARMATISTYLUS* Alexander, 1929a:204, fig. 240 (male hypopygium). Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Argentina, Chile.
- ASCENDENS* Alexander, 1945e:242, fig. 7 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- AVITUS* Alexander, 1952d:248. Type-locality: Ecuador, Tungurahua, Upper Río Topo. Distr. — Ecuador.
- AXILLISPINUS* Alexander, 1962c:155, fig. 136 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- BELlicosus* Alexander, 1929a:216, fig. 255 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina, Chile.
- BELLONA* Alexander, 1946c:539. Type-locality: Mexico, Morelos, Guernavaca, Harmsen's Place. Distr. — Mexico.
- BICAUDATUS* Alexander, 1929a:210, fig. 248 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Argentina.
- BIDIGITATUS* Alexander, 1931c:642. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- BIERIGI* Alexander, 1947c:344, fig. 20 (male hypopygium). Type-locality: Costa Rica, San Pedro de Montes de Oca. Distr. — Costa Rica.
- BINARIUS* Alexander, 1968a:31, fig. 21 (male hypopygium). Type-locality: Chile, Magallanes, Las Piedras. Distr. — Chile.
- BREVILOBATUS* Alexander, 1938i:281. Type-locality: Chile, Villa Portalcs, Pinares de Lonquimai. Distr. — Chile.
- BREVIRAMUS* Alexander, 1929a:208, fig. 244 (male hypopygium). Type-locality: Argentina, Rio Negro, Viedma. Distr. — Argentina.
- BREVISPINOSUS* Alexander, 1929a:205, fig. 239 (male hypopygium). Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Argentina, Chile.
- BROWNIANUS* Alexander, 1945a:37. Type-locality: Ecuador, Tungurahua, Minza Chica. Distr. — Ecuador. Ref. — Alexander, 1945L:393, fig. 22 (male hypopygium).



- BRUCHI Alexander, 1923b:182. Type-locality: Argentina, Palo Blanco, La Plata. Distr. — Argentina.
- CAENOSUS Alexander, 1936e:66. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CALCEATUS Alexander, 1929a:208, fig. 245 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- CANOPUS Alexander, 1952c:68, fig. 34 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Plazoleta del Yunque. Distr. — Chile.
- CAPRICORNIS Alexander, 1916a:19, pl. 1, fig. 9 (wing), pl. 4, fig. 3 (male hypopygium). Type-locality: Colombia, Magdalena, La Cumbre. Distr. — Colombia, Peru.
- CARPISHENSIS Alexander, 1949c:615, fig. 40 (male hypopygium). Type-locality: Peru, Huanuco, Carpish. Distr. — Peru, Bolivia. Ref. — Alexander, 1962e:153, fig. 138 (male hypopygium).
- CATAMARCENSIS Alexander, 1921e:292. Type-locality: Argentina, Catamarca, Punta de Balasto. Distr. — Argentina. Ref. — Alexander, 1952e:87, fig. 8 (male hypopygium).
- CAUTUS Alexander, 1947c:345, fig. 21 (male hypopygium). Type-locality: Brazil, São Paulo. Distr. — Brazil.
- CERVUS Alexander, 1929a:211, fig. 249 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Correntoso. Distr. — Argentina, Chile.
- CHIRIQUIENSIS Alexander, 1934a:72. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.
- CLADOCERUS Alexander, 1921c:163. Type-locality: Argentina, Tucumán, Arroyo Famaila. Distr. — Argentina.
- CLAVIGERUS Alexander, 1929a:219, fig. 258 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Gutiérrez. Distr. — Argentina.
- COLOSSUS Alexander, 1929a:215, figs. 109 (wing), 254 (male hypopygium). Type-locality: Argentina, Río Negro, Río Nirico. Distr. — Argentina, Chile.
- COMPACTUS Alexander, 1950a:55, figs. 33 (wing), 38 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- CONSCRIPTUS Alexander, 1938d:337. Type-locality: Colombia, Cundinamarca, Fusagasugá. Distr. — Colombia.
- DEBILIOR Alexander, 1943b:126. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- DEBILISTYLUS Alexander, 1937c:500. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DEFOEANUS Alexander, 1952e:69, fig. 35 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Inocentes Altos. Distr. — Chile.
- DICEROS Alexander, 1944i:134. Type-locality: Chile, Malleco, Caracautín, Termas Río Blanco. Distr. — Chile.
- DIDO Alexander, 1943d:138, fig. 4 (male hypopygium). Type-locality: Venezuela, Merida, Apartaderos. Distr. — Venezuela.
- DILATUS Alexander, 1951c:10, fig. 4 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.



- DIRHABDA Alexander, 1949b:496, fig. 26 (male hypopygium). Type-locality: Brazil, São Paulo, Campos do Jordão, Homem Morto. Distr. — Brazil.
- DIRUS Alexander, 1943a:242, fig. 5 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.
- DISTIFURCUS Alexander, 1952c:70, fig. 36 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Plazolcta del Yunque. Distr. — Chile.
- DIZYGUS Alexander, 1962c:156, fig. 137 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare. Distr. — Bolivia.
- DREPANUCHUS Alexander, 1929a:209, fig. 246 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- DUCTILIS Alexander, 1938e:408. Type-locality: Mexico, Chiapas, Finca Victoria. Distr. — Mexico.
- ECHO Alexander, 1952d:254. Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile. Ref. — Alexander, 1952b:114 (as 1951), fig. 13 (male hypopygium).
- EMARGINATUS Alexander, 1937d:501. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ERECTUS Alexander, 1953b:22, fig. 20 (male hypopygium). Type-locality: Chile, Chiloé Is., Aulén. Distr. — Chile.
- EUMONOSTYLUS Alexander, 1952b:112, fig. 10 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- EXEMPTUS Alexander, 1953b:19, fig. 18 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- FAGETORUM Alexander, 1929a:212, fig. 250 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile, Argentina.
- FALX Alexander, 1938e:410. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- FILIOLUS Alexander, 1952c:70, figs. 28 (ovipositor), 37 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- FILIUS Alexander, 1952c:71, figs. 29 (ovipositor), 38 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- FLAVEXEMPTUS Alexander, 1968a:31, fig. 22 (male hypopygium). Type-locality: Chile, Chaihuín. Distr. — Chile.
- FLAVIDUS Alexander, 1914b:205, pl. 9, fig. 2 (male hypopygium). Type-locality: Chile, Concepción. Distr. — Argentina, Chile. Ref. — Alexander, 1929a:202, fig. 236 (male hypopygium).
- FLEMINGI Alexander, 1950a:55, fig. 39 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- FLEXILISTYLUS Alexander, 1931c:639. Type-locality: Colombia, Magdalena, Mt. Santa Marta, La Cumbre. Distr. — Colombia.
- FLINTI Alexander, 1967c:380. Type-locality: Chile, Magallanes, Las Piedras. Distr. — Chile. Ref. — Alexander, 1968a:33, fig. 23 (male hypopygium).
- FORTIDENS Alexander, 1951c:12, fig. 6 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- FURCUS Alexander, 1951c:14, figs. 7, 7A (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.



- FUSCOLEURALIS Alexander, 1927d:317. Type-locality: Mexico, near summit of the Zontchuitz. Distr. — Mexico.
- GARGANTUA Alexander, 1941c:256. Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- GOMESI Alexander, 1942h:435, figs. 15 (wing), 19 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Ribeirão das Lajes. Distr. — Brazil.
- GRUS Alexander, 1941e:262, fig. 16 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.
- GUATEMALENSIS Alexander, 1913b:511. Type-locality: Guatemala (probably Guatemala City). Distr. — Guatemala.
- GYMNOCLADUS Alexander, 1928c:183. Type-locality: Chile, Concepción. Distr. — Chile.
- HAAGI Alexander, 1946b:245, figs. 21 (wing), 27 (male hypopygium). Type-locality: Mexico, Michoacán, Cerro Tancitaro. Distr. — Mexico.
- HONESTUS Alexander, 1923b:181. Type-locality: Argentina, Córdoba, La Granja, Alta Gracia. Distr. — Argentina.
- HYPERARMATUS Alexander, 1939f:183, fig. 22.3 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- HYSTRIX Alexander, 1939c:443, fig. 16 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- ILLECTUS Alexander, 1941g:355 (also 1946h:103). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- INARMATUS Alexander, 1939f:182, fig. 22.2 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- INCOGNITUS Alexander, 1946e:540. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- INDURABILIS Alexander, 1966c:494, fig. 18 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Caramavida. Distr. — Chile.
- INFLEXIBILIS Alexander, 1929a:203, fig. 238 (male hypopygium). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina.
- INTEGRISTYLUS Alexander, 1945j:37, fig. 16 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- ISOLATUS Alexander, 1952d:253. Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile. Ref. — Alexander, 1952b:115, fig. 14 (male hypopygium).
- JANEI Alexander, 1945c:276, fig. 21 (male hypopygium). Type-locality: Brazil, São Paulo, Amparo, Fazenda São Bento. Distr. — Brazil.
- LATEROSPINOSUS Alexander, 1929a:206, fig. 242 (male hypopygium). Type-locality: Argentina, Río Negro, Lake Gutiérrez. Distr. — Argentina.
- LAURI Alexander, 1945j:39, fig. 17 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- LEONURUS Alexander, 1951c:17, fig. 10 (male hypopygium). Type-locality: Chile, Malleco, Curacautín. Distr. — Chile.
- LERIONIS Alexander, 1945e:241, fig. 6 (male hypopygium). Type-locality: Brazil, Guanabara, Rio de Janeiro, Alto da Boa Vista. Distr. — Brazil.
- LETHAEUS Alexander, 1952b:115, fig. 15 (male hypopygium). Type-locality: Chile, Chiloé Is., Chepu. Distr. — Chile.
- LICTOR Alexander, 1938d:336. Type-locality: Colombia, Cundinamarca, Fugasugá. Distr. — Colombia.



- LUXURIOSUS Alexander, 1938d:335. Type-locality: Colombia, Usme. Distr. — Colombia.
- MAGELLANICUS Alexander, 1968a:33, fig. 24 (male hypopygium). Type-locality: Chile, Magallanes, Las Piedras. Distr. — Chile.
- MARTHAE Alexander, 1931c:641. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- MASAFUERA Alexander, 1952c:72, fig. 39 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Quebrada de las Vacas. Distr. — Chile.
- MELANOLEUCUS Alexander, 1952d:247. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- METUENDUS Alexander, 1951c:11, fig. 5 (male hypopygium). Type-locality: Chile, Malleco, Curacautín. Distr. — Chile.
- MIRACULUS Alexander, 1938c:408. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- MONOCTENUS Alexander, 1951c:15, fig. 8 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- MONOSTYLOIDES Alexander, 1951c:7, fig. 1 (male hypopygium). Type-locality: Chile, Malleco, Curacautín. Distr. — Chile.
- MONOSTYLUS Alexander, 1928c:182. Type-locality: Chile, Concepción. Distr. — Chile. Ref. — Alexander, 1947c:347, fig. 22 (male hypopygium).
- MULTIFIDUS Alexander, 1952c:73, figs. 30 (wing), 40 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Bahía Cumberland. Distr. — Chile.
- NAHUELBTAE Alexander, 1966c:496, fig. 19 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Caramavida. Distr. — Chile.
- NEODICEROS Alexander, 1966c:497, fig. 20 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Palo Botado, Contulmo. Distr. — Chile.
- NEOFACINUS Alexander, 1964b:283 (nom. nov. for *facinus* Alexander). Type-locality: Venezuela, Merida, Río Chama. Distr. — Venezuela.
facinus Alexander, 1943d:140, fig. 5 (male hypopygium) (preocc. Alexander, 1940). Type-locality: Venezuela, Merida, Río Chama.
- NEPTUNUS Alexander, 1952c:74, fig. 41 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- OBEDIENS Alexander, 1927d:316. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- OBLIVIOSUS Alexander, 1953b:20, fig. 19 (male hypopygium). Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- OLIGOTRICHUS Alexander, 1954a:43, fig. 9 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- ORION Alexander, 1914c:119, figs. 4 (wing), 9 (male hypopygium). Type-locality: Costa Rica, Alajuela. Distr. — Costa Rica.
- OTHELLO Alexander, 1941c:257, fig. 12 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpatata. Distr. — Peru, Bolivia.
- OZOTUS Alexander, 1968a:34, fig. 25 (male hypopygium). Type-locality: Chile, Río Ancoa. Distr. — Chile.
- PACIFER Alexander, 1947a:183. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica. Ref. — Alexander, 1947c:347, fig. 23 (male hypopygium).

- PAGANUS Alexander, 1942c:445, fig. 13 (male hypopygium). Type-locality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- PALA Alexander, 1941e:261, fig. 15 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru, Bolivia. Ref. — Alexander, 1962c:158, fig. 139 (male hypopygium).
- PALLATANGENSIS Alexander, 1929d:787. Type-locality: Ecuador, Chimborazo, Pallatanga. Distr. — Ecuador.
- PALLIDUS (Philippi), 1865:616 (*Erioptera?*). Type-locality: Chile, Aconcagua, Catemu. Distr. — Chile. (Identity uncertain).
- PANCHRESTUS Alexander, 1941e:260, fig. 14 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PANSUS Alexander, 1968a:34, fig. 26 (male hypopygium). Type-locality: Chile, Puerto Fui. Distr. — Chile.
- PARADICEROS Alexander, 1966c:498, fig. 21 (male hypopygium). Type-locality: Chile, Arauco, Nahuclbuta, Butamalal. Distr. — Chile.
- PARAGUAYENSIS Alexander, 1929d:787. Type-locality: Paraguay, Guaira, Villarrica. Distr. — Paraguay.
- PASTORIS Alexander, 1960:18. Type-locality: Chile, Chiloé Is., Aucar. Distr. — Chile.
- PAUCISPINOSUS Alexander, 1925a:351, fig. 20 (male hypopygium). Type-locality: Mexico, México, D.F., Desierto de los Leoncs. Distr. — Mexico.
- PECTINATUS Alexander, 1928d:220. Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile. Ref. — Alexander, 1947c:347, fig. 24 (male hypopygium).
- PECTINIFERUS Alexander, 1952c:74, fig. 42 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- PENAI Alexander, 1951c:8, fig. 2 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- PENICILLATUS Alexander, 1941e:264, fig. 17 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huasahuasi. Distr. — Peru.
- PENNATUS Alexander, 1927d:318. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- PERDEBILIS Alexander, 1943i:764, fig. 15 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PEREXTENSUS Alexander, 1951c:9, fig. 3 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- PERFIDUS Alexander, 1929a:214, fig. 253 (male hypopygium) Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina, Chile.
- PERSEUS Alexander, 1913c:201, pl. 6, fig. 4 (male hypopygium). Type-locality: Colombia, Cauca, Valle de las Papas. Distr. — Colombia.
- PERSINUOSUS Alexander, 1941c:259, fig. 13 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru. Ref. — Alexander, 1947b:339, fig. 19 (wing).
- PERTENUS Alexander, 1952d:250. Type-locality: Peru, Junín, Chanchamavo. Distr. — Peru.
- PHALLODONTUS Alexander, 1968a:35, fig. 27 (male hypopygium). Type-locality: Chile, Enco. Distr. — Chile.
- PHALLOSOMICUS Alexander, 1939c:442, fig. 15 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PIGER Alexander, 1942c:446, fig. 14 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru, Bolivia.

- PIRIONI PIRIONI Alexander, 1929a:217, figs. 110 (wing), 257 (male hypopygium). Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile, Argentina.
- ssp. OMISSUS Alexander, 1929a:218. Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile, Argentina.
- PLANITAS Alexander, 1952d:249. Type-locality: Bolivia, Cochabamba, Cristal Mayu. Distr. — Bolivia.
- PLATYPHALLUS Alexander, 1942e:261. Type-locality: Ecuador, Manabi, Palmar. Distr. — Ecuador.
- PRETIOSUS Alexander, 1929a:210, fig. 247 (male hypopygium). Type-locality: Chile, Chiloé Is., Castro. Distr. — Chile.
- PROCAX Alexander, 1946e:538. Type-locality: Mexico, Morelos, Cuernavaca. Distr. — Mexico.
- PULVINUS Alexander, 1943a:244. Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- PUSTULATUS Alexander, 1946a:139. Type-locality: Mexico, Chiapas, Nueva America. Distr. — Mexico.
- QUADRISTYLUS Alexander, 1922c:38. Type-locality: "Brazil". Distr. — Brazil.
- RAPTOR Alexander, 1947c:347, fig. 25 (male hypopygium). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil.
- RECTISPINUS Alexander, 1952c:75, fig. 43 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Miradero de Selkirk. Distr. — Chile.
- REMIGER Alexander, 1922c:39. Type-locality: "Brazil". Distr. — Brazil.
- RETORSUS Alexander, 1939a:88. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- RICHARDSI Alexander, 1929a:216, fig. 256 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile, Argentina.
- RUBIDITHORAX RUBIDITHORAX Alexander, 1929a:212, fig. 251 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Chile, Argentina.
- ssp. EXCAVATUS Alexander, 1952b:114. Type-locality: Chile, Malleco, Río Blanco. Distr. — Chile.
- SAGAX Alexander, 1938e:409. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- SAGITTARIUS Alexander, 1914b:207, pl. 9, fig. 4 (male hypopygium). Type-locality: Bolivia, Yungas de Coroico. Distr. — Peru, Bolivia.
- SCABRICORNIS Alexander, 1942i:69, fig. 38 (male hypopygium). Type-locality: Brazil, São Paulo, Cabeceiras do M'Boy Guaçu. Distr. — Brazil.
- SCHULTZEI Alexander, 1946b:246, fig. 28 (male hypopygium). Type-locality: Mexico, Guerrero. Distr. — Mexico.
- SELKIRKIANUS (Enderlein), 1940:669, fig. 40 (wing) (*Archimolophilus*). Type-locality: Chile, Juan Fernández, Masatierra. Distr. — Chile. Ref. — Alexander, 1952c:76.
- SERRULATUS Alexander, 1929a:207, fig. 243 (male hypopygium). Type-locality: Chile, Valparaiso, Marga-Marga. Distr. — Chile.
- SETOSISTYLUS Alexander, 1952d:251. Type-locality: Chile, Malleco, Curautín, Termas de Río Blanco. Distr. — Chile. Ref. — Alexander, 1952b:113, fig. 11 (male hypopygium).
- SEVERUS Alexander, 1925a:350, fig. 19 (male hypopygium). Type-locality: Mexico, México, D.F., Desierto de los Leones. Distr. — Mexico.

- SHANNONINUS Alexander, 1947a:182. Type-locality: Argentina, Tucumán. Distr. — Argentina.
- SICARIUS SICARIUS Alexander, 1929b:99. Type-locality: Peru, Junín, Tarma, Huacapistana, Río Tarma. Distr. — Peru. Ref. — Alexander, 1962c:153, fig. 140A (male hypopygium).
- ssp. PARTITUS Alexander, 1941e:265. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru. Ref. — Alexander, 1962c:153, 140B (male hypopygium).
- SPARSISPINUS Alexander, 1952d:252. Type-locality: Chile, Malleco, Curacautín. Distr. — Chile. Ref. — Alexander, 1952b:114, fig. 12 (male hypopygium).
- SPONSUS Alexander, 1954d:150. Type-locality: Bolivia, Cochabamba, Chapore, Locotal, El Limbo. Distr. — Bolivia.
- STENOPTERUS Alexander, 1935b:104, figs. 6 (wing), 7 (male hypopygium). Type-locality: Chile, Malleco, Angol. Distr. — Chile.
- STENORHABDA Alexander, 1960:20. Type-locality: Chile, Chiloé Is., Río Coluco. Distr. — Chile.
- STYLIFER Alexander, 1921e:292. Type-locality: Argentina, Catamarca, Ampajango. Distr. — Argentina.
- STYX Alexander, 1952c:76, fig. 44 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Inocentes Altos. Distr. — Chile.
- SUBAPPRESSUS Alexander, 1939f:181, fig. 22.1 (male hypopygium). Type-locality: Chile, Malleco, Curacautín, Termas de Río Blanco. Distr. — Chile.
- SUBEXEMPTUS Alexander, 1968a:95. Type-locality: Chile, Arauco, Nahuelbuta. Distr. — Chile. Ref. — Alexander, 1968b:33, fig. 28 (male hypopygium).
- SUBFALCATUS Alexander, 1940e:906. Type-locality Venezuela, Miranda, Río Chacaito. Distr. — Venezuela.
- SUBIRATUS Alexander, 1947c:348, fig. 26 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SUBLICTOR Alexander, 1939e:635, fig. 16 (male hypopygium). Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- SUBSAGAX Alexander, 1939a:87. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- SUBSTYLIFER Alexander, 1929d:786. Type-locality: Argentina, Catamarca, Bolsón. Distr. — Argentina.
- SUBTENEBRICOSUS Alexander, 1931c:640. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- TALAMANCENSIS Alexander, 1947c:350. Type-locality: Costa Rica, Talamanca, El Muerto. Distr. — Costa Rica.
- TAURUS Alexander, 1914b:206, pl. 9, fig. 1 (male hypopygium). Type-locality: Chile, O'Higgins, Rancagua. Distr. — Chile.
- TEHUELICHE Alexander, 1968a:36, fig. 29 (male hypopygium). Type-locality: Chile, Magallanes, Las Picdras. Distr. — Chile.
- TELERHABDA Alexander, 1946e:541. Type-locality: Mexico, Guanajuato, Irapuato. Distr. — Mexico.
- TENEBRICOSUS Alexander, 1916a:20, pl. 1, fig. 10 (wing), pl. 4, fig. 4 (male hypopygium). Type-locality: Colombia, Magdalena, Mt. Santa Marta, La Cumbre. Distr. — Colombia.
- TERGOSPINOSUS Alexander, 1966c:499, fig. 22 (male hypopygium). Type-locality: Chile, Arauco, Nahuelbuta, Pichinahuel. Distr. — Chile.



- TERNARIUS Alexander, 1929a:205, fig. 241 (male hypopygium). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Chile, Argentina.
- TETRACANTHUS Alexander, 1929a:203, fig. 237 (male hypopygium). Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Argentina.
- TITAN Alexander, 1928b:58. Type-locality: Chile, Valparaíso, Marga-Marga. Distr. — Chile.
- TRIDENS Alexander, 1952c:77, fig. 45 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Bahía Cumberland. Distr. — Chile.
- TRIDIGITATUS Alexander, 1947c:351, fig. 27 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- TRIFIBRA Alexander, 1954a:42, fig. 8 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- TRIPARCUS Alexander, 1944f:33, fig. 6 (male hypopygium). Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- TUCUMANUS Alexander, 1929a:97. Type-locality: Argentina, Tucumán, Parque Aconquija. Distr. — Bolivia, Argentina.
- UNIFORMIS (Blanchard), 1852:343 (*Erioptera*). Type-locality: Chile, Coquimbo, La Serena. Distr. — Chile. Identity unknown.
- VARIATUS Alexander, 1952c:78, fig. 46 (male hypopygium). Type-locality: Chile, Juan Fernández, Masafuera, Inocentes Bajos. Distr. — Chile.
- WALKERI Alexander, 1931c:640. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- YUNQUENSIS Alexander, 1952c:78, fig. 47 (male hypopygium). Type-locality: Chile, Juan Fernández, Masatierra, Plazoleta del Yunque. Distr. — Chile.

Subgenus RHYNCHOMOLOPHILUS Alexander

- Molophilus*, subg. RHYNCHOMOLOPHILUS Alexander, 1965a:407. Type-species, *perrostratus* Alexander (orig. des.).
- FERROSTRATUS Alexander, 1965a:407. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru. Ref. — Alexander, 1966c:499.

Subgenus TRICHOMOLOPHILUS Alexander


- Molophilus*, subg. TRICHOMOLOPHILUS Alexander, 1936d:768. Type-species, *multisetosus* Alexander (orig. des.). Ref. — Alexander, 1947c:344.
- CELATOR Alexander, 1942h:436, fig. 20 (male hypopygium). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- MULTISETOSUS Alexander, 1936d:768. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- TENTATOR Alexander, 1947c:352, fig. 28 (male hypopygium). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo, Sítio Bonfim, Serra dos Órgãos. Distr. — Brazil.

Genus NEOGNOPHOMYIA Alexander

- Gnophomyia*, subg. NEOGNOPHOMYIA Alexander, 1926c:391. Type-species, *Gnophomyia hirsuta* Alexander (orig. des.).
- Neognophomyia*; Alexander, 1947b:81, fig. 16.



- ADARA Alexander, 1949c:602, fig. 27 (male hypopygium). Type-locality: Ecuador, Tungurahua, Baños, Río Blanco. Distr. — Ecuador.
- RISECTA (Alexander), 1920a:72 (*Gnophomyia*). Type-locality: Brazil, Amazonas, Parintins. Distr. — Brazil.
- BISETOSA Alexander, 1944g:307, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- CITRIPES Alexander, 1945i:257. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- COCHLEARIS Alexander, 1945a:31. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- COLOMBICOLA Alexander, 1931c:636. Type-locality: Colombia, Magdalena, Mt. Santa Marta, Vista Nieve. Distr. — Colombia.
- CONSOCIATA Alexander, 1942c:259. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador.
- CRASSISTYLA Alexander, 1966c:483. Type-locality: Peru, Cuzco, Quincemil along the Marcapata River. Distr. — Peru.
- CUZCOENSIS Alexander, 1966c:484. Type-locality: Peru, Cuzco, Quincemil along the Marcapata River. Distr. — Peru.
- DEBILITATA Alexander, 1949c:604, fig. 28 (male hypopygium). Type-locality: Peru, Loreto, Contamana. Distr. — Peru.
- HELMCONIAE Alexander, 1945b:57. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- HIRSUTA (Alexander), 1913b:523, fig. 30 (wing) (*Gnophomyia*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil.
- HOSTICA Alexander, 1943i:760, fig. 12 (male hypopygium). Type-locality: Peru, Junín, Tarma, Tulumayo Valley. Distr. — Peru.
- IMMACULIPENNIS (Alexander), 1926c:392 [*Gnophomyia* (*Neognophomyia*)]. Type-locality: Paraguay, San Bernardino. Distr. — Brazil, Paraguay.
- INTERRUPTA Alexander, 1944a:52. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- LATIFASCIA (Alexander), 1926c:392 [*Gnophomyia* (*Neognophomyia*)]. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- MONOPHORA Alexander, 1941b:146. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- OBTUSILAMINA Alexander, 1952a:131. Type-locality: Peru, Huanuco, Fundo Sinchono, Cordillera Azul. Distr. — Peru.
- PANAMENSIS Alexander, 1930b:116. Type-locality: Panama, Tonosí, Los Santos. Distr. — Panama.
- PAPRZYCKIANA Alexander, 1944g:306, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru. Ref. — Alexander, 1962c:139, fig. 116 (wing).
- PERVICAX (Alexander), 1914b:208, pl. 9, fig. 7 (wing) (*Gnophomyia*). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- FINCKERTI Alexander, 1962c:139, figs. 117 (wing), 120 (male hypopygium). Type-locality: Bolivia, Santa Cruz, General Saavedra. Distr. — Bolivia.
- PRODUCTISSIMA Alexander, 1944h:382. Type-locality: Costa Rica, Talamanca, Pedregoso. Distr. — Costa Rica.
- SCAPHA Alexander, 1945a:33. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.



- SCAPHOIDES Alexander, 1952a:131. Type-locality: Peru, Huanuco, Fundo Sinchono, Cordillera Azul. Distr. — Peru.
- SCHILDI Alexander, 1945h:288. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica. Ref. — Alexander, 1947b:81, fig. 16 (wing).
- SETLOBATA Alexander, 1949c:605, fig. 29 (male hypopygium). Type-locality: Peru, Loreto, Cerro Azul. Distr. — Peru.
- SPARSISETA Alexander, 1945i:256. Type-locality: Peru, Junin, Satipo, Jauja. Distr. — Peru.
- SPECTRALIS Alexander, 1944a:51. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- TRINITATIS Alexander, 1927c:277. Type-locality: Trinidad, Port of Spain. Distr. — Trinidad.

Genus NEOLIMNOPHILA Alexander

- Limnophila*, subg. NEOLIMNOPHILA Alexander, 1920h:37. Type-species, *ultima* Osten Sacken (orig. des.) (Nearctic).
- Neolimnophila*; Alexander, 1965d:72.
Ref. — Alexander, 1947b:66.
- ANDICOLA Alexander, 1942c:424. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.

Genus NEOPHILIPPIANA Alexander

- Philippiana* Alexander, 1929a:175 (preocc. Wagner, 1897). Type-species, *egregia* Alexander (orig. des.).
- NEOPHILIPPIANA Alexander, 1964b:282 (nom. nov. for *Philippiana* Alexander). Type-species, *Philippiana egregia* Alexander (aut.).
- EGREGIA (Alexander), 1929a:177, pl. 5, fig. 97 (wing), pl. 10, figs. 216-218 (male hypopygium) (*Philippiana*). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Chile, Argentina. Ref. — Alexander, 1947b:67, fig. (wing).

Genus QUECHUAMYIA Alexander

- QUECHUAMYIA Alexander, 1943h:495. Type-species, *phantasma* Alexander (orig. des.).
- PHANTASMA Alexander, 1943h:496, figs. 6-9 (wing), 11 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.

Genus RHABDOMASTIX Skuse

- RHABDOMASTIX Skuse, 1890a:828. Type-species, *ostensackeni* Skuse (mon.) (Australasian). Ref. — Alexander, 1947c:318, fig. 1 (wing).

Subgenus RHABDOMASTIX Skuse

- ALTICOLA Alexander, 1944g:310. Type-locality: Peru, Ayacucho, La Mar, Yanamonte. Distr. — Peru.



- CHILOTA Alexander, 1929a:183, fig. 102 (wing). Type-locality: Chile, Chiloé, Is., Ancud. Distr. — Chile.
- ILLUDENS Alexander, 1914b:210, pl. 9, fig. 6 (venation). Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia.
- ISABELLA Alexander, 1927d:315. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico.
- LONGITEREBRATA Alexander, 1938e:405. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- LUTEOLA Alexander, 1944g:308. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- MEXICANA Alexander, 1938e:404. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- PERGLABRATA Alexander, 1962c:152, fig. 134 (wing). Type-locality: Bolivia, Santa Cruz, General Saavedra. Distr. — Bolivia.
- PERUVIANA Alexander, 1926d:387. Type-locality: Peru, Lima. Distr. — Peru.
- POSTICATA Alexander, 1929a:183, figs. 103 (wing), 222 (male hypopygium). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- SATIOENSIS Alexander, 1944g:309. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- SEPTEMTRIONALIS Alexander, 1914c:119, pl. 2, fig. 3 (wing). Type-locality: Costa Rica, Alajuela. Distr. — Costa Rica.
- TANTILLA Alexander, 1938e:440. Type-locality: Colombia, Meta, Restrepo. Distr. — Colombia.
- UNIPUNCTA Alexander, 1944j:158, fig. 3 (wing). Type-locality: Venezuela, Trujillo. Distr. — Venezuela.

Subgenus SACANDAGA Alexander

- SACANDAGA Alexander, 1911b:349 (as genus). Type-species, *flava* Alexander (orig. des.) (Nearctic).
- CAPARAOENSIS Alexander, 1944f:32, fig. 3 (wing). Type-locality: Brazil, Minas Gerais, Alto Caparaó. Distr. — Brazil.
- FUMIPENNIS Alexander, 1939d:99, fig. 9 (wing). Type-locality: Lesser Antilles: Dominica, Sylvania Estate. Distr. — Dominica.
- INTERMEDIA Alexander, 1929a:182, fig. 101 (wing). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Chile, Argentina.
- PARVULA Alexander, 1947c:321 (nom. nov. for *parva* Alexander). Type-locality: Jamaica, Blue Mts., Yallahs Valley. Distr. — Dominican Republic, Jamaica.
- parva* Alexander, 1913b:508, fig. 9 (wing) (*Sacandaga*; preocc. Siebke, 1863). Type-locality: Jamaica, Blue Mts., Yallahs Valley. Ref. — Alexander, 1964a:74, figs. 75 (wing), 76 (male hypopygium).
- PLAUMANNI Alexander, 1947c:321, figs. 1 (wing), 2 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SYNCLERA Alexander, 1928d:219. Type-locality: Chile, Valparaiso, Margamarga. Distr. — Chile.

Genus SIGMATOMERA Osten Sacken

- SIGMATOMERA Osten Sacken, 1869:137. Type-species, *flavipennis* [sub. mon., Osten Sacken in Loew 1873:ix (additions to vol. IV)]. Refs. — Alexander, 1930a (biology); 1947b:68.



Subgenus AUSTROLIMNOBIA Alexander

- AUSTROLIMNOBIA Alexander, 1922e:239 (as genus). Type-species, *spectabilis* Alexander (orig. des.) (Australian).
- Gnophomyia*, subg. *Astelobia* Edwards, 1923:297. Type-species, *Tipula rufa* Hudson (orig. des.) (New Zealand).
- BULLOCKI (Alexander), 1935b:103, fig. 5 (wing) (*Astelobia*). Type-locality: Chile, Malleco, Angol. Distr. — Chile.
- MAGNIFICA (Alexander), 1913b:522, fig. 31 (wing) (*Gnophomyia*). Type-locality: Mexico, Morelos, Cuernava. Distr. — Mexico.
- MAIAE (Alexander), 1929a:178, figs. 89 (wing), 219 (male hypopygium) (*Astelobia*). Type-locality: Chile, Chiloé Is., Ancud. Distr. — Chile.
- PLAUMANNIANA Alexander, 1938a:32 [*Sigmatomera* (*Astelobia*)]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- WOYTKOWSKIANA Alexander, 1941e:241. Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.

Subgenus EUFURINA Alexander

- Furina* Jaennickc, 1867:318 (preocc. Duméril, 1853). Type-species, *Limnobia rufithorax* Wiedemann (mon.).
- Sigmatomera*, subg. EUFURINA Alexander, 1946f:307 (nom. nov. for *Furina* Jaennicke). Type-species, *Limnobia rufithorax* Wiedemann (aut.).
- RUFITHORAX (Wiedemann), 1828:548 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil, Bolivia. Ref. — Alexander, 1962c:136, figs. 114 (wing), 118 (male hypopygium).

Subgenus SIGMATOMERA Osten Sacken

- AEQUINOCTIALIS Alexander, 1937c:496. Type-locality: Ecuador, Napo-Pastaza, Hdaila(?), Río Anzu (= R. Ansupi). Distr. — Ecuador.
- AMAZONICA Westwood, 1881:366, figs. 3a (entire insect), 3b (head), 3c (wing), 3d (leg). Type-locality: Brazil, Amazonia. Distr. — Brazil. Refs. — Alexander, 1920c:142; 1930a:158, figs. 2 (ovipositor), 3 (wing).
- ANCUSTIROSTRIS Alexander, 1947b:69, fig. 4 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- APICALIS Alexander, 1914e:246, pl. 3, fig. 9 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana. Ref. — Alexander, 1930a:158, fig. 5 (wing).
- BEEBEI Alexander, 1950a:52, figs. 28 (wing), 34 (male hypopygium). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- FELIX Alexander, 1956b:120. Type-locality: Peru, Cuzco, Santa Isabel. Distr. — Peru.
- FLAVIPENNIS Osten Sacken in Loew, 1873:ix [additions to vol. IV]. Type-locality: "Mexico". Distr. — Mexico. Refs. — Alexander, 1930a:158, fig. 8 (wing); 1947b:67, fig. 3 (wing).
- CEJSKESANA Alexander, 1946f:305. Type-locality: Surinam, Paramaribo. Distr. — Surinam.
- CCULTA Alexander, 1914b:209, pl. 9, fig. 5 (wing). Type-locality: Paraguay, Asunción, Villa Morra. Distr. — Brazil, Paraguay. Refs.



- Alexander, 1930a:158, fig. 7 (wing); Riedel, 1921. biology (as *flavipennis*).
- PICIPENNIS Alexander, 1937b:243. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- SEGUYI Alexander, 1930a:159, fig. 6 (wing). Type-locality: "Costa Rica". Distr. — Costa Rica.
- SHANNONIANA Alexander, 1930a:158, figs. 1 (ovipositor), 4 (wing). Type-locality: Brazil, Bahia. Distr. — Brazil. Ref. — Alexander, 1930a (biology).
- VARICORNIS Alexander, 1936a:18. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.

Genus STYRINGOMYIA Loew

- STYRINGOMYIA Loew, 1845:6. Type-species, *venusta* Loew (fossil in copal) (mon.).
- Styringia* Berendt, 1845:57, misspell. Ref. — Alexander, 1947c:354.
- AMERICANA Alexander, 1914e:231, pl. 4, fig. 1 (wing). Type-locality: Guyana, Mallali. Distr. — British Honduras, Costa Rica, Colombia, Venezuela, Guyana, Surinam, Ecuador, Bolivia. Ref. — Alexander, 1945i:241, fig. 8 (male hypopygium).
- DORSOLINEATA Alexander, 1945n:419. Type-locality: Ecuador, El Oro, Morro Morro. Distr. — Ecuador.
- MYSTICA Alexander, 1945i:261, fig. 7 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru. Ref. — Alexander, 1947c:355, fig. 29 (wing).
- PAULISTA Alexander, 1946c:7, figs. 5 (wing), 8 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- SIMPLEX Alexander, 1945i:263, fig. 9 (male hypopygium). Type-locality: Guyana, Bartica. Distr. — Guyana, Peru, Bolivia.

Genus TASIOCERA Skuse

- TASIOCERA Skuse, 1890a:815. Type-species, *tenuicornis* Skuse (Alexander, 1920g:52) (Australasian).

Subgenus DASYMOLOPHILUS Goetghebuer


- Molophilus*, subg. DASYMOLOPHILUS Goetghebuer in Goetghebuer & Tonnoir, 1920:132. Type-species, *Erioptera murina* Meigen (mon.) (Western Palaearctic).
- BORACEAE Alexander, 1954a:41, fig. 7 (male hypopygium). Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil.
- BREVICORNIS Alexander, 1929a:198, figs. 108 (wing), 235 (male hypopygium). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.

Genus TEUCHOLABIS Osten Sacken

- TEUCHOLABIS Osten Sacken, 1859:222. Type-species, *complexa* Osten Sacken (mon.) (Nearctic). Refs. — Alexander, 1927c:17; 1946g:375.

Subgenus EUPARATROPESA Alexander

- Teucholabis*, subg. EUPARATROPESA Alexander, 1946g:378. Type-species, *Paratropesa jactans* Alexander (orig. des.).



- AMATRIX Alexander, 1945c:268, fig. 13 (wing) [*Teucholabis (Paratropesa)*]. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- AMOENA (Alexander), 1922c:47 (*Paratropesa*). Type-locality: "Venezuela". Distr. — Venezuela.
- ANGUSTIFASCIA Alexander, 1968c:217. Type-locality: Peru, Cuzco, Quince-mil. Distr. — Peru.
- FASCIOLARIS (Wiedemann), 1828:552 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil.
- FUMIDAPICALIS FUMIDAPICALIS Alexander, 1942c:425, fig. 7 (wing) [*Teucholabis (Paratropesa)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- ssp. LAETABILIS Alexander, 1952a:123 [*Teucholabis (Paratropesa)*]. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- IIETEROPODA Alexander, 1943a:227 [*Teucholabis (Paratropesa)*]. Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Peru.
- INVENUSTA Alexander, 1946g:382, fig. 1 (wing). Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica.
- JACTANS (Alexander), 1913e:447, fig. 10 (wing) (*Paratropesa*). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.
- LAETIFICA Alexander, 1948b:139. Type-locality: Peru, Loreto, Yarinacocha, near Pucallpa on Upper Ucayali River. Distr. — Peru.
- LINDNERI Alexander, 1933:42, figs. 1 (wing), 2 (male hypopygium) [*Teucholabis (Paratropesa)*]. Type-locality: Bolivia, Santa Cruz, El Cairo. Distr. — Bolivia.
- MARTINEZI Alexander, 1962c:141, figs. 121 (wing), 124 (male hypopygium). Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- PRAENOBILIS Alexander, 1945i:246 [*Teucholabis (Paratropesa)*]. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- SANGUIOLENTA Alexander, 1938a:35 [*Teucholabis (Paratropesa)*]. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- XYSTOPHANES (Alexander), 1921c:165 (*Paratropesa*). Type-locality: Argentina, Tucumán, Quebrada Famaila. Distr. — Argentina.

Subgenus EUTEUCHOLABIS Alexander

- Teucholabis*, subg. EUTEUCHOLABIS Alexander, 1946g:378. Type-species, *paradoxa* Alexander (orig. des.).
- NEPENTHE Alexander, 1946g:384, fig. 3 (wing). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- PARADOXA Alexander, 1913e:445, pl. 16, fig. 8 (wing). Type-locality: Peru, Callanga. Distr. — Peru.

Subgenus PARATROPESA Schiner

- PARATROPESA Schiner, 1866:932. Type-species, *singularis* Schiner (orig. des.).
- CHALYBEIA Alexander, 1927e:18. Type-locality: Colombia, El Líbano. Distr. — Colombia.
- COLLARIS (Osten Sacken), 1887b:190 (*Paratropesa*). Type-locality: "Brazil". Distr. — Brazil, Peru. Ref. — Alexander, 1913e:447, pl. 16, fig. 9 (wing).
- LANEANA Alexander, 1959:447, fig. 1 (male hypopygium). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.



- NEOCOLLARIS Alexander, 1944k:606. Type-locality: Ecuador, Santiago-Zamora, Rio Zamora, Zumbi. Distr. — Ecuador.
- NIGROCOXALIS Alexander, 1936d:764. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NODULIFERA Alexander, 1949a:103. Type-locality: Peru, Huanuco, Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:593, fig. 1 (male hypopygium).
- PARACOLLARIS Alexander, 1945i:248. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- PLACABILIS Alexander, 1941c:335, fig. 4 (wing). Type-locality: Costa Rica, Talamanca, Pedregoso, Valley of the General. Distr. — Costa Rica.
- PRAEUSTA (Osten Sacken), 1886a:8 (*Paratropesa*). Type-locality: Mexico, Jalapa. Distr. — Mexico. Ref. — Alexander, 1927e:19.
- SINGULARIS (Schiner), 1868:46 (*Paratropesa*). Type-locality: Venezuela (as "Colombia"). Distr. — Venezuela. Ref. — Alexander, 1927e:17.
- SUBCOLLARIS Alexander, 1940a:288. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador.

Subgenus TEUCHOLABIS Osten Sacken

- AEQUINIGRA Alexander, 1941g:351 (also 1946h:98) Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- AMAPANA Alexander, 1959:449, fig. 2 (male hypopygium). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.
- AMBLYPIHALLOS Alexander, 1966b:695. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- ANALIS Alexander, 1954d:146. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- ANGUSTAPICALIS Alexander, 1938c:329. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- ANNULATA Williston, 1896a:290. Type-locality: Lesser Antilles: St. Vincent. Distr. — Guyana, St. Vincent.
- ANTHRACINA Alexander, 1921d:85. Type-locality: Peru, Loreto, Napo River. Distr. — Venezuela, Ecuador, Peru. Ref. — Alexander, 1949c:593, fig. 2 (male hypopygium).
- ARGENTEA Alexander, 1927e:21. Type-locality: Brazil, Santa Catarina, Blumenau. Distr. — Brazil.
- ATAHUALPA Alexander, 1968a:25, fig. 12 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- ATRIPENNIS Alexander, 1940a:289. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Rio Pastaza. Distr. — Ecuador.
- ATROLATA Alexander, 1948b:140. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- AUDAX Alexander, 1913a:44, fig. d (wing). Type-locality: Panama, Canal Zone. Distr. — Panama.
- AZUAYENSIS Alexander, 1945p:764. Type-locality: Ecuador, Azuay, Cuenca. Distr. — Ecuador.




- BIACIFERA Alexander, 1949b:486, figs. 17 (wing), 21 (male hyogygium). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- BIDENTIFERA BIDENTIFERA Alexander, 1966b:696. Type-locality: Honduras. Lancetilla. Distr. — Honduras.
- ssp. MELANOPHALLUS Alexander, 1968b:92. Type-locality: Ecuador, Balao Chico, Río Frío. Distr. — Ecuador.
- BIGLADIUS Alexander 1942c:429, fig. 2 (male hypopygium). Type-locality: Peru, Huanuco, Afilador. Distr. — Peru.
- BIRAMOSA Alexander, 1940a:291. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Río Pastaza. Distr. — Ecuador.
- BREVISETOSA Alexander, 1941e:246, figs. 3 (wing), 6 (male hypopygium). Type-locality: Peru, Junín, Tarma, Carpapata. Distr. — Peru.
- BRUNERI Alexander. 1926b:227. Type-locality: Cuba, Oriente, Sierra Macstra, Pico Turquino. Distr. — Cuba.
- CATHARINENSIS Alexander, 1931a:117. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- CHALYBEIVENTRIS (Loew), 1861:33 (*Rhamphidia*). Type-locality: "Cuba". Distr. — Cuba, Jamaica. Ref. — Alexander, 1964a:69, figs. 70 (wing), 77 (male hypopygium).
- CINEREICEPS Alexander, 1928a:115. Type-locality: Mexico, Pena Blanca. Distr. — Mexico.
- CIRCUMSCRIPTA Alexander, 1945h:279. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- COCKERELLAE Alexander, 1915b:442, fig. 2 (wing). Type-locality: Guatemala, Quirigua. Distr. — Mexico, Guatemala, Costa Rica, Venezuela.
- COLOMELANIA Alexander, 1968b:91. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- CUNEIFORMIS Alexander, 1947a:178. Type-locality: Guyana, Bartica. Distr. Guyana.
- CYBELE Alexander, 1954d:147. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- DAMPFI Alexander, 1927d:308. Type-locality: Mexico, summit of one of the mountains of the Moshbiquil Range. Distr. — Mexico.
- DECORA Alexander, 1920a:71. Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Peru, Brazil.
- DEDECORA Alexander, 1936d:766. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DENUDA Alexander, 1943h:492, fig. 4 (male hypopygium). Type-locality: Ecuador, El Oro, Morro Morro, Las Pinas. Distr. — Ecuador.
- DESDEMONA Alexander, 1949a:106. Type-locality: Peru, Loreto, Pucallpa. Distr. — Peru. Ref. — Alexander, 1949c:593, fig. 4 (male hypopygium).
- DIANA Alexander, 1944k:608. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- DILEUCA Alexander, 1947a:179. Type-locality: Guatemala, Quirigua. Distr. Guatemala.
- DIFLACA Alexander, 1945h:281. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Panama.
- DISTIFURCA Alexander, 1943b:125. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

- DUCALIS Alexander, 1949a:104. Type-locality: Peru, Huanuco, Sinchona. Distr. — Peru, Bolivia. Ref. — Alexander, 1949c:593, fig. 3 (male hypopygium).
- DUIDENSIS Alexander, 1931b:15, fig. 10 (wing). Type-locality: Venezuela, Mt. Duida. Distr. — Venezuela.
- EGENS Alexander, 1952a:124. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- ELISSA Alexander, 1948b:144. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.
- FLAVITHORAX (Wiedemann), 1821:43 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil, Peru. Refs. — Alexander, 1927e:25; Enderlein, 1912:70, fig. S1 (wing).
- FLAVOFIMBRIA Alexander, 1968b:92. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- FOERSTERI Alexander, 1952e:91, figs. 6 (wing), 8 (male hypopygium). Type-locality: Argentina, Misiones. Distr. — Argentina.
- FORMOSISSIMA Alexander, 1946g:385, fig. 2 (wing). Type-locality: Costa Rica, Turrialba. Distr. — Costa Rica.
- FULGENS Alexander, 1913e:440, pl. 16, fig. 2 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- FULVINOTA Alexander, 1962c:143, fig. 126 (male hypopygium) Type-locality: Bolivia, Espíritu, Río Yacuma. Distr. — Bolivia.
- FURVA Alexander, 1930b:112. Type-locality: Guatemala, San José, Lago Petén. Distr. — Guatemala.
- FUSCOAPICALIS Alexander, 1937d:612. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- GALATEA Alexander, 1949b:484, figs. 16 (wing), 20 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- GOWDEYI GOWDEYI Alexander, 1928g:25, fig. 1 (male hypopygium). Type-locality: Jamaica, Hope Gardens. Distr. — Cuba, Jamaica. Ref. — Alexander, 1964a:70, fig. 71 (wing).
- ssp. NIGROTERRMINALIS Alexander, 1932:386. Type-locality: Cuba, Pinar del Río, Sierra Rangel. Distr. — Cuba.
- GRACILIS Osten Sacken, 1886a:7. Type-locality: Mexico, Orizaba. Distr. — Mexico, Panama.
- buscki Alexander, 1915b:442, fig. 3 (wing). Type-locality: Panama, Porto Bello.
- HERA Alexander, 1949d:260. Type-locality: Peru, Junín, Tarma, Utcuyacu. Distr. — Peru. Ref. — Alexander, 1949c:593, fig. 5 (male hypopygium).
- HILARIS Alexander, 1913e:443, pl. 16, fig. 6 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- HOMILACANTHA Alexander, 1968c:218. Type-locality: Ecuador, San Mateo, Esmeraldas. Distr. — Ecuador.
- HONDURENSIS Alexander, 1941c:336, figs. 5 (wing), 7 (male hypopygium). Type-locality: Honduras, Minas de Oro. Distr. — Honduras.
- HYPOMELA Alexander, 1948b:142. Type-locality: Peru, Yahuar Mayo. Distr. — Peru.
- IDIOPHALLUS Alexander, 1942c:435, fig. 6 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- IMMACULIPLURA Alexander, 1946g:399, fig. 7 (wing). Type-locality: Puerto Rico, Villalba. Distr. — Puerto Rico.
- INCA Alexander, 1968a:26, fig. 13 (male hypopygium). Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.



- INEPTA INEPTA Alexander, 1942c:434, fig. 5 (male hypopygium). Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- ssp. BISETOSA Alexander, 1945a:31. Type-locality: Ecuador, Napo-Pastaza, Río Huagra Yacu. Distr. — Ecuador.
- INULTA Alexander, 1936d:765. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INVARIPES Alexander, 1936a:19. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- JALISCANA Alexander, 1946g:394, fig. 11 (male hypopygium). Type-locality: Mexico, Santa Cruz, Jalisco, near Zapotiltéc. Distr. — Mexico.
- JIVARO Alexander, 1944k:610. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- JOCOSA Alexander, 1913e:440, pl. 16, fig. 3 (wing). Type-locality: Bolivia, Yungas de Coroico. Distr. — Colombia, Venezuela, Ecuador, Peru, Brazil, Bolivia, Argentina.
- JUCUNDA Alexander, 1913e:441, pl. 16, fig. 4 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- LAETA Alexander, 1913e:442, pl. 16, fig. 5 (wing). Type-locality: Bolivia, Cuticucho, Yungas de Songo Valley. Distr. — Peru, Bolivia.
- LADIS Alexander, 1966b:697. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- LAIS Alexander, 1946g:397, fig. 13 (male hypopygium). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- LATEROSPINOSA Alexander, 1937d:615. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LATIBASALIS Alexander, 1968b:93. Type-locality: Ecuador, Tena, Libertad. Distr. — Ecuador.
- LAUTA Alexander, 1949d:256. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru, Bolivia. Ref. — Alexander, 1949c:593, fig. 6 (male hypopygium).
- LAXA Alexander, 1946g:359, figs. 6 (wing), 12 (male hypopygium). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- LEONORA Alexander, 1938c:328. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- LERIDENSIS Alexander, 1934a:69. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.
- LETHE Alexander, 1946g:388, figs. 4 (wing), 9 (male hypopygium). Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- LINEPLEURA Alexander, 1949d:257. Type-locality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru. Ref. — Alexander, 1949c:594, fig. 7 (male hypopygium).
- LIPONEURA Alexander, 1937d:610. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LIPOPHLEPS Alexander, 1937d:611. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LONGISETOSA Alexander, 1941e:244, figs. 2 (wing), 5 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- LONGISPINA Alexander, 1938b:357. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- LUDICRA Alexander, 1949d:254. Type-locality: Peru, Loreto, Cerro Azul. Distr. — Peru. Ref. — Alexander, 1949c:594, fig. 8 (male hypopygium).
- LUGUBRIS Alexander, 1914e:234, pl. 4, fig. 2 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana.

- LUTEICOLOR Alexander, 1942c:431, fig. 3 (male hypopygium). Type-locality: Peru, Huanuco, Afilador. Distr. — Ecuador, Peru.
- MANNIANA Alexander, 1946g:390, fig. 5 (wing). Type-locality: Honduras Lombardía. Distr. — Honduras.
- MARTICOLA Alexander, 1931c:633. Type-locality: Colombia, Magdalena, Mt. Santa Marta. Distr. — Colombia.
- MEGAPHALLUS Alexander, 1968c:219. Type-locality: Colombia, Mount Rondono. Distr. — Colombia.
- MELANOCEPHALA (Fabricius), 1794:241 (*Tipula*). Type-locality: "South America". Distr. — Guyana, French Guiana, Peru, Brazil. Ref. — Alexander, 1914e:233, pl. 3, fig. 7 (wing).
- MENDAX Alexander, 1920b:5. Type-locality: Brazil, Pará, Prata. Distr. — Surinam, Peru, Brazil.
- MINIATA Alexander, 1930b:114. Type-locality: Panama, Changuinola. District. Distr. — Panama.
- MINUTA Alexander, 1926a:168, fig. 6 (wing). Type-locality: Mexico, Veracruz. Distr. — Mexico.
- MOLESTA Osten Sacken, 1886a:6. Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- MORIONELLA (Schiner), 1868:47 (*Paratropesa*). Type-locality: "South America." Distr. — Venezuela. Refs. — Alexander, 1927e:22; 1944j:154.
- MULTISPINOSA Alexander, 1944j:154, fig. 5 (male hypopygium). Type-locality: Venezuela, East Bolivar, Paraitepuy. Distr. — Venezuela.
- MUNDA Alexander, 1913e:444, pl. 16, fig. 7 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- MYERSI Alexander, 1926b:228. Type-locality: Cuba, Soledad. Distr. — U.S.A. (Florida), Cuba.
- MYTHICA Alexander, 1926d:393. Type-locality: Paraguay, Santa Barbara. Distr. — Paraguay.
- NEBULIPENNIS Alexander, 1928g:27. Type-locality: Jamaica, St. Andrew. Cinchona. Distr. — Jamaica. Ref. — Alexander, 1964a:70, fig. 72 (wing).
- NEOLERIDENSIS Alexander, 1940a:292. Type-locality: Ecuador, Tungurahua, Baños. Distr. — Ecuador.
- NEOSALVA Alexander, 1943a:229. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- NIGROCLAVARIA Alexander, 1946a:128. Type-locality: Mexico, Chiapas, Zapote. Distr. — Mexico.
- NIGROCORPORIS Alexander, 1937d:613. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NIGROCOSTATA Alexander, 1939c:438, figs. 6 (wing), 12 (male hypopygium). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NIGROPOSTICA Alexander, 1938b:358. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- NIGROSIGNATA Alexander, 1931a:118. Type-locality: Cuba, Trinidad Mts. Distr. — Cuba. Ref. — Alexander, 1932:387, fig. 16 (wing).
- NOCTURNA Alexander, 1941b:143. Type-locality: Venezuela, Carabobo, San Esteban. Distr. — Venezuela.
- OMISSA Alexander, 1921c:166. Type-locality: Argentina, Tucumán, Arroyo Famailla. Distr. — Argentina.



- OMISSINERVIS Alexander, 1921d:86. Type-locality: Peru, Loreto, Yurimaguas. Distr. — Peru.
- OTEROI Alexander, 1936b:878, fig. 2 (wing). Type-locality: Cuba, Sierra Rangel, Las Ánimas. Distr. — Cuba.
- PABULATORIA Alexander, 1920c:141. Type-locality: Guatemala, Gualán. Distr. — Guatemala.
- PARAXANTHA Alexander, 1962c:145, fig. 122 (wing). Type-locality: Bolivia, Santa Cruz, General Saavedra. Distr. — Bolivia.
- PARISHIANA Alexander, 1929e:406. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- PATENS Alexander, 1939a:82. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico.
- PERANGUSTA Alexander, 1939a:83. Type-locality: Mexico, Chiapas, Santa Isabel. Distr. — Mexico, Venezuela.
- VERBASALIS Alexander, 1949d:259. Type-locality: Peru, Loreto, Pucallpa. Distr. — Peru. Ref. — Alexander, 1949c:594, fig. 9 (male hypopygium).
- PERLATA Alexander, 1943b:124. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- FILIPES (Walker), 1856:440 (*Geranomyia*). Type-locality: "Amazonian Region."
- FLATYPHALLUS Alexander, 1945h:282. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Mexico, Costa Rica.
- PLEURALIS Alexander, 1913a:45. Type-locality: Guatemala, Aguna. Distr. — Guatemala. Ref. — Alexander, 1915b:444, fig. 4 (wing).
- PLEUROLINEA Alexander, 1959:450, fig. 3 (male hypopygium). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.
- PODAGRA Alexander, 1952a:128. Type-locality: Guatemala, Chicacao, El Naranjo. Distr. — Guatemala.
- POLITA Osten Sacken, 1887b:189. Type-locality: "Brazil": Distr. — Brazil. Ref. — Alexander, 1927e:23.
- FORTORICANA Alexander, 1936b:880, fig. 3 (wing). Type-locality: Puerto Rico, Villalba, El Semille. Distr. — Puerto Rico.
- PROJECTA Alexander, 1949b:492, fig. 24 (male hypopygium). Type-locality: Brazil, São Paulo, Jiquiá. Distr. — Brazil.
- PULCHELLA Alexander, 1913a:44, figs. b (wing), l (male hypopygium). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil. Ref. — Alexander, 1959:455, fig. 6 (male hypopygium).
- RECTANGULARIS Alexander, 1952a:126. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- RECTISPINA Alexander, 1968b:91. Type-locality: Ecuador, Tena, Libertad. Distr. — Ecuador.
- RETUSA Alexander, 1935a:327. Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.
- RHABDOPHORA Alexander, 1942e:256. Type-locality: Ecuador, Manabí, Palmar. Distr. — Ecuador.
- ROSTRATA Enderlein, 1912:68, fig. Q1. Type-locality: Costa Rica. Distr. — Costa Rica, Panama.
- RUBRICEPS Alexander, 1937d:614. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SUBROSTRATA Alexander, 1962c:146, figs. 123 (wing), 125 (male hypopygium). Type-locality: Bolivia, Espiritu, Río Yacuma. Distr. — Bolivia.



- RUFULA Alexander, 1946g:393, fig. 10 (male hypopygium). Type-locality: Mexico, Venodia. Distr. — Mexico.
- SACKENI Alexander, 1913a:42, fig. a (wing). Type-locality: El Salvador, Sonsonate. Distr. — Mexico, Guatemala, El Salvador, Ecuador, Peru.
- SALTI Alexander, 1930c:738. Type-locality: Colombia, Magdalena, Río Frío. Distr. — Colombia.
- SALVA Alexander, 1942c:427, fig. 1 (male hypopygium). Type-locality: Peru, Huanuco, Pumahuasi. Distr. — Peru.
- SANGUINEA Alexander, 1938f:253. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SCABROSA Alexander, 1938b:360. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- SCAPULARIS (Macquart), 1838:77(73) (*Rhamphidia*). Type-locality: "Brazil". Distr. — Brazil.
- SCHINERI Enderlein, 1912:71. Type-locality: "South America". Distr. — South America. Ref. — Alexander, 1927e:26.
flavithorax Wiedemann of Schiner, 1868:43.
- SEPOSITA Alexander, 1941g:353 (also 1946h:100). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.
- SERRULIFERA Alexander, 1945h:284. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- SIGMOIDEA Alexander, 1949b:490, figs. 19 (wing), 23 (male hypopygium). Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantareira. Distr. — Brazil.
- SIMPLEX (Wiedemann), 1828:549 (*Limnobia*). Type-locality: "Brazil". Distr. — Brazil.
- SPICA Alexander, 1943b:123. Type-locality: Ecuador, Napo-Pastaza, Abitagua, Cunibunda. Distr. — Ecuador.
- SPINIGERA Schiner, 1868:44. Type-locality: "Venezuela". Distr. — Venezuela, Colombia, Peru. Refs. — Alexander, 1927e:24; 1943c:24.
- STADELMANNI Alexander, 1945L:392, fig. 20 (male hypopygium). Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- STRICTISPINA Alexander, 1968b:92. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- STRUMOSA Alexander, 1942e:254. Type-locality: Ecuador, Manabi, Palmar. Distr. — Ecuador.
- STYGHICA Alexander, 1914e:234, pl. 3, fig. 6 (wing). Type-locality: Guyana, Bartica. Distr. — Guyana, Brazil.
- SUBANTHRACINA Alexander, 1949a:105. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador. Ref. — Alexander, 1949c:594, fig. 10 (male hypopygium).
- SUBARGENTEA Alexander, 1948b:143. Type-locality: Brazil, São Paulo, Boracéia. Distr. — Brazil. Ref. — Alexander, 1966c:489, fig. 15 (male hypopygium).
- SUBFURVA Alexander, 1966b:698. Type-locality: Nicaragua, Villa Somoza. Distr. — Nicaragua.
- SUBGALATEA Alexander, 1959:452, fig. 4 (male hypopygium). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.
- SUBJOCOSA Alexander, 1942c:432, fig. 4 (male hypopygium). Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- SUBLERIDENSIS Alexander, 1938h:439. Type-locality: Panama, Chiriqui, Finca Lerida. Distr. — Panama.



- SUBMOLESTA Alexander, 1930b:115. Type-locality: Mexico, Nayarit, between Portesuelo and Ixtlán. Distr. — Mexico.
- SUBMUNDA Alexander, 1941e:242, figs. 1 (wing), 4 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru.
- SUBPATENS Alexander, 1966b:699. Type-locality: Honduras, Lancetilla. Distr. — Honduras.
- SUBPULCHELLA Alexander, 1949b:454, fig. 5 (male hypopygium). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil.
- SUBRUBICEPS Alexander, 1945a:29. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- SULTANA Alexander, 1938b:355. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- TACTILIS Alexander, 1952a:127. Type-locality: Guatemala, Chicacao, El Naranjo. Distr. — Guatemala.
- TAINO Alexander, 1964a:71, figs. 73 (wing), 78 (male hypopygium). Type-locality: Jamaica, St. Thomas, Portland Gap. Distr. — Jamaica.
- TALAMANCANA Alexander, 1939e:624. Type-locality: Costa Rica, Talamanca, San Juan Miramar. Distr. — Costa Rica.
- TARTARUS Alexander, 1949b:488, figs. 18 (wing), 22 (male hypopygium). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- TRIFASCIATA TRIFASCIATA Enderlein, 1912:69, fig. R¹ (nom. nov. for *bifasciata* Fabricius). Type-locality: "South America". Distr. — Panama, Guyana, Colombia, Venezuela.
- bifasciata* Fabricius, 1805:31 (*Tipula*; preocc. Schrank, 1781). Type-locality: "South America".
- ssp. XANTICLES Alexander, 1946g:387. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- TRISTIS Alexander, 1913e:439, pl. 16, fig. 1 (wing). Type-locality: Peru, Madre de Dios, Callanga. Distr. — Peru.
- TULLOCHI Alexander, 1945e:238. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- TURRIALBENSIS Alexander, 1945h:285. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.
- UNICINGULATA Alexander, 1944c:43, figs. 3 (wing), 6 (male hypopygium). Type-locality: Venezuela, Merida, San Diego. Distr. — Venezuela.
- VENEZUELENSIS (Macquart), 1846:147(19), pl. 2, figs. 7, 7 a (*Limnobia*). Type-locality: "South America". Distr. — Venezuela, Colombia.
- VOLENTIS Alexander, 1952a:125. Type-locality: Bolivia, Cochabamba, Yungas de Cristal Mayu. Distr. — Bolivia.
- WIGHTI Alexander, 1939d:126, figs. 8 (wing), 17 (male hypopygium). Type-locality: Jamaica, Balaclava. Distr. — Jamaica. Ref. — Alexander, 1964a:73, fig. 74 (wing).
- WILHELMINAE Alexander, 1947a:176. Type-locality: Surinam, Wilhemina Mts. Distr. — Surinam.
- XANTIA Alexander, 1952d:246. Type-locality: Bolivia, Santa Cruz, Bucna Vista, Río Yapacani. Distr. — Bolivia.

Genus TOXORHINA Loew

TOXORHINA Loew, 1850:36 (also 1851:400). Type-species, *fragilis* Loew (Osten Sacken, 1869:113).

Toxorrhina, emend.

Ref. — Alexander, 1947c:356.



Subgenus CERATOCHEILUS Wesché

- CERATOCHEILUS Wesché, 1910:358 (as genus). Type-species, *winnsampsoni* Wesché (Brunetti, 1920:230). Ref. — Alexander, 1947c:356.
- AMERICANA (Alexander), 1913a:49, figs. e (wing), j (ovipositor) (*Ceratocheilus*). Type-locality: Panama, Culebra. Distr. — Costa Rica, Panama, Colombia, Guyana, Peru, Brazil.
- ATRITARSIS Alexander, 1943e:211. Type-locality: Ecuador, Santiago-Zamora, Río Zamora, Zumbi. Distr. — Ecuador.
- CHIAPASENSIS Alexander, 1938e:410. Type-locality: Mexico, Chiapas, Finca Belén. Distr. — Mexico.
- LEUCOMELANOPUS (Enderlein), 1912:78 (*Aporosa*). Type-locality: Colombia, Hacienda Pehlke. Distr. — Colombia.
- LEUCOSTENA Alexander, 1937c:502. Type-locality: Surinam, King Frederick William Falls, Courantijne River. Distr. — Surinam. Ref. — Alexander, 1947c:355, fig. 30 (wing).
- MACULIPENNIS Alexander, 1936a:22. Type-locality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil.
- NIVETARSIS (Alexander), 1922c:49 (*Ceratocheilus*). Type-locality: Colombia, Boca del Condoto. Distr. — Panama, Colombia.
- PROLONGATA Alexander, 1938f:259. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- REVULSA REVULSA Alexander, 1954d:151. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- ssp. MACRORHYNCHIA Alexander, 1954d:152. Type-locality: Bolivia, Cochabamba, Chapare, Locotal, El Limbo. Distr. — Bolivia.
- SUPERSTES Alexander, 1942c:447, figs. 8 (wing), 15 (male hypopygium). Type-locality: Peru, Junín, Tarma, Huacapistana. Distr. — Peru. Ref. — Alexander, 1947c:355, fig. 31 (wing).
- VULSA Alexander, 1952a:134. Type-locality: Peru, Junín, Chanchamayo. Distr. — Peru.

Subgenus TOXORRHINA Loew

- ATRIPES Alexander, 1922c:48. Type-locality: Colombia, Condoto. Distr. — Colombia.
- BRASILIENSIS (Westwood), 1835:683 (*Limnobiaorhynchus*). Type-locality: "Brazil". Distr. — Guyana, Peru, Brazil, Paraguay. (See *westwoodi* Brunetti).
- CENTRALIS Alexander, 1913a:52, fig. i (wing) (*Toxorrhina*). Type-locality: "Surinam". Distr. — Mexico, Guatemala, Costa Rica, Panama, Guyana, Surinam, Brazil.
- CURVATA Alexander, 1938b:361. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- DISTALIS Alexander, 1936b:881, fig. 4 (wing). Type-locality: Bahamas, New Providence. Distr. — Bahamas.
- DOMINGENSIS Alexander, 1937a:186, fig. 4 (wing). Type-locality: Hispaniola: Dominican Republic, Sanchez. Distr. — Dominican Republic, Cuba.
- FLAVIDA Alexander, 1913a:51, fig. h (wing) (*Toxorrhina*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil, Peru.



- FRAGILIS Locw, 1851:401, pl. 2, figs. 16 (antenna), 17 (body, female), 18 (venation), 22 (head, female). Type-locality: "Puerto Rico". Distr. — Puerto Rico. Ref. — Alexander, 1932:378, fig. 19 (wing).
- INFUMIPENNIS Alexander, 1942j:31 (nom. nov. for *fumipennis* Alexander). Type-locality: Lesser Antilles: Dominica, Sylvania Estate. Distr. — Dominica.
- fumipennis* Alexander, 1939d:100 (preocc. Alexander, 1936). Type-locality: Dominica, Sylvania Estate.
- JAMAICENSIS Alexander, 1964a:83, figs. 92 (wing), 96 (male hypopygium). Type-locality: Jamaica, St. Thomas, Corn Puss Gap. Distr. — Jamaica.
- LATAMERA Alexander, 1968b:96. Type-locality: Ecuador, Huagra Yacu. Distr. — Ecuador.
- LONGICOLLIS Pierre, 1924:80, figs. 6-12 (body, wing). Type-locality: Venezuela, Bolivar, San Fernando de Apure. Distr. — Venezuela.
- MENDOSA Alexander, 1935a:330. Type-locality: Brazil, Rio de Janeiro. Rezende, Campo Belo. Distr. — Brazil.
- MERIDIONALIS Alexander, 1913a:51, fig. g (wing) (*Toxorhina*). Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.
- NIGRIVENA Alexander, 1939e:636, fig. 8 (wing). Type-locality: Costa Rica, Talamanca, San Juan Miramar. Distr. — Costa Rica.
- PERGRACILIS Alexander, 1944b:181, fig. 13 (male hypopygium). Type-locality: Peru, Ayacucho, La Mar, Ayna. Distr. — Venezuela, Peru, Bolivia.
- PHIORACAENA Alexander, 1966a: 227. Type-locality: Ecuador, Napo-Pastaza. Libertad. Distr. — Ecuador.
- POLYCANTHA Alexander, 1940e:907, fig. 6 (wing). Type-locality: Venezuela, Miranda, Río Chacaito. Distr. — Venezuela.
- SCAPANIA Alexander, 1966a:228. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- STENOPHALLUS Alexander, 1937c:503. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Venezuela, Brazil.
- TRILOBATA Alexander, 1938e:411. Type-locality: Mexico, Chiapas, Vergel. Distr. — Mexico. Ref. — Alexander, 1947c:355, fig. 32 (wing).
- VIOLACEIPENNIS Alexander, 1937a:187, fig. 5 (wing). Type-locality: Cuba, Sierra del Cobre, Loma del Gato. Distr. — Cuba. Ref. — Alexander, 1964a:85, fig. 97 (male hypopygium).
- WESTWOODI Brunetti, 1920:229 (nom. nov. for *brasiliensis* Westwood). Based on female only. See Brunetti, 1918:300-304, discussion. Type-locality: "Brazil".
- brasiliensis* Westwood, 1835:638 (*Limnobiaorhynchus*; part; female only). Type-locality: "Brazil".

Genus TRENTEPOHLIA Bigot

- TRENTEPOHLIA Bigot, 1854:456, 473. Type-species, *trentepohlii* (Wiedemann) (mon.) (Oriental). Ref. — Alexander, 1947b:69-12, figs.

Subgenus MONGOMA Westwood

- MONGOMA Westwood, 1881:364 (as genus). Type-species, *fragillima* Westwood (mon.).



ERRANS Alexander, 1944d:321. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.

Subgenus NEOMONGOMA Alexander

- Trentepohlia*, subg. NEOMONGOMA Alexander, 1919a:140. Type-species, *Mongoma disjuncta* Alexander (orig. des.).
- DISJUNCTA (Alexander), 1913b:500, fig. 22 (wing) (*Mongoma*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil.
- FUSCOTERMINALIS Alexander, 1949b:481, fig. 15 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil. Ref. — Alexander, 1947b:67, fig. 5 (wing).
- MESONOTALIS Alexander, 1949b:482. Type-locality: Brazil, São Paulo, Guarujá. Distr. — Brazil.
- SORDIDIPENNIS Alexander, 1943f:302, fig. 11 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- SUBERECTA Alexander, 1939b:208. Type-locality: Ecuador, Napo-Pastaza, Abitagua. Distr. — Ecuador.
- ZERNYI Alexander, 1942g:180. Type-locality: Brazil, Pará, Santarém, Taparinhá. Distr. — Brazil.

Subgenus PARAMONGOMA Brunetti

- PARAMONGOMA Brunetti, 1911:295 (as genus). Type-species, *Cylindrotoma albitarsis* Doleschall (orig. des.) (Oriental).
- Mongomella* Enderlein, 1912:61. Type-species, *Mongoma pallida* Williston (orig. des.) (Neotropical).
- AMATRIX Alexander, 1942g:181. Type-locality: Brazil, Rio de Janeiro, Serra do Itatiaia, "Südabhang". Distr. — Brazil.
- BROMELIADICOLA (Alexander), 1912d:415, fig. (wing) (*Mongoma*). Type-locality: "Costa Rica". Distr. — Costa Rica. Ref. — Picado, 1913:356, pl. 13 (biology).
- CALLIOPE Alexander, 1944g:304. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru, Bolivia.
- CHIRIQUIANA Alexander, 1934a:68. Type-locality: Panama, Chiriqui, Finca Parada, Distr. — Panama.
- CONCUMBENS Alexander, 1942i:66, fig. 35 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- CONSCRIPTA Alexander, 1949a:102. Type-locality: Peru, Huanuco, Carpish. Distr. — Peru, Bolivia.
- CUBITALIS Alexander, 1931a:116. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- DISPARILIS Alexander, 1944b:178. Type-locality: Peru, Junín, Satipo, Jauja. Distr. — Peru.
- FITZLERI Alexander, 1947e:65. Type-locality: Venezuela, Táchira, East Táchira. Distr. — Venezuela.
- DOMINICANA Alexander, 1947b:71. Type-locality: Lesser Antilles: Dominica. Distr. — Puerto Rico, Dominica.
- EXTENSA (Alexander), 1913b:501, fig. 11 (wing) (*Mongoma*). Type-locality: Panama, Culcra. Distr. — Panama, Guyana, Brazil.
- FAUSTINA Alexander, 1945d:277. Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil.



- FEMORATA Alexander, 1921d:88. Type-loeality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- FLAVELLA Alexander, 1921d:89. Type-loeality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- FUSCIPES Alexander, 1921d:88. Type-loeality: Brazil, Amazonas, Tefé. Distr. — Brazil.
- FUSCOLIMBATA Alexander, 1950a:53, fig. 29 (wing). Type-locality: Venezuela, Aragua, Rancho Grande. Distr. — Venezuela.
- GENICULATA (Alexander), 1914e:247, pl. 4, fig. 7 (wing) (*Mongoma*). Type-locality: Guyana, Bartica. Distr. — Guyana, Brazil.
- LAUDABILIS Alexander, 1942c:442. Type-loeality: Peru, Huanuco, Piedras Grandes. Distr. — Peru.
- LEUCOXENA (Alexander), 1915a:29 (*Mongoma*). Type-locality: Mexieo, Veraeruz, Córdoba. Distr. — Mexico. Ref. — Alexander, 1915b:444, fig. 7 (wing).
- LONGIFUSA (Alexander), 1913b:502, fig. 10 (wing) (*Mongoma*). Type-loeality: Brazil, Pará, Igarapé-Açu. Distr. — Ecuador, Brazil.
- MANEA (Williston), 1896a:293 (*Mongoma*). Type-loeality: Lesser Antilles: St. Vineent. Distr. — St. Vineent.
- METATARSATA (Alexander), 1915b:444, fig. 6 (wing) (*Mongoma*). Type-locality: Panama, Trinidad River. Distr. — Panama, Brazil.
- MONTIVAGA Alexander, 1949a:102. Type-locality: Peru, Huanuco, Saria-pampa. Distr. — Peru, Bolivia.
- NEOGAMA Alexander, 1943f:303, fig. 12 (wing). Type-loeality: Brazil, Espírito Santo, Morro Argolas. Distr. — Brazil.
- NIVETARSIS (Alexander), 1913b:501, fig. 13 (wing) (*Mongoma*). Type-locality: Puerto Rico, El Yunque. Distr. — Jamaiea, Puerto Rieo. Refs. — Alexander, 1932:375, fig. 17 (wing); 1964a:67, fig. 69 (venation).
- PALLIDA (Williston), 1896a:293 (*Mongoma*). Type-loeality: Lesser Antilles: St. Vineent. Distr. — Brazil, St. Vineent.
- PALLIPES (Alexander), 1914e:247, pl. 4, fig. 8 (wing) (*Mongoma*). Type-loeality: Guyana, Mallali. Distr. — Guyana, Brazil.
- PETULANS Alexander, 1937b:244. Type-loeality: Brazil, Rio de Janeiro, Rezende, Campo Belo. Distr. — Brazil. Ref. — Alexander, 1947b:67, fig. 7 (wing).
- RORAIMICOLA Alexander, 1935a:329. Type-locality: Venezuela, Mt. Roraima. Distr. — Venezuela, Brazil.
- SORORCULA Alexander, 1919a:141, fig. 15 (wing). Type-loeality: Panama, Darién, between Río Capeti and Río Yape. Distr. — Panama.
- SUBLEUCOXENA Alexander, 1940b:160. Type-loeality: Mexieo, Chiapas, San Cristóbal. Distr. — Mexieo.
- SUFFUSCIPES Alexander, 1947f:101. Type-locality: Venezuela, Aragua, Choroní. Distr. — Venezuela.
- TETHYS Alexander, 1949a:101. Type-loeality: Peru, Huanuco, Fundo Sinchono. Distr. — Peru.
- TUCUMANA Alexander, 1936d:762. Type-locality: Argentina, Tucumán. Distr. — Argentina.

Subgenus PROMONGOMA Alexander

Trentepohlia, subg. PROMONGOMA Alexander, 1938b:353. Type-species, *mirabilis* Alexander (orig. des.).



MIRABILIS Alexander, 1938b:354, fig. 8 (wing). Type-locality: Eeuador. Napo-Pastaza, Abitagua, 1200m. Distr. — Eeuador. Ref. — Alexander, 1947b:67, fig. 6 (wing).

LITERATURE CITED

ABREU, D. E. SANTOS

1923. Monografía de los Limonidos de las Islas Canarias. *Mem. R. Acad. Cienc. Artes Barcelona* (3)18(4):1-164, 25 figs., 4 color. pls. with 33 figs.

ALDRICH, J. M.

1905. A catalogue of North American Diptera. *Smiths. Misc. Coll.* 46(2):1-680.

ALEXANDER, e. P.

1911a. Synonymical, and other notes on the Tipulidae (Diptera). *Psyche* 18:192-203; 1 pl.

1911b. Notes on two Tipulidae (Dipt.) *Ent. News* 22:349-354; 4 figs.

1912a. A peculiar new crane-fly from Porto Rico (Tipulidae: Diptera). *Psyche* 19:63-66; 1 pl. (3 figs.).

1912b. New species of *Furcomyia* (Tipulidae). *Canad. Ent.* 44:333-342, 361-364, 1 pl. (16 figs.).

1912c. On the tropical American Rhipidiidae (Tipulidae, Dipt.). *Bull. Brooklyn Ent. Soc.* 8:6-17; 1 pl. (9 figs.).

1912d. A Bromeliad-inhabiting crane-fly (Tipulidae, Dipt.). *Ent. News* 23:415-417; 1 text fig.

1912e. A new tropical *Gonomyia* (Tipulidae, Dipt.). *Ent. News* 23:418-420; 4 text figs.

1912f. New Neotropical Tipulinae (Tipulidae, Dipt.). *Ann. Ent. Soc. America* 5:343-365; 3 pls. (23 figs.).

1912g. A revision of the genus *Brachypremna* Osten Sacken (Tipulidae, Diptera). *Jour. New York Ent. Soc.* 20:225-236; 1 pl. (6 figs.), 1 text fig.

1912h. New African Tipulidae. *Canad. Ent.* 44:83-88; 2 figs.

1913a. New neotropical Antoehini (Tipulidae, Diptera). *Psyche* 20:40-54; pl. 2 (13 figs.).

1913b. A synopsis of part of the neotropical crane-flies of the subfamily Limnobiinae. *Proc. U. S. Nat. Mus.* 44:481-549, pls. 65-68 (42 figs.).

1913c. Report on a collection of crane-flies (Tipulidae, Dipt.) from Colombian Andes, taken by Mr. John Thomas Lloyd. *Jour. New York Ent. Soc.* 21:193-212; 3 pls. (26 figs.), 2 photographs.


1913d. The neotropical Tipulidae in the Hungarian National Museum (Diptera). — I. *Ent. News* 24:404-412; 1 pl. (10 figs.).

1913e. The neotropical Tipulidae in the Hungarian National Museum (Diptera). — II. *Ent. News* 24:439-449; 1 pl. (10 figs.).


1914a. New or little-known neotropical Hexatomini (Tipulidae, Diptera). *Psyche* 21:33-45, 1 pl. (8 figs.).

1914b. The neotropical Tipulidae in the Hungarian National Museum (Diptera). — III. *Ent. News* 25:205-215, 1 pl. (13 figs.).

1914e. The crane-flies collected in Costa Rica by Dr. P. P. Calvert (Tipulidae, Diptera). *Jour. New York Ent. Soc.* 22:116-124; 1 pl. (10 figs.).




- 1914d. A revision of the American species of *Tanyremna* Osten Sacken and *Megistocera* Wiedemann (Tipulidae, Diptera). *Jour. New York Ent. Soc.* 22:205-218; 1 pl. (5 figs.).
- 1914e. On a collection of crane-flies from British Guiana (Tipulidae, Diptera). *Trans. Amer. Ent. Soc.* 40:223-225, pls. 3-4 (20 figs.).
- 1914f. The neotropical Tipulidae in the Hungarian National Museum (Diptera). — IV. *Ent. News* 25:351-363; 1 pl. (18 figs.).
- 1914g. New or little-known crane-flies from the United States and Canada. Tipulidae, Diptera. *Proc. Acad. Nat. Sci. Philadelphia* 66:579-606, 3 pls. (32 figs.).
- 1915a. A second Bromeliad-inhabiting crane-fly (Tipulidae, Diptera). *Ent. News* 26:29-30.
- 1915b. Description of new species of crane-flies from Central America. *Proc. U. S. Nat. Mus.* 48:441-444, pl. 24 (7 figs.).
- 1915e. Two new crane-flies from Porto Rico (Tipulidae, Diptera). *Insec. Inscit. Menst.* 3:104-107.
- 1915d. New or little-known crane-flies from the United States and Canada: Tipulidae, Diptera. Part 2. *Proc. Acad. Nat. Sci. Philadelphia* 67:458-514, 6 pls. (87 figs.).
- 1916a. New or little-known crane-flies from Colombia, Ecuador, and Peru (Tipulidae, Diptera). *Trans. Amer. Ent. Soc.* 42:1-32; 5 pls. (43 figs.).
- 1916b. New North American species of the genus *Gonomyia* Meigen (Tipulidae, Diptera). *Canad. Ent.* 48:316-325.
- 1916c. New species of crane-flies from the West Indies (Tipulidae, Dip.). *Ent. News* 27:343-347, 6 text-figs.
- 1916d. New or little-known crane-flies from the United States and Canada: Tipulidae, Ptychopteridae, Diptera. Part 3. *Proc. Acad. Nat. Sci. Philadelphia* 68:486-549, 7 pls. (98 figs.).
- 1917a. New Nearctic crane-flies (Tipulidae, Diptera). Pt. II. *Canad. Ent.* 49:22-31, 61-64, 199-211, pl. 12.
- 1917b. The crane flies of South Africa in the South African Museum (Diptera, Tipulidae). Pt. I. *Ann. South African Mus.* 17:139-180, 5 pls. (67 figs.), 2 text figs.
- 1919a. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). I. *Jour. New York Ent. Soc.* 27:132-154; 1 pl. (16 figs.).
- 1919b. The crane-flies of New York. Part I. Distribution and taxonomy of the adult flies. *Mem. New York (Cornell) Agr. Expt. Sta.* 25:763-993; 26 pls. (345 figs.), 12 text figs.
- 1919e. New Nearctic crane-flies (Tipulidae, Diptera). Part VIII. *Canad. Ent.* 51:191-199.
- 1919d. The biology of the North American crane-flies (Tipulidae, Diptera). IV. The genus *Dicranoptycha* Osten Sacken. *Pomona College Jour. Ent. and Zool.* 11:67-74; 2 pls. (14 figs.).
- 1920a. Undescribed Tipulidae collected by Mr H. S. Parish in Brazil (Dipt.). *Ent. News* 31:70-75.
- 1920b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). II. *Jour. New York Ent. Soc.* 28:1-13.
- 1920e. New or little-known crane-flies from tropical America (Tipulidae, Diptera). *Canad. Ent.* 52:141-144.
- 1920d. In Skottsberg, *The natural history of Juan Fernandez and Easter Island. Diptera — Tipulidae.* *Zool.* 3:25-27.



- 1920e. The crane-flies collected by the Swedish expedition (1895-1896) to southern Chile and Tierra del Fuego (Tipulidac, Diptera). *Arkiv för Zoologi* 13:1-32; 2 pls. (20 figs.).
- 1920f. Undescribed crane-flies from Argentina (Tipulidae, Diptera). *Ent. News* 31:215-221.
- 1920g. New or little-known crane-flies in the Queensland Museum (Tipulidae, Diptera). *Mem. Queensland Mus.* 7:52-63.
- 1920h. Undescribed Tipulidae (Diptera) from western North America. *Proc. California Acad. Sci.* 10:35-36.
- 1920i. The crane-flies of New York. Part II. Biology and Phylogeny. *Mem. New York (Cornell) Agr. Expt. Sta.* 38:699-1133; 87 pls. (540 figs.).
- 1921a. Undescribed crane-flies from Argentina (Tipulidae, Diptera). Part II. *Ent. News* 32:72-76.
- 1921b. Undescribed crane-flies from Argentina (Tipulidae, Dipt.). Part III. *Ent. News* 32:175-179.
- 1921c. Undescribed Eriopterine crane-flies from Argentina (Tipulidae, Diptera). *Canad. Ent.* 53:162-166.
- 1921d. New or little-known crane-flies from the Amazonian region. *Proc. Acad. Nat. Sci. Philadelphia* 73:39-103; 1 pl. (12 figs.).
- 1921e. Undescribed crane-flies from Argentina (Tipulidae, Diptera). Part IV. *Ent. News* 32:292-295.
- 1921f. New species of crane-flies from North Queensland (Tipulidae, Diptera). *Canad. Ent.* 53:205-211.
- 1921g. On new or little-known Tipulidac (Diptera). VII. Australasian species. *Ann. Mag. Nat. Hist.* (8)9:546-563.
- 1921h. New or little known Tipulidae. V. Ethiopian species. *Ann. Mag. Nat. Hist.* (8)8:161-175.
- 1922a. Undescribed species of Costa Rican flies belonging to the family Tipulidae in the United States National Museum. *Proc. U. S. Nat. Mus.* 60:1-7.
- 1922b. Undescribed crane-flies from Argentina (Dipt.: Tipulidae). Part V. *Ent. News* 33:207-210.
- 1922c. New or little-known exotic Tipulidae (Diptera). *Trans. Royal Ent. Soc. London* 1922:34-64.
- 1922d. Undescribed crane-flies in the Paris Museum (Tipulidae, Diptera). Part III. *Bull. Mus. d'Hist. Nat.* 1921:303-306, 346-350, 539-542; 1922:73-75.
- 1922e. Undescribed crane-flies (Tanyderidae and Tipulidae) in the South Australian Museum. *Rec. South Australian Mus.* 2(2):223-270.
- 1922f. New or little-known Tipulidae (Diptera). X. Australasian species. *Ann. Mag. Nat. Hist.* (9)9:505-524.
- 1922g. New or little-known species of Australian Tipulidae (Diptera). I. *Proc. Linn. Soc. New South Wales* 1922:583-590.
- 1922h. Four undescribed species of *Limnobia* from the Oriental Region (Diptera, Tipulidae). *Insec. Inscit. Menst.* 10:75-80.
- 1923a. Undescribed species of *Eriocera* and *Penthoptera* from tropical America (Tipulidae, Diptera). *Ent. News* 34:17-20.
- 1923b. Undescribed crane-flies from Argentina (Dipt.: Tipulidae). Part VI. *Ent. News* 34:181-186.
- 1923c. Undescribed crane-flies from Argentina (Dipt.: Tipulidac). Part VII. *Ent. News* 34:309-313.




- 1924a. Undescribed crane-flies from Argentina (Dipt.: Tipulidae). Part VIII. *Ent. News* 35:61-65.
- 1924b. Undescribed species of the genus *Tanyremna* Osten Sacken (Dipt.: Tipulidae). *Ent. News* 35:289-291.
- 1924c. New or little-known Tipulidae (Diptera). XXI. Australasian species. *Ann. Mag. Nat. Hist.* (9)13:359-380.
- 1924d. Undescribed species of Japanese crane flies (Tipulidae, Diptera). Part IV. *Ann. Ent. Soc. Amer.* 17:59-74.
- 1924e. New or little-known Tipulidae (Diptera). XX. Australasian species. *Ann. Mag. Nat. Hist.* (9)13:177-194.
- 1924f. New or little-known Tipulidae (Diptera). XXII. Australasian species. *Ann. Mag. Nat. Hist.* (9)13:499-523.
- 1925a. Studies on the crane-flies of Mexico, Part I. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 18:341-362; pls. 14-15 (26 figs.).
- 1925b. New or little-known Tipulidae (Diptera). XXIX. Australasian species. *Ann. Mag. Nat. Hist.* (9)16:65-81.
- 1926a. Studies on the crane-flies of Mexico, Part II. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 19:158-179; pls. 11-12 (19 figs.).
- 1926b. Undescribed species of crane-flies from Cuba and Jamaica (Tipulidae, Dipt.). *Jour. New York Ent. Soc.* 34:223-230.
- 1926c. New or little-known Tipulidae in the collection of the Deutsches Entomologisches Institut (Dipt.). *Ent. Mitt.* 15:386-392; 2 text figs.
- 1926d. New species of crane-flies from South America, Part I. (Family Tipulidae, Order Diptera). *Ann. Ent. Soc. Amer.* 19:378-394.
- 1926e. New or little-known Tipulidae (Diptera). XXXII. Australasian species. *Ann. Mag. Nat. Hist.* (9)18:169-193; 16 text figs.
- 1927a. Undescribed species of Tipuline crane-flies from tropical America. *Insec. Inscit. Menst.* 14(1926):159-167.
- 1927b. Genotypes of the Chilean Tipuloidea described by Philippi. *Rev. Chil. Hist. Nat.* 31:15.
- 1927c. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). III. *Jour. New York Ent. Soc.* 35:265-278.
- 1927d. Studies on the crane-flies of Mexico, Part III. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 20:301-318.
- 1927e. New or little-known crane-flies. Part II. *Encycl. Ent., Diptera* 4:17-27.
- 1928a. Studies on the crane-flies of Mexico. Part IV. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 21:101-119.
- 1928b. Records and descriptions of neotropical crane-flies (Tipulidae, Diptera). IV. *Jour. New York Ent. Soc.* 36:47-59.
- 1928c. Undescribed species of crane-flies from Chile. *Ent. News* 39:180-183.
- 1928d. New species of Tipulidae from Chile. *Rev. Chil. Hist. Nat.* 31:217-221.
- 1928e. Undescribed species of crane-flies from Valdivia (Chile). *Rev. Chil. Hist. Nat.* 31:245-249.
- 1928f. New species of crane-flies from South America. Part II. (Family Tipulidae, Order Diptera). *Ann. Ent. Soc. Amer.* 21:623-641.



- 1928g. The crane-flies of Jamaica. *Catalogus Insectorum Jamaicaensis*. *Dept. Sci. & Agr., Ent. Bul.* 4(3):19-29; 2 text figs.
- 1928h. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). V. *Jour. New York Ent. Soc.* 36:355-367.
- 1928i. New or little-known species of the genus *Tipula* from Chile (Tipulidae, Diptera). *Rev. Chil. Hist. Nat.* 32:276-286; 1 pl. (13 figs.).
- 1928j. New or little-known Tipulidae from eastern Asia. Part II. *Philippine Jour. Sci.* 35:455-489; 2 pls. (30 figs.).
- 1929a. *Diptera of Patagonia and South Chile*. Part I — Crane-flies. *British Mus. Nat. Hist.* 1929, 240 pp.; 12 pls. (259 figs.), 17 text figs, 1 map.
- 1929b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). VI. *Jour. New York Ent. Soc.* 37:89-99.
- 1929c. New or little-known species of the genus *Gnophomyia* Osten Saeken from Ecuador and Peru (Family Tipulidae, Order Diptera). *Rev. Chil. Hist. Nat.* 33:164-168; 1 pl. (4 figs.).
- 1929d. New species of crane-flies from South America. Part III. *Ann. Ent. Soc. Amer.* 22:768-788.
- 1929e. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). VII. *Jour. New York Ent. Soc.* 37:395-407.
- 1929f. New or little-known Tipulidae from the Philippines (Diptera). V. *Philippine Jour. Sci.* 40:239-271; 18 figs.
- 1930a. The genus *Sigmatomera* Osten Saeken with observations on the biology by Raymond C. Shannon. *Encycl. Ent., Diptera* 5:155-162; 8 text figs.
- 1930b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). VIII. *Jour. New York Ent. Soc.* 38:109-120.
- 1930c. New species of crane-flies from South America. Part IV. (Family Tipulidae, Order Diptera). *Ann. Ent. Soc. Amer.* 23:721-740.
- 1931a. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). IX. *Jour. New York Ent. Soc.* 39:109-122.
- 1931b. New or little-known Tipulidae in the American Museum of Natural History (Diptera). I. *Amer. Mus. Novitates* 491:1-16; 18 text figs.
- 1931c. New species of crane-flies from South America. Part V. (Family Tipulidae, Order Diptera). *Ann. Ent. Soc. Amer.* 24:622-642.
- 1931d. Crane flies of the Baltic Amber (Diptera). *Bernstein-Forschungen*, [Heft] 2:135 pp.; 168 figs.
1932. The crane-flies of Puerto Rico. *Jour. Dept. Agr. Puerto Rico* 16:349-387; 6 pls. (26 figs.).
1933. Die Ausbeute der deutschen Chaco-Expedition 1925-26 Diptera. XXXVIII. Tipulidae. *Konowia* 12:41-46; 2 text figs.
- 1934a. New or little-known Tipulidae from Chiriqui, Panama (Order Diptera). *Ann. Ent. Soc. Amer.* 27:55-73.
- 1934b. Tipulidae, in Curran, *The Diptera of Kartabo, Bartica District, British Guiana*. *Bull. Amer. Mus. Nat. Hist.* 66:303-305; 1 pl. (9 figs.).
- 1934c. Tipuloidae, in Curran, *The Families and Genera of North American Diptera*. 1934. Pp. 28-60, 4 pls. (44 figs.), 5 text figs.
- 1934d. Records and descriptions of Chilean Tipulidae (Order Diptera). Part I. *Rev. Chil. Hist. Nat.* 38:173-179; 1 text fig.



- 1935a. New species of crane-flies from South America, Part VI. (Family Tipulidae: Order Diptera). *Ann. Ent. Soc. Amer.* 28:313-331.
- 1935b. Records and descriptions of Chilean Tipulidae (Order Diptera). Part II. *Rev. Chil. Hist. Nat.* 39:99-105; 7 text figs.
- 1936a. Records and descriptions of Brazilian Tipulidae (Dipt.). Part I. *Rev. Ent.* 6:10-23.
- 1936b. New or little-known species of West Indian Tipulidae (Diptera). I. *Jour. Agr. Univ. Puerto Rico* 20:877-882; 4 text figs.
- 1936c. Undescribed species of *Brachypremna* and *Tanypremna* (Tipulidae, Diptera). *Rev. Chil. Hist. Nat.* 40:298-305; 1 pl. (4 figs.).
- 1936d. New species of crane-flies from South America. Part VII. (Family Tipulidae, Order Diptera). *Ann. Ent. Soc. Amer.* 29:749-769.
- 1936e. New or little-known Tipulidae from tropical America (Diptera). Part I. *Konowia* 16("1937"):55-66
- 1936f. New or little-known species of Tipulidae from eastern Asia (Diptera). XXX. *Philippine Jour. Sci.* 60:165-204; 45 figs.
- 1936g. The Diptera of the territory of New Guinea. V. Family Tipulidae. Part III. *Proc. Linn. Soc. New South Wales* 61:322-340; 44 figs.
- 1937a. New or little-known species of West Indian Tipulidae (Diptera). II. *Jour. Agr. Univ. Puerto Rico* 21:179-190; 1 pl. (11 figs.).
- 1937b. Records and descriptions of Brazilian Tipulidae (Dipt.). Part II. *Rev. Ent.* 7:233-246.
- 1937c. New or little-known Tipulidae (Diptera). LIV. Neotropical species. *Ann. Mag. Nat. Hist.* (10)20:481-504.
- 1937d. New species of crane-flies from South America. Part. VIII. (Tipulidae, Diptera). *Ann. Ent. Soc. Amer.* 30:598-617.
- 1937e. New or little-known species of West Indian Tipulidae (Diptera). III. *Jour. Agr. Univ. Puerto Rico* 21:523-534; 1 pl. (10 figs.).
- 1937f. Undescribed species of South American Tipulidae (Diptera) Part I. *Rev. Chil. Hist. Nat.* 41:218-227.
- 1938a. New or little-known crane-flies. Pt. III. *Encycl. Ent., Diptera* 9:23-36.
- 1938b. New or little-known Tipulidae (Diptera). LIV. Neotropical species. *Ann. Mag. Nat. Hist.* (11)1:336-362; 8 text figs.
- 1938c. Records and descriptions of Brazilian Tipulidae (Dipt.). Part III. *Rev. Ent.* 8:318-331.
- 1938d. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). X. *Jour. New York Ent. Soc.* 46:327-337.
- 1938e. Studies on the crane-flies of Mexico. Part V. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 31:303-412.
- 1938f. Records and descriptions of Brazilian Tipulidae (Dipt.). Part IV. *Rev. Ent.* 9:247-260.
- 1938g. New or little-known Tipulidae (Diptera). LV. Neotropical species. *Ann. Mag. Nat. Hist.* (11)2:416-438.
- 1938h. Records and descriptions of Tipulidae from tropical America (Diptera). Part I. *Rev. Ent.* 9:426-441.
- 1938i. Records and descriptions of Chilean Tipulidae. Part III. *Rev. Chil. Hist. Nat.* 42:276-282.
- 1938j. New or little-known Tipulidae from eastern Asia (Diptera). XXXVI. *Philippine Jour. Sci.* 66:93-134; 3 pls. (46 figs.).



- 1939a. Studies on the crane-flies of Mexico. Part VI. *Ann. Ent. Soc. Amer.* 32:70-90.
- 1939b. New or little-known Tipulidae (Diptera). LVI. Neotropical species. *Ann. Mag. Nat. Hist.* (11)3:186-209.
- 1939c. Records and descriptions of Brazilian Tipulidae (Dipt.). Part V. *Rev. Ent.* 10:428-444; 16 text figs.
- 1939d. New or little-known species of West Indian Tipulidae (Diptera). Part IV. *Jour. Agr. Univ. Puerto Rico* 23:91-130; 2 pls. (18 figs.).
- 1939e. Records and descriptions of Tipulidae from tropical America (Diptera). Part II. *Rev. Ent.* 10:622-637; 16 text figs.
- 1939f. Records and descriptions of Chilean Tipulidae (Order Diptera). Part IV. *Rev. Chil. Hist. Nat.* 43:176-184; 3 text figs.
- 1940a. New or little-known Tipulidae (Diptera). LVII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)5:275-297.
- 1940b. Studies on the crane-flies of Mexico. Part VII. (Order Diptera, superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 33:140-161.
- 1940c. Records and descriptions of Neotropical crane-flies. XI. *Jour. New York Ent. Soc.* 48:105-116.
- 1940d. New or little-known Tipulidae (Diptera). LVIII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)6:194-222; 9 text figs.
- 1940e. Records and descriptions of Tipulidae from tropical America (Diptera). Part III. *Rev. Ent.* 11:894-908; 10 text figs.
- 1940f. Records and descriptions of Brazilian Tipulidae (Dipt.). Part VI. *Rev. Ent.* 11:382-397; 15 text figs.
- 1940g. New or little-known Tipulidae from eastern Asia (Diptera). XLI. *Philippine Jour. Sci.* 71:39-76; 50 figs.
- 1941a. New species of crane-flies from South America. Part IX. (Tipulidae, Diptera). *Ann. Ent. Soc. Amer.* 34:231-254.
- 1941b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XII. *Jour. New York Ent. Soc.* 49:139-148.
- 1941c. Records and descriptions of Tipulidae from tropical America (Diptera). Part IV. *Rev. Ent.* 12:322-337; 7 text figs.
- 1941d. New or little-known Tipulidae (Diptera). LIX. Neotropical species. *Ann. Mag. Nat. Hist.* (11)8:105-132; 15 text figs.
- 1941e. New or little-known Tipulidae (Diptera). LX. Neotropical species. *Ann. Mag. Nat. Hist.* (11)8:240-265; 17 text figs.
- 1941f. New or little-known Tipulidae (Diptera). LXI. Neotropical species. *Ann. Mag. Nat. Hist.* (11)8:313-344; 11 text figs.
- 1941g. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XIII. *Jour. New York Ent. Soc.* 49:345-356.
- 1941h. Undescribed species of South American Tipulidae (Diptera). Part II. *Rev. Chil. Hist. Nat.* 44:258-263.
- 1942a. New or little-known Tipulidae (Diptera). LXII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)9:219-245; 11 text figs.
- 1942b. New or little-known Tipulidae (Diptera). LXIII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)9:338-370; 16 text figs.
- 1942c. New or little-known Tipulidae (Diptera). LXIV. Neotropical species. *Ann. Mag. Nat. Hist.* (11)9:424-448; 15 text figs.
- 1942d. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XIV. *Jour. New York Ent. Soc.* 50:135-146.
- 1942e. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XV. *Jour. New York Ent. Soc.* 50:251-262.

- 1942f. New or little-known Tipulidae (Diptera). LXV. Neotropical species. *Ann. Mag. Nat. Hist.* (11) 9:759-790; 7 text figs.
- 1942g. Records and descriptions of Brazilian Tipulidae (Dipt.). Part VII. *Rev. Ent.* 13:166-182; 12 text figs.
- 1942h. Records and descriptions of Brazilian Tipulidae (Dipt.). Part VIII. *Rev. Ent.* 13:421-437; 21 text figs.
- 1942i. New or little-known Tipulidae (Diptera) from São Paulo, Brazil. Part I. *Papéis Avulsos do Dept. de Zool.* 2:39-77; 38 text figs.
- 1942j. The Oriental Tipulidae in the collection of the Indian Museum. Part II. *Rec. Indian Mus.* 44:29-72; 21 figs.
- 1942k. The Diptera or true flies of Connecticut. First Fascicle. *State Geol. and Nat. Hist. Surv. Connecticut Bull.* 64:1-509; 55 figs. (with subfigs.), 4 pls.
- 1943a. New or little-known Tipulidae (Diptera). LXVI. Neotropical species. *Ann. Mag. Nat. Hist.* (11)10:217-245; 5 text figs.
- 1943b. New species of crane-flies from South America. Part X. (Tipulidae: Diptera). *Ann. Ent. Soc. Amer.* 36:103-127.
- 1943c. New or little-known Tipulidae from Venezuela (Diptera). Part I. *Bol. Ent. Venezolana* 2:17-26.
- 1943d. New or little-known Tipulidae from Venezuela (Diptera). Part II. *Bol. Ent. Venezolana* 2:125-144; 5 text figs.
- 1943e. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XVI. *Jour. New York Ent. Soc.* 51:199-212.
- 1943f. Records and descriptions of Brazilian Tipulidae (Dipt.). Part IX. *Rev. Ent.* 14:291-306; 14 text figs.
- 1943g. New or little-known Tipulidae (Diptera). LXVII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)10:522-552; 11 text figs.
- 1943h. Records and descriptions of Tipulidae from tropical America (Diptera). Part V. *Rev. Ent.* 14:485-502; 19 text figs.
- 1943i. New or little-known Tipulidae (Diptera). LXVIII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)10:738-765; 14 text figs.
- 1944a. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XVII. *Jour. New York Ent. Soc.* 52:45-57.
- 1944b. New or little-known Tipulidae (Diptera). LXIX. Neotropical species. *Ann. Mag. Nat. Hist.* (11)11:154-182; 13 text figs.
- 1944c. New or little-known Tipulidae from Venezuela (Diptera). III. *Bol. Ent. Venezolana* 3:35-50; 8 text figs.
- 1944d. New species of crane-flies from South America. Part XI. (Tipulidae, Diptera). *Ann. Ent. Soc. Amer.* 37:298-322.
- 1944e. A noteworthy undescribed crane-fly from tropical America (Diptera, Tipulidae). *Bull. Brooklyn Ent. Soc.* 39:92-93.
- 1944f. Records and descriptions of Brazilian Tipulidae (Dipt.). Part X. *Rev. Ent.* 15:19-34; 6 text figs.
- 1944g. New or little-known Tipulidae (Diptera). LXX. Neotropical species. *Ann. Mag. Nat. Hist.* (11)11:284-311; 7 text figs.
- 1944h. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XVIII. *Jour. New York Ent. Soc.* 52:369-383.
- 1944i. New or insufficiently-known crane-flies from Chile (Family Tipulidae, Order Diptera). Part I. *Bol. Sanidad Vegetal* 3:117-138; 11 text figs.
- 1944j. New or little-known Tipulidae from Venezuela (Diptera). IV. *Bol. Ent. Venezolana* 3:143-160; 5 text figs.



- 1944k. New or little-known Tipulidae (Diptera). LXXI. Neotropical species. *Ann. Mag. Nat. Hist.* (11)11:586-616.
- 1944L. New or little-known Tipulidae from Venezuela (Diptera). Part V. *Bol. Ent. Venezuelana* 3:171-192; 5 text figs.
- 1944m. Records and descriptions of Brazilian Tipulidae (Dipt.). Part XI. *Rev. Ent.* 15:292-312; 8 text figs.
- 1945a. New or little-known Tipulidae (Diptera). LXXII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)12:8-38.
- 1945b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XIX. *Jour. New York Ent. Soc.* 53:49-61.
- 1945c. New or little-known Tipulidae (Diptera) from São Paulo, Brasil — Part II. *Papéis Avulsos do Dept. de Zool.* 6:241-282; 22 text figs.
- 1945d. New species of crane-flies from South America. Part XII. (Diptera: Tipulidae). *Ann. Ent. Soc. Amer.* 38:256-280.
- 1945e. Records and descriptions of Brazilian Tipulidae (Dipt.). Parts XII and XIII. *Rev. Ent.* 16:210-243; 7 text figs.
- 1945f. New or insufficiently-known crane-flies from Chile (Family Tipulidae, Order Diptera). Part II. *Agr. Tecnica* 5:5-23; 12 text figs.
- 1945g. New or little-known Tipulidae from Venezuela (Diptera). Part VI. *Bol. Ent. Venezuelana* 4:59-80; 6 text figs.
- 1945h. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XX. *Jour. New York Ent. Soc.* 53:279-291.
- 1945i. New or little-known Tipulidae (Diptera). LXXIII. Neotropical species. *Ann. Mag. Nat. Hist.* (11)12:234-264; 9 text figs.
- 1945j. New or little-known Tipulidae (Diptera) from São Paulo, Brasil. Part III. *Papéis Avulsos do Dept. de Zool.* 7:1-44; 17 text figs.
- 1945k. Notes on the tropical American species of the genus *Tipula* Linnaeus (Tipulidae, Diptera). Part I. *Rev. Ent.* 16:330-356; 9 text figs.
- 1945L. Records and descriptions of Tipulidae from tropical America (Diptera). Part VI. *Rev. Ent.* 16:373-393; 22 text figs.
- 1945m. Notes on the tropical American species of the genus *Tipula* Linnaeus (Tipulidae, Diptera). Part II. *Rev. Ent.* 16:415-440; 25 text figs.
- 1945n. New or little-known Tipulidae (Diptera). LXXIV. Neotropical species. *Ann. Mag. Nat. Hist.* (11)12:390-419.
- 1945o. New or little-known Tipulidae (Diptera). LXXV. Neotropical species. *Ann. Mag. Nat. Hist.* (11)12:579-609.
- 1945p. New or little-known Tipulidae (Diptera). LXXVI. Neotropical species. *Ann. Mag. Nat. Hist.* (11)12:734-765.
- 1946a. Studies on the crane-flies of Mexico. Pt. VIII. (Order Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 39:119-139.
- 1946b. Records and descriptions of Mexican crane-flies (Dipt. Tipulidae). Pt. I. *Anales de la Escuela Nacional Ciencias Biológicas* 4 (for 1945):213-254; 28 text figs.
- 1946c. Undescribed species of Tipulidae from the State of São Paulo, Brasil (Diptera). *Livro Homenagem a R. F. d'Almeida, São Paulo* 1:1-10; 1 pl. (8 figs.).




- 1946d. Notes on the tropical American species of the genus *Tipula* Linnaeus (Tipulidae, Diptera). Pt. III. *Rev. Ent.* 17:172-201; 7 text figs.
- 1946e. Studies on the crane-flies of Mexico. Pt. IX. (Diptera, Superfamily Tipuloidea). *Ann. Ent. Soc. Amer.* 39:522-541.
- 1946f. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXI. *Jour. New York Ent. Soc.* 54:293-307.
- 1946g. Notes on the tropical American species of Tipulidae (Diptera) I. The genus *Teucholabis* Osten Saken. *Rev. Ent.* 17:375-400; 14 text figs.
- 1946h. New or little-known crane-flies. Part IV. *Encycl. Ent., Diptera* 10:91-104. (Note: This paper is an exact copy of 1941g).
- 1946i. Records and descriptions of North American crane flies (Diptera). Part VI. Tipuloidea of Arizona, New Mexico and Trans-Pecos Texas. I. *Amer. Mid. Nat.* 35:484-531.
- 1946j. Undescribed species of crane-flies from the eastern United States and Canada (Dipt.: Tipulidae). Part X. *Ent. News* 57:245-252.
- 1947a. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXII. *Jour. New York Ent. Soc.* 55:173-184.
- 1947b. Notes on the tropical American species of Tipulidae (Diptera). II. The primitive Eriopterini: *Sigmatomera*, *Trentepohlia*, *Gnophomyia*, *Neognophomyia*, *Gonomyia*, and allies. *Rev. Ent.* 18:65-100; 31 text figs.
- 1947c. Notes on the tropical American species of Tipulidae (Diptera). III. The specialized Eriopterini: *Rhabdomastix*, *Cryptolabis*, *Erioptera*, *Molophilus*, *Styringomyia*, *Toxorhina*, and allies. *Rev. Ent.* 18:317-360; 32 text figs.
- 1947d. New or little-known Tipulidae from Venezuela (Diptera). Pt. VII. *Bol. Ent. Venezolana* 6:37-54; 5 text figs.
- 1947e. New or little-known Tipulidae from Venezuela (Diptera). Pt. VIII. *Bol. Ent. Venezolana* 6:55-74; 14 text figs.
- 1947f. New or little-known Tipulidae from Venezuela (Diptera). Pt. IX. *Bol. Ent. Venezolana* 6:74-105; 13 text figs.
- 1948a. Notes on the tropical American species of Tipulidae (Diptera). IV. The primitive Hexatomini: *Paradelphomyia*, *Austrolimnophila*, *Epiphragma*, *Lecteria*, *Polymera*, and allies. *Rev. Ent.* 19:149-190; 33 text figs.
- 1948b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXIII. *Jour. New York Ent. Soc.* 56:137-148.
- 1948c. Notes on the tropical American species of Tipulidae (Diptera). V. The specialized Hexatomini: *Limnophila*, *Shannonomyia*, *Gynoplistia*, *Hexatoma*, *Atarba*, *Elephantomyia*, and allies. *Rev. Ent.* 19:509-556; 36 text figs.
- 1949a. New species of crane-flies from South America. Pt. XIII. (Diptera: Tipulidae). *Ann. Ent. Soc. Amer.* 42:101-119.
- 1949b. New or little-known Tipulidae (Diptera) from São Paulo, Brasil. Pt. IV. *Rev. Ent.* 20:461-497; 26 text figs.
- 1949c. Records and descriptions of Tipulidae from tropical America (Diptera). Pt. VII. *Rev. Ent.* 20:589-616; 40 text figs.
- 1949d. Records and descriptions of Neotropical crane-flies (Diptera). XXIV. *Jour. New York Ent. Soc.* 57:253-265.

- 1950a. The Tipulidae (Order Diptera) of Rancho Grande, north-central Venezuela. *Zoologica* 35:33-56; 39 text figs.
- 1950b. Notes on the tropical American species of Tipulidae (Diptera). VI. The tribe Limoniini, genus *Limonia*: Subgenera *Limonia*, *Neolimnobia*, *Discobola*, and *Rhipidia*. *Rev. Ent.* 21:161-221; 42 text figs.
- 1951a. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXV. *Jour. New York Ent. Soc.* 59:99-110.
- 1951b. Notes on the tropical American species of the genus *Tipula* Linnaeus (Tipulidae, Diptera). Part IV. *Rev. Ent.* 22:265-314; 24 text figs.
- 1951c. New or insufficiently-known crane-flies from Chile (Family Tipulidae, Order Diptera). *Agr. Tecnica* 11:5-19; 10 text figs.
- 1951d. New or little-known Tipulidae from the Andes Mountains (Diptera). *Rev. Chil. Ent.* 1:5-14; 1 pl. (6 figs.).
- 1952a. New species of crane-flies from South America. Part XIV. *Ann. Ent. Soc. Amer.* 45:119-135.
- 1952b. New or insufficiently-known crane-flies from Chile (Family Tipulidae, order Diptera). *Agr. Tecnica* 11(1951):99-118; 15 text figs.
- 1952c. Los Insectos de las Islas Juan Fernandez. 5. Tipulidae (Diptera). *Rev. Chil. Ent.* 2:35-80; 47 text figs.
- 1952d. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXVI. *Jour. New York Ent. Soc.* 60:245-254.
- 1952e. New or little-known Tipuloidea (Diptera) from Argentina. Part I. *Acta Zool. Lilloana* 10(1951):75-94; 9 text figs.
- 1952f. The oriental Tipulidac in the collection of the Indian Museum. Part III. *Rec. Indian Mus.* 50:321-358; 15 figs. with subfigs.
- 1953a. Notes on the Tipulidae of Ecuador (Order Diptera). Part II. *Rev. Ecuat. Ent. Paras.* 1(2):6-15; 8 text figs.
- 1953b. New or insufficiently-known crane-flies from Chile (Family Tipulidac, Order Diptera). Part V. *Agr. Tecnica* 13:1-23; 20 text figs.
- 1953c. Notes on the Tipulidae of Ecuador (Order Diptera). Part III. *Rev. Ecuat. Ent. Paras.* 1(3):29-41; 9 text figs.
- 1953d. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXVII. *Jour. New York Ent. Soc.* 61:147-157.
- 1953e. New species of cranc-flies from South America. Part XV. (Diptera: Tipulidae). *Ann. Ent. Soc. Amer.* 46:599-617; 9 text figs.
- 1953f. Notes on the Tipulidae of Ecuador (Order Diptera). Part IV. *Rev. Ecuat. Ent. Paras.* 1(4):77-95; 1 pl. (9 figs.).
- 1953g. Records and descriptions of Chilean crane-flies, Tipulidae (Diptera). Part I. *Rev. Chil. Ent.* 3:7-21; 12 text figs.
- 1954a. Records and descriptions of Brazilian crane-flies (Diptera, Tipulidae). *Rev. Brasil. Ent.* 1:31-46; 9 text figs.
- 1954b. Notes on the Tipulidae of Ecuador (Order Diptera). Part V. *Rev. Ecuat. Ent. Paras.* 2:51-67; 2 pls. (23 figs.).
- 1954c. Notes on the Tipulidae of Ecuador (Order Diptera). Part VI. *Rev. Ecuat. Ent. Paras.* 2:231-246; 2pls. (18 figs.).
- 1954d. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXVIII. *Jour. New York Ent. Soc.* 62:139-152.
- 1956a. Undescribed crane-flies from tropical America (Diptera, Tipulidae). Part I. *Bull. Brooklyn Ent. Soc.* 51:22-25.



- 1956b. Records and descriptions of Neotropical crane-flies (Tipulidae, Diptera). XXIX. *Jour. New York Ent. Soc.* 63(1955):111-122.
1958. New or little-known South American crane-flies (Diptera, Tipulidae). *Rev. Chil. Ent.* 5:193-204; 10 text figs.
1959. Insecta Amapaensia. Diptera: Tipulidac. *Studia Ent.* 2:445-455; 6 text figs.
1960. New exotic crane-flies (Tipulidae: Diptera). Part I. *Ent. News* 71:15-20.
- 1962a. Undescribed species of Nematoceros Diptera. Part XI. *Bull. Brooklyn Ent. Soc.* 57:33-39.
- 1962b. New or little-known Tipuloidca (Diptera) from Argentina. II. *Acta Zool. Lilloana* 18:335-358; 23 text figs.
- 1962c. Beiträge zur Kenntnis der Insektenfauna Boliviens XVII. Diptera II. The crane-flies (Tipulidae, Diptera). *Veröff. Zool. Staatssamml. München* 7:9-159; 140 text figs.
- 1962d. The crane-flies of the Galapagos Islands. *Opuscula Zool.* (Munich) 61:1-5; 6 text figs.
- 1964a. The crane-flies of Jamaica (Diptera, Tipulidae). *Bull. Inst. Jamaica Sci. Ser.* 14, pp. 86; 97 text figs.
- 1964b. New or little-known Tipulidae from eastern Asia (Diptera). LIV. *Philippine Jour. Sci.* 93:277-324; 5 pls. (54 figs.).
- 1965a. New species of crane-flies from tropical America. (Diptera: Tipulidae). I. *Jour. Kansas Ent. Soc.* 38:401-407.
- 1965b. Undescribed or little-known tropical American Tipulidae (Diptera). Part I. *Studia Ent.* 8:337-352; 8 text figs.
- 1965c. New subgenera and species of crane-flies from California (Diptera: Tipulidae). *Pacific Insects* 7:333-386; 33 figs.
- 1965d. In *A Catalog of the Diptera of America north of Mexico*, by Stone, Sabrosky, Wirth. Foot and Coulson, pp. i-iv, 1-1696 (Tipulidae, pp. 16-90).
- 1966a. Records and descriptions of Tipulidac from South America (Diptera). *Papéis Avulsos do Dept. de Zool., S. Paulo* 18(art. 18): 221-229.
- 1966b. New species of crane-flies from tropical America (Diptera: Tipulidae). II. *Jour. Kansas Ent. Soc.* 39:695-705.
- 1966c. Undescribed or little-known tropical American Tipulidae (Diptera). Part II. *Studia Ent.* 9:465-500; 22 text figs.
- 1967a. Notes on the Tropical American species of Tipulidae (Diptera). VII. The tribe Limoniini, genus *Limonia*, concluded; *Helius*, *Orimarga* and others; Tribe Pediciini, Subfamily *Cylindrotominae*. *Studia Ent.* 10:277-352; 78 text figs.
- 1967b. Undescribed or little-known Tropical American Tipulidae (Diptera). Part III. *Studia Ent.* 10:449-472; 15 text figs.
- 1967c. New species of crane-flies from tropical America (Diptera: Tipulidae). III. *Jour. Kansas Ent. Soc.* 40:373-381.
- 1967d. The crane flies of California. *Bull. California Insect Survey*, 1967:1-263; 1 pl., 524 figs., 106 maps.
- 1968a. New or little-known Tipulidae from Chile and Peru (Diptera: Tipulidae). Part I. *Rev. Chil. Ent.* 6:21-36, figs.
- 1968b. Records and descriptions of Tipulidae from South America (Diptera). II. *Papéis Avulsos do Dept. de Zool., S. Paulo*, 21(art. 8):87-97.



- 1968c. New species of crane flies from Tropical America. IV. (Diptera: Tipulidae). *Jour. Kansas Ent. Soc.* 41:210-220.
- BELLARDI, L.
 1859. *Saggio di ditteologia messicana. Parte I. Pt. 1*, 80 pp., 2 pls. Torino. (Also in *Mem. R. Accad. Sci. Torino*, 19:201-277, 2 pls.; 1861).
 1861. *Saggio di ditteologia messicana. Parte II. Pt. 2*, 99 pp., 2 pls. Torino. (Also in *Mem. R. Accad. Sci. Torino*, 21:103-199, 2 pls.; 1864).
- BERENDT, G. C.
 1945. *Die im Bernstein befindlichen organischen Reste*, 1:57.
- BERGROTH, E.
 1915. Some tipulid synonymy. *Psyche* 22:54-59.
- BIGOT, J. M. F.
 1854. Essai d'une classification générale et synoptique de l'ordre des Insectes Diptères. IIIc. *Ann. Soc. Ent. France* (3)2:447-482.
 1888 (1891). *Miss. Scient. Cap Horn, Zool., Diptera*, 6:1-45; 4 pls.
- BLANCHARD, E.
 1852. Orden IX. Dipteros, pp. 327-468, in C. Gay, ed., *Historia fisica y politica de Chile, Zoologia* 7, 471 pp., illus.
- BRÈTHES, JUAN
 1909. Dipteros e Himenopteros de Mendoza. *Anales Mus. Nac. Buenos Aires* (3)19:85-105.
- BRUCH, CARLOS
 1939a. Contribución al conocimiento de los Tipulidos Argentinos (Diptera). *Physis* 17:1-28; 11 pls. (83 figs.).
 1939b. Catalogo de los Tipulidos de la Republica Argentina. *Physis* 17:29-55.
- BRUNETTI, E.
 1911. Revision of the Oriental Tipulidae with descriptions of new species. *Rec. Indian Mus.* 6:231-314.
 1912. Diptera Nematocera (excluding Chironomidae and Culicidae). Diptera, [Vol. 1], 581 pp., 12 pls., 44 figs., [in Blanford, W. T., ed.]. *Fauna of British India*. London.
 1918. Revision of the Oriental Tipulidae with descriptions of new species. Part II. *Rec. Indian Mus.* 15:255-340, 2 pls., (35 figs.), 7 text figs.
 1920. Catalogue of Oriental and south Asiatic Nematocera. *Rec. Indian Mus.* 17:1-300.
- BYERS, G. W.
 1966. A crane fly new to the known fauna of Hawaii (Diptera: Tipulidae). *Jour. Kansas Ent. Soc.* 39:708-711.
- COQUILLET, D. W.
 1910. The Type-species of the North American genera of Diptera. *Proc. U. S. Natl. Mus.* 37:499-647.
- CURRAN, C. H.
 1934. *The Families and Genera of North American Diptera*, 512 pp., 235 figs., 2 pls., New York. (revised 1964).



CURTIS, J.

1825. *British entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland.* 2(1825): pls. 51-98. London.
1833. *Idem*, Vol. 10: pls. 434-481. London.

DALE, J. C.

1842. Descriptions, &c., of a few rare or undescribed species of British Diptera, principally from the collection of J. C. Dale, Esq., M. A., F. L. S., &c. *Ann. Mag. Nat. Hist.* 8:430-433.

EDWARDS, F. M.

1914. A revision of the Tipulid genus *Styryngomyia* Lw. *Trans. Ent. Soc. London* 1914:206-227; 81 figs.
1918. Two new Diptera from Trinidad. *Ann. Mag. Nat. Hist.* (9) 1:424-425; fig.
1919. Diptères Nematocères. in "Mission du service géographique de l'Armée pour la mesure d'un arc de Méridien Equatorial en Amérique du Sud sous le contrôle scientifique de l'Académie des Sciences, 1899-1906." 10, fasc. 2:143-162; 26 figs.
1923. A preliminary revision of the crane-flies of New Zealand (Anisopodidae, Tanyderidae, Tipulidae). *Trans. New Zealand Inst.* 54:265-352; 10 pls. (180 figs.). (wings, male hypopygia).
1927. A Tipulid apparently mimicking a mosquito. *Proc. Ent. Soc. London* 2:45-46.
1931. Some suggestions on the classification of the genus *Tipula* (Diptera, Tipulidae). *Ann. Mag. Nat. Hist.* (10)8:73-82.
1934. Oxford University Expedition to British Guiana, 1929. Diptera Nematocera. *Ann. Mag. Nat. Hist.* (10)14:632-635.

ENDERLEIN, G.

1906. Eine neue Fliegengattung von den Falklands-Inseln. 9. Beitrag zur Kenntnis der antarktischen Landarthropoden. *Zool. Anzeig.* 29:70-72; 2 figs.
1912. Studien über die Tipuliden, Limoniiden, Cylindrotomiden und Ptychopteriden. *Zool. Jahrb., Abt. f. System., Geog. u. Biol. Tiere* 32:1-88, 51 figs.
1917. Dipterologische Studien. XVI. *Zool. Anzeig.* 49:57-71; 10 figs.
1940. Die Dipterenfauna der Juan-Fernandez-Inseln und der Osterinsel. In Skottsberg, C. — *The Natural History of Juan Fernandez and Easter Islands*, *Zool.* 3, 60:643-680; 49 figs.

FABRICIUS, J. C.

1787. *Mantissa insectorum sistens species nuper detectas*, 2, 382 pp. Hafniae [= Copenhagen].
1794. *Entomologia systematica emendata et aucta*, 4, 472 pp. Hafniae [= Copenhagen].
1805. *Systema Antliatorum secundum ordines, genera, species*, 373 pp., + 30 pp. Brunsvigae [= Brunswick].

FREEMAN, P.

1962. Diptera Nematocera from Gough Island, South Atlantic. *Proc. r. Ent. Soc. London* 31 (B):77-80; 2 figs.



GOETGHEBUER, M. & A. L. TONNOIR

1920. Catalogue raisonné des Tipulidae de Belgique, Eriopterini. *Bull. Soc. Ent. de Belgique* 2:104-112, 131-147, 5 figs., 2 pls. (32 figs.).

GUÉRIN-MÉNEVILLE, F. E.

1831. Pls. 20-21 (= livrs. 23-24, part), *In* Duperrey, L. I., ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille. Zoologie, Atlas, Insects*, 21 pls. Paris.
[1838]. Première division. Crustacés, arachnides et insects. Div. 1, pp. 1-216, 217-319 (=livrs. 25, 26), *in* Duperrey, L. I., ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille. Zoologie* 2(2). Paris

HALDAY, A. H.

1833. Catalogue of Diptera occurring about Holywood in Downshire. *Ent. Mag.* (London) 1:147-180.
1856. Addenda and corrigenda. Pp. xi-xv. *In* *Insecta Britannica Diptera* Vol. 3 (by Walker), xxiv + 352 pp.; pls. 21-30. London.

HENDEL, F.

1908. Nouvelle classification des mouches à deux ailes (Diptera L.), d'après un plan tout nouveau par J. G. Meigen, Paris, an VIII (1800 v.s.). Mit cinem Kommentar. *Verh. zool.-bot. Ges. Wien* 58:43-69.

HUNTER, W. D.

1900. A catalogue of the Diptera of South America. Part I. (Bibliography: Nemocera). *Trans. Amer. Ent. Soc.* 26:260-298.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. Internatl. Comm. Zool. Nomencl., *Bull. Zool. Nomencl.* 20:339-342.

JACOBS, J. C.

1900. Diptères. *Ann. Soc. Ent. de Belgique* 44:106, 107.
1906. *Resultats Voy. du S. Y. Belgica. Zool. Diptères*, 67-71 pp.; pls. 3-5 (color).

JAENNICKE, F.

1867. Neue exotische Dipteren. *Senckenb. Naturf. Gesell. Abhandl.* 6:311-408; pls. 43-44.

JOHNSON, C. W.

1913. The Dipteran fauna of Bermuda. *Ann. Ent. Soc. Amer.* 6:443-452; 2 figs.

KERTÉSZ, K.

1902. *Catalogus dipterorum hucusque descriptorum* 2:359 pp. Lipsiae, Budapestini [= Leipzig, Budapest].

LATREILLE, P. A.

1809. *Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata.* Vol. 4, 399 pp.; 4 pls.
1810. *Considérations générales sur l'ordre naturel des animaux.* 444 pp. Paris.



LEPELETIER, A. L. M., J. G. A. SERVILLE, & F. E. GUÉRIN-MÉNEVILLE

1828. Entomologie, ou histoire naturelle des Crustacés, des Arachnides et des Insectes, Vol. 10:345-833. In Société de gens de Lettres, de Savans et d'Artistes, *Encyclopédie méthodique. Histoire naturelle*. Paris.

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae. Ed. 10. Vol. 1*, 824 pp. Stockholm

LOEW, H.

1845. (Dipterologischer Beitrag) (I). *K. Friedrich-Wilhelms-Gymnasium zu Posen, Öffentl. Prüf. d. Schüler*, pp. 1-52; 1 pl.

1850. Dipterologische Beiträge. Vierter Theil. *K. Friedrich-Wilhelms-Gymnasium zu Posen, Öffentl. Prüf. d. Schüler*, pp. 1-40; 1 pl.

1851. Beschreibung einiger neuen Tipularia terricola. *Linnaea Ent.* 5:385-406; pl. 2.

1861. Diptera aliquot in insula Cuba collecta. *Wien. Ent. Monatschr.* 5:33-43.

1863. Diptera Americac septentrionalis indigena. Centuria quarta. *Berlin. Entomol. Zeitschr.* 7:275-326.

1864. *Diptera Americac septentrionalis indigena 1*; 266 pp. (reprinting Centuries 1-5).

1865. Ueber die bisher beschriebenen europäischen Anisomera-Arten. *Zeitschr. Gesammt. Naturwissenschaften* 24:424.

1869. Diptera Americac septentrionalis indigena. Centuria octava. *Berlin. Entomol. Zeitschr.* 13:1-52.

1872. *Diptera Americac septentrionalis indigena 2*:300 pp. (reprinting Centuries 6-10).

1873. Monographs of the Diptera of North America. Pt. III. *Smithsonian Misc. Coll.* 11(3):1-351 (pub. 256), illus.

MACQUART, J.

1834. *Histoire naturelle des Insectes, Diptères 1*:578 pp. 12 pls. Paris.

1838a. Diptères exotiques nouveaux ou peu connus. *Mém. Soc. Roy. des Sci., de l'Agr. et des Arts, Lille 1838(2)*:9-225, 25 pls. (Also published separately as Vol. 1, Pt. 1, pp. 5-221, 25 pls., Paris, 1838).

1838b. *Diptères exotiques nouveaux ou peu connus 1(2)*:5-207, 14 pls. Paris. (Also published in *Mém. Soc. Roy. des Sci., de l'Agr. et des Arts, Lille 1838(3)*:121-323, 14 pls., 1839).

1839. In Webb, P. B. & S. Berthelot, eds., *Histoire naturelle des Iles Canaries*. Vol. 2, pt. 2: Zoologie, (Sect. 6): Entomologie, 119 pp., 7 pls. Paris.

MATSUMURA, S.

1916. *Thousand Insects of Japan. Additamenta*. Vol. 2, pp. 185-474; pls. 16-25. Tokyo.

MEIGEN, H. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*. 40 pp. Paris.

1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Illiger's Mag. f. Insektenkunde* 2:259-281.

1818. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 1, xxxvi + 333 pp., pls. 1-11. Aachen.

1830. Same. Vol. 6, iv + 401 pp.; pls. 55-66. Hamm.



MEIJERE, J. C. H. DE

- 1911a. Studien über südostasiatischen Dipteren. V. *Tijd. v. Ent.* 54:21-79; pls. 1-5.
1911b. The same. VI. *Tijd. v. Ent.* 54:258-432; pls. 18-22.
1913. The same. VII. *Tijd. v. Ent.* 56:317-354; pls. 15-17.
1916. Beiträge zur Kenntnis der Dipteren-Larven und Puppen. *Zool. Jahrb. Sys.* 40:177-322; 14 pls. (181 figs.).
1946. *Oxydiscus* (Diptera) changed in *Oxyrhiza*. *Entomologische Berichten* 12:68.

MILLER, DAVID

1945. Generic name changes in Diptera. *Proc. R. Ent. Soc. London* (B)14:72.

NEEDHAM, J. G.

1908. Report of the entomological field station conducted at Old Forge, N. Y., in the summer of 1905. *New York State Mus. Rep.* 61(vol. 2):156-248; figs. 2-16, pls. 4-32.

OSTEN SACKEN, C. R.

1859. New genera and species of North American Tipulidae with short palpi, with an attempt at a new classification of the tribe. *Proc. Acad. Nat. Sci. Philadelphia 1859*:197-256, 1 fig., 2 pls.
1861. Description of nine new North American Limnobiaceae. *Proc. Acad. Nat. Sci. Philadelphia 1861*:287-292.
1865. Description of some new genera and species of North American Limnobia. Part I. *Proc. Ent. Soc. Philadelphia 4*:224-242.
1869. Monographs of the Diptera of North America. Part IV. Smithsonian Inst., *Smithsn. Misc. Collect.* 8 (= pub. 219): 1-345, 7 figs., 4 pls.
1878. Catalogue of the described Diptera of North America. Smithsonian Inst., *Smithsn. Misc. Collect.* 16(2):1-276.
1886a. Diptera, Vol. I (part). Pp. 1-24, 25-48, 49-72, 73-104, 105-128, pls. 1-2 (cont.). In Godman, F. D. & O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, vol. 1, 378 pp. 6 pls. London.
1886b. Studies on Tipulidae. Part I. Review of the published genera of the Tipulidae longipalpi. *Berlin. Ent. Ztschr.* 30:153-188.
1887a. Diptera, Vol. I (part, concl.). Pp. 129-160, 161-176, 177-208, 209-216, pl. 3. In Godman, F. D. and O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, vol. 1, 378 pp. 6 pls. London.
1887b. Studies on Tipulidae. Part II. Review of the published genera of the Tipulidae brevipalpi. *Berlin. Ent. Ztschr.* 31:163-242.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *K.-k. Zool.-Bot. Gesell. Wien. Verhandl.* 15 (Abhandl.): 595-782, 7 pls.

PICADO, C.

1913. Les bromeliacées epiphytes. *Bull. Sci. France et Belgique* 47 (3):215-360. (Ref. 356-357, color pl. 13, figs. 1, 1a, 2, 4; text figs. 50, 51).

PIERRE, C.

1924. Tipulidae nouveaux. *Encycl. Entomologique, Diptera*, 1, fasc. 2:79-93, 60 figs.

- REED, EDWIN C.
1888. Catalogo de los insectos dipteros de Chile. *Anal. Univ. Chile* 73:i-v, 1-46.
- REED, EDWIN P.
1929. Notable dimorfismo sexual de un Tipúlido Chileno. *Rev. Chil. Hist. Nat.* 33:384-385, fig.
- RIEDEL, M. P.
1921. Dr. L. Zürcher's Dipterenausbeute aus Paraguay. *Nematocera polyncura* (Familie Tipulidae). *Arch. für Naturgeschichte* 87, Abt. A. Heft 3:123-127, 6 figs.
- KÖDER, V. VON
1885. Dipteren von der Insel Portorico. *Stett. Ent. Zeitg.* 46:337-349.
1886. Dipteren von den Cordilleren in Columbien. *Stett. Ent. Zeitg.* 47:257-270.
- ROGERS, J. S.
1927a. Notes on the life history, distribution and ecology of *Dio-trepha mirabilis* Osten Sacken. *Ann. Ent. Soc. Amer.* 20:23-36; 9 figs.
1927b. Notes on the biology of *Atarba picticornis* Osten Sacken — Tipulidae — Diptera. *Florida Entomologist*, 10:49-55; 6 figs.
1933. Contributions toward a knowledge of the natural history and immature stages of the crane-flies. I. The genus *Polymera* Wiedemann. *Occas. Pap. Mus. Zool., Univ. Michigan*, 268:1-13; pl. 1 (adult, immature stages), pl. 2, 8 figs. (immature stages).
1949. The life history of *Megistocera longipennis* (Macquart) (Tipulidae, Diptera), a member of the neuston fauna. *Occas. Pap. Mus. Zool., Univ. Michigan*, 521:1-17, 2 pls., 7 figs.
- RONDANI, CAMILLO
1842. Note sur un nouveau genre d'insecte Diptère subaptère (*Pterelachisus Bertei*). *Revue Zool.* 5:243 [Magas. Zool (Guérin), 12, No. 106 (color. fig.)].
1850. Osservazioni sopra alquante specie di Esapodi Ditteri del Musco Torinese. *Nuov. Annali delle Scienze Naturali e Rendiconto Bologna* (3)2:165-197.
- ROSSI, F. W.
1848. *Sstematisches Verzeichniss der zweiflügelichten Insecten (Diptera) des Erzherzogthumes Österreich.* iii-x, 86 pp. Wien.
- SAY, THOMAS
1823. Descriptions of dipterous insects of the United States. *Jour. Acad. Nat. Sci. Philadelphia* 3:9-54, 73-104.
1829. Descriptions of North American Dipterous insects. *Jour. Acad. Nat. Sci. Philadelphia* 6:149-178. (Also published in J. L. Leconte ed. *The complete writings of Thomas Say on the entomology of North America*, Vol. 2. 814 pp., 1859. New York).
- SCHINER, I. R.
1863. Vorläufiger Commentar zum dipterologischen Theile der Fauna Austriaca V. (concl.). *Wien. Ent. Monatschr.* 7:217-226.
1866. Bericht über die von der Weltumseglungsreise der k. Fregatte Novara mitgebrachten Dipteren. *K.-k. Zool. Bot. Gesell. Wien. Verhand.* 16 (Abhandl.): 927-934.


1868. Diptera (Art. 1), 388 pp., 4 pls. In *Reise der österreichischen Fregatte Novara um die Erde*, Zool. Vol. 2, Abt. 1.
- SINTENIS, F.
1889. Über *Limnophila pilicornis* Zett. *Sitzber. Naturf. Gesell. Univ. Dorpat* 8:396-398.
- SKUSE, F. A. A.
1890a. Diptera of Australia. Part VII. The Tipulidae brevivalpi. *Proc. Linn. Soc. New South Wales* (1889) (2)2 (whole ser., 13): 757-892, 4 pls. (70 figs.). (Preprint Sept. 25, 1889).
1890b. Diptera of Australia. Part VIII. The Tipulidae longivalpi. *Proc. Linn. Soc. New South Wales* (1890) (2)5:53-139, 3 pls. (52 figs.).
- SPEISER, P.
1909. 4. Orthorhapha, pp. 31-65, figs. 1-15. In *Sjöstedt, Y., ed., Wissenschaftliche Ergebnisse der schwedischen zoologischen Expedition nach dem Kilimandjaro*. Vol. 2, Pt. 10: Diptera, 202 pp., 19 figs., 1 pl. Stockholm.
- STEPHENS, J. F.
1829. *A systematic catalogue of British insects: Being an attempt to arrange all the hitherto discovered indigenous insects in accordance with their natural affinities*. Vol. 2, 388 pp. London.
- STUARDO ORTIZ, C.
1946. *Catalogo de los dipteros de Chile*, 253 pp. Santiago.
- VANE-WRIGHT, R. T.
1967. A re-assessment of the genera *Holorusia* Loew (*Ctenacrosceles* Enderlein), *Ischnotoma* Skuse and *Zelandotipula* Alexander (Diptera. Tipulidae), with notes on their phylogeny and zoogeography. *Jour. Nat. Hist.* 1(4):511-549, 43 figs.
- WALKER, FRANCIS
1837. Descriptions, etc., of the insects collected by Capt. P. P. King, R. N., F. R. S., in the survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. London for 1836*, 17:331-359.
1848. *List of the specimens of dipterous insects in the collection of the British Museum*. 1:1-229. British Museum (Nat. Hist.). London.
1856. Diptera (concl.). Vol. 1, pp. 415-474. In (Saunders, W. W., ed.), *Insecta Saundersiana*. London
1861. Characters of undescribed Diptera in the collection of W. W. Saunders (concl.). *Trans. Ent. Soc. London* (1858-1861), (2)5:297-334.
- WESCHÉ, W.
1910. On the new Tipulid subfamily Ceratocheilinae. *Jour. Linn. Soc. London, Zool.* 30:355-360, pl. 49.
- WESTWOOD, J. O.
1835. Insectorum novorum exoticorum (ex ordine dipterorum) descriptionis. *London and Edinb. Phil. Mag. and Jour. Sci. ser. 3*, 6:280-281, 447-449.
1840. Order XIII. Diptera Aristotle (Antliata Fabricius, Alteriptera Clairv.), pp. 125-128, 129-144, 145-158, in his *An introduction to the modern classification of insects. Synopsis of the genera of British insects*, 158 pp. London.



1881. Notae Dipterologicae. No. 5. Descriptions of new species of Exotic Tipulidae, with an annotated summary of species belonging to the same family, previously described. *Trans. Ent. Soc. London*, Pt. 3:363-385, pls. 17-19, with 14 figs.
- WIEDEMANN, C. R. W.
 1820. *Diptera exotica*. (Ed. 1), Pt. I, pp. i-xix + 1-42, 1 fig. Kiel.
 1821. *Diptera exotica*. (Ed. 1), Pt. II, pp. i-iv + 43-50 (cont.). Kiel.
 1824. *Munus rectoris in Acadcmina Christiana Abertina aditurus analceta entomologica ex Museo Regio Havniensi*, 60 pp., 1 pl. Kiliae [= Kiel].
 1828. *Aussereuropäische zwelflügelige Insekten*. 1:xxxii + 608 pp., 7 pls.
 1830. The same. 2:xii + 684 pp., 5 pls.
- WILLISTON, S. W.
 1896a. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896:253-446, 6 pls. (170 figs.).
 1896b. *Manual of the families and genera of North American Diptera*. Ed. 2. 167 pp.
 1900-01. Supplement (part). Pp. 217-248, 1900; pp. 249-264, 265-272, 273-296, 297-328, 329-332, pls. 4-5 + pl. 6, figs. 1-6, 1901. In Godman, F. D. and O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, vol. 1, 378 pp., 6 pls. London.
- WILP, F. M. VAN DER
 1881. Amerikaansche Diptera. *Tijdschr. v. Ent.* 24:141-168, pl. 15.
 1891. Eenige uitlandsche Diptera. *Tijdschr. v. Ent.* 34:193-218, 2 pl.
- ZETTERSTEDT, J. W.
 1837. Conspectus familiarum generum et specierum dipterorum, in fauna insectorum Lapponica descriptorum. *Isis* (Oken's) 1837:28-67.
 1838. Dipterologia Scandinaviae. Sect. 3: Diptera, pp. 477-868. In *Insecta Lapponica*. vi + 1,140 pp. Leipzig.

Index


aberrans (Schiner), 1868, Limonia (Peripheroptera)	4.72
Abitagua Alexander, 1944	4.93
abitaguae Alexander, 1945, Braehypremna	4.6
abltagual (Alexander), 1941 (Erioptera), Cheilotrichia (Empeda) .	4.119
abnormis Alexander, 1914, Lecteria (Psaronius)	4.106
abortiva Alexander, 1966, Shannonomyia (Shannonomyia)	4.114
abortiva Alexander, 1914, Tipula (Eumierotipula)	4.21
abra Alexander, 1949, Shannonomyia (Shannonomyia)	4.114
abselssa Alexander, 1946, Tipula (Lunatipula)	4.32
absona Alexander, 1937, Tipula (Eumierotipula)	4.21
abstrusa Alexander, 1929, Limnophila (Limnophila)	4.108
aeanthias Alexander, 1941, Gonomyia (Progonomyia)	4.141
aeanthostylus Alexander, 1944, Helius	4.45
aeacipitrina Alexander, 1946, Tipula (Eumierotipula)	4.21
aeacumulatrix Alexander, 1951, Tipula (Eumierotipula)	4.21
aeacanthoides Alexander, 1929, Elnoretta	4.8
aeacricula Alexander, 1931, Gnophomyia	4.128



<i>acrisissima</i> Alexander, 1944, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>aeroleuca</i> Alexander, 1964, <i>Orlmarga</i> (<i>Diotrepha</i>)	4.81
<i>acuminata</i> Alexander, 1921, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>aeuminata</i> (Alexander), 1921 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Caenoglo-</i> <i>china</i>)	4.47
<i>aeunai</i> (Alexander), 1928 (<i>Erioceera</i>), <i>Hexatoma</i> (<i>Erioceera</i>)	4.100
<i>aeuproducta</i> Alexander, 1962, <i>Limonia</i> (<i>Dicranomyia</i>)	4.49
<i>aeutergata</i> Alexander, 1939, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.89
<i>aeutiloba</i> Alexander, 1937, <i>Dolichochepeza</i> (<i>Megistomastix</i>)	4.8
<i>acutissima</i> (Alexander), 1940 (<i>Oxydiseus</i>), <i>Paradelphomyia</i> (<i>Oxyr-</i> <i>hiza</i>)	4.109
<i>acutistyla</i> (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.41
<i>adara</i> Alexander, 1949, <i>Neognophomyia</i>	4.153
<i>adela</i> Alexander, 1949, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>adjusta</i> Alexander, 1949, <i>Gnophomyia</i>	4.128
<i>adspersa</i> (Wiedemann), 1818 (<i>Limnobia</i>), <i>Epiphragma</i> (<i>Epiphrag-</i> <i>ma</i>)	4.95
<i>adumbrata</i> Alexander, 1946, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>adunea</i> Alexander, 1921, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
<i>advena</i> (Alexander), 1928 (<i>Cryptolabis</i>) <i>Baeoura</i>	4.119
<i>aedon</i> Alexander, 1947, <i>Tipula</i> (<i>Eumierotipula</i>)	4.21
<i>aenigmatia</i> Alexander, 1929, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>aequabilis</i> (Alexander), 1923 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.60
<i>aequalis</i> Alexander, 1916, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
<i>aequatorialis</i> (Alexander), 1944 (<i>Oxydiseus</i>), <i>Paradelphomyia</i> (<i>Oxyrhiza</i>)	4.109
<i>aequidens</i> Alexander, 1938, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>aequinigra</i> Alexander, 1941, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.159
<i>aequinoetlalis</i> Alexander, 1937, <i>Sigmatomera</i> (<i>Sigmatomera</i>)	4.156
<i>acquispinosa</i> Alexander, 1926, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>aequitatorialis</i> Alexander, 1940, <i>Tipula</i> (<i>Microtipula</i>)	4.32
<i>aetherea</i> (Alexander), 1916 (<i>Erioceera</i>), <i>Hexatoma</i> (<i>Erioceera</i>)	4.100
<i>aetherea</i> (Bigot), 1888 (<i>Isehnotherix</i>), <i>Atarba</i> (<i>Isehnotherix</i>)	4.88
<i>affabilis</i> Alexander, 1952, <i>Limonia</i> (<i>Dicranomyia</i>)	4.49
<i>affabilis</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.32
<i>affinis</i> (Bellardi), 1859 (<i>Tipula</i>), <i>Nephrotoma</i>	4.16
<i>agglomerata</i> (Alexander), 1926 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ..	4.74
<i>aglaia</i> Alexander, 1945, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>aglala</i> Alexander, 1945, <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
<i>aglossa</i> Alexander, 1962, <i>Tipula</i> (<i>Eumierotipula</i>)	4.21
<i>agrippina</i> Alexander, 1945, <i>Tipula</i> (<i>Eumierotipula</i>)	4.21
<i>akestra</i> Alexander, 1965, <i>Tipula</i> (<i>Microtipula</i>)	4.32
<i>albicentra</i> Alexander, 1962, <i>Pedicia</i> (<i>Triephyphona</i>)	4.84
<i>albiditarsis</i> Alexander, 1953, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>alblfasciata</i> Macquart, 1838, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.21
<i>albimana</i> (Wiedemann), 1830 (<i>Tipula</i>), <i>Braehypremna</i>	4.6
<i>albitarsis</i> (Osten Sacken), 1887 (<i>Rhamphidia</i>), <i>Helius</i>	4.45
<i>albitarsis</i> Williston, 1896, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>alboapicalis</i> Alexander, 1928, <i>Limonia</i> (<i>Dicranomyia</i>)	4.49
<i>alboeostata</i> Macquart, 1846 (<i>Tipula</i>)	4.44
<i>albogeniculata</i> Alexander, 1926, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>albogeniculatus</i> Alexander, 1936, <i>Helius</i>	4.45
<i>albogenualis</i> Alexander, 1939, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>Aldrovandia</i> Enderlein, 1912	4.10




alcetro Alexander, 1945, Tipula (Microtipula)	4.32
alfaroi (Alexander), 1922 (Diceratomyia), Limonia (Diceratomyia) .	4.49
alleni (Alexander), 1913 (Pachyrhina), Nephrotoma	4.16
almeidai Alexander, 1946, Atarba (Atarba)	4.85
altandina Alexander, 1956, Limonia (Diceratomyia)	4.50
alticola (Alexander), 1925 (Erioptera), Cheilotrichia (Empeda) ..	4.119
alticola Alexander, 1944, Cryptolabis (Cryptolabis)	4.120
alticola Alexander, 1925, Elephantomyia (Elephantomyia)	4.94
alticola Alexander, 1944, Rhabdomastix (Rhabdomastix)	4.154
altivolans Alexander, 1942, Gonomyia (Progonomyia)	4.141
amabilis Alexander, 1928, Atarba (Atarba)	4.85
<i>Amalopsis</i> Haliday, 1856	4.84
amapana Alexander, 1959, Teucholabis (Teucholabis)	4.159
amara Alexander, 1951, Tipula (Microtipula)	4.32
amaryllis Alexander, 1944, Limonia (Caenolimonia)	4.48
amatrix Alexander, 1945, Teucholabis (Euparatropesa)	4.158
amatrix Alexander, 1942, Trentepohlia (Paramongoma)	4.169
amazona Alexander, 1921, Gonomyia (Paralipophleps)	4.140
amazonensis (Alexander), 1912 (Rhipidia), Limonia (Rhipidia) ..	4.75
amazonica (Alexander), 1920 (Diceratomyia), Limonia (Neo- limonia)	4.70
amazonica (Alexander), 1912 (Microtipula), Tipula (Microtipula) .	4.32
amazonica Westwood, 1881, Sigmatomera (Sigmatomera)	4.156
amazonicola (Alexander), 1920 (Eriocera), Hexatoma (Eriocera) .	4.100
ambigua Alexander, 1929, Limonia (Diceratomyia)	4.50
amblyphallos Alexander, 1966, Teucholabis (Teucholabis)	4.159
amblythrix Alexander, 1967, Tipula (Eumicrotipula)	4.21
amblytylos Alexander, 1964, Limonia (Geratomyia)	4.60
ambrosiana Alexander, 1962, Limonia (Idioglochina)	4.68
americana Alexander, 1914, Styringomyia	4.157
americana (Alexander), 1913 (Ceratocheilus), Toxorhina (Ce- ratocheilus)	4.167
amoena (Alexander), 1922 (Paratropesa), Teucholabis (Eupara- tropesa)	4.158
amoenalis Alexander, 1945, Limonia (Geratomyia)	4.60
amoenicornis Alexander, 1922, Tipula (Microtipula)	4.33
amphacanthus Alexander, 1952, Molophilus (Molophilus)	4.144
amphileuca Alexander, 1941, Epiphragma (Epiphragma)	4.95
Amphineurus Skuse, 1890	4.117
<i>Amphinome</i> Melgen, 1800	4.47
amphion Alexander, 1952, Tipula (Eumicrotipula)	4.21
amphionis Alexander, 1952, Limonia (Diceratomyia)	4.50
analis Alexander, 1954, Teucholabis (Teucholabis)	4.159
anax Alexander, 1942, Limonia (Diceratomyia)	4.50
andalgala Alexander, 1919, Tipula (Eumicrotipula)	4.21
<i>andensis</i> (Alexander), 1916 (Trimicra), Erioptera (Trimicra)	4.126
andicola Alexander, 1913, Gonomyia (Gonomyia)	4.132
andicola (Alexander), 1923 (Eriocera), Hexatoma (Eriocera)	4.100
andicola (Alexander), 1912 (Furcomyia), Limonia (Diceratomyia) .	4.50
andicola Alexander, 1942, Neolimnophila	4.154
andina Alexander, 1913, Erioptera (Erioptera)	4.122
andina Alexander, 1916, Orimarga (Orimarga)	4.82
andina Bréthes, 1909, Tipula (Eumicrotipula)	4.21
andinalta Alexander, 1956, Limonia (Diceratomyia)	4.50



andromache Alexander, 1950, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.21
anduzeana Alexander, 1941, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
anduzeana Alexander, 1943, <i>Limonia</i> (<i>Geranomyia</i>)	4.60
anduzei Alexander, 1940, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
anerasta Alexander, 1934, <i>Molophilus</i> (<i>Molophilus</i>)	4.144
angelica Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
angolensis Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.21
angusta Alexander, 1945, <i>Brachypremna</i>	4.6
angustalbus Alexander, 1953, <i>Helius</i>	4.45
angustapicalis Alexander, 1938, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.159
angusticincta (Alexander), 1921 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.60
angustifascia Alexander, 1968, <i>Teucholabis</i> (<i>Euparatropesa</i>)	4.158
angustifasciata (Alexander), 1922 (<i>Peripheroptera</i>), <i>Limonia</i> (<i>Peripheroptera</i>)	4.72
angustifrons (Alexander), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ...	4.75
angustilineata Alexander, 1926, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
angustior Alexander, 1936, <i>Molophilus</i> (<i>Eumolophilus</i>)	4.143
angustipennis Alexander, 1928, <i>Atarba</i> (<i>Atarba</i>)	4.85
angustirostris Alexander, 1947, <i>Sigmatomera</i> (<i>Sigmatomera</i>)	4.156
angustissima Alexander, 1931, <i>Elephantomyia</i> (<i>Elephantomyia</i>) ..	4.94
angustissima Alexander, 1928, <i>Gonomyia</i> (<i>Idiocera</i>)	4.134
animula Alexander, 1966, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
anisacantha Alexander, 1964, <i>Limonia</i> (<i>Geranomyia</i>)	4.60
<i>annulata</i> (Enderlein), 1912 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>) ..	4.62
<i>annulata</i> Williston, 1896, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.159
annulicornis Alexander, 1921, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.95
annulicornis (Enderlein), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ..	4.74
annulipes (Philippi), 1865 (<i>Tipula</i>), <i>Leptotarsus</i> (<i>Leptotarsus</i>) ...	4.11
annulipes Williston, 1896, <i>Erioptera</i> (<i>Erioptera</i>)	4.122
anserina Alexander, 1943, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
<i>antarctica</i> (Walker), 1836 (<i>Gonomyia</i>), <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>antarctica</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.21
antennata Alexander, 1953, <i>Aphrophila</i>	4.118
antennata (Alexander), 1930 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ...	4.100
antenniferus (Alexander), 1942 (<i>Tanyremna</i>), <i>Leptotarsus</i> (<i>Tanyremnella</i>)	4.15
anthina Alexander, 1945, <i>Limonia</i> (<i>Geranomyia</i>)	4.60
anthonympha Alexander, 1928, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.21
anthracina Alexander, 1937, <i>Atarba</i> (<i>Atarba</i>)	4.85
anthracina Alexander, 1921, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.159
anthracopoda Alexander, 1938, <i>Limonia</i> (<i>Neolimnobia</i>)	4.69
anticalba Alexander, 1939, <i>Polymera</i> (<i>Polymera</i>)	4.111
antimenus Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.144
antinympa (Alexander), 1942 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Icriomastax</i>)	4.8
Antocha Osten Sacken, 1859	4.44
antrotichia Alexander, 1967, <i>Limonia</i> (<i>Rhipidia</i>)	4.74
apacheana (Alexander), 1946 (<i>Erioptera</i>), <i>Empedomorpha</i>	4.122
aperta Alexander, 1926, <i>Atarba</i> (<i>Atarba</i>)	4.85
aphanta Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
aphrodite Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.74



Aphrophila Edwards, 1923	4.118
apicalis Macquart, 1838, Ozodicera (Dihexaclonus)	4.18
apicalis Alexander, 1914, Sigmatomera (Sigmatomera)	4.156
apicata (Alexander), 1914 (Dicranomyia), Limonia (Caenoglochina)	4.47
apicalba Alexander, 1921, Erioptera (Erioptera)	4.123
apicidens Alexander, 1951, Molophilus (Molophilus)	4.144
apicinigra Alexander, 1927, Erioptera (Mesocyphona)	4.124
apicispinosa Alexander, 1934, Atarba (Atarba)	4.85
apicularis Alexander, 1945, Gnophomyia	4.128
apiculatus Alexander, 1968, Amphineurus (Rhamphoneurus)	4.118
aplecta Alexander, 1946, Tipula (Trichotipula)	4.40
<i>apocilla</i> (Philippi), 1865 (Limnophila), Erioptera (Trimicra)	4.126
apollyon Alexander, 1951, Tipula (Microtipula)	4.33
<i>Aporosa</i> Macquart, 1839	4.60
appendens (Enderlein), 1912 (Macromastix), Tipula (Microtipula) .	4.33
appendiculata Alexander, 1943, Gonomyia (Gonomyia)	4.132
appendigera Alexander, 1945, Brachyremna	4.6
apposita Alexander, 1942, Limonia (Dicranomyia)	4.50
apressospinus Alexander, 1952, Molophilus (Molophilus)	4.144
apressus Alexander, 1929, Molophilus (Molophilus)	4.144
aprica Alexander, 1962, Atarba (Ischnothrix)	4.88
apterogynce Philippi, 1865, Tipula (Eumicrotipula)	4.22
araguae Alexander, 1947, Shannonomyia (Shannonomyia)	4.114
atacama Alexander, 1912, Tipula (Eumicrotipula)	4.22
arajuno Alexander, 1945, Brachyremna	4.6
arajuno Alexander, 1945, Gonomyia (Lipophleps)	4.135
araucana (Alexander), 1929 (Holorusia), Ischnotoma (Ischnotoma)	4.9
araucania Alexander, 1928, Limnophila (Limnophila)	4.107
araucania Alexander, 1928, Tipula (Eumicrotipula)	4.22
araucanus Alexander, 1929, Molophilus (Molophilus)	4.144
araucariae Alexander, 1945, Heliis	4.45
Araucolimnophila Alexander, 1939	4.106
Araucomyia Alexander, 1929	4.11
arawak Alexander, 1964, Orimarga (Diotrepha)	4.81
arawak Alexander, 1964, Polymera (Polymera)	4.111
archangelica Alexander, 1942, Limonia (Neolimnobia)	4.69
<i>Archimolophilus</i> Enderlein, 1940	4.143
arcuaria Alexander, 1937, Brachyremna	4.6
arcuaria Alexander, 1945, Elephantomyia (Elephantomyia)	4.94
arciferus Alexander, 1952, Molophilus (Molophilus)	4.144
areuata Alexander, 1914, Gnophomyia	4.128
areuata (Alexander), 1913 (Peripheroptera), Limonia (Periphe- roptera)	4.72
arecuna Alexander, 1931, Limonia (Geranomyia)	4.60
arecuna Alexander, 1931, Tipula (Eumicrotipula)	4.22
arepuchoensis Alexander, 1962, Tipula (Eumicrotipula)	4.25
argentea Alexander, 1927, Teucholabis (Teucholabis)	4.159
argenteoplura Alexander, 1913, Orimarga (Orimarga)	4.82
argenteiceps Alexander, 1929, Shannonomyia (Shannonomyia)	4.114
argentina Alexander, 1944, Erioptera (Symplecta)	4.126
argentina (Alexander), 1919 (Eriocera), Hexatoma (Eriocera) ...	4.100
argentina (Alexander), 1912 (Furcomomyia), Limonia (Dicranomyia)	4.50
argentina (Wulp), 1881 (Ozodicera), Tipula (Pectinotipula)	4.39



argentinensis Alexander, 1923, Cryptolabis (Procryptolabis)	4.122
argentinensis Alexander, 1920, Gonomyia (Progonomyia)	4.141
argentinensis (Alexander), 1920 (Geranomyia), Limonia (Geranomyia)	4.60
argentinicola (Alexander), 1921 (Orolimnophila), Atarba (Isehnotherix)	4.88
argutula Alexander, 1944, Gnophomyia	4.128
<i>Arhipidia</i> Alexander, 1912	4.74
ariadne Alexander, 1942, Limonia (Dieranomyia)	4.50
airanhae Alexander, 1945, Tipula (Microtipula)	4.33
armatipennis Alexander, 1912, Tipula (Microtipula)	4.33
armatistylus Alexander, 1929, Molophilus (Molophilus)	4.144
armigera Alexander, 1945, Limnophila (Limnophila)	4.108
armillaris (Fabricius), 1805 (Tipula), Lecteria (Lecteria)	4.106
armillata Alexander, 1916, Tipula (Eumierotipula)	4.22
<i>Arrhenica</i> Osten Saeken, 1859	4.99
asarotica Alexander, 1962, Tipula (Eumierotipula)	4.22
aseendens Alexander, 1945, Molophilus (Molophilus)	4.144
aspera Alexander, 1943, Gonomyia (Gonomyia)	4.132
aspilota Alexander, 1967, Limonia (Rhipidia)	4.74
associans (Walker), 1861 (Tipula), Zelandotipula	4.41
assueta Alexander, 1943, Limonia (Geranomyia)	4.60
<i>Astelobia</i> Edwards, 1923	4.156
asteria Alexander, 1951, Tipula (Eumierotipula)	4.22
<i>Astrolabis</i> Osten Saeken, 1865	4.84
astuta Alexander, 1965, Gnophomyia	4.128
atacama Alexander, 1912, Tipula (Eumierotipula)	4.22
atahualpa Alexander, 1963, Teucholabis (Teucholabis)	4.159
atacama Alexander, 1912, Tipula (Eumierotipula)	4.22
Atarba Osten Saeken, 1869	4.85
atmophora Alexander, 1929, Cryptolabis (Cryptolabis)	4.120
atribasis (Alexander), 1914 (Diotrepha), Orimarga (Diotrepha) ..	4.81
atripennis Alexander, 1940, Teucholabis (Teucholabis)	4.159
atripes Alexander, 1922, Toxorhina (Toxorhina)	4.167
atrirostris (Alexander), 1945 (Macromastix), Leptotarsus (Longurio)	4.11
atritarsis Alexander, 1949, Limonia (Dieranomyia)	4.50
atritarsis Alexander, 1943, Toxorhina (Ceratocheilus)	4.167
atroapicalis Alexander, 1934, Shannonomyia (Shannonomyia)	4.114
atroapicata Alexander, 1938, Gonomyia (Progonomyia)	4.141
atrolata Alexander, 1948, Teucholabis (Teucholabis)	4.159
atroscapa Alexander, 1949, Tipula (Eumierotipula)	4.22
atrosignata Alexander, 1941, Hexatoma (Eriocera)	4.100
atrosignata Alexander, 1934, Limonia (Peripheroptera)	4.72
atrostyla Alexander, 1942, Limonia (Dieranomyia)	4.50
atroterminata Alexander, 1943, Epiphragma (Epiphragma)	4.95
atrovelutina (Alexander), 1928 (Microtipula), Tipula (Eumierotipula)	4.22
attenuata Alexander, 1920, Ozodicera (Ozodicera)	4.19
Atypophthalmus Brunetti, 1911	4.47
audax Alexander, 1913, Teucholabis (Teucholabis)	4.159
aurantiaca Alexander, 1944, Aphrophila	4.118
aurantionota Alexander, 1939, Hexatoma (Eriocera)	4.100
aurantiothorax (Alexander), 1953 (Longurio), Leptotarsus (Tany-	




premna)	4.14
aurantiothorax Alexander, 1941, Limonia (Dieranomyia)	4.50
aurieomata Alexander, 1942, Tipula (Eumierotipula)	4.22
aurieosta Alexander, 1939, Epiphragma (Epiphragma)	4.95
auricularis Alexander, 1942, Tipula (Mierotipula)	4.33
aurigena Alexander, 1944, Limonia (Neogloehina)	4.68
austera Alexander, 1950, Limonia (Neolimonia)	4.70
australis Alexander, 1923, Braehypremna	4.6
austroandina Alexander, 1929, Limonia (Geranomyia)	4.60
austroandina Alexander, 1929, Limonia (Peripheroptera)	4.72
austroandina Alexander, 1929, Tipula (Eumierotipula)	4.22
austrolathraea Alexander, 1930, Shannonomyia (Shannonomyia) ..	4.114
Austrolimnobia Alexander, 1922	4.156
Austrolimnophila Alexander, 1920	4.89
austronymphica (Alexander), 1943 (Erioptera), Cheilotrichia (Empeda)	4.119
austrovolens Alexander, 1951, Tipula (Mierotipula)	4.33
avara Alexander, 1944, Limonia (Geranomyia)	4.60
avitus Alexander, 1952, Molophilus (Molophilus)	4.144
axierasta Alexander, 1952, Limonia (Dieranomyia)	4.50
axillaris Alexander, 1921, Gnophomyia	4.128
axillispinus Alexander, 1962, Molophilus (Molophilus)	4.144
aymara Alexander, 1962, Gonomyia (Gonomyia)	4.132
aymara Alexander, 1912, Tipula (Mierotipula)	4.33
<i>Aymaramyia</i> Alexander, 1943	4.118
aysenensis Alexander, 1944, Pedieia (Trieypfona)	4.84
azrael (Alexander), 1943 (Gnophomyia), Eugnophomyia	4.127
aztea Alexander, 1925, Tipula (Eumierotipula)	4.22
azuayensis Alexander, 1945, Teueholabis (Teueholabis)	4.159
baeckstroemi Alexander, 1920, Tipula (Eumierotipula)	4.22
Baeoura Alexander, 1924	4.119
bahiana Alexander, 1930, Orimarga (Orimarga)	4.82
bahiensis Alexander, 1930, Limonia (Geranomyia)	4.60
balloui Alexander, 1938, Tipula (Eumierotipula)	4.22
balzapambae Alexander, 1941, Gonomyia (Progonomyia)	4.141
banksi Alexander, 1927, Elephantomyia (Elephantomyia)	4.94
banksiana Alexander, 1945, Gnophomyia	4.128
banksiana Alexander, 1939, Limonia (Geranomyia)	4.60
banosensis Alexander, 1945, Limonia (Rhipidia)	4.74
bariloehensis Alexander, 1929, Cryptolabis (Procryptolabis)	4.122
bariloehensis Alexander, 1929, Shannonomyia (Shannonomyia) ...	4.114
barretoi Alexander, 1923, Tipula (Eumierotipula)	4.22
basalis (Alexander), 1921 (Rhabdomastix), Cheilotrichia (Empeda)	4.119
basillea Alexander, 1921, Braehypremna	4.6
basispinosa Alexander, 1936, Gonomyia (Lipophleps)	4.135
basistylata Alexander, 1928, Limonia (Caenogloehina)	4.48
batesi Alexander, 1945, Gonomyia (Lipophleps)	4.135
batesi (Alexander), 1921 (Pentoptera), Hexatoma (Eriocera) ...	4.100
batesi Alexander, 1939, Shannonomyia (Shannonomyia)	4.114
bathomees Alexander, 1962, Tipula (Eumierotipula)	4.22
beatrice Alexander, 1945, Limonia (Geranomyia)	4.60
beebeana Alexander, 1950, Erioptera (Erioptera)	4.123
beebeana Alexander, 1950, Hexatoma (Eriocera)	4.100



beebei Alexander, 1950, Sigmatomera (Sigmatomera)	4.156
bellardiana Alexander, 1926, Tipula (Pterelachisus)	4.40
Bellardina Edwards, 1931	4.20
bellicosus Alexander, 1929, Molophilus (Molophilus)	4.144
bellingieri Alexander, 1964, Limonia (Rhipidia)	4.74
bellona Alexander, 1946, Molophilus (Molophilus)	4.144
bequaertiana Alexander, 1938, Hexatoma (Eriocera)	4.100
berthae Alexander, 1948, Atarba (Ischnothrix)	4.88
bezzianus (Alexander), 1921 (Tipula), Leptotarsus (Tanypremna) .	4.14
biacifera Alexander, 1949, Teucholabis (Teucholabis)	4.160
biaculeata Alexander, 1953, Ozodicera (Dihexaclonus)	4.18
biargentata Alexander, 1929, Limonia (Geranomyia)	4.60
biarmata Alexander, 1931, Gynoplistia (Gynoplistia)	4.98
bicaudatus Alexander, 1929, Molophilus (Molophilus)	4.144
bicincta (Alexander), 1921 (Geranomyia), Limonia (Geranomyia)	4.60
bicinctipes Alexander, 1913, Erioptera (Mesocyphona)	4.124
bicolor (Philippi), 1865 (Ctedonia), Gynoplistia (Gynoplistia)	4.98
bicomifera Alexander, 1943, Limonia (Dicranomyia)	4.50
bicornuta Alexander, 1927, Gonomyia (Lipophleps)	4.135
bidentata Alexander, 1968, Aphrophila	4.118
bidentifera Alexander, 1966, Teucholabis (Teucholabis)	4.160
bidigitatus Alexander, 1931, Molophilus (Molophilus)	4.144
bierigi Alexander, 1947, Molophilus (Molophilus)	4.144
bievexa Alexander, 1967, Erioptera (Mesocyphona)	4.124
bifasciata (Fabricius), 1805 (Tipula), Teucholabis (Teucholabis) .	4.166
bifasciolata Alexander, 1937, Gonomyia (Progonomyia)	4.141
bifida Alexander, 1926, Gonomyia (Paralipophleps)	4.140
bifidaria Alexander, 1942, Austrolimnophila (Austrolimnophila) ..	4.89
bifiligerata Alexander, 1932, Gonomyia (Lipophleps)	4.135
bifilosa Alexander, 1948, Atarba (Atarba)	4.85
bifurcata Alexander, 1947, Hexatoma (Eriocera)	4.100
bifurcifer Alexander, 1939, Gonomyia (Gonomyia)	4.132
bifurcula Alexander, 1931, Atarba (Atarba)	4.85
bifurcula Alexander, 1946, Gonomyia (Gonomyia)	4.132
bigladiata Alexander, 1942, Limonia (Dicranomyia)	4.50
bigladius Alexander, 1942, Teucholabis (Teucholabis)	4.160
bigoti (Alexander), 1913 (Furcomyia), Limonia (Zelandoglochina)	4.80
bigotiana Alexander, 1920, Tipula (Eumicrotipula)	4.22
bilimeki Alexander, 1941, Tipula (Microtipula)	4.33
bimaculata Enderlein, 1912, Ozodicera (Ozodicera)	4.19
bimucronata Alexander, 1938, Limonia (Neolimonia)	4.70
binarius Alexander, 1968, Molophilus (Molophilus)	4.144
binigrocincta Alexander, 1962, Epiphragma (Epiphragma)	4.95
bipectinata Alexander, 1965, Gnophomyia	4.128
bipectinata (Williston), 1896 (Rhipidia), Limonia (Rhipidia)	4.74
bipendula Alexander, 1966, Atarba (Atarba)	4.85
biproducta Alexander, 1966, Atarba (Atarba)	4.85
biprolata Alexander, 1967, Tipula (Microtipula)	4.33
bipunctata (Philippi), 1865 (Ctedonia), Gynoplistia (Gynoplistia) .	4.98
birama Alexander, 1941, Gonomyia (Gonomyia)	4.132
biramosa Alexander, 1940, Teucholabis (Teucholabis)	4.160
bisatra (Alexander), 1962 (Holorusia), Zelandotipula	4.41
bisecta (Alexander), 1920 (Gnophomyia), Neognophomyia	4.153
bisetosa Alexander, 1944, Neognophomyia	4.153

bisetosa Alexander, 1945, Teucholabis (Teucholabis)	4.162
bispinifer Alexander, 1921, Ozodicera (Ozodicera)	4.19
bispinosa Alexander, 1921, Gonomyia (Lipophleps)	4.135
bitergatus Alexander, 1950, Helius	4.45
bitribula Alexander, 1967, Tipula (Microtipula)	4.33
bituberculata (Macquart), 1838 (Limnobia), Hexatoma (Eriocera)	4.100
bivera Alexander, 1921, Ctenolimnophila (Ctenolimnophila)	4.93
<i>blanchardi</i> (Alexander), 1913 (Furcomomyia), Limonia (Dicranomyia)	4.56
blaseri Alexander, 1942, Tipula (Microtipula)	4.33
bogotana Alexander, 1938, Tipula (Eumicrotipula)	4.22
bogotensis Alexander, 1938, Shannonomyia (Shannonomyia)	4.114
boliviana Alexander, 1930, Atarba (Atarba)	4.85
boliviana (Alexander), 1930 (Erioptera), Cheilotrichia (Empeda) .	4.119
boliviana Alexander, 1930, Limonia (Dicranomyia)	4.50
boliviana Alexander, 1947, Nephrotoma	4.16
boliwijsis Alexander, 1930, Elephantomyia (Elephantomyia)	4.94
boliwijsis(Alexander), 1946 (Pectinotipula), Tipula (Pectinotipula)	4.39
boraceae Alexander, 1954, Tasiocera (Dasymolophilus)	4.157
borburatana Alexander, 1941, Gonomyia (Lipophleps)	4.135
borgmeieri (Alexander), 1942 (Tanypremna), Leptotarsus (Tany- premna)	4.14
borinquena Alexander, 1968, Limonia (Dicranomyia)	4.50
borinquensis Alexander, 1950, Limonia (Neolimonia)	4.70
brachycantha Alexander, 1945, Limonia (Caenolimonia)	4.48
brachynema Alexander, 1936, Limonia (Rhipidia)	4.75
brachyneura Alexander, 1962, Polymera (Polymera)	4.111
Brachypremna Osten Sacken, 1886	4.5
braconides (Enderlein), 1912 (Eriocera), Hexatoma (Eriocera) ..	4.100
bradleyi Alexander, 1929, Austrolimnophila (Austrolimnophila) ..	4.89
brasiliae (Alexander), 1935 (Macromastix), Leptotarsus (Longurio)	4.11
brasiliensis Alexander, 1913, Erioptera (Mesocyphona)	4.125
brasiliensis (Westwood), 1835 (Limnobiorthynchus), Limonia (Ge- ranomyia)	4.61
brasiliensis (Westwood), 1835 (Limnobiorthynchus), Toxorhina (Toxorhina)	4.167
<i>brasiliensis</i> (Westwood), 1835 (Limnobiorthynchus), Toxorhina ..	4.168
brasiliensis (Wiedemann), 1828 (Megistocera), Tipula (Microtipula)	4.33
brethesiana Alexander, 1929, Tipula (Eumicrotipula)	4.22
brevicoma Alexander, 1944, Tipula (Eumicrotipula)	4.22
brevicornis Alexander, 1929, Atarba (Atarba)	4.85
brevicornis Alexander, 1929, Tasiocera (Dasymolophilus)	4.157
brevicubitalis Alexander, 1947, Limonia (Dicranomyia)	4.50
brevicula Alexander, 1926, Gonomyia (Gonomyia)	4.132
brevicula Alexander, 1931, Shannonomyia (Shannonomyia)	4.114
brevigenua Alexander, 1945, Brachypremna	4.6
brevigliadia Alexander, 1962, Limonia (Dicranomyia)	4.50
brevihirsutus (Alexander), 1934 (Macromastix), Leptotarsus (Arau- comyia)	4.11
brevilobatus Alexander, 1938, Molophilus (Molophilus)	4.144
brevilyra Alexander, 1966, Atarba (Ischnothrix)	4.88
brevinervis Alexander, 1929, Shannonomyia (Shannonomyia)	4.114
brevipetalia Alexander, 1950, Limonia(Rhipidia)	4.74
breviramosa Alexander, 1936, Limonia (Rhipidia)	4.74
breviramus Alexander, 1929, Molophilus (Molophilus)	4.144



<i>brevisector</i> (Alexander), 1935 (<i>Isehnothrix</i>), <i>Atarba</i> (<i>Ischnothrix</i>)	4.88
<i>brevisector</i> Alexander, 1936, <i>Leeteria</i> (<i>Psaronius</i>)	4.106
<i>brevisetosa</i> Alexander, 1968 (<i>Neophilippiana</i>), <i>Erioptera</i> (<i>Eriopterella</i>)	4.123
<i>brevisetosa</i> Alexander, 1941, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>brevispica</i> Alexander, 1962, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>brevispinosus</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.144
<i>brevispinula</i> Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>brevissima</i> Alexander, 1944, <i>Atarba</i> (<i>Atarba</i>)	4.85
<i>brevissima</i> Alexander, 1926, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
<i>breviterebra</i> Alexander, 1944, <i>Brachypremna</i>	4.6
<i>brevitibia</i> (Alexander), 1920 (<i>Psaronius</i>), <i>Leeteria</i> (<i>Psaronius</i>)	4.106
<i>breviuseula</i> (Alexander), 1929 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
<i>brevivena</i> (Osten Sacken), 1869 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Dieranomyia</i>)	4.50
<i>breviventris</i> (Wiedemann), 1821 (<i>Tipula</i>), <i>Brachypremna</i>	4.6
<i>bromeliadieola</i> (Alexander), 1912 (<i>Mongoma</i>), <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.169
<i>browni</i> (Alexander), 1940 (<i>Maeromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.11
<i>browniana</i> Alexander, 1940, <i>Tipula</i> (<i>Eumierotipula</i>)	4.23
<i>brownianus</i> Alexander, 1945, <i>Molophilus</i> (<i>Molophilus</i>)	4.144
<i>bruehi</i> Alexander, 1920, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
<i>bruchi</i> Alexander, 1923, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>bruchi</i> Alexander, 1926, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>bruchi</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
<i>bruehiana</i> Alexander, 1929, <i>Limonia</i> (<i>Rhipidia</i>)	4.74
<i>bruesi</i> Alexander, 1945, <i>Tipula</i> (<i>Mierotipula</i>)	4.33
<i>bruneri</i> (Alexander), 1928 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
<i>bruneri</i> Alexander, 1926, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>bruneriana</i> Alexander, 1937, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>brunneicornis</i> Alexander, 1916, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>brunneipennis</i> Alexander, 1938, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.94
<i>brunneipes</i> (Williston), 1900 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
<i>bulbibasis</i> Alexander, 1944, <i>Gnophomyia</i>	4.128
<i>bulbifera</i> Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>bulbifera</i> Alexander, 1948, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.89
<i>bullocki</i> (Alexander), 1931 (<i>Maeromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.11
<i>bullocki</i> (Alexander), 1935 (<i>Astelobia</i>), <i>Sigmatomera</i> (<i>Austrolimnobia</i>)	4.156
<i>bullockiana</i> Alexander, 1938, <i>Limonia</i> (<i>Dieranomyia</i>)	4.51
<i>bullockiana</i> (Alexander), 1939 (<i>Limnophila</i>), <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>buscki</i> Alexander, 1913, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>buscki</i> Alexander, 1915, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>bustilloi</i> Alexander, 1938, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>byersiana</i> Alexander, 1968, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.89
<i>cabralensis</i> Alexander, 1945, <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
<i>eacoxcna</i> Alexander, 1929, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
<i>caeminis</i> Alexander, 1945, <i>Nephrotoma</i>	4.17
<i>Caenoglochina</i> Alexander, 1964	4.47
<i>Caenolimonia</i> Alexander, 1967	4.48
<i>caenusus</i> Alexander, 1936, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>caesia</i> Alexander, 1937, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114



calceatus Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
caliptera Say, 1823, Erioptera (<i>Mesocyphona</i>)	4.124
calliargon Alexander, 1939, <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
callinota Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
calliope (Alexander), 1945 (<i>Holorusia</i>) <i>Ischnotoma</i> (<i>Icriomastax</i>) .	4.8
calliope (Alexander), 1945 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i>	
premna)	4.14
calliope Alexander, 1944, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.169
callisto Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
<i>Caloptera</i> Guérin-Ménéville, 1831	4.99
caloptera Osten Sacken, 1887, <i>Gnophomyia</i>	4.128
calopus Walker, 1856, <i>Lecteria</i> (<i>Lecteria</i>)	4.106
calverti Alexander, 1914, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
calverti (Alexander), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>)	4.74
calvicornis (Edwards), 1919 (<i>Prionocera</i>), <i>Zelandotipula</i>	4.41
calypso (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.41
caminaria (Wiedemann), 1828 (<i>Limnobia</i>), <i>Hexatoma</i> (<i>Eriocera</i>) .	4.100
campa Alexander, 1914, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
Campbellomyia Alexander, 1925	4.93
cana (Walker), 1848 (<i>Limnobia</i>), Erioptera (<i>Symplecta</i>)	4.126
cancellata Alexander, 1962, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
candida Alexander, 1912, <i>Brachypremna</i>	4.6
candidipes (Alexander), 1923 (<i>Pentoptera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
candiditarsis Alexander, 1937, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
canopus Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
cantareirae Alexander, 1945, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
caparaocensis Alexander, 1944, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
caparaocensis Alexander, 1944, <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
caparaonus (Alexnader), 1944 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i>	
gurio)	4.12
capitella Alexander, 1945, <i>Atarba</i> (<i>Ischnothrix</i>)	4.88
capitonus Alexander, 1945, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
capniopterus Alexander, 1945, <i>Helius</i>	4.45
capnora (Alexander), 1921 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Neoglochina</i>)	4.68
capra Alexander, 1947, <i>Limonia</i> (<i>Dicranomyia</i>)	4.50
capricornis Alexander, 1916, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
captiosa Alexander, 1945, <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
capucina Alexander, 1947, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
carbonaria Alexander, 1931, <i>Aphrophila</i>	4.118
carbonipes (Alexander), 1938 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i>	
premna)	4.14
caribaea Alexander, 1932, <i>Limonia</i> (<i>Neolimonia</i>)	4.70
carizona Alexander, 1913, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
carpapatae Alexander, 1941, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
carpishensis Alexander, 1949, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
carrerai Alexander, 1943, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
carrerai Alexander, 1942, <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
carreranus (Alexander), 1945 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i>	
gurio)	4.12
carrerella Alexander, 1956, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
carunculata Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
cassandra Alexander, 1945, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
castroensis Alexander, 1929, <i>Amphincurus</i> (<i>Amphineurus</i>)	4.117



catamarcae Alexander, 1929, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
catamarcana Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
catamarcensis Alexander, 1941, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
catamarcensis Alexander, 1921, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
catamarcensis (Alexander), 1920 (<i>Holorusia</i>), <i>Zelandotipula</i>)	4.41
catharinae Alexander, 1931, <i>Polymera</i> (<i>Polymerodes</i>)	4.113
catharinensis Alexander, 1931, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
caudifera Alexander, 1945, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
cautinensis Alexander, 1941, <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
cautus Alexander, 1947, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
cavernicola Alexander, 1964, <i>Polymera</i> (<i>Polymera</i>)	4.111
cayutuensis Alexander, 1944, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.92
celator Alexander, 1946, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
celator Alexander, 1942, <i>Molophilus</i> (<i>Trichomolophilus</i>)	4.152
celestior Alexander, 1956, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
celestis Alexander, 1940, <i>Erioptera</i> (<i>Erioptera</i>)	4.123
celestissima (Alexander), 1945 (<i>Austrolimnophila</i>), <i>Limnophila</i>	
(<i>Habrolimnophila</i>)	4.107
celestissima (Alexander), 1953 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.41
centralis Alexander, 1913, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.167
Ceoneura Alexander, 1924	4.11
Ceratocheilus Wesché, 1910	4.167
cerbereana Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
cerbercana Alexander, 1945, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
cerberus (Alexander), 1927 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.61
cermatoleuca Alexander, 1967, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
cerogama Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.33
cerrita (Alexander), 1945 (<i>Tanyremna</i>), <i>Leptotarsus</i> (<i>Tanyremna</i>)	4.114
certa Alexander, 1952, <i>Gnophomyia</i>	4.128
certhia (Alexander), 1916 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>) .	4.61
cervaria Alexander, 1938, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
cervus Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
chacopata Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
chalybeia Alexander, 1927, <i>Teucholabis</i> (<i>Paratropesa</i>)	4.158
chalybeiventris Alexander, 1949, <i>Hexatoma</i> (<i>Eriocera</i>)	4.100
chalybeiventris (Loew), 1861 (<i>Rhamphidia</i>), <i>Teucholabis</i> (<i>Teu-</i>	
<i>cholabis</i>)	4.160
chanca Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
chaparensis Alexander, 1962, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
charmosyne Alexander, 1967, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
charon Alexander, 1937, <i>Limnophila</i> (<i>Limnophila</i>)	4.109
Cheilotrichia Rossi, 1848	4.119
chiapasensis Alexander, 1927, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.132
chiapasensis Alexander, 1938, <i>Toxorhina</i> (<i>Ceratocheilus</i>)	4.167
chicana Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
Chilelimnophila Alexander, 1968	4.92
chilena Alexander, 1928, <i>Aphrophila</i>	4.118
chilensis Blanchard, 1852 (<i>Limnophila</i>)	4.117
chilensis (Alexander), 1913 (<i>Furcomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
chilensis Alexander, 1929, <i>Stibadocera</i>	4.44
chilensis Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23
chillanica Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.23


chiloeana Alexander, 1953, Austrolimnophila (Austrolimnophila) .	4.90
chilocana Alexander, 1967, Limonia (Rhipidia)	4.75
chiloensis Alexander, 1929, Edwardsomyia	4.93
chilota Alexander, 1929, Cryptolabis (Cryptolabis)	4.120
chilota Alexander, 1929, Gnophomyia	4.128
chilota (Alexander), 1929 (Tricyphona), Pedicia (Tricyphona) ..	4.84
chilota Alexander, 1929, Rhabdomastix (Rhabdomastix)	4.155
chilota Alexander, 1929, Tipula (Eumicrotipula)	4.23
chimbrazicola Alexander, 1945, Limonia (Dicranomyia)	4.51
chimbrazo (Alexander), 1945 (Holorusia), Zelandotipula	4.41
chionomera Alexander, 1945, Orimarga (Orimarga)	4.82
chionopoda Alexander, 1943, Elephantomyia (Elephantomyia) ...	4.94
chionopus Alexander, 1943, Orimarga (Orimarga)	4.82
chiriquiana Alexander, 1945, Elephantomyia (Elephantomyia)	4.94
chiriquiana Alexander, 1934, Trentepohlia (Paramongoma)	4.169
chiriquiensis Alexander, 1941, Elephantomyia (Elephantomyia) ..	4.94
chiriquiensis Alexander, 1934, Molophilus (Molophilus)	4.145
chiriquiensis Alexander, 1941, Polymera (Polymera)	4.111
chlorotica (Philippi), 1865 (Limnobia), Limonia (Dicranomyia) ..	4.51
chrysoptera (Walker), 1856 (Limnobia), Hexatoma (Eriocera) ...	4.101
chrysopteroides (Alexander), 1922 (Eriocera), Hexatoma (Eriocera)	4.101
chryso stigma (Alexander), 1944 (Macromastix), Leptotarsus	
(Longurio)	4.12
cineticornis Alexander, 1916, Atarba (Atarba)	4.86
cinetiventris Alexander, 1967, Elephantomyia (Elephantomyia) ..	4.94
cineracea Philippi, 1865, (Limnophila)	4.117
cinerea (Philippi), 1865 (Polymoria)	4.117
cinereiceps Alexander, 1928, Teucholabis (Teucholabis)	4.160
cinereinota (Alexander), 1913 (Geranomyia), Limonia (Geranomyia)	4.61
einereipennis Alexander, 1937, Ozodicera (Ozodicera)	4.19
einereipennis Alexander, 1926, Polymera (Polymera)	4.111
eingara Alexander, 1948, Shannonomyia (Shannonomyia)	4.114
elrce Alexander, 1945, Atarba (Atarba)	4.86
eirelnata Osten Sacken, 1886, Epiphragma (Epiphragma)	4.96
eircumscripta Alexander, 1945, Teucholabis (Teucholabis)	4.160
circumscripta (Loew), 1863 (Paehyrrhina), Nephrotoma	4.17
cisatlantica Alexander, 1937, Hexatoma (Cladolipes)	4.99
cisatlanticus (Alexander), 1937 (Xenotipula), Leptotarsus (Xeno-	
tipula)	4.16
cisatlanticus Alexander, 1938, Protohelius	4.83
cithariformis Alexander, 1967, Tipula (Microtipula)	4.33
citripes Alexander, 1945, Neognophomyia	4.153
eivica Alexander, 1939, Limonia (Geranomyia)	4.61
cladoerus Alexander, 1921, Molophilus (Molophilus)	4.145
Cladolipes Loew, 1865	4.99
cladophora Alexander, 1920, Erioptera (Erioptera)	4.123
cladophoroides Alexander, 1921, Erioptera (Erioptera)	4.123
claribasis Alexander, 1945, Limonia (Dicranomyia)	4.51
clarkiana Alexander, 1929, Tipula (Eumicrotipula)	4.23
claudia Alexander, 1945, Epiphragma (Epiphragma)	4.96
clausa Alexander, 1939, Polymera (Polymera)	4.111
clavaria Alexander, 1945, Tipula (Eumicrotipula)	4.23
clavifera Alexander, 1943, Gonomyia (Lipophleps)	4.136
clavigera Alexander, 1931, Limonia (Dicranomyia)	4.51
clavigerus Alexander, 1929, Molophilus (Molophilus)	4.145



clavistyla Alexander, 1956, Limonia (Dicranomyia)	4.51
clementis Alexander, 1945, Limonia (Geranomyia)	4.66
clitillaria Alexander, 1929, Elephantomyia (Elephantomyia)	4.94
clotho (Alexander), 1944 (Tanypremna), Leptotarsus (Tanypremna)	4.14
clymenc Alexander, 1945, Brachypremna	4.6
coangustata (Alexander), 1944 (Erioptera), Cheilotrichia (Empeda)	4.119
cochabambae Alexander, 1945, Limonia (Peripheroptera)	4.72
cochlearis Alexander, 1945, Neognophomyia	4.153
cockerclae Alexander, 1915, Teucholabis (Teucholabis)	4.160
coheri Alexander, 1964, Limonia (Dicranomyia)	4.51
collaris (Osten Sacken), 1887 (Paratropesa), Teucholabis (Para-	
tropesa)	4.158
colombiana (Alexander), 1937 (Helobia), Erioptera (Symplecta) ..	4.126
colombicola Alexander 1931, Neognophomyia	4.153
colombicola Alexander, 1929, Tipula (Microtipula)	4.33
colomelania Alexander, 1968, Teucholabis (Teucholabis)	4.160
colophallus Alexander, 1966, Limnophila (Limnophila)	4.108
colossus Alexander, 1929, Molophilus (Molophilus)	4.145
columbiana Alexander, 1913, Atarba (Atarba)	4.86
columbiana (Alexander), 1919 (Eriocera), Hexatoma (Eriocera) .	4.101
columbianus (Enderlein), 1912 (Pehlkea), Leptotarsus (Pehlkea) .	4.13
comantis Alexander, 1948, Austrolimnophila (Austrolimnophila) ..	4.90
combostena Alexander, 1967, Limonia (Caenolimonia)	4.48
commelina Alexander, 1946, Limonia (Rhipidia)	4.75
commina Alexander, 1967, Limonia (Dicranomyia)	4.51
compacta Alexander, 1941, Gonomyia (Progonomyia)	4.141
compactus Alexander, 1950, Molophilus (Molophilus)	4.145
complacita Alexander, 1942, Limonia (Dicranomyia)	4.51
complexa Alexander, 1950, Limonia (Rhipidia)	4.75
complicata (Alexander), 1922 (Rhabdomastix), Cheilotrichia	
(Empeda)	4.119
concinna (Philippi), 1865 (Tipula), Ischnotoma (Ischnotoma)	4.9
concinna (Williston), 1896 (Diotrepha), Orimarga (Diotrepha) ...	4.81
concumbens Alexander, 1942, Trentepohlia (Paramongoma)	4.169
condensa Alexander, 1938, Gonomyia (Neolipophleps)	4.139
conica (Alexander), 1914 (Rhipidia), Limonia (Rhipidia)	4.75
conifera Alexander, 1941, Gonomyia (Paralipophleps)	4.140
conjuncta (Alexander), 1914 (Pentoptera), Hexatoma (Eriocera)	4.101
conjuncta Alexander, 1913, Polymera (Polymerodes)	4.113
conjunctoides Alexander, 1920, Polymera (Polymerodes)	4.113
connivens Alexander, 1946, Gonomyia (Gonomyia)	4.132
Conorhipidia Alexander, 1914	4.74
conquisita Alexander, 1941, Limonia (Geranomyia)	4.61
conspersa (Enderlein), 1912 (Dactylolabis), Paralimnophila	4.110
conscriptus Alexander, 1938, Molophilus (Molophilus)	4.145
consociata Alexander, 1942, Neognophomyia	4.153
consonata Alexander, 1940, Tipula (Eumicrotipula)	4.23
conspersa (Enderlein), 1912 (Dactylobasis), Paralimnophila	4.110
conspicillata Alexander, 1945, Tipula (Eumicrotipula)	4.23
consularis (Osten Sacken), 1886 (Pachyrrhina), Nephrotoma	4.17
contemplata Alexander, 1951, Tipula (Microtipula)	4.34
contorta Alexander, 1941, Limonia (Geranomyia)	4.61




<i>contradistineta</i> Alexander, 1936, Limonia (Caenolimonia)	4.49
<i>contristans</i> Alexander, 1945, Limonia (Dicranomyia)	4.51
<i>corallina</i> Alexander, 1938, Limonia (Neolimnobia)	4.69
<i>cordillerense</i> Alexander, 1913, Epiphragma (Epiphragma)	4.96
<i>cordillerensis</i> (Alexander), 1913 (Dicranomyia), Limonia (Neolimonia)	4.70
<i>cornigera</i> (Alexander), 1914 (Eriocera), Hexatoma (Eriocera) ...	4.101
<i>corolcoensis</i> Alexander, 1962, Hexatoma (Eriocera)	4.101
<i>coronaria</i> Alexander, 1940, Tipula (Eumicrotipula)	4.23
<i>coronata</i> Alexander, 1944, Aphrophila	4.118
<i>correntosana</i> Alexander, 1929, Limonia (Dicranomyia)	4.56
<i>corrientesana</i> Alexander, 1962, Ozodicera (Ozodicera)	4.19
<i>eortesi</i> Alexander, 1952, Cryptolabis (Cryptolabis)	4.120
<i>costalis</i> Alexander, 1913, Erioptera (Mesocyphona)	4.124
<i>costalis</i> (Williston), 1896 (Rhipidia), Limonia (Rhipidia)	4.79
<i>costalis</i> Say, 1823, Tipula (Yamatotipula)	4.40
<i>costaloides</i> (Alexander), 1920 (Rhipidia), Limonia (Rhipidia) ...	4.75
<i>costaricensis</i> Alexander, 1939, Dicranoptycha	4.45
<i>costaricensis</i> (Alexander), 1916 (Geranomyia), Limonia (Geranomyia)	4.61
<i>costaricensis</i> (Alexander), 1922 (Adelphomyia), Paradelphomyia (Oxyrhiza)	4.109
<i>costaricensis</i> Alexander, 1922, Tipula (Eumicrotipula)	4.23
<i>coxialis</i> Alexander, 1944, Gnophomyia	4.128
<i>cramptoni</i> (Alexander), 1928 (Eriocera), Hexatoma (Eriocera) ..	4.101
<i>cramptoni</i> (Alexander), 1912 (Rhipidia), Limonia (Rhipidia)	4.75
<i>crassicornis</i> Alexander, 1964, Shannonomyia (Shannonomyia)	4.115
<i>crassipyga</i> (Alexander), 1928 (Tricyphona), Pedicia (Tricyphona)	4.84
<i>crassistyla</i> Alexander, 1966, Neognophomyia	4.153
<i>crassistyla</i> Alexander, 1965, Tipula (Microtipula)	4.34
<i>craverii</i> Bellardi, 1859, Tipula (Bellardina)	4.20
<i>creper</i> Alexander, 1926, Helius	4.46
<i>crepera</i> Alexander, 1952, Tipula (Eumicrotipula)	4.23
<i>crepuscula</i> Alexander, 1921, Gonomyia (Lipophleps)	4.136
<i>crinita</i> Alexander, 1941, Gonomyia (Gonomyia)	4.132
<i>crystalta</i> (Alexander), 1946 (Holorusia), Zelandotipula	4.41
<i>crystalta</i> Alexander, 1945, Tipula (Eumicrotipula)	4.24
<i>cristifera</i> (Alexander), 1945 (Holorusia), Zelandotipula	4.41
<i>eristobtusa</i> (Alexander), 1946 (Holorusia), Zelandotipula	4.41
<i>croceiapicalis</i> Alexander, 1962, Limonia (Dicranomyia)	4.51
<i>croceibasis</i> Alexander, 1942, Limonia (Periphoptera)	4.72
<i>crossospila</i> Alexander, 1929, Tipula (Eumicrotipula)	4.24
<i>Cryptolabis</i> Osten Sacken, 1859	4.120
<i>crystallina</i> (Alexander), 1938 (Tanypremna), Leptotarsus (Tanypremna)	4.15
<i>crystalloptera</i> Alexander, 1921, Polymera (Polymera)	4.111
<i>Ctedonia</i> Philippi, 1865	4.98
<i>Ctenolimnophila</i> Alexander, 1921	4.93
<i>ctenonycha</i> Alexander, 1929, Limnophila (Limnophila)	4.109
<i>Ctenophora</i> Meigen, 1803	4.7
<i>ctenophora</i> Alexander, 1921, Gonomyia (Lipophleps)	4.136
<i>ctenopyga</i> Alexander, 1940, Tipula (Microtipula)	4.34
<i>cubana</i> Alexander, 1931, Gonomyia (Lipophleps)	4.136



cubana Alexander, 1929, Limonia (Geranomyia)	4.61
cubense Alexander, 1929, Epiphragma (Epiphragma)	4.96
cubensis (Alexander), 1928 (Megistomastix), Dolichozepeza (Megistomastix)	4.8
cubensis (Alexander), 1926 (Eriocera), Hexatoma (Eriocera)	4.101
cubensis Alexander, 1932, Orimarga (Orimarga)	4.82
cubitalis Alexander, 1931, Trentepohlia (Paramongoma)	4.169
cucullata Alexander, 1945, Atarba (Atarba)	4.86
cultrata Alexander, 1941, Gonomyia (Lipophleps)	4.136
cuneiformis Alexander, 1947, Teucholabis (Teucholabis)	4.160
curinao Alexander, 1941, Tipula (Eumicrotipula)	4.24
curraniana (Alexander), 1945 (Gnophomyia), Eugnophomyia	4.127
curraniana Alexander, 1944, Limonia (Neoglochina)	4.68
curta (Alexander), 1925 (Erioptera), Cheilotrichia (Empeda)	4.119
curtispina Alexander, 1962, Limonia (Dicranomyia)	4.58
curvata Alexander, 1938, Toxorhina (Toxorhina)	4.167
cuzcoensis Alexander, 1967, Limonia (Neolimonia)	4.70
cuzcoensis Alexander, 1966, Neognophomyia	4.153
cybele Alexander, 1954, Teucholabis (Teucholabis)	4.166
cyclomera Alexander, 1951, Tipula (Eumicrotipula)	4.24
cydippe Alexander, 1947, Tipula (Bellardina)	4.20
cygniformis Alexander, 1953, Ozodicera (Ozodicera)	4.19
Cylindrotominae	4.44
cynara Alexander, 1942, Limonia (Peripheroptera)	4.72
cynotis Alexander, 1945, Epiphragma (Epiphragma)	4.96
cynthia Alexander, 1945, Erioptera (Mesocyphona)	4.124
cytherea Alexander, 1942, Limonia (Rhipidia)	4.75
damicoi Alexander, 1942, Limonia (Geranomyia)	4.61
dampfi Alexander, 1927, Erioptera (Erioptera)	4.123
dampfi (Alexander), 1925 (Dicranomyia), Limonia (Dicranomyia)	4.51
dampfi Alexander, 1925, Nephrotoma	4.17
dampfi Alexander, 1926, Orimarga (Orimarga)	4.82
dampfi Alexander, 1939, Shannonomyia (Shannonomyia)	4.115
dampfi Alexander, 1927, Teucholabis (Teucholabis)	4.160
dampfiana Alexander, 1938, Gonomyia (Euptilostena)	4.132
dampfiana Alexander, 1946, Tipula (Lunatipula)	4.32
darlingtoni Alexander, 1939, Dolichozepeza (Megistomastix)	4.8
darlingtoni (Alexander), 1937 (Gnophomyia), Eugnophomyia	4.127
darlingtoniana Alexander, 1939, Tipula (Eumicrotipula)	4.24
dasycera Alexander, 1948, Atarba (Atarba)	4.86
Dasymolophilus Goetghebuere, 1920	4.157
dasyphallus Alexander, 1949, Gonomyia (Gonomyia)	4.132
debilior Alexander, 1943, Molophilus (Molophilus)	4.145
debilis Alexander, 1928, Gonomyia (Gonomyia)	4.133
debilistylus Alexander, 1937, Molophilus (Molophilus)	4.145
debillitata Alexander, 1949, Neognophomyia	4.153
decasbila (Wiedemann), 1828 (Limnobia)	4.117
decens Alexander, 1942, Tipula (Microtipula)	4.34
deceptrix Alexander, 1945, Limonia (Neolimonia)	4.70
decincta Alexander, 1962, Elephantomyia (Elephantomyia)	4.94
decisa (Alexander), 1914 (Gnophomyia), Ctenolimnophila (Ctenolimnophila)	4.93
decolorata (Alexander), 1935 (Habromastix), Tipula (Microtipula)	4.34




<i>decora</i> Alexander, 1920, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>decorata</i> (Philippi), 1865 (<i>Tipula</i>), <i>Ischnotoma</i> (<i>Ischnotoma</i>)	4.9
<i>dedeora</i> Alexander, 1936, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>defoeanus</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>delectabilis</i> Alexander, 1965, <i>Gnophomyia</i>	4.128
<i>delectata</i> Alexander, 1940, <i>Tipula</i> (<i>Eumlerotipula</i>)	4.24
<i>delectissima</i> Alexander, 1954, <i>Austrolimnophila</i> (<i>Medlophragma</i>) ..	4.92
<i>dellberata</i> Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>dellcata</i> Alexander, 1913, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>dellcatula</i> (Philippi), 1865 (<i>Lachnoeera</i>), <i>Atarba</i> (<i>Ischnothrix</i>) ..	4.88
<i>dellcatula</i> Osten Sacken, 1887, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>dellciosa</i> Alexander, 1934, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>delpontei</i> (Alexander), 1929 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Ischnotoma</i>)	4.9
<i>deludens</i> (Alexander), 1939 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>) ..	4.119
<i>denuda</i> Alexander, 1943, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>deplexa</i> Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.60
<i>desdemona</i> Alexander, 1949, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>destituta</i> (Alexander), 1942 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>) .	4.119
<i>destituta</i> (Alexander), 1945 (<i>Oxydiscus</i>), <i>Paradelphomyia</i> (<i>Oxyrhiza</i>)	4.109
<i>destricta</i> Alexander, 1939, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>destricta</i> Alexander, 1940, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>detecta</i> Alexander, 1926, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>devexa</i> Alexander, 1937, <i>Dolichocheza</i> (<i>Megistomastix</i>)	4.8
<i>devota</i> Alexander, 1929, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>diabolica</i> Alexander, 1944, <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>diaantha</i> Alexander, 1945, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>dianantha</i> Alexander, 1966, <i>Gnophomyia</i>	4.128
<i>diananthophora</i> Alexander, 1941, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>diananthos</i> Alexander, 1921, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>diadema</i> Alexander, 1929, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>diadexia</i> Alexander, 1967, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>diana</i> Alexander, 1944, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
<i>diargyria</i> Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>dlazi</i> Alexander, 1937, <i>Gnophomyia</i>	4.128
<i>dicax</i> Alexander, 1941, <i>Limonia</i> (<i>Neolimonia</i>)	4.70
<i>diceros</i> Alexander, 1944, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>dichrooptera</i> Alexander, 1950, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
<i>Dieranomyia</i> Stephens, 1829	4.49
<i>Dicranoptycha</i> Osten Sacken, 1859	4.45
<i>Dicranota</i> Zetterstedt, 1838	4.84
<i>dictena</i> Alexander, 1942, <i>Gnophomyia</i>	4.128
<i>dictyophora</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>dictyoptera</i> Alexander, 1922, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
<i>didactyla</i> Alexander, 1942, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>dido</i> Alexander, 1943, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>didolos</i> Alexander, 1962, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>diffusa</i> Alexander, 1921, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>diffusior</i> Alexander, 1968, <i>Paralimnophila</i>	4.110
<i>digitifera</i> Alexander, 1945, <i>Atarba</i> (<i>Ischnothrix</i>)	4.88
<i>digitiformis</i> Alexander, 1941, <i>Gnophomyia</i>	4.128
<i>Dihexaionus</i> Enderlein, 1912	4.18
<i>dilatistyla</i> Alexander, 1966, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>dilatatus</i> Alexander, 1951, <i>Molophilus</i> (<i>Molophilus</i>)	4.145
<i>dileuca</i> Alexander, 1947, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160




dillmani Alexander, 1952, Gynoplistia (Gynoplistia)	4.98
dilucida Alexander, 1954, Limonia (Geranomyia)	4.67
dimidiata (Alexander), 1923 (Eriocera), Hexatoma (Eriocera) ..	4.104
dimorpha Alexander, 1929, Tipula (Eumicrotipula)	4.24
dinematophora Alexander, 1943, Atarba (Atarba)	4.86
Diotrepha Osten Sacken, 1878	4.81
diplaca Alexander, 1945, Teucholabis (Teucholabis)	4.160
diplacantha Alexander, 1966, Gonomyia (Paralipophleps)	4.140
dirhabda Alexander, 1949, Molophilus (Molophilus)	4.146
dirhabdophora Alexander, 1951, Tipula (Microtipula)	4.34
Dirhipis Enderlein, 1917	4.98
dirus Alexander, 1943, Molophilus (Molophilus)	4.146
dis Alexander, 1944, Limonia (Peripheroptera)	4.72
discalis Alexander, 1929, Polymera (Polymera)	4.112
Discobola Osten Sacken, 1865	4.59
discophora Alexander, 1951, Tipula (Microtipula)	4.34
disjuncta (Alexander), 1913 (Mongoma), Trentepohlia (Neomon- goma)	4.169
disparilis Alexander, 1946, Limonia (Geranomyia)	4.62
disparilis Alexander, 1944, Trentepohlia (Paramongoma)	4.169
dispellens (Walker), 1861 (Tipula), Brachypremna	4.6
dissoluta Alexander, 1929, Limonia (Dicranomyia)	4.51
distalis Alexander, 1936, Toxorhina (Toxorhina)	4.167
distans (Osten Sacken), 1859 (Dicranomyia), Limonia (Dicranomyia)	4.52
distela Alexander, 1967, Limonia (Rhipidia)	4.75
distifurca Alexander, 1943, Teucholabis (Teucholabis)	4.160
distifurcula Alexander, 1943, Gnophomyia	4.128
distifurcus Alexander, 1952, Molophilus (Molophilus)	4.146
distinctior Alexander, 1952, Elephantomyia (Elephantomyia)	4.94
distinervis Alexander, 1940, Helius	4.46
ditzleri Alexander, 1947, Trentepohlia (Paramongoma)	4.169
diura Alexander, 1942, Limonia (Dicranomyia)	4.52
diwa (Schiner), 1868 (Limnobia), Limonia (Neolimnobia)	4.70
divaricata (Alexander), 1939 (Erioptera), Cheilotrichia (Empeda)	4.119
diversa (Osten Sacken), 1859 (Dicranomyia), Limonia (Dicranomyia)	4.52
diversigladia Alexander, 1942, Limonia (Dicranomyia)	4.52
diversipes (Alexander), 1921 (Limnophila), Austrolimnophila (Limnophilella)	4.92
diversipes Alexander, 1941, Brachypremna	4.6
diversipes Alexander, 1939, Cryptolabis (Cryptolabis)	4.120
divisa Alexander, 1929, Limonia (Dicranomyia)	4.52
dizygus Alexander, 1962, Molophilus (Molophilus)	4.146
dolorosa Alexander, 1931, Limonia (Dicranomyia)	4.52
Dolichocheza Curtis, 1825	4.8
dolorosa Alexander, 1922, Gonomyia (Progonomyia)	4.141
domballah Alexander, 1939, Limonia (Neolimonia)	4.71
domestica (Osten Sacken), 1859 (Rhipidia), Limonia (Rhipidia) ..	4.75
domingensis Alexander, 1939, Dolichocheza (Megistomastix)	4.8
domingensis(Alexander), 1916 (Eriocera), Hexatoma (Eriocera) .	4.101
domingensis (Alexander), 1916 (Geranomyia), Limonia (Gera- nomyia)	4.61
domingensis Alexander, 1937, Toxorhina (Toxorhina)	4.167
dominicæ Alexander, 1939, Polymera (Polymera)	4.111
dominicana Alexander, 1939, Limonia (Geranomyia)	4.62



<i>dominicana</i> Alexander, 1947, <i>Trentcophlia</i> (<i>Paramongoma</i>)	4.169
<i>dominicensis</i> Alexander, 1939, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
<i>dorsolineata</i> Alexander, 1945, <i>Styngomyia</i>	4.157
<i>dorsolobata</i> Alexander, 1945, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>dotata</i> Alexander, 1945, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>dotalis</i> Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
<i>drepanuchus</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>dubia</i> Alexander, 1943, <i>Aymaramyia</i>	4.119
<i>ducalis</i> Alexander, 1949, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>ductilis</i> Alexander, 1938, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>duidae</i> Alexander, 1931, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>duidensis</i> Alexander, 1931, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
<i>duidensis</i> Alexander, 1931, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>duplex</i> Alexander, 1937, <i>Gnophomyia</i>	4.128
<i>durabilis</i> Alexander, 1946, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.134
<i>durangensis</i> Alexander, 1953, <i>Nephrotoma</i>	4.17
<i>duseni</i> (Alexander), 1920 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
<i>duseni</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>duurvoorti</i> Alexander, 1928, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>ecalcarata</i> Alexander, 1947, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.120
<i>echo</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>cdax</i> Alexander, 1948, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>edwardsi</i> Bellardi, 1859, <i>Tipula</i> (<i>Bellardina</i>)	4.20
<i>edwardsina</i> Alexander, 1929, <i>Valdiviana</i>	4.41
<i>Edwardsomyia</i> Alexander, 1929	4.93
<i>effecta</i> Alexander, 1944, <i>Ozodicera</i> (<i>Dihexaclonus</i>)	4.18
<i>effera</i> Alexander, 1929, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>efferox</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>effeta</i> Alexander, 1921, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>efficax</i> Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>efficiens</i> Alexander, 1943, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>effulta</i> Alexander, 1941, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>egae</i> (Alexander), 1921 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Caenoglochina</i>) ..	4.48
<i>egens</i> Alexander, 1952, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>egregia</i> (Alexander), 1929 (<i>Philippiana</i>), <i>Neophilippiana</i>	4.154
<i>eiseni</i> Alexander, 1913, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>eiseni</i> (Alexander), 1912 (<i>Furcomyia</i>), <i>Limonia</i> (<i>Neolimonia</i>) ...	4.71
<i>elegans</i> (Fabricius), 1805 (<i>Tipula</i>), <i>Nephrotoma</i>	4.17
<i>elegantior</i> (Alexander), 1945 (<i>Tanyproctna</i>), <i>Leptotarsus</i> (<i>Tanyproctna</i>)	4.14
<i>elegantula</i> (Alexander), 1913 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>elegantula</i> (Williston), 1896 (<i>Pachyrrhina</i>), <i>Nephrotoma</i>	4.17
<i>Elephantomyia</i> Osten Sacken, 1859	4.93, 4.94
<i>Elephantomyia</i> Alexander, 1938	4.95
<i>eliana</i> Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
<i>eliana</i> Alexander, 1953, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
<i>elissa</i> Alexander, 1948, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>einora</i> Alexander, 1929, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
<i>einora</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>elnorae</i> (Alexander), 1929 (<i>Ctedonia</i>), <i>Gynoplistia</i> (<i>Gynoplistia</i>) ..	4.99
<i>Elnoretta</i> Alexander, 1929	4.8



<i>elongata</i> Alexander, 1943, <i>Orimarga</i> (<i>Diotrepha</i>)	4.81
<i>elongata</i> Macquart, 1838 (<i>Pachyrhina</i>)	4.44
<i>elquiensis</i> (Blanchard), 1852 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>eluta</i> (Schiner), 1868 (<i>Tipula</i>), <i>Zelandotipula</i>	4.43
<i>emarginata</i> Alexander, 1945, <i>Gnophomyia</i>	4.128
<i>emarginatus</i> Alexander, 1937, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>emerita</i> Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>eminens</i> Alexander, 1944, <i>Nephrotoma</i>	4.17
<i>Empeda</i> Osten Sacken, 1869	4.119
<i>Empedomorpha</i> Alexander, 1916	4.122
<i>empelia</i> Alexander, 1962, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>enderleinana</i> Alexander, 1929, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>enderleini</i> (Alexander), 1913 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>enixa</i> Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>enixoides</i> Alexander, 1962, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>epicosma</i> Alexander, 1926, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
<i>epione</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>Epiphragma</i> Osten Sacken, 1859	4.95
<i>epiphragmoides</i> (Alexander), 1913 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Limnophilella</i>)	4.92
<i>erectus</i> Alexander, 1953, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>cremnoptera</i> Alexander, 1967, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
<i>Eriocera</i> Macquart, 1838	4.99
<i>Erioptera</i> Meigen, 1803	4.122
<i>Eriopterella</i> Alexander, 1929	4.123
<i>Eriopterini</i>	4.117
<i>eriopteroides</i> Alexander, 1926, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>erostrata</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>errabunda</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>errans</i> Alexander, 1944, <i>Trentepohlia</i> (<i>Mongoma</i>)	4.169
<i>erubescens</i> Alexander, 1944, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
<i>erythraea</i> (Osten Sacken), 1886 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>erythrocephala</i> (Fabricius), 1805 (<i>Tipula</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
<i>esau</i> Alexander, 1947, <i>Limonia</i> (<i>Neoglochina</i>)	4.68
<i>esmeralda</i> Alexander, 1952, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>espinozai</i> (Alexander), 1938 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
<i>eucrypta</i> (Alexander), 1953 (<i>Longurio</i>), <i>Leptotarsus</i> (<i>Longurio</i>) ..	4.12
<i>eudorae</i> (Alexander), 1913 (<i>Peripheroptera</i>), <i>Limonia</i> (<i>Peripherop-</i> <i>terae</i>)	4.72
<i>Eufurina</i> Alexander, 1946	4.156
<i>Eugnophomyia</i> Alexander, 1947	4.126
<i>Eulimnophila</i> Alexander, 1919	4.110
<i>Eumicrotipula</i> Alexander, 1922	4.21
<i>Eumolophilus</i> Alexander, 1921	4.143
<i>eumonostylus</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>Euparatropesa</i> Alexander, 1946	4.157
<i>cuphrosyne</i> Alexander, 1945, <i>Erioptera</i> (<i>Mesocycphona</i>)	4.124
<i>Euptilostena</i> Alexander, 1938	4.132
<i>eurygramma</i> (Alexander), 1928 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Gerano-</i> <i>myia</i>)	4.62
<i>euryleon</i> Alexander, 1950, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>eurymera</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>euryphallus</i> Alexander, 1967, <i>Limonia</i> (<i>Rhipidia</i>)	4.75




<i>euryptera</i> Alexander, 1953, <i>Limonia</i> (<i>Peripheroptera</i>)	4.72
<i>eurystyla</i> Alexander, 1953, <i>Ozodicera</i> (<i>Ozodieera</i>)	4.19
<i>eutacniata</i> (Bigot), 1888 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Austro-</i> <i>limnophila</i>)	4.90
<i>cuterpe</i> (Alexander), 1945 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Icriomastix</i>)	4.8
<i>Euteucholabis</i> Alexander, 1946	4.158
<i>eutheta</i> Alexander, 1935, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
<i>evanescens</i> Alexander, 1948, <i>Polymera</i> (<i>Polymicrodes</i>)	4.113
<i>exaeta</i> (Alexander), 1927 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>excavatus</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>excelsior</i> Alexander, 1944, <i>Limonia</i> (<i>Neolimnobia</i>)	4.70
<i>excessiva</i> Alexander, 1926, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
<i>excordis</i> (Alexander), 1951 (<i>Gnophomyia</i>), <i>Eugnophomyia</i>	4.127
<i>exemptus</i> (Alexander), 1936 (<i>Habromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i> <i>gurio</i>)	4.12
<i>exemptus</i> Alexander, 1953, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>exercita</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>exilis</i> Alexander, 1916, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24
<i>expansa</i> Alexander, 1938, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>expleta</i> Alexander, 1962, <i>Tipula</i> (<i>Eumlerotipula</i>)	4.24
<i>expressa</i> Alexander, 1937, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
<i>exquisita</i> (Alexander), 1914 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ...	4.101
<i>extensa</i> Alexander, 1914, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>extensa</i> Alexander, 1921, <i>Ozodieera</i> (<i>Ozodicera</i>)	4.19
<i>extensa</i> (Alexander), 1913 (<i>Mongoma</i>), <i>Trentepohlia</i> (<i>Paramon-</i> <i>goma</i>)	4.169
<i>extenuata</i> Alexander, 1945, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>extranea</i> Alexander, 1944, <i>Limonia</i> (<i>Dieranomyia</i>)	4.52
<i>extraordinarius</i> Alexander, 1938, <i>Amphineurus</i> (<i>Amphineurus</i>) ...	4.117
<i>fabricii</i> Alexander, 1913, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>facinus</i> Alexander, 1943, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
<i>factiosa</i> Alexander, 1943, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>fagetorum</i> (Alexander), 1929 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Isehnotoma</i>)	4.9
<i>fagetorum</i> Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
<i>fagetorum</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>falcicula</i> Alexander, 1962, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>falcifer</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>falcifera</i> Alexander, 1921, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>falklandiea</i> (Enderlein), 1906 (<i>Zalusa</i>), <i>Limonia</i> (<i>Zalusa</i>)	4.79
<i>falx</i> Alexander, 1938, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
<i>farri</i> Alexander, 1951, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>farri</i> Alexander, 1964, <i>Limonia</i> (<i>Dicranomyia</i>)	4.52
<i>farriana</i> Alexander, 1964, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>farriana</i> Alexander, 1964, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
<i>fasciata</i> (Guérin-Ménéville), 1830 (<i>Evaniptera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>fasciata</i> (Williston), 1900 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>fasciolaris</i> (Wiedemann), 1828 (<i>Limnobia</i>), <i>Teucholabis</i> (<i>Eupara-</i> <i>tropesa</i>)	4.158
<i>fascipennis</i> Osten Sacken, 1887, <i>Gnophomyia</i>	4.129
<i>fassliana</i> (Alexander), 1962 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
<i>fatidica</i> Alexander, 1940, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.24




faustina Alexander, 1945, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.169
fazi Alexander, 1928, <i>Tipula</i> (<i>Eumierotipula</i>)	4.24
feliciana Alexander, 1945, <i>Tipula</i> (<i>Mierotipula</i>)	4.34
felix Alexander, 1943, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
felix Alexander, 1950, <i>Limonia</i> (<i>Neoglochina</i>)	4.68
felix Alexander, 1956, <i>Sigmatomera</i> (<i>Sigmatomera</i>)	4.156
femoranigra Alexander, 1913, <i>Erioptera</i> (<i>Mesoccyphona</i>)	4.124
femorata Alexander, 1921, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
ferax Alexander, 1937, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
feriata (Alexander), 1929 (<i>Limnophila</i>), <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
ferocia Alexander, 1936, <i>Tipula</i> (<i>Mierotipula</i>)	4.35
ferruginea (Fabricius), 1805 (<i>Tipula</i>), <i>Nephrotoma</i>	4.17
ferruginea Williston, 1900, <i>Gnophomyia</i>	4.128
fessa Alexander, 1944, <i>Gnophomyia</i>	4.128
fidelis (Alexander), 1929 (<i>Isechnothrix</i>), <i>Atarba</i> (<i>Isechnothrix</i>)	4.88
fiibrigi Alexander, 1922, <i>Atarba</i> (<i>Atarba</i>)	4.86
fiibrigi Alexander, 1941, <i>Tipula</i> (<i>Mierotipula</i>)	4.35
fieldi Alexander, 1966, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
fieldi Alexander, 1967, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
fieldianus (Alexander), 1956 (<i>Longurio</i>), <i>Leptotarsus</i> (<i>Tanyremna</i>)	4.14
filicauda (Alexander), 1925 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Dieranomyia</i>)	4.53
filiforme Alexander, 1940, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
filiformis Alexander, 1929, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
filigera Walker, 1856 (<i>Tipula</i>)	4.44
filiolus Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
filius Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
fissilis Alexander, 1926, <i>Limonia</i> (<i>Atypophthalmus</i>)	4.47
<i>Flabellifera</i> , authors	4.7
flabellifera Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
flabellifera (Loew), 1851 (<i>Ptilogyna</i>), <i>Phaeclodoera</i>	4.20
flabelliformis Alexander, 1934, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
flagrans (Alexander), 1928 (<i>Gnophomyia</i>), <i>Eugnophomyia</i>	4.127
flammeinota (Alexander), 1930 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
flammeipennis Alexander, 1942, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
flammeithorax (Alexander), 1949 (<i>Gnophomyia</i>), <i>Eugnophomyia</i>	4.127
flavaperta Alexander, 1941, <i>Limonia</i> (<i>Dieranomyia</i>)	4.52
flavella Alexander, 1921, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
flavexemptus Alexander, 1968, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
flavibasis Alexander, 1916, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
flavicauda (Bigot), 1888 (<i>Poecilcostola</i>)	4.117
flavida (Williston), 1900 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
flavieornis (Alexander), 1914 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
flavicosta (Alexander), 1928 (<i>Diotrepha</i>), <i>Orimarga</i> (<i>Diotrepha</i>) .	4.81
flavida (Williston), 1900 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
flavida (Philippi), 1865 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dieranomyia</i>)	4.52
flavida Alexander, 1913, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.167
flavidula Alexander, 1952, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
flavidula Alexander, 1940, <i>Tipula</i> (<i>Eumierotipula</i>)	4.24
flavidula Alexander, 1914, <i>Molophilus</i> (<i>Molophilus</i>)	4.146
flavipennis (Philippi), 1865 (<i>Ctedonia</i>), <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
flavipennis Osten Saeken, 1873, <i>Sigmatomera</i> (<i>Sigmatomera</i>)	4.156
flavipennis Philippi, 1865 (<i>Tipula</i>)	4.44
flavithorax (Wiedemann), 1821 (<i>Limnobia</i>), <i>Teucholabis</i> (<i>Teucho-</i>	

labis)	4.161
<i>flavithorax</i> Wiedemann of Sehiner, 1868, Teucholabis	4.165
<i>flavoannulata</i> Jacobs, 1900, Tipula (Eumierotipula)	4.24
<i>flavofaseialis</i> (Alexander), 1924 (Dieranomyia), Limonia (Dieranomyia)	4.52
<i>flavofimbria</i> Alexander, 1968, Teucholabis (Teucholabis)	4.161
<i>flavogenualls</i> (Alexander), 1945 (Holorusia), Zelandotipula	4.42
<i>flavopolita</i> Alexander, 1942, Tipula (Mierotipula)	4.35
<i>flebilis</i> Alexander, 1944, Gnophomyia	4.128
<i>flemingi</i> Alexander, 1950, Molophilus (Molophilus)	4.146
<i>flexillstylus</i> Alexander, 1931, Molophilus (Molophilus)	4.146
<i>flinti</i> Alexander, 1967, Molophilus (Molophilus)	4.146
<i>fluxa</i> Alexander, 1941, Limonia (Geranomyia)	4.62
<i>foersteri</i> Alexander, 1952, Teucholabis (Teucholabis)	4.161
<i>foersteriana</i> Alexander, 1967, Tipula (Eumierotipula)	4.24
<i>foreeps</i> Alexander, 1941, Gonomyia (Progonomyia)	4.141
<i>formosissima</i> Alexander, 1946, Teucholabis (Teucholabis)	4.161
<i>forsteri</i> Alexander, 1962, Tipula (Eumierotipula)	4.24
<i>forsteriana</i> Alexander, 1962, Limonia (Geranomyia)	4.62
<i>forsteriana</i> (Alexander), 1962 (Holorusia), Zelandotipula	4.42
<i>fortibasis</i> Alexander, 1966, Gonomyia (Lipophleps)	4.136
<i>fortieornis</i> Alexander, 1947, Atarba (Atarba)	4.86
<i>fortieornis</i> Alexander, 1937, Shannonomyia (Shannonomyia)	4.114
<i>fortidens</i> Alexander, 1951, Molophilus (Molophilus)	4.146
<i>fortior</i> Alexander, 1951, Tipula (Eumierotipula)	4.25
<i>fragilis</i> Loew, 1851, Toxorhina (Toxorhina)	4.168
<i>fragosus</i> Alexander, 1937, Helius	4.46
<i>fraudentula</i> Alexander, 1928, Tipula (Eumierotipula)	4.25
<i>frauenfeldi</i> (Sehiner), 1868 (Tipula), Ischnotoma (Ischnotoma) ..	4.9
<i>fucgiensis</i> Alexander, 1920, Tipula (Eumierotipula)	4.25
<i>fulgens</i> Alexander, 1913, Teucholabis (Teucholabis)	4.161
<i>fuliginosa</i> (Sehiner), 1868 (Pentoptera), Hexatoma (Eriocera) ..	4.104
<i>fulvinota</i> Alexander, 1962, Teucholabis (Teucholabis)	4.161
<i>fulvistigma</i> Alexander, 1953, Limonia (Peripheroptera)	4.72
<i>fulvithorax</i> Alexander, 1966, Hexatoma (Eriocera)	4.101
<i>fulvithorax</i> (Philippi), 1865 (Pettusa), Limonia (Zelandogloehina)	4.80
<i>fumibasalis</i> Alexander, 1937, Limonia (Peripheroptera)	4.73
<i>fumicosta</i> (Alexander), 1921 (Diotrepha), Orimarga (Diotrepha) .	4.81
<i>fumidapicalis</i> Alexander, 1942, Teucholabis (Euparatropesa)	4.158
<i>fumipennis</i> (Alexander), 1919 (Rhamphidia), Helius	4.45
<i>fumipennis</i> Loew, 1851, Ozodieera (Dihexaclonus)	4.18
<i>fumipennis</i> Alexander, 1939, Rhabdomastix (Saeandaga)	4.155
<i>fumipennis</i> Alexander, 1939, Toxorhina (Toxorhina)	4.168
<i>fumipennis</i> Alexander, 1912 (Tipula)	4.44
<i>fumipes</i> Alexander, 1928, Elephantomyia (Elephantomyia)	4.94
<i>fumosa</i> (Alexander), 1912 (Furcomyia), Limonia (Neogloehina) .	4.68
<i>funebri</i> (Alexander), 1922 (Gnophomyia), Eugnophomyia	4.127
<i>funerea</i> (Alexander), 1938 (Gnophomyia), Eugnophomyia	4.127
<i>funerula</i> Alexander, 1929, Orimarga (Orimarga)	4.82
<i>fureifera</i> (Alexander), 1944 (Holorusia), Zelandotipula	4.42
<i>Furcomyia</i> Meigen, 1818	4.49
<i>fureula</i> Alexander, 1962, Gonomyia (Gonomyia)	4.133
<i>furcus</i> Alexander, 1951, Molophilus (Molophilus)	4.146




<i>Furina</i> Jaennicke, 1867	4.156
furor Alexander, 1944, Limonia (Geranomyia)	4.62
furva Alexander, 1930, Teucholabis (Teucholabis)	4.161
fusca (Jaennicke), 1867 (Ctedonia), Gynoplistia (Dirhipis)	4.98
fusca Wiedemann, 1828, Polymera (Polymera)	4.111
fuscifusus Alexander, 1929, Amphineurus (Rhamphoneurus)	4.118
fuscipennis (Alexander), 1914 (Psaronius), Lecteria (Psaronius) ..	4.106
fuscipes (Alexander), 1919 (Rhamphidia), Helius	4.45
fuscipes Alexander, 1921, Trentepohlia (Paramongoma)	4.170
fuscitarsis (Alexander), 1919 (Tanypremna), Leptotarsus (Tany- premna)	4.14
fuscitarsis Alexander, 1937, Polymera (Polymera)	4.111
fuscivena Alexander, 1923, Erioptera (Mesocyphona)	4.124
fuscoanalis Alexander, 1946, Ctenolimnophila (Ctenolimnophila) ..	4.93
fuscoapicalis Alexander, 1944, Atarba (Atarba)	4.86
fuscoapicalis Alexander, 1937, Teucholabis (Teucholabis)	4.161
fuscocostalis Alexander, 1947, Gnophomyia	4.129
fuscocostata (Alexander), 1959 (Gnophomyia), Eugnophomyia ...	4.127
fuscodiscalis Alexander, 1938, Erioptera (Mesocyphona)	4.124
fuscodhalterata Alexander, 1929, Austrolimnophila (Austrolimno- phila)	4.90
fuscolimbata Alexander, 1950, Trentepohlia (Paramongoma)	4.170
fuscolineata Alexander, 1929, Limonia (Dicranomyia)	4.56
fuscopleuralis Alexander, 1927, Molophilus (Molophilus)	4.147
fuscostigmalis (Alexander), 1966 (Limnophila), Shannonomyia (Shannonomyia)	4.115
fuscostigmata (Alexander), 1929 (Holorusia), Ischnotoma (Ischno- toma)	4.9
fuscoterminalis Alexander, 1949, Trentepohlia (Neomongoma)	4.169
fuscovenosa Alexander, 1928, Cryptolabis (Cryptolabis)	4.121
gaigei Alexander, 1928, Epiphragma (Epiphragma)	4.96
galapagoensis Alexander, 1962, Limonia (Dicranomyia)	4.53
galatea Alexander, 1949, Teucholabis (Teucholabis)	4.161
galbipes Alexander, 1967, Limonia (Caenolimonia)	4.49
gargantua Alexander, 1941, Molophilus (Molophilus)	4.147
gaudens (Alexander), 1922 (Geranomyia), Limonia (Geranomyia)	4.62
geijskesana Alexander, 1946, Sigmatomera (Sigmatomera)	4.156
geijskesi Alexander, 1945, Brachypremna	4.6
<i>gelatifer</i> Edwards, 1934, Limonia (Geranomyia)	4.65
geminata Alexander, 1943, Atarba (Ischnothrix)	4.88
gemula Alexander, 1946, Gonomyia (Gonomyia)	4.133
geniculata Alexander, 1915, Polymera (Polymera)	4.111
geniculata (Alexander), 1914 (Mongoma), Trentepohlia (Para- mongoma)	4.170
gentilis (Alexander), 1943 (Tanypremna), Leptotarsus (Tany- premnella)	4.15
genuale Alexander, 1934, Epiphragma (Epiphragma)	4.96
<i>Geotipula</i> Savchenko, 1964	4.40
Geranomyia Haliday, 1833	4.59
gesneri (Enderlein), 1912 (Aldrovandia), Leptotarsus (Aldrovandia)	4.11
gibbera (Alexander), 1916 (Dicranomyia), Limonia (Dicranomyia)	4.53
gillottae Alexander, 1928, Gonomyia (Lipophleps)	4.136

<i>gilvipennis</i> Alexander, 1928, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
<i>glabrieristata</i> Alexander, 1939, <i>Nephrotoma</i>	4.17
<i>glabripennis</i> (Alexander), 1948 (<i>Gnophomyia</i>), <i>Eugnophomyia</i> ...	4.127
<i>glabrispina</i> Alexander, 1926, <i>Gonomyia</i> (<i>Ncolipophleps</i>)	4.139
<i>glabristyla</i> Alexander, 1946, <i>Tipula</i> (<i>Microtipula</i>)	4.37
<i>glabristylatus</i> Alexander, 1929, <i>Amphineurus</i> (<i>Rhamphineurus</i>) ..	4.118
<i>glabritergata</i> Alexander, 1965, <i>Gnophomyia</i>	4.129
<i>gladiator</i> Alexander, 1919, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
<i>gladiator</i> Alexander, 1914, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>glaphyoptera</i> Philippi, 1865, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>glauca</i> (Alexander), 1916 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>) ..	4.62
<i>globulicornis</i> (Alexander), 1924 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>globulicornis</i> Alexander, 1968, <i>Shannonomyia</i> (<i>Shannonomyia</i>) ...	4.115
<i>glochinosoides</i> (Alexander), 1922 (<i>Peripheroptera</i>), <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
<i>gloriosa</i> (Alexander), 1912 (<i>Fureomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>glossophora</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>Gnophomyia</i> Osten Sacken, 1859	4.127
<i>golbaehi</i> (Alexander), 1962 (<i>Gnophomyia</i>), <i>Eugnophomyia</i>	4.127
<i>gomesi</i> Alexander, 1942, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>gomesiana</i> Alexander, 1945, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>Gonomyia</i> Osten Sacken, 1869	4.131
<i>Gonomyella</i> Alexander, 1917	4.141
<i>Gonomyia</i> Meigen, 1818	4.131, 4.132
<i>Gonomyina</i> Alexander, 1946	4.134
<i>gowdeyi</i> Alexander, 1932, <i>Limonia</i> (<i>Discobola</i>)	4.59
<i>gowdeyi</i> Alexander, 1928, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>goyazanus</i> (Alexander), 1940 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurlo</i>)	4.12
<i>goyazensis</i> Alexander, 1940, <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>goyazicola</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.34
<i>gracilicorne</i> Alexander, 1916, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>gracillior</i> Alexander, 1944, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
<i>gracilipes</i> (Walker), 1836 (<i>Tipula</i>), <i>Zelandotipula</i>	4.42
<i>gracilirama</i> Alexander, 1940, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
<i>gracilirama</i> Alexander, 1941, <i>Ozodicera</i> (<i>Dihexaclonus</i>)	4.18
<i>gracilis</i> (Osten Sacken), 1886 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.101
<i>gracilis</i> Osten Sacken, 1886, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>gracilis</i> (Westwood), 1835 (<i>Hemiteina</i>), <i>Ozodicera</i> (<i>Ozodicera</i>) ..	4.19
<i>graphica</i> Schiner, 1868, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>grisea</i> Alexander, 1913, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>griscipennis</i> Loew, 1851, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
<i>grossa</i> Alexander, 1938, <i>Limonia</i> (<i>Neoglochina</i>)	4.69
<i>grus</i> Alexander, 1941, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>guarani</i> Alexander, 1914, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>guatemalensis</i> (Alexander), 1916 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>guatemalensis</i> Alexander, 1913, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>guatemalensis</i> (Alexander), 1926 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.41
<i>guato</i> Alexander, 1912, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>guayaquilensis</i> Alexander, 1938, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
<i>guerrerensis</i> Alexander, 1946, <i>Limonia</i> (<i>Rhipidia</i>)	4.76



guerreroensis Alexander, 1940, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
guerreroensis Alexander, 1940, <i>Tipula</i> (<i>Microtipula</i>)	4.35
guianensis Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.62
guianensis Alexander, 1934, <i>Ozodicera</i> (<i>Dihexaelonus</i>)	4.18
guimaraesi (Alexander), 1942 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
gulosa Alexander, 1943, <i>Ericptera</i> (<i>Mesocyphona</i>)	4.124
<i>guttata</i> (Philippi), 1865 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.51
<i>guttiecllula</i> Alexander, 1936, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>guttistigma</i> Alexander, 1941, <i>Pedicia</i> (<i>Trieyphona</i>)	4.84
<i>guttulatissima</i> Alexander, 1913, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
<i>guzmani</i> (Alexander), 1953 (<i>Holorusia</i>), <i>Isehnotoma</i> (<i>Isehnotoma</i>)	4.10
<i>Gymnastes</i> Brunetti, 1911	4.142
<i>gymnocera</i> (Alexander), 1938 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
<i>gymnoeladus</i> Alexander, 1928, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>Gynoplistia</i> Westwood, 1835	4.98
<i>haagi</i> Alexander, 1946, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>Habrolimnophila</i> Alexander, 1968	4.107
<i>haemorrhoea</i> (Osten Sacken), 1886 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.101
<i>haitensis</i> Alexander, 1939, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
<i>halophila</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>halterata</i> Alexander, 1966, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
<i>haploa</i> Alexander, 1926, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>haploides</i> Alexander, 1938, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>harpax</i> Alexander, 1952, <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>harpysia</i> Alexander, 1946, <i>Dicranoptyeha</i>	4.45
<i>hatschbachi</i> Alexander, 1954, <i>Helius</i>	4.46
<i>hazela</i> Alexander, 1929, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
<i>hedymopa</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.25
<i>helenae</i> Alexander, 1948, <i>Atarba</i> (<i>Isehnotherix</i>)	4.88
<i>heliconiae</i> Alexander, 1945, <i>Neognophomyia</i>	4.153
<i>helios</i> (Alexander), 1949 (<i>Holorusia</i>), <i>Isehnotoma</i> (<i>Ieriomastax</i>) ..	4.9
<i>Helius</i> Lepelletier & Serville, 1828	4.45
<i>Helobia</i> Lepelletier & Serville, 1828	4.126
<i>helophila</i> Alexander, 1916, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>Hemiteina</i> Westwood, 1835	4.17
<i>hera</i> Alexander, 1949, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>hesione</i> Alexander, 1950, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>heteracantha</i> Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>heterodaetyla</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>heteromera</i> Alexander, 1942, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
<i>heteropoda</i> Alexander, 1943, <i>Teucholabis</i> (<i>Euparatropesa</i>)	4.158
<i>heteroxipha</i> Alexander, 1942, <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>hexamelania</i> Alexander, 1951, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>Hexatoma</i> Latreille, 1809	4.99
<i>hilaris</i> Alexander, 1945, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>hilaris</i> Alexander, 1913, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>hirsuta</i> (Alexander), 1913 (<i>Gnophomyia</i>), <i>Neognophomyia</i>	4.153
<i>hirsutinota</i> Alexander, 1943, <i>Limonia</i> (<i>Geranomyia</i>)	4.62
<i>hirsutipes</i> Alexander, 1929, <i>Mesolimnophila</i>	4.109

<i>hirsutissima</i> Alexander, 1962, <i>Limonia</i> (<i>Dieranomyia</i>)	4.53
<i>hirticornis</i> Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>hirticornis</i> (Fabricius), 1805 (<i>Chironomus</i>), <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>hirtilobata</i> Alexander, 1939, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>hirtistylata</i> Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>histrionica</i> Alexander, 1945, <i>Erioptera</i> (<i>Mesocycphona</i>)	4.124
<i>histrionica</i> Schiner, 1868, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.96
<i>histrionica</i> Alexander, 1943, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>histrionica</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>hodgei</i> (Alexander), 1939 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tanypremna</i>)	4.14
<i>hoffmani</i> Alexander, 1927, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
<i>hoffmani</i> Alexander, 1936, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
<i>hoffmaniana</i> Alexander, 1947, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>hoffmaniana</i> Alexander, 1937, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
<i>homilacantha</i> Alexander, 1968, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>homochroa</i> (Alexander), 1926 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Ischnotoma</i>)	4.9
<i>hondurensis</i> Alexander, 1941, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>honesta</i> Alexander, 1940, <i>Polymera</i> (<i>Polymera</i>)	4.111
<i>honestus</i> Alexander, 1923, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>hoogstraali</i> Alexander, 1946, <i>Gonomyia</i> (<i>Idioera</i>)	4.135
<i>hoogstraali</i> Alexander, 1940, <i>Tipula</i> (<i>Tripletipula</i>)	4.40
<i>horni</i> (Alexander), 1926 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
<i>horrenda</i> Alexander, 1944, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>horribilis</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>horridus</i> (Alexander), 1941 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tanypremna</i>)	4.14
<i>hostica</i> Alexander, 1938, <i>Limonia</i> (<i>Dieranomyia</i>)	4.53
<i>hostica</i> Alexander, 1943, <i>Neognophomyia</i>	4.153
<i>hostifica</i> Alexander, 1937, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>huacapistanae</i> Alexander, 1941, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>huanca</i> Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>huanucensis</i> (Alexander), 1953 (<i>Longurio</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
<i>humerosa</i> Alexander, 1942, <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>humidicola</i> Alexander, 1929, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
<i>humilis</i> Alexander, 1931, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.94
<i>hylonympha</i> Alexander, 1929, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
<i>hylonympha</i> Alexander, 1929, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>hyperarmatus</i> Alexander, 1939, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>hyperphallus</i> Alexander, 1938, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>hyperplatys</i> Alexander, 1944, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>hypoerita</i> Alexander, 1935, <i>Limonia</i> (<i>Neolimnobia</i>)	4.70
<i>hypomela</i> Alexander, 1948, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>hystrix</i> Alexander, 1939, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>icasta</i> Alexander, 1941, <i>Tipula</i> (<i>Microtipula</i>)	4.35
<i>Teriomastax</i> Enderlein, 1912	4.8
<i>Idioera</i> Dale, 1842	4.134
<i>Idioglochina</i> Alexander, 1929	4.68
<i>Idioneura</i> Philippi, 1865	4.126
<i>idoneurus</i> (Alexander), 1924 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Ceoneura</i>)	4.11
<i>idiophallus</i> Alexander, 1942, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.161
<i>idonea</i> Alexander, 1939, <i>Atarba</i> (<i>Atarba</i>)	4.86



ignara Alexander, 1967, Limonia (Dicranomyia)	4.53
ignava Alexander, 1943, Shannonomyia (Shannonomyia)	4.115
ignithorax Alexander, 1929, Atarba (Ischnothrix)	4.88
iguazuensis (Alexander), 1928 (Microtipula), Tipula (Eumicrotipula)	4.25
illectus Alexander, 1941, Molophilus (Molophilus)	4.147
illepida Alexander, 1944, Limonia (Dicranomyia)	4.53
illicis Alexander, 1921, Gonomyia (Gonomyia)	4.133
illudens Alexander, 1945, Brachypremna	4.6
illudens Alexander, 1914, Rhabdomastix (Rhabdomastix)	4.155
illuminata Alexander, 1967, Limonia (Rhipidia)	4.76
imitabilis Alexander, 1942, Limonia (Dicranomyia)	4.53
imitans Alexander, 1913, Epiphragma (Epiphragma)	4.96
immaculata Alexander, 1913, Erioptera (Mesocyphona)	4.124
immaculipennis (Alexander), 1926 (Gnophomyia), Neognophomyia	4.153
immaculipes Alexander, 1935, Limonia (Neolimnobia)	4.70
immaculipes Alexander, 1941, Epiphragma (Epiphragma)	4.97
immaculipleura Alexander, 1946, Teucholabis (Teucholabis)	4.161
immaculosa Alexander, 1929, Limonia (Dicranomyia)	4.57
immerens Alexander, 1944, Tipula (Eumicrotipula)	4.25
immerita Alexander, 1929, Limonia (Geranomyia)	4.62
immodica Alexander, 1938, Limonia (Neolimonia)	4.71
immorsa Alexander, 1945, Tipula (Eumicrotipula)	4.25
impatiens Alexander, 1951, Tipula (Microtipula)	4.35
impedita Alexander, 1946, Gonomyia (Lipophleps)	4.136
imperturbata Alexander, 1951, Limonia (Neoglochina)	4.69
improperata Alexander, 1936, Limonia (Rhipidia)	4.76
inaequiarmata Alexander, 1962, Tipula (Eumicrotipula)	4.25
inaequicineta Alexander, 1941, Epiphragma (Epiphragma)	4.97
inacquidens Alexander, 1946, Tipula (Eumicrotipula)	4.25
inaequilobata Alexander, 1937, Tipula (Microtipula)	4.35
inaequipctinata Alexander, 1929, Limonia (Rhipidia)	4.76
inaequispinosa Alexander, 1940, Limonia (Geranomyia)	4.62
inaequituberculata Alexander, 1929, Limonia (Geranomyia)	4.62
inarmata Alexander, 1928, Tipula (Microtipula)	4.35
inarmatus Alexander, 1939, Molophilus (Molophilus)	4.147
inca Alexander, 1968, Teucholabis (Teucholabis)	4.161
inca Alexander, 1912, Tipula (Eumicrotipula)	4.25
incisurata Alexander, 1938, Atarba (Atarba)	4.86
incognitus Alexander, 1946, Molophilus (Molophilus)	4.147
incommoda (Osten Sacken), 1887 (Peripheroptera), Limonia (Peripheroptera)	4.73
incommoda (Osten Sacken), 1887 (Peripheroptera), Limonia (Peripheroptera)	4.73
incommodus Alexander, 1928, Limonia (Peripheroptera)	4.73
incompleta (Alexander), 1944 (Tanyremna), Leptotarsus (Tanyremna)	4.14
incondita Alexander, 1951, Tipula (Eumicrotipula)	4.25
inculta Alexander, 1929, Limnophila (Limnophila)	4.108
indefensa Alexander, 1939, Limonia (Dicranomyia)	4.53
indomita Alexander, 1938, Limonia (Neolimonia)	4.71
indotata Alexander, 1966, Gonomyia (Paralipophleps)	4.140
indurabilis Alexander, 1966, Molophilus (Molophilus)	4.147
inepta Alexander, 1942, Teucholabis (Teucholabis)	4.162
ineptus Alexander, 1938, Helius	4.46



<i>inermis</i> Alexander, 1914, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>infamosa</i> Alexander, 1937, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>infernalis</i> (Alexander), 1947 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
<i>infestiva</i> (Alexander), 1929 (<i>Gynoplistia</i>), <i>Paralimnophila</i>	4.110
<i>infida</i> Alexander, 1941, <i>Tipula</i> (<i>Mierotipula</i>)	4.35
<i>infidelis</i> Alexander, 1929, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.90
<i>infidelis</i> Alexander, 1951, <i>Tipula</i> (<i>Eumierotipula</i>)	4.25
<i>infinita</i> Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.25
<i>inflexibilis</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>infueata</i> Alexander, 1927, <i>Limonia</i> (<i>Neogloehina</i>)	4.69
<i>infumata</i> (Philippi), 1865 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dieranomyia</i>) ...	4.53
<i>infumipennis</i> Alexander, 1942, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
<i>ingens</i> Alexander, 1945, <i>Limonia</i> (<i>Neogloehina</i>)	4.69
<i>ingenua</i> Alexander, 1945, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>ingloria</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>ingrata</i> (Alexander), 1928 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>innubens</i> Alexander, 1942, <i>Tipula</i> (<i>Eumierotipula</i>)	4.26
<i>inornata</i> Alexander, 1913, <i>Polymera</i> (<i>Polymera</i>)	4.112
<i>inornatipes</i> Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>inornatipes</i> Alexander, 1925, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>inquieta</i> Alexander, 1943, <i>Limnophila</i> (<i>Limnophilella</i>)	4.92
<i>inquisita</i> Alexander, 1942, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>insanus</i> Alexander, 1952, <i>Amphineurus</i> (<i>Rhamphoneurus</i>)	4.118
<i>insidiosus</i> (Alexander), 1949 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i> <i>gurio</i>)	4.12
<i>insignifica</i> (Alexander), 1912 (<i>Furcomyia</i>), <i>Limonia</i> (<i>Dieranomyia</i>)	4.53
<i>insignis</i> (Loew), 1851 (<i>Aporosa</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>insolabilis</i> Alexander, 1946, <i>Limonia</i> (<i>Dicranomyia</i>)	4.53
<i>insolabilis</i> Alexander, 1944, <i>Tipula</i> (<i>Mierotipula</i>)	4.32
<i>instrenua</i> (Alexander), 1942 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>)	4.119
<i>insularis</i> (Williston), 1896 (<i>Limnobia</i>), <i>Limonia</i> (<i>Neogloehina</i>) ...	4.69
<i>insularis</i> (Johnson), 1913 (<i>Limnophila</i>)	4.117
<i>integriloba</i> Alexander, 1943, <i>Atarba</i> (<i>Ischnothrix</i>)	4.88
<i>integristigma</i> Alexander, 1940, <i>Brachypremna</i>	4.6
<i>integristylus</i> Alexander, 1945, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
<i>interperata</i> Alexander, 1943, <i>Tipula</i> (<i>Mierotipula</i>)	4.35
<i>interecepta</i> Alexander, 1947, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>interlineata</i> Alexander, 1934, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
<i>intermedia</i> (Alexander), 1927 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.102
<i>intermedia</i> (Walker), 1848 (<i>Limnobia</i>), <i>Limonia</i> (<i>Geranomyia</i>) ..	4.63
<i>intermedia</i> Alexander, 1929, <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
<i>interrupta</i> Alexander, 1944, <i>Neognophomyia</i>	4.153
<i>interspersa</i> Alexander, 1953, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>interstitialis</i> Alexander, 1940, <i>Limonia</i> (<i>Caenolimonia</i>)	4.49
<i>inulta</i> Alexander, 1936, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162
<i>invalida</i> (Alexander), 1916 (<i>Dicranomyia</i> , <i>Limonia</i> (<i>Dicranomyia</i>))	4.53
<i>invariegata</i> Alexander, 1921, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.124
<i>invariegatus</i> Alexander, 1933, <i>Helius</i>	4.46
<i>invaripennis</i> Alexander, 1941, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>invaripes</i> Alexander, 1939, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>invaripes</i> (Alexander), 1936 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premna</i>)	4.14
<i>invaripes</i> Alexander, 1936, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162




invenusta Alexander, 1946, Teucholabis (Euparatropesa)	4.158
invigilans Alexander, 1951, Tipula (Eumicrotipula)	4.26
iquitosensis Alexander, 1945, Erioptera (Mesocyphona)	4.125
iris Alexander, 1929, Austrolimnophila (Austrolimnophila)	4.90
irrorata (Philippi), 1865 (Polymoria), Paralimnophila	4.110
isabella Alexander, 1927, Rhabdomastix (Rhabdomastix)	4.155
Ischnothrix Bigot, 1888	4.88
Ischnotoma Skuse, 1890	4.8, 4.9
isolatus Alexander, 1952, Molophilus (Molophilus)	4.147
itaiayana Alexander, 1944, Brachypremna	4.6
itaiayana Alexander, 1944, Limonia (Dicranomyia)	4.54
itaiayensis Alexander, 1944, Tipula (Eumicrotipula)	4.26
ixion (Alexander), 1949 (Macromastix), Leptotarsus (Longurio) ..	4.12
jacobsiana Alexander, 1929, Tipula (Eumicrotipula)	4.26
jactans (Alexander), 1913 (Paratropeza), Teucholabis (Euparatropesa)	4.158
jaenickeana Alexander, 1929, Tipula (Eumicrotipula)	4.26
jaffueli Alexander, 1929, Erioptera (Eriopterella)	4.123
jaffueli Alexander, 1928, Shannonomyia (Shannonomyia)	4.115
jaliscana Alexander, 1946, Teucholabis (Teucholabis)	4.162
jamaicae Alexander, 1939, Limonia (Geranomyia)	4.66
jamaicana Alexander, 1964, Gonomyia (Gonomyia)	4.133
jamaicensis Alexander, 1926, Limonia (Neolimonia)	4.71
jamaicensis Alexander, 1928, Tipula (Yamatotipula)	4.40
jamaicensis Alexander, 1964, Toxorhina (Toxorhina)	4.168
jejuna Alexander, 1916, Gonomyia (Gonomyia)	4.133
jesseana Alexander, 1945, Ozodicera (Dihexaclonus)	4.18
jivaro Alexander, 1944, Teucholabis (Teucholabis)	4.162
jivaro Alexander, 1916, Tipula (Eumicrotipula)	4.26
Jivaromyia Alexander, 1943	4.143
jivaronis Alexander, 1943, Tipula (Microtipula)	4.35
joana Alexander, 1929, Austrolimnophila (Austrolimnophila)	4.90
jocosa Alexander, 1913, Teucholabis (Teucholabis)	4.162
jocularis Alexander, 1948, Hexatoma (Eriocera)	4.102
jordanensis Alexander, 1949, Tipula (Microtipula)	4.35
jordanica Alexander, 1949, Limnophila (Limnophila)	4.108
jorgenseni (Alexander), 1919 (Dicranomyia), Limonia (Dicranomyia)	4.54
jovialis Alexander, 1945, Cryptolabis (Cryptolabis)	4.121
juarezi Alexander, 1946, Gonomyia (Gonomyia)	4.133
jubilans Alexander, 1940, Tipula (Eumicrotipula)	4.26
jubilata Alexander, 1943, Cryptolabis (Cryptolabis)	4.121
jubilata Alexander, 1938, Limonia (Rhipidia)	4.76
jucunda Alexander, 1913, Teucholabis (Teucholabis)	4.162
jujuyensis (Alexander), 1920 (Holorusia), Ischnotoma (Icriomastax) ..	4.9
jujuyensis (Alexander), 1924 (Dicranomyia), Limonia (Dicranomyia)	4.54
juliana Alexander, 1937, Hexatoma (Eriocera)	4.102
junincensis Alexander, 1942, Limonia (Rhipidia)	4.76
juniniana (Alexander), 1951 (Gnophomyia), Eugnophomyia	4.127
juquiana Alexander, 1945, Gonomyia (Lipophleps)	4.136
juquicola Alexander, 1945, Epiphragma (Epiphragma)	4.97



juquiella Alexander, 1945, Tipula (Microtipula)	4.35
juquiensis Alexander, 1945, Elephantomyia (Elephantomyia)	4.94
jurata Alexander, 1936, Hexatoma (Eriocera)	4.102
jurator Alexander, 1948, Epiphragma (Epiphragma)	4.97
justa Alexander, 1938, Gnophomyia	4.129
justa Alexander, 1937, Shannonomyia (Shannonomyia)	4.115
justoides Alexander, 1938, Gnophomyia	4.129
juturna (Alexander), 1945 (Holorusia), Zelandotipula	4.42
juventa Alexander, 1937, Tipula (Eumicrotipula)	4.26
kadeni (Alexander), 1941 (Tanypremna), Leptotarsus (Tanypremna)	4.14
kaieturana Alexander, 1930, Limnophila (Limnophila)	4.108
kaicturensis (Alexander), 1914 (Eriocera), Hexatoma (Eriocera) .	4.102
karma Alexander, 1945, Brachypremna	4.6
kathema Alexander, 1965, Tipula (Eumicrotipula)	4.26
kerteszi Alexander, 1914, Limnophila (Limnophila)	4.107
kerteszi Alexander, 1929, Gnophomyia	4.129
klagesi Alexander, 1945, Tipula (Microtipula)	4.35
klausma Alexander, 1962, Tipula (Eumicrotipula)	4.25
knabi Alexander, 1913, Erioptera (Mesocyphona)	4.125
knabi (Alexander), 1912 (Furcomyia), Limonia (Dicranomyia) ..	4.54
knabiana (Alexander), 1916 (Geranomyia), Limonia (Geranomyia)	4.63
kraussi Alexander, 1946, Tipula (Trichotipula)	4.40
kuehlhorni Alexander, 1962, Tipula (Eumicrotipula)	4.26
kuscheli Alexander, 1952, Shannonomyia (Shannonomyia)	4.115
kuscheli Alexander, 1967, Tipula (Eumicrotipula)	4.23
kuscheliana Alexander, 1952, Limonia (Dicranomyia)	4.54
labecula Alexander, 1942, Limonia (Dicranomyia)	4.54
lachesis Alexander, 1942, Limonia (Neolimonia)	4.71
?Lachnocera Philippi, 1865	4.88
lachrymalis (Alexander), 1916 (Geranomyia), Limonia (Geranomyia)	4.63
lachrymosa Alexander, 1919, Gnophomyia	4.129
laetitarsis (Alexander), 1922 (Geranomyia), Limonia (Geranomyia)	4.63
laddeyana Alexander, 1944, Atarba (Atarba)	4.86
laddeyi Alexander, 1943, Cryptolabis (Cryptolabis)	4.121
laddeyi Alexander, 1942, Hexatoma (Eriocera)	4.102
laeta Alexander, 1913, Teucholabis (Teucholabis)	4.162
lactabilis Alexander, 1952, Teucholabis (Euparatropesa)	4.158
laetifera Alexander, 1948, Teucholabis (Euparatropesa)	4.158
laetipleura Alexander, 1938, Erioptera (Erioptera)	4.123
laetitarsis Alexander, 1937, Limonia (Rhipidia)	4.76
laetiventris Alexander, 1945, Brachypremna	4.6
laevis (Alexander), 1914 (Holorusia), Zelandotipula	4.42
lagotis Alexander, 1942, Tipula (Microtipula)	4.35
laidis Alexander, 1966, Teucholabis (Teucholabis)	4.162
lais Alexander, 1950, Limonia (Rhipidia)	4.76
lais Alexander, 1946, Teucholabis (Teucholabis)	4.162
laistes Alexander, 1967, Limonia (Dicranomyia)	4.54
laneana Alexander, 1959, Teucholabis (Paratropesa)	4.158



lanei Alexander, 1945, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
lanci Alexander, 1942, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
lanei Alexander, 1942, <i>Ozodicera</i> (<i>Dihexaclonus</i>)	4.18
languidula Alexander, 1942, <i>Tipula</i> (<i>Microtipula</i>)	4.36
lanigera Alexander, 1928, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
lankesteri Alexander, 1946, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
lapazensis Alexander, 1962, <i>Limonia</i> (<i>Dicranomyia</i>)	4.54
larga Alexander, 1946, <i>Tipula</i> (<i>Bellardina</i>)	4.20
larotypa Alexander, 1926, <i>Helius</i>	4.46
larotypa (Alexander), 1929 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Ischnotoma</i>)	4.9
lassula Alexander, 1940, <i>Limonia</i> (<i>Rhipidia</i>)	4.75
lassula (Alexander), 1940 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
lata Alexander, 1943, <i>Gnophomyia</i>	4.129
latamera Alexander, 1968, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
lataurata Alexander, 1937, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104
latebra Alexander, 1967, <i>Limonia</i> (<i>Dicranomyia</i>)	4.54
lateralis Alexander, 1938, <i>Helius</i>	4.46
lateropolita Alexander, 1938, <i>Nephrotoma</i>	4.17
laterosetosa Alexander, 1931, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
laterospina Alexander, 1962, <i>Atarba</i> (<i>Atarba</i>)	4.86
laterospinosa Alexander, 1937, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162
laterospinosus Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
lathraea (Alexander), 1926 (<i>Pilaria</i>), <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
latibasalis Alexander, 1968, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162
latibasis Alexander, 1967, <i>Limonia</i> (<i>Idioglochina</i>)	4.68
laticincta Alexander, 1919, <i>Gnophomyia</i>	4.129
laticostata Alexander, 1940, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
laticostata Alexander, 1937, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
laticostata Alexander, 1937, <i>Tipula</i> (<i>Microtipula</i>)	4.36
latifascia (Alexander), 1926 (<i>Gnophomyia</i>), <i>Neognophomyia</i>	4.153
latifolia Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
latilobata Alexander, 1949, <i>Gnophomyia</i>	4.129
latilutea Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
latispina Alexander, 1942, <i>Limonia</i> (<i>Dicranomyia</i>)	4.54
latistyla Alexander, 1926, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
latitudinis Alexander, 1954, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
latissima Alexander, 1966, <i>Gnophomyia</i>	4.129
latissima (Alexander), 1922 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
laudabilis Alexander, 1942, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
laudanda Alexander, 1938, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
lauri Alexander, 1945, <i>Molophilus</i> (<i>Molophilus</i>)	4.147
lauta Alexander, 1949, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162
lawlori Alexander, 1934, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
laxa Alexander, 1946, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.162
Lecteria Osten Sacken, 1887	4.105, 4.106
leda Alexander, 1945, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
legata Alexander, 1948, <i>Lecteria</i> (<i>Psaronius</i>)	4.106
legitima (Alexander), 1928 (<i>Microtipula</i>), <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
<i>Leiponeura</i> Skuse, 1890	4.135
lemniscata Alexander, 1931, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
lemniscata Alexander, 1929, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
lenta Alexander, 1941, <i>Tipula</i> (<i>Microtipula</i>)	4.36
lemniscatus (Alexander), 1929 (<i>Habromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i>	



gurio)	4.12
lenitatis Alexander, 1946, Shannonomyia (Shannonomyia)	4.115
lentina Alexander, 1928, Shannonomyia (Shannonomyia)	4.115
lentoides (Alexander), 1913 (Limnophila), Shannonomyia (Shannonomyia)	4.115
leonardi Alexander, 1932, Shannonomyia (Shannonomyia)	4.115
leonensis Alexander, 1946, Erioptera (Mesocyphona)	4.125
leonora Alexander, 1938, Teucholabis (Teucholabis)	4.162
leonura Alexander, 1941, Gonomyia (Lipophleps)	4.136
leonurus Alexander, 1951, Molophilus (Molophilus)	4.147
Leptotarsus Guérin-Méneville, 1838	4.10, 4.11
leridensis Alexander, 1934, Teucholabis (Teucholabis)	4.162
lerionis Alexander, 1945, Molophilus (Molophilus)	4.147
lessepsi (Osten Saeken), 1886 (Eriocera), Hexatoma (Eriocera) ..	4.102
letalii Alexander, 1936, Tipula (Microtipula)	4.36
lethaeus Alexander, 1952, Molophilus (Molophilus)	4.147
lethe Alexander, 1946, Teucholabis (Teucholabis)	4.162
leucomelanopus (Enderlein), 1912 (Aporosa), Toxorhina (Ceratoheillus)	4.167
leucopasta Alexander, 1927, Erioptera (Mesocyphona)	4.125
leucopeza Alexander, 1929, Gynoplistia (Gynoplistia)	4.99
leucopeza Alexander, 1940, Polymera (Polymera)	4.112
leueoplaea (Alexander), 1921 (Gnophomyia), Eugnophomyia	4.127
leucoplaca Alexander, 1936, Helius	4.46
leucoplaca (Alexander), 1924 (Tanyremna), Leptotarsus (Tanyremnodes)	4.16
leueopoda Alexander, 1953, Dieranomyia	4.45
leueoseelis Alexander, 1950, Limonia (Neoglochina)	4.69
leucostena Alexander, 1937, Toxorhina (Ceratoheillus)	4.167
leucostigma Alexander, 1937, Limnophila (Limnophila)	4.107
leucostropha Alexander, 1966, Polymera (Polymerodes)	4.113
leueoxena (Alexander), 1915 (Mongoma), Trentepohlia (Paramongoma)	4.170
lewisii Alexander, 1964, Limonia (Dieranomyia)	4.54
liberta (Osten Saeken), 1859 (Dieranomyia), Limonia (Dieranomyia)	4.54
liehnophora Alexander, 1962, Limonia (Rhipidia)	4.76
liehyana Alexander, 1945, Tipula (Microtipula)	4.36
liehyella Alexander, 1947, Limonia (Peripheroptera)	4.73
liehyi Alexander, 1943, Limonia (Geranomyia)	4.63
lietor Alexander, 1938, Molophilus (Molophilus)	4.147
ligulata Alexander, 1929, Tipula (Eumicrotipula)	4.26
ligulipeniellata Alexander, 1946, Tipula (Eumicrotipula)	4.26
limai (Alexander), 1945 (Tanyremna), Leptotarsus (Tanyremna)	4.15
limbinervis Alexander, 1941, Limonia (Neoglochina)	4.69
Limnobia Meigen, 1818	4.47
Limnobia Westwood, 1835	4.60
Limnophila Maequart, 1834	4.106, 4.107
Limnophilella Alexander, 1919	4.92
Limonia Meigen, 1803	4.47
Limoniinae	4.44
Limoniini	4.44
Limoniodes Alexander, 1938	4.11
lindneri Alexander, 1933, Teucholabis (Euparatropesa)	4.158



<i>lincata</i> (Enderlein), 1912 (Geranomyia), Limonia (Geranomyia) .	4.63
<i>lincicollis</i> (Blanchard), 1852 (Limnobia), Limonia (Dicranomyia)	4.54
<i>lineipleura</i> Alexander, 1949, Teucholabis (Teucholabis)	4.162
<i>liponcura</i> Alexander, 1937, Teucholabis (Teucholabis)	4.162
<i>Lipophleps</i> Bergroth, 1915	4.135
<i>lipophleps</i> Alexander, 1937, Teucholabis (Teucholabis)	4.162
<i>Lipsothrix</i> Loew, 1873	4.143
<i>lissomelania</i> Alexander, 1967, Limonia (Peripheroptera)	4.73
<i>litigiosa</i> Alexander, 1928, Limnophila (Limnophila)	4.107
<i>liturata</i> (Enderlein), 1912 (Psaronius), Lacteria (Psaronius)	4.106
<i>livida</i> (Say), 1829 (Limnobia), Limonia (Dicranomyia)	4.54
<i>lloydi</i> (Alexander), 1913 (Oromyia), Atarba (Ischnothrix)	4.88
<i>lloydi</i> Alexander, 1913, Limnophila (Limnophila)	4.107
<i>lobuliferus</i> Alexander, 1938, Helius	4.46
<i>longibasis</i> Alexander, 1946, Tipula (Eumicrotipula)	4.26
<i>longicollis</i> Bigot, 1888, Limonia (Zelandoglochina)	4.80
<i>longicollis</i> Pierre, 1924, Toxorhina (Toxorhina)	4.168
<i>longifurcata</i> (Alexander), 1940 (Erioptera), Cheilotruchia (Empeda)	4.119
<i>longifusa</i> Alexander, 1944, Ctenolimnophila (Abitagua)	4.93
<i>longifusa</i> (Alexander), 1913 (Mongoma), Trentepohlia (Par-	
mongoma)	4.170
<i>longimana</i> (Fabricius), 1805 (Tipula), Ozodicera (Ozodicera)	4.19
<i>longipennis</i> (Alexander), 1923 (Eriocera), Hexatoma (Eriocera) .	4.102
<i>longipennis</i> (Macquart), 1838 (Tipula), Megistocera	4.16
<i>longipes</i> (Philippi), 1865 (Erioptera?), Amphineurus (Amphi-	
neurus)	4.117
<i>longipes</i> (Fabricius), 1805 (Tipula), Leptotarsus (Tanypremna) .	4.14
<i>longiradialis</i> Alexander, 1939, Cryptolabis (Cryptolabis)	4.121
<i>longiradialis</i> Alexander, 1929, Shannonomyia (Shannonomyia) ...	4.115
<i>longirostris</i> Williston, 1896, Elephantomyia (Elephantomyia) ...	4.94
<i>longisector</i> Alexander, 1944, Ozodicera (Dihexaclonus)	4.18
<i>longisetosa</i> Alexander, 1941, Teucholabis (Teucholabis)	4.162
<i>longispina</i> (Alexander), 1922 (Rhipidia), Limonia (Rhipidia)	4.76
<i>longispina</i> Alexander, 1938, Teucholabis (Teucholabis)	4.162
<i>longisquama</i> (Alexander), 1937 (Erioptera), Cheilotruchia (Empeda)	4.120
<i>longissimus</i> (Enderlein), 1912 (Stegasmonotus), Leptotarsus (Ta-	
nypremna)	4.14
<i>longistyla</i> (Alexander), 1914 (Eriocera), Hexatoma (Eriocera) ...	4.102
<i>longitarsis</i> (Macquart), 1846 (Tipula), Zelandotipula	4.42
<i>longiterebra</i> Alexander, 1943, Gnophomyia	4.129
<i>longiterebrata</i> Alexander, 1938, Rhabdomastix (Rhabdomastix) ..	4.155
<i>longitergata</i> Alexander, 1945, Atarba (Atarba)	4.86
<i>longitergata</i> Alexander, 1943, Gnophomyia	4.129
<i>longiventris</i> (Alexander), 1913 (Dicranomyia), Limonia (Dicra-	
nomyia)	4.54
<i>Longurio</i> Loew, 1869	4.11
<i>longurio</i> Alexander, 1938, Limonia (Rhipidia)	4.76
<i>longurioides</i> Alexander, 1954, Tipula (Eumicrotipula)	4.26
<i>lopesi</i> Alexander, 1942, Hexatoma (Eriocera)	4.102
<i>luctifica</i> Alexander, 1940, Tipula (Microtipula)	4.36
<i>luctuosa</i> (Osten Sacken), 1859 (Gnophomyia), Eugnophomyia ..	4.127
<i>ludibunda</i> Alexander, 1927, Limonia (Neolimonia)	4.71
<i>ludicra</i> Alexander, 1949, Teucholabis (Teucholabis)	4.162



<i>ludoviciana</i> Alexander, 1919, <i>Tipula</i> (<i>Yamatotipula</i>)	4.40
<i>lugubris</i> Alexander, 1914, <i>Tcueholabis</i> (<i>Teueholabis</i>)	4.162
<i>Lunatipula</i> Edwards, 1931	4.32
<i>lunigera</i> Alexander, 1929, <i>Erioptera</i> (<i>Erioptera</i>)	4.123
<i>luquilloensis</i> Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>lustralis</i> Alexander, 1945, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>lustralis</i> (Alexander), 1936 (<i>Tipula</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
<i>lustralis</i> Alexander, 1938, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>lutea</i> (Philippi), 1865 (<i>Polymoria</i>), <i>Mesolimnophila</i>	4.109
<i>luteiannulata</i> Alexander, 1939, <i>Elephantomyia</i> (<i>Elephantomyia</i>) .	4.94
<i>luteiangularis</i> Alexander, 1962, <i>Limonia</i> (<i>Dicranomyia</i>)	4.54
<i>luteiceps</i> Alexander, 1927, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>luteicolor</i> Alexander, 1968, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
<i>luteicolor</i> Alexander, 1942, <i>Teueholabis</i> (<i>Teueholabis</i>)	4.163
<i>luteicosta</i> Alexander, 1934, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>luteilimbata</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.36
<i>luteimana</i> Alexander, 1938, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>luteipennis</i> (Osten Saeken), 1859 (<i>Limnophila</i>), <i>Pseudolimnophila</i>	4.113
<i>luteistigma</i> (Alexander), 1940 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
<i>luteitarsis</i> Alexander, 1964, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
<i>lutcivena</i> (Alexander), 1946 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
<i>luteola</i> Alexander, 1946, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>luteola</i> Alexander, 1944, <i>Rhabdomastix</i> (<i>Rhabdomastix</i>)	4.155
<i>lutzi</i> (Alexander), 1912 (<i>Fureomyia</i>), <i>Limonia</i> (<i>Neoglochina</i>) ...	4.69
<i>luxuriosa</i> Alexander, 1929, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>luxuriosus</i> Alexander, 1938, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
<i>lycaon</i> Alexander, 1952, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>lydia</i> Alexander, 1950, <i>Limonia</i> (<i>Dicranomyia</i>)	4.54
<i>lyra</i> (Alexander), 1952 (<i>Limnophila</i>), <i>Chiliclimnophila</i>	4.93
<i>lyriformis</i> Alexander, 1966, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>lyriformis</i> Alexander, 1942, <i>Tipula</i> (<i>Microtipula</i>)	4.36
<i>machaeria</i> Alexander, 1921, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>machupichuana</i> Alexander, 1953, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
<i>macintyreana</i> Alexander, 1941, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
<i>macintyreii</i> Alexander, 1937, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>macintyreii</i> Alexander, 1938, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
<i>macquarti</i> (Enderlein), 1912 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ...	4.102
<i>macracantha</i> Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>macracantha</i> Alexander, 1952, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
<i>macracantha</i> Alexander, 1940, <i>Ozodieera</i> (<i>Dihexaelonus</i>)	4.18
<i>macrauehenia</i> Alexander, 1956, <i>Limonia</i> (<i>Geranomyia</i>)	4.63
<i>macrocera</i> (Alexander), 1914 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.102
<i>macroptera</i> (Philippi), 1865 (<i>Idioneura</i>), <i>Erioptera</i> (<i>Symplecta</i>) .	4.126
<i>maeropyga</i> Alexander, 1953, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
<i>macrorhyneha</i> Alexander, 1954, <i>Toxorhina</i> (<i>Ceratocheilus</i>)	4.167
<i>macrosterna</i> (Alexander), 1912 (<i>Pachyrhina</i>), <i>Tipula</i> (<i>Microtipula</i>)	4.36
<i>macrotrichiata</i> (Alexander), 1922 (<i>Microtipula</i>), <i>Tipula</i> (<i>Eumicrotipula</i>)	4.167
<i>maeswaini</i> Alexander, 1940, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.26
<i>macta</i> Alexander, 1945, <i>Limonia</i> (<i>Geranomyia</i>)	4.136
<i>maculata</i> (Fabricius), 1805 (<i>Tipula</i>), <i>Epiphragma</i> (<i>Epiphragma</i>) .	4.63
<i>maulipennis</i> Alexander, 1936, <i>Toxorhina</i> (<i>Ceratocheilus</i>)	4.96
	4.167



madida Alexander, 1928, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
maestia Alexander, 1921, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
maestitia Alexander, 1914, <i>Gnophomyia</i>	4.129
magellanica Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
<i>magellanicola</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
magellanicus Alexander, 1968, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
magiea Alexander, 1945, <i>Gnophomyia</i>	4.129
magistra Alexander, 1942, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
magistrata Alexander, 1949, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
magniareuata Alexander, 1949, <i>Gnophomyia</i>	4.129
magnificia (Alexander), 1914 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.102
magnificia (Alexander), 1913 (<i>Gnophomyia</i>), <i>Sigmatomera</i> (<i>Austro-</i> <i>limnobia</i>)	4.156
majae (Alexander), 1929 (<i>Astelobia</i>), <i>Sigmatomera</i> (<i>Austrolimno-</i> <i>bia</i>)	4.156
Maietta Alexander, 1929	4.143
majuseula Alexander, 1940, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
maldonadoi Alexander, 1952, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
maldonadoi (Alexander), 1953 (<i>Longurio</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premnella</i>)	4.15
maligna Alexander, 1945, <i>Limonia</i> (<i>Dieranomyia</i>)	4.54
malitiosa Alexander, 1942, <i>Limonia</i> (<i>Dieranomyia</i>)	4.54
manabiana Alexander, 1942, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
manabiana Alexander, 1953, <i>Limonia</i> (<i>Geranomyia</i>)	4.61
manea (Alexander), 1921 (<i>Psaronius</i>), <i>Leetertia</i> (<i>Psaronius</i>)	4.106
manea (Williston), 1896 (<i>Mongoma</i>), <i>Trentepohlia</i> (<i>Paramongoma</i>) ..	4.170
mandator Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.36
manicata Alexander, 1929, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
manieatus (Osten Sacken), 1887 (<i>Tanypremnella</i>), <i>Leptotarsus</i> (<i>Ta-</i> <i>nypremnella</i>)	4.14
mannheimsi Alexander, 1960, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.90
manniana Alexander, 1946, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
manniana Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.36
mantissa Alexander, 1966, <i>Austrolimnophila</i> (<i>Limnophilella</i>)	4.92
margarita Alexander, 1965, <i>Ctenophora</i> (<i>Pselliophora</i>)	4.7
margaritae Alexander, 1954, <i>Hexatoma</i> (<i>Eriocera</i>)	4.102
marmoripennis Rondani, 1850, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
marthae Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
marthae Alexander, 1931, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
martieola Alexander, 1931, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
martinbrowni Alexander, 1946, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
martinezi Alexander, 1958, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.90
martinezi Alexander, 1962, <i>Teucholabis</i> (<i>Eupararopesa</i>)	4.158
masafuerae Alexander, 1952, <i>Limonia</i> (<i>Dieranomyia</i>)	4.55
masafuerae Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
masatierrae Alexander, 1952, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.115
mattogrossae Alexander, 1913, <i>Leetertia</i> (<i>Leetertia</i>)	4.106
maya Alexander, 1927, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
maya (Alexander), 1912 (<i>Tipula</i>), <i>Ischnotoma</i> (<i>Neotipula</i>)	4.10
mayanymphica (Alexander), 1946 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>)	4.120
mecoglossa Alexander, 1967, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
meconeura Alexander, 1962, <i>Limonia</i> (<i>Caenolimonia</i>)	4.49



meeophallus Alexander, 1966, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
<i>mediatlantica</i> Freeman, 1962, <i>Limonia</i> (<i>Dieranomyia</i>)	4.52
mediocompressa Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.36
medioicornis (Alexander), 1944 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premnella</i>)	4.15
mediodontata Alexander, 1944, <i>Tipula</i> (<i>Eumierotipula</i>)	4.27
medioligula Alexander, 1945, <i>Nephrotoma</i>	4.17
<i>Mediophragma</i> Alexander, 1954	4.92
mediotubereolata Alexander, 1949, <i>Gnophomyia</i>	4.129
megaeera (Alexander), 1943 (<i>Tanypremnna</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premnella</i>)	4.15
megalops Alexander, 1942, <i>Pseudolimnophila</i>	4.113
megalopyga Alexander, 1967, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
megalyla Alexander, 1945, <i>Tipula</i> (<i>Mierotipula</i>)	4.36
megaphallus Alexander, 1921, <i>Atarba</i> (<i>Atarba</i>)	4.87
megaphallus Alexander, 1968, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>Megarhina</i> Lepeletier & Serville, 1828	4.45
megarhopala Alexander, 1946, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>Megistoeera</i> Wiedemann, 1828 (<i>Maekistoeera</i>)	4.16
<i>Megistomastix</i> Alexander, 1912	4.8
melampodia Alexander, 1940, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
melanaantha Alexander, 1954, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.136
melanaera (Wiedemann), 1828 (<i>Limnobia</i>), <i>Hexatoma</i> (<i>Erioera</i>)	4.102
melanaria Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.76
melaneholica (Alexander), 1922 (<i>Gnophomyia</i>), <i>Eugnophomyia</i> ..	4.127
melanocephala (Fabricius), 1794 (<i>Tipula</i>), <i>Teucholabis</i> (<i>Teucho-</i> <i>labis</i>)	4.163
melanoera (Alexander), 1925 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Diera-</i> <i>nomyia</i>)	4.55
melanoleucus Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
melanolitha (Alexander), 1927 (<i>Erioera</i>), <i>Hexatoma</i> (<i>Erioera</i>) ..	4.103
melanomera Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.87
melanonota Alexander, 1968, <i>Hexatoma</i> (<i>Erioera</i>)	4.103
melanophallus Alexander, 1938, <i>Helius</i>	4.46
melanophallus Alexander, 1968, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.160
melanosterna Alexander, 1945, <i>Polymera</i> (<i>Polymera</i>)	4.112
melanoxantha Alexander, 1962, <i>Nephrotoma</i>	4.17
melaxantha Alexander, 1943, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
melaxantha Alexander, 1934, <i>Limonia</i> (<i>Caenolimonia</i>)	4.49
melica Alexander, 1929, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
melina (Alexander), 1922 (<i>Erioera</i>), <i>Hexatoma</i> (<i>Erioera</i>)	4.103
mendax Alexander, 1920, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
mendica Alexander, 1929, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.114
mendoza Alexander, 1935, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
mephistophelica Alexander, 1947, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
meridensis Alexander, 1940, <i>Limonia</i> (<i>Caenolimonia</i>)	4.49
meridiana Edwards, 1919, <i>Tipula</i> (<i>Eumierotipula</i>)	4.27
meridieola Alexander, 1943, <i>Limonia</i> (<i>Dieranomyia</i>)	4.55
meridionalis Alexander, 1913, <i>Elephantomyia</i> (<i>Elephantomyia</i>) ..	4.94
meridionalis Alexander, 1913, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
merita Alexander, 1938, <i>Atarba</i> (<i>Atarba</i>)	4.87
merklei Alexander, 1928, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.90
mesamericana (Alexander), 1944 (<i>Pselliophora</i>), <i>Ctenophora</i>	



(Pselliophora)	4.7
mesocera Alexander, 1929, Atarba (Ischnothrix)	4.89
Mesocyphona Osten Sacken, 1869	4.123
Mesolimnophila Alexander, 1929	4.109
mesonotalis Alexander, 1949, Trentepohlia (Neomongoma)	4.169
mesophragma Alexander, 1928, Shannonomyia (Shannonomyia) ..	4.115
mesophragmoides Alexander, 1937, Shannonomyia (Shannonomyia)	4.115
mesotricha Alexander, 1944, Limonia (Neoglochina)	4.69
mesoxantha (Osten Sacken), 1886 (Eriocera), Hexatoma (Eriocera)	4.103
metae Alexander, 1938, Limonia (Neoglochina)	4.68
mctatarsata (Alexander), 1915 (Mongoma), Trentepohlia (Para-	
mongoma)	4.170
methodica Alexander, 1931, Gonomyia (Gonomyia)	4.133
metuendus Alexander, 1951, Molophilus (Molophilus)	4.148
mexicana Alexander, 1926, Atarba (Atarba)	4.87
mexicana Alexander, 1946, Dicranota (Rhaphidolabis)	4.84
mexicana Alexander, 1916, Gonomyia (Gonomyia)	4.133
mexicana (Bellardi), 1861 (Aporosa), Limonia (Geranomyia) ...	4.64
mexicana (Alexander), 1940 (Oxydiscus), Paradelphomyia (Oxy-	
rhiza)	4.109
mexicana Alexander, 1938, Rhabdomastix (Rhabdomastix)	4.155
mexicana (Macquart), 1846 (Pachyrhina), Nephrotoma	4.17
michaelseni Alexander, 1928, Austrolimnophila (Austrolimnophila)	4.90
michoacana Alexander, 1946, Tipula (Lunatipula)	4.32
micracantha Alexander, 1926, Gonomyia (Paralipophleps)	4.140
micracanthus Alexander, 1945, Helius	4.46
microcephala (Wulp), 1881 (Tipula), Zelandotipula	4.42
microcera (Alexander), 1943 (Tanypremna), Leptotarsus (Tany-	
premnella)	4.15
micromera Alexander, 1945, Gonomyia (Paralipophleps)	4.140
micromyia Alexander, 1920, Erioptera (Erioptera)	4.123
microphaea Alexander, 1939, Limonia (Geranomyia)	4.64
microphallus Alexander, 1944, Atarba (Atarba)	4.87
microptera Alexander, 1945, Erioptera (Symplecta)	4.126
microscola Alexander, 1967, Limonia (Dicranomyia)	4.55
microserrata Alexander, 1938, Gonomyia (Gonomyia)	4.133
microsoma Alexander, 1944, Limonia (Dicranomyia)	4.55
microsomoides Alexander, 1954, Limonia (Dicranomyia)	4.55
microspilota Alexander, 1943, Austrolimnophila (Austrolimnophila)	4.90
microspilota Alexander, 1928, Tipula (Eumicrotipula)	4.27
microsticta Alexander, 1929, Austrolimnophila (Austrolimnophila)	4.90
microsticta (Alexander), 1921 (Rhipidia), Limonia (Rhipidia) ...	4.76
microstictula Alexander, 1929, Polymera (Polymera)	4.112
Microtipula Alexander, 1912	4.32
militaris Alexander, 1952, Limonia (Geranomyia)	4.64
miniata Alexander, 1929, Limonia (Zelandoglochina)	4.80
miniata Alexander, 1930, Teucholabis (Teucholabis)	4.163
minuta Alexander, 1926, Teucholabis (Teucholabis)	4.163
minutior Alexander, 1942, Polymera (Polymera)	4.112
minutipennis Alexander, 1929, Shannonomyia (Shannonomyia) ..	4.115
minutissima Alexander, 1945, Polymera (Polymerodes)	4.113
mirabilis (Alexander), 1914 (Rhamphidia), Helius	4.46
mirabilis (Osten Sacken), 1878 (Diotrepha), Orimarga (Diotrepha)	4.81
mirabilis Alexander, 1938, Trentepohlia (Promongoma)	4.171




miraculus Alexander, 1938, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
miranda (Alexander), 1921 (<i>Rhamphidia</i>), <i>Helius</i>	4.46
miranda (Alexander), 1944 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premna</i>)	4.14
miranha Alexander, 1913, <i>Tipula</i> (<i>Eumierotipula</i>)	4.27
misera Alexander, 1921, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
miscranda Alexander, 1945, <i>Limonia</i> (<i>Dieranomyia</i>)	4.55
mistura Alexander, 1942, <i>Limonia</i> (<i>Dieranomyia</i>)	4.55
mithras Alexander, 1949, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
mithrodatus Alexander, 1951, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
mitua Alexander, 1916, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
moeoa Alexander, 1913, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
moetezuma Alexander, 1928, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
moetzumae Alexander, 1925, <i>Tipula</i> (<i>Eumierotipula</i>)	4.27
modica Alexander, 1927, <i>Erioptera</i> (<i>Mesoeocyphona</i>)	4.125
molcsta Osten Sacken, 1886, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
molinae Alexander, 1929, <i>Gnophomyia</i>	4.129
molophiloides Alexander, 1943, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>Molophilus</i> Curtis, 1833	4.143, 4.144
monacantha Alexander, 1943, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
monacantha Alexander, 1937, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
Mongoma Westwood, 1881	4.168
<i>Mongomella</i> Enderlein, 1912	4.169
monilifera Loew, 1851, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
moniliferoides Alexander, 1920, <i>Tipula</i> (<i>Eumierotipula</i>)	4.27
moniliformis Röder, 1886, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
moniliger Alexander, 1967, <i>Limonia</i> (<i>Neoglochina</i>)	4.69
monnula Alexander, 1962, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
monoecra Alexander, 1967, <i>Tipula</i> (<i>Mierotipula</i>)	4.36
monoetenus Alexander, 1951, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
monophaea Alexander, 1946, <i>Gnophomyia</i>	4.129
monophora Alexander, 1941, <i>Neognophomyia</i>	4.153
<i>Monorhipidia</i> Alexander, 1912	4.74
monosticta Alexander, 1948, <i>Polymera</i> (<i>Polymera</i>)	4.112
monostictula (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
monostyloides Alexander, 1951, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
monostylus Alexander, 1928, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
monoxantha Alexander, 1944, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
monticola Alexander, 1917, <i>Antocha</i>	4.45
montivaga Alexander, 1949, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
mordax Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
mordax Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
morelosensis (Alexander), 1946 (<i>Oxydiscus</i>), <i>Paradelphomyia</i> (<i>Oxyrhiza</i>)	4.109
morgana Alexander, 1944, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
morionella (Shiner), 1868 (<i>Limnobia</i>), <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
morphea Alexander, 1946, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
mulfordi Alexander, 1945, <i>Tipula</i> (<i>Mierotipula</i>)	4.36
muliercula Alexander, 1942, <i>Limonia</i> (<i>Dieranomyia</i>)	4.55
mulsa (Alexander), 1916 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Dieranomyia</i>) .	4.55
multiannulata Alexander, 1937, <i>Erioptera</i> (<i>Erioptera</i>)	4.123
multiarmata Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.87
multiarmata Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80



<i>multidentata</i> Alexander, 1931, <i>Aphrophila</i>	4.118
<i>multiermis</i> Alexander, 1942, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.19
<i>multifida</i> (Alexander), 1926 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>multifidus</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
<i>multiguttata</i> (Alexander), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ..	4.77
<i>multiguttula</i> Alexander, 1939, <i>Hexatoma</i> (<i>Eriocera</i>)	4.103
<i>multimoda</i> Alexander, 1938, <i>Tipula</i> (<i>Microtipula</i>)	4.36
<i>multinodosa</i> Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
<i>multipecta</i> Alexander, 1939, <i>Austrolimnophila</i> (<i>Limnophilella</i>)	4.92
<i>multipectata</i> (Fabricius), 1805 (<i>Tipula</i>), <i>Limonia</i>	4.81
<i>multipectata</i> Alexander, 1938, <i>Orimarga</i> (<i>Orimarga</i>)	4.82
<i>multipectigera</i> Alexander, 1945, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>multiramosa</i> Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>multisetosus</i> Alexander, 1936, <i>Molophilus</i> (<i>Trichomolophilus</i>)	4.152
<i>multisignata</i> Alexander, 1936, <i>Limonia</i> (<i>Neoglochina</i>)	4.69
<i>multispicata</i> Alexander, 1938, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>multispinosa</i> Alexander, 1944, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>multivolus</i> Alexander, 1945, <i>Helius</i>	4.46
<i>munda</i> Alexander, 1913, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>musciicola</i> Alexander, 1943, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>muscosa</i> (Enderlein), 1912 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Neolimnobia</i>)	4.70
<i>mutata</i> Alexander, 1935, <i>Limonia</i> (<i>Dicranomyia</i>)	4.55
<i>muzocensis</i> Alexander, 1938, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>myctera</i> Alexander, 1967, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
<i>myersi</i> Alexander, 1926, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>myersiana</i> Alexander, 1929, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
<i>myersiana</i> (Alexander), 1930 (<i>Diotrepha</i>), <i>Orimarga</i> (<i>Diotrepha</i>) .	4.81
<i>myersiana</i> Alexander, 1931, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
<i>myersiellus</i> Alexander, 1935, <i>Helius</i>	4.46
<i>myriatricha</i> Alexander, 1942, <i>Tipula</i> (<i>Microtipula</i>)	4.36
<i>myriosticta</i> Alexander, 1941, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>myrtea</i> (Alexander), 1922 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.103
<i>mystica</i> Alexander, 1931, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
<i>mystica</i> Alexander, 1945, <i>Styringomyia</i>	4.157
<i>mythica</i> Alexander, 1945, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
<i>mythica</i> Alexander, 1926, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>nacrea</i> (Alexander), 1913 (<i>Limnophila</i>), <i>Shannonomyia</i> (<i>Shanno-</i>	
<i>nomyia</i>)	4.116
<i>nahuelbutae</i> Alexander, 1966, <i>Gnophomyia</i>	4.129
<i>nahuelbutae</i> (Alexander), 1945 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Lon-</i>	
<i>gurio</i>)	4.12
<i>nahuelbutae</i> Alexander, 1966, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
<i>nahuclicola</i> Alexander, 1958, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.90
<i>naiquatana</i> Alexander, 1940, <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
<i>napoensis</i> (Alexander), 1921 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Caenoglo-</i>	
<i>china</i>)	4.48
<i>napoensis</i> Alexander, 1921, <i>Tipula</i> (<i>Microtipula</i>)	4.33
<i>navarinoensis</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.27
<i>neanthina</i> Alexander, 1954, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
<i>nearcuata</i> Alexander, 1940, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
<i>nebulicincta</i> Alexander, 1941, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
<i>nebulipennis</i> Alexander, 1928, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.163
<i>nebulosa</i> (Bellardi), 1861 (<i>Tipula</i>), <i>Epiphragma</i> (<i>Epiphragma</i>) ..	4.97



neetarea Alexander, 1944, Gnophomyia	4.129
nefasta Alexander, 1944, Limonia (Dieranomyia)	4.55
neglecta Alexander, 1936, Limonia (Rhipidia)	4.77
neivai Alexander, 1940, Ozodicera (Dihexaelonus)	4.18
neivai Alexander, 1940, Tipula (Eumicrotipula)	4.27
nematocera Alexander, 1939, Limnophila (Limnophila)	4.108
nemorivaga Alexander, 1929, Limnophila (Limnophila)	4.107
neobifida Alexander, 1952, Gonomyia (Paralipophleps)	4.140
neoclausula Alexander, 1967, Polymera (Polymera)	4.112
neocollaris Alexander, 1944, Teucholabis (Paratropesa)	4.159
neodiceros Alexander, 1966, Molophilus (Molophilus)	4.148
neofacinus Alexander, 1964, Molophilus (Molophilus)	4.148
neofaleifera Alexander, 1943, Gonomyia (Neolipophleps)	4.139
neogama Alexander, 1943, Trentepohlia (Paramongoma)	4.170
neogaudens Alexander, 1942, Limonia (Geranomyia)	4.64
neogaudens Alexander, 1945, Orimarga (Orimarga)	4.82
Neoglochina Alexander, 1967	4.68
Neognophomyia Alexander, 1926	4.152
neolenta Alexander, 1945, Tipula (Microtipula)	4.36
neoleridensis Alexander, 1940, Teucholabis (Teucholabis)	4.163
Neolimnobia Alexander, 1929	4.69
Neolimnophila Alexander, 1965	4.154
neolimnophiloides Alexander, 1942. Ctenolimnophila (Campbellomyia)	4.93
Neolimonia Alexander, 1964	4.70
Neolipophleps Alexander, 1947	4.139
Neomongoma Alexander, 1919	4.169
neonumenius Alexander, 1930, Limonia (Geranomyia)	4.64
neoparilis Alexander, 1962, Limonia (Geranomyia)	4.64
neopentheres Alexander, 1930, Limonia (Geranomyia)	4.64
Neophilippiana Alexander, 1964	4.154
neorepanda Alexander, 1964, Limonia (Caenolimonia)	4.49
neosaga Alexander, 1947, Hexatoma (Eriocera)	4.103
neosalva Alexander, 1943, Teucholabis (Teucholabis)	4.163
Neotipula Alexander, 1940	4.10
neotropica Alexander, 1940, Lipsothrix	4.143
nepenthe Alexander, 1946, Teucholabis (Euteucholabis)	4.158
nephele Alexander, 1948, Epiphragma (Epiphragma)	4.97
Nephrotoma Meigen, 1803	4.16
Nephrotomodes Alexander, 1946	4.32
neptunus Alexander, 1952, Molophilus (Molophilus)	4.148
nestor Alexander, 1947, Gonomyia (Lipophleps)	4.137
nethis Alexander, 1965, Tipula (Eumicrotipula)	4.27
neuquenensis Alexander, 1952, Austrolimnophila (Austrolimnophila)	4.91
neuquenensis Alexander, 1962, Valdiviana	4.41
neurotrichia (Alexander), 1962 (Holorusia), Zelandotipula	4.42
nicoya Alexander, 1944, Tipula (Microtipula)	4.36
nigra (Latreille), 1809 (Hexatoma), Hexatoma (Eriocera)	4.102
nigra (Wiedemann), 1828 (Limnobia), Hexatoma (Eriocera)	4.99
nigricola (Walker), 1861 (Limnobia), Eugnophomyia	4.127
nigrina (Wiedemann), 1828 (Limnobia), Gnophomyia	4.129
nigripleura (Alexander), 1919 (Geranomyia), Limonia (Geranomyia)	4.64
nigripleuralis Alexander, 1945, Epiphragma (Epiphragma)	4.97
nigriscapa Alexander, 1946, Tipula (Eumicrotipula)	4.27



nigrivena Alexander, 1939, Toxorhina (Toxorhina)	4.168
nigroabdominalis (Alexander), 1936 (Habromastix), Tipula (Micro- tipula)	4.36
nigroapicalis Alexander, 1941, Orimarga (Orimarga)	4.83
nigrochalybea (Alexander), 1922 (Eriocera), Hexatoma (Eriocera)	4.103
nigroclavaria Alexander, 1946, Teucholabis (Teucholabis)	4.163
nigrocorporis Alexander, 1937, Teucholabis (Teucholabis)	4.163
nigrocostata Alexander, 1939, Teucholabis (Teucholabis)	4.163
nigrocoxalis Alexander, 1936, Teucholabis (Paratropesa)	4.159
nigrofemorata Alexander, 1936, Braehypremna	4.6
nigrolincata (Enderlein), 1912 (Empeda), Cheilotrichia (Empeda)	4.120
nigrolutea (Bellardi), 1859 (Tipula), Nephrotoma	4.17
nigromarginata Alexander, 1938, Ozodicera (Ozodicera)	4.19
nigroplagiata Alexander, 1939, Epiphragma (Epiphragma)	4.97
nigropostica Alexander, 1938, Teucholabis (Teucholabis)	4.163
nigrostrata Alexander, 1938, Limonia (Rhipidia)	4.77
nigrosetosa (Alexander), 1945 (Holorusia), Zelandotipula	4.42
nigrosignata Alexander, 1931, Teucholabis (Teucholabis)	4.163
nigroterminalis Alexander, 1932, Teucholabis (Teucholabis)	4.161
nigrotincta Alexander, 1945, Limonia (Caenolimonia)	4.49
nigrovariegata Alexander, 1928, Tipula (Microtipula)	4.36
nimbifera Alexander, 1943, Gnophomyia	4.129
nimbinervis Alexander, 1946, Tipula (Eumicrotipula)	4.28
nimbipennis Alexander, 1945, Polymera (Polymera)	4.112
niobe Alexander, 1947, Tipula (Microtipula)	4.36
nitens (Schiner), 1868 (Peripheroptera), Limonia (Peripheroptera) ..	4.73
nitra (Alexander), 1945 (Holorusia), Isehnotoma (Icriomastax) ..	4.9
niveitarsis Alexander, 1915, Orimarga (Orimarga)	4.83
niveitarsis Alexander, 1913, Polymera (Polymera)	4.112
niveitarsis (Alexander), 1922 (Ceratocheilus), Toxorhina (Cerato- cheilus)	4.167
niveitarsis (Alexander), 1913 (Mongoma), Trentepohlia (Paramon- goma)	4.170
nobilissima Alexander, 1941, Limonia (Rhipidia)	4.77
noeivus (Alexander), 1953 (Longurio) Leptotarsus (Longurio) ...	4.12
noctivagans Alexander, 1914, Ozodicera (Ozodicera)	4.19
nocturna Alexander, 1941, Teucholabis (Teucholabis)	4.163
nodulifera Alexander, 1929, Limonia (Zelandoglochina)	4.80
nodulifera Alexander, 1940, Polymera (Polymera)	4.112
nodulifera Alexander, 1949, Teucholabis (Paratropesa)	4.159
nodulosa Alexander, 1939, Atarba (Atarba)	4.87
nolens Alexander, 1965, Tipula (Eumicrotipula)	4.28
nonnullus Alexander, 1966, Amphineurus (Rhamphoneurus)	4.118
nordenskjoldi Alexander, 1920, Tipula (Eumicrotipula)	4.28
nothofagctorum Alexander, 1929, Tipula (Eumicrotipula)	4.28
nothofagctorum Alexander, 1929, Amphineurus (Rhamphoneurus) ..	4.118
nothofagi Alexander, 1929, Limonia (Dicranomyia)	4.55
notoria Alexander, 1942, Tipula (Eumicrotipula)	4.28
novaleonensis Alexander, 1940, Tipula (Eumicrotipula)	4.28
novatrix Alexander, 1942, Tipula (Eumicrotipula)	4.28
nubifera Wulp, 1881, Tipula (Eumicrotipula)	4.28



nubilosa Alexander, 1967, Limonia (Rhipidia)	4.77
nudicornis (Macquart), 1838 (Tipula), Ischnotoma (Icriomastax) .	4.9
nudipennis Alexander, 1964, Shannonomyia (Shannonomyia) ...	4.116
nugatoria Alexander, 1943, Limonia (Geranomyia)	4.64
nullus Alexander, 1966, Amphineurus (Rhamphoneurus)	4.118
numenius (Alexander), 1913 (Geranomyia), Limonia (Geranomyia)	4.64
nympha Alexander, 1943, Austrolimnophila (Austrolimnophila) ...	4.91
nymphica (Alexander), 1928 (Erioptera), Cheilotrichia (Empeda) .	4.120
oaxacensis (Alexander), 1946 (Oxydiscus), Paradelphomyia (Oxyrhiza)	4.109
obediens Alexander, 1927, Molophilus (Molophilus)	4.148
obirata Alexander, 1940, Tipula (Eumicrotipula)	4.28
obliquefasciata Macquart, 1846, Tipula (Bellardina)	4.20
obliterata Alexander, 1913, Lecteria (Psaronius)	4.106
obliviosus Alexander, 1953, Molophilus (Molophilus)	4.148
obscura (Fabricius), 1805 (Tipula), Lecteria (Psaronius)	4.106
obscura Macquart, 1838, Polymera (Polymera)	4.112
obscurata (Blanchard), 1852 (Limnobia), Erioptera (Trimicra) ...	4.126
obscuricincta Alexander, 1940, Tipula (Eumicrotipula)	4.28
obsoleta (Williston), 1900 (Eriocera), Hexatoma (Eriocera)	4.103
obtusilamina Alexander, 1952, Neognophomyia	4.153
obtusiloba Alexander, 1944, Atarba (Ischnothrix)	4.89
obtusiloba Alexander, 1937, Dolichoceza (Megistomastix)	4.8
obtusirostris Alexander, 1962, Limonia (Dicranomyia)	4.54
obtusistyla Alexander, 1940, Gonomyia (Gonomyia)	4.134
obtusistylus Alexander, 1925, Limonia (Dicranomyia)	4.55
occellata Alexander, 1914, Sigmatomera (Sigmatomera)	4.156
ocellata Alexander, 1942, Limonia (Rhipidia)	4.77
ocellata (Enderlein), 1912 (Icriomastax), Ischnotoma (Icriomastax)	4.9
ocellata (Röder), 1886 (Limnobia), Limonia	4.81
ocellifera (Alexander), 1915 (Eriocera), Hexatoma (Eriocera) ...	4.103
ochracea Macquart, 1834, Ozodicera (Ozodicera)	4.19
ochricauda (Alexander), 1925 (Erioptera), Cheilotrichia (Empeda)	4.120
adontomera Alexander, 1967, Tipula (Eumicrotipula)	4.28
ofella Alexander, 1953, Limonia (Zelandoglochina)	4.80
ogloblini Alexander, 1935, Hexatoma (Eriocera)	4.103
ohausiana (Enderlein), 1912 (Eriocera), Hexatoma (Eriocera) ...	4.103
ohlini (Alexander), 1920 (Dicranomyia), Limonia (Dicranomyia)	4.55
oiticicai Alexander, 1948, Limnophila (Limnophila)	4.108
oligotrichus Alexander, 1954, Molophilus (Molophilus)	4.148
oliviae (Alexander), 1915 (Peripheroptera), Limonia (Peripherop- tera)	4.73
olssoni (Alexander), 1919 (Gnophomyia), Shannonomyia (Shanno- nomyia)	4.116
olssoniana Alexander, 1943, Tipula (Eumicrotipula)	4.28
ominosa Alexander, 1926, Gonomyia (Progonomyia)	4.141
ominosa Alexander, 1938, Polymera (Polymera)	4.112
omissa (Alexander), 1912 (Furcomyia), Limonia (Dicranomyia) ..	4.55
omissa Alexander, 1921, Teucholabis (Teucholabis)	4.163
omissinervis (Alexander), 1913 (Diotrepha), Orimarga (Diotrepha)	4.81
omissinervis Alexander, 1921, Teucholabis (Teucholabis)	4.164
omissistyla Alexander, 1929, Limonia (Zelandoglochina)	4.80
omissivena (Alexander), 1922 (Dicranomyia), Limonia (Dicrano-	



myia)	4.55
omissus Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
omnilutea Alexander, 1967, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
ona Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
oneris Alexander, 1956, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
onerosa (Alexander), 1928 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.55
onoma Alexander, 1950, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
opilio (Osten Sacken), 1886 (<i>Tanyremna</i>), <i>Leptotarsus</i> (<i>Tanyremna</i>)	4.14
opinator Alexander, 1950, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
opipara Alexander, 1951, <i>Tipula</i> (<i>Microtipula</i>)	4.36
optabilis Alexander, 1921, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
opulens Alexander, 1945, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
opulenta Alexander, 1951, <i>Hexatoma</i> (<i>Eriocera</i>)	4.103
ordinaria Alexander, 1945, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
ordinaria (Osten Sacken), 1886 (<i>Pachyrrhina</i>), <i>Nephrotoma</i>	4.17
<i>Oreomyza</i> Pokorny, 1887	4.40
<i>oreomyzoides</i> Alexander, 1926, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
oreonympha Alexander, 1928, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
oreonympha Alexander, 1929, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
oresitropha (Alexander), 1927 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>)	4.120
Orimarga Osten Sacken, 1869	4.81, 4.82
orion Alexander, 1914, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
orizabensis Alexander, 1941, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
ornaticornis Alexander, 1939, <i>Hexatoma</i> (<i>Eriocera</i>)	4.103
ornaticornis Wulp, 1891, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
ornatipennis (Blanchard), 1852 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.55
ornatipes (Alexander), 1941 (<i>Tanyremna</i>), <i>Leptotarsus</i> (<i>Tanyremna</i>)	4.15
oroensis Alexander, 1943, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.91
<i>Orolimnophila</i> Alexander, 1921	4.88
<i>Oromyia</i> Alexander, 1913	4.88
orophila (Alexander), 1913 (<i>Limnophila</i>), <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
orophila (Alexander), 1914 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.42
orthogonia Alexander, 1945, <i>Limonia</i> (<i>Caenolimonia</i>)	4.49
orthomera Alexander, 1937, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
orthomeroides Alexander, 1939, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
orthorhabda Alexander, 1940, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
ortoni Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.37
osculata Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.28
ostensackeni Skuse, 1890, <i>Gnophomyia</i>	4.129
ostentator Alexander, 1946, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
osterhouti (Alexander), 1912 (<i>Furcomyia</i>), <i>Limonia</i> (<i>Caenolimonia</i>)	4.49
oteroi Alexander, 1936, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
othello Alexander, 1941, <i>Molophilus</i> (<i>Molophilus</i>)	4.148
otiosa Alexander, 1949, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
ovaliformis Alexander, 1946, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
ovalis Alexander, 1949, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
<i>Orydiscus</i> Meijere, 1913	4.109
oxymera Alexander, 1943, <i>Gnophomyia</i>	4.130
oxyphallus Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>Oxyrhiza</i> Meijere, 1946	4.109
<i>Ozodicera</i> Macquart, 1834	4.17
ozotus Alexander, 1968, <i>Molophilus</i> (<i>Molophilus</i>)	4.148



pabulatoria Alexander, 1920, Teueholabis (Teucholabis)	4.164
pachyrhampha Alexander, 1967, Erioptera (Mesoeocyphona)	4.125
<i>Pachyrhina</i> Macquart, 1834	4.16
<i>Pachyrhina</i> , <i>Pachyrhina</i> , <i>Pachyrhyna</i> , error or emends.	4.16
pachyspila (Alexander), 1928 (Gynoplistia), Paralimnophila	4.110
pacifer Alexander, 1947, Molophilus (Molophilus)	4.148
pacifera Alexander, 1936, Austrolimnophila (Austrolimnophila) ..	4.91
paganus Alexander, 1942, Molophilus (Molophilus)	4.149
pala Alexander, 1941, Molophilus (Molophilus)	4.149
pala Alexander, 1939, Tipula (Microtipula)	4.37
palaeogama Alexander, 1944, Tipula (Microtipula)	4.37
palenca Alexander, 1944, Tipula (Eumicrotipula)	4.28
<i>Pales</i> Meigen, 1800	4.16
palitans Alexander, 1965, Tipula (Eumicrotipula)	4.28
pallatangae Alexander, 1929, Limonia (Rhipidia)	4.77
pallatangensis Alexander, 1929, Molophilus (Molophilus)	4.149
pallens (Philippi), 1865 (Limnophila)	4.117
<i>pallida</i> (Williston), 1896, Limonia (Geranomyia)	4.63
<i>pallida</i> (Williston), 1896 (Mongoma), Trentepohlia (Paramongoma)	4.170
<i>pallida</i> (Macquart), 1838 (Limnobia), Limonia	4.81
pallidapex Alexander, 1943, Atarba (Atarba)	4.87
pallidapex Alexander, 1929, Gnophomyia	4.130
pallidapex Alexander, 1945, Limonia (Geranomyia)	4.64
pallidibasis Alexander, 1921, Orimarga (Orimarga)	4.83
pallidineuris Macquart, 1846, Tipula (Eumicrotipula)	4.28
<i>pallidineris</i> Alexander, 1914, Tipula (Eumicrotipula), error	4.28
pallidipes Alexander, 1926, Helius	4.46
pallidipes (Alexander), 1926 (Eriocera), Hexatoma (Eriocera) ...	4.103
pallidisignata Alexander, 1928, Tipula (Eumicrotipula)	4.28
pallidistyla Alexander, 1942, Austrolimnophila (Austrolimnophila)	4.91
palliditerga Alexander, 1942, Limonia (Dicranomyia)	4.55
<i>pallidula</i> Alexander, 1929, Limonia (Geranomyia)	4.63
pallidus (Philippi), 1865 (Erioptera?), Molophilus	4.149
pallipes (Alexander), 1920 (Psaronius), Lecteria (Psaronius)	4.106
pallipes Alexander, 1964, Polymera (Polymera)	4.111
pallipes (Alexander), 1914 (Mongoma), Trentepohlia (Paramon-	
goma)	4.170
pallitarsis (Alexander), 1937 (Tanypremna), Leptotarsus (Pehlkea)	4.13
pallitarsis (Alexander), 1929 (Gynoplistia), Paralimnophila	4.110
paloides Alexander, 1940, Tipula (Microtipula)	4.37
pammelas (Alexander), 1922 (Gnophomyia), Eugnophomyia	4.127
pamphaea (Alexander), 1929 (Limnophila), Shannonomyia (Shan-	
nonomyia)	4.115
pampoecila Alexander, 1922, Limonia (Dicranomyia)	4.55
panamensis Alexander, 1934, Helius	4.46
panamensis Alexander, 1930, Neognophomyia	4.153
panamensis Alexander, 1922, Ozodicera (Dihexaclonus)	4.18
panehrestus Alexander, 1941, Molophilus (Molophilus)	4.149
pansus Alexander, 1968, Molophilus (Molophilus)	4.149
pantherina Alexander, 1941, Tipula (Eumicrotipula)	4.29
paprzyckiana Alexander, 1944, Neognophomyia	4.153
paprzyckii (Alexander), 1941 (Tipula), Ischnotoma (Neotipula) ..	4.10
paprzyckii Alexander, 1941, Limonia (Caenolimonia)	4.49



paracollaris Alexander, 1945, <i>Teucholebis</i> (<i>Paratropesa</i>)	4.159
Paradelphomyia Alexander, 1936	4.109
paradiceros Alexander, 1966, <i>Molophilus</i> (<i>Molophilus</i>)	4.149
paradoxa Alexander, 1913, <i>Teucholebis</i> (<i>Euteucholebis</i>)	4.158
paracnsis Alexander, 1920, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.141
paraguayana Alexander, 1952, <i>Austrolimnophila</i> (<i>Mediophragma</i>)	4.92
paraguayana Alexander, 1929, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
paraguayanus (Riedel), 1921 (<i>Tipulodina</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
paraguayensis (Alexander), 1935 (<i>Tipula</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.12
paraguayensis Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.149
paraguayensis Alexander, 1929, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
paraguayicola Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.34
Paragymnastes Alexander, 1922	4.142
paralenta Alexander, 1950, <i>Tipula</i> (<i>Microtipula</i>)	4.37
Paralimnophila Alexander, 1921	4.110
Paralipophleps Alexander, 1947	4.140
paramoensis Alexander, 1944, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
Paramongoma Brunetti, 1911	4.169
paranensis Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
pararia Alexander, 1965, <i>Tipula</i> (<i>Microtipula</i>)	4.37
Paratropesa Schiner, 1866	4.158
paraxantha Alexander, 1962, <i>Teucholebis</i> (<i>Teucholebis</i>)	4.164
parilis Alexander, 1945, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
parinermis Alexander, 1940, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
parishi (Alexander), 1913 (<i>Teucholebis</i>), <i>Gonomyia</i> (<i>Gonomyia</i>)	4.134
parishi (Alexander), 1914 (<i>Dicranomyia</i> , <i>Limonia</i> (<i>Dicranomyia</i>))	4.52
parishi Alexander, 1920, <i>Polymera</i> (<i>Polymerodes</i>)	4.113
parishi Alexander, 1912, <i>Tipula</i> (<i>Microtipula</i>)	4.37
parishiana Alexander, 1929, <i>Teucholebis</i> (<i>Teucholebis</i>)	4.164
parrai Alexander, 1946, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.121
parrai Alexander, 1940, <i>Tipula</i> (<i>Bellardina</i>)	4.21
partitus Alexander, 1941, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
parva Osten Sacken, 1859, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.125
parva (Alexander), 1913 (<i>Sacandaga</i>), <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
parvicellula Alexander, 1968, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
parviceps (Speiser), 1909 (<i>Tipula</i>), <i>Zelandotipula</i>	4.42
parvidens Alexander, 1944, <i>Helius</i>	4.46
parviloba Alexander, 1929, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
parvimacula (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
parvipuncta Alexander, 1938, <i>Orimarga</i> (<i>Orimarga</i>)	4.83
parviseta Alexander, 1941, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
parvispinosa Alexander, 1928, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
parvula Alexander, 1947, <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
pastazicola Alexander, 1938, <i>Limonia</i> (<i>Neolimonia</i>)	4.71
pastazina Alexander, 1944, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
pastoris Alexander, 1960, <i>Molophilus</i> (<i>Molophilus</i>)	4.149
patagonica (Alexander), 1928 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Limnophilella</i>)	4.92
patagonica Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
patagonica Alexander, 1929, <i>Tonnoiromyia</i>	4.83
patens (Alexander), 1940 (<i>Ischnothrix</i>), <i>Atarba</i> (<i>Ischnothrix</i>)	4.89
patens Alexander, 1938, <i>Hexatoma</i> (<i>Eriocera</i>)	4.103
patens Alexander, 1939, <i>Teucholebis</i> (<i>Teucholebis</i>)	4.164



patruelis Alexander, 1930, Gonomyia (Progonomyia)	4.142
patruelis (Alexander), 1924 (Dicranomyia), Limonia (Dicranomyia)	4.56
paucilobata Alexander, 1940, Limonia (Caenoglochina)	4.48
pauciseta Alexander, 1949, Gnophomyia	4.130
paucispinosus Alexander, 1925, Molophilus (Molophilus)	4.149
paulista Alexander, 1946, Styringomyia	4.157
paulistae Alexander, 1943, Ctenolimnophila (Campbellomyia)	4.93
paulseni (Philippi), 1865 (Tipula), Leptotarsus (Araucomyia) ...	4.11
paulus Alexander, 1947, Limonia (Rhipidia)	4.77
pectinata (Wiedemann), 1828 (Tipula), Ozodicera (Ozodicera)	4.19
pectinatus Alexander, 1928, Molophilus (Molophilus)	4.149
pectinella (Alexander), 1940 (Tipula), Ischnotoma (Neotipula) ..	4.10
pectiniferus Alexander, 1952, Molophilus (Molophilus)	4.149
Pectinotipula Alexander, 1920	4.39
pedestris Alexander, 1952, Limonia (Dicranomyia)	4.56
pedica Alexander, 1947, Gonomyia (Lipophleps)	4.137
Pedicia Latreille, 1809	4.84
Pediciini	4.84
pediformis Alexander, 1962, Tipula (Eumicrotipula)	4.29
<i>Peditia</i> , error	4.84
<i>Pehlkea</i> Enderlein, 1912	4.13
pelates Alexander, 1962, Limonia (Dicranomyia)	4.56
penai Alexander, 1965, Elephantomyia (Xenoelephantomyia)	4.95
penai (Alexander), 1952 (Holorusia), Ischnotoma (Ischnotoma) ..	4.10
penai Alexander, 1951, Molophilus (Molophilus)	4.149
penai Alexander, 1953, Pedicia (Tricyphona)	4.84
penana Alexander, 1967, Gynoplistia (Gynoplistia)	4.99
penana Alexander, 1966, Limnophila (Limnophila)	4.108
penana Alexander, 1956, Limonia (Dicranomyia)	4.56
penana Alexander, 1951, Tipula (Microtipula)	4.37
penata (Alexander), 1965 (Holorusia), Ischnotoma (Neotipula) ..	4.10
pendulifera Alexander, 1949, Cryptolabis (Cryptolabis)	4.121
penicillatus Alexander, 1941, Molophilus (Molophilus)	4.149
penitus (Alexander), 1944 (Macromastix), Leptotarsus (Araucomyia)	4.11
pennatus Alexander, 1927, Molophilus (Molophilus)	4.149
pennifera Alexander, 1945, Limonia (Dicranomyia)	4.56
pennipes Alexander, 1921, Molophilus (Eumolophilus)	4.143
pentheres (Alexander), 1928 (Geranomyia), Limonia (Geranomyia)	4.65
<i>Pentoptera</i> Schiner, 1863	4.100
penumbrosa Alexander, 1939, Shannonomyia (Shannonomyia)	4.116
peracuta Alexander, 1928, Gonomyia (Paralipophleps)	4.140
peracutior Alexander, 1949, Gnophomyia	4.130
peralta Alexander, 1945, Limonia (Dicranomyia)	4.56
peramoena Alexander, 1945, Limonia (Peripheroptera)	4.73
perangusta Alexander, 1939, Teucholabis (Teucholabis)	4.164
perangustula Alexander, 1938, Tipula (Microtipula)	4.37
perarmata (Alexander), 1921 (Rhipidia), Limonia (Rhipidia)	4.77
perbasalis Alexander, 1949, Teucholabis (Teucholabis)	4.164
percara Alexander, 1958, Austrolimnophila (Austrolimnophila) ..	4.91
percelestis Alexander, 1953, Limonia (Zelandoglochina)	4.80
percompressa Alexander, 1944, Tipula (Microtipula)	4.37
percomptaria Alexander, 1945, Tipula (Microtipula)	4.37



percupida (Alexander), 1941 (Erioptera), Cheilotrichia (Empeda)	4.120
perdebilis Alexander, 1949, Gnophomyia	4.130
perdebilis Alexander, 1943, Molophilus (Molophilus)	4.149
perdecora (Alexander), 1914 (Eriocera), Hexatoma (Eriocera)	4.103
perdelecta Alexander, 1940, Limonia (Peripheroptera)	4.73
perdelecta Alexander, 1942, Tipula (Microtipula)	4.37
perdistalis Alexander, 1943, Limonia (Dicranomyia)	4.56
perdistans Alexander, 1967, Cryptolabis (Cryptolabis)	4.121
perdistinctus (Alexander), 1942 (Tanypremna), Leptotarsus (Tanypremnella)	4.15
perenensis (Alexander), 1923 (Eriocera), Hexatoma (Eriocera)	4.103
perexcelsior Alexander, 1962, Limonia (Dicranomyia)	4.56
prexigua Alexander, 1941, Hexatoma (Eriocera)	4.103
perexquisita Alexander, 1938, Gymnastes (Paragymnastes)	4.143
perextensus Alexander, 1951, Molophilus (Molophilus)	4.149
perfecta (Alexander), 1928 (Geranomyia), Limonia (Geranomyia)	4.65
perfestiva Alexander, 1951, Hexatoma (Eriocera)	4.103
perfidus Alexander, 1929, Molophilus (Molophilus)	4.149
perflaveola Alexander, 1927, Limonia (Dicranomyia)	4.56
perfuga Alexander, 1944, Ozodicera (Dihexaclonus)	4.18
perglabrata Alexander, 1962, Rhabdomastix (Rhabdomastix)	4.155
perglabratus (Alexander), 1953 (Longurio), Leptotarsus (Longurio)	4.13
pergracilis Alexander, 1943, Limnophila (Limnophila)	4.108
pergracilis Alexander, 1944, Toxorhina (Toxorhina)	4.168
perincisa Alexander, 1948, Atarba (Atarba)	4.87
Peripheroptera Schiner, 1866	4.72
perirrorata (Alexander), 1928 (Gynoplistia), Paralimnophila	4.110
perjovialis Alexander, 1944, Tipula (Eumicrotipula)	4.29
perlaeta (Alexander), 1922 (Eriocera), Hexatoma (Eriocera)	4.103
perlata Alexander, 1944, Gnophomyia	4.130
perlata Alexander, 1943, Teucholabis (Teucholabis)	4.164
perlaticosta Alexander, 1941, Tipula (Microtipula)	4.37
perlimbata Alexander, 1942, Austrolimnophila (Austrolimnophila)	4.91
permagica Alexander, 1947, Gnophomyia	4.130
permonstrata (Alexander), 1935 (Limnophila), Shannonomyia (Roraimomyia)	4.114
pernobilis Alexander, 1941, Limonia (Neoglochina)	4.69
perornatus (Alexander), 1929 (Tanypremna), Leptotarsus (Tanypremna)	4.15
perpallens Alexander, 1964, Orimarga (Orimarga)	4.83
Perpallidus Alexander, 1942, Helius	4.46
perpulchra (Alexander), 1914 (Eriocera), Hexatoma (Eriocera)	4.103
perpuncticosta Alexander, 1950, Limonia (Dicranomyia)	4.56
perrara Alexander, 1936, Hexatoma (Eriocera)	4.103
perreducta (Alexander), 1938 (Gynoplistia), Paralimnophila	4.110
perretracta Alexander, 1942, Limonia (Dicranomyia)	4.56
perrostratus Alexander, 1965, Molophilus (Rhynchomolophilus)	4.152
persancta Alexander, 1938, Epiphragma (Epiphragma)	4.97
perserena Alexander, 1946, Limonia (Dicranomyia)	4.56
perseus Alexander, 1913, Molophilus (Molophilus)	4.149
persessilis Alexander, 1939, Austrolimnophila (Austrolimnophila)	4.91
persevera Alexander, 1949, Gnophomyia	4.130
persimilis (Alexander), 1920 (Teucholabis), Gonomyia (Gonomyia)	4.134



persimplex Alexander, 1950, Limonia (Rhipidia)	4.77
persinuosus Alexander, 1941, Molophilus (Molophilus)	4.149
perstrangalia (Alexander), 1962 (Holorusia), Zelandotipula	4.43
perstudiosa Alexander, 1937, Tipula (Eumicrotipula)	4.29
persuffusa Alexander, 1967, Limonia (Rhipidia)	4.77
pertenuis Alexander, 1952, Molophilus (Molophilus)	4.149
pertruncata Alexander, 1962, Limonia (Dicranomyia)	4.56
peruviana Alexander, 1922, Gonomyia (Progonomyia)	4.142
peruviana (Alexander), 1914 (Eriocera), Hexatoma (Eriocera) ..	4.103
peruviana Alexander, 1926, Rhabdomastix (Rhabdomastix)	4.155
peruviana (Alexander), 1914 (Holorusia), Zelandotipula	4.43
pervenustus Alexander, 1953, Helius	4.46
perversa Alexander, 1937, Hexatoma (Eriocera)	4.103
pervia Alexander, 1966, Hexatoma (Eriocera)	4.104
pervicax (Alexander), 1914 (Gnophomyia), Neognophomyia	4.153
pervincta Alexander, 1936, Limonia (Zelandoglochina)	4.80
petalina Alexander, 1947, Epiphragma (Epiphragma)	4.97
petalura Alexander, 1953, Tipula (Eumicrotipula)	4.29
petentis Alexander, 1949, Gnophomyia	4.130
petiolaris Alexander, 1940, Tipula (Eumicrotipula)	4.29
petronia Alexander, 1942, Gonomyia (Lipophleps)	4.137
petulans Alexander, 1937, Trentepohlia (Paramongoma)	4.170
Phacelodocera Enderlein, 1912	4.20
phaedra (Alexander), 1949 (Macromastix), Leptotarsus (Longurio)	4.13
phaeostigma Alexander, 1962, Pedicia (Tricyphona)	4.84
phaeostigmata Alexander, 1943, Shannonomyia (Shannonomyia)	4.116
phaeoanthera Alexander, 1944, Epiphragma (Epiphragma)	4.97
phaeton (Alexander), 1945 (Holorusia) Isehnotomyia (Icriomastax)	4.9
phalangioides Alexander, 1945, Tipula (Eumicrotipula)	4.29
phallacantha Alexander, 1942, Ozodicera (Ozodicera)	4.19
phalodontus Alexander, 1968, Molophilus (Molophilus)	4.149
phallosomicus Alexander, 1939, Molophilus (Molophilus)	4.149
phallostena Alexander, 1968, Cryptolabis (Cryptolabis)	4.121
phantasma Alexander, 1943, Quechuamyia	4.154
phaon Alexander, 1950, Limonia (Rhipidia)	4.77
phasmatis Alexander, 1945, Helius	4.46
phatta (Philippi), 1865 (Limnobia), Limonia (Dicranomyia)	4.56
philippiana Alexander, 1920, Tipula (Eumicrotipula)	4.29
<i>Philippiana</i> Alexander, 1929	4.154
<i>philippii</i> (Alexander), 1913 (Geranomyia), Limonia (Geranomyia)	4.67
philomela Alexander, 1945, Gonomyia (Lipophleps)	4.137
phoracaena Alexander, 1966, Toxorhina (Toxorhina)	4.168
phoroctenia Alexander, 1921, Gonomyia (Lipophleps)	4.137
phragmophora Alexander, 1937, Shannonomyia (Shannonomyia) .	4.116
phrixus Alexander, 1953, Brachypremna	4.7
phylax (Alexander), 1953 (Longurio), Leptotarsus (Tanypremna)	4.15
piabilis Alexander, 1944, Limonia (Dicranomyia)	4.52
piatrix Alexander, 1942, Hexatoma (Eriocera)	4.104
piatrix Alexander, 1944, Ozodicera (Ozodicera)	4.19
pictipennis (Philippi), 1865 (Ctedonia), Gynoplistia (Gynoplistia)	4.99
pictipennis Alexander, 1937, Sigmatomera (Sigmatomera)	4.157
pictipennis Walker, 1837, Tipula (Eumicrotipula)	4.29
pictipes Osten Sacken, 1887, Brachypremna	4.7
pictiventris Alexander, 1945, Brachypremna	4.7



<i>pictiventris</i> Alexander, 1945, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.94
<i>picturata</i> (Alexander), 1929 (<i>Ischnothrix</i>), <i>Atarba</i> (<i>Ischnothrix</i>)	4.89
<i>picturellus</i> (Alexander), 1941 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tanypremna</i>)	4.15
<i>pietatis</i> Alexander, 1943, <i>Limonia</i> (<i>Dicranomyia</i>)	4.56
<i>piger</i> Alexander, 1942, <i>Molophilus</i> (<i>Molophilus</i>)	4.149
<i>Pilaria</i> Sintenis, 1888	4.110
<i>pilipes</i> (Fabricius), 1787 (<i>Tipula</i>), <i>Erioptera</i> (<i>Trimicra</i>)	4.126
<i>pilipes</i> (Walker), 1856 (<i>Geranomyia</i>), <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
<i>pilosipennis</i> Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.80
<i>pilosipes</i> (Alexander), 1929 (<i>Philippiana</i>), <i>Erioptera</i> (<i>Erioptrella</i>)	4.123
<i>pilosispina</i> Alexander, 1936, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
<i>pilulifera</i> Edwards, 1919, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
<i>pinckerti</i> Alexander, 1962, <i>Neognophomyia</i>	4.153
<i>pinodes</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.56
<i>pirioni</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>pirioni</i> (Alexander), 1928 (<i>Gynoplistia</i>), <i>Paralimnophila</i>	4.110
<i>pirioni</i> Alexander, 1928, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
<i>piro</i> Alexander, 1914, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
<i>placabilis</i> Alexander, 1941, <i>Teucholabis</i> (<i>Paratropesa</i>)	4.159
<i>placata</i> Alexander, 1965, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.20
<i>plagifera</i> (Alexander), 1943 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
<i>planitas</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>platensis</i> (Alexander), 1923 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.91
<i>platensis</i> (Alexander), 1923 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.65
<i>platensis</i> (Alexander), 1923 (<i>Limnophila</i>), <i>Paralimnophila</i>	4.110
<i>platymera</i> Alexander, 1939, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
<i>platymera</i> Alexander, 1943, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>platymerella</i> Alexander, 1947, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>platymerina</i> Alexander, 1944, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>platymeroides</i> Alexander, 1952, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>platyphallus</i> Alexander, 1942, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>platyphallus</i> Alexander, 1945, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
<i>platyptera</i> (Alexander), 1929 (<i>Tricyphona</i>), <i>Pedicia</i> (<i>Tricyphona</i>)	4.85
<i>platytergata</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
<i>plaumanni</i> Alexander, 1936, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104
<i>plaumanni</i> Alexander, 1947, <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
<i>plaumanniana</i> Alexander, 1938, <i>Sigmatomera</i> (<i>Austrolimnobia</i>)	4.156
<i>plaumannina</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.37
<i>plebeius</i> Alexander, 1954, <i>Helius</i>	4.46
<i>Plectromyia</i> Osten Sacken, 1869	4.84
<i>Plettusa</i> Philippi, 1865	4.60
<i>pleuralis</i> (Williston), 1896 (<i>Atarba</i>), <i>Gonomyia</i> (<i>Paralipophleps</i>)	4.140
<i>pleuralis</i> (Alexander), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>)	4.78
<i>pleuralis</i> Alexander, 1913, <i>Polymera</i> (<i>Polymera</i>)	4.112
<i>pleuralis</i> Alexander, 1913, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
<i>pleurolinea</i> Alexander, 1959, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
<i>pleurolineata</i> Alexander, 1929, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>pleurostriata</i> Alexander, 1939, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
<i>pleurotaeniata</i> Alexander, 1942, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.135
<i>plumbeicolor</i> Alexander, 1942, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104
<i>plumbeicolor</i> Alexander, 1938, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
<i>plumbeinota</i> Alexander, 1940, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104



plumbeipleura (Alexander), 1916 (Geranomyia), Limonia (Geranomyia)	4.65
plumbeithorax Alexander, 1921, Tipula (Microtipula)	4.37
pluto Alexander, 1939, Pseudolimnophila	4.113
plutoides Alexander, 1948, Pseudolimnophila	4.113
pluvialis Alexander, 1929, Limonia (Dicranomyia)	4.56
podacantha Alexander, 1944, Gnophomyia	4.130
podagra Alexander, 1952, Teucholabis (Teucholabis)	4.164
polingi Alexander, 1946, Gonomyia (Euptilostena)	4.132
polita Osten Sacken, 1887, Teucholabis (Teucholabis)	4.164
polycantha Alexander, 1940, Toxorhina (Toxorhina)	4.168
polyelada Alexander, 1943, Limonia (Rhipidia)	4.78
polydonta Alexander, 1944, Erioptera (Erioptera)	4.123
polygrapha Alexander, 1939, Limonia (Dicranomyia)	4.51
<i>Polymeda</i> Meigen, 1800	4.122
Polymera Wiedemann, 1820	4.111
Polymerodes Alexander, 1920	4.113
<i>Polymoria</i> Philippi, 1865	4.109
<i>Polymoria</i> Philippi of Alexander, 1924	4.89
<i>Polypraesidia</i> Miller, 1943	4.109
polyspilota Alexander, 1937, Austrolimnophila (Austrolimnophila)	4.91
polystieta (Philippi), 1865 (Limnobia), Limonia (Dicranomyia) ..	4.57
polystonyx Alexander, 1968, Limonia (Dicranomyia)	4.57
polytricha Alexander, 1945, Erioptera (Erioptera)	4.123
pomerantzi Alexander, 1964, Shannonomyia (Shannonomyia)	4.116
pontifex Alexander, 1967, Tipula (Microtipula)	4.37
porrecta Alexander, 1950, Limonia (Neolimonia)	4.71
porrecta Alexander, 1951, Tipula (Microtipula)	4.37
porteri Alexander, 1929, Gnophomyia	4.130
porteri (Alexander), 1929 (Holorusia), Ischnotoma (Ischnotoma) .	4.10
porteri (Alexander), 1919 (Dicranomyia), Limonia (Idioglochina)	4.68
porterianus (Alexander), 1936 (Tanypremna), Leptotarsus (Tany-	
premna)	4.15
portorieana Alexander, 1936, Teucholabis (Teucholabis)	4.164
portoricensis (Alexander), 1912 (Megistomastix), Dolichopeza	
(Megistomastix)	4.8
portorieensis Alexander, 1932, Erioptera (Mesocyphona)	4.125
postea Alexander, 1929, Gynoplistia (Gynoplistia)	4.99
posticata (Alexander), 1939 (Gnophomyia), Eugnophomyia	4.127
posticata Alexander, 1929, Rhabdomastix (Rhabdomastix)	4.155
postnotalis (Alexander), 1929 (Holorusia), Ischnotoma (Ischnotoma)	4.10
praeellens Alexander, 1944, Limonia (Dicranomyia)	4.57
praelara Alexander, 1928, Limonia (Neoglochina)	4.69
praenobilis Alexander, 1945, Teucholabis (Euparatropesa)	4.158
praepostera (Alexander), 1925 (Dicranomyia), Limonia (Dicra-	
nomyia)	4.57
praeusta (Osten Saeken), 1886 (Paratropesa), Teucholabis (Para-	
tropesa)	4.159
praevia Alexander, 1952, Limonia (Dicranomyia)	4.57
pratti Alexander, 1950, Limonia (Rhipidia)	4.78
pretiosa (Osten Saeken), 1886 (Eriocera), Hexatoma (Eriocera) .	4.104
pretiosa Alexander, 1945, Tipula (Microtipula)	4.37
pretiosus Alexander, 1929, Molophilus (Molophilus)	4.150



primitiva Alexander, 1943, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.94
primogenia Alexander, 1948, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.94
prindlei Alexander, 1938, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
pritchardi Alexander, 1940, <i>Tipula</i> (<i>Microtipula</i>)	4.37
proavita (Alexander), 1940 (<i>Tanypremna</i>), <i>Leptotarsus</i> (<i>Tany-</i> <i>premna</i>)	4.15
problematica (Alexander), 1945 (<i>Holorusia</i>), <i>Ischnotoma</i> (<i>Ischno-</i> <i>toma</i>)	4.10
problematica Alexander, 1943, <i>Jivaromyia</i>	4.143
procella Alexander, 1944, <i>Limnophila</i> (<i>Limnophila</i>)	4.107
procax Alexander, 1946, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
procericornis Alexander, 1944, <i>Atarba</i> (<i>Atarba</i>)	4.87
procericornis Edwards, 1919, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
Procryptolabis Alexander, 1923	4.122
proctigerica Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.78
proctotricha Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.37
producta Alexander, 1919, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
productellus Alexander, 1944, <i>Helius</i>	4.46
productissima Alexander, 1944, <i>Neognophomyia</i>	4.153
profana Alexander, 1941, <i>Limonia</i> (<i>Rhipidia</i>)	4.78
profuga Alexander, 1937, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
profusa Alexander, 1945, <i>Orimarga</i> (<i>Diotrepha</i>)	4.82
Progonomyia Alexander, 1920	4.141
projecta Alexander, 1941, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
projecta Alexander, 1949, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
proliferata Alexander, 1940, <i>Limonia</i> (<i>Rhipidia</i>)	4.78
prolixicornis Alexander, 1927, <i>Polymera</i> (<i>Polymera</i>)	4.112
prolixisterna Alexander, 1941, <i>Tipula</i> (<i>Microtipula</i>)	4.37
prolixistylus Alexander, 1930, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
prolongata Alexander, 1940, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
prolongata Alexander, 1938, <i>Toxorhina</i> (<i>Ceratocheilus</i>)	4.167
Promongoma Alexander, 1938	4.170
propatula Alexander, 1947, <i>Gnophomyia</i>	4.130
propera Alexander, 1949, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
propinqua Alexander, 1936, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104
propitia Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.30
proseni Alexander, 1954, <i>Limonia</i> (<i>Rhipidia</i>)	4.78
Protohelius Alexander, 1928	4.83
protudens Alexander, 1952, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
protuberans Alexander, 1946, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
providens Alexander, 1950, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.116
provocator Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
proxima (Bellardi), 1859 (<i>Tipula</i>), <i>Nephrotoma</i>	4.17
Psaronius Enderlein, 1912	4.106
Pselliophora Osten Sacken, 1886	4.7
Pseudolimnophila Alexander, 1919	4.113
Psiloconopa Zetterstedt, 1838	4.126
psittacina Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.29
pterelachisus Rondani, 1842	4.40
pubescens (Alexander), 1913 (<i>Erioptera</i>), <i>Cheilotrichia</i> (<i>Empeda</i>)	4.120
puella (Williston), 1896 (<i>Atarba</i>), <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
puer Alexander, 1913, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137




<i>pugilis</i> Alexander, 1943, <i>Limonia</i> (<i>Neolimnobia</i>)	4.70
<i>pugnax</i> Alexander, 1946, <i>Limonia</i> (<i>Caenogloehina</i>)	4.48
<i>pulchella</i> (Alexander), 1914 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.61
<i>pulehella</i> (Rondani), 1850 (<i>Paehyrhina</i>), <i>Nephrotoma</i>	4.17
<i>pulehella</i> Alexander, 1913, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.164
<i>pulehrieornis</i> Alexander, 1914, <i>Polymera</i> (<i>Polymera</i>)	4.112
<i>pulchripes</i> (Alexander), 1922 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.104
<i>pullatipes</i> Alexander, 1951, <i>Hexatoma</i> (<i>Eriocera</i>)	4.104
<i>pullipes</i> Alexander, 1937, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
<i>pulvinaris</i> Alexander, 1945, <i>Gnophomyia</i>	4.130
<i>pulvinus</i> Alexander, 1943, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>pumila</i> Alexander, 1942, <i>Ozodicera</i> (<i>Dihexaelonus</i>)	4.18
<i>punetatissima</i> (Wiedemann), 1828 (<i>Limnobia</i>), <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>punetifrons</i> (Maequart), 1838 (<i>Paehyrhina</i>), <i>Nephrotoma</i>	4.17
<i>punetipennis</i> (Philippi), 1865 (<i>Polymoria</i>), <i>Austrolimnophila</i> (<i>Aus-</i> <i>trolimnophila</i>)	4.91
<i>punctipennis</i> (Alexander), 1914 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ..	4.78
<i>punetipennis</i> Alexander, 1914, <i>Orimarga</i> (<i>Orimarga</i>)	4.83
<i>punetiscuta</i> Alexander, 1922, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>punetoria</i> Alexander, 1929, <i>Limonia</i> (<i>Rhipidia</i>)	4.78
<i>punoensis</i> Alexander, 1945, <i>Limonia</i> (<i>Dieranomyia</i>)	4.57
<i>pupillata</i> Alexander, 1913, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.97
<i>pusilla</i> (Bigot), 1888 (<i>Trieyphona</i>)	4.85
<i>pustulatus</i> Alexander, 1946, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>pyensoni</i> Alexander, 1939, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
<i>pygmaea</i> (Alexander), 1914 (<i>Psaronius</i>), <i>Leetoria</i> (<i>Psaronius</i>)	4.106
<i>pygmaeus</i> (Alexander), 1914 (<i>Maeromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.13
<i>quadricincta</i> Alexander, 1927, <i>Erioptera</i> (<i>Erioptera</i>)	4.123
<i>quadricollis</i> Alexander, 1945, <i>Tipula</i> (<i>Microtipula</i>)	4.37
<i>quadrifidus</i> Alexander, 1926, <i>Helius</i>	4.46
<i>quadrifureata</i> Alexander, 1928, <i>Erioptera</i> (<i>Mesoeypfona</i>)	4.125
<i>quadrigladiata</i> Alexander, 1944, <i>Limonia</i> (<i>Dieranomyia</i>)	4.57
<i>quadrilineata</i> (Maequart), 1838 (<i>Paehyrhina</i>), <i>Nephrotoma</i>	4.17
<i>quadrimaeulata</i> Bellardi, 1859 (<i>Tipula</i>)	4.44
<i>quadrisetosa</i> Alexander, 1951, <i>Tipula</i> (<i>Eumierotipula</i>)	4.29
<i>quadristylus</i> Alexander, 1922, <i>Molophilus</i> (<i>Molophilus</i>)	4.150
<i>quadrituberculata</i> Alexander, 1937, <i>Limonia</i> (<i>Dieranomyia</i>)	4.57
<i>quaesita</i> Alexander, 1938, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>quasimodo</i> Alexander, 1950, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>quasimodo</i> Alexander, 1943, <i>Braehypremna</i>	4.7
<i>Quechuamyia</i> Alexander, 1943	4.154
<i>queribunda</i> Alexander, 1941, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.133
<i>quiehua</i> Alexander, 1916, <i>Tipula</i> (<i>Eumierotipula</i>)	4.30
<i>quinquecincta</i> Alexander, 1927, <i>Erioptera</i> (<i>Erioptera</i>)	4.123
<i>quinqueplagiata</i> Alexander, 1931, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>rabelloi</i> (Alexander), 1953 (<i>Longurio</i>), <i>Leptotarsus</i> (<i>Longurio</i>) ..	4.13
<i>rabula</i> Alexander, 1940, <i>Limonia</i> (<i>Geranomyia</i>)	4.65
<i>ramus</i> Alexander, 1943, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.137
<i>rapax</i> (Alexander), 1921 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Caenogloehina</i>)	4.48
<i>rapida</i> Alexander, 1946, <i>Limonia</i> (<i>Dieranomyia</i>)	4.57
<i>raptor</i> Alexander, 1947, <i>Molophilus</i> (<i>Molophilus</i>)	4.150




rara (Alexander), 1929 (Gynoplistia), Paralimnophila	4.110
rastriformis Alexander, 1946, Gonomyia (Lipophleps)	4.137
ravida (Alexander), 1925 (Dicranomyia), Limonia (Dicranomyia)	4.57
<i>reciproca</i> (Walker), 1848 (Limnobia), Erioptera (Trimicra)	4.126
<i>reciproca</i> Alexander, 1946, Tipula (Eumicrotipula)	4.30
recisa (Alexander), 1927 (Geranomyia), Limonia (Geranomyia) ..	4.65
recondita (Alexander), 1921 (Geranomyia), Limonia (Geranomyia)	4.65
rectangula Alexander, 1966, Gonomyia (Gonomyia)	4.134
rectangularis Alexander, 1954, Atarba (Ischnothrix)	4.89
rectangularis Alexander, 1952, Teucholabis (Teucholabis)	4.164
rectangulus Alexander, 1951, Tipula (Microtipula)	4.38
rectispina Alexander, 1947, Helius	4.47
rectispina Alexander, 1968, Teucholabis (Teucholabis)	4.164
rectispinus Alexander, 1952, Molophilus (Molophilus)	4.150
rectus Alexander, 1945, Helius	4.47
recurvata Alexander, 1942, Cryptolabis (Cryptolabis)	4.121
recurvata Alexander, 1941, Gonomyia (Lipophleps)	4.137
rediviva Alexander, 1942, Limonia (Peripheroptera)	4.73
redunca Alexander, 1967, Tipula (Eumicrotipula)	4.30
reedi Alexander, 1934, Tipula (Eumicrotipula)	4.30
refuga Alexander, 1944, Limonia (Geranomyia)	4.65
regifica (Alexander), 1916 (Dicranomyia), Limonia (Dicranomyia)	4.57
regina (Alexander), 1914 (Tanypremna), Leptotarsus (Pehlkea) .	4.13
regina Alexander, 1926, Polymera (Polymera)	4.112
regius Alexander, 1944, Helius	4.47
regnatix Alexander, 1943, Gnophomyia	4.130
regressa Alexander, 1950, Tipula (Microtipula)	4.38
regulus (Alexander), 1943 (Tanypremna), Leptotarsus (Pelkea) .	4.13
relata Alexander, 1944, Limonia (Geranomyia)	4.65
religiosa Alexander, 1946, Atarba (Atarba)	4.87
religiosa Alexander, 1946, Tipula (Trichotipula)	4.40
remiger Alexander, 1922, Molophilus (Molophilus)	4.150
remigera Alexander, 1946, Gonomyia (Gonomyia)	4.134
<i>remingtoni</i> Alexander, 1946, Limonia (Geranomyia)	4.63
remota Alexander, 1926, Gonomyia (Gonomyia)	4.134
<i>repanda</i> Alexander, 1928, Limonia (Caenolimonia)	4.49
repentina Alexander, 1929, Limonia (Dicranomyia)	4.57
resplendens Alexander, 1946, Tipula (Eumicrotipula)	4.30
restricta Alexander, 1943, Atarba (Atarba)	4.87
reticulata (Alexander), 1912 (Furcomyia), Limonia (Dicranomyia)	4.57
retractior Alexander, 1943, Austrolimnophila (Limnophilella)	4.92
retrorsa Alexander, 1962, Tipula (Microtipula)	4.38
retrorsus Alexander, 1939, Molophilus (Molophilus)	4.150
rectusa Alexander, 1935, Teucholabis (Teucholabis)	4.164
reverentia Alexander, 1949, Hexatoma (Ericera)	4.104
revulsa Alexander, 1954, Toxorhina (Ceraticheilus)	4.167
reyesi Alexander, 1946, Gonomyia (Lipophleps)	4.137
Rhodomastix Skuse, 1890	4.154
rhadnophora Alexander, 1942, Teucholabis (Teucholabis)	4.164
rhadnostyla Alexander, 1964, Gonomyia (Lipophleps)	4.138
<i>Rhamphidia</i> Meigen, 1830	4.45
Rhamphoncurus Alexander, 1929	4.117



Rhaphidolabis Osten Sacken, 1869	4.84
Rhipidia Meigen, 1818	4.74
rhoda Alexander, 1939, Limonia (Periphroptera)	4.73
Rhynchomolophilus Alexander, 1965	4.152
richardsi Alexander, 1929, Molophilus (Molophilus)	4.150
riedeliana (Enderlein), 1917 (Dirhipis), Gynoplistia (Dirhipis) ...	4.98
ringens (Alexander), 1935 (Holorusia), Zelandotipula	4.43
riveti Edwards, 1919, Tipula (Eumicrotipula)	4.30
roraima Alexander, 1931 Limnophila (Limnophila)	4.108
roraimac Alexander, 1931, Limonia (Neolimonia)	4.72
roraimella Alexander, 1935, Hexatoma (Eriocera)	4.104
roraimensis Alexander, 1935, Shannonomyia (Shannonomyia) ...	4.116
roraimicola Alexander, 1931, Limnophila (Limnophila)	4.108
roraimicola Alexander, 1935, Trentepohlia (Paramongoma)	4.170
Roraimomyia Alexander, 1935	4.113
rostrata antillarum Alexander, 1929, Limonia (Geranomyia)	4.65
rostrata Enderlein, 1912, Teucholabis (Teucholabis)	4.164
rostrifera Alexander, 1946, Dicranota (Rhaphidolabis)	4.84
rounds Alexander, 1939, Cryptolabis (Cryptolabis)	4.121
rubecula Alexander, 1944, Limnophila (Limnophila)	4.108
rubella Alexander, 1926, Palaria	4.110
rubicundula Alexander, 1921, Gnophomyia	4.130
rubicundus (Alexander), 1922 (Rhamphidia), Helius	4.47
rubidithorax Alexander, 1929, Molophilus (Molophilus)	4.150
rubiginosa Alexander, 1931, Limonia (Geranomyia)	4.65
rubriceps Alexander, 1937, Teucholabis (Teucholabis)	4.164
rubroatra Alexander, 1962, Teucholabis (Teucholabis)	4.164
rucana Alexander, 1946, Tipula (Eumicrotipula)	4.30
rufescens (Loew), 1851 (Aporosa), Limonia (Geranomyia)	4.65
ruficornis (Macquart), 1838 (Cylindrotoma), Hexatoma (Eriocera)	4.104
rufirostris Bigot, 1888, Tipula (Eumicrotipula)	4.30
rufithorax (Wiedemann), 1828 (Limnobia), Sigmatomera (Eufurina)	4.156
rufostigma (Macquart), 1838 (Tipula), Ischnotoma (Ischnotoma)	4.10
rufula Alexander, 1946, Teucholabis (Teucholabis)	4.165
runa Alexander, 1947, Gonomyia (Gonomyina)	4.134
runtunensis Alexander, 1940, Tipula (Eumicrotipula)	4.30
rupununi Alexander, 1945, Hexatoma (Eriocera)	4.104
rutristylus Alexander, 1968, Amphineurus (Rhamphineurus)	4.118
sabethoides Edwards, 1927, Molophilus (Eumolophilus)	4.143
Sacandaga Alexander, 1911	4.155
sackeni Williston, 1896, Epiphragma (Epiphragma)	4.97
sackeni Alexander, 1913, Teucholabis (Teucholabis)	4.165
saga Alexander, 1940, Hexatoma (Eriocera)	4.104
sagax Alexander, 1938, Molophilus (Molophilus)	4.150
sagitta Alexander, 1949, Gnophomyia	4.130
sagittarius Alexander, 1914, Molophilus (Molophilus)	4.150
sagittoides Alexander, 1951, Gnophomyia	4.130
salmani Alexander, 1927, Gonomyia (Gonomyia)	4.134
salome (Alexander), 1945 (Tanypremna), Leptotarsus (Tanypremna)	4.15
saltatrix (Alexander), 1944 (Tanypremna), Leptotarsus (Tanypremna)	4.15
salti Alexander, 1930, Teucholabis (Teucholabis)	4.165



saluatoria Alexander, 1945, Tipula (Eumicrotipula)	4.28
salva Alexander, 1942, Teucholabis (Teucholabis)	4.165
sana Alexander, 1936, Gonomyia (Lipophleps)	4.138
sanctaecatharinae (Alexander), 1945 (Tanypremna), Leptotarsus (Tanypremna)	4.15
sanctaccruzae (Alexander), 1920 (Dicranomyia), Limonia (Dicra- nomyia)	4.57
sanctacmartae (Alexander), 1919 (Pentoptera), Hexatoma (Eriocera)	4.104
sanctaeartae Alexander, 1930, Limonia (Neolimonia)	4.72
sandersi Alexander, 1931, Gonomyia (Lipophleps)	4.138
sanguinea Alexander, 1938, Teucholabis (Teucholabis)	4.165
sanguinolenta Alexander, 1938, Teucholabis (Euparatropesa)	4.158
sanguinolentus (Alexander), 1921 (Rhamphidia), Helius	4.47
sanus Alexander, 1929, Amphineurus (Rhamphoneurus)	4.118
sappho Alexander, 1943, Brachypremna	4.7
sappho Alexander, 1943, Epiphragma (Epiphragma)	4.98
sariapampae Alexander, 1967, Tipula (Eumicrotipula)	4.30
satipoana Alexander, 1945, Limonia (Geranomyia)	4.65
satipoensis Alexander, 1944, Rhabdomastix (Rhabdomastix)	4.155
satrapa Alexander, 1952, Tipula (Eumicrotipula)	4.30
saturata Alexander, 1927, Erioptera (Mesocyphona)	4.125
saturata Alexander, 1937, Gonomyia (Progonomyia)	4.142
saturnina Alexander, 1943, Orimarga (Orimarga)	4.83
saxicola Alexander, 1923, Gonomyia (Progonomyia)	4.142
sayi Alexander, 1911, Tipula (Yamatotipula)	4.40
scabricornis Alexander, 1942, Molophilus (Molophilus)	4.150
scabrifolia Alexander, 1967, Erioptera (Mesocyphona)	4.125
scabriseta Alexander, 1938, Orimarga (Orimarga)	4.83
scabristyla Alexander, 1968, Limonia (Neolimonia)	4.72
scabrosa Alexander, 1944, Atarba (Atarba)	4.87
scabrosa Alexander, 1938, Teucholabis (Teucholabis)	4.165
scaenalis Alexander, 1951, Limonia (Caenoglochina)	4.48
scapania Alexander, 1966, Toxorhina (Toxorhina)	4.168
scapha Alexander, 1945, Neognophomyia	4.153
scaphoides Alexander, 1952, Neognophomyia)	4.154
scaphula Alexander, 1945, Tipula (Microtipula)	4.38
scapularis (Macquart), 1838 (Rhamphidia), Teucholabis (Teucholabis)	4.165
scaramuzzai Alexander, 1937, Shannonomyia (Shannonomyia)	4.116
scelerata Alexander, 1945, Gonomyia (Lipophleps)	4.138
sclerosa Alexander, 1948, Polymera (Polymera)	4.112
scelesta Alexander, 1945, Tipula (Microtipula)	4.38
<i>Scepasma</i> Enderlein, 1917	4.98
schachovskoyana Alexander, 1960, Gynoplistia (Gynoplistia)	4.99
schachovskoyi Alexander, 1952, Tipula (Eumicrotipula)	4.30
schadeana Alexander, 1935, Gonomyia (Neolipophleps)	4.139
schadei Alexander, 1935, Cryptolabis (Cryptolabis)	4.121
schadci Alexander, 1926, Limnophila (Limnophila)	4.108
schadei Alexander, 1929, Limonia (Rhipidia)	4.78
schauseana Alexander, 1966, Hexatoma (Eriocera)	4.104
schildeana Alexander, 1966, Hexatoma (Eriocera)	4.104
schildeana Alexander, 1945, Tipula (Microtipula)	4.38



sehildi Alexander, 1945, Helius	4.47
sehildi Alexander, 1945, Neognophomyia	4.154
schindleri Alexander, 1962, Limonia (Dieranomyia)	4.57
sehincri (Alexander), 1922 (Eriocera), Hexatoma (Eriocera)	4.104
schinceri (Osten Saeken), 1887 (Peripheroptera), Limonia (Peri- pheroptera)	4.73
schineri Enderlein, 1912, Teucholabis (Teucholabis)	4.165
sehineri (Alexander), 1934 (Holorusia), Zelandotipula	4.43
sehineriana (Alexander), 1928 (Holorusia), Isehnnotoma (Ischno- toma)	4.10
sehizomera Alexander, 1940, Tipula (Bellardina)	4.21
sehultzei Alexander, 1946, Molophilus (Molophilus)	4.150
schunkeana Alexander, 1948, Austrolimnophila (Limnophilella) ..	4.92
sehwarzi (Alexander), 1912 (Rhipidia), Limonia (Rhipidia)	4.78
sehwarzmaierana Alexander, 1942, Ozodieera (Ozodicera)	4.20
sehwarzmaieri Alexander, 1940, Tipula (Mierotipula)	4.38
sciasma Alexander, 1950, Limonia (Neoglochina)	4.69
seimitar Alexander, 1914, Gonomyia (Lipophleps)	4.138
seimitar Alexander, 1942, Limonia (Dieranomyia)	4.57
scolopax (Alexander), 1913 (Geranomyia), Limonia (Geranomyia)	4.66
scriptella Alexander, 1940, Tipula (Eumierotipula)	4.30
seulleniana Alexander, 1965, Hexatoma (Eriocera)	4.104
scutata Alexander, 1939, Atarba (Atarba)	4.87
scutellata (Alexander), 1929 (Isehnothrix), Atarba (Isehnothrix)	4.89
seespita Alexander, 1936, Gonomyia (Lipophleps)	4.138
seelusa (Alexander), 1928 (Limnophila), Shannonomyia (Shanno- nomyia)	4.116
seguyi Alexander, 1930, Sigmatomera (Sigmatomera)	4.157
segnipes (Alexander), 1953 (Longurio), Leptotarsus (Tanypre- nella)	4.16
sejugata Alexander, 1939, Limonia (Rhipidia)	4.78
selanderi Alexander, 1956, Tipula (Trichotipula)	4.40
selkirki (Alexander), 1920 (Dicranomyia), Limonia (Dicranomyia)	4.57
selkirkiana Alexander, 1952, Shannonomyia (Shannonomyia)	4.116
selkirkianus (Enderlein), 1940 (Arehimolophilus), Molophilus (Molophilus)	4.150
semiflava Alexander, 1949, Cryptolabis (Cryptolabis)	4.121
semirufa (Alexander), 1927 (Eriocera), Hexatoma (Eriocera)	4.104
semivulpina Alexander, 1944, Tipula (Eumicrotipula)	4.30
senaria Alexander, 1943, Gonomyia (Lipophleps)	4.138
separata (Alexander), 1921 (Geranomyia), Limonia (Geranomyia)	4.66
seposita Alexander, 1929, Limonia (Dieranomyia)	4.57
seposita Alexander, 1941, Teucholabis (Teucholabis)	4.165
septemhastata Alexander, 1962, Tipula (Mierotipula)	4.38
septempunctata Alexander, 1939, Shannonomyia (Shannonomyia)	4.116
septemtrionalis Alexander, 1914, Rhabdomastix (Rhabdomastix) .	4.155
septemtrionis Alexander, 1940, Ozodieera (Ozodieera)	4.20
septuosa Alexander, 1962, Epiphragma (Epiphragma)	4.98
sepulchralis Alexander, 1922, Cryptolabis (Cryptolabis)	4.121
serena Alexander, 1922, Gonomyia (Progonomyia)	4.142
serotina (Alexander), 1923 (Geranomyia), Limonia (Geranomyia)	4.66
serotinellus (Alexander), 1928 (Maeromastix), Leptotarsus (Lon- gurio)	4.13
serpentina Alexander, 1938, Gonomyia (Gonomyia)	4.134



serratiloba Alexander, 1950, Limonia (Dieranomyia)	4.57
serrilobata Alexander, 1941, Tipula (Eumierotipula)	4.30
serrimarga Alexander, 1944, Pedicia (Tricyphona)	4.85
serristyla Alexander, 1945, Epiphragma (Epiphragma)	4.98
serrulatus Alexander, 1929, Molophilus (Molophilus)	4.150
serrulifera Alexander, 1949, Elephantomyia (Elephantomyia)	4.95
serrulifera Alexander, 1945, Teucholabis (Teucholabis)	4.165
serval Alexander, 1937, Tipula (Eumierotipula)	4.30
sensoris (Alexander), 1953 (Longurio), Leptotarsus (Longurio) ..	4.13
seticornis Alexander, 1946, Atarba (Isehnothrix)	4.89
setieornis Maequart, 1846 (Tipula)	4.44
setilobata Alexander, 1964, Atarba (Atarba)	4.87
setilobata Alexander, 1949, Neognophomyia	4.154
setosistylus Alexander, 1952, Molophilus (Molophilus)	4.150
setosivena Alexander, 1929, Gonomyia (Progonomyia)	4.142
setulipennis Alexander, 1928, Limonia (Zelandoglochina)	4.80
setulistyla Alexander, 1938, Elephantomyia (Elephantomyia)	4.95
severa Alexander, 1943, Ctenolimnophila (Campbellomyia)	4.93
severus Alexander, 1925, Molophilus (Molophilus)	4.150
sexeineta Alexander, 1942, Tipula (Microtipula)	4.38
shannoni (Alexander), 1914 (Rhipidia), Limonia (Rhipidia)	4.78
shannoniana (Alexander), 1929 (Holorusia), Isehnotoma (Isehno- toma)	4.10
shannoniana Alexander, 1930, Sigmatomera (Sigmatomera)	4.157
shannonina Alexander, 1929, Valdiviana	4.41
shannoninus Alexander, 1947, Molophilus (Molophilus)	4.151
Shannonomyia Alexander, 1929	4.113, 4.114
sibyllina Alexander, 1929, Limonia (Dieranomyia)	4.58
sica Alexander, 1941, Limonia (Caenoglochina)	4.48
sicarius Alexander, 1929, Molophilus (Molophilus)	4.151
Sigmatomera Osten Sacken, 1869	4.155, 4.156
sigmoidea Alexander, 1949, Teucholabis (Teucholabis)	4.165
silvai (Alexander), 1929 (Holorusia), Isehnotoma (Isehnotoma) ..	4.10
similis Williston, 1900, Braehypremna)	4.7
simillima (Alexander), 1912 (Fureomyia), Limonia (Dieranomyia)	4.58
simplex (Walker), 1856 (Ptilogyna), Ozodicera (Ozodicera)	4.20
simplex Alexander, 1945, Styringomyia	4.157
simplex (Wiedemann), 1828 (Limnobia), Teucholabis (Teuchola- bis)	4.165
simplicieornis Alexander, 1938, Limonia (Rhipidia)	4.78
simulans coneinna (Williston), 1893 (Limnobia), Limonia (Diera- nomyia)	4.58
singularis Alexander, 1944, Limonia (Caenoglochina)	4.48
singularis (Sehiner), 1868 (Paratropesa), Teucholabis (Paratro- pesa)	4.159
sinuosa (Alexander), 1926 (Holorusia), Zelandotipula	4.43
slossonae Alexander, 1914, Gonomyia (Progonomyia)	4.142
smilodon Alexander, 1940, Tipula (Microtipula)	4.38
smithi Alexander, 1912, Tipula (Microtipula)	4.38
smythiana Alexander, 1942, Limonia (Dieranomyia)	4.58
socialis Alexander, 1949, Gnophomyia	4.130
solatrix (Osten Sacken), 1859 (Limnophila), Epiphragma (Epi- phragma)	4.98
somnifera Alexander, 1943, Limonia (Caenoglochina)	4.48



songoana Alexander, 1962, Tipula (Eumierotipula)	4.30
songoensis (Alexander), 1962 (Holorusia), Zelandotipula	4.43
spora Alexander, 1944, Shannonomyia (Shannonomyia)	4.116
sordipennis Alexander, 1943, Trentepohlia (Neomongoma)	4.169
sordidipes Alexander, 1943, Cryptolabis (Cryptolabis)	4.121
sordidipes Alexander, 1962, Elephantomyia (Elephantomyia)	4.95
sordidipes Alexander, 1938, Polymera (Polymera)	4.112
soreula Alexander, 1919, Trentepohlia (Paramongoma)	4.170
sparsipila Alexander, 1962, Tipula (Microtipula)	4.38
sparipunctata Alexander, 1940, Shannonomyia (Shannonomyia) ..	4.116
sparsiseta Alexander, 1945, Neognophomyia	4.154
sparsipinus Alexander, 1952, Molophilus (Molophilus)	4.151
sparsissima (Alexander), 1929 (Limnophila), Shannonomyia (Shan- nonomyia)	4.116
sparsituber Alexander, 1962, Limonia (Dicranomyia)	4.58
spatulata Alexander, 1929, Cryptolabis (Cryptolabis)	4.122
spatulifera Alexander, 1928, Tipula (Eumierotipula)	4.30
speciosa (Alexander), 1914 (Eriocera), Hexatoma (Eriocera)	4.104
speciosa Alexander, 1953, Orimarga (Orimarga)	4.83
spectralis Alexander, 1944, Neognophomyia	4.154
sphaerulifera Edwards, 1919, Tipula (Eumierotipula)	4.23
spica Alexander, 1943, Teucholabis (Teucholabis)	4.165
splcata Alexander, 1921, Gonomyia (Lipophleps)	4.138
spleulistyla Alexander, 1950, Gonomyia (Gonomyia)	4.134
spilophaca Alexander, 1942, Ozodleera (Dihexaelonus)	4.18
spilota Wiedemann, 1828, Tipula (Eumierotipula)	4.30
spinbasis Alexander, 1943, Gnophomyia	4.130
spinicauda Alexander, 1919, Tipula (Microtipula)	4.38
spinicaudata Alexander, 1937, Austrolimnophila (Austrolimnophila)	4.91
spinigera Schiner, 1868, Teucholabis (Teucholabis)	4.165
spinistyla Alexander, 1926, Gonomyia (Paralipophleps)	4.140
spniterga Alexander, 1948, Gonomyia (Lipophleps)	4.138
spnituber Alexander, 1950, Atarba (Ischnothrix)	4.89
spinulosa Alexander, 1945, Limnophila (Limnophila)	4.108
splendida Alexander, 1913, Erioptera (Mesocyphona)	4.125
splendidula Alexander, 1930, Limonia (Dicranomyia)	4.58
sponsus Alexander, 1954, Molophilus (Molophilus)	4.151
sprucei Alexander, 1943, Limonia (Rhipidia)	4.78
squamigera Alexander, 1929, Maletta	4.143
stadelmanni Alexander, 1945, Teucholabis (Teucholabis)	4.165
<i>Stegasmonotus</i> Enderlein, 1912	4.13
steinbachl Alexander, 1946, Tipula (Eumicrotipula)	4.30
stellata Alexander, 1946, Gonomyia (Gonomyia)	4.134
stenochorema Alexander, 1962, Gnophomyia	4.130
stenoglossa Alexander, 1942, Tipula (Eumierotipula)	4.30
stenooleuca Alexander, 1956, Limonia (Geranomyia)	4.66
stenomera Alexander, 1964, Limonia (Geranomyia)	4.62
stenophallus Alexander, 1941, Gnophomyia	4.130
stenophallus Alexander, 1944, Limonia (Geranomyia)	4.66
stenophallus Alexander, 1937, Toxorhina (Toxorhina)	4.168
stenoptera Alexander, 1949, Polymera (Polymera)	4.112
stenopterus Alexander, 1935, Molophilus (Molophilus)	4.151
stenorhabda Alexander, 1960, Molophilus (Molophilus)	4.151
stenostyla (Alexander), 1953 (Longurio), Leptotarsus (Longurio)	4.13



sternohirsuta Alexander, 1941, Tipula (Microtipula)	4.38
sternolobata Alexander, 1958, Austrolimnophila (Austrolimnophila)	4.91
Stibadocerina Alexander, 1929	4.44
<i>stictica</i> (Blanchard), 1852 (Limnobia), Limonia (Dicranomyia) ..	4.56
<i>stigmatica</i> (Blanchard), 1852 (Limnobia), Limonia (Dicranomyia)	4.58
<i>stigmatica</i> (Philippi), 1865 (Limnophila), Shannonomyia (Shannonomyia)	4.116
<i>stigmosa</i> Alexander, 1930, Atarba (Atarba)	4.87
stoica Alexander, 1941, Limonia (Geranomyia)	4.66
stolida Alexander, 1934, Hexatoma (Eriocera)	4.104
stonei Alexander, 1938, Limonia (Rhipidia)	4.78
strangalia (Alexander), 1926 (Holorusia), Zelandotipula	4.43
striatipennis Alexander, 1928, Gynoplistia (Dirhipis)	4.98
striatipennis Alexander, 1941, Ozodicera (Ozodicera)	4.20
strictispina Alexander, 1968, Teucholabis (Teucholabis)	4.165
strigilis Alexander, 1926, Gonomyia (Neolipophleps)	4.139
strohmi Alexander, 1945, Ozodicera (Ozodicera)	4.20
strumosa Alexander, 1942, Teucholabis (Teucholabis)	4.165
stuardoi Alexander, 1952, Limonia (Dicranomyia)	4.58
stupens (Walker), 1861 (Limnobia), Gnophomyia	4.130
stygia (Alexander), 1927 (Erioptera), Cheilotruchia (Empeda) ...	4.120
stygia Alexander, 1914, Teucholabis (Teucholabis)	4.165
stygicornis Alexander, 1962, Limonia (Neolimnobia)	4.70
stygipes (Alexander), 1929 (Gynoplistia), Paralimnophila	4.110
stylacuta Alexander, 1947, Gnophomyia	4.131
stylifer Alexander, 1921, Molophilus (Molophilus)	4.151
stylobtusa Alexander, 1967, Limonia (Geranomyia)	4.66
<i>Styringia</i> Berendt, 1845	4.157
Styringomyia Loew, 1845	4.157
styx (Alexander), 1954 (Longurio), Leptotarsus (Longurio)	4.13
styx Alexander, 1952, Molophilus (Molophilus)	4.151
suavissima Alexander, 1951, Tipula (Eumicrotipula)	4.30
subacuminata Alexander, 1967, Limonia (Caenoglochina)	4.48
subamoena Alexander, 1942, Limonia (Peripheroptera)	4.73
subandicola Alexander, 1947, Limonia (Dicranomyia)	4.58
subandina Alexander, 1913, Limonia (Dicranomyia)	4.58
<i>subandina</i> (Alexander), 1919, Limonia (Peripheroptera)	4.72
subandina Philippi, 1865, Tipula (Eumicrotipula)	4.31
subanthracina Alexander, 1949, Teucholabis (Teucholabis)	4.165
subaperta Alexander, 1945, Atarba (Atarba)	4.85
subapicalis (Alexander), 1941 (Tanypremna), Leptotarsus (Tanypremnodes)	4.16
subapicularis Alexander, 1945, Gnophomyia	4.131
subappendiculata Alexander, 1954, Gonomyia (Gonomyia)	4.134
subappressus Alexander, 1939, Molophilus (Molophilus)	4.151
subarcuata Alexander, 1946, Gnophomyia	4.131
subargentea Alexander, 1948, Teucholabis (Teucholabis)	4.165
subarmata Alexander, 1914, Tipula (Microtipula)	4.38
subaymara Alexander, 1962, Tipula (Microtipula)	4.38
subbrevicula Alexander, 1947, Gonomyia (Gonomyia)	4.134
subcana Edwards, 1919, Tipula (Eumicrotipula)	4.31
subcollaris Alexander, 1940, Teucholabis (Paratropesa)	4.159
subconcinna Alexander, 1940, Orimarga (Diotrepha)	4.82
subcostalis (Alexander), 1922 (Rhipidia), Limonia (Rhipidia) ..	4.78



subeostata Alexander, 1919, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
subeostata Alexander, 1954, <i>Orimarga</i> (<i>Orimarga</i>)	4.83
subeynthia Alexander, 1967, <i>Erioptera</i> (<i>Mesoeyphona</i>)	4.125
subdentata Alexander, 1952, <i>Atarba</i> (<i>Atarba</i>)	4.87
subdola (Alexander), 1913 (<i>Dieranomyia</i>), <i>Limonia</i> (<i>Dieranomyia</i>)	4.58
subdomita Alexander, 1945, <i>Limonia</i> (<i>Neolimonia</i>)	4.72
subduleis Alexander, 1937, <i>Erioptera</i> (<i>Mesoeyphona</i>)	4.125
subeffeta Alexander, 1951, <i>Tipula</i> (<i>Microtipula</i>)	4.38
subenixa Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.98
subereeta Alexander, 1939, <i>Trentepohlia</i> (<i>Neomongoma</i>)	4.169
subevaneseens Alexander, 1962, <i>Braehypremna</i>	4.7
subevaneseens Alexander, 1940, <i>Erioptera</i> (<i>Mesoeyphona</i>)	4.124
subexemptus Alexander, 1968, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
subfaleatus Alexander, 1940, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
subfaleifera Alexander, 1922, <i>Gonomyia</i> (<i>Neolipophleps</i>)	4.139
subflavida Alexander, 1929, <i>Limonia</i> (<i>Dieranomyia</i>)	4.58
subflebilis Alexander, 1949, <i>Gnophomyia</i>	4.131
subfureifera (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
subfurva Alexander, 1966, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.165
subfuseata (Alexander), 1921 (<i>Limnophila</i>), <i>Paralimnophila</i>	4.110
subgalatea Alexander, 1959, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.165
subgaudens Alexander, 1941, <i>Limonia</i> (<i>Geranomyia</i>)	4.66
subglabrata Alexander, 1951, <i>Tipula</i> (<i>Eumierotipula</i>)	4.31
subgracilis Alexander, 1939, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
subhistrio Alexander, 1966, <i>Erioptera</i> (<i>Mesoeyphona</i>)	4.125
subhumilis Alexander, 1938, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.95
subhyalina Alexander, 1913, <i>Gnophomyia</i>	4.131
subinermis Alexander, 1939, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.138
subinfuseata Williston, 1896, <i>Tipula</i> (<i>Microtipula</i>)	4.38
subinsignis (Alexander), 1916 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Gerano-</i>	
<i>myia</i>)	4.66
subiratus Alexander, 1947, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
subjoeosa Alexander, 1942, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.165
sublaevis (Alexander), 1938 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
subleridensis Alexander, 1938, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.165
subleucoxena Alexander, 1940, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
sublietor Alexander, 1939, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
subligulata Alexander, 1941, <i>Tipula</i> (<i>Eumierotipula</i>)	4.31
sublima (Alexander), 1914 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
sublustralis Alexander, 1966, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.138
submolesta Alexander, 1930, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
submunda Alexander, 1941, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
submutata Alexander, 1942, <i>Limonia</i> (<i>Dieranomyia</i>)	4.58
subneglecta Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.77
subnimbifera Alexander, 1966, <i>Gnophomyia</i>	4.131
subobliterata Alexander, 1946, <i>Gnophomyia</i>	4.131
subocellata Alexander, 1964, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
subpacifera Alexander, 1942, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.91
subparilis Alexander, 1962, <i>Limonia</i> (<i>Geranomyia</i>)	4.66
subpatens Alexander, 1966, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
subpectinata (Williston), 1896 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>) ..	4.78
subpentheres Alexander, 1943, <i>Limonia</i> (<i>Geranomyia</i>)	4.66
subprotrusa Alexander, 1968, <i>Orimarga</i> (<i>Diotrepha</i>)	4.82



subpulchella Alexander, 1949, Teucholabis (Teucholabis)	4.166
subbravida (Alexander), 1928 (Dicranomyia), Limonia (Dicranomyia)	4.58
subrecisa Alexander, 1932, Limonia (Geranomyia)	4.66
subremota Alexander, 1938, Gonomyia (Gonomyia)	4.134
subreticulata Alexander, 1943, Limonia (Dicranomyia)	4.58
subrubricps Alexander, 1945, Teucholabis (Teucholabis)	4.166
subsaga Alexander, 1941, Hxatoma (Eriocera)	4.105
subsagax Alexander, 1939, Molophilus (Molophilus)	4.151
subsaturata Alexander, 1942, Gonomyia (Progonomyia)	4.142
subscimitar Alexander, 1945, Gonomyia (Lipophleps)	4.138
subserotina (Alexander), 1921 (Geranomyia), Limonia (Geranomyia)	4.66
subsimilis Alexander, 1921, Brachypremna	4.7
subsolatrix Alexander, 1939, Epiphragma (Epiphragma)	4.98
subspcciosa Alexander, 1962, Orimarga (Orimarga)	4.83
substolidia Alexander, 1939, Hxatoma (Eriocera)	4.105
substylifer Alexander, 1929, Molophilus (Molophilus)	4.151
subsuperba Alexander, 1926, Polymera (Polymera)	4.112
subtarda (Alexander), 1935 (Holorusia), Zelandotipula	4.43
subtartarus Alexander, 1946, Orimarga (Orimarga)	4.83
subtecta Alexander, 1964, Tipula (Microtipula)	4.33
subtenebricosus Alexander, 1931, Molophilus (Molophilus)	4.151
subterminalis Alexander, 1966, Aphrophila	4.118
subterminalis Alexander, 1927, Gonomyia (Lipophleps)	4.138
subterminalis (Alexander), 1921 (Rhipidia), Limonia (Rhipidia)	4.78
subtribulator Alexander, 1943, Gonomyia (Lipophleps)	4.138
subtroilus Alexander, 1966, Gonomyia (Lipophleps)	4.138
subunicolor Alexander, 1948, Gonomyia (Gonomyia)	4.134
subuniformis Alexander, 1945, Brachypremna	4.7
subvafra Alexander, 1962, Limonia (Rhipidia)	4.78
subvictor Alexander, 1948, Austrolimnophila (Limnophilella)	4.92
subvirescens Alexander, 1930, Limonia (Geranomyia)	4.66
subvirgo Alexander, 1951, Tipula (Yamatotipula)	4.40
subvittata Alexander, 1938, Ozodicera (Ozodicera)	4.20
succentiva Alexander, 1941, Limonia (Rhipidia)	4.78
suffuscipes Alexander, 1947, Trentepohlia (Paramongoma)	4.170
sulphurcllus (Alexander), 1938 (Tanypremna) Leptotarsus (Limoniodes)	4.11
sultana Alexander, 1938, Teucholabis (Teucholabis)	4.166
sumptuosa Alexander, 1942, Limonia (Geranomyia)	4.66
superarmata Alexander, 1942, Limonia (Rhipidia)	4.79
superarmata Alexander, 1942, Ozodicera (Dihexaclonus)	4.18
superba Alexander, 1913, Polymera (Polymera)	4.112
supernumeraria Alexander, 1921, Elephantomyia (Elephantomyia)	4.95
superstes Alexander, 1942, Toxorhina (Ceratocheilus)	4.167
supplementa Alexander, 1945, Pseudolimnophila	4.113
supplicata Alexander, 1943, Atarba (Ischnothrix)	4.89
surinamensis Alexander, 1947, Erioptera (Mesocophona)	4.125
surinamica Alexander, 1946, Limonia (Rhipidia)	4.79
susurra Alexander, 1945, Erioptera (Erioptera)	4.123
sustrina (Alexander), 1941 (Erioptera), Cheilotrichia (Empeda) ..	4.120
sybarita Alexander, 1945, Limonia (Rhipidia)	4.79
sybaritica Alexander, 1947, Epiphragma (Epiphragma)	4.98



<i>sycopnanta</i> Alexander, 1945, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>sylvania</i> Alexander, 1939, <i>Limonia</i> (<i>Geranomyia</i>)	4.66
<i>Sympleeta</i> Meigen, 1830	4.126
<i>synebroa</i> Alexander, 1929, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>synelera</i> Alexander, 1928, <i>Rhabdomastix</i> (<i>Sacandaga</i>)	4.155
<i>syndactyla</i> Alexander, 1942, <i>Orimarga</i> (<i>Diotrepha</i>)	4.82
<i>synempora</i> Alexander, 1929, <i>Valdiviana</i>	4.41
<i>tabescens</i> (Enderlein), 1912 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>tabida</i> (Enderlein), 1912 (<i>Tipula</i>)	4.44
<i>taciturna</i> Alexander, 1945, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.122
<i>tactilis</i> Alexander, 1952, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
<i>taenioptera</i> (Wiedemann), 1828 (<i>Limnobia</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>tafiensis</i> Alexander, 1962, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.138
<i>taino</i> Alexander, 1964, <i>Teucholabis</i> (<i>Teueholabis</i>)	4.166
<i>talamancana</i> Alexander, 1939, <i>Teucholabis</i> (<i>Teueholabis</i>)	4.166
<i>talamancensis</i> Alexander, 1947, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>tancitaro</i> Alexander, 1946, <i>Tipula</i> (<i>Microtipula</i>)	4.38
<i>tantilla</i> Alexander, 1916, <i>Erioptera</i> (<i>Mesoeyphona</i>)	4.124
<i>tantilla</i> Alexander, 1938, <i>Rhabdomastix</i> (<i>Rhabdomastix</i>)	4.155
<i>tanymetra</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>Tanypremna</i> Osten Sacken, 1886	4.13
<i>Tanypremnella</i> Alexander, 1938	4.15
<i>Tanypremnodes</i> Alexander, 1924	4.16
<i>tarda</i> (Alexander), 1935 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
<i>tarmae</i> Alexander, 1943, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>tarsalba</i> Alexander, 1928, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.95
<i>tartarus</i> Alexander, 1946, <i>Orimarga</i> (<i>Orimarga</i>)	4.83
<i>tartarus</i> Alexander, 1949, <i>Teucholabis</i> (<i>Teueholabis</i>)	4.166
<i>Tasiocera</i> Skuse, 1890	4.157
<i>tasioceroides</i> Alexander, 1948, <i>Polymera</i> (<i>Polymerodes</i>)	4.113
<i>tatei</i> Alexander, 1929, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>tatei</i> Alexander, 1931, <i>Limonia</i> (<i>Geranomyia</i>)	4.66
<i>tau</i> Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.26
<i>taurus</i> Alexander, 1914, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>tecta</i> Alexander, 1926, <i>Tipula</i> (<i>Microtipula</i>)	4.33
<i>tehueche</i> Alexander, 1929, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.81
<i>tehueche</i> Alexander, 1968, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>tehueche</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>teinoterga</i> Alexander, 1967, <i>Limonia</i> (<i>Dicranomyia</i>)	4.58
<i>teleneura</i> Alexander, 1944, <i>Gnophomyia</i>	4.131
<i>taurus</i> Alexander, 1914, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>temperata</i> Alexander, 1940, <i>Tipula</i> (<i>Microtipula</i>)	4.38
<i>tempestiva</i> (Alexander), 1943 (<i>Gnophomyia</i>), <i>Eugnophomyia</i>	4.127
<i>tenebricosus</i> Alexander, 1916, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>tenebrosa</i> (Walker), 1856 (<i>Limnobia</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>tenella</i> (Philippi), 1865 (<i>Polymoria</i>)	4.117
<i>tentator</i> Alexander, 1947, <i>Molophilus</i> (<i>Trichomolophilus</i>)	4.152
<i>tenuieineta</i> Alexander, 1921, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.122
<i>tenuicula</i> Enderlein, 1912, <i>Tipula</i> (<i>Microtipula</i>)	4.38
<i>tenuilobata</i> Alexander, 1942, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.91
<i>tenuilobata</i> Alexander, 1940, <i>Tipula</i> (<i>Microtipula</i>)	4.38
<i>tenuipalpis</i> Alexander, 1928, <i>Limonia</i> (<i>Zelandoglochina</i>)	4.81
<i>tenuipes</i> (Say), 1823 (<i>Limnobia</i>), <i>Pilaria</i>	4.110
<i>tenuis</i> (Wulp), 1885 (<i>Tipula</i>), <i>Megistocera</i>	4.16
<i>tenuissima</i> (Alexander), 1929 (<i>Ischnothrix</i>), <i>Atarba</i> (<i>Ischnothrix</i>)	4.89




<i>tenuissima</i> Alexander, 1944, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.95
<i>tcphronota</i> Alexander, 1942, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>tergata</i> (Alexander), 1926 (<i>Rhaphidolabis</i>), <i>Dicranota</i> (<i>Plectromyia</i>)	4.84
<i>tergifera</i> Alexander, 1953, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.91
<i>tergofimbriata</i> Alexander, 1928, <i>Gonomyia</i> (<i>Lipohleps</i>)	4.138
<i>tergospinosus</i> Alexander, 1966, <i>Molophilus</i> (<i>Molophilus</i>)	4.151
<i>ternarius</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.152
<i>terpischore</i> Alexander, 1951, <i>Tipula</i> (<i>Microtipula</i>)	4.39
<i>terribilis</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.39
<i>terrifica</i> Alexander, 1942, <i>Ozodicera</i> (<i>Dihexaclonus</i>)	4.18
<i>tersa</i> Alexander, 1949, <i>Gonomyia</i> (<i>Lipohleps</i>)	4.138
<i>tersa</i> Alexander, 1928, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>tersoides</i> Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>tesselata</i> Alexander, 1943, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>testacea</i> (Philippi), 1865 (<i>Plettusa</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.67
<i>tethys</i> Alexander, 1949, <i>Trentepohlia</i> (<i>Paramongoma</i>)	4.170
<i>tetracaena</i> Alexander, 1966, <i>Gnophomyia</i>	4.131
<i>tetracantha</i> Alexander, 1945, <i>Atarba</i> (<i>Atarba</i>)	4.87
<i>tetracanthus</i> Alexander, 1929, <i>Molophilus</i> (<i>Molophilus</i>)	4.152
<i>tetracradus</i> Alexander, 1967, <i>Helius</i>	4.47
<i>tetraleuca</i> Alexander, 1937, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>Teucholabis</i> Osten Sacken, 1859	4.157, 4.159
<i>teucholaboides</i> (Alexander), 1913 (<i>Peripheroptera</i>), <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
<i>textrina</i> Alexander, 1967, <i>Limonia</i> (<i>Neolimnobia</i>)	4.70
<i>thalia</i> Alexander, 1945, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.125
<i>thalia</i> Alexander, 1945, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>Thambeta</i> Williston, 1896	4.81
<i>thamia</i> Alexander, 1950, <i>Atarba</i> (<i>Atarba</i>)	4.86
<i>thamyris</i> Alexander, 1950, <i>Limonia</i> (<i>Dicranomyia</i>)	4.58
<i>thaumasta</i> Alexander, 1953, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.20
<i>thaumastopodus</i> Alexander, 1913, <i>Molophilus</i> (<i>Molophilus</i>)	4.143
<i>theobromina</i> Edwards, 1919, <i>Tipula</i> (<i>Bellardina</i>)	4.21
<i>theresiae</i> Alexander, 1945, <i>Limnophila</i> (<i>Limnophila</i>)	4.108
<i>thioptera</i> Alexander, 1941, <i>Limonia</i> (<i>Peripheroptera</i>)	4.73
<i>thiosema</i> Alexander, 1921, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>thixis</i> Alexander, 1967, <i>Limonia</i> (<i>Dicranomyia</i>)	4.58
<i>tholopa</i> (Alexander), 1931 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>thoracica</i> Alexander, 1913, <i>Polymera</i> (<i>Polymera</i>)	4.112
<i>thyestes</i> Alexander, 1953, <i>Brachypremna</i>	4.7
<i>thysbe</i> Alexander, 1942, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>tibialis</i> (Loew), 1851 (<i>Aporosa</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.66
<i>tibialis</i> Alexander, 1922, <i>Polymera</i> (<i>Polymera</i>)	4.112
<i>tigriventris</i> Alexander, 1942, <i>Elephantomyia</i> (<i>Elephantomyia</i>) ..	4.95
<i>tijucanus</i> (Alexander), 1943 (<i>Macromastix</i>), <i>Leptotarsus</i> (<i>Longurio</i>)	4.13
<i>tijucensis</i> Alexander, 1943, <i>Tipula</i> (<i>Microtipula</i>)	4.39
<i>timens</i> Alexander, 1950, <i>Limonia</i> (<i>Geranomyia</i>)	4.67
<i>tinguana</i> Alexander, 1943, <i>Orimarga</i> (<i>Orimarga</i>)	4.83
<i>Tipula</i> Linnaeus, 1758	4.20
<i>Tipulinae</i>	4.5
<i>tirsias</i> Alexander, 1949, <i>Gnophomyia</i>	4.131
<i>tiresias</i> Alexander, 1950, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>titan</i> Alexander, 1928, <i>Molophilus</i> (<i>Molophilus</i>)	4.152

titicacae (Alexander), 1944 (Pectinotipula), Tipula (Pectinotipula)	4.39
titicacana Alexander, 1945, Limonia (Dieranomyia)	4.58
toleranda Alexander, 1954, Gnophomyia	4.131
tolimensis Alexander, 1941, Tipula (Eumicrotipula)	4.30
Tonnoiroomyia Alexander, 1926	4.83
topoensis Alexander, 1952, Gonomyia (Neolipophleps)	4.139
topoensis Alexander, 1951, Tipula (Microtipula)	4.39
torrida Alexander, 1932, Limonia (Dieranomyia)	4.51
torticornis Alexander, 1928, Limonia (Zelandoglochina)	4.81
tortilis Alexander, 1968, Paralimnophila	4.110
torulosa Alexander, 1968, Limonia (Dicranomyia)	4.59
tovarensis Alexander, 1944, Limonia (Rhipidia)	4.74
tovarensis Alexander, 1947, Tipula (Eumicrotipula)	4.31
townsendi (Alexander), 1914 (Eriocera), Hexatoma (Eriocera)	4.105
townsendi (Alexander), 1916 (Geranomyia), Limonia (Geranomyia)	4.67
Toxorhina Loew, 1850	4.166, 4.167
<i>Toxorhina</i> , emend.	4.166
tragica Alexander, 1946, Limonia (Dicranomyia)	4.59
tranquilla (Alexander), 1922 (Eriocera), Hexatoma (Eriocera)	4.105
transfasciatus (Alexander), 1938 (Tanyremna), Leptotarsus (Tanyremnella)	4.16
transitoria Alexander, 1941, Limonia (Geranomyia)	4.67
transluceida (Alexander), 1912 (Furcomyia), Limonia (Caenolimonia)	4.49
transversa Alexander, 1949, Gnophomyia	4.131
travassosanus (Alexander), 1942 (Macromastix), Leptotarsus (Longurio)	4.13
travassosi Alexander, 1945, Cryptolabis (Cryptolabis)	4.122
travassosi Alexander, 1943, Orimarga (Diotrepha)	4.82
tremula Alexander, 1929, Austrolimnophila (Austrolimnophila)	4.91
Trentepohlia Bigot, 1854	4.168
triaeculeata Alexander, 1938, Gonomyia (Gonomyia)	4.134
trialbocincta Alexander, 1941, Limonia (Neoglochina)	4.69
trialbosignata Alexander, 1941, Hexatoma (Eriocera)	4.105
triangularis Alexander, 1945, Brachypremna	4.6
triangularis Alexander, 1945, Erioptera (Mesocyphona)	4.125
triangularis (Alexander), 1927 (Pilaria), Shannonomyia (Shannonomyia)	4.117
triangulifera (Alexander), 1921 (Psaronius), Lecteria (Psaronius)	4.106
triatra (Alexander), 1962 (Holorusia), Zelandotipula	4.43
triatrata Alexander, 1968, Gnophomyia	4.131
tribulator Alexander, 1942, Gonomyia (Lipophleps)	4.138
triceps Alexander, 1947, Gnophomyia	4.131
tricepoides Alexander, 1966, Gnophomyia	4.131
trichomera Alexander, 1939, Limonia (Geranomyia)	4.67
Trichomolophilus Alexander, 1936	4.152
trichoneura (Alexander), 1962 (Holorusia), Zelandotipula	4.43
trichoprocta Alexander, 1945, Tipula (Microtipula)	4.39
<i>trichopus</i> (Philippi), 1865 (Limnophila), Erioptera (Trimicra)	4.126
Trichotipula Alexander, 1915	4.40
tricineta (Alexander), 1913 (Dicranomyia), Limonia (Neolimnobia)	4.70
tricornis Alexander, 1949, Gnophomyia	4.131
<i>tricornis</i> (Alexander), 1925 (Dicranomyia), Limonia (Dicranomyia)	4.52



Tricyphona Zetterstedt, 1837	4.84
tridens Alexander, 1952, Molophilus (Molophilus)	4.152
tridentata (Alexander), 1925 (Erioptera), Cheilitrichia (Empeda)	4.120
tridenticulata Alexander, 1967, Cryptolabis (Cryptolabis)	4.122
tridigitata Alexander, 1943, Limonia (Rhipidia)	4.79
tridigitatus Alexander, 1947, Molophilus (Molophilus)	4.152
triemarginata Alexander, 1935, Tipula (Eumicrotipula)	4.31
trifasciata (Röder), 1885 (Eriocera), Hexatoma (Eriocera)	4.105
trifasciata Enderlein, 1912, Teucholabis (Teucholabis)	4.166
trifibra Alexander, 1954, Molophilus (Molophilus)	4.152
triguttata Alexander, 1921, Ozodicera (Dihexaclonus)	4.18
trihastata Alexander, 1945, Tipula (Microtipula)	4.39
trilobata Alexander, 1938, Toxorhina (Toxorhina)	4.168
trilobifera Alexander, 1967, Limonia (Dicranomyia)	4.59
trimaculata Macquart, 1838, (Tipula)	4.44
trimedia Alexander, 1966, Maietta	4.143
trimclania Alexander, 1942, Limonia (Peripheroptera)	4.74
Trimicra Osten Sacken, 1861	4.126
trina Wiedemann, 1824, (Tipula)	4.44
trinidadensis (Alexander), 1912 (Pachyrhina), Tipula (Microtipula)	4.39
trinigrina Alexander, 1940, Limonia (Peripheroptera)	4.74
trinitatis Alexander, 1931, Limonia (Dicranomyia)	4.59
trinitatis Alexander, 1927, Neognophomyia	4.154
trinitatis Alexander, 1941, Tipula (Microtipula)	4.39
trionyx Alexander, 1949, Gonomyia (Lipophleps)	4.138
tripallens Alexander, 1942, Ozodicera (Dihexaclonus)	4.18
triparcus Alexander, 1944, Molophilus (Molophilus)	4.152
tripcetinata Alexander, 1931, Limonia (Rhipidia)	4.79
Triplicitipula Alexander, 1965	4.40
trisetigera Alexander, 1949, Gnophomyia	4.131
trispilota Alexander, 1942, Tipula (Eumicrotipula)	4.31
trispinifer Alexander, 1940, Ozodicera (Ozodicera)	4.20
trispinigera Alexander, 1945, Orimarga (Orimarga)	4.83
trispinosa Alexander, 1921, Gonomyia (Neolipophleps)	4.139
tristella Alexander, 1929, Limonia (Geranomyia)	4.67
tristillata Alexander, 1929, Gynoplistia (Gynoplistia)	4.99
tristillata Alexander, 1929, Tipula (Eumicrotipula)	4.31
tristis (Loew), 1851 (Aporosa), Limonia (Geranomyia)	4.67
tristis Alexander, 1913, Teucholabis (Teucholabis)	4.166
trituberculata Alexander, 1929, Limonia (Dicranomyia)	4.59
Trochobola Osten Sacken, 1869	4.59
trogodyta Edwards, 1918, Erioptera (Mesocyphona)	4.125
troglophila Alexander, 1929, Limonia (Dicranomyia)	4.59
troilus Alexander, 1949, Gonomyia (Lipophleps)	4.138
tropicalis Alexander, 1913, Cryptolabis (Cryptolabis)	4.122
trunculata (Alexander), 1962 (Holorusia), Ischnotoma (Ischnotoma)	4.9
tuber Alexander, 1944, Gnophomyia	4.131
tuber (Alexander), 1953 (Limnophila), Shannonomyia (Shannonomyia)	4.117
tuberculata Alexander, 1925, Gonomyia (Gonomyia)	4.133
tuberculifera Alexander, 1943, Atarba (Atarba)	4.87
tuberculifera (Alexander), 1953 (Holorusia), Zelandotipula	4.43
tucumana (Alexander), 1946 (Pectinotipula), Tipula (Pectinotipula)	4.39



tucumana Alexander, 1936, Trentepohlia (Paramongoma)	4.170
tucumanensis Alexander, 1945, Tipula (Microtipula)	4.39
tucumanus Alexander, 1929, Molophilus (Molophilus)	4.152
tullochi Alexander, 1945, Teucholabis (Teucholabis)	4.166
tulumayoensis Alexander, 1944, Limonia (Geranomyia)	4.67
tumidibasis Alexander, 1938, Limonia (Geranomyia)	4.67
tungurahuana Alexander, 1945, Gnophomyia	4.131
tunguraguana Alexander, 1940, Tipula (Eumicrotipula)	4.31
tunguraguensis Alexander, 1940, Austrolimnophila (Austrolimnophila)	4.91
tungurahuensis Alexander, 1945, Atarba (Atarba)	4.88
turbida (Alexander), 1928 (Geranomyia), Limonia (Geranomyia)	4.67
turrialbae Alexander, 1945, Erioptera (Mesocyphona)	4.125
turrialbensis Alexander, 1945, Teucholabis (Teucholabis)	4.166
turrifera Alexander, 1931, Limonia (Rhipidia)	4.75
turritella Alexander, 1943, Limonia (Rhipidia)	4.79
tyranna Alexander, 1946, Tipula (Eumicrotipula)	4.31
uberis Alexander, 1939, Limonia (Geranomyia)	4.67
umbra Alexander, 1948, Shannonomyia (Shannonomyia)	4.117
umbrata (Meijere), 1911 (Dicranomyia), Limonia (Atypophthalmus)	4.47
umbricolor Alexander, 1937, Limonia (Geranomyia)	4.67
umbrifera Alexander, 1937, Ozodicera (Dihexaelonus)	4.18
umbrosa Alexander, 1938, Cryptolabis (Cryptolabis)	4.122
undulata Bellardi, 1861, Limnophila (Limnophila)	4.108
uniatra (Alexander), 1945 (Holorusia), Zelandotipula	4.43
unicincta Alexander, 1962, Elephantomyia (Elephantomyia)	4.95
unicingulata Alexander, 1944, Teucholabis (Teucholabis)	4.166
unicolor Osten Sacken, 1887, Brachypremna	4.7
unicolor Alexander, 1913, Gonomyia (Gonomyia)	4.134
unidentata (Alexander), 1925 (Erioptera), Cheilotrichia (Empeda)	4.120
uniformis Alexander, 1920, Brachypremna	4.7
uniformis (Alexander), 1914 (Rhamphidia), Helius	4.47
uniformis (Blanchard), 1852 (Erioptera), Molophilus (Molophilus)	4.152
uniguttatus (Alexander), 1953 (Longurio), Leptotarsus (Tanypremna)	4.15
unilateralis Alexander, 1931, Atarba (Atarba)	4.88
unipectinata (Williston), 1896 (Rhipidia), Limonia (Rhipidia) ..	4.79
unipuncta Alexander, 1944, Rhabdomastix (Rhabdomastix)	4.155
unipunctata Alexander, 1921, Polymera (Polymera)	4.112
unispinifera Alexander, 1938, Limonia (Geranomyia)	4.67
unistriata Alexander, 1941, Tipula (Eumicrotipula)	4.31
urania Alexander, 1944, Erioptera (Erioptera)	4.123
urophora Alexander, 1938, Tipula (Microtipula)	4.39
ursula Alexander, 1951, Tipula (Eumicrotipula)	4.31
usta (Osten Sacken), 1886 (Pachyrrhina), Nephrotoma	4.17
uxor Alexander, 1942, Limonia (Rhipidia)	4.79
uxoria Alexander, 1946, Tipula (Trichotipula)	4.40
vafra Alexander, 1941, Limonia (Rhipidia)	4.79
valdiviana Philippi, 1865, Tipula (Eumicrotipula)	4.31
Valdiviana Alexander, 1929	4.41
valida (Loew), 1851 (Aporosa), Limonia (Geranomyia)	4.67



<i>vallicola</i> Alexander, 1947, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.122
<i>vana</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.57
<i>vanderwulpi</i> (Alexander), 1912 (<i>Tipula</i>), <i>Zelandotipula</i>	4.42
<i>varia</i> (Wiedemann), 1828 (<i>Limnobia</i>), <i>Epiphragma</i> (<i>Epiphragma</i>)	4.98
<i>variata</i> Alexander, 1931, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
<i>variatus</i> Alexander, 1952, <i>Molophilus</i> (<i>Molophilus</i>)	4.152
<i>varicornis</i> Alexander, 1913, <i>Atarba</i> (<i>Atarba</i>)	4.88
<i>varicornis</i> Alexander, 1936, <i>Sigmatomera</i> (<i>Sigmatomera</i>)	4.157
<i>variegata</i> Alexander, 1937, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>variegata</i> (Walker), 1836 (<i>Geranomyia</i>), <i>Limonia</i> (<i>Geranomyia</i>) ..	4.67
<i>variicalcarata</i> Alexander, 1929, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
<i>variicosta</i> Alexander, 1934, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>variispina</i> Alexander, 1938, <i>Atarba</i> (<i>Atarba</i>)	4.88
<i>variitibia</i> Alexander, 1936, <i>Brachypremna</i>	4.7
<i>varineura</i> (Bigot), 1888 (<i>Nephrotoma</i>), <i>Tipula</i> (<i>Eumicrotipula</i>) ..	4.31
<i>varipes</i> Alexander, 1939, <i>Cryptolabis</i> (<i>Cryptolabis</i>)	4.122
<i>varipes</i> Alexander, 1929, <i>Gynoplistia</i> (<i>Gynoplistia</i>)	4.99
<i>varitarsis</i> Alexander, 1929, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.91
<i>vehemens</i> Alexander, 1949, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.138
<i>velasteguii</i> Alexander, 1938, <i>Limonia</i> (<i>Neolimonia</i>)	4.72
<i>velutina</i> Alexander, 1916, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>venatrix</i> Alexander, 1952, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>venavitta</i> Alexander, 1940, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>veneris</i> Alexander, 1952, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>venezolanus</i> Alexander, 1950, <i>Protohelius</i>	4.83
<i>venezuelensis</i> (Alexander), 1944 (<i>Pselliophora</i>), <i>Ctenophora</i> (<i>Psel-</i> <i>liophora</i>)	4.7
<i>venezuelensis</i> (Macquart), 1846 (<i>Limnobia</i>), <i>Teucholabis</i> (<i>Teuchola-</i> <i>bis</i>)	4.166
<i>venezuolana</i> Alexander, 1950, <i>Paradelphomyia</i> (<i>Oxyrhiza</i>)	4.110
<i>venosa</i> (Philippi), 1865 (<i>Limnophila</i>)	4.117
<i>venustior</i> Alexander, 1950, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>venustipes</i> Alexander, 1926, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.125
<i>verecunda</i> (Philippi), 1865 (<i>Limnophila</i>)	4.117
<i>vernalis</i> (Philippi), 1865 (<i>Limnobia</i>), <i>Limonia</i> (<i>Dicranomyia</i>) ..	4.59
<i>versuta</i> Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.67
<i>verticillata</i> Alexander, 1948, <i>Polymera</i> (<i>Polymera</i>)	4.113
<i>vicina</i> (Alexander), 1916 (<i>Rhipidia</i>), <i>Limonia</i> (<i>Rhipidia</i>)	4.76
<i>victor</i> (Alexander), 1919 (<i>Limnophila</i>), <i>Austrolimnophila</i> (<i>Limno-</i> <i>philella</i>)	4.92
<i>vilis</i> Alexander, 1931, <i>Gnophomyia</i>	4.131
<i>villaricae</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>villaricensis</i> Alexander, 1929, <i>Limonia</i> (<i>Geranomyia</i>)	4.67
<i>vindex</i> Alexander, 1941, <i>Gonomyia</i> (<i>Lipophleps</i>)	4.139
<i>vindicta</i> Alexander, 1943, <i>Limonia</i> (<i>Geranomyia</i>)	4.67
<i>violaceipennis</i> Alexander, 1937, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
<i>virescens</i> (Loew), 1851 (<i>Aporosa</i>), <i>Limonia</i> (<i>Geranomyia</i>)	4.68
<i>virescens</i> (Philippi), 1865 (<i>Plettusa</i>), <i>Limonia</i> (<i>Geranomyia</i>) ...	4.67
<i>virgilia</i> Alexander, 1951, <i>Tipula</i> (<i>Microtipula</i>)	4.39
<i>virgo</i> Osten Sacken, 1886, <i>Tipula</i> (<i>Yamatotipula</i>)	4.40
<i>virgulata</i> Williston, 1900, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.40
<i>virgulativentris</i> (Enderlein), 1912 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>)	4.31
<i>viridella</i> Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.105
	4.68



<i>viridinervis</i> Alexander, 1934, <i>Aphrophila</i>	4.118
<i>viridula</i> Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.64
<i>virilis</i> (Alexander), 1916 (<i>Dicranomyia</i>), <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>vitripennis</i> Alexander, 1942, <i>Gnophomyia</i>	4.131
<i>vittigera</i> (Philippi), 1865 (<i>Tipula</i>), <i>Ischnotoma</i> (<i>Ischnotoma</i>)	4.10
<i>vittinervis</i> Alexander, 1937, <i>Dolichocheza</i> (<i>Mcgistomastix</i>)	4.8
<i>vittinervis</i> Alexander, 1936, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>vivasberthieri</i> Alexander, 1938, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>)	4.92
<i>vivasberthieri</i> (Alexander), 1944 (<i>Gnophomyia</i>), <i>Eugnophomyia</i> ..	4.127
<i>vivasberthieri</i> Alexander, 1940, <i>Limonia</i> (<i>Peripheroptera</i>)	4.74
<i>vivida</i> (Alexander), 1941 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
<i>vocator</i> Alexander, 1944, <i>Shannonomyia</i> (<i>Shannonomyia</i>)	4.117
<i>volens</i> Alexander, 1944, <i>Tipula</i> (<i>Microtipula</i>)	4.39
<i>volentis</i> Alexander, 1952, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
<i>voracis</i> Alexander, 1948, <i>Atarba</i> (<i>Ischnothrix</i>)	4.89
<i>vorax</i> Alexander, 1941, <i>Limonia</i> (<i>Caenoglochina</i>)	4.48
<i>votiva</i> Alexander, 1944, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>vulpes</i> (Alexander), 1945 (<i>Holorusia</i>), <i>Zelandotipula</i>	4.43
<i>vulsa</i> Alexander, 1952, <i>Toxorhina</i> (<i>Ceratocheilus</i>)	4.167
<i>vultuosa</i> Alexander, 1946, <i>Tipula</i> (<i>Trichotipula</i>)	4.40
<i>walkeri</i> Alexander, 1930, <i>Limonia</i> (<i>Geranomyia</i>)	4.68
<i>walkeri</i> Alexander, 1931, <i>Molophilus</i> (<i>Molophilus</i>)	4.152
<i>weberi</i> Alexander, 1951, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>weiseri</i> Alexander, 1920, <i>Gonomyia</i> (<i>Progonomyia</i>)	4.142
<i>weiseriana</i> Alexander, 1929, <i>Limonia</i> (<i>Dicranomyia</i>)	4.59
<i>westwoodi</i> Brunetti, 1920, <i>Toxorhina</i> (<i>Toxorhina</i>)	4.168
<i>westwoodi antillarum</i> Alexander, 1932, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.95
<i>wetmorcana</i> Alexander, 1947, <i>Tipula</i> (<i>Bellardina</i>)	4.21
<i>whitei</i> Alexander, 1930, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.125
<i>wiedemanni</i> Alexander, 1933, <i>Hexatoma</i> (<i>Eriocera</i>)	4.105
<i>willistoni</i> (Alexander), 1913 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.105
<i>wighti</i> Alexander, 1939, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
<i>wilhelminae</i> Alexander, 1947, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
<i>williamsoni</i> Alexander, 1912, <i>Brachypremna</i>	4.7
<i>williamsoni</i> (Alexander), 1923 (<i>Eriocera</i>), <i>Hexatoma</i> (<i>Eriocera</i>) ..	4.105
<i>willinki</i> Alexander, 1962, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.31
<i>willistoni</i> Alexander, <i>Elephantomyia</i> (<i>Elephantomyia</i>)	4.93
<i>willistoniana</i> Alexander, 1929, <i>Limonia</i> (<i>Rhipidia</i>)	4.79
<i>winthemi</i> Alexander, 1922, <i>Erioptera</i> (<i>Psiloeonopa</i>)	4.126
<i>withycombei</i> Alexander, 1929, <i>Erioptera</i> (<i>Mesocyphona</i>)	4.125
<i>witteana</i> Alexander, 1942, <i>Ozodicera</i> (<i>Ozodicera</i>)	4.20
<i>wittei</i> Alexander, 1920, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.32
<i>wolffhuegeli</i> Alexander, 1939, <i>Limnophila</i> (<i>Araucolimnophila</i>)	4.107
<i>woytkowskiana</i> Alexander, 1941, <i>Sigmatomera</i> (<i>Austrolimnobia</i>)	4.156
<i>woytkowskiana</i> Alexander, 1942, <i>Tipula</i> (<i>Eumicrotipula</i>)	4.32
<i>wygodzinskyi</i> Alexander, 1948, <i>Austrolimnophila</i> (<i>Austrolimnophila</i>) ..	4.92
<i>wygodzinskyi</i> Alexander, 1962, <i>Gonomyia</i> (<i>Gonomyia</i>)	4.134
<i>xantha</i> Alexander, 1952, <i>Teucholabis</i> (<i>Teucholabis</i>)	4.166
<i>xanthomela</i> Alexander, 1939, <i>Epiphragma</i> (<i>Epiphragma</i>)	4.98

xanthomela Alexander, 1962, Limonia (Caenolimonia)	4.49
xanthoplaea (Alexander), 1921 (Geranomyia), Limonia (Geranomyia)	4.68
xanthoptera Alexander, 1929, Austrolimnophila (Austrolimnophila)	4.92
xanthostoma Loew, 1851, Ozodieera (Dihexaelonus)	4.18
xantieles Alexander, 1946, Teucholabis (Teucholabis)	4.166
Xenoelephantomyia Alexander, 1965	4.95
Xenotipula Alexander, 1921	4.16
xystophanes (Alexander), 1921 (Paratropeza), Teucholabis (Euparatropesa)	4.158
Yamatotipula Matsumura, 1916	4.40
yanamontea Alexander, 1945, Tipula (Eumicrotipula)	4.32
yungasensis Alexander, 1962, Tipula (Eumicrotipula)	4.32
yungasicola (Alexander), 1962 (Holorusia), Zelandotipula	4.43
yunqueana Alexander, 1952, Limonia (Dieranomyia)	4.59
yunquensis Alexander, 1956, Limonia (Geranomyia)	4.68
yunquensis Alexander, 1952, Molophilus (Molophilus)	4.152
Zalusa Enderlein, 1906	4.79
zamorae (Alexander), 1945 (Holorusia), Zelandotipula	4.43
Zelandoglochina Alexander, 1924	4.80
Zelandotipula Alexander, 1922	4.41
zeltale Alexander, 1928, Tipula (Eumicrotipula)	4.32
zernyana Alexander, 1949, Shannonomyia (Shannonomyia)	4.117
zernyi Alexander, 1942, Trentepohlia (Neomongoma)	4.169
zeteki Alexander, 1945, Tipula (Microtipula)	4.39
zeugmata Alexander, 1951, Tipula (Eumicrotipula)	4.32
zikanellus (Alexander), 1953 (Longurio), Leptotarsus (Longurio)	4.13
zikani (Alexander), 1936 (Holorusia), Ischnotoma (Icriomastax)	4.9
zikaniana Alexander, 1937, Ozodieera (Ozodicera)	4.20
zonalis Alexander, 1926, Tipula (Microtipula)	4.39
zonata (Osten Sacken), 1886 (Eriocera), Hexatoma (Eriocera) ..	4.105
zotzil Alexander, 1928, Tipula (Eumicrotipula)	4.32





SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

5

FAMILY TANYDERIDAE

CHARLES P. ALEXANDER AND MABEL M. ALEXANDER

Amherst, Massachusetts

The family Tanyderidae includes 36 species distributed in 11 genera, of which *Macrochile* Loew (type, *spectrum* Loew) is known as fossil in the Baltic Amber (Upper Eocene). The remaining genera, with the number of species, are *Protoplasa* Osten Sacken (1), eastern Nearctic; *Tanyderus* Philippi (1), *Araucoderus* Alexander (1), and *Neoderus* Alexander (1), southern Neotropical; *Peringueyomyia* Alexander (1), southern Ethiopian; *Radinocerus* Handlirsch (11), *Mischoderus* Handlirsch (5), *Nothoderus* Alexander (1), and *Eutanyderus* Alexander (1), Australasian, including New Zealand (*Mischoderus*); and *Protanyderus* Handlirsch, with 3 species in the western Nearctic and 8 in the eastern Palaearctic regions.

The immature stages of four of the genera now are known and all conform closely to a uniform type, both in the larvae and pupae [*Protoplasa fitchii* (Osten Sacken): Alexander, 1930b (larva, pupa), Crampton, 1930a (pupa), 1930b (larva); *Protanyderus margarita* Alexander: Knight, 1963 (larva), 1964 (pupa); *Protanyderus vipio* (Osten Sacken): Rose, 1963 (larva); *Peringueyomyia barnardi* Alexander: Wood, 1952 (larva, pupa); *Eutanyderus wilsoni* Alexander: Hinton, 1966 (pupal spiracular gills)]. The early stages are aquatic or virtually so, pupation taking place in the sandy soil along stream margins. The three Neotropical genera are among the rarest of all known Diptera and the discovery of their early stages, particularly those of the type of the family, *Tanyderus*, would be of particular interest.

Genus TANYDERUS Philippi

TANYDERUS Philippi, 1865:780. Type-species, *pictus* Philippi (mon.).*PICTUS* Philippi, 1865:781, fig. 57 with subfigs. (entire insect, wing, head).

Type-locality: Chile, Valdivia, Los Ulmos. Distr. — Chile (Concepción, south to Llanquihue; Arauco; Valdivia [Alexander, 1935]). Refs. — Alexander, 1927 (venation); Williams, 1933: fig. 34 (wing). Type apparently lost.

Genus ARAUCODERUS Alexander

ARAUCODERUS Alexander, 1929:228. Type-species, *Tanyderus gloriosus* Alexander (mon.).

GLORIOSUS (Alexander), 1920:5, fig. (wing) (*Tanyderus*). Type-locality: Chile, Rio Aysén. Distr. — Chile, Argentina (Tucumán). Refs. — Alexander, 1927:9, fig. 6 (venation), 1929:228. Type in Riksmuseum, Stockholm.

Genus NEODERUS Alexander

Tanyderus, subg. NEODERUS Alexander, 1927:10. Type-species, *patagonicus* Alexander (mon.).

Ref. — Alexander, 1928:228 (as genus).

PATAGONICUS (Alexander), 1913:331, figs. 1, 2 (head), 3 (wing) (*Tanyderus*).

Type-locality: Argentina, Patagonia, Latitude Cove. Distr. — Argentina. Ref. — Alexander, 1927:10, fig. 8 (venation), 19 (head). Type in U.S. National Museum, n° 14919.

LITERATURE CITED

ALEXANDER, C. P.

1913. A revision of the South American Dipterous insects of the family Ptychopteridae. *Proc. U. S. Nat. Mus.* 44:331-335, 3 figs.

1920. A new genus and species of net-winged midge (Blepharocridae) and an undescribed species of Tanyderidae (Diptera). *Arkiv för Zool.* 13(7):5-7, fig.

1927. Diptera. Fam. Tanyderidae, in P. Wytzman: *Genera Insectorum* 189:13 pp., pl. Bruxelles.

1929. Tanyderidae, in *Diptera of Patagonia and South Chile*, Part I:i-xvi, 1-240, 12 pls. with 259 figs.

1930a. *Tanyderus pictus* Philippi (Fam. Tanyderidae, Ord. Diptera). *Rev. Chilena Hist. Nat.* 34:110-113.

1930b. Observations on the dipterous family Tanyderidae. *Proc. Linn. Soc. New South Wales* 55:221-230, pl. 5 (larva), pl. 6 (pupa).

1935. The distribution of *Tanyderus pictus* Ph. (Fam. Tanyderidae, Ord. Diptera). *Rev. Chilena Hist. Nat.* 39:86-87.

CRAMPTON, G. C.

1930a. Some anatomical details of the pupa of the archaic Tanyderid Dipteran *Protoplasa fitchii* O.S. *Proc. Ent. Soc. Washington* 32:83-98, 3 pls. with 17 figs.

1930b. A comparison of the more important structural details of the larva of the archaic Tanyderid Dipteran *Protoplasa fitchii* with other holometabola from the standpoint of phylogeny. *Bull. Brooklyn Ent. Soc.* 25:239-258, 4 pls. with 33 figs.

HINTON, H. E.

1966. The spiracular gill of the fly *Eutanyderus* (Tanyderidae). *Australian Jour. Zool.* 14:365-369, 2 figs., 1 pl. (pupa).

KNIGHT, A. W.

1963. Description of the Tanyderid larva *Protanyderus margarita* Alexander from Colorado. *Bull. Brooklyn Ent. Soc.* 58:99-102, 5 figs.

1964. Description of the Tanyderid pupa *Protanyderus margarita* Alexander from Colorado. *Ent. News* 75:237-241, 4 figs.

PEUS, FRITZ

1958. Tanyderidae, in Erwin Lindner, *Die Fliegen der palaearktischen Region*. Lief. 200:1-9, figs. 1-9.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verhandl. k. k. zool.-bot. Ges. Wien* 15:595-782, 7 pls.

ROSE, J. H.

1963. Supposed larva of *Protanyderus vipio* (Osten Sacken) discovered in California (Diptera: Tanyderidae). *Pan-Pacific Ent.* 39:272-275, 2 figs.

WILLIAMS, I. W.

1933. The external morphology of the primitive Tanyderid Dipteron *Protoplasa fitchii* O.S., with notes on the other Tanyderidae. *Jour. N. Y. Ent. Soc.* 41:1-35, text-fig. 1, 7 pls. with 38 figs.

WOOD, H. G.

1952. The crane-flies of the South-West Cape (Diptera, Tipuloidea). *Ann. South African Mus.* 39:1-327, 105 figs.

INDEX

Araucoderus Alexander, 1929	5.1
gloriosus (Alexander), 1920 (Tanyderus), Araucoderus	5.2
Neoderus Alexander, 1927	5.2
patagonicus (Alexander), 1913 (Tanyderus), Neoderus	5.2
pictus Philippi, 1865, Tanyderus	5.1
Tanyderus Philippi, 1865	5.1



Faint, illegible text at the top of the page.

Faint, illegible text in the upper middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.

Faint, illegible text in the middle section.







SciELO

SUPERFAMILY PSYCHODOIDEA

Family Tanyderidae

Family Psychodidae

Family Nymphomyiidae





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
OF THE UNITED STATES

6A

FAMILY PSYCHODIDAE

SUBFAMILIES BRUCHOMYIINAE, TRICHOMYIINAE,
SYCORACINAE AND PSYCHODINAE

D. A. DUCKHOUSE


Department of Zoology, University of Adelaide, South Australia

Apart from Phlebotominae, which have been studied extensively, at present the described neotropical Psychodidae are few in number and less well known than those of any other region. The indications are, however, that the fauna is large and very distinctive with a high proportion of genera and subgenera restricted to the neotropics. Of 20 non-phlebotomine genera listed below, 10 are unknown from any other region. The most important papers on this remarkable fauna are those of Tonnoir (1929), Barretto (1950, 1954a, 1954b, 1954c, 1956, 1961), Fairchild (1952), Quate (1963) and Duckhouse (1968). Several genera occur in Mexican amber (Quate, 1961, 1963b).

The Bruchomyiinae, which are the most plesiomorphic subfamily, are concentrated in northern Argentina, Paraguay and southern Brazil. Species of Bruchomyiinae are often taken with *Phlebotomus*, and a study of the immature stages of *Bruchomyia argentina* (Satchell, 1953c) has suggested a closer actual relationship with Phlebotominae than is shown by adult morphology alone. Some authors have accordingly united the two subfamilies (e.g. Fairchild, 1955) but I have preferred to follow Satchell by keeping them apart, thus giving expression to the considerable imaginal differences.

Sycoracinae have rarely been taken in the neotropics. However, this is more probably due to their small size and reclusive habits than to an actual scarcity. At present the subfamily contains a single genus, *Sycorax*. Several European species of *Sycorax* have aselliform, rheophile larvae (Feuerborn, 1932; Jung, 1956) and at least one is haematophagous (Desportes, 1942; Parrot, 1951). It is likely that the neotropical species are similar in these respects.

The Trichomyiinae contain a single genus, *Trichomyia*. Although the type species of *Trichomyia* has a wood-boring larva (Keilin & Tate, 1937) it should not be assumed that the neotropical species all share this habit. Probably the larger species, e.g. *T. edwardsi*, do, but from experience elsewhere I suspect that the small species whose palpi are



3-segmented, with the sensory rods occurring in a pit on segment 1 (e.g. *pedicellata*, *vargasi*) will be found to breed amongst the soft decomposing material contained in tree-holes.

The Psychodinae are by far the largest and most diversified subfamily, including numerous small and mostly unicolorous species, generally encountered in moist woods or forests where is an abundance of decomposing plant material. Some few species are widely distributed or cosmopolitan (e.g. *Telmatoscopus albipunctatus*, *Psychoda surcoufi*). Immatures often occur amongst layers of fallen leaves at the edges of streams or pools, or in tree-holes, while the adults rest nearby, on the overhanging vegetation, on tree trunks or in dark crevices. Immatures of most neotropical species probably resemble those of the European forms described by Satchell (1947, 1949). One obvious exception is the peculiar genus *Maruina*, first described by Müller (1895), in which the larva has a row of ventral suckers enabling it to attach to wet rocks in swift-flowing streams or by waterfalls.

The following abbreviations have been used:

A.M.N.H. — American Museum of Natural History, New York.

B.M. (N.H.) — British Museum (Natural History), London.

Dept. Paras., São Paulo — Department of Parasitology, Faculty of Medicine, University of São Paulo.

M.C.Z. — Museum of Comparative Zoology, Cambridge, Massachusetts.

U.S.N.M. — United States National Museum, Washington, D.C.

Z.M.B. — Zoologisches Museum, Berlin.

Subfamily BRUCHOMYIINAE

Genus BRUCHOMYIA Alexander

- BRUCHOMYIA Alexander, 1920:403. Type-species, *argentina* Alexander (orig. des.). Ref. — Fairchild, 1952:274-279.
- ALMEIDAI Barretto & d'Andretta, 1946:68-71 (♀); figs. 10-13. Type-locality: Brazil, Estado de São Paulo, Franca. Holotype ♀ in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Franca; Jaraguá). Ref. — Barretto, 1954:343-346; figs. 9-13 (♂).
- ARGENTINA Alexander, 1920:405 (♂); figs. 1-9. Type-locality: Argentina, Córdoba prov., La Granja, Alta Gracia. Holotype ♂, in Alexander collection. Distr. — Argentina (Córdoba prov., La Granja). Refs. — Tonnoir, 1940:212; figs. 5E, 7A, B (♂); Satchell, 1953c:1-12; figs. 1-14 (immatures).
- BRASILIENSIS Alexander, 1940:795-796 (♂); figs. 1, 3. Type-locality: Brazil, Ceará, Crato. Holotype ♂ in Alexander collection. Distr. — Brazil (Ceará, Crato).
- FUSCA Barretto, 1950:348-350 (♂); figs. 6-8. Type-locality: Brazil, São Paulo, Serra da Cantareira. Holotype ♂ in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira; Rio de Janeiro, Boa Vista). Ref. — Fairchild, 1952:276-279 (♂, ♀); figs. 46-52.
- PERUVIANA Alexander, 1929:6 (♀). Type-locality: Peru, Chanchamayo. Colonia Perené. Holotype ♀ in U.S.N.M., n° 41581. Distr. — Peru (Chanchamayo, Colonia Perené).



- PLAUMANNI Alexander, 1944:313-315 (♂); figs. 1-2. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂ in Alexander collection. Distr. — Brazil (Santa Catarina, Nova Teutônia).
- SHANNONI Alexander, 1929:4-6 (♂, ♀); figs. 1-2. Type-locality: Peru, Department of Lima, Verrugas Cañon. Holotype ♂ in U.S.N.M., n° 41582. Distr. — Peru (Dept. Lima, Verrugas Cañon; Rimac Valley). Ref. — Fairchild, 1952:279 (♂, ♀); figs. 53-56.
- UNICOLOR Barretto, 1950:347-348 (♂); figs. 14-16. Type-locality: Brazil, São Paulo, Jaraguá. Holotype ♂ in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Jaraguá).

Genus NEMOPALPUS Macquart

- NEMOPALPUS Macquart, 1838a:102. Type-species, *flavus* Macquart (mon.). Refs. — Macquart, 1838b; Fairchild, 1952:260-274 (includes key); Fairchild, 1955:182-188; Jung, 1963.
- Nemopalpus* Macquart, 1838b:85 (*lapsus*).
- ANTHILLARUM Fairchild, 1952:270-272 (♂); figs. 16-21, 32, 38. Type-locality: Dominican Republic, Sabana del Mar, Cacao Plantation, 100 ft. Holotype ♂, in Gorgas Memorial Laboratory, Panama. Distr. — Dominican Republic (Sabana del Mar, 100-650 ft.).
- ARROYO De Leon, 1950:14-16 (♂, ♀); figs. 12-18. Type-locality: Guatemala, Mazatenango. Holotype ♂, presumably in De Leon collection or University of San Carlos, Guatemala. Distr. — Guatemala (Mazatenango), Mexico (Michoacán, basin of R. Tepalcatepec), Panama (Chiriqui Prov.). Ref. — Fairchild, 1952:264-268 (♂, ♀); figs. 1-11, 30, 31, 39.
- BREVINERVIS Barretto & d'Andretta, 1946:60-62 (♂); figs. 1-6. Type-locality: Brazil, São Paulo. Holotype ♂ in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo).
- DAMPFIANUS Alexander, 1940:796-798 (♂); figs. 2-4. Type-locality: Mexico, Chiapas, Finca Vergel. Holotype ♂, in Alexander collection. Distr. — Mexico (Chiapas, Finca Vergel; Palenque). Ref. — Fairchild, 1952:272-274 (♀); figs. 12, 13, 22-26, 37.
- DISSIMILIS Barretto & d'Andretta, 1946:62-64 (♀); figs. 7-8. Type-locality: Brazil, Estado do Espírito Santo, São João. Holotype ♀ in Dep. Paras., São Paulo. Distr. — Brazil (Espírito Santo, São João; São Paulo, Jaraguá). Ref. — Barretto, 1950:341-343 (♂); figs. 1-5.
- IMMACULATUS Freeman, 1949:86-87 (♂); figs. 1-3. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype, in B. M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).
- MOPANI De Leon, 1950:9-14 (♂, ♀); figs. 2, 5-11. Type-locality: Guatemala, Petén, Canchacán. Holotype ♂, presumably in De Leon collection or University of San Carlos, Guatemala. Distr. — Guatemala (Petén, Canchacán), Mexico (Chiapas, Palenque). Ref. — Fairchild, 1952:263-264 (♂, ♀); figs. 14, 15, 33-36.
- MORALES De Leon, 1950:16-20 (♂, ♀); figs. 19-20. Type-locality: Guatemala, Dept. de Suchitepéquez, Mazatenango. Holotype ♂, presumably in De Leon collection or University of San Carlos, Guatemala. Distr. — Guatemala (Dept. Suchitepéquez, Mazatenango). Ref. — Fairchild, 1952:262.



- PALLIPES (Shannon & Del Ponte), 1927:733-734 (♂) (*Bruchomyia*). Type-locality: Argentina, Misiones, Cataratas del Iguazú. Holotype ♀, in U.S.N.M., n° 41580. Distr. — Argentina (Misiones, Cataratas del Iguazú). Refs. — Alexander, 1929:8; Fairchild, 1952:262.
- PILIPES Tonnoir, 1922:130-134 (♂, ♀); figs. 1-8. Type-locality: Paraguay, San Bernardino. Holotype ♂, in Vienna Museum. Distr. — Paraguay (San Bernardino), Argentina (Iguazú Falls), Brazil (São Paulo, Bateia; Rio de Janciro, Campo Belo; Santa Catarina, Nova Teutônia). Refs. — Alexander, 1929:8; Freeman, 1949:87-88; figs. 4-7.
- maculipennis* Barretto & d'Andretta, 1946:64-66 (♀); fig. 9. Type-locality: Brazil, Estado de São Paulo, Bateia.
- voxans* Alexander, 1940:798 (♂); fig. 5. Type-locality: Brazil, Rio de Janeiro, Campo Belo.
- SZILADYI Tonnoir, 1940:206-208 (♂); figs. 1, 3, 4. Type-locality: Costa Rica, Suiza de Turrialba. Holotype ♂, in Hungarian National Museum. Distr. — Costa Rica (Suiza de Turrialba), Panama (Bocas del Toro Prov.). Ref. — Fairchild, 1952:268-270 (♂, ♀); figs. 27-29.

Subfamily TRICHOMYIINAE

Genus TRICHOMYIA Haliday

- TRICHOMYIA Haliday, 1839:745. Type-species, *urbica* Haliday (mon.). Refs. — Satchell, 1956:147-148; Duckhouse, 1965:334.
- Eubonetia* Vargas & Diaz Nájera, 1953:155-156. Type-species, *Trichomyia cirrata* Coquillett (mon.).
- Lepria* Enderlein, 1937:112. Type-species, *squamosa* Enderlein (orig. des.).
- Termitodipteron* Holmgren, 1905:533-536; figs. 4-6. Type-species, *wasmanni* Holmgren (mon.).
- ARMATA Barretto, 1954b:127-129 (♂, ♀); figs. 1-12. Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♂, in Dep. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- BARRETTO, *nom. nov.* for *Eubonetia fairchildi* Barretto, 1954a, which is a junior homonym of *Trichomyia fairchildi* Vargas & Diaz Nájera, 1953. Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♀, Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira). Ref. — Barretto, 1954a:85-87 (♀); figs. 5-9.
- fairchildi* Barretto, 1954a:85-87 (*Eubonetia*; preocc. Vargas & Diaz Nájera, 1953). Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal.
- BRASILENSIS Satchell, 1956:152-153 (♂); fig. 5A-E. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).
- CIRRATA Coquillett, 1902:137 (♀). Type-locality: Mexico, Tabasco, Frontera. Holotype ♀, in U.S.N.M., n° 6309. Distr. — Mexico (Tabasco, Frontera). Refs. — Satchell, 1956:149-151 (♂, ♀); fig. 2A-H; Quate & Wirth, 1951:163.



- EOUTINHOI** (Barretto), 1954a:87-89 (♀); figs. 10-15 (*Eubonetia*). Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♀ in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- EATONI** Satchell, 1956:151-152 (♀); fig. 4A-E. Type-locality: Jamaica, Moneague. Holotype ♀ in B.M. (N.H.). Distr. — Jamaica (Moneague). Ref. — Duekhouse, 1965.
- EDWARDSI** Tonnoir, 1929:30-31 (♂); figs. 66-70. Type-locality: Argentina, Neuquén province, Puerto Blest. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén province, Puerto Blest). Ref. — Satchell, 1956:151 (♀); fig. 3A, B (dubious identification).
- FAIRCIELDI** Vargas & Diaz Nájera, 1953:154; figs. 5, 8. Type-locality: Mexico, Chiapas, Cerro Hueytepae, San Cristóbal Las Casas. Location of type unknown. Distr. — Mexico (Chiapas, Cerro Hueytepae).
- flavescens* Von Roser. Not neotropical.
- INERMIS** Barretto, 1954b:129-130 (♂); figs. 13-17. Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♂, in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- MALDONADOI** (Vargas), 1953:128-129 (♀); figs. 8-11 (*Eubonetia*). Type-locality: Mexico, Estado de Chiapas, Palenque, Río Chacamae. Holotype ♀, in Instituto de Salubridad y Enfermedades Tropicales. Distr. — Mexico (Chiapas, Palenque).
- PEDICILLATA** Satchell, 1956:153-154 (♂, ♀); fig. 6A-G. Type-locality: Panama, Almirante, Bocas. Holotype ♂, in B.M. (N.H.). Distr. — Panama (Almirante, Bocas), Canal Zone (Forst Reserve, Cruces Trail). Ref. — Duckhouse, 1965:343.
- SQUAMOSA** (Enderlein), 1937:112 (♀); fig. 1 (*Lepria*). Type-locality: "Costa Rica". Holotype ♀, in Z.M.B. Distr. — Costa Rica. Refs. — Satchell, 1956:148; Quate, 1963a:182-183; fig. 1(a)-(f).
- TRAVASSOSI** (Barretto), 1954a:91-94 (♀); figs. 22-26 (*Eubonetia*). Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♀, in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- urbica* Haliday. Not neotropical.
- VARGASI** (Barretto), 1954a:83-85 (♂); figs. 1-4 (*Eubonetia*). Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♂, in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- VAZI** (Barretto), 1954a:89-91 (♀); figs. 16-21 (*Eubonetia*). Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♀, in Dept. Paras., São Paulo. Distr. — Brazil (São Paulo, Serra da Cantareira).
- WASMANNI** (Holmgren), 1905:533-534 (♂); figs. 4-6 (*Termitodipteron*). Type-locality: Peru, Carabaya Province, Chaquimayo, in nest of *Eutermes rotundiceps* Holmgren. Location of type unknown. Distr. — Peru (Carabaya Province, Chaquimayo). Refs. — Speiser, 1906:717; Edwards, in Tonnoir, 1929: footnote, p. 2.



Subfamily SYCORACINAE

Genus SYCORAX Haliday

- SYCORAX Haliday, 1839:745. Type-species, *silacea* Haliday (mon.). Ref. — Duckhouse, 1965:332.
- ASSIMILIS Barretto, 1956:71-73 (♂, ♀); figs. 1-8. Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♂, in Dept. Paras., São Paulo, Distr. — Brazil (São Paulo, Serra da Cantareira).
- CHILENSIS Tonnoir, 1929:31-32 (♂, ♀); figs. 71-75. Type-locality: Chile, Llanquihue, Peulla. Holotype ♂, in B.M. (N.H.). Distr. — Chile (Llanquihue, Peulla; Chiloé, Isla de Chiloé).
- SATCHELLI Barretto, 1956:73-75 (♂, ♀); figs. 9-18. Type-locality: Brazil, São Paulo, Serra da Cantareira, Horto Florestal. Holotype ♂, in Dep. Paras., São Paulo, Distr. — Brazil (São Paulo, Serra da Cantareira).
- sp. Fairchild, 1955:184; fig. 14 (wing). Locality: Panama.

Subfamily PSYCHODINAE

Genus ALEPIA Enderlein

- ALEPIA Enderlein, 1937:94-95. Type-species, *scripta* Enderlein (orig. des.). Refs. — Quate, 1963a:192; Duckhouse, 1968:31.
- ALBICOLLARE (Enderlein), 1937:104 (♂, described as ♀) (*Chirolepia*). Type-locality: Bolivia, Sara Province, 600-700 m. Holotype ♂, in Z.M.B. Distr. — Bolivia (Sara province), Paraguay. Ref. — Quate, 1963a: 193-194 (♂); fig. 10(a)-(h).
- BISUBULATA Duckhouse, 1968:31-33 (♂); figs. 12-16. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).
- HIRTIVENTRIS (Tonnoir), 1920:151-154 (♂); fig. 3 (*Psychoda*). Type-locality: Brazil, Amazon, Gurupá. Location of type not know. Distr. — Brazil (Amazon, Gurupá). N. COMB.
- hirtiventris* Enderlein, 1937:86 (lapsus).
- SCRIPTA Enderlein, 1937:95 (♂, described as ♀). Type-locality: Bolivia, Yungas. Holotype ♂, in Z.M.B. Distr. — Bolivia (Yungas). Ref. — Quate, 1963a:192 (♂); fig. 9(a)-(e).
- sp. Duckhouse, 1968:33-34 (♀); figs. 17-22. Locality: Brazil, Santa Catarina, Nova Teutônia.

Genus ARISEMUS Satchell

- Telmatoscopus*, subg. ARISEMUS Satchell, 1955:88. Type-species, *maculosus* Satchell (orig. des.). Ref. — Botosaneanu & Vaillant, 1970:177-178.
- BOXI Satchell, 1955:89-90 (♂); fig. 3A-D. Type-locality: West Indies, Windward Is., St. Lucia, Bar de l'Isle. Holotype ♂, in B.M. (N.H.). Distr. — West Indies (Windward Is., St. Lucia).

HEXADACTYLUS Botosaneanu & Vaillant, 1970:180-181 (♂); plate 1, fig. 8; plate 2, figs. 1-10. Type-locality: West Indies, Cuba, Rio Caburny, 550 m. No type material is cited. Distr. — West Indies (Cuba, Rio Caburny).

MACULOSUS Satchell, 1955:88-89 (♂, ♀); fig. 2a-g. Type-locality: West Indies, Jamaica, Moncaguc. Holotype ♂, in B.M. (N.H.). Distr. — West Indies (Jamaica, Moneague).

TETRADACTYLUS Botosaneanu & Vaillant, 1970:181; plate 3, figs. 1-4. Type-locality: West Indies, Cuba, Sierra Sagua Barracoa, 420 m. No type material is cited. Distr. — West Indies (Cuba, Sierra Sagua Barracoa).

Genus BRUNETTIA Annandale

BRUNETTIA Annandale, 1910:141-142. Type-species, *Diplonema superstes* Annandale (Brunetti, 1911:310). Ref. — Duckhouse, 1966:187-188.

ALBONOTATA (Brunetti), 1908:373 (♀); fig. 7 (*Psychoda*). Type-locality: India, Calcutta. Holotype ♀, in Zoological Survey of India, Calcutta, n° 5901/18. Distr. — India, Ceylon, Africa, Malaya, Formosa, Jamaica (Hopedale).

duripuncta Curran, 1928:30 (♂, ♀) (*Psychoda*). Type-locality: Jamaica, Hopedale. (Quate, 1962:169, doubts this synonymy).

duripunctata Quate, 1962:169 (lapsus).

Genus CHIROLEPIA Enderlein

CHIROLEPIA Enderlein, 1937:103. Type-species, *maculipennis* Enderlein (orig. des.). Ref. — Duckhouse, 1968:33.

MACULIPENNIS Enderlein, 1937:103 (♀). Type-locality: Bolivia, Yungas. Holotype ♀, in Z.M.B. Distr. — Bolivia (Yungas). Rcfs. — Quate, 1963a:192 (♀); fig. 10(i)-(l); Duckhouse, 1968:33.

Genus DICTYOCAMPSA Enderlein

DICTYOCAMPSA Enderlein, 1937:100. Type-species, *guttata* Enderlein (orig. des.). Ref. — Quate, 1963a:187-188.

GUTTATA Enderlein, 1937:100 (♂). Type-locality: Brazil, Colônia Hansa (= Corupá). Holotype ♂, in Z.M.B. Distr. — Brazil (Colônia Hansa). Ref. — Quate, 1963a:188 (♂); fig. 6(a)-(d).

Genus KUPARA Rapp

KUPARA Rapp, 1945:310. Type-species, *albipeda* Rapp (mon.). Ref. — Fairchild, 1951:17.

ALBIPEDA Rapp, 1945:311 (♀); fig. 5(a)-(c). Type-locality: Canal Zone, Barro Colorado Island. Holotype ♀, in A.M.N.H. Distr. — Canal Zone (Barro Colorado Island).



Genus MARUINA Müller

- MARUINA Müller, 1895:480-481. Type-species, *pilosella* Müller (Enderlein, 1937:109). Refs. — Quate & Wirth, 1951; Vaillant, 1963; Duckhouse, 1966:186-187.
- HIRTA Johannsen, 1938:224-225 (♂, ♀ immatures). Type-locality: Puerto Rico, Guaynabo River. Holotype ♂, in Dept. of Entomology, Cornell University. Distr. — Puerto Rico (Guaynabo and Yunez rivers). Ref. — Quate & Wirth, 1951:154 (♂, pupa); plate 1, figs. 2, 2A; plate 2, fig. 1.
- PILOSELLA Müller, 1895:480-481 (♂, ♀, immatures); figs. 2-5, 8, 9, 11-13, 18-20. Type-locality: Brazil, Santa Catarina, Itajaí, Blumenau?; No type specimen in known. Distr. — Brazil (Itajaí, Blumenau?; São Paulo, Rancharia, Usina Caiuá), Argentina (Tucumán province, Dept. Monteros, Quebrada de Los Sosa Acherál). Ref. — Quate & Wirth, 1951 (♂, ♀, immatures); plate 1, figs. 3, 3A; plate 2, figs. 2, 3.
- sp. A Quate & Wirth, 1951:162. Locality: Mexico, Contreras, near Mexico City (pupae in U.S.N.M.).
- SPINOSA Müller, 1895:480-481 (♀, immatures); figs. 6, 10, 14-17. Type-locality: Brazil, Santa Catarina (Itajaí, Blumenau). No type specimen is known. Distr. — Brazil (Itajaí, Blumenau). Ref. — Quate & Wirth, 1951:161-2.

Genus NEMONEURA Tonnoir

- NEMONEURA Tonnoir, 1929:27. Type-species, *Psychoda punctata* Philippi (orig. des.). Refs. — Satchell, 1953b:417-418; Quate, 1965a:466.
- DEALBATA Tonnoir, 1929:29 (♂); figs. 64-65. Type-locality: Chile, Llanquihue, Casa Pangué. Holotype ♂, in B.M. (N.H.). Distr. — Chile (Llanquihue, Casa Pangué; Chiloé, Isla de Chiloé), Argentina (Neuquén, Puerto Blest).
- PUNCTATA (Philippi), 1865:631 (*Psychoda*). Type-locality: Chile, Valdivia province. No type specimen is known. Distr. — Chile (Valdivia province; Llanquihue, Peulla; Chiloé, Isla de Chiloé). Ref. — Tonnoir, 1929:27-29 (♀); figs. 61-63.

Genus PARASETOMIMA Duckhouse

- PARASETOMIMA Duckhouse, 1968:29-31. Type-species, *umbella* Duckhouse (mon.).
- UMBELLA Duckhouse, 1968:31 (♂); figs. 1-8. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).

Genus PERICOMA Walker

- PERICOMA Walker, 1856:256. Type-species, *Trichoptera trifasciata* Meigen (Coquillett, 1910:587). Refs. — Quate, 1963a; Duckhouse, 1966:155-156.



Some of Enderlein's genera given below have been tentatively recognized as subgenera (Quate, 1963a), but until the group is better known I prefer to leave all species under the one heading).

Desmioza Enderlein, 1937:99. Type-species, *Pericoma edwardsi* Tonnoir (orig. des.).

Didicrum Enderlein, 1937:105. Type-species, *Pericoma griseata* Tonnoir (orig. des.).

Mecysmia Enderlein, 1937:108. Type-species, *Mecysmia schoenemanni* Enderlein = *simplex* Tonnoir (orig. des.).

Podolepria Enderlein, 1937:105. Type-species, *Pericoma inornata* Tonnoir (orig. des.).

Synmormia Enderlein, 1937:99. Type-species, *Pericoma chilensis* Tonnoir (orig. des.).

Syntomolaba Enderlein, 1937:100. Type-species, *Pericoma complicata* Tonnoir (orig. des.).

ALFAROANA (Dyar), 1926a:110 (♂, ♀) (*Psychoda*). Type-locality: Costa Rica, Alajuela. Holotype ♂, in U.S.N.M., n° 29392. Distr. — Costa Rica (Alajuela). Ref. — Del Rosario, 1936:145-146.

AMPLIPENNA (Knab), 1914:103-104 (♂) (*Psychoda*). Type-locality: West Indies, Cuba, San Antonio de los Baños. Holotype ♂, in U.S.N.M., n° 15933. Distr. — West Indies (Cuba, San Antonio de los Baños). Ref. — Del Rosario, 1936:145-146.

BRASILENSIS Duckhouse, 1968:34 (♀); figs. 23-28. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♀, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).

BUSEKANA (Dyar), 1926a:109 (♂) (*Psychoda*). Type-locality: West Indies, Martinique. Holotype ♂, in U.S.N.M., n° 29391. Distr. — West Indies (Martinique). Ref. — Del Rosario, 1936:145-146.

CAUDATA Satchell, 1955:90-92 (♂); fig. 4A-E. Type-locality: Jamaica, Runaway Bay. Holotype ♂, in B.M. (N.H.). Distr. — Jamaica (Runaway Bay).

CHILENSIS Tonnoir, 1929:14-15 (♂, ♀); figs. 19-22. Type-locality: Chile, Chiloé, Isla de Chiloé, Ancud. Holotype ♂, in B.M. (N.H.). Distr. — Chile (Chiloé, Isla de Chiloé).

COMPLICATA Tonnoir, 1929:15-16 (♂); figs. 23-27. Type-locality: Argentina, Neuquén province, Lake Nahuel Huapi, Puerto Blest. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Puerto Blest).

CONTIGUA Tonnoir, 1929:24-25 (♂, ♀); figs. 51-53. Type-locality: Argentina, Neuquén province, Puerto Blest. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Puerto Blest), Chile (Llanquihue, Casa Pangué).

EDWARDSI Tonnoir, 1929:12-14 (♂, ♀); figs. 14-18. Type-locality: Argentina, Neuquén province, Bariloche. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Bariloche). Ref. — Quate, 1963a:183; fig. 2(a)-(g).


EQUALIS Tonnoir, 1934:71 (nom. nov. for *speciosa* Tonnoir). Type-locality: Chile, Llanquihue, Casa Pangué. Holotype ♀, in B.M. (N.H.). Distr. — Chile (Llanquihue, Casa Pangué). Ref. — Tonnoir, 1929:11-12 (♀); fig. 13.



- speciosa* Tonnoir, 1929:11-12 (preocc. Mcunier, 1905). Type-locality: Chile, Llanquihuc, Casa Pangue.
- aequalis* Enderlein, 1937:103 (*Thrysocanthus*) (*lapsus*).
- FENESTRATA Tonnoir, 1929:20 (♀); figs. 37-39. Type-locality: Chile, Llanquihue, Casa Pangue. Holotype ♀, in B.M. (N.H.). Distr. — Chile (Llanquihue, Casa Pangue). Refs. — Enderlein, 1937:92 (in *Synseodais*); Quate, 1963a:189.
- GRABHAMANA (Dyar), 1926a:110 (♂, ♀) (*Psychoda*). Type-locality: West Indies, Jamaica, Kingston. Holotype, in U.S.N.M., n° 29393. Distr. — West Indies (Jamaica, Kingston). Ref. — Del Rosario, 1936:145-146.
- GRISEATA Tonnoir, 1929:22 (♀); figs. 42-45. Type-locality: Chile, Chiloé, Isla de Chiloé, Ancud. Holotype ♀, in B.M. (N.H.). Distr. — Chile (Chiloé, Isla de Chiloé). Ref. — Quate, 1963a:184-185.
- INCOMPLETA (Knab), 1914:105-106 (♂) (*Psychoda*). Type-locality: Panama, Canal Zone, Tabernilla. Holotype ♂, in U.S.N.M., n° 15936. Distr. — Canal Zone (Tabernilla). Ref. — Del Rosario, 1936:145-146.
- INORNATA Tonnoir, 1929:23-24 (♂, ♀); figs. 46-50. Type-locality: Argentina, Neuquén province, Puerto Blest. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Puerto Blest; Bariloche). Ref. — Quate, 1963a:185.
- NIVEOPUNCTATA Tonnoir, 1929:18-20 (♂, ♀); figs. 32-36. Type-locality: Chile, Chiloé, Isla de Chiloé, Castro. Holotype ♂, in B.M. (N.H.). Distr. — Chile (Chiloé, Isla de Chiloé). Ref. — Quate, 1963a:185.
- PALLIDULA Tonnoir, 1929:16-18 (♂, ♀); figs. 28-31. Type-locality: Argentina, Lake Correntoso. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Lake Correntoso). Ref. — Quate, 1963a:185.
- PICTIPENNIS Tonnoir, 1920:155-157 (♀). Type-locality: Jamaica, Kingston. Holotype ♀, in B.M. (N.H.). Distr. — Jamaica (Kingston). Ref. — Satchell, 1955:93.
- SIMPLEX Tonnoir, 1929:25-26 (♂); figs. 54-57. Type-locality: Argentina, Neuquén province, Bariloche. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Bariloche), Chile (Llanquihue, Casa Pangue). Ref. — Quate, 1963a:184-185; figs. 3(a)-(g).
- schoenemanni* Enderlein, 1937:108 (*Mecysmia*). Type-locality: Chile, Contulmo. Ref. — Quate, 1963a:185.
- SQUAMITARSIS Tonnoir, 1929:21 (♀); figs. 40-41. Type-locality: Chile, Llanquihue, Casa Pangue. Holotype ♀, in B.H. (N.H.). Distr. — Chile (Llanquihue, Casa Pangue).
- TRICOLOR (Knab), 1914:105 (♀) (*Psychoda*). Type-locality: Panama, Coscajar River. Holotype ♀, in U.S.N.M., n° 15935. Distr. — Panama (Coscajar River). Ref. — Del Rosario, 1936:145-146.
- VIDUATA Tonnoir, 1929:26 (♀); figs. 58-60. Type-locality: Chile, Chiloé, Isla de Chiloé, Ancud. Holotype ♀, in B.M. (N.H.). Distr. — Chile (Chiloé, Isla de Chiloé). Ref. — Quate, 1963a:185.

Genus PHILOSEPEDON Eaton

- PHILOSEPEDON Eaton, 1904:57. Type-species, *Psychoda humeralis* Meigen (orig. des.). Ref. — Quate, 1959c:448.



Subgenus PHILOSEPEDON Eaton

INTERDICTA (Dyar), 1928:88 (♂, ♀); plate 4, figs. 5-7 (*Psychoda*). Type-locality: United States, New York, Ithaca. Holotype ♂ (?), in U.S.N.M., n° 40516. Distr. — United States, West Indies (St. Lucia, Castries). Ref. — Quate, 1955:227-229 (♂, ♀); fig. 77a-c.

Subgenus EURYGARKA Quate, n. status

EURYGARKA Quate, 1959c:450 (as genus). Type-species, *Psychoda helcis* Dyar (orig. des.).

HELICIS (Dyar), 1929:64 (♂, ♀) (*Psychoda*). Type-locality: West Indies, Cuba, Central Jaronú. Holotype ♂, in U.S.N.M., n° 41186. Distr. — United States (Florida; Alabama), Cuba (Central Jaronú). Ref. — Quate, 1955:235 (♂, ♀); fig. 83a-d.

Subgenus QUATIELLA Botosaneanu & Vaillant, n. status

QUATIELLA Botosaneanu & Vaillant, 1970:173-174 (as genus) Type-species, *Philosepedon leonhardti* Vaillant (orig. des.).

CUBANA (Botosaneanu & Vaillant), 1970:174-176 (♂, ♀); plate 1, figs. 1-7. (*Quatiella*). Type-locality: West Indies, Cuba, Province de Las Villas, Sierra Escambray, Rio Caburny, 550 m. No type material is cited. Distr. — West Indies (Cuba, Province de las Villas. Sierra Escambray).

FUMATA (Knab), 1914:104-105 (♂) (*Psychoda*). Type-locality: Mexico, Córdoba. Holotype ♂, in U.S.N.M., n° 15934. Distr. — Mexico (Córdoba). Ref. — Del Rosario, 1936:132-134 (♂); plate 5, figs. 7-10. N. COMB.

PLAUMANNI (Duckhouse), 1968:39 (♂), figs. 49-53 (*Psychoda*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia). N. COMB.

SPATHIPENIS (Duckhouse), 1968:37 (♂); figs. 44-48 (*Psychoda*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia). N. COMB.

Genus PLATYPLASTINX Enderlein

PLATYPLASTINX Enderlein, 1937:107. Type-species, *solor* Enderlein (orig. des.). Ref. — Quate, 1963a:195.

SOLOX Enderlein, 1937:107. Type-locality: "Costa Rica". Holotype ♀, in Z.M.B. — Ref. — Quate, 1963a:195 (♀); fig. 11(a)-(d).

Genus PSYCHODA Latreille

PSYCHODA Latreille, 1796:152. Type-species, *Tipula phalaenoides* Linnaeus (sub. mon., Latrille, 1802:424). Ref. — Duckhouse, 1966:194.




- AITKENI Quate, 1959a:218-219 (♀); figs. 10-11. Type-locality: West Indies, Trinidad, Chaguaramas. Holotype ♀, in U.S.N.M. Distr. — West Indies (Trinidad, Chaguaramas), Canal Zone, Panama.
- ALABANGENSIS Del Rosario, 1936:566 (♂); plate 1, figs. 17-20. Type-locality: Philippine Is., Luzon, Rizal Province, Alabang. Holotype ♂, in Philippine Bureau of Science. Distr. — Philippines, India, Ceylon, Borneo, Formosa, New Guinea, Micronesia; Trinidad (Tucker Valley). Ref. — Quate, 1959b:472; figs. 21a-c (♀).
- ichthycerca* Quate, 1959b:472 (♀); fig. 21a-c. Type-locality: Micronesia, Yap Is., Yap.
- ALTERNATA Say, 1824:358. Type-locality: U.S.A., Pennsylvania, Philadelphia. No type-specimen is known. Distr. — All zoogeographical regions. Mexico (Orizaba; Córdoba), Canal Zone (Ft. Kobbe), West Indies (Puerto Rico, Bayamon; Trinidad; Jamaica), Chile (Chiloé, Puerto Montt). Refs. — Quate, 1955:218-222 (♂, ♀); fig. 72a-c; Duckhouse, 1966:195-197; fig. 146; Satchell, 1947:60-63 (larva); figs. 47-52.
- ALTERNICULA Quate, 1955:222 (♂, ♀); fig. 73a-c. Type-locality: U.S.A., Florida, W. Palm Beach, Morrison Field. Holotype ♀, in U.S.N.M. Distr. — U.S.A., Mexico (S.L. Potosí, Ciudad Valles), Canal Zone (Ft. Kobbe), West Indies (Trinidad, Tucker Valley).
- CINEREA Banks, 1894:331 (♂, ♀). Type-locality: U.S.A., New York, Long Island, Sea Cliff. Holotype, in Museum of Comparative Zoology, Harvard, U.S.A. Distr. — All zoogeographical regions. West Indies (Puerto Rico, Rio Piedras), Chile (Santiago), Juan Fernández Is. (Masatierra, Cumberland Bay). Refs. — Duckhouse, 1966:208-209 (♂, ♀); figs. 204-207; Satchell, 1947:56-58 (larva); figs. 13-19.
- COCHILEARIA Satchell, 1950:181-183 (♂, ♀); figs. 4A-F. Type-locality: Fiji Is., Suva, Botanical Gardens. Holotype ♂, in B.M. (N.H.). Distr. — Fiji, Samoa, Micronesia, Borneo, Philippines; Canal Zone (Ft. Kobbe), West Indies (Trinidad).
- DIVARICATA Duckhouse, 1968:37 (♂, ♀); figs. 34-42. Type-locality: Brazil. Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).
- GEHRKEAE Quate, 1959a:216 (♀); figs. 6-7. Type-locality: West Indies, Trinidad, Tucker Valley. Holotype ♀, in U.S.N.M. Distr. — West Indies (Trinidad, Tucker Valley).
- grisescens* Tonnoir. Not neotropical.
- MASATIERRENSIS Satchell, 1952:106-107 (♀); fig. 1A-F. Type-locality: Juan Fernández Is., Masatierra, Bahía Cumberland. Location of type unknown. Distr. — Juan Fernández Is. (Masatierra, Bahía Cumberland; Masafuera).
- PARTHENOGENETICA Tonnoir, 1940:53-54 (♀). Type-locality: "Europe". No type-specimen is known. Distr. — Europe, Japan, Korea, New Zealand, southern Australia, South Georgia, Kerguelen I., Macquarie I., Campbell I.; Juan Fernández Is. Ref. — Tonnoir, 1922:80; fig. 11A (♀, as *Psychoda severini* Tonn.); Duckhouse, 1962:428.
- phalacnoides* Tonnoir. Not neotropical.
- PLAESIA Quate, 1959b:461-462 (♂, ♀); fig. 16a-d. Type-locality: Micronesia, Bonin Is., Chichi Jima, Omura. Holotype ♂, in Kyushu Univer-



- sity. Distr. — Micronesia (Bonin Is., Carolinc Is.); Canal Zone (Ft. Kobbe). Ref. — Quate, 1959a:219.
- PLATIOBATA** Tokunaga, 1957:65-66 (♀); figs. 34-36. Type-locality: Formosa, Szechuan, nr. Chaochow. Location of type unknown. Distr. — Formosa, Philippine Is., Borneo, New Guinea; West Indies (Trinidad; Jamaica). Ref. — Quate, 1965b:888 (♂, ♀); fig. 28a-b.
- PSEUDOCOMPAR** Tonnoir, 1929:7-8 (♂); figs. 5-9. Type-locality: Argentina, Lake Correntoso. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Lake Correntoso).
- SAVAIENSIS** Edwards, 1928:74 (♂); fig. 12(a)-(c). Type-locality: Samoa, Savaii, Salailua. Holotype ♂, in B.M. (N.H.). Distr. — Samoa, Cook Is., Micronesia, New Britain, Bougainville, New Guinea, Borneo, Sarawak, Philippines, Ryukyu Is., Taiwan, Japan, Hawaiian Is., U.S.A. Nicaragua (Rama, Zelaya), Costa Rica (Higuito, San Mateo; San Pedro de Montes de Oca), Panama (Trinidad Rio), Tobago I., Canal Zone (Ft. Claiton; Ft. Kobbe; Barro Colorado I.; Mojinga Swamp), West Indies (Puerto Rico; Trinidad). Ref. — Satchell, 1953a:183-185 (♂, ♀); fig. 3A-G (as *rarotongensis*).
lucia Quate, 1954:349-350 (♂, ♀); figs. 28-31. Type-locality: West Indies, St. Lucia, Castries.
rarotongensis Satchell, 1953a:183-185 (♂, ♀); fig. 3A-G. Type-locality: Cook Is., Rarotonga, Wigmores Waterfall.
- SCOTINA** Quate, 1959:218 (♀); figs. 8-9. Type-locality: Canal Zone, Fort Claiton. Holotype ♀, in U.S.N.M. Distr. — Canal Zone (Fort Claiton), West Indies (Trinidad, Tueker Valley).
- SIMILLIMA** Tonnoir, 1929:8-9 (♂); figs. 10-12. Type-locality: Argentina, Neuquén province, Bariloche. Holotype ♂, in B.M. (N.H.). Distr. — Argentina (Neuquén, Bariloche).
- sp. Dyar, 1926b:103-105; plate 1, fig. 1 (as *phalaenoides* Tonnoir). Localities: New Mexico (Las Cruces; Las Vegas), Mexico (Córdoba), Guatemala (Dep. Sololá, Finea San Basilio), Venezuela (Dep. Aragua, Maracay), Chile (Angol). Probably Dyar was dealing with several or many species. Ref. — Quate, 1955:212-213.
- SUREOUFI** Tonnoir, 1922:74-76 (♂, ♀); fig. 9. Type-locality: France, Lamballe. Holotype in Natural History Museum, Paris. Distr. — Europe, southern Australia, New Zealand, Campbell I., Macquarie I., Antipodes Is., Chile (Chiloé, Isla de Chiloé, Aneud; Puerto Montt). Ref. — Duekhouse, 1966:211-213.
subimmaculata Tonnoir, 1929:6-7 (♀); figs. 1-4. Type-locality: Chile, Chiloé, Isla de Chiloé, Aneud.
- ZETOSCOTA** Quate, 1959a:214 (♂); figs. 1-2. Type-locality: Canal Zone, Fort Claiton. Holotype ♂, in U.S.N.M. Distr. — Canal Zone (Ft. Claiton), West Indies (Trinidad, Tucker Valley).

Genus SYNTOMOZA Enderlein

- SYNTOMOZA** Enderlein, 1937:88-89. Type-species, *niveitarsis* Enderlein (orig. des.). Ref. — Quate, 1963a:185-186.
Lepidiella Enderlein, 1937:89. Type-species, *lanuginosa* Enderlein (orig. des.). Ref. — Quate, 1963a:187.



- CERVI (Satchell), 1955:92-93 (♂, ♀); fig. 5A-E (*Pericoma*). Type-locality: West Indies, Windward Is., St. Lucia, Bar de l'Isle. Holotype ♂, in B.M. (N.H.). Distr. — West Indies, Windward Is., St. Lucia.
- LANUGINOSA (Enderlein), 1937:89-90 (*Lepidiella*). Type-locality: Bolivia, Yungas. Holotype ♂, in Z.M.B. Distr. — Bolivia (Yungas). Ref. — Quate, 1963a:187 (♂); fig. 5(a)-(f).
- languginosa* Quate, 1963a:187 (lapsus).
- NIVERITARSIS Enderlein, 1937:89 (♂, as ♀). Type-locality: Peru, Dept. Cuzco, Callanga, Valle Piñi-Piñi, 1500 m. Holotype ♂, in Z.M.B. Distr. — Peru (Dept. Cuzco, Callanga). Ref. — Quate, 1963a:186 (♂, ♀); fig. 4(a)-(i).

Genus TELMATOSCOPIUS Eaton

- TELMATOSCOPIUS Eaton, 1904:58-59. Type-species, *Pericoma morula* Eaton (Tonnoir, 1933:70). Ref. — Duckhouse, 1966:174, 216-217 (Quate's designation of *Pericoma advena* Eaton, as type-species, 1965c:93, is invalid).

Subgenus CLOGMIA Enderlein

- Telmatoscopus*, subg. CLOGMIA Enderlein, 1937:87. Type-species, *Psychoda albipunctata* Williston (orig. des.) (as *albipennis*, lapsus). Refs. — Quate, 1963a:189; Quate & Quate, 1967:79.
- ALBIPUNCTATUS (Williston), 1893:113-114 (*Psychoda*). Type-locality: West Indies, Cuba, Havana. Holotype ♂, in U.S.N.M. Distr. — Tropicopolitan. In neotropics: Mexico (Tabasco, Frontera; Oaxaca, Tuxtepec; Yucatan; Córdoba), Costa Rica (La Suiza de Turrialba; Alajuela), Panama, West Indies (Bahamas, Nassau and New Providence I.; Cuba, Baracoa; Jamaica; Haiti; Puerto Rico; Tobago; Trinidad), Guyana (Georgetown), Colombia (?), Brazil (Rio de Janeiro), Paraguay (?), Argentina (Paso de los Libres, Corrientes). Refs. — Efflatoun, 1921:22-32 (immatures); Quate, 1955:185-187; figs. 53 (♂, ♀), 92 (pupa).
- albipuncta* Dyar, 1926c:149 (lapsus) (*Psychoda*).
- erecta* Curran, 1926:102 (*Psychoda*). Type-locality: West Indies, Jamaica, Hope Gardens.
- albopunctata* Curran, 1934:309, 525 (lapsus) (*Psychoda*).
- albipennis* Enderlein, 1937:87 (lapsus).
- giesi* Vargas, 1953:130-131 (♀); figs. 5-7 (*Syntomolaba*). Type-locality: Mexico, Yucatán, Mérida.

Subgenus NEUROSYSTASIS Satchell

- Telmatoscopus*, subg. NEUROSYSTASIS Satchell, 1955:86. Type-species, *terminalis* Satchell (mon.).
- TERMINALIS Satchell, 1955:86-88 (♂, ♀); figs. 1A-F. Type-locality: West Indies, Jamaica, Moneague. Holotype ♂, in B.M. (N.H.). Distr. — West Indies (Jamaica, Moneague).



Unplaced species of TELMATOSCOPIUS

- CALCARATUS Duckhouse, 1968:34-36 (♀); figs. 29-33. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Holotype ♂, in B.M. (N.H.). Distr. — Brazil (Santa Catarina, Nova Teutônia).
- FURCATUS (Kincaid), 1899:34 (*Pericoma*). Type-locality: U.S.A., Washington, Pullman. Holotype ♂, in Kincaid Collection (University of Washington?). Distr. — Widespread in U.S.A.; British Columbia, Brazil (Santa Catarina, Nova Teutônia). Refs. — Quate, 1955: 169-170 (♂, ♀); fig. 41a-f; Duckhouse, 1968:36.

Genus THRYSOCANTHUS Enderlein

- THRYSOCANTHUS Enderlein, 1937:103. Type-species, *Psychoda stellunata* Loew (orig. des.). Ref. — Quate, 1963a:189.
- RUBIGINOSUS Quate, 1963a:191 (♂); fig. 8(a)-(g). Type-locality: "Paraguay". Holotype ♂, in Z.M.B. Distr. — Paraguay.
- STELLULATUS (Loew), 1854:1 (♂) (*Psychoda*). Type-locality: "Brazil". Holotype ♂, in Z.M.B., n° 10899. Distr. — Brazil. Refs. — Enderlein, 1937:103; Quate, 1963a:189-190 (♂); fig. 7 (c)-(j).

Genus TONNOIRA Enderlein

- TONNOIRA Enderlein, 1937:106. Type-species, *pelliticornis* Enderlein (orig. des.). Ref. — Quate, 1963a:189.
- PELLITICORNIS Enderlein, 1937:106. Type-locality: Peru, Callanga. Holotype ♀, in Z.M.B. Distr. — Peru (Callanga). Ref. — Quate, 1963a:189 (♀); fig. 7(a)-(d).

UNRECOGNIZED SPECIES

- ALBITARSIS Williston, 1896:284 (♂, ♀); fig. 52 (*Pericoma*; procc. Banks, 1895). Type-locality: West Indies, St. Vincent. Holotype in B.M. (N.H.). Ref. — Dyar, 1926c:149 (*willistoni*, as nom. nov.).
- ANGUSTIPENNIS Williston, 1896:284 (♀); fig. 51 (*Psychoda*). Type-locality: West Indies, St. Vincent. Holotype ♀, in B.M. (N.H.).
- ANTENNALIS Williston, 1896:283 (♂) (*Psychoda*). Type-locality: West Indies, St. Vincent. Holotype ♂, in B.M. (N.H.).
- ATRASETA Rapp, 1945:310 (♀); fig. 4 (*Psychoda*). Type-locality: Canal Zone, Barro Colorado I. Holotype ♀, in A.M.N.H. Ref. — Fairchild, 1951:17.
- BREVITARSA Rapp, 1945:310 (♀); fig. 3(a)-(c) (*Psychoda*). Type-locality: Canal Zone, Barro Colorado I. Holotype ♀, in A.M.N.H. Ref. — Fairchild, 1951:17.
- DUBIA Bigot, 1888:4 (*Psychoda*). Type-locality: Chile, Cape Horn. No type-specimen is known. Ref. — Tonnoir, 1929:4.
- EBURNA Rapp, 1945:309 (♀); fig. 1(a)-(c) (*Psychoda*). Type-locality: Canal Zone, Barro Colorado I. Holotype ♀, in A.M.N.H. Ref. — Fairchild, 1951:17.



- FIMBRIATISSIMA Blanchard, 1852:351 (*Psychoda*). Type-locality: "Chile". Types, Muséum National d'Histoire Naturelle, Paris, n° 1275. Ref. — Tonnoir, 1929:4.
- GUIANICA Curran, 1934:309 (♂) (*Psychoda*). Type-locality: Guyana, Bartica District, Kartabo. Holotype ♂, in A.M.N.H., n° 201272.
- HYALINATA Blanchard, 1852:351-2 (*Psychoda*). Type-locality: Chile. Types, Muséum National d'Histoire Naturelle, Paris, n° 1277. Ref. — Tonnoir, 1929:4.
- MACULOSA Rapp, 1945:309-310 (♀); fig. 2(a)-(e) (*Psychoda*). Type-locality: Canal Zone, Barro Colorado I. Holotype ♀, in A.M.N.H. Ref. — Fairehild, 1951:17.
- MINUTISSIMA Enderlein, 1940:673 (♀); fig. 45 (*Psychoda*). Type-locality: Juan Fernández Is., Masatierra. Location of type unknown.
- NIGROPUNETATA Sehiner, 1868:29 (*Pericoma*). Type-locality: "Brazil". No type-specimen is known.
- NOTATA Blanchard, 1852:351 (*Psychoda*). Type-locality: "Chile". Types, Muséum National d'Histoire Naturelle, Paris, n° 1276. Ref. — Tonnoir, 1929:4.
- PALLENS Williston, 1896:283 (♂); figs. 50, 50a (*Psychoda*). Type-locality: West Indies, St. Vincent. Holotype ♂, in B.M. (N.H.). Ref. — Quate, 1963:184.
- PULLA Rondani, 1864:90 (*Psychoda*). Type-locality: "Chile". No type-specimen is known. Ref. — Tonnoir, 1929:4.
- PUNCTATELLA Townsend (nec Williston), 1897:19-20 (*Psychoda*). Type-locality: Mexico, Veraacruz State, Paso de Telaya. Location of type unknown.
- SEPTEMPUNCTATA Philippi, 1865:631 (*Psychoda*). Type-locality: Chile, Santiago. No type-specimen is known. Ref. — Tonnoir, 1934: 79-80.
- SILVESTRII Holmgren, 1905:530-532 (♂); figs. 1-3 (*Termitadelphos*). Type-locality: Peru, Carabaya prov., Chaquimayo. No type-specimen is known.
- TENELLA Philippi, 1865:631-632 (*Psychoda*). Type-locality: Chile, Santiago. No type-specimen is known. Ref. — Tonnoir, 1929:4.
- URSULA Müller, 1895:480-481 (larva); fig. 1 (*Marina*). Type-locality: Brazil (Itajaí, Blumenau?). No type-specimen is known. Ref. — Quate & Wirth, 1951:162-163.
- VENUSTA Weyenbergh, 1886:132 (♂, ♀); fig. 19 (*Psychoda*). Type-locality: "Argentina". No type specimen is known. Ref. — Tonnoir, 1929:4.

FOSSIL SPECIES

Genus TRICHOMYIA Haliday

- †ANTIQUARIA Quate, 1961:949-950 (♂); figs. 1, 2. Type-locality: Mexico, Chiapas, Las Cruces Landslide, locality B-5103. Late Oligocene or early Miocene amber fossil. Holotype ♂, in Univ. Calif. Mus. Paleo. n° 12685. Distr. — Mexico (Chiapas, Las Cruces Landslide).
- †DECLIVIVENA Quate, 1963b:112 (♀); fig. 1g. Type-locality: Mexico, Chiapas, Simojovel area, locality B-4119. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12748. Distr. — Mexico (Chiapas, Simojovel area).



- † *DISCALIS* Quate, 1963b:112 (♀), fig. 2A; plate 15, fig. 3. Type-locality: Mexico, Chiapas, Simojovel area, locality B-7043. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12749. Distr. — Mexico (Chiapas, Simojovel area).
- † *GLOMEROSA* Quate, 1963b:113-114 (♂); figs. 2b, c. Type-locality: Mexico, Chiapas, Las Cruces Landslide, locality B-5100. Late Oligocene or early Miocene amber fossil. Holotype ♂, in Univ. Calif. Mus. Paleo. N° 12816. Distr. — Mexico (Chiapas, Las Cruces Landslide).
- † *MECOCERA* Quate, 1963b:110-111 (♀); figs. 1a-c; plate 15, fig. 1. Type-locality: Mexico, Chiapas, Simojovel area, Pabuchil, locality B-7461. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. n° 13074. Distr. — Mexico (Chiapas, Simojovel area, Pabuchil).
- † *SMITHI* Quate, 1963b:112 (♀); figs. 1d-f; plate 15, fig. 2. Type-locality: Mexico, Chiapas, Simojovel area, locality B-1402. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12745. Distr. — Mexico (Chiapas, Simojovel area).
- † sp. Quate, 1963b:111-112 (♀). Locality: Mexico, Chiapas, Simojovel area, locality B-1402. Late Oligocene or early Miocene amber fossil.
- † sp. Quate, 1963b:112 (♀). Locality: Mexico, Chiapas, Ixhuatan. Late Oligocene or early Miocene amber fossil.

Genus *BRUNETTIA* Annandale

- † *HURDI* Quate, 1961:950-951 (♀); fig. 3; plate 108, fig. 1. Type-locality: Mexico, Chiapas, Las Cruces Landslide, locality B-5103. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12671. Distr. — Mexico (Chiapas, Las Cruces Landslide).

Genus *TELMATOSCOPIUS* Eaton

Unplaced species of *TELMATOSCOPIUS*

- † *HURDI* Quate, 1963b:114 (♂); fig. 3; plate 15, fig. 5. Type-locality: Mexico, Chiapas, Simojovel area, locality B-5345. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12901. Distr. — Mexico (Chiapas, Simojovel area).

Genus *PHILOSEPEDON* Eaton

Subgenus *PHILOSEPEDON* Eaton

- † *LABECCULA* Quate, 1963b:116 (♀); plate 15, fig. 7. Type-locality: Mexico, Chiapas, Simojovel area, locality B-4117. Late Oligocene or early Miocene amber fossil. Holotype ♀, in Univ. Calif. Mus. Paleo. N° 12904. Distr. — Mexico (Chiapas, Simojovel area).
- † *MEXICANA* Quate, 1963b:114-116 (♂); figs. 4a-c; plate 15, fig. 6. Type-locality: Mexico, Chiapas, Simojovel area, Cerro Balumtum, locality B-7041. Late Oligocene or early Miocene amber fossil. Holotype ♂, in Univ. Calif. Mus. Paleo. N° 12902. Distr. — Mexico (Chiapas, Simojovel area, Cerro Balumtum).



Genus PSYCHODA Latreille

- † USITATA Quate, 1963b:116 (♂); fig. 4d-f; plate 15, fig. 8. Type-locality: Mexico, Chiapas, Simojovel area, Pabuchil, locality B-7461. Late Oligocene or early Miocene amber fossil. Holotype ♂, in Univ. Calif. Mus. Paleo. N^o 13076. Distr. — Mexico (Chiapas, Simojovel area, Pabuchil).
- † sp. Quate, 1963b:116-117 (♀). Locality: Mexico, Chiapas, Simojovel area, Pabuchil.

LITERATURE CITED

ALEXANDER, C. P.

1920. A new subfamily of tanyderid flies (Diptera). *Ann. ent. Soc. Am.* 13:402-406.
1929. A revision of the American two-winged flies of the psychodid subfamily Bruchomyiinae. *Proc. U. S. nat. Mus.* 75:1-9.
1940. Further observations on the psychodid subfamily Bruchomyiinae (Diptera). *Rev. Ent.*, Rio de Janeiro 11:793-799.
1944. Two undescribed species of psychodid Diptera from tropical America. *Rev. Ent.*, Rio de Janeiro, 15:313-317.

ANNANDALE, N.


1910. A new genus of psychodid Diptera from the Himalayas and Travancore. *Rec. Indian Mus.* 5:141-142.

BANKS, N.

1894. Some Psychodidae from Long Island, N. Y. *Can. Ent.* 26: 329-333.
1895. Notes on *Psychoda*. *Can. Ent.* 27:324.

BARRETTO, M. P.

1950. Contribuição para o estudo dos Bruchomyiinae brasileiros, com as descrições de duas novas espécies (Diptera, Psychodidae). *Papéis Avulsos Dep. Zool. S. Paulo* 9:341-350.
- 1954a. Sobre o gênero *Eubonetia* Vargas e Díaz-Nájera com a descrição de novas espécies do Brasil (Diptera, Psychodidae). *Rev. bras. Ent.* 2:81-95.
- 1954b. Novas espécies de *Trichomyia* Hal. do Brasil (Diptera, Psychodidae). *Folia clin. biol.* 21:127-137.
- 1954c. Sobre o gênero *Maruina* F. Müller, 1895 (Diptera, Psychodidae). *Rev. bras. Ent.* 2:61-69.
1956. Sobre o gênero *Sycorax* Hal., com as descrições de duas novas espécies do Brasil (Diptera, Psychodidae). *Folia clin. biol.* 26: 71-75.
1961. Subfamílias e gêneros neotropicais da família Psychodidae Big., 1854 (Diptera). *Papéis Avulsos Dep. Zool. S. Paulo* 14: 211-225.



BARRETTO, M. P. & M. A. V. D'ANDRETTA

1946. Observações sobre a subfamília Bruchomyiinae Alexander, 1920 com a descrição de quatro novas espécies. *Livro de homenagem a E. F. d'Almeida* 6:55-75.

BIGOT, J. M. F.

1888. Diptères. In Gauthier, V., *Mission Scientifique du Cap Horn 1882-1883*, 6(2):1-4.

BLANCHARD, E.

1852. In Gay, C., *Historia fisica y politica de Chile* 7:351.

BOTOSANEANU, I. & F. VAILLANT

1970. Trois Diptères Psychodidae nouveaux de Cuba. *Trav. Lab. Hydrobiol.* 61:173-183.

BRUNETTI, E.

1908. Indian Psychodidae. *Rec. Ind. Mus.* 2(4):369-384, pl. XXIV.
1911. New Oriental Nemocera. *Rec. Ind. Mus.* 4:259-316.

COQUILLET, D. W.

1902. New orthorrhaphous Diptera from Mexico and Texas. *Jl. N. Y. ent. Soc.* 10:136-141.
1910. The type-species of the North American genera of Diptera. *Proc. U. S. nat. Mus.* 37:499-622.

CURRAN, C. H.

1926. New Diptera from Jamaica. Appendix, in Gowday, C. C., *Catalogus Insectorum Jamaicensis. Dept. Agr. Jamaica Ent. Bull.* 4:102.
1928. Insects of Porto Rico and the Virgin Islands. Diptera or two-winged flies. Pt 1, pp. 1-118, 39 figs., in *New York Academy of Sciences, Scientific Survey of Puerto Rico and the Virgin Islands*, Vol. 11. New York.
1934. The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. Am. Mus. nat. Hist.* 66:287.

DE LEON, J. R.

1950. Tres nuevas especies de *Nemopalpus* de Guatemala. *University of San Carlos, Guatemala, publication* 150:3-29.

DEL ROSARIO, F.

1936. The American species of *Psychoda* (Diptera: Psychodidae). *Philipp. J. Sci.* 59:85-148.

DESPORTES, C.

1942. *Forcipomyia velox* Winn. et *Sycorax silacea* Curtis, vecteurs d'*Icosiella neglecta* (Diesing), filaire commune de la grenouille verte. *Ann. Parasit. hum. comp.* 19:53-68.



DUCKHOUSE, D. A.

1962. Some British Psychodidae (Diptera, Nematocera): descriptions of species and a discussion on the problem of species pairs. *Trans. R. ent. Soc. Lond.* 114:403-436.
1965. Psychodidae (Diptera, Nematocera) of Southern Australia, subfamilies Bruchomyiinae and Trichomyiinae. *Trans. R. ent. Soc. Lond.* 117:329-343.
1966. Psychodidae (Diptera, Nematocera) of Southern Australia: subfamily Psychodidae. *Trans. R. ent. Soc. Lond.* 118:153-220.
1968. Psychodidae (Diptera, Nematocera) collected by Mr. Plaumann in Southern Brazil. *Proc. R. ent. Soc. Lond. (B)* 37:29-40.

DYAR, H. G.

- 1926a. Some apparently new American psychodids (Diptera, Psychodidae). *Insector Inscit. menstr.* 14:107-111.
- 1926b. Three psychodids from the Glacier National Park. *Insector Inscit. menstr.* 14:103-106.
- 1926c. A note in Psychodidae. *Insector Inscit. menstr.* 14:148-149.
1928. American Psychodidae — II (Diptera). *Proc. ent. Soc. Wash.* 30:87-89, 2 pl.
1929. American Psychodidae (Diptera) III. *Proc. ent. Soc. Wash.* 31:63-64.

EATON, A. E.

1904. New genera of European Psychodidae. *Ent. mon. Mag.* (2) 15:55-59.

EDWARDS, F. W.

1928. *Insects of Samoa, Part 6, Diptera* (British Museum, London).

EFFLATOUN, H. C.


1921. The life-history of *Telmatoscopus meridionalis* Eaton (Diptera). *Bull. Soc. ent. Egypte* 1920:22-34.

ENDERLEIN, G.

1937. Klassifikation der Psychodiden (Dipt.). *Deutsh. ent. Z.* 1936: 81-112.
1940. Die Dipterenfauna der Juan-Fernandez-Inseln und der Oster-Insel. In Skottsberg, C., *The Natural History of Juan Fernandez and Easter Island* 3:673. Uppsala.

FAIRCHILD, G. B.

1951. Some nomenclatorial notes on Psychodidae (Diptera). *Bull. Brooklyn ent. Soc.* 46:10-18.
1952. Notes on *Bruchomyia* and *Nemopalpus* (Diptera, Psychodidae). *Ann. ent. Soc. Am.* 45:259-280.
1955. The relationships and classification of the Phlebotominae (Diptera, Psychodidae). *Ann. ent. Soc. Am.* 48:182-196.



FEUERBORN, H. J.

1932. Die Psychodide *Maruina indica* sp. n. und ihre Beziehung zu den Blepharoceriden. Zugleich einige Mitteilungen über die Metamorphose von *Sycorax* und über Chitin als Klabemittel. *Arch. Hydrobiol. Plankt.* 11:55-128.

FREEMAN, P.

1949. A key to the South American species of *Nemopalpus* (Dipt., Psychodidae) with the description of a new species. *Ent. mon. Mag.* 85:85-88.

HALIDAY, A. H.

1839. In Curtis, J., *British Entomology*. London:745.

HOLMGREN, N.

1905. Zwei neue Termitomastiden aus Südamerika. *Zool. Anz.* 29: 529-536.

JOHANNSEN, O. A.

1938. New species of Nematocera from Puerto Rico. *J. Agric. Univ. P. Rico* 22:219-225.

JUNG, H. F.

1956. Beiträge zur Biologie, Morphologie und Systematik der europäischen Psychodiden (Diptera). *Dtsch. ent. Z.* 3:97-257.
1963. On *Nemopalpus flavus* Macquart (Diptera, Psychodidae — Bruchomyiinae). *Annotnes. zool. jap.* 36:45-53.

KEHLIN, D. & P. TATE.

1937. A comparative account of the larvae of *Trichomyia urbica* Curtis, *Psychodes albipennis* Zett., and *Phlebotomus argentipes* Ann. & Brun. (Diptera: Psychodidae). *Parasitology* 29:247-258.

KINCAID, T.

1899. The Psychodidae of the Pacific Coast. *Ent. News* 10:30-37.

KNAB, F.

1914. New moth-flies (Psychodidae) bred from Bromeliaceae and other plants. *Proc. U. S. Nat. Mus.* 46:103-106.

LATREILLE, P. A.

1796. *Précis des caractères générales des Insectes, disposés dans un ordre naturel*, 179 pp. Paris "An V".
1802. *Histoire naturelle, générale et particulière des Crustacés et des Insectes*, tome troisième (Vol. 3) [Vol. 95], 468 pp. in C.S. Sonnini, ed., *Histoire Naturelle par Buffon*. Paris, "An X".

LOEW, H.

1854. Neue Dipteren. In *Neue Beiträge zur Kenntnis der Dipteren* 2:1.



MACQUART, M.

- 1838a. In Webb, P.B. and S. Berthelot, *Hist. Nat. Iles Canaries. Ent., 1, Diptera*:101-102.
1838b. Diptères exotiques nouveaux ou peu connus. *Mém. Soc. Roy. Sci. Agric. Arts. Lille 1, Pt. 2*:85-86; Plate 12, figs. 1, 1a, 1b.

MEUNIER, F.

1905. Monographie des Psychodidae de l'ambre de la Baltique. *Annls. hist.-nat. Mus. nat. hung. 3*:235-255.

MÜLLER, F.

1895. Contribution towards the history of a new form of larvae of Psychodidae (Diptera) from Brazil. *Trans. ent. Soc. Lond. 1895*: 479-482.

PARROT, L.

1951. Notes sur les Phlébotomes. LXI. A propos de classification. *Archs. Inst. Pasteur Algér. 29*:23-45.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool.-bot. Ges. Wien 15*:595-782.

QUATE, L. W.

1954. A revision of the Psychodidae of the Hawaiian Islands (Diptera). *Proc. ent. Soc. Hawaii 15*:335-356.
1955. A revision of the Psychodidae (Diptera) in America North of Mexico. *Univ. Calif. Publ. Ent. 10*:103-273.
1959a. Taxonomy of neotropical Psychodidae (Diptera) 1. *Psychoda* species of West Indies and Central America with a key to Trinidad species. *Pan-Pacif. Ent. 35*:213-220.
1959b. Insects of Micronesia. Diptera: Psychodidae. *Insects Micronesia 12*:435-484.
1959c. Classification of the Psychodini (Psychodidae: Diptera). *Ann. ent. Soc. Am. 52*:444-451.
1961. Fossil Psychodidae (Diptera: Insecta) in Mexican amber; Part 1. *J. Paleont. 35*:949-951; Plate 108.
1962. Psychodidae (Diptera) at the Zoological Survey of India. *Proc. Hawaii. ent. Soc. 18*:155-188.
1963a. Review of G. Enderlein's non holarctic genera of Psychodidae and description of a new species (Diptera). *Trans. R. ent. Soc. Lond. 115*:181-196.
1963b. Fossil Psychodidae in Mexican amber, Part. 2. *J. Paleont. 37*:110-118.
1965a. The psychodid flies of the south temperate zone. *Int. Congr. Ent.*, London, 1964:466-467.
1965b. A taxonomic study of Philippine Psychodidae (Diptera). *Pacif. Insects 7*:815-902.



1965c. Family Psychodidae, pp. 91-97, in A. Stone et. al., A catalog of the Diptera of America North of Mexico. U. S. Department of Agriculture, *Agric. Handbook* 276:iv + 1696 pp., Washington, D. C.

QUATE, L. W. & S. QUATE

1967. A monograph of Papuan Psychodidae, including *Phlebotomus* (Diptera). *Pacif. Insects Mon.* 15:1-216.

QUATE, L. W. & W. W. WIRTH.

1951. A taxonomic revision of the genus *Maruina* (Diptera: Psychodidae). *Wasmann J. Biol.* 9:151-161.

RAPP, W. F.

1945. New Psychodidae from Barro Colorado Island. *Jl. N. Y. ent. Soc.* 53:309-311; Plate 3.

RONDANI, C.

1864. Dipterorum species et genera aliqua exotica revisa et annotata, novis, nonnullis descriptis. *Arch. Zool. Anat. di Canestrini* 3:1-99.

SATCHELL, G. H.

1947. The larvae of the British species of *Psychoda* (Diptera: Psychodidae). *Parasitology* 33:51-69.

1949. The early stages of the British species of *Pericoma* Walker (Diptera: Psychodidae). *Trans. R. ent. Soc. Lond.* 100:411-447.

1950. New species of Fijian Psychodidae (Diptera). *Proc. R. ent. Soc. Lond. (B)* 19:176-185.

1953a. New and little known Samoan Psychodidae and a new species from Rarotonga. *Proc. R. ent. Soc. Lond. (B)* 22:181-188.

1953b. The Australian Psychodidae (Diptera), Part I. *Aust. J. Zool.* 1:357-418.

1953c. On the early stages of *Bruchomyia argentina* Alexander (Diptera: Psychodidae). *Proc. R. ent. Soc. Lond. (A)* 28:1-12.

1955. Two new subgenera of Psychodidae (Diptera) from Jamaica, with descriptions of five new species. *Ann. Mag. nat. Hist. (12)* 8: 85-93.

1956. On the genus *Trichomyia* Haliday (Diptera: Psychodidae), with descriptions of four new species. *Proc. R. ent. Soc. Lond. (B)* 25:147-156.

SAY, T.

1824. *In* Kcating: *Narrative of an expedition to source of St. Peter's River* 2. Appendix: 358. Philadelphia.

SCHINER, J. R.

1868. *Reise der österreichischen Fregatte Novara um die Erde.* Zoologische Theil, 2 (1) Diptera. Vienna.



SHANNON, R. C. & E. DEL PONTE

1927. Cuatro notas sobre especies nuevas de dipteros nematóceros, hematófagos o no, de la República Argentina. *Rev. Inst. bact.*, B. Aires 4 (Tercera nota): 733-734.

SPEISER, P.

1906. Über die systematische Stellung der Dipterenfamilie Termitomastidae. *Zool. Anz.* 30:716-718.

TOKUNAGA, M.

1957. Moth flies from Formosa (Psychodidae, Diptera). *Scient. Rep. Saikyo Univ.* 9:53-77.

TONNOIR, A. L.

1920. Contribution a l'étude des Psychodidae. *Ann Soc. ent. Belg.* 60:149-157.
1922. Notes sur le genre *Nemopalpus* (Dipt. Psychodidae) et description d'une espèce nouvelle. *Ann. Soc. ent. Belg.* 62:125-136.
1929. *Diptera of Patagonia and South Chile*. 2, Fascicle 1, Psychodidae. London, British Museum (Nat Hist.).
1933. Descriptions of remarkable Indian Psychodidae and their early stages, with a theory of the evolution of the ventral suckers of dipterous larvae. *Rec. Indian Mus.* 35:53-75.
1934. Notes synonymiques sur quelques Psychodidae (Diptera). *Bull. Anns. Soc. r. ent. Belg.* 74:69-82.
1940. Sur un remarquable organe sexuel secondaire chez certains mâles du genre *Nemopalpus* Macq. avec description d'une espèce nouvelle et d'une autre peu connue (Diptera, Psychodidae). *Int. Congr. Ent.*, Madrid, 1935, 1:203-213.

TOWNSEND, C. H. T.

1897. Contributions from the New Mexico Biological Station. 2. On a collection of Diptera from the lowlands of the Rio Nautla, in the State of Vera Cruz. *Ann. Mag. nat. Hist.* 20:19-33.

VAILLANT, F.

1963. Le *Marina* d'Amérique du Nord (Dipt. Psychodidae). *Bull. Soc. ent. Fr.* 68:71-92.

VARGAS, L.

1953. Sobre alguns Psychodidae de Mexico. *Eubonetia maldonadoi* n.sp. y *Syntomolaba giesi* n. sp. *Rev. Inst. Salubr. Enferm. trop. Mex.* 13:127-132.

VARGAS, L. & A. DÍAZ NÁJERA

1953. *Trichomyia fairchild* n. sp., y *Eubonetia* n. gen. (Diptera: Psychodidae). *Rev. Inst. Salubr. Enferm. trop., Méx.* 13:153-158.



WALKER, F.

1856. Family VIII. Phlebotomidae. *Ins. Brit. Dipt.* 3:253-263.

WEYENBERGH, H.

1886. Dipterologische fragmenten. *Tijdschr. Ent.* 29:126.

WILLISTON, S. W.

1893. The North American Psychodidae. *Ent. News.* 5:113-114.

1896. On the Diptera of St. Vincent (West Indies) *Trans. ent. Soc. Lond.* 1896:253.


INDEX

<i>aequalis</i> Enderlein, 1937 (Thrysocanthus), Pericoma (lapsus)	6A.10
<i>aitkeni</i> Quate, 1959, Psychoda	6A.12
<i>alabangensis</i> Del Rosario, 1936, Psychoda	6A.12
<i>albicollare</i> (Enderlein), 1937 (Chirolepia), Alepia	6A.6
<i>albipeda</i> Rapp, 1945, Kupara	6A.7
<i>albipennis</i> Enderlein, 1937, Telmatoscopus (lapsus)	6A.14
<i>albipuncta</i> Dyar, 1926 (Psychoda), Telmatoscopus (lapsus)	6A.14
<i>albipunctatus</i> (Williston), 1893 (Psychoda), Telmatoscopus	6A.14
<i>albitarsis</i> Williston, 1896, Pericoma	6A.15
<i>albonotata</i> (Brunetti), 1908 (Psychoda), Brunettia	6A.7
<i>albopunctata</i> Curran, 1934 (Psychoda), Telmatoscopus (lapsus) ..	6A.14
<i>Alepia</i> Enderlein, 1937	6A.6
<i>alfaroana</i> (Dyar), 1926 (Psychoda), Pericoma	6A.9
<i>almeidai</i> Barretto & d'Andretta, 1946, Bruchomyia	6A.2
<i>alternata</i> Say, 1824, Psychoda	6A.12
<i>alternicula</i> Quate, 1955, Psychoda	6A.12
<i>amplipenna</i> (Knab), 1914 (Psychoda), Pericoma	6A.9
<i>angustipennis</i> Williston, 1896, Psychoda	6A.15
<i>antennalis</i> Williston, 1896, Psychoda	6A.15
<i>antillarum</i> Fairchild, 1952, Nemopalpus	6A.3
<i>argentina</i> Alexander, 1920, Bruchomyia	6A.2
<i>Arisemus</i> Satchell, 1955	6A.6
<i>armata</i> Barretto, 1954, Trichomyia	6A.4
<i>arroyoi</i> De Leon, 1950, Nemopalpus	6A.3
<i>assimilis</i> Barretto, 1956, Sycorax	6A.6
<i>atraseta</i> Rapp, 1945, Psychoda	6A.15
<i>barrettoi</i> Duckhouse, <i>nom. nov.</i> , Trichomyia	6A.4
<i>bisubulata</i> Duckhouse, 1968, Alepia	6A.6
<i>boxi</i> Satchell, 1955, Telmatoscopus	6A.6
<i>brasiliensis</i> Duckhouse, 1968, Pericoma	6A.9
<i>brasiliensis</i> Satchell, 1956, Trichomyia	6A.4
<i>brasiliensis</i> Alexander, 1940, Bruchomyia	6A.2
<i>brevinervis</i> Barretto & d'Andretta, 1946, Nemopalpus	6A.3
<i>brevitarsa</i> Rapp, 1945, Psychoda	6A.15
<i>Bruchomyia</i> Alexander, 1920	6A.2
<i>Bruchomyiinae</i>	6A.2
<i>Brunettia</i> Annandale, 1910	6A.7
<i>busckana</i> (Dyar), 1926 (Psychoda), Pericoma	6A.9

6A.25




<i>calcaratus</i> Duckhouse, 1968, <i>Telmatoscopus</i>	6A.15
<i>caudata</i> Satchell, 1955, <i>Pericoma</i>	6A.9
<i>cervi</i> (Satchell), 1955 (<i>Pericoma</i>), <i>Syntomoza</i>	6A.14
<i>chilensis</i> Tonnoir, 1929, <i>Pericoma</i>	6A.9
<i>chilensis</i> Tonnoir, 1929, <i>Sycorax</i>	6A.6
<i>Chirolepia</i> Enderlein, 1937	6A.7
<i>cinerea</i> Banks, 1894, <i>Psychoda</i>	6A.12
<i>cirrata</i> Coquillett, 1902, <i>Trichomyia</i>	6A.4
<i>Clogmia</i> Enderlein, 1937	6A.14
<i>cochlearia</i> Satchell, 1950, <i>Psychoda</i>	6A.12
<i>complicata</i> Tonnoir, 1929, <i>Pericoma</i>	6A.9
<i>contigua</i> Tonnoir, 1929, <i>Pericoma</i>	6A.9
<i>coutinhoi</i> (Barretto), 1954 (<i>Eubonetia</i>), <i>Trichomyia</i>	6A.5
<i>cubana</i> (Botosaneanu & Vaillant), 1970 (<i>Quatiella</i>), <i>Philosepedon</i>	6A.11
<i>dampfianus</i> Alexander, 1940, <i>Nemopalus</i>	6A.3
<i>dealbata</i> Tonnoir, 1929, <i>Nemoneura</i>	6A.8
<i>Desmioza</i> Enderlein, 1937	6A.9
<i>Dictyocampa</i> Enderlein, 1937	6A.7
<i>Didicrum</i> Enderlein, 1937	6A.9
<i>dissimilis</i> Barretto & d'Andretta, 1946, <i>Nemopalus</i>	6A.3
<i>divaricata</i> Duckhouse, 1968, <i>Psychoda</i>	6A.12
<i>dubia</i> Bigot, 1888, <i>Psychoda</i>	6A.15
<i>duripuncta</i> (Curran), 1928 (<i>Psychoda</i>), <i>Brunettia</i>	6A.7
<i>duripunctata</i> Quate, 1962 (<i>Psychoda</i>), <i>Brunettia</i> (<i>lapsus</i>)	6A.7
<i>eatoni</i> Satchell, 1956, <i>Trichomyia</i>	6A.5
<i>eburna</i> Rapp, 1945, <i>Psychoda</i>	6A.15
<i>edwardsi</i> Tonnoir, 1929, <i>Pericoma</i>	6A.9
<i>edwardsi</i> Tonnoir, 1929, <i>Trichomyia</i>	6A.5
<i>cqualis</i> Tonnoir, 1934, <i>Pericoma</i>	6A.9
<i>erecta</i> Curran, 1926 (<i>Psychoda</i>), <i>Telmatoscopus</i>	6A.14
<i>Eubonetia</i> Vargas & Díaz Nájera, 1953	6A.4
<i>Eurygarka</i> Quate, 1959	6A.11
<i>fairchildi</i> (Barretto), 1954 (<i>Eubonetia</i>), <i>Trichomyia</i>	6A.4
<i>fairchildi</i> Vargas & Díaz Nájera, 1953, <i>Trichomyia</i>	6A.5
<i>fenestrata</i> Tonnoir, 1929, <i>Pericoma</i>	6A.10
<i>fimbriatissima</i> Blanchard, 1852, <i>Psychoda</i>	6A.16
<i>flavescens</i> Von Roser	6A.5
<i>fumata</i> (Knab), 1914 (<i>Psychoda</i>), <i>Philosepedon</i>	6A.11
<i>furcatus</i> (Kincaid), 1899 (<i>Pericoma</i>), <i>Telmatoscopus</i>	6A.15
<i>fusca</i> Barretto, 1950, <i>Bruchomyia</i>	6A.2
<i>gehrkeae</i> Quate, 1959, <i>Psychoda</i>	6A.12
<i>giesi</i> Vargas, 1953 (<i>Syntomolaba</i>), <i>Telmatoscopus</i>	6A.14
<i>grabhamana</i> (Dyar), 1926 (<i>Psychoda</i>), <i>Pericoma</i>	6A.10
<i>griseata</i> Tonnoir, 1929, <i>Pericoma</i>	6A.10
<i>grisescens</i> Tonnoir	6A.12
<i>guianica</i> Curran, 1934, <i>Psychoda</i>	6A.16
<i>guttata</i> Enderlein, 1937, <i>Dictyocampa</i>	6A.7



<i>helicis</i> (Dyar), 1929 (Psychoda), Philosepedon	6A.11
<i>hertiventris</i> Enderlein, 1937 (Psychoda), Alepia (lapsus)	6A.6
<i>hexadactylus</i> Botosaneanu & Vaillant, 1970, Arisemus	6A.7
<i>hirta</i> Johannsen, 1938, Maruina	6A.8
<i>hirtiventris</i> (Tonnoir), 1920 (Psychoda), Alepia	6A.6
<i>hyalinata</i> Blanchard, 1852, Psychoda	6A.16
<i>ichthycerca</i> Quate, 1959, Psychoda	6A.12
<i>immaculatus</i> Freeman, 1949, Nemopalpus	6A.3
<i>incompleta</i> (Knab), 1914 (Psychoda), Pericoma	6A.10
<i>inermis</i> Barretto, 1954, Trichomyia	6A.5
<i>inornata</i> Tonnoir, 1929, Pericoma	6A.10
<i>interdicta</i> (Dyar), 1928 (Psychoda), Philosepedon	6A.11
Kupara Rapp, 1945	6A.7
<i>languginosa</i> Quate, 1963, Syntomoza (lapsus)	6A.14
<i>lanuginosa</i> (Enderlein), 1937 (Lepidiella), Syntomoza	6A.14
<i>Lepidiella</i> Enderlein, 1937	6A.13
<i>Lepria</i> Enderlein, 1937	6A.4
<i>lucia</i> Quate, 1954, Psychoda	6A.13
<i>maculipennis</i> Barretto & d'Andretta, 1946, Nemopalpus	6A.4
<i>maculipennis</i> Enderlein, 1937, Chirolepia	6A.7
<i>maculosa</i> Rapp, 1945, Psychoda	6A.16
<i>maculosus</i> Satchell, 1955, Telmatoscopus	6A.7
<i>maldonadoi</i> (Vargas), 1953 (Eubonetia), Trichomyia	6A.5
<i>Maruina</i> Müller, 1895	6A.8
<i>masatierrensis</i> Satchell, 1952, Psychoda	6A.12
<i>Mecysmia</i> Enderlein, 1937	6A.9
<i>minutissima</i> Enderlein, 1940, Psychoda	6A.16
<i>mopani</i> De Leon, 1950, Nemopalpus	6A.3
<i>moralesi</i> De Leon, 1950, Nemopalpus	6A.3
<i>Nemopalpus</i> Macquart, 1838 (lapsus)	6A.3
<i>Nemoneura</i> Tonnoir, 1929	6A.8
<i>Nemopalpus</i> Macquart, 1938	6A.3
<i>Neurosystasis</i> Satchell, 1955	6A.14
<i>nigropunctata</i> Schiner, 1868, Pericoma	6A.16
<i>niveitarsis</i> Enderlein, 1937, Syntomoza	6A.14
<i>niveopunctata</i> Tonnoir, 1929, Pericoma	6A.10
<i>notata</i> Blanchard, 1852, Psychoda	6A.16
<i>pallens</i> Williston, 1896, Psychoda	6A.16
<i>pallidula</i> Tonnoir, 1929, Pericoma	6A.10
<i>pallipes</i> (Shannon & Del Ponte), 1927 (Bruchomyia), Nemopalpus	6A.4
<i>Parasetomima</i> Duckhouse, 1968	6A.8
<i>parthenogenetica</i> Tonnoir, 1940, Psychoda	6A.12
<i>pedicillata</i> Satchell, 1956, Trichomyia	6A.5
<i>pelliticornis</i> Enderlein, 1937, Tonnoira	6A.15
<i>Pericoma</i> Walker, 1856	6A.8
<i>peruviana</i> Alexander, 1929, Bruchomyia	6A.2
<i>phalaenoides</i> Tonnoir	6A.12
<i>Philosepedon</i> Eaton, 1904	6A.10,
	6A.11



<i>pictipennis</i> Tonnoir, 1920, <i>Pericoma</i>	6A.10
<i>pilipes</i> Tonnoir, 1922, <i>Nemopalpus</i>	6A.4
<i>pilosella</i> Müller, 1895, <i>Maruina</i>	6A.8
<i>plaesia</i> Quate, 1959, <i>Psychoda</i>	6A.12
<i>platilobata</i> Tokunaga, 1957, <i>Psychoda</i>	6A.13
<i>Platyplastinx</i> Enderlein, 1937	6A.11
<i>plaumanni</i> Alexander, 1944, <i>Bruchomyia</i>	6A.3
<i>plaumanni</i> (Duckhouse), 1968 (<i>Psychoda</i>), <i>Philosepedon</i>	6A.11
<i>Podolepria</i> Enderlein, 1937	6A.9
<i>pseudocompar</i> Tonnoir, 1929, <i>Psychoda</i>	6A.13
<i>Psychoda</i> Latreille, 1796	6A.11
<i>Psychodinae</i>	6A.6
<i>pulla</i> Rondani, 1864, <i>Psychoda</i>	6A.16
<i>punctata</i> (Philippi), 1865 (<i>Psychoda</i>), <i>Nemoneura</i>	6A.8
<i>punctatella</i> Townsend, 1897, <i>Psychoda</i>	6A.16
<i>Quatiella</i> Botosaneanu & Vaillant, 1970	6A.11
<i>rarotongensis</i> Satchell, 1953, <i>Psychoda</i>	6A.13
<i>rubiginosus</i> Quate, 1963, <i>Thrysocanthus</i>	6A.15
<i>satchelli</i> Barretto, 1956, <i>Sycorax</i>	6A.6
<i>savaiiensis</i> Edwards, 1928, <i>Psychoda</i>	6A.13
<i>schoenemanni</i> Enderlein, 1937 (<i>Mecysmia</i>), <i>Pericoma</i>	6A.10
<i>scotina</i> Quate, 1959, <i>Psychoda</i>	6A.13
<i>scripta</i> Enderlein, 1937, <i>Alepia</i>	6A.6
<i>septempunctata</i> Philippi, 1865, <i>Psychoda</i>	6A.16
<i>shannoni</i> Alexander, 1929, <i>Bruchomyia</i>	6A.3
<i>silvestrii</i> Holmgren, 1905, <i>Termitadelphos</i>	6A.16
<i>simillima</i> Tonnoir, 1929, <i>Psychoda</i>	6A.13
<i>simplex</i> Tonnoir, 1929, <i>Pericoma</i>	6A.10
<i>solox</i> Enderlein, 1937, <i>Platyplastinx</i>	6A.11
<i>sp. A</i> Quate & Wirth, 1951, <i>Maruina</i>	6A.8
<i>spathipenis</i> (Duckhouse), 1968 (<i>Psychoda</i>), <i>Philosepedon</i>	6A.11
<i>sp. Duckhouse</i> , 1968, <i>Alepia</i>	6A.6
<i>sp. Dyar</i> , 1926 (as <i>phalaenoides</i>), <i>Psychoda</i>	6A.13
<i>sp. Fairchild</i> , 1955, <i>Sycorax</i>	6A.6
<i>speciosa</i> Tonnoir, 1929, <i>Pericoma</i>	6A.10
<i>spinosa</i> Müller, 1895, <i>Maruina</i>	6A.8
<i>squamitarsis</i> Tonnoir, 1929, <i>Pericoma</i>	6A.10
<i>squamosa</i> (Enderlein), 1937 (<i>Lepria</i>), <i>Trichomyia</i>	6A.5
<i>stellulatus</i> (Loew), 1854 (<i>Psychoda</i>), <i>Thrysocanthus</i>	6A.15
<i>subimmaculata</i> Tonnoir, 1929, <i>Psychoda</i>	6A.13
<i>surcoufi</i> Tonnoir, 1922, <i>Psychoda</i>	6A.13
<i>Sycoracinae</i>	6A.6
<i>Sycorax</i> Haliday, 1839	6A.6
<i>Synmormia</i> Enderlein, 1937	6A.9
<i>Syntomolaba</i> Enderlein, 1937	6A.9
<i>Syntemoza</i> Enderlein, 1937	6A.13
<i>sziladyi</i> Tonnoir, 1940, <i>Nemopalpus</i>	6A.4
<i>Telmatoscopus</i> Eaton, 1904	6A.14
<i>tenella</i> Philippi, 1865, <i>Psychoda</i>	6A.16
<i>terminalis</i> Satchell, 1955, <i>Telmatoscopus</i>	6A.14



<i>Termitodipteron</i> Holmgren, 1905	6A.4
tetradactylus Botosaneanu & Vaillant, 1970, Arisemus	6A.7
Thrysocanthus Enderlein, 1937	6A.15
Tonnoira Enderlein, 1937	6A.15
travassosi (Barretto), 1954 (Eubonetia), Trichomyia	6A.5
Trichomyia Haliday, 1839	6A.4
Trichomyiinae	6A.4
tricolor (Knab), 1914 (Psychoda), Pericoma	6A.10
umbella Duckhousc, 1968, Parasetomima	6A.8
unicolor Barretto, 1950, Bruchomyia	6A.3
urbica Haliday	6A.5
ursula Müller, 1895, Maruina	6A.16
vargasi (Barretto), 1954 (Eubonetia), Trichomyia	6A.5
vazi (Barretto), 1954 (Eubonetia), Trichomyia	6A.5
venusta Weyenbergh, 1886, Psychoda	6A.16
verans Alexander, 1940, Nemopalpus	6A.4
viduata Tonnoir, 1929, Pericoma	6A.10
wasmanni (Holmgren), 1905 (Termitodipteron), Trichomyia	6A.5
zetoscota Quate, 1959, Psychoda	6A.13

INDEX OF FOSSIL SPECIES

† antiquaria Quate, 1961, Trichomyia	6A.16
† declivivena Quate, 1963, Trichomyia	6A.16
† discalis Quate, 1963, Trichomyia	6A.17
† glomerosa Quate, 1963, Trichomyia	6A.17
† hurdi Quate, 1961, Brunettia	6A.17
† hurdi Quate, 1963, Telmatoscopus	6A.17
† labecula Quate, 1963, Philosepedon	6A.17
† mecocera Quate, 1963, Trichomyia	6A.17
† mexicana Quate, 1963, Philosepedon	6A.17
† smithi Quate, 1963, Trichomyia	6A.17
† sp. Quate, 1963, Psychoda	6A.18
† sp. Quate, 1963, Trichomyia	6A.17
† usitata Quate, 1963, Psychoda	6A.18





SciELO

DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA. SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

7

FAMILY NYMPHOMYIIDAE

F. P. IDE

*Department of Zoology, University of Toronto
Toronto, Canada*

These are small, delicate, elongate flies with short, clubbed antennae, vestigial mouthparts, eyes holoptic centrally and narrow wings fringed with long setae. Two asiatic species (*Nymphomyia alba* Tokunaga, 1932) from Japan and one from India (Brundin, 1966), and one North American (*Palaeodipteron walkeri* Ide, 1965) from New Brunswick, Canada, are known.

The larval stage is unknown, but is thought to live in rapids in the vicinity of which the Japanese species was taken. The adults of the North American species were taken from rapid water generally as paired, de-alated individuals with eggs attached to the female by stalks. The pupa, unlike that of typical Nematocera, has short, anteriorly-directed antennal sheaths (Tokunaga, 1959; Ide, 1965).

The family is placed in a separate suborder, Archidiptera, by Rohdendorf (1960).

No species is known south of Canada in the Western Hemisphere. An undescribed fly from Brazil [included by Ide in the North American catalogue, as belonging to Nymphomyiidae], however, resembles the Nymphomyiidae superficially but agrees more closely with *Ipomyia* Colless (1964). This West Australian species is tentatively placed in the family Cecidomyiidae by Colless with the observation that it may have close affinities with the Nymphomyiidae.

LITERATURE CITED

BRUNDIN, L.

1966. Transatlantic relationships and their significance, as evidenced by chironomid midges. *K. svenska V. A. Handl.* 11(1):1-472 (p. 457).

COLLESS, D. H.

1964. An extraordinary Australian cecidomyiid, possibly related to the Nymphomyiidae (Diptera; Nematocera). *Proc. Roy. Entom. Soc. London (B)* 34(11-12):145-149.

IDE, F. P.

1965. A fly of the archaic family Nymphomyiidae from North America. *Canad. Entom.* 97(5):496-507.



ROHDENDORF, B. B.

1960. Neue Angaben über das System der Dipteren. *Verh. XI. Int. Kongr. Entom., Wien* 1:153-158.

TOKUNAGA, M.

1932. A remarkable dipterous insect from Japan, *Nymphomyia alba*, gen. et sp. nov. *Annot. Zool. Jap.* 13:559-569.

1959. *Illustrated insect larvae of Japan*. Hokuryukan Ltd., Tokyo.

ADDITIONAL REFERENCES

TOKUNAGA, M.

1935. A morphological study of a nymphomyiid fly. *Philipp. J. Sci.* 56:127-214.

1936. The central nervous system, tracheal system and digestive system of a nymphomyiid fly. *Ibid.* 59:189-216.





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(Formerly Departamento de Zoologia, Secretaria da Agricultura,
São Paulo)A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

8

FAMILY BLEPHARICERIDAE

(BLEPHAROCERIDAE)

CHARLES L. HOGUE

Los Angeles County Museum of Natural History
Los Angeles, California

Although the family Blephariceridae, or the *net-winged midges* has attracted the attention of numerous authors since its discovery in 1842, it remains poorly known even in regards to its basic internal classification and biology. Aside from the European fauna, with which our knowledge is most advanced, many discoveries of new species, stage associations and functional morphology are being made by today's students. This is certainly the case with Neotropical blepharicerid fauna, being the least studied of all the zoogeographic regions, in spite of the fact that, at present, 62 regional species in five genera are recognized, being a sizable percentage of the world's 520 species and 21 genera. Almost certainly however this is an unreal figure, considering the need to reexamine the disproportionately large number of species in the genus *Kelloggina* (= *Dimorphotarsa*). These were mostly described by Lutz (1920) and Lane & d'Andretta (1956), the former's material being almost entirely the immature stages, the latter's adults. Since roughly the same area of southeast Brazil is the range of all these species, numerous synonymies are suspected. If all the species are found to be valid, *Kelloggina*, an entirely Neotropical genus, would constitute the largest in the family.

It is not surprising that the immature stages are sooner encountered than the adults. The former are a conspicuous element of the epilithic fauna of torrential waters. The larvae (four instars) and pupae adhere in groups or singly to smooth rock surfaces in very swiftly flowing portions of streams and shallow rivers, usually at the brink or at the base of cataracts and small waterfalls. Algae and diatoms browsed from the substratum constitute the larva's food. The much more seldom seen adults rest on vegetation, under fallen logs and overhanging rocks, in hollow trees and in other protected, shady places. In flight they move swiftly and during peak emergences, numbers may be seen hovering in the spray below waterfalls or darting between perches and capturing



preys Normal mandibulate females feed on the blood of other small Diptera. Males, and those females of a few species which lack mandibles, probably take other natural liquids or do not feed.

The best summaries of general information on Blephariceridae is available in the works of Alexander (1963) and Mannheims (1935). A particularly important contribution to the family's classification and zoogeography was made by Alexander (1958) who is followed here with the exception of the separation of *Dimorphotarsa* from *Kelloggina*.

All citations to the 1920 work of Lutz, unless otherwise noted, pertain properly to the Portuguese version which has precedence over the German.

Type repositories are known for nearly all the regional species and are included in the listings. The following abbreviations have been used:

AMNH	American Museum of Natural History, New York, N.Y.
BMNH	British Museum (Natural History), London, England
BZM	Berliner Zoologisches Museum, Berlin, Germany
CU	Cambridge University, Cambridge, England
DEI	Deutsches Entomologisches Institut, Berlin, Germany
FSP	Fauldade de Saúde Pública (Departamento de Epidemiologia), Universidade de São Paulo, Brasil.
IMZ	Istituto e Museo di Zoologia, Università di Torino, Italy
IOC	Instituto Oswaldo Cruz, Rio de Janeiro, Guanabara, Brasil
KSR	Kungliga Svenska Riksmuseet, Stockholm, Sweden
MZUSP	Museu de Zoologia, Universidade de São Paulo, Brasil
NMW	Naturhistorisches Museum, Vienna, Austria
USNM	United States National Museum, Washington, D.C.

Subfamily EDWARDSININAE

Genus EDWARDSINA Alexander

EDWARDSINA Alexander, 1920:2. Type-species, *chilensis* Alexander (mon.).

ARGENTINENSIS Alexander, 1922:10. Type-locality: Argentina, Mendoza, Rio Diamante. Distr. — Argentina (Mendoza). Ref. — Edwards, 1929: 47-48, pl. VII, figs. 35-36 (♂, ♀ genitalia). Type, Alexander's collection.

CHILENSIS Alexander, 1920:3, fig. 1 (wing.). Type-locality: Chile, Aisén, Rio Aisén. Distr. — Chile (Aisén). Ref. — Edwards, 1929:47 (redescription). Type, KSR.

CHILOTA, Edwards, 1929:50. Type-locality: Chile, Chiloé, Isla de Chiloé, 3 mi. s.w. of Castro. Distr. — Chile (Chiloé). Type, BMNH.

DENTULOSA Edwards, 1934:185. Type-localities (syntypes): Chile, Maule, Cauquenes; O'Higgins, Ranagua. Distr. — Chile (Maule, O'Higgins). Type, BZM, BMNH.

DISPAR Edwards, 1929:52, pls. V-VI, figs. 2, 4-5, 14, 18, 25-26, 31-34 (wing, ♂ and ♀ head, genitalia and hind tarsus, pupal and larval structures). Type-locality: Argentina, Rio Negro, Cerro Runge, w. of



San Carlos de Bariloche. Distr. — Argentina (Rio Negro). Type, BMNH.

FUSCIPES Edwards, 1929:48, pl. V, figs. 8-10 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué, near summit of Rosales Pass. Distr. — Argentina (Rio Negro), Chile (Llanquihue). Type, BMNH.

GRACILIS Edwards, 1929:56, text-figs. 3a-b, pls. V-VI, figs. 3, 15-17, 20, 22, 27-28 (♂ and ♀ head, genitalia and hind tarsus, larval and pupal structures). Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina (Rio Negro), Chile (Llanquihue). Type, BMNH.

IMPERATRIX Alexander, 1961:126. Type-locality: Argentina, Neuquen, Lago Lacar, 650 m. Distr. — Argentina (Neuquen). Type, Alexander's collection.

LUTEIPLEURA Alexander, 1953b:99. Type-locality: Chile, Malleco, Curacautin. Distr. — Chile (Malleco). Type, Alexander's collection.

NIGRA Edwards, 1929:60, text-fig. 18c, pls. V-VI, figs. 2, 6-7, 11-13, 21, 23-24, 31-34 (wing, ♂ and ♀ head, genitalia and hind tarsus, larval and pupal structures). Type-locality: Argentina, Rio Negro, foot of Sicra Ventana, between San Carlos de Bariloche and Lake Gutiérrez. Distr. — Argentina (Rio Negro). Ref. — Anthon & Lyneborg, 1968 (larval structures). Type, BMNH.

STIGMATICA Edwards, 1929:51, pls. V-VI, figs. 1, 19 (wing, ♀ genitalia). Type-locality: Chile, Chiloé, Isla de Chiloé, Ancud. Distr. — Chile (Chiloé). Type, BMNH.

SP. "A" Mannheims, 1938:330, fig. 2 (pupa and larva). Locality: Chile, (?Aisén), Rio Pasena (approximately 300 m from mouth), near Puerto Puyuhuapi. Specimens in DEI.

Subfamily PALTOSTOMINAE

Genus *ELPORIA* Edwards

ELPORIA Edwards, 1915:203. Type-species, *Kelloggina barnardi* Edwards (orig. des.).

ELNORAE (Edwards), 1929:67, pls. VII-VIII, figs. 50, 54-55 (♂ genitalia, hind tarsus) (*Curupira*). Type-locality: Argentina, Misiones, side streams of Iguazú Falls. Distr. — Argentina (Misiones). Type, BMNH.

Genus *KELLOGGINA* Williston

Snowia Williston, 1893:119 (preocc. Neumoegan, 1884). Type-species, *rufescens* Williston (orig. des.).

Curupira Osten Sacken, 1895:162 (preocc. Distant, 1888). Type-species, *Paltostoma torrentium* Müller (orig. des.).

Sackeniella Williston, 1896:270 (nom. nov. for *Snowia* Williston, but preocc. Neunier, 1895). Type-species, *Snowia rufescens* Williston (aut.).

KELLOGGINA Williston, 1907:1 (nom. nom. for *Snowia* Williston). Type-species, *Snowia rufescens* Williston (aut.).



- Dimorphotarsa* Lutz, 1920 (Portuguese version): 29. Type-species, *fascibranchia* Lutz (sub. des.; Alexander, 1958:820).
- Dimorphotaenia* Lutz, 1920 (German translation): 26 (lapsus for *Dimorphotarsa* Lutz).
- BOCAINAE (Lutz), 1920:30, pl. 1, figs. 14-20 (egg, pupal and larval structures, ♀ head, thorax and leg) (*Curupira*). Type-locality: Brazil, São Paulo, Serra da Bocaina (exact locality unknown). Distr. — Brazil (São Paulo). Ref. — Lane & d'Andretta, 1956:180, fig. 1 (♂ genitalia) (redescription). Type, IOC.
- BREVICORNIA (Edwards), 1929:66, pls. VII-VIII, figs. 51-53 (♂ genitalia and hind tarsus) (*Curupira*). Type-locality: Argentina, Misiones, side streams of Iguazu Falls. Distr. — Argentina (Misiones). Type, BMNH.
- BREVIVECTIS (Lutz), 1920:35, pl. 7, figs. 98-105 (pupal and larval structures, ♂ head and thorax, ♀ head, thorax and abdomen) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, Salto do Pirai, near Joinville, and Rio dos Bugres, near São Bento do Sul. Distr. — Brazil (Santa Catarina). Type, IOC.
- CATAGUASI Lane & d'Andretta, 1956:181, fig. 2 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Cachoeira Simões, 1150 m. Distr. — Brazil (Rio de Janeiro). Type, FSP.
- CHERENTESI Lane & d'Andretta, 1956:183, fig. 3 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia, Cachoeira Bonita, 2100 m. Distr. — Brazil (Rio de Janeiro). Type, FSP.
- CHILENA (Alexander), 1952:93 (*Curupira*). Type-locality: Chile, Santiago, Santiago. Distr. — Chile (Santiago). Type, Alexander's collection.
- DISTICHA (Lutz), 1920:34, pls. 5-6, figs. 77-78 (pupal and larval structures, ♂, ♀ hind legs and ♀ thorax and abdomen) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, Rios Paulo-Mirim and Hansa (Corupá). Distr. — Brazil (Santa Catarina). Type, IOC.
- EDWARDSI Lane & d'Andretta, 1956:184, fig. 4 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Rio Campo Belo, 750 m. Distr. — Brazil (Rio de Janeiro). Type, FSP.
- FASCIBRANCHIA (Lutz), 1920:29, pl. 1, figs. 1-13 (pupal and larval structures, ♂, ♀ head and hind legs) (*Dimorphotarsa*). Type-localities (syntypes): Brazil, Rio de Janeiro, "... small stream with many cataracts which issues from the Serra da Estrêla by the station Raiz da Serra of the Leopoldina Company, and in the Rio Macacu near the falls by the railway from Nova Friburgo". Distr. — Brazil (RJ). Type, IOC.
- GARCIANA (Lutz), 1920:35, pl. 7, figs. 106-111 (pupal and larval structures, ♂, ♀ thoraces and abdomens, whole ♂) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, Rios Garcia and Jordão (Blumenau), Salto do Pirai, and near São Bento do Sul. Distr. — Brazil (Santa Catarina). Type, IOC.
- GOMEZI Lane & d'Andretta, 1956:186, fig. 5 (♂ genitalia). Type-locality: Brazil, Minas Gerais, Caparaó, Córrego Candeias, 2100 m. Distr. — Brazil (Minas Gerais). Type, FSP; paratypes, BMNH, USNM, IOC, MZUSP.

- GRANULIPUPA (Lutz), 1920:31, pls. 2-3, figs. 31-45 (egg, pupal and larval structures, ♂ thorax, leg, antenna, ♀ leg (*Curupira*). Type-locality: Brazil, São Paulo, Serra da Bocaina (exact locality unknown). Distr. — Brazil (São Paulo). Type, IOC.
- GUAIANASI Lane & d'Andretta, 1956:187, fig. 6 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Alto do Itatiaia, Cachoeira das Flores, 2500 m. Distr. — Brazil (Rio de Janeiro). Type, FSP; paratypes, BMNH, USNM.
- GUARANI Lane & d'Andretta, 1956:188, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, Juquiá, Salto do Açungul, 100 m. Distr. — Brazil (São Paulo). Type, FSP; paratypes, BMNH, USNM, IOC, MZUSP.
- HIRTIPUPA (Lutz), 1920:31, pl. 2, figs. 21-30 (egg, pupal and larval structures, ♂ leg, ♀ head, leg) (*Curupira*). Type-localities (syntypes): Brazil, São Paulo, Serra da Bocaina ("various streams"); Minas Gerais, Pacau, Serra da Mantiqueira. Distr. — Brazil (Minas Gerais, São Paulo). Type, IOC.
- HORRENS (Lutz), 1920:32, pls. 3-4, figs. 46-53 (pupal and larval structures) (*Curupira*). Type-locality: Brazil, São Paulo, Serra da Bocaina (exact locality unknown). Distr. — Brazil (São Paulo). Type, IOC.
- INCERTA (Lutz), 1920:33, pl. 4, fig. 66 (pupal structures) (*Curupira*). Type-locality: Brazil, São Paulo, Serra da Bocaina (exact locality unknown). Distr. — Brazil (São Paulo). Type, IOC.
- LAEMMERTI Lane & d'Andretta, 1956:190, fig. 8 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Rio Campo Belo, 1800 m. Distr. — Brazil (Rio de Janeiro). Type, FSP; paratypes, BMNH, USNM, IOC, MZUSP.
- LORENZI (Lutz), 1920:37 (*Dimorphotarsa*). Type-locality: Brazil, Santa Catarina, Engano. Distr. — Brazil (Rio de Janeiro, Santa Catarina). Ref. — Lane & d'Andretta, 1956:191 (redescription), fig. 8 (♂ genitalia). Type, IOC.
- MOCHLURA (Lutz), 1920:32, pls. 4, 7, figs. 54-65, 112-113 (pupal and larval structures, ♂, ♀ whole) (*Curupira*). Type-localities (syntypes): Brazil, Guanabara, Rio de Janeiro (Corcovado, Serra da Tijuca, Morin districts); São Paulo, Serra da Bocaina. Distr. — Brazil (Guanabara, São Paulo). Refs. — Lane & d'Andretta, 1956:193 (redescription), fig. 11 (♂ genitalia); Pinto, 1933:177, fig. 1 (♂ genitalia). Type, IOC.
- MUELLERI (Lutz), 1920:35, pl. 6, figs. 95-97 (egg, pupal and larval structures) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, Rios Jordão and Garcia. Distr. — Brazil (Santa Catarina). Type, IOC.
- PLURIPUNCTATA (Lutz), 1920:34, pl. 6, figs. 88-90 (pupal and larval structures, ♂ thorax and abdomen) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, Hansa (= Corupá), Rios Paulo and Paulo-Mirim, and Salto do Pirai, near Joinville. Distr. — Brazil (Santa Catarina). Type, IOC.
- RABELLOI Lane & d'Andretta, 1956:195, fig. 11 (♂ genitalia). Type-locality: Brazil, São Paulo, Ilha de São Sebastião, Ihabela, 300 m. Distr. — Brazil (São Paulo). Type, FSP.



- RUFESCENS (Williston), 1893:120 (*Snowia*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara). Type, ?
- SHANNONI Lane & d'Andretta, 1956:195, fig. 12 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Cachoeira Simões, 1500 m. Distr. — Brazil (Rio de Janeiro, São Paulo). Type, FSP; paratypes, BMNH, USNM, IOC, MZUSP.
- SPINIVECTIS (Lutz), 1920:35, pl. 6, figs. 91-94 (pupal and larval structures) (*Curupira*). Type-localities (syntypes): Brazil, Santa Catarina, near Hansa (= Corupá), Rios Paulo and Paulo-Mirim. Distr. — Brazil (Santa Catarina). Type, IOC.
- TAMOIOI Lane & d'Andretta, 1956:198, fig. 13 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Rio Campo Belo, 750 m. Distr. — Brazil (Rio de Janeiro). Type, FSP.
- TAPUIASI Lane & d'Andretta, 1956:198, fig. 14 (♂ genitalia). Type-locality: Brasil, Minas Gerais, Caparaó, Cachoeira da Fumaça, 1800 m. Distr. — Brazil (Minas Gerais). Type, FSP.
- TETRAGONURA (Lutz), 1920:36 (*Dimorphotarsa*). Type-locality: Brazil, Santa Catarina, near São Bento do Sul, Rio dos Bugres. Distr. — Brazil (Santa Catarina). Type, IOC.
- TETRASTICHA (Lutz), 1920:33, pl. 5, figs. 68-76 (pupal and larval structures, ♀ thorax, leg) (*Dimorphotarsa*). Type-locality: Brazil, Santa Catarina, near Blumenau, Rio Garcia. Distr. — Brazil (Santa Catarina). Note — Probably = *torrentium* Müller, in part. Type, IOC.
- TIMBIRA Lane & d'Andretta, 1956:201, fig. 15 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Cachoeira Bonita, 2100 m. Distr. — Brazil (Rio de Janeiro). Type, FSP.
- TORRENTIUM (Müller), 1881a:47-85, 147-151, pls. IV-VII (all stages) (*Pal-tostoma*). Type-localities (syntypes): Brazil, Santa Catarina, Rios Garcia, Jordão and Caeté. Distr. — Brazil (Santa Catarina). Note. — Müller's material contained more than one species (Lutz, 1912: 83); at least one of these is recognizable as one subsequently redescribed and renamed by Lutz (1920). However, the taxonomic actions required to establish the identity of *torrentium* and those species confused with it must await a revisionary study. Refs. — Müller, 1880:37-42 (additional on dimorphic female eye structure); Osten Saeken, 1880:130-132, 1881:206 (criticism of Müller's description of dimorphic females); Müller, 1881b:225-226 (reaction to Osten Saeken's criticism). Types, apparently all lost.
- TRAVASSOSI Lane & d'Andretta, 1956:202, fig. 16 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão, Eugênio Lefevre, 1200 m. Distr. — Brazil (São Paulo). Type, FSP.
- WILLISTONI Lane & d'Andretta, 1956:202, fig. 17 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Itatiaia, Rio Campo Belo, 1800 m, and Maromba, 100 m. Distr. — Brazil (Rio de Janeiro). Type, FSP; paratypes, BMNH, USNM, IOC, MZUSP.

Genus LIMONICOLA Lutz

- LIMONICOLA Lutz, 1928:67. Type-species, *plurivectis* Lutz (mon.).
- LEUCOPTERA Edwards, 1929:73, pl. VII, figs. 39, 41, 49 (♂ head, wing, leg). Type-locality: Peru, Lima, Verrugas Canyon, approximately 5000 feet. Distr. — Peru (Lima). Type, BMNH.

PLURIVECTIS Lutz, 1928:67, pl. 10, figs. 103, pl. 11, fig. 104 (egg, pupal and larval structures, ♂ whole, ♀ claws). Type-locality: Venezuela, Aragua, near Maracay, Río Limón. Distr. — Venezuela (Aragua). Type, ?IOC.

Genus PALTOSTOMA Schiner

- PALTOSTOMA Schiner, 1866:931. Type-species, *superbiens* Schiner (orig. des.). Ref. — Schiner, 1868:27 (redescription).
- ARGYROCINCTA Curran, 1927:1. Type-locality: Puerto Rico, Río Grande. Distr. — Puerto Rico. Ref. — Edwards, 1929:68 (redescription; erroneously cited from "Costa Rica"). Type, AMNH.
- BELLARDII Bezzi, 1913:63. Type-locality: "Mexico" (exact locality unknown). Distr. — Mexico. Type, IMZ.
- DELECTATA Alexander, 1953a:47. Type-locality: Costa Rica, San José, Rivas, Río Buena Vista. Distr. — Costa Rica (San José). Type, Alexander's collection.
- LOBATA Edwards, 1929:70, pls. VII-VIII, figs. 38, 43, 46-47, 58-59 (wing, ♂ genitalia, leg, ♀ mouthparts, leg). Type-locality: Peru, Lima, Verrugas Canyon, approximately 5000 feet. Distr. — Peru (Lima). Type, BMNH.
- PARVICEPS Alexander, 1953a:46. Type-locality: Peru, Junín, Carpapata, Tarma, 2600 m. Distr. — Peru (Junín). Type, Alexander's collection.
- SALTANA Edwards, 1929:71, pl. VII, figs. 37, 42, 48, 62-65 (pupal and larval structures, ♀ wing, mouthparts, leg, genitalia). Type-locality: Argentina, Salta, Quebrada San Lorenzo. Distr. — Argentina (Salta). Type, BMNH.
- SCHINERI Williston, 1896:269, pl. VIII, fig. 27 (♂ genitalia, wing). Type-locality: St. Vincent, West Indies. Distr. — St. Vincent. Type, BMNH.
- SCHINERI Williston of Scott, 1915:181-195, 197-202, pls. IX-XI, figs. 1-15 (pupal and larval structures, ♂ wing, head, leg, genitalia) and Lamb, 1915:195-197, pl. XI, figs. 16-17. Locality: Trinidad, Diego Martin. Specimens in BMNH and CU. Note. — A misidentification, probably an undescribed species. Ref. — Edwards, 1929:68 (redescription); Stone, 1966:294 (key).
- SHANNONI Edwards, 1929:69, pls. VII-VIII, figs. 44-45, 56-57 (♂ wing, leg, genitalia). Type-locality: Peru, Lima, Verrugas Canyon, approximately 5000 feet. Distr. — Peru (Lima). Type, BMNH.
- SUPERBIENS Schiner, 1866:931. Type-locality: "Colombia" (exact locality unknown). Distr. — Colombia. Ref. — Schiner, 1868:28 (redescription). Type, NMW.
- SP. A Stone, 1966:294. Locality: Dominica, Fond Figes River and Springfield Estate. Specimens in USNM. Note. — larvae only.
- SP. B Stone, 1966:294. Locality: Dominica, Fond Figes River. Specimens in USNM. Note. — Larvae only.
- SP. C Stone, 1966:294. Locality: Dominica, Antrim, 1000 feet. Specimens in USNM. Note. — Probably adults of species A or B of Stone.



LITERATURE CITED

ALEXANDER, C. P.

1920. A new genus and species of net-winged midge (Blepharoceridae) and an undescribed species of Tanyderidae (Diptera). *Arkiv för Zoologi* 13:1-7.
1922. An undescribed species of net winged midge from Argentina (Blepharoceridae, Diptera). *Ent. News* 33:10-11.
1952. Undescribed species of nematoceros Diptera. Pt. 1. *Bull. Brooklyn ent. Soc.* 47:88-94.
- 1953a. Undescribed species of nematoceros Diptera. Part II. *Bull. Brooklyn Ent. Soc.* 48(2):41-49.
- 1953b. Undescribed species of nematoceros Diptera. Part IV. *Bull. Brooklyn ent. Soc.* 48(4):97-103.
1958. Geographic distribution of the net-winged midges (Blepharoceridae, Diptera). *Proc. 10th Intern Congr. Ent. (1956)* 1:813-838, figs. 1-23, maps I-III.
1961. Undescribed species of nematoceros Diptera. Part X. *Bull. Brooklyn ent. Soc.* 56:121-127.
1963. Guide to the insects of Connecticut. Part. VI. The Diptera or true flies of Connecticut. Eighth fascicle [part]. Family Blepharoceridae. *Bull. Conn. State Geol. & Nat. Hist. Survey* 93:29-71, pls. 9-13.

ANTHON, H. & L. LYNEBORG

1968. The cuticular morphology of the larval head capsule in Blepharoceridae (Diptera). *Spolia Zool. Mus. Haun.* 27:1-56, figs.

BEZZI, M.

1913. Blepharoceridi italiani con descrizione di una nuova forma e di due specie esotiche. *Boll. Soc. Ent. Ital.* 44:1-114, figs. 1-18.

CURRAN, C. H.

1927. New Neotropical and Oriental Diptera in the American Museum of Natural History. *Amer. Mus. Nov.* 245:1-9.

EDWARDS, F. W.

1915. On *Elporia*, a new genus of blepharicerid flies from South Africa. *Ann. Mag. Nat. Hist. (8)* 16(93):203-215, 22 figs.
1929. Blepharoceridae, in British Museum (Natural History), *Diptera of Patagonia and South Chile* 2:1-75, figs. 1-3, pls. V-VIII.
1934. Four new Diptera in the collection of the Berlin Zoological Museum. *Stylops* 3:185-186.

KELLOGG, V. L.

1913. Diptera. Fam. Blepharoceridae., In P. Wytzman, *Genera Insectorum* 56:1-15, figs. 1-28.

LAMB, C. G.

1915. VI. The female imago, in Scott, 1915. *Ann. Mag. Nat. Hist. (8)* 15:195-197, pl. XI, figs. 16-17.



LANE, J. & C. D'ANDRETTA, JR.

1956. Brazilian Blepharoceridae (Diptera, Nematocera). *Ann. Mag. Nat. Hist.* (12)9:177-204, figs. 1-17.

LUTZ, A.

1912. Contribuição para o estudo da biologia dos dipteros hematofagos. I. Sobre as partes bucaes dos nematoceros que sugam sangue. *Mem. Inst. Oswaldo Cruz* 4:75-83.
1920. Dipteros da familia Blepharoceridae, observados no Brasil. *Mem. Inst. Oswaldo Cruz* 12:21-40 (Portuguese version), 16-40 (German version), pls. 1-7, figs. 1-113.
1928. Blepharoceridae. *Estudios de Zoologia y Parasitologia Venezolanas*, pp. 65-92, 2 pls. and table. Caracas.

MANNHEIMS, B. J.

1935. Beiträge zur Biologie und Morphologie der Blepharoceriden (Dipt.). *Zool. Forschungen* 2:i-iv, 1-115, 69 figs.
1939. Über das Vorkommen der Gattung *Curupira* in Manschukuo nebst Beschreibung der Entwicklungsstadien zweier neuer Blepharoceriden aus Anatolien und süd-Chile. *Arb. Morph. Taxon. Ent.* 5:328-332.

MÜLLER, F.

1879. [Meeting of Nov. 5, 1879: a photographic copy of some drawings of a very curious dipterous insect]. *Trans. Ent. Soc. London (Proc.)* 1879: L.
1880. *Paltostoma torrentium*. Eine Mücke zweigestaltigen Weibchen. *Kosmos* 7:37-42.
- 1881a. A metamorphose de um insecto diptero. *Arch. Mus. Nac.*, Rio de Janeiro 4:47-85, 147-151, pls. IV-VII.
- 1881b. On female dimorphism of *Paltostoma torrentium*. *Ent. Monthly Mag.* 17:225-226.

OSTEN SACKEN, C. R.

1880. Dr. Fritz Müller's discovery of a case of female dimorphism among Diptera (Blepharoceridae). *Ent. Monthly Mag.* 17:130-132.
1881. Dimorphism of female Blepharoceridae. *Ent. Monthly Mag.* 17:206.
1891. Synopsis of the described genera and species of the Blepharoceridae. *Berl. Ent. Zeit.* 36:407-411.
1895. Contributions to the study of the family Liponeuridae Loew (Blepharoceridae Loew, olim.) *Berl. Ent. Zeit.* 40:148-169.

PINTO, C.

1933. Características do hypopygio e asa de *Curupira mochlura* Lutz, 1920 (Diptera, Blepharoceridae). *Rev. Ent.*, Rio de Janeiro 3: 177-179, fig. 1.

SCHINER, J. R.

1866. Bericht über die von der Weltumseglungsreise der k. Fregatte Novara mitgebrachten Dipteren. *Verh. zool.-bot. Ges. Wien* 16: 927-934.



1868. Diptera (Art. 1), 388 pp., 4 pls., in [B. von Wüllersdorf-Urbair, in chargel. *Reise der österreichischen Fregatte Novara um die Erde*. Zool. (1,B). Wien.

SCOTT, H.

1915. The early stages of *Paltostoma schineri* Williston (Diptera, Blepharoceridae). *Ann. Mag. Nat. Hist.* (8)15:181-195, 197-202, pls. IX-XI, figs. 1-15.

STONE, A.

1966. Blephariceridae of Dominica (Diptera). *Proc. Ent. Soc. Wash.* 68:294.

WILLISTON, S. W.


1893. Diptera Brasiliana. III. A new genus of Blepharoceridae. *Kansas Univ. Quart.* 1:119-120.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London 1896*:253-446, pls. ciii-cxiv.

1907. Dipterological notes. *Jour. N. Y. ent. Soc.* 15:1-2.

INDEX

argentinensis Alexander, 1922, Edwardsina	8.2
argyrocineta Curran, 1927, Paltostoma	8.7
bellardii Bezzi, 1913, Paltostoma	8.7
bocainae (Lutz), 1920 (Curupira), Kelloggina	8.4
brevicornia (Edwards), 1929 (Curupira), Kelloggina	8.4
brevivectis (Lutz), 1920 (Curupira), Kelloggina	8.4
cataguasi Lane & d'Andretta, 1956, Kelloggina	8.4
cherentesi Lane & d'Andretta, 1956, Kelloggina	8.4
chilena (Alexander), 1952 (Curupira), Kelloggina	8.4
chilensis Alexander, 1920, Edwardsinina	8.2
chilota Edwards, 1929, Edwardsinina	8.2
Curupira Osten Sacken, 1895	8.3
delectata Alexander, 1953a, Paltostoma	8.7
dentulosa Edwards, 1934, Edwardsinina	8.2
Dimorphotaenia Lutz, 1920	8.4
Dimorphotarsa Lutz, 1920	8.4
dispar Edwards, 1929, Edwardsinina	8.2
disticha (Lutz), 1920 (Curupira), Kelloggina	8.4
edwardsi Lane & d'Andretta, 1956, Kelloggina	8.4
Edwardsinina Alexander, 1920	8.2
elnorae (Edwards), 1929 (Curupira), Elporia	8.3
Elporia Edwards, 1915	8.3
fascibranchia (Lutz), 1920 (Dimorphotarsa), Kelloggina	8.4
fuscipes Edwards, 1929, Edwardsinina	8.3
garciana (Lutz), 1920 (Curupira), Kelloggina	8.4
gomezi Lane & d'Andretta, 1956, Kelloggina	8.4



<i>gracilis</i> Edwards, 1929, Edwardsinina	8.3
<i>granulipupa</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>guaianasi</i> Lane & d'Andretta, 1956, Kelloggina	8.5
<i>guarani</i> Lane & d'Andretta, 1956, Kelloggina	8.5
<i>hirtipupa</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>horrens</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>imperatrix</i> Alexander, 1961 Edwardsinina	8.3
<i>incerta</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>Kelloggina</i> Williston, 1907	8.3
<i>laemmerti</i> Lane & d'Andretta, 1956, Kelloggina	8.5
<i>leucoptera</i> Edwards, 1929, Limonicola	8.6
<i>Limonicola</i> Lutz, 1928	8.6
<i>lobata</i> Edwards, 1929, Paltostoma	8.7
<i>lorenzi</i> (Lutz), 1920 (Dimorphotarsa), Kelloggina	8.5
<i>luteipleura</i> Alexander, 1953b, Edwardsinina	8.3
<i>mochlura</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>muelleri</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>nigra</i> Edwards, 1929, Edwardsinina	8.3
<i>Paltostoma</i> Schiner, 1866	8.7
<i>Paltostominae</i>	8.3
<i>parviceps</i> Alexander, 1953a, Paltostoma	8.7
<i>pleurivectis</i> Lutz, 1928, Limonicola	8.7
<i>pluripunctata</i> (Lutz), 1920 (Curupira), Kelloggina	8.5
<i>rabelloi</i> Lane & d'Andretta, 1956, Kelloggina	8.5
<i>rufescens</i> (Williston), 1893 (Snowia), Kelloggina	8.6
<i>Sackeniella</i> Williston, 1896	8.3
<i>saltana</i> Edwards, 1929, Paltostoma	8.7
<i>schineri</i> Williston, 1896, Paltostoma	8.7
<i>schineri</i> Williston, of Scott, 1915, Paltostoma	8.7
<i>shannoni</i> Edwards, 1929, Paltostoma	8.7
<i>shannoni</i> Lane & d'Andretta, 1956, Kelloggina	8.6
<i>Snowia</i> Williston, 1893	8.3
sp. "A" Mannheims, 1938, Edwardsina	8.3
sp. A Stone, 1966, Paltostoma	8.7
sp. B Stone, 1966, Paltostoma	8.7
sp. C Stone, 1966, Paltostoma	8.7
<i>spinivectis</i> (Lutz), 1920 (Curupira), Kelloggina	8.6
<i>stigmatica</i> Edwards, 1929, Edwardsinina	8.3
<i>superbiens</i> Schiner, 1866, Paltostoma	8.7
<i>tamoioi</i> Lane & d'Andretta, 1956, Kelloggina	8.6
<i>tapuiasi</i> Lane & d'Andretta, 1956, Kelloggina	8.6



tetragonura (Lutz), 1920 (Dimorphotarsa), Kelloggina	8.6
tetrasticha (Lutz), 1920 (Dimorphotarsa), Kelloggina	8.6
timbira Lane & d'Andretta, 1956, Kelloggina	8.6
torrentium (Müller), 1881a (Paltostoma), Kelloggina	8.6
travassosi Lane & d'Andretta, 1956, Kelloggina	8.6
willistoni Lane & d'Andretta, 1956, Kelloggina	8.6



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

10

FAMILY DIXIDAE

ALAN STONE

Entomology Research Division, Agric. Res. Serv.
U.S. Department of Agriculture
Washington, D.C.

The family Dixidae comprises rather small, slender, nearly bare nematoceros flies found in the vicinity of the streams or pools in which their larvae develop. The adults rest on rocks or foliage near the water or swarm over the water. The larvae rest at the water's edge in an inverted U-shape and the pupae are attached to the substratum slightly above the water surface. The family has been treated rather fully by Nowell (1951) (biology, anatomy, taxonomy, catalogue), and Lane (1953) revised the group in the Neotropics as a subfamily of the Culicidae.

Genus DIXELLA Dyar & Shannon

Dixa, subg. DIXELLA Dyar & Shannon, 1924:201. Type-species, *Dixa lirio* Dyar & Shannon (mon). [*Dixella* was considered as subgenus of *Dixa* in the North American catalogue. Eds.]

Dixa, subg. *Paradixa* Tonnoir, 1924:222. Type-species, *Dixa neozelandica* Tonnoir (Edwards, 1932:9).

ANDEANA (Lane), 1942:94, fig. 5, pl. 2, fig. 1 (wing, male genitalia) (*Dixa*). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru.

ARGENTINA (Alexander), 1920:294 (*Dixa*). Type-locality: Argentina, Cordoba, La Granja, Alta Gracia. Distr. — Argentina (Cordoba, San Juan, Salta). Ref. — Lane, 1953:45, figs. 19-20 (wing, male genitalia).

ATRA (Lane), 1942:93, pl. 1, fig. 5 (wing) (*Dixa*). Type-locality: Panama, Gatun Lake, Cano Saddle. Distr. — Panama, Argentina.

CHAFADENSIS (Lane), 1939:99, figs. 1-3 (wing, male genitalia) (*Dixa*). Type-locality: Brazil, Mato Grosso, Ponce, Pontinha. Distr. — Brazil (Mato Grosso, São Paulo).

10.1



- CLAVULUS (Williston), 1896:298, fig. 73 (wing) (*Dixa*). Type-locality: St. Vincent (B.W.I.). Distr. — British West Indies.
- HOFFMANI (Lane), 1942:95, pl. 2, fig. 2 (wing) (*Dixa*). Type-locality: Puerto Rico, Pueblo Viejo. Distr. — Puerto Rico.
- LIMAI (Santos), 1940:21, figs. 1-7 (wing, larva, pupa) (*Dixa*). Type-locality: Brazil, Minas Gerais, Uberaba. Distr. — Brazil (Mato Grosso, Bahia, Minas Gerais, São Paulo). Ref. — Lane, 1953:56, figs. 36-39 (wing, male genitalia, pupa).
- LIRIO (Dyar & Shannon), 1924:200 (*Dixa*). Type-locality: Panama, Canal Zone, Monte Lirio. Distr. — Panama. Ref. — Lane, 1953:40, fig. 15 (male genitalia).
- PAULISTANA (Lane, Forattini & Rabello), 1955:113, figs. 1-5 (wing, male genitalia, pupa, larva) (*Dixa*). Type-locality: Brazil, São Paulo, Capital, Campo Experimental da Água Funda, Departamento de Produção Animal. Distr. — Brazil (São Paulo), Argentina (Misiones).
- PERUVIANA (Edwards), 1931:261 (*Dixa*). Type-locality: Peru, Lima, Verugas. Distr. — Peru.
- SHANNONI (Lane), 1942:88 (*Dixa*). Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica. Ref. — Lane, 1953:51, figs. 26-27 (wing, male genitalia).
- TORRENTIA (Lane), 1939:101, figs. 4-5 (palpus, male genitalia) (*Dixa*). Type-locality: Brazil, São Paulo, Juquiá, Salto do Poço Grande. Distr. — Brazil (Rio de Janeiro, São Paulo). Ref. — Lane, 1953:55, figs. 7, 34, 35 (male, wing, male genitalia).
- TRINITENSIS (Lane), 1943:162, figs. 1-2 (wing, male genitalia) (*Dixa*). Type-locality: Trinidad, near Piaseo (B.W.I.). Distr. — Trinidad (B.W.I.), Brazil (Bahia). Ref. — Lane, 1953:42, figs. 16-18 (wing, male genitalia, pupa, larva).
- VENEZUELENSIS (Lane), 1942:91, fig. 3, pl. 1, fig. 4 (wing, male genitalia) (*Dixa*). Type-locality: Venezuela, Maracay. Distr. — Costa Rica, Venezuela.
- WYGODZINSKYI (Lane), 1945:444, figs. 1-5 (wing, pupa, larva) (*Dixa*). Type-locality: Brazil, São Paulo, Campos do Jordão, Vila Capivari. Distr. — Brazil.

Genus NOTHODIXA Edwards

Dixa, subg. NOTHODIXA Edwards, 1920:101. Type-species, *Dixa campbelli* Alexander (orig. des.).

ATROVITTATA (Edwards), 1930:105, pl. 10, fig. 19 (male genitalia) (*Dixa*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile (Chiloé, Llanquihue, Concepción). Ref. — Lane, 1953:39, fig. 12 (wing).

CHILENSIS (Alexander), 1913:177, fig. (wing) (*Dixa*). Type-locality: Chile, Concepción. Distr. — Argentina (Rio Negro), Chile (Concep-



ción, *Aeoneagua*). Ref. — Lane, 1953:36, figs. 8-9 (wing, male genitalia).

ENSIFERA (Edwards), 1930:103, fig. 5, pl. 10, figs. 13, 17 (male and female genitalia) (*Dixa*). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile (Llanquihue, Chiloe). Ref. — Lane, 1953:39, figs. 13, 14 (wing, male genitalia).

NITIDA (Edwards), 1930:102, pl. 10, figs. 14, 15, 16, 18 (male genitalia, female abdomen) (*Dixa*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina (Rio Negro), Chile (Llanquihue). Ref. — Lane, 1953:83, pl. 1, fig. 1 (wing).

LITERATURE CITED

ALEXANDER, C. P.

1913. A new species of *Dixa* from Chile (Dixidae, Dipt.). *Ent. News* 24:176-177, 1 fig.

1920. A new species of *Dixa* from Argentina (Dixidae, Dipt.). *Ent. News* 31:294.

DYAR, H. G. & R. C. SHANNON

1924. Some new species of American *Dixa* Meigen (Diptera, Culicidae). *Insec. Insc. Menstr.* 12:193-201.

EDWARDS, F. W.

1930. Dixinae, in *Diptera of Patagonia and South Chile Part II*, fasc. 3:100-106, text-fig. 5, pl. 10, figs. 13-19. British Museum (Nat. Hist.).

1931. New Neotropical nematocerous Diptera. *Ann. & Mag. Nat. Hist.* (10)7:255-261.

1932. Diptera. Fam. Culicidae, in P. Wystman: *Genera Insectorum* 194:258 pp., Bruxelles.

LANE, J.

1939. Notes on non hematophagous Culicidae. *Bol. Biol.* (n.s.) 4:99-113, figs. 1-11.

1942. Dixinae e Chaoborinae. Revisão das espécies neotrópicas (Diptera, Culicidae). *Rev. de Ent.* 13:81-148, figs. 1-7, pls. 1-4.

1943. Aditamento e corrigenda ao meu trabalho sobre Dixinae e Chaoborinae (Dipt.). *Rev. de Ent.* 14:162-166, figs. 1-2.

1945. Uma nova espécie de *Dixa* brasileira. *Rev. de Ent.* 16:444-446, figs. 1-5.

1953. Subfamily Dixinae, pp. 31-59, figs. 4, 7, 17-39, in *Neotropical Culicidae*, vol. I. São Paulo.

LANE, J., O. P. FORATTINI & E. X. RABELLO


1955. *Dixa (Dixella) paulistana*, n. sp. *Arq. Fac. Hig. Saúde Públ. Univ. São Paulo* 9(1-2):113-117, figs. 1-5.

NOWELL, W. R.

1951. The dipterous family Dixidae in Western North America (Insecta: Diptera). *Microentomology* 16:187-271, figs. 74-88.

SANTOS, O. B. DOS

1940(1939). Sobre uma nova espécie de *Dixaella* [sic]. *Tese Fac. Odont. e Farm. Minas Gerais*, 22 pp., 8 figs.



TONNOIR, A. L.

1924. New Zealand Dixidae (Dipt.). *Rec. Cant. Mus.* 2(4):221-233.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London*, pp. 263-308, 346-435, 439-446, pls. VII-XIV.

INDEX

andecana (Lane), 1942 (Dixa), Dixella	10.1
argentina (Alexander), 1920 (Dixa), Dixella	10.1
atra (Lane), 1942 (Dixa), Dixella	10.1
atrovittata (Edwards), 1930 (Dixa), Nothodixa	10.2
chapadensis (Lane), 1939 (Dixa), Dixella	10.1
chilensis (Alexander), 1913 (Dixa), Nothodixa	10.2
clavulus (Williston), 1896 (Dixa), Dixella	10.2
Dixella Dyar & Shannon, 1924	10.1
ensifera (Edwards), 1930 (Dixa), Nothodixa	10.3
hoffmani (Lane), 1942 (Dixa), Dixella	10.2
limai (Santos), 1940 (Dixa), Dixella	10.2
lirio (Dyar & Shannon), 1924 (Dixa), Dixella	10.2
nitida (Edwards), 1930 (Dixa), Nothodixa	10.3
Nothodixa Edwards, 1930	10.2
<i>Paradixa Tonnoir</i> , 1924	10.1
paulistana (Lane, Forattini & Rabello), 1955 (Dixa), Dixella ..	10.2
peruviana (Edwards), 1931 (Dixa), Dixella	10.2
shannoni (Lane), 1942 (Dixa), Dixella	10.2
torrentia (Lane), 1939 (Dixa), Dixella	10.2
trinitensis (Lane), 1943 (Dixa), Dixella	10.2
venczuelensis (Lane), 1942 (Dixa), Dixella	10.2
wygodzinskyi (Lane), 1942 (Dixa), Dixella	10.2



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

11

FAMILY CHAOBORIDAE

ALAN STONE

*Entomology Research Division, Agric. Res. Serv.
U.S. Department of Agriculture
Washington, D.C.*

This family of mostly small, fragile, non-biting midges is closely related to the mosquitoes and is often treated as a subfamily of the Culicidae. The larvae live in quiet-water habitats varying from lakes to the water held in bamboo nodes or plant axils. At times the adults may occur in immense numbers, the swarms creating a nuisance. Lane & Aitken (1956) treated the bionomics of a number of species in Trinidad. Most of the Neotropical species are described and keyed by Lane (1953).

Subfamily CORETHRELLINAE

Genus CORETHRELLA Coquillett

CORETHRELLA Coquillett, 1902:191. Type-species, *Corcthra brakeleyi* Coquillett (orig. des.).

ANANAGOLA Dyar, 1926:150. Type-locality: Panama, Canal Zone, Gatun Lake. Distr. — Trinidad (B.W.I.), Panama, Peru (Loreto), Venezuela. Ref. — Lane, 1953:83, figs. 51, 52 (pupa, larva).

inca Lane, 1939a:103. Type-locality: Peru, Loreto, Iquitos.

APPENDICULATA Grabham, 1906:343, figs. 1-2 (larva). Type-locality: Jamaica, Kingston (B.W.I.). Distr. — U.S.A. (North Carolina to Florida), Grenada, Jamaica, Martinique, Hispaniola (Santo Domingo), Puerto Rico, Panama, Brazil (Goiás, Amazonas), Argentina (Tucumán). Ref. — Lane, 1942:113.

arboREALIS Shannon & Del Ponte, 1928:102. Type-locality: Argentina, Tucumán, Vipos.

confusa Lane, 1939a:109. Type-locality: Brazil, Goiás, Inhumas.

BLANDA Dyar, 1928:79. Type-locality: Panama, Las Sabañas. Distr. — Panama, Trinidad (B.W.I.). Ref. — Lane & Aitken, 1957:530 (female).



11.1


- BROMELICOLA Lane, 1939a:108. Type-locality: Brazil, São Paulo, Juquiá, Poço Grande. Distr. — Brazil (Bahia, Espírito Santo, São Paulo). Ref. — Lane, 1942:107, pl. 3, fig. 4; pl. 4, fig. 1 (pupa, larva).
- CARDOSOI Lane, 1942:118. Type-locality: Brazil, São Paulo, Casa Grande. Distr. — Brazil (Bahia, Rio de Janeiro, São Paulo), Trinidad (B.W.I.).
- DOWNSI Lane, 1943:164. Type-locality: Trinidad, Montevideo (B.W.I.). Distr. — Trinidad (B.W.I.), Panama. Refs. — Lane, 1953:77, fig. 49 (pupa); Lane & Aitken, 1957:530, fig. 1 (male genitalia).
- DYARI Lane, 1942:125. Type-locality: Panama, Canal Zone, Darien. Distr. — Panama.
- EDWARDSI Lane, 1942:126. Type-locality: Brazil, Mato Grosso. Distr. — Brazil (Mato Grosso, Minas Gerais).
- FLAVITIBIA Lane, 1939a:392. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Mexico (Chiapas), Trinidad (B.W.I.), Panama, Brazil (São Paulo). Ref. — Lane & Aitken, 1957:531, figs. 2-4 (male genitalia, pupa, larva).
- FULVA Lane, 1939a:107, figs. 9-10 (male genitalia). Type-locality: Brazil, São Paulo, Capital, Perus. Distr. — Brazil (Bahia, São Paulo). Ref. — Lane, 1953:69, figs. 42-45 (wing, male genitalia, pupa, larva).
- INFUSCATA Lane, 1939b:389. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil (São Paulo). Ref. — Lane, 1942:103, pl. 3, fig. 6 (pupa).
- IZQUIERDOI Vargas, 1952:59, fig. 2 (male genitalia, wing). Type-locality: Mexico, Tabasco, Tenosique. Distr. — Mexico (Tabasco).
- JENNINGSI Lane, 1942:106. Type-locality: Panama, Canal Zone. Distr. — Trinidad (B.W.I.), Costa Rica, Panama, Colombia. Ref. — Lane & Aitken, 1957:533, figs. 5-7 (male genitalia, pupa, larva).
- KUMMI Lane, 1942:121. Type-locality: Brazil, Bahia. Distr. — Brazil (Bahia), Argentina (Salta).
- LANEANA Vargas, 1946:64, 2 pls. (wing, head, male genitalia). Type-locality: Mexico, Nuevo León, Monterrey. Distr. — U.S.A. (s. California), Mexico (Nuevo León). Ref. — Belkin & McDonald, 1955:82, pl. 26 (larva, pupa).
- LOPESI Lane, 1942:117. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Japuiba. Distr. — Brazil (Rio de Janeiro, São Paulo, Santa Catarina).
- MACULATA Lane, 1939a:104, fig. 7 (male genitalia). Type-locality: Peru, Loreto, Iquitos. Distr. — Cuba, Peru (Loreto).
- MELANICA Lane & Aitken, 1957:535, figs. 8-10 (male genitalia, pupa, larva). Type-locality: Trinidad, Arima Valley, St. Patrick Estate (B.W.I.). Distr. — Trinidad (B.W.I.), Brazil (Amazonas).
- PILLOSA Lane, 1939a:108. Type-locality: Brazil, São Paulo, Juquiá, Poço Grande. Distr. — Brazil (São Paulo).



- PUELLA Shannon & Del Ponte, 1928:101 (error as *arborealis*). Type-locality: Argentina, Jujuy, Cañitas Viejo, Ledesma. Distr. — Argentina (Jujuy).
- QUADRIVITTATA Shannon & Del Ponte, 1928:101. Type-locality: Argentina, Salta, Tres Pozos, near Embareación. Distr. — Panama, Brazil (Amazonas, Mato Grosso, Bahia, Espírito Santo), Argentina (Salta, Chaco). Refs. — Lane, 1942:122 (female, male); Lane & Cerqueira, 1958:561, fig. 1 (pupa).
- SELVICOLA Lane, 1939b:391, fig. 3 (male terminalia). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Mexico (Chiapas), Trinidad (B.W.I.), Panama, British Guiana, Brazil (Amazonas, Bahia, Espírito Santo, Rio de Janeiro, São Paulo).
- SIMILANS Lane & Aitken, 1957:537. Type-locality: Trinidad, Tucker Valley (B.W.I.). Distr. — Trinidad (B.W.I.).
- STONEI Lane, 1942:119. Type-locality: Panama, Porto Bello Bay, Caldera Is. Distr. — Panama.
- STRIATA Lane, 1942:110. Type-locality: Brazil, São Paulo, Palmeira. Distr. — Brazil (Espírito Santo, São Paulo).
- TARSATA Lane, 1942:102. Type-locality: Brazil, Bahia, Camassari. Distr. — Trinidad (B.W.I.), Brazil (Bahia). Refs. — Lane, 1953:71, fig. 46 (male genitalia); Lane & Aitken, 1957:537, figs. 11, 12 (pupa, larva).
- TRAVASSOSI Lane, 1942:105. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil (Mato Grosso).
- TRIPUNETATA Lane, 1942:120. Type-locality: Puerto Rico, Carolina. Distr. — Trinidad (B.W.I.), Puerto Rico, Brazil (São Paulo). Ref. — Lane & Aitken, 1957:538.
- VITTATA Lane, 1939b:390, fig. 2 (male genitalia). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil (Espírito Santo, São Paulo).
- WHARTONI Vargas, 1952:57, fig. 1 (male genitalia). Type-locality: Mexico, Comalealeo. Distr. — Mexico (Tabasco).
- WHITMANI Lane, 1942:108, pl. 3, fig. 5, pl. 4, fig. 2 (pupa, larva). Type-locality: Brazil, Espírito Santo. Distr. — Brazil (Amazonas, Espírito Santo, São Paulo). Ref. — Lane, 1953:79, fig. 50 (pupa).

Genus LUTZOMIOPS Lane

- Corethrella*, subg. LUTZOMIOPS Lane, 1942:127. Type-species, *Corethrella nigra* Lane (orig. des.) = *davisi* (Shannon & Del Ponte).
- ALTIOLA (Lane), 1939a:110, fig. 11 (male genitalia) (*Corethrella*). Type-locality: Brazil, Goiás, Anápolis. Distr. — Brazil (Goiás, Rio de Janeiro, São Paulo), Argentina (Misiones).
- AMAZONIEUS (Lane), 1939a:110 (*Corethrella*). Type-locality: Brazil, Rondônia, Pôrto Velho. Distr. — Trinidad (B.W.I.), Costa Rica, Panama, Brazil (Rondônia, Espírito Santo, São Paulo).
- BARRETOI (Lane), 1942:132 (*Corethrella*). Type-locality: Brazil, São Paulo, Casa Grande. Distr. — Brazil (São Paulo).
- COUTINHOI (Lane), 1942:131 (*Corethrella*). Type-locality: Brazil, São Paulo, Palmeira. Distr. — Brazil (São Paulo).



- DAVISI (Shannon & Del Ponte), 1928:102 (*Corethrella*). Type-locality: Argentina, Tucumán, 5 milles w. of Concepción. Distr. — Panama, Brazil (Bahia, Mato Grosso, Minas Gerais, São Paulo), Argentina (Tucumán). Ref. — Lane, 1953:94, figs. 53-54 (male genitalia, pupa).
- nigra* Lane, 1939a:105, fig. 8 (male genitalia) (*Corethrella*). Type-locality: Brazil, Mato Grosso, Maracaju.
- IRIDESCENTS (Lane), 1939b:389 (*Corethrella*). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil (Amazonas, São Paulo). Ref. — Lane & Cerqueira, 1958:584, figs. 3-4 (male terminalia, pupa).
- JUQUIANUS (Lane), 1939a:109 (*Corethrella*). Type-locality: Brazil, São Paulo, Juquiá, Poço Grande. Distr. — Trinidad (B.W.I.), Brazil (São Paulo). Ref. — Lane & Aitken, 1957:539, figs. 13-16 (pupa, larva).
- LUTZI (Lane), 1942:128 (*Corethrella*). Type-locality: Brazil, São Paulo, Jaraguá. Distr. — Brazil (São Paulo).
- MANAOSENSIS Lane & Cerqueira, 1958:565. Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil (Amazonas).
- NIGRESCENS (Lane), 1942:129 (*Corethrella*). Type-locality: Brazil, Rio de Janeiro, Magé. Distr. — Trinidad (B.W.I.), Brazil (Rio de Janeiro, São Paulo).
- PALLIDUS (Lane), 1942:128 (*Corethrella*). Type-locality: Panama, Porto Bello. Distr. — Panama.
- PERUVIANUS (Lane), 1939a:110 (*Corethrella*). Type-locality: Peru, Loreto, Iquitos. Distr. — Panama, Peru (Loreto).
- SIANNONI (Lane), 1939a:104, fig. 6a (male genitalia) (*Corethrella*). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru (Loreto), Brazil (Amazonas). Ref. — Lane & Cerqueira, 1958:566, fig. 5 (pupa).

Subfamily CHAOBORINAE

Genus SAYOMYIA Coquillett

- SAYOMYIA Coquillett, 1903:190. Type-species: *Corethra punctipennis* Say (orig. des.). [*Sayomyia* was considered as subgenus in the North American Catalogue. Eds.].
- ANTILLUM (Knab), 1913:121 (*Chaoborus*). Type-locality: Cuba, Habana, Antonio de los Baños. Distr. — Cuba (Habana), Peru (Loreto), Brazil (Amazonas). Ref. — Lane, 1953:103, fig. 56 (male genitalia).
- AUSTRALIS (Shannon & Del Ponte), 1928:99 (*Chaoborus*). Type-locality: Argentina, Tucumán, Finca Santa Barbara. Distr. — Trinidad (B.W.I.), Brazil (Mato Grosso, Minas Gerais, Rio de Janeiro), Argentina (Salta, Jujuy, Tucumán, Chaco).
- BRASILIENSIS (Theobald), 1901:302 (*Corethra*). Type-locality: "Brazil". Distr. — Puerto Rico, Nicaragua, Panama, Trinidad (B.W.I.), Vene-



zuela, Peru (Loreto), Brazil (Mato Grosso, Bahia, Minas Gerais, Rio de Janeiro, São Paulo). Ref. — Lane, 1942:141, pl. 3, fig. 3, pl. 4, figs. 3-5 (wing, male genitalia, larva, pupa).

festivus Dyar & Shannon, 1924:215 (*Chaoborus*). Type-locality: Panama, Canal Zone, Matachin.

antunesi Lane, 1939b:392, fig. 4 (male genitalia) (*Chaoborus*). Type-locality: Brazil, São Paulo, Juquiá.

ELNORAE (Shannon & Del Ponte), 1928:100 (*Chaoborus*). Type-locality: Argentina, Salta, Tres Pozos, near Embarcación. Distr. — Brazil (São Paulo), Argentina (Salta). Ref. — Lane, 1942:145, fig. 7 (male genitalia).

PUNCTIPENNIS (Say), 1823:43 (*Corethra*). Type-locality: U.S.A., Pennsylvania. Distr. — Saskatchewan to Quebec, s. to Colorado, Texas, and Tennessee, Mexico (Jalisco). Ref. — Cook, 1956:32, figs. 13D, 14E, 15B, 16A-B, 17C-F-M, 18H-I (wing, male genitalia, female genitalia, larva, pupa).

SOUZAI (Lane), 1939a:112, fig. 12 (male genitalia) (*Chaoborus*, as *sousai*). Type-locality: Brazil, Pará, Santarém. Distr. — Panamá, Trinidad (B.W.I.), Brazil (Pará, Minas Gerais), Argentina (Tucumán). Ref. — Lane & Aitken, 1957:542, figs. 17-19 (pupa, larva).

Genus EDWARDSOPS Lane

Chaoborus, subg. EDWARDSOPS Lane, 1942:134. Type-species, *Chaoborus unicolor* Lane (orig. des.).

The species of this genus have been keyed by Lane & Heredia (1956).

BOLIVIENSIS Lane & Heredia, 1956:206, fig. 1 (wing). Type-locality: Bolivia, El Paylon. Distr. — Bolivia.

BREVISECTOR (Edwards), 1930:533, fig. 2b (wing) (*Chaoborus*). Type-locality: Brazil, Amazonas, Manaus. Distr. — Venezuela, Brazil (Amazonas), Bolivia.

MAGNIFICUS (Lane), 1942:138, pl. 3, fig. 2 (wing) (*Chaoborus*). Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Peru (Loreto), Bolivia, Brazil (Mato Grosso). Ref. — Lane, 1953:107, figs. 57, 58 (wing, male genitalia).

STONEI (Lane), 1942:137, fig. 6, pl. 3, fig. 1 (male genitalia, wing) (*Chaoborus*). Type-locality: Venezuela, Zulia, Maracaibo. Distr. — Venezuela (Zulia), Brazil (Pará).

UNICOLOR (Lane), 1942:135, fig. 4, pl. 2, fig. 5 (male genitalia, wing) (*Chaoborus*). Type-locality: Brazil, Mato Grosso, Salobra, margins of River Miranda. Distr. — Brazil (Mato Grosso).

LITERATURE CITED

BELKIN, J. N. & W. A. McDONALD

1955. A population of *Corethrella laneana* from Death Valley, with descriptions of all stages and discussion of the Corethrellini (Diptera, Culicidae). *Bull. So. Calif. Acad. Sci.* 54:82-96, pl. 26.



COOK, E. F.

1956. The Nearetie Chaoborinae (Diptera: Culicidae). *Univ. Minnesota Agr. Expt. Sta. Techn. Bull.* 218:102 pp., 34 figs.

COQUILLET, D. W.

1902. New forms of Culicidae from North America. *Jour. N. Y. Ent. Soc.* 10:191-194.
1903. A new culicid genus related to *Corethra*. *Canad. Ent.* 35:189-190.

DYAR, H. G.

1926. Notes on *Corethrella appendiculata* Grabham (Diptera, Culicidae). *Insec. Insc. Menstr.* 14:150.
1928. A new *Corethrella* from Panama (Diptera, Culicidae). *Ent. News* 39:79-80.

DYAR, H. G. & R. C. SHANNON

1924. The American Chaoborinae. *Insec. Insc. Menstr.* 12:201-216.

EDWARDS, F. W.

1930. Notes on exotic Chaoborinae, with descriptions of new species (Diptera, Culicidae). *Ann. & Mag. Nat. Hist.* (10)6:528-540, 2 figs.

GRABHAM, M.

1906. A new *Corethrella* from Jamaica. *Ent. News* 17:343-345, figs. 1-2.

KNAB, F.

1913. A new Cuban *Chaoborus*. *Insec. Insc. Menstr.* 1:121-122.

LANE, J.

- 1939a. Notes on non hematophagous Culicidae. *Bol. Biol.* (n.s.) 4:99-113, figs. 1-11.
1939b. Non hematophagous Culicidae (second paper). *Bol. Biol.* (n.s.) 4:386-393, figs. 1-4.
1942. Dixinae e Chaoborinae. Revisão das espécies neotrópicas (Diptera, Culicidae). *Rev. Entom.* 13:81-148, figs. 1-7, pls. 1-4.
1943. Aditamento e corrigenda ao meu trabalho sobre Dixinae e Chaoborinae (Dipt.). *Rev. Entom.* 14:162-166, figs. 1-2.
1953. Chaoborinae, pp. 63-109, in *Neotropical Culicidae*, vol. 1. São Paulo.

LANE, J. & T. H. G. AITKEN

- 1957 (1958). Chaoborinae from Trinidad, B.W.I. (Diptera, Culicidae). *Ann. Ent. Soc. Amer.* 49:530-543, figs. 1-19.

LANE, J. & N. L. CERQUEIRA

1958. Amazonian Chaoborinae (Diptera, Culicidae). *Studia Entom.* 1:561-566, figs. 1-5.



LANE, J. & R. L. HEREDIA

1956. A new Neotropical *Edwardsops*. *Rev. Brasil. Entom.* 4:205-208, fig. 1.

SHANNON, R. C. & E. DEL PONTE

1928. Los Culieidos de la Argentina. *Rev. Inst. Bact.* 5(1):28-140, figs. 1-14.

THEOBALD, F. V.

1901. A monograph of the *Culicidae* or mosquitoes. British Museum (Nat. Hist.), vol. 2: pp. 1-391, figs. 152-318.

VARGAS, L.

1946. *Corethrella* (*Corethrella*) *laneana* n. sp. (Diptera, Culicidae), procedente de Monterrey, N.L. *Rev. Inst. Salubr. y Enferm. Trop.* 12:63-67, 2 pls.

1952. Dos nuevas especies Mexicanas de *Corethrella* (Diptera, Culicidae). *Rev. Soc. Mex. Hist. Nat.* 13:57-62, figs. 1-2.

INDEX

<i>alticola</i> (Lane), 1939 (<i>Corethrella</i>), <i>Lutzomiops</i>	11.3
<i>amazonicus</i> (Lane), 1939 (<i>Corethrella</i>), <i>Lutzomiops</i>	11.3
<i>ananaeola</i> Dyar, 1926, <i>Corethrella</i>	11.1
<i>antillum</i> (Knab), 1913 (<i>Chaoborus</i>), <i>Sayomyia</i>	11.4
<i>antunesi</i> Lane, 1939 (<i>Chaoborus</i>), <i>Sayomyia</i>	11.5
<i>appendiculata</i> Grabham, 1906, <i>Corethrella</i>	11.1
<i>arboREALIS</i> Shannon & Del Ponte, 1928, <i>Corethrella</i>	11.1
<i>australis</i> (Shannon & Del Ponte), 1928 (<i>Chaoborus</i>), <i>Sayomyia</i> ..	11.4
<i>barrettoi</i> (Lane), 1942 (<i>Corethrella</i>), <i>Lutzomiops</i>	11.3
<i>blanda</i> Dyar, 1928, <i>Corethrella</i>	11.1
<i>boliviensis</i> Lane & Heredia, 1956, <i>Edwardsops</i>	11.5
<i>brasilienSIS</i> (Theobald), 1901 (<i>Corethra</i>), <i>Sayomyia</i>	11.4
<i>breviseetor</i> (Edwards), 1930 (<i>Chaoborus</i>), <i>Edwardsops</i>	11.5
<i>bromelicola</i> Lane, 1939, <i>Corethrella</i>	11.2
<i>cardosoi</i> Lane, 1942, <i>Corethrella</i>	11.2
<i>Chaoborinae</i>	11.4
<i>confusa</i> Lane, 1939, <i>Corethrella</i>	11.1
<i>Corethrella</i> Coquillett, 1902	11.1
<i>Corethrellinae</i>	11.1
<i>coutinhoi</i> (Lane), 1942 (<i>Corethrella</i>), <i>Lutzomiops</i>	11.3
<i>davisi</i> (Shannon & Del Ponte), 1928 (<i>Corethrella</i>), <i>Lutzomiops</i> ..	11.4
<i>downsi</i> Lane, 1943, <i>Corethrella</i>	11.2
<i>dyari</i> Lane, 1942, <i>Corethrella</i>	11.2
<i>edwardsi</i> Lane, 1942, <i>Corethrella</i>	11.2
<i>Edwardsops</i> Lane, 1942	11.5
<i>elnorae</i> (Shannon & Del Ponte), 1928 (<i>Chaoborus</i>), <i>Sayomyia</i> ..	11.5



<i>festivus</i> Dyar & Shannon, 1924 (Chaoborus), Sayomyia	11.5
<i>flavitibia</i> Lane, 1939, Corethrella	11.2
<i>fulva</i> Lane, 1939, Corethrella	11.2
<i>inea</i> Lane, 1939, Corethrella	11.1
<i>infusca</i> Lane, 1939, Corethrella	11.2
<i>iridescens</i> (Lane), 1939 (Corethrella), Lutzomiops	11.4
<i>izquierdoi</i> Vargas, 1952 Corethrella	11.2
<i>jenningsi</i> Lane, 1942, Corethrella	11.2
<i>juquianus</i> (Lane), 1939 (Corethrella), Lutzomiops	11.4
<i>kummi</i> Lane, 1942, Corethrella	11.2
<i>laneana</i> Vargas, 1946, Corethrella	11.2
<i>lopesi</i> Lane, 1942, Corethrella	11.2
<i>lutzi</i> (Lane), 1942 (Corethrella), Lutzomiops	11.4
Lutzomiops Lane, 1942	11.3
<i>maculata</i> Lane, 1939, Corethrella	11.2
<i>magnificus</i> (Lane), 1942 (Chaoborus), Edwardsops	11.5
<i>manaosensis</i> Lane & Cerqueira, 1958, Lutzomiops	11.4
<i>melanica</i> Lane & Aitken, 1957, Corethrella	11.2
<i>nigra</i> Lane, 1939 (Corethrella), Lutzomiops	11.4
<i>nigrescens</i> (Lane), 1942 (Corethrella), Lutzomiops	11.4
<i>pallidus</i> (Lane), 1942 (Corethrella), Lutzomiops	11.4
<i>peruvianus</i> (Lane), 1939 (Corethrella), Lutzomiops	11.4
<i>pillosa</i> Lane, 1939, Corethrella	11.2
<i>puella</i> Shannon & Del Ponte, 1928, Corethrella	11.3
<i>punctipennis</i> (Say), 1823 (Corethra), Sayomyia	11.5
<i>quadrivittata</i> Shannon & Del Ponte, 1928, Corethrella	11.3
Sayomyia Coquillett, 1903	11.4
<i>sclicola</i> Lane, 1939, Corethrella	11.3
<i>shannoni</i> (Lane), 1939 (Corethrella), Lutzomiops	11.4
<i>similans</i> Lane & Aitken, 1957, Corethrella	11.3
<i>souzai</i> (as <i>sousai</i>) (Lane), 1939 (Chaoborus), Sayomyia	11.5
<i>stonei</i> Lane, 1942, Corethrella	11.3
<i>stonei</i> Lane, 1942 (Chaoborus), Edwardsops	11.5
<i>striata</i> Lane, 1942, Corethrella	11.3
<i>tarsata</i> Lane, 1942, Corethrella	11.3
<i>travassosi</i> Lane, 1942, Corethrella	11.3
<i>tripunctata</i> Lane, 1942, Corethrella	11.3
<i>unicolor</i> (Lane), 1942 (Chaoborus), Edwardsops	11.5
<i>vittata</i> Lane, 1939, Corethrella	11.3
<i>whartoni</i> Vargas, 1952, Corethrella	11.3
<i>whitmani</i> Lane, 1942, Corethrella	11.3



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

13

FAMILY THAUMALEIDAE
(ORPHNEPHILIDAE)


ALAN STONE

Entomology Research Division, Agric. Res. Serv.
U.S. Department of Agriculture
Washington, D.C.

This is a small family of rarely encountered flies represented in the region by a single genus known from Southern Argentina, Chile, New Zealand, and Australia. Nothing has been published on the biology of species of *Austrothaumalea*, but it is probably much like that of the other genera of the family. The habits of a North American species, *Thaumalea americana* Bezzi, have been described by Leathers (1922). The larvae are found in cold streams on rocks over which a thin layer of water flows. They feed on organic matter on the rocks. The pupae are found on wet moss, dead leaves, or in mud; the adults are found on vegetation near the larval habitat.

Genus AUSTROTHAUMALEA Tonnoir

- AUSTROTHAUMALEA Tonnoir, 1927:109. Type-species, *neozelandica* Tonnoir (orig. des.).
- APICALIS Edwards, 1930:113, fig. 6b, pl. 11, fig. 26. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile (Llanquihue).
- CHILENSIS Edwards, 1930:111, fig. 6a, pl. 11, fig. 24. Type-locality: Chile, Chiloe Is., Ancud. Distr. — Chile.
- HALTERALIS Edwards, 1930:114, pl. 11, fig. 22. Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.
- NUDIPENNIS Edwards, 1930:113, fig. 6e, pl. 11, fig. 23. Type-locality: Chile, Chiloe Is., Ancud. Distr. — Chile.
- SETIPENNIS Edwards, 1930:113, pl. 11, fig. 21. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.



13.1



LITERATURE CITED

EDWARDS, F. W.

1930. Thaumalcidae (Orphnephilidae). *Diptera of Patagonia and South Chile*, Part II, fasc. 3:111-114, text-fig. 6, pl. 11, figs. 20-26. British Museum (Nat. Hist.).

LEATHIERS, A. L.

1922. Ecological study of aquatic midges and some related insects with special reference to feeding habits. *Bull. U.S. Bur. Fisheries* 38:1-61, 57 figs.

TONNOIR, A. L.

1927. Description of new and remarkable New Zealand Diptera. *Rec. Canterbury Mus.* 3(2):101-112, 4 figs.



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
OF THE UNITED STATES

14

FAMILY CERATOPOGONIDAE

WILLIS W. WIRTH

*Systematic Entomology Laboratory, Agricultural Research Service, USDA
c/o U. S. National Museum, Washington, D. C. 20560*


The "biting midges" may be found in large numbers in nearly any aquatic or semiaquatic habitat in all regions of the world. Because of their small size (usually 1-4 mm) they have been little collected and are poorly known. Adult habits are diverse but most species are adapted to some type of blood-sucking. Species of *Culicoides*, *Lasiohelea*, and *Leptoconops* suck vertebrate blood and some are notorious pests, especially in beach or mountain resort areas. Some *Culicoides* are known vectors of diseases including onchocerciasis of horses and cattle, blue-tongue of sheep, horse-sickness, several human filariases, and *Haemoproteus* and *Leucocytozoon* diseases of birds. Some *Atrichopogon* and *Forcipomyia* are ectoparasitic on larger insects, while many genera of Ceratopogoninae are predaceous on other smaller insects. Some genera (e. g. *Dasyhelea*) are exclusively flower-visiting, feeding on nectar, while many others thus supplement their diet or have non-haematophagous species. The immature stages may be terrestrial under bark or on wet or damp wood (*Forcipomyia*); semiaquatic on wet algae-covered soil, wood, or rocks (*Atrichopogon*, *Dasyhelea*, etc.); in wet, decomposing, plant material such as cactus stems, banana stalks or leaf compost (*Culicoides*, etc.); or aquatic in mud or wet sand on lake, pond, or stream margins, salt marshes, tree holes, or water-holding plants (most genera).

The foremost pioneer in the study of Neotropical Ceratopogonidae was Adolpho Lutz (1912-1914) working at the Instituto Oswaldo Cruz in Rio de Janeiro, Brazil. J. W. S. Macfie (1932-1953), for many years the foremost international specialist in this family, described many Neotropical species. From 1945 to 1961 John Lane of the Universidade de São Paulo revised many genera of Brazilian biting midges. Lima (1928-1937), Barretto (1944), Barbosa (1943-1953), and Forattini (1953-1960) in Brazil; Fox (1942-1955) in Puerto Rico; Floch and Abonnene (1942-1950) in French Guiana; Ortiz (1950-1969) in Venezuela; Vargas (1944-1960) in Mexico; and Wirth and Blanton (1963-1972) in the U.S.A. published series of revisions of the bloodsucking species of *Culicoides*.

Important historic collections of Neotropical Ceratopogonidae are housed in the British Museum (Natural History) in London (Macfie), Instituto Oswaldo Cruz in Rio de Janeiro (Lutz and Lima), Universidade de São Paulo (Lane and Forattini), Instituto Nacional de Higiene in Caracas (Ortiz), and the U. S. National Museum in Washington, D. C. (Wirth and Blanton).

REFERENCES: Carter, Ingram & Macfie, 1920-1922 (anatomy, techniques, biology, immature stages, revision of West African species); Edwards, 1926

14.1



(classification, generic revision, British species); Goetghebuer & Lenz, 1933-1934 (revision Palaearctic species); Mayer, 1934 (classification of immature stages); Thomsen, 1937 (revision of immature stages of North American species); Macfie, 1940d (revision and key to world genera); Johannsen, 1943b (synopsis of genera of Americas); Wirth, 1952e (general classification, anatomy, economics, techniques, biology of immature stages, revision of California species); Fox, 1955 (catalog of American bloodsucking species); Wirth, 1956a (ceratopogonid pollinators of *Hevea* rubber trees); Tokunaga & Murachi, 1959 (generic classification, revision of Micronesian species); Fallis & Bennett, 1961 (ceratopogonid vectors of blood parasites); Kettle, 1962, 1969 (bionomics and control); Downes, 1971 (evolution of adult feeding habits).

Subfamily LEPTOCONOPINAE

Genus LEPTOCONOPS Skuse

- LEPTOCONOPS Skuse, 1889: 288. Type-species, *stygius* Skuse (mon.).
Tersesthes Townsend, 1893: 370. Type-species, *torrens* Townsend (orig. des.).
Mycterotypus Noé, 1905: 114. Type-species, *bezzii* Noé (Carter, 1921: 3).
Mycteromyia Lutz, 1912:24. Error for *Mycterotypus* Noé.
Protersesthes Kieffer, 1921b: 107. Type-species, *Tersesthes brasiliensis* Lutz (orig. des.).
Refs.: Carter, 1921 (world revision); Cavalieri & Chiossone, 1966a: 56 (key to American species).

Subgenus HOLOCONOPS Kieffer

- HOLOCONOPS Kieffer, 1918: 135 (as genus). Type-species, *Leptoconops kerteszi* Kieffer (orig. des.).
BEQUAERTI (Kieffer), 1925b: 405 (*Holoconops*, as *becquaerti*). Type-locality: Honduras, Puerto Castilla. Distr. — U.S.A. (Florida, Texas), circum-Caribbean. Refs. — Bequaert, 1924: 193 (biology); Painter, 1926: 257, figs. 1, 4, 7 (larva, pupa) (biology); Wirth, 1951:281 (distribution); Linley, 1965b: 37 (ovarian cycle); Davies & Linley, 1965: 117 (breeding sites, Jamaica and Honduras).
hondurensis Hoffman, 1926: 135, fig. 1 (antenna, male genitalia, claw, wing, spermathecae, palpus). Type-locality: Honduras, Puerto Castilla.

Subgenus LEPTOCONOPS Skuse

- BRASILIENSIS (Lutz), 1913: 66, pl. 8, fig. 18 (habitus) (*Tersesthes*). Type-locality: Brazil, Amazonas, Tocantins. Distr. — Brazil. Ref. — Lane, 1945: 359, fig. 1 (wing).
CASALI Cavalieri & Chiossone, 1966a: 45, figs. 1-15 (antenna, tarsus, head, tibial comb, palpus, wing, abdomen, spermathecae, claw). Type-locality: Argentina, Prov. San Luis, route Villa Saladillo to La Petra crossing Río Quinto. Distr. — Argentina, Uruguay. Ref. — Chiossone, 1969: 17, figs. 1-11 (mouthparts) (distribution).
CHILENSIS Forattini, 1958: 38, figs. 1-5 (wing, eye separation, antenna, palpus, spermathecae). Type-locality: Chile, Arica, Valle Chaca. Distr. — Chile.



PETROCCHIAE Shannon & Del Ponte, 1927:734, figs. 1-2 (female, palpus). Type-locality: Argentina, Tucumán, Río Tapia. Distr. — Argentina (Tucumán, Catamarca, San Luís). Ref. — Cavaliere & Chiossone, 1966a: 51, figs. 16-31 (many parts) (redescribed, distribution).

VENEZUELENSIS Ortiz, 1952a: 165, figs. 1-9 (wing, eye separation, antenna, palpus, spermathecae). Type-locality: Venezuela, Terr. Amazonas, San Carlos de Rio Negro. Distr. — Venezuela.

Subgenus MEGACONOPS Wirth & Atchley

Leptoconops, subg. MEGACONOPS Wirth & Atchley, 1973:18. Type-species, *floridensis* Wirth (orig. des.).

FLORIDENSIS Wirth, 1951:282, figs. 1-7 (wing, legs, palpus, antenna, claws). Type-locality: U.S.A., Florida, Santa Rosa Island. Distr. — U.S.A. (Florida), Jamaica, Colombia.

Subfamily FORCIPOMYIINAE

Genus FORCIPOMYIA Meigen

FORCIPOMYIA Meigen, 1818:73, 75. Type-species, *Tipula bipunctata* Linnaeus. Generic name cited in specific synonymy; needs action by I.C.Z.N. to validate, with *bipunctata* as type-species.

Tetraphora Philippi, 1865:630. Type-species, *fusca* Philippi (mon.).

Euprojoannisia Brèthes, 1914:155. Type-species, *plantensis* Brèthes (orig. des.). Refs. — Ingram & Macfie, 1931:157 (key to Chilean species); Macfie, 1939a:142 (key to Neotropical species); Wirth, 1956b:356 (review of species parasitic on insects); Saunders, 1956:657 (revision of subgenera based on immature stages); Saunders & Bowman, 1956:1 (role in cacao pollination in Costa Rica); Saunders, 1959:33 (collecting methods, role as cacao pollinators).

Subgenus BLANTONIA Wirth & Dow

Forcipomyia, subg. BLANTONIA Wirth & Dow, 1971:289. Type-species, *caribbea* Wirth & Dow (orig. des.).

CARIBBEA Wirth & Dow, 1971:291, figs. 1-13 (male, female, many parts). Type-locality: Jamaica, Negril, Westmorland Parish. Distr. — U.S.A. (Florida), British Honduras, Guyana, Trinidad, West Indies (Jamaica).

Subgenus CALOFORCIPOMYIA Saunders

Forcipomyia, subg. CALOFORCIPOMYIA Saunders, 1956:680. Type-species, *caerulea* Saunders (orig. des.).

CAERULEA Saunders, 1956:681, fig. 9 (all stages). Type-locality: Brazil, Rio de Janeiro, reared from larvae on rotten wood in forest. Distr. — Brazil.

EUKOSMA Macfie, 1939a:148, fig. 5 (palpus, wing, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

FURCIFERA Macfie, 1940e:920, fig. 1 (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



NIGRESCENS Macfie, 1939a:156, figs. 9-10 (antenna, palpus, wing, spermatheca, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

VARICOLOR Saunders, 1956:683, fig. 10 (all stages). Type-locality: Brazil, Rio de Janeiro, reared from larva on moist rotting wood in forest. Distr. — Brazil.

Subgenus EUFORCIPOMYIA Malloch

EUFORCIPOMYIA Malloch, 1915b:312 (as genus). Type-species, *hirtipennis* Malloch (orig. des.).

Forcipomyia, subg. *Proforcipomyia* Saunders, 1956:662. Type-species, *wirthi* Saunders (orig. des.).

Ref. — Saunders, 1956:772 (revision, immature stages).

BROMELIAE Saunders, 1956:665, fig. 3 (all stages). Type-locality: Brazil, Rio de Janeiro, larvae in terrestrial bromeliads. Distr. — Brazil.

CALCARATA (Coquillett), 1905:64 (*Ceratopogon*). Type-locality: México, Tabasco, Frontera. Distr. — U.S.A. (Mississippi to Florida and Virginia), Mexico.

FALCIFERA Saunders, 1959:39, fig. 3 (all stages). Type-locality: Trinidad, Las Hermanas Plantation, near San Rafael. Distr. — Trinidad.

INDECORA Kieffer, 1914:269. Type-locality: South Africa, Stellenbosch. Distr. — Pantropical, Trinidad.

ingrami Carter, 1919:290. Type-locality: Gold Coast, Accra. Ref. — Macfie, 1938:160 (Trinidad record).

LONGISPINA Saunders, 1956:669, fig. 5 (all stages). Type-locality: Brazil, Rio de Janeiro, larvae under dead leaves, on wet rock face. Distr. — Brazil.

LOTUS (Williston), 1896:282, plate 9, fig. 47 (wing) (*Ceratopogon*). Type-locality: St. Vincent. (notes: of 4 syntypes in the British Museum (Nat. Hist.), 2 are in subgenus *Lasiohelea*, and 2 are in *Euforcipomyia*; the wing figured by Williston is of a *Euforcipomyia*).

MORTUIFOLII Saunders, 1959:35, fig. 1 (all stages). Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.

QUASHINGRAMI Macfie, 1939a:164, fig. 12 (palpus, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Trinidad, Costa Rica. Ref. — Macfie, 1944:297 (Trinidad, from cacao flowers).

SETIGERA Saunders, 1959:38, fig. 2 (all stages). Type-locality: Trinidad, Maracas Valley. Distr. — Trinidad.

SPATULIFERA Saunders, 1956:667, fig. 4 (all stages). Type-locality: Brazil, São Paulo, Santos, larvae on underside of banana leaves lying in grass. Distr. — Brazil.

Subgenus FORCIPOMYIA Meigen

ABERCROMBYI Macfie, 1938:161, fig. 4 (male genitalia). Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.

ANNULATIPES Macfie, 1939a:154, fig. 8 (palpus, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.




- ARGENTEOLA Macfie, 1939a:146, fig. 4 (antenna, palpus, wing, leg, spermatheca).
Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
Ref. — Lane, 1947c:163, fig. 2 (male genitalia) (male, immature stages, distribution).
- BICOLOR Lutz, 1914:89. Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil.
- CALIGINOSELLA Wirth, nom. nov.
caliginosa Ingram & Macfie, 1931:167, fig. 19a (male genitalia). Type-locality: Argentina, Bariloche. Distr. Argentina. Preoccupied in *Forcipomyia* by *Lasiohalea caliginosa* Ingram & Macfie, 1924.
- CHILENSIS (Philippi), 1865:601 (*Ceratopogon*). Type-locality: Chile, Santiago. Distr. — Chile. Ref. — Ingram & Macfie, 1931:163, figs. 15-17 (wing, antenna, tibial spine, male genitalia).
- DELPONTEI Cavalieri, 1962a:169, figs. 1-4 (wing, antenna, palpus, spermatheca). Type-locality: Argentina, Tierra del Fuego, El Tristen. Distr. — Argentina.
- DESERTENSIS Wirth & Hubert, 1960:639, figs. 1, 22 (wing, palpus, spermatheca, male genitalia). Type-locality: U.S.A., California, Los Angeles Co., San Dimas Canyon, reared from *Opuntia* cactus. Distr. — U.S.A. (California), Mexico (Baja California).
- DISCOLORIPES Macfie, 1939a:159, fig. 11 (palpus, antenna). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DUBIA Macfie, 1939a:162. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- EDWARDSIANA Wirth, nom. nov.
edwardsi Ingram & Macfie, 1931:160, figs. 15, 17 (wing, antenna). Type-locality: Argentina, Rio Negro, Lago Gutiérrez. Distr. — Argentina. Preoccupied in *Forcipomyia* by *Apelma edwardsi* Saunders, 1925.
- FLAVA (Williston), 1896:280, pl. 9, fig. 42 (wing, tarsus) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent).
- FLAVIFEMORIS Macfie, 1940a:24. Type-locality: Guyana, New River. Distr. — Guyana.
- FUSCA (Philippi), 1865:630 (*Tetrachora*). Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Wirth, 1955c:235 (type redescribed).
patagonica Ingram & Macfie, 1931:165, fig. 18 (male genitalia). Type-locality: Argentina, Bariloche.
- GENUALIS (Loew), 1866:128 (Cent. 6, no. 1) (*Ceratopogon*). Type-locality: "Cuba". Distr. — U.S.A. (Louisiana, Georgia, Florida), Bermuda, West Indies, Central and South America to Brazil and Galápagos Islands.
propinqua Williston, 1896:279, pl. 9, fig. 41 (wing, tarsus) (*Ceratopogon*). Type-locality: St. Vincent.
raleighi Macfie, 1938:160, fig. 3 (male genitalia). Type-locality: Trinidad, St. Augustine. Refs. — Saunders, 1956:660, fig. 1 (all stages; larvae under rotting wood); Wirth, 1956a:248 (pollinating *Hevea* rubber trees).
squamosa Lutz, misident.; Floch & Abonnenc, 1942c:1, fig. 1 a-f (wing, palpus, tarsus, male genitalia) (French Guiana).
- LACRIMATORII Macfie, 1939a:161, fig. 11 (palpus, antenna). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- MACULOSA Ingram & Macfie, 1931:159, figs. 14, 16, 17 (wing, palpus, antenna). Type-locality: Argentina, Rio Negro, Lago Gutiérrez. Distr. — Argentina.



- MULTIPICTA Ingram & Macfie, 1931:157, figs. 14, 17 (wing, antenna). Type-locality: Argentina, Rio Negro, Lago Nahuel Huapi. Distr. — Argentina, Chile.
- NUNCUPATA Macfie, 1949:110. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- ORNATIPENNIS Macfie, 1939a:151, fig. 6 (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Costa Rica.
- PICTONI Macfie, 1938:161, fig. 5 (male genitalia). Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.
- PIROSKYI Cavalieri, 1962a:172, figs. 5-8 (wing, antenna, palpus, spermathecae). Type-locality: Argentina, Tierra del Fuego, Rio Tristen, Ushuaia. Distr. — Argentina.
- PLATENSIS (Brèthes), 1914:156 (*Euprojoannisia*). Type-locality: Argentina, General Urquiza. Distr. — Argentina.
- POULAINAE Ingram & Macfie, 1931:227. Type-locality: Argentina, Buenos Aires. Distr. — Argentina, Brazil. Ref. — Macfie, 1939a:144 (notes, Brazil).
- Punctumalbum* (Kieffer), of Kieffer, 1917:297, fig. 1 (antenna) (recorded from Paraguay). Not Neotropical.
- SETOSICRUS (Kieffer), 1906:357 (*Ceratopogon*). Type-locality: Chile, Concepción. Distr. — Chile.
- SEXVITTATA Wirth, 1956a:248, fig. 2 (palpus, tibial spine, male genitalia). Type-locality: Costa Rica, Los Diamantes.
- SPATULIGERA Macfie, 1949:111. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- SQUAMITHORAX Clastrier, 1972:170, fig. 1 (female), fig. 2 (male). Type-locality: French Guiana, Cayenne, Cabassou Forest. Distr. — French Guiana.
- SQUAMITIBIA Lutz, 1914:88, pl. 8, fig. 4, pl. 9, fig. 4 (wing, leg.). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil, Mexico. Refs. — Macfie, 1939a:145, fig. 3 (antenna, palpus, wing, spermatheca); Macfie, 1949:111, fig. 2 (male genitalia).
- TENUISQUAMIPES Wirth, 1952a:89, fig. 1 d-g (palpus, tibial spine, antenna, spermathecae). Type-locality: Juan Fernández Islands, Masatierra, Plazoleta del Yunque. Distr. — Juan Fernández Islands.
- TOWNSENDI Knab, 1915:111. Type-locality: Peru, Matucana. Distr. — Peru.
- UTAE Knab, 1915:109. Type-locality: Peru, Matucana. Distr. — Peru.
- VARIENNIS Wirth & Williams, 1957:8. Type-locality: Bermuda. Distr. — U.S.A. (Florida, Texas), Bermuda, Puerto Rico, Guatemala.
- WYGODZINSKYI Cavalieri, 1961:17, figs. 1-4 (wing, antenna, palpus, spermathecae). Type-locality: Argentina, Tierra del Fuego, Ushuaia. Distr. — Argentina.
- ZONOGASTER Ingram & Macfie, 1931:168, fig. 19b (male genitalia). Type-locality: Chile, Puerto Varas. Distr. — Chile.

Subgenus LASIOHELEA Kieffer

Centrorhynchus Lutz, 1914:62 (as genus). Type-species, *stylifer* Lutz (orig. des.). Preoccupied by *Centrorhynchus* Steven or Fischer Waldheim 1829 (Coleoptera) and *Centrorhynchus* Luche 1911 (Vermes).



- LASIOHELEA Kieffer, 1921b:115 (as genus). Type-species, *Atrichopogon pilosipennis* Kieffer (orig. des.) = *Forcipomyia velox* (Winnertz).
Refs. — Edwards, 1922:166 (discussion); Lane, 1945:359 (key); Saunders, 1964:463 (revision, all stages). Females suck vertebrate blood; larvae live on wet wood or moss.
- ATTENUATA Saunders, 1964:469, fig. 2 (all stages). Type-locality: Costa Rica, La Lola, larvae in moss on clay bank of drainage ditch. Distr. — Costa Rica.
- CORNUTA Saunders, 1964:464, fig. 1 (all stages). Type-locality: Costa Rica, Siquirres, Finca Theobroma, larvae on mossy, rotting wood in deep shade. Distr. — Costa Rica, Trinidad, Brazil (Santos).
- INSIGNIFORCEPS Macfie, 1939a:165, fig. 13 (wing, tarsal spine, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INTERMEDIA Saunders, 1964:468, fig. 2 f-i (larva, pupa, antenna, male genitalia). Type-locality: Costa Rica, Siquirres, Finca Theobroma, on mossy log in deep shade. Distr. — Costa Rica.
- QUASICORNUTA Saunders, 1964:467, fig. 2 a-e (larva, pupa, male genitalia, palpus, antenna). Type-locality: Costa Rica, Siquirres, Finca Theobroma, on mossy rotting wood in shade. Distr. — Costa Rica.
- SALTENSIS (Cavalieri), 1962b:360, figs. 1-5 (wing, antenna, spermatheca, palpus) (*Lasiohelea*). Type-locality: Argentina, Salta, road from Tartagal to Santa Victoria. Distr. — Argentina.
- STYLIFERA (Lutz), 1913:63, pl. 8, figs. 12, 22 (wing, palpus) (*Centrorhynchus*). Type-locality: Brazil, São Paulo, Tatuí (lectotype desig. Lane, 1945:362). Distr. — Costa Rica, Venezuela, Trinidad, Brazil. Refs. — Lane 1945:362, fig. 5 (antenna) (redescribed); Lane, 1947c:161, fig. 1 (larva; above water line on side of cement water tank); Ortiz, 1952b:241, figs. 1-7 (wing, head, palpus, spermatheca) (Venezuela); Macfie, 1953:97, fig. 1 (male genitalia) (Costa Rica); Wirth, 1956a:246 (redescribed; Costa Rica, pollinating *Hevea*).

Subgenus LEPIDOHELEA Kieffer

- LEPIDOHELEA Kieffer, 1917:364 (as genus). Type-species, *Ceratopogon chrysocephus* Kieffer (mon.).
- BRASILIENSIS Macfie, 1939a:153, fig. 7 (palpus, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- KUANOSCELES Macfie, 1939a:150, fig. 6 (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Subgenus METAFORCIPOMYIA Saunders

- Forcipomyia*, subg. METAFORCIPOMYIA Saunders, 1956:685. Type-species, *Forcipomyia cerifera* Saunders.
- CERIFERA Saunders, 1956:685, fig. 11 (all stages). Type-locality: Brazil, Rio de Janeiro, reared from larvae under rotting wood and under bark of logs in forest. Distr. — Brazil.
- PLUVIALIS Malloch, 1923:5. Type-locality: U.S.A., Maryland, Glen Echo. Distr. — Eastern North America (Ontario to Louisiana), Panama.




Subgenus MICROHELEA Kieffer

- MICROHELEA Kieffer, 1917:364 (as genus). Type-species, *Atrichopogon microtominus* Kieffer (desig. Kieffer, 1921:7, pointing out his orig. des. of *A. tropicus* Kieffer was a lapsus). = *Forcipomyia fuliginosa* (Meigen).
- Phasmidohelea* Mayer, 1937:233 (as genus). Type-species, *crudelis* Mayer (orig. des. = *Forcipomyia ixodooides* (Ficbrig-Gertz).
Refs. — Mayer, 1938:13, figs. 1-2 (female, mouthparts) (discussion); Séguy, 1941:86 (key to species); Wirth, 1972:564 (revision of Neotropical species); Wirth, 1971:229 (revision of "stick-ticks"). Biology. — Females suck blood from caterpillars, sawfly larvae, and stick insects.
- AMAZONICA Wirth, 1971:241, fig. 4 m-p (maxilla, mandible, palpus, spermathecae). Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- ERIOPHORA (Williston), 1896:279, pl. 9, fig. 40 (tarsus, antenna, palpus) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — U.S.A. (Florida), West Indies, Mexico, Central America to Panama. Refs. — Baker, 1907:117 (Cuba, on larva of *Melanchroia geometroides*); Wirth, 1972:571, figs. 27-31 (palpus, mandible, spermathecae, parameres).
- FULIGINOSA (Meigen); 1818:86 (*Ceratopogon*). Type-locality: Europe. Distr. — Worldwide. Refs. — Wirth, 1956b:357; 1971:233; 1972:567 (many figures; redescription; synonymy; habits, females suck blood from caterpillars and sawfly larvae).
crudelis Knab, 1914:66. Type-locality: Mexico, La Oaxaqueña, near Santa Lucrecia, attacking caterpillars. Preoccupied in *Forcipomyia* by *crudelis* Karsch, 1886.
tropicus Kieffer, 1917:297 (*Ceratopogon*). Type-locality Costa Rica, Higuito, San Mateo.
- GALAPAGENSIS (Coquillett), 1901b:372 (*Ceratopogon*). Type-locality: Galápagos Islands, Albermarle Island, Tagus Cove. Distr. — Galápagos Islands. Ref. — Wirth, 1969:573, figs. 1-2 (male genitalia, spermathecae) (redescription, synonymy); Wirth, 1972:572, figs. 32-36 (palpus, mandible, spermathecae, parameres).
- INSIGNIPALPIS Macfie, 1949:109, fig. 1 (palpus). Type-locality: Mexico, Chiapas, San Cristóbal north of Aurora. Distr. — Brazil, Colombia, Costa Rica, Mexico, Panama. Ref. — Wirth, 1972:573, figs. 37-41 (palpus, mandible, spermathecae, parameres).
- IXODOIDES (Ficbrig-Gertz), 1928:290, 5 figs. (*Ceratopogon*). Type-locality: Paraguay. Distr. — Brazil, Colombia, Costa Rica, Panama, Paraguay. Ref. — Wirth, 1971:236, fig. 4 a-d (palpus, maxilla, mandible, spermathecae) (redescription; distribution; synonymy).
crudelis Mayer, 1937:233, figs. 3-4 (mouthparts, tibial comb) (*Phasmidohelea*). Type-locality: Costa Rica, San José, Hamburg Farm, on phasmid *Crenozylus spinosus*. Preoccupied in *Forcipomyia* by *crudelis* Karsch and *crudelis* Knab.
wagneri Séguy, 1941:85, figs. 1-6 (female, leg, palpus, tarsus, claws) (*Phasmidohelea*). Type-locality Brazil, Rio de Janeiro, Organ Mountain, sucking blood of *Bacteria* (Phasmidae).
mayeri Forattini & Lane, 1955:4, figs. 1-4 (palpus, tarsus, antenna, spermatheca, egg). Unnecessary new name for *crudelis* Mayer 1937.
- LUTEISQUAMOSA Wirth, 1972:575, figs. 52-53 (palpus, spermathecae). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

- MICROTOMA (Kieffer), 1917:299, fig. 2 (palpus) (*Atrichopogon*). Type-locality: Paraguay, Asunción.
- MINISQUAMOSA Wirth, 1972:570, figs. 21-26 (palpus, mandible, spermathecae, aedeagus, parameres). Type-locality: British Honduras, Augustine. Distr. — Brazil, British Honduras, Costa Rica, El Salvador.
- OBESA Lima, 1928:85, figs. 1-5 (wing, body, antenna, palpus). Type-locality: Brazil, Rio Negro, São Gabriel, on phasmid. Distr. — Brazil, Colombia, Costa Rica, Panama. Refs. — Lane, 1947c:163 (female redescribed; Brazil, on phasmid); Wirth, 1971:239, figs. 1, 3, 4 e-h (habitus, many parts) (on phasmids, habits, distribution; female redescribed).
- PENULTIMATA Wirth, 1972:573, figs. 42-46 (palpus, mandible, spermathecae, parameres). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Central America and West Indies to Colombia and Brazil.
- SQUAMOSA Lutz, 1914:87, pl. 8, fig. 6, pl. 9, figs. 2-3 (wing, abdomen, eggs). Type-locality: Brazil, Rio de Janeiro, Mangueiros. Refs. — Ingram & Macfie, 1931:226 (redescribed, Argentina); Floch & Abonnenc, 1942c:1, fig. 1 a-f (wing, palpus, tarsus, male genitalia) (French Guiana); Wirth, 1972:569, figs. 16-20 (palpus, mandible, spermathecae, parameres).
- WILLISTONI Wirth, 1971:242, fig. 4 i-l (palpus, mandible, spermathecae, parameres). Type-locality: Puerto Rico, El Verde, Luquillo Mts., from phasmid. Distr. — West Indies (Jamaica, Puerto Rico), Brazil.

Subgenus PHYTOHELEA Remm

- Forcipomyia*, subg. PHYTOHELEA Remm, 1971:189. Type-species, *Lasiohelea bromelicola* Lutz (orig. des.).
- Apelma* Kieffer, authors
- Trichohelea* Goetghebuer, authors
- Ref. — Saunders, 1925:252 (all stages, habits, as *Apelma*; Saunders, 1956:695 (revision, all stages, as *Trichohelea*).
- ANTIGUENSIS Saunders, 1956:700, fig. 17 (all stages). Type-locality: Antigua, Falmouth Harbour, from larvae in epiphytic bromeliads. Distr. — West Indies (Antigua).
- BROMELICOLA (Lutz), 1914:84, pl. 8, fig. 1 (wing) (*Lasiohelea*). Type-locality: Brazil, Rio de Janeiro, Ilha do Governador. Distr. — Brazil, French Guiana. Refs. — Saunders, 1925:263, fig. 4 (all stages) (Brazil, from small bromeliads on rocks in Rio bay); Floch & Abonnenc, 1942c:2, fig. 1 i (wing, palpus, tarsus) (French Guiana); Lane, 1945:363, figs. 2, 3 (male genitalia) (lectotype designation; redescribed).
- CARIBBEANA Saunders, 1956:696, fig. 15 (all stages). Type-locality: Trinidad, Mount St. Benedict, from larvae in fallen epiphytic bromeliad in forest. Distr. — Trinidad, Tobago, Guyana.
- EDWARDSI (Saunders), 1925:260, fig. 3 (male, female) (*Apelma*). Type-locality: Brazil, Bahia; Rio de Janeiro, Paquetá; and Pernambuco; syntypes from larvae in terrestrial bromeliads. Distr. — Brazil.
- LOCOSA Saunders, 1956:701, fig. 18 (all stages). Type-locality: Trinidad, Mount St. Benedict, reared from *Bromelia pinguin*. Distr. — Trinidad.
- KEILINI (Saunders), 1925:265, fig. 5 (all stages) (*Apelma*). Type-locality: Brazil, Pernambuco, Dois Irmãos, from terrestrial bromeliads in woods.



- MAGNA (Saunders), 1925:266, fig. 6 (larva, pupa) (*Apelma*). Type-locality: Brazil, Pernambuco, Dois Irmãos, from terrestrial bromeliads in woods. Distr. — Brazil.
- OLIGARTIRA Saunders, 1956:698, fig. 16 (all stages). Type-locality: Puerto Rico, Corcega Beach, from larvae in leaf axils of *Bromelia pinguin*. Distr. — West Indies (Puerto Rico, Montserrat, Dominica), Guyana.

Subgenus PTEROBOSCA Macfie

- PTEROBOSCA Macfie, 1932:266 (as genus). Type-species, *Ceratopogon aescnosuga* Meijere (Macfie, 1940d:16).
Refs. — Johannsen, 1950:142 (key to American species); Wirth, 1956b:362 (review, check list).
- FARRI Wirth, 1966:29, figs. 1-5 (wing, antenna, palpus, tarsus, spermathecae). Type-locality: Jamaica, St. Thomas, Morant Bay, from wing of *Chrysopa*. Distr. — West Indies (Jamaica).
- FUSICORNIS (Còquille), 1905:63 (*Ceratopogon*). Type-locality: U.S.A., Florida, Biscayne Bay. Distr. — U.S.A. (Louisiana, Florida), Bermuda, West Indies (Puerto Rico), Brazil. Refs. — Johannsen, 1951:117 (records; synonym: *macfie*); Wirth, 1956b:362 (redescribed; distr.).
fuscicornis, error.
macfie Lima, 1937b:616, pls. 1-2 (head, wing, body, antenna, palpus, claws, spermatheca) (*Pterobosca*). Type-locality: Brazil, Rio de Janeiro, Tijuca, Alto da Boa Vista, on Odonata.
- INCUBANS (Macfie), 1937b:111 (*Pterobosca*). Type-locality: British Honduras, Cayo District, Mtn. Pine Ridge, on *Argia ulmeca* Calvert. Distr. — Mexico, British Honduras, Colombia, Argentina, West Indies (Puerto Rico). Refs. — Wirth, 1956b:363 (distribution); Cavalieri & Chiossone, 1965:65, figs. 1-18 (wing, tarsus, antenna, palpus, mandible, tibial comb, spermathecae, pharyngeal armature, eggs) (Argentina record).

Subgenus RHYNCHOFORCIPOMYIA Wirth & Dow

- Forcipomyia*, subg. RHYNCHOFORCIPOMYIA Wirth & Dow, 1972:863. Type-species, *Forcipomyia messersmithi* Wirth & Dow (orig. des.).
- BRACHYRHYNCHUS Wirth & Dow, 1972:867, fig. 3 a-1 (male, female, many parts). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia, Trinidad.
- CYLINDRICA Wirth & Dow, 1972:865, fig. 2 a-m (male, female, many parts). Type-locality: El Salvador, Cuscatlán, San Pedro Perulapán. Distr. — El Salvador.
- DORSALIS Wirth & Dow, 1972:870, fig. 5 g-n (male, female, many parts). Type-locality: Mexico, Sinaloa, 22 mi E Concordia. Distr. — El Salvador, Honduras, Mexico.
- GUAMAI Wirth & Dow, 1972:869, fig. 4 h-l (female, many parts). Type-locality: Brazil, Pará, Belém. Distr. — Brazil, Panama.
- MESSERSMITHI Wirth & Dow, 1972:864, fig. 1 a-n (male, female, many parts). Type-locality: El Salvador, Cuscatlán, San Pedro Perulapán. Distr. — Mexico through Central America to Colombia, West Indies (Jamaica).



PURACENSIS Wirth & Dow, 1972:867, fig. 4 a-g (female, many parts). Type-locality: Colombia, Cauca, Páramo de Purace. Distr. — Colombia.

ZETEKI Wirth & Dow, 1972:870, fig. 5 a-f (female, many parts). Type-locality: Panama, C. Z., Barro Colorado Island. Distr. — Colombia, Panama.

Subgenus SYNTHYRIDOMYIA Saunders

Forcipomyia, subg. SYNTHYRIDOMYIA Saunders, 1956:688. Type-species, *Lasiohelea acidicola* Tokunaga (orig. des.).

SANCTAELARAE Wirth, 1952a:90, fig. 1 h (male genitalia). Type-locality, Juan Fernández Islands, Santa Clara, El Corral. Distr. — Juan Fernández Islands. Ref. — Wirth, 1957:399 (Masatierra Island).

TENUIFORCEPS Macfie, 1939a:167, fig. 15 (palpus, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — U.S.A. (California, New Mexico), British Honduras, El Salvador, Trinidad, Brazil. Ref. — Dow & Wirth, 1972:199, fig. 11 a-m (many parts) (redescribed, distribution).

Subgenus THYRIDOMYIA Saunders

TRYRIDOMYIA Saunders, 1925:268 (as genus). Type-species, *palustris* Saunders (orig. des.).

Refs. — Wirth, 1970:429 (revision and key to Neotropical species); Dow & Wirth, 1972:177 (revision and key to Nearctic species). Biology. — Adults pollinating *Hevea* and *Theobroma*, etc.; larvae found in rotting plant materials and in ulcerating tree wounds.

COLOMBIAE Wirth, 1970:435, fig. 3 (antenna, palpus, wing, tarsus, spermatheca, male genitalia). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Mexico, Costa Rica, Panama, Colombia, Trinidad.

JAMAICENSIS Wirth, 1970:436, fig. 4 (antenna, palpus, wing, spermatheca, male genitalia). Type-locality: Jamaica, Hardwar Gap. Distr. — West Indies (Jamaica, Puerto Rico).

JIPAJAPAE Wirth, 1970:437, fig. 5 (antenna, palpus, wing, spermatheca, male genitalia). Type-locality: Panama, C. Z., Gamboa. Distr. — Costa Rica, Panama, Colombia, Ecuador, Trinidad.

NANA (Macfie), 1939a:171 (*Lasiohelea*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Mexico, Central and South America to Southern Brazil, West Indies. Refs. — Macfie, 1944:297, fig. a (male genitalia) (Trinidad, from cacao); Wirth, 1956a:247 (redescribed; pollinating *Hevea*; records); Wirth, 1970:432, fig. 1 a-f (wing, antenna, palpus, spermatheca, male genitalia) (redescribed).

NODOSA Saunders, 1959:43, fig. 5 (all stages). Type-locality: Costa Rica, Finca Theobroma near Siquirres. Distr. — U.S.A. (Arizona, Kansas, Mississippi), Mexico, Costa Rica, Colombia. Ref. — Dow & Wirth, 1972:186, fig. 4 (many parts) (redescribed; distribution).

SINUOSA Dow & Wirth, 1972:189, fig. 6 (many parts). Type-locality: Mexico, Sinaloa, 22 mi E Concordia, 3000 ft. Distr. — Mexico.

UNIVESICULA Macfie, 1939a:170. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



Subgenus TRICHOHELEA Goetghebuer

- Apelma* Kieffer, 1919:64. Type-species, *auronitens* Kieffer (desig. Macfie, 1940:16). Preoccupied by *Apelma* Billberg 1820, Coleoptera.
- TRICHOHELEA Goetghebuer, 1920:39. Type-species, *tonnoiri* Goetghebuer (mon.).
- Forcipomyia*, subg. *Neoforcipomyia* Tokunaga, in Tokunaga & Murachi, 1959:200. Type-species, *Ceratopogon pectinunguis* Meijere (orig. des.).
Refs. — Wirth, 1956b:361 (habits, list of species as *eques* group); Saunders, 1964:473 (diagnosis, all stages); Wirth & Messersmith, 1971:15 (revision and key to Nearctic species; synonymy).
- AERONAUTICA Macfie, 1935c:265, fig. (tarsus). Type-locality: Guyana, Tumatumari, on butterfly, *Catoblepia xanthus* L. Distr. — Venezuela, Guyana, French Guiana. Ref. — Wirth, 1956b:361 (distribution).
- DANAISI (Floch & Abonnenc), 1949:3 (*Lasiohelea*); also 1950a:72, fig. 2 (wing, antenna, spermatheca, palpus). Type-locality: Venezuela, Aragua, Rancho Grande, on butterfly *Danais eresimus*. Distr. — Venezuela.
- GONIOGNATHA Wirth & Messersmith, 1971:20, fig. 4 (antenna, palpus, mandible, maxilla, spermatheca, tarsus, male genitalia). Type-locality: Mexico, Sonora, 5 mi W Álamos. Distr. — U.S.A. (Arizona, Texas, Kansas), Mexico.
- INTREPIDA Macfie, 1936:228. Type-locality: Peru, Iquitos, on wing of dragonfly. Distr. — Peru.
- LIMNETIS Ingram & Macfie, 1931:169, fig. 19 (male genitalia). Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- MEXICANA Wirth, 1956b:361. Type-locality: Mexico, San Luis Potosí, Tamazunchale, on butterfly, *Pyrrhogyra otolais neis*. Distr. — Mexico. Ref. — Bauer, 1961:92 (habits).
- OPILIONIVORA (Lane), 1947c:159 (*Lasiohelea*). Type-locality: Brazil, São Paulo, Juquiá, sucking blood from Pbalangidae. Distr. — Brazil.
- SHANNONI (Ingram & Macfie), 1931:171 (*Lasiohelea*). Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Argentina.
- TRINIDADENSIS Saunders, 1964:476, fig. 5 (all stages). Type-locality: Trinidad, St. Augustine, reared from moss on tree root. Distr. — Trinidad.

Subgenus WARMKEA Saunders

- Forcipomyia*, subg. WARMKEA Saunders, 1956:671. Type-species, *Forcipomyia bicolor* Saunders (orig. des.).
- AERIA Saunders, 1956:675, fig. 7 (pupa, spermatheca, palpus, male genitalia). Type-locality: Puerto Rico, Mayaguez, reared from larvae in epiphytic bromeliads. Distr. — West Indies (Puerto Rico).
- LESLIEI Wirth, nom. nov.
bicolor Saunders, 1956:672, fig. 6 (all stages). Type-locality: Puerto Rico, Mayaguez, reared from larvae in leaf axils of *Alocasia macrorrhiza*. Distr. — West Indies (Puerto Rico). Preoccupied in *Forcipomyia* by *bicolor* Lutz 1914.
- LOURIEI (Macfie), 1935a:49 (*Lasiohelea*). Type-locality: Brazil, Piauí, Tutóia. Distr. — Brazil.



- NARTHEKOPHORA Macfie, 1939a: 166, fig. 14 (palpus, wing, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SPINOSA Saunders, 1956:676, fig. 7 (pupa, male genitalia, spermatheca, larva). Type-locality: Puerto Rico, Mayaguez, reared from larvae in leaf axils of *Alocasia macrorrhiza*. Distr. — West Indies (Puerto Rico).
- TERRESTRIS Saunders, 1964:479, fig. 6 (all stages). Type-locality: Trinidad, Las Hermanas Plantation near San Rafael. Distr. — Trinidad.
- TUBERCULATA Saunders, 1956:677, fig. 7 (pupa, spermatheca, male genitalia). Type-locality: Trinidad, North Range, reared from terrestrial *Bromelia pinquin*. Distr. — Trinidad.

Genus ATRICHOPOGON Kieffer

- Didymophleps* Weyenbergh, 1883:108. Type-species, *hortorum* Weyenbergh (mon.). Position uncertain.
- Ceratopogon*, subg. ATRICHOPOGON Kieffer, 1906:53. Type-species, *Ceratopogon exilis* Coquillett (Coquillett, 1910:512) = *levis* (Coquillett).
- Lophomyidium* Cordero, 1929:94. Type-species, *uruguayense* Cordero (orig. des.). Refs. — Ingram & Macfie, 1931:173 (key to Chilean species); Macfie, 1939a:173 (key to Neotropical species); Ewen & Saunders, 1958:671 (revision based on immature stages).
- ADAMSONI Macfie, 1937a:4. Type-locality: Trinidad, St. Augustine. Distr. — Mexico, Trinidad. Ref. — Macfie, 1949:112 (redescribed).
- ALBINENSIS Ingram & Macfie, 1931:228. Type-locality: Argentina, Bahia Blanca. Distr. — Costa Rica, Argentina.
- ALTIVOLANS Macfie, 1949:113, fig. 3 (abdominal armature). Type-locality: Mexico, Chiapas, between Santa Julia and Santa Rita north of Huixtla. Distr. — Mexico.
- ASSIMILIS Ingram & Macfie, 1931:176. Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina.
- AURICOMA Kieffer, 1917:300, fig. 3 (antenna). Type-locality: Colombia, Aracataca. Distr. — Colombia.
- BIFIDUS Ewen, in Ewen & Saunders, 1958:694, fig. 10 (all stages). Type-locality: Brazil, Rio de Janeiro, Niterói, above water line on partly submerged bamboo.
- BRASILIENSIS Macfie, 1939a:184. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- BREVIPALPIS Macfie, 1944:298, fig. 1 b, c (palpus, spermatheca). Type-locality: Trinidad, from moss on cacao trunk. Distr. — Trinidad.
- CARIBBEANUS Ewen, in Ewen & Saunders, 1958:683, fig. 5 a-j (all stages). Type-locality: West Indies, Tobago, larvae on wood in stream. Distr. — West Indies (Tobago).
- CHILENSIS Ingram & Macfie, 1931:175. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Brazil, Argentina, Chile.
- COLUMBIANUS Kieffer, 1917:303, fig. 7 (antenna). Type-locality: Colombia, Aracataca. Distr. — Colombia, Brazil. Ref. — Macfie, 1939a:177 (redescribed; Brazil).
- COSTALIS Macfie, 1939a:179. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



- COSTARICAE Macfie, 1953:98. Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- CRYPTOGAMUS Macfie, 1939a:182. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DEPILIS Macfie, 1939a:178. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DIDYMOTHECAE Macfie, 1953:99. Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- ECHINODES Macfie, 1939a:194. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- EUCNEMUS Macfie, 1939a:185. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FIEBRIGI Kieffer, 1917:302, fig. 4 (male antenna). Type-locality: Paraguay. Distr. — Paraguay.
- FIMBRIATUS (*gordoni* var.), Macfie, 1939a:194. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FLAVICAUDAЕ Macfie, 1939a:181, fig. 16 (wing, antenna, palpus). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
pilosior (*flavicaudae* var.), Macfie, 1939a:182. Type-locality: Brazil, Santa Catarina, Nova Teutônia.
- FLAVIPES Lutz, 1914:90, pl. 8, fig. 5 (wing). Type-locality: not stated. Distr. — Brazil. Ref. — Lane, 1945:359 (notes on type).
- FLUMINEUS Macfie, 1935a:52. Type-locality: Brazil, Piauí, Tutóia. Distr. — Brazil, Guyana. Ref. — Macfie, 1939a:177 (notes).
- FUSCULUS (Coquillett), 1901a:605 (*Ceratopogon*). Type-locality: U.S.A., New Jersey, Riverton. Distr. — North America (Saskatchewan to Ontario, s. to California and Florida), Costa Rica, Brazil. Refs. — Wirth, 1952e:118, fig. 3 (wing, palpus, leg, antenna, male genitalia) (redescription); Wirth, 1956a:242 (pollinating *Hevea*; Costa Rica, Brazil).
- GLABER Macfie, 1935a:50, fig. 1 (abdomen). Type-locality: Brazil, Piauí, Tutóia. Distr. — Costa Rica, Trinidad, Brazil. Refs. — Macfie, 1938:162, fig. 6 (male genitalia) (redescribed); Wirth, 1956a:243 (redescribed; pollinating *Hevea*).
- GLOBALIFER Macfie, 1939a:191. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- GORDONI Macfie, 1938:164, fig. 7 (palpus). Type-locality: Trinidad, St. Augustine. Distr. — Costa Rica, Trinidad, Brazil. Refs. — Macfie, 1939a:193 (notes); Macfie, 1940c:182 (distribution).
- GUIANENSIS Macfie, 1940c:184. Type-locality: Guyana, Kaieteur. Distr. — Guyana.
- HARRISI Macfie, 1938:163. Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.
- HORTORUM (Weyenbergh), 1883:110, figs. 1-3 (wing, antenna) (*Didymophleps*). Type-locality: Argentina, Córdoba. Distr. — Argentina. Position uncertain.
- INCULTUS Ewen. in Ewen & Saunders, 1958:699, fig. 12 (all stages). Type-locality: Costa Rica, Siquirres, reared from moss. Distr. — Costa Rica.
- INSIGNIPALPIS Macfie, 1940b:73. Type-locality: Brazil, Soledade and Simão (Northeast Brazil). Distr. — Brazil.

- INSIGNIVENTRIS Macfie, 1935a:51, fig. 2 (abdominal armature). Type-locality: Brazil, Piauí, Tutóia. Distr. — Trinidad, Brazil. Ref. — Macfie, 1937a:2, fig. (male genitalia).
- LACAJAE Macfie, 1953:100, fig. 2 (male genitalia). Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- LITURATUS (Williston), 1896:281 (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent).
- LONGICORNIS Ewen, in Ewen & Saunders, 1958:700, fig. 13 (all stages). Type-locality: Costa Rica, Siquirres, larvae on rotting wood. Distr. — Costa Rica.
- MACULIPENNIS Clastrier, 1968:89, fig. 2 (wing, antenna, palpus, male and female genitalia). Type-locality: French Guiana, Cayenne, Cabassou Forest. Distr. — French Guiana.
- MENDOZAE Ingram & Macfie, 1931:229. Type-locality: Argentina, Mendoza. Distr. — Argentina.
- NANUS Macfie, 1940c:184. Type-locality: Guyana, Mazaruni, High Forest. Distr. — Guyana.
- NEBULOSUS Macfie, 1939a:192. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NOVAETEUTONIAE Macfie, 1939a:189. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Costa Rica, Brazil.
- NUBECULOSUS Macfie, 1949:113. Type-locality: Mexico, Chiapas, El Vergel. Distr. — Mexico.
- OBFUSCATUS Ingram & Macfie, 1931:173. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile.
- OBNUBILUS Ingram & Macfie, 1931:175. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- OBSCURUS Ewen, in Ewen & Saunders, 1958:685; fig. 6 a-1 (all stages). Type-locality: Puerto Rico, Mayaguez, larvae on wet dead leaves bordering stream. Distr. — West Indies (Puerto Rico).
- PACHYCNEMUS Macfie, 1953:101. Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- FALLIDIPES Kieffer, 1917:301. Type-locality: Paraguay, San Bernardino. Distr. — Mexico, Costa Rica, Paraguay. Refs. — Macfie, 1949:114 (redescribed); Macfie, 1953:98 (redescribed).
- PALPALIS Macfie, 1939a:196, fig. 21 (palpus, antenna, abdominal armature, spermatheca, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PECTINATUS Macfie, 1939a:195, fig. 17 (palpus, antenna, abdominal armature, spermatheca). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PERUVIANUS Kieffer, 1917:302, fig. 5 (male antenna). Type-locality: Peru, Callanga. Distr. — Peru.
- PICEIVENTRIS Kieffer, 1917:301. Type-locality: Paraguay, San Bernardino. Distr. Paraguay.
- REMIGATUS Ewen, in Ewen & Saunders, 1958:687, fig. 7 a-j (all stages). Type-locality: Brazil, Petrópolis, larvae on wet sticks over stream. Distr. — Brazil.



- RUSTICUS Macfie, 1939a:190. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- SANCTAECLARAE Macfie, 1949:114. Type-locality: Mexico, Chiapas, Puerto Ocos, near Hacienda Santa Clara. Distr. — Mexico.
- SANCTILAURENTII Kieffer, 1917:303, fig. 6 (antenna). Type-locality: Paraguay, San Lorenzo. Distr. — Paraguay.
- SAUNDERSI Ewen, in Ewen & Saunders, 1958:691, fig. 9 a-j (all stages). Type-locality: Puerto Rico, Mayaguez, on wood in stream. Distr. — West Indies (Puerto Rico).
- SEQUAX (Williston), 1896:282, plate 9, fig. 48 (wing) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent). N. COMB.
- TAENIATUS Macfie, 1939a:186, fig. 20 (wing, antenna, palpus, spermatheca). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- THERSITES (Williston), 1896:280, pl. 9, fig. 44 (wing) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent).
thirsites, error, Williston, 1896:277.
- TRINIDADENSIS Macfie, 1937a:5, fig. (male genitalia). Type-locality: Trinidad, St. Augustine. Distr. — Trinidad.
- TUBERCULATUS Ewen, in Ewen & Saunders, 1948:709, fig. 17 (all stages). Type-locality: Trinidad, Maracas, larvae on rotting cacao leaves. Distr. — Trinidad.
- UMBRATILIS Macfie, 1935a:51. Type-locality: Brazil, Piauí, Tutóia. Distr. — Guyana, Brazil. Ref. — Macfie, 1940c:183 (redescribed).
- URUGUAYENSIS (Cordero), 1929:95, figs. 1-12 (male, antenna, wing, palpus, mouthparts, legs, claws, male genitalia) (*Lophomyidium*). Type-locality: Uruguay, Tacuarembó. Distr. — Uruguay.
- UTRICULARIUS Macfie, 1953:102. Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- WARMKEI Wirth, 1956a:243, fig. 1 (female, palpus, mandible, male genitalia). Type-locality: Puerto Rico, Mayaguez, in *Hevea* flowers. Distr. — U.S.A. (Florida), West Indies (Puerto Rico).
- WOODFORDI Macfie, 1938:162. Type-locality: Trinidad, St. Augustine. Distr. — Costa Rica, Guyana, Brazil, Trinidad. Refs. — Macfie, 1939a:117 (notes); Macfie, 1939c:183 (redescribed).

Subfamily DASYHELEINAE

Genus DASYHELEA Kieffer

- DASYHELEA Kieffer, 1911b:5. Type-species, *halophila* Kieffer (mon.).
Pseudoculicoides Malloch, 1915b:309. Type-species, *Ceratopogon mutabilis* Coquillett (orig. des.).
Ref. — Ingram & Macfie, 1931:177 (key to Chilean species).
- ALBOPICTA Ingram & Macfie, 1931:187, fig. 22c (male genitalia). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina, Chile.
- ANDENSIS Ingram & Macfie, 1931:180, fig. 22a (male genitalia). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina.
- ARGENTINENSIS Kieffer, 1925c:92. Type-locality: Argentina. Distr. — Argentina.



- AUSTRALIS Wirth, 1952a:92, fig. 1 a-c (wing, palpus, male genitalia). Type-locality: Juan Fernández Islands, Masatierra, Bahía Cumberland, Grutas de los Patriotas. Distr. — Juan Fernández Islands.
- BAHAMENSIS (Johnson), 1908:71 (*Ceratopogon*). Type-locality: Bahamas, Andros Island, Mangrove Cay. Distr. — U.S.A. (Florida), West Indies (Bahamas).
- CACTORUM Wirth & Hubert, 1960:642, figs. 3, 24 (wing, palpus, spermatheca, male genitalia). Type-locality: México, Baja California, San Felipe, reared from *Lophocereus cactus*. Distr. — U.S.A. (Arizona, California), Mexico (Baja California).
- CALVESCENS Macfie, 1938:157, fig. (male genitalia). Type-locality: Hawaii, Oahu, Hanauma Bay and Waianae Coast (syntypes). Distr. — Hawaiian Islands, Mexico (Revillagigedo Islands).
- CHILENSIS Ingram & Macfie, 1931:186. Type-locality: Chile, Casa Pangué. Distr. — Chile.
- COLUMBIANA (Kieffer), 1917:304, fig. 3 (antenna) (*Culicoides*). Type-locality: Colombia, Aracataca. Distr. — Colombia.
- FILIBRANCHIA (Lutz), 1914:85, fig. (pupa) (*Ceratopogon*). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil. Ref. — Lane, 1945:364, figs. 6-11 (antenna, wing, male genitalia, pupa).
- FLAVICAUDA Macfie, 1939a:201. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- HIPPOLYTAE Macfie, 1935a:54. Type-locality: Brazil, Piauí, Tutóia. Distr. — Brazil.
- HIRTIPES (Kieffer), 1917:305, fig. 9 (antenna) (*Culicoides*). Type-locality: Peru, Callanga. Distr. — Peru.
- LACUSTRIS Ingram & Macfie, 1931:190. Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina.
- LUTEOGRISEA Wirth & Williams, 1957:10. Type-locality: Bermuda, Spittal Pond. Distr. — U.S.A. (Florida to Texas), Bermuda, West Indies (Bahamas).
- MACULATA Macfie, 1943:119, fig. 1 (male genitalia). Type-locality: Bahamas, New Providence, Lake Cunningham. Distr. — West Indies (Bahamas).
- MONTICOLA Ingram & Macfie, 1931:188, fig. 24 (male genitalia, spermatheca). Type-locality: Chile, Casa Pangué. Distr. — Chile.
- PARACINCTA Wirth, 1969:576, figs. 4-6 (male genitalia). Type-locality: Galápagos Islands, Isla Santa Cruz, Academy Bay. Distr. — Galápagos Islands.
- PARAHYBAE Macfie, 1940b:74. Type-locality: Brazil, Simão, Double (?). Distr. — Brazil.
- PATAGONICA Ingram & Macfie, 1931:182, fig. 22 b (male genitalia). Type-locality: Argentina, Rio Negro, Lake Gutiérrez. Distr. — Argentina.
- PAULISTANA Forattini & Rabello, 1957:245, figs. 11-16 (all stages). Type-locality: Brazil, São Paulo, Parque da Água Funda. Distr. — Brazil.
- PENTALINEATA Wirth & Hubert, 1960:644, figs. 4, 26 (wing, palpus, spermatheca, male genitalia, scutum). Type-locality: Mexico, Baja California, San Felipe, reared from *Lophocereus cactus*. Distr. — U.S.A. (California), Mexico (Baja California).
- PENTHESILEAE Macfie, 1935:55. Type-locality: Brazil, Piauí, Tutóia. Distr. — Brazil.
- PHILOTHERMA Macfie, 1939a:202, fig. 18 (spermatheca, male genitalia). Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.



- PUMILA Macfie, 1939a:202, fig. 18 (spermatheca, male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PUSILLA (Lutz), 1913:65 (*Centrorhynchus*). Type-locality: Brazil, Serra da Boacaina, Bonito. Distr. — Brazil. Ref. — Lane, 1945:365, figs. 12, 13 (leg, wing) (types redescribed); Forattini & Rabello, 1957:241, figs. 1-10 (all stages).
- PYGMAEA (Williston), 1896:278, pl. 9, fig. 37 (wing) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent).
- PYRSONOTA Macfie, 1953: 104, fig. 5 (male genitalia). Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- REYNOLDSI Ingram & Macfie, 1931:185, fig. 23 d-f (male genitalia). Type-locality: Chile, Tierra del Fuego, Río Grande, Estancia Viamonte. Distr. — Chile.
- SANCTAEMARTHAE (Kieffer), 1917:306 (*Culicoides*). Type-locality: Colombia, Santa Marta. Distr. — Colombia.
- SCISSURAE Macfie, 1937:15, fig. 5 (male genitalia). Type-locality: Trinidad, St. Augustine. Distr. — Neotropical, Bermuda. Ref. — Macfie, 1953:103 (redescribed, Costa Rica).
- SHANNONI Ingram & Macfie, 1931:183, fig. 23 a-c (male genitalia). Type-locality: Chile, Ensenada. Distr. — Chile.
- SPATHICERCUS Wirth, 1969:577, figs. 8-9 (male genitalia, female abdomen). Type-locality: Galápagos Islands, Isla Fernandina, Punta Espinosa. Distr. — Galápagos Islands.
- THALESTRIS Macfie, 1935:55. Type-locality: Brazil, Piauí, Tutóia. Distr. — Brazil, Guyana, Trinidad. Ref. — Macfie, 1937:12, fig. 4 (male genitalia, spermatheca) (redescribed, Trinidad).
- VILLOSIPES (Kieffer), 1917:305, fig. 10 (antenna) (*Culicoides*). Type-locality: Paraguay. Distr. — Paraguay. N. COMB.
- VIRIDANS Forattini & Rabello, 1957:247, figs. 19-22 (wing, antenna, palpus, spermatheca). Type-locality: Brazil, Rio de Janeiro, km 47, Estrada Rio-São Paulo. Distr. — Brazil.

Genus PARADASYHELEA Macfie

- PARADASYHELEA Macfie, 1940:17. Type-species, *Dasyhelea brevipalpis* Ingram & Macfie (orig. des.).
Refs. — Wirth & Lee, 1959:114 (revision); Wirth & Blanton, 1969b:97 (notes).
- BREVIPALPIS (Ingram & Macfie), 1931:178, fig. 20 (antenna, palpus, male genitalia), fig. 21 (wing) (*Dasyhelea*). Type-locality: Argentina, Rio Negro, Lake Nahuel Huapi. Distr. — Argentina.

Subfamily CERATOPOGONINAE

Tribe CULICOIDINI

Genus CULICOIDES Latreille

- CULICOIDES Latreille, 1809:251. Type-species, *punctatus* Latreille (mon).
Refs. — Lima, 1937a:411 (key to Neotropical species); Adamson, 1939:79 (habits, Trinidad); Fox, 1946:249 (key to Caribbean species); Barbosa, 1947:3 (revision, key to Neotropical species); Macfie, 1948:69

(key to Caribbean species); Vargas, 1949:191 (list of world species); Ortiz, 1951c:8 (key to Venezuelan species); Ortiz & León, 1955:564 (review, key to Ecuadorean species); Aitken, 1957:23 (habits, Trinidad); Forattini, 1957:161 (revision of Neotropical species); Forattini, Rabello, & Pattoli, 1957:312 (larval habitats, photos); Forattini, Rabello, & Pattoli, 1958:1 (larval habitats, biology, adult habits of Brazilian species); Vargas, 1958:75 (list of Mexican species); Vargas & Garza, 1959:203 (key to subgenera, list of world species); Wirth & Blanton, 1959:237 (revision and key to Panama species); Vargas, 1960:35 (key to subgenera); Cavalieri & Chiossonc, 1966b:145 (notes on Argentine species); Wirth & Lee, 1967:1 (species of Colombian Paramos); Davies, 1967:39 (larval habits, mangrove species, Jamaica); Gutsevich, Garcia, & Gonzalez, 1969:1 (distribution; Cuba); Vargas, 1969:783 (disease transmission); Ortiz, 1969:31 (group classification); Linley & Davies, 1971:264 (biology and control in Caribbean area).

Subgenus ANILOMYIA Vargas

Culicoides, subg. ANILOMYIA Vargas, 1960:37. Type-species. *Culicoides covagarciai* Ortiz (orig. des.).

Refs. — Wirth & Blanton, 1956b:58 (rev., key, as *covagarciai* group); Wirth & Blanton, 1970d:141 (revision and key to *nigrigenus* group).

CHRYSONOTUS Wirth & Blanton, 1956b:226, fig. 9 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Chiriquí, El Volcán. Distr. — Panama. Refs. — Forattini, 1957:256, fig. 31 (wing, thorax, palpus, antenna, eyes, male genitalia); Wirth & Blanton, 1959:312, fig. 22 (wing, scutum, palpus, tibial comb, spermathecae, male genitalia).

COVAGARCIAI Ortiz, 1950c:457, plates 5, 6 (wing, scutum, spermathecae, palpus, antenna, male genitalia). Type-locality: Venezuela, D.F., Caracas. Distr. — Panama, Venezuela. Refs. — Wirth & Blanton, 1956b:216, fig. 3 (wing, palpus, spermathecae, male genitalia); Forattini, 1957:273, fig. 37 (wing, thorax, palpus, antenna, eyes, male genitalia); Wirth & Blanton, 1959:304, fig. 18 (wing, scutum, palpus, tibial comb, spermathecae, male genitalia) (synonym: *beebei*).

beebei Fox, 1952:366, fig. 4 (wing, palpus, spermathecae, tibial comb). Type-locality: Venezuela, Rancho Grande.

DECOR (Williston), 1896:281, plate 9, fig. 35 (wing) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (Dominica, St. Vincent, St. Lucia). Ref. — Wirth & Blanton, 1970d:145, fig. 2 (antenna, palpus, eyes, wing, leg, spermathecae, male genitalia).

DOMINICANUS Wirth & Blanton, 1970d:146, fig. 3 (antenna, palpus, eyes, wing, spermathecae, leg, male genitalia). Type-locality: Dominica, Pont Cassé. Distr. — West Indies (Dominica).

EFFERUS Fox, 1952:365, fig. 3 (wing, antenna, palpus, tibial comb, spermathecae). Type-locality: Peru, Rio Charape. Distr. — Honduras, Nicaragua, Panama, Peru. Refs. — Wirth & Blanton, 1956a:217, fig. 4 (wing, palpus, spermathecae, male genitalia); Forattini, 1957:266, fig. 35 (wing, thorax, palpus, antennae, eyes, male genitalia); Wirth & Blanton, 1959:299, fig. 15 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

FARRI Wirth & Blanton, 1970d:148, fig. 4 (antenna, wing, palpus, eye separation, leg, spermathecae, male genitalia). Type-locality: Jamaica, Hardwar Gap. Distr. — West Indies (Jamaica).



- HAYESI Matta, 1967:75, figs. 1-10 (all stages). Type-locality: Honduras, Dist. Central, La Tigra. Distr. — Mexico, Honduras. Ref. — Wirth & Blanton, 1970d:149 (notes).
- LUTEALARIS Wirth & Blanton, 1956b:225, fig. 8 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Chiriquí, El Volcán. Distr. — Costa Rica, Panama. Refs. — Forattini, 1957:260, fig. 32 (wing, palpus, male genitalia); Wirth & Blanton, 1959:310, fig. 21 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- MARSHI Wirth & Blanton, 1956b:220. Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama. Ref. — Wirth & Blanton, 1959:302, fig. 17 (wing, thorax, palpus, tibial comb, spermathecae).
- METAGONATUS Wirth & Blanton, 1956b:221, fig. 6 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Cerro Campana. Distr. — Nicaragua, Panama. Refs. — Forattini, 1957:38 (wing, thorax, palpus, antenna, eye separation, male genitalia); Wirth & Blanton, 1959:306, fig. 19 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- NIGRIGENUS Wirth & Blanton, 1956b:222, fig. 7 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Mexico, Panama, Colombia, Trinidad. Refs. — Forattini, 1957:262, fig. 33 (wing, thorax, palpus, antenna, eye separation; male genitalia); Wirth & Blanton, 1959:308, fig. 20 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- POPAYANENSIS Wirth & Lee, 1967:10, fig. 3, plate 1, fig. 4 (wing, antenna, palpus, eyes, tibial comb, spermathecae, male genitalia). Type-locality: Colombia, Cauca, Purace. Distr. — Colombia.
- ROSTRATUS Wirth & Blanton, 1956b:218, fig. 5 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Cerro Campana. Distr. — Panama. Refs. — Forattini, 1957:268, fig. 36 (wing, thorax, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:301, fig. 16 (wing, mesonotum, palpus, tibial comb, spermathecae, male genitalia).

Subgenus AVARITIA Fox

- Culicoides*, subg. AVARITIA Fox, 1955:218. Type-species, *Ceratopogon obsoletus* Meigen (orig. des.).
Ref. — Wirth & Lee, 1967:4 (diagnosis; revision of *andicola* group).
- ANDICOLA Wirth & Lee, 1967:5, fig. 1 a-c, plate 1, fig. 1 (wing, antenna, palpus, spermathecae). Type-locality: Colombia, Cauca, Páramo de Purace, Lago Buey. Distr. — Colombia.
- ORJUELAI Wirth & Lee, 1967:6, fig. 1 d-g, plate 1, fig. 2 (wing, eyes, antenna, palpus, spermathecae). Type-locality: Colombia, Cauca, Páramo de Purace. Distr. — Colombia.
- PURACENSIS Wirth & Lee, 1967:7, fig. 2, a-f, plate 1, fig. 3 (wing, antenna, eye separation, palpus, spermathecae, male genitalia). Type-locality: Colombia, Páramo de Purace, Lago Buey. Distr. — Colombia.
- PUSILLODES Wirth & Blanton, 1955a: 104, fig. 3 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Guatemala to Panama. Refs. — Forattini, 1957:287, fig. 45 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:290, fig. 11 (wing, palpus, thorax, tibial comb, spermathecae, male genitalia).



PUSILLUS Lutz, 1913:52, plate 7, fig. 4 (wing). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Mexico to Brazil, West Indies, Galápagos Islands. Refs. — Macfie, 1938:165, fig. 8 a-c (male genitalia); Forattini, 1957:284, fig. 44 (wing, thorax, palpus, eye separation, antenna, male genitalia); Wirth & Blanton, 1959:292, fig. 12 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

Subgenus BELTRANMYIA Vargas

Culicoides, subg. BELTRANMYIA Vargas, 1953:34. Type-species, *Culicoides crepuscularis* Malloch (orig. des.).

CREPUSCULARIS Malloch, 1915a:303, plate 20, figs. 7, 16, plate 22, fig. 7, plate 23, fig. 3 (wing, thorax, antenna, male genitalia). Type-locality: U.S.A., Illinois, Dubois. Distr. — North America (British Columbia to Quebec, south to California, and Florida), Bermuda, Mexico, Honduras, Costa Rica.

Subgenus CULICOIDES Latreille

ELUTUS Macfie, 1948:75, fig. 1 (wing). Type-locality: Mexico, Chiapas, El Carrizal. Distr. — Mexico to Panama. Refs. — Wirth & Blanton, 1956g:58, fig. 1 (wing, palpus, spermathecae, male genitalia); Forattini, 1957:250, fig. 29 (wing, thorax, palpus, eye separation, antenna, male genitalia); Wirth & Blanton, 1959:295, fig. 13 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

tristriatulus Hoffman, of Vargas, 1945:45 (Mexico records; fig. male genitalia).

LUTEOVENUS Root & Hoffman, 1937:156, figs. 5, 28 (wing, male genitalia). Type-locality: Mexico, D. F., San Jacinto. Distr. — Mexico to Panama. Refs. — Wirth & Blanton, 1956b:214, fig. 2 (wing, palpus, spermathecae, male genitalia); Forattini, 1957:253, fig. 30 (wing, thorax, palpus, antenna, eyes, male genitalia); Wirth & Blanton, 1959:297, fig. 14 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

NEOPULICARIS Wirth, 1955b:355, fig. 1 (wing, palpus, male genitalia). Type-locality: U.S.A., Texas, Kerrville. Distr. — U.S.A. (Texas), Mexico to Costa Rica. Ref. — Wirth & Blanton, 1969a:229, fig. 9 (antenna, palpus, wing, eye separation, spermathecae, male genitalia).

yukonensis Hoffman, of Vargas, 1945:45 (Mexico record).

Subgenus DIPHAOMYIA Vargas

Culicoides, subg. DIPHAOMYIA Vargas, 1960:40. Type-species, *Culicoides baueri* Hoffman (orig. des.).

BLANTONI Vargas & Wirth, 1955:33, figs. a-d (wing, palpus, male genitalia, spermathecae). Type-locality: Mexico, Tamaulipas, Ciudad Mante. Distr. — U.S.A. (Texas), Mexico (Morelos, Sinaloa, Tamaulipas).

EVANSI Wirth & Blanton, 1959:342, fig. 35 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Chiriquí, El Volcán. Distr. — Panama.

HAEMATOPOTUS Malloch, 1915a:302. Type-locality: U.S.A., Illinois, Urbana. Distr. — U.S.A., southern Canada, Mexico. Ref. — Vargas, 1945:44 (Mexico records).



- IRIARTEI Fox, 1952:368, fig. 6 (wing, palpus, spermathecae, tibial comb). Type-locality: Venezuela, Zulia, La Salina. Distr. — Honduras, Panama, Venezuela. Ref. — Wirth & Blanton, 1959:344, fig. 36 (wing, palpus, tibial comb, spermathecae, male genitalia) (synonymy, distribution).
- vargasi* Wirth & Blanton, 1953a:74, fig. 5 (thorax, wing, palpus, male genitalia). Type-locality: Panama, Los Santos, Las Tablas.
- baueri* Hoffman, of authors; Ortiz & Mirsa, 1951:599, figs. 9-12, 31 (wing, thorax, spermathecae, antenna, palpus) (Venezuela); Forattini, 1957:475, fig. 118 (wing, thorax, palpus, antenna, eyes, male genitalia) (synonymy, distribution).
- bahueri*, error for *baueri* Hoffman, of Ortiz, 1951.
- MARINKELLEI Wirth & Lee, 1967:13, fig. 4 a-f, plate 1, fig. 5 (wing, antenna, eye separation, palpus, spermathecae, male genitalia). Type-locality: Colombia, Cauca, Páramo de Purace, Lago Buey. Distr. — Colombia.

Subgenus DRYMODESMYIA Vargas

- Culicoides*, subg. DRYMODESMYIA Vargas, 1960:40. Type-species, *Culicoides copiosus* Root & Hoffman (orig. des.).
Ref. — Wirth & Hubert, 1960:646 (revision, key, as *copiosus* group).
- ARIZONENSIS Wirth & Hubert, 1960:655, figs. 14, 35 (wing, palpus, spermathecae, male genitalia). Type-locality: U. S. A., Arizona, Maricopa Co., Morristown. Distr. — U. S. A. (Arizona, California), Mexico (Baja California).
- BAKERI Vargas, 1954:27, fig. A, 1-5 (wing, palpus, antenna, male genitalia). Type-locality: Mexico, D.F., Lomas de Chapultepec. Distr. — Mexico. Ref. — Wirth & Hubert, 1960:652, fig. 31 (palpus, spermathecae, male genitalia) (redescribed from type).
- BORINQUENI Fox & Hoffman, 1944:110, fig. 4 (wing). Type-locality: Puerto Rico, Palmas Abajo, reared from tree hole. Distr. — West Indies (Jamaica, Puerto Rico). Refs. — Fox, 1949:30, fig. 2 (wing, palpus, spermathecae, male genitalia); Forattini, 1957:368, fig. 74 (wing, palpus, male genitalia, pupa); Wirth & Hubert, 1960:658, fig. 18 (wing); Linley, 1965b:57, figs. (pupa) (Jamaica).
species 2; Fox, 1942:417, fig. 4 (pupa).
- COPIOSUS Root & Hoffman, 1937:171, fig. 26 (male genitalia). Type-locality: Mexico, D.F., San Jacinto. Distr. — U.S.A. (California to Texas), Mexico (Baja California, Sonora to D.F.). Ref. — Wirth & Hubert, 1960:657, figs. 16, 37 (wing, palpus, spermathecae, male genitalia).
- INSOLATUS Wirth & Hubert, 1960:654, figs. 13, 34 (wing palpus, spermathecae, male genitalia). Type-locality: Mexico, Baja California, San Felipe. Distr. — Mexico (Baja California).
- JAMAICENSIS Edwards, 1922:165, fig. 10 (wing) (as *loughnani* var). Type-locality: Jamaica, Kingston. Distr. — U.S.A. (s. Texas), Mexico and Central America to Venezuela, West Indies. Refs. — Wirth & Blanton, 1959:339, fig. 34 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Wirth & Hubert, 1960:649, fig. 7 (wing).
loughnani Edwards (misidentification); Ortiz & Mirsa, 1952a:271, figs. 39-45 (palpus, antenna, wing, spermathecae, male genitalia); in part, of Forattini, 1957:412, fig. 93.



- LOUGHNANI Edwards, 1922:165, fig. 9 (wing). Type-locality: Jamaica, Kingston. Distr. — U.S.A. (Texas, Florida), West Indies (Cuba, Bahamas, Jamaica). Ref. — Wirth & Hubert, 1960:649, figs. 6, 27 (wing, palpus, spermathecae, male genitalia).
- PANAMENSIS Barbosa, 1947:22, plate 1, fig. 4 (palpus), plate 3, fig. 2 (male genitalia), plate 9, fig. 2 (thorax). Type-locality: Panama, C.Z., Barro Colorado Island, from *Heliconia* bracts. Distr. — Mexico, Guatemala, Panama, West Indies (Jamaica). Ref. — Wirth & Blanton, 1959:334, fig. 32 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- alambiculorum* Macfie, 1948:81, fig. 5 (wing). Type-locality: Mexico, Chiapas, El Vergel.
- PILOSUS Wirth & Blanton, 1959:332, fig. 31 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- POIKILONOTUS Macfie, 1948:82, fig. 7 (wing). Type-locality: Mexico, Chiapas, El Vergel. Distr. — Mexico to Panama, Venezuela. Refs. — Forattini, 1957:407, fig. 91 (wing, palpus, male genitalia); Wirth & Blanton, 1959:337, fig. 33 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- cacozelus* Macfie, 1948:85, fig. 10 (wing). Type-locality: Mexico, Chiapas, El Vergel.
- hertigi* Wirth & Blanton, 1953b:229, fig. 1 (wing, palpus, thorax, male genitalia). Type-locality: Panama, Coclé, Río Hato.
- RYCKMANI Wirth & Hubert, 1960:656, figs. 15, 36 (wing, palpus, spermathecae, male genitalia). Type-locality: U.S.A., California, Los Angeles Co., San Dimas Canyon. Distr. — U.S.A. (Arizona, California, Texas), Mexico (Baja California).
- SITIENS Wirth & Hubert, 1960:652, figs. 10, 30 (wing, palpus, spermathecae, male genitalia). Type-locality: U.S.A., California, Los Angeles Co., San Dimas Canyon, reared from *Opuntia* cactus. Distr. — U.S.A. (Arizona, California), Mexico (Baja California).
- TORRIDUS Wirth & Hubert, 1960:654, figs. 12, 33 (wing, palpus, thorax, spermathecae, male genitalia). Type-locality: Mexico, Baja California, San Felipe. Distr. — Mexico (Baja California).
- WIRTHOMYIA Vargas, 1953b:227, figs. (wing, male genitalia). Type-locality: Mexico, Guerrero, Iguala. Distr. — Mexico (Guerrero).

Subgenus GLAPHIROMYIA Vargas

- Culicoides*, subg. GLAPHIROMYIA Vargas, 1960:40. Type-species, *Culicoides scopus* Root & Hoffman (orig. des.).
- SCOPUS Root & Hoffman, 1937:170, figs. 24, 36 (wing, male genitalia). Type-locality: Mexico, D.F., San Jacinto. Distr. — Mexico, Panama. Ref. — Wirth & Blanton, 1959:347, fig. 37 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

Subgenus HOFFMANIA Fox


- Culicoides*, subg. HOFFMANIA Fox, 1948:21. Type-species, *Culicoides inamollae* Fox & Hoffman (orig. des.) = *insignis* Lutz.



Refs. — Ortiz, 1950c:437 (review of Neotropical species); Wirth & Blanton, 1956f:305 (revision of *guttatus* group); Wirth & Blanton, 1956c:95 (revision of *hylas* group); Wirth & Blanton, 1968a:201 (revision of *hylas* group).

- AITKENI Wirth & Blanton, 1968a:214, fig. 7 (wing, antenna, palpus, legs, spermathecae). Type-locality: Trinidad, Macqueripe. Distr. — Trinidad.
- BIMACULATUS Floch & Abonnenc, 1942b:3, fig. 1 a (wing, palpus). Type-locality: French Guiana, Cayenne. Distr. — French Guiana.
- BRASILIANUS Forattini, 1956b:81, figs. 1-5 (wing, antenna, eye separation, palpus, thorax). Type-locality: Brazil, São Paulo, São Vicente. Distr. — Brazil.
- COUTINHOI Barretto, 1944:96, fig. 13 (male antenna, palpus, genitalia), plate 13, fig. 5 (wing). Type-locality: Brazil, São Paulo, Juquiá, Prainha. Distr. — French Guiana, Brazil. Refs. — Barbosa, 1947:13, plate 2, fig. 2 (palpus), plate 9, fig. 1 (thorax); Wirth & Blanton, 1956f:314, fig. 2 (wing, palpus, spermathecae, male genitalia) (redescribed).
- DIABOLICUS Hoffman, 1925:294, plate 1, fig. 7, plate 2, fig. 12 (wing, thorax). Type-locality: Panama, Cabima, "bites like the devil". Distr. — Mexico to Panama. Refs. — Vargas, 1944:163, figs. 3-8 (male genitalia, legs, antenna, palpus) (Mexico); Fox, 1948:24, fig. 5 (palpus, male genitalia) (Panama); Wirth & Blanton, 1956f:316, fig. 3 (wing, palpus, spermathecae, male genitalia) (distribution); Wirth & Blanton, 1959:109, figs. 2, 7 (habitus, parts) (synonymy, distribution, redescribed).
- FILARIFER Hoffman, 1939:172, figs. 1-3 (wing, thorax, palpus). Type-locality: Mexico, Chiapas, El Vergel, Huixtla. Distr. — Mexico, Central America.
- FLAVIVENULA Lima, 1937a:418, fig. 4 (palpus). Type-locality: Brazil, Rio de Janeiro, Japuiba, Angra dos Reis. Distr. — Brazil. Refs. — Forattini, 1954a:135, figs. 1, 2 (wing, palpus, male genitalia); Forattini, 1957:235, fig. 24 (wing, thorax, palpus, antenna, male genitalia) (distribution); Wirth & Blanton, 1956f:318, fig. 9 (wing, palpus, spermathecae) (topotypes redescribed).
- FOXI Ortiz, 1950:461. Type-locality: Puerto Rico, Campo Tortuguero (Fox, 1949:31, as *guttatus*). Distr. — Central and South America from Honduras to Argentina (Corrientes), West Indies (Jamaica, Puerto Rico, Dominica). Refs. — Ortiz, 1951c:4, plate 2 (male genitalia, wing, palpus, spermathecae, thorax); Wirth & Blanton, 1956f:309, fig. 1 (wing, spermathecae, palpus, male genitalia) (distribution); Forattini, 1957:205, fig. 15 (wing, thorax, antenna, palpus, male genitalia) (Brazil distribution); Wirth & Blanton, 1959:283, fig. 8 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonymy; Panama, distribution).
- diabolicus* Hoffman, misidentification; Floch & Abonnenc, 1942:2, fig. 2a (wing, palpus) (French Guiana).
- guttatus* (Coquillett), of authors
- GUTTATUS (Coquillett), 1904a: 35 (*Ceratopogon*). Type-locality: Brazil, São Paulo, Serra da Cantareira. Distr. — Brazil. Refs. — Lima, 1937a:416, fig. 1 (wing) (Brazil); Barretto, 1944:91, fig. 9 (male palpus, genitalia), plate 1, fig. 2 (wing) (male described); Forattini, 1955:103, fig. 1 (wing, palpus, thorax, eye separation, antenna, male genitalia); Wirth & Blanton, 1956f:312, fig. 7 (wing) (type redescribed); Forattini, 1957:208, fig. 16 (many parts).
- HELICONIAE Fox & Hoffman, 1944:108, fig. 1 (wing). Type-locality: Venezuela, Maracay, reared from bromeliads. Distr. — Honduras to Colombia, Ve-

- nezucla, Trinidad, West Indies (Grenada). Ref. — Wirth & Blanton, 1968:205, fig. 2 (antenna, palpus, wing, legs, spermathecae, male genitalia, pupa) (redescription, distribution, synonymy).
- species* 3, Fox, 1942:418, figs. 19, 24 (pupa).
- rozeboomi* Barbosa, 1947:26, figs. (palpus, thorax, wing, male genitalia).
Type-locality: Trinidad.
- HYLAS** Macfie, 1940a:26, fig. 2 (wing). Type-locality: Guyana, New River. Distr. — Mexico to Brazil and Peru — Refs. — Forattini, 1957:243, figs. 26-27 (wing, thorax, palpus, antenna, eye separation, male genitalia); Wirth & Blanton, 1959:276, fig. 5 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Wirth & Blanton, 1968:203, fig. 1 (antenna, palpus, wing, legs, eye separation, spermathecae, male genitalia).
- IGNACIOI** Forattini, 1957:215, fig. 17 (wing, thorax, eye separation, antenna, palpus, male genitalia). Type-locality: Brazil, São Paulo, Boracéia, Saleópolis. Distr. — Brazil.
- INSIGNIS** Lutz, 1913:51, plate 7, fig. 3 (wing). Type-locality: Brazil, Rio de Janeiro and Bahia. Distr. — U.S.A. (Florida), Mexico to Brazil, Argentina (Corrientes), West Indies. Refs. — Lima, 1937a:415, fig. 3 (palpus) (notes on Lutz collection); Wirth & Blanton, 1956f:319, fig. 6 (wing, palpus, spermathecae, male genitalia) (synonymy); Forattini, Rabello & Pattoli, 1956:195, figs. 1-7 (larva, pupa) (immature stages, Brazil); Forattini, 1957: 223, figs. 20-22 (wing, thorax, palpus, eye separation, antenna, male genitalia, larva, pupa); Wirth & Blanton, 1959:285, fig. 9 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Linley, 1965b:57, figs. (pupa) (Jamaica).
- inamollae* Fox & Hoffman, 1944:110, fig. 2 (wing). Type-locality: Puerto Rico, Mayaguez.
- painteri* Fox, 1946:257, fig. 10 (wing). Type-locality: Honduras, Puerto Castilla.
- LUTZI** Lima, 1937a:419, fig. 5 b (palpus). Type-locality: Brazil, Pará, Abaeté. Distr. — Guyana, Brazil. Refs. — Wirth & Blanton, 1956f:318, fig. 8 (wing); Forattini, 1957:237, fig. 25 (wing, thorax, palpus, eye separation, antenna, palpus, male genitalia).
- MARUIM** Lutz, 1913:48, plate 7, fig. 1 (wing), plate 8, figs. 14, 16, 19 (egg, pupa, palpus). Type-locality: Brazil, Rio de Janeiro, São Paulo, Bahia, Trinidad, mangrove zone. Distr. — Trinidad, Venezuela, French Guiana, Brazil. Refs. — Forattini & Galvão, 1955:225, figs. 1-3 (wing, palpus, antenna, thorax, male genitalia) (redescribed from types); Wirth & Blanton, 1956f:323, fig. 4 (wing, palpus, spermathecae, male genitalia) (syn.: *recifei*).
- recifei* Barbosa, 1947:25. Type-locality: Brazil, Pernambuco, Recife.
- lutzi* Lima: Floch & Abonnenc, 1942a:2, fig. 2b (wing, palpus) (French Guiana); Floch & Abonnenc, 1942b:2, fig. 1c (wing, palpus, male genitalia).
- insignis* Lutz: Barbosa, 1944:259, figs. 1-4 (wing, palpus, male genitalia) (Brazil).
- OCUMARENSIS** Ortiz, 1950c:455, plates 3, 4 (antenna, palpus, spermathecae, thorax, wing, male genitalia). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Venezuela, Ecuador. Ref. — Ortiz & León, 1954:571 (redescribed, Ecuador).



- PALPALIS Macfie, 1948:78, fig. 3 (wing, palpus). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico to Peru and Venezuela. Ref. — Wirth & Blanton, 1968:207, fig. 3 (wing, antenna, palpus, legs, spermathecae, male genitalia).
- contubernalis* Ortiz & León, 1954:574, figs. 24-27 (wing, antenna, palpus, spermathecae) (*rozeboomi* var.) Type-locality: Ecuador, San Miguel de los Colorados.
- POLYPORI Wirth & Blanton, 1968a:212, fig. 6 (wing, antenna, palpus, legs, spermathecae, male genitalia). Type-locality: Panama, C. Z., Barro Colorado I., from bracket fungus. Distr. — Honduras to Colombia and Brazil.
- luteovenus* Root & Hoffman; Barbosa, 1947:21 (Panama, from bracket fungus).
- PSEUDODIABOLICUS Fox, 1946:256, fig. 1 (wing). Type-locality: Trinidad, Camuto Village. Distr. — Mexico to Colombia and Brazil, Trinidad.
- RUIZI Forattini, 1954b:189, figs. 1-2 (wing, thorax, palpus, male genitalia). Type-locality: Brazil, Goiás, Aruanã. Distr. — Brazil. Refs. — Wirth & Blanton, 1956f:310, fig. 10 (wing, spermathecae, palpus); Forattini, 1957:201, fig. 13 (wing, eye separation, antenna, palpus, thorax).
- TRAVASSOSI Forattini, 1957:198, fig. 12 (eye separation, thorax, wing, palpus, antenna, male genitalia). Type-locality: Brazil, Pará, Serra do Cachimbo. Distr. — Brazil (Pará, Mato Grosso).
- TRINIDADENSIS Hoffman, 1925:286, plate 2, fig. 6 (wing). Type-locality: Trinidad, Port of Spain, Caronia River. Distr. — Nicaragua, Panama, Trinidad, West Indies (Bahamas, Cuba). Refs. — Wirth & Blanton, 1956f:324, fig. 5 (wing, palpus, spermathecae, male genitalia) (synonyms: *diminutus*, *oliveri*); Forattini, 1957:231, fig. 23 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:287, fig. 10 (wing, thorax, tibial comb, palpus, spermathecae, male genitalia).
- oliveri* Fox & Hoffman, 1944:108, fig. 5 (male genitalia). Type-locality: Haiti, Mariani.
- wokei* Barbosa, 1947 (Nov.): 28, fig. (palpus, male genitalia) (preoccupied by *wokei* Fox 1947 (June). Type-locality: Panama, Canal Zone, Balboa (lectotype selected, Wirth & Blanton, 1956e:325).
- diminutus* Barbosa, 1951:163 (nom. nov. for *wokei* Barbosa, not Fox).
- VERECUNDUS Macfie, 1948:76, fig. 2 (wing). Type-locality: Mexico, Chiapas, El Vergel. Distr. — Mexico to Ecuador. Ref. — Wirth & Blanton, 1968:209, fig. 4 (antenna, wing, palpus, legs, spermathecae, male genitalia).
- XANIFER Wirth & Blanton, 1968a:210, fig. 5 (wing, antenna, palpus, legs, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Honduras, Costa Rica, Panama.

Subgenus MACFIELLA Fox

- Culicoides*, subg. MACFIELLA Fox, 1955:217. Type-species, *Ceratopogon phlebotomus* Williston (orig. des.).
- PHLEBOTOMUS (Williston), 1896:281, plate 9, fig. 46 (wing, palpus) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — Mexico to Ecuador and Brazil, West Indies. Refs. — Painter, 1927:258, figs. 3, 8 (larva, pupa) (biology); Fox, 1942:419, fig. 1 (pupa); Wirth & Blanton, 1953c:114, figs. 1-2 (wing, male genitalia); (synonym: *amazonius*; distribution); Forattini, 1957:455, figs. 110-111 (pupa, wing, antenna, palpus, thorax,

eye separation, male genitalia); Wirth & Blanton, 1959:469, fig. 90 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

amazonius Macfie, 1935a:52, fig. 3 (male genitalia). Type-locality: Brazil, Piauí, Tutóia.

WILLISTONI Wirth & Blanton, 1953c:116, figs. 3-6 (wing, tarsus, palpus, male genitalia). Type-locality: Panama, Coclé, Rio Hato. Distr. — Panama. Refs. — Forattini, 1957:460, fig. 112 (wing, thorax, palpus, antenna, eye separation, male genitalia); Wirth & Blanton, 1959:471, fig. 91 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

Subgenus MONOCULICOIDES Khalaf

Culicoides, subg. MONOCULICOIDES Khalaf, 1954:39. Type-species, *Ceratopogon nubeculosus* Meigen (orig. des.).

VARIIPENNIS (Coquillett), 1901a:602 (*Ceratopogon*). Type-locality: U.S.A., Virginia, Richmond.

ssp. OCCIDENTALIS Wirth & Jones, 1957: 21. Type-locality: U.S.A., California, Lake Co., Borax Lake. Distr. — North America (British Columbia to California and Arizona), Mexico (Baja California).

ssp. SONORENSIS Wirth & Jones, 1957:18. Type-locality: U.S.A., Arizona, Cochise Co., St. David. Distr. — U.S.A. (California to Washington and Tennessee), Mexico (Guerrero, D.F., Nuevo León, Puebla, Sonora).

Subgenus OECTACTA Poey

OECTACTA Poey, 1851:238 (as genus). Type-species, *fulvens* Poey (mon.).

Psychophaena Philippi, 1865:628. Type-species, *pictipennis* Philippi (mon.) = *venezuelensis* Ortiz & Mirsa.

Haematomyidum Goeldi, 1905:137. Type-species, *paraensis* Goeldi (orig. des.).

Cotocripus Brèthes, 1912:451. Type-species, *caridei* Brèthes (mon.).


Culicoides, subg. *Mataemyia* Vargas, 1960:43. Type-species, *Culicoides mojingaensis* Wirth & Blanton (orig. des.).

ACOTYLUS Lutz, 1913:69, plate 8, figs. 11, 21 (wing, palpus). Type-locality: Brazil, Mato Grosso, Rio Tapajós, Salto Augusto. Distr. — Mexico, Panama, Colombia, Venezuela, Brazil (Pará, Mato Grosso). Refs. — Ortiz & Mirsa, 1952a:257, figs. 9, 15, 16 (palpus, wing, spermathecae) (synonym: *panamericanus*); Forattini, 1957:445, fig. 107 (wing, thorax, eye separation, palpus, antenna, male genitalia); Wirth & Blanton, 1959:367, fig. 46 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

panamericanus Fox, 1947:30, figs. 3, 4 (wing, palpus). Type-locality: Mexico, D.F., San Jacinto.

ALAHIALINUS Barbosa, 1952:11, plate 5, fig. 2 (palpus), plate 7 (wing). Type-locality: Ecuador, Esmeraldas, Limones. Distr. — Ecuador, Panama. Refs. — Ortiz & León, 1954:571, figs. 508 (wing, antenna, palpus, spermathecae); Forattini, 1957:316, fig. 53 (wing, thorax, eye separation, antenna, palpus); Wirth & Blanton, 1959:398, fig. 59 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

ALMIRANTEI Wirth & Blanton, 1959:454, fig. 84 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.



- ANDINUS Wirth & Lee, 1967:17, figs. 5 c-g, plate 1, fig. 7 (wing, palpus, male genitalia). Type-locality: Colombia, Cauca, El Tambo, Finca Carpen-
teria. Distr. — Colombia.
- ARCHBOLDI Wirth & Blanton, 1970c:39, fig. 1 (wing, palpus, antenna, eye sepa-
ration, leg, spermathecae, male genitalia). Type-locality: Dominica,
Clarke Hall. Distr. — West Indies (Dominica, Trinidad).
- AUREUS Ortiz, 1951b:585, figs. 37-42 (palpus, scutum, wing, male genitalia).
Type-locality: Venezuela, Yaraucuy, San Felipe. Distr. — Panama, Vene-
zuela, Trinidad. Refs. — Forattini, 1957:478, fig. 119 (wing, thorax,
palpus, antenna, eye separation, male genitalia) (synonym: *miyamotoi*);
Wirth & Blanton, 1959:447, fig. 81 (wing, scutum, palpus, tibial comb,
spermathecae, male genitalia).
- miyamotoi* Wirth & Blanton, 1953b:231, fig. 3 (wing, thorax, palpus, male
genitalia). Type-locality: Panama, C. Z., Loma Borracho.
- AVILAENSIS Ortiz & Mirsa, 1951:593, figs. 1-4, 26 (wing, palpus, antenna, thorax,
spermathecae). Type-locality: Venezuela, Miranda, Los Chorros. Distr.
— Venezuela. Ref. — Forattini, 1957:364, fig. 73 (wing, palpus, antenna).
- AZUREUS Wirth & Blanton, 1959:377, fig. 50 (wing, thorax, palpus, tibial comb,
spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro,
Almirante. Distr. — Panama.
- BALSAPAMBENSIS Ortiz & León, 1954:579, figs. 30-31 (wing, palpus) (*pifanoi* ssp.).
Type-locality: Ecuador, Bolivar, Balsapamba. Distr. — Ecuador, Panama.
Refs. — Forattini, 1957:333, fig. 60 (wing, palpus, eye separation);
Wirth & Blanton, 1959:414, fig. 66 (wing, thorax, palpus, tibial comb,
spermathecae, male genitalia).
- BAMBUSICOLA Lutz, 1913:62, plate 7, fig. 10 (wing). Type-locality: Brazil, Rio
de Janeiro, Petrópolis (designated by Forattini, 1957:321). Distr. —
Colombia, Brazil. Refs. — Barretto, 1944:95, fig. 12, plate 2, fig. 4
(wing, male antenna, palpus, genitalia); Forattini, 1957:321, figs. 55-56
(wing, thorax, antenna, palpus, eye separation, male genitalia, pupa)
(synonym: *bahiensis*; larvae in bamboo).
- bahiensis* Barbosa, 1947:11, plate 2, fig. 5 (palpus), plate 4, fig. 2 (male
genitalia). Type-locality: Brazil, Bahia, from bamboo.
- BARBOSAI Wirth & Blanton, 1956e:161, fig. 2 (wing, thorax, palpus, spermathecae,
tibial comb, male genitalia). Type-locality: Panama, C.Z., Mojinga
Swamp. Distr. — U.S.A. (Florida), Panama, Ecuador, West Indies
(Bahamas, Cuba, Jamaica). Refs. — Lewis, 1958:723, figs. 21-23 (female
internal reproductive tract) (habits, Jamaica); Linley, 1965b:57, figs.
(pupa); Wirth & Blanton, 1959:400, fig. 60 (wing, thorax, palpus, tibial
comb, spermathecae, male genitalia).
- BIRABENI Cavaliere, 1966:59, figs. 1-10 (wing, thorax, antenna, legs, tibial comb,
palpus, eye separation, spermathecae). Type-locality: Venezuela, Sucre,
Cariaguaito. Distr. — Venezuela.
- CAMPOSI Ortiz & León, 1954:580, figs. 34-37 (wing, thorax, palpus, antenna).
Type-locality: Ecuador, Pichincha, Zapallo Grande, San Miguel de los
Colorados. Distr. — Panama, Ecuador. Ref. — Wirth & Blanton,
1959:356, fig. 41 (wing, thorax, palpus, tibial comb, spermathecae, male
genitalia) (synonym: *fairchildi*).
- fairchildi* Wirth & Blanton, 1955a:102, fig. 2 (wing, palpus, spermathecae,
male genitalia). Type-locality: Panama, C.Z., Madden Dam.

- CANCER Hogue & Wirth, 1968:2, figs. 1-13 (all stages). Type-locality: Costa Rica, Puntarenas, Golfo de Nicoya, Boca de Barranca, from crab holes. Distr. — Costa Rica.
- CAPRILESI Fox, 1952:364, fig. 1 (wing, palpus, eye separation, tibial comb, spermathecae, antenna). Type-locality: Venezuela, Mt. Marahuaca. Distr. — Panama, Venezuela. Refs. — Forattini, 1957:315, fig. 52 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:456, fig. 85 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonym: *kintzi*).
- kintzi* Wirth & Blanton, 1953a:72, fig. 2 (wing, palpus, male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp.
- pachymerus* Lutz; Ortiz, 1958:39, figs. 1-5 (legs, antenna, palpus, spermatheca) (Venezuela).
- CARIDEI (Brèthes), 1912:451, fig. (female) (*Cotocripus*). Type-locality: Argentina, Buenos Aires, Berazategui. Distr. — Brazil, Uruguay, Argentina. Refs. — Lane, 1945:366, figs. 14-15 (wing, antenna) (synonym: *setifer*); Wirth, 1960b:44 (redescribed; synonymy).
- setifer* Lutz, 1913:64, plate 8, fig. 13 (wing) (*Centrorhynchus*). Type-locality: Brazil, Rio Grande do Sul, Santa Vitória do Palmar.
- CARPENTERI Wirth & Blanton, 1953a:72, fig. 3 (wing, thorax, palpus, male genitalia). Type-locality: Panama, C.Z., Madden Dam. Distr. — Panama. Ref. — Wirth & Blanton, 1959:358, fig. 42 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- CARSOMELAS Wirth & Blanton, 1955a:100, fig. 1 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama. Ref. — Forattini, 1957:404, fig. 89 (wing, palpus, male genitalia); Wirth & Blanton, 1959:365, fig. 45 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- CASTILLAE Fox, 1946:251, figs. 2-3 (wing, thorax). Type-locality: Honduras, Puerta Castilla. Distr. — Guatemala to Panama, Ecuador, Venezuela. Refs. — Forattini, 1957:499, fig. 127 (wing, palpus, male genitalia); Wirth & Blanton, 1959:416, fig. 67 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonym: *gibsoni*).
- gibsoni* Wirth, 1952d:246, fig. 1 d-f (wing, spermatheca, palpus). Type-locality: Guatemala, Chimaltenango, San Pedro Yepocapa.
- CAUCAENSIS Wirth & Lee, 1967:19, figs. 6 a-d, plate 1, fig. 8 (wing, antenna, eye separation, palpus, spermathecae). Type-locality: Colombia, Cauca, Páramo Purace. Distr. — Colombia.
- CRUCIFER Clastrier, 1968:85, fig. 1 (wing, head, antenna, palpus, tibial comb, male and female genitalia). Type-locality: French Guiana, Cayenne, reared from tree hole. Distr. — Trinidad, French Guiana, Guyana.
- DARLINGTONAE Wirth & Blanton, 1971:39, fig. 2 (antenna, palpus, wing, eye separation, spermathecae, male genitalia). Type-locality: Trinidad, Tamana Caves. Distr. — Trinidad.
- DASYOPHRUS Macfie, 1940a:27. Type-locality: Guyana, New River. Distr. — Venezuela, Guyana, Brazil. Refs. — Wirth & Blanton, 1956d:186, fig. 1 (wing, palpus, spermathecae, tibial comb, male genitalia); Forattini, 1957:503, fig. 130 (wing, palpus, male genitalia).
- DEBILIPALPIS Lutz, 1913:60, plate 7, fig. 8 (wing). Type-locality: Brazil, São Paulo, Serra da Bocaina. Distr. — U.S.A. (Maryland to Florida), Panama to Argentina, Trinidad. Refs. — Macfie, 1937:7 (redescribed; Trinidad); Floch & Abonnenc, 1942a:3, fig. 2 d (wing, palpus); Wirth &

Blanton, 1959:442, fig. 79 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Forattini, 1957:383, fig. 81 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1971:34 (redescribed).

germanus Macfie; Forattini, 1957:381 (Argentina records).

- DENISAE Clastrier, 1971:290, fig. 2 a-e (wing, eye separation, palpus, spermathecae, male genitalia). Type-locality: French Guiana, Forêt du Gallion, reared from moist soil in depression in fallen tree trunk. Distr. — French Guiana.
- DICROURUS Wirth & Blanton, 1955b:123, fig. 2 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, C.Z., Loma Borracho. Distr. — Panama. Refs. — Forattini, 1957:354, fig. 68 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:386, fig. 54 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- DISCREPANS Ortiz & Mirsa, 1951:595, figs. 5-6, 13, 27 (wing, spermathecae, palpus, antenna). Type-locality: Venezuela, Miranda, Los Chorros. Distr. — Venezuela. Ref. — Forattini, 1957:353, fig. 67 (wing, thorax, palpus, antenna).
- DONAJII Vargas, 1954:28, fig. B, 1-5 (wing, antenna, palpus, male genitalia). Type-locality: Mexico, Oaxaca. Distr. — Mexico. Ref. — Forattini, 1957:375, fig. 77 (wing, thorax, palpus, antenna, male genitalia).
- EADSI Wirth & Blanton, 1971:34, fig. 1 (male, female parts). Type-locality: U.S.A., Texas, Cameron Co., reared from tree hole. Distr. — U.S.A. (Texas), Mexico (Nayarit, Sonora, San Luis Potosí, Yucatan).
- EQUATORIENSIS Barbosa, 1952:13, plates 3-5 (wing, palpus) (*debilipalpis* var.). Type-locality: Ecuador, Santo Domingo, Colorados. Distr. — Ecuador.
- EUBLEPHARUS Macfie, 1948:86. Type-locality: Guyana, New River. Distr. — Venezuela, Guyana. Ref. — Ortiz, 1952d:129, figs. 1-4 (wing, palpus, spermatheca, male genitalia).
- FERNANDEZI Ortiz, 1954:223, figs. 20-23 (wing, antenna, palpus, spermathecae). Type-locality: Venezuela, Miranda, Los Chorros. Distr. — Venezuela. Ref. — Forattini, 1957:351, fig. 66 (wing, palpus, antenna).
- FIELDI Wirth & Blanton, 1956a:50, fig. 3 (wing, palpus, spermatheca, tibial comb, male genitalia). Type-locality: Honduras, Lancetilla. Distr. — Honduras, Panama. Ref. — Wirth & Blanton, 1959:427, fig. 72 (wing, palpus, tibial comb, spermathecae, male genitalia).
- FLOCHABONNENCI Ortiz & Mirsa, 1952a:267, figs. 19, 20, 25 (palpus, wing, spermatheca). Type-locality: Venezuela, Miranda, Los Chorros. Distr. — Venezuela, Ecuador. Ref. — Ortiz & León, 1954:574, figs. 9-12, 14, 29 (wing, antenna, palpus, spermatheca, male genitalia).
- lichyi* Floch & Abonnenc; Ortiz & Mirsa, 1951:601, figs. 14-15 (wing, spermatheca, palpus) (Venezuela).
- FLORENCIAE Messersmith, 1972:167, fig. 1 a-f (antenna, eye separation, palpus, tibial comb, spermathecae, wing). Type-locality: Colombia, Caquetá, Florencia, Rio Hacho. Distr. — Colombia.
- FLORIDENSIS Beck, 1951:135, figs. a-b (male genitalia). Type-locality: U.S.A., Florida, Sarasota Co., Englewood. Distr. — U.S.A. (Florida), West Indies (Bahamas), Bermuda.
- FLUVIALIS Macfie, 1940a:25, fig. 1 (wing). Type-locality: Guyana, New River. Distr. — Honduras, Panama, Venezuela, Guyana. Refs. — Ortiz, 1951c:2,

plate 1 (wing, palpus, spermathecae); Ortiz & Mirsa, 1952a:264, figs. 8, 12-14 (palpus, antenna, wing, male genitalia); Wirth & Blanton, 1959:407, fig. 63 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

FORATTINI Ortiz, 1961:211, figs. 1-4 (antenna, palpus, tibial comb, spermathecae), photos 1-3 (wing, thorax, spermathecae). Type-locality: Venezuela, Guarico, Altigracia de Orituco. Distr. — Venezuela.

FURENS (Poey), 1851:238 (*Oecacta*). Type-locality: Cuba. Distr. — U.S.A. (coastal, Massachusetts to Texas), Mexico and Central America (both coasts), West Indies, south to Ecuador and Brazil (coastal). Refs. — Bequaert, 1924:197 (habits; Honduras); Painter, 1927:256, figs. 2, 5 (pupa) (biology, Honduras); Hoffman, 1925:287, plate 1, fig. 2 (thorax), plate 2, fig. 7 (wing) (distribution, description, synonym: *maculithorax*); Macfie, 1937:10, fig. (male genitalia) (Trinidad); Barretto, 1944:89, fig. 8 a-e, plate 1, fig. 1 (wing, antenna, palpus, male genitalia); Wirth, 1952c:95, figs. 11-18 (immature stages); Woke, 1954:61 (biology, Panama); Blanton, Graham, & Keenan, 1955:13 (biology and control, Panama); Lewis, 1958:724, figs. 1-20, 24 (anatomy of female reproductive tract) (biology; Jamaica); Wirth & Blanton, 1956e:159, fig. 1 (wing, thorax, tibial comb, spermathecae, male genitalia) (redescription); Forattini, 1957:448, figs. 108-109 (wing, thorax, eye separation, antenna, palpus, male genitalia, larva, pupa); Wirth & Blanton, 1959:404, figs. 3, 62 (female, wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

maculithorax Williston, 1896:277, plate 9, fig. 36 (wing) (*Ceratopogon*). Type-locality: St. Vincent.

GABALDONI Ortiz, 1954:221, figs. 16-19 (wing, antenna, palpus, spermathecae). Type-locality: Venezuela, Yaracuy, San Felipe. Distr. — Guatemala to Panama, Venezuela. Ref. — Wirth & Blanton, 1959:431, fig. 74 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

germanus Macfie; Wirth, 1955a:111, fig. 1 (male genitalia) (Guatemala).

GALINDOI Wirth & Blanton, 1953a:73, fig. 4 (wing, thorax, spermathecae, palpus, male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama. Refs. — Forattini, 1957:361, fig. 71 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:349, fig. 38 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

GERMANUS Macfie, 1940a:27. Type-locality: Guyana, New River. Distr. — Guyana. Ref. — Wirth & Blanton, 1956d:188, fig. 2 (wing, spermathecae, palpus, tibial comb) (type redescribed).

GINESI Ortiz, 1951b:586, figs. 26-28 (palpus, antenna, wing, spermathecae). Type-locality: Venezuela, Yaracuy, San Felipe. Distr. — Panama, Venezuela. Refs. — Forattini, 1957:395, fig. 85 (wing, thorax, palpus, antenna, eye separation, male genitalia); Wirth & Blanton, 1959:450, fig. 82 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

GLABELLUS Wirth & Blanton, 1956a:47, fig. 1 (wing, thorax, palpus, spermathecae, tibial comb, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Honduras, Nicaragua, Panama, Brazil, Trinidad. Ref. — Wirth & Blanton, 1959:429, fig. 73 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

GLABRIOR Macfie, 1940a:27 (*debilipalpis* var.). Type-locality: Guyana, New River. Distr. — Honduras to Colombia and Brazil. Ref. — Wirth & Blanton, 1959:452, fig. 83 (wing, palpus, tibial comb, spermathecae, male genitalia) (synonym: *grahambelli*).

- grahambelli* Forattini, 1956a:35, figs. 1-3 (wing, eyes, palpus). Type-locality: Panama, Cerro Cefa.
- GORGASI Wirth & Blanton, 1953b:232, fig. 4 (wing, thorax, palpus, male genitalia). Type-locality: Panama, Los Santos, Las Tablas. Distr. — Panama. Refs. — Forattini, 1957:370, fig. 75 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:402, fig. 61 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- GUADELOUPENSIS Floch & Abonnenc, 1950b:2, fig. 1 a-g (wing, palpus, eye separation, head, spermatheca, legs, buccal armature). Type-locality: Guadeloupe, St. Claude. Distr. — West Indies (Guedeloupe).
- GUERRAI Wirth & Blanton, 1971:41, fig. 3 (antenna, palpus, eye separation; wing, spermatheca, male genitalia). Type-locality: Trinidad, Tucker Valley. Distr. — Trinidad.
- GUYANENSIS Floch & Abonnenc, 1942a: 5, fig. 3 (wing, antenna, palpus, leg, male genitalia). Type-locality: French Guiana, Cayenne. Distr. — Panama, French Guiana, Brazil, Trinidad. Refs. — Forattini & Rabello, 1956:43, figs. 1, 2 (larva, pupa); Forattini, 1957:466, figs. 114-116 (larva, pupa, wing, thorax, palpus, antenna, eye separation, male genitalia); Wirth & Blanton, 1959:370, fig. 47 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonyms: *recifensis*, *stubalensis*).
- recifensis* Barbosa, 1943:263, figs. 6, 8 (palpus, wing). Type-locality: Brazil, Recife, Estrada dos Remédios e Boa Viagem.
- stubalensis* Fox, 1946:254, figs. 7-8 (wing, thorax). Type-locality: Trinidad, Stubal's Bay.
- stellifer* (Coquillett); Macfie, 1937:12 (Trinidad record).
- HOFFMANI Fox, 1946:251, figs. 6, 9 (wing, thorax). Type-locality: Trinidad, Camuto Village. Distr. — Panama, West Indies, Jamaica, Puerto Rico, Virgin Islands, Trinidad. Refs. — Fox, 1949:29, fig. 1 (palpus, spermathecae, male genitalia) (Puerto Rico); Wirth & Blanton, 1959:436, fig. 76 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Linley & Kettle, 1964:129, figs. (larva, pupa) (Jamaica); Wirth & Blanton, 1971:36 (redescribed).
- HORTICOLA Lutz, 1913:61, plate 7, fig. 9 (wing). Type-locality: Brazil, São Paulo, Tatuí, and Formosa, Serra da Bocaina, on horses. Distr. — Venezuela, French Guiana, Brazil. Refs. — Floch & Abonnenc, 1942a:4, fig. 2 e (wing, palpus); Forattini, 1957:318, fig. 54 (wing, thorax, eye separation, antenna, palpus, male genitalia, from types).
- IMITADOR Ortiz, 1953b:807, figs. 1-20 (wing, palpus, eye separation, antenna, spermathecae, male genitalia). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Panama, Venezuela. Ref. — Wirth & Blanton, 1959:438, fig. 77 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- INSINUATUS Ortiz & León, 1954:577, figs. 15-19 (wing, antenna, palpus, eye separation, spermathecae). Type-locality: Ecuador, Napo-Pastaza, Nucuvo Rocafuerte. Distr. — Ecuador, Brazil. Ref. — Forattini, 1957:373, fig. 76 (wing, thorax, palpus, antenna, eye separation).
- LANEI Ortiz, 1950b:431, fig. (wing, palpus, thorax, male genitalia). Type-locality: Panama, Cerro Zefa. Distr. — Honduras, Panama, Venezuela, Brazil, Trinidad. Refs. — Forattini, 1957:472, fig. 117 (wing, thorax, eye separation, palpus, male genitalia); Wirth & Blanton, 1959:374, fig. 49 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

- LEONI Barbosa, 1952:17, figs. (palpus, wing). Type-locality: Ecuador, Santo Domingo. Distr. — Ecuador. Refs. — Wirth & Blanton, 1956a:46, fig. 2 (wing, palpus, spermathecae, tibial comb, male genitalia); Forattini, 1957:488, fig. 123 (wing, palpus, antenna, eye separation).
- LEOPOLDOI Ortiz, 1951b:579, figs. 16-20 (wing, palpus, antenna, thorax). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Panama, Venezuela, Brazil. Refs. — Ortiz & Mirsa, 1951:596, figs. 7, 22, 28 (wing, palpus, male genitalia); Forattini, 1957:490, fig. 124 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:418, fig. 68 (wing, thorax, palpus, tibial comb, spermatheca, male genitalia).
- LICHYI Floch & Abonnenc, 1949:1, fig. 1 (wing, palpus). Type-locality: Venezuela, Río Borburata. Distr. — Venezuela. Ref. — Floch & Abonnenc, 1950:69 (again as n. sp.).
- LIMAI Barretto, 1944:99, fig. 14, plate 3, fig. 6 (wing, antenna, palpus, male genitalia). Type-locality: Brazil, São Paulo, Moji das Cruzes. Distr. — Honduras, Panama, Colombia to Ecuador and Brazil, Trinidad. Refs. — Forattini, 1957:335, fig. 61 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:352, fig. 39 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- LIMONENSIS Ortiz & León, 1954:576, figs. 20-23 (wing, antenna, palpus, spermathecae). Type-locality: Ecuador, Esmeraldas, Limones. Distr. — Ecuador, Venezuela, Brazil. Ref. — Forattini, 1957:378, fig. 78 (wing, eye separation, thorax, antenna, palpus).
- LOPESI Barretto, 1944:102, fig. 15, plate 4, fig. 7 (wing, male antenna, palpus, male genitalia). Type-locality: Brazil, São Paulo, Moji das Cruzes. Distr. — Brazil, Argentina. Ref. — Forattini, 1954d:223, figs. 1-2 (wing, eye separation, palpus, thorax, male genitalia); Forattini, 1957:325, fig. 58 (wing, thorax, eye separation, antenna, palpus, male genitalia).
- LUGLANI Jones & Wirth, 1958:89, figs. 1-6 (wing, palpus, thorax, spermathecae, male genitalia). Type-locality: U. S. A., Texas, Kerr County, USDA Exp. Sta. Distr. — U. S. A. (Arizona to Texas), Mexico (Baja California, Sonora, Tamaulipas).
- LYRINOTATUS Wirth & Blanton, 1955b:126, fig. 3 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Nicaragua, Panama. Refs. — Forattini, 1957:435, fig. 103 (wing, palpus, male genitalia); Wirth & Blanton, 1959:391, fig. 56 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- MACROSTIGMA Wirth & Blanton, 1953b:230, fig. 2 (wing, palpus, male genitalia). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama. Refs. — Forattini, 1957:311, fig. 51 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:395, fig. 58 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- MAGNIPALPIS Wirth & Blanton, 1953:76, fig. 6 (wing, palpus, spermathecae, male genitalia). Type-locality: Panama, Cerro Campana. Distr. — Panama. Refs. — Forattini, 1957:366, fig. 72 (wing, palpus, male genitalia); Wirth & Blanton, 1959:360, fig. 43 (wing, palpus, tibial comb, spermathecae, male genitalia).
- MARTINEZI Wirth & Blanton, 1970c:43, fig. 3 (wing, antenna, palpus, eye separation, spermathecae, male genitalia). Type-locality: Trinidad, Macqueripe. Distr. — Trinidad.



- MELLEUS (Coquillett), 1901a:604 (*Ceratopogon*). Type-locality: U. S. A., Florida, Lake Worth. Distr. — U.S.A. (Atlantic and Gulf Coasts; intertidal) West Indies (Bahamas).
- MIRSÆ Ortiz, 1953a:781, figs. 1-2 (wing). Type-locality: Venezuela, Lara, Duaca. Distr. — Panama, Venezuela, Trinidad. Refs. — Forattini, 1957:497, fig. 126 (wing); Wirth & Blanton, 1959:446, fig. 80 (wing, thorax, palpus, tibial comb, spermathecae).
- MOHAVE Wirth, 1952c:187, fig. 19 (wing, palpus, male genitalia). Type-locality: U.S.A., California, San Bernardino Co., Vidal. Distr. — U. S. A. (California, Arizona), Mexico.
- MOJINCAENSIS Wirth & Blanton, 1953b:232, fig. 5 (wing, scutum, palpus, male genitalia). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama. Ref. — Wirth & Blanton, 1959:379, fig. 51 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- MONTICOLA Wirth & Lee, 1967:15, figs. 5 a-d, plate 1, fig. 6 (wing, antenna, eye separation, spermathecae, palpus). Type-locality: Colombia, Cauca, El Tambo, Finca Carpentería. Distr. — Colombia.
- NEGHMEI Vargas, 1955:673, fig. (wing, palpus, male genitalia). Type-locality: Mexico, Puebla, Atlixco. Distr. — Mexico.
- OENOXIUS Fox, 1952:365, fig. 2 (wing eye separation, antenna, tibial comb.) Type-locality: Venezuela, Mount Marahuaca. Distr. — Venezuela. Ref. — Forattini, 1957:303, fig. k (wing, antenna, palpus, eye separation).
- PACHYMERUS Lutz, 1914:83, plate 8, fig. 8 plate 9, fig. 1 (wing, leg.). Type-locality: Brazil, Rio Negro, Manaus. Distr. — Guatemala, Panama, Brazil. Ref. — Wirth & Blanton, 1959:459, fig. 86 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- PAMPOIKILUS Macfie, 1948:79, fig. 4 (wing). Type-locality: Mexico, Chiapas, El Vergel. Distr. — Mexico, Panama, Venezuela. Refs. — Forattini, 1957:398, fig. 86 (wing, palpus, antenna, male genitalia) (synonym: *dominicii*); Wirth & Blanton, 1959:324, fig. 27 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- dominicii* Ortiz, 1951c:7, plate 1 (wing, palpus, male genitalia). Type-locality: Venezuela, Miranda, Ocumare del Tuy.
- PARAENSIS (Goeldi), 1905:137 (*Haematomyidium*). Type-locality: Brazil, Pará, Belém. Distr. — U. S. A. (Pennsylvania to Mississippi and Florida), Central and South America to Argentina, West Indies. Refs. — Barretto, 1944:92, fig. 10, plate 2, fig. 3 (wing, male genitalia); Romaña & Wygodzinsky, 1950:29 (transmission of *Mansonella ozzardi* filariasis in Argentina); Forattini, 1957:426, fig. 100 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:440, fig. 78 (wing, thorax, tibial comb, palpus, spermathecae, male genitalia).
- lahillei* Ichès, 1906:264, figs. 1-3 (antenna, palpus, abdomen) (*Ceratopogon*). Type-locality: Argentina, Chaco Austral.
- undecimpunctatus* Kieffer, 1917:307. Type-locality: Argentina, San Pablo.
- PATULIPALPIS Wirth & Blanton, 1959:421, fig. 69 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Cerro Campana. Distr. — Panama.
- PAUCIENFUSCATUS Barbosa, 1947:23, plate 1, fig. 3, plate 9, fig. 6 (palpus, thorax). Type-locality: Brazil, Amazonas, Manaus. Distr. — Panama; Colombia, Brazil, Trinidad. Refs. — Forattini, 1957:345, fig. 64 (wing,


- thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:381, fig. 52 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- PHAEONOTUS** Wirth & Blanton, 1959:326, fig. 28 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- PIFANOI** Ortiz, 1951b:588, figs. 43-47 (wing, palpus, thorax, spermathecae). Type-locality: Venezuela, Maracay, San Felipe. Distr. — Honduras to Panama, Venezuela, Trinidad. Refs. — Forattini, 1957:342, fig. 63 (wing, palpus, male genitalia); Wirth & Blanton, 1959:384, fig. 53 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonym: *tricoloratus*).
- tricoloratus* Wirth & Blanton, 1953b:233, fig. 6 (wing, thorax, palpus, male genitalia). Type-locality: Panama, Panama Prov., Pacora.
- PROPINQUUS** Macfie, 1948:81, fig. 6 (wing, male genitalia). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico. Ref. — Forattini, 1957:410, fig. 92 (wing, male genitalia).
- PROPRIIPENNIS** Mcfie, 1948:84, fig. 9 (wing). Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico to Panama, Venezuela. Refs. — Forattini, 1957:442, fig. 106 (wing, thorax, eye separation, palpus, male genitalia); Wirth & Blanton, 1959:412, fig. 65 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- PULCHRIPENNIS** Macfie, 1939a:200, fig. 19 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Forattini, 1957:496.
- QUASIPARAENSIS** Clastrier, 1971:286, fig. 1 a-f (wing, eye separation, antenna, palpus, spermathecae, male genitalia). Type-locality: French Guiana, Cayenne, Cabassou Forest, reared from palm stipe. Distr. — French Guiana.
- RANGELI** Ortiz & Mirsa, 1952:126, plate 1, figs. 1-4, plate 2, fig. 11 (palpus, antenna, wing, eye separation, spermathecae). Type-locality: Venezuela, Los Chorros. Distr. — Panama, Ecuador, Venezuela. Refs. — Forattini, 1957:502, fig. 128 (wing, antenna, palpus); Wirth & Blanton, 1959:423, fig. 70 (wing, palpus, tibial comb, spermathecae).
- RETICULATUS** Lutz, 1913:49, plate 7, fig. 2, plate 8, figs. 15, 17, 20 (wing, palpus, male, larva). Type-locality: Brazil, São Paulo: Santos, Rio de Janeiro and Bahia, larvae in crab holes. Distr. — Panama, Brazil. Refs. — Forattini, 1957:432, fig. 102 (wing, thorax, eye separation, palpus, antenna, male genitalia); Wirth & Blanton, 1959:393, fig. 57 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia); Forattini, Rabello & Pattoli, 1960:1, figs. 1, 2 (larva, pupa) (biology, from crab holes).
- RODRIGUEZI** Ortiz, 1968:67, figs. 1, 2 (wing, eye separation). Type-locality: Venezuela, Yaracuy, Yumare. Distr. — Venezuela.
- SPURIUS** Wirth & Blanton, 1959:433, fig. 75 (wing, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, C. Z., Fort Davis. Distr. — Guatemala, Honduras, Panama.
- TAMBOENSIS** Wirth & Lee, 1967:20, fig. 6 e-j, plate 1, figs. 9-10 (wing, antenna, eye separation, palpus, spermathecae, male genitalia). Type-locality: Colombia, Cauca, El Tambo, Finca Carpentaria. Distr. — Colombia.
- TENULOBUS** Wirth & Blanton, 1959:342, fig. 40 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.



- TETRATHYRIS Wirth & Blanton, 1959:409, fig. 64 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama, Brazil.
- TRANSFERRANS Ortiz, 1953a:801, figs. 3-7 (wing, antenna, palpus, spermatheca). Type-locality: Venezuela, Amazonas Terr., head of Rio Orinoco. Distr. — Panama, Ecuador, Venezuela. Ref. — Wirth & Blanton, 1959:424, fig. 71 (wing, thorax, palpus, tibial comb, spermatheca, male genitalia).
- eublepharus* Macfie; Ortiz, 1952d:129, figs. 1-4 (wing, palpus, spermatheca, male genitalia) (Venezuela); Forattini, 1957:493, fig. 125 (wing, thorax, antenna, palpus, eye separation, male genitalia).
- TRILINEATUS Fox, 1946:250, figs. 5, 11 (wing, thorax). Type-locality: Virgin Islands, St. Thomas, Red Hook. Distr. — West Indies (Puerto Rico, Virgin Islands, Dominica, Barbados). Refs. — Wirth & Blanton, 1956d:189, fig. 3 (wing, thorax, spermathecae, tibial comb, palpus); Forattini, 1957:389, fig. 83 (wing, thorax, eye separation, antenna, palpus, male genitalia).
- UNIRADIALIS Wirth & Blanton, 1953a:70, fig. 1 (wing, palpus, male genitalia). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama. Refs. — Forattini, 1957:302, fig. 48 (wing, palpus, male genitalia); Wirth & Blanton, 1959:462, fig. 87 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- VENEZUELENSIS Ortiz & Mirsa, 1950:137, figs. 1-8, 4 plates (wing, antenna, palpus, spermathecae, head, male genitalia). Type-locality: Venezuela, Caracas, Los Chorros. Distr. — Panama, Venezuela, Brazil, Chile, Argentina. Refs. — Wirth, 1955c:234 (synonym: *pictipennis*, *ortizi*); Forattini, 1957:463, fig. 113 (wing, thorax, antenna, palpus, eye separation, male genitalia); Wirth & Blanton, 1959:372, fig. 48 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- pictipennis* Philippi, 1865:628 (*Psychophaena*). Type-locality: Chile, Santiago, San Cristóbal. Preoccupied in *Culicoides* by *pictipennis* (Staeger) 1839).
- ortizi* Fox, 1952:366, fig. 5 (wing, palpus, antenna, tibial comb, spermathecae). Type-locality: Venezuela, Zulia, La Salina.
- WOKEI Fox, 1947 (April):90, figs. 1-2 (wing, palpus). Type-locality: Panama, C. Z., Balboa, West Bank. Distr. — Panama. Refs. — Forattini, 1957:357, fig. 70 (wing, thorax, eye separation, antenna, palpus, male genitalia); Wirth & Blanton, 1959:363, fig. 44 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia) (synonym: *aethionotus*).
- aethionotus* Wirth & Blanton, 1955b:121, fig. 1 (wing, palpus, male genitalia; spermathecae). Type-locality: Panama, Darien, Patino Point.

Subgenus SELFIA Khalaf

- Culicoidis*, subg. SELFIA Khalaf, 1954:38. Type-species, *Culicoides hieroglyphicus* Malloch (orig. des.).
- HIEROGLYPHICUS Malloch, 1915a:297. Type-locality: U. S. A., Arizona, Graham Mountains, Ash Creek. Distr. — U.S.A. (California, to South Dakota and Texas), Mexico (Sonora). Ref. — Atchley, 1970:242, figs. 4-7, 9, 116-148 (male, female, pupa, many parts) (biosystematics).



Unplaced species of *Culicoides*

- ALBOMACULA Root & Hoffman, 1937:164, figs. 17, 29 (wing, male genitalia). Type-locality: Mexico, D.F., San Jacinto. Distr. — Mexico.
- ALVAREZI Ortiz, 1957:161, figs. 1-4 (wing, antenna, palpus, spermathecae). Type-locality: Venezuela, Miranda, Los Teques, Parque El Encantado. Distr. — Venezuela.
- ANTEFURCATUS Wirth & Blanton, 1959:315, fig. 23 (wing, thorax, tibial comb, palpus, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- ANTUNESI Forattini, 1954c:315, figs. 1-2 (wing, thorax, palpus, male genitalia). Type-locality: Brazil, Goiás, Aruanã. Distr. — Brazil (Goiás, Mato Grosso, São Paulo). — Ref. — Forattini, 1957:485, fig. 122 (thorax, wing, eye separation, palpus, antenna, male genitalia).
- ARUBAE Fox & Hoffman, 1944:109, fig. 3 (wing). Type-locality: Aruba Island, West Indies, reared from crab holes. Distr. — U. S. A. (Texas), Mexico (Tamaulipas), Panama, Venezuela, Aruba Island; in crab holes. Refs. — Forattini, 1957:438, fig. 104 (wing, thorax, eye separation, antenna, palpus, male genitalia), 105 (pupa); Wirth & Blanton, 1959:464, fig. 88 (wing, spermathecae, thorax, palpus, tibial comb, male genitalia). sp. 1, Fox, 1942:420, figs. 2, 5 (pupa).
variipennis (Coquillett), of authors (Venezuela records).
- BENARROCHI Ortiz & Mirsa, 1952:126 (as *benarrochei*). Type-locality: Venezuela, Caraballeda, Dept. Vargas, D. F. Distr. — Venezuela, Brazil. Ref. — Forattini, 1957:506, fig. 131 (wing, eye separation, antenna, palpus).
- BREDINI Wirth & Blanton, 1970c:41, fig. 2 (wing, antenna, palpus, eye separation, leg, spermathecae, male genitalia). Type-locality: Dominica, Clarke Hall. Distr. — West Indies (Dominica).
- BRICENOI Ortiz, 1951a:445, figs. 1-5 (wing, palpus, male genitalia). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Venezuela, Ecuador. Ref. — Forattini, 1957:348, fig. 65 (wing, male genitalia).
- COCHISENSIS Wirth & Blanton, 1967:216, figs. 26-33 (wing, leg, palpus, eye separation, spermathecae, male genitalia). Type-locality: U.S.A., Arizona, Santa Cruz, Sycamore Canyon. Distr. — U.S.A. (Arizona, Utah), Mexico (Baja California).
- COMMATIS Wirth & Blanton, 1959:321, fig. 26 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- CRESCENTIS Wirth & Blanton, 1959:317, fig. 24 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Mexico (Chiapas), Nicaragua, Panama.
- DAEDALOIDES Wirth & Blanton, 1959:330, fig. 30 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- DAEDALUS Macfie, 1948:83, fig. 8 (wing, male genitalia). Type-locality: Mexico, Chiapas, El Vergel. Distr. — U.S.A. (Arizona), Mexico, Honduras, Panama, Venezuela. Refs. — Forattini, 1957:401, fig. 88 (wing, palpus, antenna, male genitalia); Wirth & Blanton, 1959:319, fig. 25 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).
- DAMPFI Root & Hoffman, 1937:169, figs. 23, 35 (wing, male genitalia). Type-locality: Mexico, D.F., San Jacinto. Distr. — Mexico.
- DAVIESI Wirth & Blanton, 1968b:251, figs. 1-7 (antenna, palpus, wing, leg, spermatheca, eye separation, thorax). Type-locality: Guyana, Rupununi Savannah, Ike Swamp. Distr. — Guyana.



- DUNNI Wirth & Blanton, 1959:328, fig. 29 (wing, thorax, palpus, tibial comb, spermathecae). Type-locality: Panama, Chiriquí, El Volcán. Distr. — Panama.
- FLUVIATILIS (Lutz), 1914:4, plate 8, fig. 7 (wing) (*Johannseniella*). Type-locality: Brazil, Rio Negro, S. Gabriel. Distr. — Brazil, Trinidad. Ref. — Lane, 1945:363, fig. 16 (antenna) (types redescribed; in *Lasiohelea*). *scorzai* Ortiz, 1956:93, figs. 1-4 (wing, palpus, spermathecae). Type-locality: Venezuela, Bolivar, Rio Guayaraca.
- pachymerus* Lutz; Forattini, 1957:305, fig. 49 (wing, eye separation, antenna, palpus, thorax) (Brazil records).
- OKLAHOMENSIS Khalaf, 1952:355, (*villosipennis* ssp.). Type-locality: U.S.A., Oklahoma, Wichita Refuge. Distr. — U.S.A. (Arizona to Texas), Mexico (Baja California, Sonora). Ref. — Wirth & Blanton, 1967:223, figs. 52-59 (wing, antenna, palpus, eye separation, leg, spermathecae, male genitalia).
- OUSAIRANI Khalaf, 1952:354, fig. 5 (antenna, palpus, male genitalia). Type-locality: U.S.A., Oklahoma, Wichita Refuge. Distr. — U.S.A. (Pennsylvania to Florida, Oklahoma and Texas), Mexico (Nuevo León). Ref. — Wirth & Blanton, 1967:225, figs. 60-67 (wing, antenna, palpus, eye separation, spermathecae, leg, male genitalia).
- POLYSTICTUS Kieffer, 1921c:181. Type-locality: Paraguay. Distr. — Paraguay.
- STIGMALIS Wirth, 1952d:245, fig. 1 a-c (wing, palpus, spermathecae). Type-locality: Guatemala, Chimaltenango, San Pedro Yepocapa. Distr. — Mexico, Guatemala, Panama. Refs. — Forattini, 1957:308, fig. 50 (wing, male genitalia); Wirth & Blanton, 1959:467, fig. 89 (wing, thorax, palpus, tibial comb, spermathecae, male genitalia).

Tribe CERATOPOGONINI

Genus ALLUAUDOMYIA Kieffer

- ALLUAUDOMYIA Kieffer, 1913:12. Type-species, *imparunguis* Kieffer (mon.). Six un-named new Neotropical species in the collection of the U. S. National Museum in Washington, D. C.

Genus BAEOHELEA Wirth & Blanton

- BAEOHELEA Wirth & Blanton, 1970b:95. Type-species, *nana* Wirth & Blanton (orig. des.).
- NANA Wirth & Blanton, 1970b:95, fig. 1 (antenna, head, palpus, wing, legs, tarsi, claws, spermatheca, male genitalia). Type-locality: Dominica, Pont Cassé. Distr. — Colombia, Ecuador, West Indies (Dominica).

Genus BRACHYPOGON Kieffer

- BRACHYPOGON Kieffer, 1899:69. Type-species, *Ceratopogon vitiosus* Winnertz (orig. des.). Ref. — Wirth & Blanton, 1970b:93 (revision).
- FUSCIVENOSUS (Lutz), 1914:94, plate 8, fig. 3 (wing) (*Palpomyia*). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — U.S.A. (Florida), Mexico, Panama, French Guiana, Brazil, West Indies (Jamaica, Virgin Islands). Refs. — Floch & Abonnenc, 1942c:4, fig. 1h, 2h (wing, palpus, male genitalia).



genitalia) (French Guiana); Lane, 1945:370, fig. 17 (male genitalia) (redescribed from types; in *Parabrezza*); Wirth & Blanton, 1970b:101, fig. 4 (antenna, palpus, wing, spermatheca, male genitalia).

IMPAR (Johansen), 1938:223 (*Ceratopogon*). Type-locality: Puerto Rico, El Yunque trail. Distr. — West Indies (Puerto Rico). Ref. — Johansen, 1943a:780, fig. 9 (wing).

PARAENSIS Wirth & Blanton, 1970b:99, fig. 3 (male, female head, antenna, palpus, legs, wing, tarsi, claws, spermatheca, male genitalia). Type-locality: Brazil, Pará, Rio Paru, Missão Tiriós. Distr. — Brazil.

Genus LEPTOHELEA Wirth & Blanton

LEPTOHELEA Wirth & Blanton, 1970a:12. Type-species, *micronyx* Wirth & Blanton (orig. des.).

MICRONYX Wirth & Blanton, 1970a:12, fig. 3 (antenna, palpus, wing, legs, tarsi, claws, spermatheca, tibial comb). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia.

Tribe STILOBEZZIINI

Genus ECHINOHELEA Macfie

ECHINOHELEA Macfie, 1940c:187. Type-species, *ornatipennis* Macfie (orig. des.).

MACFIEI Lane, 1948:228. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.

ORNATIPENNIS Macfie, 1940c:188, fig. 4 (male genitalia). Type-locality: Guyana, Mazaruni. Distr. — Mexico, Guyana. Ref. — Wirth, 1965a:249 (Mexico, pollinating *Hevea*).

RICHARDSI Macfie, 1940c:189. Type-locality Guyana, Mazaruni. Distr. — Guyana, Brazil. Ref. — Lane, 1948:228 (Brazil).

SMARTI Macfie, 1940c:190. Type-locality: Guyana, Mazaruni. Distr. — Guyana, Brazil. Ref. — Lane, 1948:227, fig. 4 (male genitalia) (Brazil).

Genus FITKAUHELEA Wirth & Blanton

FITKAUHELEA Wirth & Blanton, 1970a:7. Type-species, *amazonica* Wirth & Blanton (orig. des.).

AMAZONICA Wirth & Blanton, 1970a:9, fig. 1 (male, female antennae, palpus, female head, wing, legs, tarsi, claws, mandible, tibial comb, spermathecae, male genitalia). Type-locality: Brazil, Amazonas, Rio Maruiá. Distr. — Brazil.


Genus MACRUROHELEA Ingram & Macfie

MACRUROHELEA Ingram & Macfie, 1931:203. Type-species, *caudata* Ingram & Macfie (orig. des.).

Ref. — Wirth, 1965a:46 (revision, key).

CAUDATA Ingram & Macfie, 1931:205, figs. 27, 28a (abdomen, male genitalia). Type-locality: Chile, Casa Pangué. Distr. — Chile.

KUSCHELI Wirth, 1965a:49, figs. 1-2 (wing, antenna). Type-locality: Chile, Valparaíso, Algarrobo. Distr. — Chile.



SETOSA Wirth, 1965a:49, fig. 3 (male genitalia). Type-locality: Chile, Atacama, Vegas de San Andrés. Distr. — Chile.

THORACICA Ingram & Macfie, 1931:206, figs. 28b, 29 (wing, male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

Genus MONOHELEA Kieffer

MONOHELEA Kieffer, 1917:294. Type-species, *hieroglyphica* Kieffer (orig. des.). Refs. — Lane, 1948:224 (key); Lane & Wirth, 1964:209 (revision, key; Neotropical species); Wirth & Williams, 1964:302 (key, wing photos, figures North American species, including some Neotropical species).

Subgenus ISTHMOHELEA Ingram & Macfie

ISTHMOHELEA Ingram & Macfie, 1931:208 (as genus). Type-species, *disjuncta* Ingram & Macfie (orig. des.).

Ref. — Wirth & Blanton, 1972a:173 (diagnosis, subgeneric status).

DISJUNCTA (Ingram & Macfie), 1931:209, figs. 30-31 (wing, male genitalia) (*Isthmohelea*). Type-locality: Chile, Casa Pangué. Distr. — Chile.

SHANNONI Wirth & Blanton, 1972a:175, fig. 1 a-k (many parts, female). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

Subgenus MONOHELEA Kieffer

BALBOA Lane & Wirth, 1964:225, figs. 20, 29 (male genitalia). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama.

BLANTONI Lane & Wirth, 1964:217, figs. 10, 22 (male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.

BRASILIENSIS Lane, 1948:226, fig. 3 (wing). Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. — Brazil. Ref. — Lane & Wirth, 1964:229, fig. 37 (male genitalia).

CEBACOI Lane & Wirth, 1964:218, figs. 15, 24 (male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama, Brazil.

CHIAPASI Lane & Wirth, 1964:219, figs. 9, 23 (male genitalia). Type-locality: Nicaragua, Villa Somoza. Distr. — Nicaragua, Panama, Colombia.

CHIRUSI Lane & Wirth, 1964:218, figs. 8, 21 (male genitalia). Type-locality: Panama, Coclé, Puerto Obaldia. Distr. — Nicaragua, Panama.

COLOMBIAE Lane & Wirth, 1964:220, figs. 11, 25 (male genitalia). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia.

CUNASI Lane & Wirth, 1964:232, fig. 38 (male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.

FAIRCHILDII Lane & Wirth, 1964:214, figs. 5, 6 (male genitalia). Type-locality: Panama, Bocas del Toro, Rio Changuena. Distr. — Panama.

FUSCIPENNIS Lane & Wirth, 1964:221. Type-locality: Colombia, Rio Raposo. Distr. — Colombia.

GUAIMIESI Lane & Wirth, 1964:227, fig. 27 (male genitalia). Type-locality: Panama, Coclé, Anton. Distr. — Panama.



- GUIANAE Wirth, 1953b:150, fig. 19 k (male genitalia). Type-locality: Guyana, Mazaruni. Distr. — Trinidad, Guyana. Ref. — Lane & Wirth, 1964:224, fig. 16 (male genitalia).
- HIEROGLYPHICA Kieffer, 1917:312, fig. 13 (antenna, tarsi). Type-locality: Paraguay. Distr. — Guyana, Brazil, West Indies (Dominica, Trinidad). Refs. — Macfie, 1937:17, fig. 6 (wing, legs, claws) (Trinidad); Macfie, 1940a:30, fig. 4 (male genitalia) (Guyana); Lane & Wirth, 1964:231, fig. 36 (male genitalia).
hieroglyphica (error), Lane, 1948:226, fig. 2 (male genitalia).
- JOHANNSENI Wirth, 1953b:153, figs. 18g, k, l, 19a (wing, leg, spermathecae, male genitalia). Type-locality: U.S.A., Virginia, Falls Church. Distr. — U.S.A. (Michigan and Iowa to Maryland and Florida), Panama. Ref. — Lane & Wirth, 1964:214, figs. 1-4, 14 (wing, spermathecae, leg, male genitalia).
- LANEI Wirth, 1953b:142, figs. 18b, j, 19e (wing, leg, male genitalia). Type-locality: U.S.A., Florida, Miami. Distr. — U.S.A. (Florida), Costa Rica, West Indies (Bahamas). Ref. — Lane & Wirth, 1964:230, figs. 12, 33 (male genitalia).
- MACULIPENNIS (Coquillett), 1905:64 (*Ceratopogon*). Type-locality: U.S.A., Florida, Jacksonville. Distr. — U.S.A. (Texas to Georgia and Florida), Mexico, Guatemala, Panama. Ref. — Lane & Wirth, 1964:227, figs. 13, 30 (male genitalia).
- MAYERI Ortiz, 1950:202, plate 2 (wing, antenna, leg, tarsus, palpus, claws). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Costa Rica, Panama, Colombia, Venezuela, Trinidad. Ref. — Lane & Wirth, 1964:230, fig. 34 (male genitalia).
- MULTILINEATA (Lutz), 1914:93, plate 8, fig. 2, plate 9, figs. 6, 6a (wing, abdomen) (*Palpomyia*). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Panama, Guyana, Brazil, West Indies (Puerto Rico). Ref. — Lane, 1945:368, figs. 18-19 (male genitalia) (type lost, neotype selected from Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 42; redescribed); Lane & Wirth, 1964:224, figs. 19, 26 (male genitalia).
- OCUMARE Ortiz, 1952c:254, figs. 1-3 (wing, palpus, antenna). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Venezuela.
- ORNATA Wirth, 1953b:144, fig. 18d (male genitalia). Type-locality: U.S.A., Florida, Escambia Co., Santa Rosa Island. Distr. — U.S.A. (Florida), Panama. Ref. — Lane & Wirth, 1964:226, fig. 31 (male genitalia).
- PANAMENSIS Lane & Wirth, 1964:221, figs. 7, 17 (male genitalia). Type-locality: Panama, Darien, Patino Point. Distr. — Mexico (Guerrero), Panama, West Indies (St. John).
- PONCAI Lane & Wirth, 1964:228, fig. 35 (male genitalia). Type-locality: Panama, Panama Prov., Tocumén. Distr. — Costa Rica, Panama.
- STONEI Wirth, 1953b:148, figs. 18e, 19i (wing, male genitalia). Type-locality: U.S.A., Louisiana, Baton Rouge. Distr. — U.S.A. (Connecticut and Tennessee to Louisiana and Florida), Costa Rica, Panama, Colombia, Trinidad, West Indies (Cuba). Ref. — Lane & Wirth, 1964:222, figs. 18, 28 (male genitalia).
- URRAGAI SI Lane & Wirth, 1964:231, fig. 32 (male genitalia). Type-locality: Panama, C. Z., Fort Kobbe, Camaron. Distr. — Panama, Guyana, Trinidad.
hieroglyphica Kieffer; Macfie, 1937:18 (Trinidad); Macfie, 1940:30 (Guyana).
macfie Wirth (in part); Wirth, 1953b:143 (Trinidad and Guyana records).



Genus PARABEZZIA Malloch

- PARABEZZIA Malloch, 1915a:358. Type-species, *petiolata* Malloch (orig. des.)
Diaphanobezzia Ingram & Macfie, 1931:223. Type-species; *pellucida* Ingram & Macfie (orig. des.).
Refs. — Wirth, 1952b:22 (revision; synonym: *Diaphanobezzia*); Wirth, 1965b:215 (revision, key).
- BLANTONI Wirth, 1965b:220, fig. 21 (wing). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- BRUNNEA Wirth, 1965b:220, figs. 1, 22 (wing, tarsus). Type-locality: Panama, Panama Prov., Tocumén. Distr. — Panama.
- COSTALIS Wirth, 1965b:221, figs. 2, 3, 23 (wing, male genitalia). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.
- FUSCIPENNIS Wirth, 1965b:223. Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.
- JAMAICENSIS Wirth, 1965b:224, figs. 10, 26 (wing, tarsus). Type-locality: Jamaica, St. Thomas, Bath. Distr. — West Indies (Jamaica).
petiolata Malloch; Wirth, 1952b:23 (Jamaica records).
- PANAMENSIS Wirth, 1965b:225, figs. 11-12 (male genitalia). Type-locality: Panama, C.Z., Fort Kobbe, Camaron. Distr. — Panama.
- PELLUCIDA (Ingram & Macfie), 1931:223, figs. 33, 34b-g (wing, antenna, halter, male genitalia) (*Diaphanobezzia*). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina. Ref. — Wirth, 1952b:26 (combination).
- SPANGLERI Wirth, 1965b:227, fig. 28 (wing). Type-locality: Puerto Rico, Maricao Fish Hatchery. Distr. — West Indies (Puerto Rico).
- UNGUIS Wirth, 1965b:228. Type-locality: U.S.A., Arizona, Pima Co., Sabino Canyon. Distr. — U.S.A. (Arizona, Texas), Mexico (Sonora), Panama.

Genus PARASTILOBEZZIA Wirth & Blanton

- PARASTILOBEZZIA Wirth & Blanton, 1970a:10. Type-species, *leei* Wirth & Blanton (orig. des.).
- LEEI Wirth & Blanton, 1970a:10, fig. 2 (antenna, palpus, wing, legs, tarsi, claws, mandible, spermathecae, tibial comb). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia.

Genus STILOBEZZIA Kieffer

- STILOBEZZIA Kieffer, 1911:118. Type-species, *festiva* Kieffer (orig. des.). Refs. — Wirth, 1953a:57 (revision); Ingram & Macfie, 1931:191 (key to Chilean species); Lane & Forattini, 1961:83 (key to Neotropical species).

Subgenus ACANTHOHELEA Kieffer

- ACANTHOHELEA Kieffer, 1917:198 (as genus). Type-species. *pruinosa* Kieffer (mon.).
- GUIANAE (Macfie), 1940:28 (*Acanthohelea*). Type-locality: Guyana, New River. Distr. — Guyana.



Subgenus EUKRAIOHELEA Ingram & Macfie

- EUKRAIOHELEA Ingram & Macfie, 1921:347 (as genus). Type-species, *africana* Ingram & Macfie (des. Macfie, 1940d: 22).
- AMNIGENA (Macfie), 1935:56 (*Eukraiohelea*). Type-locality: Brazil, Piauí, Tutóia. Distr. — Panama, Brazil, West Indies (Jamaica).
- DORSOFASCIATA (Lutz), 1914:96, plate 8, fig. 9, plate 9, fig. 7 (wing, abdomen) (*Palpomyia*). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil.
- ELEGANTULA (Johannsen), 1907:109 (*Bezzia*). Type-locality: U.S.A., Kansas, Douglas Co., Kansas River. Distr. — U.S.A. (Kansas, Louisiana, Florida), West Indies (Jamaica, Puerto Rico).
- MACULITIBIA Lane & Forattini, 1956:207, figs. 1-2 (wing, male genitalia). Type-locality: Panama, Mandinga. Distr. — Panama.
- SUBSESSILIS Kieffer, 1917:311. Type-locality: Paraguay. Distr. — Paraguay.

Subgenus NEOSTILOBEZZIA Goetghebuer

- Stilobezzia*, subg. NEOSTILOBEZZIA Goetghebuer, 1934:53. Type-species. *Ceratopogon ochraceus* Winnertz (des. Wirth, 1953:63).
- ATRICHOPOGON Lane & Forattini, 1956:208, figs. 3-4 (wing, male genitalia). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama.
- BICINCTIPES Ingram & Macfie, 1931:195. Type-locality: Chile, Casa Pangué. Distr. — Chile.
- EDWARDSI Ingram & Macfie, 1931:198, fig. 26 e, f (male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- FURVA Ingram & Macfie, 1931:200. Type-locality: Argentina, Rio Negro, Lake Correntoso. Distr. — Argentina, Chile.
- HIRSUTA Ingram & Macfie, 1931:201, fig. 26 c (male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- MAIA Lane & Forattini, 1958:204, figs. 2-3 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- MANAOSSENSIS Lane & Forattini, 1958:205, figs. 4-5 (wing, male genitalia). Type-locality: Brazil, Amazonas, Manaus. Distr. — Brazil.
- NIGERRIMA Ingram & Macfie, 1931:196. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- ORNATA Lane & Forattini 1958:206, fig. 1 (wing). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- ORNATICRUS Ingram & Macfie, 1931:194. Type-locality: Chile, Ensenada. Distr. — Chile.
- PATAGONICA Ingram & Macfie, 1931:196, fig. 26a (male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- RAVA Ingram & Macfie, 1931:203, fig. 26d (male genitalia). Type-locality: Chile, Casa Pangué. Distr. — Chile.
- SANCTIBERNARDINI Kieffer, 1917:308. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- SUCCINEA Ingram & Macfie, 1931:200, figs. 25c (male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.



- TIBIALIS Lane & Forattini, 1956:209, figs. 5-6 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- VARIA Ingram & Macfie, 1931:191, fig. 25 (male genitalia). Type-locality: Chile, Ancud. Distr. — Argentina, Chile.

Subgenus *STILOBEZZIA* Kieffer

- ALBICOXA Lane & Forattini, 1956:210, figs. 7-8 (wing, male genitalia). Type-locality: Panama, Panama Prov., Puerto Farallón. Distr. — Panama.
- ALBOCINCTA Kieffer, 1917:309. Type-locality: Paraguay. Distr. — Paraguay.
- AMERICANA Kieffer, 1917:310. Type-locality: Paraguay, Asunción. Distr. — Paraguay.
- ANTENNALIS (Coquillett), 1901a:606 (*Ceratopogon*). Type-locality: U.S.A., District of Columbia. Distr. — U.S.A. (west to Winconsin and Texas), Panama. Ref. — Wirth, 1953a:68, fig. 12 c (male genitalia).
- BECKAE Wirth, 1953a:69, fig. 11 f (male genitalia). Type-locality: U.S.A., Florida, Citrus Co., Crystal River. — Distr. — U.S.A. (Florida, Mississippi, Panama, Peru).
- BICOLOR Lane, 1947a:208, fig. 5 (male genitalia). Type-locality: Brazil, São Paulo, Osasco. Distr. — Panama, Brazil. Refs. — Forattini & Rabello, 1956:49, fig. 5 (pupa); Lane & Forattini, 1958:208 (Panama).
- BIMACULATA Lane & Forattini, 1956:211, figs. 9, 10 (wing, male genitalia). Type-locality: Panama, Coclé, Aguadulce. Distr. — Panama, Trinidad, Brazil.
- BISPINOSA Kieffer, 1917:310, fig. 12 (tarsus). Type-locality: Paraguay. Distr. — Brazil, Paraguay. Ref. — Lane, 1947:209 (redescribed, Brazil).
- BLANTONI Lane & Forattini, 1956:211, figs. 11-12 (wing, male genitalia). Type-locality: Panama, Panama Prov., Tocumén. Distr. — Panama.
- CARIBE Lane & Forattini, 1958:208, fig. 6 (male genitalia). Type-locality: Panama, C.Z., Camarón. Distr. — Panama.
- CHIACONI Macfie, 1938:166, fig. 9 (male genitalia). Type-locality: Trinidad, St. Augustine. Distr. — Trinidad, Brazil. Ref. — Lane & Forattini, 1961:89, fig. 4 (pupa).
- COQUILLETTI Kieffer, 1917:308 (nom. nov. for *Ceratopogon pictus* Coquillett, not Meigen). Distr. — U.S.A. (Illinois and Maryland to Louisiana and Florida), West Indies, Central and South America to Brazil. Refs. — Wirth, 1953a:75, fig. 11 h (male genitalia); Lane & Forattini, 1956:213, figs. 15-16 (wing, male genitalia); Lane & Forattini, 1961:90, fig. 3 (pupa).
- pictus* Coquillett, 1905:607 (*Ceratopogon*). Type-locality: U.S.A., Virginia.
- CORACINA Kieffer, 1917:311. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay, Brazil. Ref. — Lane, 1947a:211 (redescribed).
- DIMINUTA Lane & Forattini, 1958:209, figs. 7-8 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- DRYADUM Macfie, 1940c:186. Type-locality: Guyana, Mazaruni, High Forest. Distr. — Guyana, Panama. Ref. — Lane & Forattini, 1958:210, fig. 9 (male genitalia).
- DUBITANS Lane, Forattini & Rabello, 1955:85, figs. 8-9 (male genitalia, pupa). Type-locality: Brazil, São Paulo, Capital, Parque da Água Funda. Distr. — Brazil.

- DURETI Lane & Forattini, 1958:210, figs. 10-11 (wing, male genitalia). Type-locality: Brazil, São Paulo, Ribeirão Preto. Distr. — Brazil.
- ESMERALDA Lane & Forattini, 1958:211, figs. 12-13 (wing, male genitalia). Type-locality: Panama, Coclé, Aguadulce. Distr. — Panama.
- FEMORALIS Lane & Forattini, 1956:212, figs. 13-14 (wing, male genitalia). Type-locality: Panama, C.Z., Loma Borracha. Distr. — Panama.
- FIEBRIGI Kieffer, 1917:309. Type-locality: Paraguay. Distr. — Brazil, Paraguay. Refs. — Lane, 1947a:199, fig. 1 (male genitalia); Forattini & Rabello, 1956:45, figs. 3-4 (larva, pupa).
- GLAUCA Macfie, 1939a:204, fig. 22 (male genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Costa Rica, Panama, Brazil, West Indies (Jamaica), U.S.A. (Virginia, South Carolina, Louisiana). Refs. — Wirth, 1953a:77, fig. 12h (male genitalia); Lane & Forattini, 1958:211, fig. 14 (male genitalia); Lane, Forattini, & Rabello, 1955:84, fig. 5 (pupa) (syn.; *fluminensis*).
- fluminensis* Lane, 1947:210. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Japuíba.
- GRANDIS Lane & Forattini, 1958:213, fig. 15 (wing). Type-locality: Panama, Panama Prov., Tocumén.
- KIEFFERI Lane, 1947a:205, fig. 8 (wing). Type-locality: Brazil, Minas Gerais, Governador Valadares. Distr. — Panama, Brazil. Ref. — Lane & Forattini, 1958:215, fig. 16 (male genitalia).
- MACFIEI Lane, 1947a:213, fig. 6 (male genitalia). Type-locality: Brazil, São Paulo, Osasco. Distr. — Brazil.
- MACULATA Lane, 1947a:207, fig. 4 (male genitalia). Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. — Panama, Venezuela, Brazil. Ref. — Ortiz, 1950a:429, figs. (female).
- MODESTA Lane, 1947a:206, fig. 9 (wing). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil.
- NIGROFLAVA Lane & Forattini, 1958:215, figs. 17-18 (wing, male genitalia). Type-locality: Costa Rica, San Mateo, Hguito. Distr. — Costa Rica, Panama, West Indies (Puerto Rico).
- OBSCURA Lane & Forattini, 1958:216, figs. 19-20 (wing, male genitalia). Type-locality: Panama, Panama Prov., Tocumén. Distr. — Panama.
- PALLESCENS Lane & Forattini, 1958:218, figs. 21-22 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama, Brazil.
- PANAMENSIS Lane & Forattini, 1958:218, figs. 23-24 (wing, male genitalia). Type-locality: Panama, C.Z., Camarón. Distr. — Panama, Brazil, West Indies (Puerto Rico). Ref. — Forattini, Rabello & Pattoli, 1960:3, fig. e (pupa).
- PAULISTENSIS Lane, 1947a:200, fig. 2 (male genitalia). Type-locality: Brazil, São Paulo, Casa Grande. Distr. — Panama, Brazil, West Indies (Puerto Rico).
- PUNCTIPES Wirth, 1953a:79, fig. 11 d (male genitalia). Type-locality: U.S.A., Florida, Citrus Co., Crystal River. Distr. — U.S.A. (Florida), Mexico (Tamaulipas).
- PUNCTULATA Lane, 1947a:204. Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. — Panama, Brazil. Ref. — Lane & Forattini, 1958:220, fig. 25 (male genitalia).



- RABELLOI Lane, 1947a:203, figs. 3, 7 (wing, male genitalia). Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. U.S.A. (Louisiana, Georgia, Florida), Panama, Brazil. Ref. — Wirth, 1953a:81, fig. 11a (male genitalia).
- SCUTATA Lane & Forattini, 1961:92, figs. 1-2 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- SILVICOLA Macfie, 1940c:185, fig. 3 (male genitalia). Type-locality: Guyana, Mazaruni, High Forest. Distr. — Guyana.
- SIMILANS Lane & Forattini, 1956:214, figs. 17-18 (wing, male genitalia). Type-locality: Panama, Coclé, Aguadulce. Distr. — Panama.
- SIMPLEX Lane & Forattini, 1958:222, figs. 26-27 (wing, male genitalia). Type-locality: Panama, Panama Prov., Tocumén. Distr. — Panama.
- TRANSVERSA Lane & Forattini, 1958:222, figs. 28-29 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- TRAVASSOSI Lane, 1947a:210. Type-locality: Brazil, Espírito Santo, Santa Teresa. Distr. — Brazil.
- VENEZUELENSIS Ortiz, 1950d:199, figs. 1-6 (wing, tarsi, palpus, abdomen) (*glauca* var.). Type-locality: Venezuela, Miranda, Ocumare del Tuy. Distr. — Venezuela. Ref. — Ortiz, 1963:307, figs. 1-5 (female redescribed, figured).
- WIRTHI Lane & Forattini, 1956:214, figs. 19-20 (wing, male genitalia). Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Panama.
- WYGODZINSKYI Lane, 1947a:212. Type-locality: Brazil, São Paulo, Serra da Cantareira, Cuca. Distr. — Panama, Brazil. Ref. — Lane, Forattini & Rabello, 1955:85, figs. 6-7 (male genitalia, pupa).

Tribe HETEROMYIINI

Genus CERATOBIZZIA Kieffer

- CERATOBIZZIA Kieffer, 1917:296. Type-species, *fallax* Kieffer (orig. des.).
- FALLAX Kieffer, 1917:326, fig. 19 (antenna). Type-locality: Paraguay. Distr. — Brazil, Paraguay. Ref. — Lane, 1948:232, fig. 6 (male genitalia).

Genus CLINOHELEA Kieffer

- CLINOHELEA Kieffer, 1917:295. Type-species, *Ceratopogon variegatus* Winnertz (orig. des.).
- Refs. — Lane, 1944:249 (revision, key to Neotropical species); Lane & Duret, 1954:247 (key to Neotropical species).
- ALBOPENNIS Lane, 1944:259, fig. 2 (wing). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil, Argentina.
- ARGENTINA Lane & Duret, 1954:248. Type-locality: Argentina, Corrientes, Ita Ibate. Distr. — Argentina.
- BARRETTOI Lane & Duret, 1954:249. Type-locality: Brazil, São Paulo, Ribeirão Preto. Distr. — Brazil.
- DAMASCENOI Lane & Duret, 1954:250. Type-locality: Brazil, Pará, Belém. Distr. — Brazil, Peru.



- HORACIOI Lane, 1944:257, figs. 5, 7 (male genitalia). Type-locality: Brazil, São Paulo, S. Roque. Distr. — Honduras to Panama, Brazil, Argentina.
- NEIVAI Lane, 1944:252, figs. 3, 4 (male genitalia). Type-locality: Brazil, São Paulo, S. Roque. Distr. — Panama, Brazil.
- NIGRIPES Macfie, 1939a:205, fig. 23 (wing, antenna). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Costa Rica, Panama, Brazil, Argentina. Ref. — Lane, 1946a:210, fig. 8 (male genitalia).
- RUBRICEPS Kieffer, 1917:319. Type-locality: Paraguay. Distr. — Brazil, Paraguay. Ref. — Lane, 1944:250, fig. 1 (wing).
- SALTANENSIS Lane & Duret, 1954:252. Type-locality: Argentina, Corrientes, Santo Tomé. Distr. — Brazil, Argentina.
- TOWNESI Lane, 1944:254. Type-locality: Brazil, Mato Grosso, Pontal dos Morros Azuis. Distr. — Brazil.
- TOWNSENDI Lane, 1944:256. Type-locality: Brazil, São Paulo, S. Roque. Distr. — Peru, Brazil, Argentina.
- WYGODZINSKYI Lane, 1948:231. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil.

Genus HETEROMYIA Say

- HETEROMYIA Say, 1825: plate 35 (1859:79). Type-species, *fasciata* Say (mon.).
- Pachyleptus* Walker, 1856:426. Type-species, *fasciatus* Walker (mon.).
Ref. — Duret & Lane, 1955:35 (key to Neotropical species).
- ANTEQUERAE (Lynch Arribáizaga), 1893:227 (*Pachyleptus*). Type-locality: Argentina, Antequera Island, Rio Paraná. Distr. — Argentina.
- BEJARANOI Duret & Lane, 1955:36, figs. 2, 3 (wing, male genitalia). Type-locality: Argentina, Chaco (as Presidente Perón Prov.), Capitán Colari. Distr. — Argentina.
- CASTANEA Lane, 1946a:214. Type-locality: Brazil, São Paulo, Severínia. Distr. — Brazil.
- CHAQUENSIS Duret & Lane, 1955:37, fig. 1 (wing). Type-locality: Argentina, Chaco (as Presidente Perón Prov.), Capitán Solari. Distr. — Argentina.
- CLAVATA Williston, 1900:225. Type-locality: Mexico, Vera Cruz. Distr. — Mexico.
- CORRENTINA Duret & Lane, 1955:39, fig. 4 (male genitalia). Type-locality: Argentina, Chaco (as Presidente Perón Prov.), Tacuarí. Distr. — Panama, Argentina.
- KIEFFERI Lane, 1946a:213, fig. 7 (spermathecae). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil.
- LAMPROGASTER Edwards, 1933:87. Type-locality: Argentina, San José. Distr. — Argentina.
- NIGRA Kieffer, 1917:326. Type-locality: Paraguay. Distr. — Brazil, Paraguay. Refs. — Lane, 1946a:211, figs. 6, 7 (wing, spermathecae); Lane, 1961c:40, fig. 1 (male genitalia).
- fasciatus* Walker, 1856:426 (*Pachyleptus*) (preoccupied by *Heteromyia fasciatus* Say). Type-locality: South America.
- ORELLANA (Roback), 1957:1, figs. A-F (wing, leg, tibial comb, antenna, spermathecae (*Palpomyia*)). Type-locality: Peru, Iquitos. Distr. — Peru.
- N. COMB.



RUFA Kieffer, 1917:325, fig. 18 (palpus). Type-locality: Colombia, Baranquilla. Distr. — Panama, Colombia.

caloptera Kieffer, 1919:192. Unnecessary nom. nov. for *rufa* (Loew) of Malloch, 1915.

Genus NEUROHELEA Kieffer

NEUROHELEA Kieffer, 1925:112. Type-species, *Ceratopogon luteitarsis* Meigen (mon.).

OEDIDACTYLA Ingram & Macfie, 1931:212. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

TURGIDIPES Ingram & Macfie, 1931:214. Type-locality: Chile, Mechuque Island. Distr. — Chile.

Genus PELLUCIDOMYIA Macfie

PELLUCIDOMYIA Macfie, 1939c:99. Type-species, *ugandae* Macfie (mon.).

Macfiehelea Lane, 1946a:208. Type-species, *oliveirai* Lane (orig. des.).

Refs. — Lane, 1956b:435 (revision, key, as *Macfiehelea*); Wirth, 1960a:1 (revision, key; synonym: *Macfiehelea*).

BLANTONI (Lane), 1956b:435 (*Macfiehelea*). Type-locality: Panama, C. Z., Mojinga Swamp. Distr. — Panama.

LANEI Wirth & Ratanaworabhan, 1971:446, fig. 1 (many parts). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia, West Indies (Jamaica).

OLIVEIRAI (Lane), 1946a:209, fig. 5 (wing) (*Macfiehelea*). Type-locality: Brazil, Minas Gerais, Governador Valadares. Distr. — Brazil.

WIRTHI (Lane), 1956b:437 (*Macfiehelea*). Type-locality: Panama, C.Z., Mindi Dairy. Distr. — Mexico to Colombia. Ref. — Wirth & Ratanaworabhan, 1971:446, fig. 1 (many parts) (redescribed).

Tribe SPHAEROMIINI

Ref. — Wirth, 1962:272 (revision of North American species, key to genera).

Genus JOHANNSENYMYIA Malloch

JOHANNSENYMYIA Malloch, 1915a:332. Type-species, *halteralis* Malloch (Wirth, 1952e:211) = *argentata* (Loew).

Dicrohelea Kieffer, 1917:363. Type-species, *Palpomyia filicornis* Kieffer (Macfie, 1940d:26).

Ref. — Wirth, 1962:276 (synonymy).

BLANTONI (Lane & Wirth), 1961:81 (*Dicrohelea*). Type-locality: Panama, La Jolla. Distr. — Panama.

LANEI Wirth, 1965c:4. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



Genus LANEHELEA Wirth & Blanton

- LANEHELEA Wirth & Blanton, 1972b:433. Type-species, *leei* Wirth & Blanton (orig. des.).
- LEEI Wirth & Blanton, 1972b:434, fig. 1 a-1 (female, many parts). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia.
- SPINIFEMUR Wirth & Blanton, 1972b:436, 2 a-1 (female, many parts). Type-locality: Colombia, Valle, Rio Raposo. Distr. — Colombia.

Genus MALLOCHOHELEA Wirth

- MALLOCHOHELEA Wirth, 1962:278. Type-species, *Johannsenomyia albibasis* Malloch (orig. des.).
- AENIPES (Macfie), 1940c:193 (*Bezzia*). Type-locality: Guyana, Mazaruni. Distr. — Guyana.
- NEMORALIS (Macfie), 1940c:190, fig. 5 (male genitalia) (*Johannsenomyia*). Type-locality: Guyana, Mazaruni. Distr. — Guyana.
- NIGRIPES (Macfie), 1939a:216 (*Johannsenomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PULLATA (Wirth), 1952e:213, fig. 25j (tarsus) (*Johannsenomyia*). Type-locality: U.S.A., California, Imperial Co., El Centro. Distr. — U.S.A. (California to Texas), Mexico (Tamaulipas), Brazil.
- bicellii* Lane, 1961a:449 (*Nilobezzia*). Type-locality: Brazil, Amapá, Rio Felício.

Genus NEOBEZZIA Wirth & Ratanaworabhan

- NEOBEZZIA Wirth & Ratanaworabhan, 1972b:477. Type-species, *clavipes* Wirth & Ratanaworabhan (orig. des.).
- Ref. — Wirth & Ratanaworabhan, 1972:476 (revision, key).
- ALBITARSIS Wirth & Ratanaworabhan, 1972b:478, fig. 1 (female; many parts). Type-locality: Brazil, Pará, Rio Paru do Oeste, Igarapé Okueima. Distr. — Brasil.
- AMNICOLA (Macfie), 1940a:30 (*Bezzia*). Type-locality Guyana, New River. Distr. — British Honduras, Costa Rica, Panama, Colombia, Peru, Guyana, Brazil. Refs. — Mayer, 1959:230, fig. 1 (pupa); Wirth & Ratanaworabhan, 1972b:481, fig. 2 (many parts).
- BLANTONI Wirth & Ratanaworabhan, 1972b:482, fig. 3 (female, many parts). Type-locality: Panama, C.Z., Gamboa, Rio Agua Salud. Distr. — Nicaragua, Panama, Colombia, Brazil.
- BRASILIAE (Lane), 1961c:37 (*Macropeza*). Type-locality: Brazil, Brasília, D.F. Distr. — Brazil.
- CLAVIPES Wirth & Ratanaworabhan, 1972b:485, fig. 4 (female, many parts). Type-locality: Brazil, Amazonas, Rio Cuieiras, Igarapé Cachoeira. Distr. — Panama, Brazil.
- COSTARICAE Wirth & Ratanaworabhan, 1972b:486, fig. 5 (female, many parts). Type-locality: Costa Rica, Puntarenas Prov., Palmar Sur. Distr. — Costa Rica, Panama.
- FITTKAUI Wirth & Ratanaworabhan, 1972b:489, fig. 6 (female, many parts). Type-locality: Brazil, Amazonas, Rio Preto, Tiririca. Distr. — Brazil.



Genus NILOBEZZIA Kieffer

NILOBEZZIA Kieffer, 1921:24. Type-species, *armata* Kieffer (mon.).

Dicrobezzia Kieffer, of Lane, 1948:232.

Refs. — Lane, 1948:232 (key, as *Dicrobezzia*); Lane, 1961:449 (key).

ATELES (Macfie), 1940b:78 (*Bezzia*). Type-locality: Brazil, Mundo Novo. Distr. — Brazil. N. COMB.

NEOTROPICA (Macfie), 1940b:78 (*Bezzia*). Type-locality Brazil, Mundo Novo. Distr. — Brazil.

PARAENSIS (Lane), 1958:33 (*Bezzia*). Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.

SCHWARZII (Coquillett), 1901a:605 (*Ceratopogon*). Type-locality: U.S.A., Texas, Sharpsburg. Distr. — U.S.A. (Texas to Florida, Maryland, New York), Panama, Brazil, West Indies (Cuba). Ref. — Wirth, 1962:285, figs. 12-14 (tarsus, male genitalia) (redescribed, distribution; synonym: *brasiliensis*).

banksi Gerry, 1933:94 (*Bezzia*). Type-locality: Cuba, Soledad, Cienfuegos.

brasiliensis Lane, 1958:28 (*Bezzia*). Type-locality: Brazil, Rio de Janeiro, Itaguaí.

Tribe STENOXENINI

Ref. — Wirth & Ratanaworabhan, 1972a:1369 (revision, keys).

Genus PARYPHOCONUS Enderlein

PARYPHOCONUS Enderlein, 1912:57. Type-species, *angustipennis* Enderlein (orig. des.).

Refs. — Lane, 1956a:299 (revision, key); Lane, 1961b:453 (key); Wirth & Ratanaworabhan, 1972a:1369 (revision, key).

AEMULUS Macfie, 1940c:180, fig. 1 (male genitalia). Type-locality: Guyana, Marzaruni, High Forest. Distr. — Guyana, Brazil. Ref. — Lane, 1946a:206, fig. 7 (spermatheca).

AMAPAENSIS Lane, 1961a:450, fig. 1 (wing). Type-locality: Brazil, Terr. Amapá, Serra do Navio. Distr. — Brazil.

ANGUSTIPENNIS Enderlein, 1912:57, figs. 1-4 (wing, tarsus, head, thorax). Type-locality: Brazil, Santa Catarina. Distr. — Mexico, Colombia, Peru, Venezuela, Brazil, Argentina. Refs. — Macfie, 1939b:5, plate 1, fig. 3 (wing); Lane, 1946a:202, figs. 2, 7 (wing, spermatheca); Kieffer, 1917:332, fig. 21 (tarsus); Mayer, 1952:583, figs. 1-2 (egg); Wirth & Ratanaworabhan, 1972a:1371, fig. 1 (many parts).

ANOMALICORNIS Kieffer, 1917:333, fig. 22 (antenna, tarsus). Type-locality: Colombia, Santa Marta, and Paraguay, Puerto Max. Distr. — Colombia, Mexico, Honduras, Venezuela, Paraguay.

BARRETOI Lane, 1946a:203, figs. 4, 7 (wing, spermatheca). Type-locality: Brazil, Goiás, Corumbá, Fazenda Monjolino. Distr. — Mexico, Venezuela, Brazil.

BATESI Lane, 1961b:455, fig. 2 (wing). Type-locality: Brazil, Brasília, D.F. Distr. — Brazil.

- ENDERLEINI Lane, 1956a:301. Type-locality: Brazil, Mato Grosso, Bodoquena. Distr. — Brazil.
- FUSCUS Lane, 1946a:206, figs. 3, 7 (wing, spermatheca). Type-locality: Brazil, Goiás, Corumbá. Distr. — Brazil.
- GOIANENSIS Lane, 1961b:456, fig. 3 (wing). Type-locality: Brazil, D.F., Brasília. Distr. — Brazil.
- GRANDIS Macfie, 1939b:6, plate 1, fig. 2 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- KIEFFERI Lane, 1956a:302. Type-locality: Brazil, Goiás, Corumbá. Distr. — Brazil.
- LANEI Wirth, 1959b:236, fig. 3 (male genitalia). Type-locality: Brazil, Rio Madeira, Igarapé das Três Casas, from pupa in vegetation of *Polygonum*. Distr. — Brazil. Ref. — Mayer, 1959:233, fig. 4 (pupa).
- MACFIEI Lanc, 1946a:203, figs. 1, 7 (wing, spermatheca). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- MAYERI Wirth, 1959b:236. Type-locality: Brazil, Rio Madeira, Igarapé das Três Casas, from pupa in vegetation of *Polygonum*. Distr. — Brazil. Ref. — Mayer, 1959:232, fig. 3 (pupa).
- NEOTROPICALIS (Lane), 1948:229 (*Macropeza*). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- NIGRIPES Macfie, 1939b:8, plate 1, fig. 4 (wing). Type-locality: Argentina, Misiones, Puerto Aguirre. Distr. — Guyana, Brazil, Argentina.
- NUBIFER Macfie, 1939b:5, plate 1, fig. 1 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Panama, Brazil, Uruguay, Bolivia.
- OLIVEIRAI Lane, 1956:303. Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.
- PAULISTENSIS Lane, 1961b:457. Type-locality: Brazil, São Paulo, Ribeirão Preto, Tamanduá. Distr. — Brazil.
- SONORENSIS Wirth & Ratanaworabhan, 1972a:1374, fig. 5 a-c (antenna, wing, spermatheca). Type-locality: Mexico, Sonora, Alamos. Distr. — U.S.A. (Oklahoma), Mexico (Sonora).
- SUBFLAVUS Macfie, 1940a:23. Type-locality: Guyana, New River. Distr. — Colombia, Guyana, Brazil.
- TERMINALIS (Coquillett), 1904b:90 (*Ceratopogon*). Type-locality: Nicaragua, San Marcos. Distr. — British Honduras to Panama, Peru, Guyana, Brazil.
guyanae Macfie, 1940c:180. Type-locality: Guyana, Mazaruni.
- TRAVASSOSI Lane, 1956a:304. Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.
- UNIMACULATUS Macfie, 1940c:179. Type-locality: Guyana, Tukeit, Gorge of Potaro. Distr. — Guyana, Brazil.
- WIRTHI Lanc, 1961b:458. Type-locality: Brazil, D.F., Brasília. Distr. — Brazil.
- WYGODZINSKYI Lane, 1946:208, fig. 7 (spermatheca). Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. — Brazil.

Genus STENOXENUS Coquillett

- STENOXENUS Coquillett, 1899:61. Type-species, *johnsoni* Coquillett (mon.).
Refs. — Macfie, 1939a:141 (key); Lane, 1961b:452 (revision, key to Neotropical species); Wirth & Ratanaworabhan, 1972a:1378 (revision and key).



- ARCUATUS Wirth & Ratanaworabhan, 1972a:1379, fig. 9 f (wing). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.
- BLANTONI Wirth & Ratanaworabhan, 1972a:1379, fig. 8 a-e (antenna, palpus, wing, spermatheca, abdomen). Type-locality: Panama, Darien, Santa Fe. Distr. — Panama.
- BRSILIENSIS Macfie, 1939a:140, fig. 2 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Argentina.
- CARRERAI Lane, 1956:305, fig. 1 (wing) (as *carreirai*). Type-locality: Brazil, Goiás, Jataí. Distr. — Colombia, Brazil.
- DIMORPHUS Kieffer, 1909:46. Type-locality: Peru, Callanga. Distr. — Peru.
- EXCENTRICUS Lane, 1961b:453, fig. 1 (wing). Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.
- FULVUS Johannsen, 1927:70. Type-locality: Peru, Putumayo Dist., La Chorerra. Distr. — Peru.
- JOHNSONI Coquillett, 1899:61. Type-locality: U.S.A., New Jersey, Delaware Water Gap. Distr. — Southeastern U.S.A., Mexico to Panama.
- LANEI Wirth & Ratanaworabhan, 1972a:1381, fig. 11 a-d (antenna, palpus, wing, spermatheca, abdomen). Type-locality: Panama, Darien, Santa Fe. Distr. — Panama.
- LIMPIDUS Wirth & Ratanaworabhan, 1972a:1383, fig. 9 g (wing). Type-locality: Costa Rica, Higuito, San Mateo. Distr. — Costa Rica.
- MARCINALIS Wirth & Ratanaworabhan, 1972a:1385, fig. 9 h (wing). Type-locality: Panama, C.Z., Mojinga Swamp. Distr. — Panama.
- NICER Lane, 1948:230, fig. 5 (male genitalia) (as *nigrus*). Type-locality: Brazil, Goiás, Corumbá, Fazenda Monjolinho. Distr. — Brazil.
- PARAENSIS Lanc, 1956a:306, fig. 2 (wing). Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.
- SETICER Macfie, 1939a:138, fig. 1 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — West Indies (Jamaica, Dominican Republic), Colombia, Brazil.

Tribe PALPOMYIINI

Genus BEZZIA Kieffer

- BEZZIA Kieffer, 1899:69. Type-species, *Ceratopogon ornatus* Meigen (orig. des.). Ref. — Lane, 1958:25 (revision, key to Neotropical species).
- ACANTHODES Macfie, 1940c:192. Type-locality: Guyana, Mazaruni. Distr. — Guyana, Brazil.
- ALBUQUERQUEI Lane, 1961c:43. Type-locality: Brazil, D.F., Brasília. Distr. — Brazil.
- BREVICORNIS (Kieffer), 1917:328, fig. 20 (*Allobezzia*). Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- CARIOCA Lane, 1958:30. Type-locality: Brazil, Rio de Janeiro, Itaguaí. Distr. — Brazil.
- COLORADENSIS Wirth, 1952e:238. Type-locality: U.S.A., California, Imperial Co., Westmorland. Distr. — U.S.A. (California), West Indies (St. John).
- CONCOLORIPES Macfie, 1940a:31. Type-locality: Guyana, New River. Distr. — Guyana.

- FLAVIDA (Johannsen), 1943a:761, figs. 20, 22 (antenna, thorax) (*Ceratobezzia*). Type-locality: Guyana, Bertina. Distr. — Guyana. N. COMB.
- FLUMINENSIS Lane, 1948:236. Type-locality: Brazil, Rio de Janeiro, Estrada Rio-São Paulo, km 47. Distr. — Brazil. Ref. — Lane, 1958:31, fig. 1 (male genitalia).
- GIBBERA (Coquillett), 1905:60 (*Ceratopogon*). Type-locality: Cuba, Cayamas. Distr. — U.S.A. (Florida), West Indies (Cuba).
- COIANENSIS Lane, 1961c:44. Type-locality: Brazil, Brasília, D.F. Distr. — Brazil.
- LENKOI Lane, 1958:31, figs. 2-3 (wing, male genitalia). Type-locality: Brazil, São Paulo, Barueri. Distr. — Panama, Brazil.
- MAZARUNI Macfie, 1940c:193. Type-locality: Guyana, Mazaruni. Distr. — Guyana.
- PULCHRIPES Kieffer, 1917:330. Type-locality: Paraguay. Distr. — Brazil, Paraguay. Refs. — Mayer, 1959:232, fig. 2 (pupa); Wirth, 1959b:234, fig. 1 (male genitalia).
- PUNCTIPENNIS (Williston), 1896:278, plate 9, fig. 39 (wing) (*Ceratopogon*). Type-locality: St. Vincent. Distr. — U.S.A. (California, Nevada, Florida), West Indies, Central and South America.
- SETULOSA (Loew), 1861:312 (Cent. 1, no. 8) (*Ceratopogon*). Type-locality: U.S.A., District of Columbia. Distr. — North America (California to Ontario and Florida), West Indies (Jamaica).
- SNOWI Lane, 1958:34, fig. 4 (genitalia). Type-locality: Guatemala. Distr. — Guatemala.
- SUBFUSCA Macfie, 1939a:218. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Panama, Brazil.
- TELMATOPHILA Macfie, 1940b:77. Type-locality: Brazil, Riacho do Atalho. Distr. — Brazil. Ref. — Lane, 1958:35, fig. 5 (male genitalia).
- VENUSTULA (Williston), 1896:278, plate 9, fig. 38 (*Ceratopogon*). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent, Trinidad).

Genus PACHYHELEA Wirth

- PACHYHELEA Wirth, 1959a:50. Type-species, *Ceratopogon magnus* Coquillett (orig. des.) = *pachymera* (Williston).
- PACHYMERA (Williston), 1900:224 (*Ceratopogon*). Type-locality: Mexico, Medellín near Vera Cruz. Distr. — U.S.A. (Texas), Mexico to Argentina. Refs. — Wirth, 1959a:52 (redescribed; synonyms: *albidiventris*, *latifemoris*); Lane, 1961:42 (Brazil).
- albidiventris* Kieffer, 1917:316 (*Sphaeromias*). Type-locality: Colombia, Barranquilla.
- latifemoris* Ingram & Macfie, 1931:231 (*Johannsenomyia*). Type-locality: Argentina, Buenos Aires, San Isidro. Ref. — Macfie, 1940b:75 (redescribed).

Genus PALPOMYIA Meigen

- PALPOMYIA Meigen, 1818:82. Type-species, *Ceratopogon flavipes* Meigen (mon.). Refs. — Ingram & Macfie, 1931:215 (key to Chilean species); Macfie, 1939a:207 (key to Neotropical species); Lane, 1947b:446 (key); Lane, 1960:381 (revision, key to Neotropical species).



- ACULEATA Ingram & Macfie, 1931:215. Type-locality: Chile, Ancud. Distr. — Chile.
- ALMEIDAI Lane, 1946b:219, figs. 1-2 (wing, abdomen) (*Sphaeromyias*). Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- APRICANS Ingram & Macfie, 1931:219, fig. 32a-c (male genitalia). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina, Chile.
- BARRETTOI Lane, 1947b:442. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Marombá. Distr. — Brazil.
- BOLIVIENSIS Kieffer, 1917:324, fig. 17 (tarsus). Type-locality: Bolivia, Coroico. Distr. — Bolivia.
- BRASILIENSIS Macfie, 1939a:213. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CALLANGANA Kieffer, 1917:323, fig. 16 (tarsus). Type-locality: Peru, Callanga. Distr. — Peru.
- CARIOCA Lane, 1960:384. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Agulhas Negras. Distr. — Brazil.
- CARRERAI (Lane), 1948:235 (*Dicrobezzia*, as *carreirai*). Type-locality: Brazil, São Paulo, São Miguel. Distr. — Brazil.
- CASTANEA Macfie, 1939a:211. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CATARINENSIS Lane, 1960:385. Type-locality: Brazil, Santa Catarina, Florianópolis, Rio Tavares. Distr. — Brazil.
- CHILENSIS Ingram & Macfie, 1931:218. Type-locality: Chile, Ensenada. Distr. — Brazil, Chile.
- COLUMBIANA Kieffer, 1917:323, fig. 15 (tarsus). Type-locality: Colombia, Sierra San Lorenzo. Distr. — Colombia.
- CONIFERA Macfie, 1939a:214. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Lane, 1947b:439 (redescribed).
- COROICOENSIS Wirth, nom. nov.
boliviensis Kieffer, 1917:331 (*Jenkinsia*). Type-locality: Bolivia, Coroico. Distr. — Bolivia. Preoccupied in *Palpomyia* by *boliviensis* Lane, 1917:324.
- CRASSICRUS Kieffer, 1917:321. Type-locality: Paraguay. Distr. — Brazil, Paraguay.
- GUARANI Lane, 1946b:221, figs. 3-9 (wing, abdomen, male genitalia, pupa). Type-locality: Brazil, São Paulo, Cantareira. Distr. — Brazil.
- INERMICOLLIS Kieffer, 1917:323. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.
- JOHANNSENI (Lane), 1948:233 (*Dicrobezzia*). Type-locality: Brazil, São Paulo, Itatinga. Distr. — Brazil.
- LACUSTRIS Lane, Forattini & Rabello, 1955:83, figs. 3-4 (male genitalia, pupa). Type-locality: Brazil, São Paulo, Capital, Parque da Água Funda. Distr. — Brazil.
- LIMNOCHARES (Macfie), 1940b:76, fig. 1 (male genitalia) (*Sphaeromyias*). Type-locality: Brazil, Canafistula. Distr. — Brazil. N. COMB.
- LONGICORNIS (Williston), 1896:280, plate 9, fig. 43 (wing, antenna) (*Ceratopogon*) (preoc. by *Ceratopogon longicornis* Waltl 1837). Type-locality: St. Vincent. Distr. — West Indies (St. Vincent). Needs new name but only after comparison with other named species.

- LUTZI Lane, 1947b:440. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo Distr. — Brazil.
- MACULICRUS Ingram & Macfie, 1931:230. Type-locality: Argentina, Misiones, Posadas. Distr. — Brazil, Argentina.
- MELICHROA Macfie, 1939a:212. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- MICROCHELA Kieffer, 1917:321. Type-locality: Colombia, Sierra San Lorenzo. Distr. — Colombia.
- NIGROFLAVA Lane, 1947b:446. Type-locality: Brazil; São Paulo, Campos do Jordão. Distr. — Brazil.
- NIGROSCUTELLATA Lane, 1947b:444. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil.
- OLIVEIRAI Lane, 1947:443. Type-locality: Brazil, Minas Gerais, Governador Valadares. Distr. — Brazil.
- PAMPANA Lane, 1960:386. Type-locality Brazil, Rio Grande do Sul, Arroio Grande. Distr. — Brazil.
- PARAENSIS Lane, 1960:387. Type-locality: Brazil, Pará, Cachimbo. Distr. — Brazil.
- PATAGONICA Ingram & Macfie, 1931:222. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.
- PAULISTENSIS Lane, 1947b:445. Type-locality: not stated. Distr. — Panama, Brazil.
- SCABRA (Coquillett), 1905:62 (*Ceratopogon*). Type-locality: Mexico, Tabasco, Frontera. Distr. — Mexico.
- SENIOR Ingram & Macfie, 1931:222 (*apricans* var.). Type-locality: Chile, Castro, Distr. — Chile.
- SORDIDIPES Macfie, 1939a:209. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Refs. — Lane, 1947:440, fig. 1 (male genitalia); Lane, Forattini & Rabello, 1955:81, fig. 1 (pupa).
- SPINIFEMUR (Lane), 1948:234, fig. 8 (male genitalia) (*Dicrobezzia*). Type-locality: Brazil, São Paulo, Osasco. Distr. — Brazil.
- SPINOSA Lutz, 1914:193, plate 9, fig. 5 (leg.). Type-locality: Brazil, Rio de Janeiro, Manguinhos. Distr. — Brazil. Ref. — Lane, 1945:371 (redescribed from Lutz's type material).
- SUBFUSCULA Ingram & Macfie, 1931:216, fig. 32d, 34a (male genitalia). Type-locality: Chile, Casa Pangue. Distr. — Argentina, Chile.
- TAMOIOI Lane, 1960:388. Type-locality: Brazil, Rio de Janeiro, Itatiaia, Fazenda Penido. Distr. — Brazil.
- TENUICRUS Kieffer, 1917:320. Type-locality: Paraguay, San Lorenzo. Distr. — Paraguay.
- UMBELLA Macfie, 1939a:208. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Lane, 1960:389, fig. 1 (male genitalia).
- VERSICOLOR Macfie, 1939a:215. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- WIRTHI Lane, Forattini & Rabello, 1955:82, fig. 2 (pupa). Type-locality: Brazil, São Paulo, Capital, Parque da Água Funda. Distr. — Brazil.

Unplaced Species of Ceratopogonidae

- Ceratopogon* CLARIPENNIS Lynch-Arribáizaga, 1893:229. Type-locality: Argentina, Buenos Aires, Navarro. Distr. — Argentina.



Excluded from Ceratopogonidae

Ceratopogon blanchardi Ichés, 1906:266, figs. 4, 5 (female, antenna). Type-locality: Argentina, Chaco Austral. Belongs in subfamily Tanypodinae in Chironomidae.

LITERATURE CITED

- ADAMSON, A. M.
1939. Observations on biting sandflies (Ceratopogonidae) in Trinidad, British West Indies. *Trop. Agr.* 16:79-81.
- AITKEN, T. H. G.
1957. Pestiferous Trinidad sandflies. *Jour. Trinidad Field Nat. Club* 1957:23-26.
- ATCHLEY, W. R.
1970. A biosystematic study of the subgenus *Selfia* of *Culicoides* (Diptera: Ceratopogonidae). *Kansas Univ. Sci. Bull.* 49:181-336, figs.
- BAKER, C. F.
1907. Remarkable habits of an important predaceous fly (*Ceratopogon eriophorus* Will.). *U.S. Dept. Agr. Bur. Ent. Bull.* 67:117-118.
- BARBOSA, F. A. SIMÕES
1943. Descrição de "*Culicoides recifensis*" n. sp. e do macho de "*Culicoides reticulatus*" Lutz (Diptera, Chironomidae). *Rev. Brasil. Biol.* 3:261-264, figs.
1944. "*Culicoides insignis*" Lutz, com a descrição do hipopígio (Diptera, Chironomidae). *Rev. Brasil. Biol.* 4:259-261, figs.
1947. *Culicoides* (Diptera: Heleidae) da região neotrópica. *An. Soc. de Biol. Pernambuco* 7:3-30, figs.
1951. A change of specific name in the genus *Culicoides* (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 53:163.
1952. *Novos subsídios para o conhecimento dos Culicoides neotrópicos* (Diptera: Heleidae). Univ. Recife Thesis, Imprensa Industrial, Recife, pp. 1-21, figs.
1953. *Novos subsídios para o conhecimento dos Culicoides neotrópicos* (Diptera: Heleidae). *Pub. Avuls. do Inst. Aggeu Magalhães, Recife* 2:11-42, figs.
- BARRETTO, M. P.
1944. Sobre o gênero "*Culicoides*" Latreille, 1809, com a descrição de três novas espécies (Diptera Ceratopogon.). *An. Fac. Med. Univ. São Paulo* 20:89-105, figs.
- BAUER, D. L.
1961. Midge-flies biting butterflies. *Jour. Lepidopt. Soc.* 15:91-92.
- BECK, E. C.
1951. A new *Culicoides* from Florida (Diptera, Ceratopogonidae). *Florida Ent.* 34:135-136, figs.
- BEQUAERT, J.
1924. Report of an entomological trip to the Truxillo Division, Honduras, to investigate the sand-fly problem. *United Fruit Co. 13th Ann. Rept.* 1924:193-206.



BLANTON, F. S., O. H. GRAHAM, & C. M. KEENAN

1955. Notes on *Culicoides furens* (Poey) at Fort Kobbe, Canal Zone. *Mosquito News* 15:13-18.

BRÈTHES, J.

1912. Descripción de un nuevo género y especie nueva de Chironomidae. *Anal. Mus. nac. Buenos Aires* 15:451-453, 1 fig.

1914. Description de six Cécidomyiidae (Dipt.) de Buenos Aires. *Anal. Mus. nac. Hist. Nat.*, Buenos Aires 26:151-156, figs.

CARTER, H. F.

1919. New West African Ceratopogoninae. *Ann. Trop. Med. & Parasit.* 12:289-302, 1 plate.

1921. A revision of the genus *Leptoconops*, Skuse. *Bull. Ent. Res.* 12:1-28, figs.

CARTER, H. F., A. INGRAM & J. W. S. MACFIE

1920-1922. Observations on the Ceratopogonine midges of the Gold Coast with descriptions of new species. I-IV. *Ann. Trop. Med. & Parasit.* 14:187-274, 309-331; 15:177-212, 5 plates.

CAVALIERI, F.

1961. Notas sobre Ceratopogonidae I. *Forcipomyia* (F.) *wygodzinski* sp. n. para Tierra del Fuego. *Rev. Soc. Ent. Argentina* 23:17-19, figs.

1962a (1961). Notas sobre Ceratopogonidae (Diptera, Nematocera). II. Sobre dos nuevas especies de *Forcipomyia* para Tierra del Fuego, *Forcipomyia* (F.) *delpontei* n. sp. y F. (F.) *piroskyi* n. sp. *Acta Zool. Lilloana* 18:169-175, figs. (publ. May 23, 1961).

1962b (1961). Notas sobre Ceratopogonidae (Diptera, Nematocera). III. Sobre un nuevo diptero hematofago para Argentina, *Lasiohelea saltensis* n.sp. y notas sobre las especies neotropicales de *Lasiohelea*. *Acta Zool. Lilloana* 18:359-365, figs. (publ. May 23, 1961).

1966. Notas sobre Ceratopogonidae (Diptera, Nematocera) V. Descripción de una nueva especie de Venezuela: *Culicoides* (*Oecacta*) *birabeni* sp. n. *Physis* 26:59-63, figs.

CAVALIERI, F., & I. F. CHIOSSONE

1965. Notas sobre Ceratopogonidae (Diptera, Nematocera). IV. "*Pterobosca*" Macfie, nuevo género para Argentina con la redescrpcion de "*P. incubans*" Macfie. *Neotropica* 11:64-70, figs.

1966a. Datos sobre la familia de dipteros hematofagos Leptoconopidae en la Argentina, con la descripción de *Leptoconops* (*Leptoconops*) *casali* sp. n. y la redescrpcion de *L. (L.) petrocchia* Shannon y Del Ponte, 1927. *Physis* 26:43-57, figs.

1966b. Sobre el conocimiento actual del género hematofago *Culicoides* Latreille, 1809, en la Argentina (Diptera, Ceratopogonidae). *Physis* 26:145-153, figs.

CHIOSSONE, I. F.

1969. Sobre algunos datos adicionales de *Leptoconops* (*L.*) *casali* Cavalieri y Chiossone, 1966, con la descripción del aparato bucal (Diptera: Leptoconopidae). *Neotropica* 15:17-20, figs.

CLASTRIER, J.

1968. Deux Ceratopogonidae nouveaux de la Guyane Française (Diptera, Ceratopogonidae). *Arch. Inst. Pasteur Guyane Française et de l'Inini*, 21:85-92, figs.

1971. Deux nouveaux *Culicoides* (Diptera, Ceratopogonidae) de la Guyane Française. *Ann. Parasit. Humaine et Comparée* 46:285-294, figs.
1972. *Forcipomyia* (*F.*) *squamithorax* n. sp. de la Guyane Française (Dipt. Ceratopogonidae). *Bull. Soc. Ent. France* 77:170-176, figs.
- COQUILLET, D. W.
1899. A new dipterous family related to the Chironomidae. *Ent. News* 10:60-61, 1 fig.
- 1901a. New Diptera in the U. S. National Museum. *Proc. U. S. Nat. Mus.* 23:593-618.
- 1901b. Papers from the Hopkins Stanford Galapagos Expedition, 1898-1899. II. Entomological results (2): Diptera. *Proc. Washington Acad. Sci.* 3:371-379.
- 1904a. A new *Ceratopogon* from Brasil. *Jour. New York Ent. Soc.* 12:35-36.
- 1904b. New Diptera from Central America. *Proc. Ent. Soc. Washington* 6:90-98.
1905. New nematoceros Diptera from North America. *Jour. New York Ent. Soc.* 13:56-69.
- CORDERO, E. H.
1929. Contribución al estudio de los Dípteros de Uruguay, I. *Lophomyidum uruguayense* n. gen., n. sp. nueva Ceratopogonina hematófaga. *An. Mus. Hist. Nat. Montevideo* (2)3:93-108, figs.
- CURTIS, J.
1829. *British Entomology*. Vol 6, plates 242-289. London.
- DAVIES, J. B.
1967. The distribution of sandflies (*Culicoides* spp.) breeding in a tidal mangrove swamp in Jamaica and the effect of tides on the emergence of *C. furens* Poey and *C. barbosa* (Wirth and Blanton). *W. Ind. Med. Jour.* 16:39-50, figs.
- DAVIES, J. B. & J. R. LINLEY
1965. Observations on the breeding sites of the sandfly *Leptoconops bequaerti* in the Montego Bay area of Jamaica, with a note on one breeding site in Honduras. *Caribbean Jour. Sci.* 5:117-128, figs.
- DOW, M. I. & W. W. WIRTH
1972. Studies on the genus *Forcipomyia*, 2. The Nearctic species of the subgenera *Thyridomyia* and *Synthyridomyia* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 65:177-201, figs.
- DOWNES, J. A.
1971. The ecology of blood-sucking Diptera: an evolutionary perspective. pp. 230-258, in *Ecology and Physiology of Parasites*, ed. A. M. Fallis. Toronto Univ. Press.
- DURET, J. P. & J. LANE
1955. Novas *Heteromyia* da Argentina (Diptera. Ceratopogonidae). *Dusenía* 6:35-40, figs.
- EDWARDS, F. W.
1922. On some Malayan and other species of *Culicoides*, with a note on the genus *Lasiohelea*. *Bull. Ent. Res.* 13:161-167, 1 plate.



1926. On the British biting midges (Diptera, Ceratopogonidae). *Trans. Ent. Soc. London* 74:389-426, 3 figs., 2 plates.
1933. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera (Fortsetzung) XXXIX-XLI. Cecidomyiidae, Chironomidae, Ceratopogonidae. *Konowia* 12:86-88.
- ENDERLEIN, G.
1912. *Paryphoconus*, eine neue Chironomiden-Gattung aus Brasilien. *Stett. Ent. Ztg.* 73:57-60, figs.
- EWEN, A. B. & L. G. SAUNDERS
1958. Contributions toward a revision of the genus *Atrichopogon* based on characters of all stages (Diptera, Heleidae). *Canad. Jour. Zool.* 36:671-724, figs.
- FALLIS, A. M. & G. F. BENNETT
1961. Ceratopogonidae as intermediate hosts for *Haemoproteus* and other parasites. *Mosquito News* 21:21-27, figs.
- IEBRIG-GERTZ, C.
1928. Un Diptère ectoparasite sur un Phasme: *Ceratopogon ixodoides* n. sp. *Ann. Parasit. hum. comp.* 6:284-290, figs.
- FLOCH, H. & E. ABONNENC
- 1942a. Ceratopogonides hematophages de la Guyane Française. *Publ. Inst. Pasteur de la Guyane et du Territ. de L'Inini* 37:1-10, figs.
- 1942b. Ceratopogonides hematophages de la Guyane Française — II. *Publ. Inst. Pasteur de la Guyane et du Territ. de L'Inini* 49:1-5, figs.
- 1942c. Ceratopogonides divers de la Guyane Française — III. *Publ. Inst. Pasteur de la Guyane et du Territ. de L'Inini* 55:1-5, figs.
1949. Sur les Ceratopogonides du Venezuela. Description de deux espèces nouvelles: *Culicoides lichyi* et *Lasiohelea danaisi*. *Publ. Inst. Pasteur de la Guyane et du Territ. de L'Inini* 196:1-5, figs.
- 1950a. Ceratopogonides nouveaux du Venezuela: *Culicoides lichyi* n. sp. et *Lasiohelea danaisi* n. sp. *Bol. Ent. Venezolana* 8:69-75, figs.
- 1950b. Les *Culicoides* de la Guadeloupe. Description de *Culicoides guadeloupensis* n. sp. *Publ. Inst. Pasteur de la Guyane et du Territ. de L'Inini* 203:1-5, figs.
- FORATTINI, O. P.
1953. O *Culicoides trinidadensis* Hoffman 1925 (Diptera, Ceratopogonidae). *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 7:123-126, figs.
- 1954a. Redescrção de *Culicoides flavivenula* Costa Lima 1937 (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 1:135-138, figs.
- 1954b. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) do Brasil Central. *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 8:189-192, figs.
- 1954c. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) do Estado de Goiaz, Brasil. *Folia Clin. et Biol.* 21:315-319, figs.
- 1954d. Redescrção de *Culicoides lopesi* Barretto, 1944, *C. limai* Barretto 1944, e *C. bambusicola* Lutz, 1913 (Diptera, Ceratopogonidae). *Folia Clin. et Biol.* 22:223-232, figs.
1955. Redescrção de *Culicoides guttatus* (Coquillett, 1904) e *C. insignis* Lutz, 1913 (Diptera, Ceratopogonidae). *Folia Clin. et Biol.* 23:103-114, figs.



- 1956a. A new *Culicoides* from Panama (Diptera, Ceratopogonidae). *Proc. Ent. Soc. Washington* 58:35-36, figs.
- 1956b. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) de São Vicente, Estado de São Paulo, Brasil. *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 10:81-84.
1957. *Culicoides* da Região Neotropical (Diptera, Ceratopogonidae). *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 11:161-526, 132 figs.
1958. Nova espécie de *Leptoconops* s. str. da região neotropical (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 8:37-41, figs.
- FORATTINI, O. P. & A. GALVÃO
1955. Redescrição de *Culicoides maruim* Lutz, 1913 e *C. reticulatus* Lutz, 1913 (Diptera, Ceratopogonidae). *Rev. Brasil. Malar. e Doenças Trop.* 7:225-233, figs.
- FORATTINI, O. P. & J. LANE
1955. Sôbre uma *Forcipomyia* que suga Phasmidae (Diptera, Nematocera, Ceratopogonidae). *Rev. Brasil. de Ent.* 4:1-6, figs.
- FORATTINI, O. P. & E. X. RABELLO
1956. As formas imaturas de *Culicoides guyanensis* Floch e Abonnenc, 1942 e de algumas espécies de *Stilobezzia* (Diptera, Ceratopogonidae). *Rev. Brasil. de Ent.* 6:43-49, figs.
1957. Dados sôbre *Dasyhelea* Kieffer, 1911 do Brasil (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 7:241-250.
- FORATTINI, O. P., E. X. RABELLO & D. PATTOLI
1956. Nota sobre a larva e pupa de *Culicoides insignis* Lutz, 1913 (Diptera, Ceratopogonidae). *Rev. Brasil. de Ent.* 4:195-198, figs.
1957. A brief note on breeding places of *Culicoides* in São Vicente, Brazil. *Mosquito News* 17:312-313, figs.
1958. *Culicoides* da Região Neotropical (Diptera, Ceratopogonidae) II. Observações sobre biologia em condições naturais. *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 12:1-52, figs.
1960. Sobre as formas imaturas de *Culicoides reticulatus* Lutz e de *Stilobezzia panamensis* Lane e Forattini (Diptera, Ceratopogonidae). *Rev. Brasil. de Ent.* 9:1-4, figs.
- FOX, I.
1942. The respiratory trumpet and anal segment of the pupae of some species of *Culicoides* (Diptera: Ceratopogonidae). *Puerto Rico Jour. Pub. Health Trop. Med.* 17:412-425, figs.
1946. A review of the species of biting midges or *Culicoides* from the Caribbean Region. *Ann. Ent. Soc. Amer.* 39:248-258, figs.
1947. Two new Central American biting midges or *Culicoides* (Diptera: Ceratopogonidae). *Kuba* 3:90-91, figs.
1948. *Hoffmania*, a new subgenus in *Culicoides* (Diptera: Ceratopogonidae). *Proc. Biol. Soc. Washington* 61:21-28, 1 plate.
1949. Notes on Puerto Rican biting midges or *Culicoides* (Diptera: Ceratopogonidae). *Bull. Brooklyn Ent. Soc.* 44:29-34, figs.
1952. Six new Neotropical species of *Culicoides*. *Ann. Ent. Soc. Amer.* 45:364-368, figs.
1955. A catalogue of the bloodsucking midges of the Americas (*Culicoides*, *Leptoconops*, and *Lasiohelea*) with keys to the subgenera and Nearctic species, a geographic index, and bibliography. *Jour. Agr. Univ. Puerto Rico* 39:214-285.

- FOX, I. & W. A. HOFFMAN
 1944. New neotropical biting sandflies of the genus *Culicoides* (Diptera: Ceratopogonidae). *Puerto Rico Jour. Pub. Health & Trop. Med.* 20:108-111, figs.
- GERRY, B.
 1933. Four new species of Chironomidae from the Greater Antilles. *Psyche* 40:94-97.
- GOELDI, E.
 1905. Os Mosquitos no Pará. *Mem. Mus. Goeldi* 4:1-154, 21 plates.
- GOETGHEBUER, M.
 1920. Ceratopogoninae de Belgique. *Mém. Mus. Roy. d'Hist. Nat. Belgique* 8:1-116, figs.
- GOETGHEBUER, M. & F. LENZ
 1933-1934. Heleidae (Ceratopogonidae). *Fam.* 13a, pp. 1-48, figs. 1-42, plates 1-6 (lfg. 77), 1933; pp. 49-133, figs. 43-177, plates 7-12 (lfg. 78), 1934. In Lindner, E., ed. *Die Fliegen der Palaearktischen Region*, vol. 3. Stuttgart.
- GUTSEVICH, A. Y., A. I. GARCIA & R. GONZALEZ BROCHE
 1969. Resultado del estudio sobre los jejenes hematofagos (Diptera Ceratopogonidae) en Cuba. *Torreia* (n. ser.) 16:1-8.
- HOFFMAN, W. A.
 1926. Two new species of American *Leptoconops* (Diptera, Chironomidae). *Bull. Ent. Res.* 17:133-136, 1 fig.
 1939. *Culicoides filariferus*, new species. *Puerto Rico Jour. Pub. Health & Trop. Med.* 15:172-174.
- HOGUE, C. L. & W. W. WIRTH
 1968. A new Central American sand fly breeding in crab holes (Diptera, Ceratopogonidae). *Los Angeles Co. Mus. Contr. Sci.* 152:1-7, figs.
- ICHES, L.
 1906. Sobre cinco Dipteros nuevos del Chaco Austral. *Bol. Ministr. Agr. Argentine Rep.* 6:262-273.
- INGRAM, A. & J. W. S. MACFIE
 1931. Ceratopogonidae. *Diptera of Patagonia & S. Chile*, part II, fasc. 4, pp. 155-232, 34 figs. London.
- JOHANSEN, O. A.
 1907. Some new species of Kansas Chironomidae. *Univ. Kansas Sci. Bull.* 4:109-112.
 1927. The genus *Stenozenus* (Chironomidae, Diptera). *Ent. News* 38:70-72.
 1938. New species of Nemocera from Puerto Rico. *Jour. Agr. Univ. Puerto Rico* 22:219-225.
 1943a. Two new species of American Ceratopogonidae (Diptera). *Ann. Ent. Soc. Amer.* 36:761-762.
 1943b. A generic synopsis of the Ceratopogonidae (Heleidae) of the Americas, a bibliography, and a list of the North American species. *Ann. Ent. Soc. Amer.* 36:763-791, 3 plates.
 1950. A new *Pterobosca* from Florida with a key to American species. *Florida Ent.* 33:141-144.
 1951. Notes on North American species of *Pterobosca*. *Florida Ent.* 34:116-117.



JOHNSON, C. W.

1908. The Diptera of the Bahamas, with notes and description of one new species. *Psyche* 15:69-80.

JONES, R. H. & W. W. WIRTH

1958. New records, synonymy, and species of Texas *Culicoides* (Diptera, Heleidae). *Jour. Kansas Ent. Soc.* 31:81-91.

KETTLE, D. S.

1962. The bionomics and control of *Culicoides* and *Leptoconops* (Diptera, Ceratopogonidae = Heleidae). *Ann. Rev. Ent.* 7:401-418.
1969. The ecology and control of blood-sucking ceratopogonids. *Acta Tropica* 26:235-248.

KHALAF, K. T.

1952. The *Culicoides* of the Wichita Refuge, Oklahoma. Taxonomy and seasonal incidence (Diptera, Heleidae). *Ann. Ent. Soc. Amer.* 45:348-358, figs.
1954. The speciation of the genus *Culicoides* (Diptera, Heleidae). *Ann. Ent. Soc. Amer.* 47:34-51.

KIEFFER, J. J.

1899. Description d'un nouveau genre et tableau des genres européens de la famille des Chironomides (Dipt.). *Bull. Soc. Ent. France* 1899:66-70, 1 fig.
1906. Diptera. Fam. Chironomidae. Fasc. 42, 78 pp., 4 plates. In Wytzman, P., ed., *Genera Insectorum*. Bruxelles.
1909. Considérations sur le genre *Stenozenus* Coq. avec description d'une espèce nouvelle. *Ann. Mus. Nat. Hungarici* 7:46-49.
1911a. Description de nouveaux Chironomides de l'Indian Museum de Calcutta. *Rec. Indian Mus.* 6:113-177, 2 plates.
1911b. Nouvelles descriptions de Chironomides obtenus d'éclosion. *Bull. Soc. Hist. Nat. Metz* 27:1-60.
1913. Nouvelle étude sur les Chironomides de l'Indian Museum de Calcutta. *Rec. Indian Mus.* 9:119-197, 2 plates.
1914. South African Chironomidae. *Ann. S. African Mus.* 10:259-270.
1917. Chironomides d'Amérique conservés au Musée National Hongrois de Budapest. *Ann. Mus. Nat. Hungarici* 15:292-364, figs.
1918. Chironomides d'Afrique et d'Asie conservés au Muséum National Hongrois de Budapest. *Ann. Mus. Nat. Hungarici* 16:31-136, figs.
1919a. Chironomides d'Europe conservés au Musée National Hongrois de Budapest. *Ann. Mus. Nat. Hungarici* 17:1-160, figs.
1919b. Observations sur les Chironomides (Dipt.) décrits par J. R. Malloch. *Bull. Soc. Ent. France* 1919:191-194.
1921a. Chironomides de l'Afrique équatoriale (1re partie). *Ann. Soc. Ent. France* 90:1-56, 2 plates.
1921b. Sur quelques Diptères piqueurs de la tribu des Ceratopogoninae. *Arch. Inst. Pasteur Afrique du Nord* 1:107-115, figs.
1921c. Description de quelques Chironomides exotiques. *Ann. Soc. Sci. Bruxelles* 40:181-186.
1925a. Diptères (Nematocères piqueurs): Chironomidae Ceratopogoninae. *Faune de France* 11:1-139, figs.
1925b. Nouveaux genres et nouvelles espèces de Chironomides piqueurs.

Arch. Inst. Pasteur d'Algérie 3:405-430, figs.

1925c. Chironomides de la République Argentine. *Ann. Soc. Sci. Bruxelles* 44:73-92.

KNAB, F.

1914. Ceratopogoninae sucking the blood of caterpillars. *Proc. Ent. Soc. Washington* 16:63-66.

1915. New Ceratopogoninae from Peru. *Ins. Ins. Mens.* 3:109-111.

LANE, J.

1944. As espécies Neotrópicas do gênero *Clinohelca* Kieffer, 1917 (Diptera, Ceratopogonidae). *Rev. Ent.* 15:249-261, figs.

1945. Redescrção de Ceratopogonídeos Neotrópicos (Diptera: Ceratopogonidae). *Rev. Ent.* 16:357-372, figs.

1946a. New Neotropical Ceratopogonidae (Heleidae) (Diptera, Nematocera). *Rev. Ent.* 17:202-215, figs.

1946b. Duas espécies novas de Ceratopogonídeos Brasileiros (Diptera Ceratopogonidae (Heleidae). *Liv. Hom. R. Ferreira Almeida*, (art. 22), pp. 219-225, figs.

1947a. Espécies Brasileiras de *Stilobezzia* (Dipt. Ceratopogonidae) e *Zygoneura stonei* nov. nom. (Dipt. Mycetophilidae). *Rev. Ent.* 18:197-214, figs.

1947b. Novas espécies de *Palpomyia* do Brasil (Diptera, Ceratopogonidae). *Rev. Ent.* 18:438-447, figs.

1947c. A biologia e taxonomia de algumas espécies dos grupos *Forcipomyia* e *Culicoides* (Diptera, Ceratopogonidae) (Heleidae). *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 1:159-170, figs.

1948. Novos Ceratopogonídeos do Brasil (Diptera, Ceratopogonidae (Heleidae). *Arq. Fac. Hig. Saúde Púb. Univ. São Paulo* 1:225-239, 3 plates.

1956a. On "*Paryphoconus*" and "*Stenoxenus*" (Diptera, Ceratopogonidae). *Rev. Brasil. Biol.* 61:299-308, figs.

1956b. The genus *Macfiehelea* Lane, 1946. *Rev. Brasil. Biol.* 16:435-438.

1958. On Neotropical *Bezzia*. *Rev. Brasil. Ent.* 8:25-36, figs.

1960. Additional data on *Palpomyia* (Diptera, Ceratopogonidae). *Rev. Brasil. Biol.* 20:381-389, figs.

1961a. Insecta Amapaensia, Diptera: Ceratopogonidae and Anisopodidae. *Studia Ent.* 4:449-452, figs.

1961b. The genera *Stenoxenus* and *Paryphoconus* in the Neotropics (Diptera: Ceratopogonidae). *Studia Ent.* 4:452-458, figs.

1961c. Further notes on Neotropical Ceratopogonidae (Diptera). *Rev. Brasil. Biol.* 21:37-44, figs.

LANE, J. & M. P. DURET

1954. *Clinohelca* da Argentina e do Brasil (Diptera, Ceratopogonidae). *Dusenía* 5:247-254, figs.

LANE, J. & O. P. FORATTINI

1956. Neotropical *Stilobezzia* Kieffer, 1911. I. Nine new Panamanian species (Diptera, Nematocera, Ceratopogonidae). *Rev. Brasil. Malar. Doenças Trop.* 8:207-226, figs.

1958. Neotropical *Stilobezzia*. 2. Fourteen new species, chiefly from Panama (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 8:203-224, figs.



1961. Neotropical *Stilobezzia* Kieffer, 1911. 3. Key for the adults of this genus and description of one species (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 10:83-94, figs.
- LANE, J., O. P. FORATTINI & E. X. RABELLO
1955. Biologia e espécies novas de *Palpomyia* e *Stilobezzia* (Diptera, Nematocera, Ceratopogonidae). *Dusenía* 6:81-88, figs.
- LANE, J., & W. W. WIRTH
1961. A new Neotropical species of *Dicrohelea* (Diptera, Ceratopogonidae). *Rev. Brasil. Ent.* 10:81-82.
1964. The biting midge genus *Monohoelea* Kieffer in the Neotropical Region (Diptera, Ceratopogonidae). *Studia Ent.* 7:209-236, figs.
- LATREILLE, P. A.
1809. *Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita*. Vol. 4, 399 pp., 4 plates. Paris and Strasbourg.
- LIMA, A. DA COSTA
1928. Ceratopogoninos ectoparasitos de phasmídeos. *Mem. Inst. Oswaldo Cruz, Suppl.* 3:84-85, 2 plates.
1937a. Chave das espécies de *Culicoides* da região neotropical (Diptera: Ceratopogonidae). *Mem. Inst. Oswaldo Cruz* 32:411-422, figs.
1937b. Primeira espécie americana do género *Pterobosca* (Diptera: Ceratopogonidae). *Mem. Inst. Oswaldo Cruz* 32:615-616, 2 plates.
- LEWIS, D. J.
1958. Some observations on Ceratopogonidae and Simuliidae (Diptera) in Jamaica. *Ann. Mag. Nat. Hist.* (13)1:721-732, figs.
- LINLEY, J. R.
1965a. Descriptions of the pupae of *Culicoides barbosai*, *C. insignis*, and *C. borinqueni* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 58:57-63, figs.
1965b. The ovarian cycle and egg stage in *Leptoconops (Holoconops) bequaerti* (Kieff.) (Diptera, Ceratopogonidae). *Bull. Ent. Res.* 56:37-56, figs.
- LINLEY, J. R. & J. B. DAVIES
1971. Sandflies and tourism in Florida and the Bahamas and Caribbean area. *Jour. Econ. Ent.* 64:264-278.
- LINLEY, J. R. & D. S. KETTLE
1964. A description of the larvae and pupae of *Culicoides furens* Poey, and *Culicoides hoffmani* Fox (Diptera: Ceratopogonidae). *Ann. Mag. Nat. Hist.* (Ser. 13) 7: 129-149, figs.
- LOEW, H.
1866. Diptera Americae septentrionalis indigena. Centuria sexta. *Berlin. Ent. Ztschr.* 9: 127-186.
- LUTZ, A.
1913. Contribuição para o estudo das Ceratopogoninas hematofagas do Brasil. Parte sistemática. Segunda Memória. *Mem. Inst. Oswaldo Cruz* 5: 45-73, 3 plates.
1914. Contribuição para o conhecimento das Ceratopogoninas do Brasil. Terceira memória. Aditamento terceiro e descrição de espécies que não sugam sangue. *Mem. Inst. Oswaldo Cruz* 6: 81-99, 2 plates, 2 figs.

LYNCH ARRIBÁLZAGA, F.

1893. Dipterologia Argentina (Chironomidae). *Bol. Acad. Nac. Cienc. Córdoba* 13: 211-258.

MACFIE, J. W. S.

1932. Ceratopogonidae from the wings of dragonflies. *Tijdschr. Ent.* 75: 265-283, figs.
1934. Report on a collection of Ceratopogonidae from Malaya. *Ann. Trop. Med. Parasit.* 28: 177-194, figs.
- 1935a. Ceratopogonidae (Dipt.) from the River Amazon. *Stylops* 4:49-56, fig.
- 1935b. A correction to my paper entitled "Ceratopogonidae (Dipt.) from the river Amazon". *Stylops* 4: 89.
- 1935c. A new ceratopogonid (Dipt.) from British Guiana. *Stylops* 4: 265, 1 fig.
1936. Four species of Ceratopogonidae (Diptera) from the wings of insects. *Proc. Roy. Ent. Soc. London* (B) 5:227-230, 1 fig.
- 1937a. Ceratopogonidae from Trinidad. *Ann. Mag. Nat. Hist.* (10)20: 1-18, fig.
- 1937b. Notes on Ceratopogonidae (Diptera). *Proc. R. Ent. Soc. London* (B) 6: 111-118.
1938. Notes on Ceratopogonidae (Diptera). *Proc. Roy. Ent. Soc. London* (B) 7: 157-166, figs.
- 1939a. A report on a collection of Brazilian Ceratopogonidae (Dipt.). *Rev. Ent.* 10: 137-219, figs.
- 1939b. A key to the species of Ceratopogonidae akin to *Macropeza* Mg. (Diptera). *Trans. Roy. Ent. Soc. London*, 89: 1-12, figs., 1 plate.
- 1939c. *Ruvenzori Expedition 1934-5*. Vol. 1, n.º 5. Ceratopogonidae. With an appendix by Dr. B. De Meillon. pp. 81-107, figs. British Museum, London.
- 1940a. A report on a collection of Ceratopogonidae (Diptera) from British Guiana. *Ent. Mo. Mag.* 76: 23-32, figs.
- 1940b. Ceratopogonidae (Diptera) from North-eastern Brazil. *Proc. Roy. Ent. Soc. London* (B) 9: 73-79, figs.
- 1940c. Ceratopogonidae (Diptera) from British Guiana and Trinidad. *Proc. Roy. Ent. Soc. London* (B) 9:179-195, figs.
- 1940d. The genera of Ceratopogonidae. *Ann. Trop. Med. Parasit.* 34: 13-30.
- 1940d. *Forcipomyia furcifera*, sp. n. (Dipt. Ceratopogonidae). *Rev. Ent.* 11: 920-922, figs.
1943. A new species of *Dasyhelea* (Diptera) from the Bahama Islands. *Proc. Roy. Ent. Soc. London* (B) 12: 119-120, figs.
1944. Ceratopogonidae collected in Trinidad from cacao flowers. *Bull. Ent. Res.* 35: 297-300, figs.
1948. Some species of *Culicoides* (Diptera, Ceratopogonidae) from the state of Chiapas, México. *Ann. Trop. Med. Parasit.* 42: 67-87, figs.
1949. Notes on Ceratopogonidae. *Proc. Roy. Ent. Soc. London* (B) 18: 109-115, figs.
1953. Ceratopogonidae from Costa Rica (Diptera). *Beitr. zur Ent.* 3: 95-105, figs.



MALLOCH, J. R.

- 1915a. The Chironomidae, or midges, of Illinois, with particular reference to the species occurring in the Illinois River. *Bull. Illinois St. Lab. Nat. Hist.* 10: 275-543, 24 plates.
- 1915b. Some additional records of Chironomidae for Illinois and notes on other Illinois Diptera. *Bull. Illinois St. Lab. Nat. Hist.* 11: 305-363, 5 plates.
1923. A new species of *Forcipomyia* from the eastern United States (Diptera: Ceratopogonidae). *Ent. News* 34: 4-5.

MATTA, J. F.

1967. A new species of *Culicoides* (Diptera, Ceratopogonidae) from Honduras. *Florida Ent.* 50: 75-77, figs.

MAYER, K.

1934. Die Metamorphose der Ceratopogonidae (Dipt.). *Arch. für Naturges.* (Leipzig) 3: 205-288, figs.
1937. Beobachtungen über blutsaugende Ceratopogoniden. *Arb. Morph. Tax. Ent. Berlin* 4: 231-234, figs.
1938. Ceratopogoniden als Phasmenparasiten. *Rev. Ent.* 9: 13-15.
1952. Die *Macropeza* Gruppe der Heleiden (Diptera: Heleidae). *Beitr. zur Ent.* 2: 582-585, figs.
1959. Die Puppen brasilianischer Heleiden (Diptera). *Deutsche Ent. Ztshr.* (n. F.) 6: 230-233, figs.

MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau.* 40 pp. Paris.
1818. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* 1: 1-333, 11 plates.

MESSERSMITH, D. H.

1972. A new species of *Culicoides* from Colombia (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Washington* 74: 165-169, figs.

MIRSA, M., & I. ORTIZ

1952. Estudios en *Culicoides* (Diptera Ceratopogonidae) 10. Sobre la hembra de *Culicoides dominicii* Ortiz, 1951. *Bol. Lab. Clin. Luis Razetti* 12: 475-478, figs.

NOÉ, G.

1905. Un nuovo genere appartenente alla famiglia Chironomidae. *Rend. Accad. Lincei* 14: 114-120, figs.

ORTIZ, I.

- 1950a. Estudios en *Culicoides*. I. Sobre la hembra de *Stilobezzia maculata* Lane, 1947 (Diptera, Ceratopogonidae). *Rev. Sanid. Asist. Social* 15: 429-430, 1 plate.
- 1950b. Estudios en *Culicoides*. II. Diptera, Ceratopogonidae. *Culicoides lanei* n. sp. de Panama. *Rev. Sanid. Asist. Social* 15: 431-433, 1 plate.
- 1950c. Estudios en *Culicoides*. IV. Revisión de las especies americanas del sub-género *Hoffmania* Fox 1948, con la descripción de dos nuevas especies. *Rev. Sanid. Asist. Social* 15: 437-460, figs.
- 1950d. Estudios en *Culicoides*. V. Informes sobre una nueva especie y lista de los machos cuyas genitalias son conocidas. *Rev. Sanid. Asist. Social* 15: 461-465.

- 1950e. Descripción de *Stilobezzia glauca venezuelensis* n. vr. y *Monohela mayeri* n. sp. *Mem. Soc. Cien. Nat. La Salle* 10: 199-204, figs.
- 1951a. Estudios en *Culicoides* (Diptera, Ceratopogonidae). 6. *Culicoides bricenoi* n. sp. *Bol. Lab. Clin. Luis Razetti* 11: 442-448, figs.
- 1951b. Estudios en *Culicoides* (Diptera, Ceratopogonidae). 9. Sobre los caracteres diferenciales entre *Culicoides paraensis* (Goeldi, 1905), *C. stelleri* (Coquillett, 1901), y *C. lanei* (Ortiz, 1950). Descripción de cuatro nuevas especies con la redescrición de algunas otras poco conocidas. *Rev. Sanid. Asist. Social* 16: 573-591, figs.
- 1951c. Estudios sobre *Culicoides* (Diptera, Ceratopogonidae). VII. *Culicoides dominicii* n. sp. y clave para el reconocimiento de las especies venezolanas. *Nov. Cient. Zool. Ser. Mus. Hist. Nat. La Salle*, 5: 1-12, figs.
- 1952a. Los insectos del genero *Leptoconops* Skusc, 1889 en Venezuela. Descripción de una nueva especie: *Leptoconops (L.) venezuelensis* y redescrición de la hembra de *Leptoconops (L.) torrens* (Townsend, 1893). *Rev. Sanid. Asist. Social* 17: 163-171, figs.
- 1952b. Nota sobre la presencia en Venezuela de un nuevo Ceratopogonido hematofago: *Lasiohelea styliifer* (Lutz, 1913). *Rev. Sanid. Asist. Social* 17: 241-244, figs.
- 1952c. Apuntes en Ceratopogonidae (Diptera Nematocera). 3. Sobre una nueva especie de *Monohela* Kieffer 1917 (*Monohela ocumare* n. sp.). *Rev. Sanid. Asist. Social* 17: 253-256, figs.
- 1952d. Nota sobre la presencia de *Culicoides eublepharus* Macfie, 1948 (Diptera, Ceratopogonidae) en Venezuela — descripción de la armadura genital del macho. *Acta Cient. Venezolana* 3:129, figs.
- 1953a. Nueva contribucion al conocimiento de los caracteres morfologicos externos de las hembras americanas del genero *Culicoides* Latr. (Diptera, Ceratopogonidae) con una espermatheca. Descripción de dos nuevas especies de Venezuela: *Culicoides transferrans* (*C. eublepharus* Ortiz 1952) y *Culicoides mirsae*. *Rev. Sanid. Asist. Social* 18:797-806, figs.
- 1953b. Sobre una nueva especie del genero *Culicoides* (Diptera Ceratopogonidae) proxima de *C. horticola* Lutz 1913. *Rev. Sanid. Asist. Social* 18:807-812, figs.
1954. Sobre dos nuevos dipteros hematofagos del genero *Culicoides* (Nematocera, Ceratopogonidae). *Arch. Venezolanos Patol. Trop. y Parasit. Med.* 2:221-226.
1956. Sobre un pequeño díptero de la region del Auyantepuy. *Bol. Venez. Lab. Clin.* 1:93-96, figs.
1957. Nuevos representantes hematofagos de los generos *Culicoides* (Diptera: Ceratopogonidae) y *Simulium* (Diptera: Simuliidae) de Venezuela. *Bol. Venezol. Lab. Clin.* 2:161-168, figs.
1958. Sobre *Johannseniella fluvialis* Lutz, 1914, y *Culicoides pachymerus* Lutz, 1914 (Diptera: Ceratopogonidae). *Bol. Venez. Lab. Clin.* 3:37-43.
1961. Descripción de una nueva especie de *Culicoides* (*C. forattinii* sp. n.) del subgenero *Oeacta* Poey, 1851, de Venezuela. *Rev. Brasil. Ent.* 10:211-215.
1963. Estudios en Ceratopogonidae (Diptera: Nematocera). *Rev. Venez. Sanid. Asist. Soc.* 28:307-310.
1968. Los Dipteros hematofagos del género *Culicoides* en Venezuela (Diptera, Ceratopogonidae). Part I. Hembras da una sola espermateca con la descripción de una nueva especie (*C. rodriguezi* n. sp.). *Rev. Inst. Nac. Higiene* 1:65-71.



1969. Sistematización agrupativa, identificación específica de los *Culicoides* (Diptera: Ceratopogonidae) Neotropicales por medio de nuevo tipo simplificado de clave. *Rev. Inst. Nac. Higiene Caracas* 11:31-44.
- ORTIZ, I. & L. A. LEON
 1955. Los *Culicoides* (Diptera: Ceratopogonidae) de la República del Ecuador. *Bol. Inf. Cient. Nac.* 7:564-590, figs.
- ORTIZ, I. & M. MIRSA
 1950. Estudios en *Culicoides* (Diptera, Ceratopogonidae). 8 — Sobre una nueva especie del grupo *haematopotus*: *Culicoides venezuelensis* n. sp. *Arch. Venezolanas Patol. Trop. y Parasit. Med.* 2:137-144, figs.
 1951. Estudios en *Culicoides*. Descripción de dos nuevas especies: *Culicoides avilaensis* y *C. discrepans*, y del macho de *C. leopoldoi* Ortiz, 1951. Redescrpcion de *C. limai* Barretto, 1944, *C. baueri* Hoffman 1925, *C. lichi* Floch & Abonnenc, 1949, y *C. pusillus* Lutz, 1913. *Rev. Sanid. Asist. Social* 16:593-605, figs.
 1952. *Culicoides* de Venezuela — Redescrpcion de 10 especies con la descripción de algunos sexos no conocidos. *Rev. Sanid. Asist. Social* 17:257-279, figs.
- PAINTER, R. H.
 1926. The biology, immature stages, and control of the sandflies (biting Ceratopogonidae) at Puerto Castilla, Honduras. *Ann. Rept. Med. Dept. United Fruit Co.* 1926:245-262, 1 plate.
- PHILIPPI, R. A.
 1865. Aufzählung der chilenischen Dipteren. *Verh. K.-k. Zool.-Bot. Gesell. Wien*, 15:595-782, 7 plates.
- POEY, F.
 1851. *Memorias sobre la historia natural de las Isla de Cuba, acompañadas de sumarios latinos y extractos en francés.* Vol. 1, 463 pp., 34 plates. Habana.
- REMM, H.
 1971. On the fauna of Heleidae of South Primorye (Ussuri Land), pp. 182-220. In: *Living Nature of the Far East*, 239 pp. Akad. Nauk Estonskoi S.S.R. In Russian.
- ROBACK, S. S.
 1957. A new *Palpomyia* from Peru (Diptera: Heleidae). *Notulae Nat. Acad. Nat. Sci. Philadelphia* 297, 2 pp., figs.
- ROMAÑA, C. & P. WYGODZINSKY
 1950. Acerca de la transmisión de *Mansonella ozzardi* (Manson) (*Filaria tucumana* Biglieri y Araoz). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3:29-34.
- ROOT, F. M. & W.A. HOFFMAN
 1937. The North American species of *Culicoides*. *Amer. Jour. Hyg.* 25: 150-176, 8 plates.
- SAUNDERS, L. G.
 1924. On the life history and the anatomy of the early stages of *Forcipomyia* (Diptera, Nemat., Ceratopogoninae). *Parasit.* 16:164-213, 3 plates.
 1925. On the life history, morphology and systematic position of *Apelma* Kieff. and *Thyridomyia* n. g. (Diptera, Nemat. Ceratopogoninae). *Parasit.* 17:252-277, 1 plate.



1956. Revision of the genus *Forcipomyia* based on characters of all stages (Diptera, Ceratopogonidae). *Canad. Jour. Zool.* 34:657-705, figs.
1959. Methods for studying *Forcipomyia* midgcs, with special reference to cacao-pollinating species (Diptera, Ceratopogonidae). *Canad. Jour. Zool.* 37:33-51, figs.
1964. New species of *Forcipomyia* in the *Lasiohelea* complex described in all stages (Diptera, Ceratopogonidae). *Canad. Jour. Zool.* 42:463-482, figs.
- SAUNDERS, L. G. & G. F. BOWMAN
1956. *Cacao pollination in Costa Rica*. Banco de Costa Rica, San José, Costa Rica. 13 pp.
- SAY, T.
1825. *American entomology, or descriptions of the insects of North America*. Vol. 2, 121 pp. plates 19-36. Philadelphia.
- SÉGUY, E.
1941. Quelques Cératopogonides vulnérantes parasites des insectes. *Rev. Française d'Ent.* 8:82-88, figs.
- SHANNON, R. C. & E. DEL PONTE
1927. Cuatro notas sobre especies nuevas de Dipteros, Nematoceros, hematófagos o no, de la Republica Argentina. *Rev. Inst. Bacteriologico* 4: 724-736.
- SKUSE, F. A. A.
1889. Diptera of Australia. Part VI. The Chironomidae. *Proc. Linn. Soc. New South Wales*, (2)4:215-311, plates 11-14.
- THOMSEN, L. C.
1937. Part. V. Ceratopogonidae. pp. 57-80, plates 10-18. In: Johannsen, O. A. & L. C. Thomsen, Aquatic Diptera. Parts IV and V. *Cornell Univ. Agr. Exp. Sta. Mem.* 210:1-80, 18 plates.
- TOKUNAGA, M. & E. K. MURACHI
1959. Diptera: Ceratopogonidae. pp. 103-434, figs. In: B. P. Bishop Mus. *Insects of Micronesia*, 12:1-434, figs. Honolulu.
- TOWNSEND, C. H. T.
1893. An interesting blood-sucking gnat of the family Chironomidae. *Psyche* 6:369-371, 1 plate.
- VARGAS, L.
1944. *Culicoides diabolicus* en Mexico. Caracteres del macho. *Rev. Inst. Salub. Enf. Trop.* 5:163-169, figs.
1945. Nota sobre Ceratopogonidos y *Culicoides*. *Rev. Inst. Salub. Enf. Trop.* 6:41-49, figs.
1949. Lista de los *Culicoides* del mundo (Diptera, Heleidae). *Rev. Soc. Mexicana Hist. Nat.* 10:191-218.
- 1953a. *Beltranmyia* n. subg. de *Culicoides* (Insecta: Heleidae). *Rev. Inst. Salub. Enf. Trop.* 13:33-36.
- 1953b. *Culicoides wirthomyia* n. sp. y *Culicoides stigmalis* Wirth 1952 (Insecta, Diptera). *Rev. Inst. Salub. Enf. Trop.* 13:227-233, figs.
1954. Dos nuevas especies de *Culicoides* Mexicanos. *Rev. Inst. Salub. Enf. Trop.* 41:25-32, figs.



1955. *Culicoides neghmei* n. sp. *Bol. Lab. Clin. Luis Razetti*, 43-44:673-676, figs.
1958. Los *Culicoides* o jejenes de Mexico (Insecta: Diptera); claves graficas para la determinacion de los subgeneros. *Rev. Inst. Salub. Enf. Trop.* 18:75-80, figs.
1960. The subgenera of *Culicoides* of the Americas (Diptera, Ceratopogonidae). *Rev. Biol. Trop.* 8:35-47, figs.
1969. Los *Culicoides* como transmisores de enfermedades. *Gaceta Medica de Mexico* 99:782-787.
- VARGAS, L. & A. GARZA
1959. Nueva lista de *Culicoides* del mundo (Insecta: Diptera). *Rev. Inst. Salub. Enf. Trop.* 19:203-244, figs.
- VARGAS L. & W. W. WIRTH
1955. *Culicoides blantoni* n. sp. (Diptera, Heleidae). *Rev. Inst. Salub. Enf. Trop.* 15:33-37, figs.
- WALKER, R.
1856. Diptera. Vol. 1, pp. 415-474. In: Saunders, W. W., ed. *Insecta Saundersiana*. London.
- WEYENBERGH, H.
1883. Die Gattung *Didymophleps* m. *Stett. Ent. Zeit.* 1883:108-111.
- WILLIAMS, R. W.
1964. Observations on habitats of *Culicoides* larvae in Trinidad, W. I. (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 57:462-466.
- WILLISTON, S. W.
1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896:253-446, plates 8-14.
1900. Supplement (part), pp. 217-248. In Godman, F. D. & O. Salvin, eds. *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, vol. 1, 378 pp., 6 plates. London.
- WINNERTZ, J.
1852. Beitrag zur Kenntniss der Gattung *Ceratopogon* Meigen. *Linnaea Ent.* 6:1-80, 8 plates.
- WIRTH, M. O., W. W. WIRTH & F. S. BLANTON
1968. Plant materials as breeding places of Panama *Culicoides* (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Washington* 70:132.
- WIRTH, W. W.
1951. A new biting midge of the genus *Leptoconops* from Florida, with new records of other American species (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 53:281-284, figs.
- 1952a. Los insectos de las Islas Juan Fernandez. 7. Heleidae and Tendipedidae. *Rev. Chilena Ent.* 2:87-104, figs.
- 1952b. The status of the genus *Parabezzia* Malloch (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 54:22-26, figs.
- 1952c. The immature stages of two species of Florida salt marsh sand flies (Diptera, Heleidae). *Florida Ent.* 35:91-100, figs.



- 1952d. Two new species of anthropophilic *Culicoides* from Guatemala (Diptera: Heleidae). *Jour. Parasit.* 38:245-247, figs.
- 1952e. The Heleidae of California. *Univ. California Pubs. Ent.* 9:95-266, figs.
- 1953a. Biting midges of the heleid genus *Stilobezzia* in North America. *Proc. U. S. Nat. Mus.* 103:57-85, figs.
- 1953b. American biting midges of the heleid genus *Monohalea*. *Proc. U. S. Nat. Mus.* 103:135-153, figs.
- 1955a. Report on a collection of biting midges of the genus *Culicoides* from Guatemala (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 57:109-117, figs.
- 1955b. Three new species of *Culicoides* from Texas (Diptera: Heleidae). *Jour. Washington Acad. Sci.* 45:355-359, figs.
- 1955c. The biting midge genera *Psychophaena* and *Tetrachora* of Philippi 1865. *Rev. Chilena Ent.* 4:233-235.
- 1956a. The heleid midges involved in the pollination of rubber trees in America. *Proc. Ent. Soc. Washington* 58:241-250, figs.
- 1956b. New species and records of biting midges ectoparasitic on insects (Diptera, Heleidae). *Ann. Ent. Soc. Amer.* 49:356-364, figs.
1957. Los insectos de las Islas Juan Fernandez. 32. Heleidae and Tendipedidae (Diptera) (Supplementary Report). *Rev. Chilena Ent.* 5:399-401.
- 1959a. *Pachyhelca*, a new genus of American Ceratopogonidae related to *Palpomyia* (Diptera). *Bull. Brooklyn Ent. Soc.* 54:50-52.
- 1959b. New species and records of Heleidae from Brazil. *Deutsche Ent. Ztschr.* 6:234-237, figs.
- 1960a. The genus *Pellucidomyia* Macfie (Diptera, Ceratopogonidae). *Bull. Brooklyn Ent. Soc.* 55:1-3.
- 1960b. The correct status of *Cotocripus caridei* Brèthes, a South American biting midge (Diptera: Ceratopogonidae). *Proc. Washington Ent. Soc.* 62:44.
1962. A reclassification of the *Palpomyia-Bezzia-Macropoeza* groups, and a revision of the North America Sphaeroniini (Diptera, Ceratopogonidae). *Ann. Ent. Soc. Amer.* 55:272-287, figs.
1963. New synonyms in American *Culicoides* (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Washington* 65:68.
- 1965a. Two new species of *Macrurohelea* from Chile. *Pan-Pacific Ent.* 41:46-50, figs.
- 1965b. A revision of the genus *Parabezzia* Malloch (Diptera, Ceratopogonidae). *Proc. Ent. Soc. Washington* 67:215-230, figs.
- 1965c. A new *Johannsenomyia* from Brazil (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Washington* 67:4.
1966. A New Jamaican blood-sucking midge from lacewings (Diptera, Ceratopogonidae). *Proc. Ent. Soc. Washington* 68:29-32, figs.
1969. New species and records of Galapagos Diptera. *Proc. California Acad. Sci.* (4)36:571-594, figs.
1970. The Neotropical *Forcipomyia* midges of the subgenus *Thyridomyia* Saunders (Diptera, Ceratopogonidae). *Studia Ent.* 13:429-440, figs.



1971. A review of the "stick-ticks", Neotropical biting midges of the *Forcipomyia* subgenus *Microhelea* parasitic on walking stick insects (Diptera: Ceratopogonidae). *Ent. News* 82:229-245, figs.
1972. The Neotropical *Forcipomyia* (*Microhelea*) species related to the caterpillar parasite *F. fuliginosa* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 65:564-577.

WIRTH, W. W. & W. R. ATCHLEY

1973. A review of the North American *Leptoconops* (Diptera: Ceratopogonidae). *Grad. Stud. Texas Tech. Univ.* 5:1-57, figs.

WIRTH, W. W. & BLANTON

- 1953a. Studies in Panama *Culicoides* (Diptera: Heleidae): I, Descriptions of six new species. *Jour. Washington Acad. Sci.* 43:69-76, figs.
- 1953b. Studies in Panama *Culicoides* (Diptera, Heleidae). II. — Descriptions of six additional new species. *Jour. Parasit.* 39:229-236, figs.
- 1953c. Studies in Panama *Culicoides* (Diptera, Heleidae). III. — A new species related to *phlebotomus* (Williston). *Ent. News* 64:113-120, figs.
- 1955a. Studies in Panama *Culicoides* (Diptera, Heleidae). IV. Description of three new species. *Bull. Brooklyn Ent. Soc.* 50:100-106, figs.
- 1955b. Studies in Panama *Culicoides* (Diptera, Heleidae). V. Descriptions of three new species of the subgenus *Oecacta* Poey. *Bull. Brooklyn Ent. Soc.* 50:121-127, figs.
- 1956a. Studies in Panama *Culicoides* (Diptera, Heleidae). IX. Two new species related to *leoni* Barbosa and *reevesi* Wirth. *Bull. Brooklyn Ent. Soc.* 51:45-52, figs.
- 1956b. Studies in Panama *Culicoides* VII. The species of the *pulicaris* and *cova-garciai* groups (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 58:211-227, figs.
- 1956c. Studies in Panama *Culicoides* (Diptera: Heleidae). VI. The *hylas* group of the subgenus *Hoffmania*. *Jour. Washington Acad. Sci.* 46:95-99, figs.
- 1956d. Redescriptions of four species of Neotropical *Culicoides* of the *debilipalpis* group (Diptera: Heleidae). *Jour. Washington Acad. Sci.* 46:186-190, figs.
- 1956e. A new species of salt-marsh sand fly from Florida, the Bahamas, Panama and Ecuador: its distribution and taxonomic differentiation from *Culicoides furens* (Poey) (Diptera, Heleidae). *Florida Ent.* 39:157-162, figs.
- 1956f. Studies in Panama *Culicoides*. VIII. The Neotropical species of the *guttatus* group of the subgenus *Hoffmania* (Diptera, Heleidae). *Proc. Ent. Soc. Washington* 58:305-326, figs.
1959. Biting midges of the genus *Culicoides* from Panama (Diptera: Heleidae). *Proc. U. S. Nat. Mus.* 109:237-482, figs.
1967. The North American *Culicoides* of the *guttipennis* group (Diptera: Ceratopogonidae). *Florida Ent.* 50:207-232, figs.
- 1968a. A revision of the Neotropical biting midges of the *hylas* group of *Culicoides* (Diptera, Ceratopogonidae). *Florida Ent.* 51:201-215, figs.
- 1968b. A new *Culicoides* species from Guyana (Diptera, Ceratopogonidae). *Florida Ent.* 51:251-252, figs.
- 1969a. North American *Culicoides* of the *pulicaris* group (Diptera: Ceratopogonidae). *Florida Ent.* 52:207-243, figs.



- 1969b. A new Nearctic species of the genus *Paradasyhelea* Macfie (Diptera: Ceratopogonidae). *Pan-Pac. Ent.* 45:97-100, figs.
- 1970a. New genera of Neotropical Ceratopogonidae (Diptera). *Florida Ent.* 53:7-14, figs.
- 1970b. Notes on *Brachypogon* Kieffer (Diptera, Ceratopogonidae), a new species, and two new Neotropical genera of the tribe Ceratopogonini. *Florida Ent.* 53:93-104, figs.
- 1970c. New species of Neotropical *Culicoides* (Diptera: Ceratopogonidae). *Florida Ent.* 53:39-45, figs.
- 1970d. A review of the *Culicoides nigrigenus* group, with two new species (Diptera: Ceratopogonidae). *Ent. News* 81:141-151, figs.
1971. New Neotropical sandflies of the *Culicoides debilipalpis* group (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Washington* 73:34-43, figs.
- 1972a. A new Patagonian biting midge of the genus *Monohclea*, *Isthmohclea* (Diptera: Ceratopogonidae). *Florida Ent.* 55:173-176, figs.
- 1972b. *Lanehelea*, a new Neotropical genus of Sphaeromiini (Diptera: Ceratopogonidae). *Studia Ent.* 15:433-438, figs.

WIRTH, W. W. & M. I. DOW

1971. Studies on the genus *Forcipomyia* III. *Blantonina*, a new subgenus in the *Trichohlela* complex (Diptera: Ceratopogonidae). *Florida Ent.* 54:289-295, figs.
1972. Studies on the genus *Forcipomyia* 4. *Rhynchoforcipomyia*, a new Neotropical subgenus in the *Trichohlela* complex (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 65:862-872, figs.

WIRTH, W. W. & A. A. HUBERT

1960. Ceratopogonidae (Diptera) reared from cacti, with a review of the *copiosus* group of *Culicoides*. *Ann. Ent. Soc. Amer.* 53:639-658, figs.

WIRTH, W. W. & R. H. JONES

1957. The North American subspecies of *Culicoides variipennis* (Diptera, Heleidae). *U. S. Dept. Agr. Tech. Bull.* 1170:1-35, figs.

WIRTH, W. W. & D. J. LEE

1959. The genus *Paradasyhelea* Macfie, with descriptions of two new species from eastern Australia (Ceratopogonidae). *Bull. Brooklyn Ent. Soc.* 54:114-121, figs.

WIRTH, W. W. & V. H. LEE

1967. New species of *Culicoides* from high altitudes in the Colombian Andes (Diptera: Ceratopogonidae). *Proc. U. S. Nat. Mus.* 124:1-22, 2 plates.

WIRTH, W. W. & D. H. MESSERSMITH

1971. Studies on the genus *Forcipomyia*. 1. The North American parasitic midges of the subgenus *Trichohlela* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 64:15-26, figs.

WIRTH, W. W. & N. C. RATANAWORABHAN

1971. Notes on Neotropical *Pellucidomyia* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 64:446-448, figs.
- 1972a. A revision of the tribe Stenoxenini (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 65:1368-1388, figs.



1972b. *Neobezzia*, a new Neotropical biting midge genus of the tribe Sphaeromiini (Diptera: Ceratopogonidae). *Jour. Kansas Ent. Soc.* 45:476-490, figs.

WIRTH, W. W. & R. W. WILLIAMS

1957. The biting midges of the Bermuda Islands, with descriptions of five new species (Diptera: Heleidae). *Proc. Ent. Soc. Washington* 59:5-14, figs.


1964. New species and records of North American *Monohalea* (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Amer.* 57:302-310, figs.

WOKE, P.


1954. Observations on Central American biting midges (Diptera, Heleidae). *Ann. Ent. Soc. Amer.* 47:61-74, figs.

INDEX

abercrombyi Macfie, 1938, Forcipomyia	14.4
acanthodes Macfie, 1940, Bezzia	14.52
Acanthohelea Kieffer, 1917	14.42
acotylus Lutz, 1913, Culicoides	14.27
aculeata Ingram & Macfie, 1931, Palpomyia	14.54
adamsoni Macfie, 1937, Atrichopogon	14.13
aemulus Macfie, 1940, Paryphoconus	14.50
aenipes (Macfie), 1940 (Bezzia), Mallochohelea	14.49
aeria Saunders, 1956, Forcipomyia	14.12
aeronautica Macfie, 1935, Forcipomyia	14.12
aethionotus Wirth & Blanton, 1955, Culicoides	14.36
aitkeni Wirth & Blanton, 1968, Culicoides	14.24
alahialinus Barbosa, 1952, Culicoides	14.27
alambiculatorum Macfie, 1948, Culicoides	14.23
albicoxa Lane & Forattini, 1956, Stilobezzia	14.44
albidiventris Kieffer, 1917 (Sphaeromias), Pachyhelea	14.53
albinensis Ingram & Macfie, 1931, Atrichopogon	14.13
albitarsis Wirth & Ratanaworabhan, 1972, Neobezzia	14.49
albocincta Kieffer, 1917, Stilobezzia	14.44
albomacula Root & Hoffman, 1937, Culicoides	14.37
albopennis Lane, 1944, Clinohelea	14.46
albopicta Ingram & Macfie, 1931, Dasyhelea	14.16
albuquerquei Lane, 1961, Bezzia	14.52
Alluaudomyia Kieffer, 1913	14.38
almeidai Lane, 1946, Palpomyia	14.54
almirantei Wirth & Blanton, 1959	14.27
altivolans Macfie, 1949, Atrichopogon	14.13
alvarezi Ortiz, 1957, Culicoides	14.37
amapaensis Lane, 1961, Paryphoconus	14.50
amazonica Wirth, 1971, Forcipomyia	14.8
amazonica Wirth & Blanton, 1970, Fittkauhelea	14.39
amazonius Macfie, 1935, Culicoides	14.27
americana Kieffer, 1917, Stilobezzia	14.44
annicola (Macfie), 1940 (Bezzia), Neobezzia	14.49
amnigena (Macfie), 1935 (Eukraiohelea), Stilobezzia	14.43
andensis Ingram & Macfie, 1931, Dasyhelea	14.16
andicola Wirth & Lee, 1967, Culicoides	14.20
andinus Wirth & Lee, 1967, Culicoides	14.28



angustipennis Enderlein, 1912, Paryphoconus	14.50
Anilomyia Vargas, 1960	14.19
annulatifera Macfie, 1939, Forcipomyia	14.4
anomalicornis Kieffer, 1917, Paryphoconus	14.50
antefurcatus Wirth & Blanton, 1959, Culicoides	14.37
antennalis (Coquillett), 1901 (Ceratopogon), Stilobezzia	14.44
antequerae (Lynch Arribálzaga), 1893 (Pachyleptus), Heteromyia ..	14.47
antiguensis Saunders, 1956, Forcipomyia	14.9
antunesi Forattini, 1954, Culicoides	14.37
<i>Apelma</i> , authors	14.9
<i>Apelma</i> Kieffer, 1919	14.12
apricans Ingram & Macfie, 1931, Palpomyia	14.54
archboldi Wirth & Blanton, 1970, Culicoides	14.28
arcuatus Wirth & Ratanaworabhan, 1972, Stenoxenus	14.52
argenteola Macfie, 1939, Forcipomyia	14.5
argentina Lane & Duret, 1954, Clinohoelea	14.46
argentinensis Kieffer, 1925, Dasyhelea	14.16
arizonensis Wirth & Hubert, 1960, Culicoides	14.22
arubae Fox & Hoffman, 1944, Culicoides	14.37
assimilis Ingram & Macfie, 1931, Atrichopogon	14.13
ateles (Macfie), 1940 (Bezzia), Nilobezzia	14.50
Atrichopogon Kieffer, 1906	14.13
atrichopogon Lane & Forattini, 1956, Stilobezzia	14.43
attenuata Saunders, 1964, Forcipomyia	14.7
aureus Ortiz, 1951, Culicoides	14.28
auricoma Kieffer, 1917, Atrichopogon	14.13
australis Wirth, 1952, Dasyhelea	14.17
Avaritia Fox, 1955	14.20
avilaensis Ortiz & Mirsa, 1951, Culicoides	14.28
azureus Wirth & Blanton, 1959, Culicoides	14.28
Baeohoelea Wirth & Blanton, 1970	14.38
bahamensis (Johnson), 1908 (Ceratoopogon), Dasyhelea	14.17
bahiensis Barbosa, 1947, Culicoides	14.28
bahuerei, error	14.22
bakeri Vargas, 1954, Culicoides	14.22
balboa Lane & Wirth, 1964, Monohelea	14.40
balsapambensis Ortiz & León, 1954, Culicoides	14.28
bambusicola Lutz, 1913, Culicoides	14.28
banksi Gerry, 1933 (Bezzia), Nilobezzia	14.50
barbosai Wirth & Blanton, 1956, Culicoides	14.28
barrettoi Lane, 1946, Paryphoconus	14.50
barrettoi Lane, 1947, Palpomyia	14.54
barrettoi Lane & Duret, 1954, Clinohoelea	14.46
batesi Lane, 1961, Paryphoconus	14.50
baueri authors, Culicoides	14.22
beckae Wirth, 1953, Stilobezzia	14.44
becquaerti, error	14.2
beebei Fox, 1952, Culicoides	14.19
bejaranoi Duret & Lane, 1955, Heteromyia	14.47
Beltranmyia Vargas, 1953	14.21
benarrochei, error	14.37
bennarochi Ortiz & Mirsa, 1952, Culicoides	14.37
bequaerti (Kieffer), 1925 (Holoconops), Leptoconops	14.2
Bezzia Kieffer, 1899	14.52
bicellii (Lane), 1961 (Nilobezzia), Mallochohelea	14.49
bicinctipes Ingram & Macfie, 1931, Stilobezzia	14.43



<i>bicolor</i> Lutz, 1914, Forcipomyia	14.5
<i>bicolor</i> Saunders, 1956, Forcipomyia	14.12
<i>bifidus</i> Ewen, 1958, Atrichopogon	14.13
<i>bicolor</i> Lane, 1947, Stilobezzia	14.44
<i>bimaculata</i> Lane & Forattini, 1956, Stilobezzia	14.44
<i>bimaculatus</i> Floch & Abonnenc, 1942, Culicoides	14.24
<i>birabeni</i> Cavalieri, 1966, Culicoides	14.28
<i>bispinosa</i> Kieffer, 1917, Stilobezzia	14.44
<i>blanchardi</i> Ichès, 1906 (Cerato-pogon), not Ceratopogonidae	14.56
<i>blantoni</i> (Lane), 1956 (Macfiehelea), Pellucidomyia	14.48
<i>blantoni</i> Lane & Forattini, 1956, Stilobezzia	14.44
<i>blantoni</i> (Lane & Wirth), 1961 (Dicrohelea), Johannsenomyia	14.48
<i>blantoni</i> Lane & Wirth, 1964, Monohelea	14.40
<i>blantoni</i> Vargas & Wirth, 1955, Culicoides	14.21
<i>blantoni</i> Wirth, 1965, Parabezzia	14.42
<i>blantoni</i> Wirth & Ratanaworabhan, 1972, Neobezzia	14.49
<i>blantoni</i> Wirth & Ratanaworabhan, 1972, Stenoxenus	14.52
<i>Blantonia</i> Wirth & Dow, 1971	14.3
<i>boliviensis</i> Kieffer, 1917 (Jenkinsia), Palpomyia	14.54
<i>boliviensis</i> Kieffer, 1917, Palpomyia	14.54
<i>borinquèni</i> Fox & Hoffman, 1944, Culicoides	14.22
<i>Brachypogon</i> Kieffer, 1899	14.38
<i>brachyrhynchus</i> Wirth & Dow, 1972, Forcipomyia	14.10
<i>brasiliae</i> (Lane), 1961 (Macropeza), Neobezzia	14.49
<i>brasilianus</i> Forattini, 1956, Culicoides	14.24
<i>brasiliensis</i> Lane, 1958 (Bezzia), Nilobezzia	14.50
<i>brasiliensis</i> Lane, 1948, Monohelea	14.40
<i>brasiliensis</i> (Lutz), 1913 (Tersesthes), Leptoconops	14.2
<i>brasiliensis</i> Macfie, 1939, Atrichopogon	14.13
<i>brasiliensis</i> Macfie, 1939, Forcipomyia	14.7
<i>brasiliensis</i> Macfie, 1939, Palpomyia	14.54
<i>brasiliensis</i> Macfie, 1939, Stenoxenus	14.52
<i>bredini</i> Wirth & Blanton, 1970, Culicoides	14.37
<i>brevicornis</i> (Kieffer), 1917 (Allobezzia), Bezzia	14.52
<i>brevipalpis</i> (Ingram & Macfie), 1931 (Dasyhelea), Paradasyhelea	14.18
<i>brevipalpis</i> Macfie, 1944, Atrichopogon	14.13
<i>bricenoi</i> Ortiz, 1951, Culicoides	14.37
<i>bromeliae</i> Saunders, 1956, Forcipomyia	14.4
<i>bromelicola</i> (Lutz), 1914 (Cerato-pogon), Forcipomyia	14.9
<i>brunnea</i> Wirth, 1965, Parabezzia	14.42
<i>cacozelus</i> Macfie, 1948, Culicoides	14.23
<i>cactorum</i> Wirth & Hubert, 1960, Dasyhelea	14.17
<i>caerulea</i> Saunders, 1956, Forcipomyia	14.3
<i>calcarata</i> (Coquillett), 1905 (Cerato-pogon), Forcipomyia	14.4
<i>caliginosa</i> Ingram & Macfie, 1931	14.5
<i>caliginosella</i> Wirth, nom. nov., Forcipomyia	14.5
<i>callangana</i> Kieffer, 1917, Palpomyia	14.54
<i>Caloforcipomyia</i> Saunders, 1956	14.3
<i>caloptera</i> Kieffer, 1919, Heteromyia	14.48
<i>calvescens</i> Macfie, 1938, Dasyhelea	14.17
<i>composi</i> Ortiz & León, 1954, Culicoides	14.28
<i>cancer</i> Hogue & Wirth, 1968, Culicoides	14.29
<i>caprilesi</i> Fox, 1952, Culicoides	14.29
<i>caribbea</i> Wirth & Dow, 1971, Forcipomyia	14.3
<i>caribbeana</i> Saunders, 1956, Forcipomyia	14.9
<i>caribbeanus</i> Ewen, 1958, Atrichopogon	14.13

caribe Lane & Forattini, 1958, <i>Stilobezzia</i>	14.44
caridei (Brèthes), 1912 (<i>Cotocripus</i>), <i>Culicoides</i>	14.29
carioca Lane, 1958, <i>Bezzia</i>	14.52
carioca Lane, 1960, <i>Palpomyia</i>	14.54
carpenteri Wirth & Blanton, 1953, <i>Culicoides</i>	14.29
carrerai (Lane), 1948 (<i>Dicrobezzia</i>), <i>Palpomyia</i>	14.54
carrerai Lane, 1956, <i>Stenoxenus</i>	14.52
carsiomelas Wirth & Blanton, 1955, <i>Culicoides</i>	14.29
casali Cavalieri & Chiossone, 1966, <i>Leptoconops</i>	14.2
castanea Lane, 1946, <i>Heteromyia</i>	14.47
castanea Macfie, 1939, <i>Palpomyia</i>	14.54
castillae Fox, 1946, <i>Culicoides</i> ..	14.29
catarinensis Lane, 1960, <i>Palpomyia</i>	14.54
caucaensis Wirth & Lee, 1967, <i>Culicoides</i>	14.29
caudata Ingram & Macfie, 1931, <i>Macrurohelea</i>	14.39
cebacoï Lane & Wirth, 1964, <i>Monohelea</i>	14.40
<i>Centrorhynchus</i> Lutz, 1914	14.6
<i>Ceratobezzia</i> Kieffer, 1917	14.46
<i>Ceratopogoninae</i>	14.18
<i>Ceratopogonini</i>	14.38
<i>cerifera</i> Saunders, 1956, <i>Forcipomyia</i>	14.7
chaconi Macfie, 1938, <i>Stilobezzia</i>	14.44
chaquensis Duret & Lane, 1955, <i>Heteromyia</i>	14.47
chiapasi Lane & Wirth, 1964, <i>Monohelea</i>	14.40
chilensis Forattini, 1958, <i>Leptoconops</i>	14.2
chilensis Ingram & Macfie, 1931, <i>Atrichopogon</i>	14.13
chilensis Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.17
chilensis Ingram & Macfie, 1931, <i>Palpomyia</i>	14.54
chilensis (Philippi), 1865 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.5
chirusi Lane & Wirth, 1964, <i>Monohelea</i>	14.40
chrysonotus Wirth & Blanton, 1956, <i>Culicoides</i>	14.19
claripennis Lynch Arribáizaga, 1893, <i>Ceratopogon</i> . Position Uncertain	14.55
clavata Williston, 1900, <i>Heteromyia</i>	14.47
clavipes Wirth & Ratanaworabhan, 1972, <i>Neobezzia</i>	14.49
<i>Clinhelea</i> Kieffer, 1917	14.46
cochisensis Wirth & Blanton, 1967, <i>Culicoides</i>	14.37
colombiae Lane & Wirth, 1964, <i>Monohelea</i>	14.40
colombiae Wirth, 1970, <i>Forcipomyia</i>	14.11
coloradensis Wirth, 1952, <i>Bezzia</i>	14.52
columbiana (Kieffer), 1917 (<i>Culicoides</i>), <i>Dasyhelea</i>	14.17
columbiana Kieffer, 1917, <i>Palpomyia</i>	14.54
columbianus Kieffer, 1917, <i>Atrichopogon</i>	14.13
commatis Wirth & Blanton, 1959, <i>Culicoides</i>	14.37
concoloripes Macfie, 1940, <i>Bezzia</i>	14.52
conifera Macfie, 1939, <i>Palpomyia</i>	14.54
<i>contubernalis</i> Ortiz & León, 1954, <i>Culicoides</i>	14.26
copiosus Root & Hoffman, 1937, <i>Culicoides</i>	14.22
coquilletti Kieffer, 1917, <i>Stilobezzia</i>	14.44
coracina Kieffer, 1917, <i>Stilobezzia</i>	14.44
cornuta Saunders, 1964, <i>Forcipomyia</i>	14.7
coroicoensis Wirth, nom. nov., <i>Palpomyia</i>	14.54
correntina Duret & Lane, 1955, <i>Heteromyia</i>	14.47
costalis Macfie, 1939, <i>Atrichopogon</i>	14.13
costalis Wirth, 1965, <i>Parabezzia</i>	14.42
costalis Wirth, 1965, <i>Parabezzia</i>	14.14
costaricae Macfie, 1953, <i>Atrichopogon</i>	14.49
costaricae Wirth & Ratanaworabhan, 1972, <i>Neobezzia</i>	14.49
<i>Cotocripus</i> Brèthes, 1912	14.27



<i>coutinhoi</i> Barretto, 1944, <i>Culicoides</i>	14.24
<i>covagarciai</i> Ortiz, 1950, <i>Culicoides</i>	14.19
<i>crassicus</i> Kieffer, 1917, <i>Palpomyia</i>	14.54
<i>crepuscularis</i> Malloch, 1915, <i>Culicoides</i>	14.21
<i>crescentis</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.37
<i>crucifer</i> Clastrier, 1968, <i>Culicoides</i>	14.29
<i>crudelis</i> Knab, 1914, <i>Forcipomyia</i>	14.8
<i>crudelis</i> Mayer, 1937 (<i>Phasmidohelea</i>), <i>Forcipomyia</i>	14.8
<i>cryptogamus</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>Culicoides</i> Latreille, 1809	14.18, 14.21
<i>Culicoidini</i>	14.18
<i>cunasi</i> Lane & Wirth, 1964, <i>Monohoelea</i>	14.40
<i>cylindrica</i> Wirth & Dow, 1972, <i>Forcipomyia</i>	14.10
<i>daedaloides</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.37
<i>daedalus</i> Macfie, 1948, <i>Culicoides</i>	14.37
<i>damascenoi</i> Lane & Duret, 1954, <i>Clinohoelea</i>	14.46
<i>dampfi</i> Root & Hoffman, 1937, <i>Culicoides</i>	14.37
<i>danaisi</i> (Floch & Abonnenc), 1949 (<i>Lasiohelea</i>), <i>Forcipomyia</i>	14.12
<i>darlingtonae</i> Wirth & Blanton, 1970, <i>Culicoides</i>	14.29
<i>Dasyhelea</i> Kieffer, 1911	14.16
<i>Dasyheleinae</i>	14.16
<i>dasyophrus</i> Macfie, 1940, <i>Culicoides</i>	14.29
<i>daviesi</i> Wirth & Blanton, 1968, <i>Culicoides</i>	14.37
<i>debilipalpis</i> Lutz, 1913, <i>Culicoides</i>	14.29
<i>decor</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.19
<i>delpontei</i> Cavalieri, 1962, <i>Forcipomyia</i>	14.5
<i>denisae</i> Clastrier, 1971, <i>Culicoides</i>	14.30
<i>depilis</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>desertensis</i> Wirth & Hubert, 1960, <i>Forcipomyia</i>	14.5
<i>diabolicus</i> Hoffman, 1925, <i>Culicoides</i>	14.24
<i>diabolicus</i> , authors, <i>Culicoides</i>	14.24
<i>Diaphanobezzia</i> Ingram & Macfie, 1931	14.42
<i>Dicrobezzia</i> , authors	14.50
<i>Dicrohelea</i> Kieffer, 1917	14.48
<i>dicourus</i> Wirth & Blanton, 1955, <i>Culicoides</i>	14.30
<i>Didymophleps</i> Weyenbergh, 1883	14.13
<i>didymothecae</i> Macfie, 1953, <i>Atrichopogon</i>	14.14
<i>diminuta</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.44
<i>diminutus</i> Barbosa, 1951, <i>Culicoides</i>	14.26
<i>dimorphus</i> Kieffer, 1909, <i>Stenoxenus</i>	14.52
<i>Diphaomyia</i> Vargas, 1960	14.21
<i>discoloripes</i> Macfie, 1939, <i>Forcipomyia</i>	14.5
<i>discrepans</i> Ortiz & Mirsa, 1951, <i>Culicoides</i>	14.30
<i>disjuncta</i> (Ingram & Macfie), 1931 (<i>Isthmohelea</i>), <i>Monohoelea</i>	14.40
<i>dominicanus</i> Wirth & Blanton, 1970, <i>Culicoides</i>	14.19
<i>dominicii</i> Ortiz, 1951, <i>Culicoides</i>	14.34
<i>donajii</i> Vargas, 1954, <i>Culicoides</i>	14.30
<i>dorsalis</i> Wirth & Dow, 1972, <i>Forcipomyia</i>	14.10
<i>dorsofasciata</i> (Lutz), 1914 (<i>Palpomyia</i>), <i>Stilobezzia</i>	14.43
<i>dryadum</i> Macfie, 1940, <i>Stilobezzia</i>	14.44
<i>Drymodesmyia</i> Vargas, 1960	14.22
<i>dubia</i> Macfie, 1939, <i>Forcipomyia</i>	14.5
<i>dubitans</i> Lane, Forattini & Rabello, 1955, <i>Stilobezzia</i>	14.44
<i>dunni</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.38
<i>dureti</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.45



<i>Echinohelea</i> Macfie, 1940	14.39
<i>eadi</i> Wirth & Blanton, 1970, <i>Culicoides</i>	14.30
<i>echinodes</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>edwardsi</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.5
<i>edwardsi</i> Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.43
<i>edwardsi</i> (Saunders), 1925 (<i>Apelma</i>), <i>Forcipomyia</i>	14.9
<i>edwardsiana</i> Wirth, nom. nov., <i>Forcipomyia</i>	14.5
<i>efferus</i> Fox, 1952, <i>Culicoides</i>	14.19
<i>elegantula</i> (Johannsen), 1907 (<i>Bezzia</i>), <i>Stilobezzia</i>	14.43
<i>elutus</i> Macfie, 1948, <i>Culicoides</i>	14.21
<i>enderleini</i> Lane, 1956, <i>Paryphoconus</i>	14.51
<i>equatoriensis</i> Barbosa, 1952, <i>Culicoides</i>	14.30
<i>eriophora</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.8
<i>esmeralda</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.45
<i>eublepharus</i> Macfie, 1948, <i>Culicoides</i>	14.30
<i>eublepharus</i> , authors, <i>Culicoides</i>	14.36
<i>eucnemus</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>Euforcipomyia</i> Malloch, 1915	14.4
<i>eukosma</i> Macfie, 1939, <i>Forcipomyia</i>	14.3
<i>Eukraiohelea</i> Ingram & Macfie, 1921	14.43
<i>Euprojoannisia</i> Brèthes, 1915	14.3
<i>evansi</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.21
<i>excentricus</i> Lane, 1961, <i>Stenoxenus</i>	14.52
<i>fairchildi</i> Lane & Wirth, 1964, <i>Monohhelea</i>	14.40
<i>fairchildi</i> Wirth & Blanton, 1955, <i>Culicoides</i>	14.28
<i>falcifera</i> Saunders, 1959, <i>Forcipomyia</i>	14.4
<i>fallax</i> Kieffer, 1917, <i>Ceratobezzia</i>	14.46
<i>farri</i> Wirth, 1966, <i>Forcipomyia</i>	14.10
<i>farri</i> Wirth & Blanton, 1970, <i>Culicoides</i>	14.19
<i>fasciatus</i> Walker, 1856 (<i>Pachyleptus</i>), <i>Heteromyia</i>	14.47
<i>femoralis</i> Lane & Forattini, 1956, <i>Stilobezzia</i>	14.45
<i>fernandezi</i> Ortiz, 1954, <i>Culicoides</i>	14.30
<i>fiebrigi</i> Kieffer, 1917, <i>Atrichopogon</i>	14.14
<i>fiebrigi</i> Kieffer, 1917, <i>Stilobezzia</i>	14.45
<i>fieldi</i> Wirth & Blanton, 1956, <i>Culicoides</i>	14.30
<i>filarifer</i> Hoffman, 1939, <i>Culicoides</i>	14.24
<i>filibranchia</i> (Lutz), 1914 (<i>Ceratopogon</i>), <i>Dasyhelea</i>	14.17
<i>fimbriatus</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>Fittkauhelea</i> Wirth & Blanton, 1970	14.39
<i>fittkai</i> Wirth & Ratanaworabhan, 1972, <i>Neobezzia</i>	14.49
<i>flava</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.5
<i>flavicauda</i> Macfie, 1939, <i>Dasyhelea</i>	14.17
<i>flavicauda</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>flavida</i> (Johannsen), 1943 (<i>Ceratobezzia</i>), <i>Bezzia</i>	14.53
<i>flavifemoris</i> Macfie, 1940, <i>Forcipomyia</i>	14.5
<i>flavipes</i> Lutz, 1914, <i>Atrichopogon</i>	14.14
<i>flavivenula</i> Lima, 1937, <i>Culicoides</i>	14.24
<i>flochabonnenci</i> Ortiz & Mirsa, 1952, <i>Culicoides</i>	14.30
<i>florenciae</i> Messersmith, 1972, <i>Culicoides</i>	14.30
<i>floridensis</i> Beck, 1951, <i>Culicoides</i>	14.30
<i>floridensis</i> Wirth, 1951, <i>Leptoconops</i>	14.3
<i>fluminensis</i> Lane, 1948, <i>Bezzia</i>	14.53
<i>fluminensis</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>flumineus</i> Macfie, 1935, <i>Atrichopogon</i>	14.14
<i>fluvialis</i> Macfie, 1940, <i>Culicoides</i>	14.30
<i>fluviatilis</i> (Lutz), 1914 (<i>Johannseniella</i>), <i>Culicoides</i>	14.38



forattinii Ortiz, 1961, <i>Culicoides</i>	14.31
Forcipomyia Meigen, 1818	14.3, 14.4
Forcipomyiinae	14.3
foxi Ortiz, 1950, <i>Culicoides</i>	14.24
fuliginosa (Meigen), 1818 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.8
fulvus Johannsen, 1927, <i>Stenoxenus</i>	14.52
furens (Poey), 1851 (<i>Oecacta</i>), <i>Culicoides</i>	14.31
furcifera Macfie, 1940, <i>Forcipomyia</i>	14.3
furva Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.43
fusca (Philippi), 1865 (<i>Tetrastoma</i>), <i>Forcipomyia</i>	14.5
<i>fusicornis</i> , error, <i>Forcipomyia</i>	14.10
fuscipennis Lane & Wirth, <i>Monohalea</i>	14.40
fuscipennis Wirth, 1965, <i>Parabozzia</i>	14.42
fuscivenosus (Lutz), 1914 (<i>Palpomyia</i>), <i>Brachypogon</i>	14.38
fuscus (Coquillett), 1901 (<i>Ceratopogon</i>), <i>Atrichopogon</i>	14.14
fuscus Lane, 1946, <i>Paryphoconus</i>	14.51
fusicornis (Coquillett), 1905 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.10
gabaldoni Ortiz, 1954, <i>Culicoides</i>	14.31
galapagensis (Coquillett), 1901 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.8
galindoi Wirth & Blanton, 1953, <i>Culicoides</i>	14.31
genualis (Loew), 1866 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.5
germanus Macfie, 1940, <i>Culicoides</i>	14.31
<i>germanus</i> , authors, <i>Culicoides</i>	14.30, 14.31
gibbera (Coquillett), 1905 (<i>Ceratopogon</i>), <i>Bezzia</i>	14.53
<i>gibsoni</i> Wirth, 1952, <i>Culicoides</i>	14.29
ginesi Ortiz, 1951, <i>Culicoides</i>	14.31
glabellus Wirth & Blanton, 1956, <i>Culicoides</i>	14.31
glaber Macfie, 1935, <i>Atrichopogon</i>	14.14
glabrior Macfie, 1940, <i>Culicoides</i>	14.31
<i>Glaphiromyia</i> Vargas, 1960	14.23
glauca Macfie, 1939, <i>Stilobezzia</i>	14.45
globulifer Macfie, 1939, <i>Atrichopogon</i>	14.14
goianensis Lane, 1961, <i>Bezzia</i>	14.53
goianensis Lane, 1961, <i>Paryphoconus</i>	14.51
goniognatha Wirth & Messersmith, 1971, <i>Forcipomyia</i>	14.12
gordoni Macfie, 1938, <i>Atrichopogon</i>	14.14
gorgasi Wirth & Blanton, 1953, <i>Culicoides</i>	14.32
<i>grahambelli</i> Forattini, 1956, <i>Culicoides</i>	14.32
grandis Lane & Forattini, 1958, <i>Stilobezzia</i>	14.45
grandis Macfie, 1939, <i>Paryphoconus</i>	14.51
guadeloupensis Floch & Abonnenc, 1950, <i>Culicoides</i>	14.32
guaimiesi Lane & Wirth, 1964, <i>Monohalea</i>	14.40
guamai Wirth & Dow, 1972, <i>Forcipomyia</i>	14.10
guarani Lane, 1947, <i>Palpomyia</i>	14.54
guerrai Wirth & Blanton, 1970, <i>Culicoides</i>	14.32
guianae (Macfie), 1940, (<i>Acanthohalea</i>), <i>Stilobezzia</i>	14.42
<i>guianae</i> Macfie, 1940, <i>Paryphoconus</i>	14.51
guianae Wirth, 1953, <i>Monohalea</i>	14.41
guianensis Macfie, 1940, <i>Atrichopogon</i>	14.14
guttatus (Coquillett), 1904 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.24
<i>guttatus</i> authors, <i>Culicoides</i>	14.24
guyanensis Floch & Abonnenc, 1942, <i>Culicoides</i>	14.32
<i>Haematomyidium</i> Goeldi, 1905	14.27
haematopotus Malloch, 1915, <i>Culicoides</i>	14.21
harrisi Macfie, 1938, <i>Atrichopogon</i>	14.14




hayesi Matta, 1967, Culicoides	14.20
heliconiae Fox & Hoffman, 1944, Culicoides	14.24
hertigi Wirth & Blanton, 1953, Culicoides	14.23
Heteromyia Say, 1825	14.47
Heteromyiini	14.46
hieroglyphica authors, Monohelea	14.41
hieroglyphica Kieffer, 1917, Monohelea	14.41
hieroglyphicus Malloch, 1915, Culicoides	14.36
hippolytae Macfie, 1935, Dasyhelea	14.17
hirsuta Ingram & Macfie, 1931, Stilobezzia	14.43
hirtipes (Kieffer), 1917 (Culicoides), Dasyhelea	14.17
hoffmani Fox, 1946, Culicoides	14.32
Hoffmania Fox, 1948	14.23
Holoconops Kieffer, 1918	14.2
hondurensis Hoffman, 1926, Leptoconops	14.2
horacioi Lane, 1944, Clinohela	14.47
horticola Lutz, 1913, Culicoides	14.32
hortorum (Weyenbergh), 1883 (Didymophleps), Atrichopogon	14.14
hyeroglyphica, error, Monohelea	14.41
hylas Macfie, 1940, Culicoides	14.25
ignacioi Forattini, 1957, Culicoides	14.25
imitator Ortiz, 1953, Culicoides	14.32
impar (Johannsen), 1938 (Ceratopogon), Brachypogon	14.39
inamollae Fox & Hoffman, 1944, Culicoides	14.25
incubans (Macfie), 1937 (Pterobosca), Forcipomyia	14.10
incultus Ewen, 1958, Atrichopogon	14.14
indecora Kieffer, 1914, Forcipomyia	14.4
inermicollis Kieffer, 1917, Palpomyia	14.54
ingrami Carter, 1919, Forcipomyia	14.4
insigniforceps Macfie, 1939, Forcipomyia	14.7
insignipalpis Macfie, 1940, Atrichopogon	14.14
insignipalpis Macfie, 1949, Forcipomyia	14.8
insignis authors, Culicoides	14.25
insignis Lutz, 1913, Culicoides	14.25
insigniventris Macfie, 1935, Atrichopogon	14.15
insinuatus Ortiz & León, 1954, Culicoides	14.32
insolatus Wirth & Hubert, 1960, Culicoides	14.22
intermedia Saunders, 1964, Forcipomyia	14.7
intrepida Macfie, 1936, Forcipomyia	14.12
iriartei Fox, 1952, Culicoides	14.22
Isthmohelea Ingram & Macfie, 1931	14.40
ixodooides (Fiebrig-Gertz), 1928 (Ceratopogon), Forcipomyia	14.8
jamaicensis Edwards, 1922, Culicoides	14.22
Jamaicensis Wirth, 1970, Forcipomyia	14.11
Jamaicensis Wirth, 1965, Parabezzia	14.42
Japajapae Wirth, 1970, Forcipomyia	14.11
Jocosa Saunders, 1956, Forcipomyia	14.9
Johannseni (Lane), 1948 (Dicrobezzia), Palpomyia	14.54
Johannseni Wirth, 1953, Monohelea	14.41
Johannsenomyia Malloch, 1915	14.48
Johnsoni Coquillett, 1899, Stenoxenus	14.52
kelini (Saunders), 1915 (Apelma), Forcipomyia	14.9
Kiefferi Lane, 1946, Heteromyia	14.47
Kiefferi Lane, 1956, Paryphoconus	14.51



<i>kiefferi</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>kintzi</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.29
<i>kuanosceles</i> Macfie, 1939, <i>Forcipomyia</i>	14.7
<i>kuscheli</i> Wirth, 1965, <i>Macrurohelea</i>	14.39
<i>lacajae</i> Macfie, 1953, <i>Atrichopogon</i>	14.15
<i>lacrimatorii</i> Macfie, 1939, <i>Forcipomyia</i>	14.5
<i>lacustris</i> Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.17
<i>lacustris</i> Lane, Forattini & Rabello, 1955, <i>Palpomyia</i>	14.54
<i>lahillei</i> Ichas, 1906 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.34
<i>lamprogaster</i> Edwards, 1933, <i>Heteromyia</i>	14.47
<i>Lanhelea</i> Wirth & Blanton, 1972	14.49
<i>lanei</i> Ortiz, 1950, <i>Culicoides</i>	14.32
<i>lanei</i> Wirth, 1965, <i>Johannsenomyia</i>	14.48
<i>lanei</i> Wirth, 1953, <i>Monhelea</i>	14.41
<i>lanei</i> Wirth, 1959, <i>Paryphoconus</i>	14.51
<i>lanei</i> Wirth & Ratanaworabhan, 1971, <i>Pellucidomyia</i>	14.48
<i>lanei</i> Wirth & Ratanaworabhan, 1972, <i>Stenoxenus</i>	14.52
<i>Lasiohelea</i> Kieffer, 1921	14.6, 14.7
<i>latifemoris</i> Ingram & Macfie, 1931 (<i>Johannsenomyia</i>), <i>Pachyhelea</i> ...	14.53
<i>leei</i> Wirth & Blanton, 1972, <i>Lanhelea</i>	14.49
<i>leei</i> Wirth & Blanton, 1970, <i>Parastilobezzia</i>	14.42
<i>lenkoi</i> Lane, 1958, <i>Bezzia</i>	14.53
<i>leoni</i> Barbosa, 1952, <i>Culicoides</i>	14.33
<i>leopoldoi</i> Ortiz, 1951, <i>Culicoides</i>	14.33
<i>Lepidohelea</i> Kieffer, 1917	14.7
<i>Leptoconopinae</i>	14.2
<i>Leptoconops</i> Skuse, 1889	14.2
<i>Leptohelea</i> Wirth & Blanton, 1970	14.39
<i>lesliei</i> Wirth, nom. nov., <i>Forcipomyia</i>	14.12
<i>lichyi</i> Floch & Abonnenc, 1949, <i>Culicoides</i>	14.33
<i>lichyi</i> , authors, <i>Culicoides</i>	14.30
<i>limai</i> Barretto, 1944, <i>Culicoides</i>	14.33
<i>limnetis</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.12
<i>limnochares</i> (Macfie), 1940 (<i>Sphaeromias</i>), <i>Palpomyia</i>	14.54
<i>limonensis</i> Ortiz & León, 1954, <i>Culicoides</i>	14.33
<i>limpidus</i> Wirth & Ratanaworabhan, 1972, <i>Stenoxenus</i>	14.52
<i>lituratus</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Atrichopogon</i>	14.15
<i>longicornis</i> Ewen, 1958, <i>Atrichopogon</i>	14.15
<i>longicornis</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Palpomyia</i>	14.54
<i>longispina</i> Saunders, 1956, <i>Forcipomyia</i>	14.4
<i>lopesi</i> Barretto, 1944, <i>Culicoides</i>	14.33
<i>Lophomyidium</i> Cordero, 1929	14.13
<i>lotus</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.4
<i>loughnani</i> Edwards, 1922, <i>Culicoides</i>	14.23
<i>loughnani</i> , authors, <i>Culicoides</i>	14.22
<i>louriei</i> (Macfie), 1935 (<i>Lasiohelea</i>), <i>Forcipomyia</i>	14.12
<i>luglani</i> Jones & Wirth, 1958, <i>Culicoides</i>	14.33
<i>lutealaris</i> Wirth & Blanton, 1956, <i>Culicoides</i>	14.20
<i>luteisquamosa</i> Wirth, 1972, <i>Forcipomyia</i>	14.8
<i>luteogrisea</i> Wirth & Williams, 1957, <i>Dasyhelea</i>	14.17
<i>luteovenus</i> Root & Hoffman, 1937, <i>Culicoides</i>	14.21
<i>luteovenus</i> , authors, <i>Culicoides</i>	14.26
<i>lutzi</i> Lima, 1937, <i>Culicoides</i>	14.25
<i>lutzi</i> , authors, <i>Culicoides</i>	14.25
<i>lutzi</i> Lane, 1947, <i>Palpomyia</i>	14.55
<i>lyrinotatus</i> Wirth & Blanton, 1955, <i>Culicoides</i>	14.33


<i>Macfiella</i> Fox, 1955	14.26
<i>Macfichelea</i> Lane, 1946	14.48
<i>macfieii</i> Lane, 1946, <i>Paryphoconus</i>	14.51
<i>macfieii</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>macfieii</i> Lane, 1948, <i>Echinohelea</i>	14.39
<i>macfieii</i> Lima, 1937 (<i>Pterobosca</i>), <i>Forcipomyia</i>	14.10
<i>macfieii</i> , authors, <i>Monohelea</i>	14.41
<i>macrostigma</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.33
<i>Macurohelea</i> Ingram & Macfie, 1931	14.39
<i>maculata</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>maculata</i> Macfie, 1943, <i>Dasyhelea</i>	14.17
<i>maculicrus</i> Ingram & Macfie, 1931, <i>Palpomyia</i>	14.55
<i>maculipennis</i> Clastrier, 1968, <i>Atrichopogon</i>	14.15
<i>maculipennis</i> (Coquillett), 1905 (<i>Ceratopogon</i>), <i>Monohelea</i>	14.41
<i>maculithorax</i> Williston, 1896 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.31
<i>maculitibia</i> Lane & Forattini, 1956, <i>Stilobezzia</i>	14.43
<i>maculosa</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.5
<i>magna</i> (Saunders), 1925 (<i>Apelma</i>), <i>Forcipomyia</i>	14.10
<i>magnipalpis</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.33
<i>maia</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.43
<i>Mallochohelea</i> Wirth, 1962	14.49
<i>manaosensis</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.43
<i>marginalis</i> Wirth & Ratanaworabhan, 1972, <i>Stenoxenus</i>	14.52
<i>marinkellei</i> Wirth & Lee, 1967, <i>Culicoides</i>	14.22
<i>marshi</i> Wirth & Blanton, 1956, <i>Culicoides</i>	14.20
<i>martinezi</i> Wirth & Blanton, 1970, <i>Culicoides</i>	14.33
<i>maruim</i> Lutz, 1913, <i>Culicoides</i>	14.25
<i>Mataemyia</i> Vargas, 1960	14.27
<i>mayeri</i> Forattini & Lane, 1956, <i>Forcipomyia</i>	14.8
<i>mayeri</i> Ortiz, 1950, <i>Monohelea</i>	14.41
<i>mayeri</i> Wirth, 1959, <i>Paryphoconus</i>	14.51
<i>mazaruni</i> Macfie, 1940, <i>Bezzia</i>	14.53
<i>Megaconops</i> Wirth & Atchley, 1973	14.3
<i>melleus</i> (Coquillett), 1901 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.34
<i>mellichroa</i> Macfie, 1939, <i>Palpomyia</i>	14.55
<i>mendozae</i> Ingram & Macfie, 1931, <i>Atrichopogon</i>	14.15
<i>messersmithi</i> Wirth & Dow, 1972, <i>Forcipomyia</i>	14.10
<i>Metaforcipomyia</i> Saunders, 1956	14.7
<i>metagonatus</i> Wirth & Blanton, 1956, <i>Culicoides</i>	14.20
<i>mexicana</i> Wirth, 1956, <i>Forcipomyia</i>	14.12
<i>microchela</i> Kieffer, 1917, <i>Palpomyia</i>	14.55
<i>Microhelea</i> Kieffer, 1917	14.8
<i>micronyx</i> Wirth & Blanton, 1970, <i>Leptohelea</i>	14.39
<i>microtoma</i> (Kieffer), 1917 (<i>Atrichopogon</i>), <i>Forcipomyia</i>	14.9
<i>minisquamosa</i> Wirth, 1972, <i>Forcipomyia</i>	14.9
<i>mirsae</i> Ortiz, 1953, <i>Culicoides</i>	14.34
<i>miyamotoi</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.28
<i>modesta</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>mohave</i> Wirth, 1952, <i>Culicoides</i>	14.34
<i>mojingaensis</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.34
<i>Monoculicoides</i> Khalaf, 1954	14.27
<i>Monohelea</i> Kieffer, 1917	14.40
<i>monticola</i> Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.17
<i>monticola</i> Wirth & Lee, 1967, <i>Culicoides</i>	14.34
<i>mortuifolii</i> Saunders, 1959, <i>Forcipomyia</i>	14.4
<i>multilineata</i> (Lutz), 1914 (<i>Palpomyia</i>), <i>Monohelea</i>	14.41
<i>multipecta</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.6




<i>Nycteromyia</i> Lutz, 1912	14.2
<i>Mycterotypus</i> Noé, 1905	14.2
<i>nana</i> (Macfie), 1939 (Lasiohelea), Forcipomyia	14.11
<i>nana</i> Wirth & Blanton, 1970, Bacohelea	14.38
<i>nanus</i> Macfie, 1940, Atrichopogon	14.15
<i>narthekophora</i> Macfie, 1939, Forcipomyia	14.13
<i>nebulosus</i> Macfie, 1939, Atrichopogon	14.15
<i>neghmei</i> Vargas, 1955, Culicoides	14.34
<i>neivai</i> Lane, 1944, Clinohela	14.47
<i>memoralis</i> (Macfie), 1940 (Johannsenomyia), Mallochohelea	14.49
<i>Neobezzia</i> Wirth & Ratanaworabhan, 1972	14.49
<i>Neoforcipomyia</i> Tokunaga, 1959	14.12
<i>neopulicaris</i> Wirth, 1955, Culicoides	14.21
<i>Neostilobezzia</i> Goetghebuer, 1934	14.43
<i>neotropica</i> (Macfie), 1940 (Bezzia), Nilobezzia	14.50
<i>neotropicalis</i> (Lane), 1948 (Macropeza), Paryphoconus	14.51
<i>Neurohelea</i> Kieffer, 1925	14.48
<i>niger</i> Lane, 1948, Stenoxenus	14.52
<i>nigerrima</i> Ingram & Macfie, 1931, Stilobezzia	14.43
<i>nigra</i> Kieffer, 1917, Heteromyia	14.47
<i>nigrescens</i> Macfie, 1939, Forcipomyia	14.4
<i>nigrigenus</i> Wirth & Blanton, 1956, Culicoides	14.20
<i>nigripes</i> Macfie, 1939, Clinohela	14.47
<i>nigripes</i> (Macfie), 1939 (Johannsenomyia), Mallochohelea	14.49
<i>nigripes</i> Macfie, 1939, Paryphoconus	14.51
<i>nigroflava</i> Lane, 1947, Palpomyia	14.55
<i>nigroflava</i> Lane & Forattini, 1958, Stilobezzia	14.45
<i>nigroscutellata</i> Lane, 1947, Palpomyia	14.55
<i>Nilobezzia</i> Kieffer, 1921	14.50
<i>nodosa</i> Saunders, 1959, Forcipomyia	14.11
<i>novaetuntoniae</i> Macfie, 1939, Atrichopogon	14.15
<i>nubeculosus</i> Macfie, 1949, Atrichopogon	14.15
<i>nubifer</i> Macfie, 1939, Paryphoconus	14.51
<i>nuncupata</i> Macfie, 1949, Forcipomyia	14.6
<i>obesa</i> Lima, 1928, Forcipomyia	14.9
<i>obfuscatus</i> Ingram & Macfie, 1931, Atrichopogon	14.15
<i>obnoxius</i> Fox, 1952, Culicoides	14.34
<i>obnubilus</i> Ingram & Macfie, 1931, Atrichopogon	14.15
<i>obscura</i> Lane & Forattini, 1958, Stilobezzia	14.45
<i>obscurus</i> Ewen, 1958, Atrichopogon	14.15
<i>occidentalis</i> Wirth & Jones, 1957, Culicoides	14.27
<i>ocumare</i> Ortiz, 1952, Monohelea	14.41
<i>ocumarensis</i> Ortiz, 1950, Culicoides	14.25
<i>Oeacta</i> Poey, 1851	14.27
<i>oedidactyla</i> Ingram & Macfie, 1931, Neurohelea	14.48
<i>oklahomensis</i> Khalaf, 1952, Culicoides	14.38
<i>oligarthra</i> Saunders, 1956, Forcipomyia	14.10
<i>oliveirai</i> Lane, 1947, Palpomyia	14.55
<i>oliveirai</i> Lane, 1956, Paryphoconus	14.51
<i>oliveirai</i> (Lane), 1946 (Macfiehelea), Pellucidomyia	14.48
<i>oliveri</i> Fox & Hoffman, 1944, Culicoides	14.26
<i>opilionivora</i> (Lane), 1947 (Lasiohelea), Forcipomyia	14.12
<i>orellana</i> (Roback), 1957 (Palpomyia), Heteromyia	14.47
<i>orjuelai</i> Wirth & Lee, 1967, Culicoides	14.20
<i>ornata</i> Lane & Forattini, 1958, Stilobezzia	14.43




ornata Wirth, 1953, <i>Monohoelea</i>	14.41
ornaticrus Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.43
ornatipennis Macfie, 1940, <i>Echinohelea</i>	14.39
ornatipennis Macfie, 1939, <i>Forcipomyia</i>	14.6
ortizi Fox, 1952, <i>Culicoides</i>	14.36
ousairani Khalaf, 1952, <i>Culicoides</i>	14.38
pachynermus Macfie, 1953, <i>Atrichopogon</i>	14.15
Pachyhelea Wirth, 1959	14.53
<i>Pachyleptus</i> Walker, 1856	14.47
pachymera (Williston), 1900 (<i>Ceratopogon</i>), <i>Pachyhelca</i>	14.53
pachymerus Lutz, 1914, <i>Culicoides</i>	14.34
<i>pachymerus</i> , authors, <i>Culicoides</i>	14.29, 14.38
<i>painteri</i> Fox, 1946, <i>Culicoides</i>	14.25
pallescens Lane & Forattini, 1958, <i>Stilobezzia</i>	14.45
pallidipes Kieffer, 1917, <i>Atrichopogon</i>	14.15
palpalis Macfie, 1939, <i>Atrichopogon</i>	14.15
palpalis Macfie, 1948, <i>Culicoides</i>	14.26
Palpomyia Meigen, 1818	14.53
Palpomyiini	14.52
pampana Lane, 1960, <i>Palpomyia</i>	14.55
pampokilus Macfie, 1948, <i>Culicoides</i>	14.34
panamensis Barbosa, 1947, <i>Culicoides</i>	14.23
panamensis Lane & Forattini, 1958, <i>Stilobezzia</i>	14.45
panamensis Lane & Wirth, 1964, <i>Monohoelea</i>	14.41
panamensis Wirth, 1965, <i>Parabezzia</i>	14.42
<i>panamericanus</i> Fox, 1947, <i>Culicoides</i>	14.27
Parabezzia Malloch, 1915	14.42
paracincta Wirth, 1969, <i>Dasyhelea</i>	14.17
Paradasyhelea Macfie, 1940	14.18
paraensis (Goeldi), 1805 (<i>Haematomyidium</i>), <i>Culicoides</i>	14.34
paraensis Lane, 1956, <i>Stenoxenus</i>	14.52
paraensis (Lane), 1958 (<i>Bezzia</i>), <i>Nilobezzia</i>	14.50
paraensis Lane, 1960, <i>Palpomyia</i>	14.55
paraensis Wirth & Blanton, 1970, <i>Brachypogon</i>	14.39
parahybae Macfie, 1940, <i>Dasyhelea</i>	14.17
Parastilobezzia Wirth & Blanton, 1970	14.42
Paryphoconus Enderlein, 1912	14.50
patagonica Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.17
<i>patagonica</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.5
patagonica Ingram & Macfie, 1931, <i>Palpomyia</i>	14.55
patagonica Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.43
patulipalpis Wirth & Blanton, 1959, <i>Culicoides</i>	14.34
paucienfuscatus Barbosa, 1947, <i>Culicoides</i>	14.34
paulistana Forattini & Rabello, 1957, <i>Dasyhelea</i>	14.17
paulistensis Lane, 1947, <i>Palpomyia</i>	14.55
paulistensis Lane, 1961, <i>Paryphoconus</i>	14.51
paulistensis Lane, 1947, <i>Stilobezzia</i>	14.45
pectinatus Macfie, 1939, <i>Atrichopogon</i>	14.15
pellucida (Ingram & Macfie), 1931 (<i>Diaphanobezzia</i>), <i>Parabezzia</i>	14.42
Pellucidomyia Macfie, 1939	14.48
pellucidomyia Macfie, 1939	14.17
pentalineata Wirth & Hubert, 1960, <i>Dasyhelea</i>	14.17
penthesileae Macfie, 1935, <i>Dasyhelea</i>	14.17
penultimata Wirth, 1972, <i>Forcipomyia</i>	14.9
peruvianus Kieffer, 1917, <i>Atrichopogon</i>	14.15
<i>petiolata</i> , authors, <i>Parabezzia</i>	14.42
petrocchiaie Shannon & Del Ponte, 1927, <i>Leptoconops</i>	14.3



<i>phaeonotus</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.35
<i>Phasmidohelea</i> Mayer, 1937	14.8
<i>philotherma</i> Macfie, 1953, <i>Dasyhelea</i>	14.17
<i>phlebotomus</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.26
<i>Phytohelea</i> Remm, 1971	14.9
<i>piceiventris</i> Kieffer, 1917, <i>Atrichopogon</i>	14.15
<i>pictipennis</i> Philippi, 1865 (<i>Psychophaena</i>), <i>Culicoides</i>	14.36
<i>pictoni</i> Macfie, 1938, <i>Forcipomyia</i>	14.6
<i>pictus</i> Coquillett, 1905 (<i>Ceratopogon</i>), <i>Stilobezzia</i>	14.44
<i>pifanoi</i> Ortiz, 1951, <i>Culicoides</i>	14.35
<i>pilosior</i> Macfie, 1939, <i>Atrichopogon</i>	14.14
<i>pilosus</i> Wirth & Blanton, 1959, <i>Culicoides</i>	14.23
<i>piroskyi</i> Cavalieri, 1962, <i>Forcipomyia</i>	14.6
<i>platensis</i> (Brèthes), 1915, (<i>Euprojoannisia</i>), <i>Forcipomyia</i>	14.6
<i>pluvialis</i> Malloch, 1923, <i>Forcipomyia</i>	14.7
<i>poikilonotus</i> Macfie, 1948, <i>Culicoides</i>	14.23
<i>polypori</i> Wirth & Blanton, 1968, <i>Culicoides</i>	14.26
<i>polystictus</i> Kieffer, 1921, <i>Culicoides</i>	14.38
<i>poncai</i> Lane & Wirth, 1964, <i>Monohelea</i>	14.41
<i>popayanensis</i> Wirth & Lee, 1967, <i>Culicoides</i>	14.20
<i>poulaineae</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.6
<i>Proforcipomyia</i> Saunders, 1956	14.4
<i>propinqua</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.5
<i>propinquus</i> Macfie, 1948, <i>Culicoides</i>	14.35
<i>propriipennis</i> Macfie, 1948, <i>Culicoides</i>	14.35
<i>Protersesthes</i> Kieffer, 1921	14.2
<i>Pseudoculicoides</i> Malloch, 1915	14.16
<i>pseudodiabolicus</i> Fox, 1946, <i>Culicoides</i>	14.26
<i>Psychophaena</i> Philippi, 1865	14.27
<i>Pterobosca</i> Macfie, 1932	14.10
<i>pulchripennis</i> Macfie, 1939, <i>Culicoides</i>	14.35
<i>pulchripes</i> Kieffer, 1917, <i>Bezzia</i>	14.53
<i>pullata</i> (Wirth), 1952 (<i>Johannsenomyia</i>), <i>Mallochohelea</i>	14.49
<i>pumila</i> Macfie, 1939, <i>Dasyhelea</i>	14.18
<i>punctipennis</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Bezzia</i>	14.53
<i>punctipes</i> Wirth, 1953, <i>Stilobezzia</i>	14.45
<i>punctulata</i> Lane, 1947, <i>Stilobezzia</i>	14.45
<i>punctumalbum</i> , authors, <i>Forcipomyia</i>	14.6
<i>puracensis</i> Wirth & Lee, 1967, <i>Culicoides</i>	14.20
<i>puracensis</i> Wirth & Dow, 1972, <i>Forcipomyia</i>	14.11
<i>pusilla</i> (Lutz), 1913 (<i>Centrorhynchus</i>), <i>Dasyhelea</i>	14.18
<i>pusilloides</i> Wirth & Blanton, 1955, <i>Culicoides</i>	14.20
<i>pusillus</i> Lutz, 1913, <i>Culicoides</i>	14.21
<i>pygmaea</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Dasyhelea</i>	14.18
<i>pyrsonota</i> Macfie, 1953, <i>Dasyhelea</i>	14.18
<i>quasicornuta</i> Saunders, 1964, <i>Forcipomyia</i>	14.7
<i>quasiingrami</i> Macfie, 1939, <i>Forcipomyia</i>	14.4
<i>quasiparaensis</i> Clastrier, 1931, <i>Culicoides</i>	14.35
<i>rabelloi</i> Lane, 1947, <i>Stilobezzia</i>	14.46
<i>raleighi</i> Macfie, 1938, <i>Forcipomyia</i>	14.5
<i>rangeli</i> Ortiz & Mirsa, 1952, <i>Culicoides</i>	14.35
<i>rava</i> Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.43
<i>recifei</i> Barbosa, 1947, <i>Culicoides</i>	14.25
<i>recifensis</i> Barbosa, 1943, <i>Culicoides</i>	14.32
<i>remigatus</i> Ewen, 1958, <i>Atrichopogon</i>	14.15



<i>reticulatus</i> Lutz, 1913, <i>Culicoides</i>	14.35
<i>reynoldsi</i> Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.18
<i>Rhynchoforcipomyia</i> Wirth & Dow, 1972	14.10
<i>richardsi</i> Macfie, 1940, <i>Echinohelea</i>	14.39
<i>rodriguezii</i> Ortiz, 1968, <i>Culicoides</i>	14.35
<i>rostratus</i> Wirth & Blanton, 1956, <i>Culicoides</i>	14.20
<i>rozeboomi</i> Barbosa, 1947, <i>Culicoides</i>	14.25
<i>rubriceps</i> Kieffer, 1917, <i>Clinohelca</i>	14.47
<i>rufa</i> Kieffer, 1917, <i>Heteromyia</i>	14.48
<i>ruizi</i> Forattini, 1954, <i>Culicoides</i>	14.26
<i>rusticus</i> Macfie, 1939, <i>Atrichopogon</i>	14.16
<i>ryckmani</i> Wirth & Hubert, 1960, <i>Culicoides</i>	14.23
<i>saltanensis</i> Lane & Duret, 1954, <i>Clinohelca</i>	14.47
<i>saltensis</i> (Cavaliere), 1962 (<i>Lasiohelea</i>), <i>Forcipomyia</i>	14.7
<i>sanctaeclarae</i> Macfie, 1949, <i>Atrichopogon</i>	14.16
<i>sanctaeclarae</i> Wirth, 1952, <i>Forcipomyia</i>	14.11
<i>sanctaemarthae</i> Kieffer, 1917 (<i>Culicoides</i>), <i>Dasyhelea</i>	14.18
<i>sanctibernardini</i> Kieffer, 1917, <i>Stilobezzia</i>	14.43
<i>sanctilaurentii</i> Kieffer, 1917, <i>Atrichopogon</i>	14.16
<i>saundersi</i> Ewen, 1958, <i>Atrichopogon</i>	14.16
<i>scabra</i> (Coquillett), 1905 (<i>Ceratopogon</i>), <i>Palpomyia</i>	14.55
<i>schwarzii</i> (Coquillett), 1901 (<i>Ceratopogon</i>), <i>Nilobezzia</i>	14.50
<i>scissurae</i> Macfie, 1937, <i>Dasyhelea</i>	14.18
<i>scopus</i> Root & Hoffman, 1937, <i>Culicoides</i>	14.23
<i>scorzai</i> Ortiz, 1956, <i>Culicoides</i>	14.38
<i>scutata</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.46
<i>Selfia</i> Khalaf, 1954	14.36
<i>sentior</i> Ingram & Macfie, 1931, <i>Palpomyia</i>	14.55
<i>sequax</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Atrichopogon</i>	14.16
<i>setifer</i> Lutz, 1913 (<i>Centrorhynchus</i>), <i>Culicoides</i>	14.29
<i>setiger</i> Macfie, 1939, <i>Stenoxenus</i>	14.52
<i>setigera</i> Saunders, 1959, <i>Forcipomyia</i>	14.4
<i>setosa</i> Wirth, 1965, <i>Macrurohelea</i>	14.40
<i>setosierus</i> (Kieffer), 1906 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.6
<i>setulosa</i> (Loew), 1861 (<i>Ceratopogon</i>), <i>Bezzia</i>	14.53
<i>sexvittata</i> Wirth, 1956, <i>Forcipomyia</i>	14.6
<i>shannoni</i> Ingram & Macfie, 1931, <i>Dasyhelea</i>	14.18
<i>shannoni</i> (Ingram & Macfie), 1931 (<i>Lasiohelea</i>), <i>Forcipomyia</i>	14.12
<i>shannoni</i> Wirth & Blanton, 1972, <i>Monohelca</i>	14.40
<i>silvicola</i> Macfie, 1940, <i>Stilobezzia</i>	14.46
<i>similans</i> Lane & Forattini, 1956, <i>Stilobezzia</i>	14.46
<i>simplex</i> Lane & Forattini, 1958, <i>Stilobezzia</i>	14.46
<i>sinuosa</i> Dow & Wirth, 1972, <i>Forcipomyia</i>	14.11
<i>sitiens</i> Wirth & Hubert, 1960, <i>Culicoides</i>	14.23
<i>smartii</i> Macfie, 1950, <i>Echinohelea</i>	14.39
<i>snowi</i> Lane, 1958, <i>Bezzia</i>	14.53
<i>sonorensis</i> Wirth & Jones, 1957, <i>Culicoides</i>	14.27
<i>sonorensis</i> Wirth & Ratanaworabhan, 1972, <i>Paryphoconus</i>	14.51
<i>sordidipes</i> Macfie, 1939, <i>Palpomyia</i>	14.55
<i>sp. 2</i> , Fox, <i>Culicoides</i>	14.22
<i>sp. 3</i> , Fox, <i>Culicoides</i>	14.25
<i>spangleri</i> Wirth, 1965, <i>Parabezzia</i>	14.42
<i>spathicercus</i> Wirth, 1969, <i>Dasyhelea</i>	14.18
<i>spatulifera</i> Saunders, 1956, <i>Forcipomyia</i>	14.4
<i>spatuligera</i> Macfie, 1949, <i>Forcipomyia</i>	14.6
<i>Sphaeromiini</i>	14.48



spinifemur (Lane), 1948 (Dicrobezzia), Palpomyia	14.55
spinifemur Wirth & Blanton, 1972, Lanehelea	14.49
spinosa Lutz, 1914, Palpomyia	14.55
spinosa Saunders, 1956, Forcipomyia	14.13
spurius Wirth & Blanton, 1959, Culicoides	14.35
squamithorax Clastrier, 1972, Forcipomyia	14.6
squamitibia Lutz, 1914, Forcipomyia	14.6
squamosa Lutz, 1914, Forcipomyia	14.9
squamosa, authors, Forcipomyia	14.5
stellifer, authors, Culicoides	14.32
Stenoxcnini	14.50
Stenoxenus Coquillett, 1899	14.51
stigmalis Wirth, 1952, Culicoides	14.38
Stilobezzia Kieffer, 1911	14.42, 14.44
Stilobezziini	14.39
stonei Wirth, 1953, Monohelea	14.41
stubalensis Fox, 1946, Culicoides	14.32
stylifera (Lutz), 1913 (Centrorhynchus), Forcipomyia	14.7
subflavus Macfie, 1940, Paryphoconus	14.51
subfusca Macfie, 1939, Bezzia	14.53
subfuscula Ingram & Macfie, 1931, Palpomyia	14.55
subsessilis Kieffer, 1917, Stilobezzia	14.43
succinea Ingram & Macfie, 1931, Stilobezzia	14.43
Synthyridomyia Saunders, 1956	14.11
taeniatus Macfie, 1939, Atrichopogon	14.16
tamboensis Wirth & Lee, 1967, Culicoides	14.35
tamoioi Lane, 1960, Palpomyia	14.55
telmatophila Macfie, 1940, Bezzia	14.53
tenuicrus Kieffer, 1917, Palpomyia	14.55
tenuiforceps Macfie, 1939, Forcipomyia	14.11
tenuilobus Wirth & Blanton, 1959, Culicoides	14.35
tenuisquamipes Wirth, 1952, Forcipomyia	14.6
terminalis (Coquillett), 1904, Paryphoconus	14.51
terrestris Saunders, 1964, Forcipomyia	14.13
<i>Tersesthes</i> Townsend, 1893	14.2
<i>Tetraphora</i> Philippi, 1865	14.3
tetrathyris Wirth & Blanton, 1959, Culicoides	14.36
thalestris Macfie, 1935, Dasyhelea	14.18
thersites (Williston), 1896 (Ceratopogon), Atrichopogon	14.16
<i>thirsistes</i> , error	14.16
thoracica Ingram & Macfie, 1931, Macrurohelea	14.40
Thyridomyia Saunders, 1925	14.11
tibialis Lanc & Forattini, 1956, Stilobezzia	14.44
torridus Wirth & Hubert, 1960, Culicoides	14.23
townesi Lane, 1944, Clinohela	14.47
townsendi Knab, 1915, Forcipomyia	14.6
townsendi Lane, 1944, Clinohela	14.47
transferrans Ortiz, 1953, Culicoides	14.36
transversa Lane & Forattini, 1958, Stilobezzia	14.46
travassosi Forattini, 1957, Culicoides	14.26
travassosi Lane, 1956, Paryphoconus	14.51
travassosi Lane, 1947, Stilobezzia	14.46
<i>Trichohelea</i> , authors	14.9
Trichohelea Goetghebuer, 1920	14.12
<i>tricoloratus</i> Wirth & Blanton, 1953, Culicoides	14.35
trilineatus Fox, 1946, Culicoides	14.36



<i>trinidadensis</i> Hoffman, 1925, <i>Culicoides</i>	14.26
<i>trinidadensis</i> Macfie, 1937, <i>Atrichopogon</i>	14.16
<i>trinidadensis</i> Saunders, 1964, <i>Forcipomyia</i>	14.12
<i>tristriatulus</i> , authors, <i>Culicoides</i>	14.21
<i>tropicus</i> Kieffer, 1917 (<i>Ceratopogon</i>), <i>Forcipomyia</i>	14.8
<i>tuberculata</i> Saunders, 1956, <i>Forcipomyia</i>	14.13
<i>tuberculatus</i> Ewcn, 1948, <i>Atrichopogon</i>	14.16
<i>turgidipes</i> Ingram & Macfie, 1931, <i>Neurohelea</i>	14.48
<i>umbella</i> Macfie, 1939, <i>Palpomyia</i>	14.55
<i>umbratilis</i> Macfie, 1935, <i>Atrichopogon</i>	14.16
<i>undecimpunctatus</i> Kieffer, 1917, <i>Culicoides</i>	14.34
<i>unguis</i> Wirth, 1965, <i>Parabozzia</i>	14.42
<i>unimaculatus</i> Macfie, 1940, <i>Paryphoconus</i>	14.51
<i>uniradialis</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.36
<i>univesicula</i> Macfie, 1939, <i>Forcipomyia</i>	14.11
<i>urraeaisi</i> Lanc & Wirth, 1964, <i>Monohela</i>	14.41
<i>uruguayensis</i> (Cordero), 1929 (<i>Lophomyidium</i>), <i>Atrichopogon</i>	14.16
<i>utae</i> Knab, 1915, <i>Forcipomyia</i>	14.6
<i>utricularius</i> Macfie, 1953, <i>Atrichopogon</i>	14.16
<i>vargasi</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.22
<i>varia</i> Ingram & Macfie, 1931, <i>Stilobezzia</i>	14.44
<i>varicolor</i> Saunders, 1956, <i>Forcipomyia</i>	14.4
<i>variipennis</i> (Coquillett), 1901 (<i>Ceratopogon</i>), <i>Culicoides</i>	14.27
<i>variipennis</i> , authors, <i>Culicoides</i>	14.37
<i>variipennis</i> Wirth & Williams, 1957, <i>Forcipomyia</i>	14.6
<i>venezuelensis</i> Ortiz, 1952, <i>Leptoconops</i>	14.3
<i>venezuelensis</i> Ortiz, 1950, <i>Stilobezzia</i>	14.46
<i>venezuelensis</i> Ortiz & Mirsa, 1950, <i>Culicoides</i>	14.36
<i>venustula</i> (Williston), 1896 (<i>Ceratopogon</i>), <i>Bezzia</i>	14.53
<i>verecundus</i> Macfie, 1948, <i>Culicoides</i>	14.26
<i>versicolor</i> Macfie, 1939, <i>Palpomyia</i>	14.55
<i>villosipes</i> (Kieffer), 1917 (<i>Culicoides</i>), <i>Dasyhelea</i>	14.18
<i>viridans</i> Forattini & Rabello, 1957, <i>Dasyhelea</i>	14.18
<i>wagneri</i> Séguy, 1941 (<i>Phasmodhelea</i>), <i>Forcipomyia</i>	14.8
<i>Warmkea</i> Saunders, 1956	14.12
<i>warmkei</i> Wirth, 1956, <i>Atrichopogon</i>	14.16
<i>willistoni</i> Wirth, 1971, <i>Forcipomyia</i>	14.9
<i>willistoni</i> Wirth & Blanton, 1953, <i>Culicoides</i>	14.27
<i>wirthi</i> Lane, 1961, <i>Paryphoconus</i>	14.51
<i>wirthi</i> (Lanc), 1956 (<i>Macfichelea</i>), <i>Pellucidomyia</i>	14.48
<i>wirthi</i> Lane & Forattini, 1956, <i>Stilobezzia</i>	14.46
<i>wirthi</i> Lane, Forattini & Rabello, 1955, <i>Palpomyia</i>	14.55
<i>wirthomyia</i> Vargas, 1953, <i>Culicoides</i>	14.23
<i>wokei</i> Barbosa, 1947, <i>Culicoides</i>	14.26
<i>wokei</i> Fox, 1947, <i>Culicoides</i>	14.36
<i>woodfordi</i> Macfie, 1938, <i>Atrichopogon</i>	14.16
<i>wygodzinskyi</i> Cavalleri, 1960, <i>Forcipomyia</i>	14.6
<i>wygodzinskyi</i> Lane, 1948, <i>Clinohela</i>	14.47
<i>wygodzinskyi</i> Lane, 1946, <i>Paryphoconus</i>	14.51
<i>wygodzinskyi</i> Lane, 1947, <i>Stilobezzia</i>	14.46
<i>xanifer</i> Wirth & Blanton, 1968, <i>Culicoides</i>	14.26
<i>yukonensis</i> , authors, <i>Culicoides</i>	14.21
<i>zeteki</i> Wirth & Dow, 1972, <i>Forcipomyia</i>	14.11
<i>zonogaster</i> Ingram & Macfie, 1931, <i>Forcipomyia</i>	14.6



IMPrensa OFICIAL DO ESTADO
SÃO PAULO - BRASIL
1974





SciELO



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

16

FAMILY SIMULIIDAE

M. A. VULCANO

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

Adult simuliids are commonly known as *jejenes*, *mosquitos negritos*, *alazanes*, etc. (Spanish-speaking countries), and *piuns* or *borrachudos* (Brazil). Their biology is not yet sufficiently known in the Neotropical region, and many obscure points are awaiting a more thorough study. The females, after a blood-meal, search for an appropriate place for oviposition, usually in quiet places of rivers and streams. The larvae need well-aerated running waters and are apneustic, although they show non-functional respiratory stigmata in the thorax and in each abdominal segment. Pupae are usually enclosed in silken cocoons spun by the last-instar larvae, but some instances exist of free pupae. Only adult females are hematophagous, being strictly diurnal in habit.

As these flies are vectors of several diseases (e.g., onchocerciasis, *Leucocytozoon* infections), and due to their immense numbers, being a plague to man and domestic animals, several studies from the public health viewpoint were undertaken in several countries.

Some 15 genera and more than 1,000 species exist in the world; the Neotropical region contributes to that amount with 235 species, distributed in 5 genera.

Genus AUSTROSIMULIUM Tonnoir

AUSTROSIMULIUM Tonnoir, 1925:230. Type-species, *Simulium australense* Schiner (orig. des.).

Subgenus PARAUSTROSIMULIUM Wygodzinsky & Coscarón

Austrosimulium, subg. PARAUSTROSIMULIUM Wygodzinsky & Coscarón, 1962:244. Type-species, *Simulium anthracinum* Bigot (mon.).

ANTHRACINUM (Bigot), 1888:15 (*Simulium*). Type-locality: Argentina, Tierra del Fuego, Orange Bay. Distr. — Chile (Aysén, Magallanes, Chiloe I.), Argentina (Rio Negro, Tierra del Fuego). Ref. — Wygodzinsky & Coscarón, 1962:242, figs. 1-2, 3a, 3l-o, 4a-1, 4l-o.

moorei Silva Figueroa, 1917:30, pl. 1, figs. 16-18 (*Simulia*). Type-locality: Chile, Aysén, Península Taitao.

16.1

Genus CNEPHIA Enderlein

- CNEPHIA Enderlein, 1921a:199. Type-species, *Simulium pecuarum* Riley (orig. des.).
- AGUIRREI (Dalmat), 1949:544, figs. 10-18 (*Simulium*). Type-locality: Guatemala, Acatenango, Río San Diego, Finca San Diego. Distr. — Mexico (Chiapas), Guatemala (Chimaltenango, Acatenango). Refs. — Dalmat, 1951:48; Vargas & Díaz Nájera, 1952:53, 4 figs.
- ATROPARVA (Edwards), 1931:148, fig. 13f (*Simulium*). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Neuquén), Chile. Ref. — Vargas & Díaz Nájera, 1951:165, figs. 56-58.
- ATZOMPENSIS Díaz Nájera, 1962:273, pl. 2, figs. 6-10. Type-locality: Mexico, Morelos, Cañada de Atzompa. Distr. — Mexico (Morelos).
- DISSIMILIS (Edwards), 1931:145, figs. 8a, 10g, 12f, 13e (*Simulium*). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Neuquén), Chile (Llanquihue). Ref. — Coscarón & Wygodzinsky, 1962:310, figs. 126-184.
- GRENIERI Vargas & Díaz Nájera, 1948b:325, figs. 12-24. Type-locality: Mexico, Veracruz, Tlanepaquila, 1500 m. Distr. — Mexico (Veracruz, Oaxaca).
- GYNANDRA (Edwards), 1931:147 (*Simulium*). Type-locality: Argentina, Río Negro, L. Correntoso. Distr. — Argentina (Río Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:166, figs. 59-65.
- MIXENSIS Díaz Nájera, 1962:274, pl. 4, figs. 16-20. Type-locality: Mexico, Oaxaca, Totontepec Mixe. Distr. — Mexico (Oaxaca).
- MONTANA (Philippi), 1865:633 (*Simulium*). Type-locality: Chile, Chacabuco, Catemú. Distr. — Chile.
- PACHECOLUNAI (León), 1945:67, fig. 3 (*Simulium*; as *pacheco-lunai*). Type-locality: Guatemala, Sololá, Totonicapán, Cumbre María Tecún. Distr. — Mexico (Chiapas), Guatemala (Sololá, Totonicapán, Chimaltenango, El Quiché). Ref. — Vargas & Díaz Nájera, 1954:67, figs. 17-30.
- ROBLESI (León), 1943-97, fig. 1 (*Simulium*). Type-locality: Guatemala, between Sololá and Totonicapán, María Tecún. Distr. — Guatemala (San Marcos, Quezaltenango, Huehuetenango, Sololá, Totonicapán). Ref. — Vargas & Díaz Nájera, 1954:64, figs. 40-54.

Genus GIGANTODAX Enderlein

- GIGANTODAX Enderlein, 1925:205. Type-species, *bolivianum* Enderlein (orig. des.).
- ABALOSI Wygodzinsky, 1958:143, figs. 109-141. Type-locality: Peru, Chacamasa, 2100 m. Distr. — Peru.
- ANTARCTICUM (Bigot), 1888:15 (*Simulium*). Type-locality: Argentina, Tierra del Fuego, Orange Bay. Distr. — Argentina (Neuquén, Río Negro, Tierra del Fuego), Chile (Magallanes). Refs. — Edwards, 1931:132, figs. 12b, 12g; Smart, 1945:499.
- AQUAMARENSE (León), 1945:72, figs. 10-11 (*Simulium*). Type-locality: Guatemala, Cuenca Agua Amarga, 1800 m. Distr. — Guatemala (Que-



- zaltenango). Ref. — Dalmat, 1955:102, figs. 10-12, 127, 129, 279, 319, 359, 396.
- ARAUCANUM (Edwards), 1931:142, fig. 13d (*Simulium*). Type-locality: Chile, [Llanquihue, Chiloe I.], Ancud. Distr. — Chile (Llanquihue, Chiloe I.), Argentina (Rio Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:155, figs. 31-36.
- BOLIVIANUM Enderlein, 1925:205. Type-locality: Bolivia, Cucsta de Cillunticara, 3500 m. Distr. — Peru, Bolivia. Ref. — Wygodzinsky, 1958:140, figs. 87-108.
- BONORINORUM Coscarón & Wygodzinsky, 1962:284, figs. 1-53 (as *bonorinum*). Type-locality: Argentina, Tierra del Fuego, Ushuaia. Distr. — Argentina (Tierra del Fuego).
- BROPHYI (Edwards), 1931:133, figs. 10c, 10d, 13b (*Simulium*). Type-locality: Argentina, Rio Negro, L. Nahuel Huapí, eastern end. Distr. — Argentina (Rio Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:156, figs. 37-39.
- CHILENSE (Philippi), 1865:634 (*Simulium*). Type-locality: Chile, Valdivia. Distr. — Chile (Valdivia, Llanquihue, Chiloe I.), Argentina (Rio Negro, Neuquén). Ref. — Edwards, 1931:139, fig. 12d
- FEMINEUM (Edwards), 1931:135, figs. 7b, 11a-b, 12c, 12e, 12h (*Simulium*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Chile (Llanquihue, Chiloe I.), Argentina (Rio Negro, Neuquén).
- FULVESCENS (Blanchard) in Gay, 1852:353 (*Simulium*). Type-locality: Chile, Coquimbo. Distr. — Chile (Chiloe I., Concepción, Llanquihue, Coquimbo), Argentina (Rio Negro, Neuquén). Ref. — Edwards, 1931:137, figs. 10f, 12a.
- HORCOTIANI Wygodzinsky, 1949b:326, pls. 1-3. Type-locality: Argentina, Tucumán, Tafi del Valle. Distr. — Argentina (Tucumán). Ref. — Wygodzinsky, 1951a:199, figs. 1-26.
- IGNICULUM Coscarón & Wygodzinsky, 1962:296, figs. 71-107. Type-locality: Argentina, Tierra del Fuego, Ushuaia. Distr. — Argentina (Tierra del Fuego).
- KUSCHELI Wygodzinsky, 1952:81, figs. 1-23. Type-locality: Chile, Juan Fernandez Is., Masafuera, Qucbrada de las Casas. Distr. — Juan Fernandez Is. Ref. — Coscarón & Wygodzinsky, 1962:308, figs. 112-125.
- MARGINALE (Edwards), 1931:140, fig. 13c (*Simulium*). Type-locality: Chile, [Llanquihue, Chiloe I.], Ancud. Distr. — Chile (Llanquihue, Chiloe I.), Argentina (Rio Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:160, figs. 44-47.
- NIGRESCENS (Edwards), 1931:133, fig. 13a (*Simulium*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Argentina (Rio Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:161.
- PENNIPUNCTUM Enderlein, 1933:274. Type-locality: Peru, Cuzco. Distr. — Peru (Cuzco).
- RUFESCENS (Edwards), 1931:134, figs. 10a-b (*Simulium*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Argentina (Rio Negro, Neuquén), Chile.
- RUFINOTUM (Edwards), 1931:140, fig. 10e (*Simulium*). Type-locality: Argentina, Rio Negro, L. Gutierrez. Distr. — Argentina (Rio Negro, Neuquén), Chile (Llanquihue, Chiloe I.). Ref. — Vargas & Díaz Nájera, 1951:162, figs. 48-50.



- SHANNONI (Edwards), 1931:137 (*Simulium*). Type-locality: Argentina, Rio Negro, L. Correntoso. Distr. — Argentina (Rio Negro, Neuquén).
- WITTMER Wygodzinsky, 1951a:202, figs. 27-44. Type-locality: Argentina, Rio Negro, San Carlos de Bariloche, Arroyo de la Cascata, "arriba del Hotel Cerro Catedral". Distr. — Argentina (Rio Negro, Tierra del Fuego). Ref. — Coscarón & Wygodzinsky, 1962:307, figs. 108-111.
- WRIGHT (Vargas, Martínez Palacios & Díaz Nájera), 1944:37, pl. 1, figs. 1-3, pl. 2, figs. 4-7 (*Simulium*). Type-locality: Mexico, D.F., Desierto de los Leones, 3200 m. Distr. — Mexico (D.F., Chiapas, Oaxaca, Veracruz), Guatemala (Chimaltenango, El Quiché, San Marcos, Sololá, Totonicapán, Huehuetenango, Suchitepéquez). Refs. — Vargas, Martínez Palacios & Díaz Nájera, 1946:165, figs. 138, 147; M.A.V. d'Andretta & C. d'Andretta, 1947a:23, 32 figs.
- vargasi* León, 1945:69, figs. 5-6. Type-locality: Guatemala, Totonicapán, Cumbre María Tecún.

Genus LUTZSIMULIUM C. d'Andretta & M. A. V. d'Andretta

- LUTZSIMULIUM C. d'Andretta & M.A.V. d'Andretta, 1947:401 (1948a:668). Type-species, *cruzi* C. d'Andretta & M.A.V. d'Andretta (orig. des.) = *hirticosta* (Lutz). N. SYN.
- HIRTICOSTA (Lutz), 1909:135 (*Simulium*). Type-locality: Brazil, São Paulo, Serra da Bocaina [Faz. do Bonito]. Distr. — Brazil (São Paulo, Rio de Janeiro, Paraná). Ref. — Lutz, 1910:226, fig. 16.
- cruzi* C. d'Andretta & M.A.V. d'Andretta, 1947:401, figs. 1-35 (1948a:668). Type-locality: Brazil, São Paulo, Bragança Paulista. Ref. — C. d'Andretta & M.A.V. d'Andretta, 1948b:639, figs. 1-38. N. SYN.
- PERNIGRUM (Lutz), 1910:263, figs. 19-20 (*Simulium*; nom. nov. for *montanum* of Lutz). Type-locality: Brazil, São Paulo, Serra da Bocaina [Faz. do Bonito]. Distr. — Brazil (Rio de Janeiro, São Paulo). Ref. — M.A.V. d'Andretta & C. d'Andretta, 1952:312, figs. 1-49. N. COMB.
- montanum* Philippi of Lutz, 1909:134 (*Simulium*).

Genus SIMULIUM Latreille

- SIMULIUM Latreille, 1802:426. Type-species, *Rhagio colombaschensis* Fabricius (mon.).
- Melusina*, authors, not Meigen.
- Simulia*, emend.
- Simulium*, subg. *Eusimulium* Roubaud, 1906a:521. Type-species, *aureum* Fries (mon.).
- Chirostilbia* Enderlein, 1921a:199. Type-species, *flavifemur* Enderlein (orig. des.).
- Wilhelmia* Enderlein, 1921a:199. Type-species, *Atractocera lineata* Meigen (orig. des.).



- Friesia* Enderlein, 1922:69 (preocc. McDunnough, 1912). Type-species, *Neocrmannia tristigata* Enderlein (orig. des.).
- Byssodon* Enderlein, 1925:209. Type-species, *Simulium forbesi* Malloch (orig. des.) = *meridionale* Riley.
- Pternaspatha* Enderlein, 1930:88. Type-species, *nigristrigata* Enderlein (orig. des.).
- Notolepria* Enderlein, 1930:95. Type-species, *Simulium exiguum* Roubaud (orig. des.).
- Dasypelmoza* Enderlein, 1933:375. Type-species, *Simulium varipes* Philippi (orig. des.).
- Acropogon* Enderlein, 1933:276. Type-species, *barbatipes* Enderlein (orig. des.).
- Ectemnaspis* Enderlein, 1933:281. Type-species, *maeca* Enderlein (orig. des.) = *bicoloratum* Malloch.
- Psilopelmia* Enderlein, 1933:283. Type-species, *rufidorsum* Enderlein (orig. des.) = *escomeli* Roubaud.
- Thyrsopelma* Enderlein, 1933:284. Type-species, *brasiliensis* Enderlein (orig. des.).
- Trichodagnia* Enderlein, 1933:288. Type-species, *latitarsis* Enderlein (orig. des.) = *chalcocoma* Knab.
- Hemicnetha* Enderlein, 1934:190. Type-species, *mexicana* Enderlein (orig. des.) = *paynei* Vargas.
- Psaroniocompsa* Enderlein, 1934:192. Type-species, *opaliniifrons* Enderlein (orig. des.).
- Cryptectemnia* Enderlein, 1936:114. Type-species, *latialx* Enderlein (orig. des.) = *orsovae* Smart.
- Pliodasina* Enderlein, 1936:124. Type-species, *guttata* Enderlein (orig. des.).
- Simulium*, subg. *Hearlea* Rubzov, 1940:116, 126. *Nomen nudum*, type-species as below but not described.
- Simulium*, subg. *Hearlea* Vargas, Martínez Palacios & Díaz Nájera, 1946:104, 106. Type-species, *canadense* Hearle (orig. des.).
- Simulium*, subg. *Neosimulium* Rubzov, 1940:116, 124, 130. *Nomen nudum*, type-species as below but not described.
- Simulium*, subg. *Neosimulium* Vargas, Martínez Palacios & Díaz Nájera, 1946:103, 108. Type-species, *vittatum* Zetterstedt (orig. des.).
- Simulium*, subg. *Lanea* Vargas, Martínez Palacios & Díaz Nájera, 1946:103, 107. Type-species, *haematopotum* Malloch (orig. des.).

Different opinions exist concerning the arrangement of the species of *Simulium* into subgenera, but none so far definite, and for this reason we have not adopted subgeneric divisions, but only listed above the different names proposed as subgenera. Refs. — Vargas, 1945c (morphol., biol., anat., catal.); Vargas & Díaz Nájera, 1957b (rev., Mexican species); Stone, 1963 (annotated list of genus-group names).

ADOLFOLUTZI Wygodzinsky, 1951c:234, figs. 1-5, 50, 52, 98-103. Type-locality: Argentina, Tucumán, Raco, 800 m. Distr. — Argentina (Salta,




- Tucumán, Catamarca).
- ALBICINCTUM (Enderlein), 1933:277 (*Acropogon*). Type-locality: "Upper Peru". Distr. — Peru.
- ALBILINEATUM (Enderlein), 1936:115 (*Pternaspatha*). Type-locality: Peru, Callanga. Distr. — Peru.
- ALBOPICTUM Lane & Porto, 1940:193. Type-locality: Brazil, Paraná, Cambará. Distr. — Brazil (Paraná). Ref. — Lane & Vulcano, 1943: 438, fig. 21.
- ALTICOLUM (Dyar & Shannon), 1927:24, pl. 4, fig. 47 (*Eusimulium*). Type-locality: Mexico, Chihuahua, Sierra Madre, Río Piedras Verdes. Distr. — Mexico (Chihuahua).
- AMAZONICUM Goeldi, 1905:138. Type-locality: Brazil, Amazonas, Tefé, Purus, and Acre. Distr. — Puerto Rico, Venezuela, British Guiana, Ecuador, Peru, Brazil (Amazonas and Acre, south to São Paulo), Paraguay. Refs. — Smart, 1942:46, fig. 1; Briceño-Irrogorry, Vargas & Díaz Nájera, 1953:243, 1 pl.
- exiguum* Roubaud of Lutz, 1910:234, 266.
- minusculum* Lutz, 1910:253, figs. 31-32, 34. Type-locality: Brazil, Minas Gerais, Lassance.
- minutum* Surcouf & González Rincones, 1911:290 (nom. nov. for *exiguum* of Lutz; preocc. Lugger, 1897).
- nitidum* Malloch, 1912:652. Type-locality: Peru, Huancabamba.
- lutzi* Knab, 1913:154 (nom. nov. for *minutum* Surcouf & González Rincones).
- lutzi* Malloch, 1914:14 (nom. nov. for *minutum* Surcouf & González Rincones; preocc. Knab, 1913).
- ANAMARIAE Vulcano, 1962:246, figs. 1-55, 59 (as *ana-mariae*). Type-locality: Brazil, São Paulo, Serra da Bocaina, Fazenda do Bonito. Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo).
- ANDUZEI Vargas & Díaz Nájera, 1948b:328, figs. 25-31. Type-locality: Mexico, Veracruz, Jalacingo, 1980 m. Distr. — Mexico (D.F., Veracruz, Oaxaca). Ref. — Vargas & Díaz Nájera, 1949:295, figs. 38, 40.
- ANGRENSE Pinto, 1931:683, figs. 9, 14-15. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis. Distr. — Brazil (Rio de Janeiro).
- ANNULATUM Philippi, 1865:634. Type-locality: Chile, Santiago, Valle San Ramón. Distr. — Chile (Santiago, Tacna).
- ANTILLARUM Jennings, 1915:200. Type-locality: West Indies, St. Croix and Jamaica, Hope River near Kingston and Roaring River. Distr. — St. Croix, Jamaica, Guadeloupe, Cuba, Mexico (Chiapas), Venezuela. Ref. — Díaz Nájera, 1961:79, figs. 1-26.
- ANTONII Wygodzinsky, 1953b:325, figs. 30-43. Type-locality: Bolivia, Cochabamba, Chapare, 3150 m. Distr. — Bolivia.
- ANTUNESI (Lane & Porto), 1940:189, fig. 1 (*Eusimulium*). Type-locality: Brazil, Bahia, Pirajá. Distr. — Brazil (Bahia). Ref. — Lane & Vulcano, 1943:436, fig. 13.
- ARGENTATUM (Enderlein), 1936:125 (*Thyrsopelma*). Type-locality: "Upper Peru". Distr. — Peru.
- ARGUS Williston, 1893:253. Type-locality: U.S.A., California, Argus Mountain. Distr. — Canada, U.S.A. (California), Mexico (D.F., Durango, Guanajuato, Guerrero, Hidalgo, Jalisco, Michoacán, Morelos, Nuevo León, Oaxaca, Puebla, San Luis Potosí, Sonora, Veracruz). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:171, figs. 134, 154.

- kamloopsi* Hearle, 1932:12, fig. A. Type-locality: Canada, British Columbia, Kamloops.
- AUREUM (Fries), 1824:16 (*Simulia*). Type-locality: Scandinavia, Esperöd. Distr. — Europe; Mexico (Chiapas, Durango, Guanajuato, Michoacán, Morelos, Veracruz), Guatemala (Chimaltenango, Huehuetenango, Quezaltenango, El Quiché, San Marcos, Sololá, Totonicapán). Refs. — Dalmat, 1955:116, figs. 8_s, 19-21, 136-138, 268, 282, 322, 362, 369, 399; Vargas & Díaz Nájera, 1957b:164-166, figs. 45-53.
- donovani* Vargas, 1943b:359, figs. 34-36. Type-locality: Mexico, Chiapas, Pueblo de Chamula, 2350 m.
- diazii* León, 1945:5, fig. 7. Type-locality: Guatemala, María Tecún.
- AURPELLITUM Enderlein, 1933:286. Type-locality: "Paraguay, 'hothenau'". Distr. — Paraguay.
- AURISTRIATUM Lutz, 1910:245. Type-locality: Brazil, Minas Gerais, Juiz de Fora and Alfenas; São Paulo, Serra da Bocaina. Distr. — Brazil (Minas Gerais, São Paulo, Mato Grosso, Rio de Janeiro, Guanabara). Ref. — M.A.V. d'Andretta & C. d'Andretta, 1949:55, figs. 1-42.
- infuscatum* Lutz, 1910:236 (*nec* Lutz, 1909). Type-locality: Brazil, São Paulo, Serra da Bocaina, Fazenda do Bonito, 1600 m., and Itanhaem; Rio de Janeiro, Petrópolis, Caxambu de Cima.
- AYROSAI Vargas, 1945a:74, pl. 7, fig. 4, pl. 8, figs. 4, 7, pl. 9, fig. 5, pl. 10, figs. 3-4. Type-locality: Mexico, D.F., Desierto de los Leones. Distr. — Mexico (México, D.F., Puebla). Refs. — Vargas, Martínez Palacios & Díaz Nájera, 1946:106, 111, 164, 181, 184, 185, fig. 125; Vargas & Díaz Nájera, 1957b:166, 167, 170, 173, figs. 65-72.
- BAIENSE Pinto, 1931:685, figs. 16-18. Type-locality: Brazil, Bahia, Nazaré. Distr. — Brazil (Bahia).
- BARBATIPES (Enderlein), 1933:277 (*Acropogon*). Type-locality: Bolivia, La Paz, Luribay. Distr. — Peru, Bolivia. Ref. — Wygodzinsky, 1953b:321, 29 figs.
- BENJAMINI Dalmat, 1952:339, figs. 1-7. Type-locality: Guatemala, El Petén, Río Sokchá, Poctún. Distr. — Guatemala (El Petén, Alta Verapaz). Ref. — Dalmat, 1955:121, figs. 8-9, 22-24, 139-141, 275, 283, 325.
- BICOLORATUM Malloch, 1913:649, pl. 46, figs. 6-7. Type-locality: Peru, Río Charape. Distr. — Peru, Venezuela, Bolivia, Colombia. Ref. — Wygodzinsky, 1953b:327, figs. 44-58.
- macca* Enderlein, 1933:281 (*Ectemnaspis*; preocc. Enderlein, 1933:276, as *Dasytelmoxa*). Type-locality: "Peru".
- mollis* Vargas, 1943a:142 (nom. nov. for *Ectemnaspis macca* Enderlein).
- BIVITTATUM Malloch, 1914:31, pl. 5, fig. 7. Type-locality: U.S.A., New Mexico, Las Vegas. Distr. — U.S.A. (New Mexico), Mexico (Zacatecas). Ref. — Vargas & Díaz Nájera, 1958:13, 15, 17, 18, 20, 22, figs. 1-16.
- BOTULIBRANCHIUM Lutz, 1910:256, 263, fig. 1. Type-locality: Brazil, Rio de Janeiro, Morro do Observatório, Petrópolis. Distr. — ?Venezuela (D.F.), Brazil (Minas Gerais, Rio de Janeiro, São Paulo). Ref. — M.A.V. d'Andretta & C. d'Andretta, 1947b:145, figs. 1-48.
- tulibranchium* Lutz, 1922:92 (error).
- BRACHYCLADUM Lutz & Pinto *in* Pinto, 1931:690, figs. 19-25. Type-locality: Brazil, Pernambuco, Tapera, and Bahia, São Félix. Distr. — Brazil (Pernambuco, Bahia).

- BREVI-FURCATUM Lutz, 1910:259, 263, figs. 14-15. Type-locality: Brazil, São Paulo, Serra da Bocaina, Fazenda do Bonito. Distr. — Brazil (Rio de Janeiro, São Paulo, Paraná), Argentina (Misiones). Ref. — Vulcano, 1962:255, figs. 56, 58, 60-95.
- aequifurcatum* Lutz, 1910:259. Type-locality: unknown [Brazil].
- BRICENOI Vargas, Martínez Palacios & Díaz Nájera, 1946:115, figs. 8-15 (as *bricenoi*). Type-locality: Mexico, México, Dos Ríos, 2645 m. Distr. — Mexico (México, D.F., Durango, Guanajuato, Veracruz). Ref. — Vargas & Díaz Nájera, 1957b:174, 175, 177-179, figs. 179-187.
- BURCHI Dalmat, 1951:42, figs. 16-23. Type-locality: Guatemala, El Quiché, Nebaj, Río Micovez. Distr. — Guatemala (El Quiché, Huehuetenango). Ref. — Dalmat, 1955:239, figs. 94-96, 214-216, 251, 308, 348, 386, 423.
- BUSTOSI Vargas, Martínez Palacios & Díaz Nájera, 1946:138, figs. 64-70, 120. Type-locality: Mexico, Veracruz, Cruz Blanca, Acajete. Distr. — Mexico (Veracruz, Oaxaca). Ref. — Vargas & Díaz Nájera, 1957b:190, 193, 194, 196, 198, 247, 260, 264, figs. 393-400.
- CALLIDUM (Dyar & Shannon), 1927:12-13, 16, fig. 41 (*Eusimulium*; nom. nov. for *ochraceum* of Malloch). Type-locality: Mexico, Veracruz, Córdoba. Distr. — Mexico (Chiapas, Colima, Guerrero, Hidalgo, Jalisco, México, Michoacán, Morelos, Oaxaca, Quintana Roo, Puebla, San Luis Potosí, Veracruz), Guatemala (Chimaltenango, Sololá, Quezaltenango, San Marcos, Altaverapaz, Bajaverapaz, El Progreso, Santa Rosa, Jutiapa, Escuintla, El Quiché, Suchitepéquez, Izabal, Zacapa, Retalhuleu, Chiquimula, Huehuetenango), British Honduras. Refs. — Vargas, 1942a:231, figs. 1, 8; Vargas, 1943b:363, figs. 28-29; Vargas, Martínez Palacios & Díaz Nájera, 1946:102, 107, 112, 162, 166, 184, 187.
- ochraceum* Walker of Malloch, 1914:30, pl. 2, fig. 5, pl. 5, fig. 8.
- mooseri* Dampf, 1927:127, figs. 1-3a (*Eusimulium*). Type-locality: Mexico, Oaxaca, Tiltepec.
- CANADENSE Hearle, 1932:14 (as *virgatum* ssp.). Type-locality: Canada, British Columbia, Kamloops. Distr. — Canada (British Columbia), U.S.A., Mexico (D.F., Durango, Mexico, Oaxaca, Puebla, Veracruz). Refs. — Hearle, 1929:48 (as *virgatum*), figs. A-G; Hearle, 1935:15, figs. 1, 4.
- virgatum* Coquillett of Malloch, 1914:48, figs. A-G.
- CAPRICORNE León, 1945:71, figs. 8-9 (as *capricornis*). Type-locality: Guatemala, Totonicapán, and Río Samalá, Km 169 "proximo a la ciudad". Distr. — Mexico (Chiapas, Veracruz), Guatemala (Chimaltenango, Totonicapán, Guatemala, Sacatepéquez, Altaverapaz, El Quiché, Sololá, Huehuetenango). Ref. — Dalmat, 1955:244, figs. 97-99, 217-219, 252, 309, 349, 387, 424.
- CAROLINAE León, 1945:73, figs. 12-13. Type-locality: Guatemala, Quezaltenango, Salto de Zunil, 1900 m. Distr. — Mexico (Chiapas, Oaxaca, Veracruz), Guatemala (Chimaltenango, Guatemala, Altaverapaz, Bajaverapaz, Sololá, Suchitepéquez). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:122, figs. 29-34.
- CATARINENSE Pinto, 1931:695, figs. 26-29. Type-locality: Brazil, Santa Catarina, São Bento. Distr. — Brazil (Santa Catarina).
- CAUCHENSE Floch & Abonnenc, 1946c:1, figs. 1-2. Type-locality: French Guiana, crique Patawa, "bassin de Caux e crique Blanche, bassin d'Iracoubou". Distr. — French Guiana.

- CHALCOCOMA Knab, 1914c:85. Type-locality: Peru, Tincocochoca, 210 m. Distr. — Peru, Bolivia. Refs. — Vargas & Díaz Nájera, 1951:142, figs. 74-75; Vargas & Díaz Nájera, 1953b:139.
- latitarsis* Enderlein, 1933:288 (*Trichodagnia*). Type-locality: Peru, Callanga, Mamara, Arequipa and Chanchamayo; Bolivia, Mapiri, Lorenzopata and Sarapiuni.
- chalcocomense* Vargas, 1945c:125 (error).
- CHILIANUM Philippi in Rondani, 1864:90. Type-locality: "Chile". Distr. — Chile.
- CLARKI Fairchild, 1940:710, figs. 10, 18, 25, 30. Type-locality: Panama, Coclé, El Valle River. Distr. — Panama (Cocié, Darién), Venezuela (Miranda). Ref. — Vargas & Díaz Nájera, 1948a:75, figs. 1-2.
- CLAVIBRANCHIUM Lutz, 1910:257. Type-locality: Brazil, Rio de Janeiro, Petrópolis, Morro do Observatório, 1800 m. and Estação Capivari (between Barra Mansa and Angra dos Reis); São Paulo, Serra da Bocaina, [Fazenda do Bonito], 1200 m. Distr. — Brazil (Rio de Janeiro, São Paulo).
- CONTRERENSE Díaz Nájera & Vulcano, 1962:94, figs. 15-45. Type-locality: Mexico, D.F., Los Dinamos Contreras. Distr. — Mexico (Morelos, Oaxaca).
- COSTALIMAI Vargas, Martínez Palacios & Díaz Nájera, 1946:108, 111, 140-142, 151, 162, 172-173, 185, 187, figs. 71-73. Type-locality: Mexico, Morelos, Lagunas de Zempoala. Distr. — Mexico (Morelos, Oaxaca).
- jacumbae* Dyar & Shannon of Vargas, Díaz Nájera & Martínez Palacios, 1943:289, figs. 6-7.
- COSTARICENSE Smart, 1944:132 (nom. nov. for [*Acropogon*] *rufidorsum* Enderlein; as *costaricensis*). Type-locality: "Costa Rica". Distr. — Costa Rica, West Indies, Peru.
- rufidorsum* Enderlein, 1936:119 (*Acropogon*; preocc. Enderlein, 1933 [*Psilopelmia*]).
- costaricence* Vargas, 1945b:5 (nom. nov. for *rufidorsum* Enderlein; preocc. Smart, 1944).
- CUNEATUM (Enderlein), 1936:124 (*Psaroniocompsa*). Type-locality: "Paraguay". Distr. — Paraguay.
- DALMATI Vargas & Díaz Nájera, 1948b:337, figs. 47-56. Type-locality: Mexico, Veracruz, Cumbres de Acultzingo. Distr. — Mexico (Veracruz, Morelos). Ref. — Díaz Nájera & Vulcano, 1962:92, figs. 46-56.
- DANDRETTAI Vargas, Martínez Palacios & Díaz Nájera, 1946:124, figs. 35-40 (as *dandrettai*). Type-locality: Mexico, México, La Cima Salazar, 3005 m. Distr. — Mexico (México, D.F., Oaxaca, Veracruz). Ref. — Vargas & Díaz Nájera, 1948b:342, figs. 62-63.
- DEACOSTINI Coscarón & Wygodzinsky, 1962:322, figs. 185-236. Type-locality: Argentina, Tierra del Fuego, Tierra Mayor. Distr. — Argentina (Tierra del Fuego).
- DELATORREI Dalmat, 1950:137, figs. 1-7. Type-locality: Guatemala, Totonicapán, Río Samalá. Distr. — Mexico (Chiapas), Guatemala (Huehuetenango, Chimaltenango, Totonicapán, El Quiché, Sololá, San Marcos). Ref. — Dalmat, 1955:253, figs. 103-105, 223-225, 258, 311, 351, 389, 426.
- DELEONI Vargas, 1945a:72, pl. 7, fig. 3, pl. 8, fig. 5, pl. 9, fig. 2, pl. 10, figs. 5-6. Type-locality: Mexico, D.F., Desierto de los Leones.



- Distr. — Mexico (Chiapas, D.F., Durango, México, Oaxaca, Puebla, Veracruz). Refs. — Vargas & Díaz Nájera, 1957b:167, 168, 171, 172, 174, figs. 116-124; Díaz Nájera & Vulcano, 1962:92, figs. 57-61.
- DELPONTEIANUM** Wygodzinsky, 1961:30, figs. 1-16. Type-locality: Argentina, Buenos Aires, La Plata, La Granja, Arroyo "El Gato". Distr. — Argentina (Buenos Aires).
- DIAZNAJERAI** Vargas, 1943a:136 (nom. nov. for [*Odagnia*] *tenuifrons* Enderlein). Type-locality: "Mexico". Distr. — Mexico.
- tenuifrons* Enderlein, 1936:130 (*Odagnia*; preocc. Enderlein, 1921).
- DINELLI** (Joan), 1912:363 (sep., p. 5), figs. 1-11, pl. 1 (*Melusina*). Type-locality: Argentina, Tucumán, Montes de Apachiri. Distr. — Guatemala, Colombia, Peru, Venezuela, Ecuador, Bolivia, Brazil (Santa Catarina), Argentina (Tucumán, Salta, Jujuy). Ref. — Wygodzinsky, 1950:77, pls. 1-9.
- bipunctatum* Malloch, 1912:650. Type-locality: Peru, Charape.
- miniata* Enderlein, 1933:275 (*Dasypelmoza*). Type-locality: Bolivia, Mapiro, Lorenzopata, Sarampiuni, 700 m, San Ernesto, Sorata, 2300 m, San Carlos, 800 m, La Paz; Peru, Rosalina, Urubamba River, Cuzco.
- miniata* Enderlein, 1934:194-195 (*Trichodagnia*; preocc. Enderlein, 1933). Type-locality: Bolivia, Yungas de la Paz, 1000 m, San Antonio and Songo; Peru, Cuzco, Callanga, Valle de Pifiipiñi, 1500 m, and Vilcanota; Colombia, Tierra Caliente; Ecuador, Tambillo.
- ochraceum* Walker of Lane & Porto, 1939:171 (*Eusimulium*).
- martinezi* Vargas, 1943a:136 (nom. nov. for *miniata* Enderlein, 1934).
- DISTINCTUM** Lutz, 1910:241. Type-locality: Brazil, São Paulo, Serra da Bocaina, [Fazenda do Bonito]; Rio de Janeiro, Petrópolis, Caxambu de Cima; Minas Gerais, Juiz de Fora. Distr. — Brazil (Minas Gerais, Rio de Janeiro, São Paulo), ?Argentina (Córdoba). Ref. — C. d'Andretta, 1954:63, 6 figs.
- pertinax* Kollar of Lane & Porto, 1939:169 (part).
- DIVERSIBRANCHIUM** Lutz, 1910:238, 263, fig. 8. Type-locality: Brazil, São Paulo, Serra da Bocaina, [Fazenda do Bonito]. Distr. — Brazil (São Paulo).
- DIVERSIFURCATUM** Lutz, 1910:258, fig. 12. Type-locality: Brazil, São Paulo, Serra da Bocaina, [Fazenda do Bonito]; Rio de Janeiro, Petrópolis, Capivari. Distr. — Brazil (Bahia, Rio de Janeiro, São Paulo).
- DOWNSI** Vargas, Martínez Palacios & Díaz Nájera, 1946:126, figs. 41-44, 128. Type-locality: Mexico, Chiapas, Soconusco, Finca el Rosario. Distr. — Mexico (Chiapas, Oaxaca), Guatemala (Chimaltenango, Guatemala, Escuintla, Santa Rosa, Altaverapaz, Bajaverapaz, El Petén, Huehuetenango, El Quiché, Jalapa, Jutiapa, Zacapa, El Progreso, Retalhuleu, San Marcos, Sololá, Suchitpeque), Venezuela (Miranda).
- ochraceum* Walker of Hoffmann, 1931:215, figs. 3, 4, 8, 12, 14, 15 (*Eusimulium*).
- samboni* Jennings of Dampf, 1943:34.
- DUGESI** Vargas, Martínez Palacios & Díaz Nájera, 1946:129, 167, figs. 45-48. Type-locality: Mexico, Tamaulipas, Xicoténcatl, Río Guayalejo. Distr. — México (San Luis Potosí, Tamaulipas), Guatemala (Escuintla, San Marcos). Ref. — Vargas & Díaz Nájera, 1957b:182, 187, 188, 189, figs. 322-328.




- EARLEI Vargas, Martínez Palacios & Díaz Nájera, 1946:118, 177, figs. 16-21, 131, 141. Type-locality: Mexico, Morcros, Temixco. Distr. — Mexico (Collma, Michoacán, Morelos, Nuevo León, Oaxaca, Veracruz), Guatemala (Chimaltenango, Escuintla, Jutiapa, El Progreso, Santa Rosa, Suchitépéquez).
- earli* Vargas & Díaz Nájera, 1957b: 175, 177, 180, figs. 188-195 (error).
- ECUADORIENSE (Enderlein), 1934:193 (*Trichodagmia*). Type-locality: Ecuador, Pichincha, Tambillo. Distr. — Ecuador (Pichincha). Ref. — Bequaert, 1945:113.
- aequatoriense* Vargas, 1945c:109 (error).
- ecuadoriense* León & Wygodzinsky, 1953:273 (error).
- ENCISOI Vargas & Díaz Nájera, 1949:292-295, figs. 29-37. Type-locality: Mexico, Hidalgo, San Miguel Regla. Distr. — Mexico (Hidalgo, Michoacán). Ref. — Vargas & Díaz Nájera, 1957b:180, 181, 182, 233, 263, figs. 287-293.
- ESCOMELI Roubaud, 1909:428-429. Type-locality: Peru, Arequipa, Tingo. Distr. — Peru (Lima, Arequipa), Ecuador (Imbabura, Azuay, Cotopaxi, Pichincha), Chile (Arica, Tacna). Refs. — Knab, 1914b:18-21, fig. 1; Vargas & Díaz Nájera, 1953a:17-21, figs. 1-8.
- rufidorsum* Enderlein, 1933:283 (*Psilopelmia*). Type-locality: "Upper Peru".
- ESTEVEZI Vargas, 1945a:71, pl. 7, figs. 2, 5, pl. 8, fig. 3, pl. 9, fig. 1. Type-locality: Mexico, Oaxaca, Ixtlán, San Felipe del Agua, 1600 m. Distr. — Mexico (Chiapas, D.F., Guerrero, Morelos, Oaxaca, Veracruz). Refs. — Díaz Nájera & Vulcano, 1962:92, 93, figs. 62-66.
- canadense* Hearle of Vargas, 1943a:133-143, figs. 2-3.
- ETHELAE Dalmat, 1950:143, figs. 8-14. Type-locality: Guatemala, Sololá, Río los Arcos, near Los Encuentros. Distr. — Mexico (Chiapas), Guatemala (Sololá, Quezaltenango, Huehuetenango, Chimaltenango, Guatemala). Refs. — Dalmat, 1951:52, 1955:258, figs. 106-108, 226-228, 254, 312, 352, 390, 427.
- EXIGUUM Roubaud, 1906c:108. Type-locality: Venezuela, Alto Sarare. Distr. — Mexico (Morelos), Panama (Canal Zone), Colombia, Ecuador (Esmeralda, Los Ríos, Napo, Pastaza), Venezuela (Lara, Barinas, Amazonas, Aragua, Yaracuy, Carabobo, Miranda, Mérida), Peru, Bolivia (Gran Chaco, Covende, Chapare), ?Brazil (São Paulo, Rio de Janeiro), Argentina (Tucumán, Salta, Jujuy, Misiones). Refs. — Vargas, 1943b:363, figs. 1-9; Wygodzinsky, 1951b:214.
- glaucophthalmum* Knab, 1914d:123. Type-locality: Peru, Santa Clara, Chosica.
- delpontei* Paterson & Shannon, 1927:739, 742. Type-locality: Argentina, Salta, Embarcación and Jujuy, Zapla, Ledesma.
- FALCULATUM (Enderlein), 1929:327, fig. 2 (*Friestia*). Type-locality: México, Veracruz, Boca del Monte. Distr. — Mexico (Veracruz).
- coffea* Vargas, 1945b:4 (unnecessary change of name).
- FIGUEROA Smart, 1944:133 (nom. nov. for *simile* Silva Figueroa). Type-locality: Chile, Santiago, Apoquindo. Distr. — Chile (Santiago), Argentina (Rio Negro, Neuquén). Ref. — Wygodzinsky, 1958:133.
- simile* Silva Figueroa, 1917:33, figs. 19-20 (preocc. Malloch, 1914). Type-locality: Chile, Santiago, Apoquindo.
- FLAVIFEMUR (Enderlein), 1921b:79 (*Chirostibia*). Type-locality: "Brazil". Distr. — Brazil. Ref. — Stone, 1962:207.

- FLAVIPICTUM Knab, 1914c:83. Type-locality: Peru, Río Charape. Distr. — Peru.
- flavopictum* Pinto, 1931:744 (error).
- FLAVOPUBESCENS Lutz, 1910:248, figs. 17-18. Type-locality: Brazil, Rio de Janeiro, Itatiaia, 2400 m. and Agulhas Negras. Distr. — Brazil (Rio de Janeiro, São Paulo). Ref. — Lane & Vulcano, 1943:434, figs. 8, 18.
- FREEMANI Vargas & Díaz Nájera, 1949:289, figs. 19-28. Type-locality: Mexico, Oaxaca, Etla, San Pablo. Distr. — Mexico (Guanajuato, Guerrero, Oaxaca). Ref. — Vargas & Díaz Nájera, 1957b:175, 177-179, figs. 196-204.
- GANALESSENSE Vargas, Martínez Palacios & Díaz Nájera, 1946: 114, 165, figs. 1-7. Type-locality: Mexico, San Luis Potosí, Guerrero, Hacienda de Ganales. Distr. — Mexico (Morelos, San Luis Potosí, Tabasco). Ref. — Vargas & Díaz Nájera, 1957b:163, figs. 29-36.
- ganalesense* Nettel, 1952:439 (error).
- GAUDEATUM Knab, 1914c:81. Type-locality: Peru, Cañon de Verrugas. Distr. — Peru.
- gandatum* Iriarte, 1946:460 (error).
- GONZALESII Vargas & Díaz Nájera, 1953c:235, figs. 1, 3, 5-7, 9 (nom. nov. for *exiguum* of Knab). Type-locality: Not stated [types were not selected]. Distr. — Mexico (Chiapas, Colima, Guerrero, Hidalgo, Jalisco, Morelos, Nayarit, Oaxaca, Puebla, San Luis Potosí, Tabasco, Veracruz), Guatemala (Altaverapaz, Sololá, Chimaltenango, Zacatepéquez, Guatemala, Escuintla, Izabal, Huehuetenango, El Petén, El Quiché, Chiquimula, Jutiapa, Zacapa, San Marcos, Suchitepéquez). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:171, fig. 153.
- exiguum* Roubaud of Knab, 1913:154 (part).
- GORIROSSIAE Vargas & Díaz Nájera, 1957b:199, figs. 143-151 (as *gorirossi*; emend. by Vargas & Díaz Nájera, 1959:106). Type-locality: Mexico, Chiapas, Mariscal, Finca Guadalupe Zajú. Distr. — Mexico (Chiapas, Oaxaca, Veracruz). Ref. — Díaz Nájera & Vulcano, 1962: 92, 101, 102, figs. 67-71.
- GUERRERENSE Vargas & Díaz Nájera, 1956:51, figs. 1-8. Type-locality: Mexico, Guerrero, Ayutla. Distr. — Mexico (Guerrero).
- pulverulentum* Knab of Vargas, 1943b:366, pl. 2, figs. 10-18.
- GUIANENSE Wise, 1911:248. Type-locality: British Guiana, Essequibo River. Distr. — British Guiana, Venezuela (Bolívar). Refs. — Wise, 1912:43; Smart, 1940:4, figs. 1-2, 1942:47, figs. 1a-b.
- tarsale* Williston of Floch & Abonnenc, 1946a: pl. 2.
- GUTTATUM (Enderlein), 1936:124 (*Phodasina*). Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Paraguay (La Cordillera).
- HAEMATOPOTUM Malloch, 1914:62. Type-locality: Mexico, Veracruz, Santa Lucrecia; Cuba, Cayamas. Distr. — Mexico (Chiapas, Colima, Guerrero, Jalisco, Michoacán, Morelos, Nayarit, Oaxaca, Puebla, Sinaloa, Tabasco, Veracruz), Guatemala (Chimaltenango, Escuintla, Santa Rosa, Altaverapaz, Izabal, Huehuetenango, Guatemala, Chiquimula, El Progreso, Zacapa, Quezaltenango, San Marcos, El Petén), Cuba, Puerto Rico, Venezuela (Bolívar), British Guiana, Colombia.

- pseudohaematopum* Hoffmann, 1930b:295, figs. 3, 10. Type-locality: Mexico, Chiapas, Pueblo Nuevo.
- quadrivittatum* Fairchild of Dunn, 1934:177.
- boydi* León, 1945:76, fig. 16. Type-locality: Guatemala, Guatemala, Palín, Hacienda "El Llano".
- HECHTI Vargas, Martínez Palacios & Díaz Nájera, 1946:142, figs. 79-82. Type-locality: Mexico, D.F., Los Dinamos Contreras. Distr. — Mexico (D.F., Morelos). Ref. — Vargas & Díaz Nájera, 1957b:190, 193, 195-196, 198, 260, figs. 410-416.
- sayi* Dyar & Shannon of Vargas, Díaz Nájera & Martínez Palacios, 1943:288.
- HINMANI Vargas, Martínez Palacios & Díaz Nájera, 1946:120, figs. 22-28, 130, 158. Type-locality: Mexico, México, Los Remedios. Distr. — Mexico (Chiapas, Colima, México, Oaxaca, Veracruz).
- HIRTUPA Lutz, 1910:260, 263-264, fig. 21. Type-locality: Brazil, Minas Gerais, Lassance. Distr. — Brazil (Minas Gerais).
- HOFFMANNI Vargas, 1943a:138 (nom. nov. for *angustitarsis* Enderlein). Type-locality: Peru, Calca. Distr. — Peru.
- angustitarsis* Enderlein, 1933:289, 290 (*Trichodagmia*; preocc. Lunds-tröm, 1911). Type-locality: Peru, Calca.
- HUAIYAYACU Wygodzinsky, 1953a:310, figs. 81-127. Type-locality: Argentina. Tucumán, Arroyo Matadero (between Raco and Siambón). Distr. — Argentina (Salta, Tucumán, Jujuy).
- IGNESCENS Roubaud, 1906b:107. Type-locality: Ecuador, Napo. Distr. — Ecuador (Imbabura, Cotopaxi, Pichincha, Napo). Bolivia (Mapiří, Yungas), Peru, Argentina (Tucumán, Salta, Buenos Aires).
- rubiginosum* Enderlein, 1933:287 (*Chirostilbia*). Type-locality: Bolivia. Lorenzopata, Mapiří, San Carlos.
- INAEQUALE (Paterson & Shannon). 1927:739 (*Eusimulium*). Type-locality: Argentina, Jujuy, Zapla. Distr. — Argentina (Tucumán, Salta, Jujuy), Bolivia (Yungas). Ref. — Wygodzinsky, 1953a:304, figs. 37-80.
- INCERTUM Lutz, 1910:259. Type-locality: Brazil, Minas Gerais, Lassance; São Paulo, Salto Grande, Rio Paranapanema. Distr. — Brazil (Minas Gerais, São Paulo). Ref. — Pinto, 1931:755, fig. 8j.
- INCRUSTATUM Lutz, 1910:243, figs. 5-6. Type-locality: Brazil, Rio de Janeiro, Petrópolis and Mendes; São Paulo, Serra da Bocaina, [Fazenda do Bonito], and São Paulo: Rio Pacaembu and Rio Anhangabaú; Minas Gerais, Alfenas; Pernambuco. Distr. — Brazil (Pernambuco, Bahia, Minas Gerais, Rio de Janeiro, São Paulo), Paraguay, Colombia, Ecuador, Venezuela. Rcf. — Wygodzinsky, 1915b:217, figs. 67-68; Vargas & Díaz Nájera, 1953b:137.
- opalifons* Enderlein, 1934:192 (*Psaroniocompsa*). Type-locality: Paraguay, La Cordillera, San Bernardino.
- INEXORABLE Schrottky, 1909:63. Type-locality: Argentina and Brazil, Rio Paraná. Distr. — Paraguay, Brazil, Argentina. Ref. — C. d'Andretta & M.A.V. d'Andretta, 1950:196.
- IRACOUBOENSE Floch & Abonnenc, 1946c:7, figs. 4-5. Type-locality: French Guiana, Bassin d'Iracoubo, Crique Blanche, and Oyapock River, Cafésoca. Distr. — French Guiana. Ref. — Vargas & Díaz Nájera, 1951:150.
- IRIARTEI Vargas, Martínez Palacios & Díaz Nájera, 1946:144, 173, figs. 74-78, 148. Type-locality: Mexico, D.F., Los Dinamos Contreras. Distr.

- Mexico. (D.F., Durango, Guerrero, Oaxaca, Veracruz). Ref. — Vargas & Díaz Nájera, 1957b:191-192, 195, 198, figs. 417-424.
- hirtipes* Fries of Vargas, 1943a: 143, fig. 1.
- ITAUNENSIS C. d'Andretta & González B., 1964:106, figs. 13-19. Type-locality: Brazil, São Paulo, Itatinga, Fazenda Itaúna. Distr. — Brazil (São Paulo).
- JACOBSI Dalmat, 1953:40, figs. 9-11. Type-locality: Guatemala, Escuintla, Palín, Finca el Llano, Río Michatoya. Distr. — Guatemala (Escuintla).
- JACUMBAE Dyar & Shannon, 1927:44, figs. 113-114. Type-locality: U.S.A., California, Jacumba Springs. Distr. — U.S.A. (California), Mexico (Chiapas, Hidalgo, Jalisco, Michoacán, Oaxaca, Veracruz), Guatemala (Bajaverapaz, Chimaltenango, Guatemala, Jalapa, El Progreso, Quezaltenango, Sololá, El Quiché, Zacapa). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:146, figs. 83-86, 150.
- guatemalensis* León, 1945:75, fig. 15. Type-locality: Guatemala, Chimaltenango, Yepocapa.
- JOBBINSI Vargas, Martínez Palacios & Díaz Nájera, 1946:147, figs. 87-90. Type-locality: Mexico, Chiapas, Soconusco, Aldea El Naranjo. Distr. — Mexico (Chiapas, Oaxaca, Veracruz), Guatemala (Altaverapaz, Bajaverapaz, Chimaltenango, Guatemala, Huehuetenango, Quezaltenango, San Marcos, Sololá, Suchitepéquez, Totonicapán, Sacatepéquez). Ref. — Dalmat, 1955:213, figs. 76-78, 196-198, 249, 293-294, 302, 342, 380, 417.
- JOHNSONI Vargas & Díaz Nájera, 1957b:203, figs. 152-160. Type-locality: Mexico, Veracruz, Cascadas de Teocelo. Distr. — Mexico (Michoacán, Oaxaca, Veracruz). Ref. — Díaz Nájera & Vulcano, 1962:99, figs. 72-75.
- JUAREZI Vargas & Díaz Nájera, 1957b:205, figs. 161-169. Type-locality: Mexico, Oaxaca, Ixtlán, El Estudiante. Distr. — Mexico (Michoacán, Oaxaca). Ref. — Díaz Nájera & Vulcano, 1962:100, figs. 76-80.
- JUJUYENSE Paterson & Shannon, 1927:739, 741. Type-locality: Argentina, Jujuy, Zapla. Distr. — Bolivia, Ecuador (Esmeraldas, Cotopaxi), Argentina (Jujuy, Mendoza, Tucumán, Salta, Buenos Aires). Ref. — Wygodzinsky, 1951b:207, figs. 1-51.
- mendozanum* Enderlein, 1936:123 (*Psaroniocompsa*). Type-locality: Argentina, Mendoza.
- JUNDIAIENSIS C. d'Andretta & González B., 1964:108, figs. 20-28. Type-locality: Brazil, São Paulo, Jundiá, Bauru, Caxambu. Distr. — Brazil (São Paulo).
- KOMFI Dalmat, 1951:38, figs. 8-15. Type-locality: Guatemala, Chimaltenango, Acatenango, Aldea Los Pajales, Río Ciprés. Distr. — Guatemala (Chimaltenango). Ref. — Dalmat, 1955:218, figs. 79-81, 199-201, 303, 343, 381, 418.
- LAHILLEI (Paterson & Shannon), 1927:740 (*Eusimulium*). Type-locality: Argentina, Tucumán, Quebrada de Lules. Distr. — Argentina (Tucumán, Salta, Jujuy, Córdoba, Catamarca). Refs. — Wygodzinsky, 1949a:306, 7 pls.; Vargas & Díaz Nájera, 1951:141, figs. 14-22.
- LANEPORTOI Vargas, 1941:118 (as *lane-portoi*; nom. nov. for *pilosum* Lane & Porto). Type-locality: Brazil, Mato Grosso, Chapada, Ponce, 600 m. Distr. — Brazil (Mato Grosso, Paraná).
- lane-porteroi* Smart, 1945:507 (error).



- pilosum* Lane & Porto, 1940:190, figs. 2-5 (preocc. Knowlton & Rowe, 1934). Type-locality: Brazil, Mato Grosso, Chapada, Ponce, 600 m.
- LARVISPINOSUM León, 1948:5, figs. 1-19. Type-locality: Guatemala, Chicacao, Finca Olas de Mocá. Distr. — Mexico (Chiapas), Guatemala (Chimaltenango, Suchitepéquez, Sololá). Ref. — Díaz Nájera & Vulcano, 1962:101, figs. 81-84 (revalidation of sp.).
- LASSMANNI Vargas, Martínez Palacios & Díaz Nájera, 1946:149, 174, figs. 91-97, 123. Type-locality: Mexico, México, Los Remedios. Distr. — Mexico (D.F., Durango, Guanajuato, Veracruz).
- LITICALX (Enderlein), 1933:291 (*Trichodagmia*). Type-locality: Peru, Callanga, Rosalina, Río Urubamba. Distr. — Peru.
- LATIDIGITUS (Enderlein), 1936:129 (*Trichodagmia*). Type-locality: Colombia, Muza. Distr. — Colombia.
- LIMAY Wygodzinsky, 1958:123, figs. 1-19, 22-30. Type-locality: Argentina, Neuquén, Paso Limay, "orilla del Río Limay". Distr. — Argentina (Neuquén).
- LIMBATUM Knab, 1915a:280. Type-locality: British Guiana, Rupununi River. Distr. — British Guiana, Venezuela (Bolívar).
- LONGITHALLUM Díaz Nájera & Vulcano, 1961:221, figs. 1-27. Type-locality: Mexico, Morelos, Tepoztlán. Distr. — Mexico (Jalisco, Morelos).
- LURYBAYAE Smart, 1944:132 (nom. nov. for *angustifrons* Enderlein). Type-locality: Bolivia, La Paz, Luribay. Distr. — Bolivia.
- angustifrons* Enderlein, 1933:288 (*Trichodagmia*; preocc. Enderlein, 1921). Type-locality: Bolivia, La Paz, Luribay.
- MACCA (Enderlein), 1933:276 (*Dasypelmoza*). Type-locality: Peru, Valley of Lares, Rosalina, Río Urubamba. Distr. — Venezuela, Peru.
- maccum* Vargas, 1943a:141 (error).
- MAJOR Lane & Porto, 1940:192, figs. 6-9. Type-locality: Brazil, Mato Grosso, Chapada, Ponce. Distr. — Brazil (Mato Grosso, Goiás, São Paulo). Ref. — Lane & Vulcano, 1943:438, fig. 19.
- magnum* Porto, 1940:383 (error).
- MANGABEIRAI Vargas, 1945a:76, pl. 7, fig. 1, pl. 8, figs. 1-2, pl. 9, figs. 3-4. Type-locality: Mexico, México, Salazar, La Cima, 3005 m. Distr. — Mexico (D.F., México, Michoacán, Veracruz). Refs. — Vargas, Martínez Palacios & Díaz Nájera, 1946:168, fig. 156; Vargas & Díaz Nájera, 1957b:184, 186-187, 189, 204, 231, 235-236, 258, figs. 336-342.
- MANICATUM (Enderlein), 1933:288 (*Trichodagmia*). Type-locality: "Upper Paraguay". Distr. — Paraguay, Peru.
- MARATHRUMI Fairchild, 1940:715, figs. 3, 15, 21, 34. Type-locality: Panama, Lajas River, Chorreras Falls. Distr. — Mexico (Chiapas), Panama.
- MARONIENSE Floch & Abonnenc, 1946b:9, pls. 8-9. Type-locality: French Guiana, Sinnamary, Crique Coeur Maroni. Distr. — French Guiana.
- MARQUESTI Vargas & Díaz Nájera, 1957a: 37, figs. 29-42. Type-locality: Mexico, D.F., Desierto de los Leones. Distr. — Mexico (D.F., México, Chihuahua, Morelos). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:176, fig. 145.
- vandalicum* Dvar & Shannon of Vargas, 1945a: 78.
- MATTEBRANCHIA Anduze, 1947:374, fig. 1. Type-locality: Venezuela, Aragua, Colonia Tovar. Distr. — Venezuela (Aragua, Trujillo, D.F., Mérida). Ref. — Vargas & Díaz Nájera, 1948b:334, figs. 41-46.

- MEDIOVITTATUM** Knab, 1915b:77. Type-locality: U.S.A., Texas, Arlington Distr. — U.S.A. (Texas), Mexico (Chihuahua, San Luis Potosí, Tamaulipas, Veraeruz). Refs. — Vargas & Díaz Nájera, 1954:69, figs. 31-39; Vargas & Díaz Nájera, 1957b:183, 185, 187-188, 230, 254, 258, 264, figs. 343-349.
- MENCHACAI** Vargas & Díaz Nájera, 1957b:208, figs. 170-178, pl. 20bis, figs. A-B. Type-locality: Mexico, Oaxaca, San Felipe del Agua. Distr. — Mexico (Oaxaca). Ref. — Díaz Nájera & Vuleano, 1962:101, figs. 85-88.
- MERIDIONALE** Riley, 1887:513, pl. 7, figs. 2, 3a-c, 4-6, pl. 8, figs. 4, 6. Type-locality: U.S.A., Mississippi, Lake View. Distr. — U.S.A. (Mississippi, New Mexico), Mexico (Tamaulipas). Refs. — Dyar & Shannon, 1927:32, pl. 5, figs. 63a, 64, 131-132; Stone, 1952:94.
- occidentale* Townsend, 1891:107. Type-locality: U.S.A., New Mexico, Río Grande Valley.
- tamaulipense* Townsend, 1897:171. Type-locality: México, Tamaulipas.
- METALLIUM** Bellardi, 1859:13. Type-locality: "Mexico". Distr. — Mexico (México, Tamaulipas, Chiapas, Chihuahua, Colima, Durango, Guerrero, Hidalgo, Jalisco, Michoacán, Morelos, Nayarit, Nuevo León, Oaxaca, Puebla, San Luis Potosí, Sinaloa, Sonora, Tabasco, Veraeruz), Guatemala (Altaverapaz, Bajaverapaz, Chimaltenango, Guatemala, Izabal, Jutiapa, Santa Rosa, Totonicapán, El Quiché, Quezaltenango, Sacatepéquez, Sololá, Retalhuleu, Zacapa, Huehuetenango, Chiquimula, Jalapa, Esequintla, El Progreso, Suchitepéquez, El Petén, San Marcos), Costa Rica, Panama (Coclé), Trinidad, Venezuela (Carabobo, Aragua, Trujillo, Anzoátegui, Mérida, Yaracuy, Lara, Cojedes, Monagas, Bolívar, Sucre, Miranda, Falcón, D.F., Maracaib); Ecuador. Refs. — Faurehild, 1940:712, figs. 13, 17, 26, 38; Vargas, Martínez Palacios & Díaz Nájera, 1946:175, figs. 143, 157.
- avidum* Hoffmann, 1930a:51, figs. 1-4. Type-locality: Mexico, Chiapas, Soconuseo, Finca La Granja, 700 m.
- MEXICANUM** Bellardi, 1862:6. Type-locality: Mexico, Veraeruz, Tuxpán, Orizaba. Distr. — Mexico (Chiapas, México, Oaxaca, Veracruz), Guatemala (Altaverapaz, Chimaltenango, Sololá, Izabal, Huehuetenango, Escuintla, Chiquimula, Zaeapa, Santa Rosa, Suchitepéquez, Quezaltenango, Guatemala), Panama (Coclé), Trinidad, Tobago I., Colombia, Venezuela (Aragua, Monagas, Miranda, Barinas), Bolivia. Refs. — Dyar & Shannon, 1927:15, fig. 46; Bequaert, 1934:216, fig. 99.
- aureopunctatum* Malloch, 1914:27. Type-locality: Guatemala, Altaverapaz, Livingston, Cacao, Treee Aguas.
- placidum* Knab, 1915a:281. Type-locality: West Indies, Trinidad, Arima River.
- lugubre* Lutz & Nuñez Tovar, 1928:46, pl. 4, fig. 2, pl. 6, fig. 6. Type-locality: Venezuela, Aragua, Maracaib.
- turgidum* Hoffmann, 1930b:293, 298, 299, 300, 303, figs. 6, 13 (*Eusimulium*). Type-locality: Mexico, Chiapas, Soconuseo, Finca Santa Anita.
- MICROBRANCHIUM** Dalmat, 1949:538, figs. 1-9. Type-locality: Guatemala, Sololá, Río los Arcos, near Los Encuentros. Distr. — Guatemala (Huehuetenango, Quiché, Quezaltenango, Totonicapán, Sololá). Refs. — Dalmat, 1951:50, 1955:268, figs. 8, 112-114, 232-234, 257, 314, 354, 392, 429.



- NEMORALE Edwards, 1931:152, figs. 8b, 10h-i, 13g. Type-locality: Argentina, Río Negro, L. Correntoso. Distr. — Argentina (Río Negro, Neuquén). Ref. — Vargas & Díaz Nájera, 1951:138, figs. 8-10.
- NIGRICOORNIS Dalmat, 1950:148, figs. 15-21. Type-locality: Guatemala, Chimaltenango, Acatenango, Finca Tehuyá Luch, Río Laguneta. Distr. — Guatemala (Chimaltenango, Acatenango). Ref. — Dalmat, 1955: 273, figs. 115-117, 235-236, 315, 355.
- NIGRIFEMUR (Enderlein), 1936:118 (*Acropogon*). Type-locality: Chile, Cauquenes. Distr. — Chile.
- NIGRIMANUM Macquart, 1838:88 (as *nigrimana*). Type-locality: Brazil, São Paulo. Unrecognized.
- NIGRISTRIGATUM (Enderlein), 1930:88, fig. 3 (*Pternaspatha*). Type-locality: Argentina, Neuquén. Distr. — Argentina (Neuquén, Tierra del Fuego, Mendoza, Río Negro), Chile (Magallanes). Refs. — Vargas & Díaz Nájera, 1953b:142, figs. 4-11; Edwards, 1931:154.
- edwardsi* Enderlein, 1933:279 (*Aeropogon*; unnecessary change of name for *nigristrigatum* of Edwards).
- NOGUEIRAI C. d'Andretta & González B., 1964:103, figs. 1-12. Type-locality: Brazil, São Paulo, Aldeinha (Estrada de Ferro Sorocabana). Distr. — Brazil (São Paulo).
- OBESUM Vulcano, 1959a:351, figs. 1-44. Type-locality: Brazil, São Paulo, Serra da Bocaina, Fazenda do Bonito. Distr. — Brazil (Río de Janeiro, São Paulo).
- obesum* Lutz, 1922:91. *Nomen nudum*.
- OCHOAI Vargas, Martínez Palacios & Díaz Nájera, 1946:131, figs. 49-52, 152. Type-locality: Mexico, Veraeruz, Tierra Blanca. Distr. — Mexico (Veraeruz). Ref. — Vargas & Díaz Nájera, 1957b:182, 187, 188, 189, 264, figs. 350-356.
- OCHIRACEUM Walker, 1861:332. Type-locality: "Mexico". Distr. — Mexico (Chiapas, Oaxaca, Veraeruz), Guatemala (Altaverapaz, Bajaverapaz, Guatemala, Quezaltenango, El Progreso, Chimaltenango, Sololá, Santa Rosa, Jutiapa, Esequintla, Suchltepéquez, Saatepéquez, El Petén, Huehuetenango, El Quiché, Chiquimula, Jalapa, Retalhuleu, San Marcos), Panama (Coclé, Colón), Jamaica, Colombia, Venezuela (Aragua, Mérida, Trujillo, Miranda), Ecuador, Bolivia. Refs. — Hoffmann, 1931:217, figs. 3-5, 12, 14-15; Fairchild, 1940:706, figs. 6, 11, 23, 35.
- rubrithorax* Lutz of Lutz & Nuñez Tovar, 1928: pl. 6, fig. 2 (part).
- lutzianum* Pinto, 1931:748, figs. 39-40 (nom. nov. for *rubrithorax* of Lutz & Nuñez Tovar).
- ORBITALE Lutz, 1910:231, figs. 35, 36, 39, 40 (nom. nov. for *nigrimanum* of Lutz). Type-locality: Brazil, São Paulo, Río Tietê, Salto de Itapura, and Franca, Fazenda Borda do Mato; Benjamin Constant, Río Paraíba. Distr. — Brazil (Minas Gerais, Río de Janeiro, São Paulo, Paraná, Santa Catarina), Paraguay, Argentina (Misiones). Ref. — M.A.V. d'Andretta & C. d'Andretta, 1946:85, pls. 1-9, pl. 10, figs. 1-4.
- nigrimanum* Macquart of Lutz, 1909:137.
- brasiliense* Enderlein, 1933:284 (*Thyrsopeima*). Type-locality: "Brazil".
- paraguayense* Schrottky of Lane & Vulcano, 1943:437.
- OYAPOCKENSE Floch & Abonnenc, 1946c:4, fig. 3. Type-locality: French Guiana, Oyapoek, Maripa, "près du saut Cafésoca".

- PANAMENSE Fairchild, 1940:705, figs. 12, 20, 29, 39. Type-locality Panama, Coclé, El Valle, Puente River. Distr. — Panama (Coclé, Darién), Costa Rica. Ref. — Vargas & Díaz Nájera, 1954:57, figs. 1-16.
- PARAGUAYENSE Schrottky, 1909:63. Type-locality: Paraguay, and Argentina, Alto Paraná. Distr. — Paraguay, Argentina (Misiones, Corrientes), Brazil (Bahia, Minas Gerais, Rio de Janeiro, São Paulo), Venezuela (Aragua, D.F.). Ref. — Wygodzinsky, 1953a:298, figs. 25-27, 29-36.
- PARANENSE Schrottky, 1909:63. Type-locality: Paraguay, Alto Paraná. Distr. — Paraguay.
- PARRAI Vargas, Martínez Palacios & Díaz Nájera, 1946:151, 175, figs. 98-105, 146. Type-locality: Mexico, Veracruz, Cascada de Teocelo. Distr. — Mexico (Oaxaca, Veracruz), Guatemala (Chimaltenango, Sololá). Ref. — Vargas & Díaz Nájera, 1948b:336.
- PAYNEI Vargas, 1942a:246 (nom. nov. for *mexicana* Enderlein). Type-locality: "Mexico". Distr. — Mexico (Chiapas, Colima, Durango, Guerrero, Jalisco, México, Michoacán, Morelos, Nuevo León, Oaxaca, Puebla, San Luis Potosí, Veracruz, Sinaloa), Guatemala (Chimaltenango, Santa Rosa, Totonicapán, Sacatepéquez, Huehuetenango, Jutiapa, El Quiché, Altaverapaz, Guatemala, Chiquimula). Refs. — Vargas & Díaz Nájera, 1953b:145, 1957b:232, 240, pl. C, figs. 231-238. *mexicana* Enderlein, 1934:190 (*Hemicnetha*; preocc. Bellardi, 1862). Type-locality: "Mexico".
- mathesoni* Vargas, 1943b:360, figs. 37-43. Type-locality: Mexico, Morelos, Temisco, 1400-1500 m. Ref. — Vargas, 1945a:78.
- bilimekae* Smart, 1944:132 (nom. nov. for *mexicana* Enderlein).
- acatenangoensis* Dalmat, 1951:31, figs. 1-17. Type-locality: Guatemala, Chimaltenango, Acatenango, Finca La Esperanza Pérez, Río La-drillera.
- PERFLAVUM (Roubaud), 1906b:517 (*Eusimulium*). Type-locality: Brazil, São Paulo. Distr. — Brazil (Minas Gerais, Guanabara, Rio de Janeiro, São Paulo), Venezuela (Mérida). Ref. — C. d'Andretta, 1956:33, figs. 1-11.
- PERTINAX Kollar in Pohl & Kollar, 1832:19, figs. 14, 14b. Type-locality: Brazil [probably Rio de Janeiro, Guanabara]. Distr. — Brazil (Rondônia, Mato Grosso, Bahia, Rio de Janeiro, Guanabara, São Paulo, Paraná), Argentina (Misiones, Córdoba), Paraguay. Ref. — C. d'Andretta & M.A.V. d'Andretta, 1950:193, figs. 1-53.
- venustum* Say of Lutz, 1909:136 (*nec pupa*) (part).
- distinctum* Lutz of Pinto, 1931:669, fig. 5b (part).
- paraguayense* Schrottky of Pinto, 1931:715, figs. 6a, 7g, 33-34.
- lutziana* Enderlein, 1933:291 (*Trichodagmia*; preocc. Pinto, 1931). Type-locality: Paraguay, La Cordillera, San Bernardino.
- lutzianum* Enderlein of Rubzov, 1940:128.
- PHILIPPIANUM Pinto, 1931:729 (nom. nov. for *tarsatum* Philippi). Type-locality: Chile, Santiago, Corral. Distr. — Chile (Santiago).
- tarsatum* Philippi, 1865:634 (preocc. Macquart, 1846). Type-locality: Chile, Santiago, Corral.
- PINTOI M.A.V. d'Andretta & C. d'Andretta, 1946:101, pl. 10, figs. 5-8, pls. 11-19. Type-locality: Brazil, São Paulo, Salto de Piracicaba. Distr. — Brazil (Minas Gerais, São Paulo). Ref. — Vargas & Díaz Nájera, 1951:137.
- guyanense* Wise of Floch & Abonnenc, 1946a: fig. 2.
- orbitale* Lutz, 1910:233 (*nec p. 231*) (part).

- orbitale* Lutz of Pinto, 1931:706 (part).
- PRODEXARGENTEUM (Enderlein), 1936:116 (*Pternaspatha*; as *prodex-argenteum*). Type-locality: "Upper Peru". Distr. — Peru.
- FRUINOSUM Lutz, 1910:250, figs. 38, 41, 42. Type-locality: Brazil, Minas Gerais, Lassance, Rio das Velhas. Distr. — Brazil (Goíás, Minas Gerais, Mato Grosso, São Paulo), Venezuela (Barinas), Paraguay. Ref. — Vulcano, 1959b:33, figs. 1-117.
- PSEUDOCALLIDUM Díaz Nájera, 1965:25, figs. 1-31. Type-locality: Mexico, México, Avándaro, Valle de Bravo, 1800 m. Distr. — Mexico (Chiapas, México, Oaxaca).
- PUGETENSE (Dyar & Shannon), 1927:23, figs. 121-123 (*Eusimulium*). Type-locality: U.S.A., Washington, Seattle. Distr. — U.S.A. (Washington). Ref. — Vargas & Díaz Nájera, 1957b:164, 165, 231, 262, pl. B, figs. 54-64.
- PUIGI Vargas, Martínez Palacios & Díaz Nájera, 1946:153, figs. 106-109. Type-locality: Mexico, Chiapas, Mariscal, Aldea El Oriente, 2350 m. Distr. — México (Chiapas, Veracruz, Oaxaca). Ref. — Vargas & Díaz Nájera, 1957b:191, 192, 194, 196, 199, 230, 252, 261, 264, figs. 479-485.
- PULCHRUM Philippi, 1865:633. Type-locality: Chile, Santiago. Distr. — Chile (Santiago).
- PULVERULENTUM Knab, 1914a:177. Type-locality: British Honduras, Punta Gorda. Distr. — Mexico (Chiapas, Guerrero, Hidalgo, Michoacán, Nayarit, Oaxaca, San Luis Potosí, Veracruz), Guatemala (Chimaltenango, Escuintla, Huehuetenango, Izabal, Jutiapa, Progreso, Retalhuleu, San Marcos, Santa Rosa, Sololá, Suchitepéquez, Zacapa, Chiquimula), British Honduras, El Salvador, Panama, Venezuela (Miranda, Anzoategui, D.F.). Ref. — Vargas & Díaz Nájera, 1957b:256, figs. 239-246.
- PUNCTATVENTRIS (Enderlein), 1936:118 (*Acropogon*). Type-locality: Central Chile, Cortoderal. Distr. — Chile.
- QUADRIFIDUM Lutz, 1917:66. Type-locality: Brazil, Rondônia, Estrada de Ferro Madeira-Mamoré. Distr. — Brazil (Rondônia). Ref. — Lutz, 1910:255, figs. 31-32 (as *amazonicum*).
amazonicum Goeldi of Lutz, 1910:255, figs. 31-32.
- QUADRISTRIGATUM Enderlein, 1933:285. Type-locality: Brazil, Paraná. Distr. — Brazil (Paraná), Paraguay. Ref. — Enderlein, 1936:123 (*Psaroniocompsa*).
- QUADRIVITTATUM Loew, 1862:56. Type-locality: "Cuba". Distr. — Mexico (Campeche, Chiapas, Oaxaca, Puebla, Veracruz), Cuba, Costa Rica, Jamaica, Puerto Rico, Panama (Canal Zone), Venezuela (Trujillo, Falcón), Ecuador (Pichincha, Los Ríos). Refs. — Vargas, 1945a: 79, pl. 12, figs. 1-4; Vargas & Díaz Nájera, 1957b:162, 163, 164, 211-212, 238, 252, 255, 262, figs. 37-44; Bradt, 1932:77, figs. 1-5.
metallicum Bellardi of Malloch, 1914:49, pl. 6, fig. 4.
quadrivittatum Loew of Malloch, 1914:61.
mallochi Enderlein, 1925:208 (*Wilhelmia*; nom. nov. for *quadrivittatum* of Malloch).
haematopotum Malloch of Fairchild, 1940:714, figs. 2, 16, 27, 33.
fairchildi Vargas, 1942b:458 (nom. nov. for *haematopotum* of Malloch).
- RIVETI Roubaud, 1906c:107. Type-locality: Ecuador, Napo. Distr. —

- Ecuador (Imbabura, Pichincha, Tungurahua, Chimborazo, Napo). Refs. — Wygodzinsky, 1953b:332, figs. 74-89; León & Wygodzinsky, 1953:277.
- ROMANAI** Wygodzinsky, 1951c:229, figs. 51, 53-97 (as *romañai*). Type-locality: Argentina, Tucumán, Cerro San Javier, Parque Aconquija. Distr. — Argentina (Tucumán, Salta).
- ROROTAENSE** Floch & Abonnenc, 1946b:4, pls. 4-7. Type-locality: French Guiana, Plateau Rorota. Distr. — French Guiana.
- RUBRITHORAX** Lutz, 1909:132. Type-locality: Brazil, São Paulo, Serra da Bocaina, [Fazenda do Bonito], 1500 m, and Batatais. Distr. — Venezuela (Aragua, Yaracuy, Mérida, Miranda), Brazil (Minas Gerais, Rio de Janeiro, São Paulo), Argentina (Misiones). Ref. — Vulcano, 1958:227, figs. 1-50.
- RUIZI** Vargas & Díaz Nájera, 1948b:330, figs. 37-40. Type-locality: Mexico, Veracruz, Tequila. Distr. — Mexico (México, Oaxaca, Veracruz). Ref. — Vargas & Díaz Nájera, 1957b:191, 193, 195, 197-198, 235, 252, 261, 265, figs. 486-492.
- SAMBONI** Jennings, 1915:199. Type-locality: Panama, Canal Zone, Empire, Comacho River. Distr. — Mexico (Chiapas, Tabasco, Oaxaca, Veracruz), Guatemala (El Petén, San Marcos, Sololá, Totonicapán, Escuintla, Altaverapaz, Santa Rosa), British Honduras, Panama (Canal Zone, Chucumaque, Panama, Coclé), Venezuela (Maracay, Miranda). Refs. — Vargas & Díaz Nájera, 1948b:343, figs. 60-61, 1954:62; Dalmat, 1955:151, figs. 40-42, 157-159, 264, 289, 329, 369, 406.
- colvini* Dalmat, 1952:344. Type-locality: Guatemala, San Marcos, "in a stream between Malacatán and Ayutla".
- SANGUINEUM** Knab, 1915:279. Type-locality: Colombia, Boca de Arquia, Río Atrato. Distr. — Panama (Darién), Venezuela (Anzoategui, Mérida, Amazonas), Colombia, British Guiana.
- SCHOENEMANNI** (Enderlein), 1933:277 (*Acropogon*). Type-locality: Chile, Cauquenes. Distr. — Chile.
- SCUTELLATUM** (Lane & Porto), 1940:192, figs. 10-13 (*Eusimulium*). Type-locality: Colombia, Restrepo. Distr. — Colombia, Venezuela, Brazil (São Paulo). Ref. — Lane & Vulcano, 1943:434, fig. 6.
- SCUTISTRIATUM** Lutz, 1909:133. Type-locality: Brazil, Rio de Janeiro, Itaguaí and Duque de Caxias, Xerém. Distr. — Brazil (Goiás, Minas Gerais, Rio de Janeiro, São Paulo). Ref. — Lane & Vulcano, 1943:436, fig. 17.
- SERIATUM** Knab, 1914c:84. Type-locality: Peru, Santa Eulalia, Chosica. Distr. — Peru.
- SICUANI** Smart, 1944:135 (nom. nov. for *limbatum* Enderlein, 1933). Type-locality: Peru, Sicuaní, Calca, Valley of Lares, Rosalina, Río Urubamba. Distr. — Peru.
- limbatum* Enderlein, 1933:282 (*Ectemnaspis*; preocc. Knab, 1915; Enderlein, 1921). Type-locality: Peru, Sicuaní, Calca, Valley of Lares, Rosalina, Río Urubamba.
- incaicum* Vargas, 1945b:4 (nom. nov. for *limbatum* Enderlein, 1933).
- SIMPLICICOLOR** Lutz, 1910:251, figs. 37, 43. Type-locality: Brazil, Rondônia, Estrada de Ferro Madeira-Mamoré. Distr. — Brazil (Rondônia).
- hebeticolor* Lutz, 1910:267 (error).
- SMARTI** Vargas, 1946:330, figs. 1-4. Type-locality: Mexico, Chiapas, Finca El Vergel, Arroyo El Rubí, 1000 m. Distr. — Mexico (Chiapas, Oaxa-

- ca), Guatemala (Chlmaltcnango, Huehuetenango, Sololá, Suchitpéquez, Sacatepéquez, El Quiché). Ref. — Dalmat, 1955:199, figs. 67-69, 187-189, 243, 299, 339, 377, 414.
- SOLARI Stone, 1948:402, figs. 2, 4, 6, 8, 10. Type-locality: U.S.A., Texas, San Sabas River, Menard County. Distr. — U.S.A. (Texas, New Mexico), Mexico (México, Nuevo León, San Luis Potosí, Tamaulipas). Ref. — Vargas & Díaz Nájera, 1957b:175, 176, 178, 180, figs. 255-262.
- virgatum* Coquillett of Stains & Knowlton, 1943:274.
- virgatum* Coquillett of Vargas, Martínez Palacios & Díaz Nájera, 1946:180, figs. 161-162.
- SPADICIDORSUM (Enderlein), 1934:194 (*Trichodagmia*). Type-locality: Brazil, São Paulo. Distr. — Brazil (São Paulo).
- SPINIBRANCHIUM Lutz, 1910:248, 263-264, fig. 13. Type-locality: Brazil, São Paulo, Anhembi, Rio Tietê; Minas Gerais, Lassance. Distr. — Brazil (Minas Gerais, São Paulo).
- subpallidum* Lutz, 1910:247, 263, 264. Type-locality: Brazil, São Paulo, Anhembi, Rio Tietê and Minas Gerais, Lassance (part).
- SPINIFER Knab, 1914c:81. Type-locality: Peru, Cañon de Verrugas. Distr. — Peru, Ecuador. Ref. — Vargas & Díaz Nájera, 1951:153, figs. 66-69.
- STRIGATUM (Enderlein), 1933:289 (*Trichodagmia*). Type-locality: Peru, Mamara, Rosalina, Río Urubamba. Distr. — Peru.
- STRIGIDORSUM (Enderlein), 1933:279 (*Acropogon*). Type-locality: Peru, Juliaca. Distr. — Peru.
- STRIGINOTUM (Enderlein), 1933:284 (*Thyrsopelma*). Type-locality: Bolivia, Mapiri, Sarampiuni. Distr. — Bolivia.
- SURCLAIVIBRANCHIUM Lutz, 1910:260, fig. 33. Type-locality: Brazil, Rondônia, Estrada de Ferro Madeira-Mamoré. Distr. — Brazil (Rondônia).
- SUBNIGRUM Lutz, 1910:239, fig. 7. Type-locality: Brazil, São Paulo, Rio Pacaembu and Rio Anhangabaú; Rio de Janeiro, Mendes and Petrópolis. Distr. — Trinidad, Venezuela (D.F., Miranda, Bolívar), Colombia, Brazil (Rio de Janeiro, São Paulo, Paraná). Refs. — Lane & Vulcano, 1943:138, fig. 18; Porto, 1940:383.
- SURPALLIDUM Lutz, 1910:247, fig. 13 (part). Type-locality: Brazil, São Paulo, Anhembi, Rio Tietê; Minas Gerais, Benjamin Constant, Lassance. Distr. — Brazil (Minas Gerais, São Paulo).
- TARSALE Williston, 1896:268, figs. 25, 25a. Type-locality: West Indies, St. Vincent. Distr. — West Indies (St. Vincent, Guadeloupe). Ref. — Floch & Abonnenc, 1946b:20, pl. 10.
- pulchrum* Philippi of Johannsen, 1903:376.
- clavipes* Malloch, 1914:40, pl. 2, fig. 5. Type-locality: West Indies, Guadeloupe.
- TARSATUM Macquart, 1846:20. Type-locality: Colombia [Nouvelle Grenade]. Distr. — Colombia.
- TEMASCALENSE Díaz Nájera & Vulcano, 1962:102, figs. 85-105. Type-locality: Mexico, Michoacán, Temascal, 2800 m. Distr. — Mexico (Michoacán, Oaxaca).
- TENUIPES Knab, 1914b:21. Type-locality: Chile, Valle de Lluta. Distr. — Chile.
- TOWNSENDI Malloch, 1912:651, pl. 46, fig. 9. Type-locality: Peru, Río Charape. Distr. — Peru.



- TRAVASSOSI M.A.V. d'Andretta & C. d'Andretta, 1947b:161, figs. 49-125.
Type-locality: Brazil, Rio de Janeiro, Angra dos Rcis. Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo).
botulibranchium Lutz of Pinto, 1931:688 (part).
- TRICORNIS León, 1945:68, fig. 4. Type-locality: Guatemala, Totonicapán, Sololá, María Tecún. Distr. — Mexico (Chiapas), Guatemala (Totonicapán, Sololá, Chimaltenango, Huehuetenango, San Marcos, Guatemala). Ref. — Vargas, Martínez Palacios & Díaz Nájera, 1946:155, figs. 110-116, 122, 151.
- TRIVITTATUM Malloch, 1914:25, 30, 39, 68. Type-locality: Mexico, Tamaulipas, Tampico. Distr. — U.S.A. (Texas), Mexico (Chiapas, Chihuahua, Coahuila, Colima, Guanajuato, Guerrero, Jalisco, Michoacán, Morelos, Nuevo León, Oaxaca, Puebla, San Luis Potosí, Tamaulipas, Veracruz), Guatemala (Altavcrapaz). Refs. — Dyar & Shannon, 1927:37, figs. 78-79; Vargas, Martínez Palacios & Díaz Nájera, 1946:169, fig. 127; Dalmat, 1955:156, figs. 43-45, 160, 162, 266, 290, 330.
distinctum Malloch, 1913:133 (preocc. Lutz, 1910). Type-locality: U.S.A., Texas, Devils River.
- TUBEROSUM (Lundström), 1911:14, fig. 10 (*Melusina*). Type-locality: "Lapponia". Distr. — Europe, Asia; Alaska, Canada, U.S.A. (Florida, California, Texas), Mexico (Durango). Ref. — Vargas & Díaz Nájera, 1957a:35, figs. 15-28.
- URUBAMBANUM Enderlein, 1933:286. Type-locality: Peru, Rosalina, Río Urubamba, Unini, Río Ucayali. Distr. — Peru.
- VARIANS Lutz, 1909: 141. Type-locality: Brazil, São Paulo, Río Grande, near Franca. Distr. — Brazil (São Paulo).
- VARIPES Philippi, 1865:634. Type-locality: Chile, Valdivia. Distr. — Chile (Valdivia, Llanquihuc, Concepción), Argentina (Río Negro, Neuquén). Refs. — Enderlein, 1929:327, fig. 1; Vargas & Díaz Nájera, 1951:139, figs. 11-13.
- VERACRUZANUM Vargas, Martínez Palacios & Díaz Nájera, 1946:133, figs. 53-57, 155. Type-locality: Mexico, Veracruz, Río Sedcño. Distr. — Mexico (Chiapas, Colima, Durango, Jalisco, Michoacán, México, Morelos, Nayarit, Oaxaca, Veracruz), Guatemala (Chimaltenango, Guatemala, Sacatepéquez, Escuintla, Santa Rosa, Altavcrapaz, Huehuetenango, El Quiché, Zacapa, Quezaltenango, Sololá, Totonicapán, Bajaverapaz, San Marcos). Refs. — Dalmat, 1955:159, figs. 46-48, 163-165, 263, 291, 331, 370, 407; Vargas & Díaz Nájera, 1957b: 246, figs. 273-279.
- VERSICOLOR Lutz & Nuñez Tovar, 1928:47, pl. 4, fig. 1, pl. 6, fig. 7. Type-locality: "Venezuela". Distr. — Venezuela (Bolívar). Ref. — C. d'Andretta & M.A.V. d'Andretta, 1946:308.
- VIOLACESCENS (Enderlein), 1933:285 (*Odagmia*). Type-locality: "Mexico". Distr. — Mexico, Colombia, Venezuela (D.F.).
- VIRGATUM Coquillett, 1902:97. Type-locality: U.S.A., New Mexico, Las Vegas, Hot Springs. Distr. — U.S.A. (New Mexico), Mexico (Chiapas, Chihuahua, Jalisco, Michoacán, Oaxaca, Sinaloa, San Luis Potosí, Tabasco, Veracruz), Guatemala (Chimaltenango, Sololá, Guatemala, Sacatepéquez, Escuintla, Santa Rosa, Bajaverapaz, Huehuetenango, El Quiché, Chiquimula, Jutiapa, El Progreso, Altavcrapaz, Suchitepéquez, Totonicapán, San Marcos, Quezalte-

- nango), Panama. Refs. — Dyar & Shannon, 1927:39, figs. 82-83, 126-128; Fairchild, 1940:718, figs. 5, 7, 32; Vargas, Martínez Palacios & Díaz Nájera, 1946:105, figs. 132, 161-162.
- cinereum* Bellardi, 1859:13 (p.occ. Macquart, 1834). Type-locality: Mexico, Morelia.
- tephrodes* Speiser, 1904:148 (nom. nov. for *cinereum* Bellardi).
- hippovororum* Malloch, 1914:28. Type-locality: Mexico, Chihuahua, Sierra Madre, Río Piedras Verdes.
- rubicundulum* Knab, 1914a:178. Type-locality: Mexico, Veracruz, Córdoba; U.S.A., New Mexico, Las Vegas, Hot Springs.
- chiapanense* Hoffmann, 1930b:295, figs. 2-9 (as *virgatum* ssp.). Type-locality: Mexico, Chiapas, Soconusco, Finca Santa Anita.
- VITTATUM (Zetterstedt), 1838:803 (*Simulia*). Type-locality: "Laponia". Distr. — Europe; Canada, U.S.A., Mexico (D.F., Tamaulipas). Refs. — Dyar & Shannon, 1927:28, figs. 74-75, 106-108; Vargas, 1942a:243.
- WALTERVITMERI Wygodzinsky, 1958:129, figs. 31-48. Type-locality: Argentina, Neuquén, Paso Limay, "orilla del Río Limay". Distr. — Argentina (Neuquén).
- WOLCOTTI Fox, 1953:138, figs. 5-6, 8-10. Type-locality: Puerto Rico, Henry Barracks. Distr. — Puerto Rico.
- WOLFFHUEGELI (Enderlein), 1922:70 (*Wilhelmia*). Type-locality: Argentina, Tucumán. Distr. — Argentina (Tucumán, Salta, Jujuy, Mendoza, Neuquén, Buenos Aires). Ref. — Wygodzinsky, 1950:88, pls. 10-15.
- haarupianum* Enderlein, 1936:127 (*Trichodagmia*). Type-locality: Argentina, Mendoza.
- jujuyense* Paterson & Shannon of Vargas & Díaz Nájera, 1951:132.
- WUAYARAKA Ortiz C., 1957:163, pls. 2-3, 1 photograph. Type-locality: Venezuela, Bolívar, Auyantepuy, Río Guayaraka. Distr. — Venezuela (Bolívar).
- YEPOCAPENSE Dalmat, 1949:548, figs. 19-28. Type-locality: Guatemala, Chimaltenango, Finca Niágara, Río Sacayá, San Pedro, Yepocapa. Distr. — Mexico (Chiapas, Durango), Guatemala (Altaverapaz, Chimaltenango, Huehuetenango, El Quiché, Suchitepéquez). Refs. — Vargas & Díaz Nájera, 1954:60; Dalmat, 1955:204, figs. 70-72, 190-192, 244, 300, 340, 378, 415.
- ardeni* Dalmat, 1953:35, figs. 1-8. Type-locality: Guatemala, Alta-verapaz, San Pedro Carcha, Río Tzunutz.
- ZEMPOALENSE Vargas, Martínez Palacios & Díaz Nájera, 1946:136, 170, figs. 58-63, 133. Type-locality: Mexico, Morelos, Lagunas de Zempoala. Distr. — Mexico (Durango, Morelos, Oaxaca, Veracruz). Ref. — Vargas & Díaz Nájera, 1957b:183, 184, 187, 188, 189, 237, 247, 260, 264, pl. B, figs. 380-386.

UNPLACED SPECIES

- INFUSCATUM Lutz, 1909:137 (*Simulium*; as *venustum* var.). Type-locality: Brazil, Rio de Janeiro, and São Paulo.
- SUBVIRIDE Lutz & Machado, 1915:46 (*Simulium*). *Nomen nudum*.



LITERATURE CITED

D'ANDRETTA, C., JR.

1954. Sobre *Simulium distinctum* Lutz 1910 (Diptera Simuliidae). *An. Fac. Farm. Odontol. Univ. São Paulo* 12:63-66, 6 figs.
 (1956). Redescricao de *Simulium perflavum* Roubaud, 1906. *An. Fac. Farm. Odontol. Univ. São Paulo* 13 (1955):33-38, 11 figs.

D'ANDRETTA, C., JR. & M. A. V. D'ANDRETTA

1946. Corrigenda ao capitulo Simuliidae dos "Estudios de zoologia y parasitologia venezolanas" do Prof. A. Lutz. *Rev. Brasil. Biol.* 6(2):307-308.
 1947. Espécies neotropicais da família Simuliidae Schiner (Diptera Nematocera). II. *Lutzsimulium cruzi*, n. gen. e n. sp. e nova concepção da nervação das asas dos simulídeos. *Mem. Inst. Oswaldo Cruz* 44(3):401-412, 35 figs.
 1948a. Espécies neotropicais da família Simuliidae Schiner (Diptera Nematocera). II. *Lutzsimulium cruzi*, n. gen. e n. sp. e nova concepção da nervação das asas dos simulídeos. [Reprinted] *Mem. Inst. Oswaldo Cruz* 45(3):668-677, 35 figs.
 1948b. Espécies neotropicais da família Simuliidae Schiner (Diptera Nematocera). IV. *Lutzsimulium cruzi* Andr. & Andr., 1946, descrição do alótipo macho e da larva. *Arq. Zool. S. Paulo* 5:637-647, 38 figs.
 1950. Espécies neotropicais da família Simuliidae Schiner (Diptera Nematocera). VI. Redescricao de *Simulium pertinax* Kollar 1832. *Papéis Avulsos Dep. Zool. S. Paulo* 9:193-213, 53 figs.

D'ANDRETTA, C., JR. & D. GONZÁLEZ B.

- [1964]. Três novas espécies brasileiras de Simuliidae (Diptera Nematocera). *Mem. Inst. Butantan* 30(1960-62):103-116, 28 figs.

D'ANDRETTA, M. A. V. & C. D'ANDRETTA JR.

1946. As espécies neotropicais da família Simulidae Schiner, 1864. (Diptera-Nematocera). I. *Simulium (Eusimulium) orbitale* Lutz, 1910, *Simulium (Eusimulium) pintoi* n. sp. e *Simulium nigrimanum* Macquart, 1837, sp. inquirendae [sic]. *Mem. Inst. Oswaldo Cruz* 43:85-152, 19 pls., 2 maps.
 1947a. Redescricao de *Gigantodax wrighti* (Vargas, Martínez & Díaz [sic], 1944) (Diptera Simulidae). *Papéis Avulsos Dep. Zool. S. Paulo* 8:23-37, 32 figs.
 1947b. Espécies neotropicais da família Simulidade Schiner (Diptera Nematocera). III. *Simulium botulibranchium* Lutz, 1910 e *travassosi*, n. sp. *Papéis Avulsos Dep. Zool. S. Paulo* 8:145-180, 125 figs.
 1949. Espécies neotropicais da família "Simuliidae" Schiner (Diptera, Nematocera). V. Redescricao do "*Simulium auristriatum*" Lutz, 1910 e "*Simulium (Eusimulium) venustum infuscata*" Lutz, 1909 var. inquirendae [sic]. *Rev. Brasil. Biol.* 9(1):55-66, 42 figs.

1952. Espécies neotropicais da família Simuliidae (Diptera Nemato-cera). VII. Redescrção de *Lutzsimulium pernigrum* (Lutz, 1910), n. comb. e considerações sôbre o gênero. *Papéis Avulsos Dep. Zool. S. Paulo* 10:307-324, 49 figs.

ANDUZE, P. J.

1947. *Simulium matteabanchia*. *Bol. Acad. Cienc. Fis., Matem. y Nat.*, Caracas 10(30):374-376, 1 unnumbered fig.

BELLARDI, L.

1859. *Saggio di ditteologia messicana. Parte 1*. Pt. 1:80 pp., 2 pls. Torino.

1861a. *Idem, Mem. R. Accad. Sci. Torino* 19:201-277, 2 pls.

1862. *Saggio di ditteologia messicana, appendice. Appendice: 28 pp., 1 pl. Torino.*

1864b. *Saggio di ditteologia messicana. Appendice. Mem. R. Accad. Sci. Torino* 21:200-225, 1 pl.

BEQUAERT, J. C.

1934. Part III. Notes on the black-flies or Simuliidae, with special reference to those of the *Onchocerca* region of Guatemala, pp. 175-224, figs. 93-99, in R. P. Strong, J. H. Sandground, J. C. Bequaert & M. Muñoz Ochoa: *Onchocerciasis, with special reference to the Central American form of the disease. Contr. Dept. Trop. Med. & Inst. Trop. Biol. and Med.* 6:234 pp., 103 figs., 2 tables, 1 map. Harvard University Press, Cambridge.

1945. Dr. Luis Vargas on American black-flies — a review with critical notes (Diptera). *Bull. Brooklyn Ent. Soc.* 40(4):111-115.

EIGOT, E.

1888. Diptères, in *Mission Scient. Cap Horn, Zool.* 6:45 pp., 4 pls.

BLANCHARD, E.

1852. Diptera, in C. Gay: *Historia fisica y política de Chile, Zool.* 7: 353 pp.

BRADT, S.

1932. Notes on Puerto Rican black flies. Nota sobre la mosca negra en Puerto Rico. *Jour. Publ. Health Trop. Med.* 8(1):69-81, 2 pls.

BRICEÑO-IRAGORRY, L., L. VARGAS & A. DÍAZ NÁJERA

1953. Algunos caracteres morfológicos de *Simulium amazonicum* Goeldi 1905 (Insecta, Diptera). *Rev. Inst. Sal. Enf. Trop.*, México 13(3):243-245, 1 pl.

COQUILLET, D. W.

1902. New Diptera from North America. *Proc. U. S. Nat. Mus.* (1903) 25:83-126.



COSCARÓN, S. & P. WYGODZINSKY

1962. Simuliidae (Diptera, Insecta) de Tierra del Fuego. Patagonia e Islas Juan Fernandez. *Acta Zool. Lilloana* 18:281-333, 236 figs.

DALMAT, H. T.

1949. New species of Simuliidae (Diptera) from Guatemala. I. *Ann. Ent. Soc. Amer.* 42(4):538-553, 7 pls.
1950. New species of Simuliidae (Diptera) from Guatemala. II. *Ann. Ent. Soc. Amer.* 43(1):137-151, 5 pls.
1951. Notes on the Simuliidae (Diptera) of Guatemala, including descriptions of three new species. *Ann. Ent. Soc. Amer.* 44(1):31-58, 6 pls.
1952. Descriptions of two new species of *Simulium* (Diptera, Simuliidae) from Guatemala. *Ann. Ent. Soc. Amer.* 45(2):339-347, 1 pl.
1953. Simuliidae (Diptera) of Guatemala: Descriptions of *Simulium* (*Dyarella*) *ardeni* n. sp. and *Simulium* (*Lanea*) *jacobsi* n. sp. *Ann. Ent. Soc. Amer.* 46(1):35-42, 11 figs.
1955. The black flies (Diptera, Simuliidae) of Guatemala and their role as vectors of onchocerciasis. *Smiths. Misc. Coll.* 125(1):vii + 425 pp., 43 pls., 429 figs., maps.

DAMPF, A.

- (1928). Un simulido nuevo de Mexico (Orden Diptera, sub-orden Ne-matocera), procedente de Tiltepec, Estado de Oaxaca. *Rev. Mex. Biol.* 7(1927):125-130 (sep. 3-7).
1943. La crisalida de *Eusimulium ochraceum* (Walker 1860) (Insecta, Diptera). *Rev. Soc. Mex. Hist. Nat.* 4(1-2):33-41, pl. 2.

DÍAZ NÁJERA, A.

1961. Nota acerca de *Simulium* (*Psilopelmia*) *antillarum* Jennings, 1915. Nueva especie para Mexico. (Diptera, simuliidae [sic]). *Rev. Inst. Sal. Enf. Trop.*, México 21(1-2):79-92, 3 pls.
1962. Claves para identificar las larvas del genero *Cnephia* y descripcion de dos nuevas especies (Diptera Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 22(3-4):271-287, 26 figs.
1965. Acerca de un nuevo simulido mexicano (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 25(1):25-45, 6 pls.

DÍAZ NÁJERA, A. & M. A. VULCANO

1961. Descripcion de *Simulium* (*Psilopelmia*) *longithallum* n. sp. (Diptera, Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 21(3-4):221-235, 4 pls.
1962. Claves para identificar las larvas de simulidos del subgenero *Hearlea*, con descripcion de dos nuevas especies (Diptera, Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 22(1-2):91-144, 18 pls., 1 illustrated key.

DUNN, L. H.

1934. Entomological investigations in the Chiriqui region of Panama. *Psyche* 41:166-183.

16.26



- DYAR, H. G. & R. C. SHANNON
 1927. The North American two-winged flies of the family Simuliidae. *Proc. U. S. Nat. Mus.* 69(10):1-54, 134 figs., 7 pls.
- EDWARDS, F. W.
 1931. Simuliidae, in *British Museum (Natural History): Diptera of Patagonia and South Chile* 2(4):121-154, figs. 1-13. London.
- ENDERLEIN, G.
 1921a. Das System der Kriebelmücken (Simuliidae). *Deutsche Tierärztl. Wochenschr.* 29:197-200.
 1921b. Neue aussereuropäischen Simuliiden. *Sitz. Ges. Nat. Freund. Berlin* 4-5:77-81.
 1922. Weitere Beiträge zur Kenntniss der Simuliiden. *Konowia* 1: 67-76.
 1925. Weitere Beiträge zur Kenntniss der Simuliiden und ihrer Verbreitung. *Zool. Anz.* 62:201-211.
 1929. Über einige neotropische Simuliiden der Genus *Friesia* (Diptera). *Deutsch. Ent. Zeitschr.* (1929):327-328, 2 figs.
 1930. Der heutige Stand der Klassifikation der Simuliiden. *Arch. f. Klassifikatorische u. Phylogenetische Ent.* 1:77-97, 7 figs.
 1933. Weiterer Ausbau des Systems der Simuliiden (Diptera). *Deutsch. Ent. Zeitschr.* 1933:273-292.
 1934. Aussereuropäische Simuliiden aus dem Wiener Museum. *Sitz. Ges. Naturf. Freund. Berlin* 1934:190-195.
 1936. Simuliologica. I. *Sitz. Ges. Naturf. Freund. Berlin* 1936:113-130.
- FAIRCHILD, G. B.
 1940. Notes on the Simuliidae of Panama. *Ann. Ent. Soc. Amer.* 33(4):701-719, 39 figs.
- FLOCH, H. & E. ABONNENC
 1946a. Simulidés de la Guadeloupe: *S. antillarum* Jennings et *S. tarsale* Williston. *Publ. Inst. Pasteur Guyane Franc. et Terr. de l'Inini* 130:1-6, 2 pls.
 1946b. Simulides de la Guyane Française. I. *S. guianense* Wise 1911. *S. Rorotaense* n. sp., *S. Maroniense* n. sp. *Publ. Inst. Pasteur Guyane Franç. et Terr. de l'Inini* 136:1-20, 10 pls.
 1946c. Simulides de la Guyane Française (II). *S. cauchense*, n. sp., *S. oyapockense* n. sp., *S. iracouboense* n. sp. *Publ. Inst. Pasteur Guyane Franç. et Terr. de l'Inini* 137:1-19, 6 figs.
- FOX, I.
 1953. Notes on Puerto Rican Simuliidae from light traps. *Proc. Ent. Soc. Washington* 55(3):135-142, 10 figs.
- FRIES, B. F.
 1824. *Observaciones entomologicae. Pars I*: 20 pp., 1 pl. Lund.



GOELDI, E. A.

1905. Os mosquitos no Pará. *Mem. Mus. Hist. Nat. Ethnogr. (Museu Goeldi)* 4:1-154, 144 figs., 5 pls.

HEARLE, E.

1929. A remarkable simuliid pupa. Notes on *Simulium virgatum* in British Columbia. *Proc. Ent. Soc. Brit. Columbia* 25:48-54, figs. A-G, 2 pls.
1932. The blackflies of British Columbia (Simuliidae, Diptera). *Proc. Ent. Soc. Brit. Columbia* 29:5-19, 7 figs.
1935. Notes on *Simulium canadense* Hearle and *S. virgatum* Coquillett and its varieties. *Canad. Ent.* 67:15-19, illus.

HOFFMANN C., C.

- 1930a. Un *Simulium* nuevo de la zona cafetera de Chiapas. *An. Inst. Biol. Mex.* 1(1):51-53, 4 figs.
- 1930a. Un *Simulium* nuevo de la zona cafetera de Chiapas. *An. Inst. Biol. Mex.* 1(4):293-306, 15 figs.
1931. Los simúlidos de la region onchocercosa de Chiapas. Segunda parte. Los estados larvales. *An. Inst. Biol. Mex.* 2(3):207-218. 18 figs.

IRIARTE, D. R.

1946. La familia Simuliidae en Venezuela. *Bol. Laborat. Clínica "Luis Razetti"*, Caracas 15(21-22):401-479, figs. 8-17, 5 pls., 2 maps.

JENNINGS, A.

1915. Two new species of Simuliidae from tropical America. *Proc. Ent. Soc. Washington* 17(4):199-200.

JOAN, T.

1912. Nota sobre un díptero ponzoñoso. *Bol. Minist. Agric. Buenos Aires* 14:363-385, 11 figs., 2 pls. (sep. 25 pp.).

JOHANNSEN, O. A.

1903. Part 6. Aquatic nematoceros Diptera, pp. 328-441, pls. 32-50. in J. G. Needham, A. D. MacGillivray, O. A. Johannsen & K. C. Davis: Aquatic insects of New York State. *Bull. N. Y. State Mus.* 68 (= Ent. 18): 199-517, 26 figs., 52 pls. (= *Bull. N. Y. State Univ.* 295).

KNAB, F.

1913. A note on some American Simuliidae. *Insec. Insc. Menstr.* 1(12): 154-156.
- 1914a. New data and species in Simuliidae (Diptera). *Insec. Insc. Menstr.* 2(12):177-180.
- 1914b. Simulididae de Chile septentrional. *An. Zool. Aplicada Chile* 1(7):17-22, pl.



- 1914c. Simuliidae of Peru. *Proc. Biol. Soc. Washington* 27:81-86.
 1914d. Supplementary notes on Peruvian Simuliidae. *Proc. Biol. Soc. Washington* 27:123-124.
 1915a. Some new Neotropical Simuliidae. *Bull. Ent. Res.* 6(3):279-282.
 1915b. A new *Simulium* from Texas (Diptera, Simuliidae). *Insec. Insc. Menstr.* 3(1-4):77-78.

KOLLAR, V.

1832. In I. E. Pohl & V. Kollar: *Vorzüglich lästige Insecten*, 20 pp., 1 pl. in I. E. Pohl: *Reise im Innern von Brasilien*. Wien.

LANE, J. & C. E. PORTO

1939. Simuliídeos da região neotrópica: O gênero *Eusimulium*. *Bol. Biol.* (n.s.), São Paulo 4(2):168-176, 7 figs.
 1940. Simuliídeos da região neotropical. III. Descrição de novas espécies dos gêneros "*Simulium*" e "*Eusimulium*". *Arq. Inst. Biol.*, São Paulo 11:189-195, 16 figs.

LANE, J. & M. A. VULCANO

1943. A armadura bucal dos simuliídeos [sic] e seu valor taxonômico (Diptera, Simuliidae). *Rev. Entomol.*, Rio de Janeiro 14(3):430-440, 28 figs.

LATREILLE, P. A.

- [1802]. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes* 3:xii + 468 pp. Paris, "An X".

LEÓN, J. R. DE

1943. Preliminares para la descripción de cinco nuevas especies de simúlidos en Guatemala. *Bol. Sanit. Guatemala* 51:94-101, 7 figs., 1 map.
 1945. Nuevas especies de simúlidos en la región occidental de Guatemala. *Bol. Sanit. Guatemala* 52:66-77, 16 figs., 1 photograph, 1 map (separate, pp. 1-12).
 1948. *Apuntes para una monografía sobre los Simulidae de Guatemala. Un nuevo simúlido de la región occidental de Guatemala*, 23 pp., 19 figs. Universidad de San Carlos de Guatemala, Instituto de Investigaciones Científicas, Departamento de Investigaciones Científicas, Guatemala.

LEÓN, L. A. & P. WYGODZINSKY

1953. Los simúlidos del Ecuador (Diptera), su importancia en medicina humana. *Bol. Inform. Cient. Nacion.*, Quito 6(57):269-288, 7 figs.



LOEW, H.

1862. Monographs of the Diptera of North America. Part. I. *Smiths. Misc. Coll.* 6(1):xxiv + 221 pp., 12 figs., 2 pls.

LUNDSTRÖM, C.

1911. Beiträge zur Kenntniss der Dipteren Finlands. VII. Melusini-
dae (Simuliidae). *Acta Soc. Fauna et Flora Fenn.* 34(12):1-23,
1 pl.

LUTZ, A.

1909. Contribuição para o conhecimento das especies brasileiras do
gênero "*Simulium*". *Mem. Inst. Oswaldo Cruz* 1(2):124-146.
1910. Segunda contribuição para o conhecimento das especies bra-
zileiras do genero "*Simulium*". *Mem. Inst. Oswaldo Cruz* 2(2):
213-267, pls. 18-21.
1917. Terceira contribuição para o conhecimento das especies brazi-
leiras do genero *Simulium*. O piúm do norte (*Simulium amaz-
onicum*). *Mem. Inst. Oswaldo Cruz* 9(1):63-67, pl. 25.
1922. Zoologia medica. Nematoceros hematophagos não pertencen-
tes aos culicídeos. *A Folha Medica*, Rio de Janeiro 3:89-92.

LUTZ, A. & A. MACHADO

1915. Viagem pelo São Francisco e por alguns dos seus afluentes entre
Pirapora e Joazeiro. *Mem. Inst. Oswaldo Cruz* 7(1-2):5-49, pls.
1-18.

LUTZ, A. & M. NUÑEZ TOVAR

1928. *Estudios de zoologia y parasitologia venezolanas*, 133 pp., 26 pls.
Rio de Janeiro.

MACQUART, J.

1838. Diptères exotiques nouveaux ou peu connus. Part. 1. *Mém.
Soc. Roy. des Sci., de l'Agric. et des Arts Lille* 1838(2):9-225, 25 pls.
(also published separately as Vol. 1, part 1, pp. 5-221, 25 pls.,
Paris, 1838).
1846. Diptères exotiques nouveaux ou peu connus. Supplément [I].
Mém. Soc. Roy. des Sci., de l'Agric. et des Arts Lille (1845) 1844:
133-364, 20 pls. (also published separately as Supplément [I], pp.
5-238, 20 pls., Paris, 1846).

MALLOCH, J. R.

1912. One new genus and eight new species of dipterous insects in the
United States National Museum collection. *Proc. U. S. Nat. Mus.*
(1913) 43:649-658, pl. 46.
1913. A new species of *Simulium* from Texas. *Proc. Ent. Soc. Was-
hington* 15:133-134.
1914. American black flies or buffalo gnats. *U. S. Dept. Agric., Bur.
Entomol. Techn. Series* 26:1-72, 6 pls.



NETTEL, F. R.

1952. Revisión del problema entomológico de la oncocercosis y plan para la erradicación de *Simulium ochraceum* Walker. *Med. Rev. Mex.* 32(660):438-441.

ORTIZ c., I.

1957. Nuevos representantes hematofagos de los generos *Culicoides* (Diptera, Ceratopogonidae) y *Simulium* (Diptera: Simuliidae) de Venezuela. *Bol. Venez. Laborat. Clinica "Luis Razetti"*, Caracas 2(3-4):161-168, 3 pls., 2 photographs.

PATERSON, G. & R. C. SHANNON

1927. Los simúlidos del noroeste argentino. *Rev. Inst. Bacteriol. Depart. Nac. Hig.* 4(7):737-742.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool. bot. Ges. Wien* 15:595-782, pls. 23-29.

PINTO, C.

1931. Simuliidae da America Central e do Sul. 7º *Reun. Soc. Argent. Patol. Reg. Norte* 60:661-763.

PORTO, C.

1940. Simuliideos da região neotropical. *Arq. Zool. São Paulo* 1(12): 383-385.

RILEY, C. V.

1887. Report of the entomologist. *U. S. Dept. Agric. Com. Agric. Rept.* 1886:459-592, 11 pls.

RONDANI, C.

1864. Diptorum species et genera aliqua exotica revisa et annotata novis nonnullis [sic] descriptis. *Arch. per la Zool. l'Anat. e la Fisiol.*, Modena 3(1):23-99, pl. 5.

ROUBAUD, M. E.

- 1906a. Aperçus nouveaux, morphologiques et biologiques sur les diptères piqueurs du groupe des Simulies. *C. R. Acad. Roy. Sci. Paris* 1906:519-521 (sep. 1-3).

- 1906b. Insectes Diptères, Simulies nouvelles ou peu connus. *C. R. Acad. Roy. Sci.* 7:517-522.

- 1906c. Simulies nouvelles de l'Amérique du Sud. *Bull. Mus. Hist. Nat.* 2:106-109 (sep. 1-4).

1909. Description d'une Simulie nouvelle du Peru. *Bull. Soc. Pathol. exot.* 20:428-431.



RUBZOV, I. A.

1940. Moshki (Sem. Simuliidae) [Black-flies. Fam. Simuliidae], in Zoologicheskii Institut Akademii Nauk SSSR: *Fauna SSSR* (Novaia Seria n° 23). *Nasekomye Dvukrylye* 6(6):533 pp., 93 figs. (Reprinted, 1963, *Fauna SSSR* (Novaia Seria n° 64), *Nasekomye Dvukrylye* 6(6):860 pp., 424 pls.). Izdatelstvo Akademii Nauk SSSR, Leningrad.

SCHROTTKY, C.

1909. Drei neue blutsaugende Dipteren aus Paraguay. *Zeitschr. Wiss. Insekt. Biol.* 5:61-63.

SILVA FIGUEROA, C.

1917. Dos nuevos simúlidos de Chile. *Bol. Mus. Nacion. Chile* 10:28-35, 1 pl., figs. 16-20.

SMART, J.

1940. Simuliidae (Diptera) from British Guiana and the Lesser Antilles. *Trans. R. ent. Soc. London* 90:1-11, 4 pls., 3 figs.
1942. Notes on Simuliidae (Diptera). *Proc. R. ent. Soc. London* (B) 11(3):46-50, 1 pl.
1944. Notes on Simuliidae (Diptera). II. *Proc. R. ent. Soc. London* (B) 13(11-12):131-136.
1945. The classification of the Simuliidae (Diptera). *Trans. R. ent. Soc. London* 95(8):463-532.

SPEISER, P.

1904. Zur Nomenklatur blutsaugender Diptera. *Amerikas Insektenboerse* 2:148.

STAINS, G. S. & G. F. KNOWLTON

1943. A taxonomic and distribution study of Simuliidae of western United States. *Ann. Ent. Soc. Amer.* 36(2):259-280, 133 figs.

STONE, A.

1948. *Simulium virgatum* Coquillett and a new related species (Diptera, Simuliidae). *Jour. Washington Acad. Sci.* 38(12):399-404, 10 figs.
1952. The Simuliidae of Alaska. *Proc. Ent. Soc. Washington* 54(2):69-96.
1962. Notes on the types of some Simuliidae (Diptera) described by Enderlein. *Ann. Ent. Soc. Amer.* 55:206-209.
1963. An annotated list of genus-group names in the family Simuliidae (Diptera). *U. S. Dept. Agric. Techn. Bull.* 1284:28 pp.

SURCOUF, J. M. & R. GONZÁLEZ RINCONES

1911. *Essai sur les diptères vulnérants de Venezuela*, 320 pp.



TONNOIR, A. L.

1925. Australasian Simuliidae. *Bull. Ent. Res.* 15(3):213-265, 13 pls.

TOWNSEND, C. H. T.

1891. A new Simulium from southern New Mexico. *Psyche* 6:106-107.

1897. Diptera from the lower Rio Grande or Tamaulipan region of Texas. *I. Jour N. Y. Ent. Soc.* 5:171-172.

VARGAS, L.

1941. *Simulium lane-portoi* n. n. (Simuliidae, Dipt.) y lista de simúlidos mexieanos. *Rev. Inst. Sal. Enf. Trop.*, Méxio 2(1):115-122.

1942a. Notas sobre la terminalia de algunos simulidos de Mexico. *S. (E.) paynei* n. n. Vargas 1942. *Rev. Inst. Sal. Enf. Trop.*, Méxio 3(3):229-249, 15 figs.

1942b. *Simulium fairchildi* Vargas 1942 n. n. de Panama. (Diptera Simuliidae). *Rev. Med. Mex.* 22(420):458-459.

1943a. Nombres y datos nuevos de simulidos del Nuevo Mundo. *Rev. Soc. Mex. Hist. Nat.* 4(3-4):135-146, pls. 6-7.

1943b. Nuevos datos sobre simulidos mexicanos (Dipt. Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, Méxio 4(4):359-370, 6 pls.

1945a. Cuatro nuevas especies y otros datos sobre simulidos de Mexico. *Rev. Soc. Mex. Hist. Nat.* 6(1-2):71-82, pls. 7-12.

1945b. Tres nuevos nombres propuestos para simulidos del Nuevo Mundo. *Rev. Med. Trop. Parasitol.*, Habana 11:4-5.

1945e. Simulidos del Nuevo Mundo. *Monogr. Inst. Sal. y Enf. Trop.*, México 1:vi + 241 pp., 10 pls.

1946. Contribucion al conocimiento de los simulidos de la region onco-cereosica de Chiapas, Mexico. *Simulium (Eusimulium) smarti* n. sp. (Diptera Simuliidae). *Puerto Rico Jour. Publ. Health. Med.* 21:330-335, 5 figs.

VARGAS, L. & A. DÍAZ NÁJERA

1948a. *Simulium clarki* Fairehild (1940) y *Simulium lutzianum* Pinto (1931). Estructuras genitales. *Rev. Inst. Sal. Enf. Trop.*, Méxio 9(1):75-79, 5 figs.

1948b. Nuevas especies de simulidos de Mexico y consideraciones diversas sobre especies ya descritas. *Rev. Inst. Sal. Enf. Trop.*, México 9(4):321-369, 12 pls.

1949. Claves para identificar las pupas de los simulidos de Mexico. Descripción de *Simulium (Dyarella) freemani* n. sp., de *Simulium (Neosimulium) enciso* n. sp. y referencias adicionales sobre *S. anduzei* y *S. ruizi*. *Rev. Inst. Sal. Enf. Trop.*, México 10(4):283-319, 10 pls.

1951. Notas sobre sistematica y morfologia de simulidos. *Rev. Soc. Mex. Hist. Nat.* 12(1-4):123-172, 18 pls.

1952. Descripción de la larva de *Cnephia aguirrei* (Dalmat 1949). *Rev. Soc. Mex. Hist. Nat.* 13(1-4):53-56, 4 figs.

- 1953a. Nota sobre simulidos del Ecuador. *Rev. Ecuat. Ent. Parasitol.* 1(4):17-21, 8 figs.
- 1953b. Nota sobre el examen de tipos de simulidos descritos por prof. G. Enderlein. *Rev. Inst. Sal. Enf. Trop.* 13(2):137-149, 3 pls.
- 1953c. *Simulium* (*Notolepria*) *gonzalesi* n. sp. (Insecta Diptera). *Rev. Inst. Sal. Enf. Trop.*, México 13(3):235-239, 2 pls.
1954. Algunas consideraciones morfológicas y de nomenclatura relativas a simulidos americanos (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 14(2):57-72, 8 pls.
1956. *Simulium* (*Hemicnetha*) *guerrerense* n. sp. (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 16(3):51-56, 15 figs.
- 1957a. Simulidos de Durango y *Simulium* (*Simulium*) *marquezi* n. sp. (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 17(1):33-41, 6 pls.
- 1957b. Simulidos mexicanos. *Rev. Inst. Sal. Enf. Trop.*, México 17(4):143-399, 65 pls.
1958. Nota sobre *Simulium* (*Psilopelmia*) *bivittatum* Malloch 1914 (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 18(1):13-30, 2 pls.
1959. Claves graficas para identificar generos y subgeneros de larvas, pupas y adultos de simulidos (Diptera: Simuliidae). *Rev. Inst. Sal. Enf. Trop.* 19(1):105-114.

VARGAS, L., A. DÍAZ NÁJERA & A. MARTÍNEZ PALACIOS

1943. Tres simulidos nuevos para Mexico. *Rev. Inst. Sal. Enf. Trop.*, México 4(3):287-290, 2 pls.

VARGAS, L., A. MARTÍNEZ PALACIOS & A. DÍAZ NÁJERA

1944. *Simulium* (*Gigantodax*) *wrighti* n. sp. en Mexico (Diptera Simuliidae). *Rev. Inst. Sal. Enf. Trop.*, México 5(1):37-41, 2 pls.
1946. Simulidos de Mexico. Datos sobre sistematica y morfologia. Descripcion de nuevos subgeneros y especies. *Rev. Inst. Sal. Enf. Trop.*, México 7(3):95-192, 25 pls.

VULCANO, M. A. (= M. A. V. D'ANDRETTA)

1958. Redescrição do *Simulium rubrithorax* Lutz, 1909, e descrição do alotipo ♂ (Diptera, Simuliidae). *Papéis Avulsos Dep. Zool. São Paulo* 13:227-240, 50 figs.
- 1959a. Descrição de *Simulium obesum*, sp. n. (Diptera, Simuliidae). *Papéis Avulsos Dep. Zool. São Paulo* 13:351-360, 44 figs.
- 1959b. Descrição do alotipo de *Simulium pruinosum* Lutz, 1910 e caracteres adicionais da fêmea. *Mem. Inst. Oswaldo Cruz* 57(1):33-43, 10 pls.
1962. Simulídeos da Serra da Bocaina. Observações bionômicas, revalidação de *Simulium brevifurcatum* Lutz e descrição de uma espécie nova, *S. ana-mariae* (Diptera, Simuliidae). *Papéis Avulsos Dep. Zool. São Paulo* 15:239-272, 89 figs., 1 map, 1 photograph.



WALKER, F.

1861. Characters of undescribed Diptera in the collection of W. W. Saunders [concl.]. *Trans. Ent. Soc. London* (n.s.) 5(1858-1861): 297-334.

WILLISTON, S. W.

1893. List of Diptera of the Death Valley Expedition, pp. 253-259, in C. V. Riley: *The Death Valley Expedition. A biological survey of parts of California, Nevada, Arizona, and Utah. Part II. 4. Report on a small collection of insects made during the Death Valley Expedition.* *North Amer. Fauna* 6:235-268.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896:253-446, pls. 8-14.

WISE, K. S.

1911. The Simuliidae of British Guiana. *Jour. Roy. Agric. Comm. Soc. British Guiana* 1(3):248-254.

1912. The itanimi fly. A new *Simulium* in British Guiana. *Jour. Trop. Med. Hyg.* 15:43.

WYGODZINSKY, P.

- 1949a. Contribuciones al conocimiento de los Simuliidae argentinos. I. Introducion. Redescrpcion de *Simulium lahillei* Paterson & Shannon, 1927 (Diptera). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 2(3):303-319, 7 figs.

- 1949b. Contribuciones al conocimiento de los Simuliidae argentinos II. *Gigantodax horcotiani* sp. n., de la Provincia de Tucumán (Diptera). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 2(3):325-333, 3 figs.

1950. Contribuciones al conocimiento de los Simuliidae argentinos. III. *Simulium dinellii* (Joan, 1912) y *Simulium wolffhügeli* (Enderlein, 1922). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(1):75-97, 15 figs.

- 1951a. Sobre *Gigantodax horcotiani* Wygodzinsky, 1949 y *Gigantodax wittmeri* sp. n. (Diptera, Simuliidae). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(2):199-206, 44 figs.

- 1951b. Sobre *Simulium jujuyense* Paterson y Shannon, 1927, *Simulium exiguum* Roubaud, 1906, y *Simulium opalinifrons* (Enderlein, 1934). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(2):207-220, 80 figs.

- 1951c. Sobre *Simulium ignescens* Roubaud, 1906, *Simulium romañai* sp. n. y *Simulium adolfolutzi* sp. n. (Diptera, Simuliidae). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(2):221-236, 103 figs.

1952. Los insectos de las Islas Juan Fernandez. Simuliidae (Diptera). *Rev. Chilena Entomol.* 2:81-85, 23 figs.

- 1953a. Sobre algunos simulidos argentinos (Diptera). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(3):293-320, 127 figs.

- 1953b. Sobre algunos simulidos de los países andinos (Diptera). *An. Inst. Med. Reg. Univ. Nac. Tucumán* 3(3):321-337, 104 figs.

1958. Notas y descripciones de "Simuliidae" patagónicas y andinas (Diptera). *Acta Zool. Lilloana* 16:121-148, 141 figs.



1959. Diptera, Simuliidae, in J. F. R. Bejarano, E. Del Ponte & R. N. Orfila, eds.: *Primeras Jornadas Entomoepidemiologicas Argentinas* 2:563-564. Buenos Aires.

1961. Sobre una nueva especie de *Simulium* de la Provincia de Buenos Aires (Simuliidae, Diptera). *Acta Kravsi, Cuad. del Inst. Nac. Microbiol.*, Buenos Aires 2:30-36, 66 figs.

WYGOZINSKY, P. & S. COSCARÓN

1962. On the relationship and zoogeographical significance of *Austrosimulium anthracinum* (Bigot), a blackfly from southern South America. *Pacific Insects* 4(1):235-244, 4 figs.

ZETTERSTEDT, J. W.

1838. *Dipterologia Scandinaviae*. Sect. 3: Diptera, pp. 477-868, in his *Insecta Lapponica*, vi + 1140 pp. Lipsiae [Leipzig].

INDEX

abalosi Wygodzinsky, 1958, Gigantodax	16.2
acatenangoensis Dalmat, 1951, Simulium	16.18
Acropogon Enderlein, 1933	16.5
adolfolutzi Wygodzinsky, 1951, Simulium	16.5
aequatoriense Vargas, 1945, Simulium (error)	16.11
aegifurcatum Lutz, 1910, Simulium	16.8
aguirrei (Dalmat), 1949 (Simulium), Cnephia	16.2
albicinctum (Enderlein), 1933 (Acropogon), Simulium	16.6
albilineatum (Enderlein), 1936 (Pternaspatha), Simulium	16.6
albopictum Lane & Porto, 1940, Simulium	16.6
alticolum (Dyar & Shannon), 1927 (Eusimulium), Simulium	16.6
amazonicum Goeldi, 1905, Simulium	16.6
amazonicum Goeldi of Lutz, 1910, Simulium	16.19
anamariae Vulcano, 1962, Simulium	16.6
anduzei Vargas & Diaz Nájera, 1948, Simulium	16.6
angrense Pinto, 1931, Simulium	16.6
angustifrons (Enderlein), 1933 (Trichodagmia), Simulium	16.15
angustitarsis (Enderlein), 1933 (Trichodagmia), Simulium	16.13
annulatum Philippi, 1865, Simulium	16.6
antarcticum (Bigot), 1888 (Simulium), Gigantodax	16.2
anthracinum (Bigot), 1888 (Simulium), Austrosimulium	16.1
antillarum Jennings, 1915, Simulium	16.6
antonii Wygodzinsky, 1953, Simulium	16.6
antunesi (Lane & Porto), 1940 (Eusimulium), Simulium	16.6
aquamarense (León), 1945 (Simulium), Gigantodax	16.2
araucaium (Edwards), 1931 (Simulium), Gigantodax	16.3
ardeni Dalmat, 1953, Simulium	16.23
argentatum (Enderlein), 1936 (Thyrsopelma), Simulium	16.6
argus Williston, 1893, Simulium	16.6
atoparva (Edwards), 1931 (Simulium), Cnephia	16.2
atompensis Diaz Nájera, 1962, Cnephia	16.2
aureopunctatum Malloch, 1914, Simulium	16.16



<i>aureum</i> (Frics), 1824 (Simulia), Simulium	16.7
<i>auripellitum</i> Enderlein, 1933, Simulium	16.7
<i>auristriatum</i> Lutz, 1910, Simulium	16.7
<i>Austrosimulium</i> Tonnoir, 1925	16.1
<i>avidum</i> Hoffmann, 1930, Simulium	16.16
<i>ayrosai</i> Vargas, 1945, Simulium	16.7
<i>baiense</i> Pinto, 1931, Simulium	16.7
<i>barbatipes</i> (Enderlein), 1933 (Acropogon), Simulium	16.7
<i>benjamini</i> Dalmat, 1952, Simulium	16.7
<i>bicoloratum</i> Malloch, 1913, Simulium	16.7
<i>bilimekae</i> Smart, 1944, Simulium	16.18
<i>bipunctatum</i> Malloch, 1912, Simulium	16.10
<i>bivittatum</i> Malloch, 1914, Simulium	16.7
<i>bolivianum</i> Enderlein, 1925, Gigantodax	16.3
<i>bonorinorum</i> Coscarón & Wygodzinsky, 1962, Gigantodax	16.3
<i>botulibranchium</i> Lutz, 1910, Simulium	16.7
<i>botulibranchium</i> Lutz of Pinto, 1931, Simulium	16.22
<i>boydi</i> León, 1945, Simulium	16.13
<i>brachycladum</i> Lutz & Pinto, 1931, Simulium	16.7
<i>brasiliense</i> (Enderlein), 1933 (Thysopelma), Simulium	16.17
<i>brevifurcatum</i> Lutz, 1910, Simulium	16.8
<i>bricenoi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.8
<i>brophyi</i> (Edwards), 1931 (Simulium), Gigantodax	16.3
<i>burchi</i> Dalmat, 1951, Simulium	16.8
<i>bustosi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.8
<i>Byssodon</i> Enderlein, 1925	16.5
<i>callidum</i> (Dyar & Shannon), 1927 (Eusimulium), Simulium	16.8
<i>canadense</i> Hearle, 1932, Simulium	16.8
<i>canadense</i> Hearle of Vargas, 1943, Simulium	16.11
<i>capricorne</i> León, 1945, Simulium	16.8
<i>carolinae</i> León, 1945, Simulium	16.8
<i>catarinense</i> Pinto, 1931, Simulium	16.8
<i>cauchense</i> Floch & Abonnenc, 1946, Simulium	16.8
<i>chalcocoma</i> Knab, 1914, Simulium	16.9
<i>chalcocomense</i> Vargas, 1945, Simulium (error)	16.9
<i>chilapanense</i> Hoffmann, 1930, Simulium	16.23
<i>chilense</i> (Philippi), 1865 (Simulium), Gigantodax	16.3
<i>chilianum</i> Philippi, 1864, Simulium	16.9
<i>Chirostilbia</i> Enderlein, 1921	16.4
<i>cinereum</i> Bellardi, 1859, Simulium	16.23
<i>clarki</i> Fairchild, 1940, Simulium	16.9
<i>clavibranchium</i> Lutz, 1910, Simulium	16.9
<i>clavipes</i> Malloch, 1914, Simulium	16.21
<i>Cnephia</i> Enderlein, 1921	16.2
<i>coffae</i> Vargas, 1945, Simulium	16.11
<i>colvini</i> Dalmat, 1952, Simulium	16.20
<i>contrerense</i> Díaz Nájera & Vulcano, 1962, Simulium	16.9
<i>costalimai</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.9
<i>costaricense</i> Vargas, 1945, Simulium (error)	16.9
<i>costaricense</i> Smart, 1944, Simulium	16.9
	16.37

<i>crusi</i> d'Andretta & d'Andretta, 1947, Lutzsimulium	16.4
<i>Cryptectemnia</i> Enderlein, 1936	16.5
<i>cuneatum</i> (Enderlein), 1936 (Psaroniocompsa), Simulium	16.9
<i>dalmati</i> Vargas & Díaz Nájera, 1948, Simulium	16.9
<i>dandrettai</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.9
<i>Dasypelmoza</i> Enderlein, 1933	16.5
<i>deagostinii</i> Coscarn & Wygodzinsky, 1962, Simulium	16.9
<i>delatorrei</i> Dalmat, 1950, Simulium	16.9
<i>deleoni</i> Vargas, 1945, Simulium	16.9
<i>delpontei</i> Paterson & Shannon, 1927, Simulium	16.11
<i>delponteianum</i> Wygodzinsky, 1961, Simulium	16.10
<i>díazi</i> León, 1945, Simulium	16.7
<i>diazajerai</i> Vargas, 1943, Simulium	16.10
<i>dinellii</i> (Joan), 1912 (Melusina), Simulium	16.10
<i>dissimilis</i> (Edwards), 1931 (Simulium), Cnephia	16.2
<i>distinctum</i> Lutz, 1910, Simulium	16.10
<i>distinctum</i> Malloch, 1913, Simulium	16.22
<i>distinctum</i> Lutz of Pinto, 1931, Simulium	16.18
<i>diversibranchium</i> Lutz, 1910, Simulium	16.10
<i>diversifurcatum</i> Lutz, 1910, Simulium	16.10
<i>donovani</i> Vargas, 1943, Simulium	16.7
<i>downsi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium.	16.10
<i>dugesi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium.	16.10
<i>earlei</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium.	16.11
<i>earli</i> Vargas & Díaz Nájera, 1957, Simulium (error)	16.11
<i>Ectemnaspis</i> Enderlein, 1933	16.5
<i>ecuadoriense</i> (Enderlein), 1934 (Trichodagmia), Simulium	16.11
<i>edwardsi</i> (Enderlein), 1933 (Acropogon), Simulium	16.17
<i>encisoii</i> Vargas & Díaz Nájera, 1949, Simulium	16.11
<i>cquadoriense</i> León & Wygodzinsky, 1953, Simulium (error)	16.11
<i>escomeli</i> Roubaud, 1906, Simulium	16.11
<i>estevezi</i> Vargas, 1945. Simulium	16.11
<i>ethelae</i> Dalmat, 1950, Simulium	16.11
<i>Eusimulium</i> Roubaud, 1906	16.4
<i>exiguum</i> Roubaud, 1906, Simulium	16.11
<i>exiguum</i> Roubaud of Knab, 1913, Simulium	16.12
<i>exiguum</i> Roubaud of Lutz, 1910, Simulium	16.6
<i>fairchildi</i> Vargas, 1942, Simulium	16.19
<i>fulcatum</i> (Enderlein), 1929 (Friesia), Simulium	16.11
<i>femineum</i> (Edwards), 1931 (Simulium), Gigantodax	16.3
<i>figuroa</i> Smart, 1944, Simulium	16.11
<i>flavifemur</i> (Enderlein), 1921 (Chirostilbia), Simulium	16.11
<i>flavipictum</i> Knab, 1914, Simulium	16.12
<i>flavopictum</i> Pinto, 1931, Simulium (error)	16.12
<i>flavopubescens</i> Lutz. 1910. Simulium	16.12
<i>freemani</i> Vargas & Díaz Nájera, 1949, Simulium	16.12
<i>Friesia</i> Enderlein, 1922	16.5
<i>fulvescens</i> (Blanchard), 1852 (Simulium), Gigantodax	16.3
<i>ganalesense</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.12



<i>ganalesense</i> Nettel, 1952, <i>Simulium</i> (error)	16.12
<i>gandeatum</i> Iriarte, 1946, <i>Simulium</i> (error)	16.12
<i>gaudcatum</i> Knab, 1914, <i>Simulium</i>	16.12
<i>Gigantodax</i> Enderlein, 1925	16.2
<i>glaucophthalmum</i> Knab, 1914, <i>Simulium</i>	16.11
<i>gonzalesi</i> Vargas & Díaz Nájera, 1953, <i>Simulium</i>	16.12
<i>gorirossiae</i> Vargas & Díaz Nájera, 1957, <i>Simulium</i>	16.12
<i>grenieri</i> Vargas & Díaz Nájera, 1948, <i>Cnephia</i>	16.2
<i>guatemalensis</i> León, 1945, <i>Simulium</i>	16.14
<i>guerrrense</i> Vargas & Díaz Nájera, 1956, <i>Simulium</i>	16.12
<i>gulanense</i> Wise, 1911, <i>Simulium</i>	16.12
<i>guttatum</i> (Enderlein), 1936 (<i>Pliodasina</i>), <i>Simulium</i>	16.12
<i>guyanense</i> Floch & Abonnenc, 1946, <i>Simulium</i> (error)	16.18
<i>Eynangra</i> (Edwards), 1931 (<i>Simulium</i>), <i>Cnephia</i>	16.2
<i>haarupianum</i> (Enderlein), 1936 (<i>Trichodagmia</i>), <i>Simulium</i>	16.23
<i>haematopotum</i> Malloch, 1914, <i>Simulium</i>	16.12
<i>haematopotum</i> Malloch of Fairchild, 1940, <i>Simulium</i>	16.19
<i>Hearlea</i> Rubzov, 1940 (nomen nudum)	16.5
<i>Hearlea</i> Vargas, Martínez Palacios & Díaz Nájera, 1946	16.5
<i>hebeticolor</i> Lutz, 1910, <i>Simulium</i> (error)	16.20
<i>hechti</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.13
<i>Hemicnetha</i> Enderlein, 1934	16.5
<i>hinmani</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.13
<i>hippovororum</i> Malloch, 1914, <i>Simulium</i>	16.23
<i>hirticosta</i> (Lutz), 1909 (<i>Simulium</i>), <i>Lutzsimulium</i>	16.4
<i>hirtipes</i> Fries of Vargas, 1943, <i>Simulium</i>	16.14
<i>hirtipupa</i> Lutz, 1910, <i>Simulium</i>	16.13
<i>hoffmanni</i> Vargas, 1943, <i>Simulium</i>	16.13
<i>horcotiani</i> Wygodzinsky, 1949, <i>Gigantodax</i>	16.3
<i>huairayacu</i> Wygodzinsky, 1953, <i>Simulium</i>	16.13
<i>ignescens</i> Roubaud, 1906, <i>Simulium</i>	16.13
<i>igniculum</i> Coscarón & Wygodzinsky, 1962, <i>Gigantodax</i>	16.3
<i>inaequale</i> (Paterson & Shannon), 1927 (<i>Eusimulium</i>), <i>Simulium</i>	16.13
<i>inaicum</i> Vargas, 1945, <i>Simulium</i>	16.20
<i>incaicum</i> Vargas, 1945, <i>Simulium</i>	16.13
<i>incertum</i> Lutz, 1910, <i>Simulium</i>	16.13
<i>incrustatum</i> Lutz, 1910, <i>Simulium</i>	16.13
<i>inexorabile</i> Schrottky, 1909, <i>Simulium</i>	16.13
<i>infuscatum</i> Lutz, 1909, <i>Simulium</i> (sp. inquirenda)	16.23
<i>infuscatum</i> Lutz, 1910, <i>Simulium</i>	16.7
<i>iracoboense</i> Floch & Abonnenc, 1946, <i>Simulium</i>	16.13
<i>iriartei</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.13
<i>itaunensis</i> d'Andretta & González B., 1964, <i>Simulium</i>	16.14
<i>jacobsi</i> Dalmat, 1953, <i>Simulium</i>	16.14
<i>jacumbae</i> Dyar & Shannon, 1927, <i>Simulium</i>	16.14
<i>jacumbae</i> Dyar & Shannon of Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.9
<i>jobbinsi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.14
<i>johnsoni</i> Vargas & Díaz Nájera, 1957, <i>Simulium</i>	16.14
	16.39

juarezi Vargas & Díaz Nájera, 1957, <i>Simulium</i>	16.14
jujuyense Paterson & Shannon, 1927, <i>Simulium</i>	16.14
<i>jujuyense</i> Paterson & Shannon of Vargas & Díaz Nájera, 1951, <i>Simulium</i>	16.23
jundialensis d'Andretta & González B., 1964, <i>Simulium</i>	16.14
<i>kamloopsi</i> Hearle, 1932, <i>Simulium</i>	16.7
kompi Dalmat, 1951, <i>Simulium</i>	16.14
kuscheli Wygodzinsky, 1952, <i>Gigantodax</i>	16.3
lahillei (Paterson & Shannon), 1927 (<i>Eusimulium</i>), <i>Simulium</i>	16.14
<i>Lanea</i> Vargas, Martínez Palacios & Díaz Nájera, 1946	16.5
<i>lane-porteroi</i> Smart, 1945, <i>Simulium</i> (error)	16.14
laneportoi Vargas, 1941, <i>Simulium</i>	16.14
larvispinosum León, 1948, <i>Simulium</i>	16.15
lassmanni Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.15
laticalx (Enderlein), 1933 (<i>Trichodagmia</i>), <i>Simulium</i>	16.15
latidigitus (Enderlein), 1936 (<i>Trichodagmia</i>), <i>Simulium</i>	16.15
<i>latitarsis</i> (Enderlein), 1933 (<i>Trichodagmia</i>), <i>Simulium</i>	16.9
limay Wygodzinsky, 1958, <i>Simulium</i>	16.15
<i>limbatum</i> (Enderlein), 1933 (<i>Ectemnaspis</i>), <i>Simulium</i>	16.20
<i>limbatum</i> Knab, 1915, <i>Simulium</i>	16.15
longithallum Díaz Nájera & Vulcano, 1961, <i>Simulium</i>	16.15
<i>lugubre</i> Lutz & Nuñez Tovar, 1928, <i>Simulium</i>	16.16
lurybayae Smart, 1944, <i>Simulium</i>	16.15
lutzi Knab, 1913, <i>Simulium</i>	16.6
<i>lutzi</i> Malloch, 1914, <i>Simulium</i>	16.6
<i>lutzi</i> ana (Enderlein), 1933 (<i>Trichodagmia</i>), <i>Simulium</i>	16.18
<i>lutzi</i> anum Enderlein of Rubzov, 1940, <i>Simulium</i>	16.18
<i>lutzi</i> anum Pinto, 1931, <i>Simulium</i>	16.17
<i>Lutzsimulium</i> d'Andretta & d'Andretta, 1947	16.4
<i>macca</i> (Enderlein), 1933 (<i>Dasypelmoza</i>), <i>Simulium</i>	16.15
<i>macca</i> (Enderlein), 1933 (<i>Ectemnaspis</i>), <i>Simulium</i>	16.7
<i>maccum</i> , Vargas, 1943, <i>Simulium</i> (error)	16.15
major Lane & Porto, 1940, <i>Simulium</i>	16.15
<i>magnum</i> Porto, 1940, <i>Simulium</i> (error)	16.15
<i>mallochi</i> (Enderlein), 1925 (<i>Wilhelmia</i>), <i>Simulium</i>	16.19
mangabeirai Vargas, 1945, <i>Simulium</i>	16.15
manicatum (Enderlein), 1933 (<i>Trichodagmia</i>), <i>Simulium</i>	16.15
marathrumi Fairchild, 1940, <i>Simulium</i>	16.15
marginale (Edwards), 1931 (<i>Simulium</i>), <i>Gigantodax</i>	16.3
maroniense Floch & Abonnenc, 1946, <i>Simulium</i>	16.15
marquesi Vargas & Díaz Nájera, 1957, <i>Simulium</i>	16.15
<i>martinezi</i> Vargas, 1943, <i>Simulium</i>	16.10
<i>mathesoni</i> Vargas, 1943, <i>Simulium</i>	16.18
matteabranchia Anduze, 1947, <i>Simulium</i>	16.15
mediovittatum Knab, 1915, <i>Simulium</i>	16.16
<i>Melusina</i> , authors	16.4
menchacai Vargas & Díaz Nájera, 1957, <i>Simulium</i>	16.16
<i>mendozanum</i> (Enderlein), 1936 (<i>Psaroniocompsa</i>), <i>Simulium</i>	16.14



meridionale Riley, 1887, <i>Simulium</i>	16.16
metallicum Bellardi, 1859, <i>Simulium</i>	16.16
<i>metallicum</i> Bellardi of Malloch, 1914, <i>Simulium</i>	16.19
<i>mexicana</i> (Enderlein), 1934 (<i>Hemicnetha</i>), <i>Simulium</i>	16.18
mexicanum Bellardi, 1862, <i>Simulium</i>	16.16
microbranchium Dalmat, 1949, <i>Simulium</i>	16.16
<i>miniata</i> (Enderlein), 1933 (<i>Dasypelmoza</i>), <i>Simulium</i>	16.10
<i>miniata</i> (Enderlein), 1934 (<i>Trichodagma</i>), <i>Simulium</i>	16.10
<i>minuscolum</i> Lutz, 1910, <i>Simulium</i>	16.6
<i>minutum</i> Surcouf & González Rincones, 1911, <i>Simulium</i>	16.6
mixensis Díaz Nájera, 1962, <i>Cnephia</i>	16.2
<i>mollis</i> Vargas, 1943, <i>Simulium</i>	16.7
montana (Philippi), 1865 (<i>Simulium</i>), <i>Cnephia</i>	16.2
montanum Philippi of Lutz, 1909, <i>Lutzsimulium</i>	16.4
<i>moorei</i> (Silva Figueroa), 1917 (<i>Simulia</i>), <i>Austrosimulium</i>	16.1
<i>mooseri</i> (Dampf), 1927 (<i>Eusimulium</i>), <i>Simulium</i>	16.8
nemorale Edwards, 1931, <i>Simulium</i>	16.17
<i>Neosimulium</i> Rubzov, 1940 (nomen nudum)	16.5
<i>Neosimulium</i> Vargas, Martínez Palacios & Díaz Nájera, 1946 ...	16.5
nigrescens (Edwards), 1931 (<i>Simulium</i>), <i>Gigantodax</i>	16.3
nigricornis Dalmat, 1950, <i>Simulium</i>	16.17
nigrifemur (Enderlein), 1936 (<i>Acropogon</i>), <i>Simulium</i>	16.17
nigrimanum Macquart, 1838, <i>Simulium</i>	16.17
<i>nigrimanum</i> Macquart of Lutz, 1909, <i>Simulium</i>	16.17
nigristrigatum (Enderlein), 1930 (<i>Pternaspatha</i>), <i>Simulium</i>	16.17
<i>nitidum</i> Malloch, 1912, <i>Simulium</i>	16.6
nogueirai d'Andretta & González B., 1964, <i>Simulium</i>	16.17
<i>Notolepria</i> Enderlein, 1930	16.5
<i>obesum</i> Lutz, 1922, <i>Simulium</i> (nomen nudum)	16.17
<i>obesum</i> Vulcano, 1959, <i>Simulium</i>	16.17
<i>occidentale</i> Townsend, 1891, <i>Simulium</i>	16.16
ochoai Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.17
ochraceum Walker, 1861, <i>Simulium</i>	16.17
<i>ochraceum</i> Walker of Hoffmann, 1931, <i>Simulium</i>	16.10
<i>ochraceum</i> Walker of Lane & Porto, 1939, <i>Simulium</i>	16.10
<i>ochraceum</i> Walker of Malloch, 1914, <i>Simulium</i>	16.8
<i>ochraceum</i> Walker of Malloch, 1914, <i>Simulium</i>	16.13
<i>opaliniifrons</i> (Enderlein), 1934 (<i>Psaroniocompsa</i>), <i>Simulium</i> ...	16.17
orbitale Lutz, 1910, <i>Simulium</i>	16.18
orbitale Lutz, 1910, <i>Simulium</i> (part)	16.18
orbitale Lutz of Pinto, 1931, <i>Simulium</i>	16.19
oyapockense Floch & Abonnenc, 1946, <i>Simulium</i>	16.17
pachecolunai (León), 1945 (<i>Simulium</i>), <i>Cnephia</i>	16.2
panamense Fairchild, 1940, <i>Simulium</i>	16.18
paraguayense Schrottky, 1909, <i>Simulium</i>	16.18
<i>paraguayense</i> Schrottky of Pinto, 1931, <i>Simulium</i>	16.18
paranense Schrottky, 1909, <i>Simulium</i>	16.1
Paraustrosimulium Wygodzinsky & Coscarón, 1962	16.18
parrai Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.18
paynei Vargas, 1942, <i>Simulium</i>	16.18
	16.41

<i>pennipunctum</i> Enderlein, 1933, <i>Gigantodax</i>	16.3
<i>perflavum</i> (Roubaud), 1906 (<i>Eusimulium</i>), <i>Simulium</i>	16.18
<i>pernigrum</i> (Lutz), 1910 (<i>Simulium</i>), <i>Lutzsimulium</i>	16.4
<i>pertinax</i> Kollar, 1832, <i>Simulium</i>	16.18
<i>pertinax</i> Kollar of Lane & Porto, 1939, <i>Simulium</i>	16.10
<i>philippianum</i> Pinto, 1931, <i>Simulium</i>	16.18
<i>pilosum</i> Lane & Porto, 1940, <i>Simulium</i>	16.15
<i>pintoi</i> d'Andretta & d'Andretta, 1946, <i>Simulium</i>	16.18
<i>placidum</i> Knab, 1915, <i>Simulium</i>	16.16
<i>Pliodasina</i> Enderlein, 1936	16.5
<i>prodexargenteum</i> (Enderlein), 1936 (<i>Pternaspatha</i>), <i>Simulium</i> ...	16.19
<i>pruinoseum</i> Lutz 1910, <i>Simulium</i>	16.19
<i>Psaroniocompsa</i> Enderlein, 1934	16.5
<i>pseudocallidum</i> Diaz Nájera, 1965, <i>Simulium</i>	16.19
<i>pseudohaematopotum</i> Hoffmann, 1930, <i>Simulium</i>	16.13
<i>Psilopelmia</i> Enderlein, 1933	16.5
<i>Pternaspatha</i> Enderlein, 1930	16.5
<i>pugetense</i> (Dyar & Shannon), 1927 (<i>Eusimulium</i>), <i>Simulium</i> ...	16.19
<i>puigi</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i> .	16.19
<i>pulchrum</i> Philippi, 1865, <i>Simulium</i>	16.19
<i>pulchrum</i> Philippi of Johannsen, 1903, <i>Simulium</i>	16.21
<i>pulverulentum</i> Knab, 1914, <i>Simulium</i>	16.19
<i>pulverulentum</i> Knab of Vargas, 1943, <i>Simulium</i>	16.12
<i>punctativentris</i> (Enderlein), 1936 (<i>Acropogon</i>), <i>Simulium</i>	16.19
<i>quadrifidum</i> Lutz, 1917, <i>Simulium</i>	16.19
<i>quadristrigatum</i> Enderlein, 1933, <i>Simulium</i>	16.19
<i>quadrivittatum</i> Loew, 1862, <i>Simulium</i>	16.19
<i>quadrivittatum</i> Fairchild of Dunn, 1934, <i>Simulium</i>	16.13
<i>quadrivittatum</i> Loew of Malloch, 1914, <i>Simulium</i>	16.19
<i>riveti</i> Roubaud, 1906, <i>Simulium</i>	16.19
<i>roblesi</i> (León), 1943 (<i>Simulium</i>), <i>Cnephia</i>	16.2
<i>romanai</i> Wygodzinsky, 1951, <i>Simulium</i>	16.20
<i>rorotaense</i> Floch & Abonnenc, 1946, <i>Simulium</i>	16.20
<i>rubicundulum</i> Knab, 1914, <i>Simulium</i>	16.23
<i>rubiginosum</i> (Enderlein), 1933 (<i>Chirostilbia</i>), <i>Simulium</i>	16.13
<i>rubrithorax</i> Lutz, 1909, <i>Simulium</i>	16.20
<i>rubrithorax</i> Lutz of Lutz & Nuñez Tovar, <i>Simulium</i>	16.17
<i>rufescens</i> (Edwards), 1931 (<i>Simulium</i>), <i>Gigantodax</i>	16.3
<i>rufidorsum</i> (Enderlein), 1936 (<i>Acropogon</i>), <i>Simulium</i>	16.9
<i>rufidorsum</i> (Enderlein), 1933 (<i>Psilopelmia</i>), <i>Simulium</i>	16.11
<i>rufinotum</i> (Edwards), 1931 (<i>Simulium</i>), <i>Gigantodax</i>	16.3
<i>ruizi</i> Vargas & Díaz Nájera, 1948, <i>Simulium</i>	16.20
<i>samboni</i> Jennings, 1915, <i>Simulium</i>	16.20
<i>samboni</i> Jennings of Dampf, 1943, <i>Simulium</i>	16.10
<i>sanguineum</i> Knab, 1915, <i>Simulium</i>	16.20
<i>sayi</i> Dyar & Shannon of Vargas, Martínez Palacios & Díaz Nájera,	
1943, <i>Simulium</i>	16.13
<i>schoenemanni</i> (Enderlein), 1933 (<i>Acropogon</i>), <i>Simulium</i>	16.20

scutclatum (Lane & Porto), 1940 (Eusimulium), Simulium	16.20
scutistriatum Lutz, 1909, Simulium	16.20
seriatum Knab, 1914, Simulium	16.20
shannoni (Edwards), 1931 (Simulium), Gigantodax	16.4
sicuanii Smart, 1944, Simulium	16.20
<i>simile</i> Silva Figucroa, 1917, Simulium	16.11
simplicicolor Lutz, 1910, Simulium	16.20
<i>Simulia</i> , emend.	16.4
Simulium Latreille, 1802	16.4
smartii Vargas, 1946, Simulium	16.20
solarii Stone, 1948, Simulium	16.21
spadicidorsum (Enderlein), 1934 (Trichodagnia), Simulium	16.21
spinibranchium Lutz, 1910, Simulium	16.21
spinifer Knab, 1914, Simulium	16.21
strigatum (Enderlein), 1933 (Trichodagnia), Simulium	16.21
strigidorsum (Enderlein), 1933 (Acropogon), Simulium	16.21
striginotum (Enderlein), 1933 (Thyrsopelema), Simulium	16.21
subclavibranchium Lutz, 1910, Simulium	16.21
subnigrum Lutz, 1910, Simulium	16.21
subpallidum Lutz, 1910, Simulium	16.21
<i>supallidum</i> Lutz, 1910, Simulium (part)	16.21
<i>subviride</i> Lutz & Machado, 1915, Simulium (nomen nudum)	16.23
<i>tamaulipense</i> Townscoed, 1897, Simulium	16.16
tarsale Williston, 1896, Simulium	16.21
<i>tarsale</i> Williston of Floch & Abonnenc, 1946, Simulium	16.12
tarsatum Macquart, 1846, Simulium	16.21
<i>tarsatum</i> Philippi, 1865 Simulium	16.18
temascalense Díaz Nájera & Vulcano, 1962, Simulium	16.21
tenuifrons (Enderlein), 1936 (Odagnia), Simulium	16.10
tenuipes Knab, 1914, Simulium	16.21
<i>tephrodes</i> Speiser, 1904, Simulium	16.23
<i>Thyrsopelema</i> Enderlein, 1933	16.5
townsendi Malloch, 1912, Simulium	16.21
travassosi d'Andretta & d'Andretta, 1947, Simulium	16.22
<i>Trichodagnia</i> Enderlein, 1933	16.5
tricornis León, 1945, Simulium	16.22
trivittatum Malloch, 1914, Simulium	16.22
tuberosum (Lundström), 1911 (Melusina), Simulium	16.22
<i>turgidum</i> (Hoffmann), 1930 (Eusimulium), Simulium	16.16
urubambanum Enderlein, 1933, Simulium	16.22
<i>vandalicum</i> Dyar & Shannon of Vargas, 1945, Simulium	16.15
vargasii León, 1945, Gigantodax	16.4
varians Lutz, 1900, Simulium	16.22
varipes Philippi, 1865, Simulium	16.22
venustum Say of Lutz, 1909, Simulium	16.18
veracruzianum Vargas, Martínez Palacios & Díaz Nájera, 1946, Simulium	16.22
versicolor Lutz & Nuñez Tovar, 1928, Simulium	16.22
	16.43

<i>violascens</i> (Enderlein), 1933 (Odagnia), <i>Simulium</i>	16.22
<i>virgatum</i> Coquillett, 1902, <i>Simulium</i>	16.22
<i>virgatum</i> Coquillett of Malloch, 1914, <i>Simulium</i>	16.8
<i>virgatum</i> Coquillett of Stains & Knowlton, 1943, <i>Simulium</i>	16.21
<i>virgatum</i> Coquillett of Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.21
<i>vittatum</i> (Zetterstedt), 1838 (<i>Simulia</i>), <i>Simulium</i>	16.23
<i>walterwittmeri</i> Wygodzinsky, 1958, <i>Simulium</i>	16.23
<i>Wilhelmia</i> Enderlein, 1921	16.4
<i>wittmeri</i> Wygodzinsky, 1951, <i>Gigantodax</i>	16.4
<i>wolcottii</i> Fox, 1953, <i>Simulium</i>	16.23
<i>wolffhuegeli</i> (Enderlein), 1922 (<i>Wilhelmia</i>), <i>Simulium</i>	16.23
<i>wrighti</i> (Vargas, Martínez Palacios & Díaz Nájera), 1944 (<i>Simu-</i> <i>lium</i>), <i>Gigantodax</i>	16.4
<i>wuyaraka</i> Ortiz C., 1957, <i>Simulium</i>	16.23
<i>yepocapense</i> Dalmat, 1949, <i>Simulium</i>	16.23
<i>zempoalense</i> Vargas, Martínez Palacios & Díaz Nájera, 1946, <i>Simulium</i>	16.23



17

SUPERFAMILY ANISOPODOIDEA

Family Anisopodidae





DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

17

FAMILY ANISOPODIDAE

(RHYPHIDAE, PHRYNEIDAE, SYLVICOLIDAE)

NELSON PAPAVERO

*Departamento de Zoologia, Secretaria da Agricultura
São Paulo, Brasil*

The Anisopodidae are relatively well represented in the Neotropical region. Adults occur in small swarms or on tree trunks. Immature stages have been studied by Müller (1922) and are found in decaying organic matter. Edwards (1928) monographed the family and provided some morphological elements for the study of the male genitalia (1930). Lane & d'Andretta (1958) revised the genera *Sylvicola* (as *Anisopus*) and *Olbogaster*.

Subfamily ANISOPODINAE

Genus SYLVICOLA Harris

SYLVICOLA Harris, 1776:100. Type-species, *brevis* Harris (Coquillett, 1910:610) = *fenestralis* (Scopoli).

Phryne Meigen, 1800:16. Type-species, *Tipula fuscata* Fabricius (Coquillett, 1910:589). Suppressed by I.C.Z.N., 1963:339.

Anisopus Meigen, 1803:264. Type-species, *fuscus* Meigen (Coquillett, 1910:507) = *fuscata* (Fabricius).

Rhyphus Latreille, 1804:188. Type-species, *Tipula fenestralis* Scopoli (as *fenestrarum*; mon.).

ALBIAPEX (Lane & d'Andretta), 1958:499, figs. 6, 15 (*Anisopus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Território do Amapá, Rio de Janeiro, São Paulo). N. comb.

ALBICORNIS (Edwards), 1923:487, pl. 16, fig. 6 (*Anisopus*). Type-locality: Jamaica, Cinchona. Distr. — Jamaica. N. comb.

ANDINUS (Edwards), 1930:118, pl. 11, figs. 30-32 (*Anisopus*). Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Chile, Argentina. N. comb.

Published under the auspices of the "Fundação de Amparo à Pesquisa do Estado de São Paulo."

- ANNULIFERUS (Edwards), 1923:485, pl. 16, fig. 3 (*Anisopus*). Type-locality: Brazil, São Paulo. Distr. — Brazil (Minas Gerais and Rio de Janeiro, south to Santa Catarina). N. COMB.
- ANNULIPES (Edwards), 1919:152 (*Anisopus*). Type-locality: "Ecuador". Distr. — Ecuador. N. COMB.
- APICATUS (Edwards), 1919:151, fig. 8 (*Anisopus*). Type-locality: "Ecuador". Distr. — Ecuador. N. COMB.
- ARGENTINUS (Edwards), 1928:16 (*Anisopus*). Type-locality: Argentina, Salta, 2500 m. Distr. — Argentina. N. COMB.
- ATER (Edwards), 1919:152 (*Anisopus*). Type-locality: "Ecuador". Distr. — Ecuador. N. COMB.
- BORACEAE (Lane & d'Andretta), 1958:501, fig. 9 (*Anisopus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo). N. COMB.
- CAIUASI (Lane & d'Andretta), 1958:502, fig. 16 (*Anisopus*). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Minas Gerais, São Paulo). N. COMB.
- DOLOROSUS (Williston), 1896:298, pl. 10, fig. 74 (*Rhyphus*). Type-locality: St. Vincent, B.W.I. Distr. — St. Vincent (B.W.I.). N. COMB.
- FASCIATUS (Roeder), 1886:260 (*Rhyphus*). Type-locality: Colombia, Cordillera, Páramo, 3200 m. Distr. — Colombia. N. COMB.
- FLUMINENSIS (Lane & d'Andretta), 1958:504, figs. 4, 10 (*Anisopus*). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Distr. — Brazil (Rio de Janeiro). N. COMB.
- FUSCIPENNIS (Macquart), 1838:80, pl. 11, fig. 21 (*Rhyphus*). Type-locality: "Chile". Distr. — Chile, Juan Fernandez Is. (Masatierra, Masafuera), Argentina. Refs. — Edwards, 1930:117, pl. 11, figs. 27, 29; Hennig, 1955:21. N. COMB.
- GRANDIS (Lane & d'Andretta), 1958:505, figs. 8, 14 (*Anisopus*). Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil (Minas Gerais, Rio de Janeiro, São Paulo). N. COMB.
- GUTTATUS (Schiner), 1868:48 (*Rhyphus*). Type-locality: "Brazil". Distr. — Brazil (Mato Grosso to Rio de Janeiro, south to Santa Catarina). N. COMB.
- HYALINUS (Lane & d'Andretta), 1958:508, figs. 3, 13 (*Anisopus*). Type-locality: Puerto Rico, Yauco-Lares Road, km 29. Distr. — Puerto Rico. N. COMB.
- INCASICUS (Lane & d'Andretta), 1958:509, fig. 17 (*Anisopus*). Type-locality: Trinidad, B.W.I. Distr. — Trinidad, Puerto Rico. N. COMB.
- INFUMATUS (Knab), 1912:112 (*Anisopus*). Type-locality: "Trinidad". Distr. — Trinidad. N. COMB.
- LUTEATUS (Edwards), 1923:487 (*Anisopus*). Type-locality: British Guiana, Mt. Roraima, 8600 feet. Distr. — British Guiana. N. COMB.

- MAJOR (Edwards), 1930:118, pl. 11, figs. 33-34 (*Anisopus*). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Chile, Argentina. N. COMB.
- MARMORATUS (Edwards), 1923:484, pl. 16, fig. 2 (*Anisopus*). Type-locality: Colombia, Bogotá. Distr. — Colombia. N. COMB.
- ORNATUS (Edwards), 1923:483, pl. 16, fig. 1 (*Anisopus*). Type-locality: "Venezuela". Distr. — Venezuela. N. COMB.
- PAUPERATUS (Edwards), 1923:485, pl. 16, fig. 5 (*Anisopus*). Type-locality: "Venezuela". Distr. — Venezuela. N. COMB.
- PICTURATUS (Knab), 1912:111 (*Anisopus*). Type-locality: Costa Rica, Cartago, 1500 m (bred from epiphytic bromeliads). Distr. — Costa Rica. N. COMB.
- QUADRIVITTATUS (Edwards), 1928:18 (*Anisopus*). Type-locality: Costa Rica, Cartago, Turrialba, 1608 m. Distr. — Costa Rica. N. COMB.
- TIBIALIS (Edwards), 1923:486 (*Anisopus*). Type-locality: Mexico, "région montagnaise des environs de Toluca". Distr. — Mexico, Costa Rica. N. COMB.
- TUCUMANUS (Lane & d'Andretta), 1958:510, figs. 7, 11 (*Anisopus*). Type-locality: Argentina, Tucumán, San Miguel de Tucumán. Distr. — Argentina. N. COMB.
- VANIPES (Curran), 1934:309 (*Rhyphus*). Type-locality: British Guiana, Bartica, Kartabo. Distr. — British Guiana. N. COMB.
- WYGODZINSKYI (Lane & d'Andretta), 1958:512, figs. 2, 12 (*Anisopus*). Type-locality: Argentina, Tucumán, Cerro de San Javier, 1000 m. Distr. — Argentina. N. COMB.

Genus CARRERAIA Corrêa

- CARRERAIA Corrêa, 1947:97. Type-species, *Olbogaster edwardsi* Carrera (orig. des.).
- EDWARDSI (Carrera), 1941:196, figs. 1-4 (*Olbogaster*). Type-locality: Brazil, Paraná, Curitiba. Distr. — Brazil.

Genus LOBOGASTER Philippi

- LOBOGASTER Philippi, 1865:632. Type-species, *paradoxus* Philippi (mon.).
- PARADOXUS Philippi, 1865:632, pl. 24, fig. 16. Type-locality: "Chile". Distr. — Chile. Refs. — Edwards, 1923:488, figs. 2a-b (male genitalia); Reed, 1937:26, fig. 2 (redescriptions).
- philippii* Schiner, 1868:23. Type-locality: "Chile".

Genus OLBOGASTER Osten Sacken

- OLBOGASTER Osten Sacken, 1886:20. Type-species, *Rhyphus taeniatus* Bellardi (Coquillett, 1910:579).




- ALMEIDA Corrêa, 1946:142, figs. 1-6. Type-locality: Brazil, São Paulo, Itanhacm, Praia Grande, Fazenda Rondônia. Distr. — Brazil (Guanabara, São Paulo).
- ANTILLARUM Lane & d'Andretta, 1958:516, fig. 19. Type-locality: St. Croix I., Christiansted. Distr. — St. Croix I.
- CARIOCA Lane & d'Andretta, 1958:516, fig. 22. Type-locality: Brazil, Rio de Janeiro, Teresópolis, Parque Nacional da Serra dos Órgãos. Distr. — Brazil (Rio de Janeiro).
- CHAVANTESI Lane & d'Andretta, 1958:518, fig. 20. Type-locality: Brazil, Minas Gerais, Sapucaí-Mirim, Cidade Azul. Distr. — Brazil (Minas Gerais).
- CINCTA Kertész, 1902:4, figs. a-b. Type-locality: Peru, Callanga. Distr. — Peru (Callanga).
- COGNATA Osten Sacken, 1886:21. Type-locality: Costa Rica, Irazu. Distr. — Costa Rica, Panama (Canal Zone).
- FULVA Edwards, 1928:22. Type-locality: "Puerto Rico". Distr. — Puerto Rico.
- GUATEMALTECA Lane & d'Andretta, 1958:519. Type-locality: Guatemala, Cayuga. Distr. — Guatemala (Cayuga).
- MEXICANA Lane & d'Andretta, 1958:520. Type-locality: Mexico, Tuxtepec. Distr. — Mexico (Tuxtepec).
- PANAMENSIS Lane & d'Andretta, 1958:521, fig. 23. Type-locality: Panama, Canal Zone, Fort Kobbe. Distr. — Panama (Canal Zone).
- SACKENI Edwards, 1915:502. Type-locality: Mexico, Vera Cruz, Atoyac. Distr. — U.S.A. (Texas, Florida, South Carolina), Mexico.
sp. Williston, 1901:229, pl. 4, fig. 6. Locality: Mexico, Vera Cruz, Atoyac.
- SCALARIS (Wiedemann), 1830:618 (*Rhyphus*). Type-locality: Brazil, Bahia (as U.S.A., Georgia, error; cf. Edwards, 1928:23, footnote no. 1). Distr. — Brazil (Bahia, Mato Grosso, Santa Catarina), Paraguay. Ref. — Corrêa, 1946: figs. 7-8 (male genitalia).
quinquefasciata Enderlein, 1909:65 (*Anisopus*). Type-locality: Brazil, Santa Catarina.
- SIMILANS Lane & d'Andretta, 1958:523, fig. 26. Type-locality: Panama, Bocas del Toro, Almirante. Distr. — Costa Rica, Panama.
- SURINAMENSIS Lane & d'Andretta, 1958:524, fig. 28. Type-locality: Surinam, Moengo. Distr. — Surinam.
- TAENIATA (Bellardi), 1862:5, pl. 3, fig. 15 (1864:202) (*Rhyphus*). Type-locality: "Mexico". Distr. — U.S.A. (Texas, Florida), Mexico.
- TAMOIOI Lane & d'Andretta, 1958:524. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara).
- TRINIDADENSIS Edwards, 1928:23. Type-locality: Trinidad, B.W.I. Distr. — Trinidad.

Subfamily MYCETOBIIINAE

Genus MYCETOBIA Meigen

- MYCETOBIA Meigen, 1818:229. Type-species, *pallipes* Meigen (Westwood, 1840:127).



- ?FULVA Philippi, 1865:626. Type-locality: Chile, Corral. Distr. — Chile.
STONEI Lane & d'Andretta, 1958:527. Type-locality: Costa Rica, San Pedro de los Montes de Oca. Distr. — Costa Rica.

LITERATURE CITED

- BELLARDI, L.
1862. *Saggio di Ditterologia Messicana*, Append.: 28 pp., 1 pl. Torino.
1864. Idem. *Mem. R. Accad. delle Sci. Torino* 21:200-225, 1 pl.
- CARRERA, M.
1941. Notas sôbre o gênero *Olbiogaster* Osten Sacken, 1886, com a descrição de uma nova espécie (Fam. Anisopodidae, Dipt.). *Papéis Avulsos Dep. Zool. S. Paulo* 1:193-202, 4 figs.
- COQUILLET, D. W.
1910. The type-species of the North American genera of Diptera. *Proc. U. S. Nat. Mus.* 37:499-647.
- CORRÊA, R. R.
1946. Descrição de *Olbiogaster almeidai*, nova espécie de Anisopodidae do Brasil. *Livro Homenagem R. Ferreira D'Almeida*: 141-147, 8 figs.
1947. Sôbre alguns anisopódidas da América do Sul. Descrição de *Carreraia*, n. gen. *Papéis Avulsos Dep. Zool. S. Paulo* 8:97-107, 11 figs.
- CURRAN, C. H.
1934. The Diptera of Kartabo, Bartica District, British Guiana. *Bull. Amer. Mus. Nat. Hist.* 66:287-532, 54 figs.
- EDWARDS, F. W.
1915. Three new species of the dipterous genus *Olbiogaster*, O.-S., in the British Muscum collection. *Ann. & Mag. Nat. Hist.* (8) 16:502-505.
1919. Diptères. Nematocères. *Mission Arc Méridien Equat.* 10(2): 143-162, figs.
1923. Notes on the dipterous family Anisopodidae. *Ann. & Mag. Nat. Hist.* (9)12:475-493, 3 figs., pl. 16.
1928. Fam. Protorhyphidae, Anisopodidae, Pachyneuridae, Trichoce-ridae, in P. Wytzman: *Genera Insectorum* 190:1-26, 2 pls.
1930. Bibionidae, Scatopsidae, Cecidomyiidae, Culicidae, Thaumalcidae. Anisopodidae, in *Diptera of Patagonia and South Chile* 2(3): 114-119, pl. 11.
- ENDERLEIN, G.
1909. *Olbiogaster quinquefasciata*, eine neue Rhyphidae aus Süd-Brasilien. *Stett. ent. Zeit.* 71 (1910):65-66.



HARRIS, M.

1776. *An exposition of British Insects. Decad IV*: 100-138, pls. 31-40, London.

HENNIG, W.

1955. Los insectos de las Islas Juan Fernandez. 16. Phryncidae, Helomyzidae, Lonchaeidae, Piophilidae, Anthomyzidae und Muscidae. (Diptera). *Rev. Chil. Entom.* 4:21-34.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.* 20:339-342.

KERTÉSZ, K.

1902. Über die Familie Rhyphidcn. *Természetrzaji Füzetek* 25:4-6, figs. a-b.

KNAB, F.

1912. New species of Anisopodidae (Rhyphidae) from tropical America (Diptera: Nematocera). *Proc. Biol. Soc. Washington* 25:111-113.

LANE, J. & C. D'ANDRETTA JR.

1958. Neotropical Anisopodidae (Diptera, Nematocera). *Studia Entom.* 1:497-528, 29 figs.

LATREILLE, P. A.

1804. Tableau méthodique des Insectes, pp. 129-200, in Société de Naturalistes et d'Agriculteurs: *Nouveau dictionnaire d'histoire naturelle, appliqué aux arts, principalement à l'agriculture et à l'économie rurale et domestique* 24(3): Tableau méthodique d'histoire naturelle, 238 pp., 5 pls. Paris.

MACQUART, J.

1838. *Diptères exotiques nouveaux ou peu connus* 1(1):5-221, 25 pls., Paris.

MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp., Paris.

1803. Versuch einer neuen Gattungseinteilung der europäischen zweiflügelige Insekten. *Mag. f. Insektenkunde* 2:259-281.

1818. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 1:xxxvi+333 pp., pls. I-II, Aachen.

MÜLLER, M.

1922. *Rhyphus* und *Mycetobia* mit besonderer Berücksichtigung des larvalen Darmes. *Arch. f. Naturg.* 88A (2):1-44, 36 figs.

OSTEN SACKEN, C. R.

1886. *Biologia Centrali-Americana, Diptera 1*: viii+378 pp., 6 pls.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool. bot. Ges. Wien* 15:595-782, pls. 23-29.

REED, E. P.

1937. *Sobro Lobogaster paradoxus* Ph. Rev. Chil. Hist. Nat. 41:25-27, fig. 2.

ROEDER, V. V.

1886. Dipteren von den Cordilleren in Columbien, gesammelt durch Herrn Dr. Alphons Stübel. *Stett. ent. Zeit.* 47:257-270.

SCHINER, J. R.

1868. *Reise der österreichische Fregatte Novara um die Erde*, Zool. Theil, Diptera: 388 pp., 4 pls. K. K. Hof- und Staatsdruckerei, Wien.

WESTWOOD, J. O.

1840. *An introduction to the modern classification of insects. Synopsis of the genera of British Insects*, 158 pp., London.

WIEDEMANN, C. R. W.

1830. *Aussereuropäische zweiflügelige Insekten* 2:684 pp., 3 pls.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. ent. Soc. London* (1896):253-446, pls. 8-14.

1901. *Biologia Centrali-Americana, Diptera 1*:229.

INDEX

albiapex (Lane & d'Andretta), 1958 (Anisopus), Sylvicola	17.1
albicornis (Edwards), 1923 (Anisopus), Sylvicola	17.1
almeidai Corrêa, 1946, Olbiogaster	17.4
andinus (Edwards), 1930 (Anisopus), Sylvicola	17.1
Anisopodinae	17.1
<i>Anisopus</i> Meigen, 1803	17.1
annuliferus (Edwards), 1923 (Anisopus), Sylvicola	17.2
annulipes (Edwards), 1919 (Anisopus), Sylvicola	17.2
antillarum Lane & d'Andretta, 1958, Olbiogaster	17.4
apicatus (Edwards), 1919 (Anisopus), Sylvicola	17.2
argentinus (Edwards), 1928 (Anisopus), Sylvicola	17.2
ater (Edwards), 1919 (Anisopus), Sylvicola	17.2
boraceae (Lane & d'Andretta), 1958 (Anisopus), Sylvicola	17.2
caiuasi (Lanc & d'Andretta), 1958 (Anisopus), Sylvicola	17.2
carioca Lanc & d'Andretta, 1958, Olbiogaster	17.4



<i>Carreraia</i> Corrêa, 1947	17.3
<i>chavantesi</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>cincta</i> Kertész, 1902, <i>Olbiogaster</i>	17.4
<i>cognata</i> Osten Sacken, 1886, <i>Olbiogaster</i>	17.4
<i>dolorosus</i> (Williston), 1896 (<i>Rhyphus</i>), <i>Sylvicola</i>	17.2
<i>edwardsi</i> (Carrera), 1941 (<i>Olbiogaster</i>), <i>Carreraia</i>	17.3
<i>fasciatus</i> (Roeder), 1886 (<i>Rhyphus</i>), <i>Sylvicola</i>	17.2
<i>fluminensis</i> (Lane & d'Andretta), 1958 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>fulva</i> Edwards, 1928, <i>Olbiogaster</i>	17.4
? <i>fulva</i> Philippi, 1865, <i>Mycetobia</i>	17.5
<i>fuscipennis</i> (Macquart), 1838 (<i>Rhyphus</i>), <i>Sylvicola</i>	17.2
<i>grandis</i> (Lanc & d'Andretta), 1958 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>guatmalteca</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>guttatus</i> (Schiner), 1868 (<i>Rhyphus</i>), <i>Sylvicola</i>	17.2
<i>hyalinus</i> (Lane & d'Andretta), 1958 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>incasicus</i> (Lane & d'Andretta), 1958 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>infumatus</i> (Knab), 1912 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>Lobogaster</i> Philippi, 1865	17.3
<i>luteatus</i> (Edwards), 1923 (<i>Anisopus</i>), <i>Sylvicola</i>	17.2
<i>major</i> (Edwards), 1930 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>marmoratus</i> (Edwards), 1923 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>mexicana</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>Mycetobia</i> Meigen, 1818	17.4
<i>Mycetobiinae</i>	17.4
<i>Olbiogaster</i> Osten Sacken, 1886	17.3
<i>ornatus</i> (Edwards), 1923 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>panamensis</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>paradoxus</i> Philippi, 1865, <i>Lobogaster</i>	17.3
<i>pauperatus</i> (Edwards), 1923 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>philippii</i> Schiner, 1868, <i>Lobogaster</i>	17.3
<i>Phryne</i> Meigen, 1800	17.1
<i>picturatus</i> (Knab), 1912 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>quadrivittatus</i> (Edwards), 1928 (<i>Anisopus</i>), <i>Sylvicola</i>	17.3
<i>quinquefasciata</i> (Enderlein), 1909 (<i>Anisopus</i>), <i>Olbiogaster</i>	17.4
<i>Rhyphus</i> Latrille, 1804	17.1
<i>sackeni</i> Edwards, 1915, <i>Olbiogaster</i>	17.4
<i>scalaris</i> (Wiedemann), 1830 (<i>Rhyphus</i>), <i>Olbiogaster</i>	17.4
<i>similans</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>sp.</i> Williston, 1901, <i>Olbiogaster</i>	17.4
<i>stonci</i> Lane & d'Andretta, 1958, <i>Mycetobia</i>	17.5
<i>surinamensis</i> Lane & d'Andretta, 1958, <i>Olbiogaster</i>	17.4
<i>Sylvicola</i> Harris, 1776	17.1



taeniata (Bellardi), 1862 (Rhyphus), Olbiogaster	17.4
tamoioi Lane & d'Andretta, 1958, Olbiogaster	17.4
tibialis (Edwards), 1923 (Anisopus), Sylvicola	17.3
trinidadensis Edwards, 1923, Olbiogaster	17.4
tucumanus (Lane & d'Andretta), 1958 (Anisopus), Sylvicola	17.3
varipes (Curran), 1934 (Rhyphus), Sylvicola	17.3
wygodzinskyi (Lane & d'Andretta), 1958 (Anisopus), Sylvicola ..	17.3





18

SUPERFAMILY BIBIONOIDEA

Family Bibionidae

MUSEU DE ZOOLOGIA - USP
BIBLIOTECA
RECEBIDO em





DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

18

FAMILY BIBIONIDAE *

D. ELMO HARDY

*Senior Professor and Chairman
Department of Entomology
University of Hawaii
Honolulu, Hawaii*

Approximately 650 species of Bibionidae are known over the world. 169 species and 4 subspecies are now known from the Neotropical region. The genus *Dilophus* is still rather poorly known and the total number of species for the family in the region will surely be over 250. The Neotropical species belong in three subfamilies and represent eight genera.

These flies are herbivores and feed predominantly as scavengers on decaying plant material. No information is available concerning the habits or biologies of Neotropical species. The adults of *Dilophus* are often collected in flowers where they are apparently nectar feeders. *Bibio* and *Plecia* are more commonly taken hovering, during their mating swarms.

The Neotropical *Plecia* and *Penthetria* were revised by Hardy (1945).

Subfamily HESPERININAE

Genus HESPERINUS Walker

HESPERINUS Walker, 1848:81. Type-species, *brevifrons* Walker (mon.).

Spodius Loew, 1858:101. Type-species, *imbecillus* Loew (mon.).

CONJUNGENS Schincr, 1868:23. Type-locality: "Brazil". Distr. — Brazil.


Subfamily PLECIINAE

Genus PENTHETRIA Meigen

PENTHETRIA Meigen, 1803:264. Type-species, *funebis* Meigen (sub. mon., Meigen, 1804:104).

Eupeitenus Macquart, 1838a:88 (1838:84). Type-species, *Penthetria atra* Macquart (mon.) = *heteroptera* (Say).

* Published with the approval of the Director of the Hawaii Agricultural Experiment Station as Miscellaneous Paper No. 185.



- APPENDICULA Hardy, 1945:384, pl. 29, figs. 122a-c. Type-locality: Costa Rica, Volcán Poas. Distr. — Costa Rica.
- DISTINCTA Hardy, 1945:385, pl. 29, figs. 123a-b. Type-locality: Guatemala, Antigua. Distr. — Guatemala.
- HEROS Say, 1829:154. Type-locality: "Mexico". Distr. — Mexico. Ref. — Hardy, 1959:441.
- MEXICANA (Hardy), 1937:199 (*Plecia*). Type-locality: Mexico, Guadalajara. Distr. — Mexico. Ref. — Hardy, 1945:387, pl. 30, figs. 125a-b.
- NIGERRIMA (Bellardi), 1859:14 (*Plecia*). Type-locality: Mexico, Orizaba. Distr. — Mexico, Ecuador, Colombia. Ref. — Hardy, 1945:388, pl. 30, figs. 126a-b.
- NIGRITA Perty, 1830:180, pl. 36, fig. 1. Type-locality: Brazil, Piauí. Distr. — Venezuela, Colombia, Ecuador, Peru, Brazil.


Genus PLECIA Wiedemann

- PLECIA Wiedemann, 1828:72. Type-species, *Hirtea fulvicollis* Fabricius (Blanchard, 1840:576).
- Rhinoplecia* Bellardi, 1859:16. Type-species, *rostrata* Bellardi (mon.).
- Penthera* Philippi, 1865:639. Type-species, *nigra* Philippi (mon.).
- ALACRIS Curran, 1934:310. Type-locality: British Guiana, Bartica, Kartabo. Distr. — British Guiana, Paraguay. Ref. — Hardy, 1945:395, pl. 30, figs. 128a-d.
- AMERICANA Hardy, 1940:15, figs. 1a-b. Type-locality: Southern U.S.A., Florida, Pensacola. Distr. — Southern U.S.A., Mexico. Ref. — Hardy, 1945:395, pl. 30, figs. 129a-d.
- AVICEPHALIFORMA Hardy, 1940:16, figs. 2a-d. Type-locality: Brazil, Minas Gerais, Diamantina. Distr. — Brazil. Ref. — Hardy, 1945:396, pl. 30, figs. 130a-d.
- BIARMATA Hardy, 1942b:105. Type-locality: Costa Rica, Crontena. Distr. — Costa Rica. Ref. — Hardy, 1945:397, pl. 30, figs. 131a-b.
- BICOLOR Bellardi, 1859:16. Type-locality: Mexico, Cordova. Distr. — Mexico, Panama (Canal Zone). Ref. — Hardy, 1945:398, pl. 30, figs. 132a-c.
- BIFORMIS Hardy, 1942b:106. Type-locality: "Trinidad". Distr. — Trinidad. Ref. — Hardy, 1945:399, pl. 31, figs. 133a-d.
- BRAZILIANA Hardy, 1942b:106. Type-locality: Brazil, Mato Grosso. Distr. — Brazil. Ref. — Hardy, 1945:400, pl. 31, figs. 134a-c.
- COLLARIS (Fabricius), 1805:54 (*Hirtea*). Type-locality: "America meridionalis". Distr. — Widespread over Neotropical region. Refs. — Hardy, 1945:401, pl. 31, figs. 135a-e; 1953:351, fig. 2a.
- thoracica* Fabricius, 1805:163 (*Laphria*). Type-locality: "America meridionalis".
- CONFUSA Loew, 1848:109. Neotype-locality: Brazil, Bahia. Distr. — Widespread over Neotropical region. Refs. — Edwards, 1931a:75; Hardy, 1945:402, pl. 31, figs. 136a-d. (Change of name for *ruficollis* Fabricius).
- ruficollis* Fabricius, 1805:53 (*Hirtea*) [nec *ruficollis* Fabricius, 1781:410 (*Tipula*)]. Type-locality: "America meridionalis".

- COSTALIS* Wiedemann, 1830:618. Type-locality: Brazil, São Paulo (?), Caçapava ["Cassapawa im inneren Brasilien"]. Distr. — Brazil. Ref. — Hardy, 1945:404.
- CURVISTYLATA* Hardy, 1942b:107. Type-locality: Mexico, Cuernavaca. Distr. — Mexico. Ref. — Hardy, 1945:405, pl. 31, figs. 137a-d.
- DENTATA* Hardy, 1942b:107. Type-locality: Nicaragua, Chontales. Distr. — Nicaragua. Ref. — Hardy, 1945:405, pl. 31, figs. 138a-d.
- DISCOLOR* Wulp, 1881:143. Type-locality: "Argentina". Distr. — Argentina. Ref. — Hardy, 1945:406.
- DISPARIS* Hardy, 1942b:108. Type-locality: Colombia, Upper Putumayo River. Distr. — Colombia. Ref. — Hardy, 1945:406, pl. 31, figs. 139a-b.
- ECUADORENSIS* *ECUADORENSIS* Hardy, 1942b:108. Type-locality: "Ecuador". Distr. — Ecuador, Colombia. Ref. — Hardy, 1945:406-407, pl. 32, figs. 140a-c.
- ssp. *MICANS* Hardy, 1942b:108. Type-locality: Ecuador, Guayaquil. Ref. — Hardy, 1945:407, pl. 32, figs. 141a-b.
- EDWARDSI* Hardy, 1940:17, figs. 3a-b. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Jussara. Distr. — Brazil. Ref. — Hardy, 1945:408, pl. 32, figs. 142a-b.
- FEMORATA* Macquart, 1838a:86. Type-locality: "Brazil". Distr. — Brazil.
- FUNEBRIS* (Fabricius), 1805:54 (*Hirtea*). Type-locality: "America meridional". Distr. — Supposedly widespread over the Neotropical region. Ref. — Hardy, 1959:446.
- GIBBOSA* Hardy, 1942b:109. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay, Argentina. Ref. — Hardy, 1945:409, pl. 32, figs. 143a-e.
- GRISEA* Edwards, 1938:322. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Argentina. Refs. — Hardy, 1945:410, pl. 32, figs. 144a-b; 1953:353, fig. 3a.
- IMPENSA* Hardy, 1957:238, figs. 1-2. Type-locality: "Mexico". Distr. — Mexico. Ref. — Hardy, 1961:86, figs. 3a-c.
- IMPERIALIS* Schiner, 1868:22. Type-locality: "Colombia". Distr. — Colombia. Refs. — Hardy, 1945:411, figs. 145a-b; 1956:86. (First available name for *costalis* Walker).
- costalis* Walker, 1858:422. (nec Wiedemann, 1830:618).
- IMPILOSA* Hardy, 1940:17, figs. 4a-b. Type-locality: Brazil, Mato Grosso. Distr. — Brazil. Ref. — Hardy, 1945:412, pl. 33, figs. 146a-b.
- INCURVATA* Hardy, 1942b:110. Type-locality: Guadeloupe (W.I.). Distr. — West Indies. Ref. — Hardy, 1945:413, pl. 33, figs. 147a-c.
- LATERALIS* Hardy, 1940:18, figs. 5a-c. Type-locality: Mexico, Hidalgo, San Miguel. Distr. — Mexico. Ref. — Hardy, 1945:413, pl. 33, figs. 148a-c.
- LINDNERI* Edwards, 1931a:75. Type-locality: Bolivia, Santa Cruz de la Sierra, El Cairo. Distr. — Bolivia, Peru, Brazil. Refs. — Hardy, 1945:414, pl. 33, figs. 149a-c; 1961:90.
- LOPESI* Hardy, 1940:19, figs. 6a-c. Type-locality: Brazil, São Paulo. Distr. — Brazil. Ref. — Hardy, 1945:415, pl. 33, figs. 150a-c.

- MACULATA Hardy, 1942b:110. Type-locality: Trinidad, Mayaro, East Coast-Distr. — West Indies. Ref. — Hardy, 1945:416, pl. 33, figs. 151a-b.
- MARGINATA Edwards, 1920:148. Type-locality: Ecuador, Tulcan. Distr. — Ecuador. Ref. — Hardy, 1945:417.
- MAURA Walker, 1837:336. Type-locality: Brazil, Minas Gerais. Distr. — Brazil, Argentina. Ref. — Hardy, 1945:417, pl. 33, figs. 152a-d.
velutina Macquart, 1846:21. Type-locality: Brazil, Minas Gerais.
- NEARCTICA Hardy, 1940:20, figs. 7a-c. Type-locality: U.S.A., Texas, Galveston. Distr. — Southern U.S.A., Mexico, Costa Rica, Guatemala, Honduras. Ref. — Hardy, 1945:418, pls. 33-34, figs. 153a-c.
- NIGRA NIGRA (Philippi), 1865:639 (*Penthera*). Type-locality: Chile, "prope Corral". Distr. — Chile. Ref. — Hardy, 1945:419, pl. 34, figs. 154a-c.
- ssp. ARGENTINA Hardy, 1953:353, fig. 4a. Type-locality: Argentina, El Infernillo. Distr. — Argentina.
- NITIDICOLLIS Edwards, 1931b:260. Type-locality: Brazil, Paraná, Castro-Distr. — Brazil. Ref. — Hardy, 1945:420, pl. 34, figs. 155a-c.
- NITIDIPES Edwards, 1920:147. Type-locality: Ecuador, Casitagua. Distr. — Ecuador. Ref. — Hardy, 1945:421, pl. 34, figs. 156a-d.
- PANAMAENSIS Hardy, 1942b:111. Type-locality: Panama, Canal Zone, Gatun Lake. Distr. — Panama. Ref. — Hardy, 1945:422, pl. 34, figs. 157a-d.
- PARACOLLARIS Hardy, 1961:88, figs. 4a-d. Type-locality: Colombia, Buenaventura. Distr. — Colombia.
- PARVISTYLATA Hardy, 1942b:112. Type-locality: Guatemala, Antigua. Distr. — Guatemala. Ref. — Hardy, 1945:423, pl. 34, figs. 158a-b.
- PERPLEXA Hardy, 1942b:112. Type-locality: "Cuba". Distr. — Cuba, Trinidad (W.I.), Argentina. Ref. — Hardy, 1945:423, pl. 34, figs. 159a-d.
- PERSIMILIS Hardy, 1940:21, figs. 8a-c. Type-locality: Brazil, São Paulo. Distr. — Brazil. Ref. — Hardy, 1945:424, pl. 35, figs. 160a-e.
- PERTINENS Hardy, 1942b:112. Type-locality: Mexico, Guerrero, Venta de Zopilote. Distr. — Mexico, Guatemala, Honduras. Ref. — Hardy, 1945:425, pl. 35, figs. 161a-g.
- PERUVIANA Hardy, 1961:89, figs. 5a-b. Type-locality: Peru, Chanchamayo-Distr. — Peru.
- PICTIPENNIS Edwards, 1931b:260. Type-locality: Brazil, Paraná, Castro-Distr. — Brazil. Ref. — Hardy, 1945:426, pl. 35, figs. 162a-c.
- var. *rufovittata* Edwards, 1938:323. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Ref. — Hardy, 1945:427.
- PLAGIATA Wiedemann, 1824:11. Type-locality: "Brazil". Distr. — Widespread throughout the Neotropical region. Refs. — Hardy, 1945:427, pl. 35, figs. 163a-d; 1953:355, fig. 5a.
- heteroptera* Macquart, 1846:21. Type-locality: Colombia, Santa Fe de Bogotá.
- vittata* Bellardi (nec Wiedemann), 1862:4.
- bellardii* Townsend, 1912:289.
- PLATYSTILA Hardy, 1957:239, figs. 3-4. Type-locality: "Venezuela". Distr. — Venezuela.

- PRUINOSA Hardy, 1940:22, figs. 9a-b. Type-locality: Cuba, Pico Turquino. Distr. — Cuba, Jamaica. Ref. — Hardy, 1945:428, pl. 35, figs. 164a-b.
- PUNCTULATA Hardy, 1940:23, figs. 10a-b. Type-locality: Brazil, Minas Gerais, Diamantina. Distr. — Brazil. Ref. — Hardy, 1945:429, pl. 35, figs. 165a-b.
- QUADRIVITTATA Williston, 1900:222. Type-locality: Mexico, Guerrero, Rincon. Distr. — Mexico. Ref. — Hardy, 1945:430, pl. 36, figs. 166a-c.
- QUASIMACULATA Hardy, 1961:90, figs. 6a-c. Type-locality: "Haiti". Distr. — Haiti.
- RECTIORA Hardy, 1942b:113. Type-locality: Brazil, Maranhão, São Luiz Ricamar. Distr. — Brazil, Paraguay. Ref. — Hardy, 1945:431, pl. 36, figs. 167a-c.
- ROSTELLATA Loew, 1858:109. Type-locality: "Brazil". Distr. — Widespread in the Neotropical region, apparently more common in Brazil and Argentina. Refs. — Hardy, 1945:432, pl. 36, figs. 168a-d; 1953:356.
- rostrata* Bellardi, 1859:15.
- RUFIMARGINATA Hardy, 1940:24, figs. 11a-b. Type-locality: Panama, Paraiso. Distr. — Panama. Ref. — Hardy, 1945:433, pl. 36, figs. 169a-b.
- RUFISUTELLA Hardy, 1940:24, figs. 12a-b. Type-locality: Peru, Moyobamba region. Distr. — Peru. Ref. — Hardy, 1945:433, pl. 36, figs. 170a-b.
- RUFITHORAX RUFITHORAX Walker, 1848:116. Type-locality: "Jamaica". Distr. — Jamaica, Mexico, Honduras, Guatemala. Ref. — Hardy, 1945:434, pl. 36, figs. 171a-b.
- ssp. CONCAVA* Hardy, 1942b:114. Type-locality: Guatemala, Alta V. Paz. Ref. — Hardy, 1945:435, pl. 37, figs. 172a-f.
- RUGOSA Hardy, 1940:25, figs. 13a-c. Type-locality: Colombia, Bet. Quercmal. Distr. — Colombia. Ref. — Hardy, 1945:436, pl. 37, figs. 173a-c.
- SEMINITENS Edwards, 1938:321. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Hardy, 1945:437, pl. 37, figs. 174a-c.
- SERRATA Hardy, 1942b:114. Type-locality: Paraguay, Villarica. Distr. — Paraguay. Ref. — Hardy, 1945:438, pl. 37, figs. 175a-c.
- SIMILIS Rondani, 1850:193. Type-locality: Brazil, São Sebastião (?). Distr. — Brazil, Argentina. Refs. — Hardy, 1945:439, pl. 37, figs. 176a-e; 1953:356, figs. 6a-d.
- minor* Jaennicke, 1867:318.
- TRILOBATA Hardy, 1942b:114. Type-locality: British Guiana, upper Courantyne River, King Frederick William IV Falls. Distr. — British Guiana. Ref. — Hardy, 1945:440, pl. 37, figs. 177a-d.
- UBERTA Hardy, 1942b:115. Type-locality: Honduras, Lombardia. Distr. — Mexico, Honduras. Ref. — Hardy, 1945:441, pl. 37, figs. 178a-e.
- VARABILIS Hardy, 1942b:115. Type-locality: Panama, Canal Zone, Barro Colorado Is. Distr. — Guatemala, Panama. Ref. — Hardy, 1945:441, pl. 38, figs. 179a-c.



- VITTATA Wiedemann, 1828:75. Type-locality: "Brazil". Distr. — Brazil.
Ref. — Hardy, 1945:442, pl. 38, figs. 180a-c.
- XENIA Hardy, 1942b:109. Type-locality: Costa Rica, Higuito, San Mateo.
Distr. — Costa Rica. Ref. — Hardy, 1945:443, pl. 38, figs. 181a-e.

Subfamily BIBIONINAE

Genus BIBIO Geoffroy

- BIBIO Geoffroy, 1762:568. Type-species, *Tipula hortulana* Linnaeus (Latreille, 1810:442). Validated by I.C.Z.N., 1957:86.
- Pullata* Harris, 1776:77. Type-species, *funesta* Harris = *pomonae* (Fabricius) (Coquillett, 1910:598).
- Hirtea* Fabricius, 1798:551 (nec Scopoli, 1763) (no type des.).
- ALBIPENNIS Say, 1823:78. Type-locality: U.S.A., Pennsylvania. Distr. — Recorded from northern Mexico. This species is common throughout the Nearctic region. Ref. — Hardy, 1945:451, pl. 38, fig. 183a.
- CRIORHINUS Bellardi, 1859:17. Type-locality: "Mexico". Distr. — Mexico. Ref. — Hardy, 1945:462.
- DISPAR Schiner, 1868:20. Type-locality: "Colombia". Distr. — Colombia, Argentina. Ref. — Hardy, 1953:347.
- FULIGINUS Bellardi, 1859:19. Type-locality: "Mexico". Distr. — Mexico. *fuliginosus* Aldrich, 1905:166. (Misspelling of *fuliginus*).
- INTERMEDIUS Rondani, 1850:193. Type-locality: "Venezuela". Distr. — Venezuela.
- NUDIOCULATUS Macquart, 1847:26. Type-locality: "Brazil". Distr. — Brazil. *nudioculatus* Kertész, 1902:143. (Misspelling of *nudioculatus*).
- PERUVIANUS Edwards, 1935:20. Type-locality: Peru, Oroya. Distr. — Peru.
- PICEUS Bellardi, 1859:17. Type-locality: "Mexico". Distr. — Mexico.
- SUBAEQUALIS Rondani, 1868:39. Type-locality: Argentina, Buenos Aires. Distr. — Brazil, Argentina. Ref. — Hardy, 1953:347.
- SUPERFLUUS Schiner, 1868:20. Type-locality: "Colombia". Distr. — Colombia.
- VICINUS Arribáizaga F.L., 1878:297. Type-locality: Argentina, Baradero. Distr. — General through the Neotropical region from Mexico to Argentina.
- dubius* Bellardi, 1859:18 [preocc. by *dubius* Germar, 1837 (Giebel, 1858:221)].
- stonei* Hardy, 1942c:142. Type-locality: "Mexico". Ref. — Hardy, 1953:348.
- WULPI Hardy, 1953:349. Type-locality: "Argentina". Distr. — Argentina. (Change of name for *abbreviatus* Wulp).
- abbreviatus* Wulp, 1881:145 (nec Loew, 1864:54).
- XANTHOPUS Wiedemann, 1828:80. Type-locality: U.S.A., "New York". Distr. — General over United States and probably occurs in Mexico, identified with a query from northern Mexico. Ref. — Hardy, 1945:491, fig. 204a.
- canadensis?* Bellardi, 1859:17.



Genus BIBIONELLUS Edwards

- BIBIONELLUS Edwards, 1935:19. Type-species, *tibialis* Edwards (orig. des.).
- ACZELI Hardy, 1953:345, figs. 1a-c. Type-locality: Argentina, Jujuy, Palpala. Distr. — Argentina.
- BARRETOI Lane & Forattini, 1948:569. Type-locality: Brazil, Goiás, Corumbá. Distr. — Brazil. Ref. — Hardy, 1953:344.
- PAULISTENSIS Lane & Forattini, 1948:571. Type-locality: Brazil, São Paulo, Pitangueiras. Distr. — Brazil. Ref. — Hardy, 1953:344.
- TIBIALIS Edwards, 1935:19. Type-locality: Bolivia, Yungas de Coroico. Distr. — Bolivia. Refs. — Lanc & Forattini, 1948:573; Hardy, 1953:344.

Genus BIBIODES Coquillett

- BIBIODES Coquillett, 1904b:171. Type-species, *halteralis* Coquillett (orig. des.).
- HALTERALIS Coquillett, 1904b:171. Type-locality: California, San Mateo Co. Distr. — Baja California, southwest United States, Mexico. Ref. — Hardy, 1945:496.

Genus ENICOSCOLUS Hardy

- ENICOSCOLUS Hardy, 1961:81. Type-species, *dolichocephalus* Hardy (orig. des.).
- BRACHYCEPHALUS Hardy, 1961:82, fig. 1. Type-locality: Mexico, Morelos, Yautepec. Distr. — Mexico.
- DOLICHOCEPHALUS Hardy, 1961:82, figs. 2a-e. Type-locality: Mexico, Morelos, Tepotztlan. Distr. — Mexico.

Genus DILOPHUS Meigen

- Philia* Meigen, 1800:20. Type-species, *Tipula febrilis* Linnaeus (Coquillett, 1910:588). Suppressed by I.C.Z.N., 1963:339.
- DILOPHUS Meigen, 1803:264. Type-species, *Tipula febrilis* Linnaeus (Latreille, 1810:442).
- Acanthocnemis* Blanchard, 1852:355 (no type des.).
- Triploctenia* Enderlein, 1934:181. Type-species, *Dilophus tenuis* Meigen (orig. des.).
- Tridicroctena* Enderlein, 1934:181. Type-species, *Dilophus africanus* Becker (orig. des.).
- Dactylodiscia* Enderlein, 1934:181. Type-species, *Dilophus hiemalis* Becker (orig. des.).
- Cnemidoctenia* Enderlein, 1934:181. Type-species, *Dilophus crassierus* Lundström (orig. des.).
- ANOMALUS (Hardy), 1942a:127, figs. 1a-b (*Philia*). Type-locality: Mexico, San Rafael. Distr. — Mexico.



- ANTARCTICUS (Walker), 1837:336 (*Bibio*). Type-locality: Argentina, Port Famine. Distr. — Chile, Argentina. Ref. — Hardy, 1953:360.
- ARENARIUS Edwards, 1935:23. Type-locality: Chile, Punta Arenas. Distr. — Chile.
- ATER (Philippi), 1865:638 (*Acanthocnemis*). Type-locality: Chile, Santiago. Distr. — Chile.
- BALFOURI Edwards, 1935:20. Type-locality: Colombia, Villavicencio, Quatiquia River. Distr. — Colombia.
- BICOLOR Macquart, 1838:89 (nec Wiedemann, 1821:34), an unrecognizable species. Recorded from Uruguay, Montevideo.
- BICOLORIPES Edwards, 1938:324. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil, Argentina. Ref. — Hardy, 1953:361, figs. 7a-b.
- BRAZILENSIS (Hardy), 1948:36 (*Philia*). Type-locality: Brazil, Rio de Janeiro, Trecsópolis. Distr. — Brazil. (Change of name for *occipitalis* Edwards).
- occipitalis* Edwards, 1935:21 (preocc. by *occipitalis* Coquillett, 1904:20).
- CALCARATUS Edwards, 1930:83. Type-locality: Argentina, Puerto Blest. Distr. — Argentina. Ref. — Hardy, 1953:359.
- CARBONARIUS (Philippi), 1865:638 (*Acanthocnemis*). Type-locality: Chile, Aconcagua. Distr. — Chile.
- carbonarius* Stuardo, 1946:59 (lapsus).
- CASTANIPES (Bigot), 1888:17 (*Acanthocnemis*). Type-locality: "Cape Horn". Distr. — Cape Horn. Ref. — Hardy, 1953:359.
- COLLARIS Guérin-Méneville, 1843:535, pl. 93, fig. 7. Type-locality: Uruguay, Rio de la Plata. Distr. — Uruguay.
- thoracicus* Guérin-Méneville, 1843:535.
- DICHROUS Edwards, 1938:329. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DISTINGUENDUS Edwards, 1935:23. Type-locality: Chile, Chillan. Distr. — Chile.
- DORSALIS (Philippi), 1865:639 (*Acanthocnemis*). Type-locality: Chile, Santiago. Distr. — Chile, Uruguay, Argentina. Ref. — Hardy, 1953:362, fig. 8a.
- thoracicus* Philippi, 1865:637 (*Acanthocnemis*) (nec Say, 1823, nec Guérin, 1843).
- nigripennis* Philippi, 1865:637 (*Acanthocnemis*).
- EDWARDSI (Hardy), 1948:36 (*Philia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. (Change of name for *minor* Edwards).
- minor* Edwards, 1938:330 (nec *minor* Strobl, 1900:92).
- ELEPHAS Edwards, 1935:22. Type-locality: Peru, Cuzco. Distr. — Peru.
- EPHIPPIUM (Philippi), 1865:638 (*Acanthocnemis*). Type-locality: Chile, Santiago. Distr. — Chile.
- FLAVICORNIS Edwards, 1938:325. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FLAVIFEMUR Edwards, 1930:85. Type-locality: Chile, Casa Panguc. Distr. — Chile. Ref. — Hardy, 1953:359.



- FLAVIHALTER Edwards, 1930:82. Type-locality: Argentina, Lago Gutierrez.
Distr. — Argentina. Ref. — Hardy, 1953:359.
- FLAVISTIGMA Edwards, 1935:23. Type-locality: Chile, Magallanes Territory.
Distr. — Chile.
- FLAVITARSIS Edwards, 1938:327. Type-locality: Brazil, Santa Catarina,
Nova Teutônia. Distr. — Brazil.
- FULVIMACULA (Walker), 1848:116 (*Plectia*). Type-locality: "Venezuela".
Distr. — Venezuela, Ecuador. Refs. — Hardy, 1945:408; 1956:87.
flavimaculata Hunter, 1900:297 (erroneous spelling).
- FUMOSUS Coquillett, 1904a:91. Type-locality: Nicaragua, Granada. Distr.
— Nicaragua.
- GAGATINUS (Philippi), 1865:638 (*Acanthocnemis*). Type-locality: Chile,
Catemu. Distr. — Chile.
gagatinus Hunter, 1900:295 (lapsus).
- GIGANTEUS Macquart, 1846:21, pl. 2, fig. 2. Type-locality: "Nova Grana-
da". Distr. — Colombia.
- GLOBOSUS (Hardy), 1942a:128, figs. 2a-d (*Philia*). Type-locality: Brazil,
São Paulo, Capital, Cantareira. Distr. — Brazil.
- GOLBACHI (Hardy), 1953:364, figs. 9a-c (*Philia*). Type-locality: Argentina,
Formosa, Mision Laishi. Distr. — Argentina.
- GRACILIPES Edwards, 1930:85. Type-locality: Argentina, Lago Gutierrez.
Distr. — Argentina. Ref. — Hardy, 1953:365, fig. 10a.
- HYALPENNIS (Blanchard), 1852:358 (*Acanthocnemis*). Type-locality: Chile,
Coquimbo. Distr. — Chile. Ref. — Edwards, 1930:85.
- IMMACULIPENNIS (Blanchard), 1852:358 (*Acanthocnemis*). Type-locality:
Chile, Coquimbo. Distr. — Chile.
- INCONNEXUS (Hardy), 1961:94, figs. 9a-d (*Philia*). Type-locality: Nor-
thern Venezuela. Distr. — Venezuela.
- INTERRUPTUS Edwards, 1930:86. Type-locality: Chile, Mochuque Is. Distr.
— Chile. Ref. — Hardy, 1953:360.
- LATERALIS (Philippi), 1865:638 (*Acanthocnemis*). Type-locality: Chile, Santi-
ago. Distr. — Ecuador, Chile. Ref. — Edwards, 1920:149.
- LUCIFER Schiner, 1868:18. Type-locality: "Colombia". Distr. — Colombia.
- LUTEICOLLIS (Philippi), 1865:637 (*Acanthocnemis*). Type-locality: "Chile".
Distr. — Chile.
- LUTEUS Edwards, 1930:88. Type-locality: Chile, Casa Pangué. Distr. —
Chile. Ref. — Hardy, 1953:360.
- MACRORHINUS Macquart, 1838b:178. Type-locality: "Chile". Distr. — Chile.
- MACULATUS Bellardi, 1859:19. Type-locality: "Mexico". Distr. — Mexico.
- MACULIPENNIS Blanchard, 1852:354. Type-locality: Chile, Carelmapu. Distr.
— Chile. Ref. — Edwards, 1930:80.
- MEGACANTHUS Edwards, 1938:326. Type-locality: Brazil, Santa Catarina,
Nova Teutônia. Distr. — Brazil.
- MELANARIUS Wulp, 1881:146. Type-locality: "Mexico". Distr. — Mexico,
Costa Rica.
- MICROCERUS Edwards, 1935:24. Type-locality: Argentina, Buenos Aires.
Distr. — Argentina. Ref. — Hardy, 1953:366.
- MINIMUS (Hardy), 1942a:129, figs. 3a-b (*Philia*). Type-locality: Costa
Rica, Higuito, San Mateo. Distr. — Costa Rica.




- MINUTUS Bellardi, 1862:7. Type-locality: "Mexico". Distr. — Mexico.
- NIGRIPES Blanchard, 1852:354. Type-locality: Chile, Coquimbo. Distr. — Chile. Ref. — Edwards, 1930:84.
- NUBILIPENNIS Edwards, 1935:22. Type-locality: Peru, valley of Lari. Distr. — Peru.
- OBSCURUS (Blanchard), 1852:357 (*Acanthocnemis*). Type-locality: Chile, Coquimbo. Distr. — Chile.
- ORNATUS (Hardy), 1942a:130, figs. 4a-b (*Philia*). Type-locality: Guatemala, Guatemala city. Distr. — Guatemala.
- PALLENS (Blanchard), 1852:357 (*Acanthocnemis*). Type-locality: Chile, Chiloe. Distr. — Colombia, Chile. Ref. — Edwards, 1930:80.
- PALIDIPENNIS Philippi, 1865:636. Type-locality: "Chile". Distr. — Chile.
- PATAGONICUS Edwards, 1930:85. Type-locality: Argentina, Bariloche. Distr. — Argentina. Ref. — Hardy, 1953:367.
- PAULSENI Philippi, 1865:636. Type-locality: Chile, Santiago. Distr. — Chile.
- PECTORALIS Wiedemann, 1828:76. Type-locality: Uruguay, Montevideo. Distr. — Uruguay, Argentina. Ref. — Hardy, 1953:368, fig. 11a.
- PERUENSIS (Hardy), 1948:36 (*Philia*). Type-locality: Peru, Cuzco. Distr. — Peru. (Change of name for *clavipes* Edwards).
- clavipes* Edwards, 1935:24 (nec Duda, 1930:33).
- PHILIPPI (Hardy), 1959:470 (*Philia*). Type-locality: Chile, Aconcagua. Distr. — Chile. (Change of name for *bimaculatus* Philippi). Ref. — Hardy, 1959:470.
- bimaculatus* Philippi, 1865:638 (nec Walker, 1858:422).
- PICTUS Schiner, 1868:19. Type-locality: "Colombia". Distr. — Colombia.
- PILIFERUS Edwards, 1935:21. Type-locality: Peru, Mamara. Distr. — Peru.
- PLAUMANNI Edwards, 1938:328. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- QUADRIDENS (Hardy), 1953:370, figs. 12a-c (*Philia*). Type-locality: Argentina, Buenos Aires, San Isidro. Distr. — Argentina.
- QUINQUESPINAE (Hardy), 1961:96, figs. 10a-d (*Philia*). Type-locality: Cuba, Cayamas. Distr. — Cuba, Puerto Rico, Saba (Dutch West Indies).
- RHYNCHOPS Coquillett, 1904a:91. Type-locality: Nicaragua, Granada. Distr. — Nicaragua.
- RUBRICOLLIS (Blanchard), 1852:356 (*Acanthocnemis*). Type-locality: Chile, Coquimbo. Distr. — Chile.
- RUBRIPES (Philippi), 1865:639 (*Acanthocnemis*). Type-locality: "Chile". Distr. — Chile.
- RUFIPES Blanchard, 1852:355. Type-locality: Chile, Coquimbo. Distr. — Chile.
- SAYI (Hardy), 1959:471 (*Philia*). Type-locality: U.S.A., Pennsylvania. Distr. — Common throughout south and central U.S.A. and extends into Mexico. (Change of name for *orbatus* Osten Sacken).
- orbatus* Osten Sacken, 1859:70 (nec Wiedemann, 1828:77).
- SEGREGATUS (Hardy), 1961:98, figs. 11a-c (*Philia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



- SERENUS (Hardy), 1953:371, figs. 13a-c (*Philia*). Type-locality: Argentina, Villa Angostura. Distr. — Argentina.
- SIMILIS Rondani, 1868:40. Type-locality: Argentina, Buenos Aires. Distr. — Argentina. Ref. — Hardy, 1953:358.
- STYGIUS Say, 1829:155. Type-locality: "Mexico". Distr. — Mexico.
- TAPIR Schiner, 1868:18. Type-locality: "Colombia". Distr. — Colombia.
- TESTACEIPES Blanchard, 1852:355. Type-locality: Chile, Coquimbo. Distr. — Ecuador, Chile. Ref. — Edwards, 1920:149.
- TETRACANTHUS Edwards, 1930:87. Type-locality: Argentina, Bariloche. Distr. — Argentina. Ref. — Hardy, 1953:360.
- TRISULCATUS Macquart, 1838a:88. Type-locality: "Brazil". Distr. — Brazil, Argentina. Ref. — Hardy, 1953:373, figs. 14a-c.
- VALDIVIANUS Philippi, 1865:636. Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Edwards, 1930:80.
- VARICEPS (Hardy), 1942a:133, fig. 5a (*Philia*). Type-locality: Panama, Tabernilla. Distr. — Panama, Peru (?).
- VENULATUS Johnson, 1920:426. Type-locality: Jamaica, Newton. Distr. — Jamaica.
- VILLOSUS Edwards, 1936:21. Type-locality: Ecuador, Loja. Distr. — Ecuador.
- VITATUS Philippi, 1865:636. Type-locality: Chile, Valdivia. Distr. — Ecuador, Chile. Ref. — Edwards, 1930:80.

LITERATURE CITED

- ALDRICH, J. M.
1905. A Catalogue of North American Diptera. *Smiths. Misc. Coll.* 46:1-680.
- ARRIBÁLZAGA, F. L.
1878. Notas dipterológicas sobre los Bibionites del Baradero. *El Naturalista Argentino* 1:295-299.
- BELLARDI, L.
1859. *Saggio di ditteologia messicana* 1:14-19.
1862. *Saggio di ditteologia messicana*, Append., p. 4.
- BIGOT, J.
1888. *Mission Scientifique au Cap Horn*, Zool. 6:17.
- BLANCHARD, E.
1852. Diptera, in Gay's *Historia Fisica y Politica de Chile, Fauna Chilena, Insectos*, Zool. 7:353-358.
- COQUILLET, D. W.
1904a. New Diptera from Central America. *Proc. Ent. Soc. Washington* 6:91.
1904b. New North American Diptera. *Proc. Ent. Soc. Washington* 6:171.



1904c. in Baker, C. F.: Diptera. Reports on Californian and Nevada Diptera. I. *Invertebrata Pacifica* 1:20.

1910. The type-species of the North American Genera of Diptera. *Proc. U.S. Nat. Mus.* 37:499-647.

CURRAN, C. H.

1934. The Diptera of Kartabo, Bartica District, British Guiana. *Bull. Amer. Mus. Nat. Hist.* 46(3):310.

DUDA, O.

1930. 4. Bibionidae, in Lindner, E.: *Die Fliegen der palaarktischen Region* 2(1):1-75.

EDWARDS, F. W.

1920. Diptères Nematocères. *Miss. Arc MÉR. Equat., Paris* 10(2):147-149.

1930. Bibionidae, in *Diptera of Patagonia and South Chile* 2(3):77-88.

1931a. Die Ausbeute der deutschen Chaco-Expedition 1925/26 — Diptera. XXIII. Bibionidae. *Könovia* 10(1):75-76.

1931b. New Neotropical Nematocerous Diptera. *Ann. & Mag. Nat. Hist.* (10)7:260-261.

1935. New Neotropical Bibionidae (Diptera). *Stylops* 4(1):19-24.

1938. Bibionidae (Diptera) collected by Mr. F. Plaumann in Brazil. *Ann. & Mag. Nat. Hist.* (11)2:321-330.

ENDERLEIN, G.

1934. Dipterologia. II. *Sitzber. Ges. Naturf. Freunde, Berlin* 2:181.

FABRICIUS, J. C.

1771. *Species Insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin* 2:410.

1798. *Entomologia systematica*, Suppl.: 551.

1805. *Systema Antliatorum, secundum ordines, genera, species*, pp. 53-54, 163. *Brunsvigae*.

GEOFFROY, E. L.

1764. *Histoire abrégée des Insectes* 2:571.

GERMAR, E. F.

1837. *Fauna Insectorum Europae* 19. *Insectorum Protogeae specimen sistens insecta carborum fossilium*, pp. 1-25. Halle.

GIEBEL, C. G. A.

1856. *Fauna der Vorwelt, mit steter Berücksichtigung der lebenden Thiere* 2:221.

GUÉRIN-MÉNEVILLE, F. E.

1843. *Iconographie du règne animal de G. Cuvier*, Paris 7:535.

HARDY, D. E.

1937. New Bibionidae (Diptera) from Nearctic America. *Proc. Utah Acad. Sci.* 14:199.



1940. Studies in New World *Plecia* (Bibionidac-Diptera) Part I. *Journ. Kansas Ent. Soc.* 13(1):15-27.
- 1942a. Studies in New World *Philia* (Bibionidac) Part I. *Journ. Kansas Ent. Soc.* 15(4):127-134.
- 1942b. Studies in New World *Plecia* (Diptera-Bibionidac) Part II. *Can. Ent.* 74:106-116.
- 1942c. Notes on Diptera in Snow Entomological Collection. *Journ. Kansas Ent. Soc.* 15(4):142.
1945. Revision of Nearctic Bibionidac including Neotropical *Plecia* and *Penthetria*. *Univ. Kansas Sci. Bull.* 30(2):367-547.
1948. Homonymy notes in the Bibionidae. *Journ. Kansas Ent. Soc.* 21(1):36.
1953. The Argentine Bibionidae (Diptera). *Acta Zool. Lilloana* 12:343-376.
1956. The Walker types of Bibionidae (Diptera). *Journ. Kansas Ent. Soc.* 29(3):85-91.
1957. New Neotropical Bibionidac in the Naturhistorisches Museum, Vienna. *Ann. Naturhist. Mus. Wien* 61:238-240.
1959. Catalogue of the Neotropical Bibionidae. *Acta Zool. Lilloana* 17:437-476.
1961. Notes and descriptions of exotic Bibionidae. *Proc. Ent. Soc. Washington* 63(2):81-99.

HARRIS, M.

1776. *An exposition of English Insects...*, London, p. 77.

HUNTER, W. D.

1900. Catalogue of the Diptera of South America. Part. I. — Bibliography and Nemocera. *Trans. Amer. Ent. Soc.* 26:294-297.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE


1957. Opinion 441. Validation under Plenary Powers of the names of five genera in the Order Diptera (Class Insecta) published in 1762 by Geoffroy (E.L.) in the work entitled "Histoire abrégée des insectes qui se trouvent aux environs de Paris" ("Opinion" supplementary to "Opinion" 228). pp. 83-120 (= pt. 6), in Hemming, F., ed., *Opinions and declarations rendered by the International Commission of Zoological Nomenclature* 15:530 pp., London.
1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.* 20:339-342.

JAENNICKE, F.

1867. Neue exotische Dipteren. *Abhandl. Senckenb. naturf. Ges.* 6:318.

JOHNSON, C. W.

1920. A revised list of the Diptera of Jamaica. *Bull. Amer. Mus. Nat. Hist.* 41(8):426.



KERTÉSZ, C.

1902. *Catalogus Dipteriorum* 1:135-156.

LANE, J. & O. P. FORATTINI

1948. Duas novas espécies de *Bibionellus* Edwards, 1935 (Diptera, Bibionidae). *Rev. Ent.*, Rio de Janeiro 19(3):569-574.

LOEW, H.

1858. Ueber einige neue Fliegengattungen. *Berl. ent. Zeitschr.* 2:101-109.

1864. *Diptera americanae septentrionalis indigena*. Centuria Quinta. *Berl. ent. Zeitschr.* 8:49-104.

MACQUART, J.

1838a. *Diptères exotiques, nouveaux ou peu connus* 1(1):84-89.

1838b. *Idem* 1(2):178.

1844. *Idem*, Suppl. 1:148-150.

1846. *Idem*, Suppl. 1:20-22.

1847. *Idem*, Suppl. 2:26.

MEIGEN, J. G. [W.]

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, Paris, p. 20.

1803. Versuch einer neuen Gattungs Eintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2:264.

1804. *Klassifikation und Beschreibung der europäischen zweiflügeligen Insekten (Diptera Linn.)*, Braunschweig 1:104.

OSTEN SACKEN, C. R.

1891. in Le Conte: *The Complete Writings of Thomas Say* 2:70.

PERTY, M.

1830. *Delectus Animalium articulorum, quae in itinere per Brasiliam...*, p. 180.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool. bot. Ges. Wien* 15:635-639.

RONDANI, C.

1850. Osservazioni sopra alcune specie di esapodi ditteri del Museo Torinese. *R. Accad. Sci. Ist. Bologna, Nuovi Ann. Sci. Nat.* 2:193.

1868. *Diptera alicae in America meridionali lecta a Prof. S. Strobel annis 1866 et 1867*. *Annuario della Soc. dei Nat. Modena*: 39-40.

SAY, T.

1823. Descriptions of dipterous insects of the United States. *Journ. Philad. Acad. Sci.* 3:78.

1829. Descriptions of North American dipterous insects. *Journ. Philad. Acad. Sci.* 6:154-155.

- SCHINER, J. R.
 1868. *Reise der österreichischen Fregatte Novara. Zool. Theil, I. Diptera*: 16-23.
- STROBL, G.
 1900. Spanische Dipteren. *Wien. Ent. Ztg.* 19:92.
- STUARDO, C.
 1946. *Catálogo de los dipteros de Chile*, 250 pp.
- TOWNSEND, C. H. T.
 1912. On the Diptera of Baja California, including some species from adjacent regions. *Can. Ent.* 44:289.
- WALKER, F.
 1837. Descriptions, etc., of the insects collected by Capt. P. King in the survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. London* 17:336.
 1848. *List of specimens of dipterous insects in the collection of the British Museum* 1:81-116.
 1858. Characters of undescribed Diptera in the collection of W. W. Saunders. *Trans. Ent. Soc. London* (n. s.) 5:422.
- WIEDEMANN, C. R. W.
 1821. *Diptera exotica* 1:34.
 1824. *Munus rectoris in Academia Christiana Albertina aditurus. Analecta entomologica ex Museo Regio Havniensi*, p. 11. Kiliae.
 1828. *Aussereuropäische zweifügelige Insekten* 1:72-81.
 1830. *Idem* 2:618.
- WILLISTON, S. W.
 1900. *Biologia Centrali-Americana, Insecta. Diptera 1, Suppl.*: 221-223.
- WULF, F. M. VAN DER
 1881. Amerikaansche Diptera. *Tijdschr. v. Ent.* 24:143-146.

INDEX

<i>abbreviatus</i> Wulp, 1881, <i>Bibio</i>	18.6
<i>Acanthocnemis</i> Blanchard, 1852	18.7
<i>aczelii</i> Hardy, 1954, <i>Bibionellus</i>	18.7
<i>alacris</i> Curran, 1934, <i>Plecia</i>	18.2
<i>albipennis</i> Say, 1823, <i>Bibio</i>	18.6
<i>americana</i> Hardy, 1940, <i>Plecia</i>	18.2
<i>anomalus</i> (Hardy), 1942 (<i>Philia</i>), <i>Dilophus</i>	18.7
<i>antarcticus</i> Edwards, 1935, <i>Dilophus</i>	18.8
<i>appendicula</i> Hardy, 1945, <i>Penthetria</i>	18.2
<i>arenarius</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>argentina</i> Hardy, 1953, <i>Plecia</i> , subspecies of <i>nigra</i> Philippi	18.4
<i>ater</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>avicephaliforma</i> Hardy, 1940, <i>Plecia</i>	18.2




<i>balfouri</i> Edwards, 1935, <i>Dilophus</i>	18.8
<i>barrettoi</i> Lane & Forattini, 1948, <i>Bibionellus</i>	18.7
<i>bellardii</i> Townsend, 1912, <i>Plecia</i>	18.4
<i>biarmata</i> Hardy, 1942, <i>Plecia</i>	18.2
<i>Biblio</i> Geoffroy, 1762	18.6
<i>Bibiodes</i> Coquillett, 1904	18.7
<i>Bibionellus</i> Edwards, 1935	18.7
<i>Bibioninae</i>	18.6
<i>bicolor</i> Bellardi, 1859, <i>Plecia</i>	18.2
<i>bicolor</i> Macquart, 1838, <i>Dilophus</i>	18.8
<i>bicoloripes</i> Edwards, 1938, <i>Dilophus</i>	18.8
<i>biformis</i> Hardy, 1942, <i>Plecia</i>	18.2
<i>bimaculatus</i> Philippi, 1865, <i>Dilophus</i>	18.10
<i>brachycephalus</i> Hardy, 1961, <i>Enicoscolus</i>	18.7
<i>brazilana</i> Hardy, 1942, <i>Plecia</i>	18.2
<i>brazilensis</i> (Hardy), 1948 (<i>Philia</i>), <i>Dilophus</i>	18.8
<i>calcaratus</i> Edwards, 1930, <i>Dilophus</i>	18.8
<i>canadensis</i> Bellardi, 1859, <i>Biblio</i>	18.6
<i>carbonarius</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>castanipes</i> (Bigot), 1888 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>clavipes</i> Edwards, 1935, <i>Dilophus</i>	18.10
<i>Cnemidoctenia</i> Enderlein, 1934	18.7
<i>collaris</i> (Fabricius), 1805 (<i>Hirtea</i>), <i>Plecia</i>	18.2
<i>collaris</i> Guérin-Ménéville, 1843, <i>Dilophus</i>	18.8
<i>concava</i> Hardy, 1942, <i>Plecia</i> , subspecies of <i>rufithorax</i> Walker ...	18.5
<i>confusa</i> Loew, 1848, <i>Plecia</i>	18.2
<i>conjungens</i> Schiner, 1868, <i>Hesperinus</i>	18.1
<i>carbonarius</i> Stuardo, 1946, <i>Dilophus</i> (misspell.)	18.8
<i>costalis</i> Walker, 1858, <i>Plecia</i>	18.3
<i>costalis</i> Wiedemann, 1830, <i>Plecia</i>	18.3
<i>criorhinus</i> Bellardi, 1859, <i>Biblio</i>	18.6
<i>curvistylata</i> Hardy, 1942, <i>Plecia</i>	18.3
<i>Dactylodiscia</i> Enderlein, 1934	18.7
<i>dentata</i> Hardy, 1942, <i>Plecia</i>	18.3
<i>dichrous</i> Edwards, 1938, <i>Dilophus</i>	18.8
<i>Dilophus</i> Meigen, 1803	18.7
<i>discolor</i> Wulp, 1881, <i>Plecia</i>	18.3
<i>dispar</i> Schiner, 1868, <i>Biblio</i>	18.6
<i>disparis</i> Hardy, 1942, <i>Plecia</i>	18.3
<i>distincta</i> Hardy, 1945, <i>Penthetria</i>	18.2
<i>distinguendus</i> Edwards, 1935, <i>Dilophus</i>	18.8
<i>dolichocephalus</i> Hardy, 1961, <i>Enicoscolus</i>	18.7
<i>dorsalis</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>dubius</i> Bellardi, 1859, <i>Biblio</i>	18.6
<i>ecuadorensis</i> Hardy, 1942, <i>Plecia</i>	18.3
<i>edwardsi</i> Hardy, 1940, <i>Plecia</i>	18.3
<i>edwardsi</i> (Hardy), 1948 (<i>Philia</i>), <i>Dilophus</i>	18.8
<i>elephas</i> Edwards, 1938, <i>Dilophus</i>	18.8
<i>Enicoscolus</i> Hardy, 1961	18.7
<i>ephippium</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>Eupeitenus</i> Macquart, 1838	18.1



<i>femorata</i> Macquart, 1838, Plecia	18.3
<i>flavicornis</i> Edwards, 1938, Dilophus	18.8
<i>flavifemur</i> Edwards, 1930, Dilophus	18.8
<i>flavihalter</i> Edwards, 1930, Dilophus	18.8
<i>flavimaculata</i> Hunter, 1900, Dilophus (misspell.)	18.9
<i>flavistigma</i> Edwards, 1935, Dilophus	18.9
<i>flavitarsis</i> Edwards, 1938, Dilophus	18.9
<i>fuliginus</i> Bellardi, 1859, Bibio	18.6
<i>fuliginosus</i> Aldrich, 1905, Bibio (misspell.)	18.6
<i>fulvamacula</i> (Walker), 1848 (Plecia), Dilophus	18.9
<i>fumosus</i> Coquillett, 1904, Dilophus	18.9
<i>funebri</i> (Fabricius), 1805 (Hirtca), Plecia	18.3
<i>gagantinus</i> Hunter, 1900, Dilophus (misspell.)	18.9
<i>gagatinus</i> (Philippi), 1865 (Acanthocnemis), Dilophus	18.9
<i>gibbosa</i> Hardy, 1942, Plecia	18.3
<i>giganteus</i> Macquart, 1846, Dilophus	18.9
<i>globosus</i> (Hardy), 1942 (Philia), Dilophus	18.9
<i>golbachi</i> (Hardy), 1953 (Philia), Dilophus	18.9
<i>gracilipes</i> Edwards, 1930, Dilophus	18.9
<i>grisea</i> Edwards, 1938, Plecia	18.3
<i>halteralis</i> Coquillett, 1940, Bibiodes	18.7
<i>heros</i> Say, 1829, Penthetria	18.2
<i>Hesperininae</i>	18.1
<i>Hesperinus</i> Walker, 1848	18.1
<i>heteroptera</i> Macquart, 1846, Plecia	18.4
<i>Hirtea</i> Fabricius, 1798	18.6
<i>hyalipennis</i> (Blanchard), 1852 (Acanthocnemis), Dilophus	18.9
<i>immaculipennis</i> (Blanchard), 1852 (Acanthocnemis), Dilophus	18.9
<i>impensa</i> Hardy, 1957, Plecia	18.3
<i>imperialis</i> Schiner, 1868, Plecia	18.3
<i>impilosa</i> Hardy, 1940, Plecia	18.3
<i>inconnexus</i> (Hardy), 1961 (Philia), Dilophus	18.9
<i>incurvata</i> Hardy, 1942, Plecia	18.3
<i>intermedius</i> Rondani, 1850, Bibio	18.6
<i>interruptus</i> Edwards, 1930, Dilophus	18.9
<i>lateralis</i> Hardy, 1940, Plecia	18.3
<i>lateralis</i> (Philippi), 1865 (Acanthocnemis), Dilophus	18.9
<i>lindneri</i> Edwards, 1931, Plecia	18.3
<i>lopesi</i> Hardy, 1940, Plecia	18.3
<i>lucifer</i> Schiner, 1868, Dilophus	18.9
<i>luteicollis</i> (Philippi), 1865 (Acanthocnemis), Dilophus	18.9
<i>luteus</i> Edwards, 1930, Dilophus	18.9
<i>macrorhinus</i> Macquart, 1838, Dilophus	18.4
<i>maculata</i> Hardy, 1942, Plecia	18.9
<i>maculatus</i> Bellardi, 1859, Dilophus	18.9
<i>maculipennis</i> Blanchard, 1852, Dilophus	18.4
<i>marginata</i> Edwards, 1920, Plecia	18.4
<i>maura</i> Walker, 1837, Plecia	18.9
<i>megacanthus</i> Edwards, 1938, Dilophus	18.9



<i>melanarius</i> Wulp, 1881, <i>Dilophus</i>	18.9
<i>mexicana</i> (Hardy), 1937 (<i>Plecia</i>), <i>Penthetria</i>	18.2
<i>micans</i> Hardy, 1942, <i>Plecia</i> , subspecies of <i>ecuadorensis</i> Hardy ...	18.3
<i>microcerus</i> Edwards, 1935, <i>Dilophus</i>	18.9
<i>minimus</i> (Hardy), 1942 (<i>Philia</i>), <i>Dilophus</i>	18.9
<i>minor</i> Edwards, 1938, <i>Dilophus</i>	18.8
<i>minor</i> Jaennlecke, 1867, <i>Plecia</i>	18.5
<i>minutus</i> Bellardi, 1862, <i>Dilophus</i>	18.10
<i>nearctica</i> Hardy, 1940, <i>Plecia</i>	18.4
<i>nigerrima</i> (Bellardi), 1859 (<i>Plecia</i>), <i>Penthetria</i>	18.2
<i>nigra</i> (Philippi), 1865 (<i>Penthera</i>), <i>Plecia</i>	18.4
<i>nigripennis</i> (Philippi), 1865 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.8
<i>nigripes</i> Blanchard, 1852, <i>Dilophus</i>	18.10
<i>nigrita</i> Perty, 1830, <i>Penthetria</i>	18.2
<i>nitidicollis</i> Edwards, 1931, <i>Plecia</i>	18.4
<i>nitidipes</i> Edwards, 1920, <i>Plecia</i>	18.4
<i>nodioculatus</i> Kertész, 1902, <i>Bibio</i> (misspell.)	18.6
<i>nubilipennis</i> Edwards, 1935, <i>Dilophus</i>	18.10
<i>nudiceulatus</i> Maequart, 1847, <i>Bibio</i>	18.6
<i>obscurus</i> (Blanchard), 1852 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.10
<i>occipitalis</i> Edwards, 1935, <i>Dilophus</i>	18.8
<i>orbatus</i> Osten Sacken, 1859, <i>Dilophus</i>	18.10
<i>ornatus</i> (Hardy), 1942 (<i>Philia</i>), <i>Dilophus</i>	18.10
<i>pallens</i> (Blanchard), 1852 (<i>Acanthocnemis</i>), <i>Dilophus</i>	18.10
<i>pallidipennis</i> Philippi, 1865, <i>Dilophus</i>	18.10
<i>panamaensis</i> Hardy, 1942, <i>Plecia</i>	18.4
<i>paracollaris</i> Hardy, 1961, <i>Plecia</i>	18.4
<i>patagonicus</i> Edwards, 1930, <i>Dilophus</i>	18.10
<i>parvistylata</i> Hardy, 1942, <i>Plecia</i>	18.4
<i>paulistensis</i> Lane & Forattini, 1948, <i>Bibionellus</i>	18.7
<i>paulseni</i> Philippi, 1865, <i>Dilophus</i>	18.10
<i>pectoralis</i> Wiedemann, 1828, <i>Dilophus</i>	18.10
<i>Penthera</i> Philippi, 1865	18.2
<i>Penthetria</i> Meigen, 1803	18.1
<i>perplexa</i> Hardy, 1942, <i>Plecia</i>	18.4
<i>persimilis</i> Hardy, 1940, <i>Plecia</i>	18.4
<i>pertinens</i> Hardy, 1942, <i>Plecia</i>	18.4
<i>perucnsis</i> (Hardy), 1948 (<i>Philia</i>), <i>Dilophus</i>	18.10
<i>peruviana</i> Hardy, 1961, <i>Plecia</i>	18.4
<i>peruvianus</i> Edwards, 1935, <i>Bibio</i>	18.6
<i>Philia</i> Meigen, 1800	18.7
<i>philippii</i> (Hardy), 1959 (<i>Philia</i>), <i>Dilophus</i>	18.10
<i>piceus</i> Bellardi, 1859, <i>Bibio</i>	18.6
<i>pictipennis</i> Edwards, 1931, <i>Plecia</i>	18.4
<i>pictus</i> Schiner, 1868, <i>Dilophus</i>	18.10
<i>piliferus</i> Edwards, 1935, <i>Dilophus</i>	18.10
<i>plagiata</i> Wiedemann, 1824, <i>Plecia</i>	18.4
<i>platystyla</i> Hardy, 1957, <i>Plecia</i>	18.4
<i>plaumanni</i> Edwards, 1938, <i>Dilophus</i>	18.10
<i>Plecia</i> Wiedemann, 1828	18.2
<i>Pleciinae</i>	18.1



pruinosa Hardy, 1940, Plecia	18.5
<i>Pullata</i> Harris, 1776	18.6
punctulata Hardy, 1940, Plecia	18.5
quadridens (Hardy), 1953 (Philia), Dilophus	18.10
quadrivittata Williston, 1900, Plecia	18.5
quasimaculata Hardy, 1961, Plecia	18.5
quinquespinae (Hardy), 1961 (Philia), Dilophus	18.10
rectiora Hardy, 1942, Plecia	18.5
<i>Rhinoplecia</i> Bellardi, 1859	18.2
rhynchops Coquillett, 1904, Dilophus	18.10
rostellata Loew, 1858, Plecia	18.5
<i>rostrata</i> Bellardi, 1859, Plecia	18.5
rubricollis (Blanchard), 1852 (Acanthocnemis), Dilophus	18.10
rubripes (Philippi), 1865 (Acanthocnemis), Dilophus	18.10
<i>ruficollis</i> (Fabricius), 1805 (Histera), Plecia	18.2
rufimarginata Hardy, 1940, Plecia	18.5
rufipes Blanchard, 1852, Dilophus	18.10
rufiscutella Hardy, 1940, Plecia	18.5
rufithorax Walker, 1848, Plecia	18.5
rufovittata Edwards, 1938, Plecia, variety of pictipennis Edwards.	18.4
rugosa Hardy, 1940, Plecia	18.5
sayi (Hardy), 1959 (Philia), Dilophus	18.10
segregatus (Hardy), 1961 (Philia), Dilophus	18.10
seminitens Edwards, 1938, Plecia	18.5
serrata Hardy, 1942, Plecia	18.5
serenus (Hardy), 1953 (Philia), Dilophus	18.11
similis Rondani, 1868, Dilophus	18.11
similis Rondani, 1850, Plecia	18.5
<i>Spodius</i> Loew, 1868	18.1
<i>stonei</i> Hardy, 1942, Bibio	18.6
stygius Say, 1829, Dilophus	18.11
subaequalis Rondani, 1868, Bibio	18.6
superfluus Schiner, 1868, Bibio	18.6
tapir Schiner, 1868, Dilophus	18.11
testaceipes Blanchard, 1852, Dilophus	18.11
tetracanthus Edwards, 1930, Dilophus	18.11
<i>thoracica</i> (Fabricius), 1805 (Laphria), Plecia	18.2
<i>thoracicus</i> Guérin-Ménéville, 1843, Dilophus	18.8
<i>thoracicus</i> Philippi, 1865 (Acanthocnemis), Dilophus	18.8
tibialis Edwards, 1935, Bibionellus	18.7
<i>Tridicroctena</i> Enderlein, 1934	18.7
trilobata Hardy, 1942, Plecia	18.5
<i>Triploctenia</i> Enderlein, 1934	18.7
trisulcatus Macquart, 1838, Dilophus	18.11
uberta Hardy, 1942, Plecia	18.5
valdivianus Philippi, 1865, Dilophus	18.11
varabilis Hardy, 1942, Plecia	18.5
variceps (Hardy), 1942 (Philia), Dilophus	18.11

<i>velutina</i> Macquart, 1846, Plecia	18.4
<i>venulatus</i> Johnson, 1920, Dilophus	18.11
<i>vicinus</i> Arribáizaga, 1878, Bibio	18.1
<i>villosus</i> Edwards, 1935, Dilophus	18.11
<i>vittata</i> Bellardi, 1862, Plecia	18.4
<i>vittata</i> Wiedemann, 1828, Plecia	18.6
<i>vittatus</i> Philippi, 1865, Dilophus	18.11
<i>wulpi</i> Hardy, 1953, Bibio	18.6
<i>xanthopus</i> Wiedemann, 1828, Bibio	18.6
<i>xenia</i> Hardy, 1942, Plecia	18.6





SciELO



SUPERFAMILY MYCETOPHILOIDEA

Family Diadocidiidae

Family Ditomyiidae

Family Keroplatidae

Family Lygistorrhinidae

Family Mycetophilidae


Family Sciaridae

Family Hyperoscelididae

Family Scatopsidae

Family Cecidomyiidae

Family Perissommatidae





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

19 A

FAMILY DIADOCIDIIDAE

N. PAPAVERO

Museu de Zoologia
Universidade de São Paulo

Three genera occur in South America, two of which only tentatively referred to this family. Larvae of *Diadocidia* live in mucus tubes in wood.

Genus DIADOCIDIA Ruthe

DIADOCIDIA Ruthe, 1821: 1210. Type-species, *flavicans* Ruthe (orig. des., as gen. n., sp. n.) = *ferruginosa* (Meigen).

Macronevra Macquart, 1834: 146. Type-species, *winthemii* Macquart (mon.) = *ferruginosa* (Meigen).

Subgenus ADIDOCIDIA Lastovka & Matile

Diadocidia, subg. ADIDOCIDIA Lastovka & Matile, 1972: 218. Type-species, *valida* Mik (orig. des.).

NIGRIPALPIS Edwards, 1940: 443. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Syntypes ♂ ♀, BMNH. Lastovka & Matile (1972: 218) have tentatively referred this species to their subgenus *Adidocidia*; I have examined a male topotype, and can confirm their assumption. The male genitalia is similar to that of *borealis* Coquillett — the hypandrium is intermediary in shape between *borealis* and *valida* Mik; the apex of the dististylus is very similar to *borealis*, but there are two very strong black spines in addition to the bifid apex. Freeman's (1951: 11) record of this species from Chile (Marga Marga) may refer to a new species.

sp. Edwards, 1940: 44. Chile, Valparaiso, Marga Marga. ♂, BMNH.

Genus HETEROTRICHA Loew

HETEROTRICHA Loew, 1850: 33. Type-species, *hirta* Loew (mon.).

Palaeoheterotricta Meunier, 1904: 62. Type-species, *grandis* Meunier (mon.).

19A.1

Representatives of this genus are known from the Baltic amber, Chile, southern Brazil, South Africa, Tasmania and New Zealand. Edwards (1925) and Shaw (1953) considered it as a Sciaridae. Tonnoir & Edwards (1927), Frey (1942), Colless (1963) and Hennig (1954, 1973) as probably a Diadocidiidae. The situation of this genus is still uncertain.

- EQUALIS Freeman, 1951: 28, fig. 8 (♂ genitalia). Type-locality: Chile, Chiloé, Castro. Type ♂, BMNH.
- NUDICOSTALIS Freeman, 1951: 28, fig. 7 (♂ genitalia). Type-locality: Chile, Chiloé, Meehuque I. Type ♂, BMNH.
- SEMINUDA Freeman, 1951: 27, fig. 6 (♂ genitalia), 270 (wing). Type-locality: Chile, Chiloé, Ancud. Distr. — Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- MARGINATA Edwards, 1940: 444. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Genus PTEROGYMNUS Freeman

PTEROGYMNUS Freeman, 1951: 11. Type-species, *elongatus* Freeman (orig. des.).

Pterogymnus was originally included in Diadocidiidae. Hennig (1954: 306) very judiciously observed that it could very well be a Bolitophilidae. Colless (1963) and Hennig (1973) preferred to leave it in Diadocidiidae again.

ELONGATUS Freeman, 1951: 12, fig. 271 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Type ♀, BMNH.

LITERATURE CITED

COLLESS, D. H.

1963. New species of *Ohakunea* Edwards and a related new genus with notes on the relationships of *Heterotricha* Loew (Diptera). *Proc. Linn. Soc. N. S. W.* (1962) 87: 303-308, 2 figs.

EDWARDS, F. W.

1925. Mycetophilidae and Eibionidae (Diptera) in the collections of the South African Museum. *Ann. S. Afr. Mus.* 19: 601-616, 3 figs.
1940. New neotropical Mycetophilidae (IV) (Diptera). *Revta Ent.* 11 (1-2): 440-465, pls. 18-19.

FREEMAN, P.

1951. Mycetophilidae, in British Museum (Natural History), *Diptera of Patagonia and South Chile* 3: 1-133, 49 pls. London.

FREY, R.

1942. Entwurf einer neuen Klassifikation der Mückenfamilie Sciaridae. *Notulae ent.* 22: 5-45, 12 figs.

19A.2



HENNIG, W.

1954. Flügelgeäder und System der Dipteren unter Berücksichtigung der aus dem Mesozoikum beschriebenen Fossilien. *Beitr. Ent.* 4: 246-388, 272 figs.
1973. Diptera (Zweiflügler), pp. 1-337, in Helmeke, J. G., D. Starek & H. Wermuth (eds.), *Handbuch der Zoologie. Eine Naturgeschichte der Stämme des Tierreichs gegründet von Willy Kükenthal*, 4 (2) 2/31.

LASTOVKA, P. & L. MATILE

1972. Révision des *Diadoecidia* holaretiques (Dipt. Mycetophilidae). *Ann. Soc. Ent. France* (n.s.) 8 (1): 205-223, 28 figs.

LOEW, H.

1850. Ueber den Bernstein und die Bernsteinfauna. *Programm K. Realschule zu Meseritz 1850*: 1-44. [Also separately published, Berlin, 1850].

MACQUART, J.

1834. *Histoire naturelle des Insectes. — Diptères*, 1: 578 pp., 12 pls., in [N. E. Roret, ed.] (Collection des suites à Buffon). Paris.

MEUNIER, F.

1904. Monographie des Cecidomyidae, des Scleridae, des Mycetophilides et des Chironomides de l'ambre de la Baltique. *Ann. Soc. sci. Bruxelles* 28: 19-92.

RUTHE, J. F. VON


1831. Einige Bemerkungen und Nachträge zu Meigen's "Systematische Beschreibung der europäischen zweiflügeligen Insecten". *Isis* 11: 1203-1222.

SHAW, F. R.

1953. A review of some of the more important contributions to our knowledge of the systematic relationships of the Scleridae (Diptera). *Proc. Hawaii. Ent. Soc.* 15: 25-32.

TONNOIR, A. L. & F. W. EDWARDS

1927. New Zealand fungus gnats (Diptera, Mycetophilidae). *Trans. N. Z. Inst.* 57: 747-876, pls. 58-80.





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

19 B

FAMILY DITOMYIIDAE

N. PAPAVERO


Museu de Zoologia
Universidade de São Paulo

Approximately 100 species known in the world, except in the Ethiopian and Oriental regions. Larvae of *Ditomyia* and *Symmerus* (holarctic) live in hard species of Polyporaceae and in decaying wood, respectively. Nothing is known about the biology of the neotropical species.

A new arrangement of the neotropical genera is here proposed; they may be recognized by the following key:

- | | |
|---|---------------------------------|
| 1. Transverse vein r-m (ta) present; alula present | 2 |
| Transverse vein r-m (ta) absent, obliterated by the fusion, for a short distance, of M with R _s ; alula absent | 5 |
| 2. 4 scutellar bristles | 3 |
| 2 scutellar bristles | 4 |
| 3. Sc parallel to R, apically obsolescent; squama pilose (Chile-Patagonia; Australian region) <i>Australosymmerus</i> Freman | |
| Sc short, strong, curved up to meet the Costa; squama bare (Patagonia) | <i>Neocrionisca</i> , gen. n. |
| 4. Anterior tibia without spines along its length; wings with macro- and microtrichiae on membrane (Chile-Patagonia) | <i>Nothosymmerus</i> , gen. n. |
| Anterior tibia with several well-developed spines or bristles along its length (Guiano-Brazilian subregion; north to Mexico) | |
| | <i>Austrosymmerus</i> , gen. n. |
| 5. Eyes reniform or oval, never emarginated above antennae | 6 |
| Eyes strongly emarginated above antennae, forming a bridge as in Sciaridae; base of fork of M evanescent; antennae with cylindrical segments; R ₄ relatively short (New Zealand; southern South America) | <i>Nervijuncta</i> Marshall |
| 6. Palpi 3-segmented; antennae with flagellar segments projecting into ventral filamentose expansions; R ₄ moderately long (Brazil: São Paulo) | <i>Calliceratomyia</i> Lane |
| Palpi 2-segmented; projections of antennal flagellum not filamentose (Southern Brazil: São Paulo to Santa Catarina) | |
| | <i>Rhipidita</i> Edwards |

19B.1



Genus AUSTRALOSYMMERUS Freeman

- Centrocnemis* Philippi, 1865: 619 (preocc. Signoret, 1852). Type-species, *stigmatica* Philippi (mon.).
- Australomyia* Freeman, 1951: 7 (nom. nov. for *Centrocnemis* Philippi, but preocc. Schmid, 1949). Type-species, *Centrocnemis stigmatica* Philippi (aut.).
- AUSTRALOSYMMERUS Freeman, 1954: 39 (nom. nov. for *Australomyia* Freeman). Type-species, *Centrocnemis stigmatica* Philippi (aut.).
- BIVITTATUS (Freeman), 1951: 9, fig. 4 (♂ genitalia) (*Australomyia*). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro, Neuquén), Chile (Osorno, Cautín, Llanquihue, Chiloé). Type ♂, BMNH.
- SIMPLEX (Freeman), 1951: 8, figs. 3 (♂ genitalia), 299 (wing) (*Australomyia*). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro). Type ♂, BMNH.
- STIGMATICUS (Philippi), 1865: 619, pl. 23, fig. 7 (whole insect) (*Centrocnemis*). Type-locality: Chile, Valdivia. Distr. — Argentina (Río Negro, Neuquén, Tierra del Fuego), Chile (Valparaíso to Chiloé). Type?

Genus AUSTROSYMMERUS, new

Differs from *Australosymmerus* and *Neocrionisca* by the possession of only 2 scutellar bristles; from *Nothosymmerus* by the presence of bristles or spines on the entire length of the anterior tibia. Distr. — Guiano-Brazilian subregion, north to Mexico.

Type-species, *Centrocnemis bisetosus* Edwards.

- BIFASCIATUS (Williston), 1900: 217, pl. 4, figs. 1 (whole insect), 1a (head, lateral) (*Plesiastina*). Type-locality: Mexico, Guerrero, Sierra de las Aguas Escondidas, 7000 feet. Distr. — Mexico (Guerrero). Syntypes ♂, BMNH. N. COMB.
- BISETOSUS (Edwards), 1940: 442 (*Centrocnemis*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Ref. — Lane, 1946: 339, fig. 1 (♂ genitalia). Syntypes ♂ ♀, BMNH. N. COMB.
- BORORO (Lane), 1947: 341, fig. 8 (♂ genitalia) (*Centrocnemis*). Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP. N. COMB.
- GUARANI (Lane), 1947: 341, fig. 4 (♂ genitalia) (*Centrocnemis*). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Distr. — Brazil (Goiás, São Paulo, Santa Catarina). Type ♂, MZSP. N. COMB.
- GUAYANASI (Lane), 1947: 340, fig. 2 (♂ genitalia) (*Centrocnemis*). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo). Type ♂, MZSP. N. COMB.
- LENKOI (Lane), 1959: 105, fig. 12 (♂ genitalia) (*Australosymmerus*). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP. N. COMB.

PEDIFER (Edwards), 1940: 443 (*Centrocnemis*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Syntypes ♂ ♀, BMNH. N. comb.

TUPI (Lane), 1947: 341, fig. 3 (♂ genitalia) (*Centrocnemis*). Type-locality: Brazil, Goiás, Corumbá. Distr. — Brazil (Goiás, São Paulo), Argentina (Misiones). Type ♂, MZSP. N. comb.

Genus NEOCRIONISCA, new

Wing very similar to that of *Crionisca* Colless, from Australia (Colless, 1970: 95, pl. 1, fig. G), but pleurotergites bare.

Type-species, *Neocrionisca collessi*, sp. n.

NEOCRIONISCA COLLESSI, sp. n.

♀ — Head dark, pollinose, with scattered greyish hairs; labella reddish-yellow, the palpi lighter. First two antennal segments yellowish, flagellum dark brown.

Mesonotum with pale pollinose ground color, on which three wide longitudinal brown stripes, slightly shining, are superimposed: the median stripe beginning large on the anterior margin and posteriorly tapering but not reaching the hind margin; the lateral stripes with approximately the same width throughout, beginning distinctly before the middle of the mesonotum and running almost to the hind margin. Scutellum, postnotum and pleura brown.

Legs yellowish; front and middle tarsi brownish; setae black, tibial spurs dark brown.

Wings clear, with a brown patch from the base of Rs through ta (r-m) and another originating at basal half of R₁ and running and narrowing in width to Costa. Much paler patches may be seen somewhat distally on Rs, M₂, M₃ and M₄. Halteres brownish-yellow.

Abdominal tergites and sternites shiny, brown, with yellow band apically; the eighth tergite predominantly yellow, with brown basal band. Visible part of terminalia pale yellowish. Vestiture long, yellow.

Holotype ♀, ARGENTINA, Tierra del Fuego, iii.1948 (Budin), in the Museu de Zoologia, Universidade de São Paulo.

Genus NOTHOSYMMERUS, new

Can be immediately recognized by the following characters: only 2 scutellar bristles; anterior tibia without spines or bristles along its length; wings with both macro- and microtrichiac.

Type-species, *Platyura insolita* Walker.

INSOLITUS (Walker), 1837: 335 (*Platyura*). Type-locality: "Chile". Distr. — Argentina (Santa Cruz, Tierra del Fuego), Chile. Ref. — Freeman, 1951: 9, fig. 5 (♂ genitalia), 299 (wing). Type ♂, BMNH. N. comb.
incerta Bigot, 1888: 16 (*Ditomyia*). Type-locality: "Cape Horn". Type, MNHN.



Genus CALLICERATOMYIA Lane

- CALLICERATOMYIA Lane, 1946: 244. Type-species, *pectinata* Lane (orig. des.).
PECTINATA Lane, 1946: 244, fig. 1 (antenna), 2 (wing), 3 (♂ genitalia).
Type-locality: Brazil, São Paulo, Campos do Jordão, 1600 m.
Type ♂, MZSP.

Genus NERVIJUNCTA Marshall

- NERVIJUNCTA Marshall, 1896: 254, 265. Type-species, *nigrescens* Marshall (mon.).
CONJUNCTA (Freeman), 1951: 7, figs. 1 (♂ genitalia), 297 (wing) (*Ditomyia*).
Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
LAFFOONI Lane, 1952: 139 (nom. nov. for *marshalli* Lane). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo, Santa Catarina).
marshalli Lane, 1948: 250, figs. 5 (♂ genitalia), 6 (9th tergite), 7 (mesosome) (preocc. Edwards, in Tonnoir & Edwards, 1927).
Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
marshally Lane, 1952: 131, lapsus.

Genus RHIPIDITA Edwards

- RHIPIDITA Edwards, 1940: 411. Type-species, *fusca* Edwards (orig. des.).
FUSCA Edwards, 1940: 442. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, BMNH.
NIGRA Lane, 1952: 132, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.

UNPLACED DITOMYIIDAE

- MEXICANA Giglio-Tos, 1890: [3], pl. 2, fig. 7 (wing) (*Ditomyia*). Type-locality: Mexico, Orizaba. Type, IMZUT.
ZONATA Giglio-Tos, 1890: [2], pl. 2, fig. 6 (wing) (*Ditomyia*). Type-locality: Mexico, Orizaba. Type, IMZUT.
Ienis Johannsen, 1910: 229 (*Symmerus*; unnecessary nom. nov. for *zonata* Giglio-Tos).

ABBREVIATIONS

- BMNH — British Museum (Natural History), London
IMZUT — Istituto e Museo di Zoologia, Università di Torino
MNHN — Muséum National d'Histoire Naturelle, Paris
MZSP — Museu de Zoologia, Universidade de São Paulo



LITERATURE CITED

- LIGOT, J. M. F.
 1888. Diptères, pp. 1-45, pls. 1-4, in Ministère de la Marine et de l'Instruction Publique, *Mission scientifique au Cap Horn, 1882-1883*, 5 (Zoologie; 2me. Partie), Paris, "1891".
- COLLESS, D. H.
 1970. The Mycetophilidae (Diptera) of Australia. Part 1. Introduction, key to subfamilies, and review of Ditomyiinae. *J. Aust. Ent. Soc.* 9 (2): 83-99, 25 figs., 1 pl.
- EDWARDS, F. W.
 1940. New neotropical Mycetophilidae (IV) (Diptera). *Revta Ent., Rio de J.* 11 (1-2): 440-465, 8 figs.
- FREEMAN, P.
 1951. Mycetophilidae, in British Museum (Natural History), *Diptera of Patagonia and South Chile*, 3: 1-133, 49 pls. London.
 1954. Los insectos de las Islas Juan Fernandez. 13. Mycetophilidae, Sciaridae, Cecidomyiidae and Scatopsidae (Diptera). *Revta chil. Ent.* (1953) 3: 23-40, 2 pls., 1 fig. .
- GIGLIO-TOS, E.
 1890. Nuove specie di Ditteri del Museo Zoologico di Torino. II. *Boll. Mus. Zool. Anat. comp. R. Univ. Torino* 5 (84): 1-4, pl. 2.
- JOHANNSEN, O. A.
 1910. The fungus gnats of North America. The Mycetophilidae of North America. Part I. *Maine agric. Exp. Stn Bull.* (1909) (2) 172: 209-276, 3 pls. (i. e., figs. 46-107).
- LANE, J.
 1946. A new genus of Neotropical Mycetophilidae (Diptera, Nemocera). *Revta Ent.* 17 (1-2): 244-247, 3 figs.
 1947. New Brazilian Mycetophilidae (Diptera, Nemocera). *Revta Ent.* 17 (1946): 339-360, 14 figs.
 1948. "Mycetophilidae" do Brasil (Diptera, Nemocera). *Revta bras. Biol.* 8 (2): 247-254, 15 figs.
 1952. Some new Brazilian Mycetophilidae (Diptera, Nematocera). *Dusenja* 3 (2): 131-140, 5 figs.
 1959. Insecta Amapaensia. — Diptera: Mycetophilidae. *Studia ent.* 2 (1-4): 105-118, 14 figs.
- MARSHALL, P.
 1896. New Zealand Diptera: N° 2. — Mycetophilidac. *Trans. Proc. N. Z. Inst.* 28 [= N. S., 11]: 250-309, pls. 8-13.
- PHILIPPI, R. A.
 1865. Aufzählung der chilenischen Dipteren. *Verh. zool.-bot. Ges. Wien* 15: 595-782, pls. 23-29.



WALKER, F.

1837. Descriptions, etc., of the insects collected by Cpt. P. P. King, R. N., F. R. S., in the survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. Lond.* 17: 331-359.

WILLISTON, S. W.

1900. Supplement [part], pp. 217-248, in F. D. Godman & O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera*, 1: 378 pp., 6 pls. London.

INDEX

<i>Australomyia</i> Freeman, 1951	19B.2
<i>Australosymmerus</i> Freeman, 1954	19B.2
<i>Austrosymmerus</i> , gen. n.	19B.2
<i>bifasciatus</i> (Williston), 1900 (Plesiastina), <i>Austrosymmerus</i>	19B.2
<i>bisetosus</i> (Edwards), 1940 (Centrocnemis), <i>Austrosymmerus</i>	19B.2
<i>bivittatus</i> (Freeman), 1951 (<i>Australomyia</i>), <i>Australosymmerus</i> ..	19B.2
<i>bororo</i> (Lane), 1947 (Centrocnemis), <i>Austrosymmerus</i>	19B.2
<i>Calliceratomyia</i> Lane, 1946	19B.4
<i>Centrocnemis</i> Philippi, 1865	19B.2
<i>collessi</i> , sp.n., <i>Neocrionisca</i>	19B.3
<i>conjuncta</i> (Freeman), 1951 (<i>Ditomyia</i>), <i>Nervijuncta</i>	19B.4
<i>fusca</i> Edwards, 1940, <i>Rhipidita</i>	19B.4
<i>guarani</i> (Lane), 1947 (Centrocnemis), <i>Austrosymmerus</i>	19B.2
<i>guayanasi</i> (Lane), 1947 (Centrocnemis), <i>Austrosymmerus</i>	19B.2
<i>incerta</i> (Bigot), 1888 (<i>Ditomyia</i>), <i>Nothosymmerus</i>	19B.3
<i>insolitus</i> (Walker), 1837 (<i>Platyura</i>), <i>Nothosymmerus</i>	19B.3
<i>laffooni</i> Lane, 1952, <i>Nervijuncta</i>	19B.4
<i>lenis</i> Johannsen, 1910 (<i>Symmerus</i>)	19B.4
<i>lenkoi</i> (Lane), 1959 (<i>Australosymmerus</i>), <i>Austrosymmerus</i>	19B.2
<i>marshalli</i> Lane, 1948, <i>Nervijuncta</i>	19B.4
<i>marshalli</i> , error	19B.4
<i>mexicana</i> Giglio-Tos, 1890 (<i>Ditomyia</i>)	19B.4
<i>Neocrionisca</i> , gen. n.	19B.3
<i>Nervijuncta</i> Marshall, 1896	19B.4
<i>nigra</i> Lane, 1952, <i>Rhipidita</i>	19B.4
<i>Nothosymmerus</i> , gen. n.	19B.3
<i>pectinata</i> Lane, 1946, <i>Calliceratomyia</i>	19B.4
<i>pedifer</i> (Edwards), 1940 (Centrocnemis), <i>Austrosymmerus</i>	19B.3
<i>Rhipidita</i> Edwards, 1940	19B.4
<i>simplex</i> (Freeman), 1951 (<i>Australomyia</i>), <i>Australosymmerus</i>	19B.2
<i>stimagicus</i> (Philippi), 1865 (Centrocnemis), <i>Australosymmerus</i>	19B.2
<i>tupi</i> (Lane), 1947 (Centrocnemis), <i>Austrosymmerus</i>	19B.3
<i>zonata</i> Giglio-Tos, 1890 (<i>Ditomyia</i>)	19B.4

19B.6







MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
 (FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)
 A CATALOGUE OF THE DIPTERA OF THE AMERICAS
 SOUTH OF THE UNITED STATES

19 C

FAMILY KEROPLATIDAE

(CEROPLATIDAE, INCL. MACROCERIDAE)

N. PAPAVERO

Museu de Zoologia
Universidade de São Paulo


Keroplattid larvae spin mucus webs and some (*Arachnocampa* from New Zealand, *Orfelia fultoni* from North America) are luminescent. Larvae of *Keroplatus* are found in or on wood or on fungi, being primarily fungus-spore feeders. Those of *Macrocera* and *Orfelia* are found under rocks, in and under logs, or in soil around roots; larvae of *Orfelia* are primarily predatory; some species of *Macrocera* at least feed on both fungus spores and small animals. Those of *Neoditomyia* live in mucus webs under overhanging rocks and tree trunks, feeding on insects caught in these webs. Some species of *Orfelia* secrete oxalic acid droplets on the webs.

Distribution records include both published data and unpublished information obtained from Lane's collection, now in the Museu de Zoologia da Universidade de São Paulo. When the type repository is known, it is added to the respective species. Abbreviations used are:

- AMNH — American Museum of Natural History, New York.
 BMNH — British Museum (Natural History), London.
 CAS — California Academy of Sciences, San Francisco.
 IML — Instituto Miguel Lillo, Tucumán.
 MZSP — Museu de Zoologia, Universidade de São Paulo.
 STUT — Staatliches Museum für Naturkunde, Stuttgart-Ludwigsburg.
 USNM — United States National Museum, Washington, D. C.
 WIEN — Naturhistorisches Museum, Wien.

I am very much obliged to Dr. Loïc Matile (Muséum National d'Histoire Naturelle, Paris) for many suggestions and corrections.

19C.1



Subfamily KEROPLATINAE

Genus CEROTELION Rondani

- CEROTELION Rondani, 1856: 191. Type-species, *Platyura laticornis* Meigen (orig. des.) = *lineatus* (Fabricius).
Refs. — Lane, 1958b: 147 (key to species); Matile & Burghel-
-Balacescu, 1969: 82 (generic status).
- BORACENSIS (Lane), 1950a: 42 (*Ceroplatus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♀, MZSP. N. COMB.
- DURETI (Lane), 1958: 147, fig. 2 (♂ genitalia) (*Keroplatus*). Type-locality: Argentina, Buenos Aires, Buenos Aires. Type ♂, MZSP. N. COMB.
- ENDERLEINI (Lane), 1948: 443, fig. 4 (♂ genitalia) (*Ceroplatus*). Type-locality: Brazil, São Paulo, Juquiá. Type ♂, MZSP, N. COMB.
- FLAVICORNIS (Freeman), 1951: 20 (*Ceroplatus*). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile (Llanquihue, Chiloé). Type ♂, BMNH. N. COMB.
- FUNEREUS (Freeman), 1951: 19 (*Ceroplatus*). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Neuquén, Río Negro), Chile (Llanquihue). Type ♂, BMNH. N. COMB.
- NIGRICANS (Lane), 1948: 444 (*Ceroplatus*). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♀, MZSP. N. COMB.

Genus DOLICHODACTYLA Freeman

- DOLICHODACTYLA Freeman, 1951: 18. Type-species, *flavithorax* Freeman (orig. des.).
- FLAVITHORAX Freeman, 1951: 18, fig. 273 (wing). Type-locality: Chile, Llanquihue, Peulla. Type ♀, BMNH.

Genus EUCEROPLATUS Edwards

- Ceroplatus*, subg. EUCEROPLATUS Edwards, 1929: 174. Type-species, *notaticoxa* Senior-White (orig. des.).
Ref. — Matile, 1970: 779 (generic status).
- BELLULUS (Williston), 1900: 219, pl. 4, figs. 3 (whole insect), 3a (head) (*Ceroplatus*). Type-locality: Mexico, Tabasco, Teapa. Type ♂, BMNH. N. COMB.
- SINGULARIS (Lane), 1948: 445, fig. 10 (♂ genitalia) (*Ceroplatus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP. N. COMB.

Genus HETEROPTERNA Skuse

- HETEROPTERNA Skuse, 1888: 1129. Type-species, *macleayi* Skuse (mon.).
Refs. — Lane, 1948: 446 (key to spp.); Matile, 1971: 270 (generic status).
- ABDOMINALIS (Lane), 1948: 447, fig. 7 (♂ genitalia) (*Ceroplatus*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.

MAJOR (Curran), 1928: 29 (*Cerotelion*). Type-locality: "Jamaica". Ref. — Edwards, 1940: 447. Type, AMNH.

TETRALEUCA (Edwards), 1940: 446, pl. 18, fig. 3 (wing) (*Ceroplatus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Rio de Janeiro to Santa Catarina). Type ♂, BMNH.

TRILEUCA (Edwards), 1940: 447 (*Ceroplatus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Trinidad, Brazil (Ceará, Rio de Janeiro, Santa Catarina), Argentina (Jujuy). Type ♂, BMNH.

Genus ISONEUROMYIA Brunetti

ISONEUROMYIA Brunetti, 1912: 66. Type-species, *annandalei* Brunetti (mon.). Ref. — Lane, 1948 (partial key to spp.).

ARGENTEOTOMENTOSA (Kertész), 1909: 138, figs. 2a (♂ genitalia), 2b (wing) (*Platyura*). Type-locality: Peru, Callanga. Distr. — Peru, Brazil (Rio de Janeiro). Type ♂, lost. N. comb.

ATRA (Lane), 1948: 456, fig. 14 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.

BICINGULATA (Edwards), 1940: 448 (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Rio de Janeiro to Santa Catarina). Ref. — Lane, 1948: 453, fig. 11 (♂ genitalia). Type ♂, BMNH. N. comb.

BORINQUENT (Lane), 1950a: 49, fig. 10 (♂ genitalia) (*Platyura*). Type-locality: Costa Rica, San José. Type ♂, USNM. N. comb.

BREVINERVUS (Shaw), 1940: 804 (*Platyura*). Type-locality: "Costa Rica". N. comb.

ELEGANTULA (Williston), 1900: 218, pl. 4, fig. 2 (whole insect) (*Platyura*). Type-locality: Mexico, Guerrero, Omiltemi, 8000 feet. Type ♀, BMNH. N. comb.

FALCAOI (Lane), 1961b: 1, fig. 1 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Minas Gerais, Lagoa Santa (Gruta da Lapinha). Type ♂, MZSP. N. comb.

FLAVOFASCIATA (Edwards), 1940: 449 (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♀, BMNH. N. comb.

FORCIPATA (Kertész), 1909: 137, figs. 1a (♂ genitalia), 1b (wing) (*Platyura*). Type-locality: Peru, Callanga. Type ♂, lost. N. comb.

GOIANENSIS (Lane), 1948: 454, fig. 12 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Goiás, Corumbá. Type ♂, MZSP. N. comb.

GRISEOFASCIATA (Edwards), 1933: 304 (*Platyura*). Type-locality: Bolivia, La Paz, Mapi, Sarampiuni. Distr. — Bolivia, Brazil (Espírito Santo, Rio de Janeiro). Refs. — Lane, 1950a: 48; 1961b: 2. Type ♂, Dresden.

LEOMI Lane, 1959b: 110, fig. 3 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.

LENKOI Lane, 1961a: 246. Type-locality: Brazil, Amapá, Serra do Navio. Type ♀, MZSP.

- LOPESI (Lane), 1950a: 50 (*Platyura*). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro (Grajaú). Type ♀, MZSP.
- LUTEA (Freeman), 1951: 22 (*Platyura*). Type-locality: Chile, Chiloé, Castro. Distr. — Chile (Ñuble, Llanquihue, Chiloé). Type ♀, BMNH.
- NIGROFASCIATA (Freeman), 1951: 22 (*Platyura*). Type-locality: Chile, Valparaíso, Marga-Marga. Type ♂, USNM. N. comb.
- PAULISTANA (Lane), 1948: 455, fig. 13 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, Osasco. Type ♂, MZSP. N. comb.
- SESHIFORMIS (Edwards), 1940: 448 (*Platyura*). Type-locality: Santa Catarina, Nova Teutônia. Type ♀, BMNH. N. comb.
- TIMBIRA Lane, 1956: 128, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão (Cidade Azul). Type ♂, MZSP.
- TUCUMANA Lane, 1958b: 149, fig. 4 (♂ genitalia). Type-locality: Argentina, Tucumán, Lacavera. Type ♂, IML.
- XANTHOCERA (Edwards), 1931b: 256 (*Platyura*). Type-locality: Brazil, Pará. Distr. — Brazil (Pará, Goiás). Type ♀, BMNH. N. comb.

Genus KEROPLATUS BOSC

KEROPLATUS Bosc, 1792: 42. Type-species, *tipuloides* Bosc (mon.).

Ceroplatus, emend.

Ref. — Lane, 1958b: 145 (key to spp.).

- CARIBAI Lane, 1950a: 40, fig. 7 (♂ genitalia). Type-locality: Panama, Canal Zone, Barro Colorado I. Distr. — Panama, Trinidad. Type ♂, USNM.
- FIEBRIGI Edwards, 1934: 357. Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Paraguay, Argentina (Misiones). Syntypes ♂ ♀, WIEN.
- GOLBACHI Lane, 1958b: 145, fig. 1 (♂ genitalia). Type-locality: Argentina, Salta, Urundel. Type ♂, IML.
- MEXICANUS Lane, 1948: 437. Type-locality: Mexico, Oaxaca, Tuxtepec. Distr. — Mexico (Oaxaca), Panama. Type ♀, MZSP.
- TOWNSENDI Lane, 1948: 438, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Itaquaquecetuba. Distr. — Brazil (Pará, São Paulo, Santa Catarina). Type ♂, MZSP.

Genus MICRAPEMON Edwards

- MICRAPEMON Edwards, 1925: 523. Type-species, *Platyura parva* Williston (orig. des.).
- MAJUSCULUM (Edwards), 1940: 450 (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH. N. comb.
- PARVUM (Williston), 1896: 257 (*Platyura*). Type-locality: St. Vincent, W. I. Distr. — St. Vincent, Venezuela (Caracas), Brazil (Rio de Janeiro, São Paulo, Santa Catarina), Argentina (Tucumán). Refs. — Edwards, 1940: 450; Lane, 1950a: 55, fig. 15 (♂ genitalia). Syntypes ♂, BMNH.

Genus NEOCEROPLATUS Edwards, stat. n.

Ceroplastus, subg. NEOCEROPLATUS Edwards, 1941: 304. Type-species, *minimax* Edwards (orig. des.).

LAUROI (Lane), 1961: 243, fig. 1 (♂ genitalia) (*Keroplastus*). Type-locality: Brazil, Amapá, Rio Felício. Type ♂, MZSP. N. COMB.

MINIMAX (Edwards), 1934: 358 (*Ceroplastus*). Type-locality: Costa Rica, Farm La Caja, 8 km w. of San José. Distr. — Costa Rica, Peru (Loreto), Brazil (Goiás, São Paulo). Refs. — Edwards, 1941: 304; Lane, 1948: 440. Type ♂, Hamburg (lost). N. COMB.

PAICOENAI (Lane), 1950: 43, fig. 8 (♂ genitalia) (*Ceroplastus*; as *pacoenai*, in caption for fig., *lapsus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP. N. COMB.

Genus NEODITOMYA Lane & Sturm

NEODITOMYA Lane & Sturm, 1958: 199. Type-species, *colombiana* Lane & Sturm (orig. des.).

Refs. — Coher, 1963: 24 [footnote; synonymy with *Orfelina* (*Taulyrpa*)]; Sturm, 1973 (behavior of larvae).

AEROSPICATOR (Jackson), 1974: 240 (*Orfelina*). Type-locality: British Honduras, Cayo District, Blancaneaux Cave. Type ?♂, USNM. N. COMB.

ANDINA Lane & Sturm, 1958: 201, figs. 1 (♂ genitalia), 2 (larva), 3 (larval head), 4-8 (larva spinning a catching thread). Type-locality: Colombia, Sierra de la Macarena, s. of Villavicencio, Río Zanza, 400-500 m. Type ♂, MZSP.

COLOMBIANA Lane & Sturm, 1958: 204, figs. 9 (wing), 10 (♂ genitalia), 11 (pupa), 12 (larval head). Type-locality: Colombia, Resina (w. Andes, between Garzón and Florencia), 1600-2000 m. Type ♂, MZSP.

TROGOPHILA Matile, 1978: 369, figs. 1 (habitus), 2 (♂ genitalia), 6-11 (larvae). Type-locality: Cuba.

Genus ORFELIA Costa

Zelmira Meigen, 1800: 16. Type-species, *Platyura fasciata* Meigen (Coquillett, 1910: 621). Suppressed ICZN, 1963: 339.

ORFELIA Costa, 1857: 448. Type-species, *Platyura fasciata* Meigen (Hardy, 1960: 200).

Platyura, authors.

Subgenus APYRTULA Edwards

Platyura, subg. *Apyrtula* Edwards, 1940: 451, *nomen nudum*.

Platyura, subg. APYRTULA Edwards, 1941: 306. Type-species, *abbrevinervis* Edwards (orig. des.).

ABBREVINERVIS (Edwards), 1941: 306 (*Platyura*). Type-locality: Brazil, Mato Grosso, Salobra. Type ♂, BMNH. N. COMB.



SPATULATA (Edwards), 1940: 451 (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Syntypes ♂ ♀, BMNH.

Subgenus LOPYRUTA Edwards

Platyura, subg. LOPYRUTA Edwards, 1929: 172. Type-species, *fasciventris* Williston (orig. des.).

FASCIVENTRIS (Williston), 1896: 258, pl. 8, fig. 11 (wing) (*Platyura*). Type-locality: St. Vincent, W. I. Syntypes ♀, BMNH. N. comb.

MATHESONI (Lane), 1961b: 3 (*Platyura*). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♀, MZSP. N. comb.

Subgenus LYPRAUTA Edwards

Platyura, subg. LYPRAUTA Edwards, 1931a: 77. Type-species, *defecta* Edwards (orig. des.).

Ref. — Lane, 1950a: 63 (partial key to spp.).

ALTICOLA (Lane), 1948: 450, fig. 9 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo), Argentina (Salta, Catamarca, Buenos Aires). Type ♂, MZSP. N. comb.

AMAZONENSIS (Lane), 1958a: 211, fig. 3 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Amazonas, Manaus (Igarapé do Mariano). Type ♂, MZSP. N. comb.

CHACOENSIS (Edwards), 1931a: 77 (*Platyura*). Type-locality: Argentina, Chaco, Tacaaglé. Distr. — Brazil (Rio de Janeiro to Santa Catarina), Argentina (Chaco). Type ♀, STUT. N. comb.

DEFECTA (Edwards), 1931a: 77 (*Platyura*). Type-locality: Grenada (W. I.), Balthazar. Type ♂, BMNH. N. comb.

KNABI (Lane), 1950a: 64, fig. 20 (♂ genitalia) (*Platyura*). Type-locality: Costa Rica, Puntarenas, nr. Aramuz. Type ♂, USNM. N. comb.

NUBILAPEX (Edwards), 1940: 452 (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Trinidad, Brazil (Amapá, Rio de Janeiro, São Paulo, Santa Catarina). Syntypes ♂ ♀, BMNH. N. comb.

PARAENSIS (Lane), 1958a: 212, fig. 5 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Pará, Fordlândia, Rio Tapajós. Type ♂, MZSP. N. comb.

ZETEKI (Lane), 1950a: 54, fig. 21 (♂ genitalia) (*Platyura*). Type-locality: Panama, Canal Zone, Quarry Heights. Distr. — Panama, Trinidad. Type ♂, USNM. N. comb.

Subgenus NEOPLATYURA Malloch

NEOPLATYURA Malloch, 1928: 600, 601 (as genus). Type-species, *Platyura setiger* Johannsen (orig. des.).

Ref. — Lane, 1958b: 148 (key to spp.).

ACZELI (Lane), 1958b: 149, fig. 3 (♂ genitalia) (*Platyura*). Type-locality: Argentina, Salta, Tonono. Type ♂, IML. N. comb.



ANTUNESI (Lane), 1959b: 107 (*Platyura*). Type-locality: Brazil, Amapá, Serra do Navio. Type ♀, MZSP. N. comb.

IGNOBILIS (Williston), 1896: 257, pl. 8, fig. 9 (wing) (*Platyura*). Type-locality: St. Vincent, W. I. Distr. — St. Vincent, Panama to southern Brazil and Argentina (Misiones). Ref. — Lane, 1950a: 51, fig. 11 (♂ genitalia). Type ♂, BMNH. N. comb.

REGULARIS (Edwards), 1934: 355 (*Platyura*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Distr. — Brazil (Goiás, Rio de Janeiro, São Paulo). Type ♀, Halle. N. comb.

SAPARAI (Lane), 1950b: 140, fig. 2 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, São Paulo (Morumbi). Distr. — Trinidad, Brazil (Amapá, São Paulo). Type ♂, MZSP. N. comb.

Subgenus ORFELIA Costa

ANOMALA (Lane), 1960: 376, fig. 1 (♂ genitalia) (*Platyura*). Type-locality: Trinidad, Navy Base. Type ♂, MZSP. N. comb.

MAIAPENAI (Lane), 1950b: 143, fig. 4 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, São Paulo (Morumbi). Type ♂, MZSP. N. comb.

Subgenus PLAUTYRA Edwards

Platyura, subg. PLAUTYRA Edwards, 1941: 304. Type-species, *macilentata* Lynch Arribálzaga (orig. des.).

Ref. — Lane, 1950a: 65 (partial key to spp.).

AITKENI (Lane), 1961b: 4 (*Platyura*). Type-locality: Trinidad, Chaguaramas. Type ♂, MZSP. N. comb.

AUTUMNALIS (F. Lynch Arribálzaga), 1892: 433 (59) (*Platyura*). Type-locality: Argentina, Buenos Aires, Baradero and Las Conchas. Distr. — Brazil (Rio Grande do Sul), Argentina. N. comb.

BATESI (Edwards), 1931b: 256 (*Platyura*). Type-locality: Brazil, Pará. Type ♀, BMNH.

EURMEISTERI (Edwards), 1934: 255 (*Platyura*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Type ♀, Halle. N. comb.

INCASICA (Lane), 1950a: 68 (*Platyura*). Type-locality: Peru, Loreto, Iquitos. Distr. — Trinidad, Peru (Loreto), Brazil (Pará, São Paulo). Type ♀, USNM. N. comb.

MACILENTA (F. Lynch Arribálzaga), 1892: 432 (58) (*Platyura*). Type-locality: Argentina, Chaco, near Formosa. Distr. — Brazil (Goiás, São Paulo), Argentina. Ref. — Lane, 1948: 449. N. comb.

SALOBRENSIS (Edwards), 1941: 305, pl. 15, fig. 2 (wing) (*Platyura*). Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Trinidad, Panama, Brazil (Mato Grosso). Type ♂, BMNH. N. comb.

SHANNONI (Lane), 1948: 449 (*Platyura*). Type-locality: Brazil, Rio de Janeiro, Teresópolis. Type ♀, MZSP. N. comb.

TAMOYOI (Lane), 1950a: 66, fig. 22 (♂ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, Campos do Jordão, Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP. N. COMB.

Subgenus PSEUDOPLATYURA Skuse

PSEUDOPLATYURA Skuse, 1888: 1180 (as genus). Type-species, *dux* Skuse (mon.).

SOUZAI (Lane), 1959a: 184, fig. 2 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Minas Gerais, Serra do Cipó. Type ♂, MZSP. N. COMB.

Subgenus PYRATULA Edwards

Platyura, subg. PYRATULA Edwards, 1929: 167. Type-species, *zonata* Zetterstedt (orig. des.).

PARAGUAYANA (Edwards), 1934: 356 (*Platyura*). Type-locality: Paraguay, La Cordillera, San Bernardino, Distr. — Brazil (Goiás), Paraguay. Type ♂, WIEN. N. COMB.

Subgenus PYRTAULA Edwards

Platyura, subg. PYRTAULA Edwards, 1929: 166. Type-species, *agricolae* Marshall (orig. des.).

BIGUTTATA (Freeman), 1951: 23 (*Platyura*). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Chubut), Chile (Llanquihue). Type ♂, BMNH. N. COMB.

CONSIMILIS (Freeman), 1951: 25 (*Platyura*). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH. N. COMB.

DELPONTEI (Lane), 1963: 4, fig. 1 (♂ genitalia) (*Platyura*). Type-locality: Argentina, Tierra del Fuego, Lago Fagnano. Type ♂, MZSP. N. COMB.

FERRUGINEA (Freeman), 1951: 25 (*Platyura*). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Chubut), Chile (Llanquihue, Chiloé). Type ♂, BMNH. N. COMB.

INAEQUALIS (Freeman), 1951: 24 (*Platyura*). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro), Chile (Chiloé). Type ♂, BMNH. N. COMB.

INFUMATA (Freeman), 1951: 23 (*Platyura*). Type-locality: Chile, Chiloé, Puerto Montt. Type ♀, EMNH. N. COMB.

QUASILINEATA (Edwards), 1919: 144, fig. 5 (wing) (*Platyura*). Type-locality: Ecuador, El Angel, 3020m. Type ♀, Mus. Nat. Hist. Nat., Paris (Matile exam.). N. COMB.

TRIANGULIFERA (Freeman), 1951: 25 (*Platyura*). Type-locality: Chile, Chiloé, Puerto Montt. Distr. — Chile (Llanquihue, Chiloé). Type ♂, BMNH. N. COMB.



Subgenus RYPATULA Edwards

Platyura, subg. RYPATULA Edwards, 1929: 167. Type-species, *brevis* Tonnoir (orig. des.).

CHILENSIS (Freeman), 1951: 21 (*Platyura*). Type-locality: Chile, Llanquihuc, Puculla. Type ♂, BMNH. N. COMB.

Subgenus TAULYRPA Edwards

Platyura, subg. TAULYRPA Edwards, 1929: 171. Type-species, *Cerotelion vespiformis* Enderlein (orig. des.).

AMAPAENSIS (Lane), 1959b: 108, fig. 2 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP. N. COMB.

MONOCHROMA (Edwards), 1933: 304 (*Platyura*). Type-locality: Peru, Pasco, Pichis, Puerto Bermúdez. Distr. — Peru (Pasco), Brazil (São Paulo). Ref. — Lane, 1950a: 56, fig. 16 (♂ genitalia). Type ♂, Dresden. N. COMB.

VESPIFORMIS (Enderlein), 1910: 147 (*Cerotelion*). Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Goiás, Santa Catarina). Ref. — Lane, 1950a: 55, fig. 14 (♂ genitalia). Type ♂, Stettin. N. COMB.

Subgenus XENOPLATYURA Malloch

XENOPLATYURA Malloch, 1928: 600, 601 (as genus). Type-species, *Platyura conformis* Skuse (orig. des.).
Ref. — Lane, 1961b: 7 (key to spp.).

BICELLII (Lane), 1961a: 245 (*Platyura*). Type-locality: Brazil, Amapá, Rio Felício. Type ♀, MZSP. N. COMB.

COHERI (Lane), 1950b: 142, fig. 3 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Goiás, Corumbá. Distr. — Trinidad, Brazil (Goiás), Argentina (Tucumán). Type ♂, MZSP. N. COMB.

GALINDOI (Lane), 1961b: 7, fig. 4 (♀ terminalia) (*Platyura*). Type-locality: Panama, Tapagra. Type ♀, MZSP. N. COMB.

MANAOSSENSIS (Lane), 1961b: 9 (*Platyura*). Type-locality: Brazil, Amazonas, Manaus (Igarapé do Francês). Type ♂, MZSP. N. COMB.

TAPUIAI (Lane), 1950a: 53, fig. 13 (♂ genitalia) (*Platyura*). Type-locality: Brazil, Minas Gerais, Lassance. Distr. — Trinidad, Brazil (Pará, Minas Gerais, Rio de Janeiro, São Paulo), Argentina (Chaco). Type ♂, MZSP. N. COMB.

Genus PLACOCERATIAS Enderlein

PLACOCERATIAS Enderlein, 1910: 149. Type-species, *bimaculipennis* Enderlein (orig. des.).

BARRETOI (Lane), 1948: 440, fig. 2 (♂ genitalia) (*Ceroplastus*). Type-locality: Brazil, Goiás, Corumbá. Distr. — Brazil (Goiás, Minas Gerais). Type ♂, MZSP. N. COMB.

BIMACULIPENNIS Enderlein, 1910: 149. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Mato Grosso, Santa Catarina). Type ♂, Stettin.



- FLUCITHORAX Enderlein, 1910: 152. Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.
- IMITANS (Lane), 1948: 442, fig. 3 (♂ genitalia) (*Ceroplastus*). Type-locality: Brazil, São Paulo, Porto Cabral. Type ♂, MZSP. N. comb.
- LONGIMANUS (Williston), 1896: 258, pl. 8, fig. 12 (wing) (*Ceroplastus*). Type-locality: St. Vincent, W. I. Distr. — St. Vincent, Trinidad, Panama, Peru. Type ♂, BMNH. N. comb.
- UARACUI (Lane), 1950a: 46, fig. 9 (♂ genitalia) (*Ceroplastus*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo), Argentina (Tucumán). Type ♂, MZSP. N. comb.

Genus PLATYROPTILON Westwood

- Platyura*, subg. PLATYROPTILON Westwood, 1849: 231. Type-species, *miersii* Westwood (mon.).
- Ref. — Duret, 1975 (key to spp.).
- DURETI Lane, 1956: 122, figs. 1 (wing), 2 (♂ genitalia). Type-locality: Brazil, Minas Gerais, Juiz de Fora. Type ♂, MZSP.
- MIERSII (Westwood), 1849: 231, pl. 23, figs. 3 (whole insect), 3a-b (head), 3c (antenna), 3d (wing), 3e-f (legs) (*Platyura*). Type-locality: "Brazil". Distr. — Brazil (São Paulo), Argentina (Tucumán, Misiones, Buenos Aires). Ref. — Duret, 1975: 291, figs. 1-5 (wing, claw, ♂ genitalia).
- MISIONENSIS Duret, 1975: 295, figs. 8-11 (antenna, wing, ♂ genitalia). Type-locality: Argentina, Misiones, San Ignacio, Arroyo Barra Bonita. Type ♂, Coll. Duret.
- RAMICORNIS Enderlein, 1910: 145. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro to Santa Catarina). Refs. — Edwards, 1934: 360; Lane, 1947: 345; Duret, 1975: 294, figs. 6-7 (♂ genitalia).
- ZERNYI Edwards, 1934: 358. Type-locality: Brazil, Pará, Belém. Distr. — Peru, Brazil (Pará). Ref. — Lane, 1962a: 10, fig. 1 (♂ genitalia). Type ?♀, WIEN.

Genus PROCEROPLATUS Edwards


- PROCEROPLATUS Edwards, 1925: 523. Type-species, *Platyura pictipennis* Williston (orig. des.).
- Calliplatyura* Malloch, 1928: 600 (as *Calloplatyura*, p. 601). Type-species, *Platyura elegans* Coquillett (orig. des.). Refs. — Lane, 1956 (key); Matile, 1973: 923 (generic status).
- BORGMEIERI (Shaw), 1940: 803 (*Platyura*). Type-locality: Costa Rica, San Mateo. N. comb.
- CATHARINAE (Edwards), 1932: 139 (*Platyura*). Type-locality: Brazil, Santa Catarina, Bom Retiro. Distr. — Brazil (Goiás, Rio de Janeiro, São Paulo, Santa Catarina). Type ♂, BMNH. N. comb.

- GUAYANASI (Lane), 1950a: 59, fig. 17 (δ genitalia) (*Platyura*). Type-locality: Brazil, São Paulo, São Paulo (Santo Amaro). Type δ , MZSP. N. comb.
- IAUNAI (Lane), 1956: 126, figs. 4-5 (wing, δ genitalia) (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Type δ , BMNH. N. comb.
- KERTESZI (Lane), 1956: 125 (*Platyura*; nom. nov. for *elegans* Kertész). Type-locality: Peru, Vilcanota. Distr. — Peru (Loreto), Brazil (Amazonas, Pará, Amapá). Refs. — Lane, 1956: 125, fig. 2 (wing); 1959: 108, fig. 1 (δ genitalia). N. comb.
- elegans* Kertész, 1901: 404, pl. 20, fig. 2 (wing) (*Platyura*; preocc. Coquillett, 1895). Type-locality: Peru, Vilcanota. Type δ , lost.
- PARAMARIBOENSIS (Edwards), 1934: 356, fig. 1 (wing) (*Platyura*). Type-locality: Surinam, Paramaribo. Distr. — Costa Rica, Trinidad, Surinam. Type φ , WIEN. N. comb.
- PICTIPENNIS (Williston), 1896: 257, pl. 8, fig. 10 (wing) (*Platyura*). Type-locality: St. Vincent, W. I. Distr. — St. Vincent, Trinidad. Ref. — Edwards, 1925: 523. Syntypes δ φ , BMNH. N. comb.
- FUNCTICOXALIS (Edwards), 1940: 451, pl. 18, fig. 4 (wing) (*Platyura*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Amazonas, Amapá, São Paulo, Santa Catarina), Argentina (Chaco). Syntypes δ φ , BMNH. N. comb.
- RABELLOI (Lane), 1956: 127, fig. 6 (wing) (*Platyura*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Peru (Tingo Maria), Brazil (São Paulo). Type φ , MZSP. N. comb.
- STONEI (Lane), 1950a: 60, fig. 18 (δ genitalia) (*Platyura*). Type-locality: "Jamaica". Type δ , USNM. N. comb.
- TERENOI (Lane), 1950a: 61, fig. 19 (δ genitalia) (*Platyura*). Type-locality: Brazil, Mato Grosso. Ref. — Lane, 1956: 127, fig. 7 (wing). Type δ , MZSP. N. comb.
- TRINIDADENSIS (Lane), 1960: 379, fig. 3 (δ genitalia) (*Platyura*). Type-locality: Trinidad, Navy Base. Type δ , MZSP. N. comb.
- VARIVENTRIS (Edwards), 1931a: 76 (*Platyura*). Type-locality: Bolivia, Villa Montes. Type φ , STUT.
- VENUSTIPENNIS (Shaw), 1940: 804 (*Platyura*). Type-locality: Costa Rica, San Mateo. N. comb.
- VILLASBOASI (Lane), 1961b: 5, fig. 2 (wing), 3 (δ genitalia) (*Platyura*). Type-locality: Brazil, Pará, Cachimbo. Type δ , MZSP. N. comb.
- VITTATUS (Fisher), 1941: 8 (*Platyura*). Type-locality: Costa Rica. N. comb.

Subfamily MACROCERINAE

Genus MACROCERA Meigen

Euphrosyne Meigen, 1800: 16. Type-species, *Macrocera lutea* Meigen (Coquillett, 1910: 542). Suppressed ICZN, 1963: 339.



- MACROCERA Meigen, 1803: 261. Type-species, *lutea* Meigen (Curtis, 1837: pl. 637).
- ANTENNATA Freeman, 1951: 15, fig. 272 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Santiago, Concepción, Llanquihue, Chiloé). Type ♂, BMNH.
- ARMATA Freeman, 1951: 17. Type-locality: Argentina, Tierra del Fuego, Lago Yuvín. Type ♂, BMNH.
- CHILENA Lane, 1962b: 528, fig. 1 (wing). Type-locality: Chile, Llanquihue, Los Muermos. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, CAS.
- CONCINNA Williston, 1896: 255, pl. 8, fig. 7 (wing). Type-locality: St. Vincent, W. I. Syntypes ♂ ♀, BMNH.
- FLAVESCENS Freeman, 1951: 15. Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- FLAVITHORAX Freeman, 1951: 16. Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.
- FLAVOBRUNNEA Freeman, 1951: 14. Type-locality: Chile, Chiloé, Puerto Montt. Type ♂, BMNH.
- FUMIDAPEX Freeman, 1954: 25. Type-locality: Juan Fernandez Is., Masafuera, Quebrada de las Casas, 1800 m. Type ♂, Universidad de Chile, Santiago.
- FUNEREA Freeman, 1951: 14. Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Juan Fernandez Is. (Masatierra), Chile (Llanquihue, Chiloé), Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- GUAIANASI Lane, 1950a: 36. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♀, MZSP.
- GUARANI Lane, 1950a: 37, figs. 3 (wing), 4 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Goiás, São Paulo, Paraná). Type ♂, MZSP.
- HORRIDA Freeman, 1951: 15. Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro, Chubut), Chile (Llanquihue). Type ♂, BMNH.
- INAEQUALIS Freeman, 1951: 16. Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Chiloé). Type ♂, BMNH.
- KAINGANGI Lane, 1950a: 34, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- MATILLI, nom. nov. for *nobilis* Edwards. Type-locality: Colombia, Cordillera Central, Paso del Quindín, 3000 m.
- nobilis* Edwards, 1934: 354 (preocc. Johnson, 1922). Type-locality: Colombia, Cordillera Central, Paso del Quindín, 3000 m. Type ♀, WIEN.
- PERPICTULA Edwards, 1940: 445, pl. 18, fig. 2 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina (Misiones). Type ♂, EMNH.



- PICTA Freeman, 1951: 13. Type-locality: Chile, Chiloé, Puntra. Type ♂, BMNH.
- PLAUMANNI Edwards, 1940: 445. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Golás to Rio de Janeiro and Santa Catarina), Argentina (Tucumán, Corrientes, Misiones). Ref. — Lane, 1950a: 35, fig. 2 (♂ genitalia). Syntypes ♂ ♀, BMNH.
- SHANNONI Lane, 1947: 344. Type-locality: Brazil, Mato Grosso, Maracaju. Type ♂, MZSP.
- TAMOYOI Lane, 1950a: 39, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.
- TESTACEA Philippi, 1865: 617. Type-locality: "Chile".
- THOMSONI E. Lynch Arribálzaga, 1882: 116 (non. nov. for *fascipennis* Thomson). Type-locality: "Patagonia".
- fascipennis* Thomson, 1869: 448 (preocc. Staeger, 1840). Type-locality: "Patagonia".
- UNICA Fisher, 1939: 227, pl. 13, figs. 1-2 (wing, tip of fore tibia). Type-locality: Costa Rica, San José. Type ♂, Academy of Natural Sciences, Philadelphia.
- UNIDENS Edwards 1931b: 255. Type-locality: Peru, Lima, San Bartolomé. Type ♂, BMNH.
- VALDIVIANA Philippi, 1865: 617. Type-locality: Chile, Valdivia, Corral.

Genus PARAMACROCERA Edwards

- PARAMACROCERA Edwards, 1927: 778. Type-species, *brevicornis* Edwards (orig. des.)
- ANOMALA Freeman, 1951: 18. Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.

Unplaced Keroplatidae

- MIRA Tarwid, 1936: 40, fig. 8 (♂ genitalia) [*Zelmira* (*Neoplatyura*)]. Type-locality: ?Peru (Coll. Jelski).
- OBSCURUS Philippi, 1865: 618, pl. 23, fig. 8 (whole insect) (*Ceroplatus*). Type-locality: Chile, Valdivia.
- RUBENS Wiedemann, 1828: 60 (*Platyura*). Type-locality: "Brazil".
- SUBANNULATA Philippi, 1865: 620 (*Platyura*). Type-locality: "Chile".

LITERATURE CITED

- BOSCH, L. A. G.
 1792. *Keroplatus*. *Actes Soc. Hist. Nat. Paris* 1: 42-43, 1 fig.



BRUNETTI, E.

1912. Diptera Nematocera (excluding Chironomidae and Culicidae), in [W. T. Blanford, ed.] *Fauna of British India. Diptera* [Vol. 1], 581 pp., 44 figs., 12 pls. London.

COHER, E. I.

1963. Asian *Macrocera* Meigen, 1803 (Diptera: Mycetophilidae), with remarks on the status of the genus and related genera. *Bull. Brooklyn Ent. Soc.* (n.s.) 58 (1): 23-36, 2 pls.

COQUILLET, D. W.

1910. The type-species of the North American genera of Diptera. *Proc. U. S. Natl. Mus.* 37: 499-647.

COSTA, A.

1857. Contribuzione alla fauna ditterologica italiana. [Giambattista Vico] *Giorn. Sci.*, Napoli 2: 438-460.

CURRAN, C. H.

1928. In C. C. Gowdey, *Catalogus insectorum jamaicensis*. *Ent. Bull. Dept. Agric. Jamaica*, Kingston 4: 44 + 3 pp., 2 figs.

CURTIS, J.

1837. *British entomology: Being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 13: pls. 626-673. London.

DURET, J. P.

1975. Notas sobre el genero *Platyroptilon* Westwood, 1849 (Diptera, Mycetophilidae). *Rev. Soc. Ent. Argent.* (1973-1974) 34 (3-4): 289-297, 11 figs.

EDWARDS, F. W.

1919. Diptères, Nématocères. *Miss. Arc MÉR. AmÉR. Sud*, Paris 10: 143-162, figs.
1925. British fungus-gnats (Diptera, Mycetophilidae). With a revised generic classification of the family. *Trans. Ent. Soc. London* 1925: 505-662, pls. 49-61.
1927. In A. L. Tonnoir & F. W. Edwards, *New Zealand fungus gnats* (Diptera, Mycetophilidae). *Trans. Proc. N. Z. Inst.* 57: 747-878, pls. 58-80.
1929. Notes on the Ceroplatinae, with descriptions of new Australian species (Diptera, Mycetophilidae). *Proc. Linn. Soc. N. S. W.* 54: 162-175, 4 figs.
- 1931a. Die Ausbeute der deutschen Chaco-Expedition 1925/26. — Diptera. XXIII. Bibionidae and XXIV. Mycetophilidae. *Konowia* 10 (1): 75-78.



- 1931b. New neotropical nematoceros Diptera. *Ann. Mag. Nat. Hist.* (10) 7: 255-261.
1932. New Brazilian Mycetophilidae (Diptera). *Rev. Ent.* 2 (2): 138-149, 6 figs.
1933. New neotropical Mycetophilidae (II) (Dipt.). *Rev. Ent.* 3 (3): 303-322.
1934. New neotropical Mycetophilidae (III). *Rev. Ent.* 4 (3): 354-372, 5 figs.
1940. New neotropical Mycetophilidae (IV) (Diptera). *Rev. Ent.* 11 (1-2): 440-465, 2 pls.
1941. Mycetophilidae (Diptera) collected by the expedition to Mato Grosso of the Brazilian Zoological Club, in July 1939. *Rev. Ent.* 12 (1-2): 303-314, pl. 15.

ENDERLEIN, G.

1910. Neue Gattungen und Arten aussereuropäischer Fliegen. *Stettin. ent. Ztg* (1911) 72: 135-209, 4 figs.

FISHER, E. G.

1939. Costa Rican Mycetophilidae (Diptera). *Trans. Amer. Ent. Soc.* 65: 227-236, 1 pl.
1941. Notes on Costa Rican Mycetophilidae (Diptera). *Ent. News* 52: 8.

FREEMAN, P.

1951. Mycetophilidae, in *British Museum (Natural History), Diptera of Patagonia and South Chile 3*: 1-138, 355 figs. London.
1954. Los insectos de las islas Juan Fernandez. 13. Mycetophilidae, Sciaridae, Cecidomyiidae and Scatopsidae (Diptera). *Rev. Chil. Ent.* (1953) 3: 23-40, 3 pls.

HARDY, D. E.

1960. Diptera: Nematocera-Erachycera (except Dolichopodidae), in E. C. Zimmermann, ed., *Insects of Hawaii 10*: 368 pp., 120 figs. Honolulu.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.* 20: 339-342.

JACKSON, J. F.

1974. Goldschmidt's dilemma resolved: Notes on the larval behavior of a new neotropical web-spinning mycetophilid. *Amer. Midl. Nat.* 92 (1): 240-245, 4 figs.

KERTÉSZ, K.

1901. Neue und bekannte Dipteren in der Sammlung des Ungarischen National-Museums. *Termész. Füzetek* 24: 403-432, pl. 20.



1909. Vier neue südamerikanische Mycetophiliden. *Ann. Mus. Nat. Hung.* 7: 137-142, 2 figs.

LANE, J.

1947. New Brazilian Mycetophilidae (Diptera, Nematocera). *Rev. Ent.* (1946) 17 (3): 339-360, 14 figs.
1948. Ceroplatinae da região neotropical (Diptera, Mycetophilidae). *Rev. Ent.* 19 (3): 437-458, 14 figs.
- 1950a. Neotropical Ceroplatinae (Diptera, Mycetophilidae). *Dusenía* 1 (1): 32-69, 22 figs.
- 1950b. Additional neotropical Ceroplatinae (Diptera, Mycetophilidae). *Dusenía* 1 (3): 139-144, 4 figs.
1956. Further notes on neotropical "Keroplatinae" (Diptera, Mycetophilidae). *Rev. Bras. Biol.* 16 (1): 121-128, 8 figs.
- 1958a. On Amazonian Mycetophilidae (Dipt., Nematocera). *Studia Ent.* 1 (1-2): 209-216, 8 figs.
- 1958b. Mycetophilidae, chiefly from Argentina. *Proc. 10th Int. Congr. Ent.* (Montreal, 1956) 1: 143-162, 17 figs.
- 1959a. Note on neotropical "Mycetophilidae" (Diptera, Nematocera). *Rev. Bras. Biol.* 19 (2): 183-190, 5 figs.
- 1959b. Insecta Amapaensia. — Diptera: Mycetophilidae. *Studia Ent.* 2 (1-4): 105-118, 14 figs.
1960. Mycetophilidae from Trinidad, B. W. I. (Diptera, Nematocera). *Studia Ent.* 3 (1-4): 375-384, 9 figs.
- 1961a. Insecta Amapaensia. — Diptera: Mycetophilidae. (Second contribution). *Studia Ent.* 4: 243-256, 7 figs.
- 1961b. Further new neotropical Mycetophilidae (Diptera, Nematocera). *Rev. Bras. Ent.* 10: 1-15, 5 figs.
- 1962a. A brief note on neotropical Mycetophilidae (Diptera, Nematocera). *Rev. Bras. Biol.* 22 (1): 9-16, 4 figs.
- 1962b. Mycetophilidae from Chile (Dipt.). *Studia Ent.* 5: 527-548, 8 figs.
1963. Insecta Patagonica (Diptera — Mycetophilidae). *Rev. Soc. Cient. Argent.* (1962) 25 (1-4): 3-16, 3 figs.

LANE, J. & H. STURM

1958. A new genus of "Ditomyiinae". Description of two new species with bionomical notes (Diptera, Mycetophilidae). *Rev. Bras. Biol.* 18 (2): 199-207, 12 figs.

LYNCH ARIBÁLZAGA, E.

1882. Catálogo de los dípteros hasta ahora descritos que se encuentran en las Repúblicas del Río de la Plata. *Bol. Acad. Nac. Cienc. Córdoba* 4: 109-152.

LYNCH ARIBÁLZAGA, F.

1892. Dipterología argentina: Mycetophilidae. *Bol. Acad. Nac. Cienc. Córdoba* 12: 377-436, 2 pls. [Also separately published, 1892, 75 pp., 2 pls., Buenos Aires].

19C.16



MALLOCH, J. R.

1928. Notes on Australian Diptera. N° xvii. *Proc. Linn. Soc. N. S. W.* 53: 598-617, 12 figs.

MATILE, L.

1970. Diptères Mycetophilidae du Cameroun et de République Centrafricaine. I. Keroplatinae. *Bull. Inst. Fond. Afr. Noire (A)* 32: 773-816, 61 figs.
1971. Caractères écologiques et morphologiques des larves de Mycetophilidae (Diptera) au niveau supraspécifique. *Proc. XIII Int. Congr. Ent.* (Moscow, 1968) 1: 270.
1973. Diptères Mycetophilidae du Cameroun et de République Centrafricaine. IV. Keroplatinae (2e. note). *Bull. Inst. Fond. Afr. Noire (A)* 35: 917-946.
1978. Un keroplatine cavernicole de Cuba (Dipt., Mycetophilidae). In: Acad. Sci. Rep. Soc. Roumanie, *Résultats des explorations biospéologiques cubano-roumaines à Cuba*, vol. 2. Bucarest.

MATILE, L. & A. BURCHELE-BALACESCU

1969. Note sur les genres *Keroplatus* et *Cerotelion* et description de *Cerotelion racovitzai* n. sp. (Dipt. Mycetophilidae). *Bull. Soc. Ent. France* 74: 82-86, 2 figs.

MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp. Paris.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2: 259-281.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool.-bot. Ges. Wien* 15: 595-782, 7 pls.

RONDANI, C.

1856. *Dipterologiae Italicae prodromus. Vol. 1: Genera Italica ordinis dipterorum ordinatim disposita et distincta et in familias et stirpes aggregata*, 228 pp. Parma.

SHAW, F.

1940. Some new Mycetophilidae from Costa Rica (Dipt.). Part I. *Rev. Ent.* 11 (3): 803-808, 8 figs.

SKUSE, A. A.

1888. Diptera of Australia. *Proc. Linn. Soc. N. S. W.* (2) 3: 17-145, pls. 1-2; 657-726, pl. 11; 1123-1222, pls. 31-32.

STURM, H.

1973. Fanggespinste und Verhalten des Larves von *Neoditomyia andina* und *colombiana* Lane (Diptera, Mycetophilidae). *Zool. Anz.* 191 (1-2): 61-68.



TARWID, K.

1936. Notatka w sprawie tożsamości systematycznej rodzajów: *Zelmira* Meig. (= *Platyura* Meig.) i *Asindulum* Latr. (Diptera, Fungivoridae). [Note sur l'identité systématique des genres: *Zelmira* Meig. (= *Platyura* Meig.) et *Asindulum* Latr.]. *Sprawozd. z posiedzen Towarzystwa Naukowego Warszawsk.* [C. R. Soc. Sci. Varsovie] (4, Nauk biol.) 29 (1-3): 35-44, 10 figs.

WESTWOOD, J. O.

1849. Diptera nonnulla exotica descripta. *Trans. Ent. Soc. London* 5: 231-236, pl. 23 (figs. 3-9).

WIEDEMANN, C. R. W.

1828. *Aussereuropäische zweiflügelige Insekten* 1: xxxii + 608 pp., 7 pls. Hamm.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896: 253-439, pls. 8-14.
1900. Supplement [part], pp. 217-248, in F. D. Godman & O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera* 1: 378 pp., 6 pls. London.

INDEX

abbrevinervis (Edwards), 1941 (Platyura), Orfelia (Apyrtula)	19C.5
abdominalis (Lane), 1948 (Ceroplatus), Heteropterna	19C.2
aczeli (Lane), 1958 (Platyura), Orfelia (Neoplatyura)	19C.6
aerospicator (Jackson), 1974 (Orfelia), Neoditomyia	19C.5
aikeni (Lane), 1961 (Platyura), Orfelia (Platyura)	19C.7
alticola (Lane), 1948 (Platyura), Orfelia (Lyprauta)	19C.6
amapaensis (Lane), 1959 (Platyura), Orfelia (Taulyrpa)	19C.9
amazonensis (Lane), 1958 (Platyura), Orfelia (Lyprauta)	19C.6
andina Lane & Sturm, 1958, Neoditomyia	19C.5
anomala (Lane), 1960 (Platyura), Orfelia (Orfelia)	19C.7
anomala Freeman, 1951, Paramacrocera	19C.13
antennata Freeman, 1951, Macrocera	19C.12
antunesi (Lane), 1959 (Platyura), Orfelia (Neoplatyura)	19C.7
<i>Apyrtula</i> Edwards, 1940	19C.5
<i>Apyrtula</i> Edwards, 1941	19C.5
argenteotomentosa (Kertész), 1909 (Platyura), Isoneuromyia	19C.3
armata Freeman, 1951, Macrocera	19C.12
atra (Lane), 1948 (Platyura), Isoneuromyia	19C.3
autumnalis (Lynch Arribálzaga), 1890 (Platyura), Orfelia (Platyura)	19C.7
barrettoi (Lane), 1948 (Ceroplatus), Placoceratias	19C.9
batesi (Edwards), 1931 (Platyura), Orfelia (Platyura)	19C.7
bellulus (Williston), 1900 (Ceroplatus), Euceroplatus	19C.2
bicellii (Lane), 1961 (Platyura), Orfelia (Xenoplatyura)	19C.9

19C.18



bicingulata (Edwards), 1940 (Platyura), Isoneuromyia	19C.3
biguttata (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
bimaculipennis Enderlein, 1910, Placoceratias	19C.9
boracensis (Lane), 1950 (Ceroptatus), Cerotelion	19C.2
borgmeieri (Shaw), 1940 (Platyura), Proceroplatus	19C.10
borinqueni (Lane), 1950 (Platyura), Isoneuromyia	19C.3
brevinervus (Shaw), 1940 (Platyura), Isoneuromyia	19C.3
burmeisteri (Edwards), 1934 (Platyura), Orfelia (Platyura)	19C.7
<i>Calliplatyura</i> Malloch, 1928	19C.10
<i>Calloplatyura</i> , lapsus	19C.10
caribai Lane, 1950, Keroplatus	19C.4
catharinae (Edwards), 1932 (Platyura), Proceroplatus	19C.10
<i>Ceroptatus</i> , emend.	19C.4
Cerotelion Rondani, 1856	19C.2
chacocensis (Edwards), 1931 (Platyura), Orfelia (Lyprauta)	19C.6
chilena Lane, 1962, Macrocera	19C.12
chilensis (Freeman), 1951 (Platyura), Orfelia (Rypatula)	19C.9
coheri (Lane), 1950 (Platyura), Orfelia (Xenoplattyura)	19C.9
colombiana Lane & Sturm, 1958, Neoditomyia	19C.5
concinna Williston, 1896, Macrocera	19C.12
consimilis (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
defecta (Edwards), 1931 (Platyura), Orfelia (Lyprauta)	19C.6
delpontei (Lane), 1963 (Platyura), Orfelia (Pyrtaula)	19C.8
Dolichodactyla Freeman, 1951	19C.2
dureti (Lane), 1958 (Keroplatus), Cerotelion	19C.2
dureti Lane, 1956, Platyroptilon	19C.10
<i>elegans</i> (Kertész), 1901 (Platyura), Proceroplatus	19C.11
elegantula (Williston), 1900 (Platyura), Isoneuromyia	19C.3
enderleini (Lane), 1948 (Ceroptatus), Cerotelion	19C.2
Euceroptatus Edwards, 1929	19C.2
<i>Euphrosyne</i> Meigen, 1800	19C.11
falcaoi (Lane), 1961 (Platyura), Isoneuromyia	19C.3
fascipennis Thomson, 1869, Macrocera	19C.13
fasciventris (Williston), 1896 (Platyura), Orfelia (Lapyrta)	19C.6
ferruginea (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
fielbrigi Edwards, 1934, Keroplatus	19C.4
flavescens Freeman, 1951, Macrocera	19C.12
flavicornis (Freeman), 1951 (Ceroptatus), Cerotelion	19C.2
flavithorax Freeman, 1951, Dolichodactyla	19C.2
flavithorax Freeman, 1951, Macrocera	19C.12
flavobrunnea Freeman, 1951, Macrocera	19C.12
flavofasciata (Edwards), 1940 (Platyura), Isoneuromyia	19C.3
forcipata (Kertész), 1909 (Platyura), Isoneuromyia	19C.3
fumidapex Freeman, 1954, Macrocera	19C.12
funerea Freeman, 1951, Macrocera	19C.12
funereus (Freeman), 1951 (Ceroptatus), Cerotelion	19C.2
fuscithorax Enderlein, 1910, Placoceratias	19C.10



galindoi (Lane), 1961 (Platyura), Orfelia (Xenoplatyura)	19C.9
goianensis (Lane), 1948 (Platyura), Isoneuromyia	19C.3
golbachi Lane, 1958, Keroplatus	19C.4
griseofasciata (Edwards), 1933 (Platyura), Isoneuromyia	19C.3
gualanasi Lane, 1950, Macrocera	19C.12
guarani Lane, 1950, Macrocera	19C.12
guayanasi (Lane), 1950 (Platyura), Proceroplatus	19C.11
Heteropterna Skuse, 1888	19C.2
horrida Freeman, 1951, Macrocera	19C.12
iaunai (Lane), 1956 (Platyura), Proceroplatus	19C.11
iconi Lane, 1959, Isoneuromyia	19C.3
ignobilis (Williston), 1896 (Platyura), Orfelia (Neoplatyura)	19C.7
imitans (Lane), 1948 (Keroplatus), Placoceratias	19C.10
inaequalis Freeman, 1951, Macrocera	19C.12
inaequalis (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
incasica (Lane), 1950 (Platyura), Orfelia (Plautyura)	19C.7
infumata (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
Isoneuromyia Brunetti, 1912	19C.3
kaingangi Lane, 1950, Macrocera	19C.12
Keroplatinae	19C.2
Keroplatus Bosc, 1792	19C.4
kerteszi (Lane), 1956 (Platyura), Proceroplatus	19C.11
knabi (Lane), 1950 (Platyura), Orfelia (Lyprauta)	19C.6
Lapyruta Edwards, 1929	19C.6
lauroi (Lane), 1961 (Keroplatus), Neoceroplatus	19C.5
lenkoi Lane, 1961, Isoneuromyia	19C.3
longimanus (Williston), 1896 (Ceroptatus), Placoceratias	19C.10
lopsi (Lane), 1950 (Platyura), Isoneuromyia	19C.4
lutea (Freeman), 1951 (Platyura), Isoneuromyia	19C.4
Lyprauta Edwards, 1931	19C.6
macilentia (Lynch Arribálzaga), 1890 (Platyura), Orfelia (Plau- tyra)	19C.7
Macroccra Meigen, 1803	19C.11, 19C.12
Macrocerinae	19C.11
maiapenai (Lane), 1950 (Platyura), Orfelia (Orfelia)	19C.7
major (Curran), 1928 (Cerotelion), Heteropterna	19C.3
majusculum (Edwards), 1940 (Platyura), Micrapemon	19C.4
manaosensis (Lane), 1961 (Platyura), Orfelia (Xenoplatyura)	19C.9
mathesoni (Lane), 1961 (Platyura), Orfelia (Lapyruta)	19C.6
matilei, nom. nov., Macrocera	19C.12
mexicanus Lane, 1948, Keroplatus	19C.4
Micrapemon Edwards, 1925	19C.4
miersii (Westwood), 1849 (Platyura), Platyroptilon	19C.10
minimax (Edwards), 1934 (Ceroptatus), Neoceroplatus	19C.5
mira Tarwid, 1936 [Zelmira (Neoplatyura)]	19C.13
misionensis Duret, 1975, Platyroptilon	19C.10
monochroma (Edwards), 1933 (Platyura), Orfelia (Taulyrpa)	19C.9



<i>Neoceroplatus</i> Edwards, 1941	19C.5
<i>Neoditomyia</i> Lane & Sturm, 1958	19C.5
<i>Neoplatyura</i> Malloch, 1928	19C.6
<i>nigricans</i> (Lane), 1948 (<i>Ceroplatus</i>), <i>Cerotelion</i>	19C.2
<i>nigrofasciata</i> (Freeman), 1951 (<i>Platyura</i>), <i>Isoneuromyia</i>	19C.4
<i>nobilis</i> Edwards, 1934, <i>Macrocera</i>	19C.12
<i>nubilapex</i> (Edwards), 1940 (<i>Platyura</i>), <i>Orfelia</i> (<i>Lyprauta</i>)	19C.6
<i>obscurus</i> Philippi, 1865 (<i>Ceroplatus</i>)	19C.13
<i>Orfelia</i> Costa, 1857	19C.5, 19C.7
<i>paicoenai</i> , lapsus	19C.5
<i>paicoenai</i> (Lane), 1950 (<i>Ceroplatus</i>), <i>Neoceroplatus</i>	19C.5
<i>paraensis</i> (Lane), 1958 (<i>Platyura</i>), <i>Orfelia</i> (<i>Lyprauta</i>)	19C.6
<i>paraguayana</i> (Edwards), 1934 (<i>Platyura</i>), <i>Orfelia</i> (<i>Pyratula</i>)	19C.8
<i>Paramacrocera</i> Edwards, 1927	19C.13
<i>paramariboensis</i> (Edwards), 1934 (<i>Platyura</i>), <i>Proceroplatus</i> ...	19C.11
<i>parvum</i> (Williston), 1896 (<i>Platyura</i>), <i>Micrapemon</i>	19C.4
<i>paulistana</i> (Lane), 1948 (<i>Platyura</i>), <i>Isoneuromyia</i>	19C.4
<i>perpictula</i> Edwards, 1940, <i>Macrocera</i>	19C.12
<i>picta</i> Freeman, 1951, <i>Macrocera</i>	19C.13
<i>pictipennis</i> (Williston), 1896 (<i>Platyura</i>), <i>Proceroplatus</i>	19C.11
<i>Placoceratias</i> Enderlein, 1910	19C.9
<i>Platyroptilon</i> Westwood, 1849	19C.10
<i>Platyura</i> , authors	19C.5
<i>plumanni</i> Edwards, 1940, <i>Macrocera</i>	19C.13
<i>Plautyra</i> Edwards, 1941	19C.7
<i>Proceroplatus</i> Edwards, 1925	19C.10
<i>Pseudoplatyura</i> Skuse, 1888	19C.8
<i>puncticoxalis</i> (Edwards), 1940 (<i>Platyura</i>), <i>Proceroplatus</i>	19C.11
<i>Pyratula</i> Edwards, 1929	19C.8
<i>Pyrtaula</i> Edwards, 1929	19C.8
<i>quasilineata</i> (Edwards), 1919 (<i>Platyura</i>), <i>Orfelia</i> (<i>Pyrtaula</i>)	19C.8
<i>rabelloi</i> (Lane), 1956 (<i>Platyura</i>), <i>Proceroplatus</i>	19C.11
<i>ramicornis</i> Enderlein, 1910, <i>Platyroptilon</i>	19C.10
<i>regularis</i> (Edwards), 1934 (<i>Platyura</i>), <i>Orfelia</i> (<i>Neoplatyura</i>)	19C.7
<i>rubens</i> Wiedemann, 1828 (<i>Platyura</i>)	19C.13
<i>Rypatula</i> Edwards, 1929	19C.9
<i>salcbrensis</i> (Edwards), 1941 (<i>Platyura</i>), <i>Orfelia</i> (<i>Plautyra</i>)	19C.7
<i>saparai</i> (Lane), 1950 (<i>Platyura</i>), <i>Orfelia</i> (<i>Neoplatyura</i>)	19C.7
<i>sesiiformis</i> (Edwards), 1940 (<i>Platyura</i>), <i>Isoneuromyia</i>	19C.4
<i>shannoni</i> Lane, 1947, <i>Macrocera</i>	19C.13
<i>shannoni</i> (Lane), 1948 (<i>Platyura</i>), <i>Orfelia</i> (<i>Plautyra</i>)	19C.7
<i>singularis</i> (Lane), 1948 (<i>Ceroplatus</i>), <i>Euceroplatus</i>	19C.2
<i>souzai</i> (Lane), 1959 (<i>Platyura</i>), <i>Orfelia</i> (<i>Pseudoplatyura</i>)	19C.8
<i>spatulata</i> (Edwards), 1940 (<i>Platyura</i>), <i>Orfelia</i> (<i>Apyrtula</i>)	19C.6
<i>stonei</i> (Lane), 1950 (<i>Platyura</i>), <i>Proceroplatus</i>	19C.11
<i>subannulata</i> Philippi, 1865 (<i>Platyura</i>)	19C.13



tamoyoi Lane, 1950, Macrocera	19C.13
tamoyoi (Lane), 1950 (Platyura), Orfelia (Plautyra)	19C.8
tapuaii (Lane), 1950 (Platyura), Orfelia (Xenoplatyura)	19C.9
Taulyrpa Edwards, 1929	19C.9
teranoi (Lane), 1950 (Platyura), Proceroplatus	19C.11
testacea Philippi, 1865, Macrocera	19C.13
tetrалеuca (Edwards), 1940 (Ceroplatus), Heteropterna	19C.3
thomsoni Lynch Arribálzaga, 1882, Macrocera	19C.13
timbira Lane, 1956, Isoneuromyia	19C.4
townsendi Lane, 1948, Keroplatus	19C.4
triangulifera (Freeman), 1951 (Platyura), Orfelia (Pyrtaula)	19C.8
trileuca (Edwards), 1940 (Ceroplatus), Heteropterna	19C.3
trinidadensis (Lane), 1960 (Platyura), Proceroplatus	19C.11
troglophila Matile, 1978, Neoditomyia	19C.5
tucumana Lane, 1958, Isoneuromyia	19C.4
uaracui (Lane), 1950 (Ceroplatus), Placoceratias	19C.10
unica Fisher, 1939, Macrocera	19C.13
unidens Edwards, 1931, Macrocera	19C.13
valdiviana Philippi, 1865, Macrocera	19C.13
variventris (Edwards), 1931 (Platyura), Proceroplatus	19C.11
venustipennis (Shaw), 1940 (Platyura), Proceroplatus	19C.11
vespiformis (Enderlein), 1910 (Cerotelion), Orfelia (Taulyrpa)	19C.9
villasboasi (Lane), 1961 (Platyura), Proceroplatus	19C.11
vittatus (Fisher), 1941 (Platyura), Proceroplatus	19C.11
xanthocera (Edwards), 1931 (Platyura), Isoneuromyia	19C.4
Xenoplatyura Malloch, 1928	19C.9
<i>Zelmira</i> Meigen, 1800	19C.5
zernyi Edwards, 1934, Platyroptilon	19C.10
zeteki (Lane), 1950 (Platyura), Orfelia (Lyprauta)	19C.6



Composto e impresso em 1978, nas oficinas da
EMPRESA GRÁFICA DA REVISTA DOS TRIBUNAIS S.A.
Rua Conde de Sarzedas, 38 — Tel. 36-6958 (PBX)
CEP 01512 — São Paulo, SP, Brasil,





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO

(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

19 D

FAMILY LYGISTORRHINIDAE

N. PAPAVERO

Museu de Zoologia
Universidade de São Paulo

Thompson (1975), in his review of the American species, discusses the family status and the geographical distribution of this group. Only one genus (considered as subgenus by Thompson) occurs in the Americas.

Genus PROBOLAEUS Williston

PROBOLAEUS Williston, 1896: 261. Type-species, *singularis* Williston (mon.).

Aphanizophleps Enderlein, 1910: 201. Type-species, *coxata* Enderlein (orig. des.).

Lygistorrhina, authors, not Skuse.

BARRETOI (Lane), 1947: 346 (*Lygistorrhina*). Type-locality: Brazil, Goiás, Corumbá. Type ♀, MZSP. N. comb.

BRASILIENSIS (Edwards), 1932: 139 (*Lygistorrhina*). Type-locality: Brazil, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro to Santa Catarina). Type, ?BMNH. N. comb.

CERQUEIRAI (Lane), 1958: 209, figs. 1 (wing), 2 (♂ genitalia) (*Lygistorrhina*). Type-locality: Brazil, Amazonas, Manaus, Igarapé do Mariano. Distr. — Brazil (Amazonas). Type ♂, MZSP. N. comb.

COXATUS (Enderlein), 1910: 203, fig. 4 (wing) (*Aphanizophleps*). Type-locality: Brazil, Santa Catarina. Type ♂, Stettin. N. comb.

EDWARDSI (Lane), 1947: 347 (*Lygistorrhina*). Type-locality: Brazil, São Paulo, Ferraz de Vasconcelos. Type ♀, MZSP. N. comb.

SINGULARIS Williston, 1896: 261, pl. 8, fig. 15 (wing), 15a (head), 15b (mouthparts), 15c (genitalia). Type-locality: St. Vincent (W. I.). Ref. — Johannsen, 1909: 93 (descr., distr.). Type ♂, BMNH.

URICHI (Edwards), 1912: 204 (*Lygistorrhina*). Type-locality: Trinidad, Diego Martin. Distr. — Trinidad, Brazil (Amapá). Ref. — Lane, 1959: 106. Type ♂, BMNH. N. comb.

19D.1



LITERATURE CITED

EDWARDS, F. W.

1912. *Lygistorrhina urichi*, a new mycetophilid from Trinidad. *Ann. Mag. nat. Hist.* (8) 10: 203-204.
 1932. New Brazilian Mycetophilidae (Diptera). *Revta Ent.* 2: 138-149, 6 figs.

ENDERLEIN, G.

1910. Neue Gattungen und Arten aussereuropäischer Fliegen. *Stettin. ent. Ztg* 72 (1911): 135-209.

JOHANNSEN, O. A.

1909. Diptera, Fam. Mycetophilidac. *Gen. Insect.* 93: 1-141.

LANE, J.

1947. New Brazilian Mycetophilidae (Diptera, Nemocera). *Revta Ent.* 17 (1946): 339-360, 14 figs.
 1958. On Amazonian Mycetophilidac (Diptera, Nematocera). *Studia ent.* 1: 209-216, 8 figs.
 1959. Insecta Amapaensia. — Diptera: Mycetophilidae. *Studia ent.* 2 (1-4): 105-118, 14 figs.

THOMPSON, F. C.

1975. Notes on the genus *Lygistorrhina* Skuse with the description of the first nearctic species (Diptera: Mycetophiloidea). *Proc. ent. Soc. Wash.* 77 (4): 434-445, 14 figs.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. ent. Soc. Lond.* 1896: 253-449, pls. 8-14.



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
 (FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)
 A CATALOGUE OF THE DIPTERA OF THE AMERICAS SOUTH
 OF THE UNITED STATES

19 E

FAMILY MYCETOPHILIDAE

(FUNGIVORIDAE)

N. PAPAVERO

Museu de Zoologia
Universidade de São Paulo

Adult mycetophilids are mostly caught in moist, wooded places, or in light- and Shannon- or Malaise traps. Larvae feed on decaying wood, mosses, but mostly on several groups of fungi. Very little is known about the biology of the family in the neotropical region.

Over 750 species have been described from the neotropics, especially by Edwards, Freeman, and Lane. Many more are expected to occur, and most species are only known from the type-localities.

Abbreviations for museums are the same employed in fascicie 19C. I am greatly indebted to Dr. Loïc Matile (MHNP) for many corrections and suggestions.

Subfamily MANOTINAE

Ref. — Tuomikoski, 1966b.

Genus MANOTA Williston

MANOTA Williston, 1896: 260. Type-species, *defecta* Williston (mon.).


Aphanizophleps Enderlein, 1910: 201. Type-species, *coxata* Enderlein (orig. des.). This genus was placed in the synonymy of *Manota*. Lane (1947: 345) synonymized it with *Lygistorrhina*. This last wrong placement was followed by me in catalogue 19D, whence the name should be deleted.

COXATA (Enderlein), 1910: 203, fig. 4 (wing) (*Aphanizophleps*). Type-locality: Brazil, Santa Catarina. Type ♂, Stettin. N. comb.

DEFECTA Williston, 1896: 260, pl. 8, fig. 14 (whole insect). Type-locality: St. Vincent, W. I. Syntypes ♂, BMNH.

PALPALS Lane, 1948b: 247, figs. 1 (palpus), 2 (antenna), 3 (wing), 4 (♂ genitalia). Type-locality: Brazil, Goiás, Corumbá. Type ♂, MZSP.

19E.1



Subfamily MYCETOPHILINAE

Tribe EXECHIINI

Ref. — Tuomikoski, 1966a (gen. rev.).

Genus ALLODIOPSIS Tuomikoski

ALLODIOPSIS Tuomikoski, 1966a: 171. Type-species, *Rhymosia rustica* Edwards (orig. des.).

Subgenus GYMNOGONIA Tuomikoski

Allodiopsis, subg. GYMNOGONIA Tuomikoski, 1966a: 171. Type-species, *Rhymosia sintenisi* Lackschewitz (orig. des.).

CINEREA (Freeman), 1951: 91, fig. 176 (♂ genitalia) (*Rhymosia*). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Ñuble, Llanquihue). Ref. — Tuomikoski, 1966a: 172. Type ♂, BMNH. N. COMB.

Genus BORACEOMYIA Lane

BORACEOMYIA Lane, 1948a: 231. Type-species, *edwardsi* Lane (orig. des.) = *goianensis* (Lane).

Exechia, authors, not Winnertz.

ACARIENSIS (Lane), 1951a: 251, fig. 4 (♂ genitalia) (*Exechia*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP. N. COMB.

AITKENI (Lane), 1960a: 384, fig. 9 (♂ genitalia) (*Exechia*). Type-locality: Trinidad, Chaguaramas. Type ♂, MZSP. N. COMB.

BORGMEIERI (Lane), 1951a: 252, fig. 14 (♂ genitalia) (*Exechia*). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP. N. COMB.

BORINQUENSIS (Lane), 1951a: 254, fig. 15 (♂ genitalia) (*Exechia*). Type-locality: Costa Rica, San José. Type ♂, USNM. N. COMB.

CAJUENSIS (Lane), 1951a: 252, fig. 5 (♂ genitalia) (*Eyechia*, misspell.). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP. N. COMB.

EMBUENSIS (Lane), 1956b: 148, fig. 1 (♂ genitalia) (*Exechia*). Type-locality: Brazil, São Paulo, Embu. Type ♂, MZSP. N. COMB.

GOIANENSIS (Lane), 1947a: 356, figs. 5 (♂ genitalia), 13 (wing) (*Exechia*). Type-locality: Brazil, Goiás, Corumbá. Distr. — Trinidad, Peru (Tingo Maria), Brazil (Goiás, São Paulo), Argentina (Salta). Ref. — Lane, 1956b: 147 (syn.). Type ♂, MZSP. N. COMB.

- edwardsi* Lane, 1948a: 231. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♀, MZSP.
- PAULISTENSIS (Lane), 1947a: 357, fig. 10 (wing) (*Exechia*). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Ref. — Lane, 1951a: 250, fig. 3 (♂ genitalia). Type ♀, MZSP. N. comb.

Genus EXECHIOPSIS Tuomikoski

- EXECHIOPSIS Tuomikoski, 1966a: 177. Type-species, *Exechia subulata* Winnertz (orig. des.).

Subgenus XENEXECHIA Tuomikoski

- Exechiopsis*, subg. XENEXECHIA Tuomikoski, 1966a: 179. Type-species, *Exechia perspicua* Johannsen (orig. des.).
- BIFIDA (Freeman), 1951: 90, fig. 172 (♂ genitalia) (*Exechia*). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH. N. comb.
- BREVIFURCATA (Freeman), 1951: 90, figs. 174 (♂ genitalia), 293 (wing) (*Exechia*). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Llanquihue, Valparaíso). Type ♂, BMNH. N. comb.
- EXTENSA (Freeman), 1951: 91, fig. 171 (♂ genitalia) (*Exechia*). Type-locality: Chile, Llanquihue, Casa Pangué. Type ♂, BMNH. N. comb.
- FUNEREA (Freeman), 1951: 91, fig. 175 (♂ genitalia) (*Exechia*). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH. N. comb.
- FURCILLA (Freeman), 1954: 29, fig. 1 (♂ genitalia) (*Exechia*). Type-locality: Juan Fernandez Isl., Masafuera, Quebrada de las Casas. Distr. — Juan Fernandez Isl. (Masafuera, Masatierra). Type ♂, Universidad de Chile, Santiago. N. comb.
- SETIGERA (Freeman), 1951: 89, fig. 170 (♂ genitalia) (*Exechia*). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH. N. comb.
- TRUNCATA (Freeman), 1951: 90, fig. 173 (♂ genitalia) (*Exechia*). Type-locality: Chile, Llanquihue, Casa Pangué. Type ♂, BMNH. N. comb.

Genus BREVICORNU Marshall

- BREVICORNU Marshall, 1896: 306. Type-species, *flavum* Marshall (Tuomikoski, 1966a: 184).
- Allodia*, authors, not Winnertz.
- ARAUCANIENSE (Lane), 1962b: 539, fig. 6 (♂ genitalia) (*Allodia*). Type-locality: Chile, Ñuble, 40 km e. of San Carlos. Distr. — Chile (Ñuble, Bío Bío). Type ♂, CAS. N. comb.




- ARGENTINENSE (Lane), 1958b: 162, fig. 6 (♂ genitalia) (*Allodia*). Type-locality: Argentina, Tumucán, Tafi del Valle. Type ♂, IML. N. comb.
- CALLIDUM (Johannsen), 1912: 319 (*Allodia*). Type-locality: U.S.A., Washington. Distr. — U.S.A. (Washington, Wyoming), Costa Rica. N. comb.
- COMPLEXUM (Lane), 1948a: 234, fig. 1 (♂ genitalia) (*Allodia*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo), Argentina (Santa Cruz, Corrientes, Misiones). Ref. — Lane, 1956b: 150, fig. 2 (♂ genitalia). Type ♂, MZSP. N. comb.
- ERULIENSE (Lane), 1956b: 151, fig. 3 (♂ genitalia) (*Allodia*). Type-locality: Brazil, Rio de Janeiro, Itatiaia. Type ♂, MZSP. N. comb.
- HUMILE (Lane), 1948a: 233 (*Allodia*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Ref. — Lane, 1956b: 152, fig. 4 (♂ genitalia). Type ♀, MZSP. N. comb.
- SIMILE (Freeman), 1951: 92, fig. 177 (♂ genitalia) (*Allodia*). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Santa Cruz, Tierra del Fuego), Chile (Ñuble, Llanquihue). Type ♂, BMNH. N. comb.

Genus NEALLODIA Edwards

- NEALLODIA Edwards, 1932: 146. Type-species, *flavida* Edwards (mon.) = *brevicornis* (Enderlein).
- BREVICORNIS (Enderlein), 1910a: 204 (*Allodia*). Type-locality: Brazil, Santa Catarina. Distr. — Trinidad, Brazil (Mato Grosso, São Paulo, Santa Catarina), Argentina (Salta), Paraguay. Type ♀, Stettin.
- flavida* Edwards, 1932: 146. Type-locality: Brazil, Rio de Janeiro, Petrópolis.
- brasiliiana* Lane, 1947b: 448, figs. 1 (wing), 2 (♂ genitalia) (*Trichonta*). Type-locality: Brazil, São Paulo, São Paulo (Ipiranga). Type ♀, MZSP.

Genus RYMOSIA Winnertz

- RYMOSIA Winnertz, 1863: 810. Type-species, *Mycetophila discoidea* Meigen (Johannsen, 1909: 102) = *fasciata* (Meigen).
- Rhymosia*, emend.
- AIROSAI Lane, 1947a: 358. Type-locality: Brazil, Amazonas, Maués. Distr. — Panama, Brazil (Amazonas, Pará, Rio de Janeiro). Ref. — Lane, 1958a: 213, fig. 7 (♂ genitalia). Type ♂, MZSP.
- DAMASCENOI Lane, 1958a: 214, fig. 8 (♂ genitalia). Type-locality: Brazil, Pará, Rio Tapajós, Fordlândia. Distr. — Brazil (Amazonas, Pará). Type ♂, MZSP.
- PARAENSIS Lane, 1958a: 215. Type-locality: Brazil, Pará, Rio Tapajós, Fordlândia. Type ♀, MZSP.



PILOSA Edwards, 1931: 259. Type-locality: Peru, Lima, Verrugas. Type ♀, BMNH.

WORONTZOWI Lane, 1947a: 359. Type-locality: Brazil, Amazonas, Maués. Type ♀, MZSP.

UNPLACED EXECHIINI

BORACENSIS Lane, 1962a: 14, fig. 3 (♂ genitalia) (*Allodia*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP. — Near *Brevicornu*, but posterior fork almost below anterior fork.

GRACILE Kertész, 1901a: 403, pl. 20, fig. 1 (wing) (*Dynatosoma*). Type-locality: Peru, Callanga. Type ♀, Budapest.

Tribe MYCETOPHILINI

Genus EPICYPTA Winnertz

EPICYPTA Winnertz, 1863: 909. Type-species, *Mycetophila scatophora* Perris (Johannsen, 1909: 110).

Delopsis Skuse, 1890: 623. Type-species, *flavipennis* Skuse (mon.).

Plastacephala Enderlein, 1910a: 175. Type-species, *planiventris* Enderlein (orig. des.).

Subgenus BOSCMYIA Lane

Epicypta, subg. BOSCMYIA Lane, 1954c: 115. Type-species, *lenguasi* Lane (orig. des.).

ARAPASCOI Lane, 1954c: 117, fig. 1 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

BARAI Lane, 1954c: 119, fig. 2 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

BOSCH (Lane), 1948a: 266, figs. 17, 33, 53 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.

FORATTINI Lane, 1960b: 482, fig. 1 (♂ genitalia). Type-locality: Brazil, Mato Grosso, Bataguaçu, Rio Três Barras. Type ♂, MZSP.

IUMENSIS Lane, 1954c: 120, fig. 3 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Amapá, Santa Catarina). Type ♂, BMNH.

LENGUASI Lane, 1954c: 121, fig. 4 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

MAUESENSIS Lane, 1954c: 124, fig. 6 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

MILTONI (Lane), 1948a: 270 (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Ref. — Lane, 1954: 123, fig. 5 (♂ genitalia). Type ♀, MZSP.



- OPAI EI Lane, 1954c: 125, fig. 7 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- TRISTIS (Lane), 1948a: 271 (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Pará, São Paulo). Ref. — Lane, 1954c: 126, fig. 8 (♂ genitalia). Type ♀, MZSP.
- lugubris* Lane, 1948a: 272 (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♀, MZSP.
- WILLIAMS I Lanc, 1960b: 484, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, São Miguel Arcanjo (Turvo dos Hilários). Type ♂, MZSP.

Subgenus CALLICYPTA Lane

- Epicypsa*, subg. CALLICYPTA Lane, 1954c: 127. Type-species, *Delopsis lunata* Lane (orig. des.).
- AMAPAENSIS Lane, 1961a: 254, fig. 7 (wing). Type-locality: Brazil, Amapá, Rio Felício. Type ♀, MZSP.
- ANAMBESI Lane, 1960b: 485, fig. 3 (♂ genitalia). Type-locality: Brazil, Mato Grosso, Bataguáçu, Rio Caraguatá. Type ♂, MZSP.
- BACAIRISI Lane, 1954c: 129, fig. 9 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- BICELLII Lane, 1961a: 255, fig. 6 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- CHIRIGUANAI Lane, 1954c: 130, fig. 10 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- COHERI Lane, 1954c: 132, fig. 11 (♂ genitalia). Type-locality: Brazil, São Paulo, Salcsópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- COUASI Lane, 1954c: 133, fig. 12 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- INORNATA Lane, 1948a: 237, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- IRANCHESI Lane, 1960b: 485, fig. 4 (♂ genitalia). Type-locality: Trinidad, Navy Base. Type ♂, MZSP.
- JOHANNSENI (Lane), 1947a: 354 (*Delopsis*). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- taunayi* Lane, 1948a: 249, figs. 9, 47 (♂ genitalia) (*Mycetophila*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- KUICURUSI Lane, 1954c: 135, fig. 13 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Amapá, Santa Catarina). Type ♂, BMNH.
- LUNATA (Lane), 1948a: 267, figs. 21, 55 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de



- Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.
- MACUNAI Lane, 1954c: 136, fig. 14 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- ORNATA (Lane), 1948a: 265, figs. 16, 51 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- PAULISTENSIS Lane, 1948a: 236, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Costa Rica, Trinidad, Brazil (São Paulo). Type ♂, MZSP.
- PUNCTULATA (Lane), 1947a: 355 (*Delopsis*). Type-locality: Brazil, Mato Grosso, Maracaju. Type ♂, MZSP.
- RABELLOI (Lane), 1948a: 275, figs. 23, 58 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SIUSISI Lane, 1954c: 137, fig. 15 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- TACANHAPESI Lane, 1960b: 486, fig. 5 (♂ genitalia). Type-locality: Trinidad, Navy Base. Distr. — Trinidad, Brazil (Amapá). Type ♂, MZSP.
- TRINIDADENSIS Lane, 1960b: 487, fig. 6 (♂ genitalia). Type-locality: Trinidad, Sangre Grande, Mandillán Road. Type ♂, MZSP.

Subgenus EPICYPTA Winnertz

- ACZELI Lane, 1958b: 154, fig. 5 (♂ genitalia). Type-locality: Argentina, Salta, Embarcación. Distr. — Trinidad, Brazil (Goiás, São Paulo), Argentina (Corrientes, Salta, Chaco, Tucumán). Type ♂, IML.
- AGUARENSIS Lane, 1951b: 329, fig. 1 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- BAPIANENSIS Lane, 1951b: 332, fig. 4 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- BRASILLANA (Edwards), 1932: 147 (*Delopsis*). Type-locality: Brazil, São Paulo, Rio Claro; Rio de Janeiro, Petrópolis. Distr. — Trinidad, Brazil (Goiás, Rio de Janeiro, São Paulo, Santa Catarina), Argentina (Salta). Refs. — Lane, 1947b: 454, fig. 4 (♂ genitalia); 1951b: 338, fig. 10 (♂ genitalia). Syntypes, BMNH.
- BRASILIENSIS (Enderlein), 1910a: 171 (*Mycetophila*). Type-locality: Brazil, Santa Catarina. Distr. — Panama, Brazil (São Paulo, Santa Catarina). Refs. — Lane, 1947b: 455, fig. 5 (♂ genitalia); 1951b: 340, fig. 12 (♂ genitalia). Type ♂, Stettin.
- CURUPIENSIS Lane, 1961a: 253, fig. 5 (♂ genitalia). Type-locality: Brazil, Amapá, Rio Felício. Type ♂, MZSP.
- DUANASI Lane, 1951b: 337, fig. 9 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Panama, Brazil (Santa Catarina). Type ♂, BMNH.



- GOIANENSIS (Lane), 1947b: 456 (*Delopsis*). Type-locality: Brazil, Goiás, Corumbá. Distr. — Panama, Brazil (Goiás), Argentina (Misiones). Type ♀, MZSP.
- GUANENSIS Lane, 1951b: 38, fig. 11 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Costa Rica, Brazil (Amapá, Santa Catarina), Argentina (Tucumán). Type ♂, BMNH.
- GUARANI Lane, 1951b: 331, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.
- IAUNENSIS Lanc, 1951b: 333, fig. 5 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Panama, Brazil (Santa Catarina). Type ♂, BMNH.
- JARUENSIS Lane, 1951b: 334, fig. 6 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- LUEDERWALDTI (Enderlein), 1910a: 170 (*Mycetophila*). Type-locality: Brazil, Santa Catarina. Ref. — Lane, 1951b: 340. Type ♀, Stettin.
- MACULIPENNIS (Enderlein), 1910a: 178 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Ref. — Lane, 1948a: 277, figs. 24, 38, 61 (♂ genitalia). Syntypes ♀, Stettin.
- MAUPIENSIS Lane, 1951b: 340. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♀, BMNH.
- NIROHIALTERATA (Enderlein), 1910a: 179 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Ref. — Lane, 1951b: 329, fig. 2 (♂ genitalia). Syntypes ♀, Stettin.
- OEDIPUS Edwards, 1934: 365. Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Panama, Brazil (São Paulo, Santa Catarina), Paraguay. Type ♀, WIEN.
- PERUANA Lane, 1962a: 14, fig. 4 (♂ genitalia). Type-locality: Peru, Tingo Maria, Monson Valley. Type ♂, CAS.
- PLANIVENTRIS (Enderlein), 1910a: 176 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Distr. — Costa Rica, Brazil (Mato Grosso, Santa Catarina). Ref. — Lane, 1951b: 336, fig. 7 (♂ genitalia). Syntypes ♂ ♀, Stettin.
- var. FLAVIPES (Enderlein), 1910a: 178 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Syntypes ♀, Stettin.
- SCHILDI Lane, 1956b: 156, fig. 7 (♂ genitalia). Type-locality: Costa Rica, San Mateo, Higuito. Type ♂, USNM.
- TACUENSIS Lane, 1951b: 336, fig. 8 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Unplaced EPICYPTA

- ANCYLOFORMANS (Holmgren), 1907: 1-7 (*Mycetophila*). Type-locality: Peru.
- CAPITATA (Enderlein), 1910a: 180 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Syntypes ♀, Stettin.



- INSIPENS (Williston), 1896: 264, pl. 8, fig. 19 (wing) (*Mycetophila*). Type-locality: St. Vincent, W. I. Syntypes ♀, BMNH.
- LONGICALCAR (Enderlein), 1910a: 179 (*Plastacephala*). Type-locality: Brazil, Santa Catarina. Syntypes ♂ ♀, Stettin.
- MACULA (Enderlein), 1910a: 172 (*Mycetophila*). Type-locality: Brazil, Santa Catarina. Type ♀, Stettin.
- ORNATIDORSUM (Enderlein), 1910a: 173 (*Mycetophila*). Type-locality: Brazil, Santa Catarina. Syntypes ♂ ♀, Stettin.

Genus MYCETOPHILA Meigen

- Fungivora* Meigen, 1800: 16. Type-species, *Tipula agarici* Villiers (as Olivier; Coquillett, 1910: 545). Suppressed ICZN, 1963: 339.
- MYCETOPHILA Meigen, 1803: 263. Type-species, *Tipula agarici* Villiers (Johannsen, 1909: 116).

Subgenus ABMYCETA Lane

- Mycetophila*, subg. ABMYCETA Lane, 1955: 400. Type-species, *galibisi* Lane (orig. des.).
- ACARISI Lane, 1958b: 157, fig. 9 (♂ genitalia). Type-locality: Argentina, Tucumán, Tafi del Valle. Type ♂, IML.
- ANHANGAENSIS Lane, 1961a: 251, fig. 3 (♂ genitalia). Type-locality: Brazil, Amapá, Río Felício. Type ♂, MZSP.
- ARAUASI Lane, 1958c: 400, fig. 3 (♂ genitalia). Type-locality: Trinidad, Navy Base. Type ♂, MZSP.
- ARGENTINA Lane, 1958b: 157, fig. 10 (♂ genitalia). Type-locality: Argentina, Santa Cruz, Lago Argentino. Distr. — Argentina (Santa Cruz, Tierra del Fuego). Type ♂, IML.
- BARRETOI (Lane), 1947b: 453 (*Delopsis*). Type-locality: Brazil, São Paulo, Cajuru. Distr. — Trinidad, Brazil (São Paulo). Ref. — Lane, 1958c: 400, fig. 4 (♂ genitalia). Type ♀, MZSP.
- BORGMEIERI Edwards, 1932: 149. Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil (Amapá, São Paulo, Paraná), Argentina (Catarinense, Tucumán). Syntypes ♂ ♀, BMNH.
- BRIDGESI Lane, 1963: 13, fig. 3 (♂ genitalia). Type-locality: Argentina, Tierra del Fuego, Lago Fagnano. Type ♂, MZSP.
- CANICULA Freeman, 1951: 127, fig. 258 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Santa Cruz), Chile (Bío Bío). Type ♂, BMNH.
- CARIBAI Lane, 1955: 402, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- CARIPUNAI Lane, 1955: 403, fig. 8 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.



- CARRERAI Lane, 1958c: 401, fig. 5 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- CAYUENSIS Lane, 1955: 404, fig. 9 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- CLAROVITTATA Freeman, 1951: 102, figs. 192 (♂ genitalia), 193 (♀ terminalia), 312 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Bío Bío, Ñuble, Llanquihue). Type ♂, BMNH.
- CLAVIGERA Freeman, 1951: 125, fig. 253 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Santa Cruz, Chubut, Tierra del Fuego), Chile (Osorno, Bío Bío, Ñuble, Llanquihue). Type ♂, BMNH.
- CONIFERA Freeman, 1951: 125, fig. 252 (♂ genitalia), 344 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Ñuble, Llanquihue). Type ♂, BMNH.
- CONSTRICTA Freeman, 1951: 128, fig. 261 (♂ genitalia), 349 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Santa Cruz, Tierra del Fuego), Chile (Ñuble, Llanquihue). Type ♂, BMNH.
- COSTARICENSIS Lane, 1958c: 402, fig. 6 (♂ genitalia). Type-locality: Costa Rica, San Mateo, Higuito. Type ♂, USNM.
- COXIPONESI Lane, 1958c: 402, fig. 7 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- DEMACURI Lane, 1955: 405, fig. 10 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- DICHAETA Freeman, 1951: 126, fig. 257 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tucumán), Chile (Ñuble, Bío Bío). Type ♂, BMNH.
- DISTINCTA Freeman, 1951: 104, figs. 197 (♂ genitalia), 198-199 (♀ terminalia), 316 (wing). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro), Chile (Ñuble, Llanquihue). Type ♂, BMNH.
- ERAMANENSIS Lane, 1955: 406, fig. 11 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- FLAVOLUNATA Freeman, 1951: 122, figs. 246 (♂ genitalia), 341 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Santa Cruz), Chile (Bío Bío, Llanquihue). Type ♂, BMNH.
- FLEXISETA Freeman, 1951: 105, figs. 203 (♂ genitalia), 318 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Ñuble). Type ♂, BMNH.
- FORATTINI Lane, 1955: 407, fig. 12 (♂ genitalia). Type-locality: Brazil, São Paulo, Presidente Veneslau (Fazenda Guanabara). Distr. — Trinidad, Brazil (Amapá, São Paulo, Santa Catarina). Type ♂, MZSP.

- FUNEREA Freeman, 1951: 107, fig. 206 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Osorno, Llanquihue). Type ♂, BMNH.
- GALIBISI Lane, 1955: 408, fig. 13 (♂ genitalia). Type-locality: Brazil, São Paulo, Salcsópolis, Estação Biológica de Boracéia. Distr. — Brazil (Amapá, São Paulo, Santa Catarina). Type ♂, MZSP.
- GOLBACHI Lane, 1958b: 159, fig. 12 (♂ genitalia). Type-locality: Argentina, Tucumán, Tafi del Valle. Distr. — Argentina (Tucumán, Tierra del Fuego), Chile (Osorno). Type ♂, IML.
- GRADATA Freeman, 1951: 130, figs. 353 (wing), 265 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- GUATENSIS Lane, 1955: 408, fig. 14 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- HUMBOLDTI (Lane), 1948a: 273, figs. 19, 52 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- ICOSI Lane, 1958c: 404, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- JURII Lanc, 1955: 412, fig. 18 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- JURUNENSIS Lane, 1955: 410, fig. 15 (♂ genitalia). Type-locality: Brazil, São Paulo, São Miguel Areanjo (Turvo). Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.
- KAINGANGI Lane, 1956b: 158, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- KUNASENSIS Lane, 1955: 413, fig. 16 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- LACUNA Freeman, 1951: 123, figs. 248 (♂ genitalia), 342 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Bio Bio, Ñuble, Llanquihue, Magallanes). Type ♂, BMNH.
- LAIANASI Lane, 1958c: 404, fig. 9 (♂ genitalia). Type-locality: Trinidad, Macqueripe, Tucker Valley. Type ♂, MZSP.
- LATIVITTA Freeman, 1951: 129, figs. 264 (♂ genitalia), 352 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- NUBLENSIS Lane, 1962b: 545, fig. 8 (♂ genitalia). Type-locality: Chile, Ñuble, 40 km e. of San Carlos. Type ♂, CAS.
- PARANOTATA Freeman, 1954: 39 (nom. nov. for *notata* Freeman). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Ñuble, Llanquihue).
- notata* Freeman, 1951: 126, fig. 256 (♂ genitalia) (preocc. Stannius, 1831, now in *Zygomysia*). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.

- PARAPICALIS Freeman, 1954: 39 (nom. nov. for *apicalis* Freeman). Type-locality: Argentina, Río Negro, Lago Gutiérrez.
- apicalis* Freeman, 1951: 124, fig. 250 (♂ genitalia) (preocc. Meigen, 1838, now in *Rymosia*, as synonym of *fenestralis* Meigen). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- PATAGONESI Lane, 1958c: 406, fig. 10 (♂ genitalia). Type-locality: Argentina, Tucumán, Tafi del Valle. Type ♂, IML.
- PECTINATA Freeman, 1951: 125, figs. 254 (♂ genitalia), 346 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- PELLUCIDA Freeman, 1951: 103, figs. 195-196 (♂ genitalia), 313-314 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Valparaiso to Magallanes). Type ♂, BMNH.
- PENICULATA Freeman, 1951: 107, figs. 207 (♂ genitalia), 321 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- PIRAPESI Lane, 1958c: 406, fig. 11 (♂ genitalia). Type-locality: Argentina, Santa Cruz, Lago Argentino. Distr. — Argentina (Santa Cruz, Tierra del Fuego). Type ♂, MZSP.
- PLAUMANNI Lane, 1958c: 407, fig. 12 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, MZSP.
- PLURIPUNCTATA (Lane), 1948a: 274, figs. 20, 60 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.
- PUNENSIS Lane, 1955: 415, fig. 17 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- RUFA (Lane), 1948a: 264 (*Delopsis*; preocc. Macquart, 1826). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, MZSP. If a valid species, requires a new name.
- SINUATA Freeman, 1951: 130, figs. 266 (♂ genitalia), 354 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- SOLITA Freeman, 1951: 124, fig. 251 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- SPINOSA Freeman, 1951: 129, figs. 262 (♂ genitalia), 350 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Bío Bío, Ñuble, Llanquihue). Type ♂, BMNH.
- SUBBRUNNEA Freeman, 1951: 131, figs. 269 (♂ genitalia), 355 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Osorno). Type ♂, BMNH.
- SUBFUMOSA Freeman, 1954: 32, pl. 2, h (♂ genitalia). Type-locality: Juan Fernandez Isl., Masatierra, Yunque. Type ♂, Universidad de Chile, Santiago.



- SUBRECTA Freeman, 1954: 39 (nom. nov. for *recta* Freeman). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Tucumán, Río Negro), Chile (Chiloé).
- recta* Freeman, 1951: 124, figs. 249 (♂ genitalia), 343 (wing) (preocc. Johannsen, 1912). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- SUBVITTATA Freeman, 1954: 40 (nom. nov. for *subvittata* Freeman). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro, Tierra del Fuego).
- trivittata* Freeman, 1951: 131, fig. 268 (♂ genitalia) (preocc. Staeger, 1840, now in *Erechia*). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- TAPINIRAI Lane, 1955: 415 (nom. nov. for *confusa* Lane). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Goiás, São Paulo).
- trivittata* Freeman, 1951: 131, fig. 268 (♂ genitalia) (preocc. Staeger, dzicki, 1886). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- TAPUIAI Lane, 1958b: 160, fig. 14 (♂ genitalia). Type-locality: Argentina, Tucumán, Lacavera. Type ♂, IML.
- TRAVASSOSI (Lane), 1948a: 268, figs. 18, 54 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- TRIORDINATA Freeman, 1954: 40 (nom. nov. for *triseriata* Freeman). Type-locality: Argentina, Río Negro, Bariloche.
- triseriata* Freeman, 1951: 128, figs. 260 (♂ genitalia), 348 (wing) (preocc. Bukovski, 1949). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- TUCUMANA Lane, 1958b: 160, fig. 16 (♂ genitalia). Type-locality: Argentina, Tucumán, Tafi del Valle. Distr. — Argentina (Tucumán, Tierra del Fuego), Chile (Ñuble). Type ♂, IML.
- TUPANENSIS Lane, 1961a: 252, fig. 4 (♂ genitalia). Type-locality: Brazil, Amapá, Rio Felício. Type ♂, MZSP.
- UACUPISI Lane, 1958c: 408, fig. 13 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- UAICENSIS Lane, 1955: 415, fig. 19 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- UBOYASI Lane, 1955: 416, fig. 20 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Trinidad, Brazil (Santa Catarina). Type ♂, BMNH.
- VERBERIFERA Freeman, 1951: 130, figs. 267 (♂ genitalia), 355 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Osorno). Type ♂, BMNH.
- verbifera* Lane, 1962b: 547, *lapsus*.
- WINNERTZI Lane, 1948b: 253, figs. 12, 44 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.



- XAMASENSIS Lane, 1955: 417, fig. 21 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- YURIMAUESI Lane, 1955: 419, fig. 22 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Subgenus MYCETOPHILA Meigen

- ACARENSIS Lane, 1952b: 423, figs. 1-2 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- AMPLIPENNIS Freeman, 1951: 115, figs. 225-226 (♂ genitalia), 330 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- ARMATURA Freeman, 1951: 118, figs. 231-232 (♂ genitalia), 334 (wing). Type-locality: Chile, Chiloé, Punta. Distr. — Chile (Ñuble, Chiloé). Type ♂, BMNH.
- BANHUMAI Lane, 1952b: 424, figs. 3-4 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- BASALIS Freeman, 1951: 120, figs. 239-240 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Tucumán, Santa Cruz, Río Negro). Type ♂, BMNH.
- BIFIDA Freeman, 1954: 39 (nom. nov. for *fasciata* Freeman). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue).
- fasciata* Freeman, 1951: 121, figs. 242-243 (♂ genitalia), 339 (wing) (preocc. Meigen, 1804, now in *Rymosia*). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- BISETOSA Freeman, 1951: 108, fig. 208 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- BORACENSIS Lane, 1948a: 257, figs. 15, 49 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- BREVI-FURCATA Freeman, 1951: 108, fig. 209 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- BRUNNESCENS Freeman, 1951: 118, figs. 233-234 (♂ genitalia), 335 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Ñuble, Llanquihue, Chiloé). Type ♂, BMNH.
- CATHARINAE Edwards, 1932: 148. Type-locality: Brazil, Santa Catarina, Bom Retiro. Type ♀, BMNH.
- CONJUNCTA Freeman, 1951: 117, figs. 227, 229 (♂ genitalia), 332 (wing). Type-locality: Argentina, Río Negro, Lago Nahuel Huapí. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Ñuble, Llanquihue, Chiloé). Type ♂, BMNH.



- CURIAENSIS Lane, 1952b: 425, figs. 5-6 (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- DUANENSIS Lane, 1952b: 426, figs. 7-8 (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- FREEMANI Lane, 1948a: 256, fig. 14 (δ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type δ , MZSP.
- FUSCENS Freeman, 1951: 119, figs. 227 (δ genitalia), 337 (wing). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Rio Negro, Tierra del Fuego), Chile (Osorno, Llanquihue). Type δ , BMNH.
- GUANASI Lane, 1958b: 156, fig. 7 (δ genitalia). Type-locality: Argentina, Tucumán, Tafí del Valle, 2000 m. Type δ , IML.
- GUARAIASI Lane, 1952b: 427, figs. 9-10 (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- IHERINGI Lane, 1948a: 260. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Ref. — Lane, 1952b: 433. Type φ , MZSP.
- ILLITA Freeman, 1951: 122, figs. 244 (δ genitalia), 340 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type δ , BMNH.
- JUINENSIS Lane, 1952b: 427, figs. 11-12 (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Venezuela (Trujillo), Brazil (Santa Catarina), Argentina (Tucumán). Type δ , BMNH.
- MATHESONI Lane, 1948a: 252, figs. 8, 43 (δ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type δ , MZSP.
- MULTIPLEX Freeman, 1951: 13, figs. 220 (δ genitalia), 221 (φ terminalia), 327 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Tucumán, Río Negro). Type δ , BMNH.
- NERVITACTA Freeman, 1951: 116, figs. 227 (δ genitalia), 331 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type δ , BMNH.
- PASSEI Lane, 1952b: 430, figs. 13-14 (δ genitalia). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis (Juçaral). Type δ , MZSP.
- PICEA Freeman, 1951: 119, figs. 235-236 (δ genitalia), 336 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Osorno). Type δ , BMNH.
- SHAMANENSIS Lane, 1952b: 430, figs. 15-16 (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- SHANNONI Lane, 1948a: 254, figs. 7, 29, 45 (δ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type δ , MZSP.
- SPINIPES Freeman, 1951: 116, figs. 228 (anterior tibia), 332 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Angol). Type δ , BMNH.



- SPLENDIDA Lane, 1948a: 247, fig. 34 (wing), 28, 37, 39 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- STONEI Lane, 1952b: 431 (nom. nov. for *edwardsi* Lane). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia.
- edwardsi* Lane, 1948a: 248, figs. 4 (♂ genitalia), 32 (wing) (preocc. Lundström, 1913). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SUBCAPITATA Freeman, 1954: 39 (nom. nov. for *subfasciata* Freeman). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue, Ñuble, Chiloé).
- subfasciata* Freeman, 1951: 120, figs. 226, 241 (♂ genitalia) (preocc. Meigen, 1818, now in *Leia*). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- TACUENSIS Lane, 1952b: 432, figs. 17-18 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- THERESAE Edwards, 1932: 148. Type-locality: Brazil, Espírito Santo, Santa Teresa. Distr. — Brazil (Espírito Santo, São Paulo). Ref. — Lane, 1948a: 255, figs. 13, 48 (♂ genitalia). Type ♀, BMNH.
- TOBASI Lane, 1958c: 397, fig. 1 (♂ genitalia). Type-locality: Argentina, Neuquén, Mallo, 1700 m. Type ♂, MZSP.
- TRIANGULIFERA Freeman, 1951: 110, figs. 214 (♂ genitalia), 323 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- UAIANAI Lane, 1952b: 433, figs. 19-20 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Costa Rica, Brazil (São Paulo). Type ♂, MZSP.
- VAURASI Lane, 1958c: 398, fig. 2 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Trinidad, Panama, Brazil (Bahia, Santa Catarina). Type ♂, BMNH.

Subgenus OROMYCETA Lane

- Mycetophila*, subg. OROMYCETA Lane, 1955: 395. Type-species, *javaesi* Lane (orig. des.).
- ARAUCANA Lane, 1962b: 542, fig. 7 (♂ genitalia). Type-locality: Chile, Ñuble, 50 km e. of San Carlos. Type ♂, CAS.
- ARECUNAI Lane, 1955: 396, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- FERRAZI Lane, 1955: 397, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- JAVAESI Lane, 1955: 397, fig. 4 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Costa Rica, Trinidad, Peru, Brazil (Rio de Janeiro, Santa Catarina), Argentina (Salta). Type ♂, BMNH.



- SALTANENSIS Lane, 1955: 398, fig. 5 (♂ genitalia). Type-locality: Argentina, Salta, Urundel. Type ♂, IML.
- TUCUNENSIS Lane, 1955: 399, fig. 6 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- WYGODZINSKYI (Lane), 1947b: 457, figs. 7 (wing), 6 (♂ genitalia) (*Delopsis*). Type-locality: Brazil, Rio de Janeiro, km 47 of the road Rio-São Paulo. Distr. — Costa Rica, Brazil (Rio de Janeiro, Santa Catarina), Argentina (Salta). Type ♂, MZSP.

Unplaced and unrecognized MYCETOPHILA

- ANGUSTIFURCA Enderlein, 1940: 672, figs. 43 (wing), 44 (♂ genitalia). Type-locality: Juan Fernandez Isl., Masatierra. Ref. — Freeman, 1954: 32, pl. 2, g (♂ genitalia).
- APICATA Philippi, 1865: 622. Type-locality: Chile, Valdivia.
- ATRICORNIS Philippi, 1865: 622. Type-locality: Chile, Santiago.
- AVICULARIA Freeman, 1951: 109, fig. 210 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- BIFILA Freeman, 1951: 114, figs. 222 (♂ genitalia), 328 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- BLANCHARDI Enderlein, 1910a: 174 (nom. nov. for *punctipennis* Blanchard). Type-locality: Chile, Carelmapu.
punctipennis Blanchard, 1852: 345 (preocc. Stannius, 1831). Type-locality: Chile, Carelmapu.
- CHLOROCHROA Freeman, 1951: 117, figs. 230 (♂ genitalia), 333 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- COGNATA Philippi, 1865: 621. Type-locality: Chile, Valdivia.
- CORNUTA Freeman, 1951: 112, figs. 217 (♂ genitalia), 326 (wing). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- DIGITALIS Freeman, 1951: 123, fig. 247 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- DOLOSA Williston, 1896: 264. Type-locality: St. Vincent, W. I. Type ♂, BMNH.
- DOMINICA Curran, 1927: 1. Type-locality: Dominica, Laudet, 2100 ft. Type ♀, AMNH.
- FASCIPENNIS Philippi, 1865: 621. Type-locality: Chile, Santiago.
- FLABELLIFERA Freeman, 1951: 112, figs. 219 (♂ genitalia), 325 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- FLAVITHORAX Freeman, 1951: 126, fig. 255 (♂ genitalia). Type-locality: Chile, Llanquihue, Puerto Varas. Type ♂, BMNH.
- HETERONEURA Philippi, 1865: 622. Type-locality: Chile, Corral.



- INSECTA Freeman, 1951: 109, fig. 211 (♂ genitalia). Type-locality: Argentina, Rio Negro, Lago Gutiérrez. Type ♂, BMNH.
- LORIGERA Freeman, 1951: 105, figs. 202 (♂ genitalia), 317 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Type ♂, BMNH.
- MERDIGERA Knab & Zwaluwenburg, 1918: 140, pl. 8, figs. 1-3 (whole insect, cocoon). Type-locality: Puerto Rico, Aibonito. Type ♂, USNM.
- MESORPIINA Speiser, 1911: 238. Type-locality: Peru.
- NIGRESCENS Freeman, 1951: 110, figs. 213 (♂ genitalia), 322 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Type ♂, BMNH.
- NIGRIVENTRIS Philippi, 1865: 623. Type-locality: Chile, Valdivia.
- NODULOSA Williston, 1896: 264, pl. 8, fig. 20 (wing). Type-locality: St. Vincent, W. I. Syntypes ♂, BMNH.
- OBSCURIPENNIS Blanchard, 1852: 346. Type-locality: Chile, Coquimbo.
- ORDINISETA Freeman, 1951: 106, figs. 204 (♂ genitalia), 310 (wing). Type-locality: Argentina, Rio Negro, Lago Correntoso. Distr. — Argentina (Rio Negro), Chile (Llanquihue). Type ♂, BMNH.
- ORNATIPENNIS Blanchard, 1852: 344. Type-locality: Chile, Coquimbo.
- PALLIDA Johnson, 1919: Type-locality: Jamaica.
- PUCARAENSIS Duret, 1973b: 471, figs. 1-5 (wing, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará (Parque Nacional Lanin), 600 m. Type ♂, Coll. Duret.
- RAMOSA Freeman, 1951: 114, fig. 223 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Type ♂, BMNH.
- SEEGOI Duret, 1973a: 57, figs. 1 (wing), 2 and 5 (♂ genitalia). Type-locality: Argentina, Neuquén, Pucará (Parque Nacional Lanin). Type ♂, Coll. Duret.
- SIMPLEX Freeman, 1951: 121, figs. 245 (♂ genitalia), 336 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina (Rio Negro), Chile (Chiloé). Type ♂, BMNH.
- SIMPLICISTILA Freeman, 1951: 127, fig. 259 (♂ genitalia). Type-locality: Argentina, Rio Negro, Lago Correntoso. Type ♂, BMNH.
- SUBFUSCA Freeman, 1951: 102, figs. 189 (♂ genitalia), 190-191 (♀ terminalia), 311 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Type ♂, BMNH.
- TALARIS Freeman, 1951: 115, figs. 223 (♂ genitalia), 329 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Type ♂, BMNH.
- UNICORNUTA Freeman, 1951: 120, figs. 237 (♂ genitalia), 337 (wing). Type-locality: Chile, Valparaiso, Marga Marga. Distr. — Argentina (Rio Negro), Chile (Valparaiso). Type ♂, BMNH.
- UNISERIES Freeman, 1951: 106, figs. 205 (♂ genitalia), 320 (wing). Type-locality: Argentina, Rio Negro, Lago Gutiérrez. Type ♂, BMNH.
- VELIGERA Freeman, 1951: 111, figs. 212 (♂ genitalia), 216-217 (♀ terminalia), 324 (wing). Type-locality: Argentina, Rio Negro, Bariloche. Type ♂, BMNH.



VIRIDIPES Freeman, 1951: 104, figs. 200 (♂ genitalia), 201 (♀ terminalia), 316 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.

VITICOLLIS Blanchard, 1852: 345. Type-locality: Chile, Coquimbo.

VITIPENNIS Freeman, 1951: 129, figs. 263 (♂ genitalia), 351 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.

VOLTANS Lynch Arribálzaga, 1892: 429 (1892: 55). Type-locality: Argentina, Misiones.

Genus PHRONIA Winnertz

PHRONIA Winnertz, 1863: 857. Type-species, *rustica* Winnertz (Johannsen, 1909: 96) = *exigua* (Zetterstedt).

LONGINERVIS Freeman, 1951: 93, figs. 179 (♂ genitalia), 296 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Tucumán, Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.

Genus PLATUROCYPTA Enderlein

PLATUROCYPTA Enderlein, 1910b: 76. Type-species, *limbatifemur* Enderlein (orig. des.).

Neoepicypta Coher, 1950a: 114. Type-species, *Mycetophila punctum* Stannius (orig. des.).

NEOTROPICALIS Lane, 1947b: 450, fig. 3 (♀ terminalia). Type-locality: Brazil, São Paulo, Cajuru. Distr. — Trinidad, Brazil (São Paulo, Santa Catarina). Ref. — Lane, 1948a: 238, fig. 10 (♂ genitalia). Type ♀, MZSP.

Genus PLEUROGYMNUS Freeman

PLEUROGYMNUS Freeman, 1951: 92. Type-species, *fuscus* Freeman (orig. des.).

FUSCUS Freeman, 1951: 93, figs. 178 (♂ genitalia), 295 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Bío Bío). Type ♂, BMNH.

Genus SCEPTONIA Winnertz

SCEPTONIA Winnertz, 1863: 907. Type-species, *Mycetophila nigra* Meigen (Johannsen, 1909: 113).

BICOLORATA Lane, 1948a: 239, fig. 28 (wing). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Ref. — Lane, 1956b: 154, fig. 5 (♂ genitalia). Type ♀, MZSP.



- LONGICORNIS Enderlein, 1910a: 183. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Goiás, Rio de Janeiro, São Paulo, Santa Catarina). Ref. — Lane, 1947b: 452. Syntypes ♀, Stettin.
- ORNATIFEMORA Fisher, 1939: 235, pl. 13, figs. 7-8 (♂ genitalia). Type-locality: Costa Rica, San José. Type ♂, ANSP.
- FAIAGUENSIS Lane, 1956b: 155, fig. 6 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, MZSP.

Genus TRICHONTA Winnertz

- TRICHONTA Winnertz, 1863: 847. Type-species, *Mycetophila melanura* Staeger (Johannsen, 1909: 94).
- FASCIATA Freeman, 1951: 97, figs. 188-189 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Tucumán, Río Negro, Tierra del Fuego). Type ♂, BMNH.
- FUNEREA Freeman, 1951: 96, figs. 186-187 (♂ genitalia), 310 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- MAJOR Freeman, 1951: 95, figs. 182-183 (♂ genitalia), 308 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- NUBILPENNIS Freeman, 1951: 94, figs. 180-181 (♂ genitalia), 309 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- SIMILIS Freeman, 1951: 96, fig. 185 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- SPINIGERA Freeman, 1951: 95, fig. 184 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.

Genus ZYGOMYIA Winnertz

- ZYGOMYIA Winnertz, 1863: 901. Type-species, *Mycetophila vara* Staeger (Johannsen, 1909: 112).
- Ref. — Lane, 1951a (key to spp.).
- AGUARENSIS Lane, 1951a: 244, fig. 9 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- ARGENTINA Lane, 1961b: 13, fig. 5 (♂ genitalia). Type-locality: Argentina, Tucumán, Lacavera. Type ♂, MZSP.
- AURANTIACA Edwards, 1934: 365. Type-locality: Costa Rica, Farm La Caja, 8 km w. of San José. Type ♀, Hamburg.
- BICOLOR Edwards, 1934: 366. Type-locality: Bolivia, Mapiro, Lorenzopata. Distr. — Bolivia, Brazil (São Paulo). Refs. — Lane, 1948a: 240, fig. 3 (♂ genitalia), 1951a: 249, fig. 2 (♂ genitalia). Type ♀, Dresden.



- BRASILIANA Lane, 1947b: 452. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo, Santa Catarina). Ref. — Lane, 1951a: 241, fig. 6 (♂ genitalia). Type ♀, MZSP.
- CHAVANTESI Lane, 1951a: 239, fig. 1 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, MZSP.
- FREEMANI Lane, 1951a: 248, fig. 12 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- GOLBACHI Lane, 1961b: 14, fig. 6 (♂ genitalia). Type-locality: Argentina, Tucumán, Lacavera. Type ♂, MZSP.
- HEROS Lane, 1951a: 248, fig. 13 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- HERTELI Lane, 1951a: 244, fig. 10 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica da Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.
- LAFOONI Lane, 1951a: 246, fig. 11 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- MODESTA Lane, 1948a: 244, figs. 11, 57 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- ORNATIPENNIS Lane, 1948a: 241, figs. 5 (♂ genitalia), 35 (mesosome), 42 (9th tergite). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo), Argentina (Misiones). Type ♂, MZSP.
- PLAUMANNI Lane, 1951a: 243, fig. 8 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- TAPUIAI Lane, 1951a: 241, fig. 7 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, MZSP.

Subfamily SCIOPHILINAE

Tribe GNORISTINI

Genus AUSTROSYNAPHA Tonnoir

- AUSTROSYNAPHA Tonnoir, 1929: 607. Type-species, *hirta* Tonnoir (orig. des.). Ref. — Duret, 1973c: 76 (partial key to spp.).
- ARGENTINENSIS Duret, 1973c: 69, figs. 2-3 (♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.
- BEJARANOI Duret, 1973c: 69, figs. 4-5 (♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.
- BIDENTATA Duret, 1975a: 311, fig. 1 (♂ genitalia). Type-locality: Argentina, Neuquén, Pucará, Parque Nacional Lanín, Lago Lácar. Type ♂, Coll. Duret.



- DENTATA Freeman, 1951: 72, figs. 139-140 (δ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén). Type δ , BMNH.
- FILICAUDA (Edwards), 1934: 364 (*Synapha*). Type-locality: Paraguay, La Cordillera, San Bernardino. Type δ , WIEN.
- FORCIPATA Freeman, 1951: 71, fig. 137 (δ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén), Chile (Ñuble, Llanquihue). Type δ , BMNH.
- FREEMANI Lane, 1960a: 382, fig. 6 (δ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type δ , MZSP.
- FURCATA Lane, 1954a: 109, fig. 13 (δ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Jaraguá). Type δ , MZSP.
- IBARRAI Duret, 1973c: 71, figs. 6-7 (δ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type δ , Coll. Duret.
- LAMELLATA Freeman, 1951: 73, fig. 144 (δ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén), Chile (Llanquihue). Type δ , BMNH.
- MARTINEZI Duret, 1973c: 71, figs. 8-9 (δ genitalia). Type-locality: Argentina, Neuquén, Isla Victoria. Type δ , Coll. Duret.
- NAUMANNI Duret, 1973c: 73, figs. 11-12 (δ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type δ , Coll. Duret.
- PAULISTANA Lane, 1956a: 123, fig. 8 (δ genitalia). Type-locality: Brazil, São Paulo, Embu. Type δ , MZSP.
- PROSENI Duret, 1973c: 74, fig. 13 (δ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type δ , Coll. Duret.
- QUADRISSETOSA Freeman, 1951: 72, figs. 138 (δ genitalia), 287 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén). Type δ , BMNH.
- REDUCTA Freeman, 1951: 70, fig. 135 (δ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Neuquén, Chubut), Chile (Ñuble, Llanquihue, Chiloé). Type δ , BMNH.
- ROSSI Lane, 1962b: 534, fig. 3 (δ genitalia). Type-locality: Chile, Osorno, 20 km e. Puyehué. Type δ , CAS.
- SCHAJOVSKOYI Duret, 1973c: 74, figs. 14-15 (δ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type δ , Coll. Duret.
- SPINATA Lane, 1954a: 108, fig. 12 (δ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type δ , MZSP.
ssp. SOUZAI Lane, 1960a: 382, fig. 7 (δ genitalia). Type-locality: Brazil, Minas Gerais, Serra do Cipó. Type δ , MZSP.
- SPINIFERA Freeman, 1951: 73, figs. 141-142 (δ genitalia). Type-locality: Chile, Chiloé, Castro. Distr. — Argentina (Río Negro, Neuquén), Chile (Ñuble, Chiloé). Type δ , BMNH.
- TRUNCATA Freeman, 1951: 73, fig. 143 (δ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Neuquén), Chile (Bío Bío, Ñuble). Type δ , BMNH.

- UNGUICULATA Freeman, 1951: 71, fig. 136 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- YIVOFFI Duret, 1973c: 76, figs. 16-18 (♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.

Genus COELOSIA Winnertz


- COELOSIA Winnertz, 1863: 796. Type-species, *Boletina flava* Staeger (Johannsen, 1909: 86).
- FLAVITHORAX Freeman, 1951: 64, fig. 285 (wing). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro). Type ♂, BMNH.
- NEOTROPICA Lane, 1959b: 111, fig. 5 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- SP. Freeman, 1951: 64. Argentina, Río Negro: Bariloche and Puerto Blest. ♀, BMNH.

Genus DZIEDZICKIA Johannsen

- Hertwigia* Dziedzicki, 1885: 166 (preocc. Schmidt, 1880). Type-species, *marginata* Dziedzicki (mon.).
- DZIEDZICKIA Johannsen, 1909: 44 (nom. nov. for *Hertwigia* Dziedzicki). Type-species, *Hertwigia marginata* Dziedzicki (aut.).
- Sytemna*, authors, not Winnertz.
- ABSYRTA Lane, 1954a: 96, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- ARMATA Freeman, 1951: 65, figs. 127 (♂ genitalia), 283 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.
- ASPASIA Lane, 1954a: 96, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- BASALIS Freeman, 1951: 66, fig. 126 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Type ♂, BMNH.
- BEATRICEA Lane, 1960a: 380, fig. 4 (♂ genitalia). Type-locality: Trinidad, Navy Base. Type ♂, MZSP.
- CAIPORA Lane, 1961a: 248. Type-locality: Brazil, Amapá, Rio Felício. Type ♀, MZSP.
- CLIO Lane, 1954a: 97, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- COHERI Lane, 1954a: 97, fig. 4 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- CRYPTURA Edwards, 1941: 308. Type-locality: Brazil, Mato Grosso, Salobra. Distr. — Brazil (Mato Grosso to São Paulo). Type ♂, BMNH.



- DIANA Lane, 1959b: 112, fig. 6 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- DUBITANS Lane, 1954a: 98, fig. 5 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- EDWARDSIANA Lane, 1954a: 99. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♀, MZSP.
- ENDYMION Lane, 1959b: 113, fig. 8 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- FIEBRIGI Edwards, 1934: 364. Type-locality: Paraguay, La Cordillera, San Bernardino. Type ♂, WIEN.
- FLAVONIGRA Lane, 1954a: 100. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♀, BMNH.
- FREEMANI Lane, 1954a: 100, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Juquiá. Type ♂, MZSP.
- FUNEREA Freeman, 1951: 66, fig. 128 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- GALINDOI Lane, 1961b: 10. Type-locality: Panama, Darién, Camp Tacarcuna. Type ♀, MZSP.
- GLORIOSA Lane, 1959b: 114, fig. 8 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- HYPSEIPLE Lane, 1954a: 101, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo), Argentina (Misiones). Type ♂, MZSP.
- IARA Lane, 1961a: 248. Type-locality: Brazil, Amapá, Igarapé do Sucuriçu. Type ♀, MZSP.
- INTERMEDIA Lane, 1954a: 102, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Costa Rica (San Mateo), Brazil (Amapá, São Paulo). Type ♂, MZSP.
- IONICA Lane, 1959b: 115, fig. 9 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- JANUS Lane, 1959b: 115, fig. 10 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- JURUPARI Lane, 1961a: 249. Type-locality: Brazil, Amapá, Rio Felício. Type ♂, MZSP.
- LIRIOPE Lane, 1954a: 102. Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- MEDEA Lane, 1954a: 103, fig. 9 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Distr. — Brazil (Goiás, São Paulo), Argentina (Misiones). Type ♂, MZSP.
- METALLICA Lane, 1954a: 103, fig. 10 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- NIGRA Fisher, 1939: 229. Type-locality: Costa Rica, San José. Type ♂, ANSP.




- OLAMPENSIS Lane, 1961a: 250, fig. 2 (♂ genitalia). Type-locality: Brazil, Amapá, Rio Amaparl. Type ♂, MZSP.
- PUBERICORNIS Edwards, 1933: 308. Type-locality: Bolivia, Mapiri, Lorenzopata. Type ♂, Dresden.
- SACI Lane, 1961b: 11. Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♀, MZSP.
- VARIABILIS Lane, 1954a: 104, fig. 11 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, MZSP.

Genus SCHNUSEA Edwards

- SCHNUSEA Edwards, 1933: 307. Type-species, *sessilis* Edwards (orig. des.). Ref. — Lane & Coher, 1950 (key to spp.).
- AGUARASI Lane & Coher, 1950: 279, figs. 1-3 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- BAREI Lane & Coher, 1950: 281, figs. 4-6 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- CALABII Lane & Coher, 1950: 281, figs. 7-9 (♂ genitalia). Type-locality: Brazil, São Paulo, Cajuru. Type ♂, MZSP.
- DESANEI Lane & Coher, 1950: 282, figs. 10-12 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Distr. — Brazil (Goiás, São Paulo), Argentina (Misiones). Type ♂, MZSP.
- SESSILIS Edwards, 1933: 308. Type-locality: Peru, mouth of Pachitea River, 150 m. Distr. — Peru, Brazil (Amapá, Goiás), Argentina (Salta). Type ?, Dresden.

Genus SYNAPHA Meigen

- SYNAPHA Meigen, 1818: 227. Type-species, *fasciata* Meigen (mon.).
- FLAVIPALPIS Freeman, 1951: 69, figs. 130 (midtibial sense organ), 133 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- FULVA Lane, 1954a: 105. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil (Amapá, Rio de Janeiro). Type ♂, MZSP.
- FUMIPENNIS Freeman, 1951: 69, figs. 129 (midtibial sense organ), 134 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Distr. — Argentina (Chubut), Chile (Llanquihue). Type ♂, BMNH.
- FUNEREA Freeman, 1951: 68, figs. 131 (midtibial sense organ), 132 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro, Chubut, Tierra del Fuego), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- IDALIS Lane, 1954a: 106. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.



- JUNO Lane, 1954a: 106. Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- MICHELBACHERI Lane, 1962b: 533, fig. 2 (♂ genitalia). Type-locality: Chile, Ñuble, 40 km e. of San Carlos. Type ♂, CAS.
- MORENOI Lane, 1963: 7, fig. 2 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche (Llao-Llao). Type ♂, MZSP.
- NIGRESCENS Lane, 1954a: 107. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.
- RUFESCENS Edwards, 1932: 140. Type-locality: Brazil, Rio de Janeiro, Petrópolis and Rio de Janeiro. Syntypes ♂, BMNH.

Unplaced Gnoristini

- LATICORNIS Enderlein, 1910a: 163 (*Syntemna*). Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.

Tribe LEIINI

(Tetragoneurini; incl. Selkirkiini Enderlein, 1940: 670).

Genus APHRASTOMYIA Coher & Lane


- APHRASTOMYIA Coher & Lane, 1949: 485. Type-species, *cramptoni* Coher & Lane (orig. des.).
- CERQUEIRAI Lane, 1961b: 10. Type-locality: Brazil, Amazonas, Manaus (Igarapé do Francês). Type ♀, MZSP.
- CRAMPTONI Coher & Lane, 1949: 486, figs. 1 (head), 2 (antenna), 3 (thoracic sclerites), 4 (wing), 5 (8th abdominal tergite), 6 (8th abdominal sternite), 7-9 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SHANNONI Lane, 1956a: 119, fig. 1 (♂ genitalia). Type-locality: Peru, Loreto, Iquitos. Distr. — Peru (Loreto), Brazil (Amapá). Type ♂, USNM.

Genus DOCOSIA Winnertz

- DOCOSIA Winnertz, 1863: 802. Type-species, *Mycetophila sciarina* Meigen (Johannsen, 1909: 92).
- CUZCOENSIS Edwards, 1933: 321. Type-locality: Peru, Cuzco, 3500-4200 m. Type ♂, Dresden.
- PAMMELA Edwards, 1933: 322. Type-locality: Peru, Río Urubamba, 3000 m; Paraguay, Hohenau (Syntypes ♂ ♀, Dresden); Paraguay, La Cordillera, San Bernardino (on *Caesaria silvestris* inflorescences) (Syntypes ♂ ♀, WIEN).

Genus LEIA Meigen

- Leia* Meigen, 1818: 253. Type-species, *fascipennis* Meigen (Curtis, 1837: pl. 645).



Leja, error or emend.

Lia, emend.

Glaphyroptera Winnertz, 1863: 781 (preocc. Heer, 1852). Type-species, *Leia fascipennis* Meigen (Coquillett, 1910: 547).

Neoglaphyroptera Osten Sacken, 1878: 10 (nom. nov. for *Glaphyroptera* Winnertz). Type-species, *Leia fascipennis* Meigen (aut.).

Ref. — Shaw & Lane, 1950 (key to spp.).

ACZELI Lane, 1959a: 186. Tkp-species, Argentina, Tucumán, Lacavera. Type ♀, MZSP.

ALBICOXA Enderlein, 1910a: 190. Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.

AMABILIS (Williston), 1900: 219 (*Neoglaphyroptera*). Type-locality: Mexico, Morelos, Cuernavaca, and Veracruz, near Medellín. Syntypes ♂ ♀, BMNH.

AMAPAENSIS Lane, 1959b: 116, fig. 13 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Distr. — Brazil (Pará, Amapá). Type ♂, MZSP.

ANDIRAI Lane, 1950: 130. Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (Rio de Janeiro, São Paulo), Argentina (Buenos Aires). Type ♀, MZSP.

APICALIS (Kertész), 1902: 581 (*Neoglaphyroptera*). Type-locality: Peru, Callanga. Type ♂, Budapest.

APINAGEI Lane, 1950: 125, fig. 13 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Distr. — Brazil (São Paulo), Argentina (Misiones). Type ♂, MZSP.

ARUACI Lane, 1950: 122, figs. 8-11 (♂ genitalia). Type-locality: Brazil, Goiás, Corumbá. Type ♂, MZSP.

AXILLIPUNCTUM Enderlein, 1910a: 198. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Pará, Amapá, Santa Catarina), Argentina (Misiones). Ref. — Lane, 1959a: 186, fig. 5 (♂ genitalia). Type ♀, Stettin.

BILUNULA Wiedemann, 1828: 65. Type-locality: "Brazil". Distr. — Brazil, Argentina (Tucumán, Buenos Aires).

BIPARTITA (Lynch Arribáizaga), 1892: 409 (1892: 35), pl. 1, fig. 19 (whole insect) (*Glaphyroptera*). Type-locality: Argentina, Buenos Aires: Baradero and Las Conchas. Distr. — Argentina (Buenos Aires).

BIPUNCTATA Fisher, 1939: 231, pl. 13, fig. 5 (♂ genitalia). Type-locality: Costa Rica, San José. Type ♂, ANSP.

BIVITTATA Say, 1829: 152 (1859: 351). Type-locality: U.S.A., Indiana. Distr. — U.S.A. (Minnesota to Maine, south to Kansas and North Carolina), Costa Rica.

ssp. TROPICALIS Fisher, 1939: 233. Type-locality: Costa Rica. Type ♂, ANSP.



- ssp. PUNCTIFORMIS Fisher, 1939: 234. Type-locality: Costa Rica. Type ♂, ANSP.
- BORGMEIERI Lane, 1959b: 117, fig. 14 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.
- CAYAPOI Lane, 1950: 127. Type-locality: Brazil, Goiás, Corumbá. Type ♀, MZSP.
- CINCTICAUDA Enderlein, 1910a: 192. Type-locality: Brazil, Santa Catarina. Syntypes ♂, Stettin.
- CIRCUMFERA Fisher, 1939: 231. Type-locality: Costa Rica, San José. Type ♂, ANSP.
- COLLARIGERA Enderlein, 1910a: 194. Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.
- COMPLETA (Kertész), 1902: 577 (*Neoglaphyoptera*). Type-locality: Peru, Callanga; Bolivia, Songo. Syntypes ♂, Budapest.
- CONCINNA (Williston), 1896: 259 (*Neoglaphyoptera*). Type-locality: St. Vincent, W. I. Distr. — Mexico to Panama, St. Vincent. Syntypes ♀, BMNH.
- COSTARICENSIS Fisher, 1940: 8 (nom. nov. for *analis* Fisher). Type-locality: Costa Rica, San José.
 analis Fisher, 1939: 232, pl. 13, fig. 3 (♂ genitalia) (preocc. Meigen, 1818, now in *Boletina*). Type-locality: Costa Rica, San José. Type ♂, ANSP.
- DICHROMA Freeman, 1951: 78. Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Chile (Llanquihue, Chiloé). Type ♀, BMNH.
- DISGREGA Edwards, 1933: 318. Type-locality: Peru, Río Urubamba, 3000 m. Type ♂, Dresden.
- DIVERSICORNIS (Kertész), 1902: 576 (*Neoglaphyoptera*). Type-locality: Peru, Callanga. Distr. — Peru, Brazil (Mato Grosso). Type ♂, Budapest.
?fuscicornis Edwards, 1941: 309. Type-locality: Brazil, Mato Grosso, Salobra. Type ♀, BMNH.
- DIVERSIPES Edwards, 1933: 310. Type-locality: Bolivia, Mapiri, Cordilleras, 4000-5000 m. Distr. — Bolivia (La Paz), Argentina (Salta). Ref. — Lane, 1959a: 187. Type ♂, Dresden.
- ELEGANS (Kertész), 1902: 579 (*Neoglaphyoptera*). Type-locality: Peru, Callanga. Distr. — Peru, Brazil (Rio de Janeiro). Type ♂, Budapest.
- FALCULATA Edwards, 1933: 313. Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil (São Paulo, Paraná), Argentina (Buenos Aires). Type ♂, BMNH.
- FASCIATA (Kertész), 1902: 574, fig. (wing) (*Neoglaphyoptera*). Type-locality: Peru, Vilcanota and Callanga; Bolivia, Suapi. Distr. — Peru, Bolivia, Brazil (Rio de Janeiro, São Paulo, Paraná, Santa Catarina), Paraguay (La Cordillera), Argentina (Buenos Aires). Syntypes ♂ ♀, Budapest.
- FISHERAE Shaw, 1950: 113, fig. 2 (♂ genitalia). Type-locality: Brazil, Paraná, Rio Negro.

- FLAVOSCUTELLATA (Lynch Arribálzaga), 1892: 410 (1892: 36) (*Glaphyoptera*).
Type-locality: Argentina, Buenos Aires, Las Conchas. Distr. —
Brazil (Minas Gerais, São Paulo), Paraguay (La Cordillera),
Argentina (Buenos Aires).
- GUAYCURUSI Lane, 1950: 124, fig. 12 (♂ genitalia). Type-locality: Brazil,
Santa Catarina, Nova Teutônia. Type ♂, MZSP.
- HALTERATA (Kertész), 1902: 575 (*Neoglaphyoptera*). Type-locality: Peru,
Vilcanota and Callanga. Distr. — Peru, Paraguay (La Cordillera),
Brazil (São Paulo, Rio de Janeiro, Goiás), Argentina (Misiones).
Syntypes ♂, Budapest.
- ?*triciniventris* Enderlein, 1910a: 196. Type-locality: Brazil, Santa
Catarina. Type ♀, Stettin.
- INCOMPLETA (Curran), 1928: 13, fig. 2 (wing) (*Boletina*). Type-locality:
Puerto Rico, Adjuntas. Distr. — Puerto Rico, Brazil (Mato Grosso,
Minas Gerais). Type ♀, AMNH.
- biamputata* Edwards, 1933: 317. Type-locality: Brazil, Minas Gerais,
Araguaí. Type ♂, BMNH.
- salobrensis* Edwards, 1941: 310. Type-locality: Brazil, Mato Grosso,
Salobra. Type ♀, BMNH.
- INTERRUPTA (Kertész), 1901b: 495 (*Neoglaphyoptera*). Type-locality: Peru,
Callanga. Type ♂, Budapest.
- JUNAI Lane, 1950: 128. Type-locality: Brazil, São Paulo, Salesópolis, Estação
Biológica de Boracéia. Type ♀, MZSP.
- LEUCOCERA Edwards, 1933: 311. Type-locality: Bolivia, Mapiri, Sarampuni,
700 m. Distr. — Peru, Bolivia. Type ♂, Dresden.
- MALLEOLUS Freeman, 1954: 28, pl. 2, fig. 3, f (♂ genitalia). Type-locality:
Juan Fernandez Isl., Masatierra, Bahía Cumberland, Grutas de los
Patriotas. Type ♂, Universidad de Chile, Santiago.
- MONOLEUCA Edwards, 1933: 312. Type-locality: Brazil, São Paulo. Distr. —
Brazil (Goiás, Rio de Janeiro, São Paulo, Rio Grande do Sul),
Argentina, Misiones. Type ♂, BMNH.
- MUCHTLERI Curran, 1928: 14, fig. 3 (wing). Type-locality: Puerto Rico,
Adjuntas. Type ♂, AMNH.
- NIGRICAUDA Edwards, 1933: 318. Type-locality: Bolivia, Mapiri, San Carlos,
800 m. Type ♂, Dresden.
- NIGROSPLeniATA (Lynch Arribálzaga), 1892: 411 (1892: 37) (*Glaphyoptera*).
Type-locality: Argentina, Buenos Aires: Baradero and Las Conchas.
- NITENS (Williston), 1896: 259, pl. 8, fig. 13 (wing) (*Neoglaphyoptera*).
Type-locality: St. Vincent, W. I. Syntypes ♂ ♀, BMNH.
- OBLECTABILIS (Loew), 1869: 146 (Cent. 9, n° 31) (*Glaphyoptera*). Type-
locality: U.S.A., "Middle States". Distr. — U.S.A. (Wisconsin
to Maine, south to Kansas and North Carolina), Costa Rica. Ref.
— Fisher, 1939: 231.
- OLIVEIRAI Lane, 1959a: 188, fig. 4 (♂ genitalia). Type-locality: Brazil, Pará,
Cachimbo. Distr. — Brazil (Pará, Amapá). Type ♂, MZSP.



- PARANENSIS Edwards, 1933: 313. Type-locality: Brazil, Paraná, Rio Negro. Distr. — Brazil (Goiás, Rio de Janeiro, São Paulo, Paraná), Argentina (Misiones). Ref. — Lane, 1950: 122, figs. 4-7 (♂ genitalia). Type ♂, BMNH.
- PAULENSIS Edwards, 1933: 312. Type-locality: Brazil, São Paulo. Distr. — Brazil (Amapá, Rio de Janeiro, São Paulo, Rio Grande do Sul), Panama. Ref. — Lane, 1950: 121, figs. 1-3 (♂ genitalia). Type ♂, BMNH.
- PEDIFERA Edwards, 1933: 316. Type-locality: Venezuela, Ciudad Bolívar. Type ♂, BMNH.
- PICTICORNIS (Kertész), 1902: 578 (*Neoglaphyoptera*). Type-locality: Peru, Callanga. Distr. — Panama, Brazil (Amapá, Goiás, Minas Gerais, São Paulo). Type ♂, Budapest.
- PLAUMANNI Lane, 1959a: 189. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♀, MZSP.
- ?FOECILOPTERA Philippi, 1865: 623. Type-locality: Chile, Valdivia.
- PUNCTATA Bellardi, 1861: 5, pl. 3, fig. 3 (whole insect). Type-locality: Mexico, Tuxpango, near Orizaba. Type ♂, Università di Torino.
- RUFITHORAX Freeman, 1951: 78, fig. 153 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Type ♂, BMNH.
- SCHMIDTI Fisher, 1939: 230, pl. 13, fig. 4 (♂ genitalia). Type-locality: Costa Rica, San José. Type ♂, ANSP.
- SCHNUSEI Edwards, 1933: 314. Type-locality: Bolivia, Mapiri, San Antonio, 1000 m. Distr. — Bolivia (La Paz), Brazil (São Paulo). Type ♂, Dresden.
- SETOSICAUDA Edwards, 1933: 316. Type-locality: Bolivia, Mapiri, San Antonio, 1000 m. Type ♂, Dresden.
- SMITHI Edwards, 1933: 318. Type-locality: Mexico, Tabasco, Frontera. Distr. — Mexico (Tabasco), Costa Rica. Type ♂, BMNH.
- SPINIFERA Edwards, 1933: 316. Type-locality: Bolivia, Mapiri, San Carlos, 800 m. Distr. — Peru, Bolivia. Type ♂, Dresden.
- STONEI Lane, 1958b: 152 (nom. nov. for *incompleta* Edwards). Type-locality: Bolivia, Mapiri, Sarampiuni, 700 m.
incompleta Edwards, 1933: 314 (preocc. Curran, 1928). Type-locality: Bolivia, Mapiri, Sarampiuni, 700 m. Type ♂, Dresden.
- SUBMACULIPENNIS Freeman, 1954: 39 (nom. nov. for *maculipennis* Freeman). Type-locality: Chile, Valparaiso, Marga Marga.
maculipennis Freeman, 1951: 77, figs. 151-152 (♂ genitalia), 304 (wing) (preocc. Say, 1824). Type-locality: Chile, Valparaiso, Marga Marga. Type ♂, BMNH.
- TRUNCATOVENOSA Enderlein, 1910a: 199. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Santa Catarina), Argentina (Salta). Type ♂, Stettin.



Genus LEIELLA Enderlein

- LEIELLA** Enderlein, 1910a: 200. Type-species, *ochreocalcar* Enderlein (orig. des.).
Ref. — Lane, 1954d (key to spp.).
- ARNAUDI** Lane, 1962d: 536, fig. 4 (♂ genitalia). Type-locality: Chile, Osorno, 10 km e. of Puyehué. Distr. — Chile (Osorno), Argentina (Río Negro). Type ♂, CAS.
- BICOLOR** (Fisher), 1939: 234 (*Leia*). Type-locality: Costa Rica, San José. Type ♂, ANSP.
- CATHARINENSIS** Lane, 1954d: 451, fig. 2 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- DISTINCTA** Freeman, 1951: 79, fig. 303 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- FULVA** Lane, 1954d: 452, fig. 3 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- JARUSI** Lane, 1952d: 138. Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♀, MZSP.
- OCCHREOCALCAR** Enderlein, 1910a: 200, fig. 3 (wing). Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Santa Catarina), Argentina (Chaco, Salta, Tucumán). Type ♀, Stettin.
- SHANONNI** Lane, 1954d: 453, fig. 4 (♂ genitalia). Type-locality: Panama, Sabanas. Type ♂, USNM.
- UNICINCTA** Edwards, 1933: 319. Type-locality: Peru, mouth of Río Pachitea, 150 m. Distr. — Peru (Cuzco), Brazil (Goiás). Refs. — Lane, 1952a: 137, fig. 5 (♂ genitalia); 1954d: 450, fig. 1 (♂ genitalia). Type ♀, Dresden.
- ZONALIS** Edwards, 1931: 259. Type-locality: Guyana, Berbice. Distr. — Guyana, Argentina (Santa Fé, Chaco, Salta, Tucumán). Type ♀, BMNH.

Genus MEGOPHTHALMIDIA Dziedzicki

- MEGOPHTHALMIDIA** Dziedzicki, 1889: 404, 525 (as *Megophtalmidia*, pp. 526, 532). Type-species, *zugmeyeriae* Dziedzicki (mon.) = *crassicornis* (Curtis).
- BICOLOR** Edwards, 1934: 363. Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Brazil (Santa Catarina), Paraguay. Syntypes ♂ ♀, WIEN.
- DIVERGENS** Edwards, 1932: 141. Type-locality: Brazil, Santa Catarina, Bom Retiro. Distr. — Peru (Junín), Brazil (Santa Catarina), Argentina (Tucumán). Type ♂, BMNH.
- EDWARDSIANA** Lane, 1954b: 106, fig. 1 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.



- EQUATORIANA Lane, 1962a: 11, fig. 2 (♂ genitalia). Type-locality: Ecuador, Pichincha, 105 km s. of Cayamba, 2470 m. Type ♂, CAS.
- FREEMANI Lane, 1954b: 108, fig. 2 (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- NIGRA Freeman, 1951: 82, fig. 291 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- PLAUMANNI Lane, 1954b: 109. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Genus PARALEIA Tonnoir

- PARALEIA Tonnoir, 1929: 611. Type-species, *fulvescens* Tonnoir (orig. des.).
- Selkirkius* Enderlein, 1940: 670. Type-species, *nephrodops* Enderlein (orig. des.). Ref. — Freeman, 1954: 26 (syn.).
- ANTARCTICA (Bigot), 1888: 12, pl. 3, fig. 1 (wing) (*Boletina*). Type-locality: "Cap Horn". Distr. — Argentina (Río Negro, Tierra del Fuego). Ref. — Freeman, 1951: 75, fig. 146 (♂ genitalia), 306 (wing). Syntypes ♂ ♀, MNHN.
- CASTANEA Freeman, 1951: 77, fig. 150 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- CAUDATA Edwards, 1933: 310. Type-locality: Bolivia, Mapiri, Cordilleras, 4000-5000 m. Type ♂, Dresden.
- CINEREA Freeman, 1951: 76, figs. 147 (♂ genitalia), 306 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- FALCIGERA Freeman, 1951: 76, fig. 148 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Type ♂, BMNH.
- FUNEREA Freeman, 1951: 76, fig. 149 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- NEPHRODOPS (Enderlein), 1940: 671, figs. 41-42 (wing, ♂ genitalia) (*Selkirkius*). Type-locality: Juan Fernandez Isl., Masatierra. Ref. — Freeman, 1954: 28, pl. 2, figs. c, d (♂ genitalia).
- NUBILPENNIS (Walker), 1837: 334 (*Leia*). Type-locality: Straits of Magellan, Port Famine. Distr. — Juan Fernandez Isl., Ecuador, Chile (Osorno, Magallanes), Argentina (Santa Cruz, Chubut, Río Negro, Tierra del Fuego). Ref. — Freeman, 1951: 75, figs. 145 (♂ genitalia), 305 (wing). Type, BMNH.
- PERUVIANA Edwards, 1933: 309. Type-locality: Peru, Mamara. Distr. — Peru, Argentina (Río Negro). Type ♂, Dresden.
- SP. Freeman, 1951: 77. Type-locality: Argentina, Río Negro, Puerto Blest. ♀, BMNH.



Genus PROCYCLONEURA Edwards

- PROCYCLONEURA Edwards, 1932: 140. Type-species, *paranensis* Edwards (mon.).
- FURCATA Freeman, 1951: 80, fig. 154 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Tucumán, Río Negro). Type ♂, BMNH.
- MOROSA Edwards, 1933: 320. Type-locality: Peru, Mamara. Syntypes ♂ ♀, Dresden.
- PARANENSIS Edwards, 1932: 140. Type-locality: Brazil, Paraná, Río Negro. Distr. — Brazil (Paraná), Argentina (Misiones). Syntypes ♂ ♀, BMNH.
- SIMILIS Freeman, 1951: 81, fig. 155 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue, Osorno). Type ♂, BMNH.

Genus RONDANIELLA Johannsen

- RONDANIELLA Johannsen, 1909: 66. Type-species, *Leia variegata* Winnertz (orig. des.).
- SORORCULA (Loew), 1869: 147 (Cent. 9, n° 32) (*Leia*). Type-locality: U.S.A., New York. Distr. — Canada (British Columbia), U.S.A. (Wisconsin to Maine, south to Iowa and North Carolina), Mexico (Jalisco). Ref. — Lane, 1962a: 11.
- abbreviata* Loew, 1869: 147 (Cent. 9, n° 33) (*Leia*). Type-locality: U.S.A., "Middle States".

Genus TETRAGONEURA Winnertz

- TETRAGONEURA Winnertz, 1846: 18. Type-species, *distincta* Winnertz (Johannsen, 1909: 34, 35) = *sylvatica* (Curtis).
- ANDINA Lane, 1962b: 537, fig. 5 (♂ genitalia). Type-locality: Chile, Ñuble, 40 km e. of San Carlos. Type ♂, CAS.
- ARAUENSIS Lane, 1952c: 402, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.
- ARDEICEPS Freeman, 1951: 85, figs. 161 (♂ genitalia), 289 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Chubut, Tierra del Fuego). Type ♂, BMNH.
- BACILLIGER Edwards, 1932: 144, fig. 5 (♂ genitalia). Type-locality: Brazil, Paraná, Río Negro. Distr. — Peru, Brazil (Paraná). Type ♂, BMNH.
- ?BECKERI (Enderlein), 1910a: 168 (*Parastemma*). Type-locality: Brazil, Santa Catarina. Type ♀, Stettin.
- BIFIDA Freeman, 1951: 85, fig. 162 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.



- BORGMEIERI Edwards, 1932: 142, fig. 1 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Distr. — Brazil (Rio de Janeiro, São Paulo), Argentina (Corrientes). Type ♂, BMNH.
- BURENARI Lane, 1952c: 403, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Saiesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- CAETENSIS Lane, 1952c: 404, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- CALOPUS (Bigot), 1888: 14, pl. 3, fig. 3 (whole insect) (*Sciophila*). Type-locality: "Cap Horn".
var. MINOR Enderlein, 1910a: 155. Type-locality: Brazil, Santa Catarina. Syntypes ♂, Stettin.
- CONCURRENS Edwards, 1933: 321. Type-locality: Bolivia, La Paz. Type ♂, Dresden.
- DENTATA Freeman, 1951: 85, fig. 160 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- DERUNEI Lane, 1952c: 405, fig. 4 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo), Argentina (Tucumán). Type ♂, MZSP.
- ERAMANAI Lane, 1952c: 405, fig. 5 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- FANIPENNIS Lane, 1952c: 406, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Amapá, São Paulo). Type ♂, MZSP.
- FLAVICAUDA Edwards, 1941: 310. Type-locality: Brazil, Mato Grosso, Salobra. Type ♂, BMNH.
- FLAVICINCTA Freeman, 1951: 84, fig. 158 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- GALEA Freeman, 1951: 84, fig. 159 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén), Chile (Osorno, Llanquihue). Type ♂, BMNH.
- GUAJAENSIS Lane, 1952c: 406, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- HUARPEI Lane, 1952c: 407, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- IAUNAI Lane, 1952c: 408, fig. 9 (♂ genitalia). Type-locality: Costa Rica, San José. Distr. — Costa Rica, Peru. Type ♂, USNM.
- NIGRIPALPIS Freeman, 1951: 87, fig. 168 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- NOCTICOLOR Edwards, 1932: 146, fig. 6 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Type ♂, BMNH.
- PECTINATA Freeman, 1951: 88, figs. 169 (♂ genitalia), 290 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.



- PERUANA Kertész, 1909: 141. Type-locality: Peru, Callanga. Type ♂, Budapest.
- POLLUX Freeman, 1951: 86, fig. 164 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Argentina (Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.
- RIVETI (Edwards), 1919: 146, fig. 6 (wing) (*Megophthalmidia*). Type-locality: Ecuador, El Pelado, 4151 m. Syntypes ♂ ♀, MHNH (examined by Dr. L. Matile). N. comb.
- SIMILLIMA Freeman, 1954: 39 (nom. nov. for *similis* Freeman). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue).
- similis* Freeman, 1951: 86, fig. 166 (♂ genitalia) (preocc. Garrett, 1925). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- SIMPLEX Edwards, 1932: 143, fig. 2 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Rio de Janeiro. Ref. — Lane, 1952c: 408. Syntypes ♂, BMNH.
- SIMPLICIPES Freeman, 1951: 87, fig. 167 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Osorno, Llanquihue). Type ♂, BMNH.
- SIMPLICISTILA Freeman, 1951: 89, fig. 163 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangue. Type ♂, BMNH.
- SINUATA Freeman, 1951: 83, fig. 157 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Neuquén, Tierra del Fuego), Chile (Osorno, Llanquihue, Chiloé). Type ♂, BMNH.
- SPINATA Edwards, 1932: 144, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo; Rio de Janeiro, Petrópolis. Syntypes ♂, BMNH.
- TIEIALIS Freeman, 1951: 86, fig. 165 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Chubut, Tierra del Fuego), Chile (Bío Bío). Type ♂, BMNH.
- VOGELI Edwards, 1932: 144, fig. 4 (♂ genitalia). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Type ♂, BMNH.

Genus THORACOTROPIS Freeman

- THORACOTROPIS Freeman, 1951: 81. Type-species, *cypriformis* Freeman (orig. des.).
- CYPRIFORMIS Freeman, 1951: 81, figs. 156 (♂ genitalia), 292 (wing). Type-locality: Chile, Chiloé, Ancud. Type ♂, BMNH.

Tribe MYCOMYINI

Ref. — Coher, 1959 (rev.).



Genus ECHINOPODIUM Freeman

ECHINOPODIUM Freeman, 1951: 45. Type-species, *nigricoxa* Freeman (orig. des.).

Ref. — Duret, 1976 (key to spp.).

ARAUCANUM Duret, 1974: 102, figs. 13-15 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Chiloé, Llanquihue, Aisén, Malleco, Osorno, Valdivia, Arauco, Cautín). Type ♂, Coll. Duret.

ARGENTINENSE Duret, 1975b: 77, figs. 1-3 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.

ARMATUM Duret, 1976: 180, figs. 4-6 (midcoxal process, ♂ genitalia). Type-locality: Chile, Cautín, Los Coigües, Lago Villarica. Type ♂, CNC.

AYSENSE Duret, 1976: 181, fig. 7 (midcoxal process). Type-locality: Chile, Aisén, Puerto Cisnes (72°40'W, 44°45'S). Distr. — Chile (Aisén, Osorno). Type ♂, CNC.

BEJARANOI Duret, 1974: 103, figs. 16-20 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Ñuble). Type ♂, Coll. Duret.

BIFURCATUM Duret, 1974: 105, figs. 21-24 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén, Osorno). Type ♂, Coll. Duret.

CEKALOVICI Duret, 1976: 181, figs. 8-10 (midcoxal process, ♂ genitalia). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcáhue. Distr. — Chile (Cautín, Chiloé). Type ♂, CNC.

CHILENSE Duret, 1976: 183, figs. 11-13 (midcoxal process, ♂ genitalia). Type-locality: Chile, Cautín, Los Coigües, Lago Villarica. Type ♂, CNC.

CHILOENSE Coher, 1959: 129, figs. 126-127 (♂ genitalia). Type-locality: Chile, Chiloé, Butalcura. Distr. — Argentina (Río Negro), Chile (Chiloé, Magallanes). Type ♂, USNM.

COHERI Duret, 1974: 105, figs. 25-27 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén, Malleco, Llanquihue, Chiloé). Type ♂, Coll. Duret.

CORONATUM Duret, 1976: 184, figs. 14-16 (midcoxal process, ♂ genitalia). Type-locality: Chile, Cautín, Los Coigües, Lago Villarica. Type ♂, CNC.

CURVISPINA Freeman, 1951: 47, figs. 94-95 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro), Chile (Aisén, Llanquihue, Chiloé). Type ♂, BMNH.

DIGITALIS Freeman, 1951: 47, figs. 87 (midcoxal process), 91-93 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Neuquén, Río Negro), Chile (Llanquihue, Chiloé). Ref. — Duret, 1974: 98, figs. 1-3 (midcoxal process, ♂ genitalia). Type ♂, BMNH.



- ELEGANS** Duret, 1976: 184, figs. 17-19 (midcoxal process, ♂ genitalia). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcahue. Distr. — Chile (Ñuble, Chiloé). Type ♂, CNC.
- FLAGELLATUM** Freeman, 1951: 48, figs. 86 (midcoxal process), 96-98 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro, Neuquén), Chile (Aisén, Malleco, Llanquihue, Chiloé). Ref. — Duret, 1974: 98, figs. 4-9 (palpus, midcoxal process, tibial spur, ♂ genitalia). Type ♂, BMNH.
- FREEMANI** Duret, 1974: 106, figs. 28-31 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén, Malleco, Cautín). Type ♂, Coll. Duret.
- GENTILH** Duret, 1975b: 77, figs. 4-6 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Chiloé). Type ♂, Coll. Duret.
- INSOLITUM** Duret, 1976: 185, figs. 20-22 (midcoxal process, ♂ genitalia). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcahue. Distr. — Chile (Cautín, Chiloé). Type ♂, CNC.
- LANEI** Duret, 1974: 107, figs. 32-34 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén). Type ♂, Coll. Duret.
- MINOR** Duret, 1975b: 79, figs. 7-10 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén). Type ♂, Coll. Duret.
- MUNROEI** Duret, 1976: 185, figs. 23-25 (midcoxal process, ♂ genitalia). Type-locality: Chile, Aisén, Puerto Cisnes (72°40'W, 44°45'S). Type ♂, CNC.
- NEUQUENSE** Duret, 1975b: 81, figs. 11-14 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.
- NIGRESCENS** Duret, 1974: 109, figs. 35-38 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Chiloé, Magallanes). Type ♂, Coll. Duret.
- NIGRICOXA** Freeman, 1951: 46, figs. 82 (mouthparts), 83 (leg), 84 (midcoxal process), 88-90 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Chubut, Neuquén, Río Negro, Tierra del Fuego), Chile (Malleco, Ñuble, Curicó, Cautín, Aisén, Llanquihue, Chiloé, Magallanes). Ref. — Duret, 1974: 101, fig. 10 (midcoxal process). Type ♂, BMNH.
- PATAGONICUM** Duret, 1976: 187, figs. 26-29 (midcoxal process, ♂ genitalia). Type-locality: Chile, Cautín, Los Coigües, Lago Villarrica. Distr. — Chile (Cautín, Chiloé). Type ♂, CNC.
- PENAI** Duret, 1976: 189, figs. 30-32 (midcoxal process, ♂ genitalia). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcahue. Type ♂, CNC.
- FILOSUM** Duret, 1976: 189, fig. 33 (midcoxal process). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcahue. Type ♂, Universidad de Chile, Santiago.
- PUCARAENSE** Duret, 1974: 109, figs. 39-41 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina



- (Neuquén), Chile (Aisén, Malleco, Osorno, Chiloé). Type ♂, Coll. Duret.
- RIONEGRENSE** Coher, 1959: 130, figs. 130-131 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Aisén, Linares, Malleco, Curicó, Ñuble, Llanquihue, Chiloé, Magallanes). Ref. — Duret, 1974: 101, figs. 11-12 (midcoxal process, ♂ genitalia). Type ♂, USNM.
- SERGIOTI** Duret, 1974: 110, figs. 42-45 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén, Chiloé). Type ♂, Coll. Duret.
- SPINOSUM** Duret, 1974: 111, figs. 46-48 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Aisén, Chiloé). Type ♂, Coll. Duret.
- STONEI** Duret, 1974: 113, figs. 49-52 (midcoxal process, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Distr. — Argentina (Neuquén), Chile (Malleco, Osorno, Ñuble, Llanquihue, Chiloé). Type ♂, Coll. Duret.
- STRIATUM** Duret, 1974: 115, figs. 53-57 (midcoxal process, tibial spur, ♂ genitalia). Type-locality: Argentina, Neuquén, Pucará. Type ♂, Coll. Duret.
- STUARDOI** Duret, 1976: 190, figs. 34-36 (midcoxal process, ♂ genitalia). Type-locality: Chile, Aisén, Puerto Cisnes (72°40'W, 44°45'S). Type ♂, CNC.
- VILLARRICAENSE** Duret, 1976: 191, figs. 37-39 (midcoxal process, ♂ genitalia). Type-locality: Chile, Cautín, Los Coigües, Lago Villarrica. Type ♂, CNC.
- VOCKEROTHI** Duret, 1976: 192, figs. 40-43 (midcoxal process, ♂ genitalia). Type-locality: Chile, Chiloé, Isla Chiloé, Dalcahue. Distr. — Chile (Aisén, Chiloé). Type ♂, CNC.

Genus MYCOMYA Rondani


- MYCOMYA** Rondani, 1856: 194. Type-species, *Sciophila marginata* Meigen (mon.).
- Mycomya*, emend.
- Cnephaeophila* Philippi, 1865: 618. Type-species, *fenestralis* Philippi (mon.).
- Sciophila*, authors, not Meigen.
- ALEXANDERI** Coher, 1950b: 565, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- AMERICANA** (Schiner), 1868: 14 (*Sciophila*). Type-locality: "South America". Type ?, WIEN.
- ANSATA** Freeman, 1951: 38, figs. 33-35 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.



- ARETHUSA Coher, 1952: 216, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- AUSTROBLIQUA Coher, 1959: 73, figs. 72-73 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo). Type ♂, MZSP.
- BASINERVA Freeman, 1951: 37, figs. 29-32 (♂ genitalia), 80 (midcoxal process). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- BEQUAERTI Coher, 1950b: 577, fig. 13 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- BIFIDA Freeman, 1951: 34, figs. 18 (♂ genitalia), 78 (midcoxal process). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- BORACENSIS Coher, 1959: 80, figs. 84-85 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, Coll. Coher.
- BORINQUENSIS Coher, 1959: 88, figs. 97-99 (♂ genitalia). Type-locality: Puerto Rico, Maricao. Distr. — Puerto Rico, Brazil (Mato Grosso, Goiás, São Paulo). Type ♂, USNM.
- BREVIFURCATA Enderlein, 1910a: 162. Type-locality: Brazil, Santa Catarina. Syntypes ♀, Stettin.
- BRONTES Coher, 1952: 217, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Cajuru. Distr. — Brazil (Goiás, São Paulo). Type ♂, MZSP.
- CAMPESTRA Coher, 1959: 89, figs. 100-101 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- CARRERA Coher, 1950b: 572, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- CHILENSIS (Blanchard), 1852: 347 (*Sciophila*). Type-locality: Chilc, Coquimbo. Distr. — Argentina (Río Negro), Chile (Coquimbo, Valparaiso, Nuble, Bío Bío, Malleco, Osorno, Llanquihue, Chiloé). Ref. — Freeman, 1951: 32, figs. 10, 12 (♂ genitalia), 75 (midcoxal process).
- CITRINA Shaw, 1940: 806, fig. 4 (♂ genitalia). Type-locality: Costa Rica, Rivas, Valley of the General, 2925 ft. Ref. — Coher, 1959: 76, figs. 76-77 (♂ genitalia).
- CLAVATA (Lynch Arribáizaga), 1892: 417 (1892: 43) (*Sciophila*). Type-locality: Argentina, Buenos Aires, Chacabuco.
- CLETA Coher, 1952: 217, fig. 3 (♂ genitalia). Type-locality: Argentina, Salta, Urundel. Type ♂, MZSP.
- CONNECTA Coher, 1959: 86, figs. 95-96 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, Coll. Coher.
- COXALIS Freeman, 1951: 43, figs. 57-58. Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.



- CRAMPTONI Coher, 1950b: 569, fig. 5 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- CYLINDRICA Freeman, 1951: 41, figs. 50-52 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Distr. — Argentina (Río Negro), Chile (Ñuble, Osorno, Llanquihue). Type ♂, BMNH.
- DIVISA Freeman, 1951: 45, figs. 68-70 (♂ genitalia). Type-locality: Chile, Llanquihue, Puerto Varas. Distr. — Argentina (Río Negro), Chile (Ñuble, Bío Bío, Osorno, Llanquihue, Chiloé). Type ♂, BMNH.
- DRYOPE Coher, 1952: 219, fig. 4 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- DUMETA Coher, 1959: 85, figs. 92-94 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- EDWARDSII Coher, 1950b: 575, fig. 11 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- EPACRA Coher, 1959: 77, figs. 78-79 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo). Type ♂, MZSP.
- EXCERPTA Coher, 1959: 79, figs. 82-83 (♂ genitalia). Type-locality: Brazil, São Paulo. Type ♂, MZSP.
- FALCIFERA Freeman, 1951: 41, figs. 47-49 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- FENESTRALIS (Philippi), 1865: 618 (*Cnephacophila*). Type-locality: Chile, Santiago. Ref. — Freeman, 1951: 3 (in *Mycomya*).
- FERRAZI Coher, 1950b: 579, fig. 15 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- FLAVESCENS Freeman, 1951: 38, figs. 40-41 (♂ genitalia), 75 (midcoxal process). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- FORCIPATA Freeman, 1951: 42, figs. 53-54 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Distr. — Argentina (Río Negro), Chile (Ñuble). Type ♂, BMNH.
- FREEMANI Coher, 1950b: 570, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- FUNEBRIS Freeman, 1951: 38, figs. 36-37 (♂ genitalia). Type-locality: Chile, Chiloé, Ancud. Type ♂, BMNH.
- FUSCICORNIS Freeman, 1951: 36, figs. 24-25 (♂ genitalia), 81 (midcoxal process). Type-locality: Chile, Chiloé, Puntra. Distr. — Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- HAMADRYAS Coher, 1952: 220, fig. 5 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Santo Amaro). Distr. — Brazil (Goiás, São Paulo), Argentina (Salta). Type ♂, MZSP.



- HANSONI Coher, 1950b: 566, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- INERMIS Freeman, 1951: 40, figs. 44-46 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- INFIRMA (Lynch Arribáizaga), 1892: 420 (1892: 46) (*Sciophila*). Type-locality: Argentina, Prov. of Buenos Aires.
- INFUSCATA Freeman, 1951: 39, figs. 42-43 (♂ genitalia). Type-locality: Chile, Angol. Type ♂, BMNH.
- IPHIS Coher, 1952: 220, fig. 6 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
- IRENE Coher, 1952: 222, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Distr. — Brazil (São Paulo), Argentina (Salta). Type ♂, MZSP.
- JAFFUELENSIS Freeman, 1951: 43, figs. 59-60 (♂ genitalia). Type-locality: Chile, Valparaiso, Marga Marga. Type ♂, BMNH.
- LAMELLATA Freeman, 1951: 35, figs. 19 (♂ genitalia), 76 (midcoxal process). Type-locality: Argentina, Río Negro, Puerto Blest. Distr. — Argentina (Río Negro), Chile (Chiloé). Type ♂, BMNH.
- LANEI Coher, 1950b: 572, fig. 9 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- LONGISTILA Freeman, 1951: 42, figs. 55-56 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- LUTEA Enderlein, 1910a: 161. Type-locality: Brazil, Santa Catarina. Syntypes ♀, Stettin.
var. DORSIMACULA Enderlein, 1910a: 162. Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.
- MANTERI Coher, 1950b: 564, fig. 1 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- MATHESONI Coher, 1950b: 578, fig. 14 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- MERIDIONALIS Johannsen, 1909: 180 (nom. nov. for *diluta* Williston). Type-locality: St. Vincent. W. I.
diluta Williston, 1896: 263, pl. 8, fig. 17 (wing) (*Sciophila*; proocc. Zetterstedt, 1860). Type-locality: St. Vincent, W. I. Syntypes ♂, BMNH.
- MIDAS Coher, 1952: 223, fig. 8 (♂ genitalia). Type-locality: Brazil, São Paulo, Cajuru. Type ♂, MZSP.
- MULTISETA Coher, 1959: 83, figs. 90-91 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (São Paulo). Type ♂, Coll. Coher.



- NICOTHOE Coher, 1952: 224, fig. 9 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Morumbi). Type ♂, MZSP.
- OCHRACEA Freeman, 1951: 34, figs. 15, 17 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Distr. — Argentina (Río Negro), Chile (Valparaiso, Ñuble, Bío Bío, Llanquihue, Chiloé). Type ♂, BMNH.
- PAUPERCULA Coher, 1959: 99, figs. 104-105 (♂ genitalia). Type-locality: Chile, Valparaiso, Marga Marga. Type ♂, USNM.
- PECTINATA Freeman, 1951: 43, figs. 61-63 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego). Type ♂, BMNH.
- PERUVIANA Edwards, 1931: 258. Type-locality: Peru, Lima, Verrugas. Type ♀, BMNH.
- PONTIS Coher, 1959: 75, figs. 74-75 (♂ genitalia). Type-locality: Brazil, São Paulo, Cajuru. Distr. — Brazil (São Paulo). Type ♂, MZSP.
- PORTOBLEST Coher, 1959: 98, figs. 102-103 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Type ♂, USNM.
- SAMESTERI Coher, 1950b: 576, fig. 12 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SETIFERA Freeman, 1951: 45, figs. 71-74 (♂ genitalia). Type-locality: Chile, Chiloé, Castro. Distr. — Chile (Osorno, Chiloé). Type ♂, BMNH.
- SHANNONI Coher, 1950b: 574, fig. 10 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SHAWI Coher, 1950b: 568, fig. 4 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SIMILLIMA Freeman, 1951: 33, fig. 11 (9th abdominal tergite). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- SIMPLA Coher, 1959: 82, figs. 88-89 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SPINIFERA Freeman, 1951: 36, figs. 22-23 (♂ genitalia), 79 (midcoxal process). Type-locality: Chile, Chiloé, Ancud. Type ♂, BMNH.
- SPINOSA Freeman, 1951: 36, figs. 26-28 (♂ genitalia), 81 (midcoxal process). Type-locality: Chile, Chiloé, Punta. Type ♂, BMNH.
- SUBEPACRA Coher, 1959: 78, figs. 80-81 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- SUBFUSCA Freeman, 1951: 33, figs. 14, 16 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.
- TANTALOS Coher, 1959: 71, figs. 70-71 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.



- TANTILLA (Loew), 1869: 140 (Cent. 9, n° 21) (*Sciophila*). Type-locality: U.S.A., Nebraska. Distr. — S. Canada, south to New Mexico and North Carolina; Costa Rica.
- TAURUS Freeman, 1951: 44, figs. 64-67 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Bío Bío, Llanquihue). Type ♂, BMNH.
- THEOBALDI Coher, 1959: 81, figs. 86-87 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, Coll. Coher.
- TRAVERI Coher, 1950b: 571, fig. 7 (♂ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type ♂, MZSP.
traveriae Coher, 1959: 70, emend.
- TRICHOPS Freeman, 1951: 35, figs. 20-21 (♂ genitalia), 77 (midcoxal process). Type-locality: Chile, Chiloé, Ancud. Type ♂, BMNH.
- VALDIVIANA (Philippi), 1865: 624 (*Sciophila*). Type-locality: Chile, Corral. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Ref. — Freeman, 1951: 39, figs. 38-39 (♂ genitalia), 81 (midcoxal process).

Genus NEOEMPHERIA Osten Sacken

- Empheria* Winnertz, 1863: 783 (preocc. Hagen, 1856). Type-species, *Sciophila striata* Meigen (Coquillett, 1910: 537).
- NEOEMPHERIA Osten Sacken, 1878: 9 (nom. nov. for *Empheria* Winnertz). Type-species, *Sciophila striata* Meigen (aut.).
- Pleonazoneura* Enderlein, 1910a: 156. Type-species, *johannseni* Enderlein (orig. des.).
- Neurocompsa* Enderlein, 1910a: 158. Type-species, *ornatipennis* Enderlein (orig. des.).
- ACIDOTI Coher, 1959: 27, pl. 2 (wing), figs. 16-17 (♂ genitalia). Type-locality: Peru, Loreto, Cerro Azul. Type ♂, Coll. Coher.
- APICALIS Kertész, 1909: 140. Type-locality: Peru, Callanga. Type ♀, Budapest.
- BIDENTATA Coher, 1959: 20, pl. 1 (wing), figs. 2-3 (♂ genitalia). Type-locality: Argentina, Salta, Embarcación. Type ♂, IML.
- BIFIDA Coher, 1959: 31, pl. 2 (wing), figs. 20-21 (♂ genitalia). Type-locality: Argentina, Salta, Aguaray. Type ♂, IML.
- BIFLAGELLATA Edwards, 1940b: 127, fig. 12B (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, BMNH.
- BILOBATA Edwards, 1940b: 113, fig. 3B (♂ genitalia), pl. 1, fig. 11 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- BIPECTINATA Edwards, 1940b: 110, fig. 1B (♂ genitalia), pl. 1, fig. 2 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Goiás, Santa Catarina). Type ♂, BMNH.



- BORGMEIERI Edwards, 1940b: 113. Type-locality: Brazil, Santa Catarina, Bom Retiro. Type ?♀, BMNH.
- BRADLEYI Edwards, 1940b: 118, fig. 6E (♂ genitalia) (as *maculipennis* ssp.). Type-locality: Peru, Putumayo, La Sombra. Type ♂, Cornell University.
- BRASILIENSIS Coher, 1959: 56, pl. 4 (wing), figs. 60-61 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- BREVICAUDA Edwards, 1940b: 112, pl. 1, fig. 8 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♀, BMNH.
- CINCTA Shaw, 1940: 806, fig. 4 (♂ genitalia). Type-locality: Costa Rica, San Ramón, Valley of the General, 2800 ft. Ref. — Coher, 1959: 39, pl. 3 (wing).
- COMES Coher, 1959: 44, pl. 4 (wing), figs. 41-42 (♂ genitalia). Type-locality: Costa Rica, San Mateo, Higuito. Type ♂, MZSP.
- COSTALIMAI Edwards, 1940b: 126, fig. 12A (♂ genitalia), pl. 2, fig. 23 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- COSTARICENSIS Coher, 1959: 49, pl. 4 (wing), figs. 50-51 (♂ genitalia). Type-locality: Costa Rica, San Mateo, Higuito. Type ♂, MZSP.
- DEFLETA Coher, 1959: 53, pl. 4 (wing), figs. 56-57 (♂ genitalia). Type-locality: Brazil, Goiás, Corumbá. Type ♂, MZSP.
- DZIEDZICKII Coher, 1959: 44, pl. 3 (wing), figs. 39-40 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- ECUADORENSIS Coher, 1959: 21, pl. 1 (wing), figs. 4-5 (♂ genitalia). Type-locality: Ecuador, Napo-Pastaza, Abitagua. Type ♂, USNM.
- ENDERLEINI Edwards, 1940b: 122, fig. 9B (♂ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina), Argentina (Salta). Type ♂, Stettin.
- EVANESCENS (Enderlein), 1910a: 158 (*Pleonazoneura*; as *johannseni* var.). Type-locality: Brazil, Santa Catarina. Ref. — Edwards, 1940b: 121, fig. 7B (♂ genitalia) (as sp.). Type ♂, Stettin.
- FACETA Coher, 1959: 48, pl. 4 (wing), figs. 47-49 (♂ genitalia). Type-locality: Argentina, Salta, Aguaray. Type ♂, IML.
- FALLAX Coher, 1959: 33, pl. 2 (wing), figs. 24-25 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo. Distr. — Brazil (São Paulo), Argentina (Jujuy). Type ♂, Coll. Coher.
- FLAVICORNIS Edwards, 1940b: 123, fig. 10B (♂ genitalia), pl. 2, fig. 21 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- FLAVICOXA Edwards, 1940b: 115, fig. 4E (♂ genitalia), pl. 2, fig. 15 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.



- FORMOSENSIS (Lynch Arribáizaga), 1892: 416 (1892: 42) (*Sciophila*). Type-locality: Argentina, Chaco, near Formosa.
- GLOCHIS Coher, 1959: 22, pl. 1 (wing), figs. 6-7 (♂ genitalia). Type-locality: Panama, Canal Zone, Barro Colorado. Distr. — Costa Rica, Panama. Type ♂, USNM.
- GOIANA Coher, 1959: 38, pl. 3 (wing), figs. 31-32 (♂ genitalia). Type-locality: Brazil, Goiás, Corumbá. Type ♂, MZSP.
- HORRENS Coher, 1959: 25, pl. 1 (wing), fig. 8 (tergite 8), 9 (sternite 8), 10-11 (♂ genitalia). Type-locality: Argentina, Salta, Aguaray. Type ♂, IML.
- JAMAICENSIS Coher, 1961: 129, figs. 1-6. Type-locality: Jamaica.
- JOHANNSENI (Enderlein), 1910a: 156 (*Pleonazoneura*). Type-locality: Brazil, Santa Catarina. Ref. — Edwards, 1940: 126, fig. 11B (♂ genitalia). Lectotype ♂, Stettin.
- JUGALIS Coher, 1959: 47, pl. 4 (wing), figs. 45-46 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Morumbi). Type ♂, Coll. Coher.
- KAESTNERI Edwards, 1940b: 121, fig. 8A (♂ genitalia). Type-locality: Brazil, Santa Catarina. Distr. — Brazil (São Paulo, Santa Catarina), Argentina (Salta, Tucumán). Type ♂, Stettin.
- LANEI Edwards, 1940b: 118, fig. 6B (♂ genitalia), pl. 2, fig. 17 (wing). Type-locality: Brazil, São Paulo, Jiquiá. Distr. — Panama (Canal Zone), Brazil (São Paulo). Type ♂, BMNH.
- LARIFUGA Coher, 1959: 35, pl. 2 (wing), figs. 27-28 (♂ genitalia). Type-locality: Argentina, Salta, Aguaray. Distr. — Peru (Loreto), Argentina (Salta). Type ♂, IML.
- LEVIR Coher, 1959: 39, pl. 3 (wing), figs. 33-34 (♂ genitalia). Type-locality: Peru, Loreto, Cerro Azul. Type ♂, Coll. Coher.
- LINDNERI Edwards, 1940b: 125, fig. 11A (♂ genitalia), pl. 1, fig. 6 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- LONGISETA Coher, 1959: 37, pl. 3 (wing), figs. 29-30 (♂ genitalia). Type-locality: Argentina, Salta, Aguaray. Distr. — Brazil (São Paulo), Argentina (Salta). Type ♂, IML.
- LUEDERWALDTI Edwards, 1940b: 121, fig. 9A (♂ genitalia), pl. 2, fig. 20 (wing). Type-locality: Brazil, Santa Catarina. Distr. — Brazil (São Paulo, Santa Catarina). Type ♂, Stettin.
- LUTZI Edwards, 1940b: 128, pl. 1, fig. 4 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Ref. — Coher, 1959: 51, figs. 52-53 (♂ genitalia). Type ♀, BMNH.
- MACULIPENNIS Williston, 1896: 262, pl. 8, fig. 16 (wing). Type-locality: St. Vincent, W. I. Ref. — Edwards, 1940b: 118, fig. 6D (♂ genitalia), pl. 2, fig. 18 (wing). Syntypes ♂, BMNH.



- MUELLERI Edwards, 1940b: 116, fig. 6C (δ genitalia), pl. 2, fig. 16 (wing).
Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. —
Brazil (São Paulo, Santa Catarina). Type δ , BMNH.
- NEIVAI Edwards, 1940b: 109, fig. 1A (δ genitalia), pl. 1, fig. 1 (wing).
Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. —
Panama, Guyana, Brazil (São Paulo, Santa Catarina). Type δ ,
BMNH.
- ORNATIPENNIS (Enderlein), 1910: 158, fig. 1 (wing) (*Neurocompsa*). Type-
locality: Brazil, Santa Catarina. Distr. — Brazil (Bahia, São
Paulo, Santa Catarina), Argentina (Salta). Ref. — Edwards,
1940b: 119, fig. 7A (δ genitalia), pl. 2, fig. 19 (wing). Type δ ,
Stettin.
- PANAMENSIS Coher, 1959: 19, pl. 1 (wing), fig. 1 (δ genitalia). Type-
locality: Panama, Canal Zone, Barro Colorado. Type δ , USNM.
- PAULENSIS Coher, 1959: 55, pl. 4 (wing), figs. 58-59 (δ genitalia). Type-
locality: Brazil, São Paulo, Campos do Jordão. Type δ , MZSP.
- FEREIRAI Edwards, 1940b: 129, pl. 1, fig. 5 (wing) (1941: 307, again as
n. sp.). Type-locality: Brazil, Mato Grosso, Salobra. Type φ ,
BMNH.
- PHILIPSI Shaw, 1951: 148 (nom. nov. for *neivai* Shaw). Type-locality: Costa
Rica, Valley of the General, Pedregoso, 2300 ft.
neivai Shaw, 1940 [Dec.]: 808, fig. 7 (δ genitalia) (preocc. Edwards,
1940 [April]). Type-locality: Costa Rica, Valley of the General,
Pedregoso, 2300 ft.
- PILOSA Edwards, 1940b: 124, fig. 10D (δ genitalia). Type-locality: Brazil,
Santa Catarina, Nova Teutônia. Type δ , BMNH.
- PLAUMANNI Edwards, 1940b: 114, fig. 4A (δ genitalia), pl. 2, fig. 13 (wing).
Distr. — Brazil (São Paulo, Santa Catarina). Type δ , BMNH.
var. ? Edwards, 1940b: 115, pl. 2, fig. 14 (wing). φ from Nova
Teutônia.
- PORTORICENSIS Coher, 1959: 41, pl. 3 (wing), figs. 37-38 (δ genitalia). Type-
locality: Puerto Rico, Mayaguez. Type δ , USNM.
- PUNCTICOXA Edwards, 1940b: 115, fig. 5 (δ genitalia). Type-locality: Brazil,
Santa Catarina, Nova Teutônia. Distr. — Brazil (Mato Grosso,
São Paulo, Santa Catarina), Argentina (Salta, Tucumán). Ref. —
Coher, 1959: 24, pl. 1 (wing). Type δ , BMNH.
- RABELLOI Coher, 1959: 52, pl. 4 (wing), figs. 54-55 (δ genitalia). Type-
locality: Brazil, São Paulo, Salesópolis, Estação Biológica de
Boracéia. Type δ , MZSP.
- ROSTRATA Edwards, 1940b: 124, fig. 10C (δ genitalia). Type-locality: Brazil,
Santa Catarina, Nova Teutônia. Type δ , BMNH.
- SEPARATA Coher, 1959: 32, pl. 2 (wing), figs. 22-23 (δ genitalia). Type-
locality: Costa Rica, San Mateo, Higuito. Type δ , MZSP.
- SHANNONI Edwards, 1940b: 125, fig. 11C (δ genitalia), pl. 2, fig. 22 (wing).
Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ ,
BMNH.

- SIMPLEX Edwards, 1940b: 112, fig. 3A (δ genitalia), pl. 1, fig. 10 (δ wing), pl. 1, fig. 9 (φ wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Ref. — Coher, 1959: 16. Type δ , BMNH.
- SMARTI Edwards, 1940b: 123, fig. 9C (δ genitalia). Type-locality: Guyana, Mazaruni. Type δ , BMNH.
- SOCIA Coher, 1959: 45, pl. 4 (wing), figs. 43-44 (δ genitalia). Type-locality: Brazil, Goiás, Corumbá. Type δ , MZSP.
- SPINOSA Edwards, 1940b: 111, fig. 2 (δ genitalia), pl. 1, fig. 7 (wing). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- SUBCLAVATA Edwards, 1940b: 121, fig. 8B (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- SUBFALLAX Coher, 1959: 34, pl. 2 (wing), fig. 26 (δ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type δ , MZSP.
- SUBHORRENS Coher, 1959: 26, pl. 1 (wing), figs. 12 (tergite 8), 13 (sternite 8), 14 (δ genitalia). Type-locality: Costa Rica, San Matco, Higuito. Type δ , MZSP.
- SUBLEVIR Coher, 1959: 40, pl. 3 (wing), figs. 35-36 (δ genitalia). Type-locality: Panama, Canal Zone, Summit. Type δ , USNM.
- TETRAPHAEA Shaw, 1940: 807, fig. 6 (δ genitalia). Type-locality: Costa Rica, Valley of the General, Rivas, 2875 ft. Ref. — Coher, 1959: 21, pl. 1 (wing).
- UNISPINOSA Edwards, 1940b: 123, fig. 10A (δ genitalia). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type δ , BMNH.
- VARIPENNIS (Lynch Arribáizaga), 1892: 423 (1892: 49) (*Empheria*). Type-locality: Argentina, Misiones, Santa Ana. Refs. — Edwards, 1940b: 116, fig. 6A (δ genitalia); Coher, 1959: 18, pl. 1 (wing). *variipennis*, lapsus.
- VOGELI Edwards, 1940b: 127, fig. 12C (δ genitalia), pl. 1, fig. 3 (wing). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Type δ , BMNH.
- ZETEKI Coher, 1959: 28, pl. 2 (wing), figs. 18-19 (δ genitalia). Type-locality: Panama, Canal Zone, Barro Colorado. Type δ , USNM.

Tribe SCIOPHILINI

Genus ACNEMIA Winnertz

- ACNEMIA Winnertz, 1863: 798. Type-species, *Leia nitidicollis* Meigen (Johannsen, 1909: 63).
- Agaricobia* Philippi, 1865: 626. Type-species, *fulvicollis* Philippi (mon.).
- FLAVICOXA Freeman, 1951: 60, fig. 116 (δ genitalia). Type-locality: "Chile". Type δ , BMNH.
- FREEMANI Lane, 1956a: 121, fig. 4 (δ genitalia). Type-locality: Brazil, São Paulo, Campos do Jordão. Type δ , MZSP.



- FULVICOLLIS (Philippi), 1865: 626, pl. 24, fig. 11 (whole insect) (*Agaricobia*). Type-locality: Chile, Santiago, near Santiago. Distr. — Argentina (Neuquén), Chile (Valparaiso, Santiago, Llanquihue). Ref. — Freeman, 1951: 58, fig. 114 (♂ genitalia).
- FUNEREA Freeman, 1951: 59, fig. 115 (♂ genitalia). Type-locality: Argentina, Río Negro, Bariloche. Type ♂, BMNH.
- VITTATA Freeman, 1951: 59, fig. 274 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Type ♀, BMNH.

Genus ALLOCOTOCERA Mik

- Eurycera* Dziedzicki, 1885: 169 (preocc. Laporte, 1833). Type-species, *flava* Dziedzicki (mon.) = *pulchella* (Curtis).
- ALLOCOTOCERA Mik, 1886: 102 (nom. nov. for *Eurycera* Dziedzicki). Type-species, *Eurycera flava* Dziedzicki (aut.) = *pulchella* (Curtis).
- COXIPONENSIS Lane, 1952a: 134, fig. 2 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.
- FLAVICOXA Freeman, 1951: 51, figs. 102 (♂ genitalia), 275 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Argentina (Río Negro), Chile (Osorno, Llanquihue, Chiloé). Type ♂, BMNH.
- GLANDULARIS Edwards, 1940a: 453. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♂ ♀, BMNH.
- NIGRICOXA Freeman, 1951: 52, figs. 103-104 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Nahuel Huapi. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Llanquihue). Type ♂, BMNH.

Genus ANEURA Marshall

- ANEURA Marshall, 1896: 256, 287. Type-species, *boletinoides* Marshall (mon.).
- BASALIS Freeman, 1951: 53, fig. 106 (♂ genitalia). Type-locality: Argentina, Río Negro, Puerto Blest. Type ♂, BMNH.
- LCNGISTILA Freeman, 1951: 52, figs. 105 (♂ genitalia), 277 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.

Genus APHELOMERA Skuse

- APHELOMERA Skuse, 1888: 1206. Type-species, *sydneiensis* Skuse (mon.).
- ARMATA Freeman, 1951: 61, fig. 120 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Gutiérrez. Type ♂, BMNH.
- CRISTATA Freeman, 1951: 60, figs. 117-119 (♂ genitalia), 278 (wing). Type-locality: Argentina, Río Negro, Lago Correntoso. Distr. — Argentina (Río Negro, Tierra del Fuego), Chile (Ñuble, Llanquihue). Type ♂, BMNH.



INERMIS Freeman, 1951: 61. Type-locality: Argentina, Rio Negro, Lago Correntoso. Type ♀, BMNH.

SUBCOMPLETA Edwards, 1940a: 462. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Genus CLUZOBRA Edwards

CLUZOBRA Edwards, 1940a: 463. Type-species, *Acnemia binocellaris* Edwards (orig. des.).

AITKENI Lane, 1956a: 123, fig. 6 (♂ genitalia). Type-locality: Trinidad, Arena Forest. Type ♂, MZSP.

BINOCELLARIS (Edwards), 1934: 362 (*Acnemia*). Type-locality: Paraguay, La Cordillera, San Bernardino. Distr. — Paraguay, Argentina (Chaco). Syntypes ♂ ♀, WIEN, BMNH.

BUTLERI Lane, 1959b: 118, fig. 11 (♂ genitalia). Type-locality: Brazil, Amapá, Serra do Navio. Type ♂, MZSP.

EDWARDSI Lane, 1960a: 381, fig. 5 (♂ genitalia). Type-locality: Trinidad, Navy Base. Distr. — Panama, Trinidad. Type ♂, MZSP.

FASCIPENNIS Edwards, 1940a: 464. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Goiás, Santa Catarina), Argentina (Misiones). Ref. — Lane, 1948b: 254, figs. 14-15 (♂ genitalia). Type ♀, BMNH.

FUSCIPENNIS Edwards, 1940a: 464. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

LANEI Edwards, 1941: 308. Type-locality: Brazil, Mato Grosso, Salobra. Type ♂, BMNH.

PLAUMANNI Edwards, 1940a: 463. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (São Paulo, Santa Catarina). Syntypes ♂ ♀, BMNH.

Genus EUDICRANA Loew


EUDICRANA Loew, 1869: 142 (Cent. 9, n° 23). Type-species, *obumbrata* Loew (mon.).

BASINERVA Freeman, 1951: 49, figs. 99 (♂ genitalia), 302 (wing). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Argentina (Rio Negro), Chile (Llanquihue). Type ♂, BMNH.

CLARIPENNIS Edwards, 1931: 258. Type-locality: Peru, Lima, Verrugas. Type ♀, BMNH.

ELEGANS Lane, 1948a: 251, figs. 8-10 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.

PALLIDA Freeman, 1951: 51, fig. 101 (♂ genitalia). Type-locality: "Chile". Type ♂, BMNH.



- SIMILIS Freeman, 1951: 50, fig. 100 (♂ genitalia). Type-locality: Chile, Valparaíso, Marga Marga. Type ♂, BMNH.
- SPLENDENS Lane, 1948a: 253, figs. 11-13 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Distr. — Brazil (Rio de Janeiro, São Paulo). Type ♂, MZSP.
- VITATA Edwards, 1931: 257. Type-locality: Peru, Junín, Colonia Perené. Distr. — Trinidad, Peru. Ref. — Lane, 1960a: 383, fig. 8 (♂ genitalia). Type ♀, BMNH.

Genus LEPTOMORPHUS Curtis

- LEPTOMORPHUS Curtis, 1831: pl. 365. Type-species, *walkeri* Curtis (orig. des.).
- Diomonus* Walker, 1848: 87. Type-species, *nebulescens* Walker (mon.).
- FASCICULATUS Edwards, 1933: 305. Type-locality: Bolivia, Mapiri, Sarampiuni, 700 m. Distr. — Peru, Bolivia, Brazil (Mato Grosso, São Paulo). Type ♂, Dresden.
- FEMORATUS Edwards, 1933: 306. Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Peru, Brazil (Rio de Janeiro). Type ♂, BMNH.
- NEVAI Edwards, 1940a: 452. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina (Misiones). Syntypes ♂ ♀, BMNH.

Genus MEGALOPELMA Enderlein

- MEGALOPELMA Enderlein, 1910a: 165. Type-species, *planiceps* Enderlein (orig. des.).
- CELLULARIS Edwards, 1940a: 458. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♀, BMNH.
- FRAUDULENTA (Williston), 1896: 263, pl. 8, fig. 18 (wing) (*Phthinia*). Type-locality: St. Vincent, W. I. Syntypes ♂, BMNH.
- PLANICEPS Enderlein, 1910a: 166, fig. 2 (wing). Type-locality: Brazil, Santa Catarina. Type ♀, Stettin.
- PLATYURA Edwards, 1940a: 459. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina (Tucumán). Syntypes ♂ ♀, BMNH.

Genus MONOCLONA Mik

- Staegeria* Wulp, 1876: xlix (preocc. Rondani, 1856). Type-species, *Sciophila halterata* Staeger (mon.) = *rufilatera* (Walker).
- MONOCLONA Mik, 1886: 279 (nom. nov. for *Staegeria* Wulp). Type-species, *Sciophila halterata* Staeger (aut.) = *rufilatera* (Walker).
- ABNORMALIS Fisher, 1939: 229. Type-locality: Costa Rica, San José. Type ♂, ANSP.



- BICOLOR (Enderlein), 1910a: 180 (*Acnemia*). Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.
- vittidorsum* Enderlein, 1910a: 182 (*Acnemia*). Type-locality: Brazil, Santa Catarina. Type ♂, Stettin.
- DIGITATA Edwards, 1940a: 460. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Trinidad, Peru, Brazil (Santa Catarina), Paraguay (La Cordillera). Syntypes ♂ ♀, BMNH.
- MACULATA Edwards, 1933: 306. Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil (Rio de Janeiro), Paraguay (La Cordillera). Type ♂, BMNH.
- NIGRIVENTRIS Edwards, 1941: 461. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- TAPICAREI Lane, 1952a: 136, fig. 4 (♂ genitalia). Type-locality: Brazil, São Paulo, São Paulo (Cantareira). Type ♂, MZSP.
- TRIFASCIATA Edwards, 1940a: 461. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.

Genus NEURATELIA Rondani

- NEURATELIA Rondani, 1856: 195. Type-species, *Mycetophila nemoralis* Meigen (orig. des.).
- SAPAICI Lane, 1952a: 135, fig. 3 (♂ genitalia). Type-locality: Brazil, São Paulo, Salesópolis, Estação Biológica de Boracéia. Type ♂, MZSP.

Genus NOVAKIA Strobl

- NOVAKIA Strobl, 1893: 162. Type-species, *scatopsiformis* Strobl (mon.).
- DISTINCTA Fisher, 1939: 234. Type-locality: Costa Rica, San José. Type ♂, ANSP.

Genus PARATRIZYGIA Tonnoir

- PARATRIZYGIA Tonnoir, 1929: 605. Type-species, *conformis* Tonnoir (orig. des.).
- INFUSCATA Freeman, 1951: 63, figs. 124-125 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- SETIFERA Freeman, 1951: 62, figs. 121-122 (♂ genitalia), 279 (wing). Type-locality: Argentina, Río Negro, Bariloche. Distr. — Argentina (Río Negro), Chile (Llanquihue, Chiloé). Type ♂, BMNH.
- SPINULOSA Freeman, 1951: 62, fig. 123 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Type ♂, BMNH.



Genus PARVICELLULA Marshall

- PARVICELLULA Marshall, 1896: 256, 284. Type-species, *triangula* Marshall (mon.).
- FLABELLIFERA Freeman, 1951: 55, fig. 110 (♂ genitalia). Type-locality: Argentina, Río Negro, Lago Correntoso. Type ♂, BMNH.
- PRODUCTA Freeman, 1951: 55, fig. 111 (♂ genitalia). Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Argentina (Río Negro), Chile (Llanquihue). Type ♂, BMNH.
- QUADRIPECTINATA Edwards, 1940a: 461. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- SHANNONI Lane, 1956a: 120, fig. 3 (♂ genitalia). Type-locality: Panama, Canal Zone, Barro Colorado I. Type ♂, USNM.

Genus PHTHINIA Winnertz

- PHTHINIA Winnertz, 1863: 779. Type-species, *humilis* Winnertz (Johannsen, 1909: 83).
- FASCIATA Freeman, 1951: 54, fig. 109 (♂ genitalia). Type-locality: Chile, Llanquihue, Peulla. Type ♂, BMNH.
- FLAGELLATA Freeman, 1951: 53, figs. 107 (♂ genitalia), 276 (wing). Type-locality: Argentina, Río Negro, Lago Nahuel Huapi, eastern end. Type ♂, BMNH.
- FURCATA Freeman, 1951: 54, fig. 108 (♂ genitalia). Type-locality: Chile, Chiloé, Ancud. Type ♂, BMNH.

Genus SCIOPHILA Meigen

- SCIOPHILA Meigen, 1818: 245. Type-species, *hirta* Meigen (Curtis, 1837: pl. 641).
- Lasiosoma* Winnertz, 1863: 748. Type-species, *Sciophila pilosa* Meigen (Coquillett, 1910: 558) = *hirta* Meigen.

Most of the species placed here are unrecognized, and may belong to many different genera.

- ABERRANS Philippi, 1865: 625. Type-locality: Chile, Santiago, Santiago, Mt. S. Cristóval.
- ANTARCTICA Walker, 1837: 334. Type-locality: Straits of Magellan, Port Famine. Ref. — Freeman, 1951: 56, figs. 112 (♂ genitalia), 284 (wing). Type ♂, BMNH.
- AUSTRALIS Philippi, 1865: 625. Type-locality: Chile, Valdivia Prov.
- CILIATA Edwards, 1940a: 456. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♂ ♀, BMNH.
- CINCTICORNIS Edwards, 1940a: 458. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina (Misiones). Syntypes ♂ ♀. BMNH.

- CINCTIFEMUR Shaw, 1940: 808, fig. 9 (♂ genitalia). Type-locality: Costa Rica, Valley of the General, Pedregoso, 2075 ft.
- FIDELIS Edwards, 1940a: 457. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- FRACTINERVIS Edwards, 1940a: 457. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♂ ♀, BMNH.
- OBSOLETA Blanchard, 1852: 347. Type-locality: Chile, Valparaiso.
- OCREATA Philippi, 1865: 625. Type-locality: Chile, Santiago, Santiago.
- PARANENSIS (Lynch Arribáizaga), 1892: 413 (1892: 39) (*Lasiosoma*). Type-locality: Argentina, Entre Rios: Paraná; Buenos Aires: Baradcro, Las Conchas and Navarro.
- POPOCATEPETLI Bellardi, 1859: 11. Type-locality: Mexico, Volcan Popocatepetl, 3800 m. Type ♀, Università di Torino.
- PRAECOX Philippi, 1865: 624. Type-locality: Chile, Santiago, Santiago.
- PUSILLA Philippi, 1865: 625. Type-locality: Chile, Valdivia Prov.
- THORACICA Philippi, 1865: 624. Type-locality: Chile, Corral.
- TRISTIS Blgot, 1888: 13, pl. 9, fig. 2 (whole insect). Type-locality: Cape Horn.
- VERNALIS Philippi, 1865: 624. Type-locality: Chile, Santiago, Santiago.

Genus STENOPHRAGMA Skuse

- Homaspis* Skuse, 1888: 1131, 1191 (preocc. Foerster, 1868). Type-species, *meridiana* Skuse (mon.).
- STENOPHRAGMA Skuse, 1890: 612 (nom. nov. for *Homaspis* Skuse). Type-species, *Homaspis meridiana* Skuse (aut.).
- FUSCA Edwards, 1934: 362. Type-locality: Peru, Cuzco, 4000-4200 m. Type ?♂, Dresden.
- HUMERALIS Edwards, 1934: 360. Type-locality: Paraguay, La Cordillera, San Bernardino. Type ♂, WIEN.
- INTERMEDIA Edwards, 1940a: 455. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Peru (Lima), Brazil (Santa Catarina). Type ♂, BMNH.
- LONGIFURCATA Freeman, 1951: 57, fig. 282 (wing). Type-locality: Argentina, Río Negro, Puerto Blest. Type ♀, BMNH.
- MORIGENEA Edwards, 1940a: 455. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Type ♂, BMNH.
- NIGRICAUDA Edwards, 1940a: 454. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♂ ♀, BMNH.
- OCHRACEA Freeman, 1951: 58, fig. 281 (wing). Type-locality: Argentina, Río Negro, Bariloche. Type ♀, BMNH.
- PLEURALIS Edwards, 1934: 361. Type-locality: Bolivia, La Paz. Distr. — Peru, Bolivia. Type ♂, Dresden.



Genus TRIZYGIA Skuse

- TRIZYGIA Skuse, 1888: 1137, 1204. Type-species, *flavipes* Skuse (mon.).
NITENS Edwards, 1940a: 462. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Syntypes ♂ ♀, BMNH.

UNPLACED MYCETOPHILIDAE

- CHILENSIS Philippi, 1865: 620, pl. 23, fig. 9 (head, lateral view) (*Gnoriste*).
Type-locality: Chile, Valdivia Prov.
OBSCURIVENTRIS Bigot, 1888: 13 (*Boletina*). Type-locality: Cape Horn. Type ♂, MNHN.

LITERATURE CITED

BELLARDI, L.

1859. *Saggio di ditteologia messicana 1*: 80 pp., 2 pls., Torino. [Also in *Mem. R. Accad. Sci. Torino* 19: 201-277, 2 pls., 1861].
1862. *Saggio di ditteologia messicana. Appendice*, 28 pp., 1 pl. Torino. [Also in *Mem. R. Accad. Sci. Torino* 21: 200-225, 1 pl., 1864].

BIGOT, J. M. F.

1888. Diptères, pp. 1-45, pls. 1-4, in Ministère de la Marine et de l'Instruction Publique, *Mission Scientifique du Cap Horn, 1882-1883*, 6 (Zool., 2). Paris, "1891".

BLANCHARD, E.

1852. Orden IX. Dipteros, pp. 327-468, in C. Gay, ed., *Historia fisica y política de Chile. Zoología* 7: 471 pp. Paris.

COHER, E. I.

1949. A note on synonymy in Mycetophilidae and *Neopeicypta*, n. n. (Dipt.). *Rev. Ent.* 20: 172.
1950a. Correccion: A new genus of fungus gnats (Diptera: Mycetophilidae). *Rev. Ent.* 21: 114.
1950b. Neotropical *Mycomyia*. I. (Diptera, Mycetophilidae). *Rev. Ent.* 21 (3): 561-580, 15 figs.
1952. Neotropical *Mycomyia*. II. (Diptera: Mycetophilidae). *Dusenía* 3 (3): 215-224, figs.
1959. A synopsis of American Mycomyini with description of new species (Diptera — Mycetophilidae). *Ent. Amer.* (1958) (n. s.) 38: 1-155, 17 pls.
1961. A Jamaican *Neoempheria* Osten Sacken, 1878 (Diptera: Mycetophilidae). *Bull. Brooklyn Ent. Soc.* 56: 128-130, 6 figs.

COHER, E. I. & J. LANE

1949. A new neotropical genus of "Mycetophilidae" (Diptera, Nematocera). *Rev. Brasil. Biol.* 9 (4): 485-488, 9 figs.

19E.54

COQUILLET, D. W.

1910. The type-species of the North American genera of Diptera. *Proc. U. S. Natl. Mus.* 37: 499-647.

COSTA, A.

1857. Contribuzione alla fauna ditterologica italiana. [G. Vico] *Giorn. Sci.*, Napoli 2: 438-460.

CURRAN, C. H.

1927. Four new American Diptera. *Amer. Mus. Nov.* 275: 1-4, 3 figs.
1928. Insects of Porto Rico and the Virgin Islands. Diptera or two-winged flies. Part 1, pp. 1-118, 39 figs., in New York Academy of Sciences, *Scientific Survey of Puerto Rico and The Virgin Islands*, 11. New York.

CURTIS, J.

1831. *British Entomology: Being illustrations and descriptions of the genera of insects found in Great Britain and Ireland* 8: pls. 338-383. London, "1823-1840".
1837. *Idem* 13: pls. 626-673. London.

DURET, J. P.

- 1973a. Una nueva especie de *Mycetophila* argentina (Diptera — Mycetophilidae). *Neotropica* 19 (59): 57-61, 3 figs.
1973b. *Mycetophila pucaraensis*, una nueva especie de la Patagonia (Diptera, Mycetophilidae). *Physis*, B. Aires (C) 32 (85): 471-474, 5 figs.
1973c. Nuevas especies argentinas de *Austrosynapha* Tonnoir, 1929 (Diptera — Mycetophilidae). *Rev. Soc. Ent. Argent.* 34 (1-2): 65-78, 18 figs.
1974. Nuevas especies argentinas del genero *Echinopodium* Freeman, 1951 (Diptera, Mycetophilidae). *Physis*, B. Aires (C) 33 (86): 93-117, 57 figs.
1975a. *Austrosynapha bidentata* una especie nueva de la Argentina. (Diptera — Mycetophilidae). *Rev. Soc. Ent. Argent.* (1973-1974) 34 (3-4): 311-312, 1 fig.
1975b. Cuatro especies nuevas del genero *Echinopodium* Freeman, 1951 (Diptera, Mycetophilidae). *Physis*, B. Aires (C) 34 (88): 75-81, 14 figs.
1976. Notas sobre el genero *Echinopodium* Freeman, 1951, en Chile, con la descripcion de catorce especies nuevas (Diptera, Mycetophilidae). *Physis*, B. Aires (C) 35 (90): 173-195, 43 figs.

DZIEDZICKI, H.

1885. Przyczynek do fauny owadów dwukrzylych. Rodzage nowe: *Hertwigia*, nov. gen., *Eurycera*, nov. gen. i gatunki rodzajów: *Boletina*, *Sciophila*. *Pamięt. Fizyograficzny* 5 (3): 164-194, pls. 4-9.
1889. Revue des espèces européennes du genre *Phronia* Winnertz, avec la description des deux genres nouveaux: *Macrobrachius* et

Megophthalmidia. *Horae Soc. Ent. Ross.* 23: 404-532, 17 figs., pls. 12-21.

EDWARDS, F. W.

1919. Diptères, Nématocères. *Mission Arc Mér. Amér. Sud*, Paris 10: 143-162, figs.
1931. New neotropical nematoceros Diptera. *Ann. Mag. Nat. Hist.* (10) 7: 255-261.
1932. New Brazilian Mycetophilidae (Diptera). *Rev. Ent.* 2 (2): 138-149, 6 figs.
1933. New neotropical Mycetophilidae (II). *Rev. Ent.* 3 (3): 303-322.
1934. New neotropical Mycetophilidae (III). *Rev. Ent.* 4 (3): 354-372, 5 figs.
1940a. New neotropical Mycetophilidae (IV) (Diptera). *Rev. Ent.* 11 (1-2): 440-465, 2 pls.
1940b. Neotropical *Neoempheria* (Diptera, Mycetophilidae). *Nov. Zool.* 42 (1): 107-129, pls. 1-2.
1941. Mycetophilidae (Diptera) collected by the expedition to Mato Grosso of the Brazilian Zoological Club, in July 1939. *Rev. Ent.* 12 (1-2): 303-314, pl. 15.

ENDERLEIN, G.

- 1910a. Neue Gattungen und Arten aussereuropäischen Fliegen. *Stettin. ent. Ztg* (1911) 72: 135-209, 4 figs.
1910b. Diptera, Mycetophilidae [of the Seychelles]. *Proc. Linn. Soc. London* (2) 14: 59-81.
1940. Die Dipterenfauna der Juan-Fernandez-Inseln und der Osterinsel, in Skottsberg, *Nat. Hist. of Juan Fernandez and Easter Island*, *Zool.* 3: 643-680, 49 figs. Uppsala.

FISHER, E. G.

1939. Costa Rican Mycetophilidae (Diptera). *Trans. Ent. Soc. Amer.* 65: 227-236, pl. 13.
1941. Notes on Costa Rican Mycetophilidae (Diptera). *Ent. News* 52: 8.

FREEMAN, P.

1951. Mycetophilidae, in British Museum (Natural History), *Diptera of Patagonia and South Chile* 3: 138 pp., 355 figs. London.
1954. Los insectos de las islas Juan Fernandez. 13. Mycetophilidae, Sciaridae, Cecidomyiidae and Scatopsidae (Diptera). *Rev. Chil. Ent.* (1953) 3: 23-40, 3 pls.

GIGLIO-TOS, E.

1890. Nuove specie di ditteri del Museo Zoologico di Torino. *Boll. Mus. Zool. Anat. Comp. Torino* 5 (84): 1-4.

HOLMGREN, N.

1907. Monographische Bearbeitung einer schalentragenden Mycetophilidenlarve (*Mycetophidla ancyliiformans* n. sp.). *Z. wiss. Zool.*, Leipzig 38: 1-77, 5 pls., 1 table.



INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.* 20: 339-342.

JOHANNSEN, O. A.

1909. Diptera, Fam. Mycetophilidae, in P. Wytzman, ed., *Genera Insectorum* 93: 141 pp., 7 pls. Bruxelles.
1912. The fungus gnats of North America. The Mycetophilidae of North America. Part III. *Bull. Maine Agric. Expt. Sta.* (1911) 196: 249-328, 5 pls. [i. e., figs. 98-245].

JOHNSON, C. W.

1919. A revised list of the Diptera of Jamaica. *Bull. Amer. Mus. Nat. Hist.* 41 (8): 421-449.

KERTÉSZ, K.

- 1901a. Neue und bekannte Dipteren in der Sammlung des Ungarischen National-Museums. *Termész. Füzetek* 24: 403-432, pl. 20.
- 1901b. *Neoglaphyoptera interrupta* n. sp. ♂. *Termész. Füzetek* 24: 495-496.
1902. Neue *Neoglaphyoptera*-Arten aus Süd-Amerika. *Termész. Füzetek* 25: 573-581.
1909. Vier neue südamerikanische Mycetophilidae. *Ann. Mus. Natl. Hung.* 7: 137-142, 2 figs.

KNAB, F. & R. H. VAN ZWALUWENBURG

1918. A second *Mycetophila* with dung bearing larva (Diptera; Mycetophilidae). *Ent. News* 29: 138-142, pl. 8.

LANE, J.

- 1947a. New Brazilian Mycetophilidae (Diptera, Nemocera). *Rev. Ent.* (1946) 17 (3): 339-360, 14 figs.
- 1947b. Novos Mycetophilinae do Brasil (Diptera, Mycetophilidae). *Rev. Ent.* 18 (3): 448-458, 7 figs.
- 1948a. Mycetophilinae de Boracéia, São Paulo (Diptera, Mycetophilidae). *Rev. Ent.* 19 (1-2): 231-278, 61 figs.
- 1948b. "Mycetophilidae" do Brasil (Diptera, Nematocera). *Rev. Brasil. Biol.* 8 (2): 247-254, 15 figs.
1950. New Brazilian "*Leia*" Meigen, 1818 (Diptera, Mycetophilidae). *Rev. Brasil. Biol.* 10 (1): 121-131, 17 figs.
- 1951a. Neotropical *Zygomyia* and *Exechia* (Diptera, Mycetophilidae). *Dusenía* 2 (4): 237-254, 15 figs.
- 1951b. Neotropical *Epicrypta* (Diptera, Nemocera, Mycetophilidae). *Dusenía* 2 (5): 327-340, 12 figs.
- 1952a. Some new Brazilian Mycetophilidae (Diptera, Nematocera). *Dusenía* 3 (2): 131-140, 5 figs.
- 1952b. Neotropical *Mycetophila* (Diptera, Mycetophilidae). *Dusenía* 3 (6): 421-434, 19 figs.



- 1952c. Neotropical "*Tetragoneura*" (Diptera, Mycetophilidae, Sciophilinae). *Rev. Brasil. Biol.* 12 (4): 401-409, 9 figs.
- 1954a. Neotropical Gnoristini (Diptera, Mycetophilidae). *Rev. Brasil. Ent.* 1: 93-112, 13 figs.
- 1954b. The genus *Megophthalmidia* in Brazil (Diptera, Mycetophilidae, Sciophilinae). *Rev. Brasil. Ent.* 2: 105-110, 2 figs.
- 1954c. Revision of the genus *Epicypa* Winnertz, 1863 in the neotropical region (Diptera, Mycetophilidae). *Rev. Brasil. Ent.* 2: 111-138, 15 figs.
- 1954d. Revision of "*Leiella*" Enderlein, 1911 (Diptera, Nematocera, Mycetophilidae). *Rev. Brasil. Biol.* 14 (4): 449-453, 4 figs.
1955. Neotropical *Mycetophila* Meigen excluding those of the Chilean subregion (Diptera: Mycetophilidae). *Trans. R. Ent. Soc. London* 106 (10): 393-420, 22 figs.
- 1956a. New Neotropical Sciophilinae (Diptera, Mycetophilidae). *Dusenia* 7 (3): 119-124, 8 figs.
- 1956b. New Mycetophilinae from the neotropics (Diptera, Mycetophilidae). *Rev. Brasil. Ent.* 5: 147-160, 8 figs.
- 1958a. On Amazonian Mycetophilidae (Dipt., Nematocera). *Studia Ent.* 1 (1-2): 209-216, 8 figs.
- 1958b. Mycetophilidae, chiefly from Argentina. *Proc. 10th Int. Congr. Ent.* (Montreal, 1956) 1: 143-162, 17 figs.
- 1958c. Further new "*Mycetophila*" (Diptera, Mycetophilidae, Mycetophilinae). *Rev. Brasil. Biol.* 18 (4): 397-408.
- 1959a. Note on neotropical "Mycetophilidae" (Diptera, Nematocera). *Rev. Brasil. Biol.* 19 (2): 183-190, 5 figs.
- 1959b. Insecta Amapaensia. — Diptera: Mycetophilidae. *Studia Ent.* 2: 105-118, 14 figs.
- 1960a. Mycetophilidae from Trinidad, B. W. I. (Diptera, Nematocera). *Studia Ent.* 3: 375-384, 9 figs.
- 1960b. Additional new neotropical *Epicypa* (Diptera, Mycetophilidae). *Studia Ent.* 3: 481-488, 6 figs.
- 1961a. Insecta Amapaensia. — Diptera: Mycetophilidae (Second contribution). *Studia Ent.* 4: 243-256, 7 figs.
- 1961b. Further new neotropical Mycetophilidae (Diptera, Nematocera). *Rev. Brasil. Ent.* 10: 1-15, 6 figs.
- 1962a. A brief note on neotropical Mycetophilidae. *Rev. Brasil. Biol.* 22 (1): 9-16, 4 figs.
- 1962b. Mycetophilidae from Chile (Dipt.). *Studia Ent.* 5: 527-548, 8 figs.
1963. Insecta Patagonica (Diptera — Mycetophilidae). *Rev. Soc. Cient. Argent.* (1962) 25 (1-4): 3-16, 3 figs.


LANE, J. & E. I. COHER

1950. The genus "*Schnusea*" Edwards, 1933 (Diptera, Mycetophilidae, Sciophilinae). *Rev. Brasil. Biol.* 10 (3): 279-283, 12 figs.

LOEW, H.

1850. Dipterologische Beiträge. Vierter Theil. *Öffentl. Prüf. d. Schüler, Friedrich-Wilhelms Gymnasium, zu Posen 1850: 1-40, 1 pl.*
1869. Diptera Americae septentrionalis indigena. Centuria octava. *Berl. ent. Zeit.* 13: 1-32. [Also 1872. *Diptera Americae septentrionalis indigena* 2 (Cent. 6-10), 300 pp., Berlin].

19E.58



LYNCH ARRIBÁLZAGA, F.

1892. *Dipterologia argentina: Mycetophilidae; Complemento a los Mycetophilidae. Bol. Acad. Nac. Cienc. Córdoba 12: 377-436; 2 pls.; 471-480. [Also separately published, as *Dipterologia argentina. Mycetophilidae*, 75 pp., 2 pls., Buenos Aires, 1892].*

MACQUART, J.

1834. *Histoire naturelle des insectes. Diptères 1: 578 pp., 12 pls. In N. E. Roret, ed., Collection des suites à Buffon. Paris.*

MARSHALL, P.

1896. New Zealand Diptera: N° 2. — Mycetophilidae. *Trans. Proc. N. Z. Inst. 28* (= n. s., 11): 250-309, pls. 8-13.

MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp. Paris.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. Insektenkunde 2: 259-281.*
1804. *Klassifikation und Beschreibung der europäischen zweiflügeligen Insekten (Diptera Linn.) 1* (1): xxviii + 142 pp., 8 pls.; (2): vi + pp. 153-314, pls. 9-15. Braunschweig.
1818. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten 1: xxxvi + 333 pp., 11 pls. Aachen.*

MIK, J.

1886. Dipterologische Miscellen I, II, III. *Wien. ent. Ztg 5: 101-102, 276-279, 317-318.*

OSTEN SACKEN, C. R.

1878. Catalogue of the described Diptera of North America (Ed. 2). *Smithson. Misc. Coll. 16* [2 (= publ. 270)]: 1-276.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool.-bot. Ges. Wien 15: 595-782, 7 pls.*

RONDANI, C.

1856. *Dipterologiae Italicae Prodromus. Vol. 1: Genera Italica ordinis dipteriorum ordinatim disposita et distincta et in familias et stirpes aggregata*, 228 pp. Parma.

SAY, T.

1829. Descriptions of North American dipterous insects. *Jour. Acad. Nat. Sci. Philad.* (1829-1830) 6: 149-178. [Also 1859, *The Complete Writings of Thomas Say...*].

SCHINER, I. R.

1868. *Diptera*, 388 pp., 4 pls., in B. von Wüllerstorff-Urbair, in charge, *Reise der österreichische Fregatte Novara. Zool. 2* (1, B). Wien.



SHAW, F.

1940. Some new Mycetophilidae from Costa Rica (Dipt.). Part I. *Rev. Ent.* 11 (3): 803-808, 8 figs.
1950. A new Brazilian "*Leia*" (Diptera, Mycetophilidae). *Rev. Brasil. Biol.* 10 (1): 113-114, 1 fig.
1951. Some notes on synonymy of the Mycetophilidae (Diptera). *Psyche* 58: 148.

SHAW, F. & J. LANE

1950. Key for neotropical "*Leia*" Meigen, 1818 (Diptera, Mycetophilidae). *Rev. Brasil. Biol.* 19 (2): 253-256.

SKUSE, F. A. A.

1888. Diptera of Australia. Part III. The Mycetophilidae. *Proc. Linn. Soc. N. S. W.* (2) 3 [= whole ser., 12]: 1123-1222, pls. 31-32.
1890. Diptera of Australia. Nematocera. Supplement II. *Proc. Linn. Soc. N. S. W.* (2) 5 [= whole ser., 14]: 595-640, pl. 19.

SPEISER, P.

1911. Zur Kenntnis aussereuropäischer Dipteren. *Wiesbaden Jahrb. Ver. Naturkunde* 64: 237-261.

STONE, P. G.

1941. The generic names of Meigen 1800 and their proper application. (Diptera). *Ann. Ent. Soc. Amer.* 34: 404-418.

STROBL, P. G.

1893. Beiträge zur Dipteren-Fauna des österreichischen Littorale. *Wien. ent. Ztg* 12: 29-42, 74-80, 89-108, 121-136, 160-170.

TONNOIR, A. L.

1929. Australian Mycetophilidae. Synopsis of the genera. *Proc. Linn. Soc. N. S. W.* 54: 584-614, 7 figs., pls. 22-23.

TUOMIKOSKI, R.

- 1966a. Generic taxonomy of the Exechiini (Dipt., Mycetophilidae). *Ann. Ent. Fenn.* 32 (2): 159-194.
- 1966b. On the subfamily Manotinae (Edw.) (Dipt., Mycetophilidae). *Ann. Ent. Fenn.* 32 (3): 211-223, 6 figs.

WALKER, F.

1837. Descriptions, etc., of the insects collected by Cpt. P. P. King, R. N., F. R. S., in the survey of the Straits of Magellan. Diptera. *Trans. Linn. Soc. London* 17: 331-359.

WIEDEMANN, C. R. W.

1828. *Aussereuropäische zweiflügeligen Insekten* 1: xxxii + 608 pp., 7 pls. Hamm.

19E.60



WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896: 253-446, pls. 8-14.
1900. Supplement (part), pp. 217-248, in F. D. Godman & O. Salvin, eds., *Biologia Centrali-Americana. Zoologia-Insecta-Diptera* 1: 378 pp., 6 pls. London.

WINNERTZ, J.

1846. Beschreibung einiger neuen Gattung aus der Ordnung der Zweiflügl. *Stettin. ent. Ztg* 7: 11-20, 2 pls.
1863. Beitrag zu einer Monographie der Pilzmücken. *Verh. zool. bot. Ges. Wien* 13: 637-964, 4 pls.

WULP, F. M. VAN DER

1876. Verslag van de dertigste zomervergadering der Nederlandsche Entomologische Vereeniging gehouden te Amsterdam op Zaterdag 24 Julij 1875. *Tijdschr. v. Ent.* 19 (spec. pagination): i-liv, 2 figs.

INDEX

<i>abbreviata</i> (Loew), 1869 (Leia), Rondaniella	19E.33
<i>aberrans</i> Philippi, 1865, Sciophila	19E.52
<i>Abmyceta</i> Lane, 1955	19E.9
<i>abnormalis</i> Fisher, 1939, Monoclona	19E.50
<i>absyrtia</i> Lane, 1954, Dziedzickia	19E.23
<i>acarensis</i> Lane, 1952, Mycetophila (Mycetophila)	19E.14
<i>acariensis</i> (Lane), 1951 (Exechia), Boraceomyia	19E.2
<i>acarisi</i> Lane, 1958, Mycetophila (Abmyceta)	19E.9
<i>acidoti</i> Coher, 1959, Neoempheria	19E.43
<i>Acnemia</i> Winnertz, 1863	19E.47
<i>aczeli</i> Lane, 1958, Epicypta (Epicypta)	19E.7
<i>aczeli</i> Lane, 1959, Leia	19E.27
<i>Agaricobia</i> Philippi, 1865	19E.47
<i>aguarasi</i> Lane & Coher, 1950, Schnusea	19E.25
<i>aguarensis</i> Lane, 1951, Epicypta (Epicypta)	19E.7
<i>aguarensis</i> Lane, 1951, Zygomomyia	19E.20
<i>airosal</i> Lane, 1947, Rymosia	19E.4
<i>aitkeni</i> Lane, 1956, Cluzobra	19E.49
<i>aitkeni</i> (Lane), 1960 (Exechia), Boraceomyia	19E.2
<i>albicoxa</i> Enderlein, 1910, Leia	19E.27
<i>alexanderi</i> Coher, 1950, Mycomyia	19E.38
<i>Allocotocera</i> Mik, 1886	19E.48
<i>Allodia</i> , authors	19E.3
<i>Allodiopsis</i> Tuomikoski, 1966	19E.2
<i>amabilis</i> (Williston), 1900 (Neoglaphyoptera), Leia	19E.27
<i>amapaensis</i> Lane, 1961, Epicypta (Callicypta)	19E.6
<i>amapaensis</i> Lane, 1959, Leia	19E.27
<i>americana</i> (Schiner), 1868 (Sciophila), Mycomyia	19E.38
<i>amplipennis</i> Freeman, 1951, Mycetophila (Mycetophila)	19E.14
<i>analis</i> Fisher, 1939, Leia	19E.28

19E.61



anambesi Lane, 1950, <i>Epicypa</i> (<i>Callicypa</i>)	19E.6
ancyliformans (Holmgren), 1907 (<i>Mycetophila</i>), <i>Epicypa</i>	19E.8
andina Lane, 1962, <i>Tetragoneura</i>	19E.33
andirai Lane, 1950, <i>Leia</i>	19E.27
Aneura Marshall, 1896	19E.48
angustifurca Enderlein, 1940, <i>Mycetophila</i>	19E.17
anhangaensis Lane, 1961, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.9
ansata Freeman, 1951, <i>Mycomya</i>	19E.38
antarctica (Bigot), 1888 (<i>Boletina</i>), <i>Paraleia</i>	19E.32
antarctica Walker, 1837, <i>Sciophila</i>	19E.52
<i>Aphanizophleps</i> Enderlein, 1910	19E.1
Aphelomera Skuse, 1888	19E.48
Aphrastomyia Coher & Lane, 1949	19E.26
<i>apicalis</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>apicalis</i> (Kertész), 1902 (<i>Neoglyphyoptera</i>), <i>Leia</i>	19E.27
<i>apicalis</i> Kertész, 1909, <i>Neoempheria</i>	19E.43
<i>apicata</i> Philippi, 1865, <i>Mycetophila</i>	19E.17
<i>apinagei</i> Lane, 1950, <i>Leia</i>	19E.27
<i>arapascoi</i> Lane, 1954, <i>Epicypa</i> (<i>Bosomyia</i>)	19E.5
<i>arauasi</i> Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.9
<i>araucana</i> Lane, 1962, <i>Mycetophila</i> (<i>Oromyceta</i>)	19E.16
<i>araucaniense</i> (Lane), 1962 (<i>Allodia</i>), <i>Brevicornu</i>	19E.3
<i>araucanum</i> Duret, 1974, <i>Echinopodium</i>	19E.36
<i>arauensis</i> Lane, 1952, <i>Tetragoneura</i>	19E.33
<i>ardeiceps</i> Freeman, 1951, <i>Tetragoneura</i>	19E.33
<i>arecunai</i> Lane, 1955, <i>Mycetophila</i> (<i>Oromyceta</i>)	19E.16
<i>arethusa</i> Coher, 1952, <i>Mycomya</i>	19E.39
<i>argentina</i> Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.9
<i>argentina</i> Lane, 1961, <i>Zygomyia</i>	19E.20
<i>argentense</i> Duret, 1975, <i>Echinopodium</i>	19E.36
<i>argentense</i> (Lane), 1958 (<i>Allodia</i>), <i>Brevicornu</i>	19E.4
<i>argentinensis</i> Duret, 1972, <i>Austrosynapha</i>	19E.21
<i>armata</i> Freeman, 1951, <i>Aphelomera</i>	19E.48
<i>armata</i> Freeman, 1951, <i>Dziedzickia</i>	19E.23
<i>armatum</i> Duret, 1976, <i>Echinopodium</i>	19E.36
<i>armatura</i> Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.14
<i>arnaudi</i> Lane, 1962, <i>Leiella</i>	19E.31
<i>aruaci</i> Lane, 1950, <i>Leia</i>	19E.27
<i>aspasia</i> Lane, 1954, <i>Dziedzickia</i>	19E.23
<i>atricornis</i> Philippi, 1865, <i>Mycetophila</i>	19E.17
<i>aurantiaca</i> Edwards, 1934, <i>Zygomyia</i>	19E.20
<i>australis</i> Philippi, 1865, <i>Sciophila</i>	19E.52
<i>austrobligua</i> Coher, 1959, <i>Mycomya</i>	19E.39
<i>Austrosynapha</i> Tonnoir, 1929	19E.21
<i>avicularia</i> Freeman, 1951, <i>Mycetophila</i>	19E.17
<i>axillipunctum</i> Enderlein, 1910, <i>Leia</i>	19E.27
<i>aysense</i> Duret, 1976, <i>Echinopodium</i>	19E.36
<i>bacairisi</i> Lane, 1954, <i>Epicypa</i> (<i>Callicypa</i>)	19E.6
<i>bacilliger</i> Edwards, 1932, <i>Tetragoneura</i>	19E.33
<i>banhumai</i> Lane, 1952, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.14
<i>bapianensis</i> Lane, 1951, <i>Epicypa</i> (<i>Epicypa</i>)	19E.7
<i>barai</i> Lane, 1954, <i>Epicypa</i> (<i>Bosomyia</i>)	19E.5



bari Lane & Coher, 1950, Schnusea	19E.25
barrettoi (Lane), 1947 (Delopsis), Mycetophila (Abmyceta)	19E.9
basalis Freeman, 1951, Anoura	19E.48
basalis Freeman, 1951, Dziedzickia	19E.23
basalis Freeman, 1951, Mycetophila (Mycetophila)	19E.14
basinerva Freeman, 1951, Eudicrana	19E.49
basinerva Freeman, 1951, Mycomya	19E.39
beatricea Lane, 1960, Dziedzickia	19E.23
beckeri (Enderlein), 1910 (Parastemma), Tetragoneura	19E.33
bejaranoi Duret, 1972, Austrosynapha	19E.21
bejaranoi Duret, 1974, Echinopodium	19E.36
bequaerti Coher, 1950, Mycomya	19E.39
<i>biamputata</i> Edwards, 1933, Leia	19E.29
bicellii Lane, 1961, Epicypa (Callicypa)	19E.6
bicolor Fisher, 1939, Leiella	19E.31
bicolor Edwards, 1934, Megophthalmidia	19E.31
bicolor (Enderlein), 1910 (Acnemia), Monoclona	19E.51
bicolor Edwards, 1934, Zygomya	19E.20
bicolorata Lane, 1948, Sceptonia	19E.19
bidentata Coher, 1959, Neoempheria	19E.43
bidentata Duret, 1975, Austrosynapha	19E.21
bifida Coher, 1959, Neoempheria	19E.39
bifida (Freeman), 1951 (Exechia), Exechiopsis (Xenexechia)	19E.3
bifida Freeman, 1951, Mycetophila (Mycetophila)	19E.14
bifida Freeman, 1951, Mycomya	19E.43
bifida Freeman, 1951, Tetragoneura	19E.33
bifila Freeman, 1951, Mycetophila	19E.17
biflagellata Edwards, 1940, Neoempheria	19E.43
bifurcatum Duret, 1974, Echinopodium	19E.36
bilobata Edwards, 1940, Neoempheria	19E.43
bilunula Wiedemann, 1828, Lea	19E.27
binocellaris (Edwards), 1934 (Acnemia), Cluzobra	19E.49
bipartita (Lynch Arribalzaga), 1892 (Glaphyroptera), Leia	19E.27
bipectinata Edwards, 1940, Neoempheria	19E.43
bipunctata Fisher, 1949, Leia	19E.27
bisetosa Freeman, 1951, Mycetophila (Mycetophila)	19E.14
bivittata Say, 1829, Leia	19E.27
blanchardi Enderlein, 1910, Mycetophila	19E.17
boracensis Coher, 1959, Mycomya	19E.39
boracensis Lane, 1962 (Allodia)	19E.5
boracensis Lane, 1948, Mycetophila (Mycetophila)	19E.14
Boraceomyia Lane, 1948	19E.2
borgmeieri Edwards, 1932, Mycetophila (Abmyceta)	19E.9
borgmeieri Edwards, 1940, Neoempheria	19E.44
borgmeieri Edwards, 1932, Tetragoneura	19E.34
borgmeieri (Lane), 1951 (Exechia), Boraceomyia	19E.2
borgmeieri Lane, 1959, Leia	19E.28
borinquensis Coher, 1959, Mycomya	19E.39
borinquensis (Lane), 1951 (Exechia), Boraceomyia	19E.2
boscii (Lane), 1948 (Delopsis), Epicypa (Bosomyia)	19E.5
Bosomyia Lane, 1954	19E.5
bradleyi Edwards, 1940, Neoempheria	19E.44
<i>brasiliana</i> (Lane), 1947 (Trichonta), Neallodia	19E.4
brasiliana (Edwards), 1932 (Delopsis), Epicypa (Epicypa)	19E.7



brasiliana Lane, 1947, Zygomysia	19E.21
brasiliensis Coher, 1959, Neoempheria	19E.44
brasiliensis (Enderlein), 1910 (Mycetophila), Epicypta (Epicypta)	19E.7
brevicauda Coher, 1959, Neoempheria	19E.44
brevicornis (Enderlein), 1910 (Aliodia), Cordyla	19E.4
Brevicornu Marshall, 1896	19E.3
brevifurcata Enderlein, 1910, Mycomysa	19E.39
brevifurcata (Freeman), 1951 (Exechia), Exechiopsis (Xenexechia)	19E.3
brevifurcata Freeman, 1951, Mycetophila (Mycetophila)	19E.14
bridgesi Lane, 1963, Mycetophila (Abmyceta)	19E.9
brontes Coher, 1952, Mycomysa	19E.39
brunnescens Freeman, 1951, Mycetophila (Mycetophila)	19E.14
burenari Lane, 1952, Tetragoneura	19E.34
butleri Lane, 1959, Cluzobra	19E.49
caetensis Lane, 1952, Tetragoneura	19E.34
caibii Lane & Coher, 1950, Schnusea	19E.25
caipora Lane, 1961, Dziedzickia	19E.23
cajuensis (Lane), 1951 (Exechia), Boraceomyia	19E.2
Callicypta Lane, 1954	19E.6
callidum (Johannsen), 1912 (Allodia), Brevicornu	19E.4
calopus (Bigot), 1888 (Sciophila), Tetragoneura	19E.34
campestra Coher, 1959, Mycomysa	19E.39
canicula Freeman, 1951, Mycetophila (Abmyceta)	19E.9
capitata (Enderlein), 1910 (Plastacephala), Epicypta	19E.8
caribai Lane, 1955, Mycetophila (Abmyceta)	19E.9
caripunai Lane, 1955, Mycetophila (Abmyceta)	19E.9
carrerai Coher, 1950, Mycomysa	19E.39
carrerai Lane, 1958, Mycetophila (Abmyceta)	19E.10
castanea Freeman, 1951, Paraleia	19E.32
catharinae Edwards, 1932, Mycetophila (Mycetophila)	19E.14
catharinensis Lane, 1954, Leiella	19E.31
caudata Edwards, 1933, Paraleia	19E.32
cayapoi Lane, 1950, Leia	19E.28
cayuensis Lane, 1955, Mycetophila (Abmyceta)	19E.10
cekalovici Duret, 1976, Echinopodium	19E.36
cellularis Edwards, 1940, Megalopelma	19E.50
cerqueirai Lane, 1961, Aphrastomyia	19E.26
chavantesi Lane, 1951, Zygomysia	19E.21
chilense Duret, 1975, Echinopodium	19E.36
chilensis (Blanchard), 1852 (Sciophila), Mycomysa	19E.39
chilensis Philippi, 1865 (Gnoriste)	19E.54
chiloense Coher, 1959, Echinopodium	19E.36
chiriguanai Lane, 1954, Epicypta (Callicypta)	19E.6
chlorochroa Freeman, 1951, Mycetophila	19E.17
ciliata Edwards, 1940, Sciophila	19E.52
cincta Shaw, 1940, Neoempheria	19E.44
cinticauda Enderlein, 1910, Leia	19E.28
cincticornis Edwards, 1940, Sciophila	19E.52
cinctifemur Shaw, 1940, Sciophila	19E.53
cinerea Freeman, 1951, Paraleia	19E.32



cinerea (Freeman), 1951 (Rhymosia), Allodiopsis (Gymnogonia)	19E.2
circumfera Fisher, 1939, Leia	19E.28
citrina Shaw, 1940, Mycomya	19E.39
claripennis Edwards, 1931, Eudicrana	19E.49
clarovittata Freeman, 1951, Mycetophila (Abmyceta)	19E.10
clavata (Lynch Arribálzaga), 1892 (Sciophila), Mycomya	19E.39
clavigera Freeman, 1951, Mycetophila (Abmyceta)	19E.10
cleta Coher, 1952, Mycomya	19E.39
clio Lane, 1954, Dziedzickia	19E.23
Cluzobra Edwards, 1940	19E.49
<i>Cnephaeophila</i> Philippi, 1865	19E.38
Coelosia Winnertz, 1863	19E.23
cognata Philippi, 1865, Mycetophila	19E.17
coheri Duret, 1974, Echinopodium	19E.36
coheri Lane, 1954, Dziedzickia	19E.23
coheri Lane, 1954, Epicypta (Callicypta)	19E.6
collarigera Enderlein, 1910, Leia	19E.28
comes Coher, 1959, Neoempheria	19E.44
completa (Kertész), 1902 (Neoglaphyroptera), Leia	19E.28
complexum (Lane), 1948 (Allodia), Brevicornu	19E.4
concinna (Williston), 1896 (Neoglaphyroptera), Leia	19E.28
concurrans Edwards, 1933, Tetragoneura	19E.34
<i>confusa</i> Lane, 1948, Mycetophila (Abmyceta)	19E.13
conifera Freeman, 1951, Mycetophila (Abmyceta)	19E.10
conjuncta Freeman, 1951, Mycetophila (Mycetophila)	19E.14
connecta Coher, 1959, Mycomya	19E.39
constricta Freeman, 1951, Mycetophila (Abmyceta)	19E.10
cornuta Freeman, 1951, Mycetophila	19E.17
coronatum Duret, 1976, Echinopodium	19E.36
costalimai Edwards, 1940, Neoempheria	19E.44
costariensis Coher, 1959, Neoempheria	19E.44
costaricensis Fisher, 1940, Leia	19E.28
costariensis Lane, 1958, Mycetophila (Abmyceta)	19E.10
couasi Lane, 1954, Epicypta (Callicypta)	19E.6
coxalis Freeman, 1951, Mycomya	19E.39
coxata (Enderlein), 1910 (Aphanizophleps), Manota	19E.1
coxiopensis Lane, 1952, Allocotocera	19E.48
coxiopensis Lane, 1958, Mycetophila (Abmyceta)	19E.10
cramptoni Coher, 1950, Mycomya	19E.40
cramptoni Coher & Lane, 1949, Aphrastomyia	19E.26
cristata Freeman, 1951, Aphelomera	19E.48
cryptura Edwards, 1941, Dziedzickia	19E.23
curiaensis Lane, 1952, Mycetophila (Mycetophila)	19E.15
curiopiensis Lane, 1961, Epicypta (Epicypta)	19E.7
curvispina Freeman, 1951, Echinopodium	19E.36
cuzaensis Edwards, 1933, Docosia	19E.26
cylindrica Freeman, 1951, Mycomya	19E.40
cypriformis Freeman, 1951, Thoracotropls	19E.35
damascenoi Lane, 1958, Rymosia	19E.4
defecta Williston, 1896, Manota	19E.1
deflecta Coher, 1959, Neoempheria	19E.44
<i>Delopsis</i> Skuse, 1890	19E.5



demacuri Lane, 1955, Mycetophila (Abmyceta)	19E.10
dentata Freeman, 1951, Austrosynapha	19E.22
dentata Freeman, 1951, Tetragoneura	19E.34
derunei Lane, 1952, Tetragoneura	19E.34
desanei Lane & Coher, 1950, Schnusea	19E.25
diana Lane, 1959, Dziedzickia	19E.24
dichaeta Freeman, 1951, Mycetophila (Abmyceta)	19E.10
dichroma Freeman, 1951, Leia	19E.28
digitalis Freeman, 1951, Echinopodium	19E.36
digitalis Freeman, 1951, Mycetophila	19E.17
digitata Edwards, 1940, Monoclona	19E.51
<i>diluta</i> (Williston), 1896 (Sciophila), Mycomya	19E.41
<i>Diomonus</i> Walker, 1848	19E.50
disgrega Edwards, 1933, Leia	19E.28
distincta Freeman, 1951, Leiella	19E.31
distincta Freeman, 1951, Mycetophila (Abmyceta)	19E.10
distincta Fisher, 1939, Novakia	19E.51
divergens Edwards, 1932, Megophthalmidia	19E.31
diversicornis (Kertész), 1902 (Neoglaphyoptera), Leia	19E.28
diversipes Edwards, 1933, Leia	19E.28
divisa Freeman, 1951, Mycomya	19E.40
Docosia Winnertz, 1863	19E.26
dolosa Williston, 1896, Mycetophila	19E.17
dominica Curran, 1927, Mycetophila	19E.17
dorsimacula Enderlein, 1910, Mycomya	19E.41
dryope Coher, 1952, Mycomya	19E.40
Juanasi Lane, 1951, Epicypa (Epicypa)	19E.7
duanensis Lane, 1952, Mycetophila (Mycetophila)	19E.15
dubitans Lane, 1954, Dziedzickia	19E.24
dumeta Coher, 1959, Mycomya	19E.40
Dziedzickia Johannsen, 1909	19E.23
diedzickii Coher, 1959, Neoempheria	19E.44
Echinopodium Freeman, 1951	19E.36
ecuadorensis Coher, 1959, Neoempheria	19E.44
edwardsi Lane, 1948, Boraceomyia	19E.3
edwardsi Lane, 1960, Cluzobra	19E.49
edwardsi Lane, 1948, Mycetophila (Mycetophila)	19E.16
edwardsiana Lane, 1952, Dziedzickia	19E.24
edwardsiana Lane, 1954, Megophthalmidia	19E.31
edwardsii Coher, 1960, Mycomya	19E.40
elegans Duret, 1976, Echinopodium	19E.37
elegans Lane, 1948, Eudicrana	19E.49
elegans (Kertész), 1902 (Neoglaphyoptera), Leia	19E.28
embuensis (Lane), 1956 (Exechia), Boraceomyia	19E.2
<i>Empheria</i> Winnertz, 1863	19E.43
enderleini Edwards, 1940, Neoempheria	19E.44
endymion Lane, 1959, Dziedzickia	19E.24
epacra Coher, 1959, Mycomya	19E.40
Epicypa Winnertz, 1863	19E.7
equatoriana Lane, 1962, Megophthalmidia	19E.32
eramalai Lane, 1952, Tetragoneura	19E.34
eramansensis Lane, 1955, Mycetophila (Abmyceta)	19E.10




eruliense (Lane), 1956 (Allodia), Brevicornu	19E.4
Eudicrana Loew, 1869	19E.49
Eurycera Dziedzicki, 1885	19E.48
evanescens (Enderlein), 1910 (Pleonazoneura), Neoempheria	19E.44
excerpta Coher, 1959, Mycomya	19E.40
Exechia, authors	19E.2
Exechiini	19E.2
Exechiopsis Tuomikoski, 1966	19E.3
extensa (Freeman), 1951 (Exechia), Exechiopsis (Xenexechia)	19E.3
faceta Coher, 1959, Neoempheria	19E.44
falcifera Freeman, 1951, Mycomya	19E.40
falcigera Freeman, 1951, Paraleia	19E.32
falculata Edwards, 1933, Leia	19E.28
fallax Coher, 1959, Neoempheria	19E.44
faniipennis Lane, 1952, Tetragoneura	19E.34
fasciata Freeman, 1951, Mycetophila (Mycetophila)	19E.14
fasciata (Kertész), 1902 (Neoglyphyoptera), Leia	19E.28
fasciata Freeman, 1951, Phthinia	19E.52
fasciata Freeman, 1951, Trichonta	19E.20
fasciculatus Edwards, 1933, Leptomorphus	19E.50
fascipennis Edwards, 1940, Cluzobra	19E.49
fascipennis Philippi, 1865, Mycetophila	19E.17
femoratus Edwards, 1933, Leptomorphus	19E.50
fenestralis (Philippi), 1865 (Cnephaeophila), Mycomya	19E.40
ferrazi Coher, 1950, Mycomya	19E.40
ferrazi Lane, 1955, Mycetophila (Oromyceta)	19E.16
fidelis Edwards, 1940, Sciophila	19E.53
fielbrigi Edwards, 1934, Dziedzickia	19E.24
filicauda (Edwards), 1934 (Synapha), Austrosynapha	19E.22
fisherae Shaw, 1950, Leia	19E.28
flabellifera Freeman, 1951, Mycetophila	19E.17
flabellifera Freeman, 1951, Parvicellula	19E.52
flagellata Freeman, 1951, Phthinia	19E.52
flagellatum Freeman, 1951, Echinopodium	19E.37
flavescens Freeman, 1951, Mycomya	19E.40
flavicauda Edwards, 1941, Tetragoneura	19E.34
flavicincta Freeman, 1951, Tetragoneura	19E.34
flavicornis Edwards, 1940, Neoempheria	19E.44
flavicoxa Edwards, 1940, Neoempheria	19E.44
flavicoxa Freeman, 1951, Acnemia	19E.47
flavicoxa Freeman, 1951, Allocotocera	19E.48
flavida Edwards, 1932, Neallodia	19E.4
flavipalpis Freeman, 1951, Synapha	19E.25
flavipes (Enderlein), 1910 (Plastacephala), Epicypta (Epicypta)	19E.8
flavithorax Freeman, 1951, Coelosia	19E.23
flavithorax Freeman, 1951, Mycetophila	19E.17
flavolunata Freeman, 1951, Mysetophila (Abmyceta)	19E.10
flavonigra Lane, 1954, Dziedzickia	19E.24
flavoscutellata (Lynch Arribálzaga), 1892 (Glyphyoptera), Leia	19E.29
flexiseta Freeman, 1951, Mycetophila (Abmyceta)	19E.10
forattinii Lane, 1960, Epicypta (Bosomyia)	19E.5
forattinii Lane, 1955, Mycetophila (Abmyceta)	19E.10



<i>forcipata</i> Freeman, 1951, <i>Austrosynapha</i>	19E.22
<i>forcipata</i> Freeman, 1951, <i>Mycomya</i>	19E.40
<i>formosensis</i> (Lynch Arribázaga), 1892 (<i>Sciophila</i>), <i>Neoempheria</i>	19E.45
<i>fractinervis</i> Edwards, 1940, <i>Sciophila</i>	19E.53
<i>fraudulenta</i> (Williston), 1896 (<i>Phthinia</i>), <i>Megalopelma</i>	19E.50
<i>freemani</i> Coher, 1950, <i>Mycomya</i>	19E.40
<i>freemani</i> Duret, 1974, <i>Echinopodium</i>	19E.37
<i>freemani</i> Lane, 1956, <i>Acnemia</i>	19E.47
<i>freemani</i> Lane, 1960, <i>Austrosynapha</i>	19E.22
<i>freemani</i> Lane, 1954, <i>Dziedzickia</i>	19E.24
<i>freemani</i> Lane, 1954, <i>Megophthalmidia</i>	19E.32
<i>freemani</i> Lane, 1948, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
<i>freemani</i> Lane, 1951, <i>Zygomya</i>	19E.21
<i>fulva</i> Lane, 1954, <i>Leiella</i>	19E.31
<i>fulva</i> Lane, 1954, <i>Synapha</i>	19E.23
<i>fuivicolis</i> (Philippi), 1865 (<i>Agaricobia</i>), <i>Acnemia</i>	19E.48
<i>fumipennis</i> Freeman, 1951, <i>Synapha</i>	19E.25
<i>funebis</i> Freeman, 1951, <i>Mycomya</i>	19E.40
<i>funerea</i> Freeman, 1951, <i>Acnemia</i>	19E.48
<i>funerea</i> Freeman, 1951, <i>Dziedzickia</i>	19E.24
<i>funerea</i> (Freeman), 1951 (<i>Exechia</i>), <i>Exechiopsis</i> (<i>Xenexechia</i>)	19E.3
<i>funerea</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11
<i>funerea</i> Freeman, 1951, <i>Paraleia</i>	19E.32
<i>funerea</i> Freeman, 1951, <i>Synapha</i>	19E.25
<i>funerea</i> Freeman, 1951, <i>Trichonta</i>	19E.20
<i>Fungivora</i> Meigen, 1800	19E.9
<i>furcata</i> Lane, 1954, <i>Austrosynapha</i>	19E.22
<i>furcata</i> Freeman, 1951, <i>Phthinia</i>	19E.52
<i>furcata</i> Freeman, 1951, <i>Procydoneura</i>	19E.33
<i>furcilla</i> (Freeman), 1954 (<i>Exechia</i>), <i>Exechiopsis</i> (<i>Xenexechia</i>)	19E.3
<i>fusca</i> Edwards, 1934, <i>Stenophragma</i>	19E.53
<i>fusceens</i> Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
<i>fuscicornis</i> Edwards, 1941, <i>Leia</i>	19E.28
<i>fuscicornis</i> Freeman, 1951, <i>Mycomya</i>	19E.40
<i>fuscipennis</i> Edwards, 1940, <i>Cluzobra</i>	19E.49
<i>fuscus</i> Freeman, 1951, <i>Pleurogymnus</i>	19E.19
<i>galea</i> Freeman, 1951, <i>Tetragoneura</i>	19E.34
<i>galibisi</i> Lane, 1955, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11
<i>galindoi</i> Lane, 1961, <i>Dziedzickia</i>	19E.24
<i>gentilii</i> Duret, 1975, <i>Echinopodium</i>	19E.37
<i>glandularis</i> Edwards, 1940, <i>Allocotocera</i>	19E.48
<i>Glaphyroptera</i> Winnertz, 1863	19E.27
<i>glochis</i> Coher, 1959, <i>Neoempheria</i>	19E.45
<i>gloriosa</i> Lane, 1959, <i>Dziedzickia</i>	19E.24
<i>Gnoristini</i>	19E.21
<i>goiana</i> Coher, 1959, <i>Neoempheria</i>	19E.45
<i>goianensis</i> (Lane), 1947 (<i>Delopsis</i>), <i>Epicypa</i> (<i>Epicypa</i>)	19E.8
<i>goianensis</i> (Lane), 1946 (<i>Exechia</i>), <i>Boraceomyia</i>	19E.2
<i>golbachi</i> Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11
<i>golbachi</i> Lane, 1961, <i>Zygomya</i>	19E.21
<i>gracile</i> Kertész, 1901 (<i>Dynatosoma</i>)	19E.5
<i>gradata</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11



guaiaensis Lane, 1951, <i>Tetragoneura</i>	19E.34
guanasi Lane, 1958, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
guanensis Lane, 1951, <i>Epicypa</i> (<i>Epicypa</i>)	19E.8
guaraiasi Lane, 1952, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
guarani Lane, 1951, <i>Epicypa</i> (<i>Epicypa</i>)	19E.8
guatensis Lane, 1955, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11
guaycurusi Lane, 1950, <i>Leia</i>	19E.29
<i>Gymnognonia</i> Tuomikoski, 1966	19E.2
halterata (Kertész), 1902 (<i>Neoglaphyroptera</i>), <i>Leia</i>	19E.29
hamadryas Coher, 1952, <i>Mycomya</i>	19E.40
hansoni Coher, 1950, <i>Mycomya</i>	19E.41
heros Lane, 1951, <i>Zygomyia</i>	19E.21
herteli Lane, 1951, <i>Zygomyia</i>	19E.21
<i>Hertwigia</i> Dziedzicki, 1885	19E.23
heteroneura Philippi, 1865, <i>Mycetophila</i>	19E.17
<i>Homaspis</i> Skuse, 1888	19E.53
horrens Coher, 1959, <i>Neocempheria</i>	19E.45
huarpesi Lane, 1952, <i>Tetragoneura</i>	19E.34
humboldti (Lane), 1948 (<i>Delopsis</i>), <i>Mycetophila</i> (<i>Abmyceta</i>) ...	19E.11
humeralis Edwards, 1934, <i>Stenophragma</i>	19E.53
humile (Lane), 1948 (<i>Allodia</i>), <i>Brevicornu</i>	19E.4
hypspilic Lane, 1954, <i>Dziedzickia</i>	19E.24
iara Lane, 1961, <i>Dziedzickia</i>	19E.24
iaunai Lane, 1952, <i>Tetragoneura</i>	19E.34
iaunensis Lane, 1951, <i>Epicypa</i> (<i>Epicypa</i>)	19E.8
ibarrai Duret, 1973, <i>Austrosynapha</i>	19E.22
icosi Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.11
idalis Lane, 1954, <i>Synapha</i>	19E.25
iheringi Lane, 1948, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
illita Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
incompleta Curran, 1928, <i>Leia</i>	19E.29
<i>incompleta</i> Edwards, 1933, <i>Leia</i>	19E.30
inermis Freeman, 1951, <i>Aphelomera</i>	19E.49
inermis Freeman, 1951, <i>Mycomya</i>	19E.41
infirma (Lynch Arribáizaga), 1892 (<i>Sciophila</i>), <i>Mycomya</i>	19E.41
infuscata Freeman, 1951, <i>Mycomya</i>	19E.41
infuscata Freeman, 1951, <i>Paratrizygia</i>	19E.51
inornata Lane, 1948, <i>Epicypa</i> (<i>Callicypa</i>)	19E.6
insecta Freeman, 1951, <i>Mycetophila</i>	19E.18
insipiens (Williston), 1896 (<i>Mycetophila</i>), <i>Epicypa</i>	19E.9
insolitum Duret, 1976, <i>Echinopodium</i>	19E.37
intermedia Lane, 1954, <i>Dziedzickia</i>	19E.24
intermedia Edwards, 1940, <i>Stenophragma</i>	19E.53
interrupta (Kertész), 1902 (<i>Neoglaphyroptera</i>), <i>Leia</i>	19E.29
ionica Lane, 1959, <i>Dziedzickia</i>	19E.24
iphis Coher, 1952, <i>Mycomya</i>	19E.41
iranchesii Lane, 1960, <i>Epicypa</i> (<i>Callicypa</i>)	19E.6
irene Coher, 1952, <i>Mycomya</i>	19E.45
iumentis Lane, 1954, <i>Epicypa</i> (<i>Bosomyia</i>)	19E.5



jaffuelensis Freeman, 1951, Mycomya	19E.41
jamaicensis Coher, 1961, Neoempheria	19E.45
janus Lane, 1959, Dzedzickia	19E.24
jaruensis Lane, 1951, Epicypa (Epicypa)	19E.8
jaruai Lane, 1952, Leiella	19E.31
javaesi Lane, 1955, Mycetophila (Oromyceta)	19E.16
johannseni (Enderlein), 1910 (Pleonazoneura), Neoempheria	19E.45
johannseni (Lane), 1947 (Delopsis), Epicypa (Callicypa)	19E.6
jugalisi Coher, 1959, Neoempheria	19E.45
juinensis Lane, 1952, Mycetophila (Mycetophila)	19E.15
junai Lane, 1950, Leia	19E.29
juno Lane, 1954, Synapha	19E.26
jurii Lane, 1955, Mycetophila (Abmyceta)	19E.11
jurunensis Lane, 1955, Mycetophila (Abmyceta)	19E.11
jurupari Lane, 1961, Dzedzickia	19E.24
kaestneri Edwards, 1940, Neoempheria	19E.45
kaingangi Lane, 1956, Mycetophila (Abmyceta)	19E.11
kuilcurusi Lane, 1954, Epicypa (Callicypa)	19E.6
kunasensis Lane, 1955, Mycetophila (Abmyceta)	19E.11
lacuna Freeman, 1951, Mycetophila (Abmyceta)	19E.11
lafooni Lane, 1951, Zygomya	19E.21
lalanasi Lane, 1958, Mycetophila (Abmyceta)	19E.11
lamellata Freeman, 1951, Austrosynapha	19E.22
lamellata Freeman, 1951, Mycomya	19E.41
lanei Coher, 1950, Mycomya	19E.41
lanei Duret, 1974, Echinopodium	19E.37
lanei Edwards, 1941, Cluzobra	19E.49
lanei Edwards, 1940, Neoempheria	19E.45
larifuga Coher, 1959, Neoempheria	19E.45
Lasiosoma Winnertz, 1863	19E.52
laticornis Enderlein, 1910 (Syntemna)	19E.26
lativitta Freeman, 1951, Mycetophila (Abmyceta)	19E.11
Leia Meigen, 1818	19E.26
Leiella Enderlein, 1910	19E.31
Leini	19E.26
Leja, error or emend.	19E.27
lenguasi Lane, 1954, Epicypa (Bosomyia)	19E.5
Leptomorphus Curtis, 1831	19E.50
leucocera Edwards, 1933, Leia	19E.29
levir Coher, 1959, Neoempheria	19E.45
Lia, emend.	19E.27
lindneri Edwards, 1940, Neoempheria	19E.45
liriopae Lane, 1954, Dzedzickia	19E.24
longicalcar (Enderlein), 1910 (Plastacephala), Epicypa	19E.9
longicornis Enderlein, 1910, Sceptonia	19E.20
longifurcata Freeman, 1951, Stenophragma	19E.53
longinervis Freeman, 1951, Phronia	19E.19
longiseta Coher, 1959, Neoempheria	19E.45
longistila Freeman, 1951, Aneura	19E.48
longistila Freeman, 1951, Mycomya	19E.41



<i>lorigera</i> Freeman, 1951, <i>Mycetophila</i>	19E.18
<i>luederwaldti</i> Edwards, 1940, <i>Neoempheria</i>	19E.45
<i>luederwaldti</i> (Enderlein), 1910 (<i>Plastacephala</i>), <i>Epicyptha</i> (<i>Epicyptha</i>)	19E.8
<i>lugubris</i> (Lane), 1948 (<i>Delopsis</i>), <i>Epicyptha</i> (<i>Boscmyia</i>)	19E.6
<i>lunata</i> (Lane), 1948 (<i>Delopsis</i>), <i>Epicyptha</i> (<i>Callicyptha</i>)	19E.6
<i>lutea</i> Enderlein, 1910, <i>Mycomya</i>	19E.41
<i>lutzi</i> Edwards, 1940, <i>Neoempheria</i>	19E.45
<i>macula</i> (Enderlein), 1910 (<i>Mycetophila</i>), <i>Epicyptha</i>	19E.9
<i>maculata</i> Edwards, 1933, <i>Monoclona</i>	19E.51
<i>maculipennis</i> (Enderlein), 1910 (<i>Plastacephala</i>), <i>Epicyptha</i> (<i>Epicyptha</i>)	19E.8
<i>maculipennis</i> Freeman, 1951, <i>Leia</i>	19E.30
<i>maculipennis</i> Williston, 1896, <i>Neoempheria</i>	19E.45
<i>macunai</i> Lane, 1954, <i>Epicyptha</i> (<i>Callicyptha</i>)	19E.7
<i>major</i> Freeman, 1951, <i>Trichonta</i>	19E.20
<i>malleolus</i> Freeman, 1954, <i>Leia</i>	19E.29
<i>Manota</i> Williston, 1896	19E.1
<i>Manotinae</i>	19E.1
<i>manteri</i> Coher, 1950, <i>Mycomya</i>	19E.41
<i>martinezi</i> Duret, 1972, <i>Austrosynapha</i>	19E.22
<i>mathesoni</i> Coher, 1950, <i>Mycomya</i>	19E.41
<i>mathesoni</i> Lane, 1948, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
<i>mauesensis</i> Lane, 1954, <i>Epicyptha</i> (<i>Boscmyia</i>)	19E.5
<i>maupiensis</i> Lane, 1951, <i>Epicyptha</i> (<i>Epicyptha</i>)	19E.8
<i>medea</i> Lane, 1954, <i>Dziedzickia</i>	19E.24
<i>Megalopelma</i> Enderlein, 1910	19E.50
<i>Megophthalmidia</i> Dziedzicki, 1889	19E.31
<i>merdigera</i> Knab & Zwaluwenburg, 1918, <i>Mycetophila</i>	19E.18
<i>meridionalis</i> Johannsen, 1910, <i>Mycomya</i>	19E.41
<i>mesorphina</i> Speiser, 1911, <i>Mycetophila</i>	19E.18
<i>metallica</i> Lane, 1954, <i>Dziedzickia</i>	19E.24
<i>micelbacheri</i> Lane, 1962, <i>Synapha</i>	19E.26
<i>midas</i> Coher, 1952, <i>Mycomya</i>	19E.41
<i>miltoni</i> (Lane), 1948 (<i>Delopsis</i>), <i>Epicyptha</i> (<i>Boscmyia</i>)	19E.5
<i>minor</i> Duret, 1975, <i>Echinopodium</i>	19E.37
<i>minor</i> Enderlein, 1910, <i>Tetragoneura</i>	19E.34
<i>modesta</i> Lane, 1948, <i>Zygomyia</i>	19E.21
<i>Monoclona</i> Mik, 1886	19E.50
<i>monoleuca</i> Edwards, 1933, <i>Leia</i>	19E.29
<i>morenoi</i> Lane, 1963, <i>Synapha</i>	19E.26
<i>morigenea</i> Edwards, 1940, <i>Stenophragma</i>	19E.53
<i>morosa</i> Edwards, 1933, <i>Procycloneura</i>	19E.33
<i>muchtleri</i> Curran, 1928, <i>Leia</i>	19E.29
<i>muelleri</i> Edwards, 1940, <i>Neoempheria</i>	19E.46
<i>multiplex</i> Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
<i>multiseta</i> Coher, 1959, <i>Mycomya</i>	19E.41
<i>munroei</i> Duret, 1976, <i>Echinopodium</i>	19E.37
<i>Mycetophila</i> Meigen, 1803	19E.14
<i>Mycetophilinae</i>	19E.2
<i>Mycetophilini</i>	19E.5
<i>Mycomya</i> Rondani, 1856	19E.38



<i>Mycomyia</i> , emend.	19E.38
Mycomyini	19E.35
naumanni Duret, 1973, Austrosynapha	19E.22
Nealodia Edwards, 1932	19E.4
neivai Edwards, 1940, Leptomorphus	19E.50
neivai Edwards, 1940, Neoempheria	19E.46
neivai Shaw, 1940, Neoempheria	19E.46
Neoempheria Osten Sacken, 1878	19E.43
<i>Neoglaphyoptera</i> Osten Sacken, 1878	19E.27
<i>Neopicypta</i> Coher, 1950	19E.19
neotropica Lane, 1959, Coelosia	19E.23
neotropicalis Lane, 1947, Platurocypta	19E.19
nephrodops (Enderlein), 1910 (Selkirkius), Paraleia	19E.32
nervitacta Freeman, 1951, Mycetophila (Mycetophila)	19E.15
acuquense Duret, 1975, Echinopodium	19E.37
Neuratelia Rondani, 1856	19E.51
<i>Neurocompsa</i> Enderlein, 1910	19E.43
nicothoe Coher, 1952, Mycomyia	19E.42
nigra Fisher, 1939, Dziedzickia	19E.24
nigra Freeman, 1951, Megophthalmidia	19E.32
nigrescens Duret, 1974, Echinopodium	19E.37
nigrescens Freeman, 1951, Mycetophila	19E.18
nigrescens Lane, 1954, Synapha	19E.26
nigricauda Edwards, 1933, Leia	19E.29
nigricauda Edwards, 1940, Stenophragma	19E.53
nigricoxa Freeman, 1951, Allocotocera	19E.48
nigricoxa Freeman, 1951, Echinopodium	19E.37
nigripalpis Freeman, 1951, Tetragoneura	19E.34
nigriventris Edwards, 1940, Monoclona	19E.51
nigriventris Philippi, 1865, Mycetophila	19E.18
nigrohalterata (Enderlein), 1910 (Plastacephala), Epicypta (Epicypta)	19E.8
nigrospleniata (Lynch Arribáizaga), 1892 (Glaphyoptera), Leia	19E.29
nitens Edwards, 1940, Trizygia	19E.54
nitens (Williston), 1896 (Neoglaphyoptera), Leia	19E.29
nocticolor Edwards, 1932, Tetragoneura	19E.34
nodulosa Williston, 1896, Mycetophila	19E.18
notata Freeman, 1951, Mycetophila (Abmyceta)	19E.11
Novakia Strobl, 1893	19E.51
nubilipennis Freeman, 1951, Trichonta	19E.20
nubilipennis (Walker), 1837 (Leia), Paraleia	19E.32
nublensis Lane, 1962, Mycetophila (Abmyceta)	19E.11
oblectabilis Loew, 1869, Leia	19E.29
obscuripennis Blanchard, 1852, Mycetophila	19E.18
obscuriventris Bigot, 1838 (Boletina)	19E.54
obsoleta Blanchard, 1852, Sciophila	19E.53
ochracea Freeman, 1951, Mycomyia	19E.42
ochracea Freeman, 1951, Stenophragma	19E.53
ochreocalcar Enderlein, 1910, Leiella	19E.31
ocreata Philippi, 1865, Sciophila	19E.53
oedipus Edwards, 1934, Epicypta (Epicypta)	19E.8



oiampensis Lane, 1961, Dziedzickia	19E.25
oliveirai Lane, 1959, Leia	19E.29
opaiei Lane, 1954, Epicypta (Bosomyia)	19E.6
ordiniseta Freeman, 1951, Mycetophila	19E.18
ornata Lane, 1948, Epicypta (Callicypta)	19E.7
ornatidorsum (Enderlein), 1910 (Mycetophila), Epicypta	19E.9
ornatifemora Fisher, 1939, Sceptonia	19E.20
ornatipennis Blanchard, 1852, Mycetophila	19E.18
ornatipennis (Enderlein), 1910 (Neurocompsa), Neoempheria	19E.46
ornatipennis Lane, 1948, Zygomylia	19E.21
Oromyceta Lane, 1955	19E.16
paiaiguensis Lane, 1956, Sceptonia	19E.20
pallida Freeman, 1951, Eudicrana	19E.49
pallida Johnson, 1919, Mycetophila	19E.18
palpalis Lane, 1948, Manota	19E.1
pammela Edwards, 1933, Docosia	19E.26
panamensis Coher, 1959, Neoempheria	19E.46
paraensis Lane, 1958, Rymosia	19E.4
Paraleia Tonnoir, 1929	19E.32
paranensis Edwards, 1933, Leia	19E.30
paranensis Edwards, 1932, Procycloneura	19E.33
paranensis (Lynch Arribálzaga), 1892 (Lasiosoma), Sciophlia	19E.53
paranotata Freeman, 1954, Mycetophila (Abmyceta)	19E.11
parapicalis Freeman, 1954, Mycetophila (Abmyceta)	19E.12
Paratrizygia Tonnoir, 1929	19E.51
Parvicellula Marshall, 1896	19E.52
passei Lane, 1952, Mycetophila (Mycetophila)	19E.15
patagonesi Lane, 1958, Mycetophila (Abmyceta)	19E.12
patagonicum Duret, 1976, Echinopodium	19E.37
paulensis Coher, 1959, Neoempheria	19E.46
paulensis Edwards, 1933, Leia	19E.30
paulistana Lane, 1956, Austrosynapha	19E.22
paulistensis Lane, 1948, Epicypta (Callicypta)	19E.7
paulistensis (Lanc), 1946 (Ezechia), Boraceomyia	19E.3
paupercula Coher, 1959, Mycomya	19E.42
pectinata Freeman, 1951, Mycetophila (Abmyceta)	19E.12
pectinata Freeman, 1951, Mycomya	19E.42
pectinata Freeman, 1951, Tetragoneura	19E.34
pedifera Edwards, 1933, Leia	19E.30
pellucida Freeman, 1951, Mycetophila (Abmyceta)	19E.12
penai Duret, 1976, Echinopodium	19E.37
peniculata Freeman, 1951, Mycetophila (Abmyceta)	19E.12
pereirai Edwards, 1940, Neoempheria	19E.46
peruana Kertész, 1909, Tetragoneura	19E.35
peruana Lane, 1962, Epicypta (Epicypta)	19E.8
peruviana Edwards, 1931, Mycomya	19E.42
peruviana Edwards, 1933, Paraleia	19E.32
philipsi Shaw, 1951, Neoempheria	19E.46
Phronia Winnertz, 1863	19E.19
Phthinia Winnertz, 1863	19E.52
picea Freeman, 1951, Mycetophila (Mycetophila)	19E.15
picticornis (Kertész), 1902 (Neoglaphyoptera), Leia	19E.30
pilosa Edwards, 1940, Neoempheria	19E.46



<i>pilosa</i> Edwards, 1931, <i>Rymosia</i>	19E.5
<i>pllosum</i> Duret, 1976, <i>Echinopodium</i>	19E.37
<i>pirapeis</i> Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>planiceps</i> Enderlein, 1910, <i>Megalopelma</i>	19E.50
<i>planiventris</i> (Enderlein), 1910 (<i>Plastacephala</i>), <i>Epicypa</i> (<i>Epicypa</i>)	19E.8
<i>Plastacephala</i> Enderlein, 1910	19E.5
<i>Platurocypta</i> Enderlein, 1910	19E.19
<i>platyura</i> Edwards, 1940, <i>Megalopelma</i>	19E.50
<i>plaumanni</i> Edwards, 1940, <i>Cluzobra</i>	19E.49
<i>plaumanni</i> Edwards, 1940, <i>Neoempheria</i>	19E.46
<i>plaumanni</i> Lane, 1959, <i>Leia</i>	19E.30
<i>plaumanni</i> Lane, 1958, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>plaumanni</i> Lane, 1954, <i>Megophthalmidia</i>	19E.32
<i>plaumanni</i> Lane, 1951, <i>Zygomyia</i>	19E.21
<i>Pleonazoneura</i> Enderlein, 1910	19E.43
<i>pleuralis</i> Edwards, 1934, <i>Stenophragma</i>	19E.53
<i>Pleurogymnus</i> Freeman, 1951	19E.19
<i>pluripunctata</i> (Lane), 1948 (<i>Delopsis</i>), <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>poeciloptera</i> Philippi, 1865, <i>Leia</i>	19E.30
<i>pollux</i> Freeman, 1951, <i>Tetragoneura</i>	19E.35
<i>pontis</i> Coher, 1959, <i>Mycomya</i>	19E.42
<i>popocatepetli</i> Bellardi, 1859, <i>Sciophila</i>	19E.53
<i>portoblest</i> Coher, 1959, <i>Mycomya</i>	19E.42
<i>portoricensis</i> Coher, 1959, <i>Neoempheria</i>	19E.46
<i>praecox</i> Philippi, 1865, <i>Sciophila</i>	19E.53
<i>Procycloneura</i> Edwards, 1932	19E.33
<i>producta</i> Freeman, 1951, <i>Parvicellula</i>	19E.52
<i>proseni</i> Duret, 1973, <i>Austrosynapha</i>	19E.22
<i>pubericornis</i> Edwards, 1933, <i>Dziedzickia</i>	19E.25
<i>pucaraense</i> Duret, 1974, <i>Echinopodium</i>	19E.37
<i>pucaraensis</i> Duret, 1973, <i>Mycetophila</i>	19E.18
<i>punctata</i> Bellardi, 1859, <i>Leia</i>	19E.30
<i>puncticoxa</i> Edwards, 1940, <i>Neoempheria</i>	19E.46
<i>punctiformis</i> Fisher, 1939, <i>Leia</i>	19E.28
<i>punctipennis</i> Blanchard, 1852, <i>Mycetophila</i>	19E.17
<i>punctulata</i> (Lane), 1947 (<i>Delopsis</i>), <i>Epicypa</i> (<i>Callicypa</i>)	19E.7
<i>punensis</i> Lane, 1955, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>pusilla</i> Philippi, 1865, <i>Sciophila</i>	19E.53
<i>quadripectinata</i> Edwards, 1940, <i>Parvicellula</i>	19E.52
<i>quadrisetosa</i> Freeman, 1951, <i>Austrosynapha</i>	19E.22
<i>rabelloi</i> Coher, 1959, <i>Neoempheria</i>	19E.46
<i>rabelloi</i> (Lane), 1948 (<i>Delopsis</i>), <i>Epicypa</i> (<i>Callicypa</i>)	19E.7
<i>ramosa</i> Freeman, 1951, <i>Mycetophila</i>	19E.18
<i>recta</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.13
<i>reducta</i> Freeman, 1951, <i>Austrosynapha</i>	19E.22
<i>riogrense</i> Coher, 1959, <i>Echinopodium</i>	19E.38
<i>riveti</i> (Edwards), 1919 (<i>Megophthalmidia</i>), <i>Tetragoneura</i>	19E.35
<i>Rondaniella</i> Johannsen, 1909	19E.33
<i>rossi</i> Lane, 1962, <i>Austrosynapha</i>	19E.22
<i>rostrata</i> Edwards, 1940, <i>Neoempheria</i>	19E.46
<i>rufa</i> (Lane), 1948 (<i>Delopsis</i>), <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12



<i>sinuata</i> Freeman, 1951, <i>Tetragoneura</i>	19E.35
<i>siusisi</i> Lane, 1954, <i>Epicypta</i> (<i>Callicypta</i>)	19E.7
<i>smartii</i> Edwards, 1940, <i>Neoempheria</i>	19E.47
<i>smithi</i> Edwards, 1933, <i>Leia</i>	19E.30
<i>socia</i> Coher, 1959, <i>Ncoempheria</i>	19E.47
<i>solita</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>sororcula</i> (Loew), 1869 (<i>Leia</i>), <i>Rondaniella</i>	19E.33
<i>souzai</i> Lane, 1960, <i>Austrosynapha</i>	19E.22
sp. Freeman, 1951, <i>Coelosia</i>	19E.23
sp. Freeman, 1951, <i>Paraleia</i>	19E.32
<i>spinata</i> Lane, 1954, <i>Austrosynapha</i>	19E.22
<i>spinata</i> Edwards, 1932, <i>Tetragoneura</i>	19E.35
<i>spinifera</i> Edwards, 1933, <i>Leia</i>	19E.39
<i>spinifera</i> Freeman, 1951, <i>Austrosynapha</i>	19E.22
<i>spinifera</i> Freeman, 1951, <i>Mycomya</i>	19E.42
<i>spinigera</i> Freeman, 1951, <i>Trichonta</i>	19E.20
<i>spinipes</i> Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.15
<i>spinosa</i> Edwards, 1940, <i>Neoempheria</i>	19E.47
<i>spinosa</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>spinosa</i> Freeman, 1951, <i>Mycomya</i>	19E.42
<i>spinosum</i> Duret, 1974, <i>Echinopodium</i>	19E.38
<i>spinulosa</i> Freeman, 1951, <i>Paratrizygia</i>	19E.51
<i>splendens</i> Lane, 1948, <i>Eudicrana</i>	19E.50
<i>splendida</i> Lane, 1948, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.16
<i>Staegeria</i> Wulp, 1876	19E.50
<i>Stenophragma</i> Skuse, 1888	19E.53
<i>stonei</i> Duret, 1974, <i>Echinopodium</i>	19E.38
<i>stonei</i> Lane, 1958, <i>Leia</i>	19E.30
<i>stonei</i> Lane, 1952, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.16
<i>striatum</i> Duret, 1974, <i>Echinopodium</i>	19E.38
<i>stuardoi</i> Duret, 1976, <i>Echinopodium</i>	19E.38
<i>subbrunnea</i> Freeman, 1951, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>subcapitata</i> Freeman, 1954, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.16
<i>subclavata</i> Edwards, 1940, <i>Neoempheria</i>	19E.47
<i>subcompleta</i> Edwards, 1940, <i>Aphelomera</i>	19E.49
<i>subepacra</i> Coher, 1959, <i>Mycomya</i>	19E.42
<i>subfallax</i> Coher, 1959, <i>Neoempheria</i>	19E.47
<i>subfasciata</i> Freeman, 1951, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.16
<i>subfumosa</i> Freeman, 1954, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.12
<i>subfusca</i> Freeman, 1951, <i>Mycetophila</i>	19E.18
<i>subfusca</i> Freeman, 1951, <i>Mycomya</i>	19E.42
<i>subhorrens</i> Coher, 1959, <i>Neoempheria</i>	19E.47
<i>sublevir</i> Coher, 1959, <i>Neoempheria</i>	19E.47
<i>submaculipennis</i> Freeman, 1954, <i>Leia</i>	19E.30
<i>subrecta</i> Freeman, 1954, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.13
<i>subvittata</i> Freeman, 1954, <i>Mycetophila</i> (<i>Abmyceta</i>)	19E.13
<i>Synapha</i> Meigen, 1818	19E.25
<i>Syntenna</i> , authors	19E.23
.	
<i>tacanhapesi</i> Lane, 1960, <i>Epicypta</i> (<i>Callicypta</i>)	19E.7
<i>tacuensis</i> Lane, 1951, <i>Epicypta</i> (<i>Epicypta</i>)	19E.8
<i>tacuensis</i> Lane, 1952, <i>Mycetophila</i> (<i>Mycetophila</i>)	19E.16
<i>talaris</i> Freeman, 1951, <i>Mycetophila</i>	19E.18



rufescens Edwards, 1932, Synapha	19E.26
rufithorax Freeman, 1951, Lcia	19E.30
<i>Ekymosia</i> , emend.	19E.4
Rymosia Winnertz, 1863	19E.4
saci Lane, 1961, Dziedzickia	19E.25
salobrensis Edwards, 1941, Leia	19E.29
saltanensis Lane, 1955, Mycetophila (Oromyceta)	19E.17
samesteri Coher, 1950, Mycomya	19E.42
sapaici Lanc, 1952, Neuratelia	19E.51
Sceptonia Winnertz, 1863	19E.19
schajovskoi Duret, 1973, Austrosynapha	19E.22
schildi Lane, 1956, Epicypta (Epicypta)	19E.8
schmidti Fisher, 1939, Leia	19E.30
Schnusea Edwards, 1933	19E.25
schnusei Edwards, 1933, Leia	19E.30
<i>Sciophila</i> , authors	19E.38
Sciophila Meigen, 1818	19E.52
Sciophilinae	19E.21
Sciophilini	19E.47
Selkirkiini	19E.26
<i>Selkirkius</i> Enderlein, 1940	19E.32
separata Coher, 1959, Neoempheria	19E.46
sergioi Duret, 1974, Echinopodium	19E.38
sergioi Duret, 1973, Mycetophila	19E.18
sessilis Edwards, 1933, Schnusea	19E.25
setifera Freeman, 1951, Mycomya	19E.42
setifera Freeman, 1951, Paratrizygia	19E.51
setigera (Freeman), 1951 (Exechia), Exechiopsis (Xencxechia)	19E.3
setosicauda Edwards, 1933, Leia	19E.30
shamanensis Lane, 1952, Mycetophila (Mycetophila)	19E.15
shannoni Coher, 1950, Mycomya	19E.42
shannoni Edwards, 1940, Neoempheria	19E.46
shannoni Lane, 1956, Aphrastomyia	19E.26
shannoni Lane, 1954, Leiella	19E.31
shannoni Lane, 1948, Mycetophila (Mycetophila)	19E.15
shannoni, Lane, 1956, Parvicellula	19E.52
shawi Coher, 1950, Mycomya	19E.42
simile (Freeman), 1951 (Allodia), Brevicornu	19E.4
similis Freeman, 1951, Eudicrana	19E.50
similis Freeman, 1951, Procycloneura	19E.33
<i>similis</i> Freeman, 1951, Tetrgoneura	19E.35
similis Freeman, 1951, Trichonta	19E.20
simillima Freeman, 1951, Mycomya	19E.42
simillima Freeman, 1954, Tetrgoneura	19E.35
simplicis Coher, 1959, Mycomya	19E.42
simplex Edwards, 1940, Neoempheria	19E.47
simplex Edwards, 1932, Teragoneura	19E.35
simplex Freeman, 1951, Mycetophila	19E.18
simplicipes Freeman, 1951, Tetrgoneura	19E.35
simplicistila Freeman, 1951, Mycetophila	19E.18
simplicistila Freeman, 1951, Tetrgoneura	19E.35
sinuata Freeman, 1951, Mycetophila (Abmyceta)	19E.12



tantalos Coher, 1959, Mycomya	19E.42
tantilla (Loew), 1869 (Sciophila), Mycomya	19E.43
tapicarei Lane, 1952, Monoclona	19E.51
tapinirai Lane, 1955, Mycetophila (Abmyceta)	19E.13
tapuiari Lane, 1958, Mycetophila (Abmyceta)	19E.13
tapuiari Lane, 1951, Zygomysia	19E.21
taunayi (Lane), 1948 (Mycetophila), Epicypta (Callicypta)	19E.6
taurus Freeman, 1951, Mycomya	19E.43
Tetragoneura Winnertz, 1846	19E.33
Tetragoneurini	19E.26
tetraphaea Shaw, 1940, Neoempheria	19E.47
theobaldi Coher, 1959, Mycomya	19E.43
theresae Edwards, 1932, Mycetophila (Mycetophila)	19E.16
thoracica Philippi, 1865, Sciophila	19E.53
Thoracotropis Freeman, 1951	19E.35
tibialis Freeman, 1951, Tetragoneura	19E.35
tobasi Lane, 1958, Mycetophila (Mycetophila)	19E.16
travassosi (Lane), 1948 (Delopsis), Mycetophila (Abmyceta)	19E.13
traveriae, emend.	19E.43
traveri Coher, 1950, Mycomya	19E.43
triangulifera Freeman, 1951, Mycetophila (Mycetophila)	19E.16
Trichonta Winnertz, 1863	19E.20
trichops Freeman, 1951, Mycomya	19E.43
tricinctiventris Enderlein, 1910, Leia	19E.29
trifasciata Edwards, 1940, Monoclona	19E.51
trinidadensis Lane, 1960, Epicypta (Callicypta)	19E.7
trioridinata Freeman, 1954, Mycetophila (Abmyceta)	19E.13
triseriata Freeman, 1951, Mycetophila (Abmyceta)	19E.13
tristis Bigot, 1888, Sciophila	19E.53
tristis (Lane), 1948 (Delopsis), Epicypta (Bosemyia)	19E.6
trivittata Freeman, 1951, Mycetophila (Abmyceta)	19E.13
Trizygia Skuse, 1888	19E.54
tropicalis Fisher, 1939, Leia	19E.27
truncata Freeman, 1951, Austrosynapha	19E.22
truncata (Freeman), 1951 (Exechia), Exechiopsis (Xenexechia)	19E.3
truncatovenosa Enderlein, 1910, Leia	19E.30
tucumana Lane, 1958, Mycetophila (Abmyceta)	19E.13
tucumensis Lane, 1955, Mycetophila (Oromyceta)	19E.17
tupanensis Lane, 1961, Mycetophila (Abmyceta)	19E.13
uacupisi Lane, 1958, Mycetophila (Abmyceta)	19E.13
uaianai Lane, 1952, Mycetophila (Mycetophila)	19E.16
uaicensis Lane, 1955, Mycetophila (Abmyceta)	19E.13
uboyasi Lane, 1955, Mycetophila (Abmyceta)	19E.13
unguiculata Freeman, 1951, Austrosynapha	19E.23
unicincta Edwards, 1933, Leiella	19E.31
unicornuta Freeman, 1951, Mycetophila	19E.18
uniseriata Freeman, 1951, Mycetophila	19E.18
unispinosa Edwards, 1940, Neoempheria	19E.47
valdiviana (Philippi), 1865 (Sciophila), Mycomya	19E.43
variabilis Lane, 1954, Dziedzickia	19E.25
varipennis, lapsus	19E.47



<i>varipennis</i> (Lynch Arribálzaga), 1892 (Empheria), Neoempheria	19E.47
<i>vaurasi</i> Lane, 1958, Mycetophila (Mycetophila)	19E.16
<i>veligera</i> Freeman, 1951, Mycetophila	19E.18
<i>verberifera</i> Freeman, 1951, Mycetophila (Abmyceta)	19E.13
<i>verbifera</i> , lapsus	19E.13
<i>vernalis</i> Philippi, 1865, Sciophila	19E.53
<i>villaricaense</i> Duret, 1976, Echinopodium	19E.38
<i>viridipes</i> Freeman, 1951, Mycetophila	19E.19
<i>vittata</i> Freeman, 1951, Acnemia	19E.48
<i>vittata</i> Edwards, 1931, Eudicrana	19E.50
<i>vitticollis</i> Blanchard, 1852, Mycetophila	19E.19
<i>vittidorsum</i> (Enderlein), 1910 (Acnemia), Monoclona	19E.51
<i>vittipennis</i> Freeman, 1951, Mycetophila	19E.19
<i>vockerothi</i> Duret, 1976, Echinopodium	19E.38
<i>vogeli</i> Edwards, 1940, Neoempheria	19E.47
<i>vogeli</i> Edwards, 1932, Tetragoneura	19E.35
<i>volitans</i> Lynch Arribálbaga, 1892, Mycetophila	19E.19
<i>williamsi</i> Lane, 1960, Epicypta (Boscmyia)	19E.6
<i>winnertzi</i> Lane, 1948, Mycetophila (Abmyceta)	19E.13
<i>worontzowi</i> Lane, 1947, Rymosia	19E.5
<i>wygodzinskyi</i> (Lane), 1947 (Delopsis), Mycetophila (Oromyceta)	19E.17
<i>xamasensis</i> Lane, 1955, Mycetophila (Abmceta)	19E.14
<i>Xenexechia</i> Tuomikoski, 1966	19E.3
<i>yivoffi</i> Duret, 1973, Austrosynapha	19.23
<i>yurimauesi</i> Lane, 1955, Mycetophila (Abmyceta)	19E.14
<i>zeteki</i> Coher, 1959, Neoempheria	19E.47
<i>zonalis</i> Edwards, 1931, Leiella	19E.31
<i>Zygomya</i> Winnertz, 1863	19E.20



Composto e impresso em 1978, na oficina da
EMPRESA GRÁFICA DA REVISTA DOS TRIBUNAIS S/A
Rua Conde de Sarzedas, 38 — Tel. 36-6958 (PBX)
01512 São Paulo, SP, Brasil.





DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

21

FAMILY HYPEROSCELIDIDAE
(CORINOSCELIDIDAE)

EDWIN F. COOK

*Professor**Department of Entomology, Fisheries and Wildlife
University of Minnesota
St. Paul, Minnesota 55101*

This very small family of flies has until fairly recently been included with the Scatopsidae. The family is composed of three species from the northern hemisphere, four species from New Zealand, and four species from the west coast of South America. The larva and pupa of one species from New Zealand have been described, but nothing of the biology of the American species.

Refs. — Hardy & Nagatomi, 1960:263; Tollet, 1959:132; Tonnoir, 1927: 291 (larval disc).

Subfamily HYPEROSCELIDINAE

Genus CANTHYLOSCELIS Edwards

CANTHYLOSCELIS Edwards, 1922:267. Type-species, *antennata* Edwards (orig. dcs.).

Subgenus ARAUCOSCELIS Edwards

Canthyloscelis, subg. ARAUCOSCELIS Edwards, 1930:90. Type-species, *pectinata* Edwards (orig. dcs.).

PECTINATA Edwards, 1930:90, fig. 4 [the legend reads "*antennata* sp. n. (lapsus); should be corrected to *pectinata* (cf. Edwards, 1934: 185)], pl. 9, figs. 1-2. Type-locality: Argentina, Rio Negro, Puerto Blest. Distr. — Chile (Llanquihue), Argentina (Rio Negro).

Subgenus CHILISCELIS Tollet

Canthyloscelis, subg. CHILISCELIS Tollet, 1959:146. Type-species, *valdiviana* Tollet (orig. des.).

Paper n.º 6105, Scientific Journal Series, Minnesota Agric. Experiment Station, St. Paul, Minnesota 55101.

21.1

MUSEU DE ZOOLOGIA - USP

SciELO

- APICATA Edwards, 1934:186. Type-locality: Chile, Comudes. Distr. — Chile.
- PICUPENNIS Edwards, 1930:92 (*Araucoscelis*). Type-locality: Argentina, Rio Negro, Laguna Frias, on road towards Rosales Pass. Distr. — Argentina (Rio Negro).
- VALDIVIANA Tollet, 1959:147, figs. 1-6. Type-locality: Chile, Valdivia, Shoshucno. Distr. — Chile (Valdivia, Araueo, Llanquihue, Chiloe I.).

LITERATURE CITED

EDWARDS F. W.

1922. Preliminary note on a new genus of Scatopsid flies from New Zealand. *Ann. & Mag. Nat. Hist.* (9)9:267-269.
1930. Bibionidae, Scatopsidae, Ceidomyiidae, Culicidae, Thaumalidae (Orphnophilidae), Anisopodidae (Rhyphidae), in *Diptera of Patagonia and South Chile* 2(3):77-119, pls. 9-11.
1934. Four new Diptera in the collection of the Berlin Zoological Museum. *Stylops*, London 3:185-187.

HARDY D. E. & A. NAGATOMI


- 1960. An unusual new Nematocera from Japan (Diptera) and a new family name. *Pacific Insects* 2(3):263-267, figs. 1a-j.

TOLLET, R.

1959. Note systématique sur les Corynoseelidae fam. nov. (Diptera) du globe et description d'un Corynoscelidae nouveau de l'hémisphère austral. *Bull. Ann. Soc. Roy. Ent. Belg.* 95:132-153, 6 figs.

TONNOIR, A. L.

1927. An acephalous larva of a nematocerous dipteran: the early stages of *Canthyloscelis*. *Ann. & Mag. Nat. Hist.* (9)19:291-300.



DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

22

FAMILY SCATOPSIDAE

- EDWIN F. COOK

*Professor**Department of Entomology, Fisheries, and Wildlife
University of Minnesota
St. Paul, Minnesota 55101*

This small family of small, usually dark colored flies is very poorly known in the Neotropical region. As is true for this group in the rest of the world, there is scarcely anything known about the biology of species of this family in this great region. Exclusively Neotropical species have no immature stages described. There have been no revisional studies of Neotropical species and many unrecognized species await description.

Subfamily SCATOPSINAE

Tribe SCATOPSINI

Genus SCATOPSE Geoffroy

SCATOPSE Geoffroy, 1762:450. Type-species, *Tipula notata* Linnaeus (I.C.Z.N., 1957:86). Conserved as of 1762 under suspension of the Rules (I.C.Z.N., 1957:86)

Reichertella Enderlein, 1912:268. Type-species, *Scatopse femoralis* Meigen (orig. des.; misident.) = *nigra* Meigen.


Rhaeboza Enderlein, 1936:55. Type-species, *Scatopse fuscipes* Meigen (orig. des.).

Masatierra Enderlein, 1940:666. Type-species, *ferruginea* Enderlein (orig. des.) = *fuscipes* Meigen.

Ref. — Cook, 1957 (rev.).

Paper n.º 6105, Scientific Journal Series, Minnesota Agricultural Experiment Station, St. Paul, Minnesota 55101.

22.1



- COSTARICANA* Duda, 1928:286, text-figs. 17, 18 (male terminalia, wing). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Mexico (Vera Cruz, Córdoba), Costa Rica.
- FUSCIPES* Melgen, 1830:314. Type-locality: "Europe". Distr. — Cosmopolitan. Ref. — Cook, 1957:608, figs. 3a, b, d, n, 5d, f, 6b, g (male genitalia, spiracular sclerite, female 7th sternum, female genitalia, male tergum and sternum 7, wing, tergum 8 of male).
- parvula* Blanchard, 1852:359. Type-locality: Chile, Coquimbo.
- peruana* Enderlein, 1912:275, fig. 7 (wing) (*Reichertella*). Type-locality: Peru, Mollendo.
- ferruginea* Enderlein, 1940:666 (*Masatierra*). Type-locality: Juan Fernandez Is., Masatierra.
- coutinhoi* Rabello & Forattini, 1962:303, figs. 1-4 (wing, haltere, male and female genitalia, larva, and pupa) (*Rhcxoza*). Type-locality: Brazil, São Paulo, Salesópolis.
- MARITIMA* Duda, 1928:278, text-figs. 9, 10 (leg. part of wing). Type-locality: Mexico, Vera Cruz. Distr. — Mexico (Vera Cruz).
- NOTATA* (Linnaeus), 1758:588 (*Tipula*). Type-locality: "Europa". Distr. — Cosmopolitan. Refs. — Lyall, 1929:630 (larval disc); Cook, 1957:598, figs. 1a, 2g, 3c, 4a, 5c, 6e (male genitalia, male metathoracic tarsi, spiracular sclerite, female genitalia, male 7th tergum and sternum, wing).
- carolina* Blanchard, 1852:359. Type-locality: Chile, Chilloc I., Ancud [= San Carlos]. Ref. — Edwards, 1930:94.
- carbonaria* Philippi, 1865:640. Type-locality: Chile, Santiago, Corral. Ref. — Edwards, 1930:94.
- TUBIFERA* Edwards, 1930:95, pl. 1X, figs. 6, 8 (wing, male genitalia). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile (Llanquihue).

Genus *COLOBOSTEMA* Enderlein

- COLOBOSTEMA* Enderlein, 1926:140. Type-species, *oldenbergi* Enderlein (orig. des.) = *tristis* (Zetterstedt).
Ref. — Cook, 1956 (rev.).
- ROTUNDUM* Cook, 1956:332, figs. 1f, 2e (male genitalia, male 7th tergum and sternum). Type-locality: Panama, Tocumen. Distr. — Panama, Costa Rica (San Mateo).

Genus *HOLOPLAGIA* Enderlein

- HOLOPLAGIA* Enderlein, 1912:267. Type-species, *Scatopse transversalis* Loew (orig. des.).
- BRUNI* Spelser, 1921:82. Type-locality: Brazil, Santa Catarina, Blumenau. Distr. — Brazil (Santa Catarina).
- GUAMENSIS* (Johannsen), 1946:187, figs. 1a-e [wing (the wing figured is that of *Scatopse fuscipes* and not that of *H. guamensis*), male genitalia, female genitalia] (*Scatopse*). Type-locality: "Guam". Distr. — Hawaii, Truk, Ghana, Accra; Panama (Canal Zone, Colón, Summit). Ref. — Cook, 1957:610, figs. 7c, e, j, k (male and female genitalia, male 7th tergum, wing).

Tribe RHEGMOCLEMATINI

Genus RHEGMOCLEMA Enderlein

- RHEGMOCLEMA Enderlein, 1912:276. Type-species, *rufithorax* Enderlein (orig. des.).
- Aldrovandiella* Enderlein, 1912:278. Type-species, *Scatopse halterata* Meigen (orig. des.).
Ref. — Cook, 1955 (rev.).
- BLANTONI Cook, 1955:363, pl. IV, i, j, k (male tergum and sternum 7, male genitalia, wing). Type-locality: Panama, Bocas del Toro Prov., Almirante. Distr. — Panama (Bocas del Toro).
- CAUDATUM (Duda), 1928:289, text-fig. 20 (male terminalia), pl. IV, fig. 5 (wing) (*Aldrovandiella*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).
- SETIFERA (Edwards), 1930:95, pl. IX, figs. 3, 4 (wing, male genitalia) (*Scatopse*). Type-locality: Argentina, Rio Negro, L. Correntoso. Distr. — Argentina (Rio Negro).

Genus RHEGMOCLEMINA Enderlein

- RHEGMOCLEMINA Enderlein, 1936:55. Type-species, *Scatopse vaginata* Lundström (orig. des.).
Ref. — Cook, 1955 (rev.).

Subgenus RHEGMOCLEMINA Enderlein

- CLAVICRUS (Duda), 1928:282, pl. IV, fig. 4 (wing) (*Scatopse*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba). Ref. — Cook, 1955:358, pl. Ie, IIe (male genitalia, male tergum and sternum 7).
- DIABOLICA (Duda), 1928:285, text-figs. 15, 16 (male terminalia, wing) (*Scatopse*). Type-locality: Mexico, Huizquilcan. Distr. — Mexico (Huizquilcan, Chapingo).

Subgenus NEORHEGMOCLEMINA Cook

- RhegmoCLEMINA*, subg. NEORHEGMOCLEMINA Cook, 1955:358. Type-species, *parvum* Cook (orig. des.).
- ABRUPTINERVIS (Duda), 1928:279, pl. IV, fig. 2 (wing) (*Scatopse*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica, Panama. Ref. — Cook, 1955:360, pls. Id, IIh (male genitalia, male tergum and sternum 7).
- ASYMMETRICUM Cook, 1955:360, pls. Ig, IIg, IIIe (male genitalia, male tergum and sternum 7, wing). Type-locality: Jamaica, Ocho Rios. Distr. — Jamaica (Ocho Rios).
- CONTRACTA (Edwards), 1930:96, pl. IX, figs. 5, 7 (male genitalia, wing) (*Scatopse*). Type-locality: Chile, Llanquihue, Peulla. Distr. — Chile (Llanquihue: Peulla, Casa Pangue).
- TRICHIONEURA (Duda), 1928:288, text-fig. 19 (wing) (*Scatopse*). Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).



?WILLISTONI (McAtee), 1921:124 (*Scatopse*; nom. nov. for *pygmaea* Williston). Type-locality: British West Indies, St. Vincent. Distr. — British West Indies (St. Vincent).

pygmaea Williston, 1896:269 (*Scatopse*; preocc. Loew, 1864). Type-locality: British West Indies, St. Vincent.

Tribe SWAMMERDAMELLINI

Genus SWAMMERDAMELLA Enderlein

SWAMMERDAMELLA Enderlein, 1912:277. Type-species, *Scatopse brevicornis* Meigen (orig. des.).

Ref. — Cook, 1956 (rev.).

APPROXIMATA Edwards, 1924:571. Type-locality: Trinidad, St. Augustine. Distr. — Trinidad (St. Augustine).

MARGINATA Cook, 1956:24, figs. 1d, 2d, 3g, 4e, 5h (male genitalia, male 7th segment, wing, male 6th segment, female genitalia). Type-locality: U.S.A., New Mexico, Las Vegas. Distr. — U.S.A. (New Mexico, Nevada, California), Mexico (Guadalajara).

MOJINGAE Cook, 1956:29, figs. 5j, k (male 7th segment, male genitalia). Type-locality: Panama, Canal Zone, Mojinga Swamp, Fort Sherman. Distr. — Panama (Canal Zone).

Genus RHEXOZA Enderlein

RHEXOZA Enderlein, 1936:55. Type-species, *zacheri* Enderlein (orig. des.) = *subnitens* (Verrall).

Ref. — Cook, 1956 (rev.).

AURICULARIS (Duda), 1928:283, text-figs. 12, 13, 14 (wing, male genitalia) (*Scatopse*). Type-locality: Mexico, Chapultepec. Distr. — Mexico (Chapultepec).

DAMPFI (Duda), 1928:280, text-fig. 11, pl. IV, fig. 3 (male genitalia, wing) (*Scatopse*). Type-locality: Mexico, Popotea. Distr. — U.S.A. (Texas), Mexico. Ref. — Cook, 1956:10, figs. 5b, c, e (male genitalia, male 7th and 8th terga and sterna, female genitalia).

LOBATA Cook, 1956:8, figs. 1g, 2f (male genitalia, male segment 7). Type-locality: Mexico, Córdoba, Vera Cruz. Distr. — Mexico (Córdoba, Vera Cruz).

PANAMENSIS Cook, 1956:8, figs. 1c, 2c, 3c (male genitalia, male segment 7, female genitalia). Type-locality: Panama, Canal Zone, Ancon. Distr. — Panama (Canal Zone).

Subfamily PSECTROSCIARINAE

Genus PSECTROSCIARA Kieffer

PSECTROSCIARA Kieffer in Enderlein, 1911:192 (also Kieffer, 1912:192). Type-species, *mahensis* Kieffer (orig. des.) = *brunnipes* (Brunetti).

Ref. — Cook, 1958 (rev.).



- CALIFORNICA (Colc), 1912:150, fig. 85 (adult) (*Scatopse*). Type-locality: U.S.A., California, Laguna Beach. Distr. — U.S.A. (Arizona, California), Mexico (Córdoba). Ref. — Cook, 1958:593, figs. 1f, g, 3a, b, c, 5a (meso- and metathoracic 1st tarsi, male genitalia, female genitalia).
- mexicana* Duda, 1928:294, pl., fig. 25 [male genitalia (misidentified as female genitalia)] (as *scatopsiformis* subsp.). Type-locality: Mexico, Chapingo. Ref. — Cook, 1958:593.
- JAMAICENSIS Cook, 1958:595, figs. 4g, h (female genitalia, spermatheca). Type-locality: Jamaica, Ocho Rios. Distr. — Jamaica (Ocho Rios).
- ROSSI Cook, 1958:595, figs. 3d, e (male genitalia). Type-locality: Chile, Coquimbo, Laguna Dam. Distr. — Chile (Coquimbo).
- SCATOPSIFORMIS Enderlein, 1912:281, fig. 14 (wing). Type-locality: "Costa Rica". Distr. — Costa Rica (La Suiza de Turrialba, La Caja, San José), Panama (Espino, Garachine, Tocumen, Chorrera, Pt. Mensabe, Los Santos Prov.). Ref. — Cook, 1958:593, figs. 2b, c, 5e, f, g (male genitalia, female genitalia, spermatheca, thorax).
- SERRATA Cook, 1958:594, figs. 5b, d (spermatheca, female genitalia). Type-locality: U.S.A., Texas, Los Fresnos. Distr. — U.S.A. (Texas), Mexico.

Genus ANAPAUSIS Enderlein

- ANAPAUSIS Enderlein, 1912:278. Type-species, *Scatopse soluta* Loew (orig. des.).
- Ref. — Cook, 1965 (rev.).
- FUSCINERVIS (Edwards), 1930:97, pl. X, figs. 11, 12 (male genitalia, wing) (*Psectrosciara*). Type-locality: Chile, Llanquihue, Casa Pangue. Ref. — Cook, 1965:16, figs. 42, 43, 45 (male terminalia, wing).

Subfamily ECTAETIINAE

Genus ECTAETIA Enderlein

- ECTAETIA Enderlein, 1912:279. Type-species, *Scatopse clavipes* Loew (orig. des.).
- CORNUTA Edwards, 1930:98, pl. X, figs. 9, 10 (wing, male genitalia). Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile (Llanquihue).
- LASIOPA Duda, 1928:294. Type-locality: Costa Rica, La Suiza de Turrialba. Distr. — Costa Rica (La Suiza de Turrialba).

UNRECOGNIZED AND UNPLACED SPECIES

- FULVITARSIS Macquart of Lynch Arribálzaga, 1878:298 (*Scatopse*). Locality: Argentina, Buenos Aires, Baradero. Inadequately described for recognition.
- HYALINATA Philippi, 1865:640 (*Scatopse*). Type-locality: "Chile". Unrecognizable.
- PALLIDIPES Lynch Arribálzaga, 1878:298 (*Scatopse*). Type-locality: Argentina, Buenos Aires, Baradero. Unrecognizable.
- TRANSATLANTICA Philippi, 1865:640 (*Scatopse*). Type-locality: Chile, Santiago. Unrecognizable — may be an *Ectaetia*.



LITERATURE CITED

BLANCHARD, E.

1852. Orden IX. Dipteros, pp. 327-468 in C. Gay, ed.: *Historia Fisica y Politica de Chile, Zoologia* 7:471 pp. Paris.

COLE, F. R.

1912. Some Diptera of Laguna Beach. *Ann. Rpt. Pomona Col. Laguna Mar. Lab.* 1:150-162, 10 figs.

COOK, E. F.

1955. A contribution toward a monograph of the family Scatopsidae (Diptera). Part I. A revision of the genus *Rhegmoclema* Enderlein (= *Aldrovandiella* Enderlein) with particular reference to the North American species; Part II. The genera *Rhegmoclemina* Enderlein, *Parascatopse*, n. g., and a new species of *Rhegmoclema*. *Ann. Ent. Soc. Amer.* 48:240-251, figs. 2-3, pls. 1-5; 351-364, pls. 1-4.
1956. A contribution toward a monograph of the Scatopsidae (Diptera). Part III. The genus *Rhexoxa* Enderlein; Part IV. The genus *Swammerdamella* Enderlein; Part V. The genus *Colobostema* Enderlein. *Ann. Ent. Soc. Amer.* 49:1-12, figs. 1-5; 15-29, figs. 1-6; 325-332, figs. 1-3.
1957. A contribution toward a monograph of the Scatopsidae (Diptera). Part VI. The genus *Scatopse* Geoffroy and *Holoplagia* Enderlein. *Ann. Ent. Soc. Amer.* (1956) 49:593-611, 7 pls.
1958. A contribution toward a monograph of the Scatopsidae (Diptera). Part VII. The genus *Psectrosciara* Kieffer. *Ann. Ent. Soc. Amer.* 51:587-595, 5 figs.
1965. A contribution toward a monograph of the Scatopsidae (Diptera). Part IX. The genera *Aspistes* and *Arthria* (Diptera: Scatopsidae: Aspistinae). *Ann. Ent. Soc. Amer.* 58:713-721, figs. 1-36.

DUDA, O.

1928. Beitrag zur Kenntniss der aussercuropäischen Scatopsiden (Dipt.). *Konowia* 7:259-297, 25 figs., 1 pl.

EDWARDS, F. W.

1924. New species of nematocerous Diptera from Fiji and Trinidad. *Ann. & Mag. Nat. Hist.* (9)14:571.
1930. *Diptera of Patagonia and South Chile* 2(3):88-98, pls. IX-X.

ENDERLEIN, G.

1911. Die phyletischen Beziehungen der Lycoriiden (Sclariden) zu den Fungivoriden (Mycetophiliden) und Itonididen (Cecidomyiiden) und ihre systematische Gliederung. *Arch. f. Naturgesch.* 77 (Heft 1, suppl. 3): 116-201, 2 figs., 2 pls.
1912. Zur Kenntniss der Zygophthalmen. Über die Gruppierung der Sclariden und Scatopsiden. *Zool. Anz.* 40:261-282, 15 figs.
1926. Zur Kenntniss der Scatopsiden. *Zool. Anz.* 68:137-142.



1936. 22. Ordnung: Zweiflügler, Diptera. Abt. 16, 259 pp., 317 figs. (= Lfg. 2, part), in P. Bröhmer, P. Ehrmann & G. Ulmer, eds.: *Die Tierwelt Mitteleuropas 6: Insekten, III Teil*. Leipzig.
1940. In C. J. F. Skottsberg: *Natural History of Juan Fernandez* 3:666.

GEOFFROY, E. L.

1762. *Histoire abrégée des insectes qui se trouvent aux environs de Paris* 2:690 pp., 11 pls. Paris.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE

1957. Opinion 441. Validation under the Plenary Powers for five genera in the Order Diptera (Class Insecta) published in 1762 by Geoffroy (E.L.) in the work entitled "Histoire abrégée des Insectes qui se trouvent aux environs de Paris" ("Opinion" supplementary to "Opinion" 228), pp. 83-120 (= pt. 6), in F. Hemming, ed.: *Opinions and declarations rendered by the International Commission on Zoological Nomenclature* 15:530 pp. London.

JOHANNSEN, O. A.

1946. Insects of Guam II. *Bull. Bishop Museum* 189:187.

KIEFFER, J. J.

1912. Beschreibung neuer Sciariden von den Seychellen-Inseln (Anhang), pp. 192-194, pl. 9, figs. 11-14, in G. Enderlein: The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner, M. A. Vol. IV. N^o XIII. Diptera, Sciaridae. *Trans. Linn. Soc. London* (2) *Zool.* 15:181-194, pl. 9.

LINNAEUS, C.

1758. *Systema Naturae per regna tria naturae*. Ed. 10. 1:824 pp. Holmiae (= Stockholm).

LYNCH ARRIBÁZAGA, F.

1878. Notas dipterológicas sobre los Bibionites del Baradero. *El Naturalista Argentino* 1(10):295-299.

MCATEE, W. L.

1921. District of Columbia Diptera: Scatopsidae. *Proc. Ent. Soc. Wash.* 23:120-124.

MEIGEN, J. W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten* 6: iv + 401 pp., pls. 55-66. Hamm.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. K. K. zool.-bot. Ges. Wien* 15 (Abh.): 595-782, 7 pls.



RABELLO, E. N. & O. P. FORATTINI

1962. Sobre uma nova espécie do gênero *Rhexoza* Enderlein. *Papéis Avulsos Dep. Zool. São Paulo* 15:303-309, figs. 1-4.

SPEISER, P.


1921. Ueber einige Scatopsiden (Dipt.). *Schr. physik. Ges. Königsberg* 61-62:82-84.

WILLISTON, S. W.

1896. On the Diptera of St. Vincent (West Indies). *Trans. Ent. Soc. London* 1896:253-446, pls. 8-14.

INDEX

<i>abruptinervis</i> (Duda), 1928 (Scatopse), <i>Rhegmoclemina</i>	22.3
<i>Aldrovandiella</i> Enderlein, 1912	22.3
<i>Anapausis</i> Enderlein, 1912	22.5
<i>approximata</i> Edwards, 1924, <i>Swammerdamella</i>	22.4
<i>asymmetricum</i> Cook, 1955, <i>Rhegmoclemina</i>	22.3
<i>auricularis</i> (Duda), 1928 (Scatopse), <i>Rhexoza</i>	22.4
<i>blantoni</i> Cook, 1955, <i>Rhegmoclema</i>	22.3
<i>brauni</i> Speiser, 1921, <i>Holoplagia</i>	22.2
<i>californica</i> (Cole), 1912 (Scatopse), <i>Psectrosciara</i>	22.5
<i>carbonaria</i> Philippi, 1865, Scatopse	22.2
<i>carolina</i> Blanchard, 1852, Scatopse	22.2
<i>caudatum</i> (Duda), 1928 (<i>Aldrovandiella</i>), <i>Rhegmoclema</i>	22.3
<i>clavicus</i> (Duda), 1928 (Scatopse), <i>Rhegmoclemina</i>	22.3
<i>Colobostema</i> Enderlein, 1926	22.2
<i>constricta</i> (Edwards), 1930 (Scatopse), <i>Rhegmoclemina</i>	22.3
<i>cornuta</i> Edwards, 1930, <i>Ectaetia</i>	22.5
<i>costaricana</i> Duda, 1928, Scatopse	22.2
<i>couthoi</i> (Rabello & Forattini), 1962 (<i>Rhexoza</i>), Scatopse	22.2
<i>dampfi</i> (Duda), 1928 (Scatopse), <i>Rhexoza</i>	22.4
<i>diabolica</i> (Duda), 1928 (Scatopse), <i>Rhegmoclemina</i>	22.3
<i>Ectaetia</i> Enderlein, 1912	22.5
<i>Ectaetiinae</i>	22.5
<i>ferruginea</i> (Enderlein), 1940 (<i>Masatierra</i>), Scatopse	22.2
<i>fulvitaris</i> Macquart of Lynch Arribálzaga, 1878, Scatopse	22.5
<i>fuscinervis</i> (Edwards), 1930 (<i>Psectrosciara</i>), <i>Anapausis</i>	22.5
<i>fuscipes</i> Meigen, 1830, Scatopse	22.2
<i>guamensis</i> (Johannsen), 1946 (Scatopse), <i>Holoplagia</i>	22.2
<i>Holoplagia</i> Enderlein, 1912	22.2
<i>hyalinata</i> Philippi, 1865, Scatopse	22.5



<i>jamaicensis</i> Cook, 1958, <i>Psectroscliara</i>	22.5
<i>lasiopa</i> Duda, 1928, <i>Ectactia</i>	22.5
<i>lobata</i> Cook, 1956, <i>Rhexoza</i>	22.4
<i>marginata</i> Cook, 1956, <i>Swammerdamella</i>	22.4
<i>maritima</i> Duda, 1928, <i>Scatopse</i>	22.2
<i>Masatierra</i> Enderlein, 1940	22.1
<i>mexicana</i> Duda, 1928, <i>Psectroscliara</i>	22.5
<i>mojingae</i> Cook, 1956, <i>Swammerdamella</i>	22.4
<i>Neorhegmoclemina</i> Cook, 1955	22.3
<i>notata</i> (Linnaeus), 1758 (<i>Tipula</i>), <i>Scatopse</i>	22.2
<i>pallidipes</i> Lynch Arribálzaga, 1878, <i>Scatopse</i>	22.5
<i>panamensis</i> Cook, 1956, <i>Rhexoza</i>	22.4
<i>parvula</i> Blanchard, 1852, <i>Scatopse</i>	22.2
<i>peruana</i> (Enderlein), 1912 (<i>Reichertella</i>), <i>Scatopse</i>	22.2
<i>Psectroscliara</i> Kieffer, 1911	22.4
<i>Psectroscliarinae</i>	22.4
<i>pygmaea</i> (Williston), 1896 (<i>Scatopse</i>), <i>Rhegmoclemina</i>	22.4
<i>Reichertella</i> Enderlein, 1912	22.1
<i>Rhaeboza</i> Enderlein, 1936	22.1
<i>Rhegmoclema</i> Enderlein, 1912	22.3
<i>Rhegmoclematini</i>	22.3
<i>Rhegmoclemina</i> Enderlein, 1936	22.3
<i>Rhexoza</i> Enderlein, 1936	22.4
<i>rossi</i> Cook, 1958, <i>Psectroscliara</i>	22.5
<i>rotundum</i> Cook, 1956, <i>Colobostema</i>	22.2
<i>Scatopse</i> Geoffroy, 1762	22.1
<i>scatopsiformis</i> Enderlein, 1912, <i>Psectroscliara</i>	22.5
<i>Scatopsinae</i>	22.1
<i>Scatopsini</i>	22.1
<i>serrata</i> Cook, 1958, <i>Psectroscliara</i>	22.5
<i>setifera</i> (Edwards), 1930 (<i>Scatopse</i>), <i>Rhegmoclema</i>	22.3
<i>Swammerdamella</i> Enderlein, 1912	22.4
<i>Swammerdamellini</i>	22.4
<i>transatlantica</i> Philippi, 1865, <i>Scatopse</i>	22.5
<i>trichioneura</i> (Duda), 1928 (<i>Scatopse</i>), <i>Rhegmoclemina</i>	22.3
<i>tubifera</i> Edwards, 1930, <i>Scatopse</i>	22.2
? <i>willistoni</i> (McAtee), 1921 (<i>Scatopse</i>), <i>Rhegmoclemina</i>	22.4





DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

23

FAMILY CECIDOMYIIDAE

RAYMOND J. GAGNÉ

*Entomology Research Division, Agr. Res. Serv.
U. S. Department of Agriculture
Washington, D. C.*

As with most families of Diptera in the Neotropics, much work remains to be done with the Cecidomyiidae. Only 360 species are known, belonging to 129 genera in three subfamilies. Of these subfamilies, the two smaller ones, the Lestremiinae and the Porricondylinae, are especially poorly known. The larvae of these groups live in decaying organic material. The larvae of most species of the large subfamily Cecidomyiinae are found in living plants where the majority cause conspicuous deformities. The larvae of a few genera are predaceous on Acarina or Insecta, or are mycophagous.

Fortunately, most of the species have been reared from known hosts. Besides providing additional biological data, knowledge of the host permits easy acquisition of more specimens of a given species, and allows one to associate sexes and adult and immature forms. In many cases, the immature forms show the best characters for separation of taxa.

General references for the Cecidomyiidae are: Pritchard (1947, 1948 and 1951) for the Lestremiinae, Panelius (1965) and Wyatt (1967) for the Porricondylinae, and Harris (1966b) and Möhn (1961b) for the Cecidomyiidi and Asphondyliidi, respectively.

I have included the known host data for each species and a separate host index. The plant host names were checked in Index Kewensis for spelling and current placement. Any names not found therein are placed in quotation marks. Animal host names were checked for accuracy by specialists of the Insect Identification and Parasite Introduction Research Branch of the U. S. Department of Agriculture.

Thanks are due my wife, Sarah, for help in cataloguing and proof-reading the manuscript.

Subfamily LESTREMIINAE



Tribe LESTREMIINI

Genus ANARETE Haliday

ANARETE Haliday, 1833:156. Type-species, *candidata* Haliday (mon.).

Microcerata Felt, 1908b:309. Type-species, *Micromya corni* Felt (orig. des.).

23.1



BUSCKI (Felt), 1915c:198 (*Microcerata*). Type-locality: Cuba, Baracoa.
Distr. — U.S.A., Cuba, Puerto Rico. Ref. — Pritchard, 1951:261.

Genus LESTREMIA Macquart

LESTREMIA Macquart, 1826:173 (separate, 1826:115). Type-species, *cinerea* Macquart (mon.).

CINEREA Macquart, 1826:173, pl. 4, fig. 3. Type-locality: northern France.
Distr. — North America, Chile, Europe. Ref. — Pritchard, 1951:244.

fusca Meigen, 1830:309. Type-locality: Europe.

NIGRA Blanchard, 1852:349. Type-locality: Chile, Chiloe, Carelmapu and Chiloe. Distr. — Chile.

Tribe MICROMYINI

Genus CORINTHOMYIA Felt

CORINTHOMYIA Felt, 1911h:35. Type-species, *Campylomyza hirsuta* Felt (orig. des.) = *brevicornis* (Felt).

BREVICORNIS (Felt), 1907:1 (*Campylomyza*). Type-locality: U.S.A., New York, Nassau. Distr. — Canada, U.S.A., Chile, Europe. Ref. — Pritchard, 1947:39.

Genus MICROMYA Rondani

MICROMYA Rondani, 1840:23. Type-species, *lucorum* Rondani (mon.).

Micromyia, emend.

Ceratomyia Felt, 1911h:33. Type-species, *johannseni* Felt (orig. des.).

JOHANNSENI (Felt), 1911h:33 (*Ceratomyia*). Type-locality: Mexico, Jalisco, Ocotlán. Distr. — U.S.A. (Minn.), Mexico, Costa Rica (unpub.). Ref. — Pritchard, 1947:61.

Genus PSADARIA Enderlein

PSADARIA Enderlein, 1940:668. Type-species, *pallida* Enderlein (orig. des.).

PALLIDA Enderlein, 1940:669, figs. 36-39. Type-locality: Juan Fernandez Is., Masafuera. Distr. — Juan Fernandez Is.

Genus TRICHOPTEROMYIA Williston

TRICHOPTEROMYIA Williston, 1896:255. Type-species, *modesta* Williston (mon.).

MODESTA Williston, 1896:255, pl. 8, figs. 6, 6a, 6b. Type-locality: St. Vincent. Distr. — U.S.A., Dominica (unpub.), St. Vincent, Mexico (unpub.), England. Ref. — Pritchard, 1947:59.



Subfamily PORRICONDYLINAE

Tribe ASYNAPTINI

Genus ASYNAPTA Loew

Cecidomyia, subg. ASYNAPTA Loew, 1850:21, 39. Type-species, *longicollis* Loew (Karsch, 1877:14). The three species listed below are cambium miners under the bark of citrus, cotton, and mango, respectively.

CITRINAE Felt, 1932:117. Type-locality: Puerto Rico, Isabela. Distr. — Puerto Rico.

GOSSYPH (Coquillett), 1905:200 (*Porricondyla*). Type-locality: Barbados. Distr. — Montserrat, Barbados. N. COMB.

MANGIFERAE Felt, 1909:299. Type-locality: Barbados. Distr. — Barbados.

Genus CAMPTOMYIA Kieffer

CAMPTOMYIA Kieffer, 1894a:86 (also 1894b:323). Type-species, *erythromma* Kieffer (orig. des., as gen. n., sp. n.).

PARRISHI (Felt), 1915a:152 (*Porricondyla*). Type-locality: Guyana, Bartica. Distr. — Guyana. N. COMB.

Tribe HETEROPEZINI

Genus MIASTOR Meinert

MIASTOR Meinert, 1864:42. Type-species, *metroloas* Meinert (mon.). Unnamed sp., Williston, 1896:254, pl. 8, figs. 4, 4a. Distr. — St. Vincent. Probably not *Miastor*.

Tribe PORRICONDYLINI

Genus HOLONEURUS Kieffer

Holoneura Kieffer, 1894a:84 (preocc. Tetens, 1891). Type-species, *cincta* Kieffer (Kieffer, 1894b:316).

HOLONEURUS Kieffer, 1895a:115 (nom. nov. for *Holoneura* Kieffer). Type-species, *Holoneura cincta* Kieffer (aut.).

OCCIDENTALIS Felt, 1911i:190. Type-locality: Panama, Canal Zone, Paraisé. Distr. — Panama.

Tribe WINNERTZIINI

Genus WINNERTZIA Rondani

Asinapta, subg. WINNERTZIA Rondani, 1860:290. Type-species, *Cecidomyia lugubris* Winnertz (orig. des.).

Unnamed sp., Williston, 1896:254, pl. 8, fig. 3. Distr. — St. Vincent.



UNPLACED GENERA OF PORRICONDYLIINAE

Genus FELTOMYINA Alexander

Feltomyia Alexander, 1936:12 (preocc. Kieffer, 1913a). Type-species, *polymera* Alexander (orig. des.).

FELTOMYINA Alexander, 1937:60 (nom. nov. for *Feltomyia* Alexander). Type-species, *Feltomyia polymera* Alexander (aut.).

POLYMERA (Alexander), 1936:14 (*Feltomyia*). Type-locality: Panama, Chiriqui, Potrerillos. Distr. — Panama.

Genus JOHNSONOMYIA Felt

JOHNSONOMYIA Felt, 1908b:417. Type-species, *rubra* Felt (orig. des.).

BRASILIENSIS Felt, 1915a:153. Type-locality: Brazil, Pará, Igarapé-Açu. Distr. — Brazil.

CINCTA Felt, 1912c:103. Type-locality: Guatemala, Polochic R. Distr. — Guatemala.

Subfamily CECIDOMYIINAE

Supertribe OLIGOTROPHIDI

Genus ANGEIOMYIA Kieffer & Herbst

ANGEIOMYIA Kieffer & Herbst, 1909:124. Type-species, *spinulosa* Kieffer & Herbst (orig. des., as gen. n., sp. n.).

SPINULOSA Kieffer & Herbst, 1909:124, fig. 5. Type-locality: Chile. Host — *Hydrangea scandens* Poepp (Saxifragaceae). Distr. — Chile.

Genus AUSTROLAUTHIA Brèthes

AUSTROLAUTHIA Brèthes, 1914:153. Type-species, *spagazzini* Brèthes (orig. des.).

SPEGAZZINI Brèthes, 1914:153, fig. 3. Type-locality: Argentina, Buenos Aires, Buenos Aires. Host — *Euphorbia serpens* H. B. & K. (Euphorbiaceae). Distr. — Argentina.

Genus CALMONIA Tavares

CALMONIA Tavares, 1917b:173. Type-species, *urostigmatis* Tavares (orig. des.).

UROSTIGMATIS Tavares, 1917b:174, pl. 10, figs. 10-11; pl. 11, fig. 9. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Urostigma* sp. (Moraceae). Distr. — Brazil.

Genus CALOPEDILA Kieffer

CALOPEDILA Kieffer, 1913b:49. Type-species, *Rhopalomyia herbsti* Kieffer (orig. des.).



HERBSTI (Kieffer), 1903:226 (*Rhopalomyia*). Type-locality: Chile, Concepción, vic. Concepción. Host — *Baccharis bogotensis* H. B. & K., and *B. rosmarinifolia* Hook. & Arn. (Compositae). Distr. — Chile.

Genus DASINEURA Rondani

DASINEURA Rondani, 1840:17. Type-species, *obscura* Rondani (Rondani, 1856:200).

Dasyneura Agassiz & Loew, 1846:11, emend. (preocc. Saunders, 1842).

Perrisia Rondani, 1846:371. Type-species, *Cecidomyia urticae* Perris (orig. des.).

BRAZILIENSIS (Tavares), 1922:12 (*Perrisia*). Type-locality: Brazil. Host — *Protium heptaphyllum* March (Burseraceae). Distr. — Brazil. N. comb.

CHILENSIS (Kieffer & Herbst), 1909:121, fig. 3 (*Perrisia*). Type-locality: Chile, Concepción, Concepción. Host — *Baccharis rosmarinifolia* Hook. & Arn. (Compositae). Distr. — Chile. N. comb.

EUGENIAE Felt, 1912c:106. Type-locality: U.S.A., Florida, Key West. Host — *Eugenia buxifolia* Phil. (Myrtaceae). Distr. — U.S.A. (Florida), Puerto Rico.

GARDOQUIAE Kieffer & Herbst, 1905:65. Type-locality: Chile, Concepción, Concepción. Host — *Gardoquia gilliesi* R. Grah. (Labiatae). Distr. — Chile. Ref. — Kieffer & Herbst, 1906:224, fig. 1; pl. 1, figs. 1-2.

GRACILICORNIS (Kieffer & Herbst), 1905:64 (*Perrisia*). Type-locality: Chile, Concepción, vic. Concepción. Host — undet. shrub. Distr. — Chile. Ref. — Kieffer & Herbst, 1906:225, fig. 2. N. comb.

SUBINERMIS (Kieffer & Herbst), 1911:696 (*Perrisia*). Type-locality: Chile, Valparaíso, Valparaíso. Host — *Baccharis rosmarinifolia* Hook. & Arn. (Compositae). Distr. — Chile. N. comb.

Unnamed spp.: Kieffer & Herbst, 1906:228, figs. 3-4; host, "*Baldoa fragrans* Gay"; distr., Chile, Concepción, Concepción, and Kieffer & Herbst, 1911:697; host, *Baccharis rosmarinifolia* Hook. & Arn. (Compositae); distr., Chile, Valparaíso, Valparaíso.

Genus GUAREPHILA Tavares

GUAREPHILA Tavares, 1909:18. Type-species, *albida* Tavares (mon.).

ALBIDA Tavares, 1909:18, pl. 2, figs. 7-10; pl. 6, fig. 5. Type-locality: Brazil, Rio Grande do Sul, vic. São Leopoldo. Host — *Guarea trichiloides* L. (Meliaceae). Distr. — Brazil.

Genus MISOSPATHA Kieffer

MISOSPATHA Kieffer, 1913b:48. Type-species, *Rhopalomyia globifex* Kieffer & Jörgensen (orig. des.).

GLOBIFEX (Kieffer & Jörgensen), 1910:364, figs. 2, 2a (*Rhopalomyia*). Type-localities: Argentina, Mendoza, near Chacras de Coria (11 km s. Mendoza), Pedregal (15 km e. Mendoza), Potrerillos and San Ignacio; San Juan, near Cancete (n. Mendoza). Host — *Baccharis salicifolia* Pers. (Compositae). Distr. — Argentina.



- OREIPLANA (Kieffer & Jörgensen), 1910:441 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, Blanco Encallada. Host — *Verbena seriphoides* Gill. & Hook. (Verbenaceae). Distr. — Argentina.
- PROSOPIDIS (Kieffer & Jörgensen), 1910:427, fig. 51 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, near Chacras de Coria (11 km s. Mendoza) and La Paz, and San Luis, nr. Alto Pencoso. Host — *Prosopis alpataco* Phil. (Leguminosae). Distr. — Argentina.
- TRICYCLAE (Kieffer & Jörgensen), 1910:440, fig. 61 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, near Blanco Encallada and San Ignacio. Host — *Tricycla spinosa* Cav. (Nyctaginaceae). Distr. — Argentina.
- VERBENAE (Kieffer & Jörgensen), 1910:441 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, Cordillera Mendoza. Host — *Verbena aspera* Gill. & Hook. (Verbenaceae). Distr. — Argentina.

Genus OLIGOTROPHUS Latreille

- OLIGOTROPHUS Latreille, 1805:288. Type-species, *Tipula juniperina* Linnaeus (mon.).
- EUGENIAE Kieffer & Herbst, 1909:125, fig. 6. Type-locality: Chile, Concepción, Concepción. Host — *Myrceugenia stenophylla* Berg (Myrtaceae). Distr. — Chile.
- LYCHCOLA Kieffer & Jörgensen, 1910:409. Type-locality: Argentina, Mendoza, "Cordillera de Mendoza". Host — *Lycium chilense* Bert. (Solanaceae). Distr. — Argentina.
- NECTANDRAE Kieffer & Jörgensen, 1910:442. Type-locality: Paraguay. Host — *Nectandra megapotamica* Mez (Lauraceae). Distr. — Paraguay.

Genus HAPLUSIA Karsch

- HAPLUSIA Karsch, 1877:15, 16. Type-species, *plumipes* Karsch (orig. des.).
- PLUMIPES Karsch, 1877:16, fig. 1. Type-locality: Brazil, Bahia. Host — unknown. Distr. — Brazil.
- Unnamed sp., Williston, 1896:254, pl. 8, fig. 5; host unknown; distr., St. Vincent.

Genus HAPLOPALPUS Rübsaamen

- HAPLOPALPUS Rübsaamen, 1916a:471. Type-species, *serjaneae* Rübsaamen (mon.).
- SERJANAE Rübsaamen, 1916a:473, figs. 49-53. Type-locality: Peru, Madre de Dios, Auristella on Rio Acre. Host — *Serjanea* sp. (Sapindaceae): Distr. — Peru.

Genus JANETIELLA Kieffer

- JANETIELLA Kieffer, 1898:23. Type-species, *Cecidomyia thymi* Kieffer (Coquillett, 1910:556).



ACUTICAUDA Kieffer & Herbst, 1906:227. Type-locality: Chile, Concepción, vic. Concepción. Host — undet. shrub. Distr. — Chile. Ref. — Kieffer & Herbst, 1905:64.

MONTIVAGA Kieffer & Jörgensen, 1910:27, fig. 56. Type-locality: Argentina, Mendoza, "Cordillera de Mendoza". Host — *Senecio mendocinus* Phil. (Compositae). Distr. — Argentina.

SCHWARZI (Felt), 1911:191 (*Lasiopteryx*). Type-locality: Panama, Canal Zone, Paraíso. Host — dead *Ficus* sp. (Moraceae). Distr. — Panama. N. COMB.

Genus LYCIOMYIA Kieffer & Jörgensen

LYCIOMYIA Kieffer & Jörgensen, 1910:412. Type-species, *gracilis* Kieffer & Jörgensen (mon.).

GRACILIS Kieffer & Jörgensen, 1910:412, fig. 40. Type-locality: Argentina. Host — *Lycium gracile* Meyen (Solanaceae). Distr. — Argentina.

Genus PANTELOLA Kieffer

PANTELOLA Kieffer, 1913b:49. Type-species, *Rhopalomyia haasi* Kieffer (orig. des.).

BEDEGUARIS (Kieffer & Jörgensen), 1910:403, fig. 32 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, "Cordillera de Mendoza". Host — *Lycium chilense* Bert. (Solanaceae). Distr. — Argentina.

LIPPIAE (Kieffer & Jörgensen), 1910:401, fig. 31 (*Rhopalomyia*). Type-locality: Argentina, Mendoza, "Cordillera de Mendoza". Host — *Lippia foliosa* Phil. (Verbenaceae). Distr. — Argentina.

Genus PERNETTYELLA Kieffer & Herbst

PERNETTYELLA Kieffer & Herbst, 1909:126. Type-species, *longicornis* Kieffer & Herbst (orig. des., as gen. n., sp. n.).

LONGICORNIS Kieffer & Herbst, 1909:126, fig. 7. Type-locality: Chile, Concepción, Concepción. Host — *Pernettya furens* Klotzsch. (Ericaceae). Distr. — Chile.

Genus PHAENOLAUTHIA Kieffer

PHAENOLAUTHIA Kieffer, 1912a:2. Type-species, *Ledomyia obscuripennis* Kieffer (orig. des.).

Lasiopteryx Felt, nec Stephens.

MANIHOT (Felt), 1912e:144 (*Lasiopteryx*). Type-locality: St. Vincent. Host — *Manihot utilisima* Pohl (Euphorbiaceae). Distr. — St. Vincent. N. COMB.

Genus PROMIKIOLA Kieffer & Herbst

PROMIKIOLA Kieffer & Herbst, 1911:700. Type-species, *rubra* Kieffer & Herbst (orig. des., as gen. n., sp. n.).



RUBRA Kieffer & Herbst, 1911:700, fig. 5. Type-locality: Chile, Valparaíso, Valparaíso. Host — *Colliguaya odorifera* Mohina (Euphorbiaceae).
Distr. — Chile.

Genus PSEUDOMIKIOLA Brèthes

PSEUDOMIKIOLA Brèthes, 1917b:411. Type-species, *lippiac* Brèthes (orig. des.).

LIPPIAE Brèthes, 1917b:413, figs. 1-6. Type-locality: Argentina, Entre Rios, Concepción del Uruguay. Host — *Lippia* sp. (Verbenaceae).
Distr. — Argentina.

Genus RIVERAELLA Kieffer & Herbst

RIVERAELLA Kieffer & Herbst, 1911:698. Type-species, *colliguayae* Kieffer & Herbst (orig. des., as gen. n., sp. n.).

COLLIGUAYAE Kieffer & Herbst, 1911:698, figs. 3, 4. Type-locality: Chile, Valparaíso, Valparaíso. Host — *Colliguaya odorifera* Mohina (Euphorbiaceae). Distr. — Chile.

Unnamed sp. Kieffer & Herbst, 1911:701. Host — *Colliguaya odorifera* Mohina (Euphorbiaceae). Distr. — Chile, Valparaíso, Valparaíso.

Genus SCHEUERIA Kieffer & Herbst

SCHEUERIA Kieffer & Herbst, 1909:120. Type-species, *longicornis* Kieffer & Herbst (orig. des., as gen. n., sp. n.).

AGLOMERATA Kieffer, 1913a:39. Type-locality: Chile, Concepción, Concepción. Host — *Baccharis eupatorioides* Hook. & Arn. (Compositae).
Distr. — Chile. Ref. — Kieffer & Herbst, 1909:120, fig. 2.

LONGICORNIS Kieffer & Herbst, 1909:120, fig. 1. Type-locality: Chile, Concepción, Concepción. Host — *Baccharis eupatorioides* Hook. & Arn. (Compositae). Distr. — Chile.

Genus ULEIA Rübsaamen

ULEIA Rübsaamen, 1905:85. Type-species, *clusiae* Rübsaamen (mon.).

CLUSIAE Rübsaamen, 1905:85. Type-locality: Brazil. Host — *Clusia* sp. (Guttiferae). Distr. — Brazil.

Genus XYLOPERRISIA Kieffer

XYLOPERRISIA Kieffer, 1913a:69. Type-species, *Perrisia azarae* Kieffer (mon.).

AZARAE (Kieffer), 1903:227 (*Perrisia*). Type-locality: Chile, Concepción, Concepción. Host — *Gaultheria vernalis* Kunze (Ericaceae).
Distr. — Chile.



Supertribe ASPHONDYLIIDI

Tribe ASPHONDYLIINI

Genus ASPHONDYLIA Loew

Cecidomyia, subg. ASPHONDYLIA Loew, 1850:21, 37. Type-species, *sarothamni* Loew (Karsch, 1877:15).

Gisonobasis Rübсаamen, 1916a:432. Type-species, *tournefortiae* Rübсаamen (Rübсаamen, 1916b:12).

Most species in this genus cause galls in the flower parts, seeds or fruit of a wide array of plants. The others cause galls on stems, leaves or buds.

Subgenus ASPHONDYLIA Loew

ACHATOCARPI Möhn, 1959:304, pl. 37, figs. 19-20. Type-locality: El Salvador, La Libertad, San Diego, c. La Libertad. Host — *Achatocarpus nigricans* Trlana (Amarantaceae). Distr. — El Salvador.

AJALLAI Möhn, 1959:366, pl. 41, figs. 120-122. Type-locality: El Salvador, La Libertad, Los Chorros. Host — *Vernonia canescens* H.B. & K. (Compositae). Distr. — El Salvador.

ANACELIAE Möhn, 1959:355, pl. 40, figs. 103-105. Type-locality: El Salvador, La Libertad, Cumbre, Hacienda Chilata. Host — *Bidens squarrosa* H.B. & K. (Compositae). Distr. — El Salvador.

ANDIRAE Möhn, 1959:314, pl. 38, figs. 35-37. Type-locality: El Salvador, La Libertad, San Diego, e. La Libertad. Host — *Andira inermis* H. B. & K. (Leguminosae). Distr. — El Salvador.

ANNONAE Möhn, 1959:309, pl. 38, figs. 26-28. Type-locality: El Salvador, Santa Ana, n. El Roblarón. Host — *Anona cherimola* Mill. (Anonaceae). Distr. — El Salvador.

ARDISIAE Möhn, 1959:333, pl. 39, figs. 67-68. Type-locality: El Salvador, La Libertad, n. El Cimarrón. Host. — *Ardisia paschalis* J. D. Smith (Myrsinaceae). Distr. — El Salvador.

BOERHAAVIAE Möhn, 1959:302, pl. 37, figs. 16-18. Type-locality: El Salvador, San Vicente, nr. Rio San Felipe. Host — *Boerhaavia erecta* L. (Nyctaginaceae). Distr. — El Salvador.

BUDDLEIA Feit, 1935:2. Type-locality: U.S.A., Texas, Austin. Host — *Buddleia americana* L. and *B. racemosa* Torr. (Loganiaceae). Distr. — U.S.A. (Texas), El Salvador. Ref. — Möhn, 1960a:209, pl. 23, figs. 26-29.

CABEZASAE Möhn, 1959:312, pl. 38, figs. 32-34. Type-locality: El Salvador, La Libertad, Los Choros. Host — *Crotalaria mucronata* Desv. (Leguminosae). Distr. — El Salvador.

CALEAE Möhn, 1959:357, pl. 41, figs. 106-108. Type-locality: El Salvador, San Salvador, Picacho, below remains of cloud forest. Host — *Calea integrifolia* (DC) Hemsl. (Compositae). Distr. — El Salvador.

CAMARAE Möhn, 1959:340, pl. 40, figs. 78-81. Type-locality: El Salvador, Cuscatlán, 27 km e. San Martin. Host — *Lantana camara* L. (Verbenaceae). Distr. — El Salvador.



- CESTRUM Möhn, 1959:344, pl. 40, figs. 86-87. Type-locality: El Salvador, La Libertad, Los Chorros. Host — *Cestrum nocturnum* L. (Solanaceae). Distr. — El Salvador.
- CISSIS Möhn, 1959:324, pl. 39, figs. 51-52. Type-locality: El Salvador, La Paz, about 8 km before Herradura. Host — *Cissus sicyoides* L. (Vitaceae). Distr. — El Salvador.
- CLEMATIDIS Felt, 1935:3. Type-locality: U.S.A., Texas, Austin. Hosts — *Clematis grossa* Benth., *C. dioica* L., and *C. drummondii* Torr. & Gray (Ranunculaceae). Distr. — U.S.A. (Texas), El Salvador. Ref. — Möhn, 1960a:206, pl. 23, figs. 20-22.
- CORBULAE Möhn, 1960a:197, pl. 22, figs. 1-3. Type-locality: El Salvador, Ahuachapan, nw. Las Chinamas. Host — *Eupatorium* sp. (Compositae). Distr. — El Salvador.
- CORDIAE Möhn, 1959:338, pl. 40, figs. 75-77. Type-locality: El Salvador, La Libertad, w. Ateos. Host — *Cordia alba* Roem. & Schult. (Boraginaceae). Distr. — El Salvador.
- CURATELLAE Möhn, 1959:330, pl. 39, figs. 62-63. Type-locality: El Salvador, Chalatenango, se. La Palma. Host — *Curatella americana* L. (Dilleniaceae). Distr. — El Salvador.
- ERYTHROXYLIS Möhn, 1959:315, pl. 39, figs. 38-39. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Erythroxylum mexicanum* H. B. & K. (Linaceae). Distr. — El Salvador.
- GODMANIAE Möhn, 1959:346, pl. 40, figs. 89-91. Type-locality: El Salvador, Chalatenango, w. Chalatenango. Host — *Godmania aesculifolia* (H. B. & K.) Standley (Bignoniaceae). Distr. — El Salvador.
- GUAREAE Möhn, 1959:316, pl. 38, figs. 40-42. Type-locality: El Salvador, Usulután, forest of San Marcos Lempa. Host — *Guarea* sp. (Meliaceae). Distr. — El Salvador.
- HELICTERIS Möhn, 1959:327, pl. 39, figs. 56-58. Type-locality: El Salvador, La Paz, nw. Olocuilta. Host — *Helicteres guazumaefolia* H. B. & K. (Sterculiaceae). Distr. — El Salvador.
- HERCULESI Möhn, 1959:363, pl. 41, figs. 117-119. Type-locality: El Salvador, Chalatenango, se. La Palma. Host — *Vernonia canescens* H. B. & K. (Compositae). Distr. — El Salvador.
- HYPTIS Möhn, 1960a:206, pl. 22, fig. 19. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Hyptis verticillata* Jacq. (Labiatae). Distr. — El Salvador.
- INDIGOFERAE Möhn, 1960a:204, pl. 22, figs. 15-16. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Indigofera suffruticosa* Mill. (Leguminosae). Distr. — El Salvador.
- JATROPHAE Möhn, 1959:319, pl. 38, figs. 46-47. Type-locality: El Salvador, La Paz, 8 km before Herradura. Host — *Jatropha curcas* L. (Euphorbiaceae). Distr. — El Salvador.
- LIPPIAE Möhn, 1959:342, pl. 40, figs. 82-84. Type-locality: El Salvador, San Salvador, e. San Marcos. Host — *Lippia cardiostegia* Benth. (Verbenaceae). Distr. — El Salvador.
- LONCHOCARPI Möhn, 1960a:205, pl. 22, figs. 17-18. Type-locality: El Salvador, Sonsonate, n. Los Cobanos. Host — *Lonchocarpus minimiflorus* J. D. Smith (Leguminosae). Distr. — El Salvador.

- MALVAVISCI Möhn, 1959:325, pl. 39, figs. 53-55. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Malvaviscus arboreus* Cav (Malvaceae). Distr. — El Salvador.
- MELANTHERAE Möhn, 1959:360, pl. 41, figs. 112-114. Type-locality: El Salvador, San Miguel, n. El Delirio. Host — *Melanthera nivea* (L.) Small (Compositae). Distr. — El Salvador.
- OCIMI Möhn, 1959:343, pl. 40, fig. 85. Type-locality: El Salvador, Morazan, sw. Jocoro. Host — *Ocimum micranthum* Willd. (Labiatae). Distr. — El Salvador.
- PALACIOSI Möhn, 1959:310, pl. 38, figs. 29-31. Type-locality: El Salvador, Santa Ana, nw. Ciudad Arce. Host — *Crotalaria longirostrata* Hook. & Arn. (Leguminosae). Distr. — El Salvador.
- PARASITICOLA Möhn, 1960:202, pl. 22, figs. 10-13. Type-locality: El Salvador, Ahuachapan, nr. Ahuachapan. Host — *Struthanthus marginatus* (Desr.) Blume (Loranthaceae). Distr. — El Salvador.
- PORTULACAE Möhn, 1959:308, pl. 37, figs. 24-25. Type-locality: El Salvador, La Libertad, San Diego, e. La Libertad. Host — *Portulaca oleracea* L. (Portulacaceae). Distr. — El Salvador.
- PSYCHOTRIAE Möhn, 1959:349, pl. 40, figs. 95-96. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Psychotria pubescens* Sw. (Rubiaceae). Distr. — El Salvador.
- RANDIAE Möhn, 1959:350, pl. 40, figs. 97-99. Type-locality: El Salvador, La Libertad, se. Ciudad Arce. Host — *Randia aculeata* L. (Rubiaceae). Distr. — El Salvador.
- RONDELETIAE Möhn, 1959:352, pl. 40, figs. 100-102. Type-locality: El Salvador, San Salvador, Picacho, below remains of cloud forest. Host — *Rondeletia strigosa* (Benth.) Hemsl. (Rubiaceae). Distr. — El Salvador.
- RUELLIAE Möhn, 1959:347, pl. 40, figs. 92-94. Type-locality: El Salvador, Santa Ana, on bank of Lake Coatepeque. Host — *Ruellia albicaulis* Bert. (Acanthaceae). Distr. — El Salvador.
- SALVADORENSIS Möhn, 1959:358, pl. 41, figs. 109-111. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Hymenostephium cordatum* (Blake) Hook. & Arn. (Compositae). Distr. — El Salvador.
- SCOPARIAE Möhn, 1959:345, pl. 40, fig. 88. Type-locality: El Salvador, Chalatenango, e. Santa Barbara. Host — *Scoparia dulcis* L. (Scrophulariaceae). Distr. — El Salvador.
- SERJANIAE Möhn, 1959:322, pl. 38, figs. 48-50. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Serjanea goniocarpa* Radlk. (Sapindaceae). Distr. — El Salvador.
- TABERNAEMONTANAE Möhn, 1959:333, pl. 39, figs. 69-71. Type-locality: El Salvador, Sonsonate, n. Los Cobanos. Host — *Tabernaemontana amygdalifolia* Jacq. (Apocynaceae). Distr. — San Salvador.
- TALINI Möhn, 1959:305, pl. 37, figs. 21-23. Type-locality: El Salvador, La Libertad, Cumbre, Hacienda Chilata. Host — *Talinum paniculatum* Jacq. (Portulacaceae). Distr. — El Salvador.
- THEVETIAE Möhn, 1959:337, pl. 39, figs. 72-74. Type-locality: El Salvador, San Vicente, km 54, se. Santo Domingo. Host — *Thevetia plumeriaefolia* Benth. (Apocynaceae). Distr. — El Salvador.



- TITHONIAE** Möhn, 1960a:200, pl. 22, figs. 6-7. Type-locality: El Salvador, La Libertad, e. Sitio del Niño. Host — *Tithonia rotundifolia* (Mill.) Blake (Compositae). Distr. — El Salvador.
- TOURNEFORTIAE** (Rübsaamen), 1916a:433, figs. 2, 3b (*Gisonobasis*). Type-localities: Brazil, Acre, vicinity of São Francisco on Acre R.; Peru, Madre de Dios, Auristella on Acre R. Hosts — *Tournefortia volubilis* L. and *T. angustiflora* Ruiz & Pav. (Boraginaceae). Distr. — El Salvador, Brazil, Peru. Ref. — Möhn, 1960a:211, pl. 23, figs. 30-32.
- TRICHILLAE** Möhn, 1959:318, figs. 43-45. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Trichilia* sp. (Meliaceae). Distr. — El Salvador.
- TRIXIDIS** Möhn, 1959:362, pl. 41, figs. 115-116. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Trixis radialis* O. Ktze. (Compositae). Distr. — El Salvador.
- VINCENTI** Felt, 1911b:109. Type-locality: St. Vincent. Hosts — *Jussiaea suffruticosa* L. and *J. linifolia* Vahl. (Onagraceae). Distr. — St. Vincent, El Salvador. Ref. — Möhn, 1960a:208, pl. 23, figs. 23-25.
- WALTHERIAE** Möhn, 1959:328, pl. 39, figs. 59-61. Type-locality: El Salvador, Santa Ana, e. Tablón del Coco. Host — *Waltheria indica* L. (Sterculiaceae). Distr. — El Salvador.
- XIMENIAE** Möhn, 1959:301, pl. 37, figs. 1-15. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Ximenia americana* L. (Olacaceae). Distr. — El Salvador.
- XYLOSMATIS** Möhn, 1959:331, pl. 39, figs. 64-66. Type-locality: El Salvador, Santa Ana, se. Candelaria, Hacienda Cortez. Host — *Xylosma flexuosum* Hemsl. (Flacourtiaceae). Distr. — El Salvador.
- ZACATECHICHI** Möhn, 1960a:199, pl. 22, figs. 4-5. Type-locality: El Salvador, San Vicente, km 54, se. Santo Domingo. Host — *Calea zacatechichi* Schlecht. (Compositae). Distr. — El Salvador.
- ZEXMENIAE** Möhn, 1960a:201, pl. 22, figs. 8-9. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Zexmenia frutescens* (Mill.) Blake (Compositae). Distr. — El Salvador.
- Unnamed sp., Möhn, 1960a:213. Host — *Casearia banquitana* E. H. L. Krause (Samydaceae). Distr. — El Salvador, Sonsonate, km 5 on road to Los Cobanos.

Subgenus EOASPHONDYLIA Möhn

- Asphondylia**, subg. EOASPHONDYLIA Möhn, 1960a:214. Type-species, *convolvuli* Möhn (orig. des.).
- AYENIAE** Möhn, 1960a:216, pl. 24, figs. 49-65. Type-locality: El Salvador, Santa Ana, road to Metapán, ne. Santa Ana. Host — *Ayenia pusilla* L. (Sterculiaceae). Distr. — El Salvador.
- CONVOLVULI** Möhn, 1960a:214, pl. 23, figs. 33-46; pl. 24, figs. 47-48. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Ipomoea arborescens* (H. B.) Don (Convolvulaceae). Distr. — El Salvador.

UNPLACED SPECIES OF *Asphondylia*

- ALTANI Felt, 1915c:206. Type-locality: Nicaragua, Carazo, San Marcos. Host — unknown. Distr. — Nicaragua.
- ATRIPLICIS (Townsend), 1893:1021 (*Cecidomyia*). Type-locality: U.S.A., New Mexico, Las Cruces. Host — *Atriplex canescens* (Pursh.) Nutt. (Chenopodiaceae). Distr. — U.S.A. (southwestern), Mexico (Baja California).
- ATTENUATATA Felt, 1909:300. Type-locality: British West Indies (Antigua?). Host — *Clerodendron aculeatum* Griseb. (Verbenaceae). Distr. — British West Indies (Antigua?).
- BACCHARIS Kieffer & Herbst, 1905:65. Type-locality: Chile, Concepción, vic. San Vicente. Host — *Baccharis* sp. (Compositae). Distr. — Chile. Ref. — Kieffer & Herbst, 1906:230.
- BAHIENSIS Tavares, 1917b:156, fig. 4 (B, C); pl. 8, figs. 5-6. Type-locality: Brazil, Bahia, Salvador. Host — sp. of Rubiaceae. Distr. — Brazil.
- BORRERIAE Rübssaamen, 1905:74. Type-locality: Brazil, Guanabara, Rio de Janeiro, Arpoador. Hosts — *Diodia* sp. and *Borreria* sp. (Rubiaceae). Distr. — Brazil.
- CRASSIPALPIS Kieffer & Jörgensen, 1910:365, figs. 3-4. Type-localities: Argentina, Mendoza, near Pedregal and north of Chacras de Corla (11 km s. Mendoza); San Juan, near Cancete (n. Mendoza). Host — *Baccharis salicifolia* Pers. (Compositae). Distr. — Argentina.
- HIERONYMI (Weyenbergh), 1875:164 (*Lasioptera*). Type-locality: Argentina. Host — unknown. Distr. — Argentina.
- MONACHIA Osten Sackén, 1869:299. Type-locality: U.S.A., New York. Hosts — *Solidago* spp. (Compositae). Distr. — U.S.A., Mexico.
- OPUNTIAE Felt, 1908b:377. Type-locality: U.S.A., Texas, Sinton. Host — *Opuntia* sp. (Cactaceae). Distr. — U.S.A. (southwestern), Mexico.
- PARVA Tavares, 1917b:159, fig. 4(A). Type-locality: Brazil, Bahia, Salvador. Host — sp. of Rubiaceae. Distr. — Brazil.
- PATTERSONI Felt, 1911d:301. Type-locality: St. Vincent. Host — *Citharexylum quadrangulare* Jacq. (Verbenaceae). Distr. — St. Vincent.
- ROCHAE Tavares, 1918b:68, pl. 3, figs. 9-11. Type-locality: Brazil, Ceará, nr. Fortaleza. Host — *Jussiaea* sp. (Onagraceae). Distr. — Brazil.
- SICCAE Felt, 1908b:376. Type-locality: Jamaica. Host — *Phyllanthus distichus* Muell. Arg. (Euphorbiaceae). Distr. — Jamaica.
- STACHYTARPHETA Barnes, 1932:482. Type-locality: Trinidad. Host — *Stachytarpheta cajanensis* Vahl. (Verbenaceae). Distr. — Trinidad.
- STRUTHANTHI (Rübssaamen), 1916a:432, fig. 1 (*Gisonobasis*). Type-locality: Brazil, Ceará, Serra do Baturité. Host — *Struthanthus* sp. (Loranthaceae). Distr. — Brazil. N. comb.
- SULPHUREA Tavares, 1909:14, pl. 1, fig. 1; pl. 5, fig. 6. Type-locality: Brazil, Rio Grande do Sul, vic. São Leopoldo. Host — *Smilax* sp. (Liliaceae). Distr. — Brazil.
- SWAEDICOLA Kieffer & Jörgensen, 1910:435. Type-localities: Argentina, Mendoza, La Paz, Potrerillos, and San Ignacio; San Juan, nr. Cancete (n. Mendoza). Host — *Suaeda divaricata* Moq. (Chenopodiaceae). Distr. — Argentina.



ULEI Rübsaamen, 1908a:172. Type-locality: Brazil, Rio de Janeiro, Palmeiras. Host — *Mikania* sp. (Compositae). Distr. — Brazil.

Unnamed sp., Tavares, 1909:8, pl. 3, fig. 10. Host — *Mikania amara* Willd. (Compositae). Distr. — Brazil, Rio Grande do Sul, vic. São Leopoldo.

Genus BRUGGMANNIELLA Tavares

BRUGGMANNIELLA Tavares, 1909:19. Type-species, *braziliensis* Tavares (mon.).

BRAZILIENSIS Tavares, 1909:20, pl. 1, figs. 2-5; pl. 3, fig. 11-12. Type-locality: Brazil, Rio Grande do Sul, São Leopoldo. Host — *Sorocea ilicifolia* Miq. (Urticaceae). Distr. — Brazil. Ref. — Möhn, 1963b:5, figs. 94-112.

Genus HEMIASPHONDYLIA Möhn

HEMIASPHODYLIA Möhn, 1960a:229. Type-species, *mimosae* Möhn (orig. des.).

MIMOSAE Möhn, 1960a:230, pl. 27, figs. 121-137. Type-locality: El Salvador, La Libertad, n. El Cimarrón. Host — *Mimosa albida* Humb. & Bonpl. (Leguminosae). Distr. — El Salvador.

Genus HEMIBRUGGMANNIELLA Möhn

HEMIBRUGGMANNIELLA Möhn, 1961a:6. Type-species, *Bruggmanniella oblita* Tavares (orig. des.).

OBLITA (Tavares), 1920a:34, figs. 1-3 (*Bruggmanniella*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — sp. of Anacardiaceae, prob. *Schinus* sp. Distr. — Brazil. Ref. — Möhn, 1961a:7, pl. 2, figs. 21-29.

Genus HETERASPHONDYLIA Möhn

HETERASPHONDYLIA Möhn, 1960a:227. Type-species, *salvadorensis* Möhn (orig. des.).

SALVADORENSIS Möhn, 1960a:228, pl. 26, figs. 111-114; pl. 27, figs. 115-120. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Piper* sp. (Piperaceae). Distr. — El Salvador.

Genus PERASPHONDYLIA Möhn

PERASPHONDYLIA Möhn, 1960a:218. Type-species, *reticulata* Möhn (orig. des.).

RETICULATA Möhn, 1960a:219, pl. 24, figs. 66-70; pl. 25, figs. 71-82. Type-locality: El Salvador, Chalatenango, w. Chalatenango. Hosts — *Eupatorium odoratum* L., *Eupatorium* sp. (Compositae). Distr. — El Salvador.

Genus RHOASPHONDYLIA Möhn

RHOASPHONDYLIA Möhn, 1960a:224. Type-species, *Oxasphondylia friburgensis* Tavares (orig. des.).

FRIBURGENSIS (Tavares), 1917b:126, fig. A (5, 6), pl. 7, figs. 6-8 (*Oxasphondylia*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Hosts — *Baccharis dracunculifolia* DC., *B. schultzei* Baker, and *B. trinervis* Pers. (Compositac). Distr. — El Salvador, Brazil. Ref. — Möhn, 1960a:225, pl. 26, figs. 93-110.

Genus SCIASPHONDYLIA Möhn

SCIASPHONDYLIA Möhn, 1960a:221. Type-species, *salviae* Möhn (orig. des.).

SALVIAE Möhn, 1960a:222, pl. 25, figs. 83-90; pl. 26, figs. 91-92. Type-locality: El Salvador, Chalatenango, 15 km. from La Palma. Host — *Salvia* sp. (Labiatae). Distr. — El Salvador.

Genus TAVARESOMYIA Möhn

TAVARESOMYIA Möhn, 1961a:9. Type-species, *Schizomyia mimosae* Tavares (orig. des.).

MIMOSAE (Tavares), 1925:28, fig. 9 (*Schizomyia*). Type-locality: Brazil, Ceará. Host — *Mimosa caesalpiniaefolia* Benth. (Leguminosae). Distr. — Brazil. Ref. — Möhn, 1961a:9, pl. 2, figs. 30-32.

Genus XENASPHONDYLIA Felt

XENASPHONDYLIA Felt, 1915c:202. Type-species, *albipes* Felt (orig. des.).

ALBIPES Felt, 1915c:202, fig. 5. Type-locality: Dominican Republic, Santo Domingo, San Francisco Mts. Host — unknown. Distr. — Dominican Republic. Ref. — Möhn, 1963b:3, figs. 85-93.

Genus ZALEPIDOTA Rübsaamen

ZALEPIDOTA Rübsaamen, 1908:38. Type-species, *piperis* Rübsaamen (orig. des., as gen. n., sp. n.).

Ozobia Kieffer, 1913a:97. Type-species, *tavaresei* Kieffer (orig. des.).

Oxasphondylia Felt, 1915c:204. Type-species, *reticulata* Felt (orig. des.).

PIPERIS Rübsaamen, 1908b:38, fig. 13. Type-locality: Brazil, Guanabara, Rio de Janeiro, Tijuca. Host — *Piper* sp. (Piperaceae). Distr. — Brazil. Refs. — Tavares, 1925:34, figs. 11-12; pl. 1, figs. 9-10; Möhn, 1963b:12, figs. 113-126.

RETICULATA (Felt), 1915c:205, figs. 7-9 (*Oxasphondylia*). Type-locality: Guatemala, Alta Vera Paz, Trece Aguas. Host — *Piper* sp. (Piperaceae). Distr. — Guatemala. Ref. — Möhn, 1963b:21, figs. 131-137.



TAVARESII (Kieffer), 1913a:98 (*Ozobia*). Type-locality: Brazil, Rio Grande do Sul, vic. São Leopoldo. Host — *Piper* sp. (Piperaceae). Distr. — Brazil. Refs. — Tavares, 1909:24, pl. 2, fig. 1; pl. 4, figs. 24-26, pl. 5, figs. 10-13; Möhn, 1963b:15, figs. 127-130.

Tribe POLYSTEPHINI

Genus POLYSTEPHA Kieffer

POLYSTEPHA Kieffer, 1897:11. Type-species, *quercus* Kieffer (orig. des.).

LAPALMAE Möhn, 1960b:335, pl. 48, figs. 11-12. Type-locality: El Salvador, Chalatenango, road to La Palma, ca. 15 km before La Palma. Host — *Quercus hondurensis* Trel. (Fagaceae). Distr. — El Salvador.

SALVADORENSIS Möhn, 1960b:333, pl. 48, figs. 1-10. Type-locality: El Salvador, Chalatenango, road to La Palma, ca. 15 km before La Palma. Host. — *Quercus eugeniaefolia* Liebm. (Fagaceae). Distr. — El Salvador.

Tribe SCHIZOMYIINI

Genus AMELIOMYIA Möhn

AMELIOMYIA Möhn, 1960a:237. Type-species, *zexmeniae* Möhn (orig. des.).

ZEXMENIAE Möhn, 1960a:237, pl. 28, figs. 163-172. Type-locality: El Salvador, Santa Ana, south of Candelaria. Host — *Zexmenia frutescens* (Mill.) Blake (Compositae). Distr. — El Salvador.

Genus BRUGGMANNIA Tavares

BRUGGMANNIA Tavares, 1906:81. Type-species, *braziliensis* Tavares (mon.).

FELTOMYIA Kieffer, 1913:100. Type-species, *Uleela pisonifolia* Felt (orig. des.).

BRAZILIENSIS Tavares, 1906:82, fig. 4. Type-locality: Brazil, Rio Grande do Sul, São Leopoldo. Host — *Myrsine* sp. (Myrsinaceae). Distr. — Brazil. Refs. — Tavares, 1909:22, pl. 6, figs. 8-9; pl. 7, fig. 10; Möhn, 1962b:211, figs. 1-18.

PISONIAE (Felt), 1912a:174 (*Bruggmanniella*). Type-locality: St. Vincent. Host — *Pisonia nigricans* Sw. (Nyctaginaceae). Distr. — St. Vincent. N. COMB.

PISONIFOLIA (Felt), 1912b:353 (*Uleela*). Type-locality: St. Vincent. Host — *Pisonia nigricans* Sw. (Nyctaginaceae). Distr. — U.S.A. (Fla.), St. Vincent. Ref. — Möhn, 1962:217, figs. 19-37.

PSYCHOTRIAE Möhn, 1960b:352, pl. 51, figs. 71-78; pl. 52, figs. 79-91. Type-locality: El Salvador, La Libertad, San Diego, e. La Libertad. Host — *Psychotria carthaginensis* Jac. (Rubiaceae). Distr. — El Salvador.

PUSTULANS Möhn, 1960b:348, pl. 50, figs. 49-52; pl. 51, figs. 53-70. Type-locality: El Salvador, Ateos, w. Ateos. Hosts — *Pisonia macranthocarpa* J. D. Smith and *P. aculeata* L. (Nyctaginaceae). Distr. — El Salvador.

RANDIAE Möhn, 1960b:354, pl. 52, figs. 92-100; pl. 53, figs. 101-109. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Randia spinosa* (Jacq.) Karst. (Rubiaceae). Distr. — El Salvador.

Genus BURSERAMYIA Möhn

BURSERAMYIA Möhn, 1960a:232. Type-species, *burserae* Möhn (orig. des.).

BURSERAE Möhn, 1960a:233, pl. 28, figs. 138-153. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño, w. lava fields. Host — *Bursera gummifera* L. (Burscraceae). Distr. — El Salvador.

Genus MACROPORPA Rübsaamen

MACROPORPA Rübsaamen, 1916a:437. Type-species, *peruviana* Rübsaamen, by present designation.

PERUVIANA Rübsaamen, 1916a:437, figs. 4-8. Type-locality: Peru, Madre de Dios, Aurlstella on Rio Acre. Host — sp. of Malpighiaceae. Distr. — Peru. Ref. — Möhn, 1962:225, figs. 41-55.

ULEI Rübsaamen, 1916a:441, figs. 9-12. Type-locality: Brazil, Acre, San Francisco on Rio Acre. Host — sp. of Lauraceae. Ref. — Möhn, 1962:229.

Genus PISONIAMYIA Möhn

PISONIAMYIA Möhn, 1960b:338. Type-species, *armeniae* Möhn (orig. des.).

ARMENIAE Möhn, 1960b:339, pl. 48, figs. 13-17; pl. 49, figs. 18-30. Type-locality: El Salvador, La Libertad, w. Ateos. Host — *Pisonia macranthocarpa* J. D. Smith (Nyctaginaceae). Distr. — El Salvador.

MEXICANA (Felt), 1911e:465 (*Bruggmanniella*). Type-locality: Mexico, Tamaulipas, Tampico. Host — *Pisonia aculeata* L. (Nyctaginaceae). Distr. — Mexico.

SIMILIS Möhn, 1960b:342, pl. 49, figs. 31-33; pl. 50, figs. 34-38. Type-locality: El Salvador, La Libertad, w. Ateos. Host — *Pisonia macranthocarpa* J. D. Smith (Nyctaginaceae). Distr. — El Salvador.

Genus PISPONDYLIA Möhn

PISPONDYLIA Möhn, 1960b:344. Type-species, *salvadorensis* Möhn (orig. des.).

SALVADORENSIS Möhn, 1960b:345, pl. 50, figs. 39-48. Type-locality: El Salvador, La Libertad, w. Ateos. Host — *Pisonia macranthocarpa* J. D. Smith (Nyctaginaceae). Distr. — El Salvador.

Genus PLACOCHELA Rübsaamen

PLACOCHELA Rübsaamen, 1916b:4. Type-species, *Epidosis nigripes* Loew (orig. des.).



SERJANIAE Möhn, 1960a:235, pl. 28, figs. 154-162. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Serjanea goniocarpa* Radlk. (Sapindaceae). Distr. — El Salvador.

Genus PROASPHONDYLIA Felt

PROASPHONDYLIA Felt, 1915c:203. Type-species, *brasiliensis* Felt (orig. des.).
BRASILIENSIS Felt, 1915c:203, fig. 6. Type-locality: Brazil, Pernambuco, Bonito. Host — unknown. Distr. — Brazil. Ref. — Möhn, 1962:232, figs. 62-73

Genus SCHIZOMYIA Kieffer

SCHIZOMYIA Kieffer, 1889:183. Type-species, *galiorum* Kieffer (Coquillett, 1910:604).
IPOMOEAE Felt, 1910b:160. Type-locality: St. Vincent. Host — *Ipomoea* sp. (Convolvulaceae). Distr. — St. Vincent.
MANIHOTI Tavares, 1925:22, figs. 3-4, 5a, 6-8; pl. 1, fig. 3. Type-locality: Brazil, Ceará. Host — *Manihot utilissima* Pohl. (Euphorbiaceae). Distr. — Brazil.
STACHYTARPHETAE Barnes, 1932:483. Type-locality: Trinidad. Host — *Stachytarpheta cajanensis* Vahl. (Verbenaceae). Distr. — Trinidad.

Genus STEPHOMYIA Tavares

STEPHOMYIA Tavares, 1916:54. Type-species, *eugeniae* Tavares (orig. des.).
EUGENIAE Tavares, 1916:54, figs. 7 (D, E), 13-14. Type-locality: Brazil, Guanabara, Rio de Janeiro, Botanical Gardens. Host — *Eugenia* sp. (Myrtaceae). Distr. — U.S.A. (Fla.), Brazil. Ref. — Möhn, 1962:234, figs. 74-84.
eugeniae Felt, 1913c:175 (*Cystodiplosis*). Type-locality: U.S.A., Florida, Key West. N. SYN., N. COMB.


UNPLACED GENERA OF ASPONDYLIIDI

Genus ANASPHONDYLIA Blanchard

ANASPHONDYLIA Blanchard, 1939:173 (preocc. Tavares, 1920). Type-species, *aspidospermae* Blanchard (orig. des.). This name, a junior homonym of *Anasphondylia* Tavares, is not changed here because the genus may prove to be a synonym of another genus of *Asphondyliidi*.
ASPIDOSPERMAE Blanchard, 1939:173, pl. 10, figs. a-e. Type-locality: Argentina, Santa Fé, San Cristobal. Host — *Aspidosperma quebrachoblanco* Schlecht. (Apocynaceae). Distr. — Argentina.

Genus ANASPHONDYLIA Tavares

ANASPHONDYLIA Tavares, 1920a:38. Type-species, *myrtacea* Tavares (orig. des.).



MYRTACEA Tavares, 1920b:38, pl. 3, figs. 10-13. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — sp. of Myrtaceae. Distr. — Brazil. Ref. — Tavares, 1922:25.

Genus METASHONDYLIA Tavares

METASPHONDYLIA Tavares, 1918a:41. Type-species, *squamosa* Tavares (orig. des.).

SQUAMOSA Tavares, 1918a:42, pl. 1, fig. 6; pl. 2, figs. 6, 7, 9, 10. Type-locality: Brazil, Bahia, Salvador. Host — sp. of Malvaceae. Distr. — Brazil.

UNPLACED SPECIES OF ASPHONDYLIDI

CLAVATA Tavares, 1920b: pl. 3, fig. 14 (*Oxasphondylia*). Type-locality: Brazil, Bahia, Salvador. Host — sp. of Myrtaceae. Distr. — Brazil. Ref. — Tavares, 1921:93

ITUENSIS Tavares, 1917b:139, pl. 4, figs. 1-2; fig. A (1-4) (*Oxasphondylia*). Type-locality: Brazil, São Paulo, Salto de Itu. Host — *Porophyllum* sp. (Compositae). Distr. — Brazil.

Supertribe LASIOPTERIDI

Genus ALYCAULUS Rübsaamen

ALYCAULUS Rübsaamen, 1916a:476. Type-species, *mikaniae* Rübsaamen (mon.).

MIKANIAE Rübsaamen, 1916a:476, figs. 55-60. Type-locality: Peru, Madre de Dios, nr. Auristella on Rio Acre. Host — *Mikania* sp. (Compositae). Distr. — Peru.

TRILOBATA Möhn, 1964b:583, pl. 10, figs. 145-146. Type-locality: El Salvador, Cuscatlan, ne. Cojutepeque. Host — *Mikania scandens* Willd. (Compositae). Distr. — El Salvador.

Genus ASTEROMYIA Felt

ASTEROMYIA Felt, 1910c:348. Type-species, *Cecidomyia carbonifera* Felt (orig. des.) = *euthamiae* Gagné.

GUTIERREZIAE Felt, 1916:179. Type-locality: U.S.A., Utah, Salt Lake City. Hosts — *Gutierrezia microcephala* (DC.) Gray, *G. sarothrae* (Pursh.) Britt. & Rusby, and *Gymnosperma glutinosa* Less. (Compositae). Distr. — U.S.A. (southwestern), Mexico (unpub.).

Genus BACCHAROMYIA Tavares

BACCHAROMYIA Tavares, 1917b:129. Type-species, *ramosina* Tavares (orig. des.).

RAMOSINA Tavares, 1917b:130, fig. B (1-6). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Baccharis genistelloides* Pers. (Compositae). Distr. — Brazil.



Genus BRACHYLASIOPTERA Möhn

- BRACHYLASIOPTERA Möhn, 1964b:576. Type-species, *rotunda* Möhn (orig. des.).
- GILBERTIAE Möhn, 1964b:579, pl. 8, fig. 129. Type-locality: El Salvador, La Libertad, nr. Los Chorros. Host — *Gilbertia arborea* (L.) March (Araliaceae). Distr. — El Salvador.
- ROTUNDA Möhn, 1964b:576, pl. 7, figs. 113-117, pl. 8, figs. 118-128. Type-locality: El Salvador, La Libertad, nr. San Diego, e. La Libertad. Host — *Pseudocalymma macrocarpa* (J. D. Sm.) Sandw. (Bignoniaceae). Distr. — El Salvador.

Genus CAMPTONEUROMYIA Felt

- CAMPTONEUROMYIA Felt, 1908b:334. Type-species, *Dasineura virginica* Felt (Coquillett, 1910:518).
- MERIDIONALIS Felt, 1910a:269. Type-locality: British West Indies, St. Vincent. Host — *Ipomoea* sp. (Convolvulaceae), inquiline in galls of *Schizomyia ipomoeae*. Distr. — St. Vincent.

Genus DIALERIA Tavares

- DIALERIA Tavares, 1918b:78. Type-species, *styracis* Tavares (orig. des.).
- STYRACIS Tavares, 1918b:78, pl. 3, fig. 13. Type-locality: Brazil, Bahia, Caitité. Host — *Styrax* sp. (Styracaceae), inquiline in galls of *Styraxdiploisis caetitensis*. Distr. — Brazil.

Genus DOLICHOLABIS Tavares

- DOLICHOLABIS Tavares, 1918b:72. Type-species, *lantanae* Tavares (orig. des.).
- INSIGNIS Tavares, 1922:13, figs. 2-6. Type-locality: Brazil. Host — *Prothium heptaphyllum* (Burseraceae), inquiline in galls of *Dasineura braziliensis*. Distr. — Brazil.
- LANTANAE Tavares, 1918b:73, pl. 3, figs. 1-6; pl. 4, fig. 1. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Lantana* sp. (Verbenaceae), inquiline in galls of *Schismatodiploisis lantanae*. Distr. — Brazil.

Genus EPILASIOPTERA Möhn

- EPILASIOPTERA Möhn, 1964b:573. Type-species, *iresinis* Möhn (orig. des.).
- IRESINIS Möhn, 1964b:573, pl. 6, figs. 98-105; pl. 7, figs. 106-112. Type-locality: El Salvador, La Libertad, Cumbre, Hacienda Chilata. Host — *Iresine calea* (Ibañez) Standley (Amaranthaceae). Distr. — El Salvador.

Genus GERALDESIA Tavares

- GERALDESIA Tavares, 1917b:134. Type-species, *eupatorii* Tavares (orig. des.).
EUPATORIUM Tavares, 1917b:134, fig. C (1-5). Type-locality: Brazil, Rio de Janeiro. Host — *Eupatorium* sp. (Compositae). Distr. — Brazil.

Genus ISOLASIOPTERA Möhn

- ISOLASIOPTERA Möhn, 1964b:584. Type-species, *eupatoriensis* Möhn (orig. des.).
EUPATORIENSIS Möhn, 1964b:585, pl. 10, figs. 147-153; pl. 11, figs. 154-162. Type-locality: El Salvador, Sonsonate, forest, km 48 on road to Sonsonate. Hosts — *Eupatorium morifolium* Mill. and *Eupatorium* sp. (Compositae). Distr. — El Salvador.

Genus LOBOLASIOPTERA Möhn

- LOBOLASIOPTERA Möhn, 1964b:580. Type-species, *media* Möhn (orig. des.).
MEDIA Möhn, 1964b:581, pl. 9, figs. 130-142; pl. 10, figs. 143-144. Type-locality: El Salvador, La Libertad, e. La Libertad, nr. San Diego. Host — *Pseudocalymma macrocarpa* (J. D. Sm.) Sandw. (Bignoniaceae). Distr. — El Salvador.

Genus NEOLASIOPTERA Felt

- Meunieria* Rübsaamen, 1905:137 (preocc. Kieffer, 1904). Type-species, *dalechampia* Rübsaamen (orig. des., as gen. n., sp. n.).
NEOLASIOPTERA Felt, 1908b:330. Type-species, *Lasioptera vitinea* Felt (Coullett, 1910b:575).
Meunieriella Kieffer, 1909:35 (nom. nov. for *Meunieria* Rübsaamen). Type-species, *Meunieria dalechampia* Rübsaamen (aut.). N. SYN.
Luisieria Tavares, 1922:43. Type-species, *fariae* Tavares (orig. des.). Most species in this genus form galls on stems, petioles or leaf midveins on a wide array of plants. A few are found in the seeds of Compositae.

Subgenus NEOLASIOPTERA Felt

- AESCHYNOMENSIS (Brèthes), 1918:1, figs. 1-2 (*Lasioptera*). Type-locality: Argentina, Buenos Aires. Host — *Aeschynomene montevidensis* Vog. (Leguminosae). Distr. — Argentina. N. COMB.
AMARANTHI Möhn, 1964a:47, pl. 1, figs. 1-14. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Amaranthus spinosus* L. (Amaranthaceae). Distr. — El Salvador.
APHELANDRAE Möhn, 1964a:90, pl. 16, figs. 220-221. Type-locality: El Salvador, La Libertad, n. El Cimarón. Host — *Aphelandra deppeana* Cham. & Schlecht. (Acanthaceae). Dist. — El Salvador.



- BORRERIAE** Möhn, 1964a:93, pl. 17, figs. 230-238. Type-locality: El Salvador, San Salvador, c. San Marcos. Host — *Borreria verticillata* (L.) Meyer (Rubiaceae). Distr. — El Salvador.
- BRICKELLIAE** Möhn, 1964b:560, pl. 3, figs. 42-52. Type-locality: El Salvador, Chalatenango, ca. 15 kms. from La Palma. Host — *Brickellia pacayensis* Coult. (Compositae). Distr. — El Salvador.
- CALEAE** Möhn, 1964b:556, pl. 2, figs. 15-29. Type-locality: El Salvador, La Libertad, nr. Los Chorros. Host — *Calea integrifolia* (DC.) Hemsl. (Compositae). Distr. — El Salvador.
- GAMARAE** Möhn, 1964b:565, pl. 4, figs. 66-71; pl. 5, figs. 72-73. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Lantana camara* L. (Verbenaceae). Distr. — El Salvador.
- CAPSICI** Möhn, 1964a:78, pl. 12, figs. 161-168; pl. 13, figs. 169-170. Type-locality: El Salvador, La Libertad, nr. San Diego, e. La Libertad. Host — *Capsicum baccatum* L. (Solanaceae). Distr. — El Salvador.
- CELTIS** Möhn, 1964b:553, pl. 1, figs. 1-13; pl. 2, fig. 14. Type-locality: El Salvador, Santa Ana, nr. San Diego. Host — *Celtis iguanea* (Jacq.) Sarg. (Urticaceae). Distr. — El Salvador.
- CEREI** (Rübsaamen), 1905:81, (*Lasioptera*). Type-locality: Brazil, Rio de Janeiro, Cabo Frio. Host — *Cereus setaceus* Salm-Dyck (Cactaceae). Distr. — Brazil. N. COMB.
- CESTRI** Möhn, 1964a:80, pl. 13, figs. 171-180; pl. 14, figs. 181-184. Type-locality: El Salvador, Santa Ana, nr. San Jose, Hacienda San Jose. Host — *Cestrum aurantiacum* Lindl. and *C. lanatum* Mart. & Gal. (Solanaceae).
- CISSAMPELI** Möhn, 1964a:55, pl. 4, figs. 44-54. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Cissampelos pareira* L. (Menispermaceae). Distr. — El Salvador.
- CLEMATICOLA** Möhn, 1964a:52, pl. 3, figs. 23-43. Type-locality: El Salvador, Santa Ana, nw. Ciudad Arce. Host — *Clematis dioica* L. (Ranunculaceae). Distr. — El Salvador.
- COMBRETI** Möhn, 1964a:68, pl. 9, figs. 112-120; pl. 10, figs. 121-124. Type-locality: El Salvador, La Libertad, n. El Cimarón. Host — *Combretum farinosum* H. B. & K. (Combretaceae). Distr. — El Salvador.
- COMPOSITARUM** Möhn, 1964b:563, pl. 4, figs. 53-65. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Hymenostephium cordatum* Blake and *Tridax procumbens* L. (Compositae). Distr. — El Salvador.
- CORDIAE** Möhn, 1964a:71, pl. 10, figs. 127-133; pl. 11, fig. 134. Type-locality: El Salvador, La Libertad, se. Ciudad Arce. Host — *Cordia cana* Mart. & Gal. (Boraginaceae). Distr. — El Salvador.
- DALECHAMPIAE** (Rübsaamen), 1905:13 (*Meunieria*). Type-locality: Brazil, Rio de Janeiro, Palmeiras. Host — *Dalechampia ficifolia* Lam. (Euphorbiaceae). Distr. — Brazil. N. COMB.
- DENTATA** Möhn, 1964b:567, pl. 5, figs. 74-75. Type-locality: El Salvador, Santa Ana, nr. San José. Host — *Eupatorium* sp. (Compositae). Distr. — El Salvador.
- ERIGERONTIS** (Felt), 1907b:163 (*Choristoneura*). Type-locality: U.S.A., New York, Albany. Host — *Erigeron canadensis* L. (Compositae). Distr. — U.S.A., El Salvador. Ref. — Möhn, 1964b:568, pl. 5, figs. 76-88.

- ERYTHRONYLI Möhn, 1964a:62, pl. 7, figs. 86-94. Type-locality: El Salvador, Sonsonate, n. Los Cobanos. Host — *Erythroxyllum mexicanum* H. B. & K. (Lineaceae). Distr. — El Salvador.
- ENIGUA Möhn, 1964a:82, pl. 14, figs. 185-191; pl. 15, figs. 192-193. Type-locality: El Salvador, La Libertad, n. El Cimarrón. Host — *Solanum umbellatum* Mill. (Solanaceae). Distr. — El Salvador.
- FARIAE (Tavares), 1922:44, figs. 11-16 (*Luisieria*). Type-locality: Brazil, Bahia, Salvador. Host — undet. plant. Distr. — Brazil.
- GRACILFORCEPS (Kieffer & Jörgensen), 1910:429, figs. 53-54 (*Lasioptera*). Type-locality: Argentina. Host — *Prosopis strombulifera* Benth. (Leguminosae). Distr. — Argentina. N. COMB.
- GRANDIS Möhn, 1964a:87, pl. 16, figs. 209-210. Type-locality: El Salvador, San Vicente, forest, km 81 nr. Rio San Felipe. Host — *Pseudocalymma macrocarpa* (J. D. Sm.) Sandw. (Bignoniaceae). Distr. — El Salvador.
- HELIOCARPI Möhn, 1964a:66, pl. 9, fig. 109. Type-locality: El Salvador, La Libertad, e. San Juan Higinio. Host — *Heliocarpus donnellsmithii* Rose (Tiliaceae). Distr. — El Salvador.
- HETEROTHALAMI (Kieffer & Jörgensen), 1910:399, figs. 28-30 (*Lasioptera*). Type-locality: Argentina, Mendoza, La Paz; San Juan, Cancete (n. Mendoza). Host — *Heterothalamus spartioides* Hook. & Arn. (Compositae). N. COMB.
- HYPTIS Möhn, 1964a:74, pl. 11, figs. 137-150; pl. 12, fig. 151. Type-locality: El Salvador, Santa Ana, nw. Ciudad Arce. Host — *Hyptis suaveolens* (L.) Poit. (Labiatae). Distr. — El Salvador.
- INDIGOFERAE Möhn, 1964a:60, pl. 6, figs. 70-83; pl. 7, figs. 84-85. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Indigofera suffruticosa* Mill. (Leguminosae). Distr. — El Salvador.
- INGAE Möhn, 1964a: 57, pl. 4, figs. 55-58; pl. 5, figs. 59-69. Type-locality El Salvador, La Libertad, Cumbre, se. San Julian, Hacienda San Julian. Host — *Inga leptoloba* Schlecht. & *I. spuria* H. & B. (Leguminosae). Distr. — El Salvador.
- INTERRUPTA (Kieffer & Jörgensen), 1910:375, fig. 10 (*Lasioptera*). Type-locality: Argentina, Mendoza, nr. Chacras de Coria (11 km s. Mendoza). Host — *Baccharis "subulata" Hook* (Compositae). Distr. — Argentina. N. COMB.
- IRESINIS Möhn, 1964a:50, pl. 2, figs. 15-28. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Iresine celosia* L. (Amaranthaceae). Distr. — El Salvador.
- LAPALMAE Möhn, 1964a:94, pl. 18, figs. 239-252. Type-locality: El Salvador, Chalatenango, se. La Palma. Host — *Calea zacatechichi* Schlecht. (Compositae). Distr. — El Salvador.
- MARTELLI Nijveldt, 1967:125, figs. 1-10. Type-locality: Mexico, Mexico, Chapingo. Host — *Agave sp.* (Agavaceae). Distr. — Mexico.
- MELANTHERAE Möhn, 1964a:97, pl. 18, figs. 253-254. Type-locality: El Salvador, Consonate, nv. Salcoatitán. Host — *Melanthera nivea* (L.) Small (Compositae). Distr. — El Salvador.
- MERREMIAE Möhn, 1964a:70, pl. 10, figs. 125-126. Type-locality: El Salvador, La Libertad, nr. San Diego, e. La Libertad. Host — *Merremia umbellata* (L.) Hallier (Convolvulaceae). Distr. — El Salvador.
- MONTICOLA (Kieffer & Herbst), 1909:124, fig. 4 (*Lasioptera*). Type-locality: Chile, "Pennaloben" (Peñalolén). Host — *Asteriscium polycephalum* (Clos) Gill. & Hook. (Umbelliferae). Distr. — Chile.



- ODONTONEMAE Möhn, 1964a:91, pl. 17, figs. 222-229. Type-locality: El Salvador, Sonsonate, nr. Sonsonate, between Armenia and Izalco. Host — *Odontonema "stricta" (Nees) Kuntze* (Acanthaceae). Distr. — El Salvador.
- ORNATICORNIS (Kieffer & Jörgensen), 1910:368, figs. 5, 5a, 5b (*Lasioptera*). Type-locality: Argentina, Mendoza, nr. Chacras de Coria (11 km. s. Mendoza). Host — *Baccharis salicifolia* Pers. (Compositae). Distr. — Argentina. N. COMB.
- PARVULA Möhn, 1964b:558, pl. 2, figs. 30-35; pl. 3, figs. 36-41. Type-locality: El Salvador, La Libertad, Boquerón Volcano, 1650 m. Host — *Calea integrifolia* (DC.) Hemsl. (Compositae). Distr. — El Salvador.
- PIRIQUETA (Felt), 1917:193 (*Lasioptera*). Type-locality: Puerto Rico, Mayaguez. Host — *Piriqueta ovata* Urban (Turneraceae). Distr. — Puerto Rico. N. COMB.
- PORTULACAE (Cook), 1906:251 (*Cecidomyia*). Type-locality: Cuba. Host — *Portulaca oleracea* L. (Portulacaceae). Distr. — Cuba, St. Vincent. N. COMB.
- portulacae* Felt, 1911j:84 (*Lasioptera*). Type-locality: St. Vincent.
- PSEUDOCALYMMAE Möhn, 1964a:84, pl. 15, figs. 194-208. Type-locality: El Salvador, La Libertad, nr. San Diego, e. La Libertad. Host — *Pseudocalymma macrocarpa* (J. D. Sm.) Sandw. (Bignoniaceae). Distr. — El Salvador.
- SALVADORENSIS Möhn, 1964a:88, pl. 16, figs. 211-219. Type-locality: El Salvador, San Miguel, s. El Delirio. Host — sp. of Bignoniaceae. Distr. — El Salvador.
- SALVIAE Möhn, 1964a:76, pl. 12, figs. 152-160. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Salvia occidentalis* Sw. (Labiatae). Distr. — El Salvador.
- SENECIONIS Möhn, 1964a:98, pl. 19, figs. 255-269. Type-locality: El Salvador, La Libertad, n. El Cimarrón. Host — *Senecio kermesinus* Hemsl. (Compositae). Distr. — El Salvador.
- SERJANIAE Möhn, 1964a:64, pl. 7, figs. 95-96; pl. 8, figs. 97-108. Type-locality: El Salvador, La Libertad, nw. Quezaltepeque, La Toma forest. Host — *Serjanea racemosa* Schum. (Sapindaceae). Distr. — El Salvador.
- SIDAE Möhn, 1964a:67, pl. 9, figs. 110-111. Type-locality: El Salvador, Chalatenango, nr. Citalá, on bank of Lempa R. Host — *Sida rhombifolia* L. (Moraceae). Distr. — El Salvador.
- TOURNEFORTIAE Möhn, 1964a:73, pl. 11, figs. 135-136. Type-locality: El Salvador, La Libertad, ne. Sitio del Niño. Host — *Tournefortia volubilis* L. (Boraginaceae). Distr. — El Salvador.
- TRIDENTIFERA (Kieffer & Jörgensen), 1910:398, fig. 27 (*Lasioptera*). Type-locality: Argentina, Mendoza, La Paz; San Juan, Cancete (n. Mendoza). Host — *Heliotropium curassavicum* L. (Boraginaceae). Distr. — Argentina. N. COMB.
- URVILLEAE (Tavares), 1909:25, pl. 3, figs. 3-4 (*Lasioptera*). Type-locality: Brazil, Rio Grande do Sul, vic. São Leopoldo. Host — *Urvillea ulmacea* H. B. & K. (Sapindaceae). Distr. — Brazil. N. COMB.
- VERBESINAE Möhn, 1964:100, pl. 19, figs. 270-272; pl. 20, figs. 273-283. Type-locality: El Salvador, Chalatenango, nr. La Palma. Hosts — *Verbesina fraseri* Hemsl., *V. sublobata* Benth., and *V. turbacensis* H. B. & K. (Compositae). Distr. — El Salvador.

VERNONIENSIS Möhn, 1964a:103, pl. 20, figs. 284-285. Type-locality: El Salvador, La Libertad, se. Ciudad Arce. Host — *Vernonia* sp. (Compositae). Distr. — El Salvador.

Subgenus DILASIOPTERA Möhn

Neolasioptera, subg. DILASIOPTERA Möhn, 1964b:570. Type-species, *serrata* Möhn (orig. des.).

FERRATA Möhn, 1964b:571, pl. 5, figs. 89-91; pl. 6, figs. 92-97. Type-locality: El Salvador, Chalatenango, se. La Palma. Host — *Calea zacatechichi* Schlecht. (Compositae). Distr. — El Salvador.

Genus NEUROLASIOPTERA Brèthes

NEUROLASIOPTERA Brèthes, 1922:138. Type-species, *baezi* Brèthes (mon.). This genus is probably a synonym of *Neolasioptera*.

BAEZI Brèthes, 1922:138. Type-locality: Argentina, Entre Rios, Paraná. Host — *Teucrium inslatum* Sw. (Labiatae). Distr. — Argentina.

Genus PHYSALIDICOLA Brèthes

PHYSALIDICOLA Brèthes, 1917a:240. Type-species, *argentata* Brèthes (orig. des.). This genus is probably a synonym of *Neolasioptera*.

ARGENTATA Brèthes, 1917a:240, figs. 1-3. Type-locality: Argentina, Buenos Aires, Buenos Aires. Host — *Physalis viscosa* L. (Solanaceae). Distr. — Argentina.

Genus TROTTERIA Kieffer

Choristoneura Rübsaamen, 1892:342 (preocc. Lederer, 1859). Type-species, *Cecidomyia obtusa* Loew (mon.).

TROTTERIA Kieffer, 1901:561 (nom. nov. for *Choristoneura* Rübsaamen). Type-species, *Cecidomyia obtusa* Loew (aut.).

CARYAE (Felt), 1907a:14 (*Choristoneura*). Type-locality: U.S.A., New York, Albany. Host — unknown. Distr. — U.S.A., Dominica (unpub.).

LINDNERI Möhn, 1963a:1, pl. 1, figs. 1-8. Type-locality: El Salvador, San Salvador, San Salvador. Host — *Indigofera suffruticosa* (Leguminosae), inquiline in galls of *Asphondylia indigoferae*. Distr. — El Salvador.

SALVADORENSIS Möhn, 1963a:3, pl. 1, figs. 9-17. Type-locality: El Salvador, San Vicente, forest nr. Rio San Felipe, km 81. Host — *Ayenia pusilla* (Sterculiaceae) and *Boerhaavia erecta* (Nyctaginaceae), inquiline in galls of *Asphondylia ayeniae* and *A. boerhaaviae*, respectively. Distr. — El Salvador.

UNPLACED SPECIES OF LASIOPTERIDI

Adults unknown

CORDOBENSIS Kieffer & Jörgensen, 1910:363, fig. 1 (*Lasioptera*). Type-locality: Argentina, Cordoba, nr. Arias and La Carlota. Host — *Baccharis coridifolia* DC. (Compositae). Distr. — Argentina.



Supertribe CECIDOMYIIDI

Tribe ANADIPLOSINI

All known members of this tribe cause globular, hairy, leaf or stem galls.

Genus ANADIPLOSIS Tavares

- ANADIPLOSIS Tavares, 1916:41. Type-species, *pulchra* Tavares (orig. des.).
CAETETENSIS Tavares, 1920c:70, fig. 5. Type-locality: Brazil, Bahia, Caitité.
Host — a sp. of Leguminosae. Distr. — Brazil.
PROCERA Tavares, 1920c:66, fig. 4; pl. 1, figs. 1-6, 11-12, 17. Type-locality:
Brazil, Bahia, Salvador, Itaparica I. Host — *Mimosa* sp. (Legu-
minosae). Distr. — Brazil.
PULCHRA Tavares, 1916:42, figs. 3-6. Type-locality: Brazil, Rio de Janeiro,
Nova Friburgo. Host — *Mimosa* sp. (Leguminosae). Distr. —
Brazil.
VENUSTA Tavares, 1916:46, fig. 7(A-B)-9. Type-locality: Brazil, Rio de
Janeiro, Nova Friburgo. Host — *Machaerium* sp. (Leguminosae).
Distr. — Brazil.

Genus RUEBSAAMENODIPLOSIS Tavares

- RUEBSAAMENODIPLOSIS Tavares, 1920c:60-61. Type-species, *Uleela machaerii*
Kieffer (orig. des.).
MACHAERI (Kieffer), 1913a:101 (*Uleela*). Type-locality: "Brazil". Host —
Machaerium sp. (Leguminosae). Distr. — Brazil. Ref. — Rüb-
saamen, 1908a:120, fig. 2.

Genus ULEELA Rübsaamen

- ULEELA Rübsaamen, 1908a:120. Type-species, *dalbergiae* Rübsaamen (orig.
des., as gen. n., sp. n.).
DALBERGIAE Rübsaamen, 1908a:121, fig. 1. Type-locality: Brazil, Guanabara,
Rio de Janeiro, Jacarepaguá. Host — *Dalbergia* sp. (Legumi-
nosae). Distr. — Brazil.
DEPRESSA Kieffer, 1913a:101. Type-localities: Brazil, Amazonas, Belém and
Juruá Miri; Guanabara, Rio de Janeiro, Floresta da Tijuca; Santa
Catarina, Minas and Teresópolis. Host — *Polypodium crassifo-
lium* L. (Polypodiaceae). Distr. — Brazil. Ref. — Rübsaamen,
1908b:48.
GLOBULIFEX Kieffer, 1913a:101. Type-locality: Brazil, Rio de Janeiro, Serra
dos Órgãos. Host — *Neea* sp. (Nyctaginaceae). Distr. — Brazil.
Ref. — Rübsaamen, 1908b:25.
LIGNICOLA Kieffer, 1913a:101. Type-locality: Brazil, Rio de Janeiro, Serra
do Macaé. Host — *Dalbergia* sp. (Leguminosae). Distr. — Brazil.
Ref. — Rübsaamen, 1905:133.
LONGICAUDA Kieffer, 1913a:101. Type-locality: Brazil, Guanabara, vic. Rio
de Janeiro. Host — *Neea* sp. (Nyctaginaceae). Distr. — Brazil.
Ref. — Rübsaamen, 1908b:26.

- LONGISETA Kieffer, 1913a:101. Type-locality: Brazil, Amazonas, Marari, Juruá. Host — *Neea* sp. (Nyctaginaceae). Distr. — Brazil. Ref. — Rübssaamen, 1908b:23.
- MICRURA Kieffer, 1913a:101. Type-locality: Brazil, Santa Catarina. Host — *Neea* sp. (Nyctaginaceae). Distr. — Brazil. Ref. — Rübssaamen, 1908b:26.
- NEEANA Kieffer, 1913a:101. Type-locality: Brazil. Host — *Neea* sp. (Nyctaginaceae). Distr. — Brazil. Ref. — Rübssaamen, 1908a:121, fig. 3.
- RUEBSAAMENI Kieffer, 1913a:101. Type-locality: Brazil, Santa Catarina, Pedras Grandes. Host — *Dalbergia* sp. (Leguminosae). Distr. — Brazil. Ref. — Rübssaamen, 1905:133.

Tribe APHIDOLETINI

Genus THRIPSOBREMIA Barnes

- THRIPSOBREMIA Barnes, 1930:331. Type-species, *liothrips* Barnes (mon.)
- LIOTHIRIPS Barnes, 1930:331, fig. 1. Type-locality: Trinidad. Host — *Liothrips urichi* Karny (Insecta: Thysanoptera: Phlaeothripidae). Distr. — Trinidad.

Tribe CLINODIPLOSINI

Genus CLÉITODIPLOSIS Tavares

- CLÉITODIPLOSIS Tavares, 1921:108. Type-species, *Necrophlebia graminis* Tavares (mon.).
- GRAMINIS (Tavares), 1916:37, figs. 1-2 (*Necrophlebia*). Type-locality: Brazil, Bahia and Rio de Janeiro. Host — *Paspalum conjugatum* Berg. (Gramineae). Distr. — Brazil. Ref. — Tavares, 1921:109, figs. 5(1-2)-6.

Genus CLINODIPLOSIS Kieffer

- CLINODIPLOSIS Kieffer, 1895b:cclxxx. Type-species, *Diplosis cilicrus* Kieffer (orig. des.).
- CHLOROPHORAE Rübssaamen, 1905:82. Type-locality: Brazil, Rio de Janeiro, Fábrica. Host — *Chlorophora tinctoria* Gaudich. (Urticaceae). Distr. — Brazil.

Genus HOUARDODIPLOSIS Tavares

- HOUARDODIPLOSIS Tavares, 1925:13. Type-species, *rochae* Tavares (orig. des.).
- ROCHAE Tavares, 1925:14, figs. 1-2, pl. 1, fig. 4. Type-locality: Brazil, Ceará. Host — *Combretum leprosum* Mart. (Combretaceae). Distr. — Brazil.



Genus HYPERDIPLOSIS Felt

- HYPERDIPLOSIS Felt, 1908b:405. Type-species, *Cecidomyia lobata* Felt (orig. des.).
- AMERICANA Felt, 1911i:192. Type-locality: Panama, Canal Zone, Paraiso. Host — reared from dead *Ficus* sp. (Moraceae). Distr. — Panama.
- COFFEEAE Felt, 1911d:305. Type-locality: St. Vincent. Host — *Coffea libERICA* Bull (Rubiaceae). Distr. — St. Vincent.
- EUPATORIUM Felt, 1911b:110. Type-locality: St. Vincent. Host — *Eupatorium* sp. (Compositae). Distr. — St. Vincent.
- PRODUCTA Felt, 1912a:177. Type-locality: St. Vincent. Host — *Stachytarpheta dichotoma* Vahl. (Verbenaceae). Distr. — St. Vincent.

Genus IATROPHOBIA Rübsaamen

- IATROPHOBIA Rübsaamen, 1916a:469. Type-species, *Clinodiplosis brasiliensis* Rübsaamen (mon.).
- Jatrophia* authors, unjustified emend.
- BRASILIENSIS (Rübsaamen), 1908a:156 (*Clinodiplosis*). Type-localities: Brazil: Amazonas, Fortaleza and Juruá Miri; Rio de Janeiro, Mauá and Palmeiras; Santa Catarina, Tubarão. Hosts — *Manihot aipi* Pohl, *M. dichotoma* Ule, and *M. utilisissima* Pohl (Euphorbiaceae). Distr. — St. Vincent, Tobago, Trinidad, Costa Rica (unpub.), Guyana, Surinam and Brazil.
- manihot* Felt, 1910a:268 (*Cecidomyia*). Type-locality: St. Vincent.

Genus PARALLELODIPLOSIS Rübsaamen

- PARALLELODIPLOSIS Rübsaamen, 1910:287 (also 1911:120). Type-species, *Diplosis galliperda* Löw (mon.).
- CATTLEYAE (Molliard), 1903:165, pl. 2, figs. 1-18 (*Cecidomyia*). Type-locality: Brazil. Hosts — *Cattleya* spp. and *Laelia* spp. (Orchidaceae). Distr. — U.S.A., Mexico (unpub.), Guatemala, Costa Rica, Columbia and Brazil.
- cattleyae* Felt, 1908b:412 (*Clinodiplosis*). Type-locality: U.S.A., Massachusetts, South Natick.

Genus SCHISMATODIPLOSIS Rübsaamen


- Eudiplosis* Tavares, 1916:51 (preocc. Kieffer, 1894). Type-species, *parva* Tavares (orig. des.).
- SCHISMATODIPLOSIS Rübsaamen, 1916a:467. Type-species, *Clinodiplosis lantanae* Rübsaamen (mon.).
- Autodiplosis* Tavares, 1920b:106 (nom. nov. for *Eudiplosis* Tavares). Type-species, *Eudiplosis parva* Tavares (aut.). N. SYN.
- BAHIENSIS (Tavares), 1917b:145, pl. 6, figs. 1-2; pl. 8, figs. 1-2; pl. 9, figs. 1-4, 9-11 (*Eudiplosis*). Type-locality Brazil, Bahia, Salvador. Host — sp. of Compositae. Distr. — Brazil. N. COMB.



- CEARENSIS (Tavares), 1917b:153, pl. 9, figs. 5-7 (*Eudiplosis*). Type-locality: Brazil, Ceará, Fortaleza. Host — unknown. Distr. — Brazil. N. COMB.
- IHERINGI (Tavares), 1925:45, figs. 13-14 (*Autodiplosis*). Type-locality: Brazil, Santa Catarina, Joinville. Host — *Aegiphila arborscens* Vahl. (Verbenaceae). Distr. — Brazil. N. COMB.
- LANTANAE (Rübsaamen), 1908a:151 (*Clinodiplosis*). Type-locality: Brazil, Rio de Janeiro, Cabo Frio and Palmeiras; Santa Catarina, Tubarão. Host — *Lantana* sp. (Verbenaceae). Distr. — Brazil.
- MARCEITAE (Tavares), 1917a:22, figs. 1-5 (*Eudiplosis*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Marcetia* sp. (Melastomataceae). Distr. — Brazil. N. COMB.
- PARVA (Tavares), 1916:51, fig. 7(C only), 10 (*Eudiplosis*). Type-locality: Brazil, Bahia, Salvador. Host — sp. of Papilionaceae. Distr. — Brazil. N. COMB.
- PULCHRA (Tavares), 1917b:pl. 10, figs. 3-7; pl. 11, fig. 6 (*Eudiplosis*). Type-locality: Brazil, Bahia, Salvador. Host — *Lantana* sp. (Verbenaceae). Distr. — Brazil. Ref. — Tavares, 1918b:31. N. COMB.
- RUBIAE (Tavares), 1918a:70, pl. 3, fig. 12 (*Eudiplosis*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — sp. of Rubiaceae. Distr. — Brazil. N. COMB.
- Unnamed sp., Tavares, 1918a:64, figs. 1-2; pl. 1, fig. 8. Host — *Solanum* sp. (Solanaceae). Distr. — Brazil, Bahia, vic. Salvador.

Tribe CONTARINIINI
Genus CONTARINIA Rondani

- Cecidomyia*, subg. CONTARINIA Rondani, 1860:289. Type-species, *Tipula loti* De Geer (orig. des.).
- GOSSYPI Felt, 1908a:210. Type-locality: Antigua. Hosts — *Gossypium* spp. (Malvaceae). Distr. — U.S.A. (Fla.), Tortola, St. Croix, Antigua, Montserrat and Barbados. Ref. — Barnes, 1949:94.
- LYCOPERSICI Felt, 1911d:303. Type-locality: St. Vincent. Host — *Lycopersicon esculentum* Mill. (Solanaceae). Distr. — Dominica, St. Lucia, St. Vincent, Barbados, Grenada, British Honduras and Guyana. Also, U.S.A. (Hawaii). Ref. — Barnes, 1946a:78.
- MEDICAGINIS Kieffer, 1895c:150. Type-locality: France. Host — *Medicago sativa* L. (Leguminosae). Distr. — Peru (unpub.), Europe. Ref. — Barnes, 1946b:43.
- SORGHICOLA (Coquillett), 1899:81 (*Diplosis*). Type-locality: U.S.A., Texas, College Station. Hosts — *Sorghum* spp. Distr. — U.S.A. (Calif. to Va, s. to Fla., and Hawaii), Dominican Republic (unpub.), Puerto Rico, Virgin Is (U.S.), St. Vincent, Curaçao, Trinidad, Venezuela, Brazil (unpub.) and Argentina. Also Africa, Aden, Indonesia and Australia. Ref. — Harris, 1964.
- palposa* Blanchard, 1958:423, figs. a-g. Type-locality: Argentina.



Genus SPHAERODIPLOSIIS Rübssaamen

- SPHAERODIPLOSIIS Rübssaamen, 1916a:461. Type-species, *dubia* Rübssaamen (mon.).
- DUBIA Rübssaamen, 1916a:461, figs. 35-37. Type-locality: Brazil. Host — undet. shrub. Distr. — Brazil.

Tribe HORMOMYIINI

Genus EPIHORMOMYIA Felt

- EPIHORMOMYIA Felt, 1915a:155. Type-species, *auripes* Felt (mon.).
- AURIPES Felt, 1915a:156. Type-locality: Guyana, Demerara, "Mallat". Host — unknown. Distr. — Guyana.

Tribe LESTODIPLOSINI

Genus ARTHROCNO DAX Rübssaamen

- ARTHROCNO DAX Rübssaamen, 1895:189. Type-species, *vitis* Rübssaamen (Coquillett, 1910b:510).
- ABDOMINALIS (Felt), 1911c:128 (*Endaphis*). Type-locality: Peru, Piura, Piura. Host — undet. Acarina. Distr. — Peru.
- CONSTRUCTUS Felt, 1914b:481. Type-locality: Puerto Rico, Rio Piedras. Host — "*Tetranychus bimaculatus*" (Acarina: Tetranychidae). Distr. — Puerto Rico.
- INSULARIS (Felt), 1913a:305 (*Mycodiplosis*). Type-locality: Puerto Rico, Rio Piedras. Host — *Tetranychus tumidus* Banks (Acarina: Tetranychidae). Distr. — Puerto Rico. N. comb.
- MERIDIONALIS Felt, 1912a:176. Type-locality St. Vincent. Host — *Eriophyes* sp. (Acarina: Eriophyidae). Distr. — St. Vincent.
- Unnamed sp., Wolcott, 1936:332 (misident. as *macrofilus* Felt). Host — found on mite infested cattle feed. Distr. — Puerto Rico, Rio Piedras.

Genus LESTODIPLOSIIS Kieffer

- LESTODIPLOSIIS Kieffer, 1894a:84. Type-species, *Cecidomyia pictipennis* Ferris (Kieffer, 1895b:cclxxx).
- Coprodiplosis* Kieffer, 1894a:84. Type-species, *cryphali* Kieffer (mon.).
- BRAZILIENSIS (Tavares), 1920b:113 (*Coprodiplosis*). Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Mimosa* sp. (Leguminosae), predator or inquiline in galls of *Anadiplosis pulchra* Tavares. Distr. — Brazil.
- PERUVIANA Felt, 1911a:10. Type-locality: Peru, Piura, Piura. Host — *Pinnaspis temporaria* Ferris (Insecta: Homoptera: Diaspididae). Distr. — Peru.
- PICTURATA Felt, 1915a:156. Type-locality: Guyana, Essequibo, Bartica. Host — Unknown. Distr. — Guyana.



UNPLACED GENERA OF CECIDOMYIIDI

Genus ALEXOMYIA Felt

ALEXOMYIA Felt, 1921a:141. Type-species, *ciliata* Felt (orig. des.).

CILIATA Felt, 1921a:141. Type-locality: Brazil, Pará, Prata. Host — unknown. Distr. — Brazil.

Genus ALLODIPLOSIS Kieffer & Jörgensen

ALLODIPLOSIS Kieffer & Jörgensen, 1910:389. Type-species, *crassus* Kieffer & Jörgensen (mon.).

CRASSUS Kieffer & Jörgensen, 1910:389, figs. 20-24. Type-locality Argentina, Mendoza. Host — *Gourliea decorticans* Gill. (Leguminosae). Distr. — Argentina.

Genus ANDIRODIPLOSIS Tavares

ANDIRODIPLOSIS Tavares, 1920b:93. Type-species, *bahiensis* Tavares (mon.).

BABIENSIS Tavares, 1920b:94, pl. 3, figs. 1-2. Type-locality: Brazil, Bahia, Salvador. Host — *Andira* sp. (Leguminosae). Distr. — Brazil.

Genus APODIPLOSIS Tavares

APODIPLOSIS Tavares, 1922:37. Type-species, *praecox* Tavares (orig. des.).

PRAECOX Tavares, 1922:35, figs. 7-8. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Psychotria* sp. (Rubiaceae). Distr. — Brazil.

Genus ASTRODIPLOSIS Felt

ASTRODIPLOSIS Felt, 1913b:217. Type-species, *speciosa* Felt (orig. des.).

SPECIOSA Felt, 1913b:218. Type-locality: Guatemala, Puerto Barrios. Host — undet. vine. Distr. — Guatemala.

Genus AUSTRODIPLOSIS Brèthes

AUSTRODIPLOSIS Brèthes, 1914:154. Type-species, *argentina* Brèthes (orig. des.).

ARGENTINA Brèthes, 1914:154, fig. 4. Type-locality: Argentina, Buenos Aires, "General Urquiza". Host — unknown. Distr. — Argentina.

Genus CENTRODIPLOSIS Kieffer & Jörgensen

CENTRODIPLOSIS Kieffer & Jörgensen, 1910:405. Type-species, *crassipes* Kieffer & Jörgensen (present des.).



CRASSIPES Kieffer & Jörgensen, 1910:406, figs. 34-35. Type-locality: Argentina, Mendoza, nr. Chacras de Coria (11 km s. Mendoza). Host — *Lycium chilense* Bert. (Solanaceae). Distr. — Argentina.

Genus CHARIDIPOLOSIS Tavares

CHARIDIPOLOSIS Tavares, 1918a:80. Type-species, *concinna* Tavares (orig. des.).

CONCINNA Tavares, 1918a:81, pl. 4, figs. 2-12. Type-locality: Brazil, Bahia. Host — unknown. Distr. — Brazil.

Genus CLEODIPOLOSIS Felt

CLEODIPOLOSIS Felt, 1922:1. Type-species, *aleyrodici* Felt (orig. des.).

ALEYRODICI Felt, 1922:1. Type-locality: Panama, Panamá. Host — *Lecanoides giganteus* Quaintance & Baker (Insecta: Homoptera: Aleyrodidae). Distr. — Panama.

Genus COMPSODIPOLOSIS Tavares

COMPSODIPOLOSIS Tavares, 1909:14. Type-species, *luteoalbida* Tavares (mon.).

FRIBURGENSIS Tavares, 1915:150, figs. 3-5; pl. 6, figs. 8-12. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Styrax* sp. (Styracaceae). Distr. — Brazil.

HUMILIS Tavares, 1915:154, figs. 6-7; pl. 6, fig. 1. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Styrax* sp. (Styracaceae). Distr. — Brazil.

ITAPARICANA Tavares, 1922:41, figs. 9-10. Type-locality: Brazil, Bahia, Salvador, Itaparica I. Host — undet. plant. Distr. — Brazil.

LUTEALBIDA Tavares, 1909:14, pl. 2, figs. 2-6. Type-locality: Brazil, Rio Grande do Sul, vic. São Leopoldo. Host — *Smilax* sp. (Liliaceae). Distr. — Brazil.

TRISTIS Tavares, 1915:156, fig. 8; pl. 6, fig. 15. Type-locality: Brazil, Rio de Janeiro, Nova Friburgo. Host — *Styrax* sp. (Styracaceae). Distr. — Brazil.

Genus COQUILLETOMYIA Felt

COQUILLETOMYIA Felt, 1908b:398. Type-species, *Mycodiplosis lobata* Felt (orig. des.).

KNABI Felt, 1912d:154. Type-locality: Costa Rica, Limon, Limon. Host — unknown. Distr. — Costa Rica.

Genus CTENODACTYLOMYIA Felt

CTENODACTYLOMYIA Felt, 1915c:199. Type-species, *watsoni* Felt (orig. des.).

WATSONI Felt, 1915c:199, figs. 2-4. Type-locality: U.S.A., Florida, Palm Beach. Host — *Coccoloba laurifolia* Jacq. (Polygonaceae). Distr. — U.S.A., (Florida), Puerto Rico.



Genus CYSTODIPLOSI Kieffer & Jörgensen

CYSTODIPLOSI Kieffer & Jörgensen, 1910:395. Type-species, *longipennis* Kieffer & Jörgensen (mon.).

LONGIPENNIS Kieffer & Jörgensen, 1910:395, fig. 25. Type-locality: Argentina, Mendoza, Chacras de Corla (11 km s. Mendoza). Host — *Grabowskiya obtusa* Arn. (Solanaceae). Distr. — Argentina.

Genus DACTYLODIPLOSI Rübsaamen

DACTYLODIPLOSI Rübsaamen, 1916a:452. Type-species, *heisteriae* Rübsaamen (mon.).

HEISTERIAE Rübsaamen, 1916a:454, figs. 24-28, 29a, 30-31. Type-localities: Peru, Madre de Dios, Auristilla on Rio Acre; Brazil, Acre, nr. San Francisco on Rio Acre. Host — *Heisteria cyanocarpa* Poepp & Endl. (Olacaceae). Distr. — Brazil, Peru.

Genus DELPHODIPLOSI Felt

DELPHODIPLOSI Felt, 1915a:155. Type-species, *cinctipes* Felt (orig. des.).

CINCTIPES Felt, 1915a:155. Type-locality: Guyana, Essequibo, Bartica. Host — unknown. Distr. — Guyana.

Genus DIADIPLOSI Felt

DIADIPLOSI Felt, 1911h:54. Type-species, *cocci* Felt (orig. des.).

BUSCKI Felt, 1914a:122. Type-locality: Puerto Rico. Host — sp. of Coccidae (Insecta: Homoptera). Distr. — Puerto Rico.

COCCI Felt, 1911h:55. Type-locality: St. Vincent. Host — *Saissetia nigra* (Nietn.) (Insecta: Homoptera: Coccidae).

PSEUDOCOCCI Felt, 1921b:225. Type-locality: Guyana. Host — *Dysmicoccus brevipes* (Ckll.) (Insecta: Homoptera: Pseudococcidae). Distr. — Guyana.

Genus EROSMYIA Felt

EROSMYIA Felt, 1911h:49. Type-species, *mangiferae* Felt (orig. des.).

Mangodiplosis Tavares, 1918a:47. Type-species, *mangiferae* Tavares (orig. des.). N. SYN.

MANGIFERAE Felt, 1911h:49. Type-locality: St. Vincent. Host — *Mangifera indica* L. (Anacardiaceae). Distr. — St. Vincent, Brazil.

mangiferae Tavares, 1918a:48, pl. 2, figs. 1-5 (*Mangodiplosis*). Type-locality: Brazil, Bahia, Salvador. N. SYN., N. COMB.

Genus FRAUENFELDIELLA Rübsaamen

FRAUENFELDIELLA Rübsaamen, 1905:122. Type-species, *coussapoe* Rübsaamen (orig. des., as gen. n., sp. n.).



COUSSAPOAE Rübsaamen, 1905:122. Type-localities: Brazil, Guanabara, Rio de Janeiro, Gávea; Amazonas, Juruá Miri. Host — *Coussapoa* sp. (Urticaceae). Distr. — Brazil.

Genus GNESIODIPLOSIS Tavares

GNESIODIPLOSIS Tavares, 1917b:162. Type-species, *itaparicae* Tavares (orig. des.).

ITAPARICAE Tavares, 1917b:162, pl. 8, figs. 3-4; pl. 9, figs. 2-5. Type-locality: Brazil, Bahia, Salvador, Itaparica I. Host — sp. of Rubiaceae. Distr. — Brazil.

Genus JORGENSENIA Kieffer

JORGENSENIA Kieffer, 1913a:148. Type-species, *Centrodiplosis falcigera* Kieffer & Jörgensen (orig. des.).

FALCIGERA (Kieffer & Jörgensen), 1910:411, figs. 37-39 (*Centrodiplosis*). Type-locality: Argentina, Mendoza. Host — *Lycium gracile* Meyen (Solanaceae). Distr. — Argentina.

Genus KALODIPLOSIS Felt

KALODIPLOSIS Felt, 1915b:229. Type-species, *Dicrodiplosis multifila* Felt (orig. des.).

MULTIFILA (Felt), 1908b:300 (*Dichrodiplosis* [sic]). Type-locality: Puerto Rico. Host — unknown. Distr. — Puerto Rico.

Genus KARSCHOMYIA Felt

KARSCHOMYIA Felt, 1908b:398. Type-species, *Mycodiplosis viburni* Felt (orig. des.).

TOWNSENDI Felt, 1912d:155. Type-locality: Peru, Jaen, Charapa R. Host — unknown. Distr. — Peru.

Genus LIEBELIOLA Kieffer & Jörgensen

LIEBELIOLA Kieffer & Jörgensen, 1910:428. Type-species, *prosopidis* Kieffer & Jörgensen (mon.).

PROSOPIDIS Kieffer & Jörgensen, 1910:428, fig. 52. Type-locality: Argentina, Mendoza and San Juan. Host — *Prosopis strombulifera* Benth. (Leguminosae). Distr. — Argentina.

Genus LOBODIPLOSIS Felt

LOBODIPLOSIS Felt, 1908b:397. Type-species, *Mycodiplosis acerina* Felt (orig. des.).

COCCIDARUM Felt, 1911g:195. Type-locality: St. Vincent. Host — *Planococcus citri* (Risso) (Insecta: Homoptera: Pseudococcidae). Distr. — St. Vincent.



PSEUDOCOCCI Felt, 1933:87. Type-locality: Hawaii, Oahu, Konia. Host — *Dysmicoccus brevipes* (Cockerell) (Insecta: Homoptera: Pseudococcidae). Distr. — Hawaii, Mexico.

SPINOSA Felt, 1909:301. Type-locality: Antigua, St. Johns. Host — unknown. Distr. — Antigua.

Genus LOPESIA Rübsaamen

LOPESIA Rübsaamen, 1908b:29. Type-species, *brasiliensis* Rübsaamen (mon.).

BRASILIENSIS Rübsaamen, 1908b:30, figs. 10-12. Type-localities: Brazil, Guanabara, vic. Rio de Janeiro; Santa Catarina. Host — *Ossaea* sp. (Melastomaceae). Distr. — Brazil.

Genus MACHAERIOBIA Rübsaamen

MACHAERIOBIA Rübsaamen, 1916a:448. Type-species, *brasiliensis* Rübsaamen (mon.).

BRASILIENSIS Rübsaamen, 1916a:448, figs. 18-23. Type-locality: "Brazil". Host — *Machaerium* sp. (Leguminosae). Distr. — Brazil.

Genus MEGAULUS Rübsaamen

MEGAULUS Rübsaamen, 1916a:461. Type-species, *sterculiae* Rübsaamen (mon.).

STERCULIAE Rübsaamen, 1916a:463, figs. 38-42. Type-locality: Brazil, Acre, São Francisco on Rio Acre and Bolivia, Pando, Cobija. Host — *Sterculia* sp. (Sterculiaceae). Distr. — Brazil.

Genus NEOBAEOMYZA Brèthes

NEOBAEOMYZA Brèthes, 1914:151. Type-species, *maculata* Brèthes (orig. des.).

AUSTRALIS Brèthes, 1914:152, fig. 2. Type-locality: Argentina, Buenos Aires. Host — unknown. Distr. — Argentina.


MACULATA Brèthes, 1914:151, fig. 1. Type-locality: Argentina, Buenos Aires. Host — unknown. Distr. — Argentina.

Genus OLEISICOCCUS Borgmeier

OLEISICOCCUS Borgmeier, 1931:186. Type-species, *costalimai* Borgmeier (orig. des.) = *coccidivora* (Felt).

Phagodiplosis Blanchard, 1938:345. Type-species, *haywardi* Blanchard (orig. des.). N. SYN.

COCCIDIVORA (Felt), 1911f:549 (*Mycodiplosis*). Type-locality: Jamaica, Kingston. Hosts — *Pulvinaria urbicola* Ckll. and *Saissetia coffeae* (Walker) (Insecta: Homoptera: Coccidae). Distr. — Jamaica, Brazil. Refs. — Parnell, 1966:51, figs. 1-11; Harris, 1966a:51.



costalimai Borgmeier, 1931:186, figs. 1-10. Type-locality: Brazil, São Paulo.

HAYWARDI (Blanchard), 1938:345, figs. a-f (*Phagodiplosis*). Type-locality: Argentina, Entre Rios, Concordia. Host — *Saissetia coffeae* (Walker). Distr. — Argentina. N. COMB. This species may be a synonym of *O. coccidivora* (Felt).

Genus OURADIPLOSIS Felt

OURADIPLOSIS Felt, 1915a:154. Type-species, *aurata* Felt (orig. des.).

AURATA Felt, 1915a:154. Type-locality: Brazil, Pará, Igarapé-Açu. Host — unknown. Distr. — Brazil.

Genus PLESIODIPLOSIS Kieffer

PLESIODIPLOSIS Kieffer, 1912a:1. Type-species, *Clinodiplosis chilensis* Kieffer & Herbst (orig. des.).

CHILENSIS (Kieffer & Herbst), 1906:231 (*Clinodiplosis*). Type-locality: Chile, Concepción, vic. Concepción. Host — unknown. Distr. — Chile.

Genus PSEUDENDAPHIS Barnes

PSEUDENDAPHIS Barnes, 1954:772. Type-species, *maculans* Barnes (mon.).

MACULANS Barnes, 1954:772. Type-locality: Trinidad. Host — *Toxoptera aurantii* (Fonsc.) (Insecta: Homoptera: Aphididae). Distr. — Trinidad.

Genus ROCHADIPLOSIS Tavares

ROCHADIPLOSIS Tavares, 1917a:33. Type-species, *tibouchinae* Tavares (orig. des.).

TIBOUCHINAE Tavares, 1917a:34, figs. 6-7; pl. 4, fig. 5; pl. 5, figs. 1-10. Type-locality: Brazil. Host — *Tibouchina* sp. (Melastomaceae). Distr. — Brazil.

Genus SCHIZODIPLOSIS Kieffer

SCHIZODIPLOSIS Kieffer, 1913c:233. Type-species, *Clinodiplosis floralis* Kieffer & Herbst (orig. des.).

FLORALIS (Kieffer & Herbst), 1906:233, pl. 1, figs. 5-6 (*Clinodiplosis*). Type-locality: Chile, Concepción, Concepción. Host — unknown. Distr. — Chile.

Genus SCOPODIPLOSIS Felt

SCOPODIPLOSIS Felt, 1915c:209. Type-species, *speciosa* Felt (orig. des.).

SPECIOSA Felt, 1915c:210. Type-locality: Paraguay, La Cordillera, San Bernardino. Host — unknown. Distr. — Paraguay



Genus SILVESTRINA Kieffer

- SILVESTRINA Kieffer, 1912b:172. Type-species, *Arthrocnodax silvestrii* Kieffer (orig. des.).
- SILVESTRII (Kieffer), 1910:133 (*Arthrocnodax*). Type-locality: Union of South Africa, Capetown. Host — *Pseudaulacaspis pentagona* (Targ.) (Insecta: Homoptera: Diaspididac). Distr. — Brazil (São Paulo), South Africa.

Genus STYRAXDIPLOSIIS Tavares

- STYRAXDIPLOSIIS Tavares, 1915:147. Type-species, *caetitensis* Tavares (orig. des.).
- Styracodiplosis* Tavares, 1918b:77. Unjustified emend.
- CAETITENSIS Tavares, 1915:147, figs. 1-2. Type-locality: Brazil, Bahia, Caitité. Host — *Styrax* sp. (Styracaceae). Distr. — Brazil.
- CEARENSIS Tavares, 1925:20, pl. 1, fig. 8. Type-locality: Brazil, Ceará. Host — *Croton hemiargyreus* Muell. Arg. (Euphorbiaceae). Distr. — Brazil.

Genus TETRADIPOSIIS Kieffer & Jörgensen

- TETRADIPOSIIS Kieffer & Jorgensen, 1910:421. Type-species, *sexdentatus* Kieffer & Jörgensen (mon.).
- SEXDENTATUS Kieffer & Jörgensen, 1910:421, figs. 46-49. Type-locality: Argentina, Mendoza. Hosts — *Prosopis alata* Phil. & *P. campestris* Griseb. (Leguminosae). Distr. — Argentina.

Genus TOXOMYIA Felt

- TOXOMYIA Felt, 1911d:302. Type-species, *fungicola* Felt (orig. des.).
- FUNGICOLA Felt, 1911d:302. Type-locality: St. Vincent. Host — *Puccinia* sp. (Basidiomycetes: Uredinales). Distr. — St. Vincent.
- RUBIDA Felt, 1911g:194. Type-locality: St. Vincent. Host — *Uromyces pisi* (DC.) Oth (Basidiomycetes: Uredinales). Distr. — St. Vincent.

Genus TRISOPSIS Kieffer

- TRISOPSIS Kieffer, 1912b:171. Type-species, *oleae* Kieffer (orig. des.).
- EIFIDA Brèthes, 1914:154, fig. 4. Type-locality: Argentina, Buenos Aires, Buenos Aires. Host — unknown. Distr. — Argentina.

UNPLACED SP. OF CECIDOMYIID

The following five species will be placed in their proper genera in a forthcoming review of the coccid-eating Cecidomyiidae by K. M. Harris.



- COCCI Felt, 1913a:304 (*Karschomyia*). Type-locality: Puerto Rico, Patillas. Host — *Saccharicoccus sacchari* (Ckll.) (Insecta: Homoptera: Pseudococcidae). Distr. — Puerto Rico.
- COCCIDARUM Cockerell, 1892:181, 4 figs. (*Diplosis*). Type-locality: Jamaica, Kingston. Hosts — *Planococcus citri* (Risso) and *Saissetia coffeae* Walker) (Insecta: Homoptera: Coccoidea). Distr. — Jamaica.
- COCCIDARUM Felt, 1911f:548 (*Dicrodiplosis*). Type-locality: Puerto Rico, Mayaguez. Host — "a coccid" (Insecta: Homoptera).
- GUATEMALENSIS Felt, 1938:43 (*Dicrodiplosis*). Type-locality: Guatemala. Host — "mealy bugs" (Insecta: Homoptera). Distr. — Guatemala.
- PULVINARIAE Felt, 1912a:175 (*Mycodiplosis*). Type-locality: St. Vincent. Host — *Protopulvinaria pyriformis* (Ckll.) (Insecta: Homoptera: Coccidae). Distr. — St. Vincent.

UNPLACED SPECIES OF CECIDOMYIIDAE

Adults known

- FLAVIDA Blanchard, 1852:350 (*Cecidomyia*). Type-locality: Chile, Coquimbo, Campos de la Serena. Distr. — Chile.
- FURCATA Philippi, 1865:631 (*Lasioptera*). Type-locality: Chile, Santiago, Santiago. Host — unknown. Distr. — Chile.
- GOLLMERI Karsch, 1880:297, pl. 6, fig. 8 (*Cecidomyia*). Type-locality: Venezuela, Caracas. Host — unknown. Distr. — Venezuela.
- PALLIPES Philippi, 1865:630 (*Lasioptera*). Type-locality: Chile, Santiago, Santiago. Host — unknown. Distr. — Chile.
- PICTIPES Williston, 1896:253 (*Diplosis*). Type-locality: St. Vincent. Host — unknown. Distr. — St. Vincent.
- UMBRA Walker, 1856:421 (*Cecidomyia*). Type-locality: South America. Host — unknown. Distr. — South America.
- UROSTIGMATIS Tavares, 1917b:169, pl. 10, figs. 9, 12; pl. 11, figs. 7,8 (*Asteromyia*). Type-locality: Brazil, Bahia, Salvador. Host — *Urostigma* sp. (Moraceae). Distr. — Brazil.
- Unnamed spp., 3 under *Diplosis*: Williston, 1896:254, pl. 8, fig. 2. Hosts — unknown. Distr. — St. Vincent.

Galls only known

- AVICENNIAE Cook, 1909:145 (*Cecidomyia*). Type-locality: Cuba. Host — *Avicennia nitida* Jacq. (Verbenaceae). Distr. — Cuba.
- BICONICA Cook, 1906:251 (*Cecidomyia*). Type-locality: Cuba, Oriente, Santiago de Cuba. Host — undet. plant. Distr. — Cuba.
- CAYAMASENSIS Cook, 1906:251 (*Cecidomyia*). Type-locality: Cuba, Santa Clara, Cayamas. Host — undet. plant. Distr. — Cuba.
- CECROPIAE Cook, 1909:145 (*Cecidomyia*). Type-locality: Cuba. Host — *Cecropia obtusa* Tréc. (Urticaceae). Distr. — Cuba.
- COCCOLOBAE Cook, 1909:145, pl. 41, fig. 2 (*Cecidomyia*). Type-locality: Cuba. Host — *Coccoloba uvifera* L. (Polygonaceae). Distr. — Cuba.



- EUPATORIUM Cook, 1909:144, pl. 40, fig. 2 (*Cecidomyia*). Type-locality: Cuba.
Host — *Eupatorium villosum* Sw. (Compositae). Distr. — Cuba.
- FICI Cook, 1909:145, pl. 41, fig. 3 (*Cecidomyia*). Type-locality: Cuba.
Host — *Ficus* sp. (Moraceae). Distr. — Cuba.
- MAZAINA Cook, 1906:252 (*Cecidomyia*). Type-locality: Cuba, Oriente, Santiago de Cuba. Host — undet. plant. Distr. — Cuba.
- PISONIAE Cook, 1909:144, pl. 40, fig. 1 (*Cecidomyia*). Type-locality: Cuba.
Host — *Pisonia* sp. (Nyctaginaceae). Distr. — Cuba.
- TORREANA Cook, 1906:252 (*Cecidomyia*). Type-locality: Cuba, Oriente, Santiago de Cuba. Host — undet. plant. Distr. — Cuba.

LITERATURE CITED

AGASSIZ, L. & H. LOEW

1846. Nomina systematica generum dipterorum. [Pt. 4], 42 pp. In Agassiz, L., *Nomenclator zoologicus*. Fasc. 9/10; Titulum et praefationem operis, etc. Soloduri [Solothurn, Switzerland].

ALEXANDER, C. P.

1936. The maximum number of antennal segments in the order Diptera, with the description of a new genus of Cecidomyiidae. *Bull. Brook. Entomol. Soc.* 31:12-14.
1937. Change of name in Diptera. *Ibid.* 32:60.

BARNES, H. F.

1930. A new thrips-eating gall midge, *Thripsobremia liothrips*, gen. et sp. n. (Cecidomyiidae). *Bull. Entomol. Res.* 21:231-232.
1932. Notes on Cecidomyiidae. *Ann & Mag. Nat. Hist.* (10) 9:475-484.
- 1946a. *Gall midges of economic importance. Vol. 1. Root and vegetable crops.* Crosby Lockwood & Son, Ltd., London. 104 pp.
- 1946b. *Gall midges of economic importance. Vol. 2. Fodder crops.* Crosby Lockwood & Son, Ltd., London. 160 pp.
1949. *Gall midges of economic importance. Vol. 6. Miscellaneous crops.* Crosby Lockwood & Son, Ltd., London. 229 pp.
1954. Gall-midge larvae as endoparasites, including the description of a species parasitising aphids in Trinidad, B.W.I. *Bull. Entomol. Res.* 45:769-775.

BLANCHARD, E.

1852. *Insectos. Hist. fisica y política de Chile. Zoologia.* VII:1-471.

BLANCHARD, E. E.

1938. Descripciones y anotaciones de dipteros Argentinos. *An. Soc. cient. Argent.* 126:345-386.
1939. Descripcion del cecidomiido productor de la agalla del quebracho blanco. *Rev. Chil. Hist. Nat.* 42:173-176.
1958. *Contarinia palposa* sp. nov., parasita del sorgo granifero. *Rev. Invest. Agr.* 12:423-425.

BORGMEIER, T.

1931. Eine neue zoophage Itonididengattung aus S. Paulo. (Diptera, Itonididae). *Rev. Entomol.* 1:184-191.



BRÉTHES, J.

1914. *Description de six Cécidomyiidae (Dipt.) de Buenos Aires*. Coni Hermanos, Peru, 6 pp. Also 1915: *An. Mus. Nac. Hist. Nat. Buenos Aires* 26:151-156.
- 1917a. Sur une cécidie de *Physalis viscosa*: description de la cécidie et de la Cécidomyie. *Physis* 3:239-241.
- 1917b. Description d'une cécidie et de sa Cécidomyie d'une *Lippia* d'Entre Rios. *Ibid.* 3:411-413.
1918. Description de la galle et de la cécidomyie d'*Aeschynomene montevidensis*. *Ibid.* 4:312-313.
1922. Himenópteros y Dípteros de varias procedencias. *An. Soc. cient. Argent.* 93:119-146.

COCKERELL, T. D. A.

1892. A cecid bred from Coccidae. *Entomologist* 25:180-182.
1895. *Cecidomyia atriplicis* n. sp. *Amer. Nat.* 29:766-767.

COOK, M. T.

1906. Algunas agallas de Cuba producidas por insectos. *Inf. an. Estac. Cent. Agron.* 1:247-252.
1909. Some insect galls of Cuba. *2nd Rept. Estac. Cent. Agron., Cuba*:143-146, pls. 40-41.

COQUILLET, D. W.

1899. A cecidomyiid injurious to seeds of sorghum. *U.S. Dep. Agr., Div. Entomol. Bull.* 18:81-82.
1905. A new cecidomyiid on cotton. *Can. Entomol.* 37:200.
1910. The type-species of the North American genera of Diptera. *Proc. U.S. Nat. Mus.* 37:499-647.

ENDERLEIN, G.

1940. Die Dipterenfauna der Juan-Fernandez-Inseln und der Oster-Insel. *Nat. Hist. Juan Fernandez & Easter Island* 3(Zool.):643-680.

FELT, E. P.

- 1907a. *New species of Cecidomyiidae*. Albany, N.Y. 53 pp.
- 1907b. New species of Cecidomyiidae. *N.Y. State Mus. Bull.* 110:97-165.
- 1908a. *Contarinia gossypii* n. sp. *Entomol. News* 19:210-211.
- 1908b. Appendix D. *N.Y. State Mus. Bull.* 124:286-422.
1909. New species of West Indian Cecidomyiidae. *Entomol. News* 20:299-302.
- 1910a. West Indian Cecidomyiidae. *Ibid.* 21:268-270.
- 1910b. *Schizomyia ipomoeae* n. sp. *Ibid.* 21:160-161.
- 1910c. Gall midges of *Aster*, *Carya*, *Quercus* and *Salix*. *J. Econ. Entomol.* 3:347-356.
- 1911a. A new *Lestodiplosis*. *Entomol. News* 22:10-11.
- 1911b. Two new gall midges (Dipt.). *Ibid.* 22:109-111.
- 1911c. *Endaphis* Kieff. in the Americas (Dipt.). *Ibid.* 22:128-129.
- 1911d. Four new gall midges (Dipt.). *Ibid.* 22:301-305.
- 1911e. Hosts and galls of American gall midges. *J. Econ. Entomol.* 4:451-484.



- 1911f. New species of gall midges. *Ibid.* 4:546-560.
 1911g. Two new gall midges. *Can. Entomol.* 43:194-196.
 1911h. A generic synopsis of the Itonidac. *J. N.Y. Entomol. Soc.* 19:31-62.
 1911i. Three new gall midges (Dipt.). *Ibid.* 19:190-193.
 1911j. A new species of *Lasioptera* with observations on certain homologies. *Psyche* 18:84-86.
 1912a. New West Indian gall midges (Dipt.). *Entomol. News* 23:173-177.
 1912b. Observations on *Uleela* Rübs. (Dipt.). *Ibid.* 23:353-354.
 1912c. New Itonididac (Dipt.). *J. N.Y. Entomol. Soc.* 20:102-107.
 1912d. New gall midges or Itonidac (Dipt.). *Ibid.* 20:146-156.
 1912e. *Lasiopteryx manihot*, n. sp. (Diptera). *Can. Entomol.* 44:144.
 1913a. Three new gall midges (Diptera). *Ibid.* 45:304-308.
 1913b. Descriptions of gall midges (Diptera). *J. N. Y. Entomol. Soc.* 21:213-219.
 1913c. *Cystodiplosis eugeniae* n. sp. (Dipt.). *Entomol. News* 24:175-176.
 1914a. New gall midges (Itonididae). *Insecutor Inscitiae Menstruus* 2:117-123.
 1914b. *Arthrocnodax constricta* n. sp. *J. Econ. Entomol.* 7:481.
 1915a. New South American gall midges. *Psyche* 22:152-157.
 1915b. New North American gall midges. *Can. Entomol.* 47:226-232.
 1915c. New genera and species of gall midges. *Proc. U.S. Nat. Mus.* 48:195-211.
 1915d. Appendix: a study of gall midges. II. *N.Y. State Mus. Bull.* 175:79-213, pls. 1-15.
 1916. New western gall midges. *J. N.Y. Entomol. Soc.* 24:175-196.
 1917. New gall midges. *Ibid.* 25:193-196.
 1921a. Three new sub-tropical midges (Itonididae, Dipt.). *Entomol. News* 32:141-143.
 1921b. A new *Diadiplosis*. *Zoologica, Scient. Cont. N.Y. Zool. Soc.* 3:225-226.
 1922. A new Cecidomyiid parasite of the white fly. *Proc. U.S. Nat. Mus.* 61:1-2.
 1932. A new citrus cambium miner from Puerto Rico. *Univ. Puerto Rico J. Agr.* 16:117-118.
 1933. A new enemy of the pincapple mealybug and a list of gall midges enemies of mealybugs. *J. N.Y. Entomol. Soc.* 41:87-89.
 1935. New species of gall midges from Texas. *J. Kans. Entomol. Soc.* 8:1-8.
 1938. A new species of gall midge predacious on mealy bugs. *Proc. Haw. Entomol. Soc.* 10:43.

HALIDAY, A. H.

1833. Catalogue of Diptera occurring about Holywood on Downshire. *Entomol. Mag.* (London) 1:147-180.

HARRIS, K. M.

1964. The sorghum midge complex (Diptera, Cecidomyiidae). *Bull. Entomol. Res.* 55:233-247.



- 1966a. [Untitled note]. *Proc. Roy. Entomol. Soc. Lond.* 41:51.
 1966b. Gall midge genera of economic importance (Diptera: Cecidomyiidae). Part 1: Introduction and subfamily Cecidomyiinae; supertribe Cecidomyiidi. *Trans. R. Entomol. Soc. Lond.* 118:313-358.

KARSCH, F.


1877. *Revision der Gallmücken*. E.E. Brunn's Verlag Münster i W. 58 pp., 1 pl.
 1880. Neue Zoocecidien und Cecidozoën. Taf. VI, VII. *Z. Naturwissensch.* 53:288-309.

KIEFFER, J. J.

1889. Neue Beiträge zur Kenntniss der Gallmücken. *Entomol. Nachr.* 15:183-194.
 1894a. Description de quelques larves de Cécidomyes. *Feuille Jeunes Nat.* 24:83-88, 119-121, 147-152, 185-189.
 1894b. Sur le groupe *Epidosis* de la famille des Cecidomyidae. *Ann. Soc. Entomol. France* 63:311-350, 2 pls.
 1895a. Ueber moosbewohnende Gallmückenlarven. *Entomol. Nachr.* 21:113-123.
 1895b. (Genres nouveaux dans le groupe des *Diplosis* (Dipt.,) *Ann. Soc. Entomol. France* 63(1894):cclxxx.
 1895c. Diagnose de deux Cecidomyies nouvelles. *Misc. Entomol.* 3:150.
 1897. *Meine Antwort an der Herrn Zeichenlehrer Rübsaamen*. Paulinus-Druckerei, Trier. 21 pp.
 1898. Synopse des Cécidomyies d'Europe et d'Algérie décrites jusqu'à ce jour. *Bull. Soc. Hist. Nat. Metz* 20:1-64.
 1901. Synopsis des zoocécides d'Europe. *Ann. Soc. Entomol. France* 70:233-579.
 1903. Descriptions de cécidomyies nouvelles de Chili. *Rev. Chil. Hist. Nat.* 7:226-228.
 1909. Contributions à la connaissance des insectes gallicoles. *Bull. Soc. Hist. Nat. Metz* 26:1-35
 1910. Cécidomyies parasites de *Diaspis* sur le Mûrier. *Boll. Lab. Zool. Portici* 4:128-133.
 1912a. *Neue Gallmücken-Gattungen*. Bitsch. 2 pp.
 1912b. Description de quatre nouveaux insectes exotiques. *Boll. Lab. Zool. Portici* 6:171-175.
 1913a. Diptera. Fam. Cecidomyidae. *Genera Insectorum*, Fasc. 152:1-346, 15 pls.
 1913b. Glanures diptérologiques. *Bull. Soc. Hist. Nat. Metz* 28:45-55.
 1913c. Nouvelle contribution a la connaissance des Cécidomyies. *Marcellia* 11(1912):219-235.

KIEFFER, J. J. & P. HERBST

1905. Ueber Gallen und Gallenerzeuger aus Chile. *Z. wissenschaftl. Insektenbiol.* 1:63-66.
 1906. Description de galles et d'insectes gallicoles du Chili. *Ann. Soc. Sci. Bruxelles* 30:223-236, 1 pl.
 1909. Ueber einige neue Gallen und Gallenerzeuger aus Chile. *Centr. Bakteriolog. & Parasitol.* (2)23:119-126.
 1911. Über Gallen und Gallentiere aus Chile. *Ibid.* 29:696-704.



KIEFFER, J. J. & P. JÖRGENSEN.

1910. Gallen und Gallentiere aus Argentinien. *Centr. Bakteriolog. Parasitenk. und Infektionskr.* 27:362-444.

LATREILLE, P. A.

1805. *Histoire naturelle, générale et particulière, des Crustacés et des Insectes*, Tome quatorzième [Vol. 14]. [Vol. 106], 432 pp., pls. 104-112. In Sonnini, C.S., ed., *Histoire naturelle par Buffon*. Paris, An XIII.

LOEW, H.

1850. Dipterologische Beiträge. Vierter Theil. *K. Friedrich-Wilhelms-Gymnasiums zu Posen, Öffentl. Prüf. d. Schüler 1850:1-40*, 1 pl.

MACQUART, J.

1826. Insects Diptères du nord de la France. Tipulaires. *Soc. d'Amateurs des Sci., de l'Agr. et des Arts, Lille, Rec. des Trav.* 1823/1824:59-224, 4 pls.

MEGEN, J. W.

1830. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten*. Vol. 6, iv + 401 pp., pls. 55-66. Hamm.

MEINERT, F.

1864. *Miastor metrolas*: Yderligere oplysning om den af Prof. Nie. Wagner nyligt beskrevne insektlarve, som formerer sig ved spiredannelse. *Naturhist. Tidsskr.* (3)3:37-43.

MÖHN, E.

1959. Gallmücken (Diptera, Itonididae) aus El Salvador. 1. Teil. *Senck. Biol.* 40:297-368.
- 1960a. Gallmücken (Diptera, Itonididae) aus El Salvador. 2. Teil. *Ibid.* 41:197-253.
- 1960b. Gallmücken (Diptera, Itonididae) aus El Salvador. 3. Teil. *Ibid.* 41:333-358.
- 1961a. Neue Asphondyliidi-Gattungen (Diptera, Itonididae). *Stuttgarter Beitr. Naturk.* Nr. 49:1-14.
- 1961b. Gallmücken (Diptera, Itonididae) aus El Salvador. 4. Zur Phylogenie der Asphondyliidi der neotropischen und holarktischen Region. *Senck. Biol.* 42:131-330.
1962. Studien über Neotropische Gallmücken (Diptera, Itonididae). *Brotéria* 31:211-239.
- 1963a. Gallmücken (Diptera, Itonididae) aus El Salvador. 5. Teil, Lasiopteridi. *Stuttgarter Beitr. Naturk.* Nr. 116:1-7.
- 1963b. Studien über neotropische Gallmücken (Diptera, Itonididae). 1. Teil (Fortsetzung). *Brotéria* 32:3-23.
- 1964a. Gallmücken (Diptera, Itonididae) aus El Salvador. 6. Teil. Lasiopteridi. *Deut. Entomol. Zeit.* 11:47-143.
- 1964b. Gallmücken (Diptera, Itonididae) aus El Salvador. 7. Teil: Lasiopteridi. *Beitr. Entomol.* 14:553-600.

MOLLIARD, M.

1903. La galle du *Cecidomyia cattleyae* n. sp. *Marcellia* 1:165-171, 1 pl.



NIJVELDT, W.

1967. *Neolasioptera martelli* sp. n., a gall midge injuring Agave in Mexico. *Entomol. Berichten* 27:125-127.

OSTEN SACKEN, C. R.

1869. Biological notes on Diptera (galls on *Solidago*). *Trans. Amer. Entomol. Soc.* 2:299-303.

PANELIUS, S.

1965. A revision of the European gall midges of the subfamily Porricondyliinae (Diptera: Itonididae). *Acta Zool. Fenn.* 113:1-157.

PARNELL, J. R.

1966. Observations on the larval morphology and life history of *Oleisicoccus costalimai* Borgmeier (Diptera: Cecidomyiidae) in Jamaica. *Proc. Roy. Ent. Soc. Lond.* 41:51-54.

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. Zool. bot. Ges. Wien* 15:595-782.

PRITCHARD, A. E.

1947. North American gall midges of the tribe Micromyini. *Entomol. Amer.* 27:1-84, 2 pls.

1948. The North American gall midges of the tribes Catostrichini and Catochini (Diptera: Itonididae (Cecidomyiidae)). *Ann. Entomol. Soc. Amer.* 40(1947):662-671.

1951. The North American gall midges of the tribe Lestremiini; Itonididae (Cecidomyiidae); Diptera. *Univ. Calif. Pubs. Ent.* 8:239-275.

RONDANI, C.

1840. *Memoria per servir alla ditteologia italiana*. No. 2: *Sopra alcuni nuovi generi di insetti ditteeri*. Parma. 28 pp., 1 pl.

1846. Compendio della seconda memoria ditteologica (publicata 1840) con alcune aggiunte et correzioni. *Soc. Agric. & Accad. Sci. Ist. Bologna, Nuovi Ann. Sci. Nat. Rend.* (2)6:363-376, 1 pl.

1856. *Dipterologiae Italicae prodromus*. Vol. 1: *Genera Italica ordinis dipterorum ordinatum disposita et distincta et in familias et stires aggregata*. Parmae [Parma]. 228 pp.

1860. Stirpis cecidomyarum. Genera revisa. Nota undecima, pro dipterologia italica. *Atti Soc. Ital. Sci. Nat. (1859-1860)* 2:286-294, 1 pl.


RÜBSAAMEN, E. H.

1892. Die Gallmücken des Königl. Museums für Naturk. zu Berlin. *Berlin. Entomol. Zeitschr.* 37:321-411, pls. 7-18.

1895. Cecidomyidenstudien. *Entomol. Nachr.* 21:177-194.

1905. Beiträge zur Kenntnis aussereuropäischer Zooecidien. II Beitrag: Gallen aus Brasilien und Peru. *Marcellia* 4:65-85, 115-138.

- 1908a. Beiträge zur Kenntnis aussereuropäischer Zooecidien. III Beitrag. Gallen aus Brasilien und Peru. *Ibid.* 6:110-173.



- 1908b. Beiträge zur Kenntnis aussereuropäischer Zooecidien. III. Beitrag: Gallen aus Brasilien und Peru. *Ibid.* 7:15-79.
1910. Ueber deutsche Gallmücken und Gallen. *Z. wissenschaft. Insektenbiol.* 6:283-289.
1911. Ueber deutsche Gallmücken und Gallen. *Ibid.* 7:120-125.
- 1916a. Beitrag zur Kenntnis aussereuropäischer Gallmücken. *Ges. Naturforsch. Freunde Berlin 1915*:431-481.
- 1916b. Cecidomyidenstudien V. Revision der deutschen Asphondyliarien. *Ibid.* 1916:1-16.

TAVARES, J. S.

1906. Descrição de uma *Cecidomyia* nova do Brazil, pertencente a um genero novo. *Brotéria*, série zoológica 5:81-84.
1909. Contributio prima ad cognitionem cecidologiae Braziliae. *Ibid.* 5-28, 8 pls.
1915. As cecidias das plantas do genero *Styrax* no Brazil. *Ibid.* 3:145-159, pl. 6.
1916. Cecidomyias novas do Brazil. *Ibid.* 14:36-57.
- 1917a. As cecidias do Brazil que se criam nas plantas da familia das Melastomataceae. *Ibid.* 15:18-49, pls. 1-5.
- 1917b. Cecidias brazileiras que se criam em plantas das familias das Compositae, Rubiaceae, Tiliaceae, Lythraceae e Artocarpaceae. *Ibid.* 15:113-181, pls. 6-11.
- 1918a. Cecidologia Brazileira. Cecidias que se criam nas plantas das familias das Verbenaceae, Euphorbiaceae, Malvaceae, Anacardiaceae, Labiatae, Rosaceae, Anonaceae, Ampelidaceae, Bignoniaceae, Aristolochiaceae e Solanaceae. *Ibid.* 16:21-68, 2 pls.
- 1918b. Cecidomyias novas do Brazil. *Ibid.* 16:68-84, 2 pls.
- 1920a. O genero *Bruggmaniella* Tav. com descrição de uma espécie nova e a clave dichotomica des gêneros das Asphondyliariae. *Ibid.* 18:33-42.
- 1920b. Cecidologia Brazileira. Cecidias que se criam em plantas das familias das Leguminosae, Sapotaceae, Lauraceae, Myrtaceae, Punicaceae, Aurantiaceae, Malpighiaceae, Sapindaceae, Umbelliferae, Loranthaceae, Apocynaceae, Urticaceae, Salicaceae e Gramineae. *Ibid.* 18:82-125, 1 pl.
- 1920c. Anadiplosariae, nouvelle tribu de Cecidomyinae (Dipt.). *Marcellia* 17:57-73.
1921. Cecidologia brazileira. Cecidias que se criam em plantas das familias das Leguminosae, Sapotaceae, Lauraceae, Myrtaceae, Punicaceae, Aurantiaceae, Malpighiaceae, Sapindaceae, Umbelliferae, Loranthaceae, Apocynaceae, Urticaceae, Salicaceae e Gramineae. *Brotéria*, série zoológica 19:76-112, 1 pl.
1922. Cecidologia brazileira. As restantes familias. *Ibid.* 20:5-48, pls. 11-19.
1925. Nova contribuição para o conhecimento da Cecidologia Brazileira. *Ibid.* 22:5-55, pls. 1-4.

TOWNSEND, C. H. T.

1893. Fleshy cecidomyiid twig gall on *Atriplex canescens*. *Amer. Nat.* 27:1021.



WALKER, F.

1856. *Diptera*. Vol. 1, pp. 415-474. In *Insecta Saundersiana*: London, "1856".

WEYENBERGH, H.

1875. *Ann. Agr. Republ. Argentina* 3:164 (not seen).

WILLISTON, S. W.

1896. XI. On the *Diptera* of St. Vincent (West Indies). *Trans. Entomol. Soc. Lond.* 1896:253-446.

WOLCOTT, G. N.

1936. *Insectae Borinquenses*, a revised, annotated checklist of the insects of Puerto Rico. *Univ. Puerto Rico Jour. Agr.* 20:1-600.

WYATT, I. J.


1967. Pupal paedogenesis in the *Cecidomyiidae* (*Diptera*). 3. A reclassification of the *Heteropezini*. *Trans. R. Entomol. Soc. Lond.* 119:71-98.

INDEX

abdominalis (Felt), 1911 (Endaphis), <i>Arthrochodax</i>	23.30
achatoearpi Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
acuticauda Kieffer & Herbst, 1906, <i>Janetiella</i>	23.7
aeschynomensis Brèthes, 1918 (<i>Lasioptera</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.21
agglomerata Kieffer, 1913, <i>Scheueria</i>	23.8
ajallai Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
albida Tavares, 1909, <i>Guarephila</i>	23.5
albipes Felt, 1915, <i>Xenasphondylia</i>	23.15
<i>Alexomyia</i> Felt, 1921	23.31
aleyrodici Felt, 1922, <i>Cleodiplosis</i>	23.32
<i>Allodiplosis</i> Kieffer & Jörgensen, 1910	23.31
altani Felt, 1915, <i>Asphondylia</i>	23.13
<i>Alycaulus</i> Rübsaamen, 1916	23.19
amaranthi Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.21
<i>Ameliomyia</i> Möhn, 1960	23.16
americana Felt, 1911, <i>Hyperdiplosis</i>	23.28
anaceliae Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
<i>Anadiplosini</i>	23.26
<i>Anadiplosis</i> Tavares, 1916	23.26
Anarete Haliday, 1833	23.1
<i>Anasphondylia</i> Blanchard, 1939	23.18
<i>Anasphondylia</i> Tavares, 1920	23.18
andirae Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
<i>Andriodiplosis</i> Tavares, 1920	23.31
<i>Angeiomyia</i> Kieffer & Herbst, 1909	23.4
annonae Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
aphelandrae Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.21
<i>Aphidoletini</i>	23.27
<i>Apodiplosis</i> Tavares, 1922	23.31
ardisiae Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9



<i>argentata</i> Brèthes, 1917, <i>Physalidicola</i>	23.25
<i>argentina</i> Brèthes, 1914, <i>Austrodiplosis</i>	23.31
<i>armeniae</i> Möhn, 1960, <i>Pisoniamyia</i>	23.17
<i>Arthrocnodax</i> Rübсаamen, 1895	23.30
<i>Asphondylia</i> Loew, 1850	23.9
<i>Asphondyliidi</i>	23.9
<i>Asphondyliini</i>	23.9
<i>aspidospermae</i> Blanchard, 1939 (<i>Anasphondylia</i>)	23.18
<i>Asteromyia</i> Felt, 1910	23.19
<i>Astrodiplosis</i> Felt, 1913	23.31
<i>Asynapta</i> Loew, 1850	23.3
<i>Asynaptini</i>	23.3
<i>atriplicis</i> (Townsend), 1893 (<i>Cecidomyia</i>), <i>Asphondylia</i>	23.13
<i>attenuatata</i> Felt, 1909, <i>Asphondylia</i>	23.13
<i>aurata</i> Felt, 1915, <i>Ouradiplosis</i>	23.36
<i>auripes</i> Felt, 1915, <i>Epihormomyia</i>	23.30
<i>australis</i> Brèthes, 1914, <i>Neobaeomyza</i>	23.35
<i>Austrodiplosis</i> Brèthes, 1914	23.31
<i>Austrolauthia</i> Brèthes, 1914	23.4
<i>Autodiplosis</i> Tavares, 1920	23.28
<i>avicenniae</i> Cook, 1909 (<i>Cecidomyia</i>)	23.38
<i>ayeniae</i> Möhn, 1960, <i>Asphondylia</i> (<i>Eoasphondylia</i>)	23.12
<i>szarae</i> (Kieffer), 1903 (<i>Perrisia</i>), <i>Xyloperissia</i>	23. 8
<i>baccharis</i> Kieffer & Herbst, 1905, <i>Asphondylia</i>	23.13
<i>Baccharomyia</i> Tavares, 1917	23.19
<i>baezi</i> Brèthes, 1922, <i>Neurolasioptera</i>	23.25
<i>bahiensis</i> Tavares, 1920, <i>Andirodiplosis</i>	23.31
<i>bahiensis</i> Tavares, 1917, <i>Asphondylia</i>	23.13
<i>bahiensis</i> (Tavares), 1917 (<i>Eudiplosis</i>), <i>Schismatodiplosis</i>	23.28
<i>bedeguaris</i> (Kieffer & Jörgensen), 1910 (<i>Rhopalomyia</i>), <i>Panteliola</i>	23.7
<i>biconica</i> Cook, 1906 (<i>Cecidomyia</i>)	23.38
<i>bifida</i> Brèthes, 1914, <i>Trisopsis</i>	23.37
<i>boerhaaviae</i> Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
<i>borreriae</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>borreriae</i> Rübсаamen, 1905, <i>Asphondylia</i>	23.13
<i>Brachylasioptera</i> Möhn, 1964	23.20
<i>brasiliensis</i> Felt, 1915, <i>Proasphondylia</i>	23.18
<i>brasiliensis</i> (Rübсаamen), 1908 (<i>Clinodiplosis</i>), <i>Iatrophobia</i>	23.28
<i>brasiliensis</i> Rübсаamen, 1908, <i>Lopesia</i>	23.35
<i>brasiliensis</i> Rübсаamen, 1916, <i>Machaeriobia</i>	23.35
<i>brasiliensis</i> Felt, 1915, <i>Johnsonomyia</i>	23.4
<i>brasiliensis</i> Tavares, 1906, <i>Bruggmannia</i>	23.16
<i>brasiliensis</i> Tavares, 1909, <i>Bruggmanniella</i>	23.14
<i>brasiliensis</i> Tavares, 1922 (<i>Perrisia</i>), <i>Dasineura</i>	23.5
<i>brasiliensis</i> (Tavares), 1920 (<i>Coprodiplosis</i>), <i>Lestodiplosis</i>	23.30
<i>brevicornis</i> (Felt), 1907 (<i>Campylomyza</i>), <i>Corinthomya</i>	23.2
<i>brickelliae</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>Bruggmannia</i> Tavares, 1906	23.16
<i>Bruggmanniella</i> Tavares, 1909	23.14
<i>buddleia</i> Felt, 1935, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.9
<i>burserae</i> Möhn, 1960, <i>Burseramyia</i>	23.17
<i>Burseramyia</i> Möhn, 1960	23.17
<i>buscki</i> (Felt), 1915 (<i>Microcerata</i>), <i>Anarete</i>	23.2
<i>buscki</i> Felt, 1914, <i>Diadiplosis</i>	23.33



cabezasae Möhn, 1959, Asphondylia (Asphondylia)	23.9
caetensis Tavares, 1920, Anadiplosis	23.26
caetensis Tavares, 1915, Styraxdiplosis	23.37
calcac Möhn, 1959, Asphondylia (Asphondylia)	23.9
calcae Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
Calmonia Tavares, 1917	23.4
Calopedilla Kieffer, 1913	23.4
camarac Möhn, 1959, Asphondylia (Asphondylia)	23.9
camarae Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
Camptomylia Kieffer, 1894	23.3
Camptoneuromylia Felt, 1908	23.20
capsici Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
caryae (Felt), 1907 (Choristoneura), Trotteria	23.25
cattleyae Felt, 1908 (Clinodiplosis), Parallelodiplosis	23.28
cattleyae (Molliard), 1902 (Cecidomyia), Parallelodiplosis	23.28
cayamasensis Cook, 1906 (Cecidomyia)	23.38
cearensis (Tavares), 1917 (Eudiplosis), Schismatodiplosis	23.29
cearensis Tavares, 1925, Styraxdiplosis	23.37
Cecidomyiidi	23.26
Cecidomyiinae	23.4
cecropiae Cook, 1909 (Cecidomyia)	23.38
celtis Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
Centrodiplosis Kieffer & Jörgensen, 1910	23.31
Ceratomyia Felt, 1911	23.2
cerei (Rübsaamen), 1905 (Lasioptera), Neolasioptera (Neolasiop- tera)	23.22
cestri Möhn, 1959, Asphondylia (Asphondylia)	23.10
cestri Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
Charidiplosis Tavares, 1918	23.32
chilensis (Kieffer & Herbst), 1909 (Perrisia), Dasineura	23.5
chilensis (Kieffer & Herbst), 1906 (Clinodiplosis), Plesioidiplosis	23.36
chlorophorae Rübsaamen, 1905, Clinodiplosis	23.27
Choristoneura Rübsaamen, 1892	23.25
ciliata Felt, 1921, Alexomyia	23.31
cincta Felt, 1915, Johnsonomyia	23.4
cinctipes Felt, 1915, Delphodiplosis	23.33
cinerea Macquart, 1826, Lestremia	23.2
cissampeli Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
cissi Möhn, 1959 Asphondylia (Asphondylia)	23.10
citrinae Felt, 1932, Asynapta	23.3
clavata Tavares, 1920 (Oxasphondylia)	23.19
clematicola Möhn, 1964, Neolasioptera (Neolasioptera)	23.22
Cleitodiplosis Tavares, 1921	23.27
clematidis Felt, 1935, Asphondylia (Asphondylia)	23.10
Cleodiplosis Felt, 1922	23.32
Clinodiplosini	23.27
Clinodiplosis Kieffer, 1895	23.27
clusiae Rübsaamen, 1905, Uleia	23.8
cocci Felt, 1911, Diadiplosis	23.33
cocci Felt, 1913 (Karshomyia)	23.38
coccidarum Cockerell, 1892 (Diplosis)	23.38
coccidarum Felt, 1911 (Dicrodiplosis)	23.38
coccidarum Felt, 1911, Lobodiplosis	23.34
coccidivora (Felt), 1911 (Mycodiplosis), Oleisicoccus	23.35
coccolobac Cook, 1909 (Cecidomyia)	23.38



<i>coffea</i> Felt, 1911, <i>Hyperdiplosis</i>	23.28
<i>colliguayae</i> Kieffer & Herbst, 1911, <i>Riveraella</i>	23.8
<i>combreti</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>compositarum</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>Compsodiplosis</i> Tavares, 1909	23.32
<i>concinna</i> Tavares, 1918, <i>Charidiplosis</i>	23.32
<i>constrictus</i> Felt, 1914, <i>Arthrocnodax</i>	23.30
<i>Contarinia</i> Rondani, 1860	23.29
<i>Contariniini</i>	23.29
<i>convolvuli</i> Möhn, 1960, <i>Asphondyllia</i> (<i>Eoasphondyllia</i>)	23.12
<i>Coquilletomyia</i> Felt, 1908	23.32
<i>Coprodiplosis</i> Kieffer, 1894	23.30
<i>corbulae</i> Möhn, 1960, <i>Asphondyllia</i> (<i>Asphondyllia</i>)	23.10
<i>cordiae</i> Möhn, 1959, <i>Asphondyllia</i> (<i>Asphondyllia</i>)	23.10
<i>cordiae</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>cordobensis</i> Kieffer & Jørgensen, 1910 (<i>Lasioptera</i>)	23.25
<i>Corinthomyia</i> Felt, 1911	23.2
<i>costalimai</i> Borgmeier, 1931, <i>Oleiscoccus</i>	23.36
<i>coussapoeae</i> Rübsaamen, 1905, <i>Frauenfeldella</i>	23.34
<i>crassipalpis</i> Kieffer & Jørgensen, 1910, <i>Asphondyllia</i>	23.13
<i>crassipes</i> Kieffer & Jørgensen, 1910, <i>Centrodiplosis</i>	23.32
<i>crassus</i> Kieffer & Jørgensen, 1910, <i>Allodiplosis</i>	23.31
<i>Ctenodactylomyia</i> Felt, 1915	23.32
<i>curatellae</i> Möhn, 1959, <i>Asphondyllia</i> (<i>Asphondyllia</i>)	23.10
<i>Cystodiplosis</i> Kieffer & Jørgensen, 1910	23.33
<i>Dactylodiplosis</i> Rübsaamen, 1916	23.33
<i>dalbergiae</i> Rübsaamen, 1908, <i>Uleela</i>	23.26
<i>dalechampiae</i> (Rübsaamen,) 1905 (<i>Meunieria</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>Dasineura</i> Rondani, 1840	23.5
<i>Dasyneura</i> Agassiz & Loew, 1846	23.5
<i>Delphodiplosis</i> Felt, 1915	23.33
<i>dentata</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>depressa</i> Kieffer, 1913, <i>Uleela</i>	23.26
<i>Diadiplosis</i> Felt, 1911	23.33
<i>Dialeria</i> Tavares, 1918	23.20
<i>Dilasioptera</i> Möhn, 1964	23.25
<i>Dolicholabis</i> Tavares, 1918	23.20
<i>dubia</i> Rübsaamen, 1916, <i>Sphaerodiplosis</i>	23.30
<i>Eoasphondyllia</i> Möhn, 1960	23.12
<i>Epihormomyia</i> Felt, 1915	23.30
<i>Epilasioptera</i> Möhn, 1964	23.20
<i>erigerontis</i> (Felt), 1907 (<i>Choristoneura</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.22
<i>Erosomyia</i> Felt, 1911	23.33
<i>erythroxyli</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.23
<i>erythroxyli</i> Möhn, 1959, <i>Asphondyllia</i> (<i>Asphondyllia</i>)	23.10
<i>Eudiplosis</i> Tavares, 1916	23.28
<i>eugeniae</i> Felt, 1912, <i>Dasineura</i>	23.5
<i>eugeniae</i> Felt, 1913 (<i>Cystodiplosis</i>), <i>Stephomyia</i>	23.18
<i>eugeniae</i> Kieffer & Herbst, 1909, <i>Oligotrophus</i>	23.6



eugeniae Tavares, 1916, Stephomyia	23.18
eupatoriensis Möhn, 1964, Isolasioptera	23.21
eupatorii Cook, 1909 (Cecidomyia)	23.39
eupatorii Felt, 1911, Hyperdiplosis	23.28
eupatorii Tavares, 1917, Geraldesia	23.21
exigua Möhn, 1964 Neolasioptera (Neolasioptera)	23.23
falcigera (Kieffer & Jörgensen), 1910 (Centrodiplosis), Jorgensenia	23.34
fariae (Tavares), 1922 (Luisieria), Neolasioptera (Neolasioptera)	23.23
<i>Feltomyia</i> Alexander, 1936	23.4
<i>Feltomyia</i> Kieffer, 1913	23.16
<i>Feltomyia</i> Alexander, 1937	23.4
fici Cook, 1909 (Cecidomyia)	23.39
flavida Blanchard, 1852 (Cecidomyia)	23.38
floralis (Kieffer & Herbst), 1906 (Clinodiplosis), Schizodiplosis...	23.36
Frauenfeldiella Rübsaamen, 1905	23.33
friburgensis Tavares, 1915, Compsodiplosis	23.32
friburgensis (Tavares), 1917 (Oxasphondylia), Rhoasphondylia ..	23.15
fungicola Felt, 1911, Toxomyia	23.37
furcata Philippi, 1865 (Lasioptera)	23.38
<i>fusca</i> Meigen, 1830, Lestremia	23.2
gadoquiae Kieffer & Herbst, 1905, Dasineura	23.5
Geraldesia Tavares, 1917	23.21
gilibertiae Möhn, 1964, Brachylasioptera	23.20
<i>Gisonobasis</i> Rübsaamen, 1916	23.9
globifex (Kieffer & Jörgensen), 1910 (Rhopalomyia), Misospatha..	23.5
globulifex Kieffer, 1913, Uleela	23.26
Gnesiodiplosis Tavares, 1917	23.34
godmaniae Möhn, 1959, Asphondylia (Asphondylia)	23.10
gollmeri Karsch, 1880 (Cecidomyia)	23.38
gossypii (Cpquillet), 1905 (Porricondyla), Asynapta	23.3
gossypii Felt, 1908, Contarinia	23.29
gracilicornis (Kieffer & Herbst), 1905 (Perrisia), Dasineura ...	23.5
graciliforceps (Kieffer & Jörgensen), 1910 (Lasioptera), Neola-	
sioptera (Neolasioptera)	23.23
gracilis Kieffer & Jörgensen, 1910, Lyciomyia	23.7
graminis (Tavares), 1916 (Necrophlebia), Cleitodiplosis	23.27
grandis Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
guareae Möhn, 1959, Asphondylia (Asphondylia)	23.10
Guarephila Tavares, 1909	23.5
guatemalensis Felt, 1938 (Dicrodiplosis)	23.38
gutierreziae Felt, 1916, Asteromyia	23.19
Haplopalpus Rübsaamen, 1916	23.6
Haplusia Karsch, 1877	23.6
haywardi Blanchard, 1938 (Phagodiplosis), Oleiscoccus	23.36
heisteriae Rübsaamen, 1916, Dactylodiplosis	23.33
helicteris Möhn, 1959, Asphondylia (Asphondylia)	23.10
heliocarpus Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
Hemiasphondylia Möhn, 1960	23.14
Hemibruggmaniella Möhn, 1961	23.14
herbsti (Kieffer), 1903 (Rhopalomyia), Calopedilla	23.5



herculesi Möhn, 1959, Asphondylia (Asphondylia)	23.10
Heterasphondylia Möhn, 1960	23.14
Heteropezini	23.3
heterothalami (Kieffer & Jörgensen), 1910 (Lasioptera), Neolasioptera (Neolasioptera)	23.23
hieronymi (Weyenbergh), 1875 (Lasioptera), Asphondylia	23.13
<i>Holoneura</i> Kieffer, 1894	23.3
Holoneurus Kieffer, 1895	23.3
Hormomyiini	23.30
Houardodiplosis Tavares, 1925	23.27
humilis Tavares, 1915, Compsodiplosis	23.32
Hyperdiplosis Felt, 1908	23.28
hyptis Möhn, 1960, Asphondylia (Asphondylia)	23.10
hyptis Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
Iatrophobia Rübsaamen, 1916	23.28
iheringi (Tavares), 1925 (Autodiplosis), Schismatodiplosis	23.29
indigoferae Möhn, 1960, Asphondylia (Asphondylia)	23.10
indigoferae Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
ingae Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
insignis Tavares, 1922, Dolicholabis	23.20
insularis (Felt), 1913 (Mycodiplosis), Arthrocnodax	23.30
interrupta (Kieffer & Jörgensen), 1910 (Lasioptera), Neolasioptera (Neolasioptera)	23.23
ipomoeae Felt, 1910, Schizomyia	23.18
iresinis Möhn, 1964, Epilasioptera	23.20
iresinis Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
Isolasioptera Möhn, 1964	23.21
itaparicae Tavares, 1917, Gnesiodiplosis	23.34
itaparicana Tavares, 1922, Compsodiplosis	23.32
ituensis Tavares, 1917 (Oxasphondylia)	23.19
Janetiella Kieffer, 1898	23.6
jatrophae Möhn, 1959, Asphondylia (Asphondylia)	23.10
<i>Jatrophobia</i> , emend.	23.28
johannseni (Felt), 1911 (Ceratomyia), Micromyia	23.2
Johnsonomyia Felt, 1908	23.4
Jorgensenia Kieffer, 1913	23.34
Kalodiplosis Felt, 1915	23.34
Karschomyia Felt, 1908	23.34
knabi Felt, 1912, Coquillettomyia	23.32
lantanae (Rübsaamen), 1908 (Clinodiplosis), Schismatodiplosis ...	23.29
lantanae Tavares, 1918, Dolicholabis	23.20
lapalmae Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
lapalmae Möhn, 1960, Polystepha	23.16
<i>Lasiopteryx</i> Felt, misident	23.7
Lestodiplosini	23.30
Lestodiplosis Kieffer, 1894	23.30
Lestremia Macquart, 1826	23.2
Lestremiinae	23.1
Lestremiini	23.1



Liebeliola Kieffer & Jörgensen, 1910	23.34
lignicola Kieffer, 1913, Uleela	23.26
lindneri Möhn, 1963, Trotteria	23.25
liothripis Barnes, 1930, Thripsobremia	23.27
lippiae Brèthes, 1917, Pseudomikiola	23.8
lippiae (Kieffer & Jörgensen), 1910 (Rhopalomyia), Panteliola ...	23.7
lippiae Möhn, 1959, Asphondylia (Asphondylia)	23.10
Lobodiplosis Felt, 1908.	23.34
Lobolasioptera Möhn, 1964	23.21
lonchocarpi Möhn, 1960, Asphondylia (Asphondylia)	23.10
longicauda Kieffer, 1913, Uleela	23.26
longicornis Kieffer & Herbst, 1909, Pernettyella	23.7
longicornis Kieffer & Herbst, 1909, Scheueria	23.8
longipennis Kieffer & Jörgensen, 1910, Cystodiplosis	23.33
longiseta Kieffer, 1913, Uleela	23.27
Lopesia Rübsaamen, 1908	23.35
Luisieria Tavares, 1922	23.21
lutealbida Tavares, 1922, Compsodiplosis	23.32
lycicola Kieffer & Jörgensen, 1910, Oligotrophus	23.6
Lyciomyia Kieffer & Jörgensen, 1910	23.6
lycopersici Felt, 1911, Contarinia	23.29
machaerii (Kieffer), 1913 (Uleela), Ruebsaamenodiplosis	23.26
Machaerobia Rübsaamen, 1916	23.35
macroflus Wolcott nec Felt, Arthrocnodax	23.30
Macroporpa Rübsaamen, 1916	23.17
maculans Barnes, 1954, Pseudendaphis	23.36
maculata Brèthes, 1914, Neobaeomyza	23.35
malvavisci Möhn, 1959, Asphondylia (Asphondylia)	23.11
mangiferae Felt, 1909, Asynapta	23.3
mangiferae Felt, 1911, Erosomyia	23.33
mangiferae Tavares, 1918 (Mangodiplosis), Erosomyia	23.33
Mangodiplosis Tavares, 1918	23.33
manihot Felt, 1910 (Cecidomyia), Iatrophobia	23.28
manihot (Felt), 1912 (Lasiapteryx), Phaenolauthia	23.7
manihoti Tavares, 1925, Schizomyia	23.18
marcetiae (Tavares), 1917 (Eudiplosis), Schismatodiplosis	23.29
martelli Nijveldt, 1967, Neolasioptera (Neolasioptera)	23.23
mazaina Cook, 1906 (Cecidomyia)	23.39
media Möhn, 1964, Lobolasioptera	23.21
medicaginis Kieffer, 1895, Contarinia	23.29
Megaulus Rübsaamen, 1916	23.35
melantherae Möhn, 1959, Asphondylia (Asphondylia)	23.11
melantherae Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
meridionalis Felt, 1912, Arthrocnodax	23.30
meridionalis Felt, 1910, Camptoneuromyia	23.20
merrimiae Möhn, 1964, Neolasioptera (Neolasioptera)	23.23
Metasphondylia Tavares, 1918	23.19
Meunieria Rübsaamen, 1905	23.21
Meunieriella Rübsaamen, 1909	23.21
mexicana (Felt), 1911 (Bruggmanniella), Pisoniamyia	23.17
Miastor Meinert, 1864	23.3
Microcerata Felt, 1908	23.1



Micromyza Rondani, 1840	23.2
<i>Micromyza</i> , emend.	23.2
Micromyziini	23.2
micrura Kieffer, 1913, Ulecla.	23.27
mikaniae Rübbsaamen, 1916, Alycaulus	23.19
mimosae Möhn, 1960, Hemiasphondylia	23.14
mimosae (Tavares), 1925 (Schizomyia), Tavaresomyia	23.15
Misospatha Kieffer, 1913	23.5
modesta Williston, 1896, Trichopteromyia	23.2
monacha Osten Sacken, 1869, Asphondylia	23.13
monticola (Kieffer & Herbst), 1909 (Lasioptera), Neolasioptera (Neolasioptera)	23.23
montivaga Kieffer & Jörgensen, 1910, Janetiella	23.7
multifila (Felt), 1908 (Dichrodiplosis [<i>sic</i>]), Kalodiplosis	23.34
myrtacea Tavares, 1920, Anasphondylia	23.19
nectandrae Kieffer & Jörgensen, 1910, Oligotrophus	23.6
neeana Kieffer, 1913, Ulecla	23.27
Neobaemyza Brèthes, 1914	23.35
Neolasioptera Felt, 1908	23.21
Neurolasioptera Brèthes, 1922	23.25
nigra Blanchard, 1852, Lestremia	23.2
oblita (Tavares), 1920 (Bruggmanniella), Hemibruggmanniella ...	23.14
occidentalis Felt, 1911, Holoneurus	23.3
ocimi Möhn, 1959, Asphondylia (Asphondylia)	23.11
odontonemae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
Oleiscoccus Borgmeier, 1931	23.35
Oligotrophidi	23.4
Oligotrophus Latreille, 1805	23.6
opuntiae Felt, 1908, Asphondylia	23.13
oreiplana (Kieffer & Jörgensen), 1910 (Rhopalomyia), Misospatha.	23.6
ornaticornis (Kieffer & Jörgensen), 1910 (Lasioptera), Neolasioptera (Neolasioptera)	23.24
Ouradiplosis Felt, 1915	23.36
<i>Oxasphondylia</i> Felt, 1915	23.15
<i>Ozobia</i> Kieffer, 1913	23.15
palaciosi Möhn, 1959, Asphondylia (Asphondylia)	23.11
pallida Enderlein, 1940, Psadaria	23.2
pallipes Phillippi, 1865 (Lasioptera)	23.38
palposa Blanchard, 1958, Contarinia	23.29
Panteliola Kieffer, 1913	23.7
Parallelodiplosis Rübbsaamen, 1910	23.28
parasitica Möhn, 1960, Asphondylia (Asphondylia)	23.11
parrishi (Felt), 1915 (Porricondylia), Camptomysia	23.3
parva Tavares, 1917, Asphondylia	23.13
parva (Tavares), 1916 (Eudiplosis), Schismatodiplosis	23.29
parvula Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
pattersoni Felt, 1911, Asphondylia	23.13
Perasphondylia Möhn, 1960	23.14
Pernettyella Kieffer & Herbst, 1909	23.7
<i>Perrisia</i> Rondani, 1846	23.5




<i>peruviana</i> Felt, 1911, <i>Lestodiplosis</i>	23.30
<i>peruviana</i> Rübсаamen, 1916, <i>Macroporpa</i>	23.17
<i>Phaenolauthia</i> Kieffer, 1912	23.7
<i>Phagodiplosis</i> Blanchard, 1938	23.35
<i>Physalidicola</i> Brèthes, 1917	23.25
<i>pictipes</i> Williston, 1896 (<i>Diplosis</i>)	23.38
<i>picturata</i> Felt, 1915, <i>Lestodiplosis</i>	23.30
<i>piperis</i> Rübсаamen, 1908, <i>Zalepidota</i>	23.15
<i>piriqueta</i> (Felt), 1917 (<i>Lasioptera</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.24
<i>pisoniae</i> Cook, 1909 (<i>Cecidomyia</i>)	23.39
<i>pisoniae</i> (Felt), 1912 (<i>Bruggmanniella</i>), <i>Bruggmannia</i>	23.16
<i>pisonifolia</i> (Felt), 1912 (<i>Uleela</i>), <i>Bruggmannia</i>	23.16
<i>Pisoniomyia</i> Möhn, 1960	23.17
<i>Pisphondylia</i> Möhn, 1960	23.17
<i>Placochela</i> Rübсаamen, 1916	23.17
<i>Plesioidiplosis</i> Kieffer, 1912	23.36
<i>plumipes</i> Karsch, 1877, <i>Haplusia</i>	23.6
<i>polymera</i> (Alexander), 1936 (<i>Feltomyia</i>), <i>Feltomyina</i>	23.4
<i>Polystepha</i> Kieffer, 1897	23.16
<i>Polystephini</i>	23.16
<i>Porricondylinae</i>	23.3
<i>Porricondylini</i>	23.3
<i>portulacae</i> (Cook), 1906 (<i>Cecidomyia</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.24
<i>portulacae</i> (Felt), 1911 (<i>Lasioptera</i>), <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.24
<i>portulacae</i> Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.11
<i>praecox</i> Tavares, 1922, <i>Apodiplosis</i>	23.31
<i>Proasphondylia</i> Felt, 1915	23.18
<i>procera</i> Tavares, 1920, <i>Anadiplosis</i>	23.26
<i>producta</i> Felt, 1912, <i>Hyperdiplosis</i>	23.28
<i>Promikiola</i> Kieffer & Herbst, 1911	23.7
<i>prosopidis</i> Kieffer & Jorgensen, 1910, <i>Liebeliola</i>	23.34
<i>prosopidis</i> (Kieffer & Jørgensen), 1910 (<i>Rhopalomyia</i>), <i>Misopatha</i>	23.6
<i>Psadaria</i> Enderlein, 1940	23.2
<i>Pseudendaphis</i> Barnes, 1954	23.36
<i>pscudocalymmae</i> Möhn, 1964, <i>Neolasioptera</i> (<i>Neolasioptera</i>)	23.24
<i>pseudococci</i> Felt, 1921, <i>Diadiplosis</i>	23.33
<i>pseudococci</i> Felt, 1933, <i>Lobodiplosis</i>	23.35
<i>Pseudomikiola</i> Brèthes, 1917	23.8
<i>psychotriae</i> Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.11
<i>psychotriae</i> Möhn, 1960, <i>Bruggmannia</i>	23.16
<i>pulchra</i> Tavares, 1916, <i>Anadiplosis</i>	23.26
<i>pulchra</i> (Tavares), 1917 (<i>Eudiplosis</i>), <i>Schismatodiplosis</i>	23.29
<i>pulvinae</i> Felt, 1912 (<i>Mycodiplosis</i>)	23.38
<i>pustulans</i> Möhn, 1960, <i>Bruggmannia</i>	23.16
<i>ramosina</i> Tavares, 1917, <i>Baccharomyia</i>	23.19
<i>randiae</i> Möhn, 1959, <i>Asphondylia</i> (<i>Asphondylia</i>)	23.11
<i>randiae</i> Möhn, 1960, <i>Bruggmannia</i>	23.17
<i>reticulata</i> (Felt), 1915 (<i>Oxasphondylia</i>), <i>Zalepidota</i>	23.15
<i>reticulata</i> Möhn, 1960, <i>Perasphondylia</i>	23.14
<i>Rhoasphondylia</i> Möhn, 1960	23.15

Riveraella Kieffer & Herbst, 1911	23.8
Rochadiplosis Tavares, 1917	23.36
rochae Tavares, 1918, Asphondylia	23.13
rochae Tavares, 1925, Houardodiplosis	23.27
rondeletiae Möhn, 1959, Asphondylia (Asphondylia)	23.11
rotunda Möhn, 1964, Brachylasioptera	23.20
rubiae (Tavares), 1918 (Eudiplosis), Schismatodiplosis	23.29
rubida Felt, 1911, Toxomyia	23.37
rubra Kieffer & Herbst, 1911, Promikiola.	23.8
ruebsaameni Kieffer, 1913, Uleela	23.27
Ruebsaamenodiplosis Tavares, 1920	23.26
ruelliae Möhn, 1959, Asphondylia (Asphondylia)	23.11
salvadorensis Möhn, 1959, Asphondylia (Asphondylia)	23.11
salvadorensis Möhn, 1960, Heterasphondylia	23.34
salvadorensis Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
salvadorensis Möhn, 1960, Pisphondylia	23.17
salvadorensis Möhn, 1960, Polystepha	23.16
salvadorensis Möhn, 1963, Trotteria	23.25
salviae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
salviae Möhn, 1960, Sciasphondylia	23.15
Scheueria Kieffer & Herbst, 1909	23.8
Schismatodiplosis Rübsaamen, 1916	23.28
Schizodiplosis Kieffer, 1913	23.36
Schizomyia Kieffer, 1889	23.18
Schizomyiini	23.16
Sciasphondylia Möhn, 1960	23.15
schwarzi (Felt), 1911 (Lasiopteryx), Janetella	23.7
scopariae Möhn, 1959, Asphondylia (Asphondylia)	23.11
Scopodiplosis Felt, 1915	23.36
senecionis Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
serjaneae Rübsaamen, 1916, Haplopalpus	23.6
serjaniae Möhn, 1959, Asphondylia (Asphondylia)	23.11
serjaniae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
serjaniae Möhn, 1960, Placochela	23.18
serrata Möhn, 1964, Neolasioptera (Dilasioptera)	23.25
sexdentatus Kieffer & Jörgensen, 1910, Tetradiplosis	23.37
siccae Felt, 1908, Asphondylia	23.13
sidae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
silvestrii (Kieffer), 1910 (Arthrocnodax), Silvestrina	23.37
Silvestrina Kieffer, 1912	23.37
similis Möhn, 1960 Pisoniamyia	23.17
sorghicola (Coquillett), 1899 (Diplosis), Contarinia	23.29
speciosa Felt, 1913, Astrodiplosis	23.31
speciosa Felt, 1915, Scopodiplosis	23.36
spgazzeni Brèthes, 1914, Austrolauthia	23.4
Sphaerodiplosis Rübsaamen, 1916	23.30
spinosa Felt, 1909, Lobodiplosis	23.35
spinulosa Kieffer & Herbst, 1909, Angeliomyia	23.4
squamosa Tavares, 1918, Metasphondylia	23.19
stachytarpheta Barnes, 1932, Asphondylia	23.13
stachytarphetae Barnes, 1932, Schizomyia	23.18
sterculiae Rübsaamen, 1916, Megaulus	23.35
Stephomyia Tavares, 1916	23.18



struthanthi (Rübsaamen), 1916 (Gisonobasis) Asphondylia	23.13
styracis Tavares, 1918, Dialeria	23.20
<i>Styracodiplosis</i> Tavares, 1918	23.37
<i>Styraxdiplosis</i> Tavares, 1915	23.37
subinermis (Kieffer & Herbst), 1911 (Perrisia), Dasineura	23.5
sulphurea Tavares, 1909, Asphondylia	23.13
suaedicola Kieffer & Jörgensen, 1910, Asphondylia	23.13
tabernaemontanae Möhn, 1959, Asphondylia (Asphondylia)	23.11
talini Möhn, 1959, Asphondylia (Asphondylia)	23.11
tavaresi (Kieffer), 1913 (Ozobia), Zalepidota	23.16
Tavaresomyia Möhn, 1961	23.15
Tetradiplosis Kieffer & Jörgensen, 1910	23.37
thevetiae Möhn, 1959, Asphondylia (Asphondylia)	23.11
Thripsobremia Barnes, 1930	23.27
tibouchinae Tavares, 1917, Rochadiplosis	23.36
tithoniae, Möhn, 1960, Asphondylia (Asphondylia)	23.12
torreana Cook, 1906 (Cecidomyia)	23.39
tournefortiae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
tournefortiae (Rübsaamen), 1916 (Gisonobasis), Asphondylia (Asphondylia)	23.12
townsendi Felt, 1912, Karschomyia	23.34
Toxomyia Felt, 1911	23.37
trichilliae Möhn, 1959, Asphondylia (Asphondylia)	23.12
Trichopteromyia Williston, 1896	23.2
tricyclae (Kieffer & Jörgensen), 1910 (Rhopalomyia), Misospatha	23.6
tridentifera (Kieffer & Jörgensen), 1910, Neolasioptera (Neolasioptera)	23.19
trilobata Möhn, 1964, Alycaulus	23.19
Trisopsis Kieffer, 1912	23.37
tristis Tavares, 1922, Compsodiplosis	23.32
trixidis Möhn, 1959, Asphondylia (Asphondylia)	23.12
Trotteria Kieffer, 1901	23.25
Uleela Rübsaamen, 1908	23.26
ulei Rübsaamen, 1908, Asphondylia	23.14
ulei Rübsaamen, 1916, Macroporpa	23.17
Uleia Rübsaamen, 1905	23.8
umbra Walker, 1856 (Cecidomyia)	23.38
urostigmatis Tavares, 1917 (Asteromyia)	23.38
urostigmatis Tavares, 1917, Calmonia	23.4
urvilleanae (Tavares), 1909 (Lasioptera), Neolasioptera (Neolasioptera)	23.24
venusta Tavares, 1916, Anadiplosis	23.26
verbenae (Kieffer & Jörgensen), 1910 (Rhopalomyia), Misospatha	23.6
verbesinae Möhn, 1964, Neolasioptera (Neolasioptera)	23.24
vernoniensis Möhn, 1964, Neolasioptera (Neolasioptera)	23.25
vincenti Felt, 1911, Asphondylia (Asphondylia)	23.12



waltheriae Möhn, 1959, Asphondylia (Asphondylia)	23.12
watsoni Felt, 1915, Ctenodactylomyia	23.32
Winnertzia Rondani, 1860	23.3
Winnertzini	23.3
Xenasphondylia Felt, 1915	23.15
ximeniae Möhn, 1959, Asphondylia (Asphondylia)	23.12
Xyloperrisia Kieffer, 1913	23.8
xylosmatis Möhn, 1959, Asphondylia (Asphondylia)	23.12
zacatechichi Möhn, 1960, Asphondylia (Asphondylia)	23.12
Zalepidota Rübsaamen, 1908	23.15
zexmeniae Möhn, 1960, Ameliomyia	23.16
zexmeniae Möhn, 1960, Asphondylia (Asphondylia)	23.12

HOST INDEX

ANIMAL HOSTS


Acarina, undet	23.30
Coccidae, sp. of	23.33
"coccid"	23.38
Dysmicoccus brevipesus	23.34,35
Eriophyes sp.	23.30
Lecanoideus giganteus	23.32
Liothrips urichi	23.27
"mealy bugs"	23.38
Pinnaspis temporaria	23.30
Planococcus citri	23.34
Protopulvinaria pyriformis	23.38
Pseudaulacaspis pentagona	23.37
Pulvinaria urbicola	23.35
Saccharicoccus sacchari	23.38
Saissetia coffeae	23.35, 36,38
S.nigra	23.33
Tetranychus bimaculatus	23.30
T.tumidus	23.30
Toxoptera aurantii	23.36

PLANT HOSTS

Achatocarpus nigricans (Amaranthaceae)	23.9
Aegiphila arborescens (Verbenaceae)	23.29
Aeschynomene montevidensis (Leguminosae)	23.21
Agave sp. (Agavaceae)	23.23
Amaranthus spinosus (Amaranthaceae)	23.21
Anacardiaceae, sp. of.	23.14
Andira inermis (Leguminosae)	23.9
Andira sp. (Leguminosae)	23.31




<i>Anona cherimola</i> (Anonaceae)	23.9
<i>Aphelandra deppeana</i> (Acanthaceae)	23.21
<i>Ardisia paschalis</i> (Myrsinaceae)	23.9
<i>Aspidosperma quebrachoblanco</i> (Apocynaceae)	23.18
<i>Asteriscium polycephalum</i> (Umbelliferae)	23.23
<i>Atriplex canescens</i> (Chenopodiaceae)	23.13
<i>Avicennia nitida</i> (Verbenaceae)	23.38
<i>Ayenia pusilla</i> (Sterculiaceae)	23.12,25
<i>Baccharis bogotensis</i> (Compositae)	23.5
<i>B.coridifolia</i> (Compositae)	23.25
<i>B.dracunculifolia</i> (Compositae)	23.15
<i>B.eupatorioides</i> (Compositae)	23.8
<i>B.genistelloides</i> (Compositae)	23.19
<i>B.rosmarinifolia</i> (Compositae)	23.5
<i>B.salicifolia</i> (Compositae)	23.5,13,24
<i>B.schultzii</i> (Compositae)	23.15
<i>B.subulata</i> (Compositae)	23.23
<i>B.trinervis</i> (Compositae)	23.15
<i>Baccharis</i> sp. (Compositae)	23.13
<i>Bidens squarrosa</i> (Compositae)	23.9
Bignoniaceae, sp. of.	23.24
<i>Boerhaavia erecta</i> (Nyctaginaceae)	23.9,25
<i>Borreria verticillata</i> (Rubiaceae)	23.22
<i>Borreria</i> sp. (Rubiaceae)	23.13
<i>Brickellia pacayensis</i> (Compositae)	23.22
<i>Buddleia americana</i> (Loganiaceae)	23.7
<i>Buddleia racemosa</i> (Loganiaceae)	23.7
<i>Bursera grummifera</i> (Burseraceae)	23.17
<i>Calea integrifolia</i> (Compositae)	23.9,22,24
<i>C.zacatechichi</i> (Compositae)	23.12,23,25
<i>Capsicum baccatum</i> (Solanaceae)	23.22
<i>Casearia banquitana</i> (Samydaceae)	23.12
<i>Cattleya</i> spp. (Orchidaceae)	23.28
<i>Cecropia obtusa</i> (Urticaceae)	23.38
<i>Celtis iguanea</i> (Urticaceae)	23.22
<i>Cereus setaceus</i> (Cactaceae)	23.22
<i>Cestrum aurantiacum</i> (Solanaceae)	23.22
<i>C.lanatum</i> (Solanaceae)	23.22
<i>C.nocturnum</i> (Solanaceae)	23.10
<i>Chlorophora tinctoria</i> (Urticaceae)	23.27
<i>Cissampelos pareira</i> (Menispermaceae)	23.22
<i>Cissus sicyoides</i> (Vitaceae)	23.10
<i>Citharexylum quadrangulare</i> (Verbenaceae)	23.13
citrus	23.3
<i>Clematis dioica</i> (Ranunculaceae)	23.10,22
<i>C.drummondii</i> (Ranunculaceae)	23.10
<i>C.grossa</i> (Ranunculaceae)	23.10
<i>Clerodendron aculeatum</i> (Verbenaceae)	23.13
<i>Clusia</i> sp. (Guttiferae)	23.8
<i>Coccoloba laurifolia</i> (Polygonaceae)	23.32
<i>C.uivifera</i> (Polygonaceae)	23.38
<i>Coffea liberica</i> (Rubiaceae)	23.28



<i>Colliguaya odorifera</i> (Euphorbiaceae)	23.8
<i>Combretum farinosum</i> (Combretaceae)	23.22
<i>C.leprosum</i> (Combretaceae)	23.27
Compositae, sp. of.	23.21,28
<i>Cordia alba</i> (Boraginaceae)	23.10
<i>C.cana</i> (Boraginaceae)	23.22
cotton	23.3
<i>Coussapoa</i> sp. (Urticaceae)	23.33
<i>Crotalaria longirostrata</i> (Leguminosae)	23.11
<i>C.mucronata</i> (Leguminosae)	23.9
<i>Croton hemiargyreus</i> (Euphorbiaceae)	23.37
<i>Curatella americana</i> (Dilleniaceae)	23.10
<i>Dalbergia</i> sp. (Leguminosae)	23.26,27
<i>Dalechampia ficifolia</i> (Euphorbiaceae)	23.22
<i>Diodia</i> sp. (Rubiaceae)	23.13
<i>Erigeron canadensis</i> (Compositae)	23.22
<i>Erythroxylum mexicanum</i> (Linaceae)	23.10,23
<i>Eugenia buxifolia</i> (Myrtaceae)	23.5
<i>Eugenia</i> sp. (Myrtaceae)	23.18
<i>Eupatorium morifolium</i> (Compositae)	23.21
<i>E.odoratum</i> (Compositae)	23.14
<i>E.villosum</i> (Compositae)	23.39
<i>Eupatorium</i> sp. (Compositae)	23.10,14,21,22,28
<i>Euphorbia serpens</i> (Euphorbiaceae)	23.4
<i>Ficus</i> sp. (Moraceae)	23.7,28,39
<i>Gardoquia gilliesi</i> (Labiatae)	23.5
<i>Gaultheria vernalis</i> (Ericaceae)	23.8
<i>Gilibertia arborea</i> (Araliaceae)	23.20
<i>Godmania aesculifolia</i> (Bignoniaceae)	23.10
<i>Gossypium</i> spp. (Malvaceae)	23.29
<i>Gcurllea decorticans</i> (Leguminosae)	23.31
<i>Grabowskiya obtusa</i> (Solanaceae)	23.33
<i>Guarea trichiloides</i> (Meliaceae)	23.5
<i>Guarea</i> sp. (Meliaceae)	23.10
<i>Gutierrezia microcephala</i> (Compositae)	23.10
<i>G.sarothrae</i> (Compositae)	23.19
<i>Gymnosperma glutinosa</i> (Compositae)	23.19
<i>Heisteria cyanocarpa</i> (Olacaceae)	23.33
<i>Helicteres guazumaefolia</i> (Sterculiaceae)	23.10
<i>Heliocarpus donnellsmithii</i> (Tiliaceae)	23.23
<i>Heliotropium curassavicum</i> (Boraginaceae)	23.24
<i>Heterothalamus spartioides</i> (Compositae)	23.23
<i>Hydrangea scandens</i> (Saxifragaceae)	23.4
<i>Hymenostephium cordatum</i> (Compositae)	23.11,22
<i>Hyptis suaveolens</i> (Labiatae)	23.23
<i>H.verticillata</i> (Labiatae)	23.10



<i>Indigofera suffruticosa</i> (Leguminosae)	23.10, 23.25
<i>Inga leptoloba</i> (Leguminosae)	23.23
<i>I. spuria</i> (Leguminosae)	23.23
<i>Ipomoea arborescens</i> (Convolvulaceae)	23.12
<i>Ipomoea</i> sp. (Convolvulaceae)	23.18,20
<i>Iresine calea</i> (Amaranthaceae)	23.20
<i>I. celosia</i> (Amaranthaceae)	23.23
<i>Jatropha curcas</i> (Euphorbiaceae)	23.10
<i>Jussieua linifolia</i> (Onagraceae)	23.12
<i>J. suffruticosa</i> (Onagraceae)	23.12
<i>Jussieua</i> sp. (Onagraceae)	23.13
<i>Laelia</i> spp. (Orchidaceae)	23.28
<i>Lantana camara</i> (Verbenaceae)	23.9,22
<i>Lantana</i> sp. (Verbenaceae)	23.20,29
Lauraceae, sp. of.	23.17
Leguminosae, sp. of.	23.26
<i>Lippia cardiostegia</i> (Verbenaceae)	23.10
<i>L. foliosa</i> (Verbenaceae)	23.7
<i>Lippia</i> sp. (Verbenaceae)	23.8
<i>Lonchocarpus minimiflorus</i> (Leguminosae)	23.10
<i>Lycium chilense</i> (Solanaceae)	23.6,7,32
<i>L. gracile</i> (Solanaceae)	23.7,34
<i>Lycopersicum esculentum</i> (Solanaceae)	23.29
<i>Machaerium</i> sp. (Leguminosae)	23.26,35
Malpighiaceae, sp. of.	23.17
Malvaceae, sp. of.	23.19
<i>Malvaviscus arboreus</i> (Malvaceae)	23.11
<i>Mangifera indica</i> (Anacardiaceae)	23.33
mango	23.3
<i>Manihot aipi</i> (Euphorbiaceae)	23.28
<i>M. dichotoma</i> (Euphorbiaceae)	23.28
<i>M. utilissima</i> (Euphorbiaceae)	23.7,18,28
<i>Marcetia</i> sp. (Melastomataceae)	23.29
<i>Medicago sativa</i> (Leguminosae)	23,29
<i>Melanthera nivea</i> (Compositae)	23.11,23
<i>Merremia umbellata</i> (Convolvulaceae)	23.23
<i>Mikania amara</i> (Compositae)	23.14
<i>M. scandens</i> (Compositae)	23.19
<i>Mikania</i> sp. (Compositae)	23.14,19
<i>Mimosa albida</i> (Leguminosae)	23.14
<i>M. caesalpiniaefolia</i> (Leguminosae).	23.15
<i>Mimosa</i> sp. (Leguminosae)	23.26,30
<i>Myrceugenia stenophylla</i> (Myrtaceae)	23.6
Myrtaceae, sp. of.	23.6
<i>Myrsine</i> sp. (Myrsinaceae)	23.16
<i>Nectandra megapotamica</i> (Lauraceae)	23.6
<i>Neea</i> sp. (Nyctaginaceae)	23.26,27



<i>Ocimum micranthum</i> (Labiatae)	23.11
<i>Odontonema stricta</i> (Acanthaceae)	23.24
<i>Opuntia</i> sp. (Cactaceae)	23.13
<i>Ossaea</i> sp. (Melastomataceae)	2335
Papilionaceae, sp. of.	23.29
<i>Paspalum conjugatum</i> (Gramineae)	23.27
<i>Pernettya furens</i> (Ericaceae)	23.7
<i>Phyllanthus distichus</i> (Euphorbiaceae)	23.13
<i>Physalis viscosa</i> (Solanaceae)	23.25
<i>Piper</i> sp. (Piperaceae)	23.14,15,16
<i>Piriqueta ovata</i> (Turneraceae)	23.24
<i>Pisonia aculeata</i> (Nyctaginaceae)	23.17
<i>P. macranthocarpa</i> (Nyctaginaceae)	23.6,17
<i>P. nigricans</i> (Nyctaginaceae)	23.16
<i>Pisonia</i> sp. (Nyctaginaceae)	23.39
<i>P. macranthocarpa</i> (Nyctaginaceae)	23.16,17
<i>P. nigricans</i> (Nyctaginaceae)	23.16
<i>Pisonia</i> sp. (Nyctaginaceae)	23.39
<i>Polypodium crassifolium</i> (Polypodiaceae)	23.26
<i>Porophyllum</i> sp. (Compositae)	23.19
<i>Portulaca oleracea</i> (Portulacaceae)	23.11,24
<i>Prosopis alpataco</i> (Leguminosae)	23.6,37
<i>P. campestris</i> (Leguminosae)	23.37
<i>P. strombulifera</i> (Leguminosae)	23.23,34
<i>Protium heptaphyllum</i> (Bursaceae)	23.5,20
<i>Pseudocalymma macrocarpa</i> (Bignoniaceae)	23.20, 21,23,24
<i>Psychotria carthaginensis</i> (Rubiaceae)	23.16
<i>P. pubescens</i> (Rubiaceae)	23.11
<i>Psychotria</i> sp. (Rubiaceae)	23.31
<i>Puccinia</i> sp. (Basidiomycetes)	23.37
<i>Quercus eugeniaefolia</i> (Fagaceae)	23.16
<i>Q. hondurensis</i> (Fagaceae)	23.16
<i>Randia aculeata</i> (Rubiaceae)	23.11
<i>R. spinosa</i> (Rubiaceae)	23.17
<i>Rondeletia strigosa</i> (Rubiaceae)	23.11
Rubiaceae, sp. of.	23.13,29,34
<i>Ruellia albicaulis</i> (Acanthaceae)	23.11
<i>Salvia occidentalis</i> (Labiatae)	23.24
<i>Salvia</i> sp. (Labiatae)	23.15
<i>Scoparia culcis</i> (Scrophulariaceae)	23.11
<i>Senecio kermesinus</i> (Compositae)	23.24
<i>S. mendocinus</i> (Compositae)	23.7
<i>Serjanea gonlocarpa</i> (Sapindaceae)	23.11,18
<i>S. racemosa</i> (Sapindaceae)	23.24
<i>Serjanea</i> sp. (Sapindaceae)	23.6
<i>Sida rhombifolia</i> (Malvaceae)	23.24
<i>Smilax</i> sp. (Liliaceae)	23.13,32
<i>Solanum umbellatum</i> (Solanaceae)	23.23



<i>Solanum</i> sp. (Solanaceae)	23.29
<i>Solanum</i> sp. (Solanaceae)	23.29
<i>Solidago</i> spp. (Compositae)	23.13
<i>Sorghum</i> spp. (Gramineae)	23.29
<i>Sorocea illicifolia</i> (Urticaceae)	23.14
<i>Stachytarpheta cajanensis</i> (Verbenaceae)	23.13,18
<i>S.dichotoma</i> (Verbenaceae)	23.28
<i>Sterculia</i> sp. (Sterculiaceae)	23.35
<i>Struthanthus marginatus</i> (Loranthaceae)	23.11
<i>Struthanthus</i> sp. (Loranthaceae)	23.13
<i>Styrax</i> sp. (Styracaceae)	23.20,32,37
<i>Suaeda divaricata</i> (Chenopodiaceae)	23.13
<i>Tabernaemontanae amydaelifolia</i> (Apocynaceae)	23.11
<i>Teucrium inslatum</i> (Labiatae)	23.25
<i>Thevetia plumeriaefolia</i> (Apocynaceae)	23.11
<i>Tibouchina</i> sp. (Melastomataceae)	23.36
<i>Tithonia rotundifolia</i> (Compositae)	23.12
<i>Tournefortia angustiflora</i> (Boraginaceae)	23.12
<i>T.volubilis</i> (Boraginaceae)	23.12,24
<i>Trichilia</i> sp. (Meliaceae)	23.12
<i>Tricycla spinosa</i> (Nyctaginaceae)	23.6
<i>tridax procumbens</i> (Compositae)	23.22
<i>Trixis radialis</i> (Compositae)	23.12
<i>Uromyces pisi</i> (Basidiomycetes)	23.37
<i>Urostigma</i> sp. (Moraceae).	23.4,38
<i>Urvillea ulmacea</i> (Sapindaceae)	23.24
<i>Verbena aspera</i> (Verbenaceae)	23.6
<i>V.seriphoides</i> (Verbenaceae)	23.6
<i>Verbesina fraseri</i> (Compositae)	23.24
<i>V.sublobata</i> (Compositae)	23.24
<i>V.turbacensis</i> (Compositae)	23.24
<i>Vernonia canescens</i> (Compositae)	23.9,10
<i>Vernonia</i> sp. (Compositae)	23.24
<i>Waltheria indica</i> (Sterculiaceae)	23.12
<i>Ximenia americana</i> (Olacaceae)	23.12
<i>Xylosma flexuosum</i> (Flacourtiaceae)	23.12
<i>Zexmenia frutescens</i> (Compositae)	23.12,16







MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

23a

FAMILY PERISSOMMATIDAE

NELSON BERNARDI

Museu de Zoologia, Universidade de São Paulo

This small family was described by Colless (1962) to include the new Australian genus *Perissomma*. In the original description, Colless included two species, and in 1969 he described three additional species, bringing the number of known species to five, namely *fusca*, *mcalpinei*, *dobrotworskyi*, *bellissima* and *congrua*. The first four are Australian, the fifth Chilean. The position of the Perissommatidae among the Nematocera presents as yet some problems, but it was included by its author in the Bibionomorpha, a procedure accepted by Hennig (1968). The two works of Colless (1962, 1969) include illustrations of adults and immatures, besides morphological, biological and cytogenetic information. The only Chilean species was reported by Colless to "differ rather markedly from all the Australian species, although it is clearly congeneric with them". Adults of *P. fusca* have been collected in or near pine forests, those of *P. mcalpinei* are supposed to be natives of rain forest and wet sclerophyll in cool climates; *P. dobrotworskyi* was collected in dry forest. Larvae of *P. fusca* have been found in fungi of the genus *Boletus*, which is commonly associated with *Pinus radiata*; those of *P. mcalpinei* were found to live in leaf litter and have no apparent association with fungi.

Genus PERISSOMMA Colless

PERISSOMMA Colless, 1962: 520. Type-species, *fusca* Colless, 1962, from Australia (orig. des.).

CONGRUA Colless, 1969: 727, figs. 1A (dorsal view of male terminalia), 4C (wing) (holotype male; female unknown). Type-locality: Chile, Cautin, Los Paraguas.

LITERATURE CITED

COLLESS, D. H.

1962. A new genus and family of Diptera (Nematocera: Perissommatidae). *Austr. J. Zool.* 10: 519-535, 4 figs.

23A1



1969. The genus *Perissomma* (Diptera: Perissomatidae), with new species from Australia and Chile. *Austr. J. Zool.* 17: 719-728, 4 figs.

HENNIG, W.

1968. Kritische Bemerkungen über den Bau der Flügelwurzel bei den Dipteren und die Frage nach der Monophylie der Nematocera. *Stutt. Beitr. Naturk.* 193: 1-23, 13 figs.



24

SUBORDER BRACHYCERA





SUPERFAMILY TABANOIDEA

Family Xylophagidae

Family Xylomyidae

Family Stratiomyidae

Family Pelecorhynchidae

Family Tabanidae

Family Rhagionidae

Family Pantophthalmidae





MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

24

FAMILY XYLOPHAGIDAE

MAURICE T. JAMES

*Department of Entomology, Washington
State University, Pullman, Washington*

The limits of this family, as treated here, are the same as those accepted for the Nearctic catalogue. As so defined, the taxon is an unsatisfactory one, as shown by Stuckenberg (1973), who believes, for example, that *Atherimorpha* should be placed in the Rhagionidae. However, with the present state of our knowledge, no better arrangement is available.

Adults frequent forested or wooded areas and feed on sap, nectar of flowers, or other liquid matter. Life histories and immature stages of the Neotropical forms are virtually unknown, but in parts of the world where such information is available, larvae are predators or scavengers, occurring in soil rich in decaying vegetable matter, under bark of trees, or in decaying logs.

Genus ATHERIMORPHA White

ATHERIMORPHA White, 1915:41. Type-species, *vernalis* White (mon.). Ref. — Malloch, 1932:206 (key).

Philippoleptis Malloch, 1931:276 (as subgenus of *Atherimorpha*). Type-species, *Leptis praefica* Philippi (orig. des.).

ALBOHIRTA Malloch, 1932:208. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.

ATRIFEMUR Malloch, 1932:210. Type-locality: Chile, Llanquihue, Casa Pangué. Distr. — Chile.

CLARIPENNIS (Philippi), 1865:772 (*Leptis*). Type-locality: Chile. Distr. — Chile.

EDWARDSI Malloch, 1932:212. Type-locality: Chile, Puerto Montt. Distr. — Chile.

FLAVOLATERALIS Malloch, 1932:213. Type-locality: Argentina, Rio Negro, Bariloche. Distr. — Argentina.



- FUSCA Malloch, 1932:213. Type-locality: Chile, Chiloé Island, Puntra. Distr. — Chile.
- FUSCICOXA Malloch, 1932:215. Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- GRISEA (Philippi), 1865:774 (*Leptis*). Type-locality: Chile. Distr. — Chile.
- HIRTULA (Bigot), 1887:116 (*Leptis*). Type-locality: Chile. Distr. — Chile.
- IMITANS Malloch, 1932:211. Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile.
- NEMORALIS (Philippi), 1865:772 (*Leptis*). Type-locality: Chile, Valdivia. Distr. — Chile.
- PRAEFICA (Philippi), 1865:772 (*Leptis*). Type-locality: Chile. Distr. — Chile. Ref. — Malloch, 1932:232 (redescr.).
- macrochaeta* (Bigot), 1889:325 (*Psilocephala*). Type-locality: Chile.
- pilosula* (Bigot), 1889:326 (*Psilocephala*). Type-locality: Chile.
- praefita*, error.
- praefixa*, error.
- SCUTELLARIS Malloch, 1932:214. Type-locality: Chile, Concepción. Distr. — Chile. May be female of *grisea* (Philippi) (Edwards, footnote to Malloch, 1932:214).
- TRIANGULARIS Malloch, 1932:234. Type-locality: Chile, Valparaiso, Perales. Distr. — Chile.

Genus AUSTROLEPTIS Hardy

- AUSTROLEPTIS Hardy, 1919:126. Type-species, *rhyphoides* Hardy (orig. des.).
- ATRICEPS Malloch, 1932:203. Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.
- FULVICEPS Malloch, 1932:202. Type-locality: Chile, Llanquihue, Casa Pangue. Distr. — Chile.

Genus HETEROSTOMUS Bigot

- HETEROSTOMUS Bigot, 1857:283. Type-species, *curvipalpis* Bigot (mon.).
- Heterostomyia* Bigot, 1879:186 (*nom. nov.* for *Heterostomus* Bigot, erroneously stated to be preoccupied).
- Heterostomatomyia* (emend).
- CURVIPALPIS Bigot, 1857:285. Type-locality: Chile. Distr. — Chile.
- inflexipalpis* (Bigot), 1879:186 (*Heterostomyia*). *Nom. nud.*
- flexipalpis* (Bigot), 1879:186 (*Heterostomyia*). *Nom. nud.*

Genus RACHICERUS Walker

- RACHICERUS Walker, 1854:103. Type-species: *fulvicollis* Walker (mon.). Ref. — Carrera, 1945 (key).
- Rhachicerus* (emend).
- BELLUS Osten Sacken, 1886:62. Type-locality: Panama, Bugaba. Distr. — Panama. Ref. — Carrera, 1945 (quotes orig. descr.).
- LANEI Carrera, 1940:389. Type-locality: Brazil, São Paulo, Alto da Serra. Distr. — Brazil. Ref. — Carrera, 1945:128 (further descr.).
- LOPESI Carrera, 1940:393. Type-locality: Brazil, Rio de Janeiro, Corcovado. Distr. — Brazil.



- MARCUSI Carrera, 1940:395. Type-locality: Brazil, São Paulo, Alto da Serra. Distr. — Brazil. Ref. — Carrera, 1945 (further descr.).
- NIGRIPALPIS Loew, 1874:378. Type-locality: Mexico. Distr. — Mexico. Ref. — Carrera, 1945 (quotes orig. descr.).
- OLIVERIOI Carrera, 1940:391. Type-locality: Brazil, São Paulo, Bananal. Distr. — Brazil. Ref. — Carrera, 1945 (further descr.).
- PICTICORNIS Kertész, 1923:89. Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica. Ref. — Carrera, 1945 (quotes orig. descr.).
- SHANNONI Carrera, 1945:125. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- VARIPES Loew, 1863:5 (Centuria 3, no. 7). Type-locality: Cuba. Distr. — Cuba. Ref. — Carrera, 1945 (quotes orig. descr.).

Unrecognized species

- VETUSTUS Walker, 1854:106 (*Xylophagus*). Type-locality: Brazil, Pará. Distr. — Brazil. (Generic reference with a query; most certainly not a *Xylophagidae*).

APPENDIX

Genus STENIMANTIA Enderlein

- STENIMANTIA Enderlein, 1932:270. Type-species, *Xylophagus carbonarius* Philippi (orig. des.).
- CARBONARIA (Philippi), 1865:727 (*Xylophagus*). Type-locality: "Chile". Distr. — Chile. Note: This genus should be transferred to the Chiromyzinae, in the Stratiomyidae (Fasc. 26); Hennig (1955:427) saw the specimen studied by Enderlein, and assigned the genus to the Chiromyzinae.

LITERATURE CITED

BIGOT, J. M. F.

1857. Diptères nouveaux provenant du Chili. *Ann. Soc. Entomol. France* (3) 5:277-308.
1879. Diptères nouveaux ou peu connus. 11^e Partie. XVI. Curiae Xylophagidarum et Stratiomyidarum. *Ann. Soc. Entomol. France* (5) 9:183-234.
1887. Diptères nouveaux ou peu connus. Leptidi, Muscidi. *Bull. Soc. Zool. France* 12:97-118, 581-617.
1889. Diptères nouveaux ou peu connus. 35^e Partie. XLIV: Therevidi. *Ann. Soc. Entomol. France* (6) 9:321-328.

CARRERA, M.

1940. Novas espécies do gênero *Rachicerus* Walker, 1854 (Dipt.). *Arq. Zool. Est. São Paulo* 1:387-394.
1945. Contribuição ao conhecimento do gênero *Rachicerus* Walker, 1854 (Diptera, Rachiceridae Handlirsch 1908). *Papéis Avulsos Dept. Zool. São Paulo* 5:119-134.



- ENDERLEIN, G.
1932. Zwei biogeographisch interessante neue Xylophagidengattungen der südlichen Hemisphär. *Zool. Anz.* 99:269-271.
- HARDY, G. H.
1919. Australian Rhyphidae and Leptidae. *Pap. Roy. Soc. Tasmania* 1919:117-129.
- HENNIG, W.
1935. Bemerkungen zur Synonymie einiger Gattungen der niederen Brachycera. *Beitr. z. Ent.* 5(3-4):426-428.
- KERTÉSZ, K.
1923. Vorarbeiten zu einer Monographie der Notacanthen. XLV-L. *Ann. Mus. Nat. Hungarici* 20:85-129.
- LOEW, HERMANN
1863. *Diptera Americae septentrionalis indigena. Centuria tertia.* *Berl. Entomol. Ztschr.* 7:1-55.
1874. Neue Nordamerikanische Diptera. *Berl. Entomol. Ztschr.* 18: 378-384.
- MALLOCH, J. R.
1931. Notes on Australian Diptera. XXVIII. *Proc. Linn. Soc. New South Wales* 56:273-276.
1932. Rhagionidae, Therevidae, Lonchopteridae, in British Museum (Natural History): *Diptera of Patagonia and South Chile* 5(3):199-235.
- OSTEN SACKEN, C. R.
1886. *Diptera*, Vol. I (part, concl.), in Godman, F. D., and Salvin, O., eds., *Biologia Centrali-Americana, Insecta, Diptera* 1: 378 pp.
- PHILIPPI, R. A.
1865. Aufzählung der chilenischen Diptera. *Verh. zool.-bot. Ges. Wien* 15:595-782.
- STUCKENBERG, B. R.
1973. The Athericidae, a new family in the lower Brachycera (Diptera). *Ann. Natal. Mus.* 21:649-673.
- WALKER, F.
1854. *List of the specimens of dipterous insects in the collection of the British Museum.* Part 5, pp. 1-330. London.
- WHITE, A.
1915. The Diptera Brachycera of Tasmania. Part I. Families Lep-tidae, Stratiomyidae, Nemestrinidae, and Cyrtidae. *Papers & Proc. Roy. Soc. Tasmania* 1914 (1915):35-74.



INDEX

albohirta Malloch, 1932, Atherimorpha	24.1
Atherimorpha White, 1915	24.1
atriceps Malloch, 1932, Austroleptis	24.2
atrifemur Malloch, 1932, Atherimorpha	24.1
Austroleptis Hardy, 1916	24.2
bellus Osten Sacken, 1886, Rachicerus	24.2
carbonaria (Philippi), 1965 (Xylophagus), Stenimantia	24.3
claripennis (Philippi), 1865 (Leptis), Atherimorpha	24.1
curvipalpis Bigot, 1857, Heterostomus	24.2
edwardsi Malloch, 1932, Atherimorpha	24.1
flavolateralis Malloch, 1932, Atherimorpha	24.1
flexipalpis (Bigot), 1879 (Heterostomyia), Heterostomus	24.2
fulviceps Malloch, 1932, Austroleptis	24.2
fusca Malloch, 1932, Atherimorpha	24.2
fuscicoxa Malloch, 1932, Atherimorpha	24.2
grisea (Philippi), 1865 (Leptis), Atherimorpha	24.2
<i>Heterostomatomyia</i> , emend.	24.2
Heterostomus Bigot, 1857	24.2
<i>Heterostomyia</i> Bigot, 1857	24.2
hirtula (Bigot), 1887 (Leptis), Atherimorpha	24.2
imitans Malloch, 1932, Atherimorpha	24.2
inflexipalpis (Bigot), 1879 (Heterostomyia), Heterostomus	24.2
lanei Carrera, 1940, Rachicerus	24.2
lopesi Carrera, 1940, Rachicerus	24.2
macrochaeta (Bigot), 1889 (Psilocephala), Atherimorpha	24.2
marcusi Carrera, 1940, Rachicerus	24.3
nemoralis (Philippi), 1865 (Leptis), Atherimorpha	24.2
nigripalpis Loew, 1874, Rachicerus	24.3
oliverioi Carrera, 1940, Rachicerus	24.3
<i>Philippoleptis</i> Malloch, 1931	24.1
picticornis Kertész, 1923, Rachicerus	24.3
vilosula (Bigot), 1889 (Psilocephala), Atherimorpha	24.2
praefica (Philippi), 1865 (Leptis), Atherimorpha	24.2
<i>praefita</i> , error, Atherimorpha	24.2
<i>praefixa</i> , error, Atherimorpha	24.2
Rachicerus Walker, 1854	24.2
<i>Rhachicerus</i> , emend.	24.2



scutellaris Malloch, 1932, Atherimorpha	24.2
shannoni Carrera, 1945, Rachicerus	24.3
Stenimantia Enderlein, 1932	24.3
triangularis Malloch, 1932, Atherimorpha	24.2
varipes Loew, 1863, Rachicerus	24.3
vetustus Walker, 1854, Xylophagus	24.3







MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
 (Formerly Departamento de Zoologia, Secretaria da Agricultura, São Paulo)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
 SOUTH OF THE UNITED STATES

25

FAMILY XYLOMYIDAE

MAURICE T. JAMES

*Department of Entomology, Washington
 State University, Pullman, Washington*

The generic and subgeneric classification used here follows that of Nagatomi and Tanaka (1971), although a critical study of the Neotropical species may show that some are improperly placed. The genus *Arthropeina* Lindner is accepted as valid, as it is by Nagatomi and Tanaka, although Lindner (1969) synonymized it with *Xylomy(i)a*.

Except for *Solva pallipes*, the life histories and immature stages of the Neotropical Xylomyidae are unknown. In parts of the world where pertinent information is available, the larvae of known species live under bark of trees and are predators or scavengers.

Ref. — Nagatomi & Tanaka, 1971 (key to genera and subgenera of world).

Genus ARTHROPEINA Lindner

ARTHROPEINA Lindner, 1949:789. Type-species, *fulva* Lindner (mon.).
 FULVA Lindner, 1949:789. Type-locality: not stated, presumably Brazil.
 Distr. — Brazil.

Genus SOLVA Walker

Subula Meigen, 1820:15 (preocc. Schumacher, 1817). Type-species, *Xylophagus varius* Meigen (Rondani, 1856:172). A manuscript name cited in specific synonymy.

SOLVA Rondani, 1861:11 (nom. nov. for *Subula* Meigen). Type-species, *Xylophagus varius* Meigen (aut.).

Subulaomyia Williston, 1896 (as *Subula Omyia*; nom. nov. for *Subula* Meigen but deleted in *Corrigenda*, p. 4). Type-species, *Xylophagus varius* Meigen (aut.).

Subulonia Enderlein, 1913:545. Type-species, *truncativena* Enderlein (mon.).



Hanaia Enderlein, 1920:281. Type-species, *Xylophagus marginatus* Meigen (orig. des.).

Subgenus SOLVA Walker

- BRASILIANA (Lindner), 1949:791 (*Subulonia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FLAVICOXIS (Enderlein), 1921:172 (*Hanaia*). Type-locality: Paraguay, San Bernardino. Distr. — Brazil, Paraguay.
- PALLIPES (Loew), 1863:6 (Centuria 3, no. 9) (*Subula*). Type-localities: Illinois, Wisconsin. Distr. — Canada, U.S.A.; "Middle America".
- TRUNCATIVENA (Enderlein), 1913:545 (*Subulonia*). Type-locality: Costa Rica. Distr. — Costa Rica.

Subgenus MACROCEROMYS Bigot

- MACROCEROMYS Bigot, 1877:73. Type-species, *fulviventris* Bigot (mon.). *Xylomya* Auct., not Rondani.
- AMERICANA (Wiedemann), 1821:51 (*Xylophagus*). Type-locality: North America. Distr. — U.S.A.; Mexico.
- ELONGATA (Osten Sacken), 1886:62 (*Subula*). Type-locality: Guatemala, Capetillo. Distr. — Guatemala.
- FULVIVENTRIS Bigot, 1879:187. Type-locality: Mexico. Distr. — Mexico. *fulviventris* Bigot, 1899:1xxiii (*nomen nudum*).
- PRISTA Enderlein, 1913:541. Type-locality: Mexico, Chiapas. Distr. — Mexico.

LITERATURE CITED

- BIGOT, J. M. F.
1877. (Diagnoses de nouveaux-genres de Diptères). *Bull. Soc. Entomol. France* 1877:lxiii-lxiv.
1879. Diptères nouveaux ou peu connus. 11^e Partie. XVI. Curiae Xylophagidarum et Stratiomyidarum. *Ann. Soc. Entomol. France* (5)9:183-234.
- ENDERLEIN, G.
1913. Dipterologische Studien. V. Zur Kenntnis der Familie Xylophagidae. *Zool. Anz.* 42:532-552.
1920. Ord. Diptera, Fliegen Zweiflügler, pp. 265-315. In Brohmer, P., *Fauna von Deutschlands*, Ed. 2, 477 pp. Leipzig.
1921. Dipterologische Studien. Ueber die phyletische älteren Stratiomyiidensubfamilien (Xylophaginae, Chiromyzinae, Solvinae, Beridinae, und Coenomyiinae). *Mitt. Zool. Mus. Berlin* 10: 153-214.
- LINDNER, E.
1949. Neotropische Stratiomyiiden des Britischen Museum in London. *Ann. Mag. Nat. Hist.* (12)1:782-821, 851-891.




1969. Fünfter Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Stuttgarter Beiträge zur Naturkunde* 203:1-14.
- LOEW, H.
1863. *Diptera Americae septentrionalis indigena. Centuria tertia.* *Berl. Entomol. Ztschr.* 7:1-55.
- MEIGEN, J. W.
1820. *Systematische Beschreibung der bekannten europäischen zweiflügeligen Insekten.* Vol. 2, x+365 pp.
- NAGATOMI, A. & A. TANAKA
1971. The Solvidae of Japan. *Mushi* 45:101-146.
- OSTEN SACKEN, C. R.
1886. *Diptera*, Vol. 1 (part, concl.), in Godman, F. D., and Salvin, O., eds., *Biologia Centrali-Americana, Insecta-Diptera* 1:378 pp.
- RONDANI, C.
1856. *Dipterologia Italicae prodromus. Vol. 1. Genera Italica ordinis dipterorum ordinatim disposita et distincta et in familias et stirpas aggregata.* 228 pp. Parmae (Parma).
1861. *Dipterologia Italicae prodromus. Vol. 4. Species Italicae ordinis dipterorum in genera characteribus definita, ordinatim collectae, methodo analitica [sic!] distinctae, et novis vel minus cognitis descriptis. Pars tertia: Muscidae, Tachininarum complementum.* 174 pp. Parmae (Parma).
- WALKER, F.
1837. Descriptions of Diptera collected by Capt. P. P. King, R. N., F. R. S., in the survey of the Straits of Magellan. *Trans. Linn. Soc. London* 17:331-359.
1859. Catalogue of dipterous insects collected at Makassar in Celebes, by Mr. A. R. Wallace, with descriptions of new species. *Jour. Proc. Linn. Soc. London, Zool.* 4:90-144.
- WIEDEMANN, C. R. W.
1821. *Diptera exotica.* Pars I. Pp. xix+244 pp. Kiliae.
- WILLISTON, S. W.
1896. *Manual of the families and genera of North American Diptera.* Ed. 2, 167 pp. New Haven, Conn.

INDEX

americana (Wiedemann), 1821 (Xylophagus), Solva	25.2
Arthropeina Lindner, 1949	25.1
brasiliانا (Lindner), 1949 (Subulonia), Solva	25.2



<i>elongata</i> (Osten Sacken), 1886 (Subula), Solva	25.2
<i>flavicoxis</i> Enderlein, 1921 (Hanauia), Solva	25.2
<i>fulva</i> Lindner, 1949, Arthropeina	25.1
<i>fulviventris</i> Bigot, 1899, Solva	25.2
<i>fulviventris</i> Bigot, 1879, Solva	25.2
<i>Hanauia</i> Enderlein, 1920	25.2
<i>Macroceromys</i> Bigot, 1877	25.2
<i>pallipes</i> (Loew), 1863 (Subula), Solva	25.2
<i>prista</i> Enderlein, 1913, Solva	25.2
Solva Walker, 1859	25.1, 25.2
<i>Subula</i> Meigen, 1820	25.1
<i>Subulaomyia</i> Williston, 1896	25.1
<i>Subulonia</i> Enderlein, 1913	25.1
<i>truncativena</i> (Enderlein), 1913 (Subulonia), Solva	25.2
<i>Xylomya</i> Authors	25.2



MUSEU DE ZOOLOGIA, UNIVERSIDADE DE SÃO PAULO
(FORMERLY DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO)

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

26

FAMILY STRATIOMYIDAE

MAURICE T. JAMES

*Department of Entomology, Washington
State University, Pullman, Washington*

The soldier flies, or Stratiomyidae, occur throughout the warmer temperate and tropical regions of the world, but they are most common in moist to humid tropical climates. Adults are often found resting on vegetation, visiting flowers, or near over-ripe fruits or decaying vegetable materials. Many larvae are aquatic, many are plant scavengers, some are predatory, and a few attack living plants.

Though much work has been done on the Neotropical fauna, there have been no comprehensive studies except, to an extent, locally or on the subfamily level. Consequently, in the present catalogue many generic references must be considered as tentative, especially in such "catch-all" genera as *Beris*, *Sargus*, *Stratiomys*, and *Odontomyia*. Undoubtedly, many species referred to these genera do not belong there.

The subfamily placement of *Nemotelus* and *Cyphomyia* and related genera is at variance with that used in the Nearctic catalogue, but the present usage seems to be a more natural one. With some exceptions, divisions of subfamilies into tribes and of genera into subgenera has not been attempted; satisfactory references on these taxonomic levels must await further study.

Subfamily CHIROMYZINAE

Refs. — Hardy, 1920; Enderlein, 1921; Bezzi, 1922.


Genus BARBIELLINIA Bezzi

BARBIELLINIA Bezzi, 1922:121. Type-species, *hirta* Bezzi (orig. des.).

Chiromyza, Authors, not Wiedemann.

ANNULIPES (Enderlein), 1921:159 (*Chiromyza*). Type-locality: Brazil, Espírito Santo. Distr. — Brazil. N. comb.

26.1



- HIRTA Bezzi, 1922:122. Type-locality: Brazil, São Paulo, Termas de Lindóia, Serra Negra. Distr. — Brazil (Guanabara, São Paulo).
- LINEATA (Enderlein), 1921:158 (*Chiromyza*). Type-locality: Brazil, no locality and São Paulo. Distr. — Brazil (Goiás, Minas Gerais, São Paulo). N. comb.
- MURCICORNIS (Enderlein), 1921:159 (*Chiromyza*). Type-locality: Brazil, São Paulo. Distr. — Brazil. N. comb.
- PARVICORNIS (Enderlein), 1921:159 (*Chiromyza*). Type-locality: Brazil, no locality and São Paulo. Distr. — Brazil. N. comb.

Genus CHIROMYZA Wiedemann

- CHIROMYZA Wiedemann, 1820:19. Type-species, *vittata* Wiedemann (Brauer, 1882:86).
- Xenomorpha* Macquart, 1838:197 (separate, p. 193). Type-species, *leptiformis* Macquart (mon.).
- BREVICORNIS (Lindner), 1949:787 (*Xenomorpha*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. N. comb.
- ENDERLEINI (Lindner), 1949:788 (*Xenomorpha*). Type-locality: "Brazil". Distr. — Brazil. N. comb.
- LEPTIFORMIS (Macquart), 1838:197 (separate, p. 193) (*Xenomorpha*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Venezuela to Brazil (Espírito Santo, Guanabara).
- NIGRA Bezzi, 1922:121. Type-locality: "Ecuador". Distr. — Ecuador.
- OCHRACEA Wiedemann, 1820:20. Type-locality: "Brazil". Distr. — Peru, Bolivia, Brazil (Pará to Santa Catarina).
- fuseana* Wiedemann, 1821:115. Type-locality: "Brazil".
- spp.* Williston, 1888:244. Brazil (Mato Grosso, Chapada, and Guanabara, Rio de Janeiro).
- STYLICORNIS (Enderlein), 1921:165 (*Xenomorpha*). Type-locality: "Brazil". Distr. — Brazil. N. comb.
- TENUICORNIS (Lindner), 1949:737 (*Xenomorpha*). Type-locality: Guyana, Upper Courantyne River, King Fredrick William IV Falls. Distr. — Guyana. N. comb.
- TRISTRIGATA (Enderlein), 1921:161 (*Xenomorpha*). Type-localities: Peru, Callanga, and Bolivia, Songo. Distr. — Peru, Bolivia. N. comb.
- VIRIDIS Bezzi, 1922:120 (as *vittata* ssp.). Type-locality: Brazil, São Paulo. Distr. — Peru, Brazil (Goiás to São Paulo). N. status.
- VITTATA Wiedemann, 1820:20. Type-locality: "Brazil". Distr. — Brazil (Minas Gerais, São Paulo), Paraguay.

Genus HYLOROPS Enderlein

- HYLOROPS Enderlein, 1921:179. Type-species, *philippii* Enderlein (orig. des.).
- PHILIPPI Enderlein, 1921:179. Type-locality: Chile, Concepción. Distr. — Chile.

Genus HYLORUS Philippi

- HYLORUS Philippi, 1865:728. Type-species, *krausei* Philippi (mon.).



AUSTRALIS Aubertin, 1930:99. Type-locality: Argentina, Chubut. Distr. — Argentina.

KRAUSEI Philippi, 1865:728. Type-locality: Chile, ?Casa Pangue. Distr. — Chile.

Genus MESOMYZA Enderlein

MESOMYZA Enderlein, 1921:162. Type-species, *interrupta* Enderlein (orig. des.).

INTERRUPTA Enderlein, 1921:162. Type-locality: "Brazil". Distr. — Brazil.

SERICEA Enderlein, 1921:162. Type-locality: Bolivia, Songo. Distr. — Bolivia.

TENUICORNIS Enderlein, 1921:162. Type-localities: Brazil, São Paulo, Cantareira; Espírito Santo. Distr. — Brazil.

Genus NONACRIS Walker

NONACRIS Walker, 1856:7. Type-species, *transequa* Walker (mon.).

BREVICORNIS Enderlein, 1921:165. Type-locality: "Brazil". Distr. — Brazil.

CHILENSIS Lindner, 1943:91. Type-locality: Chile, Aysén, Calbuco. Distr. — Chile.

LONGICORNIS Enderlein, 1921:164. Type-locality: "Colombia". Distr. — Colombia.

NIGRIVENTRIS Enderlein, 1921:165. Type-localities: Colombia, no locality, and Ecuador, Santa Inez. Distr. — Colombia, Ecuador.

PARTITIFRONS Enderlein, 1921:165. Type-locality: "Brazil". Distr. — Brazil.

SCUTELLARIS Enderlein, 1921:164. Type-locality: Colombia, Bogotá. Distr. — Colombia.

TRANSEQUA Walker, 1856:7. Type-locality: "South America". Distr. — Colombia, Venezuela.

Genus TANA Reed

Lagarus Philippi, 1865:728 (preocc. Chaudoir, 1838). Type-species, *paulseni* Philippi (mon.).

TANA Reed, 1888:284 (nom. nov. for *Lagarus* Philippi). Type-species, *Lagarus paulseni* Philippi (aut.).

Lagarinus Enderlein, 1913a:250, 251. Type-species, *paradoxus* Enderlein (orig. des.).

Ref. — Curran, 1936:281 (Generic status of *Lagarus*).

PARADOXA (Enderlein), 1913a:252 (*Lagarinus*). Type-locality: "Chile". Distr. — Chile.

PAULSENI (Philippi), 1865:729 (*Lagarus*). Type-locality: "Chile". Distr. — Chile. Ref. — Enderlein, 1913 (Quotes original descr.).

Unplaced and extralimital species

despectus Walker, 1856:6 (*Inopus*). Type-locality not stated; given by Bezzi as South America, but Nagatomi and Yukawa say that this is an error and that this species is actually Australian.

Chiromyzinae, ?n. sp., Lindner, 1933b:325. Rio de Janeiro, Itatiaia.

Subfamily BERIDINAE

Ref. — Enderlein, 1921 (rev.).

Genus ACTINA Meigen

ACTINA Meigen, 1804:116. Type-species, *Beris nitens* Latreille (Rondani, 1863:87; 1864:87).

ELEGANS Schiner, 1868:71. Type-locality: "Colombia". Distr. — Colombia.

Genus ACTINOMYIA Lindner

ACTINOMYIA Lindner, 1949:793. Type-species, *novaeaeutoniae* Lindner (mon.). *Actinomyia*, error.

NOVAEAEUTONIAE Lindner, 1949:793, 794. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus ALLOGNOSTA Osten Sacken

ALLOGNOSTA Osten Sacken, 1883:297. Type-species, *Beris fuscitarsis* Say (Coquillett, 1910:505).

LIMBATIVESTRIS Enderlein, 1921:185. Type-locality: Paraguay, San Bernardino. Distr. — Paraguay.

Genus ARCHISTRATIOMYS Enderlein

ARCHISTRATIOMYS Enderlein, 1913b:548. Type-species, *Beris luctifera* Philippi (orig. des.).

BRASILIENSIS Enderlein, 1921:190. Type-locality: Brazil, São Paulo. Distr. — Brazil (Rio de Janeiro to Santa Catarina).

ITATIAYENSIS Lindner, 1928:236. Type-locality: Brazil, Rio de Janeiro, Itatiaia Mts. Distr. — Brazil.

LUCTIFERA (Philippi), 1865:729 (*Beris*). Type-locality: Chile, Valdivia, Corral. Distr. — Chile (Valparaiso, Valdivia). Ref. — Aubertin, 1930 (suppl. descr., fig.).

philippii Bigot, 1879:199 (*Exaireta*). Type-locality: "Chile".

RUFIPALPIS (Wiedemann), 1830:619 (*Xylophagus*). Type-locality: "Mexico". Distr. — Mexico.

Genus BENHAMYIA Miller

Huttonella Enderlein, 1921:195 (procc. Pfeiffer, 1855). Type-species *Exaireta alpina* Hutton (orig. des.).

BENHAMYIA Miller, 1945:72 (nom. nov. for *Huttonella* Enderlein). Type-species, *Exaireta alpina* Hutton (aut.).

ENGELI (Lindner), 1943:97 (*Huttonella*). Type-locality: Chile, Aysén, Calbuco. Distr. — Chile.

NORDENHOLZI (Lindner), 1943:98 (*Huttonella*). Type-locality: Chile, Aysén, Calbuco. Distr. — Chile.



Genus BERIDOPS Enderlein

- BERIDOPS Enderlein, 1913b:550. Type-species, *Beris trichonota* Philippi (orig. des.) = *maculipennis* Blanchard.
- MACULIPENNIS (Blanchard), 1852:400 (*Beris*). Type-locality: "Chile". Distr. — Chile, Argentina. Refs. — Aubertin, 1930 (suppl. descr., fig.), Lindner, 1943 (suppl. descr.).
- trichonota* Philippi, 1865:730 (*Beris*). Type-locality: Chile, Valdivia.

Genus BERIS Latreille *

- BERIS Latreille, 1802:447. Type-species, *Stratiomys sexdentata* Fabricius (mon.) = *chalybeata* (Forster).
- ALBITARSIS Macquart, 1846:175 (separate, p. 47). Type-localities: "Colombia, Venezuela".
- CAERULESCENS Schiner, 1868:71. Type-locality: "Colombia". Distr. — Colombia.
- FLAVISPINOSA Macquart, 1855:38 (separate, p. 58). Type-locality: "South America". Distr. — South America.
- GUERINII Macquart, 1846:176 (separate, p. 48). Type-locality: "South America". Distr. — South America.
- MODESTA Philippi, 1865:730. Type-locality: Chile, Valdivia, Los Ulmos. Distr. — Chile.
- VIRIDIVENTRIS Philippi, 1865:731. Type-locality: Not stated, presumably Chile. Distr. — Chile.

Genus BERISMYIA Giglio-Tos

- BERISMYIA Giglio-Tos, 1891:2, 1892:6. Type-species, *fusca* Giglio-Tos (mon.). Ref. — McFadden, 1971a (synonymy).
- Antissops* Enderlein, 1914a:314. Type-species, *denticulata* Enderlein (orig. des.).
- BARBIELINII (Bezzi), 1908a:472 (*Allognosta*). Type-locality: Brazil, São Paulo. Distr. — Brazil (São Paulo to Santa Catarina). Position uncertain.
- BEZZII (Lindner), 1949:796 (*Antissops*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Lindner, 1969:2 (suppl. descr.).
- FUSCA Giglio-Tos, 1892:7. Type-locality: "Mexico". Distr. — Mexico (Veracruz, Morelos, Chiapas), Costa Rica.
- denticulata* Enderlein, 1914a:315 (*Antissops*). Type-locality: "Costa Rica".
- NIGROFEMORATA Williston, 1900:230. Type-locality: Mexico, Guerrero, Omitemc. Distr. — Mexico (Michoacan, Morelos, Puebla, Guerrero).

Genus DRAYMONIA Aubertin

- Chilota* Aubertin, 1930:102 (preocc. Michaelsen, 1899). Type-species, *nitida* Aubertin (orig. des.).

* Generic position of species referred to this genus uncertain.



DRAYMONIA Aubertin, 1932:284 (nom. nov. for *Chilota* Aubertin). Type-species, *Chilota nitida* Aubertin (aut.).

NITIDA (Aubertin), 1930:102 (*Chilota*). Type-locality: Chile, Casa Pangue. Distr. — Chile, Patagonia.

Genus EXAERETINA Enderlein

EXAERETINA Enderlein, 1921:194. Type-species, *auricoma* Enderlein (orig. des.).

AURICOMA Enderlein. 1921:195. Type-locality: "Chile". Distr. — Chile.

Genus HADRESTIA Thomson

HADRESTIA Thomson, 1869:453. Type-species, *aenea* Thomson (mon.).

AENEA Thomson, 1869:454. Type-locality: "Patagonia". Distr. — Patagonia.

MINOR Brèthes, 1924:104. Type-locality: Chile, Chiloé Island. Distr. — Chile (north to Valdivia); Argentina (Tierra del Fuego). Ref. — Aubertin, 1930 (suppl. descr., fig.).

Genus HEMIMEGACIS Enderlein

HEMIMEGACIS Enderlein, 1921:209. Type-species, *maculifrons* Enderlein (orig. des.).

MACULIFRONS Enderlein, 1921:209. Type-locality: Bolivia, Cillutincara. Distr. — Bolivia.

Genus HETERACANTHIA Macquart

HETERACANTHIA Macquart, 1849:347 (separate, p. 43). Type-species, *ruficornis* Macquart (mon.).

RUFICORNIS Macquart, 1849:348 (separate, p. 44). Type-locality: Venezuela, Caracas. Distr. — Mexico (north to Nayarit and Veracruz) to Brazil (south to Santa Catarina), Trinidad. Ref. — Enderlein, 1921 (suppl. descr.).

mexicana Giglio-Tos, 1891:1, 1892:8. Type-locality: "Mexico".

VIOLACEIVENTRIS Enderlein, 1921:201. Type-locality: "?Brazil". Distr. — ?Brazil.

Genus MACROMERACIS Enderlein

MACROMERACIS Enderlein, 1921:211. Type-species, *Beris thoracica* Philippi (orig. des.).

ELONGATA Aubertin, 1930:104. Type-locality: Chile, Chiloé Island, Castro. Distr. — Chile (Valparaiso to Chiloé Island).

LONGICORNIS (Philippi), 1865:731 (*Beris*). Type-locality: Chile, Valdivia. Distr. — Chile (Valparaiso to Aysén.). Ref. — James, 1939a (suppl. descr.).

LUTEIVENTRIS (Philippi), 1865:731 (*Beris*). Type-locality: not stated, presumably Chile. Distr. — Chile.

THORACICA (Philippi), 1865:730 (*Beris*). Type-locality: Chile, Corral. Distr. — Chile (Santiago to Aysén). Ref. — Lindner, 1943 (suppl. descr.).



Genus MICROHADRESTIA Lindner

- MICROHADRESTIA Lindner, 1943:96. Type-species, *minuta* Lindner (mon.).
MINUTA Lindner, 1943:96. Type-locality: Chile, Chiloé, Olmué. Distr. — Chile (Valdivia to Chiloé Island).

Genus NEOBERIS Lindner

- NEOBERIS Lindner, 1949:797. Type-species, *brasiliana* Lindner (mon.).
BRASILIANA Lindner, 1949:797, 798. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus NEOCHORISOPS Lindner

- NEOCHORISOPS Lindner, 1935:397. Type-species, *brasiliensis* Lindner (mon.).
BRASILIENSIS Lindner, 1935:398. Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Juçara. Distr. — Brazil.
PLAUMANNI Lindner, 1949:792. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus NEOEXAIRETA Osten Sacken

- Exaireta* Sehlnert, 1867:309 (preocc. Hübner, 1820, as *Exaereta*). Type-species, *Xylophagus spiniger* Wiedemann (mon.).
NEOEXAIRETA Osten Sacken, 1878:44 (nom. nov. for *Exaireta* Schiner). Type-species, *Xylophagus spiniger* Wiedemann (aut.).
Neoexaereta, emendation.
Exaereta, emendation.
Exairete, error.
EUPODATA (Bigot), 1879:198 (*Exaireta*). Type-locality: "Chile". Distr. — Chile.
HYACINTHINA (Bigot), 1879:199 (*Exaireta*). Type-locality: "Chile". Distr. — Chile.
LONGICORNIS (Schiner), 1868:73 (*Exaireta*). Type-locality: "Chile". Distr. — Chile.
PHILIPPI (Rondani), 1863:88 (*Beris*). Type-locality: "Chile". Distr. — Chile. Ref. — Lindner, 1943 (suppl. deser.).

Genus OPLACHANTHA Rondani

- OPLACHANTHA Rondani, 1863:87. Type-species, *Beris mexicana* Bellardi (orig. des.).
Oplacantha, emend.
Hoplacantha, emend.
ANNULIPES Enderlein, 1921:207. Type-locality: Venezuela, Caracas. Distr. — Venezuela.
CINCTICORNIS Enderlein, 1921:204. Type-locality: Brazil, Santa Catarina. Distr. — Brazil.
FORMOSA Enderlein, 1921:207. Type-locality: Colombia, Cordilleras. Distr. — Colombia.




- IUBATA Enderlein, 1921:208. Type-locality: Peru, Vilcanota. Distr. — Colombia, Peru, Bolivia.
- LANUGINOSA Enderlein, 1921:205. Type-locality: Argentina, Salta. Distr. — Argentina.
- LATIFRONS Enderlein, 1921:206. Type-locality: Peru, Cuzco, Marcapata. Distr. — Peru.
- LIMBATA Bigot, 1879:196. Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico.
- MEXICANA (Bellardi), 1859:20, 1861:227 (*Beris*). Type-locality: "Mexico". Distr. — Mexico (San Luis Potosí, Veracruz, Chiapas), Guatemala, Costa Rica, Brazil (Guanabara, Rio de Janeiro, São Paulo).
- bellula* Williston, 1888:245 (*Beris*). Type-locality: Brazil, Guanabara, Rio de Janeiro.
- NIGERRIMA Lindner, 1949:795. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PERUANA Enderlein, 1921:208. Type-locality: Peru, Cagon. Dist. — Peru.
- PULCHELLA Williston, 1888:245. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil.
- SIMILIS Enderlein, 1921:207. Type-locality: Chile, Colchagua, Rengo. Distr. — Chile.
- SUBCRASSICALX Enderlein, 1921:205. Type-locality: Peru, Cuzco, Callanga. Distr. — Peru.
- TRICOLOR (Wiedemann), 1828:83 (*Beris*). Type-locality: "Brazil". Distr. — Brazil (São Paulo).
- TRISTIS Enderlein, 1921:204. Type-locality: Brazil, São Paulo. Distr. — Brazil.
- VIRIATA Enderlein, 1921:206. Type-locality: Peru, Cuzco, Callanga. Distr. — Peru.

Subfamily SARGINAE

Genus ACROCHAETA Wiedemann

- ACROCHAETA Wiedemann, 1830:42. Type-species, *fasciata* Wiedemann (mon.).
- Compsosoma* Brauer, 1882:79, 87. Type-species, *chalconota* Brauer (orig. des.).
- ADUSTA Lindner, 1949:813. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- BREVISETA Walker, 1854:69. Type-locality: Brazil, Pará. Distr. — Brazil.
- CHALCONOTA (Brauer), 1882:79 (*Compsosoma*). Type-locality: "Venezuela". Distr. — Venezuela, ?Costa Rica. Ref. — Lindner, 1936 (suppl. descr., ?color variant).
- DEGENERATA Lindner, 1949:812. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- DIMIDIATA Lindner, 1949:811. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- EHEMON (Walker), 1849:515 (*Sargus*). Type-locality: Brazil, Pará. Distr. — Brazil.
- ELEGANS Perty, 1833:184, tab. xxxvi. Type-locality: "Brazil". Distr. — Brazil.



FASCIATA Wiedemann, 1830:42. Type-locality: "Brazil". Distr. — Mexico (north to Veracruz) to Brazil (Santa Catarina). Ref. — Giglio-Tos, 1892:16 (redescr.).

flaveola Bigot, 1879:205 (*Histiodroma*). Type-locality: "Mexico".

ssp. MEXICANA Lindner, 1949:814. Type-locality: "Mexico". Distr. — Mexico.

GORGONA Lindner, 1949:814. Type-locality: Colombia, Gorgona Island. Distr. — Colombia.

HATERIUS (Walker), 1849:516 (*Sargus*). Type-locality: "Venezuela". Distr. — Venezuela.

LONGISETA Walker, 1854:69. Type-locality: Brazil, Pará. Distr. — Brazil.

LONGIVENTRIS Enderlein, 1914b:595. Type-locality: Brazil, Santa Catarina. Distr. — Brazil. Ref. — Lindner, 1935 (suppl. descr.).

PICTA (Brauer), 1882:80 (*Comptosoma*). Type-locality: "Venezuela". Distr. — Venezuela.

PLANA (Walker), 1856:82 (*Chrysochlora*). Type-locality: "Colombia". Distr. — Colombia.

TINCTA (Walker), 1856:82 (*Chrysochlora*). Type-locality: "Colombia". Distr. — Colombia.

Genus CEPHALOCHRYSA Kertész

CEPHALOCHRYSA Kertész, 1912:99. Type-species, *Sargus hovas* Bigot (orig. des.).

Isosargus James, 1935b:273. Type-species, *Chrysonotus nigricornis* Loew (orig. des.).

TEXANA (Melander), 1904:19 (*Sargus*). Type-locality: U.S.A., Texas, Austin. Distr. — U.S.A., Mexico.

Genus EUMENOGASTRINA Enderlein

EUMENOGASTRINA Enderlein, 1914b:589. Type-species, *angusta* Enderlein (orig. des.).

ANGUSTA Enderlein, 1914b:589. Type-localities: Mexico, Chiapas; Costa Rica. Distr. — Mexico, Costa Rica.

ssp. CUPRARIA Enderlein, 1914b:590 (as *angusta* var., status uncertain). Type-locality: Costa Rica, no locality. Distr. — Costa Rica.

Genus HIMANTOLOBA McFadden

HIMANTOLOBA McFadden, 1970:274. Type-species, *Sargus flavopilosus* Bigot (orig. des.).

Chrysonotus Authors (not Loew, 1855, not Swainson, 1837).

Chrysochroma Authors (not Williston, 1896).

Chrysonotomyia Authors (not Hunter, 1900).

ALBIPES (Townsend), in Adams, 1903:31 (*Chrysochroma*). Type-locality: Mexico, Guito. Distr. — Mexico (Veracruz to Jalisco and Chiapas).

ANALIS (Williston), 1888:251 (*Chrysonotus*). Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil.



- DIETROA Sehiner, 1866:63 (*Chrysonotus*). Type-locality: "Colombia". Distr. — Colombia.
- FLAVIVENTRIS (Lindner), 1935:403 (*Chrysochroma*). Type-locality: Brazil, Rio de Janeiro, Petrópolis. Distr. — Brazil.
- FLAVONIGRA (Lindner), 1928:237 (*Sargus*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara to Santa Catarina). Refs. — Carrera & Lane, 1945:128 (deser. of ♀); Lindner, 1969 (suppl. deser.).
- FLAVOPILOSA (Bigot), 1879:227 (*Chrysonotus*). Type-locality: "Mexico". Distr. — Mexico (north to Nayarit and Veraeruz) to Costa Rica.
- aeneiventris* Giglio-Tos, 1892:25 (*Chrysonotus*). Type-locality: Mexico, Orizaba.
- FULVITHORAX Bigot, 1879:228 (*Chrysonotus*). Type-locality: "Amazonia". Distr. — Brazil.
- JAMESI (Lindner), 1969:5 (*Sargus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. N. comb.
- NIGRIFEMORATA (Maequart), 1847:47 (*Sargus*). Type-locality: "Mexico". Distr. — Mexico (Guerrero), Costa Rica.
- pulchra* Williston, 1900:233 (*Chrysochroma*). Type-locality: Mexico, Guerrero, Chilpancingo and Amula.
- SPLENDENS (Schiner), 1868:62 (*Chrysonotus*). Type-locality: "South America". Distr. — South America.
- SUPERBA (Lindner), 1949:803 (*Pedicella*). Type-locality: Guyana, Upper Courantine River, King Frederick William IV Falls. Distr. — Guyana, Peru. Ref. — Lindner, 1951:248 (suppl. deser.). N. comb.
- VENUSTULA (Lindner), 1936:156 (*Chrysochroma*). Type-locality: "Costa Rica". Distr. — Costa Rica.

GENUS MEROSARGUS LOEW


- MEROSARGUS Loew, 1855:144. Type-species, *Sargus obscurus* Wiedemann (orig. des.). Refs. — James, 1941a (key); James & McFadden, 1971 (partial revision).
- Coenosargus* Enderlein, 1914b:587. Type-species, *azureus* Enderlein (orig. des.).
- Aloipha* Enderlein, 1914b:584. Type-species, *cyaneoscutellata* Enderlein (orig. des.).
- Aloiphina* Lindner, 1949:810. Type-species, *stigmatica* Lindner (mon.).
- ABANA Curran, 1932b:4. Type-locality: Guatemala, Pochuta, Santa Emilia. Distr. — Mexico (Veraeruz), Guatemala.
- ACUTUS James, in James & McFadden, 1971:35. Type-locality: Ecuador, Baños. Distr. — Ecuador.
- AENEUS (Lindner), 1949:808 (*Coenosargus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- AKREI James & McFadden, 1971:7. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Mexico (Oaxaca) to Brazil (Amapá).
- ALBIDUS McFadden, in James and McFadden, 1971:8. Type-locality: Mexico, Sinaloa. Distr. — Mexico.



- ALBIFACIES James, 1941b:248. Type-locality: Mexico, D. F., Mexico. Distr. — Mexleo (D. F., Morelos).
- ALBOPICTUS James, *in* James & McFadden, 1971:27. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- ALTICOLA McFadden, *in* James & McFadden, 1971:47. Type-locality: Mexico, Chiapas, San Cristóbal de las Casas. Distr. — Mexico.
- ALTIFRONS James, *in* James & McFadden, 1971:40. Type-locality: Venezuela, Rancho Grande nr. Maracay. Distr. — Venezuela.
- AMETHYSTINUS (Lindner), 1969:7 (*Aloipha*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- AMPULLA James, *in* James & McFadden, 1971:4. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- ANDINUS James, *in* James and McFadden, 1971:28. Type-locality: Peru, Cuzco, Quincemil. Distr. — Ecuador, Peru.
- ANGULATUS McFadden, *in* James & McFadden, 1971:17. Type-locality: Mexico, Hidalgo, Jacala. Distr. — Peru.
- ANGUSTUS James, *in* James & McFadden, 1971:29. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- ANTENNATUS Sehiner, 1868:63. Type-locality: "South America". Distr. — "South America".
- ANTICUS Curran, 1932b:2. Type-locality: Guatemala, Moca Guatalon. Distr. — Mexico (Veracruz) to Venezuela and Argentina (Salta).
- APICALIS Lindner, 1935:404. Type-locality: Venezuela (?), Demerara. Distr. — Venezuela (?).
- ARCUATUS James, *in* James & McFadden, 1971:33. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador, Peru.
- ATRIANNULATUS McFadden, *in* James & McFadden, 1971:42. Type-locality: Mexico, Puebla, Tezuitlan. Distr. — Mexico.
- AUREONITENS James, *in* James & McFadden, 1971:40. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- AURIVENA James, *in* James & McFadden, 1971:35. Type-locality: Panama, Cerro Campana. Distr. — Panama.
- ssp. AUSTRALIS James, *in* James & McFadden, 1971:36. Type-locality: Colombia, Meta, Restrepo. Distr. — Colombia, Venezuela, Peru.
- AZUREUS (Enderlein), 1914b:587 (*Coenosargus*). Type-locality: Brazil, Santa Catarina. Distr. — Canal Zone to Guyana, Brazil (Santa Catarina), and Peru. Ref. — Lindner, 1929 (suppl. deser.).
- BANKSI James, 1936a:53. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Costa Rica, Canal Zone, Panama, Peru.
- BARBATUS James, *in* James & McFadden, 1971:20. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone, Panama.
- BASALIS James, *in* James & McFadden, 1971:3. Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica.
- BEAMERI James, 1941e:107. Type-locality: U.S.A., Arizona, Baboquivari Mts. Distr. — U.S.A. (Arizona) to Mexico (Chiapas).
- BEQUAERTI Curran, 1928a:18. Type-locality: Guatemala, Porto Barrios. Distr. — Guatemala, Honduras.
- BITAENIATUS Lindner, 1949:816. Type-locality: Guyana, Essequibo R., Moraballi Creek. Distr. — Guyana.



- BITUBERCULATUS Schiner, 1868:64. Type-locality: "South America". Distr. — "South America".
- BIVITTATUS James, in James & McFadden, 1971:10. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- BRACHIATUS James, in James & McFadden, 1971:24. Type-locality: Costa Rica, Cartago, La Suiza de Turriabla. Distr. — Costa Rica, Panama.
- BRUNNEUS Lindner, 1933b:328. Type-locality: "Brazil". Distr. — Brazil.
- BRUNNIPES McFadden, in James & McFadden, 1971:13. Type-locality: Mexico, Veracruz, Jalapa. Distr. — Mexico (Morelos, Veracruz, Chiapas).
- BULBIFRONS Williston, 1900:235. Type-locality: Mexico, Guerrero, Xucumanatlan. Distr. — Mexico.
- C-NIGRUM (Lindner), 1951:247 (*Pedicella*). Type-locality: Peru, San Martín, Tingo Maria. Distr. — Ecuador, Peru.
- CALCEOLATUS Bigot, 1879:229. Type-locality: "Mexico". Distr. — Mexico.
- CINGULATUS Schiner, 1868:62. Type-locality: "South America". Distr. — Mexico (north to Veracruz) to Brazil (south to Santa Catarina).
- CITRINELLUS Lindner, 1949:815. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CITRINUS James, in James & McFadden, 1971:47. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- COMPLICATUS James, in James & McFadden, 1971:43. Type-locality: Colombia, Meta, Villavicencio. Distr. — Colombia, Venezuela.
- CONCINNATUS Williston, 1900:235. Type-locality: Mexico, Guerrero, Omilteme. Distr. — Mexico.
- CONOPSOIDES James, in James & McFadden, 1971:45. Type-locality: Ecuador, Pastaza, Pompeya. Distr. — Ecuador.
- CONTORTUS James, in James & McFadden, 1971:4. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- CONVEXIFRONS McFadden, in James & McFadden, 1971:37. Type-locality: Mexico, Puebla, Tezuitlan. Distr. — Mexico (Hidalgo, Puebla), Costa Rica, Ecuador.
- CORIACEUS Giglio-Tos, 1891:4 (1892:21). Type-locality: Mexico, Orizaba. Distr. — Mexico (Morelos, Orizaba), Guatemala, Costa Rica. Ref. — Lindner, 1936:155 (suppl. descr.).
- cercaceus* Williston, 1900:235; ?lapsus.
- COXALIS Lindner, 1949:817. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- bituberculatus* Schiner of Lindner, 1935:405, misident.
- CYANEOSCUTELLATUS (Enderlein), 1914b:585 (*Aloipa*). Type-locality: Mexico, Chiapas. Distr. — Mexico, Guatemala.
- CYRTOMETOPUS James, in James & McFadden, 1971:28. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- DISSIMILIS Giglio-Tos, 1891:3 (1892:20). Type-locality: Mexico, Orizaba. Distr. — Mexico.
- DIVISUS James, in James & McFadden, 1971:11. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.



- DORSALIS Lindner, 1969:7. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- ELATUS Curran, 1932b:5. Type-locality: Canal Zone, France Field. Distr. — Mexico (Tabasco), Costa Rica, Panama, Canal Zone.
- ELONGATUS James, in James & McFadden, 1971:13. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Canal Zone, Ecuador, Peru.
- ETHELIA Curran, 1932b:6. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- EUNOMUS James, 1967a:8. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.
- FESTIVUS Williston, 1888:250. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara to Santa Catarina).
- FLAVICEPS McFadden, in James & McFadden, 1971:38. Type-locality: Mexico, Chiapas, Palenque Ruins. Distr. — Mexico (Veracruz, Chiapas, Oaxaca).
- FLAVISSIMUS James, in James & McFadden, 1971:37. Type-locality: Peru, San Martín, Tingo Maria. Distr. — Peru.
- FLAVITARSIS (Lindner), 1969:8 (*Aloipha*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- FRATERNUS Bigot, 1879:228. Type-locality: "Mexico". Distr. — Mexico.
- FRONTATUS Schiner, 1868:63. Type-locality: "Colombia". Distr. — Colombia.
- FROSTI James, 1941a:307. Type-locality: Ecuador, Baños. Distr. — Ecuador.
- FUMPENNIS James, in James & McFadden, 1971:42. Type-locality: Ecuador, Tandapi. Distr. — Ecuador.
- GEMINATUS James, in James & McFadden, 1971:29. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- COLBACHI James, in James & McFadden, 1971:50. Type-locality: Argentina, Tucumán, Burruyacu Distr. Distr. — ?Peru, Argentina.
- COWDEYI Curran, 1928a:18. Type-locality: Colombia, Rio Frio. Distr. — Guatemala to Brazil (south to Santa Catarina); Trinidad.
- GRACILIOR James, in James & McFadden, 1971:19. Type-locality: Ecuador, Tcma, Libertad. Distr. — Ecuador, Peru.
- GRACILIS Williston, 1888:249. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Venezuela, Ecuador, Peru, Brazil (south to São Paulo).
- GRANULOSUS James, in James & McFadden, 1971:20. Type-locality: Ecuador, Tandapi. Distr. — Ecuador.
- HANSONI James, in James & McFadden, 1971:12. Type-locality: Panama, Cerro Campana. Distr. — Costa Rica, Panama.
- HOFFMANNI Lindner, 1935:405. Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil.
- HYALOPTERUS Giglio-Tos, 1891:3 (1892:19). Type-locality: Mexico, Orizaba. Distr. — Mexico (Veracruz), Ecuador, Peru.
- IRWINI James & McFadden, 1971:6. Type-locality: Mexico, Chiapas, Huixtla. Distr. — Mexico, El Salvador.
- LACRIMOSUS James, 1971:34. Type-locality: Peru, Cuzco, Quincemil. Distr. — Peru.
- LAMPNOTUS James, 1941a:308. Type-locality: Panama, Cabima. Distr. — Costa Rica, Canal Zone, Panama.



- LANIGER James, *in* James & McFadden, 1971:48. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador.
- LATEROMACULATUS James, *in* James & McFadden, 1971:15. Type-locality: Ecuador, Tena, Libertad. Distr. — Ecuador, Peru.
- LINEARIS James, *in* James & McFadden, 1971:20. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- LONGIPES McFadden, *in* James & McFadden, 1971:9. Type-locality: Mexico, Chiapas, Rayon. Distr. — Mexico.
- LURIDUS Loew, 1855:145. Type-locality: "Venezuela". Distr. — Venezuela.
- LUTZI Curran, 1932b:3. Type-locality: Guyana, Tukeit. Distr. — Canal Zone, Guyana.
- LYRICUS James & McFadden, 1971:11. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Mexico (Veracruz, Chiapas, Oaxaca), Canal Zone.
- SSP. ECUADORENSIS James & McFadden, 1971:11. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador.
- SSP. INCANUS James & McFadden, 1971:11. Type-locality: Peru, Cuzco, Quincemil. Distr. — Costa Rica, Colombia, Peru.
- MEGALOPYGE James, *in* James & McFadden, 1971:32. Type-locality: Peru, Cuzco, Quincemil. Distr. — Ecuador, Peru.
- MELANOTHORAX McFadden, *in* James & McFadden, 1971:17. Type-locality: Mexico, Chiapas, San Cristobal. Distr. — Mexico.
- MIRABILIS James, *in* James & McFadden, 1971:22. Type-locality: Guatemala, Guatemala City. Distr. — Guatemala.
- NEBULIFER James, *in* James & McFadden, 1971:49. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Canal Zone, Panama, Ecuador, Peru.
- NIGRIBASIS James, *in* James & McFadden, 1971:41. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Peru.
- NOTATUS Lindner, 1949:807. Type-locality: Colombia, Gorgona Island. Distr. — Colombia.
- OBSCURUS (Wiedemann), 1820:35 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil.
- OBTUSIPENNIS (Lindner), 1949:809 (*Coenosargus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- OPALIGER Lindner, 1931:306. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara to São Paulo).
- ORIZABAE Giglio-Tos, 1891:3 (1892:20). Type-locality: Mexico, Orizaba. Distr. — Mexico.
- PALLIDUS McFadden, *in* James & McFadden, 1971:17. Type-locality: Mexico, Veracruz, Jalapa. Distr. — Mexico.
- PALLIFRONS Curran, 1932b:3. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- PANAMENSIS McFadden, *in* James & McFadden, 1971:9. Type-locality: Panama, Bocas del Toro. Distr. — Costa Rica, Panama, Ecuador, Peru.
- PAR Curran, 1932b:7. Type-locality: Guatemala, Moca, Guatallon. Distr. — Mexico (Veracruz, Chiapas), El Salvador, Canal Zone.



- PENAI James, in James & McFadden, 1971:25. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador.
- PERUVIANUS James, in James & McFadden, 1971:51. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Ecuador, Peru.
- PETIOLATUS (Macquart), 1838:202 (separate, p. 206) (*Sargus*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- PICTIPES James, in James & McFadden, 1971:30. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Ecuador, Peru.
- PICTITHORAX Curran, 1933:1. Type-locality: Guatemala, Moca, Guatemala. Distr. — Guatemala.
- PRODUCTUS James, in James & McFadden, 1971:33. Type-locality: Venezuela, Rancho Grande near Maracay. Distr. — Venezuela.
- PSEUDOLYRICUS James, in James & McFadden, 1971:6. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador.
- PULCHER James, in James & McFadden, 1971:7. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- QUADRATUS James, in James & McFadden, 1971:44. Type-locality: Ecuador, Zamora, Jumboe River. Distr. — Ecuador.
- QUADRIFASCIATUS Lindner, 1941:178. Type-locality: Peru, Sivia. Distr. — Colombia, Ecuador, Peru.
- ROBUSTUS McFadden, in James & McFadden, 1971:39. Type-locality: Mexico, Oaxaca, "RinAntonio". Distr. — Mexico.
- ROSSI James, in James & McFadden, 1971:5. Type-locality: Peru, San Martín, Tingo Maria. Distr. — Peru.
- ROTUNDATUS Curran, 1932b:4. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Costa Rica, Canal Zone.
- SCHILDI James, in James & McFadden, 1971:10. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica.
- SCROBICULUS McFadden, in James & McFadden, 1971:46. Type-locality: Mexico, Veracruz, Jalapa. Distr. — Mexico (Veracruz, San Luis Potosí).
- SEXMACULATUS McFadden, in James & McFadden, 1971:16. Type-locality: Mexico, Veracruz, Jalapa. Distr. — Mexico.
- SEXNOTATUS James, 1941a:306. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil.
- SMARAGDIFERUS (Bigot), 1879:226 (*Macrosargus*). Type-locality: "Mexico". Distr. — Mexico.
- SPATULATUS Williston, 1900:235. Type-locality: Mexico, Guerrero, Omilteme. Distr. — Mexico.
- STAMINEUS (Fabricius), 1805:253 (*Scaeva*). Type-locality: "South America". Distr. — Mexico (north to Nayarit and San Luis Potosí) to Peru and Brazil (Minas Gerais).
- STIGMATICUS (Lindner), 1949:810 (*Aloiphina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil. Ref. — Lindner, 1969 (suppl. descr.).
- SUBINTERRUPTUS (Bellardi), 1859:44, 1861:244 (*Sargus*). Type-locality: "Mexico". Distr. — Mexico (Sinaloa).
- SUBOBSCURUS Lindner, 1969:6. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



- SUPERNITENS James, in James & McFadden, 1971:31. Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica.
- TAENIATUS (Wiedemann), 1830:37 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil (Pará, São Paulo).
- TANGENS James, in James & McFadden, 1971:31. Type-locality: Ecuador, Napo, Coca. Distr. — Ecuador, Peru.
- TELFORDI James, in James & McFadden, 1971:48. Type-locality: Puerto Rico, El Verde, Caribbean National Forest. Distr. — Puerto Rico.
- TENERICOSUS Lindner, 1929:259. Type-locality: Brazil, Pará, Taperinha near Santarém. Distr. — Brazil.
- TENUICORNIS James, in James & McFadden, 1971:39. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- TERMINALIS James, in James & McFadden, 1971:36. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone, Ecuador.
- TRANSVERSUS McFadden, in James & McFadden, 1971:15. Type-locality: Mexico, Veraacruz, Catemaco. Distr. — Mexico, Peru.
- TRIANGULATUS McFadden, in James & McFadden, 1971:19. Type-locality: Mexico, Chiapas, Yerba Buena. Distr. — Mexico.
- TRIPARTITUS James, in James & McFadden, 1971:52. Type-locality: Peru, Cuzco, Quincemil. Distr. — Ecuador, Peru.
- tripartitis*, error.
- TRISTIS Loew, 1855:145. Type-locality: "Venezuela". Distr. — Venezuela.
- TRITAENIATUS Lindner, 1929:259. Type-locality: Brazil, Pará, Taperinha near Santarém. Distr. — Ecuador, Peru, Brazil (Amazon Basin).
- VARICORNIS James, in James & McFadden, 1971:51. Type-locality: Costa Rica, Cartago, La Suiza de Turrialba. Distr. — Costa Rica.
- VARIERUS James, in James & McFadden, 1971:23. Type-locality: Ecuador, Pastaza, Pompeya. Distr. — Ecuador.
- VENTRALIS McFadden, in James & McFadden, 1971:46. Type-locality: Mexico, Chiapas, San Cristóbal. Distr. — Mexico.
- VERTEBRATUS James, in James & McFadden, 1971:14. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Panama, Peru.
- VIRIDIS McFadden, in James & McFadden, 1971:8. Type-locality: Mexico, Veraacruz, Lake Catemaco. Distr. — Mexico, Costa Rica.
- ZETEKI James, in James & McFadden, 1971:44. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.

Genus MICROCHRYSA Loew

- MICROCHRYSA Loew, 1855:146. Type-species, *Musca polita* Linnaeus (orig. des.).
- BICOLOR (Wiedemann), 1830:41 (*Sargus*). Type-locality: "Puerto Rico". Distr. — Mexico (north to Veracruz and Nayarit) to Brazil (south to Santa Catarina), Paraguay, and Argentina (Misiones, Salta); Cuba, Haiti, Dominican Republic, Puerto Rico, Jamaica, Trinidad, Tobago. Ref. — James, 1967a (suppl. descr.).
- abdominalis* James, 1936a:51. Type-locality: Canal Zone, Ancon.
- bruesi* Johnson, 1919:426. Type-locality: Jamaica, Kingston, Liguanea Plain.



- ssp. *MACULATA* Lindner, 1949:819 (as var.). Type-localities: Argentina, Misiones, Alto Paraná, Bemberg, and Brazil, Santa Catarina, Nova Teutônia. Distr. — Argentina, Brazil.
- DISPAR* Schiner, 1868:61. Type-locality: "South America". Distr. — South America.
- LATIFRONS* (Williston), 1900:234 (*Chrysochroma*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. — Mexico.
- MACULA* (Fabricius), 1805:258 (*Sargus*). Type-locality: "South America". Distr. — South America.
- NOVA* Giglio-Tos, 1891:2, 1892:102. Type-locality: "Mexico". Distr. — Mexico.

Genus *PTECTICUS* Loew

- PTECTICUS* Loew, 1855:142. Type-species, *Sargus testaceus* Fabricius (orig. des.). Ref. — James, 1941a (key).
- Plecticus*, error.
- ADUSTUS* Lindner, 1949:804. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- AFFINIS* Schiner, 1868:65. Type-locality: "South America". Distr. — Brazil (Mato Grosso).
- CILIATUS* James, 1941a:302. Type-locality: Peru, Rio Putumayo, Puerto Alfonso. Distr. — Peru.
- CONCINNUS* Williston, 1900:238. Type-localities: Mexico, Guerrero, Amuia, Omilteme, and Xucumanatlan. Distr. — Mexico.
- CONOPSODES* Schiner, 1868:66. Type-locality: "Colombia". Distr. — Colombia.
- CYANIFRONS* (Rondani), 1848:43, 75 (*Sargus*). Type-locality: "Brazil". Distr. — Mexico (north to Nuevo Leon and Tamaulipas) to Brazil (south to Santa Catarina); Cuba, Puerto Rico, Trinidad.
- nigrifrons* Enderlein, 1914b:584. Type-locality: Mexico, Chiapas (as *testaceus* var.).
- FICULINUS* Lindner, 1940:193. Type-locality: Costa Rica, Hamburg Farm. Distr. — Costa Rica.
- FIGLINUS* Osten Sacken, 1886:24. Type-locality: Guatemala, San Gerónimo. Distr. — Mexico (Guerrero, Jalisco, Chiapas); Guatemala.
- FLAVICEPS* Bigot, 1879:230 (as *Plecticus, lapsus*). Type-locality: "Mexico". Distr. — Mexico.
- GIGLIOTOSI* McFadden, 1971b:412 (nom. nov. for *trivittatus* Say of Giglio-Tos). Type-locality: "Mexico". Distr. — Mexico.
- trivittatus* Say of Giglio-Tos, 1891:3, 1892:24, misident.
- ILLUSTRIS* Schiner, 1868:64. Type-locality: "South America". Distr. — South America.
- INOPS* Lindner, 1949:803. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INVERSUS* Curran, 1934:321. Type-locality: Guyana, Kamakusa. Distr. — Guyana, Peru, Brazil (south to São Paulo).
- LANEI* James, 1941a:301. Type-locality: Brazil, São Paulo, Batéia. Distr. — Brazil (São Paulo, Paraná).



- MACULATUS Williston, 1900:238. Type-locality: Mexico, N. Yucatan. Distr. — Mexico.
- NERULIFER James, 1941a:303. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- NIGROPYGHIALIS Lindner, 1931:305. Type-locality: Bolivia, Santa Cruz de la Sierra. Distr. — Bolivia, Guyana, Brazil (Amazonas).
- NITIDIPENNIS Loew, 1855:143. Type-locality: "Venezuela". Distr. — Venezuela.
- NUBILIPENNIS James, 1938:202. Type-locality: Colombia, Boyaca, Muzo. Distr. — Panama to Peru.
- POMACEUS Loew, 1855:144. Type-locality: "Chile". Distr. — Chile.
- SACKENII Williston, 1885:124. Type-locality: U.S.A., Florida. Distr. — U.S.A., widespread; Mexico (south to Chiapas). Ref. — McFadden, 1967 (larva).
- TENUIS Schiner, 1868:66. Type-locality: "Colombia". Distr. — Colombia.
- TERMINALIS James, 1938:202. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia.
- TESTACEUS (Fabricius), 1805:257 (*Sargus*). Type-locality: "South America". Distr. — Mexico (north to Tamaulipas) to Brazil (south to Santa Catarina) and Argentina (south to Córdoba); Trinidad. Ref. — Lindner, 1926 (larva).

Genus SARGUS Fabricius *

- SARGUS Fabricius, 1798:549, 566. Type-species, *Musca cupraria* Linnaeus (Latreille, 1810:442). Alleged preoccupation by *Sargus* Klein, a pre-Linnaean name.
- Geosargus* Bezzi, 1907:53 (unjustified new name for *Sargus* Fabricius). Type-species, *Musca cupraria* Linnaeus (aut.).
- Sargus*, subg. PEDICELLINA James, 1952:225. Type-species, *Sargus notatus* Wiedemann (orig. des.) = *fasciatus* Fabricius.
- ALCHIDAS Walker, 1849:517. Type-locality: "Jamaica". Distr. — Jamaica. Ref. — Johnson, 1894 (suppl. descr.).
- ATROBASIS (James), 1941b:247 (*Geosargus*). Type-locality: Mexico, Guerrero, Iguala. Distr. — Mexico.
- BAGOSUS Walker, 1849:518. Type-locality: "Jamaica". Distr. — Jamaica.
- BRASILIENSIS Wiedemann, 1830:37. Type-locality: "Brazil". Distr. — Canal Zone, Brazil (Rio de Janeiro, Guanabara). Ref. — Lindner, 1935 (suppl. descr.).
- CAESIUS Bellardi, 1859:40, 1861:240. Type-localities: Mexico, Guerrero, Omilteme, Xucumanatlan, Sierra de las Aguas Escondidas, and Anganguco. Distr. — Mexico (Nuevo Leon to Durango and Guerrero).
- filiformis* Giglio-Tos, 1891:4 (1892:28). Type-locality: "Mexico".
- CHRYSOCOMUS (James), 1941b:247 (*Geosargus*). Type-locality: Mexico, D. F., Mexico City. Distr. — Mexico (D. F., Michoacán).

* Undoubtedly, some species listed here belong to other genera of Sarginae. At least two subgenera can be recognized. Correct generic and subgeneric assignment can be made only after further studies have been completed.

- CIRCUMCINCTUS (James), 1941a:303 (*Geosargus*). Type-locality: Brazil, Rio de Janeiro, Tinguá. Distr. — Brazil (Rio de Janeiro to São Paulo).
- CLAVATUS Walker, 1854:93. Type-locality: Brazil, Pará. Distr. — Mexico to Brazil (Pará).
- var., Bellardi, 1859:41, 1861:241. Mexico, Michoacán, Morelia.
- CLAVIVENTRIS Rondani, 1850:190. Type-locality: "Venezuela". Distr. — Venezuela.
- CONCINNUS Osten Sacken, 1886:23. Type-locality: Guatemala, San Gerónimo. Distr. — Guatemala.
- CONTRACTUS Walker, 1854:319. Type-locality: Brazil, Pará, Santarém. Distr. — Brazil.
- FASCIATUS Fabricius, 1805:259. Type-locality: "South America". Distr. — U.S.A. (north to New York and Michigan) to Brazil (south to Santa Catarina) and Paraguay; West Indies (Cuba, Santo Domingo, Haiti, Puerto Rico, Jamaica, Granada, Dominica, Trinidad). Ref. — James, 1967a (suppl. descr., as *notatus*).
- notatus* Wiedemann, 1830:34. Type-locality: "Brazil".
- coarctatus* Macquart, 1838:207 (separate, p. 203). Type-locality: "Brazil" (Macquart says Brazil or Chile).
- sallei* Bellardi, 1859:43, 1861:243. Type-locality: "Mexico".
- lucens* Loew, 1866:7 (Cent. 7, no. 1). Type-locality: "Cuba".
- tricolor* Loew, 1866:8 (Cent. 7, no. 12). Type-locality: "Cuba".
- lunus* Bigot, 1879:187 (?*Macrosargus*), nomen nudum or lapsus for *lucens*.
- schwarzi* Curran, 1928a:16. Type-locality: Puerto Rico, Cayey.
- FLAVILATUS* James, nom. nov. for *lateralis* James.
- lateralis* James, 1936a:52 (*Isosargus*; preocc. Macquart, 1834). Type-locality: Canal Zone, Barro Colorado Island.
- ILLUMINATUS (Lindner), 1949:815 (*Geosargus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- JUCUNDUS Walker, 1850:xvii. Type-locality: Brazil, Pará. Distr. — Brazil.
- LATERALIS Macquart, 1834:262. Type-locality: "Cuba". Distr. — Cuba.
- LATUS Bellardi, 1859:41, 1861:241. Type-locality: "Mexico". Distr. — Mexico.
- LINEARIS Walker, 1854:318. Type-locality: "Mexico". Distr. — Mexico.
- LUCIDUS (Lindner), 1949:805 (*Geosargus*). Type-localities: British Honduras, Rio Grande, and Guyana, Mazaruni. Distr. — British Honduras, Guyana.
- MACQUARTH Perty, 1833:185, pl. 37, fig. 2. Type-locality: Brazil, Bahia. Distr. — Brazil.
- MELLEUS Rondani, 1848:104. Type-locality: "Brazil". Distr. — Brazil.
- MULTICOLOR (Lindner), 1949:806 (*Geosargus*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- OPULENTUS Walker, 1854:319. Type-locality: Brazil, Pará. Distr. — Brazil.
- PLEURITICUS Loew, 1866:8 (Cent. 7, no. 13). Type-locality: "Cuba".
- RUFIBASIS (Bigot), 1879:225 (*Macrosargus*). Type-locality: "South America". Distr. — South America.

* *Flavilatus* = yellow side.

- SAPHIREUS Bigot, 1887:28. Type-locality: "Cuba". Distr. — Cuba.
sapphireus, emendation or error.
- SPECIOSUS Macquart, 1846:184 (separate, p. 56). Type-locality: Mexico,
 Yucatan, Merida. Distr. — Mexico (north to Sinaloa and Veracruz)
 to Costa Rica. Ref. — Lindner, 1936 (suppl. descr.).
aureus Bellardi, 1859:42, 1861:242. Type-locality: "Mexico".
splendens Bigot, 1879:224. Type-locality: "Mexico".
- TENUIVENTRIS (Bigot), 1879:225 (*Macrosargus*). Type-locality: "Brazil,
 Amazonia". Distr. — Brazil.
- THORACICUS Macquart, 1834:261. Type-locality: "South America". Distr. —
 Mexico (north to Veracruz) to Brazil (south to Santa Catarina).
- VERSICOLOR Bellardi, 1862:13, 1864:212. Type-locality: Mexico, Oaxaca. Distr.
 — Mexico.
- VIOLACEUS Jaenicke, 1867:326. Type-locality: "Brazil". Distr. — Brazil.

Subfamily RHAPHIOCERINAE

Refs. — James, 1943a, 1966b; McFadden, 1970b.

Genus ANALCOCEROIDES Hollis

- ANALCOCEROIDES Hollis, 1962:558. Type-species, *Campeprosopa longicornis*
 Williston (orig. des.).
- LONGICORNIS (Williston), 1900:239 (*Campeprosopa*). Type-locality: Mexico,
 Guerrero, Omilteme. Distr. — Mexico.

Genus ANALCOCERUS Loew

- ANALCOCERUS Loew, 1855:140. Type-species, *atriceps* Loew (orig. des.).
- ATRICEPS Loew, 1855:140. Type-locality: "Brazil". Distr. — Brazil (Minas
 Gerais to Santa Catarina).
- ORBITALIS James, 1943a:368. Type-locality: Brazil, Santa Catarina, Nova
 Teutônia. Distr. — Brazil.
- TAURUS James, 1943a:370. Type-locality: Brazil, São Paulo, S. Sebastião.
 Distr. — Brazil (São Paulo).

Genus ANISOSCAPUS McFadden

- ANISOSCAPUS McFadden, 1970b:317. Type-species, *chlorovittatus* McFadden
 (orig. des.).
- CHLOROVITTATUS McFadden, 1970b:317. Type-locality: Mexico, Oaxaca, Matias
 Romero. Distr. — Mexico.
- OGHRACEUS McFadden, 1970:318. Type-locality: Mexico, Nayarit, Compostela.
 Distr. — Mexico.

Genus AULOCEROMYIA Lindner

- AULOCEROMYIA Lindner, 1969:8, 9. Type-species, *vespiformis* Lindner (mon.).
- VESPIFORMIS Lindner, 1969:10. Type-locality: Brazil, Santa Catarina, Nova
 Teutônia. Distr. — Brazil.



Genus BASENTIDEMA Macquart

- BASENTIDEMA Macquart, 1838:201 (separate, p. 197). Type-species, *syrphoides* Macquart (mon.).
- COERULESCENS James, 1943a:376. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- HORTULANA (Wiedemann), 1830:32 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil (Rio de Janeiro, Guanabara, São Paulo).
- SYRPHOIDES Macquart, 1838:201 (separate, p. 197). Type-locality: "Brazil". Distr. — Brazil.

Genus BRACHYTHRIX McFadden

- BRACHYTHRIX McFadden, 1970b:318. Type-species, *dileucostigma* McFadden (orig. des.).
- DILEUCOSTIGMA McFadden, 1970b:320. Type-locality: Mexico, Nuevo Leon, Cañon del Diente. Distr. — Mexico.

Genus CYCLOPHLEPS James

- CYCLOPHLEPS James, 1943a:375. Type-species, *tenebrifera* James (orig. des.).
- TENEBRIFERA James, 1943a:375. Type-locality: Peru, Iquitos. Distr. — Peru.

Genus DICRANOPHORA Macquart

- DICRANOPHORA Macquart, 1834:255. Type-species, *Sargus fureifer* Wiedemann (Macquart, 1838:99).
- AFFINIS Williston, 1888:251. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- ASTUTA Williston, 1888:250. Type-locality: Not stated [Brazil]. Distr. — presumably Brazil.
- BREVIFURCA James, 1943a:371. Type-locality: Paraguay, Villarica. Distr. — Paraguay.
- FURCIFERA (Wiedemann), 1824:40 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil (Guanabara).
- PICTA Macquart, 1834:255. Type-locality: French Guiana, Mana R. Distr. — Brazil.

Genus DOLICHTHRIX McFadden

- DOLICHTHRIX McFadden, 1970b:318. Type-species, *melanothorax* McFadden (orig. des.).
- MELANOTHORAX McFadden, 1970b:318. Type-locality: Mexico, Oaxaca, Tehuantepec. Distr. — Mexico.

Genus HEPTOZUS Lindner

- HEPTOZUS Lindner, 1949:799. Type-species, *ecuadorianus* Lindner (mon.).
- ECUADORIANUS Lindner, 1949:800. Type-locality: N. Ecuador. Distr. — Ecuador.



HANSONI James, 1966b:679. Type-locality: Panama, Cerro Campana. Distr.
— Canal Zone, Panama, Ecuador.

Genus HISTIODROMA Schiner

- HISTIODROMA Schiner, 1868:68. Type-species, *Sargus inermis* Wiedemann (orig. des.).
FASCIPENNIS James, 1943a:373. Type-locality: Brazil, São Paulo, Cotia. Distr.
— Brazil (São Paulo, Santa Catarina).
INERMIS (Wiedemann), 1830:31 (*Sargus*). Type-locality: "Brazil". Distr.
— Brazil (Guanabara).
TRICOLOR James, 1940b:121. Type-locality: Peru, Chanchamayo, La Merced,
Rio Toro. Distr. — Peru.

Genus HOPLISTES Macquart

- HOPLISTES Macquart, 1834:253. Type-species, *Sargus bispinosus* Wiedemann (orig. des.).
BISPINOSUS (Wiedemann), 1830:32 (*Sargus*). Type-locality: "Brazil". Distr.
— Guyana, Brazil.
PLEURALIS James, 1949:1. Type-locality: Brazil, Rio Grande do Norte, Natal.
Distr. — Brazil.

Genus LYSOZUS Enderlein

- LYSOZUS Enderlein, 1914b:579. Type-species, *columbianus* Enderlein (orig. des.).
COLUMBIANUS Enderlein, 1914:580. Type-locality: Colombia, Hacienda Pehlke. Distr. — Colombia.

Genus NEANALCOCERUS James

- NEANALCOCERUS James, 1943a:370. Type-species, *Analcocerus hortulanus* Williston (orig. des.).
HORTULANUS (Williston), 1900:238 (*Analcocerus*). Type-locality: Mexico, N. Yucatan. Distr. — Mexico (Guerrero, Yucatan).

Genus PEZODONTINA Lindner

- PEZODONTINA Lindner, 1949:800. Type-species, *guianae* Lindner (mon.).
GUIANAE Lindner, 1949:801. Type-localities: Guyana, Upper Courantyne R., Essequibo R., and Cuyuni R., high forest. Distr. — Guyana.

Genus PHANEROZUS James

- PHANEROZUS James, 1966b:676. Type-species, *rettenmeyeri* James (orig. des.) = *pampinus* (Osten Sacken).
PAMPINUS (Osten Sacken), 1886:25 (*Rhaphiocera*). Type-locality: Panama, Volcan de Chiriqui. Distr. — Costa Rica, Canal Zone, Panama.
rettenmeyeri James, 1966b:677. Type-locality: Canal Zone, Barro Colorado Island.



Genus RHAPHIOCERA Macquart

- RHAPHIOCERA Macquart, 1834:253. Type-species, *Sargus armatus* Wiedemann (mon.). Ref. — Lindner, 1931.
- ARMATA (Wiedemann), 1830:29 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil (Mato Grosso to Espírito Santo and Rio Grande do Sul).
ssp. BORGMEIERI Lindner, 1931:310 (as var.). Type-locality: Brazil, Santa Catarina, Hansa (= Corupá). Distr. — Brazil.
ssp. HANSAE Lindner, 1931:310 (as var.). Type-locality: Brazil, Santa Catarina, Hansa (= Corupá). Distr. — Brazil.
ssp. MINAENSIS Lindner, 1931:310 (as var.). Type-locality: Brazil, Minas Gerais. Distr. — Brazil.
ssp. STEGLMAYRI Lindner, 1931:310 (as var.). Type-locality: Brazil, Rio Grande do Sul. Distr. — Brazil.
ssp. TRANSHANSAE Lindner, 1931:310 (as var.). Type-locality: Brazil, Rio Grande do Sul. Distr. — Brazil.
ssp. ZIKANI Lindner, 1935:399 (as var.). Type-locality: Brazil, Rio de Janeiro, Itatiaia.
- CALOPTERA Osten Sacken, 1886:26. Type-locality: Mexico, Veracruz, Paso del Macho. Distr. — Mexico (Veracruz, Chiapas).
- CRUCIANA Lindner, 1951:246. Type-locality: Peru, La Merced, Rio Chanchamayo. Distr. — Peru.
- FALLENI (Perty), 1830:184 (*Sargus*). Type-locality: "Brazil". Listed by Kertész as questionably a synonym of *armata*.
- FASCIPENNIS Lindner, 1949:802. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- HOPLISTES (Wiedemann), 1830:30 (*Sargus*). Type-locality: Not stated. Distr. — Brazil.
pomaceus Wiedemann of Macquart, 1834:254 (*Hoplistes*). No locality, misident.
- LIMBATA Lindner, 1949:801. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- MIPARTITA Macquart, 1846:182 (separate, p. 54). Type-locality: "Colombia". Distr. — Colombia.
bipartita Walker, 1854:79, error.
- ORNATA Macquart, 1846:183 (separate, p. 55). Type-locality: "Colombia". Distr. — Colombia, Ecuador, Bolivia.
- PICTA Wulp, 1879:22. Type-locality: "Brazil". Distr. — Brazil.
- SANCTIPauli Lindner, 1931:310. Type-locality: Brazil, São Paulo. Distr. — Brazil.

Subfamily CLITELLARIINAE

Tribe ANTISSINI

Genus ANTISSA Walker

- ANTISSA Walker, 1854:63. Type-species, *Clitellaria cuprea* Walker (orig. des.). Ref. — Kertész, 1908, as *Cyanauges* (suppl. descr.).
Exodontha Rondani, 1856:169. Type-species, *pedemontana* Rondani (orig. des.) = *dubia* (Zetterstedt). N. SYN.



- Cyanauges Rondani*, 1863:88. Nomen nudum.
- Cyanauges Philippi*, 1865:732. Type-species, *valdivianus* Philippi = *Oplacantha valdiviana* Rondani (mon.).
- Parantissa Enderlein*, 1914c:10. Type-species, *Cyanauges ruficornis* Schiner (orig. des.).
- RUFICORNIS (Schiner), 1868:54 (*Cyanauges*). Type-locality: "Chile". Distr. — Chile.
- VALDIVIANA (Rondani), 1863:88 (*Oplacantha*). Type-locality: Chile, Valdivia. Distr. — Chile. Ref. — Lindner, 1943 (suppl. descr.).
- valdiviana* Philippi, 1865:732 (*Cyanauges*). Type-localities: Chile, Valdivia and Santiago.
- VILLOSA (Lindner), 1969:3 (*Exodontha*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Tribe CYPHOMYIINI

Ref. — James, 1940a (Key Neotropical genera, except *Abasanistus*).

Genus ABASANISTUS Kertész

- ABASANISTUS Kertész, 1923:90. Type-species, *rubicornis* Kertész (orig. des.). Ref. — Kertész, 1923.
- PAULSENI (Philippi), 1865:732 (*Cyelogaster*). Type-locality: Chile, Santiago, Andes. Distr. — Chile.
- RUBRICEPS (Philippi), 1865:732 (*Cyelogaster*). Type-locality: Chile, Santiago. Distr. — Chile. Ref. — Kertész, 1923:93 (descr., fig.).
- RUBRICORNIS Kertész, 1923:92. Type-locality: Chile, Rancagua. Distr. — Chile.

Genus CHORDONOTA Gerstaecker

- CHORDONOTA Gerstaecker, 1857:311. Type-species, *Cyphomyia inermis* Wiedemann (mon.). Refs. — James, 1940 (review); Iide, 1963a (larva).
- ATERRIMA James, 1940a:123. Type-locality: Paraguay, Villarica. Distr. — Brazil (Mato Grosso, Paraná, Santa Catarina), Paraguay. Ref. — Carrera & Lanc, 1945 (possible synonymy).
- FUSCIPENNIS Bellardi, 1862:11, 1864:210. Type-locality: Mexico, Veracruz, Playa Vicente. Distr. — Mexico (Veracruz, San Luis Potosí).
- GENICULATA Lindner, 1936:157. Type-locality: Costa Rica. Distr. — Honduras, Costa Rica.
- INERMIS (Wiedemann), 1830:55 (*Cyphomyia*). Type-locality: "Brazil". Distr. — Brazil (Mato Grosso to Espírito Santo and Santa Catarina). Ref. — Iide, 1963a (extensive descr., figs., adult and larva).
- LEIOPHTHALMA Williston, 1896:302. Type-locality: "St. Vincent". Distr. — St. Vincent, Granada, Colombia.
- NIGRA Williston, 1888:255. Type-locality: Brazil, Mato Grosso, Chapada Distr. — Peru, Brazil (Mato Grosso).



Genus CYPHOMYIA Wiedemann

- CYPHOMYIA Wiedemann, 1819:54. Type-species, *auriflamma* Wiedemann (Brauer, 1882:87; 1882:31). Ref. — James, 1940a (rev. Neotropical species), James, 1953b (rev. Argentine species), James, 1957a (larva).
- Rondania* Jaenicke, 1867:324 (1868:16) (preocc. Robineau-Desvoidy, 1850). Type-species, *Clitellaria chalybea* Wiedemann (orig. des.).
- Neorondania* Osten Sacken, 1878:50 (nom. nov. for *Rondania* Jaenicke). Type-species, *Clitellaria chalybea* Wiedemann (aut.).
- Gyneuryptaria* Enderlein, 1914b:604. Type-species, *Cyphomyia pilosissima* Gerstaecker (orig. des.).
- ABANA Curran, 1929:5. Type-locality: Brazil, Amazon. Distr. — Brazil.
- ACUMINATA James, 1940a:132. Type-locality: Jamaica, Castleton, Botanical Gardens. Distr. — Jamaica.
- ACEZELI James, 1953b:314. Type-locality: Argentina, Tucumán, Parque Aconquija. Distr. — Argentina (Tucumán, Salta).
- AFFINIS Gerstaecker, 1857:289. Type-locality: Venezuela. Distr. — Venezuela.
- ALBICAPUT (Walker), 1849:521 (*Clitellaria*). Type-localities: Brazil, Pará, and no data. Distr. — Brazil. Position uncertain. Gerstaecker says synonym of *leucocephala*.
- ALEITARSIS (Fabricius), 1805:80 (*Stratiomys*). Type-locality: "South America". Distr. — Mexico (north to Nuevo Leon and Tamaulipas) to Peru and Brazil (Pará); Trinidad.
- fenestrata* Macquart, 1846:176 (separate, p. 48). Type-locality: Mexico, Yucatan, Merida.
- cyanispinis* Bigot, 1875:487. Type-locality: Brazil, "Amazonia".
- ALBOMACULATA (Macquart), 1834:193 (separate, p. 189) (*Odontomyia*). Type-locality: Haiti, Port au Prince. Distr. — Santo Domingo, Haiti, Jamaica. Ref. — Lindner, 1949 (suppl. descr.).
- ALTIFRONS James, 1939a:603. Type-locality: Colombia, Bogotá. Distr. — Colombia, Brazil (Amazonas), Peru, ?Bolivia.
- ANDROGYNA Osten Sacken, 1886:34. Type-locality: Panama, Volcán de Chiriquí, Bugaba. Distr. — Panama, Colombia, Ecuador.
- chiriquina* Lindner, 1935:409. Type-locality: Panama, Chiriquí.
- AURIFLAMMA Wiedemann, 1819:54. Type-locality: "Brazil". Distr. — Mexico (Veracruz) to Uruguay, Paraguay and Argentina (Misiones). Ref. — Iide, 1963b (suppl. descr., figs.).
- cyanea* Fabricius of Macquart, 1834:242. Cayenne (misident).
- coerulea* Macquart, 1834. Expl. Planch. 4, tab. VI, fig. 3.
- AURIFRONS Wiedemann, 1830:56. Type-locality: Not stated. Distr. — Ecuador, Brazil (Goiás to Santa Catarina); Argentina (Misiones).
- BANKSI James, 1937:149. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Mexico (Veracruz), Canal Zone, Colombia.
- BICARINATA Williston, 1900:244. Type-localities: Mexico, Guerrero, Xucumanatlan, and Morelos, Cuernavaca. Distr. — Mexico (Veracruz, Morelos, Michoacan, Guerrero). Ref. — McFadden & James, 1969 (redefinition, speciation).



- BREVIS James, 1940a:139. Type-locality: Cuba, Soledad, near Cienfuegos. Distr. — Cuba.
- CHALYBEA (Wiedemann), 1821:30 (*Clitellaria*). Type-locality: Virgin Islands, St. Thomas. Distr. — Cuba, Dominican Republic, Puerto Rico, Virgin Islands, Antigua, St. Kitts, Jamaica, ?Mexico.
- anchialus* Walker, 1849:522 (*Clitellaria*). Type-locality: "Jamaica".
- CHRYSODOTA Perty, 1833:184. Type-locality: "Brazil". Distr. — Costa Rica to Ecuador, Guyana, and Brazil.
- CLARIPENNIS Macquart, 1847:45 (separate, p. 29). Type-locality: "Brazil". Distr. — Brazil.
- COPRATES (Walker), 1849:523 (*Clitellaria*). Type-locality: Not given. Distr. — Unknown, probably South America.
- COSTALIS Walker, 1837:341. Type-locality: Brazil, São Paulo. Distr. — Brazil.
- CYANEA (Fabricius), 1794:265 (*Stratiomys*). Type-locality: "French Guyana". Distr. — Colombia to Bolivia, Brazil, and Paraguay.
- DISPAR Schiner, 1868:54. Type-locality: "Colombia". Distr. — Colombia.
- DOMINICANA James, 1967a:2. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.
- ECUADORENSIS Enderlein, 1914b:601. Type-locality: Ecuador, Archidona. Distr. — Colombia, Ecuador, Peru.
- ERECTA McFadden, in McFadden & James, 1969:315. Type-locality: Mexico, Durango, Durango. Distr. — U.S.A. (Arizona, Texas); Mexico (south to Oaxaca).
- ERECTISPINIS Lindner, 1935:408. Type-locality: Colombia, Hacienda Pehlke. Distr. — Colombia.
- FASCIPES Walker, 1854:17. Type-locality: Brazil, Pará, Santarém. Distr. — Panama to Peru and Brazil (Pará).
- FASSLI Lindner, 1949:862. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia.
- FERRUGINEA Enderlein, 1914b:603. Type-locality: Ecuador, Baños. Distr. — Ecuador.
- FLAVIMANA Gerstaecker, 1857:291. Type-locality: "Brazil". Distr. — Guyana, Brazil.
- FLAVIPENNIS Enderlein, 1914b:602. Type-locality: Ecuador, Santa Inez. Distr. — Ecuador.
- FORMOSA James, 1940a:138. Type-locality: Colombia, Boyaca, Muzo. Distr. — Colombia, ?Brazil ("Amazon").
- GENICULATA Gerstaecker, 1857:290. Type-locality: "New Granada". Distr. — ?Colombia.
- COLBACHI James, 1953b:310. Type-locality: Argentina, Tucumán, Villa Padre Monti. Distr. — Argentina.
- GRACHICORNIS Gerstaecker, 1857:295. Type-locality: "South Brazil". Distr. — Brazil (Rio Grande do Norte and Minas Gerais to Santa Catarina), Paraguay, Argentina (Santiago del Estero). Ref. — Lindner, 1926 (larva).
- lanuginosa* Gerstaecker, 1857:297. Type-locality: "Brazil".
- HELVIPENNIS Enderlein, 1914b:601. Type-locality: Peru, Chanchamayo. Distr. — Ecuador, Peru.



- HYBRIDA Gerstaecker, 1857:307. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara, São Paulo).
- IMITANS Curran, 1925:244. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara, São Paulo).
- JAMESI Lindner, 1949:861. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LASIOPTHALMA Williston, 1896b:301. Type-locality: St. Vincent. Distr. — St. Vincent, Granada. Not *Cyphomyia lasiophthalma* Williston, 1900.
- LEUCOCEPHALA Wiedemann, 1819:55. Type-locality: "Brazil". Distr. — Peru, Brazil (north to Mato Grosso and Goiás), Paraguay, Argentina (Misiones, Tucumán). Ref. — Iide, 1963b (suppl. deser. & figs., adult and larva).
- LONGICORNIS (Walker), 1857:120 (*Acanthina*). Type-locality: "Amazon Region". Distr. — Brazil.
- MACULATA (Fabricius), 1805:85 (*Stratiomys*). Type-locality: "South America". Distr. — South America.
- MARGINATA Loew, 1866a:148 (Centuria 6, no. 31). Type-locality: "Cuba". Distr. — U.S.A. (Florida); Cuba, San Domingo, Haiti, Puerto Rico, Costa Rica. Ref. — McFadden, 1967 (larva).
scutellata Cresson, 1919:174. Type-locality: "Costa Rica".
- NEIVAI James, 1940:147. Type-locality: "Amazon". Distr. — Brazil.
- NIGRITARSIS Enderlein, 1914b:600. Type-localities: Peru, Loreto, Pebas; Brazil, Pará. Distr. — Brazil, Peru.
- NUBILPENNIS James, 1939a:600. Type-locality: Guatemala, Cayuga. Distr. — Mexico (Veracruz), Guatemala.
- OBSCURA (Jaenicke), 1867:325 (*Rondania*). Type-locality: "Mexico". Distr. — Mexico.
- OCHRACEA Giglio-Tos, 1891:2, 1892:10. Type-locality: "Mexico". Distr. — Mexico.
- ORNATA Walker, 1850:xvii. Type-locality: Brazil, Pará. Unidentifiable (type lost).
- PIETA Schiner, 1868:53. Type-locality: "South America". Distr. — Canal Zone, Panama.
- PILOSISSIMA Gerstaecker, 1857:292. Type-locality: "Mexico". Distr. — Mexico (Coahuila to Guerrero). Refs. — James, 1957a (larva); McFadden, 1967a (larva).
- PLANIFRONS James, 1939a:602. Type-locality: Colombia, Bogotá. Distr. — Colombia, ?Brazil ("Amazon"), Peru.
- PULCHELLA Gerstaecker, 1857:309. Type-locality: "Brazil". Distr. — Costa Rica to Peru, Guyana and Brazil; Trinidad.
maculata Fabricius of Wiedemann (in part), 1830:57.
- REGULARIS Curran, 1929:5. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- RUBRA Loew, 1866a:148 (Centuria 6, no. 30). Type-locality: "Cuba". Distr. — Cuba.
- SCALARIS Bigot, 1875:487. Type-locality: "Mexico". Distr. — Mexico.



- SCHWARZI James, 1940a:135. Type-locality: Colombia, Western Cordillera, Cali District. Distr. — Colombia.
- SHANNONI James, 1939a:601. Type-locality: Panama, Gatún Lake, Cano Saddle. Distr. — Panama (Cano Saddle, Gatún Lake).
- SIMPLEX Walker, 1860:268. Type-locality: "Mexico". Distr. — Mexico (Chiapas, Veracruz); Guatemala.
- bequaerti* James, 1938:198. Type-locality: Guatemala, Pochuta, Santa Emilia.
- SOUZALOPESI Iide, 1967c:226. Type-locality: Brazil, Pará, Belém, Santa Bárbara. Ref. — Iide, 1967c (Biology, larva, figs.).
- SPECIOSA Lindner, 1951:249. Type-locality: Peru, San Martín, Tingo María. Distr. — Peru.
- SULCIFRONS Curran, 1929:5. Type-locality: Mexico, Veracruz, La Bucna Ventura. Distr. — Mexico, Costa Rica.
- TOMENTOSA Gerstaecker, 1857:294. Type-locality: "Mexico". Distr. — Mexico (Nuevo Leon to Puebla). Ref. — McFadden & James, 1969 (suppl. descr. as *similis*).
- similis* Bellardi, 1859:23, 1861:223. Type-locality: "Mexico".
- UNICOLOR (Walker), 1854:61, 312 (*Clitellaria*). Type-locality: Brazil, Amazonas, Tefé (as Ega). Distr. — Colombia, Peru, Brazil (Amazonas).
- cianocephala* Gerstaecker, 1857:274. Type-locality: "South America".
- VAMPES Gerstaecker, 1857:283. Type-localities: "Mexico, Costa Rica, Venezuela". Distr. — Mexico (Veracruz) to Colombia and Venezuela. Ref. — James, 1937 (suppl. descr.).
- VERTICALIS Gerstaecker, 1857:287. Type-localities: "Surinam, Guyana". Distr. — Ecuador, Peru, Surinam, Guyana, Brazil.
- VIOLACEA Macquart, 1855:60 (separate, p. 40). Type-locality: "Brazil". Distr. — Colombia, Ecuador, Brazil (Pará).
- WIEDEMANNI Gerstaecker, 1857:302. Type-locality: Brazil, Pará. Distr. — Mexico to northern Argentina, Trinidad.
- albitarsis* Fabricius of Wiedemann, 1824, misident.
- ?flavispinis* Macquart, 1855:60 (separate, p. 40). Type-locality: "South America".
- peruana* Lindner, 1933b:329 (as subspecies). Type-locality: Peru, Iquitos, San Roque.
- ssp. BURMEISTERI Lindner, 1949:863. Type-locality: "Amazonia". Distr. — Brazil.
- WILLISTONI (Enderlein), 1914b:604 (*Gyneuryptaria*; nom. nov. for *lasiophthalma* Williston). Distr. — Mexico (Sinaloa to Guerrero).
- lasiophthalma* Williston, 1900:244 (preocc. Williston, 1896). Type-localities: Mexico, Jalisco, San Blás, and Guerrero, Acaguizotla, Dos Arroyas, and Tierra Colorada.
- XANTHOBASIS James, 1940a:134. Type-locality: "Amazon" (?Brazil). Distr. — ?Brazil.
- YPSILON James, 1940a:148. Type-locality: unknown. Distr. — Unknown, probably Brazil.
- ZERNYI Lindner, 1929:264. Type-locality: Brazil, Pará, Tapcrinha near Santarém. Distr. — Brazil (Pará to Rio de Janeiro).

Genus DICYPHOMA James

- DICYPHOMA James, 1937:151. Type-species, *Cyphomyia schaefferi* Coquillett (orig. des.). Ref. — James, 1962b (review, adults and larvae).
- RYCKMANI James, 1962b:18. Type-locality: Mexico, Oaxaca. Distr. — Mexico.
- SCHAEFFERI (Coquillett), 1904:32 (*Cyphomyia*). Type-locality: U.S.A., Texas, presumably Brownsville. Distr. — U.S.A. (north to Kansas) to Panama. Refs. — James, 1937, 1962b (suppl. descr.); McFadden, 1967 (larva).

Genus LABOCERINA Enderlein

- LABOCERINA Enderlein, 1914b:579. Type-species, *Stratiomys atrata* Fabricius (orig. des.).
- Labacerino* Enderlein, 1914b:615 (error).
- Labocerino*, error.
- ATRATA (Fabricius), 1805:83 (*Stratiomys*). Type-locality: "South America". Distr. — Mexico (Veraacruz) to Colombia and Guyana.
- carbonaria* Bellardi, 1862:11, 1864:210 (*Chordonota*). Type-locality: Mexico, Veraacruz, Tuxpango, near Orizaba.

Unplaced Species

- FLAVITARSIS Enderlein, 1914b:613 (*Lasiopa*). Type-locality: "Costa Rica".
- VITTATA Wulp, 1881:155 (*Lasiopa*). Type-locality: "Argentina".

Tribe CLITELLARIINI

Genus ABAVUS Enderlein

- ABAVUS Enderlein, 1914e:23. Type-species, *priscus* Enderlein (orig. des.).
- MACULATUS Lindner, 1933a:204. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- MULTISIGNATUS Lindner, 1949:874. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PRISCUS Enderlein, 1914e:24. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (Rio de Janeiro to Santa Catarina).

Genus ACROPELTATES Kertész

- ACROPELTATES Kertész, 1923:118. Type-species, *fasciata* Kertész (orig. des.).
- DIVERSICORNIS Kertész, 1923:121. Type-locality: "Colombia". Distr. — Colombia.
- FASCIATA Kertész, 1923:120. Type-localities: Peru, Urubamba River, Meshagua and Ucayali River, Unini. Distr. — Colombia.
- KERTESZI Szilády, 1929:250. Type-locality: Costa Rica, Cartago, Turrialba. Distr. — Costa Rica.



Genus ADOXOMYIA Bezzi

- Adoxomyia* Kertész, 1907:499. Nomen nudum.
- ADOXOMYIA Bezzi, 1908b:75. Type-species, *Clitellaria dahlia* Meigen (orig. des.). Refs. — Kertész, 1923 (rev., keys); James & McFadden, 1969 (rev. Nearctic spp., adults and larvae).
- Euclitellaria* Kertész, 1923:96, 101. Type-species, *Clitellaria heminopla* Wiedemann (orig. des.).
- Clitellaria*, authors, not Meigen.
- FENESTRATA (Macquart), 1846:54 (separate, p. 182) (*Ephippium*). Type-locality: Mexico, Yucatan, Merida.
- MARGINATA James, in James & McFadden, 1969:273. Type-locality: U.S.A., Arizona, Tucson. Distr. — U.S.A. (Arizona), Mexico (Sonora).
- MICHENERI James, 1950:71. Type-locality: U.S.A., Texas, Carrizo Springs. Distr. — U.S.A. (Texas), Mexico (Hidalgo).
- NIGRINA Lindner, 1949:873. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- REGULARIS James, in James & McFadden, 1969:269. Type-locality: U.S.A., Arizona, Baboquivari Mts. Distr. — U.S.A. (Arizona), Mexico (Durango).
- WEYRAUCHI Lindner, 1951:258. Type-locality: Peru, Lima, Atocongo. Distr. — Peru.

Genus ANAPOSASMA Lindner

- ANAPOSASMA Lindner, 1969:12. Type-species, *amoenum* Lindner (mon.).
- AMOENUM Lindner, 1969:12. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus BRACHYCARA Thomson

- BRACHYCARA Thomson, 1869:460. Type-species, *ventralis* Thomson (mon.). Ref. — James, 1962a (key to known species).
- Euryneurasoma* Johnson, 1913:51. Type-species, *slossonae* Johnson (orig. des.).
- Neurota* Curran, 1931:2. Type-species, *Sargus bicolor* Wiedemann (orig. des., misident.) = *slossonae* (Johnson).
- DIGTATA James, 1966:537. Type-locality: Galápagos Islands, Santa Cruz Island, Academy Bay. Distr. — Galápagos Islands.
- MACULATA (James), 1953:2 (*Euryneurasoma*). Type-locality: Bahamas, South Bimini Island. Distr. — U.S.A. (Florida), Bahamas, Bermuda, Cuba, Antigua.
- SLOSSONÆ (Johnson), 1913:51 (*Euryneurasoma*). Type-locality: U.S.A., Florida, Biscayne Bay. Distr. — U.S.A. (Florida), Bahamas, Bermuda, Cuba, Puerto Rico, Dominica.

Genus CALOPARYPHUS James

- Euparyphus*, subg. CALOPARYPHUS James, 1939c:49. Type-species, *Oxycera crotchi* Osten Sacken (orig. des.).



DECEMMACULATUS (Osten Sacken), 1886:40 (?*Euparhyphus*). Type-locality: Mexico, Northern Sonora. Distr. — U.S.A. (Arizona), Mexico (Sonora).

Genus DIAPHOROSTYLUS Kertész

- DIAPHOROSTYLUS Kertész, 1908:335. Type-species, *Euryneura nasica* Williston (by present designation). Ref. — Kertész, 1908 (revision and key).
- FLAVIPES Kertész, 1908:337. Type-locality: Peru, Callanga. Distr. — Costa Rica to Peru.
- INTERRUPTUS James, 1939:597. Type-locality: Panama, Alajuelo. Distr. — Costa Rica, Panama.
- NASICUS (Williston), 1888:252 (*Euryneura*). Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil.
- SIGNATIPES Kertész, 1908:338. Type-localities: Peru, Urubamba River, Meshagua, and Ucayali River, Unini. Distr. — Peru.

Genus DIEURYNEURA James

- DIEURYNEURA James, 1937:152. Type-species, *callosa* James (orig. des.) = *stigma* (Giglio-Tos).
- STIGMA (Giglio-Tos), 1891:2 (1892:13) (*Clitellaria*). Type-locality: "Mexico". Distr. — Southwest U.S.A., northern Mexico. Ref. — McFadden, 1967 (larva, as *obscura*).
- obscura* Coquillett, 1902:98 (*Aoehletus*). Type-localities: U.S.A., California, Los Angeles County, and Arizona, Sulphur Spring Valley.
- callosa* James, 1937:153. Type-locality: U.S.A., Arizona, Baboquivari Mts.

Genus DITYLOMETOPA Kertész

- DITYLOMETOPA Kertész, 1923:114. Type-species, *elegans* Kertész (orig. des.).
- ELEGANS Kertész, 1923:115. Type-localities: Peru, Vileanota, and Brazil, Espírito Santo. Distr. — Peru, Brazil (Pará to Espírito Santo).

Genus EUPARYPHUS Gerstaecker

- EUPARYPHUS Gerstaecker, 1857:314. Type-species, *Cyphomyia elegans* Wiedemann (mon.).
- Euparhyphus*, authors (error or emendation).

Subgenus EUPARYPHUS Gerstaecker

- CARBONARIUS Giglio-Tos, 1891:2 (also 1892:12, 1893:109). Type-locality: "Mexico". Distr. — Southwestern U.S.A. to Guatemala.
- ELEGANS (Wiedemann), 1830:58 (*Cyphomyia*). Type-locality: "Mexico". Distr. — Mexico.
- ELONGATULUS Williston, 1900:249. Type-localities: Mexico (Guerrero, Omilteme and Xucumanatlan). Distr. — Mexico.
- TRICOLOR Osten Sacken, 1886:40. Type-locality: Mexico (Sonora). Distr. — U.S.A. (Arizona), Mexico (Sonora).



Subgenus ACHLETUS Osten Sacken

- ACHLETUS Osten Sacken, 1886:33 (as genus). Type-species, *cinctus* Osten Sacken (mon.).
- BISTRATUS Williston, 1896b:300. Type-locality: St. Vincent. Distr. — St. Vincent.
- BRASILIENSIS Lindner, 1949:876. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CINCTUS Osten Sacken, 1886:38. Type-locality: Mexico, Northern Sonora. Distr. — Western U.S.A.; Mexico (south to San Luis Potosí). Ref. — McFadden, 1967 (larva).
- HAMIFER James, 1940b:120. Type-locality: Costa Rica, San José, La Caja. Distr. — Costa Rica.
- PERUVIANUS Lindner, 1941:184. Type-locality: Peru, Ayacucho, Cangallo. Distr. — Peru.

Genus EURYNEURA Schiner

- EURYNEURA Schiner, 1867:308 (1868:56). Type-species, *Stratiomys fascipennis* Fabricius (orig. des.). Refs. — Kertész, 1908 (revision, key); Iide, 1962b (Brazilian species, larva, transl. of Kertész key).
- ELEGANS Williston, 1888:252. Type-locality: not stated. Distr. — Mexico (Guerrero) to Brazil (south to São Paulo) and Argentina (Tucumán). Ref. — Iide, 1968b (redeser.).
- FASCIPENNIS (Fabricius), 1805:373 (*Stratiomys*). Type-locality: "South America". Correction of *fasciata*, error. Distr. — Peru, Brazil (Pará).
- fasciata* Fabricius, 1805:84 (*Stratiomys*). Type-locality: "South America"
- KERTESZI Iide, 1968b:266. Type-locality: Brazil, São Paulo. Distr. — Brazil (São Paulo, Santa Catarina).
- MEXICANA Kertész, 1908:327. Type-localities: Mexico, Veracruz, Orizaba and Acultzingo. Distr. — Mexico, Costa Rica.
- PERUANA Kertész, 1908:329. Type-locality: "Peru". Distr. — Peru.
- PROPINQUA Schiner, 1868:57. Type-locality: "Colombia". Distr. — U.S.A. (Arizona) to Colombia and Venezuela. Ref. — Giglio-Tos, 1892 (suppl. deser.).
- PYGMAEA (Bellardi), 1862:12, 1864:211 (*Clitellaria*). Type-locality: Mexico, Veracruz, Tuxpango, near Orizaba. Distr. — Mexico to Brazil (south to Santa Catarina) and Paraguay.
- ROBUSTA Kertész, 1908:332. Type-locality: "Brazil". Distr. — Brazil (Rio de Janeiro to Rio Grande do Sul), Argentina (Tucumán). Ref. — Iide, 1968b (redeser., larva).
- RUFIFRONS Curran, 1934:314. Type-locality: Guyana, Bartica, Kartabo. Distr. — Guyana.

Genus GLARIOPSIS Lindner

- GLARIOPSIS Lindner, 1935:412. Type-species, *brasiliana* Lindner (mon.).
- BRASILIANA Lindner, 1935:413. Type-locality: Brazil, Rio de Janeiro, Angra, Juçaral. Distr. — Brazil.

Genus GLARIS Kertész

GLARIS Kertész, 1923:107. Type-species, *mexicana* Kertész (orig. des.).
MEXICANA Kertész, 1923:108. Type-locality: "Mexico". Distr. — Mexico.

Genus GRYPOMYIA Kertész

GRYPOMYIA Kertész, 1923:112. Type-species, *gracilis* Kertész (orig. des.).
GRACILIS Kertész, 1923:113. Type-locality: Peru, Cuzco, Vilcanota. Distr. — Peru.

Genus HOMALARTHRIA Lindner

HOMALARTHRIA Lindner, 1933b:332. Type-species, *nigra* Lindner (orig. des.).
NIGRA Lindner, 1933b:332. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.

Genus LEUCOPTILUM James

LEUCOPTILUM James, 1943b:380. Type-species, *plaumanni* James (orig. des.).
BASSLERI James, 1943b:382. Type-locality: Peru, Lower Rio Tapiche. Distr. — Peru.
PLAUMANNI James, 1943b:380. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Rio de Janeiro to Santa Catarina).
niveipenicillata Lindner, 1949:872 (*Adoxomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia.

Genus MERINGOSTYLUS Kertész

MERINGOSTYLUS Kertész, 1908:334. Type-species, *schineri* Kertész (mon.).
CHIQUITANUS Lindner, 1930:275. Type-locality: Bolivia, Chiquitos. Distr. — Bolivia.
SCHINERI Kertész, 1908:334. Type-locality: "Brazil". Distr. — Brazil (Rio de Janeiro to Santa Catarina).
flavipes Kertész, 1908:335 (*Euryneura*). Schiner's manuscript name published by Kertész.

Genus MONACANTHELLA James

MONACANTHELLA James, 1936a:49. Type-species, *gracilis* James (orig. des.).
GRACILIS James, 1936a:50. Type-locality: Costa Rica, Hamburg Farm. Distr. — Costa Rica.

Genus PACHYPTILUM Lindner

PACHYPTILUM Lindner, 1969:13. Type-species, *hennigi* Lindner (mon.).
HENNIGI Lindner, 1969:13. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



Genus PLATOPSOMYIA James

PLATOPSOMYIA James, 1937:154. Type-species, *flavida* James (orig. des.).

FLAVIDA James, 1937:154. Type-locality: Cuba, Habana, Santiago de las Vegas. Distr. — Cuba.

Genus PROGRYPOMYIA Lindner

PROGRYPOMYIA Lindner, 1949:871. Type-species, *nigra* Lindner (mon.).

NIGRA Lindner, 1949:872. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Unplaced Species

LIBURNA Walker, 1849:528 (*Oxycera*). Type-locality: "Jamaica".

Subfamily CHRYSOCHLORINAE

Genus CACOSIS Walker

CACOSIS Walker, 1856:83. Type-species, *Sargus niger* Wiedemann (mon.).

GRANDIS Schiner, 1868:67. Type-locality: "South America". Distr. — South America.

NIGER (Wiedemann), 1830:28 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil.

Genus CHAETOSARGUS Roeder

Drasteria Brauer, 1882:78 (preocc. Hübner, 1816). Type-species, *robusta* Brauer (orig. des.).

CHAETOSARGUS Roeder, 1894:169. Type-species, *Sargus hirticornis* Wiedemann (mon.).

Parahermetia Lindner, 1928:241. Type-species, *seitzi* Lindner (mon.).

HIRTICORNIS (Wiedemann), 1830:28 (*Sargus*). Type-locality: "Brazil". Distr. — Brazil (Espírito Santo to Rio Grande do Sul). Ref. — Iide, 1967a (descr.).

ROBUSTUS (Brauer), 1882:79 (*Drasteria*). Type-locality: Brazil, Bahia. Distr. — Brazil.

SECUNDUS (Albuquerque), 1955:132 (*Parahermetia*). Type-locality: Brazil, São Paulo, Umuarama, Campos do Jordão. Distr. — Brazil.

SEITZI (Lindner), 1928:241 (*Parahermetia*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.

Genus CHRYSOCHLORINA James

CHRYSOCHLORINA James, 1939b:33. Type-species, *Sargus vespertilio* Fabricius (orig. des.). Refs. — James, 1939b (key); Iide, 1965 (immature forms); Iide, 1966 (Brazilian species); Iide, 1968a (definition of genus).

Chrysochlora Authors, not Latreille. Note: The type-species of *Chrysochlora* is the only originally included species, *Sargus amethys-*



- tinus* Fabricius. Enderlein's designation of *Sargus vespertilio* Fabricius as the type is therefore, invalid.
- ALBIPES James, 1939:36. Type-locality: Brazil, Goiás. Distr. — Brazil (Mato Grosso to Minas Gerais and Santa Catarina).
- ?*sp.*, probably *albipes*. Iide, 1965; immature forms.
- BEZZIANA Iide, 1966:97. Type-locality: Brazil, São Paulo, Ipiranga. Distr. — Brazil.
- CASTANEA (Maequart), 1838:202 (separate, p. 198) (*Chrysochlora*). Type-locality: "Guyana". Distr. — Canal Zone, Guyana, Brazil (Território de Roraima, Território de Rondônia, Pará). Ref. — Iide, 1967b (suppl. descr.).
- COSTALIMAI Iide, 1966:94. Type-locality: Brazil, São Paulo, Juquiá. Distr. — Brazil (Espírito Santo, São Paulo). Ref. — Iide, 1967b (suppl. descr.).
- CURRANI Iide, 1966:105. Type-locality: Brazil, Pará, Cachimbo. Distr. — Peru, Brazil (Amazonas, Rio Negro, Pará, Rio de Janeiro).
- FASCIATA (Thomson), 1869:460 (*Chrysochlora*). Type-locality: Galápagos Islands. Distr. — Galápagos Islands. Ref. — James, 1966a (as *dubia*; suppl. descr.); Iide, 1971:503, figs. 23-33; systematic position; suppl. descr.).
- dubia* Curran, 1932c:348. Type-locality: Galápagos Islands, Floreana, Postoffice Bay, Seaside, Ranvik.
- FEMORALIS (Curran), 1929:4 (*Chrysochlora*). Type-locality: Colombia, Magdalena, Rio Frio. Distr. — Colombia.
- FLAVESCENS (James), 1937:151 (*Chrysochlora*). Type-locality: Mexico, Yucatán. Distr. — Mexico.
- FROSTI James, 1939b:35. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone, Panama, Colombia.
- INCOMPLETA (Curran), 1929:3 (*Chrysochlora*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Bolivia; Brazil (Pará to Mato Grosso and São Paulo).
- MACULIVENTRIS (Rondani), 1850:190 (*Chrysochlora*). Type-locality: "Venezuela".
- PLURICOLOR (Bigot), 1879:209 (*Chrysochlora*). Type-locality: "Brazil". Distr. — Brazil (Minas Gerais to São Paulo).
- PULCHRA (Williston), 1900:243 (*Chrysochlora*). Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico (Jalisco to Morelos and Oaxaca).
- PURPUREA (Walker), 1860:271 (*Chrysochlora*). Type-locality: "Mexico". Distr. — Mexico.
- QUADRILINEATA (Bigot), 1887:26 (*Chrysochlora*). Type-locality: "Cuba".
- SIMILIS (Macquart), 1855:64 (separate, p. 44) (*Chrysochlora*). Type-locality: "South America". Distr. — Brazil (Mato Grosso to Santa Catarina).
- VARIA (Curran), 1929:3 (*Chrysochlora*). Type-locality: Guyana, Bartica. Distr. — Honduras to Bolivia, Guyana, and Brazil (south to Mato Grosso).
- VESPERTILIO (Fabricius), 1805:259 (*Sargus*). Type-locality: "South America". Distr. — Mexico to Bolivia and Brazil (South to Santa Catarina).
- willistoni* Curran, 1929:2 (*Chrysochlora*). Type-locality: Brazil, Mato Grosso, Chapada.



Genus PELAGOMYIA Williston

PELAGOMYIA Williston, 1896a:48 (no species), 1896b:299. Type-species, *albitalis* Williston (mon.).

ALBITALIS Williston, 1896b:299. Type-locality: St. Vincent (B.W.I.). Distr. — St. Vincent.

albitatus Iide, 1971:497, error.

ILLUCENS James, 1967a:4. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.

Genus TRICHOERCOCERA Lindner

TRICHOERCOCERA Lindner, 1928:243. Type-species, *ptecticoides* Lindner (mon.).

PTECTICOIDES Lindner, 1928:243. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil (Guanabara, São Paulo).

Subfamily HERMETIINAE

Genus AMPHILECTA Brauer

AMPHILECTA Brauer, 1882:80, 86. Type-species, *superba* Brauer (orig. des.).

SUPERBA Brauer, 1882:80. Type-locality: "Congorh?". Distr. — Brazil.

Genus CHAETOHERMETIA Lindner

CHAETOHERMETIA Lindner, 1929:261. Type-species, *apicalis* Lindner (mon.).

AENEA Lindner, 1949:869. Type-locality: Guyana, Essequibo River, Moraballi Creek. Distr. — Guyana.

APICALIS Lindner, 1929:261. Type-locality: Brazil, Pará, Tapcrinha, near Santarém. Distr. — Brazil.

Genus HERMETIA Latreille

HERMETIA Latreille, 1804:192. Type-species, *Musca illucens* Linnaeus (mon.). Refs. — Curran, 1934 (partial key); James, 1935a (partial key); James & Wirth, 1967 (*aurata* group); James, 1967b (*comstocki* group).

Acrodesmia Enderlein, 1914c:3. Type-species, *luederwaldti* Enderlein (orig. des.) = *albitarsis* Fabricius.

AENEIPENNIS Giglio-Tos, 1892:11 (as *flavipes* var.). Type-localities: "Brazil and Mexico". Distr. — Brazil, Mexico.

ALBITARSIS Fabricius, 1805:63. Type-locality: "South America". Distr. — Central Mexico (north to San Luis Potosí) to Paraguay and Southern Brazil (Santa Catarina); Trinidad. Ref. — Lindner, 1965 (larva).

apicalis Wiedemann, 1830:25. Type-locality: "South America".

bimaculata Macquart, 1834:228. Type-locality: "Brazil". (Synonymy by Williston, with a query).



- planifrons* Macquart, 1846:178 (separate, p. 50). Type-locality: Mexico, Yucatán, Merida.
- luederwaldti* Enderlein, 1914c:3 (*Acrodesmia*). Type-locality: Brazil, Santa Catarina.
- AMPULLA James, 1938:196. Type-locality: Brazil, Rio Madeira. Distr. — Brazil.
- ANTHIDIUM James, in James & Wirth, 1967:5. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- AURATA Bellardi, 1859:27, 1861:227. Type-locality: Mexico, Michoacán, Morelia. Distr. — Mexico (Sinaloa to San Luis Potosí and Oaxaca).
- AURINOTATA Lindner, 1935:411. Type-locality: Guyana, Georgetown. Distr. — Guyana. Ref. — Lindner, 1949:870 (redscr.).
- EEEBEI Curran, 1934:317. Type-locality: Guyana (Kartabo). Distr. — Guyana.
- BURMEISTERI Lindner, 1949:868. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- CALLIFERA Lindner, 1928:240. Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- CERIA Williston, 1900:242. Type-locality: Mexico, Durango, Ventanas. Distr. — Mexico.
- CERIOGASTER Williston, 1888:246. Type-locality: not stated. Distr. — Presumably Brazil.
- CHRYSOPILA Loew, 1872:56 (Centuria 10, no. 11). Type-locality: U.S.A., Texas. Distr. — U.S.A. (north to Kansas and Colorado), Mexico (Coahuila, Nuevo Leon, Tamaulipas).
- CINGULATA Greene, 1940:153. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.
- COARCTATA Macquart, 1846:178 (separate, p. 50). Type-locality: "Mexico". Distr. — Mexico (north to Sinaloa) to southern Brazil (Santa Catarina).
- COMSTOCKI Williston, 1885:125. Type-locality: U.S.A., Arizona. Distr. — Southwestern U.S.A. (Texas to New Mexico); Mexico (Nuevo Leon, Sonora).
- ssp. MEXICANA James, 1967b:63. Type-locality: Mexico, Morelos, Yautepec. Distr. — Mexico (North to Veracruz and Michoacán) to El Salvador.
- CONCINNA Williston, 1900:241. Type-localities: U.S.A., New Mexico, Arizona. Distr. — Southwestern U.S.A. (New Mexico to California); Mexico (Chihuahua to Jalisco). Ref. — McFadden, 1967 (larva).
- CONDOR Lindner, 1951:256. Type-locality: Peru, San Martín, Tingo Maria. Distr. — Peru.
- CONJUNTA James, in James & Wirth, 1967:10. Type-locality: Mexico, Isthmus of Tehuantepec (?Oaxaca). Distr. — Mexico (Morelos, ?Oaxaca).
- CORNITHORAX (Lindner), 1928:238 (*Acrodesmia*). Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- CRABRO Osten Sacken, 1886:29. Type-locality: Guatemala, El Reposo. Distr. — Guatemala, Honduras.



- CURRANI Lindner, 1949:864. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- EISENI Townsend, 1895:594. Type-locality: Mexico, Baja California del Sur, San José del Cabo and El Toste. Distr. — Mexico.
- FIMBRIATA (Lindner), 1931:307 (*Aerodesmia*). Type-locality: Brazil, Espírito Santo, Porto do Cadi. Distr. — Brazil.
- FLAVIPES Wiedemann, 1830:26. Type-locality: "Brazil". Distr. — Honduras to Guyana, Peru, and Brazil (south to São Paulo). Ref. — Lindner, 1949 (suppl. descr.).
andreas Walker, 1849:515 (*Sargus*). Type-locality: not given.
setigera James, 1938:198. Type-locality: Brazil, Rio Madcira.
- FLAVOSCUTATA Bigot, 1879:201. Type-locality: "Mexico". Distr. — Mexico (Veracruz, ?Oaxaca, Campeche, Yucatan). Ref. — James, 1967b (suppl. descr.).
- FORMICA Osten Sacken, 1886:32. Type-locality: Panama, Bugaba. Distr. — Panama.
- FULVA Walker, 1854:22. Type-locality: Brazil, Pará. Distr. — Brazil.
- GENICULATA Macquart, 1855:61 (separate, p. 41). Type-locality: "South America". Distr. — South America.
- GONCALVESI Albuquerque, 1955:129. Type-locality: Brazil, Bahia, Salvador. Distr. — Brazil.
- HUNTERI Coquillett, 1909:212. Type-localities: U.S.A., Texas, Hondo and Cotulla. Distr. — Southwest U.S.A., Mexico (Chihuahua, Baja California).
- ILLUCENS (Linnaeus), 1758:589 (*Musca*). Type-locality: "South America". Distr. — U.S.A. (widespread) to Bolivia and Northern Argentina; Bermuda and throughout the West Indies; Old World tropics, in general; southern Europe; Pacific Islands. Spread by commerce.
leucopa Linnaeus, 1767:983 (*Musca*). Type-locality: "America".
mucens, error.
rufiventris Fabricius, 1805:63. Type-locality: "South America" (not *rufiventris* Walker).
pellucens Macquart, 1834: plate 4, figs. 11-12, and legend. (Lapsus).
nigrifacies Bigot, 1879:200. Type-locality: "Mexico".
nigritibia Enderlein, 1914c:9. Type-locality: Brazil, Santa Catarina. (as var.).
- IMPRESSA James, in James & Wirth, 1967:13. Type-locality: Mexico, Morelos, Temixco. Distr. — Mexico.
- ITATAIENSIS Lindner, nom. nov. for *flavipes* Lindner.
flavipes Lindner, 1933b:330 (preocc. Wiedemann, 1830). Type-locality: Brazil, Espírito Santo, Alegre.
- LATIVENTRIS Bellardi, 1859:27, 1861:227. Type-locality: Mexico, Tamaulipas, Tampico. Distr. — U.S.A. (Texas to Arizona) to Panama.
var. A. Bellardi, 1862:8, 1864:207. Mexico, Veracruz, Tuxpango near Orizaba.
- MELANDERI James & Wirth, 1967:8. Type-locality: U.S.A., Arizona, Globc. Distr. — U.S.A. (Arizona, New Mexico), Mexico (Sinaloa).
- MITIS Curran, 1934:318. Type-locality: Guyana, Bartica Distr., Kartabo. Distr. — Guyana.



- NANA Lindner, 1935:410. Type-localities: Brazil, São Paulo, and Guanabara, Rio de Janeiro. Distr. — Brazil (São Paulo, Guanabara).
- PANAMENSIS Greene, 1940:151. Type-locality: Canal Zone, Mandingo, Distr. — Panama, Canal Zone.
- PTEROCAUSTA Osten Sacken, 1886:33. Type-locality: Panama, Bugaba. Distr. — Canal Zone.
- PULCHRA Wiedemann, 1830:24. Type-locality: "Surinam". Distr. — Surinam and Guyana to Brazil (south to Santa Catarina).
- REINHARDI James, 1935a:169. Type-locality: U.S.A., Texas, Dilley (not Oilley, as published). Distr. — U.S.A. (Texas to Arizona), Mexico (Nuevo Leon, San Luis Potosí).
reinharti, error.
- RELICTA Osten Sacken, 1886:30. Type-locality: Panama, Bugaba. Distr. — Panama.
- RUFITARSIS Macquart, 1846:177 (separate, p. 49). Type-locality: "Colombia". Distr. — Colombia.
- SCUTELLATA Macquart, 1855:61 (separate, p. 41). Type-locality: "South America, shores of the Amazon River". Distr. — Brazil.
- SEXMACULATA Macquart, 1834:229. Type-locality: Puerto Rico. Distr. — U.S.A. (north to Tennessee), Cuba, Puerto Rico, Dominican Republic, Brazil (Natal).
nucis James, 1935a:166. Type-locality: Cuba, Santo Tomas, P. de Zapata.
- SPIEHOES Curran, 1934:318. Type-locality: Guyana, Bartica. Distr. — Kartabo. Distr. — Guyana.
- SUBPELLUCIDA James & Wirth, 1967:11. Type-locality: U.S.A., Arizona, Globe. Distr. — U.S.A. (Arizona), Mexico (Sonora, Sinaloa, Nayarit, Baja California).
- TEEVANI Curran, 1934:316. Type-locality: Guyana, Bartica. Distr. — Kartabo. Distr. — Guyana.
- VARIPENNIS Bigot, 1879:202. Type-locality: "Brazil". Distr. — Brazil.
variipennis, emendation, or error.
- VIRGATA Lindner, 1949:866. Type-localities: Brazil, Santa Catarina, Nova Teutônia and São Paulo, Jaguará.

Genus PATAGIOMYIA Lindner

- PATAGIOMYIA Lindner, 1933b:331. Type-species, *cyphomyioides* Lindner (orig. des.).
- CYPHOMYIOIDES Lindner, 1933b:331. Type-locality: Peru, Mishuyacu. Distr. — Colombia (Amazonas), Peru.

Subfamily STRATIOMYINAE

Tribe STRATIOMYINI

Genus ANOPISTHOCRANIA Lindner

- ANOPISTHOCRANIA Lindner, 1935:406. Type-species, *zonata* Lindner (mon.).
- ZONATA Lindner, 1935:407. Type-locality: Brazil, Rio de Janeiro, Maromba, Itatiaia. Distr. — Brazil (Rio de Janeiro, Santa Catarina). Ref. — Lindner, 1969 (suppl. deser.).



Genus CHLOROMELAS Enderlein

- CHLOROMELAS Enderlein, 1914b:607. Type-species, *Odontomyia heteroneura* Macquart (orig. des.) = *cuprina* (Wiedemann).
- BARBATA Lindner, 1935:410. Type-locality: Brazil, São Paulo. Distr. — Brazil.
- BRUNNEIPICTA James, 1953b:316. Type-locality: Argentina, Salta, Ruiz de los Llanos. Distr. — Argentina (Salta, Tucumán).
- COATI Lindner, 1951:254. Type-locality: Peru, San Martín, Tingo Maria, Río Huallaga. Distr. — Peru.
- CORDOBAENSIS (Lindner), 1951:253 (*Hedriodiscina*). Type-locality: Argentina, Córdoba, Capilla de Montc. Distr. — Argentina.
- ?sp. James, 1953:317. Argentina, Tucumán, Tacanas.
- CUPRINA (Wiedemann), 1830:69 (*Stratiomys*). Type-locality: Brazil, Rio Grande do Sul, Caçapava do Sul. Distr. — Brazil.
- heteroneura* Macquart, 1838:190 (separate, p. 186) (*Odontomyia*). Type-locality: "Brazil".
- ssp. GORGONEA Lindner, 1949:863 (as form). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- VIRGATA Lindner, 1969:11. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus HEDRIODISCUS Enderlein

- HEDRIODISCUS Enderlein, 1914b:608. Type-species, *Odontomyia brevifacies* Macquart (orig. des.). Ref. — James (1953b) (key to Argentine species).
- Hedriodiscina* Enderlein, 1914b:609. Type-species, *Stratiomys chloraspis* Wiedemann (James, 1953:318).
- ADBITUS Lindner, 1949:820. Type-locality Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- BREVIFACIES (Macquart), 1855:63 (separate, p. 43) (*Odontomyia*). Type-locality: "South America". Distr. — Colombia and Guyana to Argentina (Jujuy, Santa Fe).
- CHLORASPIS (Wiedemann), 1830:68 (*Stratiomys*). Type-locality: "Brazil". Distr. — Bolivia to Uruguay and Argentina (south to Santa Fe).
- CURRANI (James), 1932:6 (as *Odontomyia truquii* var.). Type-locality: U.S.A., Arizona, Baboquivari Mts., Kits Peak, Rincon. Distr. — U.S.A. (Texas to California), Mexico (south to Michoacán).
- DORSALIS (Fabricius), 1805:82 (*Stratiomys*). Type-locality: West Indies. Distr. — U.S.A. (Texas, Florida); Mexico to Brazil (south to São Paulo); Cuba, Dominican Republic, Haiti, Puerto Rico, Jamaica.
- EUCHLORUS (Gerstaecker), 1857:328 (*Stratiomys*). Type-locality: "Mexico". Distr. — Mexico (Durango, Jalisco).
- affinis* Bellardi, 1859:35, 1861:235 (*Odonthomyia*). Type-locality: Mexico, Puebla.
- RUSTRIO (Walker), 1851:77 (*Odontomyia*). Type-locality: "Brazil". Distr. — Brazil.

- HUMILIS (Lindner), 1929:265 (*Hedriodiscina*). Type-locality: Brazil, Pernambuco, Recife. Distr. — Brazil.
- HYDROLUPUS (Lindner), 1949:859 (*Hedriodiscina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- INFRAPALLIDUS (Lindner), 1951:253 (*Hedriodiscina*). Type-locality: "Brazil". Distr. — Brazil.
- JACARELLUS (Lindner), 1949:853 (*Hedriodiscina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- LEFEBVREI (Macquart), 1838:193 (separate, p. 189) (*Odontomyia*). Type-locality: "Mexico". Distr. — Mexico (Durango to Michoacán and Puebla).
- lefeburei*, error or emendation.
- flavifasciatus* Macquart, 1849:357 (separate, p. 53) (*Odontomyia*). Type-locality: "Mexico".
- LINEATUS (Fabricius), 1805:83 (*Stratiomys*). Type-locality: "South America". Distr. — Peru. Ref. — Lindner, 1941 (generic reference).
- NUDIFRONS James, 1953b:321. Type-locality: Argentina, Jujuy, Palpalá. Distr. — Argentina.
- PALLIDIVENTRIS (Macquart), 1846:53 (separate, p. 181) (*Odontomyia*). Type-locality: "Brazil". Distr. — Brazil.
- PRASINA Jaenicke, 1867:324 (*Odontomyia*). Type-locality: "Mexico". Distr. — Mexico. Kertész considers this a syn. of *lefeburei*.
- PULCHER (Wiedemann), 1824:28 (*Stratiomys*). Type-locality: Brazil. Distr. — Costa Rica to Uruguay, and Argentina (south to Mendoza).
- buccphala* Gerstaecker, 1857:326 (*Odontomyia*). Type-locality: Uruguay, Montevideo.
- tremoleranus* Aldrich, 1928:1 (*Odontomyia*). Type-locality: Uruguay, Rio Negro, Menafra.
- QUADRILINEATUS (Macquart), 1834:248 (*Odontomyia*). Type-locality: "South America". Distr. — Colombia to Brazil (south to Paraná). Ref. — Gerstaecker, 1857 (suppl. descr.).
- sp. Lindner, 1929:266 (*Hedriodiscina*). Brazil, Pernambuco, Recife.
- sp. James, 1953b:321. Argentina. Presidente Perón.
- SUPERPICTUS (Lindner), 1949:857 (*Hedriodiscina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- TORTUGELLUS (Lindner), 1949:851 (*Hedriodiscina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- TRIVITTATUS (Say), 1829:160 (*Stratiomys*). Type-locality: "Mexico". Distr. — U.S.A. (widespread) to Colombia; Dominican Republic.
- tritaeniatus* Bellardi, 1859:38, 1861:238 (*Odontomyia*). Type-localities: Mexico, vicinity of Mexico City; Morelos, Cuantla.
- ssp. LEUCOGASTER James, 1933:69 (*Odontomyia*). Type-locality: U.S.A., Texas, Galveston. Distr. — U.S.A. (Arkansas to California) to Colombia.
- TRUQUH (Bellardi), 1859:34, 1861:234 (*Odonthomyia*). Type-locality: Mexico, Morelos, Cuernavaca.
- TURACELLUS (Lindner), 1949:855 (*Hedriodiscina*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



VERTEBRATUS (Say), 1824:369 (*Odontomyia*). Type-locality: U.S.A., Northwest Territories. Distr. — Canada, U.S.A., Mexico (Veracruz, Michoacán).

Genus HOPLITIMYIA James

HOPLITIMYIA James, 1934:443. Type-species, *Stratiomys constans* Loew (orig. des.). Ref. — James, 1939b (revision and key).

Subgenus HIMANTOMYIA James

Hoplitimyia, subg. HIMANTOMYIA James, 1939b:38. Type-species, *clavata* James (orig. des.).

CLAVATA James, 1939b:39. Type-locality: Ecuador, Baños. Distr. — Ecuador.

Subgenus HOPLITIMYIA James

ALEUS (Walker), 1849:526 (*Acrochacta*). Type-locality: "Brazil". Distr. — Brazil (São Paulo, Paraná).

BIMACULATA (Bellardi), 1862:10, 1864:209 (*Stratiomys*). Type-locality: Mexico, Yucatán, Cosamaloapan. Distr. — U.S.A. (Texas), Mexico (Nuevo Leon to Chiapas and Yucatán).

FASCIATA (Fabricius), 1787:331 (*Stratiomys*). Type-locality: "South America". Distr. — Mexico (north to Chihuahua and Taumalipas) to Brazil and Argentina (south to Buenos Aires); Trinidad.

subalba Walker, 1854:43 (*Stratiomys*). Type-locality: Brazil, Pará.

gerstaeckeri Bellardi, 1859:31, 1861:231 (*Stratiomys*). Type-locality: "Mexico"

MUTABILIS (Fabricius), 1787:331 (*Stratiomys*). Type-locality: "America". Distr. — U.S.A. (Texas, Arizona to Argentina, south to Buenos Aires).

variabilis Gmelin, 1792:2834 (*Musca*). Type-locality: French Guiana, Cayenne.

sericeiventris Rondani, 1848:102 (*Stratiomys*). Type-locality: "Brazil".

SEMILUNA James, 1939:41. Type-locality: Argentina, Mendoza. Distr. — Argentina (Tucumán, Catamarca, Mendoza, Cordoba).

Genus LABOSTIGMINA Enderlein

LABOSTIGMINA Enderlein, 1930:70. Type-species, *Odontomyia occipitalis* Johnson (orig. des.). Ref. — James, 1939b (part, *Udamacantha*).

Udamacantha Enderlein, 1914b:613. Type-species, *Stratiomys inermis* Wiedemann (orig. des.). N. SYN.

BONARIENSIS (Brèthes), 1922:139 (*Udamacantha*). Type-locality: Argentina, Buenos Aires, Mosconi. Distr. — Argentina.

CLYPEATA (Bigot), 1879:215 (*Odontomyia*). Type-locality: Brazil, Amazonia. Distr. — Brazil. Ref. — Lindner, 1930 (suppl. descr.).

COMMA (Williston), 1900:247 (?*Odontomyia*). Type-locality: Mexico, Guerrero, Dos Arroyos. Distr. — Mexico.

CONCINNATA (Williston), 1900:247 (*Odontomyia*). Type-locality: Mexico, Guerrero, Venta de Zapilote. Distr. — Mexico.

- ELEGANS (Macquart), 1838:191 (separate, p. 187) (*Odontomyia*). Type-locality: "Chile". Distr. — Chile (Atacama to Concepción); Argentina (Mendoza, Rio Negro).
- cruciata* Macquart, 1838:192 (separate, p. 188) (*Odontomyia*). Type-locality: "Chile".
- FENESTRATA (Thomson), 1869:458 (*Odontomyia*). Type-locality: Argentina, Buenos Aires. Distr. — Brazil (São Paulo), Argentina.
- FRATELLA (Williston), 1900:246 (*Odontomyia*). Type-locality: Mexico, Jalisco, Guadalajara. Distr. — Mexico.
- GUIANAE (Lindner), 1949:860 (*Eulalia*). Type-locality: Guyana, Georgetown. Distr. — Guyana.
- INERMIS (Wiedemann), 1830:71 (*Stratiomys*). Type-localities: Uruguay, Montevideo, and Brazil, Rio Grande do Sul, Caçapava do Sul. Distr. — Bolivia, Uruguay, Argentina (south to Buenos Aires).
- MACULIFRONS (Walker), 1849:536 (*Odontomyia*). Type-locality: "Honduras". Distr. — Mexico (north to Veracruz and Nayarit) to Honduras.
- PROTRUDENS (James), 1939b:45 (*Udamacantha*). Type-locality: Argentina, Córdoba. Distr. — Argentina.
- QUADRIMACULATA (Bellardi), 1859:37, 1861:237 (*Odonthomyia*). Type-locality: "Mexico". Distr. — Mexico (Durango to Michoacán and Puebla). Ref. — James, 1941b (suppl. descr.).
- femorata* Bellardi, 1859:37, 1861:237 (*Odonthomyia*). Type-locality: Mexico, Toluca.
- RUBRICORNIS (Macquart), 1846:181 (separate, p. 53) (*Odontomyia*). Type-locality: Mexico, Yucatán, Merida. Distr. — Mexico (Veracruz to Yucatán).
- VRIDIS (Bellardi), 1859:36, 1861:236 (*Odonthomyia*). Type-locality: Mexico, Morelos, Cuantla. Distr. — Mexico. Not *viridis* Bellardi of North American authors. Ref. — James, 1970 (suppl. descr.).

Genus METABASIS Walker

- METABASIS Walker, 1856:81. Type-species, *rostratus* Walker (mon.).
- ROSTRATUS Walker, 1856:81. Type-locality: "Brazil". Distr. — Brazil.

Genus ODONTOMYIA Meigen *

- Eulalia* Meigen, 1800:21. Type-species, *Musca hydroleon* Linnaeus (Coquillett, 1910:54). Suppressed by I.C.Z.N., 1963:339.
- ODONTOMYIA Meigen, 1803:265. Type-species, *Musca hydroleon* Linnaeus (Westwood, 1840:130). Ref. — James, 1957a (key to species of West Indies).
- Odonthomyia*, emendation.
- ALBOMACULATA Macquart, 1838:189 (separate, p. 193). Type-locality: Santo Domingo, Port au Prince. Distr. — Dominican Republic, Haiti.
- ALTIFRONS Wulp, 1888:363. Type-locality: Argentina, Córdoba. Distr. — Argentina.

* Undoubtedly some species referred here belong to other genera.



- ANCHORATA Bigot, 1879:216. Type-locality: "Chile". Distr. — Chile.
- ARANEIFERA Schiner, 1868:58. Type-locality: "Chile". Distr. — Chile.
- ATERRIMA Walker, 1856:78. Type-locality: "Brazil". Distr. — Brazil.
- BAHAMENSIS (James), 1953a:4 (*Eulalia*). Type-locality: Bahamas, South Bimini Island. Distr. — Bahama.
- DISCOLORATA James, 1936c:548. Type-locality: U.S.A., Texas, Donna. Distr. — U.S.A. (Texas), Mexico (Sonora and Tamaulipas to Yucatán), ?Costa Rica.
- DISSIMILIS Bellardi, 1859:35, 1861:235. Type-locality: Mexico, "contorni di Messico". Distr. — Mexico.
- EMARGINATA Macquart, 1838:194 (separate, p. 190). Type-locality: "Mexico". Distr. — Mexico.
- FASCIATA Macquart, 1834:249. Type-locality: "South America". Distr. — South America.
- FASCIFRONS Macquart, 1849:354 (separate, p. 50). Type-locality: "Chile".
- FOVEIFRONS Thomson, 1869:458. Type-locality: Brazil, Guanabara, Rio de Janeiro. Distr. — Brazil.
- ICAE (Lindner), 1941:180 (*Eulalia*). Type-locality: Peru, Ica, Victoria. Distr. — Peru. Ref. — Lindner, 1951 (suppl. descr.).
- INAEQUALIS Loew, 1866a:143 (Centuria 6, no. 24). Type-locality: Canada, Hudson's Bay Territory. Distr. — Canada, U.S.A., Mexico (south to Morcos).
- confusa* James, 1936c:544. Type-locality: U.S.A., Colorado, Alamosa.
- LIMAE Guérin-Ménéville, 1831:291. Type-locality: Peru, Lima. Distr. — Peru.
- MEXICANA Johnson, 1895:271. Type-locality: Mexico, Tehuantepec. Distr. — Mexico.
- NITIDICEPS Wulp, 1888:365. Type-locality: Argentina, Córdoba. Distr. — Argentina.
- PACHYCEPHALA Schiner, 1868:60. Type-locality: "Colombia". Distr. — Colombia.
- PACHYCEPS Bigot, 1879:215. Type-locality: Brazil, Pará. Distr. — Brazil.
- PERUVIANA Macquart, 1855:63 (separate, p. 43). Type-locality: "Peru". Distr. — Peru.
- RUFIPES Loew, 1866a:144 (Centuria 6, no. 25). Type-locality: "Cuba". Distr. — U.S.A. (Florida), Cuba, Bahamas.
- SPP. Williston, 1888:256. No locality given.
- SUBCUPRATA (Walker), 1854:47 (*Stratiomys*). Type-locality: Brazil, Pará, Santarém. Distr. — Brazil.
- VANDERWULPIANA Brèthes, 1907:283 (nom. nov. for *maculifrons* Wulp). Type-locality: Argentina, Córdoba.
- maculifrons* Wulp, 1888:364 (preocc. Walker, 1849). Type-locality: Argentina, Córdoba.
- VICINA Macquart, 1834:192 (separate, p. 188). Type-locality: "Cuba". Distr. — Cuba.
- VITTATA Macquart, 1850:357 (separate, p. 53). Type-locality: "Brazil". Distr. — Brazil.



Genus PROMERANISA Walker

- PROMERANISA Walker, 1854:305. Type-species, *vittata* Walker (mon.).
CYLINDRICORNIS Williston, 1888:253. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil (Mato Grosso) to Argentina.
NASUTA (Macquart), 1849:351 (separate, p. 47) (*Stratiomyia*). Type-locality: "Chile". Distr. — Costa Rica to Chile and Argentina. Ref. — Williston, 1888 (suppl. descr.).
VITTATA Walker, 1856:306. Type-locality: Brazil, Pará, Santarém.

Genus RHINGIOPSIS Röder

- RHINGIOPSIS Röder, 1886:138. Type-species, *tau* Röder (mon.). Ref. — Curran, 1932a (key).
ENDERLEINI Lindner, 1926:100 (adult, larva). Type-locality: "North Argentina". Distr. — Argentina.
JAMESI Barretto, 1947:439, fig. 1. Type-locality: Brazil, São Paulo, Avanhandava, Barra Mansa. Distr. — Brazil (São Paulo).
LANEI Barretto, 1947:441, figs. 2-3. Type-locality: Brazil, São Paulo, São Paulo, Serra da Cantarcira, Chapadão. Distr. — Brazil (São Paulo).
NASUTA Enderlein, 1914b:605. Type-locality: Brazil, Mato Grosso, Cuiabá. Distr. — Brazil.
ROSTRATA (Wiedemann), 1830:68 (*Stratiomys*). Type-locality: "Brazil". Distr. — Brazil (Mato Grosso to Santa Catarina); Paraguay.
TAU Röder, 1886:138. Type-locality: "Brazil". Distr. — French Guyana to Brazil (south to Guanabara).

Genus STRATIOMYELLA James

- STRATIOMYELLA James, 1953b:321. Type-species, *nana* James (orig. des.).
NANA James, 1953b:322. Type-locality: Argentina, Tucumán, Concepción. Distr. — Argentina (Jujuy, Tucumán, Presidente Perón, Santa Fe).

Genus STRATIOMYS Geoffroy *

- STRATIOMYS Geoffroy, 1762:475. Type-species, *Musca chameleon* Linnaeus (Latreille, 1810). Conserved as of 1762 under suspension of the Rules (I.C.Z.N., 1957b:123).
Stratiomyia, emend.
CONICA Fabricius, 1805:81. Type-locality: "South America". Distr. — South America.
CONSTRACTA Walker, 1860:269. Type-locality: "Mexico". Distr. — Mexico.
pinguis Walker, 1860:270. Type-locality: "Mexico".
CONVEXA Wulp, 1881:154. Type-locality: "Argentina". Distr. — Mexico (north to Veracruz), Guatemala, Bolivia, Paraguay, Argentina (Salta to Buenos Aires). Ref. — Lindner, 1926 (larva).
connexa, error.

* Undoubtedly some species referred here belong to other genera.



- FENESTRATA Gerstaecker, 1857:327. Type-locality: "Mexico". Distr. — Mexico.
- GONIOPHORA Say, 1829:161. Type-locality: "Mexico". Distr. — Mexico.
- LEUCOPSIS Wiedemann, 1830:74. Type-locality: Brazil, Rio Grande do Sul, Caçapava do Sul. Distr. — Brazil.
- LONGIFRONS Rondani, 1848:101. Type-locality: "Brazil". Distr. — Brazil.
- MELANOPSIS Wiedemann, 1830:70. Type-locality: Brazil, Rio Grande do Sul, Caçapava do Sul. Distr. — Brazil.
- FALLIPES Fabricius, 1781:147. Type-locality: "America, probably South America". (Given as "in America boreali" in Fabricius, 1781, but as "Südamerika" in Fabricius, 1805). Distr. — ?South America.
- PELLUCIDA Rondani, 1848:102. Type-locality: "Brazil". Distr. — Brazil.
- PRINCEPS Gerstaecker, 1857:321. Type-locality: "Mexico". Distr. — Mexico.
- VELUTINA Bigot, 1879:213. Type-locality: "Chile". Distr. — Chile.
- VIRENS Wiedemann, 1830:73. Type-locality: "Brazil". Distr. — Brazil.

Genus ZUERCHERIA Lindner

- ZUERCHERIA Lindner, 1926:102. Type-species, *malachitis* Lindner (mon.). Ref. — James, 1938 (key).
- BEQUAERTI (Curran), 1932a:1 (*Rhingiopsis*). Type-locality: Mexico, Yucatán, Chichén Itzá. Distr. — Mexico.
- FESTIVA (Walker), 1854:43, 63 (*Stratiomys*). Type-locality: Brazil, Pará. Distr. — Guyana, Brazil (Amazon basin). Ref. — Lindner, 1965 (suppl. descr., larva).
- stratiomyaeformis* Lindner, 1935:408. Type-locality: Guyana, Issororo.
- MALACHITIS Lindner, 1926:102. Type-locality: "Presumably Paraguay". Distr. — Bolivia, Paraguay. Ref. — James, 1938:200 (suppl. descr.).
- FUSTULOSA James, 1938:201. Type-locality: Colombia, Aracataca. Distr. — Colombia.

Tribe MYXOSARGINI

Ref. — James, 1942 (revision).

Genus MELANOCHROA Röder

- Melanochroa* Brauer, 1882:83, 88. Nomen nudum.
- MELANOCHROA Röder, 1886:139. Type-species, *dubia* Röder (mon.).
- DUBIA Röder, 1886:139. Type-locality: "Brazil" Distr. — Brazil (Guanabara). *dubia* "Schiner", Brauer, 1882:88. Nomen nudum.

Genus MYXOSARGUS Brauer

- MYXOSARGUS Brauer, 1882:77, 88. Type-species, *fasciatus* Brauer (orig. des.).
- BRAUERI Williston, 1888:254. Type-locality: Brazil, Mato Grosso, Chapada. Distr. — Brazil.
- FASCIATUS Brauer, 1882:78. Type-locality: Mexico, Veracruz, Orizaba. Distr. — Mexico (records from U.S.A. erroneous).



- GYPHIUS Enderlein, 1914b:612. Type-locality: Mexico, Chiapas. Distr. — Mexico.
- PANAMENSIS Curran, 1929:4. Type-locality: Canal Zone (Barro Colorado Island). Distr. — Costa Rica to Panama.
- PILOSUS James, 1942:59. Type-locality: U.S.A., Arizona, Tempe. Distr. — U.S.A. (Ariz.) to Canal Zone.
- ssp. QUATEMALAE James, 1942:60. Type-locality: Guatemala, Amatitlán. Distr. — Guatemala.
- SCUTELLATUS Williston, 1900:251. Type-locality: Mexico, Tabasco, Teapa. Distr. — Mexico.

Genus NOTHOMYIA Loew

- NOTHOMYIA Loew, 1869:4 (Centuria 8, no. 4, note). Type-species, *scutellata* Loew (Brauer, 1882:88; 1882:32).
- Berisargus* Lindner, 1933a:201. Type-species, *borgmeieri* Lindner (mon.).
- BORGMEIERI (Lindner), 1933a:202 (*Berisargus*). Type-locality: Brazil, São Paulo, Alto da Serra (= Paranapiacaba). Distr. — Brazil (Guarnabara, São Paulo).
- CALOPUS Loew, 1869:5 (Centuria 8, no. 5). Type-locality: "Cuba". Distr. — U.S.A. (Florida), Cuba, Puerto Rico.
- FASCIATIPENNIS (Lindner), 1935:401 (*Berisargus*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Juçaraí.
- INTENSICA (Curran), 1928a:18 (*Merosargus*). Type-locality: Jamaica, Cinchona. Distr. — Jamaica.
- LONGISETOSA (Lindner), 1933b:328 (*Berisargus*). Type-locality: Brazil, Rio de Janeiro, Itatiaia. Distr. — Brazil.
- LOPESI (Lindner), 1935:401 (*Berisargus*). Type-locality: Brazil, Rio de Janeiro, Angra dos Reis, Juçaraí. Distr. — Brazil (Rio de Janeiro to Santa Catarina).
- METALLICA (Wiedemann), 1830:60 (*Oxycera*). Type-locality: West Indies (St. Thomas). Loew is probably correct in suggesting that this species may be a *Nothomyia*.
- NIGRA James, 1942:55. Type-locality: Dominican Republic, Valle Nuevo, S.E. of Constanza. Distr. — Dominican Republic.
- PARVICORNIS James, 1939a:598. Type-locality: Costa Rica, San Carlos. Distr. — Costa Rica to Panama.
- SCUTELLATA Loew, 1869:4 (Centuria 8, no. 4). Type-locality: "Cuba". Distr. — Cuba.
- VIRIDIS Hine, 1911:301. Type-locality: U.S.A., Ohio, Sandusky, Cedar Point (probably adventitious). Distr. — U.S.A. (Ohio); Puerto Rico.

Tribe NEMOTELINI

Genus NEMOTELUS Geoffroy

- NEMOTELUS Geoffroy, 1762:450, 542. Type-species, *Musca pantherina* Linnaeus (I.C.Z.N., 1957a:85). Conserved as of 1762 under Suspension of the Rules (I.C.Z.N., 1957a:85). Refs. — Kertész, 1914 (rev. South American species); James, 1941b (rev. Mexican species).
- Nematotelus*, emend.



- ACUTIROSTRIS Loew, 1863:8 (Centuria 3, no. 13). Type-locality: "Cuba".
Distr. — Cuba, Bahamas, ?Galapagos Is.
- ALBIBROSTRIS Macquart, 1850:359 (separate, p. 55). Type-locality: U.S.A.,
Virginia. Distr. — U.S.A. (Virginia to Arizona), Bahama Islands.
whchelevi Mclander, 1903:182. Type-locality: U.S.A., Texas.
- ALBIVENTRIS Thomson, 1869:462. Type-locality: Galápagos Ids., erroneously
given as Manilla (sic!). Distr. — Galápagos Ids.
- BASALIS Kertész, 1914:528. Type-locality: Peru, Lima, Baranca. Distr. —
Peru.
- CATHARISTIS Lindner, 1930:276. Type-locality: Argentina, Corrientes, San
José. Distr. — Argentina.
catharistis Lindner, 1926:97 (nomen nudum).
- CONGRUENS Kertész, 1914:528. Type-locality: Chile, Arica. Distr. — Peru,
Chile. Ref. — Lindner, 1941 (suppl. descr.).
- DAMPFI James, 1941b:243. Type-locality: Mexico, Texcaltenco. Distr. —
Mexico.
- DIEHLI Lindner, 1941:184. Type-locality: Peru, Cayara. Distr. — Peru.
- DISJUNCTUS Curran, 1925:244. Type-locality: Peru, Lima.
- FLAVICORNIS Johnson, 1894:272. Type-locality: Jamaica, Kingston. Distr.
— U.S.A. (Florida), Bahamas, Haiti, Jamaica.
- FLAVOCINGULATUS Kertész, 1914:522. Type-locality: Argentina, Neuquén.
Distr. — Argentina (Cordoba to Rio Negro).
- GLABER Loew, 1872:56 (Centuria 10, no. 10). Type-locality: U.S.A., Texas.
Distr. — U.S.A. to Panama.
- KANSSENSIS TRINOTATUS Melander, 1903:180 (as species). Type-locality:
U.S.A., Texas, Austin. Distr. — U.S.A. (Oklahoma, Texas),
Mexico (south to Morelos).
- MACULATUS Kertész, 1914:521. Type-locality: Argentina, Neuquén, Piedra
Pinta. Distr. — Argentina (Salta to Neuquén).
- MONENSIS Curran, 1928a:16. Type-locality: Puerto Rico, Mona Island.
Distr. — Puerto Rico.
- NEOTROPICUS Curran, 1925:243. Type-locality: Peru, Lima.
- NEUQUENENSIS Kertész, 1914:518. Type-locality: Argentina, Neuquén. Distr.
— Argentina (Salta to Neuquén).
- NIGER Bigot, 1879:233. Type-locality: "Chile". Distr. — Chile (Aconcagua);
Paraguay; Argentina south to Rio Negro.
fasciatifrons Lynch Arribálzaga, 1881:190. Type-locality: Argentina,
Rio Negro.
- NIGRICORNIS Kertész, 1914:527. Type-locality: Chile, Arica. Distr. — Chile.
- NIGRINUS Fallén, 1817:6. Type-locality: "Sweden". Distr. — Europe;
Alaska to Quebec, south to Hidalgo, Mexico. South American
records of this species, as *unicolor* Loew, are probably erroneous.
- NUDUS Kertész, 1914:525. Type-locality: Colombia, Bolívar, Baranquilla.
Distr. — Colombia to Brazil (Pernambuco).
- PERUVIANUS Kertész, 1914:519. Type-locality: Peru, Cuzco. Distr. — Peru.
- POLYPOSUS Say, 1829:160. Type-locality: "Mexico". Distr. — Mexico
(Veracruz).

- ROEDINGERI Lindner, 1941:183. Type-locality: Peru, Ica, near Victoria. Distr. — Peru.
- RUFICORNIS Bigot, 1879:234. Type-locality: "Chile". Distr. — Chile.
- RUFODOMINIALIS Cole, 1923:459. Type-locality: Mexico, Sonora, Tepoca Bay Distr. — Southwestern U.S.A. (New Mexico to California), Mexico (Sonora, Baja California). Ref. — James, 1936b (suppl. descr.).
- RUGOSUS James, 1941b:243. Type-locality: Mexico, Mexico, Llanos de Salazar. Distr. — Mexico.
- SINGULARIS Macquart, 1846:185 (separate, p. 57). Type-locality: "Colombia". Distr. — Colombia.
- SORDIDUS Kertész, 1914:524. Type-locality: Peru, Cuzco. Distr. — Peru.
- SP. James, 1967a:7. Dominica.
- TENUISTYLUS Hanson, 1958:1379. Type-locality: U.S.A., New Mexico, Las Cruces. Distr. — Western U.S.A.; Mexico (Baja California).
- THOMAE Curran, 1928a:15. Type-locality: Virgin Islands, St. Thomas. Distr. — Virgin Islands.
- TITSCHACKI Lindner, 1941:182. Type-locality: Peru, Sivia. Distr. — Peru.
- VARIABLES Hanson, 1958:1368. Type-locality: U.S.A., Florida, Hibernia. Distr. — southern U.S.A. (Florida to California); northern Mexico.

Subfamily PACHYGASTRINAE

Refs. — Kertész, 1916 (key to genera of World); Lindner, 1964 (key to Neotropical genera); James, 1967e (key to Argentine genera).

Genus ACANTHINOMYIA Hunter

- Acanthina* Wiedemann, 1830:50 (preoce. Fischer, 1807). Type-species, *elongata* Wiedemann (Brauer, 1882:86).
- ACANTHINOMYIA Hunter, 1901:129 (nom. nov. for *Acanthina* Wiedemann). Type-species, *Acanthina elongata* Wiedemann (aut.). Ref. — Kertész, 1914 (revision, key).
- Acanthinia*, error.
- BEQUAERTI Curran, 1925:338. Type-locality: Brazil, Amazonas, Rio Branco, Vista Alegre. Distr. — Brazil (Amazonas, Pará).
- CENTOR Curran, 1934:313. Type-locality: Guyana, Bartica, Kartabo. Distr. — Guyana.
- ELONGATA (Wiedemann), 1824:30 (*Clitellaria*). Type-locality: "South America". Distr. — Costa Rica to Guyana, Peru, and Brazil (south to São Paulo); Trinidad.
- LONGA (Wiedemann), 1830:51 (*Acanthina*). Type-locality: "Brazil". Distr. — Brazil (Rio de Janeiro to Santa Catarina); Argentina (Misiones).
- METALLICA Curran, 1934:312. Type-locality: Guyana, Bartica, Kartabo. Distr. — Guyana.
- PIANA (Walker), 1854:305 (*Acanthina*). Type-locality: Brazil, Pará, Santarém. Distr. — Brazil. (Generic position uncertain).



PULCHELLA James, 1936a:54. Type-locality: Mexico, Veracruz, Córdoba.
Distr. — Mexico.

Genus ARTEMITA Walker

ARTEMITA Walker, 1854:61. Type-species, *Clitellaria amenides* Walker (Brauer, 1882:). Refs. — Kertész, 1914 (revision); James, 1971 (South American species).

Hypocerina Bigot, 1879:207. (Nomen nudum).

Pseggomma Enderlein, 1914a:304. Type-species, *Artemita argentea* Walker (orig. des.).

AMENIDES (Walker), 1849:523 (*Clitellaria*). Type-locality: Not given. Distr. — Canal Zone to Guyana, Bolivia, and Brazil (south to Santa Catarina).

ARGENTEA Osten Sacken, 1886:41. Type-localities: "Costa Rica, Caché, and Colombia, no locality". Distr. — Mexico to Peru.

aurcostrata Enderlein, 1914a:304 (*Pseggomma*). Type-locality: "Costa Rica".

AURATA Macquart, 1846:179 (separate, p. 51). Type-locality: "Colombia". Distr. — Panama, Colombia, Ecuador, Peru.

aurata Bigot, 1879:207. Type-locality: "Colombia".

BANKSI James, 1971:66. Type-locality: Canal Zone, Barro Colorado Island. Distr. — Canal Zone.

BELLARDI Giglio-Tos, 1891:2, 1892:12. Type-locality: Mexico, Tchuacán. Distr. — Mexico.

BICOLOR Kertész, 1914:482. Type-localities: "Bolivia, Coroico, and Brazil, Espírito Santo". Distr. — Panama, Canal Zone, Surinam, Bolivia, Brazil (south to Santa Catarina), Argentina (Misiones); Trinidad.

BRASILIANA Lindner, 1964:10. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

CONVEXA (Walker), 1854:304 (*Acanthina*). Type-locality: Brazil, Pará, Santarém. Distr. — Costa Rica to Peru and Brazil (south to Santa Catarina).

porcata Walker, 1857:120 (*Acanthina*). Type-locality: "Amazon Region".

HEROGLYPHICA (Wiedemann), 1830:50 (*Acanthina*). Type-locality: "Brazil". Distr. — Mexico (north to Puebla) to Brazil (south to Santa Catarina).

halala Walker, 1849:523 (*Clitellaria*). Type-locality: "Honduras".

ornata Macquart, 1846:179 (separate, p. 51) (*Acanthina*). Type-locality: "Colombia".

marmorata Gerstaecker, 1857:37 (*Acanthina*). Type-locality: "Brazil".

LATIFRONS James, 1971:62. Type-locality: Peru, Madre de Dios, Avispas. Distr. — Ecuador, Peru.

NANA (Bellardi), 1862:9, 1864:208 (*Acanthina*). Type-locality: Mexico, Veracruz, Tuxpango near Orizaba. Distr. — U.S.A. (Texas), Mexico (Tamaulipas to Nayarit and Oaxaca), Dominica. Ref. — James, 1967a (as *Artemita inornata*, suppl. descr.).

inornata Williston, 1900:249 (*Acanthina*). Type-locality: Mexico, Veracruz, Atoyac.

- PERUVIANA Kertész, 1914:486. Type-locality: Peru, Callanga. Distr. — Peru.
PODEXARGENTEUS Enderlein, 1914a:304. Type-locality: Brazil, Santa Catarina.
Distr. — Brazil (São Paulo, Santa Catarina), Paraguay.

Genus BLASTOCERA Gerstaecker

- BLASTOCERA Gerstaecker, 1857:341. Type-species, *speciosa* Gerstaecker
(mon.). Ref. — Kertész, 1908 (suppl. descr.).
SPECIOSA Gerstaecker, 1857:342. Type-locality: "Brazil". Distr. — Canal
Zone to Argentina and Brazil (south to Santa Catarina).
atra Wandolleck, 1897:216. Type-locality: St. Cruz.

Genus BORBORIDEA Kertész

- BORBORIDEA Kertész, 1916:155. Type-species, *megaspis* Kertész (orig. des.).
MEGASPIS Kertész, 1916:157. Type-locality: Brazil, Solidade (?). Distr. —
Brazil (Santa Catarina), Argentina (La Rioja). Ref. — Lindner,
1949 (suppl. descr.).
MICROCEPHALUM (Kraft & Cook), 1961:10 (*Zabrachia*). Type-locality: U.S.A.,
Texas, Marathon. Distr. — U.S.A. (Texas), Mexico (Sonora,
Vcracruz).

Genus BRACHYODINA Lindner

- BRACHYODINA Lindner, 1949:878-879. Type-species, *niveioscula* Lindner
(mon.).
NIVEIOSCULA Lindner, 1949:879. Type-locality: Guyana, Mazaruni. Distr. —
Guyana.

Genus CAENACANTHA Wulp

- CAENACANTHA Wulp, 1885:58. Type-species, *bipartita* Wulp (?mon.).
BIPARTITA Wulp, 1885:59. Type-locality: "Colombia". Distr. — Colombia.

Genus CACTOBIA James

- CACTOBIA James, 1966c:109. Type-species, *opuntiae* James (orig. des.).
OPUNTIAE James, 1966c:111. Type-locality: Mexico, Guanajuato, San Miguel
de Allende. Distr. — Mexico (Guanajuato, Aguascalientes).

Genus CHALCIDOMORPHINA Enderlein

- CHALCIDOMORPHINA Enderlein, 1914a:298. Type-species, *aurata* Enderlein
(orig. des.).
Chalcomorphina, error.
AURATA Enderlein, 1914a:299. Type-locality: Colombia, Hacienda Pehlke.
Distr. — Mexico (Morelos), Colombia, Peru. Ref. — Lindner,
1951 (suppl. descr., fig.).
PLANES James, 1967a:20 (*Chalcomorphina*). Type-locality: Dominica, Clarke
Hall. Distr. — Dominica.



Genus CHARISINA Lindner

- CHARISINA Lindner, 1951:262. Type-species, *angustifrons* Lindner (mon.).
Ref. — Lindner, 1964 (redescribed as new).
ANGUSTIFRONS Lindner, 1951:262. Type-locality: Brazil, Santa Catarina,
Nova Teutônia. Distr. — Brazil. Ref. — Lindner, 1964 (redescribed
as new).

Genus CHLAMYDONOTUM Lindner

- CHLAMYDONOTUM Lindner, 1949:882. Type-species, *nigreradiatum* Lindner
(mon.). Ref. — James, 1967c (key).
DENTATUM James, 1967e:111. Type-locality: Argentina, Tueumán,
Burruyacu, Villa Padre Monti. Distr. — Argentina.
NIGRERADIATUM Lindner, 1949:883. Type-locality: Brazil, Santa Catarina,
Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina
(Tueumán).
PALLIPES Lindner, 1951:261. Type-locality: Brazil, Santa Catarina, Nova
Teutônia. Distr. — Brazil. Generic reference uncertain; possibly
Praomyia.

Genus CHOROPHTHALMYIA Lindner

- CHOROPHTHALMYIA Lindner, 1964:6. Type-species, *brevicornis* Lindner
(mon.).
Chorophthalmus Lindner, 1964:18 (lapsus; Lindner, 1969:5).
BREVEORNIS Lindner, 1964:6, 18. Type-locality: Brazil, Santa Catarina, Nova
Teutônia. Distr. — Brazil.

Genus CLARISSA Lindner

- CLARISSA Lindner, 1964:7, 19. Type-species, *pallipes* Lindner (mon.).
PALLIPES Lindner, 1964:19. Type-locality: Brazil, Santa Catarina, Nova
Teutônia. Distr. — Brazil.

Genus COSMARIOMYIA Kertész

- COSMARIOMYIA Kertész, 1914:552. Type-species, *argyrosticta* Kertész (orig.
des.).
Lophoteles Authors, not Loew.
ARGYROSTICTA Kertész, 1914:554. Type-locality: Brazil, São Paulo. Distr. —
Brazil (São Paulo, Santa Catarina), Argentina (Salta, Tueumán).
PALLIDIPENNIS (Williston), 1900:250 (*Lophoteles*). Type-locality: Mexico,
Veraacruz, Atoyac. Distr. — U.S.A. (?Ohio), Mexico.
plumula Loew of Enderlein, 1914a:310 (*Lophoteles*), misident.

Genus CYCLOTASPIS Lindner

- CYCLOTASPIS Lindner, 1964:5, 20. Type-species, *inornata* Lindner (mon.).
INORNATA Lindner, 1964:20. Type-locality: Brazil, Santa Catarina, Nova
Teutônia. Distr. — Brazil.



Genus CYNIPIMORPHA Brauer

CYNIPIMORPHA Brauer, 1882:75, 87. Type-species, *billmecki* Brauer (as *billmecki*) (mon.). Ref. — Kertész, 1908 (suppl. descr.).

Cynipomorpha, emendation.

BILIMEKI Brauer, 1882:75. Type-localities: Mexico, Veracruz, Orizaba, and Morelos, Cuernavaca (as *billmecki*, lapsus). Distr. — Mexico (San Luis Potosí, Veracruz, Morelos).

Genus DACTYLACANTHA Lindner

DACTYLACANTHA Lindner, 1964:6, 12. Type-species, *plumanni* Lindner (mon.).

PLAUMANNI Lindner, 1964:12. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus DACTYLODEICTES Kertész

DACTYLODEICTES Kertész, 1914:544. Type-species, *amazonicus* Kertész (orig. des.).

AMAZONICUS Kertész, 1914:545. Type-locality: "Brazil", no locality. Distr. — Brazil.

DEPRESSUS James, 1967a:16. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.

INSULARIS James, 1966a:539. Type-locality: Galápagos Islands, Santa Cruz Island, Academy Bay. Distr. — Galápagos Islands.

LOPESI Lindner, 1964:5, 15. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Canal Zone, Trinidad, Brazil.

Genus DIASTROPHTHALMUS Lindner

DIASTROPHTHALMUS Lindner, 1949:885. Type-species, *flavimannus* Lindner (mon.).

FLAVIMANUS Lindner, 1949:886. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus ECCHAETOMYIA Lindner

ECCHAETOMYIA Lindner, 1949:887. Type-species, *nigrovittata* Lindner (mon.).


NIGROVITTATA Lindner, 1949:888. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil (Santa Catarina), Argentina (Tucumán).

Genus EICOCHALCIDINA Lindner

EICOCHALCIDINA Lindner, 1964:6, 14. Type-species, *marginalis* Lindner (mon.).

GOLBACHI James, 1967c:116. Type-locality: Argentina, Tucumán, Burreyacu, Villa Padre Monti. Distr. — Argentina.

MARGINALIS Lindner, 1964:14. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.



Genus EIDALIMUS Kertész

- EIDALIMUS Kertész, 1914:546. Type-species, *annulatus* Kertész (orig. des.).
Pyelomyia Kertész, 1916:189. Type-species, *annulata* Kertész (orig. des.).
ANGUSTIFRONS James, 1967c:101. Type-locality: Argentina, Tucumán, Quebrada la Toma. Distr. — Argentina.
ANNULATUS Kertész, 1914:547. Type-locality: Peru, Urubamba River, Umuhuankiali. Distr. — Peru.
annulatus Kertész, 1916:190 (*Pyelomyia*). Type-localities: Peru, Callanga and Vilcanota.
FLAVICORNIS Lindner, 1969:4. Type-locality: ?Brazil, ?Santa Catarina, Caraguatá River. Distr. — ?Brazil.
HENSHAWI (Malloch), 1917:338 (*Eupachygaster*). Type-locality: U.S.A., Illinois, Savoy. Distr. — U.S.A., Cuba.

Genus ENGICERUS Lindner

- ENGICERUS Lindner, 1964:7, 16. Type-species, *major* Lindner (mon.).
MAJOR Lindner, 1964:16. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus EUCYNIPIMORPHA Malloch

- EUCYNIPIMORPHA Malloch, 1915:312. Type-species, *Cynipimorpha minuta* Williston (orig. des.).
MINUTA (Williston), 1900:252 (*Cynipimorpha*). Type-locality: Mexico, Guerrero, Chilpancingo. Distr. — Mexico (Sonora, Guerrero) to Canal Zone.

Genus EUPACHYGASTER Kertész

- EUPACHYGASTER Kertész, 1911:31. Type-species, *Pachygaster tarsalis* Zetterstedt (orig. des.).
ALEXANDERI (Brèthes), 1922:140 (*Vappo*). Type-locality: Argentina. Distr. — Argentina (Tucumán, Mendoza). Rcf. — Blanchard, 1923 (suppl. descr., larva); James, 1967c (suppl. descr.).
VILLOSA James, 1939a:606. Type-locality: Costa Rica, San Mateo, Higuito. Distr. — Costa Rica. (Generic position uncertain).

Genus GOWDEYANA Curran

- GOWDEYANA Curran, 1928b:15. Type-species, *mirabilis* Curran (mon.).
MIRABILIS Curran, 1928b:15. Type-locality: Jamaica, Hope Gardens. Distr. — Jamaica.

Genus *Hexacraspis* Enderlein

- Hexacraspis* Enderlein, 1914a:295. Type-species, *Diphysa sexspinoso* Macquart (orig. des.).
sexspinoso (Macquart), 1846:174 (separate, p. 46) (*Diphysa*). Type-locality: "Brazil"; in error; a South African species, Not Neotropical.

Genus HYPSELOPHRUM Kertész

HYPSELOPHRUM Kertész, 1909:381. Type-species, *cyphomyioides* Kertész (mon.).

CYPHOMYIOIDES Kertész, 1909:382. Type-locality: Peru, Callanga. Distr. — Peru.

Genus LASIODEICTES Lindner

LASIODEICTES Lindner, 1964:7, 17. Type-species, *niger* Lindner (mon.).

NIGER Lindner, 1964:17. Type-localities: Brazil, Santa Catarina, Rio Caraguatá and Nova Teutônia. Distr. — Brazil.

Genus LIGYROMYIA Kertész

LIGYROMYIA Kertész, 1916:200. Type-species, *columbiana* Kertész (orig. des.).

COLOMBIANA Kertész, 1916:201. Type-locality: Colombia, Sierra S. Lorenzo. Distr. — Colombia to Brazil (south to Santa Catarina) and Argentina (Tucumán); Trinidad. Ref. — Lindner, 1933a (suppl. descr.).

Genus LYPROTEMYIA Kertész

LYPROTEMYIA Kertész, 1909:391. Type-species, *formicaeformis* Kertész (mon.).

FORMICAEFORMIS Kertész, 1909:391. Type-locality: Peru, Meshagua, Urubamba River. Distr. — Peru.

Genus MANOTES Kertész

MANOTES Kertész, 1916:180. Type-species, *plana* Kertész (orig. des.). Ref. — James, 1967c (key).

FLAVIPES James, 1967c:105. Type-locality: Argentina, Formosa, Pirané. Distr. — Argentina.

HYALINA James, 1967c:108. Type-locality: Argentina, Tucumán, Burrayacu, Villa Padre Montl. Distr. — Argentina (Salta, Tucumán).

LATIMANUS James, 1967c:106. Type-locality: Mexico, Puebla, Acatlán. Distr. — Mexico.

PLANA Kertész, 1916:181. Type-locality: Paraguay, Asunción. Distr. — Mexico (Morelos), Bolivia, Brazil (Santa Catarina), Paraguay, Argentina (Tucumán). Ref. — James, 1967c (suppl. descr.).

nigrobasis Lindner, 1949:886 (*Borboridea*). Type-locality: Brazil, Santa Catarina, Nova Teutônia.

Genus MERISTOCERA Lindner

MERISTOCERA Lindner, 1964:13. Type-species, *aurca* Lindner (mon.).

AUREA Lindner, 1964:13. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

aurata, error.



LATICORNIS James, 1967a:15. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.

Genus MYIOCAVIA Lindner

MYIOCAVIA Lindner, 1949:883-884. Type-species, *tomentosa* Lindner (mon.).

TOMENTOSA Lindner, 1949:884. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus MYCTEROCERA James

MYCTEROCERA James, 1967c:117. Type-species, *contigua* James (orig. des.).

CONTIGUA James, 1967c:118. Type-locality: Argentina, Tucumán, Burreyacu, Villa Padre Monti. Distr. — Argentina.

Genus NEOCANTHINA Kertész

NEOCANTHINA Kertész, 1914:477. Type-species, *fasciata* Kertész (orig. des.).

FASCIATA Kertész, 1914:479. Type-locality: Brazil, Amazon. Distr. — Brazil.

Genus NEOCHAUNA Williston

Chauna Loew, 1847:370 (preocc. Illiger, 1811). Type-species, *variabilis* Loew, 1847:370 (mon.).

NEOCHAUNA Williston, 1896:48 (nom. nov. for *Chauna* Loew). Type-species, *Chauna variabilis* Loew (aut.).

VARIABILIS (Loew), 1847:370 (*Chauna*). Type-locality: "Cuba". Distr. — Mexico (Chiapas) to Brazil (Mato Grosso); Cuba.

ferruginca Gerstaecker, 1857:340 (*Chauna*). Type-locality: "Cuba".

Genus NETROGRAMMA Lindner

NETROGRAMMA Lindner, 1964:5, 17. Type-species, *consona* Lindner (mon.).

CONSONA Lindner, 1964:17. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus PACHYGASTER Meigen

PACHYGASTER Meigen, 1803:266. Type-species, *Nemotelus ater* Panzer (orig. des.).

WIRTHI James, 1967a:18. Type-locality: Dominica, Hillsborough Estate. Distr. — Dominica, Antigua (Generic position uncertain).

Genus PANACRIS Gerstaecker

PANACRIS Gerstaecker, 1857:346. Type-species, *lucida* Gerstaecker (mon.). Ref. — Kertész, 1908 (key, revision).

BREVISETA Lindner, 1964:8. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

LUCIDA Gerstaecker, 1857:347. Type-locality: French Guiana, Cayenne. Distr. — French Guiana, Brazil (Mato Grosso, Santa Catarina).

- MAXIMA Kertész, 1908:367. Type-localities: Brazil, Espírito Santo and Santa Catarina. Distr. — Brazil (Minas Gerais, Espírito Santo, Santa Catarina). Ref. — Lindner, 1949 (suppl. descr.).
- MERODONTA Kertész, 1908:366. Type-localities: Peru, mouth of Pachitea R. and Meshagua, Urubamba River; Mexico; Costa Rica. Distr. — Mexico to Peru and Bolivia.
- NIGRIBASIS Lindner, 1949:877. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- PICTIPENNIS Kertész, 1908:365. Type-locality: Peru, Meshagua, Urubamba River. Distr. — Peru.
- PROXIMA Kertész, 1908:368. Type-localities: Peru, Urubamba River and Unini, Ucayali River. Distr. — Canal Zone to Peru. Possibly a synonym of *lucida* Gerstaecker.

Genus PARACANTHINOMYIA Lindner

- PARACANTHINOMYIA Lindner, 1964:4, 9. Type-species, *argyrostriata* Lindner (orig. des.).
- ARGYROSTRIATA Lindner, 1964:9. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus PAREIDALIMUS Lindner

- PAREIDALIMUS Lindner, 1964:3, 6. Type-species, *Eidalimus vitrisetosus* Lindner (mon.).
- ARGENTEUS James, 1967e:100. Type-locality: Argentina, Tucumán, Quebrada la Toma. Distr. — Argentina.
- RYCKMANI James, 1965:907. Type-locality: Mexico, Puebla, Acatlán. Distr. — Mexico.
- VITRISETOSUS (Lindner), 1935:398 (*Eidalimus*). Type-locality: Brazil, Santa Catarina, Jaraguá. Distr. — Brazil (Santa Catarina), Argentina (Tucumán, Salta). Ref. — James, 1967c (suppl. descr.).
- punctifer* Malloch of Lindner, 1949:890 (figs. 39-40), 891, misident.

Genus PEDINOCERA Kertész

- PEDINOCERA Kertész, 1909:392. Type-species, *longicornis* Kertész (mon.).
- LONGICORNIS Kertész, 1909:393. Type-locality: Peru, Ucayali River, Unini. Distr. — Peru, Brazil (Santa Catarina).

Genus PELTINA Lindner

- Peltina* Lindner, 1964:7. Type-species, *fuliginosa* Lindner. Nomen nudum.
- PELTINA Lindner, 1969:4. Type-species, *fuliginosa* Lindner (mon.).
- CAZIERI (James), 1953a:5 (*Pachygaster*). Type-locality: Bahama Islands, South Bimini. Distr. — British West Indies, Cuba.
- FULIGINOSA Lindner, 1969:4. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
- fuliginosa* Lindner, 1964:7. Nomen nudum.



Genus PLATYLOBIUM Lindner

- PLATYLOBIUM Lindner, 1933b:326. Type-species, *zurstrasseni* Lindner (orig. des.).
ZURSTRASSENI Lindner, 1933b:326. Type-locality: Peru, Iquitos, Mishuyacu.

Genus POPANOMYIA Kertész

- POPANOMYIA Kertész, 1909:393. Type-species, *femoralis* Kertész (mon.).
FEMORALIS Kertész, 1909:394. Type-locality: Peru, Meshagua, Urubamba River. Distr. — Peru, Brazil (Santa Catarina).

Genus PROEGNOMYIA Kertész

- PROEGNOMYIA Kertész, 1914:467. Type-species, *metallica* Kertész (orig. des.).
Proegnemomyia, error.
METALLICA Kertész, 1914:468. Type-locality: Peru, Urubamba River. Distr. — Guyana, Peru.

Genus PSEPHOCERA Enderlein

- PSEPHOCERA Enderlein, 1914a:300. Type-species, *flavipes* Enderlein (orig. des.).
CALLOSA James, 1967a:13. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.
COGNATA (Lindner), 1949:881 (*Pyelomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
FLAVIPES Enderlein, 1914a:301. Type-locality: Brazil, Santa Catarina. Distr. — Brazil (São Paulo, Santa Catarina).
MARGINATA James, 1967a:12. Type-locality: Dominica, Clarke Hall. Distr. — Dominica.
METZI Johnson, 1919:426. Type-locality: Jamaica, Port Antonio. Distr. — Jamaica.
MODESTA (Lindner), 1949:881 (*Pyelomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.
SP. Lindner, 1964:18. Brazil, Santa Catarina, Nova Teutônia.
SUPERBA (Lindner), 1949:879 (*Pyelomyia*). Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus PSEUDOCYPHOMYIA Kertész

- PSEUDOCYPHOMYIA Kertész, 1916:149. Type-species, *mimetica* Kertész (orig. des.).
MIMETICA Kertész, 1916:150. Type-locality: "Brazil, Amazon". Distr. — Brazil.

Genus SPANIOMYIA Kertész

- SPANIOMYIA Kertész, 1914:462. Type-species, *Artemita pulchripennis* Brauer (orig. des.). Ref. — McFadden, 1971c (review, key).



- OBESA (Walker), 1860:270 (*Clitellaria*). Type-locality: "Mexico". Distr. — Mexico. Ref. — McFadden, 1971c (redeser. of type, figs.).
- PULCHRIFENNIS (Brauer), 1882:74 (*Artemita*). Type-locality: "Brazil". Distr. — Brazil (Guanabara to Santa Catarina).
- SOROR Lindner, 1964:8. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus SPYRIDOPA Gerstaecker

- SPYRIDOPA Gerstaecker, 1857:344. Type-species, *tarsalis* Gerstaecker (mon.). Ref. — Kertész, 1908 (redeser.) (as *Spyripoda*).

Spyripoda, error.

Spyridota, error.

- TARSALIS Gerstaecker, 1857:345. Type-locality: "Brazil". Distr. — Peru, Brazil (Guanabara, Santa Catarina).

Genus STROBILASPIS Lindner

- STROBILASPIS Lindner, 1949:888-889. Type-species, *nigrimana* Lindner (mon.).
- NIGRIMANA Lindner, 1949:890. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus SYNAPTOCHAETA Lindner

- SYNAPTOCHAETA Lindner, 1964:6, 15. Type-species, *digitata* Lindner (mon.).
- DIGITATA Lindner, 1964:15. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus TEGOCERA Lindner

- TEGOCERA Lindner, 1964:6, 13. Type-species, *jamesi* Lindner (mon.).
- JAMESI Lindner, 1964:13. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus THOPOMYIA Kertész


- THOPOMYIA Kertész, 1916:171. Type-species, *dichroa* Kertész (orig. des.).
- DICHROA Kertész, 1916:172. Type-locality: "Peru". Distr. — Peru.

Genus TRIGONOCERINA Lindner

- TRIGONOCERINA Lindner, 1964:5, 11. Type-species, *flaviventris* Lindner (orig. des.).
- FLAVIVENTRIS Lindner, 1964:5, 11. Type-locality: Brazil, Santa Catarina, Nova Teutônia. Distr. — Brazil.

Genus VITTIGER Kertész

- VITTIGER Kertész, 1909:395. Type-species, *schnusei* Kertész (mon.).
- SCHNUSEI Kertész, 1909:396. Type-locality: Peru, mouth of Pachitca R. Distr. — Peru, Brazil (Santa Catarina).



Genus ZABRACHIA Coquillett

- ZABRACHIA Coquillett, 1901:585. Type-species, *polita* Coquillett (orig. des.).
Ref. — Kraft & Cook, 1961 (revision); James, 1965 (partial key,
key to larvae).
- CORNUTA Kraft & Cook, 1961:9. Type-locality: U.S.A., Arizona, Globe.
Distr. — U.S.A. (Arizona), Mexico (Sinaloa).
- STOICHOIDES James, 1965:905. Type-locality: Mexico, Oaxaca, Oaxaca. Distr.
— Mexico.

Unplaced Species

- SCUTELLARIS Walker, 1854:313 (*Clitellaria*). Type-locality: Brazil, Pará,
Santarém. A pachygastrine, according to Kertész.

LITERATURE CITED

ADAMS, C. F.

1903. Dipterological contributions. *Kansas Univ. Sci. Bull.* 2:21-47.

ALBUQUERQUE, D. DE O.

1955. Duas novas espécies de Hermetiinae do Brasil (Diptera, Stratiomyidae). *Rev. Brasil. Entomol.* 3:129-135.

ALDRICH, J. M.

1928. New Diptera or two-winged flies from South America. *Proc. U. S. Nat. Mus.* 74(1):1-25.

AUBERTIN, D.

1930. Stratiomyidae, in *British Museum (Natural History): Diptera of Patagonia and South Chile* 5(2):93-105.
1932. Note on the name of a genus in the family Stratiomyidae, in *British Museum (Natural History): Diptera of Patagonia and South Chile* 5(3):284.

BARRETTO, M. P.

1947. Estudos sobre "Stratiomyidae" brasileiros. I. Duas novas espécies de "*Rhingiopsis*" Röder, 1886 (Diptera). *Rev. Brasil. Biol.* 7: 439-443, 3 figs.

BELLARDI, L.

1859. *Saggio di Ditterologia Messicana. 2. Parte I.* 77 pp. (Also 1861, pp. 201-277).
1862. *Saggio di Ditterologia Messicana. 2. Appendice*, 28 pp. Torino. (Also 1864:202-227).

BEZZI, M.

1907. Nomenklatorisches über Dipteren. *Wien. Entomol. Ztg.* 26:51-56.
- 1908a. Eine neue brasilianische Art der Diptere ngattung *Allognosta* O. S. *Deut. Entomol. Ztschr.* 1908:471-475.
- 1908b. Nomenklatorisches über Dipteren. *Wien. Entomol. Ztg.* 27: 74-84.



1922. On the South American species of the dipterous genus *Chiro-myza* Wied. *Ann. Entomol. Soc. Amer.* 15:117-124.
- BIGOT, J. M. F.
1875. Diptères nouveaux ou peu connus. 5^e partie. VII: Espèces nouvelles du genre *Cyphomyia*. *Ann. Soc. Entomol. France* (5) 5:483-488.
1879. Diptères nouveaux ou peu connus. 11^e Partie. XVI. Curiae Xylophagidarum et Stratiomyidarum. *Ann. Soc. Entomol. France* (5)9:183-234.
1887. Diptères nouveaux ou peu connus. 31^e Partie. XXXII. Descriptions de nouvelles espèces de Stratiomyidi et de Conopsidi. *Ann. Soc. Entomol. France* (6)7:20-46.
- BLANCHARD, E.
1852. Orden IX, Dipteros, pp. 327-468, in Gay, C., Ed., *Historia Fisica y Política de Chile. Zoología*, Vol. 7, 471 pp. Paris.
- BLANCHARD, E. E.
1923. Apuntes sobre dos dípteros argentinos. *Physis* 6:319-323.
- BRAUER, F.
1882. Die Zweiflügler des Kaiserlichen Museums zu Wien. II. *Denkschr. K. Akad. Wiss. Wien, Math.-Nat. Cl.* 44(1):59-110.
- BRÉTHES, J.
1907. Catálogo de los dípteros de las repúblicas de La Plata. *An. Mus. Nac. Buenos Aires* (3, ix) 16:277-305.
1922. Himenópteros y dípteros de varias procedencias. *An. Soc. Cient. Argentina* 93:119-146.
1924. Sur quelques diptères chiliens. *Rev. Chilena Hist. Nat.* 28: 104-111.
- CARRERA, M. & J. LANE
1945. Diptera de Caiobá (Est. do Paraná) (Diptera, Stratiomyiidae e Tabanidae). *Arg. Mus. Paranaense* 4:127-136.
- COLE, F. R.
1923. Expedition of the California Academy of Sciences to the Gulf of California in 1921. II. General Report. *Proc. Calif. Acad. Sci.* (4)12:457-481.
- COQUILLET, D. W.
1901. Original descriptions of Diptera. In Needham, J. G., & Betten, C., *Bull. N. Y. State Museum* 47:383-612 (pp. 585-6).
1902. New Diptera from North America. *Proc. U. S. Nat. Mus.* 25: 83-126.
1904. Diptera from southern Texas with descriptions of new species. *J. N. Y. Entomol. Soc.* 12:31-35.
1909. A new stratiomyid from Texas. *Canad. Entomol.* 41:212.
1910. The type-species of the North American genera of Diptera. *Proc. U. S. Nat. Mus.* 37:499-647.
- CRESSON, E. T. JR.
1919. Dipterological notes and descriptions. *Proc. Acad. Nat. Sci. Philadelphia* 1919:171-194.



CURRAN, C. H.

1925. New American Diptera, I-II. *Ann. Mag. Nat. Hist.* (9)16:243-253, 338-354.
- 1928a. Insects of Porto Rico and the Virgin Islands. Diptera or Two-winged flies. Pt. 1, pp. 1-118. *In* New York Academy of Sciences, *Scientific Survey of Puerto Rico and the Virgin Islands*. Vol. 11.
- 1928b. Records and descriptions of Diptera, mostly from Jamaica. *Bull. Agric. & Sci. Jamaica* 4, suppl. 4:14-30.
1929. New Diptera in the American Museum of Natural History. *Amer. Mus. Novitates* 339:1-13.
1931. First supplement to the "Diptera of Porto Rico and the Virgin Islands". *Amer. Mus. Novitates* 456:1-23.
- 1932a. New North American Diptera, with notes on others. *Amer. Mus. Novitates* 526:1-13.
- 1932b. New American Diptera. *Amer. Mus. Novitates* 534:1-15.
- 1932c. Diptera (Excl. of Tipulidae and Culicidae). The Norwegian Zoological Expedition to the Galapagos Islands 1925, conducted by Alf Wollebaek. *Nyt Mag. Naturvidenskaberne, B* 71:347-366.
1933. Two new Diptera from Guatemala. *Amer. Mus. Novitates* 643:1-2.
1934. The Diptera of Kartabo, Bartica District, British Guiana, with descriptions of new species from other British Guiana localities. *Bull. Amer. Mus. Nat. Hist.* 66:287-532.
1936. The generic status of *Lagarus* Philippi (Stratiomyidae). *Rev. Chilena Hist. Nat.* 60:281-282.

ENDERLEIN, G.

- 1913a. Dipterologische Studien. III. Über *Lagarinus* nov. gen., eine isoliert stehende Fliegengattung. *Zool. Anz.* 42:250-253.
- 1913b. Dipterologische Studien. V. Zur Kenntnis der Familie Xylophagidae. *Zool. Anz.* 42:532-552.
- 1914a. Dipterologische Studien. VIII. Zur Kenntnis der Stratiomyiden-Unterfamilien mit 2ästiger Media Paehygasterinae, Lophotelinae und Prosopochrysiniae. *Zool. Anz.* 43:289-315.
- 1914b. Dipterologische Studien. IX. Zur Kenntnis der Stratiomyiden mit 3ästiger Media und ihre Gruppierung. A. Formen, bei denen der 1. Cubitalast mit der Discoidalzelle durch Querader verbunden ist oder sie nur in einem Punkte berührt (Subfamilien: Geosaraginae, Analcocerinae, Stratiomyiinae). *Zool. Anz.* 43:577-615.
- 1914c. Dipterologische Studien. X. Zur Kenntnis der Stratiomyiden mit 3ästiger Media und ihre Gruppierung. B. Formen, bei denen der 1. Cubitalast mit der Discoidalzelle eine Strecke verschmolzen ist (Familien: Hermetiinae, Clitellariinae). *Zool. Anz.* 44:1-25.
1921. Dipterologische Studien. Ueber die phylotische älteren Stratiomyidensubfamilien (Xylophaginae, Chiromyzinae, Solvinae, Beridinae, und Coenomyiinae). *Mitt. Zool. Mus. Berlin* 10:153-214.
1930. Dipterologische Studien. XX. *Deut. Entomol. Ztsehr.* 1930: 65-71.



FABRICIUS, J. C.

1781. *Species insectorum exhibentes eorum differentias specificas, synonyma, auctorum, loca natalia, metamorphosin*. Vol. 2, 517 pp. Hamburgi et Kilonii (Hamburg and Kiel).
1787. *Mantissa inscctorum sistens species nuper detectas*. Vol. 2, 382 pp. Hafniae (Copenhagen).
1794. *Entomologia systematica emendata et aucta*. Vol. 4, 472 pp. Hafniae (Copenhagen).
1798. *Supplementum entomologiae systematicae*. 572 pp. Hafniae (Copenhagen).
1805. *Systema antliatorum, secundum ordines, genera, species*. 373 pp. Brunsvigae (Brunswick).

FALLÉN, C. F.

1817. *Stratiomyidae Svcciac*. 14 pp. Lundae (Lund).

FREY, R.

1934. Diptera Brachycera von den Sunda-Inseln und Nord-Australien. *Rev. Suisse Zool.* 41: 299-339, 6 figs.

GMELIN, J. F.

1790. *Caroli à Linné, Systema naturae per regna tria naturae*, Ed. 13. Vol. 1: *Regnum Animale*, 5:2225-3020. Lipsiae (Leipzig).

GEOFFROY, E. L.

1762. *Histoire abrégée des insectes qui se trouvent aux environs de Paris*. Vol. 2, 690 pp. Paris. Published anonymously, reissued in 1764 with Geoffroy as author.

GERSTAECKER, A.

1857. Beitrag zur Kenntnis exotischer Stratiomyiden. *Linnaea Entomol.* 11:261-350.

GIGLIO-TOS, E.

1891. Nuove specie di ditteri del Museo Zoologico di Torino. *Boll. Mus. Zool. Anat. Comp. Univ. Torino* 6 (102) 1-4.

GREENE, C. T.

1940. Two new species of the genus *Hermetia* (Stratiomyiidae — Diptera). *Proc. Entomol. Soc. Washington* 42:150-155.

GUÉRIN-MÉNEVILLE, F. E.

1831. Zoologie, Atlas, Insectes, in Duperrey, L. I., ed., *Voyage autour du monde sur la corvette de sa majesté La Coquille*. Paris.

HANSON, W. J.

1958. A revision of the subgenus *Melanonemotelus* of America north of Mexico (Diptera, Stratiomyidae). *Univ. Kansas Sci. Bull.* 38:1351-1391.

HARDY, G. H.

1920. A revision of the Chiromyzini (Diptera). *Proc. Linn. Soc. New South Wales* 45:532-542.



IIIENE, J. S.

1911. A new species of *Nothomyia*. *Ohio Naturalist* 11:301-302.

HOLLIS, D.

1962. New and little known Stratiomyidae (Diptera, Brachycera) in the British Museum. *Ann. Mag. Nat. Hist.* (13) 5:557.

HUNTER, W. D.

1900. A catalogue of the Diptera of South America. Part II, Homodactyla and Mydadae. *Trans. Amer. Entomol. Soc.* 27:121-131.

IBBE, P.

- 1963a. Contribuição ao conhecimento do gênero *Chordonota* Gers-taecker, 1857 (Diptera, Stratiomyidae). *Mem. Inst. Oswaldo Cruz* 61:401-409.
- 1963b. Contribuição ao conhecimento do gênero *Cyphomyia* Wiede-mann, 1819 (Diptera, Stratiomyidae). *Mem. Inst. Oswaldo Cruz* 61:25-39.
1965. Contribuição ao conhecimento das formas imaturas de uma espécie do gênero *Chrysochlorina* James, 1939 (Diptera, Stratio-myidae). *Rev. Brasil. Biol.* 25:429-438.
1966. Estudo sobre as espécies brasileiras do gênero *Chrysochlorina* James, 1939 (Diptera, Stratiomyidae). *Arq. Zool.* 14:69-113.
- 1967a. Contribuição ao conhecimento do gênero *Chaetosargus* Röder, 1894 (Diptera, Stratiomyidae). *Rev. Brasil. Biol.* 27:63-70.
- 1967b. Estudos complementares sobre duas espécies brasileiras do gênero *Chrysochlorina* James (Diptera, Stratiomyidae). *Studia Entomol.* 10:473-480.
- 1967c. Estudo sobre uma nova espécie amazônica do gênero *Cypho-myia* Wiedemann (Diptera, Stratiomyidae). *Atas do Simpósio sobre a Biota Amazônica* 5 (Zoologia): 225-238.
- 1968a. Estudo sobre "*Chrysochlorina amethystina*" (Fabricius, 1805) e suas relações com as espécies brasileiras do gênero "*Chry-sochlorina*" James, 1939 (Diptera, Stratiomyidae). *Rev. Brasil. Biol.* 28:363-368.
- 1968b. Contribuição ao conhecimento das espécies brasileiras do gênero "*Euryneura*" Schiner, 1867 (Diptera, Stratiomyidae). *Rev. Brasil. Biol.* 28:251-272.
1971. Estudos sobre os Stratiomyidae [sic] da coleção do United States National Museum. II: O gênero *Pelagomyia* Williston, 1896 (Insecta, Diptera). *Rev. Brasil. Biol.* 31(4):497-506, 33 figs.

INTERNATIONAL COMMISSION ON ZOOLOGICAL NOMENCLATURE


- 1957a. Opinion 441. Validation under the Plenary Powers of the names for five genera in the order Diptera (Class Insecta) published in 1762 by Geoffroy. In Hemming, F., ed., *Opinions and Declarations rend. Internatl. Comm. Zool. Nomencl.* 15:83-120.
- 1957b. Opinion 442. Validation under the Plenary Powers of the generic name "*Stratiomys*" Geoffroy, 1762 (Class Insecta, Order Diptera). In Hemming, F., Ed., *Opinions and Declarations rend. Internatl. Comm. Zool. Nomencl.* 15:121-162.
1963. Opinion 678. The suppression under the Plenary Powers of the pamphlet published by Meigen, 1800. *Bull. Zool. Nomencl.* 20: 339-342.

JAENNICKE, F.

1867. Neue exotische Dipteren. *Abhandl. Senckenb. Naturf. Ges.* 6: 311-406.

JAMES, M. T.

1932. New Stratiomyidae in the American Museum of Natural History. *Amer. Mus. Novitates* 571:1-7.
1933. New Stratiomyidae in the Snow Entomological Collection. *J. Kansas Entomol. Soc.* 6:66-71.
1934. *Hoplitimyia*, a new genus of Stratiomyidae. *Ann. Entomol. Soc. Amer.* 27:443-444.
- 1935a. The genus *Iermetia* in the United States (Diptera, Stratiomyidae). *Bull. Brooklyn Entomol. Soc.* 30:165-170.
- 1935b. A review of the Nearctic Geosarginae (Diptera, Stratiomyidae). *Canad. Entomol.* 47:267-275.
- 1936a. New and little-known neotropical Stratiomyidae (Diptera) in the Museum of Comparative Zoology. *Psychic* 43:49-55.
- 1936b. Notes on *Nemotelus* (Dipt., Stratiomyidae). *Bull. Brooklyn Entomol. Soc.* 31:86-91.
- 1936c. The genus *Odontomyia* in America north of Mexico (Diptera, Stratiomyidae). *Ann. Entomol. Soc. Amer.* 29:517-550.
1937. Some new and little-known neotropical and subtropical Stratiomyidae. *Bull. Brooklyn Entomol. Soc.* 32:149-155.
1938. New and little known neotropical Stratiomyidae. *Rev. Entomol.* 8:196-203.
- 1939a. Neotropical flies of the family Stratiomyidae in the United States National Museum. *Proc. U. S. Nat. Mus.* 86(3065):595-607.
- 1939b. Studies in Neotropical Stratiomyidae. I. The American species formerly referred to *Chrysochlora* Latreille. II. The Genus *Hoplitimyia* James. III. The genus *Udamacantha* Enderlein. *J. Kansas Entomol. Soc.* 12:32-46.
- 1939c. The species of *Euparyphus* related to *crotchii* O. S. (Diptera, Stratiomyidae). *Pan-Pacific Entomol.* 15:49-56.
- 1940a. Studies in neotropical Stratiomyidae (Diptera). IV. The genera related to *Cyphomyia* Wiedemann. *Rev. Entomol.* 11:119-149.
- 1940b. Two new neotropical Stratiomyidae. *Arb. Morph. Taxon. Entomol.*, Berlin-Dahlem 7:120-122.
- 1941a. A preliminary study of the New World Geosarginae (Dipt., Stratiomyidae). *Lloydia* 4:300-309.
- 1941b. New species and records of Mexican Stratiomyidae (Diptera). *An. Escuela Nac. Cien. Biol.* 2:241-249.
- 1941c. Notes on the Nearctic Geosarginae (Diptera: Stratiomyidae). *Entomol. News* 52:105-108.
1942. A review of the Myxosargini (Diptera, Stratiomyidae). *Pan-Pac. Entomol.* 18:49-60.
- 1943a. Studies in Neotropical Stratiomyidae. V. The classification of the Rhaphiocerinae. *Ann. Entomol. Soc. Amer.* 36:365-379.
- 1943b. Studies in neotropical Stratiomyidae. VI. A new genus related to *Adoxomyia*. *Ann. Entomol. Soc. Amer.* 36:380-382.
1949. Some African and Brazilian Stratiomyidae (Diptera). *Amer. Mus. Novitates* 1386:1-7.



1950. Some new and poorly-known Adoxomyiinae (Diptera, Stratiomyidae) from the Southwest. *J. Kansas Entomol. Soc.* 23:71-73.
1952. The Ethiopian genera of Sarginae, with descriptions of new species. *J. Washington Acad. Sci.* 42:220-226.
- 1953a. The Stratiomyidae (Diptera) of Bimini, British West Indies. *Amer. Mus. Novitates* 1613:1-6.
- 1953b. A preliminary review of the Argentine genera and species of Stratiomyidae (Diptera). Part I. Stratiomyinae. *Acta Zool. Lilloana* 13:307-326.
- 1957a. The genus *Eulalia* in Florida and the West Indies. *Florida Entomol.* 40:15-18.
- 1957b. The larva of *Cyphomyia* (Diptera, Stratiomyidae) and its significance in classification. *Ann. Entomol. Soc. Amer.* 50:639-641.
- 1962a. Diptera: Stratiomyidae, Calliphoridae. *Insects of Micronesia*, 13:75-127.
- 1962b. The genus *Dicyphoma* James (Diptera: Stratiomyidae). *Ann. Entomol. Soc. Amer.* 55:15-20.
1965. Contributions to our knowledge of the nearctic Paehygasterinae (Diptera: Stratiomyidae). *Ann. Entomol. Soc. Amer.* 58:902-908.
- 1966a. The Stratiomyidae (Diptera) of the Galapagos Islands. *Proc. Calif. Acad. Sci.* (4)33:535-541.
- 1966b. The genera of Rhabdioerinae with the elongated first antennal segment (Diptera: Stratiomyidae). *J. Kansas Entomol. Soc.* 39:676-681.
- 1966e. A new genus of paehygasterine Stratiomyidae reared from caetus (Diptera). *J. Kansas Entomol. Soc.* 39:109-112.
- 1967a. Bredin-Arebold-Smithsonian Biological Survey of Dominica. 5. Family Stratiomyidae (Diptera). *Proc. U. S. Nat. Mus.* 123 (3622):1-22.
- 1967b. The *Hermetia comstocki* group. (Diptera: Stratiomyidae). *Pan-Pac. Entomol.* 43:61-64.
- 1967e. A preliminary review of the Argentine genera and species of Stratiomyidae (Diptera). Part 2. Paehygasterinae. *Acta Zool. Lilloana* 21:95-121.
1970. A new species, correction of synonymy, and new records of nearctic Stratiomyidae (Diptera). *Proc. Entomol. Soc. Washington* 72:327-332.
1971. The South American species of *Artemita*. *J. Kansas Entomol. Soc.* 44:59-70.

JAMES, M. T., & M. W. MCFADDEN

1969. The genus *Adoxomyia* in America north of Mexico (Diptera: Stratiomyidae). *J. Kansas Entomol. Soc.* 42:260-276.
1971. The genus *Merosargus* in Middle America and the Andean Subregion (Diptera: Stratiomyidae). *Melandria* 7:1-76.

JAMES, M. T., & W. W. WIRTH

1967. The species of *Hermetia* of the *aurata* group (Diptera: Stratiomyidae). *Proc. U. S. Nat. Mus.* 123(3603):1-19.



JOHNSON, C. W.

1894. List of the Diptera of Jamaica with descriptions of new species. *Proc. Acad. Nat. Sci. Philadelphia* 1894:271-281.
1895. A review of the *Stratiomyia* and *Odontomyia* of North America. *Trans. Amer. Entomol. Soc.* 22:227-278.
1913. Insects of Florida. I. Diptera. *Bull. Amer. Mus. Nat. Hist.* 32: 37-90.
1919. A revised list of the Diptera of Jamaica. *Bull. Amer. Mus. Nat. Hist.* 41:421-449.

KERTÉSZ, K.

1907. Ein neuer Dipteren-Gattungsname. *Ann. Mus. Nat. Hungarici* 5:499.
1908. Vorarbeiten zu einer Monographie der Notacanthen. I-XI. *Ann. Mus. Nat. Hungarici* 6:321-374.
1909. Vorarbeiten zu einer Monographie der Notacanthen. XII-XXII. *Ann. Mus. Nat. Hungarici* 7:369-397.
1911. Ueber die generische Hinzugehörigkeit der bis jetzt beschriebenen *Pachygaster*-Arten, pp. 29-32, in Severin, G., sec., 1er *Congrès international d'entomologie*. Vol. 2: Mem., 520 pp. Bruxelles.
1912. The Percy Sladen Trust Expedition to the Indian Ocean in 1905, under the leadership of Mr. J. Stanley Gardiner, M. A. Volume IV. No. VI. Diptera, Stratiomyidae. *Trans. Linn. Soc. London* (2) *Zool.* 15:95-99.
1914. Vorarbeiten zu einer Monographie der Notacanthen. XXIII-XXXV. *Ann. Mus. Nat. Hungarici* 12:449-557.
1916. Vorarbeiten zu einer Monographie der Notacanthen. XXXVI-XXXVIII. *Ann. Mus. Nat. Hungarici* 14:123-218.
1923. Vorarbeiten zu einer Monographie der Notacanthen. XLV-L. *Ann. Mus. Nat. Hungarici* 20:85-129.

KRAFT, K. J., & E. F. COOK

1961. A revision of the Pachygasterinae (Diptera, Stratiomyidae) of America north of Mexico. *Misc. Publ. Entomol. Soc. Amer.* 3:1-24.

LATREILLE, P. A.

1802. Histoire naturelle, générale et particulière, des crustacés et des insectes, Tome troisième, 468 pp. In Sonnini, C. S., ed., *Histoire naturelle par Buffon*. Paris.
1804. Tableau méthodique des insectes. In Société de Naturalistes et d'Agriculteurs, *Nouveau dictionnaire d'histoire naturelle*. Vol. 24, sect. 3, Tableaux méthodiques d'histoire naturelle, pp. 129-200.

LINDNER, E.

1926. Dr. L. Zürchers Dipteren-Ausbeute aus Paraguay: Stratiomyiden. *Arch. für Naturges.* 1926:94-103.
1928. Die von Prof. Dr. A. Seitz in Brasilien gesammelten Stratiomyiden (Ins. Dipt.). *Senckenbergiana* 10:235-244.



1929. Ergebnisse einer zoologischen Sammelreise nach Brasilien insbesondere in das Amazonasgebiet, ausgeführt von Dr. H. Zerny. *Ann. Naturh. Mus. Wien* 43:257-268.
1930. Die Ausbeute der deutschen Chaco-Expedition 1925/26. Diptera. XIV. Stratiomyidae. *Konowia* 8:273-284 (1929).
1931. Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Rev. Entomol.* 1:304-312.
- 1933a. Zweiter Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Rev. Entomol.* 3:199-205.
- 1933b. Neotropische Stratiomyiden des Senckenberg-Museums (Diptera). *Senckenbergiana* 15:325-334.
1935. Dritter Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Rev. Entomol.* 5:396-413.
1936. Stratiomyiden von Costa Rica (Dipt.). *Stettiner Entomol. Zeitung* 97:153-158.
1940. Stratiomyiden von Costa Rica (Nachtrag). *Zool. Anz.* 132:193-194.
1941. Stratiomyiden (Dipt.) in *Beiträge zur Fauna Perus*. 1:177-188.
1943. Südhilensische Stratiomyiden (Dipt.). *Ann. Naturh. Mus. Wien* 53(2):89-100.
1949. Neotropische Stratiomyiden des Britischen Museums in London. *Ann. Mag. Nat. Hist.* (12)1:782-821, 851-891.
1951. Viertes Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Rev. Entomol.* 22:245-264.
1964. Beitrag zur Kenntnis der neotropischen Pachygasterinae (Stratiomyiidae, Dipt.). *Stuttgarter Beiträge zur Naturkunde* 129:1-22.
1965. Stratiomyiden aus dem Amazonas-gebiet. *Amazoniana* 1:84-86.
1969. Fünftes Beitrag zur Kenntnis der südamerikanischen Stratiomyidenfauna (Dipt.). *Stuttgarter Beiträge zur Naturkunde* 203:1-14.

LINNAEUS, C.

1758. *Systema naturae per regna tria naturae*. Ed. 10, vol. 1: 824 pp. Holmiae (Stockholm).
1767. *Systema naturae per regna tria naturae*. Ed. 12. Vol. 1, part 2, pp. 533-1327. Holmiae (Stockholm).

LOEW, HIERMANN

1847. *Chauna* genus novum. *Stett. Entomol. Z.* 8:370.
1855. Einige Bemerkungen über die Gattung *Sargus*. *Verhandl. Zool. Bot. Ver. Wien* 5:131-148.
1863. Diptera Americae systemtrionalis indigena. Centuria tertia. *Berl. Entomol. Ztschr.* 7:1-55.
- 1866a. Diptera Americae septentrionalis indigena. Centuria sexta. *Berl. Entomol. Ztschr.* 9:127-186.
- 1866b. Diptera Americae septentrionalis indigena. Centuria septima. *Berl. Entomol. Ztschr.* 10:1-54.
1869. Diptera Americae septentrionalis indigena. Centuria octava. *Berl. Entomol. Ztschr.* 13:1-52.



1872. *Diptera Americae septentrionalis indigena. Centuria decima.*
Berl. Entomol. Ztschr. 16:49-115.

LYNCH ARRIBÁLZAGA, E.

1881. Neue Dipteren aus dem südlichen Gebiet der Pampa. *Stettin.*
Entomol. Zeitung 42:189-192.

MACQUART, J.

1834. *Histoire naturelle des Insectes. Diptères* 1:578 pp., in N. E.
Roret, ed.: *Collection des Suites à Buffon.* Paris.

1838. *Diptères exotiques nouveaux ou peu connus. 1re partie. Mém.*
Soc. Roy. Sci., Agric., Arts Lille 1838:9-225. Also published as
separate, 1:205-221.

1846. *Diptères exotiques nouveaux ou peu connus. 1er. supplément.*
Mém. Soc. Roy. Sci., Agric., Arts Lille (1845) (1844):133-364. Also
published separately as *Supplément (I)*, pp. 5-238.

1847. *Diptères exotiques nouveaux ou peu connus. 2e supplément.*
Mém. Soc. Sci., Agr., Arts Lille 1846:21-120 (separate, 5-104).

1850. *Diptères exotiques nouveaux ou peu connus. 4e supplément.*
Mém. Soc. Roy. Sci., Agric., Arts Lille 1849:309-479. Also published
separately as *Supplément 4*, pp. 5-161.

1855. *Diptères exotiques nouveaux ou peu connus. 5e supplément.*
Mém. Soc. Sci., Agr., Arts Lille 1854:25-156 (separate, 5-136).

MALLOCH, J. R.

1915. A revision of the North American Paehygasterinae with
unspined scutellum (Diptera). *Ann. Entomol. Soc. Amer.* 8:305-321.

1917. A preliminary classification of Diptera, exclusive of Pupipara,
based upon larval and pupal characters, with keys to imagines
in certain families. *Bull. Ill. State Lab. Nat. Hist.* 12:161-407.

MCFADDEN, M. W.

1967. Soldier fly larvae in America north of Mexico. *Proc. U. S. Nat.*
Mus. 121(3569):1-72.

1970a. Notes on the synonymy of *Chrysochroma* Williston and a new
name for the species formerly referred to *Chrysochroma* (Diptera:
Stratiomyidae). *Proc. Entomol. Soc. Wash.* 72:274.

1970b. New Rhabdiocerinae from Mexico, with a key to known
genera (Diptera: Stratiomyidae). *Ann. Entomol. Soc. Amer.* 63:
316-320.

1971a. A note on the synonymy of *Antissops denticulata* Enderlein.
Proc. Entomol. Soc. Washington 73:51.

1971b. Synonymies and combinations for Mexican soldier flies
(Diptera: Stratiomyidae). *Proc. Entomol. Soc. Washington* 73:
411-413.

1971c. A review of the genus *Spaniomyia* (Diptera: Stratiomyidae).
Ann. Entomol. Soc. Amer. 64:863-864.

MCFADDEN, M. W., & M. T. JAMES

1969. The *Cyphomyia bicarinata* species group (Diptera: Stratio-
myidae). *J. Kansas Entomol. Soc.* 42:313-320.



MEIGEN, J. W.

1800. *Nouvelle classification des mouches à deux ailes (Diptera L.) d'après un plan tout nouveau*, 40 pp. Paris.
1803. Versuch einer neuen Gattungseintheilung der europäischen zweiflügeligen Insekten. *Mag. f. Insektenkunde* 2:259-281.
1804. *Klassifikation und Beschreibung der europäischen zweiflügeligen Insekten (Diptera Linn.)*. Erster Band, Abt. 1, pp. xxviii + 152. Braunschweig.

MELANDER, A. L.

1903. A review of the North American species of *Nemotelus*. *Psyche* 10:171-183.
1904. Notes on North American Stratiomyidae. *Canad. Entomol.* 36: 14-24.

MILLER, D.

1945. Generic name changes in Diptera. *Proc. Royal Entomol. Soc. London (B)* 14:72.

NAGATOMAI, A., & J. YUKAWA

1968. The genus *Inopus (Metoponia, Altermetoponia)* (Diptera: Stratiomyidae). *Pacific Insects* 10:521-528.

OSTEN SACKEN, C. R.

1883. Synonymica concerning exotic Dipterology. No. II. *Berl. Entomol. Z.* 27:295-298.
1878. Catalogue of the described Diptera of North America (Ed. 2). *Smithsn. Misc. Collect.* 16(2):1-276.
1886. Diptera, Vol. I (part., conel.), in Godman, F. D., and Salvin, O., eds., *Biologia Centrali-Americana, Insecta-Diptera* 1:378 pp.

PERTY, M.

1833. *Insecta Brasiliensia*, pp. 125-224, pl. 25-40, in his *Delectus animalium articulorum quae in itinere per Brasiliam annis MDCCCXVII-MDCCCXX jussu et auspiciis Maximiliani Josephi I Bavariae regis augustissimi peracto collegerunt Dr. J. B. de Spix et Dr. C. F. Ph. de Martius*, 44 + 224 pp., 40 pl. Monachii (Munich). "1830-1834".

PHILIPPI, R. A.

1865. Aufzählung der chilenischen Dipteren. *Verh. zool. bot-Ges. Wien* 15:595-782.

RÖDER, V. V.

1886. Ueber drei neue Gattungen der Notacanthen. *Entomol. Nachr.* 12:137-140.
1894. *Chaetosargus* nov. gen. Dipteroorum. *Wien. Entomol. Zeitung* 13:169.

RONDANI, C.

1848. *Esame di varie specie d'insetti ditteri brasiliani*. *Studi Entomol.*, Turin 1:63-112.



1850. Osservazioni sopra alquante specie di cesapodi ditteri del Musco Torinese. *Nuovi Ann. e Rend. Accad. Sci. Ist. e Soc. Agric. Bologna* (3)2:165-197.
1856. *Dipterologiae Italicae prodromus. Vol. 1. Genera Italica ordinis dipterorum ordinatim disposita et distincta et in familias et stirpas aggregata.* 228 pp. Parmae (Parma).
1863. *Diptera exotica revisa et annotata*, 99 pp., 1 pl., Modena (also published as *Dipterorum species et genera aliqua exotica revisa et annotata novis nonnullis descriptis. Arch. Zool. Anat. Fis. Modena* 3:1-99, 1864).

SAY, T.

1824. Appendix. Part I. — Natural History. 1. Zoology. E. Class Insecta, pp. 268-387. In Keating, W. H., *Major Long's Second Expedition*, Vol. 2, 459 pp. Philadelphia.
1829. Descriptions of North American dipterous insects. *J. Acad. Nat. Sci. Philadelphia* 6:149-178.

SCHINER, I. R. (J. R.)

1867. Zweiter Bericht über die von der Weltumsegelungsreise [sic] der k. Fregatte Novara mitgebrachten Dipteren. *Verhandl. K.-K. Zool.-Bot. Gesell. Wien* 17:355-412.
1868. Diptera (Art. 1) 388 pp. in Wüllerstorff-Urbair, B. von, *Reise der österreichischen Fregatte Novara, Zool.*, Vol. 2, Abt. 1, Sect. B. Wien.

SZILÁDY, Z.

1929. Notacanthén-Studien. I., II. *Ann. Mus. Nat. Hungarici* 26: 72, 250.

THOMSON, C. G.

1869. 6. Diptera. Species nova descripsit. Pp. 443-614. In K. Svenska Vetenskaps-Akademién, *Kongliga svenska fregatten Eugénies resa omkring jorden. Pt. 2: Zoologic, 1, Insekter*, 617 pp. Stockholm, "1868".

TOWNSEND, C. H. T.

1895. Notes on the Diptera of Baja California. *Proc. Calif. Acad. Sci.* (2)4:593-620.

WALKER, F.

1837. Descriptions of Diptera collected by Capt. P. P. King, R. N., F. R. S., in the survey of the Straits of Magellan. *Trans. Linn. Soc. London* 17:331-359.
1849. *List of the specimens of dipterous insects in the collection of the British Museum.* Vol. 3, pp. 485-697. London.
1850. Characters of undescribed diptera in the British Museum. *Newman's Zoology* 8 (Appendix) (Art. 5):xcv-xcix.
1854. *List of the specimens of dipterous insects in the collection of the British Museum.* Part 5, pp. 1-330. London.
1856. *Insecta Saundersiana, or characters of undescribed insects in the collection of W. W. Saunders.* Diptera, 473 pp. London.



1857. Characters of undescribed Diptera in the collection of W. W. Saunders, Esq., F. R. S., *Trans. Entomol. Soc. London* (n.s.) 4:119-158.

1860. Characters of undescribed Diptera in the collection of W. W. Saunders, Esq., F. R. S., *Trans. Entomol. Soc. London* (n.s.) 5:268-334.

WANDOLLECK, B.

1897. *Blastocera atra*, eine neue Diptere aus St. Cruz. *Wiener Entomol. Z.* 16:216.

WIEDEMANN, C. R. W.

1819. Brasilianische Zweiflügler. *Zool. Mag.* 1(3):40-56.

1820. *Nova dipterorum genera iconibus illustrata*. 23 pp. Kiliae (Kiel).

1821. *Diptera exotica*. Pars I. Pp. xix + 244 pp. Kiliae.

1824. *Munus rectoris in Academia Christiana Albertina aditurus analecta entomologia ex Museo Regio Havniensi*. 60 pp. Kiliae (Kiel).

1828. *Aussereuropäische zweiflügelige Insekten*. I. xxxii + 608 pp.

1830. *Aussereuropäische zweiflügelige Insekten*. II. Pp. xli + 684 pp.

WILLISTON, S. W.

1885. Notes and descriptions of North American Xylophagidae and Stratiomyidae. *Canad. Entomol.* 17:121-128.

1888. Diptera Brasilliana, ab H. H. Smith collecta. Part I. — Stratiomyidae, Syrphidae. *Trans. Amer. Entomol. Soc.* 15:243-292.

1896a. *Manual of the families and genera of North American Diptera*. Ed. 2, 167 pp. New Haven, Conn.

1968b. Diptera of St. Vincent. *Trans. Entomol. Soc. London* 1896: 299-302.

1900. Supplement (Part), pp. 249-264, 265-272, in Godman, F. D., and Salvin, O., eds., *Biologia Centrali-Americana. Zoologia-Insecta Diptera* 1:378 pp.

WULP, F. M. VAN DER

1891. Amcrikaansche Diptera. *Tijdschr. Entomol.* 24:141-168.

1885. On exotic Diptera. *Notes Leyden Mus.* 7:1-15, 57-86.

1888. Nieuwe Argentijnsche Diptera van wijlen Prof. H. Weyenbergh Jr. *Tijdschr. v. Entomol.* 31:361-376.

INDEX

abana Curran, 1929, <i>Cyphomyia</i>	26.25
abana Curran, 1932, <i>Merosargus</i>	26.10
Abasanistus Kertész, 1923	26.24
Abavus Enderlein, 1914	26.29
abditus Lindner, 1949, <i>Hedriodiscus</i>	26.40
abdominalis James, 1936, <i>Microchrysa</i>	26.16




<i>Acanthina</i> Wiedemann, 1830	28.49
<i>Acanthinia</i> , error	26.49
<i>Acanthinomyia</i> Hunter, 1901	26.49
<i>Acrochaeta</i> Wiedemann, 1830	26.8
<i>Acrodcsmia</i> Enderlein, 1914	26.36
<i>Acropeltates</i> Kertész, 1923	26.29
<i>Actina</i> Meigen, 1804	26.4
<i>Actinomia</i> , error	26.4
<i>Actinomyia</i> Lindner, 1949	26.4
<i>acuminata</i> James, 1940, <i>Cyphomyia</i>	26.25
<i>acutirostris</i> Loew, 1863, <i>Nemotelus</i>	26.48
<i>acutus</i> James, 1971, <i>Merosargus</i>	26.10
<i>aczeli</i> James, 1953, <i>Cyphomyia</i>	26.25
<i>Adoxomyia</i> Kertész, 1907	26.30
<i>Adoxomyia</i> Bezzi, 1908	26.30
<i>adusta</i> Lindner, 1949, <i>Acrochaeta</i>	26.8
<i>adustus</i> Lindner, 1949, <i>Ptecticus</i>	26.17
<i>acnea</i> Lindner, 1949, <i>Chaetohermctia</i>	26.36
<i>aenea</i> Thomson, 1869, <i>Hadrestia</i>	26.6
<i>aeneipennis</i> Giglio-Tos, 1892, <i>Hermetia</i>	26.36
<i>acneiventris</i> (Giglio-Tos), 1892 (<i>Chrysonotus</i>), <i>Himantoloba</i>	26.10
<i>aeneus</i> (Lindner), 1949 (<i>Coenosargus</i>), <i>Merosargus</i>	26.10
<i>affinis</i> (Bellardi), 1859 (<i>Odonthomyia</i>), <i>Hedriodiscus</i>	26.40
<i>affinis</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.25
<i>affinis</i> Schiner, 1868, <i>Ptecticus</i>	26.17
<i>affinis</i> Williston, 1888, <i>Dicranophora</i>	26.21
<i>akrei</i> James & McFadden, 1971, <i>Merosargus</i>	26.10
<i>albicaput</i> (Walker), 1849 (<i>Clitellaria</i>), <i>Cyphomyia</i>	26.25
<i>albidus</i> McFadden, 1971, <i>Merosargus</i>	26.10
<i>albifacies</i> James, 1941, <i>Merosargus</i>	26.11
<i>albipes</i> James, 1939, <i>Chrysochlorina</i>	26.35
<i>albipes</i> (Townsend), 1903 (<i>Chrysochroma</i>), <i>Himantoloba</i>	26.9
<i>aibirostris</i> Macquart, 1850, <i>Nemotelus</i>	26.48
<i>aibitalis</i> Williston, 1896, <i>Pelagomyia</i>	26.36
<i>albitalus</i> , error, <i>Pelagomyia</i>	26.36
<i>aibitarsis</i> Fabricius, 1805, <i>Hermetia</i>	26.36
<i>aibitarsis</i> (Fabricius), 1805 (<i>Stratiomys</i>), <i>Cyphomyia</i>	26.25
<i>albitarsis</i> Fabricius of Wledemann, 1824, <i>Cyphomyia</i>	26.28
<i>albitarsis</i> Macquart, 1846, <i>Beris</i>	26.5
<i>albiventrts</i> Thomson, 1869, <i>Nemotelus</i>	26.48
<i>albomaculata</i> (Macquart), 1834 (<i>Odontomyia</i>), <i>Cyphomyia</i>	26.25
<i>albomaculata</i> Macquart, 1838, <i>Odontomyia</i>	26.43
<i>albopictus</i> James, 1971, <i>Merosargus</i>	26.11
<i>alchidas</i> Walker, 1849, <i>Sargus</i>	26.18
<i>aleus</i> (Walker), 1849 (<i>Acrochaeta</i>), <i>Hoplitomyia</i>	26.42
<i>alexanderi</i> Brèthes, 1922 (<i>Vappo</i>), <i>Eupachygaster</i>	26.54
<i>Allognosta</i> Osten Sacken, 1883	26.4
<i>Aloipha</i> Enderlein, 1914	26.10
<i>Aloiphina</i> Lindner, 1949	26.10
<i>aiticola</i> McFadden, 1971, <i>Merosargus</i>	26.11
<i>altifrons</i> James, 1939, <i>Cyphomyia</i>	26.25
<i>altifrons</i> James, 1971, <i>Merosargus</i>	26.11
<i>altifrons</i> Wulp, 1888, <i>Odontomyia</i>	26.43
<i>amazonicus</i> Kertész, 1914, <i>Dactylodictes</i>	26.53




amcnides (Walker), 1849 (Clitellaria), Artemita	26.50
amethystinus (Lindner), 1969 (Aloipha), Merosargus	26.11
amocnem Lindner, 1969, Anapospasma	26.30
Amphilecta Brauer, 1882	26.36
ampulla James, 1938, Hermetia	26.37
ampulla James, 1971, Merosargus	26.11
Analcoccroides Hollis, 1962	26.20
Analeocerus Loew, 1855	26.20
analis (Williston), 1888 (Chrysonotus), Himantoloba	26.9
Anapospasma Lindner, 1969	26.30
anchorata Bigot, 1879, Odontomyia	26.44
anchorata Bigot, 1879, Odontomyia	26.44
andinus James, 1971, Merosargus	26.11
<i>andreas</i> (Walker), 1849 (Sargus), Hermetia	26.38
androgyna Osten Sacken, 1886, Cyphomyia	26.25
angulatus McFadden, 1971, Merosargus	26.11
angusta Enderlein, 1914, Eumenogastrina	26.9
angustifrons James, 1967, Eidalimus	26.54
angustifrons Lindner, 1951, Charisina	26.52
angustus James, 1971, Merosargus	26.11
Anisoscopus McFadden, 1970	26.20
annulatus Kertész, 1914, Eidalimus	26.54
<i>annulatus</i> (Kertész), 1916 (Pyclomyia), Eidalimus	26.54
annulipes Enderlein, 1921, Oplaeantha	26.7
annulipes (Enderlein), 1921 (Chiromyza), Barbiellinia	26.1
Anopisthocrania Lindner, 1935	26.39
antennatus Sehiner, 1868, Merosargus	26.11
anthidium James, 1967, Hermetia	26.37
anticus Curran, 1932, Merosargus	26.11
Antissa Walker, 1854	26.23
Antissini	26.23
<i>Antissops</i> Enderlein, 1914	26.5
Aochletus Osten Sacken, 1886	26.32
apicalis Lindner, 1929, Chactohermetia	26.36
apicalis Lindner, 1935, Merosargus	26.11
<i>apicalis</i> Wiedemann, 1830, Hermetia	26.36
araneifera Sehiner, 1868, Odontomyia	26.44
Archistratiomys Enderlein, 1913	26.4
arcuatus James, 1971, Merosargus	26.11
argentea Osten Sacken, 1886, Artemita	26.50
argenteus James, 1967, Pareidalimus	26.57
armata (Wiedemann), 1830 (Sargus), Rhapsioeera	26.23
argyrosticta Kertész, 1914, Cosmariomyia	26.52
argyrostriata Lindner, 1964, Paracanthinomyia	26.57
Artemita Walker, 1854	26.50
astuta Williston, 1888, Dieranophora	26.21
aterrima James, 1940, Chordonota	26.24
aterrima Walker, 1856, Odontomyia	26.44
<i>atra</i> Wandolleck, 1897, Blastocera	26.51
atrata (Fabricius), 1805 (Stratiomys), Labocerina	26.29
atriannulatus McFadden, 1971, Merosargus	26.11
atriceps Loew, 1855, Analocerus	26.20
atrobasis (James), 1941 (Geosargus), Sargus	26.18
Aulocromyia Lindner, 1969	26.20



<i>aurata</i> , error, Meristocera	26.55
<i>aurata</i> Bellardi, 1861, Hermetia	26.37
<i>aurata</i> Bigot, 1879, Artemita	26.50
<i>aurata</i> Enderlein, 1914, Chalcidomorpha	26.51
<i>aurata</i> Macquart, 1846, Artemita	26.50
<i>aurea</i> Lindner, 1864, Meristocera	26.55
<i>aureonitens</i> James, 1971, Merosargus	26.11
<i>aureostriata</i> (Enderlein), 1914 (Pseggomma), Artemita	26.50
<i>aureus</i> Bellardi, 1859, Sargus	26.20
<i>auricoma</i> Enderlein, 1921, Exaeretina	26.6
<i>auriflamma</i> Wiedemann, 1819, Cyphomyia	26.25
<i>aurifrons</i> Wiedemann, 1830, Cyphomyia	26.25
<i>aurinotata</i> Lindner, 1935, Hermetia	26.37
<i>aurivena</i> James, 1971, Merosargus	26.11
<i>australis</i> Aubertin, 1930, Hylorus	26.3
<i>australis</i> James, 1971, Merosargus	26.11
<i>azureus</i> (Enderlein), 1914 (Coenosargus), Merosargus	26.11
<i>bagosus</i> Walker, 1849, Sargus	26.18
<i>bahamensis</i> (James), 1953 (Eulalia), Odontomyia	26.44
<i>banksi</i> James, 1936, Merosargus	26.11
<i>banksi</i> James, 1949, Cyphomyia	26.25
<i>banksi</i> James, 1971, Artemita	26.50
<i>barbata</i> Lindner, 1935, Chloromelas	26.40
<i>barbatus</i> James, 1971, Merosargus	26.11
<i>Barbiellinia</i> Bezzi, 1922	26.1
<i>barbiellinii</i> Bezzi, 1908, Berismyia	26.5
<i>basalis</i> James, 1971, Merosargus	26.11
<i>basalis</i> Kertész, 1914, Nematelus	26.48
<i>Basentidema</i> Macquart, 1838	26.21
<i>bassleri</i> James, 1943, Leucoptilum	26.33
<i>beameri</i> James, 1941, Merosargus	26.11
<i>beebei</i> Curran, 1934, Hermetia	26.37
<i>bellaardii</i> Giglio-Tos, 1891, Artemita	26.50
<i>bellula</i> (Williston), 1888 (Beris), Oplachantha	26.8
<i>Benhamyia</i> Miller, 1945	26.4
<i>bequaerti</i> Curran, 1925, Acanthinomyia	26.49
<i>bequaerti</i> Curran, 1928, Merosargus	26.11
<i>bequaerti</i> (Curran), 1932 (Rhingiopsis), Zuercheria	26.46
<i>bequaerti</i> James, 1938, Cyphomyia	26.28
Beridinae	26.4
<i>Beridops</i> Enderlein, 1913	26.5
<i>Beris</i> Latreille, 1802	26.5
<i>Berisargus</i> Lindner, 1933	26.47
<i>Berismyia</i> Giglio-Tos, 1891	26.5
<i>bezziana</i> Hide, 1966, Chrysochlorina	26.35
<i>bezzii</i> (Lindner), 1949 (Antissops), Berismyia	26.5
<i>bicarinata</i> Williston, 1900, Cyphomyia	26.25
<i>bicolor</i> Kertész, 1914, Artemita	26.50
<i>bicolor</i> (Wiedemann), 1830 (Sargus), Microchrysa	26.16
<i>billmecki</i> , error, Cynipimorpha	26.53
<i>billmecki</i> Brauer, 1882, Cynipimorpha	26.53
<i>bimaculata</i> Bellardi, 1862 (Stratiomys), Hoplitimyia	26.42
<i>bimaculata</i> Macquart, 1834, Hermctia	26.36




<i>bipartita</i> , error, <i>Rhaphiocera</i>	26.23
<i>bipartita</i> Wulp, 1885, <i>Caenacantha</i>	26.51
<i>bispinosus</i> (Wiedemann), 1830 (<i>Sargus</i>), <i>Hoplistes</i>	26.22
<i>bistriatus</i> Willston, 1896, <i>Euparyphus</i>	26.32
<i>bitaenlatus</i> Lindner, 1949, <i>Merosargus</i>	26.11
<i>bituberculatus</i> Lindner, 1935, <i>Merosargus</i> , misidentified	26.12
<i>bituberculatus</i> Schiner, 1868, <i>Merosargus</i>	26.20
<i>bivittatus</i> James, 1971, <i>Merosargus</i>	26.12
<i>Blastocera</i> Gerstaecker, 1857	26.51
<i>bonariensis</i> (Brèthes), 1922 (<i>Udamacantha</i>), <i>Labostigmia</i>	26.42
<i>Borboridea</i> Kertész, 1916	26.51
<i>borgmeieri</i> Lindner, 1931, <i>Rhaphiocera</i>	26.23
<i>borgmeieri</i> (Lindner), 1933 (<i>Berisargus</i>), <i>Nothomyia</i>	26.47
<i>brachiatatus</i> James, 1971, <i>Merosargus</i>	26.12
<i>Brachycara</i> Thomson, 1869	26.30
<i>Brachyodina</i> Lindner, 1949	26.51
<i>Brachythrix</i> McFadden, 1970	26.21
<i>brasilliana</i> Lindner, 1935, <i>Glarlopsis</i>	26.32
<i>brasilliana</i> Lindner, 1949, <i>Neoberis</i>	26.7
<i>brasilliana</i> Lindner, 1964, <i>Artemita</i>	26.50
<i>brasillensis</i> Enderlein, 1921, <i>Archistratiomys</i>	26.4
<i>brasiliensis</i> Lindner, 1949, <i>Euparyphus</i>	26.32
<i>brasiliensis</i> Lindner, 1935, <i>Neochorisops</i>	26.7
<i>brasiliensis</i> Wiedemann, 1830, <i>Sargus</i>	26.18
<i>braueri</i> Willston, 1888, <i>Myxosargus</i>	26.46
<i>brevicornis</i> Enderlein, 1921, <i>Nonacris</i>	26.3
<i>brevicornis</i> (Lindner), 1949 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>brevicornis</i> Lindner, 1964, <i>Chorophthalmyla</i>	26.52
<i>brevifacies</i> (Macquart), 1855 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.40
<i>brevifurca</i> James, 1943, <i>Dicranophora</i>	26.21
<i>brevis</i> James, 1940, <i>Cyphomyia</i>	26.26
<i>brevisetata</i> Lindner, 1964, <i>Panacris</i>	26.56
<i>brevisetata</i> Walker, 1854, <i>Acrochaeta</i>	26.8
<i>bruesi</i> Johnson, 1919, <i>Microchrysa</i>	26.16
<i>brunneipicta</i> James, 1953, <i>Chloromelas</i>	26.40
<i>brunneus</i> Lindner, 1933, <i>Merosargus</i>	26.12
<i>brunnipes</i> McFadden, 1971, <i>Merosargus</i>	26.12
<i>bucephala</i> (Gerstaecker), 1857 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>bulbifrons</i> Willston, 1900, <i>Merosargus</i>	26.12
<i>burmeisteri</i> Lindner, 1949, <i>Cyphomyia</i>	26.28
<i>burmeisteri</i> Lindner, 1949, <i>Hermetia</i>	26.37
<i>c-nigrum</i> (Lindner), 1951 (<i>Pedicella</i>), <i>Merosargus</i>	26.12
<i>Cacosis</i> Walker, 1856	26.34
<i>Caetobia</i> James, 1966	26.51
<i>Caenacantha</i> Wulp, 1885	26.51
<i>caerulea</i> Schiner, 1868, <i>Beris</i>	26.5
<i>caesius</i> Bellardi, 1859, <i>Sargus</i>	26.18
<i>calceolatus</i> Bigot, 1879, <i>Merosargus</i>	26.12
<i>callifera</i> Lindner, 1928, <i>Hermetia</i>	26.37
<i>callosa</i> James, 1937, <i>Dicuryneura</i>	26.31
<i>callosa</i> James, 1967, <i>Psephiocera</i>	26.58
<i>Caloparyphus</i> James, 1939	26.30
<i>caloptera</i> Osten Sacken, 1886, <i>Rhaphiocera</i>	26.23



<i>calopus</i> Loew, 1869, <i>Nothomyia</i>	26.47
<i>carbonaria</i> (Beilardi), 1862 (<i>Chordonota</i>), <i>Laboccrina</i>	26.29
<i>carbonarius</i> Giglio-Tos, 1891, <i>Euparyphus</i>	26.31
<i>castanca</i> (Macquart), 1838 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>catharistis</i> Lindner, 1926, <i>Nemotelus</i>	26.48
<i>catharistis</i> Lindner, 1930, <i>Nemotelus</i>	26.48
<i>cazieri</i> (James), 1953 (<i>Pachygaster</i>), <i>Peltina</i>	26.57
<i>Cephalochrysa</i> Kertész, 1912	26.9
<i>centor</i> Curran, 1934, <i>Acanthinomyia</i>	26.40
<i>cercuceus</i> Williston, 1936, <i>Merosargus</i>	26.12
<i>ceria</i> Williston, 1900, <i>Hermetia</i>	26.37
<i>ceriogaster</i> Williston, 1888, <i>Hermetia</i>	26.37
<i>Chaetohermetia</i> Lindner, 1929	26.36
<i>Chaetosargus</i> Roeder, 1894	26.34
<i>Chalcidomorphina</i> Enderlein, 1914	26.51
<i>Chalcomorphina</i> , error	26.51
<i>chaiconota</i> (Brauer), 1882 (<i>Composoma</i>), <i>Acrochaeta</i>	26.8
<i>chalybea</i> (Wiedemann), 1924 (<i>Clitellaria</i>), <i>Cyphomyia</i>	26.26
<i>Charisina</i> Lindner, 1951	26.52
<i>Chauna</i> Loew, 1847	26.56
<i>chilensis</i> Lindner, 1943, <i>Nonacris</i>	26.3
<i>Chilota</i> Aubertin, 1930	26.5
<i>chiquitanus</i> Lindner, 1930, <i>Meringostylus</i>	26.33
<i>chiriquina</i> Lindner, 1935, <i>Cyphomyia</i>	26.25
<i>Chiromyza</i> Authors	26.1
<i>Chiromyza</i> Wiedemann, 1820	26.2
<i>Chiromyzinae</i>	26.1
<i>Chlamydonotum</i> Lindner, 1949	26.52
<i>chioraspis</i> (Wiedemann), 1830 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.40
<i>Chioromcias</i> Enderlein, 1914	26.40
<i>chiorovittatus</i> McFadden, 1970, <i>Anisoscopus</i>	26.20
<i>Chordonota</i> Gerstaecker, 1857	26.24
<i>Chorophthalmomyia</i> Lindner, 1964	26.52
<i>Chorophthalmus</i> , error	26.52
<i>Chrysochlora</i> Authors	26.34
<i>Chrysochlorina</i> James, 1939	26.34
<i>Chrysochlorinae</i>	26.34
<i>Chrysochroma</i> Authors	26.9
<i>chrysocomus</i> (James), 1941 (<i>Geosargus</i>), <i>Sargus</i>	26.18
<i>chrysodota</i> Perty, 1833, <i>Cyphomyia</i>	26.26
<i>Chrosonotomyia</i> Authors	26.9
<i>Chrysonotus</i> Authors	26.9
<i>chrysopila</i> Loew, 1872, <i>Hermetia</i>	26.37
<i>ciliatus</i> James, 1941, <i>Pteticus</i>	26.17
<i>cincticornis</i> Enderlein, 1921, <i>Opiachantha</i>	26.7
<i>cinctus</i> Osten Sacken, 1886, <i>Euparyphus</i>	26.32
<i>cinguiata</i> Greene, 1940, <i>Hermetia</i>	26.37
<i>cinguiatus</i> Schiner, 1868, <i>Merosargus</i>	26.12
<i>circumcinctus</i> (James), 1941 (<i>Geosargus</i>), <i>Sargus</i>	26.19
<i>citrinellus</i> Lindner, 1949, <i>Merosargus</i>	26.12
<i>citrinus</i> James, 1971, <i>Merosargus</i>	26.12
<i>ciaripennis</i> Macquart, 1947, <i>Cyphomyia</i>	26.26
<i>Ciarissa</i> Lindner, 1964	26.52
<i>ciavata</i> James, 1939, <i>Hoplitimyia</i>	26.42




clavatus Walker, 1854, Sargus	26.19
elaviventris Rondani, 1850, Sargus	26.19
<i>Clitellaria</i> Authors	26.30
Clitellariinae	26.23
Clitellariini	26.29
elypeata (Bigot), 1879 (Odontomyia), Labostigmina	26.42
coarctata Macquart, 1846, Hermetia	26.37
coarctatus Macquart, 1838, Sargus	26.19
coati Lindner, 1951	26.40
<i>Coenosargus</i> Enderlein, 1914	26.10
coerulea Macquart, 1834, Cyphomyia	26.25
coeruloseens James, 1943, Basentidema	26.21
cognata (Lindner), 1949 (Pyclomyia), Psephiocera	26.58
colombiana Kertész, 1916, Ligyromyia	26.55
columbianus Enderlein, 1914, Lysozus	26.22
comma (Williston), 1900 (?Odontomyia), Labostigmina	26.42
complicatus James, 1971, Merosargus	26.12
<i>Compsosoma</i> Brauer, 1882	26.8
comstocki Williston, 1885, Hermetia	26.37
concinna Williston, 1900, Hermetia	26.37
concinna (Williston), 1900 (Odontomyia), Labostigmina	26.42
concinna Williston, 1900, Merosargus	26.12
concinna Osten Saeken, 1886, Sargus	26.19
concinna Williston, 1900, Ptecticus	26.17
condor Lindner, 1951, Hermetia	26.37
confusa James, 1936, Odontomyia	26.44
congruens Kertész, 1914, Nematelus	26.48
conica Fabricius, 1805, Stratiomys	26.45
conjuncta James, 1967, Hermetia	26.37
conneza, error, Stratiomys	26.45
conopsoides James, 1971, Merosargus	26.12
conopsoides Sehiner, 1868, Ptecticus	26.17
consona Lindner, 1964, Nitrogramma	26.56
constricta Walker, 1860, Stratiomys	26.45
contigua James, 1967, Myctrocera	26.56
contortus James, 1971, Merosargus	26.12
contractus Walker, 1854, Sargus	26.19
convexa (Walker), 1854 (Acanthina), Artemita	26.50
convexa Wulp, 1881, Stratiomys	26.45
convexifrons McFadden, 1971, Merosargus	26.12
coprates (Walker), 1849 (Clitellaria), Cyphomyia	26.26
cordobacensis (Lindner), 1951 (Hedriodiseina), Chloromelas	26.40
coriaceus Giglio-Tos, 1891, Merosargus	26.12
cornithorax (Lindner), 1928 (Aerodesmia), Hermetia	26.37
cornuta Kraft & Cook, 1961, Zabrachia	26.60
Cosmariomyia Kertész, 1914	26.52
costalimai Iide, 1967, Chrysochlorina	26.35
costalis Walker, 1837, Cyphomyia	26.26
coxalis Lindner, 1949, Merosargus	26.12
erabro Osten Saeken, 1886, Hermetia	26.37
eruciana Lindner, 1951, Rhapsiocera	26.23
cruciata (Macquart), 1838 (Odontomyia), Labostigmina	26.43
cupraria Enderlein, 1914, Eumenogastrina	26.9
euprina (Wiedemann), 1830 (Stratiomys), Chloromelas	26.40




currani Iide, 1966, Chrysochlorina	26.35
currani (James), 1932 (Odontomyia), Hedriodiscus	26.40
currani Lindner, 1949, Hermetia	26.38
<i>Cyanauges</i> Philippi, 1865	26.24
<i>Cyanauges</i> Rondani, 1863	26.24
<i>cyanea</i> Macquart, 1834, Cyphomyia	26.25
<i>cyanea</i> (Fabricius), 1794 (Stratiomys), Cyphomyia	26.26
<i>cyaneoscutellatus</i> (Enderlein), 1914 (Aloipha), Merosargus	26.12
<i>cyanocephala</i> Gerstaecker, 1857, Cyphomyia	26.28
<i>cyanifrons</i> (Rondani), 1848 (Sargus), Ptecticus	26.17
<i>cyanispinis</i> Bigot, 1875, Cyphomyia	26.25
<i>Cyclophleps</i> James, 1943	26.21
<i>cylindricornis</i> Williston, 1888, Promeranisa	26.45
<i>Cyclotaspis</i> Lindner, 1864	26.52
<i>Cynipomorpha</i> Brauer, 1882	26.53
<i>Cynipomorpha</i> , emendation	26.53
<i>Cyphomyia</i> Wiedemann, 1819	26.25
<i>Cyphomyiini</i>	26.24
<i>cyphomyioides</i> Kertész, 1909, Hypselophrum	26.55
<i>cyphomyioides</i> Lindner, 1933, Pataglomys	26.39
<i>cyrtonetopus</i> James, 1971, Merosargus	26.12
<i>Dactylacantha</i> Lindner, 1964	26.53
<i>Dactylodeictes</i> Kertész, 1914	26.48
<i>dampfii</i> James, 1941, Nematelus	26.48
<i>decemmaculatus</i> (Osten Sacken), 1886 (?Euparyphus), Caloparyphus	26.31
<i>degenerata</i> Lindner, 1949, Acrochaeta	26.8
<i>dentatum</i> James, 1967, Chlamydonotum	26.52
<i>denticulata</i> (Enderlein), 1914 (Antlssops), Berlsmyia	26.5
<i>depressus</i> James, 1967, Dactylodeictes	26.53
<i>despectus</i> Walker, 1856 (Inopus)	26.3
<i>Diaphorostylus</i> Kertész, 1908	26.31
<i>Diastrophthalmus</i> Lindner, 1949	26.53
<i>dichroa</i> Kertész, 1916, Thopomyia	26.59
<i>dichroa</i> (Schiner), 1868 (Chrysonotus), Himantoloba	26.10
<i>Dicranophora</i> Macquart, 1834	26.21
<i>Dicyphoma</i> James, 1937	26.29
<i>diehli</i> Lindner, 1941, Nematelus	26.48
<i>Dieuryneura</i> James, 1937	26.31
<i>digitata</i> Lindner, 1964, Synaptochaeta	26.59
<i>digitata</i> James, 1966, Brachycara	26.30
<i>dileucostigma</i> McFadden, 1970, Brachytrix	26.21
<i>dimidiata</i> Lindner, 1949, Acrochaeta	26.8
<i>discolorata</i> James, 1936, Odontomyia	26.44
<i>disjunctus</i> Curran, 1925, Nematelus	26.48
<i>dispar</i> Schiner, 1868, Cyphomyia	26.26
<i>dispar</i> Schiner, 1868, Microchrysa	26.17
<i>dissimilis</i> Bellardi, 1859, Odontomyia	26.44
<i>dissimilis</i> Giglio-Tos, 1891, Merosargus	26.12
<i>Ditylometopa</i> Kertész, 1923	26.31
<i>diversicornis</i> Kertész, 1923, Acropeltates	26.29
<i>divisus</i> James, 1971, Merosargus	26.12



<i>Dolichothrix</i> McFadden, 1970	26.21
<i>dominicana</i> James, 1967, <i>Cyphomyia</i>	26.26
<i>dorsalis</i> (Fabricius), 1805 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.40
<i>dorsalis</i> Lindner, 1969, <i>Merosargus</i>	26.13
<i>Drasteria</i> Brauer, 1882	26.34
<i>Draymonia</i> Aubertin, 1932	26.5, 26.6
<i>dubia</i> Brauer, 1882, <i>Melanochroa</i>	26.46
<i>dubia</i> Curran, 1932 (<i>Pelagomyia</i>), <i>Chrysochlorina</i>	26.35
<i>dubia</i> Röder, 1886, <i>Melanochroa</i>	26.46
<i>Ecehactomyia</i> Lindner, 1949	26.53
<i>echemon</i> Walker, 1849, <i>Acrochaeta</i>	26.8
<i>ecuadorensis</i> Enderlein, 1914, <i>Cyphomyia</i>	26.26
<i>ecuadorensis</i> James & McFadden, 1971, <i>Merosargus</i>	26.14
<i>ecuadorianus</i> Lindner, 1949, <i>Heptozus</i>	26.21
<i>Eicochalcidina</i> Lindner, 1964	26.53
<i>Eidalimus</i> Kertész, 1914	26.54
<i>eiseni</i> Townsend, 1895, <i>Hermetia</i>	26.38
<i>elatus</i> Curran, 1932, <i>Merosargus</i>	26.13
<i>elegans</i> Kertész, 1923, <i>Ditylometopa</i>	26.31
<i>elegans</i> (Macquart), 1838 (<i>Odontomyia</i>), <i>Labostigmina</i>	26.43
<i>elegans</i> Perty, 1833, <i>Acrochaeta</i>	26.8
<i>elegans</i> Schiner, 1863, <i>Actina</i>	26.4
<i>elegans</i> (Wiedemann), 1830 (<i>Cyphomyia</i>), <i>Euparyphus</i>	26.31
<i>elegans</i> Williston, 1888, <i>Euryneura</i>	26.32
<i>elongata</i> Aubertin, 1930, <i>Macromeracis</i>	26.6
<i>elongata</i> (Wiedemann), 1824 (<i>Clitellaria</i>), <i>Acanthinomyia</i>	26.49
<i>elongatulus</i> Williston, 1900, <i>Euparyphus</i>	26.31
<i>elongatus</i> James, 1971, <i>Merosargus</i>	26.13
<i>emarginata</i> Macquart, 1838, <i>Odontomyia</i>	26.44
<i>enderleini</i> Lindner, 1926, <i>Rhingiopsis</i>	26.45
<i>enderleini</i> (Lindner), 1949 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>engeli</i> (Lindner), 1943 (<i>Huttonella</i>), <i>Benhamyia</i>	26.4
<i>Engicerus</i> Lindner, 1964	26.54
<i>ereeta</i> McFadden, 1969, <i>Cyphomyia</i>	26.26
<i>erectispinis</i> Lindner, 1935, <i>Cyphomyia</i>	26.26
<i>ethelia</i> Curran, 1932, <i>Merosargus</i>	26.13
<i>euchlorus</i> (Gerstaecker), 1857 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.40
<i>Euclitellaria</i> Kertész, 1923	26.30
<i>Eycynipimorpha</i> Malloch, 1915	26.54
<i>Eulalia</i> Meigen, 1800	26.43
<i>Eumenogastrina</i> Enderlein, 1914	26.9
<i>eunomus</i> James, 1967, <i>Merosargus</i>	26.13
<i>Eupachygaster</i> Kertész, 1911	26.54
<i>Euparhyphus</i> , error or emendation	26.31
<i>Euparyphus</i> Gerstaecker, 1857	26.31
<i>eupodata</i> (Bigot), 1879 (<i>Exaireta</i>), <i>Neoexaireta</i>	26.7
<i>Euryneura</i> Schiner, 1867	26.32
<i>Euryneurasoma</i> Johnson, 1913	26.30
<i>Exaereta</i> , emend.	26.7
<i>Exaeretina</i> Enderlein, 1921	26.6
<i>Exaireta</i> Schiner, 1867	26.7
<i>Exairete</i> , error	26.7
<i>Exodontha</i> Rondani, 1856	26.23



<i>falieni</i> (Perty), 1833 (Sargus), Rhaphiocera	26.23
<i>fasciata</i> (Fabricius), 1787 (Stratiomys), Hoplitomyia	26.42
<i>fasciata</i> (Fabricius), 1805 (Stratiomys), Euryneura	26.32
<i>fasciata</i> Kertész, 1914, Neoacanthina	26.56
<i>fasciata</i> Kertész, 1923, Aeropeltates	26.29
<i>fasciata</i> Macquart, 1834, Odontomyia	26.44
<i>fasciata</i> (Thomson), 1867 (Chrysochlora), Chrysochlorina	26.35
<i>fasciata</i> Wiedemann, 1830, Aerochaeta	26.9
<i>fasciatifrons</i> Lyneh Arribáizaga, 1881, Nematelus	26.48
<i>fasciatipennis</i> (Lindner), 1935 (Ecrisargus), Nothomyia	26.47
<i>fasciatus</i> Brauer, 1882, Myxosargus	26.46
<i>fasciatus</i> Fabricius, 1805, Sargus	26.19
<i>fascifrons</i> Macquart, 1849, Odontomyia	26.44
<i>fascipennis</i> (Fabricius), 1805 (Stratiomys), Euryneura	26.32
<i>fascipennis</i> James, 1943, Histiodroma	26.22
<i>fascipennis</i> Lindner, 1949, Rhaphiocera	26.23
<i>fascipes</i> Walker, 1854, Cyphomyia	26.26
<i>fassli</i> Lindner, 1949, Cyphomyia	26.26
<i>femorialis</i> (Curran), 1929 (Chrysochlora), Chrysochlorina	26.35
<i>femorialis</i> Kertész, 1909, Popanomyia	26.58
<i>femorata</i> (Bellardi), 1859 (Odonthomyia), Labostigmina	26.43
<i>fenestrata</i> Gerstaecker, 1857, Stratiomys	26.46
<i>fenestrata</i> Macquart, 1846, Cyphomyia	26.25
<i>fenestrata</i> (Macquart), 1846 (Ephippium), Adoxomyia	26.30
<i>fenestrata</i> (Thomson), 1869 (Odontomyia), Labostigmina	26.43
<i>ferruginea</i> Enderlein, 1914, Cyphomyia	26.26
<i>ferruginea</i> (Gerstaecker), 1857 (Chauna), Neochauna	26.56
<i>festiva</i> (Walker), 1854 (Stratiomys), Zuercheria	26.46
<i>festivus</i> Williston, 1888, Merosargus	26.13
<i>fiuinius</i> Lindner, 1940, Ptecticus	26.17
<i>figlius</i> Osten Sacken, 1886, Ptecticus	26.17
<i>fitiformis</i> Giglio-Tos, 1891, Sargus	26.18
<i>fimbriata</i> (Lindner), 1931 (Aerodesmia), Hermetia	26.38
<i>flaveola</i> Bigot, 1879 (Histiodroma), Acrochaeta	26.9
<i>fiavescens</i> (James), 1937 (Chrysochlora), Chrysochlorina	26.35
<i>fiaviceps</i> Bigot, 1879, Ptecticus	26.17
<i>fiaviceps</i> McFadden, 1971, Merosargus	26.13
<i>fiavicornis</i> Johnson, 1894, Nematelus	26.48
<i>fiavicornis</i> Lindner, 1969, Eidaimus	26.54
<i>fiavida</i> James, 1937, Platopsomyia	26.34
<i>flavifasciatus</i> (Macquart), 1849 (Odontomyia), Hedriodiscus	26.41
<i>flaviatus</i> James, nom. nov., Sargus	26.19
<i>flavimana</i> Gerstaecker, 1857, Cyphomyia	26.26
<i>flavimanus</i> Lindner, 1949, Diastrophthaimus	26.53
<i>flavipennis</i> Enderlein, 1914, Cyphomyia	26.26
<i>flavipes</i> Enderlein, 1914, Psephiocera	26.58
<i>flavipes</i> James, 1967, Manotes	26.55
<i>flavipes</i> Kertész, 1908, Diaphorostylus	26.31
<i>flavipes</i> Lindner, 1933, Hermetia	26.38
<i>flavipes</i> Kertész, 1908 (Euryneura), Meringostylus	26.33
<i>flavipes</i> Wiedemann, 1830, Hermetia	26.38
<i>flavispinis</i> Macquart, 1855, Cyphomyia	26.28
<i>flavispinosa</i> Macquart, 1855, Beris	26.5
<i>fiavissimus</i> James, 1971, Merosargus	26.13
<i>fiavitaris</i> (Lindner), 1969 (Aloipha), Merosargus	26.13



<i>flavitaris</i> Enderlein, 1914 (<i>Lasiopa</i>)	26.29
<i>flaviventris</i> (Lindner), 1935 (<i>Chrysochroma</i>), <i>Himantoloba</i>	26.10
<i>flaviventris</i> Lindner, 1964, <i>Trigonocerina</i>	26.59
<i>flavoecingulatus</i> Kertész, 1914, <i>Nemotelus</i>	26.48
<i>fiavonigra</i> (Lindner), 1928 (<i>Sargus</i>), <i>Himantoloba</i>	26.10
<i>flavopilosa</i> (Bigot), 1879 (<i>Chrysonotus</i>), <i>Himantoloba</i>	26.10
<i>flavoscutata</i> Bigot, 1879, <i>Hermetia</i>	26.38
<i>formica</i> Osten Sacken, 1886, <i>Hermetia</i>	26.38
<i>formicaeformis</i> Kertész, 1909, <i>Lyprotomyia</i>	26.55
<i>formosa</i> Enderlein, 1921, <i>Oplachantha</i>	26.7
<i>formosa</i> James, 1940, <i>Cyphomyia</i>	26.26
<i>foveifrons</i> Thomson, 1869, <i>Odontomyia</i>	26.44
<i>fratella</i> (Williston), 1900 (<i>Odontomyia</i>), <i>Labostigmia</i>	26.43
<i>fraternus</i> Bigot, 1879, <i>Merosargus</i>	26.13
<i>frontatus</i> Schiner, 1868, <i>Merosargus</i>	26.13
<i>frosti</i> James, 1939, <i>Chrysochlorina</i>	26.35
<i>frosti</i> James, 1941, <i>Merosargus</i>	26.13
<i>fuliginosa</i> Lindner, 1964, <i>Peltina</i>	26.57
<i>fuliginosa</i> Lindner, 1969, <i>Peltina</i>	26.57
<i>fulva</i> Walker, 1854, <i>Hermetia</i>	26.38
<i>fulvithorax</i> (Bigot), 1879 (<i>Chrysonotus</i>), <i>Himantoloba</i>	26.10
<i>fumipennis</i> James, 1971, <i>Merosargus</i>	26.13
<i>furcifera</i> (Wiedemann), 1824 (<i>Sargus</i>), <i>Dicranophora</i>	26.21
<i>fusca</i> Giglio-Tos, 1892, <i>Berismyia</i>	26.5
<i>fuscana</i> Wiedemann, 1821, <i>Chiromyza</i>	26.2
<i>fuscipennis</i> Bellardi, 1862, <i>Chordonota</i>	26.24
<i>geminatus</i> James, 1971, <i>Merosargus</i>	26.13
<i>geniculata</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.26
<i>geniculata</i> Lindner, 1936, <i>Chordonota</i>	26.24
<i>geniculata</i> Macquart, 1855, <i>Hermetia</i>	26.38
<i>Geosargus</i> Bezzi, 1907	26.18
<i>gerstaeckeri</i> (Bellardi), 1859 (<i>Stratiomys</i>), <i>Hoplitimyia</i>	26.42
<i>gigliotosi</i> McFadden, 1971, <i>Ptecticus</i>	26.17
<i>glaber</i> Loew, 1872, <i>Nemotelus</i>	26.48
<i>Glaripsis</i> Lindner, 1935	26.32
<i>Glaris</i> Kertész, 1923	26.33
<i>golbachii</i> James, 1953, <i>Cyphomyia</i>	26.26
<i>golbachii</i> James, 1967, <i>Eicochalcidina</i>	26.53
<i>golbachii</i> James, 1971, <i>Merosargus</i>	26.13
<i>goncalvesi</i> Albuquerque, 1955, <i>Hermetia</i>	26.38
<i>goniophora</i> Say, 1829, <i>Stratiomys</i>	26.46
<i>gorgona</i> Lindner, 1949, <i>Aerochaeta</i>	26.9
<i>gorgonea</i> Lindner, 1949, <i>Chloromelas</i>	26.40
<i>Gowdeyana</i> Curran, 1928	26.54
<i>gowdeyi</i> Curran, 1928, <i>Merosargus</i>	26.13
<i>gracilicornis</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.26
<i>gracillor</i> James, 1971, <i>Merosargus</i>	26.13
<i>gracilis</i> James, 1936, <i>Monacanthella</i>	26.33
<i>gracilis</i> Kertész, 1923, <i>Grypomyia</i>	26.33
<i>gracilis</i> Williston, 1888, <i>Merosargus</i>	26.13
<i>grandis</i> Walker, 1856, <i>Cacosis</i>	26.34
<i>granulosus</i> James, 1971, <i>Merosargus</i>	26.13
<i>gryphus</i> Enderlein, 1914, <i>Myxosargus</i>	26.47



Grypomyia Kertész, 1923	26.33
guatemalae James, 1942, Myxosargus	26.47
guerinii Macquart, 1846, Beris	26.5
gulanae (Lindner), 1949 (Euiatia), Labostigmina	26.43
gulanae Lindner, 1949, Pezodontina	26.22
<i>Gyneuryptaria</i> Enderlein, 1914	26.25
Hadrestia Thomson, 1869	26.6
<i>halala</i> (Walker), 1849 (Citeiaria), Artemita.....	26.50
hamifer James, 1940, Euparyphus	26.32
hansae Lindner, 1931, Rhaphiocera	26.23
hansoni James, 1966, Heptozus	26.22
hansoni James, 1971, Merosargus	26.13
haterius Walker, 1849, Acrochaeta	26.9
<i>Hedriodiscina</i> Enderlein, 1914	26.40
Hedriodiscus Enderlein, 1914.....	26.40
helicivipennis Enderlein, 1914, Cyphomyia	26.26
Hemimegacis Enderlein, 1921	26.6
hennigi Lindner, 1969, Pachyptilium	26.33
henshawi (Malloch), 1917 (Eupachygaster), Eidalimus	26.54
Heptozus Lindner, 1949	26.21
Hermetia Latreille, 1804	26.36
Hermetiinae	26.36
Heteracanthia Macquart, 1849	26.6
<i>heteroneura</i> (Macquart), 1838 (Odontomyia), Chloromelas	26.40
<i>Hexacraspis</i> Enderlein, 1914	26.54
hieroglyphica (Wiedemann), 1830 (Acanthina), Artemita	26.50
Himantoloba McFadden, 1970	26.9
Himantomyia James, 1939	26.42
hirta Bezzi, 1922, Barbiellinia	26.2
hirticornis (Wiedemann), 1830 (Sargus), Chactosargus	26.34
Histiodroma Schiner, 1868	26.22
histrion (Walker), 1851 (Odontomyia), Chloromelas	26.40
hoffmanni Lindner, 1935, Merosargus	26.13
Homalarthria Lindner, 1933	26.33
<i>Hoplacantha</i> , emend.	26.7
Hoplites Macquart, 1834	26.22
hoplites (Wiedemann), 1830 (Sargus), Rhaphiocera	26.23
Hoplitiomyia James, 1934	26.42
hortulana (Wiedemann), 1830 (Sargus), Basentidema	26.21
hortulanus (Williston), 1900 (Analocerus), Neanaicocerus	26.22
humilis (Lindner), 1929 (Hedriodiscina), Hedriodiscus	26.41
hunteri Coquillett, 1909, Hermetia	26.38
<i>Huttonella</i> Enderlein, 1921	26.4
hyacinthina (Bigot), 1879 (Exaireta), Neocxaereta	26.7
hyalina James, 1967, Manotes	26.55
hyaiopterus Giglio-Tos, 1891, Merosargus	26.13
hybrida Gerstaecker, 1857, Cyphomyia	26.27
hydroiopus Lindner, 1849 (Hedriodiscina), Hedriodiscus	26.41
Hylorops Enderlein, 1921	26.2
Hylorus Philippi, 1865	26.2
<i>Hypocerina</i> Bigot, 1879	26.50
Hypselephrum Kertész, 1909	26.55



<i>ieae</i> (Lindner), 1941 (Eulalia), <i>Odontomyia</i>	26.44
<i>illucens</i> James, 1967, <i>Pelagomyia</i>	26.36
<i>illucens</i> (Linnaeus), 1758 (Musea), <i>Hermetia</i>	26.38
<i>illuminatus</i> (Lindner), 1948 (Geosargus), <i>Sargus</i>	26.19
<i>illustris</i> Sehiner, 1868, <i>Pteetieus</i>	26.17
<i>imitans</i> Curran, 1925, <i>Cyphomyia</i>	26.27
<i>imprensa</i> James, 1967, <i>Hermetia</i>	26.38
<i>inaequalis</i> Loew, 1866, <i>Odontomyia</i>	26.44
<i>ineanus</i> James & McFadden, 1971, <i>Merosargus</i>	26.14
<i>incompleta</i> (Curran), 1929 (Chrysochlora), <i>Chrysochlorina</i>	26.35
<i>inermis</i> (Wiedemann), 1830 (Cyphomyia), <i>Chordonota</i>	26.24
<i>inermis</i> (Wiedemann), 1830 (Sargus), <i>Hlstiodroma</i>	26.22
<i>inermis</i> (Wiedemann), 1830 (Stratiomys), <i>Labostigmina</i>	26.43
<i>infrapallidus</i> (Lindner), 1951 (Hedriodiseina), <i>Hedriodiseus</i>	26.41
<i>inops</i> Lindner, 1949, <i>Pteetieus</i>	26.17
<i>inornata</i> Lindner, 1964, <i>Cyelotaspis</i>	26.52
<i>inornata</i> (Williston), 1900 (Acanthina), <i>Artemita</i>	26.50
<i>insularis</i> James, 1966, <i>Daetylodeietes</i>	26.53
<i>intensica</i> (Curran), 1928 (Merosargus), <i>Nothomyia</i>	26.47
<i>interrupta</i> Enderlein, 1921, <i>Mesomyza</i>	26.3
<i>interruptus</i> James, 1939, <i>Diaphorostylus</i>	26.31
<i>inversus</i> Curran, 1934, <i>Pteetieus</i>	26.17
<i>irwini</i> James and McFadden, 1971, <i>Merosargus</i>	26.13
<i>Isosargus</i> James, 1935	26.9
<i>itatiaiensis</i> Lindner, nom. nov., <i>Hermetia</i>	26.38
<i>itatiayensis</i> Lindner, 1928, <i>Archistratiomys</i>	26.4
<i>iubata</i> Enderlein, 1921, <i>Oplachantha</i>	26.8
<i>jacarellus</i> (Lindner), 1949 (Hedriodiseina), <i>Hedriodiseus</i>	26.41
<i>jamesi</i> Barretto, 1947, <i>Rhingiopsis</i>	26.45
<i>jamesi</i> Lindner, 1949, <i>Cyphomyia</i>	26.27
<i>jamesi</i> Lindner, 1964, <i>Tegoeera</i>	26.59
<i>jamesi</i> (Lindner), 1969 (Sargus), <i>Himantoloba</i>	26.10
<i>jucundus</i> Walker, 1850, <i>Sargus</i>	26.19
<i>kansensis</i> , ssp. of	26.48
<i>kerteszi</i> Iide, 1968, <i>Euryneura</i>	26.32
<i>kerteszi</i> Szilády, 1929, <i>Aeropeltates</i>	26.29
<i>krausei</i> Philippi, 1865, <i>Hylorus</i>	26.3
<i>Labocerino</i> , error	26.29
<i>Labocerina</i> Enderlein, 1914	26.29
<i>Labocerino</i> , error	26.29
<i>Labostigmina</i> Enderlein, 1930	26.42
<i>laerimosus</i> James, 1971, <i>Merosargus</i>	26.13
<i>Lagarinus</i> Enderlein, 1913	26.3
<i>Lagarus</i> Philippi, 1865	26.3
<i>lampronotus</i> James, 1941, <i>Merosargus</i>	26.13
<i>lanei</i> Barretto, 1947, <i>Rhingiopsis</i>	26.45
<i>lanei</i> James, 1941, <i>Pteetieus</i>	26.17
<i>laniger</i> James, 1971, <i>Merosargus</i>	26.14
<i>lanuginosa</i> Enderlein, 1921, <i>Oplachantha</i>	26.8
<i>lanuginosa</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.26



<i>Lasiodeictes</i> Lindner, 1964	26.55
<i>lasiophthalma</i> Williston, 1896, <i>Cyphomyia</i>	26.27
<i>lasiophthalma</i> Williston, 1900, <i>Cyphomyia</i>	26.28
<i>lateralis</i> (James), 1936 (<i>Isosargus</i>), <i>Sargus</i>	26.19
<i>lateralis</i> Macquart, 1934, <i>Sargus</i>	26.19
<i>lateromaculatus</i> James, 1971, <i>Merosargus</i>	26.14
<i>laticornis</i> James, 1967, <i>Meristocera</i>	26.56
<i>latifrons</i> Enderlein, 1921, <i>Oplachantha</i>	26.8
<i>latifrons</i> James, 1971, <i>Artemita</i>	26.50
<i>latifrons</i> (Williston), 1900 (<i>Chrysochroma</i>), <i>Microchrysa</i>	26.17
<i>latimanus</i> James, 1967, <i>Manotes</i>	26.55
<i>lativentris</i> Bellardi, 1859, <i>Hermetia</i>	26.38
<i>latus</i> Bellardi, 1859, <i>Sargus</i>	26.19
<i>lefeburei</i> Authors, <i>Hedriodiscus</i>	26.41
<i>lefeburei</i> (Macquart), 1838 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>leciophthalma</i> Williston, 1896, <i>Chordonota</i>	26.24
<i>leptiformis</i> (Macquart), 1838 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>leucocephala</i> Hoffmansegg, 1819, <i>Cyphomyia</i>	26.27
<i>leucogaster</i> (James), 1933 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>leucopa</i> (Linnaeus), 1767 (<i>Musca</i>), <i>Hermetia</i>	26.38
<i>leucopis</i> Wiedemann, 1830, <i>Stratiomys</i>	26.46
<i>Leucoptilum</i> James, 1943	26.33
<i>liburna</i> Walker, 1849 (<i>Oxycera</i>)	26.34
<i>Ligyromyia</i> Kertész, 1916	26.55
<i>limae</i> Guérin, 1830, <i>Odontomyia</i>	26.44
<i>limbata</i> Bigot, 1879, <i>Oplachantha</i>	26.8
<i>limbata</i> Lindner, 1949, <i>Rhaphiocera</i>	26.23
<i>limbiventris</i> Enderlein, 1921, <i>Allognosta</i>	26.4
<i>linearis</i> James, 1971, <i>Merosargus</i>	26.14
<i>linearis</i> Walker, 1854, <i>Sargus</i>	26.19
<i>lineata</i> (Enderlein), 1921 (<i>Chiromyza</i>), <i>Barbiellinia</i>	26.2
<i>lineatus</i> (Fabricius), 1805 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.41
<i>longa</i> (Wiedemann), 1830 (<i>Acanthina</i>), <i>Acanthiomyia</i>	26.49
<i>longicornis</i> Enderlein, 1921, <i>Nonacris</i>	26.3
<i>longicornis</i> Kertész, 1909, <i>Pedinoera</i>	26.57
<i>longicornis</i> (Philippi), 1865 (<i>Beris</i>), <i>Macromeracis</i>	26.6
<i>longicornis</i> (Schiner), 1868 (<i>Exalreta</i>), <i>Neoxalreta</i>	26.7
<i>longicornis</i> (Walker), 1857 (<i>Acanthina</i>), <i>Cyphomyia</i>	26.27
<i>longicornis</i> (Williston), 1900 (<i>Campeprosopa</i>), <i>Analcoceroides</i> ...	26.20
<i>longifrons</i> Rondani, 1848, <i>Stratiomys</i>	26.46
<i>longipes</i> McFadden, 1971, <i>Merosargus</i>	26.14
<i>longiseta</i> Walker, 1854, <i>Aerochaeta</i>	26.9
<i>longisetosa</i> (Lindner), 1835 (<i>Berisargus</i>), <i>Nothomyia</i>	26.47
<i>longiventris</i> Enderlein, 1914, <i>Aerochaeta</i>	26.9
<i>lopesi</i> (Lindner), 1935 (<i>Berisargus</i>), <i>Nothomyia</i>	26.47
<i>lopesi</i> Lindner, 1964, <i>Dactylodeictes</i>	26.53
<i>Lophoteles</i> authors	26.52
<i>lucens</i> Loew, 1866, <i>Sargus</i>	26.19
<i>lucida</i> Gerstaecker, 1857, <i>Panaeris</i>	26.56
<i>lucidus</i> (Lindner), 1949 (<i>Geosargus</i>), <i>Sargus</i>	26.19
<i>luctifera</i> (Philippi), 1865 (<i>Beris</i>), <i>Archistratiomys</i>	26.4
<i>luederwaldti</i> (Enderlein), 1914 (<i>Aerodesmia</i>), <i>Hermetia</i>	26.37
<i>lunus</i> (Bigot), 1879 (? <i>Macrosargus</i>), <i>Sargus</i>	26.19
<i>luridus</i> Loew, 1855, <i>Merosargus</i>	26.14



luteiventris (Philippi), 1865 (Beris), <i>Macromeracis</i>	26.6
lutzi Curran, 1932, <i>Merosargus</i>	26.14
<i>Lyprotomyia</i> Kertész, 1909	26.55
lyricus James & McFadden, 1971, <i>Merosargus</i>	26.14
<i>Lysozus</i> Enderlein, 1914	28.22
macquartii Perty, 1933, <i>Sargus</i>	26.19
<i>Macromeracis</i> Enderlein, 1921	26.6
macula (Fabricius), 1805 (<i>Sargus</i>), <i>Microchrysa</i>	26.17
maculata (Fabricius), 1805 (<i>Stratiomys</i>), <i>Cyphomyia</i>	26.27
<i>maculata</i> Fabricius of Wiedemann part, 1830, <i>Cyphomyia</i>	26.27
maculata (James), 1953 (<i>Euryneurasoma</i>), <i>Brachycara</i>	26.30
maculata Lindner, 1949, <i>Microchrysa</i>	26.17
maculatus Kertész, 1914, <i>Nemotelus</i>	26.48
maculatus Lindner, 1933, <i>Abavus</i>	26.29
maculatus Williston, 1900, <i>Ptecticus</i>	26.12
maculifrons Enderlein, 1921, <i>Hemimegacis</i>	26.6
maculifrons (Walker), 1849 (<i>Odontomyia</i>), <i>Labostigmia</i>	26.43
<i>maculifrons</i> Wulp, 1888, <i>Odontomyia</i>	26.44
maculipennis (Blanchard), 1852 (Beris), <i>Beridops</i>	26.5
maculiventris (Rondani), 1850 (<i>Chrysochlora</i>), <i>Chrysochlorina</i> ..	26.35
major Lindner, 1964, <i>Engicerus</i>	26.54
malachitis Lindner, 1926, <i>Zuercheria</i>	26.46
<i>Manotes</i> Kertész, 1916	26.55
marginalis Lindner, 1964, <i>Eicochalcidina</i>	26.53
marginata James, 1967, <i>Psephiocera</i>	26.58
marginata James, 1969, <i>Adoxomyia</i>	26.30
marginata Loew, 1866, <i>Cyphomyia</i>	26.27
<i>marmorata</i> (Gerstaecker), 1857 (<i>Acanthina</i>), <i>Artemita</i>	26.50
maxima Kertész, 1908, <i>Panacris</i>	26.57
megalopyge James, 1971, <i>Merosargus</i>	26.14
megaspis Kertész, 1916, <i>Borboridea</i>	26.51
melanderi James & Wirth, 1967, <i>Hermetia</i>	26.38
<i>Melanochroa</i> Brauer, 1882	26.46
<i>Melanochroa</i> Röder, 1886	26.46
melanopsis Wiedemann, 1830, <i>Stratiomys</i>	26.46
melanothorax McFadden, 1970, <i>Dolichothrix</i>	26.21
melanothorax McFadden, 1971, <i>Merosargus</i>	26.14
melleus Rondani, 1848, <i>Sargus</i>	26.19
<i>Meringostylus</i> Kertész, 1908	26.33
<i>Meristocera</i> Lindner, 1964	26.55
<i>Merosargus</i> Loew, 1855	26.10
<i>Mesomyza</i> Enderlein, 1921	26.3
<i>Mctabasis</i> Walker, 1856	26.43
metallica Curran, 1934, <i>Acanthinomyia</i>	26.49
metallica Kertész, 1914, <i>Proegmenomyia</i>	26.58
metallica (Wiedemann), 1830 (<i>Oxycera</i>), <i>Nothomyia</i>	26.47
mctzi Johnson, 1919, <i>Psephiocera</i>	26.58
mexicana (Bellardi), 1859 (Beris), <i>Oplachantha</i>	26.8
<i>mexicana</i> Gigio-Tos, 1891, <i>Heteracanthia</i>	26.6
mexicana James, 1967, <i>Hermetia</i>	26.37
mexicana Johnson, 1895, <i>Odontomyia</i>	26.44
mexicana Kertész, 1908, <i>Euryneura</i>	26.32
mexicana Kertész, 1923, <i>Glaris</i>	26.33




mexicana Lindner, 1949, <i>Acrochaeta</i>	26.9
micheneri James, 1950, <i>Adoxomyia</i>	26.30
microcephalum (Kraft & Cook), 1961 (<i>Zabrachia</i>), <i>Borboridea</i> ...	26.51
<i>Microchrysa</i> Loew, 1855	26.16
microdonta Kertész, 1908, <i>Panacris</i>	26.57
<i>Microhadrestia</i> Lindner, 1943 ..	26.7
mimetica Kertész, 1916, <i>Pseudocyphomyia</i>	26.58
minacensis Lindner, 1931, <i>Rhaphiocera</i>	26.23
minor Brèthes, 1924, <i>Hadrestia</i>	26.6
minuta Lindner, 1943, <i>Microhadrestia</i>	26.7
minuta (Williston), 1900 (<i>Cynipimorpha</i>), <i>Eucynipimorpha</i>	26.54
miptitata Macquart, 1846, <i>Rhaphlocera</i>	26.23
mirabilis Curran, 1928, <i>Gowdeyana</i>	26.54
mirabilis James, 1971, <i>Merosargus</i>	26.14
mitis Curran, 1934, <i>Hermetia</i>	26.38
modesta (Lindner), 1949 (<i>Pyclomyia</i>), <i>Psephiocera</i>	26.58
modesta Philippi, 1865, <i>Beris</i>	26.5
<i>Monacanthella</i> James, 1936	26.33
monensis Curran, 1928, <i>Nemotelus</i>	26.48
<i>mucens</i> , error, <i>Hermetia</i>	26.38
multicolor (Lindner), 1949 (<i>Gcosargus</i>), <i>Sargus</i>	26.19
multisignatus Lindner, 1949, <i>Abavus</i>	26.29
muricicornis (Enderlein), 1921 (<i>Chiromyza</i>), <i>Barbiellinia</i>	26.2
mutabilis (Fabricius), 1787 (<i>Stratiomys</i>), <i>Hoplitimyia</i>	26.42
<i>Mycterocera</i> James, 1967	26.56
<i>Myiocavia</i> Lindner, 1949	26.56
<i>Myxosargini</i>	26.46
<i>Myxosargus</i> Brauer, 1882	26.46
nana (Bellardi), 1862 (<i>Acanthina</i>), <i>Artemita</i>	26.50
nana James, 1953, <i>Stratiomyella</i>	26.45
nana Lindner, 1935, <i>Hermetia</i>	26.39
nasicus (Williston), 1888 (<i>Euryneura</i>), <i>Diaphorostylus</i>	26.31
nasuta Enderlein, 1926, <i>Rhingiopsis</i>	26.45
nasuta (Macquart), 1849 (<i>Stratiomyia</i>), <i>Promeranisa</i>	26.45
<i>Neanalococcus</i> James, 1943	26.22
nebulifer James, 1941, <i>Ptecticus</i>	26.18
nebulifer James, 1971, <i>Merosargus</i>	26.14
neivai James, 1940, <i>Cyphomyia</i>	26.27
<i>Nemotelini</i>	26.47
<i>Nematotelus</i> , emend.	26.47
<i>Nemotelus</i> Geoffroy, 1762	26.47
<i>Neoacanthina</i> Kertész, 1914	26.56
<i>Neoberis</i> Lindner, 1949	26.7
<i>Neochauna</i> Williston, 1896	26.56
<i>Neochorisops</i> Lindner, 1935	26.7
<i>Neoeuxaereta</i> , emend.	26.7
<i>Neoeuxaereta</i> Osten Sacken, 1879	26.7
<i>Neorondania</i> Osten Sacken, 1878	26.25
neotropicus Curran, 1925, <i>Nemotelus</i>	26.48
<i>Netrogramma</i> Lindner, 1964	26.56
neuquenensis Kertész, 1914, <i>Nemotelus</i>	26.48
<i>Neurota</i> Curran, 1931	26.30
niger Bigot, 1879, <i>Nemotelus</i>	26.48




<i>niger</i> Lindner, 1964, <i>Lasiodeictes</i>	26.55
<i>niger</i> (Wiedemann), 1830 (<i>Sargus</i>), <i>Caecosis</i>	26.34
<i>nigerrima</i> Lindner, 1949, <i>Opiachantha</i>	26.8
<i>nigra</i> Bezzi, 1922, <i>Chiromyza</i>	26.2
<i>nigra</i> James, 1942, <i>Nothomyia</i>	26.47
<i>nigra</i> Lindner, 1933, <i>Homaiarthria</i>	26.33
<i>nigra</i> Lindner, 1949, <i>Progrypomyia</i>	26.34
<i>nigra</i> Williston, 1888, <i>Chordonota</i>	26.24
<i>nigreradiatum</i> Lindner, 1949, <i>Chlamydonotum</i>	26.52
<i>nigribasis</i> James, 1971, <i>Merosargus</i>	26.14
<i>nigribasis</i> Lindner, 1949, <i>Panacris</i>	26.57
<i>nigricornis</i> Kertész, 1914, <i>Nemoteius</i>	26.48
<i>nigrifacies</i> Bigot, 1879, <i>Hermetia</i>	26.38
<i>nigrifemorata</i> (Macquart), 1847 (<i>Sargus</i>), <i>Himantoloba</i>	26.10
<i>nigrifrons</i> Enderlein, 1914, <i>Ptecticus</i>	26.17
<i>nigrimana</i> Lindner, 1949, <i>Strobilaspis</i>	26.59
<i>nigrina</i> Lindner, 1949, <i>Adoxomyia</i>	26.30
<i>nigrinus</i> Fallén, 1817, <i>Nemoteius</i>	26.43
<i>nigritarsis</i> Enderlein, 1914, <i>Cyphomyia</i>	26.27
<i>nigritibia</i> Enderlein, 1914, <i>Hermetia</i>	26.38
<i>nigriventris</i> Enderlein, 1921, <i>Nonaeris</i>	26.3
<i>nigrobasalis</i> (Lindner), 1949 (<i>Borboridea</i>), <i>Manotes</i>	26.55
<i>nigrofemorata</i> Williston, 1900, <i>Berismyia</i>	26.5
<i>nigropygialis</i> Lindner, 1931, <i>Ptecticus</i>	26.18
<i>nigrovittata</i> Lindner, 1949, <i>Eechactomyia</i>	26.53
<i>nitida</i> (Aubertin), 1930 (<i>Chilota</i>), <i>Draymonia</i>	26.6
<i>nitidiceps</i> Wulp, 1888, <i>Odontomyia</i>	26.44
<i>nitidipennis</i> Loew, 1855, <i>Ptecticus</i>	26.18
<i>niveloscula</i> Lindner, 1949, <i>Brachyodina</i>	26.51
<i>niveipenicillata</i> (Lindner), 1949 (<i>Adoxomyia</i>), <i>Leucoptilum</i>	26.33
<i>Nonaeris</i> Walker, 1856	26.3
<i>nordenholzi</i> (Lindner), 1943 (<i>Huttonella</i>), <i>Benhamyia</i>	26.4
<i>notatus</i> Wiedemann, 1930, <i>Sargus</i>	26.19
<i>notatus</i> Lindner, 1949, <i>Merosargus</i>	26.14
<i>Nothomyia</i> Loew, 1869	26.47
<i>nova</i> Giglio-Tos, 1891, <i>Microchrysa</i>	26.17
<i>novaetetoniac</i> Lindner, 1949, <i>Actinomyia</i>	26.4
<i>nubilipennis</i> James, 1938, <i>Ptecticus</i>	26.18
<i>nubilipennis</i> James, 1939, <i>Cyphomyia</i>	26.27
<i>nucis</i> James, 1935, <i>Hermetia</i>	26.39
<i>nudifrons</i> James, 1953, <i>Hedriodiscus</i>	26.41
<i>nudus</i> Kertész, 1914, <i>Nemoteius</i>	26.48
<i>obesa</i> (Walker), 1860 (<i>Clitellaria</i>), <i>Spaniomyia</i>	26.59
<i>obscura</i> (Coquillett), 1902 (<i>Aochietus</i>), <i>Dieuryneura</i>	26.31
<i>obscura</i> (Jaenicke), 1867 (<i>Rondania</i>), <i>Cyphomyia</i>	26.27
<i>obscurus</i> (Wiedemann), 1830 (<i>Sargus</i>), <i>Merosargus</i>	26.14
<i>obtusipennis</i> (Lindner), 1949 (<i>Coenosargus</i>), <i>Merosargus</i>	26.14
<i>ochracea</i> Giglio-Tos, 1891, <i>Cyphomyia</i>	26.27
<i>ochracea</i> Wiedemann, 1820, <i>Chiromyza</i>	26.2
<i>ochraceus</i> McFadden, 1970, <i>Anisoscapus</i>	26.20
<i>Odonthomyia</i> , emendation	26.43
<i>Odontomyia</i> Mcigen, 1803	26.43
<i>opaliger</i> Lindner, 1931, <i>Merosargus</i>	26.14



<i>Oplacantha</i> , emendation	26.7
<i>Oplacantha</i> Rondani, 1863	26.7
<i>opulentus</i> Walker, 1854, <i>Sargus</i>	26.19
<i>opuntiae</i> James, 1966, <i>Caetobia</i>	26.51
<i>orbitalis</i> James, 1943, <i>Analcocerus</i>	26.20
<i>orizabae</i> Giglio-Tos, 1891, <i>Merosargus</i>	26.14
<i>ornata</i> (Macquart), 1846 (<i>Acanthina</i>), <i>Artemita</i>	26.50
<i>ornata</i> Macquart, 1846, <i>Rhaphiocera</i>	26.23
<i>ornata</i> Walker, 1850, <i>Cyphomyia</i>	26.27
<i>pachycephala</i> Schiner, 1868, <i>Odontomyia</i>	26.44
<i>pachycephala</i> Bigot, 1879, <i>Odontomyia</i>	26.44
<i>Pachygaster</i> Meigen, 1803	26.56
<i>Pachygastrinae</i>	26.49
<i>Pachyptilum</i> Lindner, 1969	26.33
<i>pallidipennis</i> (Williston), 1900 (<i>Lophoteles</i>), <i>Cosmariomyia</i>	26.52
<i>pallidiventris</i> (Macquart), 1846 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>pallidus</i> McFadden, 1971, <i>Merosargus</i>	26.14
<i>pallifrons</i> Curran, 1932, <i>Merosargus</i>	26.14
<i>pallipes</i> Fabricius, 1781, <i>Stratiomys</i>	26.46
<i>pallipes</i> Lindner, 1951, <i>Chlamydonotum</i>	26.52
<i>pallipes</i> Lindner, 1964, <i>Clarissa</i>	26.52
<i>pampinus</i> (Osten Sacken), 1886 (<i>Rhaphiocera</i>), <i>Phanerozus</i>	26.22
<i>Panaeris</i> Gerstaecker, 1857	26.56
<i>panamensis</i> Curran, 1929, <i>Myxosargus</i>	26.47
<i>panamensis</i> Greene, 1940, <i>Hermetia</i>	26.39
<i>panamensis</i> McFadden, 1971, <i>Merosargus</i>	26.14
<i>par</i> Curran, 1932, <i>Merosargus</i>	26.14
<i>Paraecanthinomyia</i> Lindner, 1964	26.57
<i>paradoxa</i> (Enderlein), 1913 (<i>Lagarinus</i>), <i>Tana</i>	26.3
<i>Parahermetia</i> Lindner, 1928	26.34
<i>Parantissa</i> Enderlein, 1914	26.24
<i>Pareidalimus</i> Lindner, 1964	26.57
<i>partitifrons</i> Enderlein, 1921, <i>Nonacris</i>	26.3
<i>parvicornis</i> (Enderlein), 1921 (<i>Chiromyza</i>), <i>Barbiellinia</i>	26.2
<i>parvicornis</i> James, 1939, <i>Nothomyia</i>	26.47
<i>Patagiomyia</i> Lindner, 1933	26.39
<i>paulseni</i> (Philippi), 1865 (<i>Cyclogaster</i>), <i>Abasanistus</i>	26.24
<i>paulseni</i> (Philippi), 1865 (<i>Lagarus</i>), <i>Tana</i>	26.3
<i>Pedicellina</i> James, 1952	26.18
<i>Pedinocera</i> Kertész, 1909	26.57
<i>Pelagomyia</i> Williston, 1896	26.36
<i>pellucens</i> Macquart, 1834, <i>Hermetia</i>	26.38
<i>pellucida</i> Rondani, 1848, <i>Stratiomys</i>	26.46
<i>Peltina</i> Lindner, 1964	26.57
<i>Peltina</i> Lindner, 1969	26.57
<i>penai</i> James, 1971, <i>Merosargus</i>	26.15
<i>peruana</i> Enderlein, 1921, <i>Oplacantha</i>	26.8
<i>peruana</i> Kertész, 1908, <i>Euryncura</i>	26.32
<i>peruana</i> Lindner, 1933, <i>Cyphomyia</i>	26.28
<i>peruviana</i> Kertész, 1914, <i>Artemita</i>	26.51
<i>peruviana</i> Macquart, 1855, <i>Odontomyia</i>	26.44
<i>peruvianus</i> James, 1971, <i>Merosargus</i>	26.15
<i>peruvianus</i> Kertész, 1914, <i>Nemotelus</i>	26.48




peruvianus Lindner, 1941, Euparyphus	26.32
petiolatus (Macquart), 1838 (Sargus), Merosargus	26.15
Pezodontina Lindner, 1949	26.22
Phanerozus James, 1966	26.22
<i>philippii</i> (Bigot), 1879 (Exaircta), Archistratiomys	26.4
philippii Enderlein, 1921, Hylorops	26.2
philippii (Rondani), 1863 (Beris), Neoexaircta	26.7
pieta (Brauer), 1882 (Compsosoma), Aerochaeta	26.9
pieta Macquart, 1834, Dieranophora	26.21
pieta Schiner, 1868, Cyphomyia	26.27
pieta Wulp, 1879, Rhapsiocera	26.23
pietipennis Kertész, 1908, Panaeris	26.57
pictipes James, 1971, Merosargus	26.15
pictithorax Curran, 1933, Merosargus	26.15
pilosissima Gerstaecker, 1857, Cyphomyia	26.27
pilosus James, 1942, Myxosargus	26.47
<i>pinguis</i> Walker, 1860, Stratiomys	26.45
plana (Brauer), 1882 (Compsosoma), Aerochaeta	26.9
plana Kertész, 1916, Manotes	26.55
plana (Walker), 1936 (Acanthina), Acanthinomyia	26.49
plancs James, 1967, Chaleidomorphina	26.51
planifrons James, 1939, Cyphomyia	26.27
<i>planifrons</i> Macquart, 1846, Hermetia	26.37
Platopsomyia James, 1937	26.34
Platylobium Lindner, 1933	26.58
plaumanni James, 1943, Leucoptilum	26.33
plaumanni Lindner, 1949, Neochorisops	26.7
plaumanni Lindner, 1964, Daetylacantha	26.53
<i>Plecticus</i> , error	26.17
pleuralis James, 1949, Hoplistes	26.22
pleuriticus Loew, 1866, Sargus	26.19
<i>plumula</i> Authors (Lophoteles), Cosmariomyia	26.52
pluricolor (Bigot), 1879 (Chrysochlora), Chrysochlorina	26.35
podexargenteus Enderlein, 1914, Artemita	26.51
polyposus Say, 1829, Nematelus	26.48
pomaceus Loew, 1855, Ptecticus	26.18
<i>pomaceus</i> Wiedemann of Macquart, 1834, Rhapsiocera	26.23
Popanomyia Kertész, 1909	26.58
<i>porcata</i> (Walker), 1857 (Acanthina), Artemita	26.50
prasina (Jaenicke), 1867 (Odontomyia), Hedriodiscus	26.41
princeps Gerstaecker, 1857, Stratiomys	26.46
priseus Enderlein, 1914, Abavus	26.29
productus James, 1971, Merosargus	26.15
Proegmenomyia Kertész, 1914	26.58
<i>Proegmenomyia</i> , error	26.58
Progrypomyia Lindner, 1949	26.34
Promeranisa Walker, 1854	26.45
propinqua Schiner, 1868, Euryneura	26.32
protudens (James), 1939 (Udamacantha), Labostigmina	26.43
proxima Kertész, 1908, Panaeris	26.57
<i>Pseptomma</i> Enderlein, 1919	26.50
Psephiocera Enderlein, 1914	26.58
Pseudoeyphomyia Kertész, 1916	26.58
pseudolyricus James, 1971, Merosargus	26.15



<i>ptecticoides</i> Lindner, 1928, <i>Trichocereocera</i>	26.36
<i>Ptecticus</i> Loew, 1855	26.17
<i>pterochausta</i> Osten Sacken, 1886, <i>Hermetia</i>	26.39
<i>pulchella</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.27
<i>pulehlla</i> James, 1936, <i>Acanthinomyia</i>	26.50
<i>pulehlla</i> Williston, 1888, <i>Oplachantha</i>	26.8
<i>puleher</i> James, 1971, <i>Merosargus</i>	26.15
<i>puleher</i> (Wiedemann), 1824 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.41
<i>pulehra</i> Wiedemann, 1830, <i>Hermetia</i>	26.39
<i>pulehra</i> (Williston), 1900 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>pulehra</i> (Williston), 1900 (<i>Chrysochroma</i>), <i>Himantoloba</i>	26.10
<i>pulehripenis</i> (Brauer), 1882 (<i>Artemita</i>), <i>Spaniomyia</i>	26.59
<i>punctifer</i> Lindner, <i>Pareidalimus</i>	26.57
<i>purpurea</i> (Walker), 1860 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>pustulosa</i> James, 1938, <i>Zuccheria</i>	26.46
<i>Pyelomyia</i> Kertész, 1916	26.54
<i>pygmaea</i> (Bellardi), 1862 (<i>Clitellaria</i>), <i>Euryneura</i>	26.32
<i>quadratus</i> James, 1971, <i>Merosargus</i>	26.15
<i>quadrifasciatus</i> Lindner, 1941, <i>Merosargus</i>	26.15
<i>quadrilineata</i> (Bigot), 1887 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>quadrilineatus</i> (Macquart), 1834 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>quadrinaculata</i> (Bellardi), 1859 (<i>Odonthomyia</i>), <i>Labostigmina</i> ..	26.43
<i>regularis</i> Curran, 1929, <i>Cyphomyia</i>	26.27
<i>regularis</i> James, 1969, <i>Adoxomyia</i>	26.30
<i>reinhardi</i> James, 1935, <i>Hermetia</i>	26.39
<i>reinhardi</i> , error, <i>Hermetia</i>	26.39
<i>relicta</i> Osten Sacken, 1886, <i>Hermetia</i>	26.39
<i>rettenmeyer</i> James, 1966, <i>Phanerozus</i>	26.22
<i>Rhaphiocera</i> Macquart, 1934	26.23
<i>Rhaphiocerinae</i>	26.20
<i>Rhingiopsis</i> Röder, 1886	26.45
<i>robusta</i> Kertész, 1908, <i>Euryncura</i>	26.32
<i>robustus</i> Brauer, 1882, <i>Chaetosargus</i>	26.34
<i>robustus</i> McFadden, 1971, <i>Merosargus</i>	26.15
<i>roedingeri</i> Lindner, 1941, <i>Nemotelus</i>	26.49
<i>Rondania</i> Jaenicke, 1867	26.25
<i>rossi</i> James, 1971, <i>Merosargus</i>	26.15
<i>rostrata</i> (Wiedemann), 1830 (<i>Stratiomys</i>), <i>Rhingiopsis</i>	26.45
<i>rostratus</i> Walker, 1856, <i>Metabasis</i>	26.43
<i>rotundatus</i> Curran, 1932, <i>Merosargus</i>	26.15
<i>rubra</i> Loew, 1866, <i>Cyphomyia</i>	26.27
<i>rubriceps</i> (Phillipi), 1865 (<i>Cyclogaster</i>), <i>Abasanistus</i>	26.24
<i>rubricornis</i> Kertész, 1923, <i>Abasanistus</i>	26.24
<i>rubricornis</i> (Macquart), 1846 (<i>Odontomyia</i>), <i>Lagostigmina</i>	26.43
<i>rufibasis</i> (Bigot), 1879 (<i>Macrosargus</i>), <i>Sargus</i>	26.19
<i>ruficornis</i> Bigot, 1879, <i>Nemotelus</i>	26.49
<i>ruficornis</i> Macquart, 1849, <i>Heteracanthia</i>	26.6
<i>ruficornis</i> (Schiner), 1868 (<i>Cyanauges</i>), <i>Antissa</i>	26.24
<i>rufifrons</i> Curran, 1934, <i>Euryneura</i>	26.32
<i>rufipalpis</i> Wiedemann, 1830 (<i>Xylophagus</i>), <i>Archistratiomys</i>	26.4
<i>rufipes</i> Loew, 1866, <i>Odontomyia</i>	26.44
<i>rufitarsis</i> Macquart, 1846, <i>Hermetia</i>	26.39
<i>rufiventris</i> Fabricius, 1805, <i>Hermetia</i>	26.38



<i>rufoabdominalis</i> Cole, 1923, <i>Nemotelus</i>	26.49
<i>rugosus</i> James, 1941, <i>Nemotelus</i>	26.49
<i>ryckmani</i> James, 1962, <i>Dicyphoma</i>	26.29
<i>ryckmani</i> James, 1965, <i>Parcidalinus</i>	26.57
<i>sackenii</i> Williston, 1885, <i>Ptecticus</i>	26.18
<i>sallei</i> Bellardi, 1859, <i>Sargus</i>	26.19
<i>sanctipauli</i> Lindner, 1931, <i>Rhaphiocera</i>	26.23
<i>saphireus</i> Bigot, 1887, <i>Sargus</i>	26.20
<i>sapphircus</i> , error or emendation, <i>Sargus</i>	26.20
<i>Sarginae</i>	26.8
<i>Sargus</i> Fabricius, 1798	26.18
<i>scalaris</i> Bigot, 1875, <i>Cyphomyia</i>	26.27
<i>schacfferi</i> (Coquillett), 1904 (<i>Cyphomyia</i>), <i>Dicyphoma</i>	26.29
<i>schildi</i> James, 1971, <i>Merosargus</i>	26.15
<i>schineri</i> Kertész, 1908, <i>Meringostylus</i>	26.33
<i>schnusci</i> Kertész, 1909, <i>Vittiger</i>	26.59
<i>schwarzi</i> Curran, 1928, <i>Sargus</i>	26.19
<i>schwarzi</i> James, 1940, <i>Cyphomyia</i>	26.28
<i>scrobiculus</i> McFadden, 1971, <i>Merosargus</i>	26.15
<i>scutellaris</i> Enderlein, 1921, <i>Nonacris</i>	26.3
<i>scutellaris</i> (Walker), 1854 (<i>Clitellaria</i>)	26.60
<i>scutellata</i> Cresson, 1919, <i>Cyphomyia</i>	26.27
<i>scutellata</i> Loew, 1869, <i>Nothomyia</i>	26.47
<i>scutellata</i> Macquart, 1834, <i>Hermetia</i>	26.39
<i>scutellatus</i> Williston, 1900, <i>Myxosargus</i>	26.47
<i>secundus</i> (Albuquerque), 1955 (<i>Parahermetia</i>), <i>Chaetosargus</i>	26.34
<i>seitzii</i> (Lindner), 1928 (<i>Parahermetia</i>), <i>Chaetosargus</i>	26.34
<i>semiluna</i> James, 1939, <i>Hoplitimya</i>	26.42
<i>sericea</i> Enderlein, 1921, <i>Mesomyza</i>	26.3
<i>sericeiventris</i> (Rondani), 1848 (<i>Stratiomys</i>), <i>Hoplitimya</i>	26.42
<i>setigera</i> James, 1938, <i>Hermetia</i>	26.38
<i>sexmaculata</i> Macquart, 1934, <i>Hermetia</i>	26.39
<i>sexmaculatus</i> McFadden, 1971, <i>Merosargus</i>	26.15
<i>sexnotatus</i> James, 1941, <i>Merosargus</i>	26.15
<i>sezspinosa</i> (Macquart), 1846 (<i>Diphysa</i>), <i>Hexacraspis</i>	26.54
<i>shannoni</i> James, 1939, <i>Cyphomyia</i>	26.28
<i>signatipes</i> Kertész, 1908, <i>Diaphorosstylus</i>	26.31
<i>similis</i> Bellardi, 1859, <i>Cyphomyia</i>	26.28
<i>similis</i> Enderlein, 1921, <i>Oplachantha</i>	26.8
<i>similis</i> (Macquart), 1855 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>simplex</i> Walker, 1860, <i>Cyphomyia</i>	26.28
<i>singularis</i> Macquart, 1846, <i>Nemotelus</i>	26.49
<i>slossonae</i> (Johnson), 1913 (<i>Euryneurasoma</i>), <i>Brachyera</i>	26.30
<i>smaragdiferus</i> (Bigot), 1879 (<i>Macrosargus</i>), <i>Macrosargus</i>	26.15
<i>sordidus</i> Kertész, 1914, <i>Nemotelus</i>	26.49
<i>soror</i> Lindner, 1964, <i>Spaniomyia</i>	26.59
<i>souzalopesi</i> Iide, 1967, <i>Cyphomyia</i>	26.28
<i>Spaniomyia</i> Kertész, 1914	26.58
<i>spatulatus</i> Williston, 1900, <i>Merosargus</i>	26.15
<i>speciosa</i> Gerstaecker, 1857, <i>Blastocera</i>	26.51
<i>speciosa</i> Lindner, 1915, <i>Cyphomyia</i>	26.28
<i>speciosus</i> Macquart, 1846, <i>Sargus</i>	26.20



<i>sphcodes</i> Curran, 1934, <i>Hermetia</i>	26.39
<i>splendens</i> Bigot, 1879, <i>Sargus</i>	26.20
<i>splendens</i> (Schiner), 1868 (<i>Chrysonotus</i>), <i>Himantoloba</i>	26.10
<i>Spyridopa</i> Gerstaecker, 1857	26.59
<i>Spyridota</i> , error	26.59
<i>Spyripoda</i> , error	26.59
<i>stamineus</i> (Fabricius), 1805 (<i>Seaeva</i>), <i>Merosargus</i>	26.15
<i>stieglmayeri</i> Lindner, 1931, <i>Rhaphiocera</i>	26.23
<i>stigma</i> (Giglio-Tos), 1891 (<i>Clitellaria</i>), <i>Dieuryneura</i>	26.31
<i>stigmaticus</i> (Lindner), 1949 (<i>Aloiphina</i>), <i>Merosargus</i>	26.15
<i>stoichoides</i> James, 1965, <i>Zabrachia</i>	26.60
<i>stratiomyaeformis</i> Lindner, 1935, <i>Zuercheria</i>	26.46
<i>Stratiomyia</i> , emendation	26.45
<i>Stratiomyella</i> James, 1953	26.45
<i>Stratiomyinae</i>	26.39
<i>Stratiomyini</i>	26.39
<i>Stratiomys</i> Geoffroy, 1762	26.45
<i>Strobilaspis</i> Lindner, 1949	26.59
<i>stylicornis</i> (Enderlein), 1921 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>subalba</i> (Walker), 1854 (<i>Stratiomys</i>), <i>Hoplitimyia</i>	26.42
<i>subclassicalx</i> Enderlein, 1921, <i>Oplachantha</i>	26.8
<i>subcuprata</i> (Walker), 1854 (<i>Stratiomys</i>), <i>Odontomyia</i>	26.44
<i>subinterruptus</i> (Bellardi), 1859 (<i>Sargus</i>), <i>Merosargus</i>	26.15
<i>subobseurus</i> Lindner, 1969, <i>Merosargus</i>	26.15
<i>subpellucida</i> James and Wirth, 1967, <i>Hermetia</i>	26.39
<i>suleifrons</i> Curran, 1929, <i>Cyphomyia</i>	26.28
<i>superba</i> Brauer, 1882, <i>Amphilecta</i>	26.36
<i>superba</i> (Lindner), 1949 (<i>Pedicella</i>), <i>Himantoloba</i>	26.10
<i>superba</i> (Lindner), 1949 (<i>Pyelomyia</i>), <i>Psephiocera</i>	26.58
<i>supernitens</i> James, 1971, <i>Merosargus</i>	26.16
<i>superpietus</i> (Lindner), 1949 (<i>Hedriodiscina</i>), <i>Hedriodiscus</i>	26.41
<i>Synaptochaeta</i> Lindner, 1964	26.59
<i>syphoides</i> Macquart, 1838, <i>Basentidema</i>	26.21
<i>taeniatus</i> (Wiedemann), 1830 (<i>Sargus</i>), <i>Merosargus</i>	26.16
<i>Tana</i> Reed, 1888	26.3
<i>tangens</i> James, 1971, <i>Merosargus</i>	26.16
<i>tarsalis</i> Gerstaecker, 1857, <i>Spyridopa</i>	26.59
<i>tau</i> Röder, 1886, <i>Rhingiopsis</i>	26.45
<i>taurus</i> James, 1943, <i>Analcocerus</i>	26.20
<i>teevani</i> Curran, 1934, <i>Hermetia</i>	26.39
<i>Tegocera</i> Lindner, 1964	26.59
<i>telfordi</i> James, 1971, <i>Merosargus</i>	26.16
<i>tenebrifera</i> James, 1943, <i>Cyclophleps</i>	26.21
<i>tenebricosus</i> Lindner, 1929, <i>Merosargus</i>	26.16
<i>tenuicornis</i> Enderlein, 1921, <i>Mesomyza</i>	26.3
<i>tenuicornis</i> James, 1971, <i>Merosargus</i>	26.16
<i>tenuicornis</i> Lindner, 1949 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>tenuis</i> Schiner, 1868, <i>Ptecticus</i>	26.18
<i>tenuistylus</i> Hanson, 1958, <i>Nemotelus</i>	26.49
<i>tenuiventris</i> (Bigot), 1879 (<i>Macrosargus</i>), <i>Sargus</i>	26.20
<i>terminalis</i> James, 1938, <i>Ptecticus</i>	26.18
<i>terminalis</i> James, 1971, <i>Merosargus</i>	26.16
<i>testaceus</i> (Fabricius), 1805 (<i>Sargus</i>), <i>Ptecticus</i>	26.18



<i>texana</i> (Melander), 1904 (Sargus), <i>Cephalochrysa</i>	26.9
<i>thomae</i> Curran, 1928, <i>Nemotelus</i>	26.49
<i>Thopomyia</i> Kertész, 1916	26.59
<i>thoracica</i> (Philippi), 1865 (Beris), <i>Maeromeracis</i>	26.6
<i>thoracicus</i> Macquart, 1835, <i>Sargus</i>	26.20
<i>tincta</i> (Walker), 1856 (<i>Chrysochlora</i>), <i>Acrochaeta</i>	26.9
<i>titschaki</i> Lindner, 1941, <i>Nemotelus</i>	26.49
<i>tomentosa</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.28
<i>tomentosa</i> Lindner, 1949, <i>Myiocavia</i>	26.56
<i>tortugellus</i> (Lindner), 1949 (<i>Hedriodiscina</i>), <i>Hedriodiscus</i>	26.41
<i>transequa</i> Walker, 1856, <i>Nonacris</i>	26.3
<i>transhansae</i> Lindner, 1931, <i>Rhaphiocera</i>	26.23
<i>transversus</i> McFadden, 1971, <i>Merosargus</i>	26.16
<i>tremoleranus</i> (Aldrich), 1928 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.41
<i>triangulatus</i> McFadden, 1971, <i>Merosargus</i>	26.16
<i>Trichocereocera</i> Lindner, 1928	26.36
<i>trichonota</i> (Philippi), 1865 (Beris), <i>Beridops</i>	26.5
<i>tricolor</i> James, 1940, <i>Histiodroma</i>	26.22
<i>tricolor</i> Loew, 1866, <i>Sargus</i>	26.19
<i>tricolor</i> Osten Sacken, 1886, <i>Euparyphus</i>	26.31
<i>tricolor</i> (Wiedemann), 1828 (Beris), <i>Oplachantha</i>	26.8
<i>Trigonocerina</i> Lindner, 1964	26.59
<i>trinitatus</i> Melander, 1903, <i>Nemotelus</i>	26.48
<i>tripartitis</i> , error, <i>Merosargus</i>	26.16
<i>tripartitus</i> James, 1971, <i>Merosargus</i>	26.16
<i>tristis</i> Enderlein, <i>Oplachantha</i>	26.8
<i>tristis</i> Loew, 1855, <i>Merosargus</i>	26.16
<i>tristrigata</i> (Enderlein), 1921 (<i>Xenomorpha</i>), <i>Chiromyza</i>	26.2
<i>tritacniatus</i> (Bellardi), 1859 (<i>Odonthomyia</i>), <i>Hedriodiscus</i>	26.41
<i>tritacniatus</i> Lindner, 1929, <i>Merosargus</i>	26.16
<i>trivittatus</i> Say of Giglio-Tos, 1891, <i>Pteticus</i>	26.17
<i>trivittatus</i> (Say), 1829 (<i>Stratiomys</i>), <i>Hedriodiscus</i>	26.41
<i>truquii</i> (Bellardi), 1859 (<i>Odonthomyia</i>), <i>Hedriodiscus</i>	26.41
<i>turacellus</i> (Lindner), 1949 (<i>Hedriodiscina</i>), <i>Hedriodiscus</i>	26.41
<i>Udamacantha</i> Enderlein, 1914	26.42
<i>unicolor</i> Authors, <i>Nemotelus</i>	26.89
<i>unicolor</i> (Walker), 1854 (<i>Clitellaria</i>), <i>Cyphomyia</i>	26.28
<i>valdiviana</i> (Philippi), 1865 (<i>Cyanauges</i>), <i>Antissa</i>	26.24
<i>valdiviana</i> (Rondani), 1863 (<i>Oplachantha</i>), <i>Antissa</i>	26.24
<i>vanderwulpiana</i> Brèthes, 1907, <i>Odontomyia</i>	26.44
<i>varia</i> (Curran), 1929 (<i>Chrysochlora</i>), <i>Chrysochlorina</i>	26.35
<i>variabilis</i> (Gmelin), 1790 (<i>Musca</i>), <i>Hoplitimyia</i>	26.42
<i>variabilis</i> Hanson, 1958, <i>Nemotelus</i>	26.49
<i>variabilis</i> (Loew), 1847 (<i>Chauna</i>), <i>Neochauna</i>	26.56
<i>varicornis</i> James, 1971, <i>Merosargus</i>	26.16
<i>varocerus</i> James, 1971, <i>Merosargus</i>	26.16
<i>variipennis</i> , emendation or error, <i>Hermetia</i>	26.39
<i>variipennis</i> Bigot, 1879, <i>Hermetia</i>	26.39
<i>varipes</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.28
<i>velutina</i> Bigot, 1879, <i>Stratiomys</i>	26.46
<i>ventralis</i> McFadden, 1971, <i>Merosargus</i>	26.16



<i>venustula</i> (Lindner), 1936 (<i>Chrysochroma</i>), <i>Himantoioba</i>	26.10
<i>versicolor</i> Bellardi, 1962, <i>Sargus</i>	26.20
<i>vertebratus</i> James, 1971, <i>Merosargus</i>	26.16
<i>vertebratus</i> (Say), 1824 (<i>Odontomyia</i>), <i>Hedriodiscus</i>	26.42
<i>verticalis</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.28
<i>vespertilio</i> (Fabricius), 1805 (<i>Sargus</i>), <i>Chrysochlorina</i>	26.35
<i>vespiformis</i> Lindner, 1969, <i>Auloeceromyia</i>	26.20
<i>vicina</i> Macquart, 1834, <i>Odontomyia</i>	26.44
<i>villosa</i> James, 1939, <i>Eupachygaster</i>	26.54
<i>villosa</i> (Lindner), 1969 (<i>Exodontha</i>), <i>Antissa</i>	26.24
<i>violacea</i> Macquart, 1855, <i>Cyphomyia</i>	26.28
<i>violaceiventris</i> Enderlein, 1921, <i>Heteracantha</i>	26.6
<i>violaceus</i> Jaenicke, 1867, <i>Sargus</i>	26.20
<i>virens</i> Wiedemann, 1830, <i>Stratiomys</i>	26.46
<i>virgata</i> Lindner, 1949, <i>Hermetia</i>	26.39
<i>virgata</i> Lindner, 1969, <i>Chloromelas</i>	26.40
<i>virjata</i> Enderlein, 1921, <i>Oplacantha</i>	26.8
<i>viridis</i> (Bellardi), 1859 (<i>Odonthomyia</i>), <i>Labostigmina</i>	26.43
<i>viridis</i> Bezzi, 1922, <i>Chiromyza</i>	26.2
<i>viridis</i> Hine, 1911, <i>Nothomyia</i>	26.47
<i>viridis</i> McFadden, 1971, <i>Merosargus</i>	26.16
<i>viridiventris</i> Philippi, 1865, <i>Beris</i>	26.5
<i>vitrisetosus</i> (Lindner), 1935 (<i>Eidalimus</i>), <i>Parcidalimus</i>	26.57
<i>vittata</i> Macquart, 1850, <i>Odontomyia</i>	26.44
<i>vittata</i> Walker, 1856, <i>Promerania</i>	26.45
<i>vittata</i> Wiedemann, 1820, <i>Chiromyza</i>	26.2
<i>vittata</i> Wulp, 1881 (<i>Lasiopa</i>)	26.29
<i>Vittiger</i> Kertész, 1909	26.59
<i>weyrauchi</i> Lindner, 1951, <i>Adoxomyia</i>	26.30
<i>wheeleri</i> Melander, 1803, <i>Nemotelus</i>	26.48
<i>wiedemanni</i> Gerstaecker, 1857, <i>Cyphomyia</i>	26.28
<i>willistoni</i> (Curran), 1929 (<i>Chrysochiora</i>), <i>Chrysochlorina</i>	26.35
<i>willistoni</i> (Enderlein), 1914 (<i>Gyneuryparia</i>), <i>Cyphomyia</i>	26.28
<i>wirthli</i> James, 1967, <i>Pachygaster</i>	26.56
<i>xanthobasis</i> James, 1940, <i>Cyphomyia</i>	26.28
<i>Xenomorpha</i> Macquart, 1838	26.2
<i>zeta</i> James, 1940, <i>Cyphomyia</i>	26.28
<i>zabrachia</i> Coquillett, 1901	26.60
<i>zernyi</i> Lindner, 1929, <i>Cyphomyia</i>	26.28
<i>zeteki</i> James, 1971, <i>Merosargus</i>	26.16
<i>zikani</i> Lindner, 1935, <i>Rhaphiocera</i>	26.23
<i>zonata</i> Lindner, 1935, <i>Anopisthocrania</i>	26.39
<i>Zuercheria</i> Lindner, 1926	26.46
<i>zurstrasseni</i> Lindner, 1933, <i>Platyobium</i>	26.58





DEPARTAMENTO DE ZOOLOGIA, SECRETARIA DA AGRICULTURA, SÃO PAULO

A CATALOGUE OF THE DIPTERA OF THE AMERICAS
SOUTH OF THE UNITED STATES

27

FAMILY PELECORHYNCHIDAE

CORNELIUS B. PHILIP

U.S. Department of Health, Education, and Welfare, Public Health Service, National Institutes of Health, National Institute of Allergy and Infectious Diseases, Rocky Mountain Laboratory, Hamilton, Montana 59849


Neotropical species, which have been recorded only from Chile, have earlier been catalogued (e. g. Kröber, 1934) in the family Tabanidae under the genus *Coenura* Bigot, monotypic for *C. longicauda* Bigot. As Mackerras (1955) has reaffirmed, the species should be transferred to the genus *Pelecorhynchus* Macquart for which he and Fuller (1942) erected the present family for this holantaretic group of flies, though only the genus was used by Stuardo Ortiz (1946) in his catalogue of Chilean Diptera. Enderlein (1922, 1925) had previously proposed a subfamily Pelecorhynchinae for this group of flies, which is presently tenable in view of the correlative Nearctic subfamily Bequaertomyiinae Philip (1965). The Family is phylogenetically a primitive one as most recently attested by Hennig (1966).

The Neotropical members of the family Pelecorhynchidae share with antipodean species a distinctive, compact group of adult characters, as quoted from Mackerras & Fuller (1942), and substantiated by larval and pupal characters:

"*Adult.* — Robust, ornate flies. Antennae subulate, porrect; third segment with eight clearly defined annuli. Thorax (and head) with long silky hairs, forming conspicuous tufts. Tibial spurs present. Wings with C continuous round margin; Rs long; cell R₁ wide, of medium length; M 4-branched; cell Cu₁ open, vein 1A sinuous; alulae developed. Ante-squame small, with a conspicuous tuft; squame very small or absent; when present it is attached only at extreme lateral end of squamal ridge. Abdomen with segments 1 to 4 normal, remainder tubular, telescopic, very extensible. Male hypopygium prominent; aedeagus with separated elements. Female with two-segmented cerci."

According to these authors, the Chilean species are not morphologically distinct from the Australian *fuseoniger* group which is differentiated only by tinctorial characters as confirmed in comparative study by the author. Although plant pollen has not been seen on Chilean specimens, meager reports indicate that flower visitation by adults has

27.1



been observed similar to habits reported for Australian species. Hematophagous proclivity has not been noted in either region. Distribution below, additional to the type-locality, is provided by provinces in Chile. The types of all but *P. kroeberi* (Lind.) have been seen by the author.

Subfamily PELECORHYNCHINAE

Genus PELECORHYNCHIUS Macquart

PELECORHYNCHIUS Macquart, 1850:332 (sep., p. 28). Type-species, *maculipennis* Macquart (mon.) = *personatus* (Walker).

Coenura Bigot, 1857:286. Type-species, *longicauda* Bigot (mon.).

Caenopygna Thomson, 1869:449 (*Caenopygna* (sic) of authors). Type-species, *maculipennis* Thomson (mon.) = *personatus* (Walker).

ELEGANS (Philippi), 1865:727 (*Coenura*). Type-locality: Chile, "prope Llico". Distr. — Chile (Maule, Concepción, Malleco, Colehagua, Llico, Nahuelbuta, "Blico" [Llico?]). Type ♂, fragments now in Santiago Nat. Hist. Mus.

var. *biguttatus* (Philippi), 1865:726 (*Coenura*, as species). Type-locality: Chile, Valdivia. Distr. — Chile (Valdivia, Chiloe, Angol, Llanquihue, Santo Domingo, "Futa"). Type? ♀, fragments now in Santiago Nat. Hist. Mus. NEW STATUS.

albopunctata Schiner, 1868:77 (*Coenura*). Type-locality: "Chile". Type ♂, Vienna Nat. Hist. Mus.

darwini Ricardo, 1900:102. Type-locality: Chile, Chiloe, "S.W. coast of South America". Type ♀, in British Museum (Natural History).

KROEBERI (Lindner), 1925:23 (*Coenura*). Type-locality: Chile, Magallanes, Punta Arenas. Distr. — Chile (Magallanes). Type ♀, destroyed in Hamburg.

LONGICAUDA (Bigot), 1857:287 (*Coenura*). Type-locality: "Chile". Distr. — Chile (Angol, Bio Bio, Nahuelbuta, Arauco, Colehagua). Type ♀, loc.?

PENAI Peckham, 1967:554. Type-locality: Chile, Arauca. Other distr. — Chiloe.

VULPES (Macquart), 1850:23 (*Pangonia*). Type-locality: "Chile". Distr. — Chile (Concepción, Valdivia, Coquimbo, Angol). Type ♂, Paris Mus.

aurantiacus Ricardo, 1900:103. Type-locality: "Chile". Type ♀, British Museum (Natural History).

XANTHOPLEURA (Philippi), 1865:726 (*Coenura*). Type-locality: Chile, Valdivia. Distr. — Chile (Valdivia, Chiloe, Cautín, Ñuble). Type ♀, Santiago Nat. Hist. Mus.

LITERATURE CITED

BIGOT, J.

1857. Diptères nouveaux provenant du Chili. *Ann. Soc. ent. France* (3) 5:42-72.

- ENDERLEIN, G.
 1922. Ein neues Tabanidensystem. *Mitt. zool. Mus. Berlin* 10(2):333-351.
 1925. Studien an blutsaugenden Insekten. I. Grundlagen eines neuen Systems der Tabaniden. *Mitt. zool. Mus. Berlin* 11 (2):255-409.
- HENNIG, W.
 1966. The Diptera fauna of New Zealand as a problem in systematics and zoogeography. *Pacific Insects Monograph* 9:81 pp. (transl. P. Wygodzinsky).
- KRÖBER, O.
 1934. Catálogo dos Tabanidae da América do Sul e Central, incluindo o México e as Antilhas. *Rev. Ent.* 4(2-3):222-276, 291-333.
- LINDNER, E.
 1925. Neue exotische Dipteren (Rhagionidae et Tabanidae). *Konowia* 4:20-24.
- MACKERRAS, I. M.
 1955. The classification and distribution of Tabanidae (Diptera). II. History: Morphology: Classification: Subfamily Pangoniinae. *Austral. J. Zool.* 3:439-511.
- MACKERRAS, I.M. & M.E. FULLER
 1942. The genus *Pelecorhynchus* (Diptera, Tabanoidea). *Proc. Linn. Soc. N. S. Wales* 67:9-76. [Key to South American species].
- MACQUART, J.
 1850. *Diptères exotiques nouveaux ou peu connus*. 4e. supplément. *Mém. Soc. des. Sci., de l'Agric. et des Arts, Lille* 1849:309-465. (Also publ. separately as Supplément IV, pp. 5-161, Paris. ?1850).
- PECHUMAN, I. L.
 1967. A new *Pelecorhynchus* from Chile (Diptera: Tabanoidea: Pelecorhynchidae). *Canad. Ent.* 99:554-555.
- PHILIP, C. B.
 1965. Family Pelecorhynchidae, in A. Stone, C. W. Sabrosky, W. W. Wirth, R.H. Foote, and J.R. Coulson (eds.): *A Catalog of the Diptera of America North of Mexico*. Agriculture Research Service, U.S.D.A., Agriculture Handbook n° 276. Washington, D.C. Govt. Printing Office, 1956, p. 319.
- PHILIPPI, R. A.
 1865. Aufzählung der ehilenisehen Dipteren. *Verh. zool. bot. Ges. Wien* 15:595-782.
- RICARDO, G.
 1900. Notes on the Pangoniinae of the Family Tabanidae in the British Museum collection. *Ann. Mag. Nat. Hist.* (7) 5:97-121.
- SEHNER, J. R.
 1868. Diptera [Art. 1], 388 pp. in *Reise der österreichischen Fregatte Novara*, Zool. 2(1B), Wien.
- STUARDO ORTIZ, C.
 1946. *Catálogo de los Dipteros de Chile*, 250 pp. Minist. Agr., Santiago.
- THOMSON, C. G.
 1869. *Kongl. Svenska Fregatten Eugénies Resa omkring Jorden*, Diptera: 68-614.
- WALKER, F.
 1868. *List Dipt. Coll. Brit. Mus.* 1:1-229.



INDEX

<i>albopunctata</i> (Schiner), 1868 (Coenura), Pelecorhynchus	27.2
<i>aurantiacus</i> Ricardo, 1900, Pelecorhynchus	27.2
<i>biguttatus</i> (Philippi), 1865 (Coenura), Pelecorhynchus var.	27.2
<i>Caenopnyga</i> , error for <i>Caenopygna</i> , q.v.	27.2
<i>Caenopnygna</i> Thompson, 1869	27.2
<i>Coenura</i> Bigot, 1857	27.2
<i>darwini</i> Ricardo, 1900, Pelecorhynchus	27.2
<i>elegans</i> (Philippi), 1865 (Coenura), Pelecorhynchus	27.2
<i>kroeberi</i> (Lindner), 1925 (Coenura), Pelecorhynchus	27.2
<i>longicauda</i> (Bigot), 1857 (Coenura), Pelecorhynchus	27.2
Pelecorhynchinae	27.2
Pelecorhynchus Macquart, 1850	27.2
<i>penai</i> Pechuman, 1967, Pelecorhynchus	27.2
<i>vulpes</i> (Macquart), 1850 (Pangonia), Pelecorhynchus	27.2
<i>xanthopleura</i> (Philippi), 1865 (Coenura), Pelecorhynchus	27.2







SciELO