

SYSTEMATICS OF *OSMORHIZA* RAF. (APIACEAE: APIOIDEAE)¹

PORTER P. LOWRY II² AND ALMUT G. JONES³

ABSTRACT

The genus *Osmorhiza* comprises ten species and two subspecies of woodland umbellifers native to temperate Asia and the Americas. Quantitative evaluation of nine morphological characters supports the recognition of two distinct subgenera: *Glycosma*, with a single species, *O. occidentalis*, and *Osmorhiza*, with the remaining nine species. The typical subgenus is further divided into three sections, each with three species. A complete synonymy and detailed description is provided for each taxon, along with an analytical key to the subgenera, sections, species, and subspecies. Phytogeographic data indicate that western North America is the center of distribution and diversity, and possibly the center of origin for *Osmorhiza*, although members of the genus are found in eastern North America as well as Asia and Central and South America. The amphitropical disjunctions between North and South America observed in the ranges of *O. chilensis* and *O. depauperata* (both sect. *Nudae*) are most likely the result of long-distance dispersal by migratory birds. By contrast, the disjunct populations of these taxa that occur in the Great Lakes region and eastern North America probably represent remnants of a more widespread, pre-Pleistocene distribution. The three species comprising *Osmorhiza* sect. *Osmorhiza* (the North American *O. claytonii* and *O. longistylis*, and the Asian *O. aristata*) provide an example of the classical eastern North American–eastern Asian pattern of disjunction. Although these taxa are very similar morphologically, the populations probably represent members of a relatively stable floristic assemblage that has persisted in these areas for a long period; the possibility that there has been any recent contact between these disjunct taxa through Beringia seems remote. The members of *Osmorhiza* sect. *Mexicanae* (*O. brachypoda*, *O. mexicana*, and *O. glabrata*) exhibit a somewhat broken distribution from northern California to central Argentina, and possibly represent derivatives of a common ancestral taxon that migrated southward along a chain of mountain tops with suitable temperate climatic conditions at some time during the Pliocene or Pleistocene. Ethnobotanical evidence indicates that at least five species of *Osmorhiza* were used for medicinal purposes by native Indian peoples of North America. The roots and greens of two species were also consumed as food by both North and South American Indians.

TAXONOMIC HISTORY

Osmorhiza (Apiaceae: Apioideae) is a small genus of perennial woodland herbs with representatives in temperate Asia and the Americas. Members of the genus typically flower in the spring, setting fruit by mid-summer. These fruits (schizocarps), which are armed with retrorse bristles in all but one species, are well adapted

for epizoochory. Plants of *Osmorhiza*, commonly known as Sweet Cicely, may be quite abundant in some areas, but they rarely form uninterrupted stands, even under the most favorable of conditions.

Osmorhiza was first recognized as a distinct genus by Rafinesque in 1818, but it was not until the following year (Rafinesque, 1819a) that this

¹ This paper resulted from research done by the first author for an M.S. thesis at the University of Illinois. We are grateful to D. A. Young for advice and suggestions. Special thanks go to L. Constance for valuable information and comments, and to S. C. Lowry for assistance in compiling phytogeographic data and preparing the base maps. We thank R. L. Hartman and two anonymous reviewers for helpful comments and suggestions. H. Heine is gratefully acknowledged for preparing the Latin diagnoses. Several persons provided living material and/or pressed specimens: D. E. Brink, P. K. Gorovoy, M. Mahu, R. H. Shan, A. L. Skvortsov, R. W. Sterner, M. J. Warnock, and O. Zöllner. We also thank the following persons for advice and information: A. W. Ghent, V. I. Grubov, R. Moberg, M. Muñoz S., and G. Wagenitz. The curators of the following herbaria generously provided the loan of specimens, photographs of types, and/or study facilities to examine collections in situ: ARIZ, AUA, BLH, BM, BR, CAS, CR, CS, DAO, DUKE, F, G, G-DC, GA, GH, GOET, ILL, ILLS, JEPS, K, KANU, LAF, LE, LL, MICH, MISSA, MO, MONT, MONTU, NMC, NY, OKL, ORE, OSC, P, PH, RM, RSA, SGO, SMU, TENN, TEX, UC, UPS, US, UWM, WIS, WTU, and Z. This work was supported in part by a Sigma Xi Grant-in-Aid of Research, an Illinois Academy of Sciences Research Grant, and the Robert Emerson Memorial Grant from the University of Illinois for 1978. An extensive list of representative specimens, a copy of which is on permanent deposit in MO, is available from the first author on request.

² Missouri Botanical Garden, P.O. Box 299, St. Louis, Missouri 63166, and Department of Biology, Washington University, St. Louis, Missouri 63130.

³ Department of Plant Biology, University of Illinois, 505 South Goodwin, Urbana, Illinois 61801.

name was validly published. Prior to that, specimens of *Osmorhiza* had been referred to three other genera: *Chaerophyllum* L. (Thunberg, 1784; Persoon, 1805), *Myrrhis* Miller (Michaux, 1803; Sprengel, 1813), and *Scandix* L. (Muhlenberg, 1813). During the next 80 years many new species were described in the genus. However, it was not until 1888, when Coulter and Rose published their "Revision of North American Umbelliferae," that an attempt was made to clarify the taxonomy of *Osmorhiza*. Twelve years later Coulter and Rose (1900) revised their previous treatment of the North American members of the genus, recognizing 12 taxa, including three new species. Mathias and Constance (1944), in their review of the Umbelliferae for the "North American Flora," listed eight species and two varieties of *Osmorhiza*.

Osmorhiza was treated for the first time on a world-wide basis by Constance and Shan (1948), who recognized the conspecificity of the North and South American populations now included in *O. chilensis* and *O. depauperata*. These authors included all the Asian populations of the genus under *O. aristata*, and described one new species, *O. bipatriata*. However, they acknowledged that "newer techniques" (i.e., biosystematic techniques) were not used in their study; they based their conclusions entirely on the examination and comparison of herbarium specimens.

Work on the current study was begun in 1974, at which time only two eastern North American species, *O. claytonii* and *O. longistylis*, were considered. Later, the scope of our work was expanded to a systematic investigation of the entire genus. Toward this end several approaches have been used, including (1) quantitative evaluation of nine morphological characters, (2) mapping of each species, (3) comparative examination of available type material, and (4) a broad survey of the literature on the genus. In addition, an analysis of electrophoretic banding patterns of water soluble seed proteins for most *Osmorhiza* species was conducted (Lowry & Young, 1979; Lowry, 1980).

In addition to the nearly 400 collections of *Osmorhiza* on deposit in ILL, approximately 8,500 specimens from 48 herbaria were studied. Photographs of type specimens unavailable for loan were received from LE and SGO. Voucher specimens for the more than 200 collections made by the first author are deposited in ILL, along with photographs of all type material examined.

The quantitative morphological data presented here support a revised taxonomy for *Osmorhiza* in which ten species and one subspecies are recognized. Two well delimited subgenera are distinguished, one of which is further divided into three sections. These conclusions are further supported by seed protein data (Lowry & Young, 1979; Lowry, 1980). Phytogeographic evidence indicates that western North America is the center of distribution and diversity, and possibly the center of origin for *Osmorhiza*, although members of the genus have migrated to and survived in eastern North America as well as in Asia and in Central and South America. Nomenclatural problems within the genus are dealt with, and a brief discussion of the ethnobotany is given.

NOMENCLATURE

Generic names. Species of *Osmorhiza* were originally placed in three other genera: *Chaerophyllum* L., *Myrrhis* Miller, and *Scandix* L. In 1818 Rafinesque published three alternative generic names for these taxa: *Washingtonia*, *Osmorhiza*, and *Gonatherus*. These were, however, invalidly published under Art. 34.1(a) of the "International Code of Botanical Nomenclature" (Voss et al., 1983), because they were not accepted by the author in the original publication. They may also be nomina nuda because they appear not to meet the requirements of Art. 41.2 of the "Code" (cf. Lowry & Jones, 1978; Lowry, 1985). In the same year Nuttall (1818) validly published the name *Uraspermum* for this taxon. Rafinesque (1819a), however, rejected *Uraspermum* Nutt., considering it too similar to (i.e., an incorrectly spelled later homonym of) *Urospermum* Scopoli, a genus of Asteraceae, and validated the name *Osmorhiza* with a reference to the description of *Uraspermum* Nutt. (Lowry & Jones, 1977, 1978). This interpretation has been followed by nearly all subsequent authors, although most have incorrectly considered Rafinesque (1819b) as the first place of valid publication for the name *Osmorhiza*. *Uraspermum* Nutt., however, cannot be treated as a later homonym since it is not spelled exactly like *Urospermum* Scopoli (cf. Art. 64.1 of the "Code"). Strict application of the rules of nomenclature would thus appear to require accepting the name *Uraspermum* Nutt. for the taxa currently recognized in *Osmorhiza*. In order to avoid this clearly undesirable change and to maintain no-

menclatural stability, a proposal has been made to conserve the generic name *Osmorhiza* against *Uraspermum* (Lowry, 1985).

Subgeneric names. Two subgenera are recognized within *Osmorhiza*, one with nine species, including the type for the genus (i.e., the type of *O. claytonii*), and one with a single species, *O. occidentalis*. When *O. occidentalis* was first described it was placed in a new, monotypic genus, *Glycosma*, by Nuttall (in Torrey & Gray, 1840), who provided both generic and specific descriptions. Drude (in Engler & Prantl, 1897), following Torrey (1859), judged this species to be an *Osmorhiza*, but placed it in a separate subgenus, citing Nuttall's generic name *Glycosma* as the basionym; the correct name for this subgenus is: *Osmorhiza* subg. *Glycosma* (Nutt.) Drude in Engl. & Prantl.

Sectional names. Three sections are recognized within *Osmorhiza* subg. *Osmorhiza* (Lowry & Jones, 1979b; Lowry, 1980). Constance and Shan (1948) published names for all three of these: (1) *Aristatae*, (2) *Mexicanae*, and (3) *Nudae*, although not validly, because none of them was accompanied by a Latin description or diagnosis or a reference to a previously and effectively published Latin description or diagnosis of the same taxon (cf. Arts. 32.1 and 36.1 of the "Code"). Because the section *Aristatae* includes the type of the genus, its epithet is a synonym for *Osmorhiza* sect. *Osmorhiza*. Constance and Shan's sectional names *Mexicanae* and *Nudae* are validated herein.

Specific epithets. Rafinesque (1830: 249) published three new species: *Osmorhiza dulcis*, *O. vilosa* [sic], and *O. cordata*. This publication reads as follows: "*Osmorhiza dulcis*, Raf. 1817 (*Myrrhis claytoni* Mx?) Sweet Sisily [sic]. Root fusiform, with a sweet smell and taste, near aniseed [sic], edible, carminative, expectorant, demulcent, useful for coughs with *Malva*, for flatulent bowels with *Heracleum*, Eq. to *Angelica*. Children are fond of this root, may be poisoned by mistaking for it, two sp. of the same genus or *Myrrhis* Auct. called *Poison* or *Bastard Sisily* [sic], distinctive by the roots less aromatic, foliage the same, but in *O. dulcis* base of the folioles acute, in my *O. vilosa* [sic] or *M. longistylis* obtuse, in *O. cordata* Raf. cordate. These last produce, when eaten, effects similar to those of the virulent Umbellate" Traditionally, placement of these names in synonymy has been based solely on this reference; *O. dulcis* under *O. claytonii*, and *O. vilosa* and *O. cordata* under *O.*

longistylis. The type specimen of *O. dulcis* (PH!) is, however, clearly a plant of *O. longistylis*, not *O. claytonii* as suggested by Rafinesque; *O. dulcis* is therefore a synonym of *O. longistylis*. The correct placement of the names *O. vilosa* and *O. cordata* may be inferred from Rafinesque's reference itself. *Osmorhiza dulcis* is said to have more aromatic roots than *O. vilosa* and *O. cordata*; the roots of *O. longistylis* are more aromatic than those of *O. claytonii* (Lowry & Jones, 1979a). Since *O. vilosa* is a synonym for *O. longistylis*, by inference *O. vilosa* and *O. cordata* belong in synonymy under *O. claytonii*. Rafinesque probably confused *O. claytonii* and *O. longistylis* when suggesting the synonymy for his new names.

Rydberg (1894) validly published the combination *Osmorhiza aristata* (Thunb.) Rydb. Nearly all subsequent authors have incorrectly attributed this combination to Makino and Yabe (in Makino, 1903), however; the only apparent exceptions are Constance (1972), Lowry and Jones (1979a), and Lowry (1980).

Accurate determination of the holotype of *Osmorhiza mexicana* Griseb. was possible only with the kind assistance of Professor Dr. G. Wagenitz (GOET). According to Wagenitz (pers. comm.), Grisebach usually marked voucher specimens he considered new by placing an "m" (for *mih*), after the new name, while other vouchers were marked with "Gr." Examination of the authentic material from the Grisebach Herbarium at GOET shows that *Schaffner 37* is marked "*Osmorhiza mexicana m*" in Grisebach's hand, while two other specimens (*Mandon 594* and *Lorentz & Hieronymus 668*) are marked "*Osmorhiza mexicana Gr.*" Thus, the Schaffner collection has been annotated as the holotype, whereas the other specimens are paratypes.

Lectotypes are designated here for four specific epithets: *O. brevistylis* DC., *O. depauperata* Philippi, *O. laxa* Royle, and *Myrrhis renjifoana* Philippi. There are several syntypes for each of these names, and extensive search in the literature has not revealed previous lectotypification for any of them.

The name *Myrrhis longistylis* Torrey (1824: 310) was published with the following type information: "In wet meadows near Albany, New York. Tracy. Near Geneva, N. Y. Paine. June. Near Hudson, N. Y. Alsop, & c." These specimens would be syntypes, although none of them has been located. An authentic collection by Paine marked "*Myrrhis n. sp.*" in Torrey's hand (at

NY) is therefore designated as the neotype for *Myrrhis longistylis* Torrey.

ETHNOBOTANY

Species of *Osmorhiza* were used in a variety of ways by native peoples in many parts of North America, and by at least one tribe in South America. Available information indicates that *O. brachypoda*, *O. chilensis*, *O. claytonii*, *O. longistylis*, and *O. occidentalis* were used for medicinal purposes, while *O. chilensis* and probably *O. depauperata* were also consumed as food. Many uses of *Osmorhiza* by Indians of North America were reviewed by French (1971).

Perhaps the most widely used species was *Osmorhiza chilensis*. In their review of the ethnobotany of the Karok and other Indians of northern California, Schenck and Gifford (1952: 386–387) reported: "The root of this plant is one of the most important medicines. It can be dried and kept in the house. The medicine requires a formula (charm) always, but it is used for almost everything. The house is smoked with it, if there has been illness in the house. The root is thrown on the fire at dances. If put under the pillow at night, it keeps sickness away. For headache, a little piece of the root is chewed. If a person is grieving over a lost relative, medicine is made from the root (with formula) and the mourner is bathed with the medicine. A piece of the root is carried as protection against the 'devil.' In the spring the young tops are eaten as greens. It is very good luck to find it growing in a place where it has never been seen before. Georgia Orcutt [a local Indian woman] says this is the only herb that is dried and kept on hand in the house. It is good for 'everything.'" Gunther (1945) reported that the Swinomish Indians of northwestern Washington chewed the roots of *Osmorhiza brevipes* (= *O. chilensis*) as a powerful love charm. She also noted that the Lummi and Skagit tribes of the same area did not seem to use *Osmorhiza*. Schneider (1906) indicated that the seeds of "*O. longistylis*" (certainly *O. chilensis*) were used by California Indians (probably the Paiutes) for medicinal purposes, primarily as a cough remedy. According to Steedman (1930: 480), the Thompson Indians of interior British Columbia ate the roots of *Osmorhiza chilensis*; she stated that "they are known for their delicate flavor and are especially appetizing because of their aromatic character."


Other species of *Osmorhiza* were used by In-

dians of western North America as cough remedies. The Paiutes prepared a decoction from the roots of *O. brachypoda* (Schneider, 1906), and several tribes from both Canada and the U.S. also used *O. occidentalis* for this purpose (Uphof, 1968).

Osmorhiza longistylis was used by many peoples of the midwestern U.S. According to Smith (1928: 249), the Meskwaki (i.e., Fox) Indians used this species for a variety of medicinal purposes: "It is chiefly used as an eye remedy. It is horse medicine, too, and the root is grated and mixed with salt for distemper. When hunting, they fed a pony with the root and he was thus enabled to catch the buffalo. Specimen 5154 of the Dr. Jones collection is the leaves of *Osmorhiza longistylis* and the bark of *Gleditsia triacanthos* mixed to make a tea which is drunk to regain flesh and strength." Gilmore (1919: 107) wrote: "The Omaha and Ponca say that horses were so fond of the roots of *Washingtonia [longistylis]* that if one whistled to them, while holding out the bag of roots, the horses came trotting up to get a taste, and so could easily be caught. An Omaha said that the roots were pounded up to make poultices to apply to boils. A Winnebago medicine-man reported the same treatment for wounds. A Pawnee said that a decoction of the roots was taken for weakness and general debility."

Smith (1932: 391) indicated that the Ojibwe (i.e., Ojibwa) Indians apparently did not distinguish between *Osmorhiza longistylis* and the closely related *O. claytonii*. He stated: "A tea for making parturition easier is prepared from the roots. The licorice flavor of the tea is said to be good for a sore throat." *Osmorhiza claytonii* was used by the Menomini Indians of northern Wisconsin to gain weight (Smith, 1923).

Osmorhiza was also used by Indians in southern South America. According to Mrs. Natalie Goodall of Harberton, Argentina (in a letter to Mrs. Helen Sharsmith, dated May 12, 1965, a copy of which was obtained from Dr. Lincoln Constance), *Osmorhiza* "was one of the three plants eaten by the [Yahgan] Indians besides fungi The owwunim [a Yahgan name] was eaten raw—leaves, stems, and roots, or the roots were baked in the fire My mother-in-law says that she often saw the Indians eating it as a child. Now it is hard to say, as there are only about four Yahgans left, two of them work in Harberton, but have forgotten their language." It is likely that the Yahgans did not distinguish between



FIGURES 1-4. Mean, standard deviation, range of variation, and sample size of quantitative morphological characters in *Osmorhiza* spp.—1. Style length (mm).—2. Stylopodium height (mm).—3. Fruit length (mm).—4. Fruit appendage length (mm).

Osmorhiza depauperata and *O. chilensis*, both of which grow throughout southern South America, often together. If they did, however, Mrs. Goodall's comments probably apply to *O. depauperata*, a specimen of which (*Y. Mexia* 7925), collected in Tierra del Fuego, Argentina, has written on its label "Awanim (Yeagan)."





The widespread use of these plants by native North and South Americans would suggest that other Indian groups, whose ethnobotany has not been studied, also used species of *Osmorhiza*. It is even possible that *O. aristata* was used by the people of Asia, although no record of this has been found.

MORPHOLOGY

Nine morphological characters, selected for their potential diagnostic value, were evaluated for all 11 taxa of *Osmorhiza*. Representative specimens were selected for study from across the geographic range of each taxon. The quantitative value for each character was obtained by taking the average of four measurements per specimen (occasionally two or three). Statistical significance of the data was determined using

contingency table analysis solved by chi square. We tested the null hypothesis that departures from random expectation are attributable to chance alone (Woolf, 1968). Mean, standard deviation, and range of variation of quantitative characters are presented in Figures 1-8. Sample sizes are given in parentheses. Comparison of values from the geographically disjunct areas of *Osmorhiza chilensis*, *O. depauperata*, and *O. aristata* is given in Tables 2-4, respectively.

1. *Style length*. This is a diagnostic character for the members of *Osmorhiza* sect. *Osmorhiza* (*O. longistylis*, *O. aristata*, and *O. claytonii*) but is of little taxonomic value for the other species (Figs. 1, 9a-c). Statistical analysis of style length values for members of section *Osmorhiza* yielded a chi square significant far beyond the 0.0005 probability level. The members of this section can readily be separated when this character is used in conjunction with geographic distribution. *Osmorhiza aristata*, with intermediate style length values, occurs only in Asia, whereas *O. longistylis* and *O. claytonii*, both of which are found in eastern North America, show no overlap in their ranges of values (Lowry, 1976; Lowry & Jones, 1979a; Ostertag & Jensen, 1980). Sev-



FIGURES 5–8. Mean, standard deviation, range of variation, and sample size of quantitative morphological characters in *Osmorhiza* spp.—5. Total hermaphrodite flowers per 1° umbel.—6. Total staminate flowers per 1° umbel.—7. Hermaphrodite ray length from 1° umbels (mm).—8. Pedicel length of hermaphrodite flowers (mm).

eral authors have relied entirely on style length to distinguish *O. longistylis* from *O. claytonii* (e.g., Deam, 1940; Steyermark, 1963).

2. *Stylopodium height*. The high-conic stylopodia of *Osmorhiza chilensis* are distinctive, and separate it from the other members of the section *Nudae*, which have low-conic to depressed stylopodia (Figs. 2, 9i–k). Statistical analysis of this character for the members of *O. sect. Nudae* resulted in a highly significant chi square. Furthermore, *O. glabrata* is distinguishable from the other members of the section *Mexicanae* by its fairly high, conic stylopodia (Fig. 2).

3. *Fruit length*. This character is diagnostic for the taxa included in *Osmorhiza sect. Nudae* (Figs. 3, 9i–k). These species are, in fact, distinguishable almost exclusively on the basis of fruit characters, including length and shape of the schizocarps, and size and shape of the stylopodium. Statistical analysis of fruit length measurements for the members of *O. sect. Nudae* yielded a highly significant chi square ($P \ll 0.0005$).

4. *Fruit appendage length*. The two subgenera of *Osmorhiza* are separable on the basis of presence or absence of caudate appendages on the base of the mericarps. All ten taxa comprising the typical subgenus have appendages of varying lengths, while they are lacking entirely from the fruits of *O. occidentalis* (*O. subg. Glycosma*), with a few rare exceptions (Figs. 4, 9). Within the typical subgenus, *O. purpurea* is distinct from the other members of the section *Nudae* by having short appendages. Similarly, *O. mexicana* subsp. *bipatriata* has shorter appendages than the other taxa in *O. sect. Mexicanae*.

Constance and Shan (1948) pointed out that the caudate appendages, along with the retrorse bristles on the fruits of most species, play an important role in dispersal. It is interesting to note that *O. chilensis* and *O. depauperata*, both of which have wide disjunctions in their ranges, have conspicuously armed fruits. By contrast, *O. occidentalis*, which lacks both appendages and bristles, has a continuous and somewhat restricted range in western North America. Similarly, *O. mexicana* subsp. *bipatriata*, with essentially

TABLE 1. Comparison of percentage of hermaphrodite flowers per 1° and 2° umbels in *Osmorhiza* species.

	1° Umbel			2° Umbel			Direction of Change
	Total ♀ Fls./ Umbel	Total ♂ Fls./ Umbel	% ♀ Fls.	Total ♀ Fls./ Umbel	Total ♂ Fls./ Umbel	% ♀ Fls.	
<i>O. occidentalis</i>	14.05	137.61	9.3	21.73	84.37	20.5	+
<i>O. aristata</i>	19.32	22.24	46.5	11.84	19.24	38.1	-
<i>O. claytonii</i>	13.94	8.20	63.0	10.68	10.32	50.9	-
<i>O. longistylis</i>	18.80	4.92	79.3	15.52	36.39	29.9	-
<i>O. glabrata</i>	20.73	32.50	39.9	11.60	21.40	35.2	-
<i>O. mexicana</i> subsp. <i>mexicana</i>	13.35	11.47	54.1	12.08	14.46	45.5	-
<i>O. mexicana</i> subsp. <i>bipatriata</i>	5.40	55.00	8.9	5.50	39.78	12.1	+
<i>O. brachypoda</i>	19.38	11.97	61.8	13.72	11.44	54.5	-
<i>O. chilensis</i>	16.49	5.10	76.4	11.39	7.84	59.2	-
<i>O. purpurea</i>	14.96	6.46	69.8	9.75	4.71	67.0	-
<i>O. depauperata</i>	13.20	3.04	81.3	9.98	2.67	79.0	-

glabrous fruits and very short appendages, has a very narrow range in Texas and northern Mexico.

5. *Total hermaphrodite flowers per 1° umbel.* Species of *Osmorhiza* are andromonoecious, a feature common to many genera of Apiaceae; their inflorescences contain both hermaphrodite and functionally staminate flowers. The staminate flowers have well-developed stylopodia that secrete nectar, but lack styles and functional ovaries, and consequently do not develop fruit (Lowry & Jones, 1979a; Lowry, 1980; see also Bell, 1971). The total number of hermaphrodite flowers produced in the primary umbels is quite uniform among *Osmorhiza* species (Fig. 5). Plants of each species produce similar numbers of fruits in their primary umbels (see below for the significance of this character).

6. *Total staminate flowers per 1° umbel.* This character is diagnostic in several ways. The two subgenera of *Osmorhiza* may be separated on the basis of this character; plants of *O. occidentalis* (*O.* subg. *Glycosma*) produce, on the average, more than twice as many staminate flowers per umbel as those of any taxon in *Osmorhiza* subg. *Osmorhiza* (Fig. 6). Furthermore, the members of *O.* sect. *Osmorhiza* are easily distinguishable from each other on the basis of this character. Values for the Asian *O. aristata* are intermediate between those for the North American *O. claytonii* and *O. longistylis* (Fig. 6). Statistical analysis of data for these three taxa yielded a very large chi square ($P \ll 0.0005$).

7. *Percentage of hermaphrodite flowers per 1°*

and 2° umbels. The ratio of hermaphrodite to staminate flowers differs between the primary and secondary umbels in *Osmorhiza* species (Table 1). In members of the typical subgenus, the primary umbel has a higher percentage of hermaphrodite flowers than does the secondary umbel. Therefore, the primary umbels of these plants contribute relatively more to the gene pool of the next generation through their ovules than do the secondary umbels. The only exception to this pattern in *O.* subg. *Osmorhiza* is *O. mexicana* subsp. *bipatriata*, for which there is a slight increase in the percentage of hermaphrodite flowers from primary to secondary umbel. This may be the result of the remarkably low number of hermaphrodite flowers in its primary umbels. This pattern of increasing "maleness" (or decreasing "femaleness") with successively later flowering umbels is correlated with protandry in the hermaphrodite flowers of many apiaceous taxa, including *Ligusticum canadense* (L.) Britt. and *Daucus carota* L. (Bell, 1971), and *Osmorhiza longistylis* (Torrey) DC. (Robertson, 1888), as well as the other species of *Osmorhiza* subg. *Osmorhiza*.

By contrast, *O. occidentalis* (subg. *Glycosma*) shows a strong trend toward increasing "femaleness" from primary to later-flowering umbels (Table 1). In these populations, the percentage of hermaphrodite flowers is more than twice as high in the secondary as compared to the primary umbel. Schlessman (1978, 1982) has shown that a similar increasing percentage of hermaphrodite flowers (expressed as a decreasing percentage of

TABLE 2. Morphological comparison of populations from the four major areas of distribution for *Osmorhiza chilensis*—mean, standard deviation, and sample size.

Character	Western North America			Great Lakes Region			Northeastern North America			Southern South America		
	\bar{x}	s.d.	N	\bar{x}	s.d.	N	\bar{x}	s.d.	N	\bar{x}	s.d.	N
Style length (mm)	0.61	0.08	38	0.57	0.06	6	0.64	0.06	9	0.69	0.14	24
Stylopodium ht. (mm)	0.42	0.09	38	0.38	0.04	6	0.43	0.07	9	0.37	0.08	24
Fruit length (mm)	16.66	2.71	36	16.42	0.93	6	18.74	1.47	9	14.97	1.72	22
Fruit appendage length (mm)	5.45	1.37	36	5.02	0.83	6	6.38	1.01	9	4.92	1.17	22
Total ♀ fls. per 1° umbel	17.58	6.02	38	13.00	6.60	6	15.44	6.33	9	16.50	7.53	24
Total ♂ fls. per 1° umbel	3.97	4.33	38	4.00	3.69	6	3.11	3.06	9	9.37	6.58	24
♀ ray length (mm) in 1° umbels	61.78	20.01	38	58.67	17.90	6	79.78	24.76	9	65.11	19.19	23
♀ pedicel length (mm)	9.65	3.26	38	11.39	3.39	6	10.78	3.83	9	9.00	3.51	29

staminate flowers) occurs in the protogynous species of tuberous lomatiums. Thus, although no observations have been made for *O. occidentalis*, circumstantial evidence suggests that it may be protogynous.

8. *Hermaphrodite ray length in 1° umbels.* Two types of rays are typically produced in the umbels of *Osmorhiza* species, those whose umbellets contain both hermaphrodite and staminate flowers (hermaphrodite rays), and those with umbellets composed entirely of staminate flowers (staminate rays). Hermaphrodite rays are generally rather stiff when the fruits reach maturity, ranging from 20 to 120 mm long, while staminate rays wither after flowering, and rarely exceed 10 mm. Hermaphrodite ray length is a useful diagnostic feature for distinguishing *O. aristata* from *O. longistylis* and *O. claytonii* (Fig. 7). This character is also useful for separating *O. brachypoda* from the other members of *O. sect. Mexicanae* (Fig. 7).

9. *Pedicel length of hermaphrodite flowers.* This is an excellent character for separating *O. aristata* from its North American relatives (Fig. 8). The pedicels in this species are, on the average, more than twice as long ($P \ll 0.0005$) as in plants of *O. longistylis* and *O. claytonii*.

10. *Comparison of disjunct populations of Osmorhiza species.* Comparison of values for morphological characters from the four geographically disjunct areas of *O. chilensis* (western North America, the Great Lakes region, northeastern North America, and southern South America) shows a remarkable similarity among the populations (Table 2). While a number of

populations in the Great Lakes area, the northeast, and South America appear to have undergone some divergence, there is no doubt as to their conspecificity with populations from western North America, where the species possibly originated.

Similarly, populations of *O. depauperata* from these same disjunct areas are morphologically very similar to one another (Table 3). While some divergence has occurred in South America, the specimens examined from this area are clearly within the range of variation found within the species in western North America.

Constance and Shan (1948) recognized two varieties of *Osmorhiza aristata*, the variety *laxa*, from southwestern China and the Himalayan region, and the typical variety from the Altai region and eastern Asia. These varieties were distinguished on the basis of leaf characters; plants from Sachalin and Siberia show a maximum of leaf dissection, those from Japan are intermediate, and those from the Himalayas have leaves that are much less divided. Although this variation in leaf dissection is indeed recognizable, there is no apparent correlation with other morphological characters. Quantitative values for these characters for populations from eastern Asia and the Himalayan region are quite similar, and do not support the recognition of infraspecific taxa within *O. aristata* (Table 4).

PHYTOGEOGRAPHY

Nearly 8,000 herbarium specimens were examined to determine the limits of geographic

TABLE 3. Morphological comparison of populations from the four major areas of distribution for *Osmorhiza depauperata*—mean, standard deviation, and sample size.

Character	Western North America			Great Lakes Region			Northeastern North America			Southern South America		
	\bar{x}	s.d.	N	\bar{x}	s.d.	N	\bar{x}	s.d.	N	\bar{x}	s.d.	N
Style length (mm)	0.46	0.07	33	0.46	0.10	7	0.46	0.09	7	0.55	0.08	9
Stylopodium ht. (mm)	0.27	0.06	33	0.31	0.08	7	0.27	0.06	7	0.29	0.06	9
Fruit length (mm)	13.74	1.56	32	14.11	0.83	7	14.79	0.86	7	12.28	1.68	8
Fruit appendage length (mm)	4.98	0.93	32	4.75	0.47	7	5.21	0.56	7	4.00	0.76	8
Total ♀ fls. per 1° umbel	13.21	4.21	33	11.33	1.75	6	13.86	2.41	7	12.22	3.46	9
Total ♂ fls. per 1° umbel	3.09	3.30	33	0.05	1.23	6	2.57	1.40	7	4.22	5.02	9
♀ ray length (mm) in 1° umbels	49.17	15.69	32	42.50	10.07	6	50.50	20.19	7	55.33	10.79	9
♀ pedicel length (mm)	14.34	3.56	33	14.50	3.36	7	16.74	4.32	7	15.05	2.72	10

distribution for *Osmorhiza* species. Distribution was mapped for each taxon by placing dots on a base map; one dot (or occasionally two) for counties in the U.S., and one dot per locality for other areas. A number of important collections reported by Hultén (1947), Constance and Shan (1948), Shishkin (1950), and Liu (1977) not available for study were mapped as circles. In general, the ranges obtained are in agreement with those given by other authors, notably Hultén (1947), Constance and Shan (1948), Constance (1963), Wood (1972), and Marquis and Voss (1981).

OSMORHIZA SECT. *OSMORHIZA*. Two species, *O. claytonii* and *O. longistylis*, are widely distributed in eastern North America and are sympatric over much of their ranges (Lowry, 1976; Lowry & Jones, 1979a). Both occur on the Gaspé Peninsula, in Nova Scotia, and in southern Quebec and Ontario, and reach their eastern limits

south along the Atlantic Coast. In the south, *O. claytonii* ranges from the southern Appalachian Mountains of North Carolina and Tennessee to the Ozark Plateau, approximately along the 35th parallel. In the west, this species reaches its limits along a line close to the 100th meridian, with extensions into north-central Nebraska and southwestern Manitoba (Fig. 10).

The range of *O. longistylis* extends farther to the south, reaching from the Piedmont Upland of central Georgia and South Carolina to Arkansas, eastern Oklahoma, and adjacent Texas. In the west, this species is found throughout the upper Great Plains, and has its limits along the base of the Rocky Mountains in Colorado, Wyoming, Montana, and Alberta (Fig. 11).

The third member of this section, *O. aristata*, occurs only in Asia, ranging from Sachalin and the lower Amur basin, through Japan, Korea, Taiwan, central and southern China, to the Hi-

TABLE 4. Morphological comparison of populations of *Osmorhiza aristata* from eastern Asia and the Himalayan Region—mean, standard deviation, and sample size.

Character	Eastern Asia			Himalayan Region		
	\bar{x}	s.d.	N	\bar{x}	s.d.	N
Style length (mm)	1.91	0.18	25	1.78	0.22	5
Stylopodium height (mm)	0.52	0.07	25	0.51	0.09	5
Fruit length (mm)	17.59	1.61	23	20.00	1.68	4
Fruit appendage length (mm)	6.27	0.99	23	7.91	1.82	4
Total ♀ fls. per 1° umbel	19.30	6.42	26	19.60	11.26	5
Total ♂ fls. per 1° umbel	23.04	9.73	25	17.25	7.23	4
♀ ray length (mm) in 1° umbels	71.09	14.17	23	80.29	26.18	4
♀ pedicel length (mm)	14.63	3.45	24	23.92	4.84	4




FIGURE 9. Drawings of the fruit of *Osmorhiza* species.—a. *O. aristata*.—b. *O. claytonii*.—c. *O. longistylis*.—d. *O. occidentalis*.—e. *O. glabrata*.—f. *O. mexicana* subsp. *mexicana*.—g. *O. mexicana* subsp. *bipatriata*.—h. *O. brachypoda*.—i. *O. purpurea*.—j. *O. chilensis*.—k. *O. depauperata*.




FIGURE 10. Geographic distribution of *Osmorhiza claytonii*.




FIGURE 11. Geographic distribution of *Osmorhiza longistylis*.




FIGURE 12. Geographic distribution of *Osmorhiza aristata*.

malayas of Bhutan, Nepal, India, and Pakistan. Disjunct populations occur in the Altai region of central U.S.S.R. and have been reported by Shishkin (1950) from the Caucasus Mountains of southeastern U.S.S.R. (Fig. 12).

The species of *Osmorhiza* sect. *Osmorhiza* provide an excellent example of the well known eastern North American—eastern Asian pattern of disjunction, which is shared by many members of the temperate deciduous forests of the two regions (Hara, 1952, 1956, 1972; Li, 1952, 1972; Constance, 1972; Wood, 1972; Ablaev et al., 1974). This pattern has long been recognized, e.g., by Linnaeus (1750; cf. Graham, 1966) and Thunberg (1784), but Asa Gray (1859) was the first botanist to examine it in detail. Gray indicated that many plants (including *Osmorhiza*) from eastern North America and Japan appear to be conspecific, and he also pointed out their conspicuous absence from western North America. To explain this phenomenon, he developed a hypothesis involving migration and exchange of species between North America and Asia across the Bering Strait, followed by their elimination from western North America and northeastern Asia during the Pleistocene.

Paleobotanical evidence confirms the widespread occurrence throughout much of North America and Eurasia of a number of genera, particularly woody ones, now restricted to the eastern parts of both continents. Wolfe (1969) indicated that certain members of the Tertiary flora of the Pacific Northwest survived changing climatic conditions, especially the shift from wet to dry summers, in relict habitats, particularly the mountains of southern Oregon and northern California. Wood (1972: 112) suggested that many plants occupying other regions of the Northern Hemisphere survived in a similar manner. He then stated: "The largest, and ecologically most complex, of the relict areas are eastern Asia and eastern North America; those of western North America and Europe-Asia Minor are smaller, and many genera that formerly occurred in them have disappeared." Wood concluded that the pattern of formerly widespread genera being restricted in distribution to eastern Asia and eastern North America by "orogenic movements, gradual climatic cooling, volcanism, and the Pleistocene glaciations seems to be well established."

The three species of *Osmorhiza* sect. *Osmorhiza* exhibit remarkable morphological similar-



FIGURE 13. Geographic distribution of *Osmorhiza purpurea*.

ity and have been regarded as conspecific by a number of authors (e.g., Gray, 1859; Clarke, 1879; Kuntze, 1891; Boivin, 1968). Constance and Shan (1948) suggested that this similarity may be ascribed to one of two factors: either there has been a relatively recent contact between the Asian and North American populations through Beringia, or these species have differentiated from a once widespread common ancestral population at a very slow rate. While Wood's reasoning, as outlined above, tends to support the latter, Li (1972) took a different view, arguing that the observed morphological similarities between populations of eastern Asia and eastern North America may be the result of geographic and ecological similarities between these areas. He pointed out that the two areas lie at roughly the same latitude, are situated in the same relative position on their respective continents, are tempered by ocean currents from the south, and share features of their topography, soil, temperature ranges, precipitation distributions, etc. Li believed that these morphological similarities were probably the result of parallel evolution, and may

not always accurately reflect an underlying phylogenetic relationship.

If, however, one regards eastern Asia and eastern North America as more or less stable, mesic refugia, it is possible that the species occupying these areas may have remained relatively unchanged both morphologically and genetically over long periods. Species of *Osmorhiza* appear to be stable taxa, with little or no indication of natural hybridization, and no "rapidly evolving" species. Rather than postulating the unlikely event of gene exchange in the recent past between the disjunct populations of this section, it seems more plausible to us to consider them parts of a relatively stable floristic assemblage that has persisted in eastern Asia and eastern North America over a rather long period.

OSMORHIZA SECT. *NUDAE*. All three species of this section occur in northwestern North America, and two of them, *O. chilensis* and *O. depauperata*, have disjunct populations in the Great Lakes area, northeastern North America, and southern South America.

Osmorhiza purpurea has a more restricted range, occurring more or less continuously along the Pacific Coast, from the redwood forests of northwestern California to Kodiak Island in southwestern Alaska. Inland, populations are found through the Cascade Mountains of Oregon and Washington eastward to the Rocky Mountains of Idaho, northwestern Montana, and extreme southwestern Alberta, and in much of southern British Columbia (Fig. 13).

Osmorhiza depauperata occurs sporadically in western North America, from the Warner Mountains of northeastern California, through Oregon, Washington, and central British Columbia, to the northern "pan-handle" and the Kenai Peninsula in Alaska. This species is much more commonly represented in the Rocky Mountains, where it ranges from southern Arizona and New Mexico northward to British Columbia, Alberta, and the extreme southwestern part of the Mackenzie District. Disjunct populations of *O. depauperata* are recorded from the Black Hills of South Dakota and the Cypress Hills in Alberta and Saskatchewan, and scattered populations are found across the Prairie Provinces of Canada in central Alberta, eastern Saskatchewan, and in Riding Mountain National Park, Manitoba. In the Great Lakes region, populations occur along the northern shore of Lake Superior, on Manitou Island in Lake Michigan, near Lake Nipigon in Ontario, and on Charlton Island in the James




FIGURE 14. Geographic distribution of *Osmorhiza depauperata*.

Bay Region (over 600 km to the northeast). In the east, *O. depauperata* ranges from Vermont and Lake Saint John in Québec, through the Gaspé Peninsula, Anticosti and Mingan Islands, to eastern Nova Scotia, Newfoundland, and southeastern Labrador. This species is also represented in South America, where it extends from Tierra del Fuego northward through the Andes of southern Argentina, reaching its northern limit at Termas de Chillán in Prov. Ñuble, Chile (Fig. 14).

Osmorhiza chilensis shows a similar overall pattern of distribution, and is sympatric with *O. depauperata* throughout much of its range, although it is much less common in the southern Rocky Mountains, particularly in Arizona and New Mexico. By contrast, populations of *O. chilensis* are common along the west coast, ranging from southern California, through Oregon and Washington, to Vancouver Island, the Queen Charlotte Islands, and southeastern Alaska. Occurrence of this species has also been reported from the coast of southwestern Alaska, with its range extending as far west as Unalaska Island in the Aleutian Chain (Hultén, 1947). With the exception of two populations in south-central Al-

berta, *O. chilensis* appears to be absent from the Prairie Provinces of southern Canada. In the Great Lakes area, it is widely distributed along the western and southern shores of Lake Superior, eastward to Lake Huron and the Georgian Bay; in eastern North America the range extends from northern New Hampshire and Maine, through the Gaspé Peninsula, New Brunswick, and Nova Scotia, to Newfoundland. South American populations occur from the Tierra del Fuego through the Andes and along the central Chilean coast northward to Prov. Aconcagua, Chile (Fig. 15).

Two well known types of disjunction are exemplified by *O. chilensis* and *O. depauperata*; amphitropical disjunctions between North and South America, and disjunctions between western and eastern North America.

Similarities between the floras of temperate western North America and southern South America were recognized over 100 years ago by Gray and Hooker (1880). Since then, many botanists have dealt with this subject (e.g., Engler, 1882; Reiche, 1907; DuRietz, 1940; Campbell, 1944; Constance, 1963; Raven, 1963; and Moore, 1972). Constance (1963) suggested that *Osmo-*




FIGURE 15. Geographic distribution of *Osmorhiza chilensis*.

rhiza chilensis and *O. depauperata* may have reached South America by a step-wise migration through the tropics along a route now marked by the members of the section *Mexicanae*. Raven listed about 130 species or species-pairs, including *O. chilensis* and *O. depauperata*, that exhibit patterns of disjunction between temperate North and South America. He also discussed the important factors concerning this pattern, which he summarized as follows: "(1) North and South American populations are closely related; (2) the plants are almost without exception self-compatible and often autogamous; (3) they constitute an unbalanced assemblage entirely unrepresentative of the floras of the two extratropical areas; (4) they grow almost exclusively in open communities, not in woodland or scrub associations; (5) there are no corresponding cases among terrestrial vertebrates and very few among the insects; and (6) the floras of the two areas have been distinct since at least the middle Cretaceous and are still very distinct at present." Raven (1963: 166) concluded that the only reasonable explanation to account for these facts is that "at least the great majority of the plants reached their disjunct areas by long-distance dispersal rela-

tively recently." He noted further that, for temperate species, the Pliocene or Pleistocene were the most likely times for this kind of dispersal, and that the majority of species involved migrated from north to south. The distribution patterns of *O. chilensis* and *O. depauperata* fit very well into this overall syndrome. There is no doubt that the North and South American populations of each of these species are very closely related. Although a few South American populations of both taxa seem to have undergone some morphological divergence, the majority of them appear to be virtually identical to their North American counterparts. Conversely, a number of morphological variants of both species that occur in North America seem to be absent from South America.

Observations made as part of this study suggest that *Osmorhiza* species are facultatively autogamous, a condition that would permit even a single propagule to establish a new population. Both *O. chilensis* and *O. depauperata* produce fruit armed with caudate appendages and numerous retrorse bristles, making them well adapted for epizoochory. Furthermore, the amphitropical pattern of disjunction for these taxa

corresponds closely to the migration routes of a number of bird species (Raven, 1963). Although many populations of both *O. chilensis* and *O. depauperata* occur in forest communities, these plants are also quite common in more open situations, making them accessible to a variety of potential dispersers. When taken together, these facts strongly suggest that the North American–South American disjunction found in *O. chilensis* and *O. depauperata* has resulted from long-distance dispersal by migratory birds, and that the direction of migration has been from north to south.

The western North America–eastern North America pattern of disjunction observed in *O. chilensis* and *O. depauperata* is somewhat less striking (Figs. 14, 15). Nevertheless, this type of disjunction has received considerable attention since Fernald (1924, 1925, 1926, 1933, 1935) developed his “nunatak” hypothesis to explain it (see also Marquis & Voss, 1981). Fernald believed that primarily arctic and western Cordilleran taxa were able to survive in eastern North America during the Pleistocene in unglaciated areas (nunataks) located around Lake Superior, on the Gaspé Peninsula, in the Long Range of Newfoundland, and in the Torngat Mountains of Labrador. Applying Willis’s (1922) “age and area” hypothesis, Fernald (1925: 243) interpreted the failure of these eastern disjuncts to extend their ranges following the glaciations as evidence of their “antiquity,” stating that “at the close of the Pleistocene they were already too old and conservative to pioneer, although they were able to linger as localized relicts in their special undisturbed crannies and pockets.”

Many of Fernald’s arguments have not withstood the test of time. All of his nunataks were in fact glaciated (Wood, 1972), and there is no reason why the plants involved could not have migrated to their present sites following the Pleistocene (Schofield, 1969, and references therein). Furthermore, the concept of senescence of species is no longer tenable (cf. Wood, 1972).

An alternative explanation for this pattern of disjunction is given by Stebbins (1935), who argued that each species involved migrated eastward at the end of the Pleistocene along the front of the receding ice sheet, becoming progressively more rare to the east. However, this hypothesis does not, by itself, explain the restriction of the majority of these taxa to the Great Lakes area and the northeast.

Schofield (1969: 200) adopted a somewhat

modified version of Fernald’s nunatak hypothesis, which he summarized as follows: “The most plausible explanation of their disjunctions is that the eastern representatives are remnants of a more widespread flora of the past, possibly of pre-Pleistocene arctic-alpine distribution in North America. The Pleistocene glaciations can be assumed to have eliminated the northcentral portion of the range, but since habitats were available in northeastern and western North America the species survived, probably south of the glacial boundary, but possibly in nunataks or coastal refuges, moving to their present sites following retreat of the ice sheet but being eliminated from their Pleistocene refugium by the encroaching vegetation and by a succession toward a more mesophytic temperate vegetation.” To this we would add the observation that these disjunct taxa are most likely restricted to their present localities by a variety of ecological factors: they tend to occur in areas where there is reduced competition from the dominant eastern boreal taxa (Rune, 1954), and where climatic conditions, particularly snowfall and moisture availability during the spring, are more similar to those of western North America.

Schofield’s hypothesis, as modified above, seems to be the most plausible to explain the western North American–eastern North American pattern of disjunction observed in *Osmorhiza chilensis* and *O. depauperata*. Nowhere in the literature have we found any suggestion that long-distance dispersal may have played a role in producing this type of disjunction; at least in the case of *Osmorhiza*, this type of dispersal would seem unlikely because the distribution of these plants does not seem to be correlated with the migration routes of any birds. Movement over shorter distances, however (e.g., from the Rocky Mountains into Saskatchewan and Manitoba), may be the result of epizoochory following the Pleistocene glaciations.

OSMORHIZA SECT. *MEXICANAE*. The representatives of this section form a “three-link chain” extending from the southwestern United States to central Argentina and Chile. The northernmost link is represented by *Osmorhiza brachypoda*, which has a relatively restricted distribution, ranging from the mountains of southwestern California northward through the Coast Range to Mount Diablo in Alameda County, and through the Sierra Nevada to Nevada and Sierra Counties, California. Disjunct populations occur in the Mazatzal Mountains of central Arizona




FIGURE 16. Geographic distribution of *Osmorhiza brachypoda*.

(Fig. 16), although they have not been collected since 1938.

Osmorhiza mexicana has an extended, although interrupted, range from southwestern Texas to northern Argentina. *Osmorhiza mexicana* subsp. *bipatriata* is known from only three localities: Madera del Carmen, Coahuila, and Cerro Potosí, Nuevo León, Mexico, and Mount Livermore, Jeff Davis Co., Texas. It is very possible, however, that this taxon also occurs on some of the other, as yet unexplored, mountains of Coahuila and Nuevo León. *Osmorhiza mexicana* subsp. *mexicana* reaches its northern limit in the Sierra Mohinora of Chihuahua and occurs together with subspecies *bipatriata* on Cerro Potosí. The typical subspecies ranges southward through the mountains of Mexico, Guatemala, and Costa Rica, into South America, where it occurs from the Páramo de Ruiz in Tolima, Colombia, through the Cordillera Oriental of southern Peru and adjacent Bolivia, and the mountains of northwestern Argentina, reaching its southern limit in the Sierra Grande of Córdoba, Argentina (Fig. 17).

Osmorhiza glabrata is restricted to the central Andes, ranging over a distance of less than 750 km from Prov. Santiago, Chile, to southern Neuquén, Argentina (Fig. 18).

The members of *Osmorhiza* sect. *Mexicanae* form a more or less continuous chain between northern California and central Argentina, "substantially bridging the gap which separates the discrete populations of the amphitropical taxa" in North and South America (Constance, 1963: 113). As Constance pointed out, however, it is




FIGURE 17. Geographic distribution of *Osmorhiza mexicana*.—*O. mexicana* subsp. *mexicana* (●).—*O. mexicana* subsp. *bipatriata* (□).

difficult to imagine a polytopic origin of both *O. chilensis* and *O. depauperata* from an extant or extinct member of the section *Mexicanae* both to the north and south of the equator.

Although one may be tempted to think of *O. brachypoda* and *O. glabrata* as northern and southern derivatives, respectively, of *O. mexicana*, there is no evidence to support this idea. A more plausible hypothesis is that all three species are derivatives of a once widespread common ancestral population whose distribution spanned the tropics, much as *O. mexicana* does today. According to Constance (1963), the




FIGURE 18. Geographic distribution of *Osmorhiza glabrata*.

mountains of Mexico and Central America were uplifted during the Pliocene or Pleistocene, orogenic movements that would have provided a chain of mountain tops with temperate climatic conditions suitable for the survival of such ancestral plants.

OSMORHIZA SUBG. GLYCOSMA. The sole member of this subgenus, *O. occidentalis*, is quite common in the somewhat drier areas throughout much of the western United States and adjacent Canada. It is widely distributed in the Rocky Mountains, ranging from southern Utah and southwestern Colorado northward to extreme southwestern Alberta and southern British Columbia, with eastward extensions into the Big Horn Mountains of Wyoming and the Big Snowy Mountains of Montana. To the west, the range of *O. occidentalis* extends through the mountains of the Great Basin in Nevada, eastern Oregon, and southwestern Idaho, to the northern Sierra Nevada, and the Coast Ranges of northern California and Oregon. This species also occurs in

the Olympic and Cascade Mountains of Washington, the Blue Mountains of Oregon, Washington, and Idaho, and in the Willamette Valley and western foothills of the Cascade Mountains in Oregon, but it has not been recorded in the central and eastern parts of the Cascades.

No major disjunctions occur in the range of *O. occidentalis*, but this is not surprising when one considers how poorly adapted its fruits seem to be for long-distance dispersal; the schizocarps are relatively large; they lack caudate appendages and retrorse bristles (Fig. 9d).

Constance and Shan (1948) erroneously included *Osmorhiza glabrata* in their section "*Glycosmae*" (= subg. *Glycosma*), leaving the group with a large North American–South American disjunction that is difficult to explain. If the affinities of *O. glabrata* are, however, recognized as lying with members of the section *Mexicanae*, the problem is eliminated.

The phytogeographic evidence presented here indicates that North America is the center of diversity, distribution, and possibly also the center of origin for the genus *Osmorhiza*. Although none of the presumably related genera (e.g., *Myrrhis*, *Chaerophyllum*, *Scandix*) occurs in the area, both subgenera of *Osmorhiza*, and eight of ten species, are represented here. Furthermore, *O.* subg. *Glycosma*, which appears to have retained many ancestral characters, is restricted to western North America, and all three sections of the typical subgenus have members in this continent.

SYSTEMATIC TREATMENT

***Osmorhiza* Raf.**, Amer. Monthly Mag. & Crit. Rev. 4: 192. Jan. 1819, nom. cons. prop.
TYPE: *Myrrhis claytonii* Michaux [= *Osmorhiza claytonii* (Michaux) C. B. Clarke].

Washingtonia Raf., Amer. Monthly Mag. & Crit. Rev. 2: 176. 1818, nom. nud., non *Washingtonia* H. Wendland (1879), nom. cons.

Osmorhiza Raf., Amer. Monthly Mag. & Crit. Rev. 2: 176. 1818, nom. nud.

Gonatherus Raf., Amer. Monthly Mag. & Crit. Rev. 2: 176. 1818, nom. nud.

Uraspermum Nutt., Gen. Amer. pl. 192. 1818, nom. rejic. prop.

Osmorhiza Raf., J. Phys. Chim. Hist. Nat. Arts 89: 157. 1819, nom. cons. prop. (cf. Lowry, 1985).

Spermatura Reichb., Consp. Reg. Veg. 141. 1828.

Glycosma Nutt. in Torrey & A. Gray, Fl. N. Amer. 1: 639. 1840.

Schudia Molina ex C. Gay, Fl. Chilena 143. 1874, pro syn.

Elleimataenia Koso-Poljansky, Bull. Soc. Imp. Naturalistes Moscou 29: 164. 1916.

Plants andromonoecious, slender to robust, perennial, herbaceous, aromatic, caulescent, dying back to a basal rosette of leaves; stems erect to spreading-ascending or decumbent, solitary to densely clustered, branching, fistulose, pubescent to glabrous. Roots fusiform, thick, fascicled, diffusely to extensively branched, surmounted by a branched caudex. Leaves alternate, basal and cauline, membranaceous, bipinnate or 2-3-ternate; leaflets lanceolate to orbicular, serrate to pinnately divided, with mucronate ultimate segments; petiole bases sheathing the stem, with their membranous margins densely ciliate to glabrous. Umbels twice compound, loose to somewhat constricted; peduncles terminal and lateral, erect to ascending or spreading, usually exceeding the leaves; involucre wanting, or composed of 1-several narrow, foliaceous, ciliate bracts; rays ascending to widely divaricate or reflexed, slender, unequal, the peripheral ones being longer; umbellets few to numerous, often of two kinds, those bearing hermaphrodite flowers or a mixture of hermaphrodite and staminate flowers (referred to as "hermaphrodite" umbellets), and those bearing only staminate flowers (referred to as "staminate" umbellets); involucrel wanting, or of several linear to ovate, acuminate, ciliate, spreading to reflexed bractlets; pedicels ascending to widely divaricate, those of the hermaphrodite flowers longer than those of the staminate flowers. Hermaphrodite flowers in each umbellet borne peripherally to the staminate flowers (if any), sometimes slightly irregular; calyx wanting; corolla white, greenish white, or yellow, or tinged with green, pink, or purple, the petals spatulate to ovate, the apex with an inflexed appendage; anthers about 0.5 mm long, smaller in the staminate flowers, inflexed before anthesis, spreading as the flower opens; styles spreading to divaricate, variable in length depending on the

species, but wanting in the staminate flowers; stylopodium conic to depressed, sometimes with a conspicuous disc, often nectariferous; carpophore 2-cleft from one-quarter to nearly one-half of its length. Fruit a schizocarp, deep brown to black at maturity, linear to oblong, fusiform to clavate, beaked to obtuse at the apex, sometimes constricted just below the stylopodium, shallowly to deeply concave furrowed, slightly compressed laterally, the ribs filiform, equidistant, moderately to sparingly hispid with retrorse bristles, or glabrous, the base rounded or with two caudate appendages; oil tubes (vittae) obscure or wanting; seeds subterete or unequally pentagonal in cross section, the face shallowly concave or sulcate.

Osmorhiza is clearly distinct from the monotypic European genus *Myrrhis* (apparently its closest relative) on the basis of the following characters: the leaves are 2-3-ternate or bipinnate (they are 2-4 times pinnatisect in *Myrrhis*), the umbels are definite in number (numerous in *Myrrhis*), the rays are glabrous (densely pubescent in *Myrrhis*), and the fruit are relatively small and lack winged ribs (fruit larger and strongly winged in *Myrrhis*).

Drude (1897), in his classic treatment of the Umbelliferae, placed *Osmorhiza* in his subfamily Apioideae, tribe Scandicineae, subtribe Scandicinae, along with the clearly related genera *Chaerophyllum* L., *Myrrhis* Miller, and *Scandix* L. A number of other genera, many of which are likely related to *Osmorhiza*, but some of which clearly are not, were also included in this subtribe. Bentham (1867) placed *Osmorhiza* in his series Heterosciadiae, tribe Ammineae, subtribe Scandicineae, along with the three genera mentioned above, as well as some others.

ANALYTICAL KEY

- 1a. Fruit glabrous, lacking caudate appendages; staminate umbellets 3-10 per umbel; staminate flowers (75-)90-225 per umbel; plants very robust; stems (1-)3-6(-8), densely clustered; leaves bipinnate; corolla yellow to greenish yellow (*Osmorhiza* subg. *Glycosma*)
One species 1. *Osmorhiza occidentalis*
- 1b. Fruit hispid with retrorse bristles (glabrous in *O. mexicana* subsp. *bipatriata*), with caudate appendages; staminate umbellets 0-4(-6) per umbel; staminate flowers 0-90(-125) per umbel; plants slender to rather stout; stems 1-3(-5), not densely clustered; leaves 2-3-ternate; corolla white, or tinged with green, pink, or purple (*Osmorhiza* subg. *Osmorhiza*)
- 2a. Involucrel conspicuous, composed of 1-6 spreading to reflexed, foliaceous bractlets.
- 3a. Styles (including stylopodium) 1-3.6 mm long; stylopodium high-conic; involucre composed of (1-)2-3(-5) bracts (*Osmorhiza* sect. *Osmorhiza*)
- 4a. Pedicels of the hermaphrodite flowers (5-)9-30 mm long; rays 4.5-11 cm long; fruit linear-clavate, obtuse or abruptly acute at the apex; plants restricted to Asia 2. *Osmorhiza aristata*

- 4b. Pedicels of the hermaphrodite flowers 4–10(–15) mm long; rays 1.5–8 cm long; fruit oblong-fusiform, strongly acute to attenuate at the apex; plants of North America.
- 5a. Styles 1–1.5(–1.7) mm long; flowers 4–7(–8) per umbellet; staminate flowers (2–)7–17(–23) per umbel; umbels loose and uncongested; roots rank-smelling, or sometimes weakly anise-scented 3. *Osmorhiza claytonii*
- 5b. Styles 2–3.6 mm long; flowers (7–)9–18 per umbellet; staminate flowers (23–)35–75(–86) per umbel; umbels dense and congested; roots with a strong, sweet, anise-like smell 4. *Osmorhiza longistylis*
- 3b. Styles 0.5–1.2 mm long; stylopodium low-conic to somewhat depressed; involucre wanting, or sometimes composed of 1–2(–3) reduced bracts (*Osmorhiza* sect. *Mexicanae*)
- 6a. Leaves laciniately lobed or pinnatifid; styles (incl. stylopodium) (0.9–)1–2 mm long; stylopodium 0.4–0.8 mm long; plants restricted to the central Andes of Chile and Argentina 5. *Osmorhiza glabrata*
- 6b. Leaves coarsely serrate to pinnately lobed or divided at the base, never lacinate; styles (incl. stylopodium) 0.5–1.2 mm long; stylopodium 0.2–0.5 mm long; plants occurring in North, Central, and South America.
- 7a. Pedicels of the hermaphrodite flowers 1–4.5 mm long, ascending; plants restricted to southern California and central Arizona 7. *Osmorhiza brachypoda*
- 7b. Pedicels of the hermaphrodite flowers (2–)3.5–7.5 mm long, spreading to spreading-ascending; plants occurring from southwestern Texas and northern Mexico to northern Argentina 6. *Osmorhiza mexicana*
- 8a. Fruit (10–)11–20 mm long, moderately to densely hispid with retrorse bristles, the appendages (1.5–)2–10 mm long; staminate flowers (0–)5–25(–35) per umbel; plants rather widespread from northern Mexico southward 6a. *Osmorhiza mexicana* subsp. *mexicana*
- 8b. Fruit 9–11(–12) mm long, glabrous, or occasionally with a few bristles toward the base, the appendages lacking, or to 1.8 mm long; staminate flowers (33–)40–70(–125) per umbel; plants restricted to three localities in southwestern Texas and adjacent Mexico 6b. *Osmorhiza mexicana* subsp. *bipatriata*
- 2b. Involucel wanting, or very rarely composed of 1(–2) minute bractlets (*Osmorhiza* sect. *Nudae*)
- 9a. Fruit linear-fusiform to linear-oblong, beaked at the apex; rays and pedicels spreading-ascending.
- 10a. Stylopodium high-conic to somewhat gibbous, lacking a disc, (0.2–)0.3–0.6 mm long; fruit tapering below the stylopodium, not constricted, 12–21(–25) mm long, the appendages 2.5–8.5(–10) mm long 8. *Osmorhiza chilensis*
- 10b. Stylopodium strongly depressed, with a conspicuous disc, 0.1–0.4 mm long; fruit constricted below the stylopodium, 8–15(–16) mm long, the appendages 1–5 mm long 9. *Osmorhiza purpurea*
- 9b. Fruit clavate, obtuse at the apex; rays and pedicels strongly divaricate to nearly reflexed 10. *Osmorhiza depauperata*

I. *Osmorhiza* Raf. subg. *Glycosma* (Nutt.) Drude. *Glycosma* Nutt. in Torrey & A. Gray, Fl. N. Amer. 1: 639. 1840. *Myrrhis* § [sect.] *Glycosma* (Nutt.) A. Gray, Proc. Amer. Acad. Arts 7: 346. 1868. *Osmorhiza* subg. *Glycosma* (Nutt.) Drude in Engl. & Prantl, Nat. Pflanzenfam. 3⁸: 153. 1897. *Washingtonia* subg. *Glycosma* (Nutt.) Coult. & Rose, Contr. U.S. Natl. Herb. 7: 62. 1900. *Osmorhiza* sect. *Glycosmae* (Nutt.) Constance & Shan, Univ. Calif. Publ. Bot. 23: 112. 1948, nom. nud. TYPE: *Glycosma occidentalis* Nutt. [= *Osmorhiza occidentalis* (Nutt.) Torrey].

bels; staminate umbellets 3–10 per umbel; staminate flowers (75–)90–225 per umbel; corolla yellow to greenish yellow. Fruit glabrous, without caudate appendages.

When *Osmorhiza occidentalis* was first described, Nuttall (in Torrey & Gray, 1840) placed it in a new, monotypic genus *Glycosma*. Plants belonging to this genus were distinguished from those of *Osmorhiza* by having unappendaged, glabrous fruit. Torrey (1859), Drude (1897), and Coulter and Rose (1900) united these genera, although in the latter two treatments *Glycosma* was retained as a separate subgenus.

Osmorhiza occidentalis is quite distinct from the other members of the genus. However, the morphological differences between *O. occidentalis* and the other species of *Osmorhiza* are much less pronounced than those between either of these groups and related genera such as *Myrrhis*, *Chaerophyllum*, and *Scandix*. Furthermore, *Os-*

Plants robust; stems densely clustered, (1–)3–6(–8), often glaucous. Root system deep, extensively branched. Leaves bipinnate; petiole bases finely ciliate to glabrous. Primary umbel with fewer hermaphrodite and more staminate flowers than the secondary and later flowering um-




FIGURE 19. Geographic distribution of *Osmorhiza occidentalis*.

morhiza, as treated here, is considered to represent a distinct monophyletic group. For these reasons, *Glycosma occidentalis* is included in *Osmorhiza* and placed in the monotypic subgenus *Glycosma*.

1. *Osmorhiza occidentalis* (Nutt.) Torrey. *Glycosma occidentalis* Nutt. in Torrey & A. Gray, Fl. N. Amer. 1: 639. 1840. *Osmorhiza occidentalis* (Nutt.) Torrey, Bot. Mex. bound. surv. 71. 1859. *Myrrhis occidentalis* (Nutt.) A. Gray, Proc. Amer. Acad. Arts 7: 346. 1868. *Washingtonia occidentalis* (Nutt.) Coult. & Rose, Contr. U.S. Natl. Herb. 7: 67. 1900. TYPE: U.S.A. Oregon: "Western side of the Blue Mountains," Nuttall s.n. [lectotype, NY! (designated by Coulter & Rose, Contr. U.S. Natl. Herb. 7: 67. 1900); isolectotypes, BM!, GH!].

Myrrhis bolanderi A. Gray, Proc. Amer. Acad. Arts 7: 346. 1868. *Glycosma bolanderi* (A. Gray) A. Gray, Proc. Amer. Acad. Arts 8: 386. 1872. *Osmorhiza occidentalis* var. *bolanderi* (A. Gray) Jepson, Madroño 1: 120. 1922. TYPE: U.S.A. California: Mendocino Co., Lambert's Lake, Bolander 6525 [lectotype, GH! (designated by Coulter & Rose, Contr. U.S. Natl. Herb. 7: 68. 1900); isolectotypes, K!, MO!, NY!].

Glycosma ambiguum A. Gray, Proc. Amer. Acad. Arts 8: 386. 1872. *Osmorhiza ambigua* (A. Gray) Coult. & Rose, Rev. N. Amer. Umbell. 119. 1888. *Myrrhis ambigua* (A. Gray) E. Greene, Fl. francisc.

332. 1892. *Washingtonia ambigua* (A. Gray) Coult. & Rose, Contr. U.S. Natl. Herb. 7: 69. 1900. TYPE: U.S.A. Oregon: Marion Co., foot of the Cascade Mts., Wolford's Orchard, Silver Creek, Hall 217 [lectotype, GH! (designated by Coulter & Rose, Contr. U.S. Natl. Herb. 7: 69. 1900); isolectotypes, F!, GOET!, K!, MO!, NY!].

Glycosma maxima Rydb., Bull. Torrey Bot. Club 40: 67. 1913. TYPE: U.S.A. Utah: Juab Co., Mt. Nebo, Rydberg & Carlton 7585 (holotype, NY!; isotype, RM!).

Plants robust, (3-)4-12 dm high; stems (1-)3-6(-8), erect or slightly ascending at the base, villos to hirsute just below the nodes, villosulous to glabrous elsewhere, and often glaucous. Root system deep, extensively branched, the roots with a strong, heavy anise-like fragrance. Leaves bipinnate, oblong to ovate, (8-)10-22 cm long, hirsutulous or more often glabrous; leaflets broadly lanceolate to ovate, 2-10(-12) cm long, 1-4(-5) cm wide, acute, serrate and incised or lobed at the base; petioles 5-25(-30) cm long. Umbels rather constricted; peduncles 2-4(-5), terminal and lateral, 6-18(-22) cm long; involucre wanting, or occasionally composed of 1-2 minute, linear, foliaceous, ciliate, spreading bracts; rays spreading-ascending to nearly erect, (2.8-)3-8 (-9.5) cm long; umbellets (3-)5-15 per umbel, (1-)3-10(-12) of them producing only staminate flowers; involucler wanting, or sometimes of 1 (-2) minute, linear-lanceolate, acuminate, ciliate bractlets; pedicels (7-)9-22(-25) per hermaphrodite umbellet, (6-)8-16(-18) per staminate umbellet, spreading to ascending, those of the hermaphrodite flowers (2-)2.5-7(-10) mm long, those of the staminate flowers 2-4.5(-6) mm long. Hermaphrodite flowers (1-)2-6 per umbellet, (1-)8-20(-45) per umbel, staminate flowers (6-)8-20(-22) per hermaphrodite umbellet, (73-)100-200(-225) per umbel; corolla yellow to greenish yellow, rather showy; styles plus stylopodium (0.7-)0.9-1.4(-1.8) mm long, stylopodium (0.1-)0.3-0.6 mm long, low-conic, with a conspicuous disc; carpophore cleft about one-third of its length. Fruit linear-fusiform, slightly constricted below the apex, rather deeply concave furrowed, (12-)13-22 mm long, the ribs glabrous throughout, the caudate appendages lacking, or very rarely to 1.5 mm long. $n = 11$ (Bell & Constance, 1957). Figures 9d and 19.

Flowering period. May to early July.

Habitat. Moist to rather dry forests, thickets, and open slopes.

Common names. Bald cicely, Mountain

sweet cicely, Sheep cicely, Sierra sweet cicely, Sweetanise, Sweetroot, Western sweet cicely, Western sweetroot.

Representative specimens. U.S.A. CALIFORNIA: Alpine Co., Red Lake, 2,440 m, *Johnson 130* (CS, NY, UC); Humboldt Co., NW slope of Black Mt., 760 m, *Tracy 8810* (CAS, JEPS, MO, NY, UC); Modoc Co., S slope of Eagle Peak, 2,440 m, *Alexander & Kellogg 5101* (UC); Mono Co., Sweetwater Canyon Creek, Sweetwater Mts., 2,440 m, *Alexander & Kellogg 3924* (JEPS, MO, UC); Nevada Co., S of Donner Pass, 2,600 m, *Heller 7183* (CAS, MICH, MO, NY, P, RM, UC); Sierra Co., *Lemmon 89* (MO, NY); *Rose 34375* (K); Siskiyou Co., Shackelford Creek, 1,220 m, *Butler 1668* (CAS, MO, NMC, RM, UC); Tehama Co., 1 km S of Lassen Chalet, *Lowry 849* (ILL, UC). COLORADO: Garfield Co., 10 km W of Triangle Park, *Klinger & Blumquist*, 10 July 1959 (CS); Gunnison Co., Ruby, *Baker 723* (MO, NY, UC); La Plata Co., Bob Creek, W La Plata Mts., 3,050 m, *Baker et al. 177* (MICH, MO, NY, RM, UC); Larimer Co., Rabbit Ears, *Goodding 1563* (MO, NY, RM, UC); Montezuma Co., 19 km N of Mancos, *Colyer 26* (CS); Pitkin Co., Maroon Lake, White River Natl. Forest, 2,900 m, *R. A. Nelson 7720* (CS); Rio Blanco Co., 1 km SW of Wilson Creek Camp, 2,350 m, *S. Tabar & J. Walker 368* (CS); Routt Co., Mts. S of Steamboat Springs, 2,595 m, *Porter 5960* (MO, NY, RM, SMU, TEX, UC). IDAHO: Bannock Co., Mint Creek Canyon, S of Pocatello, *Lingenfelter 685* (NY, UC, WTU); Blaine Co., Boulder Creek Canyon, 2,440 m, *Thompson 14092* (CAS, MICH, MO, NY, UC, WTU); Bonneville Co., 17.5 km SW of Victor, *Lowry 1118* (ILL, UC); Cassia Co., Black Pine Mts., 2,135 m, *N. H. Holmgren 3798* (NY, UC); Elmore Co., Dog Mt., N of Pine, 2,440 m, *Hitchcock & Muhllick 8726* (NY, UC, WTU); Fremont Co., above Blair Lake, Centennial Mts., 2,665 m, *Lowry 2609* (ILL, MONT, MONTU, UC); Lemhi Co., Quartzite Mt., *Hitchcock 14218* (MO, NY, WTU); Owyhee Co., 3 km S of Silver City, *Baker 8248* (NY, WTU); Teton Co., S of Victor, *Christ 5302* (NY); Twin Falls Co., 1.5 km S of Magic Mt. Ski Area, *Holmgren & Holmgren 6013* (UC). MONTANA: Beaverhead Co., Red Rock Lakes, *Lowry 1325* (ILL, MO, UC); Cascade Co., *Hawkins s.n.* (MONT); Fergus Co., Half Moon Ridge, Big Snowy Mts., *Hitchcock 16083* (CAS, MO, MONT, NY, RM, UC, WTU); Gallatin Co., Bridger Mts., *Rydberg & Bessey 4597* (K, MONT, RM, UC); Glacier Co., Midvale, *Umbach 389* (CAS, MONT, WIS); Granite Co., 3 km W of Skalkaho Rd. Summit, *Hitchcock & Muhllick 14479* (MO, NY, UC, WTU); Missoula Co., Missoula, *Kirkwood 1228* (CAS, MO, MONT, UC); Park Co., 3 km S of Livingston, *Booth 62156* (MONT); Silver Bow Co., Humbug Spires, *Lowry 2886, 2913* (ILL, MONT, MONTU, UC). NEVADA: Elko Co., Ruby Mts., *Gentry & Davidse 1806* (ILL, NY, RM, TEX, UC); Humboldt Co., Pine Forest Range, *Holmgren & Reveal 1202* (MICH, NY, TEX, UC); Lander Co., Kingston Canyon, Toiyabe Mts., *Tidestrom 10889* (MO); Nye Co., N Kawich Range, 2,045 m, *Beatley & Reveal 11221* (CAS, NY, UC); Washoe Co., Hunter Creek, W of Reno, 1,830 m, *Kennedy 1861* (CAS, MO, NY, RM, UC); White Pine Co., Snake Range, *Holmgren & Reveal 1097* (NY, TEX, UC). OREGON: Baker

Co., Powder River, *Cusick 1819* (JEPS, UC); Clackamas Co., *Goodding & Evinger*, 27 June 1927 (OSC); Clatsop Co., Saddle Mt., 715 m, *Chambers 3461* (OSC, NY); Crook Co., 3 km W of Ochoco Summit, *Kruckeberg 2153* (RM, UC, WTU); Deschutes Co., *Ireland 2663* (ORE); Grant Co., 7 km S of Long Creek, *Lowry 1094* (ILL, MO, NY, OSC, UC); Harney Co., Steens Mt., *Lowry 894* (ILL, OSC, UC); Jackson Co., Green Springs Mt., *Constance et al. 3629* (NY, SMU, UC); Klamath Co., 1.5 km N of Keno, *Peck 9426* (CAS, MO, NY); Lake Co., Crane Mt., *Thompson 13236* (CAS, MO, NY, WTU); Marion Co., Silver Creek, *Hall 217* (F, GH, GOET, K, MO, NY); Tillamook Co., *Chambers 4096* (OSC); Washington Co., Gales Creek, near Forest Grove, *Thompson 615* (CAS, MO, WTU); Wheeler Co., Wolf Mt., *Cronquist 7540* (CAS, K, NY, RM, UC, WTU). UTAH: Cache Co., Spring Hollow Canyon, 1,615 m, *Maguire 13739* (RM, UC); Garfield Co., 16 km E of Cedar Breaks, 2,450 m, *Graham 8679* (MO); San Juan Co., Abajo Mts., *Goodman & Hitchcock 1399* (MO); Summit Co., W Fork Bear River, Uintah Mts., *Payson & Payson 4936* (CAS, MO, RM, UC); Utah Co., Provo, 2,440 m, *M. E. Jones 5587* (MO, NY, RM, UC). WASHINGTON: Asotin Co., Blue Mts., above Indian Tom Cr., *Cronquist 5900* (MICH, NY, SMU, UC); Chelan Co., Tumwater Canyon, near Leavenworth, *Thompson 8451* (CAS, MO, NY, UC, WTU); Clallam Co., Mt. Angeles, 1,525 m, *Thompson 7470* (CAS, K, MO, UC, WTU); Jefferson Co., Hurricane Ridge, *Kuramoto*, 18 July 1966 (ILL); Kittitas Co., Upper Cle Elum River, *Kruckeberg 2976* (CAS, NY, RM, UC, WTU); Okanogan Co., Billy Goat Pass, 1,525 m, *Thompson 10881* (MO, NY, WTU); Spokane Co., Mt. Carleton, *Kraeger 282* (NY, UC); Yakima Co., Mt. Aix, 1,830 m, *Thompson 15056* (CAS, MICH, MO, NY, SMU, UC, WTU). WYOMING: Big Horn Co., Big Horn Mts., 2,835 m, *Gierisch 1790* (CS); Carbon Co., Lost Creek, Medicine Bow Mts., *Porter 4076* (CAS, RM, SMU, TEX, UC); Sublette Co., Middle Piney Lake, 2,500 m, *Porter 5034* (CAS, MO, RM, SMU, TEX); Teton Co., 5 km W of Teton Pass, *Lowry 1123* (ILL, UC); Uinta Co., Teton Mts., *Nelson & Nelson 6472* (BM, ILL, K, MO, NY).

CANADA. ALBERTA: Chief Mt. Int'l. Hwy., Waterton Lakes Nat'l. Park, 1,375 m, *Breitung 15930* (NY, UC); Red Rock Canyon, Waterton Lakes, 1,525 m, *de Vries 2102* (DAO); Mountain Hill, W of Pincher Creek, *Moss 56* (DAO); W of Beaver Mines, *Moss 825* (DAO). BRITISH COLUMBIA: near int'l. boundary, between Columbia River and Kettle River, 1,220 m, *Macoun 64628* (NY); near Ainsworth, 1,525 m, *Macoun*, 10 July 1890 (NY); Lightning Lake, Manning Prov. Park, 1,220 m, *Beamish & Vingtmann 60770* (DAO, ILL); km 30, Flathead Rd., *Bell & Davidson 150* (DAO); 9 km SW of Rossland, *Calder et al. 9460* (DAO); Mt. Apex, SW of Penticton, *Calder & Savile 11750* (DAO, UC); 1.5 km E of Phoenix, *Calder & Savile 33076* (DAO); 21 km W of Kaslo, 900 m, *McCabe 6572* (UC); 3 km N of Howell Creek Bridge on Flathead Rd., 1,525 m, *Taylor & Ferguson 1018* (DAO, UC); 13 km N of Flathead Customs, *Taylor & Ferguson 2039* (DAO).

II. *Osmorhiza* Raf. subg. *Osmorhiza*

Osmorhiza subg. *Euosmorhiza* Drude in Engl. & Prantl, Nat. Pflanzenfam. 3⁸: 153. 1897. nom. illeg.

Scandix subg. *Uraspermum* (Nutt.) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916.

Plants slender to rather stout; stems not densely clustered, 1–3(–5), never glaucous. Root system shallow to deep, diffusely branched. Leaves 2–3-ternate; petiole bases moderately to densely ciliate. Primary umbel with more hermaphrodite and fewer staminate flowers than the secondary and later flowering umbels; staminate umbellets 0–4(–6) per umbel; staminate flowers 0–90(–125) per umbel; corolla white, or variously tinged with green, pink, or purple. Fruit hispid with retrorse bristles, or occasionally glabrous, with short to very long caudate appendages (rarely lacking).

IIa. *Osmorhiza* Raf. sect. *Osmorhiza*

Uraspermum § [sect.] *Osmorhiza* (Raf.) Kuntze, Lexicon 582. 1904, pro parte.

Scandix 2. [sect.] *Urascandix* Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916, pro parte.

Osmorhiza sect. *Aristatae* Constance & Shan, Univ. Calif. Publ. Bot. 23: 112. 1948, nom. nud.

Involucre composed of (1–)2–3(–5) linear to lanceolate, reflexed bracts; involucrel conspicuous, composed of 3–6 reflexed bractlets. Styles (including stylopodium) 1–3.6 mm long, stylopodium 0.4–0.8 mm long, high-conic, lacking a disc.

There is no doubt that the three members of *Osmorhiza* sect. *Osmorhiza* (*O. claytonii*, *O. aristata*, and *O. longistylis*) are very closely related, and constitute a monophyletic assemblage; many authors (e.g., Gray, 1859; Clarke, 1879; Kuntze, 1891; Boivin, 1968) have even treated them as conspecific. These taxa, however, appear to represent distinct, natural populations. The Asian *O. aristata* is intermediate between the North American *O. claytonii* and *O. longistylis* for many characters but is clearly distinct for a number of others, including pedicel length and fruit shape.

The North American representatives of this section are completely separable from each other by many characters, including style length, total number of flowers per umbel, number of staminate flowers per umbellet, and pollen grain morphology (Lowry, 1976; Lowry & Jones, 1979a). Every one of the more than 2,250 herbarium specimens examined was clearly referable to one or the other of the two species; not a single individual exhibiting an intermediate combination of characters has been found. While pollen flow is likely in the numerous sympatric

populations, there is no indication of any natural hybridization or gene flow between *O. claytonii* and *O. longistylis*.

Palynological evidence also supports the treatment of *Osmorhiza claytonii* and *O. longistylis* as distinct at the specific level (Lowry, 1976; Lowry & Jones, 1979a). The pollen grains of the two species have significantly different ratios of polar axis length to equatorial diameter (P/E ratio). Those of *O. claytonii* are generally prolate in shape, i.e., their P/E ratios are between 1.33 and 2.00 (Erdtman, 1969; Kapp, 1969). By contrast, the grains of *O. longistylis* are perprolate, with P/E ratios greater than 2.00.

2. *Osmorhiza aristata* (Thunb.) Rydb. *Chaerophyllum aristatum* Thunb., Fl. Jap. 119. 1784. *Myrrhis aristata* (Thunb.) Sprengel, Pl. Umbell. 29. 1813. *Uraspermum aristatum* (Thunb.) Kuntze, Revis. gen. pl. 1: 270. 1891, pro parte. *Osmorhiza aristata* (Thunb.) Rydb., Bot. surv. Nebr. 3: 37. 1894, pro parte (exclusive of North American populations). *Scandix aristata* (Thunb.) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916. TYPE: Japan. *Thunberg s.n.* (holotype, UPS!).

Osmorhiza laxa Royle, Ill. bot. Himal. 233, pl. 52, fig. 1. 1839. *Washingtonia laxa* (Royle) Koso-Polj. in Fedtschenko, Fl. Asiat. Ross. 15: 52. 1920. *Washingtonia longistylis* var. *laxa* (Royle) Koso-Polj. in Fedtschenko, Fl. Asiat. Ross. 15: 52. 1920, pro syn. *Osmorhiza aristata* var. *laxa* (Royle) Constance & Shan, Univ. Calif. Publ. Bot. 23: 130. 1948. TYPE: India. Punjab: "Simore [Sirmur] in the Himalayan Mountains," *Royle s.n.* [lectotype (designated herein), K!; isolectotype, K!].

Osmorhiza japonica Siebold & Zucc., Abh. Math.-Phys. Cl. Königl. Bayer. Akad. Wiss. IV. 2: 203. 1843. Based on *Chaerophyllum aristatum* Thunb.

Osmorhiza amurensis F. Schmidt ex Maxim., Pr. fl. Amur. 129. 1859. *Scandix amurensis* (Maxim.) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916. *Washingtonia amurensis* (Maxim.) Koso-Polj. in Fedtschenko, Fl. Asiat. Ross. 15: 50. 1920, pro syn. TYPE: U.S.S.R. Khabarovsk Krai: on the lower Amur River, near the vicinity of the mouth of the Dondon, at Dshare, in deciduous woodland places, scattered, *K. I. Maximowicz*, 18 July 1855 [holotype, LE; photograph of holotype, ILL!; plus two authentic specimens (possibly isotypes), K!, LE, photographs at ILL!].

Osmorhiza claytonii (Michaux) C. B. Clarke in Hook., Fl. Brit. India 2: 690. 1879, pro parte (exclusive of *M. claytonii* Michaux and *M. longistylis* Torrey).

Osmorhiza aristata var. *montana* Makino, J. Jap. Bot. 2: 7. 1918. *Osmorhiza montana* (Makino) Makino, J. Jap. Bot. 5: 28. 1928. *Osmorhiza amuren-*

sis var. *montana* (Makino) Kitagawa, Rep. Inst. Sci. Res. Manchoukuo 2: 279. 1938. TYPE: Japan. Honshu: Totigi, Nikko, Shimotsuke, *T. Makino s.n.* (The holotype has not been located.)

Washingtonia claytonii subsp. *orientalis* Koso-Polj. in Fedtschenko, Fl. Asiat. Ross. 15: 51. 1920. (No type material has been located.)

Washingtonia claytonii subsp. *occidentalis* Koso-Polj. in Fedtschenko, Fl. Asiat. Ross. 15: 51. 1920. (No type material has been located.)

Plants rather stout, 3–8(–10) dm high; stems 1–2(–3), erect to ascending, villous to glabrate. Root system rather deep, spreading, with an anise-like scent. Leaves 2–3-ternate, deltoid to broadly ovate, 7–20 cm long, hirsutulous (especially along the veins) to glabrescent; leaflets oblong-oval to ovate-deltoid, (1.5–)2.5–9 cm long, 1–6 cm wide, obtuse to acuminate, coarsely serrate, incised, sparsely lobed to deeply pinnatifid at the base; petioles 5–25 cm long. Umbels loose; peduncles 2–3(–4), terminal and lateral, (3.5–)5–25 cm long; involucre composed of 1–3(–5) linear to lanceolate, foliaceous, ciliate, reflexed bracts, or sometimes wanting, each (1–)2–10(–25) mm long, (0.3–)0.5–1.3(–4.5) mm wide; rays spreading to ascending, (3.5–)4.5–11 cm long; umbellets 3–6 per umbel; involucre of (3–)4–5 linear to lanceolate, acuminate, ciliate to hirsutulous bractlets, each (1–)2.5–10(–11) mm long, 0.5–1.7(–2) mm wide, strongly reflexed; pedicels (4–)5–15 per umbellet, spreading, those of the hermaphrodite flowers (5–)9–30(–33) mm long, those of the staminate flowers 2.5–9(–10) mm long. Hermaphrodite flowers 2–6(–7) per umbellet, (10–)15–30(–38) per umbel, staminate flowers (2–)3–7(–11) per umbellet, (7–)15–40(–50) per umbel; corolla white, somewhat showy; styles plus stylopodium 1.5–2.2(–2.4) mm long, stylopodium (0.4–)0.45–0.7 mm long, high-conic, lacking a disc; carpophore cleft about to the middle. Fruit linear-clavate, obtuse or abruptly acute at the apex, concave furrowed, (13–)15–22(–23) mm long, the ribs sparingly to moderately hispid with retrorse bristles, especially toward the base, the caudate appendages (4.5–)5.5–11 mm long. $n = 11$ (Wanscher, 1932). Figures 9a and 12.

Flowering period. April to early June.

Habitat. Moist woods, at lower and middle elevations.

Common names. Ihari (Nepalese), Hsiang Kên Ts'ao Shu (Chinese), Miyama-yabu-ninjin, Nagajirami, Nagazirami, Onaga-yabu-ninjin, Yabu-ninjin, Yabu-ninjin (all Japanese).

Representative specimens. CHINA. GUIZHOU: *Cavalerie & Fortunat* 2961 (K, P, UC). HUBEI: *Henry* 5789 (BM, K, P); S of Wushan, *Wilson* 1044, *pars* (K, NY, P); Paok'ang, *Wilson* 1044, *pars* (P). JIANGSU: P'anghuang-chen, *Chiao & Cheo* 3541 (NY). JIANGXI: Lushan Mts., *Chung & Sun* 295 (NY). JILIN: Manchuria, *Komarov* 1158 (BM, K, P). SICHUAN: Pao-hsing-hsien, *Chu* 3336, 3495 (BM); Tchen-keou-tin, 1,400 m, *Farges* 72 (K, P, UC); Cheto Valley, Kangting (Tachienlu) Dist., 3,100 m, *Smith* 10963 (BM); Kiala, *Soulié* 1143 (P); S of Wushan, *Wilson* 1044, *pars* (K). XIZANG: Rongshar Valley, N of Mt. Everest, 2,900 m, *Hingston* 183 (K); 28°25'N, 97°55'E, 3,250 m, *Kingdon-Ward* 10092 (BM); Valley of Lilung Chu, between Charko and Lilung, 29°04'N, 93°56'E, 3,100 m, *Ludlow et al.* 4460 (BM); Peding, Tsangpo Valley, 29°30'N, 94°20'E, 3,000 m, *Ludlow et al.* 4533 (BM); Pe, 29°31'N, 94°54'E, 2,960 m, *Ludlow et al.* 5320 (BM). YUNNAN: Ma-eulchan, 3,000 m, *Delavay* 3902 (P); Yungning, *Handel-Mazzetti* 7049 (N—not seen; reported in Constance & Shan, 1948); N of Mengzi, 2,600 m, *Henry* 10233 (K, MO, NY). ZHEJIANG: W of Tien-mu, *Hu* 1654 (UC).

JAPAN. HOKKAIDO: Hakodate-si, Jesso Island, near Hakodate, *Albrecht s.n.* (K); Hakodate, *Faurie* 473 (P); Iwanai-tyo, Iwanai, *Faurie* 7024 (P); Kamikawa-tyo, foot of Mt. Kuro-dake, *Hiroe* 6628 (UC); Minami-huramo-tyo, Mt. Tomamu, *Hiroe* 6730 (UC); Monbetu-tyo, Monbetu, *Faurie* 709 (K, P); Rebun-tyo, Rebun Island, *Hiroe* 7611 (UC); Risiri-tyo, Mt. Rishiri, Rishiri Island, *Hiroe* 7541 (UC); Rubesibe-tyo, Onneyu, *Okamoto*, 18 Aug. 1958 (UC); Sapporo-si, Sapporo, *Tokubuchi*, 26 June 1891 (MO, NY); Yubari-si, foot of Mt. Yubari, *Hiroe* 6630 (UC). HONSHU: Akita-ken, Yokobori, *Yushun*, 23 July 1905 (NY); Aomori-ken, Moura, W coast of Natsudomari-hanto, *Mimoro et al.* 3551 (MO); Gunma-ken, Ikao, *Lyle*, May 1908 (BM); Hukusima-ken, Asakawa, *Mizushima*, 27 Apr. 1952 (UC); Ibaraki-ken, foot of Mt. Tsukuba, *Furuse*, 18 Apr. 1956 (UC); Iwate-ken, Nagamachi, *Ishiba*, 16 May 1926 (UC); Kanagawa-ken, Yamakita, *Faurie* 3303 (P); Kyoto-hu, Mt. Otoko, *Hiroe* 13565 (NY, UC, WIS); Miyagi-ken, Ninomaru, *Mori & Yashima*, 24 May 1973 (MO); Nagano-ken, Mt. Kiso-ontake, *Okuhara*, 28 Aug. 1955 (UC); Nara-ken, Mt. Kasuga, *Hiroe* 16236 (UC); Niigata-ken, Niigata, *Faurie* 79 (P); Osaka-hu, Mt. Chihaya, 600 m, *Hiroe* 13159 (NY, UC, WIS); Saitama-ken, Karisaka Pass, *Kobayashi*, 3 Aug. 1961 (UC); Siga-ken, foot of Mt. Ibuki, *Hiroe* 68 (K, NY, UC, WIS); Sizuoka-ken, ascent way of Fujinomiya, Mt. Fuji, 2,000 m, *Hiroe* 12683 (UC); Tokyo-to, Komae, *Suzuki* 77007 (UC); Totigi-ken, Lake Kirikomi, Nikko City, *Ono & Kobayashi*, 4 Aug. 1963 (UC); Toyama-ken, Yatsuo village, 17 km SW of Toyama, *Kirino* 172 (MO); Yamagata-ken, Kabuto-iwa, *Ohashi et al.* 708124 [sic] (BR, MO); Yamaguchi-ken, Hakusan Shrine, base of Castle Mt., 41 km W of Hiroshima, *Charette* 1680 (UC). KYUSHU: Kagosima-ken, *Masamune*, 4 Apr. 1923 (NY); Kumamoto-ken, Aso Volcanos, Kamisikimi Takamori, 650 m, *Tokio* 374 (WIS); Nagasaki-ken, Nagasaki, *Faurie* 3302 (BM, P); Oita-ken, Mt. Yuhu, 1,200 m, *Tokio* 1009 (WIS). SHIKOKU: Kagawa-ken, Mt. Ohtaki, 800 m, *Hiroe* 15500 (UC).

KOREA. CHEJU DO: Cheju-do Island, *Jaquet* 881 (K). KANGWON DO: Ullung Island, Kyongsang-Pukto, *Chung* 2290 (MICH); Mt. Odae, *Chung* 2728 (MICH). KYONGGI DO: Kwangnung, *Chung* 2614, 7404 (MICH).

BHUTAN. Chalimarphe Timpu, 2,290 m, *Cooper 1405* (BM); Drugge Dzong, 3,050 m, *Ludlow et al. 16209* (BM).

NEPAL. PĀLPA: Lukarban Khola, W of Beni, 3,000 m, *Stainton et al. 451* (BM); Lete, S of Tukucha, Kali Gandahi Valley, 3,000–3,200 m, *Stainton et al. 1034*, 5600 (BM).

INDIA. HIMACHAL PRADESH: Raiengarb Mut, 2,135 m, *Gamble 26799* (K); Swajau Maida, Parbatti Valley, 3,050 m, *Nath 123* (NY); Kulu-Lahoul, Punjab, *Drummond 23122, 23131, 23132* (K, UC); Nagzuda, Simla Hills, 2,250 m, *Hooker f. & Thomson*, 9 June 1849 (K, P). JAMMU: Jangla, 2,750 m, *Dudgeon & Kenoyer 386* (MO); Jammu, 2,000 m, *Hooker f. & Thomson*, 31 May 1868 (K). KASHMIR: Pahlgam, E Liddar River, 43 km N of Anantnag [Islamabad], 2,290 m, *Dickason 845* (MICH); Pahlgam, 2,200 m, *Stewart 9271* (NY). UTTAR PRADESH: below Budhi, Byans, Kumaun, 2,600 m, *Duthie 5595* (BM, K); Bamon Valley, 2,135 m, *Duthie*, 15 May 1897 (P); Jaunsar Bahar, near Kinani Pani, 2,600 m, *Gamble 1136* (K); Jaunsar, Chaelipup, *Gamble 23589* (K); Kedar Kantah Mts., *Jacquemont 830* (P); Kumaon, Lahai, 2,285 m, *Strachey & Winterbottom s.n.* (BM, BR, K); Garhwal, *Thomson 1254* (K).

PAKISTAN. KASHMIR: Kishenganga Valley, Rd. to Nanga Parbat via the Gangabal Lakes, Keran, 1,850 m, *Stewart & Stewart 17544* (NY, UC).

U.S.S.R. ALTAYSKIY KRAY: Teletskoe Ozero Tulkuy, *Koshurnikova & Vishniovskya*, 19 July 1927 (NY). KHABAROVSKIY KRAY: lower Amur River, at Dshare, *Maximowicz*, 18 July 1855 (K, LE). PRIMORSKIY KRAY: Kedrovaja Padj Reservation, SW of Vladivostok, *Gorovoy 8* (UC); S Ussuriysk, Pos'yet Dist., *Saberkin 890* (NY). SACHALIN: Sachalin Island, *Schmidt s.n.* (K).

Royle (1839) considered Himalyan populations of *Osmorhiza* conspecific with short-styled populations in North America now treated as *O. claytonii* (Michaux) C. B. Clarke, and accepted the name *O. brevistylis* DC. for them. Similarly, several authors (e.g., Gray, 1859; Hayata, 1911, 1912) incorrectly applied the name *O. longistylis* (Torrey) DC. to Asian populations of *Osmorhiza*.

Constance and Shan (1948) treated all the Asian representatives of *Osmorhiza* under *O. aristata*, although they distinguished two varieties on the basis of characters of the leaves. Quantitative evaluation of morphological characters, however, does not support the recognition of infraspecific taxa within *O. aristata* (Table 4).

3. *Osmorhiza claytonii* (Michaux) C. B. Clarke. *Myrrhis claytonii* Michaux, Fl. bor.-amer. 1: 170. 1803. *Chaerophyllum claytonii* (Michaux) Persoon, Syn. sp. pl. 1: 320. 1805. *Osmorhiza claytonii* (Michaux) C. B. Clarke in Hook., Fl. Brit. India 2: 690. 1879, sensu stricto (exclusive of *M. longistylis* Torrey

and *O. laxa* Royle). *Washingtonia claytonii* (Michaux) Britt. in Britt. & Brown, Ill. fl. 2: 530. 1897. *Scandix claytonii* (Michaux) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916. TYPE: U.S.A. "In montibus Alleghanis," A. Michaux s.n. (holotype, P!).

Scandix dulcis Muhlenb., Cat. pl. 31. 1813. *Myrrhis dulcis* (Muhlenb.) D. Eaton, Man. bot. 326. 1818. *Uraspermum dulce* (Muhlenb.) Farwell, Amer. Midl. Naturalist 9: 273. 1925. Based on *Myrrhis claytonii* Michaux.

Uraspermum hirsutum Bigelow, Fl. boston. 112. 1824. TYPE: U.S.A. Massachusetts: "Woods on the Concord turnpike," *Bigelow s.n.* (No type material has been located.)

Osmorhiza brevistylis DC., Prodr. 4: 232. 1830. *Myrrhis brevistylis* (DC.) Dietr., Syn. pl. 2: 984. 1840. *Uraspermum aristatum* α [subsp.] *brevistyle* (DC.) Kuntze, Revis. gen. pl. 1: 270. 1891. *Osmorhiza aristata* var. *brevistylis* (DC.) Boivin, Phytologia 17: 104. 1968. TYPE: U.S.A. New York: Range Co., West Point, *Torrey s.n.*, 1828 [lectotype (designated herein), G-DC!; isolectotypes (2), G-DC!].

Osmorhiza villosa Raf., Med. fl. 2: 249. 1830 (as "villosa"). (No type material has been located.)

Osmorhiza cordata Raf., Med. fl. 2: 249. 1830. (No type material has been located.)

Uraspermum aristatum α [subsp.] *brevistyle* [var.] *subintegrifoliolum* Kuntze, Revis. gen. pl. 1: 270. 1891. (No type material has been located.)

Uraspermum dulce var. *laevicaule* Farwell, Amer. Midl. Naturalist 9: 273. 1925. TYPE: U.S.A. Michigan: Oakland Co., Pontiac, *O. A. Farwell 5267* [lectotype, BLH! (designated by McVaugh et al., Bull. Cranbrook Inst. Sci. 34: 79. 1953); isolectotype, GH].

Osmorhiza claytonii f. *brevipilosa* Salamun, nom. inval. Although this name was provided with a Latin diagnosis and citation of a type specimen, it was never effectively published, having only been proposed in Salamun's dissertation (1950: 82).

Plants rather stout, 4–8(–10) dm high; stems 1–2(–3), erect to ascending, villous, villosulous, or sometimes essentially glabrous. Root system shallow, more or less horizontally spreading, the roots fibrous and limber, rank smelling, or sometimes weakly anise-scented. Leaves 2–3-ternate, more or less broadly ovate, 10–30 cm long, hirsutulous; leaflets ovate to lanceolate, (3–)4–8 cm long, 1.5–3 cm wide, acute or acuminate, serrate-dentate, often parted or divided at the base; petioles 5–15 cm long. Umbels loose; peduncles 2–3(–4), terminal and lateral, 3–10(–13) cm long; involucre wanting, or often composed of 1–2(–3) minute, linear-lanceolate, foliaceous, ciliate, reflexed bracts; rays ascending, 2–8(–10) cm long; umbellets 3–5 per umbel; involucl of 3–5 linear-lanceolate, attenuate, ciliate bractlets, each 2–5

(-6) mm long, 0.4-1 mm wide, strongly reflexed; pedicels 4-7(-8) per umbellet, ascending, those of the hermaphrodite flowers (4-)6-12(-15) mm long, those of the staminate flowers (3-)3.5-5.5 (-7) mm long. Hermaphrodite flowers 2-5(-7) per umbellet, (6-)10-23(-31) per umbel, staminate flowers 0-5(-6) per umbellet, (2-)7-17(-23) per umbel; corolla white, inconspicuous; styles (including stylopodium) 1-1.5(-1.7) mm long, stylopodium 0.5-0.75(-0.8) mm long, high-conic, lacking a conspicuous disc; pollen prolate in shape; carpophore cleft about one-fourth of its length. Fruit oblong-fusiform, tapering to a short, attenuate beak at the apex, concave furrowed, (10-)12-22(-25) mm long, the ribs sparingly to moderately hispid with retrorse bristles, especially toward the base, the caudate appendages 4.5-8.5 mm long. $n = 11$ (Bell & Constance, 1957). Figures 9b and 10.

Flowering period. April to early June.

Habitat. Dense to open deciduous forests, tending toward the more moist, lower areas.

Common names. Clayton sweetroot, Hairy sweet cicely, Sweet jarvil, Woolly sweet cicely.

Representative specimens. U.S.A. ALABAMA: Madison Co., Huntsville, *Baker*, 23 May 1897 (MO, NY). ARKANSAS: Independence Co., *Thomas* 38940 (TENN). CONNECTICUT: Hartford Co., Southington, *Bissell* 83 (NY). DELAWARE: Newcastle Co., near Wilmington, *Canby s.n.* (NY). GEORGIA: Union Co., *Duncan* 22356 (GA). ILLINOIS: Bureau Co., *Evers* 80102 (ILLS); Champaign Co., Brownfield Woods, N of Urbana, *Lowry* 1206 (ILL); Crawford Co., *Evers* 43236 (ILLS); Hancock Co., *M. J. Warnock* 215 (ILL); Lake Co., Waukegan, *Umbach* 5355 (UC); La Salle Co., Starved Rock State Park, *Greenman et al.* 27 (GH, NY, UC); Lee Co., 5 km NE of Franklin Grove, *G. N. Jones* 15839 (ILL, MO); Macon Co., Spittler Woods, Mt. Zion, *Lowry* 545 (ILL, MO, UC); McLean Co., Funk's Grove, S of Bloomington, *Lowry* 1140 (ILL); Piatt Co., Allerton Park, near Monticello, *Lowry* 211 (ILL, MO). INDIANA: Adams Co., 3 km W of Geneva, *Deam* 50233 (WIS); Fountain Co., Portland Arch, *Lowry* 1208 (ILL); Lake Co., 6 km N of Schneider, *Salamun*, 5 Aug. 1947 (ILL, UC); Morgan Co., 4.5 km N of Martinsville, *Heiser & Smith*, 19 May 1950 (ILL, MO, UC); Parke Co., Turkey Run St. Park, *Salamun*, 6 Aug. 1947 (ILL, MO). IOWA: Clayton Co., Pike's Peak, *McGregor, Shimek*, 8 Aug. 1922 (UC); Dickinson Co., 3 km W of Milford, *Thorne* 12907 (UC); Hardin Co., Iowa Falls, *Shimek*, 27 July 1922 (NY); Poweshiek Co., Grinnell, *M. E. Jones* 146 (GOET, NY). KANSAS: Doniphan Co., *Stephens* 58060 (KANU); Leavenworth Co., *A. S. Hitchcock* 701 (GH, NY, RM). KENTUCKY: Carter Co., Carter Caves, *Brown* 3940 (NY); Greenup Co., 4.5 km from Boyd Co. line, *Smith et al.* 3580 (F, GH, NY, US); Madison Co., Berea, *McFarland* 4294 (MO). MAINE: Aroostook Co., Fort Fairfield, *Fernald* 51 (GH, MO, UC, US); Knox Co., Camden, *LeBean*, 13 Aug.

1913 (WIS); Somerset Co., Skowhegan, *Eaton*, 30 June 1903 (LL, NY). MARYLAND: Alleghany Co., Cumberland, *Shriver s.n.* (NY); Howard Co., Ellicott City, *Arsène*, 26 July 1916 (MO). MASSACHUSETTS: Berkshire Co., Florida, Deerfield River, *Fernald & Long* 10088 (GH); Middlesex Co., Malden, *Manning*, 6 June 1881 (NY). MICHIGAN: Arenac Co., Mud Lake, *Sharp et al.*, 20 June 1961 (MICH); Baraga Co., Big Limestone Mt., L'Anse, *Fassett* 21037 (WIS); Cheboygan Co., Monroe Lake, *Ehlers* 374 (GH, MO, US); Emmet Co., 3 km W of Mackinac City, *McVaugh* 9430 (MICH, MO); Gogebic Co., Gogebic Lake, *Fassett* 19868 (F, MO, NY, WIS); Keweenaw Co., Isle Royale, *McFarlin* 2149 (MONT); Schoolcraft Co., 8 km W of Gulliver, *Salamun*, 19 July 1946 (ILL, MO). MINNESOTA: Clearwater Co., Floating Bog Bay, *Grant* 2885 (MO, NY, UC); Cook Co., Mineral Center, *Rosendahl & Butters* 4590 (GH, NY); Saint Louis Co., Duluth, *Lakela* 2071 (MO); Wabasha Co., 10 km N of Reed's Landing, *M. J. Warnock* 1463 (ILL). MISSOURI: Dallas Co., 8 km SW of Bennett Springs, *Conrad* 3496 (MO); Jackson Co., Kansas City, *MacKenzie*, 16 May 1896 (MO, NY); Lincoln Co., Mill Creek, NE of Silex, *Steyermark* 25981 (F, MO); Mercer Co., 5 km NE of Saline, *Palmer & Steyermark* 41324 (MO, NY). NEBRASKA: Cherry Co., NE of Valentine, *Churchill* 4489 (MO, NY); Cuming Co., 2 km SSE of Beemer, *Churchill* 5516 (NY); Dixon Co., Ponca, *Clements* 2564 (US); Washington Co., 4 km NW of Fort Calhoun, *Churchill* 5328 (MO). NEW HAMPSHIRE: Grafton Co., N Woodstock, *Fernald* 386 (BM, BR, CAS, DUKE, G, GH, K, KANU, MO, MONTU, NY, PH, RM, SMU, TENN, TEX, UC, US). NEW JERSEY: Bergen Co., W of Englewood, *Wilson*, 19 July 1915 (NY); Middlesex Co., Plainfield, *Tweedy s.n.* (GOET). NEW YORK: Chautauqua Co., Bemus Pt., Lake Chautauqua, *Churchill*, 2 Aug. 1896 (MO); Greene Co., Big Hollow, *Barnhart* 2344 (NY); Herkimer Co., 2.5 km SW of Dart Lake, *Smith* 2504 (WIS); Monroe Co., Webster, *Matthews* 4164 (UC); Rensselaer Co., Brunswick, *House* 26381 (MO, TEX). NORTH CAROLINA: Haywood Co., 1 km NW of Blue Ridge Parkway on Hwy. 276, *Lowry* 1153 (ILL); Macon Co., 1 km SW of Swain Co. line on Hwy. 19, *Lowry* 1152 (ILL, UC); McDowell Co., Curtis Creek, Blue Ridge Parkway, 825 m, *Feddema* 3028 (RM); Mitchell Co., 1 km S of Hwy. 226 on Penland Rd., *Lowry* 1155 (ILL, NY, UC); Yancey Co., 6 km NNW of Swiss, *Lowry* 1154 (ILL, NY, UC). NORTH DAKOTA: Cass Co., Fargo, *Shunk s.n.* (MONT); Grand Forks Co., *Facey s.n.* (reported in *Rhodora* 74: 393. 1972); Richland Co., Leonard, *Stevens* 1343 (UC). OHIO: Athens Co., Athens, *Abbot* 12 (NY); Cuyahoga Co., Cleveland, *Greenman* 710 (GH, MO); Scioto Co., Camp Gordon, Friendship, *Demaree* 10648 (CAS, GH, MO). PENNSYLVANIA: Armstrong Co., *Wahl* 5315 (PH); Berks Co., 3 km W of Upper Black Eddy, *Fogg* 8665 (MO, PH); Cambria Co., *Wahl* 9325 (PH); Juniata Co., 1.5 km NE of Peru Mills, *Wahl* 10730 (PH, UC); Warren Co., *Pohl* 2461 (PH). SOUTH CAROLINA: McCormick Co., *Radford* 31734 (KANU). SOUTH DAKOTA: Brookings Co., Warren's Woods, *Williams & Thordaer*, 17 June 1893 (MO); Roberts Co., Big Stone Lake, *Over* 14409 (US). TENNESSEE: Davidson Co., Joelton, *Svenson* 93 (GA, GH); Grainger Co., *Sharp* 43562 (TENN); Greene Co., Paint Creek, 1 km N of French Broad River, *Bufford et al.* 18119 (MO); Knox Co., Lane's Creek, *Sharp & Herster* 584 (NY).

VERMONT: Bennington Co., Red Mt., Arlington, *Seymour* 21637 (MO); Caledonia Co., Peacham, *Blanchard s.n.* (MO, NY, RM); Windsor Co., Norwich, *Brown*, 20 June 1946 (SMU). VIRGINIA: Fairfax Co., Dead Run, E of Plummer's Island, *Mathias* 1394 (CAS, GH, MO, NY, RM, UC, US); Smythe Co., Chatham Hill Gap, Walker Mt., *Small*, 13 June 1892 (F, GH, K, MO, P, UC). WEST VIRGINIA: Cabell Co., Pleasant Valley, *Williams & Gilbert* 445 (F, GH, MO, NY, SMU); Ohio Co., Wheeling, *Mertz* 1042 (F). WISCONSIN: Bayfield Co., *Koch* 6881 (KANU); Forest Co., 13 km SE of Crandon, *Stearns*, 23 June 1946 (NY); Green Co., 7 km W of Monticello, *Salamun*, 10 July 1947 (ILL, MO); Outagamie Co., Appleton, *Chandler*, 18 May 1896 (UC); Trempealeu Co., NW of Trempealeu, *M. J. Warnock* 1470 (ILL).

CANADA. MANITOBA: 6.5 km N of Moon Lake, Riding Mt. Nat'l. Park, *Mosquin* 6066 (DAO). NEW BRUNSWICK: Albert Co., *Roberts* 64-1655 (DAO); Charlotte Co., Grand Manan, *Weatherby & Weatherby* 7327 (GH, US); Westmorland Co., *Dore* 45235 (DAO). NOVA SCOTIA: Cape Breton Co., George River, *Bissell & Linder* 22050 (GH); Inverness Co., Mabou, *Robinson* 229 (NY); Victoria Co., Dingwall, Aspy Bay, *Churchill*, 7 July 1909 (MO). ONTARIO: Algoma Dist., *Jenkins* 4909 (DAO); Carleton Co., March Twp., *Cody & Calder* 480 (DAO, MO); Grey Co., 1.5 km SW of Meadford, *Soper & Shields* 4603 (MO); Manitoulin Dist., Manitoulin Island, *Salamun*, 20 July 1946 (ILL, MO); Muskoka Dist., *Minshall* 3076 (DAO); Oxford Co., 8 km N of Thamesford, *Soper & Shields* 4506 (MO); Rainy River Dist., 8 km below Rainy River, *Garton* 8670 (DAO); Russell Co., 1.5 km SE of Cumberland, *Cody & Calder* 540 (DAO, MO); Thunder Bay Dist., N bank of Pigeon River at Middle Falls, *Garton* 2062 (DAO, MO). PRINCE EDWARD ISLAND: Kings Co., Bear River, *Fernald & St. John* 11140 (GH, UC, US). QUÉBEC: Charlevoix Co., *Cayouette* 57-247 (DAO); Gaspé Co., *Lemieux* 14506 (DAO); Labelle Co., Bellerive, *Lucien & Eloi* 248 (UC); Lac-Saint-Jean Co., *Bouchard* 70-520 (DAO, G); Matane Co., Mt. Nicolabert, *Joffre, Fernald & Pease* 25208 (GH); Richmond Co., Cleveland, *Chamberlain & Knowlton*, 26 July 1923 (GH, MO); Terrebonne Co., Lac Tremblant, *Churchill*, 8 Aug. 1922 (GH, MO).

Osmorhiza claytonii is fairly uniform throughout its range. Although two varieties have been recognized on the basis of quantity of pubescence (Farwell, 1925; Salamun, 1950), examination of a large number of specimens shows that this character is highly variable in *O. claytonii*, and neither variety is upheld in this interpretation (Lowry, 1976; Lowry & Jones, 1979a).

4. *Osmorhiza longistylis* (Torrey) DC. *Myrrhis longistylis* Torrey, Fl. U.S. 310. 1824. *Osmorhiza longistylis* (Torrey) DC., Prodr. 4: 232. 1830. *Uraspermum aristatum* β [subsp.] *longistyle* (Torrey) Kuntze, Revis. gen. pl. 1: 270. 1891, pro parte (with two varieties: *laciniatum* and *subintegrifoliolum*). *Washingtonia longistylis* (Torrey) Britt. in Britt.

& Brown, Ill. fl. 2: 530. 1897. *Scandix longistylis* (Torrey) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916. *Osmorhiza aristata* var. *longistylis* (Torrey) Boivin, Phytologia 17: 104. 1968. TYPE: "In wet meadows near Albany, N.Y. Tracy. Near Geneva, N.Y. Paine. June. Near Hudson, N.Y. Alsop & c." (None of these syntypes has been located; only an authentic specimen collected by Paine was found, and is herein designated the neotype.) Canada. Québec: Montreal, *Paine s.n.* (neotype, NY!).

- Osmorhiza dulcis* Raf., Med. fl. 2: 249. 1830. *Myrrhis dulcis* (Raf.) Raf., Good book 53. 1840, pro syn., non *Scandix dulcis* Muhlenb. TYPE: U.S.A. "Mts Alleghy [sic]," *Rafinesque s.n.* [lectotype, PH! (designated by Lowry & Jones, Amer. Midl. Naturalist 101: 26. 1979); possible syntypes, G!, PH!].
- Chaerophyllum dulce* Fischer ex Steud., Nom. bot. 1: 339. 1841, non *Scandix dulcis* Muhlenb. (No type material has been located.)
- Osmorhiza claytonii* (Michaux) C. B. Clarke in Hook., Fl. Brit. India 2: 690. 1879, pro parte (exclusive of *M. claytonii* Michaux and *O. laxa* Royle).
- Osmorhiza aristata* (Thunb.) Rydb., Bot. surv. Nebr. 3: 37. 1894, pro parte (exclusive of Asian populations).
- Osmorhiza longistylis* var. *villicaulis* Fern., Rhodora 10: 52. 1908. *Washingtonia longistylis* var. *villicaulis* (Fern.) Coult. & Rose, Contr. U.S. Natl. Herb. 12: 443. 1909. *Uraspermum aristatum* var. *villicaulis* (Fern.) Farwell, Pap. Michigan Acad. Sci. 1: 96. 1931. TYPE: U.S.A. Pennsylvania: Lancaster Co., On the Conestoga, near Binkley's Bridge, in limestone, *A. A. Heller*, 21 June 1901 (holotype, GH!; isotypes, F!, G!, US!).
- Osmorhiza longistylis* var. *brachycoma* S. F. Blake, Rhodora 25: 110. 1923. *Washingtonia longistylis* var. *brachycoma* (S. F. Blake) House, Bull. New York State Mus. 254: 529. 1924. TYPE: U.S.A. Maryland: Montgomery Co., slope in woods, vicinity of Cabin John, *S. F. Blake* 6902 (holotype, US!; isotypes, GH!, TEX!).
- Osmorhiza longistylis* var. *imbarbata* Salamun, Amer. Midl. Naturalist 47: 253. 1952. TYPE: U.S.A. South Dakota: Lawrence Co., wooded bank of Spearfish Creek in Spearfish Canyon, approx. ¼ mi. N of Bridal Veil Falls, *A. L. Thorne*, 12 Aug. 1949 (holotype, UWM!; isotype, WIS!).

Plants rather stout, 6–10(–12) dm high; stems 1–2(–3), erect, densely pilose to villous, or often glabrous. Root system rather deep, tending toward vertical orientation, the roots carnosous (breaking with a snap), with a sweet, anise-like scent. Leaves 2–3-ternate, broadly ovate, 8–25 cm long, sparsely hirsutulous (especially along the veins) to glabrescent; leaflets ovate to oblong-lanceolate, (3–)4–10 cm long, 2–5 cm wide, acute, serrulate-dentate, often incised or parted at the

base; petioles 5–16 cm long. Umbels loose; peduncles 2–4, terminal and lateral, 5–13 cm long; involucre composed of 1–3(–4) linear to lanceolate, foliaceous, ciliate, reflexed bracts, each 5–10(–15) mm long, 1–1.5 mm wide; rays ascending, 1.5–5(–7.5) cm long; umbellets 4–6(–8) per umbel, (0–)1–3 of them producing only staminate flowers; involucre of 4–6 linear-lanceolate to ovate, acuminate, ciliate bractlets, each 2.5–6(–7) mm long, 0.7–1.8 mm wide, often strongly reflexed; pedicels (7–)9–18 per hermaphrodite umbellet, 3–13 per staminate umbellet, ascending, those of the hermaphrodite flowers 4–8(–9) mm long, those of the staminate flowers 3–6(–8) mm long. Hermaphrodite flowers (2–)3–5(–7) per umbellet, (8–)15–30(–33) per umbel, staminate flowers (3–)4–10(–14) per hermaphrodite umbellet, (23–)35–75(–86) per umbel; corolla white, showy; styles (including stylopodium) 2–3.6 mm long, stylopodium (0.4–)0.5–0.75(–0.8) mm long, high-conic, lacking a disc; pollen prolate in shape; carpophore cleft about one-third of its length. Fruit oblong-fusiform, acute at the apex, concave furrowed, (10–)15–21(–22) mm long, the ribs sparsely to moderately hispid with retrorse bristles, especially toward the base, the caudate appendages 4–8 mm long. $n = 11$ (Wanscher, 1932; Bell & Constance, 1957). Figures 9c and 11.

Flowering period. April to early June.

Habitat. Dense to open deciduous forests, tending toward the somewhat drier, upland sites.

Common names. Aniseroot, Longstyle sweetroot, Smoother sweet cicely.

Representative specimens. U.S.A. ALABAMA: Lauderdale Co., Murchison 171 (AUA); Tuscaloosa Co., Warrior River above Hurricane Creek, Harper 144 (F, GH, MO, NY, US). ARKANSAS: Benton Co., Sulphur Springs, Demaree 4974 (GH, MO, ORE); Fulton Co., Salem, Demaree 26020 (UC); Hot Springs Co., Dripping Springs, Scully 938 (UC, US). COLORADO: Boulder Co., Boulder Creek, Boulder, 1,615 m, Weber, 7 July 1954 (CAS, UC); Larimer Co., Spring Canyon, Osterhout 344 (F, RM). CONNECTICUT: Fairfield Co., vicinity of Green's Farm, Pollard 37 (US); Newcastle Co., Mt. Cuba, Commons, 25 June 1875 (MO). DISTRICT OF COLUMBIA: Washington, Pennell 15005 (PH). GEORGIA: Burke Co., Shell Bluff, Pyron & McVaugh 2494 (GA); Decatur Co., Anderson, 30 May 1923 (RM). ILLINOIS: Champaign Co., Brownfield Woods, N of Urbana, Lowry 1207 (ILL); Cook Co., Riverdale, Greenman 2627 (GH, MO); Du Page Co., Naperville, Umbach, 8 June 1897 (MO); Grundy Co., Evers 72569 (ILLS); Jo Daviess Co., Apple River Canyon State Park, G. N. Jones 15859 (ILL, MO); Macon Co., Spittler Woods, Mt. Zion, Lowry 546 (ILL, MO, NY, UC); McLean Co., Funk's Grove, S of Bloomington, Lowry 1141 (ILL); Piatt Co., Allerton Park, near Monticello, Lowry

540 (ILL). INDIANA: Blackford Co., Hartford, Deam 1078 (NY); Lawrence Co., Bedford, Kriebel 1879 (DUKE, GH); Marion Co., Indianapolis, Friesner 16672 (GH, MO, NY, SMU). IOWA: Appanoose Co., Sedan, Shimek, 15 May 1902 (WIS); Decatur Co., Fitzpatrick & Fitzpatrick, 24 May 1897 (F, GH, NY); Dubuque Co., NW of Luxembourg, Shimek, 2 July 1929 (UC); Emmet Co., Ft. Defiance St. Park, Hayden 9426 (MO); Poweshiek Co., Grinnell, M. E. Jones 145 (CAS, GOET, NY, RM, UC). KANSAS: Cowley Co., Winfield, White s.n. (MO); Crawford Co., Holland 381 (KANU); Douglas Co., N of Baldwin, Croat 116 (MO); Greenwood Co., Stephens 2925 (KANU, SMU); Miami Co., 5 km SSE of Fontana, Brooks et al. 11958 (KANU, MO); Riley Co., Norton 700 (GH, MO, NY, P, RM, US). KENTUCKY: Fayette Co., Lexington, Short s.n. (GH, NY, UC); Greenup Co., Big Woods, 4.5 km from Boyd Co. line, Smith et al. 3581 (GH, NY, US); Henry Co., 0.5 km N of Sand Ripple Creek, Gentry 932 (NY); Union Co., McCotrey School, Shacklett 296 (GH, NY, SMU). MAINE: Aroostook Co., Ft. Fairfield, Fernald 2020 (GH); Hancock Co., Somersville, Rand, 18 July 1898 (UC); Kennebec Co., Vassalboro, Chamberlain s.n. (GH). MARYLAND: Montgomery Co., near Cabin John, Blake 6902 (GH, TEX, US). MASSACHUSETTS: Berkshire Co., New Marlboro, Churchill, 13 June 1919 (GH, MO); Essex Co., Manchester, Chamberlain s.n. (NY); Suffolk Co., Waverley, Andrews, 16 July 1892 (ILL). MICHIGAN: Emmet Co., 3 km S of Good Hart, McVaugh 9282 (MICH, MO); Houghton Co., Calumet, Minns s.n. (GH); Ingham Co., E Lansing, Lowry 1142 (ILL); Monroe Co., 5.5 km SE of Milan, Robertson 109 (MICH). MINNESOTA: Chisago Co., Center City, Taylor s.n. (NY, RM, UC, US); Clearwater Co., Bear Pt., Moyle 207 (GH, NY, UC, US); Pipestone Co., Pipestone Nat'l. Mon., Moore 23192 (ILL). MISSISSIPPI: Desoto Co., Kral 8582 (LAF); Oktibbeha Co., Lott, 28 Apr. 1940 (MISSA); Tate Co., Pullen 64276 (GA, LAF). MISSOURI: Franklin Co., Arboretum, Gray's Summit, Sharp 201 (MO); Grundy Co., 20 km W of Spickard, Steyermark 11036 (MO); Harrison Co., W of Blythedale, Steyermark 10976 (MO); Osage Co., S of Meta, Steyermark 69719 (MO); Phelps Co., 6.5 km SE of St. James, Steyermark 22179 (UC); Wright Co., 3 km SE of Cedar Gap, Steyermark 23671 (F, MO, UC). MONTANA: Cascade Co., Lower Falls of the Missouri, Williams 275 (MONT, US); Fallon Co., Stephens 23282 (KANU); Gallatin Co., Bozeman, Blankinship, 18 Aug. 1898 (MO, MONT); Stillwater Co., Absarokee, Hawkins, 15 June 1918 (MONT, WIS). NEBRASKA: Antelope Co., Neligh, Harper, 30 May 1888 (WIS); Blaine Co., Halsey, Pool, 21 June 1912 (MO); Richardson Co., 1.5 km N of Barada, Reynolds 3085 (MO, UC); Sioux Co., Stephens 16324 (KANU). NEW HAMPSHIRE: Cheshire Co., Walpole, Fernald 417 (GH); Grafton Co., Lebanon, Kennedy s.n. (GH). NEW JERSEY: Burlington Co., Vincetown, Long 9790 (GH); Gloucester Co., Fogg 8564 (PH). NEW YORK: Delaware Co., N Harpersfield, Topping 166 (ILL, US); Erie Co., Buffalo, Kinnicutt s.n. (NY); St. Lawrence Co., Hermon, Phelps 1615 (GH, US); Suffolk Co., Cold Spring Harbor, Banker 2830 (NY). NORTH CAROLINA: Buncombe Co., 1.5 km N of Warren Wilson College, Lowry 1144 (ILL); Lee Co., 3 km E of Moncure, Wood 857 (MO). NORTH DAKOTA: Benson Co., Peninsula of Lake Ibsen, Lunell, 8 July 1901 (NY, RM, US); Golden Valley Co., Ste-

phens 49996 (KANU); Pembina Co., 3 km S of Wallaha, *Willenbring* 687 (MO); Richland Co., Abercrombie, *Bergman* 1769 (MO, RM, UC). OHIO: Coshocton Co., *Moldenke* 12516 (UC); Hamilton Co., Madisonville, Cincinnati, *Lowry* 547 (ILL, MO, UC); Preble Co., Devil's Backbone, near Camden, *Cobbe* 104 (CAS, G, UC); Scioto Co., Camp Gordon, Friendship, *Demaree* 10647 (CAS, GH, MO, SMU, UC). OKLAHOMA: Canadian Co., Devil's Canyon, *Goodman* 5060 (OKL, UC); Cherokee Co., Camp Egan, E of Tahlequah, *Goodman* 6811 (ILL, OKL); Creek Co., Sapulpa, *Bush* 1059 (MO, NY). PENNSYLVANIA: Berks Co., *Berkheimer* 16799 (PH); Centre Co., *Wahl* 17229 (PH); Elk Co., Benezett, *Wahl* 612 (GH, PH); Lancaster Co., Conestoga, near Binkley's Bridge, *Heller*, 21 June 1901 (F, G, GH, US); Lebanon Co., *Wherry*, 23 May 1959 (PH); Somerset Co., *Pohl* 5574 (PH). RHODE ISLAND: Kent Co., Warwick, *Congdon*, 3 June 1877 (CAS, MO, NY). SOUTH CAROLINA: Lancaster Co., *Bozeman et al.* 8812 (AUA, CAS). SOUTH DAKOTA: Brookings Co., Brookings, *Thornber*, 4 July 1894 (ARIZ, MO, UC); Harding Co., W Short Pines, *Visher* 459 (F); Lawrence Co., NW of Whitewood, *D'Arcy* 5761 (MO). TENNESSEE: Cheatham Co., 2.5 km SE of Ashland City, *Kral* 26796 (UC); Davidson Co., Nashville, *Gattinger*, 28 Mar. 1886 (F, NY, US); Knox Co., Knoxville, *Ruth* 442 (MO); Lauderdale Co., *Sharp* 12115 (TENN). TEXAS: Tarrant Co., near Trinity, *Ruth* 601 (NY, PH, US). VERMONT: Addison Co., Hancock, *Dutton*, 21 June 1914 (GH, MO); Lamoille Co., *Knowlton*, 18 July 1916 (PH). VIRGINIA: Fairfax Co., Dead Run, E of Plummer's Island, *Mathias* 1392 (CAS, GH, MO, NY, UC, US); James City Co., 1 km W of Williamsburg, *Grimes* 3601 (NY); Surry Co., Claremont Wharf, *Fernald & Long* 8386 (GH, MO, NY, US). WEST VIRGINIA: Berkeley Co., E of Martinsburg, *A. G. Jones* 4218 (ILL, MO); Monongalia Co., Monongahela River, *Millspaugh* 173 (NY); Pendleton Co., Snowy Mt., *Rydberg* 9122 (NY, PH). WISCONSIN: Brown Co., Green Bay, *Schuetz s.n.* (F); Iowa Co., Blue Mounds, *Clikenian et al.*, 5 June 1932 (UC); Walworth Co., *Salamun*, 12 June 1948 (ILL, MO). WYOMING: Big Horn Co., mouth of Dry Medicine Lodge Canyon, *Dueholm* 9509 (RM); Converse Co., 16 km W of Douglas, *Porter* 4511 (RM, WTU); Sheridan Co., Tongue Creek, N of Big Horn Coal Mine, 1,100 m, *Brink* 1366 (ILL, MO).

CANADA. ALBERTA: Edmonton Dist., *Moss* 6363 (DAO); Medicine Hat Dist., *Macoun* 858 (GH). MANITOBA: Portage-la-Prairie Dist., 4 km N of Portage-la-Prairie, *Boivin & Breitung* 6501 (DAO); Winnipeg Dist., Winnipeg, *Johnson* 1044 (NY); Thalberg, *Krivda*, 20 June 1960 (NY); Ft. Ellice, *Macoun & Herriot* 77116 (NY). NEW BRUNSWICK: Charlotte Co., St. Andrews, *Malte* 843/29 (GH, US); Kent Co., Bass River, *Fowler*, 5 July 1873 (GOET, WIS). NOVA SCOTIA: Cumberland Co., Clay, Green Bay, *Roland* 41583 (GH). ONTARIO: Elgin Co., St. Thomas, *Macoun* 81729 (F); Hastings Co., Ox Pt., near Point Anne, *Soper & Shields* 4656 (MO); Manitoulin Dist., S of Little Current, Manitoulin Island, *Fernald & Pease* 3448 (GH, US); Northumberland Co., Presqu'île Park, *Soper & Shields* 4644 (MO); Rainy River Dist., *Garton* 8862 (DAO); Thunder Bay Dist., Middle Falls on Hwy. 61, *Garton* 2061 (MO, DAO). QUÉBEC: Bonaventure Co., Restigouche River, Matapedia, *Collins & Fernald s.n.* (GH); Gaspé Co., *Dansereau* 180 (DAO); Missisquoi Co., *Perron*

66-210 (DAO); Papineau Co., *Charlebois* 1401 (DAO); Temiscamingue Co., Baie Girard, *M.-Victorin* 8546 (GH, NY); Vaudreuil Co., *Gervais & Lavigne*, 4 June 1959 (G). SASKATCHEWAN: Assiniboia Dist., 8 km S of Estevan, *Boivin & Perron* 11823 (DAO); Lake Centre Dist., Outlook, *Boivin & Alex* 9848 (DAO); Moose Jaw Dist., Moose Jaw Creek, *Macoun* 858 (NY); Qu'Appelle Dist., Qu'Appelle River, near mouth of Lake Katepwe, *Boivin & Dore* 7591 (DAO); Wood Mountain Dist., 1.5 km S of Willowbunch, *Boivin & Gillett* 8838 (DAO); Cypress Hills, *Breitung* 4719 (DAO, MO).

Clarke (1879) included all representatives of *Osmorhiza* sect. *Osmorhiza*, including *O. longistylis*, under the name *O. claytonii*. Similarly, MacMillan (1892) included *O. longistylis* in his broad concept of *Myrrhis aristata*.

Three varieties of *Osmorhiza longistylis* have been described on the basis of quantity of indument (Fernald, 1908; Blake, 1923; Salamun, 1952). This character, however, was found to be highly variable, and thus none of these infraspecific taxa is upheld (Lowry, 1976; Lowry & Jones, 1979a).

Iib. *Osmorhiza* Raf. sect. *Mexicanae* Constance & Shan ex Lowry & Jones, sect. nov.
TYPE: *O. mexicana* Griseb.

Uraspermum § *Osmorhiza* (Raf.) Kuntze, *Lexicon* 582. 1904, pro parte.

Scandix 2. [sect.] *Urascandix* Koso-Polj., *Bull. Soc. Imp. Naturalistes Moscou* 29: 143. 1916, pro parte.

Osmorhiza sect. *Mexicanae* Constance & Shan, *Univ. Calif. Publ. Bot.* 23: 112. 1948, nom. nud.

Involucrum deficiens vel interdum ab 1-3 bracteis parvis constitutum; bractee lineares vel lanceolatae, patentes vel reflexae; involucellum generaliter conspicuum, ab 1-6 bracteolis patentibus vel reflexis constitutum. Styli cum stylopodio 0.5-1.2 mm longi, stylopodium 0.2-0.8 mm longum, depresso-conicum vel depressum, saepe cum disco conspicuo.

Involucre wanting, or sometimes composed of 1-3 small, linear to lanceolate, spreading to reflexed bracts; involucl generally conspicuous, composed of 1-6 spreading to reflexed bractlets. Styles plus stylopodium 0.5-1.2 mm long, stylopodium 0.2-0.8 mm long, low-conic to somewhat depressed, often with a conspicuous disc.

5. *Osmorhiza glabrata* Philippi. *Linnaea* 28: 653. 1856. *Uraspermum glabratum* (Philippi) Kuntze, *Revis. gen. pl.* 1: 270. 1891. TYPE: Chile. Bío-Bío: Santa Barbara, *C. Gay* 1514 (holotype, SGO; photograph of holotype, ILL!; 2 possible syntypes, P!).

Myrrhis renjifoana Philippi, *Anales Univ. Chile* 85: 725. 1894. *Elleimataenia renjifoana* (Philippi)

Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 164. 1916. TYPE: Chile. Ñuble: Valle de las Nieblas, near Termas de Chillán, *F. Philippi 2029* [lectotype, SGO (no. 053472) (designated herein); photographs of lectotype, ILL!, UC!; isolectotype, SGO (no. 041594); photograph of isolectotype, ILL!].

Plants rather stout, 1–5(–6) dm high; stems 1–2(–3), erect to ascending, villous to hirsutulous, or often glabrate. Root system rather deep. Leaves ternate-bipinnate, deltoid to broadly ovate, 5–10(–13) cm long, glabrous to hirsutulous along the veins and rachis; leaflets ovate, (0.6–)1–2.5 cm long, 0.5–1.3 cm wide, acute or acuminate, laciniately lobed to parted or pinnatifid, with linear to linear-lanceolate ultimate divisions; petioles 4–12(–15) cm long. Umbels loose to somewhat congested; peduncles 2–3, terminal and lateral, 4–15 cm long; involucre wanting; rays stiffly ascending, (1–)1.5–8(–9.5) cm long; umbellets (2–)3–13 per umbel, 1–5(–6) of them producing only staminate flowers; involucre of (0–)1–2 minute, linear, acuminate, ciliate bractlets; pedicels (3–)4–10 per hermaphrodite umbellet, 2–7(–9) per staminate umbellet, stiffly ascending to nearly erect, those of the hermaphrodite flowers (2–)2.5–7(–9) mm long, those of the staminate flowers (1.5–)2.3–5.5(–6.5) mm long. Hermaphrodite flowers (2–)3–5 per umbellet, (9–)15–30(–38) per umbel, staminate flowers (1–)2–7(–9) per hermaphrodite umbellet, (7–)15–60(–82) per umbel; corolla white, rather inconspicuous; styles (including stylopodium) (0.9–)1–2 mm long, stylopodium 0.4–0.8 mm long, conic, often with a conspicuous disc; carpophore cleft about one-third of its length. Fruit linear-fusiform, acute at the apex, concave furrowed, (13–)14–20(–25) mm long, the ribs glabrous to moderately hispid with retrorse bristles, especially toward the base, the caudate appendages (0.5–)1–6 mm long. Figures 9e and 18.

Flowering period. November to January (with one collection flowering in April).

Habitat. Seasonally moist *Nothofagus* forests to open, grassy slopes.

Common names. Glabrate sweet cicely, Andean sweet cicely.

Representative specimens. ARGENTINA. NEUQUÉN: Dept. Alumine, Parque Nac. Lanin, *Eskuche 934* (UC, WIS); Dept. Huiliches, Lago Huechu-Lauquen, *Castellanos 20507* (UC); Dept. Minas, Lagunas Epu-lauquén, 1,600 m, *Boelcke et al. 10978* (UC); Cordillera del Viento, 2,000 m, *Boelcke et al. 11634* (UC); Paso del Macho, 2,280 m, *Boelcke et al. 13955* (UC).

CHILE. BÍO-BÍO: Sierra Velluda, *Poeppig 905* (BM, BR,

P). COLCHAGUA: San Fernando, Termas Vegas del Flaco, *Montero O. 1214* (UC); Cordillera de Colchagua, *Pirian 159* (GH); Termas Vegas del Flaco, 2,500 m, *Zöllner 6430* (UC). CURICÓ: hills NE of Los Baños, the Quebrada, 2,500 m, *Aravena 33301* (UC). MALLECO: Volcán Longuimay, 1,500 m, *Constance & Sparre 3580* (BM, F, K, MICH, MO, UC, US). ÑUBLE: Cordillera de Chillán, *Germain s.n.* (BM, G, K); Termas de Chillán, *Jaffuel 2035, 2837, 3715, 3717* (GH); Chillán, *Philippi 3469* (UC), *s.n.* (photographs in F, NY, UC); Baños de Chillán, *Philippi & Borchers s.n.* (BM), *Werdermann 1571* (NY). SANTIAGO: Río Yeso, Lag. Pinquenes, 2,500 m, *Biese 800* (NY); Maipo, 2,700 m, *Claude-Joseph 2964* (US); Valdes-Tál (Volcántal), 2,800 m, *Grandjot s.n.* (MO, UC); valley of the Maipo River, near the Baths of Colima, 2,500 m, *Zöllner 9715* (ILL). TALCA: Laguna Maule, 2,400 m, *Zöllner 5824* (ILL, UC).

Constance and Shan (1948: 120) included *Osmorhiza glabrata* in their section "*Glycosmae*" [= subg. *Glycosma*], stating that it "is remarkable for its close similarity to *O. occidentalis*." However, the evidence presented here indicates that the affinities of *O. glabrata* lie with *O. mexicana* and *O. brachypoda* rather than with *O. occidentalis*. Constance and Shan's misinterpretation of the relationships of *O. glabrata* most likely was due to the very limited amount of material available to them; they cite only nine collections of this species.

In their discussion of *Osmorhiza glabrata*, Constance and Shan (1948: 120) stated, "The single specimen which caused the most trouble in our study was *Pennell 12487*, from Baños de Chillán, Chile. This plant has the ascending rays and long styles of *O. glabrata*, but combines these characteristics with appendaged, bristly fruit and subentire leaflets. We can only suggest that the plant in question may be the result of interbreeding between *O. glabrata* and either *O. chilensis* or *O. obtusa* [= *O. depauperata*], all three of which occur in this area." We have tried to locate this specimen, with no success. In any case, Pennell's collection is probably much less of an anomaly than Constance and Shan suggest; appendaged, bristly fruits are not at all uncommon in *O. glabrata*. Furthermore, the occurrence of subentire leaflets in this specimen may represent an intermediate between *O. glabrata* and *O. mexicana* subsp. *mexicana*. The existence of such a specimen would not be surprising because the two taxa involved are closely related, and this would seem a more reasonable explanation than assuming intersectional hybridization involving *O. glabrata* and either *O. chilensis* or *O. depauperata*.

Clos (1848) erroneously interpreted plants referable to *O. glabrata* as being conspecific with the European species *Myrrhis odorata* (L.) Scopoli.

6. *Osmorhiza mexicana* Griseb., Abh. Königl. Ges. Wiss. Göttingen 24: 147. 1879. *Washingtonia mexicana* (Griseb.) Rose, Contr. U.S. Natl. Herb. 8: 337. 1905. TYPE: Mexico. *Schaffner 37* (holotype, GOET!; 2 isotypes, P!). [The holotype is marked in Grisebach's hand "*Osmorhiza mexicana m(ihi)*," his annotation for the original material. Several paratypes in GOET are marked by Grisebach "*Osmorhiza mexicana* Gr.," his annotation for authentic material other than the holotype, according to G. Wagenitz, GOET.]

Uraspermum aristatum β [subsp.] *brevistyle* var. *laciniatum* Kuntze, Revis. gen. pl. 1: 270. 1891. (No type material has been located.)

Plants slender to rather stout, stems 1–2, erect to ascending. Leaves 2–3-ternate, villous or pilose, especially on the veins below; leaflets ovate to ovate-oblong, acute to acuminate, coarsely serrate to divided at the base. Umbels loose to rather open; involucre wanting, or often composed of 1–2 linear, foliaceous, ciliate bracts; rays spreading-ascending; involucre of 1–4 linear, acuminate, ciliate bractlets. Styles (including stylopodium) 0.5–1 mm long, stylopodium low-conic to somewhat depressed, often with a disc; carpophore cleft about one-fourth of its length. Fruit variable in characters depending on the subspecies.

At their morphological extremes, the two subspecies of *Osmorhiza mexicana* (subsp. *mexicana* and subsp. *bipatriata*, see below) are very distinct. A number of truly intermediate collections, however, mark a transition between these taxa. For example, two specimens from Cerro Potosí, Nuevo León, Mexico (*C. H. Mueller 2231* and *R. A. Schneider 1108*) have fruit with retrorse bristles and short caudate appendages, but are in most other respects similar to specimens of the subspecies *bipatriata*. Typical representatives of both subspecies also occur on Cerro Potosí. Another collection, *G. C. Rzedowski 22915*, from Hidalgo, exhibits a similar intermediate combination of characters. Also, a number of collections from northern Mexico clearly referable to the subspecies *mexicana* have remarkably short fruit, indicating a certain simi-

ilarity to the subspecies *bipatriata*. Furthermore, many individuals of otherwise typical *O. mexicana* subsp. *bipatriata* observed on Cerro Potosí (*Lowry & M. J. Warnock 3182, 3188*) have fruit armed with a few bristles toward the base, while others have completely glabrous fruit.

The strong similarity and intermediacy observed between the two subspecies of *Osmorhiza mexicana* has not gone unnoticed in the past. Constance and Shan (1948: 121) stated that *O. mexicana* subsp. *bipatriata* "was not included in the 'North American Flora,' because only the Mexican specimens had come to the attention of Mathias and Constance at the time that account was published (1944), and these had been regarded as somewhat aberrant representatives of *O. mexicana* (subsp. *mexicana*)."

Although Constance and Shan (1948) described the taxon *Osmorhiza bipatriata* as a distinct species they, too, recognized its strong resemblance to the plants of *O. mexicana*, citing two of the intermediates mentioned above (*Mueller 2231* and *Schneider 1108*).

6a. *Osmorhiza mexicana* Griseb. subsp. *mexicana*

Plants slender to rather stout, 4–8(–10) dm high; stems hirsutulous throughout. Root system rather deep, the roots weakly to rather strongly anise-scented. Leaves 2–3-ternate, broadly ovate to deltoid, 5–12(–15) cm long, villous or pilose, especially below; leaflets ovate, (1.5–)2–4 cm long, 1.5–2.5 cm wide, acute to acuminate, coarsely serrate to incised and pinnately lobed at the base; petioles 6–14 cm long. Umbels rather loose; peduncles 2–3(–4), terminal and lateral, 7–15 cm long; involucre wanting, or sometimes composed of 1–2 linear, foliaceous, ciliate bracts, each (2–)5–10(–13) mm long, 0.3–1 mm wide; rays spreading-ascending, 2.2–10(–11) cm long; umbellets 2–5(–7) per umbel; involucre of (1–)2–4 linear, acuminate, ciliate bractlets, each (1.5–)2.5–9.5 mm long, 0.3–1 mm wide, spreading to reflexed; pedicels (2–)5–11 per umbellet, spreading-ascending, those of the hermaphrodite flowers (1–)2–8.5(–12) mm long, those of the staminate flowers 2–5(–6) mm long. Hermaphrodite flowers (2–)3–6 per umbellet, (8–)12–32(–41) per umbel, staminate flowers (0–)2–7 per umbellet, (0–)5–25(–35) per umbel; corolla white or greenish white, somewhat showy; styles (including stylopodium) 0.6–1 mm long, stylopodium 0.2–0.5(–0.6) mm long, low-conic to somewhat depressed, often with a conspicuous disc.

Fruit linear-oblong, tapering to a short beak at the apex, concave furrowed, (10–)11–20 mm long, the ribs moderately to densely hispid with retrorse bristles, especially toward the base, the caudate appendages (1.5–)2–8.5(–10) mm long. $n = 11$ (Bell & Constance, 1966; Constance et al., 1976). Figures 9f and 17.

Flowering period. Late May to July (Mexico); July (Central America); November and December (South America).

Habitat. Moist forests, at middle and higher elevations.

Common names. Mexican sweet cicely, Mexican sweetroot, Cilandrillo (Mexican Indian; Oaxaca).

Representative specimens. ARGENTINA. CATAMARCA: Dept. Andalgalá, Comné Esquina Grande, *Jørgensen* 1812 (F, GH, MO, US). CÓRDOBA: Dept. Calamuchita, Sierra Arhala de Córdoba, *Hieronymus*, 3 Dec. 1878 (UC); El Vallecito, Sierra Grande, 2,000 m, *A. T. Hunziker* 8651 (UC, WTU); La Cumbrecita, 1,500 m, *Roig* 17767 (UC). JUJUY: Dept. Santa Bárbara, Sierra El Centinela, 1,950 m, *Cabrera et al.* 17299 (UC, WIS). SALTA: Dept. Guachipas, Alemania, 1,300 m, *Venturi* 9810 (GH, K, MO). TUCUMÁN: Questa del Garabatal, Sierra del Tucumán, *Lorentz & Hieronymus* 863 (GOET); near la Cienega, *Lorentz & Hieronymus* 668 (GOET).

BOLIVIA. COCHABAMBA: N of Yungas, 3,200 m, *Buchtien* 659 (GH, NY, US). LA PAZ: Prov. Murillo, upper Valle de Zongo, 3,500 m, *Solomon* 5228 (MO), *Tate* 194 (NY); near Sorata, 3,100 m, *Mandon* 594 (BM, G, GH, GOET, K, NY, P, UC).

COLOMBIA. TOLIMA: Paramó de Ruiz, 3,500 m, *Lehmann* 3074 (BM, K, US), *Pennell* 3101 (NY, US).

COSTA RICA. ALAJUELA: San Juan, Cerrido, *Brenes*, 21 Jan. 1906 (NY). CARTAGO: Volcán de Irazú, 2,750 m, *Kuntze* 2270 (K, NY); 3,000 m, *Pittier & Durand* 4073 (US); Chicué, Irazú, 2,750 m, *J. León* 2680 (CR); Près du sommet de l'Irazú, 3,000 m, *Tonduz* 4273 (BR, CR); Sabana larga, S of Volcán de Irazú, 2,900 m, *Pittier* 44 (K), 200 (BR, CR, US). SAN JOSÉ: Cerro de las Vueltas, 2,800 m, *Standley & Valerio* 43592 (K, US), 43669 (US).

GUATEMALA. CHIMALTENANGO: Chichavac, 2,500 m, *Skutch* 499 (CAS, MICH); Volcán de Agua, 2,450 m, *Johnston* 809 (F). HUEHUETENANGO: near Hacienda de Chaucol, Sierra de los Cuchumatanes, 3,350 m, *Nelson* 3645 (GH, NY, US), *Skutch* 1210 (F, GH); near Tunima, Sierra de los Cuchumatanes, 3,450 m, *Steyermark* 48383 (F, UC); Cerro Chémal, Sierra de los Cuchumatanes, 3,750 m, *Steyermark* 50307 (F). QUEZALTENANGO: Volcán Santo Tomás, *Steyermark* 34714 (F, UC); 5 km N of Ostuncalco, Sierra Madre, 2,600 m, *Williams et al.* 25467 (GH, NY, US). SACATEPEQUEZ: Volcán de Agua, above Santa María, *Bell & Duke* 16987 (UC). SAN MARCOS: San Luis, 3.8 km W of Ixchiguan on road to Tacana, 3,450 m, *Beaman* 3247 (GH, UC, US). SOLOLÁ: Volcán Tolimán, side facing Volcán Atitlán, 2,900 m, *Steyermark* 47581 (F).

MEXICO. CHIAPAS: near San Cristobal, *Nelson* 3188

(US). CHIHUAHUA: La Rocha, Sierra Mohinora, 2,300 m, *Correll & Gentry* 23138 (GH, LL), *Nelson* 4864 (GH, US); 16 km SW of Guadalupe y Calvo, Cerro Mohinora, *Straw & Forman* 2013 (UC). DISTRITO FEDERAL: Miguel-Hidalgo Park, 3,200 m, *Bell & Duke* 16778 (GH, K, MICH, MO, NY, TEX, UC); Desierto de los Leones, *Kenoyer* 562 (MICH), 9 July 1938 (ARIZ), *MacDaniels* 83 (F), *Sharp & Gilly* 22 (MICH); Contreras, *Lyonnet* 1623 (US); Sierra de Ajusco, 2,450 m, *Pringle* 6615 (BM, BR, CAS, F, G, GH, GOET, K, MO, NY, P, PH, RM, UC, US); Cuarto Dinamo, 3,100 m, *A. Ventura A.* 3586 (G); Los Dinamos, 3,000 m, *A. Ventura A.* 3492 (G). HIDALGO: 3.8 km SW of Cabrera on Hwy. 105, 2,800 m, *Bell & Duke* 16813 (MICH, TEX, UC); El Chico Nat'l. Park, 16 km NE of Pachuca, 2,750 m, *Weller* 582 (UC); Real del Monte, 2,800 m, *Hernandez X.* 464 (LL), *Sharp* 44593 (UC); Cerro de las Ventanas, 6 km N of Pachuca, 2,900 m, *Rzedowski* 26813 (CAS, MICH). JALISCO: Nevado de Colima, 2,700 m, *McVaugh* 10047 (MICH, UC), 12853 (K, MICH, SMU, UC, US), *Gregory & Eiten* 295 (GH, MICH, MO). MÉXICO: Valleé de Mexico, *Bourgeau* 781 (BR, GH, K, UC, US); Méson Viejo, Temascaltepec, *Hinton* 1308 (F, GH, K, US); Las Cruces, Temascaltepec, 3,260 m, *Hinton* 4899 (BM, K, US), *Pringle* 5208 (MO, US); Crucero-Agua Blanca, Temascaltepec, *Hinton et al.* 8324 (ARIZ, BM, G, GH, K, LL, MO, NY, PH, TEX, UC, WTU); Vaqueria del Jacal, 3,000 m, *Liebermann* 12240 (F, MO, US); Ixtapaluca, above Río Frio, 3,000 m, *Sharp & Gilly* 69 (MICH). MICH-OACAN: *Arsène s.n.* (CAS). MORELOS: 9.25 km W of Tres Cumbres, 2,800 m, *Steingraeber & Steingraeber* 150 (UC, WIS). NUEVO LEÓN: Cerro El Infernillo, S of Galeana, 2,750 m, *Bell & Duke* 16563 (UC); Hwy. 51, 40 km S of junction with Hwy. 54, 2,400 m, *Lowry & Warnock* 3181a (MO); Ojo de Agua, foot of Cerro Potosí, 2,000 m, *Sharp* 45711 (NY, UC). OAXACA: Cerro de Cabeza de Vaca, trail to Cerro San Felipe, 2,750 m, *Bell & Duke* 16884 (UC); Sierra San Felipe, 3,000 m, *Pringle* 5547 (F, US); Cordillera de Oaxaca, 2,750 m, *Galeotti* 2751 (G, K); 21 km N of Ixtlán de Juárez, 2,800 m, *Weller* 566 (UC). PUEBLA: Orizaba, 3,000 m, *Liebmann* 12241 (F, GH, US), *Müller* 1722 (GH, NY), *Rose & Hay* 5726 (US), *Standley* 12241 (UC); 3,650 m, *Seaton* 195 (F, GH, US); Popocatepetl, *Rose & Hay* 6249 (US); Esperanza, *Purpus* 7456 (MO, NY, UC, US). SAN LUIS POTOSÍ: Sierra de Alvarez, Sierra Madre Oriental, 2,300 m, *Palmer* 127 (US), *Pennell* 17883 (PH, US). VERACRUZ: near El Puerto, 2,350 m, *Sharp* 44680 (GH, NY).

PERU. CUZCO: Ollantiatambo, 3,000 m, *Cook & Gilbert* 747 (US); Veronica, between Cuzco and Machupichu, 3,500 m, *Rauh & Hirsch* P1026 (UC); between Cachupampa and Chile-chile, 2,500 m, *Vargas* 9697 (GH, K, MO, UC), 3,000 m, *Vargas* 1352 (F).

Weddell (1861) and Hemsley (1880) considered Central and South American populations as *Osmorhiza brevistylis* DC.

6b. *Osmorhiza mexicana* Griseb. subsp. *bipatriata* (Constance & Shan) Lowry & Jones, comb. et stat. nov. *Osmorhiza bipatriata* Constance & Shan, Univ. Calif. Publ. Bot.

23: 121. 1948. TYPE: U.S.A. Texas: Jeff Davis Co., in wet ground at spring, W branch Madera Canyon on slope of Mt. Livermore, Davis Mts., *L. C. Hinckley*, 26 July 1937 (holotype, NY!; isotypes, or possibly paratypes, because label data are not identical to those of the holotype, ARIZ!, GH!).

Plants slender, 2–7 dm high; stems sparingly hirsutulous to glabrescent. Root system somewhat shallow, with a weak anise-like scent. Leaves 2–3-ternate, ovate to broadly ovate, 4–10(–14) cm long, villous or pilose, especially on the veins below; leaflets ovate to ovate-oblong, (0.7–)1.2–4 cm long, 0.5–3 cm wide, acute to acuminate, coarsely serrate-laciniate to lobed or divided at the base; petioles 4–10(–12) cm long. Umbels loose and rather open; peduncles 1–3, terminal and often lateral, 3–15(–18) cm long; involucre wanting, or often composed of 1(–2) linear, foliaceous, ciliate bracts, each 4–10 mm long, 0.5–0.8 mm wide; rays spreading-ascending, (1.4–)1.6–6.5(–7.5) cm long; umbellets (2–)3–9 per umbel, (0–)1–4(–6) of them producing only staminate flowers; involucre of 1–4 linear, acuminate, ciliate bractlets, each (2–)3–4.5 mm long, 0.3–1 mm wide, spreading; pedicels (3–)4–20(–22) per hermaphrodite umbellet, (3–)4–17 per staminate umbellet, spreading, those of the hermaphrodite flowers (3–)4–7.5(–8) mm long, those of the staminate flowers (1.5–)2–3.5(–4) mm long. Hermaphrodite flowers 1–3 per umbellet, (2–)5–10 per umbel, staminate flowers (4–)6–21 per hermaphrodite umbellet, (33–)40–70(–125) per umbel; corolla white, or sometimes tinged with purple, pink, or green, rather inconspicuous; styles (including stylopodium) 0.5–0.75 mm long, stylopodium 0.25–0.3 mm long, low-conic, often with a conspicuous disc. Fruit linear-fusiform, tapering into a short beak at the apex, concave furrowed, 9–11(–12) mm long, the ribs glabrous, or with a few retrorse bristles at the base, the caudate appendages lacking, or sometimes to 1.8 mm long. Figures 9g and 17.

Flowering period. June and July.

Habitat. Moist, generally shaded ravines and canyons, from 2,100 to over 2,750 m.

Common names. None.

REPRESENTATIVE SPECIMENS. U.S.A. TEXAS: Jeff Davis Co., Upper Madera Canyon, NW side of Mt. Livermore, 2,300 m, *Hinckley* 408 (F, NY), 3589 (NY, UC), *B. H. Warnock & Hinckley* 4147 (NY), *B. H. Warnock* 7479 (SMU, TEX).

MEXICO. COAHUILA: NW of Campo Cinco, Madera

del Carmen, 2,400 m, *Lowry & M. J. Warnock* 3130, 3149 (ILL, MEXU, MO, NY, TEX, UC), *Fryxell* 2689 (UC), 2703 (CAS); Upper Dos Canyon, Madera del Carmen, 2,450 m, *Fryxell* 2722 (MO, UC). NUEVO LEÓN: 2.2 km below microwave tower, Cerro Potosí, 3,000 m, *Lowry & M. J. Warnock* 3188 (ILL, MEXU, MO, NY, TEX, UC); microwave tower, Cerro Potosí, 3,200 m, *McGregor et al.* 252 (UC), 339 (NY, SMU, UC); Cerro Potosí, *Schneider* 1043 (F).

The following collections exhibit combinations of characters intermediate between *O. mexicana* subsp. *mexicana* and subsp. *bipatriata*:

MEXICO. HIDALGO: Penas Largas, near Tezoantla, 2,750 m, *Rzedowski* 22915 (CAS, MICH, ORE, TEX). NUEVO LEÓN: Canyon below Las Canoas, Cerro Potosí, *Mueller* 2231 (GH); Hacienda La Jolla, 2,600 m, *Schneider* 1108 (F).

7. *Osmorhiza brachypoda* Torrey in Durand, J. Acad. Nat. Sci. Philadelphia II, 3: 89. 1855. *Myrrhis brachypoda* (Torrey) E. Greene, Fl. francisc. 332. 1892. *Washingtonia brachypoda* (Torrey) A. A. Heller, Cat. N. Amer. pl. 5. 1898. *Scandix brachypoda* (Torrey) Koso-Polj., Bull. Soc. Imp. Naturalistes Moscou 29: 143. 1916. TYPE: U.S.A. California: Nevada Co., near the banks of Deer Creek, Nevada City, *H. Pratten*, July 1851. (This specimen has not been located.)

Osmorhiza brachypoda var. *fraterna* Jepson, Fl. Calif. 2: 670. 1936. TYPE: U.S.A. California: San Bernardino Co., Arroyo Seco, San Gabriel Mts., 600 m, *F. W. Peirson* 451a (holotype, JEPS!).

Plants rather stout, 3–8 dm high; stems 1–2(–3), erect, villous to hirsutulous. Root system rather deep, the roots weakly to moderately anise-scented. Leaves 2–3-ternate, ovate to deltoid, 10–18(–20) cm long, hirsutulous, especially below and along the veins; leaflets ovate, 2–6(–8) cm long, 1–4 cm wide, acute or obtuse, coarsely serrate, incised and pinnately lobed at the base; petioles 5–18(–22) cm long. Umbels somewhat congested, peduncles 2–4, terminal and lateral, 9–18 cm long; involucre wanting, or often composed of 1–3 small (or very rarely large and leaf-like), linear-lanceolate, foliaceous, ciliate bracts; rays spreading-ascending, (3.3–)3.8–12(–12.5) cm long; umbellets 2–5 per umbel; involucre of 2–6(–7) linear to lanceolate, acute, ciliate bractlets, each (1–)2–9.5(–15) mm long, (0.3–)0.5–1.3(–5) mm wide, spreading or reflexed; pedicels (2–)4–14(–16) per umbellet, ascending, those of the hermaphrodite flowers 1–4.5(–7.5) mm long, those of the staminate flowers (2–)2.8–6(–6.5) mm long.

Hermaphrodite flowers (1–)2–10 per umbellet, (6–)10–35(–43) per umbel, staminate flowers (0–)1–6(–8) per umbellet, (0–)5–30(–40) per umbel; corolla greenish white, somewhat inconspicuous; styles (including stylopodium) (0.6–)0.7–1.2(–1.3) mm long, stylopodium 0.2–0.5(–0.6) mm long, low-conic to somewhat depressed, often with a conspicuous disc; carpophore cleft about one-third of its length. Fruit oblong-fusiform, tapering to a narrow beak at the apex, deeply concave furrowed, (12–)13–18(–20) mm long, the ribs densely hispid with retrorse bristles, especially toward the base, the caudate appendages (0.5–)0.9–3.8(–4) mm long. $n = 11$ (Constance et al., 1976). Figures 9h and 16.

Flowering period. March to early May.

Habitat. Seasonally moist forests at lower to fairly high elevations in the mountains.

Common names. California cicely, California sweet cicely, Orris root, Sweet cicely.

Representative specimens. U.S.A. ARIZONA: Gila Co., Collom Camp, base of Mazatzal Mts., 1,200 m, *Collom 866* (ARIZ, UC); Hells Canyon, Mazatzal Mts., *Harrison et al. 7815, 7830* (ARIZ). CALIFORNIA: Amador Co., 2.5 km E of Oleta, *Raven 9123* (CAS, JEPS); Contra Costa Co., Mt. Diablo, 300 m, *Bowerman 2108* (CAS, UC); Kern Co., mouth of Kern Canyon, Greenhorn Mts., *Benson 3342* (CAS, NY, RM); Los Angeles Co., Evey Canyon, N of Claremont, *Lowry 1481, 1482* (ILL); Monterey Co., Arroyo Seco Camp, 29 km NW of Greenfield, Santa Lucia Mts., 760 m, *Constance 3311* (ARIZ, CAS, G, K, MO, NY, RM, UC); Orange Co., Silverado Canyon, Santa Ana Mts., 380 m, *Munz & Harwood 3722* (NY, RM); Riverside Co., Temescal, *Hall 886* (UC); San Benito Co., Idria, *Ferris 7028* (CAS, NY); San Bernardino Co., near San Bernardino, 350 m, *Parish 4165* (BM, K, NY, UC); San Diego Co., Cuyamaca, *Abrams 3838* (BM, CAS, G, K, MO, NY, UC, Z); Santa Barbara Co., Santa Barbara, *Elmer 3879* (CAS, K, MO, NY, WIS, Z); Santa Clara Co., Alum Rock Park, *Heller 8481* (G, MO, NY, WIS); Sierra Co., Downieville, *Bigelow s.n.* (NY); Tuolumne Co., Hetch-Hetchy Trail, Yosemite Nat'l. Park, 1,500 m, *Hall & Babcock 3379* (ARIZ, NY, RM, UC); Ventura Co., Sulphur Mt. Spring, Sulphur Mts., *Abrams & McGregor 24* (CAS, G, NY, Z).

IIc. *Osmorhiza* Raf. sect. *Nudae* Constance & Shan ex Lowry & Jones, sect. nov. TYPE: *O. chilensis* Hook. & Arn.

Uraspermum § *Osmorhiza* (Raf.) Kuntze, *Lexicon* 582. 1904, pro parte.

Scandix 2. [sect.] *Urascandix* Koso-Polj., *Bull. Soc. Imp. Naturalistes Moscou* 29: 143. 1916, pro parte.

Osmorhiza sect. *Nudae* Constance & Shan, *Univ. Calif. Publ. Bot.* 23: 113. 1948, nom. nud.

Involucrum deficiens vel raro ab 1(–2) bracteis minutis constitutum; involucellum deficiens vel raro ab

1(–2) bracteolis minutis constitutum. Styli cum stylopodio 0.2–1.1 mm longi, stylopodium 0.1–0.6 mm longum, conicum vel leviter depressum, saepe cum disco conspicuo.

Involucre wanting, or rarely composed of 1(–2) minute bracts; involucler wanting, or rarely composed of 1(–2) minute bractlets. Styles (including stylopodium) 0.2–1.1 mm long, stylopodium 0.1–0.6 mm long, conic to somewhat depressed, often with a conspicuous disc.

8. *Osmorhiza chilensis* Hook. & Arn., Bot. Beechey Voy. 26. 1830. TYPE: Chile. Concepción: Concepción, *Lay & Collie*, 9–20 Oct. 1825. (This specimen has not been located.)

Osmorhiza berterii DC., *Prodr.* 4: 232. 1830. *Myrrhis berterii* (DC.) Dietr., *Syn. pl.* 2: 984. 1840. *Uraspermum berteroi* [sic] (DC.) Kuntze, *Revis. gen. pl.* 1: 270. 1891. TYPE: Chile. “in sylvis umbrosis prope Tagua-Tagua,” *C. Bertero 446* (holotype, G-DC!; isotypes, G!, GH!, P!).

Schudia chilensis Molina ex Clos in *C. Gay, Fl. chil.* 3: 143. 1848, pro syn.

Scandix clavata Banks & Sol. ex Hook. f., *Fl. antarct.* 2: 288. 1846, pro syn.

Osmorhiza nuda Torrey, *Pacific railroad rep.* 4¹: 93. 1857. *Uraspermum nudum* (Torrey) Kuntze, *Revis. gen. pl.* 1: 270. 1891. *Myrrhis nuda* (Torrey) A. A. Heller, *Cat. N. Amer. pl.* 5. 1898. *Osmorhiza divaricata* var. *nuda* (Torrey) M. E. Jones, *Bull. Montana State Univ., Biol. Ser.* 15: 42. 1910. *Scandix nuda* (Torrey) Koso-Polj., *Bull. Soc. Imp. Naturalistes Moscou* 29: 143. 1916. TYPE: U.S.A. California: Napa Co., Shady woods, *J. M. Bigelow*, 27 Apr. 1853 or 1854 [lectotype, NY! (designated by Coulter & Rose, *Contr. U.S. Natl. Herb.* 7: 66. 1900); isolectotypes, GH!, K!].

Osmorhiza berterii var. *glacilior* Philippi, *Anales Univ. Chile* 85: 726. 1894. TYPE: Chile. “In Andibus provinciae l. d.” Chihuahua, *O. Philippi*, Jan. 1887 (holotype, SGO; photograph of holotype, ILL!).

Osmorhiza divaricata Nutt. ex Britt. in *Britt. & Brown, Ill. fl.* 2: 531. 1897, pro syn. *Washingtonia divaricata* (Nutt. ex Britt.) Britt. in *Britt. & Brown, Ill. fl.* 2: 531. 1897. *Scandix divaricata* (Nutt. ex Britt.) Koso-Polj., *Bull. Soc. Imp. Naturalistes Moscou* 29: 143. 1916. *Osmorhiza nuda* var. *divaricata* (Nutt. ex Britt.) Jepson, *Madroño* 1: 119. 1923. *Uraspermum divaricata* [sic] (Nutt. ex Britt.) Farwell, *Amer. Midl. Naturalist* 12: 70. 1930. TYPE: U.S.A. “Oregon woods,” *Nuttall s.n.* [lectotype, NY! (designated by Coulter & Rose, *Contr. U.S. Natl. Herb.* 7: 65. 1900); plus two authentic specimens (likely syntypes): “Columbia plains,” *Nuttall s.n.* (BM!); “Columbia woods,” *Nuttall s.n.* (GH!)].

Washingtonia brevipes Coult. & Rose, *Contr. U.S. Natl. Herb.* 7: 66. 1900. *Osmorhiza brevipes* (Coult. & Rose) Suksd., *Allg. Bot. Z. Syst.* 12: 5. 1906. *Os-*

morhiza nuda var. *brevipes* (Coult. & Rose) Jepson, Madroño 1: 119. 1923. *Uraspermum* [sic] *brevipes* (Coult. & Rose) Farwell, Amer. Midl. Naturalist 12: 70. 1930. TYPE: U.S.A. California: Siskiyou Co., Mt. Shasta and vicinity, E. Palmer 2481 (holotype, US!).

Washingtonia intermedia Rydb., Mem. New York Bot. Gard. 1: 289. 1900. *Osmorhiza intermedia* (Rydb.) A. A. Heller, Mont. Coll. Agric. Sci. Stud., Bot. 1: 93. 1905. TYPE: U.S.A. Montana: Gallatin Co., Bridger Mts., 2,150 m, P. A. Rydberg & E. A. Bessey 4595 [lectotype, NY! (designated by Constance & Shan, Univ. Calif. Publ. Bot. 23: 139. 1948); isolectotypes, K!, MONT!, US!].

Uraspermum barbatum Farwell, Amer. Midl. Naturalist 12: 70. 1930. TYPE: U.S.A. Michigan: Keweenaw Co., rocky woods, Copper Harbor, O. A. Farwell 8490 [lectotype, BLH! (designated by McVaugh et al., Bull. Cranbrook Inst. Sci. 34: 79. 1953); isolectotype, GH! ("isotype" fide Constance & Shan, Univ. Calif. Publ. Bot. 23: 139. 1948)].

Plants slender to rather stout, 3–12(–15) dm high; stems 1–3(–4), erect, villous to hirsutulous, or sometimes essentially glabrous. Root system rather deep, well developed, the roots with a weak carrot- or anise-like scent. Leaves biternate, orbicular to broadly ovate, 4–18(–23) cm long, appressed-hispidulous to villosulous (especially along the veins), or sometimes nearly glabrous; leaflets ovate-lanceolate to nearly orbicular, (2–)3–8(–10) cm long, 1–5 cm wide, obtuse or acute, coarsely serrate to incised, parted or divided at the base; petioles 5–16 cm long. Umbels rather loose; peduncles 2–4, terminal and lateral, 5–25 cm long; involucre wanting, or rarely composed of 1(–2) minute (to very rarely large and leaf-like), linear-lanceolate, foliaceous, ciliate, spreading bracts; rays spreading-ascending, (1.5–)2–12(–13) cm long; umbellets 3–8 per umbel; involucre wanting, or rarely of 1(–2) minute, linear-lanceolate, foliaceous, ciliate bractlets; pedicels (2–)3–9 per umbellet, spreading-ascending, those of the hermaphrodite flowers 4–20(–25) mm long, those of the staminate flowers (1–)2–7(–9) mm long. Hermaphrodite flowers (1–)2–6 per umbellet, (5–)9–30(–39) per umbel, staminate flowers 0–4 per umbellet, 0–18(–25) per umbel; corolla greenish white (rarely pink?), rather inconspicuous; styles (including stylopodium) 0.4–1.1(–1.2) mm long, stylopodium (0.2–)0.3–0.6 mm long, high-conic to somewhat gibbous, lacking a disc; carpophore cleft about one-fourth of its length. Fruit linear-oblong, tapering to a slender beak at the apex, concave furrowed, (11–)12–21(–25) mm long, the ribs moderately to densely hispid with retrorse bristles, especially

toward the base, the caudate appendages (2–)2.5–8.5(–10) mm long. $n = 11$ (Bell & Constance, 1957; Constance et al., 1976). Figures 9j and 15.

Flowering period. April to early July (North America), November and December (South America).

Habitat. Moist forested areas to somewhat drier, open, often disturbed areas.

Common names. Common sweet cicely, Mountain sweet cicely, Mountain sweetroot, Spreading sweetroot, Sweet cicely, Sweetroot, Western sweet cicely, Western sweetroot, Wood cicely, Asta de Cabra.

Representative specimens. U.S.A. ALASKA: Hyder, McCabe 8426 (UC); Salmon River, W of Hyder, Robuck 1348 (RM); Deer Mt., 2 km E of Ketchikan, McCabe 8644 (UC); Sitka, Eastwood 966 (CAS); Todd, SE Chichagof Island, Scamman 328 (DAO, photograph); Haines Hwy., mile 31, Walsh & Moore 6113 (NY); Yakutat, Piper 4280, 4283 (UC); Shaw Island, N of Cape Douglas, Anderson 1382 (US); Unalaska Island, Eyerdam 2256 (cited in Hultén, 1947). ARIZONA: Coconino Co., Oak Creek Canyon, 1,375 m, Demaree 41249 (NY); Gila Co., under Mogollon Rim, 1,800 m, Collom 753 (MO, US); Graham Co., Mt. Graham, Peebles et al. 4483 (US). CALIFORNIA: Butte Co., Jonesville, 1,550 m, Copeland 391 (ARIZ, BM, BR, CAS, K, MO, NY, RM, UC, WIS); Fresno Co., Huntington Lake, 2,135 m, Grant 1173 (ARIZ, JEPS); Humboldt Co., Lady Bird Johnson Grove, Redwood Nat'l. Park, Lowry 791, 795 (ILL); Lake Co., Summit Lake, near Mt. Sanhedrin, Heller 5879 (CAS, G, ILL, NY, RM, UC); Lassen Co., Diamond Mt., near Susanville, 2,000 m, M. E. Jones, 28 June 1897 (BM, CAS, MO); Los Angeles Co., Lily Spring, Mt. Hawkins, San Gabriel Mts., Thorne 41257 (RSA, UC); Mendocino Co., Noyo, Constance 2518 (BM, CAS, NY, RM, UC); Modoc Co., Deep Creek, Warner Mts., 1,825 m, Constance 3103 (K, MO, NY, UC); Monterey Co., Pfeiffer St. Park, Mathias 1300 (RM, SMU, UC); Placer Co., Emigrant Gap, M. E. Jones 2734 (BM, BR, CA, MO, NY, UC); San Diego Co., Stonewall Mine, Cuyamaca Mts., 1,400 m, Parish 4421 (BM, CAS, JEPS, K, MO, NY); Santa Clara Co., W of Los Gatos, Heller 7430 (CAS, G, MO, NY, UC, WIS); Siskiyou Co., NE base of Mt. Eddy, Heller 12467 (CAS, G, ILL, MO, NY, UC, WIS); Sonoma Co., Petrified Forest, Heller 5737 (CAS, G, MO, NY, RM, UC); Tehama Co., 1 km S of Lassen Chalet, Lowry 851 (ILL); Tuolumne Co., Sullivan Creek, 1 km W of Twain Harte P.O., 1,220 m, Alexander & Kellogg 3673 (JEPS, NY, UC). COLORADO: Archuleta Co., Piedra River, 11 km N of Arboles, 2,250 m, H. D. Harrington 4108 (CS); Boulder Co., Boulder, 1,650 m, Hanson C262 (ARIZ, MO); Garfield Co., 1.5 km N of Douglas Pass, M. J. Warnock 1699 (ILL); Mesa Co., SW slope of Hightower Mt., Kufeld K-73 (CS); Moffat Co., NE of Five Springs (Douglas Mountain) overlooking Conway Draw, 2,300 m, J. Walker & S. Tabar 349 (CS); Rio Blanco Co., Cow Creek, 4.5 km from confluence with Piceance Creek Road, 2,195 m,

- W. R. Erickson & D. Bartman 688 (CS). IDAHO: Bannock Co., Mt. Putnam, *Davis* 4672 (NY); Benewah Co., 16 km S of St. Maries, *Christ* 10927 (NY, UC); Bonneville Co., 17.5 km SW of Victor, *Lowry* 1116 (ILL, MO, NY, RM, UC); Clearwater Co., E Fork Potlatch River, *Cronquist & Jones* 5977 (CAS, NY, SMU, TEX, UC); Custer Co., Morse Canyon, Lemhi Mts., *Hitchcock & Muhlick* 11275 (NY, UC); Idaho Co., Clearwater River, 6.5 km below Lowell, *Constance & Rollins* 1608 (MO, UC, WTU); Lemhi Co., Jessie Creek, 1,375 m, *Payson & Payson* 1884 (CAS, MO, NY, RM); Lewis Co., Mission Creek, *St. John et al.* 6345 (NY, WTU); Nez Perce Co., Potlatch River, *Heller & Heller* 7006 (ILL, WIS). MAINE: Aroostook Co., Ft. Kent, *MacKenzie* 3509 (MO, NY); Somerset Co., E of Knight's Pond, Squarctown, *Fassett* 3658 (WIS). MICHIGAN: Alger Co., Miner's Falls, near Munising, *Fernald & Pease* 3450 (MICH, NY); Alpena Co., Thunder Bay Island, *Dodge*, 23 June 1907 (MICH); Baraga Co., Big Limestone Mt., L'Anse, *Fassett* 21055 (WIS); Gogebic Co., 17 km NW of Ironwood, *Voss* 6238 (MICH); Keweenaw Co., Copper Harbor, *Fernald & Pease* 3452 (K, MICH, WIS); Wayne Co., Livonia, *Farwell* 9442 (BLH). MONTANA: Beaverhead Co., Red Rock Lakes, 2,000 m, *Lowry* 2580 (ILL, MONT, UC); Carter Co., 1.5 km S of Alzada, *Booth* 2661 (MONT); Flathead Co., Columbia Falls, *Williams*, 22 June 1894 (MO, NY, RM); Glacier Co., Florence Falls, Glacier Nat'l. Park, 1,615 m, *Stickney* 1764 (MONT); Lake Co., Flathead Lake, *Hitchcock* 17734 (RM, UC, WTU); Madison Co., Bear Trap Canyon, 1,415 m, *Lowry* 2560 (ILL, MO, MONT, MONTU, NY, RM, UC); Missoula Co., Grant Creek, *Kirkwood* 1230 (MONT, NY); Silver Bow Co., Humbug Spires, 2,050 m, *Lowry* 1999 (ILL). NEVADA: Elko Co., Lamoille Canyon, Ruby Mts., 2,285 m, *Holmgren* 1354 (UC); Washoe Co., Little Valley, 2,000 m, *Baker* 1363 (G, MO, NY, RM, UC). NEW HAMPSHIRE: Coos Co., Carter Notch, *Forbes s.n.* (GH); Alpine Cascade, Gorham, *Pease* 16432 (GH). NEW MEXICO: Taos Co., 14.5 km S of Talpa, *Correll & Correll* 33135 (TEX). OREGON: Clatsop Co., Lee Wooden Co. Park, *Lowry* 1310 (ILL, MO, OSC, UC); Columbia Co., 16 km NW of Scapoose, *Lowry* 593 (ILL); Douglas Co., Iverson Co. Park, *Lowry* 1319 (ILL, MO, NY, OSC, UC); Grant Co., 7 km S of Long Creek, 1,460 m, *Lowry* 1095 (ILL, MO, NY, OSC, UC); Harney Co., *Train*, 6 June 1935 (OSC); Hood River Co., Herman Creek Trail, *Lowry* 706 (ILL, MO, OSC, UC); Jefferson Co., Jack Creek, 16 km W of Camp Sherman, *Lowry* 747 (ILL, OSC); Marion Co., Silver Creek, *Hall* 216 (GOET, MO, NY); Multnomah Co., Forest Park, Portland, *Lowry* 626 (ILL, MO, OSC, UC); Tillamook Co., Bay City, *Morrill* 162 (WTU); Wallowa Co., 0.5 km S of Wallowa Lake, 1,340 m, *Constance & Jacobs* 1304 (MO, UC); Wheeler Co., 3 km NE of Ochoco Divide, *Lowry* 1092 (ILL, OSC, UC). SOUTH DAKOTA: Lawrence Co., Deadwood, *Carr* 130 (MO, NY, RM, WIS); Lead City, 1,800 m, *Rydberg* 724 (NY). UTAH: Cache Co., Logan Canyon, 1,550 m, *Maguire* 3637 (RM, UC); Salt Lake Co., City Creek Canyon, *M. E. Jones* 1852 (BM, BR, G, NY, RM); Washington Co., 5 km E of Pine Valley, 2,135 m, *Gould* 1846 (ARIZ, CAS, NY, UC). WASHINGTON: Chelan Co., Stehekin, 400 m, *Rose* 48178 (SMU, UC); Columbia Co., Godman Spring, Blue Mts., 1,750 m, *Constance et al.* 1162 (CAS, MO, UC, WIS); Grays Harbor Co., Montesano, *Heller & Heller* 3975 (G, MO, NY, UC); Jefferson Co., N Fork Quinalt River, *Lowry* 254a (ILL); King Co., Seattle, *Thompson* 10395 (CAS, MO, NY, WTU); Kittitas Co., 5 km W of Teanaway Junction, Wenatchee Mts., *Hitchcock* 17258 (RM, UC, WTU); Klickitat Co., Bingen, *Suksdorf* 10037 (BM, CAS, ILL, K, MO, NY, UC); San Juan Co., Friday Harbor, *Zeller & Zeller* 768 (CAS, K, MO, NY); Skamania Co., 50 km E of Camas, *Lowry* 713 (ILL); Spokane Co., Clarks Spring, N of Spokane, *Kreager* 44 (NY, UC, WTU); Whatcom Co., Fairhaven Coll., Bellingham, *Lowry* 629 (ILL, MO, NY, UC). WISCONSIN: Bayfield Co., 4 km S of Little Sand Bay, *Iltis & Kawano* 20503 (BM, UC). WYOMING: Campbell Co., 4 km SW of Rockypoint, *Dueholm* 2962 (RM); Carbon Co., S Brush Creek Campground, *Lowry* 1131 (ILL, MO, NY, RM, UC); Fremont Co., Sinks Canyon, Popo Agie River, W of Lander, 2,200 m, *Scott* 909 (NY, RM); Lincoln Co., Cottonwood Lake, SE of Afton, *Porter* 3776 (CAS, RM, TEX); Teton Co., 5 km W of Teton Pass, *Lowry* 1120 (ILL, RM, UC); Uinta Co., Teton Mts., *Nelson & Nelson* 6473 (G, ILL, MO, NY, RM).
- CANADA. ALBERTA: Cameron Lake, Waterton Lakes Nat'l. Park, 1,660 m, *Breitung* 16175 (DAO, UC); Chief Mt., Waterton Lakes, 1,525 m, *Breitung* 16842 (SMU, UC); Nordegg River, *Brinkman* 4072 (DAO); Olds, *Fletcher* 739 (DAO); 13 km W of Pincher Creek, *Moss* 86 (DAO). BRITISH COLUMBIA: 13 km N of Flathead Customs Sta., 1,370 m, *Taylor & Ferguson* 2044 (DAO); 5 km NNE of Nelson, *Calder & Savile* 9405 (DAO, UC); Carbonate, Selkirk Mts., 820 m, *Shaw* 219 (MO, NY, RM); Glacier Nat'l. Park, *Lowry* 1012 (ILL, MO, NY, UC); 13 km W of Revelstoke, *Lowry* 1016 (ILL); Friday Creek, S of Princeton, *Lowry* 1020 (ILL, UC); Manning Prov. Park, *Lowry* 1026a (ILL); 11 km N of Boston Bar, *Taylor & Staudt* 4197 (DAO, UC); 11 km SE of Vedder's Crossing, *Lowry* 1036 (ILL, MO, NY, UC); 3 km S of Pemberton, *Lowry* 644 (ILL); NW tip Saturna Island, Gulf Islands, *Calder & MacKay* 28775 (DAO); Cowitcham Lake, Vancouver Island, *Rosendahl & Johnson* 1857 (DAO); Ucluelet, *Macoun* 78607 (NY); 16 km NE of 150 Mile House, 1,000 m, *Calder et al.* 19030 (DAO); Bella Coola, *McCabe* 123, 1402 (UC); Reid Lake, NW of Prince George, *Eastham* 14780 (DAO); 5.5 km SW of Telkwa, *Calder et al.* 15259 (DAO); 4 km W of Skeena Crossing, 250 m, *Calder et al.* 13021 (DAO, UC); Skidegate Mission, Graham Island, *Calder* 21427 (DAO, UC); Haines Rd., km 73.5, NW of Haines, 200 m, *Taylor et al.* 1406 (DAO). NEW BRUNSWICK: Albert Co., Goshen, *Squires & Christie*, 26 June 1963 (DAO); Restigouche Co., 1 km SE of Summit Depot, *Cunningham*, 21 July 1960 (DAO). NEWFOUNDLAND: N of Doctor Hill, *Fernald & Long* 28771 (K); Frenchman's Cove, Bay of Islands, *Waghorne*, 24 June 1895 (DAO, photo ex QK). NOVA SCOTIA: Inverness Co., Brigend, *Smith et al.* 2589 (DAO, UC); Kings Co., Kentville, *Prince & Atwood* 1011 (CAS, WIS). ONTARIO: Algoma Dist., Garden River, *Fassett* 13282 (WIS); Bruce Co., Kincardine, *Anderson & Fassett* 21585 (UC, WIS); Manitoulin Dist., Excelsior, Manitoulin Island, *Fassett* 14833 (WIS); Thunder Bay Dist., Talus Lake, Sibley Twp., *Taylor et al.* 1146 (UC). QUÉBEC: Charlevoix Co., St. Chrétien, *Cinq-Mars et al.* C69-25 (DAO); Gaspé Co., Percé, *Collins & Fernald* 119 (ILL, K, LL, NY, UC); Matane Co., 8 km S of Mt. Joli, *Scoggan* 13595 (NY, UC); Rivière-du-Loup Co., Rivière-du-Loup, *Churchill*, 8 Aug. 1902 (SMU). SAS-

KATCHEWAN: Cypress Hills Park, *Breitung* 4720, 8102 (DAO, UC).

ARGENTINA. CHUBUT: Futaleufu, Lago Futalaufquen, *Burkart* 19823 (P, UC); Río Futaleufu, *Castellanos*, 24 Jan. 1945 (NY); Colonia 16 de Octubre, *Lahitte* 52212 (UC); Senguerr, Lago Fontana, *Moreno s.n.* (NY); Languiño, Pa pa Chica, *Soriano* 2488 (UC). MENDOZA: Tunuyan, Cerro de las Piedras, *Ruiz Leal* 3106 (BM, UC). NEUQUÉN: Aluminé, Lago Quillén, *Valle et al.* 3074 (K); Huiliches, Lago Huechulafquen, Parque Nac. Lanín, *Correa* 5537 (UC); Laguna Verde, Parque Nac. Lanín, 980 m, *Correa et al.* 5774 (UC); Lacár, Hua Hum, *Cabrera* 11229 (UC), *DeBarba* 1754 (BR, MO, TEX), *O'Donnell* 2331 (NY); San Martín de los Andes, *Hunziker* 6910 (UC), *Ruiz Leal & Roig* 18128 (UC), *Ruiz Leal* 20265 (UC), *O'Donnell* 2399 (NY); Los Lagos, Parque Nac. Nahuel Huapí, *DeBarba* 1548 (TEX), 2110 (K); Minas, Lagunas Epu-Lauquén, Aduana Vieja, 1,300 m, *Boelcke et al.* 10826 (UC). RIO NEGRO: Bariloche, Parque Nac. Nahuel Huapí, *Boelcke* 5246, 5659, 5842 (UC), *Boelcke & Hunziker* 3417 (UC), *Cabrera & Job* 98 (NY), *DeBarba* 95 (NY, UC), 1127 (P), *Descole* 25 (NY), *Meyer* 7456 (NY); Entre Liao-Liao y Bahía López, *Meyer* 8002 (NY). SANTA CRUZ: Lago Argentino, Parque Nac. "Los Glaciares," *Correa et al.* 2998 (UC); Cerro Mayo, *James* 3002 (BM, UC); Lago Argentino, *Ruiz Leal* 26558 (UC); Lago Buenos Aires, Río Jeinemeni, entre Lago Buenos Aires y Codo Río Mayer, 500 m, *von Platen & Greiner* 150 (MO, Z). TIERRA DEL FUEGO: Ushuaia, Estancia Harberton, *Constance et al.* 3860 (UC), *Goodall* 481 (MICH), 549 (MICH, NY), 3585 (UC); Lago Roca, *Goodall* 2439 (UC); Estancia Figue, *Ruiz Leal & Roig* 15117 (UC).

CHILE. ACONCAGUA: Los Ojos de Agua, *Bridger* 475 (K); Zapallar (El Tigre), *Johow*, Nov. 1908 (ILL), *Looser*, 13 Oct. 1953 (UC), *Möller*, 8 Dec. 1951 (UC). ARAUCO: Contulmo, Cerro Santa Elena, *Ricardi* 9248 (UC). AYSÉN: Coihaique, cercanías del Lago Seco, 750 m, *Schlegel* 2321 (F); Pto. Puyuhuapi, Río Pascue, 8 m, *Schwabe* 45 (NY). CAUTÍN: Temuco, *Elliot* 276 (BM, NY); Temuco, Truf-Truf, *Montero O.* 6371 (UC); Temuco, Fundo Huilquilco, 2 km S of Quepe, *Moore* 312 (K, UC); Vilcún, 330 m, *Hollermayer* 334 (UC); Pucón, La Península Entrada, 250 m, *Mahu* 11425 (ILL); Victoria, 16 km from Termas de Tolhuaca on rd. to Curacautin, 950 m, *Morrison & Wagenknecht* 17486 (G, K, UC); Volcán Llaima, 1,100 m, *Werdermann* 1246 (BM, CAS, F, G, K, MO, NY, UC, Z); Pass to Longuimay, 1,000 m, *Zöllner* 5617 (ILL). CHILOÉ: Ancud, Chiloé Island, *MacMillan & Erlanson* 11 (MICH); Cucao Laguna, Chiloé Island, *Philippi & Borchers*, 2 Feb. 1885 (BM). CONCEPCIÓN: Lota, *Brooke* 6964 (BM); Concepción, *D'Urville* 289 (P), *Holway* 139 (NY); probably Talcahuano, *Poeppig s.n.* (UC). LLANQUIHUE: 10 km W of Los Muermos, *Constance & Sparre* 3573 (UC); Valle Cayutue, *Rudolph* 4925 (UC). MAGALLANES: 50 km SW of Puerto Natales, rd. to Punta Arenas, 30 m, *Eyerdam et al.* 24181 (K, MO, UC); Sandy Point, *Lechler* 1186 (GOET); Estancia Vicuña, 200 m, *Moore* 2178 (K, UC); Seno Almirantazgo, ca. 8 km from mouth of Río Azopardo, 20 m, *Moore* 2307 (UC). MALLECO: Fundo Solano, Los Alpes, Cord. Nahuelbuta, 1,200 m, *Eyerdam* 10333 (F, UC); 1 km W of Agua Fria, W of Angol, Sierra Nahuelbuta, 650 m, *Hutchison* 293 (K, UC); Termas de Tolhuaca, 1,100 m, *Looser* 2746 (UC), *Solomon & Solomon* 4479 (MO),

Zöllner 10201 (MO); Parque Nac. de Nahuelbuta, *Mahu* 5767, 8272, 8417, 8733, 8734, 8735, 11451 (ILL), *Pincheira* 7881, 8218 (ILL). ÑUBLE: Termas (Baños) de Chillán, 1,750 m, *Moore* 414 (UC), *Philippi & Borchers s.n.* (BM). O'HIGGINS: La Leona, Rancaque, *Bertero* 446 (P), *Guillemin* 446 (P); Agua de la Vida, *Borchers s.n.* (GOET); Baños, Cauquenes, *Philippi*, Sept. 1896 (BM); upper valley of River Cachapoal, *Zöllner* 9892 (ILL). OSORNO: Termas de Puyehue, 350 m, *Sparre* 2128 (K). SANTIAGO: Manzano, *Looser*, 11 Nov. 1928 (UC); Quebrada de Macul, 1,100 m, *Pisano V. & Baraona L.* 1572 (UC); Los Valdes, near Río Maipo, 2,500 m, *Zöllner* 2079 (ILL). VALDIVIA: Valdivia, *Bridges* 768 (K), *DeCandolle m.* 418 (P), *Gay* 124 (P), *Hohenacher* 481 (K), *Hollermayer* 334 (BR); Cabo Blanco, Río Cruces, *Klempau s.n.* (UC); La Unión Reserva Forestal Llancacura, *Mahu* 11424 (ILL). VALPARAÍSO: Valparaíso, *Bertero* 74 (F, NY, UC—all photos ex B), *Cuming* 335 (K); Limache (Lliu-Lliu), *Garaventa H.* 2270 (UC), *Bertero*, Oct. 1828 (BM), *Bertero* 1163 (G, P), *Steudel* 1163 (P); Laguna Verde, *Garaventa H.* 2378 (UC); Granizo, Cajón Grande, foot of Cerro Campana, near Olmue, 15 km E of Limache, *Meyer* 9704 (MO, UC), *Zöllner*, Nov. 1977 (ILL); 8 km from La Dormida on trail to Las Vizcachas, 1,100 m, *Morrison* 16832 (G, K, MO, UC); Quilpué, *Zöllner* 8353 (MO).

Constance and Shan (1948) were the first to recognize the conspecificity of North and South American populations of *Osmorhiza chilensis*. Previous to that, the North American material had been referred to a variety of names, including *Osmorhiza brevipes*, *O. divaricata*, *O. intermedia*, *O. nuda*, and *Uraspermum barbatum*, while South American specimens were identified as either *O. berterii* or *O. chilensis*. However, the similarity of plants from the two areas was mentioned by Coulter and Rose (1895) and Reiche (1902).

It has been established that the names *Osmorhiza chilensis* and *O. berterii* were both published in 1830 (Constance & Shan, 1948; cf. also Jackson, 1893; Rickett, 1945). While the month of publication for *O. berterii* most likely is September, that for *O. chilensis* is not known. However, Hooker and Arnott (1833), J. D. Hooker (1846), and Gray (1854) accepted the name *O. chilensis*, placing *O. berterii* in synonymy under it, and it was on this basis that Constance and Shan (1948) adopted the former name. We have uncovered no additional evidence that would require a different interpretation.

Bongard (1833) included Alaskan populations of *Osmorhiza chilensis* in *O. brevistylis* DC. Similarly, W. J. Hooker (1840) used an expanded concept of *O. brevistylis* to include populations from western North America now included in *O. chilensis*.

9. *Osmorhiza purpurea* (Coulter & Rose) Suksd., *Allg. Bot. Z. Syst.* 12: 5. 1906. *Washingtonia purpurea* Coulter & Rose, *Contr. U.S. Natl. Herb.* 7: 67. 1900. *Osmorhiza chilensis* var. *purpurea* (Coulter & Rose) Boivin, *Naturaliste Canad.* 93: 644. 1966. TYPE: U.S.A. Alaska: Sitka, *F. V. Coville & T. H. Kearney, Jr.*, 14–17 June 1889 (holotype, US!).

Washingtonia leibergii Coulter & Rose, *Contr. U.S. Natl. Herb.* 7: 66. 1900. *Osmorhiza leibergii* (Coulter & Rose) Blankinship, *Mont. Coll. Agric. Sci. Stud., Bot.* 1: 93. 1905. TYPE: U.S.A. Washington: Chelan (formerly Okanogan) Co., Sandy slopes of Nason Creek, (branch of the Wenatchee River) 1,370–2,133 m, *J. H. Sandberg & J. B. Leiberg* 666 [holotype, US!; isotypes, BM!, BR!, CAS!, MO!, NY!, ORE!, P!, UC! (or possibly paratypes, as label data are not identical to those of the holotype)].

Plants slender, 2–6(–8) dm high; stems 1–2(–3), erect to ascending, sparingly hirsutulous to glabrescent. Root system relatively deep, well developed, the roots with a weak carrot- or anise-like scent. Leaves (1–)2–3-ternate, orbicular to deltoid or broadly ovate, 3–10(–12) cm long, sparingly hirsutulous along the veins and margins, to glabrous; leaflets lanceolate to ovate, 1.5–6(–8) cm long, 0.5–4 cm wide, acute or acuminate, coarsely serrate to incised, lobed or deeply divided at the base; petioles 5–10(–12) cm long. Umbels rather loose; peduncles 2–3(–4), terminal and lateral, 3–9(–11) cm long; involucre wanting, or very rarely consisting of 1 minute, linear-lanceolate, foliaceous, ciliate, spreading bract; rays spreading-ascending, (2.7–)3–9.5(–11) cm long; umbellets (2–)3–7 per umbel, sometimes 1–2 of them producing only staminate flowers; involucre wanting; pedicels (2–)3–9 per hermaphrodite umbellet, 3–5 per staminate umbellet (when present), spreading-ascending, those of the hermaphrodite flowers (8–)9–25(–33) mm long, those of the staminate flowers 2–5(–6) mm long. Hermaphrodite flowers (1–)2–6 per umbellet, (6–)10–22(–29) per umbel, staminate flowers 0–5 per hermaphrodite umbellet, 0–18(–24) per umbel; corolla purplish or greenish white, somewhat inconspicuous; styles (including stylopodium) (0.3–)0.4–0.8(–0.9) mm long, stylopodium 0.1–0.4 mm long, strongly depressed, with a conspicuous disc; carpophore cleft about one-fourth of its length. Fruit linear-fusiform, with an apical beak that is strongly con-

stricted below the stylopodium, concave furrowed, (7–)8–15(–16) mm long, the ribs glabrous above and slightly to moderately hispid with retrorse bristles below, the caudate appendages 1–5(–6) mm long. $2n = 22$ (Taylor & Mulligan, 1968). Figures 9i and 13.

Flowering period. Late April to July.

Habitat. Moist coniferous forests in areas with considerable, regular precipitation.

Common names. Purplish sweet cicely, Purple sweet cicely, Purple sweetroot.

Representative specimens. U.S.A. ALASKA: Deer Mt., Revillagigedo Island, *McCabe* 8644 (UC); Loring, *Chamberlain* 20 (US); Yes Bay, *Gorman* 23 (K, NY); Texas Creek Summit, NW of Hyder, *McCabe* 8934 (UC); Misty Fjords Nat'l. Mon., *Vorobik* 413 (ORE); Mt. Dewey, Wrangell Island, *Cowles* 1288 (ILL); Washington Bay, Kuiu Island, *Eyerdam* 8340 (G, SMU); Sitka, *Jepson* 489 (JEPS); Young Bay, Admiralty Island, *Robuck* 1311 (RM); Douglas Island, S of Juneau, *Trelease* 4527 (US); Juneau, *Anderson* 6322 (DAO, RM, TEX); Mt. Harris, St. Elias Mts., *Cowles* 1402 (ILL, MO); Thum Bay, Knight Island, Prince William Sound, 300 m, *Eyerdam* 3425 (ILL, K, MO, UC); Port San Juan, Evans Island, *Eyerdam* 5969 (ILL, MO); Mt. Marathon, W of Seward, *Calder* 5096 (DAO, NY, UC); Stetson Creek Valley, NW of Seward, *Calder* 6479 (DAO, WTU); Three Saints Bay, Kodiak Island, *Eyerdam* 386 (K). CALIFORNIA: Del Norte Co., Wilson Creek, between Requa and Crescent City, *Abrams & Bacigalupi* 8323 (CAS, RM); Redwood Nat'l. Park, *Lowry* 785a (ILL, UC); Siskiyou Co., *Muth* 396 (CAS). IDAHO: Bonner Co., Hope, *Dunkle* 457 (NY); Kootenay Co., Packsaddle Peak, 2,600 m, *Sandberg et al.* 845 (BM, K, MO, NY); Shoshone Co., Sohons Pass, 1,650 m, *Leiberg* 1427 (CAS, K, NY, RM, UC). MONTANA: Glacier Co., Midvale, *Umbach* 249 (MONT, RM, WIS); Park Co., 10 km W of Four Mile Ranger Sta., Boulder River Canyon, *Hitchcock* 16381 (CAS, NY, UC, WTU). OREGON: Clackamas Co., S slope of Mt. Hood, 1,250 m, *Lowry* 1073 (ILL, MO, OSC, UC); Clatsop Co., 1 km W of Washington Co. line on Hwy. 26, *Lowry* 3092 (ILL, MO, OSC, UC); Jefferson Co., E base of Three Fingerted Jack, 2,100 m, *Constance* 3465 (BM, ILL, NY, RM, UC, WIS); Klamath Co., Crater Lake, *Baker* 7126 (NY, WTU); Tillamook Co., Neahkahnie Mt., *Lowry* 910 (ILL, MO, NY, OSC, UC). WASHINGTON: Chelan Co., Stevens Pass, *Sandberg & Leiberg* 4028 (ILL, MICH, MO, NY, US, WIS, WTU); Grays Harbor Co., Colonel Bob Lookout, 1,250 m, *Thompson* 7275 (CAS, K, MO, UC, WTU); Jefferson Co., N Fork Quinault River, *Lowry* 254 (ILL); Okanogan Co., Mt. Bonaparte, *St. John et al.* 5341 (MO, UC, WTU); Snohomish Co., Perry Creek Trail, 1,000 m, *Thompson* 14712 (MICH, MO, NY, UC, WTU).

CANADA. ALBERTA: Cameron Lake, Waterton Lakes Nat'l. Park, 1,650 m, *Boivin & Gillett* 8979 (DAO). BRITISH COLUMBIA: 16 km N of Bull River, *Taylor & Ferguson* 2714 (DAO); S Fork Kaslo Creek, near Kookanee Glacier, *McCabe* 4779 (UC); Glacier Nat'l. Park, *Haber & Shchpanek* 1492A (CAS, NY); Illecillewaet Valley, Selkirk Mts., *Brown* 273 (MO, NY); Quiniscoe

Lake, Ashnola Range, *Calder et al.* 19635 (DAO, NY, UC); Manning Prov. Park, *Lowry* 1026 (ILL, UC); Copper Canyon, 800 m, *Schmidt* 51-53 (DAO); Lake Bootahnie, Marble Mts., 1,500 m, *Thompson & Thompson* 122 (WTU); Moat Lake, Forbidden Plateau, Vancouver Island, *Calder & MacKay* 32303 (DAO, UC); Indianpoint Lake, NE of Bakerville, *McCabe* 319 (UC); Mt. Fougner, Bella Coola, *Calder et al.* 20417 (DAO); Safety Cove, Calvert Island, *McCabe* 4189 (UC); Swanson Bay, Graham Reach, *McCabe* 3458 (UC); 5 km WNW of Tyee, E of Prince Rupert, *Calder et al.* 15038 (DAO); Bigsby Inlet, Moresby Island, *Calder et al.* 34908 (DAO); trail to Mercer Lake from Empire Anchorage, Graham Island, *Calder & Savile* 21494 (DAO, UC); Stewart, *Whited* 1226 (MO).

Osmorhiza purpurea is clearly a distinct species, although it is often confused with the closely related *O. chilensis*. *Osmorhiza purpurea* is the only taxon in the section *Nudae* that does not exhibit disjunctions to the Great Lakes region, northeastern North America, and southern South America. Constance and Shan (1948) noted that the fruits of this species seem equally well adapted for long-distance dispersal as those of *O. chilensis* and *O. depauperata*. *Osmorhiza purpurea*, however, has a very restricted habitat preference, occurring only in areas with substantial, regular precipitation, often at higher elevations or along the Pacific Coast, in sharp contrast with its more widespread relatives.

Coulter and Rose (1900) published two names for plants now included in *O. purpurea*: *Washingtonia leibergi* and *W. purpurea*. When these taxa were combined by Mathias and Constance (1944), they adopted the name *Osmorhiza purpurea*, and accordingly, under Art. 57 of the "Code" (Voss et al., 1983), this choice of epithet must be followed.

10. ***Osmorhiza depauperata*** Philippi, *Anales Univ. Chile* 85: 726. 1894. TYPE: Chile. Ñuble: Valle de las Nieblas, near Termas de Chillán, *F. Philippi* 2030 [lectotype (designated herein), SGO (no. 041589); photograph of lectotype, ILL!; isolectotype, SGO (no. 053461); photograph of isolectotype, ILL!].

Washingtonia obtusa Coult. & Rose, *Contr. U.S. Natl. Herb.* 7: 64. 1900. *Osmorhiza obtusa* (Coult. & Rose) Fern., *Rhodora* 4: 154. 1902. TYPE: U.S.A. Wyoming: Park Co., Ishawood (i.e., Ishawooa) Creek, *J. N. Rose* 476 (holotype, US!).

Osmorhiza obtusa var. *cupressi-montanum* Boivin, *Canad. Field-Naturalist* 65: 20. 1951. *Osmorhiza chilensis* var. *cupressimontana* (Boivin) Boivin, *Naturaliste Canad.* 93: 644. 1966. TYPE: Canada.

Saskatchewan: Cypress Hills Park, aspen woods, *A. J. Breitung* 4742 (holotype, DAO!).

Plants slender, 1.5–6.5(–8) dm high, stems 1–3(–5), erect to spreading-ascending or decumbent, sparsely to rather densely hirsutulous, or sometimes glabrescent. Root system rather deep, well developed, the roots with a weak carrot- or anise-like scent. Leaves 2–3-ternate, orbicular to broadly ovate, 4–10(–12) cm long, sparingly hispidulous to glabrous; leaflets broadly lanceolate to ovate, 1.5–5(–6) cm long, 1–3 cm wide, obtuse or acute, coarsely serrate to incised, parted, or divided at the base; petioles 3–20(–23) cm long. Umbels open, peduncles 1–3, terminal and often lateral, 5–15 cm long; involucre wanting, or rarely consisting of 1 minute (to very rarely large and leaf-like), linear-lanceolate, foliaceous, ciliate, spreading bract; rays widely spreading to nearly reflexed, (2.5–)3–9(–10) cm long; umbellets 2–6 per umbel; involucel wanting, or rarely of 1–2 small, linear-lanceolate, foliaceous, ciliate bractlets, each 0.3–2.5 mm long, 0.2–0.4 mm wide; pedicels (2–)3–7 per umbellet, widely divaricate, those of the hermaphrodite flowers (4–)8–12(–32) mm long, those of the staminate flowers 2–12(–13) mm long. Hermaphrodite flowers 2–6 per umbellet, (3–)7–20(–25) per umbel, staminate flowers 0–2(–3) per umbellet, 0–10(–15) per umbel; corolla greenish white, rather inconspicuous; styles (including stylopodium) 0.3–0.6(–0.7) mm long, stylopodium (0.1–)0.2–0.4(–0.5) mm long, low-conic to depressed, with or without a disc; carpophore cleft about one-third of its length. Fruit clavate, obtuse at the apex, concave furrowed, (9–)10–18(–19) mm long, the ribs moderately to densely hispid with retrorse bristles, especially toward the base, the caudate appendages (2.5–)3–8.5(–9) mm long. $n = 11$ (Bell & Constance, 1960; Constance et al., 1976); $2n = 22$ (Crawford & Hartman, 1972). Figures 9k and 14.

Flowering period. April to early July (North America); November and December (South America).

Habitat. Moist to fairly dry forests, woodlands, and open slopes.

Common names. Blunt-fruit sweet cicely, Blunt-fruit sweetroot, Blunt-fruited sweet cicely, Bluntseed sweetroot, Snub cicely, Sweet cicely.

Representative specimens. U.S.A. ALASKA: Denver Trail, E of Skagway, *Anderson* 1735 (K); Pt. Gustavus, Glacier Bay, *Coville & Kearney* 729 (US); Valdez, *Piper* 4288 (US); Kenai Lake, *Calder* 5688 (DAO); Cooper

Landing, W of Kenai Lake, *Anderson 6882* (RM); 12 km E of mouth of Moore River, Kenai Peninsula, *Lutz 105* (RM). ARIZONA: Apache Co., Lukachukai Mts., 2,700 m, *Goodman & Payson 2816* (CAS, MONT, MONTU, NY, UC, WTU); Cochise Co., Chiricahua Mts., *Ferris 9950* (MICH, TEX, UC); Coconino Co., Humphreys Peak, San Francisco Mts., *MacDougal 397* (ARIZ, G, NY, P, RM, UC); Pima Co., Santa Catalina Mts., *Goodman & Hitchcock 1253* (MO, NY, RM). CALIFORNIA: Modoc Co., Warner Mts., 1,850 m, *Constance 3104* (K, MO, NY, UC). COLORADO: Boulder Co., Boulder Creek, below Eldora, 2,500 m, *G. N. Jones 33299* (CS, ILL); Clear Creek Co., Georgetown, 2,500 m, *M. E. Jones 514* (BR, CAS, MISSA, NY, UC); Delta Co., Cedaredge, 2,150 m, *Baker 240* (K, RM, UC); El Paso Co., Jack Brook, 2,600 m, *Clements & Clements 236* (NY, RM); Garfield Co., S of Rifle, *Osterhout 2151* (NY, RM, WIS); Gunnison Co., Elko Park, N of Gothic, *Mathias 3264* (UC); La Plata Co., La Plata, *Baker et al. 849* (BM, K, MICH, MO, NY, RM); Mineral Co., near Pagosa Peak, 2,750 m, *Porter 3932* (CAS, RM, UC, WTU); Montezuma Co., Upper W Mancos Canyon, 3,000 m, *Baker et al. 188* (BM, G, K, MICH, MO, NY, RM); Rio Blanco Co., Black Sulphur Creek, 5.5 km SW of confluence with Swizer Gulch, 2,115 m, *W. R. Erickson 594* (CS); Summit Co., near Breckenridge, *MacKenzie 295* (MO, RM, WIS). IDAHO: Bingham Co., Wolverine Creek, Blackfoot Mts., *Holmgren & Murttala 5415* (NY, UC); Bonneville Co., 18 km SW of Victor, *Lowry 1114* (ILL); Cassia Co., Pole Canyon, SE of Burley, 2,000 m, *Holmgren & Jensen 3499* (NY, WTU); Clark Co., Centennial Mts., 2,350 m, *Lowry 2485* (ILL, MONT, UC); Custer Co., Challis Creek, 1,800 m, *Macbride & Payson 3332* (CAS, K, MO, NY, RM, UC); Fremont Co., Henry's Lake, 1,800 m, *Payson & Payson 1946* (CAS, MO, NY, RM). MICHIGAN: Keweenaw Co., Isle Royale, *McFarlin 2049* (MICH, MONT); Leelanau Co., E end of Manitou Island, *Richards 3123* (MICH). MINNESOTA: Cook Co., Mountain Lake, *Butters et al. 303* (NY). MONTANA: Beaverhead Co., Centennial Mts., 2,450 m, *Lowry 2426* (ILL, MO, MONT, MONTU, NY, RM, UC); Carbon Co., Red Lodge, *Hawkins*, 22 July 1919 (MONT); Deerlodge Co., Storm Lake, *Hitchcock & Muhlick 14845* (MO, NY, UC, WTU); Gallatin Co., Mystic Lake, *Blankinship 220* (BM, MO, MONTU); Silver Bow Co., Humbug Spires, 1,900 m, *Lowry 2269* (ILL, MONT, UC). NEVADA: Clark Co., Little Falls, Charleston Mts., 2,400 m, *Clokey 8050* (ARIZ, CAS, ILL, MO, MONTU, NY, RM, UC, WTU); Elko Co., Lamoille Canyon, 2,450 m, *Millan & McKnight 168* (NY); White Pine Co., Snake Creek, Snake Range, 2,950 m, *Holmgren & Reveal 1627* (NY). NEW MEXICO: Grant Co., Hillsboro Peak, Black Range, 3,050 m, *Metcalf 1206* (MO, NY); Lincoln Co., 11 km W of Ruidoso, 2,750 m, *Hinckley 726* (ARIZ, NY, TEX, UC); Otero Co., Cloudcroft, *Wooton*, 30 June 1899 (ILL, MO, NMC, NY, RM); Santa Fe Co., Santa Fe Canyon, E of Santa Fe, *Heller & Heller 3822* (BM, CAS, K, MO, NY, P, UC). OREGON: Deschutes Co., Paulina Creek, 2,090 m, *Leiberg 555* (K, NY, UC); Jackson Co., Wimer, *Hammond 157a* (CS); Klamath Co., Crater Lake Nat'l. Park, 1,850 m, *Heller 13469* (CAS, ILL, WTU). SOUTH DAKOTA: Lawrence Co., S of Deadwood, *Churchill 5707* (MO, NY). UTAH: Duchesne Co., 6 km S of Moon Lake, 2,350 m, *Harrison & Larsen 7612*

(MO, RM); Garfield Co., Henry Mts., 2,400 m, *McVaugh 14681* (CAS, MICH, NY, SMU, TEX); Grand Co., Post Canyon, Uintah and Ouray Ind. Res., 110 km S of Ouray, 2,650 m, *Holmgren et al. 2313* (ARIZ, CAS, ILL, NY, TEX, UC, WIS, WTU); San Juan Co., Abajo Mts., *Goodman & Hitchcock 1437* (BM, MO, NY, RM); Utah Co., Slate Canyon, E of Provo, 2,450 m, *M. E. Jones 5580* (BM, MO, NY, RM, UC); Washington Co., 4.5 km E of Pine Valley, 2,150 m, *Gould 1793* (NY). VERMONT: Chittenden Co., Mt. Mansfield, *Greenman 691* (MO). WASHINGTON: Asotin Co., *Cronquist & Jones 5860* (WTU); Chelan Co., Wenatchie region, *Brandegge 794* (NY, UC); King Co., Seattle, *Talcott*, 15 May 1891 (MICH); Okanogan Co., Myers Creek, Chesaw, *St. John et al. 5186* (MO, NY, UC, WTU). WYOMING: Albany Co., Laramie Range, 2,500 m, *Hartman 3122* (RM); Big Horn Co., 42 km W of Burgess Junction, Big Horn Mts., *Lowry 1423* (ILL); Carbon Co., S Brush Creek Campground, Medicine Bow Mts., *Lowry 1132* (ILL, MO, NY, RM, UC); Sublette Co., Middle Piney Lake, 2,600 m, *Meyer & Meyer 2402* (MO, NY, UC); Teton Co., 5 km W of Teton Pass, *Lowry 1119* (ILL, MO, NY, RM, UC).

CANADA. ALBERTA: Cypress Hills, 1,450 m, *Boivin & Gillett 8929* (DAO); Chief Mt. Hwy., Waterton Lakes Nat'l. Park, 1,450 m, *Breitung 15881* (NY, UC); W face of Sandhills, *Brinkman 5470* (DAO, SMU); Elbow River Valley, Calgary, *Moodie 1056* (NY, UC); 3 km below Canmore, 1,300 m, *Ewan 18685* (UC); Elk Island Park, E of Ft. Saskatchewan, *Turner 4498* (DAO); Battle Lake, SW of Pigeon Lake, *Turner 5907* (DAO); Elysium Pass Trail, Jasper Nat'l. Park, *Lowry 944* (ILL, UC); Geraldine Lakes Trail, Jasper, *Lowry 961* (ILL, UC); near Atauwan Cabin, Slave Lake Dist., *Brinkman 4335* (NY); SW of Spirit River, *Moss 8415* (DAO); Saddle Hills, N of Sexsmith, *Moss 9948* (WIS); Wembley, *Wallace for Jenkins 734* (DAO). BRITISH COLUMBIA: Starvation Park, W of Waterton Lakes, 1,550 m, *Calder et al. 3416* (DAO); Sheep Mt., 55 km N of Natal, *Weber 2373* (NY, UC); 7 km WSW of Wycliffe, 950 m, *Taylor et al. 2635* (DAO, UC); 5 km W of Needles, *McCabe 6663* (UC); Friday Creek, 25 km S of Princeton, *Lowry 1021* (ILL, UC); Yoho Nat'l. Park, *Lowry 1005* (ILL, MO, RM, UC); 1.5 km NW of Falkland, 600 m, *Hitchcock & Martin 7466* (NY, RM); Botanie Valley, 1,200 m, *Beamish et al. 8204* (DAO); Three Mile, on Lac La Hache-Williams Lake Rd., *Calder et al. 18970* (DAO); 24 km SW of Kleena Kleene, *McCabe 606* (UC); 21 km W of Puntataenkut Lake, W of Quesnel, 1,150 m, *Calder et al. 18212* (DAO); Tezsli, Kluskus Lakes, *Lewicki 79* (WIS); Prince George, 600 m, *Florian 94* (DAO); 25 km W of Burns Lake, *Calder et al. 12780* (DAO); 5 km S of Ft. McLeod, *Calder et al. 12445* (DAO, UC); Pine River, 67 km W of Bissett Creek Bridge, W of Dawson Creek, *Taylor & Staudt 4169* (DAO, UC); near Hudson's Hope, *Raup & Abbe 3640* (NY); Wicked River, near Peace River, 600 m, *Raup & Abbe 3878* (NY); Wolverine Ridge, near Manson Creek Village, 1,050 m, *Calder et al. 13702* (DAO); 5 km S of Takla Landing, *McCabe 7916* (UC); Bear Lake, 40 km NW of Takla Lake, *McCabe 7998* (UC); Fairy Lake, SSW of Ft. Nelson, 1,400 m, *Calder & Kukkonen 27162* (DAO, UC); Laird Hot-springs, 500 m, *Calder & Gillett 25575* (CAS, DAO, UC); mile 578, Alaska Hwy., 550 m, *Calder & Gillette 25655* (DAO). MANITOBA: Mont Dauphin, Norgate,

Boivin et al. 10682 (DAO); 13 km W of Norgate, Riding Mt. Nat'l. Park, *Löve & Löve* 6563 (DAO); Forestry Camp, Riding Mt. Nat'l. Park, *Rowe* 305 (DAO). NORTHWEST TERRITORIES: Charlton Island, James Bay, *Porsild* 4506 (DAO—photo ex CAN); Mt. Coty, 750 m, *Cody & Spicer* 11787 (NY, SMU, UC). NEW BRUNSWICK: Madawaska Co., Claire, *Eaton M-89* (LL). NEWFOUNDLAND: Dildo, Notre Dame Bay, *Fernald & Wiegand* 5952 (K, NY); Benoit's Cove, Humber Arm, Bay of Islands, *Fernald & Wiegand* 3767 (BM); Frenchman's Cove, Bay of Islands, *Waghorne* 15 (MO); Port au Port, *MacKenzie*, 30 July 1921 (NY, UC); Nameless Cove—Mistaken Cove, Straights of Belle Isle, *Wiegand et al.* 28767 (GH, US); Forteau, Labrador, *Waghorne*, 8 Aug. 1893 (MO). NOVA SCOTIA: Inverness Co., Cape St. Lawrence, *Smith et al.* 11183 (DAO). ONTARIO: Algoma Dist., *Marie-Victorin & Rolland-Germain* 27340 (CAS); Thunder Bay Dist., Mortimer Island, Slate Islands, *Hosie et al.* 2187 (UC); Pijitiwabik Bay, Lake Nipigon, *Garton* 7248 (DAO). QUÉBEC: Gaspé Co., Grand River, *Fernald s.n.* (MICH, NY); Lac St. Jean—Ouest Co., Chambord, Roberval, *Leduc L-69-68* (DAO); Matane Co., Little Metis, *Fowler* 25351 (MO, WIS); Rimouski Co., near Bic, Isle de Bic, *Forbes*, 25 June 1905 (RM); Rivière-du-Loup Co., Pt. de la Rivière du Loup, *Cayouette & Cinq-Mars*, 19 June 1944 (DAO); Saguenay Co., Rivière du Renard, Anticosti Island, *Marie-Victorin & Rolland-Germain* 27139 (MO, WIS); Niapisca, Mingan Islands, *Hamel & Genereux* 3140 (DAO). SASKATCHEWAN: Maple Creek Dist., Cypress Hills, *Hudson* 1838 (DAO); Melfort Dist., McKague, *Breitung*, 30 Jan. 1935 (DAO); Qu'Appelle Dist., Cherry Lake, SE of Indian Head, *Jones & Ledingham* 742 (DAO); Wallwort, *Breitung* 651 (DAO, MO, UC).

ARGENTINA. CHUBUT: Río Senguerr, Lago La Plata, *Krapovickas* 4241 (UC). NEUQUÉN: Los Lagos, Fortín Chacabuco, *Correa & Constance* 3788 (UC); between Lago Correntoso and Lago Escondido, *Correa & Constance* 3845 (G, K, UC); Correntoso, Lago Nahuel Huapí, *Edwards s.n.* (BM). RÍO NEGRO: Bariloche, Parque Nac. Nahuel Huapí, *Boelcke* 5405, 5436 (UC); Estacis Lago Roca, 350 m, *James* 411 (BM). SANTA CRUZ: Güer Aike, Estancia Stag River, Río Gallegos, *Tweedie* 101a, 267 (K); Lago Argentino, Brazo Norte, valle de La Cristina, 850 m, *Vervoorst* 4387 (MO). TIERRA DEL FUEGO: Ushuaia, Lago Fagnano, *Boelcke et al.* 15236 (UC), *Moore* 2845 (K); Harberton, *Constance et al.* 3861 (UC), *Goodall* 164, 1025 (UC); Bahía Aguirre, *Goodall* 1422 (UC); Laguna Victoria, Isla de Los Estados, *Goodall* 1579 (UC), *Moore* 2079 (K); Estancia Viamonte, *Goodall* 2823 (UC); Río Valdez, *Goodall* 3405 (UC); Monte de las Carpas, *Goodall* 4669 (UC); Ushuaia, *Hunziker* 6522, 6560 (UC), *Vallerini* 66 (UC), *Zöllner* 8584 (MO), *Pennington* 196 bis (Z); Monte Olivia, *Hunziker* 8206 (UC); Estancia la Esperanza, 210 m, *Mexia* 7925 (BM, G, K, MO, NY, UC), *Moore* 1540 (K); Sierra Alvaer, E of Paso Garibaldi, *Moore* 1818 (K).

CHILE. MAGALLANES: Rosvig, 200 m, *Donat* 346 (BM, CAS, K, NY); Sandy Pt., *Cunningham s.n.* (NY), *DeCandolle m.* 1186 (P); Buncombe Bay, *Commerson s.n.* (P); Punta Arenas, 20 m, *Eyerdam et al.* 24105 (G, K, UC), *Stafford* 26 (NY), *Savatier* 61, 76 (P), *McLean D.A.14* (BM), *Hatcher s.n.* (NY); Estancia Maria Cristina, 80 km NE of Punta Arenas, *Goodall*

4018 (UC); Puerto Williams, Navarino Island, *Godley* 884 (K, UC); Pecket Harbor, *Hombroon s.n.* (P); Orange Harbor, *Hyades* 472 (P), *U.S. S Pac. Expl. Exped.* (P); Port Famine, *Le Guillou s.n.* (P), *Marivault s.n.* (P); Estancia Cameron, NW shore Lago Blanco, *Moore* 2145 (K, UC); Estancia Vicuña, 200 m, *Moore* 2177 (UC); Port Galant, *Savatier s.n.* (P). ÑUBLE: Termas de Chillán, *Jaffuel* 3765 (GH, cited in Constance & Shan, 1948). VALDIVIA: Valdivia, *Lechler* 225 (GOET); *Philippi s.n.* (G).

Osmorhiza depauperata was described by Philippi (1894) from material collected in South America. At the time Constance and Shan (1948) prepared their treatment of *Osmorhiza*, they were uncertain as to the identity of this name, and tentatively placed it in synonymy under *O. chilensis*. Instead, they accepted the name *Osmorhiza obtusa* for the taxon with clavate fruit and divaricate rays and pedicels. In 1954, Constance visited the Museo Nacional de Historia Natural in Santiago, Chile, where he examined the type material of *Osmorhiza depauperata* (Constance, pers. comm.). Three years later, this name was first used for North American plants by Mathias and Constance (1957: 11), with the following footnote: "Examination of Philippi's type at Santiago, Chile, has verified the necessity of substituting this name [*O. depauperata*] for the heretofore accepted *O. obtusa* (C. & R.) Fernald." Although Philippi's type material is not available for loan, we were able to obtain photographs of the two specimens through the courtesy of Dra. Mélica Muñoz S., Curator of Botany at SGO. While examination of these photographs does not, by itself, reveal with certainty the identity of Philippi's specimens, information contained in Constance's notes on them indicates that the name *Osmorhiza depauperata* does indeed apply to the taxon earlier referred to as *O. obtusa*.

LITERATURE CITED

- ABLAEV, A. G., I. V. VASSILIEV & P. G. GOROVOL. 1974. Floristic associations between east Asia and North America: a study of the ranges of some recent and fossil higher plants in the Primorye, U.S.S.R. *Paleobotanist* 23: 49–54.
- BELL, C. R. 1971. Breeding systems and floral biology of the Umbelliferae. Pp. 93–107 in V. H. Heywood (editor), *The Biology and Chemistry of the Umbelliferae*. Academic Press, New York and London.
- & L. CONSTANCE. 1957. Chromosome numbers in Umbelliferae. *Amer. J. Bot.* 44: 565–572.
- & ———. 1960. Chromosome numbers in Umbelliferae. II. *Amer. J. Bot.* 47: 24–32.
- & ———. 1966. Chromosome numbers in Umbelliferae. III. *Amer. J. Bot.* 53: 512–520.
- BENTHAM, G. 1867. Umbelliferae. Pp. 859–931 in G.

- Bentham & J. D. Hooker (editors), *Genera Plantarum*, Volume 1. Reeve and Co., London.
- BLAKE, S. F. 1923. The forms of *Osmorhiza longistylis*. *Rhodora* 25: 110–111.
- BOIVIN, B. 1968. Flora of the Prairie Provinces. *Phytologia* 17: 57–112.
- BONGARD, M. 1833. Observations sur la végétation de l'île de Sitcha. *Mém. Acad. Imp. Sci. St. Pétersbourg, Sér. 6, Sci. Math.* 2: 119–177.
- CAMPBELL, D. H. 1944. Relations of the temperate floras of North and South America. *Proc. Calif. Acad. Sci.* 25: 139–146.
- CLARKE, C. B. 1879. Umbelliferae. Pp. 665–720 in J. D. Hooker (editor), *Flora of British India*, Volume 2. Reeve and Co., Ashford, Kent, England.
- CLOS, D. 1848. Umbeliferas. Pp. 61–145 in C. Gay, *Flora Chilena*, Volume 3, part 2. Paris, Santiago.
- CONSTANCE, L. 1963. Amphitropical relations in the herbaceous flora of the Pacific Coast of North and South America: a symposium. Introduction and historical review. *Quart. Rev. Biol.* 38: 109–116.
- . 1972. Patterns in the distribution of Japanese–American Umbelliferae. Pp. 93–99 in A. Graham (editor), *Floristics and Paleofloristics of Asia and Eastern North America*. Elsevier Publ. Co., Amsterdam, London, New York.
- & R. H. SHAN. 1948. The genus *Osmorhiza* (Umbelliferae), a study in geographic affinities. *Univ. Calif. Publ. Bot.* 23: 111–156.
- , T.-I. CHUANG & C. R. BELL. 1976. Chromosome numbers in Umbelliferae. *V. Amer. J. Bot.* 63: 608–625.
- COULTER, J. M. & J. N. ROSE. 1888. Revision of North American Umbelliferae. Published by the authors, Crawfordsville, Indiana.
- & ———. 1895. Report on Mexican Umbelliferae, mostly from the state of Oaxaca, recently collected by C. G. Pringle and E. W. Nelson. *Contr. U.S. Natl. Herb.* 3: 289–309.
- & ———. 1900. Monograph of North American Umbelliferae. *Contr. U.S. Natl. Herb.* 7: 9–256.
- CRAWFORD, D. J. & R. L. HARTMAN. 1972. Chromosome numbers and taxonomic notes for Rocky Mountain Umbelliferae. *Amer. J. Bot.* 59: 386–392.
- DEAM, C. C. 1940. *Flora of Indiana*. Dept. of Conservation, Div. of Forestry, Indianapolis.
- DIETRICH, D. 1840. *Synopsis Plantarum*, Volume 2. Voigt, Weimar.
- DRUDE, C. G. O. 1897. Umbelliferae. Pp. 63–250 in A. Engler & K. Prantl (editors), *Die Natürlichen Pflanzenfamilien*, Volume 3⁸. Engelmann, Leipzig.
- DURIETZ, G. E. 1940. Problems of bipolar plant distribution. *Acta Phytogeogr. Suec.* 13: 215–282.
- ENGLER, A. 1882. Versuch einer Entwicklungsgeschichte der Pflanzenwelt, insbesondere der Florengebiete seit der Tertiärperiode. Engelmann, Leipzig.
- ERDTMAN, G. 1969. *Handbook of Palynology*. Scandinavian Univ. Books, Munksgaard.
- FARWELL, O. A. 1925. Botanical gleanings in Michigan. II. *Amer. Midl. Naturalist* 9: 257–282.
- FERNALD, M. L. 1908. Notes on some plants of northeastern America. *Rhodora* 10: 46–55.
- . 1924. Isolation and endemism in northeastern America and their relation to the age-and-area hypothesis. *Amer. J. Bot.* 11: 558–572.
- . 1925. Persistence of plants in unglaciated areas of boreal America. *Mem. Amer. Acad. Arts* 15: 239–342.
- . 1926. Two summers botanizing in Newfoundland. *Rhodora* 28: 49–63, 74–87, 89–111, 115–129, 145–155, 161–178, 181–204, 210–225, 234–241.
- . 1933. Recent discoveries in the Newfoundland flora. *Rhodora* 35: 1–16, 47–63, 80–107, 120–140, 161–185, 203–223, 230–247, 265–283, 298–315, 327–346, 364–386, 395–403.
- . 1935. Critical plants of the Upper Great Lakes region of Ontario and Michigan. *Rhodora* 37: 197–222, 238–262, 272–301, 324–341.
- FRENCH, D. H. 1971. Ethnobotany of the Umbelliferae. Pp. 385–412 in V. H. Heywood (editor), *Biology and Chemistry of the Umbelliferae*. Academic Press, New York and London.
- GILMORE, M. R. 1919. Uses of plants by the Indians of the Missouri River region. *Smithsonian Inst. Bur. Amer. Ethnol. Annual Rep.* 33: 43–154.
- GRAHAM, A. 1966. *Plantae Rariores Camschatcensis: a translation of the dissertation of Jonas P. Halenius, 1750*. *Brittonia* 18: 131–139.
- GRAY, A. 1854. *Botany of the United States Expedition During the Years 1838–1842 Under the Command of Charles Wilkes, U.S.N. Volume 1. Phanerogamia*. Sherman, Philadelphia.
- . 1859. Diagnostic characters of new species of phanerogamous plants, collected in Japan by Charles Wright, botanist of the U.S. North Pacific exploring expedition. With observations upon the relations of the Japanese flora to that of North America, and of other parts of the northern temperate zone. *Mem. Amer. Acad. Arts n.s.* 6: 377–452.
- . 1868. Characters of new plants of California and elsewhere, principally of those collected by H. N. Bolander in the State Geological Survey. *Proc. Amer. Acad. Arts* 7: 327–401.
- & J. D. HOOKER. 1880. *The vegetation of the Rocky Mountain region and a comparison with that of other parts of the world*. U.S. Geol. Geogr. Surv. Terr., Bull. 6: 1–77.
- GREENE, E. L. 1892. *Flora Franciscana*. Part 3. Doxey and Co., San Francisco; Wesley and Son, London; Cubery and Co., San Francisco.
- GUNTHER, E. 1945. Ethnobotany of western Washington. *Univ. Wash. Publ. Anthropol.* 10: 1–61.
- HARA, H. 1952. Contributions to the study of variations in the Japanese plants closely related to those of Europe and North America. I. *J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot.* 6: 29–96.
- . 1956. Contributions to the study of variations in the Japanese plants closely related to those of Europe and North America. II. *J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot.* 6: 343–391.
- . 1972. Corresponding taxa in North America, Japan and the Himalayas. Pp. 61–72 in D. H. Valentine (editor), *Taxonomy, Phytogeography and Evolution*. Academic Press, London and New York.

- HAYATA, B. 1911. Materials for a flora of Formosa. J. Tokyo Univ. Fac. Sci. 30:1-471.
- . 1912. Icones Plantarum Formosanarum, Volume 2. Taipei.
- HEMSLEY, W. B. 1880. Biologia Centrali-Americana, Volume 1, part 6. R. H. Porter, London.
- HOOKE, J. D. 1846. I. Flora Antarctica. Part II. Botany of Fuegia, the Falklands, Kerguelen's Land, etc. Pp. 209-579 in The Botany the Antarctic Voyage of H.M. Discovery Ships Erebus and Terror, in the Years 1839-1843. Reeve Brothers, London.
- HOOKE, W. J. 1840. Flora Boreali-americana, Volume 1. Henry G. Bonn, London.
- & G. A. WALKER ARNOTT. 1833. Contributions toward a flora of South America and the islands of the Pacific. Pp. 302-367 in W. J. Hooker (editor), Botanical Miscellany, Volume 3. J. Murray, London.
- HULTÉN, E. 1947. Flora of Alaska and Yukon, Volume 7. Håkan Ohlssons Boktryckeri, Lund, Sweden.
- JACKSON, B. D. 1893. Bibliographical notes. II. 'Botany of Beechey's Voyage' and 'Flora of North America.' J. Bot. 31: 297-299.
- KAPP, R. O. 1969. Pollen and Spores. Brown and Co., Dubuque, Iowa.
- KOSO-POLJANSKY, B. 1916. Sciadophytorum systematicis lineamenta. Bull. Soc. Imp. Naturalistes Moscou 29: 93-221.
- KUNTZE, O. 1891. Revisio Generum Plantarum. Felix, Leipzig.
- LI, H.-L. 1952. Floristic relationships between eastern Asia and eastern North America. Trans. Amer. Philos. Soc. 42: 371-429.
- . 1972. Eastern Asia-eastern North America species-pairs in wide-ranging genera. Pp. 65-78 in G. Graham (editor), Floristics and Paleofloristics of Asia and Eastern North America. Elsevier Publ. Co., Amsterdam, London, New York.
- LINNAEUS, C. 1750. Plantae Rariores Camschatcensis. Uppsala. [Reprinted in Amoen. Acad. 2: 332-364. 1751.]
- LIU, T.-S. 1977. Umbelliferae. Pp. 938-974 in H.-L. Li, T.-S. Liu, T.-C. Huang, T. Koyama & C. E. DeVol (editors), Flora of Taiwan, Volume 3. Epoch Publ. Co., Ltd., Taipei.
- LOWRY, P. P., II. 1976. A biosystematic study of two species in the genus *Osmorhiza* (Umbelliferae). B.S. thesis. Univ. of Illinois, Urbana.
- . 1980. A biosystematic study and taxonomic revision of *Osmorhiza* Rafinesque (Apiaceae). M.S. thesis. Univ. of Illinois, Urbana.
- . 1985. Proposal to conserve 5941 *Osmorhiza* Raf. against *Uraspermum* Nutt. (Apiaceae). Taxon 34: 316-318.
- & A. G. JONES. 1977. Nomenclatural notes on *Osmorhiza* Rafinesque (Apiaceae) and two proposals for additions to Article 34 of the Code. Taxon 26: 567-569.
- & ———. 1978. Correction of the publication date for the generic name *Osmorhiza* Rafinesque. Taxon 27: 550.
- & ———. 1979a. Biosystematic investigations and taxonomy of *Osmorhiza* Rafinesque section *Osmorhiza* (Apiaceae) in North America. Amer. Midl. Naturalist 101: 21-27.
- & ———. 1979b. Biosystematics and taxonomy of *Osmorhiza* Rafinesque (Apiaceae: Apioideae). Bot. Soc. Amer., Misc. Ser., Publ. 157: 62 (abstract).
- & D. A. YOUNG. 1979. Water soluble seed proteins and the taxonomy of *Osmorhiza* (Apiaceae). Bot. Soc. Amer., Misc. Ser., Publ. 157: 45 (abstract).
- MACMILLAN, C. 1892. The Metaspermae of the Minnesota Valley. Rep. Minnesota Geol. Nat. Hist. Surv. Bot. Ser. 1: 1-839.
- MAKINO, T. 1903. Observations on the flora of Japan. Bot. Mag. (Tokyo) 17: 6-14.
- MARQUIS, R. J. & E. G. VOSS. 1981. Distributions of some western North American plants disjunct in the Great Lakes region. Michigan Bot. 20: 53-82.
- MATHIAS, M. E. & L. CONSTANCE. 1944. Umbelliferae. N. Amer. Fl. 28B: 43-295.
- & ———. 1957. Umbelliferae of Nevada. Contr. Flora Nevada 44: 1-60.
- MICHAUX, A. 1803. Flora Boreali-americana, Volume 1. Levrault, Paris.
- MOORE, D. M. 1972. Connections between cool temperate floras, with particular reference to southern South America. Pp. 115-138 in D. H. Valentine (editor), Taxonomy, Phytogeography and Evolution. Academic Press, London and New York.
- MUHLENBERG, H. 1813. Catalogus Plantarum Americae Septentrionalis. Hamilton, Lancaster, Pennsylvania.
- NUTTALL, T. 1818. The Genera of North American Plants, and a Catalogue of the Species, to the Year 1817, Volume 1. Published by the author, Philadelphia.
- OSTERTAG, C. P. & R. J. JENSEN. 1980. Species and population variability of *Osmorhiza longistylis* and *O. claytonii*. Ohio J. Sci. 80: 91-95.
- PERSOON, C. H. 1805. Synopsis Plantarum seu Enchiridium Botanicum, Volume 1. J. G. Cotta, Tübingen.
- PHILIPPI, R. A. 1894. Plantas nuevas Chilenas. Anales Univ. Chile 85: 699-749.
- RAFINESQUE, C. S. 1818. A review of Pursh's Flora of North America. Amer. Monthly Mag. & Crit. Rev. 2: 170-176.
- . 1819a. [Review of:] Thomas Nuttall. The Genera of North-American Plants and a Catalogue of the Species to the Year 1817. 2 Volumes. Philadelphia. 1818. Amer. Monthly Mag. & Crit. Rev. 4: 184-196.
- . 1819b. Remarques critiques et synonymiques. J. Phys. Chim. Hist. Nat. Arts 89: 256-261.
- . 1830. Medical Flora, Volume 2. Atkinson, Philadelphia.
- RAVEN, P. H. 1963. Amphitropical relationships in the floras of North and South America. Quart. Rev. Biol. 38: 151-177.
- REICHE, K. 1902. Flora de Chile, Volume 3. Imprunta, Lithographía i Encuadernacion Barcelona, Santiago.
- . 1907. Grundzüge der Pflanzenverbreitung in Chile. In A. Engler & O. Drude (editors), Die Vegetation der Erde, Volume 8. Engelmann, Leipzig.
- RICKETT, H. W. 1945. Bibliography (Umbellales,

- Cornales, Cornaceae, Nyssaceae). *N. Amer. Fl.* 28B: 317-374.
- ROBERTSON, C. 1888. Proterogynous Umbelliferae. *Bot. Gaz.* (Crawfordsville) 13: 193.
- ROYLE, J. F. 1835. Illustrations of the Botany and Other Branches of the Natural History of the Himalayan Mountains and of the Flora of Cashmere, Volume 1. W. H. Allen and Co., London.
- RUNE, O. 1954. Notes on the flora of the Gaspé Peninsula. *Svensk Bot. Tidskr.* 48: 117-136.
- RYDBERG, P. A. 1894. A revision of the nomenclature of the Nebraska Polypetalae. *Bot. Surv. Nebraska* 3: 20-39.
- SALAMUN, P. J. 1950. The interpretation of variation in populations of *Spiraea tomentosa*, *Brachyelytrum erectum*, *Osmorhiza claytoni* [sic], and *Osmorhiza longistylis*. Ph.D. dissertation. Univ. of Wisconsin, Madison.
- . 1952. A Black Hills variety of *Osmorhiza longistylis*. *Amer. Midl. Naturalist* 47: 251-253.
- SCHENCK, S. M. & E. W. GIFFORD. 1952. Karok ethnobotany. *Anthropol. Rec.* 13: 377-392.
- SCHLESSMAN, M. A. 1978. Systematics and reproductive biology of *Lomatium farinosum* (Geyer ex Hooker) Coulter & Rose (Umbelliferae). *Madroño* 25: 1-9.
- . 1982. Expression of andromonoecy and pollination of tuberous lomatiums (Umbelliferae). *Syst. Bot.* 7: 134-149.
- SCHNEIDER, A. 1906. The medicinal plants of the California Indians. *Merck's Rep.* 15: 95-96.
- SCHOFIELD, W. B. 1969. Phytogeography of northwestern North America: bryophytes and vascular plants. *Madroño* 20: 155-207.
- SHISHKIN, B. K. (editor). 1950. Flora of the U.S.S.R., Volume 16. Umbelliferae. Keter Press, Jerusalem. (English translation by R. Lavoott, 1973.)
- SMITH, H. H. 1923. Ethnobotany of the Menomini Indians. *Bull. Public Mus. Milwaukee* 4: 1-174.
- . 1928. Ethnobotany of the Meskwaki Indians. *Bull. Public Mus. Milwaukee* 4: 175-326.
- . 1932. Ethnobotany of the Ojibwe Indians. *Bull. Public Mus. Milwaukee* 4: 327-525.
- SPRENGEL, C. 1813. *Plantarum Umbelliferarum*. Hendel, Halle.
- STEBBINS, G. L., JR. 1935. Some observations on the flora of the Bruce Peninsula, Ontario. *Rhodora* 37: 63-74.
- STEEDMAN, E. V. (editor). 1930. Ethnobotany of the Thompson Indians of British Columbia. Based on field notes by J. A. Teit. *Smithsonian Inst. Bur. Amer. Ethnol. Annual Rep.* 45: 441-522.
- STEUDEL, E. G. 1841. *Nomenclator Botanicus*, Volume 1. Cotta, Stuttgart and Tübingen.
- STEYERMARK, J. A. 1963. *Flora of Missouri*. Iowa State Univ. Press, Ames.
- TAYLOR, R. L. & G. A. MULLIGAN. 1968. *Flora of the Queen Charlotte Islands. II. Cytological Aspects of the Vascular Plants*. Queen's Printers, Ottawa.
- THUNBERG, C. P. 1784. *Flora Japonica*. Müller, Leipzig.
- TORREY, J. 1824. *A Flora of the Northern and Middle Sections of the United States*, Volume 1. Swords, New York.
- . 1859. Botany of the Boundary. In W. H. Emory (editor), *Report on the United States and Mexican Boundary Survey*, Volume 2, part 1. A. O. P. Nicholson, Washington, D.C.
- & A. GRAY. 1840. *Flora of North America*, Volume 1, part 4. Wiley and Putnam, New York.
- UPHOF, J. C. TH. 1968. *Dictionary of Economic Plants*, 2nd edition. Cramer, Lehre.
- VOSS, E. G. ET AL. (editors). 1983. *International Code of Botanical Nomenclature*. *Regnum Veg.* 111. Bohn, Scheltema, and Holkema, Utrecht and Antwerpen; W. Junk Publ., The Hague and Boston.
- WANSCHER, J. H. 1932. Studies on the chromosome numbers of the Umbelliferae. *Bot. Tidsskr.* 42: 49-58.
- WEDDELL, H. A. 1861. *Chloris Andina*, Volume 2, part 14. Paris.
- WILLIS, J. C. 1922. *Age and Area*. Cambridge Univ. Press, Cambridge.
- WOLFE, J. A. 1969. Neogene floristic and vegetational history of the Pacific Northwest. *Madroño* 20: 83-110.
- WOOD, C. E., JR. 1972. Morphology and phytogeography: the classical approach to the study of disjunctions. *Ann. Missouri Bot. Gard.* 59: 107-124.
- WOOLF, C. M. 1968. *Principles of Biometry*. Van Nostrand Co., Princeton, Toronto, London, Melbourne.