

THREE NEW RECORDS OF THE GENUS *Sabatieria* Rouville, 1903 (NEMATODA, CHROMADORIDA, COMESOMATIDAE) FROM THE CANARY ISLANDS

Rodrigo Riera^{1*}, Jorge Núñez² & María del Carmen Brito²

¹ Centro de Investigaciones Medioambientales del Atlántico (CIMA SL)
Arzobispo Elías Yanes, 44, 38206 La Laguna, Tenerife, Canary Islands, Spain

² Benthos Lab, Department of Animal Biology, Faculty of Biology
University of La Laguna, 38206 La Laguna, Tenerife, Canary Islands, Spain

*corresponding author: rodrigo@cimacanarias.com

RESUMEN

Se citan, por primera vez para el archipiélago canario, tres especies de nematodos pertenecientes al género *Sabatieria*: *S. celtica* Southern, 1914, *S. longisetosa* (Kreis, 1929) y *Sabatieria* sp., recolectadas en sedimentos intermareales y submareales someros de dos playas de la isla de Tenerife. Se muestran las descripciones, figuras e información merística de cada una de las especies, así como datos autoecológicos.

Palabras clave: Nematoda, Comesomatidae, *Sabatieria*, fondos blandos, Tenerife, islas Canarias.

ABSTRACT

Three species of the genus *Sabatieria*: *S. celtica* Southern, 1914, *S. longisetosa* (Kreis, 1929) and *Sabatieria* sp., are recorded for the first time from Canary Islands, collected from intertidal and shallow subtidal soft-bottoms of the island of Tenerife. Descriptions, figures and meristic data are reported as well as autoecological data.

Key words: Nematoda, Comesomatidae, *Sabatieria*, soft-bottoms, Tenerife, Canary Islands.

INTRODUCTION

The genus *Sabatieria* Rouville, 1903 is characterized by having a cuticle ornamented with transversely punctations, lacking longitudinal rows of dots, and buccal cavity cup-shaped. Cephalic setae is larger than outer labial setae. Spicules are short. Testes (in males) and ovaries (in females) are two, opposed and outstretched. Platt [4] divided this genus in 5 groups:

- *Armata* group. Cephalic setae long ($\leq 1,7$ cephalic diameter). Subcephalic setae present. Tubular precloacal supplement.
- *Celtica* group. Amphids multispiral (3 rounds). Precloacal supplements conspicuous. Gubernacular apophysis bent.
- *Ornata* group. Posterior group of more closely spaced precloacal supplements separated from the rest.
- *Praedatrix* group. Tubular or pore-like precloacal supplements.
- *Pulchra* group. Relatively few (5-9) usually conspicuous precloacal supplements. Paired and short subcephalic setae.

This genus comprises so far 51 species:

- *Armata* group, 6 species: *S. arcuata* Wieser, 1954, *S. armata* Gerlach, 1952, *S. elongata* Jayasree & Warwick, 1977, *S. longispinosa* Lorenzen, 1972, *S. migrans* Jensen & Gerlach, 1977 and *S. supplicans* Gerlach, 1956.
- *Celtica* group, 4 species: *S. celtica* Southern, 1914, *S. furcillata* Wieser, 1954, *S. kelletti* Platt, 1983, *S. pisinna* Vitiello, 1970, and *S. strigosa* Lorenzen, 1972.
- *Ornata* group, 4 species: *S. abyssalis* (Filipjev, 1918), *S. longisetosa* (Kreis, 1929), *S. macramphis* Lorenzen, 1972 and *S. ornata* (Ditlevsen, 1918).
- *Praedatrix* group, 30 species: *S. alata* Warwick, 1973, *S. ancudiana* Wieser, 1954, *S. bitumen* Botelho, Da Silva, Morgado-Esteves & Fonsêca-Genevois, 2007, *S. conicauda* Vitiello, 1970, *S. coomansi* Chen & Vincx, 1999, *S. demani* Bresslau & Stekhoven, 1935, *S. dodecasapillata* (Kreis, 1929), *S. exilis* Botelho, Da Silva, Tosta-Sobral & Fonsêca Genevois, 2009, *S. falcifera* Wieser, 1954, *S. flecha* Pastor de Ward, 2003, *S. fidelis* Botelho, Da Silva, Tosta-Sobral & Fonsêca-Genevois, 2009, *S. granifer* Wieser, 1954, *S. heipi* Chen & Vincx, 2000, *S. intermissa* Wieser, 1954, *S. lawsi* Platt, 1983, *S. lepida* (Vitiello, 1976), *S. lucia* Muthumbi, Soetaert & Vincx, 1997, *S. lyonessa* Warwick, 1977, *S. maboyae* Gourbault & Vincx, 1990, *S. parabyssalis* Wieser, 1954, *S. paracupida* Wieser & Hopper, 1967, *S. paradoxa* Wieser & Hopper, 1967, *S. paraspiculata* Botelho, Da Silva, Morgado-Esteves & Fonsêca-Genevois, 2007, *S. praedatrix* De Man, 1907, *S. sanjosensis* Pastor de Ward, 2003, *S. spiculata* Botelho, Da Silva, Morgado-Esteves & Fonsêca-Genevois, 2007, *S. stekhoveni* Vitiello, 1970, *S. subrotundicauda* Botelho, Da Silva, Morgado-Esteves & Fonsêca-Genevois, 2007, *S. triplex* Wieser, 1954 and *S. vasicola* Vitiello, 1970.
- *Pulchra* group, 7 species: *S. breviseta* Stekhoven, 1935, *S. maboyae* Gourbault & Vincx, 1990, *S. mortensenii* (Ditlevsen, 1921), *S. propisinna* Vitiello, 1976, *S. pulchra* (Schneider, 1906) and *S. punctata* (Kreis, 1924).
- *Sabatieria aspera* Sergeeva, 1972, *S. effilata* Stekhoven, 1950, *S. possjetiva* Platonova, 1971, *S. praebosporica* Sergeeva, 1972, *S. quadripapillata* Filipjev, 1922, *S. rota* Gerlach, 1957 and *S. sarcina* (Vitiello, 1976) are considered species *inquirenda* (Platt, 1985).

During an ecological study of the soft-bottoms from two localities on the south coast of Tenerife, several specimens that belong to the genus *Sabatieria* were collected. Three species were identified, *Sabatieria celtica* Southern, 1914, *Sabatieria longisetosa* (Kreis, 1929) and *Sabatieria* sp.

MATERIAL AND METHODS

Samples were collected in the intertidal and shallow subtidal, at 3 m deep, soft-bottoms of Los Abrigos (SE Tenerife) and Los Cristianos (SW Tenerife). PVC cores of 4.5 cm of inner diameter were taken to a depth of 30 cm in the sediment. These samples were fixed with 10% formaldehyde in seawater for one day and decanted through a sieve of 63 mm mesh size, and posteriorly preserved in 70% ethanol. All specimens were mounted in glycerine gel and drawings of these were done using a camera lucida on a Leica DMLB microscope equipped with Nomarski interference contrast. All measurements are in micrometers and curve structures are measured along the arc.

Abbreviations used in the text are: a, body length divided by maximum body diameter; b, body length divided by pharyngeal length; c, body length divided by tail length; c', tail length divided by anal body diameter; cbd, corresponding body diameter; s', spicule length divided by anal body diameter; %V, position of vulva as a percentage of body length from anterior.

SYSTEMATICS

CHROMADORIDA Chitwood, 1933

CHROMADORINA Filipjev, 1928

COMESOMATIDAE Filipjev, 1918

Sabatieria Rouville, 1903

Sabatieria celtica Southern, 1914

(Figs 1 & 2; Tab. 1)

Sabatieria celtica Southern [6]: 25, fig. 8 a-d; Platt [4]: 57, figs. 6 e. 16, 18 e-g, tab. 5; Platt & Warwick [5]: 210, fig. 93; Hua & Zhang [2]: 30, fig. 4.

Sabatieria longiseta Steiner [7]: 593, fig. 22 a-d.

Meristic data and studied material.- Abrigos subtidal: june 2000, 2 males (δ 1 and δ 5) and 1 female (φ 1), october 2000, 2 females (φ 2 and φ 4), november 2000, 1 male (δ 2), april 2001, 1 female (φ 5); Cristianos subtidal: may 2000, 2 males (δ 3 and δ 4), march 2001, 1 female (φ 3).

Description.-

Male: Body slender, tapering towards both ends. Head slightly round and not set off. Cuticle annulated and ornamented with transverse rows of dots: fewer rows or large dots laterally. Amphids are 55% of the corresponding body diameter in width, multispiral, 2.5 rounds, located at 7 μ m from the anterior end. Buccal cavity conical and small. 6 inner labial setae inconspicuous. 6 outer labial setae 3 μ m long, sometimes not discernible. 4 cephalic setae 1.8-2 cephalic diameters long, situated at the anterior half of the head. Subcephalic setae 0.7-0.9 cephalic diameters long, located at 18 μ m from the anterior body end. Pharynx slender and cylindrical. Ventral gland and nerve ring not seen.

The reproductive system is diochic with two extended testis, difficult to discern. Spicules 1.3 anal diameters long, paired and arcuate, with a short central projection at the proximal end. Gubernaculum developed with a straight and narrow dorsal apophysis. 18-21 small tubular precloacal supplements, which can be easily overlooked. Tail 2.7-3.6 anal di-

ameters long, slender and cylindrical with a slightly expanded distal end. 2 caudal setae 17 μm long. Spinneret developed.

Female: Total body length slightly larger than in males (2.1-2.9 mm), with a longer tail (4.7-4.8 anal diameters). Cephalic setae less developed (20-21 μm). Outer labial and subcephalic setae lacking. The reproductive system is didelphic, with two outstretched ovaries, difficult to discern. Vulva located at 49.2-50.1% of the total body length.

Discussion.- This species presents important intraspecific variations, especially in cephalic and subcephalic setae length. *Sabatieria celtica* Southern, 1914 is characterized by having spicules proximally expanded, although Chen & Vincx [1] noticed some individuals lacking this character. Gubernacular apophysis is straight in canarian specimens, being curved gubernacular apophyses a typical character of the *celtica* group (Platt [4]).

Ecology.- In the subtidal of Los Abrigos was recorded in fine sands ($Q_{50} = 0.24$), with a very good selection ($S_0 = 0.73$). The organic matter content was 0.51% and 4.61% of carbonates percentage. In the subtidal of Los Cristianos was collected in fine sands ($Q_{50} = 0.16$), with a very good selection ($S_0 = 0.58$). The organic matter content was 0.73% and 24.96% of carbonates percentage.

Distribution.- Cosmopolitan (Platt [4]).

Sabatieria longisetosa (Kreis, 1929)

(Fig. 3; Tab. 2)

Sabatieria longisetosa.- Platt [4]: 61, fig. 17, tab. 16; Platt & Warwick [5]: 212, fig. 94 f-g.
Parasabatieria longisetosa Kreis [3]: 89, fig. 40 a-d.

Meristic data and studied material.- Abrigos subtidal: may 2000, 4 males ($\delta 1, \delta 2, \delta 3$ and $\delta 4$); Cristianos subtidal: february 2001, 1 male ($\delta 5$).

Description.-

Male: Body slender, attenuating on both anterior and posterior ends. Head round and slightly set off. Cuticle annulated and ornamented with homogeneous punctations, lateral differentiation not seen. Amphids 93% of the corresponding body diameter in width, multisprial, 3 rounds, located at 6 μm from the anterior end. Buccal cavity conical and small, without noticeable teeth. Inner labial setae lacking. 6 outer labial setae 2 μm long and 4 cephalic setae 1.4 cephalic diameters long, situated in the anterior half of the head. Subcephalic setae 9 μm long, located at 35 μm from the anterior body end. Short scattered somatic setae fairly numerous pharyngeal region but scarce in the rest of the body. Pharynx slender and cylindrical.

The reproductive system is diorchic with two opposed testes. Spicules 1.6 anal diameters long, paired and arcuate, without capitulum. Gubernaculum 0.6 anal diameters long, with a distally curve aphopysis. 14 small tubular precloacal supplements, divided in two groups (5 + 9) broaden the posteriomost. Precloacal setae 3 μm long, located at 3 μm from the cloaca. Tail 3.7-4.5 anal diameters long, cylindrical and slender with slightly swollen tail tip. 2 caudal setae 7 μm long. Spinneret developed.

Females not found.

Discussion.- This species belongs to *ornata* group, characterized by having posteriormost precloacal supplements more closely situated than the remaining ones. *Sabatieria longiseta* (Kreis, 1929) is defined by having 14 precloacal supplements and its tail length (4.1 anal diameters). Canarian individuals are larger (1.6-2.1 mm) compared to those collected in other atlantic regions (1.3-1.6 mm) and have spicules less developed in width (Platt [4]).

Ecology.- In the intertidal of Los Abrigos this species was recorded in medium sands ($Q_{50} = 0.36$), with a very good selection ($S_0 = 0.86$). The organic matter content was 1.68% and 3.8% of carbonates percentage. In the subtidal of Los Cristianos was collected in fine sands ($Q_{50} = 0.18$), with a very good selection ($S_0 = 0.73$). The organic matter content was 0.33% and 23.08% of carbonates percentage.

Distribution.- East Atlantic (Platt [4]).

Sabatieria sp.

(Fig. 4; Tab. 3)

Meristic data and studied material.- Abrigos intertidal: may 2000, 1 female ($\varphi 1$); Abrigos subtidal: december 2000, 2 females ($\varphi 2$ and $\varphi 3$).

Description.-

Males not found.

Female: Body slender, tapering towards both ends. Head round and slightly set off. Cuticle ornamented with homogeneous punctations, lateral differentiation difficult to discern. Amphids 43% of the cbd in diameter, multispiral (2.5 rounds), located at 6 μm from the anterior body end. Buccal cavity conical and small, without developed teeth. Inner labial setae lacking. Outer labial setae 3 μm long and 4 cephalic setae 1.5 cephalic diameters long, situated in the median part of the head. Subcephalic setae absent. Pharynx cylindrical and slender.

Reproductive system not discernable. Vulva located at 46.5-60% of the total body length. Tail 4.6 anal diameters long, cylindrical and filiform in its posterior part, with swollen tail tip. 2 caudal setae 10 μm long. Spinneret developed.

Discussion.- *Sabatieria* sp. can be differentiated from the remaining species of the genus in its tail length (4.5-5.4 anal diameters), amphid size (43% cbd in diameter) and outer labial setae (1.5 anal diameters long). This species has been determined to genus level due to the absence of males.

Ecology.- In the intertidal of Los Abrigos this species was colected in medium sands ($Q_{50} = 0.34-0.42$), with a very good selection ($S_0 = 0.83-0.97$). The organic matter content was 1.33% and carbonates percentage ranged from 5.30% to 5.47%. In the subtidal of the same locality was recorded in medium sands ($Q_{50} = 0.28$), with a very good selection ($S_0 = 0.75$). The organic matter content was 0.78% and carbonates percentage was 5.47%.

ACNOWKLEDGMENTS

Authors are grateful to Dr. M. Vincx and Tom Gheskire (University of Ghent, Belgium) for providing laboratory and bibliography facilities. We also acknowledge Dr. Catalina Pastor de Ward (Centro Nacional Patagónico, Argentina) for her insightful comments and suggestions. To Alessandra Prates Botelho (University of Recife, Brazil) for encouragement and interchange of ideas relating to the genus *Sabatieria*.

REFERENCES

- [1] CHEN, G. & M. VINCX. 1999. Nematodes from the Strait of Magellan and the Beagle Channel (Chile): the genus *Sabatieria* (Comesomatidae: Nematoda) with the description of *Sabatieria coomansi* n. sp. *Hydrobiologia*, 405: 95-116.
- [2] HUA, E. & Z. ZHANG, 2007. Four newly recorded free-living marine nematodes (Comesomatidae) from the East China Sea. *Journal of Ocean University of China*, 6(1): 26-32.
- [3] KREIS, H. 1929. Freilebende marine Nematoden von der Nordwestküste Frankreichs (Trébeurden Côtes du Nord). *Capita Zoology*, 7(2): 1-98.
- [4] PLATT, H.M. 1985. The free-living marine nematode genus *Sabatieria* (Nematoda, Comesomatidae). Taxonomic revision and pictorial keys. *Zoological Journal of the Linnean Society*, 82: 27-78.
- [5] PLATT, H.M. & R.M. WARWICK. 1983. *Free-living marine nematodes. Part I: British Enoplids*. Kermarck, D.M. & R.S. Barnes (eds.). Cambridge University Press. London, 307 pp.
- [6] SOUTHERN, R. 1914. Clare Island survey. Archiannelida and Polychaeta. *Proceedings of the Royal Academy*, 31-160
- [7] STEINER, G. 1916. Freilebende Nematoden aus der Barentsee. *Zoology Jb Systematics*, 39: 511-676.

TABLES AND FIGURES

	♂1	♂2	♂3	♂4	♂5	♀1	♀2	♀3	♀4	♀5
Total body length	1887	2571.4	2600	1914.3	2389.7	2959	2142.9	2789.5	2544.9	2613.5
a	31.4	38.3	39.5	38.3	36.3	52.8	42.9	50.9	52.2	53.4
b	10	11.3	16.5	14.9	16	13.8	8.8	11.8	11.4	11.1
c	10.4	12.9	17.8	14.1	13.4	13.8	10.7	13.2	12.4	12.2
Cephalic diameter	22.9	25.7	22.4	23	21.9	15.7	17.1	16.5	15.4	16.4
Inner labial setae	-	-	-	-	-	-	-	-	-	-
Outer labial setae	3.1	2.9	3	2.9	3.1	-	-	-	-	-
Cephalic setae	40	42.3	39.7	39.9	38.7	20	21.4	19.8	20.1	20.8
Subcephalic setae	17.9	18.7	16.5	15.9	15.7	-	-	-	-	-
Buccal cavity diameter	8.6	8.6	8.6	8.6	8.4	7.1	7.1	6.9	7	7.2
Amphid diameter	11.4	-	7.1	11.4	10.1	10	11.4	10.1	10	9.8
Amphid height	15.7	-	10	12.9	11.2	10.5	14.3	10.6	10.4	10.3
Amphid from anterior	7	-	12.9	8.6	9.3	6.8	6	6.5	6.4	6.4
Pharynx length	189.2	228.6	157.1	128.6	149.8	214.3	242.9	235.6	222.5	235.4
Pharynx cbd	60.7	46.4	39.3	35.7	48.7	39.3	42.9	42.8	40.1	41.5
Maximum body diameter	60.4	67.1	65.9	50	65.9	56	50	54.8	48.7	48.9
Vulva from anterior	% V					1457.1	nd	1373.9	1267.4	1309.4
Spicule length	64.3	69.3	60.7	67.8	62.9	49.2	nd	49.3	49.8	50.1
Gubernaculum length	34.5	35.7	32.1	35.7	34.8					
s'	1.3	1.3	1.3	1.4	1					
Tail length	182.1	200	146.4	135.7	178.3	214.3	200	210.1	205.9	213.4
Anal body diameter	50	71.4	45.7	50	61.9	45.7	42.9	42.3	43.4	44.0
c'	3.6	2.8	3.2	2.7	2.9	4.7	4.7	4.9	4.7	4.8
Spicule length/Tail length	0.4	0.3	0.4	0.4	0.5	0.4				

Table 1.- Measurements of *Sabatieria celtica* in μm ; nd: not discernible.

	♂1	♂2	♂3	♂4	♂5	
Total body length	1628.6	1742.9	1885.7	2085.7	1942.9	
a	32.6	32.5	35.2	42.3	38.9	Total body length a
b	7.2	7.4	8.8	4.7	8.6	
c	9.7	10.4	12.9	13.3	12.7	
Cephalic diameter	15.7	20.3	15.7	15.7	17.1	Cephalic diameter
Inner labial setae	1.8	2	2	3	2	Inner labial setae
Outer labial setae	22.9	22.9	24.3	24.3	22.9	Outer labial setae
Cephalic setae	-	-	-	-	-	Cephalic setae
Subcephalic setae	-	-	-	-	-	Subcephalic setae
Buccal cavity diameter	7.5	5.7	5.7	7.1	7.1	Buccal cavity diameter
Amphid diameter	18.6	15.7	17.1	12.9	14.3	Amphid diameter
Amphid height	21.4	17.1	24.3	18.6	17.1	Amphid height
Amphid from anterior	8.6	10	10	8.6	11.4	Amphid from anterior
Pharynx length	225	235.7	214.3	442.9	225	Pharynx length
Pharynx cbd	39.3	42.9	42.9	32.1	39.3	Pharynx cbd
Maximum body diameter	50	53.6	53.6	49.3	50	Maximum body diameter
Spicule length	54	59	53	78.4	74.6	Vulva from anterior % V
Gubernaculum length	22.9	18.6	27.1	24.3	27.1	Tail length Anal body diameter c'
s'	1.4	1.6	1.3	2.1	1.6	
Tail length	167.9	167.9	146.4	157.1	153.6	42.9
Anal body diameter	38	37	39.3	35.7	46.4	42.9
Spicule length/Tail length	4.4	4.5	3.7	4.4	3.3	5.4
	0.3	0.4	0.4	0.6	0.6	4.5

Table 2.- Measurements of *Sabatieria longisetosa* in µm.

	♀1	♀2	♀3	♀4	♀5	
Total body length	1500	2457.1	2200	25.5	24.6	
a	28					
b	6.6	7.5	8.4			
c	7.6	9.1	11			
Cephalic diameter	13.0	20.3	22.9			
Inner labial setae	-	-	-			
Outer labial setae	2.8	3	3			
Cephalic setae	19.0	22.9	27.1			
Subcephalic setae	9	10	9			
Buccal cavity diameter	5.7	8.6	8.6			
Amphid diameter	7.1	14.3	15.7			
Amphid height	8.6	14.3	14.3			
Amphid from anterior	6	4	5.7			
Pharynx length	228.6	328.6	260.7			
Pharynx cbd	46.4	82.1	78.6			
Maximum body diameter	53.6	96.4	89.3			
Vulva from anterior % V	900	1142.9	nd			
Tail length	60	46.5	nd			
Anal body diameter c'	196.4	271.4	200			
Spicule length/Tail length	4.6	5.4	4.5			

Table 3.- Measurements of *Sabatieria spinifera* in µm. nd: not discernible.




Figure 1.- *Sabatieria celtica*. Male. **A.** Anterior end. **B.** Pharynx, oesophageal bulb and cardias. Scale A = 15 µm, B = 30 µm.




Figure 2.- *Sabatieria celtica*. Male **A.** Spicule and gubernaculum. **B.** Posterior end. Scale = 20 µm.




Figure 3.- *Sabatieria longisetosa*. Male. **A.** Anterior end. **B.** Posterior end. Scale A = 20 μm , B = 26 μm .




Figure 4.- *Sabatieria* sp. Female. **A.** Anterior end. **B.** Posterior end. Scale A = 20 μm , B = 26 μm .