

and 6–10 mm. broad, the beak about 3 mm. long and mostly straight, the stipe 4–6 mm. long.

Herba 5–8 dm. alta, omnino dense appresso-tomentosa; caule crassiusculo, leviter striato, late ramoso, ramis alternatis, ascendibus; foliis 1-foliolatis, dilute viridibus, subtus paulo lutescentibus, plerumque cordatis (infimis raro subreniformibus, supremis interdum ovatis), coriaceis, valde nervatis, basi caulem subamplectentibus, apice late rotundatis, margine integris et interdum paulo revolutis, 2–8 cm. longis, 1.5–7 cm. latis; racemis terminalibus, 3–20 cm. longis; floribus pedicellatis, pedicellis 1–5 mm. longis, bracteolatis; calyce campanulato, ca. 5 mm. longo, extus dense pubescente, intus glabro, laciinis tubum subaequantibus, labio superiore lato, integro vel paululo emarginato, laciinis inferioribus late lanceolatis; corolla flava, vexillo orbiculari et emarginato (11 mm. longo, 10 mm. lato), alis carinaque oblongis (ca. 13 mm. longis et 5 mm. latis); legumine extus dense pubescente, intus molliter pubescente, subgloboso vel ellipsoideo, 8–15 mm. longo et 6–10 mm. lato, rostro 3 mm. longo et plerumque recto, 4–6 mm. stipitato.

Distribution: on sandy soil, mostly in open, pine woods, occasionally in denser, mixed woods; Wayne County, Georgia.

The type specimen (sandy soil in open, pine woods, July 4, 1943, *Duncan 5693*) is deposited at the Gray Herbarium, Cambridge, Massachusetts, U. S. A. Isotype specimens are being distributed in the *Plantae Exsiccatae Grayanae*.

The author wishes to acknowledge the valuable aid afforded by "Monograph of the Genus *Baptisia*", Annals Missouri Botanical Garden 27: 119–244, April, 1940, by M. M. Larisey; and for specimens loaned by the Missouri Botanical Garden. Thanks are gratefully given to Dr. Lyman B. Smith of the Gray Herbarium for assistance in the preparation of the manuscript and in distribution of specimens.

UNIVERSITY OF GEORGIA and
UNITED STATES PUBLIC HEALTH SERVICE.

ASTER **acadiensis** nom. nov. *A. lateriflorus* var. *tenuipes* Wiegand, RHODORA 30: 174, 1928. *A. tenuipes* Shinners, RHODORA 45: 346, 1943; non *A. tenuipes* Makino, Bot. Mag. Tokyo 12: 64, 1898.

I am indebted to Miss Marjorie W. Stone and to Mr. C. A. Weatherby for bringing the above error to my attention.—L. H. SHINNERS, Madison, Wisconsin.