

Two Lectotypes in *Charpentiera* (Amaranthaceae)
Hawaiian Plant Studies 51

Harold St. John

B. P. Bishop Museum, Honolulu, Hawaii, 96818, Box
6037, U. S. A.

The two following formae in *Charpentiera* were described by Suessenguth, each based on several collections, but without any designation of holotypes. Below the needed lectotypes are selected. This genus has recently been monographed by Sohmer (1972), but he failed to mention or dispose of these two formae, though he listed two of the syntypic specimens.

Charpentiera obovata Gaud., forma *parvifolia*
Suesseng., Fedde Repert. 44: 38, 1938.

Lectotype: Maui, Auwahi, O. Degener 8,826
(BISH).

Sohmer (1972: 299) lists Degener 8,826 as *C. obovata* Gaud., but does not cite the three other Degener collections.

Charpentiera obovata Gaud., forma *grandifolia*
Suesseng., Fedde Repert; 44: 38, 1938.

Lectotype: Hawaii Island, Kohala, O. Degener 8,824 (GH).

Of the three syntypes listed by Suessenguth, Sohmer (1972: 302) lists both the above chosen lectotype, and Degener 8,820, as *C. tomentosa* Sohmer, var. *tomentosa*, but does not mention the third one, Degener 8,830.

Literature Cited

- Sohmer, S. H., 1972. Revision of the genus
Charpentiera (Amaranthaceae). *Brittonia* 24:
283-312, figs. 1-14.
- Suessenguth, K., 1938. *Amarantaceen-Studien*. 1.
Amarantaceae aus Amerika, Asien, Australien.
Fedde Repert. Spec. Nov. 44: 36-48.