

STREPTOPOGON JUAREZII N. SP.
and
TREMATODON NORRISII N. SP.*

A. J. Sharp
Department of Botany, University of Tennessee
Knoxville, TN 37916

Streptopogon juarezii sp. nov.

Distinguitur ex ovatis, levibus aristatis foliis quae
leves margines habent et quae gemmis carent et quae cellulas
marginis quadratis inconspicuas habent.

This new species is distinguished by having elliptically
ovate, smooth, entire, aristate leaves without gemmae. The
sporophyte with an oblong tubular immersed capsule with a
short (± 1 mm) seta and a peristome of a high, ~~hyaline~~
basal membrane and 16 slender, perforate, papillose teeth.
Autoicous.

TYPE: Sobre un tronco húmedo. 21 km al oeste de
Xalapa, Veracruz, México. 1700 m elevación. Coll. L. Gil
Juarez G., No. 627, 20 VII 1976 (type, TENN; isotype, INIREB).

Trematodon norrisii sp. nov.

Distinguitur ex parvitate, ex foliis anguste lanceolatis,
ex costa percurrente, et ex perfragilibut angustis dentibus
peristomatos.

Distinguished from other small species of Trematodon by
the narrowly lanceolate (1-2 mm) leaves with percurrent
costae; capsules 2 mm in length, abruptly narrowed to slender
hypophyses, and with thin, ~~hyaline~~ walls; teeth slender,
very fragile, 200-250 μ ; spores 25-30 μ , prominently
papillose.

TYPE: Moist, diffusely lit soil bank in dense tropical
sub-deciduous forest along road to Puerto Vallarta about 4
miles north of San Juan Caxtle, Municipio Compostela,
Nayarit, México. Coll. D. H. Norris & D. J. Taranto, No. 13641,
24 July 1970 (type, TENN; isotype, HSC).

Acknowledgment. I am greatly indebted to Dr. C. P. Craig
of the University of Tennessee Classics Department who gave me
generous assistance with the preparation of the Latin
diagnoses.

*Contribution from the Botanical Laboratory, The University of Tennessee, N. Ser 563, and from the Hattori Botanical Laboratory, Nichinan, Miyazaki, Japan.