

THE  
BOTANICAL GAZETTE

*SEPTEMBER 1919*

PHYTOGEOGRAPHY OF THE EASTERN MOUNTAIN-  
FRONT IN COLORADO

I. PHYSICAL GEOGRAPHY AND DISTRIBUTION  
OF VEGETATION

CONTRIBUTIONS FROM THE HULL BOTANICAL LABORATORY 251

ARTHUR G. VESTAL

(WITH SEVENTEEN FIGURES)

**Introduction**

The plant geography of a region is the effect of the working of present and former environmental influences upon the floras and vegetation-complexes which exist and have existed within the region and in the regions adjoining. The region of present study, lying as it does in the transition belt between two great geographic divisions of North America, the Great Plains, or western part of the prairie region, and the Rocky Mountains, has some of the characters of both; others of its physical and vegetational features are transitional, intermediate; and it has certain peculiarities, differing thus from the regions on either side. Since climatic variation, differences of soil and of topography, and multiformity of vegetation-types are considerable, the plant-covering of the area is a complex of many diverse types. Descriptive accounts of the plant associations of plains and foothills have already been published (17, 18), so that the present article may deal more particularly with geographic description and geographic relations.




FIG. 1.—Map of southern Rocky Mountains, except westernmost ranges; mountain areas shaded; names of areas indicated by numbers are: 1, Laramie Mountains; 2, Medicine Bow Range; 3, low mountain area connecting Laramie and Front ranges; 4, foothills of Poudre River area; 5, Front Range; 6, Rampart Range; 7, Pike's Peak highland; 8, Park Range; 9, Saguache Range; 10, Upper Arkansas Valley (between 9 and 8); 11, low mountains; 12, Wet Mountain Valley; 13, Sangre de Cristo Range; 14, Wet and Greenhorn mountains; 15, Huerfano Park; 16, southern sedimentary plateau; 17, Culebra Range; 18, Spanish Peaks highland; 19, Raton mesas.

While in general the plains and mountains contrast rather sharply at their junction, this is not always true; the mountain-front is a transition zone in places a number of miles broad rather than a line. It is not determined alone by altitude, by topography, by character of the bedrock, or by climate; it is the resultant of all of these. For the sake of clearness the foothills may be described as the drier and less elevated (about 5800–8000 ft.) part of the mountain plateau, with vegetation composed of grassland, scattered rock pines, and a few other trees (*foothill zone*, RAMALEY 8). Except in the southern “sedimentary plateau” (fig. 1), perhaps rather to be considered part of the mountain-front area, the foothills may be said to comprise the granitic hills of the mountain-mass proper; while to the mountain-front zone may be assigned the upturned sedimentary hogbacks and longitudinal valleys, sedimentary outcrops, buttes and broken plateaus, and the mesas and upper parts of the débris-covered slope to the plains. The vegetation is of the greatest variety. The plains proper may be said to commence where the mixed soil and

vegetation of the detrital outwash from the hills is succeeded by the fine soil and mostly short-grass vegetation of the shale beds covering most of the Great Plains surface.

*Plan of presentation.*—The writer has been much influenced by the work of DAVIS (1) on the geography of the Colorado Front Range, a regional presentation and particularly relevant in this study, since the area considered is so nearly the same. DAVIS' systematic treatment avoids repeating descriptions of frequently encountered land-forms by recognizing their common features and giving each a brief characterization and a name, thus identifying them when mentioned later. Minor differences of detail are not considered in the condensed treatment thereby made possible. In a regional study, in which numerous elements form an intricate complex, this omission of detail is essential. As the physical geographer refers land-forms to types (mental counterparts of physical realities), so in a regional study of plant geography one may refer forms of vegetation to types which are the same over considerable areas. This is a common practice in ecological classification, but many studies of limited areas of vegetation have characterized the plant communities without regard to geographic orientation. If possible, local representations or variants of widespread associations should be recognized as such. The characterization of the relatively few widespread and important vegetation-types makes it possible to systematize plant geography. This systematic treatment emphasizes the common features, the resemblances of similar plant communities, but the differences, when worthy of note, can always be stated in addition. The section of this study which is here published is the systematic part, which establishes the types of topography, soil, climate, and vegetation as developed in the region or in parts of it. It will be followed by a regional section, which describes the physical and vegetational features "in their actual spatial relations," to use the words of DAVIS, and by parts dealing with general geographic and developmental relations of the vegetation.

### Physical features

The area studied is the eastern front of the Rocky Mountains in Colorado, of which the most characteristic part is the Front

Range. This has been studied by many geographers, more recently by DAVIS. The Front Range has been described by him as a sub-maturely dissected upland of crystalline rocks, elevated above the plains to the east by a long north-south monoclinical fold. The tops of most of the hills form the remains of a peneplaned surface, the result of the erosion following the uplift, with complete removal of the sedimentary layers from the raised area on the west. A few




FIG. 2.—DAVIS' block diagram of Front Range (reproduced with author's permission from 1): at right is condition following first uplift with monoclinical fold; next part shows peneplaned upland with monadnocks and cuestas (hogbacks); third shows entire region after second uplift; last block on left shows present condition, with glacier-carved range-crest, gently sloping, dissected, crystalline upland, of which lower and eastern part forms foothills, and mountain-front, with sloping crags, cuestas, and longitudinal valleys; outside may be seen debris-covered terraces and broad valleys of streams running out into plains.

monadnocks surmount the general level. The present eastward inclination of the old peneplain and its dissected character in the crystalline area, and the removal of sedimentary strata of the plains to a depth far below the foothills, are the effects of a second uplift, an uparching of the whole region, and of the subsequent cycle of erosion. Near the base of the original fold the sedimentary strata are sharply upturned against the outer granitic slopes, the ends of the resistant strata forming ridges and sloping crags (fig. 2).

The outer slope to the plains has been described by JOHNSON (5) as a *débris-apron* or composite of alluvial fans, of which the profile is that of a stream-grade, rapidly flattening into the very slight and uniform incline of the Great Plains. The graded surface is covered by unsorted rock-waste from the hills, which thins out and becomes finer in texture toward the east; it is absent from most of the surface of the plains, which is of fine grained residual soil. This grade is that of the High Plains;<sup>1</sup> the streams have very generally cut below it, especially near the mountains. The Platte and Arkansas rivers, the trunk-streams, have cut very broad valleys in the soft shales of the plains. The north-south valleys of their tributaries which parallel the mountain-front are bordered on the east by escarpments of considerable height and are notable geographic features. This recent downcutting, where working in soft shales just outside the foothills, leaves many terraces, remnants of the older and higher stream-grade levels; their covering of rock-waste preserves their flat tops. They are generally known as "mesas"; although not true mesas, the term is convenient.<sup>2</sup> Where the upper sedimentary beds consist of sandstone or limestone, extensive plateau areas with deep canyons, buttes which may be numerous or scattered, or simple escarpments may be encountered. In a few places igneous intrusions are seen as dikes or as basaltic layers capping large mesas (true mesas in this case). From these features the mountain-front zone derives its varied character; the mountain upland on the west, and the plains extending far to the east, are of less irregular structure.

Arrangement of the component ranges and smaller ridges *en echelon* is a notable feature of the easternmost line of mountains. Ranges which are in general north and south of each other are themselves oriented with the northern end a little to the west. MARVINE writes (7, p. 132):

<sup>1</sup> The distribution of the remnants of the High Plains may be seen in a map by JOHNSON in the article mentioned.

<sup>2</sup> A true mesa is a tableland capped by a more resistant stratum which keeps the top flat by retarding erosion except on the sides. The *débris*-covered terraces flanking the mountains are like a true mesa in that the rock-waste layer acts as a more resistant cap.

In traveling from the north along the zone of hogbacks lying at the base of the mountains southward, the traveler finds the mountain-slope directly west of him falling lower and lower until it becomes an insignificant ridge, and finally dies away in the plains.

Passing around the southern end of the diminishing ridge the main mountain-slope is found lying several miles to the west, and separated from the ridge by a baylike valley extending northward behind it. . . . The ridges are uplifted or anticlinal folds, the valleys depressed or synclinal folds, both dying away southward into the flatness of the plains.

The minor embayments due to echelon arrangement may be made out only in a large scale map, but the major embayments at the south end of the Rampart Range, the Pike's Peak highland, and the Greenhorn Mountains can easily be seen in fig. 1.

A more detailed view of the typical land-forms and vegetation-forms encountered in passing from mountains to plains traverses the several north-south zones in the following order: first the granitic foothills; then the transition zone of the mountain-front, with its upturned ridges, its mesas and graded slopes, and in places its plateau areas, buttes, and escarpments; and lastly the plains themselves.

#### GRANITIC FOOTHILLS

The mountain plateau is in most places submaturely dissected, the original upland level being represented only by the rounded tops of the hills (fig. 3). Slopes and summits are thinly covered with rock-waste. Occasional resistant dikes and ledges give craggy exposures of massive rock, not covered by any soil or débris. Below these, or on the sides of steeper ravines, are talus slopes of variously sized rocks, or slides of "granite-gravel."<sup>3</sup> Table I is a synopsis of topographic areas of the foothills arranged as habitats, and, correlated with these, the characteristic vegetation-types. Edaphic conditions largely determined by topography (local position in relation to surroundings, direction, amount of slope, and soil texture) have been discussed in the account of foothills vegetation (18).

This two-column form of presentation is adopted as being concise, as emphasizing relations between physiographic and ontographic features (the environment and the envired), and as permitting a more comprehensive view of the whole complex and its

<sup>3</sup> Decomposed granite in small angular fragments.

parts than can be obtained by the linear arrangement. Geographers will note that topographic *areas* rather than *land-forms* are used as the units of area of physical conditions (habitats), since land-forms, such as mesas and ravines, may include several topographic areas presenting quite diverse environmental conditions. Moreover, a single topographic area, even if physically uniform, may allow the growth within it of several more or less distinct vegetation-types.



FIG. 3.—Maturely dissected foothills near Boulder Creek: pine-sprinkled, rather than forested, surface mostly covered with dry grassland.

A brief statement concerning mountain parks may be made. These are small plains or flat valleys shut in on all sides by hills. They are not well developed in the foothills as compared with the montane zone. They are mostly formed where one of the principal eastward flowing streams is joined by tributaries from valleys opening into the park. There is a single outlet. Many of the montane parks in the Front Range contain the terminal moraines of former valley glaciers from above, and their topography is in large measure the work of ice. The slight gradient causes many

TABLE I

TOPOGRAPHIC AREAS (HABITATS) AND ASSOCIATED VEGETATION-TYPES  
IN GRANITIC FOOTHILLS COMPLEX

## TOPOGRAPHIC AREAS

The geographic mean is that presented by rather exposed and xerophytic sloping surfaces, thinly covered with rock-waste of mixed texture, rather gravelly and with surface rocks. Local departures from the general condition are as follows:

1. Exposed rock surfaces (boulders and rock-walls)
2. Rock-crevices
3. Rock-strewn detritus slopes
4. Rock-talus
5. Compacted granite-gravel floors and side-slopes
6. Loose granite-gravel floors, washes, and talus (gravel-slides)
7. Mixed-soil floors and detritus-slopes (fine soil with imbedded and superficial rock-fragments of various sizes)
8. Fine-soil floors and detritus-slopes (infrequent)
9. Less xerophytic side-slopes (mostly north-facing, mostly of considerable gradient, and best developed in valleys)
10. Narrow mesophytic ravines (best developed as small side-canyons, especially on the south side of eastward flowing main streams)
11. Stream-sides in shaded ravines
12. Stream-sides in open canyon bottoms

## VEGETATION-TYPES

The general ground-cover is mixed foothills grassland and primitive grassland, largely of grasses and herbs of the plains, with admixture of Rocky Mountain herbs, not all xerophytic. Scattered rock pines and plants of the mixed shrub association, singly or in clumps, dot the surface. In special habitats occur:

1. Xerophytic lichen association
2. *Selaginella*, shrubs of *Jamesia* and *Ribes*, rock pine
3. Mixed grassland, and mixed consocieties of primitive grassland, with higher proportion of woody plants (rock pines, mixed shrub, *Ceanothus*, *Arctostaphylos*)
4. *Artemisia frigida*-*Koeleria* consocieties of primitive grassland (18)
5. Compacted granite-gravel consocieties of primitive grassland (18) with rosette plants; *Arctostaphylos*
6. Primitive grassland, with *Geranium-Chrysopsis* consocieties, mat (rosette) consocieties of gravel-slides, etc.
7. Foothills mixed grassland, with addition of other components, *Ceanothus*, sumac, pine, etc.
8. Foothills mixed grassland, of a form approaching plains short-grass
9. Mixture of mixed shrub, rather less xerophytic mixed grassland, and pine associations, with representatives of canyon forest and scattered trees of *Pseudotsuga*
10. Mesophytic representations of mixed shrub, *Pseudotsuga*, aspen, *Symphoricarpos*, canyon forest, and mesophytic grassland associations. Mosses, *Saxifraga*, etc., in wet rock-crevices
11. *Betula*, *Alnus*, *Corylus*, and *Acer glabrum* of the canyon forest; shrubs; moist-soil herbs, as *Heracleum*, *Rumex*, etc.
12. *Populus angustifolia*, willows, etc.



meanders and oxbows in the streams, and there are in some parks small lakes in morainal depressions. The stream-sides are frequently boggy, with meadows adjoining. The parks are mostly treeless, or nearly so, and show no signs of former or impending forestation. The exposed dry flats are covered with dry grassland, its composition depending on altitude and geographic position chiefly. Differences in soil texture cause local variation of the grassland, but this is less marked and less minutely local than on the hill slopes. Certain lower areas are occupied by meadow and sedge communities, and the rolling surfaces of moraines (in montane parks) are variable in soil texture, soil moisture, and in the composition of their grassland cover; but the greater part of park floors is well drained, flat, and quite uniformly covered with dry grassland. This vegetation, in any one park, forms what might be called a crystallization of the grassland of the neighboring hills, whether in foothills or montane zone, in view of the comparative uniformity of the grassland of the flats as contrasted with that of the diversified slopes of hill topography. The lower parks have a grassland cover very like that of coarse soil in the mountain-front area or in the plains (see description of Estes Park in the regional section). The higher parks have fewer plants of the plains and more of the mountains. There is a floristic and vegetational gradation from plains grassland through the lower parks to montane grassland as seen in the higher levels. The parks thus show a steplike series of floristic and ecological changes with altitude. RAMALEY (10, 11) for some years has studied park vegetation, especially in Boulder Park at Tolland, Colorado, on South Boulder Creek.


#### TRANSITION AREA OR MOUNTAIN-FRONT ZONE

The sedimentary rocks, lying upon the granite, are upturned at the monoclinial fold, and are seen in a horizontal series of exposures of strata, the lower and older members abutting on the granitic foothills to the west, the upper formations outcropping in order toward the east. Since the tilting at the mountain-front is for considerable distances greater than  $45^{\circ}$  (locally reaching  $90^{\circ}$  and even more, resulting in overturns), the lower formations have narrower zones of outcrop than the upper strata, which dip so slightly as to cover areas many miles wide in the plains. The

narrow zone of older and lower strata contains alternating resistant and soft members, giving rise to the hogback ridges and intervening valleys already mentioned, while the newer rocks are mostly soft shales and sandstones, giving a flat or rolling topography over the surface of the plains, with occasional escarpments at the edges of stream-valleys. Both angle of dip and hardness of rock, therefore, contribute to a differentiation, in the sedimentary area outside the foothills, of a relatively narrow ridge-valley mountain-front zone from the very broad and mostly flat plains region.

Just outside the ridge-and-valley zone is the graded slope to the plains, covered with rock-débris and dissected into terraces or mesas of varying level. In places along the mountain-front the ridge-and-valley topography is absent or poorly developed, either because the troughs are not yet carved beneath the slope from the granitic hills, or because the ridges are already planed (locally) to a graded floor. The terraces are also missing from certain parts of the mountain-front. The topographic complexes of the ridge country and of the mesa country may now be described separately.

THE HOGBACK RIDGES (CUESTAS) AND INTERVENING TROUGHS (figs. 4, 5).—Two of the numerous sedimentary strata overlying the crystalline rocks are so resistant as to form ridges over great lengths of the mountain-front. These two strata are of such conspicuous geographic importance that they merit distinctive names and since many persons know them by their geological names, these will be used here in a geographic capacity. The *Fountain* sandstone, which in most places lies directly upon the granites, is very thick, and is composed of dark red, rough arkose materials, variable in texture. It is in places more resistant than the granites, so that side-gulches tributary to the east-flowing streams of the foothills are common in the granite just beneath the Fountain. Continuous troughs between the Fountain and the granite are not frequent. In many places the hard red sandstones form broad smooth-faced crags lying upon the outer foothill slope, reaching maximum size in the well known "flat-irons" south of Boulder (fig. 6). The other hard stratum is the massive gray sandstone known as the *Dakota*. It is separated from the Fountain by several less resistant strata of considerable aggregate thickness,



FIGS. 4, 5.—Upturned sedimentary ridges of mountain-front zone: fig. 4, eastward view in Perry Park, where a broad flat valley has been leveled between ridges and outer granitic foothills; floor of flat is of compacted angular fragments; vegetation is primitive grassland alternating with scrub oak; Dawson Butte in far background; fig. 5, southward view, between Golden and Morrison, of longitudinal valley inside Dakota hogback, shown on left in long curve.

and is usually seen as a bold ridge parallel to the outer slope of the foothills some distance to the east. The term "hogback" is familiarly applied to the steep Dakota cuesta.

A deep and wide trough usually extends between the Fountain crags and the Dakota cuesta. The upper part of the east-facing slope of this trough is the outcrop of a "creamy sandstone," which in places forms prominent outcrops, or even strong ridges, as at Morrison at the mouth of Bear Creek. Just east of and below the creamy sandstone is an easily eroded shale, which gives its rich red color to the deep soil of the valley. The west-facing slope, below the Dakota crest, is the outcrop of a calcareous sandstone stratum which is weathered so slowly as to be covered only by a thin soil. In certain places this limy sandstone stratum is hard enough to form a separate ridge or hogback crest.

The Dakota hogback is one of the most constant and conspicuous topographic features of the mountain-front, since it is practically everywhere harder than the strata above and below. Its top is usually quite even and straight, representing the level of a former graded surface. Its crest is quite rocky; there is no soil except in the crevices.

The present graded slope to the plains begins usually with the outer slope of the Dakota hogback, through first a layer of dark shales, then a thin limestone overlaid by soft light-colored shales, then clays and shales. Near every east-flowing stream, however, the graded slope is likely to be cut beneath by side-gulches cutting down into the dark shales, leaving a cut-off mesa with the limestone at its high western end.

Local distribution of vegetation in the mountain-front belt of upturned sedimentary rocks presents a variability apparently dependent almost entirely upon topography and soil texture, just as in the area of granitic foothills. There seem to be few if any perceptible differences in the floras of the different geological formations which can be traced to chemical differences in the substratum. It is perhaps true that cedars are more frequent in the limestone or calcareous sands of the stratum just below the Dakota, where these are exposed in gulches which notch the Dakota hogbacks, and that there are certain slight floristic differences between

granitic and sedimentary areas. This question has been discussed by RAMALEY (9), who found the two areas about the same in floras (in the Poudre mountain-front area), with *Cercocarpus* abundantly represented in the sandstone but not in granite. *Selaginella* apparently absent from the sandstone, and lichens infrequent there. Following a suggestion from COWLES, it appears to the writer that differences in rate of erosion of the substratum may explain the distribution of lichens, and perhaps *Selaginella* also. The sandstones are rather soft in the Poudre area, and wear away too rapidly for the lichens to establish themselves abundantly. The Fountain sandstone is harder in the Boulder region than elsewhere, and there at least it bears lichens almost as abundantly as do the granites. *Selaginella* is frequent in the sedimentary rocks in the Boulder area, as RAMALEY has pointed out. The writer knows of no plants which are restricted to either sedimentary or granitic areas, the only observed differences being those of relative abundance. The gulches, exposed slopes and crests, etc., of the sedimentary area are quite comparable to similar topographic situations of the granitic foothills, and have practically the same plant assemblages.

The rocky upper slopes of the Fountain, the Dakota, and other ridge-making strata, where they occur, lack soil except in crevices, and are mostly bare, except where rock pines or pinyons, shrubs of rocky situations (*Cercocarpus*, *Ribes*, *Jamesia*, etc.), and crevice plants, including many xerophytic herbs, can obtain a foothold. The west slope of hogbacks is blufflike, usually, and rocky, while the east slope is less steep (depending on the local angle of dip) and likely to be strewn with débris, as are the slopes of the harder exposures of the valley, and these have shrubby or herbaceous vegetation, sparse, and of species of rocky situations. The softer shales occupying the bottom of the valley are usually deeply buried by débris (of fine soil with imbedded rock fragments of all sizes), and support a grassland vegetation, which is luxuriant in the rainier parts of the growing season and very dry the rest of the time. A stream-bed in the bottom of the valley may be bordered by a strip of mixed shrub, *Crataegus*, oak, or canyon forest; or if dry, by scattered narrow-leaf cottonwoods and willows. Mesophytic

ravines developing in the sedimentary area support mixed-shrub, woodland, or mesophytic herbaceous growths, as in the granitic foothills. Local meadows (mesophytic grassland) are found on slopes where seepage or a high water table moistens a deep soil for at least part of the growing season.

In places the sedimentary rocks have been worn down more than is common, so that they are mostly or in part reduced to a general grade, above which the more resistant layers rise locally. This is the condition in the valleys of some of the larger streams from the foothills, and is seen at Platte Canyon, partially at Bear Creek (Morrison), and also in Perry Park (fig. 4) and the Garden of the Gods. The floor of this graded surface, especially in the Fountain exposures, is likely to be covered very thinly with small angular fragments, loose or compacted. The vegetation, as well as the soil, is very like that of gravelly floors in the foothills, being a variant of the primitive grassland association, with scattered rosette or mat plants, *Bouteloua hirsuta*, etc.

The climatic transition in the zone of upturned sedimentary strata is rapid. At Boulder and elsewhere dense cloud-banks have frequently been seen to descend to or just beneath the Fountain crags without continuing outward and downward to the plains (figs. 6, 7). The outer granitic hills and upper sedimentary slopes receive greater and more frequent precipitation than the lower slopes and adjacent mesas and plains; it may rain slightly below while it snows considerably above (cf. fig. 12); the outer and lower slopes are more exposed to wind, less cloudy, and in places less shaded from the afternoon sun by the higher granitic hills than the inner valleys and upper slopes. No exact data are available for this sudden climatic transition. Where the outcrop of sedimentary ridges and valleys is wide, as in the northern mountain-front region, the outer hogbacks are severely exposed to sun and wind, as in the open plains. Their coarse rocky soil favors woody plants; the xerophytic *Cercocarpus* shrub assemblage is here more extensively developed than anywhere else.

MESAS AND GRADED SLOPES OF THE DÉBRIS-APRON (fig. 8).—The general character of the graded slopes and their mesa-fragments has already been suggested. The mesas are of varying ages and

levels. They are described in the accounts of LEE (6), JOHNSON (5), FENNEMAN (2), SHANTZ (15), RAMALEY, ROBBINS, and DODDS (12), and VESTAL (17). The topographic parts of a mesa are: (1) the



FIGS. 6, 7.—Climatic transition at mountain-front: fig. 6, outer mountains just south of Boulder, seen from university campus; clouds beginning to form at summit of Green Mountain, while much of South Boulder Peak, at extreme left, is already obscured; snow covers the mountain slopes and fades out toward base of high mesas; roofs of distant buildings also white; fig. 7, practically same view, a little later, with upper slopes obscured; at one time it began to snow on mountains and upper mesas, and a few minutes later to rain in town; shortly afterward it changed to snow in the upper edge of town, so that the roof of the building with the short steeple at the right in midground, and of nearer houses, were well whitened, while rain still fell on the campus, less than half a mile away, and not more than 50 ft. lower; difference in elevation at mountain-front is critical as regards climatic change.

mesa-top, with flat surface covered with mixed rock-débris; (2) the edge or mesa-crest; (3) the side-slope; and (4) valleys or draws in the side-slope. The soil conditions and their effects on plant distribution have been discussed in the three articles last cited.

The débris-cover, where it has not been removed by recent erosion, extends far out into the plains. Its removal from the extensive areas of soft shales and clays marks a change from the flat terrace level to the easily eroded, gently rolling surface of much of the plains. The High Plains are extensive remnants of the old graded surface, away from the mountains.

The north-south distribution of the terraces is practically that of the mountain-front, although as conspicuous topographic forms the mesas are not so extensive. So far as effects on distribution of vegetation are concerned, the presence of the coarse mixed soil of




FIG. 8.—Table Mesa, about 7 miles north of Boulder; outlines of hills sketched in with ink; ridge DDD is Dakota hogback; Boulder mesas may be seen in figs. 6 and 7.

the detrital surfaces is the important physical condition. It permits the growth in the same small area of a rich variety of plants, representing numerous vegetation-types and different geographic elements.

PLATEAU AREAS, BUTTES, AND ESCARPMENTS (figs. 9-11).—Where the sedimentary strata are horizontal or of rather slight dip the harder layers protect the softer rocks beneath, and extensive plateau surfaces are left above the grade established by present erosion. These can be invaded only at the edge and by ravines which eat their way headward into the bluffs. Smaller elevated areas or buttes, recently or long ago cut away from plateaus by



meeting of two such ravines, are common. Older buttes are fewer and more distant from one another.



FIGS. 9, 10.—Buttes and plateau areas: fig. 9, North Table Mountain at Golden, west of Denver; this and South Table Mountain are capped with basalt; fig. 10, Fisher Peak, northern end of Raton mesas, as seen from valley of Purgatoire River, a few miles above Trinidad; upland in midground belongs to southern sedimentary plateau; vegetation is principally dry grassland with scattered pinyons and cedars and infrequent clumps of scrub oak.

The plateaus and buttes are found outside of the upturned ridge and valley zone, wherever the surface rocks are rather resistant. These resistant strata are usually the most recent and uppermost,

although the much older Dakota is at the surface over considerable areas in the plains drained by the southern tributaries of the Arkansas. For a considerable thickness above the Dakota the strata are mostly soft shales, which erode too readily to give tableland topography.

The larger through streams and their tributaries have cut below the level of the High Plains, leaving escarpments which are particularly notable near the Platte-Arkansas divide. Plum and



FIG. 11.—Buttes and plateau areas: divide between East Plum and West Plum creeks, in Castle Rock area, showing some of rhyolite buttes; one of the most imposing of these, Dawson Butte, shown in fig. 4.

Cherry creeks, running north into the Platte from the divide, and Monument, running into Fountain Creek, south to the Arkansas, have eroded deep valleys parallel to the mountain-front. Away from the mountain-front proper these valleys are bounded by lines of steep cliffs, but the west border of Monument and West Plum Creek valleys is the graded slope from the foothills, with its débris-covered terraces. Isolated buttes are present within these valleys, some of them protected by caps of igneous rocks from local outflows.

The southern part of the Sangre de Cristo Range (sometimes considered as a separate mountain chain, the Culebra Range) is flanked on the east by a sedimentary plateau which rises abruptly above the plains in a steep line of bluffs. The plateau is of sandstones mostly, of slight dip, and is much dissected by the eastward

flowing streams and their tributaries. On it rests the highland of the Spanish Peaks, and it is ribbed by resistant dikes of igneous material from two outflows, one set radiating downward from the peaks themselves. With the plateaus should be classed the high lava-capped mesas of the mountain-front and plains in the area near the Colorado–New Mexico boundary.

As in the hogback ridges, vegetation distribution in the plateau and butte areas is largely determined by soil texture and topography. Atmospheric conditions vary with exposure to wind and sun. The tops of the plateaus are covered with short-grass and




FIG. 12.—Unbroken short-grass ground cover in plains

mixed grassland over the level upland stretches of comparatively fine-textured soil. Exposed cliffs and crests, and rocky débris-slopes, afford lodging places for woody xerophytes (*Cercocarpus*, rock pines, pinyons, and cedars), with primitive grassland as the general ground-cover. The deeper and shaded parts of canyons and ravines approach a mesophytic condition, with mixed shrub and woodland vegetation.

#### PLAINS

Plains topography is typically flat or gently rolling country, with fine clay soil from a soft-shale substratum. Short-grass is the characteristic vegetation (fig. 12). Where the substratum is

sandstone the soil is more porous, with much sand; and plants of an assemblage typical of sandy soil are seen (17). Sand hills are present locally, usually to the leeward of larger streams.

Near the mountains the débris-cover, if present, considerably modifies topography, soil conditions, and vegetation. It may extend a long way into the plains, or may have been removed very near the beginning of the graded slope from the foothills.

Saline or alkaline areas are locally present. The valleys of the Arkansas and its tributaries (wet-weather streams, many of them, with trenched flood-channels) are in many places alkaline, and show prominent stands of *Sarcobatus-Chrysothamnus* vegetation.

Woody vegetation from the foothills extends locally far into the plains in rock outcrops, and along stony crests of stream-bluffs or terraces. The larger streams are bordered for many miles from the mountains by cottonwoods, usually scattered.

### Climate

The region has a continental climate, semi-arid, less so at the base of the mountains and in the foothills, with most of the rainfall in the warmer months. Wind movement, proportion of sunshine, and evaporating power of the air are high in the plains, with wide extremes of temperature; all of these features are less marked in the foothills.

The southern part of the region is warmer and drier than the northern, and with different distribution of rainfall. The rapid east-west change in elevation and topography at and near the mountain-front is accompanied by more or less considerable climatic variation; this with the local peculiarities occasioned by the elevated Platte-Arkansas divide, and the differences between areas north and south of the divide, may be seen in the summaries of climatic data for the particular subregions. These data have been taken from the summary of Climatological Data for eastern Colorado, southeastern Wyoming, and northeastern Colorado.<sup>4</sup> The facts shown in table II should be considered in the light of their

<sup>4</sup> Section 6, northeastern New Mexico, by C. E. LINNEY. Section 7, region drained by the Arkansas in Colorado, and section 8, region drained by the Platte in Colorado, by F. H. BRANDENBURG. Section 24, southeastern Wyoming, by W. S. PALMER.

determinative influence upon the vegetation; this can be done in only the barest manner in this section, but these relations are again brought out in the part on geographic relations of the vegetation.

Temperature conditions of the different parts of the region may be summarized as follows: The foothills have a lower mean temperature and shorter period without frost than either plains or mountain-front. Certain of the foothills vegetation-types and

TABLE II  
TEMPERATURE DATA

Area	Average mean temperature ° F.	Maximum temperature	Minimum temperature	Average number of days in growing season
Foothills (4).....	43.1	100	-36	99
Northern (2).....	42.0	98	-32	95
Southern (2).....	44.3	100	-36	104
Mountain-front (5, excl. Divide).....	50.5	104	-30	154
Northern (1, Boulder).....	50.9	97	-20	164
Divide (2).....	46.7	99	-33	122
Southern (4).....	50.3	104	-30	151
Plains near mountains (5).....	47.7	105	-38	138
Northern (3).....	47.1	105	-38	134
Southern (2).....	48.6	103	-32	143
Dry plains (5).....	50.4	106	-45	151
Northern (2).....	48.6	103	-45	145
Southern (3).....	51.0	106	-32	156
"Northern area" (8).....	45.8	105	-45	131
"Southern area" (11).....	49.3	106	-36	142

The number of stations for each area is given in parentheses. The mountain-front does not include the two stations of the Platte-Arkansas divide, which is so much more elevated than other parts of the mountain-front as to be much cooler. The "northern and southern areas" are respectively the northern and southern parts of the region, each extending over foothills, mountain-front, and plains.

many of the plant species are characteristic of northeastern and northwestern coniferous forest regions, are in fact southern extensions of them. The boreal character is much more evident in the higher mountains than in the foothills.

The mountain-front has the longest frostless season, the highest mean temperature, the mildest winters, and the least range in temperature extremes. Mountain-front localities are mostly comparatively sheltered; temperature inversion is common. Early spring plants flower several weeks earlier at the mountain-front than in either plains or foothills; at Boulder in spring the

season is in general 2-3 weeks in advance of that of Denver, 14 miles from the mountains.

The divide between Platte and Arkansas drainage, which should be considered in connection with the mountain-front area, has a mean temperature and frostless period intermediate between those of mountain-front and foothills areas, as it is intermediate in altitude and in vegetation.

The plains have a slightly lower mean temperature and shorter season without frost than the mountain-front area; the temperature of the dry plains at some distance from the mountains approaches that of the mountain-front more closely than that of the plains adjoining it. This difference is accompanied by a floristic one. Temperature extremes are greatest in the plains, a condition inimical to growth of woody plants.

The plains, mountain-front, and foothills in northern Colorado ("northern area") are cooler than those to the south, but the north-south differences in temperature and length of growing season due to latitude are of much smaller range and influence upon vegetation than the east-west differences due to altitude and changes of topographic character.

For purposes of comparison table III includes rainfall data for the higher parts of the mountains bordering the foothills on the west (montane zone), and for the plains of eastern Colorado bordering the region studied on the east. Annual rainfall is higher to the west, increasing with elevation, and higher also in the eastern plains, as a part of the gradual geographic increase of rainfall from the dry belt of the Great Plains eastward through the prairie region to the border of the eastern deciduous forest region. The eastern plains mark the transition from short-grass plains to the taller prairie-grass vegetation of the prairie, and are known in Colorado as "the rain belt." The driest part of the plains region lies between the rain belt and the plains near the mountains, in a zone distant from the mountains about 18-25 miles, and of a breadth 30-60 miles. It is narrowed on the west by the elevation of the Platte-Arkansas divide, and extends farther eastward in the Arkansas River Valley. It extends only a little way north into Wyoming and apparently is much narrowed on the west in extreme southern

TABLE III  
RAINFALL DATA

Area	Annual	January	February	March	April	May	June	July	August	September	October	November	December
Montane zone (6) 8250-10,265	22.58	0.80	1.10	1.96	2.75	2.85	2.01	3.38	2.70	1.61	1.64	0.80	0.90
Foothills (10) 6800-8000	16.30	0.42	0.60	1.11	2.17	2.23	1.27	2.79	2.06	1.29	1.27	0.55	0.44
Northern (6)	16.75	0.42	0.68	1.38	2.26	2.88	1.47	2.20	1.90	1.29	1.31	0.47	0.49
Southern (4)	15.63	0.44	0.49	0.69	2.03	1.25	1.18	3.67	2.29	1.27	1.23	0.68	0.37
Mountain-front (13) 5069-7373	16.25	0.40	0.68	1.07	2.00	2.59	1.79	2.41	1.81	1.23	1.11	0.53	0.54
Northern (5)	16.95	0.46	0.68	1.48	2.64	3.11	1.56	1.99	1.29	1.31	1.29	0.55	0.52
Divide (3)	16.53	0.35	0.56	0.94	2.08	2.69	2.17	2.73	1.96	0.77	1.11	0.47	0.60
Southern (5)	15.38	0.35	0.74	0.72	1.47	2.01	1.78	2.69	2.25	1.37	0.93	0.56	0.48
Plains near mountains (6) 4985-6098	14.36	0.33	0.49	0.81	1.85	2.28	1.76	2.17	1.81	1.20	0.81	0.47	0.43
Northern (3)	14.22	0.45	0.51	0.97	2.02	2.70	1.57	1.87	1.34	1.05	0.90	0.43	0.44
Southern (3)	14.54	0.20	0.64	0.65	1.07	1.86	1.94	2.47	2.27	1.34	0.72	0.51	0.43
"Northern area" (14)	16.28	0.44	0.64	1.33	2.20	2.92	1.53	2.06	1.57	1.27	1.22	0.50	0.49
"Southern area" (12)	15.25	0.35	0.59	0.69	1.46	1.72	1.56	2.96	2.27	1.33	0.93	0.59	0.43
Dry plains belt (7) 3899-5400	12.73	0.26	0.39	0.74	1.62	2.17	1.49	2.26	1.07	0.79	0.75	0.29	0.36
Eastern plains (8) 3380-5178	16.95	0.35	0.53	0.76	2.01	2.59	2.58	2.89	2.06	1.29	0.87	0.52	0.52

The northern area is the aggregate of northern foothills, mountain-front, and plains; the southern area is its southern equivalent. The number of stations is given in parentheses for each area, and the range of altitude above sea-level of the several north-south zones (in feet) is also included. The rainfall figures are in inches.

Colorado and northeastern New Mexico by the lava-capped plateaus which there extend eastward from the mountains.

As for the mountain-front and adjacent foothills and plains, the first two average about the same, the mountain-front receiving slightly more rainfall in the north and at the divide than the foothills. This may perhaps be due to the fact that in the northern part of the state, as at Boulder, the rain usually comes with east and northeast winds; and since the change of elevation is greatest at the mountain-front, more precipitation might occur there than in the foothills beyond. At Boulder the more mesophytic forms of vegetation occur more frequently and in larger areas in the sedimentary rocks of the mountain-front than in the granite foothills half a mile or a mile inside the foothills. In the southern part of the state the mountain-front is drier than the foothills, as a rule. The plains near the mountains receive almost 2 inches less rain, on the average, than foothills and mountain-front, and the dry plains to the east nearly another 2 inches less.

The "northern area" (foothills, mountain-front, and adjacent plains) receives on the average about 1 inch greater rainfall than the "southern area." Coupled with the higher temperature and greater evaporation, this results in a considerably more xerophytic vegetation south of the Platte-Arkansas divide. There is little difference in the plains, but at the mountain-front, with a difference of 1.57 inches, the vegetation to the south is markedly drier.

COOPER finds, in the chaparral region of California, that very slight differences in the original physical conditions of north and south slopes result in very marked differences in vegetation. The same principle seems to apply, in perhaps a smaller degree, in a semi-arid region like the Colorado mountain-front. It appears that differences in rainfall from place to place, or from month to month, although small in absolute amount, can be critical in their influence upon vegetation distribution. The slight differences appear to represent marginal values above or below a critical point. The difference in vegetation in two areas, moreover, is not necessarily the result of climatic difference, but is a resultant of differences in soil, topography, geographic position, and vegetational history, in addition to climate. It should not be surprising, therefore, that



areas having climates not widely dissimilar, as the plains of the rain belt and the northern foothills, should have distinctly unlike vegetation.

**MINIMUM RAINFALL.**—One factor which seems to be partly responsible for the generally xerophytic character of the entire region studied, the plains in particular, is the wide variation in the amount of rainfall from year to year. The minima have been

TABLE IV  
MINIMUM ANNUAL RAINFALL

AREA	NUMBER OF STATIONS WITH RECORDS		AVERAGE MINIMUM FOR AREA		LOWEST MINIMUM RECORDED FOR ANY STATION IN AREA	
	1893	Other years	1893	Other years	1893	Other years
Montane zone.....	—	6	—	<b>15.65</b>	16.55 Frances	<b>11.36</b> (1907) Cripple Creek
Foothills.....	4	9	—	<b>12.83</b>	<b>7.16</b> Box Elder	<b>10.93</b> (1908) Cheesman
Mountain-front and divide.....	7	11	9.12	<b>11.91</b>	<b>7.03</b> Waterdale	8.76 (1890) Table Rock
Plains near mountains.	5	6	9.39	<b>8.91</b>	7.11 Fort Collins	<b>5.04</b> (1876) Cheyenne
Dry plains belt.....	7	7	8.11	<b>7.01</b>	5.40 Greeley	<b>3.78</b> (1894) Las Animas
Eastern plains.....	5	8	10.48	<b>10.61</b>	8.30 Cheyenne Wells	<b>6.97</b> (1894) Cope

Two stations within the foothills area are exceptional as to rainfall, and have not been included in the averages. These are Salida in the Arkansas Valley above the Royal Gorge, and Westcliffe in the Wet Mountain Valley. Similarly, Canyon City at the debouchure of the Arkansas, and Raton and Las Vegas in New Mexico, have been excluded from the mountain-front area. The stations with the lowest minima have been mentioned in the table. The lowest minimum in each area, whether in 1893 or in some other year, is printed in bold face. Except for Cheyenne Wells, which is remote from the mountains, all of the stations noted as having had least rainfall in 1893 are within a limited area (in the northern part of the region), which seems to have been most severely affected by the drought of that year.

tabulated for the several parts of the region from the climatic summaries of the Weather Bureau. The year 1893 happened to be exceptionally dry, and the minima for many of the stations fall in it. Dryness in other years has been of more local prevalence. It has seemed preferable to present the data for 1893 separately from that of other years. The data for 1893 are not available for all stations in each area, and so the number of stations from which data have been used is mentioned for each area (table IV). The column presenting the average minima for the several areas (minima



of all stations for each area averaged, excluding figures for 1893) seems to express the main fact of the table, that the rainfall reaches successively lower minima downward and eastward from the mountain zone through the foothills, mountain-front, and adjacent plains to the dry plains belt, beyond which the minima rise gradually with the gradual increase of rainfall eastward into the prairie-grass region. It appears also that annual rainfall values falling considerably below the average (as much as 4 inches below) occur more frequently in the plains than in the mountain-front and foothills areas. The well known uncertainty of farming without irrigation in much of eastern Colorado, due to frequency of very dry years, indicates further that it is not the average rainfall so much as the constantly recurring minimum which determines whether or not an area can support a cultivated or natural vegetation which is other than decidedly xerophytic.

SEASONAL DISTRIBUTION OF RAINFALL.—On the whole, precipitation during the cooler months is quite low; this is not so true of the montane area just to the west of and above the foothills. The summer rainfall is greater, but in most places distributed rather unevenly. June is thus drier than either May or July over practically the entire region. The northern area near the mountain-front receives more rain in the spring and early summer months, while the southern area receives more of its rain during late summer. This difference between north and south is of far-reaching influence upon the character and distribution of vegetation. The details of seasonal distribution of rainfall are shown in the table of averages of rainfall data, and in figs. 13-16.

The northern and southern parts of the zones at and near the mountain-front are so different as to rainfall that they cannot be incorporated in single graphs. The northern parts of the zones are selected, therefore, as the more typical. The graph for the mountain-front is omitted to avoid overcrowding, but it can be seen in fig. 14. Excluding the eastern plains, the zones have the same type of rainfall, with greatest abundance in May and July, and a decline in June. The zones are successively drier with decrease of elevation, and this is almost as true for particular months as it is for the entire year. The eastern plains have higher

summer rainfall than the plains near the mountains; the distribution is similar, except that June is as rainy as May.

The graphs for foothills and plains near the mountains are repeated in fig. 14. These zones and the mountain-front have maximum rainfall in May, with a sharp decline in June, followed by slightly greater rainfall in July. Despite its less elevated position, the mountain-front receives greater spring rainfall than the foothills.




FIGS. 13, 14.—Seasonal distribution of rainfall: fig. 13, comparison of north-south zones; fig. 14, northern foothills, mountain-front, and plains.

The data for the south are not so dependable as for the north, for some of the few stations are exceptionally situated. The contrast shown with the northern area is marked, however. The rain is less abundant in spring and more abundant in July and August than to the north.

The graph shown for the "northern area" is a composite of the 3 in fig. 14, that of the "southern area" is a composite of those

in fig. 15; they contrast strongly. The northern area is characterized by the Rocky Mountain foothill type of rainfall, the southern area by the New Mexican type (WARD 19). Both of these types are described as having a single maximum, for the first in May, for the second in July-August. The northern area receives most of its rain from northeasterly winds; the southern area probably from southeasterly winds. The centrally situated Platte-Arkansas



FIGS. 15, 16.—Seasonal distribution of rainfall: fig. 15, southern foothills, mountain-front, and plains; fig. 16, northern and southern areas and divide.

divide receives rain from both directions, and has both maxima, with a higher June rainfall, partly because of its considerable elevation. Probably rain is carried from either direction past the divide, producing secondary maxima, in July in the northern area, and in April in the south.

The abundant rainfall of the divide, especially in June, forms a local rainfall type which is intermediate between that of well watered parts of the foothill zone and that of the eastern or rain-

belt plains. The divide is also cooler than most parts of the mountain-front and adjoining plains. The vegetation of the divide is likewise transitional between that of rain belt and foothills, with the more nearly mesophytic forms of grassland, and with woody plants of the foothills extending many miles eastward from the mountains.

The effects upon the vegetation of the difference in distribution of precipitation north and south of the divide are discussed in the section on geographic relations, but may be summarized briefly herewith.

TABLE V

## INFLUENCE OF SEASONAL DISTRIBUTION OF RAINFALL ON VEGETATION

## NORTHERN AREA

The greatest rainfall is in April and May.

There is greater activity of vegetation, more luxuriant growth, and greatest abundance of flowers in spring.

There are many spring-flowering plants from the mountains of rather mesophytic character, in mixture with plains plants in the mountain-front zone.

Distribution of the bunch-grass association and of the less xerophytic plants, requiring a long season of continued moisture, is limited.

The northern plains near the mountain-front flower luxuriantly in spring and early summer, but only the more xerophytic composites, etc., in late summer, in which respect the plains are more like the driest plains just east of them in late summer.

## SOUTHERN AREA

The greatest rainfall is in July and August.

There is greater activity of vegetation and more luxuriant growth in late summer.

There is absence or scarcity of spring-flowering mountain plants, and greater prevalence of plains plants in the mountain-front zone.

Distribution of bunch-grass is less restricted; there is a greater prevalence of late-flowering plants not intensely xerophytic, as some of the asters and goldenrods, etc.

The southern plains near the mountains contain fewer spring flowers, but many long-season plants absent from the dry plains and the northern plains near the mountains are present, as the annual sunflowers. In this respect the plains are more like those of the rain belt of eastern Colorado in late summer.

It is remembered that the southern area is in general drier and warmer, with somewhat more xerophytic vegetation than the northern area, and that differences in vegetation due to this cause must be distinguished as well as possible from those due to different distribution of rainfall.

Evaporating power of the air has not been subject to geographic-statistical treatment, since there are no data. It was beyond the scope of the present study to have attempted instrumental investigation on a scale large enough to be of value. There seems to be little doubt that, as a geographic factor *in regions of continental climate*, evaporating power of the air is of about the same indicator value as rainfall. It varies geographically about as does rainfall, in inverse ratio, since evaporating power is, in large measure, a function of rainfall. This same inverse ratio seems to hold in seasonal distribution as well as geographically. This may be seen from the graphs of WEAVER (20), and from data obtained by COOPER in a study of chaparral in California.

As a local factor evaporation is separately treated in the discussion of local distribution of vegetation.

For further discussion of the climatology of Colorado in relation to vegetation the reader is referred to the articles of SHANTZ (15, 16), RAMALEY (12), and ROBBINS (13, 14). Data may be had from the bulletins of the United States Weather Bureau, Colorado College, the Agricultural Experiment Station at Fort Collins, and the Bureau of Plant Industry.

### Local distribution of vegetation

PHYSICAL FACTORS.—Local physical conditions affecting plant distribution are those concerned with *stratum* and *soil*; with *topography*, especially *local position* with respect to surroundings, and *slope* of surface, as regards both steepness and direction of exposure; and with *local variation in atmospheric conditions*, as controlled primarily by topography. The variability of these factors within the region is great, and their interactions are complex. Descriptions of the soil, topography, atmospheric conditions, etc., of the different parts of the region are scattered through both systematic and regional sections of this study, and a lengthy discussion at this point would be out of place. A few references to other parts, and certain incidental comments, may here be made.

*The character of the stratum*, and some of its influences in determining soil conditions and topography, are indicated in the

account of the sedimentary area. The contrast between the granite soil of the foothills and the soil of sedimentary origin lying just outside, with its slight selective action on flora and vegetation, has also been noted. Other mentions of soils, especially as regards soil texture, are scattered.

*Topography* is systematically treated for particular regions and smaller areas by dividing each type of topographic complex into topographic areas or habitats; with each type is correlated the particular plant community or the several communities which accompany it. In the regional section will be found similar analyses of the cuesta, high mesa, mesa-terrace, and flood-plain complexes. Particular physical factors controlled by local position and by slope are mentioned in a former article (18).

*Atmospheric factors* vary locally in this region to a probably not very great extent, but even slight differences may be critical, as has been found by COOPER in the California chaparral. The factor of greatest influence upon plant life, and the one most readily measured, is the evaporating power of the air, the value of which represents the resultant of several contributing factors. Local distribution of evaporating power is believed to be controlled primarily by differences in topography, and secondarily by differences in vegetation-cover. That is to say, flatness of the land surface makes for comparative uniformity of exposure to wind and sun; hilliness causes diversity of exposure. Local water or wet-soil surfaces may lower evaporating power by contributing much water vapor to the air. Topography thus determines the *original* local distribution of evaporating power. This original local distribution is modified by vegetation-cover. In flat country the uniformity is changed. Low and open vegetation lowers evaporating power at the ground surface only slightly, but mesophytic closed forest lowers it very greatly (GATES 3, 4). In hilly country in not too humid climates the originally protected ravines and shaded or wind-sheltered slopes may develop mesophytic vegetation which still further lowers evaporation, while the originally exposed slopes and summits usually remain xerophytic. Thus, in the mountain-front region here considered, primary environmental differences

due to topography may rather thoroughly control vegetation distribution. In such cases the reaction of vegetation-cover upon local evaporation conditions may merely heighten the original topographically determined contrast between protected and exposed habitats. Topography governs local vegetation distribution through the mediative influence of a number of physical factors, of which evaporating power is one. Depending as it does upon several other factors, evaporation forms a convenient index of habitat, but is not in itself the basic controlling condition. For these reasons the writer has subordinated the influence of evaporating power upon local distribution to that of topography.

The sudden change of elevation at the mountain-front is a topographic condition affecting evaporating power. At many places the hogbacks, mesas, and outer slopes receive no direct sunlight during several hours before sunset, being shaded by the higher slopes immediately to the west. This contributes to the comparative mesophytism of certain mountain-front stations where the descent from foothills to plains is more than ordinarily abrupt.

Direction of exposure affects local atmospheric conditions and vegetation in many easily observed ways. Cloudiness and showers occur on summer afternoons much more frequently than in the mornings, as RAMALEY has noted. East-facing slopes are thus likely to be drier than west-facing slopes (the latter are less frequent east of the range-crest). As would be expected, the difference between north- and south-facing slopes is considerable, the latter being more exposed to sun and conditions favoring rapid evaporation, and with sparser, more xerophytic vegetation. In open parts of the foothills where slopes are quite gentle the north-facing slopes are not sufficiently sheltered from sun and wind to differ in vegetation from the south-facing slopes in any marked degree. Steep north slopes, or both sides of steep and narrow ravines which run down to the north, however, are quite mesophytic. In different parts of so large a territory the combinations of contrasting vegetation of north and south slopes would be expected to vary, and a few of them are listed herewith by way of illustration.



TABLE VI  
EFFECTS OF DIRECTION OF SLOPE UPON LOCAL DISTRIBUTION

Locality	Vegetation of south-facing slope	Vegetation of north-facing slope
Poudre foothills. . . . .	Grassland	Scattered rock pine, with more mesophytic vegetation infrequent
Foothills near Boulder. . .	Grassland, rock pine, mixed shrub	<i>Pseudotsuga</i> , canyon forest, rock pine, mesophytic grassland
Poudre mountain-front. . .	Grassland	<i>Cercocarpus</i> , with very scattered rock pines in rocky places; grassland in fine soil
Mountain-front near Boulder. . . . .	Grassland, mostly	Grassland with rock pine, mixed shrub, etc.
South of Golden, mountain-front. . . . .	Grassland and <i>Cercocarpus</i>	Grassland with scattered rock pine and mixed shrub
Perry Park. . . . .	Oak and grassland	Rock pine and <i>Pseudotsuga</i>
Palmer Lake. . . . .	Oak	<i>Pseudotsuga</i>
Southern mountain-front in general. . . . .	Pinyon-cedar, dry grassland, and scattered oaks	Closer and taller oak growth with rock pines

FACTORS OTHER THAN PHYSICAL CONDITIONS OF HABITAT.—

If the physical conditions which determine the habitat and all their interactions and variations were fully known, however, the local distribution of plant communities as observed would only partially be explained. Within even a very small part of the region studied correlations between physical habitats and vegetation-types must not be too closely drawn. The rock pine, for example, grows in any soil or on any slope; its presence or absence in any particular situation is not alone a matter of physical conditions there and then operative. Local distribution of vegetation-types in these partly unstable and locally very diverse situations depends also on at least three other sets of conditions: (1) range of toleration, in individual species or groups of species, of variation of physical conditions; (2) local historic factors, physical and vegetational, which have been operative in any given spot (these often cannot be determined); (3) accident of seed distribution and germination. For these reasons it seems best to characterize the vegetation-units, in most cases, from the vegetation itself rather than from habitat. There can be no question

that, in general, local variation of present physical conditions of the habitat governs to a considerable degree the distribution of plant communities, but the need of at least recognizing these other sets of factors should be emphasized. It must be further seen that, in the invasion of a new habitat, representatives from more than one plant community can be successful in establishing themselves, resulting in *mixed vegetation-types*. In fact, probably the greater part of the area studied is occupied by mixed associations or *micelia* (CLEMENTS). Even areas of established vegetation are usually open enough to permit the continual ecesis within them of new plant immigrants from quite different communities. This diversity is likely to be relatively enduring, for plant competition usually does not here operate to exclude all but a single type of dominants. The opposite relation between plants, which may be called accommodation, is as greatly in evidence. The control exerted by vegetation upon the physical environment is slight over the generally xerophytic mountain-front region.

A second factor contributing to the mixed effect is the frequent extremely local variability of physical conditions within the habitat. This might be called *mosaic variability*, and its effect a *mosaic mixture* of vegetation-types. The influence of large surface rocks partly imbedded in fine soil, allowing the growth of comparatively mesophytic plants in a rather constant interspersal with xerophytes over a considerable area, may be cited as an example.

### Vegetation-types and their distribution

Since the plant communities have been described separately in the two articles preceding this, their systematic characterization here may be condensed very considerably. A tabular view of the communities, giving some idea of their general character and of their distribution in the main geographic divisions of the region studied, is shown in table VII.

Some of the more important features of the particular associations may now be noted. Details and references to other accounts may be found in the articles preceding. The general appearance of certain vegetation-types may be seen in fig. 17.

*Lichen association*.—Lichens, especially *Rinodina*, *Lecanora*, and *Parmelia conspersa*, partly cover the dry rock surfaces, especially granites in the foothills and the craggy outcrops and loose surface rocks of the mountain-front. Rock exposures are infrequent in the plains proper.

TABLE VII  
CONSPECTUS OF ASSOCIATIONS

	Foothills	Mountain-front	Plains
Thallus vegetation	Lichen association	Lichen association	(Lichen association)
Grassland	Foothills grassland	Mixed short-grass	Short-grass
Extensive, climatic . . . . .		{Wheat-grass <i>Stipa-Aristida</i>	Wheat-grass <i>Stipa-Aristida</i>
Local, edaphic	Bunch-grass		Bunch-grass
Xerophytic . . . . .		Mesophytic grassland	Prairie-grass
Less xerophytic . . . . .	forest herb type meadow type		
		Foothills primitive grassland	Primitive grassland
Primitive . . . . .	<i>Artemisia frigida</i> con- socioes		
		Primitive bunch-grass	Primitive bunch-grass
Shrub vegetation			
Xerophytic . . . . .	<i>Cercocarpus</i> association	{ <i>Chrysothamnus-Sarcoba-</i> <i>tus</i> association <i>Cercocarpus</i> association Mixed shrub association	<i>Chrysothamnus-Sarcoba-</i> <i>tus</i> association
Xerophytic to mesophytic . . . . .	Mixed shrub association		(Local shrub communi- ties)
	<i>Arctostaphylos</i> <i>Ceanothus</i> association <i>Symphoricarpos</i>	<i>Symphoricarpos</i>	( <i>Symphoricarpos</i> )
Tree vegetation			
Coniferous			
Xerophytic . . . . .	Pinyon-cedar associa- tion	Pinyon-cedar associa- tion	
Less xerophytic . . . . .	Rock pine association	Rock pine association	
Relatively mesophytic . . . . .	<i>Pseudotsuga</i> association	( <i>Pseudotsuga</i> associa- tion)	
Deciduous			
Xerophytic to meso- phytic . . . . .	(Oak association <i>Populus-Salix</i> associa- tion)	Oak association <i>Populus-Salix</i> associa- tion	<i>Populus-Salix</i> associa- tion
Relatively mesophytic . . . . .	Canyon forest (Aspen association)	Canyon forest	

Associations with equivalent or similar representation in plains, mountain-front, and foothills areas are shown on the same horizontal line. Very local or poorly developed representation of a community in a particular zone is indicated by parentheses.

*Mixed grasslands and short-grass*.—The shallow-rooted short-grasses, *Bouteloua* and *Bulbilis*, dominate the compacted fine soil surface of most of the plains, as the well known short-grass association. *Bouteloua* alone, with admixture of plants of different physiological and geographic character, is the important element of dry




FIG. 17.—Distribution of vegetation in typical northern mountain-front area (outlet of Gregory Canyon, near Boulder): habitats and vegetation-types distinguishable are: 1, craggy exposure of Fountain sandstone; 2, rock talus, with beginning of primitive grassland; 3, stabilized slope with primitive grassland; 4, mixed grassland, in places approaching mesophytic type, mostly on northwest-facing slope; 5, more nearly xerophytic mixed grassland, with bunch-grass and ruderal elements on northeast-facing slope; 6, *Symphoricarpos*; 7, mixed shrub; 8, canyon forest; 9, *Populus angustifolia* stream-side community; 10, rock pine; 11, *Pseudotsuga*, in scattered growth, including rock pine and other components, on north-facing slopes of Green Mountain.

grassland in the débris-covered soil of the mesas and outwash-plains of the mountain-front (short-grass mixed association). The foothills mixed grassland, very similar to the mixed short-grass, is typical over the coarse surface of the granitic foothills.

*Wheat-grass.*—This taller but also shallow-rooted grass, *Agropyron Smithii*, dominates areas of loose clay in the mountain-front and plains. Its ecological character is not well understood.

*Stipa-Aristida association.*—These tufted xerophytic grasses of coarse soil occur frequently but not extensively, together or singly, with other rather deep-rooted plains xerophytes.

*Bunch-grass.*—Tufted perennial and deep-rooted grasses, depending on continued moisture, such as *Andropogon scoparius*, *A. furcatus*, *Sorghastrum nutans*, etc., are notable through most of the prairie region, almost absent in dry plains, but abundant in the rain belt of eastern Colorado; at the mountain-front and in the foothills, scatteringly in the north, but very frequent at the Platte-Arkansas divide and southward into New Mexico. In the foothills there are grasses of similar habit which mingle with the prairie bunch-grasses.

*Mesophytic grasslands.*—Mesophytic herbaceous growths are made up partly of prairie plants and partly of Rocky Mountain forest plants. The latter element is very considerable in occasional foothill ravines. Meadow growths of both foothills and mountain-front, in moist soil, with showy flowering plants like *Delphinium*, *Cerastium*, *Castilleja*, *Orthocarpus*, etc., are conspicuous in early summer, but not very frequent. The mountain-front in many places shows a mixed grassland much like that of eastern prairie, which has been called western prairie-grass. It has plants of the mixed short-grass, with components from bunch-grass and eastern prairie or forest border, with some foothills mesophytes, and a few plants characteristic of the mountain-front, like *Stipa viridula*.

*Primitive grasslands.*—Early stages of grassland developing in areas recently bared, or remaining for long in loose shifting slopes, are frequently seen. Prostrate plants with heavy taproots (rosette plants) are common. Gravel-slides in the foothills and dry stony crests of mesas, buttes, and ridges in the plains and mountain-front are the typical habitats. The *Bouteloua hirsuta*

and the *Artemisia frigida* consociates may be mentioned specially. The last is closely allied to the *Gutierrezia-Artemisia* association of the plains, very widespread, and continuing, at the expense of short-grass, with heavy grazing. In mountain-front and plains the primitive bunch-grass association, with *Panicum virgatum*, *Sporobolus cryptandrus*, *Stipa Vaseyi*, *Eriocoma*, etc., occupies sandy or loose-soil habitats recently disturbed.

*Chrysothamnus-Sarcobatus* association.—The shrubby composite, *Chrysothamnus* (rabbit-brush), and the chenopodiaceous greasewood occupy loose soil, mostly alkaline areas, on certain slopes in the mountain-front, and are particularly abundant in stream-bottoms in the southern plains.

*Cercocarpus* association.—Mountain-mahogany, of the rose family, is the only dominant in the open shrub growth of the mountain-front and outer foothills, in very dry exposed situations and usually stony soil. In the interstices between shrubs are plants of primitive grassland or mixed short-grass.

*Mixed shrub* association.—This is a heterogeneous assemblage of shrubs, ranging from xerophytic, like *Rhus trilobata*, to relatively mesophytic forms, like *Crataegus coloradensis*, in sheltered situations. The same species range through a variety of habitat conditions, and may form a community either as shrubs or trees. The mixed shrub grades into the canyon forest.

*Arctostaphylos* association.—The well known and widespread bearberry forms its characteristic mats in the foothills, mostly on compacted gravelly floors. It is more abundant in the upper foothills, in open places among the scattered pines. Its congener, *Juniperus communis sibirica*, is present but infrequent.

*Ceanothus* association.—*Ceanothus Fendleri* forms low matlike ground-cover in the lower foothills, similar to that of *Arctostaphylos*, though it is not evergreen, is of more southerly distribution, and ranges into drier and more exposed habitats. It favors the establishment of seedling mesophytes, and plays a part in revegetation of burned areas.

*Symphoricarpos* association.—The buckbrush, as it is called, occupies moist fine soil, and invades grassland in the mountain-front and foothills, as well as in the eastern prairie, in favorable

situations such as draws and seepage areas of slopes. It in turn is frequently displaced by taller woody vegetation.

*Rock-pine association.*—*Pinus scopulorum* is the important tree of the foothills. It ranges into very variable habitats, and is structurally variable in accordance. It forms infrequent close stands, but in most places is scattered, the ground between the trees being occupied by foothills mixed grassland, *Ceanothus*, *Arctostaphylos*, etc. It is frequent in rocky crests, etc., in the mountain-front, except in the south, where it is commonly replaced by pinyon. It extends very locally into the plains in broken country, on butte-crests, etc., and on the elevated Platte-Arkansas divide.

*Pinyon-cedar association.*—*Pinus edulis* and *Juniperus monosperma* are important xerophytic conifers of the southern mountain-front and lower foothills north to the Garden of the Gods, and extending into the southern plains on mesa-crests, canyon-walls, and bluffs of broad valleys. The soil is usually rocky or gravelly. The trees are low and rounded, and do not form a closed assemblage.

*Pseudotsuga association.*—*Pseudotsuga mucronata* forms the mesophytic or relatively mesophytic coniferous forest of the region, and is confined to sheltered ravines and steep north slopes in the foothills. It is infrequent at the mountain-front.

*Oak association.*—Small trees of the white-oak group, of uncertain taxonomic affinity, form dense copses or open woods in the lower foothills and in the mountain-front about as far north as Platte Canyon. In places grazing destroys the oak slowly and allows increase of grassland. The undergrowth of mesophytic oak areas is much like that of the canyon forest.

*Populus-Salix association.*—In stream-side areas of the foothills *Populus angustifolia* and 4 or 5 common willow species are frequent. Outside the mountains *Populus Sargentii*, and in the south *Populus Wislizeni*, replace the narrow-leaf cottonwood. Cottonwoods extend eastward into the plains for many miles along watercourses.

*Canyon forest.*—The deciduous trees of the foothill canyons and of ravines, etc., in the mountain-front, include *Alnus tenuifolia*, *Betula fontinalis* (these two common along mountain streams),

*Amelanchier alnifolia*, *Prunus pennsylvanica*, *P. americana*, *P. demissa*, *Crataegus coloradensis* et spp., *Robinia neomexicana* (in the south only), *Acer glabrum*, and *A. Negundo*, with a few others. A few shrubs are present and a variable undergrowth, with one typical aspect of *Viola canadensis* Rybd., *Hydrophyllum*, and *Galium*. A *Ligusticum* is very abundant in places.

*Aspen association*.—*Populus tremuloides* is restricted, in all but the highest parts of the foothills, to relatively mesophytic ravines. It does not come up abundantly following burning of the pine forest, as is true in the higher elevations and farther north.

EASTERN ILLINOIS STATE NORMAL SCHOOL  
CHARLESTON, ILL.

#### LITERATURE CITED

1. DAVIS, W. M., The Colorado Front Range. Ann. Ass. Amer. Geog. 1:21-83. 1911.
2. FENNEMAN, N. M., Geology of the Boulder district, Colorado. U.S. Geol. Surv. Bull. 265. pp. 101. 1905.
3. GATES, F. C., The relation between evaporation and plant succession in a given area. Amer. Jour. Bot. 4:161-178. 1917.
4. GLEASON, H. A., and GATES, F. C., A comparison of rates of evaporation in certain associations in central Illinois. BOT. GAZ. 53:478-491. 1912.
5. JOHNSON, W. D., The High Plains and their utilization. Ann. Repts. U.S. Geol. Surv. 21, part 4, pp. 599-741; 22, part 4, pp. 631-669. 1900, 1901.
6. LEE, W. T., The origin of the débris-covered mesas of Boulder, Colorado. Jour. Geol. 8:504-511. 1900.
7. MARVINE, A. R. [The sedimentary rocks east of the Front Range (chap. ii, pp. 93-137, in Marvin's report)], in HAYDEN, F. V., Ann. Rept. U.S. Geol. and Geog. Surv. Terr. for 1873, embracing Colorado. 718 pp. Washington, 1874.
8. RAMALEY, F., Plant zones in the Rocky Mountains of Colorado. Science 26:642-643. 1907.
9. ———, Botany of northeastern Larimer County, Colorado. Univ. Colo. Studies 5:119-131. 1908.
10. ———, Dry grassland of a high mountain park in northern Colorado. Plant World 19:249-270. 1916.
11. ———, Vascular plants of the Tolland region in Colorado. Univ. Colo. Studies 12:27-51. 1917.
12. RAMALEY, F., ROBBINS, W. W., and DODDS, G. S., Studies in mesa and foothill vegetation, I. Univ. Colo. Studies 6:11-49. 1908.



13. ROBBINS, W. W., Climatology and vegetation in Colorado. BOT. GAZ. 49:256-280. 1910.
14. ———, Native vegetation and climate of Colorado in their relation to agriculture. Bull. 224, Agric. Exp. Sta. of the Colo. Agric. Coll. 56 pp. 1917.
15. SHANTZ, H. L., A study of the mesa region east of Pike's Peak: the *Bouteloua* formation. BOT. GAZ. 42:16-47, 179-207. 1906.
16. ———, Natural vegetation as an indicator of the capabilities of land for crop production in the Great Plains area. U.S. Dept. Agric., Bur. Pl. Industry, Bull. 201. pp. 100. 1911.
17. VESTAL, A. G., Prairie vegetation of a mountain-front area in Colorado. BOT. GAZ. 58:377-400. 1914.
18. ———, Foothills vegetation in the Colorado Front Range. BOT. GAZ. 64:353-385. 1917.
19. WARD, R. DE C., Rainfall types of the United States. Geog. Review 4:131-144. 1917.
20. WEAVER, J. E., Evaporation and plant succession in southeastern Washington and adjacent Idaho. Plant World 17:273-294. 1914.