

## First valid record of *Astacilla* Cordiner, 1793 in Australia, with description of a new species (Crustacea: Isopoda: Arcturidae)

Rachael A. King

Museum Victoria, PO Box 666E, Melbourne, Vic. 3001, Australia, and Department of Zoology,  
University of Melbourne, Parkville, Victoria 3052, Australia  
Present Address: Southeastern Regional Taxonomic Center, Marine Resources Research Institute,  
PO Box 12599, Charleston, SC 29422, USA.  
kingr@mrd.dnr.state.sc.us

---

**Abstract** – A new species of arcturid isopod, *Astacilla lewtonae*, is described from Western Australia. Although other Australian species have been attributed previously to this genus, this is the first record for the genus as currently diagnosed.

### INTRODUCTION

The Arcturidae Dana, 1849 are a specialised group of marine isopods, which use the anterior four pairs of pereopods as a setose filtering complex extending from a cylindrical body usually elevated above the substrate. The family belongs to the isopod suborder Valvifera, which has recently been restricted by Poore (2001) who removed many genera traditionally placed in it to other families, Antarcturidae Poore, 2001 in particular.

The first description of an Australian arcturid isopod was of *Arcturus brevicornis* Haswell, 1881 from New South Wales. The types of this species are lost but it is probably a species of *Neastacilla*. Further species of *Arcturus* were described from New South Wales by Whitelegge (1904), *Arcturus alcornis*, *A. dentatus*, *A. nodosus*, *A. serratulus*, *A. simplicissimus*, the types of all of which are also lost. All probably belong in the Antarcturidae.

The first review of Australian arcturid taxa was that of Hale (1924), who examined many specimens and established the endemic genus *Parastacilla* for two very distinctive species. He also described two other species and placed them in *Neastacilla* Tattersall (*Neastacilla algensis* and *Neastacilla deducta*). In a further, more comprehensive, review Hale (1946) described five arcturid species from the Southern Hemisphere and placed them all in *Astacilla* (*Astacilla attenuata*, *A. macilenta*, *A. sheardi* and *A. vicaria* from Australia and *A. fusiformis* from New Zealand) arguing that the genus *Neastacilla* Tattersall was poorly differentiated from *Astacilla* and should not be recognised. Hale (1946) re-examined his two previously described Australian species, *Neastacilla algensis* Hale, 1924 and *Neastacilla deducta* Hale, 1924, moving both to *Astacilla*, and included *Astacilla marionensis* Beddard, 1886 and *Astacilla kerguelensis* Vanhöffen, 1914 from the Southern Ocean in his discussion.

Guiler (1949) was also reluctant to confirm the genus *Neastacilla* and described five new Tasmanian species of *Astacilla* (*Astacilla monoseta*, *A. inaequispinosa*, *A. unicornis*, *A. derwenti*, and *A. oculata*). Of these, only *A. monoseta* and *A. inaequispinosa* are currently valid species (Poore *et al.*, 2002).

All southern Pacific species of *Astacilla* were removed to *Neastacilla* by Kussakin (1972) who published a new diagnosis of *Neastacilla*. A thorough review of 18 Australian species of *Neastacilla* (Lew Ton, 1980) concurred with Kussakin's conclusions, finding that *Neastacilla* was a valid Pacific genus clearly separated from *Astacilla*, which had no known representatives in Australia. So, until now the family has been represented in Australia by *Amesopous* Stebbing, 1905, *Neastacilla* Tattersall, 1921, and *Parastacilla* Hale, 1924, only the last being endemic (King, 2000; Poore *et al.*, 2002).

Although *Astacilla* is almost certainly paraphyletic (King, 2001), a useful diagnosis can be written. *Astacilla* currently comprises 32 species distributed in the northern and central North Atlantic, Mediterranean, northern North Pacific, southern Africa, and in India (Table 1). The new species described here extends the range of the genus to the eastern Indian Ocean and is the first record from Australia.

Material is deposited in the Western Australian Museum, Perth (WAM) and Museum Victoria, Melbourne (NMV).

### SYSTEMATICS

#### *Astacilla* Cordiner, 1793

*Astacilla* Cordiner, 1793. –Sars, 1897: 87. –Monod, 1970: 1127–1142. –Kensley, 1983: 163–164.

Table 1 The species of *Astacilla* and their distributions.

Species	Distribution
<i>A. amblyura</i> Stebbing, 1905	Southern India (Pillai, 1963).
<i>A. arietina</i> Sars, 1882	Norway, North Atlantic (Sars, 1897; Kussakin, 1982).
<i>A. axeli</i> Castelló, 1992	Western Mediterranean (Castelló, 1997).
<i>A. bispinata</i> (Menzies & Kruczynski, 1983)	Gulf of Mexico.
<i>A. bocagei</i> Nobre, 1903	Portugal.
<i>A. bonnierii</i> Stephensen, 1915	Straits of Gibraltar; Mediterranean.
<i>A. caeca</i> Benedict, 1898	North Atlantic (Richardson, 1905; Schultz, 1969; Kussakin, 1982).
<i>A. cinguicula</i> Castelló & Carballo, 2000	Western Mediterranean.
<i>A. corniger</i> (Stebbing, 1873)	South Africa (Barnard, 1914, 1920; Kensley, 1978, 1984).
Junior synonyms include: <i>Antarcturus ornatus</i> , Tattersall, 1913; <i>Astacilla setosa</i> Vanhöffen, 1914; <i>Arcturoopsis hirsutus</i> Barnard, 1914; <i>Arcturoopsis hirsutus</i> subglaber Barnard, 1914.	
<i>A. cymodocea</i> Menzies & Glynn, 1968	Gulf of Mexico, Caribbean (Kensley & Schotte, 1989).
<i>A. depressa</i> Castelló & Poore, 1998	western Mediterranean.
<i>A. deshayesii</i> Lucas, 1849	Europe
<i>A. eminentia</i> Kensley, 1984	South Africa.
<i>A. gibbossa</i> Pillai, 1954	India (Pillai, 1963).
<i>A. glabrus</i> (Benedict, 1898)	Bering Sea, north western Pacific Ocean (Richardson, 1899, 1905, 1909; Birstein, 1963; Schultz, 1969; Kussakin, 1982).
<i>A. gorgonophila</i> Monod, 1925	north Africa; Mediterranean.
<i>A. granulata</i> (Sars, 1877)	eastern USA, United Kingdom, Norwegian Sea (Harger, 1880; Benedict, 1898; Richardson, 1905; Schultz, 1969; Kussakin, 1982). Junior synonym: <i>Astacilla americana</i> Harger, 1878.
<i>A. intermedia</i> (Goodsir, 1841)	United Kingdom, Scandinavia (Kussakin, 1982). Junior synonym: <i>Arcturus affinis</i> Sars, 1869.
<i>A. laevis</i> Castelló & Poore, 1998	western Mediterranean.
<i>A. lauffi</i> Menzies & Frankberg, 1966	Texas; Georgia (Schultz, 1969; Clark & Robertson, 1982).
<i>A. lasallae</i> Paul & Menzies, 1971	Venezuela (Kensley & Schotte, 1989).
<i>A. longicornis</i> (Sowerby, 1805) Junior synonyms include: <i>Leacia lacertosa</i> Johnston, 1825; <i>Leachia gracilis</i> Goodsir, 1841; <i>Arcturus deshayesii</i> Lucas, 1849; <i>Arcturus linearis</i> Stebbing, 1878.	United Kingdom; Scandinavia (Sars, 1897; Stephensen, 1948; Gruner, 1965; Naylor, 1972; Kussakin, 1982).
<i>A. longispina</i> (Kensley, 1978)	South Africa (Kensley, 1978; Kensley, 1984).
<i>A. marna</i> Kensley & Schotte, 1994	Dominica.
<i>A. mediterranea</i> Koehler, 1911	Mediterranean (Barnard, 1920; Kensley, 1984; Kensley, 1978; Kensley, 1984).
<i>A. monodi</i> Tattersall, 1925	Mauritania.
<i>A. paucisetosa</i> Castelló & Caballo, 2000	Western Mediterranean.
<i>A. pusilla</i> (Sars, 1873)	Scandinavia (Sars, 1897; Kussakin, 1982).
<i>A. serrata</i> Nunomura, 1998	Japan.
<i>A. spinata</i> (Menzies & Kruczynski, 1983) Junior synonym: <i>A. regina</i> Kensley, 1984; (Kensley & Schotte, 1989).	Belize, Barbados, St Lucia.
<i>A. tayronae</i> Müller, 1993	Columbia
<i>A. tranquilla</i> (Kensley, 1975)	South Africa (Kensley, 1978, 1984).
<b>Species transferred to other genera:</b>	
<i>A. anophthalmus</i> Birstein, 1963	<i>Arcturus</i> .
<i>A. attentuata</i> Hale, 1946	<i>Neastacilla</i> .
<i>A. bacillus</i> Barnard, 1920	<i>Neastacilla</i> .
<i>A. californica</i> Boone, 1918	<i>Neastacilla</i> .
<i>A. dilatata</i> Richardson, 1909	preoccupied, replacement name: <i>Neastacilla richardsonae</i> .
<i>A. dilatata</i> Sars, 1882	type species of <i>Arcturella</i> .
<i>A. diomedae</i> Benedict, 1898	<i>Neastacilla</i> .
<i>A. estadoensis</i> Schultz, 1981	<i>Neastacilla</i> .
<i>A. falclandica</i> Ohlin, 1901	type species of <i>Neastacilla</i> .
<i>A. fusiformis</i> Hale, 1946	<i>Neastacilla</i> .

Table 1 (cont.)

Species	Distribution
<i>A. giardi</i> Bonnier, 1896	<i>Arcturopsis</i> .
<i>A. inaequispinosa</i> Guiler, 1949	<i>Neastacilla</i> .
<i>A. kerguelensis</i> Vanhöffen, 1914	<i>Neastacilla</i> .
<i>A. levis</i> Thomson & Anderton, 1921	<i>Neastacilla</i> .
<i>A. macilenta</i> Hale, 1946	<i>Neastacilla</i> .
<i>A. magellanica</i> Ohlin, 1901	<i>Neastacilla</i> .
<i>A. marionensis</i> Beddard, 1886	<i>Neastacilla</i> .
<i>A. monoseta</i> Guiler, 1949	<i>Neastacilla</i> .
<i>A. polita</i> Gurjanova, 1936	<i>Neastacilla</i> .
<i>A. setosa</i> Vanhöffen, 1914	junior synonym of <i>Astacilla corniger</i> .
<i>A. sheardi</i> Hale, 1946	<i>Neastacilla</i> .
<i>A. vicaria</i> Hale, 1946	<i>Neastacilla</i> .
<i>Leachia gracilis</i> Goodsir, 1841	junior synonym of <i>Astacilla longicornis</i> .
<i>L. granulata</i> Sars, 1877	<i>Astacilla</i> .
<i>L. intermedia</i> Goodsir, 1841	<i>Astacilla</i> .
<i>L. nodosa</i> Dana, 1849	<i>Arcturella</i> .
<i>Leacia lacertosa</i> Johnston, 1825	junior synonym of <i>Astacilla intermedia</i> .

#### Type species

*Oniscus longicornis* Sowerby, 1805, subsequent designation by Fowler, 1912 (ICZN, 1986: Opinion 1369).

#### Diagnosis

Body strongly geniculate between pereonites 4 and 5, cylindrical, with no dorsal ridge. Antenna 2 slender, 2 or 3 flagellar articles, flagellum ending with claw. Maxillipedal palp with all segments free, palp article 3 similar width as article 2. Pereonite 4 elongate, male and female of similar length.

Pereopod 1 setose, with unguis (rarely without). Pereopods 2–4 present and functional as setose appendages, dactylus absent. Pereopods 2–4 cylindrical, held close to the mouthparts, able to extend past the mouthparts, setae in rows, setae as long as segment, flexion between the carpus and the propodus present. Pereopods 5–7 with 2 unguis, secondary unguis robust.

Penial plate narrow, apex simple. Male pleopod 1 with lateral notch, with 3 lateral setae, lateral setae of similar length. Male pleopod 2 with appendix masculina tapered, without ridge, distally not widened, straight, elongated.

Uropodal exopod present, 3 setae on inner ramus.

Female functional brood pouch majority made up of pereonite 4 (or 3 and 4). Oostegite 5 absent.

#### Remarks

The systematics of the genus *Astacilla* is currently unresolved, given the similarities of many species to those in the genera *Arcturella* and *Neastacilla*. In the past *Astacilla* has been successively defined by

an elongate fifth pereonite (Bate & Westwood, 1868), pereopod 1 with a claw, one pair of oostegites (Sars, 1897), and the loss of dactyls on pereopods 2 to 4 (Kussakin, 1972). Examination of several species of *Astacilla* has shown that these characters are variable and in the case of the number of oostegites, incorrect. The lack of known synapomorphic character states for many arcturid genera combined with the reluctance of many authors to examine the higher systematics of the genera has led to much confusion within arcturid taxonomy and much debate (Nordenstam, 1933; Monod, 1970; Kussakin, 1972; Schultz, 1981; Menzies & Kruczynski, 1983; Wägele, 1989; Brandt, 1991).

#### *Astacilla lewtonae* sp. nov.

Figures 1–4


#### Material examined

##### *Holotype*

Male, 7.3 mm, Western Australia, between Dampier and Port Hedland, 19°03.00'S; 119°00.00'E, 80 m, WHOI epibenthic sled, CSIRO division of Fisheries, 11 Dec 1982 (WAM C 32344).

##### *Paratypes*

Female, 6.0 mm, Western Australia, between Dampier and Port Hedland, 19°37.00'S; 118°53.00'E, 30 m, WHOI epibenthic sled, CSIRO division of Fisheries, 3 Jun 1983 (WAM C 32345). 2 females; 7.0–7.5 mm, 19°28.40'S; 118°55.10'E, 38 m, WHOI epibenthic sled, CSIRO division of Fisheries, 25 Oct 1983 (WAM C 32346). Immature male, 5 mm,



**Figure 1** *Astacilla lewtonae* n.sp., female holotype (NMV J16900): a, lateral view; b, dorsal view. Male (NMV J16904): c, lateral view; d, dorsal view. Scale = 1.0 mm.

19°29.90'S; 118°52.00'E, 37 m, WHOI epibenthic sled, CSIRO division of Fisheries, 24 Oct 1983 (NMV J16651). Manca 2, 3 mm, 19°50.00'S; 118°57.90'E, 84 m, WHOI epibenthic sled, CSIRO division of Fisheries, 29 Aug 1983 (NMV J16667). 3 females; 6.5–7.0 mm, 19°29.60'S; 118°51.70'E, 40 m, WHOI epibenthic sled, CSIRO division of Fisheries, 25 Oct 1983 (NMV J16648).

*Other material examined*

**Northern Territory**, North West end, Bommies, McCluer Island, 11°02.00'S; 132°58.00'E, 8 m, SCUBA, in sponges, Lowry, J.K., 16 Oct 1982 (NMV

J16582). 11°02.00'S; 132°58.00'E, 8 m, SCUBA, in hydroids, Lowry, J.K., 16 Oct 1982 (NMV J16935).

**South Australia**, Spencer Gulf, Douglas Bank, 32°47.18'S; 137°50.00'E, 15 m, South Australia Fisheries, Aug 1986 (NMV J16917).

**Description**

*Female*

Body geniculate and cylindrical. Anterolateral margins of head sub-truncate with medial indentation, a small rostral point evident. Fusion of head and pereonite 1 indicated by dorsolateral groove incised laterally. Head heavily tuberculate,

with 2 tuberculate dorsal elevations central and posterior to eyes, anterior tubercle with 2 apices. Pereonite 2 wider than pereonite 1, tuberculate with angular lateral margins extended. Pereonite 3 wider than pereonite 2, tuberculate with angular lateral margins extended. Pereonite 4 about 6 times longer than pereonite 3; dorsally wider than pereonite 3, tuberculate, angular lateral margins extended, with small posterior lateral extensions. Pereonites 5 to 7 progressively shorter posteriorly; heavily tuberculate, with lateral extensions. Pleon with evidence of three fused pleonites plus plotelson; total length longer than combined lengths of pereonites 5 to 7, with small anterior lateral wings, posterior angular lateral wings and rounded apex.

Eyes round, dorsolateral. Antenna 1 reaching past distal edge of second peduncular article of antenna 2; flagellum slender with aesthetascs attached distally. Antenna 2 slender, more than half as long as body; flagellum of 2 articles plus claw, lower margin without scales.

Maxilla 1 inner lobe with 3 terminal setae; outer lobe with 10 robust setae. Maxilla 2 inner lobe with 15 plumose setae; middle lobe with 4 setae; outer lobe with 3 setae. Maxillipedal endite with 10 mesial setae; palp article 2 and 3 with mesial setal rows; article 4 with mesial and lateral setal rows; article 5 with distal setae.

Pereopod 1 propodus smaller than carpus; dactylus about twice as long as wide, bearing unguis, which is longer than dactylus. Pereopods 2–4 merus to propodus with paired rows of long setae; dactylus absent; flexion between carpus and propodus present.

Pereopods 5 to 7 progressively shorter; dactylus denticulate, unguis present with secondary unguis 2/3 length of primary unguis.

Uropod exopod not reaching mid point of endopod, with 2 setae of subequal length.

Oostegites present on pereopods 1 to 4; oostegite 4 thickened, with transverse suture delimiting posterior lobe.




Figure 2 *Astacilla lewtonae* n.sp., female holotype (NMV J16900): left maxilliped; left maxillae 1 and 2; left mandible; antennae 1 and 2. Scales: a (A2) = 0.5 mm; b (A1, ♂ A1) = 0.5 mm; c (MP, MX1, MX2, MD) = 0.5 mm.




Figure 3 *Astacilla lewtonae* n.sp., female holotype (NMV J16900): pereopods 1 to 7. Scale = 0.5 mm.

#### Male

Body geniculate and cylindrical. Anterolateral lobes of head rounded with tuberculations, small rostral point evident. Head and pereonite 1 fused and with similar tuberculate ornamentation to female. Pereonite 2 and 3 tuberculate; lateral margins not greatly extended. Pereonite 4 around 8 times length of pereonite 3, tuberculate, lateral

margins not extended. Pereonites 5 to 7 tuberculate, anterolateral margins extended. Pleon length greater than combined lengths of pereonites 5 to 7, evidence of 3 fused pleonites plus pleotelson, with small anterior lateral wings, angular posterior lateral wings and rounded apex.

Eyes round and dorsolateral. Antenna 1 extending past distal edge of second peduncular

article of antenna 2; flagellum uniaarticulate and with aesthetascs attached distally and laterally along the entire length. Antenna 2 as for female.

Mouthparts as for female.

Pereopods as for female.

Pleopod 1 exopod with lateral notch and 3 plumose setae of similar lengths on posterior face. Pleopod 2 with appendix masculina straight and extending twice the length of the endopod, tapering to two filaments. Penial plate straight and simple.

**Distribution**

Australia: Western Australia, Northern Territory, South Australia; 8 to 84 metres.

**Remarks**

Although morphologically similar to Australian *Neastacilla* species, the possession of a long, straight appendix masculina is the primary reason why this species belongs in *Astacilla*. Other characteristics that support the placement of this species within *Astacilla* are the possession of flexion between the carpus and propodus and the complete lack of dactyli on pereopods 2 to 4, the presence of a dactylus on pereopod 1 and the shape of the female, which is widened and not especially elongate at pereonite 4. The dorsal and lateral sculpture of the body of this species makes it unique among the *Astacilla* species.




Figure 4 *Astacilla lewtonae* n.sp., male (NMV J16904): pleopods 1 and 2; penial plate. Female holotype (NMV J16900): distal end of uropod; a, ventral view with oostegites. Scales: a (PL1, PL2, Pe) = 0.5 mm; b (U) = 0.5 mm; c (e) = 1.0 mm.

### Etymology

This species is named for Helen Lew Ton who detected this 'strange' species in the collections of Museum Victoria.

### ACKNOWLEDGEMENTS

This work was conducted as part of a postgraduate degree at the Zoology Department, University of Melbourne and Museum Victoria, funded by an Australian Postgraduate Award. I am grateful to Dr Gary Poore (Museum Victoria) for access to the collections, advice and encouragement and to Professor Alan Myers (University College Cork) for advice on the manuscript.

### REFERENCES

- Bate, C. S. and J. O. Westwood. (1868). *A history of the British sessile-eyed Crustacea*. Volume 2. John Van Voorst, London. 536 pp.
- Beddard, F. E. (1886). Report on the Isopoda collected by HMS Challenger during the years 1873–76. Part 2. *Report of the Voyage of HMS Challenger* 17: 1–178.
- Brandt, A. (1991). Zur Besiedlungsgeschichte des antarktischen Schelfes am Beispiel der Isopoda (Crustacea, Malacostraca). *Berichte zur Polarforschung* 98: 1–240.
- Cordiner, C. (1793). *Remarkable ruins, and romantic prospects, of North Britain. With ancient monuments, and singular subjects of natural history*. Peter Mazell, London. 96 plates with letterpress.
- Dana, J. D. (1849). *Conspectus crustaceorum quae in orbis terrarum circumnavigatione, Carolo Wilkes e classe Reipublicae, Foederate Duce, lexit et descripsit*. *American Journal of Sciences and Arts* (2) 8: 424–428.
- Guiler, E. R. (1949). New species of *Astacilla* from Tasmanian waters. *Papers and Proceedings of the Royal Society of Tasmania* 1948: 45–64.
- Hale, H. M. (1924). Notes on Australian Crustacea. No. 3. *Transactions of the Royal Society of South Australia* 48: 209–225.
- Hale, H. M. (1946). Isopoda – Valvifera. *British, Australian and New Zealand Antarctic Research Expedition, 1929–1931. Reports-Series B (Zoology and Botany)* 5: 161–212.
- Haswell, W. A. (1881). On some new Australian marine Isopoda – Part II. *Proceedings of the Linnean Society of New South Wales* 6: 181–196, pls 3, 4.
- Kensley, B. (1983). *Astacilla* Cordiner, 1793 (Crustacea, Isopoda): proposed nomenclatural validation by use of plenary powers Z.N.(S.) 2319. *Bulletin of Zoological Nomenclature* 40: 163–164.
- King, R. A. (2000). Rediagnosis of the endemic southern Australian genus *Parastacilla* Hale, 1924 (Crustacea: Isopoda: Arcturidae) with descriptions of two new species. *Memoirs of the Museum of Victoria* 58: 125–136.
- King, R. A. (2001). *The systematics, evolution and biogeography of the Arcturidae (Crustacea, Isopoda)*. PhD thesis. University of Melbourne: Melbourne. 253 pp.
- Kussakin, O. G. (1972). Isopoda from the coastal zone of the Kurile Islands. III. Three new arcturids from the Middle Kuriles with taxonomic remarks on the family Arcturidae. *Crustaceana Supplement* 3: 178–189.
- Lew Ton, H. M. (1980). *Reassessment of the genus Neastacilla Tattersall (Crustacea: Isopoda: Arcturidae) and a revision of the Australasian species*. B.Sc. (Hons) thesis, Monash University, unpublished.
- Menzies, R. J. and W. L. Kruczynski. (1983). Isopod Crustacea (exclusive of Epicaridea). *Memoirs of the Hourglass Cruises* 6: 1–126.
- Monod, T. (1970). Sur quelques isopodes marins d'Australie I. Arcturidae. *Bulletin du Muséum National d'Histoire Naturelle, Paris* (2) 42: 1127–1142.
- Nordenstam, A. (1933). Marine Isopoda of the families Serolidae, Idotheidae, Pseudidotheidae, Arcturidae, Parasellidae and Stenetriidae mainly from the South Atlantic. *Further Zoological Results of the Swedish Antarctic Expedition, 1901–1903* 3: 1–284, 2 pls, errata.
- Poore, G. C. B. (2001). Isopoda Valvifera: diagnoses and relationships of the families. *Journal of Crustacean Biology* 21: 213–238.
- Poore, G. C. B., Lew Ton, H. M. and Bardsley, T. M. (2002). Suborder Valvifera Sars, 1882. in: Houston, W.W.K., and Beesley, P. (eds) *Zoological Catalogue of Australia*. 19.2a. Crustacea: Malacostraca: Syncarida, Peracarida: Isopoda, Tanaidacea, Mictacea, Thermosbaenacea, Spelaeogriffacea. CSIRO Publishing: Melbourne.
- Sars, G. O. (1897). Parts 5, 6. Idotheidae, Arcturidae, Asellidae, Ianiridae, Munnidae. Pp. 81–116, pls 33–48. In: *An account of the Crustacea of Norway with short descriptions and figures of all the species*. Vol. 2 Isopoda. Bergen Museum, Bergen.
- Schultz, G. A. (1981). Arcturidae from the Antarctic and Southern Seas (Isopoda, Valvifera) Part I. Pp. 63–94. In: *Biology of the Antarctic Seas* 10. Antarctic Research Series. Vol. 32. American Geophysical Union.
- Stebbing, T. R. R. (1905). Report on the Isopoda collected by Professor Herdman at Ceylon, in 1902. *Report to the Government of Ceylon on the Pearl Oyster Fisheries of the Gulf of Manaar, Supplementary Report* 4: 1–64, pls 61–12.
- Sowerby, J. (1805). *The British Miscellany*, part 4. London.
- Tattersall, W. M. (1921). Crustacea. Part VI. – Tanaidacea and Isopoda. *British Antarctic "Terra Nova" Expedition, Natural History Reports, Zoology* 3: 191–258, pls 191–111.
- Vanhöffen, E. (1914). Die Isopoden der Deutschen Südpolar Expedition 1901–1903. *Deutsche Südpolar Expedition 1901–1903* 25 (Zoologie) 7: 447–598.
- Wägele, J. W. (1989). Evolution and phylogenetic System der Isopoda. Stand der Forschung und neue Erkenntnisse. *Zoologica (Stuttgart)* 140: 1–262.
- Whitelegge, T. (1904). Scientific results of the trawling expedition of H.M.C.S. "Thetis" off the coast of New South Wales in February and March, 1898. Crustacea. Part IV. Isopoda. Part III. *Memoirs of the Australian Museum* 4: 405–416.