

# Discovery of the South African polyplacophoran *Stenosemus simplicissimus* (Thiele, 1906) (Mollusca, Polyplacophora, Ischnochitonidae) in the Southern Ocean

Enrico Schwabe

Zoologische Staatssammlung München, Münchhausenstrasse 21, D-81247 München, Germany, enrico.schwabe@zsm.mwn.de

**Abstract:** Recent expeditions to the Atlantic sector of the Southern Ocean have yielded valuable collections of shelf and deep water polyplacophorans. These included several specimens of *Stenosemus simplicissimus* (Thiele, 1906), a species previously known only by its holotype and type locality at the Cape of Good Hope. The new material enabled a thorough morphological redescription of the species by studying valve, perinotum, and radula characters with SEM. The new records from Shag Rocks and the eastern Weddell Sea enlarge the species' biogeographic distribution from the temperate South African region to the polar South Georgia and Weddell Sea regions. Its bathymetric range is extended from 318 m to 285-1064 m. The limited occurrence of deep-water Antarctic polyplacophorans may be caused by benthic predators that limit the expansion of non-herbivorous chitons in the Antarctic deep sea.

**Key words:** Atlantic Ocean, Antarctica, zoogeography, distribution, new records

Antarctic waters support a small number of Polyplacophora in contrast to the highly diverse fauna of other marine molluscs (Thiele 1912, Dell 1990, Numanami 1996, Sirenko and Schrödl 2001). The species discovered to date are: *Lep-tochiton kerguelensis* Haddon, 1886, *Callochiton bouveti* Thiele, 1906, *Callochiton gaussae* Thiele, 1908, *Leloupia bel-gicae* (Pelseneer, 1903), *Stenosemus exaratus* (G. O. Sars, 1878), *Stenosemus simplicissimus* (Thiele, 1906), *Tonicina zs-chau* (Pfeffer in von Martens and Pfeffer, 1886), *Nuttallo-chiton mirandus* (E. A. Smith MS, Thiele, 1906), and *Hemi-athrum setulosum* Carpenter in Dall, 1876. With the exception of *S. simplicissimus*, all species are more or less well described in earlier revisions (e.g., Thiele 1906a, 1906b, 1908, Dell 1964, Kaas and Van Belle 1985a, 1985b, 1990, Götting 1993). The recent rediscoveries of this species, which was known only from the type material, enable a detailed morphological description using scanning electron microscopy (hereafter, SEM). The description will help non-chiton specialists to separate this species from similar representatives of the genus *Callochiton* Gray, 1847. In addition, deep water polyplacophorans from Antarctica are rare and our knowledge of their biology and habitat preference is limited. Analysis of abiotic parameters at a certain depth may help getting a better understanding of how chitons interact with their environment. The present paper deals with the unknown Antarctic deep-water chiton fauna.

and eastern Weddell Seas. The material was obtained by using Agassiz trawls (AGT), bottom trawls (BT), epibenthic sledges (EBS), and Rauschert dredges (RD). When the catch reached the deck, the samples were sieved through 500 µm or 1000 µm mesh, the remainder was fixed in 4% buffered formaldehyde or 75-96% ethanol and then sorted under stereomicroscopes. Most of the polyplacophorans were found attached to hard substrates, such as cobbles. Specimens were stored in ethanol for further morphological examinations. To confirm the identification of *S. simplicissimus*, the holotype was examined.

The type material is deposited at the Natural History Museum Berlin, Germany (ZMB). The newly collected material is deposited at the British Antarctic Survey, United Kingdom (BAS), the Zoological Institute St. Petersburg, Russia (ZISP), and the Bavarian State Collection of Zoology, Germany (ZSM).


Preparation of the specimens followed Schwabe and Ruthensteiner (2001). Specimens used for SEM were partly disarticulated, enabling examination of valves, perinotum, and radula. Microsculpture and radular photographs were made on a LEO 1430VP SEM (at the ZSM). Abiotic factors were established using a conductivity-temperature-depth (CTD) data logger, or by visual inspection of the substratum on board of the research vessel.

## MATERIALS AND METHODS

Specimens of *Stenosemus simplicissimus* were collected during recent expeditions with R/V *Polarstern* (ANT XIII/3, ANT XVII/3, ANT XXI/2, and ANT XXII/3) to the Scotia

## SYSTEMATICS

Class Polyplacophora Gray, 1821  
Subclass Neoloricata Bergenhayn, 1955  
Order Chitonida Thiele, 1910  
Family Ischnochitonidae Dall, 1889



**Figure 1.** Light micrographs of the holotype of *Stenosemus simplicissimus*, ZMB Moll 59908. A, valves ii to iv *in situ*, anterior at right; B, detail of fig. 1A, showing the dorsal perinotum scales; C, valves (from left to right) viii, v, i in dorsal view, anterior at right; D, right lateral view of the tail valve, anterior at right. Scale bars: A, 5 mm; B, D, 1 mm; C, 500  $\mu$ m.

Genus *Stenosemus* von Middendorff, 1847

*Type species*: *Chiton albus* Linnaeus, 1767, by subsequent designation, Winckworth (1926: 15)

*Stenosemus simplicissimus* (Thiele, 1906)  
(Figs. 1-5)

*Ischnochiton (Chondropleura) simplicissimus* Thiele  
1906b: 335, pl. 29, figs 21-25.

Additions to the bibliography in Kaas and Van Belle (1990: 67):

*Ischnochiton simplicissimus*; Barnard 1974: 740.

*Ischnochiton (Stenosemus) simplicissimus*; Kaas and Van Belle 1980: 120; 1990: 67, fig. 27; 1998: 171.

*Ischnochiton (Chondropleura) simplicissimus*; Kiliyas 1995: 169.

*Stenosemus simplicissimus*; Sirenko 1994: 164; 2005: 36; Gutt *et al.* 2000: 40; Linse *et al.* 2006: 155 (partim).

*Type material*: ZMB Moll 59908 (partly disarticulated holotype) (Figs. 1A-D).

*Type locality*: South Africa, Cape of Good Hope, Deutsche Tiefsee-Expedition St. 113: 34°33.3'S 18°21.2'E, 318 m.

#### Additional material examined

ZSM Mol 20050857 (1 specimen - 3.4  $\times$  2.2 mm, partly disarticulated) (Figs. 2-3), Antarctica, Weddell Sea: ANT XXII-3 (ANDEEP III) St. PS 67/074-7: 71°18.60'S 13°59.11'W-71°18.38'S 13°58.17'W, 1047-1064 m, on rock (quartz-amphibolite gneiss), laying on a sandy sediment (only 5 cm thick) (salinity: 34.7 psu; water temperature: 0.5 °C; O<sub>2</sub> concentration: 5.8 ml/l; pressure: 1016.5 dBar; all data from 1004 m, measured by CTD), AGT, collected by J. M. Bohn and E. Schwabe, 20 February 2005, preserved in 96% ethanol.

ZSM Mol 20020914 (1 specimen - width 3.1 mm [curled]), Shag Rocks, South Georgia and South Sandwich Islands: ANT XIX-5 (LAMPOS) St. PS 61/169-1: 53°22.94'S 42°41.37'W-53°22.89'S 42°41.50'W,

284.3 m, RD, collected by Dr. M. Schrödl, 10 April 2002, preserved in 78% ethanol.


ZSM Mol 20008502 (1 specimen - 8.7  $\times$  4.2 mm, partly disarticulated) (Fig. 4), Antarctica, Weddell Sea: ANT XVII-3 (EASIZ III) St. 97-1: 71°06.27'S 12°50.46'W-71°06.24'S 12°49.92'W, 728-743 m, EBS, collected by Dr. M. Schrödl, 3 April 2000, preserved in 78% ethanol.

BAS (Dr. Katrin Linse) 03-802 (1 specimen - width 3.8 mm [curled]), Antarctica, Weddell Sea: ANT XXI-2 (BENDEX) St. PS 65/324-1: 72°54.52'S 19°47.74'W-72°54.55'S 19°47.30'W, 647.2-693.6 m, RD, collected by Dr. K. Linse, 3 January 2004, preserved in 96% ethanol.

BAS (Dr. K. Linse) 03-769 (1 specimen - width 3.4 mm [curled]), Antarctica, Weddell Sea: ANT XXI-2 (BENDEX) St. PS 65/297-1: 72°48.50'S 19°31.66'W-72°48.65'S 19°31.85'W, 630.8-668 m, RD, collected by Dr. K. Linse, 1 January 2004, preserved in 96% ethanol.

#### Description

Species moderately large, up to 16 mm (the largest specimen is the holotype). It is elongate, oval, with a cari-



**Figure 2.** *Stenosemus simplicissimus* (ZSM Mol 20050857), 3.4 × 2.2 mm. A-B, light micrographs, C-F, scanning electron micrographs. A, dorsal view of the complete specimen, anterior at right; B, ventral view of the complete specimen, anterior at right; C, dorsal view of the head valve; D, dorsal view of valve ii; E, dorsal view of the tail valve; F, left lateral view of the tail valve, anterior at left. Scale bars A-B: 1 mm, C-F: 500  $\mu$ m.

nated, moderately high-elevated dorsum. Dorsal elevation quotient (height/width) (of isolated valve v of the holotype): 0.51. Color of tegmentum and perinotum uniform dull white.

Tegmentum virtually smooth, except for commarginal growth marks, which occur on all valves (Figs. 1A-D, 2C-F, 4A-D) and a micro-perforation. In earlier growth stages, fine radial striation is visible in apical regions and the middle of the first valve. The head valve has a wide V-shaped posterior margin and is clearly notched in the middle (Figs. 1C, 2C, 4A). Intermediate valves (Figs. 1A, 1C, 2D, 4B)


(valve ii) to rectangular, with short and rounded side margins, and straight to slightly concave posterior margins (on both sides of the slightly protruding apex). Anterior valve margin is convex. Lateral areas are clearly elevated. Tail valve (Figs. 1C-D, 2E-F, 4C-D) is semicircular with an anteriorly directed, weakly elevated mucro that is situated in the two anterior thirds of the valve length. Postmucronal slope is steep and straight.

Articulamentum is thin and white. Apophyses (Figs. 1A, 1C-D, 2D-E, 4B-D) are well developed, rather short, wide, and medially connected by a short smooth jugal lamina. Apophyses are triangular in intermediate valves, and rectangular in tail valve. Slit formula varies from 14/1/10 (holotype) to 16/1-2/13 (8.7 mm long specimen, ZSM Mol 20008502). Slits are wide and rather long, teeth edges are slightly thickened and faintly crenulated. Slit rays are present in all valves. Eaves are spongy.

Perinotum is rather narrow (Figs. 1A-B), dorsally covered with juxtaposed, bent, conical, round-topped, and inwardly directed calcareous scales, 160-176  $\mu$ m long, 144-150  $\mu$ m in diameter on mid-perinotum, rhomboidal at the base, and weakly longitudinally striated (Kaas and Van Belle 1990). Towards the outer margin, the scales are smaller and measure 80-90 × 50-57  $\mu$ m (Fig. 3B). Marginal fringe (Fig. 3C) consists of straight, obtusely-pointed solid spicules, which are medially keeled and measure about 110 × 22  $\mu$ m. Among them are smaller,

straight, and sharp spicules, 50  $\mu$ m in length and 10  $\mu$ m in width. These spicules are situated distally either on long, very slender shafts (140 × 7  $\mu$ m) or in shorter tubs that may attain a diameter of 15  $\mu$ m (Fig. 4E). Ventrally there are radial rows of rectangular scales that measure 53 × 27  $\mu$ m (Fig. 3D).

Radula (Fig. 3D) of a partially disarticulated specimen (ZSM Mol 20008502) measures 3.3 mm, of which 1.6 mm were taken up by the radula cartilage, with 44 teeth rows of which 31 are mineralized. Central tooth is rectangular, its distal end a little wider than the base. It measures 71 × 31  $\mu$ m long and has a forward-directed simple, slightly bent cusp.



**Figure 3.** Scanning electron micrographs of *Stenosemus simplicissimus* (ZSM Mol 20050857), 3.4 × 2.2 mm. A, detail of Fig. 2F, showing the valve microsculpture; B, dorsal perinotum scales at the margin; C, marginal perinotum fringe; D, ventral perinotum scales. Scale bars: 50  $\mu$ m.

First lateral tooth is wing-shaped and covers the lower two thirds of central tooth. It may attain a length of 60  $\mu$ m and has a simple inward directed small cusp. Total length of second lateral tooth is 175  $\mu$ m; one third of its length is taken up by the squarish head with the single, sharply pointed, and inwardly curved elongate denticle. Shaft of second lateral tooth is sharply keeled on its inner side and slightly curved in the upper half. Basally the tooth is wing-shaped. First uncinial tooth is triangular in outline with a straight inner edge and a steep slightly convex, outer edge. Second uncinial tooth is very broad and S-shaped. Third uncinial tooth is extremely slender, measuring 138 × 18  $\mu$ m, with a spoon-shaped distal extension, which may attain the double width of the shaft. First marginal tooth is similar to, but slightly more slender than, the second uncinial. Second marginal tooth is arrowhead-shaped, measuring 71 × 31  $\mu$ m. Third marginal tooth is rectangular in outline, 75  $\mu$ m long, and 46  $\mu$ m wide. Its inner edge is thickened.

Ctenidia are arranged merobranchially with the longest ctenidium on each side positioned the fourth from the posterior end. Size and number of ctenidia depend on animal size. The 8.7 mm long specimen has 16 ctenidia on the right and 17 on the left side; the juvenile (3.4 mm) has 7 ctenidia on each side of the foot (Fig. 2B).

### Distribution

Known from the Cape of Good Hope (type locality), Shag Rocks near South Georgia Island and eastern Weddell Sea (this study) (Fig. 5). Bathymetrically the species lives between 284-1064 m depth (Thiele 1906b, Gutt *et al.* 2000, Sirenko and Schrödl 2001, Linse *et al.* 2006).

### DISCUSSION


Since its description, *Stenosemus simplicissimus* was never recollected. Several expeditions to the Subantarctic Marion and Prince Edward Islands failed to locate this species, although the congeneric *Stenosemus exaratus* (G. O. Sars, 1878) [reported as *Stenosemus dorsuosus* (Haddon, 1886)] was found together with the following species: *Leptochiton kerguelensis* Haddon, 1886, *Hemiathrum setulosum* Carpenter in Dall, 1876, and *Placiphorella* sp. (Branch *et al.* 1991). *Leptochiton kerguelensis* and *H. setulosum* are typical faunistic elements of the Antarctica;

the latter is *Placiphorella atlantica* (Verrill and S. I. Smith in Verrill, 1882) (pers. obs.).

*Stenosemus simplicissimus* was first rediscovered during the ANT XIII-3 (EASIZ I) – Antarctic expedition at St. 01: 71°03.10'S 11°25.50'W, at 462 m (Gutt *et al.* 2000). During subsequent expeditions, the species was found again but only in small numbers.

The Weddell Sea supports a high diversity of mollusc species and, not surprisingly, their numbers decrease with increasing depth, along with a significant decrease in total biomass (Brey and Gerdes 1997, 1998). During the ANDEEP III expedition (from the Cape Basin to Kapp Norvegia and across the Weddell Sea), 186 mollusc morphospecies (3801 specimens from 12 EBS and 19 AGT stations) were collected from depths ranging from 1000 to 4900 m (Linse *et al.* 2006).

In contrast to former expeditions in this area of comparable scope (Sirenko and Schrödl 2001), only two polyplacophoran specimens were found, the herein mentioned juvenile of *Stenosemus simplicissimus* and *Leptochiton kerguelensis* (ZSM Mol 20060001). The latter specimen was collected at station PS67/133-2 (62°46'44"S 53°02'34"W-62°46'20"S 53°04'08"W) at 1581-1582 m depth. This is the maximum depth reported for this species, which was for-



**Figure 4.** Scanning electron micrographs of *Stenosemus simplicissimus* (ZSM Mol 20008502),  $8.7 \times 4.2$  mm. A, dorsal view of the head valve; B, dorsal view of valve ii; C, dorsal view of the tail valve; D, right lateral view of the tail valve, anterior at right; E, dorsal perinotum scales, close to the margin; F, rows 3-8 of the radula. Scale bars A-D: 500  $\mu$ m, E-F: 100  $\mu$ m.


merly only known from 1335 m in the Ross Sea (Dell 1990). A bulk of stones was found during the recent expedition, a substratum that allows settlement of chiton larvae. That they could be covered by sediment can be excluded because Paul (1976) has shown that chitons are able to remove sediment layers. Is the extremely low chiton diversity and density in the Antarctic merely a function of the great depth or related abiotic environmental conditions? Temperature is unlikely to be a problem as it does not change with depth (Brey and Gerdes 1998). The increase in water pressure can also be ruled out as diverse deep sea chiton faunas are known from other regions (Sirenko 2001). Most deep water chitons are

dependent on plant remains (e.g., sunken wood) that may not found in the Antarctic deep sea. The high organic food input (Brey and Gerdes 1997) in the Weddell Sea, together with abiotic conditions should favor colonization by non-herbivorous chitons. Several carnivorous taxa such as asteroids, ophiuroids, and polychaetes are highly adapted to the conditions of the Weddell Sea (Brey and Gerdes 1997, 1998). The chitons may lack the ability to co-occur with abundant predators such as the asteroids, which feed on chitons (Seiff 1975).

In summary, this report gives a detailed description of *Stenosemus simplicissimus*, contributes the first record of the species' juvenile stage, and extends its bathymetric range and that of *Leptochiton kerguelensis*, thus providing a further piece of the unfinished puzzle documenting Antarctic deep sea communities. The material obtained to date for *S. simplicissimus* suggests that it *could be* geographically restricted to the Southern Atlantic Ocean.

#### ACKNOWLEDGMENTS

I thank Olaf Klatt (Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany), Dr. Lawrence W. Carpenter (School of Marine Science, Virginia Institute of Marine Science, Virginia, USA), Rob Middag (Koninklijk Nederlands Instituut voor Onderzoek der Zee, Den Burg, The Netherlands), Dr. Michael Thompson (School of Earth Sciences, University of Leeds, Leeds, UK), and Dr. John A. Howe (Scottish Association for Marine Science, Argyll, UK) for providing physical and chemical data of the ANDEEP III station. All involved sorting teams and the crew of R/V *Polarstern* are warmly thanked for their professional work. Dr. Katrin Linse (BAS) is especially acknowledged for providing the material reported herein and for helpful comments on an earlier version of the manuscript. Dr. Douglas Eernisse (California State University, Fullerton, USA) kindly corrected the English and provided supportive remarks. Thanks are also due to Dr. Matthias Glaubrecht and Frank



**Figure 5.** The geographical distribution of *Stenosemus simplicissimus*.

Köhler (ZMB) for the loan of type material. I especially thank two anonymous reviewers for their helpful comments and constructive criticism. The authors participation to ANDEEP III was kindly supported by GeoBioCenterLMU. This is ANDEEP contribution No. 61.

#### LITERATURE CITED

- Barnard, K. H. 1974. Contributions to the knowledge of South African marine Mollusca. Part VII. Revised fauna list. *Annals of the South African Museum* 47: 663-781.
- Branch, M. L., P. M. Arnaud, J. Cantera, and D. Gianakouras. 1991. The benthic Mollusca and Brachiopoda of subantarctic Marion and Prince Edward Islands: 1) Illustrated keys to the species, 2) Records of the 1982-1989 University of Cape Town Surveys. *South African Journal of Antarctic Research* 21: 45-64.
- Brey, T. and D. Gerdes. 1997. Is Antarctic benthic biomass really higher than elsewhere? *Antarctic Sciences* 9: 266-267.
- Brey, T. and D. Gerdes. 1998. High Antarctic macrobenthic community production. *Journal of Experimental Marine Biology and Ecology* 231: 191-200.
- Dell, R. K. 1964. Antarctic and subantarctic Mollusca: Amphineura, Scaphopoda and Bivalvia. *Discovery Report* 33: 93-250.
- Dell, R. K. 1990. Antarctic Mollusca with special reference to the fauna of the Ross Sea. *Bulletin of the Royal Society of New Zealand, Wellington* 27: 1-311.
- Götting, K.-J. 1993. Polyplacophora des Weddell-Meeress und von King George Island, Süd-Shetlands, Antarktis. *Archiv für Molluskenkunde* 122: 171-187.
- Gutt, E., B. I. Sirenko, W. E. Arnt, I. S. Smirnov, and C. De Broyer. 2000. Biodiversity of the Weddell Sea: Macrozoobenthic species (demersal fish included) sampled during the expedition ANT XIII/3 (EASIZ I) with RV "Polarstern". *Berichte zur Polarforschung* 372: 1-103.
- Kaas, P. and R. A. Van Belle. 1980. *Catalogue of Living Chitons (Mollusca: Polyplacophora)*. Dr. W. Backhuys Publisher, Rotterdam.
- Kaas, P. and R. A. Van Belle. 1985a. *Monograph of Living Chitons (Mollusca: Polyplacophora) 1, Order Neoloricata: Lepidopleurina*. E. J. Brill/W. Backhuys, Leiden, The Netherlands.
- Kaas, P. and R. A. Van Belle. 1985b. *Monograph of Living Chitons (Mollusca: Polyplacophora) 2, Suborder Ischnochitonina, Ischnochitonidae: Schizoplacinae, Callochitoninae and Lepidochitoninae*. E. J. Brill/W. Backhuys, Leiden, The Netherlands.
- Kaas, P. and R. A. Van Belle. 1990. *Monograph of Living Chitons (Mollusca: Polyplacophora) 4, Suborder Ischnochitonina: Ischnochitonidae: Ischnochitoninae (continued). Additions to vols 1, 2 and 3*. E. J. Brill, Leiden, The Netherlands.
- Kaas, P. and R. A. Van Belle. 1998. *Catalogue of Living Chitons (Mollusca, Polyplacophora). 2nd, revised edition*. Backhuys Publishers, Leiden, The Netherlands.
- Kiliyas, R. 1995. Polyplacophora-Typen und -Typoide (Mollusca) im Zoologischen Museum in Berlin. *Mitteilungen aus dem Zoologischen Museum in Berlin* 71: 155-170.
- Linse, K., E. Schwabe, and A. Brandt. 2006. Mollusca collected during ANDEEP III - preliminary results. *Berichte zur Polar- und Meeresforschung* 533: 151-160.
- Numanami, H. 1996. Taxonomic study on Antarctic gastropods collected by Japanese Antarctic research expeditions. *Memoirs of the National Institute of Polar Research (E, Biology and Medical Science)* 39: 1-244.
- Paul, A. Z. 1976. Deep-sea bottom photographs show that benthic organisms remove sediment cover from manganese nodules. *Nature* 263: 50-51.
- Schwabe, E. and B. Ruthensteiner. 2001. *Callochiton schilfi* (Mollusca: Polyplacophora: Ischnochitonidae) a new species from Indonesian waters. *Vita Marina* 47: 175-184.
- Seiff, S. R. 1975. Predation upon subtidal *Tonicella lineata* of Mussel Point, California (Mollusca: Polyplacophora). *The Veliger* 18 (Supplement): 54-56.
- Sirenko, B. I. 1994. Chitons (Polyplacophora) of the continental slope of the Kurile Islands with a brief review of deep water species of the Russian Seas. *Russian Academy of Sciences Zoological Institute, Explorations of the Fauna of the Seas* 46: 159-174.
- Sirenko, B. I. 2001. Deep-sea chitons (Mollusca, Polyplacophora) from sunken wood off New Caledonia and Vanuatu. In: Bouchet, P. and Marshall, B. A., eds., *Tropical Deep-Sea Benthos, vol. 22. Mémoires du Muséum national d'Histoire naturelle* 185: 39-71.
- Sirenko, B. I. 2005. Unusual concentration of endemics of chitons (Polyplacophora) in bathyal zone near New Caledonia. In: Poster Abstracts of the IV. International Congress of the European Malacological Societies, Oct. 10-14. 2005 in Naples (Italy). *Notiziario S.I.M., Supplemento al Bollettino Malacologico* year 23: 36.
- Sirenko, B. I. and M. Schrödl. 2001. Mollusc biodiversity and ecol-

ogy. In: W. E. Arntz, and T. Brey, eds., *The Expedition ANTARKTIS XIII/3 (EASIZ I) of "Polarstern" to the eastern Weddell Sea in 1996. Berichte zur Polar- und Meeresforschung* **402**: 85-95.

Thiele, J. 1906a. Note sur les chitons de l'Expédition Antarctique du Dr. Charcot. *Bulletin du Museum d'Histoire naturelle* **12**: 549-550.

Thiele, J. 1906b. Ueber die Chitonen der deutschen Tiefsee-Expedition. *Wissenschaftliche Ergebnisse der deutschen Tiefsee Expedition* **9**: 325-336.

Thiele, J. 1908. Die antarktischen und subantarktischen Chitonen. *Deutsche Südpolar Expedition 1901-1903, 10 (Zoologie)* **2**: 9-23.

Thiele, J. 1912. Die antarktischen Schnecken und Muscheln. *Deutsche Südpolar Expedition 1901-1903, 13 (Zoologie)* **5**: 183-285.

Winckworth, R. 1926. Notes on British Mollusca. 1. *Journal of Conchology* **18**: 13-15.

**Accepted:** 24 July 2007; **final corrections received:**

27 November 2007