



1245
4890

HARVARD UNIVERSITY



Library of the
Museum of
Comparative Zoology

Bulletin of the Museum of Comparative Zoölogy
AT HARVARD COLLEGE
VOL. XC, No. 1

STUDIES OF NEOTROPICAL ANT-PLANTS
AND THEIR ANTS

BY WILLIAM MORTON WHEELER

WITH FIFTY-SEVEN PLATES

CAMBRIDGE, MASS., U. S. A.
PRINTED FOR THE MUSEUM
OCTOBER, 1942

PUBLICATIONS
OF THE
MUSEUM OF COMPARATIVE ZOÖLOGY
AT HARVARD COLLEGE

The BULLETIN and MEMOIRS are devoted to the publication of investigations by the Staff of the Museum or of reports by specialists upon the Museum collections or explorations.

Of the BULLETIN, Vols. I to LXXXIX and Vol. XCI, No. 1 and No. 2 have appeared and of the MEMOIRS, Vol. I to LVI.

These publications are issued in numbers at irregular intervals. Each number of the Bulletin and of the Memoirs is sold separately. A price list of the publications of the Museum will be sent upon application to the Director of the Museum of Comparative Zoölogy, Cambridge, Massachusetts.

Bulletin of the Museum of Comparative Zoölogy
AT HARVARD COLLEGE
VOL. XC, No. 1

STUDIES OF NEOTROPICAL ANT-PLANTS
AND THEIR ANTS

BY WILLIAM MORTON WHEELER

WITH FIFTY-SEVEN PLATES

CAMBRIDGE, MASS., U. S. A.
PRINTED FOR THE MUSEUM
OCTOBER, 1942

CONTENTS

	PAGE
Part I. The Neotropical Ant Plants.	
Chapter 1. The Myrmecophytes of the Genus <i>Cordia</i>	3
A. The <i>Cordias</i> of the Section <i>Physoclada</i>	4
B. The <i>Cordias</i> of the Section <i>Gerascanthus</i>	9
Chapter 2. Observations on <i>Triplaris</i>	41
A. Historical	41
B. <i>Triplaris surinamensis</i> Cham. and Schl.	50
C. <i>Triplaris americana</i> Linné	54
D. <i>Triplaris auriculata</i> Meisner	63
Chapter 3. Observations on <i>Tachigalia</i>	65
Chapter 4. Observations on <i>Cecropia</i>	69
Chapter 5. Observations on Rubiaceae and Verbenaceae	88
A. <i>Duroia</i> and <i>Remijia</i>	88
B. <i>Patima formicaria</i> Johnston	91
C. <i>Clerodendron Siphonanthus</i> R.Br.	93
Chapter 6. The Myrmecophytic Acacias	94
A. The Bull-horn Acacias	94
B. <i>Acacia cavenia</i> Hooker and Arnott	116
C. The African Acacias	118
Chapter 7. Various Non-myrmecophytic Plants with Ant-inhabited Stems, etc.	119
Chapter 8. The Epiphytic Bromeliaceae and their Fauna	133
References	144
Part II. Neotropical Plant Ants	155

Part I. THE NEOTROPICAL ANT PLANTS

Chapter 1

THE MYRMECOPHYTES OF THE GENUS *CORDIA*

The more than two hundred known species of Boraginaceous trees and shrubs of the genus *Cordia*, distributed over the tropics of both hemispheres, have been divided by botanists into several groups, or sections. Two of these, the *Physoclada* (*Pilicordia*) and *Gerascanthi*, are confined to the Neotropical Region and contain several ant-plants, or myrmecophytes, of unusual interest. The most typical representative of the *Physoclada* is *Cordia nodosa* Lamarck of Northern South

¹Revised for publication by Dr. Joseph Bequaert. Published with the aid of a special gift from Mr. George R. Agassiz.

America, and perhaps the only true myrmecophyte in the *Gerascanthus* section is *C. alliodora* Ruiz and Pavon, which ranges from Northern Mexico to Bolivia and eastward over the Antilles and portions of Northern South America. The following account is concerned primarily with these two species which I have had abundant opportunity to observe in the field, the former in British Guiana, the latter in Panama.

(A) The *Cordias* of the Section *Physoclada*

Professor I. W. Bailey and I found *C. nodosa* to be common, both in the primeval forest and in the second-growth jungle about Kartabo, Kalacoon, Barakara and on Great Batavia Island in the Cuyuni River, British Guiana. This and closely allied species have been frequently described by botanists, notably by Beccari (1884-'86), Schumann (1888), Schimper (1888), Mez (1890) and Ule (1907). Though *C. nodosa* may attain a height of 10 or 12 feet, most of the specimens are small, symmetrical bushes, with verticillate branches, and growing in the shade of tall trees and on low but not inundated soil. The whole plant, including the stem and the large ovate leaves, is covered with long reddish hairs. As Bailey (1924) says, the leaves "are alternate, paired or grouped in false verticels of four. The subnodal portion of the stem, subtending each verticel of leaves, is strongly thickened and angular and usually, though not invariably, provided with a long bladder-like swelling. This pouch is jacketed internally as well as externally, by a cuticularized epidermis and numerous trichomes. It subtends the lowermost of the four leaves, in the axil of which is a small apical outlet which is not excavated by the ants. Above this leaf, the thickened and much compressed primary axis gives rise to lateral inflorescences which are inserted in the axils of the three remaining leaves."

There has been much difference of opinion concerning the morphology of these ant-inhabited cauline swellings, which always occur immediately below each node with its verticel of leaves or branches. Schimper believed that the swellings are really the lateral enlargement of the base of the petiole of the lowermost leaf, which is adnate to the main axis. Schumann regarded the inflorescence as terminal and the swellings as cauline rather than as adnate foliar structures. Mez maintained the same opinion in regard to the inflorescence, but interpreted the swellings as formed by lateral invaginations, which he believed to have been originated by small insects living in the lateral

grooves and later to have been inherited as ant-domatia. Bailey (1924) after a careful study of these structures reaches the following conclusion:

"What then is the morphological significance of these extraordinary structures? The hypotheses of Schimper, Schumann and Mez do not afford an adequate explanation of all phases of their ontogenetic and phylogenetic development. Below the subnodal hypertrophy, the stem is of normal structure. Above this level, it rapidly increases in girth. As it does so, the circumference of its concentric layers of epidermal, cortical and fibrovascular tissues becomes correspondingly enlarged, and a commodious internal cavity is concomitantly formed in the dilated core of medullary parenchyma. This chamber, unlike that of the caulinary domatium of other myrmecophytes, is characterized by being jacketed by centripetal layers of epidermal, cortical and more or less rudimentary fibrovascular tissues. There is, in other words, no indication of a compressed stem and adnate petiolar enlargement, nor of an extensive lateral invagination. The centripetal layers of epidermal, cortical and fibrovascular tissues unite with the homologous centrifugal layers only in the apical portion of the wall of the domatium which surrounds the small circular aperture.

"A detailed study of the morphology of the myrmecodomatia, during successive stages of their ontogeny, indicates very clearly that they are hypertrophied portions of the cauline axes, whose medullary cavities are jacketed by layers of invaginating tissues. The invagination does not originate, however, in a longitudinal lateral groove, but in the axil of one of the leaves of the pseudo-apical verticel. As it develops, it produces an elongate-saccate ingrowth of the epidermal, cortical and fibrovascular tissues into the rapidly enlarging core of medullary parenchyma. The absence of even a rudimentary bud in the axil of a leaf which subtends an entrance aperture suggests that this growing point may be concerned in the formation of the invagination. If it is, the invagination may be visualized as the homologue of an ingrowing lateral shoot, and its formation may be likened to what happens when one finger of a glove is retracted so that it ultimately projects inwards instead of outwards. In exceptional cases, invaginations may develop in the axils of two of the leaves of a single verticel." Prof. Bailey figures an abnormal compound domatium of this type. (1924, pl. 7, fig. 8).

It is clear from this account and indeed from the most cursory study in the field that the myrmecodomatia of *C. nodosa* are preformed structures, which cannot be regarded as galls, or as hypertrophies

produced by insects or fungi. The same statement is undoubtedly true of the corresponding structures in *C. hispidissima* D C.¹ In this species, however, which seems to be common in Amazonas and Bolivia, but was not encountered in the hylæa of British Guiana, the cauline swelling is asymmetrical, that is, one-sided. Bailey, who has studied its structure on materials collected by Dr. Orlando White and Dr. W. M. Mann on the Rio Beni, in Bolivia, finds it to be fundamentally the same as that of *C. nodosa*, though differing in morphological details. "The striking difference in their external form is due to the fact that in the case of the former plants (*hispidissima*), the subnodal enlargement of the stem is unilateral and the basal projection of the trace of the leaf which subtends the entrance aperture is set off from the central cylinder at a much lower level."

C. hispidissima has been recently observed by Bequaert (Wheeler & Bequaert, 1929, p. 28) on the Lower Rio Branco, Brazil. In his account of the domatia he agrees with Bailey and states that they contained a "varied fauna". "One swelling contained a book-scorpion (*Chelifer*) occupying the lowermost portion of the cavity which was enclosed above by a white web-like partition; another was occupied by a medium-sized spider. Other domatia contained small colonies of a Ponerine ant, *Neoponera unidentata* Mayr var. *eburneipes* Wheeler. Each of these colonies consisted of four to six workers, with eggs, larvæ and pupæ, the last enclosed in cocoons."

Chodat and Carisso (1920) describe and figure another *Cordia* (*C. chacoensis* Chod.) from Paraguay, which has true domatia somewhat resembling those of *C. nodosa* and *hispidissima*, though longer and more slender. This *Cordia* grows in moist places and is of small size like the species of the Section *Physoclada*. Unfortunately no account is given of the structure of the domatia or of their tenants.

Up to the present time the following ants have been taken in the cauline swellings of *C. nodosa* and *hispidissima* in various parts of South America:

- (1) *Neoponera unidentata* Mayr var. *eburneipes* Wheeler. In *C. nodosa* and *hispidissima*. Brazil (Bequaert); British Guiana (Wheeler).
- (2) *Pheidole* (*Heudecapheidole*) *tachigaliæ* Wheeler. In *C. nodosa*. British Guiana (Wheeler).

¹Engler and Prantl, IV, 3, 1897, p. 83 recognize only one species, *C. nodosa*, in the section *Physoclada* and regard *hispidissima* D C. and *miranda* D C. as "wohl von der vorstehenden Art nicht spezifisch verschieden."

- (3) *Crematogaster (Orthocrema) brasiliensis* Mayr var. *ludis* Forel. In *C. nodosa*. British Guiana (Wheeler).
- (4) *Allomerus 10-articulatus* Mayr subsp. *8-articulatus* Mayr. In *C. hispidissima*. Bolivia (W. M. Mann).
- (5) *Allomerus 10-articulatus* subsp. *8-articulatus* var. *exsanguis* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).
- (6) *Allomerus 10-articulatus* subsp. *8-articulatus* var. *demeraræ* Wheeler. In *C. nodosa*. British Guiana (Wheeler).
- (7) *Allomerus 10-articulatus* subsp. *8-articulatus* var. *angulatus* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).
- (8) *Solenopsis tenuis* Mayr. In *C. hispidissima*. Bolivia (W. M. Mann).
- (9) *Azteca brevicornis* Mayr var. *boliviana* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).
- (10) *Azteca delpini* Emery. In *C. nodosa*. British Guiana (Wheeler).
- (11) *Azteca duckei* Forel. In *C. nodosa*. Brazil (Ducke).
- (12) *Azteca stanleyuli* Forel. In *C. nodosa*. Brazil (J. Huber; Chodat).
- (13) *Azteca olitrix* Forel. In *C. nodosa*. Brazil (J. Huber).
- (14) *Azteca ulei* Forel var. *cordiæ* Forel. In *C. nodosa*. Brazil (Ule); British Guiana (Wheeler).
- (15) *Azteca ulei* var. *gagatina* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).
- (16) *Azteca ulei* var. *gibbifera* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).
- (17) *Azteca ulei* subsp. *nigricornis* Forel. In *C. nodosa*. Brazil (Ule).
- (18) *Azteca trigona* Emery subsp. *mediops* Forel. In *C. nodosa*. British Guiana (Wheeler).
- (19) *Azteca velox* Forel subsp. *nigriventris* Forel. In *C. nodosa*. British Guiana (Wheeler).
- (20) *Brachymyrmex heeri* Forel. In *C. nodosa*. British Guiana (Wheeler).
- (21) *Myrmelachista schumanni* Emery var. *cordincola* Wheeler. In *C. hispidissima*. Bolivia (W. M. Mann).

Most of these forms and notably the species of *Neoponera*, *Pheidole*, *Crematogaster*, *Solenopsis*, *Brachymyrmex* and *Myrmelachista* are of infrequent occurrence and found only in single domatia. The same seems to be true of the species of *Azteca*, but the forms of *Allomerus 8-articulatus* are regular, or obligate tenants. This is certainly the case in *C. nodosa* in British Guiana, where the var. *demeraræ* was found

in every one of the plants examined and usually inhabiting all of its cauline swellings. As the habits of this ant are practically unknown, I single it out for special consideration.

The genus *Allomerus* was established by Mayr in 1877 for some small yellow worker ants taken by Prof. James Trail in Northern Brazil, presumably in the cauline swellings of *Cordia* or of some other myrmecophyte. In 1904 Forel described all three phases of *S-articulatus* taken in Amazonas by E. Ule in the petiolar sacs of *Tococa setifer* Pilger and observed that the females and males were of a much darker color and larger size (6 and 5.3 mm. respectively) as compared with the workers (1.8 mm.). Misled by a false analogy with the ants of the genera *Solenopsis*, *Carebara*, etc., he surmised that the *Allomeri* might be thief-ants. This is clearly disproved by my observations. The genus, as Emery has shown (*Genera Insect.*, Myrmicinae, 1921, p. 188), belongs to the tribe Monomorini and not to the *Solenopsidiini*, which comprise so many lestoproctid species.

Allomerus S-articulatus var. *demerarae* is well known to the native Indians of British Guiana, who call it the "Kurabelli" (Hohenkerk, 1918). Its colonies are extremely populous, comprising thousands of workers and many mother queens. Each colony normally occupies all or most of the cauline swellings of a *C. nodosa* bush or tree. All the cavities are connected with one another and with the forest floor by a peculiar system of galleries or arcades, constructed on the surface of the plant by the ants and measuring about 5-10 mm. in diameter. They consist of minute particles of black, agglutinated earth built up around and supported by the long red hairs above mentioned. A single gallery starts at the ground and ascends the trunk in a straight line to the first node, where it ramifies and sends a branch along the surface of each limb of the verticil. The branching is repeated at succeeding nodes till each swelling is furnished with a gallery that runs up its side and terminates at the orifice of the cavity. The minute, pale, yellow, small-eyed workers are thus enabled to pass under a continuous carton-like roof and between the stiff hairs which support it like so many pillars, from their nests in the domatia to the ground where they forage among the dead leaves. Coccids (*Pseudococcus brevipes*) are found in some of the domatia but they are not sufficiently numerous to provide more than an insignificant portion of the food required by so large an ant-population. This consideration and the elaborate construction of the system of galleries show that while the *Cordia* furnishes most admirable domatia for the ants, it is by no means an adequate source of food.

During the rainy season (July to September 1920) I found the cauline swellings full of the Kurabelli brood in all stages, together with many males and winged females. As soon as the swellings on the youngest branches become large enough, deälated females take possession of them and begin, apparently at once, to lay eggs. And though each incipient swelling usually contains only a single young deälated female, I have on several occasions found two, three or even four such females, all caring for their eggs or young larvæ in common in the same domatium. Unlike most ants, therefore, *Allomerus* is normally pleometrotic even during the incipient stages of colony formation. After they are reared the broods of these young females undoubtedly become a part of the single large polycladic colony which possesses the whole plant.

It cannot be maintained that the Kurabelli act as an efficient protective body guard for the *Cordia*, at least so far as man and the larger animals are concerned. When one handles the plant for some time, the workers do indeed swarm over one's clothes and body and for some time keep on stinging, but their stings are so feeble that they produce merely a rather unpleasant itching and that only of parts with very thin epidermis. While it is probable that the Kurabelli may be more efficient in keeping the plant free from certain insects, it should be noted that other Formicidæ not infrequently occupy some of the domatia although most of them may be tenanted by *Allomerus*; and both Professor Bailey and I sometimes found the foliage of such plants considerably damaged by leaf-cutting ants (*Atta cephalotes*). I have also noticed leaves that had been extensively gnawed by caterpillars. The foliage of one plant tenanted by *Allomerus* was covered with Cecidomyid galls. The only other insects found associated with *C. nodosa* were termites, which on one occasion were seen to have a gallery extending up the trunk from the ground and terminating in a cauline swelling which they were occupying.

(B) The *Cordias* of the Section *Gerascanthus*

The *Cordias* of this group, unlike the *Physoclada*, are tall shrubs or trees, frequently attaining a height of 20 to 60 feet, with gray bark and coriaceous, opaque leaves, covered with dense stellate hairs beneath and sparser hairs of the same type above. The branching, except in young specimens, is much more obscurely verticillate, and though ant-inhabited cauline swellings occur in some of the forms, they are much simpler than those of the *Physoclada*, being merely

conical, pyriform or turbinate dilatations of the stem beneath and at the nodes, with large medullary cavity and without a preformed orifice. The flowers are much more numerous and showy and usually aggregated in broad, dense panicles or corymbs. As a rule, the plants grow on higher ground, in the campos or more open woods and thickets and therefore in more xerothermal situations than the *Physoclada*.

Chodat and Vischer (1920) recognize seventeen species as belonging to the *Gerascanthus* section, but many of the forms are by no means easy to distinguish, as may be inferred from their remarks: "After a thorough examination of the flowers of all the species, we have been unable to discover other characters than those derived from the size of the organs, the more or less elongate form of the ovary or of the infraovarian disc, the importance of which is in no respect greater than that resulting from an examination of the vegetative organs and the external appearance of the calyces and corollas. The evaluation of the characters derived from a comparison of the reproductive structures is rendered difficult by the fact that there is in the *Cordias*, at least of this group, a pronounced heterostyly We are, in fact, dealing with species which are but feebly defined morphologically and the taxonomy of which will require revision from time to time as observations in the field increase in number". The authors call attention to the fact that Schumann (1888) regarded all the forms of the *Gerascanthus* section as constituting but a single species.

Of the seventeen species recognized by Chodat and Vischer, nine are recorded as certainly possessing cauline swellings and as being therefore myrmecophilous, namely *C. gerascanthus* L. and its varieties, *alliodora* Ruiz and Pavon, *Rusbyi* Chodat, *Hassleriana* Chodat, *glabrata* var. *orbicularis* Chod. and Vischer, *longituba* Chod. and Vischer, *Chamissoniana* Steud., *Cuyabensis* Manso and Lhotsk. and *gerascanthoides* H B K. They failed to detect myrmecophily in *C. hypoleuca* D C. and *excelsa* D C. and in certain large-flowered species not belonging to the *Gerascanthus* section, namely *insignis* Cham., *nettoana* Taubert, *Hainkeana* Mez and *formosa* Chod. I may add, in this connection, that I have failed to find any indications of myrmecophily in *C. lutea*, which is very common in the Galapagos Islands. It has no cauline swelling and ants merely visit its clusters of showy yellow flowers for their nectar. The same is true of the beautiful, vermilion-flowered *C. sebestena* L., which is common in gardens and along road-sides in the West Indies.

Among the myrmecophilous species cited by Chodat, *C. gerascanthus* is given as the most widely distributed and best known. He distinguishes three forms or varieties of the plant: *genuina* Chod. (= *gerascanthoides* Rich. non H B K), *Martinicensis* Chod. and *micrantha* Jacq. The first is cited as common in the Antilles (Cuba, Porto Rico, Jamaica, Haiti, St. Thomas, St. John, Antigua, St. Vincent, Trinidad), the second as known from Martinique and Guadeloupe, the last from Mexico, Guatemala, Nicaragua, Costa Rica, Panama, Peru and Bolivia. This form has somewhat smaller flowers than the West Indian type (*C. gerascanthus* L. *sens. str.* or *genuina*) and has been cited by authors as *C. gerascanthus* Jacquinus.

This interpretation of *gerascanthus* is not accepted by Dr. I. M. Johnston (1924), of the Gray Herbarium, to whom I submitted specimens of the species I studied in Panama for identification. I quote his remarks on the synonymy of *gerascanthus* and *alliodora*: "In 1910 Urban, Symb. Antil. IV. 516 indicated, that, as then used, the binomial *Cordia gerascanthus* L., was incorrectly applied to the widely distributed tree with canescent, densely stellate calyces, and that the name is properly applicable to the relatively localized species of the West Indies and Southern Mexico which has glabrous or sparsely hirsute calyces and larger flowers, and which was described and current as *C. gerascanthoides* H B K. Ten years later, in his paper on *Cordia Gerascanthus* Chodat, l.c. declared Urban's interpretation of *C. gerascanthus* L. to be incorrect, and used the term in the traditional sense, applying it to the widely distributed plant with stellate calyces. Further examination of this matter has recently been made to determine the correct specific name for use by Dr. W. M. Wheeler in his publications on myrmecophytes. For the convenience of others the results of this study are here put on record. *Cordia gerascanthus* L. is based upon the Jamaican plant which Patrick Browne, l. c., described and figured under the name "Gerascanthus". Browne's illustration, showing only the floral structures, portrays a corolla of large size which has broad short obtuse lobes with conspicuous pinnate veining, a broad saucer-shaped throat, a stocky weakly ribbed calyx, and deltoid calyx lobes. These characters definitely associate Browne's plant with *C. gerascanthoides* H B K. and prohibit the use of the Linnean name for the plant with stellate calyces. It is to be also noted that not only does Grisebach, l. c. cite Browne's figure under "*C. gerascanthoides* H B K.," but he gives *C. gerascanthoides* H B K as "common in the lowlands and mountains" of Jamaica, and gives the plant with stellate calyces (under *C. gerascanthus* Jacq.) as "rare" on

that island. Browne's plant was not rare, for he speaks of it as follows. "This tree grows in many parts of Jamaica, and is generally esteemed as one of the best timber woods of the island; it rises to considerable height . . . , especially in the lowlands, where it is most common, . . . "It is significant that concerning the Jamaica occurrence of the plant with stellate calyces, Urban, l.c. (under *C. alliodora* Cham.), comments parenthetically as follows, "fortasse a cl. Wilson introducta ex cl. Stapf. in lit." Since the identity of *C. gerascanthoides* H B K. and *Gerascanthus* Browne is certain from a study of Browne's plate and description, and from distributional considerations, it is evident that *Cordia gerascanthus* L. is, indeed, improperly applied to the widely distributed plant with stellate calyces. Among its close relatives in the West Indies and Central America, *C. gerascanthus* L. is readily recognized by its large flowers, saucer-shaped throat, hirsute or glabrescent stout weakly ribbed calyx-tube, and deltoid calyx lobes. It is known only from Cuba!, Isle of Pines!, Jamaica!, southern Mexico!, and northern Central America. As Urban, Symb. Ant. IV. 516 (1910) and VIII 574 (1921), has pointed out, *Cordia alliodora* (R. & P.) Cham. is the correct name for the widely distributed plant with stellate calyces, or in other words, for the one incorrectly current as "*C. gerascanthus*". *Cordia alliodora* ranges from Mexico and the West Indies southward along the Andes to Bolivia. A number of critical species, doubtfully distinct from it, have been described from Brazil, adjacent Paraguay and Argentina".

It is clear, therefore, that at least two quite different species of *Cordia* have been confused in the literature, *C. gerascanthus* Linn. (= *gerascanthus* Browne = *gerascanthoides* H B K = *gerascanthoides* Griseb.) of the West Indies and *C. alliodora* R. and P. (= *gerascanthus* var. *micrantha* Chodat), distributed from Mexico to Bolivia and also occurring, though apparently not very abundantly, in the West Indies. Now *C. alliodora* always has preformed domatia whereas these structures do not occur in *gerascanthus* L. Chodat, who has interpreted all the cauline thickenings in the group of *Cordias* under consideration as insect galls, has simply obscured and confused the whole subject, because while both species may have occasional stem-galls produced by insects, these galls have nothing to do with the preformed myrmecodomatia which are a conspicuous and constant feature in *C. alliodora*. This matter will be discussed in greater detail in the sequel. I may add that a very competent dendrologist, Professor J. G. Jack of the Arnold Arboretum, has at my request carefully studied many specimens of *gerascanthus* L. at the Harvard Botanical Garden

at Soledad, Cuba and reports that he has utterly failed to find the slightest traces of myrmecodomatia in this species.

I admit the possibility, however, that the Panamanian *C. alliodora* may differ varietally from the Peruvian type described by Ruiz and Pavon, for the following reasons: (1.) The original figure represents the flowers accurately but does not show the cauline swelling which should be present at the base of the panicle. Of course, either the draftsman or the authors may have omitted it owing to its resemblance to a gall. (2.) The authors state that the tree blooms in July and August, whereas the Panamanian tree is leafless at that season and blooms in spring. (3.) According to Ruiz and Pavon, the Peruvian Indians use the bark and leaves as a condiment. The bark and foliage of the Panamanian tree seem to have no such properties. (4.) The following peculiarities cited by Ruiz and Pavon, do not apply to the plants I have examined: "Cum hujusmodi arbores secantur, odorem gravissimum emittunt, et penetrantissimum, qui oculorum aciem perstringit. Corticis recenter evulsi odorem valde foetidum vulpis urinæ haud multum adsimilem, spirant; postea vero Foli uti etiam Cortex, allium maxime redolent, unde nomen ab Incolis Arbor acceptit." After calling Dr. Johnston's attention to these considerations, he agrees with me that the plant usually cited as *C. Gerascanthus* Jacquinus and more recently as var. *micrantha* Chodat of that species, may differ from the typical *alliodora*. I quote from one of his letters received February 23, 1924: "In regard to *Cordia*, I must say that I am unable to give you any thoroughly satisfactory information. The plant which Chodat treats as *Cordia gerascanthus* var. *micrantha* is newly published by him, although from Chodat's citations one might obtain the impression that the trinomial was based upon a name published by Jacquin. Chodat apparently wished to show that his form *micrantha* was based upon and included the plant described and figured in his *Stirpium americanarum* 43. t. 175, fig. 16 (1763). Unfortunately, however, Jacquin's figure shows a flower 17 mm. long, whereas Chodat describes his forma *micrantha* as having corollas 1 cm. long. In other words, Jacquin's plant is the common form growing in the West Indies, and the plant which Chodat describes is the common form of Mexico and Central America. Wright's plant from Cuba is probably the same form as that figured by Jacquin. I have talked the situation over with the men at the Gray Herbarium and the general consensus of opinion appears to be that Chodat's forma *micrantha* had best be taken as covering the plant described, rather than as being typified by the cited specimen or the reference to Jacquin. With this solution of the

problem I am now inclined to agree. If it is satisfactory to you Chodat's trinomial will be applicable to your Panamanian form. The situation will undoubtedly lead to confusion since certain workers are inclined to lay more weight on the specimens cited than upon the actual description, and such people would treat Chodat's trinomial as applying to the large-flowered West Indian form of *alliodora*. This and other aspects of Chodat's work on *Cordia* has led me to believe that it has given more confusion than it can possibly give help. Certainly it has not clarified the classification of that large and difficult genus. Intercategorical priority not being recognized in botany, I can and will describe the Mexican and Central American plant as a geographical *variety* of *alliodora*, giving it a new name and one definitely applicable to your plant." Since this name is not yet published, I am using the specific name *alliodora* for the small flowered plant, with preformed myrmecodomatia and a distribution from Mexico to Bolivia.¹

The confusion between *gerascanthus* Linnaeus and *alliodora* Ruiz and Pavon has, of course, led to confusion in the writings of those who have considered their relations to the ants. I therefore take up in chronological sequence the various authors who have been concerned with the Panamanian species.

One of the earliest, if not the earliest accounts of the relations of ants to *Cordia*, is that of Ruiz and Pavon (1799), who described and figured *C. alliodora* (called the "arbol del ajo" an account of its onion-like odor) from Peru under the name *Cerdana alliodora*. They observed that "quædam exiguæ Fornicæ quarum punctio acrem intolerabilemque pruriginem diu persistentem excitat, frequenter has arbores infestantes folia devorant, ita ut vix ulla inveniri possint integra." The first part of this sentence seems to refer to some species of *Pseudomyrma*, but the latter part is erroneous, since no ants devour foliage, and if it refers to the leaf-cutting Attini, they neither sting nor occupy the cauline swellings of *Cordia*. Chodat seems to have been led by the remark of Ruiz and Pavon to a peculiar and partly erroneous interpretation of the peculiar carton dissepiments (vide infra) in the cauline swellings and of the habits of their ant-inhabitants.

Spruce (1869 [1908]) was also familiar with *C. alliodora*, which he cites under the name *gerascanthus* Jacq. He says that "it rises to a stoutish tree 30 to 40 feet, and is throughout fasciculately branched

¹Since this passage was written I find that Standley, in his works both on the Mexican (1924) and Panamanian flora (1928), has accepted *C. alliodora* as the correct name of the plant which I studied.

(branches 3-5-nate). At the point where the branches divide there is mostly a sac, inhabited by very vicious ants of the tribe called 'Tachi' by the Brazilians." These ants are undoubtedly a species of *Pseudomyrma* and, in all probability, some form of *Ps. sericea* Mayr or *Ps. gracilis* Fabr.

Seemann (1852-1857), in his early account of the Panamanian flora shows that he was acquainted with *C. alliodora* (cited as *gerascanthus* Jacq.), which, he says, is called "laurel" by the natives, who use its wood in building and cabinet making. Although he records the tree as common on the outskirts of woods in the provinces of Panama and Veragua and as growing as far north as Mazatlan and Acapulco, Mexico, he says nothing about the peculiar cauline swellings and their tenants. The same remark applies to Saldanha da Gama (1874), who cites *C. alliodora* as distributed throughout Brazil and as known under the name "lauro". It is described as a large tree of such rapid growth that it may yield planks in only eight years after germination. The wood is described as odorous, light and nonresistant to moisture and as being used for doors in the interior of houses, lintels, etc. The carpenters who saw it become very thirsty, owing to peculiar properties of the sawdust, and the shavings withdraw so much moisture from the hands that the workmen find it an unpleasant wood to handle. There is nothing in Saldanha da Gama's description to prove that his *C. alliodora* is identical with the Panamanian species.

Beccari (1884) was the first to give a clear description and figure of the cauline swellings of *C. alliodora* (cited as *gerascanthus*) from Mexico. He mentions the plant as occurring in the Sierra Padre Nolasco, at Talca, Acapulco, etc. The peculiar convoluted or travecular carton in the cavity of each swelling was figured and described, and he noticed the entrance holes made by the ants and the large coccids kept by them on the walls of the cavity. He was also quite clear in regard to the structural differences between the swellings of *alliodora* and those of the *Physoclada* (*C. nodosa*).

Schumann (1888), after a brief revision of the *Cordias* of the *gerascanthus* group, confirmed but added nothing to Beccari's statements concerning the Mexican *alliodora*.

More than twenty years ago Mr. C. H. Tyler Townsend sent me a number of cauline swellings of *alliodora* which he found near Cualata, on the slopes of the Volcan de Colima, Mexico. Referring to his letter of August 6, 1902, I find the following note: "The ant from Cualata which keeps a large red lecanoid coccid in its nests in swollen joints of the branches of a tree is probably a new species. This is a most striking

instance of the interrelations and mutual dependence of plant, ant and coccid." Professor Forel, to whom I later sent specimens of the ant, described it as *Azteca longiceps* Emery subsp. *patruelis* (see Pt. II p. 233). The specimens of the cauline swellings and Coccids, still in my collection, are precisely the same as those described below from Panama. They bear the following note in Townsend's handwriting: "July 25, 1901. Living in hollow cavities in swollen joints of tree growing 12 to 20 ft. high. Cavities in these joints are natural. Ants keep coccids inside. Small hole or holes allow exit of workers only. These holes are sometimes found grown over, the ants having failed to keep them open, in which case the dead ants are found inside. Joints are inhabited all the way from twigs to branches an inch in diameter. The ants bite."

Pittier (1908) in his work on the economic plants of Costa Rica, cites *alliodora* under the name *gerascanthus* Jacq., with the following note: "The laurel is one of the most important trees of the country as wood for construction. The trunk is straight, with white bark; the leaves are small, entire, elliptical-lanceolate; the flowers white, very fragrant and in large racemes. The wood is rather fine and hard, of a pale chestnut color and easy to work; it is reputed to be incorruptible and is used principally for flooring. The laurel grows on both slopes of the country from sea-level to 1500 m. more or less, but attains greater dimensions on the Atlantic Slope." Pittier gives the Bribri Indian name of the tree as "dze-ui." ¹

Ule (1907) observed *C. gerascanthus* Jacq. (evidently *alliodora*) at Tarapoto, Peru, in a rather dry region and, after describing the cauline swellings, remarks that "they make the impression of galls, but are thin-walled and nearly always harbor in their cavities a very vicious species of ant, *Pseudomyrma sericea* var. *cordiæ* Forel."

Dr. W. M. Mann, while on the Mulford Expedition to Bolivia during 1921-22, made observations on *C. alliodora* (probably var. *boliviana* Chod. & Visch.) and collected ants and other insects in the domatia. These structures are precisely like those of the Central American and Mexican trees.

Bailey (1924), who made a study of the structure of the cauline swellings of *C. alliodora* (cited as *gerascanthus* Jacq.) from herbarium specimens, remarks that "the domatia vary greatly in size, shape and

¹According to Boulger (1908) the wood of *C. gerascanthus* L. of the West Indies is known in commerce as "bois de Cypre", "prince-wood", "Spanish elm", "Dominica rose-wood" and "bois de Rhodes" and is "dark brown with dusky, excentric zones, open-grained, soft, durable." He says that it is used in cabinet-work. According to Maza and Roig. (1906) *C. gerascanthus* and *gerascanthoides* are called by the Cubans "capa rota", "palo de rosa", "palo de rosa del pais." The wood of *gerascanthoides* is said to be one of the most valued in Cuba, of a dark chestnut color and to be employed in rural carpentry.

distribution in different representatives of even a single species. When present, they are irregularly shaped hypertrophies of the axis of the large diffuse inflorescences or of the transitional region between cauline and floral axes. The lateral entrances or exits are not preformed apertures, as in the Physocladæ, but evidently are excavated by insects. Furthermore, the domatia are not jacketed internally by invaginated layers of epidermal, cortical, and fibrovascular tissues. As shown in fig. 3, a transverse section of a myrmecodomatium of *Cordia Gerascanthus* Jacq. (Herbarium J. D. Smith no. 4365), the inflated central cylinder surrounds a large heterogeneous medulla, the large celled, succulent core of which has dried up and has been trimmed away by the ants. In other words, these structures are quite distinct morphologically from the myrmecodomatia of the Physocladæ, and resemble those of the Ethiopian species of *Cuviera* and *Plectronia*."

Menozzi (1927) has recently published a few notes on the domatia of *C. alliodora* (cited as *gerascanthus*) and their ant-tenants, collected by H. Schmitt in the vicinity of San José, Costa Rica. He figures the domatia and describes the coccids as probably belonging to the genus *Cryptostigma*, the ants as *Crematogaster brevispinosa* Mayr, *Cryptocerus setulifer* Emery and *Azteca pittieri* Forel.

Standley (1924, 1928) gives a brief but excellent description of *C. alliodora*. He finds it to be one of the common trees of Mexico and Central America, ranging as far north as Sinaloa, and that "the forks of the young twigs are almost always enlarged by hollow swellings, which afford shelter for fierce ants, hence the name "hormiguero." The fruit is edible. A decoction of the leaves is employed as a tonic and stimulant, especially in the case of catarrh and afflictions of the lungs, and an ointment made with the pulverized seeds has been used in the West Indies as a remedy for cutaneous diseases. The fresh bark is reported to have an odor suggestive of garlic."¹

The recent study of the *Cordias* by Chodat, Vischer and Carisso is so much more extensive than that of previous writers and so significant in connection with my own observations that it must be considered in detail. These authors investigated several species in Paraguay (*C. glabrata* var. *orbicularis*, *Hassleriana*, *Chamissoniana* and *longituba* and two species, *salicifolia* and *chacoensis*, which belong to another

¹The economic value of the tree is indicated by the number of local names which it bears in various parts of its range. These are cited by Standley as "bojon", "bojon blanco", "bojon prieto" (Tabasco); "tambor" (Michoacan); "hormiguero" (Michoacan, Guerrero, Oaxaca); "amapa prieta" (Sinaloa); "palo de rosa" (Oaxaca, Cuba, Porto Rico); "palo Maria" (Guerrero); "laurel" (Panama, Costa Rica, El Salvador, Guatemala, Honduras); "Solera" (Colombia); "laurel macho" (Nicaragua); "capa prieta" (Porto Rico, Cuba); "varea", "capa voja" (Cuba); "canjara", "pardillo" (Venezuela); "arbol del ajo" (Peru); "laurel blanco" (El Salvador); "canaleta" (Colombia).

group of forest *Cordias* related to the *Physocladæ*). As *gerascanthus* and *alliodora* were not found in that country, they resorted to herbarium material for a knowledge of their peculiarities. For many ecological and morphological details of purely botanical interest the reader may be referred to the first section of the paper by Chodat and Vischer, but the second section, on the myrmecophily of the *Cordias* of the *gerascanthus* group, calls for fuller discussion. The authors state positively that the cauline swellings are not preformed structures, like those of the *Physocladæ*, but true galls, which owe their inception to infection by a Chalcidid of the genus *Eurytoma*, according to C. Ferrière's identification. Apparently, only the egg, larva, and pupa of the gall-maker were obtained. In *C. gerascanthus* Jacq. (that is *alliodora*) "the beginning of the infection occurs at the base of the inflorescence or at one of the secondary forks of the latter." And the following remark is added: "We have traced a similar development of this gall in the plant from Guatemala. We were able to ascertain the beginnings of the development of the larva and its morphogenetic action which results in a biomorphosis of rather large dimension." Their Fig. 322 shows a longitudinal section of a young gall with the *Eurytoma* egg at the base of a feeble inflorescence of *C. gerascanthus*. Their Fig. 328 is an enlarged sketch of portions of such a gall containing larvæ. Figs. 314a and b, representing similar conditions in *C. glabrata* and *longituba*, are quite unlike the typical cauline swellings of *alliodora*. Moreover, the authors confess (p. 179) that they did not follow the ulterior development of the structures in the Paraguayan species. The exit hole of the adult *Eurytoma* is supposed to form the opening through which the ants enter the gall, but no attention is paid to the relatively enormous increase in size of the structure after the minute Chalcidid has deserted it, and neither text nor figures give any clear and convincing account of the eventual developments. The authors do, indeed, describe and figure the fully-formed swelling but they say nothing about the large coccids which were probably present in the specimens they examined and had so vividly impressed Beccari and Townsend. The Genevan botanists figure the convoluted carton mass in the cavity of the fully formed swelling and find it to consist of the pollen and stellate and other hairs of the *Cordia*, starch grains from the pith, filamentous fungi and bacteria allied to *Leptothrix*. This led them to quote the sentence which I have cited from Ruiz and Pavon and to conclude as follows: "We may therefore regard these structures (i.e. the convoluted masses of carton) as fabricated by the ants themselves with materials from at least two sources: first, the

pollen of the innumerable flowers, second, the agglomerated fragments of lacerated leaves or flowers; third, fungi, perhaps intentionally introduced by the ants (acting as mycetophagous horticultural insects?).” The general results of their study are summarized by the authors in the following paragraphs:

“Thus our investigations establish a new theory of the myrmecophily of *Cordia*, based on incontestable facts. Observations made by an entomologist in the field will fortunately complete the sketch which we have given of the curious phenomenon, but in the meantime, till these observations are forthcoming, we would call the attention of biologists to the following points:

“Throughout the entire range of species of the section *Gerascanthus*, from Mexico to Paraguay and from the Guianas to Eastern Peru and Bolivia, we have demonstrated, either in the field (Paraguay) or by examination of herbarium material, the general infection of these plants by ants of the genera *Azteca* and *Pseudomyrma*, which establish their formicaries in these empty galls with materials taken from the host plant. We are not therefore concerned with ants that protect the plants from the leaf-cutters (*Atta sexdens*), according to the theories of Belt, Mueller and Schimper, since they attack, not the glands of the plant as if they were prepared for the benefit of the ants, as in the case of *Cecropia*, but essential organs, the pollen and parenchyma of the leaves.

“We fail to see any advantage these trees might derive from the presence of the ants. Their habitual presence on certain species of *Cordia* clearly shows that these plants which are so widely distributed and so abundant, suffer in no wise essentially from this attack and symbiosis, since it does not jeopardize the existence of the *Cordias* and is profitable only to the ants. In so far as the *Cordias* are concerned, there is merely a biomorphogenic reaction.”

Chodat and Vischer thought they found confirmation of their general conclusions in their study of the large thorns of the Paraguayan *Acacia carenia*, which will be discussed in connection with the Central American *Acacias* (p. 116), but Fiebrig (1909) had previously shown that the *carenia* thorns are originally inhabited by Lepidopteran larvæ (Tineids). It is difficult to understand why Chodat and Vischer should have regarded their theory as “new.” Myrmecologists have long been familiar with the fact that ants are very fond of nesting in the abandoned galls of the most diverse plants, and if the cauline swellings of the *Cordias* of the *Gerascanthus* group are nothing but galls, these plants obviously cannot be regarded as myrmecophytes. I trust that

the following observations will demonstrate that the Genevan botanists have reached erroneous conclusions from insufficient data.

I found *C. alliodora* to be a common tree on the Pacific side of the Panama Canal Zone and less abundant on the Atlantic side near Colon. There are many specimens of it on Barro Colorado Island, in Gatun Lake. From Frijoles to Ancon it is often a common component of the second growth jungle, or thickets, and in the clearings, but usually avoids moist spots and seems to show a preference for the slopes of hills. It comes into flower during the last days of February and continues to bloom profusely till about the last of March. During this portion of the dry season the large compact racemes of small white blossoms make the trees conspicuous objects in the landscape (Plate 2). Seedlings and young trees of all sizes can often be found singly or in colonies, especially about clearings and along road-sides, so that the plant can be readily studied in all stages. The largest specimens attain a height of 30 or 60 feet, but flowers are not produced till the tree is about 10 or 15 feet high. While young (below about 4 to 6 feet) it is usually very symmetrical, with the branches coming off in regular whorls at intervals along the straight, slender trunk, so that in this stage it somewhat resembles adult specimens of *C. nodosa*. Later the branches vary much more in length and direction and are less horizontal. Eventually the crown of foliage may be either irregularly pyramidal or, especially when growing in the open, more diffuse and spreading. The trunk and branches are slender and graceful, with moderately smooth, gray bark. The ovate coriaceous leaves are two to four inches long and grayish green, with rough margins. There is considerable variation, however, in the texture and surface of the leaves. The flowers have a strong odor, somewhat like that of decayed urine. They soon turn brown and persist for some time, often till the middle of April, but later fall and the same is true of the stems of the inflorescence though its base, which has a well-developed swelling, or domatium, may remain behind as a dead and dry structure for at least a year. Those who may not care to study the *Cordias* in the Panamanian jungles will find a number of fine specimens on the hospital grounds, about the reservoirs and in the gardens of private dwellings at Ancon and Balboa. An unusually fine tree nearly 40 ft. high, on the slope of the lower reservoir in Ancon, is shown in Plate 1.

An examination of fully developed trees shows that there is almost invariably an ant-inhabited domatium at each node and that the swellings grow larger successively the nearer they are to the bases of the branches, but the stoutest branches and the trunk exhibit little or

no enlargement in the corresponding regions. Here the domatia persist, nevertheless, but are concealed by a normal and very considerable growth in the thickness of the wood. These masked cauline swellings will be discussed later. It is a singular fact that the adult *C. alliodora* trees lose their leaves during the rainy season when all the other trees of the Panamanian jungle are in full foliage. During July and August the bare trunks and branches stand out as if dead among the dense green foliage of the other trees and the regular arrangement of the domatial swellings at the insertions of the branches becomes conspicuous. These trees could thus be located with the field glasses among the tree-tops visible from the laboratories at Ancon and Barro Colorado Island.

My attention, like that of Chodat and his collaborators, was at first directed to the inflorescences, by finding distinct elongate thickenings of some of the small flower-stems, and I, too, at first took these thickenings to be the initial stages in the formation of the ant-inhabited cauline swellings. In the thickenings I also occasionally found minute maggot-like larvæ but did not succeed in rearing the adult insects. They were Hymenopterous and very probably the larvæ of the Eurytomid observed by the Genevan botanists, or of some allied species. But most of the enlargements, which are only 2 or 3 mm. in diameter, contain no traces of eggs or larvæ and are filled with a uniform and undisturbed mass of brown pith. I am certain, therefore, that they are not galls but merely occasional preformed thickenings of the flower stems, in which the Eurytomids lay their eggs. In other words, these thickenings are strictly limited structures which precede the infection and are not produced by it. The Eurytomid (?) larvæ simply feed on the pith which happens to be more abundant in the thickenings than in other portions of the flower stems. That these thickenings do not become the true nodal or cauline swellings inhabited by the ants is proved by the fact that they wither and drop off after flowering and cannot therefore produce persistent leaf-bearing branches. True woody galls are, however, occasionally produced on the twigs of the tree by some unidentified insect. One vigorous young tree about 10 ft. high observed Aug. 2, 1924 on Barro Colorado Island had many of these galls, which were regular, ovoidal thickenings of the twigs below the true domatia, quite unlike them in form and about 2 to 3 cm. long and 1 cm. in diameter. They were hard and woody and contained winding passages made by some boring larva. Many of these galls were inhabited by portions of the same *Azteca longiceps* colony that occupied the domatia on the same tree.

In order to ascertain the origin of the cauline swellings it is necessary to investigate the seedlings and young *Cordias* and the suckers that often grow up from the roots of larger trees that have been felled. These juvenile stages present a very different picture from that described by Chodat and Vischer. The plants are green throughout and actively growing and, as I have stated, very symmetrical in the arrangement and length of their few branches and nodes. Each of the latter is regularly swollen and turbinate and forms a rather thin-walled, green capsule closed on all sides and varying according to its age from 5 to 20 mm. in diameter. The delicate remnants of the pith form an even layer over the wall of its large cavity, which contains no traces of any insect parasite. Nor does the great majority of juvenile trees or suckers contain any ants till they reach a height of about three to five feet. The swellings are so perfectly regular in their arrangement and in position so comparable to those of the *Cordias* of the *Physoclada* group and so constantly present, except in the youngest seedlings less than a foot in height and with only the first whorl of leaves, that no botanist and certainly no entomologist, could possibly regard them as galls. Very occasionally there may be no swelling at a node where it might be expected to appear, but this sometimes happens also in the *Physocladæ*, and such inhibitions of development do not invalidate the general conclusion that the domatia of *C. alliodora* are quite as certainly preformed structures as those of *C. nodosa*. Of course, the crucial demonstration that the domatia are preformed can come only from growing the plant from seed under controlled conditions. I made an attempt to accomplish this with seeds sent to Boston by Mr. Zetek, but they failed to germinate in the hot houses of the Bussey Institution, probably because they were sterile.

Except in the domatia at the very base of the inflorescence, which, as previously stated, may persist and dry up when the latter falls off, there is a gradual growth in the thickness of the woody walls and in the size of the enclosed cavity. The series of photographs (Plates 3, 4, 5) show this increase very clearly. The ants perforate and enter the domatia very soon after their walls begin to lignify. I have not been able to follow the details of the invasion, although I have frequently found single young dealated females of various species and notably of *Azteca longiceps* Emery, either alone or with their first brood of larvæ, in the swellings. The perforation or perforations—for there may be several—are always made in the thinner portion of the wall below the node, but there is no regularity in their position. In many cases the opening made by the entering queen closes by growth of the

plant tissues and has to be reopened by the first brood of workers. The continued growth of the domatium after its occupation must be due to the constant irritation produced either by the ants or by the numerous Coccids which attach themselves to the walls of the cavity and sink their delicate mouthparts into the plant tissues. That the Coccids may be the more potent irritants seems to be indicated by the conditions in the various Aphid and Psyllid galls of temperate regions and the Coccid galls of Australia. In the case of *C. alliodora*, the Coccids may be responsible not only for the irregular shapes assumed by many of the domatia in their later stages, but also for the unequal vigor and growth of the branches at the nodes and the general asymmetry of the older trees (Plate 6, Fig. b).

That the growth of the domatia does not continue indefinitely is shown in longitudinal sections of the nodal regions of the trunk and larger branches of old trees. The cavity ceases to enlarge when it reaches about the size shown in Plate 6, Fig. a, but the layers of xylem in its wall increase so enormously that the external swelling is obliterated. Concomitantly with this growth in the xylem the perforations or entrances to the cavity develop as long tubular galleries which traverse the whole layer of wood. In the section represented in Plate 7, Fig. b, which is slightly less than natural size, there were seven of these galleries which radiated from the chamber and opened on the surface of the bark at points several inches apart. Although even at this stage the cavities may still be inhabited by ants, the Coccids have all disappeared, probably because their food-supply has been completely shut off by the development of the very thick layer of wood between the cavity and the cambium.

The regular development of the swellings, or domatia in *C. alliodora* and other *Cordias* with such preformed structures thus presents a very interesting problem to the plant anatomist interested in phylogeny. Attention may be called in this connection to similar structures in at least one other plant in a very different genus, all the other species of which have stems of the normal unswollen type. This is the Polygonaceous genus *Eriogonum*, which comprises about 100 species in the United States west of the Mississippi. One, however, *E. inflatum* Torre, has hollow, fusiform swellings at the upper ends of the internodes of the stems and branches. I have observed this plant at Palm Springs, California, in the Mojave Desert at the foot of the San Jacinto Mountains. It is one of the few perennial species of the genus, according to Tidestrom (1925), who cites it as belonging to the "desert areas and hillsides of the Covillae and Artemisia belts" in

Southwestern Utah, Colorado, Nevada, Arizona and California. The swellings are not inhabited by ants, because all the Formicidæ in its desert environment are earth nesting forms. But there is no doubt in my mind that if the plant were to invade the tropics, certain species of stem-inhabiting ants would at once take up their dwelling in its inflations. Under these circumstances the plant would become a regular myrmecophyte like *C. alliodora*.

The following is a list of the ants that have been found associated with *C. alliodora* in Mexico, Central America, Peru and Bolivia:

- (1) *Ectatomma ruidum* Roger. Foraging on trunk and foliage. Ancon, C. Z.
- (2) *Ectatomma tuberculatum* Oliv. Foraging on trunk and foliage. Barro Colorado Is. C. Z.
- (3) *Ectatomma (Gnamptogenys) sulcatum* F. Sm. On trunk. Barro Colorado Is. C. Z.
- (4) *Neoponera crenata* Roger var. *mæsta* Mayr. In domatia. Quebrada de Oro and Barro Colorado Is. C. Z.
- (5) *Pseudomyrma alliodoræ* sp. nov. In domatia. Ancon, C. Z.
- (6) *Pseudomyrma belti* Emery subsp. *fulvescens* Emery. Types taken by Beccari from domatia. Guatemala.
- (7) *Pseudomyrma excavata* Mayr, var. *flaviventris* Forel. In domatia. Barro Colorado Is. C. Z.
- (8) *Pseudomyrma gracilis* Fabr. In domatia. Red Tank, C. Z. In domatia of *C. alliodora* var. *boliviana*? Riberalta, Bolivia (W. M. Mann).
- (9) *Pseudomyrma gracilis* var. *bicolor* Guérin. In domatia. Red Tank, Gamboa and Quebrada de Oro, C. Z.
- (10) *Pseudomyrma sericea* Mayr var. *ita* Forel. In domatia, Ancon, C. Z.
- (11) *Pseudomyrma sericea* var. *cordiæ* Forel. In domatia of *C. alliodora* var. *boliviana*? Ivon Beni & Huachi Beni, Bolivia (W. M. Mann); Bolivia (Bang); Eastern Peru (Spruce).
- (12) *Pheidole radozkowskii* Mayr, var. In domatia. Ancon, C. Z.
- (13) *Crematogaster acuta* F. In domatia. Barro Colorado Is. C. Z.
- (14) *Crematogaster (Orthocrema) brevispinosa* Mayr. In domatia. San José, Costa Rica (H. Schmitt).
- (15) *Crematogaster (Orthocrema) brevispinosa* subsp. *tumulifera* Forel. In domatia. Red Tank, C. Z.
- (16) *Crematogaster (Orthocrema) limata* F. Sm. var. In domatia. Gamboa, C. Z.

- (17) *Crematogaster (Orthocrema) limata* subsp. *parabiotica* Forel. In domatia. Red Tank, C. Z.
- (18) *Crematogaster (Orthocrema) limata* subsp. *ludio* Forel. In domatia of *C. alliodora* var. *boliviana*? Huachi Beni, Bolivia (W. M. Mann).
- (19) *Crematogaster (Orthocrema) sumichrasti* Mayr. In domatia. Quebrada de Oro, C. Z.
- (20) *Crematogaster (Orthocrema) virgula* Forel. In domatia. Red Tank, C. Z.
- (21) *Solenopsis hermione* Wheeler. In domatia. Gamboa, C. Z.
- (22) *Solenopsis laeviceps* Mayr. In domatia. Barro Colorado Is. C. Z.
- (23) *Solenopsis picca* Emery. In domatia of *C. alliodora* var. *boliviana*? Isiamas, Bolivia (W. M. Mann).
- (24) *Solenopsis ztcki* Wheeler. In domatia. Red Tank, C. Z.
- (25) *Leptothorax (Goniothorax) echinatinodis* Forel subsp. *cordincola* subsp. nov. In domatia. Red Tank, C. Z.
- (26) *Wasmannia auropunctata* Roger var. *nigricans* Emery. In domatia. Barro Colorado Is. C. Z.
- (27) *Cryptocerus (Cyathocephalus) setulifer* Emery. In domatia. Ancon, C. Z.; San José, Costa Rica (H. Schmidt).
- (28) *Cryptocerus (Cyathocephalus) pallens* Klug var. *porrasi* var. nov. In domatia. Frijoles, Red Tank and Quebrada de Oro, C. Z.
- (29) *Atta scydens* L. On foliage. Ancon, C. Z.
- (30) *Dolichoderus (Monacis) bispinosus* Oliv. In domatia. Red Tank, C. Z.
- (31) *Dolichoderus (Hypoclinca) championi* Forel subsp. *trinidadensis* Forel var. *tæniatus* Forel. In domatia. Frijoles, C. Z.
- (32) *Azteca fasciata* Emery subsp. *læta* Wheeler. In domatia. Gamboa, C. Z. (a single female).
- (33) *Azteca foreli* Emery var. *xysticola* Forel. Running on bark and nesting in trunk. Ancon, C. Z.
- (34) *Azteca instabilis* Emery var. Nesting in trunk. Barro Colorado Island, C. Z.
- (35) *Azteca bicolor* Emery var. Incipient colony in domatium. Ancon, C. Z.
- (36) *Azteca longiceps* Emery. In domatia. Ancon, Red Tank, Frijoles, Las Cascades, Barro Colorado Is. Agua Clara Reservoir, Gamboa, Corozal, etc. C. Z.
- (37) *Azteca longiceps* subsp. *balboæ* subsp. nov. In domatium. Balboa, C. Z. A single dealated female.

- (38) *Azteca longiceps* subsp. *cordineola* Forel. In "galls" of *Cordia*. Bolivia (Chodat). In domatia of *C. alliodora* var. *boliviana*? Huachi Beni, Bolivia (W. M. Mann)
- (39) *Azteca longiceps* subsp. *patruclis* Forel. In domatia. Cualata, Colima, Mexico (C. H. T. Townsend)
- (40) *Azteca pittieri* Forel. In domatia. Tumba Muerta, Panama; Red Tank, C. Z. San José, Costa Rica (H. Schmitt).
- (41) *Azteca pittieri* Forel var. *emarginatisquamis* Forel. In domatia. Costa Rica (Pittier).
- (42) *Azteca trigona* Emery. In domatia. Ancon and Las Cascades, C. Z.
- (43) *Azteca velox* Forel. In domatia. Ancon and Marajal, near Colon, C. Z.
- (44) *Tapinoma canalis* Wheeler. In domatia. Barro Colorado Is. C. Z.
- (45) *Brachymyrmex heeri* Forel var. *obscurior* Forel. In domatia. Barro Colorado Is. C. Z.
- (46) *Camponotus (Tanæmyrmex) coruscus* F. Sm. Exploring foliage. Ancon, C. Z.
- (47) *Camponotus (Myrmosphincta) 6-guttatus* Fabr. var. *bimaculatus* F. Smith. In domatia. Barro Colorado Is. C. Z.
- (48) *Camponotus (Ncomyrmamblys) novogrenadensis* Mayr. In domatia. Quebrada de Oro, C. Z. (with *Microdon* puparia).
- (49) *Camponotus (Myrmobrachys) canescens* Mayr. In domatia. Barro Colorado Is. C. Z.
- (50) *Camponotus (Myrmobrachys) brettesi* Forel. In domatia. Ancon C. Z.
- (51) *Camponotus (Myrmobrachys) lindigi* Forel. Exploring foliage. Ancon, C. Z.
- (52) *Camponotus (Myrmobrachys) brevis* Forel. In domatia. Red Tank, Gamboa, Frijoles, Las Cascades and Barro Colorado Is. C. Z.
- (53) *Camponotus (Myrmocladæcus) bidens* Mayr. In domatia. Red Tank, C. Z.
- (54) *Paratrechina longicornis* Latr. Exploring foliage. Ancon, C. Z.

Of the 54 different forms in this list 44 were found nesting in the domatia; the remaining 10 were either nesting in the trunk (*Azteca xysticola* and *instabilis*) or in the ground (the species of *Ectatomma*, *Atta*, *Paratrechina* and some species of *Camponotus*) and merely visited the foliage to attend Aphids or Coccids or to forage for insects. *Atta sexdens* was not seen to cut the leaves, which seem not to be in-

jured by this ant, probably on account of their coarse texture. The great majority of the domatia tenants occur also in dead twigs of a great variety of trees and shrubs. Probably only four of the species listed, namely, *Azteca longiceps* and its subspecies, *A. pittieri* and its var. *emarginatisquamis*, *Pseudomyrma sericea* and its varieties *ita* and *cordia*, and *Ps. alliodora*, are to be regarded as obligate tenants of the plant. Of these *A. longiceps* is the most abundant and occurs in about 85% of the cauline swellings. Forel cites *Pseudomyrma chodati* as living in "galls" of *C. longituba* in Paraguay and *Ps. sericea* var. *cordia* from an undetermined *Cordia*. If this plant belongs to the *Gerascanthus* section we may say that its members are inhabited by at least 60 different species, subspecies and varieties of ants. Indeed, the number is probably much greater, because the ants inhabiting these *Cordias* have not been intensively collected, except in a portion of the Panamanian region.

For the purpose of studying the ants of *C. alliodora* Mr. James Zetek and I adopted a method, which we also employed successfully, with obvious modifications, in dealing with other myrmecophytes (*Triplaris*, *Acacia*, *Cecropia*, *Clerodendron*, *Tillandsia*). We either cut down the tree, or when this was impracticable, lopped off large branches. Then with a pair of strong pruning scissors we cut out the cauline swellings and carried them in cloth bags to the laboratory, where they were placed in a large jar. Some chloroform was poured on the bags and the jar covered till the insects were asphyxiated. The nodes could then be cut open and their contents examined at leisure. We found that few of the ants left their nests to die in the bags and that the swellings, even when they were inhabited by different species, contained all or nearly all of their regular inhabitants. The preliminary work of ascertaining the qualitative composition of the *alliodora* biocœnose was so time-consuming that a quantitative or statistical study of the various species could not be attempted. It is to be hoped that some future student will feel inclined to undertake such an investigation at the new tropical laboratory on Barro Colorado Island.

A number of the *alliodora* ants, especially those of the genera *Crematogaster*, *Leptothorax*, *Cryptocerus*, *Tapinoma* and *Camponotus*, are very sporadic, occurring in only a few of the swellings on a tree or branch. *Azteca longiceps* is certainly the common and dominant tenant in nearly all the localities in which I collected. It usually occupies most or all of the swellings, especially those at the bases of the branches, whereas the sporadic species inhabit by preference the terminal and especially the dead and dried swellings that bore the

inflorescences of the previous dry season. Not only *A. longiceps* and *pittieri* and the varieties of *Pseudomyrma sericea* but also several of the other forms keep living Coccids in their nest-cavities. I here insert a fuller description of the habits of *A. longiceps* as the most typical of the *alliodora* tenants and append brief notes on some of the other ants.

Unlike the larger aggressive Aztecas (*trigona*, *velox*, *foreli*, *instabilis*, etc.) which either make large pendent carton nests on various trees or form compact and populous colonies in their trunks, *longiceps* is a small, timid and rather lethargic ant. This is indicated both by its toleration of other ant-tenants on the same tree and by the fact that I have sometimes cut up *Cordias* for hours without being bitten more than half a dozen times by the larger workers. The *longiceps* inhabitants of all the domatia on a tree constitute a single polycladic colony, which keeps growing and spreading by successive occupation of the new swellings as fast as they attain the proper size on the developing branches. During March and April the nests contain much brood in all stages together with many males and winged females. The domatial cavities are lined with a thin layer of brownish or blackish substance and contain a black or dark brown mass of carton made up of a network of traveculæ like those constructed by many other Aztecas that nest in plant cavities. This structure, shown in Plates 7 and 8, was seen by Beccari and Chodat and Vischer, but in their figures it is represented as grosser or more massive. It consists of very finely and uniformly triturated and agglutinated particles of wood and pith. I have failed to detect in it any pollen-grains, leaf-fragments or stellate hairs but would not deny that such substances may, perhaps, be occasionally employed by the ants in the confection of the mass. It is obviously a kind of scaffolding which subdivides the original cavity of the domatium into smaller compartments and galleries in which the brood can be spread out and more easily cared for. The mass can be readily removed in its entirety because it is rather feebly attached to the walls of the cavity. Chodat's and Vischer's contention that the ants, "dévorent une partie des feuilles et récoltent le pollen" is highly improbable. I have never seen the Aztecas visiting the flowers and they certainly do not devour the leaves. But even if this were true, and if the carton were made of leaf material as these authors maintain, the combined mass of carton in all the domatia on a tree would be too small to represent any serious damage to the plant.

In the spaces surrounding the mass of carton and sometimes almost covering the walls of the cavity are the numerous Coccids among

which at least three kinds or species may be readily distinguished. The majority are flat Lecanoid forms of a pinkish color and varying considerably in size. Among them may be found small snow-white Pseudococci, either singly or in clusters and several large subglobular shining black or red forms belonging to three species of the genus *Cryptostigma*.

These Coccids suggest interesting problems and reflections. They are present in such numbers that they must provide the ants with a copious supply of honey-dew. That the large subglobular forms (*Cryptostigma*) breed in the domatia is indicated by the fact that they are often found to be filled with eggs and owing to their size are quite unable to escape to the surface of the plant through the tenuous openings in the walls of the domatia. But whether the coccid colonies are originally established by young individuals that crawl into the swellings from the surface of the plant or are carried in by the ants, cannot be decided without further observation. Judging from what is known of some other ants (*Lasius* species, *Iridomyrmex humilis*, etc.), the latter alternative would seem to be the more probable. I have seen what I take to be the male pupæ of one of the large *Cryptostigma* species enclosed in peculiar flattened, bivalve-like cases, embedded in the central mass of carton. The Pseudococci are often destroyed by the larvæ of a small Chalcidid of the genus *Blepyrus* very much like *B. tachigaliæ* Brues which I found infesting *Pseudococcus brevipes* in the petiolar enlargements of *Tachigalia paniculata* Aublet in British Guiana.

There is also in the cavity of each domatium occupied by *A. longiceps* another singular object which has been overlooked by previous observers. The funnel-shaped lower end of the cavity is filled with a small conical plug of moist substance, which can be readily removed as a coherent mass and on examination proves to have a very complex structure, consisting of the ejected infrabuccal pellets of the ants, moulds and innumerable bacteria and Nematode worms. Its more liquid portion is probably the fæces of the ants and such honey-dew from the Coccids as happens to drain down the walls of the cavity and has not been intercepted and imbibed at its source. We may therefore regard the lowermost funnel-shaped end of the domatium cavity as a veritable public latrine or cess-pool. The Nematodes have been studied by Dr. Cobb who informs me that they are unusual from the taxonomic point of view. The moulds and bacteria which flourish in the fæcal medium of the latrines may afford an interesting study for some future investigator.

In Mexico the subsp. *patruelis* and in Bolivia the subsp. *cordicola*

seem to replace the typical *A. longiceps* as the obligate ants of *C. alliodora*. But even in Panama the typical *longiceps* may be replaced by *A. pittieri*, an ant of very similar appearance and habits. I found this to be the case in a small piece of jungle on the Tumba Muerta Road, near Las Sabanas. According to my observations, *Pseudomyrma sericea* var. *ita* is much less important as a member of the *alliodora* biocœnose. Its colonies are small and usually occupy only a few of the domatia on trees infested with *A. longiceps*. The same is true of *Ps. gracilis* and its var. *bicolor* which are more frequently found in hollow twigs of various trees and shrubs. *Ps. alliodora* is a timid form of rather sporadic occurrence. The older accounts of Ruiz and Pavon and Spruce seem to imply that *Ps. sericea* or other species of the same genus may be abundant and more formidable tenants of the *Cordias* in certain parts of South America.

The species of *Cryptocerus* are sluggish, cowardly and harmless. Both of the forms cited above, *setulifer* and *pallens* var. *porrasi* may occupy occasional swellings on trees infested with other ants. They belong to the subgenus *Cyathcephalus*, the females and soldiers of which have the top of the head peculiarly modified in the form of a large elliptical or suborbicular dish. A swelling occupied by *Cryptocerus* can be at once recognized by its entrance hole, which unlike that made by the Aztecas and *Pseudomyrmas*, is large and regularly elliptical instead of being small and round. The mother queen of the incipient colony or in older colonies one of the soldiers, constantly stands guard at the entrance and occludes it with the dish-shaped top of her head. This behavior is exactly like that of *Colobopsis*, except that in the ants of this subgenus, the portions of the head employed as a door are the truncated, orbicular front and cheeks. In the mother queen and older soldiers of *Cryptocerus* colonies the cephalic dish has been so persistently used in the manner described, that its concavity is often found thickly incrustated with grayish green foreign matter and thus comes to resemble the dull gray lichen or alga covered bark of the *Cordia*. The two species of *Cryptocerus*, like several other ants enumerated in the list on p. 25, are merely inquilines which often nest in the dead twigs and branches of other trees and shrubs.

That *Cordia alliodora* can derive little protection from the numerous ants which it harbors, is obvious from my account of their disposition and behavior. There is further evidence of this inefficiency in the great number of insects and other organisms that infest the foliage and flowers of the plant. Although Mr. Zetek and I devoted much attention to these forms, we were soon convinced that it would require

many months or even years to gain an adequate knowledge of the *alliodora* biocenose and the interrelationships of its numerous components even in so limited an area as the Canal Zone. The following list of species hitherto observed, however, with notes on some of the more interesting forms, will constitute a sufficiently formidable rebuttal of the argument that *C. alliodora* would probably perish in the struggle for existence if it failed to attract and provide quarters for battalions of ants.

Lepidoptera. At least six different species of moth caterpillars feed on the leaves of *C. alliodora* and some of them are rather destructive. One is a small leaf-miner of which I have seen only the small blister-like feeding-spots. Of another moth I have seen only the beautiful yellowish, regularly net-like cocoon spun in a folded leaf and containing an empty chrysalis. Three slender green caterpillars belonging to as many different species also remain to be reared and identified by some future observer. One of these larvæ which is fairly abundant bites a big hole in the wall of a young green domatium and hides in its cavity during the day. At night it eats the terminal leaves and spins their remains together about the domatium, leaving its frass among them. This untidy insect is very destructive to the terminal shoots of the *Cordia*. In a domatium containing one of these larvæ I found two small white Hymenopterous pupæ but did not succeed in rearing the adult parasites. The numerous caterpillars of a sixth species are about an inch long, chocolate brown with a pale mid-dorsal streak. They have been reared by Mr. Zetek who sent me specimens and photographs (Plate 9) of the cocoon, caterpillar and its method of feeding and hiding on the *Cordia* leaves. The adult moth, which is silvery white with black markings, is a Pyralid and was identified by Mr. W. Schauss as *Conchylodes salamisalis* Druce. Its life history is sketched in the following note contributed by Mr. Zetek: "The larva makes a sort of silken retreat at the base of the upper surface of the leaf by binding threads crosswise, thus forming a triangular cavity in which it can lie concealed. There is only one larva to a leaf. On one small plant, about five feet tall I saw 15 such larvæ. When ready to pupate, it spins a capsule $7/8$ in. long by $5/6$ in. in diameter, rounded at both ends and so well sealed that it is difficult to detect the sutures. After completing its cocoon it takes the larva about 36 hours to become a chrysalis. During this time, especially if the case be disturbed, the larva moves so violently inside, as actually to make the cocoon bound up an inch. The pupal stage last 12 days. Adults emerged June 12, 1923." During 1924 I found *Conchylodes* caterpillars on

many young *Cordias* at Ancon, Red Tank and Gamboa. During June the caterpillars were mature and ready to pupate and by the middle of July there were many young caterpillars of the next brood. Probably there are several generations on the young plants during the course of the year. The adult *Cordias* are immune from their attack during the rainy season, because they are leafless at that time.

Diptera. Numerous flies may be observed on the leaves of the young *Cordias* but are probably mostly chance visitors that alight on any low vegetation. Among those captured at Ancon and on Barro Colorado I recognized 3 species of Stratiomyids, 2 Syrphids, 1 Dolichopodid, 1 Micropezid, 2 Orthalids and 3 Muscids. The *Cordia* leaves are often dotted with small, spherical, densely pilose galls, which contain orange-colored Cecidomyid larvæ. I did not succeed in rearing the adults. More interesting is a species of *Microdon*, which has been described by Dr. Mann (1928) as *M. wheeleri*, since it lives with one of the ants in the domatia. March 10 I found at Red Tank, C. Z. in a *Cordia* swelling inhabited by *Crematogaster* (*Orthocrema*) *brevispinosa* var. *tumulifera* four bright yellow puparia, which were smooth and cylindrical and quite unlike any *Microdon* puparia known to me. They were carefully kept in a Petri dish till March 27, when two of the flies emerged and proved to belong to a very small species, about 7 mm. long. On emerging the hairs on the thoracic dorsum and sides and dorsum of the abdominal segments were conspicuously golden yellow and to all appearances true trichomes, but 24 hours later they had turned black. The other pair emerged March 28 and were chloroformed in the golden phase. They proved to be females whereas the first pair were males, so that I am unable to decide whether the females also blacken like the males. This is important in connection with Mann's description of the female. Unfortunately I failed to preserve the domatium in which the *Microdon* puparia were found although I kept all the ants and Coccids. The latter belonged to one of the species of *Cryptostigma* so common in the cauline swellings, especially when they are inhabited by *Azteca longiceps*. Perhaps one of the openings in the wall of the swellings may have been large enough to admit an ovipositing *Microdon*, but I failed to notice such an opening and am therefore unable to throw any light on how the *Microdon* larvæ get into or how the flies escape from the domatium. Probably the phase with the golden trichomes just after emergence is in some way connected with securing a temporary immunity of the fly from the attacks of the ants, and perhaps the latter enlarge their nest-entrances to let the flies out, but this is pure speculation and is merely

set down here for the benefit of someone who may be fortunate enough to happen on this rare and singular insect again. The problem of the food of the *M. wheeleri* larvæ is also, of course, unsolved. Donisthorpe (1912, 1923) has shown that the larva of the common European *M. mutabilis* feeds on the infrabuccal pellets which the ants cast out in their nests, and no doubt our North American *Microdon* larvæ have the same diet, but that the tropical *Microdons* may have very different habits is shown by Borgmeier (1923), who has recently found the larvæ of a Brazilian species devouring the Coccids (*Pseudococcus inquilinus*) in the nests of the fire ant (*Solenopsis sævissima* var. *picea*). Perhaps *M. wheeleri* feeds on the coccids in the *Cordia* domatia.

Hymenoptera. The following species of this order have very diverse relations to the *Cordia* and its inhabitants.

(1) A single unidentified Tenthredinid taken on the foliage at Ancon.

(2) *Dicrophrys* sp. (C. T. Brues det.) An Ichneumonid taken occasionally on the foliage of young *Cordias* at Ancon and possibly a parasite of one of the caterpillars enumerated above.

(3) *Eurytoma* sp. (?) A small larva, apparently related to the one described and figured by Chodat and Vischer, was found living in the swollen stems of the flower panicles.

(4) *Blepyrus* sp. Living as an entoparasite of *Pseudococcus* in the cauline swellings inhabited by *Azteca longiceps*. I saw only the empty puparia but they were not uncommon and are, I am inclined to believe, identical with those of *B. tachygaliæ* Brues, which I found in the same Coccids living with the social beetle *Coccidotrophus socialis* in the petioles of *Tachigalia paniculata* of British Guiana (see my paper of 1921).

(5) A Pteromalid (Brues det.) bred from flowers at Red Tank, C. Z.

(6 and 7) At Ancon I found two small mud nests attached to the twigs of *Cordia* and from them reared two solitary wasps. One of them proved to be *Eumenes infernalis* Sauss. (J. Bequaert det.), the other a *Sceliphron*. The latter was so nearly destroyed by house-ants entering the breeding cage that more precise identification was impossible.

(8) March 28 at Frijoles I found a nest of a very interesting social wasp, *Polistella picteti* var. *wheeleri* Bequaert, among the leaves of a *Cordia* about 15 feet high. The nest was about 6 inches long and consisted of four small, rather irregular, pale gray paper combs partly enveloped by the leaves to which they were broadly attached. The wasps made not the slightest attempt to sting but retreated into

their nest. The structure was very similar to that figured by Ducke (1910, p. 473, Fig. 2) for *Polistella picteti* var. *bella* v. Ihering. Concerning an allied species *P. emortualis* Sauss. of the Guianas and Brazil, he writes: "It is myrmecophilous. I have never found it alone but always in the company of ants of the genus *Dolichoderus*. The nests are very similar to those of the ants mentioned, built on the leaves of the same small branch of the tree whose other leaves harbor the nests of the latter. I have even seen a colony of nests of the wasp and of the ant closely associated on the same leaf. It is surprising to note that the ants are very aggressive, whereas the wasps, though furnished with stings take refuge in the interior of the nest when it is touched." That *P. picteti* may also be myrmecophilous is indicated by the fact that all the swellings of the *Cordia* on which the nest occurred were occupied by vigorous colonies of *Azteca longiceps*.

(9) *Ceratina* sp. A beautiful metallic blue species nesting in a dead *Cordia* branch (Red Tank, C. Z.)

(10) A bee allied to *Anthidium* taken on the foliage of a *Cordia* (Ancon, C. Z.).

(11) The only Hymenopteron taken at the *Cordia* flowers was a stingless bee, *Melipona orbignyi* var. *jenningsi* Ckll (T. D. A. Cockerell det.)

Coleoptera. The series of beetles found associated with the *Cordia* comprises a considerable number of species. The following taken during 1923 at Ancon were kindly identified by Mr. H. S. Barber:

(1) *Ladoria desarmata* Muls. (Coccinellid) on foliage.

(2) *Spermophagus lineolatus* Mots (?) (Bruchid) on foliage.

(3) *Paratenetus tropicalis* Mots. (Tenebrionid) bred from dried blossoms.

(4) *Melanophthalma cartralis* Shp. (Lathridiid) bred from dried blossoms.

(5) *Autodice nympha* Bates (Cerambycid, Fisher det.) Taken by Mr. Zetek on foliage.

(6) *Psalidonota leprosa* Boh. (Chrysomelid) on foliage.

(7) *Eustylus sexguttatus* Champ. (Curculionid, E. A. Schwarz det.) on foliage.

(8) *Deretomus palmarum* Sharp (Curculionid) on flowers. According to Mr. Barber this beetle occurs also in great numbers on the flowers of palms.

During the summer of 1924 I captured on the foliage and branches of young *Cordias* at Ancon 30 additional species, which are still unidentified. These comprise 2 Cerambycids, 1 Telephorid, 2 Lampyrids,

1 Lathridiid, 1 Cucujid, 1 Endomychid, 12 Chrysomelids, 6 Curculionids, 1 Anthibid and 3 Platypodids (boring in a recently felled tree).

To this list must be added a second species of Coccidotrophus, *C. cordia* Barber (1928), an interesting social Silvanid beetle discovered by Dr. Mann in a single domatium of a *Cordia* (probably *alliodora*) at Huachi, on the Rio Beni, Bolivia. This beetle is somewhat larger than *C. socialis* Barb. & Schwarz, which I found nesting in the petioles of *Tachigalia paniculata* in British Guiana, but evidently has very similar habits, since Dr. Mann noticed that it was living with its larvæ and with Coccids (*Pseudococcus* sp.)

Many of the beetles taken at Ancon were probably chance visitors, but one of them, the Cassidine *Psalidonota leprosa*, passes through its entire development on *C. alliodora*, which is therefore to be regarded as its host plant. The beetle was originally described from Mexico (Boheman, 1855). According to Champion (1885-94), it is a common insect in Central America "ranging from the Mexican State of Durango right down to Panama, and probably extending into the northern parts of South America." He cites it from several Panamanian localities. Its range therefore coincides with that of *C. alliodora* except that it occurs, according to Leng (1920), also in Texas, where it must have a different host plant. The greenish eggs were found in late July and early August in clusters on the lower surfaces of the leaves of young *Cordias* at Ancon and Red Tank, and the larvæ were seen feeding on the same leaf-surfaces in clusters or when older more sporadically. The larva is flattish, pale green, with a blackish-brown mid-dorsal stripe that grows broader with age, especially in the thoracic region, and a fringe of blunt, branched spines along the sides of the body. The posterior end is enlarged into a peduncled sphere beset with knobs and with the protrusible anus at the base of the peduncle on the ventral side. As in other Cassidine larvæ, the modified caudal end can be turned forward over the back and supports a great glutinous black mass of fæces and exuviae, which is not, however, large enough to conceal the insect. The pupa is naked and attached to the underside of the leaf by its posterior end. The tortoise-like adult beetle, which is nearly a centimeter long and broad, was described by Boheman as "flavotestaceus", and Champion figures it of this color, which suggested the unpleasant specific name *leprosa*. This is not inappropriate for the dried specimen but is unfortunate, because the living insect is a magnificent creature, resembling nothing so much as a large drop of liquid gold. It is therefore a very conspicuous object when sunning itself on the deep green leaves of the *Cordia*. Since I have found it on the plant

also early in the dry season, it must have several broods during the year.

Orthoptera. The following species of this order were identified by Mr. Morgan Hebard:

(1 and 2) *Stagmomantis* (Mantid), two species, immature, on young foliage of *C. alliodora*.

(3) *Cryptostilum antillarum* Redt. (Gryllid), living in an abandoned domatium.

(4) *Gryllacris* sp. (Gryllid), immature near *picta* Bruner, on foliage.

(5) *Osmilia flavolimbata* De Geer (Acridiid). At Ancon both Mr. Zetek and I repeatedly took both sexes of this grasshopper in the act of devouring the foliage of *C. alliodora*.

(6) *Coscineuta coxalis* Serv. (Acridiid). A few specimens on the foliage of young plants.

(7) *Schistocerca* sp. (Acridiid), immature.

(8) *Aidemona azteca* (Saussure) (Acridiid) One female and one immature individual.

(9) *Orphulella concinnula* (Walker). (Acridiid) One female.

(10) *Neoconocephalus* sp. (Tettigoniid), immature.

(11) *Anaulacomera* sp. (Tettigoniid), immature.

(12) *Phaneroptera paronæ* (Griffini) (Tettigoniid) immature.

(13) *Phlugis* sp. (Tettigoniid), immature.

Thysanoptera. Two species of this order have been identified by Dr. J. Douglas Hood: *Diccratothrips armatus* Bagnall and *Elaphrothrips* sp. Both are large slender species, black in the adult and bright coral red in the younger stages, occurring in numbers in some of the dead and abandoned domatia at Ancon and on Barro Colorado Island. A third, unidentified species is a minute yellowish form, common on foliage and flowers at Ancon.

Neuroptera. Two species are represented in the collection, one a species of *Chrysopa* of which Mr. Zetek on several occasions found the eggs attached to the leaves of young *Cordias*, the other a species of Hemerobiid, the larvæ of which I have taken on several occasions in the same situations. Both insects probably prey on the Aphids and Aleurodids.

Heteroptera. The following species of this order have been kindly identified by Dr. J. G. Myers and Dr. R. F. Hussey.

(1) *Proxys punctulatus* Pal. Beauv. (Pentatomid) Ancon (Myers det.)

(2) *Edessa collaris* Dall. (Pentatomid) (Myers det.) Ancon. This bug is common in all stages on the plant (except the eggs, which were

not seen). The nymphs and adults are green in life. Undoubtedly *C. alliodora* is a normal host plant of this insect.

(3) *Hypselonotus fulvus* DeG. var. *venosus* Fabr. (Coreid) Ancon. Range: Colombia, Venezuela, Surinam (Hussey det.)

(4) *Hyalijmenus pulcher* Stal (Coreid) Las Sabanas, Panama. (Myers det.)

(5) *Dysdercus ruficollis* L. (Pyrrhocorid) Ancon (Myers det.)

(6) *Dysdercus obliquus* H S. (?) (Pyrrhocorid) Ancon. Range: California to Ecuador. (Hussey det.)

(7) *Monanthia monotropidia* Stal. (Tingitid) Ancon and Red Tank. (Myers det.) This insect feeds in great numbers and in all stages on the under surfaces of the leaves of *C. alliodora* and damages them severely, causing them to turn black. It seems to have a wide distribution. Wolcott (1923) found the "nymphs and adults abundant on the underside of leaves of a small unidentified tree in the mountains north of Yanco", Porto Rico.

(8) *Macrocephalus notatus* Westw. (Phymatid) Ancon (Myers det.)

(9) *Zelus nugax* Stal. (Reduviid) Ancon (Myers det.)

(10) *Eccritotarsus splendens* Distant. (Mirid). Ancon (Myers det.)

Homoptera. The following species have been identified by Dr. Myers and Dr. W. D. Funkhauser:

(1) *Moneophora lepidior* Fowler (Cercopid) Ancon, feeding on young shoots (Myers det.)

(2) *Micrutalis balteata* Fairm. (Membracid) Ancon, feeding on young shoots. (Funkhauser det.)

(3) *Micrutalis dubia* Fowl. (?) (Membracid) Ancon, feeding on young shoots, (Myers det.)

(4) *Ceresa bubalus* (Fabr.) (Membracid) Ancon (Myers det.)

(5) *Vanduzeeia triguttata* Burm. (Membracid) Ancon (Funkhauser det.)

(6) *Brachybelus cruralis* Stal (Membracid) Ancon (Myers det.)

(7) *Oncometopia undata* (Fabr.) (Cicadellid) Ancon. Evidently feeds on *Cordia*.

(8) *Cicadella scxguttata* (Fabr.) (Cicadellid) Ancon (Myers det.)

(9) *Cicadella rufimargo* (Walk.) (?) (Cicadellid) Ancon (Myers det.)

(10) *Gypona obscurior* (Fowler) (Gyponid) Ancon (Myers det.)

(11) *Gypona postica* Walk. (Gyponid) Ancon (Myers det.)

(12) *Athysanus bicolor* Van Duzee. (Jassid) Gamboa (Myers det.)

(13) *Deltocephalus* sp. (Jassid) Ancon (Myers det.)

(14) *Protalebra* sp. (Typhlocybid) Ancon (Myers det.)

(15) *Rudia diluta* Stal. (Tropiduchid) Ancon (Myers det.)

- (16) *Cyrpoptus suavis* Stal. (Fulgorid) Ancon (Myers det.)
- (17) *Colpoptera sinuata* Burm. (Issid) Ancon (Myers det.)
- (18) *Ormenis griseoalba* Fowler. (Flatid) Gamboa (Myers det.)
- (19) *Psyllid* indet. (nymphal exuviae) Ancon (Myers det.)
- (20) *Aphis* sp. (A. C. Baker det.)

This species was taken in great numbers on *Cordia* in November 1923 by Mr. Zetek. The insects caused the leaves to crinkle and were attended by the common crazy ant, *Paratrechina longicornis* Latr.

(21) *Aleurodicus dugesii* Ckll. (?) (Aleurodid) Ancon. Also taken late in November 1923 on the leaves of *C. alliodora* (A. C. Baker det.)

Coccidæ. The following species were recognized by Dr. H. Morrison from extensive collections made in 1923 and 1924.

(1) *Akermes cordiæ* Morrison. In domatia with *Azteca longiceps* and *Cryptocerus pallens*. Red Tank and Ancon.

(2) *Coccus hesperidum* (Linn.) On leaves and twigs, attended by *Azteca velox*. Ancon.

(3) *Cryptostigma biorbiculum* Morr. In domatia with *Pseudomyrma ita* and *Azteca longiceps*. Ancon and Frijoles.

(4) *Cryptostigma secretum* Morrison. In domatia with *A. longiceps*. Ancon.

(5) *Cryptostigma reticulolaminæ* Morrison. In domatia with *A. longiceps*. Ancon and Frijoles.

(6) *Cyclolccanium hyperbaterum* Morrison. In domatia with *Azteca longiceps* and *Pseudomyrma ita*, Red Tank. and Ancon; with *Camponotus brettesi*, Barro Colorado Island, and Frijoles; with *Azteca pittieri*, Pueblo Nuevo, Panama; with *Crematogaster tumulifera*, Red Tank.

(7) *Pseudococcus brevipex* Ckll. (previously *bromeliæ* Bouché). In domatia with *Azteca longiceps*. Red Tank and Las Cascades. The Coccid occurs also in the cauline cavities of *Triplaris* and *Cecropia*.

(8) *Pseudococcus maritimus* (Ehrh.) In a young domatium of *Cordia*, without ants. Red Tank.

(9) *Pseudococcus probrevipes* Morrison. In domatia with *Pseudomyrma ita*, *Crematogaster sumichrasti* and especially *Azteca longiceps*. Ancon, Red Tank, Quebrada de Oro, Las Cascades, Barro Colorado Is. & Frijoles.

(10) *Pseudococcus* sp. immature. In domatia with *Pseudomyrma gracilis*. Red Tank.

(11) *Saissetia hemisphaerica* Tang. Covering the twigs and domatia of young *Cordias* and injuring them severely. Collected by Mr. Zetek at Ancon June 15/26.

(12) *Saissetia nigra* (Nietn.) On leaves.

(13) *Saissetia oleæ* (Bern.) On leaves.

(14) *Aspidiotus perculcanus* Doane & Hadden. On leaves.

Myriopoda. Two species of Myriopoda were found occupying old and abandoned domatia. These were identified by Dr. R. V. Chamberlin as *Orphæus brevilabiatus* Newp., a Chilopoda, taken at Red Tank, and *Orthomorpha gracilis* Koch taken at Ancon.

Arachnida. The following spiders, kindly identified by Mr. Nathan Banks, were taken on the foliage of young *Cordias* at Ancon and on Barro Colorado Island. Many of them were living in webs spun between the leaves or twigs:

(1) *Gasteracantha kochi* Butler.

(2) *Frontinella unciata* Cb.

(3) *Chrysso vexabilis* Keys. This small red and black spider is unusually common on the plant.

(4) *Runcinia magna* Keys. (?)

(5) *Selenops mexicanus* Keys.

(6) *Nephila clavipes* L.

(7) *Zygoballus tibialis* Cambr.

(8) *Dendrophautes momus* Cambr.

(9) *Teudis roseus* Cambr.

(10) *Phyale simplicicava* Cambr. (?)

(11) *Pisaurid* gen. incert. very young.

(12-14) *Phyale*, 3 species, immature.

(15) *Dictyna* sp., immature.

(16) *Philodromus* sp., immature.

(17) Nov. gen. nov. sp. near *Simonella*.

(18) *Mites*, belonging to the group commonly called "red spiders" were found by Mr. Zetek during November 1923 on both sides of the main ribs and riblets of the leaves of young *Cordias* at Ancon. "They leave a patch which is almost white,—typical of red spider injury. They were very abundant and lived on the same leaves with the Aphids."

Isopoda. A few small white Isopods related to *Oniscus* were found at Ancon inhabiting an abandoned domatium.

Nematodes. Several peculiar species taken in the latrines of the domatia have not yet been described by Dr. Cobb.

Fungi and Bacteria. In addition to the moulds which grow on the walls of the domatia there is a powdery mildew of the family Erysiphaceæ which often covers the leaves of young *Cordias*. Specimens collected by Mr. Zetek at Ancon could not be further identified

by Prof. W. H. Weston, to whom I sent them, because the fruiting bodies were absent. The bacteriology of the domatial latrines still remains to be investigated.

Algæ and Lichens. Like other tropical trees, *C. alliodora* supports a flora of lichens and algæ on its bark.

If we consider only the Arthropoda noticed in the preceding pages as visiting or infesting *Cordia alliodora* we have the following numbers of forms in each of the larger groups:

Hymenoptera (including Formicidæ)	69
Lepidoptera	6
Diptera	14
Coleoptera	39
Orthoptera	13
Thysanoptera	3
Neuroptera	2
Heteroptera	10
Homoptera (including Coccidæ)	35
Myriopoda	2
Arachnida	18
Isopoda	1

Total 212

In all probability this total of more than 200 forms or species represents only a fragment of the biocænose centering about *C. alliodora*, because my observations were confined to a very small portion of its known geographical range. The data are nevertheless sufficiently numerous and striking to admit of certain conclusions, especially as they are supported by similar observations on other ant-plants (*Tachigalis*, *Cecropia*, *Acacia*, etc.) to be considered in the sequel. There can be no doubt that *C. alliodora* has many insect enemies, which destroy or deform its foliage, injure its terminal shoots and withdraw a considerable amount of sap from its tissues. The most numerous and serious of these pests are the moth *Conchylodes salamisalis*, the Cassidine beetle, *Psalidonota leprosa*, the Pentatomid bug *Edessa collaris*, the Tingitid *Monanthia monotropidia*, a red spider, the various Membracids, Cicadellids and Jassids, the Aphis, the Aleurodid *Aleurodicus dugesii* and last but not least, the fourteen species of Coccidæ recognized by Morrison. No evidence was found to indicate that any of the 54 forms of Formicidæ attacked any of the leaf-eating species, though it was evident that they solicitously

attended many of the Coccids to the injury of the plant. Moreover, the majority of the 212 Arthropods, excluding the Formicidæ, were found on young *Cordias*, the domatia of which were as yet either uninhabited or occupied only by recently fecundated, colony-founding queens of *Azteca longiceps* and *pittieri*. Hence the plants can derive no protection from these insects in the very stages when the incidence of natural selection should be most effective. Yet none of the thousands of young and seedling *Cordias* which I examined in Panama showed the slightest signs of being killed by its insect pests. I have no hesitancy in asserting that the plant is quite as vigorous and quite as able to withstand the attacks of insects as any other common tropical tree.

Chapter 2

OBSERVATIONS ON TRIPLARIS

The diœcious Polygonaceous trees of the genus *Triplaris* and their ants made a more vivid impression than the *Cordias* on the early naturalists who visited the American tropics. That the natives and colonists had long been duly impressed is indicated by the many vernacular names they invented for the plants and their aggressive tenants. As long ago as 1849 Weddell recognized 14 species of *Triplaris* and in 1856 Meisner enumerated some 25; Hemsley in 1882-86 and Dammer in 1893, however, recognized only about 10. Probably the latter number is not far from the present estimate of actually existing species, but it must be admitted that there is as much confusion in the botanical literature in regard to *Triplaris* as to many other genera of Neotropical woody plants. The *Triplaris*es certainly appear to be more local than the *Cordias* of the *Gerascanthus* group and are confined to the warmest portions of South and Central America, where they grow by preference in low or even swampy places along streams. One species (*T. auriculata*) occurs as far north as Mexico. The genus is not represented in the West Indies, though it may occur on several of the islands adjacent to Northern South America, e.g., Trinidad. Boldingh (1914) cites *T. coriacea* Karst. as growing in several localities on Curaçao, off the coast of Venezuela.

(A.) Historical

One of the earliest accounts of *Triplaris* and its ants is that of P. Bernabé Cobo (1653), who in his "History of the New World" says that the tree is called "palo santo" and describes it as follows: "This

tree is hollow throughout from the trunk to the slenderest twigs and full of certain yellow and largish ants, so virulent that their sting is apt to bring on fever and is always exceedingly painful. Since these ants are concealed within the tree, they are not seen and this is the reason why those who do not know the secret, are not on their guard; but if a single leaf be touched, so many of the ants swarm forth from all parts of the tree as to excite wonder, and they assail the person who touches the tree and, if he does not withdraw in time, martyr him with their stings." The editor adds the scientific name of the ant as *Myrmica triplarina* and its vernacular name as "*hormiga tangarana*"; the copyist cites the name of the tree as "*Guyacum sanctum*, according to Raimondi."

In the botanical literature Aublet (1775) seems to have been the first to notice the peculiar behavior of the ants and their relation to the tree. Writing of *T. americana*, which he described and figured (p. 910, Pl. 347), he says that the natives of French Guiana call it the "sapa-hakaapolli", and adds: "The ants swarm abundantly throughout the interior of the trunk, the branches and twigs of this tree, in such a manner that when one fells it, one is at once completely covered and cruelly tormented by them, a misfortune which befell me. The only way to get rid of them is to throw oneself into the water."

In the famous work on Brazil by von Spix and von Martius (1831), I find the following note on the species of *Tococa* and *Triplaris americana*: "A number of plants, especially those of the genus *Tococa*, seem to have been designed by Nature herself as domiciles for ants. These bushes bear at the upper end of the leaf-petioles a bladder-like enlargement, in which numerous communities of small red ants nest, and the hollow branches of *Triplaris americana* L., a slender littoral tree, are often inhabited by innumerable colonies of similar creatures. Woe to him who chances to break off such a branch; a scrambling mass of viciously biting enemies pours down upon him and leaves many blisters on his skin."

Robert Schomburgk (1838) observed the same species of *Triplaris* in British Guiana. He says that it is called the "jacuna" by the Arawak Indians, the ant being the "jacuna sac", but that the Warrows call the tree the "epouchari", the Caribis the "itassi" and the colonists the "long John". I find that this name is still applied to both *T. americana* and *T. surinamensis* by the settlers in British Guiana. The following is Robert Schomburgk's general description of the former species: "The sandy banks of the inland rivers of Guiana are peopled with them; and when shrubs, stunted in growth by the poverty of the

soil, scarcely reach the height of five or six feet, the *Triplaris* overtops them forty or fifty feet. The trunk is slender and grows up straight, and its erect branches form a pyramid. As already observed, it is unisexual, and the flowers of both sexes are insignificant: those of the male last only for a few days, when they dry up; this is likewise the case with the petals of the female; the segments of the calyx however continue to grow, changing in their growth from green to white and vermillion, and become so attenuated that the branched nerves are easily perceptible. In that state they are three times as large as the fruit, which is still protected by the tube of the calyx, and the whole might in appearance be resembled to a shuttle-cock. The risps are dense, and the tree presents now a most elegant appearance. One unacquainted with the contrary, would consider the tree covered with white blossoms, tinged with red, among which the dark green leaves have only occasionally room to make themselves visible. The uncautious botanist, who, allured by the deceptive appearance, should approach the tree to pluck the blossoms, would bitterly rue his attempt. The trunk and branches of the tree are hollow, like those of the trumpet-tree (*Cecropia*), and provided between space and space with partitions, which answer to the position of the leaves on the outside.

"These hollows are inhabited by a light brownish ant, about two to three-tenths of an inch long, which inflicts the most painful bites. Its antennæ are placed near the middle of the anterior portion of the head; mandibles triangular; peduncle of the abdomen with two rings; the anus hairy and provided with a sting or piercer (*Myrmica* Latr. nova species). They fall upon their prey with the greatest virulence, and insert their mandibles almost instantly, as soon as they come in contact with any soft substance, emitting a whitish fluid; their bite causes swelling and itching for several days. If they find themselves captured, they attack and kill one another like the scorpions." The ant to which Schomburgk refers is undoubtedly a species of *Pseudomyrma* but he makes the common mistake of confusing biting and stinging; the *Pseudomyrmas* do both, but the pain is, of course, due to the sting.

Richard Schomburgk (1848) gives an even more vivid account of *T. americana*, which he encountered on the Barama River in British Guiana and, according to his estimate, may grow to a height of 60 to 80 feet. "The peculiar internal structure of the trunk and branches make it one of the most formidable of trees. The internodes are perfectly hollow and separated from one another by horizontal partitions, so that in this respect the tree resembles *Cecropia peltata*."

He correctly describes the yellowish brown *Pseudomyrmas* and their stinging habits but erroneously refers them to the genus *Cryptocerus*. "Being unacquainted with the structure of the tree and its formidable inhabitants, and ignoring the warning gesticulations of my Waraus, I was trying to break off one of its boughs, when thousands of these insects rushed out of the small round openings in the internodes, completely covered me and in the greatest fury seized my skin with their jaws and, vomiting a white liquid, buried their terrible stings in my muscles. But not only had the ants from the severed portion of the bough fallen into our corial, but thousands more poured out of the openings in the stump and rained down into the boat since the whole colony had been aroused by the shaking of the tree. A few powerful strokes of the oars carried the boat out of the neighborhood of the tree and in the twinkling of an eye the whole crew was in the water, for only thus could we escape from the savage onslaughts of the ants. Even a few tame apes and parrots were not spared. The former with wild leaps freed themselves from their tethers and jumped into the river after us, although few animals are more averse to water. The sting of this yellow-brown ant is only less painful than that of *Ponera clarata*. The swelling, inflammation and pain may persist for several days . . . After we had cleaned the ants out of the boat with considerable labor and many excruciating stings, we continued our voyage. I must confess that thereafter a secret horror crept over me whenever we passed one of the trees."

In 1849 Weddell described eight species of *Triplaris* from various parts of South America and published a key for the identification of the known forms. The following account, concealed in a footnote recently detected by Dr. J. Bequaert (1921-22), shows that Weddell was well-acquainted with the trees and their tenants: "The trunk, the branches and even the small twigs of the species of this genus are fistulose and serve as habitations for a peculiar species of ant which exhales when excited a rather agreeable odor like that of the *Cicindelids*. If one happens to touch the trunk of a *Triplaris*, and especially if one imparts a shock to it, the ants sally forth by the hundreds from the interior of the tree through the small galleries which connect the medulary canal with the outside; and if one move not away very quickly, one is covered with these dangerous hosts, whose bite is more painful than the sting of any other insect with which I am acquainted. It is a singular fact that at no matter what period of their life the *Triplaris* be examined in their forest, these ants are always sure to be encountered. It is singular also that in *Rupprechtia*,

which some authors unite with *Triplaris*, they are never found. I do not believe that this insect has been observed under conditions other than those I have noted. Its linear form is peculiarly adapted to its mode of life. I have had occasion to examine it and even to suffer from its attacks in many parts of Brazil, Bolivia and Peru, and everywhere it seemed to me to be identical. Several travellers have already made known a portion of the facts here related and have placed the *Triplaris* ant in the genus *Myrmica* of Latreille, but I am not aware that it has been given a specific name. That of *Myrmica triplarina* might be applied to it. It is usually pale brown. Its length is 6 to 7 millimeters, its width one millimeter; the abdomen is cylindrical and somewhat attenuated at the posterior end which is hairy." I believe that the ant here described is the same as the one now called *Pseudomyrma arborissanctæ* Emery, but this species has a number of local races and varieties and the brief description renders it impossible, unless Weddell's types should be discovered in some European museum to determine which of the described subspecies or varieties should bear the name *triplarina*. It seems best therefore to regard it merely as a somewhat doubtful synonym of some one of the forms, preferably of the typical *arboris-sanctæ*.

Spruce (1869, in 1908) mentions several lignescent genera of Neotropical Polygonaceæ as harboring ants in their medullar cavities—*Triplaris*, *Coccoloba*, *Campteria*, *Symmeria* and *Rupprechtia*—the last in contradiction to Weddell's statement quoted above.¹ "Not only is every lignescent *Polygonia* a habitation for ants, but the whole of the medulla of every plant, from the root nearly to the growing apex of the ramuli, is scooped out by those insects. The ants make a lodgment in the young stem of the tree or shrub, and as it increases in size and puts forth branch after branch, they extend their hollow ways through all its ramifications. They appear to belong all to a single genus, and are long and slender, with a fusiform, very fine-pointed, dark-coloured, shining abdomen, and they all sting virulently. They are known in Brazil by the name of "Tachi" or "Tacyba", and in Peru by that of "Tangarana"; and in both countries the same name is commonly applied to any tree they infest as to the ants themselves." Spruce describes three species of *Triplaris*. "*Triplaris surinamensis* Camb., a Polygonaceous tree of very rapid growth, reaching at maturity a hundred or more feet in height, and conspicuous from afar when in fruit from the abundance and bright

¹Apparently the only species of *Rupprechtia* which harbors ants in its internodes is *Jamesoni* Meisner. (Bequaert 1921-22)

red colour of its enlarged shuttle-cock-like calyces, is common all along the Amazon, both on the river banks and in marshy inland sites; and solitary trees of it are often seen standing out above the Cacao plantations. *T. Schomburgkiana* Benth., a smaller tree, grows in the same way in the Upper Orinoco and Casiquiari. These trees as well as the other arborescent Polygoneæ, have slender elongated tubular branches, often geniculate at the leaf-nodes, and nearly always with perforations, like pinholes, just within the stipule of each leaf, which are the sallyports of the garrison, whose sentinels are besides always pacing up and down the main trunk, as the incautious traveller finds to his cost when, invited by the smoothness of the bark, he ventures to lean his back against a Tachi tree." The third species of *Triplaris* allied to *surinamensis*, was observed in the cinchona forests of Chimborazo (2000–5000 ft.) and at lower levels. Spruce says that it is called by the Guyaquilians the "arbol de frios" (malaria tree), and that its presence "is a pretty sure indication of a humid site."

Huth (1887) states that H. von Ihering called his attention to the following account of the Brazilian *Triplaris nolitangere* Wedd. by J. Severiano da Fonseca (1881): "The Pao de novato is the Taixy of Pará and is also called Pau formiguero, or ant-tree. It is remarkable because it harbors in its cavities a kind of ant which is called 'novato'. These ants are yellow, as large as a 'suava' and bite painfully. They live there by the million and drive the inexperienced traveller to desperation when he attempts to fell and make use of the tall, straight trunks of the 'novato.' "

Schimper (1888) had no opportunity to study *Triplaris* in the field but examined several branches of *T. caracasana* Cham. and Schl. from the "tierra caliente" of Venezuela, a tree which is called the "palo Maria" by the natives.¹ He says: "The numerous pieces of branches sent to me show nothing that can be interpreted as an adaptation to the ants. Within the branches there is a cavity about 5 to 8 mm. broad, interrupted by diaphragms and connected, with the outside by round openings which are made by the ants. Of the latter there is almost without exception only one at each internode and usually at the upper end of a groove which runs to the next leaf below. The wall of the hollow cylinder is somewhat thinner at the groove, which is caused by the pressure of the bud and occurs also in many other plants. This thin area evidently determines the point

¹In some parts of South America (Bolivia) *Triplaris* is called the "palo santo". This and the name "palo Maria" are obviously of rather recent Christian origin. It has been claimed by some writers that the "santo" refers to Christ's blood, as symbolized by the massed crimson fruit of the tree.

of perforation, which in the great majority of cases is at the upper end of the groove and may be due to the ants selecting, when making their orifices, only those lenticels that are situated in the upper end of the internode."

Meinert (1892) published a brief note on a species of *Triplaris* (probably *caracasana*) and its ant, which he observed at La Moca, Venezuela. The ant was referred to the genus *Myrmica* but was obviously a *Pseudomyrma*. His statement that it is not confined to the *Triplaris* but occurs abundantly on other plants, has little weight, because there are many *Pseudomyrmas* which an observer unfamiliar with the species, might mistake for the true *Triplaris* inhabitant in the field.

Warming (1894), who was with Meinert in Venezuela, investigated a *Triplaris* (presumably *caracasana*) at Las Trincheras, concentrating his attention on the structure of the elliptical cleft near the upper end of each internode. This structure, as we have seen, was interpreted by Schimper as due to pressure exerted by the bud at the node below or as arising from a lenticel. Warming adopted the former interpretation but studied the internodes more closely. Like Schimper he found the wall of the internode to be thinner and to have more feebly developed fibrovascular bundles in the region of the cleft elliptical area and believed that this attenuation induces the ants to select the spot for perforation. But he noticed considerable irregularity in the perforations and their absence in some or all the internodes of whole twigs. He seems to have been the first to notice the numerous white Coccids which the ants keep in the cavities. The ants are briefly described in Warming's paper as "en lille, brun umaadeleg bidsk Art" (a little brown, intemperate, biting species) and referred to *Pseudomyrma mordax* Meinert. This, however, seems to be a *nomen nudum*. Forel (1905), who listed the ants collected by Meinert in Venezuela referred the specimens from *Triplaris* to *Ps. arboris-sanctæ* subsp. *symbiotica* Forel, a race originally described from Colombia.

Morteo (1904) studied preserved material of *Triplaris americana* which had been grown under cultivation in the East Indies, where its cavities were inhabited by a common Oriental ant, *Dolichoderus (Hypoclinca) bituberculatus* Mayr. He devoted particular attention to the peculiar cleft area near the tip of each internode, but interpreted the fissure as caused by expansion of the pith. He also believed that the ants perforate the cleft in order to feed on the pith, which he found to contain sugar. Like Warming, he found numerous Coccids in the in-

ternodal cavities. The discovery of sugar in the pith led him to regard it as a substitute for the extrafloral nectaries of other myrmecophilous plants. His observations, if correct, might be suggestive in connection with the presence of Coccids on the walls lined with the remains of the pith, but his statements in regard to the ants devouring it are not to be taken seriously, in view of the fact that his investigations were based on herbarium material.

Forel (1904) gives the following account of a species of *Triplaris* (probably *americana*) and its ant (*Ps. arboris-sanctæ* subsp. *symbiotica*) which he encountered in 1896 in the Sierra Nevada de Santa Marta, Colombia: "Having laid my hand on the trunk of a young, green tree about four meters high, I was stung and found that the aforementioned *Pseudomyrma* was present on the trunk and had caused the sting. Observing the aggressive behavior of these ants, I suspected that they bore a symbiotic relation to the tree, because other *Pseudomyrmas*, which run about on the trees, flee instead of attacking. But seeing no dry branch and no opening, I was puzzled at first. I accosted some passing Indians and induced them to fell the tree with their machetes. I then broke up its flexible and living branches and found that they had a very narrow medullary cavity. This cavity, extending from end to end of all the branches, constituted the nest which the *Pseudomyrmas* occupied in a file, one behind the other, with their males, larvæ and pupæ, so that they were just able to pass one another, notwithstanding the slenderness of their bodies. This singular habitation baffled me considerably and I asked myself how the colony-founding queen could have penetrated into this perfectly green tree, without a dead branch and apparently without an orifice. After long and vain scrutiny of all the branches, I inspected the lower portion of the trunk and there at last detected the dried and broken remains of an early twig, about 3 mm. in diameter but provided with a medullary cavity communicating with the central cavity of the trunk itself. This it was that served as an entrance and exit for the *Pseudomyrmas*." It would seem that Forel must have overlooked the small openings which were in all probability present in the tree he examined, as in all other species of *Triplaris*, near the ends of the internodes of the branches and twigs. The same author, after his description of *Pseudomyrma dendroica*, states that Prof. E. Goeldi first found this ant in the medullary cavities of young *Triplaris* trees three to four meters high, and that "having transported the *Triplaris* inhabited by this ant from the Rio Purus to the botanical garden at Pará, he observed that the *Pseudomyrma* soon took possession of one of the *Triplaris* in the garden, which had not

been inhabited before. The habits of this ant are therefore the same as those of *arborissanctæ*."

Ule (1907) describes *Triplaris Schomburgkiana* and *surinamensis* and gives some excellent figures of the former. The natives of the Amazon, he says, call these trees "arvore de tachi" or "tachiceiro", and distinguish the two species as white and black, *Schomburgkiana* having a pale and *surinamensis* a dark trunk. "The two species of *Triplaris* accordingly harbor in their interior a pale and a dark ant." *Schomburgkiana* is inhabited by *Pseudomyrma dendroica* var. *emarginata* Forel, which nests by preference in the twigs and the crown of the tree but retains a medullary gallery through the trunk, reaching to the ground and sending off lateral galleries at intervals. "The bite of this ant is very painful, burns as if one had come in contact with red-hot iron and sometimes produces blisters on the affected parts of the body. From the trunk the ants wander to the ground and there destroy all the growing vegetation within a radius of a few meters. The location of *Triplaris* trees can be detected at once by such bare spots in the forest." *T. surinamensis* is confined more to the banks of rivers and lakes. The darker ant which inhabits its internodes is *Ps. triplariidis* Forel. It does not make a cleared area around the trunk of the host tree. According to Ule, "the *Triplaris* trees are of some importance in the landscape. When they are in bloom, the male flowers first become conspicuous because they resemble great feather dusters, then the panicles of the female trees stand forth, when the fruits have developed their bright red wings, being vivid purple in *Triplaris Schomburgkiana* Benth. and rose-red in *Triplaris surinamensis*, so that the inhabitants believe that the trees are in flower. When I sailed up the Upper Amazon to Iquitos in July, the shores were everywhere brilliant with the *Triplaris* trees which with their rose-red fruits and the fresh green of the vegetation after the floods, produced an impression of spring."

Pittier (1908) has published a brief note on *Triplaris tomentosa* Wed., which he observed in Costa Rica, where it is generally called the "hormigo." It is "a small tree of the tierra caliente of the Pacific slope. It is diœcious, with the inflorescences more or less red and showy; the wood, which is of no use, is hollow and always infested with ants. In Nicoyo this tree is called "tabaco". According to Pittier, the Brunka Indians call it the "turi-svan-kra."

In 1913 I published some notes on *Triplaris americana* (under the name *Cumingiana* Fischer and Meyer) and *auriculata* Meisn. (*Macombii* Don. Smith). The former I observed in Panama, the latter in

Guatemala. My visits to Panama in 1923 and 1924 yielded much additional information in regard to *americana* and I have recently identified the ants taken in the internodes of *auriculata*. It will be advisable, therefore, to give a revised account of these plants and their tenants in the sequel.

Ule's statement that the ants inhabiting *T. Schomburgkiana* make a clearing around the base of the tree, is of interest in connection with a communication made to me by Mr. C. D. Mell. According to this student of tropical forestry, *T. americana* is very abundant in Venezuela, where the inhabitants call it "barabas" and where he has seen many specimens, each surrounded by a cleared area, "perhaps made by the ants." It is well known that two of our North American species of *Pogonomyrmex*, *barbatus* and *occidentalis*, make clearings around their nests by destroying the vegetation and that our common *Formica exsectoides* sometimes behaves similarly in our northern woodlands. There is, therefore, no reason to suppose that *Pseudomyrma dendroica* may not have developed a similar habit, though nothing of the kind has been observed in other species of the genus.

(B) *TRIPLARIS SURINAMENSIS* Cham. & Schl.

The only species of *Triplaris* which Professor I. W. Bailey and I had an opportunity to examine in British Guiana was *surinamensis*,¹ the "long John" of the colonists. It seems to be very common in Dutch Guiana, where, according to Pulle (1906), it is known to the settlers as the "mira-hoe-hoe", or "mierenhout" (ant-tree). We were unable to find it in the immediate vicinity of the tropical laboratory at Kartabo, but on August 4, 1920, a native guided us to a spot near Camaria, some seven miles up the Cuyuni River, where there was a small grove of young and vigorous "long Johns", about 20 to 30 feet high, growing on the low bank. The river, owing to the daily rains, was so high that the roots and bases of the trunks were under water and the dense undergrowth around them impeded our approach. We succeeded, nevertheless, in securing a number of boughs and placed them in the canoe—a very painful task, because all the foliage was swarming with a single very vicious species of *Pseudomyrma*. The native evinced great fear of the insects, insisting that they cause fever.¹ While the canoe was being towed the ants swarmed along the painter into our launch and tortured us all the way down the river to the laboratory.

¹The natives make the same statement in regard to the powerful Ponerine ant, *Paraponera clavata* Fabr. and in Southern Brazil and Bolivia concerning the even larger *Dinoponera grandis* Guér.

The twigs and branches of *surinamensis* exhibit the same peculiarities as those of other species as described by Schimper, Warming, Spruce and Morteo. Professor Bailey, who studied the morphology of the internodes and of the peculiar oval, slit-shaped, area at their upper ends in the material we collected, has contributed the following paragraphs and several photomicrographs of sections to illustrate his interpretation:

"The fistulose stems of *Triplaris surinamensis* are characterized, as are other representatives of the genus, by the presence of a slit-like opening in the distal portion of each internode. Are these openings, which serve as convenient entrances and exits, spontaneous structures or are they excavated by the ants? An examination of immature shoots reveals the fact that the slit-like orifices originate during the earlier stages of the enlargement of the internodes, and that they are due, in all probability, to asymmetrical growth and differentiation of the cortical, fibrovascular and medullary tissues. As in the myrmecophytic *Cecropias*, one longitudinal surface of each internode is flattened or slightly concave and is subtended by a leaf and its accompanying axillary bud. The growth and differentiation of the tissues is retarded in this side of the internode which is, in consequence, somewhat thinner and considerably weaker (Plate 10, Fig. c). This tendency toward asymmetrical development is accentuated at the distal end of the internode, below the insertion of the lateral organs of the next (higher) node, and the concomitant peripheral tension ruptures the delicate tissue in the thinner portion of the circumference of the cauline cylinder (Plate 10, Fig. a). The asymmetry is emphasized in vigorous, rapidly differentiating shoots and a sub-nodal aperture is formed in each internode. On the contrary, in stunted, slower-growing twigs, which are more nearly cylindrical, the distribution of the slit-like openings is rather sporadic. The apertures tend to become occluded by a growth of callus from the margins of the ruptured tissues (Plate 10, Fig. b) before the enlarging internodal cavities are taken possession of by the ants. Indeed, many of them are not subsequently reopened. Those that are, are characterized by having small circular apertures gnawed through the occluding callus.

"Coccids more or less numerous are found associated with the ants in the internodal cavities of *T. surinamensis*. In *Cecropia*, *Tachigalia*, *Cuviera*, *Plectronia* and various other Neotropical and Ethiopian ant-plants, the guest ants excavate pits in the denser tissues of the cauline cylinder which enable the Coccids to feed upon the softer tissues or upon a traumatically induced, thin-walled, centripetal callus. In *T.*

surinamensis the internodal chambers are jacketed by a thick, peripheral layer of medullary tissue (Plate 10, Fig. d), which remains green and physiologically active, even in stems which have formed a thick layer of secondary wood. The Coccids are able to insert their setæ into and to feed upon this medullary tissue. In other words, there is no evidence to indicate that the ants are concerned in facilitating the feeding of the Coccids. Although there is no reliable evidence at present for assuming that the imaginal ants solicit and feed upon the sugary exudates of the Coccids, an analysis of pellets fed to the larvæ indicates very clearly that the workers carve up the Coccids and feed them to the brood. (Plate 10, Fig. e.)

"The refuse of the ant-colonies, i.e. voided infrabuccal pellets, liquid fæces, fragments of malaxated insects, triturated plant-tissues, etc., is deposited at intervals along the walls of the elongated chambers. These latrines or middens give rise to luxuriant growths of delicate fungus hyphæ which are fed upon by an interesting group of structurally highly specialized Nematodes. The writer's investigations of Neotropical and Ethiopian myrmecophytes indicate that fungi are not cultivated and eaten by the ants, but that the aerial hyphæ are periodically cropped to prevent them from obstructing the chambers and interfering with the brood. That the middens in *T. surinamensis* may at times be utilized in the feeding of the brood is shown by analyses of the contents of the larval food-pouches, or trophothylaces. Not infrequently the pouches are filled with mats of hyphæ (Plate 10, Fig. f) or wads of detritus containing large numbers of Nematodes. It should not be inferred from this, however, that the imaginal ants actually cultivate the fungi, as do the Attini, or that they themselves feed upon hyphæ or Nematodes."

The fierce, aggressive ant inhabiting the *T. surinamensis* trees near Camaria, B. G. proves to be a darker and more slender race of *Pseudomyrma triplaris* Forel, which was taken by Ule, Goeldi and Huber in the same species of *Triplaris* in Brazil. I have described it as subsp. *baileyi*. The many flat Coccids of all sizes found in the internodal cavities were identified by Dr. Harold Morrison (1922) as a new genus and species, *Farinoecoccus multispinosus* Morr. and *Akermes secretus* Morr. They are sufficiently numerous to suggest that they may furnish the ants with considerable food in the form of honey-dew, especially during the rainy season. But certain observations, both negative and positive, cast doubt on this supposition. In the first place, no one has seen *Pseudomyrmas* attending Coccids or imbibing their sweet excreta. And when the twigs are broken open and the ants escape, they are

never seen to seize the Coccids and carry them away, like the species of *Azteca* and many other ants when similarly disturbed. These negative observations indicate rather that the Coccids may, while still very young, find their way into the internodes, either through the preformed clefts or through the openings made by the ants, and settle on the walls without being molested and perhaps without being noticed at first by the ants. But that the latter may utilize the Coccids, though not as dairy cattle, is shown by Professor Bailey's observations, which confirm our statements in regard to other species of *Pseudomyrminae* (Wheeler and Bailey, 1920; Bailey, 1922, 1923). As he has remarked, examination of the internodal cavities shows that there are in certain situations in the walls, latrines, or middens consisting of pellets ejected from the infrabuccal pockets of the ants and over-grown with luxuriant fungus mycelium inhabited by great numbers of Nematode worms. These latrines are strictly comparable with those of *Azteca longiceps* in the *Cordia alliodora* swellings described on p. 20. But unlike the *Azteca*, *Pseudomyrma baileyi* makes good use of the latrine materials, since the workers collect the hyphae, Nematodes, particles of infrabuccal pellets and pieces of Coccids and fashion them into food-pellets which they place in the trophothylaces of their larvæ. Since the Coccid fragments thus employed are rather numerous we suspect that the adult ants frequently devour these insects, possibly after they attain the proper dimensions, and *horribili dictu* serve up merely the remains mixed with other garbage from the latrines as the most appropriate or at least as the most available food for their progeny. The Coccids are used, therefore, as Bailey (1923) says, "for beef rather than solely as immature milk cows." Since there is a continuous growth and multiplication of Coccids, fungi and Nematodes in the internodal cavities, we are able to understand how the ants can develop and maintain large colonies in a plant which, so far as known, has neither extrafloral nectaries nor food-bodies such as we find in the myrmecophytic Acacias and Cecropias. Whether the ants also eat, instead of merely excavating, the sugary pith in the young internodes, is doubtful, but if Morteo's contention is correct, the plant itself does furnish an additional and perennial food-supply. The foregoing observations afford, I believe, at least a partial solution of the enigma which has puzzled several myrmecologists who have been unable to understand how the *Pseudomyrmas* not only survive but flourish in trees which normally grow in swampy places or on river banks where for several months each year they are isolated by high water.

At the time of our visit to Camaria, the ants in the *surinamensis*

trees were at the acme of the breeding season, the internodes of the twigs and branches being stuffed with the slender larvæ and pupæ in all stages, intermingled with recently emerged males, females and workers. The Coccids, too, were flourishing. Obviously the rainy season, which increases the sap in the tree, is most favorable to the development of the *Pseudomyrma* colonies, the Coccids, fungi, and Nematodes, and it may be confidently predicted that the Coccids will show a considerable decrease in numbers during the dry season. But then the ants can move more freely over the surrounding vegetation and secure many small, miscellaneous insects as food.

In the gardens of private residences in Georgetown, B. G., Professor Bailey and I noticed a few large *surinamensis* trees and on their trunks several different ants, but we had no opportunity for further investigations.

A few years ago Mr. H. E. Box sent me from Blairmont, Berbice, B. G. a considerable amount of material of *T. surinamensis* containing ants and Coccids. The ants proved to belong to two different subspecies of *Pseudomyrma triplaris*, which are described in the sequel (p. 184, 186) as subsp. *boxi* and *tigrina*. They resemble *baileyi* but are reddish. There were two species of Coccids, one being the same as the *Farinococcus multispinosus* Morrison, taken at Camaria, the other *Cryptostigma quinquepori* Newstead.

So far as known, therefore, *T. surinamensis* harbors in various parts of its range eight different organisms, namely, four subspecies of *Ps. triplaris*, two Coccids, at least one species of fungus and one Nematode. That this can be only a small fragment of the biocœnose which centers about *T. surinamensis* is evident from the following account of *T. americana* of which I was able to make a more thorough study.

(C) TRIPLARIS AMERICANA Linné

My first acquaintance with *T. americana* dates from December 1910 when Mr. E. D. Christophersen showed me near Empire in the Canal Zone a few of the trees growing in a swamp which is now at the bottom of Gatun Lake.¹ At that season they were not in bloom. During my

¹I had identified the plant (1913) as *T. Cumingiana* Fischer and Meyer, which is cited from the Isthmus by Seemann (1852-57) and Hemsley (1882-86). My identification was later confirmed by Professor B. L. Robinson of the Gray Herbarium. Meisner (1856) obviously redescribed the same species as *T. columbiana*. I now find that the Panamanian plant should be known as *T. americana* L., according to Standley (1925 p. 171), who describes it as the only Central American species of *Triplaris*. He states that "in Panama the tree is sometimes known as the 'palo santo' (a name given more commonly to *Erythrina glauca*), but oftener as 'guayabo hormiguero', while in Salvador it is called 'mulato' and 'palo mulato.'"

two more recent visits to Panama in 1923 and 1924 I was able to make a much closer acquaintance with the plant and its inhabitants. It is less abundant than *Cordia alliodora* and grows singly or in small clusters on low lands along the water-courses or on the lower slopes of ravines which carry a considerable amount of water during the rainy season. On my frequent trips across the Isthmus I noticed single specimens or small groups of the tree in many parts of the Zone within a mile of the railway from Frijoles to Balboa. A few scattered specimens were seen near the ruins of Old Panama in the Republic, but none was observed in the dryer savannah region somewhat further inland. Numerous accessible trees were found during 1923 in three localities: along the stream below the large, abandoned cacao plantation at Las Cascades, on the low banks of the Rio Grande between Ft. Clayton and the locks at Miraflores, and in the ravines descending from Ancon Hill behind the new administration building at Balboa. Unfortunately clearings were being made in the two localities first mentioned and a number of the trees were being felled, but at Balboa more intelligent and appreciative persons in charge of destroying the rank vegetation had carefully spared the *Triplaris* trees. In the ravine nearest the administration building there are a dozen beautiful specimens, which being within a short walk of Mr. Zetek's laboratory, could be visited at frequent intervals. During 1924 I found a few trees also in the swampy land at Marajal, near Colon, on the Atlantic side of the Isthmus.

T. americana is a small tree which rarely attains a height of more than 20 to 35 feet, with slender trunk and pyramidal crown, smooth gray bark, reddish twigs and bright green, drooping, smooth, entire, lanceolate leaves, about 5 to 8 inches long, with prominent veins and midrib. The cavities in the trunk, branches and twigs are precisely like those described for other species of the genus and the cleft elliptical areas at the distal ends of the internodes of the twigs show the same peculiarities as in the forms described by Schimper, Warming, Morteo and Bailey. The small round perforations made in the clefts by the ants are also very similar. The trees were found in bloom from February 25 till March 30. The spikes of small, sweet-scented, greenish white male flowers persist for only a short time, but in the female plant they are soon replaced by peculiar, rapidly enlarging, shuttlecock-shaped, fruits, which are at first green, then whitish and finally pure crimson. When the tree reaches this stage it is a very handsome and conspicuous object which might be advantageously introduced into the gardens of Southern Florida, California and the West Indies.

The shape of the fruits, with their long wings, indicates that they may be disseminated by the wind, like the samaras of our maples. I was unable to find any seedlings, and the smallest trees observed were not less than five or six feet high.

All the specimens of *americana* I have seen had ants nesting in their cavities, but the fauna is much more diverse than I had inferred from my meager observations of 1910, when the only ant encountered near Empire was a yellowish *Pseudomyrma* which I identified as *arboris-sanctæ* Emery. This ant has since been described by Forel as a distinct variety, *loewensohni* of the Colombian subspecies *symbiotica* of Emery's species. During 1923 and 1924 I took the following 16 species in or on the *Triplaris* in the various localities mentioned above:

- (1) *Pseudomyrma alliadora* Whlr. Balboa; Miraflores; in twigs.
- (2) *Pseudomyrma gracilis* Fabr. var. *bicolor* Guérin. Balboa; Miraflores; in twigs.
- (3) *Pseudomyrma sericea* Mayr var. *ita* Forel. Balboa; in twigs.
- (4) *Pseudomyrma triplarina* Weddell subsp. *symbiotica* Forel var. *loewensohni* Forel. Balboa; Las Cascades; Frijoles; Marajal; in twigs, trunk and branches.
- (5) *Crematogaster (Orthocrema) limata* F. Sm. subsp. *parabiotica* Forel. Miraflores; in twigs.
- (6) *Crematogaster (Orthocerema) brasiliensis* Mayr. var. *ludio* Forel. Miraflores; in twigs.
- (7) *Crematogaster (Orthocerema) brevispinosa* Mayr. var. *ampla* Forel. Balboa; Las Cascades; in twigs.
- (8) *Cryptocerus (Paraeryptocerus) minutus* Fabr. Miraflores; in twigs.
- (9) *Atta cephalotes* L. Balboa; cutting leaves and collecting flowers.
- (10) *Dolichoderus (Monacis) bispinosus* Oliv. Frijoles; visiting flowers.
- (11) *Azteca theresiæ* Forel var. *menceps* Forel. Miraflores, Balboa, Las Cascades; in twigs and branches.
- (12) *Azteca velox* Forel. Las Cascades; attending Coccids on branches.
- (13) *Tapinoma melanocephalum* Fabr. Balboa; in twigs.
- (14) *Brachymyrmex heeri* Forel var. *obscurior* Forel Balboa; in twigs.
- (15) *Brachymyrmex pictus* Mayr. subsp. *balboæ* Whlr. Balboa; in twigs.
- (16) *Camponotus (Myrmobrachys) lindigi* Forel. Balboa; attending Coccids on branches.

Four of these ants, *Atta cephalotes*, *Dolichoderus bispinosus*, *Azteca velox* and *Camponotus lindigi* merely visit the trees. The *Camponotus* and *Azteca* attend Coccids on the twigs and stems of the flower-panicles, and the *Atta* may perhaps cut and collect the fruits. The remaining 12 species were all found nesting in the fistulose twigs and

branches, but only two, *Pseudomyrma loewensohni* and *Azteca menceps* are obligates, most if not all the others occurring also in other myrmecophytes or even in the dead twigs of various trees and shrubs. The singular fact was disclosed that *Ps. loewensohni* was altogether absent from the trees in the Miraflores locality and present only in a certain number of those near Las Cascades, Balboa and Mirajal. Many trees were inhabited in great part or entirely by the Crematogasters and *A. menceps*, the latter being the most common. The other forms in the list were usually confined to single twigs or branches of trees tenanted also by this *Azteca* or one of the Crematogasters. When *Ps. loewensohni* was present it usually had complete possession of the tree. These statements may be illustrated by the distribution of the ants recorded in my note book under the date of March 26, 1923 as occupying the 12 trees in the ravine at Balboa behind the administration building.

- (1) Tree about 15 ft. high; fruiting. Twigs inhabited exclusively by *A. menceps*.
- (2) About 30 ft. high, with two trunks; flowering. Twigs inhabited by *C. ampla* and *B. obscurior*.
- (3) About 30 ft. high, beginning to fruit. Twigs occupied by *C. ampla*.
- (4) About 9 ft. high, in flower. Twigs and branches inhabited by *Ps. loewensohni* exclusively.
- (5) About 15 ft. high; beginning to fruit; leaves much eaten. Twigs tenanted by *C. ampla*.
- (6) About 20 ft. high; in flower. Tenanted throughout by *Ps. loewensohni*.
- (7) About 20 ft. high; in flower, but with several dead branches. Twigs occupied by *A. menceps*.
- (8) About 35 ft. high; beautiful specimen, flower. Tenanted by *A. menceps*.
- (9) About 25 ft. high, in flower. Twigs inhabited by *A. menceps* and *B. balboæ*.
- (10) About 35 ft. high, in flower. Tenants: *A. menceps* (abundant); *Ps. alliodoræ* (sporadic); *C. lindigi* running on trunk and foliage.
- (11) About 35 ft. high; mature fruit. Tenants: *B. balboæ* (in several branches); *Ps. bicolor* (sporadic); *C. lindigi* running on trunk.
- (12) About 30 ft. high; just passed flowering. Tenants: *C. ampla* (abundant); *T. melanocephalum* (in a few internodes); *Ps. ita* (sporadic); *C. lindigi* running on trunk and branches.

My notes on these trees are very brief and may not represent an accurate census, especially of the inhabitants of the larger specimens, because, owing to their height, I was unable to make an exhaustive inventory of their twigs and branches. Moreover, the trees were being preserved for ornamental purposes and could not be mutilated. I believe, nevertheless, that few species of ants, and only those forming very small sporadic colonies in some of the internodes, were overlooked. Quite a number of twigs from most of the trees were cut off, carried to the laboratory in bags and chloroformed so that the ant colonies could be carefully examined. It will be seen that only two of the twelve trees were inhabited by *Ps. loewensohni*, whereas five contained *A. menceps* and four *C. ampla*. One of the largest harbored none of these species. In the series of more than 20 trees between Ft. Clayton and Miraflores, those nearest the former locality were inhabited exclusively by *A. menceps*, those nearest the latter by *Crematogaster ludio*, with a few sporadic colonies of *Cryptocerus minutus*. Near Las Cascades only three among a dozen trees examined, were tenanted by *Ps. loewensohni*, and one of these, a vigorous specimen about 15 ft. high, which was felled and carefully examined, also had *A. menceps* in some of its twigs. Many of its smaller branches were quite free from ants and in part inhabited by the larvæ of a Thyridiid moth (*vide infra*, p. 60), which was common also in many of the trees in the other localities.

Since, apart from the sporadic *Ps. ita* and *bicolor*, the only ant listed that stings painfully is *Ps. loewensohni*, and since it occurred only in a small number of the trees and was even entirely absent in one locality, *T. americana*, unlike its various South American congeners (*Schomburgkiana*, *caracasana*, *surinamensis*, etc.) could usually be handled and examined with impunity. And while it is not improbable that in some Panamanian and Colombian localities the plant may be inhabited exclusively by *Ps. loewensohni* or the typical subsp. *symbiotica* or by vicious species other than those enumerated in my list, a study of the trees in the Canal Zone shows that specimens harboring only such small inoffensive ants as *A. menceps*, *C. ampla* and *ludio*, are quite as vigorous and flourishing as those inhabited by the red, stinging *Pseudomyrma*. At Balboa, in fact, this ant did not occur in the largest and finest trees. Besides the case above mentioned of *loewensohni* living in the same tree as *A. menceps*, the following observation shows that the *Pseudomyrma* is not a very aggressive protector of its tree from alien ants. On March 28 at Frijoles, I came upon a *Triplaris* about 14 ft. high which had just

been felled by a native. It was in full fruit and the large crimson panicles had been invaded for some reason by a host of the belligerent *Dolichoderus bispinosus*. Closer examination showed that the cavities of the tree were occupied by a thriving colony of *loewensohni*, the workers of which were running about quietly and without the slightest signs of hostility among the Dolichoderi. The following observation made at Balboa, March 25, indicates, perhaps, that the other ants of *T. americana* are quite as indifferent to invasions of their host tree by alien species. A few feet from the trunk of tree No. 12 at Balboa there was a large nest of *Atta cephalotes*, but all the ants had withdrawn into the depths of their galleries. That they had been working during the preceding night was shown by the great masses of *Triplaris* flowers which they had carefully collected and left in the runways leading to their nest-craters. Of course, these wilted flowers may have been merely picked up on the ground, but that they may have been culled directly from the trees is by no means improbable. I may add, in this connection, that a few of the trees near Fort Clayton, though inhabited by flourishing colonies of *A. menceps*, nevertheless had their leaves severely damaged by *Atta cephalotes*.

Before listing the miscellaneous organisms associated with *T. americana* and with its ants, it will be advisable to insert a few remarks on the two obligates, *Azteca menceps* and *Pseudomyrma loewensohni*. The types of the former were taken by Christophersen in the Canal Zone but the precise locality was not recorded. They were probably from the very trees to which he guided me in 1910. The habits of this ant are similar to those of *A. longiceps*, described above (p. 28), and the fullest development of its colonies occurs at the same season, coinciding with the anthesis and fruiting of the host-plant. Like the other *Triplaris* ants, *menceps* makes small round entrances in the elliptical cleft areas of the internodes and also destroys the partitions between them so that their cavities become continuous. The walls of the cavities also have gnawed pits containing Coccids, which are rather numerous, pinkish in color and covered with white wax. There are, moreover, definite latrines at intervals on the walls, like those of *A. longiceps* in the cauline swellings of *Cordia alliodora* and consisting of numerous pellets from the infrabuccal pockets of the ants, covered with proliferating fungus hyphæ and invaded by hosts of Nematodes and bacteria. Mites are also present occasionally. The pellets themselves consist of spores and small pieces of insects and plant tissue.

Ps. loewensohni, though a very vicious and aggressive ant, seems not to sting quite as severely as the *Pseudomyrmas* inhabiting *T.*

surinamensis, *Tachigalia paniculata* and the bull-horn Acacias. When the twigs or branches which it inhabits are split open their walls are found to be smooth and clean and of a dark brown color. To these walls the larvæ are hung by means of their hooked, curved dorsal hairs and are fed with infrabuccal pellets like the larvæ of other Pseudomyrminae. The latrines, which occur at intervals of several inches, are of much the same composition as described for *A. menceps* and *longiceps*, but moister and more glutinous. I found few Coccids in the internodes and at rather infrequent intervals. Instead of making pits, the ants gnaw narrow grooves one to three centimeters long and in these the Coccids lie in a linear series. They resemble the species cultivated by *A. menceps* but are smaller, suggesting that *loewensohni*, like *Ps. baileyi*, may occasionally devour them instead of using them as a perennial source of honey-dew. Most of my observations on these ants, however, were made at the height of the dry season and while the trees were in flower or fruit, so that the small number of Coccids may also be due in part to less favorable trophic conditions. Since the larvæ, pupæ, males and winged females of *loewensohni* were not very abundant, it is probable that, like *Ps. baileyi*, it breeds more actively during some portion of the rainy season. This was indicated during the summer of 1924 by observations in the swamp at Marajal. Here the few colonies of *loewensohni* which I examined contained more brood and more numerous sexual forms and Coccids.

The following miscellaneous organisms were found associated with *T. americana* or with its ant-inhabitants:

Lepidoptera. Among the Heterocera there are at least five species that feed on the trees. One of these is identified by Dr. W. T. M. Forbes as a Thyridiid, the greenish caterpillar of which, nearly an inch long, is very common on the twigs, devouring the pith, tunnelling through the nodal partitions and depositing masses of coarse frass at intervals. This caterpillar is so common that it must be an important agent in facilitating the occupation of the twigs by the ants. Although pupæ were obtained during the latter part of March I did not succeed in rearing the imagines.

On the trunks of some of the trees at Balboa there were communal masses of cocoons, larger than one's fist. From these Mr. Zetek reared many specimens of both sexes of two handsome reddish brown Saturniid moths, which Mr. W. Schauss has identified as *Hylesia hamata* Schauss and *H. darlingi* Dyar.

A large empty cocoon of another Saturniid was found attached

to the trunk of one of the trees and the case of a Psychid allied to *Thyridopteryx* to a twig.

The following Rhopalocera, identified by Dr. Forbes, were taken at the *Triplaris* flowers at Balboa.

- (1) *Junonia lavinia zonalis* Felder.
- (2) *Anartia jatrophae* L.
- (3) *Anosia plexippus* L.
- (4) *Anosia berenice* Cram.
- (5) *Antigonus (Systasea) crosus* Huebn.
- (6) Hesperid (unidentified).

Diptera. The only members of this order taken were a small Trypetid which was swept from the flowers and a small Asilid occasionally resting on the trunk.

Hymenoptera. The caterpillar of the Thyridiid above mentioned is parasitized by a small Hymenopterous larva which makes its cocoon in the twig cavity inhabited by the host. I failed to rear the imago.

A single large red Ichneumonid was seen ovipositing in one of the cocoon-masses of *Hylesia hamata*, but escaped.

The following were taken at the flowers: two Vespidae, *Polistes canadensis* L. and *Eumenes nana* Kirsch (J. Bequaert det.) and the following eight bees, identified by Prof. T. D. A. Cockerell (1928):—

- (1) *Megachile poculifera* Ckll.
- (2) *Apis mellifica* L.
- (3) *Apis mellifica* var. *ligustica* Spin.
- (4) *Melipona fulvipes* subsp. *triplaridis* Ckll.
- (5) *Melipona orbigny* subsp. *phenax* Ckll.
- (6) *Trigona cupira* F. Smith.
- (7) *Trigona pectoralis* D. T. subsp. *panamensis* Ckll.
- (8) *Nannotrigona testaceicornis* Lep.

All but one of these are social bees.

Coleoptera. Three species of beetles taken at *T. americana* flowers at Balboa in 1923 were identified by Mr. H. S. Barber as *Coscinoptera cingulata* Lec. (Chrysomelid, several specimens), *Hyporhagus lœvipunctatus* Thom. (Monommid, several specimens) and *Acanthoscelides* sp. nov. (Bruchid, one specimen). The last has been reared by Mr. Bridwell from pigeon-peas and the description is in MS. Three other unidentified Coleoptera were observed at Balboa, namely some small larvæ, possibly Curculionids devouring the pith in young twigs, quite a number of Coccinellid larvæ on the foliage and evidently preying on the Coccids and a species of termitophilous Staphylinid living in considerable numbers in one of the nests of the termite cited below.

In 1924, five more Coleoptera were taken on the bark and leaves but were not identified (One small Coccinellid; one Curculionid, one Anthribid and two small Tenebrionids).

Isoptera. Some of the trees at Balboa had large elliptical, ellipsoidal, black termitaria enveloping their trunks. The termites were identified by Mr. Zetek as *Nasutitermes corniger*. A dead shoot at the base of one of the trees was inhabited by a species of *Leucotermes*.

Orthoptera. Several small cockroaches determined by Mr. Hebard as *Latiblatella angustifrons* Hebard, were found hiding between the walls of the *Nasutitermes* termitaria and the bark to which they were attached. Small oothecæ which must have belonged to some other Blattid were occasionally found in the cavities of the twigs.

Thysanoptera. A large black species with greatly incrassated anterior femora was occasionally found in the twigs together with nymphal individuals.

Heteroptera. Dr. R. F. Hussey has identified the following bugs taken from the foliage:

Dysdercus ruficollis L. (Pyrrhocorid)

Zelus sp. nov. near *atripes* Champ (Reduviid)

Hypselonotus lineatus Stal. var. *neglectus* Horvath (Coreid)

Homoptera. An unidentified Aleurodid was occasionally seen on the leaves. The following Coccids have been identified by Dr. Morrison.

(1) *Akermes* sp. On twigs.

(2) *Pseudococcus brevipes* Ckll. In internodes with *Azteca menceps*. Las Cascades; with *Crematogaster ludio* at Miraflores.

(3) *Pseudococcus probrevipes* Morr. In internodes with *Azteca menceps* near Miraflores; with *Pseudomyrma loewensohni* at Balboa.

(4) *Saissetia auriculata* Morr. On twigs inhabited by *Ps. loewensohni* at Las Cascades.

(5) *Ceroplastes cirripediformis* Comst. (?) On twigs at Balboa.

Aracina. The only spider seen was *Eriophora edax* Bl. (N. Banks det.), which was running on the bark.

Acarina. The mites found in the latrines of *Azteca menceps* have not been identified.

Nematodes. The nemas occurring in the latrines of the same ant are being studied by Dr. Cobb.

Phaenogams. A large mistletoe (*Struthanthus orbicularis* H. B. K.) was found growing on two of the trees at Balboa. It is common on other trees in the Canal Zone.

Cryptogams. Here belong the moulds and bacteria that flourish in the latrines of *A. menceps* and *Ps. loewensohni*.

Adding the 52 Arthropods cited in the foregoing list to the 16 different Formicidæ we have 68 forms associated with *T. americana* and distributed as follows:

Hymenoptera	28
Lepidoptera	11
Diptera	2
Coleoptera	11
Isoptera	2
Orthoptera	2
Thysanoptera	1
Heteroptera	3
Homoptera	6
Araneina	1
Acari	1
Total	68

The trees examined were much fewer in number than those of *Cordia alliodora* and I was unable to study any very young or seedling specimens. Nevertheless the number of insects infesting *americana* seems to be sufficient to corroborate the general conclusions derived from a study of *Cordia alliodora*.

(D) *TRIPLARIS AURICULATA* Meisner

According to Standley (1922), this is probably the correct name for the tree which I called *T. Macombii* Donn. Smith in my paper of 1913. He records it as occurring in Chiapas "and perhaps elsewhere in Mexico". Donnel Smith's material was collected at Jiquilisco, Salvador. I did not at first recognize it as a *Triplaris* when I encountered it in January 1911 along the roadsides of Escuintla and Patulul in Western Guatemala. It is a larger tree than *T. americana*, attaining a height of at least 40 or 50 feet, with stouter trunk and more spreading branches and large, coarse, dark green, broadly ovate, short-petioled leaves. When first seen it was putting forth branches of long yellowish flower-spikes, covered with a deciduous sheath. The fruit, which I did not see, is described by L. Donnel Smith (1894) as pale yellow. He has described (1895) a variety *rufescens* from Mayatinango, Guatemala, with brick-red flowers and more pilose leaves. My specimens seem to agree best with those of the typical *Macombii* in the Gray Herbarium. The twigs and branches are coarser than those of *T. americana*. Re-examination shows that the twigs have the same structure and are perforated in the same manner by the ants at the elliptical cleft areas,

which look like hypertrophied lenticels but are very probably formed as in other species of *Triplaris* by local dehiscence and subsequent healing of the expanding internodes. I have not since had an opportunity to study *auriculata* which seems to have a rather restricted distribution.

The following ants were collected in the internodes of the tree in 1911:

Pseudomyrma sericea Mayr. var. *ita* Forel.

Pseudomyrma sericea var. *fortis* Forel.

Monomorium carbonarium F. Smith subsp. *cbeninum* Forel.

Azteca prorsa Wheeler.

Tapinoma ramulorum Emery subsp. *inrectum* Forel.

Of these the *Pseudomyrmas* and *Azteca* are the most abundant and characteristic, the latter being, perhaps, peculiar to the plant (obligate). The *Monomorium* and *Tapinoma* occur also in the twigs of other trees. Cutting the branches of *T. auriculata* is decidedly painful owing to the *Pseudomyrmas* which, though they form smaller colonies, sting quite as severely as *Ps. loeuensohni*. *A. prorsa* is closely related to *A. pittieri*, *longiceps* and *menceps* and seems to have very similar habits. It is timid and inoffensive.

In order to complete my enumeration of the ants inhabiting *Triplaris*, I may refer briefly to the records of those taken in undetermined species. The following have been cited in the literature merely as occurring in *Triplaris* or without further comment:

- (1) *Pseudomyrma dendroica* Forel. In hollow branches of *Triplaris* sp. Amazonas (E. Ule)
- (2) *Pseudomyrma latinoda* Mayr. Probably in *Triplaris* or *Tachigalia*. Brazil (J. Trail)
- (3) *Pseudomyrma arboris-sanctæ* Emery. In *Triplaris* sp. Bolivia (Balzan, W. M. Mann); Peru (Staudinger); Amazonas and Matto Grosso (F. Silvestri)
- (4) *Pseudomyrma arboris-sanctæ* var. *cordobensis* Forel. Probably in *Triplaris*. Argentina (C. Bruch)
- (5) *Pseudomyrma arboris-sanctæ* var. *rurrena-baquensis* Wheeler. In *Triplaris* sp. Bolivia (W. M. Mann)
- (6) *Pseudomyrma arboris-sanctæ* var. *symbiotica* Forel. In *Triplaris* (probably *americana*) Colombia (A. Forel; G. Salt)
- (7) *Pseudomyrma sericea* Mayr. var. *rubiginosa* Stitz. In *Triplaris* sp. Brazil (E. Ule)
- (8) *Azteca brevicornis* Mayr. var. *boliviana* Wheeler. In *Triplaris* sp. Bolivia (W. M. Mann)

Ps. dendroica and *latinoda* are closely related to *arboris-sanctæ* and *triplaridis* and undoubtedly have very similar habits. The Azteca was taken in the stems of a peculiar *Triplaris* which judging from Dr. Orlando White's herbarium specimens accompanying it, is quite unlike those with which I am acquainted. The only ant recorded from *T. Schomburgkiana* is *Ps. dendroica* var. *emarginata* Forel, which was taken by Ule in Amazonas (see p. 164). Morteo, as we have seen (p. 47), records the occurrence of the common Oriental *Dolichoderus* (*Hypoclinea*) *bituberculatus* as living in *T. americana* when growing under cultivation in the East Indies. Omitting this ant, there are some 30 different Formicidæ and more than 50 other organisms known to inhabit the hollow branches and foliage of the various species of *Triplaris* in different parts of the American tropics.

Chapter 3. OBSERVATIONS ON TACHIGALIA

The beautiful Caesalpinaceous trees of the genus *Tachigalia* are confined to Amazonas, Eastern Peru, the Guianas and Venezuela, and therefore have an even more restricted distribution than the species of *Triplaris*. About 20 species have been described, but the "Index Kewensis" recognizes only 13 as valid, and perhaps this number may be reduced when more material is available and the range of variation in the different forms has been thoroughly studied. The type species, at least, varies considerably even during its ontogeny, the juvenile form when growing in the shade having a very different habitus from the large tree exposed to the sun-light.

The genus was established by Aublet as long ago as 1775 for a species, *paniculata*, which he found growing along the rivers of French Guiana. He named the plant *Tachigalia* from the Carib "tachigali", "tachi" being the name employed by the Indians of the Guianas and Brazil for the stinging ants of the genus *Pseudomyrma*, which regularly inhabit the enlarged petioles of *Tachigalia* as well as the internodal cavities of *Triplaris*. Part of Aublet's description of *T. paniculata* may be quoted: "The trunk of this tree rises to a height of fifty or sixty feet or more, with a diameter of three feet. Its bark is gray, rugose, its wood is hard and whitish. It produces at its summit a large number of stout branches which spread in all directions and are laden with twigs furnished with alternate winged leaves, with two rows of opposite leaflets, the greatest number of which is six on each side. They are firm, entire, terminating in a point, smooth and green above, grayish green beneath. Their petiole is very short,

articulated to a triangular swelling five inches long and terminating in a point, accompanied at its origin by two stipules which are soon deciduous. The largest leaflets are six inches long by two and a quarter inches broad. The flowers arise at the end of the twigs and are borne in large, long panicles, the stems of which are simple and covered with flowers throughout their length." The corolla is yellow and sweet-scented and the seeds are surrounded by a flat, elliptical wing or expansion. The tree flowers and fruits in April and November. Aublet described also a *T. trigona*, which proves, however, to be a synonym of *paniculata*.

Tulasne (1844) redescribed *paniculata* and added six other forms, only two of which are now recognized as distinct, two, *ericalyx* and *sericca*, being synonyms of *paniculata*. *T. angustifolia* described by Miquel (1851) from Surinam also proves to be a synonym of Aublet's species.

None of these botanists deigned to leave us any notes on the relations of the Tachigalias to their ants. This was left to Spruce, who writes in his article submitted to the Linnean Society in 1869 but not published till 1908: "The Tachigaliæ are low-growing riparial trees, of black-water rivers, and have pinnate, often silky foliage; and small, yellow, sweet-smelling, nearly regular flowers disposed in panicles. All have trigonous petioles, which are mostly dilated at the base into a fusiform sac, tenanted by ants. *T. caripes* (*recte cavipes*, later regarded by Bentham (1870-76) as a variety of *paniculata*) grows abundantly on the banks and on inundated islands, of the Uuapes. It is a spreading tree of 30 feet, and has the ramuli, petioles and leaves clad with a fine, close, silky pubescence. The sacs of the petiole are inhabited by small black ants, whose entrance is by a little hole on the underside of the sac. *T. ptychophysca* sp. n. grows in moist sandy caatingas by the same river, and has a similar sac on the petiole."

Ule (1907), in Amazonas and Eastern Peru, observed three species of Tachigalia, which were identified by Harms (1906) as *formicarum* Harms, *paniculata* Aubl. and *spicata* Aubl. The first, from Tarapoto, Peru, proved to be a new species and is a tree 30 meters high, with yellowish flowers. Its large petiolar sacs were inhabited by *Pseudomyrma latinoda* Mayr subsp. *tachigaliæ* Forel. Ule states that "the ants live mainly in the petiolar sacs; in blooming and fruiting specimens they also settle in the axes of the large, hollow, swollen flower panicles. Only on one occasion, near Iquitos, did I observe that they had also perforated and taken possession of the twigs."

J. Huber (1909) described a *T. macrostachya*, which was collected by Ducke on the Rio Trombetas, Amazonas, and had its hollow petioles inhabited by *Ps. latinoda* var. *endophyta* Forel.

In 1921 I gave a detailed account of *T. paniculata* and its inhabitants as observed by Professor I. W. Bailey and myself at Kartabo, British Guiana, and Professor Bailey (1923) has since published a description of the anatomical peculiarities of the plant. The reader may also be referred to my paper for an account of the ants, coccids and other organisms and especially of the very interesting social Silvanid beetles, *Coccidotrophus socialis* and *Eunausibius wheeleri*, which live in the petiolar cavities of the young trees before they are occupied by the ants.

The following list includes all the known ants which I have found in the petiolar swellings of *T. paniculata*, together with those recorded by others as occurring in other species of the genus:

- (1) *Neoponera crenata* Roger. (*T. paniculata*)
- (2) *Neoponera unidentata* Mayr. (*T. paniculata*)
- (3) *Pseudomyrma damnosa* Wheeler (*T. paniculata*)
- (4) *Pseudomyrma latinoda* Mayr. var. *coronata* Wheeler (*Tachigalia* sp. ?)
- (5) *Pseudomyrma latinoda* var. *endophyta* Forel (*T. macrostachya*)
- (6) *Pseudomyrma latinoda* var. *nigrescens* Wheeler (*Tachigalia* sp.)
- (7) *Pseudomyrma latinoda* subsp. *bradleyi* Wheeler (*Tachigalia* sp.)
- (8) *Pseudomyrma latinoda* subsp. *tachigaliæ* Forel (*T. formicarum*)
- (9) *Pseudomyrma maligna* Wheeler (*T. paniculata*)
- (10) *Pseudomyrma maligna* var. *cholericæ* Wheeler (*T. paniculata*)
- (11) *Pseudomyrma maligna* var. *crucians* Wheeler (*T. paniculata*)
- (12) *Pseudomyrma picta* Stitz (*Tachigalia* sp.)
- (13) *Pseudomyrma picta* Stitz subsp. *casta* Wheeler (*Tachigalia* sp.)
- (14) *Pseudomyrma sericea* Mayr. var. *rubiginosa* Forel (*Tachigalia* sp.)
- (15) *Pheidole cramptoni* Wheeler subsp. *petiolicola* Wheeler (*T. paniculata*)
- (16) *Pheidole* (*Hendecephidole*) *tachigaliæ* Wheeler (*T. paniculata*)
- (17) *Crematogaster* (*Orthocrema*) *delitescens* Wheeler (*T. paniculata*)
- (18) *Crematogaster* (*Orthocrema*) *limata* F. Sm. subsp. *palans* Forel (*T. paniculata*)
- (19) *Solenopsis helena* Em. subsp. *hermione* Wheeler (*T. paniculata*)
- (20) *Solenopsis helena* subsp. *ultrix* Wheeler (*T. paniculata*)
- (21) *Leptothorax* (*Goniothorax*) *echinatinodis* Forel subsp. *aculeatinodis* Emery var. *pleuriticus* Wheeler (*T. paniculata*)

- (22) *Leptothorax (Goniothorax) umbratilis* Wheeler (*T. paniculata*)
- (23) *Azteca foveiceps* Wheeler (*T. paniculata*)
- (24) *Azteca tachigaliæ* Forel (*Tachigalia* sp.)
- (25) *Azteca traili* Emery (*T. paniculata*)
- (26) *Brachymyrmex heeri* Forel (*T. paniculata*)
- (27) *Brachymyrmex heeri* var. *basalis* Wheeler (*T. paniculata*)
- (28) *Camponotus (Myrmobrachys) pittieri* Forel var. *pænalis* Wheeler (*T. paniculata*)
- (29) *Camponotus (Myrmobrachys) zoc* Forel (*T. paniculata*)

Of these 29 forms probably the only ones that can be regarded as obligates are the various Pseudomyrmas and Aztecas, the remainder being merely inquilines which frequently nest in the cavities of other plants. Perhaps *A. traili* should be assigned to this latter group. In my paper on *T. paniculata* I gave a list of 28 miscellaneous organisms associated with the plant, its ants or its Coccids (*Pseudococcus brevipes*). Dr. Morrison (1922) has since described a second coccid, *Ripersia petiolicola*, which occasionally lives on the petioles but was not mentioned in my paper. As known to date, therefore, the *Tachigalia* biocœnoses embrace more than fifty different organisms. The conclusions which we reached as a result of our investigations are well summarized in the two concluding paragraphs of Professor Bailey's paper:

"The hollow foliar axes of the ant-plant *Tachigalia paniculata* are colonized by at least seven "obligatory" guest insects. The domatia of juvenile plants are taken possession of by two extraordinary social beetles, but during the subsequent development of the plants these insects are dispossessed by ants. Both the beetles and the ants avail themselves of the structural peculiarities of their host plant in a singularly efficient manner. Not only do they make use of the hollow foliar axes as convenient nesting chambers, but they feed, either directly or indirectly, upon the softer tissues in the interior of the domatia. The beetles eat the parenchyma of the wide, primary medullary rays, utilize it in the construction of their puparia, and solicit and obtain liquid carbohydrates from herds of Coccids which graze upon it. The ants also feed upon this tissue vicariously through intervention of Coccids, and utilize it in the construction of carton partitions.

"There is no evidence to indicate that the structural peculiarities of *T. paniculata* are initiated by ants or by gall forming insects, or that they originated as adaptations for attracting a defending army of ants. The relations between the host plant and the beetles and the

ants are not those of a mutually beneficial symbiosis (Belt, Schimper), but an interesting type of parasitism, in which there is a remarkable parallelism in the behavior of representatives of such widely separated groups of insects as the Hymenoptera and the Coleoptera. The nesting and feeding habits of the insects, and their relations to the coccids, are largely determined by the structure and arrangement of the various vegetative tissues during different stages in the development of the petiole and rhachis. In studying the habits of phytophagous and plant inhabiting insects, it is evident that the anatomy of the host plants deserves more careful consideration than it has received heretofore."

Chapter 4. OBSERVATIONS ON CECROPIA

The trees of the Moraceous genus *Cecropia* have been much more carefully investigated than any other myrmecophytes. This is in part due to the fact that they are so abundant and are so readily recognized, because they contrast markedly with the remaining vegetation and especially at low altitudes constitute a very characteristic feature of the landscape. They are usually rather small or medium-sized trees, with smooth, pale-barked, slender trunk and branches, the latter few in number and arranged like the branches of a candelabrum and bearing large, coarse, long-petioled palmate leaves, which are usually silvery-white beneath. When moved by the wind the lower surfaces give a touch of vitality to what may be a rather monotonous background of vegetation. Writing of this feature of the *Cecropias* and of the coppery under surfaces of the leaves of the laurels and *Chrysobalan* along the Amazon, Spruce (1869, in 1908) says: "When my boat has been floating lazily in the water, under a burning and dazzling sun, with not a breath of air stirring, I have sometimes—as my eye wandered along the endless forest margin—given vent to some such exclamation as this "How tame and monotonous!" when the coming on of a squall, by simply revealing the glowing tints of the underside of the leaves, has in a moment waked up the scene into life and beauty." The *Cecropias*, unlike *Cordia* and *Tachigalia*, are dioecious and their inflorescence is inconspicuous, consisting in both sexes of bunches of compact green catkins, or aments. The trees are known to the natives of Brazil under the names "imbauba", "imbauva" "ambauva", "embaiba" or "ambay", to the Peruvian as "ceticos", to the natives of Central America as "guarumo", and to those of the Antilles as "llagrums" or "yagram". The British colonists in South

America call them "shakewood", "trumpet" or "drum" trees. The West Indians in the employ of the Panama Canal Zone refer to them as "trompies".

In Brazil, according to Bequaert (Wheeler and Bequaert, 1929), "the Cecropias generally grow in dense grooves. Some of the species of the Amazonian Basin are commonly found in the second growth, in plantations, along roadsides and in waste-places. They may be observed right in the towns and sometimes grow out of the walls of old buildings. This would seem to indicate that the seeds are scattered in the excrement of birds and bats that feed on the fruit. Other species prefer the alluvial woods, or what is known in Brazil as "varzea", i.e. low-lying land that becomes water-logged when the rivers reach their highest level. Finally a few species thrive in the "igapo", or true inundated forest, and are the very first plants to colonize the shifting mud-banks in the river itself, as well as the new alluvial land. During much of the year these Cecropias stand in swiftly flowing water many feet deep, so that the ant-colonies that inhabit them must have their food-supply restricted to the food-bodies produced by the plant and the honey-dew furnished by the Coccids in their internodal cavities".

My own observations in British Guiana, Mexico, Central America, Cuba and Porto Rico agree with Bequaert's account, except that in these countries there seem to be no forms adapted to amphibious conditions. Though the trees are sometimes found in moist places they are more frequent on the hills or mountain sides, either in small grooves or single and intermingled with the other plants of the jungle or hylæa. When man interferes with the natural conditions, the Cecropias behave as veritable tree-weeds and rapidly take possession of the newly turned soil of railroad and ditch embankments, roadsides, clearings; etc. I was able to witness the occupation of such sites on a grand scale in Panama, during the building of the canal and relocation of the transisthmian railroad. Bequaert's suggestion that the seeds may be spread by birds and bats seems very probable.

Of all the neotropical ant-plants the Cecropias are the most widely distributed. They are common throughout tropical Mexico, Central America, the Antilles and South America except Patagonia and parts of Argentina and in some of these regions often ascend to altitudes of four or five thousand feet. Although botanists have been sufficiently interested in the genus *Cecropia* to describe about 100 species, they have failed to give us any adequate monograph or key to the various forms, so that their identification in the field is a matter of no little

difficulty. Moreover, as Bailey (1922, p. 371) has shown, the leaves of the same species undergo profound changes in form during ontogeny, and the specific characters are probably still further obscured by hybridization. Apparently only certain species are myrmecophilous, while others are never regularly inhabited by ants. There are also two genera, *Pourouma*, with about 20 species, and *Coussapoa*, with about 15 species, which are closely related to *Cecropia* and at least occasionally harbor ants, but our knowledge of these plants is very fragmentary.

The association of ants with *Cecropias* has long been known. It was briefly described by Marcgravius (1648), Piso (1658) and Ray (1688) in the seventeenth century, but excited no further inquiry till 1874, when Belt recorded his observations in Nicaragua. Since that time a number of papers, wholly or in part devoted to the *Cecropias* and their ants have been published by Fritz Müller (1876, 1880, 1883), Schimper (1888), Schumann (1889), Warming (1894), Ule (1897, 1905, 1906), Rettig (1904), H. von Ihering (1907), Fiebrig (1909), Wheeler (1908, 1913), Bailey (1922) and Wheeler and Bequaert (1929). The papers of Müller, Schimper, von Ihering and Fiebrig on *C. adenopus* L. and of Bailey on *C. angulata* are specially worthy of notice.

A general description of the structure of the *Cecropias* is given in my ant-book (1910, p. 305-310) and is reproduced by Bequaert in his article on the myrmecophytes in my "Ants of the Belgian Congo (1922). Ule (1907) has also published a brief account which I translate: "All the *Cecropias* that harbor ants have essentially the same structure which Schimper has described in detail. The internodes of the branches and twigs are hollow and separated from one another by transverse partitions. Each internode possesses a leaf, and above it there is a groove which it produces by pressure of the petiole on the axial bud while it is still enclosed. In this groove there is always a pit. A similar shallow groove also occurs in other ant-plants, but the pit is peculiar to *Cecropia*. Now at this point the *Cecropia* is always perforated by the fecundated (Azteca) queen, and this is the more easily accomplished because the wall of the pit is thin and lacks fibrovascular bundles. After the queen has slipped into the cavity or chamber through the opening, the latter, as long as no workers are present, closes through the formation of callous tissue around its edges." Later the worker progeny of the queen again open the orifice and use it permanently as the entrance or exit to the internode. "The chambers which arise in the internodes in the place of the early disappearing pith are circular in cross-section, on an average

4 to 7 cm. long and separated from one another by very thin fragile partitions, which are always perforated by the ants. In the interior of the chambers the ants build up a kind of labyrinth for their larvæ out of a brown, wax-like substance probably derived from the *Cecropia* tree. As a rule each chamber is at first inhabited by only a single female, whose cell is often all it contains. In addition there are always white Coccids, on the saccharine excretions of which the ants in part subsist. But the insects are also afforded another source of nourishment by the plant itself, for at the base of each leaf-petiole there is a hairy cushion (trichilium) in which proteid-containing food-bodies sprout and are eagerly sought by the ants. These pear- or egg-shaped structures, called "Müllerian bodies", resemble insect-eggs. New bodies are being formed continually in the place of those carried away by the ants. Favored by these conditions, the ants increase enormously in numbers on the *Cecropia* trees and pass their lives in the crown of foliage. Of certain species of *Cecropia* nearly all the individuals, of others only occasional trees are inhabited by ants. In their earliest youth the *Cecropias* are usually free from ants; they are not invaded by fecundated queens till they reach a height of some meters. The trees exposed to inundation cannot, of course, harbor ants, because they are kept away by the water."

The last remark is evidently incorrect, since the *Cecropias* standing in water, may be settled by Azteca queens reaching them before deälation. Moreover, the internodes of the trunk as well as those of the branches are inhabited by the ants, and Ule's statement in regard to the time of infestation of the young trees with the Azteca queens does not agree with my observations on many hundreds of plants. In British Guiana, Costa Rica, Guatemala and Panama I regularly found colony-founding queens in the internodes of all young trees, often in those only a foot or two feet high. Attention should also be called to the fact that Rettig has described another possible source of ant-food in the "pearl glands" on the leaves of the *Cecropias*. These structures are very similar to the Müllerian bodies of the trichilia and also contain proteids, oil and sugar.

It is interesting to note that the first observer to call attention to the Coccids and their nutritive value to the *Cecropia* ants was Thomas Belt (1874, p. 222), though he overlooked their other important source of food in the Müllerian bodies. This is clear from the following passage: "The stem of the *Cecropia*, or trumpet-tree, is hollow, and divided into cells by partitions that extend across the interior of the hollow trunk. The ants gain access by making a hole from the

outside, and then burrow through the partitions, thus getting the run of the whole stem. They do not obtain their food directly from the tree, but keep brown scale insects (Coccidæ) in the cells, which suck the juices from the tree, and secrete a honey-like fluid that exudes from a pore on the back, and is lapped up by the ants. In one cell eggs will be found, in another grubs, and in a third pupæ, all lying loosely. In another cell, by itself, a queen ant will be found, surrounded by walls made of a brown, waxy-looking substance, along with about a dozen Coccidæ to supply her with food. I suppose the eggs are removed as soon as laid for I never found any along with the queen ant. If the tree be shaken, the ants rush out in myriads and search about for the molester. The case is not like the last one (the bull-horn Acacia), where the tree has provided food and shelter for the ants, but rather one where the ant has taken possession of the tree, and brought with it the Coccidæ; but I believe that its presence must be beneficial. I have cut down some dozens of the *Cecropia* trees, and never could find one that was not tenanted by ants. I noticed three different species, all, as far as I know, confined to the *Cecropiæ*, and all farming scale-insects. As in the bull's-horn thorn, there is never more than one species of ant on the same tree."

H. von Ihering described an exceptional condition in *Azteca muelleri* Emery which inhabits *Cecropia adenopus*. In this case the ants construct a large spindle-shaped carton nest in the bole of the tree. The nest is small at first and produces no deformation of the bole, but as the ants enlarge the structure they cut away more and more of the surrounding wood to convert it into carton and the weight of the trunk and crown above the nest causes the bole to bulge at the weakened zone so that the presence of the nest is indicated from the outside. von Ihering interpreted the enlargement which is not produced by any other *Cecropia*-inhabiting *Azteca*, as a huge gall, but it is clear that it is nothing of the kind.

Both Schimper and H. von Ihering found that one of the Brazilian *Cecropias*, *C. hololeuca*, lacks the trichilia and Müllerian bodies and is not regularly inhabited by ants, and Bailey found a similar absence of these organs in *C. sciadophylla* var. *decurrens* in British Guiana, though it is occasionally inhabited by non-obligate Formicidæ. On the other hand, I found (1908) that a Porto Rican *Cecropia*, which I referred to *C. peltata* L. but which is probably *C. Urbaniana* Snetlage, possesses trichilia and coral red food-bodies, but is not inhabited by ants. I have made similar observations on the *Cecropias* of Cuba. In fact, the obligate *Cecropia* ants, the Aztecas, are absent from all

the West Indian Islands, except a few of the Windward group. It seems that Warming (1894, p. 185, Fig. 6) had previously noticed that the Cecropias of Aguadilla, Porto Rico, though not inhabited by ants, nevertheless possessed distinct but poorly developed trichilia. If I understand his remarks, he inferred that this might represent a degenerate condition due to the absence of the customary insect stimulation. Rettig (1904, p. 16, 21) states that von Meyer (Die Secretionsorgane der Pflanzen, Berlin, 1839) described trichilia in *Porouma guianensis* Aublet and growing among the hairs small granules obviously homologous with the Müllerian bodies of Cecropia.

Bequaert (Wheeler and Bequaert, 1929) has recently compiled the following list of Cecropias definitely known to bear trichilia and to be myrmecophilous:

(1) *C. adenopus* Miquel (= *C. peltata* Vellozo, non Linné) of Brazil. Its normal ant-inhabitant, *Azteca muelleri* Emery, builds in its trunk the large carton nest and produces the swelling above described. In Paraguay, according to Fiebrig (1909), the tree is tenanted by a different species, *A. alfari* var. *mixta*, which causes no modification of the bole.

(2) *C. angulata* Bailey. British Guiana (I. W. Bailey, 1922).

(3) *C. robusta* J. Huber. Common along in the lower Amazon in woods that are frequently flooded. Stated to be myrmecophilous (J. Huber, 1910, p. 61).

(4) *C. bifurcata* Huber. Lower Rio Purus, Brazil. Cited as myrmecophilous (J. Huber, 1910, p. 62).

(5) *C. lativirens* J. Huber. Upper Rio Purus, Brazil (J. Huber, 1910, p. 63).

(6) *C. paraensis* J. Huber. Lower Amazon. Myrmecophilous (J. Huber, 1910, p. 64).

(7) *C. distachya* J. Huber. Vicinity of Para. Myrmecophilous (J. Huber, 1910, p. 65).

(8) *C. lyratiloba* Miquel (*C. paludosa* Warburg). A swamp species of Southern Brazil which H. v. Ihering found inhabited by *Azteca*. It possesses trichilia.

(9) *C. sciadophylla* Martius. This species is doubtfully myrmecophilous since it possesses no trichilia at the base of the petiole. According to E. H. Snetlage (1923, p. 358), *C. juranyiana* Al. Richter is merely a variety of *C. sciadophylla*. Another variety, described by Snetlage as var. *decurrrens*, was recorded by Ule as inhabited by *Azteca emeryi* Forel. This record may be due to an error or the ants may merely have occupied the hollow stems. Professor Bailey, who

observed the same var. *decurrens* near Kartabo, British Guiana, regards it as non-myrmecophytic. It lacks trichilia and food-bodies, although it possesses a prostoma. He occasionally found certain inquiline ants in the internodes, but no Aztecas.

(10) *C. riparia* "Warburg", Snetlage (1923, p. 363). Brazil. This species is provided with trichilia at the bases of the petioles and Ule (1906) found the internodes occupied by *Azteca alfari* Emery var. *aequilata* Forel.

(11) *C. feifolia* "Warburg", Snetlage (1923, p. 365). Rio Acre, Brazil. The bases of the petioles bear trichilia and Ule mentions that the internodes are inhabited by *Azteca minor* Forel.

(12) *C. montana* "Warburg", Snetlage (1923, p. 368), of Peru, has trichilia and Ule found it inhabited by *Camponotus (Pseudocolobopsis) ulei* Forel.

(13) *C. mexicana* Hemsley. A number of ants were found by Ross (1909) in this species, although the bases of the petioles appear to lack trichilia.

(14) *C. leucocoma* Miquel (= *C. arenaria* Warburg). Manaos, Brazil. A true myrmecophilous species with trichilia and food bodies, as Bequaert (Wheeler and Bequaert, 1929) has shown.

(15) *C. obtusa* Trecul. Rio Negro and Rio Branco, Brazil. Like the preceding species, a true myrmecophyte. Studied by Bequaert (Wheeler and Bequaert, 1929).

Bequaert remarks that many other species of *Cecropia* are known to possess trichilia, e.g., *C. carbonaria* Martius and Miquel, *C. cyrtostachya* Miquel, *C. Dielsiana* Snetlage, *C. Engleriana* Snetlage, *C. Francisci* Snetlage, *C. Glaziovii* Snetlage, *C. leucophæa* Pöppig and Miquel, *C. multiflora* Snetlage, *C. palmata* Willdenow (perhaps the same as *C. obtusa* Trecul), *C. saxatilis* Snetlage, *C. Ulei* Snetlage and *C. Urbaniana* Snetlage. Some botanists assume that all such species are ant-plants. It is, however, by no means certain that the presence of trichilia alone is sufficient to give them the status of myrmecophytes. This is shown by my observations on the *Cecropias* of Porto Rico and Cuba. A detailed field and taxonomic study of many more of these trees will have to be made before we shall be able to give a satisfactory account of the prevalence and meaning of myrmecophily in the genus.

The ants at present known to nest in the various species of *Cecropia*, *Pourouma* and *Coussapoa* are enumerated in the following list:

- (1) *Neoponera stipitum* Forel. In branches of *Cecropia sciadophylla* var. *decurrens*. British Guiana (Wheeler).

- (2) *Pseudomyrma ulei* Forel. In twigs and branches of *Coussapoa* sp. Amazonas (E. Ule).
- (3) *Pheidole guildini-muelleri* For. subsp. *avia* Forel. In trunk of *Cecropia adenopus*. Brazil (Luederwaldt).
- (4) *Solenopsis saevissima* F. Sm. In dry branch of *Cecropia* sp. Brazil (Luederwaldt).
- (5) *Crematogaster (Orthocrema) quadriformis* Mayr. In *C. adenopus* with *Azteca muelleri*. Brazil (Luederwaldt).
- (6) *Crematogaster (Orthocrema) limata* F. Sm. In internodes of *C. angulata* Bailey and *C. sciadophylla* var. *decurrens*. British Guiana (Wheeler).
- (7) *Crematogaster* sp. In dead wood of *C. adenopus* with young colonies of *Azteca muelleri*. Brazil (H. von Ihering).
- (8) *Wasmannia iheringi* Forel. In small nest constructed on a *Cecropia* leaf. Brazil (H. Luederwaldt).
- (9) *Cephalotes atratus* L. In large branch of *C. sciadophylla* var. *decurrens*. British Guiana (Wheeler).
- (10) *Cryptocerus pusillus* Klug. In dry branch of *Cecropia* sp. Brazil (Luederwaldt).
- (11) *Cryptocerus* sp. In dead wood of *C. adenopus*. Paraguay (Fiebrig).
- (12) *Azteca alfari* Emery. In *Cecropia* sp. Costa Rica (A. Alfaro, Wheeler); Panama (Champion); Venezuela; Colombia (Forel).
- (13) *Azteca alfari* var. *aqualis* Forel. In *Cecropia* sp. Brazil (A. Goeldi); Mexiana Island (Hagmann); Colombia (Lallemand).
- (14) *Azteca alfari* var. *aquilata* Forel. In *C. riparia* Warb. Brazil (E. Ule, A. Goeldi, J. Huber).
- (15) *Azteca alfari* var. *arcentina* Forel. In *C. adenopus*. San Ignacio, Misiones, Argentina (L. Hauman).
- (16) *Azteca alfari* var. *curtiscapa* Forel. In *Cecropia* sp. Panama (Christophersen, Wheeler).
- (17) *Azteca alfari* var. *fumaticeps* Forel. In *C. mexicana*. Mexico (Ross). In *Cecropia* sp. Guatemala (Wheeler).
- (18) *Azteca alfari* var. *langi* Wheeler. In *Cecropia* sp. British Guiana (H. Lang).
- (19) *Azteca alfari* var. *mixta* Forel. In *C. adenopus*. Paraguay (Fiebrig); *C. lyratiloba* Miq. Brazil (Luederwaldt); *C. leucoma* Miq. Brazil (Bequaert).
- (20) *Azteca alfari* var. *ovaticeps* Forel. In *C. lyratiloba* Miq. Brazil (Luederwaldt); In *Cecropia* sp. Brazil (Goeldi, Bequaert).

- (21) *Azteca alfari* subsp. *cecropiæ* Forel. In *C. lyratiloba*. Brazil (Luederwaldt), *C. obtusa* Trecul. Brazil (Bequaert); *C. "peltata"* Dutch Guiana (G. Stahel); British Guiana (H. E. Box); *Cecropia* sp. (Goeldi, Huber and Ule); in "swamp *Cecropia*" Brazil (H. v. Ihering).
- (22) *Azteca alfari* subsp. *lucida* Forel. In *Cecropia* sp. Guatemala (Champion; Wheeler).
- (23) *Azteca alfari* subsp. *lucidula* Forel. In *C. angulata* Bailey. British Guiana (Wheeler); *C. "peltata"* Trinidad (Wheeler); *Cecropia* sp. Guatemala (Wheeler).
- (24) *Azteca alfari* subsp. *lucidula* var. *zonalis* Wheeler. In *Cecropia* sp. Panama (Wheeler).
- (25) *Azteca alfari* subsp. *tuberosa* Forel. In *Cecropia* sp. (probably) Brazil (Diaz da Rocha).
- (26) *Azteca bicolor* Emery subsp. *belti* Emery. On trunks of *C. "peltata"* Trinidad (Wheeler).
- (27) *Azteca coeruleipennis* Emery. In *Cecropia* sp. Costa Rica (Alfaro; Wheeler); Guatemala (Champion); Mexico (Schumann); *C. mexicana* Mexico (Ross).
- (28) *Azteca constructor* Emery. In *Cecropia* sp. Costa Rica (Alfaro, Tonduz, Pittier); Guatemala (Wheeler); Panama (Christophersen, Wheeler); in *C. "peltata"*, Trinidad (Wheeler).
- (29) *Azteca constructor* var. *guianæ* Wheeler. In *C. angulata* Bailey. British Guiana (Wheeler).
- (30) *Azteca coussapoæ* Forel. In branches of *Coussapoa* sp. Amazonas (E. Ule).
- (31) *Azteca delpini* Emery. In *Cecropia* sp. (probably) Matto Grosso, Brazil (Germain).
- (32) *Azteca delpini* var. *trinidadensis* Forel. In *C. "peltata"*. Trinidad (Wheeler).
- (33) *Azteca delpini* subsp. *antillana* Forel. On *Cecropia* sp. Castries, St. Lucia (Wheeler, J. C. Bradley).
- (34) *Azteca delpini* subsp. *antillana* var. *guadeloupensis* Forel. In *Cecropia* sp. (probably) Guadeloupe (Forel); Roseau, Dominica (F. Lutz).
- (35) *Azteca duroiæ* Forel. In branches of *Pourouma* sp. Amazonas (E. Ule).
- (36) *Azteca emeryi* Forel. In *Cecropia sciadophylla* Mart. Amazonas (E. Ule).
- (37) *Azteca foreli* Emery. In *Cecropia* sp. Costa Rica (Alfaro, Wheeler); Guatemala (Wheeler).

- (38) *Azteca foreli* Emery var. *ciseni* Pergande. In *Cecropia* sp. (probably *mexicana*) Colima, Mexico (Townsend).
- (39) *Azteca foreli* subsp. *championi* Forel var. *breviscapa* Forel. In *Cecropia* sp. Costa Rica (Tonduz).
- (40) *Azteca foreli* subsp. *ursina* Forel. In *Cecropia* sp. Trinidad (Wheeler).
- (41) *Azteca instabilis* F. Sm. In *Cecropia* sp. (Mexico, Colombia, French Guiana, Central America); in *C. angulata* Bailey. British Guiana (Wheeler).
- (42) *Azteca lanuginosa* Emery. In *C. hololeuca* Miq. Brazil (Luederwaldt) In *C. adenopus* Brazil (H. von Ihering); in carton nests on *Cecropia* sp. (W. Ehrhardt).
- (43) *Azteca minor* Forel. In *C. ficifolia* Warb. Amazonas (E. Ule).
- (44) *Azteca muelleri* Emery. In *C. adenopus* Brazil (F. Müller, Hetschko, H. von Ihering, Luederwaldt, J. C. Bradley).
- (45) *Azteca schimperi* Emery. In carton nest on *C. leucocoma* Miq. Manaus, Brazil (Bequaert).
- (46) *Azteca sericea* Mayr. In *Cecropia* sp. Costa Rica and Guatemala (Wheeler).
- (47) *Azteca trigona* Forel subsp. *mathildæ* Forel var. *spuria* Forel. In *Cecropia* sp. British Guiana (Wheeler).
- (48) *Azteca trigona* subsp. *mediops* Forel. In *C. angulata* Bailey. British Guiana (Wheeler).
- (49) *Azteca ulci* Forel var. *gibbifera* Forel. In trunk of *Cecropia* sp. Brazil (Luederwaldt).
- (50) *Azteca ulci* subsp. *nigricornis* Forel. In *C. adenopus*. Brazil (Luederwaldt).
- (51) *Azteca xanthochroa* Roger. In *Cecropia* sp. Costa Rica (Alfaro, Schwarz and Barber); Guatemala (Wheeler); in *C. mexicana*. Mexico (Ross).
- (52) *Azteca xanthochroa* var. *costaricensis* Wheeler. In *Cecropia* sp. Costa Rica (Alfaro).
- (53) *Azteca xanthochroa* subsp. *australis* Wheeler. In *Cecropia* sp. (very probably), Peru (Staudinger).
- (54) *Azteca xanthochroa* subsp. *salti* Wheeler. In *Cecropia* sp. Colombia (G. Salt.)
- (55) *Azteca xanthochroa* subsp. *isthmica* Wheeler. In *Cecropia* sp. Panama (Wheeler).
- (56) *Camponotus (Tanæmyrmex) bonariensis* Mayr. In dead wood of *C. adenopus*. Paraguay (Fiebrig).

- (57) *Camponotus* (*Tanæmyrmex*) *melanoticus* Emery var. *hagmanni* Forel. In *C. sciadophylla* var. *decurrentis*. British Guiana (Wheeler).
- (58) *Camponotus* (*Myrmothrix*) *balzani* Emery. In dead branches of *C. adenopus*. Brazil (H. von Ihering).
- (59) *Camponotus* (*Myrmothrix*) *rufipes* Fabr. subsp. *renaggeri* Emery. In dead wood of *C. adenopus*. Paraguay (Fiebrig).
- (60) *Camponotus* (*Myrmobrachys*) *canescens* Mayr. In trunk of *C. adenopus*. Brazil (Luederwaldt).
- (61) *Camponotus* (*Myrmobrachys*) *veznyi* Forel. Occasionally in dead wood of *C. adenopus*. Paraguay (Fiebrig).
- (62) *Camponotus* (*Pseudocolobopsis*) *ulei* Forel. In *C. montana* Warb. Peru (E. Ule).

Probably all the ants in this list, except the Aztecas, are to be regarded as merely occasional inhabitants, or inquilines. And even among the Aztecas only certain species would seem to be really obligatory tenants of the plants—namely *A. alfari*, *delpini*, *coeruleipennis*, *constructor*, *muelleri* and *xanthochroa*. Of these *A. alfari* is the Cecropia ant *par excellencæ*. Its geographical range is almost coextensive with that of the plants. It is represented by some 14 different varieties and subspecies, ranging from Mexico to Paraguay and Argentina, and is known to inhabit at least six or seven different species of Cecropia. No doubt further collecting will add considerably to the number of recognizable forms of this highly variable insect.

While the earlier observers, Fritz Müller and Schimper, were inclined to regard the Aztecas as the protectors of the Cecropias against the attacks of the Attini, or leaf-cutting ants, all the more recent authors, including H. von Ihering, Rettig, Fiebrig, Ule, Bailey and myself, have come to see in the Attini no such menace to the life or well-being of the plants. The various arguments in support of this contention have been set forth in detail, especially in the contributions of von Ihering, Fiebrig and Bailey, and need not be repeated here. That the Cecropias are centers of important biocœnoses consisting of a great number and variety of organisms, some of which are far more injurious than the Attini, was first shown by Fiebrig in his study of *C. adenopus* in Paraguay. A conspectus of the forms he recorded together with additions by von Ihering, Bailey, Zetek and others, is given in the following paragraphs:

Mammalia. Fiebrig mentions apes and bats (*Phyllostoma* sp.) as eagerly feeding on the fruit of *C. adenopus*. The most important mammal, however, is the sloth, *Bradypus tridactylus* L., which accord-

ing to Bates, H. von Ihering and others regularly devours the foliage and is *not* molested by the Aztecas. The sloth's relation to the tree is, in fact, so constant and so well known to the natives that the name "imbauba" and its derivatives really mean "slothtree". Menegaux (1909), in his monograph of the genus *Bradypus*, states, on the authority of Geay, that the sloth feeds only on *Cecropia* leaves and cites Gmelin (1788), who says that *B. tridactylus* "vicitat foliis teneris imprimis *Cecropiæ*, non bibit, imbre metuit." Observations on the sloths at the Barro Colorado and Kartabo laboratories confirm this statement in regard to the feeding habits of the animal.

Aves. Fiebrig mentions doves as occupying the *adenopus* trees and feeding on their ripe fruit and woodpeckers as devouring the *Heliothis* larvæ (*vide infra*) which live in the twigs and branches. He also observed birds' nests among the foliage and figures one of them.

Lepidoptera. No less than six different moths and butterflies feed as larvæ on the large leaves of the *Cecropias*:

(1) *Gynaecia dirce* L. The caterpillars of this beautiful Nymphalid were found feeding in great numbers on *Cecropia* at Ancon by Mr. Zetek. They are black with branched orange-colored spines. The pupa is gray and provided with two broad horns at the anterior end. Gundlach (Wolcott, 1923) had previously noted the same habit of the caterpillars in Porto Rico: "la oruga vive debajo de hoja de *Cecropia*, comiendo las nervosas gruesas."

(2) *Historis orion* Fabr. Mr. Zetek also found the caterpillars of this Nymphalid devouring the foliage of the *Cecropias* at Ancon and reared the butterflies. The caterpillar is described by Wolcott (1923) as flattish, medium gray, with white saddle and states that Gundlach notes its occurrence on *C. peltata* in Porto Rico ("la oruga se crea en la *Cecropia*).

(3) The Syntomid moth, *Correbidia terminalis* Walker, is cited by Wolcott from Porto Rico with the following note by Gundlach: "vive en la cara inferior de las hojas de *Cecropia*, formando luego im capullo poco primeroso."

(4) The Diopiid moth, *Diops ocellata* Cr., is another *Cecropia* pest which was observed by Fiebrig in Paraguay. "The hairy larva of *Diops ocellata* Cr., devours the large leaves. It is not easily detected on the white-felted undersides on account of its chalk-white, black dotted coloration. I often found this larva, which reaches a length of 4 cm., in considerable numbers on a tree."

(5) Fiebrig describes another moth caterpillar (undertermined) "of respectable dimensions" as living in the cambium layer of the *C.*

adenopus trunk. The pupa "rests beneath the bark, which at one point is perforated by a T-shaped slit that enables the beautiful chocolate brown and sulphur-yellow moth to escape. The injury to the tree caused by this caterpillar is probably unimportant."

(6) *Heliothis* sp. Fiebrig also describes in detail the habits of an undetermined boll-worm. He says: "The third caterpillar which I observed on *Cecropia* belongs to the moth 7584 (*Heliothis* sp. in litt.) and is unquestionably the most interesting. It competes with the Azteca (*alfari* var. *mixta*) and lives in the younger portions of the twigs, precisely where the Dolichoderine is encountered. Such caterpillar-infested twigs are extremely common at all seasons of the year. At certain times I found them on nearly every tree I examined and in large trees with 60-80 main and accessory twigs I have found 30-50 of them occupied by the twig-mining caterpillars." After describing the perforation of the twigs by the woodpeckers which feed on the caterpillars, Fiebrig continues: "I was long uncertain in regard to the time and place of the caterpillar invasion till I succeeded in finding their youngest stage, scarcely more than a millimeter long, in company with the Aztecas that were just founding their colonies in the uppermost chambers! Here I detected the obviously just hatched caterpillar in the completely closed internodal cavity beside the egg-laying ant-queen with her eggs and larvæ and embedded in and evidently feeding on a damp brown mass of pith which had been chewed up by the mother ant. On several occasions also I found these small caterpillars up to a length of 6 mm. in the initial chambers of both young and older trees and repeatedly in company with just-emerged Azteca workers. Later, however, when, after perforation of the dissepiments, a greater portion of the twig becomes habitable by the ants, the caterpillars and ants were no longer observed in one another's company. Often, indeed, I found them separated, the Azteca being in the outermost tip of the twig and the caterpillar below, shut off by an intact dissepiment. While, therefore, at first the ant and caterpillar live peaceably side by side—a condition the more remarkable, because the small, large-headed, strongly mandibulate caterpillars differ so markedly from the white, nearly naked Azteca larvæ in their darker color and habitus, and especially in possessing stiff bristles—later the "pugnacious" ants, which move upwards towards the tips of the branches, followed by the caterpillars, nevertheless leave the latter in full possession of the *Cecropia* twigs. Since we cannot suppose that the soft-skinned caterpillars are aggressive, we shall not be mistaken in inferring that the ants are driven out

by the masses of caterpillar frass, often spun together, which plug up the chambers and render them uninhabitable. It is also possible that the caterpillars, which are capable of emitting considerable mucus, may use their silk as a means of defense. Only after the caterpillars have completely eaten out the ambay twig, does one of their number about to pupate, gnaw a large hole in the wall of the twig at one of the former ant-entrances. This opening is used as an exit not only by the individual that made it but also by all the other moths that develop in the twig! "It would seem, therefore, that the *Heliothis* caterpillars in their younger stages are synœketes of the Azteca, that after hatching from eggs laid by the mother moth on the outer surface of the plant, they probably enter the internodes through the perforations made by the nest-founding queens and that later the peculiar habits of the caterpillar render any further synœcetism with the ants impossible. It is probable that this same *Heliothis* larva was observed by Schimper (1888). At any rate, his Fig. 3, Pl. II, shows a similar caterpillar lying on a large mass of frass in a young *adenopus* internode just above a cavity inhabited by Azteca.

(7) Another undetermined Lepidopteron caterpillar is mentioned by H. von Ihering in Eastern Brazil.

Hymenoptera. The following Chalcidids, Vespids and Apids have been found associating with *Cecropias* or their ants:

(1) *Conoaxima affinis* Brues (1922). This Chalcid was described from a specimen which I took in a *Cecropia* internode (not in an *Acacia* spine, as stated by Brues!) at Quirigua, Guatemala, in 1911. I have since taken both sexes of the species at Balboa, C. Z., and have seen the larvæ in various stages devouring the young colony-forming queens of *Azteca alfari* subsp. *lucidula* var. *canalis*.

(2) *Conoaxima aztecicida* Brues. The habits of this species, which was taken in 1920 by Professor Bailey in internodes of *C. angulata* at Kartabo, B. G., are the same as those of the preceding species. The larvæ were found destroying the young nest-founding queens of *A. constructor* and *A. alfari*. Professor Bailey's observations are recorded in the following note: "Although many of the successive internodal cavities of each young plant become inhabited, few of the queens succeed in raising a brood. When the stems are cut open, most of the chambers are found to contain dead queens. I was unable to account for this high mortality until I discovered the presence of a small scar in the callus which fills the entrance aperture. This scar within a scar indicated, of course, that some insect had emerged since the queen became sealed within her domatium. Following up this clue, I soon

found chambers—with modified callus in the apertures—which contained, in addition to the dead and frequently dismembered queen, the larva, pupa, or imago of a Hymenopterous parasite. The evidence at hand seems to indicate that the queens are parasitized before they enter their dwellings.”

(3) *Conoaxima* sp. A specimen of an undescribed species of *Conoaxima* was recently found by Dr. George Salt in the topmost internode of a young *Cecropia* at Vista Nieve, at an altitude of 5000 ft. in the Sierra Nevada de Santa Marta, Colombia. The internodes of the plant, which was only about 6 ft. high, were occupied either with solitary nest founding queens of a new subspecies of *Azteca xanthochroa* Roger, which I have called *salti*, or lower down with young colonies consisting of queens with brood or with brood and small workers. In the internode containing the *Conoaxima* was a mouldy mass, in all probability the remains of an *Azteca* queen that had been devoured by the parasite.

(4) H. von Ihering mentions a rather large Chalcid which he found in the midst of ants (*Azteca muelleri*) in *C. adenopus* at Alto da Serra, Brazil. In all probability the insect was one of the Eucharidæ, many species of which are known to parasitize ant-larvæ.

(5) The Calverts (1917) found on one of the branches of a *Cecropia* in Costa Rica a large pendent paper-covered nest which was occupied by an active colony of wasps (apparently some species of *Polybia*) for at least four months.

(6) *Trigona* sp. At Kartabo, B. G. Professor Bailey discovered a colony of a very small stingless bee, which Professor Cockerell has identified as an undescribed species, near *goeldiana*, in the internodes of a branch of *Cecropia angulata*. This nest is figured in my "Social Life among the Insects (1923)". The same bee was also found nesting in a hollow liana.

Diptera. The following five Diptera seem in their larval stages to be either scavengers in decomposing portions of the *Cecropias* or parasites of leaf-eating species.

(1) Great numbers of a Tachinid fly were bred by Mr. Zetek from the pupæ of the Chrysomelid beetle, *Calomera cayennensis* Fabr. (*vide infra*), which feeds on the foliage of the *Cecropias* at Ancon, C. Z.

(2) H. von Ihering bred a species of *Drosophila* from larvæ living in matter derived from the decomposition of vegetable excrescences in an internode of *C. adenopus*.

(3) Prof. Bailey bred numbers of a species of *Drosophila* from the male aments of *C. angulata* at Kartabo, B. G.

(4) von Ihering mentions slender Dipteron larvæ in soft, gelatinous excrescences in the upper internodes of *C. adenopus*.

(5) Fiebrig found the moist frass masses of the *Heliothis* caterpillars described above to contain Dipteron larvæ.

Colcoptera. Some six or eight different beetles have been found to attack *Cecropias*:

(1) A small unidentified Bostrichid, according to Fiebrig, feeds not only on the dead twigs of *C. adenopus* but often also on the green internodes inhabited by the Aztecas. The female beetles lay their eggs in the fresh cambium and the hatching larvæ completely destroy this layer and also bore into the wood.

(2) Fiebrig found a Cerambycid which bores both in the dry twigs and in the wood of the trunk.

(3) *Coclomera cayennensis* Fabr. According to H. von Ihering, this Chrysomelid and a similar but smaller species lay their eggs on the leaves of *C. adenopus* and there undergo their entire development. "The larvæ gnaw away the upper layers of the leaf but are never molested by the ants, notwithstanding their great injury to the foliage." Mr. Zetek has found this same beetle in all stages on the *Cecropias* at Ancon, C. Z. and has bred from its pupæ numbers of Tachinid flies (*vide supra*). The full-grown larvæ measure about 15 mm. and have the abdominal segments provided laterally with blunt-spines and the last segment developed as a large, rugulose, shovel-shaped organ. They are opaque blackish with the head, pronotum, legs and several tuberculate spots on the dorsal surface of each abdominal segment, except the last, dull orange yellow. Most of the larvæ pupated January 16, the pupæ being naked and attached to the leaves by their posterior ends. The imaginal beetles, which emerged February 20, have opaque black elytra, with the remainder of the body more shining and of a dirty fulvous tint, except the antennæ, tibiæ, tarsi and venter, which are black. The *Cecropias* are very probably the true host plants of this beetle, which has very nearly the same geographical distribution. It was originally described from French Guiana. In the Bowditch Collection of Chrysomelidæ in the Museum of Comparative Zoölogy I have seen specimens from Peru, Colombia, Venezuela, Bolivia, Panama, Yucatan and Rio de Janeiro, Brazil.

(4) Fiebrig mentions a Chrysomelid, possibly *C. cayennensis* or an allied species, as feeding on the leaves of *C. adenopus* in Paraguay.

(5) He also mentions a very common Chrysomelid larva, evidently one of the Cassidinæ, as feeding in dense droves on the foliage. It

possesses a blackish brown, chitinous, disk-shaped process, which it holds over its back when disturbed, like the similar processes, often covered with fæces, seen in other members of the group.

(6) The larva of a Hispidine Chrysomelid, according to Fiebrig, often mines the lobes of the leaves in succession, surrounding each with a gallery and eventually pupating and completing its development in a boss-shaped gall at the base of one of the lobes.

(7) A small yellow Curculionid is said by Fiebrig to visit the catkins.

(8) Great numbers of small Curculionids, possibly allied to the preceding species, were found by Professor Bailey in the male aments of *C. angulata* at Kartabo, B. G.

Orthoptera. Fiebrig observed Mantids and migratory locusts on the foliage of *C. adenopus*. The latter almost completely defoliated whole groups of the trees in the course of a few hours.

Thysanoptera. Very common, according to Fiebrig, on the under surfaces of the leaves.

Heteroptera. Fiebrig records several Lygæids, especially a pale brown and a delicate white species, as infesting the lower surfaces of the leaves, and a stout brown Pentatomid which guards its eggs on the foliage. All three of these bugs are extremely common and may occur on the same tree. They are in all probability very injurious.

Homoptera. Fiebrig mentions a prettily colored Jassid as being common in all stages on the leaves. He also found an aphid and a pinkish Coccid living with the ants in the internodes. Wolcott (1923) records *Aphis gossypii* Glover as occurring on the leaves of *C. peltata* at Lares, Porto Rico.

Acarina. A species of mite is recorded by Fiebrig as occurring among the Thysanoptera on the undersides of the leaves of *C. adenopus*.

Nematoda. A Tylenchus-like form was found by Fiebrig living among the moist detritus left by the Heliothis larvæ in the twigs.

Fungi. Moulds occur in the same situation as the Nematodes and are often found covering the bodies of young queens that have died in the internodes before establishing colonies.

This list comprises some forty different organisms, mostly associated with a single species of Cecropia (*adenopus*). Fiebrig states that he has omitted a number of forms and remarks: "When I compare them with the Hexapods occurring in Paraguay on other plants, it seems to me that the ambay is more thoroughly infested than the great majority of the woody plants of the region, and this the more astonishing, because its leaves are not enticing nor juicy and because the tree might seem

to be protected from most of its enemies by the peculiar properties of its sap."

Among the organisms enumerated in the foregoing list the Coccids occupy a constant and privileged position. All those who have studied the Cecropias carefully—Belt, Fritz Müller, Schimper, Warming, H. von Ihering, and Fiebrig—have noticed these insects and all, with the exception of Fiebrig, have regarded them as an important source of food for the Aztecas. Fiebrig's assertion that they "have no direct relations with the ants" must be attributed to faulty observation. Recently Professor Bailey has devoted special attention to the Cecropia coccids and has made some observations of interest in connection with the species occurring in other myrmecophytes. I quote from his paper (1922, p. 388) which owing to its publication in a botanical journal, may be readily overlooked by entomologists: "Having found a very close and significant relation between ants and coccids in most Ethiopian ant-plants, I devoted particular attention to the investigation of their behavior in *C. angulata*. I did not succeed in finding a single large, ant-inhabited specimen which did not contain numerous coccids. When a tree is split open the ants are as solicitous for the welfare of the coccids as they are for that of their eggs, larvæ and pupæ. They seize them in their mandibles and carry them about until some unopened portion of the plant is found where they may be deposited in safety. In artificial nests, the workers spend hours in tending and stroking the coccids, and in feeding upon their sugary exudates. In view of these facts, it cannot be doubted that the miniature milch cows are an important source of liquid carbohydrates for the ants.

"As in many of the African myrmecophytes the ants excavate pits in the walls of their domatia which enable the coccids to reach and feed upon the softer tissues of the Cecropia. Such excavations are essential, owing to the fact that the internodal, medullary cavity is entirely jacketed by a dense, horny layer of sclerenchyma. In the African ant-plants, the ants cut through to the cambium and induce the formation of a nutritive callus. In *C. angulata* the pits are not located in the sides of the internodal chamber, but in the nodal diaphragms. At the time when the ants begin their excavations, the nodal partition consists of five distinct layers. The soft internal layer, which is provided with strands of conducting tissue and which is fed upon by the coccids, is separated from the external layers of porous, medullary tissue by two layers of dense, thick-walled tissue. The ants remove the two external layers and cut circular pits in the underlying sheets of horny sclerenchyma. The coccids sit in these pits and thrust

their setæ into the succulent tissue which is thus exposed. That the pits are not made by the coccids, as suggested by von Ihering is indicated, not only by the fact that the delicate sucking mouthparts of these insects are not adapted for excavating in dense tissues, but also by the fact that I have actually observed the ants in the process of excavating them."

The preceding account of the Cecropias shows that the general picture of their relations to the ants is very similar to those of other myrmecophytes with hollow or fistulose stems or petioles, such as the Cordias, Tachigalias and Triplarises, but there are two peculiarities, the preformed pit, or prostoma of the internodes and the Müllerian bodies, which have been repeatedly cited by authors as irrefutable proof of the adaptation of the trees to the Aztecas. The prostoma, however, is now generally conceded to be merely a deepening of the groove, or rill formed by pressure of the axillary bud. Prof. Bailey found that as a rule no pit is present in *C. angulata*, but merely a broad groove, whereas in *C. sciadophylla* var. *decurrens*, which is not inhabited by Aztecas, the thinning of the internodal wall in the groove is much greater than in *angulata*. "Such facts as these," he says, "suggest that the so-called prostoma of *C. adenopus*, and of other myrmecophytic species of Cecropia, is not an adaptation for attracting ants, but is merely a structural peculiarity, produced by the pressure of the axillary bud, which is utilized by the ants in their parasitism upon the plants."

There remain then only the Muellerian bodies and possibly the pearl glands to support the view that the Cecropias have developed specific attractions or inducements for the ants. But the botanists have recently also cast doubts on the finalistic interpretation of these structures. The matter is well summarized in the following passage from Professor Bailey's paper (1922, p. 387): "These small bead-like structures, which are packed with fat and protein, are formed in large numbers in a curious cushion or mat of hairs situated at the base of each petiole. The ripe food bodies are so assiduously collected by the ants that it is almost impossible to find one *in situ*, except in young uninhabited plants. Indeed, the ants frequently trim away the surrounding hairs and dig out the immature food bodies. Schimper interpreted these so-called Müllerian corpuseles, and similar structures which occur on the leaflets of certain myrmecophytic species of Acacia, as metamorphosed glands or highly specialized allurements for attracting ants. Rettig and others, however, have called attention to the fact that such glands occur on plants that

are not frequented by ants, and it is difficult for the adherents of myrmecophily to account for such occurrences without resorting to the purely gratuitous assumption that they are survivals from former symbioses. Ule is of the opinion, in addition, that the expenditure of carbohydrates and nitrogenous substances, contained in these corpuscles, is not compensated for by the protection which the ants afford to the plants."

Chapter 5. NOTES ON RUBIACEÆ AND VERBENACEÆ

Of all the families of plants the Rubiaceæ comprise the greatest number of myrmecophytes, but curiously enough, nearly all of these are paleotropical. From the Neotropical Region only a few species of *Duroia*, *Remigia* and *Patima* are known to be associated with ants. Though the genera *Randia*, *Uncaria*, and *Psychotria* occur in both hemispheres, and in Africa and New Guinea comprise some noteworthy ant-plants, none of the American species, so far as known, has developed this peculiarity.

(A) *Duroia* and *Remigia*

The genus *Duroia*, which is confined to the Amazonian region and northern South America, includes 10 species, three of which are known to be myrmecophilous, namely *D. hirsuta* Pöppig and Endlicher, *petiolaris* J. D. Hooker and *saccifera* (Martius). The last is provided with a pair of peculiar sac-shaped domatia at the bases of the leaf-blades; the other species have no such structures but instead present swollen internodes which are inhabited by ants. A fourth species, *D. dioica* Karsten has similar swollen internodes but these are not definitely known to be inhabited by ants. Only one of the 14 known species of *Remigia*, *R. physophora* Bentham, is myrmecophilous and like *D. saccifera* possesses a pair of pouches at the leaf-bases.

Spruce (1869, in 1908, p. 396) has the following notes on *Duroia saccifera* (= *Amajoua* or *Amaioua saccifera*, incorrectly cited as "*Amaiona*") and *Remigia physophora*: "Rubiads afford a few instances of sac-bearing leaves, especially in the genus *Amaiona* (Aubl.). In caatingas of the Rio Negro, almost throughout its extent, grows *Amaiona saccifera* Mart., a small bushy tree with leaves three together, above a foot long, obovate with a minute apiculus, tapering to the base, where there are two contiguous sacs inhabited by small red

fire-ants. The fruit resembles a large plum (except that like the leaves it is harshly hairy), and when ripe is soft and edible; but long before it reaches that stage the ants crowd on it and seem to suck the juices through the pores of the cuticle. To the same order belongs *Remigia physophora* Bth., a remarkable tree found at the falls of the Uaupes, having the aspect of an Amaiona, but the dry capsules and other characters of *Cinchona* and its allies. The opposite leaves, 9 inches long, are oblong-oval, obtuse with a short apiculus, near the base abruptly panduriform and bearing a small ant sac on the midrib. All the other known species of this large genus have non-sacciferous leaves."

In 1889 Schumann also published on *Duroia* and *Remijia*. Of the former genus he studied the three species *petiolaris*, *hirsuta* and *saccifera*, but his account is too long for quotation. In *petiolaris* and *hirsuta* the internodes of the terminal flower-bearing branches form hollow, spindle-shaped thickenings distally and these are provided with a longitudinal cleft or sometimes with two clefts, one above the other, in which the ants make their entrances. The third species, *saccifera*, has, as Spruce observed, a pair of sacs at the base of the leaf on the upper side. Each sac has its own preformed entrance, which is on the upper side of the leaf blade. Neither Schumann's figure nor his description yield a very clear notion of the position of these entrances. He also gives a brief account of *Remijia physophora*, which has similar sacs but their openings are on the lower surface of the leaf-base.

Ule's description (1907) of *D. hirsuta*, which he observed in the field, is very clear. He writes: "To the family Rubiaceæ belongs *Duroia hirsuta* K. Sch., a small dioecious tree, 3 to 5 meters high, which grows both in the inundated forest and on *terra firma*. The short-petioled leaves are about 18 to 24 cm. long and 7 to 10 cm. broad and ovate. Above and along their edges they bear rather long, sparse, eventually deciduous hairs; beneath the ribs especially are always coarsely hairy. The stipules are united to form an elongate conical hood, covered basally with a circlet of long hairs, among which scattered glands persist. *Duroia* has terminal, corymbose cymes of white flowers. The terminal branches consist of an elongate internode, which is followed by a very short one and finally by the inflorescence. Now the elongate internode also develops towards its tip a bladder-like dilation which bursts open on the underside and there displays a groove the tips of which heal over. In this slit there is a perforation made by the ants. These hollow stems are always inhabited by two species of ants, *Myrmelachista nigella* Roger and *Azteca duroiæ* Forel n. sp." The

formation of the internodal cleft by dehiscence, as described by Ule, is interesting in connection with the similar formation in *Triplaris* (see p. 49).

More recently Bequaert (Wheeler and Bequaert, 1929, p. 16) has been able to study *D. saccifera* in the type locality near Manaos, Brazil, where he found it growing in dense thickets of the second growth woods. He describes the plant and its domatia as follows: "It forms either a low bush about 6 feet high, with hard, woody twigs, or a small tree, 8 to 10 feet high, with an unbranched main stem of about 5 feet. The leaves are placed in whorls of three, more rarely opposite, especially in young plants, and near the base of the branches. They measure 15 to 30 cm. in length and 5.5 to 12 cm. in greatest width and are elongate-oval, with entire margins, ending in a short slender point. The basal half is gradually attenuate, the base itself briefly rounded off. The petiole is very short and almost wholly occupied by the lateral pouches. At the tips of the branches the leaves form an elongate bud enclosed in a large bract, which drops off after the leaves develop. The entire plant is covered with hispid, somewhat whitish hairs, which are particularly long on the stem, the midrib of the leaves and the myrmecodomatia. Neither flowers nor fruit were seen." Spruce's account of these has been quoted. "The myrmecodomatia of *D. saccifera* are two pouches at the base of the leaf on each side of the petiole. The sacs are completely separated from each other by the whole width of the petiole on the upper as well as on the under side of the leaf. They are nearly symmetrical, ovate, 10 to 14 mm. long, 6 to 7 mm. wide and 4 to 6 mm. thick, and are completely closed at the lower end and along the petiole, but open at the upper end by a narrow slit in the deep, sinuous notch that divides the base of the leaf-blade from the pouch. The pouches might be regarded as having been formed by the recurving of the decurrent bases of the blade to the upper side of the leaf, and the growing fast of the free edges of the recurved portion of the petiole. If this conception of their mode of origin is correct, the outer surface of the domatia corresponds to the underside of the blade and their inner surface to the upperside. Both the inner and outer surfaces are densely covered with long, hispid hairs. Those of the inner surface converge towards the upper slit, which they partly close. The ants found in the myrmecodomatia were *Solenopsis corticalis* Forel and *Brachymyrmex heeri* Forel var. *aphidicola* Forel. It is interesting to note that in spite of the pouches being inhabited by ants, the young leaves of this plant were badly damaged by leaf-cutting ants (*Attini*), while the older leaves were eaten by caterpillars. The same

species of *Duroia* was observed at Carmo, on the right bank of the Rio Branco, September 1. In this locality the ants found in the pouches were *Azteca ulei* Forel var. *cordiæ* Forel."

The various ants which have been taken in the domatia of *Duroia* and *Remijia* are enumerated in the following list:

(1) *Solenopsis corticalis* Forel. In leaf-sacs of *Duroia saccifera*. Amazonas (Bequaert).

(2) *Allomerus 10-articulatus* Mayr. subsp. *8-articulatus*. In leaf-sacs of *Remijia physophora*. Amazonas (E. Ule).

(3) *Allomerus 10-articulatus* subsp. *7-articulatus* Mayr. In leaf-sacs of *D. saccifera* Amazonas. (E. Ule).

(4) *Azteca angusticeps* Emery. In internodes of *D. petiolaris*. Amazonas (K. Schumann).

(5) *Azteca depilis* Emery. In internodes of *D. hirsuta*. Amazonas (K. Schumann).

(6) *Azteca duroiæ* Forel. In internodes of *D. hirsuta*. Amazonas (E. Ule, A. Goeldi and J. Huber).

(7) *Azteca ulei* Forel var. *cordiæ* Forel. In leaf-sacs of *D. saccifera*. Amazonas (Bequaert).

(8) *Myrmelachista (Decamera) nigella* Roger. In internodes of *D. hirsuta*. Amazonas (E. Ule).

(9) *Myrmelachista (Decamera) schumannii* Emery. In internodes of *D. hirsuta*. Colombia (K. Schumann).

(10) *Bruchymyrmex heeri* Forel var. *aphidicola* Forel. In leaf-sacs of *D. saccifera*. Amazonas (Bequaert).

Owing to the small size of the domatia, especially the leaf-sacs, all of these ants belong to diminutive or very small species. Moreover, most of them are timid and harmless and none of them can be of any considerable use to the plants. So far as we are able to judge from the list, the ant fauna of the *Duroias* varies with the locality and consists largely of forms which occur also in *Cordia nodosa* and the Melastomaceæ, which have leaf-sacs (vide infra) much like *D. saccifera* and *Remijia physophora*, or in almost any convenient plant cavities.

(B) PATIMA FORMICARIA Johnston

Mr. H. O. Lang, while on an expedition in British Guiana during 1922, discovered another Rubiaceous myrmecophyte with fistulose stems which are regularly inhabited by ants. He kindly sent me a number of herbarium specimens of the plant together with its tenants. Dr. I. M. Johnston of the Gray Herbarium has since described it as

Patima formicaria (1924, p. 83), a species allied to *P. guianensis* Aublet, "but having taller tetragonal stems, larger long-acuminate leaves, larger flowers and apparently many-celled fruit." Mr. Lang has supplied the accompanying photographs (Plates 13-14) and the following field note: "The plant was growing in most cases as single stems like large-leaved shoots. Though this no doubt is its regular way, in a few instances there were as many as four or five together arising from the same root. They were about 6 to 9 feet high, with lemon-yellow flowers and spinach-green fruit. I found them growing in the tropical rain forest trees about 150 feet high and collected about 30 of these shoots from different places on the hill which is just at the mouth of the Merume Creek, facing the Mazaruni River. Although all the shoots of the plants were hollow, not all of them contained ants." The Gray Herbarium also possesses specimens of this plant collected by Hitchcock at Tumatumari on the Potaro River, B. G.

The specimens received from Mr. Lang show that the plant is very smooth, with long, lanceolate, opposite leaves and tetragonal fistulose stems about 1.5 to 2 cm. in diameter. The longer internodes measure about 5 to 10 cm. and have large medullary cavities. Some of the shorter internodes are more swollen in the middle, or fusiform. There are elliptical openings made by the ants just below the nodes, and the insects gnaw through the latter, thus making the internodal cavities continuous. No Coccids were present on the walls of the cavities. The following ants were taken by Mr. Lang in the living internodes:

- (1) *Neoponera carinulata* Roger.
- (2) *Pseudomyrma tenuis* Mayr.
- (3) *Crematogaster (Orthocrema) limata* F. Sm. var. *palans* Forel.
- (4) *Azteca traili* Emery.
- (5) *Camponotus (Paracolobopsis) patimæ* Wheeler.
- (6) *Camponotus (Paracolobopsis) patimæ* var. *dolentulus* Wheeler.

The first to fourth in this list are species that occur also in dead twigs of various neotropical plants. The *Camponotus* may have similar habits or it may, perhaps, prove to be an obligate tenant of the *Patima*.

Borreria verticillata L. is yet another Rubiaceous plant in the stems of which I found a number of ants in British Guiana, but as only the dead stems are inhabited, this case had best be considered in connection with a number of other nonmyrmecophytic plants. (see p. 92).

(C) CLERODENDRON SIPHONANTHUS R. Br.

No verbenaceous myrmecophytes have been recorded from the New World though there are several belonging to the genus *Clerodendron* and at least one belonging to the genus *Vitex* in the Old World tropics. Mr. Zetek and I, while examining the ornamental trees and shrubs planted many years ago on the Prado at Balboa, C. Z., came upon a specimen of the Oriental *Clerodendron Siphonanthus* R. Br., which must have been set out as a seedling. Having attained a height of about seven feet, it was beginning to die as a result of heavy infestation by scale insects. It had evidently flowered and fruited in preceding years, because there were growing near it a number of vigorous young plants from a few inches to about three feet high. On closer examination the old plant was seen to be tenanted by two species of ants, *Pseudomyrma gracilis* var. *bicolor* and *Azteca velox* subsp. *nigriventris*. The former occupied only a single branch, the latter all the remaining branches and the trunk. The insects were living in the internodes, whose medullary cavities they had made continuous by gnawing through the nodes. In the smaller branches these cavities were as much as 4 to 5 mm., in the larger branches and trunk only 2.5 to 3 mm. in diameter. The ants had made no entrances in the walls, but seemed to have gained access to the medullary cavities through the broken ends of a few of the branches. At intervals on the walls of the internodal cavity there were latrines, *i.e.* blackened accumulations of fæces and infrabuccal pellets, invaded by many Nematodes, moulds and bacteria. Dr. Cobb has studied the Nematodes. On the surfaces of the branches the ants were busily attending great numbers of Coccids, among which Mr. Zetek distinguished four species. The most abundant was identified by Dr. H. Morrison as *Saissetia hæmispherica* Targ. Two species of Aleurodidæ and two species of spiders were taken on the foliage.

I have recorded this instance of an Oriental plant, belonging to a genus known to contain several myrmecophytes and peopled by Neotropical ants, because it is the converse of Morteo's case of the *Triplaris americana* introduced into the East Indies and there acquiring one of the native ants (*Dolichoderus bituberculatus*) as a tenant. I am inclined to believe that the *Clerodendron siphonanthus* at Balboa was first attacked by the scale-insects and that these then attracted the Aztecas and induced them to settle in the cavities of the stem and branches. The two cases, that of the *Triplaris* and that of the *Clerodendron*, suggest that some interesting modifications of ant-

behavior might be obtained by introducing Old World myrmecophytes into the American tropics and *vice versa*.

Chapter 6. THE MYRMECOPHYTIC ACACIAS

Several shrubs and trees of the huge Mimosaceous genus *Acacia*, which comprises more than 600 described species, widely distributed in the tropical and subtropical regions of both hemispheres, have unusually large, paired, stipular spines which are regularly inhabited by ants. These myrmecophytes may be divided into three groups, according to their peculiar discontinuous geographical distribution, the first comprising the famous bull-horn Acacias, ranging from Northern Mexico to Colombia, with a doubtful species in Cuba, the second represented by the single species, *A. cavenia*, in Paraguay, and the third comprising some seven species in the savannahs of Eastern Africa. The bull-horn Acacias present the most interesting and intricate problem and therefore require a more detailed discussion; the Paraguayan and African forms, which are perhaps not true myrmecophytes may be dismissed with briefer discussion in the concluding paragraphs of this section.

(A) The Bull-horn Acacias

The bull-horn Acacias, known to the natives of Mexico as "cuernezuelo", "cornisuelo", "palin", or "guisache corteno" and to the Panamanians as "cuernito" or "cachito", are bushes or small trees, rarely attaining a height of 20 feet, with delicate pinnately compound leaves with one or more extrafloral nectaries on the upper surface of the petiole and usually one between each pair of pinnæ on the rachis and with huge paired spines which are filled with a pulpy medullary substance till they are full-grown. The young leaflets produce small, elongate-elliptical, yellow or whitish food-bodies ("Beltian bodies") at their tips. The ants enter the mature spines by gnawing an opening just below the tip of one of the pair and then hollow both of them out, thus producing a bifurcate domatium with a single entrance. They collect, store and eat the food-bodies and visit the extra-floral nectaries. The flowers are yellow, buff or flesh-colored, small and aggregated in dense spadix-like, cylindrical spikes or globose heads. The pod-like fruits show considerable variation in structure in the different species, being indehiscent or dehiscent on one or both sides and with the seeds in some species embedded in a

sugary, pulp-like arillus. As a rule, the species prefer rather dry, open country (savannahs), but I have found one of the Panamanian forms (*A. melanoceras*) growing in swamps or moist jungle along water-courses.

Until recently it was supposed that there were less than half a dozen authentic species of bull-horn Acacias, but Safford (1910, 1914, 1915, 1923) and Schenck (1913, 1914), after studying a considerable amount of herbarium material have recognized as many as 28. These are here listed alphabetically, with their known geographical distribution:

bucrophora Robinson. British Honduras (Plate 15).

bursaria Schenck. Western Guatemala (Plates 16-17).

campecheana Schenck. Yucatan, Sinaloa (Plate 18).

chiapensis Safford. Chiapas, Mexico.

Collinsii Safford. Chiapas, Mexico (Plates 19-22).

Cookii Safford. Eastern Guatemala (Plates 23-24).

cornigera Linné. (= *spadicigera* Schl. & Cham.) Vera Cruz, Mexico (Plates 25-29).

costaricensis Schenck. Costa Rica; Nicaragua (Plate 30).

cubensis Schenck. Northern Cuba.

dolichocephala Safford. Veracruz, Mexico (Plate 31).

donnelliana Safford. Honduras (Plates 32-33 a).

furcella Safford. Veracruz, Mexico.

globulifera Safford. Northern Yucatan (Plate 33 b.)

Hernandezii Safford. Central Mexico.

Hindsii Benth. Salvador, Guatemala, Costa Rica, Southwestern Mexico (Plates 34-36).

interjecta Schenck. Singapore (in cultivation).

multiglandulosa Schenck. Eastern Panama (Plates 37-38).

Nelsonii Safford. Guerrero, Southwestern Mexico (Plate 39).

nicoyensis Schenck. Costa Rica (Plate 40).

panamensis Schenck. Panama.

penonomensis Safford. Panama.

rossiana Schenck. Veracruz, Mexico.

sinaloensis Safford. Sinaloa, Mexico.

sphacrocephala Schlecht & Cham. Veracruz, Mexico (Plates 41-43).

tepicana Safford. Tepic, Western Mexico.

turgida Safford. Chiapas, Mexico (Plate 45).

veracruzensis Schenck. Veracruz, Mexico.

yucatanensis Schenck. Yucatan, Honduras.

An even more recent study of these plants by Standley (1922, 1928) has led to a considerable reduction in the number of species. He re-

gards *Hernandezii* and *furella* as synonyms of *cornigera*; *sinaloensis* and *tepicana* as synonyms of *Hindsii*; *chiapensis* as a synonym of *globulifera*; *yucatanensis* of *Collinsii*; and *penonomensis* of *costaricensis*. Probably *cubensis* and *interjecta* are merely garden forms of *cornigera*. Schenck's *multiglandulosa* is certainly a synonym of *melanoceras* Beurling, which was overlooked by Schenck and Safford. With these changes the number of species in the foregoing list would be reduced to 20, and this number is probably too high.¹ Unfortunately Safford did not live to complete his study of the bull-horn Acacias. I have reproduced photographs of these plants which he generously gave me several years ago. Among them is a photograph of a species with remarkably swollen and divergent spines, *A. turgida*, of which he seems to have published no description.

We have as yet no knowledge of the precise species of bull-horn Acacia occurring in Colombia. Jacquinius recorded "*A. cornigera*" from Carthagena and Beurling (1854) long ago very briefly described an *A. melanocroides* taken by Billberg in the same locality². Dr. George Salt informs me that a species grows at Bonda and Tacurina, and Forel described a subsp. *gaigei* of *Pseudomyrma spinicola*, a well known bull-horn ant as taken at Fundacion by Dr. G. M. Gaige. All these localities are in the dry Cienaga region of Northern Colombia. Probably the plant is *A. costaricensis*, the same species that occurs in the dryer parts of Panama and Costa Rica.

Both Schenck and Safford have recognized several groups among the species of bull-horn Acacias. Safford establishes four groups, which are subdivided into sections. A fifth group, the Hebacanthæ, contains several species not cited in the foregoing list, because their thorns are peculiarly flattened and not inhabited by ants. These plants are of peculiar interest as I shall endeavor to show in the sequel.

The beginnings of our ecological knowledge of the bull-horn Acacias have been often recounted. The early botanists often referred to these plants, which were first observed and figured by Hernandez (1651) and Jacquinius (1763). Commelin figured the food-bodies as early as 1697 and Plukenet in 1720. All the other early accounts such as those of Breinius (1739) and Ph. Müller (1752, 1768) and most of the later such as those of Beccari (1884-86), Schimper (1888) and Rettig (1904) are based either on herbarium material or on specimens grown in Euro-

¹Such is also the opinion of Mr. I. M. Johnston, who in a letter calling my attention to the identity of *A. yucatanensis* and *Collinsii*, remarks: "I have investigated the other reductions in this group which Standley has made and am inclined to think that he can be followed in this particular instance with safety. I think that segregation in this group of plants has become too detailed and that the total number of species is well under 20."

²See footnote on p. 101, in connection with *A. melanoceras*.

pean botanical gardens. The type species, *C. cornigera*, was described by Linné in 1737 from material grown in George Clifford's garden. Safford (1923), from whom I take this fact, says that "it was afterwards found growing in its native habitat, near Laguna Verde in the mountains of Vera Cruz, Mexico in 1820 by the botanical explorer Christian Julius William Schiede and was described 10 year later under the name *Acacia spadicigera*, which must be regarded as a synonym of Linnaeus' *Acacia cornigera*". Hernandez's plant which, he says, was called "Hoitmamaxalli", or "forked thorn" by the Aztecs was recognized by Safford and described as a distinct species, *A. hernandezii*.

F. Smith in 1862 published a note on ant-inhabited *Acacia* spines, which has been overlooked in the literature. He described the ant as "*Pseudomyrma modesta*". The specimens were collected by R. W. Stretch in Panama, the *Acacia* being in all probability *A. costaricensis* (= *penonomensis* Saff.), the ant one of the rufotestaceous forms of the species which have been more recently designated as *Ps. belti* Emery or *Ps. spinicola* Emery. The description is however too meager to admit of more precise identification.

The first naturalist to make a careful study of the *Acacias* and their ants in the field was Thomas Belt (1874). His observations were on some of the plants growing about a league from Matagalpa, Nicaragua, and refer almost certainly to the species now known as *Acacia costaricensis* Schenck. As the *locus classicus* of our knowledge of the interrelations of the plant and its ants and of the future hypothesis of myrmecophily, his account is worth quoting in extenso, especially as it contains certain statements that need correction or explanation: "Clambering down the rocks, we reached our horse and mule, and started off again, passing over dry weedy hills. One low tree, very characteristic of the dry savannahs, I have only incidentally mentioned before.¹ It is a species of acacia, belonging to the section *Gummiferæ*, with bipinnate leaves, growing to a height of fifteen or twenty feet. The branches and trunk are covered with strong curved spines, set in pairs, from which it receives the name of the bull's-horn thorn, they having a very strong resemblance to the horns of that quadruped. These thorns are hollow, and are tenanted by ants, that make a small hole for their entrance and exit near one end of the thorn, and also burrow through the partition that separates the two thorns; so that the one entrance serves for both. Here they rear their young, and in

¹This reference is to p. 23, where he is describing the army-ant raids: "Many of the smaller birds build on the branches of the bull's-horn thorn, which is always thickly covered with small stinging honey-eating ants, that would not allow the Ecitons to ascend these trees"

the wet season every one of the thorns is tenanted; and hundreds of ants are to be seen running about, especially over the young leaves. If one of these be touched, or a branch shaken, the little ants (*Pseudomyrma bicolor*, Guér.) swarm out from the hollow thorns, and attack the aggressor with jaws and sting. They sting severely, raising a little white lump that does not disappear in less than twenty-four hours.

“These ants form a most efficient standing army for the plant, which prevents not only the mammalia from browsing on the leaves, but delivers it from the attacks of a much more dangerous enemy—the leaf-cutting ants. For these services the ants are not only securely housed by the plant, but are provided with a bountiful supply of food, and to secure their attendance at the right time and place, the food is so arranged and distributed as to effect that object with wonderful perfection. The leaves are bipinnate. At the base of each pair of leaflets, on the mid-rib, is a crater-formed gland, which, when the leaves are young, secretes a honey-like liquid. Of this the ants are very fond; and they are constantly running about from one gland to another to sip up the honey as it is secreted. But this is not all; there is a still more wonderful provision of more solid food. At the end of each of the small divisions of the compound leaflet there is, when the leaf first unfolds, a little yellow fruit-like body united by a point at the base to the end of the pinnule. Examined through a microscope, this little appendage looks like a golden pear. When the leaf unfolds, the little pears are not quite ripe, and the ants are continually employed going from one to another, examining them. When an ant finds one sufficiently advanced, it bites the small point of attachment; then bending down the fruit-like body, it breaks it off and bears it away in triumph to the nest. All the fruit-like bodies do not ripen at once, but successively, so that the ants are kept about the young leaf for sometime after it unfolds. Thus the young leaf is always guarded by the ants; and no caterpillar or larger animal could attempt to injure them without being attacked by the little warriors. The fruit-like bodies are about one-twelfth of an inch long, and are about one-third of the size of the ants; so that an ant carrying one away is as heavily laden as a man bearing a large bunch of bananas. I think these facts show that the ants are really kept by the Acacia as a standing army, to protect its leaves from the attacks of herbivorous mammals and insects.

“The bull’s-horn thorn does not grow at the mines in the forest, nor are the small ants attending on them found there. They seem specially adapted for the tree, and I have seen them nowhere else.

Besides the *Pseudomyrma*, I found another ant that lives on these acacias; it is a small black species of *Crematogaster*, whose habits appear to be rather different from those of *Pseudomyrma*. It makes the holes of entrance to the thorns near the center of one of each pair, and not near the end, like the *Pseudomyrma*; and it is not so active as that species. It is also rather scarce; but when it does occur, it occupies the whole tree, to the exclusion of the other. The glands on the acacia are also frequented by a small species of wasp (*Polybia occidentalis*). I sowed the seeds of the acacia in my garden, and reared some young plants. Ants of many kinds were numerous; but none of them took to the thorns for shelter, nor the glands and fruit-bodies for food; for, as I have already mentioned, the species that attend on the thorns are not found in the forest. The leaf-cutting ants attacked the young plants, and defoliated them, but I have never seen any of the trees out on the savannahs that are guarded by the *Pseudomyrma* touched by them, and have no doubt the acacia is protected from them by its little warriors. The thorns, when they are first developed, are soft, and filled with a sweetish, pulpy substance; so that the ant, when it makes an entrance into them, finds its new house full of food. It hollows this out, leaving only the hardened shell of the thorn. Strange to say, this treatment seems to favor the development of the thorn, as it increases in size, bulging out towards the base; whilst in my plants that were not touched by the ants, the thorns turned yellow and dried up into dead but persistent prickles. I am not sure, however, that this may not have been due to the habitat of the plant not suiting it.

“These ants seem at first sight to lead the happiest of existences. Protected by their stings, they fear no foe. Habitations full of food are provided for them to commence housekeeping with, and cups of nectar and luscious fruits await them every day. But there is a reverse to the picture. In the dry season of the plains, the acacias cease to grow. No young leaves are produced, and the old glands do not secrete honey. Then want and hunger overtake the ants that have revelled in luxury all the wet season; many of the thorns are depopulated, and only a few ants live through the season of scarcity. As soon, however, as the first rains set in, the trees throw out numerous vigorous shoots, and the ants multiply again with astonishing rapidity.”

The ant mentioned in this description is almost certainly misidentified. *Pseudomyrma bicolor* does occasionally nest in *Acacia* spines but it can hardly be described as a “small” ant, its colonies

are not very populous and it does not attack and sting unless the spines are roughly seized. Belt, I feel confident, actually observed some form of *Ps. belti* or *spinicola*.

Since the appearance of his book, several observers have published accounts on the Acacias or their ants, notably Francis Darwin (1877), Beccari (1884-86), Delpino (1886-89), Schimper (1888), Emery (1892), Wheeler (1912), Wasmann (1915), Schwarz (1917), Calvert (1917), Safford (1923), Menozzi (1927). C. F. Baker, J. Bequaert, P. P. Calvert, A. Dampf, J. Zetek and others have furnished me with valuable specimens and data collected in Costa Rica, Honduras, Panama and Mexico. My own earlier observations (1913) need not be introduced in this connection, but I shall later return to some of them. Attention may here be called to the fact that my identification of *Acacia sphaerocephala* and *cornigera* seems doubtful in the light of the more recent taxonomic revisions of Schenck, Safford and Standley. The same is true of the specific identifications of Acacias by most of the other authors cited above. Thus Menozzi figures the spines of two species of Acacia as belonging to *A. spadicigera* (that is *cornuta*), but they are clearly those of *A. costaricensis* and *Hindsii*.

During 1923 and 1924 I had an opportunity to study two of the three known Panamanian Acacias, namely *A. costaricensis* (= *penonomensis* according to Standley) and *A. melanoceras* and therefore transcribe some of my field notes which I made on the general characters of these plants. The former species I found only on the dryer Pacific side of the Isthmus, in the vicinity of Las Sabanas, near Panama City and less frequently near Red Tank in the Canal Zone. It is probably more abundant in the savannah region further eastward. It is a rather coarse, sturdy bush, rarely more than five or six feet in height and often growing along trails or roadsides through the second growth jungle or in clearings. The grayish green leaves have rather few pairs of pinnae (about 8), with a row of three nectaries on the petiole and one at the junction of each pair of pinnae on the rachis. The spines are terete and moderately long and diverging, green at first, then reddish or chocolate brown. They seem to become hollow rather early, owing to drying up of their pulp-like pith. The flowers are in elongate spikes, the fruit turgid and when ripe splitting along both sides. The young leaflets bear yellow Beltian bodies, which the ants were seen to carry into the spines, while others were feeding at the foliar nectaries. Quite a number of the plants were examined and their ant and other inhabitants noted and collected. I select the following observations as the most interesting:

April 2, 1923. Observed numerous Acacias, from a foot to five feet high, along the Tumba Muerta Road, near Las Sabanas. Nearly all of them were inhabited by *Ps. spinicola* subsp. *atrox*. The spines of two plants, however, contained only *Ps. gracilis*, except a single pair on one of them which was tenanted by a colony of *Brachymyrmex heeri* var. *obscurior* and a few dead spines on the other which were inhabited by *Camponotus (Myrmobrachys) brevis*. The *gracilis* colonies were very flourishing, with much brood and numerous winged sexual forms. The workers had stored quantities of Beltian bodies in the spines, and these also contained a few Coccids and several empty *Microdon* puparia (*vide infra*).

July 2, 1924. Near Punta Paitea. Thirty Acacias growing along the trails near the military camp varied from a foot to 6 ft. in height. The thorns of 24 of them were inhabited exclusively by *Ps. atrox*, and one had no ant-inhabitants though it was quite as flourishing as the other specimens. In two bushes, only a foot high, evidently young suckers growing from the roots of a felled bush, I found the spines inhabited by isolated deilated females with brood, but without any workers, though pupæ of the latter were sometimes present. In one case the leaves of a large bush inhabited by *atrox* were much gnawed (by Chrysomelids). Five of the bushes were tenanted by *Ps. gracilis* exclusively. Owing to its timid disposition, this ant cannot be said to afford much protection to the plants. *Ps. atrox*, however, stings severely, but it does not attack unless the foliage is roughly handled.

A. melanoceras Beurling (1854, p. 123) is clearly the same as the species described as *A. multiglandulosa* by Schenck (1913) and based on material in the Berlin Herbarium collected in 1825 by J. G. Billberg in the vicinity of Porto Bello, Panama. This is evidently part of the material on which Beurling founded his species.¹ Safford has figured and redescribed the species from specimens taken by Pittier at the head of the Gatun Valley, Panama. I include a brief description of

¹I find that Standley (1928) has reached the same conclusion in regard to the identity of this plant.

The following is a transcript of Beurling's original description. "*Acacia melanoceras* n. sp. (*Acacia sphaeracephala* Cham. et Schlect. var. sec. Bentham in herbar. Reg. Acad. scient. holm.); glabra, (ex siccatione?) nigrescens, ramis teretiusculis, verruculis; aculeis stipularibus binis, in cornua basi connata divaricata recte inflata nigricantia nitida demum exrescentibus; petiolis pinnatis, ultra 20-jugis, infra jugum infimum supra excavata—foveolatis (fovea glandulis minutis confertis cupuliformibus repleta); foliis circiter 20 jugis, oblongo-linearibus, sub-falcatis, 2-3 lin. longis, $\frac{1}{2}$ lin. latis, apice (sub lente) minutissime denticulatis; floribus in capitula globosa congestis; capitulis pedicellatis (pedicellis 3-4-nis), in racemos elongatos laterales terminalesque sæpe geminatos (unum longiorem alterum breviorum) dispositis, No. 289. *Mimosa cornigera* (Billberg) (Porto Bello) In silvis ad viam versus Panama." In a foot-note Beurling adds: "Prope Sanct. Lazaro Carthagenæ a Billbergo lecta est species simillima: "No. 139. *Mimosa cornigera* (Billb.), "vix nisi colore haud nigrescente. primarum jugis paucioribus foliolisque duplo longioribus latioribus rectis diversa. (*Acacia melanoceroide*s Beul. mnscrip)t.

the living plant, which is much more beautiful than *A. costaricensis* or indeed than any of the other species I have seen growing in Central America and would be a more ornamental plant under cultivation than *A. cornigera* which is occasionally grown in tropical botanical gardens. I have encountered *A. melanoceras* which, according to Standley, "is a common shrub in the swamps and forests of the Atlantic slope," on only two occasions, once near the head-waters of the Rio Agua Salud (March 6, 1923), and once at Marajal near Colon (July 26, 1924). Both of these localities are on the moist Atlantic side of the Isthmus. On the Rio Agua Salud there were only a few bushes, each about 12 ft. high, growing on the bank of the stream very near the water; at Marajal there were several fine trees, some of which were 18 or 20 feet high, with graceful, smooth, gray-barked trunks, upright and usually growing in a cluster from the same root (Plate 37 a). The leaves are large and moss-green and have many more pinnæ than *penonomensis*. A count of 50 of them showed the number to vary from 12 to 25 pairs, with an average of 20 to 22. The young leaflets bear at their tips rather large, golden-yellow Beltian bodies. The upper surface of the petiole bears numerous (20-30) conical nectaries, and there is a single nectary on the rhachis between each pair of pinnæ. The spines are long (2 to 2½ inches), straight and diverging, smooth and terete, gradually tapering to a short, very acute point. When young they are green, then turn to a brilliant crimson and finally become rich chestnut brown. They are inhabited by a black *Pseudomyrma*, *satanica*, which is larger and even more vicious than *spenicola*. I have not seen the flowers, but the trees at Marajal were bearing clusters of mature green pods, which were straight, cylindrical, with a short terminal snout and measured 3 to 4 inches in length. The pedicel measured about three-quarters of an inch. The pod contained a firm, white, sweetish and edible pulp enveloping a row of black seeds.

According to Beurling, Safford and Standley, the flowers are in globose heads. Safford describes them as 7 mm. long, 6 mm. in diameter, solitary or in pairs, borne in the axils of the bracts on long erect branchlets composed of many nodes.

Dr. J. Bequaert has recently been able to make some observations on *A. Collinsii* (= *yucatanensis* Schenck) and its ants in Honduras. Since little has been written on this *Acacia* I reproduce the notes which he sent me:

"The ant acacia of northern Honduras (*Acacia Collinsii* Schenck) usually forms an erect bush, 6 to 15 feet high, with a main trunk 1 to 3

inches in diameter and covered with many slender, short side branches. The size and shape of the spines vary considerably on the same bush, those of the main trunk, at the foot of the branches, being much longer and twisted in horn-fashion. Young spines are either green or bright red and both types occur on the same bush. They soon become naturally hollow, without the help of the ants, which merely cut the opening below the tip and remove the dried pith.

"These acacias are particularly abundant near the coast, a short distance (one or two miles) from the shore, beyond the lagoons and mangrove swamps. They grow in dry, sandy, slightly undulating areas which appear to be ancient, fixed sand dunes and are covered with short grass and scattering shrubs and low trees. Many specimens may be seen along the main line of the Trujillo Railroad, about 15 to 25 kilometers from Puerto Castilla (near the stops known as Los Cuartos and El Canal). Only two small specimens were seen at Puerto Castilla, on the sandy levee between the sea and the mangrove swamp. At El Canal a number of them were also found in a rather dense and more humid wooded gallery along the banks of a stream. These specimens were larger and more tree-like than usual, reaching 20 to 25 feet in height, widely branched at the top, but with the main trunk, 4 to 5 inches thick, almost free of side branches, although covered with heavy and strongly twisted spines. Further inland I only observed the plant on one occasion, namely a single, bush-like individual which grew on the high, sandy, and densely wooded bank of the Rio Aguan, near Prieta, about 90 kilometers from Puerto Castilla along the railroad to Olanchita.

"The black ant (*Pseudomyrma belti*) I found only in two plants, which grew near the sea shore at Puerto Castilla. The brownish form (*Ps. belti* subsp. *bequaerti*) is the common inhabitant of all the acacias near Los Cuartos and El Canal, where it occurs as well in the bushy plants of the dry, open areas as in the more tree-like form of the forest gallery. The subsp. *bequaerti* was also found in the single bush near Prieta. I observed how these ants actively collect the Beltian bodies at the tips of the leaflets and also visit the nectary near the foot of the petiole. At the time of my observations (March), most of the plants had flower buds, but only very few were actually blooming. After much search two or three seed-pods could still be obtained. Many of the bushes had very small, young leaves, just emerged from the bud, with food-bodies still attached. These food-bodies were either yellowish white or more rarely blackish. Both forms of *Pseudomyrma* are very aggressive; their sting is painful, but the pain soon abates."

The following list comprises the various ants recorded as living in the spines or on the foliage of the bull-horn Acacias:

(1) *Pseudomyrma belti* Emery. In spines of *Acacia* sp. (probably *costaricensis*). Costa Rica (A. Alfaro; P. P. Calvert). In thorns of *A. Collinsii*; Honduras (J. Bequaert).

(2) *Pseudomyrma belti* subsp. *bequaerti* Wheeler. In spines of *A. Collinsii*. Honduras (J. Bequaert).

(3) *Pseudomyrma belti* subsp. *fellosa* Wheeler. In spines of *A. costaricensis*. Nicaragua (C. F. Baker; W. Fluck).

(4) *Pseudomyrma belti* subsp. *fulvescens* Emery. In spines of *A. Hindsii* and *Acacia* sp. Guatemala (Wheeler); *Acacia* sp. Guatemala (Schwarz and Barber); British Honduras (J. D. Johnston; C. F. Baker); Mexico (E. Palmer, D. L. Crawford, Jourdain, A. Petrunkevitch, F. Knab); In spines of *A. sphaerocephala*., Mexico (Rangel, G. N. Collins, L. G. Culvas, J. M. Cuaron); *A. Collinsii*, Mexico (G. N. Collins); *A. cornigera* Mexico (G. N. Collins); *A. Hindsii* Mexico (G. N. Collins).

(5) *Pseudomyrma belti* subsp. *obnubila* Menozzi. In spines of *A. "spadicigera"* (probably *costaricensis* or *Hindsii*) Costa Rica (H. Schmidt).

(6) *Pseudomyrma belti* subsp. *saffordi* Wheeler. In spines of *A. Collinsii*. Mexico. (G. N. Collins). *Acacia* sp. Guatemala (Wheeler).

(7) *Pseudomyrma belti* subsp. *renefica* Wheeler. In spines of *A. Hindsii*. Mexico (J. H. Batty, C. H. T. Townsend); *Acacia* sp. Mexico (C. F. Baker).

(8) *Pseudomyrma belti* subsp. *resana* Wheeler. In spines of *Acacia* sp. Mexico (F. Knab).

(9) *Pseudomyrma belti* subsp. *wasmanni* Wheeler. In spines of *A. sphaerocephala* (W. Brakmann).

(10) *Pseudomyrma brunnea* F. Smith. In spines of *A. "spadicigera"* (probably *costaricensis* or *Hindsii*) Costa Rica (H. Schmitt).

(11) *Pseudomyrma flavidula* F. Smith. In spines of *A. "spadicigera"* (probably *costaricensis* or *Hindsii*). Costa Rica (H. Schmitt).

(12) *Pseudomyrma gracilis* Fabr. In spines of *A. Hindsii* and *Acacia* sp. Guatemala (Wheeler); *A. costaricensis*. Panama (Wheeler) *A. sphaerocephala*. Mexico (Schenck).

(13) *Pseudomyrma gracilis* subsp. *bicolor* Guer. In spines of *A. costaricensis*, Panama (Wheeler).

(14) *Pseudomyrma gracilis* subsp. *mexicana* Roger. In spines of *Acacia* sp. Costa Rica (A. Alfaro).

(15) *Pseudomyrma kuenckeli* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro); *A. "spadicigera"* (probably *costaricensis* or *Hindsii*) Costa Rica (H. Schmitt).

(16) *Pseudomyrma nigrocincta* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro; P. P. Calvert).

(17) *Pseudomyrma nigropilosa* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro; P. P. Calvert).

(18) *Pseudomyrma peltata* Menozzi. In spines of *A. "spadicigera"* (probably *costaricensis* or *Hindsii*) Costa Rica (H. Schmitt).

(19) *Pseudomyrma satanica* Wheeler. In spines of *A. melanoceras*. Panama (Wheeler).

(20) *Pseudomyrma sericea* Mayr. var. *acaciarum* Wheeler. In spines of *A. costaricensis*. Panama (Wheeler).

(21) *Pseudomyrma spinicola* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro; P. Biolley).

(22) *Pseudomyrma spinicola* subsp. *atrox* Forel. In spines of *A. costaricensis*. Panama (E. D. Christophersen; Wheeler).

(23) *Pseudomyrma spinicola* subsp. *modesta* F. Sm. In spines of *Acacia costaricensis*. Panama (R. W. Stretch).

(24) *Pseudomyrma spinicola* subsp. *convarians* Forel. In spines of *Acacia* sp. Guatemala (Peper).

(25) *Pseudomyrma spinicola* subsp. *gaigei* Forel. In *Acacia* spines (very probably), Colombia (F. M. Gaige).

(26) *Pseudomyrma spinicola* subsp. *infernalis* Wheeler. In spines of *A. costaricensis*. Panama (Wheeler).

(27) *Pseudomyrma spinicola* subsp. *scelerosa* Wheeler. In spines of *A. costaricensis*. Nicaragua (C. F. Baker).

(28) *Pseudomyrma subtilissima* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro).

(29) *Solenopsis zetekii* Wheeler. Nesting in dead twig caught among spines on trunk of *A. melanoceras*. Panama (Wheeler).

(30) *Solenopsis* sp. Nesting in flower peduncles of *Acacia* sp. Guatemala (Wheeler).

(31) *Crematogaster acuta* Fabr. Running on foliage of *A. costaricensis*. Panama (Wheeler).

(32) *Crematogaster (Orthocrema) atra* Mayr. In spines of *A. veracruzensis*. Mexico (A. Dampf).

(33) *Crematogaster (Orthocrema) brevispinosa* Mayr. In spines of *Acacia* sp. Costa Rica (A. Alfaro). *A. costaricensis*. Nicaragua (E. Chamorro); *A. veracruzensis*. Mexico (Schenck).

(34) *Crematogaster (Orthocrema) brevispinosa* var. *minutior* Forel.

Nesting in accumulations of carton around spines of *A. cornigera* in botanical garden, Port of Spain, Trinidad (Wheeler).

(35) *Crematogaster* (*Orthocrema*) *brevispinosa* subsp. *timulifera* Forel. In spines of *A. Hindsii*. Guatemala (Wheeler).

(36) *Crematogaster* (*Orthocrema*) *corvina* Mayr. In spines of *A. vera-cruzensis*. Mexico (A. Dampf).

(37) *Crematogaster* (*Orthocrema*) *curvispinosa* Mayr var. *panamana* Wheeler. In spines of *A. costaricensis*. Panama (Wheeler).

(38) *Leptothorax* (*Goniothorax*) *echinatinodis* Forel subsp. *cordincola* Wheeler. In spines of *A. costaricensis*. Panama (Wheeler).

(39) *Leptothorax* (*Goniothorax*) *echinatinodis* subsp. *schmidti* Menozzi. In spines of *A. "spadicigera"* (probably *costaricensis* or *Hindsii*). Costa Rica (H. Schmidt).

(40) *Cryptocerus* (*Paracryptocerus*) *minutus* Fabr. In spines of *A. costaricensis*. Panama (Wheeler); *A. "spadicigera"* (probably *costaricensis* or *Hindsii*). Costa Rica (H. Schmidt).

(41) *Cryptocerus* (*Cyathocephalus*) *pallens* Klug. In spines of *A. Collinsii*. Mexico (G. N. Collins).

(42) *Cryptocerus* (*Cyathocephalus*) *pallens* var. *discocephalus* F. Smith. In spines of *Acacia* sp. Costa Rica (A. Alfaro).

(43) *Dolichoderus* (*Hypoclinea*) *championi* Forel var. In spines of *A. Hindsii*. Guatemala (Wheeler).

(44) *Dolichoderus* (*Hypoclinea*) *diversus* Emery var. *ficoides* n. v. In spines of *A. cornigera*. Mexico (G. N. Collins).

(45) *Azteca rector* Forel. In spines of *A. Hindsii*. Guatemala (Wheeler).

(46) *Brachymyrmex heeri* Forel var. *obscurior* Forel. In spines of *A. costaricensis*. Panama (Wheeler).

(47) *Camponotus* (*Myrmobrachys*) *planatus* Roger var. *acacia* Emery. In spines of *A. Hindsii*. Guatemala (Wheeler); *Acacia* sp. Costa Rica (A. Alfaro); *A. "spadicigera"* (probably *costaricensis* or *Hindsii*). Costa Rica (H. Schmidt).

(48) *Camponotus* (*Myrmobrachys*) *brevis* Forel. In spines of *A. costaricensis*. Panama (Wheeler).

(49) *Camponotus* (*Myrmobrachys*) *striatus* F. Smith. In spines of *Acacia* sp. Costa Rica (A. Alfaro); *A. costaricensis*. Panama (Wheeler).

(50) *Camponotus* (*Myrmocladococcus*) *mucronatus* Emery subsp. *santschii* Forel. In spines of *A. Hindsii*. Guatemala (Wheeler).

(51) *Camponotus* (*Myrmocladococcus*) *rectangularis* Emery. In spines of *Acacia* sp. Costa Rica (A. Alfaro).

(52) *Camponotus* (*Colobopsis*) sp. In spines of *Acacia* sp. Costa Rica (A. Alfaro).

(53) *Paratrechina* (*Nylanderia*) *longicornis* Latr. Running on *Acacia* sp. Costa Rica (A. Alfaro).¹

Of the various ants listed only three species can be regarded as obligate or definitely associated with the bull-horn Acacias, namely *Ps. belti*, *spinicola* and *satanica*; all the others are species or sub-species and varieties of species which more frequently nest in dead twigs of a great number of different plants. Emery did, indeed, describe *Ps. belti* from specimens taken from *Cordia alliodora* domatia, but this must have been a very exceptional occurrence, or more probably due to an erroneous label.

Two of the forms listed, *Ps. gracilis* and *Camponotus planatus* var. *acaciæ*, are of special interest. The former, though a very frequent tenant of dead twigs and *Cordia* domatia in regions where there are no Acacias, nevertheless exhibits a strong proclivity not only to inhabit the spines of these plants, wherever they are available, but also to perforate them at the same point, to visit the foliar nectaries and to collect the food-bodies. In other words, this ant has acquired all the peculiarities of behavior of the three obligate *Acacia* tenants, *Ps. belti*, *spinicola* and *satanica*. This is perhaps true also of some or all of the other *Pseudomyrmas* in the list. In my previous paper (1913) I described the founding of *gracilis* colonies in the thorns of young Acacias at Quirigua, Guatemala. In that region I found a locality in the banana plantations where *gracilis* was the only ant inhabiting the Acacias. As previously stated, it is also not infrequently the sole tenant of *A. costaricensis* in Panama. Of course, it is quite possible that the *gracilis* associated with Acacias represents a distinct physiological race derived and as yet morphologically indistinguishable from the form occurring more generally in dead twigs. At any rate, the case is interesting because it shows such a perfect adaptation to the bull-horn Acacias by particular colonies of a species which as a rule exhibits no adaptations to ant plants beyond those exhibited by most other species of *Pseudomyrma*.

Camponotus planatus var. *acaciæ*, as Alfaro (Emery, 1892) and I (1913) have found, frequently inhabits the same Acacias as *Ps. belti* and *gracilis* in Costa Rica and Guatemala. I described the mutual relations of these ants as follows: "That the *Camponotus* does not, as Emery supposed, merely take possession of thorns excavated and abandoned by the *Pseudomyrma*, was proved on one occasion when I

¹Schenk (1914) records *Pseudomyrma arboris-sanctor* as occurring in the spines of *A. sphacrocephala*, *spadicigera* and *veracruzensis* in Mexico. The validity of this identification, which was made by Reichenperger, may be seriously doubted. In all probability the specimens belonged to some reddish subspecies of *Ps. belti* or *spinicola*.

found a small group of *Camponotus* workers busily engaged in perforating a green thorn. It is probable, therefore, that the *Camponotus* queens, after their nuptial flight, seek out the acacias and enter their young thorns even when the trees are already inhabited by the *Pseudomyrma*, and that the *Camponotus* workers continue this work side by side with the *Pseudomyrmas*, both species competing for and taking possession of the thorns as fast as they attain the proper size and maturity. It is certainly extraordinary that *C. planatus*, which throughout tropical America so constantly lives in hollow twigs, should be able in widely separated localities to utilize the acacia thorns as perfectly and in precisely the same manner as the regular *Pseudomyrmas*. That the *Camponotus* is, if anything, even more adroit in its use of the extrafloral nectaries becomes apparent when one follows the ant as it moves over the leaves, for it begins with the nectary at the base of the petiole and carefully visits each in turn, whereas the foraging *Pseudomyrmas* are much more desultory and less business-like. I have not seen the *Camponotus* collecting the Beltian bodies, but I doubt not that they make quite as good use of them as of the nectar. The behavior of the two species of ants towards each other is peculiar. They seem never to quarrel, and if not too close together, pass one another on the twigs and leaves with an air of complete indifference. But when two of them happen to meet squarely face to face, each starts back suddenly and, curiously enough, the *Pseudomyrma* always recoils more vigorously than the *Camponotus*. There is something ludicrous in this behaviour because both ants are of about the same bulk, and the *Pseudomyrma* is really the more powerful and possesses a formidable sting, whereas the *Camponotus* is much less pugnacious and can defend itself only with its rather feeble mandibles and formic acid battery. But it smells rather strongly of formic acid, and I believe that this produces the more decided reaction on the part of the *Pseudomyrma*."

I regarded the mutual relations of the *Camponotus* and *Pseudomyrma* as a case of parabiosis, but Wasmann (1915, p. 128) has objected to my use of this word, because the two species of ants do not inhabit different parts of the same nest as in the case of *Crematogaster parabiatica* and *Dolichoderus (Hypoclinca) parabioticus*, to which Forel first applied the term, "but different nests (spines) on the same tree or branch." I believe, however, that all the pairs of spines that are converted into domatia on an *Acacia* may be really regarded as so many chambers of a single nest, when the ant-colony is well-developed. It is only the incipient colony that is confined to a single pair of spines.

One might, of course, invent some other term for the relations of mutual tolerance exhibited by the *Camponotus* and *Pseudomyrma*, but this seems hardly necessary. It remains to be seen whether the concept of parabiosis may be extended to include all the other cases in which two or more species of ants inhabit the domatia of the same plant (*Acacia*, *Cordia*, *Triplaris*, *Cecropia*, etc.)

Besides the ants enumerated above the various *Acacias* also harbor a considerable number of miscellaneous insects and other organisms, which are briefly noticed in the following list:

(1) *Aves*. Birds, probably of several species, not infrequently build their nests in *Acacia* bushes or trees inhabited by the stinging *Pseudomyrmas*. This was observed by Belt in Nicaragua (*A. costaricensis*) and by myself both in Guatemala (*A. Hindsii*) and Panama (*A. costaricensis*).

Hymenoptera. Dr. E. A. Schwarz (1917) bred several species parasitic on beetle larvæ from an *Acacia* trunk from Tampico, Mexico, but has cited none of them by name. I have noticed most of the following forms:

(2) Chalcidids. Bred from spider's eggs attached to twigs of *A. costaricensis*. Panama (Wheeler).

(3) Braconids. Swept from foliage of *A. costaricensis*. Panama (Wheeler).

(4) *Solitary wasp*. Nest made in an old thorn of *A. costaricensis*. Panama (Wheeler).

(5) *Polybia* sp. Nests constructed on branches of *A. Hindsii* in Guatemala (Wheeler).

(6) *Polybia* sp. Nest on *A. costaricensis*, Panama (J. Zetek).

(7) *Polybia occidentalis*. Cited by Belt as visiting the foliar nectaries of *Acacia (costaricensis)* in Nicaragua.

(8) *Halictus* sp. A small bee visiting the flowers of *A. Hindsii* in Guatemala (Wheeler).

(9) *Trigona* sp. A black stingless bee visiting the flowers of *A. costaricensis*, Panama (Wheeler).

Lepidoptera. The caterpillars of several species of moths feed on various parts of the *Acacias* notwithstanding the presence of the stinging *Pseudomyrmas*:

(10) *Adelocephala xanthochroia* Schauss. According to Carlos Hoffmann (*teste* A. Dampf) the caterpillars of this moth feed on the pulp of the seed-pods of *Acacias* at Misanhtla, Vera Cruz.

(11) Small moth caterpillars, which devour the foliage and leaflets and attach their cocoons to the twigs of *A. costaricensis* at Punta

Paitea, Panama. Several of them were observed on single plants inhabited by *Pseudomyrma atrox*.

(12) A second small moth caterpillar was found cocooning on the stem of *A. costaricensis*.

(13) A third was living in old spines of the same species of Acacia.

(14) A Psychid larva had its case attached to a twig of the same plant.

Diptera. I have observed the larvæ or pupæ of five species of this order, without being able to secure the adults:

(15) In some localities in Guatemala (Esquintla, Patulul) the upper surfaces of the leaflets of *A. "cornigera"* bore beautiful little spherical galls, 5 to 6 mm. in diameter, single or in clusters and resembling minute strawberries, as they were bright red and uniformly covered with papillæ (Plate 46 a). Each of these galls contained a Dipteran (Cecidomyid) larva and had a preformed rhapshe along which it dehisces when brown and mature, and permits the adult fly to escape. *Pseudomyrma fulvescens* is very fond of visiting these galls when young and succulent, and was often seen gnawing away the covering of papillæ, but not eating in far enough to injure the enclosed larva.

(16) A second gall of much larger size and woody texture was occasionally found on the flower-stems of *A. Hindsii* in Guatemala, but as I could secure only old and dried specimens, I am unable to make any statement in regard to the insect.

(17) A very peculiar compound woody gall (Plate 46 b) was frequently found on the twigs and especially between and around the spines of *A. costaricensis* near Panama City. It consists of a number of tubular structures, each with an opening at its summit and containing in its more enlarged basal portion a Cecidomyid larva which I did not succeed in rearing.

(18) Several of the thorns of *A. costaricensis* at Las Labanas, Panama and Red Tank, C. Z., inhabited by *Ps. gracilis*, contained empty, rather smooth, brown puparia (about 7 mm. long and 3 mm. broad) of a small *Microdon* sp. Since the openings of the thorns were only large enough to permit the slender ants to pass in or out I am unable to account for the egress of the fly after its emergence. The flies had evidently emerged normally since the anterior dorsal portion of the puparia had been broken open at the usual transverse suture. Perhaps the insects had been devoured by the ants. From the fact, however, that these puparia were encountered in more than a dozen pairs of spines on different bushes in two localities, we must infer, I believe, that the *Microdon* is a regular synœkete in the nests of *Ps. gracilis*.

(19) Among some spider's eggs in a cocoon attached to the twigs of *A. costaricensis* I found a Muscid puparium, probably that of a Tachinid.

Coleoptera. Several beetles have been recorded from Acacias, but only one has been specifically identified:

(20) *Agrilus* sp. A small bronze-green species was common on the foliage of *A. costaricensis* at Punta Paitea, Panama. Several pairs were taken *in copula*.

(21) *Agrilus* sp. A larger bronze-green form, not uncommon on the same plant in the same locality.

(22) *Agrilus* sp. Bred by Dr. E. A. Schwarz (1917) from the trunk of *Acacia* sp. from Tampico.

(23-29) Dr. Schwarz also bred seven species of Cerambycidae (one Cerambycine and six Lamiids) from the same trunk.

(30-31) At Las Sabanas, Panama two unidentified species of Chrysomelidae were found on the foliage of *A. costaricensis*. The injury to the foliage, occasionally noticed, probably results from the attacks of these or allied species.

(32) Curculionid. The larva, apparently of some weevil, was found living in the spines of *A. costaricensis* near Las Labanas, Panama, and feeding on the pith.

(33) Curculionid sp. A small black species taken on the foliage of the same Acacia.

(34) *Bruchus cinerifer* Fabr. Cited by Schenck (1914) as boring into the pods and feeding on the seeds of *A. sphaerocephala* in the Mexican *tierra calicute*.

(35) *Bruchus* sp. Possibly the same as the preceding observed by Carlos Hoffmann (*teste* A. Dampf) as occurring in the seeds of *Acacia* sp. at Misanthla, Vera Cruz. Through the holes made by this beetle the caterpillars of *Adeloccephala xanthochroia* enter the pods and feed on the pulp.

(36) *Bruchus* sp. Dr. Schwarz observed this species, which may be the same as *B. cinerifer*, infesting the flowers of an *Acacia* sp. at Tampico and ovipositing and developing in its pods "without being molested by the ants in any of its stages."

(37) *Bothrioderes* sp. A Colydiid bred by Dr. Schwarz from the trunk of *Acacia* sp. from Tampico.

(38) *Lathropus* sp. A Cucujid from the same trunk.

(39) *Trogosita* sp. An Ostomid from the same trunk.

(40) *Clerus* sp. Bred from the same trunk by Dr. Schwarz.

Orthoptera (41) *Blattid* sp. A large compressed ootheca of some

roach was taken from the cavity of a large dead spine of *A. melanoceras* at Marajal, C. Z. (Wheeler).

Isoptera. (42) *Eutermes* sp. An ellipsoidal nest of this termite was situated among the branches of an *A. melanoceras* tree at Marajal and had galleries descending from it along the trunk. When the latter were broken open a violent battle ensued between the termites and the ants (*Ps. satanica*).

(43) *Eutermes* sp. Abandoned tunnels of another termite were found on the trunk and branches of an *A. costaricensis* bush inhabited by *Ps. gracilis*.

Heteroptera (44) The only member of this order taken was a large Pentatomid, which was crawling on the foliage of *A. costaricensis* at Las Sabanas, Panama.

Homoptera. Coccidæ. Of this family I have found only three species associated with Acacias:

(45) *Coccus elongatus* (Sign) (H. Morrison det.). A few specimens taken in spines of *A. costaricensis* inhabited by *Ps. gracilis* at Pueblo Nuevo, Panama.

(46) *Pseudococcus texensis* Tinsley (H. Morrison det.). Numerous specimens taken in spines of *A. veracruzensis*, collected by Dr. A. Dampf at Vera Cruz, and inhabited by *Crematogaster atra* and *corvina*.

(47) *Pseudococcus* sp. Inhabiting the cavities of flower-peduncles of *A. Hindsii* which contained colonies of *Solenopsis* sp. Guatemala (Wheeler).

Myriopoda. Two species of this class were found inhabiting old abandoned spines of *A. costaricensis* near Panama City, namely:

(48) *Polyxenus* sp. Forming small colonies.

(49) A *Geophilus*-like form, living singly.

Araneina. At least four species of spiders were found inhabiting webs and concealing their egg-cocoons among the leaves and branches of *A. costaricensis*, and at least six in similar situations on *A. melanoceras*. The following four have been identified by Mr. N. Banks.

(50) *Eustala fuscovittata* Keys. On foliage of *A. costaricensis* at Punto Paitea, Panama.

(51) *Phyalc* sp. On the same species of Acacia in the same locality.

(52) *Acrosoma obtusispina* Keys. A large species inhabiting orb-webs among the branches of *A. melanoceras* at Marajal.

(53) *Nephila clavipes* L. Another large spider inhabiting similar webs on the same species of Acacia.

Acarina (54) Numerous mites found attached to the workers of

Camponotus brevis Forel inhabiting spines of *A. costaricensis* at Corazel, C. Z. Panama.

Although the 54 different organisms, other than ants, above listed as associated with the bullhorn Acacias have been less satisfactorily studied and identified than those recorded for *Cordia alliodora* and *Triplaris*, and undoubtedly represent only a small fragment of the actual Acacia biocœnoses, they are sufficient, nevertheless, to show, first, that these plants have plenty of natural enemies and are in this respect like other nonmyrmecophytic trees and shrubs in the tropics, and second, that the obligate ant-tenants, though more virulent than those of *Cordia alliodora* and the Cecropias, are nearly or quite as tolerant of alien ants and other insects on the same plants. I therefore reiterate my statement of 1913 that the relations existing between the Acacias and the obligate Pseudomyrmas are not properly those of symbiosis, in which the plants have adapted themselves to the ants, but those of host and parasite, in which the adaptations are solely on the part of the ants.

This contention, in its essential features, was rejected by Wasmann in his paper of 1915. Like other authors he has been greatly impressed by the coexistence of the capacious spines, abundant foliar nectaries and Beltian bodies in these Acacias, and though he is quite unable to account for these various structures, nevertheless regards the plants as having perfected them for the purpose of securing the protection of the ants. This is clear from the following quotation (p. 130): "If, therefore, we weigh the "pros and cons" of the so-called myrmecophily of the ant-acacias, I believe that we arrive at the same conclusion, which I had already reached in my previous paper (1915, p. 303, 315, 321), namely a position midway between the two extremes of an over-estimation and an underestimation of the myrmecophilous adaptations. The initiative that resulted in the occupation of the acacias by certain acacia ants of the genus *Pseudomyrma*, had its source in the latter. They selected these trees gradually with ever increasing regularity for purposes of establishing their colonies on account of the advantages accruing to their species. This specialization of the nesting instinct is, of course, to be conceived without any such anthropomorphism as that implied in supposing the ants to be aware of these advantages through intelligent reflection. Both the favorable domiciliary conditions, which the spines of these trees afford the ants, and the favorable nutritive conditions provided by the extrafloral nectaries, are therefore not a product of "natural selection" due to "adaptation" to the ants under consideration. The prerequisite physiological condi-

tions for these actual adaptations must have been developed beforehand by the laws of growth of the host-plants. But after certain species of *Pseudomyrma* had adapted themselves to these preconditions and had taken up their regular abode on the respective Acacias, it was possible, through the protection thus incidentally acquired by the host plant—even supposing this protection to be by no means considerable—that *the further development* of the extrafloral nectaries and especially of the Beltian corpuscles *was favored* to a degree which the same peculiarities would not have attained without such myrmecophily. In this sense we may still speak of a true “symbiosis” between the ant-acacias and the acacia-ants, which is now more than a mere parasitism, though it arose *originally* from a unilateral parasitism. Somewhat similar conditions are found in the symphily of many myrmecophilous and termitophilous Coleoptera, which is also more than mere parasitism, though it had its beginning in unilateral parasitism.”

Of course, there is no way of either confirming or refuting this piece of pure phylogenetic speculation, which is so much like others in which Wasmann has indulged from time to time in regard to tropical organisms that he has never observed in a living condition. Obviously the ants could not have settled in the Acacias till their spines had developed to their present dimensions, and it was these organs, so easily convertible into domatia, that induced the ants to take up their abode on the plants. This is clearly indicated by two species of *Acacia*, among the many very diverse nonmyrmecophytic forms that grow in the same general geographical region, namely the swordthorns, *A. macracantha* Humb. and Bonpl. of Mexico, *hirtipes* Safford of Guatemala, *Standleyi* Safford and *gladiata* Safford of Mexico and the spoonthorns, *cochliacantha* Humb. and Bonpl. of Ecuador and *cymbispina* Sprague and Riley of Mexico and Central America. All these plants have large spines which are too much flattened to be used as domatia by the ants. I have not seen specimens of the various species but the following data collated from the literature indicate that food-bodies and nectaries like those of the bullhorn Acacias are present in at least some of them:

A. macracantha, according to Standley (1922) has compressed spines 2.5 to 5 mm. long, but he makes no mention of food-bodies or nectaries.

In *A. Standleyi* (Plate 44), according to Safford (1914), the young leaflets are tipped with food-bodies and the main leaf-rhachis bears “a conspicuous annular nectar-gland at the base of each pair of pinnæ and usually one on the petiole, just below the lowest of these”. The spines are “3-3.5 cm. long, 6-8 mm. broad at the base, very widely

divergent, the pair separated by a thickened ridge (the persistent base of the petiole) adnate to the branch."

A. hirtipes. Nothing is said in Safford's description (1914) about food-bodies but the foliar nectaries are "dark purplish, circular, bowl-shaped, with a thick annular margin, one in the smaller leaves of the short branchlets, one borne at the base of each pair of pinnæ and an additional one on the petiole; on the larger leaves of the longer branches similar glands borne at the base of each pair of pinnæ of the upper half of the leaf, but none in the lower half except a solitary gland on the petiole." The large stipular spines are "broadly v-shaped, cinereous, puberulent, except at the points, 3-4.2 cm. long, 10 mm. broad at the cuneate base, the latter flattened but not adnate to the branch as in *A. Standleyi*."

A. gladiata is considered by Standley as possibly merely a form of *Standleyi*. The leaf-rhachis usually bears a nectary at its base and often one just below each pair of pinnæ. The leaflets are "often mucronate or tipped with a waxy apical body, as in the true myrmecophilous acacias." "Large spines very long and divergent, usually flattened and sword-like, linear-lanceolate in outline, somewhat constricted at the base, resembling certain forms of the spines of *Acacia cochliacantha* H. & B. but connate instead of separate at the base and never split or inflated, gradually narrowed toward the apex to an acute point, 35 to 52 mm. long, 5 to 8 mm. broad, glabrous, reddish or wine-colored when young, at length brown or tan-colored."

A. cymbispina. This form is usually identified as *A. cochliacantha*, originally described from Ecuador, but is specifically distinct, according to Sprague and Riley. The spines are widely divergent and compressed, greatly enlarged and spoon-shaped, that is strongly convex on one side and concave on the other. This *Acacia* was observed by the Calverts (1917), who found it growing along the Bananita River, in Costa Rica. According to their account it resembled the bull's horn *Acacia*, "but had a less woody stem or trunk and paired thorns not curved nor inclined toward each other but merely at right angles to the stem bearing them. There were no little fruit-like bodies at the tips of the young leaflets, but along the midrib of each leaf was a row of urn-shaped glands, one at the base of the twenty-seven pairs of pinnæ." There were no ants on this *Acacia*, "but it looked so much like bull's horn thorn as to suggest that it might be the starting point for the development of the latter."

These cases show that the mere presence of well-developed foliar nectaries, or of nectaries and food-bodies, does not suffice to induce the

ants to make their abode in the Acacias. The same conclusion is also reached from observation of many other plants, such as the species of *Cassia*, *Stillingia*, *Populus*, etc. which may attract ants or other insects to their nectaries, but cannot retain them as permanent occupants, because they possess no structures that can be converted into suitable domatia. It would be hazardous to maintain that *A. cochleacantha* and *gladiata* are either persisting, older, phylogenetic stages in the evolution of the bull-horn Acacias or more recent, involutory derivatives of the same. More probably the two species and their allies represent independent developments from some stock common to that of the bull-horns; and their survival as well as that of many more delicate species of *Acacia* in Mexico and Central America without ant-protection, is not calculated to strengthen the contention of Wasmann and other believers in an adaptive myrmecophily on the part of the plants. Additional evidence of the absence of such adaptation is also available from the study of some of the African Acacias and of one South American species, *A. cavienia*, to which we may now turn.

(B) ACACIA CAVENIA Hooker and Arnott

We possess three accounts of this Paraguayan *Acacia*, which really bears only a certain superficial resemblance to the bull-horns in the great enlargement of its spines and their occupation by ants.

Fiebrig (1909) describes the plant as a shrub covering considerable areas in the Chaco of Paraguay, near the Bolivian boundary, and restricted to loess-like alluvial soil which is occasionally inundated and is not inhabited by Attine ants. Though the stems are so slender as rarely to exceed a centimeter in diameter, some of their stipular spines may become very large (90 mm. long and 8 mm. broad at the base), "but only at certain seasons, apparently as a result of abundant atmospheric humidity." Fiebrig found that their pith is devoured by a Tineid caterpillar, which before pupation gnaws an exit-hole near the tip of one of the spines of a pair. This caterpillar may hollow out not only a single spine but also its fellow and a portion of the adjacent stem. At the time of the moth's emergence the head of the pupa is pushed out of the opening. Later the spines are invaded by ants (*Pseudomyrma fiebrigi* Forel), which utilize the moth's exit-hole as an entrance. The young leaves of the *Acacia* bear no Beltian bodies. No mention is made of the presence of extra-floral nectaries.

Previously, in 1896, Emery had described a number of ants taken by Dr. J. Bohls from the large spines of a Paraguayan *Acacia*, which

is almost certainly *A. cavenia*, since the specimens were collected in a locality, San Salvador, very near the Chaco in which Fiebrig made his observations. According to Emery, the species of *Pseudomyrma* made delicate galleries in the woody substance of the spines, whereas the species of *Cryptocerus* hollowed them out completely. "The opening of the *Pseudomyrma* nests was made near the tip of the spine, the openings of the other species were at a variable distance from the base and sometimes there were several."

More recently *A. cavenia* has been studied again by Chodat and Vischer (1920) at Trinidad, Concepcion, etc. in the periodically inundated country along the Rio Paraguay. Their interpretation of the enlargement of the spines differs from that of Fiebrig, as will be seen from the following quotation and an examination of their figures 324 and 325: "We have also found spines of considerable dimensions without any insect in them. One might ask, therefore, whether Fiebrig's theory is really sufficient to account for the production, or origin of these hypertrophied stipules. We suspected the existence of a morphogenic stimulus produced by the sting of an insect in some other part of the stem, and we therefore sought for such organisms in the pith of the branches. Examination confirmed our suspicions. Below the insertion of the spines (stipules) we found the larva or pupa of a Chalcidid quite analagous to the one that produces the myrmecodomatia, or galls described for the *Cordias*. How does the infection occur. How does the female oviposit? The fissures indicate that the animal may penetrate the tissues either in the region of one of the buds or of its fellow of the opposite side. But this point should be observed at the time and place of its occurrence. The infection may be deep in the pith or more superficial (see Fig. 324). It will be noticed in Fig. 324 that the larva, after devouring a portion of the pith of the stem, moves towards the base of the spine, which becomes hypertrophied. Sometimes, as in the *Cordias*, two or several (?) larvæ may be found. Moreover, the canal be means of which the larva penetrates the thorn, may be transverse and bring about a communication between the two stipules (see Fig. 325). Fiebrig had previously indicated the presence of these continuous chambers or galleries but he assigned them an inverse origin. We are confirmed in our theory by the fact that Fiebrig himself noticed that the stipules may become hypertrophied without exhibiting any localized attack in their pith or base, that is, without any internal or external indications. For us, and we have made many concordant and no discordant observations, there is always an infection of the stem before the stipules are hollowed out. We must suppose, therefore, that

the excitation starts from the stem and is transmitted either by the plant tissue or by substances secreted by the animal, or, what is even more probable and agrees with most of our observations, by the direct action of the progressing larva. It may happen also, no doubt, that the infection is caused directly at the base of the leaf, since on one occasion we found a stipular spine that contained a pupa and a kind of cocoon, without any communication existing between the spine and the gallery in the stem."

It would seem therefore, if this account be correct, that the enlarged spines of *A. cavenia* are really Hymenopterous galls and that the Tineid larva observed by Fiebrig is a secondary, or inquiline which feeds on their medullary tissue. The authors say nothing about the Tineids nor the ants which later inhabit the spines.

The following are the ants that have been collected in these organs of *A. cavenia*:

- (1) *Pseudomyrma acanthobia* Emery (J. Bohls).
- (2) *Pseudomyrma acanthobia* var. *fusca* Emery (J. Bohls).
- (3) *Pseudomyrma fiebrigi* Forel (K. Fiebrig).
- (4) *Crematogaster (Orthocrema) brevispinosa* Mayr. (J. Bohls).
- (5) *Leptothorax (Goniothorax) echinatinodis* Forel subsp. *spininodis* Mayr. (J. Bohls).
- (6) *Cryptocercus bohlsi* Emery (J. Bohls).
- (7) *Cryptocercus grandinosus* F. Smith (J. Bohls).
- (8) *Cryptocercus peltatus* Emery (J. Bohls).
- (9) *Cryptocercus pilosus* Emery (J. Bohls).
- (10) *Cryptocercus quadratus* Mayr (J. Bohls).
- (11) *Cryptocercus (Paraeryptocercus) pusillus* Klug (J. Bohls).
- (12) *Cryptocercus (Cyathoccephalus) pallens* Klug (J. Bohls).
- (13) *Myrmelachista (Decamera) nodigera* Mayr var. *flavicornis* Emery (J. Bohls).

There is no evidence that any one of these species is an obligate tenant of *A. cavenia*. Many and probably all of them are generally found nesting in dead twigs or branches of other trees or shrubs. Even apart from this fact, there is nothing to indicate any "myrmecophilous" peculiarities in the plant.

(C) The African Acacias

Some six different species of African Acacias (*A. seyal* Delile, *drepanolobium* Harms, *formicarum* Harms, *pseudofistula* Harms, *mala-cocephala* Harms, and *Bussei* Harms) have been found to possess

greatly swollen spines inhabited by ants. Their consideration in this place may be greatly abbreviated both because I am dealing only with the Neotropical ant plants and because Bequaert (1921-22) has so recently and so adequately reviewed what is known in regard to the African Acacias. Most observers (Schweinfurth, Ascherson, Keller, Sjöstedt, Glover Allen, Winkler, Schenck, H. Lang) of these plants in the field admit that the hypertrophy of their spines is really a gall-formation produced by Dipterous or Hymenopterous larvæ. Bequaert gives the following reasons for adopting this interpretation: "They are not found on all specimens of the same species of Acacia, even in one locality; while on some plants practically all the thorns are swollen, others nearby bear hardly any galls; furthermore, their size is quite variable and their shape rather irregular. Mention may still be made of the fact that, while the species of Acacia enumerated above have a rather wide distribution in eastern Central Africa, swollen thorns have been noted in only a few localities within this range."

The list of ants inhabiting the spines, as compiled by Bequaert, comprises 16 species (13 species of *Crematogaster*, one of *Cataulacus* and two of *Tetraponera*), and he adds: "As would be expected from the fortuitous production of galls on plants, none of the ants mentioned in the preceding pages seems to restrict the location of its nest to galls. They are evidently all arboreal species which are in the habit of sheltering their brood in hollow branches or cavities of trees."¹

Chapter 7

VARIOUS NON-MYRMECOPHYTIC PLANTS WITH ANT-INHABITED STEMS, ETC.

Among the vast number of Neotropical plants there are many that cannot be regarded as myrmecophytes though they furnish ants with sufficiently commodious living quarters (pseudodomatia) in their internodes or other cauline structures. Several of these plants are so frequently or regularly inhabited as to be of special interest, at least to

¹Recent researches, which could not be considered by Professor Wheeler, throw a new light on the problem of the African ant Acacias. Paoli (1929 and 1930) studied two of the species, *Acacia fistula* Schweinfurth and *A. Bussei* Harms, in Somaliland. He found that these plants produce regularly two types of thorns. Some remain slender, while others slowly swell up at the base or over most of their length. The swelling is due to normal growth and is not induced by an insect. Eventually the pith of the swollen thorn dries up and the cavity thus produced may or may not be settled by ants. The swollen thorns are not insect galls but normal productions of the plant. The African ant Acacias are true myrmecophytes, but they lack the Beltian food-bodies of the American species. Both *A. fistula* and *A. Bussei* bear small nectaries on the petioles, but these are also found on non-myrmecophilous Acacias. Paoli and Menozzi (1930) list many ants found in the thorns of the African ant Acacias. [J. Bequaert

the collector. The simplest cases are those in which the internodes become hollow by disappearance of their pith before they are occupied. In others the ants clean out the pith and in still others they occupy burrows that have been more or less extensively excavated by wood-boring beetle larvæ, caterpillars, etc. The data cited in the literature often leave doubt as to which of these conditions obtains in a particular instance. The predilection of the ants for certain species of plants with inhabitable stems is probably enhanced by the regular development of extrafloral nectaries or the frequent occurrence of particular Coccids on their shoots or foliage. In the following paragraphs I have brought together a number of records on Neotropical plants that frequently have ant-inhabited stems, etc. There are, of course, several woody plants in temperate regions, like the common elder of our Northern and the white ash of our Southern states, the English walnut and blackberry in Switzerland, etc. which are similarly tenanted by a number of Formicidæ, but these plants I have not included.¹ I have also omitted the galls of oaks and other plants, which after serving for the development of their makers, often furnish convenient habitations for small colonies of ants, especially of the genera *Leptothorax*, *Crematogaster* and *Camponotus* (subgenera *Myrmentoma* and *Colobopsis*), etc. both in Europe and North America.

Gramineæ and Cyperaceæ

Bamboos. The hollow internodes of various bamboos are sometimes inhabited by ants, but this occurrence seems to be rather local. In the large Javanese bamboos surrounding the tropical laboratory at Kartabo, B. G. I failed to find any ants, and examinations of the smaller species in that region were equally negative. Lutz, Forel, Luederwaldt (1926) and other collectors in Brazil and H. Schmitt in Costa Rica, however, have found quite a series of bamboo-inhabiting forms. The following is a list of the recorded species:

- (1) *Eciton (Acamatus) legionis* F. Smith. Abundant in bamboos (in temporary nests?). Brazil (Garbe).
- (2) *Acanthoponera dolo* Roger. Brazil (Luederwaldt).
- (3) *Neoponera crenata* Roger. Brazil (Luederwaldt). Occupying nests made by *Camponotus albo-annulatus*.

¹See Ståger's interesting paper (1917) on the stem inhabiting ants of Switzerland. He records seven different forms, belonging to the genera *Leptothorax*, *Crematogaster*, *Dolichoderus* (*Hypoclinea*) and *Camponotus* (*Colobopsis*), as living in the stems of *Rubus ulmifolius*. Three of these forms occur also in walnut twigs. In all cases the ants invade cavities previously excavated by solitary wasps (mainly Crabronids), solitary bees (*Ceratina*) and beetles.

- (4) *Neoponera crenata* var. *moesta* Mayr. Brazil (H. Luederwaldt).
- (5) *Pseudomyrma gracilis* Fabr. subsp. *mexicana* Roger. In Bambusa. Costa Rica (H. Schmitt, cited by Menozzi, 1927).
- (6) *Pheidole bambusarum* Forel. Brazil (A. Lutz; Luederwaldt).
- (7) *Pheidole guilelmi-muelleri* Mayr subsp. *avia* Forel.
- (8) *Pheidole lutzi* Forel. Brazil (A. Lutz; Luederwaldt).
- (9) *Solenopsis saevissima* F. Smith. Brazil (H. Luederwaldt).
- (10) *Solenopsis franki* Forel subsp. *idæ* Forel. Brazil (H. Luederwaldt).
- (11) *Crematogaster (Orthocrema) brevispinosa* Mayr. Costa Rica (H. Schmitt).
- (12) *Crematogaster (Orthocrema) curvispinosa* Mayr. Costa Rica (H. Schmitt).
- (13) *Crematogaster (Orthocrema) distans* Mayr. subsp. *parviceps* Forel. Brazil (Luederwaldt).
- (14) *Crematogaster (Orthocrema) lutzi* Forel. Brazil (A. Lutz).
- (15) *Crematogaster (Orthocrema) quadriiformis* Mayr. In *Guadua distorta*. Brazil (Luederwaldt).
- (16) *Cryptocerus maculatus* F. Smith. In Bambusa. Costa Rica (H. Schmitt).
- (17) *Cryptocerus (Paracryptocerus) minutus* Fabr. In Bambusa. Costa Rica (H. Schmitt).
- (18) *Cryptocerus (Cyathocephalus) segulifer* Emery. In Bambusa. Costa Rica (H. Schmitt).
- (19) *Cryptocerus (Cyathocephalus) varians* F. Smith subsp. *marginatus* Wheeler and Mann. Haiti (W. M. Mann).
- (20) *Iridomyrmex iniquus* Mayr. In bamboo: Brazil (H. Luederwaldt).
- (21) *Iridomyrmex iniquus* var. *succinea* Forel. Brazil (Luederwaldt).
- (22) *Iridomyrmex leucomelas* Forel. Brazil (Luederwaldt).
- (23) *Tapinoma atriceps* Emery. Brazil (A. Lutz; Luederwaldt).
- (24) *Tapinoma atriceps* var. *breviscapa* Forel. Brazil (A. Lutz; Luederwaldt).
- (25) *Myrmelachista (Decamera) bambusarum* Forel. Brazil (Goeldi).
- (26) *Myrmelachista (Decamera) paderevskii* Forel. Brazil (A. Lutz).
- (27) *Camponotus (Tanæmyrmex) lutzi* Forel. Brazil (A. Lutz).
- (28) *Camponotus (Myrmothrix) cingulatus* Mayr var. *bambusarum* Forel. Brazil (Goeldi; Luederwaldt).
- (29) *Camponotus (Myrmomalis) emeryodicatus* Forel. Brazil. Luederwaldt).

(30) *Camponotus (Myrmobrachys) planatus* Roger var. *acaciæ* Emery. In *Bambusa*. Costa Rica (H. Schmitt).

(31) *Camponotus (Pseudocolobopsis) alboannulatus* Mayr. In *Bambusa taquara* Nees. Brazil (Luederwaldt).

(32) *Camponotus (Hypercolobopsis) paradoxus* Mayr subsp. *janitor* Forel. Brazil (A. Lutz); in *Bambusa taquara* Nees and allied species (Luederwaldt).

Luederwaldt also records two Coccids, *Lachnodiella taquaræ* Hemp. and *Orthezia grandis* Hemp., living with *Pheidole lutzii* in the bamboo internodes.

Uniola and *Cladium*. In the Bahamas I frequently took colonies of the following ants in the culms of a large grass, *Uniola paniculata* L., and a sedge, *Cladium jamaicense* Crantz:

- (1) *Pseudomyrma elongata* Mayr.
- (2) *Pseudomyrma flavidula* F. Smith
- (3) *Macromischa splendens* Wheeler
- (4) *Crematogaster (Acrocalia) lucayana* Wheeler
- (5) *Cryptocerus (Cyathocephalus) varians* F. Smith
- (6) *Tapinoma littorale* Wheeler
- (7) *Camponotus (Colobopsis) culmicola* Wheeler

According to Stäger (1917), the regular occurrence of ants in the culms of grasses in the savannahs of Colombia was noticed by E. A. Goeldi as early as 1896. I have found various species of *Pseudomyrma*, especially *Ps. flavidula*, in the same situations in Panama and British Guiana.

Scitamineæ

Crematogaster (Orthocrema) limata F. Sm. subsp. *parabiotica* Forel, which is associated with so many different plants, is also recorded as living in the inflorescences of a species of *Costus* in Amazonas (E. Ule).

Orchidaceæ

Many of the epiphytic species of this huge family undoubtedly harbor ants either about their roots, between the plants and their support, or in the pseudobulbs, but unfortunately many of the observations lack precision and the ants have rarely been identified. Bequaert (1921-22) includes only two Neotropical genera, *Diacrium* and *Schomburgkia*, among the myrmecophytes. He remarks that "*D. bicornutum* (Hooker), of Trinidad and Guiana, has a swollen spindle-shaped stem which is normally hollow and perhaps regularly inhabited by ants

(Rodway, 1911, p. 111). Schlechter claims that even under cultivation the pseudobulbs form at their base a slit through which the ants gain access into the cavity." The pseudobulbs of several species of *Schomburgkia* are also described as hollow and ant-inhabited. I find that only the following ants have been identified as orchid tenants:

(1) *Neoponera villosa* F. Smith. In pseudobulbs of *Schomburgkia tibicinis* Bateman. Mexico (G. Mayr).

(2) *Monomorium floricola* Jerdon. In pseudobulbs of *Epidendrum imatophyllum* Lindl. Honduras (O. Ames).

(3) *Crematogaster (Orthocrema) armandi* Forel. In pseudobulbs of an orchid. Brazil (S. Moore).

(4) *Crematogaster (Orthocrema) limata* F. Sm. subsp. *parabiotica* Forel. In pseudobulbs of *Diacrium bilamellatum* Hemsl. Panama (Wheeler).

(5) *Azteca tonduzi* Forel. In pseudobulbs of an orchid. Costa Rica (Tonduz).

(6) *Azteca velox* Forel subsp. *nigriventris* Forel. In pseudobulbs of *Diacrium bicornutum*, "in constant symbiosis". Costa Rica (P. Biolley).

(7) *Dolichoderus (Monacis) bispinosus* Olivier. Nesting in a tuft of orchids. Costa Rica (P. Biolley).

Polygonaceæ

Coccoloba uvifera (L.) Jacq.—This peculiar tree, the "seagrape", is very common along the beaches of tropical Florida and the West Indian islands. As a rule it is less than 15 ft. high, but in exceptional cases may attain a height of nearly 50 ft. with a trunk more than a yard in diameter. The peculiar, leathery, shining, orbicular or even transversely elliptical leaves make it a quaint and conspicuous object among the littoral vegetation. The foliage is a source of food for a number of miscellaneous insects and the following ants not infrequently inhabit its short, hollow internodes:

(1) *Pseudomyrma elongata* Mayr. Bahamas (Wheeler).

(2) *Pseudomyrma elongata* var. *cubaensis* Forel. Cuba (Wheeler).

(3) *Cryptocerus (Cyathocephalus) varians* F. Smith. Florida (Miss Nancy B. Fairchild).

(4) *Tapinoma melanocephalum* Fabr. Cuba (Wheeler).

(5) *Myrmelachista ambigua* Forel subsp. *ramulorum* Wheeler. Porto Rico (Wolcott). The "hormiguilla", a pest of the coffee plantations.

(6) *Camponotus (Tanæmyrmex) ramulorum* Wheeler. Bahamas and Cuba (Wheeler).

(7) *Camponotus (Tanæmyrmex) ramulorum* var. *marcidus* Wheeler. Bahamas (Wheeler).

(8) *Camponotus (Tanæmyrmex) ustus* Forel. Porto Rico (Wheeler).

(9) *Camponotus (Myrmobrachys) planatus* Roger. Cuba (Wheeler).

(10) *Camponotus (Myrmosphincta) scrguttatus* Fabr. Porto Rico (Wheeler).

The following insects have been recorded by Wolcott (1923) as infesting *C. wifera* in Porto Rico:

(1) *Dichomeris zingarella* Walsh. Gelechiid moth reared by Busck.

(2) *Ctenodactylomyia watsoni* Felt. Cecidomyid fly reared from galls by R. H. van Zwaluwenburg.

(3) *Cryptocephalus perspicax* Weise. Chrysomelid beetle feeding on foliage.

(4) *Attelabus coccolobæ* Wolcott. Curculionid beetle feeding on foliage.

(5) *Exophthalmodes roseipes* Chev. Curculionid beetle feeding on young foliage.

(6) *Lachnopus curvipes* Fabr. Curculionid beetle on foliage.

(7) *Tangia* sp. Fulgorid.

(8) *Ormenis marginata* Brunnich. Fulgorid.

(9) *Ormenis pygmæa* Fabr. Fulgorid.

(10) *Cyarda* sp. Fulgorid.

(11) *Pseudococcus nipæ* Maskell. Coccid.

Coccoloba rugosa Desf. The few specimens of this plant which I saw in Porto Rico, were small trees with very slender trunk and few branches and enormous, coarse, cordate clasping leaves, (30-60 cm. broad!) among the large veins of which, on the lower side and at the base, a small yellow ant, *Iridomyrmex melleus* Wheeler builds fragile carton nests. More frequently, however, this insect inhabits the hollow twigs of a variety of plants.

In South America Spruce (1869, in 1908) observed a number of ant-inhabited *Coccoloba* species as well as several other Polygonaceous genera (*Campderia*, now included in *Coccoloba*, *Lymmeria*, *Rupprechtia* and *Triplaris*, which has been considered at length (p. 41 *et. seq.*) All these, he says, "grow in moist situations, and most of them on lands subject to inundations. Not only is every lignescent Polygonaea a habitation for ants, but the whole of the medulla of every plant, from the root nearly to the growing apex of the ramuli, is scooped out by these insects." These are species of *Pseudomyrma*, which are

known in Brazil by the name of "Tachi" or "Tacyba" and in Peru by that of "Tangarana"; and in both countries the same name is commonly applied to any tree they infest as to the ants themselves." One of the species of *Coccoloba* he mentions in particular: "Some Tachi trees seem as if they were actually trying to run away from the ever encroaching ants. *Coccoloba parimensis* Benth., found by Schomburgk in British Guiana and by myself on the Rio Uaupes, is an arbuscle with a stem 15 feet long, that tapers upwards and arches over so as finally to touch the ground, the ants all the while hollowing it out, as it stretches away apparently in the hopeless attempt to escape their invasion. Some slender *Coccolobas* climb high into the adjacent trees, not by twining but by crooking their branches and thereby hoisting themselves up; others are self-standing bushy trees, but still have the same slender geniculate branches."

Bixaceæ

Bixa orellana L. This bush or small tree, the "achiote anatto" or "anotto", reaches a height of 30 feet and is common throughout the West Indies and Central America. It is also cultivated both in these regions and in the East Indies, because the testa of its seeds yields a beautiful orange-red pigment, the famous "anatto" dye. Pittier (1908) gives a good account of the plant. "*Bixa orellana*, a tree of even elegant aspect, with rose-colored, rather conspicuous flowers and capsular fruit covered with soft spines and containing numerous seeds. The outer integument, or tests of these seeds contains a yellowish red coloring matter, which is extracted with hot water and may be decomposed, according to Chevreul, into two pigments, *burine*, which is yellow, and *orelline*, which is red.

"*Bixa orellana* seems to be indigenous through tropical America and has been cultivated like the cacao plant from the most remote times. It easily matures its fruits up to an altitude of 1200 m. The wild form, or *Achiote simarron*, with smaller leaves and less developed floral organs, is found in Costa Rica in the woods of the Pacific slope up to an altitude of about 800 m. The dye which is fast and brilliant, was used by the natives to stain their bodies and also to tint their clothes and various other objects. At the present time it is employed in coloring butter, certain kinds of cheese, sarozas and other fabrics. In the Creole Kitchen it is used to color rice.

"The word *achiote*, which designates both the tree and the red, resinous paste extracted from the outer integument of the seeds, is a

corruption of the Nahoutl *achiotl*. In the Guyanas, the Indians call the tree and its product *uruau*, whence the French *rocou* is derived. *Anatto* and *bija* are other names applied to the coloring matter."

I found this plant rather common in Panama, both in the gardens at Ancon and Balboa, and wild, especially on the Atlantic side of the Isthmus near Colon, Marajal and on Barro Colorado Island. The leaves are broad and subcordate and covered with microscopic glands beneath, while the petiole bears at its junction with the blade a distinct thickening with a pair of large nectaries, which are visited by the ants. It is not surprising to find, therefore, that the short internodes are frequently inhabited by these insects. These internodes have to be hollowed out by the ants, since they contain a persistent pith. A careful study of the plant in different localities will probably yield quite a number of different ants. The following list includes those collected in Panama and two forms recorded by Forel (1906) as taken in Costa Rica:

(1) *Pseudomyrma flavidula* F. Smith var. *capperi* Forel. In twigs, Ancon, C. Z. (Wheeler).

(2) *Pseudomyrma gracilis* Fabr. In twigs. Marajal, C. Z. (Wheeler).

(3) *Pseudomyrma clongata* Mayr. var. *taudem* Forel. In the trunk and fruits. Costa Rica (P. Biolley).

(4) *Pseudomyrma sericca* Mayr var. *ita* Forel. In trunk and leaves. Costa Rica (P. Biolley). In twigs. Barro Colorado Isl. (Wheeler).

(5) *Crematogaster (Orthocrema) brevispinosa* Mayr. Feeding at foliar nectaries. Ancon, C. Z. (Wheeler).

(6) *Crematogaster (Orthocrema) brevispinosa* var. *ampla* Forel. In twigs. Barro Colorado Isl. (Wheeler).

(7) *Leptothorax (Goniothorax) echinatinodis* Forel var. *cordincola* Wheeler. In twigs. Barro Colorado Isl. (Wheeler).

(8) *Cryptocerus (Paraeryptocerus) minutus* Fabr. In twigs. Barro Colorado Isl. (Wheeler).

(9) *Dolichoderus (Hypoclinea) championi* Forel subsp. *trinidadensis* Forel var. *taniiatus* Forel. In twigs. Marajal C. Z. (Wheeler).

(10) *Tapinoma melanoccephalum* Fabr. In twigs. Ancon, C. Z. (Wheeler).

(11) *Paratrechina longicornis* Latr. Feeding at nectaries. Ancon, C. Z. (Wheeler).

(12) *Camponotus (Myrmobrachys) zoc* Forel. In twigs. Barro Colorado Isl. (Wheeler).

(13) *Camponotus (Pseudocolobopsis) claviscapus* Forel. In twigs. Marajal, C. Z. (Wheeler).

The general insect fauna of *B. orellana* has not been studied, but is probably quite as abundant as that of many other tropical trees. Two species of Coccids, *Howardia biclavis* Comst. and *Inglisia vitrae* Cockerell, are cited by Wolcott (1923) as infesting the plant in Porto Rico.

Marcgraviaceae

The types of *Azteca longiceps* Emery subsp. *juruenssis* were taken by E. Ule in the twigs and branches of a species of *Schwartzia* (Norantea) in Amazonas.

The juvenile stages of the species of *Marcgravia* have the form of peculiar creepers which adhere to the trunks of trees or the surfaces of rocks. On each side of the reptant stem is a regular series of round-elliptical leaves which are also rooted to their support and leave very narrow spaces between the latter and their lower surfaces. In a plant running over rocks in the island of Dominica, W. I., I found numerous colonies of *Wasmannia auro-punctata* Roger utilizing these spaces for nesting purposes. I believe that Forel (1899-1900) must have taken the type specimens of his *Azteca hypophylla* in such quarters under the leaves of some species of *Marcgravia* in Colombia, since he describes the insect as making "its nest under the round leaves of a plant which climbs on tree trunks like ivy. The ant glues the edges of each leaf to the bark of the tree with carton."

Combretaceae

Terminalia catappa L. This fine tree, known as the Indian or Malabar almond and in tropical America, where it is often planted for shade, as the "almandra" is easily recognized by its peculiar shining, obovate leaves, about 15 to 20 cm. long, with narrow cordate base and short petiole, clustered at the ends of the branches. The small white flowers are in slender spikes and the drupe-like, edible fruit has the shape of an almond. The internodes of its terminal twigs are often hollowed out and inhabited by ants. I have collected the following species from the trees planted along the drives at Balboa and Fort Amador, C. Z. Panama:

- (1) *Pseudomyrma zebelli* Forel.
- (2) *Solenopsis laeviceps* Mayr var. *antoniensis* Forel.
- (3) *Solenopsis zeteki* Wheeler.
- (4) *Crematogaster (Orthocrema) brevispinosa* Mayr var. *ampla* Forel.

(5) *Cryptocerus (Paracryptocerus) minutus* Fabr.

(6) *Dolichoderus (Hypoclinaea) lutosus* F. Smith.

Although *T. catappa* is not a native of the American tropics it is nevertheless attacked by several insects, as shown by the following data collected by Wolcott (1923) in Porto Rico:

(1) *Megalopyge krugii* Dewitz. Megalopygid moth, feeding on foliage.

(2) *Oecetius kirbyi* Guilding. Psychid moth cited by Gundlach as feeding on foliage.

(3) *Attelabus sexmaculatus* Chev. Curculionid beetle, feeding on foliage.

(4) *Aspidiotus destructor* Sign. Coccid.

(5) *Pseudococcus virgatus* Ckll. Coccid.

(6) *Saissetia nigra* Nietn. Coccid.

(7) *Saissetia oleæ* Bernard. Coccid.

(8) *Chrysomphos aonidum* L. Coccid.

In the twigs of another Terminalia, *T. (Bucida) buccras* Wright, in Porto Rico, I have taken colonies of the "hormigilla", *Myrmelachista ambigua* subsp. *ramulorum*.

Bombacaceæ

Probably the twigs of several of the large arborescent species of this family, especially the various silk-cotton trees, are not infrequently inhabited by ants, but I can cite only two records in support of this statement. J. Huber and A. Goeldi found twigs of *Bombax mungaba* on the Rio Purus, Amazonas, inhabited by *Pseudomyrma sericea* Mayr., and recently Prof. Jack, of the Arnold Arboretum, has brought me from Soledad, Cuba a number of twigs of the common Ceiba tree, *Ceiba pentandra*, containing colonies of *Pseudomyrma elongata* Mayr var. *cubaensis* Forel.

Euphorbiaceæ

Mabea. About 30 species of this genus are known from Brazil and the Guianas. They are shrubs with flexuous hollow branches, and some of the species, especially *M. fistulifera* Mart. and *angustifolia* Benth., seem to be well-supplied with calyx-glands, or glands on the branches of the flower panicles (*M. occidentalis* Benth). The latex of *M. piri* Aubl. of the Guianas yields rubber, and the stems of both it and *M. fistulifera* are made into tobacco-pipes, as described in the following quotation from Spruce: "The Mabeas are still more remark-

able than the *Tachias*, the long sarmentose branches stretching away to a great length among the adjacent vegetation, although never actually twining. All *Mabeas* of the section *Taquari* have this habit, and all are infested by *Tachi* ants. The slender but tough twigs, hollowed and polished internally by ants, are a favorite material for tobacco-pipes with the Indians of the Amazon, who strip off the bark and paint and varnish the surface of the wood. These "*Taquaris*", as they are called, are commonly sold in the shops of Para. A bundle of them which I purchased there is now in the Kew Museum. The arborescent *Mabeas*, however, with tall erect trunks and paniculate inflorescence, are apparently never touched by ants."

Sapium. Also a fairly large genus, comprising 25 species, and occurring in both the Old and New World tropics. The petioles have a pair of nectaries at their junction with the leaf-blade. The latex of *S. biglandulosum* Aubl. is medicinal and is said to yield a kind of rubber, but that of some of the Panamanian species (probably *S. aucuparium* Jacq.) is, according to Standley (1928 p. 240), "chewed by boys, who place it on twigs for the purpose of catching small birds." *Azteca longiceps* Emery subsp. *sapii* Forel was taken by Ducke in the hollow stems of *S. glandulosum* Morong at Ica, Amazonas, and *Pseudomyrma caroli* Forel var. *sapii* Forel was taken by E. Ule in the hollow stems of an undetermined species of *Sapium* at Bom Fim, Jurua, Amazonas.

Alchornea. This genus contains about 30 species of frutescent or arborescent plants which, like those of the preceding genus, occur in the tropics of both hemispheres. The leaves are described as bearing two to several nectaries at the base on the lower side. *A. irucurana* Casar of the subgenus *Eualchornea* is called "*irucurana*" or "*arariba*" by the Brazilians and yields a valuable wood. The following ants have been taken in its branches by K. Fiebrig in Paraguay:

- (1) *Pseudomyrma serieca* Mayr var. *ita* Forel.
- (2) *Forelius maccooki* Forel var. *brasiliensis* Forel.
- (3) *Forelius maccooki* subsp. *fiebrigi* Forel.
- (4) *Azteca luederwaldti* Forel.

Leguminosæ

The genera of this huge family, besides *Tachigalia*, which have been cited as containing species tenanted by ants in the New World are *Sclerolobium*, *Pterocarpus* and *Platymischium*:

Sclerolobium. According to Spruce, "the species of *Sclerolobium* are

not usually riparian but one species (*S. odoratissimum* sp. nov.) is eminently so, constituting a great ornament of the shores and islands of the Rio Negro towards the mouth of the Casiquiari, and perfuming the whole breadth of the river with the abundance of its pale yellow honey-scented flowers; and it is notable that this is the only species of the genus in which I have found sacciferous petioles. The sac is large, extending upwards from the knee of the petiole to the base of the second pair of leaflets, and it has a furrow along the upper surface. I presume the ants have been induced to take up their residence on these particular plants on account of the abundance and long persistence of their honied flowers." Bequaert believes that the petiolar sac described by Spruce "is merely an insect gall which, when empty, becomes settled by ants." In two other species, *S. tinctorium* Benth and *paniculatum* Vog., Spruce observed no such domatia but merely the infestation of the flower-panicles "with little fire-ants, which, however, seemed to have their permanent habitation in the ground about or near the tree-roots, and never to perforate the leaf-stalks."

Pterocarpus. A genus comprising some 20 species of tropical trees with hard wood of considerable economic value and yellow flowers. Spruce alludes to *P. ancylocalyx* Benth as "a small tree on the banks of the Solimoes, or Upper Amazon, which has the rachis of the racemes thickened in the middle, the swelling being sometimes (but not ways) tenanted by ants." Forel (1904) cites *Pseudomyrma sericea* Mayr as having been taken by E. Ule in the swollen flower-axes and twigs of *P. uli* Harms at Jurua Miry, Amazonas.

Platymischium. A Neotropical genus comprising trees or shrubs with yellow flowers. In at least some of the 15 known species the stems are hollow or even dilated at the nodes and inhabited by ants. The following species have been recorded:

(1) *Pseudomyrma picta* Stitz var. *heterogyna* Wheeler. In *Platymischium* sp. Bolivia (W. M. Mann).

(2) *Pseudomyrma sericea* Mayr var. *longior* Forel. In *P. stipulare* Benth. Amazonas (E. Ule).

(3) *Cryptocerus* (*Paracryptocerus*) *complanatus* Guérin subsp. *ramiphilus* Forel. In *P. uli* Harms. Amazonas (E. Ule).

(4) *Azteca huberi* Forel. In *Platymischium* sp. Brazil (J. Huber).

Perhaps some of the species of the genus *Cercidium* may prove to harbor several species of ants in their twigs and branches. *Cryptocerus rohweri* Wheeler, at least, has been taken from the branches of *C. torreyanum* in Arizona.

Olacaceæ

Agonandra. In Paraguay Chodat and Vischer (1920) found two species of ants, *Crematogaster* (*Orthocrema*) *goeldii* Forel var. *chodati* Forel and *Cryptocerus eduarduli* Forel, nesting in the trunks of *A. brasiliensis* Benth. and Hook., a dioecious tree with pendent twigs, thin, elliptical, pointed leaves and small, racemose flowers. These botanists promise a fuller account of the plant and the nesting habits of the ants and merely remark (p. 193): "The material which we brought back from Paraguay for the purpose of studying the myrmecophily of the *Agonandra*, does not permit us to reach such precise (*sic!*) conclusions from this genus as those obtained from the *Cordias* and *Acacia carenia*. In fact, in the trunks of this species, the existence of medullary galleries with rather large openings excavated perpendicularly to the wood can be observed only at certain points. It seemed to us that these galleries had been excavated by some phytophagous insect, possibly a Coleopteron." It is difficult to see how "myrmecophily" can apply to this case any more than to the excavations made by *Camponotus herculeanus* L. and its various races in the wood of North American and European trees which have been previously attacked by various beetles.

Loranthaceæ

Phoradendron. In Arizona I found the stems of a mistletoe, *Ph. flavescens* Nutt. var. *villosum* Nutt., growing on live oaks (*Quercus emoryi*) to be regularly inhabited by colonies of *Crematogaster* (*Orthocrema*) *arizonensis* Wheeler. According to Schwarz (1901) the cavities tenanted by the ants are made by a Curculionid beetle of the genus *Otidocephalus*. The plant, indeed, constitutes the center of an interesting biocœnose, comprising a Coccid, *Pseudococcus phoradendri* Cockerell, which lives in the cavities with the ants, another Coccid, *Lecanium phoradendri* which, according to Schwarz, lives on the outer surface of the plant, a Coccinellid beetle, *Cephaloscymnus occidentalis* Horn, that feeds on this Coccid, a butterfly caterpillar, *Thecla halesus*, that consumes the leaves and two beetles, a Bostrychid (*Amphicerus* sp.) and a Scolytid (*Stephanoderus* sp.) that bore in the stems.

Rutaceæ

Xanthoxylon. Many years ago I found that the twigs of the common prickly ash of our Southern States, *X. clavis-Herculis* L., in the vicinity of Austin, Texas were frequently inhabited by colonies of three species

of *Pseudomyrma* (*Ps. brunnea*, *flavidula* and *pallida*), and more recently Mr. E. D. Christophersen has found two species of ants, *Pheidole gauthieri* Forel var. *oxymora* Forel and *Camponotus (Hypercolobopsis) christopherseni* Forel, living in the large, corky spines which stud the trunk of *X. panamense* P. Wils. Although I have often examined the spines of this Central American tree and have found in many of them galleries that had been eaten out by insects, I failed to find any ant-colonies. The species taken by Christophersen may have been chance occupants of such burrows.

Anacardiaceæ

Schinus. Luederwaldt (1926) records the occurrence of *Proecryptocerus subpilosus* F. Smith subsp. *lepidus* Forel in dry excrescences on *Schinus terebinthifolius* Raddi in Brazil.

Rubiaceæ

Borreria verticillata (L.) G. F. W. Mey. As previously stated (p. 92) the hollow, internodes of this low shrub are often inhabited by ants. I first observed it in a small clearing, covering less than an acre, behind the tropical laboratory at Kartabo, B. G., where it was very abundant. Its small, agglomerated flowers attracted many bees, especially *Meliponinae*, and the dead stems seemed to be even more frequently occupied by ants than those of any other plants in the vicinity. According to Schumann (1897 p. 114), *B. verticillata* is common in the warmer parts of America from Florida and Mexico to Argentina, not rare in West Africa and on the Cape Verde Islands and has been recorded from Mozambique. The following is a list of the ants taken in its internodes at Kartabo:

- (1) *Pseudomyrma acanthobia* Emery
- (2) *Pseudomyrma*
- (3) *Pseudomyrma*
- (4) *Pseudomyrma*
- (5) *Pseudomyrma gracilis* Fabr.
- (6) *Monomorium*
- (7) *Solenopsis corticalis* Forel
- (8) *Crematogaster*
- (9) *Leptothorax (Goniothorax) aculeatinodis* Emery
- (10) *Leptothorax (Goniothorax) spininodis* Mayr
- (11) *Cryptocerus maculatus* Fabr.
- (12) *Myrmelachista*
- (13) *Brachymyrmex*
- (14) *Camponotus*

Gentianaceæ

Tachia. The only Neotropical genus of this family said to be associated with ants is Aublet's *Tachia* (*Myrmecia* Schreber). In the original description of the type species, *T. guianensis*, Aublet states that "the trunk and branches, which are hollow, serve as a retreat for ants; it is for this reason that this shrub is called by the Galibis 'Tachi', which according to their report, signifies in their language "ant-nest". According to Spruce, "the pretty Gentianaceous shrubs of the genus *Tachia* have long, slender, hollow branches, that either hang down or support themselves on the branches of adjoining shrubs and trees; yet although this character is (as I suppose) an undoubted inheritance of the effects of ant-agency, it is singular that *Tachias* are now-a-days often found entirely free from ants; while the name taken by Aublet from the Tupi language, distinctly implies that in his day they were notoriously ant-infested." These remarks seem to indicate that the *Tachias* are not myrmecophytes but shrubs sometimes or perhaps only locally inhabited by ants. I have seen no record of Formicidæ from these plants.

Compositæ

The sole Neotropical representative of this huge family recorded as harboring ants is the Brazilian *Erigeron maximus* Link. H. von Ihering and Luederwaldt found its stalks inhabited by *Azteca muelleri* Emery var. *wacketi* Emery. Like the typical form of the species, this variety also makes carton nests in the trunks of *Cecropia*.

Chapter 8

THE EPIPHYTIC BROMELIACEÆ AND THEIR FAUNA

Some years ago considerable attention was devoted to a study of the epiphytic Bromeliaceæ of the Neotropical Region, but there are few references in the literature to their ant-inhabitants. Several botanists, notably Cedervall (1884), Schimper (1884, 1888, 1903), Ule (1900), Mez (1904), Wercklé (1909) and Aso (1910), investigated the anatomy and ecology of these epiphytes, and quite a number of zoölogists, including Fritz Müller (1879, 1880), Ad. Lutz (1903), Calvert (1910a, 1910b, 1911a, 1911b, 1917) Scott (1912), Knab (1912, 1913a, 1913b) and Picado (1911, 1912a, 1912b, 1913) were keenly interested in the associated faunal components.

Most of the Bromeliaceæ are epiphytes on living trees or shrubs, but some of them occasionally grow on inanimate supports, such as rocks, fences, telegraph-wires, etc., and although the majority of the genera and species are confined to the rain-forests, certain species, especially of the genus *Tillandsia*, flourish in dry, open forests or coppices or along sea-beaches. The plants lack stems and their slender roots serve merely for holding fast to the substratum. Many species, however, if they happen to be detached from their support, readily take root in the soil and continue their growth under the new conditions. The leaves grow in the form of a close rosette and are widened at their bases, which are arranged in such a manner that each can receive and retain a certain amount of water, derived from the rain, dew or mist, and both inorganic and organic detritus (windblown leaves, soil, etc.). Some large Bromeliads may thus store as much as 20 litres of water. On this account the epiphytic Bromeliaceæ have been called "tank-epiphytes" (Schimper, 1903) and may be regarded as so many diminutive pools, which in the tropics replace the ponds of more temperate regions (Picado, 1913). The flowers, often showy though evanescent, are usually borne on a long leafy or scaly axis arising from the center of the leaf-rosette. The fruits are very diverse, being in some species berries, in others dehiscent pods. Peculiar appendages in the form of asymmetrical wings or crests or pappus-like tufts of silky hairs are developed from the walls of the capsule and serve to disseminate the seeds, which are usually numerous but differ considerably in shape in the various genera.

Owing to the peculiar arrangement of the leaves and their method of retaining water and detritus, the plant, according to Picado, may be divided into an aquarium, or tank, and a terrarium; the former represented by all the water-holding leaf-bases, the latter by the dry detritus which is left over around the periphery after the older outer leaves of the rosette have decayed or broken off and therefore no longer hold water. Both the tank and the terrarium furnish habitations for a large number of organisms, the tank for many aquatic and amphibious creatures (Infusoria, fly, beetle, and dragonfly larvæ, Copepods, tree-frogs, etc.), the terrarium for many cryptobiotic forms (worms, Isopods, earwigs, Myriopods, Onychophora, etc.). Ad. Lutz believes that fully one fifth of the species of Brazilian mosquitoes, including some nosophoric (malarial) forms, regularly breed in the tanks of Bromeliaceæ and Knab has bred a number of Culicidæ from these plants in Mexico and Panama, including the large species of *Megarhinus*, which are so cannibalistic as larvæ that only one of them can

live in a single water-holding leaf-base. The foliage of the Bromeliads, of course, affords food for several caterpillars and leaf-eating beetles, bugs, etc., and opportunities for various spiders to construct their webs and prey on the small Diptera that oviposit in the tanks.

Ad. Lutz discovered that the water in the reservoirs is unusually pure and that no putrefaction of its organic contents takes place as long as it remains in contact with the plants' leaves. It has long been known that the latter are covered with peculiar shield-shaped scales (modified trichomes), which are always most abundant at the leaf-bases and have been carefully studied by Cedervall, Schacht, Schimper, Mez and Aso. Schimper showed that the water as well as its contained salts and organic substances are absorbed by the leaves through these scales and are essential to the growth of the plant. He states that "the spoon-shaped leaf-bases, which are closely applied to one another, nearly always contain even during the dry season considerable quantities of water, pieces of rotting leaves and twigs, dead animals and earthy materials of undeterminable origin. *Experiments have shown that these substances are not only utilized by the plant but are indispensable to its existence*, since the roots, under the most favorable circumstances, take up but little water to cover transpiration and besides under natural conditions usually remain quite dry on account of the umbrella-like form of the leaf-rosette."

Most epiphytic Bromeliads are strongly negatively geotropic and therefore grow in such a position that their leaves are able to acquire and retain considerable quantities of meteoric water and both mineral and organic detritus. Picado has confirmed and extended the observations of previous investigators. He has proved that the absorption of minerals from the water by the leaves is an habitual nutritive process and that these organs secrete a gummy matter (consisting of 77% bassorine and 27% arabine and diverse soluble substances), which has both an amyolytic and a proteolytic action on the organic detritus and thus renders its soluble portions assimilable by the plant. The insoluble refuse forms a kind of peat, which is free from proteids and starches and contains almost no traces of salts. Hence the purity of the water in the reservoirs and the absence of decomposition in the refuse as long as it remains in contact with the living leaves. The terrarium, which is formed by the failure of the old peripheral leaves to form adequate reservoirs, may therefore be said to consist of peat. The formation of this substance in the Bromeliads is compared by Picado with that of our northern bog-peat, which, unlike humus, is formed only under water that is constantly being renewed, "The climate, in which these

plants (the epiphytic Bromeliaceæ) grow, being very mild and the water in their tanks being very pure, since all its soluble contents are slowly and constantly diminished by absorption, the decomposition of the detritus occurs under conditions closely analogous to those that lead to the formation of peat-bogs."

The Bromeliaceæ of the type here discussed flourish most luxuriantly where they are fully exposed to the sun-light on the high branches of trees and where they would seem also to be most advantageously situated for securing their meteoric water and wind-blown detritus. But these conditions are rather unfavorable for ants, since as a rule they can find nesting sites only under the dry roots of the plants or in the terrarium. For this reason most of the water-bearing Bromeliads are avoided by the ants. There are certain species of *Tillandsias*, however, which afford them much more favorable quarters. Since Picado has very little to say about these plants, we may turn to Schimper who was led to study them closely, because he found them constituting the great bulk of the epiphytic flora in the West Indies, where he first investigated the ecology of tropical plants (1884). His earlier account is reproduced in his paper of 1888 from which I quote: "In *Tillandsia flexuosa*, which inhabits very dry, sunny situations, the tips of the leaves are brought together over the water-reservoir and spirally twisted around one another, so that the latter is entirely concealed from the direct sun-light but accessible to the rain and dew by means of the long slender canals. But the most complete protective arrangements are found in *Tillandsia bulbosa*, which also grows in sunny situations and is represented on our Plate IV. The leaves of *Tillandsia bulbosa* are spoon-shaped at their ensheathing bases, whereas the blade is cylindrical and either trough-like, with a narrow slit, or tubular, with the leaf-margins closely approximated or overlapping each other. The blade is always more or less strongly retroflected and twisted around its axis. The sheaths form an onion-like bulb which is nearly everywhere compactly closed and since they are strongly and convexly spoon-shaped and applied to one another only at their margins, contain large cavities continued above into the tubular leaf-blade and provided with only a very small orifice opening to the outside at the junction of the sheath and blade. The peripheral half of the tubular blade consists of chlorophyll-containing parenchyma and a very thin layer of aquiferous tissue; the inner side, on the contrary, is quite colorless and carpeted with exceedingly numerous, very large scales, which are sunk into a dense layer of aquiferous tissue. While young, the sheath, so far as it is covered by the other leaves, is devoid of

chlorophyll, thin and invested on both surfaces with scales which surpass in dimensions those of most of the other species and are so densely crowded that the epidermis is reduced to narrow strands. The plant entirely lacks the very strong negative geotropism characteristic of the rosettes of the epiphytic Bromeliaceæ. It occurs sometimes on the upper, sometimes on the lower side of the branches, or on perpendicular trunks, and grows in an erect horizontal or inverted position, without ever exhibiting a trace of geotropic curvature. The bulbs always contain water in their inner cavities, besides earthy materials and small dead insects, whereas the outermost cavities are free from such substances and harbor ants. That the contained water does not fall out, even in the inverted position, requires no explanation, since each chamber, except for its small upper orifice, is tightly closed on all sides. How the water manages to gain entrance, however, calls for brief explanation. If drops of water are allowed to fall on the margins of the leaf-blades, no matter whether they overlap or are merely approximated, the liquid is greedily sucked in by capillarity. The same occurs at the margins of the sheaths and at the narrow orifice at the base of the blade. By such means the cavities may be filled in a short time, and this occurs in nature in the presence of rain or dew. It should be emphasized, in case this experiment is repeated, that the first drop is less quickly imbibed, if the plant happens to have remained unmoistened for some time. The outer leaves are, in fact, usually not easily wetted and take up only a small amount of water. Not only does water enter the wetted bulbs, even when they are inverted, but they may, in any position, as shown in our figure, imbibe water and eventually conduct it into their reservoirs. The earthy materials, which are always present in the water, are derived from the small amounts of solids that are washed off by the rain from the leaves and branches of the host tree. The plant probably secures its nitrogen also from the cadavers of the ants, which are not content to remain in the peripheral cavities, but also, as we have found, make fatal incursions into the water-containing chambers. The narrow orifice at the base of the leaf-blade serves the ants as an entrance."

Schimper thus seems to have been the first to notice the regular nidification of ants in *Tillandsias*. Since 1901, when I recorded the occurrence of several species of these plants in the neighborhood of Cuernavaca, Mexico, I have been able to observe the same association on numerous occasions in several of the countries surrounding the Gulf and Caribbean. The insects live only in the species of *Tillandsia* that have the leaf-bases more or less spoon-shaped and compactly

overlapping. The species concerned are those designated in the literature as *T. utriculata* Linn., *fasciculata* Swartz, *balbisiana* Schultes, *Benthamiana* Klobsch, *aloifolia* Hooker and *flexuosa*. Unfortunately I failed to collect specimens of the various *Tillandsias* in which I found Formicidæ so that I am in most cases uncertain of the precise identification of the plants.

The ants enter and leave the *Tillandsias* either through the small preformed openings near the tips of the expanded leaf-bases, as Schimper observed, or as I have observed, through larger openings which they gnaw in the leaf-bases. Since such openings are made not only in the outermost but also in the enclosed leaves, the ants are provided with a number of convenient intercommunicating chambers. Incipient colonies of larger species and single adult colonies of minute species often confine themselves to the space between a single leaf-base and those which it covers, so that not infrequently several different species may occupy as many different chambers in the same plant and live as near neighbors, or in a kind of parœcism or parabiosis. The ants rarely enter the wet innermost portion of the plant, and I am inclined to believe that the openings which they gnaw prevent any deleterious retention of water in their chambers by serving to draw it away. More careful observations in the field, however, are required to substantiate this supposition.

Owing to the fact that Picado did not bestow special attention on the *Tillandsias*, the Hymenoptera in his conspectus of the known Bromeliaceous fauna, comprise only the two species of ants (*Odonotomachus hastatus* and *Apterostigma calverti*) observed by Calvert in some of the genera of larger Costa Rican Bromeliads. The following list of ants shows that they really constitute a considerable portion of the Bromeliad biocœnose as a whole. I include several forms recorded by Luederwaldt as occurring among or under the roots of epiphytes. Though the latter may not have been Bromeliacæ, the ants mentioned very probably live occasionally among the roots of these plants. It is obviously impossible to draw a hard and fast line between such nesting sites and the ant-gardens.

(1) *Ectatomma (Gnamptogenys) annulatum* Mayr. Among roots of epiphytes with *Pheidole angusta* Forel as neighbors. Brazil (H. Luederwaldt).

(2) *Holcoponera striatula* Mayr. Nesting between leaves of Bromeliacæ. Brazil (H. Luederwaldt).

(3) *Necoponera crenata* Roger. In dried fruits of *Bromelia fastuosa* Lindh. and in *B. epiphytica*. Brazil (Luederwaldt).

(4) *Neoponera erenata* var. *mæsta* Mayr. In the same plants. Brazil (Luederwaldt).

(5) *Neoponera villosa* Fabr. Nesting under *Bromelia epiphytica*. Brazil (Luederwaldt).

(6) *Ponera distinguenda* Emery. Nesting under roots of epiphytes. Brazil (Luederwaldt).

(7) *Ponera foeda* Forel var. *sarolta* Forel. Nesting under roots of epiphytes. Brazil (Luederwaldt).

(8) *Anochetus altisquamis* Mayr. Among roots of Bromeliaceæ. Brazil (Luederwaldt).

(9) *Odontomachus hastatus* Fabr. Between leaves of Bromeliaceæ and other epiphytes. Costa Rica (P. P. Calvert); Brazil (Luederwaldt).

(10) *Pseudomyrma elongata* Mayr. In Tillandsias. Florida (Wheeler).

(11) *Pseudomyrma gebelli* Forel. In *Tillandsia aloifolia* Hooker. Panama (Wheeler).

(12) *Pseudomyrma gracilis* Fabr. subsp. *mexicana* Roger. In *Tillandsia Benthamiana* Klotsch. Mexico (Wheeler).

(13) *Pseudomyrma sericca* Mayr var. *vimeni* Forel. Under epiphytes. Brazil (Luederwaldt).

(14) *Pheidole anastasi* Emery var. *sospes* Forel. Among roots of epiphytes. Brazil (Luederwaldt).

(15) *Pheidole angusta* Forel. Among roots of epiphytes with *Ectatomma (Gnamptogenys) annulatum* as neighbors, Brazil (Luederwaldt).

(16) *Pheidole emeryi* Mayr. Among roots of epiphytes. Brazil (Luederwaldt).

(17) *Pheidole punctatissima* Mayr. In Bromeliaceæ and in green spathes of *Dicffenbachia oerstedii*. Costa Rica (P. Biolley).

(18) *Crematogaster acuta* Fabr. In *Tillandsia aloifolia*. Panama (Wheeler).

(19) *Crematogaster (Orthoerema) arcuata* Forel var. *aruga* Forel. In *Tillandsia aloifolia*. Panama (Wheeler).

(20) *Crematogaster (Orthoerema) brevispinosa* Mayr. In tufts of Bromeliaceæ.

(21) *Crematogaster (Orthoerema) brevispinosa* var. *minutior* Forel. In *Tillandsia Benthamiana*. Mexico (Wheeler).

(22) *Crematogaster (Orthoerema) brevispinosa* var. *tumulifera* Forel. In *Tillandsia aloifolia*. Panama (Wheeler).

(23) *Crematogaster (Orthoerema) distans* Mayr. In Tillandsias. Guatemala and Costa Rica (Wheeler).

- (24) *Crematogaster (Orthocrema) limata* F. Smith subsp. *parabiotica* Forel. In *Tillandsia aloifolia*. Panama (Wheeler).
- (25) *Crematogaster (Orthocrema) montezumia* F. Smith var. *sulcata* Mayr. In ant-gardens among Tillandsias. Brazil (E. Wasmann).
- (26) *Crematogaster (Orthocrema) sculpturata* Pergande. In Tillandsias. Guatemala (Wheeler).
- (27) *Crematogaster (Orthocrema) steinheili* Forel. In Tillandsias. Bahamas (Wheeler).
- (28) *Crematogaster (Orthocrema) virgula* Forel. In Tillandsias. Porto Rico (Wheeler).
- (29) *Crematogaster (Aerocalia) lucayana* Wheeler. In Tillandsias. Bahamas (Wheeler).
- (30) *Crematogaster (Aerocalia) lucayana* subsp. *ctiolata* Wheeler. In Tillandsias. Bahamas (Wheeler).
- (31) *Monomorium carbonarium* F. Smith subsp. *ebeninum* Forel. In Tillandsias. Bahamas (Wheeler).
- (32) *Monomorium floricola* Jerdon. In Tillandsias. Florida, Bahamas, Cuba, Porto Rico (Wheeler).
- (33) *Xenomyrmex stolli* Forel subsp. *floridanus* Emery var. *lucayanus* Wheeler. In Tillandsias. Bahamas (Wheeler).
- (34) *Solenopsis corticalis* Forel subsp. *amazonensis* Forel. In Tillandsias (*Pseudocatopsis* sp.). Peru (E. Ule).
- (35) *Solenopsis dceipiens* Emery subsp. *adjecta* Emery. Nesting under roots of epiphytes. Brazil (Luederwaldt).
- (36) *Macromischa petiolata* Forel. In *Tillandsia Benthamiana*. Mexico (Wheeler).
- (37) *Leptothorax (Goniothorax) echinatinodis* Forel subsp. *dalmasi* Forel. In Tillandsias. Costa Rica (Wheeler).
- (38) *Cryptocercus aztecus* Forel. In *Tillandsia Benthamiana*. Mexico (Wheeler).
- (39) *Cryptocercus scutulatus* F. Smith. In Tillandsias. Costa Rica (Wheeler).
- (40) *Cryptocercus wheeleri* Forel. In *Tillandsia Benthamiana*. Mexico (Wheeler).
- (41) *Cryptocercus (Cyathocephalus) varians* F. Smith. In Tillandsias. Florida (Wheeler).
- (42) *Apterostigma calverti* Wheeler. Nesting in Bromeliaceæ and making fungus-gardens of insect excrement. Costa Rica (P. P. Calvert).
- (43) *Cyphomyrmex auritus* Mayr. Nesting among roots of epiphytes. Brazil (Luederwaldt).

(44) *Dolichoderus* (*Hypoelinea*) *championi* Forel subsp. *trinidadensis* Forel var. *tæniatus* Forel. In *Tillandsia aloifolia*. Panama (Wheeler).

(45) *Dolichoderus* (*Hypoelinea*) *lutosus* F. Smith. In *Tillandsias*. Costa Rica (Wheeler).

(46) *Dolichoderus* (*Monacis*) *bispinosus* Olivier. In *Tillandsia aloifolia*. Panama (Wheeler).

(47) *Iridomyrmex iniquus* Mayr var. *nigellus* Emery. In *Tillandsias*. Guatemala and Costa Rica (Wheeler).

(48) *Iridomyrmex sordescens* Wheeler. In *Tillandsias*. Costa Rica (Wheeler).

(49) *Azteca traili* Emery subsp. *tillandsiarum* Wheeler. In *Tillandsias*. British Guiana (Wheeler).

(50) *Azteca velox* Forel. In *Tillandsias*. Costa Rica (Wheeler).

(51) *Tapinoma littorale* Wheeler. In *Tillandsias*. Florida and Bahamas (Wheeler).

(52) *Tapinoma littorale* var. *cubaënsis* Wheeler. In *Tillandsias*. Cuba (Wheeler).

(53) *Myrmelachista ambigua* Forel subsp. *ramulorum* Wheeler. In *Tillandsias*. Costa Rica (Wheeler).

(54) *Myrmelachista* (*Decamera*) *zeledoni* Emery. In *Tillandsias*. Costa Rica (Wheeler).

(55) *Paratrechina longicornis* Latr. In *Tillandsia paraënsis* Mez. Brazil (E. Ule).

(56) *Paratrechina* (*Nylanderia*) *fulva* Mayr. Nesting among leaves of Bromeliaceæ. Brazil (Luederwaldt).

(57) *Camponotus* (*Tanæmyrmex*) *conspicuus* F. Smith subsp. *zonatus* Emery. In *Tillandsias*. Costa Rica (Wheeler).

(58) *Camponotus* (*Tanæmyrmex*) *ramulorum* Wheeler. In *Tillandsias*. Bahamas (Wheeler).

(59) *Camponotus* (*Tanæmyrmex*) *ramulorum* var. *mareidus* Wheeler. In *Tillandsias*. Bahamas (Wheeler).

(60) *Camponotus* (*Myrmothrix*) *cingulatus* Mayr. Among Bromeliaceæ. Brazil (Luederwaldt).

(61) *Camponotus* (*Myrmothrix*) *abdominalis* Fabr. var. *costaricensis* Forel. In *Tillandsias*. Costa Rica (Wheeler).

* (62) *Camponotus* (*Myrmothrix*) *abdominalis* subsp. *mediopallidus* Forel. In *Tillandsia Benthamiana*, Mexico (Wheeler).

(63) *Camponotus* (*Myrmobrachys*) *planatus* Roger. In *Tillandsias*. Cuba (Wheeler).

(64) *Camponotus* (*Myrmobrachys*) *planatus* var. *continentis* Forel. In *Tillandsias*. Florida (Wheeler).

(65) *Camponotus (Myrmophænus) fastigiatus* Roger subsp. *schmalzi* Emery. Nesting under epiphytic Bromeliaceæ. Brazil (Luederwaldt).

(66) *Camponotus (Myrmocladæcus) bidens* Mayr. In Tillandsias. Costa Rica (Wheeler).

(67) *Camponotus (Myrmocladæcus) rectangularis* Emery subsp. *rubroniger* Forel. In *Tillandsia Benthamiana*. Mexico (Wheeler).

(68) *Camponotus (Hypercolobopsis) paradoxus* Mayr subsp. *janitor* Forel. Among leaves of Bromeliaceæ. Brazil (H. von Ihering).

Many of the ants in this list live also in hollow twigs or in other available plant-cavities. One species, *Apterostigma calverti*, is of unusual interest, because it belongs to the fungus-growing Attini, which regularly nest in the ground, under stones or in and under rotten logs, and is therefore exceptional in having become arboreal and in constructing its gardens in the terraria of large Bromeliads with the excrement of the beetles or caterpillars that feed on their foliage. *Cyphomyrma auritus* is another fungus-growing ant which is recorded by Luederwaldt as nesting and making its gardens among the roots of epiphytes. It may, therefore, occasionally live among Bromeliads. I am inclined to believe that yet another Attine ant, *Cyphomyrma salvini* Forel, may eventually be found to have similar habits. In Panama I have taken it repeatedly under the bases of palm-petioles several feet above the ground. It might, therefore, nest about the roots of Bromeliads and other epiphytes. Its gardens consist of the collected excrement of small insects and are covered with a peculiar fungus very much like that cultivated by *C. rimosus* Spinola and its various subspecies (see Wheeler, 1907, p. 771, Pl. 50, Fig. 29).

The colonies of ants nesting in Tillandsias are singularly free from parasites or myrmecophiles. The only ant-guest which I have seen in these plants is a flat, broadly elliptical *Microdon* puparium in a colony of *Pseudomyrma gracilis* subsp. *mexicana*, taken many years ago at Cuernavaca, Mexico. The adult larva of this insect is figured in my paper of 1901.

The following table includes a list of the number of species of bromeliadicolous animals cited by Picado and the ants above enumerated:

Rotifera.....	1	Thysanoptera.....	1
Planarians.....	5	Dermaptera.....	28
Oligochæta.....	8	Plecoptera.....	1
Hirudinea.....	3	Odonata.....	1
Gastropoda.....	4	Hemiptera.....	8
Crustacea.....	9	Coleoptera.....	36
Onychophora.....	2	Lepidoptera.....	3
Myriopoda.....	16	Diptera.....	88
Arachnida.....	29	Hymenoptera (Formicidæ)...	66
Thysanura.....	1	Batrachia.....	6
Orthoptera.....	29		
		Total.....	342

This list represents, of course, only a small fraction of the total bromeliadicolous fauna, which has nowhere been exhaustively studied. Throughout great regions of Brazil, Eastern Peru and the Guianas no systematic search has been made for the animal tenants of the luxurious epiphytic flora. The majority of the recorded species (261) are insects, but whole groups of animals, such as the Infusoria and Nematodes, have received no attention. If we include the Bacteria, fungi and algæ which must be associated with the plants and their tenants the total Bromeliad biocenosis would probably be very extensive. Picado mentions Saprolegniaceæ on the aquatic insect larvæ and Werckle states that the flowers of certain *Tillandsias* are often greatly injured by smuts. There are, as we have seen, fungi in the gardens of *Apterostigma calverti* and other species no doubt invade the peaty remains of the terraria. The fauna and flora of the epiphytic vegetation may be suggested as very promising subjects for intensive investigation in some of the laboratories that have been recently established in the American tropics.¹

¹Professor Wheeler intended to expand the Chapter dealing with the Bromeliaceæ. Among other recent papers which he wanted to review are three important contributions by Miss Skwarra (1930, 1934a, 1934e) on ant plants and bromeliads of Mexico. [J. Bequaert].

REFERENCES¹

ALFARO, A.

1935. Investigaciones científicas. (San José de Costa Rica), 219 pp.
(*Cecropia* and *Acacia*, pp. 195-203).

ASO, K.

1910. Können Bromeliaceen durch die Schuppen der Blätter Salze aufnehmen? *Flora*, **100**, pp. 447-450.

AUBLET, J. B.

1775. Histoire des plantes de la Guyane Française. (Paris), 4 vols.,
xxxii + 976 + 52 + 160 pp., 392 pls.

BAILEY, I. W.

- 1922a. The Anatomy of certain plants from the Belgian Congo, with special reference to myrmecophytism. *Bull. Amer. Mus. Nat. Hist.*, **45**, pp. 585-621, 16 pls.
1922b. Notes on Neotropical ant-plants. I. *Cecropia angulata* sp. nov. *Bot. Gazette*, **74**, pp. 369-391.
1923. Notes on Neotropical ant-plants. II. *Tachigalia paniculata* Aubl. *Bot. Gazette*, **75**, pp. 27-41, 2 pls.
1924. Notes on Neotropical ant-plants. III. *Cordia nodosa* Lam. *Bot. Gazette*, **77**, pp. 39-49, 2 pls.

BARBER, H. S.

1928. A new Bolivian silvanid beetle from the myrmecodomatia of *Cordia*. *Psyche*, **35**, pp. 167-168.

BECCARI, O.

- 1884-1886. Piante ospitatrici ossia formicarie della Malesia e della Papuasias. *Malesia*, **2**, pp. 1-340, 65 pls.

BELT, T.

1874. The naturalist in Nicaragua. (London), xvi + 403 pp.

BENTHAM, G.

- 1870-1876. Leguminosae. In de Martius and Eichler, *Flora brasiliensis*. (Munich), vol. 15, pt. 2.

BEQUAERT, J.

1922. Ants in their diverse relations to the plant world. *Bull. Amer. Mus. Nat. Hist.*, **45**, pp. 333-583, 4 Pls.

BEURLING, P. J.

1854. Primitiae Florae portobellensis sive enumeratio plantarum vascularum quas juxta oppidum Portobello in isthmo panamensi Americae centralis mense Aprili anno 1826 legit Joh. Em. Billberg. *Svensk. Vet. Akad. Handl.*, 105-148.

¹Prepared by Dr. Joseph Bequaert.

- BOHEMAN, C. H.
1853. *Monographia Cassidarum*. Vol. 3, (Holmiae).
- BOLDINGH, H.
1914. *The flora of Curaçao, Aruba and Bonaire* (Leiden).
- BORGMEIER, T.
1923. Beitrag zur Biologie der Feuerameise und ihrer Gäste. *Zeitschr. Deutsch. Ver. Wiss. Kunst, São Paulo*, **3**, pp. 1-9.
- BOULGER, G. S.
1908. *Wood. A manual of the natural history and industrial applications of the timbers of commerce*. (London).
- BREYNIUS, J.
1739. *Prodromi fasciculi rariorum plantarum primus et secundus*. (Gedani), 108 + 54 pp., 32 pls. (*Acacia*, pp. 36-38).
- BRUES, C. T.
1922. *Conoaxima*, a new genus of the Hymenopterous family Eurytomidae, with a description of the larva and pupa. *Psyche*, **29**, pp. 153-158.
- BUSCALIONI, L. AND HUBER, J.
1900. Eine neue Theorie der Ameisenpflanzen. *Beihefte Bot. Zentralbl.*, **9**, pt. 2, pp. 85-88
- CALVERT, P. P.
1910a. A plant-dwelling Odonate larva. *Ent. News*, **21**, pp. 264 and 365-366.
1910b. Zoological researches in Costa Rica. *Old Penn*, **9**, pp. 165-170.
1911. Studies on Costa Rican Odonata, II-III. *Ent. News*, **22**, pp. 402-411, 449-460, 3 Pls.
- CALVERT, A. S. AND CALVERT, P. P.
1917. *A year of Costa Rican natural history*. (New York), xix + 577 pp.
- CEDERVALL, E. V.
1884. Anatomiskt-fysiologiska Undersökningar öfver Bladet hos Bromeliaceerna. *Göteborgs Kongl. Vet. Handl.*, **19**, pp. 1-55, 5 Pls.
- CHAMPION, G. C.
1893-1894, *Cassididae*. In *Biologia Centrali-Americana*. Coleoptera, vol. 6, Pt. 2.
- CHODAT, R.
1920. Revision critique des espèces de *Cordia* appartenant à la section *Gerascanthus* (Cham.) et de quelques espèces du même genre. *Bull. Soc. Bot. Genève*, (2) **12**, pp. 209-218.
- CHODAT, R. AND CARISSO, L.
1920a. Une nouvelle théorie de la myrmécophilie. *C. R. Séances Soc. Phys. Hist. Nat. Genève*, **37**, pp. 9-12.

- 1920b. La végétation du Paraguay. La myrmécophilie des Cordias de la section *Gerascanthus*. Bull. Soc. Bot. Genève, (2) **12**, pp. 172-200.
- CHODAT, R. AND VISCHER, W.
1920. La vegetation du Paraguay. Borriginacées. Bull. Soc. Bot. Genève, (2) **12**, pp. 157-171.
- COBO, P. BERNABE
1653. Historia del Nuevo Mundo. Vol. 2, Lib. 6, Cap. CXXIX, p. 126 (Palo santo).
- COCKERELL, T. D. A.
1928. Bees collected by Dr. W. M. Wheeler at the flowers of *Triplaris*. Psyche, **35**, pp. 170-172.
- COMMELYN, J.
1697-1701. Rariorum plantarum Horti Medici Amstelodamensis descriptio (Amsterdam). Vol. 1, 220 pp., 112 pls.; Vol. 2, 224 + 112 pp.
- DAMMER, U.
1893. Polygonaceae. In Engler and Prantl, Die Natürlichen Pflanzenfamilien, (Leipzig), Vol. 3, Pt. 1, Abth. a, pp. 1-36.
- DARWIN, F.
1877. On the glandular bodies on *Acacia sphaerocephala* and *Cecropia peltata* serving as food bodies. Jl. Linn. Soc. London, Bot., **15**, pp. 398-409, 1 Pl.
- DELPINO, F.
1886-1889. Funzione mirmecofila nel regno vegetabile. Prodomo d'una monografia delle piante formicarie. Mem. Acc. Bologna, (4) **7**, pp. 215-323 (1886); (4) **8**, pp. 601-659 (1887); (4) **10**, pp. 115-147 (1889).
- DONISTHORPE, H. S. J. K.
1912. Myrmecophilous notes for 1911. Ent. Record, **24**, pp. 4-10, 34-40.
1923. Myrmecophilous notes for 1922. Ent. Record, **35**, pp. 1-9.
- DUCKE, A.
1910. Revision des guêpes sociales polygames d'Amérique. Ann. Mus. Nat. Hungar., **8**, pp. 449-544.
- EMERY, C.
1892. Hormigas de Costa Rica, que viven en las Acacias. Anal. Mus. Nac. San José, (1888-1889), pp. 65-67.
1893. Studio monografico sul genere *Azteca* Forel. Mem. R. Accad. Sci. Inst. Bologna, (5) **3**, pp. 319-352, 2 Pls.

1896. Alcune forme nuove del genere *Azteca* For. e note biologiche Boll. Mus. Zool. Anat. Comp. Torino, **11**, No. 230, pp. 1-7.
1912. Le piante formicarie. *Scientia*, **12**, pp. 48-62.
- FIEBRIG, K.
1909. *Cecropia peltata* und ihr Verhältnis zu *Azteca alfari*, zu *Atta sexdens* und anderen Insekten, mit einer Notiz über Ameisen-Dornen bei *Acacia cavenia*. Biol. Centralbl., **29**, pp. 1-16, 33-55, 65-77, 5 Pls.
- FOREL, A.
- 1899-1900. Formicidae. In *Biologia Centrali-Americana*. Hymenoptera, vol. 3.
1904. In und mit Pflanzen lebende Ameisen aus dem Amazonas-Gebiet und aus Peru. Zool. Jahrb. Syst., **20**, pp. 677-707.
1905. Miscellanea myrmécologiques. Ann. Soc. Ent. Belgique, **49**, pp. 155-160.
1906. Fourmis néotropiques nouvelles ou peu connues. Ann. Soc. Ent. Belgique, **50**, pp. 225-249.
1920. Fourmis trouvées dans les galles de *Cordia* et d'*Agonandra*, etc. Bull. Soc. Bot. Genève, **12**, 201-208.
- GÜRKE, M.
1897. Borriginaceae. In Engler and Prantl, *Die Natürlichen Pflanzenfamilien*, Vol. 4, pt. 3, Abt. a, pp. 71-131.
- HARMS, H.
1906. Leguminosae (in Ule, Beiträge zur Flora der Hyalea). Verh. Bot. Ver. Prov. Brandenburg, **48**, pp. 164-165.
- HEMSLEY, W. B.
- 1882-1886. *Biologia Centrali-Americana*, Botany, vol. 3. (*Triplaris*, p. 38).
- HERNANDEZ, F.
1651. *Rerum medicarum Novae Hispaniae thesaurus seu plantarum, animalium et mineralium Mexicanorum historia*. (Rome), xiv + 950 + 22 + 90 + 6 pp.
- HOHENKERK, L. S.
1918. Botanical identifications of British Guiana trees and plants. Jl. Board Agric. Brit. Guiana, **11**, pp. 98-106, 178-185.
- HUBER, J.
1909. Materiales para a flora amazonica. VII. Plantae Duckeanae austro-guyanenses. Bol. Mus. Goeldi, **5**, (1908), pp. 387-388.
1910. Novitates florae amazoniae (I). Bol. Mus. Goeldi, **6** (1909), pp. 60-90.

HUTH, E.

1887. Myrmecophile und myrmecophobe Pflanzen. (Berlin), 27 pp., 2 Pls.

IHERING, H. VON

1891. Die Wechselbeziehungen zwischen Pflanzen und Ameisen in den Tropen. Das Ausland, 1891: 474-477.
1907. Die Cecropien und ihre Schutzameisen. Engler's Bot. Jahrb., 39, pp. 666-714, 5 Pls.

JACQUIN, N. J.

1763. Selectarum stirpium americanarum historia. (Vienna), vii + 284 pp., 183 Pls.

JOHNSTON, I. M.

1924. Taxonomic records concerning American spermatophytes. 2. New or otherwise noteworthy plants. Contrib. Gray Herb. Harvard Univ., N. S. 70, pp. 69-87.

KNAB, F.

1912. A new species of Anisopidae (Rhyphidae) from tropical America. Proc. Biol. Soc. Washington, 25, pp. 111-112.
- 1913a. Larvae of Cyphonidae in Bromeliaceae. Ent. Mo. Mag., (2) 24, pp. 54-55.
- 1913b. A new bromeliocolous *Megarhinus*. Insec. Inscit. Menstr., 1, pp. 35-36.

KOELSCH, K. A.

1908. Die Theorie der Ameisenpflanzen—Ein Irrtum der Biologie. Beil. z. Allgem. Zeitg. München, No. 8: 59-61. (Review in 1910, Zeitschr. Wiss. Insektenbiol., 6, p. 36).

LENG, C. W.

1920. Catalogue of the Coleoptera of America, north of Mexico. (Mount Vernon, N. Y.).

LINNÉ, C. VON

1737. Hortus Cliffortianus. (Amsterdam), x + iv + 501 + 16 + 36 pp. 1 Pl.

LUEDERWALDT, H.

1926. Observações biológicas sobre formigas brasileiras, especialmente do Estado de São Paulo. Rev. Mus. Paulista, 14, pp. 1-118, 5 Pls.

MANN, W. M.

1928. A new *Microdon* from Panama. Psyche, 35, pp. 168-170.

MARCGRAVIUS DE LIEBSTAD, G.

1648. *Historiae rerum naturalium Brasiliae libri octo*. (Amsterdam), viii + 300 pp.

MAZA, M. G. DE LA AND ROIG, J. T.

1916. *Flora de Cuba*. Bol. 22, Estac. Agronomica Santiago de las Vegas.

MEINERT, F.

1892. *Traek of Insektlivet i Venezuela*. Entom. Meddelelser, 3, pp. 125-144.

MEISNER, K. F.

1856. *Polygoneae*. In De Candolle, *Prodromus Systematis Naturalis Regni Vegetabilis*, Vol. 14: 171-178.

MÉNÉGAUX, A.

1909. *Contribution à l'étude des Edentés actuels, Famille des Brady-podidés*. Arch. Zool. Exp. Gén., (5) 1, pp. 277-344, 4 Pls.

MENOZZI, C.

1927. *Formiche raccolti dal Sig. H. Schmidt nei dintorni di San José de Costa Rica*. Entom. Mitt., Berlin, 16, pp. 266-277, 336-345.
1930. *Formiche della Somalia italiana meridionali*. Mem. Soc. Ent. Italiana, 9, pp. 76-130, 3 Pls.

MEZ, C.

1890. *Morphologische und anatomische Studien über die Gruppe der Cordieae*. Engler's Bot. Jahrb., 12, pp. 526-588, 2 Pls.
1904. *Physiologische Bromeliaceen-Studien*. Jahrb. Wiss. Bot., 40, pp. 157-229.

MIQUEL, F. A. W.

1850. *Stirpes surinamenses selectae*. Natuurk. Verh. Bataafsch. Maatsch. Wetensch., (2) 7, (1851), 7 + 234 pp., 65 Pls.

MOEBIUS, M.

1925. *Versuch zur Erklärung der Ameisenpflanzen*. Flora, (N. F.) 118-119, pp. 393-398.

MORRISON, H.

1922. *On some trophobiotic Coccidae from British Guiana*. Psyche, 29, pp. 132-152, 2 Pls.

MORTEO, E.

1904. *Sopra due piante formicarie*. Malpighia, 18, pp. 504-511, 2 Pls.

MÜLLER, FRITZ

1876. *Ueber das Haarkissen am Blattstiel der Imbauba (Cecropia), das Gemüsebeet der Imbauba-Ameise*. Jen. Zeitschr. Naturw., 10, pp. 281-286.

1879. Wasserthiere in den Wipfeln des Waldes. Kosmos, Leipzig, **4**, pp. 390-392.
1880. Wasserthiere im Baumwipfeln. *Elpidium bromeliarum*. Kosmos, Leipzig, **16**, pp. 386-388.
1880. Die Imbauba und ihre Beschützer. Kosmos, Leipzig, **18**, pp. 109-116.
1883. (On *Cecropia*). Blumenauer Zeitung, January.
- PAOLI, G.
1929. Strane abitazioni di una formica su acacie della Somalia. Riv. Colonie Italiane, **3**, pp. 474-485, 4 Pls.
1930. Contributo allo studio dei rapporti fra le acacie e la formiche. Mem. Soc. Ent. Italiana, **9**, pp. 131-195, 7 Pls.
- PICADO, C.
1911. Les Broméliacées épiphytes comme milieu biologique. C. R. Acad. Sci. Paris, **153**, p. 960.
- 1912a. Sur la nutrition chez les Broméliacées épiphytes. C. R. Acad. Sci. Paris, **154**, p. 607.
- 1912b. Les mares aériennes de la forêt vierge américaine. Les Broméliacées. Biologica, **2**, pp. 110-115.
1913. Les Broméliacées épiphytes considérées comme milieu biologique. Bull. Scientif. France Belgique, **47**, pp. 215-360, 19 Pls.
- PISO, G.
1658. Historia naturalis Brasiliae. (Antwerp).
- PITTIER, H.
1908. Ensayo sobre las plantas usuales de Costa Rica. (Washington, D. C.).
- PLUKENETT, L.
1696. *Almagestum botanicum*. (London), 402 pp.
1720. *Opera omnia botanica*. 6 vols.
- PULLE, A.
1906. An enumeration of the vascular plants known from Surinam, together with their distribution and synonymy. (Leiden).
- RAY, J.
- 1686-1704. *Historia plantarum*. (London), 3 Vols. (*Cecropia*, Vol. 3, p. 1373).
- RETTIG, E.
1904. Ameisenpflanzen-Pflanzenameisen. Beih. z. Bot. Centralbl., **17**, pp. 89-122.
- RODWAY, J.
1911. In the Guyana forest. Studies of nature in relation to the struggle for life. (London) 2nd Ed., 326 pp.

ROSS, H.

1909. Pflanzen und Ameisen im tropischen Mexiko. Naturwiss. Wochenschr., **8**, pp. 822-830.

RUIZ, H. AND PAVON, J.

1799. Flora peruviana et chilensis. (Madrid), Vol. 2. (*Cordia*, Vol. 2: 47; Pl. 184).

SAFFORD, W. H.

1910. Bull-horn acacias in botanical literature, with a description of two new species. Science, (N. S.) **31**, pp. 676-677.
1914. *Acacia cornigera* and its allies. Jl. Washington (D. C.) Ac. Sci., **4**, pp. 356-368.
1915. New or imperfectly known species of bull-horn acacias. Jl. Washington (D. C.) Ac. Sci., **5**, pp. 355-360.
1922. Ant acacias and acacia ants of Mexico and Central America. Smithson. Rept. for 1921: 381-394, 15 Pls.

SALDANHA DA GAMA, JOSÉ DE

1874. Notes sur quelques arbres employés dans l'industrie brésilienne. Ann. Sci. Nat., Bot., (5) **19**, pp. 210-218.

SCHENCK, H.

1913. Acaciae myrmecophilae novae. Fedde's Repertorium, **12**, p. 370.
1914. Die myrmekophilen *Acacia*-Arten. Engler's Bot. Jahrb., **50**, Suppl., pp. 449-487.

SCHIMPER, A. F. W.

1884. Ueber Bau und Lebensweise der Epiphyten Westindiens. Botan. Centralbl., **17**, pp. 192-195, 223-227, 253-258, 284-294, 319-326, 350-359, 381-389, 2 Pls.
- 1888a. Die Wechselbeziehungen zwischen Pflanzen und Ameisen im tropischen Amerika. (Jena), 95 pp., 3 Pls.
- 1888b. Die epiphytische Vegetation Amerikas. (Jena), 162 pp., 6 Pls.
1903. Plant geography upon a physiological basis. (Oxford), xxx + 839 pp.

SCHOMBURGK, RICHARD

- 1847-1848. Reisen in Britisch-Guiana in den Jahren 1840-1844. (Leipzig), 3 Vols.

SCHOMBURGK, ROBERT

1838. On the ant tree of Guiana (*Triplaris americana*). Ann. Nat. Hist., **1**, pp. 264-267.

SCHUMANN, K.

1888. Einige neue Ameisenpflanzen. Jahrb. Wiss. Bot., **19**, pp. 357-421, 2 Pls.

1889. Ueber Ameisenpflanzen. Samml. Gemeinverst. Wiss. Vorträge (Hamburg), 38 pp., 1 Pl.
1891. Rubiaceae. In Engler and Prantl, Die Natürlichen Pflanzenfamilien, Vol. 4, pt. 4: 1-156.
- SCHWARZ, E. A.
1901. On the insect fauna of the mistletoe. Proc. Ent. Soc. Washington, 4, pp. 392-394.
1917. Ants protecting *Acacia* trees in Central America. Proc. Ent. Soc. Washington, 18 (1916), p. 211.
- SCOTT, H.
1912. A contribution to the knowledge of the fauna of Bromeliaceae. Ann. Mag. Nat. Hist., (8) 10, pp. 424-431, 433-437.
- SEEMANN, B.
- 1852-1857. The botany of the voyage of H.M.S. Herald under the command of Capt. Henry Kellett during the years 1845-51. (London), 6 + 483 pp., 98 Pls.
- SEVERIANO DA FONSECA, J.
1881. Viagem ad redor do Brazil. (Vol. 1: 325).
- SKWARRA, E.
1930. Ameisen und Ameisenpflanzen im Staate Veracruz (Mexiko). 4. Wandervers. Deutsch. Entom. Kiel: 160-170, 2 Pls.
- 1934a. Oekologische Studien über Ameisen und Ameisenpflanzen in Mexiko. (Königsberg i. Pr.), 153 pp.
- 1934b. Oekologie der Lebensgemeinschaften mexikanischer Ameisenpflanzen. Zeitschr. Morph. Oekel. Tiere, 29, pp. 306-373.
- SMITH, F.
1862. Descriptions of new species of aculeate Hymenoptera, collected at Panama by R. W. Stretch, Esq., with a list of described species, and the various localities where they have previously occurred. Trans. Ent. Soc. London, (3) 1, pp. 29-44.
- SMITH, J. DONNEL
1894. Undescribed plants from Guatemala and other Central American republics, XIII. Botan. Gazette, 19, pp. 255-266.
1895. Undescribed plants from Guatemala and other Central American republics, XV. Botan. Gazette, 20, pp. 281-295.
- SNETHLAGE, E. H.
1923. Neue Arten der Gattung *Cecropia*, nebst Beiträgen zu ihrer Systematik. Notizbl. Bot. Gart. Mus. Berlin-Dahlem, 8, No. 75, pp. 357-369.
- SPIX, J. VON AND MARTIUS, C. VON
1831. Reise in Brazilien in den Jahren 1817 bis 1820. (Vol. 3, pp. 1283-1284, Melastomaceae).

SPRUCE, R.

1908. Notes of a botanist on the Amazon and Andes. (London), Vol. 1, lii + 518 pp.; Vol. 2, xii + 542 pp.

STÄGER, R.

1917. Beitrag zur Kenntnis stengelbewohnender Ameisen in der Schweiz. Rev. Suisse Zool., **25**, pp. 95-109.

STANDLEY, P. C.

1922. Trees and shrubs of Mexico (Fagaceae-Fabaceae). Contrib. U. S. Nat. Herbarium, **23**, pp. 171-515.
1924. Trees and shrubs of Mexico (Passifloraceae-Scrophulariaceae). Contrib. U. S. Nat. Herbarium, **23**, pp. 849-1312.
1927. The flora of Barro Colorado Island, Panama. Smithson. Miscell. Coll., **78**, No. 8, 32 pp.
1928. Flora of the Panama Canal Zone. Contrib. U. S. Nat. Herbarium, **27**, pp. 416, 66 pls.

TIDESTROM, I.

1925. Flora of Utah and Nevada. Contrib. U. S. Nat. Herbarium, **25**, pp. 665, 15 pls.

TULASNE, L. R.

1844. Légumineuses arborescentes de l'Amérique du Sud. Arch. Mus. Hist. Nat. Paris, **4**, pp. 65-196.

ULE, E.

1897. Symbiose zwischen *Asclepias curassavica* und einem Schmetterling nebst Beitrag zu derjenigen zwischen Ameisen und Cecropia. Ber. Deutsch Bot. Ges., **15**, pp. 385-387.
1900. Ueber weitere neue und interessante Bromeliaceen. Ber. Deutsch. Bot. Ges., **18**, pp. 318-327, 1 Pl.
1905. Wechselbeziehungen zwischen Ameisen und Pflanzen. Flora, **94**, pp. 491-497.
- 1906a. Ameisenpflanzen. Engler's Bot. Jahrb., **37**, pp. 335-352, 2 Pls.
- 1906b. Ameisenpflanzen des Amazonasgebietes. In Karsten und Schenck, Vegetationsbilder. Series 4, 14 pp., 6 Pls.
- 1907-1908. Die Pflanzenformationen des Amazonas-Gebietes. Pflanzengeographische Ergebnisse meiner in den Jahren 1900-1903 unternommenen Reisen. Engler's Bot. Jahrb., **40** (1907), pp. 114-172; (1908), pp. 398-443, 8 Pls.

WARBURG, O.

1892. Ueber Ameisenpflanzen (Myrmekophyten). Biol. Centralbl., **12**, pp. 129-142.

WARMING, E.

1894. Om et Par af Myrer beboede Traeer. Vidensk. Meddel. Naturh. Foren. Kjöbenhavn, (5) 5 (1893), pp. 173-187.

WASMANN, E.

1915. Eine neue *Pseudomyrma* aus der Ochsendornakazie in Mexiko. Tijdschr. v. Entom., 58, pp. 296-325, 4 Pls. (Nachtrag, 58, pp. 125-131).

WEDDELL, H. A.

1849. Additions à la flore de l'Amérique du Sud. Ann. Sci. Nat., Bot., (3) 13, pp. 40-113, 249-268.

WERCKLE, C.

1909. La subregión fitogeografica costarricense. Soc. Nac. Agric. Costa Rica, San José.

WHEELER, W. M.

- 1901a. *Microdon* larvae in *Pseudomyrma* nests. Psyche, 9, pp. 222-224.
1901b. Notices biologiques sur les fourmis mexicaines. Ann. Soc. Ent. Belgique, 45, pp. 199-205.
1907. The fungus-growing ants of North America. Bull. Amer. Mus. Nat. Hist., 23, pp. 669-807, 5 Pls.
1908. The ants of Porto Rico and the Virgin Islands. Bull. Amer. Mus. Nat. Hist., 24, pp. 117-158, 2 Pls.
1910. Ants, their structure, development and behavior. (New York), xxv + 663 pp.
1913. Observations on the Central American *Acacia* ants. Trans. 2d. Intern. Entom. Congr. Oxford (1912), 2, pp. 109-139.
1921. A study of some social beetles in British Guiana and of their relations to the ant-plant *Tachigalia*. Zoologica, 3, pp. 35-183, 5 Pls.

WHEELER, W. M. AND BAILEY, I. W.

1920. The feeding habits of Pseudomyrmine and other ants. Trans. Amer. Phil. Soc., (N. S.) 22, pp. 235-279, 5 Pls.

WHEELER, W. M. AND BEQUAERT, J.

1929. Amazonian myrmecophytes and their ants. Zoolog. Anzeiger (Wasmann-Festband), pp. 10-39.

WITTMACK, L.

1888. Bromeliaceae. In Engler and Prantl, Die Natürlichen Pflanzenfamilien, 2, Abt. 4, pp. 32-76 (Addenda: Nachtrag, 1897, pp. 61-69).

WOLCOTT, G. N.

1923. Insectae Portoricensis. Jl. Dept. Agric. Porto Rico, 7, pp. 1-313.

Part II. NEOTROPICAL PLANT ANTS

In the following pages I have endeavored to bring together the scattered information which has long been accumulating in our myrmecological literature concerning the Formicidae inhabiting living plants in the Neotropical Region, together with descriptions and records of the forms recently collected by Dr. W. M. Mann, Dr. J. C. Bradley, Prof. I. W. Bailey, Mr. H. O. Lang and myself. The citations of the literature are restricted to the actual records and do not cover mere taxonomic descriptions. These may be readily found by consulting the various fascicles by Professor Carlo Emery on the Formicidae in the "Genera Insectorum". In some cases, no doubt, the collected records refer to the occurrence of ants in dead twigs and do not properly belong in my list, since their authors sometimes fail to make the necessary distinction. Many species, however, which normally nest in dead twigs may occasionally inhabit cavities in living vegetable tissue. For the same reason I may have omitted a number of species which should have been included. Since, however, the list is presented merely as a convenient basis for future observations, these defects are not to be regarded too seriously. In its present form it gives a definite and suggestive picture of the known neotropical ants which have either already become exquisitely adapted and specialized for living in the various myrmecophytes that occur within their geographical range or are on the road to acquiring such an adaptation.

Subfamily PONERINAE

Genus NEOPONERA Emery

NEOPONERA CARINULATA Roger

British Guiana: Merume Mouth (H. O. Lang), nesting in the fistulose stems of *Patima formicaria* Johnston (Rubiaceae).

NEOPONERA CRENATA Roger

Wheeler, Zoologica **3**, 1921, p. 138.

British Guiana: Kartabo (Wheeler); in hollow petioles of *Tachigalia paniculata* Aubl.

Panama: Barro Colorado Island, Gatun Lake, C. Z. (Wheeler), in cauline swelling of *Cordia gerascanthus* L.

NEOPONERA STIPITUM Forel

Guatemala: Quirigua (Wheeler); in internode of a young *Cecropia*.

British Guiana: Kartabo (Wheeler), nesting in a large branch of *Cecropia sciadophylla* var *decurrens* Snetl.

NEOPONERA UNIDENTATA Mayr

Wheeler, *Zoologica* **3**, 1921, p. 138.

British Guiana: Kartabo (Wheeler); in hollow petioles of *Tachigalia paniculata* Aubl. In the same locality I have also taken this ant on several occasions in the cauline swellings of *Cordia nodosa* L.

NEOPONERA VILLOSA F. Smith

Mayr, *Verh. Zool. bot. Ges. Wien* **16**, 1866, p. 720.

Mexico: Vera Cruz at the foot of Orizaba; in the pseudo-bulbs of an orchid, *Schomburgkia tibicinis*.

Genus ANOCHETUS Mayr

ANOCHETUS (STENOMYRMEX) EMARGINATUS Fabr.

Wheeler, *Ecology* **2**, 1921, p. 96.

British Guiana: Kartabo (Wheeler); in ant-gardens about the roots of epiphytes. This ant usually nests under bark or in the cavities of dead limbs.

Genus ODONTOMACHUS Mayr

ODONTOMACHUS AFFINIS Guérin subsp. MAYRI Mann.

Mann, *Psyche* **19**, 1912, p. 39 ♀.

Brazil: Madeira Mamore R.R., Matto Grosso (W.M. Mann). In ant gardens, living in parabiosis with *Dolichoderus (Monacis) debilis* var. *rufescens*. *O. affinis* usually nests in the ground. Its habits have been studied by Borgmeier. (*Deutsch. Ver. Wiss. Kunst.* **7**, 1920, p. 31-38).

ODONTOMACHUS HASTATUS Fabr.

Calvert, *A Year of Costa Rican Natural History*, 1917, p. 231, fig.

. Costa Rica: (P. P. Calvert), frequently nesting in Bromeliads.

Subfamily PSEUDOMYRMINAE

Genus PSEUDOMYRMA F. Smith

PSEUDOMYRMA ACANTHOBIA Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896, p. 628, ♀ ♀ ♂.

Paraguay: San Salvador (J. Bohls); in woody thorns of *Acacia* (*cavenia*).

Var. FUSCATA Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896, p. 629, ♀ ♀ ♂.

Paraguay: San Salvador (J. Bohls); in woody thorns of *Acacia* (probably *cavenia*).

The other varieties and subspecies of *acanthobia* live by preference in dead twigs or the culms of grasses.

PSEUDOMYRMA ALLIODORAE sp. nov.

Plate 47, Fig. a

Worker:. Length 2.5—2.8 mm.

Head subrectangular, about one and two-thirds times as long as broad, as broad in front as behind, with feebly convex sides and very slightly sinuate posterior border. Eyes very large, elongate-elliptical, flat, more than half as long as the head, somewhat nearer the anterior than the posterior corners. Ocelli well-developed. Mandibles convex at the base, more flattened distally, with five well-developed teeth. Clypeus very short, laterally depressed, its anterior border projecting in the middle as a short, narrow lobe with straight edge and sharp corners. Frontal carinae represented by a short, convex, longitudinal ridge continuous with the clypeal lobe. Frontal groove absent. Antennae short; the scapes scarcely attaining the middle of the head, rather slender, curved, about four times as long as broad; first funicular joint one and one-half times as long as broad; remaining joints, except the last, transverse, nearly twice as broad as long. Thorax rather elongate, the pro- and mesonotum together equal to the epinotum; the promesonotal and mesoepinotal sutures pronounced, the dorsal outline at the latter rather deeply and abruptly notched. Pronotum longer than broad, slightly narrowed behind, bluntly submarginate on the sides; mesonotum rather flat, broader than long, its anterior border bluntly angular, its posterior border rounded. Epinotum nar-

rowed behind, its base feebly convex, nearly twice as long as the straight sloping declivity. Petiole from above elongate-elliptical, broadest in the middle, about one and two-thirds times as long as broad; in profile without a peduncle, nearly as high as long, evenly rounded above and slightly higher just behind the middle, the sides flattened but not marginate above, the anteroventral tooth well-developed, directed downward. Post-petiole less than twice as broad as the petiole, convex and rounded dorsally, ventrally and laterally, slightly broader than long and somewhat narrowed anteriorly. Gaster of the usual shape. Fore femora dilated and convex, somewhat more than twice as long as their median diameter.

Somewhat shining; head and thorax more subopaque. Mandibles opaque, very finely and indistinctly punctulate; head and thorax densely punctate, the punctures on the former distinctly coarser; gaster very finely and superficially punctulate.

Pilosity and pubescence white, the former short, uneven and sparse on the body, absent on the appendages; the pubescence very fine, short and uniform over the whole surface of the body and appendages, rendering them pruinose.

Dark brown; anterior portion of head, gula, sides of pronotum and posterior borders of gastric segments paler, more yellowish brown; mandibles, clypeus, anterior borders of cheeks, antennæ, tarsi and tips and bases of femora and tibiæ yellow.

Female. Length 3-3.2 mm.

Very similar to the worker but with the head longer, fully twice as long as broad, with larger eyes and ocelli. Thorax through the wing-insertions slightly narrower than the head through the eyes. Mesonotum subhexagonal, slightly broader than long, very feebly convex in profile. Pronotum more sharply submarginate on the sides than in the worker; epinotum decidedly shorter and more rounded in profile.

Petiole longer and more distinctly pedunculate. Sculpture, pilosity and color as in the worker, except that the punctuation of the head is finer and the surface, therefore, more shining. Wings hyaline and colorless, with pale yellow veins and pterostigma.

Described from numerous workers and two females. This ant was not infrequently found nesting in the cauline swellings of *Cordia gerascanthus* at Ancon, C. Z., Panama. On one occasion I took it also in a hollow twig of *Triplaris americana* near Miraflores, C. Z. It is rather closely related to *Ps. dolichopsis* Forel, *elongata* Mayr and *subtillissima* Emery, but cannot be referred to any of these species.

PSEUDOMYRMA BELTI Emery

Emery, Bull. Soc. Ent. Ital. **22**, 1890, p. 26, ♀ ♀ ♂, Pl. 6, Fig. 1; Biol. Centralbl. **11**, 1891, p. 165-168; Anal. Mus. Nac. Costa Rica, 1892, p. 66; Wheeler, Trans. 2nd Ent. Congr. Oxford, 1912, p. 115; Wasmann, Tijdschr. v. Ent. **58**, 1915, p. 304; Safford, Smiths. Rep. (1921), 1923, p. 393.

The worker measures 5-6 mm. and is black, with the mandibles, border of mouth, antennae, articulations of legs and tarsi reddish. The body is opaque and densely punctate, the abdomen more finely than the head and thorax, the epinotum and summit of petiole also with superimposed foveolae. The pilosity is pale and sparse, the pubescence grayish, appressed and sericeous. The head is ovate, scarcely longer than broad, the eyes occupying less than half its sides, the ocelli small, the anterior border of the clypeus with a narrow, median, truncated lobe, which is not dentate at the corners. Frontal groove distinct, second funicular joint as broad as long, remaining joints, except last, broader than long. The thorax resembles that of *Ps. spinicola* Emery, but the mesonotum is more convex and projecting and the epinotum is shorter and with a more obtuse angle between the base and declivity. The petiolar node and its peduncle are much shorter, the latter not longer than broad, the former with a small, acute ventral tooth. The postpetiole is broader than long, nearly twice as broad as the petiole, subtriangular, narrowed in front and convex behind.

The female measures 8-10 mm. and resembles the worker in sculpture, color and pilosity, but has the head one and one half times as long as broad, with rather straight, subparallel sides and straight posterior border. Mesonotum shining, with a blackish or dark brown streak on each side. The wings are infuscated, with dark brown veins and pterostigma.

The male was not described, but its head was figured by Emery. It has larger though less convex eyes than *spinicola*.

The species was first taken by A. Alfaro at Alajuela, Jimenez and Siberia, Costa Rica in the thorns of an *Acacia* which Wasmann believes to have been *Acacia spadicigera*, but which was probably *costaricensis*. I possess a cotype worker and numerous workers taken by Dr. P. P. Calvert from *Acacia* thorns at Santa Cruz, Guanacaste, Costa Rica and numerous workers and a female taken by Dr. J. Bequaert in thorns of *Acacia yucatanensis* Schenck at Puerto Castilla, Honduras.

Subsp. *FELLOSA* subsp. nov.

Worker. Length 5–5.5 mm.

Differing from the typical form of the species in having the sides of the pronotum more distinctly submarginate, the petiole and especially its peduncle longer, the latter stouter and with more prominent spiracles, the node higher, more convex and more sharply truncated behind, the postpetiole broader in proportion to its length and more convex. The sculpture, especially of the thorax, is coarser, the foveolae on the epinotum and petiolar node sharp and discrete. The pubescence and pilosity is longer and more conspicuous. The color is the same, except that the thorax and petiolar peduncle are red, with the dorsal surfaces of the pro-, meso- and epinotum, except their borders, black. There is usually some infuscation of the sides of the thorax, especially of the mesopleura. The mandibles, clypeus and cheeks are brownish yellow, the antennae and legs brownish red, with the femora and tibiae, except their tips and bases, dark brown or blackish.

Described from numerous specimens taken by Prof. C. F. Baker at Granada, Nicaragua, in thorns of *Acacia costaricensis*. I have also a number of specimens taken by Mr. William Fluck in another unrecorded Nicaraguan locality and also in *Acacia* thorns.

Subsp. *FULVESCENS* Emery

Emery, Bull. Soc. Ent. Ital. **22**, 1890, p. 64, ♀ ♀; Biol. Centralbl. **11**, 1891, p. 167; Anal. Mus. Nac. Costa Rica, 1892, p. 66, *nota*; Wheeler, Trans. 2nd. Intern. Congr. Ent. Oxford, 1912, p. 117; Forel, Mém. Soc. Ent. Belg. **20**, 1912, p. 22 ♀; Safford, Smithson. Rep (1921), 1923, p. 390, figs. 5 & 6 & Pl. 3.

Worker. Length 3.5–5.5 mm.

Head longer than in the preceding subspecies, fully one and one-third times as long as broad, the punctation finer so that the surface is more shining, the pilosity and pubescence somewhat sparser. Thoracic dorsum straight in profile, mesonotum rather flat; petiole and postpetiole much like those of the typical *belti*, but the postpetiole is somewhat narrower; the petiolar node varies, being as broad as long in some specimens and somewhat longer than broad in others. Yellowish or ferruginous red, the postpetiole, gaster and legs, except their articulations, usually darker.

Female. Length 6.5–8 mm.

Head fully one and two-thirds times as long as broad, parallel-sided,

the eyes more than twice as long as broad, flattened. Mandibles robust, with convex surfaces and external borders. Epinotum long, its declivity rather abrupt, shorter than the base. Sculpture, pilosity and color much as in the worker; foveolation of epinotum coarser. Mandibles red, clypeus, antennae and anterior third of head yellow; sides of mesonotum and sometimes the mesopleura castaneous; ocellar region infuscated. Wings grayish hyaline or slightly brownish, with brown veins and pterostigma.

Male. Length 6.5–8 mm.

Head, including the eyes, distinctly longer than broad, narrowed behind, with rather straight posterior border. Eyes about half as long as the sides of the head. Antennal scapes twice as long as broad; first funicular joint longer than broad. Epinotum rather long, sloping, but with distinct base and declivity. Surface of body more shining and more finely punctate than in the worker and female. Dark brown; mandibles, clypeus, scapes, first funicular joint and tarsi whitish yellow; sides of pronotum, sutures of thorax, bases and tips of femora and tibiae and the borders of the gastric segments yellowish brown. Wings like those of the female.

The types of this subspecies were found by Beccari in cauline swellings of *Cordia gerascanthus* L. from Guatemala, but the occurrence of the ant in this plant must be very exceptional, since it is the commonest and most widely distributed obligate of the various bull-thorn Acacias in Central America and Mexico. I give here a list of the localities from which I possess specimens:

Guatemala: Escuintla, Patulul, Zacapa and Quirigua (Wheeler), in thorns of *Acacia hindsii* Benth., *Acacia* sp. *bursaria* Schenck; Trece Aguas, Alta Vera Paz (Schwarz & Barber).

British Honduras: Manatee (J. D. Johnston), in thorns of *Acacia* sp.; Belize (C. F. Baker), in thorns of *Acacia* sp.

Mexico: Tampico (Ed. Palmer, D. L. Crawford, H. Jourdain); Tonola, Chiapas and Los Cocos, Vera Cruz (A. Petrunkevitch), in thorns of *Acacia* sp.; Santa Lucrecia, Vera Cruz (F. Knab); Julapa (Rangel), in thorns of *Acacia sphaeracephala*; San Sebastian, Chiapas (G. N. Collins), in thorns of *Acacia Collinsii* Saff. Pichucalco, Chiapas (G. N. Collins), in thorns of *A. cornigera* L.; Llano Grande, Chiapas (G. N. Collins), in thorns of *A. hindsii*. San Luis Potosi, probably in thorns of *A. hernandesi* Safford.

The subspecies *fulvescens* is also recorded by Dr. Safford from the following localities in Mexico: Tanquian, San Luis Potosi (L. G. Cuevas), in thorns of *Acacia sphaeroccephala*; Tampico (J. M. Cuaron)

in thorns of the same *Acacia*; San Sebastian, near Tuxtla (G. N. Collins) in thorns of *Acacia collinsi*.

Subsp. SAFFORDI subsp. nov.

Worker. As large as *fulvescens* and with the head of the same proportions, but with the coarse opaque sculpture of the typical *belti*, the petiole, especially its peduncle shorter than in either of these forms, the node slightly broader than long, the postpetiole also broader than long but only one and two-thirds times as broad as the petiolar node. Mesonotum rather convex and prominent. Dark brown; anterior portion of head, borders of orbits, two longitudinal streaks on the gula, pronotum, sides of petiole, tibiæ and tarsi paler, reddish brown.

Nine workers taken by Mr. G. N. Collins from thorns of *Acacia collinsi* Saff. at Chicoasen, Chiapas, Mexico, and received from Dr. Safford. Three small workers and an immature and imperfect female taken by Mr. Collins at Yerba Lanta, Chiapas in the same species of *Acacia* seem to belong to the same subspecies, although the workers are more uniformly dark brown. I also refer to this subspecies a series of workers from the Department of Solola, Guatemala, 3,000 feet. They have the top of the head black, but the thorax and petiole are uniformly deep brownish red.

Subsp. VENEFICA subsp. nov.

Worker. Length 3.5-4 mm.

Decidedly smaller than the subsp. *felloso*. Head of the same proportions. Epinotum longer, the base very convex and decidedly longer than the declivity into which it passes through a convex curve, without an angle. Thoracic sutures very deep and broad. Petiole as in *felloso* but the node slightly longer than broad; postpetiole a little broader than long, twice as broad as the petiolar node. Sculpture and pilosity as in *felloso*. Very dark brown, nearly black; clypeus and frontal carinae yellow; mandibles, antennae, thoracic sutures, posterior and lateral borders of pronotum, tarsi, articulations of legs and posterior borders of gastric segments pale reddish brown.

Female. Length 5 mm.

Head short, less than one and one half times as long as broad, broadest behind the eyes, narrower through the cheeks, which are straight and subparallel. Eyes less than twice as long as broad, feebly convex. Sculpture, pilosity and color as in the worker, but the pron-

tum, except its dorsal disc, the whole of the scutellum, the sides and posterior portion of the epinotum and the petiole, except the dorsal surface of its node, yellowish brown. Wings grayish hyaline, not infuscated; veins pale, the pterostigma dark brown.

Male. Length 4.-4.5 mm.

Head as broad as long, rounded but not narrowed behind. Eyes convex, more than half as long as the sides of the head. Scapes nearly two and one half times as long as broad; first funicular joint longer than broad. Epinotum low and sloping, not very convex. Postpetiole a little longer than broad, behind nearly three times as broad as the node of the slender petiole. Sculpture, pilosity and color like those of the worker, but the head and thorax more shining and more finely punctate; the antennae and legs infuscated; the wings as in the female.

Described from numerous specimens of all the phases taken by Mr. J. H. Batty in the thorns of *Acacia sinaloensis* Saff. at Escuinapa, Sinaloa, Mexico. Another series of workers and deälated females taken ay Mr. C. H. Tyler Townsend in thorns of *A. Hindsi* at Manzanillo, Colima, Mexico, may be referred to the same subspecies, though the head of the female is distinctly longer and the pale portions of the thorax and petiole are darker. Of this same subspecies, which is characterized by the small size of all the phases, I have a male, female and worker taken by Prof. C. F. Baker at Acapulco, Mexico.

Subsp. VESANA subsp. nov.

Worker. Length 3.5 mm.

Characterized by its small size, pale color and the shape of the pedicel. The petiole is very short, there is almost no peduncle and the node is high, triangular when seen from above, as broad as long; the postpetiole is nearly twice as broad as long and twice as broad as the petiolar node; rather abruptly narrowed anteriorly. Base and declivity of epinotum subequal, straight, the latter sloping. Head a little longer than in the subspecies *venefica*, punctation of head and thorax somewhat finer, foveolæ of epinotum and petiolar node less distinct. Yellowish red; mandibles and anterior portion of head yellow; legs slightly brownish, postpetiole and gaster dark brown.

A single specimen taken by Mr. Fred Knab at Cordoba, Mexico, probably on some species of *Acacia*. This form is obviously transitional to the subsp. *fulvescens* in color and sculpture, in the structure of the pedicel, but not in the shape of the head.

Subsp. BEQUAERTI subsp. nov.

Worker. Length 3.5–4.5 mm.

Head distinctly shorter than in the other forms of the species, more rounded and elliptical. Sides of pronotum scarcely submarginate. Petiole with well-developed peduncle, the node from above triangular, as long as broad, postpetiole nearly as long as broad. Surface somewhat shining and lustrous; the foveolae on the epinotum and petiolar node more indistinct than in the other subspecies. Dark brown, sutures of thorax, petiole and postpetiole, the tarsi, antennæ, greater portion of tibiæ, mandibles and anterior half of head reddish; posterior half of head feebly infuscated.

Numerous workers taken by Dr. Joseph Bequaert in thorns of *Acacia yucatanensis* Schenck at Puerto Castilla, Honduras.

Subsp. WASMANNI Wheeler

Pseudomyrma canescens Wasmann (*nom. praeocc.*) Tijdschr. Ent. **58**, 1915, p. 297 et seq., ♀ ♂; *Pseudomyrma wasmanni* Wheeler, Ecology **2**, 1921, p. 92, *nota*.

I have not been able to recognize this form among my material. Wasmann described and figured it from specimens collected by W. Brakhoven in thorns of *Acacia sphaerocephala* at Tampico, Mexico, and regarded it as an independent species. He did not compare it with the subsp. *fulvescens*, but with the typical *belti* and hence made much of the length of the head. The workers observed by Wasmann had the sides of the head sub-parallel, which is not true of any of the forms of *belti* known to me. The other characters mentioned in his description may nearly all be observed in *fulvescens*, which is rather variable. The measurements given are: Worker: 5.5–6.5 mm; Female: 7–8 mm.; Male: 7–8 mm. There is nothing in Wasmann's figures to show that his specimens will not admit of the interpretation I have given.

PSEUDOMYRMA CAROLI Forel

var. SAPII Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904, p. 688, ♀ ♀.

Brazil: Bom Fim, Jurua, Amazonas (E. Ule), in the perforated stems of *Sapium* (Euphorbiaceae).

The typical *caroli* was described from Costa Rica. I have taken it at Escuintla, Guatemala and have received many specimens of it from Manatee, British Honduras (J. D. Johnston).

PSEUDOMYRMA CHODATI Forel

Forel, Bull. Soc. Bot. Genève (2) **11**, 1920, p. 1, ♀.

Paraguay: (R. Chodat), in the cauline swellings of *Cordia longituba* Chodat.

PSEUDOMYRMA DAMNOSA Wheeler

Wheeler, Zoologica **3**, 1921, p. 139, ♀ ♀ ♂, Fig. 13. Bailey, Bot. Gazette **75**, 1923, p. 34.

British Guiana: Kartabo, Kalacoon and Penal Settlement (Wheeler); in hollow petioles of *Tachigalia paniculata* Aublet.

PSEUDOMYRMA DENDROICA Forel

Forel, Rev. Suisse Zool. **12**, 1904, p. 40, ♀ ♂; C. R. 6e Congr. Intern. Zool. Berne (1904), 1905, p. 452.

Brazil: Rio Purus, Amazonas (A. Goeldi), in hollow branches of *Triplaris*.

Var. EMARGINATA Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904, p. 684, ♀ ♀; C. R. 6e Congr. Intern. Zool. Berne (1904), 1905, p. 452.

Brazil: Marary Jurua, Amazonas (E. Ule), in hollow twigs of *Triplaris schomburgkiana* Benth.

PSEUDOMYRMA ELONGATA Mayr

Wheeler, Bull. Amer. Mus. Nat. Hist. **21**, 1905, p. 87, ♀ ♀ ♂.

Florida: Cards' Point (Wheeler), nesting in *Tillandsias*.

Var. TANDEM Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906, p. 228, ♀.

Costa Rica: El Hiquito, San Mateo (P. Biolley), in the trunk and fruit of *Bixa orellana*.

PSEUDOMYRMA FIEBRIGI Forel

Forel, Verh. Zool. bot. Ges Wien 1908, p. 383, ♂ ♀; Fiebrig, Biol. Centralbl. **29**, 1907, p. 68; Chodat and Carisso, Bull. Soc. Bot. Genève **12**, 1920, p. 189, fig. 324, 325.

Paraguay: San Bernardino (K. Fiebrig), in thorns of *Acacia cavenia* H. & A.

PSEUDOMYRMA GEBELLI Forel

Panama: Las Sabanas (Wheeler), in Tillandsias growing on manzanillo trees (*Hippomane mancinella* L.) also in branches of almendra (*Terminalia catappa*), at Ancon C. Z.

PSEUDOMYRMA GRACILIS Fabr.

Wheeler, Trans. 2nd Intern. Ent. Congr. Oxford, 1912, p. 118.

Guatemala: Quirigua (Wheeler), nesting in thorns of *Acacia*.

Panama: Chivachiva Trail, near Red Tank, C. Z. (Wheeler); Tumba Muerta Road, near Las Sabanas (Wheeler), in thorns of *Acacia penonensis* Saff.

Bolivia: Riberalta (W. M. Mann), in cauline swellings of *Cordia gerascanthus* var.

As a rule this ant nests in dead twigs. This is also true of the following variety.

Var. BICOLOR Guérin

The workers are of the same stature as the typical *gracilis*, black, with the mandibles, clypeus, cheeks and frontal carinae yellow, the thorax and petiole dull orange, with the mesonotum, a large spot and streak on the base of the epinotum, two elongate spots on the pronotum and the ventral border of the pleura black. The legs are variable, usually black, with the tips of the femora and tibiae yellow, streaked with black on the extensor surfaces. Some of the specimens lack the black spots on the pronotum and some have merely the peduncle of the petiole and the pronotum somewhat reddish and thus form a transition to the true *gracilis*. The variety resembles the subsp. *mexicana* Roger but is more slender, with the same stature as the typical *gracilis*.

Panama: Chivachiva Trail, near Red Tank, C. Z. (Wheeler), in cauline swellings of *Cordia gerascanthus* (common). Same locality in

the hollow stems of *Triplaris americana*. Tumba Muerta Road, Las Sabanas (Wheeler); in fistulose stems of *Clerodendron siphonanthus* (introduced from the Orient.)

Subsp. MEXICANA Roger

Emery, Biol. Centralbl. **11**, 1891, p. 167; Anal. Mus. Nac. Costa Rica 1892, p. 67; Wheeler, Amer. Natural. **35**, 1901, p. 527; Ann. Soc. Ent. Belg. **45**, 1901, p. 204.

Costa Rica: (A. Alfaro), in thorns of Acacia.

Mexico: Cuernavaca (Wheeler), in *Tillandsia Benthamiana*. In internodes of bamboo. Costa Rica (H. Schmitt).

PSEUDOMYRMA KUENCKELI Emery

Emery, Bull. Soc. Ent. Ital. **22**, 1890, p. 62; Anal. Mus. Nac. Costa Rica 1892, p. 67; Wheeler, Ann. Soc. Ent. Belg. **45**, 1901, p. 203.

This ant was originally described from Alajuela, Costa Rica, and as nesting in wood. In his paper of 1892, Emery mentions that he received from A. Alfaro one specimen taken in an Acacia thorn. The further fact that I took this *Pseudomyrma* in 1900 at Cuernavaca, Mexico, running in considerable numbers on the trunk and branches and living in the dry twigs of small acacias not of the bull-horn type, but probably *A. farnesiana*, shows that unlike *Ps. belti* and *spinicola*, it is only an occasional tenant of the bull-horn Acacias.

PSEUDOMYRMA LATINODA Mayr

Mayr, Verh. Zool. bot. Ges. Wien 1877, p. 877, ♀.

The type of this species was taken by Prof. James Trail in Northern Brazil, probably in some myrmecophyte (*Triplaris* or *Tachigalia*). Prof. Forel gave me a specimen from Sarayacu, Brazil, which he regarded as belonging to the typical form and which agrees well with Mayr's description. Probably Forel's *dendroica* and my *maligna* and *damnosa* may be eventually regarded as subspecies of *latinoda*, but for the present I treat them as distinct species to avoid introducing more confusion into a sufficiently intricate congeries of closely related forms.

Var. CORONATA var. nov.

Worker. Length 5—5.5 mm.

Larger and more robust than other forms of the species. Black, with the head reddish yellow and with the same peculiar markings on

the occiput as in var. *nigrescens* (*vide infra*), only more distinct. Mandibles red, blackish at the base. Antennæ and tarsi reddish yellow, somewhat paler than the head; legs dark brown or blackish, with the knees and posterior borders of the gastric segments reddish. Form of body, sculpture and pilosity very much as in *nigrescens* and with the funicular joints 2-10 very nearly as short and transverse.

Male. Length 6 mm.

Head and body of the usual shape; eyes half as long as the sides of the head; antennal scapes a little less than twice as long as broad; funiculi short, first joint a little longer than broad, remaining joints, except the last, not more than twice and mostly less than twice as long as broad. Epinotum rounded and sloping, without distinct base and declivity. Petiole and postpetiole much as in the worker.

Very smooth and shining, sparsely punctate; ocellar region subopaque, finely and densely punctate; sides of thorax indistinctly rugulose.

Pilosity like that of the worker, but somewhat less abundant and less even.

Piceous black, including the genital valves; mandibles, clypeus, anterior portion of head, antennæ, tarsi, articulations of legs, thoracic sutures, portions of pleura and borders of gastric segments, sordid or brownish yellow. Wings distinctly infuscated, with dark brown veins and pterostigma.

Described from numerous workers and four males taken by Mr. H. O. Lang at Kamakusa, British Guiana in the hollow petioles of *Tachigalia paniculata* Aublet.

Var. ENDOPHYTA Forel

Forel, Mém. Soc. Ent. Belg. **20**, 1912, p. 22, ♀.

Taken by A. Ducke on the Rio Ariramba, near the Rio Trombetas, Amazonas, Brazil, in the hollow leaf-petioles of *Tachigalia macrostachya* Huber. The worker measures 4.1-5.3 mm. and is therefore larger than the typical *latinoda* which measures only 4 mm., according to Mayr. *Endophyta* is entirely yellowish red, with a brownish spot in the ocellar region.

Var. NIGRESCENS Forel

Forel, Rev. Suisse Zool. **12**, 1904 p. 38 ♀.

The worker of this variety was taken by Prof. E. A. Goeldi at Para, Brazil, probably in some myrmecophyte (*Tachigalia*?). It averages

about the same size (4.4-4.8 mm.) as *endophyta*, but has the thorax, abdomen, femora and anterior surfaces of tibiae brown, with the mandibles, head, tarsi and posterior surfaces of the tibiae and sometimes also the anterior portion of the pronotum, reddish yellow. There is a brown spot covering the ocelli and connected with a transverse band of the same color on the occiput. This band sends off a longitudinal branch towards or to the posterior orbit of each eye.

Subsp. BRADLEYI subsp. nov.

Worker. Length 3.5—4 mm.

Differing from the subsp. *tachigalia* Forel (*vide infra*) in having the head much less deeply excised behind, with more nearly straight and parallel sides, eyes somewhat smaller, one-third as long as the sides of the head, the frontal groove much more distinct and becoming very deep and broad in front of the anterior ocellus. Mesoepinotal impression decidedly deeper, the petiolar node much thicker in front, scarcely longer than broad, subcuboidal, trapezoidal from above, its ventral surface flat and unarmed. Postpetiole from above semicircular, twice as broad as long. Fore femora slightly more dilated than in *tachigalia*.

In sculpture, pilosity and pubescence very similar to *tachigalia* but the vertex of the head somewhat more shining.

Color much darker, dark brown, with the gaster blackish; pro- and mesonotum somewhat paler; mandibles, clypeus, anterior third of head and tarsi dull whitish yellow; antennae, femora and tibiae dark brown, knees and bases of funiculi paler.

Described from nine workers taken by Prof. J. C. Bradley at Perené, Peru, in the hollow petiole of a *Tachigalia*.

Subsp. OPACIOR Forel

Ps. latinoda var. *opacior* Forel, Ann. Soc. Ent. Belg. 48, 1904 p. 170 ♀.

This form was described as a mere variety from Cuba (Coll. Ballion), but it evidently deserves to rank as a subspecies. The worker measures 4.3 mm. and is uniformly dull yellowish brown, with paler antennae and legs. The head is densely punctate and subopaque, the punctuation of the remainder of the body is also denser and coarser than in the other forms of the species. The petiolar node is broader than long. The head is more narrowed anteriorly and the anterior slope of the pronotum is less abrupt, the epinotum less rounded and more angular than in the typical *latinoda* and the var. *nigrescens*, the only forms with which Forel compares it.

Subsp. TACHIGALIE Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 686 ♀ ♀ ♂.

Forel described all three phases of this ant from specimens taken by E. Ule at Tarapoto, Amazonas, Peru, in the hollow petioles of *Tachigalia formicarum* Harms.

The worker measures 4.5-5 mm. and has the head longer and narrower than in the preceding forms, with the posterior border rather broadly and deeply excised. The petiolar node is lower and has two teeth on the ventral side. Funicular joints, except the first and last, very short and transverse. Body and legs only feebly shining and rather densely punctate, with dense yellowish pubescence, especially on the gaster. The color is a dirty yellowish brown, the anterior portion of the head, mandibles, antennæ, tarsi, borders of gastric segments, articulations and the nodes of the pedicel in part yellowish.

The female measures 9-9.5 mm. and has a longer head than the worker, the mandibles geniculate at the base and marginate externally, with two apical and no basal teeth. The clypeus has a prominent anterior lobe and median carina. The petiolar node is one and one-half times as long as broad and scarcely broader behind than in front. The surface of the body is much more shining than in the worker, rather coarsely and densely punctate. Brownish black or blackish brown, with the cheeks, clypeus, antennal foveæ, antennæ, tibiæ, tarsi, articulations and borders of the gastric segments yellowish red or yellowish. Wings tinged with blackish brown, with blackish brown veins and pterostigma.

The male measures 6.2-6.7 mm. The clypeus has a concave border but no median carina. The funicular joints are short, not twice as long as broad, the scape being of about the same length. Eyes small, shorter than their distance from the posterior corners of the head. Sculpture, pilosity and color as in the female, but the membranes of the wings are yellowish, the legs almost entirely yellow, the funiculus, except its first joint, brown.

PSEUDOMYRMA MALIGNA Wheeler

Wheeler, Zoologica 3, 1921 p. 143 ♀ ♀ ♂; Bailey, Bot. Gazette 75, 1923 p. 34.

British Guiana. Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

Var. CHOLERICA Wheeler

Wheeler, Zoologica 3, 1921 p. 146 ♀; Bailey, Bot. Gazette 75, 1923 p. 34.

British Guiana. Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

Var. CRUCIANS Wheeler

Wheeler, *Zoologica* **3**, 1921 p. 147 ♀; Bailey, *Bot. Gazette* **75**, 1923 p. 34.

British Guiana. Kartabo (Wheeler) in hollow petioles of *Tachigalia paniculata* Aubl.

PSEUDOMYRMA NIGROCINCTA Emery

Emery, *Bull. Soc. Ent. Ital.* **22**, 1890 p. 64 ♀ ♀, Pl. 6 Fig. 3; *Biol. Centralbl.* **11**, 1891, p. 166; *Anal. Mus. Nac. Costa Rica* 1892, p. 66; Wheeler, *Trans. 2nd Intern. Ent. Congr. Oxford* 1912, p. 118; Wasmann, *Tijdschr. Ent.* **58**, 1915 p. 304.

Costa Rica. Alajuela and Jimenez (A. Alfaro), in thorns of *Acacia* (*spadicigera*, according to Wasmann); Guanacaste and Turrucares (P. P. Calvert), in thorns of *Acacia* sp.

PSEUDOMYRMA NIGROPILOSA Emery

Emery, *Bull. Soc. Ent. Ital.* **22**, 1890 p. 62 ♀ Pl. 5 fig. 24; *Biol. Centralbl.* **11**, 1891 p. 168; *Anal. Mus. Nac. Costa Rica* 1892 p. 67; Wheeler, *Trans. 2nd Intern. Ent. Congr. Oxford* 1912 p. 116.

Costa Rica (A. Alfaro); in thorns of *Acacia*; Guanacaste, Santa Cruz (P. P. Calvert), in thorns of *Acacia*.

This ant more frequently nests in dead twigs.

PSEUDOMYRMA PICTA Stitz

Stitz, *Deutsch. Ent. Zeitschr.* 1913 p. 209 ♀ Fig. 2.

This species was described from worker specimens taken by E. Ule at Alto Acre, Brazil, in hollow petioles of *Tachigalia*.

It measures 5.5 mm. and resembles *Ps. sericea* Mayr in form but is more robust and the head is much larger, flattened, as broad as long, with convex sides and straight posterior border. It has two elongate diverging impressions on the front for the scapes, which reach a little beyond the middle of the head. The clypeus is strongly carinate and has a narrow median lobe, with straight entire border. The frontal groove is distinct and terminates in a deep longitudinal impression in front of the anterior ocellus. The eyes are large, elongate, two-thirds as long as the sides of the head. The thorax is much like that of *sericea* but broader, with the pronotum and base of epinotum more flattened and more sharply submarginate on the sides. The petiole and postpetiole are also similar to those of *sericea*, the postpetiole, according to Stitz being broader than the pronotum, with convex posterior border.

Slightly shining; surface covered rather uniformly with very short dense pubescence, which is somewhat longer on the gaster. Hairs sparse, much as in *sericea*, longest and most conspicuous on the abdomen.

Ferruginous yellow, with the anterior portion of the head paler; mandibles sordid brown, with blackish brown dental border. The body is spotted with black as follows: "Deepest portion of frontal groove with a short, black, longitudinal spot; on each side of the vertex, between the lateral ocellus and posterior inner orbit a narrow longitudinal stripe on each side; on the lower surface of the head behind the eye a broad longitudinal band reaching to the occipital border, most strongly developed under the eye; a very strong median band along the gula to the edge of the occiput. The pronotum has on the transverse ridge (separating its anterior and posterior faces) a round spot on each side, sometimes also one or two spots on its sides. The edges adjoining the mesonotal sutures are marked by a black band. The black color is more extensive on the epinotum; its basal surface is black, also the adjacent portion of the declivity. Among the spots on the epinotum the most prominent is an elongate ring-shaped spot, which runs obliquely from the middle of the lateral surface to the insertion of the petiole. The anterior surface of the petiolar node has three black longitudinal bands, the two lateral of which skirt its borders but pass over on the lateral surface and are broadest behind, where like the median band, they partially extend over onto the posterior surface. The gastric tergites and sternites have broad black borders. The coxae are usually dark. The middle and hind femora and tibiae have in the extensor surface at the distal end a longer or shorter, blackish brown stripe; the tarsal joints of these legs are brown with only their tips yellow."

Var. HETEROGYNA Wheeler and Mann, var. nov.

Worker. Length 4.5—5 mm.

Smaller than the type but very similar in sculpture, pilosity, color and the shape of the petiole, but the black spots are more variable, the large one in the base of the epinotum often reduced to a median longitudinal band. The postpetiole is shorter and not as broad as the pronotum and with a somewhat concave posterior border. The petiolar node is stouter and shorter, not longer than broad, its anterior surface rather flat, convex only above where it joins the abrupt posterior surface. The second funicular joint is as long as broad, the succeeding joints distinctly broader than long.

Female (deälated). Length 9 mm.

Head large, distinctly longer than broad and strongly rectangular, except in front of the eyes, where it is narrowed. Eyes more than half as long as the head. Mandibles strongly convex. Antennal scapes not reaching to the middle of the head; funicular joints as long as broad. Thorax stout, with convex mesonotum, the promesonotal suture very deeply impressed. Epinotum short, subcuboidal, with distinct base and declivity. Petiolar node subcuboidal, nearly square from above, but slightly narrowed in front. Postpetiole shaped as in the typical *picta* worker, with convex posterior border, its ventral surface very convex anteriorly. Gaster as in the worker, with pointed tip.

Black; tip of gaster, mandibles, anterior two-fifths of head, cheeks, antennæ, fore tibiæ and tarsi and terminal joints of middle and hind tarsi reddish yellow; knees and posterior borders of gastric segments reddish. The light and dark portions of the upper surface of the head are separated by a sharp, straight, transverse line. The pilosity and pubescence are much as in the worker, the punctuation more distinct, the surface somewhat more shining.

Described from eight workers and a single female taken by Dr. W. M. Mann at Cavinás, Bolivia, in the hairy cauline swelling of a species of *Platymischium*.

Subsp. *CASTA* subsp. nov.

Worker. Length 4.5—5 mm.

Uniformly opaque and ferruginous, with the extreme bases of the gastric segments blackish, a longitudinal blackish spot on the base of the epinotum and in one specimen with two black spots on the pronotum; the mandibles and anterior portion of the head yellowish, the antennal funiculi brown. Scapes reaching a little beyond the middle of the head, joints 2-4 of the funiculi as long as broad, the succeeding joints shorter. Base and declivity of epinotum subequal, the former flattened and distinctly submarginate on the sides like the pronotum. Petiolar node from above not longer than broad, stouter anteriorly than in the typical *picta*, in profile much more convex anteriorly than in both the preceding forms, the line of juncture of the antero-dorsal and posterior surfaces straight and transverse, rather sharp, the lateral surfaces flat, marginate above as in the other forms of the species. Postpetiole convex, shaped much as in *heterogyna* and bearing the same relations in size to the petiolar node.

Described from nine specimens taken by Prof. J. C. Bradley at La Sombre, Putumayo, Peru, in the hollow petioles of *Tachigalia*.

PSEUDOMYRMA SATANICA sp. nov.

(Plate 47, fig. b)

Worker. Length 5.5—6 mm.

Closely related to *spinicola* (*vide infra*) but somewhat larger. Head slightly longer than broad, somewhat depressed or flattened, broadest through the eyes, as broad through the well-developed posterior corners as at the clypeus, occipital border straight. Eyes larger and more convex than in *spinicola*, ocelli well-developed. Mandibles rather flat, with 6 subequal teeth, the external borders rather straight. Clypeus very short, convex in the middle, laterally depressed, the anterior border with a very short lobe which is irregularly excised in the middle and has on each side a triangular, flattened and not very acute tooth. Frontal carinae continued backward as longitudinal swellings, much as in *spinicola* but including a distinct frontal groove which terminates in a short, deep, longitudinal impression in front of the anterior ocellus. Antennae stout, scapes reaching nearly to a line joining the posterior orbits; second funicular joint as long as broad, remaining joints, except the last, distinctly broader than long. Thorax and pedicel much as in *spinicola*, but the mesonotum more convex and projecting and the epinotum higher, with subequal base and declivity, the posterior corners of the petiolar node more acute, subdentate, the postpetiole less narrowed and more convex in front, the tooth on the ventral side of the peduncle lacking.

Subopaque; abdomen more shining than the head and thorax; punctuation coarser than in *spinicola* and its subspecies. Mandibles rather opaque, more coarsely striato-punctate.

Pilosity yellowish, less abundant, pubescence grayish, conspicuously longer and more abundant than in *spinicola*, well-developed, even on the thorax.

Black; postpetiole and gaster dark brown, the posterior margins of the segments of the latter somewhat reddish; mandibles, clypeus and insertions of scapes brownish yellow; antennae, lateral borders of pronotum and legs reddish brown, the median portion of the scapes, femora and tibiae darker and more blackish.

Female. Length 8—8.5 mm.

Very similar to the worker, except in the shape of the head, which is subrectangular, nearly one and three-fourths times as long as broad, with nearly parallel sides and straight posterior border. Eyes large and elongate, more than two-fifths as long as the sides of the head. Clypeal lobe more rectangular and longer than in the worker, frontal

groove distinct but not terminating behind in an impression. Mandibles robust, rather flattened at the tips, with rounded, convex external borders. Thorax shaped much as in *spinicola* subsp. *scelerosa*, but the epinotum is shorter and more rounded. Posterior corners of petiolar node large, prominent, subdentate and somewhat turned outward. Postpetiole longer than broad, nearly parallel-sided behind, rounded and narrowed in front.

Sculpture, pilosity, pubescence and color much as in the worker, but the punctuation coarser, especially on the epinotum and petiolar node and the pubescence on the head and gaster longer. Wings deeply infuscated as in *spinicola* subsp. *scelerosa*, with dark-brown veins and pterostigma.

Male. Length 6.5—7 mm.

Very similar to the male of *spinicola* subsp. *scelerosa*, but the head is longer and more produced behind the eyes, the latter more convex, the first funicular joint only as long as broad, the epinotum less convex and more sloping, the petiole more convex above. Wings scarcely paler than in the female.

Described from numerous specimens taken near the headwaters of the Rio Agua Salud, C. Z., Panama, and Marajal, near Colon, C. Z. They were inhabiting the large straight thorns of two superb trees about 12 feet high of *Acacia multiglandulosa* Schenk. The ants are extremely aggressive, fiercer even than *belti* and *spinicola* and their stings are more painful. I should have attached this form as a subspecies to *spinicola* were it not that its behavior is quite different. Instead of advancing to the attack like *belti* and *spinicola*, with the gaster turned forward under the thorax, it keeps its body rigid with the gaster extended as in *Ps. gracilis* Fabr. and turns it forward to sting only after seizing the intruder with its mandibles. *Multiglandulosa* is one of the most beautiful of the bull-horn acacias. Its foliage is heavy, spreading and bright green, the large leaves each with 8-15 pinnæ, the pinnules when young tipped with golden yellow food-bodies. There are numerous extrafloral nectaries on the base of the petiole and one between each pair of pinnæ. The handsome thorns, when young, are bright crimson.

PSEUDOMYRMA SERICEA Mayr

(Plate 48, fig. a)

Forel: Zool. Jahrb. Abt. Syst. 20, 1904 p. 691 ♂ ♀; Ann. Soc. Ent. Belg. 50, 1906 p. 230.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in the swollen flower axes and twigs of *Pterocarpus ulei* Harms. Bom Lugar, Rio Purus, Amazonas (J. Huber and A. Goeldi), in the hollow twigs of *Bombax mungaba*.

Var. ACACIARUM var. nov.

Worker. Length 3—3.5 mm.

Closely related to the var. *ita* Forel of Costa Rica, but smaller. The petiole and postpetiole are similar but the dorsal surface of the former is even less convex and distinctly marginate in the sides and behind, the posterior surface sharply truncated and flat. Seen from behind the sharp angle at which these surfaces meet is straight and transverse. Postpetiole nearly twice as broad as the petiolar node. Thorax and head much like those of the var. *ita*.

Pubescence white, sericeous, much more abundant than in *ita*, so that the surface is pruinose.

Black; mandibles, clypeus, antennæ, sides of pronotum, neck, fore tibiæ and tarsi and tips of fore femora brownish yellow; a fuscous spot on each mandible, the extensor surface of the funiculus and a long black spot on the tip of the scape black.

Female. Length about 4.3 mm.

Head fully one and one-half times as long as broad. In other respects much like the worker. Wings whitish hyaline, with colorless veins and pale brown pterostigma.

Male. Length 4—4.5 mm.

Head longer than broad, elliptical. Eyes nearly two-thirds as long as its sides. Mandibles long, well-developed, deflected at the tip, their straight apical borders minutely denticulate. Scapes one and one-half times as long as broad, first funicular joint as long as broad. Thorax narrow, the dorsal surface in profile rather straight, sloping backwards, epinotum with subequal base and declivity. Petiole and postpetiole similar to those of the worker, the nodes lower and the surfaces more rounded.

Black; mandibles, clypeus, antennæ, coxae, legs and pronotum ivory yellow, the mandibles with a large black spot in the middle; clypeus infuscated in the middle, antennæ at the tips; the pronotum with a dark brown transverse streak on the dorsal surface, femora feebly infuscated in the middle. Wings colored as in female.

Several colonies of this ant were found nesting in the thorns of *Acacia penonomensis* Saff. along the Tumba Muerte Road, near Las Sabanas, in the Panamanian Republic.

Var. *CORDIÆ* Forel

(Plate 48, fig. b)

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 690 ♀ ♀; Bull. Soc. Bot. Genève (2) **11**, 1920 p. 2 ♀ ♂.

Peru: Tarapoto (E. Ule) in the cauline swellings of an undetermined *Cordia*; Eastern Peru (Spruce 3932), in cauline swellings of *Cordia gerascanthus* forma *micrantha*.

Bolivia: (Bang) in cauline swellings of *Cordia* sp.

A number of workers and three deālated females were taken by Dr. W. M. Mann in cauline swellings of *Cordia gerascanthus* var. at Ivon Beni and Huachi Beni, Bolivia.

Var. *FORTIS* Forel

(Plate 49, fig. a)

Ps. sericea, Wheeler, Trans. 2nd Intern. Ent. Congr. (1912) 1913 p. 135.

Guatemala. Patulul and Escuintla (Wheeler), in hollow twigs of *Triplaris macombii* D. Smith.

Var. *ITA* Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906, p. 230 ♀; Deutsch. Ent. Zeitschr. 1909 p. 260 ♀.

Costa Rica: San Mateo (P. Biolley) in trunk and foliage of *Bixa orellana*.

Paraguay: San Bernardino (K. Fiebrig), in branches of *Alchornea irucurana*.

Guatemala: Escuintla and Patulul (Wheeler); in hollow twigs of *Triplaris auriculata*.

Panama: Numerous colonies were taken in the cauline swellings of *Cordia gerascanthus* at Ancon, C. Z. and in twigs of *Triplaris americana* at Balboa.

The worker of this form differs from the typical *sericea* in having a smaller postpetiole and the petiole with a less convex anterodorsal surface. The latter terminates behind in a distinctly transverse border. The female (deālated) measures 5.5 to 6 mm. and is much more like the type in the shape of the petiole and postpetiole. The head is longer and its posterior border more deeply excised than in the worker.

Var. LONGIOR Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 690 ♀.

Peru: Tarapoto (E. Ule), in perforated twigs of *Platymischium stipulare* Benth.

Var. RUBIGINOSA Stitz

Stitz: Deutsch. Ent. Zeitschr. 1913 p. 211 ♀.

Brazil: Alto Acre (E. Ule), *Triplaris* sp.

PSEUDOMYRMA SPINICOLA Emery

Emery, Bull. Soc. Ent. Ital. 22, 1890 p. 64 ♀ ♀ ♂ Pl. 6 Fig. 2; Biol. Centralbl. 11, 1891 p. 166; Anal. Mus. Nac. Costa Rica 1892, p. 66; Wheeler, Trans. 2nd Intern. Ent. Congr. Oxford (1912) 1913 p. 117; Wasmann, Tijdschr. Ent. 58, 1915 p. 304.

The worker of the typical form of this species measures 4-5.5 mm. and is ferruginous red or testaceous, with the gaster towards the tip and the middle portions of the femora somewhat infuscated, the mandibles, clypeus and cheeks pale yellow. Head distinctly longer than broad, the eyes rather convex and two-fifths as long as its sides; the clypeus with a short, broad median lobe which is distinctly toothed on each side; the frontal groove rather indistinct, the frontal carinae continued back as a pair of low narrow welts which subside at the middle of the head. Antennal scapes reaching to the middle of the internal orbits; second funicular joint longer than broad, joints 3-7 as long as broad. Thoracic dorsum straight in profile, except for the projecting semicircular mesonotum and the rather pronounced mesoepinotal impression. Sides of pronotum distinctly submarginate, the base of the epinotum longer than the declivity and forming a distinct angle with it. Metathoracic stigmata prominent. Petiole with well-developed peduncle, the node rather low, rounded, longer than broad, narrowed gradually in front and abruptly behind; the antero-ventral tooth small and acute. Postpetiole nearly twice as broad as the petiolar node, slightly broader than long, narrowed in front, convex behind. Legs rather long and slender. The body is densely and finely punctate, slightly shining, covered with short whitish pubescence and with a few pale erect hairs on the dorsal surface. Mandibles shining, rather coarsely but not sharply striatopunctate.

In the female the head is fully one and one-half times as long as broad, the sides subparallel. The color is darker than in the worker,

the front, mesonotum and epinotum somewhat infuscated. Wings smoky, with piceous veins and pterostigma. Length 7-8.5 mm.

The male was seen by Emery and its head figured. I have seen no specimens of this sex belonging to the typical *spinicola*.

The types were taken by Anastasio Alfaro at Alajuela and Jimenez, Costa Rica, in the spines of an acacia (identified by Wasmann as *A. spadicigera*). I have specimens from Surubres, San Mateo, Costa Rica (P. Biolley).

Subsp. ATROX Forel

Forel, Mém. Soc. Ent. Belg. **20**, 1912 p. 24 ♀.

The worker measures 4-4.7 mm. and is somewhat smaller than the typical form of the species, with subopaque mandibles, the median lobe of the clypeus less pronounced, less emarginate and more feebly dentate. Head somewhat shorter, the eyes a trifle larger, somewhat more than one-third the length of the sides of the head. Petiole decidedly shorter, its node scarcely longer than broad, the peduncle also more abbreviated, the postpetiole somewhat narrower, scarcely twice as broad as the petiolar node. Pilosity sparser and shorter, pubescence more dilute, the punctuation less pronounced and the surface therefore somewhat more shining. The color is the same as that of the typical *spinicola*.

The type specimens were taken by Mr. Christophersen in Panama. This is the common form of the species in the Savanna country south of the city of Panama where I have often taken it near Las Sabanas and Pueblo Nuevo. The acacia in the thorns of which it lives, is probably *penonomensis* Saff.

Subsp. CONVARIANS Forel

Forel, Bull. Soc. Vaud. Sc. Nat. **49**, 1913 p. 214 ♀.

The worker of this form was taken by Peper at Patulul, Guatemala, probably on Acacia. It measures only 3.6 mm. The head is short, only one and one-fourth times as long as broad, with more convex sides and posterior border than in the typical *spinicola*, the scapes are shorter, scarcely reaching to the median transverse diameter of the head, the funicular joints, except the first and last, are broader than long. The petiolar node is much higher and shorter and the legs are shorter. The sculpture and pilosity as in the type, the color slightly paler.

Subsp. GAIGEI Forel

Forel, Deutsch. Ent. Zeitschr. 1914 p. 615 ♀.

The worker of this subspecies was taken by Dr. F. M. Gaige at Fundacion, in the Santa Marta Mountains of Colombia. It measures 4.5–5.2 mm. The postpetiole is very broad, two and one-half times as broad as the petiole, which is gradually narrowed in front and has a shorter peduncle than the typical *spinicola*. The funicular joints are broader, the head more opaque, the color of a more vivid reddish, with the gaster brown.

I have received two cotypes from Dr. Gaige, who probably took them in *Acacia* thorns.

Subsp. INFERNALIS subsp. nov.

Worker. Length 3.5–4.5 mm.

Head as in the typical *spinicola*, about one and one-third times as long as broad, with feebly convex sides and feebly excavated posterior border. Clypeal lobe broadly emarginate in the middle, with a rather large, blunt tooth on each side. Antennal scapes reaching a little beyond the middle of the internal orbits, funicular joints 3–10 as broad as long. Base and declivity of epinotum subequal, forming a distinct obtuse angle in profile. Petiolar node long and low, its posterior corners almost subdentate, the peduncle very short, without a tooth on its ventral surface. Postpetiole as long as broad, about one and two-thirds as broad as the petiolar node. Legs long.

Punctuation moderately strong and dense, uniform, the surface slightly shining; mandibles and clypeus somewhat more shining than the head.

Hairs and pubescence much as in the typical *spinicola*, the pubescence longest and most distinct on the head and gaster.

Castaneous brown; the gaster, femora and tibiae, except their tips and bases, somewhat darker; the head paler anteriorly, the mandibles, clypeus, antennae and, in some specimens, also the lateral borders of the pronotum, brownish yellow.

Female. Length 6.5–7 mm.

Head longer than in the typical *spinicola*, very nearly twice as long as broad, the sides parallel, except at the anterior corners where they are slightly dilated, the posterior border nearly straight. Eyes elongate, more flattened than in the typical *spinicola*, clypeal lobe longer and much narrower, its corners rounded, not dentate. Mandibles larger, parallel-sided, flattened near the tips, their external borders

straight. Thorax rather long and narrow, but through the wing-insertions as broad as the head. Pronotum concave on the sides, strongly submarginate above; mesonotum and base of epinotum straight in profile, the latter longer than the declivity and forming with it a rounded angle. Petiole and postpetiole similar to those of the worker, the former with a longer, lower and posteriorly more angular node, the latter distinctly longer than broad.

Sculpture, pilosity and color much as in the worker; mesonotum more shining; pronotum and borders of gastric segments yellowish or golden brown like the mandibles, clypeus, cheeks, antennæ and tarsi. Wings deeply infuscated, with very dark brown veins and pterostigma.

Malc. Length about 5 mm.

Head, including the eyes, distinctly longer than broad, oval, narrowed and rounded behind, without posterior corners. Mandibles with well-developed though scarcely denticulate apical borders. Clypeus depressed, tuberculate in the middle, concave on the sides, its anterior border with a small, median, projecting lobe. Scape about twice as long as broad, first funicular joint nearly one and one-half times, second nearly four times as long as broad, remaining joints shorter. Mesonotum convex. Epinotum small, low, convex and rounded, without distinct base and declivity. Petiole without a distinct node, about three times as long as broad, gradually narrowed anteriorly. Postpetiole distinctly longer than broad, behind about twice as broad as the petiole. Gaster long, enlarged at the tip. Legs slender.

Sculpture finer than in the female; epinotum very smooth and shining; mandibles opaque, indistinctly punctate.

Pilosity and pubescence much as in the worker and female; clypeus and mandibles very hairy.

Color as in the worker and female but the pale portions, including the genitalia, are whitish yellow. Antennæ fuscous. Wings slightly paler than in the female.

Described from numerous specimens of all three phases found living in the thorns of *Acacia penonomensis* Saff. at Las Cascades C. Z. (type locality), along the Chivachiva Trail, near Red Tank, C. Z., at Venado, C. Z. and Pueblo Nuevo, in the Panamanian Republic. This form is readily distinguished by its dark color.

Subsp. *SCELEROSA* subsp. nov.

Worker. Length 3.5—4 mm.

Head as in the subsp. *convarians*, slightly longer than broad, with

convex sides and straight or feebly convex posterior border. Clypeal lobe entire, transverse, with rather blunt corners. Eyes about two-fifths as long as the sides of the head. Antennal scapes reaching beyond the middle of the head, almost to a line joining the posterior orbits; funicular joints 2-4 a little longer than broad, 5-10 as long as broad. Pronotum less distinctly submarginate on the sides, epinotum more rounded, the angle between the base and declivity less distinct than in the typical *spinicola*. Petiole and postpetiole much the same as in the type. Legs rather long.

Punctuation finer, the surface somewhat more shining, the pilosity and pubescence somewhat finer and less abundant. The color is paler, more uniformly yellowish red, the mandibles, clypeus, legs and antennæ yellow.

Described from a number of specimens taken by Prof. C. F. Baker at Granada, Nicaragua, in the thorns of *Acacia costaricensis* Schenck.

PSEUDOMYRMA SUBTILISSIMA Emery

Emery, Bull. Soc. Ent. Ital. **22**, 1890 p. 65 ♀ ♀ Pl. 6 Fig. 7; Biol. Centralbl. **11**, 1891 p. 166; Anal. Mus. Nac. Costa Rica 1892 p. 66; Wheeler, Trans. 2nd. Intern. Ent. Congr. Oxford (1912) p. 118; Wasmann, Tijdschr. Ent. 1915 p. 305.

Costa Rica: Alajuela (A. Alfaro) in thorns of an *Acacia* (identified by Wasmann as *spadicigera*), which was inhabited also by *Ps. belti*.

PSEUDOMYRMA TENUIS Mayr

(Plate 49, fig. b)

Numerous workers of the typical form of this species were taken by Mr. H. O. Lang at Merumè-Mouth, British Guiana, in the fistulose stems of a *Patima formicaria*, Johnst. This ant nests by preference, at least in British Guiana, in dead twigs.

PSEUDOMYRMA TRIPLARIDIS Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 684 ♀ ♀ ♂; C. R. 6 me. Congr. Intern. Zool. Berne (1904) 1905 p. 452; Ann. Soc. Ent. Belg. **50**, 1906 p. 231.

Worker. Length 4.4-4.8 mm.

Mandibles shining, with a few striæ, more porrect than in *arboris-sanctæ* (*triplarina* Wedd.) Clypeus higher, carinate, more truncate than in that species. Funicular joints 3-9 more than twice as broad as long; the funiculus much thicker and shorter than in *arboris-sanctæ*

and its race *symbiotica*. Head narrower than in *arboris-sanctæ*, more as in the race *symbiotica*. Thorax less convex and broader than in *arboris-sanctæ*, rather flat as in the race *symbiotica*, but broader and somewhat more sharply marginate and with much steeper epinotal declivity. First node as in *arboris-sanctæ*, shorter and broader than in *symbiotica* and somewhat marginate anteriorly. Second node less narrowed in front.

Moderately shining; head feebly shining, more densely and more deeply punctate than in *arboris-sanctæ* and *symbiotica*. Erect pilosity as in these two forms. The appressed pubescence, however, is longer and denser; on the gaster it forms a gray investment which partially conceals the sculpture.

Blackish brown; tarsi, tibiæ, antennæ and anterior part of head ferruginous; mandibles brownish red; femora brown.

Female. Length 7.5—8 mm.

Much more slender than *arboris-sanctæ*. Mandibles longer, narrower, feebly bent, not at all angularly geniculate and without a transverse impression in the external border (both of these peculiarities obtain in *arboris-sanctæ*, though much less strongly than in *dendroica*) with about 6 teeth on the apical and three teeth on the inner border. Head longer than broad, broader behind, with straight posterior border (feebly concave in *arboris-sanctæ*). Epinotum with two sharply separated surfaces, though node as in *arboris-sanctæ*, but the second much less and very feebly narrowed in front, $1\frac{2}{3}$ to $1\frac{3}{4}$ times as broad as long.

Brownish black; legs brown; mandibles, antennæ and anterior portion of head ferruginous. Wings tinged with brownish black veins and pterostigma. Pilosity and sculpture as in the worker.

Male. Length about 5 mm.

Head rounded rectangular, broader behind, with distinct, nearly straight posterior border. Antennæ rather short; third to last funicular joints about twice as long as broad. Thorax barely broader than the head. Like the female in sculpture, pilosity and color, but the mandibles, tarsi, scapes and first funicular joint pale yellow. Nodes lower, the first below with a small tooth which is lacking in the race *symbiotica* or *arboris-sanctæ*.

Jurua Miry, Jurua, Amazonas, June 1907 (E. Ule) in the hollow twigs of *Triplaris surinamensis* Cham. (Forel) Tabatinga, Para (Goeldi and Huber), in the medullary cavities of the same tree.

Two worker cotypes of this species, given me by Prof. Forel, have been compared with the following three subspecies from British Guiana:

Subsp. BOXI subsp. nov.

Worker. Length 4—5 mm.

Very similar to the typical *triplaridis* in the shape of the head, but the epinotum has the base and declivity straight in profile and meeting at a much more distinct angle. The peduncle of the petiole is decidedly longer, the node smaller and the hook-shaped ventral tooth larger.

The punctuation is sharper and deeper so that the surface, especially of the head, is less shining than in the typical *triplaridis*. The pilosity is distinctly less abundant, the tibiæ having only a few erect hairs, the pubescence much finer and shorter on the gaster. Color paler, ferruginous brown; mandibles red, with black teeth; coxæ yellow, paler than the antennæ and legs; gastric segments with yellowish bases and apical borders.

Female. Length 6.5—7 mm.

Closely resembling the worker, but the head somewhat longer; the punctuation coarser, so that the surface, especially of the head, is subopaque. The pilosity is even sparser than in the worker, the color decidedly darker brown, but with the yellow markings of the gaster paler and more vivid. The antennæ, cheeks, clypeus, tibiæ and tarsi are yellowish brown, the wings distinctly infuscated, with strong blackish veins and pterostigma.

Male. Length 6.5 mm.

Head through the eyes as broad as long, its posterior border straight, its posterior corners rounded. Antennal scapes about $2\frac{1}{4}$ times as long as broad, the first funicular joint longer than broad, the third to 12th joints a little more than twice as long as broad, the second three times as long as broad.

Surface of body a little smoother than in the worker, pilosity almost lacking, pubescence rather long but dilute. Color much as in the worker, the wings as in the female.

Described from many workers and a few females and males taken by Mr. H. E. Box on the Blairmont Plantation, at Berbice, British Guiana, in the branches of *Triplaris surinamensis*. Among the specimens is an old, deälated mother queen, measuring fully 11 mm. with the gaster enormously distended and its sclerites widely separated as in old termite queens. This condition, of which I find no mention in the literature, I have observed occasionally also in the old mother queens of some other species of *Pseudomyrma*.

Subsp. BAILEYI subsp. nov.

(Plate 50)

Worker. Length 4.5—5 mm.

More elongate and more slender than the preceding form, head decidedly longer, fully one and one-third times as long as broad, suboblong, as broad in front as behind, with very feebly and evenly convex sides and straight posterior border. Eyes more elongate; clypeal lobe somewhat more transverse, the scapes reaching only to the second third of the internal orbits. Thorax, especially the epinotum longer and narrower; sides of pronotum more sharply submarginate; base of epinotum longer in proportion to the declivity which is more sloping and forms a more rounded angle with the base. Petiole longer than broad, the node as long as broad (distinctly broader in the typical form), postpetiole also distinctly longer in proportion to its width.

Sculpture and pubescence very much as in the type but the erect pilosity on the body distinctly more abundant.

Color somewhat darker brown, especially in the dorsal surface; mandibles, anterior portion of head and antennæ paler.

Female. Length 7—7.5 mm.

Head greatly elongated, fully one and three-fourth times as long as broad, more sharply rectangular than in the worker, with straight, parallel sides, sharp posterior corners and distinctly excised posterior border. Eyes elongate and very flat. Thorax very long, subdepressed above, broadened through the mesonotum where it is somewhat wider than the head. Pronotum sharply marginate and concave on the sides; mesonotum more convex in profile, the epinotum very long, its base straight, fully three times as long as the declivity, which is much more nearly vertical than in the worker. Petiole one and one-half times as long as broad; postpetiole behind scarcely broader than long. Gaster elongate.

Sculpture, pilosity and color as in the worker but with the following differences: the front and gular surface of the head, sides of pronotum, ventral surface of petiole and bases of gastric segments bright brownish yellow. Wings grayish hyaline, with resin-yellow pterostigma and pale yellow veins.

Described from numerous specimens taken at Camaria on the Cuyuni River, British Guiana, in the hollow branches of *Triplaris surinamensis*. This very distinct subspecies is dedicated to Prof. J. W. Bailey. It is quite as fierce and aggressive as the various forms of *triplarina* (*vide infra*), *damnosa*, *maligna*, etc. In the locality in

which it was found it was in full possession of a cluster of about a dozen trees which grow so close to the river that their roots are submerged during the rainy season.

Subsp. *TIGRINA* subsp. nov.

Worker. Length 4—5 mm.

Resembling *baileyi* in the elongation of the head, but it is only $1\frac{1}{4}$ times as long as broad and appears to be more rectangular because the sides are more nearly straight and parallel. Eyes distinctly shorter. Epinotum a little shorter but of the same shape as in *baileyi*, the petiole more abruptly truncated behind, the postpetiole distinctly shorter in proportion to its width.

Surface of body, and especially of the head, much smoother and more shining than in *baileyi*, with smaller, shallower and sparser punctures.

Hairs less abundant, pubescence shorter and finer, though abundant and partially concealing the surface on the gaster.

Color decidedly paler, yellowish red; mandibles red; posterior portion of head and the remainder of the body brownish, with yellowish sutures; the borders and bases of the gastric segments reddish yellow; antennæ and legs yellowish, with the middle portions of the femora brownish.

Male. Length 6.5—7 mm.

Head decidedly longer than in the male *boxi*, longer than broad; eyes much narrower. Surface of body more shining, the suberect pilosity on the dorsal surface and even on the tibiæ distinct, though sparse. Petiolar node lower than in *boxi*. The wings are much paler than in the other subspecies and the typical form, being grayish hyaline, with pale yellow, weak veins and pale brown pterostigma.

Described from numerous workers and a few males taken by Mr. H. E. Box on the Blairmont Plantation, Berbice, British Guiana, in the branches of *Triplaris surinamensis* (not the same tree as the one inhabited by the subsp. *boxi*).

PSEUDOMYRMA TRIPLARINA (Weddell)

Myrmica triplarina Weddell, Ann. Sc. Nat. Botan. (3) **13**, 1849 pp. 40–113, 249–268 ♀; *Pseudomyrma arboris-sanctæ* Emery, Bull. Soc. Ent. Ital. **26**, 1894 p. 147 ♀ ♀; Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 685 *nota*; Emery, Bull. Soc. Ent. Ital. **37**, 1905 p. 119 ♀; Forel, C. R. 6me Congr. Intern. Zool. Berne (1904), 1905 p. 452.

Brazil. (Weddell), in *Triplaris*; Coxipo, Matto Grosso (F. Silvestri).
 Bolivia: (Weddell, Balzan), in *Triplaris* ("palosanto"); Huachi Beni
 (W. M. Mann), in *Triplaris*.

Peru. (Weddell); Collanga (Staudinger). Tarapoto, Amazonas.

Dr. Bequaert (1921-22) has recently ascertained that Weddell, as early as 1849, discovered this ant and its relations to *Triplaris* in Brazil, Bolivia and Peru. He describes the insect as follows: "Je ne crois pas que cette insecte ait été observé dans d'autres conditions que celles que j'ai notées; sa forme linéaire est particulièrement adaptée à son genre de vie. J'ai eu l'occasion de l'examiner et même de souffrir ses atteintes dans bien des parties du Brésil, en Bolivie et au Pérou; et partout il m'a paru identique. Déjà plusieurs voyageurs ont signalé une partie des faits dont il vient d'être question, et ils ont rapporté la fourmi du *Triplaris* au genre *Myrmica* de Latreille; mais je ne sache pas qu'on lui ait donné de nom spécifique; on pourrait lui appliquer celui de *Myrmica triplarina*. Elle est ordinairement d'un brun clair. Sa longueur est de 6 ou 7 millimètres, et sa largeur de 1 millimètre; l'abdomen est cylindrique et un peu atténué vers son extrémité postérieure qui est poilue." Since there can be no doubt of the identity of this ant with Emery's *arboris-sanctæ* described in 1894, the rules of nomenclature require the substitution of the older name. This is unfortunate, both because the insect has been so long known under the name *arboris-sanctæ* and because Forel has described another species as *Ps. triplaridis*.

The following is a translation of Emery's description of *Ps. arboris-sanctæ*:

"*Worker*. Testaceous, head somewhat darker, borders of mandibles fuscous, abdomen, except the pedicel, in great part brownish, abundantly pilose and delicately pubescent, shining; head and thorax anteriorly rather shining; head a little longer than broad, scarcely narrowed in front, truncated behind, its anterior portion somewhat less finely, the posterior portion very finely, microscopically punctulate; eyes moderately large, flattened; clypeus anteriorly with a truncated lobe; mandibles smooth at the base, apically punctate-striate; antennal funiculi with joints 3-10 broader than long; thorax incised in the region of the mesometanotal suture; pronotum convex, obtusely marginate on each side; mesonotum sub-semicircular, epinotum with the sides not distinctly marginate, the angle between the base and declivity obtuse, rounded; petiole with a very short peduncle in front, posteriorly with a large node, which is transverse, nearly twice as broad as long; postpetiole transverse cupuliform. Length 5-6 mm.

Female. Testaceous throughout; first gastric segment often with two brown spots at the base; vertex around the ocelli and the thorax sometimes clouded with brown; sculpture nearly as in the worker; eyes larger, nearer to the mouth; peduncle of petiole nearly straight above, not concave, the node trapezoidal, scarcely broader than long; postpetiole less distinctly transverse than in the worker. Length 8-9 mm.

Collected by Balzon in Bolivia where it lives in a tree called "palo santo" (*Triplaris?*). On another occasion I received the same species from Tarapotas, in Amazonas. The new species is recognizable by its color, by the mandibles, which have their apical half strongly striated or striato-punctate, by the strongly transverse petiolar node of the worker, by the antennal funiculus, the second joint of which is much smaller than the first and not broader than long, by the nearly uniform punctuation of the head and the size and distribution of the punctures. All these characters are in part found in other species (some of which are still unpublished), but all are united in this species. I do not exclude the possibility that this species may be the *Tetraponera testacea* of F. Smith, but his description is too indefinite to enable one to establish its identity."

Var. CORDOBENSIS Forel

Forel, Bull. Soc. Vaud. Sc. Nat. (5) 50, 1914 p. 265 ♀.

Argentina: Cordoba (C. Bruch); probably from *Triplaris*. The worker is described as follows: "Length 6-6.5 mm. Differs from the type of the species in its larger eyes and the form of the head, the posterior border of which is narrower, less sharp and not concave. The angle between the base and declivity of the epinotum is also sharper, the former less convex and the latter higher (long). The color is reddish yellow throughout, more vivid than in the type, which has the gaster brown. In other respects identical."

Var. RURRENABAQUENSIS Wheeler & Mann, var. nov.

Worker. Length 4-5 mm.

Smaller than the type and the preceding variety and somewhat more slender, the head narrow behind as in the latter, the antennal scapes a little shorter and broader, joints 2-10 of the funiculus more transverse, the eyes larger, the clypeal lobe shorter, more rounded at the corners and less distinct than in the typical *arboris-sanctæ*. Brownish yellow, with dark brown gaster and ocellar region; mandibles,

clypeus and antennæ yellow, paler than the legs which in most specimens are as dark as the body. The pilosity, especially on the gaster, is somewhat more abundant than in the type of the species.

Described from six specimens taken by Dr. W. M. Mann at Rurrenabaque Beni, Bolivia, in the cavities of *Triplaris*.

Subsp. SYMBIOTICA Forel

Forel, Rev. Suisse Zool. **12**, 1904 p. 38 ♀ ♂; C. R. 6 me Congr. Intern. Zool. Berne (1904) 1905, p. 451; Ann. Soc. Ent. Belg. **49**, 1905 p. 158 ♀.

Colombia. *Dibulla* (A. Forel); in trunk and branches of *Triplaris* (probably *americana*).

Venezuela: Las Trincheras (Meinert), in *Triplaris*.

Forel describes the worker and male as follows:

"*Worker*. Length 4.4–4.7 mm. Reddish yellow, with the middle of the gastric segments brownish. Smaller and paler than the type. Erect pilosity much sparser, very sparse on the tibiæ and scapes. Mandibles narrower, with more oblique terminal border, more feebly sculptured, shining towards the base. Thorax subopaque, much more densely punctate, as is also the head. The pubescence is also decidedly shorter and decidedly less abundant. The promesonotal suture is more deeply impressed, forming a small notch in the thoracic dorsum. Pronotum not at all submarginate, with convex sides. Sting very short. In other respects like the type."

"*Male*. Length 5–5.5 mm. Head rather oval, mandibles sharp, with an apical tooth. Frontal groove deep. Petiolar node as long as broad, subopaque or feebly shining, punctate. Pilosity even more scattered than in the worker, except on the abdomen. Yellowish brown. Wings subhyaline, with pale veins."

Forel has described the nesting habits of this ant in the trunk and branches of *Triplaris*. More recently Dr. George Salt has sent me workers and females of this subspecies taken by him in *Triplaris* branches at Aracataca, Colombia, which is not very far from the type locality.

Var. LOEWENSOHNI Forel

(Plate 51, fig. a)

Ps. arboris-sanctæ subsp. *symbiotica* var. *panamensis* Forel, (*nom. praeocc.*) Mém. Soc. Ent. Belg. **20**, 1912, p. 22 ♀ ♀ ♂; *Ps. arboris-sanctæ* Wheeler, Trans. 2nd Intern. Ent. Congr. (1912) 1913 p. 135; *Ps. arboris-sanctæ* subsp. *symbiotica* var. *loewensohni* Forel, Soc. Vaud. Sc. Nat. **51**, 1918 p. 719.

Panama. Frijoles (Christophersen and Wheeler); in trunk and branches of *Triplaris americana*; also Las Cascades and Marajal, Balboa (Wheeler).

All three phases of this variety were taken in 1911 by Mr. Christophersen and myself near Frijoles in a region which is now at the bottom of Gatun Lake. It differs but little from the subsp. *symbiotica* as shown in Forel's description which is here translated:

"*Worker*. Length 5.5–5.7 mm. Larger than the type of the subspecies and of a more yellowish brownish yellow. Head slightly more coarsely punctate and less narrowed in front. Otherwise identical.

Female. Length 8–11 mm. The same differences as in the worker. Petiolar node a little narrower than in the typical *arboris-sanctæ*.

Male. Length 5.5–5.8 mm. Of a deeper brown color. Petiolar node lower and narrower than in the typical *symbiotica*."

During 1923 and 1924 I again took this variety in a different locality near Frijoles, near Las Cascades and at Balboa in the Canal Zone.

PSEUDOMYRMA ULEI Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 689 ♀ ♀ ♂.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in twigs and branches of *Coussapoa* sp. (Ule No. 5717).

Subfamily MYRMICINÆ

Genus PHEIDOLE Westwood

Forel, Verh. Zool. bot. Ges. Wien 1908 p. 376 ♀ ♀.

Brazil: São Paulo (A. Lutz), living in large bamboos.

PHEIDOLE CRAMPTONI Wheeler

subsp. PETIOLICOLA Wheeler

Wheeler, Zoologica **3**, 1921 p. 147 ♀.

British Guiana: Kartabo (Wheeler), in the hollow petioles of *Tachigalia paniculata* Aubl.

PHEIDOLE GAUTHIERI Forel

var. OXYMORA Forel

Forel, Mém. Soc. Ent. Belg. 19, 1912 p. 233 ♂ ♀.

Panama: (Christophersen), nesting in the spines of *Xanthoxylon* sp. (probably *panamensis* P. Wies)¹

PHEIDOLE LUTZI Forel

Forel, Ann. Soc. Ent. Belg. 49, 1905 p. 168 ♂ ♀.

Brazil: São Paulo (A. Lutz), nesting in the cavities of bamboo.

PHEIDOLE MINUTULA Mayr

Mayr, Verh. Zool. bot. Ges. Wien. 1877 p. 872 ♂ ♀ ; Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 681 ♂ ♀.

Brazil: (J. Trail), probably in the leaf-sacs of *Tococa*; Jurua Miry, Jurua, Amazonas (E. Ule), in the leaf sacs of *Tococa ulei* Pilger.I took numerous colonies of this ant in the primitive forest near Kalacoon, British Guiana, in the leaf-sacs of *Tococa aristata* Benth.

The soldier agrees more closely in the sculpture of the head with Mayr's original description than with Forel's account of the specimens taken by Ule. The humeral angles have a small, acute tooth, however, and the epinotal teeth are minute, slender and acute. In the worker also the humeri are dentate and the epinotal teeth are like those of the soldier though smaller.

The *female* (undescribed) measures 4-4.3 mm. The rugæ on the head are coarser than in the soldier and extend very nearly to the posterior border and corners. The mesonotum is flat, the teeth on the epinotum very minute, short and broad at the base. Shining, mesonotum more coarsely, abdomen more finely punctate. Brown, the head darker and more blackish, especially about the ocelli, the appendages and pleura paler, the mandibles and anterior portion of the head red, the wings distinctly brownish, with brown veins and pterostigma. Pilosity much as in the soldier.The *male* (undescribed) measures about 2.5 mm. Head and thorax opaque and very finely punctate, the former black behind, the mesonotum dark brown above, the remainder of the thorax, the mandibles,¹According to Pitter (1922) "the corky prickles of the lower third of the trunk are said to be often hollow and inhabited by ants." The type-locality of the plant is Mamei Hill, C. Z.

clypeus, abdomen and appendages sordid yellow, the wings somewhat paler than those of the female, the pilosity similar, but shorter and more appressed. The hairs on the legs are very short, bristly and oblique.

Subsp. FOLICOLA Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904, p. 681 ♂ ♀ ♀ ♂.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in the petiolar sacs of *Maieta poeppigii* Mart. and *M. guianensis* Aubl.

PHEIDOLE (HENDECAPHEIDOLE) TACHIGALIÆ Wheeler

Ph. tachigaliæ Wheeler, Zoologica 3, 1921 p. 148 ♂ ♀ ♀ ; *Ph. (Hendecapheidole) tachigaliæ* Wheeler, Amer. Mus. Novitates No. 46, 1922 p. 3.

British Guiana: Kartabo (Wheeler), in the hollow petioles of *Tachigalia paniculata* Aubl.

I have also taken a single colony of this ant at Kartabo in a cauline swelling of *Cordia nodosa* L.

Genus CREMATOGASTER Lund

CREMATOGASTER CORVINA Mayr

Mexico: Vera Cruz (A. Dampf), nesting in thorns of *Acacia vera-cruzensis* Schenck.

CREMATOGASTER SANGUINEA Roger

subsp. LUCAYANA Wheeler

C. lucayana Wheeler, Bull. Amer. Mus. Nat. Hist. 21, 1905 p. 95 ♀ ; *C. sanguinea* subsp. *lucayana* Wheeler Bull. Mus. Comp. Zool. 54, 1913 p. 490.

Bahamas: Andros and New Providence Islands (Wheeler), nesting in Tillandsias.

I found the var. *torrei* Wheeler in Cuba nesting in dead branches and twigs. Probably the typical form also prefers such nesting sites.

Var. ETIOLATA Wheeler

Wheeler, Bull. Amer. Mus. Nat. Hist. 21, 1905 p. 96 ♀ ♀.

Bahamas: Andros Island (Wheeler), nesting in Tillandsias.

CREMATOGASTER (EUCREMA) ACUTA Fabr.

Panama: Tabogilla Island (Wheeler), nesting in Tillandsias on manganillo trees (*Hippomanes mancinella*); Barro Colorado Island, Gatun Lake, C. Z. (Wheeler), in cauline swellings of *Cordia alliodora*; Tumba Muerte Road, near Las Sabanas (Wheeler), in thorns of *Acacia penonomensis*.

Large colonies of this ant nest by preference in dead branches.

CREMATOGASTER (ORTHOCREMA) ARCUATA Forel

var. ARUGA Forel

Panama: Otoque Island and Taboga Island (Wheeler), nesting in Tillandsias.

CREMATOGASTER (ORTHOCREMA) ARIZONENSIS Wheeler

Wheeler, Journ. N. Y. Ent. Soc. 20, 1912 p. 130, 132 ♂ ♀.

Arizona: Huachuca Mountains (Wheeler); nesting in stems of mistletoe (*Phoradendron flavescens* var. *villosum*).

CREMATOGASTER (ORTHOCREMA) ARMANDI Forel

Forel, Bull. Soc. Bot. Genève 12, 1920 p. 208 ♂ ♀; Chodat and Carisso, *ibid.* p. 195 fig. 326.

Brazil: Matto Grosso (Spencer Moore), in the pseudobulbs of an orchid (*Cattleya?*).

CREMATOGASTER (ORTHOCREMA) ATRA Mayr

Mexico: Vera Cruz (A. Dampf), nesting in thorns of *Acacia vera-cruzensis* Schenck.

This ant is apparently merely a subspecies of *brevispinosa* Mayr.

CREMATOGASTER (ORTHOCREMA) BRASILIENSIS Mayr

Mayr, Verh. Zool. bot. Ges. Wien 1877, p. 875 ♂.

Northern Brazil: (James Trail), probably in myrmecophytes.

British Guiana: Kamakusa (H. O. Lang), nesting among the matted roots of an undetermined epiphyte.

Bolivia: Tumupasas (W. M. Mann), nesting in the petiolar sacs of *Tococa* sp.

Var. LUDIO Forel

C. limata subsp. *ludio* Forel, Mém. Soc. Ent. Belg. **19**, 1912, p. 217 ♂; Wheeler, *Zoologica* **3**, 1921, p. 151 ♂ ♀.

C. brasiliensis var. *ludio* Forel, Bull. Soc. Vaud. Sc. Nat. **49**, 1913, p. 233.

Panama: Fort Clayton and Miraflores, C. Z. (Wheeler); common in the hollow branches of *Triplaris americana*.

British Guiana: Kartabo (Wheeler), in petioles of *Tachigalia paniculata* Aubl.; Grand Batavia Island, Cuyuni River (Wheeler), in cauline swellings of *Cordia nodosa* L.; Kamakusa (H. O. Lang), in petiolar sacs of *Tococa aristata* Benth.

Bolivia: Huachi Beni (W. M. Mann), in cauline swellings of *Cordia alliodora* var.

The specimens taken by Mr. Lang include a deälated female, which is entirely yellow. This phase of the typical *brasiliensis*, according to Forel, has a black gaster. The epinotal teeth are as long as broad at their bases and rather blunt at the tip. In the worker, as in the typical *brasiliensis*, the mesonotal carinæ are minutely dentate.

CREMATOGASTER (ORTHOCREMA) BREVISPINOSA Mayr

Emery, Biol. Centralbl. **11**, 1891 p. 168; Anal. Mus. Nac. Costa Rica 1892 p. 67; Zool. Jahrb. Abt. Syst. **9**, 1896, p. 626; Forel, Bull. Soc. Vaud. Sc. Nat. (5) **44**, 1908, p. 47; Safford, Smithson. Rep. (1921) 1923 p. 393, Pl. 15.

Costa Rica: (A. Alfaro), in *Acacia* thorns; Surubrès, near San Mateo (P. Biolley), in bromeliads.

Nicaragua: Granada (Gen. E. Chamorro), in thorns of *Acacia costaricensis*.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

Var. AMPLA Forel

Colonies, comprising all three phases of this ant, were taken near Las Cascades, C. Z. Panama, March 14, 1923, in the hollow twigs of *Triplaris americana*.

Var. MINUTIOR Forel

Wheeler, Amer. Natural. **35**, 1901 p. 526; Ann. Soc. Ent. Belg. **45**, 1901 p. 203.

Mexico: Cuernavaca (Wheeler), nesting in *Tillandsia benthamiana*.

Subsp. TUMULIFERA Forel

Forel, Bull. Soc. Vaud. Sc. Nat. (5) 44, 1908 p. 47 ♀ ♀ ♂.

Costa Rica: San José (P. Biolley), in stem of a *Clematis*.

Panama: Las Sabanas (Wheeler), in Tillandsias; Chivachiva Trail, near Red Tank, C. Z., common in cauline swellings of *Cordia alliodora*.

Guatemala: Zacapa (Wheeler), in thorns of *Acacia hindsii* and Tillandsias.

CREMATOGASTER (ORTHOCREMA) CURVISPINOSA Mayr

var. PANAMANA var. nov.

Worker. Length 1.8—2 mm.

Differing from the typical form of the species in its somewhat smaller size, somewhat more slender thorax and in having the epinotal spines more slender and tapering, though as long as in the type and curved inward in the same manner. Head very smooth and shining, as long as broad, very much rounded behind, with the eyes at the posterior third of the sides. Antennal scapes almost reaching the convex posterior border of the head. Pilosity white, long and very sparse as in the type, each epinotal spine with a long hair near the middle. Color also as in the type, piceous brown, mandibles a little paler, gaster and usually also the head black.

Female. Length 3.5—4 mm.

Head more rectangular than in the worker, with distinct posterior corners and nearly straight posterior border. Antennal scapes shorter than in the worker. Thorax short, somewhat more than twice as long as broad, narrower through the wing insertions than the head, epinotum sloping, the spines reduced to small, rather acute teeth. Petiole and postpetiole as in the worker, the anterior angles of the former a little less rounded. Gaster very small, not longer than the remainder of the body, broadest through the first segment, very acute at the tip. Sculpture and color as in the worker, pilosity more abundant on the dorsal surface of the head, thorax and gaster. Wings grayish hyaline, with pale yellow veins and pterostigma.

Numerous specimens found nesting in the thorns of *Acacia penonensis* Saff. along the Tumba Muerte Road, near Las Sabanas, Panama.

CREMATOGASTER (ORTHOCREMA) DELITESCENS Wheeler

Wheeler, Zoologica 3, 1921 p. 152 ♀ ♀.

British Guiana: Kartabo (Wheeler), in the hollow petioles of *Tachigalia paniculata* Aubl.

CREMATOGASTER (ORTHOCREMA) DISTANS Mayr

Guatemala: Zacapa (Wheeler), in Tillandsias.

Costa Rica: Alajuela (Wheeler), in Tillandsias.

CREMATOGASTER (ORTHOCREMA) GOELDII Forel

var. CHODATI Forel

Forel, Bull. Soc. Bot. Genève (2) 11, 1920 p. 7 ♀.

Paraguay: Concepcion (Chodat and Vischer), in the trunk of *Agonandra brasiliensis*, inhabited also by *Cryptocerus eduarduli* Forel.

CREMATOGASTER (ORTHOCREMA) LAEVIS Mayr

Mayr, Verh. zool. bot. Ges. Wien 1877 p. 876 ♀; Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 682 ♀.

Brazil: (James Trail), probably in myrmecophytes; Jurua Miry, Jurua, Amazonas (E. Ule), in petiolar sacs of *Maieta tococoides* Cogn.; Cachoeira, Jurua (E. Ule), in the petiolar sacs of *Maieta juruensis* Pilger.

CREMATOGASTER (ORTHOCREMA) LIMATA F. Smith

Mayr, Verh. zool. bot. Ges. Wien 1877 p. 875 ♀.

Brazil: (James Trail), probably in myrmecophytes.

British Guiana: Kartabo (Wheeler), in the internodes of young *Cecropia angulata* Bailey; in large branch of *Cecropia angulata* Bailey; in large branch of *Cecropia sciadophylla* var. *decurrens* Snethl.

Var. PALANS Forel

Wheeler, Zoologica 3, 1921 p. 151 ♀ ♀.

British Guiana: Kartabo (Wheeler), in the hollow petioles of *Tachigalia paniculata* Aublet; Merumé-Mouth (H. O. Lang), in the fistulose stems of *Patima formicaria* Johnston.

The female (undescribed) is 8-9 mm. long, yellow, with blackish funiculi, black gaster and ocellar spots, three longitudinal dark brown streaks on the mesonotum and blackish wings with dark brown veins and pterostigma. Head decidedly broader than long, slightly narrower in front than behind, with straight posterior border and convex dorsal surface. Scapes reaching to the occipital corners. Epinotum with two stout acute teeth, which are distinctly longer than broad at the base. Posterior corners of petiole acute and dentate.

The *male* (undescribed) measures 4–4.5 mm. and is colored like the worker, dark brown, with the head behind the clypeus, the mandibles, antennæ and legs yellow, the femora and tibiæ somewhat infuscated in the middle. The wings are as dark as in the female.

Subsp. PARABIOTICA Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 683 ♀; Stitz, Deutsch. Ent. Zeitschr. 1913 p. 209 ♀ ♀; Wheeler, Ecology 2, 1921 p. 96; *Crematogaster* sp., Mann, Psyche 19, 1912 p. 36.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in *Tillandsia*; Alto Acre (E. Ule), in inflorescence of *Costus* sp.; Pará (W. M. Mann), in ant gardens in parabiosis with *Dolichoderus bispinosus*.

British Guiana: Kartabo (Wheeler), in "ant gardens" in parabiosis with *Camponotus femoratus*.

Panama: Las Sabanas and Tabogilla Island (Wheeler), in *Tillandsias*; Chivachiva Trail, near Red Tank, C. Z. (Wheeler), in cauline swellings of *Cordia alliodora*.

Bolivia: Rurrenabaque (W. M. Mann), in the leaf sacs of *Tococa* (Rusby) No. 1756.

This ant has also been taken by Forel, Christophersen and myself living in parabiosis with *Dolichoderus (Monacis) parabioticus* in dead wood in Colombia, Panama and other Central American localities.

CREMATOGASTER (ORTHOCREMA) LUTZI Forel

Forel, Ann. Soc. Ent. Belg. 49, 1905 p. 167 ♀ ♀.

Brazil: São Paulo (A. Lutz), nesting in cavities of bamboo.

CREMATOGASTER (ORTHOCREMA) MONTEZUMIA F. Smith
var. SULCATA Mayr

Wasmann, Tijdschr. Ent. 48, 1905 p. 210 pl. 8 fig. 1.

Brazil: Rio Grande do Sul, in ant garden among *Tillandsias*.

CREMATOGASTER (ORTHOCREMA) SCULPTURATA Pergande

Guatemala: Zacapa (Wheeler), nesting in *Tillandsias*.

CREMATOGASTER (ORTHOCREMA) SUMICHRASTI Mayr

A single colony found nesting in a cauline swelling of *Cordia alliodora* at Quebrada de Oro, C.Z., Panama.

CREMATOGASTER (ORTHOCREMA) VIRGULA Forel

Costa Rica: Alajuela (Wheeler), nesting in Tillandsias.

Panama: Chivachiva Trail, near Red Tank, C. Z. (Wheeler), nesting in cauline swellings of *Cordia alliodora*.

Genus MONOMORIUM Mayr

MONOMORIUM CARBONARIUM F. Smith

subsp. EBENINUM Forel

Wheeler, Bull. Amer. Mus. Nat. Hist. **24**, 1908 p. 127 ♀ ♀ ♂.

Panama: Corrozal, C. Z. (Wheeler), in Tillandsias.

Guatemala: Escuintla (Wheeler), in the hollow twigs of *Triplaris auriculata*.

Porto Rico and Bahamas: (Wheeler), in Tillandsias.

MONOMORIUM FLORICOLA Jordan

Wheeler, Bull. Amer. Mus. Nat. Hist. **24**, 1908, p. 127 ♀ ♀ ; Bull. Mus. Comp. Zool. **54**, 1913, p. 485.

Bahamas: Porto Rico and Cuba (Wheeler), in Tillandsias.

Spanish Honduras: Tela (Prof. Oakes Ames), nesting in the pseudobulbs of an orchid, *Epidendrum imatophyllum* Linde.

Genus XENOMYRMEX Emery

XENOMYRMEX STOLLI Forel subsp. FLORIEANUS Emery var.

LUCAYANUS Wheeler.

Wheeler, Bull. Amer. Mus. Nat. History **21**, 1905 p. 87 ♀ .

Bahamas: Andros Island (Wheeler), nesting in Tillandsias.

Genus ALLOMERUS Mayr

ALLOMERUS DECEMARTICULATUS Mayr

Mayr, Verh. zool. bot. Ges. Wien **27**, 1877, p. 873 ♀ ; Forel, Mém. Soc. Ent. Belg. **20**, 1912 p. 3 ♀ .

Brazil: (James Trail), probably in *Tococa* or *Cordia*; Oyapoc, (A. Ducke), in dilated peduncle of *Hirtella* sp.

Mayr distinguished three species of *Allomerus* according to the number of antennal joints in the worker as *decem-*, *octo-* and *septem-*

articulatus. The discovery by Dr. Mann of a form with nine joints and Forel's observation that occasional specimens of *septemarticulatus* may have eight joints or one of the basal funicular joints partially divided, leads me to include all the forms in one species and to regard the number of joints in the worker as a subspecific character. In all the forms, so far as known, the female has ten antennal joints, the male thirteen.

Subsp. NOVEMARTICULATUS Wheeler & Mann, subsp. nov.

Worker. Length 1.2—1.3 mm.

Closely resembling the typical *octoarticulatus* in structure, sculpture, pilosity and color but the antennæ 9-jointed. Scapes reaching half the distance between the posterior orbits and occipital corners of the head. Promesonotum moderately convex, mesoepinotal impression moderately deep but acute. Epinotum a little longer than broad, the base in profile straight, shorter than the declivity and forming with it a distinct though rounded angle. Petiolar node laterally compressed, longer than broad; postpetiole a little broader, as broad as long and a little broader behind than in front.

Described from eight workers taken by Dr. W. M. Mann on the Rio Madid, Bolivia, in leaf-sacs of *Tococa*.

Subsp. OCTOARTICULATUS Mayr

Allomerus octoarticulatus Mayr, Verh. zool. bot. Gesel. Wien **27**, 1877 p. 873 ♀ ; Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 679 ♀ ♀ ♂.

Brazil: (James Trail); Para (Forel collection), in the leaf-sacs of *Remijia physophora* Benth; Marary, Amazonas (E. Ule), in the petiolar sacs of *Tococa setifera* Pilger.

Bolivia: Rio Negro, Tumupasa and Riberalta (W. M. Mann), in the cauline swellings of *Cordia hispidissima*.

The worker closely resembles the preceding form in the size of the eyes, length of the scapes and shape of the thorax, but the epinotum and petiolar node are as broad as long and the latter is not compressed laterally. The post-petiole is slightly broader than long and more rounded though a little broader than the petiolar node.

The female measures 6 mm. and has the head longitudinally rugulose. The body is reddish brown, the head and mesonotum above and the greater part of the abdomen brown.

The male measures 5.3 mm. and is colored like the female.

Var. EXSANGUIS Wheeler and Mann, var. nov.

Worker. Length 1—1.3 mm.

Of the same dimensions as the typical *octoarticulatus*, the antennal scapes of the same length, the eyes of the same size, but of a pale, whitish or ivory yellow color, with the mesonotum more depressed above, the epinotum shorter than broad and in profile more rounded with a more sloping declivity. The petiole has a distinctly shorter petiole and a more anteroposteriorly compressed node, the postpetiole shorter in proportion to its length. The erect hairs on the head and thorax seem to be shorter and less numerous.

Female (deälated). Length nearly 6 mm.

Very similar to the female of the typical *octoarticulatus*. Mandibles deep red, rather subopaque, antennæ, pronotum, mesonotum, scutellum and gaster behind the first segment reddish brown. The epinotum has a feeble projection or angle on each side.

Described from five workers and a single female taken by Dr. W. M. Mann at Esperanza, Bolivia, in a cauline swelling of *Cordia hispidissima*.

Var. DEMERARÆ var. nov.

Worker. Length 1.5—1.8 mm.

Differing from the two preceding forms of the subspecies in having the scapes somewhat longer, the eyes distinctly larger, the thorax longer, the mesonotum for the most part straight and sloping in profile, the mesoepinotal impression somewhat shallower but longer, the epinotum distinctly longer than broad, subangular in profile, but with very sloping declivity, one and one-half times as long as the base. The peduncle of the petiole is longer, the node conical in profile, seen from above as long as broad, somewhat compressed anteroposteriorly at the summit, which is transverse. Postpetiole subcircular, very nearly as long as broad and slightly broader than the petiole.

Female. Length 6—6.5 mm.

Antennal scapes reaching the posterior ocelli. Epinotum with distinct teeth, which are short and rather acute, shorter than the length of their bases. Sculpture of head finer than in the preceding forms, very indistinctly rugulose; the scutellum and epinotum smooth. The color of the body is more uniformly reddish brown, the dorsal surface of the head and thorax and the posterior portion of the gaster slightly darker in some specimens, mandibles subopaque, more shining along the borders. Wings blackish or dark brown, with dark brown veins and pterostigma.

Male. Length 5.5—6 mm.

Very similar to the male of the typical *octoarticulatus*. Brownish yellow, posterior portion of head, pronotum, except its borders, mesonotum, scutellum, posterior portion of epinotum, gaster and nodes of pedicel, base of scapes, apical portion of funiculi and middle portions of femora and tibiæ dark brown. Wings colored as in the female.

Described from many specimens of all three phases which I took at Kartabo, Kalacoon and on Grand Batavia Island in the Cuyuni River, British Guiana, in the cauline swellings of *Cordia nodosa* L. Mr. H. O. Lang has also taken the same variety in the same plant at Kamakusa, in the interior of British Guiana.

According to Hohenkerk (Journ. Board. Agric. Brit. Guiana 11, 1918 p. 99) the natives call this ant "Kurabelli". In the localities above mentioned, it inhabited most or all of the cauline swellings on every one of the numerous specimens of *C. nodosa* examined both by Prof. J. W. Bailey and myself.

Var. ANGULATUS Wheeler & Mann, var. nov.

Worker. Length 1.5 mm.

More deeply colored than the preceding forms of the subspecies, reddish yellow, head a little darker, mandibles red, their bases and teeth and the border of the clypeus blackish. Eyes smaller than in the typical form and the var. *exsanguis* but the scapes somewhat longer. Promesonotum more convex, mesoepinotal impression short and acute, deeper than in the typical *octoarticulatus*. Epinotum as long as broad, the base and declivity forming a very distinct angle in profile as in the var. *tuberculatus* Forel, but not subdentate and with the base straight. Petiolar node rising more abruptly from the peduncle than in var. *demerarae*, but less than in *tuberculatus*. Post-petiole broader than long, about one-third broader than the petiolar node, with rather straight sides.

Thirteen workers taken by Dr. W. M. Mann on the Lower Rio Madidi, Bolivia, in cauline swellings of *Cordia hispidissima*.

Var. TUBERCULATUS Forel

Allomerus 8-articulatus race *tuberculatus* Forel, Mém. Soc. Ent. Belg. 20, 1912 p. 2 ♀ ♂.

Worker. Length 2—2.3 mm.

Somewhat larger than the preceding forms. Head more nearly square, less trapezoidal, that is less narrowed in front, with less convex sides, scarcely longer than broad, the scapes slightly shorter. Mesoepinotal constriction deep. Epinotum with two small tubercles or swellings at the angles. Color less uniform, sordid yellow, head reddish or brownish yellow, a brown band or cloud on the middle of the gaster.

Male. Length 5.2 mm.

Posterior portion of head behind the eyes longer and narrower, somewhat more rectangular and less rounded than in the typical *octoarticulatus*. Epinotum with two blunt tubercles. Color slightly paler, more yellowish brown.

This form was described from specimens taken by J. Huber at Monte Verde, on the Middle Purus, Amazon Basin, in the leafsacs of a *Tococa*. I have examined cotypes received from Prof. Forel.

Var. MELANOTICUS Wheeler & Mann, var. nov.

Worker. Length 1.6—1.8 mm.

Resembling the typical *octoarticulatus* in the shape of the head but the scapes somewhat longer. The promesonotum at its posterior end descends abruptly to the mesoepinotal constriction which is very pronounced. The epinotum is as long as broad, in profile with subequal base and declivity, the former convex, the latter sloping and somewhat concave, the angles as distinctly tuberculate or subdentate as in the var. *tuberculatus*. Petiole nearly as high through the node as long, the node rising abruptly from the well-developed peduncle, which is as long as the node. The latter is transversely elliptical from above, the postpetiole very similar but slightly broader.

Deep castaneous brown; head, pro- and mesonotum and posterior portion of first gastric segment black; mandibles deep red; antennæ and legs brown, the femora darker in the middle, the tarsi and posterior borders of the gastric segments sordid yellow.

Female. Length about 6 mm.

Head smaller, the thorax longer and narrower than in the var. *demerarae*, the former nearly as long as broad, the sides of the epinotum with blunt projections instead of teeth, the median surface more concave. Sculpture of head distinctly coarser, that of the mesonotum finer, so that this region is more shining. Mandibles more coarsely striate and punctate. Pubescence gray instead of golden yellow, much more dilute.

Black; mesopleura, legs and antennæ dark brown, borders of

gastric segments golden brown. Wings dark, of the same shade as in the var. *demerarae*.

Male. Length 5.6 mm.

Head longer especially behind the eyes, than in the var. *demerarae*, thorax somewhat more slender. The epinotum as in that form, rounded and sloping, without angular swellings. Subopaque; pilosity and pubescence as in the female.

Dark brown; dorsal surface black, mandibles, articulations of antennæ and legs, tarsi, edges of gastric segments and external genital valves sordid yellowish brown. Wings colored as in the female.

Described from numerous workers, a female and a male taken by Dr. W. M. Mann at Tumupasa (type locality) and Isiamas, Bolivia, in leaf-sacs of *Tococa* sp.

Subsp. SEPTEMARTICULATUS Mayr

Allomerus septemarticulatus Mayr, Verh. zool. bot. Ges. Wien **27**, 1877 p. 874 ♀; Schumann, Die Ameisenpflanzen 1889 p. 31; Emery, Anal. Mus. Nac. Costa Rica 1892 p. 65; *A. octoarticulatus* var. *septemarticulatus* Forel, Zool. Jahrb. Abt. Syst. **20**, 1904, p. 680 ♀ ♀.

Brazil: (James Trail), probably in a myrmecophyte; Amazonas (K. Schumann), in leaf sacs of *Duroia saccifera* Spruce; São Joaquim, Rio Negro, Amazonas (E. Ule), in leaf sacs of *Duroia saccifera* Spruce.

As already stated, Forel found a variation of the number of antennal joints in some workers from seven to eight joints. The female is identical with that of *octoarticulatus*, but is a little smaller and paler.

Genus SOLENOPSIS Westwood

SOLENOPSIS CORTICALIS Forel

Wheeler, Bull. Amer. Mus. Nat. Hist. **24**, 1908 p. 131 ♀ ♀.

Porto Rico: Utuado (Wheeler), in bamboo.

Subsp. AMAZONENSIS Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 686 ♀ ♀ ♂.

Peru: Cerro de Escaler, about 1300 m. (E. Ule), nesting in a Tillandsia (*Pseudocatopsis* n. sp.).

SOLENOPSIS HELENA Emery subsp. HERMIONE Wheeler

Wheeler, Zoologica **3**, 1921 p. 157 ♀ ♀.

British Guiana: Kartabo (Wheeler), in petioles of *Tachigalia paniculata* Aubl.

Subsp. ULTRIX Wheeler

Wheeler, *Zoologica* 3, 1921 p. 157 ♂ ♀.

British Guiana: Kartabo (Wheeler), in petioles of *Tachigalia paniculata* Aubl.

SOLENOPSIS PICEA Emery

Taken by Dr. W. M. Mann at Ixiamas, Bolivia, nesting in a cauline swelling of *Cordia gerascanthus* var.

SOLENOPSIS TENUIS Mayr

Mayr, *Verh. zool. bot. Ges. Wien* 1877 p. 874 ♂ ♀.

Northern Brazil: (James Trail), probably in a myrmecophyte.

Bolivia: Ixiamas (W. M. Mann), in a cauline swelling of *Cordia hispidissima*.

SOLENOPSIS ZETEKI sp. nov.

Worker. Length 1—1.1 mm.

Head subrectangular, distinctly longer than broad, as broad in front as behind, with nearly straight sides and posterior border. Eyes very small, consisting of only four or five very minute ommatidia, situated at one fourth the distance between the anterior and posterior corners. Mandibles narrow, with oblique, 4-toothed apical borders. Clypeus moderately convex in the middle, bicarinate, but without teeth in the anterior border, which is straight, transverse and entire in the middle. Antennæ slender, the scapes reaching a little more than half way between the eyes and posterior corners of the head; first funicular joint as long as the five succeeding joints together; joints 2-7 fully twice as broad as long, the two-jointed club slender, much longer than the remainder of the funiculus, the basal about one-third as long as the terminal joint. Thorax small, much narrower and slightly shorter than the head, including the mandibles, with short mesoepinotal impression, the epinotum small, similarly rounded, without distinct base and declivity. From above, the promesonotum is nearly one and one-third times as long as broad, rounded on the sides and in front, the epinotum about one and one-fourth times as long as broad. Petiole in profile as high as long, with a short peduncle, the node rising abruptly, with a conical summit, the ventral surface convex, with a distinct tooth near the anterior end; from above, the node is subcircular, as long as broad. Postpetiole slightly broader

than the petiole, a little broader than long, convex and rounded above but decidedly lower than the petiolar node. Gaster small and narrow, the first segment elongate-elliptical, narrowed in front and behind. Legs rather slender.

Very smooth and shining throughout, with very minute, scattered, piligerous punctures.

Hairs whitish, delicate, sparse and erect or suberect on the body, on the appendages reduced to very fine appressed pubescence.

Mandibular teeth and eyes black, all the rest of the body pale yellow, the head and thorax a shade darker than the abdomen and appendages.

Described from several specimens taken from a colony that was nesting in a cauline swelling of *Cordia alliodora* on the Chivachiva Trail, near Red Tank, C. Z. Panama.

This species which is dedicated to Mr. James Zetek, resembles *sulfurea* Roger of Venezuela, *succinea* Emery of Costa Rica and *inermiceps* Wheeler and Mann of Haiti in having the clypeus unarmed, but it is much smaller and the antennal scapes are much shorter than in the species mentioned.

Genus MACROMISCHA Roger

MACROMISCHA PETIOLATA (Forel)

Leptothorax petiolatus Forel, Ann. Soc. Ent. Belg. 45, 1901, p. 129 ♀; Wheeler, *ibid.* p. 201; Amer. Natural. 35, 1901, p. 527; *Macromischa petiolata* Emery, Gen. Insect. Myrmicinae 1921, p. 247.

Mexico: Cuernavaca (Wheeler), in *Tillandsia benthamiana*.

Genus LEPTOTHORAX Mayr

Subgenus GONIOTHORAX Emery

LEPTOTHORAX (GONIOTHORAX) ECHINATINODIS Forel subsp.

ACULEATINODIS Emery var. PLEURITICUS Wheeler

Wheeler, Zoologica 3, 1921, p. 158 ♀ ♀ ♂.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aublet.

Subsp. CORDINOLA subsp. nov.

Worker. Length 2—2.3 mm.

Resembling the subsp. *dalmasi* Forel and *aculeatinodis* Emery but

considerably smaller and differing in other respects. Mandibles subopaque, striated; clypeus smooth in the middle, without median carina and with the lateral carinæ feeble. Head very shining, but finely, superficially and not very evenly striate, with longitudinal seams of coarse punctures, most numerous anteriorly, posterior corners very smooth. Thorax above rather delicately longitudinally reticulate, rugose, densely punctate on the sides, except the mesopleura, which are smoother and more shining. Epinotal spines as long as the basal surface, as in *dalmasi*, rather stout and erect, but curved outwards and downwards. Petiole and postpetiole with well-developed spinules, longitudinally rugose and punctate above. Gaster smooth and shining, its base very finely and obscurely striate. Hairs whitish, erect, blunt, abundant and of even length on the body, appendages with very short, appressed hairs or pubescence.

Piceous black, head and gaster, except at the base, darker; mandibles, scapes, basal half of funiculi, epinotal spines, tibiæ and tarsi brownish yellow. In many specimens the pronotum is paler and more reddish than the remainder of the thorax, thus approaching the coloration of the subsp. *aculeatinodis* Emery.

Numerous specimens taken from the cauline swellings of *Cordia alliodora* along the Chivachiva Trail, near Red Tank, C. Z. Panama. A few colonies were also found nesting in the thorns of *Acacia penonomensis* Saff. along the Tumba Muerte Road, near Las Sabanas, Panama.

This subspecies seems to be most closely related to the subsp. *aculeatinodis*, described from Costa Rica, but it differs considerably in sculpture, pilosity and color. Emery fails to give the dimensions of his specimens.

Subsp. DALMASI Forel

(Plate 51, fig. b)

Costa Rica: Alajuela and San Jose (Wheeler), nesting in Tillandsias and also in dead twigs.

Subsp. SPININODIS Mayr

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 626.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

LEPTOTHORAX (GONIOTHORAX) UMBRATILIS Wheeler

Wheeler, Zoologica 3, 1921 p. 160 ♂ ♀.

British Guiana: Penal Settlement, near Bartica (Wheeler), in hollow petioles of *Tachigalia paniculata* Aublet.

WASMANNIA AUROPUNCTATA Roger

Dominica W. I.: Roseau (Wheeler) under the appressed leaves of juvenile *Marcgravia*s growing over a large boulder.

Genus CEPHALOTES Latr.

CEPHALOTES ATRATUS L.

British Guiana: Kartabo (Wheeler), nesting in large branch of a *Cecropia Sciadophylla* var. *decurrens* Sn.

Genus CRYPTO CERUS F. Smith

CRYPTOCERUS AZTECUS Forel

Wheeler, Amer. Natural. 35, 1901, p. 527; Ann. Soc. Ent. Belg. 45, 1901 p. 201.

Mexico: Cuernavaca (Wheeler), in *Tillandsia benthamiana*.

CRYPTOCERUS BOHLSI Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896, p. 631 2 ♀ Fig. C.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

CRYPTOCERUS EDUARDULI Forel

Forel, Bull. Soc. Bot. Genève (2) 11, 1920 p. 4 2 ♀ ♀.

Paraguay: Concepcion (Chodat and Vischer), in trunk of *Agonandra brasiliensis*.

CRYPTOCERUS GRANDINOSUS F. Smith

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 636 2 ♀.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

CRYPTOCERUS PELTATUS Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 633 2 ♀; Fig. D.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

CRYPTOCERUS PILOSUS Emery

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 630 2 ♂; Fig. B.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

CRYPTOCERUS QUADRATUS Mayr

Emery, Zool. Jahrb. Abt. Syst. 9, 1896, p. 634 2 ♀; Fig. E.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

CRYPTOCERUS ROHWERI Wheeler

C. (Cyathocephalus) rohweri Wheeler, Proc. New. Eng. Zool. Soc. 6, 1916 p. 34 2 ♀.

Arizona: Santa Catalina Mountains (Chrisman), in limbs of palo verde (*Cercidium torreyanum*).

CRYPTOCERUS SCUTULATUS F. Smith

Costa Rica: Alajuela (Wheeler), in *Tillandsias*.

CRYPTOCERUS WHEELERI Forel

Wheeler, Amer. Natural. 35, 1901 p. 527; Ann. Soc. Ent. Belg. 45, 1901 p. 201.

Mexico: Cuernavaca (Wheeler), in *Tillandsia benthamiana*.

CRYPTOCERUS sp.

Fiebrig, Biol. Centralbl. 29, 1909 p. 38.

Paraguay: San Bernardino (K. Fiebrig), occasionally in *Cecropia peltata* L.

Subgenus PARACRYPTOCERUS Emery

CRYPTOCERUS (PARACRYPTOCERUS) COMPLANATUS Guérin

subsp. RAMIPHILUS Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904, p. 678 ♂ ♀.

Brazil: Bom Fim Jurua, Amazonas (E. Ule), in perforated twigs of *Platymischium uli* Harms.

CRYPTOCERUS (PARACRYPTOCERUS) CORDLÆ Stitz

Stitz, Deutsch. Ent. Zeitschr. 1913 p. 207 ♂ ♀. Fig. 1.

Brazil: Alto Acre (E. Ule), in *Cordia*.

CRYPTOCERUS (PARACRYPTOCERUS) MINUTUS Fabr.

Emery, Biol. Centralbl. 11, 1891 p. 168; Anal. Mus. Nac. Costa Rica 1892, p. 67; Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 234 ♀.

Costa Rica: (A. Alfaro) in thorns of *Acacia*; El Hiquito, near San Mateo (P. Biolley), in trunk of *Bixa orellana*.

Panama: Tumba Muerte Road, near Las Sabanas (Wheeler), nesting in thorns of *Acacia penonomensis*; Miraflores, C. Z. (Wheeler), in twigs of *Triplaris americana*.

This ant is most frequently found nesting in the dead twigs and branches of various trees and bushes or even in the culms of coarse grasses.

CRYPTOCERUS (PARACRYPTOCERUS) PUSILLUS Klug

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 636 ♂ ♀ ♀ ♂.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

Like *C. minutus*, this ant is most frequently found nesting in dead branches.

Subgenus CYATHOCEPHALUS Emery

CRYPTOCERUS (CYATHOCEPHALUS) PALLENS Klug

Emery, Zool. Jahrb. Abt. Syst. 9, 1896 p. 635 ♂ ♀ ♀ ♂; Safford, Smithson. Rep. (1921) 1923 p. 393.

Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

Mexico: San Sebastian, near Tuxtla (G. N. Collins), in thorns of *Acacia collinsi* Saff.

Var. DISCOCEPHALUS F. Smith

Emery, Biol. Centralbl. **11**, 1891 p. 168; Anal. Mus. Nac. Costa Rica 1892, p. 67.

Costa Rica (A. Alfaro), in thorns of Acacia.

Var. PORRASI var. nov.

(Plate 52)

Soldier. Length 4—4.5 mm.

Differing from the typical form of the species and the var. *discocephalus* and *patellaris* Mayr in having the short scale-like hairs which emerge from the foveolæ on the cephalic disc and thorax, pedicel and base of gaster much coarser and more conspicuous and with a silvery luster. The concavity of the cephalic disc is distinctly shining and its foveolæ are larger and not crowded.

Worker. Length 3.5—3.7 mm.

Characterized by having the same peculiarity of the squamiform hairs as the soldier and also by the thorax which is narrower, with the marginal tooth of the mesonotum more acute and the lateral margin of the epinotum represented by three distinct teeth. The lateral spines of the petiole and postpetiole are long, flat, curved and acute, those on the former longer than on the latter.

Female. Length 5—6 mm.

Only the disc of the head shows the enlarged squamiform hairs, the pilosity of the remainder of the body being as in the typical *pallens*. Wings heavily infuscated, not pale as in *varians*, with dark brown veins and pterostigma.

Male. Length 4—5 mm.

Very similar to the male of *varians* but the gaster is darker, as dark as the thorax, heavily shagreened and more opaque and the wings are darker, as in the female.

Dedicated to the former genial and learned president of the Panamanian Republic, Dr. Belisario Porras, and described from numerous specimens of all four phases found nesting in the cauline swellings of *Cordia alliodora* and in the dead twigs of various trees and shrubs at Quebrada de Oro (type locality), Red Tank, Frijoles and Ancon, C.Z.

The colonies of this ant are not very populous. Its habits are similar to those of *Colobopsis*. The elliptical nest-entrance is guarded by one of the soldiers, which occludes the orifice with the disk-shaped dorsal cephalic disc just as the *Colobopsis* soldier uses the truncated circular

anterior surface of the head for the same purpose. The cephalic disc in old soldiers and the mother queen of the colony often becomes coated with dirt and extraneous particles so that it closely resembles the bark of the plant.

CRYPTOCERUS (CYATHOCEPHALUS) SETULIFER Emery

(Plate 53)

Soldier. (Undescribed) Length 3.5–3.8 mm.

Resembling the female, but the posterior border of the cephalic disc sharper. This disc is broadly oval, the narrower being the posterior end. The thorax is much like that of *pallens* in shape but the pronotal crest is much less pronounced and interrupted in the middle. The gaster is shorter and more broadly elliptical. Color, sculpture and pilosity much as in the worker, except that the spots at the anterior corners of the first gastric segment are larger, more distinct and of a paler, more ivory yellow.

Male. (Undescribed) Length 3.8–4 mm.

Head, including the eyes, fully twice as broad as long, with a straight, broad posterior border and small rounded angles. Eyes and ocelli very convex. Mandibles very small, with very indistinctly and bluntly denticulate apical borders. Clypeus small and short, its anterior border entire and broadly rounded. Antennæ very long, the scapes only one and one half times as long as broad, the remaining joints much longer than broad, diminishing in length to the penultimate which is about two-thirds as long as the last joint. Thorax broad and convex, decidedly narrower than the head; humeri of pronotum rectangular; mesonotum as long as broad, subtriangular, with well-developed Mayrian furrows dividing it into three subequal convex areas. Epinotum small and short, with subequal base and declivity, meeting at a rounded right angle. Petiole and postpetiole from above subrectangular, about as long as broad, the latter at its anterior corners with short broad teeth representing those on the corresponding segment of the other phases. Gaster slender, the first segment as long as all the remaining segments together. Legs moderately long.

Head, including the mandibles, thorax, petiole and postpetiole opaque, densely and finely punctate-rugulose; gaster and legs slightly shining and merely densely punctate.

Hairs long, abundant, dull yellowish, erect, confined to the dorsal surface of the head and thorax, the tip of the gaster and flexor surfaces of the femora.

Black; antennæ dark brown; tibiæ and terminal tarsal joints reddish brown; genitalia and posterior borders of gastric segments golden brown or brownish yellow; wings colorless, hyaline, with pale brown pterostigma and yellowish veins.

Several colonies of this species were found nesting in the cauline swellings of *Cordia gerascanthus* trees just back of the laboratory at Ancon, C.Z. The soldier has the same habits as *C. pallens* and many of the individuals have the cephalic disc incrustated with foreign matter from exposure to the elements during guard duty at the oval nest entrance.

CRYPTOCERUS (CYATHOCEPHALUS) VARIANS F. Smith

(Plate 54)

Wheeler, Bull. Amer. Mus. Nat. Hist. **21**, 1905 p. 104 2 ♀ ♀ ♂.

Bahamas: New Providence and Andros Islands (Wheeler), nesting in Tillandsias.

Florida: Card's Point (Wheeler), in Tillandsias; Coconut Grove (Miss Nancy Fairchild), in twigs of *Coccoloba wifera*.

Subsp. MARGINATUS Wheeler and Mann

Wheeler and Mann, Bull. Amer. Mus. Nat. Hist. **33**, 1914 p. 39 2 ♀ ♀.

Haiti: (W. M. Mann), nesting in bamboos.

Genus APTEROSTIGMA Mayr.

APTEROSTIGMA CALVERTI Wheeler

Wheeler, Psyche **18**, 1911 p. 207 ♀ ♀; Calvert, a Year of Costa Rican Natural History 1917 p. 241 Fig.

Costa Rica: Juan Viñas (P. P. Calvert), nesting in Bromeliads.

Subfamily DOLICHODERINAE

Genus DOLICHODERUS Lund

DOLICHODERUS (HYPOCLINEA) BITUBERCULATUS Mayr

Morteo, Malpighia **18**, 1904 p. 509 ♀.

Ceylon and Buitenzorg; inhabiting twigs of *Triplaris americana* Vahl. The ant is a common East Indian species.

DOLICHODERUS (HYPOCLINEA) CHAMPIONI Forel

Subsp. TRINIDADENSIS Forel var. TAENIATUS Forel

Panama: Chivachiva Trail near Red Tank, C. Z. (Wheeler), nesting in the cauline swellings of *Cordia alliodora*; Las Sabanas (Wheeler), in Tillandsias.

The colonies in *Cordia* were very sporadic and small, those in the Tillandsias much more populous. This ant more frequently nests in dead branches.

DOLICHODERUS (HYPOCLINEA) LUTOSUS Mayr

Panama: Las Sabanas and Tabogilla Island (Wheeler), nesting in Tillandsias.

Costa Rica: Alajuela (Wheeler), in Tillandsias.

Guatemala: Escuintla (Wheeler), nesting in thorns of *Acacia bursaria* Schenck.

DOLICHODERUS (MONACIS) BISPINOSUS Olivier

Brazil: Pará (W. M. Mann), in ant gardens in parabiosis with *Crematogaster limata* subsp. *parabiotica*.

Panama: Las Sabanas (Wheeler) in Tillandsias; Chivachiva Trail, near Red Tank, C. Z., in cauline swellings of *Cordia alliodora*.

Fully developed colonies of this ant are very populous and live in dead branches and abandoned aërial termitaria. The colonies found in *Cordia* and Tillandsias were small and incipient and usually contained only a single mother queen. That pleometrosis, or the founding of a colony by a number of recently fecundated females, not infrequently occurs is indicated by my finding several associations of more than 60 deälated females all living together in single large Tillandsias.

DOLICHODERUS (MONACIS) DEBILIS Emery var. RUFESCENS Mann

Mann, Psyche 19, 1912 p. 40 ♀ ♀ ♂.

Brazil: Madeira—Mamore R.R., Matto Grosso (W. M. Mann), in ant-gardens in parabiosis with *Odontomachus affinis* subsp. *mayri*.

Genus IRIDOMYRMEX Mayr

IRIDOMYRMEX INIQUUS Mayr

var. NIGELLUS Emery

Costa Rica: San José, Alajuela and Cartago (Wheeler), nesting in Tillandsias.

Guatemala: Zacapa (Wheeler), in Tillandsias.

IRIDOMYRMEX PORDESCENS sp. nov.

Worker. Length 1.3—1.5 mm.

Head slightly longer than broad, somewhat broader behind than in front, with evenly rounded sides and straight posterior border. Eyes small, flattened, in front of the middle of the head. Mandibles with long apical margin, furnished with two acute apical teeth, the remainder finely and indistinctly denticulate. Clypeus strongly convex in the middle, its anterior border entire and nearly straight. Frontal area rather distinct, large, triangular; frontal groove obsolete; frontal carinae short, as far apart as their distance from the lateral borders of the head. Antennae rather slender, scapes extending only about twice their greatest diameter beyond the occipital corners, funicular joints distinctly longer than broad. Thorax small and compact, the pro- and mesonotum broadly and evenly convex in outline, the mesoepinotal impression short and acute, the epinotum small and short, very convex, its base rising rather abruptly from the impression, as long as the declivity with which it forms a rounded right angle, the metasternal angles produced backwards. Seen from above the pronotum is broader than long, transversely elliptical; the mesonotum narrow and rectangular, slightly longer than broad; the epinotum scarcely broader than the mesonotum and fully as broad as long. Petiole short, the scale well-developed, rather high, inclined forward, strongly compressed anteroposteriorly, rather narrow, with parallel sides and rounded, sharp and entire superior border. Gaster small, broadly elliptical, its first segment not covering the petiole.

Moderately shining; surface of body very finely sharpened.

Pilosity very delicate, whitish, represented by a few scattered hairs on the clypeus, thorax and gaster. Pubescence pale, very fine, uniformly distributed on the body and appendages, a little longer and oblique on the scapes. It is so dilute as not to obscure the shining.

Pale, sordid yellow; head slightly darker; mandibular teeth reddish.

Described from fourteen workers belonging to a single colony which I found nesting in a *Tillandsia* near San José, Costa Rica.

This species is quite distinct from any of the described Neotropical *Iridomyrmex* in its small size and the structure of the thorax.

Genus FORELIUS Emery
 FORELIUS MACCOOKI Forel
 var. BRASILIENSIS Forel

Forel, Deutsch. Ent. Zeitschr. 1909 p. 260 ♀.

Paraguay: San Bernardino (K. Fiebrig), in branches of *Alchornea urucurana*.

Subsp. FIEBRIGI Forel

Forel, Mém. Soc. Ent. Belg. 20, 1912 p. 44 ♀.

Paraguay: San Bernardino (K. Fiebrig), in branches of *Alchornea urucurana*.

Genus AZTECA Forel
 AZTECA ALFARI Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 338 ♀ fig. 48, 49;
 Boll. Mus. Zool. Anat. Torino 11, 1896, p. 3, 4 ♀, Fig. IV; Forel, Biol.
 Centr. Amer. Hymen. 3, 1899-1900 p. 112.

Of this species, which is one of the most abundant ants in the Cecropias of the Neotropical Region, Forel has described ten subspecies and varieties and I have added two varieties. I here compile the various scattered descriptions as an aid in identifying the forms. Undoubtedly many additional subspecies and varieties will be discovered in Central and South America. Emery describes the worker and female of the typical *alfari* as follows:

"*Worker*. Testaceous, scarcely shining; head slightly darker, reddish; gaster infuscated posteriorly. Subopaque, pubescent and with long hairs; scape and legs without erect hairs. Stature only slightly variable, head rather elongate, narrowed in front; eyes in front of the middle of its length; mandibles shining, nearly smooth, with nearly straight dental margin, about 8-toothed; scape scarcely reaching the posterior margin of the head in the worker minor, thorax robust, convex, mesometanotal suture deeply impressed; scale cuneiform, with rounded dorsal angle. Length $2\frac{1}{3}$ -3 mm; head of worker 1 x 0.9 mm, scape 0.8 mm."

"The female of this species is black, with the anterior portion of the head, the mandibles, antennæ, articulations of the legs, tibiæ and tarsi more or less reddish. Sculpture, pubescence and hairs as in the worker. The head is elongate rectangular, with slightly curved sides, the posterior margin entire. The posterior boundary of the eyes is

slightly in front of the posterior half of the head; the tip of the scape surpasses half the space which separates the eye from the occipital margin. The scale is higher than in the worker, with more acute border. The feebly smoky wings have reddish veins and a dark brown stigma."

The worker types were taken by A. Alfaro at Jimenez, Costa Rica, in the internodes of *Cecropia*, and I have taken it at Zent, Costa Rica, in the same hosts. Emery records the species also from Venezuela where it was taken in *Cecropia peltata* and Forel took specimens at Santa Marta, Colombia, in an unidentified *Cecropia*. He also cites the species from Bugaba, Panama (Champion).

In a more recent paper (1912) Forel says: "The forms which I called *alfari r. lucida* and *r. lucidula* should be regarded as distinct species. The former is much more dimorphic and has a large female, the head very large and elongate. The latter, on the contrary, is smaller and more monomorphic, with smaller and posteriorly more excavated head." In this paper I have retained the two forms as subspecies, since I am not certain that the differences mentioned by Forel are of specific value and since it seems wiser to keep all the forms together till the species can be more carefully studied on the basis of a much larger amount of material than either Forel or I have seen.

Var. AEQUALIS Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906 p. 239 ♀; Verh. Zool. bot. Ges. Wien **58**, 1908 p. 386 ♀ ♂; Mém. Soc. Ent. Belg. **20**, 1912 p. 51 ♀.

"*Worker*. Length 2.1-3.1 mm.

Much less dimorphic and smaller than the var. *aequilata* Forel, near the var. *ovaticeps*, but the sides of the head are less convex and the scapes are shorter; their tips distant from the occipital angles by about one-fourth their length in the large worker. The head is less excised posteriorly than in *ovaticeps*. The epinotum is even more cuboidal than in *ovaticeps*, the basal surface nearly flat and the stigmata protrude as tubercles towards its extremity. Scale rounded, thick. Color reddish yellow, a shade deeper than in *ovaticeps*. Opaque or feebly subopaque, like *ovaticeps*."

"*Female*. Length 7.5 mm.

Like the female of *mixta*, but the head narrower, longer. More reddish brown or brownish red, with brown spots. Gaster brown posteriorly."

"*Male*. Length 3 mm.

Tip of funiculus easily collapsing, shrivelled in all the specimens. Head trapezoidal posteriorly, scale somewhat thicker than in *mixta*. Color more dark brown."

Brazil: Obidos near Pará (Goeldi), in a *Cecropia*; Mexicana Island, Amazon Delta (Dr. Hagmann); Forel also cites this variety from Dibulla, Colombia (Lallemand).

Var. AEQUILATA Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 691 ♀ ♀ ♂; Ann. Soc. Ent. Belg. 50, 1906 p. 239; Ule, Ameisenpflanzen des Amazonasgebietes 1907.

"*Worker*. Length 2.6—4 mm.

Distinguished from the typical *alfari* by the form of the head. It is scarcely or only slightly broader behind than in front, otherwise as in the type, about one-fifth or one-sixth longer than broad. It is larger and more rectangular, with less convex sides than in the var. *ovaticeps* Forel from Pará. Color and sculpture as in the specific type and not as in the subsp. *lucida* Forel and *lucidula* Forel. In other respects like the specific type."

"*Female*. Length 7—7.4 mm.

Head more rectangular than in the var. *ovaticeps*, with less convex sides; otherwise the same in all particulars. Wings iridescent, suffused with yellowish brown, with brown stigma and brownish yellow veins."

"*Male*. Length 3.4 mm.

The head is more narrowed behind the eyes, with more distinct posterior border than in the male of the subsp. *lucidula*, in which it is more rounded. The color is yellow, feebly shining, but the specimens seem to be somewhat immature. Otherwise like *lucidula*, the male of which, however, is black and shining."

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in internodes of *Cecropia* (Ule No. 5587); Rio Purus (A. Goeldi and J. Huber), in *Cecropia*.

Var. CURTISCAPA Forel

"*Worker*.

Scape slightly shorter. Mesonotum more projecting, forming a boss more distinct from the pronotum than in the type of the species. The basal surface of the epinotum is also a little more elevated and more convex, so that a narrow and very distinct notch is formed in the constriction of the thorax between the mesonotum and said basal surface. In other respects like the typical *alfari*."

"*Female*. Length 7.2 mm.

Testaceous yellow, with the thorax and abdomen partly brown. In other respects like the worker.

"Panama (Christophersen), in the trunk of a *Cecropia*." I have taken this form at Corozal, C. Z. (Wheeler), in young *Cecropia*.

Var. FUMATICEPS Forel

Forel, Deutsch. Ent. Zeitschr. 1909, p. 250 ♀; Ross, Naturwiss. Wochenschr. 8, 1909, p. 829.

"*Worker*. Length 2.5—3 mm.

Head shaped as in the typical *alfari*, decidedly narrowed anteriorly, but somewhat smaller in the major worker and less deeply excised posteriorly. In the major worker the scape does not reach the occiput, in the minor worker it surpasses it only slightly. Thorax, gaster and legs as in the typical *alfari*; the scale is somewhat more rounded above. Brownish yellow; head, antennæ and mandibles dark brown, smoky.

Buenaventura, Mexico, collected by Dr. Ross in *Cecropia mexicana*. Ross cites this variety also from Zacuaparu, Mexico, in the same plant.

I have frequently taken what is evidently this same form in young *Cecropias* at Puerto Barrios and Quirigua, Guatemala. There is considerable variation in the coloring of the workers in the same colony. Usually only the larger individuals have the head dark brown and often the mesopleura are of the same color, while the smaller workers may be uniformly pale or brownish yellow.

The *female* (undescribed) measures 7.5–8 mm. and is also rather variable in color, the thorax and abdomen being dark reddish brown or blackish brown, the head red, with the front and vertex darker, the clypeus paler, the mandibles red with black borders, the scapes and legs dark brown, the funiculi brownish yellow, the gastric segments with rather broad and sharply delimited brownish yellow borders. The wings are lacking in all my specimens, which were found living in the *Cecropia* internodes, either singly or with their small colonies of workers.

Var. LANGI var. nov.

Worker. Length 2.3—3.5 mm.

Close to the var. *aequilata* Forel, but smaller, with the same color, sculpture and pubescence but much less pilosity; the mesonotum dis-

tinctly less convex in the major worker, the mesoepinotal constriction much shorter and more acute, the epinotum shorter and lower, the petiolar scale more compressed anteroposteriorly. The head is of the same shape in all the workers, similar to that of *aequilata*, but distinctly broader behind than in front and the sides are less convex, the scapes in the largest workers somewhat shorter, but even the penultimate funicular joints are as long as broad, as in *aequilata*.

Female. Length 6 mm.

Smaller than the female of *aequilata*. Black; with red mandibles, brown scapes and basitarsi and the sides of the clypeus, funiculi and remaining tarsal joints brownish yellow. Wings whitish, not suffused with yellowish brown as in *aequilata*, with pale yellow veins and dark brown pterostigma.

Male. Length 2.3—3 mm.

Also smaller than the male of *aequilata*. Head only moderately narrowed behind, with rounded posterior corners. Dark brown, thorax and top of head black; antennæ and tarsi pale brown. Wings whitish, with colorless veins and pale brown pterostigma.

Described from a number of specimens of all three phases taken by Mr. H. O. Lang at Kamakusa, British Guiana, in the internodes of *Cecropia*.

Var. MIXTA Forel

Forel, Verh. zool. bot. Ges. Wien 1908, p. 386 ♀ ♀ ♂; Fiebrig, Biol. Centralbl. 29, 1909 p. 4 et seq.

Worker. Length 2.4—3.6 mm.

Not so opaque as the type of the species, but less shining than the subsp. *lucida* Forel. Head more strongly narrowed in front than in the typical *alfari*, but less excised behind, somewhat broader. The scape is conspicuously short, like that of the var. *aequalis* Forel, scarcely surpassing the posterior fourth of the head. Scale blunt above. Basal surface of epinotum quadrangular, somewhat broader than long, in front almost as broad as behind (in the var. *aequalis* the head is scarcely broader behind than in front and the basal surface of the epinotum is much narrower in front than behind, as long as its posterior width). Darker than *aequalis*; gaster brown, vertex brownish red, otherwise yellowish red.

Female. Length 6.8 mm.

Blackish brown; funiculi, anterior border of head and tips of tarsi reddish. Head somewhat broader than in the var. *aequalis*. The scape

does not reach the posterior fourth of the head. Wings rather pale, with a brownish yellow tinge, brown stigma and brownish yellow veins. A single cubital cell as in other species."

"*Male*. Length 2.4—2.9 mm.

Head rectangular behind the eyes, not narrowed posteriorly (in the var. *aequalis* and the typical form of the species trapezoidal, narrowed posteriorly). Antennæ as in the var. *aequalis* and in the specific type; scape and first funicular joint broader than long; second joint very large, one and three-fourths times as long as broad, much broader than the first. The succeeding joints increasingly shorter and narrower, but nearly all longer than broad; the terminal joints somewhat longer (not thicker). Scale thin above and rather pointed. Black with brown appendages. Wings paler than in the female.

San Bernardino, Paraguay (Fiebrig); São Paulo, Brazil (von Ihering)
In *Cecropia peltata* L.

Var. OVATICEPS Forel

Forel, Rev. Suisse Zool. 12, 1904, p. 44 ♀ ♀ ♂.

"*Worker*. Length 2.4—3.5 mm.

A little less polymorphic than the typical *alfari*. Head slightly narrower, with more convex sides, as narrow behind as in front. The small differ little from the large workers. Color vivid yellow (the typical *alfari* is brownish yellow). The sculpture is that of the typical *alfari* but the stature is smaller than that of the subsp. *lucida* Forel, and larger than that of the subsp. *lucidula* Forel, which is even less polymorphic."

"*Female*. Length 7—7.5 mm.

Very similar to the female of the subsp. *lucidula* Forel, but the scale is higher, thinner and more acuminate and the first segment of the gaster more elongate and more attenuate anteriorly. Deep brown, with the anterior portion of the head, the legs, the antennæ and the mandibles yellowish red. Posterior borders of the gastric segments reddish. The form of the head is precisely as in *lucidula*, but the mandibles are shorter and thicker. The thorax, on the contrary, is a little narrower, the two surfaces of the epinotum less distinct from one another, forming together a more feeble convexity. The wings are lacking.

Pará (E. Goeldi)."

No doubt this, like the preceding varieties, inhabits Cecropias.

Subsp. *CECROPIÆ* Forel

Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 240 ♀; Verh. zool. bot. Ges. Wien 1908 p. 387 ♀ ♀ ♂; Mém. Soc. Ent. Belg. 20, 1912 p. 51 ♀ ♀; Stitz, Deutsch. Ent. Zeitschr. 1913 p. 212 ♀ ♂.

"*Worker*. Length 2.5—3 mm.

Color brown, varying from blackish brown to uniform reddish brown, with the mandibles and base of the scapes reddish. Moreover, the head is more trapezoidal, more enlarged posteriorly than in the other subspecies, a little longer than broad, but not much. Finally, joints 3–10 of the funiculus are a little thicker than in the typical *alfari*, rather distinctly broader than long. Apart from these characters I find none that would suffice for the establishment of a species.

Manaos, Brazil (A. Goeldi and Dr. Huber), in a *Cecropia*."

To the characterization of the worker Forel later added the following remarks and a description of the sexual phases: "Between the mesonotum and basal surface of the epinotum there is a sharp, slit-like incision, which descends abruptly from the basal surface and bears on its anterior (mesonotal) wall the two very prominent stigmata. This conformation is indicated also in other varieties (especially *mixta*) but only feebly."

"*Female*. Length 6—6.6 mm.

Color as in *mixta* and scarcely distinguishable from that form. The head is merely somewhat shorter and somewhat broader behind. Distinguishable from the specific type only by the shorter scape."

"*Male*. Length 2.8—3 mm.

Precisely like the var. *mixta*, but the antennæ thicker and stouter; the funicular joints 4–10 broader than long or at least as broad as long.

Campo Besso, near São Paulo (H. von Ihering), in a swamp *Cecropia*."

Forel records this subspecies also from Panama (Christophersen), "nesting in the cavities of a *Cecropia* trunk"; and Stitz saw specimens taken by E. Ule in Brazil. I received a fine series of workers together with branches of the *Cecropia peltata* in which they were living from Dr. G. Stabel, Director of the Agricultural Experiment Station at Paramaribo, Dutch Guiana, and all three phases taken by Mr. H. E. Box in *Cecropia peltata* at Blairmont Plantation, Berbice, British Guiana.

Subsp. LUCIDULA Forel

A. alfari race *lucidula* Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 113 ♂ ♀ ♂, Wheeler, Amer. Mus. Novitates No. 45, 1922 p. 14 ♂ ♀; *A. alfari* Bailey, Bot. Gazette 74, 1922 p. 379; *A. lucidula* Forel, Mém. Soc. Ent. Belg. 20, 1912, p. 51.

"*Worker*. Length 2.6-2.9 mm.

Much more shining, more feebly sculptured than the type. Thoracic impression broad and deep; mesonotum and epinotum slightly convex. Head narrower, slightly more elongate. Stature a little more slender. The funicular joints broader than long, especially towards the tip."

"*Female*. Length 7.5 mm.

Densely punctate, scarcely shining. Very similar to the female of *A. bicolor* and of the same black color (see Emery's figure), but the sides of the head are more convex; the head is also as broad in front as behind. Scale lower and thicker than in *A. bicolor*. Thorax a little broader. The wings are almost hyaline, with the stigma brown and the veins yellowish brown (in *A. bicolor* the wings are tinged with brownish yellow).

"*Male*. Length 2.5-2.9 mm.

Mandibles very short, pointed, not meeting in the middle. Head rounded rectangular, longer than broad. Scape broader than long, as is also the first funicular joint. Funicular joints 6-10 nearly as broad as long. Petiolar scale thin. Dorsal appendage of the median genital valvules surpassing the external valvules, which are acuminate. Wings rather long, subhyaline; in other respects as in the female.

"Hab. Guatemala, Retalhuleu (Stoll); Trinidad (Urich)."

I have seen specimens of this ant from the following localities:

British Guiana: Kartabo and Kalacoon (Wheeler), nesting in the internodes of *Cecropia angulata* Bailey.

Trinidad: Caroni and Botanical Garden, Port of Spain (Wheeler), in internodes of *Cecropia peltata*.

Var. ZONALIS var. nov.

Worker. Length 2-2 mm.

Very close to the typical *lucidula* and with the same sculpture, but smaller. Feebly polymorphic, the antennal scapes shorter, in the worker minor reaching to the occipital corners of the head, in the worker major about two-thirds the distance between the eyes and occipital corners. Pubescence distinctly more abundant on the head and thorax than in *lucidula*. Color varying from brownish yellow to

yellowish brown, the head usually somewhat darker. Mandibles usually not much darker than the head, clypeus paler.

Female. Length scarcely more than 6 mm.

Also smaller than the female of *lucidula*, of the same color and sculpture but the wings are distinctly darker and the posterior borders of the gastric segments less distinctly brownish. The head is less concave behind, with more rounded occipital corners.

All the numerous specimens which I have taken in young *Cecropias* in the Canal Zone and the Republic of Panama belong to this variety. I have specimens from the following places: Corrozal (type-locality), Ancon, Gatun, Monte Lirio, Colon, Rio Agua Salud, Red Tank and Otoque Island.

Subsp. LUCIDA Forel

A. alfari race *lucida* Forel, Biol. Centr. Amer. Hymen. **3**, 1899-1900 p. 113♀ ;
A. lucida Forel, Mém. Soc. Ent. Belg. **20**, 1912 p. 51.

Worker minor.

Identical with the worker of the preceding race (*lucidula*), but the thorax is more arcuate (more like that of the type), the color duller and the funiculi, excepting the first joint brown."

Worker major. Length 4.2 mm.

Head 1.6 mm. long and 1.25 mm. broad, very similar to Emery's figure of the major worker of *A. bicolor*, but narrower in front and a little more elongate. The scapes fail to reach the occipital angles by half their length. Funicular joints, except the penultimate, a little longer than broad. Head deeply excavated behind with feebly convex sides. Borders of gastric segments yellowish. Reddish yellow; head reddish brown, except anteriorly where it is reddish. Gaster and funiculi, except the first joint, brownish. Gastric segments brownish posteriorly. Head feebly pubescent, very shining, very feebly punctate, the punctures numerous but superficial.

Hab. Guatemala, Pantaleon 1700. (Champion) Found in the cavities of a *Cecropia* trunk."

I have taken what I regard as this subspecies in young *Cecropias* at Patulul and Escuintla, Guatemala. The workers agree well with Forel's description, except that in the major the head is rather broad, with convex sides. In none of the workers do the scapes surpass the occipital border. Two winged females from Patulul are very pale in color and are, perhaps, immature. They are yellowish brown, with the dorsal surface of the thorax and the bases of the

gastric segments darker, the head, antennæ and legs brownish yellow, the mandibles deep red. The scapes reach less than half the distance between the eyes and occipital corners. The surface is much less shining than in the worker, though the pubescence is poorly developed. The wings are tinged with yellow, the veins resin-yellow, the stigma brown. The size is unusually small, scarcely 6 mm., but the gaster is much contracted in both specimens.

Subsp. TUBEROSA Forel

Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 240 ♀.

“*Worker*. Length 2.5—3.9 mm.

Form of the head as in *aequilata*, but a little longer and only feebly excised behind. The eyes are larger than in the varieties of the typical subspecies. The mesoepinotal impression is strong; basal surface of epinotum convex, enlarged posteriorly and bearing at its extremity two large obtuse tubercles formed by the stigmata. The scale is cuneiform as in the typical *alfari* and its varieties (sharper than in the subsp. *lucida*). The penultimate joints of the funiculi are as long as broad (thicker than in the typical *alfari* and its varieties).

Sculpture and pilosity of the typical *alfari*. Brownish yellow; upper surface of head in great part, mandibles, tarsi and tibiæ brown. Sometimes the brown color extends to a part of the thorax or even of the abdomen as well as the femora.

“*Female*. Length 9 mm.

Blackish brown; anterior portion of head and antennæ reddish. Very similar to the female of the var. *aequilata*, but larger, with broader and shorter head. Scale elevated and acuminate. Wings lacking.

Ceara, Brazil (Diaz da Rocha).”

AZTECA ANGUSTICEPS Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna 1893, p. 26 ♀ figs. 61–63; Wheeler, Amer. Mus. Novitates No. 45 1922, p. 14 ♀.

Costa Rica: (type locality); Alajuela (Wheeler) nesting in Tillandsias.

Trinidad: Botanical Garden, Port of Spain (Wheeler), running on the trunks of *Cecropia peltata* L.

AZTECA BICOLOR Emery var?

What seems to be a variety of this species is represented by a single dealated female and her incipient brood of three small black workers, taken from a *Cordia alliodora* domatium at Ancon, C. Z. I hesitate to introduce a new name because the large workers are unknown and the female possesses no striking characters.

AZTECA BREVICORNIS Mayr

Liometopum brevicorne Mayr, Verh. zool. bot. Ges. Wien p. 870 ♀; Emery, Mem. R. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 345:♀, Fig. 76-78.

Brazil: Amazonas (James Trail), probably in some myrmecophyte.

Var. BOLIVIANA Wheeler & Mann, var. nov.

Worker. Length 2—2.2 mm.

Agreeing very closely with the descriptions of Mayr and Emery of the typical *brevicornis*, except in color, which is dark brown, the gaster and posterior portion of the head being black, the mandibles red, the corners of the clypeus, the antennal scapes, first funicular joint and legs sordid yellowish, the femora fuscous in the middle, the posterior borders of the gastric segments pale brown.

Six specimens taken by Dr. W. M. Mann at Huachi Beni, Bolivia, in the hollow stems of a species of *Triplaris* (White, No. 958).

AZTECA COERULEIPENNIS Emery

Emery, Mem. Re. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 330 ♀ ♀ ♂ Figs. 1-11; Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900, p. 110; Ross, Naturw. Wochenschr. 8, 1909 p. 829.

Costa Rica: Pacific Slope (A. Alfaro), in trunks of *Cecropia*; Zent, Atlantic Slope (Wheeler), in *Cecropia*.

Mexico: Atoyac in Vera Cruz (Schumann); Zacuapam (H. Ross), in *Cecropia mexicana*.

Guatemala: Cerro Zunil, 4,000-5,000 (Champion).

AZTECA CONSTRUCTOR Emery

Emery, Bull. Mus. Zool. Anat. Comp. Torino 11, 1896 p. 2 figs. II, III; Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 110; Mém. Soc. Ent. Belg. 20, 1912 p. 50 ♀; Wheeler, Amer. Mus. Novitates No. 45 p. 14 ♀ ♀.

Costa Rica: Atlantic and Pacific Slopes (A. Alfaro, Pittier, Tonduz), in *Cecropias*.

Guatemala: Quirigua (Wheeler), in young *Cecropias*.

Panama: (Christophersen); Ancon (Wheeler), in young *Cecropias*.

Trinidad: Port of Spain (Wheeler), in internodes of *Cecropia peltata*.

According to Alfaro, as quoted by Emery, *A. constructor* is "the fiercest and most aggressive of the Costa Rican species," a statement to be accepted "*cum grano salis*." The opening of the nest in the cavity of the *Cecropia* "is a fissure of 15 mm. In the cavity of the plant it constructs a nest of brown carton."

Var. GUIANÆ var. nov.

A. constructor Bailey, Bot. Gazette 74, 1922 p. 377.

Worker. Length 2-3 mm.

Smaller, head distinctly longer and somewhat more rectangular than the typical *constructor*, distinctly longer than broad, the posterior border less deeply excised; the epinotum distinctly angular in profile, the petiolar scale somewhat sharper and even more inclined forward. The color is somewhat darker, more blackish brown, the pilosity on the body somewhat less abundant and more uneven.

Female (deälated). Length 7 mm.

Also smaller than the typical *constructor*, which measures 7.5-8 mm. Head larger and broader. Surface distinctly less shining. Dark brown, not black, like the typical form; pilosity similar but pubescence grayish and much more abundant and conspicuous, especially on the head and gaster.

Described from nine workers and a single female taken at Kartabo, British Guiana, in an internode of *Cecropia angulata* Bailey.

AZTECA COUSSAPOÆ Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 700 ♀.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in twigs and branches of *Coussapoa* sp. (Ule No. 5717).

AZTECA DELPINI Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 333, ♀ fig. 30-32.

This species was originally described from Matto Grosso, Brazil (Germain). Mr. H. O. Lang has sent me workers of the typical form

of the species which he took at Kamakusa, British Guiana, in large leaf-sacs of *Tococa guianensis* Aubl. A set of larger and darker workers was taken by me at Kartabo, British Guiana in a cauline swelling of *Cordia nodosa* L.

Var. TRINIDADENSIS Forel

Wheeler, Amer. Mus. Novitates No. 45, 1922 p. 14 ♀ ♀.

Trinidad: Caroni (Wheeler), nesting in internodes of *Cecropia peltata* L.

AZTECA DEPILIS Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 339 ♀ ♀ ♂ Fig. 37-42; Schumann, Samml. gemeinverst. Wiss. Vortr. 83, 1889 p. 17.

Brazil: Amazonas (K. Schumann), from stems of *Duroia hirsuta* Schum. and leaf-sacs of *Tococa coronata* Benth.

AZTECA DUCKEI Forel

Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 243 ♀ ♀ Fig. 2.

Brazil: Barcellos, Rio Negro, Amazonas (Ducke), in cauline swellings of *Cordia nodosa* L.

AZTECA DUROLE Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 697 ♀.

Brazil: Jaburu, Upper Jurua, Amazonas (E. Ule), in swellings of twigs of *Duroia hirsuta* Sch.; Jurua (E. Ule), in the branches of *Porouma* sp.; Porto Alegre, Rio Purus, Amazonas (A. Goeldi and J. Huber), in the hollow stems of *Duroia hirsuta*.

AZTECA EMERYI Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 698 ♀ ♀.

Brazil: Cachoeira, Jurua, Amazonas (E. Ule), in the hollow internodes of *Cecropia sciadophylla* Mast.

AZTECA FASCIATA Emery

subsp. LETA subsp. nov.

Female. (deälated) Length nearly 8 mm.

Differing from the typical *fasciata* of Bolivia in the shape of the head and petiole. The former is one and one-half times as long as

broad, and regularly oblong, except for the feebly and angularly excised posterior border, the sides being straight and parallel, not feebly convex as in the typical form of the species. The petiolar node is not acuminate above as in that form, but blunt and rounded, though narrower above than below.

Yellow; head faintly reddish, with black ocellar spot; mandibles deep red. Posterior border of scutellum and a few clouds on the pleura fuscous. Each gastric segment dorsally with a transverse castaneous band and ventrally with a pair of brown spots. Last tarsal joint of all the legs blackish; middle and hind tibiæ and hind femora dark brown.

Described from a single specimen taken July 9, 1924 from a domatium of *Cordia alliodora* on Barro Colorado I., C. Z.

AZTECA FORELI Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 337 ♀ Fig. 50, 51.

The types of this species were collected by A. Alfaro at Bagaces, on the western slope of Costa Rica. I have taken it at Zent, on the Atlantic Slope of the same republic and at Patulul, Guatemala, in both places in *Cecropias*.

Var. XYSTICOLA Forel

Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 111 ♀.

Populous colonies of this vigorous and aggressive ant, originally described from Colombia, were found nesting in the trunks of old *Cordia alliodora* trees near the laboratory at Ancon, C. Z., Panama.

Subsp. CHAMPIONI Forel var. BREVISCAPA Forel

Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 112 ♀.

Costa Rica: (Tonduz), in *Cecropia*.

Subsp. URSINA Forel

Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 112 ♀.

Forel described this subspecies from specimens taken by Urich in Trinidad. I found it in the Botanical Garden at Port of Spain, running on the trunks of large *Cecropias* and probably nesting in their cavities.

AZTECA FOVEICEPS Wheeler

Wheeler, Zoologica **3**, 1921 p. 163 ♀ ♀ ♂ Fig. 16; Bailey, Bot. Gazette **75**, 1923 p. 34.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

AZTECA GOELDII Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906 p. 245 ♀ .

Brazil: Porto Alegre, Upper Rio Purus, Amazonas (J. Huber), in hollow branches of a tree (Laurineæ?), smelling strongly of onions.

AZTECA HUBERI Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906 p. 244 ♀ .

Brazil. Teffé, near Para (J. Huber), in branches of *Platymischium* sp.

AZTECA HYPOPHYLLA Forel

Forel, Biol. Centr. Amer. Hymen. **3**, 1899-1900 p. 120 *nota*. ♀ .

Colombia: San Diego (A. Forel). "Makes its nest under the round leaves of a climbing plant which from the description is very probably the juvenile stage of a species of *Marcgravia*."

AZTECA INSTABILIS (F. Smith)

Tapinoma instabile F. Smith, Trans. Ent. Soc. London (3) **1**, 1862 ♀ ; Emery, Mem. Accad. Sc. Bologna (5) **3**, 1893 p. 335, fig. 16-21; Bailey Bot. Gazette **74**, 1922 p. 379.

British Guiana: Kartabo (Wheeler), nesting in internodes of *Cecropia angulata*.

This species is known from Mexico, Central America, Colombia and French Guiana. Most of the citations of *instabilis*, however, refer to *muelleri* Emery of Brazil.

AZTECA LANUGINOSA Emery

Emery, Mem. Acad. Sc. Ist. Bologna (5) **3**, 1893, p. 341 ♀ fig. 47; H. v. Ihering, Engler's Bot. Jahrb. **39**, 1907 p. 692; Forel, Verh. Zool. bot. Ges. Wien 1908, p. 389 ♀ ♀ ♂, fig. of nest.

Brazil: Santa Catharina (type-locality); São Paulo, etc. (H. v. Ihering), on *Cecropia adenopus* Mart.; Jaragua Distr., Prov. Santa Catharina (W. Ehrhardt), carton nest on *Cecropia*.

AZTECA LONGICEPS Emery

(Plate 55, fig. a)

Emery, Mem. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 28 ♀, fig. 64, 65.

Worker major. Length 3—3.5 mm.

Head suboblong, one and one-third times as long as broad, narrowed in front of the eyes, the sides in the middle parallel and nearly straight, the posterior border distinctly but not very deeply excised. Eyes small, flat, near the anterior third of the sides of the head; ocelli absent. Clypeus swollen and somewhat advanced at the sides, longitudinally flattened in the middle, with nearly straight, entire anterior border, the posterior suture indistinct. Frontal area and groove obsolete. Mandibles rather small and convex, with 7-8 teeth. Antennal scapes reaching to half the distance between the eyes and the posterior corners of the head; funiculi but very slightly thickened towards their tips; first joint about one and one-half times as long as broad; second joint longer than broad, remaining joints, except the last, subequal, broader than long. Thorax stout and thickset, about as long as the head including the mandibles. Mesonotum convex and rounded, but not gibbous; mesoepinotal constriction pronounced; epinotum small and low, as long as broad, its base in profile rather straight, longer than the sloping declivity. Petiolar node strongly inclined forward, rather high, its summit thin but not acute, seen from behind narrowed and rounded above, its ventral surface convex behind in profile. Gaster broadly elliptical, not much longer than broad. Legs short, with the femora distinctly compressed.

Shining; mandibles finely shagreened and with a few coarse punctures along their apical borders; remainder of the body finely and superficially punctulate.

Hairs yellowish, moderately long, erect, most abundant on the upper surface of the body, sparse on the scapes and tibiae. Pubescence fine, grayish, most conspicuous on the head, gaster and appendages.

Dark reddish brown or blackish, thorax usually somewhat paler than the head and gaster; cheeks, anterior border of clypeus and tarsi, except the terminal joint, and borders of gastric segments pale brown.

Worker minor. Length 2—2.5 mm.

Head only one and one-fourth times as long as broad and more distinctly narrowed in front, its sides more rounded, its posterior border less excised. Eyes just in front of the middle of the head.

Antennal scapes reaching to two-thirds the distance separating the eyes from the posterior corners of the head; funiculi slender, joints 3-10 proportionally shorter than in the worker major.

Female. Length 5.5-6 mm.

Head one and three-fourths times as long as broad, regularly oblong, with straight, parallel sides, scarcely narrowed in front of the eyes and nearly straight posterior border. Ocelli rather small; eyes moderately large, feebly convex, as long as their distance from the corners of the clypeus, their posterior orbits well in front of the middle of the head. Mandibles large, convex at the base, flattened and deflected at the tip, with 8 subequal teeth. Clypeus as in the worker major. Thorax elliptical, through the wing-insertions a little broader than the head. Epinotum sloping, with feebly developed base and declivity, the latter short. Petiolar scale much as in the worker but higher, its upper border more produced, seen from behind trapezoidal and narrowed above. Gaster elongate, suboblong. Legs long; femora, especially the fore pair, strongly compressed.

Sculpture and color as in the worker; pilosity less abundant, but the flexor surfaces of the scapes and tibiae with a few erect hairs. Mandibles entirely black, corners of clypeus and tarsal joints, except the last, red. Wings grayish hyaline, with brown pterostigma and pale yellow veins.

Male. Length 2.5 mm.

Head, excluding the eyes, longer than broad, rounded behind. Eyes half as long as the sides of the head, moderately convex; ocelli small. Mandibles very small, not meeting, with only the terminal tooth, which is acute and acuminate. Antennæ short; scapes not longer than broad; first funicular joint broader than long; second joint longer than broad, succeeding joints, except the last, as broad as long. Thorax large, much broader than the head; epinotum small and short, very convex, without distinct base and declivity. Petiolar scale erect, anteroposteriorly compressed, with blunt, broadly rounded summit. Legs slender.

Smooth and shining. Hairs almost, pubescence quite lacking, except on the funiculi where it is very dense and erect.

Black; mandibles, antennæ, legs and genitalia piceous. Wings colored as in the female, but the pterostigma paler.

Described from numerous specimens taken in the cauline swellings of *Cordia alliodora* in the following localities in the Canal Zone: Frijoles, Ancon, Red Tank and Las Cascades. This is by far the most abundant ant in the Cordias, sometimes occupying all or most

of the cauline swellings. It is rather timid and, though sometimes inclined to bite, is not very offensive.

I have redescribed all four phases of the species, because it has been known hitherto only from Emery's figures and brief description of the dealated female taken at Alajuela, Costa Rica, by A. Alfaro. That my identification is correct is shown by the perfect agreement of the head and petiole of the female with Emery's drawings, and the fact that I possess two precisely similar females taken by Messrs. E. A. Schwarz and Barber at Livingston and Trece Aguas, Alta Vera Paz, Guatemala. Forel has seen a female from Mexico. He has also described from various parts of tropical America the following four forms of *longiceps* which I treat as subspecies:

Subsp. CORDINCOLA Forel

Forel, Bull. Soc. Bot. Genève (2) 11, 1920 p. 203 ♀.

This form was originally described from specimens taken by Bang at Cochabamba, Bolivia, in the cauline swellings of a *Cordia* (probably *alliodora*). Dr. W. M. Mann has given me five workers and a dealated female which he found in cauline swellings of *C. alliodora* var. at Ivon, Beni, and Huachi Beni, Bolivia.

In the worker major, which measures about 2.5 mm., the head is long, rather deeply excavated behind and somewhat narrowed anteriorly. The scapes extend somewhat more than two-thirds the distance between the eyes and posterior corners of the head, the penultimate funicular joints are as long as broad. The base of the epinotum passes abruptly into the short declivity. The body is reddish brown, the thorax somewhat paler than the head and gaster, the mandibles red, with dark bases. The pilosity is very sparse, absent in the scapes and legs.

The worker minor measures only 2 mm. and is very similar to the major except that the head is shorter and more narrowed anteriorly and the scapes reach further towards the posterior corners of the head.

The *female* (undescribed) measures nearly 6 mm. and is black, with dark red cheeks and clypeal border; the tarsi, except the last joint, testaceous. The head is fully twice as long as broad and deeply emarginate posteriorly. The mandibles are 8-toothed. Antennal scapes extending half the distance between the eyes and posterior corners of the head, the median joints of the funiculi distinctly broader than long. The epinotum is rounded and sloping, the petiolar scale seen

from behind with a bluntly pointed summit. The erect pilosity is sparse as in the worker and absent on the appendages.

Subsp. JURUENSIS Forel

A. longiceps var. *juruenis* Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 699, ♀ ♀ ♂.

All the phases of this form were taken by E. Ule at Jurua Miry, Jurua, Amazonas, in the twigs and branches of a Leguminose plant, *Schwartzia*. Dr. Mann has given me three major workers which he collected in cauline swellings of *Cordia alliodora* var. *b liviana* at Ivon Beni, Bolivia. I also possess a worker minor cotype given me by Prof. Forel.

The head of the major worker is one and one-half times as long as broad, distinctly emarginate behind. The antennal scapes reach somewhat more than half the distance between the eyes and the posterior corners of the head. The epinotum is subcuboidal, slightly longer than broad, the base longer than the abrupt declivity. The pilosity is somewhat more abundant than in the subsp. *cordincola* and there may be a few hairs on the scapes, but none on the legs. Reddish brown, head and gaster darker than the thorax. Mandibles red, darker at the base. Length 3.4 mm.

The worker minor measures 2–2.5 mm, and has a shorter, anteriorly more narrowed and posteriorly less deeply excised head, with the scapes reaching the posterior fourth of the head.

The female is somewhat smaller than the typical *longiceps* and the subsp. *cordincola*, measuring only 5–5.2 mm. The head is twice as long as broad. Color as in the typical form, but the pilosity feebler than in the worker. Wings brownish.

The male measures 2.9 mm. The head is somewhat longer than broad, the scape scarcely longer than the first funicular joint. Black, with the antennæ, mandibles and legs pale sordid yellow.

Subsp. PATRUELIS Forel

Forel, Verh. zool. bot. Ges. Wien 1908 p. 392 *nota* ♀ ♀ ♂.

Forel described all the phases of this subspecies from specimens which I sent him and which were taken by Mr. C. H. Tyler Townsend near Colima, Mexico, in the cauline swellings of *Cordia alliodora* var. I insert a translation of the original description.

“Length of worker 2.2—3 mm.

Worker major. Distinguished as follows from the subsp. *juruensis* Forel: Head somewhat broader, one and two-fifths times as long as broad, not broader behind than in front, with somewhat more convex sides. Eyes at the anterior third. The antennal scape reaches the posterior fourth of the head. Promesonotum gibbous, much more protuberant and higher than in *juruensis*, where it is moderately convex and hardly higher than the base of the epinotum. The latter rather low and almost flat. Constriction not deep. Petiole as in *juruensis*, but sharper above, angular in profile. Sculpture and pilosity as in *juruensis* but with more bristly hairs on the legs. Brownish, yellowish red, mandibles darker reddish, gaster brown, with yellowish borders to the segments. Legs as in *juruensis*, short, rather stout, with feebly compressed femora. Otherwise precisely as in *juruensis*.

Worker Minor. Head one and one-fourth times as long as broad. The antennal scapes reach nearly to the posterior sixth of the head. Eyes somewhat in front of the middle of the sides. In other respects like the worker major.

Female. Length 6 mm. Larger than *juruensis* and like the typical form of the species. Head 1.7 mm long and 1 mm broad, posteriorly a little broader than anteriorly. The scapes reach nearly to the posterior third of the head; funicular joints somewhat less transverse than in the typical *longiceps*. Petiole slightly thicker and above somewhat blunter.

Sculpture somewhat denser than in *juruensis* and the typical *longiceps* and therefore less shining. Pubescence somewhat denser, otherwise with the same pilosity. Color as in *juruensis*, but somewhat paler dark brown (in the typical *longiceps* darker, almost black). Posterior borders of gastric segments more broadly yellow. The whole clypeus and the cheeks reddish; mandibles and antennæ reddish brown. Wings slightly more brownish."

An examination of the cauline swellings of *Cordia* that contained the types of this subspecies shows that they were inhabited by the same large black *Coccids* as those found with the typical *longiceps* in Panama.

Subsp. SAPII Forel

Forel, Mém. Soc. Ent. Belg. 20, 1912 p. 56 ♀ ♀.

Forel described the worker and female of this subspecies from specimens taken by Ducke at San Antonio de Ica, Amazonas, Brazil, in the hollow stems of *Sapium glandulosum*. Their occurrence in this plant is not mentioned by Forel but is indicated on the label of a

female and three worker co-types which he gave me. I translate his description:

Worker. Length 1.7—2.8 mm.

Mandibles shining, punctate, very feebly and finely reticulate, slightly curved, armed with about 7 teeth. Anterior border of clypeus slightly concave and bisinuate. Head subdepressed, 0.9 mm. long and 0.7 mm. broad in the worker major, 0.65 mm. long and 0.47 mm. broad in the worker minor, of the same shape in both, rectangular, slightly broadened behind, with feebly convex sides and scarcely concave posterior border. Eyes scarcely behind the anterior third. The scape slightly surpasses the posterior fifth of the head in the minor and is almost as short as in the major worker. Second funicular joint at least as broad as long; joints 5–10 much broader than long (about one and one-half times; the last joint at least twice). Thorax large and robust; pronotum one and one-half times as broad as long, but without humeri; mesonotum large, rounded, subdepressed; the promesonotum moderately convex. Thoracic constriction feeble and narrow; basal surface of epinotum square, subdepressed, scarcely longer than the declivity. Petiole short and thick, about as in *stolli* and *polymorpha*, but its very short anterior surface is more convex. Femora quite as dilated as in *hypophylla*, but proportionally shorter; fore tibiae dilated and also compressed.

Shining; very finely punctate; punctuation very sparse. Pubescence rather long and rather dense. Erect pilosity short, scattered over the body, very sparse on the appendages, also on the tibiae and scapes.

Body and appendages uniform reddish brown. Gaster brown, with the borders of the segments sordid yellow.

Female. Length 5.2—5.4 mm.

Head absolutely rectangular, depressed, 1.2 mm. long and 0.6 mm. broad, with straight posterior border and much rounded posterior corners. Eyes at the anterior fourth. The scape reaches the third fifth of the head from the front. Thorax compressed, long, scarcely broader than the head, feebly convex anteroposteriorly. Petiolar scale cuneiform, higher than in the worker.

Brownish black; appendages, mandibles and anterior portion of head reddish. Wings feebly but very distinctly tinged with brown.

AZTECA LUEDERWALDTI Forel

Forel, Deutsch. Ent. Zeitschr. 1909 p. 261 ♀.

Paraguay: San Bernardino (K. Fiebrig), in branches of *Alchornea irucurana*.

AZTECA MINOR Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 696 ♀ ♀; Ule, Ameisenpflanzen des Amazonasgebietes 1907.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in the hollow internodes of *Cecropia ficifolia* Warb.

AZTECA MUELLERI Emery

A. instabilis Fr. Müller, Jen. Zeitschr. Naturw. **10**, 1876 p. 281 (*nec* Smith, *nec* auct.); Kosmos 8, 1880, p. 109; Huth, Monatl. Mitteil. Naturw. **4**, 1887 p. 319; Emery, Zeitschr. wiss. zool. **46**, 1888, p. 390 pl. 28 fig. 25-28 Schimper, Wechselbez. Zwischen Pflanzen Ameisen 1888; H. V. Ihering, Berl. Ent. Zeitschr. **39**, 1894 p. 379.

A. muelleri Emery, Mem. Accad. Sc. Ist. Bologna (5) **3**, 1893 p. 331 ♀ ♀ figs. 12-15; H. v. Ihering, Engler's Bot. Jahresb. **39**, 1907 p. 677 et seq.; Emery, Scientia **12**, 1912 p. 50; Stitz, Deutsch. Ent. Zeitschr. 1913 p. 212 ♀.

A. nigella Emery, Mem. R. Accad. Sc. Ist. Bologna (5) **3**, 1893 p. 332 ♀ fig. 23, 24; H. v. Ihering, Engler's Bot. Jahresb. **39**, 1907 (incipient nest form of *muelleri*).

Brazil: Santa Catharina (F. Müller, Hetschko), in *Cecropia adenopus* Mart.; Rio Grande do Sul, São Paulo, etc. (H. v. Ihering) in *Cecropia* (probably *adenopus*); Rio de Janeiro, in *Cecropia adenopus*; Alto Boa Vista, Rio (J. C. Bradley) in *Cecropia*; Alto Acre (E. Ule), in one of the Bombaceæ.

Var. WACKETI Emery

In Forel, Verh. zool. bot. Ges. Wien 1908 p. 391 *nota* ♀ ♀ ♂; H. v. Ihering, Engler's Bot. Jahrb. **39**, 1907, p. 710.

Brazil: São Paulo, Ilha Victoria and Estancia São Paulo (H. v. Ihering), making carton nests in *Cecropia*. São Paulo (H. v. Ihering), in hollow stems of *Erigeron maximus* Link.

AZTECA OLITRIX Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 693 ♀ ♀; Bull. Soc. Bot. Genève (2) **11**, 1920 p. 3 ♀.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in ant-gardens among Gesneriaceæ with galls; Para (J. Huber), in cauline swellings of *Cordia* sp.

AZTECA PITTIERI Forel

(Plate 55, fig. b)

Forel, Biol. Centr. Amer. Hymen 3, 1899-1900 p. 120 ♀ Fig. 16.

Female (undescribed). Length 5—5.5 mm.

Head oblong, one and two-thirds times as long as broad, slightly narrower in front than behind the eyes where the sides are nearly straight and parallel, posterior border feebly excised. Eyes small and rather flat, as long as their distance from the anterior corners of the head, the posterior orbits in front of its middle. Ocelli small. Mandibles rather large, convex, elongate, with 8 teeth. Clypeus with a longitudinal flattening in the middle, somewhat swollen and advanced at the sides, the anterior border entire, nearly straight, the posterior suture indistinct. Antennal scapes reaching to two-thirds the distance between the posterior orbits and the posterior corners of the head; funiculus scarcely enlarged at the tip, first joint a little longer than broad, second as long as broad, remaining joints, except the last, subequal, distinctly broader than long. Thorax elongate-elliptical, through the wing-insertions as broad as the head; epinotum rounded and sloping, without distinct base and declivity; petiolar scale erect, acuminate but not very acute. Legs long, scarcely compressed.

Sculpture much as in the worker; gaster and legs more shining and more finely punctate than the head and thorax; mandibles shining, sparsely punctate.

Hairs white, fine, not very long, but moderately abundant, erect on the body and appendages, sparser on the latter. Pubescence grayish and very fine, covering the body and appendages, most distinct on the head.

Black; anterior border of clypeus, apical borders of mandibles, except the teeth, tarsi and insertions of antennæ, red; posterior borders of gastric segments brownish. Wings grayish hyaline, with yellow veins and dark brown pterostigma.

Small colonies of this species, originally described from Buenos Aires, Costa Rica (Pittier), were found nesting in the cauline swellings of *Cordia alliodora* along the Tumba Muerte Road near Las Sabanas, in the Panamanian Republic and along the Chivachiva Trail, near Red Tank in the Canal Zone. The species seems to be rare compared with *A. longiceps*. Its relation to this species is so close that it hardly deserves more than subspecific rank. The largest workers in my collection have the head somewhat more elongate and somewhat less narrowed in front than in Forel's figure.

Var. EMARGINATISQUAMIS *Forel*

Forel, Bull. Soc. Bot. Genève (2) 11, 1920 p. 4 ♀.

This variety, described from workers taken by Pittier in Costa Rica, in the cauline swellings of *Cordia gerascanthus*, is distinguished from the type by having the petiolar scale emarginate at the summit, the eyes somewhat smaller, the legs a little more compressed and the head somewhat larger and broader behind in the major worker, with more convex sides.

AZTECA PRORSA sp. nov.

(Plate 56, fig. a)

Worker major. Length 2.5—3 mm.

Head somewhat depressed and flattened above and on the gular surface, one and one-half times as long as broad, narrower in front of the eyes, but behind them with straight, parallel sides, the posterior border rather deeply excised. Eyes small, flat, in front of the median transverse diameter of the head; ocelli absent. Mandibles rather small, convex, with 7-8 teeth. Clypeus convex in the middle and on the sides, its anterior border very feebly bisinuate, its posterior suture distinct. Frontal area and groove obsolete. Antennal scapes reaching to nearly two-thirds the distance between the eyes and the posterior corners of the head; funicular joints, except the first and last, somewhat broader than long. Thorax rather long, the mesonotum moderately convex, as long as broad; mesoepinotal constriction moderately deep, short, not very abrupt; epinotum as long as broad, its base and declivity sub-equal, the latter sloping but forming a distinct though rounded angle with the former. Petiole small, its scale inclined forward, narrow, elongate, elliptical, with blunt, rounded superior border. Gaster elongate-elliptical, the first segment short, truncated anteriorly. Legs rather long, feebly compressed, the fore femora somewhat dilated.

Mandibles shining, sparsely punctate; body and appendages densely and finely punctulate, shining, the legs somewhat less so than the body.

Hairs whitish, very delicate, erect, not very abundant on the upper surface of the body, shorter but distinct on the tibiae and especially on the scapes. Pubescence pale, fine and dilute, distinct on all parts of the body but only slightly dimming the shining surface.

Reddish brown; thorax, legs, anterior part of head, scapes, first

funicular joint and borders of gastric segments, paler; mandibles deep red, with black teeth; cheeks, anterior border of clypeus, and tarsi, except the terminal joint, yellowish.

Worker minor. Length 1.5—2.5 mm.

Head smaller than in the worker major, but with much the same proportions, slightly more narrowed anteriorly, the sides a little more rounded, the occipital border less deeply excised. The funicular joints are shorter and more transverse. Mesonotum more depressed, flatter and less projecting; mesoëpinotal constriction less pronounced; petiolar scale proportionately thicker and more obtuse.

Sculpture, pilosity and color as in the worker major, but the mandibles have a large dark brown spot externally at the base.

Described from numerous specimens which I took during December 1911, at Escuintla, Guatemala, in the hollow twigs of *Triplaris auriculata*.

This form is closely allied to *A. longiceps* but is smaller, the head is longer, more deeply excised behind and with straight sides, the scapes shorter, the epinotum more sloping and less cuboidal, the hairs finer and less abundant.

AZTECA SERICEA (Mayr)

Iridomyrmex sericeus Mayr, Sitzb. Akad. Wiss Wien **53**, 1866 p. 498, fig. 8;

Azteca sericea Emery, Mem. R. Acad. Sc. Ist. Bologna (5) **3**, 1893 p. 334
♀ fig. 25—29.

Mexico: (Heller), in pseudobulbs of the orchid *Schomburgkia tibicinis*. Also recorded from Guatemala (Stoll) and Cayenne (Forel).

I have taken workers and dealated females of this species at Zent, Costa Rica, and at Patulul and Escuintla, Guatemala, in the internodes of young *Cecropias*.

The female resembles that of *coeruleipennis* Emery, but the head is more rectangular, although of the same proportions, the sides are more nearly straight, the posterior and anterior corners sharper, the posterior border more broadly and less deeply excised, the eyes less convex, the antennal scapes somewhat longer, the petiolar scale less acute above in profile, the femora broader. The surface is shining as in *coeruleipennis*, the mandibles subopaque, shagreened and sparsely punctate. The pale erect hairs are abundant but not very long, the pubescence very indistinct. Dark reddish brown, legs and antennæ paler; mandibles red; clypeus, bases of scapes and tarsi more yellowish red.

AZTECA SCHUMANNI Emery

Emery, Mem. R. Accad. Sc. Ist. Bologna (5) **3**, 1893 p. 345 ♀ ♀ fig. 72-75.

Venezuela: Guainia River, an affluent of the Cassiquiare, in the leaf-sacs of *Hirtella guainia* Hook fil.

AZTECA STANLEYULA Forel

Forel, Bull. Soc. Bot. Genève (2) **11**, 1920 p. 2 ♀.

Brazil: Para (J. Huber), in cauline swellings of *Cordia nodosa* L.

AZTECA TACHIGALLÆ Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 701 ♀.

Peru: Cerro de Escaler, 1000 m. (E. Ule), in the hollow petioles of an undetermined *Tachigalia*.

AZTECA THERESLÆ Forel

Forel, Berlin. Ent. Zeitschr. **44**, 1899 p. 273 ♀ fig.; Ann. Soc. Ent. Belg. **43**, 1899 p. 303 ♀.

This peculiar species was described and figured from three workers taken by Princess Theresa of Bavaria on the firewood of the steamer, while she was travelling on the Lower Magdalena River in Colombia. They were probably nesting in some myrmecophyte (*Triplaris?*). As all the specimens were of the same stature Forel was at first unable to decide whether they were major or minor workers. They evidently belonged to the former caste. In 1912 Forel described the following variety from workers taken by Christophersen on *Triplaris americana* in Panama. As it is one of the commonest ants on this tree I describe all four of its phases:

Var. MENCEPS Forel

(Plate 56, fig. b)

Forel, Mém. Soc. Ent. Belg. **20**, 1912 p. 54 ♀.

Worker major. Length 3—3.5 mm.

Head large, distinctly flattened, nearly one and one fourth times as long as broad, a little narrower in front than behind, with rounded sides, deeply and angularly excised posterior border and prominent, narrow and somewhat approximated posterior corners. Eyes small, flat, their posterior orbits just in front of the median transverse diam-

eter of the head. Ocelli absent. Mandibles stout, convex, with 7-8 teeth, the two apical large, the others small, subequal. Clypeus convex in the middle and on the sides, its anterior border feebly bisinuate, its posterior suture feeble but perceptible. Frontal area obsolescent; frontal groove absent. Antennal scapes reaching to or a little beyond half the distance between the eyes and the posterior corners of the head; funicular joints 2-10 distinctly broader than long. Thorax stout, the mesonotum convex and projecting but not gibbous, subcircular; mesoëpinotal constriction pronounced but short and abrupt; epinotum small and low, rounded and sloping, the base a little longer than the declivity into which it passes without a distinct angle. Petiolar scale small and thick, its upper margin narrowed, rounded and blunt. Gaster elliptical, less than twice as long as broad, the first segment short, truncated anteriorly. Legs, especially the femora, somewhat compressed and dilated.

Shining; mandibles sparsely, remainder of body more densely and more finely punctate, the punctures superficial and indistinct.

Pilosity yellowish, erect, not abundant, absent on the front and vertex, on the gaster confined to the posterior borders of the segments and the anterior truncated surface; shorter and finer on the head than on the thorax, absent on the appendages. Pubescence pale, very short and fine, covering the whole body like a delicate bloom but scarcely obscuring the shining surface.

Dark piceous brown; thorax often somewhat paler; mandibles deep red; clypeus anteriorly and laterally paler red; legs, funiculi and posterior borders of gastric segments brown, the tibiæ and especially the tarsi paler and more testaceous.

Worker minor. Length 1.8—3 mm.

Proportions of the head very much as in the worker major, but smaller and somewhat more narrowed anteriorly, the basal joints of the funiculi much shorter and more transverse, the mesonotum much less convex and the petiolar scale somewhat blunter.

Sculpture, pilosity and color very similar, but the head and mandibles smoother and more shining; the pubescence less distinct.

Female. Length 4.8—5 mm.

Head resembling that of the worker major but nearly one and one-half times as long as broad, with less convex sides and much less deeply excised posterior border, narrower in front of than behind the eyes, the latter longer than their distance from the clypeus. Ocelli small. Antennæ much as in the worker major. Mandibles large and convex. Thorax through the wing-insertions as broad as the head; mesonotum

subhexagonal, narrowed in front. Petiolar scale strongly inclined forward, its very flat posterior surface elongate-elliptical, the superior border much less blunt than in the worker. Gaster elongate, parallel-sided.

Sculpture, pilosity and color as in the worker major, but the erect hairs on the thorax are decidedly less abundant, the pilosity is somewhat longer and the mandibles are black. Wings grayish hyaline, with dark brown veins and pterostigma.

Male. Length 2.8 mm.

Head, including the eyes, as broad as long, the eyes prominent, longer than half the sides, the ocelli prominent. Mandibles very small, but meeting with their acute tips. Clypeus convex. Antennal scapes as long as broad; first funicular joint twice as long as broad, remaining joints shorter. Thorax through the wing-insertions broader than the head; mesonotum large and protuberant, as broad as long; base and declivity of epinotum more distinct than in the worker and female. Petiole similar to that of the female but the scale is lower and erect.

Smooth and shining. Pilosity and pubescence very feebly developed. Piceous black; antennæ and tarsi yellow; femora, tibiæ and genitalia sordid or brownish yellow; wings grayish hyaline, with pale brown veins and pterostigma.

Described from many specimens taken at Ancon, Balboa, Fort Clayton, Miraflores and Las Cascades, C. Z., in the hollow twigs and branches of *Triplaris americana*.

I have referred my specimens to Forel's var. *menceps*, described from the aforementioned workers taken by Christophersen, but they may belong to the typical form of the species. The differences mentioned by Forel are slight and their value somewhat dubious, owing to the small number of workers on which he based the species. I surmise that Mr. Christophersen may have taken his specimens either at Empire, where he lived (near Las Cascades), or while he was collecting with me in 1911 at the old town of Frijoles, which is now submerged.

AZTECA TRAILI Emery

Emery, Mem. Accad. Sc. Ist. Bologna (5) 3, 1893 p. 333 ♀ ♂ fig. 33-36;
Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 692 ♀; Wheeler, Zoologica 3,
1921 p. 163 ♀.

Brazil: Amazonas (James Trail), probably in a myrmecophyte; Manaus, Amazonas (E. Ule), in leaf-sacs of *Tococa bullifera* Mart. &

Schr. Bom Fim, Jurua, Amazonas and São Joachim, Rio Negro (E. Ule), in ant gardens among Gesneriaceæ.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.; Merumé Mouth (H. O. Lang) in the fistulose *Patima formicaria* Johnston (Rubiaceæ). The specimens collected by Mr. Lang comprise several very small workers and a dealated female. The former are very dark in color but this may be due to their belonging to an incipient colony. This seems to be the condition in several other species of *Azteca* (*xanthochroa*, *muelleri*, etc.) the workers growing paler as they increase in size in older colonies.

The female, taken by Mr. Lang, measures 7 mm. The head is scarcely longer than broad, broader behind than in front, with feebly rounded sides and slightly excised posterior border. The scapes reach nearly to the posterior corners of the head. The border of the petiolar scale is much compressed, produced upward and distinctly emarginate. The color is black, with the clypeus, cheeks, gula, antennæ and tarsi brownish yellow, the mandibles red, the knees reddish. The surface is smooth and shining, but less so than in the worker, because the very fine pubescence is denser; the pilosity is like that of the worker, the scapes and legs with delicate erect hairs.

Forel described with some misgivings what he took to be the female of this species from Amazonas, but the head of his specimen was like that of *alfari* and this is certainly not like the head in my specimen or in the female of the following variety:

Var. TILLANDSIARUM var. nov.

Worker. Length 2—3.4 mm.

Very feebly polymorphic. Differing from the typical *trilli* in having the head distinctly longer than broad in all the forms, the posterior border moderately and angularly excised, the antennal scapes extending even in the largest individuals one-fourth their length beyond the posterior border of the head. Eyes small, near the middle of its sides. Thorax with nearly straight dorsal outline, the mesoëpinal impression very feeble, the mesonotum scarcely projecting even in large workers. Petiolar scale low, scarcely inclined forward, with acute upper border, the ventral surface much more convex than in the typical *trilli*.

Sculpture and pilosity as in the type. Pale reddish brown, thorax, petiole and posteronodal portion of gaster somewhat darker.

Female. Length 8.5—9.5 mm.

Head slightly longer than broad, gradually narrowed in front, broadest through the posterior corners which are rectangular, the posterior border slightly sinuate. Eyes small, as long as their distance from the anterior border of the head, their posterior orbits at its median transverse diameter. Mandibles convex, 8-toothed; clypeus with straight anterior border. Frontal area and groove obsolete. Antennal scapes not reaching the posterior border of the head, all the funicular joints distinctly longer than broad. Thorax rather robust, through the wing-insertions as broad as the head; epinotum sloping, evenly rounded, without distinct base and declivity. Petiolar scale higher than in the worker, its summit sharp, narrowed, compressed and produced upward. Gaster elongate-elliptical. Legs slender, feebly compressed.

Shining; finely punctate; mandibles very finely shagreened and coarsely and sparsely punctate.

Erect hairs delicate, abundant, rather short, uniform over the head, thorax and abdomen, shorter and somewhat less numerous on the scapes and legs. Pubescence dilute, longest and most distinct on the gaster.

Black; mandibles red, with black teeth; clypeus, cheeks, gula, antennal foveæ, antennæ, tarsi, tips of tibiæ, knees, posterior borders of gastric segments and their ventral sclerites, except at the base, brownish yellow. Wings yellowish, with brown veins and dark brown pterostigma.

Described from numerous specimens which were nesting in flourishing and very aggressive colonies in large clusters of *Tillandsias* growing on mangroves along the Cuyuni River, near the Tropical Laboratory of the New York Zoological Society at Kartabo, British Guiana.

Subsp. TOCOCÆ Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 693 ♀ ♂.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in the leaf-sacs of *Tococa guianensis* Aubl.

Bolivia: Tumupapas (W. M. Mann), in leaf-sacs of *Tococa* sp.

Var. ELATIOR Forel

Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 239 ♀ ♂.

Brazil: Bom Lugar, Rio Purus (J. Huber), in the petiolar sacs of a *Tococa*.

Peru: Iquitos and Putumayo (J. C. Bradley), in the leaf-sacs of a *Tococa*.

AZTECA TRIGONA Emery

Emery, Mem. Accad. Sc. Ist. Bologna (5) **3**, 1893 p. 346 ♀ fig. 78, 80;
A. festai Emery, Bull. Mus. Zool. Anat. Torino 1896 p. 4 ♀.

A colony of this ant was found near Las Cascades, C. Z. inhabiting all the cauline swellings of a young *Cordia alliodora* and attending numerous coccids on the stems and petioles, about which fragile carton shelters had been built. The species usually makes large pendent carton nests on the branches of various trees.

AZTECA TONDUZI Forel

Forel, Biol. Centr. Amer. Hymen, **3**, 1899-1900 p. 114 ♀ ♂.

Costa Rica: (Tonduz) in the pseudobulbs of an orchid.

Subsp. MATHILDÆ Forel var SPURIA Forel

Forel, Ann. Soc. Ent. Belg. **50**, 1906 p. 238 ♀.

British Guiana: Kartabo and Kalakoon (Wheeler), inhabiting the internodes of *Cecropia*.

Subsp. MEDIOPS Forel

A. festai subsp. *mediops* Forel, Rev. Suisse Zool. **12**, 1904 p. 43 ♀; *A. trigona* subsp. *mediops* Bailey, Bot. Gazette **74**, 1922 p. 379.

British Guiana: Kartabo (Wheeler), in *Cecropia angulata* Bailey.

AZTECA ULEI Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 694 ♀.

Brazil: Jurua Miry, Jurua, Amazonas (E. Ule), in "ant gardens" among Gesneriaceæ.

Var. CORDIÆ Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 695 ♀.

Brazil: Marary, Jurua, Amazonas (E. Ule), in the cauline swellings of *Cordia nodosa* L.

British Guiana: Kartabo and Kalacoon (Wheeler), in cauline swellings of *Cordia nodosa* L.

Bolivia: Rio Negro and Tumupasas (W. M. Mann), in cauline swellings of *Cordia hispidissima*.

The series from British Guiana comprise many workers and three dealated females. The latter (undescribed) measure 7-8 mm. and have the head if anything slightly broader than long, with straight, sub-parallel sides behind the eyes and also with parallel sides in front of the eyes, though the head is here much narrower; the posterior border being angularly excised. Eyes rather large, convex, broadly elliptical, about one-half their length from the anterior corners of the cheeks. Lateral borders of clypeus swollen. Antennæ stout, scapes not reaching to the posterior border of the head; first and second funicular joints longer than broad, remaining joints, except the last, as broad as long. Thorax robust, somewhat more than twice as long as broad, epinotum rounded and sloping, more abruptly at its posterior end. Petiolar scale convex in front, concave behind, with rather sharp superior border, the ventral portion protuberant and somewhat compressed. Gaster elliptical, about twice as long as broad, the first segment narrowed anteriorly. Legs long and stout.

Shining; mandibles sparsely and sharply punctate; punctuation of the body very fine, dense and superficial.

Pilosity yellow, short, rather abundant; oblique hairs numerous on tibiæ, few on scapes. Pubescence distinct only on the clypeus.

Black; mandibles deep red, with black teeth; corners of clypeus, funiculi, bases of scapes, tarsi, except at the base, brownish yellow. Wing-insertions pale sordid yellow.

Var. GAGATINA Wheeler and Mann var. nov.

Worker minor. Length 2.3 mm.

Differing from the other forms of *ulei* in sculpture and color, being smooth and shining and jet-black, with the apical borders of the mandibles, except the teeth, the lateral corners of the clypeus, the insertions of the antennæ, the antennal foveæ and terminal tarsal joints reddish yellow. The petiolar scale is low, with rather blunt summit, the mesonotum moderately convex.

Three specimens taken by Dr. W. M. Mann at Rurrenabaque, Bolivia, in a cauline swelling of *Cordia hispidissima*.

Var. GIBBIFERA Forel

Forel, Verh. zool. bot. Ges. Wien 58, 1908 p. 393 ♀.

Originally described from São Paulo, Brazil (H. v. Ihering). A number of workers, taken by Dr. W. M. Mann at Tumupapas, Bolivia in the cauline swellings of *Cordia hispidissima*, are referable to this variety.

Subsp. NIGRICORNIS Forel

Forel, Zool. Jahrb. Abt. Syst. 20, 1904 p. 695 ♀.

Brazil: Cachoeira, Jurua, Amazonas (E. Ule), in cauline swellings of *Cordia nodosa* L.

Subsp. ROSSI Forel

Forel, Deutsch. Ent. Zeitschr. 1909 p. 251 ♀.

Mexico: (Ross), in ant-gardens among the roots of orchids.

AZTECA VELOX Forel

Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 108, ♀ ♀ ♂, pl. 4, fig. 13.

Panama: Ancon, C. Z. (Wheeler), attending coccids on young *Cordia gerascanthus* trees; Las Cascades (Wheeler), living in twigs and branches of *Triplaris cumingiana*.

Guatemala: Patulul (Wheeler), nesting in trunk of *Acacia hindsii* Benth.

Costa Rica: Alajuela (Wheeler), nesting in Tillandsias.

Subsp. NIGRIVENTRIS Forel

Forel, Ann. Soc. Ent. Belg. 50, 1906 p. 241.

Brazil: Bom Lugar, Rio Purus (A. Goeldi and J. Huber), in cauline swellings of *Cordia nodosa* L.

Costa Rica: Esparta (P. Biolley), occurring regularly in the pseudobulbs of an orchid, *Epidendrum bicornutum* Hook.

Panama: Balboa (Wheeler), nesting in the fistulose stems of *Clerodendron siphonanthus* (introduced from the Orient).

AZTECA VIRENS Forel

Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 115 *nota* ♀.

Brazil: Parà, Amazonas (A. Goeldi), living in the green stems of a plant (undetermined).

AZTECA XANTHOCHROA (Roger)

(Plate 57, fig. a)

Liometopum (?) *xanthochroum* Roger, Berlin. Ent. Zeitschr. 7, 1863 p. 167 ♀;
Azteca xanthochroa Emery, Boll. Mus. Zool. Anat. Torino 1896 p. 2, ♀ ♀,
 fig. 1 a-d; Forel, Biol. Centr. Amer. Hymen. 3, 1899-1900 p. 115 fig. 14,
 14a; Ross, Naturw. Wochenschr. 8, 1909 p. 829.

Mexico: (type locality); Teapa in Tabasco (H. H. Smith); Zacuapam (H. Ross), in *Cecropia mexicana* Hemsl.; Palomares (A. Petrunkevitch).

Honduras: Puerto Castilla (J. Bequaert), in *Cecropia* sp.

Guatemala: Las Mercedes 3000 ft. (Champion); Quirigua, Escuintla and Puerto Barrios (Wheeler), in internodes of young *Cecropias*; Trece Aguas, Alta Vera Paz (Schwarz and Barber).

Costa Rica: (A. Alfaro), in *Cecropias*.

The singular yellow female of this species was described by Roger from Mexico in 1863. Mayr in 1866 described what he took to be the conspecific worker, but which Emery claims was *A. instabilis* Sm. var. *mexicana* Emery. I find that I have also misidentified the worker of *xanthochroa*, for my record from Coroni, Trinidad (Amer. Mus. Novitates 45, 1922 p. 15), as I find from reëxamination of the specimens, refers instead to *A. delpini* subsp. *trinidadensis* Forel. In 1896 Emery assigned workers taken by Alfaro in Costa Rica to *xanthochroa*, but a study of materials which I collected in Guatemala in 1911 shows that although his specific allocation is correct, these workers must belong to a distinct variety. I have frequently taken deëlated queens of the typical *xanthochroa* accompanied by young colonies of workers in the internodes of *Cecropias* at Quirigua and Puerto Barrios, Guatemala, and these workers are different in several significant particulars from those described by Emery. Though the largest individuals approach the pale flavotestaceous color which he describes, the mediæ and minimæ are all much darker, being dark brown. The color shows a gradual gradation from the smallest to the largest individuals. The head of the major is broader, as broad as long, distinctly narrower in front than behind and the scapes reach to the posterior corners of the head and in the minor workers further beyond these corners than in Emery's figure. Since there is no doubt that the Guatemalan workers belong to the true *xanthochroa* I believe one must regard Emery's workers as representing a distinct variety. I suggest that it be called var. *costariënsis* var. nov.

Concerning this variety Emery states on the authority of Alfaro that it differs from the other *Cecropia* Aztecas. It habitually runs with its gaster elevated. The opening of the nest is a longitudinal fissure in the *Cecropia* internode. It constructs carton septa across the internodal cavities. It is less aggressive than the other species and does not bite the hands of the collector.

I find in my collection specimens of three additional subspecies of *xanthochroa*, which are here described.

Subsp. AUSTRALIS subsp. nov.

Female. Length 10.5—11 mm; wings 10 mm.

Slightly larger than the female of the typical *xanthochroa* and differing in the following characters: Head much more narrowed behind, where it is scarcely broader than in front, the mandibles more flattened and more deflected at the tips, more shining and more coarsely punctate and without striæ. Surface of body and especially the head, more shining. Color yellow, resembling that of the typical *xanthochroa* but the brown spot on the head is more extensive, there are usually three longitudinal brown streaks on the mesonotum, the bases of the gastric segments are brown, the antennal scapes blackish, the funiculi yellow, the mandibles deep red, with blackish borders, the anterior border of the clypeus infuscated, the legs reddish, the tibiæ darker, the fore tibiæ and all the basi tarsi blackish; the wings as in the type, brownish yellow, with resin yellow veins and dark brown pterostigma. I can detect no significant differences in pilosity and pubescence.

Described from three specimens from Songo, Bolivia (type locality), and two from Callanga, Peru, purchased from Staudinger and Bang-Haas.

Subsp. ISTHMICA subsp. nov.

Worker. Length 2—4 mm.

Very similar to the minor worker of the typical form of the species and of the same dark brown color, but the mandibles and anterior portion of the head are more yellowish. The antennæ are distinctly shorter, the scapes surpassing the posterior corners of the head to a noticeably less extent and the funicular joints broader in proportion to their length. The terminal funicular joints are blackish, the petiolar scale a little higher and more compressed anteroposteriorly, the pubescence, especially on the head, more abundant and more conspicuous.

Female. (deälated) Length 6.5 mm.

Much smaller than the female of the typical *xanthochroa* which measures 9-10 mm. Head decidedly shorter, with more convex eyes and shorter antennæ; sides behind the eyes are more convex, the excavation of the posterior border less pronounced. Joints 3-10 of the funiculi distinctly broader than long, the first and second joints distinctly shorter in proportion to their length than in the typical *xanthochroa*. The petiolar scale is lower and less pointed above. Body smoother and somewhat more shining; mandibles, however, subopaque

and more sharply striate. The color is the same but the body is a shade paler, there is no brown spot on the head and the scapes and first funicular joint are yellow. The hairs on the body are blackish.

Described from a dozen workers and a single female, constituting an incipient colony which I found nesting in an internode of a young *Cecropia* near the source of the Rio Agua Salud, C. Z., Panama. To the same subspecies belong a number of workers which I took in young *Cecropias* at Culebra and Monte Sirio, C. Z., during November 1913, and a number of females taken on Barro Colorado Island, C. Z., in 1924.

AZTECA XANTHOCHROA subsp. SALTII subsp. nov.

Worker minor. Length 2.5 mm.

Smoother and more shining than the typical *xanthochroa* or the subsp. *isthmica*, the thorax, gaster, antennæ and legs being deep castaneous brown, the head black, the mandibles and corners of the clypeus red. Pilosity shorter and less abundant, especially on the head. Antennal scapes extending a distance more than their greatest diameter beyond the posterior corners of the head.

Female: (deälated). Length 10-12 mm.

Head even longer and narrower than in the subsp. *australis*, especially behind, with nearly straight sides, more flattened eyes and deeper occipital excision. Surface much smoother and more shining than in either the typical form or the other subspecies. Thorax and gaster somewhat darker, more reddish yellow; head, antennæ and femora red, the clypeus and mandibles deep red; tibiæ and metatarsi blackish red. Hairs on the body and tibiæ blackish, longer and more abundant than in *australis*, appressed pubescence on the gaster long and yellowish.

Described from five workers and nine females taken by Dr. George Salt from the internodes of young *Cecropias* at Vista Nieve, Sierra Madre de Santa Marta, Colombia, at an altitude of 5,000 feet.

Genus TAPINOMA Foerster
TAPINOMA ATRICEPS Emery

Forel, Ann. Soc. Ent. Belg. 49, 1905 p. 171 ♂.

Brazil: São Paulo (A. Lutz), nesting in bamboos.

Var. BREVISCAPA Forel

Forel, Verh. Zool. bot. Ges. Wien 1908 p. 384 ♂ ♀.

Brazil: São Paulo (A. Lutz), nesting in bamboos.

TAPINOMA CANALIS sp. nov.

Worker. Length 1—1.2 mm.

Head subrectangular, longer than broad, as broad in front as behind, with very feebly convex sides, straight posterior border and rounded posterior corners. Eyes small, flat, placed about one and one-half times their length from the anterior corners of the head. Mandibles narrow, with oblique apical borders, armed with six teeth, the first, second and fourth from the tip larger, the others minute. Frontal carinae short, parallel, further apart than their distance from the sides of the head. Clypeus convex and rounded in the middle, the anterior border straight and entire. Frontal area and groove absent. Antennal scapes not reaching the posterior corners of the head by a distance equal to their greatest diameter; first funicular joint nearly one and one-half times as long as broad, second joint short and small, broader than long, remaining joints, except the last, all broader than long, but increasing in length and breadth distally. Thorax small, feebly but distinctly impressed at the mesoëpinotal suture, the dorsal outline otherwise slightly convex, the base of the epinotum straight, forming a distinct obtuse angle with the declivity, which is decidedly longer than the base. Seen from above the pronotum is nearly twice as broad as long, the mesonotum subcircular and about as long as broad, the epinotum slightly longer than broad and laterally compressed. Petiole very small and flat, longer than broad, its scale represented by a low, semicircular swelling at its anterior end, completely concealed under the large first gastric segment. Gaster elliptical. Legs of the usual shape.

Shining, very finely and indistinctly punctate.

Hairs and pubescence pale yellow, the former very sparse, distinct only on the clypeus, mandibles and gastric segments; the pubescence rather dense, somewhat obscuring the shining surface, rather long and oblique on the scapes.

Pale brownish yellow; legs scarcely paler; mandibular teeth reddish.

Female. (deälated) Length 1.8 mm.

Head much as in the worker but the posterior corners less rounded and the posterior border very feebly excised, the eyes larger and more convex, only about half their length from the anterior corners. Scapes shorter, extending about two-thirds the distance between the eyes and posterior corners of the head. Thorax slightly narrower than the head, elongate subelliptical, depressed above; pronotum very short and transverse; mesonotum as long as broad, rounded in front; epinotum short, with very short base, passing gradually into the long sloping

declivity. Petiole as in the worker. Gaster elongate, more than three times as long as broad, parallel-sided, first segment covering the petiole and provided with a very distinct impression for its accommodation.

Subopaque and more densely punctate than the worker. Pilosity and pubescence very similar; head and thorax pale yellowish brown; gaster dark brown; mandibles, antennæ, legs and broad margins of the gastric segments brownish yellow.

Male. Length about 8 mm.

Head subrectangular, longer than broad, somewhat narrowed behind. Eyes flattened about half as long as the sides of the head. Mandibles well-developed, overlapping with their very minutely denticulate apical borders. Clypeus with straight, transverse anterior border. Antennal scapes reaching to the posterior corners of the head; basal funicular joints slightly, terminal joints considerably longer than broad. Thorax as broad as the head. Petiole very much as in the worker. Gaster short.

Sculpture and pilosity like those of the worker. Color pale sordid yellow. Wings grayish pubescent, rather opaque, with pale brownish veins.

Described from the members of a single colony inhabiting a cauline swelling of *Cordia alliodora* on Barro Colorado Island in Gatun Lake, C. Z., Panama.

This very small species seems to be quite distinct from any of the indigenous neotropical species of *Tapinoma*, such as *atriceps* Emery, *heyeri* Forel, *littorale* Wheeler and *ramulorum* Emery.

TAPINOMA MELANOCEPHALUM Fabr.

Colonies of this widely distributed tropicopolitan ant were found nesting in the twigs of *Triplaris americana* at Balboa, C. Z. Panama.

TAPINOMA RAMULORUM Emery

subsp. INRECTUM Forel

Guatemala: Escuintla (Wheeler), nesting in twigs of *Triplaris auriculata*.

TAPINOMA LITTORALE Wheeler

Wheeler, Bull. Amer. Mus. Nat. Hist. 21, 1905 p. 110 ♀ ♂.

Florida: Card's Point (Wheeler), nesting in Tillandsias.

Bahamas: Andros and New Providence Islands (Wheeler), nesting in Tillandsias.

Var. CUBAËNSIS Wheeler

Wheeler, Bull. Mus. Comp. Zool. **54**, 1913 p. 498 ♀ ♀ ♂.

Cuba: (Wheeler), nesting in *Tillandsias*.

Subfamily FORMICINAE

Genus PRENOLEPIS Mayr

PRENOLEPIS (PARATRECHINA) LONGICORNIS Latr.

Panama: Ancon, C. Z. (J. Zetek), attending aphids on foliage of *Cordia alliodora*.

PRENOLEPIS (NYLANDERIA) FULVA Mayr

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 705.

Brazil: San Joãquim, Rio Negro, Amazonas (E. Ule), nesting in *Tillandsia paraënsis* Metz.

Genus BRACHYMYRMEX Mayr

BRACHYMYRMEX HEERI Forel

Wheeler, Zoologica **3**, 1921 p. 166 ♀ ♀.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl., in cauline swellings of *Cordia nodosa*.

Var. BASALIS Wheeler

Wheeler, Zoologica **3**, 1921 p. 116 ♀.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

Var. OBSCURIOR Forel

Panama: Ancon, C. Z. (Wheeler), attending coccids on the foliage of seedling *Cordia alliodora*; Balboa C. Z., (Wheeler), nesting in the twigs and running on the trunks of *Triplaris americana*; Tumba Muerte Road, near Las Sabanas (Wheeler), nesting in the thorns of *Acacia penonomensis* Saff.

BRACHYMYRMEX PICTUS Mayr

subsp. BALBOÆ subsp. nov.

Worker. Length 1—1.2 mm.

Smaller than the typical form from Brazil, the scapes shorter, ex-

tending scarcely one-fourth their length beyond the posterior corners of the head. Honey yellow, the antennæ and legs whitish yellow; the gaster black on the dorsal side, with a broad pale yellow area embracing the posteromedian portion of the first segment and the median portion of each of the succeeding segments.

Female. Length 2.5 mm.

Smaller than the typical form of the species. Ocellar triangle black; a brown spot on the posterior portion of the mesonotum and the gaster dark brown above, with the posterior borders of the segments yellow. Wings long, colorless, with colorless veins and pterostigma.

Male. Length 0.8—1 mm.

Scapes reaching to the posterior ocelli. Head with the eyes slightly broader than long, rounded rectangular. Eyes elongate, about-two-thirds as long as the sides of the head, not very convex. Mandibles very slender. Whitish yellow; top of head and gaster, except its last segments, brown; antennal funiculi slightly infuscated at the tip. Wings as in the female.

Several colonies of this pretty form were found nesting in hollow twigs of *Triplaris americana* at Balboa, C. Z. Panama.

Genus MYRMELACHISTA Roger

MYRMELACHISTA AMBIGUA Forel

subsp. RAMULORUM Wheeler

Wheeler, Bull. Amer. Mus. Nat. Hist. **24**, 1908 p. 155 ♀ ♀ ♂.

Porto Rico and Culebra Island (Wheeler), in twigs of seagrape (*Coccoloba uvifera*) and "torchuelo" (*Bucida buceros*)

Costa Rica: Alajuela (Wheeler), in Tillandsias.

MYRMELACHISTA (DECAMERA) BAMBUSARUM Forel

Forel, Ann. Soc. Ent. Belg. **47**, 1903 p. 263 ♀.

Brazil: Corcorado, Prov. Rio Janeiro (Goeldi), in small branches of bamboo.

MYRMELACHISTA (DECAMERA) NIGELLA Roger

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 704 ♀.

Brazil: Bocca do Tejo, Jurua, Amazonas (E. Ule), in the twig swellings of *Duroia hirsuta* H. Sch.

MYRMELACHISTA (DECAMERA) NODIGERA Mayr

var. FLAVICORNIS Emery

Emery, Zool. Jahrb. Abt. Syst. **9**, 1896 p. 638 ♀ fig. F.Paraguay: San Salvador (J. Bohls), in woody thorns of *Acacia* (probably *cavenia*).

MYRMELACHISTA (DECAMERA) PADEREWSKII Forel

Forel, Verh. zool. bot. Ges. Wien 1908 p. 397 ♀ ♀ ♂.

Brazil: São Paulo (E. Lutz), nesting in bamboo.

MYRMELACHISTA (DECAMERA) SCHUMANNI Emery

Emery, Ann. Soc. Ent. France 1890 p. 70 nota ♀; Schumann, Sammlun. gemeinverst. wiss. Vortr. **83**, 1889 p. 17; Emery, Biol. Centralbl. **11**, 1891, p. 165; Anal. Mus. Nac. Costa Rica 1892 p. 65.Colombia: (K. Schumann), in cavities of *Duroia hirsuta*.

Var. CORDINCOLA Wheeler & Mann, var. nov.

Worker. Length 1.5—3 mm.

Averaging somewhat larger than the typical form of the species and differing in color. Brownish yellow, with the head and gaster brown, the latter usually darker than the former, the bases of the gastric segments paler than their posterior borders which are edged with black. Mandibles and anterior portion of head paler and more yellowish than the posterior portion. Antennæ and legs yellow, the clubs of the former and in some specimens also the middle portions of the femora feebly infuscated. Summit of petiolar node entire, its anterior surface distinctly convex, its posterior surface nearly flat.

Described from numerous specimens taken by Dr. W. M. Mann at Osunto, Bolivia, in the cauline swellings of *Cordia hispidissima*.

MYRMELACHISTA (DECAMERA) ULEI Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 704 ♀.

Peru: Cerro de Escaler, 1200 m. (E. Ule), in the swellings of the flower stems of a melastomaceous plant.

MYRMELACHISTA (DECAMERA) ZELEDONI Emery

Costa Rica: San Jose and Cartago (Wheeler), in Tillandsias.

This species was also found nesting in dead twigs in the same localities. Recorded also from Colombia (Forel).

Genus CAMPONOTUS Mayr

CAMPONOTUS (TANAEMYRMEX) BONARIENSIS Mayr

Paraguay: San Bernardino (K. Fiebrig *in litteris*), occasionally nesting in *Cecropia peltata* L.

CAMPONOTUS (MYRMOTURBA) MELANOTICUS Emery var.

HAGMANNI Forel

British Guiana: Kartabo (Wheeler), nesting in large branches of *Cecropia sciadophylla* var. *decurrans* Sn.

Costa Rica: Alajuuela (Wheeler), nesting in Tillandsias.

CAMPONOTUS (MYRMOTURBA) ULEI Forel

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 706 ♂ ♀ Ule, Ameisenpflanzen des Amazonas gebietes 1907.

Peru: Cerro de Escaler, 1300 m. (E. Ule), in the hollow internodes of *Cecropia montana* Warb. (Ule No. 6845)

CAMPONOTUS (MYRMOTHRIX) ABDOMINALIS Fabr.

var. COSTARICENSIS Forel

Costa Rica: Alajuuela (Wheeler), nesting in Tillandsias.

Var. between subsp. ESURIENS and var.

MEDIOPALLIDUS Forel

Wheeler, Amer. Natur. **35**, 1901 p. 526; Ann. Soc. Ent. Belg. **45**, 1901 p. 205.

Mexico: Cuernavaca (Wheeler), nesting in *Tillandsia benthamiana*.

CAMPONOTUS (MYRMOTHRIX) FEMORATUS Fabr.

Forel, Zool. Jahrb. Abt. Syst. **20**, 1904 p. 705 ♂ ♀; Wheeler, Ecology **2**, 1921 p. 96.

Brazil: Manaus (E. Ule), in ant gardens among *Streptocalyx* and *Codonanthe*.

British Guiana: Kartabo (Wheeler), in ant gardens in parabiosis with *Crematogaster limata* Sm. subsp. *parabiotica* Forel.

CAMPONOTUS (MYRMOTHRIX) LUTZI Forel

Forel, Ann. Soc. Ent. Belg. 49, 1905 p. 169 ♀ ♀.

Brazil: São Paulo (E. Lutz), nesting in bamboos.

CAMPONOTUS (MYRMOTHRIX) RUFIPES Fabr.

Subsp. RINGGERI Emery

Paraguay: San Bernardino (K. Fiebrig *in litteris*), occasionally nesting in *Cecropia peltata* L.

CAMPONOTUS (NEOMYRMANIBLYS) NOVOGRENADENSIS Mayr

A small colony of this ant was taken at Quebrada de Oro, C. Z., Panama, from the cavity of a cauline swelling of *Cordia alliodora*. The cavity also contained the remains of small narrow puparia of a species of *Microdon*. This ant nests much more frequently under bark and especially in the abandoned galleries of termitaria. In the latter situations, I have frequently found considerable numbers of the same *Microdon*.

CAMPONOTUS (MYRMOBRACHYS) BRETTESEI Forel

Taken on several occasions in the cauline swellings of *Cordia alliodora* near the laboratory at Ancon, C. Z. It is more frequently found nesting in dead twigs, as Forel observed in Colombia.

CAMPONOTUS (MYRMOBRACHYS) BREVIS Forel

Not uncommon at Frijoles, Red Tank and Las Cascades, C. Z., in the cauline swellings of *Cordia alliodora*. Along the Tumba Muerte Road, near Las Sabanas, Panama, I found several colonies nesting in the thorns of *Acacia penonomensis* Saff. In 1911 I found this ant nesting in *Tillandsias* on Otoque and Tabogilla Island, Panama. The female measures 6—6.5 mm. and resembles the worker major. The wings are grayish hyaline, with resin-yellow veins and pterostigma. The male measures 3.5—3.8 mm. and is black, moderately shining,

with yellow tips to the mandibles. The wings are yellowish hyaline, with the veins and pterostigma paler than in the female. As Forel observed in Colombia, *C. brevis* is most abundant in dead twigs.

CAMPONOTUS (MYRMOBRACHYS) CANESCENS Mayr

A few small colonies were found nesting in cauline swellings of *Cordia alliodora* on Barro Colorado Island, in Gatun Lake, C. Z.

CAMPONOTUS (MYRMOBRACHYS) LINDIGI Mayr

Running up and down the trunks and attending coccids on the foliage of *Triplaris americana* at Balboa, C. Z. and of *Cordia alliodora* near the laboratory at Ancon, and nesting in the ground about the roots of these same trees.

CAMPONOTUS (MYRMOBRACHYS) PLANATUS Roger

Wheeler, Bull. Amer. Mus. Nat. Hist. **21**, 1905 p. 110 Cuba: (Wright), in Tillandsias.

Var. CONTINENTIS Forel

C. planatus Emery, Bull. Soc. Ent. Ital. **22**, 1890 p. 56; Wheeler, Trans. 2nd Intern. Ent. Congr. 1912.

Florida: Card's Point (Wheeler), nesting in Tillandsias.

Costa Rica: Alajuela, etc. (A. Alfaro), living in *Acacia* thorns abandoned by *Pseudomyrma*; Alajuela (Wheeler) nesting in Tillandsias.

Guatemala: Zacapa, Quirigua, Escuintla and Patulul (Wheeler), living in thorns of *Acacia hindsii*.

Both this variety and the typical *planatus* also live under bark and in dead twigs.

CAMPONOTUS (MYRMOBRACHYS) PITTIERI Forel

var. POENALIS Wheeler

Wheeler, Zoologica **3**, 1891 p. 167 ♂ ♀.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

CAMPONOTUS (MYRMOBRACHYS) VEZENYI Forel

Paraguay: San Bernardino (K. Fiebrig *in litteris*), occasionally in *Cecropia peltata* L.

CAMPONOTUS (MYRMOBRACHYS) ZOC Forel

Wheeler, Zoologica **3**, 1921 p. 167 ♀.

British Guiana: Kartabo (Wheeler), in hollow petioles of *Tachigalia paniculata* Aubl.

CAMPONOTUS (MYRMOCLADOECUS) BIDENS Mayr

A single colony found nesting in a cauline swelling of *Cordia alliodora* on the Chivachiva Trail, near Red Tank, C. Z., Panama. This species and its allies more frequently nest in dead twigs.

CAMPONOTUS (MYRMOCLADOECUS) MUCRONATUS Emery

Costa Rica: Alajuela (Wheeler), nesting in Tillandsias.

Guatemala: Escuintla (Wheeler), nesting in thorns of *Acacia bursaria* Schenck. More common in dead twigs.

CAMPONOTUS (MYRMOCLADOECUS) STRIATUS F. Smith

Subsp. ALFAROI Emery

Several colonies of this subspecies were found nesting in thorns of *Acacia penonomensis* Saff. along the Tumba Muerte Road, near Las Sabanas, Panama. The female measures 4—4.5 mm. and has the thorax subopaque and coarsely shagreened above and finely striated on the sides. The petiolar scale is broader and thicker than in the worker. The wings are yellowish hyaline, with yellow veins and dark brown pterostigma. The male is very small, measuring only 3—3.2 mm., and is shining black throughout, except the wings, which are whitish, with paler veins and pterostigma than in the female.

This ant more frequently nests in dead twigs, as Forel observed in Colombia.

CAMPONOTUS (MYRMOCLADOECUS) RECTANGULARIS Emery

Emery, Biol. Centralbl. **11**, 1891 p. 168; Anal. Mus. Nac. Costa Rica 1892 p. 67.

Costa Rica: (A. Alfaro), in thorns of *Acacia*.

Var. RUBRONIGER Forel

Wheeler, Amer. Natural. **35**, 1901 p. 526; Ann. Soc. Ent. Belg. **45**, 1901 p. 205.

Mexico: Cuernavaca (Wheeler), nesting in *Tillandsia benthamiana*.

CAMPONOTUS (PARACOLOBOPSIS) PATIMÆ sp. nov.

(Plate 57, fig. b)

Worker major. Length 6—7 mm.

Allied to *salvini* Forel and especially *rudigenis* Emery. Head large, thick, convex, subrectangular, about one and one-fifth times as long as broad, about as broad in front as behind, with nearly straight sides and broadly but not deeply excised posterior border, the cheeks swollen and projecting anteriorly beyond the clypeus, which is longer than broad, nearly parallel-sided, convex in the middle and narrowly carinate anteriorly. Frontal area very small, triangular. Frontal groove distinct; frontal carinæ approximated in front, sigmoidal in the middle, straight and parallel behind. Eyes rather small and flat, at the posterior third of the head. Mandibles short, very convex, with four large subequal teeth. Antennal scapes slender, compressed, curved, somewhat thickened at the tips which reach to the posterior corners of the head. Thorax broad through the pronotum, which is flattened above, submarginate on the sides, rapidly narrowing posteriorly to the epinotum, which is compressed and tectiform. The promesonotal suture is very distinct, the mesoëpinotal suture nearly obsolete. In profile the whole thoracic dorsum is very evenly arcuate, though the outline of the epinotum really consists of three subequal, nearly straight lines. Petiolar scale rather thick, oval, broader above than below, beveled at the summit, with flat posterior surface, the superior border rounded, entire and rather blunt. Gaster elongate, elliptical. Legs long and stout, the fore femora somewhat enlarged.

Opaque; only the mandibles and posterior corners of the head somewhat shining. Mandibles finely punctate; head more coarsely and densely, the cheeks and clypeus also covered with shallow piligerous foveolæ. Remainder of body and the appendages very finely and densely punctulate.

Hairs golden yellow, erect or suberect, moderately long and abundant on the front, thorax, petiolar border and gaster, on the cheeks and clypeus short, stout and blunt. Legs with short, appressed hairs, scapes finely pubescent. Thorax and gaster with long, hair-like pubescence, not sufficiently dense to conceal the surface or forming a pelt as in *salvini*.

Ferruginous red; vertex of head, pleura, rarely the dorsal surface of the epinotum, dorsal border of petiole, more rarely the whole scale, and gaster, except the bases of the segments, black. The posterior borders of the gastric segments are dull yellowish. Fore coxæ and all

the femora more yellowish ferruginous, their bases, the knees, tibiæ, tarsi and antennæ dark brown.

Worker minor. Length about 4.5 mm.

Head longer than in *rudigenis*, distinctly longer than broad, with straight, subparallel sides and somewhat larger and more prominent eyes, but the latter connected with the posterior corners of the head by distinct ridges and the posterolateral surfaces of the head concave as in that species. Antennal scapes longer, very thin, extending about half their length beyond the posterior corners of the head. Thorax and petiole resembling those of the worker major, but the head is not more coarsely punctate than the remainder of the body. The whole surface is a little more shining or lustrous.

Pilosity much as in the worker major.

Black; mandibles yellow; sides of head below the eyes, a spot on each side of the pronotum, a streak on each side of the epinotum, the ventral portion of the petiole, and base of first gastric segment, ferruginous. Appendages colored as in the worker major.

Male. Length 4.5—5 mm.

Head with a trace of the worker's postocular carina on each side. Eyes very prominent. Mandibles rather well developed, with four small teeth. Antennæ long and slender. Thorax robust, much broader than the head, the mesonotum from above nearly as broad as long. Petiolar scale low, its upper border transverse, blunt, broadly excised in the middle.

Sculpture like that of the worker minor, subopaque, but the gaster more shining.

Pilosity and pubescence also similar but white and of more uneven length.

Black; mandibles, anterior border of clypeus and insertions of antennæ ferruginous yellow; thorax of the same color, but pleura and epinotum spotted with dark brown and mesonotum with three broad dark brown streaks, the median abbreviated behind. Genital appendages yellow. Legs yellowish brown, tibiæ, tarsi and a streak along the extensor surface of each femur dark brown. Wings yellowish hyaline, with resin-yellow veins and pterostigma.

Described from four major workers, one minor worker and four males taken by Mr. H. O. Lang on Merumé Creek, British Guiana, in the fistulose stems of *Patima formicaria* Johnston (Rubiaceæ).

This species is, perhaps, only a subspecies of Emery's *rudigenis* from Venezuela, Peru and Brazil, but his description is very brief and I have seen no specimens. The head of the worker minor compared

with his figure certainly shows important differences. The color is very different, but in this respect the following variety agrees more closely with Emery's species. Moreover, there is considerable color variation in both the series of specimens.

Var. DOLENTULUS var. nov.

Worker major. Length 5—5.5 mm.

Very similar to the preceding form but smaller and differing in the following particulars: The posterior corners of the head are more extensively shining, the color is black, with the mandibles, gula and borders of the head deep ferruginous red; the legs are brownish yellow, with the extensor surfaces of the femora and tibiae and the bases of the femora black, the tarsi dark brown. The posterior borders of the gastric segments are more sordid yellow. The hairs are white and the pubescence is conspicuously shorter on the thorax and gaster and the surface of these regions is somewhat less opaque and more lustrous.

Worker minor. Length 3.5—3.8 mm.

Also smaller than the worker minor of the typical form, of the same color as the worker major, but with the sides of the pronotum, a spot at the base of the first gastric segment and in some specimens a streak on each side of the epinotum reddish.

Described from six major and six minor workers taken by Mr. Lang in the same locality and in the same species of Rubiaceæ as the preceding form.

CAMPONOTUS (HYPERCOLOBOPSIS) CHRISTOPHERSENI Forel

Forel, Mém. Soc. Ent. Belg. 20, 1912 p. 85 ♂ ♀ ♀.

Panama. Mamei (Christophersen), nesting in the spines of *Xanthoxylon* (*panamensis* P. Wilson).

CAMPONOTUS (HYPERCOLOBOPSIS) PARADOXUS Mayr

subsp. JANITOR Forel

Forel, Verh. zool. bot. Ges. Wien 1908 p. 415 ♂ ♀ ♀ ♂.

Brazil: São Paulo (E. Lutz), nesting in bamboo.

EXPLANATIONS OF PLATES

PLATE 1

PLATE 1

Cordia alliodora R. and P. Reservoir at Ancon, C.Z., Panama. March 17, 1923. Photograph by J. Zetek.



PLATE 2

PLATE 2

a. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. March 17, 1923.
Photograph by J. Zetek.

b. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. March 17, 1923.
Photograph by J. Zetek.



a



b

PLATE 3

PLATE 3

a. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Smallest swellings in flower cluster. Photograph by J. Zetek.

b. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Swellings at base of flower panicles. Photograph by J. Zetek.



a



b

PLATE 4

PLATE 4

a. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Larger swellings.
Photograph by J. Zetek.

b. *Cordia alliodora* R. and P. Ancon, C.A., Panama. Old swellings.
Photograph by J. Zetek.



a



b

PLATE 5

PLATE 5

Cordia alliodora R. and P. Ancon, C.Z., Panama. Very old swellings.
Photograph by J. Zetek.



PLATE 6

PLATE 6

a. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Old swellings in longitudinal section. Photograph by J. Zetek.

b. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Section of very old swelling, showing large black coccids. Photograph by J. Zetek.



b



a

PLATE 7

PLATE 7

a. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Old swelling containing a carton nest of *Azteca longiceps*. Photograph by J. Zetek.

b. *Cordia alliodora* R. and P. Ancon, C.Z., Panama. Oldest stage, the reduced cavity completely filled with a carton nest of *Azteca longiceps*. Photograph by J. Zetek.



a



b

PLATE 8

PLATE 8

Cordia alliodora R. and P. Ancon, C.Z., Panama. Nest of *Azteca longiceps*
in an old swelling.




PLATE 9

PLATE 9

Larva and cocoons of *Conchylodes salamisalis* Druce, and leaves of *Cordia alliodora* R. and P. attacked by this pyralid. Ancon, C.Z., Panama. Photograph by J. Zetek.




PLATE 10

PLATE 10

a. *Triplaris surinamensis* Cham. and Schl. Cuyuni River, British Guiana. Section of stem at distal end of internode. Photomicrograph by I. W. Bailey.


b. *Triplaris surinamensis* Cham. and Schl. Cuyuni River, British Guiana. Section of stem showing occlusion of aperture by callus growth. Photomicrograph by I. W. Bailey.

c. *Triplaris surinamensis* Cham. and Schl. Cuyuni River, British Guiana. Section of stem about the middle of an internode. Photomicrograph by I. W. Bailey.

d. *Triplaris surinamensis* Cham. and Schl. Cuyuni River, British Guiana. Section of stem showing jacket of peripheral medullary tissue in internodal chamber. Photomicrograph by I. W. Bailey.

e. Pellet fed by the ants of *Triplaris surinamensis* to the larvae, showing pieces of coccids. Photomicrograph by I. W. Bailey.


f. Hyphae growing on the inner walls of the pouches of *Triplaris surinamensis*. Photomicrograph by I. W. Bailey.




c




f




d



e



a



p

PLATE 11

PLATE 11

Triplaris americana L. Ft. Clayton, C.Z., Panama. March 1, 1923. Photograph by J. Zetek.



PLATE 12


PLATE 12

a. *Triplaris americana* L. Ft. Clayton, C.Z., Panama. Inflorescence; leaves eaten by insects (*Atta*, etc.), although the stems are inhabited by *Azteca*. Photograph by J. Zetek.

b. *Triplaris americana* L. Ft. Clayton, C.Z., Panama. Panicles of fruit. Photograph by J. Zetek.



b




a

PLATE 13

PLATE 13

a. *Patima formicaria* Johnston. Merume Creek, British Guiana. Photograph by H. O. Lang.

b. *Patima formicaria* Johnston. Merume Creek, British Guiana. Longitudinal section of hollow stem. Photograph by H. O. Lang.



b



a

PLATE 14

PLATE 14

a. *Patina formicaria* Johnston. Merume Creek, British Guiana. Longitudinal sections of upper internodes, some of them slightly swollen. Photograph by H. O. Lang.




PLATE 15

PLATE 15

Acacia bucerophora Robinson. Photograph of type in Gray Herbarium by
W. E. Safford.



PLATE 16

PLATE 16

Acacia bursaria Schenk. Tucumán, Guatemala. Photograph by W. E. Safford.



PLATE 17

PLATE 17

Acacia bursaria Schenk. Rio de las Cañas, Guatemala. Photograph by
W. E. Safford.



PLATE 18

PLATE 18

Acacia cochliacantha Humb. and Bonpl. (= *campecheana* Schenck). Rosario Sinaloa, Mexico. Photograph by W. E. Safford.



PLATE 19

PLATE 19

Acacia Collinsii Safford (= *yucatanensis* Schenck). San Sebastian, Chiapas Mexico. Photograph by W. E. Safford.



PLATE 20

PLATE 20

Acacia Collinsii Safford (= *yucatanensis* Schenck). San Sebastian, Chiapas, Mexico. Photograph by W. E. Safford.



PLATE 21

PLATE 21

Acacia Collinsii Safford (= *yucatanensis* Schenck). Chiapas, Mexico. Photograph by W. E. Safford.



PLATE 22

PLATE 22

Acacia Collinsii Safford (= *yucatanensis* Schenck). Vicinity of Merida, Yucatan, Mexico. Photograph by W. E. Collins.



PLATE 23

PLATE 23

Acacia Cookii Safford. Alta Vera Paz, Guatemala. Photograph by W. E. Safford.



PLATE 24

PLATE 24

Acacia Cookii Safford. Alta Vera Paz, Guatemala. Photograph by W. E. Safford.




PLATE 25

PLATE 25

Acacia cornigera L. (= *Hernandezii* Safford). Rascon, San Luis Potosi, Mexico. Photograph by W. E. Safford.



PLATE 26

PLATE 26

Acacia cornigera L. (= *Hernandezii* Safford). San Martin, Chalchicua, Tamazunchale, Huasteca Region, San Luis Potosi, Mexico. Photograph by W. E. Safford.



PLATE 27

PLATE 27

Acacia cornigera L. (= *Hernandezii* Safford). Huasteca Region, San Luis Potosi, Mexico. Photograph by W. E. Safford.



PLATE 28

PLATE 28

Acacia cornigera L. Green house plant grown from seed, Washington, D. C.
Photograph by W. E. Safford.



PLATE 29

PLATE 29

a. *Acacia cornigera* L. Green house plant grown from seed, Washington, D. C. Photograph by W. E. Safford.

b. *Acacia cornigera* L. Photograph of type of *Acacia spadicigera* Schl. and Cham., in Halle Herbarium.



a



b

PLATE 30

PLATE 30

a. *Acacia costaricensis* Schenck (= *penonomensis* Safford). Alajuela
Costa Rica. Photograph by W. E. Safford.

b. *Acacia costaricensis* Schenck (= *penonomensis* Safford). Penonomé,
Panama. Photograph by W. E. Safford.



11



12

PLATE 31

.

PLATE 31

Acacia dolichocephala Safford. North of city of Vera Cruz, Mexico. Photograph by W. E. Safford.



PLATE 32

PLATE 32

Acacia donnelliana Safford. Type, San Pedro Sula, Dept. Sa. Barbara, Honduras. Photograph by W. E. Safford.



PLATE 33

PLATE 33

a. *Acacia donnelliana* Safford. Type, San Pedro Sula, Dept. Sa. Barbara, Honduras. Photograph by W. E. Safford.

b. *Acacia globulifera* Safford (= *chiapensis* Safford). Type, northern Yucatan, Mexico. Photograph by W. E. Safford.



a



b

PLATE 34

PLATE 34

Acacia Hindsii Bentham. Acapulco, Guerrero, Mexico. Photograph by
W. E. Safford.



PLATE 35

PLATE 35

Acacia Hindsii Bentham. Manzanillo, Colima, Mexico. Photograph by
W. E. Safford.




PLATE 36

PLATE 36

Acacia Hindsii Bentham (= *tepicana* Safford). Type of *tepicana*, from Tepic, Mexico. Photograph by W. E. Safford.




PLATE 37

PLATE 37

a. *Acacia melanoceras* Beurling (= *multiglandulosa* Schenck). Marajal near Colon, Panama. Photograph by D. Fairchild.

b. *Acacia melanoceras* Beurling (= *multiglandulosa* Schenck). Marajal near Colon, Panama. Photograph by J. Zetek.



PLATE 38

PLATE 38

a. *Acacia melanoceras* Beurling (= *multiglandulosa* Schenck). Marajal near Colon, Panama. Photograph by J. Zetek.

b. *Acacia melanoceras* Beurling (= *multiglandulosa* Schenck). Marajal near Colon, Panama. Photograph by D. Fairchild.



b



a

PLATE 39

PLATE 39

Acacia Nelsonii Safford. Acapulco, Guerrero, Mexico. Photograph by
W. E. Safford.



PLATE 40

PLATE 40

Acacia nicoyensis Schenk. Nicoya, Costa Rica. Photograph by W. E. Safford.



PLATE 41

PLATE 41

Acacia sphaerocephala Schlecht. and Cham. (= *veracruzensis* Schenck).
Photograph of type of *sphaerocephala* in Berlin Herbarium. Actopan, State
Vera Cruz, Mexico.



PLATE 42

PLATE 42

Acacia sphaerocephala Schlecht. and Cham. (= *veracruzensis* Schenck).
From a plant grown in Darmstadt from seeds collected in sand dunes south
of the city of Vera Cruz, Mexico. Photograph by W. E. Safford.



PLATE 43

PLATE 43

Acacia sphaerocephala Schlecht. and Cham. (= *veracruzensis* Schenk).
Dunes of Vera Cruz, Mexico. Photograph by A. Dampf.



PLATE 44

PLATE 44

Acacia Standleyi Safford. Mexico. Photograph of type by W. E. Safford.



PLATE 45

PLATE 45


Acacia turgida Safford (apparently not published). Herba Santa, Chiapas, Mexico. Photograph by W. E. Safford.




PLATE 46

PLATE 46

- a. Dipterous galls on leaflets of *Acacia* "*cornigera*". Esequintla, Patulul, Guatemala.
- b. Dipterous galls on twigs of *Acacia costaricensis*, near Panama City.



a



b

PLATE 47

PLATE 47

a. *Pseudomyrma alliodoræ* Wheeler. Head of female and worker; thorax and petiolar nodes of worker.

b. *Pseudomyrma satanica* Wheeler. Head of worker and female; thorax and petiolar nodes of worker.











PLATE 48

PLATE 48

- a. *Pseudomyrma sericea* Mayr. Thorax and petiolar nodes of worker.
- b. *Pseudomyrma sericea* var. *cordiae* Forel. Head, thorax and petiolar nodes of worker.



a



b






PLATE 49


PLATE 49

a. *Pseudomyrma sericea* var. *fortis* Forel. Thorax and petiolar nodes of worker.

b. *Pseudomyrma tenuis* Mayr. Head, thorax and petiolar nodes of worker.



11



12

PLATE 50

PLATE 50

Pseudomyrma triplaridis subsp. *baileyi* Wheeler. Head of worker and female; thorax and petiolar nodes of worker.










PLATE 51




PLATE 51

a. *Pseudomyrma triplarina* var. *loewensohni* Forel. Head of female and worker; thorax and petiolar nodes of worker.

b. *Leptothorax* (*Goniothorax*) *echinatinodis* subsp. *dalmasi* Forel. Head, thorax and petiolar nodes of worker.



a



b

PLATE 52

PLATE 52

Cryptocerus (Cyathocephalus) pallens Klug var. *porrasi* Wheeler. a, soldier;
b, head of same, dorsal view; c, worker.






PLATE 53

PLATE 53

Cryptocerus (Cyathocephalus) setulifer Emery. a, soldier; b, head of same, dorsal view; c, worker.




PLATE 54

PLATE 54

Cryptocerus (Cyathocephalus) varians F. Smith. Soldier and head of same; worker.










PLATE 55




PLATE 55

a. *Azteca longiceps* Emery. Head of female, workers and male; thorax and petiolar node of worker.

b. *Azteca pittieri* Emery. Head of female and worker; thorax and petiolar node of worker.



a



b



PLATE 56

PLATE 56

- a. *Azteca prorsa* Wheeler. Head, thorax and petiolar node of worker.
- b. *Azteca theresiae* var. *menceps* Forel. Head of female and workers; thorax and petiolar node of worker.









PLATE 57





PLATE 57

a. *Azteca xanthochroa* (Roger). Head of female and workers; thorax and petiolar node of worker.

b. *Camponotus (Paracolobopsis) patimae* Wheeler. Head, thorax and petiolar node of worker.



a



b

105

Bulletin of the Museum of Comparative Zoölogy
AT HARVARD COLLEGE
VOL. XC, No. 2

Museum of Comparative
Zoology
OCT 28 1942
HARV

THE LEPIDOPTERA OF
BARRO COLORADO ISLAND, PANAMA. No. 2

BY WILLIAM T. M. FORBES
Cornell University, Ithaca, New York

WITH EIGHT PLATES

CAMBRIDGE, MASS., U. S. A.
PRINTED FOR THE MUSEUM
OCTOBER, 1942

8

PUBLICATIONS
OF THE
MUSEUM OF COMPARATIVE ZOÖLOGY
AT HARVARD COLLEGE

The BULLETIN and MEMOIRS are devoted to the publication of investigations by the Staff of the Museum or of reports by specialists upon the Museum collections or explorations.

Of the BULLETIN, Vols. I to LXXXIX, Vol. XC, No. 1 and Vol. XCI, No. 1 and No. 2 have appeared and of the MEMOIRS, Vol. I to LVI.

These publications are issued in numbers at irregular intervals. Each number of the Bulletin and of the Memoirs is sold separately. A price list of the publications of the Museum will be sent upon application to the Director of the Museum of Comparative Zoölogy, Cambridge, Massachusetts.


Bulletin of the Museum of Comparative Zoölogy
AT HARVARD COLLEGE
VOL. XC, No. 2

THE LEPIDOPTERA OF
BARRO COLORADO ISLAND, PANAMA. No. 2

BY WILLIAM T. M. FORBES
Cornell University, Ithaca, New York

WITH EIGHT PLATES

CAMBRIDGE, MASS., U. S. A.
PRINTED FOR THE MUSEUM
OCTOBER, 1942



BY WILLIAM T. M. FORBES

TABLE OF CONTENTS

	PAGE
Introductory Remarks	268
Additional Records for Families in First Report	269
Remaining Families	286
Family Sematuridae	287
Sematura	289
Family Thyatiridae	290
Family Lasiocampidae	290
Tolyte	292
Artace	293
Euglyphis	294
Family Eupterotidae	303
Epia	305
Colla	306
Quentalia	306
Anticla	308
Tamphana	309
Zanola	309
Apatelodes	310
Olceclostera	312
Family Uraniidae	314
Urania	315
Family Epiplemidæ	315
Syngria	317
Syngriodes	318
Schidax	319
Psamathia	319
Nedusia	320
Erosia	320
Gathynia	322
Antiplecta	324
Thysanocraspeda	325

¹ Published with the aid of a special gift from Mr. George R. Agassiz.

	PAGE
Skaphion	326
Bicavernosa	326
Tricolpia	327
Epiplema	327
Family Mimallonidae	328
Mimallo	330
Cicinnus	330
Psychocampa	331
Trogoptera	332
Bedosia	332
Lacosoma	333
Zaphanta	333
Menevia	333
Alheita	334
Aleyda	335
Family Thyrididae	335
Dysodia	337
Ochrothyris	341
Herdonia	342
Hypolamprus	342
Rhodoneura	343
Zeuzerodes	348
Risama	349
Draconia	350
Family Castniidae	351
Castnia	351
Family Cossidae	352
Xyleutes	354
Cossula	357
Miacora	358
Givira	360
Langsdorfia	366
Inguromorpha	367
Family Psychidae	367
Family Zygaenidae	368
Harrisina	369

	PAGE
Family Eucleidae	370
Sibine	373
Euclea	375
Talima	379
Tanadema	380
Semyra	380
Euphobetron	383
Euprosterna	383
Narosopsis	384
Natada	385
Perola	387
Palaeophobetron	388
Epiperola	390
Dichromapteryx	391
Prolimacodes	391
Family Dalceridae	392
Acraga	393
Acragopsis	395
Family Megalopygidae	395
Microrape	397
Mesoscia	397
Norape	398
Trosia	399
Megalopyge	402
Podalia	403
Family Hepialidae	404
Dalaca	405

INTRODUCTORY REMARKS

This article continues a study of the so-called Bombycine Lepidoptera, published as Bulletin Vol. 85, No. 4 of this Museum. It was issued in August, 1939, and was illustrated with eight plates, containing a total of 66 figures, numbered consecutively. The first article reported on various families, related to the Noctuidae, namely, those which possess a thoracic tympanum. These families were the Euchromiidae, Nolidae, Arctiidae, Pericopidae, Agaristidae, Lymantriidae, Notodontidae, and Dioptidae.

This second article contains an account of fifteen families, mostly small in size, with varied and, on the whole, generalized structure. They include all the remaining so-called Bombycine Lepidoptera except the Saturnid complex, which Dr. Marston Bates had originally intended to study. Under the present circumstances further reports of this series will be indefinitely postponed.

The treatment in this report follows that of the first one exactly. The diagrams of seasonal records or flight periods are identical in form. Keys to genera and species have been included, including a few extralimital forms. Positive and probable synonyms have also been freely included. For convenience the figures continue the numbering in the first article; figs. 67-127 are arranged in plates 9-16.

Much the larger part of the material used is identical with that listed in detail in the first report, to which the reader is referred. Details of the manner of collection will consequently be given only in cases of special interest. The chief defect of this earlier material was the great gap in the summer months. This gap has been very nicely filled by a collection made by Mr. N. S. Scrimshaw from the middle of June to the first week of August, 1940. Our only serious gap is now September, though no really intensive collecting has been done in April and May. The most noticeable change in the picture, resulting from the Scrimshaw collection, is the indication that a good many species fly almost continuously, though there are quite a number with well marked flight periods. The Scrimshaw material is not all mounted as yet, but what is available seems to be a fair sample, covering the whole period, and I doubt if the complete lot will alter the picture materially, except possibly in the obscure and fragile Nolidae and Lithosiinae.

The first section of this report consequently gives the summer flight data for those families treated in the first article, based mainly on the Scrimshaw collection. Additional genera and species new to Barro Colorado Island, are inserted in the proper systematic sequence, and discussed and treated in the method followed in both papers.

Additional Records for Families in first Report
Family EUCHROMIIDAE

Diagram I

ADDITIONAL RECORDS OF BARRO COLORADO ISLAND
BOMBYCES

(Scrimshaw Records for June–August)

NAME	Month Week	June				July				Aug.				Total
		1	2	3	4	1	2	3	4	1	2	3	4	
P. aliena														0
H. stictosoma			○	○		●	○	●			●			10
G. salvini			●	●		●	●	●	●		●			33
G. colona			●	●			●	●	●					12
P. paucipuncta							○	●			●			5
M. pyrha				●			○	○			●			7
M. ethela						○			○		●			4
C. echemus				●			●	○	○					8
C. tabascensis														0
C. saron			○	●		●	○	○			○			8
C. metallescens				○		○	○				○			9
C. hercynacula														0
S. clusia				○				○			○			3
S. nox				●		○	●	○						8
S. phoenicosticta				●			○	●						6
S. anselma														0
S. afflicta				○				●			○			4
P. thoracica			●	●		○	●	●			●			45
D. correbioides														0
M. auripes				○		○	○				○			4
M. cyllarus														0
C. tiburtus				●		●	●	●			●			35
M. achrysa			○					○						2
H. restricta			○								●			3
E. obscurata														0
D. rubricineta														0
E. obscurum								○			○			2
C. affinis														0
C. e. germana				●							●			4
C. e. costinotata														0
C. e. elegans			●	○							○			5
C. e. striata											○			1

ADDITIONAL RECORDS OF BARRO COLORADO ISLAND
BOMBYCES — Continued

(Scrimshaw Records for June–August)

	NAME	Month Week	June				July				Aug.				Total
			1	2	3	4	1	2	3	4	1	2	3	4	
LITHOSIINAE	A. marg. & biop.														0
	A. sericea														0
	An. intervenata			○	○		○			●		●			8
	T. grisescens				●			○			○				4
	T. alba														0
	O. sixola						●		○		○				4
	I. opulentana														0
	G.? phelina														0
	G. paidicus														0
	L. sordida			○	●		○	●	○	●		●			15
D. falsimonia							○							1	
Di. coroides										○				1	
ARCTIINAE	R. sanaea			○				○							2
	I. hippia						○			○					2
	E. aberrans														0
	N. cotes			●			●	●	●		●				24
	A. critheis			●			○		●		●				15
	A. obscurata														0
	A. sicilia						○		○		○				3
	M. incerta			●			●	●	○	●		○			15
	M. asana			●	●		●		○	●					12
	M. laodamia						○								1
B. haemorrhoides														0	
T. imperialis							○	●		●				5	
H. maroniensis														0	
H. sobrina								○						1	
H. grandis														0	
H. catenulata				○										1	
A. minuta														0	
A. semivitrea			○				●				●			11	
B. pervenosa				●										4	
V. rosenbergi			●	○			●	●	●	●	●			33	
E. icasia			●				○							3	

ADDITIONAL RECORDS OF BARRO COLORADO ISLAND
BOMBYCES—Continued
(Scrimshaw Records for June–August)

NAME	Month Week	June				July				Aug.				Total
		1	2	3	4	1	2	3	4	1	2	3	4	
LYM. C. vinasia S. purpurascens S. corydona								○						0 1 1
	C. marmorea		●	●		○	●	○						9
	C. arema					○								0
C. rufescens						○								1
P. xylinoides						○								1
N. superciliosa			●	●		○	○							11
N. lophocera			●			●	○	●		●				13
L. concordens														0
L. mixta														0
L. maltha						●	○	●						7
T. pachydeus							●	○						
T. albosigma			○	○				○		●				5
A. multilinea														0
C. striolata														0
D. guarana				○		○								2
Di. argentilinea														0
A. divisula			○	○				●		○				6
A. rufinsulae			○											1
K. muscosula														0
D. gelduba			○			●		○						4
D. lama							●			○				4
D. perplexa														0
D. tharis			○						●					3
R. distinguenda			○				○							2
R. apella			○			○								2
M. multilinea				○		●	○	○		●				7
D. morona				○			○	●	●					8
H. indistans				●		●	○	○	○	○				8
H. plana			○	○		●	○	○	●	○				15
H. commentica			○											1
H. plusiata			○					○						2
H. colombia			●	●		○	●	●		●				15
H. vecina														0
H. flava								○						1
H. dentata			●						●	○				8
H. meona							○							1
H. nigrescens										●				2
H. sparsipennis			○	○			○	●						5
H. conspirata														0
H. vinicosta			○	○		●								5
H. subochraceum							○			○				2
Ha. curvilinea									●	○				4
Ha. repandens				○			●	●	○	○				7
Ha. simplex								○						1
A. lichyi							○	●						4
C. rarata			○						○					2
D. Sc. leucophleps			●	●		●	●	●		●				25

EUCHROMIIDAE

Homococera stictosoma. A definite summer flight. See diagram.

Gymnelia beata. Mar. 1, '33 (A.M.N.H.).

Gy. salvini. See diagram for further dates

GYMNELIA JANSONIS Butler

Gymnelia jansonis Btl., Cist. Ent., 1, 116, 1872.

Figured: Butler, Lep. Exot., 61: 17; Seitz, 2: b2.

Abdomen with orange on edges of segments (as in *beata*) and first segment pale yellow above (as in *perniciosa*), but no orange on fore wing above.

June 26 (Scr.). Costa Rica to Colombia.

Gymnelia colona. A definite flight in midsummer (see diagram). The species is presumably not limited seasonally.

Phoenicoprocta paucipuncta. See diagram.

PHOENICOPROCTA INSPERATA Walker

Poecilosoma insperata Wlk., List Lep. Ins. Br. Mus., 7, 1606, 1856; *Leucotmemis i.* Hmps., Cat. Lep. Phal., 1, 224; 8: 25, 1898; *Cosmosoma i.* Zerny, Lep. Cat., 7, 66, 1912.

Phoenicoprocta rubiventer Hmps., Cat. Lep. Phal., 1, 198, fig. 94, 1898.

Also figured: Seitz, 12: c1 (as *rubriventris*).

July 2 (Scr.).

Pheia albisigna. July 23 (Scr.)

Loxophlebia flavipicta. July 3, 9 (Scr.)

L. leucothema. June 24 (Scr.)

Mesothera pyrrha. See diagram.

Mesothera ethela. See diagram.

Chrostosoma echemus. See diagram. The summer specimens taken are too few to change the picture, and are probably stragglers.

Cosmosoma saron. See diagram.

C. metallescens. See diagram. Did not dominate over the preceding species as it did in the winter collection.

C. batesi. Mar. 23, '33 (A.M.N.H.) June 23 (Scr.), a specimen with only a black bar on the vertex.

C. xanthostictum. A poor specimen, July 30, may be this species.

C. remotum. Four specimens cover the period (Scr.)

C. teuthras. Eight specimens, June 21-Aug. 6, seem to indicate summer as the proper flight period.

C. stibostictum. Nine specimens, from June 22 to Aug. 6, again indicate a principal flight in summer.

Saurita clusia. Three stragglers (see diagram).

S. nox. See diagram. These may be stragglers, but the lack of specimens before and after suggests a distinct flight.

S. phoenicosticta. See diagram. Like the last.

S. afflicta. Four; see diagram.

Psoloptera thoracica. See diagram. There is evidently a very heavy second flight, especially in August.

Dycladia correbioides. June 24 (Scr.)

Macrocneme thyra. June 26 (Scr.)

M. thyridia. A female taken June 30 may belong here.

M. albitarsia. A specimen taken June 26 is probably this.

M. auripes. See diagram. Summer shows its fair proportion.

Calonotos tiburtus. See diagram. The material shows two separate heavy flights, in late June and third week of July.

Marccidia achrysa. See diagram.

Hypocladia restricta. See diagram.

Chrysostola augusta. Aug. 7 (Scr.)

Episcepsis lenaeus. June 26, July 1, 2 (Scr.)

E. hypoleuca. July 29, Aug. 2 (Scr.)

E. frances. Females June 22-Aug. 3, males July 11 (Scr.), Mar. 19, '33 (A.M.N.H.)

E. lamia. Mar. 1, '33 (A.M.N.H.), July 26 (Scr.)

Napata flaviceps. Feb. 10 (A.M.N.H.)

Aclytia gymamorpha. June 23-Aug. 8 (Scr.)

AGYRTA Hübner

Palpi upturned with third joint correct but very short. Shaft of antenna simple. Fore wing with R_{2-5} stalked, M_1 separate, M_2 and 3 connate, Cu_1 separate, strongly concave above and approximating somewhat to Cu_2 at border; hind wing with R and M_1 connate, M_2 a little separate, M_3 and Cu_1 connate. Tibiae close-scaled.

A striking black, blue and transparent moth. It will run in my key to *Mydromera*, from which it differs in the less convergent Cu_1 and Cu_2 of fore wing and the free M_2 of hind wing, or to *Ctenucha*, which has a longer third segment of palpus and almost invariably fully scaled wings.

AGYRTA DUX Walker

Diophtis dux Wlk., List Lep Ins. Br. Mus., 2, 327, 1854; *Agyrta d.* Hmps., p. 470, in part, not figure.

Figured: Seitz, 24: a1.

Black, with patches of metallic blue, and transparent white markings, a large triangle in and below cell of fore wing and transverse pm. band, a longitudinal streak on hind wing (entering cell, unlike *A. superba*), and streak on inner margin.

Feb. 10, Mar. 19 (A.M.N.H.). Mojinga Swamp, C.Z. (Lawler-CU). Venezuela (type), Peru, etc.

Delphyre rubricincta. See diagram.

D. atava. Six specimens, July 11–Aug. 8, suggest the principal flight is in summer.

HELIURA RHODOPHILA Walker

Eucerea rhodophila Wlk., List Lep. Ins. Br. Mus., 7, 1638, 1856; *Heliura r.* Hmps. Cat. Lep. Phal., 1, 483, fig. 267.

Also figured: Biol., 72; 19 (as *pyrrhosoma*); Ill. Het. Br. Mus., 1, 9: 4 (as *solicauda*); Seitz, 24: f4.

Coloring more brilliant than *H. balia*, the veins of fore wing crimson. Male with an enormous anal lobe, covering a paler patch on the pink abdomen. Appearance closer to *Delphyre aurorina*, but distinguished by the brighter coloring, crimson veins, cream to pink, instead of dark, middle and hind coxae and darker vertex.

June 26–July 23 (Scr.), Tela, Honduras (Bts.). Ranges to the Amazons.

H. balia. Mar. 1, '33 (A.M.N.H.)

H. maculosa. July 3 (Scr.)

Eucereon obscurum. Two. See diagram.

E. rosa. June 29, July 10, Aug. 5 (Scr.)

E. atrigutta. July 10 (Scr.)

E. latifascia. July 19, Aug. 7 (Scr.)

E. dentatum. July 15 (Scr.)

E. aeolum. June 24, July 15, Aug. 7 (Scr.)

E. flavicaput. Aug. 8 (Scr.)

E. intranotata Dgn. July 21, Aug. 6 (Scr.)

Correbia lycoides. Five; June 26–Aug. 8 (Scr.)

C. undulata. Aug. 7, 8 (Scr.)

Correbidia terminalis elegans. See diagram. Form *elegans* was relatively commoner than in the other season.

Family NOLIDAE

No Nolidae have been received; they are both obscure and fragile, and presumably did not happen to be taken.

Family ARCTIIDAE

Subfamily LITHOSIINAE

Agylla sericea. Aug. 8, females only.

Apistosia judas. July 29 (Scr.)

Antona dorsisigna. Nine specimens, June 26–Aug. 1. This species was relatively commoner than in the winter.

A. interrenata. See diagram.

Thyone griseescens. See diagram.

Thyone alba. Feb. 8, (A.M.N.H.)

Odozana sixola. See diagram.

Illice citrina. Feb. 13 (A.M.N.H.)

Lycomorphodes sordida. Rather common; see diagram.

Talara phacella. Mar. 18 (A.M.N.H.)

Dolichesia falsimonia. July 11 (Scr.)

Paraprepia fuscilingua. June 30 (Scr.)

Subfamily ARCTIINAE

Robinsonia sanae. See diagram.

Idalus hippia. Two. See diagram.

Phaeomolis lepida. Mar. 18 (A.M.N.H.)

PHAEOMOLIS OSMOPHORA Hampson

Amaxia osmophora Hmps., Cat. Lep. Phal., 3, 32, fig. 25, 1901.

A. dyuna Dr., Biol., 2, 369, 1897 (not Schaus).

? *A. pyga* Schs., Proc. Zool. Soc., 1892, 279; Am. Lep., 1: 27, 1892.

Also figured: Seitz, 45: e7.

This is very close to *P. lepida*, but can be separated as follows:

2. Subterminal and terminal spots connected; postmedial spot in cell

M_1 bifurcated *lepida*

Subterminal and terminal spots separate; pm. spot in cell M_1
rounded like the rest; all marks paler *osmophora*

I can see no difference between *Amaxia* and *Phaeomolis*, unless the genotype of *Amaxia* has R_2 as cited in Hampson's key rather than as figured. This species shows the structure of *Phaeomolis*, but if the key is wrong and the figure correct should be restored to *Amaxia*.

June 29 (Scr.). Costa Rica (type): Peru (C.U.).

Neritos cotes. See diagram. The summer flight is evidently the principal one.

Parevia griseotincta. July 11 (Scr.)

AUTOMOLIS TRIPUNCTATA Druce

Sutonocrea tripunctata Dr., Biol. Centr., Am. Lep. Het., 1, 79, 9:2, 1884.
Also figured: Seitz, 50: h1.

Structure of Section II B, but with a slight anal lobe in male. Yellow, with gray spot on thorax, ante- and post-medial bands not quite reaching inner and outer margins respectively, and three gray spots in disc.

July 8 (Scr.). Also found in Costa Rica.

A. sicilia. See diagram.

A. diluta. June 24-Aug. 5 (Scr.)

Melese incerta. See diagram. Evidently the summer flight is a well developed one.

M. asana. A good summer flight (see diagram.); as usual only 2 out of 12 are males.

M. laodamia. July 5 (Scr.)

Tricypha imperialis. See diagram.

Elysius diseiplaga. July 25 (Scr.)

Thalesa citrina. July 10 (Scr.)

Halysidota maroniensis. Feb. 29 (A.M.N.H.)

H. sobrina. July 10 (Scr.)

H. interlineata. July 11 (Scr.)

H. grandis. Feb. 28 (A.M.N.H.)

H. iridescens. June 21, Aug. 8 (Scr.)

H. catenulata. June 29 (Scr.), Mar. 10-13 (A.M.N.H.)

*Agoraea semivitre*a. The material indicates a good summer flight. See diagram.

Bituryx pervenosa. See diagram.

Virbia rosenbergi. Males were as abundant as in other months, but the three females were all taken July 1-7. (See diagram).

V. fluminea. June 24 (Scr.)

Eepantheria icasia trinitatis. June 21, July 5 (Scr.).

Family PERICOPIDAE

Hyalurga fenestra. Mar. 10, 16 (A.M.N.H.), June 24 (Scr.)

H. soroides. June 21, 26 (Scr.)

Family LYMANTRIIDAE

PHIDITIA Möschler

Tympanum absent; fore wing without accessory cell, R_1 free, R_2 , $3+4$ and 5 stalked; hind tibia with all spurs. Antennae normal for the

Lymantriidae, ocellus absent; tongue weak, palpi rather weak, oblique to middle of front; fore wing quadrid, with middle discocellular angled; hind wing with Sc definitely fused to cell for a short distance, R and M_1 stalked, lower discocellular only a little shorter than middle one.

An aberrant genus, in some ways more like the Old-World Endromiidae and the Apatelodes group, but genitally a Lymantriid, and usually put with the Lymantriids. The loss of the tympanum is extremely aberrant for the Noctuid series, but occurs in some wingless female Lymantriidae, and also in the old-world Amata (Syntomis) group. The larva is apparently unknown; it will probably have the tufted feathery hair of the Lymantriidae, but may lack the glands.

PHIDITIA CUPREA Kaye

Tarchon cuprea Kaye, Trans. Ent. Soc. London, 1901, 142, 6: 11.

Also figured: Seitz, 75: a2.

Medium sized, brown, with normal but not contrasting transverse banding, and a translucent st. lunule in cell R_5 . Hind wing with two deep scallops toward anal angle.

June 21 (Scr.). Guatemala to Guiana.

In *P. lucernaria* Walker (Seitz 74: i3), which is regional, the ground is paler and there are five large translucent st. lunules.

Sarsina purpurascens. July 11 (Scr.).

Stacterinia corydona. July 26 (Scr.)

Desmoloma chironomus. June 24, Aug. 8 (Scr.)

Family NOTODONTIDAE

A genitalic survey of this family by Mr. J. G. Franclemont shows that our present classification is very largely artificial, and that I improved it but little in the grouping published in the first portion of this report. Besides a revision of the grouping there prove to be several very close species, easily distinguished only by the genitalia, notably in the case of *Dasylophia* and the North American genus *Symmerista*. He has contributed a report on the genus *Antaea*, below, since the Barro Colorado species turns out distinct from the South American *A. juturna*. *Hapigia* also needs revision, but types must still be examined. The Barro Colorado species are probably correctly determined, but there may be more species.

The Notodontidae are adapted to humid conditions, and we note that a good many species which were taken in good numbers from

Nov. to March were absent or only appeared as stragglers the rest of the year. Examples are *Pentobesa xylinoides*, *Lepasta concordens* and *mixta*, *Crinodes striolata*, *Dasylophia guarana*, *Didugua argentilinea*, *Kalkoma muscosula*, *Disphragis perplexa*, *Rifargia distinguenda*, and *Hemiceras plusiata*, *vecina*, *flava*, *dentata*, *meona*, *conspirata* and *vinicosta*. On the other hand more specimens of *Nystalea ebalca* were taken in June–July.

Rosema deolis. June 22, July 3 (Scr.)

R. zelica auct. There are a dozen or more neotropical species now confused under this name; in the absence of males we cannot be sure which is the Barro Colorado one, but it is probably undescribed.

Calledema marmorca. See diagram. The range of dates now suggests two flights, Nov.–Mar. and May–July.

C. jocasta. July 15 (Scr.)

C. rufescens. July 1 (Scr.)

Pentobesa xylinoides. July 5 (Scr.)

Nystalea ebalca. June 21–July 27, 9 specimens (Scr.)

N. superciliosa. See diagram. In this species, as in *Calledema marmorca* and a few others there is a definite summer flight.

NYSTALEA VIRGULA Felder

N. virgula Fld., Reise Novara, Lep., 98: 3, 1874.

Also figured: Seitz, 143: i4.

Fore wing even more pointed than in *superciliosa*, brown, shaded with olive, with a longitudinal shade from base to just below apex, defined with darker brown below it toward base, but above it on outer half of wing; a double black pm. lunule beyond cell.

July 5, 1940 (Scr.) General in tropical America.

N. nyseus. July 21 (Scr.)

NYSTALEA MARONA Schaus

N. marona Schaus, Proc. U. S. Nat. Mus., 29, 233, 1906.

Not figured. (similar to *N. multiplex* Dgn., Seitz, 143: g2). Omitted from Lep. Cat.

Confused fine blackish markings on a pale gray base. Basal 2/5 mainly blackish; stronger blackish striations outlining two vague ocellate spots between pm. and inner st. lines, centering in cells R_5 and M_3 ; outer st. line single, finely and doubly dentate, unlike *N. multiplex* which has two inner st. lines, and the more basal nearly even.

Runs to 6 in my key but differs from *lophocera* and *plumipes* in the absence of striking sex-modification, larger size, more solidly dark base, vague pm. ocelli and doubly dentate outer st. line.

Aug. 8 (Scr.). Guiana (type), middle Amazons (C.U.).

N. plumipes. June 20, 21, Aug. 8 (Scr.)

N. lophocera. See diagram. There is a well-marked summer flight.

Proclymiotis joanna. July 3, 16, 25, Aug. 8 (Scr.)

Lepasta maltha. See diagram.

Strophocerus albonotatus. June 24, 30 (Scr.)

Tachuda pachydexius. See diagram.

T. albosigma. May 27 (Fried.) also June-Aug. (Scr.) See diagram.

Crinodes beskei. July 16 (Scr.)

Dasylophia terrena. June 21 (Scr.)

D. guarana auct. This is probably not the true *D. guarana*, described from S. Brazil, but only genitalic study will decide the question.

Betola aroata. June 24, 30, July 1, 9 (Scr.)

Lirimiris lignitecta. June 23 (Scr.)

DICENTRIA MARIMBA Schaus

D. marimba Schs., Proc. U. S. Nat. Mus., 57, 143, 1921.

♀ *D. limosoides*, Schs., Ann. Mag. Nat. Hist., (8) 7, 270, 1891 (not ♂).

Not figured.

Larger than *D. violascens*. Male dull brown, confusedly flecked and shaded with black and green, the most conspicuous markings being a subterminal series of black bars between veins, alternating with blackish terminal shades on the veins. Hind wing mainly white, with contrasting blackish costa. Female not at hand.

Aug. 8 (Scr.) Guatemala (type) to Guiana (C.U.)

Malocampa puella. Mar. 20, '33 (A.M.N.H.)

Farigia pallida. June 21 (Scr.)

PAULUMA Schaus (with *Euphastia* Dognin)

Somewhat related to *Disphragis*. Antenna typically half-pectinate, brush-like in *P. (E.) nubila*; basal segment smooth above, with a small tuft below. Palpi rather short, with short third segment. Fore wing with very long acc. cell, M_1 well separated; hind wing with Sc closely parallel to cell over its middle half or more, then abruptly diverging; M_3 and Cu_1 almost connate. Rather small species.

The present species will run in our key to 39, where it is distinguished by the well separated M_1 on fore wing.

PAULUMA NUBILA Druce

Rifargia nubila Dr., Ann. Mag. Nat. Hist., (7) 7, 437, 1901.

Figured: Seitz, 152: k5.

Mixed gray and brown, with a large round black patch over end of cell, in July specimen suffusing out toward outer margin; st. lines far out toward margin. 24 mm.

Mar. 18. (A.M.N.H.) Apr. 23 (Fried.), July 26 (Scr.). Ranges to Guiana.

Disphragis gelduba. Four, see diagram.

D. lama. Only 4 specimens taken, see diagram.

D. biundata. June 21, July 25 (Scr.) This is almost certainly a distinct species from the true (North American) *biundata*.

D. tharis. See diagram. The few summer specimens are not really enough to indicate a distinct summer flight.

D. matralis June 24 (Scr.) a somewhat suffused female.

Talmeca consociata. Jan. 5 (Fried.)

Rifargia demissa. July 25 (Scr.)

R. apella. See diagram.

R. distinguenda. See diagram.

Magava multilinea. See diagram.

Drugera morona. See diagram.

Rincodes flavodiscata. See diagram.

Rhuda focula. July 20 (Scr.)

BLERA CUCULLIOIDES Schaus

Chadisa (?) *cucullioides* Schs., Proc. U. S. Nat. Mus., 29, 266, 1906.

Boriza lignosa Dgn., Ann. Soc. Ent. Belge, 53, 83, 1909.

Tagela (?) *pigritia* Dgn., Het. Nouv. Am. Sud, 8, 90, 1914.

Figured: Seitz, 152: f4.

Male tawny, female dull wood brown, with ordinary lines strongly zigzag and pm. marked with black points between veins, giving a dead wood effect. Pm. dot in cell Cu_1 enlarged.

Mar. 23, '33 (A.M.N.H.). Lino, Panama (Dognin), Guiana (types).

We have all been doubtful of the generic position of this species, which does not look the least like a *Blera*, but has the widely separated Sc and R of hind wing. M_1 of fore wing arises from middle of acc. cell.

Hemiceras indistans. See diagram. In contrast to the species mentioned above, *indistans*, *plana*, *colombia* and *subochraceum* have definite summer flights, and are presumably double brooded. The evidence as to the *Hapigias* is less definite.

H. plana, *commentica*, *plusiata*, *colombia*, *flava*, *subochraceum*. See diagrams.

H. cadmia. July 3, 9 (Scr.)

H. meona. July 9 (Scr.)

H. lepida June 24, 30, (Scr.)

H. nigrescens. June 16 (Fried.) also see diagram.

HEMICERAS RUFESCENS Walker

Gadiana rufescens Wlk., List Lep. Ins. Br. Mus., **33**, 854, 1865.

H. casiaclara Dgn., Het. Nouv. Am. Sud, **25**, 14, 1924.

Figured: Seitz, 156: k1.

Light reddish buff, nearly even, with traces of the oblique brown st. shade of *conspirata* and *colorata*. Ground of hind wing white, slightly creamy, the light brown veins, stigma and terminal shade-line strongly contrasting.

Feb. 11 (A.M.N.H.). Lancetilla, Tela, Honduras (Bts.). Brazil (type *rufescens*) and Amazons (type *casiaclara*); South Brazil (C.U.). Also reported by Dyar from the Zone. (Accidentally omitted from main report).

HEMICERAS COLORATA Dognin

H. colorata Dgn., Het. Nouv. Am. Sud, **22**, 11, 1923.

Figured: Seitz, 156: g5.

Similar to *conspirata*, smaller, without the contrasting pale costa; oblique st. patch near margin rather distinct, yellowish brown than general ground, and lacking the faint violet iridescence. Hind wing as in *rufescens*.

July 26 (Scr.) a female apparently of this species. Guiana to S. Brazil.

H. walkeri. June 21 (Scr.)

H. deornata. July 25 (Scr.)

H. monegonda. June 24 (Scr.)

H. quebra. June 21, July 3 (Scr.)

Hapigia curvilinea. See diagram.

H. repandens. See diagram.

H. simplex. See diagram.

H. raatzii. July 9 (Scr.)

Antaea. Mr. J. G. Françlemont has contributed the following analysis of the group to which the Barro Colorado species belongs. One or two species now standing in *Hapigia* are very closely related.

Key to species of ANTAEA based on superficial characters

1. Subterminal line with three sinuses; color distinctly grayish 2
 Subterminal line with two sinuses; color brownish to sepia 3
2. Termen of forewing more or less even; subterminal line straight from costa to vein M_3 ; the other lines rigid *licormas*
 Termen of forewing distinctly crenulate, especially below vein M_3 ; subterminal line inwardly arcuate to vein M_3 ; the other lines somewhat waved *omana*
3. Last abdominal sternite, in situ, heavily chitinized, the excavation distinctly twice as deep as wide; the color warm sepia . . . *juturna*
 Last abdominal sternite, in situ, lightly chitinized, except the tips of the lateral projections which are well chitinized, the excavation about as deep as wide; color pinkish brown 4
4. Underside of hindwing of male covered almost entirely with velvety yellow sex scaling; upperside of hindwing fuscous *lichyi*
 Underside of hindwing of male with the yellow scaling restricted to the basal area, the remainder of the wing fuscous; upperside of hindwing black *jaraguana*

ANTAEA LICHYI Franclemont spec. nov.

Antaea juturna (pars) Auct.‡*Antaea juturna* Forbes, Bull. Mus. Comp. Zoöl., **85**, no. 4, (Lepid. Barro Colorado Is., Panama), 317, 1939.

Male, general color of head, thorax and fore wings warm pinkish brown. The basal, antemedial, median and postmedial lines straight; the basal erect, not reaching the inner margin; the antemedial erect; there is a faint indication of an erect line between the basal and antemedial lines, it extends from the inner margin to vein Cu; the median, which is interrupted by the reniform, and the postmedial lines oblique; the subterminal line straight from costa to vein Cu_1 , where it is offset inwardly, it is again offset inwardly on vein Cu_2 ; all the lines fine, light and raised, bordered exteriorly by fine dark lines; the reniform and orbicular large, inverted u-shaped, indicated by fine, pale raised lines; the orbicular sometimes closed below; the veins lightly marked with hoary; the costa pale. The hind wings fuscous above, below clothed almost entirely with velvety yellow sex scaling.

Female generally darker.

Male genitalia (fig. 67) symmetrical; the uncus short and trigonate; the socii moderate; the tegumen moderately long and broad; the

vinculum long and moderately broad; the valves approximately three times as long as wide, with pleated areas at the outer basal angle and at the apex, the costa with a long curved projection, the exact homology of which is not known, the outer basal angle with a large tuft of long scales (omitted on the left valve in the drawing); the aedoeagus long and stout, the apex bearing a projection which is produced into a spur on one side; the vesica armed with long triple branched spines, these are apparently shed during the act of copulation.

The last sternite of the abdomen excavated on its posterior margin, the two resulting lateral projections heavily chitinized at their apices.

This species is at once separable from *juturna* Cramer by its lighter color throughout, that species being more or less sepia in tone. The genitalia of *juturna* differ in that the uncus is narrow, the costal projection bears a tuft of hairs on its inner face, in none of the preparations of the genitalia of *lichyi* are any such tufts discernible, the aedoeagus bears two spines laterally near the apex, one considerably larger than the other. The last abdominal sternite of *juturna* is approximately twice as deeply excavated and the lateral projections twice as large and more extensively chitinized.

Holotype. ♂, Massif du Naiguata, D. F., Venezuela, Sept. 23, 1938 (Rene Lichy), [in Coll. Franclemont].

Allotype. ♀, Route Maracay-Choroni, Aragua, Venezuela, Sept. 19, 1937 (René Lichy), [in Coll. Franclemont].

Paratypes. 1♂, Massif du Naiguata, Venez. (René Lichy), [in Coll. Franclemont]; 1♀, Quiriquire, Monagas, Venez., [in Coll. Cornell Univ.]; 1♂, Colombia, [in Coll. Am. Mus. Nat. Hist.]; 11♂♂ 5♀♀, Barro Colorado Island, Panama, [7 in Coll. Am. Mus. Nat. Hist., 6 in Coll. Mus. Comp. Zool., 3 in Coll. Cornell Univ.]; 1♂, Lancetilla, Tela, Honduras, [in Coll. Mus. Comp. Zool.].

This species is dedicated to Mr. René Lichy, a very kind friend, who has most generously supplied the writer with many specimens of Venezuelan Heterocera. It gives me great pleasure to name this striking form for so diligent and assiduous a collector.

ANTAEA JUTURNA Cramer

Phalaena juturna Cramer, Pap. Exot., 2, 48, pl. 129, fig. E, 1780.

Antaea juturna (Cramer), Hübner, Verz. bek. Schmett., 266, 1823.

Ophisma? juturna (Cramer), Walker, Cat. Lep. Het. B. M., 14, 1373, 1858.

Antaea juturna (Cramer), Schaus, Trans. Ent. Soc. Lond., (1901), 342, 1901.

Antaea juturna (Cramer) [pars], Draudt, in Seitz, Macrolep., 6, 1049, 1934.

A specimen from Moengo, Surinam, which is in the Cornell University Collection, agrees excellently with Cramer's rather crude figure of this species. The moth, the darkest in the genus, is warm sepia in color, with fine pale lines, following the course of those in *lichyi*. The reniform is smaller than in either of the new species and is always closed below; it never extends to vein Cu. The hind wing is fuscous black above, and below the yellow sex scaling is restricted to the base of the wing, the greater part of the wing being dark fuscous.

The outstanding character of the male genitalia (fig. 68) in this species is the two spines on the side of the aedoeagus.

The last sternite of the male is quite characteristic; its deeper sinus and stouter projections, heavily chitinized, readily separate it from the two new species. This character may be easily seen by denuding the last sternite.

The species has been seen from Venezuela, British Guiana, Dutch Guiana (Surinam) and the Amazon region of Brazil. [Cornell University also has a damaged specimen from the Ellsworth collection, presumably from Fyzabad, Trinidad.—W.T.M.F.]

ANTAEA JARAGUANA Franclemont spec. nov.

Antaea juturna (pars), Auct.

‡*Antaea juturna* Draudt, in Seitz, *Macrolep.*, 6, pl. 158, fig. c-1, 1938.

Superficially this species is very close to *lichyi*; it may subsequently prove to be a southern race of that species, but in the material before the writer the two do not overlap in distribution; *lichyi* being apparently restricted to Central America and the coastal portion of northern South America, while this species occurs from the Amazon region to southern Brazil. The third species of this group, *juturna*, ranges between the two new species, overlapping *lichyi* in the north and *jaraguana* in the south; it is more distinct from either of the new species than they are from one another.

Male, the head, thorax and forewings about the same color as *lichyi*, but with a distinctly pinker cast; the lines following the same course as those of that species, but wider and somewhat more raised. The hindwings above are fuscous black, much darker than *lichyi*; the underside with the sex scaling yellowish only toward the base, the greater part of the wing fuscous.

The female of this species is unknown to me at present.

The male genitalia (fig. 69) are similar to *lichyi*, but differ from that species in that the costal projection is stouter and has two tufts

of hair, one on the inner and one on the outer face; and in that the apical projection of the aedoeagus ends in two spines; the genitalia are, as a whole, larger in size than those of *lichyi*.

The last abdominal sternite of the male is more widely excavated than that of *lichyi*.

From *juturna* this species differs in the same manner as does *lichyi*, but the costal projection of the genitalia is stouter and has the two hair tufts mentioned previously.

Type. ♂, Jaragua, Santa Catharina, Brazil, Nov. 27, 1934, [in Coll. Franclemont].

Paratypes. 4♂♂, Jaragua, Santa Catharina, Brazil, June, November and December, [in Coll. Franclemont]; 2♂♂, Below Codajos, Rio Solimoes, Brazil, 1♂, Above Obidos, Amazon, Brazil, [in Coll. Cornell Univ.].

Chliara cresus. July 25, (Scr.)

Rhapigia accipiter. July 5 (Scr.)

Cerura rarata. See diagram. It still looks as if the main flight may be between June and August. These few specimens are again females.

Family DIOPTIDAE

Scotura fulviceps leucophleps. See diagram. There is a heavy summer flight.

ZUNACETHA Walker

Structurally hardly distinct from *Scotura*, but usually with Cu_1 of fore wing connate with M_3 (usually stalked in *Scotura*); entirely different in pattern, white with fine transverse black lines on basal half of fore wing, and lines alternating with the veins on outer half. Besides the following species *Z. bugabensis* Dr. (Biol., 61: 15; Seitz, 68: 17) is regional; it lacks the yellow and black on basal half of costa and at end of cell.

ZUNACETHA ANNULATA Guerin

Lithosia annulata Guer., Icones Règne Anim., 519, 1844.

Z. bipartita Wlk., List Lep. Ins. Br. Mus., 27, 134, 1863.

Mieza nervosa Fld., Reise, Novara, Lep., 139: 43.

Also figured: Seitz, 58: k1.

Basal $2/5$ of costa and a spot in cell yellow, bordered with black, the two areas often connected. Hind wing white with a broad gray border.

June 25, 27 (Scr.) Ranges from Texas to Guiana.

Josia ena. Aug. 2, 1924 (Banks).

J. cruciata Feb. 9 (A.M.N.H.)

Remaining Families

In the following account of the remaining families of so-called Bombyces, the Scrimshaw data have been incorporated in the records and diagrams.

Most of these families have never had critical revisions, but are catalogued and figured in Seitz's "Macrolepidoptera of the World" vol. vi. In the references to figures this is abbreviated merely "Seitz" with plate, line and number of the figure on the line, counting from the left. The figures in the "Biologia Centrali-Americana, Heterocera" are cited merely as "Biol." with plate and figure numbers, but other literature is less drastically abbreviated. Bibliography is given a little more fully than in previous instalments, but references already published in the "Lepidopterorum Catalogus" are not repeated unless for some special reason. Further references differ widely from family to family and are given under the families.

INDEX OF GENERAL WORKS BY FAMILIES

Family	Biologia		Seitz		Lep. Cat.
	Text	Plates	Text	Plates ²	
Sematuridae	ii, 4-8, 325	41, 9S	S31-S37	138, 139	30(2)
[Thyatiridae]	i, 257; ii, 470		1171-1176	172	25]
Lasiocampidae	i, 200-208; ii, 428-436	21, 22, 85-87	565-628	75-88	73
Eupterotidae	i, 224-226, ii, 445-446	23 ³ , 8S	675-711	89	not out
				140-142	
Uraniidae	ii, 3-4	41	829-S30	138	30(2)
Epiplemlidae	ii, 123-128, 540	53, 99	1140-1170	170-172	30(1)
Mimallonidae	i, 227; ii, 446-448	24, 8S	635-673	87, 8S	50
Thyrididae	ii, 184-188, 545	59	1187-1213	173-175	20
Castniidae	i, 24-28; ii, 320	3, 4, 68, 69	5-19	1-8	15
Cossidae	i, 230-231; ii, 449, 450	24, 8S-90	1263-1287	181-184	29
Psychidae	i, 228-230		1177-1187	169	34
[Egeriidae]	i, 29-34; ii, 321-326	5, 69	1215-1262	176-180	31]
Zygaenidae	i, 37-40; ii, 330	6, 70	21-31	9	71 ⁵
	i, 120; ii, 393 ⁴	12, 77 ⁴			
Eucleidae	i, 210-222; ii, 439-444	22, 23, 87, 8S	1104-1139	164-166 ⁶	32
[Epipyropidae]	i, 230	24	1313-1315	168	16]
Dalceridae	i, 213, ii, 441		1303-1312	168 ⁷	16
Megalopygidae	mixed with Lasiocampidae, etc.		1071-1099	160-163 ⁸	16
Hepialidae	i, 232-234; ii, 450-452	24, 89	1289-1302	99-100	4
				185	

¹ None yet taken on Barro Colorado Id.,—regional.

² A few scattered figures on other plates.

³ *Zanola* and *Tarchon* with *Limacodidae* (i.e. *Eucleidae*).


⁴ As *Chalcosiidae*.

⁵ Also fasc. 33, containing only old-world species.

⁶ Pl. 167 not yet published.

⁷ *Dalcera* on pl. 185.

⁸ *Aidinae* on pl. 164.



SEMATURIDÆ

(Manidiidae; Uraniidae in part)

Head strong, with antennae simple and tending to be somewhat swollen before the tip, which may be fine; chaetosema very large, reaching forward to base of antenna between ocelli and eyes; eyes hairy; palpi upturned about to middle of front, with third segment lengthened in female, extremely long in female *Sematura*; eyes hairy. Fore wing (fig. 70) with costal venation crowded: R_1 free, M_1 short-stalked, M_2 above middle of cell; hind wing with small to minute frenulum; frenulum-base thickened and costal area lobed, Sc much swollen and sharply angled at a point near base, where it is connected by a brace to base of frenulum, as in the Geometridae; beyond this Sc and R approach closely along the cell or are connected by a "cross-vein" (i.e. the free fragment of R_1 , as usual); cell very short, M_3 and Cu stalked, and sometimes Cu_2 out of their base; anal area very narrow.

This is a very small family, frequently grouped with the Uraniidae, but, as I believe, more closely related to the most primitive Geometridae. They agree with the latter in wing structures, and to some extent in pattern (especially *Anurapteryx*) but I have found no sign of a tympanum. The larva is reported by Kirby¹ as "with sixteen legs; head and pro-thorax small; body with conical tubercles and curved spines"; the pupa as "enclosed in a loose cocoon at the base of a folded leaf; the sheath for the proboscis continued beyond the wing-cases, and the extremity of the body forming a short, deflexed spine." Nothing more is known of the life history of the family.

Besides the following species, which is the only one known from Barro Colorado Id., tropical America has a number of species of *Coronidia*, some of which should be found on the island. One *Anurapteryx* has been described from Arizona, and Janse considers *Apoprogenes hesperistis* of Africa to belong here, but the family is otherwise strictly neotropical. *Apoprogenes* has the venation and antennae of *Coronidia*, but differs in the naked eyes and absence of ocelli, and is, as I suspect, a parallel development of the primitive Hemitheine Geometridae.

¹ Hand-Book of the Order Lepidoptera, in Lloyd's Natural History, vol. iii, p. 55.

Diagram II

SEASONAL RECORD OF PRIMITIVE BOMBYCES AT BARRO COLORADO ISLAND

NAME	Month Week	Jan.				Feb.				Mar.				Apr.				May				June				July				Aug.	Oct.				Nov.				Dec.				Number
		1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4						
LASIOCAMPID.	E. vittabunda	●	○	○				○		●	○	○		●	○	○				○				○				○				○				○				○		25	
	E. rundala													●	○	○						●	○	○		●	○	○		●	○	○		●	○	○		51					
	E. rivulosa							○										●	○	○						●	○	○		●	○	○		●	○	○		50					
	E. amathuria	●	○	○	●					●	○			○	○	○		●				●	○	○		●	○	○		●	○	○		●	○	○		42					
	E. modesta			○	○					●	○																			●	○	○		●	○	○		14					
	E. marginata	●	○	○	○	●				○	○			●	○	○										●	○	○		●	○	○		●	○	○		45					
E. sobrina					○	○			○	○			○	○			○	○			○	○							○	○			○	○			○	○			14		
	E. attenuata	○	○	○		○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			34	
EUPTEROTID.	E. muscosa	●	○	○	○	○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			33	
	Q. veca	○	○	○	○	○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			51					
	A. antica	○	○	○	○	○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			66					
	T. marmorea	○	○	○	○	○	○			○	○			○	○			○	○			○	○			○	○			○	○			○	○			26					
EUPTEROTID.	A. lapitha	●	○	○	○																	○	○							○	○			○	○			○	○			20	
	A. tuisa	○	○	○	○																									○	○			○	○			13					
	O. reperta	○	○	○	○																	○	○							○	○			○	○			14					
	O. irrorata	○	○	○	○																	○	○							○	○			○	○			26					
EPIPIEM.	S. drepanata	●	○	○	○	●	○	○		○	○																			○	○			○	○			○	○			22	
	S. druidaria	○	○	○	○	○	○			○	○																			○	○			○	○			44					
	N. mutilaria	○	○	○	○	○	○			○	○																			○	○			○	○			12					
	E. incendiaria	○	○	○	○	○	○			○	○																			○	○			○	○			28					
	E. palpulata	○	○	○	○	○	○			○	○																			○	○			○	○			33					
MIM.	C. solvens					○																								○	○			○	○			22					
	Z. infantilis					○																								○	○			○	○			30					
	M. lantona					○								○	○															○	○			○	○			33					
THYR.	D. speculifera																																	○	○			○	○			12	
	D. confusata																																	○	○			○	○			12	
	R. hedilalis																																	○	○			○	○			11	
	R. rufigrisea																																	○	○			○	○			13	
COSSIDAE	X. comisteon	●				○																																				12	
	X. pyracmon	○	○	○	○	○	○																																			13	
	X. terraforma	○	○	○	○	○	○																																			18	
	G. necreros	○	○	○	○	○	○			○	○																															45	
	G. perfida	○	○	○	○	○	○			○	○																															71	
	L. lunifera	○	○	○	○	○	○			○	○																															34	
EUCLEIDAE	E. cippus																																									14	
	E. norba	○	○			○	○			○	○																															73	
	E. buscki	○	○			○	○			○	○																															38	
	T. rubicolor	○	○			○	○			○	○																															60	
EUCLEIDAE	Ta. mas	○	○																																							13	
	S. distincta	○	○			○	○			○	○																															21	
	E. cupreitin.	○	○			○	○			○	○																															14	
	E. elaeasa	○	○			○	○			○	○																															20	
	N. fusea	○	○			○	○			○	○																															107	
	N. micharta	○	○			○	○			○	○																															14	
EUCLEIDAE	N. l. dognini	○	○			○	○			○	○																															13	
	P. sericea																																									33	
	P. repetita	○	○			○	○			○	○																															73	
	Pa. perurnata																																									18	
EUCLEIDAE	D. didyma																																									20	
	P. trianguli.	○	○			○	○			○	○																															12	
IDAL.	A. philetera					○	○			○	○																															115	
	A. dulciola					○	○			○	○																															18	
MEG.	N. pura																																									16	
	M. opercularis	○	○			○	○																																			17	
	P. albicollis	○	○			○	○																																			34	
H.	D. assa																																									35	

○—1 capture in this week.

●—2 or more captures in

References

- Guenée, Hist. Nat. Lep. Het., 9, 16-23, 1857, with pl. 1 (as Sematuridae).
 Sharp, Cambridge Nat. Hist. Ins., 2, 419, 1899 (larva of Coronidia).
 Forbes, Jour. N. Y. Ent. Soc., 25, 47, 1917 (in key only, as Coronidiinae).
 Hampson, Novit. Zool., 25, 367-376, 1918 (as Sematuridae).
 Dalla Torre, Lep. Cat., 30 (2), 9-15, 1924 (as Coronidiinae).
 Biol., 2, 4-8, 525, pls. 41, 98; Seitz, 6, 831-837, pls. 138, 139; p. 830 (larva and pupa after Fassl).

Key to Genera

1. Tibiae spinulated; tail of hind wing spatulate and twice as long as wide; male without frenulum hook. *Sematura*
 Tibiae unarmed; tail of hind wing short or not spatulate (absent in *Anurapteryx*); male so far as examined with a functional though minute frenulum hook. *Coronidia*

SEMATURA Dalman

(*Mania* Hübner, not Treitschke¹; *Manidia* Westwood)

This genus has only three representative species, the present one from the Amazon northward, the hardly distinct *S. diana* from south Brazil, and *S. aegistus* F. from the Greater Antilles. The latter is quite distinct, with scalloped transverse banding; and it even breaks up into minor island races.

SEMATURA LUNUS Linnaeus

♂ *Phalaena Noctua lunus* L., Syst. Nat. (Ed. 10), 508, 1758; Cramer, Pap. Exot., 3, 13, 200: A, 1782.

♀ *Papilio Eques empedocles* Cr., Pap. Exot., 3, 11, 199: A, B, 1782.

♂ *Manidia selene* Guenée, Sp. Gen. Lep. Het., 9, 18, 1857.

♂ *S. actaeon* Fld., Reise Novara, Lep., 121: 5, 1874 (♂ as ♀).

Also figured: Hbn., Samml. Exot. Schm., 1, 202: 3, 4 (♀ as *empedoclaria*); Seitz, 138: c1 ♂, 4 ♀; 139: a1 ♂, 5 ♀ as *selene*).

Early Stages: Kirby, l.c.

Male with more buff ground, female with contrasting luteous post-medial stripe, etc., otherwise dull brown on luteous. Female with extremely long third segment of palpus. The spatulate tail has two large ocellate spots. All works keep *lunus* and *empedocles* as separate species, but the large number I have seen run all *lunus* ♂, *empedocles* ♀.

May 27, June 4, 22 (Fried.) Mexico to Brazil; represented by the slight race *diana* Gn. in southern Brazil.

¹ In fact *Mania* Hübner 1823, is valid over *Mania* Tr. 1825, but only the latter has had any currency in the last century.

[THYATIRIDÆ

Mostly stout-bodied, Noctuid-like moths, with Sc and R of hind wing closely parallel beyond the end of the cell, and tympanum pointing backward at rear of first segment of abdomen. Larva smooth, with head decidedly wider than high, living in a folded leaf.

Thyatira mexicana H. Edw. (Seitz, 172: i3) ranges from Arizona to Peru or further, and will certainly be found in the Canal Zone. It is mottled gray-brown, with contrasting pale patches, and the larva is probably dead-leaf brown, with oblique lateral shading. The related European *T. batis* feeds on *Rubus*.]

LASIOCAMPIDÆ

Very stout and heavy moths. Head small and retracted, palpi moderate to small, but remaining mouth parts absolutely vestigial; ocelli and chaetosema absent; antennae broadly pectinate in male and almost invariably in female also. Vestiture of body deep and fluffy, of wings deep, soft, formed of two kinds of scales, one or both of which are often much modified, frequently reduced to forked hairs;—the wings sometimes translucent as a result. Fore wing with R_4 and 3 long-stalked, rarely united, R_5 stalked with M_1 and sometimes R_3 from their base, M_2 arising near lower angle of cell and Cu_2 very close to base, often easy to mistake for 1st A, which is absent. Hind wing with humeral angle much widened, without frenulum, and supported by one or more humeral veins, arising from a humeral cell (at least in part); M_2 as in fore wing, but Cu_2 from further out on cell. All veins extremely stout, the moth frequently being able to expand its wings without letting them hang down. These so-called humerals are fully supplied by tracheae.

Larva stout and normally much flattened, with tufts from lateral lappets on body; the hair dense and fine, mostly secondary, but with the hairy warts often diffuse or obscure; secondary hair present even on mouth-parts; hooks of prolegs biordinal. Cocoon normally heavy, often double and often with the meshes filled with a gummy or powdery secretion; pupa finely hairy.

A large and world-wide family, but most at home in the old world tropics; America has few genera, but *Euglyphis* is rich in species. All structures are plastic, and tribes and genera as well as species are difficult to define.

References

- Hampson, Fauna of British India, Moths, **1**, 402, 1892 (first clear recognition of family).
- Aurivillius, D.E.Z. Iris, **7**, 121-185, 1894 (includes first key and analysis of larvae—Palearctic only).
- Dyar, Can. Ent., **30**, 2-6, 1898 (first key including neotropical genera).
- Tutt, British Lep., **2**, 434-3, 229, 1900, 1902 (monograph).
- Forbes, Ann. Ent. Soc. Amer., **3**, 110, 1910 (family definition on larval characters).
- Mosher, Bull. Ill. St. Lab. Nat. Hist., **12** (2) 123, 1916 (pupal characters) (Omitted by Lep. Cat.).
- Forbes, Lep. N. Y., (Cornell Mem. **68**) 679, 1924 (definition of family on imago and larva). (Omitted by Lep. Cat.).
- Aurivillius, Seitz' Macrolep. World, **14**, 205, 1927 (principal subdivisions of family—treated as subfamilies).
- Draudt, Seitz' Macrolep. World, **6**, 565-628, pls. 75-86, 1927-1928 (American species).
- Collier, Lep. Cat., **73**, 1-484, 1936 (full bibliography).
- Biol., **1**, 200-208; **2**, 428-431, pls. 22, 85.

Key to Genera

1. Humeral cell good-sized, rounded-trapezoidal, with the free part of Sc arising from near its apex, which diverges from the discal cell (*Lasiocampini* (fig. 72); eyes large and hairy, palpi long and beak-like; R₃ short-stalked, running to apex or outer margin *Prorifrons*
- Humeral cell small and fusiform, apparently cut out of the basal angle of the discal cell (figs. 71, 74, etc.); Sc and frequently second humeral vein also, arising directly from discal cell (*Malacosomatini*) 2
2. Hind wing with only one humeral vein (figs. 73, 74), which may have minute anterior branches; M₂ slightly separate from M₃ at origin; fore wing with R₂ and R₃ long-stalked, usually much farther than R₅ is stalked with M₁, R₄ strongly stalked with R₅ + M₁; palpi hardly exceeding front 3
- Hind wing with two humeral veins (fig. 71), the second arising at or beyond end of humeral cell and sometimes stalked with Sc; costa strongly arched, M₂ and ₃ generally short-stalked; fore wing with R₂ and ₃ usually forking about opposite the fork of R₅ from M₁; R₄ very short-stalked or free; palpi moderate *Euglyphis*

3. Body with a massive middorsal crest on disc of thorax, basal third or more of abdomen, or both. *Tolype*
 Whole body clothed with smooth fine decumbent hair and hair-scales. *Artace*

Dyar describes *Prorifrons castullux* from the Canal Zone; it is brown with five pale shaded transverse lines (Seitz., 76: e1).

TOLYPE Hübner

(including *Titya* Walker)

This genus falls into three main groups on the nature of the dorsal tufting. In the typical (mostly North American) species the disc of thorax and extreme base of abdomen have large shining spatulate scales, contrasting with the fine hairy vestiture of collar and tegulae; in the *phyllius* group the spatulate scales do not contrast, and are extended further back on the abdomen, also the vestiture of the collar and tegulae is noticeably flattened or spatulate. The residue, including the *Tityas* and many species now standing as *Tolype*, have all the hair fine, but raised in a broad crest almost the whole length of the abdomen. (Venation fig. 73.)

The normal larvae are strongly flattened, with strong sidelappets, and with a contrasting (blue) band across the posterior edge of the thorax (but see *T. synoecura*, below). They are tree feeders, and the cocoon, which is strong and flat, is formed on the bark of the tree.

1. Dorsal vestiture spatulate, very dense but nearly smooth; buff, the fore wing shaded with blackish toward apex, contrasting with a broad pale subterminal area. *nana*
 Dorsal vestiture fine, of hair or narrow flattened scales; apex of fore wing not specially darkened (*Titya*) 2
2. Ground light brown; outer margin strongly scalloped in male, at least with hair-tufts at tips of veins in female. 3
 Ground dark gray, with fine, closely paired ordinary lines, thorax with contrasting white shoulders. *primitiva*
3. The group of four pale pm. lines rather evenly spaced at costa; median area below cell not darkened, or somewhat darkened just beyond the antemedial line only; male with pale cell. *mexicana*
 The second line of the group of four obsolescent, the outermost curving in sharply to costa and fusing with the third; median area below cell much darkened in both sexes, and the veins (especially M_1 and M_3) accented with blackish *synoecura*

TOLYPE NANA Druce

? *Phalaena Bombyx silveria* Cr., Pap. Exot., 4, 134, 359: B, C, 1782.

Echedorus nanus Dr., Biol. Centr. Am., Lep. Het., 1. 212, 22: 16 ♀, 1887.

Also figured: Seitz, 79: e6 ♂, 5 ♀.

The female is rounder-winged and the dorsal half of the median area is contrasting, dark brown, usually enclosing a couple of tawny spots. *T. lemoulti* Schaus from Guiana has been mixed with this, but is larger, more buffy, with blackish apical shade less extensive, though contrasting, and the female lacks the contrasting brown median area. A few males in the National Museum from the Upper Amazons are intermediate and the difference may be only racial. Draudt thinks this may be the *silveria* of Cramer, but the figure of the latter looks to me more like *T. poggei* Schs.

Dec. 5 (Bts.) Ranges north to Mexico.

TOLYPE PRIMITIVA Walker

Poecilocampa primitiva Wlk., List Lep. Ins. Br. Mus., 6, 1476, 1855.

T. taruda Schaus, Proc. U. S. Nat. Mus., 29, 318, 1906.

Figured: Seitz, 80: c5.

Female much larger and rounder-winged than male.

Costa-Lima reports the closely related *T. proxima* as feeding on plum, pear and almost all kinds of acacias (from Mabilde).

Dec. 5 ♂, 27 ♀, July 11 (Scr.); Lancetilla, Honduras, May (Bts.). Ranges to Guiana and the Amazons.

Dyar reports *T. synoecura* Dy. from the Canal Zone, and the N. M. has specimens collected by Zetek. The types were from Mexico. The larvae are social in a hanging, pod-like nest, the cocoons formed in the nest. Larva cylindrical, mixed gray and brown, with a mixture of soft brown and stiff black-tipped hairs, the latter presumably stinging; food *Terminalia buceras*. He also reported *T. mexicana* H. S. from the Zone, but I could not verify the specimen. The larvae suggest these two species should be aberrant Malacosomas, but the pattern, venation and other conventional characters agree with *Tolype*, group *Titya*.

ARTACE Walker

A minor variant of *Tolype*, differing most conspicuously in its white color. (Venation, fig. 74). The larvae are similar to *Tolype*, but in the present species the transverse band is orange.

ARTACE CRIBRARIA Ljungh

Cossus cribraria Ljungh, Vet. Akad. Handl., 1825, 348, 2: C.

Artace punctistriga Wlk., List Lep. Ins. Br. Mus., 6, 1491, 1855.

Also figured: Beut., Bull. Am. Mus. Nat. Hist., 10, 19: 5; Holland, Moth Book, 12: 5; Seitz, 82; 6 ♂, 5 ♀.

Larva: Beut., Bull. Am. Mus. Nat. Hist., 10, 444, 1898; Mabilde, Borboletas do estado do R. G. do Sul, Porto Alegre, 1896; Costa-Lima, Terc. Catalogo dos Ins. que vivem nas plantas do Brasil, Rio Janeiro, 235, 1936.

White with numerous black dots; palpus black and white, unlike *A. rubripalpis*, with which it has sometimes been confused. Larva on wild cherry, oak and peach (U. S.) or Ivy (Brazil).

Jan. 3 (Bts.) Feb. 16, '36 (Wood), Mar. 14, '33 and 19, '36 (A.M. N.H.), June 21, July 9 (Scr.), also reported by Dyar. Ranges from southern U. S. to Brazil.

EUGLYPHIS Hübner

(*Claphe*, *Hydrias*, *Ocha*, *Nesara*)

Palpi much longer than in the preceding genera, beak-like; scaling of fore wing almost always with an under layer of trifid or quadrifid scales, more or less overlaid with simpler ones, some of which may be attached with their length oblique to the axis of the wing, varying much from species to species. Thorax with a small metallic posterior tuft, sometimes buried in the general loose hair. (Venation fig. 71).

Larvae flattened, reported from various plants; Costa Lima notes *deusta* on canella branca (*Psychotria*), *ogenes* on aroeira (*Lithraea*), jasmine, branquillo (*Sebastiania*), canelleira do matto and other plants; *rivulosa* on avocado.

A strictly neotropical genus of some 350 species. General characters are slight, but the species may be grouped by the wing scaling. The following is, I believe the first key ever attempted.

1. Thorax and inner margin of fore wing strongly contrasting with middle part of wing. 2
- Thorax and inner margin of fore wing not contrasting. 3
2. Thorax and inner margin whitish and pale brown, the disc blackish. *thyatira*
- Thorax and inner margin black, the disc brown. *vittabunda*

3. Overscaling, at least out to antemedial line, blunt-ended, spatulate or strap-shaped, decumbent 4
 Overscaling not represented as such, but in the form of many smooth-edged erect scales mixed with the normal long-toothed ones *marginata*
4. Costa of hind wing deeply dentate in male (fig. 75), less so in female; overscaling narrow, strap-shaped, dense in outer part of wing, roughly parallel to direction of veins *modesta*
 Costa of hind wing with a large broad lobe, supported by second humeral vein, scaling normal (with moderately dentate under scales, and broad strap-shaped and narrow-dentate overscales) 5
 Hind wing with costa even, or a little waved in male, not dentate; over-scaling normally oblique, crossing vein A on a strong slant or even transversely, broader and imbricated 6
5. Ante- and postmedial lines pale, even and nearly straight, the other markings obscure *sigurda*
 Ante- and postmedial lines waved and less oblique, also with a double series of whitish st. spots at middle of outer margin *defnita*
6. Pm. as well as st. line deeply waved, the former extended in in long lines on veins, the latter with teeth between veins much longer than distance between veins; brown *maria*
 Pm. line, at least, moderately sinuate and waved, when visible . . 7
7. St. line deeply and regularly scalloped; major part of wing overlaid with obliquely placed hair-scales; general effect greenish *submarginalis*
 St. line when distinct not much more deeply waved than pm.; ground brown or buff 8
8. Antemedial line evenly excurved and not or only slightly waved, double, forming the outer boundary of a usually contrasting dark brown or buff base; outer markings less conspicuous . . . 9
 Antemedial line moderately waved, much like postmedial, the basal area not contrasting 11
9. Median area narrow (1-5 length of wing), varying from cream white to brownish, the double pm. line almost as strong as the am. and cutting it off sharply on outer side *discorica*
 Median area broad, the pm. line weak, so that it gradually shades into the vaguely marked outer area 10

10. Am. line hardly retracted at fold; base black-brown, the inner side of the am. lines defined by a narrow luteous line, and pale patch over fork of Cu when present essentially independent of it.
rivulosa
Am. line sharply retracted in fold, often interrupted on A; preceded by an extensive pale patch which gradually narrows to costa; smaller and paler.....*larunda*
11. Disc of wing heavily shaded with white, giving a rather violet effect (scaling of inner border of narrower scales, though not strikingly so).....*rundala*
White scaling, when present confined to fine markings on ordinary lines and along veins.....12
12. D.d. represented by two separate black raised dots; ground of both wings buff (Nesara).....13
D.d. single, when black; ground darker.....14
13. Special st. spot in cell M_1 of male bright tawny; ground duller, dark costal markings of hind wing extending to M_1 ...*caramina*
Special st. spot in cell M_1 of fore wing deep brown in male; the ground bright straw like most of hind wing; costal dark marks of hind wing extending only down to R.....*casada*
14. Thorax and fore wing a sort of dirty buff, the hind wing pale brown; veins of fore wing finely pale, lines pale and double, with some darker shading defining them.....*lyso*
Thorax and both wings gray or brown, the ordinary lines usually single and dark, though often defined by pale dots on veins.. 15
15. Fore wing overlaid with slender pale yellow scales, giving a frosted effect; the yellow-white on ordinary lines well marked; costal part of hind wing with the same pattern.....16
Fore wing mottled with dull brown, without slender pale scaling; the ordinary lines obscure, dark, or partly dotted with yellow-white; entire hind wing with diffuse markings.....18
16. Pale on veins in the form of separate dots; hind wing with st. markings obscure.....17
Pale on veins continuous or nearly so, often extending in teeth on veins, and the am. and pm. lines typically connected by a pale line on Cu; d.d. obscure; st. dots of hind wing conspicuously pale.....*charax*
17. Darker, the dorsal half of hind wing redder; fore wing with d.d. obscure, slightly darker; st. markings regular....*melancholica*

Paler, the dorsal half of hind wing dull brown; fore wing with conspicuous black d.d. (rarely small, but always contrasting); st. retracted at cell M_2 , and marked by a contrasting blackish spot or shade.....*amathunia*

18. Discal dot obscure, st. a fairly regular series of black dots (though tending to omit M_1); no whitish shades.....*sobrina*

Discal dot conspicuous, sometimes very large; st. dots small, in a whitish shade, tending to be ocellate as far as M_1 , especially in the female, then abruptly larger and with little or no white.....*attenuata*

Group I: *Oversealing of fore wing of two sizes, that toward the inner margin definitely finer than the rest; R_1 slanting up to Sc at an angle of about 30° ; 1st hum. of hind wing long, reaching to about opposite end of cell; hind wing evenly rounded or nearly so* (Euglyphis).

EUGLYPHIS THYATIRA Druce

♂ *Lasiocampa thyatira* Dr., Biol. Centr. Am., Lep. Het., 1, 203, 22: 2, 1887.

♀ *Claphe albiplaga* Schaus, Proc. U. S. Nat. Mus., 29, 301, 1906.

E. t. form *taminata* Draudt in Seitz, Macrolep. World, 6, 601, 83: h1, 1927.

Figured: Seitz, 83: g6 (typical), h1 (*taminata*).

Blackish, with two small and a large apical cream costal patches and three partly fused dorsal ones. Female with spots small, white, blurred; a large elongate ocellate elliptical black discal dot with white center.

Var. *taminata* has a light gray thorax, and the white at middle of costa is a dash 3 times as long as wide. It is perhaps the southern race.

Dec. 27, Mar. 23 (Bts.) Mar. 1, '33 (A.M.N.H.), June 22, July 6, '40 (Scr.). Lancetilla, Honduras (Bts.) to Guiana. Var. *taminata* was described from Colombia.

EUGLYPHIS VITABUNDA Dyar

Claphe vittabunda Dyar, Proc. U. S. Nat. Mus., 47, 222, 1914.

Superficially, this species is extremely close to *C. marna* Schs., from Brazil, (fig. 83), but the genitalia (fig. 82) are of a different type. There is also an undescribed species even closer to *vittabunda* (fig. 84), but it is a little coarser looking, the hind wing much redder, and the genitalia have stouter costal hooks of the valves, the uncus has a smaller median portion and the lateral spinules on the tip of the aed-oegus are vestigial. It is figured in Seitz (83: g2 ♂, 1 ♀) as *marna*, which in fact differs from both in having the pm. line continuous

though somewhat sinuous where the black patch begins. In vitta-bunda and the new species there is a definite break.

Oct.-May (see diagram). Described from the Canal Zone, and not known authentically from elsewhere.

EUGLYPHIS RUNDALA Schaus

Claphe rundala Schs., Proc. U. S. Nat. Mus., 29, 304, 1906.

Figured: Seitz, 83: i2.

Finer scaling of inner margin hardly distinct from the rest. Male with postmedial line on costal half of fore wing represented by two diffuse whitish lines, the inner only moderately excurved, but the outer swinging almost out to the st. Female with this whole region suffused with violet gray. In the closely related *E. laronia* Druce, which is regional, only the inner pm. line is present.

Abundant (see diagram). Guatemala (Nat. Mus.) to Guiana (type) and Middle Amazons (Cornell).

Group II: *Similar, except that the oversealing of the whole wing is a similar mixture of dentate and strap-shaped scales; 1st hum. of hind wing generally shorter, especially in melancholica and charax, but overlapping base of Sc.*

[EUGLYPHIS SUBMARGINALIS Walker

Eriogaster submarginalis Wlk., List Lep. Ins. Br. Mus., 35, 1948, 1866.

Hydrius praxithea Dr., Ann. Mag. Nat. Hist. (6) 13, 181, 1894; Biol., 87: 1.

Also figured: Seitz, 82: e3 ♂, 2 ♀.

Metallic crests coppery, also present on abdomen. The abundant fine yellow hair on a gray base gives an olive effect.

Mexico to Brazil, reported from the Canal Zone by Dyar.

Larva (U. S. Nat. Mus.) of the "lappet type," quite unlike the normal Malacosoma group, but as in Tolyte. Pattern speckly in shades of wood brown, with a tendency to longitudinal striping; side tufts strong, pale; pale transverse dorsal tufts on meso- and metathorax, a thick tuft on 1st segment of abdomen, transverse tufts on 2 and 3 with velvety black patches before them; 4 to 8 each with 3 (dorsal and subdorsal) tufts, the ones on 5 larger, with a dark patch behind them. Cocoon flat, silken, pupa pubescent.]

Dyar also reports *E. maria* Schs., a brown species with pm. as well as st. line very deeply scalloped, and the amount of overscaling much less.

EUGLYPHIS DISCORICA Dyar

Claphe discorica Dyar, Proc. U. S. Nat. Mus., **47**, 222, 1914.

? *Gastropacha obtusa* H.-S., Samml. aussereur. Schm., fig. 471, 1856.

The narrow pale median area with large black discal spot on its inner border is distinctive. The area varies from cream white to pale brown, only a little lighter than the ground.

Nov. 27 (Bts.) Mar. 13 (A.M.N.H.), July 3, 21, 25 (Scr.), males; described from Taboga Id. (Dyar) and Guiana (H.-S.)

EUGLYPHIS RIVULOSA Müschler

Hydrias rivulosa Moch., Verh. k. k. zool.-bot. Ges. Wien, **27**, 675, 10: 39, 1878;
Lep. Cat. **73**, 86, 1936.

Also figured: Seitz, 84: f3 (much too dull).

Early Stages: Jörgensen, D.E.Z., Iris, **46**, 58, 1932.

There are several extremely close species in this group. I have a few specimens from the Guianas, whose genitalia agree with the Barro Colorado ones (fig. 79), but none from Paramaribo. The species from southern Brazil (which I take to be *maha* Schs., fig. 80) has practically the same pattern in the male, but the antemedial line is more deeply dentate at A, the genitalia are distinct, and the basal area in the female is extended far out in the middle. In the present lot the female is much like the male in pattern, though larger and heavier as usual. Both this species and *maha* are dimorphic, the male frequently having an oval cream patch on the brown basal area.

Larva bark-like, pale yellowish gray, with a red longitudinal stripe, blackish side-spots and other faint yellowish stripes; lappet-hairs gray; under side red with black borders. On Lauraceae. (I am not at all sure this is the true *rivulosa*; from the locality it is more likely to be *maha*, but *rivulosa* is doubtless similar).

Common (see diagram). Described from Dutch Guiana, other localities doubtful from confusion with related species.

E. larunda Dr. (Biol. 22: 7, as *laronia*) is regional and will doubtless be taken on the island. It is smaller and more delicately marked than *rivularis*, the female light gray, much like the male; and the male valves lack the tooth at the base of the costal hook (fig. 81).

EUGLYPHIS CHARAX Druce

Hydrias charax Dr., Biol. Centr. Am., Lep. Het., **2**, 433, 86: 20, 1897; *Euglyphis c.* Collier, Lep. Cat., **73**, 72. 1936.

Also figured: Seitz, 83: i6.

The amount of cream-white on the veins varies,—sometimes the lines are not connected below the cell as in the type and present specimen.

Dec. 8 (Bts.) Guatemala to Colombia (Nat. Mus.)

EUGLYPHIS AMATHURIA Druce

Hydrias amathuria Dr., Proc. Zool. Soc. London, 1890, 503; *Euglyphis a.* Collier, Lep. Cat., 73, 69, 1936.

Metallic crest high, continued on much of abdomen; overscaling rather slender, yellowish, abundant; am. and pm. lines marked mostly by the series of yellow dots, but st. with well-marked preceding blackish shades, the subcostal one and one in cell M_1 heavy.

Common, Jan.-Apr. (see diagram). Costa Rica to Ecuador.

Dyar reports *E. melancholica* Btl. from the Canal Zone. It is more olive, without the conspicuous discal dot, and the hind wing is redder. The A.M.N.H. party took it at Chiriqui.

EUGLYPHIS LYSO Druce

Hydrias lyso Dr., Biol. Centr. Am., Lep. Het., 2, 435, 87: 4, 1897.

Claphe inflata Schs., Ann. Mag. Nat. Hist. (8) 7, 371, 1911.

Also figured: Seitz, 84: h4 (*lyso* ♀); 85: i1 (*inflata*)

Terminal lunules white, contrasting; discal dot conspicuous and single in the present male, double in original figure of female; a group of three st. black spots at costa. Female with pattern more blurred and lacking the white terminal lunules.

Apr. 23 (Fried.), July 1 (Scr.). Costa Rica to Brazil.

This is the species standing as *lyso* in the Nat. Mus. and agrees with the type of *inflata*. The original figure of female *lyso* looks different.

Group III: *Identical with group II, except that the costa is broadly lobed over the tip of 2d hum.; 1st hum. relatively short, scaling soft and fine.*

EUGLYPHIS DEFINITA Schaus

Claphe definita Schs., Ann. Mag. Nat. Hist. (8) 6, 563-4, 1910; *Euglyphis d.* Seitz, 617.

Ground dull brown, markings a little blurred, m. area a little paler, except for a very vague longitudinal streak through end of cell. Costal part of hind wing darker, but not otherwise contrasting.

Feb. 3, ♂ (Bts.); Dec. 4 (Bts.), Feb. 17 (Fried.) ♀'s. Described from Costa Rica.

EUGLYPHIS SIGURDA Schaus

Claphe sigurda Schs., Ann. Mag. Nat. Hist. (8) 6, 570, 1910; *Euglyphis* s. Seitz, p. 618.

Color and soft appearance as in the preceding; lines fine, almost white, slightly converging to inner margin, the pm. line acutely curved about between R_3 and R_4 . St. marks barely traceable.

The type is male, the present specimens female, but I have no doubt of the determination.

Dec. 27 (Bts.) Mar. 22 (Fried.) Described from Costa Rica.

Group IV: R_1 still running across to Sc at an angle of about 30° ; scaling and fore wing as before; hind wing with strong basal lobe, supported by a short and heavily bifurcated 1st hum., outer part of costa deeply scalloped, R and M_1 arising widely separated.

EUGLYPHIS MODESTA Druce

Lasiocampa modesta Dr., Biol. Centr. Am., Lep. Het., 1, 203, 1887; *Phyllodesma* ? m. Kby., Cat. Lep. Het., 825, 1892; *Epicnaptera* ? m. Closs, Int. Ent. Zeit., 7, 179, 1915; *Euglyphis* m. Ddt. in Seitz, p. 618. 1928.

The doubtful references to Gastropachine genera are on account of the odd wing-form; the venation (fig. 75) is perfectly normal for *Euglyphis*, except for the development of one of the usually minute anterior branches of 1st hum. into a main fork.

Wood-brown, the ordinary lines blackish and waved, finely and partly defined with white. Median area sometimes contrastingly deep brown, as in *Malacosoma disstria* f. *sylvatica*.

Oct. 8 to Mar. 6 (see diagram). Costa Rica to southern Brazil (Cornell).

Group V: R_1 crossing to Sc at a much steeper angle, about 45° , and immediately becoming extremely close to Sc; hind wing with 1st hum. very short, extending less than half length of cell and barely overlapping base of 2d hum.; Sc arising almost at apex of cell and R long-stalked with M_1 . Scaling with no distinction in position of under and over scales, but with a proportion of scales erect and smooth-edged, without the usual marginal teeth (Nesara in part).

EUGLYPHIS MARGINATA Schaus

Ocha marginata Schs., Proc. Zool. Soc. London, 1894, 239.

Ocha libnites Dr., Biol. Centr. Am., Lep. Het., 2, 436, 87: 8, 1897.

♀ *Claphe falsifica* Dgn., Het. nouv. Am. Sud, 22, 19, 1923.

Claphe jeba Schs., Ann. Mag. Nat. Hist., (8) 6, 573, 1910.

Also figured: Seitz, 85: a7 ♂, 8 ♀, k10 ♀ (*falsifica*).

Brown, the pm. line double, arched far out, almost to st. line, but abruptly retracted in cell M_1 ; st. lunulate, black in a violet gray shade, which is stronger before and beyond, and fills the lunules. In typical *marginata* (*libnites*) there is a little darkening of cell M_1 from discal cell to pm. line and a slight basal dash in fold, but in var. *jeba* the pm. area is deep brown and the basal dash extends almost $1/3$ length of wing. Both forms normally have an oval patch over cells M_2 and M_3 . Female umber brown, rather suffusing all the markings, the hind wing mouse gray. *Falsifica* was based on its darkest condition.

Abundant (see diagram). Mexico to Venezuela.

Group VI: R_1 of fore wing and 1st hum. of hind wing like group V; scaling normal, with fine over-scaling, the vestiture of thorax normally fine and strap-like; Sc arising well back from end of cell and R rarely stalked (Nesara).

Dark species, the thoracic tuft lead-color.

EUGLYPHIS SOBRINA Schaus

Claphe sobrina Schs., Ann. Mag. Nat. Hist., (8) 6, 570, 1910.

Figured: Seitz, 84: i4.

Effect of group II. Dull brown, the overscaling broad and deep giving only a vague mottled effect; crest confined to thorax. Lines blackish, rather punctiform and more or less defined by white dots, the st. plainer and more scalloped.

Dec. 8–Apr. 10 (see diagram). Described from Costa Rica.

EUGLYPHIS ATTENUATA Schaus

Claphe attenuata Schs., Ann. Mag. Nat. Hist., (8) 6, 572, 1910.

Ocha falsa Schs., Proc. Zool. Soc. London, 1894, 238.

Claphe palma Schs., Proc. U. S. Nat. Mus., 29, 316, 1906.

Figured: Seitz, 85: k9 (*attenuata*), 8 (*palma*)

The type form has a dark m. area and extremely large black discal dot, var. *palma* is lighter, warmer brown, with a blackish patch beyond cell, *falsa* is grayer, and the type had a minute discal dot, but others from southern Brazil have the large spot. I think the forms are not local but the grayer tint of southern specimens (*falsa*) may have some meaning.

Common, Oct.–Apr. (see diagram). Costa Rica to southern Brazil.

Pale species, the glossy thoracic tuft buff.

EUGLYPHIS CASADA Schaus

Ocha casada Schs., Ann. Mag. Nat. Hist., (8) 7, 372, 1911; *Nesara c.* Ddt. in Seitz, p. 619.

Figured: Seitz, 85: e7.

Ground ochre, the fore and hind wings similar in color, markings brown, contrasting, the most conspicuous a bar from inner pm. line to st., cut by the pale element of the outer pm.

Mar. 13, 1936 (A.M.N.H.), also Chiriqui. Described from Costa Rica.

Bates took a single female of this group (Dec. 11) which certainly does not belong to *casada*. I suspect it is *E. (N.) caramina*, of which he got males in Honduras.

EUPTEROTIDÆ

Head with tongue absolutely rudimentary, but palpi usually well developed; antennae in our species pectinate in both sexes. Body stout, with deep vestiture, tympanum absent. Wings (figs. 86-89, 92) with Cu apparently trifid (almost quadrifid in a few Old World Eupterotinae); radial system in our species (Apatelodinae) normally with R_2 stalked with $_3$ and R_4 with $_5$, their stems normally more shortly stalked together, occasionally with R_2 and $_3$ completely united; hind wing with Sc rather closely parallel to cell near base, then gradually diverging and connected with cell by a cross vein. Frenulum in our species strong. Pattern complex, the antemedial and postmedial lines each represented by two lines or bands, which may be widely separated and may differ in behavior,—as in various other lower “Bombycine” families, such as the Bombycidae and Thyatiridae. Under side normally with complex and sharply defined pattern; inner margin of hind wing above normally with special markings and displayed at rest.

Egg extremely flat, disc-like; caterpillar as in the Lasiocampidae, with much fine secondary hair and obscure warts; in the *Epia* group with spatulate scales mixed with the hair, and frequently with long pencils. Pupa smooth, with a ring of sculpture on the edge of each abdominal segment; that of the Apatelodes group buried in the ground, unlike the residue of the Bombycoidea.

In appearance there is considerable difference between the Eupterotinae and Apatelodinae, but it is difficult to find any simple formula to separate them; as a rule in the Eupterotinae the frenulum is weaker and in the radial system R_3 and $_4$ are stalked the farthest. In the New

World they are represented only by *Preptos*, with two species in Central America. The larvae are also closely similar, and quite unlike the nearly naked, humped or horned larva of the Bombycidae. Only *Colla* is reported to have a Bombycid-like larva.¹ Even the row of fine dorsal hair-pencils of *Apatelodes* reappears in *Preptos*, though they are absent in the Old World Eupterotinae.

The Apatelodinae are generally treated as limited to the New World, but "*Andraca*" *albilunata*, from eastern Asia, is a normal Apatelodine, perhaps even a species of *Apatelodes*, and *Prismosticta* and *Panacela* show some Apatelodine features. Of these three types only the larva of *Panacela* is known,—it is hairy, unlike the Bombycidae.

Aside from the general references listed in the introduction, there has never been a revision of the subfamily.

Key to Genera (after Schaus in Seitz)

1. R_4 of fore wing convex and closely parallel to costa, R_2 $_3$ running parallel to them, generally a simple vein (fig. 86).....2
 - R_4 of fore wing sinuous, R_2 and $_3$ generally forking immediately after their separation from R_4 and $_5$ (fig. 92).....6
2. Apex of fore wing rounded.....3
 - Apex of fore wing marked.....4
3. Cu_1 of fore wing arising well before angle of cell; underscaling normal.....*Epia*
 - Cu_1 of fore wing close to angle of cell or connate with M_3 ; under scales with very slender bases bearing about 4 long spikes, like some species of *Euglyphis*.....*Colla*
4. Apex produced (fig. 87).....*Quentalia*
 - Apex merely right-angled (fig. 86).....5
5. Abdomen not extending beyond hind wing and without tufts.
 - Anticla*
 - Abdomen extending beyond hind wing, and with tufts on both.
 - Tamphana*
6. Apex acute; outer margin almost evenly rounded.....*Zanola*
 - Apex usually a little blunted, outer margin sharply bent or angled just below vein M_17
7. Outer margin of fore wing scalloped on dorsal half, in particular with a distinct tooth at M_3*Olceclostera*
 - Outer margin even from M_2 to anal angle.....*Apatelodes*

¹ Schade, Ent. Rundsch., 56, 65-67, 1939.

EPIA Hübner

(*Anthrocroca* Boisduval, *Hygrochroa* Felder et al., not Hübner¹)

Wing-vestiture deep and soft, the under scaling only moderately toothed; colors darker, buff, brown or olive, (venation fig. 77).

1. Male with an olive costal triangle extending down to vein M_2 and contrasting with the brown general ground. *muscosa*
 Male with the triangle when distinct extending down across M_3 ; the ground olive to bright green and not contrasting. . . . *casnonia*

EPIA MUSCOSA Butler

♀ *Anthrocroca muscosa* Btl., Trans. Ent. Soc. London, 1878, 79, 3: 5; Biol., 1, 224; Kirby, Cat. Lep. Het., 721, 1892.

♂ *A. cuneifera* Btl., Trans. Ent. Soc. London, 1878, 79, 3: 4; Biol., 1, 224 (pr. ♂).

Also figured: Seitz, 89: a2 ♂, 1 ♀.

Male with ground two shades of warm brown; an olive costal triangle, shades near margin, and dark brown pm. line most conspicuous. Female strikingly different, the olive markings confused and hind wing much redder than fore wing.

Common; flight scattering, after mid-December (see diagram). Guatemala to southern Brazil.

Epia casnonia Druce ranges from Guatemala to Peru. Northern males are dull olive with violet-gray shades, and the costal triangle rather distinctly set off; southern ones are almost evenly olive green (very bright when fresh), and the median area has fine widely separated dark lines not enclosing a triangle; the female is smaller but not easily distinguished from *E. muscosa*. Both forms were taken by Bates at Lancetilla, Honduras, and will doubtless occur on Barro Colorado Id.

COLLA WALKER (*Prismoptera* Butler)

This genus has a remarkable superficial likeness to the old world Bombycid genus *Ocinara*, which has a good deal to do with the tendency to combine the *Apatelodinae* and *Bombycidae*. In fact venation (fig. 78) and scaling are quite different, and the likeness is certainly parallelism. The ground is generally white, shaded with light gray, and so thinly scaled as to expose the iridescent membrane. Under-scaling reduced to branched hairs.

¹ *Hygrochroa* Hübner was based on a true *Apatelodes* and a European geometer; the name is now used for the geometer.

Larva of *C. jehlii* Schade nearly naked, humped with horns on 2d and 8th abdominal segments; on Ficus (Schade, l.c.)

Pupa of *C. rhodope* Dru. according to Druce dark brown, enclosed in a silvery gray silky cocoon.

COLLA COELESTIS Schaus

C. coelestis Schs., Ann. Mag. Nat. Hist., (8) 6, 414, 1910.

Figured: Seitz, 89: b8.

Small and noticeably round-winged, the wings rather strongly marked with light gray; thorax white, the posterior tuft contrasting, gray.

Jan. 27 (Bts.), June 24 (Fried.). Described from Costa Rica.

QUENTALIA Schaus

(*Carthara* auct., not Walker)

Similar to *Apatelodes*, with the same marked apex and bent or angled outer margin, but with the stem of R_2 and $_3$ closely parallel to R_4 for some distance beyond the forking of R_5 , and both convex to costa (fig. 87); frenulum short but functional.

A good sized genus, but with the species characters not well understood. The few species yet known from the Island however are clearly defined.

1. Discal lunule of upper side of fore wing a narrow oblique bar of curly raised scales, about as high as the cell; outer pm. line sinuous and excurved over M_3 ; discal lunule of hind wing below in our species with white scales. *veca*
 Discal lunule of upper side of fore wing represented by two widely separated black dots; outer line not convex over M_2 , normally concave to just above M_3 ; discal lunule of hind wing below dark . . 2
2. Base of abdomen with a contrasting transverse dark (brown or olive) band; outer pm. evenly concave from costa to M_3 , forming the outer boundary of a triangular dark patch in both sexes. . *numalia*
 Base of abdomen with two small brown tufts only; marking much less definite in male, the outer pm. whitish, obscure and nearly straight above, though scalloped below M_3 ; female with blurred markings. *ephonia*

QUENTALIA VECA Druce

Carthara veca Dr., Biol. Centr. Am., Lep. Het., 1, 224, 23: 21 ♂, 22 ♀, 1887.

Q. chromana Schs. in Seitz, Macrolep. World, 6, 690, 89: 11 ♂, 2 ♀, 1929.

Carthara reissi Mssn. in Stübel, Reisen in Sudamerika, Lep., 132, 5: 6, 1890.

Also figured: Seitz, 89: h9 ♂.

Fore wing squarer than in the other species, the outer margin nearly vertical from apex to middle angle, especially in male; pm. line double, the inner line dark and waved, the outer widely separated at costa, closest opposite cell, dark followed by pale, concave toward costa and waved below,—the am. elements similar in reverse order. Ground color variable, typically deep indian red, shaded with red-brown, in var. *chromana* Schs., deep olive brown, frequently frosted with white, in the var. which the Nat. Mus. takes to be *reissi*¹ with a contrasting olive wedge between M_3 and Cu_1 , becoming diffuse to margin.

The majority of the present lot are of v. *chromana*, with a fair block of the type coloring, one *reissi* and several transitional specimens, including several with the *reissi* wedge on the *chromana* ground. The single female is of v. *chromana*.

Common, with principal flight in Nov.—Dec. (see diagram). Ranges to Costa Rica and Ecuador.

QUENTALIA NUMALIA Schaus and Heinrich

Q. numalia S. & H. in Seitz' Macrolep. World, 6, 684, 89; f1, 1929.

The dark costal triangle is the upper half (down to M_3) of the double pm. line with the space between, filled with gray in its upper half and below with brown.

In this group there are many neotropical species absolutely identical in markings but with strikingly distinct genitalia. Names will be a little uncertain till types of the older names have been examined, but we figure a few that we believe are correctly named (figs. 90 *numalia*, 91 *surynorta*, 85 an undescribed species common in southern Brazil). The last sternite is easily uncovered by denuding and is distinctive of each species,—that of *numalia* has two long asymmetrical terminal spikes, its ninth tergite has two pairs of marginal points, about like *surynorta* from farther north, and the lower half of the valve simply tapering and short, with terminal bristles (subterminal at a sharp bend in *surynorta*).

¹ The original figure is paler than any specimens I have seen of *veca*, a common fault in hand-coloring.

Jan. 9, Feb. 6 (Fried.) Chiriqui (A.M.N.H.). Costa Rica (type and C.U.)

This species would have been included in the *Biologia* conception of *drepanoides* (p. 225), but they had no material from Panama. The variation in color that they mention probably occurs in each of the species, and is striking in the series of *surymorta* collected by Bates in Honduras.

QUENTALIA EPHONIA Stoll

Phalaena Bombyx ephonia Stoll in Cr., Suppl. Pap. Exot., 96, 19:5 (larva and pupa); 159, 35: 6, 1791.

Carthara lividia Dr., Biol. Centr. Am., Lep. Het., 1, 225, 23: 23 ♂, 24 ♀, 1887. Also figured: Seitz, 89: g6 ♂, 7 ♀.

Biology; Stoll, l.c. 96.

Male variable in ground color, the present specimen showing much more red tint than Druce's figure. The original figure of the female does not show the subapical reddish or olive patch; but agrees in the course of the lines; I believe it can only be this species.

Dec. 25 to Jan. 25, Mar. 4 (Bts. & Fried.) July-Aug. (Fairchild). Mexico to Peru and Amazons.

Larva somewhat flattened, obviously of the general type of *Olcclostera angelica*, but with clusters of long spatulate scales on thorax and penultimate segments; gray with orange head, legs and lateral stripe. Food yellow jasmine.

ANTICLA Walker

Palpi short; frenulum a little longer than in *Quentalia*; fore wing with outer margin quite even; apex marked but rounded over; sexes strikingly dimorphic. (Venation fig. 86.)

ANTICLA ANTICLA Walker

A. anticla Wlk., List. Lep. Ins. Br. Mus., 5, 1174, 1855.

A. carya Dr., Biol. Centr.-Am., Lep. Het., 1, 226, 23: 27, 1887.

Hygrochroa rutila Dr., Biol. Centr.-Am., Lep. Het., 1, 224, 1887.

Anthrocroca amycla Dr., Proc. Zool. Soc. London, 1890, 507.

Also figured: Seitz, 89: i7 ♂, 6 ♀.

Light gray-brown, the lines waved and the lines of the pairs representing the am. and pm. so widely separated as to make five almost evenly spaced lines,—lines often obsolete; outer pm. followed by two green patches at costa (one large one in the closely related *A. flavaria*),

and with some green dorsally. Hind wing somewhat redder; female with green and light brown in broad shades and almost any proportion; the hind wing bright yellow, contrasting, often with two dark outer lines.

Common (see diagram). Ranges to the Amazons at least, two specimens apparently of this from southern Brazil being both females.

TAMPHANA Schaus

Outer margin a little more oblique and sinuous; inner margin of hind wing with a large tuft a little beyond middle, projecting vertically up in resting position, much like that on the fore wing of *Apela* (Notodontidae). (Venation fig. 88.)

TAMPHANA MARMOREA Schaus

T. marmorea Schs., Proc. Zool. Soc. London, 1892, 325.

Figured: Seitz, 89: k8.

Male genitalia: Forbes, Ent. News, 52, 79, 1941 (with figure).

A complicated pattern in shades of brown, with a large triangular pale costo-apical patch and smaller rounded st. patch below M_3 . Genitalia with valves not functional as such, apparently modified into a stridulating organ.

Common (see diagram); also Amazons and S. Brazil. We have not seen the female.

ZANOLA Walker

Similar to *Apatelodes*, but the outer margin of fore wing hardly angled at M_1 , costa of hind wing slightly concave on outer part and slightly lobed at apex. Though the venation of this genus agrees with *Apatelodes*, the larva and genitalia are more like *Quentalia*.

ZANOLA VERAGO Cramer

Phalaena Bombyx verago Cr., Pap. Exot., 2, 102, 162: D, E, 1779.

Zanola difficilis Wlk., List Lep. Ins. Br. Mus., 5, 1174, 1855.

Z. harpis Dr., Biol. Centr. Am., Lep. Het., 1, 222, 23:18, 1886.

Apatelodes vivax H. Edw., Pap., 4, 77, 1884; Biol., 208.

Also figured: Seitz, 141: k3 ♂, 142: c3 ♀.

Male rusty orange with brown shadings and rather fine lines, the terminal space more yellow, also a st. lunule in M_2 and much of the base yellow. Female mostly light yellow.

Larva (U. S. Nat. Mus.) much like *Malacosoma* in build; body with broken longitudinal striping of cream, tawny, gray and black, hair loose and fine, tending to curve back, abundant but not long; the spatulate scales very long, and with conspicuous black ends,—mixed with the subdorsal tufts on thoracic and 8th abdominal segments,—twice as long as the ordinary hairs, except on prothorax. Pupa glossy, with the segmental lines of beading as in *Apatelodes*; cocoon of rough brown silk.

Nov. 22 (Bts.) Mexico to Amazons.

The larva of the closely similar *Tarchon felderi* Dr. in the U. S. Nat. Mus. has dense matted silky light brown hair, leaning back except on the prothorax where it leans forward about the head; dorsum of meso- and metathorax with dense masses of black spatulate scales, the abdominal segments with white subdorsal ones, tending to lie in longitudinal lines and mixed with shorter black ones; the 7th and 8th segments with mouse gray dorsal ones, when fresh forming a loose posterior tuft. Cocoon largely of the matted hair, decorated with the white spatulate scales.

APATELODES Packard

In this and the following genus (figs. 92 and 89) the branches R_{2-5} usually fork rather far out, and normally R_{2-3} and R_{4+5} only separate shortly before they bifurcate; the stem of R_{2+3} diverges evenly from R_5 , not being convex to the costa; outer margin usually bent or angled above M_2 , sometimes merely strongly curved over M_2 , concave above the bend, but either straight or a little convex or concave below. The abdomen lacks the white lateral dots usual in *Olceclostera*.

Egg flat, disc-like; larva with fine dense secondary hair, conspicuous in color, with a series of fine middorsal pencils, mostly short, but in *A. torrefacta* with two long anterior and one posterior one much lengthened. Pupa in ground (unlike the few known *Zanolini* and other *Eupterotidae*), shining, with the edges of the movable abdominal segments beaded.

The pattern is usually distinctive, the inner and outer elements of the am. and pm. lines widely separated, the outer pm. usually conspicuous, the st. usually conspicuous at costa, frequently running into the outer pm. below a transparent dot, and in that case obscure below the point of contact; usually with contrasting dark spots or shades before 1st am. at inner margin and beyond outer pm. at costa.

1. Outer pm. line even; outer margin even, sometimes concave. 2
Outer pm. line scalloped, the margin very faintly wavy. 3
2. Outer pm. continuous from costa to inner margin, evenly excurved;
ground gray with brown lines. *firminiana*
Outer pm. line straight below level of the cell, continuing the course
of st. toward costa, pale on a brown ground. *lapitha*
3. Ground straw; smaller, outer margin concave. *tuisa*
Ground gray, larger, outer margin slightly convex. *adrastia*

APATELODES FIRMIANA Stoll

Phalaena Geometra firminiana (*firminiana* in index) Stoll in Cr., Pap. Exot., 4, 234, 398: F, G, 1781.

Also figured: Seitz, 140: d4.

Light gray; median space and apical area darker; inner am. line with upper part running out in a dark line on Cu₂, lower part separate, vertical, just beyond the large brown inner marginal patch; outer am. line angled at fold and convex; outer pm. much more excurved than inner, st. slight, pale, without a brown spot before it. No transparent spot.

Oct. 30–Dec. 26; May 20; July 15, 25 (Scr.) Ranges to south Brazil.

APATELODES LAPITHA Druce

A. lapitha Dr., Ann. Mag. Nat. Hist., (7) 5, 511, 1900.

Figured: Seitz, 140: f4.

Brown, shading into darker at outer margin except apex, and outer half of inner margin; subbasal brown patch strong, apical a minute triangle well below costa. Inner and outer am. lines parallel, evenly excurved; inner pm. parallel to them, but twice as far apart.

Not rare (see diagram), but males only taken. Mexico to Guiana (Cornell).

APATELODES TUISA Schaus

Zanola tuisa Schs., Ann. Mag. Nat. Hist., (8) 6, 112, 1910.

Figured: Seitz, 141: c4.

Ochre, lightly dusted and shaded with brown, an ocellate white discal dot with brown center; outer pm. line most distinct; transparent spot small, with a brown dot below it; fringe brown, contrasting,—the whole width below apex, and outer edge the rest of the way.

The closely related *A. cinnomoides* from South America is a little smaller and brighter, the discal dot less ocellate, and the pm. line usually preceded by a separate series of brown dots.

Not rare (see diagram), Oct. 22–Mar. 8. Ranges north to Guatemala.

APATELODES ADRASTIA Druce

A. adrastia Dr., Biol. Centr. Am., Lep. Het., 1, 208, 22: 8, 1887.

A. diffidens H. Edw., Ent. Am., 2, 13, 1886.

A. a. costaricensis Ddt. in Seitz, Macrolep. World, 6, 693, 1929.

Also figured: Seitz, 140: a3 (as *adrastia*).

Gray, somewhat powdery, lines rather irregular, an oblique dark shade starting on am. line at costa, reaching outer pm. line at fold and accompanying it to inner margin (rarely absent or indicated only by the dark Cu_2) transparent dot outlined with blackish, especially before. There is a little variation, but I cannot see any reason for varietal names.

Larva (U.S.Nat.Mus.) yellow with fine cream to lemon yellow hair but no pencils; body with a quadrate subdorsal black spot before middle of each of the segments A2–7 and a trace on A1. Pupa normal, glossy.

Dec. 4, Jan. 28. Ranges to Mexico.

OLCECLOSTERA Butler

(*Apatelodes* in part; *Parathyris* Walker, not Hübner)

Moth identical with *Apatelodes* except for the more wavy dorsal half of outer margin; transparent spot usually larger, frequently double; abdomen normally with two conspicuous white lateral spots, sometimes defined with red-brown.

Larva totally different; flattened like a lappet-caterpillar, with short, coarse secondary hair, the dorsal pencils all very short, and with dense discolored hair on dorsum of thorax. The larva rests on the trunk of a tree. The North American "*Apatelodes*" *angelica* belongs to this genus, and is the only one whose life history is known.

1. Most of course of ante- and postmedial lines even and nearly straight; most of segments of abdomen with subventral white points, besides indications of the usual lateral ones. *irrorata*
Postmedial lines regularly waved or obsolete; only the two basal lateral spots on abdomen distinct, the subventral ones very weak or absent 2

2. First two abdominal segments with reddish lateral patches finely edged behind with white; transparent spot minute; a blackish antemedial dot on inner margin.....*nigripuncta*
The lateral spots on abdomen pure white, the first two segments otherwise unicolorous; transparent spot larger, quadrate, no antemedial black dot.....3
3. Darker, browner, male with base to antemedial line blackish, somewhat contrasting; transparent spot very large.....*reperta*
Light gray with a pinkish tinge, the base as light as any part.
amoria

OLCECLOSTERA AMORIA Druce

Oecclostera (sic) *amoria* Dr., Proc. Zool. Soc. London, 1890, 505.
Figured: Biol., 87: 22; Seitz, 141: h6.

Druce in the *Biologia* considered *O. maya* Schs. a synonym; it is now recognized as distinct, and has a blunter fore wing.

Oct. 31-June 21 (Bts. Fried. Scr.) also Lancetilla, Honduras (Bts.).
Ranges south to southern Brazil (Cornell).

OLCECLOSTERA NIGRIPUNCTA Schaus

O. nigripuncta Schs., Ann. Mag. Nat. Hist., (8) 6, 411, 1910.
Figured: Seitz, 141: g7.

This species is very close to the North American *O. angelica*, though the latter shows more the coloring of *amoria*, and has only traces of the lateral white spots. The twice waved inner pm. line is a little emphasized, especially toward costa.

Nov. 27, Dec. 2 (Bts.). Ranges to Guatemala (Nat. Mus.).

OLCECLOSTERA REPERTA Walker

Parathyris reperta Wlk., List Lep. Ins. Br. Mus., 32, 438, 1865.
Figured: Seitz, 141: h3.

Besides the dark base the inner pm. line is shaded with blackish; transparent spot larger and less notched on outer side than in the closely related *O. microps* from South America.

Female larger, the blackish shades very strong before am. and beyond pm. lines.

Common in Oct., scattering to Jan. and in midsummer (see diagram). Ranges to Brazil.

OLCECLOSTERA IRRORATA Butler

O. irrorata Btl., Trans. Ent. Soc. London, 1878, 70.

Figured: Seitz, 141: f2.

Dark brown, dusted and shaded with gray, with a brown antemedial costal shade and the apical area broadly brown; am. line dominantly pale, inner pm. line dark, outer pm. line dark, followed by pale, angled opposite the small transparent dot and waved to inner margin. Lateral abdominal spots obscure, appearing as white posterior edges of several segments.

This is the genotype of *Olceclostera*, but is a little abnormal in appearance.

Commonest in Nov.-Dec. (see diagram). Guiana and Amazons.

URANIIDAE

Head strong, with well developed palpi and mouth-parts. Body small, the moderately slender but strong legs with strong spurs. Wings ample (fig. 76), Cu apparently 3-fid, R_5 stalked with M_1 , separate from the other radials; hind wing with strong humeral angle and no frenulum, the present genus also without frenulum-base; Sc and R fused for a very short distance and then abruptly diverging; outer margin usually angled or tailed at M_3 ; inner margin narrow, with 3d A obsolescent. Auditory organ near base of abdomen, subdorsally, different in the sexes, conspicuous in the present genus.

Larva slender, with irregular secondary hair; usually brightly marked.

A small family in the tropics of both hemispheres and warmer temperate part of the Old World. The Uraniinae, which do not have even a thickening of the base of hind wing, have a curiously interrupted distribution, with closely related genera in the Neotropical Region and Madagascar, with the adjacent edge of Africa, and a couple of more distinct genera in the Indo-Australian Region. The other (*Micronia*) group are strictly Old-World. But the following *Epiplemidæ* hardly make a distinct family.

References

Guenée, Hist. Nat. Lep. Het., 9, 1-15, with pl. 1 (the Uraniinae only as *Cydimonidae*, *Uranidae*, *Nyctalemonidae*); 10, 21 (the *Microniinae* as part of the *Micronidae*).

Dalla Torre, Lep. Cat., 30 (2) 1-9 (Uraniinae).

Biol., 2, 3-4; pl. 41; Seitz, 6, 829-830, pl. 138.

Biology: MacLeay, Trans. Zool. Soc., **1**, 180, pl. 1, 1834; Gosse, Ent., **13**, 133-135, 1880; **14**, 241-245, 1881. For references to biology of Old World genera see Dalla Torre, l.c. under *Chrysidia croesus* and *madagascariensis*, *Nyctalemon patroclus*; also Hampson, Fauna of Br. India, Moths, **3**, 110, fig. 56; Lavauden, V^e Cong. Int. d'Ent., 421, 1933.

URANIA Fabricius

(*Cydimon* Dalman, *Uranidia* Westwood, *Leilus* Swainson)

Antennae slightly swollen and fusiform; hind wing with a single long tail. Coloring brilliant, black and green, sometimes with copper.

Larva cylindrical, black with white markings, the head, cervical shield and true legs reddish, and prolegs white; with some of the sparse setae clubbed. Cocoon covered with frass and bits of leaf; pupa brown, rounded, with well developed cremaster. Food Euphorbiaceae, *U. sloanus* on *Omphalea triandra* L. in Jamaica (Gosse).

URANIA FULGENS Walker

U. fulgens Wlk., List Lep. Ins. Br. Mus., **1**, 5, 1854.

Figured: Biol., 41: 16; Seitz, 138: b1.

Male green, female usually brassy; fore wing with about four green stripes before the broader median band and usually none beyond. Tail mostly black, unlike the Amazonian *U. leilus*.

The moth flies in great swarms, like others of the genus.

July-Aug. (Fairchild). Mexico to Colombia.

EPIPLEMIDÆ

Structurally distinct from the Uraniidae only in the fully developed frenulum. R_5 is occasionally only approximate to M_1 (Gathynia). Normally small and inconspicuous moths, contrasting in the extreme with *Urania*, frequently mistaken for Geometridae. Venation and wing form frequently sexually dimorphic; wing outline most often irregular, but never with a long tail at M_3 . Costa of hind wing normally bisinuate, and often with two scale-tufts; normally with large eyes and narrow front.

Larva inconspicuous, when young somewhat social in a web, later becoming cryptically colored and hiding singly. Setae single, iv and v rather close together, on the same level below spiracle; prolegs with biordinal hooks in a sharply curved band; pupa obtect, in a cocoon. The foods are various, Caprifolicaceae, Rubiaceae etc.

The family is not rare and wide-spread in both hemispheres, but is absent by exception from Europe. It takes an intense oriented light to bring out the iridescence of some species.

References

- Hampson: Fauna of British India, Moths, **2**, 9, 1892, iii, 121, 1895. (Family founded and defined, and key to oriental genera).
 Guenée: Spec. Gen. Lep. Het., **10**, 21-42 (as Micronidae).
 Hulst, Trans. Amer. Ent. Soc., **23**, 309-311, 1896 (as Strophidiinae).
 Sharp, Cambridge Nat. Hist., Ins., **2**, 368, 420, 1899.
 Janse, Moths of South Africa, **1**, 91-115, 1932 (with key to African genera).
 Biol., **2**, 123-128, 540, pls. 53, 99; Seitz, **6**, 1141-1170, pls. 170-172.
 Larva: Hampson l.c. fig. 66; Dyar, Proc. Ent. Soc. Wash., **4**, 414; **5**, 131.
 Dalla Torre, Lep. Cat., **30** (1), 1-38, 1924.

There has been no world review of the family, and the following key follows the general lines of Hampson's in the "Fauna of British India."

Key to Genera

1. Anal angle of hind wing deeply emarginate below Cu_2 , the other marginal irregularities weaker. Fore wing with R_5 and M_1 stalked, M_2 approximate; hind wing with M_2 widely separated 2
 Notch at anal angle when present sometimes broader than the ones preceding it but not deeper; general course of dorsal margin normally straight or convex from M_3 or Cu_1 to 2d A. . 3
2. Palpi short, the third segment tapering and less than three times as long as thick; front more than a third as wide as eye

Syngriodes

 Palpi moderate, the third segment cylindrical and more than four times as long as thick; front extremely narrow, less than $\frac{1}{4}$ as wide as eye. (Fig. 93) *Syngria*
3. R_2 of fore wing stalked with R_{3+4} ; middle of outer margin lobed. . 4
 R_2 of fore wing stalked with R_1 , R_3 and $_4$ separately stalked; M_2 arising widely separated from M_1 at least in fore wing; male with dorsum of hind wing modified 5
 R_2 arising free though closely parallel to R_1 and stem of R_3 and $_4$; M_2 normally arising near upper angle of cell in both wings. . 6
4. Hind wing a squarish lobe supported by both M_3 and Cu_1 , which arise from the cell close together (Fig. 95) *Skaphion*
 Hind wing with a more triangular lobe, supported by M_3 only; Cu_1 arising from cell well before M_3 *Bicavernosa*

- Hind wing right-angled at M_3 and scalloped below, M_3 and Cu_1 short-stalked. *Tricolpia*
5. Fore wing with R_5 and M_1 separate though approximate; both wings of female and fore wings of male with M_2 arising widely separated from M_1 (Fig. 100) *Gathynia*
Fore wing with R_5 and M_1 distinctly stalked, often strongly. *Antiplecta*
6. Hind wing squarish or at least with the outer margin sharply bent at M_3 or Cu_1 and the border straight from there to anal angle. 7
Hind wing rounded in outline or extended at apex, the teeth when present supported by single veins R or M_1 and sometimes M_3 , but not Cu_1 9
7. M_3 and Cu_1 of hind wing together bearing a squarish lobe which is specially marked, the margin nearly straight above as well as below it (Fig. 94) 8
Only a single tooth on either M_3 or Cu_1 , the border straight in general course to anal angle, but usually convex toward costa. *Schidax*
8. Male antennae unipectinate *Psamathia*
Male antennae simple, prismatic, about like female *Nedusia*
9. Both wings with M_2 arising from cell close to origin of M_1 at upper angle (Figs. 96, 97) *Erosia*
Hind wing with M_2 arising far below upper angle of cell 10
10. Male with a large lobe at anal angle of hind wing (in the present species covering dorsum of abdomen), with sex-scales and hair-pencil *Thysanocraspeda*
Male with inner margin of hind wing not distorted (Fig. 98). *Epilema*

SYNGRIA Guenée

Eyes enormous, four times as wide as the front in female and even wider in male. Male (fig. 93) with apex of fore wing marked, the outer margin mostly concave below it, hind wing with squarish anal notch; $Sc+R$ swollen at base, with the free parts of Sc and R arising from the abrupt end of the swelling, a little separated from M_1 , which continues the upper side of the cell; female with apex falcate, and outer margin convex except just below the apex; hind wing with the notch at anal angle deep and with a lobe at the tip of Cu_2 beyond it; venation normal, with R and M_1 stalked.

1. Larger; fore wing with am. line strongly oblique from its bend in upper part of cell to inner margin, cutting stem of Cu 5/6 way out to origin of Cu₂; pm. line only slightly bent at M₃. . . . *drepanata*
 Smaller, fore wing with am. line erect below the bend in cell, cutting Cu about 2/3 way out to origin of Cu₂; pm. line sharply angled or even offset at M₃ and strongly concave to inner margin.
druidaria

SYNGRIA DREPANATA Felder

S. drepanata Fld., Reise Novara, Lep., 128: 35, 35a, 1875; Biol., 125, 1892.

In spite of the remark in Seitz, this species is only known from the male and is a distinct species; the contrasting dark thorax and slender male abdomen are plainly shown in the original figure. It is not strongly variable, rather smooth-looking above, buff with some olive gray tint; below with the black not very extensive, but strongly contrasting on a pale buff ground.

Flight scattering, only males taken (see diagram). Chiriqui (Biol.); Amazons (type).

SYNGRIA DRUIDARIA Guenée

♂ *S. druidaria* Gn., Sp. Gen. Lep. Het., 10, 32, 16:1, 1852.

♀ *S. falcinaria* Gn. Sp. Gen. Lep. Het., 10, 33, 1852; Biol., 2, 124, 1892 (pr. syn.).

S. d. var. *derwaria* Obt, Et. Lep., 20, 226, 552: 4680, 1923.

Highly variable, the female more often dark than the male. Male specimens may be divided into the normal form, with ground powdery luteous and a large white patch before the anal angle, and var. *derwaria* with two small white preanal spots on a fuscous ground. Females are usually intermediate in color, with the two small white spots or none.

Common (see diagram). Ranges from Mexico to Brazil.

SYNGRIODES Warren

Similar to *Syngria*, smaller, the outer margin angled near middle in male and sometimes in female; R₂ sometimes from the stalk of R₃₊₄.

Material is inadequate and descriptions in this genus are desperate, so that I cannot guarantee the following determination.

SYNGRIODES INCISARIA Walker

Erosia incisaria Wlk., List Lep. Ins. Br. Mus., **23**, 842, 1861; *Syngriodes i.* Warr., Nov. Zool., **12**, 309, 1905; as genotype of *Syngriodes* Gaede in Seitz, **6**, 1149, 1936; *Epiplema i.*, Dalla Torre, Lep. Cat., **30** (1) 17, 1924. Also figured: Biol., **53**: 3 ♂?, 4 ♀; Seitz, **170**: g2 ♂, 3 ♀?

Clay color, lightly marked with fuscous; fore wing with hooked apex and strongly angled outer margin in both sexes (?). Druce figures the male with merely sinuous outer margin and Seitz the female, so there must be serious confusion. Our specimen is a female.

Feb. 10, 1936 (A.M.N.H.). Reported from Mexico to Brazil.

SCHIDAX Hübner

Fore wing with M_3 and Cu_1 connate, outer margin convex or bent at M_1 or Cu_1 ; hind wing evenly dentate on convex costal half of outer margin, then angled at M_3 or Cu_1 and straight and scalloped to anal angle. Male antennae unipectinate, female laminate so far as examined.

A small genus, more varied in wing form and appearance than in deeper structures, related to *Syngria*.

SCHIDAX SQUAMMARIA Hübner

S. squammaria Hbn., Zutr. Samml. Exot. Schm., **1**, 27, figs. 161–162, 1818; Biol., **2**, 128, 1892.

S. squammularia Hbn., Verz. bek. Schm., 315, 1825; and of authors generally. Also figured: Seitz, **170**: e3.

Ground gray, variable in tint, typically very pale and powdery with dorsal part of hind wing brown.

Dec. 12 (fresh), Jan. 4 (rubbed–Bates). Nicaragua and Cuba to South Brazil.

PSAMATHIA Walker

General characters and venation as in *Syngria* etc., but with M_3 of fore wing somewhat separate from Cu_1 . Fore wing with acute subfalcate apex and sometimes an angle at M_3 , otherwise with even margin; hind wing with bluntly angled apex and the lobe at middle of outer margin only. Venation alike in both sexes, male antennae unipectinate, female prismatic.

A small genus, hardly distinct from *Nedusia*. *P. placidaria* Wlk. is identical in pattern with the following, but has the outer margin of fore wing almost straight.

PSAMATHIA AMPLATA Warren

P. amplata Warr., Nov. Zool., **14**, 195, 1907.

Figured: Seitz, 170: b1, 2.

Gray with dark markings; pm. of both wings looped far out at middle; sometimes accented with black spots toward inner margin of hind wing, or shaded with fuscous. The present specimens are all females, those taken in Chiriqui in Feb., 1936, by Lutz & Gertsch, all males.

Dec. 2, Jan. 27, Feb. 2 (Bts.). Chiriqui to Paraguay.

I suspect that *subangulata* Warr. (Nov. Zool. vii, 123) described from St. Vincent, is merely a variant of this species.

NEDUSIA Hübner

Differs from *Psamathia* only in the simple laminate antenna of male as well as female, and should probably be combined with it. The stalking of R and M₁ of hind wing, given as a second difference by Gaede in Seitz, actually varies in both genera. (Venation fig. 94.)

NEDUSIA MUTILARIA Hübner

Nedusia mutilaria Hbn., Zutr. Samml. Exot. Schm., 30, figs. 181, 182, 1818; Verz. bek. Schm., 291, 1823; Biol., 2, 123, 1892.

N. cuticulata Gn., Sp. Gen. Lep. Het., 10, 31, 12: 5, 1852; Biol., l.c.; Seitz, Macrolep. World, 6, 1143, 1936 (pr. var.).

Erosia obliteraria Wlk., List Lep. Ins. Br. Mus., 23, 845, 1861; Seitz, l.c. (pr. syn.).

Also figured: Seitz, 170: a2 (not 173: b).

Pale gray or almost white, showing a tendency to dimorphism; pm. line on fore wing less sinuous than in *Psamathia*, on hind wing even more acute, normally reaching the marginal pattern; fine and pale, at least on hind wing, and defined on both sides with brown, more continuous than in *P. amplata*, for the most part nearly obsolete on fore wing but accented with dark brown toward costa.

Flight scattering (see diagram). The flight is too irregular to make the break in March and April significant. Mexico to mouth of Amazon.

EROSIA Guenée

Similar to *Epiplema*, third segment of palpus longer, porrect; venation the same except for the higher position of M₂ in hind wing; M₃ and Cu₁ free in fore wing, stalked in hind wing; outer margin toothed strongly at R, M₁ and M₃ of fore wing and female hind wing, but only

R and M_1 of male hind wing. Anal area of male involved in an enormous pale roll containing a mass of sex-scales.

Only a single species is now recognized, but the second one given here differs markedly in the length of palpi, and slightly in pattern.

1. Third segment of palpus very long, $2/5$ eye in male, over half in female; a larger species. *incendiata*
 Third segment of palpus only $1/4$ as long as width of eye, the whole palpus smaller; a smaller species. *palpulata*

EROSIA INCENDIATA Guenée

E. incendiata Gn., Sp. Gen. Lep. Het., 10, 35; 8: 4, 1852.

E. birostrata Gn., Sp. Gen. Lep. Het., 10, 35, 1852; *Epiplema b.* Biol., 2, 125, 1892; Seitz, 6, 1162, 1936 (pr. syn.).

♀ *E. furcillata* Fld., Reise Novara, Lep., 128: 30, 1875; Biol., l.c. (pr. syn.).

E. veninotata Warr., Nov. Zool., 12, 42, 1906; Seitz, l.c. (var. *incendiaria*).

Also figured: Seitz, 171: h7 ♂, i1, 2 ♀, 3 (*furcillata*), 4 (*veninotata*).

Male blackish, with a deep chestnut splash along interspace M_2 of hind wing and contrasting cream anal roll; ordinary lines blackish, irregular, and very close together, often fusing below cell. Hind wing below buff, contrasting. Female light brown, lines pale, somewhat defined with dark brown; both wings buff beneath. (Venation fig. 96).

Common (see diagram). Mexico to Brazil.

EROSIA PALPULATA spec. nov.

Closely similar to *E. incendiata*, but smaller; with ordinary lines more widely separated, and with shorter palpi.

Male coal black, with slight violet iridescence; front of scape and front edge of the scale ridge on vertex whitish. Fore with ordinary lines black, partly defined with chocolate brown, and the pm. outwardly with whitish also; am. strongly oblique out from costa to lower half of cell, then roundedly bent more than a right angle and oblique in to inner margin, also waved out over R and fold; pm. starting perpendicular to costa at $2/3$, excurved opposite cell and strongly retracted below, nearly parallel to am. line and rather near it from Cu_1 to Cu_2 , and connected to it by a black bar on Cu_2 ; st. represented by a black wedge-spot in cell R_5 and a tapering black streak from middle of cell M_1 , fading out very close to margin below Cu_1 , these two parts sometimes faintly connected. Whitish edging of pm. most distinct as two lunules each side of M_2 , often and st. with a whitish dot or shade before it below M_2 . Hind wing lighter and redder except over the roll; am.

and pm. lines from costa at $1/3$ and $2/3$, straight to the apparent "fold" (i.e. cell M_2) then right angled and nearly straight to inner margin; both dark, the former followed by coal black across cell M_2 , and latter interrupted the whole width of that cell; and pm. followed by a contrasting whitish bar from M_1 to M_2 ; st. represented by an almost marginal lunule from R to M_1 and a wholly separate oblique stria much farther basad in cell M_1 . T. edge coal black, even across cell M_2 . Cell M_2 brown, contrasting, from beyond am. line to t. band. Under side light luteous, including the part of the roll visible beneath, lightly flecked with blackish (Venation fig. 97). *Female* essentially like female of *incendiata*; reddish wood brown, am. line as in male; pm. fine, dark, strongly edged with whitish below M_1 , with a sharp angle at M_3 and angularly reëntrant from there to inner margin; connected to am. line by pale veins along Cu_1 and Cu_2 , but not especially close to am. line. St. with a blackish lunule parallel to the notch and a separate dot above as in male. Hind wing concolorous, with both lines dark, defined with whitish, and some pale lining on veins; am. angled out at lower angle of cell, replaced just above the angle by a brown shade, also roundly angled out in fold; pm. right-angled sharply at M_3 , then a little bowed out again at 2d A, otherwise concave to costa and inner margin; st. blackish, very closely parallel to margin from about M_1 to Cu_1 , the t.sp. with blackish flecking before the teeth but not black as a whole. Under side as in male. Expanse 30 to 35 mm.

Barro Colorado Id. Holotype male Feb. 1, 1935 (Bates); paratypes mostly females, Barro Colorado Id. Jan. 5 to early March (Bts. & Fried.). Type and most of paratypes in M.C.Z. Note the June flight of *incendiata* appears to be wanting (see diagram).

GATHYNIA Walker

Close to *Epiplima*. Front wider than in the preceding genera, about half as wide as eye; palpi weak. Fore wing (fig. 100) with R_5 and M_1 separate, though decidedly close together at base; M_2 arising about a third way down the cell; M_3 and Cu_1 arising close together. Hind wing of female with M_2 from middle of end of cell; but in male with dorsal part of wing modified into a large roll containing androconia, and even the costal venation sometimes modified and partly aborted.

The genus is found in similar species in the tropics of both hemispheres, but in the old world species I have seen, the male modifications are less extreme. Our species agree with several of the following genera in having the apex of the fore wing specially marked with black and copper.

1. Larger, ante- and postmedial lines brilliant copper in a cross light (60L 45U: 90L 75U); discal spot of hind wing obscure. . . *atriceps*
 Smaller; metallic iridescence confined to apical markings; hind wing with a black-ringed white discal dot with a plain cream dot adjacent to it. *biocellata*

GATHYNIA BIOCELLATA Warren

G. biocellata Warr., Nov. Zool., 12, 308, 1905.

Figured: Seitz, 172: e5.

Fore wing luteous, a double apical black spot, overlaid with copper; hind wing of female much darker, fuscous with copper ante- and postmedial and subterminal lines, the am. and st. preceded and pm. followed by irregularly wedge-like black spots, the st. ones almost marginal; d. d. white, black outlined, the black coppery in proper light, with a plain cream-white dot below it. Male with the tooth on R of hind wing very strong.

Dec. 26 (Bts.) 1 ♀. Described from Peru.

GATHYNIA ATRICEPS spec. nov.

Head with small eyes (for an Epiplemid), less than twice as wide as front, palpi rather weak, exceeding front; antennae prismatic; head and body smoothly scaled, with a ridge between bases of antennae; legs stocky, the hind tibia fusiform, as frequently in the family.

Male fore wing ample, the apex blunt, and outer margin bowed but not angled, much longer than inner margin, with a fovea-like structure near base of fold; Sc free; R₁ and R₂ stalked, R₃ and R₄ long-stalked, R₅ and M₁ approximate but free; M₂ about 1/3 way down on cell, mdcv being erect and ldcv angled and slanting in, M₃ and Cu₁ approximate, Cu₂ a little apart from them; 2d A sinuate opposite the fovea. Hind wing less than half as large; all but the costal region with Sc, the base of R and basal half of the free part of R involved in an enormous roll, enclosing a mass of sex-scaling; the remaining venation involved in the roll; including tip of R, more than half of M₁, and a slight thickening which may perhaps represent M₂; two free dorsal veins of uncertain homology. Cell obsolete. (Fig. 100).

Deep umber brown, shot with a slight violet glass, the body darker and head coal-black, immaculate. In most lights with the front even blacker just below the transverse ridge. Fore wing with antemedial line dull ochre, edged on both sides with black, strongly and irregularly excurved, and fading out to nothing on costal third; pm. band black, faintly edged on outer side with ochre, widest across fold, taper-

ing to both costa and inner margin; nearly obsolete at costa; and deeply narrowed or even interrupted at each vein. A double black subapical patch, extending from R_4 to M_1 and edged inwardly with ochre. Am. line followed by a blackish spot and second bar of ochre at inner margin. Iridescence very slight, best brought out by an intense illumination at 90 L or R 45 U; and viewed nearly vertically, shown most strongly by the underscaling of the pm. band, but partly masked by the blackish overscaling.

Hind wing plain umber brown with the violet gloss, with chestnut costoapical spot, oblique am. line and pm. bar across the roll. Under side dull smoky, immaculate. 24 mm.

Female fore wing venation as in male, but outer margin decidedly shorter than inner; hind wing with complete normal venation; costa deeply bisinuate with scale tufts before and beyond the deep middle notch; R and M_1 , M_3 and Cu_1 connate, M_2 arising slightly above middle of cell; outer margin toothed on veins, very strongly on R, less on M_3 , less on M_1 and slightly on the other veins.

Fore wing lighter umber brown than male, with the same markings, the apical band extending below M_1 and ochre perhaps a little stronger; hind wing with decidedly redder purple iridescence, a small white discal dot; am. band deep brown, edged with ochre, starting faintly from the discal dot, strong over fold where it is right-angled, and narrowing again a little to inner margin; pm. band similar, roundly bent opposite lower angle of cell, and only a little narrowed to costa; a browner marginal stripe, finely edged inwardly with ochre, tapering to nothing above M_1 and below Cu_1 . Under side immaculate fuscous, rather paler than male. 25mm.

Barro Colorado Id., C.Z., Panama, Male type Feb. 6, 1936, (Gertsch, Lutz, Wood, in A.M.N.H.) three male paratypes Jan. 24 and June 23 (Friedman, M.C.Z.), July 11 (Scrimshaw) all in bad condition; two female paratypes Jan. 29 (Bts.) and Feb. 13 (Friedman) in M.C.Z.

The species should perhaps be placed in *Capnophylla* Warren, on account of the enormous anal roll, but in the present nebulous condition of *Epiplemid* genera I prefer to use the older *Gathynia*. It differs from *C. albiceps* by the dark head and apparently more developed roll. It is a third larger than *C. semibrunnea* Dgn., described from Guiana.

ANTIPLECTA Warren

Very close to *Epiplema*, only differing in the low position of M_2 in fore as well as hind wings. In the present species R_2 is stalked with R_1 ,

but in the Porto Rican *ineptaria* Msch. (Forbes, Sci. Surv. P.R. xii, 72, 1930, suppl. 345, 1931) which Gaede in Seitz credits to Antiplecta, they are free. All the species I know have the male hind wing modified, an exceptional character in Epiplema.

The few known species range from Mexico and the Greater Antilles to Guiana.

ANTIPLECTA CINERASCENS Warren

A. cinerascens Warr., Proc. U. S. Nat. Mus., **30**, 400, 1906.

Not figured: (the Seitz reference to 173: e being an error).

Fore wing light buff, with a darker apical patch outlined with a broken blackish line; the ordinary lines dark brown, broken and largely obscure, the am. most distinct toward costa, the pm. forming a broad sweep on costal $3/5$ of wing, filled with dark brown, and a rounded-triangular dark brown patch on middle of inner margin; hind wing with the lines broken but well marked, the pm. most strongly bent on discal fold.

Nov. 13, Dec. 13 (Bts.), two females. Described from Guiana.

There is also a single male of a second species, the fore wing gray, immaculate except for the apical marking, the hind wing contrastingly marked in gray, dull brown and black. Nov. 4 (Bts.).

THYSANOCRASPEDA Warren

Like Epiplema except for the modified inner margin of male hind wing, which in this species takes the form of a lobe overhanging the abdomen, and fringed with long, curly hair; but typically is a roll and hair-pencil, about as in *Erosia*.

As now listed the few species are American, but the genus may turn out identical with the old-world *Dirades*.

THYSANOCRASPEDA NUDATA Warren

T. nudata Warr., Nov. Zool., **12**, 309, 1905.

Figured: Seitz, 171: g6.

Dull light red-brown, ordinary lines dark, partly edged with clay color, very faintly overlaid with copper (best visible at light 30U 90R and viewed at 80U 90R); both widening into dark brown patches at inner margin, and the pm. also thickened on the costal half; both lines irregularly excurved on hind wing, but the am. rather faint. Outer margin of fore wing with a gray patch along costal half of outer margin,

with about three black-brown dots on its inner edge and one just above it, also showing copper in a critical light. Fore wing with Sc and R₁ anastomosing, the dentation of the wings moderate and sinuation of costa of hind wing slight. The single female is the darkest specimen.

Oct. 25, Nov. 4, 25 ♂'s, Jan. 20 ♀ (Bts.). Described from Peru.

I have seen no authentic material of the species, and the original description leaves several points uncertain.

SKAPHION Gaede

Differs from *Bicavernosa* mainly in the shape of the marginal lobe of hind wing, (fig. 95) and should, I believe, be sunk.

SKAPHION LILACINA Gaede

S. lilacina Gaede in Seitz, *Macrolep. World*, 6, 1144, 170:b6, 1936.

Light powdery gray; lines fine, pale, defined with dark brown, ex-curved and angled, especially on M₃ of both wings; pm. followed by a pale shade on hind wing below M₃.

I have seen no authentic material of this species, but the specimens match the original description so far as it goes. The present species has palpi a little longer than in the following, whitish below; front black and vertex almost white.

Dec. 11, May 2♂, June 4♀ (Bts. & Fried.). Described from Venezuela and Brazil.

BICAVERNOSA Gaede

Antennae prismatic; similar to *Epiplema*; R₂ long-stalked, fore wing broadly lobed over M₃ and Cu₁ and concave above; hind wing with short teeth on R and M₃ and costa not sinuate, R and M₁ somewhat stalked, M₃ and Cu₁ arising separate.

BICAVERNOSA ALBIOCELLATA Warr.

Epiplema albiocellata Warr., *Nov. Zool.*, 6, 412, 1897.

Bicavernosa albilunata Gaede in Seitz, *Macrolep. World*, 6, 1146, 1936.

Figured: Seitz, 171: d1 (*albiocellata*); 172: a4 (*albilunata*).

Basal half of fore wing and costal half of hind wing dull mottled gray brown, scaled with violet-white and marked with brown and black; outer part of fore wing and dorsum of hind wing lighter powdery gray, contrasting on hind wing, crossed by a strong blackish pm. line on hind wing; a series of blue-white admarginal spots in a brown and

blackish marginal band on costal half of both wings. Discal lunule of hind wing only, contrasting, white, outlined except below with black.

Nov. 28, Dec. 4 (Bts.) both ♀. Described from Venezuela (*albicellata*) and Brazil (*albilunata*).

TRICOLPIA Warren

Near Bicavernosa, Cu_1 close to M_3 at origin in fore wing, which has the outer margin only slightly extended over M_3 and Cu_1 ; hind wing with a sharp angle at M_3 and slightly scalloped below, R and M_1 , M_3 and Cu_1 distinctly stalked.

The following species seems very near to the genotype, *T. acutaria* Walker, though the latter shows only a trace of the brown longitudinal stripe.

TRICOLPIA SEMISSARIA Herrich-Schaeffer

Arrhostia semissaria H.-S., Samml. Exot. Schm., 61, 39: 190, 1854; *Acidalia* s. l. c. p. 79¹: *Schidax* s. Guenée, Sp. Gen. Lep. Het., 10, 41, 1857; *S?* s. Gaede in Seitz, 6, 1148, 172: b1, 1938 (as *remissaria*).

Clay color, lightly striate with brown, the dark brown pm. line deeply sinuate on fore wing, and right-angled with two extended teeth on hind wing; a deep brown longitudinal line on fore wing from base to tip of M_3 .

Nov. 22 (Bts.), described from Surinam. This is the only specimen of which I have knowledge.

EPIPLEMA Herrich-Schaeffer

(*Callizzia* Packard, *Calledapteryx* Grote, *Erosia* auct. in part, etc.)

Antennae prismatic, pectinate in some foreign species, Front about half as wide as eye. Fore wing (fig. 98) with outer margin usually a little bent at middle; R_2 free, R_3 and R_4 , R_5 and M_1 stalked; M_2 from near upper angle of cell, M_3 separate. Hind wing usually with bisinuate costa, teeth at R and M_3 , and sometimes others; typically without sexual modifications on inner margin. R normally short-stalked, M_2 from near middle of end of cell, M_3 and Cu_1 commonly short-stalked.

A large genus covering the range of the family. Besides the following, additional species will certainly be taken on the Island. The North American *amorata*, *certiorata*, *dryopterata* and *slossoniae* belong here.

¹ It would appear that Guenée's work came out between the publication of Herrich-Schaeffer's plate and text, since each cites the other. Under a strict interpretation we should probably credit the name to Guenée, but it is always given to Herrich-Schaeffer (with practical justice).

EPIPLEMA ACUTANGULARIA Herrich-Schaeffer

E. acutangularia H.-S., Samml. Aussereur. Schm., 62, 81, fig. 324; *Erosia a.* Gn., Sp. Gen. Lep. Het., 10, 35, 1852.

Also figured: Seitz, 171: e1.

Brownish fuscous, the fore wing with separate costal and dorsal median dark brown patches. Male hind wing with special scaling in a narrow fold.

June 23 (Fried.), Lancetilla, Honduras (Bts.). Brazil.

MIMALLONIDÆ

(*Lacosomidae*, *Perophoridae*)

Body stout, head small, with pectinate antennae, almost as wide in female as in male, and with two or three exceptions abruptly narrowed at about $\frac{3}{5}$ their length to a serrate or subpectinate apex. Palpi short, not extending above middle of front, stout or weak, the third segment short save in *Lacosoma valva*; tongue obsolete. Body heavy, no tympanum; legs short and stout, hairy. Abdomen normally with a bifid terminal tuft in male, but a single one in female. Wings thick and strongly veined, fore wing with R_4 and 5 stalked, widely separated from the stalked R_2 and 3 , which arise close to R_1 before end of cell. M_2 from near middle of end of cell, or somewhat below, but well separated from M_3 ; 1st A obsolescent or lost. Hind wing with Sc and R adjacent at base for a very short distance, then Sc swinging abruptly away and not again approaching cell; M_2 from near middle of end of cell, 1st A lost.

Larva unique; the setae arranged as in the Micros, with iv and v close together below spiracle, setae single, save for four or more on outer side of proleg; crotchets in a complete circle. Head rugose, body very wide, the thorax chitinized and brilliantly marked, the abdomen soft. Young larva in a web between two leaves; later in a portable case, which is equally open at both ends, unlike the Psychidae. Case in a folded leaf or covered with frass, which in some species is said to be formed in flat blocks, ready for use. Pupa obtect, heavy, with short tongue and exposed femora.

A very curious little family, neotropical save for three species in the U. S. The adult characters and pupa are essentially of "Bombycid" type, but the larva is in every essential way a micro. Many of the species were first thought to belong to various Bombycine families, while the known larvae were generally taken for Psychids.

8. Terminal third of antenna formed of about 10 serrate segments.
Alheita
 Terminal third of antenna narrowly pectinate, in male normally
 narrowing abruptly to about 15 short-pectinate segments. 9
9. Outer margin of fore wing strongly irregular, falcate, bowed out at
 middle, and with two deep emarginations, above and below Cu_2
 *Lacosoma*
 Outer margin of fore wing less irregular, at most with a smooth
 emargination or a single scallop below Cu_2 10
10. Frenulum absent *Cicinnus*
 Frenulum weak but present *Bedosia*

MIMALLO Hübner

Female with outline of wings much less irregular than male.

A small genus with one very common species, covering the whole neotropical area; probably a direct derivative of *Cicinnus*. (Venation fig. 101.)

MIMALLO AMILIA Stoll

Attacus amilia Stoll in Cr., Pap. Exot., 3, 130, 265: D, E, 1780; *Mimallo a.*

Hübner, Verz. bek. Schm., 190, 1820; *M.a.* Berg, Horae Soc. Ent. Ross., 12, 163, 1876; Kirby, Handb. Order Lep., 4, 27.

Phalaena vorax Sepp, Surin. Vlinders, 1, 47, pl. 20, 1830.

Also figured: Seitz, 87: a1.

Larva: Sepp, Berg, Kirby ll.cc.

Blue-gray with brown markings and transparent discal lunule on fore wing.

Larva dark brown, in a case covered with chewed wood and some sand; on lower branches of guava. Sepp indicates the case as open only at the upper end, but his account may be incomplete.

Jan. 31 ♂ (Bts.), Apr. 20 and May 6 and 13 ♀ (Fried.) July-Aug. ♀ (Fairchild), Feb. 27 (A.M.N.H.). The moth is generally common from Mexico to southern Brazil, so the sporadic records suggest the main flight season was missed.

CICINNUS Blanchard

(*Saccophora*, *Perophora* Harris,—preoccupied)

This genus should probably be divided, the position of M_2 (figs. 99, 103) being more important than many characters used as generic in the family, and the few known larvae being of more than one type.

Although the frenulum is called "absent" there is in fact a minute vestige (see Comstock's "Introduction" fig. 713) and the genus is really very close to Pamea and Menevia.

A good sized genus, ranging from the U. S. to Argentine. Further species will certainly be found in the Zone.

For general accounts of *Cicinnus* larvae see especially Berg in Horae Soc. Ent. Ross., 12, 167-170 with pl. 4, 1876 (*C. despecta*); Sharp, Cambridge Nat. Hist., 6 (Ins. ii) 377-380, with figure and further references; Harris, Ins. Inj. Vegetation, 1st ed., 299, Flint, Ed. 415, with fig. 206 (*C. melsheimeri*); Beut., Bull. Am. Mus. Nat. Hist. 10, 434, 1898 (do.); Dyar, Jour. N. Y. Ent. Soc., 4, 92, 1896 (head setae of do.)

Note that *Cicinnus violacea* Sepp is a Thyridid and not a Lacosomid.

1. Discal spot transparent, wings long.....*forbesi*
 Discal spot opaque, wings short.....*solvens*

CICINNUS FORBESI Schaus

C. forbesi Schs., in Seitz Macrolep. World, 6, 644, 1928, 87:e3.

Olive-buff, markings brown, clean-cut; pm. right-angled opposite cell, double below; st. tangent to pm. at its angle, curving out to apex and anal angle, heavy toward apex, where it continues the lower part of pm. Am. line fine, transverse; a fine longitudinal line on fold to middle, and one from costa to discal spot, which is followed by a black spot.

C. volucris Schs. is closely related and regional, but has blunter wings.

Dec. 27-Apr. 23. Moengo, Surinam and French Guiana (types).

CICINNUS SOLVENS Dyar

C. solvens Dyar, Proc. U. S. Nat. Mus., 47, 253, 1914.

Smoky, powdered with blackish, the lines somewhat diffuse, blackish; pm. angled opposite cell and more or less double below; sinuous on hind wing; am. excurved, d.d. a black smudge.

Very close to *incerta* Msch. (Seitz 87: b2) but the apex less hooked.

Common (see diagram). Described from the Canal Zone.

PSYCHOCAMPA Grote & Robinson

(*Roelmana* Schaus)

Near *Cicinnus*, with the fore wing smoothly falcate as in normal *Cicinnus*, differing only in the slightly stronger frenulum and stalking

of M_1 . Schaus separates the two genera widely but I can see no difference. In both the following species there is a double transparent discal dot, unlike the type.

1. Smaller; pm. line very heavy, straight below the angle. *beta*
Larger, pm. line not strong, scalloped below the angle. *maloba*

PSYCHOCAMPA MALOBA Schaus

Roelmana maloba Schs. in Seitz Macrolep. World, 6, 671: 88:19; 1928.

Light red-brown, patchily shaded with gray; pm. line scalloped, not very strong; discal spots transparent, double.

Nov. 8 (Bts.) June 30, 1940 (Scr.). Guatemala to Brazil.

Dyar reports *P. beta* Schs. from the Zone (Seitz 87: b2). It is smaller, shaded with pink, with a dark oblique pm. line.

TROGOPTERA Herrich-Schaeffer

Fore wing squarish, with the anal angle deeply excavated in two scallops from Cu_2 to A, being more nearly longitudinal than transverse; frenulum short and weak, without a hook in male.

TROGOPTERA RUMINA Druce

T. rumina Dr., Ann. Mag. Nat. Hist., (6) 12, 1894; Seitz, p. 654, 1928.

Differing from other Trogopteras by the merely serrate apex of antenna and yellow color. There is a brown patch from pm. line to anal angle and a small spot at costa.

Mar. 26 (Fried.). Costa Rica (Druce) to Surinam (Cornell).

BEDOSIA Schaus

Hardly different from *Cicinnus*, save in habitus. The frenulum while slightly larger is still minute. In the present species the outer margin of the fore wing is notched below Cu_2 , and the hind wing regularly scalloped, but other species have an even margin. The species are close and I am not quite sure of my determination.

BEDOSIA TURGIDA Schaus

Cicinnus turgidus Schs., Ann. Mag. Nat. Hist., (8) 6, 420, 1910.

Figured: Seitz, 88; d3.

Large. Light pinkish gray, somewhat dusted with darker; pm. line angulate opposite cell, brown, connected to apex and anal angle by

powdery remnants of the st. line; pale brown dashes along basal half of fold and obliquely from costa to the transparent oblique discal dot, exactly as in *Cicinnus forbesi*.

Jan. 22, 27 ♂, Feb. 4 ♀ (Bts.) May 22 ♀ (Fried.) June 21 (Sc.).
Costa Rica to Colombia.

LACOSOMA Grote

Generally recognizable by the distinctively scalloped wing-form (see Lep. N. Y., p. 656, fig. 411). Frenulum weak but distinct. Female of present species (but not most forms) with long third segment of palpus.

Larva in a case formed between two leaves but less neat than in *Cicinnus* (Beut., Bull. Am. Mus. Nat. Hist., 10, 433, 1898.) I believe that "*Lacosoma*" *violacea* Sepp, figured with an immovable case, is a Thyrid, and apparently a miscolored Rhodogonia. Other species of the genus will surely be taken in the Zone.

LACOSOMA VALVA Schaus

L. valva Schs., Proc. U. S. Nat. Mus., 29, 329, 1905.

Figured: Seitz, 86: g4.

Dark brown, pm. line angulate as usual, darker with a fine following pale shade; am. line and d.d. faint; tips of fringe paler above middle.

Jan. 10, (Fried.), 12, 13, all ♀ (Bts.). Ranges to French Guiana.

ZAPHANTA Dyar

Antenna gradually tapering to apex; frenulum fully functional (fig. 104). A little, soft looking square-winged thing, not at all Lacosomid at first glance, and originally described near Apatelodes. The wing-form and small simply tapering antenna suggest a primitive type, but there is nothing abnormal in venation or genitalia.

ZAPHANTA INFANTILIS Dyar

Z. infantilis Dyar, Proc. Ent. Soc. Wash., 12, 85, 1910.

Figured: Seitz, 88:f6.

Light buff, base and margin of both wings more or less darkened.

Common (see diagram). Guatemala to Guiana.

MENEVIA Schaus

Differs from the species of *Cicinnus* which have mdev. long and bent, only in the fully functional frenulum (fig. 102).

MENEVIA LANTONA Schaus

Cicinnus lantona Schs., Proc. U. S. Nat. Mus., 29, 327, 1905.

Figured: Seitz, 88: f8.

Buff, shaded with gray; postmedian line dark brown, angled opposite cell; subterminal white, starting from near inner end of a longitudinal white dash below apex, tangent to pm. line, then running to outer margin above anal angle.

M. lucara is regional. It is grayer, the marginal area is broader and the genitalia different.

Common (see diagram). Ranges to Guiana (type).

ALHEITA Schaus

Terminal portion of antenna serrate, palpi short; abdomen with a bifid terminal tuft in male, a dorsal one in female. Fore wing with outer margin falcate and sinuate, not notched, venation normal; hind wing with fully developed frenulum, passing through a frenulum-hook in male.

1. Ground dull, violet-gray with contrasting pale fringes and patch at lower angle of cell of hind wing *rionica*
Ground warm brown, the fringe not paler and no pale spot on hind wing 2
2. Fore wing bluntly angled at middle of outer margin; outer line pale and even from costa to inner margin *caudina*
Fore wing merely falcate and sinuous; outer line dark, defined with pale and acutely angled opposite cell *subrubiginosa*

ALHEITA RIONICA Schaus

A. rionica Schs., in Seitz Macrolep. World, 6, 669, 88: h7, 1928.

The pm. line of fore wing is partly indicated by pale shading.

Oct. 21, ♀ (Bts.) Described from the Upper Amazons, from which Cornell also has specimens.

ALHEITA CAUDINA Schaus

Cicinnus caudina Schs., Proc. U. S. Nat. Mus., 29, 326, 1905.

Figured: Seitz, 88: i1.

The pm. is excurved to the costa and followed except toward costa by a chocolate brown area.

Jan. 9, ♀ (Fried.). Guiana (type), Upper Amazons (Cornell).

ALHEITA SUBRUBIGINOSA Dognin

Cicinnus subrubiginosa Dgn., Het. Nouv. Am. Sud, 19, 19, 1916.

Figured: Seitz, 88: i2.

Rich brown, somewhat shaded, noticeably paler on upper half of outer margin; pm. line blackish, defined with white frosting; a sharply curved subterminal shade from apex, not joining it; diffuse dark am. and discal shades. The present female is much larger, but I think unquestionably belongs to *subrubiginosa*.

Oct. 22 (Bts.). The type, from eastern Colombia, is colored like the B.C.Id. specimen, the three specimens from Venezuela are definitely paler.

ALEYDA Schaus

Apical portion of antenna short-pectinate; palpi short; fore wing rounded but with the apical portion extended, cell very short; d.d. transparent, oblique; hind wing rounded, small.

ALEYDA ACCIPITER Dognin

Cicinnus accipiter Dgn., Het. Nouv. Am. Sud, 10, 20, 1916.

Figured: Seitz, 86: g1.

Light red-brown and tawny brown, shading into gray and yellow; am. band heavy, black, excurved, pm. heavy, black and curved in from cell to inner margin and close to am. line,—above Cu_2 continued by a slight straight dark shade to apex.

Nov. 3, Jan. 4 (Bts.) May 2 (Fried.), all females. Originally described from Panama.

THYRIDIDÆ

Head prominent; ocelli and chaetosema absent; palpi usually moderate, sometimes with long third segment, tongue present. Body stout (*Thyridinae*) or slender with ample wings (*Rhodoneurinae*); legs strong, with all spurs. Fore wing with all veins present and normally separate, but occasionally R_3 and R_4 stalked (or others in some extra-regional genera); M_2 associated with Cu-stem, but generally well spaced out from M_3 ; 1st A absent in both wings; hind wing with Sc and R fused or closely parallel near middle of cell (*Thyridinae*) or along whole cell and base of R, as in *Pyralinae* and *Thyatiridae* (*Rhodoneurinae*), M_2 as in fore wing. Wing form often irregular, the pattern usually of irregular transverse striation, not clearly showing the cus-

tomary elements. Frenulum present, tympanum absent. The use of the customary pattern-names in this family is merely vaguely geographical.

A moderately small family, obviously connected to the pyralid stem, though formerly associated with the Sphingidae (*Thyridinae*) or the Drepanidae (*Rhodoneurinae*). The resemblance to the Drepanidae may really be significant, but I think the Lacosomidae and Epiplemididae, which sometimes are remarkably similar, are cases of parallelism.

Larva of normal pyraloid type; with strongly chitinized plates and tubercles, but no pattern. Prespiracular wart with 2 or 3 setae, abdomen with setae iv and v associated; subventral plate of mesothorax with 2 setae; preanal segment with a single plate bearing setae i as well as ii; prolegs with biordinal hooks in a circle. The recently studied larvae are borers in stems or make twig-galls, but Sepp's old figure of *Rhodogonia violacea* (i, pl. 30) shows it in a slender silken tube on a leaf.

References

- Boisduval: Sp. Gen. Lep. Het., 1, 488-493, 1873. (Thyris group only).
 Guenée: Ann. Soc. Ent. France, (5) 7, 275-304, 1877 (Rhodoneurinae as Siculides) (The accompanying plate was issued with Sp. Gen. Lep. Het., 10).
 Pagenstecher: D. E. Z. Iris, 5, 5-131, 1891 (Rhodoneurinae and part of Dysodia).
 Druce, Biol., 2, 184-188, 1895 (as Siculidae), also Dysodia 1, 324-326, 1889.
 Hampson: Proc. Zool. Soc., 1897, 603 (genera with keys, and catalogue of species).
 Dalla Torre: Lep. Cat., 20, 1-55, 1914 (Bibliography).
 Gaede in Seitz: Macrolep. World, 6, 1187-1213, pls. 173-175, 1936 (many figured and all catalogued).

Key to Genera

1. Hind wing with Sc fused to R about at middle of cell, then divergent well before end of cell (in American species) (fig. 105); body extremely stout, 1/3 as wide as length of fore wing (*Thyridinae*)
Dysodia
 Hind wing with Sc more or less parallel to upper side of cell and normally also to R after it diverges from cell; body moderate, about 1/6 as stout as length of wing (*Rhodoneurinae*).....2
2. Sc and R divergent before end of cell.....3
 Sc and R closely parallel to a point beyond end of cell.....4

3. R leaving cell of hind wing exactly at its end, approximate to M_1 ,
Sc diverging decidedly before it; eyes extremely large, front
close-scaled, half of second joint of palpus projecting beyond it.
Herdonia
- R leaving cell well before its end, at the same point where Sc
diverges from it (fig. 106); eyes less than twice width of front;
which is more tufted, palpi shorter *Ochrothyris*
4. Fore wing with R_3 and 4 stalked *Hypolamprus*
Fore wing with all radial branches separate (fig. 107) 5
5. Third segment of palpus much longer than second, porrect 4
Third segment of palpus upturned, usually short,—if long extending
above vertex *Rhodoneura*
6. Palpus with second segment upturned, fitting front to middle,
third a little longer and porrect 7
Palpus with either second or third segment much lengthened, ex-
tending as far as whole width of head in front of it *Risama*
7. Hind wing falcate and scalloped *Draconia*
Hind wing rounded apically, though deeply dentate below.
Zeuzerodes

DYSODIA Clemens

(Platythyris Grote and Robinson, Varnia Walker)

All veins arising separately from cell in both wings; base of M of hind wing simple, attached to an angulation of the vestigial discocellular. Hampson's figure of the venation in Fauna Br. India, Moths, 1, 368, 1892 and Proc. Zool. Soc. London, 1897, 609 fig. 5 is in error, M_2 and M_3 being present and separate in *D. ignita* as in other species.

Not a large genus, but with similar species in the tropics of both hemispheres, extending north to the warm temperate. Closely related to the holarctic genus *Thyris*. Dyar has revised the species with a key in Ins. Ins. Men., 5, 1, 37-45, 1913.

Larva forming a very messy and foul-smelling nest (whence the generic name),—reported in the U. S. on beans and Eupatorium.

1. Postmedial band of fore wing definite, oblique, the dorsal $2/3$ at
least normally nearly straight, at least on its outer side 2
Postmedial band broken into costal and dorsal patches or confused 7
2. Dorsal part of submarginal band represented only by a patch
starting below Cu_1 and resting on anal angle *thyridina*
Dorsal part of st. line tapering upward and either extending across
 Cu_1 , or represented by a separate spot above it 3

3. Ground yellowish, markings dull brown, without any tawny or red tints; pm. fine below the costal patch, upper part of st. line fine and extending right across apex. *sica*
Ground orange, the brown markings reddish. 4
4. Pm. line narrow toward inner margin, often linear. 5
Pm. line broad and widening at least a little to inner margin. 6
5. Pm. line usually rather thick; hind wing with confused markings or with st. as strong as pm. *speculifera*
Pm. line on both wings similar, fine and straight, the st. markings obscure; discal spot of hind wing usually smaller. *remie*
6. Discal spot of hind wing broad and squarish, but not reaching top of cell; ground fiery red, the markings contrastingly darker. *pyrsocoma*
Discal spot of hind wing constricted, with the upper outer angle reaching up to R; ground not much paler than markings *confusata*
7. Ground light buff with reddish striation; brown markings patchy, the pm. band of two separate costal and dorsal patches. *longalis*
Ground deep orange ochre, the brown markings tending to fuse and cover outer part of wing. *spissicornis*

DYSODIA SPECULIFERA Sepp

Bombyx speculifera Sepp, Surin. Vlinders, **3**, pl. 135, 1855; *Dysodia* s. Dyar, Ins. Ins. Men., **1**, 38, 1913.

Varnia aequalis Walker, List Lep. Ins. Br. Mus., **33**, 825, 1865; Pag., D. E. Z. Iris, **5**, 30, 1892.

Also figured: Seitz, 173: b5.

Life History: Sepp, l.c.

Ground a light clear orange-yellow, differing from the other species on the island. Larva in a rolled leaf.

Two of the specimens have the pm. line linear from the costal patch to the inner margin, in the rest it is thickened, but narrower and more even than in *pyrsocoma* and *confusata*. The National Museum consider this a mere variety, though it would run out elsewhere in Dyar's key.

Dec.-Jan.; Mar.-May (Bts., Fried., Bradley in C. U.) (see diagram). Ranges to Florida and Guiana.

DYSODIA REMIE Dyar

D. remie Dyar, Ins. Ins. Men., **1**, 44, 1913; Proc. U. S. Nat. Mus., **47**, 254, 1914.

Figured: Seitz, 173: a7 looks more like *remie* than *sica*.

A small, rather dull and plain but not very dark species.

Jan. 24, 26, (Bts.) Feb. 6, 12, Mar. 5 (Fried.). Described from the Canal Zone.

DYSODIA THYRIDINA Felder

? *Pachythyris thyridina* Fld., Reise Novara, **22**, 117: 20, 1875; *Varnia t.* Pag., D. E. Z. Iris, **5**, 31, 1892; *Dysodia t.* Gaede in Seitz, **6**, 1190, 173: a6, 1936. *Dysodia angulisola* Dyar, Ins. Ins. Men., **1**, 41, 1913; Proc. U. S. Nat. Mus., **47**, 254, 1914 (? syn. pr.); Gaede in Seitz, t.c. (? syn. pr.).

Fore wing with outer third contrastingly paler than basal 2/3, the boundary even and somewhat sinuate. Hind wing with a heavy dark band across it just before the hour-glass-shaped or double discal lunule.

Dyar left his *angulisola* as a synonym of *thyridina* in the Nat. Mus., but to me Felder's original figure looks more like *confusata*. Our species has been compared with Dyar's type.

Dec. 2 (Bts.) Feb. 24 and Apr. 7 (Fried.), June 21 (Sc.). Specimens are recorded under one or the other name from Mexico to Bolivia.

DYSODIA PYRSOCOMA Dyar

D. pyrsocoma Dyar, Ins. Ins. Men., **1**, 42, 1913.

Deeper and more brilliant orange than *speculifera*, the dark markings heavier. Our specimens are smaller than the type.

Jan. 29–May (Bts., Fried.). Costa Rica (type).

DYSODIA CONFUSATA Warren

D. confusata Warr., Nov. Zool., **15**, 332, 1908; Dyar, Ins. Ins. Men., **1**, 43, 1913.

D. c. olivescens Warr., l.c.; Dyar, l.c.

Figured: Seitz, 173: b4 (*confusata*), 2 (var. *olivescens*).

Ground deep blood red, the markings hardly darker, but purple-brown; in var. *olivescens* the ground is dull yellow, the markings umber, without the reddish tint.

Eleven specimens (see diagram) all of the typical coloring. Described from Ecuador.

DYSODIA SICA Druce

D. sica Dr., Biol. Centr. Am. Lep. Het., **1**, 325, 30: 4, 1889; Dyar, Ins. Ins. Men., **1**, 44, 1913.

Not *D. sica* of Seitz, 173: a7 (which is probably *remie*); omitted from Hampson and Lep. Cat.

The markings tend to be fine and crisp; the buff ground has a slight olive tint, and the usual fine reticulation is of the same dark brown as the markings, mixed with olive striae.

Jan. 26 (Bts.). Described from Mexico. I have seen no other specimen, but the original figure agrees.

DYSODIA LONGALIS spec. nov.

Structure wholly normal, marginal scalloping strong.

Head and prothorax dark brown, somewhat mixed with light brown and luteous, the edge of the collar finely luteous. Palpi with the lateral and dorso-medial keels on outer part of second and third segments finely edged with luteous. Rest of body orange red, mixed with luteous and brown, with a longitudinal gray-brown stripe its whole length, widest on thorax. Third segment of abdomen with some subdorsal brown, representing the usual transverse band, becoming conspicuous in the rubbed female.

Ground of wings bright straw color, the base of fore wing shaded and rest flecked with orange-red; fore wing with costa shaded with brown down to cell and R_5 and M_1 (fading out gradually), costal edge cut at five points with luteous; two transverse brown an. bands, median band broad at costa and a little narrower at inner margin, strongly constricted opposite lower angle of cell, where it is almost interrupted by areas of the straw yellow ground; st. line represented by a fusiform band from below costa to outer margin at Cu_1 and a transverse patch above anal angle, extending up to Cu_1 , and with a small separate dot above it. Hind wing similar in color, without definite bands, the yellow, reddish and brown being almost evenly mixed, but more solidly brown beyond cell and lighter toward anal angle. Discal lunule a deep horizontal V, its base roundedly swollen, its upper limb spatulate, reaching almost to upper angle of cell, and its lower limb shorter and pointed, pointing at lower angle of cell. All the light areas with scattered orange-red and brown flecking and striation, and the dark areas of fore wing with edges and striae of darker brown. Under side with the same pattern but with the orange-red elements absent and the light brown flecking dense.

I believe this species is close to *lusia* Dr. (Biol., 1, 325, 30: 6, 1889, Dyar, Ins. Ins. Men., 1, 44, 1913,—omitted from Hamps. and Lep. Cat.), but the latter according to the original figure and description is mainly orange, and there is no indication of the dorsal abdominal stripe or the lunule on hind wing. The dorsal stripe is unique in the

genus, so far as I know. In Dyar's key it runs to *D. sica* Dr., but has much broader brown markings and much warmer coloring as well as the dorsal stripe.

Barro Colorado Id., C. Z. Panama, holotype ♂ Nov. 29, 1934 (Bates), two paratypes ♀ Dec. 27, 1934, Feb. 4, 1935 (Bates) in M.C.Z.

DYSODIA SPISSICORNIS Warren

? *D. spissicornis* Warr., Nov. Zool., 15, 334, 1908.

Figured: Seitz, 173: b1.

Our specimen (a single female) is very dark, with small, strongly contrasting yellow spots on costa. The discal lunule of hind wing is very large, unlike the true *spissicornis*, which has two small windows. It may be a distinct species.

May 8 (Fried.) *D. spissicornis* was described from southern Brazil.

OCHROTHYRIS genus nov.

Antenna laminate, nearly simple; eyes not quite twice as wide as front, which is rough-scaled; palpi obliquely porrect, the second segment strongly exceeding front, third about 1/6 as long, blunt. Legs stout with the usual strong spurs, tibiae with rough spatulate scaling, the hind tibia swollen as usual in the family. Fore wing triangular, apex marked, outer margin a little excurved; all veins arising separately, R_2 and $3, 4$ and 5 approximate, M_1 a little above middle of cell; M_2 close to M_3 , above lower angle; base of M represented only by two slight folds, meeting mdev below M_1 and above M_2 respectively; hind wing with Sc curving down and becoming closely approximate to R just as it leaves cell, then diverging again, so that R just about bisects the angle between Sc and upper side of cell, which latter becomes M_1 without a break at end of cell; M_2 close to M_3 ; base of M as a strong forked fold, the upper branch attached to the moderate incurved mdev a little above middle, the latter exactly opposite M_2 ; outer margin even, bent at M_3 .

This genus differs from *Rhodoneura* mainly in the more prompt separation of Sc and R in hind wing, but also in appearance, in which it approaches the *Thyridinae*. The following species stands alone in it.

OCHROTHYRIS MESOGRAMMA spec. nov.

Between luteous and dull ochre. A pale stripe between antennae; palpi fuscous on outer side; vertex deeper and brighter ochre at sides, collar bright ochre; rest of body dorsally fuscous brown. Fore legs

fuscous on outer side, also middle and hind tarsi above; mid tibiae lightly and hind tibiae faintly obliquely banded with light fuscous. Wings of the luteous, a little shaded with ochre brown, reticulated and marked with the light fuscous brown. The principal brown markings on fore wing are a patch at basal angle, a transverse median fascia, with a hooked spur or extension into st. area opposite lower angle of cell, a weaker st. band across apex and short parallel band resting on anal angle, both irregular, and a smaller am. brown spot on costa. Hind wing with the brown median band becoming twice as wide toward inner margin, running just before the d.d., and with a st. spot riding on Cu_2 . Fringes of both wings a little darker than ground, the tips infuscated; d. dots all minute, quadrate. Female not at hand.

This pattern is definitely that of a *Dysodia*, from which the thin body and different Sc of hind wing will easily separate it.

Barro Colorado Id., C. Z. Panama; type Jan. 10, 1935 collected by Friedman, in M.C.Z., 7 paratypes, Jan. 14, Bates, June 17-20 (Friedman), 4 paratypes, June 22, 30, July 1, 1940 (Scrimshaw).

HERDONIA Walker

Separable from all other Rhodoneurinae known to me by Sc diverging from R well before end of cell, from most by R_{4+5} preserved as a distinct fold in the cell. The pattern is peculiar, and shared by Old- and New-World species.

HERDONIA BRIXIFACIES Dyar

H. brixifacies Dyar, Proc. U. S. Nat. Mus., 47, 256, 1914; Seitz, p. 1193, 1936.

Closely similar to *H. thetis* Dr. (Seitz, 173: b6), the most conspicuous pattern-elements being a double brown fascia across middle of hind wing, its outer part blackish across the cell, and followed by a large oval cream patch.

Oct. 25 (Bts.), rubbed. Presumably the normal flight period is earlier. Described from the Canal Zone.

HYPOLAMPRUS Hampson

This mere form-genus differs from *Rhodoneura* in the stalking of two radial veins. Most of the species are from the Old-world tropics.

1. Smaller; fore wing yellowish with oblique excurved transverse lines *arcuata*
Fore wing reddish with rounded gray patches *new species*
Large, fore wing buff, with irregular blackish medial patch and postmedian bar below costa (*Zeuzerodes fasciata*)

HYPOLAMPRUS ARCUATUS Pagenstecher

Siculodes arcuata Pag., D. E. Z. Iris, 5, 69, 1892.

Rhodoneura violalis arcuata Gaede in Seitz, 1212, 1936.

Rhodoneura changuinola Schaus, Ann. Mag. Nat. Hist., (8) 11, 235, 1913.

Straw yellow, with fragments of reticulation and four fine brown excurved transverse lines on fore wing, the second emphasized toward inner margin and much curved, the third running out about at middle of wing; hind wing with 3 lines, the middle one more oblique than the other two. Under side of fore wing with an am. patch of black and metallic scales in alternate bars, a smaller patch at end of cell, the dorsal half of which has yellow strap-shaped scales in place of metallic ones, and a small patch further out.

This species has nothing in common with *R. violalis* Pag.

Dec. 1, (Bts.) Described from Costa Rica and Chiriqui.

A single specimen of a species near *stellatus* Schs. was taken by Friedman, June 22. It is dull pinkish with many rather small powdery pale gray rounded patches, smaller and less regular than in *stellatus*. It is derived obviously from a different group of *Rhodoneura*, having very smooth palpi and no metallic scaling below.

RHODONEURA Guenée

(with *Brixia* Walker, *Siculodes* Guenée, etc.)

This varied genus covers half of the family. The structures vary in detail, and the appearance more. Some may have two radials shortly stalked as an individual aberration. Palpus varying in length but never with the third segment both long and porrect; wing form also variable, but never in the extreme type of *Zeuzerodes*. Many species have black and buff or metallic scaling on under side of fore wing, suggesting common types of sex-scaling, but present in both sexes.

Tropics of both hemispheres,—ranging well into the temperate of eastern Asia, but only once taken (*R. myrsusalis*) in New York.

1. Fore wing diagonally divided into a dark costal and a paler, pinker dorsal half, the boundary running from before middle of inner margin to apex.....*dimidiata*
- Fore wing with most of ground even or nearly so.....2
2. Entire body and wings spotted with black on an indian red ground.
new species

At least dorsal half of hind wing below with the striation slender or only brown.....3

3. Ground of apex of fore wing pale and evenly cut off, above and below *hedilalis*
Apex not specially marked 4
4. Under side of fore wing with a complex pattern, including black, metallic and slender cream scales and a tawny stripe from beyond middle to apex; upper side with about 5 black st. dots, approaching outer margin toward costa *thiastoralis*
Under side of fore wing without black, metallic or tawny markings 5
5. Under side of fore wing heavily striate with black, but leaving clear median and costo apical patches of the ground; abdomen extremely long *rufigrisea*
Under side of fore wing evenly marked or nearly so 6
6. Less reddish; third segment of palpus a quarter as long as second; discal spots of hind wing obscure, concolorous, a little larger than other flecking; body orange, contrasting *fulviceps*
Redder; third segment of palpus half as long as second; abdomen dull 7
7. Discal spot of hind wing transparent white, double; fore wing with a brown pm. spot on fold at the convergence of two striae. *anastomosalis*
Discal spot of hind wing dark 8
8. Wings broader; striation of vertical bars nearly the whole height of interspaces, except toward margin of hind wing; discal bar of hind wing similar to the others *myrsusalis*
Wings narrower; striations short, irregular and broken up; discal spots each of a pair of noticeably blackish dots *earneola*
- Group I: *Under side of fore wing in both sexes with black, buff and metallic special scaling; valves of male genitalia spoon-shaped.*

RHODONEURA HEDILALIS Walker

Pyralis hedilalis Wlk., List Lep. Ins. Br. Mus., 19, 895, 1859; *Siculodes h.*, Pag., D. E. Z. Iris, 5, 115, 1892; *Iza h.*, Biol., 2, 186, 1895; *Rhodoneura h.*, Hmps., Proc. Zool Soc. London, 1897, 619, 1898.

Figured: Biol., 59: 7.

Probably *Siculodes apicialis* Pag. D. E. Z. Iris, 5, 70, 1: 16, 1892; Biol., 2, 186, pr. syn.; Seitz, p. 1198, 173: e5 as good sp.

Palpus with third segment more than half second, upturned obliquely beyond vertex. Under side of fore wing with mixed black and blue-silver scaling along upper edge of cell, overlaid with pale yellow strap-

shaped scales and with a separate spot of the same beyond cell; a band of alternate black and blue bars along lower edge of cell, without overlay; and two squarish black spots between cell and fold, with the overlay covering not only the spots but the space between them.

Light indian red above with numerous partly confluent cream patches each outlined and crossed in the middle with fine blackish striae; the contrasting pale apex often enclosing a dark costal spot or lunule. Hind wings mostly pale, with similar pattern.

Dec. 1–Feb. 10 (see diagram). Ranges from the western limit of Panama (and Costa Rica if *apicialis* is the same) to British Guiana and the Upper Amazons. *Apicialis* was also reported from Peru and S. Brazil.

RHODONEURA THIASTORALIS Walker

Pyralis thiastoralis Wlk., List Lep. Ins. Br. Mus., 19, 893, 1859.

Siculodes punctum Fld., Reise Novara, Lep., 134: 7, 1875.

Also figured: Seitz, 173: g6 (as *punctum*).

Not *R. thiastoralis* Hmps., Fauna Br. India, Moths, 4, 481, 1896, which is a very distinct species, *R. rhodosticta* Swin.

Much smaller, fore wings with a falcate apex; palpi as before. Special scaling fundamentally as in *hedilalis*, much more extensive than in *rhodosticta* of the Old World, with which it has been confused. A small oval patch in place of the line at upper angle of cell, the yellow over-scaling forming parallel lines across it; a single patch below, separated from the blue-and-black barred stripe in the cell; the patch at end of cell largest.

Pale grayish pink; the striation weak and broken up, but with several rounded black dots in outer part of wing, especially five st. and 2 or 3 pm. ones on fore wing and two complete series on hind wing. Under side much like upper, but fore wing interrupted by confused brown bars and shades in the middle, an arcuate stripe from end of cell toward middle of outer margin, an anal bar, and a longitudinal orange stripe from end of cell to apex.

Oct. 31, Nov. 7, 17, Dec. 10 (Bts.) Amazons (type).

Group II: *Under side of fore wing without brilliant markings; male valves as far as examined slender, even in height, strap-like.*

RHODONEURA DIMIDIATA spec. nov.

Palpi loosely upturned beyond vertex, with third segment blunt, cylindrical, and half as long as second; fore wing with apex minutely

falcate, outer margin oblique and strongly convex on lower 2/3; much larger and longer than hind wing.

Thorax and ground of fore wing warm yellow-brown; vertex and large areas on tegulae heavily dusted with black, face red-brown; antennae light brick-red, slightly shaded above with blackish; palpi dull light brick-red, the tip of second segment darker and third segment blackish on outer and luteous on inner side. Abdomen paler and duller, the first segment on whole dorsum and second and third subdorsally heavily dusted with black.

Fore wing with costal half of the yellow-brown, toward the costa with a few blackish striae in paler and pinker flecks, and bounded below by a slightly darker vague straight streak from apex to middle of inner-margin, the extreme apex blackish; dorsal half of wing paler with a decided pink overcast, with about 15 slight gray flecks between the veins to represent the usual striae. Hind wing with less than basal third of the brown, and immaculate, bounded by a blackish band from middle of costa to near base of inner margin, the outer 2/3 strongly shaded with pink, with flecks about twice as numerous as on fore wing.

Under side of thorax concolorous, the mid tarsi and hind legs pale and spurs etc., touched up with blackish; under side of abdomen heavily dusted with gray, especially toward apex. Wings mixed yellow-brown (lighter than upper side) and pink, the posterior third of fore wing and the hind wing with many black striae, lying in the pink areas; fore wing also with four striae in cell M_1 . Discal spots not marked. 25 mm. 1 ♀.

Closely similar to the species figured in Seitz, 173: i4, as *subtransversalis*, but differing in the pink color and in the basal part of wings being darker than the outer; differing in the darker color and stronger oblique stripe from *subtransversalis* as originally described by Warren (Nov. Zool. iv, 408, 1897). In *R. ferruginea* Pag. (Seitz 173: f4) the longitudinal stripe runs to the base of the fore wing and is absent on the hind wing.

Barro Colorado Id., C. Z. Panama, Dec. 2, 1934 (Bates), holotype in M.C.Z.

RHODONEURA CARNEOLA Felder

Siculodes carneola Fld., Reise Novara, 134: 10, 1875.

Also figured: Seitz, 173: h1.

Structure as before, but with third segment of palpus more fusiform and apex of fore wing not actually falcate.

Grayish pink, finely striate with black, the striae with some tend-

ency to fuse in pairs and form X-shaped dots, never as regularly dot-like as in *R. thiastoralis* nor as bar-like as in *R. myrsusalis*. Discal dots 2 on each side of each wing, well separated, black. Under side similar, but striae on hind wing broader and more confluent, except the few marginal ones, which are as above. Black discal dots tend to become confluent. Body concolorous.

Nov. 23, Dec. 3 (Bts.). Described from the Amazons. I have seen no authentic material and the figure is amorphous, so do not guarantee the determination.

R. myrsusalis Walker (Seitz 173: e3) covers the whole warm temperate and tropical of both hemispheres, and will certainly be found in the Canal Zone.

RHODONEURA FULVICEPS Felder

Siculodes fulviceps Fld., Reise Novara, 134: 12, 1875; *Rhodoneura f.*, Hmps., Proc. Zool. Soc. London, 1897, 620, 1898.

Also figured: Seitz, 173: g7 (determination doubtful, the body being shown as concolorous).

Palpus with second segment upturned to middle of front, third obliquely porrect, only $\frac{1}{4}$ as long and pointed, not nearly reaching up to vertex. Fore wings blunt, with convex outer margin.

Pale brown with violet iridescence, the striation fine, somewhat blurred, the black edges of the striae frequently of a single line of scales; ground shaded, in the form of three broad subequal bands on basal $\frac{3}{5}$ and a half-band beyond. Body bright tawny orange with some brown shading, and a broad gray chevron on the abdomen edged with a brown line behind and shade before. Vertex bright orange.

Nov. 2, 8, 11 (Bts.). Described from Brazil.

Bates took a single specimen of a new species with heavy coal black spotting on body as well as wings, Jan. 20.

RHODONEURA RUFIGRISEA Warren

Iza rufigrisea Warr., Nov. Zool., 7, 119, 1900.

Figured: Seitz, 173: i5.

Palpus closely upturned to vertex, third segment more than half second, somewhat blunt. Male abdomen exceeding hind wings by half its length, almost half its length being formed of the 8th segment and genitalia (female not seen).

Tawny, leaning toward pale brick red, heavily striate with coal

black, the striae being partly confluent on most of wings, but paler subapically, below end of cell and along dorsum of hind wing, also leaving costa of fore wing tawny. Under side similar but with large areas over end of cell, subapically, and on dorsal part of hind wing, where the striae are thick and hardly darker brown than the ground.

Oct. 30–Jan. 8 (see diagram). Ranges to the edge of French Guiana (R. Maroni, in Nat. Mus.).

RHODONEURA ANASTOMOSALIS Pagenstecher

Siculodes anastomosalis Pag., D.E.Z. Iris, **5**, 76, 1892; *Rhodoneura a.* Druce, Biol. Centr. Am., Lep. Het., **2**, 10, 59: 6, 1895.

R. trigoniphora Hmps., Proc. Zool. Soc. London, 1897, 620, 1898.

Also figured: Seitz, 173: i3.

Easily distinguished by the two strong striae fusing into a dorsal patch. Palpus oblique, with long third segment.

I can see no difference either in the original descriptions or in National Museum specimens under the two names.

Nov. 13, Dec. 4 (Bts.). Wide-spread in tropical America.

ZEUZERODES Pagenstecher

Essentially like *Rhodoneura*, but with third segment of palpus longer than second, obliquely porrect. Fore wing triangular in general shape, long with the outer margin about as long as inner, but apex bluntly rounded; hind wing very small and triangular, the outer margin toothed on Cu_1 , excavate and scalloped above and below. Superficially the species resemble the North American genus *Meskea*, but they lack the specialized venation.

1. Dorsal portion of hind wing covered with a large whitish patch. . .

..... *umbrata*

Hind wing evenly colored with inconspicuous darker markings. . .

..... *fasciata*

ZEUZERODES FASCIATA Warren

Z. fasciata Warr., Nov. Zool., **12**, 41, 1905.

Figured: Seitz, 174: d5.

Palpi shorter than in typical *Zeuzerodes*, the third segment shorter than the second; fore wing with R_3 and 4 in the present specimen barely stalked (where they are closely approximate in *umbrata*), M_3 and Cu_1 closely approximate at base as in *umbrata*; hind wing with the apex

marked and subfalcate, the outer margin not scalloped, relatively not quite as small as in typical *Zeuzerodes*.

Buff, with normal dark brown striation, the most conspicuous markings being a blackish patch over end of cell (much larger in the Seitz figure than the present specimen) and a black bar in the fork of R_{3+4} ; abdomen with a double triangular brown spot on second segment.

Feb. 5 (Fried.). Described from Peru.

ZEUZERODES UMBRATA Schaus

Rhodoneura umbrata Schs., Ann. Mag. Nat. Hist., (8), 17, 236, 1913.

Fore wing blackish, somewhat shaded with clay color, especially toward anal angle, and with warmer brown below costa. Hind wing with costal half blackish and dorsal half mixed clay color and buff, the area beyond cell blackish, but with coarse clay striation.

Oct. 10 (Bts.) Feb. 8, Mar. 4, 22 (Fried.). Described from Costa Rica.

Z. leuconotula Pag. (*fumatilis* Pag., Seitz, 174: d3) is similar, but with the patch on the hind wing shorter and whiter.

RISAMA Walker

Similar to *Rhodoneura*, save for the long porrect third segment of palpus. Apex of hind wing somewhat marked but not at all falcate. Head with extremely large eyes, front $1/3$; tongue weak. The two species are not at all closely related.

1. Yellow and pink, hind wing with translucent spots. *aurorula*
Buff, wings opaque. *avicula*

RISAMA AURORULA Guenée

Siculodes aurorula Gn., Sp. Gen. Lep. Het., 10, Siculodides, pl. 1, fig. 4, 1857; Ann. Soc. Ent. France, (5) 7, 294, 1877; H.-S., Samml. Aussereur. Schm., fig. 402, 1858.

R. picta Wlk., List Lep. Ins. Br. Mus., 32, 519, 1865.

Also figured: Seitz, 175: b3 (as *picta*).

Front flat, palpus projecting most of its length beyond it, the third segment obliquely porrect and a little longer. Fore wing with costa bisinuate, the basal part much broadened and apical part convex, apex somewhat falcate, anal angle notched. Hind wing larger with strongly convex costa and apex rounded over.

Yellow, shaded with pink and lightly reticulated with brown in places; hind wing with a large round hyaline discal spot, followed by several small ones in a median brown band of heavier reticulations.

Mar. 25 (Fried.) July 30, 1924 (Bks). Brazil.

RISAMA AVICULA Guenée

Siculodes avicula Gn., Ann. Soc. Ent. France, (5) 7, 293, 1877.

Drepanodcs rchemensaria H. Edw., Papilio, 4, 19, 1884 (omitted from Lep. Cat.)

Aziba macropterana Dr., Biol. Centr. Am., Lep. Het., 2, 157, 59: 8, 1895.

Aziba substrigata Warr., Nov. Zool., 7, 117, 1900.

Also figured: Seitz, 175: d3 (*macropterana*), c4 (*substrigata*).

Front bulging strongly transversely, the angle between front and eyes forming a groove; palpi with first two joints very short, hardly exceeding front, the third exceedingly long and slender, five times as long, and porrect. Fore wing strongly falcate, the outer margin not notched, hind wing with costa less arched, apex more marked and outer margin less curved.

Buff, with brown striation covering the surface, but tending to gather in three oblique bands on both wings.

May 1 (Fried.) Mexico to Taboga Id. and Brazil.

DRACONIA Hübner

Similar to *Risama*, the hind as well as fore wing with falcate apex, in the present species sharper than on fore wing. Palpi rather short, third segment longer than second and porrect. The few species are large and showy.

DRACONIA ALBIAPICATA Warren

D. albiapicata Warr., Nov. Zool., 15, 331, 1908.

Figured: Seitz, 175: b4.

Light buff, finely and irregularly reticulate with two shades of brown; with a brown subcostal stripe, a narrow straight median red-brown band, widened to both ends and forked at the upper, a long-triangular costo-apical patch, enclosing two pale spots, the st. line curving sharply out along its lower boundary to apex; two st. brown patches above anal angle. St. line in the present specimen nearly straight, pointing toward outer margin above anal angle, but fading out, typically more parallel to outer margin. Hind wing with the m. stripe and two st. patches, also a minute translucent discal dot.

This is the species identified by Gaede in Seitz as *albiapicata*, but not the one standing in the Nat. Mus. under that name. This species and *patercula* Pag. (Seitz 174: g3) differ from the rest of the genus in the fine even st. line, abruptly curving out to the apex.

Dec. 6 (Bts.). Described from southern Peru.

CASTNIIDÆ

Head prominent, with well developed tongue and palpi and strongly clubbed antennae. Body stout. Legs strong. Fore wing with veins heavy but scaling loosely attached as in the true micros, the scales the largest of any Lepidoptera. Accessory cell well set off, its broad side lying on discal cell, usually with all five radials arising separately from discal or accessory cell. Base of M_3 preserved, of M_{1+2} usually lost; M_2 associated with M_3 (quadrifid); 1st A fully developed, very slightly weaker than 2d A; 3d A normally forked, the upper fork joining 2d A. Hind wing with strong frenulum; Sc abruptly diverging from cell from near base; M as in fore wing. Only two anals. Early stages essentially as in the Cossidae, borers in Monocotyledonous plants,—*Heliconia*, palms, orchids, etc. Larva sometimes with prolegs reduced, the hooks vestigial.

A rather small family of large and extremely showy moths, mostly neotropical, but with well marked relatives in the Indomalayan region (*Tascinidae*) and Australasia (*Synemoninae*). They are certainly the connecting link between the butterflies and moths, but in fundamental characters closer to the Cossidae than to the skippers. A number of species should be taken on Barro Colorado Island, but since the systematic collecting of Lepidoptera has been at night and the whole group are strict day-fliers they have been neglected. There is a full revision of the American species by Houlbert in Oberthür's *Études Lep. Comp.*, vol. 15.

CASTNIA Fabricius

I treat Houlbert's "genera", which are based on pattern-types, as species groups, and consider his tribe "Castniini" to be about a sound genus.

Wings ample; fore wing triangular with marked apex; venation normal for the family, with base of M_3 strong in both wings, M_{1+2} rudimentary, R's all arising separately from cell and accessory cell, or with R_{3+4} stalked, rarely with R_5 from base of the stalk, or (*cochrus*) with

acc. cell open between R_3 and R_4 . Upper branch of 3d A erect or recurrent. Coloring usually bright but not mimetic, with hind wing more brightly marked than fore wing and covered in sleeping position. Normal resting position with wings erect, as in the butterflies.

CASTNIA CACICA PROCERA Poisduval

[*Castnia cacica* H.-S., Samml. aüssereur. Schm., pp. 56, 79, pl. (54), fig. 143, 1854; Boisduval, Sp. Gen. Ins., Lep. Het., 1, 502, 1875.]

[*Amauta cacica* Houlb., Et. Lep. Comp., 15, 129, pl. I, fig. 46 (type), 1918.]

C. procera Bdv., Spec. Gen. Lep. Het., 1, 503, 1874; *Amauta p.* Houlb. Et. Lep. Comp., 15, 131, 438: 3780, 1918.

Also figured: Seitz, 1: b2; Preiss, Abb. Nachtschm., 9: 2, 1888 (as *cacica*).

Fore wing green-black with straight white pm. line; hind wing brown-black with red or yellow pm. band and about 4 st. spots.

July-Aug. 1923 (Fairchild). The race ranges north to Mexico, typical *cacica* occurs in Colombia.

COSSIDAE

(with *Zeuzeridae*)

Medium to very large moths, some females being the bulkiest of known Lepidoptera. Head small and retracted; mouth parts weak or rudimentary, even the palpi usually being weak; antennae of male usually pectinate (though laminate in the typical genus *Cossus*, from Europe); legs short and stout, often hairy, with short spurs, frequently the upper pair of the hind tibia lost. Wings (figs. 108-112) sphinx-like, very heavily veined; acc. cell lying broadly on discal cell, base of media preserved, almost always forked in cell, enclosing an intercalated cell, but occasionally with the upper fork vestigial; 1st A preserved, weak in some small species, frequently connected or anastomosing with second; 3d A running into second, but frequently with a free posterior branch. Hind wing usually with Sc and R parallel about to end of cell, sometimes connected by a cross-vein, which may be present or absent even in species of the same genus; base of M forked, 1st A preserved.

Larvae boring, most often in hard wood of trees, in many species taking more than a year to develop; jaws strong, held horizontally; mesothorax with one subventral seta, prespiracular wart of prothorax with 3 setae; uppermost setae of 9th segment of abdomen well separated on dorsal line, and prolegs with a circle of hooks or two transverse

bands. Pupa incomplete, partly emerging from the bore before the moth comes out.

One of the most primitive families of higher moths, but as the early stages show, on the line of the Tortricidae, and very close to the line that leads through the Castniidae to the butterflies. In strict classification this family belongs to the "Micros", but is here discussed along with the Bombyces, largely because it has historically interested the same persons.

There is substantial agreement as to its major division, but some uncertainty what rank to give to the divisions (as often in the field of Zoology). I treat the *Zeuzera* and *Cossus* groups as subfamilies, since they contrast in all features studied, even the type of egg, which is upright in the *Cossinae*, flat in the *Zeuzerinae*. The types which have 1st and 2d A connected (*Hypoptinae* of Barnes and McDunnough) agree with the *Cossinae* in more fundamental points of venation, and may be made a tribe.

References

Newman, Ent. Mag., **1**, 68, 1832 (first definite recognition of the family,—our *Cossidae* are his *Stygiidae*, *Zeuzeridae*, *Cossidae*).

Hampson, Fauna of British India, Moths, **2**, 304–314, 1892 (key to Oriental genera).

Schaus, Proc. U. S. Nat. Mus., **29**, 339, 1907 (key to American genera).

Barnes & McDunnough, Contr. Nat. Hist. Lep., **1**, (1); 1911 (monograph of North American species, with keys).

Hering, in Brohmer's Tierwelt Mitteleurop., **18**, 33, 1935 (latest key to Palaearctic genera).

There has been no general review of the family, nor, so far as I know, any key to the Ethiopian genera.

Key to Genera

1. Intercalated cell of hind wing, and in the typical group of fore wing also, more or less quadrangular, located in the lower outer angle of the discal cell, the lower branch of M much shorter and more transverse than the upper (figs. 111, 112)¹ Male antenna pectinate at base, abruptly changing into the simple apex, the basal part fitting over the eye in resting position like a Lyonetiid eyecap, female antenna simple; R₁ in all the present species arising from the very large acc. cell (free though approximate in *Zeuzera*); 1st A free (*Zeuzerinae*).....*Xyleutes*

¹ In *X. strigifera* Dyar the intercalated cell of the fore wing is lost by the complete fusion of M₁₊₂ with M₃, and in hind wing by the obsolescence of M₃, but the enormous acc. cell will indicate it as a *Xyleutes*.

- Intercalated cell of both wings (figs. 108-110), when present, fairly symmetrical, with both branches of M subequal in length and ending at outer end of discal cell; antenna of both sexes generally pectinate, gradually narrowing to apex; occasionally unipectinate, serrate, simple or laminate; R_1 usually arising from discal cell before acc. cell (*Cossinae*).....2
2. R_1 arising from acc. cell; frequently all radials free.....3
 R_1 arising from discal cell; usually with R_4 and 5 long-stalked...4
3. R_{3-5} on a common stalk.....*Trigena*
 No three radial veins from a common stalk.....*Cossula*
4. Vein 1st A of fore wing distinct and free (*Cossini*).....*Miacora*
 Vein 1st A of fore wing anastomosing or connected with $2d A^1$, sometimes reduced to hardly more than a fold in species where Cu_2 and $2d A$ are close together (*Hypoptini*).....5
5. Int. cell preserved in both wings, its upper boundary reduced to a fold or a spur on udev. in a few small species (fig. 108).....6
 Int. cell of both wings absent, there being no trace of M_{1+2} ; R and M_1 connate in our species, usually separate (fig. 110).
Inguromorpha
6. Hind wing with R and M_1 stalked, usually for a considerable distance (fig. 108).....*Givira*
 Hind wing with R and M_1 arising separately; fore wing in present species with Cu_2 and $2d A$ closely crowded (fig. 109)..*Langsdorfia*

XYLEUTES Hübner

(*Duomitus* Butler, *Endoxyla* Herrich-Schaeffer, etc.;
 with *Psychonoctua* Grote)

Head and venation figured by Hampson, Fauna of Br. India, Moths, 1, 308 (as *Duomitus*).

A showy genus of the tropical and subtropics of both hemispheres, barely reaching the United States (*X. ramosa* Schs.). *Phalaena Noctua strix* of Linnaeus, considered by him the largest of moths, belongs to this genus. (Venation fig. 112). The larvae bore in the solid wood of orange, willow, mimosa and other trees (Costa-Lima).

1. Ground dull pearl gray, usually with base of costa and discal lunule black and contrasting, striations faint.....*terrafirma*
 Ground pure white, with finer, strongly black striation.....2

¹ The connecting vein sometimes represented by a mere fold.

2. Fore wing with even transverse striation over whole surface, no brown shade below cell.....*strigifer*
 Fore wing with a longitudinal discolorous shade below cell or covering dorsal part of wing.....3
3. Ground of costal half brown, contrasting with the whitish dorsal half.....4
 Ground all of one color; outer half with a longitudinal black splash.....5
4. Expanse 50-75 mm. basal 2/5 of wing all pale below the cell.*ramosa*
 Expanse 100 mm., area of basal half of wing between Cu and 1st A largely filled with black.....*xylotribus*
5. Striation finer, e.g. with about 20 meshes in cell M₃; hind wing mostly striate.....*pyracmon*
 Striation coarser, with about 10 meshes in cell M₃; hind wing variable, but generally rather evenly pale.....*comisteon*

[XYLEUTES STRIGIFER Dyar

X. strigifer Dyar, Proc. U. S. Nat. Mus., **38**, 269, 1910; *X. strigifera* (sic) Lep. Cat., **29**, 54, 1923, and later authors.

Figured: Seitz, 181: d1 (correctly *strigifer* on plate, as *strigifera* in text).

Lancetilla, Tela, Honduras (Bts.), Guatemala (Bequaert). Ranges north to Mexico.]

XYLEUTES RAMOSA Schaus

Zeuzera ramosa Schs., Proc. Zool. Soc., 1892, 329.

Zeuzera ramuscula Dyar, Sci. Bull. Bklyn. Inst. Mus., **1**, 200, 1906; *Hamilcara* r. B. & McD., Contr. Nat. Hist. Lep., **1**, 22, 6: 2, 1911.

Zeuzera acetes Dr., Ann. Mag. Nat. Hist., (7) **7**, 436, 1901.

Also figured: Seitz, 181 e4 ♂, 2 ♀.

Fore wing with costal half light brown, shading into black, dorsal half striate with dark gray on white, the boundary sharp but highly irregular.

Jan. 31, Feb. 4, 23, Mar. 2. Arizona to Brazil and Paraguay.

XYLEUTES XYLOTRIBUS Herrich-Schaeffer

Endoxyla xylotribus H.-S. Samml. Aussereur. Schm., 58, *Zeuzera x.*, p. 78, fig.

37 ♂, 38 ♀, 1850-1858; *Z. (Endoxyla) x.*, Pack., 2d-3d Ann. Rept. Peabody Acad. Sci., p. 86, 1871.

Also figured: Seitz, 181; c1 (♂ as ♀ *xylotriba*), b3 ♀ (as ♂ *xylotriba*).

Much larger than the preceding, the black shading heavier and more irregular; the reticulated dorsal area continued, though without definite boundary, up and around the apex.

Nov. 26 (Bts.). *Lancetilla*, Honduras (Bts.). General in tropical America.

XYLEUTES COMISTEON Schaus

Zeuzera comisteon Schs., Ann. Mag. Nat. Hist., (8) 7, 628, 1911.

To be figured: Seitz, 167: e.

Markings generally coarser than in *Z. pyracmon*; the body rather generally solid black in *comisteon*, with white dorsal stripe in *pyracmon* but variable in both; hind wing varying from white to pearl gray with only under side of costa barred.

Not as common as *pyracmon* (see diagram). Described from Costa Rica.

XYLEUTES PYRACMON Cramer

Sphinx pyracmon Cr., Pap. Exot., 3, 169, 287: B, 1780; *Xyleutes p.*, Dalla Torre, Lep. Cat., 29, 68, 1923.

Duomitus pyracmonides Schs., Jour. N. Y. Ent. Soc., 9, 45, 1901; Dyar in Seitz, 1266, pr. syn. 1937.

Zeuzera putrida Percheron, Gen. Ind. Lep., 4, p. 1, 1838; Dyar l. c., pr. syn.

Cossus palmarum H.-S., Samml. aussereur. Schm., fig. 36, 1853; Kirby, Cat.

Lep. Het., 875, no. 18, 1892 (syn. *putrida*); Dyar, l. c. syn. *pyracmon*.

Zeuzera cognata Dr., Biol. Centr. Am. Lep. Het., 1, 231, 24: 6, ♀, 1887 ≠, not Walker; Dyar l. c., pr. syn.

Zeuzera fracta Walker, List Lep. Ins. Br. Mus., 7, 1542, 1856; Dyar l. c., pr. syn.

Z. lelex Dgn., Le Nat., 13, 121, 1891.

A little larger than the preceding, the male with reticulation sharp on fore wing, and hind wing more or less infuscated and with a large area of not always too contrasting reticulation. Female dimorphic, typically like male, in var. *lelex* (which is the normal female in Venezuela) with the fore wing suffused with powdery gray and reticulation obscure.

Common (see diagram). General in the Neotropical Region.

XYLEUTES TERRAFIRMA Schaus

Psychonoctua terrafirma Schs., Ann. Mag. Nat. Hist. (8) 7, 629, 1911; Dalla Torre & Strand, Lep. Cat., 24, 183, 1929.

P. nullifer Dyar, Proc. U. S. Nat. Mus., 47, 329, 1914.

Figured: Seitz, 181: f1.

Fore wing with strong discal dot and a blackish basal costal patch, strong in male, often faint or lost in female. *P. nullifer* of Dyar was based mainly on the female without the patch, though normal males were included in the original series. By some aberration the Lep. Cat. puts this group in the Psychidae, though it is not even generically distinct from Xyleutes. As a group it is the only Cossid type represented in the West Indies, where each island has a sub-race.

Males Jan. 24–Feb. 3 only (!), females Jan. 24, 26, Feb. 2, 5, 7 (see diagram). The startlingly short flight period must be of biological significance, though I found no such phenomenon in Porto Rican *X.(P.) personalis*. Representative species (or perhaps races) range from Mexico (*xuna* Dyar) to the Amazon (*albogrisea* Dognin).

COSSULA Bailey

(with *Hemipecten* Dyar, not Adams & Reeve, *Schausiana* Strand, *Hemipectrona* Schaus; *Costria* Schaus)

Antennae unipectinate or bipectinate, almost as wide in female as male; fore wing with R_1 from acc. cell, the other veins various,—most often with R_2 and 3 connate, and R_4 and R_5 separate, but in the type species with R_2 and $3, 4$ and 5 stalked; never with R_3 stalked on R_{4+5} ; int. cell symmetrical in both wings; hind wing with Sc free, R and M_1 parallel or connate, M_2 unusually far from M_3 , at least in hind wing. Only in *C. vinnea* is Sc connected to the cell before middle.

Larva of the North American *C. magnifica* boring in the trunk of oak and hickory (Bailey, Pap., 2, 93).

1. Outer margin not specially marked; fore wing with a dark median shade-spot *vinnea*
Fore wing with a contrasting dark (black, or black with brown or buff) apical patch 2
2. Ground silky blue-gray with little striation *abnoba*
Ground whitish with many fine blackish striae 3
3. Apical patch solid black; am. spot in fold large, diffusely extending up into cell *duplex*
Apical patch partly brown or buff; am. spot in fold small and normally round 4
4. Lighter part of patch brown except at costa, darker than the general ground *arpi*
Patch largely yellow with a short blackish streak on outer margin *niveogrisea*

COSSULA ARPI Schaus

Costria arpi Schs., Jour. N. Y. Ent. Soc., 9, 47, 1901.

Cossula nigripuncta Dgn., Het. nouv. Am. Sud, 12, 29, 1910; Dyar in Seitz, 1270, pr. syn.

To be figured: Seitz, 167: h.

R₂ and ₃ connate or short-stalked, R₅ well separated; M₂ arising from end of cell below the end of the base of M₃; antennae bipectinate.

Marginal patch with upper quarter cream, residue gray-brown, heavily shaded with black. Two separate black am. spots.

Jan. 25 (Bts.). Guatemala to Southern Brazil.

Dyar also reports *C. niveogrisea* Schs. (Seitz 183: a1) from the Canal Zone. It ranges from Guatemala to the Amazons.

COSSULA DUPLEX Dyar

C. duplex Dyar in Seitz, Macrolep. World, 6, 1271, 1937.

(Not Seitz, 182: g, as cited in text,—this is *Prionozygus duplex* Schaus).

M₂ arising rather above M₃; terminal branches of R all arising separately; antennae unipectinate and laminate.

Darker and somewhat more mottled than the preceding; wings blunter; a single larger black am. patch.

May 31, (Fried.) Described from the Canal Zone.

COSSULA ABNOBA Schaus

Costria abnoba Schs., Jour. N. Y. Ent. Soc., 9, 47, 1901; Lep. Cat., 29, 29 (as *adnoba* in error).

To be figured: Seitz, 167: g.

Antenna unequally bipectinate. Fore wing with R₂ and ₃ barely stalked, the rest free; base of M slightly below M₂.

Blue-gray, glossy; discal spot vague, pale with a dark center; am. black spots more or less double; apical brown patch very large, the brown part extending down to the black bar at tip of Cu₂.

May 21 (Fried.). Described from Brazil. The present specimen like the type is a female.

MIACORA Dyar

(*Toronia* Barnes & McDunnough; *Psychopsis* Dyar)

Appearance Cossus-like, at least in female. Antennae various; fore wing with R₄₊₅ stalked, and R₃ connate or stalked with them; M₂ arising below point of ending of base of M₃; int. cell symmetrical,

short in the present species. Hind wing with frenulum weak or absent, Sc and R separate, int. cell absent, the simple base of M ending at an angle in mdev.

The present species is sexually dimorphic, the male being a *Psychopsis*, though a little less extreme in the length of cell of hind wing than *P. infantilis*, the female a normal *Miacora*. With this hint I suspect *adolescens* Dyar may also be the female of *infantilis* Dyar.

MIACORA DIPHYES spec. nov.

Male. Head and thorax white, lightly flecked with gray scale-tips; front and palpi gray, third segment of palpi and antennae blackish; abdomen blackish, with contrasting pale tip; basal hair lighter gray; posterior massive tuft of thorax, small basal and large second tuft on abdomen rather dark gray. Legs dark gray, tarsi black, with a white bar at tip of each segment. Fore wing dark gray, shading into light gray along outer margin, narrowly and weakly at apex, becoming a patch over branches of Cu, where there are some white striations superposed. Costa with about four whitish bars, one at end of cell continued into a whitish transverse discal shade, the next extending diffusely across radial branches, the last two small. Some black striae, an am. one below cell of black scale-tips only; a median one across submedian space; a st. one starting from costa but almost lost in the blackish ground, then roughly parallel to outer margin to M₃, where it ends tangent to another black stria from the tip of the second pale costal bar to anal angle. Fringe barred medium gray and whitish. Hind wing brown-black, the fringe-tips pearl gray, contrasting. Under side all brown-black, the fringe as above, but basal half of fringe of hind wing barred like fore wing. 27 mm. Barro Colorado Id., C. Z. Panama, Dec. 23, 1934, Bates; holotype male in M.C.Z.

The supposed *female* is much larger and lighter: Body gray, somewhat mottled, the thorax not noticeably paler than abdomen; but face, palpi and antennae dark gray, contrasting. Fore wing whitish, flecked and striate with gray; and marked with black,—median line conspicuous, formed of a costal spot, a bar across cell and a bar across fold, almost continuous with each other; first two costal whitish bars as in male, but the first only conspicuous as a whitish cell-spot, and the second weak; outer lines essentially as in male, the pm. and st. actually joining, the st. clear to costa where it bifurcates, and the pm. sending an anterior branch down and forward to Cu₂. Hind wing mouse gray (fringes defective). Under side gray, with the dark mark-

ings of fore wing diffusely repeated, and hind wing with faint and diffuse dark striation. Under side of body and legs of the same mottly gray, the tarsi only vaguely banded in light gray and blackish. Expanse 35–40 mm. Barro Colorado Id., Feb. 22 (Bts.) Mar. 13, 16, 1936 (A.M.N.H.). The description above is from the best specimen (the Mar. 16 one). In the other two the striae are more broken, and the anterior branch of the pm. is represented by broken striae; and in one the black bar in cell is replaced by an elliptical spot. If these two sexes belong to the same species it makes a case parallel to *Prionoxyxus*. Other species of *Miacora* have a male like the female. Venation with R_3 sinuous, approaching R_4 where it leaves R_5 , R_4 and 5 stalked; M_2 opposite base of M_3 ; hind wing with R and M_1 stalked.

The male resembles the unique type of *Psychopsis infantilis* Dyar, which is much smaller, with slenderer body, also cell of hind wing longer (though hardly reaching $\frac{3}{4}$ way to outer margin). The latter is rubbed but the fringe appears not to be checkered, but solidly dark except toward anal angle. The female resembles the unique type of *Miacora adolescens* Dyar, but the latter has the st. line sinuous above, reaching the margin at Cu_2 and represented by a couple of oblique striae at anal angle; the stria before it does not anastomose at middle, the median line is thin and more continuous and the hind wing is much paler than the fore wing.

GIVIRA Walker

(*Hypopta* in large part of earlier authors; with
Lentagena, *Eugivira* etc.)

Antennae normally bipectinate (serrate in a few species); acc. cell small (fig. 108), R_4 and 5 stalked, and R_3 sometimes from their base; M_1 and M_2 very widely separated; 1st A and 2d A connected by a crossvein (rarely a mere fold) toward margin; hind wing sometimes with Sc and R connected a little before or beyond end of cell; R and M_1 stalked, usually strongly, and int. cell normal but varying much in size. In a few species the upper fork of base of M fades out, as in *Inguromorpha*, but these have Cu_2 and 2d A a normal distance apart, and R and M_1 of hind wing long-stalked.

This is much the largest genus of the family, with some 60 species, all American. The following genera are directly derived from it and lead up to the closely related Old-World family *Arbelidae* (*Lepidarbelidae*); in fact several of this group were originally described as *Arbelidae*.

1. Fore wing with the most prominent marks three parallel black stripes, in cells R_5 , M_2 , and obliquely across discal cell from costa *stypus*
Fore wing without conspicuous longitudinal stripes 2
2. Fore wing rusty orange,—in two shades on a cream base . . . *tigrata*
Fore wing dull brown or gray 3
3. Fore wing with base of inner margin white, but otherwise rather evenly and not contrastingly striate in two shades of brown . . 4
Fore wing with contrasting markings, usually black or dark gray spots or patches with pale outlines 5
4. Ground pale brown, at least in outer third; white on inner margin conspicuous, reaching almost up to 2d A *nais*
Ground dark wood-brown, the white stripe on inner margin weaker *neeceros*
5. Ground of fore wing rather evenly gray, with a few sharply defined black dots 6
Fore wing contrastingly marked all over 8
6. Fore wing with a black median spot above A as well as the am. spotting (which may rarely be obsolete) 7
Fore wing with two median black spots (at $2/5$ length of wing) at costa and in cell *nudaria*
7. A large squarish median black patch reaching from lower edge of cell to A and almost as long as wide *new species*
A small longitudinal bar along upper side of A; ground paler, and tending to be mottled *tristani*
8. Fore wing with the gray ground considerably suffused with white, particularly in antemedial area ($\sigma^{\text{♂}}$) or suffused with mottled gray, except subapically (♀) *perfida*
Fore wing with sharply defined pale-edge brown spots on a more even gray or light brown ground in all parts of wing 9
9. Disc of hind wing whitish, with contrasting fuscous pm. band on costal part of wing; fore wing with violet-gray ground, basal portion of cell Cu_1 almost wholly dark brown, and whitish patch below Cu_2 abruptly cut off by the brown anal spot . *juturna*
Hind wing wholly blackish, the darker markings not contrasting. 10
10. Ground brown, cell Cu_1 with three pale-edged transverse dark bars besides the minute one in apex *amanosa*

Ground violet gray; cell Cu_1 with only two marginal brown spots, the basal $2/3$ wholly of the gray ground; whitish patch below Cu_2 triangular, reaching anal angle *aroa*

Group I: *Male antennae serrate and fasciculate or subpectinate, less than twice as long as their bristles; ground of fore wing normally rather even gray (Lentagena)*

GIVIRA NUDARIA Schaus

Eugivira nudaria Schs., Jour. N. Y. Ent. Soc., 9, 75, 1901.

Lentagena nudara (sic) of Lep. Cat., 29, 26, 1933.

Figured: Seitz, 182: f4.

Antennae almost pectinate, the serrations though acute being almost twice as long as their coating of bristles; palpi short; fore wing without acc. cell, R_2 and 3 arising separately or shortly stalked from discal cell; int. cell very large, with its dorsal side closely parallel to Cu ; hind wing with Sc free, R and M_1 long-stalked, upper side of int. cell reduced to a very faint fold, as in *Inguromorpha*.

Ground decidedly mottled with whitish. Besides the two basal sets of spots some specimens show quite subordinate black spotting along the costa.

Jan. 25 (Bts.). Guatemala (Nat. Mus.) to British Guiana (Nat. Mus.) and southern Brazil (Cornell).

GIVIRA TRISTANI Schaus

Lentagena tristani Schs., Ann. Mag. Nat. Hist., (8) 7, 631, 1911; Dalla Torre, Lep. Cat., 29, 26, 1923; *Givira t.* Dyar in Seitz, 1280, 1937.

Figured: Seitz 182: f3.

Antennae moderately serrate; palpi minute; fore wing with R_4 and 5 stalked, R_{1-3} closely crowded; acc. cell absent and int. cell open by loss of M_{1+2} ; hind wing with Sc free, int. cell absent, base of M_3 extremely close to lower edge of cell; R and M_1 strongly stalked.

Extremely variable in size, amount of black spotting and degree of contrast in mottling of the ground, if the Nat. Mus. material is correctly associated. Supposed range from Mexico to Argentine (if *brunnea* Köh. is a synonym, as Dyar suspected).

Feb. 1 (Fried.) runty, abnormally marked and in bad condition.

There is a single specimen of another species of this group; with a large quadrate submedian spot and small st. spots, but too poor to make a type;—Jan. 30 (Bts.).

Group II: *Antennae of both sexes bipectinated, in male, and usually female, with the pectinations several times as long as their vestiture; pattern generally contrasting (Givira)*

GIVIRA NAIS Druce

Arbela nais Dr., Biol. Centr. Am., Lep. Het., 2, 450, 10: 1 ♂ 4 ♀, 1898: Dalla Torre and Strand, Lep. Cat., 28, 3, 1923; *Givira n.*, Dyar in Seitz, 1281, 1937.

Arbela naida Dyar, Proc. U. S. Nat. Mus., 44, 324, 1913; Lep. Cat., 28, 3, 1923; Dyar, l.c., pr. syn.

Hypopta albipuncta Schs., Proc. U. S. Nat. Mus., 59, 293, 1921; Dyar, l.c. pr. syn.

Also figured: Seitz, 182: f5.

Light wood brown, the base a little darker out to a vague blackish pm. shade from before apex to middle of inner margin; hind wing mouse gray, a little darker than fore wing. Inner margin of fore wing with a conspicuous white stripe, narrowing out about half way to outer margin, and lateral edge of tegulae also with a white stripe.

The type of *albipuncta* is smaller and the base out to the pm. fascia is filled with dark; also the discal lunule is white, not large but contrasting. It may turn out distinct.

This species and the next are transitional to *Lentagena*, having the plain coloring, and a narrower antenna than the residue.

Jan. 3, Feb. 1 (Bts.) Mar. 4 (Fried.) Lancetilla, Honduras (Bts.) Ranges north to Mexico.

GIVIRA NECREROS Dyar

Arbela necreros Dyar, Proc. U. S. Nat. Mus., 47, 350, 1914; D, T. & Str., Lep. Cat., 28, 3, 1923; *Givira n.*, Dyar in Seitz, 1281.

Figured: Seitz, 184: c5.

Exactly like the preceding, except for the much darker umber color and weak white streaks on tegulae and fore wing. Probably a mere color form of the preceding, but strongly dominant on the island.

Common (see diagram); described from the Canal Zone.

GIVIRA STYPUS spec. nov.

Female antennae broadly bipectinate; thorax with large bifurcated posterior tuft; abdomen with bifurcated dorsal tufts, the second one high; fore wing with R_4 and 5 not long-stalked, the rest free; int. cell rather small, but M_2 and 3 arising from its posterior edge; hind wing with a crossvein between Sc and R exactly opposite m_{5+6} . R and M_1

short-stalked; M_2 and 3 arising separately from the end of the small but distinct int. cell.

Head and thorax white, a little scaled with pale gray; head heavily shaded with gray; palpi gray above; scaling of antennae white. Thorax with a strong black median stripe, the metathorax and posterior side of the bifid tuft blackish, contrasting; abdomen blackish. Fore legs white with a little gray flecking; middle and hind femora and tibiae, including the long hair on the latter, mostly gray, a little whitish hair on tarsi. Fore wing with ground whitish but heavily shaded with brownish gray; only a little reticulation along margins. A large vague patch from M_2 to Cu_2 , and inner margin below A, brownish gray; veins toward costa infuscated, small black-brown streaks below R_3 and R_4 and a heavy one below R_5 , reaching $2/3$ way in to cell; a black-brown spot over forks of M_2 and 3 , connected to costa by an inwardly oblique stripe and to outer margin by a stripe in middle of cell M_2 ; a heavy black am. bar from costa obliquely out across cell, then extended out in heavy bars along Cu to its fork and along A almost to margin; the former connected by a black stripe to the spot over M_3 ; outer marginal reticulation forming black bars either side of veins M_3 to Cu_2 , making a broken marginal festoon. Hind wing fuscous. Beneath with markings of fore wing vague, but outer margin of both wings with the festoon formed of pairs of black bars at veins, for most of the margin; costa of hind wing mottled with whitish. Expanse 55 mm.

Type and two paratypes ♀; Barro Colorado Id., Panama, Jan. 14, 1935 and Dec. 28, 1934 (Bates) and Mar. 3, 1935 (Friedman). In M.C.Z.

What I take to be the male is much darker, dull wood brown, the body fuscous and hind wing black; the markings black, nearly lost in the dark ground, but so far as traceable like the female, only the longitudinal streaks consistently shorter and thicker. Expanse 33 mm. Jan. 26, 1935 (Bates).

This species is close to *invenusta* Schaus; but the latter is generally reticulate; and the st. markings in cells R_3 and R_4 take the form of two st. spots, each extended by a dash toward outer margin; an undescribed species from S. Brazil in the Nat. Mus. is nearer, but the details of the marginal pattern are different, and the black dashes are shorter and less regular.

GIVIRA JUTURNA Schaus

Dolecta juturna Schs., Proc. Zool. Soc. London, 1892, 328; Lep. Cat., 37;

Givira j. Dyar in Seitz, 1277.

Figured: Seitz, 182: d3.

Easily recognized by the clean-cut pattern of fore wing and contrasting, though diffuse, pale disc of hind wing.

Nov. 8 (Bts.) Mexico to Brazil.

GIVIRA AROA Schaus

Langsdorfia aroa Schs., Proc. Zool. Soc. London, 1894, 235; Lep. Cat., 26;
Givira a. Dyar in Seitz, 1277.

Figured: Seitz, 182: d2.

Bluish ash gray, with about 6 or 8 larger and many smaller black-brown spots, each finely pale edged, many pale veins, and a triangular pale area from Cu_2 to inner margin, beyond the median brown patch on inner margin. Hind wing dull fuscous with 4 or 5 rather distinct but not contrasting, paler-edged marginal spots. One specimen is smaller, with rounder and relatively smaller brown spots.

The deep brown pigment is unstable to moisture, and is likely to become yellow-brown on relaxing.

Nov. 22-Dec. 24 (Bts.); Jan. 20, 1936 (Wood- A.M.N.H.). Ranges to Peru and Venezuela.

GIVIRA AMANOSA Schaus

G. amanosa Schs., Ann. Mag. Nat. Hist., (8) 7, 629, 1911; Lep. Cat., 24.
Figured: Seitz, 183: e3.

Light wood brown, the dark spots more numerous, only a little darker, and their outlines only a little lighter. Hind wing with narrow pm. and st. pale lines, besides the usual marginal spotting.

Dec. 31 (Bts.) Mar. 2 (Fried.). Described from Costa Rica.

G. macrochir Schs. from Brazil (*Langsdorfia*) is close, but the disc of the hind wing is whitish.

GIVIRA TIGRATA Schaus

G. tigrata Schs., Ann. Mag. Nat. Hist., (8) 7, 634, 1911.
Figured: Seitz, 183: e5.

Male easily recognized by the rows of alternately lighter and darker tawny buff spots on a cream ground; hind wing pale orange, faintly reticulate. Supposed female with orange of fore wing more extensive, reducing the cream to narrow lines, and hind wing blackish, contrasting. Discal spot of fore wing a little accented, especially in female.

Males. Nov. 5 (Bts.), May 12 (Fried.), July 11 (Sc.). *Females* Jan. 4 (Bts.), May 3 (Fried.) Described from Costa Rica. The July

specimen is abnormal and may represent another species, but is not in good condition.

GIVIRA PERFIDA Schaus

G. perfida Schs., Proc. U. S. Nat. Mus., 59, 392, 1921.

Figured: Seitz, 183: h5 (unrecognizable).

Male easily recognized by the mottled appearance with considerable white in the ground; what I take to be the female is a little larger, merely mottled dark gray out to the pm. area, but with a conspicuous dark gray elliptical st. spot in the pale apex.

The types are a little larger and darker, but otherwise appear the same. *G. carisca* Schs. looks at first sight the same, but has a contrasting red-brown discal patch.

Common from Dec. to Mar. (see diagram). In the last week of Jan., 1936, the A.M.N.H. group had a heavy flight. Described from Guatemala.

LANGSDORFIA Hübner

Typical group (to which the present species belongs) with the space between Cu_2 and 2d A of fore wing extremely narrow (fig. 109), tending to crowd out 1st A and its connecting vein with 2d A; hind wing with R and M_1 separate and basally parallel. Markings mottled and confused, but with a group of am. silver spots. Antennae bipectinate in both sexes; palpi moderate, upturned, frenulum weak.

Some of the South American species intergrade with *Givira*, and I am inclined to transfer to *Givira* those that have R and M_1 of the hind wing stalked or even connate.

LANGSDORFIA LUNIFERA Dyar

L. lunifera Dyar, in Seitz, Macrolep. World, 6, 1282, 1937.

Figured: Seitz, 184: d4.

L. francki auct. in part.

Larval food: Dyar, l. c.

Dull brown, with paler shades, and an irregular darker subterminal shade. A larger silver triangle above A and a small one below it, but with the more costal silver spots of *L. francki* much reduced or lost. *L. francki* is figured in Seitz, 182: h2 ♂, 1 ♀, as *langsdorfi*.

Larva boring in pigeon pea, near the ground level (Dyar, after Moore).

Flight scattering, (see diagram). The grouping of records suggests that the main flight was in summer, and was missed, but Scrimshaw did not find it common in 1940. Mexico to Guiana.

INGUROMORPHA Henry Edwards

Close to *Langsdorfia*; fore wing (fig. 110) with acc. cell minute or absent, and both wings with int. cell absent, fore wing with R_4 and δ stalked; hind wing with Sc and R weakly connected $1/3$ way out on cell (i.e. R_1 traceable); R and M_1 rather parallel.

INGUROMORPHA POLYBIA Schaus

Langsdorfia polybia Schs., Proc. Zool Soc. London, 1892, 329.

Hypopta crassiplaga Schs., Proc. U. S. Nat. Mus., 29, 343, 1905; Lep. Cat., 22;

Dyar in Seitz, 1285, pr. syn.

Hypopta inguromorpha Schs., Proc. U. S. Nat. Mus., 29, 343, 1905; Lep. Cat.,

22; Dyar, l.c. pr. syn.

Figured: Seitz, 182: i5.

Blotchily mottled with pale brown and white, with fine but large-meshed black reticulation, gathering into a few black spots, of which a subapical black lunule and a dot at anal angle are most obvious; hind wing gray.

I. polybioides Schs. is closely similar, but the fore wing has a larger proportion of white, and the hind wing is dominantly whitish.

Jan. 8, Feb. 23 (Fried.), Feb. 9 (A.M.N.H.). Guatemala to Brazil.

PSYCHIDÆ

Male with head weak; mouth-parts reduced, including palpi, but antennae large and bipectinate, save in a few primitive species not known from the Neotropical. Body medium or long, soft; wings variable, usually with weak scaling, sometimes transparent; venation extremely variable, with base of M preserved and generally forked; 1st A preserved, anastomosing or running into 2d A; 3d A when distinct also running into 2d A, but often with free branches to inner margin; hind wing with Sc normally separate, but connected to cell, base of M forked. Legs weak, without obvious spurs.

Female, excepting a few primitive species not known from the Neotropical, reduced to a maggot-like creature, which normally never leaves the pupal shell, or only after the eggs are laid. Often without

recognizable appendages of any sort, but usually with a hair tuft at posterior end.

Eggs laid within the pupa shell; larva in a portable case with a large anterior opening through which the larva reaches out to feed or travel, and an inconspicuous posterior one through which the imago finally emerges. Neotropical species leaf-feeders.

There has never been a revision of the family in Central and northern South America;¹ specimens in collections are rare, no critical determinations have been made, and all records and specimens (save types of original descriptions) must be considered mere guesses.

Dyar reports three species from the Canal Zone, and a fourth is considered regional, and was taken by Bates at Lancetilla, Honduras. It agrees with *geyeri*, so far as our knowledge goes, but no critical comparison has been made.

1. Expanse over 25 mm.; fore wing elongate, half longer than hind wing (*Oiketeticus*) 2
 Expanse under 20 mm.; fore wing broad and rounded, but little longer than hind wing; brown species (*Platoeceticus*) 3
2. Fore wing shaded; brown with a contrasting pale oblique bar at end of cell *kirbyi* Guild. (Seitz, 169: e3, f. *platensis*, 4, f. *poeyi*)
 Both wings plain translucent fuscous. . *geyeri* Berg (Seitz, 169: e2)
3. Expanse 19 mm. Both wings with M_2 and $_3$ stalked; fore wing with R_2 out of stalk of R_{3+4} *aphidropa* Dyar
 Expanse 12 mm. Both wings with M_2 and $_3$ connate. *symmicta* Dyar

ZYGÆNIDÆ

Similar to the Eucleidae and neighboring families, but with fully developed tongue. Chaetosemas enormous, the two frequently meeting on the middorsal line, their bristles mixed with dense short scales; fore wing sometimes with all radial veins arising separately from the cell; hind wing with Sc and R various. Larva and cocoon as in the Megalopygidae, the cocoon in the type genus without a formed lid for emergence, but our genera presumably with the usual lid. Larva with the 16 normal legs only, the hooks uniordinal.

An attempt is often made to separate the American species as a separate family, Pyromorphidae, but some of our species are very close, even generically, to the more primitive Old-World Chalcosiinae. The true genus *Zygaena* is more distinct.

¹ Vazquez has just published the first instalment of a revision of the Mexican species in An. Inst. Biol. Mex., 12, 295, ff., including *Platoeceticus*, but not *Oiketeticus* as yet.

References

Besides the usual authorities the following references give revisions or critical definitions. For further bibliography see Lep. Cat., 71, cited. Earlier references generally confuse the family with the Euchromiidae.

Neumoegen and Dyar, Journ. N. Y. Ent. Soc., **2**, 63, 1894 (Nearctic species, as Anthroceridae).

Dyar, Proc. Ent. Soc. Wash., **6**, 322-329, 1902 (Nearctic species, and several larvae, as Pyromorphidae).

Fracker, Ill. Biol. Monog., **2** (1) 95, 1915 (larvae).

Mosher, Bull. Ill. State Lab. N. H., **12** (2) 44, 1916 (pupa of *Harrisina*).

Forbes, Cornell Mem., no. 68, 113 ff, 1924 (characterization of family and Nearctic species).

Hering, D. E. Z. Iris, **39**, 152-178, 1925 (revision of neotropical species).

Hering, Arch. Naturg., **88**, A (11), 1-93 (revision of oriental species).

Burgeff and Bryk, Lep. Cat., **33**, 1926 (*Zygaena* only) and **71**, 1936 (bibliography).

Key to Genera

1. Hind wing reduced, half the area of fore wing, with anal region reduced and 3d A, as well as M₁ (fig. 115) lost *Harrisina*
Hind wing ample, with all veins preserved *Malthaca* etc.¹

HARRISINA Packard

Fore wing with R₂ typically stalked, R₃ and ₄ stalked or united, R₅ free, widely separated. Larvae stout, slug-like, though with normal hooks on prolegs, social, commonly feeding in a regular rank, side by side; white or buff, with contrasting dark tubercles and sometimes transverse bands; on Vitaceae. The moths are black, often with more or less blue or green iridescence, and often with an orange collar (absent in the present species).

1. One radial lost, R₃ and ₄ being united *guatemalena*.
All veins present, R₃ and ₄ stalked 2
2. R₂ stalked, wings almost opaque *mexicana*
R₂ connate, wings translucent with black veins and a black longitudinal stripe *new species*

HARRISINA GUATEMALENA Druce

Aglaope guatemalena Dr., Biol. Centr. Am. Lep. Het., **1**, 40, 6: 10, 1884.

Also figured Seitz, 9: h5.

Oct. 16 (Bts.).

¹ Not yet known from the Canal Zone.

HARRISINA MEXICANA Schaus

H. mexicana Schs., Ent. Am., 5, 87, 1889.

Figures: Biol., 70: 11; Seitz, 9: h6.

Larva similar to the North American one which Dyar considered to belong to *H. texana*; cream white, with a brown transverse stripe on each segment, ending below in a thinner or broken lateral stripe on abdomen only, and represented by the brown cervical shield on prothorax (Nat. Mus.). Oct. 16 (Bts.). Mexico (type).

Fairchild took a single poor specimen of a third species with translucent, partly pale-scaled wings, with a stronger black longitudinal stripe as well as black veins. The Nat. Mus. has a similar specimen from Peru.

EUCLEIDÆ

(*Cochlidiidae*, *Limacodidae*)

Small stout moths with head retracted, tongue, ocelli and chaetosema absent; antennae various, palpi usually moderate, rarely long; 1st A preserved in both wings (figs. 116-121), which are short, thick and heavy-veined; fore wing with R_{3-5} normally stalked, sometimes R_2 also, Cu apparently quadrifid; hind wing with Sc connected to R by a cross vein or shortly fused, M_2 normally from near lower angle of cell, but the ldcv most often re-entrant. Base of M preserved in both wings, nearly central in fore wing (unlike Tortricidae), frequently forked. Legs stout and hairy, but usually with well developed spurs.

Egg thin and flat with few exceptions; larva characteristic, extremely stout, the head large but completely retractile in thorax, usually with only the mouth parts exposed, and lightly chitinized; prolegs absent but replaced by a series of strong suckers. Skin thick, often making the caterpillar prismatic, naked, hairy or spined, the major subdivisions of the family being defined on the armature of the caterpillar. Pupa in a short-ellipsoidal cocoon with a trap door at one end; extremely stout, hardly chitinized, with the appendages almost free and the dorsum finely clothed with many spinules.

A good sized family, found in most parts of the world, but only represented by two inconspicuous species in Europe. The larvae feed on a great variety of deciduous woody plants, very few on herbs, and none so far as I know on conifers. Many have poison-spines. The basic subdivision is on the larval structure (see Dyar, Jour. N. Y. Ent. Soc., 7, 234-244, 1899), so the placing of the many tropical genera

with unknown larva is tentative. On the whole the types with spined larvae are tropical, those with naked larvae temperate, but there are many exceptions.

References

Dyar: A series of papers on life histories in Journ. N. Y. Ent. Soc., culminating in the general account in vol. vii, pp. 234 ff.

Dyar: Journ. N. Y. Ent. Soc., **6**, 233, 1898 (classified catalogue).

Tutt: British Lepidoptera, **1**, 360-383, 1899 (a general account with much biology).

Dyar: Proc. Ent. Soc. Wash., **4**, 422, 1901 (key to Sibine—containing only 5 species!)

Dyar: Proc. U. S. Nat. Mus., **29**, 359 ff. 1906 (Key to genera and catalogue for the two Americas).

Dyar: Proc. U. S. Nat. Mus., **51**, 33, 1917 (partial key to Euclea).

Van Eecke, Lep. Cat., **32**, 1-80, 1925 (Bibliography).

Dyar: Ins. Ins. Men., **14**, 13 ff., 1926 (Keys to species of *Parasa*, *Euclea*, *Talima*, *Perola*, *Euphobetron*, *Isochaetes*).

Dyar in Seitz: Macrolep. World, **6**, 1104-1139, 1935-6 (all American species with many colored figures on pls. 164-166, and 167, not yet published; keys to *Sibine*, *Parasa*, *Euclea*, and *Natada*).

Hering and Hopp, D.E.Z. Iris, **41**, 1-9, 173-186, 1927 (with keys to *Tanadema*, *Prolimacodes*).

Hampson's Key to the Indian genera in the Fauna of British India, Moths, **1**, 371-373 and Hering's Key to the Indo-australian ones in Seitz, **10**, 668 ff. have also been considered in drawing up this key.

Key to Genera

1. Middle discocellular vein of fore wing sharply angled far below its middle, the simple base of M attached to the angle; male antenna simple, at least at base and apex. 2
- Middle discocellular angled about at middle or above unless base of M is forked or obsolete; male antenna strongly pectinate, at least on basal half. 6
2. Male antenna pectinate on middle portion, base and apex serrate, palpi upturned above vertex, long, the third segment tapering; frontal tuft in the form of a ridge between antennae. 3
- Male antenna wholly serrate or simple; head tufted on front, well below antennae; palpi obliquely upturned about to middle of front, the third segment not long. 4

3. Vertical tuft long, hind as well as fore wing with upper sector of mdcv. twice as long as lower *Semyra* (present spp.)
Vertical tuft slight; hind wing with sectors of mdcv. subequal.
Euphobetron (our species)
4. Male antenna simple, laminate; fore wing with inner margin subsinuate; small, slender, plain species *Vipsophobetron*
Male antenna serrate below; stouter species with an oblique postmedial line or abrupt change of color 5
5. Fore wing with a large triangular brown patch, extended along costa to base and more or less distinctly outlined with silver.
Prolimacodes
Fore wing bisected by a line or change of color from middle of costa to near anal angle *Dichromapteryx*
6. Media of fore wing forked, enclosing an intercalated cell, or rarely simple (fig. 113), but then with the discocellular *acutely* angled at its attachment; male antenna with pectinations becoming abruptly shorter about at middle 7
Median of fore wing simple when present (fig. 114), attached near the middle of the obtusely angled mdcv; when obscure with male antenna strongly pectinate to $\frac{3}{4}$ 8
7. Inner margin of fore wing deeply sinuate; ground rich brown, with ruffled scales giving a watered effect, and usually with minute silver or pale gold dots *Sibine*
Inner margin of fore wing not sinuate, the outer half not noticeably concave *Euclea*
8. Male antennae with pectinations abruptly decreasing or ending about half way out 9
Male antennae pectinate practically to apex 10
9. Male antennae unipectinate *Tanadema*
Male antennae bipectinate *Talima*
10. Fore wing with Cu_1 and 2 stalked (fig. 119) 11
Fore wing with Cu_1 and Cu_2 arising separately 12
11. Hind tibia with upper spurs absent 14
Hind tibia with two pairs of spurs *Perola*
12. Hind tibiae with upper spurs absent *Sisyrosea*
Hind tibia with two pairs of spurs 13
13. Second segment of palpus with a triangular tuft in front toward apex *Natada*
Palpus with vestiture of second segment rounded off, not widened at tip *Narosopsis, Euprosterna*

14. Palpi falling short of frontal tuft; angle of dev of hind wing strongly asymmetrical. *Palaeophobetron*
 Palpi exceeding frontal tuft, mdev of hind wing symmetrical. *Epiperola*

SIBINE Herrich-Schaeffer

(Empretia Clemens, with Episibine Dyar)

Like *Euclea* except for the key-difference. The pattern is distinctive, rich brown, rarely fuscous or somewhat golden; with limited silver or pale gold dots subapically at costa and antemedially below cell; the scaling patchily set at different angles, giving an effect of varied shades of color.

Larva (the Saddle-Back Caterpillar) similar to *Euclea*, but with the middle third of the body bearing rudimentary spines or none and usually with a brilliant green or green and brown saddle-mark. The species are very close and the present key will not always hold; characters of penis and anellus are more trustworthy. The first notes on genitalic characters were published in Proc. Ent. Soc. Wash., 4, 422, 1901; but little more has been put on record.

1. A metallic line (violet in male, brassy in female) from near costa subterminally, obliquely and irregularly in to inner margin near base; frequently interrupted or reduced to scattered scales in middle portion. *nesea*
 Metallic spots small to minute and generally rounded, never violet. 2
2. Hind wing contrastingly much paler than fore wing, dull luteous, with red-brown only in vicinity of fold. *ophelians*
 Hind wing brown, frequently almost as dark as fore wing, when paler with considerable brown shading. 3
3. A strong coppery or brassy gloss underlying the brown scaling of fore wing, always visible toward apex and below cell, and becoming dominant when specimen is rubbed. *hyperoche*
 All purple or brown, even in rubbed specimens. 4
4. Wings short, bluntly rounded over apex. *apicalis*
 Wings long, the apex extended. 5
5. Ground chocolate brown, the silver spotting slight. *affinis*
 Ground purplish fuscous, the silver dots strong. *joyceans*

SIBINE NESEA Stoll

Bombyx nesea Stoll in Cr., Pap. Exot., 4, 31, 305: C, 1781.

Phalaena vidua Sepp, Surin. Vlinders, pl. 6, 1852.¹

Also figured: Seitz, 164: g1 (not e).

Life History: Sepp, l.c.; discussed by Dyar in Ann. N. Y. Acad. Sci., 8, 216, 1894; Can. Ent., 29, 77, 1897; Seitz, 1111, 1935.

Easily distinguished from all others by the silver being in the form of a sinuous line.

Larva similar to the North American *S. stimulea*, gray with a large squarish green saddle, outlined in yellow, but without the central brown spot of *stimulea*; spines very long, the subdorsal ones especially so just before and beyond the saddle, but none between,—the lateral ones regular; on *Convolvulus*. Cocoon double, the outer layer loose, the inner very smooth and hard.

Dec. 5–28 (Bts.), June 21, July 5 (Scr.). Guatemala to Amazons and Bolivia.

SIBINE OPHELIANS Dyar

S. ophelians Dyar, Jour. Wash. Acad. Sci., 17, 546, 1927; Seitz, p. 1110, 1935.

Wings long, the matt band beyond the cell rather narrow. Hind wing even paler than in *joyceans*, and fore wing warmer brown.

Nov. 29 (Bts.). Guatemala to Colombia.

SIBINE HYPEROCHE Dognin

S. hyperoche Dgn., Het. nouv. Am. Sud, 8, 98, 1914.

Larva: Dyar, Ins. Ins. Men., 13, 218, 1925.

The brassy color is paler than in the type in the National Museum, and in one rubbed female covers most of the surface of the fore wing. In the male, which is fresh, it is only visible in small patches. Silver dots large and brassy. Hind wing even dark brown, with slightly darker veins.

Larva almost like *stimulea*, the saddle-spot practically confined to segment 8, the posterior yellow dots partly fused.

Jan., June (Bts. Fried.). Also in Nat. Mus., reared by Zetek. Ranges to Ecuador.

Sibine apicalis Dyar (Seitz, 164: g2) is known from Mexico to Costa Rica, and should be found in the Zone. The silver spots, while large and conspicuous, are much smaller than in the Seitz figure.

¹ See footnote on p. 246 of previous part.

SIBINE AFFINIS Möschler

S. affinis Msch., Verh. Zool.-Bot. Ges. Wien, **32**, 353, 1883; Dyar, Proc. U. S. Nat. Mus., **29**, 362, 1906.

S. horrida Dyar, Proc. U. S. Nat. Mus., **29**, 362, 1906; Seitz, 1109, as possibly syn. *affinis*.

S. horrida var. *nitens* Dyar, Proc. U. S. Nat. Mus., **29**, 362, 1906; Seitz, 1110 as possibly syn. *fusca* Stoll.

? *Bombyx fusca* Stoll in Cr., Pap. Exot., **4**, 37, 307: G, 1781.

Also figured (nominally): Seitz, 164: d6 (This figure shows none of the distinctive features of this species).

The moth is rather distinctive in having broad paler veins on the dark brown hind wing. It shows a decided crimson gloss, especially in the female.

Nov. 3-Jan. 27. Described from Guiana.

SIBINE JOYCEANS Dyar

S. joyceans Dyar., Jour. Wash. Acad. Sci., **17**, 545, 1927.

Figured: Seitz, 164: e1 (unrecognizable).

Larva: Dyar in Seitz, 1109.

Ground grayer than any of the preceding, almost coal-black, with the lines of water-banding narrow. Disc of hind wing decidedly paler; silver dots stronger than in *affinis*, the two pm. dots in contact. A female which presumably goes with this has the lower pm. dot elongated into a bar, as in *violans* Dyar.

Nov. 26, Dec. 13 ♂; Dec. 5 supposed ♀. Described from Panama.

Larva on *Lansium americanum* and other plants. Light green down to the sides, lateral horns short, absent on the saddle; light with black tip and black basal ring.

EUCLEA Hübner

(*Parasa* in part, *Metraga* in part)

Antenna broadly pectinate on basal 2/5, then practically simple, palpi obliquely upturned to middle of front; wings (fig. 113) broad, well rounded, the inner margin strongly convex toward base, then less convex or nearly straight to anal angle; R_4 strongly curved and approximate to Sc, R_2 free or short-stalked on R_{3-5} ; base of M deeply forked, enclosing an intercalary cell (rarely absent) in fore wing, in hind wing M simply forked, the upper fork running rather squarely across to $R+M_1$, the lower running out into M_2 . Hind tibia with end-spurs only.

The larva is stout, somewhat prismatic, with the subdorsal and lateral spines both long, the subdorsal ones often irregular in length but without the long central gap of Sibine. They are brilliantly striped or spotted. For (colored) figures of North American species see Jour. N. Y. Ent. Soc., 5, 10, pl. 2 (*indetermina*) 57, pl. 3 (*delphinii*) 61, pl. 4 (*chloris*).

Dyar has published several keys to the species, the latest in Seitz, p. 1115. The ones in Proc. U. S. Nat. Mus., 51, 33 and Ins. Ins. Men., 14, 77, are fragmentary, but develop some additional characters.

1. Fore wing marked with green 2
 Fore wing not marked with green, usually with silver spots or lines
 subapically near costa, and in fold antemedially (as in Sibine) . . 3
2. Chocolate brown with a warmer red-brown patch beyond the am.
 green spot, which is normally larger, extending above the white
 dot on 1st A almost up to Cu₂ *cippus*
 Gray-brown without any discolorous patch in fold, the green spots
 practically enclosed between the two white dots; scaling more
 erect *norba*
3. Apical area yellow, with contrasting brown veins, no silver
 *Talima aurora*
 Apical area concolorous, silver marks present, faint only in *E.*
 plugma, which is dark fuscous 4
4. Ground light brown with a yellow patch in fold; am. silver in the
 form of a line which is oblique outward and meets 2dA more than
 2/3 way to margin *distrakens*
 Am. silver line perpendicular to 2d A or oblique in, meeting 2d A
 less than 2/3 way out 5
5. Am. silver line very faint, linear, and crossing 2d A to inner margin
 or nearly; ground all fuscous *plugma*
 Am. silver line irregular, wholly above the contrasting pale line on
 2d A 6
6. Scaling decumbent, veins very heavily overlaid with strap-shaped
 ochre scales, the general surface also heavily dusted with ochre;
 st. silver a thick curved stripe *pallicolor*
 Considerable of scaling erect, especially in male; ground deeper
 brown, the ochre scaling limited to vicinity of veins and not ex-
 tensive; st. silver of separate dots or irregularly fused 7
7. St. silver of several conspicuous dots; am. silver a deeply bisinuate
 line, curving well in below middle of fold area *buseki*
 St. silver obscure, am. silver in the form of a single blunt tooth
 concave above and below *trichathdota*

EUCLEA CIPPUS Cramer

Bombyx cippus Cr., Pap.-Exot., 1, 84, 53: E, 1775.

Also figured: Seitz, 164: k4.

(Not *Bombyx cippus* Smith & Abbot. Nat. Hist. Lep. Ins. Georgia, 2, pl. 13, 1797, etc., not *E. monitor* Pack., Proc. Ent. Soc. Phil., 3, 337, 1864; nor *Limacodes querceti* H.-S., Lep. Exot., fig. 174, 1853; which are all *E. delphinii* of North America).

E. delphinii Bdv. is close, but is larger and rounder-winged, with more erect scaling and the larger spots more edged with white.

The first and last green spots are always distinct, the intermediate ones vary from an irregular and broken line to mere points on the veins. The outer boundary of the am. green spot is more oblique in the female than the male.

Two flight periods (see diagram): Oct. 27, Dec. 5, 6, 8, 20, 23, May 17, 25, 31, June 21, July 21. Mexico to Paraguay.

EUCLEA NORBA Druce

Sibine norba Dr., Biol. Centr. Am. Lep. Het., 1, 211, 22: 15, 1887.

Also figured: Seitz, 164: k6.

The even dull color will distinguish this species from *cippus*. The green spots are small to minute, the two end one often only a little larger; d.d. a mere black point.

Common (see diagram). Mexico to Ecuador.

EUCLEA DISTRAHENS Dyar

E. distrahens Dyar, Proc. U. S. Nat. Mus., 42, 94, 1912.

Figured: Seitz, 164: 16 (much too gray, unrecognizable).

Easily recognized from the other local species by the patchy buff and light brown pattern, with a single yellow spot in fold. This species looks more like Seitz' figure of *E. baranda* Schs. (165: a4) but the latter is darker and warmer brown; *E. diversa* Dr. (*verierux* Dyar) which Bates took in Honduras, is closely similar in coloring, but the am. silver line is zigzag and the st. one continuous.

Nov. 17, Dec. 7, Feb. 2 (Bts.). Described from Costa Rica.

EUCLEA BUSCKI Dyar

E. buscki Dyar, Proc. U. S. Nat. Mus., 42, 95, 1912.

Figured: Seitz, 165: a5.

This species in color and scaling resembles *cippus* without the green; there is some strap-like overscaling on the veins, especially post-medially opposite lower angle of cell. Am. silver mark a zig-zag, wholly above 2d A, st. series of spots, usually 4 or 5, offset out at M_1 . Female lighter, the velvety dark areas more contrasting, strap like scales more extensive postmedially and st. silver nearly continuous.

Common (see diagram). Originally described from the Canal Zone.

Dyar also reported *E. trichathdota* Dy. from the Canal Zone, but there is no longer a specimen under that name at the National Museum.

EUCLEA PLUGMA Sepp

Phalaena plugma Sepp, Surin. Vlinders, 2, pl. 84, 1853?

Metraga perplexa Walker, List Lep. Ins. Br. Mus., 5, 1129, 1855; Dyar in Seitz, 1117, 1937 (syn. pr.).

E. zygia Dr., Biol. Centr. Am. Lep. Het., 1, 216, 23: 7, 1887; *Metraga z. Dyar*, Proc. U. S. Nat. Mus., 29, 370, 1906; Dyar in Seitz, l.c. (syn. *plugma*).

♀ *E. chiriquensis* Schs., Journ. N. Y. Ent. Soc., 8, 231, 1901; syn. *perplexa* Dyar l.c.

Metraga colle Dyar, Proc. U. S. Nat. Mus., 43, 95, 1912; syn. *perplexa* Dyar, l.c.

(*Neomiresa rufa* Btl., Tr. Ent. Soc. London, 1878, 74, is listed by Dyar and Lep. Cat. as a synonym of *perplexa*, but in Seitz, 1118 is again considered distinct.)

Early stages: Sepp, l.c.; Stoll in Cr., Pap. Exot., Suppl., 102, 21: 4, 4G, 1791 (as *aterea*).

Dark gray-brown, heavily water-lined and shaded with blackish, the am. silver line very fine, obscure, running to inner margin, crossing a rather similar line along 2d A almost at right angles.

Larva green with bluish dorsum and red middorsal stripe, the sub-dorsal spines at both ends rather longer than the middle ones; on banana, cassava, and doubtless a general feeder. Sepp identified the larva figured by Stoll as *aterea* as his *plugma*; the moth of Stoll's *aterea* is of course the well known geometer.

Nov. 30, Dec. 5, 8 (Bts.), May 8, 17 (Fried.). Mexico to Guiana.

EUCLEA PALLICOLOR Dyar

E. pallicolor Dy., Proc. U. S. Nat. Mus., 29, 368, 1906.

Figured: Seitz, 164: 11 (unrecognizable).

Dyar's unique type is almost illegible, but I think this species with little doubt; the smooth appearance, heavy st. silver bar and heavy scaling with yellow are distinctive. Hind wing fuscous, contrasting.

Jan. 8 (Fried.) 1 ♀. French Guiana (type ♂).

TALIMA Walker (with *Metraga* in part)

Similar to *Euclea*, palpi perhaps a little weaker; intercalated cell of fore wing absent (fig. 114), the fork of hind wing even more asymmetrical than in *Euclea*. Pattern simpler, the normal species merely with a pm. line. But there is a transitional group to which *straminea* belongs, with intercalated cell preserved, narrower wings, and ground toward apex ochre yellow, crossed by brown veins. It includes *aurora* Dyar, reported by Dyar from the Canal Zone as *straminea*, and also *ingenuor* Dyar from Guiana and *straminea* Schs. from Mexico.

Dyar has published a key to this group (including the *straminea* group and the species formerly in *Metraga*) in *Ins. Ins. Men.*, 14, 83.

Larva apparently unknown, presumably like *Euclea*.

1. Fore wing with a yellow patch or suffused with tawny yellow at apex, and more or less obvious yellow base, crossed by brown veins *aurora*
Fore wing rather evenly brown or buff, with a pm. line parallel to outer margin 2
2. Hind wing of male buff with a brown patch at anal angle, of female more or less suffused, but with the disc pale *emilia*
Hind wing mostly red-brown, approximately concolorous with fore wing in male *rubicolor*

TALIMA RUBICOLOR Dyar

Metraga rubicolor Dyar, *Proc. U. S. Nat. Mus.*, 29, 370, 1905.

Sisyrosea parva Dyar, *Proc. U. S. Nat. Mus.*, 29, 376, 1905; *Euclea p.* Dyar, *Proc. Ent. Soc. Wash.*, 13, 106, 1911; Seitz, p. 1119, 1937, pr. syn.

♀ *Sisyrosea aphasis* Dyar, *Proc. U. S. Nat. Mus.*, 47, 250, 1914; *Ins. Ins. Men.*, 14, 83, as syn. *E. phara* Dr.; Seitz, p. 1119, as syn. *rubicolor*.

Figured: Seitz, 165: b8.

Male fore wing tawny to red brown, with darker, purple-iridescent inner margin below fold and border beyond the slightly darker pm. line. Female much lighter and duller. Light specimens show dark veins and a dark discal dot or lunule. Hind wing of male concolorous, yellow toward base and costa, of female fuscous.

Common (see diagram). Ranges to Nicaragua and Venezuela.

TALIMA EMILIA Dyar

Metraga emilia Dyar, Proc. Ent. Soc. Wash., **10**, 50, 1908; *Talima e.* Dyar, Ins. Ins. Men., **14**, 84, 1926.

? ♀ *T. varians* Dyar, Jour. Wash. Acad. Sci., **17**, 548, 1927; Seitz, 165: b6.

Similar to the last, sex for sex, except for the dominantly light hind wings.

May 6 (Fried.). Guatemala to Amazons.

TANADEMA Dyar

Like *Talima* except for the unipectinate male antennae. A female of *T. mas* in the present lot shows that the female antennae are simple.

Hering and Hopp have published a key to the species in D.E.Z. Iris., **41**, 178, 1927.

TANADEMA MAS Dyar

T. mas Dyar, Proc. U. S. Nat. Mus., **29**, 392, 1906.

Figured: Seitz, 166: i6.

Apex marked, with a white dot before it; brown, with broken and blurred blackish st. line, and pm. from cell to inner margin. Female much larger with the same marks.

Oct. to April (see diagram). Described from the Guianas.

SEMYRA Walker

Palpi upturned, the third segment triangular with a sharp apex, reaching up about to top of eye (*bella*) or well above vertex (*irena*); male antennae moderately pectinate, tapering to a thick but nearly simple base (except in *S. cardia* from Brazil), as well as the short simple apex; fore wing (fig. 118) oblong, with rather straight costa and short outer margin; R_1 nearly straight, R_2 stalked on R_{3+4} , R_5 free; base of M simple; hind wing with upper segment of mdcv. much longer than lower, though more transverse. Hind tibia with all spurs. Pattern complicated, of gray and two shades of brown, usually with the two silver marks of Euclea, but with the submedian one much nearer the base and the outer one far toward apex.

The larva is uncertain. Sepp figures a larva of the *Prolimacodes* type (*S. gibbosa*, Surin, Vlinders pl. 129, Hering and Hopp, D.E.Z. Iris, **41**, 173, 1927), naked, with a subdorsal keel rising to a sharp crest at middle; but the imaginal appearance suggests a close connec-

tion with *Euclea*. Dyar accepts the Sepp larva and places the genus next to *Prolimacodes* (Seitz, p. 1135).

Since the species are close and no key has been published it seems advisable to present the following one, including all the recognized species, except *eucharista* Dyar, based on a single female from Brazil. All have been examined except *S. gibbosa*.

Key to Neotropical Species

1. A black patch or wedge before subterminal line at inner margin . . . 2
 No such patch, the lower outer part of the wing beyond the pm.
 line only vaguely shaded 5
2. Base contrastingly darker than the outer half, with a straight
 boundary at middle (not seen) *gibbosa*
 Ground color not darkened on basal half 3
3. Male antenna pectinate to base; a silvery zigzag st. line from fold
 to inner margin and a strong yellow dash below Cu; hind wing
 almost all cream white *cardia*
 Male with base of antenna serrate or simple; st. line near inner
 margin nearly even, oblique and only faintly silvery; hind wing
 about half dark 4
4. Umber brown, the black shade before subterminal on inner quarter
 of wing a solid block, no yellow dash below Cu. (type only seen)
 *phrygia*
 Red-brown, with separate black wedges at Cu and A; a small ante-
 medial yellow dash below Cu as well as the spot in fold *paula*
5. No apical spot, no crescent, no silver; though with the golden
 iridescence beyond the am. line in fold present and pm. line as
 in *irena* *mariae*¹
 With a blackish apical patch (obscure in *veterna*), crescent (a mere
 dot in *veterna*) and strong silver in am. line 6
6. Very small and blackish, but apical spot, crescent, golden am.
 patch etc. traceable *veterna*
 Well-marked 7
7. Am. silver in the form of a group of spots, sometimes fused into
 a dentate line 8
 A continuous fine am. line from below Cu to above inner margin . 14

¹ *Euphobetron natadooides* Dyar approaches this, but has a distinct bent st. line over apex preceded by heavy blackish, a single change of color toward inner margin to represent the am. line, and pm. present near inner margin.

8. Ground of flat tints except costo-apically; base to pm. line evenly umber brown and outer 2/5 even wood brown; apical spot large, incorporating the usual wedge below it and touching the crescent, which is simple and thick; silver spot with upper half thick, but deeply bidentate. *zinie*
 More mottled, usually with gray shades and golden patches; apical patch weaker, upper part of silver mark tending to break into spots. 9
9. Pm. line crossing costal part of wing at end of cell, with a quadrate blackish patch beyond it above basal end of the st. crescent¹. 10
 Pm. line offset and much further out than dorsal half of line when distinct, often incorporating the st. brown patch. 12
10. Crescent replaced by a large st. patch over 2 cells, followed by two large wedges below (♂ type only seen). *ornata*
 St. mark smaller, curving only a little below its proper cell, and with one small spot or none below it. 11
11. All violet gray and blackish; fore wings longer and hind wings small. *lucilla*
 Much dark wood brown, pm. line distinct to inner margin; wing form normal. *irena*
12. Apical spot strongly golden-iridescent, the yellow beyond am. line conspicuous (Lesser Antilles). *coarctata*
 Apical spot blackish; generally with less tawny yellow. 13
13. With much violet-gray (Central America). *finita*
 With considerable rich brown (Paraguay). *erna*
 Very small, dull, spotty blackish on wood brown (Argentina).
frances
14. Area in fold beyond antemedial line umber. *distincta*
 This area tawny with shot-silk effect. *bella*

SEMYRA BELLA Herrich-Schaeffer

Limacodes bella H.-S., Aussereur. Schm., fig. 181, 1854. (as *Euryda* in text).
Semyra gladys Dyar, Proc. U. S. Nat. Mus., 47, 249, 1914; Seitz, Macrolep. World, p. 1136, 1937, syn. *bella*.

Also to be figured: Seitz, 167: c.

Brown, shaded with violet gray, the dark markings rich brown in special lighting, but in a perpendicular light with only the patch in fold beyond the fine am. line brilliantly coppery.

¹ The deep brown st. marking mostly in cell M1.

Feb. 24, Mar. 14, Apr. 3, 7 (Fried.). Described from Bahia (*bella*) and Panama (*gladys*).

SEMYRA DISTINCTA Möschler

Eulimacodes distincta Msch., Verh. z.-b. Ges. Wien, **27**, 672, 1878.

Darker than the preceding, the violet-gray shades a little more prominent, and the brown patches chocolate brown, mostly without iridescence, only a little deep copper on the spot beyond the am. line.

Taken in series (see diagram). Ranges from Mexico to Guiana. This may be only a color form of *bella*.

SEMYRA IRENA Dyar

S. irena Dyar, Proc. U. S. Nat. Mus., **29**, 373, 1906.

To be figured: Seitz, 167: c.

The area beyond the am. silver is narrower and the copper glow only appears at a very favorable angle.

Oct. 31, Nov. 26, Dec., May 10, 12 (Bts. Fried.). Amazons (Cornell). Described from British Guiana and S. Brazil.

EUPHOBETRON Dyar

Structure exactly like *Semyra* except as stated in key. Pattern more confused, suggesting *Phobetron* at first glance,—but there is no structural resemblance to *Phobetron*.

E. cupreitincta is the genotype. Other species now standing in *Euphobetron* have normal antennae.

EUPHOBETRON CUPREITINCTA Dyar

E. cupreitincta Dyar, Proc. U. S. Nat. Mus., **29**, 387, 1906.

To be figured: Seitz, 167: h.

Basal three fifths brown mixed with blue-gray, outer 2/5 a complex mixture of luteous, brown, blue-gray and black, with a black triangle before the middle of the waved blackish st. line.

Series (see diagram). Guatemala to Amazons.

EUPROSTERNA Dyar

Like *Natada* except for the smoothly upturned palpi. Venation fig. 116.

1. Pm. line straight, dark and pale, oblique; st. line obsolete. *elaeasa*
 Pm. line obsolete; st. straight, dark followed by a pale terminal
 area, running onto outer margin above anal angle. *caria*

EUPROSTERNA CARIA Druce

Perola caria Dr., Biol. Centr. Am., Lep. Het., **1**, 219, 23: 13, 1887.

Natada c. Dyar, Proc. U. S. Nat. Mus., **29**, 377, 1906; van Eecke, Lep. Cat., **32**, 51; Seitz, 1124.

Dull mouse gray with vague dark discal dot and straight st. band, the margin much paler. Palpus with a dense black mass of scales near tip of second segment below, homologous with the projecting tuft of *Natada*.

Nov. 23, Dec. 5, 6 (Bts.). Described from Chiriqui.

EUPROSTERNA ELAEASA Dyar

Perola elaea Dr., Biol. Centr. Am., Lep. Het., **1**, 219, 1887, in part, not type.

E. elaeasa Dyar, Proc. U. S. Nat. Mus., **29**, 377, 1906.

Natada avellana Dgn., Het. Nouv. Am. Sud., **1**, 41, 1910; Seitz, 1122, pr. syn. Figured: Seitz, 165: f1.

Scaling with water-banding like the North American *Sisyrosea textula*, and many tropical *Natadas*. Fuscous, the straight pm. line dull black followed by whitish.

Eight specimens in Oct. and scattering till end of Jan. (see diagram). Mexico to Guiana.

NAROSOPSIS Dyar

This genus is not clearly separable from *Euprosterina* (with which Dyar does not compare it). Fore wing (fig. 117) with R_2 strongly stalked (unlike *E. caria*) and mdev only moderately curved, with faint traces of a forked base of M attached to it (unlike *E. elaeasa*).

NAROSOPSIS LEUCOSPILA Dyar

N. leucospila Dyar, Proc. U. S. Nat. Mus., **42**, 96, 1912.

N. vynia Schs., Proc. U. S. Nat. Mus., **57**, 148, 1920; Dyar in Seitz, 1129, pr. syn.

Straw yellow vaguely banded with reddish buff, a st. band most distinct, with a black dot in discal fold before it and sometimes a few other black points. Hind wing below with black d.d. conspicuous.

Euprosterina notula Dgn. is similar, but much darker, dull buff.

Oct. 11, 25 Dec. (Bts.) May 7, 20 (Fried.). Described from Central America.

NATADA Walker

Frontal tuft somewhat low on the front; palpi clavate, male antennae rather evenly pectinate to near the tip, only tapering distinctly on the last dozen joints, of female simple. Legs with all spurs. Fore wing (figs. 120, 121) with R_5 free or practically so, R_2 various, base of M normally simple, to the angle of the m_{dcv} , sometimes obsolete, and in that case with m_{dcv} of various types; Hind wing with Sc anastomosing with cell, R and M_1 connate or nearly so, immediately diverging; m_{dcv} bent far below its middle. There are two principal groups in the New World, one close to *Sisyrosea* (fig. 121), the other more distinct (fig. 120), but it is probable that neither is truly congeneric with the typical (Old World) group.

Larva of *N. nasoui* (belonging to the *fusca* group) very stout, especially in front of middle, with the usual subdorsal and lateral rows of spines reduced to bristle-bearing warts; on oak and other Amentiferae. (Dyar, Jour. N. Y. Ent. Soc., 7, 61 ff., pl. 1). The larva of the other group may well be as in *Sisyrosea*, very flat, with the lateral spines long and the subdorsals minute. Some oriental larvae are more like *Euclea*,—*N. nararia* is injurious to tea.

1. Fore wing with R_2 and R_5 free (fig. 120); scaling smooth without bands of watering 2
 Fore wing with R_2 and sometimes R_5 , stalked on R_{3+4} (fig. 121); fore wing without lines, but with transverse watering 4
2. Ordinary lines straight, converging toward costa 3
 Outer line curved, inner blurred and obscure *nindla*
3. Smaller, subterminal line with a strong dark brown shade before it, but only the pale ground beyond *subpectinata*
 Larger; st. line with a distinct mouse gray shade beyond, but only a slight darkening of the ground before *fusca*
4. Head and dorsum of thorax and base of abdomen orange, contrasting with wings *incandescens*
 Thorax and dorsum of abdomen concolorous; scape with a white tuft below 5
5. Water-lining dense and regular; head concolorous, fore wing with a blackish st. spot near middle *dogmini*
 Water-lining of about five ridges, the three basal well marked; head light tawny; no blackish st. spot *michorta*

1, R_2 free; scaling smooth

NATADA FUSCA Druce

Trabala ? fusca Dr., Biol. Centr. Am., Lep. Het., **1**, 207, 22: 11, 1887.

Perola salta Dr., Ann. Mag. Nat. Hist. (7) **5**, 512, 1900; Dyar in Seitz, *Macrolep. World*, 1124, pr. syn.

Ground varying from luteous to mouse gray, the ordinary lines with the dark elements prominent in the first, the pale in the second case; converging and almost meeting at costa before apex.

Abundant (see diagram). Mexico to Venezuela.

NATADA SUBPECTINATA Dyar

N. subpectinata Dyar, Proc. U. S. Nat. Mus., **29**, 381, 1905.

N. urichia Schs., Proc. Ent. Soc. Wash., **26**, 180, 1924; Dyar in Seitz, 1124, pr. syn.

Biology: Freeman, Rept. Dept. Agr. T'dad, & Tobago, 1924, 22, 1925 (Rev. Applied Ent., **13**, 493, 1925).

Much smaller than *N. fusca*, the fore wing wholly of reddish tints. Base of M lost, represented by a faint fold, well below middle of m₁cv.

N. nucca Dgn. is similar but has much finer pure black lines, the st. somewhat sinuous. The present specimens are slightly abnormal, having more nearly the color of *daona* (Seitz, 165: g3).

Dec. 5, 9, 21 (Bts.) Feb. 7 (Fried.) Dutch Guiana (type).

NATADA NINDLA Dyar

N. nindla Dyar, Proc. U. S. Nat. Mus., **42**, 96, 1912.

Figured: Seitz, 165: i1.

Dull brown, head and thorax paler, basal half of wing vaguely darker. Fore wing with traces of a blackish pm. band from near middle of wing to inner margin and a curved blackish line from shortly before apex to well below middle of outer margin.

Aug. 8, 1940 2 ♂ (Sc.,). Described from Costa Rica.

2, *R*₂ stalked; fore wing water-lined.

NATADA INCANDESCENS Dyar

N. incandescens Dyar, Proc. U. S. Nat. Mus., **29**, 380, 1906.

Figured: Seitz, 165: h5.

Warmer brown than the following; the fore wing with some violet gloss, and vague suggestions of a darker base and sinuous pm. band. Water-banding irregular.

Oct. 18-19 (Bts.) May 6-June 6 (Fried.). Ranges to Guiana and the Amazons.

NATADA MICHORTA Dyar

N. michorta Dyar, Proc. U. S. Nat. Mus., 42, 96, 1912.

Figured: Seitz, 165: i4.

With very sparse single black scales, otherwise immaculate or with a faint trace of the dark st. spot of the following.

Principal flight in Dec. (see diagram) but with scattering specimens in Mar.-Apr., perhaps indicating a second brood. Ranges to Guatemala.

NATADA LUCENS DOGNINI Dyar

(*Amydona lucens* Wlk., List Lep. Ins. Br. Mus., 5, 1111, 1855).

N. dognini Dyar, Proc. U. S. Nat. Mus., 29, 379, 1906; Dyar in Seitz, 1125, race *lucens*.

Figured: Seitz, 165: i6.

Very like the North American *Sisyrosea textula*, but larger and darker. The Seitz figure is very bad, the wings too dark, thorax too light, tint too dull (should have a violet tint) and water-lining not shown.

Scattering, with perhaps some concentration in Feb.-Mar. (see diagram). Ariz. to Brazil.

PEROLA Walker

Similar to Natada, except for the stalked Cu_1 and $_2$ of fore wing (fig. 119); R_2 connate or slightly separate; R_5 separate; hind wing with base of M attached far above middle of mdev.

1. Fore wing dull gray, immaculate.....*repetita*
Fore wing yellow with buff-brown markings and often pink shades 2
2. Postmedial line double toward costa, pink shading strong, basal half of fore wing below fold contrastingly pale.....*villosipes*
Postmedial line single, marked by a brown spot in fold; pink limited to body, slight; inner margin of fore wing concolorous...*sericea*

PEROLA VILLOSIPES Walker

Trabala villosipes Wlk., List Lep. Ins. Br. Mus., 32, 555, 1865.

Phocodermma v. Kby, Syn. Cat. Lep. Het., 538, 1892.

Perola v. Dyar, Jour. N. Y. Ent. Soc., 6, 238, 1898, etc.

Not figured.

Fore wing with brown streaks on veins, especially above fold and before outer pm. line, the inner pm. curving sharply in at end of cell,

and running into the streak on R; outer pm. irregularly curved and dentate below cell.

Four ♂'s May 1, 2 (Fried.) 1 ♀ May 26 (Fried.) Ranges to Guiana.

PEROLA SERICEA Möscher

Asbolia sericea Msch., Verh.z-b. Ges. Wien, 27, 671, 10: 36, 1878.

Pseudasbolia s., Kby., Syn. Cat. Lep. Het., 877, 1892; *Perola s.* Dyar, Jour. N. Y. Ent. Soc., 6, 238, 1898, etc.

Also figured: Seitz, 166: f1.

Yellower and a little smaller than the preceding species, the vein-streaks more distinct toward inner margin, but not distinct below cell; pm. obscure or interrupted beyond lower angle of cell, not dentate.

Oct.-Nov. (Bts.) (see diagram). Two rubbed specimens taken by Friedman May 17, may represent a third species. Mexico to Guiana.

PEROLA REPETITA Druce

P. repetita Dr., Ann. Mag. Nat. Hist., (7) 5, 512, 1900.

Figured: Seitz, 166: b2.

Scaling of two shades of fuscous, partly with white tips, with a hint of the water-lining of the second group of Natada. Immaculate.

Common (see diagram), chief flights Oct.-Nov. and Mar.-Apr. Mexico to Guiana and Bolivia.

PALAEOPHOETRON Dyar

Similar to *Epiperola*, but frontal tuft shorter, and palpi much shorter, hardly exceeding the front. R_5 strongly stalked. Habitus distinct, the pm. line irregular or obscure and wings more pointed than *Epiperola*.

Dyar finally sunk this genus in *Epiperola*, but the general appearance is distinct, and I suspect it may really be closer to *Phobetron* than to *Epiperola*.

1. Ground bright buff, marked with brown, coal black and some white *perornata*
Luteous, shaded with blackish along costa and fold, with a large oval white spot on middle of A. *cinereum*

PALAEOPHOETRON PERORNATA Dyar

Epiperola perornata Dyar, Proc. U. S. Nat. Mus., 29, 383, 1906.

Figured: Seitz, 166: a4.

Fore wing with markings complex, the most distinctive being large rounded buff patches at apex and opposite lower angle of cell and a fine irregular pm. white line, preceded with black and followed with dark brown. Female with blunter wings than male.

Dyar never placed this species in *Palaeophobetron*, even when he recognized the genus as distinct, but it shows the essential features, and is even nearer to *Phobetron* in pattern than the other species. The larva will settle the matter.

Common, obviously double-brooded, only three specimens being taken between Nov. 8 and May 6 (see diagram). Costa Rica to Guiana.

PALAEOPHOBETRON CINEREUM spec. nov.

Cream, marked and shaded with gray. Head paler, the front darker and upper side of the flattened palpi dark brown, contrasting; thorax shading into light brown laterally; abdomen light brown, legs shaded with brown, the fore tibia with a strong fringe of blackish tipped hair scales on outer side, ending in a triangular apical black tuft which extends more than half length of tarsus. Fore wing irregularly shaded with blackish, especially along costa and very strongly along veins, leaving the costal space contrastingly pale; vague pm. and st. blackish bars extending down a short distance from costa, strongly accented on veins, vague smoky shades near base and beyond middle of cell and a gray chevron over discal bar; outer part of wing with a semicircular gray line from beyond end of cell almost out to margin, curving back and ending on Cu_2 just before anal angle; a strong blackish shade along 1st A from base to outer margin, giving off a branch on its lower side to inner margin at $2/3$, generally fading out above and below, but sharply defined below by the upper edge of an ovate shining cream spot lying on middle of 2d A, and a small st. spot on 2dA, both of these spots more clearly defined above than below; fringe dark brown, contrasting, with blackish terminal line and middle and terminal lines in the fringe; hind wing cream, shading into light brown toward inner margin, the veins dark, especially Cu with its forks; and fringe as on fore wing.

Barro Colorado Id., C. Z. Panama, holotype Dec. 31, 1934 (Bates) in M.C.Z., paratype Jan. 3, 1935, deposited in Cornell University collection.

This species is related to *P. arcuata* Dr. (Biol., 88: 9; Seitz, 166: a7) but is much paler, with relatively much larger cream spot on 2d A.

EPIPEROLA Dyar

Similar to *Perola* except for the loss of the upper spurs of hind tibia. This genus (or section) is obviously connected with *Perola*, and has the same head characters. I have not examined *E. monochroma* structurally and it may be a *Palaeophobetron*, but I think not. The pm. line is single, oblique and pale as in various *Natada* and *Perola*; R_5 almost always free.

1. Ground blackish, the pale pm. line running to apex *vafera*
Ground light gray, the pm. line from well before apex. *albimarginata*
Ground cream, mottled, the line defined with light gray. *paida*
Immaculate brownish ochreous. *monochroma*

EPIPEROLA ALBIMARGINATA Kaye

Sisyrosea albimarginata Kaye, Trans. Ent. Soc. London, 1901, 158;

Epiperola a., Dyar, Proc. U. S. Nat. Mus., 29, 383, 1906.

E. argentilinea Gaede, D.E.Z. Iris, 30, 208, 1916; Dyar in Seitz, 1126, pr. syn.

Light gray with scattered black scales, the pm. line somewhat defined with brown, and dark basal line in fringe.

Oct. 30 (Bts.) Feb. 24 (Fried.) Mar. 3 (Fried.), 12-18 (A.M.N.H.)
Ranges to Guiana and Brazil.

EPIPEROLA PAIDA Dyar

E. paida Dyar, Proc. U. S. Nat. Mus., 42, 97, 1912.

Perola osseata Schs., Proc. U. S. Nat. Mus., 57, 148, 1920; Dyar in Seitz, 1126, pr. syn.

Figured: Seitz, 165: k3 (unrecognizable if meant for this at all).

R_5 short-stalked in 3 out of 4 specimens; ground paler than the preceding, with shade before the pm. line broad and irregular and discal dot strong and black.

Oct. 19, 24, Nov. 15, Dec. 13 (Bts.). Ranges to Guatemala.

EPIPEROLA VAFERA Druce

Perola vafera Dr., Ann. Mag. Nat. Hist., (7) 5, 512, 1900.

Palaeophobetron v., Dyar, Proc. U. S. Nat. Mus., 29, 382, 1906.

Epiperola v., Dyar in Seitz, 1126.

Perola gaya Schs., Proc. U. S. Nat. Mus., 57, 148, 1920; Seitz, l.c. pr. syn.

Figured: Seitz, 165: k7.

Dark fuscous, the line fine, luteous, slightly powdery, faintly defined with darker. This is a short-winged, heavy, Natada-like species, and I see no resemblance to Paleophobetron.

Dec. 13 (Bts.) Mexico to Amazons.

Dyar also reports *E. monochroma* Dyar; it was described from further west in Panama.

DICHROMAPTERYX Dyar

Similar to Prolimacodes in build and structure; differing only in the small size and distinctive pattern, the wing being practically bisected by a straight pale shade. I cannot see the antennal difference indicated by Dyar.

DICHROMAPTERYX DIDYMA Dyar

D. didyma Dyar, Proc. U. S. Nat. Mus., 42, 99, 1912.

Distinguished from related species by the contrasting coppery dorsal crest and apical shade.

Series (see diagram). Described from Costa Rica.

PROLIMACODES SCHAUS

(*Eulimacodes* auct. not Möschler)

This is the only Neotropical genus so far as I know belonging without doubt to the smooth Eucleids, though plenty of possibly related genera have unknown larvae.

Antenna simple, with a double series of small ventral dentations, rather less than truly serrate; palpi obliquely upturned to middle of front. Fore wing with upper angle of cell somewhat extended, M attached to near lower end of mdcv. or even opposite M₂; hind wing with Sc and R distinctly separated, connected by the "cross-vein"; upper angle of cell rather retracted.

Larva of the North American *P. scapha* smooth, with a strong longitudinal subdorsal crest, which rises almost to an angle at middle of body. Lateral crest absent (shown by earlier stages to be practically fused with subdorsal one); abdominal spiracles all alike and in line. Green, the dorsum brown, contrasting. Food various trees and shrubs.

Hopp has published a key to the species (D.E.Z. Iris, 41, 174).

1. Body long, male with half its length projecting beyond abdomen; brown patch of fore wing truncate-triangular, extending to 2d A and very finely or imperfectly outlined with silver. . . *triangulifera*

Body hardly exceeding hind wings even in male; triangular patch rounded, extending down only to 1st A, strongly edged with silver, especially the basal extension along costa *scaphoides*

PROLIMACODES TRIANGULIFERA Schaus

P. triangulifera Schs., Jour. N. Y. Ent. Soc., 4, 56, 1896; Biol., 2, 444, 88: 12, 1898; Strand, Arch. Naturg., 1922A (7) 140 ♂; Hering & Hopp, D. E. Z. Iris, 41, 177, 3: 2 (♂ genitalia).

To be figured: Seitz, 167: d.

All but one of our specimens show only scattered silver scales.

Oct. 31–Jan. 11; May (see diagram). Ranges to Mexico, the closely related *P. lilalia* Dyar in Guiana.

PROLIMACODES SCAPHOIDES Hering

P. scaphoides Hrg. & Hopp, D. E. Z. Iris, 41, 177, 3: 1 (♂ genitalia).

Close to the North American *P. badia* Hbn. (*scapha* Harr.) but larger, with longer wings, a lighter ground with faint pinkish cast and different genitalia. (See original description).

Oct. 31 (Bts.) ♂; Lancetilla, Honduras, Feb. 25 (Bts. ♀). Described from Mexico.

The National Museum has a specimen of *Vipsophobeton marisa* Dr. from the Canal Zone. It is the one credited to *V. marona* in Proc. U. S. Nat. Mus., 47. It is light brown, with purple and copper iridescence, and with sinuous light brown transverse bands.

DALCERIDAE

Intermediate between the Eucleidae and Megalopygidae, but more primitive than either. Head small, but with relatively large eyes; tongue and ocelli absent, chaetosema not clearly developed, if present lost in the general loose hair. Antennae pectinate (in both sexes as a rule), but relatively extremely short, sometimes sinuously bent and with the apex knobbed. Wings (figs. 125, 126) as in Eucleidae but extremely broad, most often with accessory cell (except *Dalcera* itself and a few others); legs heavily tufted as in Eucleidae and Megalopygidae.

Larva slug-like, much like the Eucleidae, but with at least the hooks of the normal prolegs; translucent, with glass-like spines. Cocoon with a loose outer web enclosing a more or less fusiform inner cocoon

which does not have a formed trap-door (very much like the cocoon of the European genus *Zygaena*, but lighter); the outer cocoon sometimes with a formed thin place corresponding to the paired thin places of *Aidos*.

The family is small, purely American, but some of the species may be common. The larva, cocoon and acc. cell are primitive for the superfamily, but the appearance and minute antennae are specialized, so it certainly is on a line separate from both the Eucleidae and Megalopygidae. The typical *Dalcera* group are regional but have not been taken on the Island; they are large species with modified male antennae, and open acc. cell, R_{2+3} being on a separate stalk from R_{4+5} .

References

Dyar: Jour. N. Y. Ent. Soc., **6**, 231-233, 1898 (first treated as a family with key to genera).

Schaus: Proc. U. S. Nat. Mus., **29**, 331, 1906 (revised key to genera by Dyar).

Dyar: Proc. Ent. Soc. Wash., **12**, 113 ff., 1910 (key to genera; larvae on pp. 116-119).

Dyar and Strand: Lep. Cat., **16**, 27-31, 1913 (Bibliography).

Dyar: Proc. U. S. Nat. Mus., **47**, 252-253, 1914 (Panama species).

Hopp: Iris, **42**, 283-287, 1928 (with note on larva).

Seitz: Macrolep. World, **6**, 1303-1312, pl. 168 (186).

1. Accessory cell absent; with only 2 or 3 radials,— R_5 and the stalked or united R_{1+2} *Ca*
 Accessory cell present; R_1 , the long-stalked R_{2+3} and the connate or short-stalked R_{4+5} arising from it 2
2. Fore wing with cell retracted at upper angle, the acc. cell well before its end; normally with Sc and R of hind wing fusing for a long distance and abruptly diverging near end of cell *Acragopsis*
 Fore wing with cell cut off squarely at end, the ends of discal and acc. cells in line; hind wing normally with Sc and R not fusing, or if so very gradually diverging *Acraga*

ACRAGA Walker

(with *Pinconia* Moore, *Epipinconia* Dyar)

Hind wing with Sc and R extremely variable, even individually, from completely free to strongly fused, but so far as examined always separate but close along outer part of cell. The plain orange species are practically indeterminable save by male genitalia. Heinrich has

checked specimens of the present lot and found them to agree with the type of *A. philetera*, already reported by Dyar from the Zone. Of course other species may be mixed in.

Larva of *A. moorei*, according to Jones "quite white and translucent, just as if made of Venetian glass. The abdominal legs are wanting [presumably the usual series of crotchets are present] their place being indicated merely by a slight swelling of the skin. The motion of the caterpillar . . . is that of a slug"; Burmeister reports *A. flava* as green with detachable transparent spines (Desc. Rep. Argent., 5, 517); that of *A. coa* according to Dyar is ice white, with transparent skin, glistening, all tubercles and spines tipped with scarlet. The cocoon in a folded leaf, the silk of the outer cocoon (Nat. Mus.) so spun as to leave a circular gap in the middle of the upper side.

1. Cream with brown discal bar and more or less submarginal shading. *caretta*
 Orange, without darker markings 2
2. Veins contrastingly and broadly pale *coa*
 Veins concolorous *philetera*

ACRAGA PHILETERA Schaus

Anacraga philetera Schs., Ann. Mag. Nat. Hist., (8) 6, 415, 1910.

A. phileterea Dyar and Strand, Lep. Cat., 16, 30.

The orange of the fore wing is slightly deeper toward the inner margin. *A. conda* Dyar from Guiana is distinguishable only by the genitalia.

Two flights, in Nov. and Apr. (see diagram). Described from Costa Rica.

ACRAGA CARETTA Dyar

A. caretta Dyar, Proc. U. S. Nat. Mus., 38, 268, 1910; Proc. Ent. Soc. Wash., 12, 118, 1910.

Male with strong brown shading toward anal angle of fore wing and a distinct shade at anal angle of hind wing, obsolescent in female.

Nov. 23, Dec. 4, 9 (Bts.), Apr. 9 (Fried.). Ranges to Mexico.

Dyar also reports *A. coa* Schs. from the Canal Zone. It is larger and deeper orange, with the broad pale veins conspicuous, and is figured in Holland, Moth Book, 8: 6.

ACRAGOPSIS Dyar

Easily distinguished by the position of the accessory cell (fig. 126)'. The species are small. Both the following species were placed by Dyar in *Anacraga*, but show the characteristic venation of *Acragopsis*.

1. Yellow of fore wing limited to base and basal $2/5$ of inner margin.
dulciola
 All borders of fore wing shading into yellow. *mesoa*

ACRAGOPSIS DULCIOLA Dyar

Anacraga dulciola Dyar, Proc. U. S. Nat. Mus., 47, 252, 1914.

Chief flight in Oct.-Nov. (see diagram). Described from the Canal Zone.

ACRAGOPSIS MESOA Druce

Dalcera mesoa Dr., Biol. Cent.-Am., Lep. Het., 1, 213.

Dalcerides m. Dyar, Jour. N. Y. Ent. Soc., 6, 232, 1898.

Anacraga m., Dyar, Proc. Ent. Soc. Wash., 12, 119, 1910.

June 17 (Fried.) 1 ♀. Mexico.

Dyar describes *Ca anastigma* from the Canal Zone. It is cream with darker shades, and has a black point at upper angle of cell and a couple subterminally.

MEGALOPYGIDAE

Similar to the Eucleidae; chaetosema present but frequently very weak. Fore wing (figs. 122-124) without acc. cell, the American species with base of M simple (forked in the African genus *Somabrachys*), R_4 and 5 stalked farthest, R_3 normally out of their base; M_1 far below angle of cell and 2d A frequently with one or more secondary posterior branches to inner margin. Hind wing normally with Sc and R fused for most of length of cell, rarely fused for a short distance or separate and connected by a cross vein; M_1 far from R.

Egg ellipsoidal, laid on its side, larva stout, slug-like, of the general form of the Eucleidae, but with prolegs well developed and two additional pairs which bear suckers though not hooks; either clothed with tufts of poison-hairs from warts, or with these reduced and covered by dense secondary hair. Cocoon with a trap door at one end, the outer cocoon if present fitting rather closely; in the North American *M. opercularis* at least, extremely thin and weathering away.

The normal members of the family seem to divide into two series, aside from the very distinct African types, one with Sc and R fused almost to end of cell or beyond, and caterpillar with skin exposed, the other with Sc and R more shortly fused if at all and caterpillar with dense secondary hair, but the adult and larval features on present knowledge seem not to correspond. The third American group, with *Aidos* and *Brachycodilla*, has sometimes been made a separate family, with Sc and R closely parallel but not fusing at all in either sex, larva with small tufts of poison spines only and cocoon unique, with trap door, silken stem for suspension and one or two pairs of rounded depressions in the sides (varying in individuals of a single batch of *A. amanda*). The ranges of *A. amanda* and *perfusa* include the Canal Zone.

References

- Dyar: Proc. Ent. Soc. Wash., **12**, 161 ff, 1910 (key to genera).
 Dyar and Strand: Lep. Cat., **16**, 7-26, 1913.
 Hopp: Berlin (Univ.) Zool. Mus., Mitt., **13**, 209-336, 1926, with supplement in l.c., **15**, 39-51, 1929 (key to genera of Trosiinae and to species; discussion of subdivision).
 Seitz: 1071-1099, 1935.

The family is found in the two Americas and Africa, including the Palaearctic part, but does not reach Europe or Asia. The African types form a separate group, with more primitive venation and wingless female.

1. Hind wing with Sc separating from cell well before its end; larva normally densely hairy.....2
 Hind wing with Sc leaving cell just before its end or even stalked with R; larva generally with small spine-tufts.....3
2. Frenulum at least in male functional (fig. 122) caught in the frenulum-hook of fore wing; Sc and R frequently fused less than half length of cell.....*Podalia*
 Frenulum vestigial, of several bristles even in male (fig. 123); costa of hind wing much arched; Sc normally fused with cell for more than half its length.....*Megalopyge*
3. One radial vein absent; R₁ free, uncus simple.....*Microrape*
 Fore wing with all veins.....4
4. Uncus massive, hood-like, our species marked with red or tawny
 *Trosia*

- Uncus rudimentary, a mere middorsal angulation; our species pure white *Norape*
 Uncus with central process and well marked lateral lobes (socii), our species marked with gray or black on white *Mesoscia*

MICRORAPE Dyar

Small, slender white species, with simple, slender uncus, R_1 free, R_2 stalked; hind wing with Sc stalked on R. The males can easily be named by denuding the tip of the abdomen.

MICRORAPE MINUTA Druce

Carama minuta Dr., Biol. Centr. Am., Lep. Het., 1, 168, 1886.
Microrape m., Dyar, Proc. Ent. Soc. Wash., 12, 167, 1910.

White, with only the minute palpi and the fore legs gray. Uncus slender, down-curved, simple; costa of valve represented only by a minute chitinous crest, the sacculus moderate, oblong, upcurved.

Nov. 5 (Bts.). Type and Dyar's series also from Panama. Costa Rica (Nat. Mus.), Colombia (Fassl-Hopp).

MESOSCIA Hübner (with *Archylus* Walker)

Similar to *Norape*; socii in present species free, about half as long as the well developed uncus.

1. White with about 6 oblique black dashes and a few rounded spots, forming a pm. series *guttifascia*
 Four alternating bands of cream and gray, the cream bands costal and subterminal *pusilla*

MESOSCIA GUTTIFASCIA Walker

Archylus guttifascia Wlk., List Lep. Ins. Br. Mus., 17, 1719, 1856; Biol., 1, 166, 15: 14, 1886; Dyar, Proc. Ent. Soc. Wash., 12, 168, 1910.

Mesoscia g., Hopp, Mitt. Berl. Zool. Mus., 13, 227-228, fig. 20 (σ^7 genitalia). Also figured: Seitz, 160: c4.

Most of the feet, antennae and front are also black.

Dec. 23 (Bts.) Jan. 23 (Fried.). Ranges to Guiana and Brazil.

M. unifascia Dgn. is very close but by the unique ♀ type has no black dot in the cell, 3 vertical oval spots in place of the oblique dashes below lower angle of cell and a single longitudinal bar in radial area instead of two or three oblique spots.

MESOSCIA PUSILLA Stoll

Phalaena Bombyx pusilla Stoll in Cr., Pap. Exot., 4, 220, 395: G, 1782.

Mesoscia p. Hübner, Verz. bek. Schm., 194, no. 1991, 1820; Hopp, Mitt. Berl.

Zool. Soc., 13, 225, figs. 14, 15 (♂ genitalia) 1927.

Also figured: Seitz, 160: b2.

Oct. 18, Dec. 20, Feb. 3 (Bts.). Ranges to Guiana and Peru.

NORAPE Walker (*Carama* auct.)

Thorax with much loose hair, especially in male, and M_3 and Cu_1 of hind wing normally connate (fig. 124), unlike *Trosia*.

Larva stout, frequently brilliantly marked, the warts relatively small, with small dense tufts of stiff poison-hairs; hooks of prolegs divided into two separate groups with the suckers intervening. *N. cretata* was described by Dyar in Jour. N. Y. Ent. Soc., 10, 54.

A large genus, the great majority of the species immaculate white, and determinable only by the male genitalia. Only the two following species have been authenticated from the Zone by examination of males, but two very large females taken will probably belong to another species. The only means of determination is Hopp's revision in Mitt. Berlin Zool. Mus., 12.

1. Head and abdomen yellow; uncus small, but longer than wide; sacculi with a long terminal spine separated by a cleft from the broad main portion *xantholopha*
Abdomen and usually head also white; uncus merely a slight angulation; sacculi with a blunt, compressed blade-like upturned apex
. *pura*

NORAPE PURA Butler

Probably *Euproctis argyrorrhoca* Hbn., Zutr. Samml. Exot. Schm., 2, 13, 43: 245; Hopp in Seitz, 1086, as valid name for species.

Probably *Phalaena ovina* Sepp, Surin. Vlinders, 3, 233, pl. 105, 1852; Hopp, p. 282, no. 1.

Carama pura Btl., Trans. Ent. Soc. London, 1878, 64.

Norape p. Dyar, Proc. Ent. Soc. Wash., 12, 166, 1910; Hopp, Mitt. Berl. Zool. Mus., 12, 315, f. 159-161 (♂ genitalia) 1927.

Carama pruinosa Berg, An. Soc. Cienc. Argent., 1882, 276; Hopp, l.c. 317, pr. var.

Carama butleri Baker, Trans. Ent. Soc. London, 1887, 133 ff., 6: 1-3; Hopp, l.c. pr. syn.

Trosia cuthula Dyar, Proc. Ent. Soc. Wash., 12, 171, 1910.

Sulychra argentea Btl., Trans. Ent. Soc. London, 1888, 64; Hopp, l.c. pr. syn.

Also figured: Baker, Trans. Ent. Soc. London, 1887, 6: 8; Seitz, 161: b3.

Presumed larva (Sepp, l.c.) yellow, with black longitudinal stripe and yellow hair; on Panicum.

I have found this species so abundant in Surinam as to interfere with the capture of other species at light. It dies much more slowly in the cyanide bottle than the ordinary run of moths. Its abundance there is the chief reason for guessing it is the *ovina* of Sepp, whose figures would otherwise suggest a larger species.

Oct. 31-Jan. 28, June (see diagram). Common all over the Neotropical, but North American records are doubtless based on *crotata* or perhaps *virgo*.

Dyar described *N. xantholopha (corporalis* Hopp) from the Canal Zone.

TROSIA Hübner

Structurally almost identical with *Norape*, the genitalic difference not always being valid. Wings generally more pointed, thorax and wings more closely scaled, head more prominent.

1. Fore wing with a series of postmedial black spots; the red of the thoracic tufts, costa etc. scarlet to crimson 2
Fore wing without spots; body and hind wing tawny red *ribbei*
2. Pm. series of black spots without a member below A, the spot beyond the cell drawn in, making a decided angle with the next two below; ground of both wings gray *pulla*
A black spot below A, the spot opposite cell in line with those from M_3 down, though the one in M_2 if present may be farther out; hind wing scarlet 3
3. No spot of the pm. series above M_1 ; the costal red stripe rather broad and of even width to apex, though normally leaving the tip of R_5 free; wings long *punctigera*
A black spot above M_1 , wings blunter 4
4. Spots tending to be small, no spots in cells M_2 and M_3 ; costal red stripe broad, normally strong on the apical branches of R, including R_5 and sometimes part of M_1 *fallax*
Spots large, a large one in M_3 and usually M_2 also, making 8 in all; costal red weak, and usually widened subterminally *dimas*

TROSIA DIMAS Cramer¹

Phalaena Bombyx dimas Cr., Pap. Exot., **1**, 91, 59: C, 1775.

Sciathos d., Kirby, Cat. Lep. Het., 198, 1892, etc.

Trosia d., Dyar, Proc. Ent. Soc. Wash., **12**, 170, 1910, etc.

Bombyx tricolora F. Mant. Ins., **2**, 114, 1787; Dyar, Jour. N. Y. Ent. Soc., **7**, 173, 1899, (blended with other species).

T. dimas var. *fumosa* Hopp in Seitz, Macrolep. World, **6**, 1081, 1934.

T. flavida Dgn., Het. nouv. Am. Sud, **3**, 59, 1911; Hopp, l.c., var. pr.²

T. incostata Schs., Proc. U. S. Nat. Mus., **29**, 335, 1905; Hopp, l.c., syn. var. *flava*.

T. tolimata Dgn., Het. nouv. Am. Sud, **3**, 56, 1911; Hopp, l.c. syn. *flava*.

T. misda Dyar, Proc. Ent. Soc. Wash., **12**, 171, 1910 (Schs. ms.); Hopp, l.c., syn. *flava*.

T. amarilla Hopp, Deut. Ent. Zeit., 1922, 431; l.c., syn. *flava*.

Edebessa albida Dgn., Le Nat., May 15, 1905, p. 120; Hopp, l.c., var. *dimas*.

Sciathos metaleuca Dr., Ann. Mag. Nat. Hist. (7) **18**, 89, 1906; Hopp, l.c., syn. *albida*.

T. donckieri Dgn., Het. nouv. Am. Sud, **24**, 32, 1924; Hopp, l.c., syn. *albida*.

T. obsolescens Dyar, Jour. N. Y. Ent. Soc., **7**, 173, 1899; Hopp, loc., syn. *albida*.

T. rosita Schs., Proc. U. S. Nat. Mus., **57**, 145, 1920; Hopp, l.c., var. *dimas*.

T. dimas ab. *nigra* Hopp in Seitz, Macrolep. World, **6**, 1082, 1935.

Also figured: Seitz, 160: f2 (*dimas*), 3 (*fumosa*), 4 (*incostata*), 5 (*tolimata*), 6 (*amarilla*), g1 (*albida*), 2 (*rosita*), 161: b1 (*nigra*).

The three present specimens are all pinkish white females, with heavy black spots.

Oct. 31, May 19-22. Neotropical generally, ranging north to Arizona.

¹ Genitalic characters are weak in this series, and more than one species may be included here. The following forms, as represented in the Nat. Mus. could be grouped in two or even three species:

No contrasts; ground all buff, with gray spots. *flavida*
Thorax at least contrastingly of two colors, orange or red on white, or red on buff

Fore wing pure white

Apical patch of fore wing scarlet, hind wing more or less scarlet

Hind wing mainly buff. *donckieri*

Hind wing scarlet

Pm. dots of fore wing black *dimas*

Dots of fore wing red *roseipuncta*

Apical patch orange, hind wing white. *albida* (*metaleuca*)

Apical patch absent, hind wing white, very small unnamed race from Argentina

Fore wing cream to buff

With an apical red dash, fore wing generally paler, rarely buff

Larger, paler, typically cream ground (type from Guiana) *dimas* (*tricolora*)

Smaller, normally buff (type from Paraná) *misda*

No apical red dash, ground deep buff, like the darkest specimens of *misda*

Hind wing scarlet. *incostata*

Hind wing concolorous buff *tolimata*

² I believe this is a distinct species, see preceding note. Hopp calls it "*flava*" by a laps. cal.

TROSIA PUNCTIGERA AMALA Dyar

[*Phalaena Bombyx punctigera* Stoll in Cr. Suppl. Pap. Exot., 151, 34: 1, 1A, 1790; *Sciathos p.* Wlk., List Lep. Ins. Br. Mus., 3, 752, 1855; *Trosia p.*, Dyar, Proc. Ent. Soc. Wash., 12, 169, 1910.]

[*T. anax* Dgn., Het. nouv. Am. Sud., 23, 33, 1923; Hopp in Seitz, 6, 1082, pr. var.]

[Also figured: Seitz, 160: g3 ♂, 4 ♀, 5 (var. *anax*).

T. p. var. *amala* Dyar, Proc. Ent. Soc. Wash., 12, 170, 1910.

Wings normally pale pinkish gray with a distinct whiter shade below the costal stripe. The Nat. Mus. keeps the Central American specimens (race *amala*) separate from the type race from South America, but the differences are slight, if any. Var. *anax* has white hind wings, and I have not yet seen it from the Canal Zone.

Apr. 22, June 6 (Fried.), Also Tela, Honduras (Bts.). Typical *punctigera* is wide-spread in the Neotropical.

TROSIA FALLAX Felder

Isochroma fallax Fld., Reise Novara, 83: 18 ♂, 19 ♀, 1874; *Trosia f.* Dyar, Proc. Ent. Soc. Wash., 12, 170, 1910.

T. rosita Schs., Proc. U. S. Nat. Mus., 57, 145, 1920.

Also figured: Seitz, 160: g7 (not g2, which is a form of *dimas*.)

There is little or no white below the costal stripe; the two missing spots make a distinct gap in the series.

Oct. 31–Feb. 2 (Bts.), July 5 (Scr.), July–Aug. (Fairchild). Costa Rica to Ecuador, at least; a female from "Brazil" in the Nat. Mus. also appears to be this species.

TROSIA PULLA spec. nov.

Vertex red, face and palpi (which are minute) orange; antennae black, first segment white below, buff above, second white; thorax white above, with red base of collar and three pair of red spots, abdomen smoky with first two segments red. Under side of thorax red, breast orange, tibiae and tarsi blackish, also tips of fore femora; abdomen smoky with red patches under base of wings, some red hair mixed in basally, and cream terminal segment. Wings ash gray, the costa of fore wing with a broad, even but diffuse red fascia from base to apex; a series of black spots in two concave sweeps with the spot in M_3 forming the cusp: spots in R_5 , M_1 , M_3 , Cu_1 , Cu_2 , 1stA; the spot below 2dA very faint and much further out than the series. Hind wing darker

ash gray, the base and costa overlaid with red hair reaching down to R at margin; under side with the red on fore wing much more extensive, covering cell, and with red overlay on much of surface; the spots showing through faintly; hind wing with costal red stripe broad and strong, covering more than a third of cell. 38 mm.

Easily distinguished from the known species of the genus by the different series of spots on fore wing and dark gray hind wing.

Barro Colorado Id., C. Z. Panama, Oct. 10, 1934 (Bates) female holotype in M.C.Z.

TROSIA RIBBEI Druce

Sciathos ribbei Dr., Biol. Centr. Am., Lep. Het., 2, 441, 88: 1, 1898; Dyar, Proc. Ent. Soc. Wash., 12, 169, 1910, syn. *acca*.

T. acca aterrima Hopp, D. E. Z. Iris, 44, 76, 1930.

T. electra Hopp, Deut. Ent. Zeit. (Berlin), 1922, 431; Hopp in Seitz, 1081, syn. *acca*.

Also figured: Seitz, 160: e2 (as *acca*), 3 (*ribbei*), 4 (*aterrima*).

Fore wing light brown, with darker veins and more less distinct traces of a vague brown pm. band, accented on the veins; in var. *aterrima* the wings are deep gray, contrasting with the orange head, thorax and base of abdomen.

Hopp considers this a synonym of *acca* Schs. but Heinrich considers the latter distinct, based on the unique type, which has a narrow even dark pm. line and distinct genitalia. *T. ribbei* is also found at Rio.

Dec. 5, 8, 13 (Bts.) Mexico to southern Brazil.

MEGALOPYGE Hübner (with *Lagoa* Harris)

Fore wing (fig. 123) with all veins, R_2 sometimes stalked with R_{3-5} ; which are well stalked, the rest separate; anal veins normally with one or more secondary branches to inner margin, besides the free tip of 3rd A. Hind wing with Sc and R fused for a considerable distance, but separating well before end of cell. Scaling usually fine, more or less hair-like, frequently laid in curly waves (the "Flannel-Moths"); body stout with front of thorax prominent.

Formerly this genus was often divided on minor points of venation now considered unsafe; now even *Podalia* is frequently combined with it, but I believe the distinction can be made if frenulum as well as venation is considered.

Body of larva densely covered with fine woolly hair, arising from diffuse warts and concealing the poison setae; usually with a slight dorsal crest, and longer hair posteriorly. Hooks of prolegs in a chevron.

MEGALOPYGE OPERCULARIS COSTARICENSIS Schaus

[*Phalaena opercularis* Smith and Abbot, Lep. Ins. Georgia, 2, pl. 53, 1797.]

[Also figured: Holland, Moth Book, 38: 25].

[Larva and Biology: Fracker, Class. of Lep. Larvae, Ill. Biol. Monog., 2 (1) 97, 1915; Forbes, Field Tables, 90, 1906; Lep. N. Y., 102, 1924; Kellogg, Am. Insects, 383; Comstock, Introd. to Ent., 189, 607; Bishopp, The Puss Caterpillar and its Effects on Man., U. S. Dept. Agr. Circ. 288, 1923; Matheson, Med. Ent., 446, 449, fig. 201.4; etc. etc.]

Megalopyge costaricensis Schs., Ann. Mag. Nat. Hist., (8) 9, 55, 1912; Seitz, 1093, pr. race.

Figured: Seitz, 162: b5 (unrecognizable).

Base of fore wing mixed curly buff, brown and blackish hairs, the brown dominant; border light buff. Race *costaricensis* is only a little larger and lighter than the type form, and looks more like Seitz's figure of "*opercularis*" than his "*costaricensis*".

Common (see diagram). The race ranges from Costa Rica to Colombia, the species from southern U. S. to Bolivia, and is replaced by the whiter *M. amita* Schs. in southern Brazil.

PODALIA Walker

The following species are listed in *Megalopyge*, but show both frenulum and venation of *Podalia*. (See fig. 122).

1. Nearly immaculate, the male blackish with transparent wings, the female gray with small pm. white dots *bolivari*
Postmedial area at least whitish with gray streaks on and between veins; fully scaled 2
2. Basal 2/3 coal black, contrasting with outer third *thanatos*
Basal 2/3 contrasting, mouse gray, but with whitish markings at least toward costa *albicollis*

PODALIA ALBICOLLIS Walker

Gasina albicollis Wlk., List Lep. Ins. Br. Mus., 6, 1479, 1855; Druce, Biol., 1, 204¹; *Megalopyge a.*, Dyar, Proc. U. S. Nat. Mus., 38, 267, 1900.

Megalopyge vipera Schs., Jour. N. Y. Ent. Soc., 4, 58, 1896.

Megalopyge braulio Schs., Proc. U. S. Nat. Mus., 65, (7), 61, 1924.

Figured: Seitz, 161: f4 ♂, 3 ♀.

Biology (of *P. a. superba*)¹: Edw. Pap., 4, 79, 1884 (pupa); Packard, Mem. Nat. Acad. Sci., 7, 76, fig. 37 (pupa); Dyar, Proc. Ent. Soc. Wash., 14, 58, 1912 (larva).

¹ These northern citations apply primarily to *M. a. superba* H. Edw., Pap., 4, 79, 1884: *Lagoa grandis* Kby., Cat. Lep. Het., 472, 1892 (laps. cal.).

Female much larger than male, but with the same pattern. Larva with fine dense hair, white till last stage, then turning to tawny, like the North American *M. crispata*.

Not rare (see diagram). Mexico to southern Brazil, the blend zone of the two races lying in Central America.

PODALIA THANATOS Schaus

P. thanatos Schs., Proc. U. S. Nat. Mus., **29**, 339, 1905; *Repnoa t.*, Dyar, Proc. Ent. Soc. Wash., **12**, 168, 1910.

Figured: Seitz, 163: f2.

This species has blackish patches in the pale area in cell Cu_1 , pm. line regularly dentate on dorsal part of wing, and dark spots in the pale area costad; hind wing with the dark gray dorsal half contrasting. *P. dimidiata* has the deeply dentate pm. regular up to M_2 , and wholly pale hind wing; *P. arpi* has a less dentate pm. below, running into an abrupt change of color at end of cell costad, followed by slight separate dots at lower angle of cell, and an isolated st. dot at R_5 ; hind wing intermediate. *P. nigrescens* looks much like *arpi*, but has the dark area extending broadly and almost solidly to anal angle.

May 12 (Fried.). Guatemala to Guiana and Peru.

PODALIA BOLIVARI Heylaerts

♂ *Pentophora bolivari* Heyl., Comptes-rendus Soc. Ent. Belge, **28**, p. xli, 1884 (missed by Lep. Cat.); *Podalia b.*, Hopp in Seitz, 1098.

Megalopyge pellucens Dgn., Mem. Soc. Ent. Belge, **19**, 171, 1912.

♀ *Unduzia gistinda* Dyar, Proc. U. S. Nat. Mus., **47**, 252, 1914; Hopp, D. Ent. Zeit. (Berlin), 1926, syn. *bolivari*, and refs.

Unduzia phaule Dyar, Proc. U. S. Nat. Mus., **47**, 252, note, 1914.

I have seen only females.

Dec. 10, 29 (Bts.). Mexico to Venezuela and Peru.

HEPIALIDÆ

Head commonly small; mouth parts absolutely reduced, palpi short to minute; antennae typically very short and weak, but moderate in Dalaca. Body very long and slender, loosely constructed, the metathorax almost as well developed as the mesothorax. Hind wing like fore wing (figs. 127) except for minor points, the radius four- or five-branched. Fore legs long, tufted and displayed at rest; middle and

hind legs weak, the latter without spurs and with a large sex-tuft in male.

Eggs minute, extremely numerous, broadcast by the female, at least in some species. Larva slender, active, the ocelli not arranged in the usual crescent; prolegs with multiserial hooks; upper two setae of meso- and metathorax almost horizontally placed, as on abdomen, setae iv and v of abdomen both behind spiracle. Pupa long, slender, incomplete.

All the species are borers, mostly in woody plants. Some are semi-aquatic, others form a burrow in the trunks of the hardest trees, but browse on the bark around the hole.

The most primitive of larger moths, representing a separate sub-order (the *JUGATAE*) from all the preceding. As with other very ancient groups, species are widely scattered but relatively enormously common in Australasia.

References

Most of the following references concern the morphology and fundamental relationships of the family. There has been no systematic work, aside from Seitz, of major interest from the Neotropical point of view.

Kirby, *Cat. Lep. Het.* 286-292, 1892. (catalogue).

Hampson, *Fauna of British India, Moths*, 1, 315 ff., 1892 (contains a generic key).

Packard, *Monograph of the Bombycine Moths*, 1 (Mem. Nat. Acad. Sci., 7), 71-76, with figs. 31-34, 1895 (general discussion with structural figures of larvae and pupae).

Dyar, *Am. Nat.*, 29, 1066-1072, 1895 (discussion of relationships, with special reference to larva).

Eyer, *Bull. Bklyn. Ent. Soc.*, 16, 1-8, 1921 (major subdivisions as indicated by genitalia).

Eyer, *Ann. Ent. Soc. Am.*, 17, 303-305, 1924 (genital characters).

Philpott, *Trans. Ent. Soc. London*, 1925, 331-340 (wing-coupling).

Tindale, *Rec. S. Austr. Mus.*, 4, 497-536, 1938 (revision of Australian *Hepialidae*).

Issiki and Stringer, *Stylops*, 1, 73-80, 1932 (relationships and morphology of annectant forms, with references).

DALACA Walker

Male antennae bipectinate. Fore wing with apex extended but somewhat rounded over, not falcate.

In our lists this genus includes African as well as Neotropical species, but I suspect they may really not be congeneric. The type (from Chile) was not examined.

1. Two blackish spots below Cu_2 , the inner grading into the fuscous costal area. *metellus*
 No special spots below Cu_2 , and costal area not noticeably darker than dorsal. 2
2. About five striae in cell M_1 beyond the nearly uninterrupted pm. line, the alternate ones often enclosing slightly darker areas. . . . *assa*
 Only three such striae, the ground nearly even. *new species*

DALACA ASSA Druce

♂ *Dalaca assa* Dr., Biol. Centr. Am., Lep. Het., 1, 232, 24: 10, 1887.

♀ *Phassus costaricensis* Dr., Biol. Centr. Am., Lep. Het., 1, 234, 24: 4, 1887.

Also figured: Seitz, 99: c3 ♂, 4 ♂ labelled ♀, 185: c1 ♀ (*costaricensis*).

Male fore wing either light brown or salmon, the spaces between alternate striae a little darker; a black spot with brassy central dot near end of cell, rarely absent. Hind wing nearly plain, pale brown or salmon. Female much larger, brilliant salmon, the markings faint. The type of *costaricensis* is a slight variety with a blackish patch near end of cell.

Common (see diagram). Ranges to Guatemala.

A second closely related species with blunter wings, fewer striae, no brassy dot, and a dark shade over end of cell, is represented by a single specimen,—Apr. 24 (Fried.)

DALACA METELLUS Druce

Hepialus metellus Dr., Proc. Zool. Soc., 1890. 509, 43: 2; Wagner & Pfitzner, Lep. Cat., 4, 18, 1911; *Pseudophassus momus metellus* Pf. in Seitz, Macrolep. World, 6, 1301, 1938.

Hepialus momus Dyar, Proc. U. S. Nat. Mus., 47, 350, 1914 (not Druce?).

Pseudophassus metricus, v. *songoensis* Pf., Ent. Rund., 31, 110, 1914; Seitz, l.c., var. *momus metricus*.

Costa fuscous, dorsum pale, the boundary pretty sharp, irregular, extending down in a blunt triangle at middle, where it ends in the first blackish spot below Cu_2 . Pm. line fine, somewhat waved, parallel to outer margin, stronger than the other transverse markings. Like the preceding, this species has a fuscous form (*metellus*) and a red form (*songoensis*).

Two specimens, May 1, 8 (Fried.), both fuscous males. Ranges to Bolivia. Dyar formerly reported the Panama species as *momus*, but later transferred the specimen to *metellus*. Typical *momus* is much more mottled, with a good deal of cream color. If conspecific, *momus* will take "line priority" over *metellus*, but I believe we should take advantage of the option and treat *metellus* as the species name.

EXPLANATIONS OF PLATES

PLATE 9

PLATE 9

Male genitalia of *Antaea* species

Each figure is in three parts: a, the genitalia proper, opened out from the ventral side; b, aedoeagus; c. last sternite.

67. *A. lichyi* Franç. holotype (expansile tuft shown on left side only).
68. *A. juturna* (left valve omitted), one cornutus from aedoeagus slightly more enlarged.
69. *A. jaraguana* Franç. holotype (left valve omitted).











PLATE 10


PLATE 10

Venations of Manidiidae, Lasiocampidae, Uraniidae, Eupterotidae.


70. *Manidia lunus*.
71. *Euglyphis rundala*.
72. *Gloveria arizonensis*, humeral area of hind wing.
73. *Tolyte velleda*
74. *Artace cribraria*, hind wing
75. *Euglyphis modesta*, hind wing (outline of costal scaling shown by dotted line)
76. *Urania fulgens*
77. *Epia muscosa*
78. *Colla glaucescens*




70




71





72





73




74




75




76



77



78



79



80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385


PLATE 11

PLATE 11


Male genitalia of *Euglyphis* and *Quentalia*

All the figures on this plate are at the same scale. All drawings of main portions are ventral view, with the valves expanded, except in *E. marna*, where they are coalesced ventrally.

79. *Euglyphis rivulosa*, Tumatumari, British Guiana
80. *E. maha*, Nova Bremen, S. Cath., Brazil
81. *E. larunda*, Chiriqui, Panama.
82. *E. vittabunda*, Barro Colorado Id.
83. *E. marna*, Santa Catherina, Brazil
84. *Euglyphis*, new species near *vittabunda*, from South Brazil; uncus and right valve
85. *Quentalia*, new species of *drepanoides* group from South Brazil: figured in four parts: a, main part; b, aedoeagus, c, chitinized part of last sternite, d, last tergite.




79



80



81




82




84



83



84




85 a.



b.



c.



85 d

PLATE 12

PLATE 12

Structures of Eupterotidae, Apatelodinae

86. *Anticla anticla*, venation
87. *Quentalia chromana*, venation
88. *Tamphana marmorea*, venation
89. *Olceclostera amoria*, venation
90. *Quentalia numalia*, Barro Colorado Id., Male genitalia in four parts:
a, main part, with valves opened out; b. aedoeagus; c, chitinized edge
of last tergite; d, chitinized part of last sternite.
91. *Q. surynorta*, Lancetilla, Tela, Honduras, male genitalia.
92. *Apatelodes lapitha*, venation




PLATE 13

PLATE 13

Venations of Epiplemlidae and Mimallonidae

93. *Syngria druidaria* ♂
94. *Nedusia mutilaria*
95. *Skaphion lilacina*
96. *Erosia incendiata* ♂. The anal roll is not a flat surface and was cut at the point Z to flatten it.
97. *Erosia palpulata* ♂, paratype. To the same scale as fig. 96.
98. *Epiplema* species
99. *Cicinnus incerta*
100. *Gathynia atriceps* ♂, paratype
101. *Mimallo amilia* ♂.
















PLATE 14


PLATE 14

Venations of Mimallonidae and Cossidae


102. *Menevia lantona*
103. *Cicinnus forbesi*
104. *Zaphanta infantilis*
105. *Dysodia speculifera*
106. *Ochrothyris mesogramma*, paratype
107. *Rhodoneura hedilalis*
108. *Givira necreros*
109. *Langsdorfia lunifera*
110. *Inguromorpha polybia*






102




103






104




105





106





107



108



110



109




PLATE 15

PLATE 15

Venations of Cossidae, Zygaenidae, Eucleidae

111. *Zeuzera pyrina*, fore wing
112. *Xyleutes ramosa*
113. *Euclea norba*
114. *Talima rubicolor*
115. *Harrisina metallica*
116. *Euprosterna clacasa*
117. *Narosopsis leucospila*
118. *Semyra* species
119. *Perola repetita*

















PLATE 16

PLATE 16

Venations of Zygaenoidea and Hepialidae


120. *Natada fusca*
121. *Natada michorta*; costal part of fore wing
122. *Podalia orsilochus* (genotype)
123. *Megalopyge lanata*
124. *Norape pura*
125. *Acraga philetera*
126. *Acragopsis dulciola*
127. *Dalaca assa* ♂





120





121





123





122





124





126



125



127



127

Date Due

~~MAR 31 1985~~

NOV 30 1985

~~JAN 31 1985~~

ACME
BOOKBINDING CO. INC.

AUG 23 1984

100 CAMBRIDGE STREET
CHARLESTOWN, MASS.

Harvard MCZ Library



3 2044 066 303 637

