

BLM LIBRARY



88014048




# IDAHO BLM

## TECHNICAL BULLETIN

AN ILLUSTRATED GUIDE TO THE  
SENSITIVE PLANTS OF BURLEY DISTRICT  
BUREAU OF LAND MANAGEMENT

by  
Ann DeBolt



Goose Creek Milk-Vetch


Astragalus anserinus

Technical Bulletin 89-3

February 1989

**BUREAU OF LAND MANAGEMENT**  
**IDAHO STATE OFFICE**  
3380 Americana Terrace  
Boise, Idaho 83706

DL  
84.2  
L352  
no. 89-3



# 19793878

ID 88014048




GL  
84.2  
.1352  
11.89-3

# IDAHO BLM TECHNICAL BULLETIN

AN ILLUSTRATED GUIDE TO THE  
SENSITIVE PLANTS OF BURLEY DISTRICT  
BUREAU OF LAND MANAGEMENT

by  
Ann DeBolt

BLM LIBRARY  
SC-324A, BLDG. 50  
DENVER FEDERAL CENTER  
P. O. BOX 25047  
DENVER, CO 80225-0047



Goose Creek Milk-Vetch

Astragalus anserinus

Technical Bulletin 89-3

February 1989

**BUREAU OF LAND MANAGEMENT**  
**IDAHO STATE OFFICE**  
3380 Americana Terrace  
Boise, Idaho 83706



## INTRODUCTION

This Technical Bulletin was developed to familiarize Burley District field personnel with what sensitive plant species occur, or might occur, in their area. It is believed that it will help streamline the Environmental Assessment and clearance processes by providing a search image for most species, and by listing all current location data and habitat information.

No one flora covers this part of Idaho, as the "Literature Cited" section reveals. Rather, the region lies between four floristic units with distinct floras, namely, the Snake River Plains to the north, the Great Basin to the south, the Owyhee Uplift to the west, and the Albion Mountains to the east (Packard, et. al. 1979). The Salmon Falls Creek area, which is lower than lands to the east and west, appears to be a migration route for Great Basin species such as Allium anceps, Glyptopleura marginata, and Scutellaria nana. Unique edaphic (soil) conditions are prevalent throughout the district as well, and provide the habitat for most of the species of concern.

The plants are arranged in alphabetical order by genera. An index of common names has been included for those unfamiliar with scientific names. Plant nomenclature follows that of Hitchcock and Cronquist in the Flora of the Pacific Northwest (1973), and Cronquist, et. al. in the Intermountain Flora, Vols. 4 & 6 (1977, 1984). Illustrations are from a variety of sources including Flora of the Pacific Northwest, Intermountain Flora, Threatened and Endangered Vascular Plants of Oregon, "Brittonia", and the "Great Basin Naturalist".



LIST OF SENSITIVE PLANTS IN THE BURLEY DISTRICT

- Allium anceps* (Taper-Tip Onion)
- Astragalus anserinus* (Goose Creek Milk-Vetch)
- Astragalus atratus* var. *inseptus* (Mourning Milk-Vetch)
- Astragalus atratus* var. *owyheensis* (Owyhee Mourning Milk-Vetch)
- Astragalus tetrapterus* (Four-Wing Milk-Vetch)
- Castilleja christii* (Christ's Indian Paintbrush)
- Cymopterus davisii* (Davis Parsley)
- Epipactis gigantea* (Giant Helleborine)
- Eriogonum ochrocephalum* var. *sceptrum* (Ochre-Flowered Buckwheat)
- Glyptopleura marginata* (White Margined Wax Plant)
- Gymnosteris nudicaulis* (Large Flowered Gymnosteris)
- Lepidium davisii* (Davis Peppergrass)
- Mentzelia torreyi* var. *acerosa* (Torrey's Blazing Star)
- Pediocactus simpsonii* var. *robustior* (Simpson's Hedgehog Cactus)
- Scutellaria nana* (Dwarf Skullcap)
- Townsendia scapigera* (Stemless Townsendia)

ALPHABETICAL LISTING OF COMMON NAMES

- Christ's Indian Paintbrush (*Castilleja christii*)  
Davis Parsley (*Cymopterus davisii*)  
Davis Pepperygrass (*Lepidium davisii*)  
Dwarf Skullcap (*Scutellaria nana*)  
Four-Wing Milk-Vetch (*Astragalus tetrapterus*)  
Giant Helleborine (*Epipactis gigantea*)  
Goose Creek Milk-Vetch (*Astragalus anserinus*)  
Large Flowered Gymosteris (*Gymosteris nudicaulis*)  
Mourning Milk-Vetch (*Astragalus atratus* var. *inseptus*)  
Ochre-Flowered Buckwheat (*Eriogonum ochrocephalum* var. *sceptrum*)  
Owyhee Mourning Milk-Vetch (*Astragalus atratus* var. *owyheensis*)  
Simpson's Hedgehog Cactus (*Pediocactus simpsonii* var. *robustior*)  
Stemless Townsendia (*Townsendia scapigera*)  
Taper-Tip Onion (*Allium anceps*)  
Torrey's Blazing Star (*Mentzelia torreyi* var. *acerosa*)  
White Margined Wax Plant (*Glyptopleura marginata*)



LIST OF SPECIES WITH AUTHORITIES

*Allium anceps* Kellogg

*Astragalus anserinus* Atwood, Goodrich, & Welsh

*Astragalus atratus* Wats. var. *inseptus* Barneby

*Astragalus atratus* var. *owheensis* (Nels. & Macbr.) Jones

*Astragalus tetrapterus* Gray

*Castilleja christii* N. Holmgren

*Cymopterus davisii* R.L. Hartman

*Epipactis gigantea* Douglas ex Hook.

*Eriogonum ochrocephalum* Wats. var. *sceptrum* Reveal

*Glyptopleura marginata* D.C. Eat.

*Gymnosteris nudicaulis* Gooding

*Lepidium davisii* Rollins

*Mentzelia torreyi* var. *acerosa* Gray

*Pediocactus simpsonii* (Engelm.) Britt. & Rose var. *robustior* Coult.

*Scutellaria nana* A. Gray

*Townsendia scapigera* D.C. Eat.

LIST OF SPECIES BY STATUS

Federal Category 1 (C1)

*Castilleja christii*

Federal Category 2 (C2)

*Astragalus anserinus*  
*Astragalus atratus* var. *inseptus*  
*Lepidium davisii*

State Priority 1

*Glyptopleura marginata*

State Priority 2

*Allium anceps*  
*Astragalus tetrapterus*  
*Epipactis gigantea*  
*Mentzelia torreyi* var. *acerosa*

State Sensitive

*Astragalus atratus* var. *owyheensis*  
*Cymopterus davisii*  
*Gymnosteris nudicaulis*  
*Pediocactus simpsonii* var. *robustior*

State Review

*Eriogonum ochrocephalum* var. *sceptrum*  
*Scutellaria nana*  
*Townsendia scapigera*

Plant Distribution by Soil Characteristics

Dried Mudflat or Playa

Allium anceps  
Lepidium davisii

Volcanic Ash

Astragalus anserinus  
Astragalus tetrapterus  
Eriogonum ochrocephalum var. sceptrum  
Mentzelia torreyi var. acerosa

Fine Alluvial Sand

Glyptopleura marginata  
Gymnosteris nudicaulis  
Pediocactus simpsonii var. robustior  
Townsendia scapigera

Shallow, Rocky

Pediocactus simpsonii var. robustior  
Scutellaria nana

Subalpine to Alpine

Castilleja christii  
Cymopterus davisii

Shallow Clay Over Basalt

Astragalus atratus var. inseptus

Calcareous Hot/Cold Springs

Epipactis gigantea

Sagebrush Hillsides (deeper soils than var. inseptus)

Astragalus atratus var. owyheensis



Allium anceps (Taper-tip Onion)

Family: Liliaceae (Lily)

Status: Priority 2 on the State Sensitive Plant List.

Known Locations: Twin Falls County, Idaho  
NE California  
Nevada  
SE Oregon

sites in or near Burley District

T12S, R18E, Sec. 10 SESE 3.5 miles S of Rock Creek townsite

T14S, R15E, Sec. 31 E side of Salmon Falls Creek Reservoir

T15S, R15E, Sec. 5, 8

Soil Type: Heavy soils of volcanic origin in swales, where water stands in spring. Also known from at least one playa perimeter (non-alkaline).

Habitat and Ecology: A perennial found in low-lying, sparsely vegetated areas where water stands in the spring. Associated species include low sage and Ericogonum microthecum. At the playa, it is associated with basin big sage and Astragalus calycosus. In Idaho it is known from 4600 to 5050 feet, but undoubtedly spans a broader elevational range since only three sites are known. First reported for Idaho in 1979. Blooms in May and early June.

Threats: Range improvement projects


#### Key Characteristics

Flowers pinkish

Leaves two, flattened

Tepals slender, linear to lance-linear

Stamens inserted



BURLEY DISTRICT

THE UNIVERSITY OF CHICAGO  
DEPARTMENT OF CHEMISTRY  
5800 S. UNIVERSITY AVENUE  
CHICAGO, ILLINOIS 60637  
TEL: (773) 835-3100

Astragalus anserinus (Goose Creek Milk-Vetch)

Family: Fabaceae (Legume)

Status: Federal Category II (C2) and Sensitive on the BLM Sensitive Plant List.

Known Locations: Cassia County, Idaho  
Elko County, Nevada  
Box Elder County, Utah

Idaho

T16S, R21E, Sec. 33 N1/2 of SENE near Beaverdam Creek (private land)

Nevada

T47N, R70E, Sec. 29 SW 6 km S of Idaho on the UT-NV line

Utah

T14N, R19W, Sec. 15 SE 6.5 km S of UT-ID line  
.5 km E of UT-NV line at Hardister, 7 km S of ID  
25.5 km NW of Lynn


Soil Type: On white tuffaceous ash, usually of outcrops

Habitat and Ecology: A dwarf, matted perennial known at this time only from undeveloped soils of tuffaceous outcrops in the Goose Creek drainage. It was not described until 1984. Associated species include Stipa comata, Eriogonum ovalifolium, and Chrysothamnus viscidiflorus. Also known from juniper communities. Grows at elevations from 4700 to 5000 feet. It is more matted and has smaller flowers than the common A. purshii. The leaves and pods are woolly, but the hairs on A. anserinus are shorter than those of A. purshii. Blooms in May and June.


Threats: Off-road vehicles

Key Characteristics

Flowers small, 9-11 mm,  
pink-purple in color  
Dwarf, matted growth  
Tomentose herbage  
Pods compressed, curved, and  
lightly hairy  
Similar to A. purshii, but more  
diminutive with shorter hairs



BURLEY DISTRICT



Habit and details of *Astragalus anserinus*



Astragalus atratus var. inseptus (Mourning Milk-Vetch)

Family: Fabaceae (Legume)

Status: Federal Category II (C2) recommended for Threatened status.

Known Locations: Blaine, Camas, Elmore, Gooding, Lincoln, and Twin Falls  
Counties, Idaho

sites near Burley District

T6S, R13E, Sec. 20 SE Shoestring Bridge, S of Bliss

T7S, R13E, Sec. 32 Peter's Gulch near Hagerman (var. in question, specimen  
immature)

Soil Type: Shallow clay soil over basalt



Habitat and Ecology: A wiry, perennial milk-vetch endemic to the mid Snake River Plains of southern Idaho. Flowering stems are decumbent to prostrate. It is usually found in flats and on plains, but also occurs on gentle slopes. At lower elevation sites it occurs with Wyoming sage and low sage, while at higher altitudes it occurs with mountain big sage and A. longiloba. It is also associated with Allium acuminatum, Aster scopulorum and Poa sandbergii. It often occurs within the protection of sagebrush plants. The plant is most common within the Shoshone District, but should be watched for in Burley District, particularly the northwest corner on basalt flats. Collections to verify the variety are best if the fruits are mature. Blooms in May and June.

Threats: Range improvement programs, over grazing, and agricultural development

Key Characteristics

Pods red speckled, with  
leathery texture  
Flowers white, sometimes  
faintly lilac-tinged  
Leaflets 9-15, terminal  
one jointed to the stem

(see key on next page)



BURLEY DISTRICT

Astracalus atratus var. owyheensis (Owyhee Mourning Milk-Vetch)

Family: Fabaceae (Legume)

Status: Sensitive on the ELM and State Sensitive Plant Lists.

Known Locations: Elmore, Owyhee, Twin Falls Counties, Idaho  
Baker, Malheur Counties, Oregon  
Elko County, Nevada

sites in or near Burley District

T9S, R13E, Sec. 11 Salmon Falls Creek Canyon, 5 miles SW of US #30

T14S, R13E, Sec. 7 E of Devil's Creek

T14S, R15E, Sec. 7 1 mile N of Salmon Falls Dam, on bench above Salmon Falls Creek

16S, R17E, Sec. 30 S of Magic Hot Springs, on sides of Shoshone Creek Cyn.

Soil Type: Sagebrush hillsides with relatively deep, volcanic soils

Habitat and Ecology: A perennial with many very slender, often prostrate and creeping to erect stems. Found on steep hillsides and flats over basalt. usually taking shelter under and entangled in sagebrush. Found from 3,500 to 6,000 feet. Often on bluffs overlooking the Jarbidge, Bruneau, and Owyhee River canyons. Its leaves blend in with the leaves of grass and are not easily noticed. Blooms from May to July.

Threats: Range improvement programs, overgrazing, agricultural development

Key to two varieties of Astracalus atratus:

1. Leaflets all very small, narrow, and remote, the terminal one continuous with the rachis or represented by a small dilation of the rachis; pod of papery texture; sagebrush slopes.....var. owyheensis

1. Leaflets more ample and less scattered, the terminal one jointed to the rachis; pod of leathery texture; shallow clay soils where moist in spring .....var. atratus

Key Characteristics

Flowers whitish, purplish-lined  
or tinged and 8-9 mm long  
Leaflets 7-11, terminal one  
continuous with the stem  
Pod 14-20 mm long and 3-4 mm wide  
Pod of papery texture



BURLEY DISTRICT

Astragalus tetrapterus (Four-Wing Milk-Vetch)

Family: Fabaceae (Legume)

Status: Priority 2 on the State Sensitive Plant List.

Known Locations: Twin Falls County, Idaho  
Northwest Arizona  
Eastern Nevada  
Southeast Oregon  
4 counties in Southwest Utah

Idaho

T16S, R15E, Sec. 8 small drainage east of Salmon Falls Reservoir

Soil Type: Sparsely vegetated ash or sandy alkaline soils

Habitat and Ecology: A perennial milk-vetch that is known from only one site in Idaho. Apparently south-central Idaho is the northern extent of its range. It is found in coarse soils within Wyoming sage habitats as well as pinyon-juniper habitat in Utah and Nevada, from 3500 to 6500 feet. Mostly in exposed places but sometimes taking shelter under or entangled in sagebrush. Barneby states that the plant is widely dispersed but uncommon. This species is highly variable. Collections are needed to determine its range in Idaho. Seen but not collected and verified in the BLM Winnemucca District of Nevada. Blooms from late April through June.

Threats: Off-road vehicles, trampling and overgrazing by wild horses

Key Characteristics


Pods four-sided

Pods pendulous, incurved or  
coiled, usually pubescent

Pods 2-4 cm long

Flowers vary in color from white  
with lilac tinges to bright pink  
purple

Stems and leaves round in cross  
section & with pointed tips



BURLEY DISTRICT

Castilleja christii (Christ's Indian Paintbrush)

Family: Scrophulariaceae (Figwort)

Status: Federal Category I (C1) recommended for Endangered status.

Known Locations: Cassia County, Idaho

T13S, R24E, Sec. 4,9 Harrison Mountain, near the top; SE of Burley  
(Forest Service administered land)


Soil Type: Loamy gravel with quartzite and mica schist stones

Habitat and Ecology: A perennial plant endemic to Harrison Mountain in the Cache Peak Range of the Albion Mountains. It is found in grassy subalpine meadows with Trisetum spicatum, Festuca idahoensis, Solidago multiradiata, Pedicularis contorta, and Achillea millefolium. Elevation is 9000 to 9300 feet, on Sawtooth National Forest land. The species was described by Noel Holmgren in 1973, who unsuccessfully searched the neighboring mountains and adjacent peaks in the same range. It is locally abundant on Harrison Mtn. Blooms in July.


Threats: Additional radio relay or lookout facilities, and possibly grazing

Key Characteristics:

Subalpine meadow habitat  
Flowers and bracts uniquely  
yellow to yellow orange  
Plants to 18 inches high



BURLEY DISTRICT



Cymopterus davisii (Davis' Parsley)

Family: Apiaceae (Carrot)

Status: Sensitive on the State and BLM Sensitive Plant Lists.

Known Locations: Cassia County, Idaho

T13S, R24E, Sec. 1,4,9 NE near the summit of Harrison Mountain

T14S, R24E, Sec. 20 NE, 21 SW pass between Mt. Independence and Cache Peak

Soil Type: Gravelly disturbed sites or rock outcrops of granitic and quartzite substrate.

Habitat and Ecology: A low-growing, herbaceous perennial known only from the Cache Peak Range of the Albion Mountains. It is locally abundant on grassy slopes or rock outcrops of alpine areas on Harrison Mountain and Cache Peak. This plant probably does not occur on BLM lands, but it should be watched for when examining higher elevation sites. Blooms in July, with fruiting in late July through August.

Threats: Expansion of radio relay or lookout facilities

Key Characteristics


Flowers yellow

Fruits densely granular-roughened


Ultimate leaf divisions 3.5-15 mm

or more long

2-16 cm tall



BURLEY DISTRICT



A. Habit. B. Foliage leaf. C. Fruit, dorsal view. D. Fruit transection.

E. Carphophore.



Epipactis gigantea (Giant Helleborine)

Family: Orchidaceae (Orchid)

Status: Priority 2 on the State Sensitive Plant List.

Known Locations: Boise, Bonner, Boundary, Clark, Elmore, Idaho, Jerome, Owyhee, Twin Falls Counties, Idaho  
Uncommon in most of the western states in the Rocky Mountains

sites in or near Burley District

Murtaugh section of the Snake River

T10S, R18E, Sec. 3 Vinyard Creek ACEC, 12 miles NE of the city of Twin Falls, on N rim of Snake River Canyon

Soil Type: Streambanks and springs, often on calcareous sites

Habitat and Ecology: A rhizomatous orchid with one to many stems. It is restricted to streambanks, springs, and seepage areas, near thermal or cold water, often in otherwise desert regions. Often grows with monkey flowers, spike rushes, and sedges. The plant still has a broad range, but because of its vulnerable habitat, it is rapidly disappearing. It should be watched for at appropriate habitats in the Burley District. Blooms from April to July.


Threats: Development and human disturbance of cold and hot springs, livestock grazing

#### Key Characteristics

Flowers brownish-purple  
Plants up to 3 feet tall  
Leaves numerous, elliptical,  
broad, with lengthwise folds



BURLEY DISTRICT





Erigeronum ochrocephalum var. spectrum (Ochre-Flowered Buckwheat)

Family: Polygonaceae (Buckwheat)

Status: Review species on the State Sensitive Plant List.

Known Locations: Elmore, Owyhee, Twin Falls Counties, Idaho  
Malheur County, Oregon

sites in or near Burley District:

T8S, R13E, Sec. 10 SENW Yahco Creek, 8 air miles S of Hagerman  
T8S, R14E, Sec. 29 near Banbury Hot Springs



Soil Type: Loose, white, lacustrine ash deposits and heavy clays, mostly barren of vegetation

Habitat and Ecology: A perennial buckwheat of barren lacustrine slopes, typically in the Wyoming big sagebrush zone. It is usually found at elevations below 4000 feet. This variety has not been officially described by James Reveal yet, who is the current authority on Erigeronum. Collections of it are needed to determine its true taxonomic status and distribution. Blooms in June and July.

Threats: Off-road vehicles, mining exploration

#### Key Characteristics

Involucres tomentose  
Flowers cream-colored to yellow  
Taller than other similar buckweats



BURLEY DISTRICT



Gymnosteris nudicaulis (Large-Flowered Gymnosteris)

Family: Polemoniaceae

Status: Sensitive on the BLM and State Sensitive Plant Lists.

Known Locations: Elaine, Butte, Canyon, Elmore, Gem, Lincoln, Minidoka, Owyhee, Twin Falls Counties, Idaho  
Malheur County, Oregon

sites in or near Burley District

T7S, R12E, Sec. 14 west of Hagerman  
T7S, R13E, Sec. 25 3 miles south of Hagerman, 1940 record  
T7S, R15E, Sec. 24 NWNE, 27 SE, 33 NE of Wendell  
T9S, R17E, Sec. 34 1936 record, near Twin Falls  
T10S, R12E, Sec. 8 SESE, 20 NENW  
T12S, R25E, Sec. 6 1893 record, from Albion


Soil Type: Sandy to sandy loam

Habitat and Ecology: A weak-stemmed annual that grows in somewhat open, sandy areas in the basin big sage-grassland zone. Found from 2700 to 5000 feet. The plant was collected much more frequently in the early to mid-1900's than in the past 15 years. This might partly be due to its early blooming time, which can be as early as the first week of April. It may also be due to the invasion of much of its range by cheatgrass. It was once noted as common. Blooms from April to May.

Threats: Agricultural development, range improvement projects, competition with cheatgrass

Key Characteristics

Flowers showy and vary from white to yellow to lavender  
Short naked stem  
Whorl of entire leaves just beneath the flower cluster



BURLEY DISTRICT

Lepidium davisii (Davis Peppergrass)

Family: Brassicaceae (Mustard)

Status: Federal Category II (C2) recommended for Threatened status.

Known Locations: Ada, Elmore, Owyhee, Twin Falls Counties, Idaho  
Malheur County, Oregon

sites in or near Burley District

T14S, R15E, Sec. 23 NWNE, 31 NENE, 32 NESW about 4 miles S of Salmon Falls  
Dam



Soil Type: Hard bottom plays

Habitat and Ecology: A caespitose perennial found only in very hard bottom plays that are usually barren of other vegetation. They are poorly drained and often inundated with standing water early in the spring. On rare occasions a few shadscale and silver sage plants may grow in the plays. Plays are located in Wyoming big sage and fourwing saltbush habitat at 2900 to 5000 feet. One of the plays in Burley District is in low sage habitat. Populations vary in leaf shape, size, and time of flowering, and may be genetically distinct populations because of the plant's short distance dispersal mechanism and pollination vectors. Blooms from April to as late as August, depending on moisture.

Threats: Water storage pond development, spring livestock trampling, off-road vehicles, and military tanks (Boise District). Indirect threats may include siltation from range fires and rehabilitation projects.

#### Key Characteristics

Only found on plays  
White, 4-petaled flowers  
Plant can become quite woody  
Fleshy entire to toothed to  
pinnately lobed leaves



BURLEY DISTRICT

Mentzelia torreyi var. acerosa (Torrey's Blazing Star)

Family: Loasaceae (Blazing Star)

Status: Priority 1 on the State Sensitive Plant List.

Known Locations: Ada, Elmore, Gooding, Owyhee, Twin Falls Counties, Idaho  
Mono County, California  
Nevada

sites in or near Burley District

T6S, R13E, Sec. 33 NE across from mouth of Malad River, on the Snake River  
T8S, R13E, Sec. 10 SENW along Yahoo Creek, near Thousand Springs  
T8S, R14E, Sec. 29 SENW, SWSE several miles W of Buhl, on road to Banbury  
Hot Springs  
T8S, R14E, Sec. 32 historic record from 1949; 11 miles NW of Buhl  
T9S, R14E, Sec. 9 SWNE 2 miles S of Banbury Hot springs  
T9S, R14E, Sec. 10 near mouth of Mud Creek  
T9S, R15E, Sec. ? Snake River Canyon, 10 miles NW of Filer  
several sites along Salmon Falls Creek, on canyon slopes


Soil Type: Barren sandy lacustrine soil or volcanic cinder, particularly on south or west-facing slopes.

Habitat and Ecology: A caespitose perennial of barren sandy or volcanic soils from 2900 to 3500 feet in Idaho. It is typically found in the Wyoming big sage-grassland zone or shadscale zone, and often grows with white-stemmed blazing star, Indian ricegrass, and Phlox sp. It is sometimes found with Astragalus kentrophyta var. jessiae, another sensitive plant species. Southern Idaho is the northern limit of its range. In late summer and fall the plant is still easily recognized and often breaks loose, blowing around like a very small tumbleweed. Blooms May and June.

Threats: Off-road vehicles, increased agricultural development, and mining claims

Key Characteristics

Barren lacustrine slopes  
Branched spiny white  
hairy stems and leaves  
Flowers orange



BURLEY DISTRICT

Pediocactus simpsonii var. robustior (Simpson's Hedgehog Cactus)

Family: Cactaceae (Cactus)

Status: Sensitive on the BLM and State Sensitive Plant Lists.

Known Locations: Cassia, Owyhee, Twin Falls Counties, Idaho  
Colorado, Nevada, Utah, Wyoming

sites in or near Burley District

T12S, R18E, Sec. 10 SW

T14S, R20E, Sec. 21 NW, 36 NE about 16 miles SW of Oakley on Hudson Ridge

T14S, R21E, Sec. 31 NW Trapper Creek proposed RNA

T14S, R22E, Sec. 35 N edge of Middle Mountain, 6 air miles S of Oakley

T15S, R23E, Sec. 12 SW, 13 N 1/2 Graham Peak ridgeline, SE of Oakley

T15S, R29E, Sec. 16 W 1/2, 17 E 1/2 Pole Canyon proposed Research Natural  
area

T16S, R17E, Sec. 26 SW



Soil Type: Shallow rocky soils. Sometimes sandy sites

Habitat and Ecology: A typically solitary-stemmed cactus of rocky soils, benches, and canyon rims. Also known from sandy soils near the City of Rocks. It is often associated with low sage and bud sage. In Burley District it may be found with juniper, Aster scopulorum, and Haplopappus acaulis. This cactus variety is relatively widespread in southern Idaho, and even though it is somewhat protected by its habitat, it may be exploited by cactus collectors.

Threats: Commercial collectors

Key Characteristics

Only barrel cactus in  
southern Idaho  
Flowers light pink, yellowish,  
or greenish



BURLEY DISTRICT

Scutellaria nana (Dwarf Scullicap)

Family: Lamiaceae (Mint)

Status: Review species on the State Sensitive Plant List.

Known Locations: Ada, Owyhee, Twin Falls Counties, Idaho  
Northeast California  
Central Nevada  
Southeast Oregon  
Iron, Washington Counties, Utah

sites in or near Burley District

T16S, R16E, Sec. 29 SW near the Mule Creek Crossing



Soil Type: On rhyolitic gravel or shallow scabland sites associated with basalt

Habitat and Ecology: A diminutive, rhizomatous perennial in the mint family. Very striking appearance when in bloom. It grows on sites with shallow rocky soil, usually in low sage habitat, where other vegetation is sparse. Other associated species might include bluebunch wheatgrass, Penstemon deustus, and Erigeron bloomeri. It has also been found on soils sorted by stream action such as dry gravel bars along desert riparian areas. This species was put on the "Review" list at the 1988 Sensitive Plant Workshop. It has a wide range, but collections reflect that it may be uncommon. This might partly be due to its small size and harsh choice of habitat. Blooms in May and June.

Threats: None known.

Key Characteristics

Flowers cream-colored, the  
upper lip pale purplish  
Leaves elliptic, entire  
Rhizomatous  
Gravelly soils



BURLEY DISTRICT

Townsendia scapigera (Stemless Townsendia)

Family: Asteraceae (Composite)

Status: Review species on the State Sensitive Plant List.

Known Locations: Twin Falls County, Idaho  
Millard County, Utah  
California  
Nevada

Idaho

T16S, R16E, Sec. 30 N edge of Jackpot Basin


Soil Type: Dry sandy sites in the Great Basin. Ash slopes at the Idaho site.

Habitat and Ecology: A caespitose, acaulescent, biennial or short-lived perennial that grows in a broad range of habitats, from sagebrush to alpine tundra (4500-9500 feet). It has only been collected once in Idaho, in 1979. It is distinguished from T. florifer by its lack of stems. The plant apparently just gets into Idaho. Collections are needed to determine its range and extent in this state. Blooms in July.

Threats: None known

Key Characteristics

Caespitose growth form  
Flowers white to pink  
Stems lacking




BURLEY DISTRICT



LITERATURE CITED

- Atwood, N.D., S. Goodrich, S. Welsh. 1984. New Astragalus (Leguminosae) from the Goose Creek Drainage, Utah-Nevada. Great Basin Naturalist, 44(2), pp. 263-264.
- Barneby, R.C. 1964. Atlas of North American Astragalus, Vol. 1 & 2. New York Botanical Garden, Bronx, New York.
- Cronquist, A., A.H. Holmgren, N.H. Holmgren, & J. Reveal. 1972. Intermountain Flora: Vascular Plants of the Intermountain West, USA, Vols. 1, 4, 6. Hafner Publ. Co., Inc. New York & London.
- Davis, R.J. 1952. Flora of Idaho. Brigham Young Univ. Press, Provo, UT.
- DeBolt, A. & R. Rosentreter. 1988. An Illustrated Guide to the Sensitive Plants of Boise District Bureau of Land Management, 1988. BLM Tech. Bull. #88-4. 80 pp.
- Hall, J.D. & E. Lee. 1980. Inventory of Threatened and Endangered Plants - Cassia County, Idaho. Prepared for Burley District, Contract No. ID020-C10-3.
- Hartman, R.L. 1985. A new species of Cymopterus (Umbelliferae) from Southern Idaho. Brittonia, 37(1), pp. 102-105.
- Henderson, D., et. al. 1977. Endangered and Threatened Plants of Idaho. Univ. of Idaho Forest, Wildlife and Range Experiment Station. Contrib. No. 73 ISSN:0073-4586.
- Hitchcock, C.L. 1974. Flora of the Pacific Northwest: An Illustrated Manual. Univ. of Washington Press, Seattle, WA.
- Hitchcock, C.L., A. Cronquist, M. Ownbey, & J.W. Thompson. 1971. Vascular Plants of the Pacific Northwest, Vols. 1-5. Univ. of Washington Press, Seattle, WA.
- Holmgren, N.H. 1973. Five new species of Castilleja (Scrophulariaceae) from the Intermountain Region. Bull. Torr. Bot., 100(2), pp. 83-93.
- Packard, P.L., et. al. 1979. Inventory of Threatened and Endangered Plants Located in the Twin Falls Planning Unit. Prepared for Burley District, Contract No. ID-910-C19-0007. 57 pp.
- Rosentreter, R. 1986. Sensitive and Uncommon Plants in the Boise District Bureau of Land Management. BLM Tech. Bull. #86-2. 87 pp.
- Smithman, L.S. 1988. Astragalus atratus var. inseptus Training and Inventory Project. Prepared for Shoshone District. 22 pp.
- Welsh, S.L., N.D. Atwood, L.C. Higgins, & S. Goodrich. 1987. A Utah Flora. Great Basin Nat. Memoir No. 9, Brigham Young Univ., Provo, UT.



Local Descriptions of Sensitive Plants in or near Burley District

T6S, R13E, Sec. 20 SE Sec. 33 NE	Astragalus atratus var. inseptus Mentzelia torreyi var. acerosa
T7S, R12E, Sec. 14 R13E, Sec. 32 Sec. 25 R15E, Sec. 24 NWNE Sec. 27 SE Sec. 33	Gymnosteris nudicaulis Astragalus atratus var. inseptus Gymnosteris nudicaulis Gymnosteris nudicaulis Gymnosteris nudicaulis Gymnosteris nudicaulis
T8S, R13E, Sec. 10 SENW Sec. 10 SENW	Eriogonum ochrocephalum var. sceptrum Mentzelia torreyi var. acerosa
T8S, R14E, Sec. 29 SENW Sec. 29 SWSE Sec. 29 Sec. 32	Mentzelia torreyi var. acerosa Mentzelia torreyi var. acerosa Eriogonum ochrocephalum var. sceptrum Mentzelia torreyi var. acerosa
T9S, R13E, Sec. 11	Astragalus atratus var. owyheensis
T9S, R14E, Sec. 9 SWNE Sec. 10	Mentzelia torreyi var. acerosa Mentzelia torreyi var. acerosa
T9S, R15E, Sec. ?	Mentzelia torreyi var. acerosa
T9S, R17E, Sec. 34	Gymnosteris nudicaulis
T10S, R12E, Sec. 8 SESE Sec. 20 NENW	Gymnosteris nudicaulis Gymnosteris nudicaulis
T10S, R18E, Sec. 4 NENE	Epipactis gigantea
Murtaugh sec. of the Snake R.	Epipactis gigantea
T12S, R18E, Sec. 10 SW Sec. 10 SE R25E, Sec. 6	Pediocactus simpsonii var. robustior Allium anceps Gymnosteris nudicaulis
T13S, R24E, Sec. 1 Sec. 4 Sec. 4 Sec. 9 NE Sec. 9	Cymopterus davisii Cymopterus davisii Castilleja christii Cymopterus davisii Castilleja christii
T14S, R13E, Sec. 11 R15E, Sec. 7 R15E, Sec. 23 NWNE Sec. 31 Sec. 31 NENE Sec. 32 NESW R20E, Sec. 21 NW Sec. 36 NE	Astragalus atratus var. owyheensis Astragalus atratus var. owyheensis Lepidium davisii Allium anceps Lepidium davisii Lepidium davisii Pediocactus simpsonii var. robustior Pediocactus simpsonii var. robustior



T14S, R21E, Sec. 31 NW	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
R22E, Sec. 35 NESE	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
R24E, Sec. 20 NE	<i>Cymopterus davisii</i>
Sec. 21 SW	<i>Cymopterus davisii</i>
T15S, R15E, Sec. 5	<i>Allium anceps</i>
Sec. 8	<i>Allium anceps</i>
R23E, Sec. 12 SW	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
Sec. 13 N 1/2	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
R29E, Sec. 16 W 1/2	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
Sec. 17 E 1/2	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
T16S, R14E, Sec. 29 SEW	<i>Glyptopleura marginata</i>
R15E, Sec. 8	<i>Astragalus tetrapterus</i>
Sec. 7	<i>Glyptopleura marginata</i>
R16E, Sec. 29 SW	<i>Scutellaria nana</i>
Sec. 30	<i>Townsendia scapigera</i>
R17E, Sec. 30	<i>Astragalus atratus</i> var. <i>owyheensis</i>
Sec. 26 SW	<i>Pediocactus simpsonii</i> var. <i>robustior</i>
R21E, Sec. 33 N 1/2 SESE	<i>Astragalus anserinus</i>











