

INDEX

- 12S rDNA, 32, 228, 269
 16S rDNA, 32, 60, 237, 249, 269
 18S rDNA, 32, 60, 61, 158, 228, 274, 275, 285,
 304, 307, 335, 360, 366, 369, 395, 399, 402,
 468, 475
 28S rDNA, 32, 158, 278, 402, 468, 475, 522, 526
- A**
 aardvark, 638
 abdomen: function, 131; structure, 131–136
Abies, 410
Abromocomophagidae, 274
Acacia, 283
 acalyptate Diptera: ecological diversity, 540;
 taxonomy, 540
Acanthocnemidae, 391
Acaris, 99, 105–107: diversity, 101, fossils, 53,
 105–107; in-Cretaceous amber, 105, 106
Aceraceae, 413
Acerentomoidea, 113
Acherontia atropos, 585
Achias rothschildi, 541
Acontiinae, 590
Acontistoptera, 538, 539
Acrididae, 127, 205, 210
Acridoidea, 210
Acroceridae, 516, 521
Acrolophidae, 575
Acrolophus, 575
Acromis, 394
Acromyrmex, 446
Acropyga: fossil, 302, 303
 acrotergite, 121
Actias, 582
Aculeata, 415, 427, 429–463: defined, 429;
 phylogeny, 431; sting, 429–431
Adamsiana, 349
Adela, 557
Adelgidae, 293, 295, 302
Adelpha, 601
Ademosynidae, 361, 363
Adephaga, 630, 361, 363, 366–70: defining
 features, 366; hind coxae, 366
Aderidae, 390, 391: Cretaceous, 390
Adicophasma, 226
Adicophasma spinosa, 226
Adiaterothripidae, 284
Adventathrips inquilinus, 283
 aedeagus, 134
Aedes, 503, 504
Aenetus, 557
Aenigmatinae, 536
Aenne, 489
Aeolothripidae, 284, 285, 286
Aeshnoidea, 187
 African rock crawlers (see
 Mantophasmatodea)
Afroclinocera, 630
Agaonidae, 573, 616: fossil, 423
Agaoninae, 423
Agassiz, Alexander, 26
Agathiphaga, 560
Agathiphagidae, 561, 562, 567, 630
Agathis, 76
Agelaia, 439
 ages, using fossils, 37–39; using DNA, 38–40
Ageniellini, 435
Aglaeaspida, 99
Aglia, 582
Agrias, 600
Agrius (Staphylinidae), 14
Agromyzidae, 540, 623
Agrotis, 590
 Aix-en-Provence, France (Oligocene), 89
 Albertus Universität, 88
Alcides, 585
 alderflies (see Megaloptera, Sialidae)
Aleochara, 374
Aleocharinae, 374
Alepidophora peali, 525
Aleuropteryginae, 352
 Alexander, Charles P., 498
Aleyrodidae, 290
Aleyrodoidea, 290–292, 295: in Cretaceous
 amber, 292; structure, 291
Alleculidae, 389
 allopatry, 26
Alloraphidiidae, 336, 338
Alsophila pometaria, 587
Alticinae, 394
Amauris, 603
 amber, 55–61, 76, 86, 166, 270, 303, 387, 476:
 ¹⁴C dating, 55, 57; ages, 55; amino acids,
 61; ancient DNA ultrastructural
 preservation in, 58, 60; behaviors captured
 in, 57, 58; botanical origins, 55; Cenozoic
 deposits, 53, 55; chemistry/composition,
 55; copal, 55, 57; Cretaceous deposits, 53,
 54, 77–82; dating, 55; deposits, 55;
 destructive sampling, 61; Dominican, 59,
- 91; general, 57; inclusions, 57; menageries
 in, 56; Mexican, 55; parasitism in, 57;
 preservation in, 58; resinite, 55; sub-fossil
 resin, 57; symbioses in, 303; taphonomy,
 57
Ambermantis wozniaki, 259
Amblycera, 274, 278
Amblyoponini, 446, 490
Amblypygida, 99, 104, 105: in amber, 104
Amborella trichopoda, 613, 620
Ameghinoia, 450, 632
Ameletopsidae, 628
 American Museum of Natural History, 26, 87,
 91
Amesinae, 630
 ametaboly, 331
 amino acids: racemization, 61
 ammonites, 63, 642
Amorphoscelidae, 254, 257
Amphientomidae, 270
Amphiesmenoptera, 147, 468: defining
 features, 548; general, 548; stem-group,
 548, 549
Amphizoidea, 367
Ampulicidae, 451
Amydria, 557
Anabrus simplex, 209
Anacanthotermes, 244
Anactinothrips gustaviae, 283
Anagyrinus, 367
 anal veins, 129
 analogy (in characters), 28
 anamorphosis, 108, 131
Anania, 578
Anaplectidae, 231
Anareolatae, 213
Anartioptera, 147
 Anaximander, 16
Anaxyelidae, 409, 412: Cretaceous, 413
 ancestry, 29
Anchineuridae, 172
 ancient DNA, 60–61
Ancistropsyllidae, 480
Ancopteridae, 172
Ancylacheira, 382
Ancyluris inca, 598
 Andersen, Nils Møller, 317
Andesianidae, 572
Andrenelia, 419
Andreneliidae, 419

- Andrenidae, 455
Andricus vectensis, 422
Anelosimus, 467
Aneuretellus, 446
 Aneuretiniae, 446
Aneuretopsyche, 472
 Aneuretopsychidae, 472
Angaragabus, 367
Angarixyla vitimica, 410
Angarogyrus, 367
 Angarosphecidae, 452, 453
Angarosphex magnus, 453
 angiosperms, 607: ANITA group, 613; compression fossils, 609; Cretaceous flowers, 609; Cretaceous fossils, 609–10; diversity, 607; DNA phylogeny, 609; earliest fossil, 609; early growth forms, 610; evolution, 609; evolution as trees, 611; fusainized flowers and fruits, 609; fusainized remains, 609; molecular clock estimates, 38, 609; most basal species, 611; phylogenetics, 609; phylogeny, 612, 613; pollen record, 610; radiations, 5, 76, 330, 302, 607, 610; relationships to other seed plants, 607
 angiosperm radiations, significance to insects: Coleoptera, 363, 393–5, 399; Diptera, 512–3, 523; Hemiptera, 290, 298, 302–3; Thysanoptera, 287
Aniferella, 302
 Anisembiidae, 198
 Anisopodidae, 494, 498, 507, 513, 517: Cretaceous, 514
 Anisopodinae, 513
 Anisoptera, 180, 184, 185–187: Cretaceous, 187; defining features, 185
 Anisozygoptera (see Anisoptera)
 anlage (see imaginal discs)
 Annelida, 93
 Annulipalpia, 550, 551
 Anobiidae, 386, 431
 Anomalopsychidae, 554
Anopheles gambiae complex, 11
 Anophelinae, 506
 Anoplura, 274–5
 Anostostomatidae, 208
Ansorgius, 501
 ant crickets (see Myrmecophilidae)
 ant mimics (see myrmecomorphs)
 Antarcticoperlaria, 194, 195: defining features, 195–96
 anteaters and tamanduas, 638
 anteaters, as host, 272, 274
 antecostal suture, 121
 antenna: development, 123; structure, 123
 Antero Formations, Colorado (Oligocene), 88
Antheraea assamensis, 584; *mylyitta*, 584; *pernyi*, 584
 Anthicidae, 391
 Anthoboscinae, 434
 Anthocoridae, 326, 329
Anthonomus grandis, 397
 Anthophila (bees), 50, 454–563, 467, 618, 619, 638: cleptoparasitism, 458; corbicula, 461; Cretaceous, 462; cuckoo bees, 458; defining features, 454; eusocial, 461; fossil, 458, 461; fossil nests, 50, 51; phylogeny, 460, 461
 anthophily, 454, 521, 540: earliest specialized, 523; fossil pollinators, 619; generalized, 620, 621; in Brachycera, 618, 619; in Diptera, 616, 618; longest proboscis, 521; venation and hovering, 521
Anthrax, 521
Anthrenus, 387
 Anthribidae, 396
 antlions (see Neuroptera, Myrmeleontidae)
 Antliophora, 147, 468–70: defined, 468; phylogeny, 469
 ants (see Formicidae)
 Apachyidae, 220
 apatite, 42
 Aphelopinae, 433
Aphelopus, 433
 Aphididae, 89, 293, 294, 295, 301, 302: features, 294
Aphidocallis caudatus, 295
 Aphidoidea, 290–295, 467, 642: *Buchnera* bacteria symbionts, 293; classification, 292; cornicles, 293; fossils, 54, 89, 293, 295; galls, 293; heteroecy, 293; oviparae, 293; reproduction, 292; siphunculi, 293; sociality, 293; stem-group, 293; viviparae, 293
 aphids (see Aphidoidea)
Aphis maidiradicis, 301
 Aphodiinae, 378
Aphodius holderei (Scarbaeidae), 61, 642
 Aphrophoridae, 308
 Aphroteriinae, 627
 Apicotermnitinae, 248
 Apidae, 442, 454–458: extinct, 466
 Apiformes (see Anthophila)
 Apinae, 467: nests, 457, 458
 Apini, 457, 466
 Apioceridae, 521, 630
 Apiomerini, 323
 Apionidae, 396
Apis (honey bees), 9–12, 429, 454, 456, 461, 466, 541: *armbrusteri*, 461; *cerana*, 9; *dorsata*, 9, 457; *cerana nuluensis*, 11; *floreana*, 10, 458; *henshawi*, 461, 463; *koschevnikovi*, 10, 11; *mellifera*, 9, 10, 54; *mellifera capensis*, 10; *nigrocincta*, 11; relationships, 10; species concepts, 9–12
Apocephalus, 533
 Apocrita, 333, 409, 414, 415, 417, 463: mesosoma, 414; metasoma, 414; ovipositor, 430; parasitoid, 420; phylogeny, 415; propodeum, 414
 apodemes, 120
 Apoditytria, 560, 577
Apoglaeaconis ackermani, 353
Apoica, 438
 Apoidea, 430, 451–463: Cretaceous, 453; defined, 451
 apomorphy, 27
 apophysis, 120
 aposematism, 202, 385, 390, 603: in Lepidoptera, 603
 Appalachian - Asian disjunction, 237
 Appalachian mountains, 235
Apteropanopra, 476
 Apteropanorpidae, 469, 476
Apterostigma, 278
 Apterygota, 148, 150
 aquatic insects: as ancestral habit, 159; in Coleoptera, 360, 366–71, 381, 397; in Diptera, 499, 502–8; in Hemiptera, 317, 319, 321; in Mecoptera, 478; in paleopterans, 160–66, 178–87; in Polynooptera, 193, 194–6; in Trichoptera, 548–555
 Arabic teachings, 17
 Arachnida, 99, 101–7
Arachnocampa, 511
Arachnocoris, 323
 Arachnomorpha, 93, 98–107: defined, 98
 Aradidae: Cretaceous, 326, 328, 329
Aradus, 329
 Araneae, 102, 103
 Araneomorphae, 102
Arariopogon axelrodi, 522
 Araucaria, as hosts, 302, 393, 627
 Araucariaceae, 76, 396, 625, 627, 628: as hosts, 630; distributions, 627
Araucarioxylon arizonicum, 71
Araucarius, 630
Archaeofructus sinensis, 611
 Archaeopteridae
Archaeochlus, 504
 Archaeocynipidae, 422
 Archaeognatha, 12, 148–150, 628: classification, 148; defining features, 148; diversity, 148; fossils, 148, 154; mating behavior, 148
Archaeolepis mane, 556–60
Archaeopteryx, 160, 277
Archaeopteryx lithographica, 74
 Archaeoscoliiinae, 437
Archaeostephanus corae, 416
Archaeoprepona, 596
Archearadus burmensis, 328
 Archecoleoptera, 361, 362, 365
 archedictyon, 129
 Archegocimicidae, 319
Archeorhinotermes rossi, 247, 249
 Archescytinidae, 288
Archescytinopsis, 295
 archetypes, 28
Archexyela, 410
 Archexyelinae, 410
Archicnephia ornithoraptor, 506
Archiconiopteryx, 352
 Archidermaptera, 218, 220
 Archiglossata, 417
 Archihymenoptera, 410
Archimylacridia bertrandi, 233
 Archimylacrididae, 229, 233
Archiphora, 632: *patagonica*, 633; *pria*, 633; *robusta*, 633
Archips, 578
 Archipsyllidae, 269
 Archisargidae, 637
Architarbus rotundatus, 101
 Archithemistidae, 187
Architiphia rasnitsyni, 434
 Archizelmiridae, 509, 511, 637
 Archizyoptera: defined, 183
 Archodonata, 170
 Archoglossopteridae, 332
 Archostemata, 358, 360–365: features, 362; fossil record, 361, 362; relationships among families, 363, 364

INDEX

735

- Archotermopsis*, 247
 Arctiidae: caterpillar, 589; toxic host plants, 589; tymbal organs, 589
 Arctoperlaria: 194, defining features, 195
Arctotypus sinuatus, 176
 area apicalis, 213
Arega, 557
 Areolatae, 213
Areomerus, 220
 Argasidae, 106
 Argidae, 409, 410
Argyrotaenia, 578
 arista, 167, 511: convergent origins, 167, 511
 Aristolochiaceae: as hosts, 604
 Aristotle, 16
 Arixeniidae, 220, 490
 Arkansas amber (Eocene), 55
 armadillos, as hosts, 272
Armania robusta, 449
 Armaniidae, 448, 449
 Arms, Jennie M., 137
 army ants, 447
 Artematopodidae, 363
 arthrodial membranes, 128
 Arthropleona, 115
 Arthropleurida, 109
 Arthropoda, 93–118: classification, 99;
 defined, 97; invasion of land, 109–11;
 major groups, 98–118; monophyly, 98;
 relationships, 93, 107–8
 Articulata, 93
 artiodactyls, 482
Arverineura, 260
 Ascalaphidae, 336, 342, 346–7: in amber, 348;
 larva, 347, 348; roosting, 346
 Ascencio Formation, Uruguay (K/T), 92
Ascodipteron, 334, 485, 544
 Ascothoracida, 99
 Asilidae, 492, 521, 522
 Asiloidea, 521, 523
Asindulum, 511
 Asiocoelidae, 362
 Asiopteridae, 187
 Asopinae, 327, 329
 Aspidohymenidae, 172
 Aspidothoracidae, 172
 Aspidytidae, 367
 Atelestidae, 526: Cretaceous, 529
Atelestites senectus, 529
 Atelocerata, 93, 98, 99, 107, 108: defined, 107
 Ateluridae, 151
Atemele, 377
 Athericidae, 492, 522
Atheta, 374
Atopetalura, 628
Atractocerus, 387
Atrichopogon, 504, 621
Atta, 278, 443: symbiotic fungus, 445
Attacus, 584
Atticopus fimbriunguis, 102
Atteva, 557
 Attini, 277, 444: cospeciation with host
 fungus, 277; ectosymbionts of, 377; fossil,
 446
 Auchenorrhyncha, 288, 303–12, 338:
 antenna, 304; arista, 304; clypeus, 304;
 defining features, 304; earliest fossil, 307;
 infraorders, 304; monophyly, 303, 307;
 paraphyly, 288; relationships, 304–7;
 tymbal acoustic system, 304
Augochlora leptoloba, 461
 Augustine Saint, 16
 Aulacidae, 417
Aulacidea succinea, 422
 Aulaciformes, 419
Aulacita, 578
Aulacothrips dictyotus, 283
Auplopus, 435
 austral arthropods, 628–631
 austral disjunction, 41, 247, 284, 312, 313,
 478, 485, 625
 austral distributions in insects: in
 Antliophora, 485, 537, 478; in Coleoptera,
 363, 396; in Hymenoptera, 416, 431, 458; in
 Neuroptera, 341, 342, 345, 356; in
 Peloridiidae, 312–3; in Polynoaptera, 195,
 209, 211, 244, 247; in Thysanoptera, 284
 Austral Region, 396, 625: defined, 625;
 depauperate African fauna, 628
 Australembiidae, 198
 Australimyzidae, 542
Australoplatus incompertus, 397
Austrobaileya, 613, 621
 Austrobaileyaceae, 613
Austrochilus, 504
Austroconops, 504, 507, 508, 509, 630, 632:
 Cretaceous, 509; *fossilis*, 509; *macmillani*,
 508
 Austrocynipidae, 422
Austrogoniodes, 277
Austromerope poultoni, 476
 Austroniidae, 421
 Austroperlidae, 628
 Austrophasmatidae, 226
 autapomorphy, 27
 axillary sclerites, 129
 Axymyiidae, 508
Aysheaia, 94
 Azar, Dany, 80
Azteca, 302, 443, 451
- B**
Baeonotus, 505
 Baetidae, 167
 bag worms, 575
Baisomyia incognita, 506
 Baissa, Siberia (Cretaceous), 79, 80, 234, 247,
 275, 380, 398, 472, 506
 Baissidae, 419
 Baissoferidae, 551
Baissoptera brasiliensis, 339
 Baissopteridae, 336, 338, 339
 Baltic amber (Eocene), 87–88, 215, 226, 235,
 244, 271, 275, 285, 295, 302, 386, 391, 393,
 402, 403, 404, 406, 462, 463, 466, 476, 485,
 498, 504, 505, 520, 524, 536, 539, 547, 550,
 559, 572, 577, 581, 632, 633, 642
 Bardohymenidae, 172
 bark beetles: 396–8; trace fossils, 398
 bark lice (see Psocoptera)
Baronia brevicornis, 597
 Baroniinae, 597
 Barstow Formation, California (Miocene),
 45, 46
Bartonella quintana, 272
 basalare, 128
 bat, 257, 260, 272, 274, 326, 482, 583, 589,
 615, 638–639, 640: and insect ears, 639; as
 hosts, 482; as pollinators, 615; avoidance
 behavior, 583; echolocation, 639;
 ectoparasites, 641; flies, 544; fossil, 640;
 origins of, 639
 bats, insect ectoparasites of: 640–1; in
 Cimicidae, 326; in Dermaptera, 220; in
 Diptera, 543–45; in Phthiraptera (lack of),
 272–4; in Siphonaptera, 482–490
Batesia hypochlora, 601
 Bathyceridae, 465
 Batkeniidae, 183
 Batrachedridae, 577
Batrachomyia, 540
Battus philenor, 605
 bauplan, 93, 119
 Bavaria, Germany (Triassic), 71
 beaded lacewings (see Berothidae)
Beamys, 220
 bed bug, 326
 Beecher's trilobite beds, New York
 (Ordovician), 49
 bees (see Anthophila)
 beetles (see Coleoptera)
 Belidae, 396, 398
Bellicositermes, 241
 Belmont, Australia (Permian), 69
 Belmontidae, 469, 472
 Belostomatidae, 71, 268, 319, 321, 325:
 Cretaceous, 325; fossils, 47, 319; Triassic,
 325
 Bembridge Marls, Isle of Wight, UK (Eocene–
 Oligocene), 48, 89, 244, 590
 Bennettitales, 607
Beorn leggi, 97
 Bergamo, Italy (Triassic), 71
 Bering Land Bridge, 90
 Bermuda, 647
Bernaea neocomica, 290
 Berothidae, 336, 348, 353, 356, 633
 Berothinae, 356
 Bethylidae, 432
 Bethylonymidae, 430
 Betulaceae, 413
Bibiocephala, 501
 Bibionidae, 86, 493, 508
 Bibionomorpha, 496, 507–514: Cretaceous,
 511; families, 508; fossil, 509, 510, 512–14;
 larvae, 508, 511; parasitic, 512;
 phytophagous, 512; sister group, 513;
 sister group to Brachycera, 507
Bicellonycha, 385
 biogeography, 40: ecological, 40; historical,
 40; vicariance, 41
 bioluminescence, 383–386, 512: in beetles,
 383; in Diptera, 512; mechanism, 383;
 occurrence, 383
 biostratinomy, 43
 biphasic allometry, 379, 443
 birds, 4, 36, 160: dinosaur relatives, 36;
 fossils, 42, 74, 81, 85; as hosts of insect
 ectoparasites, 266, 272–80, 326, 543,
 482–90, 506, 540, 543–4
 Bishop Ussher, 23
 Bittacidae, 469, 475: nuptial feeding, 475,
 476; pretarsus, 475

- Bittacomorpha*, 499
Bittacus, 475
 Blaberidae, 231
 Blasticotomidae, 409, 410
Blatella germanica, 231
Blattabacterium, 237, 241, 242
 Blattaria, 47, 147, 199, 227, 229, 230–238:
 Cretaceous oothecae, 231; classification,
 231; fossil record, 43, 47, 231; tegmina,
 233; phylogeny, 229
 Blattellidae, 231, 235
 Blattidae, 231, 235
 Blattodea, 12, 43, 47, 147, 207, 227
 Blattopterylliidae, 223, 224
Blattogryllus karavavicus, 224
 Blattoptera, 227
Blattotermes, 244
Bledius, 374
 Blephariceridae, 493, 501
 Blephariceromorpha, 496, 501
 blood feeders, 489–491, 502, 504, 540: in
 Diptera, 499–507, 519, 522, 523, 540, 542,
 543, 545; evolution of, 489; homeothermic
 vertebrate hosts, 489; in Hemiptera, 323,
 326; mouthparts, 489; mycetomes, 489;
 occurrence in insects, 489; in Phthiraptera,
 270–80; in Siphonaptera, 480–90; Tertiary
 origins, 523
 Bluewater Creek Formation of New Mexico,
 72
 body plan (Bauplan), 28
Bohartilla kinzelbachi, 405, 406
 Bohartillidae, 405, 406
 Bojophlebiidae, 163, 164
Bolbomyia, 522
 Bolitophilidae, 511
 boll weevil, 397
 bombardier beetles, 369
 Bombini, 466
Bombus: proavus, 461; *vetustus*, 461
 Bombycoidea, 583, 584
 Bombyliidae, 492, 516, 517, 521, 523,
 525, 618, 619: fossil, 525; in amber, 525;
 larva, 521
Bombylius, 521
Bombyx mori, 584: cultural significance, 584;
 physiology, 584; silk industry, 584
 Bon-Tsagan, Mongolia (Cretaceous), 80
 book lice (see Liposcelidae)
 Boopiidae, 274
Boreallodape, 461
 Boreidae, 426, 468, 469, 475, 478–480, 488,
 630: features, 478; fossil, 479; habits, 478;
 Jurassic, 480; recent diversity, 478;
 relationship to fleas, 478, 479
 Boreoscytidae, 295
Boreus, 478, 479
 borings, 54
 Börner, Karl, 137
Borrelia recurrentis, 272
 Bostrichiformia, 386–7
 Bostrychoidea: Cretaceous, 386
 Bothrideridae, 389
 Boudreux, H. Bruce, 144
 Brachinini, 369
 Brachycentridae, 555
 Brachycera, 514–547, 618: antenna, 516;
 anthophily, 522; arista, 516; coxae and
 flight, 515; defining features, 514–516;
 evolution of adult diets, 519; flower
 feeding, 515, 519; honeydew and, 515, 523;
 infraorders, 516; Jurassic, 516, 518;
 labellum, 515, 517; larval head capsule,
 514; larval predators, 516; maxillary palps,
 515; monophyly, 514; origin of, 547;
 parasitoid, 516, 540; phylogeny, 519;
 predatory, 515, 523, 526, 529;
 pseudotracheae, 515; putative Triassic,
 516; venation, 516
 Brachycistidinae, 434
Brachypanorpa, 475
 Brachypsectridae: larva, 384
 Braconidae, 420, 427: Cretaceous, 429;
 larvae, 428
 Bradynobaenidae, 437
Bradyptericola, 581
Bradyphopha, 490, 581
 bradytely, 97
Brahmaea, 582
 Branchiopoda, 97, 99
 Branchitergalia, 162
 Branchiura, 99
Braula coeca, 544
 Braulidae, 541, 542, 544
 Brentidae, 396, 398
 bristletails (see Archaeognatha)
 Brodie, P. B., 137
 Brodiidae, 172
 Brodiopteridae, 172
 Brongniart, Charles, 137
 Brongniartiellidae, 336, 348
 brown lacewings (see Hemerobiidae)
Brownimelia clavata, 446, 450
 Brues, Charles T., 140
 Brundin, Lars, 627–31
 Buckland, William, 23
 Buffon, Comte de, 23
 bumble bees (see *Bombus*)
 Buprestidae, 358, 359, 381, 382, 415, 417:
 larva, 381; fossil, 381
Burmapygia resinata, 221
Burmazelmira aristica, 509, 510
 Burmese amber (Cretaceous), 81, 201, 234,
 247, 251, 269, 270, 290, 295, 317, 329, 387,
 398, 390, 391, 403, 500, 506, 526
Burmitembia, 199
 bursa copulatrix, 132
 bush crickets (see Orthoptera,
 Tettigonioidae)
 butterflies (see Papilioidea and
 Rhopalocera)
 Byrrhidae, 381
 Byrrhoidea, 381
- C**
 caddisflies (see Trichoptera)
 Caelifera, 202, 203, 205, 210–211:
 classification, 210; defining features, 203;
 fossil history, 210; in amber, 210;
 phylogeny, 210
Caenocholax fenyesi, 402, 406
Calcitro, 50
Caligo, 600
Callipappus, 297
 Calliphoridae, 489, 490, 492, 542, 543, 640
 Callirhipidae, 381
 Calomantispinae, as hosts, 353
 Caloneurodea, 147, 191, 193, 217, 636
 Calonymphidae, 240
 Calopterygidae, 167, 184
 Calyptrata, 542–545; defining features, 542;
 species diversity, 542
 Cambrian Explosion, 93
 camel crickets, 209
Campodea, 116
 Campodeidae, 117: in Miocene amber, 118
 Campodeomorpha, 116
Camponotus, 440, 443, 451
 Canadaphididae, 295, 302
 Canadian amber (Cretaceous), 295, 302, 329,
 403, 506, 516, 520
Canapone dentata, 446
 Cantharidae, 358, 359, 383, 384, 386
 cantharidin, 390
 Cantharoidea, 383, 384: bioluminescence,
 384–86; fossils, 386; photic organs, 384
 Canthyloscelidae, 500
 capitulum, 213
Car, 398
 Carabidae, 63, 358–59, 366, 368–70, 627, 631,
 643: ecological diversity, 368–9; fossil, 369,
 370; in amber, 370; monography, 369;
 pygidial glands, 369; subfamilies, 369;
 Triassic, 369; relationships, 369
 Carapacea, 162, 163
 Carayonemidae, 297
 Carboniferous, 40, 67–68: continental
 configurations, 68; insect fossil deposits,
 67
 Carbonopteridae, 172
Carbotriplura kukalovae, 152
 Carcinothrips, 283
 cardo, 124
 Caridae: Cretaceous, 398
 carina, 120
Carinametra burmensis, 319, 324
Carinatermes nascimbeni, 246
Carios jerseyi, 106
Cariridris bipetiolata, 449
 Carnidae, 489, 540
Carnus, 489, 490, 540
 Carpenter, Frank M., 69, 143
Carpopodus difficilis, 311
 carrion beetles (see Silphidae)
Caryedes brasiliensis, 394
 Cascade Mountains, 72, 235
Cascadilar eocenicus, 353
 Cassidini, 394
 Castniidae, 579
Casuarina, 283
Cataglyphis, 443
 Catiniidae, 363
Catocala, 582
Catopidius, 373, 490
Catotricha mesozoica, 512
 Catotrichinae, 513
 Caulopteridae, 172
Caurinus, 478
 cave crickets, 209
 caviomorph rodents, 274
 Cecidomyiidae, 57, 426, 505, 511, 512, 623:
 basal subfamilies, 513; Cretaceous, 512;
 fossil, 512; hosts, 512; in amber, 512
 Cecidosidae, 572
Cecropia, and ants, 302, 443
 Cedar Lake, Manitoba (Cretaceous), 81
Celliforma (Halictidae), 51

INDEX

737

- cellular/ultrastructural preservation, in fossils, 58, 60
- cellulose, 238
- Cenozoic, 84–85; continental configurations, 84; fossil insect deposits, 85–92; significance of, 84
- centipedes (see Chilopoda)
- Central Park (New York City), 11
- Centruroides*, 101
- Cephalocarida, 99
- Cephalorhyncha, 94
- Cephoidea, 412
- Cerambycidae, 359, 361, 392, 393, 415, 417, 440, 622, 637: diversity, 393; fossil, 393; habits, 393
- Cerambycomima*, 393
- Ceraphronoidea, 412, 415, 417, 421, 426–27
- Ceratocombidae, 318
- Ceratocombus*, 318
- Ceratophyllidae, 480, 481
- Ceratophyllum gallinae*, 482
- Ceratopogonidae, 38, 45, 489, 490, 504, 506, 508, 509, 621, 630: in Cretaceous ambers, 506, 509; stratigraphic cladistic rank correlation, 506
- cerci, 131, 161
- Cercopidae, 305, 308, 310, 540
- Cercopionidae, 308
- Cercopoidea, 308
- Cerophytidae, 391
- Cetacea, 482; hosts, 272
- Ceuthophilus*, 209
- Chaeteessa*, 254, 259, 260
- Chaeteessidae, 257, 259
- Chaeteessites*, 259
- Chagas's disease, 323
- Chalcididae: 418, 423, fossil, 424
- Chalcidoidea, 415, 423, 424: antennal sensilla, 424; defining features, 424
- Chalcoela*, 581
- Chalicodomopsis* (see *Megachile*)
- Chamaemyiidae, 426, 540
- Chaoboridae, 504
- character polarity: ontogenetic criterion, 28; outgroup comparison, 28
- charcoalified (fusainization), 49
- Charente-Maritime (France) amber (Cretaceous), 79
- Chathamiidae, 548
- Chauliodinae, 341
- Cheirolepidiaceae, 76
- Cheiromeles*, 220
- Chelicera, 98, 99, 100–107: defined, 100
- Cheliciformes, 99, 100
- Chiapas amber (Mexico) (Oligocene), 90
- Chilicola*, 461
- Chilopoda, 99, 108: Devonian, 109
- Chimaeromyrma*, 446
- Chimaeropsyllidae, 480
- Chimeromyia*, 40, 533, 536
- Chionea*, 498
- Chironomidae, 57, 489, 490, 493, 502, 504, 505, 506, 507, 627: amber fossils, 505; austral, 629; earliest fossils, 504, 505; in Cretaceous ambers, 506, 507
- Chironomus*, 491
- Chiroptera (see bat)
- Chiropteromyza*, 490
- chitin, 43–4: composition, 43
- Chloranthaceae, 613, 621
- Chlorion*, 453
- Chlorippe wilmattae*, 594
- Chloropidae, 426, 540
- Choristidae, 469, 476
- Choristotanyderus*, 497
- Chresmoda aquatica*, 74, 193
- Chresmodidae, 193, 637
- Chresmododea, 193
- Chrysidiidae, 432, 630
- Chrysididae, 430–433
- Chrysiridia*, 582, 585
- Chrysolina*, 394
- Chrysomelidae, 359, 361, 392, 393–5, 622, 637, 642: and toxins, 394; classifications, 393; fossil records, 394; trace fossil, 394
- Chrysopa*, 301, 350, 351
- Chrysoperla plorabunda*, 350
- Chrysopidae, 336, 348, 349–351, 352, 426: in amber, 352; larval, 352
- Chrysopilus*, 522
- cibarium, 124, 288
- Cicadaprobsbolidae, 308
- cicadas (see Cicadoidea)
- Cicadellidae, 57, 305, 308, 311, 312: Cretaceous, 311
- Cicadelloidea, 307, 308–12
- Cicadidae, 305, 307, 308, 310: acoustic system, 308; Cretaceous, 307; periodical, 308; tymbals, 308; tympana, 308
- Cicadoidea: nymphs, 307, 308
- Cicadomorpha, 304–8: defining features, 304
- Cicindelinae, 368, 369, 370
- Ciidae, 391
- Cimbicidae, 409, 410, 411
- Cimex lectularius*, 326: phallus, 328
- Cimicidae, 326, 328, 329, 490, 640
- Cimicoidea, 326, 329
- Cimicomorpha, 319, 321, 323, 328, 330, 622, 637: in Cretaceous amber, 328, 330
- Citheronia*, 584
- Cixiidae, 268, 304, 312
- cladistic biogeography (see vicariance biogeography)
- cladistics: congruence (character), history, 27, 28
- Cladochaeta*, 14, 135, 136
- Cladochoristidae, 469, 548
- cladograms, 31, 37: general, 35; polytomies, 37
- Cladothrips*, 465
- Clairborne Formation, Arkansas (Eocene), 86
- Clambidae, 381
- Clarkia, Idaho (Miocene), 60
- classification, 34
- Clastopteridae, 308
- Clatrotitan andersoni*, 216
- Cleistogaster*, 417
- cleptoparasitism, 277, 283, 284, 377, 540
- Cleradinae, 327, 490
- Cleridae, 358, 359, 391
- Cleroidea, 387
- click beetles (see Elateroidea)
- Clitemnestra*, 453
- Clothodidae, 197
- Clothonopsis miocenica*, 198
- clypeus, 122
- Clystopsenella*, 630
- Cnemidolestodea, 189
- Cnesia*, 505
- Cobboldia*, 62
- Coccidae, 296, 297, 299, 300, 426: feeding styles, 297
- Coccinellidae, 388: chemical defenses, 388; in amber, 388; larva, 388
- Coccoidea, 295–301, 494: archaeococcoids, 297; classifications, 297; commercial uses, 297; features, 298; fossils, 299–300; hamulohalteres, 297; in Cretaceous amber, 299, 300; labium structure, 297; male eye reduction, 298; Mesozoic, 299; neococcoids, 298; parthenogenesis, 297; paternal genome elimination (PGE), 298; phylogeny, 298; stem-group, 300
- cochineal dye, 297
- Cockerell, Theodore D. A., 140
- coelacanth, 226
- Coelolepida, 560, 567
- Coelophoridae, 576, 577
- Coenagrionidae, 184
- coevolution, 277, 624
- Coleoptera, 12, 54, 60, 147, 357–399, 467, 630, 642: ant inquilines, 372, 374, 377, 387; aquatic, 360, 366–371, 381, 397; bioluminescence, 383–86; Cretaceous, 363; cryptonephridic excretory system, 386; defining features, 357; diversification with angiosperms, 399; diversity, 357; dung beetles, 378, 397; early fossils, 360; ectoparasites, 372, 373, 380; edaphic, 371; elytra, 357; features of social insect inquilines, 377; fossil borings, 54, 398; hind wing folding, 357; horn growth, 379; Jurassic, 363; largest size, 378; larvae, 359, 364, 365, 367, 368, 378, 384, 391, 399, 400, 402, 406; male horns, 379; minute size, 373; mycangia, 388, 397; oldest stem-group, 360; parental behavior, 373, 389, 394; parthenogenesis, 374; Permian, 360, 362; phylogeny, 361; phytophagy, 393–96, 399; plastron, 370; prothorax, articulation, 357; relationships among suborders, 361; semi-aquatic, 360, 370, 371, 381; social insect inquilines, 372; sociality, 397; stridulation, 367, 378; suborders (reviewed), 360; Triassic, 363; urogomphi, 372
- Coleorrhyncha, 288, 312–14: as sister group to Heteroptera, 313; austral disjunctions, 313; defining features, 312; distribution, 312, 313; fossils, 313; relationships, 313
- Collembola, 12, 111, 113–116, 154, 628: classification, 114–16; defined, 113; spring mechanism, 113, 114; fossil record, 116
- colleteral glands, 230
- Colletidae, 454, 455
- Collocalliinae, 278
- collophore, 113
- Coloburiscidae, 628
- Colydiidae, 391
- Colymbothetidae, 367
- Commentry, France (Carboniferous), 137, 168, 172, 227
- communal behavior, 464: in Embioidea, 191, 197; in Hymenoptera, 435, 466; in Thysanoptera, 283
- comparative method, 119
- compression fossils, 43–44, 63

- Comstock-Needham system, 128
 concretions, 44–45
 condyle, 124
Condylognatha, 147, 262
Coniopterygidae, 336, 348, 351–352, 426:
 Cretaceous, 352, 353
Conoderinae, 395
Conopidae, 492, 540, 541
 continental drift: Gondwana, 627; Pangaea,
 627; sea floor spreading, 627
 convergence, 31, 32
 cooloola monster (see *Cooloolinae*)
Cooloolinae, 207
 Coope, Russell, 61
Copelatus, 369
Copepoda, 99
Coprinisphaera, 50, 380
Coptoclava longipoda, 368
Coptoclavidae, 367, 368
Coptopsyllidae, 480
 copulation, 134, 135: in *Cimicidae*, 323; in
 Dictyoptera, 256; in Odonata, 179, 182; in
 Phasmatodea, 212; and nuptial feeding, 476
 corbiculate bees, 466
 Cordaitales, 173
Cordulecerus, 346
 Cordulegastroidea, 187
Corduliidae, 628
Coreidae, 327, 329
Coreoidea, 329
Corethrellidae, 489, 490, 504
Coriphagus, 401
Corixidae, 319, 321
Cornitermes, 247
Corydalidae, 336, 340
Corydalinae, 341
Corydaloididae, 172
Corydalus, 340
Coscinocera, 584
Cosmopterigidae, 576
 cospeciation, 278, 277–80: of hosts and
 parasites, 278; of lice with hosts, 278
Cossidae, 579
Coccoidea, 579
Cossoninae, 630
 costal vein, 129
 Cow Branch Formation, Virginia (Triassic),
 72
 coxa, 125, 126
 coxopodites, 131
Crabronidae, 451, 467
 Crampton, Guy C., 140, 142
Cramptonomyia
Craneopsyllinae, 485
Cranothrips, 284
Cratomorphus, 385
Cratomyia macrorrhyncha, 83, 523, 617, 618
Cratosyrops, 353
Creaphis theodora, 292, 293
 Creede Formation, Colorado (Oligocene), 88
Crematogaster, 302, 443
Cretaceogaster pygmaeus, 516
Cretaceometra brasiliensis, 319
 Cretaceous, 76–84, 448, 449, 607–37:
 angiosperm radiations, 302, 330, 607–613;
 continental configurations, 83; drift, 627;
 fossil insect deposits, 77–84; Gondwana
 fragmentation, 627; K/T extinctions, 76,
 635–7; modern families of insects, 607
 Cretaceous ambers, 77–82, 323, 329, 391,
 498: Agapa River (Siberia), 80; Alava
 (Spain), 76; Canadian, 295, 302, 329, 403,
 506, 516, 520; Charente-Maritime
 (France), 79; Jordon, Kheta River (Siberia),
 80; Lebanon, 80; Maimecha River (Siberia),
 80; Myanmar (Burma), 80–81; New Jersey,
 82, 166, 234, 235, 246, 259, 285, 287, 290,
 295, 300, 302, 329, 339, 381, 383, 398, 632;
 Paris Basin (France), 79; Taimyr Peninsula
 (Siberia), 80
Cretamyzidae, 295
Cretaphormia fowleri, 545
Cretatermes carpenteri, 250
Cretatermitinae, 247
Cretephilatites kourios, 416
Cretevaniidae, 419
Cretocar luzzi, 398
Cretomyrma, 446
Cretomyzidae, 302
Cretoperipatus burmiticus, 94, 95
Cretophormia fowleri, 546
Cretothrips antiquus, 285, 286, 287
Cretotrigona prisca, 458, 461, 462, 466, 619
Cricetomys, 220
 crickets, 201 (see also Orthoptera, Ensifera,
 Gryllidae)
 criterion of continuation (homology), 29
 criterion of position (homology), 29
 criterion of special similarity (homology), 29
Crocinae, 342
Crosaphis, 514
 crown group, 39
Crowsoniella relicta, 363
Crowsoniellidae, 363, 364
 Crustacea, 45, 98, 99, 467
 Crustaceomorpha, 93, 107: defined, 98;
 major groups, 99
 cryptobiosis, 97
Cryptocephalinae, 51, 394: in amber, 393;
 larva, 393
Cryptocercidae, 231, 235–238
Cryptocercus, 228, 231, 235–238, 240, 241,
 252, 378, 476, 630: allogrooming, 237;
 antenna, 237; behavior, 235–6; *clevelandi*,
 236; cryptic species, 236; *darwini*, 236;
 distribution, 236, 237; *garciae*, 236;
 monogamy, 237; parental care, 237;
primarius, 236; proctodeal trophallaxis,
 236; *punctulatus*, 235, 236; relationships to
 other Dictyoptera, 237; *relictus*, 236; social
 behavior, 237; symbiotic bacteria in, 237;
 symbiotic protists in, 237, 241; *wrighti*, 236
Cryptochaetidae, 540
Cryptophagidae, 391
cryptopleuron, 203, 224
cryptorrhynchine, 396
Cryptoses, 581
Cryptothlea, 576
Ctenocephalides felis, 482
Ctenophthalmidae, 481, 485
Ctenoplectrella viridiceps, 463
Cucujidae, 389, 391
Cucujiformia, 361, 387–99
Cucujooidea, 387–91: Cretaceous, 391; fossil,
 391
Culex territans, 502
Culicidae, 490, 502, 504, 506: Cretaceous,
 506; earliest, 506; fossil, 506, 507; larva,
 503; most primitive, 506; oldest, 506;
 phylogeny, 506; proboscis, 504
Culicinae, 506
Culicoides, 504: Cretaceous, 506; *yooosti*, 509
Culicomorpha, 489, 496, 502–507: ambers,
 506; blood feeding, 502; fossil record, 505;
 disease vectors, 502; in Cretaceous amber,
 506, 507, 509; Jurassic, 505; malaria, 502;
 relationships, 504; Triassic, 505
Cuneocoridae, 317
Cupedidae, 363, 365: Cretaceous, 366;
 Triassic, 365
Curculionidae, 359, 361, 392, 396, 398, 422,
 637: in amber, 399
Curculionoidea, 363, 393, 395–99, 622:
 Cretaceous, 398; Cretaceous galleries, 398;
 fossil record, 63, 397–99; Jurassic, 397, 398;
 larvae, 395; rostrum of, 395
Curetis regula, 597
Cuterebrinae, 543
 Cuvier, Georges L. C. F. D., 23
Cycadothrips, 284, 285, 287
 cycads, 284, 608
Cyclopes, 638
Cyclorrhapha, 514, 516, 528, 531–547:
 ancestral adult diet, 522; and angiosperm
 radiations, 515; anthophilous, 522; basal
 families, 532, 534; cardia, 516;
 circumversion of male terminalia, 532;
 Cretaceous stem-group, 533;
 development, 532; early, 536; fossil larvae,
 533; fossil record, 523; muscle plaques,
 529, 533; oldest, 533; origin of, 547;
 peritrophic membrane, 516;
 pseudotracheae, 515; puparium, 532;
 saprophagous larvae, 514; saprophagy,
 514; stalk-eyed, 541; Tertiary radiation,
 526, 547
Cyclotornidae, 580
Cydia deshaiesiana, 572
Cydinidae, 327, 329
Cylindrachetidae, 211, 628, 630
Cylindrotomidae, 498
Cynipidae, 422, 623
Cynipini, 422
Cynipoidea, 415, 418, 421–422: Cretaceous,
 422
Cypselosomatidae, 547
Cyrenoherethinae, 356
 Cytochrome b (Cytb) gene, 32
 Cytochrome Oxidase I (COI) gene, 32
 Cytochrome Oxidase II (COII) gene, 32
- D**
Dacetom, 445
Dacetonini, 444
Dactylopiidae, 301
Dactylopius, 297, 301
Daiopterix olgae, 562
 Dakota Formation, Arizona (Cretaceous), 51,
 52
Dakulosphaira vitifoliae, 293
Dalceridae, 579, 580
 damselflies (see Odonata, Zygoptera)
Danainae, 602, 604
Danaus plexippus, 601, 603
 Dark Ages, 16
 Darlington, Philip J., 630, 631
 Darwin, Charles R., 24–25, 137, 643

INDEX

739

- Dasyleptidae, 148, 149, 150
Dasyleptus, 149: *brongniarti*, 150; *sharovi*, 150
Dasymutilla, 429
 Dasypodidae, 638
Daviscardia, 557
 DDT, 648–50
 deathwatch beetles (see Anobiidae)
 Declinidae, 381
Delitzschala bitterfeldensis, 170
Deltotrichonympha, 240
Dendroleon septemmontanus, 346
 Dendrothripinae, 284
 dengue, 502
 Denmark Hill, Queensland (Triassic)
Dennysus, 278
Depressaria, 557, 576
 Derbidae, 494
 Dermaptera, 12, 147, 191, 217–222, 640, 648: cerci, 218; Cretaceous, 218; defining features, 217; fossils, 220–222; hind wings, 190, 217; in amber, 221, 222; Jurassic, 220; phylogeny, 221
 Dermapteridae, 218
Dermatobia hominis, 543
Dermestes, 387
 Dermestidae: Cretaceous, 386, 387
 Dermodermaptera, 220
 Dermoptera, as hosts, 274, 275, 482
 desert locust (see *Schistocerca gregaria*)
 Deuterophlebiidae, 501
 Devescovinidae, 240
 Devonian, 66: insect deposits, 66–67; significance of, 66, 152–4
Devonohexapodus bocksbergensis, 99, 111, 112
 Diadocidiidae, 511
 diagenesis, 43
 Diamminae, 434
 diapause, 335
 Diaphanopteroidea, 147, 160, 169, 172–73
 Diapriidae, 418, 420, 421, 630
 Diaspididae, 296, 297, 299, 300, 301
 Diathemidae, 169, 171
 Diathemoptera, 170
 Dichenotomidae, 269
 Dicliptera, 147, 169, 170–71
 Dicondylia, 147, 150
 dicondylic mandibles, 150, 153
 Dictyoptera, 12, 147, 188, 191, 207, 227–260: claval furrow in forewing, 228; ootheca, 230, 231; ovipositor, 230, 233; phylogeny, 229; proventriculus, 228, 229; relationships, 228; stem-group, 227, 228, 231; stem-group roachoids, 233; tegmina, 190
Dicyndomia julianalis, 581
 digger wasps (see Apoidea)
 Dignatha, 99, 108
Dilar septentrionalis, 353
 dilarid clade, 353
 Dilaridae, 336, 348, 353
 Dilarinae, 353
 Dinmore Formation, Queensland (Eocene), 92
 Dinmore, Queensland (Triassic)
 Diopsidae, 29, 30, 88, 492, 541, 632: fossil, 543
 Diopinae, 587
 Diploglossata, 220
 Diplolepidini, 422
 Diplopoda, 99
Diploptera punctata, 231
 Diplura, 12, 111, 116–18, 154, 628: development, 116; fossil record, 117–18; relationships, 116–17
 Diprionidae, 409, 410
 Dipsocoromorpha: 317, 318, 321
 Diptera, 12, 147, 491–547, 640, 642: anthophilous, 516, 521, 523, 529, 537; antlered, 541; aquatic, 505; arista, 511, 515; as disease vectors, 491; blood feeding, 496, 502, 522, 540, 542; carrion breeding, 542; cephalopharyngeal skeleton, 514; cleptoparasitic, 540; crop, 516; dung breeding, 542; earliest, 497; ecological diversity, 491; ectoparasites, 543; evolution of larval breeding sites, 541; extreme sexual dimorphisms, 541; filtering larvae, 502; genetics, 491; halteres, 494; in experimental biology, 491; infraorders, 497; labellum, 495, 515, 517; land crab commensals, 541; largest, 516; larvae, 493; larval head capsule/evolution of, 514, 515; larval predators, 537, 540; leaf miners, 540; major evolutionary episodes, 547; major works, 491; malaria, 502; microbial grazers, 516; molecular dating, 547; mouthparts, 495; Nematocera paraphyly, 496–497; neotenic, 505; parasites, 540; parasitoids, 516, 517, 537, 540, 543; Permian stem-groups, 497; planidium, 517, 543; primitive venation, 497; torrenticolous, 501; Triassic, 497; see also Brachycera, Cyclorrhapha
 Dipterocarpaceae, 55
 Discolomidae: in amber, 391
 Dismorphiinae, 597
 Ditaxineuridae, 183
 Ditomyidae, 511
 Ditrysia 561, 565, 573–602, 637: defined, 573; tympanal organs, 583
 divergence via host plants, 624: ecological significance, 636; host races, 624; major lineages/groups, 628
 divine origins of life, 21
 Dixidae, 504, 505
Dlusskyidris zherichini, 446
 DNA lability, 60
 DNA sequences, 29–32: alignment, 30; analysis, 30; codon structure, 30; commonly sequenced genes, 31, 32; deletions, 30; gaps, 30; insertions, 30; multiple substitutions (“hits”), 31; substitutions, 31
Doa, 582
 Dobberlin, Germany (Jurassic), 560
Dobbertinia reticulata, 341
 dobsonflies (see Megaloptera, Corydalidae)
 Dobzhansky, Theodosius, 26
 Doidae, 587
 Dolichoderinae, 302, 443, 446, 467, 638
Dolichoderus, 302, 402, 443, 451
Dolichoformica helferi, 446, 449
 Dolichopodidae, 492, 517, 526, 529, 530: Cretaceous ambers, 526; earliest, 529; Microphorinae, 526
 domatia, 440
Dominibythus, 431
 Dominican amber (Miocene), 90, 91–92, 249, 271, 302, 319, 351, 386, 387, 391, 393, 406, 463, 464, 485, 536, 575, 590
Donacia, 393
 Dopa Decarboxylase (DDC) gene, 32
 Dorset, England (Jurassic), 556
 Dorylinae, 238, 377, 446, 467, 638: ectosymbionts in, 377
Dorylus, 443
Dorythrips, 284
 dragonflies (see Odonate, Epiprocta)
 Drepanicinae, 353
 Drepanidinae, 615
 Drepanoidea, 583
 Drepanosiphinae, 295
 Drilidae, 384
 dromaeosaurs, 36, 277
 Dromopoda, 99
Drosophila, 7–8, 26, 280, 334, 335, 491, 541, 644: *mauritiana*, 7; *melanogaster*, 7; *melanogaster* complex, 7, 8; *paulistorum*, 8; *sechellia*, 7; *simulans*, 7; species concepts, 7–8; *virilis*, 541; *willistoni* species groups, 8
Drosophila melanogaster, 333, 491, 532, 540: metamorphosis, 532
 Drosophilidae, 135, 136, 334, 442, 540, 541, 643: ecological diversity, 540; Hawaiian, 644; species diversity, 540
 Dryinidae, 433, 537: fossil larva, 433; in Dominican amber, 433
Dryocoelus australis, 647
 Dryomyzidae, 540
 Dryopidae, 381
Dunatothrips, 283
Dunbaria fascipennis, 170
Duncanovelia extensa, 317, 323
 dung beetles, 378
 Dunstaniidae, 308
 dustywings (see Coniopterygidae)
 Dynastinae, 379
Dyseriocrania, 557
 Dytiscidae, 46, 62, 358, 359, 367: in amber, 369

E

- earwigs (see Dermaptera)
-
- Eberhard, William G., 135
-
- Eccoptarthridae, 398
-
- ecdysial cleavage line, 120
-
- Ecdysozoa, 93, 94
-
- echidna, 638
-
- echinoderms, 642
-
- Echinophthiridae, 272, 273, 275
-
- Eciton*
- , 443, 447
-
- Ecitoninae, 238, 377, 446: ectosymbionts in, 377
-
- Eckfeld, Germany (Eocene), 87
-
- Ectognatha, 99, 111
-
- ectoparasites (of vertebrates), 489, 543, 638: and mammal rediations, 488; ctenidia, 486; defined, 489; evolution of, 489; homeothermic vertebrate hosts, 489; Jurassic, 473; by Coleoptera, 372, 373; by Dermaptera, 220; by Diptera, 543–5; by mecopteroids (extinct), 473, 479; by mites, 106; by Phthiraptera, 270–80; by Siphonaptera, 480–89.
-
- See also*
- : bats

- ectosymbionts, 377, 533
 edentates, 482
 Eichstätt, Germany (Jurassic), 42, 74
Eickwortapis, 461
 Elachistidae, 576
 elaiosomes, 213
 Elasmini, 427
 Elateridae, 358, 359, 382, 383:
 bioluminescence, 383
 Elateriformia, 361, 380–86
 Elateroidea, 380, 382, 381–83: clicking
 mechanism, 382; Cretaceous, 382; fossil
 record, 382
 Elcanidae, 208, 210, 637
Electrapis, 466
Electribius, 633
 Electrinocelliinae, 338
 Electrococcidae, 299
Electrococcus canadensis, 299
Electrolictus antiquus, 461
Electromyrmecoccus, 303
Electrosania cretica, 537
 Electrotomidae, 409, 411
 Elektraphididae, 295, 302
 Elenchidae, 406
 elephantiasis, 502
 Elias, Scott, 61
 Ellidae, 508
 Ellipsocidae, 270
 Ellipura, 111, 112
 elm (see Ulmaceae)
 Elmidae, 381
 Elmo, Kansas (Permian), 69, 175, 362
 Elongation factor 1 α (EF1 α) gene, 32
 Embiididae, 198–99
 Embiidina (see Embioidea)
 Embioidea, 12, 147, 189–191, 196–199, 432:
 blood sinuses, 196; classification, 196;
 communal behavior, 197; defining
 features, 196–97; earliest fossil, 198; in
 amber, 198
 Embioptera (see Embioidea)
 Embolemidae, 433
 Emeraldellida, 99
 Emesinae, 323
 Empididae, 618: Cretaceous ambers, 530,
 531
 Empidoidea, 526, 621, 630: Cretaceous, 529,
 530, 531; Jurassic, 526; sister group to
 Cyclorrhapha, 526
Empis, 529
 Empusidae, 257, 260
 encephalatides, 502
 endangered insects, 647, 650: and
 endangered habitats, 648; and pesticides,
 647; carrion-feeding, 648
 Enderleinellidae, 275
 Endopterygota (see Holometabola)
 Engisopteridae, 172
 Enicocephalidae, 268: Cretaceous, 315–317
 Enicocephalomorpha, 315–17, 321
Enicocephalus, 317
 Ennominae, 588
 Ensifera, 202, 203–205, 208–210, 215:
 Cretaceous, 210; defining features, 203;
 fossil history, 208; ovipositor, 210;
 phylogeny, 208
 Entognatha, 12, 99, 111–118, 135:
 fertilization, 112; relationships, 112
 entognath, 111, 123
 entomology, 16–27, 137–145
 Eobelidae, 398
 Eocene, 85–88, 637, 640: insect deposits, 637;
 significance for insects, 640
 Eodelopterum priscum, 333
 Eodermaptera, 218
 Eoditomyidae, 497, 511
 Eoichneumonidae, 429
 Eomacropis glaesaria, 461
 Eomatsucoccus: andrewi, 299; *casei*, 299;
 sukachevae, 299
 Eomerope, 476
 Eomeropidae, 469, 476
 Eoprotoneura hyperstigma, 185
 Eoptychopteridae: Cretaceous, 499, 500
 Eosentomoidea, 113
 Eotapinoma: E. macalpinei, 446, 449; *E. zherichinius*, 446
 Eotettigarcta, 308
 Eotrechus, 317
 Eoxenos laboulbeni, 401
 Ephedra, 608
 Ephemeridae, 161, 271
 Ephemeropsis melanurus, 165, 166
 Ephemeroptera, 12, 63, 147, 157, 160–7, 628:
 arista, 167; Branchitergalia, 162;
 Carapacea, 162; classification, 162; costal
 brace, 160, 164; Cretaceous, 165, 166;
 defining features, 160; fossils, 63,
 163–167; Furcatergalia, 162; naiad, 162,
 166; Pannota, 162; phylogeny, 163;
 Pisciforma, 162; Schistonota, 162;
 Setisura, 162; stem-group, 163, 164;
 subimago, 162
 Ephialtitoidea, 415, 416: Cretaceous, 416;
 Jurassic, 416
 epidemic typhus, 272
 Epidermoptera, 218, 221
Epides, 582
Epimartyia, 557
 epimeron, 125
 epimorphosis, 116
Epiophlebia: E. laidlawi, 185; *E. superstes*, 185
 Epiophlebiidae, 185
 Epiophlebioptera, 184
 epipharynx, 124
Epiphile, 601
Epipompilus, 630
 epiproct, 131
 Epiprocta, 175, 180, 184–187
 Epiipyropidae, 580
 episternum, 125
 epistomal sulcus, 120, 122, 224
Eptesicus, 639
 Erasipteridae, 174
Erasipterooides valentini, 177
Erasmia, 578
 Eremiaphilidae, 254, 257
 Eremochaetidae, 637
 Eremonera, 526: earliest, 526; evolution of
 adult diets, 528; evolution of larval diets,
 528; Mesozoic, 527; phylogeny, 528
 Erinaceidae, 638
 Eriococcidae, 296, 299
 Eriocraniidae, 568
Eristalis, 540
 Erythrolchnia, 385
 Escherich, Karl L., 140
 Eteoanisoptera, 185
 Euaculeata, 430
 Eubleptidae, 172
 Eubleptidodea, 172
 Eubleptoptera, 169, 172
Euchauliodes, 341
 Euchaulioididae, 336, 341
Eucheira socialis, 597
 Euchelicerata, 99
 Eucinetidae, 361, 381
Euclea, 578, 580
 Eucnemidae: Cretaceous, 382, 383
 Eucoilidae, 420
 Eucrustacea, 99
 Euephemeroptera, 164
 Eugeronidae, 173
 Eugeroppteridae, 174
Egeropteron, 174
Euglossa moronei, 461
 Euglossini, 456, 461, 466
 Euherbstiinae, 455
Euhoplopsyllus glacialis, 482
 Euhymenoptera, 409, 413–414, 463: defining
 features, 413
Eulaema meriana, 456
Eulonchus, 521
 Eulophidae, 427
Eumaeus godartii, 597
 Eumalacostraca, 99
 Eumastacidae: 205, in amber, 210
 Eumastacoidea, 211
 Eumegasecoptera, 169, 171, 172
 Eumeninae, 435, 437
 Eumenetola, 147, 188, 192
 Euparagiinae, 437
Eupatorium, 614
 Euphasmatodea: defining features, 213
Eupithecia, 586
 Euplectoptera, 163, 164
Euptychia insolita, 601
Eurhynheria, 578
Eurybia patrona, 600
 Eurypterida, 99, 100
Euselasia, 598, 623
 eusociality, 238, 252, 283, 463, 464–467: in
 thrips, 283
 Eustheniidae, 196, 628
 Euthemistidae, 187
 eutherian mammals, as hosts, 275
 Euthymenoptera, 413
 Euzygentoma: defined, 151, 152
 Evaniidae, 417, 418, 419
 Evanioformes, 419
 Evanioidea, 415, 417–419
Evergestis, 578
 eversible vesicles, 117, 131, 132, 149
 evolutionary stasis, 100
 evolutionary success of insects (see species
 diversity in insects)
 exite, 159
 exoskeleton, 646
 extensor muscles, 126
 extinctions, 36–37; modern, 647–650
 extinction rates, 636–8, 647
 extrinsic muscles, 126
 eye structure, 124

INDEX

741

- F**
 Fabricius, J. C., 19, 21
 Fagaceae, 237, 422, 610
 Fagales, 422
 Falkland Islands (Devonian), 67
 Falsiformicidae, 437, 637
 Fatjanopteridae, 338
 Fauriellidae, 284
 Fedtschenkiinae, 435
 feeding damage, 52
 Felsenstein Zone, 32
 female genitalia, 136
 femur, 125
Feniseca tarquinius, 598
 Fergana Valley (Madygen Formation, Triassic), 71
 Fergusoninidae, 540
Ficus, 283, 423
 Fideliinae, 455
 Fideliini, 630
 Figitidae, 422
 figs, 616; fig wasps (Agaonidae), 616
 fire flies (see Lampyridae)
 firebrat (see *Petrobius*)
 Fisher, Sir Ronald, 26
 fishflies (see Chauliodinae)
 fleas (see Siphonaptera)
Fletcheriana triassica, 309
 flexion lines, in wing, 131
 flexor muscles, 126
 flies (see Diptera)
 flight, 155–157, 195; asynchronous flight muscles, 157; indirect flight muscles, 156; origins, 158–160, 195; synchronous flight muscles, 157
Florinemestrius, 523, 616
 Florissant Formation, Colorado (Eocene-Oligocene), 86, 87, 139, 310, 339, 345, 386, 393, 525, 545, 546, 590, 593, 594, 642
 flowers: fossils, 50
Folindusia kemaensis, 551, 554
 folk taxonomy, 15
 fontanelle, 247
Fontecilla, 349
 Foraminitermitinae, 248
Forcipomyia, 504, 507
 Foremost Formation, Canada (Cretaceous), 82
 Forficulidae, 218, 222
Formica, 300, 443, 451
 Formicidae, 18, 301, 437, 440, 467, 638; castes, 443; chemical communication, 443; cocoon, 441; Cretaceous, 448, 450, 452; eusociality, 447; fossil, 85, 303, 446, 448; and Hemiptera, 300–2; metapleural gland, 441; phylogeny, 447; proventriculus, 301; structure, 441
 Formicinae, 302, 443, 446, 467, 638: Cretaceous, 451
 fossil, 36–41, 63, 92, 523, 631: dating and ages, 62–65; defined, 62
 fossil insect deposits, 65–92
 fossil preservation, 43–63
 fossilization, insect: amino acids, 61; calcium phosphate, 46; carbonates, 44; celestite, 45; cellular preservation, 48, 59, 60; charcoalification, 49–50; compressions/impressions, 43; concretions, 44; coprolites, 50; encapsulation, 49; frass, 50; frozen remains, 61; fusainization, 49; goethite, 46; gypsum, 46; ichnofossils (see trace fossils); marcasite, 49; mineralized replication, 45; of chitin, 61; of epicuticular waxes, 44; of protein, 44; of spores/pollen in guts, 44; Orsten preservation, 45; pack rat middens, 61; permineralization, 45; phosphatization, 46; Pleistocene beetles, 63; proteins, 44; pyrite, 49; pyritization, 49
 fossorial forelegs, 29, 31
 fossorial insects, 31
Franklinothrips, 284
 frog hoppers (see Cercopoidea)
 frogs, 489, 540, 502
 fruit bats, 615
 fruit flies (see Tephritidae, Drosophilidae, *Drosophila*)
 Fulgoroidea: Cretaceous, 306, 311, 312
 Fulgoromorpha, 304, 306, 312: defining features, 304
 fungus gnats (see Sciaroidea)
 Fur Formation, Denmark (Paleocene), 85, 590
 Furcatergalia, 162, 163
 furculum, 114
- G**
 galea, 124
 Galerucinae, 394
 gall wasps (see Cynipidae)
Galleria, 581
Galloisiana, 222
 galls, 54, 283, 289; fossil, 49, 53, 54
 gall-making insects: in Diptera, 512–3, 540–1; in Hymenoptera, 422; in Lepidoptera, 572; in Sternorrhyncha, 289, 293; in Thysanoptera, 283–4
 Gaspé Bay, Canada (Devonian), 66
 Gasterophilinae, 543
 Gasteruptiidae, 417, 418, 630
 Gaston, Kevin, 13
Gastrotheus, 628
 Gastrotricha, 93, 94
 Geadephaga (see Carabidae)
Geijera, 283
 Gelastocoridae, 319, 321
 Gelechiidae, 576
 Gelechioidea, 201–2: coumarins and, 576; diversity, 576; fossil, 577; larval predators, 577; toxic host plants, 576
Genaphis, 295
 genes, 32
 genitalic structure: 132–136, complexity in, 135–136
 Geocoridae, 327
 Geocorisaee, 319
 geographical extinctions, 339, 345, 355, 466, 501, 545, 632, 633, 634: and competitive exclusion, 634; and paleoclimatic change, 634; and vertebrates, 634; in amber, 632; in Baltic amber, 632; in Quaternary insects, 642
 Geological Time Scale, 62, 64
Gelothrips, 284
 Geomagnetic Polarity Timescale (GPTS), 65
 Geometridae, 583, 585, 586: aptery, 587; fossil, 588; in amber, 568; larvae, 587
 Geometroidea, 585–89
Geomydoecus, 277, 280
 Geomyidae, as hosts, 278, 279
Geoscaphius, 237
 Geotrupidae, 379
Geranomyia, 498
 Geraridae, 216
Gerarus danielsi, 216
 Germar, Ernst F., 137
 Gerocynipidae, 422
Gerontoformica cretacica, 449
 Geroptera: defining features, 147, 160, 174
 Gerridae, 193, 268, 317, 324; fossil, 324
 Gerromorpha, 193, 317–19: families of, 317; fossils, 324; monophly, 317; pelagic, 317; phylogeny, 321; pretarsal structure, 317, 323
 giant lacewings (see Polystoechotidae)
 Giebel, Christoph G. A., 137
 gigantism, 178
 Gigantopteridaceae, 52
 Gigasiricidae, 409, 413
 Gilboa, New York (Devonian), 66
 Gill Theory, 159
Glabellula brunnifrons, 525
Glaesoconis baliopteryx, 352
Glaucoma xerxes, 649
 glossa, 124
 Glossata, 29, 562–564: basal families, 564; oldest, 566
 Glosselytridae, 332
 Glosselytrodea, 147, 158–60, 331–333, 337, 360, 636
Glossina oligocena, 544, 545
 Glossinidae, 87, 326, 490, 492, 543, 544, 545, 632: and sleeping sickness, 5; fossil, 545
 Glossophaginae, 615
 Glossopteridae, 332
 Glossopteris seed ferns, 52
 Glossosomatidae, 549, 550
 glow worms, 511
 gnathal pouch, 112
 gnathobase, 98
 Gnetales, 607, 609: Cretaceous, 611; pollinating insects, 608
Gnetum, 608, 609
Goliathus, 378
 Gomphidae, 182
 Gomphoidea, 187
 gonangulum, 134, 229
 gonapophyses, 132, 177, 229
 Gondwana, 41, 76, 627: continental configurations, 626; Cretaceous drift, 627
 Goniodidae, 275
 gonocoxae, 132, 135, 177, 229
 gonoplac, 134, 229
 gonoplacs, 229
 gonopore, 133
 gonostyli, 132, 135, 161
Gracillaria, 557
 Gracillariidae, 52, 573, 575: Cretaceous leaf mine, 575; fossil mines, 575
 Gracillarioidea, 14, 575–76
 Gramineae, 412
 Grammosmylidae, 336
 graptolites, 63, 642
 grasses, 247, 638

grasshoppers (see Orthoptera, Caelifera, Acridoidea)
 grasshoppers, 201
 Grassy Lake, Alberta, 82
Grauvogelia arvilleriana, 497
 Greeks (Ancient), 16
 green lacewings (see Chrysopidae)
 Green River Formation, U.S., 85, 590
 Grès-à-Voltzia, France (Triassic), 71, 497
Gryea, 573
 griffenflies (see Protodonata)
 Grimen, Germany (Jurassic), 74, 560
Grimmenaphis, 295
 Gripopterygoidea, 195
Gromphadorhina portentosa, 231
 Grube Messel, 44, 406
 Grylacrididae, 205, 208, 209
 Grylacridinae, 208
 Gryllida, 190
 Gryllidae, 204, 205, 209, 643: Cretaceous, 204, 209; foretibial tympanum, 204; mirror in wing, 204; wing, 204
Grylloblatta: 222, 223, *washoa*, 225
Grylloblattella, 222
 Grylloblattidae, 44
Grylloblattina, 222
 Grylloblattodea, 12, 147, 188, 191, 222–224, 225, 630: defining features, 222; fossils, 223; stem-group, 224
 Grylloidea, 209
 Grylotalpidae, 205, 208, 209, 628
 gula, 123
 gypsy moth, 589
 Gyrinidae, 358, 367
 Gyropidae, 274

H

Haeckel, Ernst, 137, 138
Haematobia, 489
Haematomyzus, 273, 274: *elephantis*, 274: *hopkinsi*, 274: *porci*, 274
 Haematopinidae, 273, 275
Haetera macleannanii, 600
 Hagen-Vorhalle, Germany, 67, 177
 Haglidae, 208
 Hagloidea, 208, 209
Haidomyrmex cerberus, 446, 449, 452
 Haldane, J. B. S., 26
 Halictidae, 453, 461, 463, 465, 614
 Halictinae, 2, 455, 467
 Halictophagidae, 401
Halictus savenyei, 461, 465
 Halimococcidae, 297
 Haliplidae, 367
Hallucigenia, 94
Halobates ruffoi, 319
Halobates, 317
 halteres, 494: convergent origins, 494; function, 494
 Halteria, 402
Hamadryas, 602
Hamamelis, 237
 Hamophthiriidae, 274, 275
 hamuli, 408
 Handlirsch, Anton, 137, 139
 Hanidae, 172
 haplodiploidy, 397
 Haplodeninae, 346
 Haplodiidae, 313

Haptopoda, 99
 Harpalinae, 368
 Hartigiini, 412
 Harvestmen (see Opilliones)
 Hawaiian Islands, 643–44, 647: conveyor belt biogeography, 645; endemism, 643
 head segmentation: criteria, 123
 head structure, 121–124
 Hebridae, 317, 321
 hedgehogs, 638
 Hedyliidae, 583, 590, 591: relationships, 590
 Heer, Oswald, 89
Heidea cretacea, 290
Heledodora, 94
 Heleomyzidae, 542
 Heliconiinae, 277, 600, 604
Heliconius, 277, 602, 605, 624: *cyclone*, 601; *erato*, 605; *numata*, 605; mimicry rings, 605; polymorphism, 605
 Helicopsychidae, 550
Heliozela, 557
 hellgrammites (see Megaloptera, Corydalidae)
 Helodidae, 381
 Heloridae, 418, 421
 hematophagy (see blood feeders)
 Hemerobiidae, 336, 348, 351, 426
 Hemerobiiformia, 336, 342, 348–356: defined, 348
Hemeroplanes, 586
 Hemimeridae, 220, 490
 Hemimerina, 220
Hemimerus, 220
 hemimetabolism, 331, 332
 Hemiplebioidea, 632
 Hemiptera, 12, 147, 193, 287–330, 622, 630, 642: alimentary filter system, 288; feeding, 288; honeydew, 288; mouthparts, 264, 288; Permian origins, 288; suborders, 288
 Hennig, Willi, 26, 27, 80, 143, 144, 145
 Hennigmatidae, 499
 Hepialoidea, 569
 Heptageniidae, 161
 Heptapsogasteridae, 273, 275, 277
 herbivory: see phytophagous insects
 Hermatobatidae, 317
 Hesperiidae, 591, 592, 595
Hesperoboreus, 478
 Heterobathmiidae, 560, 561, 562
Heterocephalus glaber, 465
 Heteroceridae, 381
Heterodoxus longitarsus, 275
 Heterogynaidae, 451
Heterojapyx, 116
 Heteroneura, 560
 Heterophlebiidae, 187
 Heteroptera, 48, 288, 314–30, 622, 640: blood-feeding in, 323, 326; defining features, 314, 315; earliest fossils, 315; fossils, 315, 318, 320, 325–30; infraorders, 315, 321; myrmecomorphy, 323; parental care, 319, 323; phylogeny, 321; phytophagy, 326, 330; relationships, 315; scent gland, 314; spider-web cleptoparasites, 328; traumatic insemination in, 326; wings, 268
Heterosarus eickworti, 461
 Heterothripidae, 284, 285
Hexagenia, 162

Hexapoda, 87–88, 98, 99, 111: defined, 111; early fossils, 111; hierarchical classification, 111
 higher termites, 247
 high-resolution CT scanning, 47, 259
Hilara, 529
 Hilarimorphidae: Cretaceous, 525
Hilarimorphites yeatsi, 525
 Hinton, Howard E., 142
Hippeococcus, 301
 Hippoboscidae, 272, 492, 543
 Hispinae, 394
 Histeridae, 372
Hodotermes, 244
 Hodotermitidae, 240, 242, 244, 246, 249, 250, 251: alates, 250; diet, 244; fossils, 244, 246
Hodotermopsis, 247
Holcoroebeus nigrimontanus, 380
Holcoroebeus, 380
Holcorpa, 475, 476
 Holodonata, 147, 174–75, 180
 Holometabola, 12, 147, 188, 333–35: adaptationist, 334; defined, 331; genital discs, 333; imaginal discs, 331; larva, 331; larva control of development, 335; origin of, 333–335; pairs, 333; pronymph, 333; similarities between, 334
 Holoptilinae, 326
Homalocnemis, 526, 630
 Homiopteridae, 173
 homology, 28–31: behavioral, 30; criteria, 29, 30; molecular, 29; primary, 29; secondary, 29; topological, 29
 homoplasy (defined), 31
 Homoptera, 287
 honey bees (see *Apis*)
 honeycreepers, 615
 honeydew, 288, 289, 290, 293, 297, 300–2, 327
 honeyeaters, 615
 Hooker, Sir Joseph, 626
 Hoplocarida, 99
 Hoplopleuridae, 273, 275
Hoplodarium, 313
 Hormaphididae, 467
 Hormaphidinae, 293, 465
 hornets (see Vespidae)
 horntails (see Siricidae)
 host races: and cryptic species, 624; conditions for, 624; examples, 624
 hover wasps (see Stenogastrinae)
 hovering: ecological significance, 618
 human bot, 543
 humans, as hosts, 274
 humeral plate, 129, 130
 hummingbirds, 615
 Hyatt, Alpheus, 137
 Hybophthiridae, 274, 275
 hybrids, 8
 Hydradephaga, 361, 363, 366–68, 357–407: Mesozoic, 368
 Hydraenidae, 371, 372
 Hydrobiosidae, 550
 Hydrometridae, 317, 319, 321, 324: Cretaceous, 324
 Hydrophilidae, 363, 372
 Hydrophiloidea, 372
 Hydropsychoidea, 549, 551
 Hydroptilidae, 549, 551
 Hydroscaphidae, 370

INDEX

743

Hygrobiidae, 367
Hymenaea (Leguminosae), 55, 91
Hymenoepimecis argyraphaga, 427
 Hymenopodidae, 256, 257, 260
 Hymenoptera, 12, 147, 407–409, 413, 463, 623, 630, 642; cenchri, 408; defined, 408; diversity, 407; ectoparasitoids, 408; endoparasitoids, 408; hamuli, 408, 409; haplodiploid, 408; hyperparasitoid, 408; parasitoidism, 409, 413; parasitoids, 408; phylogeny, 409; superparasitism, 426; Triassic, 410
 Hymenopterida, 147
Hymenopus coronatus, 255
 hyperparasitoids, 408
Hypocliffea cuspidata, 301
 Hypodermatinae, 543
 hypognath, 121
 Hypoperlida, 44
 hypopharynx, 124
 hypoproct, 132
 hypotheses, 93
 Hyracoidea, 482
 Hystricopsyllidae

I
 Ibalidae, 422
Icaronycteris index (Chiroptera), 85
 ice crawlers (see Grylloblattodea)
 Ichneumonidae, 418, 427, 630; larva, 428
 Ichneumonoidea, 415, 418, 423, 427; fossils, 428, 429; symbiotic viruses, 427
 ichnofossils (see trace fossils)
Idea, 601
 Idiogastra, 413
 Idiolothropiniae, 283
Illacme plenipes, 109
 imaginal discs, 331, 333–334, 379
 impression fossils, 43
 inchworms (see Geometridae)
Incurvaria, 557
 Incurvarioidea, 371–99, 572
 Indriidae, as hosts, 275
Inka minuta, 299
 Inkaidae, 299
 Inocelliidae, 336, 338
 inquilines, 326
 insect, 119, 144: classification, 147; defined, 137; ecological significance, 4–6; flight, 67; orders, 137; ordinal relationships, 146; species diversity, 11–15, 650
 insect diversification: end-Permian extinctions, 636; families over time, 636; rates of first and last appearances, 636; significance of Cretaceous, 636; taxic analyses, 636
 insect fossilization, 42; cuticle, 43; epicuticular waxes, 44; gut, 44; of chitin, 43; of protein, 44; of spores and pollen, 44; Pleistocene elytra, 63
 insect morphology, 119–137
 insect phylogeny, 30, 31, 40, 146, 154: Aculeata, 431; Anthophila, 460, 461; Antliophora, 461; Apocrita, 415; Blattaria, 231; Brachycera, 519; Caelifera, 210; Coccoidea, 298; Coleoptera, 361; Culicidae, 506; Dermaptera, 221; Dictyoptera, 229; Ensifera, 208; Ephemeroptera, 163; Eremoneura, 528;

Formicidae, 447; Heteroptera, 321; Hymenoptera, 409; spindle phylogenies, 636; Isoptera, 249, 251; Lepidoptera, 560, 561; Mandibulata, 108; nematocerous Diptera, 496; Neuroptera, 150; Neuropterida, 336; Odonata, 163, 169; Odonatoptera, 180, 181; Orthopterida, 202; Panarthropoda, 93; Panorpida, 469; Paraneoptera, 263; Plecoptera, 194; Polynooptera, 191; Symphytans, 409; Trichoptera, 544, 550; Thysanoptera, 286; Zoraptera, 199–201
 insect sociality, 237, 238, 283, 293, 372, 377, 397, 464, 533
 insectivores, 482
 Institut und Museum für Geologie und Paläontologie, Göttingen, 88
 Integripalpia, 548, 550
 intercalary segment, 123
 intertidal, 504
 intima: proctodeal trophallaxis, 240
 intrinsic muscles, 126
 Introverta, 94
Iotatubothrips, 283
Ips, 396
 Ipswich Series, Australia (Triassic), 72
Iridomyrmex ants, 443, 580
Ironomyia, 536
 Ironomyiidae, 536, 632
 Ischnocera, 274, 275, 277
 Ischnopsyllidae, 481, 482, 640
 islands faunas, 647: and dispersal ability, 644; and niche expansion, 645; fragility, 647; Galapagos Islands, 643; Hawaiian, 644; reasons for endemic diversity, 644–45
 Isle of Wight, UK (Eocene-Oligocene), 47: Bembridge Marls, 47; marcasite, 49; pyrite, 49; pyritization, 49
 Isophlebiidae, 187
 Isoptera, 12, 51, 55, 147, 238–251, 467, 638: Alates, 250; ancestral, 252; and humification, 241; and mineralization, 241; arboreal nest, 249; biomass, 241; castes, 238; cellulose consumption, 241; Cretaceous fossil nest, 248; diet, 238; diversity, 242; ecological significance, 241; evolutionary trends in, 250; fecundity, 238; fossil nests, 51, 248; fossilized workings, 55; fossils, 242; lower termites, 242; methane production of, 241; mounds, 242; nasute soldiers, 250; neotenic reproductive, 238; origins, 251; phylogeny, 249, 251; physogastry, 238, 240; sex-linked translocations, 238; sociality, 238; soil excavation, 242; soil feeders, 241; symbiotic bacteria of, 238; symbiotic protists of, 238, 240–1
 Isotomidae, 116
 isotope dating: ^{14}C (carbon), 63; ^{40}K - ^{40}Ar , 63; ^{235}U – ^{207}Pb , 63
 Issidae, 2, 306
 Issyk-Kul', Kazakhstan (Triassic), 71
 Ithomiinae, 614
 Ithoniidae, 336, 348, 349
 Izarra, Spain (Miocene), 45

J
Jantarostylops kinzelbachi, 406
 Japygidae, 118

Japygomorpha, 116
 Jarmilidae, 164
 Jascopidae, 312
 Javelina Formation, 247
 Jeannel, René, 627, 631
Jeholodens, 81
Jersicoccus kurthi, 299
Jersimantis, 259
 Jerusalem crickets, 209
 Johnston's organ, 137
 Jordanian amber (Cretaceous), 80
 jugal bar, 192
 Juglandaceae, 410
Juraconiopteryx zherichini, 352
Juraphis, 295
 Jurassic, 73–76: continental configurations, 73; insect deposits, 74–76; significance of, 73
 Jurassic Park, 60
Jurina marginata (Jurinidae), 333
 Jurinidae, 332
 juvenile hormone (JH), 334, 379

K
 K/T event, 635: Chicxulub crater, 635; insect family survivorship, 637
Kalligramma, 74
 Kalligrammatidae, 75, 336, 347, 348, 637
Kalotermes swinhoei, 247
 Kalotermitidae, 238, 240, 242, 247, 248, 250, 251; Cretaceous fossil nest, 248; diversity, 247; fossils, 247; pseudergates, 238
 Kaltanidae, 469, 472
 Kansasiidae, 171
Karabasia evansi, 313
 Karabasiidae, 313
 Karabastau Formation, 75
 Karajassidae, 312
Karanabis kiritshenkoi, 317
Karataothrips jurassica, 285, 286
 Karatau, Kazakhstan, 74, 75, 220, 270, 380, 397, 398, 416, 526, 560
Karataublatta longicaudata, 233
 Karatawiidae, 187
Karoophasma bielouwensis, 225
 Karschielloidea, 218
 katydids (see Orthoptera, Tettigonioidae)
Kazachothrips triassicus, 285, 286
Keithia luzzii, 299
Kelita, 458
Kemperala, 189
 Kennedyidae, 183
 Keroplatidae, 511: Cretaceous, 512; larva, 511
Kerria lacca: historical significance, 297
 Keuper Basin (Triassic), 71
 king crickets, 209
 Kinorhyncha, 93, 94
 Kishenehn Basin, Montana (Oligocene), 88
Kladothrips, 283, 467: *harpophyllae*, 284
 Klapálek, Franz, 143
 Klondike Mountain Formation (Eocene), 86
 Kojutellitinae, 419
 Kokiriidae, 555
 Königsberg, 88
 Koonwarra, Australia (Cretaceous), 84, 319, 485, 487
Koruga, 240

- Koshelevka Formation, 69
 Kotá Formation, India (Jurassic), 76
 Kotujellitinae, 419
Kovalevomyia lacrimosa, 505
 Krausse, Anton, 140
 Kristensen, Niels P., 144
Krombeinictus nordenae, 453
 Kukalová-Peck, Jarmila, 144
Kyromyrmex neffi, 446, 449, 451
- L**
 La Brea, California, 61, 62, 90
 La Cabrua, Spain, 79
 La Voulte-sur-Rhône, France (Jurassic), 49
 Labandeira, Conrad, C., 52
 labial palpus, 124
Labiococcus joosti, 299
 labium structure, 124
 labrum, 122, 123
 lacewings (see Neuroptera)
 Lachesillidae, 270
 lacinia, 124
 Lacturidae: aposematic, 579
 lacustrine (deposits/preservation), 42
 ladybugs (see Coccinellidae)
 Laemobothriidae, 274
Laetilia coccidovora, 301, 581
 Lagerstätten: Konservat, 42; Konzentrat, 42
 Laiyang Formation, 81
 Lalacidae, 311, 312
 Lamarck, Jean Baptiste, 23
Lamproblatta, 237
 Lampyridae, 383, 384, 386: bioluminescence, 384–385; femme fatale, 385; in amber, 386; light signals, 385; photic organs, 385
 Languriidae, 490
Laphria, 522
Larropsis, 453
 larva, 331, 333–335: fossil, 346, 348, 352, 365, 377, 378, 381, 384, 386, 388, 391, 393, 533, 566, 575, 581, 588; in Coleoptera, 359, 364; 394; in Diptera, 493, 503, 505, 511, 515, 521; in Lepidoptera, 565, 574, 580, 585, 589, 595, 597–601; in Neuroptera, 342, 344, 347, 349, 350; in Siphonaptera, 484; in Strepsiptera, 401; in Trichoptera, 551
 larval cases: fossils, 394
 Las Hoyas, Spain (Cretaceous), 79
Lasia, 521
Lasiocampa, 582
 Lasiocampidae, 583
 Lasiochilidae, 329
Lasioglossum, 14
Lasiohelea, 504
Lasius neoniger, 301
Lasius, 443: *neoniger*, 301
 Latah Formation, Idaho (Miocene), 89
 Lathiceridae, 210
 Lathridiidae, 391
Latimeria, 97
 Latreille, Pierre André, 21
Latrodetus, 103
 Laurasia, 76
Laurhervasian setacea, 344
 leaf beetles (see Chrysomelidae)
 leaf hoppers (see Cicadelloidea)
 leaf mines, 52, 575: fossil, 52
 leaf-cutter ants (see Attini)
 leaf-cutter bees (see *Megachile*)
 leaf-insects, 211
 Lebanese amber (Cretaceous), 80, 247, 270, 285, 295, 312, 317, 398, 499, 506
Lebanophlebotomus, 499
 Lebiinae, 369
 leg structure, 125–128
 Leidy, Joseph, 240
 Leiiodidae, 335, 371, 372
Lemmatophora typa, 196
 Lemmatophoridae, 194, 195, 196
 lemurs, as hosts, 275
 Lentulidae, 210
 Lepiceridae, 370
 Lepidodendrales, 173
Lepidophthirus macrorhini, 272
 Lepidopsocidae, 332
 Lepidoptera, 12, 147, 333, 469, 548, 555–606, 618, 622, 642: ant-enticing organs, 598; aptery, 572, 583; aquatic, 578; austral distributions, 562, 572; basal relationships, 558, 560–2; bat detection, 583, 589; blood feeding, 590; brachypterous, 586; butterfly mimics, 576; caterpillar crochets, 570; Cretaceous, 560, 562, 572; crochets, 565, 570; cyanide production, 579–580; detritivorous, 581; earliest fossil, 556; ectoparasitic larvae, 577; ectoparasitoid larvae, 580; evolution of larval diets, 558; external feeders on plants, 573; female reproductive systems, 569–71; fossil leaf mine, 572; fossilized wing scales, 570; frenulum, 572; galls, 572; head features, 567; Heteroneuran, 571; in Cretaceous amber, 563; Jurassic, 556, 560, 561; largest in amber, 588; leaf mine fossil, 572; leaf miners, 558, 567; leaf-mining, 572; Mesozoic fossils, 556–61, 562, 566; mimics, 579; monographs, 556; monotrysian, 570–2; most diverse phytophages, 556; myrmecophily, 599; parthenogenesis, 575; phylogeny, 558, 561; popularity, 555; predators, 558, 598; primitively mandibulate, 561–62; proboscis functional morphology, 562; proboscis structure, 563, 564, 567; prolegs, 563, 581; pupa, 577; pupae, 568; radiation with angiosperms, 555; reproductive system structure, 569, 571; retinaculum, 571, 594; root borers, 579; scale structure, 567–9; silk production, 584; silken galleries, 569; smallest, 572; snake mimics, 586; stem, 579; triangula, 580; tympanal organs, 586, 583, 585, 586; wasp mimics, 579; wing scale colors, 567; wing scale structure, 568, 569
Lepidothrix pilifera, 151
 Lepidotrichidae, 150, 151
 Lepiotaceae, 278, 444
 Lepismatidae, 151
Leptinotarsa decemlineata, 648
Leptocola stanleyana, 255
 Leptoconops, 504, 507
Leptopanorpa, 475
 Leptophlebiidae, 628
 Leptopodidae, 319, 326
 Leptopodomorpha, 319, 321, 326
Leptosalda, 326
 Leptosomidae, 634
Leptospironympha, 240
Leptychoptera, 500
Leshmania: brasiliensis, 499;
donovani, 499
 leshmaniasis, 499
 Lesotho, South Africa (Stormberg Series: Triassic), 72
 Lestidae, 184
 Lestremiinae, 513
Leto, 557
 Letopalopteridae, 338
 Leucospidae, 418, 423, 424
 Levantine amber belt, 80
 Liadopsyllidae, 289
Liadotauius maior, 550, 554
 Liadytidae, 367
Liassocicada, 308
 Liassophilidae, 469, 472
 Liassophlebiidae, 187
Liassopsychodina, 497, 499
 Liayang Formation (Cretaceous), 77
Libanophlebotomus lutfallahi, 500
 Libellulidae, 180, 182
Libellulum longialata, 187
 Libelluloidea, 187
 Libytheinae, 600
 lice (see Phthiraptera)
Lichanothrips, 283
 lignocellulose, 238, 241
 ligula, 124
 Limacodidae, 579: caterpillar, 580
Limenitis archippus, 603
 Limnephilidae, 550
 Limnichidae, 381
Limnoporus, 324
Limonia, 494, 517
 Limoniidae, 498
Limothrips denticornis, 281
 Limulodinae, 374
Limulus polyphemus, 97, 100
 Lindroth, Carl, 61
 lineages, 37: crown-group, 40; stem-groups, 40
 Linnaeus, Karl, 17, 19
 Linognathidae, 273, 275
 Liomopteridae, 194, 196
 Liopteridae, 422
 Liposcelidae, 266, 267, 269, 270: as sister group to lice, 266, 269; Cretaceous, 269; habits, 266; structure, 269; wings, 269
Liposcelis, 266, 267, 269: *bostrychophila*, 266, 269; *formicaria*, 266; *myrmecophila*, 266; *prenolepidis*, 266
Liquidambar, 237
Lirimis, 582
Lirioidendron, 237, 353
Lithomantis carbonarius, 171
Lithoneura lameerei, 163, 164
Lithophotina, 260
Lithopsyche antiqua, 593
Lithoserix, 437
Lithoserix williamsi, 413
 Litophlebiidae, 164
 living fossils, 244
Lobophasma redeling huysensis, 226
Locusta migratoria, 206
 Locustavidae, 210
 Locustopseidae, 210, 215, 637
 locusts (see Orthoptera, Caelifera, Acridoidea)

INDEX

745

- Lomamyia*, 356
Lonchopteridae: Cretaceous, 532, 533, 534
Lonchopterites prisca, 534
Lonchopteromorpha asetocella, 534
 London Clay (Eocene), 49
 long-horned beetles (see Cerambycidae)
Lophioneurida, 285, 286, 287
Lophocoronidae, 560
 Lord Howe Island stick insect, 647
Loricifera, 93, 94
 lorises, as hosts, 277
Lorisicola, 277
 Los Angeles Natural History Museum, 89
 Los Rastros Formations, Argentina (Triassic), 72
 louse cospeciation, 279
Lucanidae, 379
Lucilia, 543
Luteitermes prisca, 246
 Lutz, Frank, 11
Lutzomyia, 499
Lycaenidae, 592, 593, 594, 597–99, 649:
 Caterpillars, 597; extinct, 649;
 myrmecophily, 599; relation to
 Riodinidae, 597
Lycidae, 358, 359, 383, 384, 386
Lyctocoridae, 326
Lyctocoris, 326
 Lyell, Sir Charles, 23, 24
Lygaeidae, 327
Lygaeoidea, 327, 329
Lygistorrhinidae, 38, 39, 511
Lymantria dispar, 589
Lymantriidae, 589
 Lyme disease, 272
Lymexylidae, 387, 402: Cretaceous, 387; in
 amber, 387
- M**
Maamingidae, 421
Macaranga, and ants, 302
Macherotidae, 308
Machiloides, 628
Machiloides group, 150
Macroglossinae, 615
Macroglossum, 584
Macrolepidoptera, 581
Macromiidae, 187
Macroscelidea, as hosts, 274
Macrosoma, 590–91
Macrotermes, 51, 241, 242: *falciger*, 242;
 mounds, 242
Macrotermitinae, 248
Macroveliidae, 317, 321
Macroxyela, 410
Macrozamia cycads, 284
 Madygen Formation, 71, 497
Madygenius, 410
Magicicada, 307, 308
Magnolia latahensis, 60, 237
Magnoliaceae, 621
Magnoliidae, 621
 Magothy Formation, New Jersey
 (Cretaceous), 82
Maimetsha, 417
Maindroniidae, 150
Malacopsyllidae, 480
Malacosoma, 583
Malacostraca, 99
Malaicoccus formicarii, 302
 malaria: sickle cell anemia, 502; sickle cell
 trait, 502; vectors, 502
 male genitalia (in insect), 9, 134, 228, 477
 male genitalia complexity, 135
Mallophaga, 274
Mallophorina, 522
Malmopsyllidae, 289
 mammalian radiations and insects, 638–42:
 insectivores, 638; specialized ant and
 termite feeders, 638
 mammals: fossils, 37; orders, 37
 mandible, 124; articulation, 124; structure,
 124
Mandibulata, 99, 107–18: phylogeny, 108
 mandibulate hypothesis, 93
Manidae, 638
Manlayinae (see Baissidae)
Mantidae, 257, 260: mimicry, 256
 mantises (see Mantodea)
Mantispa, 354
 mantispid lacewings (see Mantispidae)
Mantispidae, 336, 348, 353–354: Cretaceous,
 354; in amber, 354
Mantispidiptera, 355: *henryi*, 354
Mantispinae, 353
 Mantodea, 12, 147, 193, 252–260:
 adaptations, 257; bat predation of, 257;
 camouflage, 255; Cretaceous, 229, 258,
 259; diversity, 252, 253; ear function, 257;
 ear structure, 257; eye, 255; flower mimics,
 255; femoral brush, 254, 255; foreleg
 structure, 254; fossils, 257–260; in amber,
 259; mating, 256; metathoracic ear, 257;
 polymorphisms, 256; pseudovein, 259;
 raptorial fore legs, 254, 255; relationships,
 229, 259; sexual cannibalism, 256;
 structure, 254; Tertiary radiations, 260;
 vision, 255
Mantoidea, 257, 252
Mantoididae, 257, 259
Mantophasmatodea, 12, 14, 147, 188, 191,
 224–226, 628, 632: defined, 224; fossils,
 226
Marattiiales, 173
Marella, 98
Marellomorpha, 98, 99
Margarodidae, 296, 297, 298, 302:
 marsupium, 298
Margas Verdes Formation, Argentina, 92
Marquettia americana, 87
 marsupials: biogeography, 274, 485, 486, 634
Martinssonia, 99
 Martynov, Andreas V., 139, 140, 141
Masarinae, 437, 454
 mass extinctions, 37, 635: 24 my periodicity,
 635; and insects, 635; changes in sea
 levels, 635; climate change, 635;
 conditions of determining, 635; defined,
 635; drifting continents, 635; fern, 635;
 iridium layers, 635; K/T event, 635;
 microtekrites, 635; shocked quartz, 635;
 spike, 635; volcanism, 635
Mastotermes, 89, 92, 229, 230, 242–245, 250:
anglicus, 244; castes, 243; *darwiniensis*,
 230, 240, 243, 244, 628; *electrodominicinus*,
 60, 242, 244, 245; *electromexicus*, 242;
 fossils, 89, 244, 245
Mastotermitidae, 242–245, 250,
 251: extinctions, 244; fossils, 242, 244, 245
 mating behavior, 8
 mating swarms, 530
Matsucoccidae, 298, 299
 Matsuda, Ryuichi, 143
 maxilla structure, 124
 maxillary palpus, 124
Maxillopoda, 94, 99
 maximum likelihood, 32–33
 mayflies (see Ephemeroptera)
 Mayr, Ernst, 26
 Mazon Creek, Illinois (Carboniferous), 44,
 45, 67, 68, 118, 216, 335
*Mazothairo*s, 170
 mealy bugs (see Coccoidea)
Mecoptera, 12, 147, 469, 470: Mesozoic, 472;
 nuptial trophallaxis, 476; recent diversity,
 474–480; relationship to Diptera, 472
Mecopterida, 468, 469, 470–491, 547:
 Cretaceous, 470; defined, 474; early
 history, 470; in Permian, 472; larval eye,
 475; Mesozoic ectoparasites, 472–4,
 480–91; primitive features, 470
Mecopteroidea, 468
 medial plate, 130
 medial veins, 129
 Medicine Hat, Alberta (Cretaceous), 81
 Mediterranean floras, 78
Medullosoales, 173
Megachile, 435, 458, 461: *Chalicodomopsis*,
 461; cut leaves, 461; *glaesaria*, 461;
rotundata, 435
Megachilidae, 454, 455, 456, 459, 463
Megachilinae, 435
Megakhosaridae, 223
Megaleurodes megocellata, 304
Megalodontesidae, 409, 412
Megalodontidae (see Megalodontesidae)
Megalopinae, 393
Megaloptera, 147, 335, 336, 340–341; fossils,
 341
Megalopyga, 578
Megalyra troglodytes, 417
Megalyridae, 417, 630, 632
Megalyroidea, 415, 416
Meganeura monyi, 176
Meganeuridae, 175, 176
Meganeuropsis permiana (syn. *americana*),
 69, 175, 176
Meganomiinae, 455
Megarhyssa, 427
Megasecoptera, 147, 160, 169, 171, 172:
 defining features, 171–72; nymphs, 172
Megaselia (Phoridae), 14, 538
Megasoma, 379
Megaspilidae, 426
Megathon zwicki, 501
Megatypus schucherti, 175, 178
Megeniophalini, 317
Meiatermes, 245, 246: *araripina*, 246;
bertrani, 245, 246
Meinertellidae, 148
Meioneurites spectabilis, 348
 Melander, Axel L., 140
Melandryidae, 391
Melanothripidae, 284
Melieria calligrapha, 546
Melikertes stilbonotus, 466

- Melinaea*, 602
Meliphagidae, 615
Meliponinae, 58, 464; nests, 465
Meliponini, 461, 466, 619
Melissites trigona, 466
Melittia, 578
Melittidae, 454, 455, 456
 melittophiles, 377
Meloidae, 390
Melyridae, 387, 391
Membracidae: 305, in amber, 310
Membracoidea, 307
 Mená, France (Paleocene), 85
Menga tertiaria, 403, 404
Mengeidae, 402
Mengenillidae, 399, 401, 402
Menoponidae, 274, 278
 Merian, Maria Sibylla, 19, 20
 meroistic orioles, 132
Merope tuber, 476
Meropeidae, 469, 471, 476
Merothripidae, 283, 284, 285
Mesaaulacinus, 417
Mesephemeridae, 164
Mesithonidae, 336, 349
Mesochorista, 471
Mesochria neotropica, 513
Mesococcus, 299
Mesoheimerobius, 351
Mesopanorpa, 471
Mesoplectopteridae, 164
Mesopsychidae, 469, 472
Mesoraphidia luzzii, 339; *pterostigmatis*, 338
Mesoraphidiidae, 336, 338, 339
Mesorhopalosoma, 436
Mesorhyphus, 514
Mesorussus taimyrensis, 414
Mesoserphus karavavicus, 421
Mesothaumalea, 505
Mesothelae, 102
Mesotitanoptera, 215
Mesovelitiidae: Cretaceous, 317, 321, 323
 Mesozoic, insect deposits, 70–84
Mesozoicaphididae, 295
 Messel, Germany (Eocene), 87
Metallyticidae, 257, 259
Metallyticus, 259
 metamorphism, 119
Metapterygota, 147, 153, 158, 166: defined, 158; defining features, 166–167
 methanogens, 240
Metoecis, 581
Metopina goeleti, 536
Metriini, 369
 Mexican “jumping beans”, 572
 Mfwangano Island, Kenya (Miocene), 46
Miastor, 331, 334
 Michener, Charles D., 26, 584
Microcoryphia (see Archaeognatha)
Microdiplatys campodeiformis, 220
Microdontinae, 537
Microhodotermes, 244
Micromalthidae: in Cretaceous amber, 365
Micromalthus debilis, 331, 334, 363, 364, 365: amphitokous females, 364; arrhenotokous females, 364; larvae, 364
Microphor, 516, 517
Microphorinae: Cretaceous, 529
Micropterigidae, 560, 562, 564, 568, 621, 623: Cretaceous, 562; fossils, 562
Micropterix, 557, 583
Microptysmatidae, 469, 548
Microsporidae, 370
Microstigmus, 467
Microthoraciidae, 275
Micrura, 99
 Midco, Oklahoma (Permian), 69, 171, 175, 362
 Middlesex County, New Jersey (Cretaceous), 82
 migratory locust (see *Locusta migratoria*)
Miliichiidae, 58, 540
 milkweed butterflies (see Danainae)
 millipedes (see Diplopoda)
Milnesiidae, 97
Milnesium, 96, 97; *swolenskyi*, 96, 97; *tardigradum*, 96
Mimallonoidea, 583
Mimetaster, 98
Mimetica, 207
 mimicry, 537, 602–606: aculeate wasp models, 604; aposematism, 603; Batesian, 602, 603; butterflies, 603; Diptera, 604–5; Lepidoptera, 601; Müllerian, 602; Orthoptera, 605; Pompilidae, 605; Reduviidae, 605; rings, 602; Vespidae, 604
 mineralized replication, 45–49
Minyorussus luzzii, 414
 Miocene, 89–90
Miomoptera, 147, 331–333, 341, 636; as stem-group Paraneoptera, 332
Miotermes, 244
Miridae, 323, 326, 329
Mischocyttarus, 437
Mischoptera nigra, 172
Mischopteridae, 172
Misthodotidae, 163, 164
 mites (see Acari)
Mitophlebiidae, 183
Mixotricha paradoxa, 240
Mnesarchaeidae, 569
 Mo-Clay, Denmark 85, 317
Moegistorrhynchus longirostris, 521, 618, 619
 mole crickets (see Gryllotalpidae)
 molecular clock, 38, 237, 609: overestimation of age, 38
Molinodus variabilis, 554
Molossidae, 220
Molothrus as hosts, 277
 Molteno Formation, South Africa (Triassic), 72, 497
Mongoloraphidia eklices, 336
 monocondylid mandibles, 124, 148
 monographic revisions, 14
 monophly, 27
 monotremes, as hosts, 272, 482, 486
 Montceau-les-Mines, France (Carboniferous), 67
 Montsec, Spain (Cretaceous), 79
 Monura, 149
Moraceae, 423
Moravia grandis, 171
Moravocoleus, 362
Moravohymenidae, 172
Mordellidae: Cretaceous, 389, 391
 Morgan, Thomas, H., 7, 26
Mormon cricket (see *Anabrus simplex*)
Mormotomyia hirsuta, 489, 541, 542
Mormotomyidae, 489, 490, 541, 544, 640
Morphinae, 600
Morpho 600: achilles, 601
 morphology, 119, 144: comparative, 119; functional, 119
Morphopsis, 594
 Morrison Formation (Jurassic: west North Amer.), 76
 mosquitoes (see Culicidae)
 moth lacewings (see Ithonidae)
 moths (see Lepidoptera)
 Mount Flora, Grahamland, Antarctica (Jurassic), 76
 mouthpart structure, 122–124, 646
 Mt. Crosby in Queensland (Triassic), 73, 497
 multituberculates, 486
Munroessa, 578
Musca vetustissima, 378
Muscidae, 426, 490, 493, 542
 muscles, 126, 157
 Museum of Comparative Zoology, Harvard University, 86, 87, 88, 139
Mutillidae, 429, 435
Mycetobiinae, 513
 mycetocytes, 242, 288
Mycetophilidae, 508, 511
Mydidae, 521, 630
Mygalomorphae, 102
Myiodactylidae, 344
Myiodactylus, 344
Myiacrididae, 233
Mylothrites pluto, 139
Mymaridae, 424
Mymaromma (see *Palaeomymar*)
Mymarommataidae, 425: Cretaceous amber, 425
Mymarommatoidea, 415, 424
Mymarothrips, 284
Myodocopa, 99
 Myriapoda, 98, 99, 108–9: relationships, 108
Myrmecia, 429, 446, 630, 632
Myrmeciinae, 446, 447, 630
Myrmicinae, 638
Myrmecobius, 638
Myrmecocystus, 443
Myrmecolacidae, 402, 406
Myrmecolax, 406
 myrmecomorphs: 256, 323 fossilized, 440
Myrmecophagidae, 638
 myrmecophiles, 377, 440
Myrmecophilidae, 207, 209
 myrmecophytes, 440
Myrmeleontidae, 336, 342, 345–346, 516: Cretaceous, 346; larva, 346
Myrmeleontiformia 336, 342–8: defining features, 342
Myrmicinae, 302, 443
Mystacinobia zelanica, 489, 490, 544, 640
Mystacocarida, 99
Mythicomyiinae, 517, 525, 618
Myxophaga, 360, 361, 363, 370–71
Myzus persicae, 648

INDEX

747

N

- Nabidae, 322, 329
- NADH Dehydrogenase (subunit 1) (ND1) gene, 32
- Nadipteridae, 497, 499
- Nahecarida, 99
- naked mole rat (see *Heterocephalus glaber*)
- Nallachiinae, 353
- Nallachius americanus*, 353
- Namkungia*, 222
- Namurodiaphidae, 172
- Namurotypus*, 167, 177: *sippeli*, 177
- Nannochorista*, 472, 477, 479: *neotropica*, 477
- Nannochoeristidae, 475, 477, 478: Cretaceous, 478; fossil record, 478; recent diversity, 478; relationships, 478
- Nannotanyderus*, 499
- Nanoraphidia electroburmica*, 338, 339
- Nanosella fungi*, 373
- nasutes, 249
- Nasutitermes*: 239, 240, 249 *electrodominicu*s, 250
- Natal, South Africa (Beaufort Series, Permian), 69
- Natural History Museum, London, 81, 89, 91
- natural selection, 25
- Naucoridae, 71, 319, 321: fossils, 71
- Naupheta cinerea*, 231
- Necrotauliidae, 548
- Necrotaulus tener*, 549
- nectar-feeding bats, 615
- Nectarinidae, 615
- Necymylacrididae, 233
- Neduba extincta*, 648
- Neelipleona, 115
- Nel, André, 144
- Nemapalpus*, 630
- nematocerous Diptera: evolution of blood-feeding, 496; phylogeny, 496
- Nematoda, 93, 94
- Nematoida, 94
- Nematomorpha, 93, 94
- Nemestrinidae, 516, 523, 616, 618, 619
- Nemomychidae, 395, 398
- Nemoptera bipennis*, 344
- Nemopteridae, 88, 336, 342, 344, 633
- Nemopterinae, 342
- Neocorynura*, 461: *electra*, 461
- Neodermaptera, 218, 221
- Neodiptera, 507
- Neohymenoptera, 409, 410
- Neolepidoptera, 560, 568, 569
- Neolinognathidae, 274, 275
- Neopanorpa*, 475
- Neopetaliidae, 628
- Neophyllaphis*, 302
- Neopseustidae, 564
- Neoptera, 147, 157, 188–189: defining features, 188; evolutionary success, 188
- Neorthophlebiidae, 475
- neosomy, 485
- neoteny, 505
- Neottiophilum*, 540
- Nepidae, 319, 321
- Nepomorpha: airstraps, 319; families of, 319; fossil record, 319, 320; predatory behavior, 319; relationships, 319, 321

- Nepticulidae, 52, 572: fossil mines, 572; fossil, 572
- Nepticuloidea, 572
- Nesomachilis*, 628

- Neuroptera, 73, 147, 335, 341–356: classification, 341–2; defining features, 341; fossils, 149, 341–56; phylogeny, 336

- Neuropterida, 12, 147, 332, 335–357, 360: defining features, 335; phylogeny, 336

- Neuropteris* seed ferns, 52, 233

- Nevorthidae, 336, 341, 342

- Nevorthiformia, 336, 341–2: fossil, 342; primitive features, 342

- Nevorthus*, 341, 342: *fallax*, 342

- New Jersey amber (Cretaceous), 166, 234, 235, 246, 259, 285, 287, 290, 295, 300, 302, 329, 339, 381, 383, 398, 632

- New Synthesis, 26

- Newark Supergroup rift basins, 72, 75

- Newcastle Coal Measures, Australia (Permian), 69

- Nicoletiidae, 150

- Nicrophorus*, 373, 650: *americanus*, 650; *tomentosus*, 373; *vespillo*, 373

- Nitidulidae, 388, 391

- Nitidulina*, 391

- Niwratia elongata*, 485

- Nocticolidae, 231

- Noctua*, 582

- Noctuidae, 178, 589: tympanum, 583

- Noctuoidea, 560, 583, 587–90: monophly, 587

- Nogueirapis*, 463

- Nomadinae, 458

- nomenclature: 34–36; author names, 34; combinations, 35; homonyms, 34;

- International Code of Zoological, 34, 36; principle of priority, 34; synonyms, 34;

- type species, 35; typification, 35; zoological ranks, 33

- Nosybinae, 356

- notal wing processes, 128

- Noteridae, 367

- Nothofagus*, 297, 562, 625, 626: as coccoid hosts, 562; biogeography, 625; distribution, 625

- Nothomyrmecia macrops*, 446, 630, 632

- Notiophilus biguttatus*, 369

- Notiothauma reedi*, 476

- Notocampa*, 628

- Notocupoides triassicus*, 362

- Notodontidae, 590: larvae, 589; larval, 589

- Notonectidae, 319, 321

- Notonemouridae, 195

- Notoneumouridae, 628

- Notoptera, 222

- notum, 121

- numbat, 638

- nuptial feeding, 203, 529

- Nycteribiidae*, 543, 640

- Nymphaceae, 394, 621: as hosts, 622; pollination, 621

- Nymphaeales*, 613, 621

- Nymphalidae, 14, 583, 593, 594, 599–602, 614: caterpillars, 601; diversity, 600; major groups, 600

- Nymphes*, 344

- Nymphidae, 336, 342, 344

- Nymphitidae, 336, 348

- Nymphomyia succina*, 505

- Nymphomyiidae, 501, 505, 506: fossil, 505

- Nyrmina, 356

- Nyssonini, 453

O

- Oboriphlebiidae, 164

- Oboro, Czech Republic (Permian), 69

- Obrienia*, 399

- Obrieniidae, 397

- Obtectomera, 560, 577

- ocelli, 124

- Ochteridae, 319, 321

- Ochteroidea, 319

- Ocoa chilensis*, 522

- Ocoidata, 522

- Odonatoptera, 12, 43–44, 147, 167, 174, 178–187,

- 628: arista, 167; classification, 183–7;

- copulation wheel, 179; copulation, 131,

- 134, 155; copulation, 179, 182; defined,

- 179; early history, 182–87; fossils, 162;

- Jurassic, 184, 187; male genitalia, 182;

- naiads, 181; nymphal labium, 181;

- phylogeny, 180; prehensile mask, 181;

- secondary genitalia, 182; subordinal

- relationships, 180; venation, 181

- Odonatoptera, 147, 157, 173–187; defined,

- 174; gigantism, 178; phylogeny, 180, 181

- Odontotermes obesus*, 132

- Oecophoridae, 576, 577

- Oecophylla* ants, 46, 47: behavior, 46, 47;

- fossils, 46, 47

- Oedemeridae, 390, 391

- Oedischiiidae, 207

- Oedoparena glauca*, 540

- Oeningen, Germany (Miocene), 54, 89, 382

- Oenosandridae, 587

- Oestridae, 492, 493, 543

- Oestroidea, 62, 543, 640: fossilized in hosts , 62

- Oiketicus*, 576

- Okanongan Highlands, Bristish Columbia (Eocene), 86

- Okham's Razon, 32

- Olbiogaster perezi*, 514

- Olbiogastrinae, 513

- Oligochlora eickworti*, 461, 463

- oligophagy, 283

- Oligophrynidiae, 516

- Omalisidae, 384

- Omalium*, 374

- Omamina*, 365

- Omethidae, 384

- Omma*, 363

- Ommatidae, 363, 364

- ommatidium, 124

- Ommexechidae, 210

- onchocerciasis, 502

- Oncopeltus*, 327

- Oncothrips*, 283, 465, 467

- Oniscigastridae, 628

- Onthophagus*, 378

- ontogenetic criterion (of character polarity), 28

- Onychomachilis fischeri*, 150, 152

- Onychophora, 93, 94, 95, 98: fossils, 94;

- phylogenetic position, 94

- Onychotrechus*, 317

oothecae, 83, 230, 231, 242; fossil, 83, 231, 235; in Dictyoptera, 230, 231, 252
 operculum, 193, 212, 213, 226, 262
Opelia, 532, 533, 630; *nigra*, 534
Opetiala shatalkani, 533
 Opetiidae, 532, 533
 Opiliones, 99, 103
 opistognath, 121
 Opisthothelae, 102
 Opostegidae, 572
Opsiphanes, 601
 Orapa, Botswana (Cretaceous), 83
 orb webs, 102
 orchid bees (see Euglossini)
 orchids, 615; and orchid bees, 616; and pseudocopulation, 615; pollination, 615
 orders (insect), phylogeny, 146; classification, 147
Oreadomyia, 505
 Oreopteridae, 187
 Örsten preservation 45, 96
Ornithoptera, 595
 Orstenocarida, 99
 Ortheziidae, 296, 297, 298, 299
Orthophlebia, 471
 Orthophlebiidae, 470, 471, 475
 Orthoptera, 12, 147, 189, 191, 201, 202–211, 622, 628, 642; Aposematic coloration, 202; calling behaviour, 203–4; crypsis, 206–7; cryptopleuron, 203; defining features, 203; fossils, 208–11; plague locusts, 206; tympana, 203, 204
 Orthopterida, 147, 188, 202; defining features, 193; phylogeny, 191, 202
 Orussidae, 409, 411, 413, 630
 Orussoidea, 412
Orycteropus afer, 638
Osmia, 455
 osmylid lacewings, 348
 Osmylidae, 336, 348, 349
 Osmylitidae, 336
Osmylops, 344
 Osmylopsychopsidae, 336, 344
Osmylus fulvicephalus, 349
 Ostracoda, 99
 Otitidae, 30, 541; fossil, 546
 outgroup comparison (of character polarity), 158
 ovary structure, 132
 ovarioles, 132
 ovipositor, 127, 132–134, 137; evolution, 133
 oviscapt, 133
 ovoviparity, 230
 Owen, Richard, 26
 owlflies (see Ascalaphidae)
 Oxymonadidae, 240
Oxymonas, 240

P
 Pachyneuridae, 508
Pachysandra, 237
Pachysystrophus rohweri, 619
 Pachytrochidae, 270
 pack rat middens, 62
 paederin, 374
 Paederinae, 374
Paederus, 374
 paedogenesis, 331
 paedomorphosis, 220, 252

Palaeoboreus: baissicus, 479; *zherikhini*, 479, 480, 488
 Palaeocynipidae, 422
 Palaeodictyoptera, 147, 170; defining features, 170
 Palaeodictyopterida, 147, 157, 168–173, 623; classification, 169; defining features, 170; nymphs, 169; Paleozoic herbivory, 173
Palaeogryyon musesebecki, 426
Palaeokoenenia mordax, 104
Palaeoleon ferrogeneticus, 346
 Palaeoleontidae, 346
Palaeomyimar, 424
Palaeomyrmecolax, 406
Palaeopsylla: baltica, 485; *dissimilis*, 485; in Baltic amber, 485; *klebsiana*, 485
 Palaeoptera, 157–187
Palaeotis, 634
 Palaeophatidae, 572
 Paleocene: insect deposits, 85, 637; insects, 637
Paleoculicis minutus, 506
 paleoentomology, 144
Paleomelitta nigripennis, 461
 Paleontinidae, 309
 Paleontological Institute, Moscow, 71
 paleontology, 36–41; 144
 paleopterous, 157
Paleotermopsis, 247
 Paleozoic: insect deposits, 3, 38, 45, 53, 63, 65–6, 69; significance of, 65
 Paleozoic roachoids, 36–7, 231
Paleothygramma tenuicornis, 217
 Paleothygrammatidae, 217
 Pallopteridae, 540
Palophagus, 393
 Palpigrada, 99
 Pamphagidae, 210
 Pamphagodidae, 210
 Pamphiliidae, 409, 411, 412
 Pamphilioidea, 412
 Panamanian Land Bridge, 92
 Panarthropoda, 93, 94; phylogeny, 93
 Pancrustacea, 107, 108; defined, 107
 Panderaptera, 218
Panesthia, 237
 Panesthiinae, 241
 pangolins, 638
 Panhexapoda, 99, 111
 Pannota, 162
 panoistic ovarioles, 132
Panorpa, 475, 476
 Panorpida, 147; ancestral, 468; defined, 468; monophly, 468; mouthparts, 495; phylogeny, 469; relationship to Hymenoptera, 407, 468
 Panorpidae: fossil, 469, 475, 476; male terminalia, 475; nuptial feeding/trophallaxis, 475
Panorpodes, 475
 Panorpodidae, 469, 475
 panorpoid orders, 468
 Pantophthalmidae, 516
 Pantopoda (Chelicerata), 45
 Paoliidae, 147, 160, 188, 189
Papaipema, 582

paper wasps (see Polistinae)
Papilio dardanus, 603, 604; *glaucus*, 605; *polyxenes*, 596
 Papilionidae, 556, 595; caterpillar, 596; major groups, 597; osmeterium, 595, 596
 Papilioninae, 597
 Papilionoidea, 555, 556, 591, 594, 618; ant-tended caterpillars, 599, 600; chrysalis, 596; color genetics, 605; color pattern, 605; cremaster, 594; families, 594; faunistic treatments, 595; field guides, 594; fossil, 592, 594, 595; in amber, 594, 595; mimicry, 604; predatory larvae, 598
 Parabasalea, 240
 Paracolletinae, 455
 Paracolletini, 630
Paradixia, 504
 paraglossa, 124
Parahemiphlebia mickoleiti, 187
Parahygrobia natans, 368
 Parahygrobiidae, 367
 parallel cladogenesis (see cospeciation)
 Paralogidae, 176
 Paramegasecoptera, 172
 parameres, 134, 135
 Paraneoptera, 12, 147, 160, 188, 261–330; feeding habits, 261, 263; monophly, 261; mouthparts, 261, 264; phylogeny, 263; relationships among orders, 261; wing coupling mechanisms, 261, 265; wing venation, 268
 paranotal lobes, 158
 Parapamphiliidae, 412
 Paraphrynoveliidae, 317, 321
 paraphyly, 27
 Paraplectoptera, 193, 196
Paraponera clavata, 430, 446
 Paraponerini, 446
 paraprocts, 131
 Parasialidae, 336, 341
 parasitic wasps (see Apocrita)
 parasitism, 413
 parasitoids, 79, 204, 301, 323, 371, 543; in Coleoptera, 374, 377, 380, 389, 390; in Diptera, 516–7, 521, 536, 537, 540, 543; in Hymenoptera, 408, 413–29; in Lepidoptera, 558, 572, 577, 580; in Mantispidae, 354; in Strepsiptera, 399–406
Parasphearia boleiriana, 241
Parastizopus armaticeps, 389
Parastylotermes, 247
Parategeticula, 572, 573
 parental care, 325, 330; in Arachnida, 101; in Blattaria, 237, 252; in Coleoptera, 373, 378, 394; in Dermaptera, 217; in Diptera, 507; in Hemiptera, 312, 319, 323, 325; in Mantodea, 252
 Paris Basin amber (Cretaceous), 79
 Parnassiinae, 597
 Paroryssidae, 409, 414
Parotermes, 247
 parsimony: 32 Camin-Sokal, 32; Dollo, 32; general, 32; Wagner, 32
 parthenogenesis, 266, 290, 297
 Paskapoo Formation, Alberta (Paleocene), 85
 Passalidae, 378
Passiflora, 277
 Passifloraceae: as hosts, 604

INDEX

749

passion vine butterflies (see *Heliconius*)
Pauliniidae, 210
Paurometabola (see *Polyneoptera*)
Pauropoda, 99, 108, 110
Paussinae, 368, 369
Paussini, 369
Pearman's organ, 265, 266
Pecaroecidae, 275
Pedicia, 498
Pedicinidae, 275
Pediculidae, 275
Pediculus humanus, 62, 270–272, 274, 275, 280; on mummies, 62
pedipalps, 135
Pelecinidae, 421
Pelecorhynchidae, 522
Peloridiidae, 312, 317, 630; *See also Coleorrhyncha*
Peloridora, 312
Pemphiginae, 54, 293, 295, 465, 467
 penguins, as hosts, 277
 penis, 132, 134, 135
Pentastomida, 45, 93, 99
Pentatomidae, 268, 327, 329, 330
Pentatomoidea, 327, 329
Pentatomorpha, 321, 326–330, 622, 637: Cretaceous, 330
Pentoxyton, 607
Pepsis, 429
Peradeniidae, 421
Pergidae, 409, 410, 411, 630
period gene, 8
Peripatidae, 94
Peripatopsidae, 94
Peripatus, 94
Periplaneta americana, 231
Perissomatidae, 508, 630
Perlidae, 195
Permaeschidae, 183
Permagrionidae, 183
Permaleurodes, 299
Permelcanidae, 208, 210
Permembia, 199
Permeplallidae, 183
 Permian, 44, 53, 68–70; fossil insect deposits, 65, 77, 79; significance of, 68, 70, 178; paleomap, 70
Permila, 497
Perilampidae, 420
Permithonidae, 336, 341
Permoberotha villosa, 333
Permoberothidae, 332, 336, 337
Permozentropidae, 469, 472
Permochoristidae, 469, 471, 472, 479
Permocupedidae, 362
Permocupoides, 362
Permoestidae, 183
Permopanorpidae, 469, 472
Permoplectoptera, 163, 164
Permopsocida, 269, 285
Permopsocidae, 269
Permorphidiidae, 208, 338
Permotanyderidae, 469, 472, 497
Permothemistida, 169, 170
Permotipula, 497
Permotipulidae, 469, 472, 497
 Permo-Triassic Extinctions (End Permian Event), 69, 71
Petaluridae, 182, 628

Petaluroidea, 187
 Petrified Forest, Arizona (Triassic), 72
Petrobius, 148
Petromantis, 471
Petropsychops superba, 344
Phaetempis lebanensis, 530
Phalangiotarbida, 99
phallomeres, 228
Phallothrips, 283
Pharnacia kirbyi, 211
Phasmatodea, 12, 147, 189–191, 211–215, 622; aptery, 212; area apicalis, 213; classification, 213; defining features, 211; eggs, 214; fossils, 215; in amber, 215; mimicry, 211; operculum, 212, 213; parthenogenesis, 213; vomer, 211
Phasmida (see *Phasmatodea*)
Phasmomimidae, 208
Pheidole, 14, 443
Phenacoleachia, 297
Phenacoleachiidae, 296, 297, 298
 phenetics, 26
Phengodidae, 383, 385
Philopotamidae, 550
Philopteridae, 275
Phlaeothripidae, 280, 283, 284, 285, 286, 287
Phlaeothripine, 467
Phlebotominae, 490, 500: as disease vectors, 499; Cretaceous, 499; earliest, 499
Phlebotomites, 499
Phloeidae, 327
Pholidota, 482, 638
Phoridae, 538–540: ecological diversity, 534; fossil, 539; in Cretaceous amber, 539; *Megaelia*, 534, 538; parasitoid, 533, 536; *Prioriphora*, 536, 539; species diversity, 533; stem-group, 536
Phormia, 491
 phosphatization, 46, 47, 96
Phosphatocopida, 99
 phosphoenolpyruvate carboxykinase (PepCk) gene, 32
Photinus, 385
Photuris, 385
 phragma, 120
Phrixothrix, 385
Phryganeidae, 555
Phryganopsychidae, 555
Phthiraptera, 12, 62, 147, 270–280, 490, 640: ages, 276–77; as disease vectors, 272; classification, 274–5; cowbird hosts, 277; cuckoo hosts, 277; dispersal, 272, 280; effects of host extinctions on, 277; egg laying in, 272; eggs, 272; fossils, 275–76; founder effects, 280; high rates of nucleotide substitutions, 280; host age estimation, 276; host colonization in, 276, 277; host use patterns, 277; hosts, 272–74; on mummies, 275; Pleistocene remains, 275; putative Cretaceous, 276
 phyletic gradualism, 11
Phylloblatta gallica, 227
Phylloblattidae, 227, 229, 233
Phyllobrotica, 624
Phyllocarida, 99
Phyllopoda, 99
Phyllostomidae, 615
Phylloxeridae, 293, 295

phylogenetic analyses, 31: Bayesian methods, 38; general, 31; taxon completeness, 31
 phylogeny, 15, 27, 31, 40: defined, 31, 32; general, 31, 32; significance of fossils, 40
Phymatinae, 323
Physalaemus pustulosus, 206
Phyocephala, 541
Phytalmia, 541
Phytophaga, 361, 392–399, 622, 637: Monophly, 393
phytophagan Beetles, 393
phytophagous insects, 622–25, 642: and allopatric divergence, 625; and angiosperms, 623; Paleozoic, 173; and plant toxins, 624
phytotelmata, 317
Piagetella, 274
Pieridae, 592, 597: and Brassicaceae, 597
Pieris, 596: *brassicae*, 597; *rapae*, 597
Piesmatidae, 329, 330
Pinaceae, 55, 88, 410
Pincombea, 295
Pincombeidae, 295
Pinites succinifer, 297
 pinnipeds, 482: as hosts, 272
Pinus, 410
Piophilidae, 540
Pipunculidae, 537, 539
Pisciformia, 162, 163
Piscindusia, 52, 551
Plagiocephalus latifrons, 30, 541
 plague, 480, 482: and flea proventriculus, 482; bubonic, 482; historical significance, 480; pneumonic, 482; *Yersinia*, 482
planidia, 517, 543
Planivora, 512
 planktonic foraminiferans, 63, 642
Planoneoptera, 192
 plant feeding (see *phytophagous insects*)
 plant galls: fossils, 50, 52, 53
 plant lice (see *Psylloidea*)
Plasmodium falciparum, 502
Plastoceridae, 384
Plataspididae, 327
Plattenkalke deposits, 74, 82
Platycoelostoma, 297
Platygastridae, 425
Platygastroidea, 415, 425–6: Cretaceous, 426; defining features, 425
Platypedia primigenia, 310
Platypezidae, 532: Cretaceous, 537
 platypodid, 58
Platypodinae, 387, 396, 397, 467: fossils, 387; in amber, 397
Platypyllus castor, 373, 490
Platystoechotes, 349
Platystomatidae, 541
 pleasing lacewings (see *Dilaridae*)
Plebeia, 457, 465
Plecoptera, 12, 147, 188, 191, 194–196, 554, 628: classification, 194; defining features, 194; fossils, 195–96; nymphs, 195; phylogeny, 194
Plecopterida, 192–201: defining features, 192
Plectrotarsidae, 555
Pleidae, 319, 321
 Pleistocene dispersal, 642–43

Pleistocene insects, 61, 90
Pleotomus, 385
Plesiometra argyra, 427
 plesiomorphy, 27
 pleural coxal process, 125
 pleural wing process, 128
 pleuron, 125
 plica orales, 112
 Pliny the Elder, 16
 Pliocene, 90
 Plokiophilidae, 323, 328
 Plumariidae, 430, 632
Plusiotes resplendens, 378
Plutella xylostella, 648
 Pneumoroidea, 210
 pocket gopher-louse cospeciation, 278
 pocket gophers, as hosts, 277, 279
Podocarpus, 302, 625: as hosts, 302
 Podonominae, 627
 Poduridae, 114
Pogonomyrmex, 443
Polanskiella, 450, 632
Polistes, 429
 Polistinae, 437, 438
 pollination, 5, 540, 613–622; age of
 specialized pollination, 616; ages of insect
 pollinators, 616; amount by insects, 615;
 and fossilized pollen meals, 616; and
 hovering, 617; and long proboscis, 616;
 and origins of bees, 619; bee, 619; by bats,
 615; by birds, 615; chronology of
 pollination features, 616; defined, 613;
 droplets, 620; earliest pollinators, 621;
 ecological advantages, 615; examples of
 specialized, 615, 616; faunas of pollinators,
 621; fossil evidence, 616–20; fossilized
 pollinators, 619; in basal angiosperms,
 620; in cycads, 620; in gnetaleans, 620; in
 grasses, 621; in primitive living
 angiosperms, 620–22; insect, 613, 614;
 pollination features, 616; pollinators, 618;
 significance of Brachycera, 618;
 specialized floral structures, 616;
 syndromes, 615; wind, 615
 Polyctenidae, 272, 326, 329, 490, 640
 Polycyellidae, 332
 Polydnaviruses, 427
Polyergus, 442, 443
 Polyneoptera, 147, 188–260: defining
 features, 189; orders of, 191; phylogeny,
 191; wing venation, 190
 Polyorthoptera, 193
 Polyphaga, 371–99, 630: diversity, 371;
 infraorders, 371; monophly, 371
 Polyphagidae, 231, 235, 237
 polyphagy, 283
 polyphyly, 27
Polystoechotes, 349
 Polystoechotidae, 336, 342, 348, 349, 633
 polytene chromosomes, 8
 polytrophic overioles, 132
 Pompilidae, 429, 433, 434, 435, 604, 630:
 Cretaceous, 435
Pompilopterus, 435
 pond skaters (see Gerromorpha)
 Ponerinae, 446
 poneroid grade of ants, 449, 450: Cretaceous,
 435
 Pongidae, as hosts, 275

Ponopterix axelrodi, 235
Pontomyia, 548
Populus (poplars), 52
Poroterpes, 247, 628
 Porphyry: Tree of, 16
Porrerus dominicanus, 346
 Porricondylinae, 513
 postmentum, 124
 postnotum, 121
 postoccipital suture, 122
Postopsyllidium, 289
 Potrerillos Formation, Argentina (Triassic), 72
 potter wasps (see Eumeninae)
 Praeaulacidae, 419
Praeichneumon, 429
 Praeichneumonidae, 429
Praemordella, 389
 preantennal segment, 123
 Praesiricidae, 409
 Precambrian Period, 93
 prehalter, 499
 prementum, 124
Premnoxylon, 53, 54
 pretarsus, 125, 127, 475
Priapula, 93
 Priapulida, 94
 primates, 482
 principle of connections (St. Hilaire), 28
 Prionoglariidae, 269, 270
Prionomyrmex, 450, 632
Prioriphora, 40, 536, 539: *casei*, 539; in
 Cretaceous amber, 539
 Priscaenigmatidae, 336, 338
 Priscaenigmatomorpha, 338
Pristapenesia, 431
 Probnidae, 196
 Proboscidea, 482
 Procyclariiformes, as hosts, 278
 Proceropidae, 308
Prochaerododis enigmaticus, 260
Prochoristella, 472
 Procramptonyidae, 508, 509
 proctodeal trophallaxis, 235
 Proctorenyidae, 421
 Proctotrupidae, 421
 Proctotropoidea, 415, 421: Jurassic, 421
 Proctotropomorpha, 415, 418, 419–427:
 defining features, 421
 Prodoxidae: diets, 572; evolution of
 monoelectric, 573; monophagy, 573;
 pollination of yuccas, 572–73; specialized
 maxillae, 573
Prodoxus, 573
Podryas persephone, 87, 590, 593
 prognath, 121
 Progonocimicidae, 313, 314, 637
 Progonophlebiidae, 187
Promantispa similis, 355
Promegalomus, 351
 pronymph, 333, 334
Propalosoma, 436
 Proparagryllacrididae, 215
Proplebeia dominicana, 60, 461, 464
 propodeum, 414
Proscolia, 437
 Proscopiidae, 203, 205, 211, 224, 628
Proscorpius osborni, 101, 102
Prosoeca, 523
Prosphyracephala, 543

Prostomidae, 390: Cretaceous, 390
Prosyntexis gouleti, 413
 Protanisoptera: defined, 183
 protein: fossilization, 42
Proteinus, 374
Protelencholax, 406
 Protelytroptera, 192, 220, 221
Protembia permiana, 198
 Protembriaria, 198
 Protembidiidae, 198
Protensis antennata, 526, 527
Proteresa, 167: *americanum*, 165;
 permianum, 164
 Protoreismatidae, 163, 164
 Protobiellinae, 356
Probombus, 466: *indecisus*, 462
 Protocoeloptera, 361, 362
 Protodermaptera, 218, 221
 Protodiplatyidae, 218, 220
 Protodonata, 147, 174–178
 Protohymenidae, 172
 Protohymenoptera, 169, 171, 172
Protolepis cuprealata, 575
Protoligoneura limai, 165
Protolithurgus, 461: *ditomeus*, 461
Protomelecta brevipennis, 458
 Protomeropeidae, 548
 Protomyrmeleontidae, 183
Protomonomastus, 616
 Protoneuroptera, 332
 Protoperlaria, 194, 196
 Protoplecidae, 508
Protopone, 446
 Protopsylidiidae, 289, 290, 637
 Protoptera, 188
Protreogeton, 526
 Protorhynchidae, 514
 Protorthoptera, 147, 173, 188: Paleozoic
 herbivory, 173
Protoscelis, 394
Protosceloides, 394
Protosolpuga carbonaria, 103
 Protozygoptera (see Archizygoptera)
 Protura, 12, 111–113, 154
 proventriculus, 228, 237, 474, 479: in
 Dictyoptera, 228; in fleas, 474, 479; plicae,
 228; structure, 228
Psacodromeus guttata, 367
Psaronius tree ferns, 53
 Pselaphidae, 627
 Pselaphinae, 374
 Pselaphognatha, 99
 Psephenidae, 359, 381
 Pseudococcidae, 296, 298–303: and nomadic
 ants, 301–2
Pseudococcites eocaenicus, 406
Pseudohymen, 171
Pseudolarix, 88
Pseudomacromia sensibilis, 187
Pseudomegamerus, 394
 Pseudomorphinae, 368
Pseudomyrmecinae, 446
Pseudoperla, 215
Pseudophengodes, 385
 Pseudophyllinae, 207
 Pseudopolycentropodidae, 469, 637: arista,
 472; halteres, 472; in Burmese amber, 472;
 long proboscis, 472; possible diet, 472;
 vestigial labial palps, 472

INDEX

751

- Pseudopolycentropus*, 471, 472, 494, 617
Pseudopsis, 14
Pseudoregma bambucicola, 293
Pseudoscorpionida, 99, 103
Pseudosphinx, 603
Pseudostigmatidae, 184
Pseudotettigonia amoena, 206
Psocidae, 270
Psocidiidae, 269
Psocodea, 147, 261–3, 276, 332
Psocoptera, 12, 147, 199, 261–270, 332:
 antennal rupturing mechanism, 266;
 aptery, 266; brachyptery, 266–67;
 cibarium, 266; classification, 266; feeding
 habits, 266; in Cretaceous amber, 269; in
 Permian, 269; mating behavior, 266;
 nymphal stages, 332; oldest fossils, 270;
 Paleozoic paraphyly, 269; paraphyly of,
 266; parthenogenesis, 266; Pearman's
 organ, 265; sister group to lice, 266;
 sitophore sclerites, 263; structure, 265,
 266; viviparity, 266; water vapor
 absorption, 263
Psychidae, 572: fossil cases, 575; larval cases,
 576
Psychodidae, 489, 492, 498, 500: Cretaceous,
 500
Psychodomorpha, 496, 498–501
Psychomorpha, 582
Psychopsidae, 85, 336, 342, 633
Psychotanyderus, 500
Psyllipsocidae, 270
Psyloidea, 268, 291
pteralia, 128, 129
Pteromalidae, 420
Pteropinae, 615
Pteroptyx, 385
 pterosaurs, as hosts, 276
pterostigma, 131
pterothorax, 128
Pterygota, 147, 155, 157: defining features,
 155
Pthiridae, 275
Pthirus pubis, 274
Ptiliidae, 373
Ptilocerus ochraceus, 326
Ptilodactylidae, 381
Ptinidae: fossils, 62
Ptininae, 386
Ptychopteridae, 492, 498, 499
Pulex, 485, 486: in amber, 485; *irritans*, 481;
larimerius, 485, 486; Miocene, 486
Pulicidae, 480, 481
Puliciphora, 540
Pulicoidea, 485
 punctuated equilibrium, 11, 26
Pupipara, 490
Purbeck Group, England (Cretaceous), 78
Purbeck, 533
Putoidae, 296, 297, 298, 299
Pycnogonida, 99, 100
Pygidicranidae, 221
Pygidicranoidea, 218, 221
pygidium, 131
 pygmy grasshoppers (see *Tetridoidea*)
Pyralidae, 301, 569, 580–81, 608
Pyrearinus termilluminans, 384
Pyrgomorphoidea, 205, 210
Pyrotidae, 540
 pyritization, 49
Pyrochroidae, 390
Pyrocoelia, 385
Pyrophoridae, 384
Pyrophorus, 384
Pyrrhocoroidea, 327, 329
- Q**
 quartz, 45, 635
Quaternary Period, 642
Queensland, Australia, 92, 244
Quercus (oaks), 52, 53, 353
Quercy, France (Eocene), 46
Quinarians, 21
 quinoline, 213
- R**
 races, 9
 radial veins, 129, 183
 radiations (evolutionary), 37, 53–54
Radoboj, Croatia (Miocene), 89
Rangomaramidae, 511
Ranzovius, 323
Raphidia funerata, 339
Raphidiidae, 336, 338, 339
Raphidiomimidae, 229, 234, 637
Raphidiomimula, 234
Raphidiomorpha, 338
Raphidioptera, 75, 147, 335, 336, 337–340,
 633: classification, 337–8; Cretaceous, 339;
 defined, 337; fossils, 338; Jurassic, 338
Rapisma, 349
Rapismatidae, 336, 349
Raptophasma, 226
 raptorial forelegs, 254, 255, 367: convergence
 in, 356
Rasnickytipidae, 422
Rasnitsyn, Alexander, 144
Ratemiidae, 275
Ray, John, 17, 19
Redbank Formation, Queensland
 (Paleocene), 92
Redivivini, 456
Reduviidae, 268, 323, 327, 329
Regiata scutra, 258
Remane, A., 29
Remipedia, 99
Renova Formation, Montana (Oligocene),
 88
 repugnatorial glands, 109
resilin, 470, 482, 563
resinate, 55, 57
Reticulitermes flavipes, 240
Rhachibermissa splendida, 355, 634
Rhachiberothidae, 336, 348, 355–356, 634:
 Cretaceous, 355, 634
Rhaetomyiidae, 505
Rhagio, 522
Rhagionidae, 489, 490, 492, 516, 522, 523,
 524: blood feeding, 522; Cretaceous, 524
Rhagoletis pomonella, 624
Rhagophthalmidae, 384
Rhagovelia, 317, 323
Rhamphomyia, 529
Rhaphidophoridae, 208, 628
Rhetinothripidae, 285
Rheumatobates, 317
Rhinophoridae, 543
- Rhinotermitidae**, 242, 247, 249, 250, 251:
 diversity, 247; fossils, 247; oldest, 249
Rhipidoblattina katavica, 234
Rhipiphoridae, 390, 391, 402: in amber, 390,
 403
Rhipipterygidae, 211
Rhodites vetus, 422
Rhodnius, 323
Rhombocoleidae, 362
Rhopalocera, 590, 591: Eocene, 590; fossil,
 590; in amber, 590; Paleocene, 590;
 recency of, 590
Rhopalopsyllidae, 485, 486
Rhopalopsyllus, 485
Rhopalosomatidae, 433, 436
Rhyacophilidae, 550
Rhynchosciara, 491
Rhyncophthirina, 274
Rhynie Chert, 49, 66, 116, 152
Rhynie, Scotland, 49, 66, 116, 153
Rhyniella praecursor (Collembola), 66, 116,
 152, 154
Rhyniognatha hirsti, 40, 66, 152–154, 160,
 646: phylogenetic position, 154
Rhyparochromidae, 327
Rhysodinae, 368, 369, 370: in amber,
 368
Riciniidae, 274, 306
Ricinulei, 99, 106
Rickettsia prowazekii, *typhi*, 272
Riodinidae, 594, 597, 598, 623: caterpillars,
 599–600; myrmecophily, 599
Riversleigh, Queensland (Miocene), 46
 RNA Polymerase II (Pol II) gene: calcium
 phosphate, 46
 roaches (see *Blattaria*)
 roachoids, 227
Robinjohnia, 497
Robinjohniidae, 469, 472, 497
Robopus, 385
 rock crawlers (see *Mantophasmatodea*)
Rodentia, 465
Rohdendorf, Boris B., 37, 74, 143
Romaleidae, 210
Roman Empire, 16
Ronheim, Germany (Oligocene), 46
Rophalis, 342
Roproniidae, 421
Rosa, 422
Rosaceae, 412
Rothschild, Miriam, 480
Rothschildia, 584
Rott, Germany (Oligocene), 463
Roussiana, 466
Rubielos de Mora, Spain (Miocene), 89,
 465
Ruby River Basin (Oligocene), 88
- S**
 Saint-Hilaire, Etienne Geoffroy, 23, 28
 Sakhalin Island amber (Paleocene), 85
Saldidae, 319
Saldonia, 319
Salicaceae, 413
 salivarium, 124
 salivary canals, 288
Salpingidae, 391
Samia cynthia, 584, 585
Sanctacarida, 99

Santana Formation (Crato member), Brazil, 47, 48, 83, 152, 165, 166, 185, 187, 235, 380, 381, 382, 398, 449, 523, 562, 611, 617, 618; assorted fossils, 46; diversity, 53; general, 82–83; preservation, 46
Santanantis axelrodi, 258, 259
Sapyga, 435
Sapyginae, 435
Sarcophagidae, 493, 543
Saturniidae, 584, 585
Saurodetes vrsanskyi, 275, 276
Saurophthirus longipes, 472, 473
 Savage Canyon Formation, California (Miocene), 90
 sawflies (see Symphyta)
Saxonagrionidae, 184
Scala Naturae, 16
 scale insects, 297
Scalidorhyncha, 94
Scaphidiinae, 374
Scaptomyza, 644
Scarabaeidae, 378, 490: dung rolling, 378; significance among Egyptians, 378
Scarabaeiformia, 377: monophyly, 377
Scarabaeinae, 378
Scarabaeoidea, 358, 359, 361, 377–380: Cretaceous, 380; fossil, 380; larva, 378; male horns, 379; monophyly, 377; relationships in, 377, 424
 scarabs (see Scarabaeoidea)
Scathophagidae, 492, 542
Scatopsidae, 500
Scatopoidea, 501, 500
Scelionidae, 425
Sceliphron, 48
Scenopinidae, 521, 618
 Scheuchzer, Johann J., 22, 23
Schistocerca gregaria, 206
Schistonota, 162
 schistosomiasis, 540
Schizocoleidae, 361, 362, 365
Schizodactylidae, 208
Schizomida, 99, 106
Schizophora, 539–547, 637: acalyptates, 539; compression fossil, 545, 546; defined, 539; fossil, 545; in amber, 547; origin of, 547; ptilinum, 539; sexual dimorphisms, 542; Tertiary radiation of, 546
Schizophoridae, 363
Schizopteridae, 317, 318
Schizoramia, 93, 107
 schizoramian hypothesis, 93
Schizura, 589
 Schlee, Dieter, 80
Schlettereriinae, 416
Sciadocera rufomaculata, 537
Sciadoceridae, 533, 536, 630
Sciariidae, 511
Sciaroidea, 508, 511: Cretaceous, 509
Sciomyzidae, 540
Scirtidae, 363, 381
Scirtoidea, 380
Sciuridae, as hosts, 274
 sclerites, 120
Sclerogibbidae, 432
Scolebythidae, 431, 630, 632: Cretaceous, 432
Scoliidae, 437
 Scolytinae (bark beetles), 54, 387, 396, 397: fossils, 387, 396; in amber, 397

Scolytus, 398
 scorpionflies, 468
 scorpions, 99, 101–102, 319, 474, 642: Cretaceous, 102; Silurian, 102
Scaptiidae, 391
 Scudder, Samuel H., 85, 137, 139
Scutellaridae, 327
 scutellum, 125
 scutum, 125
Scydmaenidae, 372: Cretaceous, 372
Scyphindusia, 551
Scytohymenidae, 172
 sea birds, as hosts, 278
 sea scorpions, 100
 seabirds, 279
 secondary segmentation: in insects, 120
 Seitz, Adalbert, 555
Sematuridae, 585
Semenoviidae, 218
 semiaquatic insects, 499
 semisocial, 464
 semi-species, 8
Sepulcidae, 409, 412
 sericulture, 584
Serphtidae, 424–425, 637
Serphtoidea, 415, 424–425
Serritermitidae, 247, 250
Sesiidae, 579
Sesioidae, 579
Setisura, 162
 Seward Peninsula, Alaska (Pliocene), 90
 sexual cannibalism, 103
 sexual dimorphism, 379, 542, 644,
 sexual selection, 136, 379
 Shaanxi Province, 71
 Shangu Formation, 71
 Shanwang Formation, China (Miocene), 89
Sheimiodea, 198
Sialidae, 340
Sialis, 340
 Siberia, 80, 398
 Siberian amber (Cretaceous), 270, 293, 295, 299, 391, 572
 siderite, 44
Sidneyiida, 99
Sierolomorpha, 434
Sierolomorphidae, 434
Sikhotealinidae, 363
 silkworms (see *Bombyx*)
 silkworm moths (see *Bombycoidea*)
 silky lacewings (see *Psychopsidae*)
Silphidae, 358, 372, 373: parental care, 373
Silphopsyllus, 373
 silverfish (see *Zygentoma*)
 Simpson, George Gaylord, 26
Simuliidae, 84, 489, 490, 493, 502, 504, 505: Cretaceous amber, 506; labral fans, 505; larva, 505
Simulimima grandis, 505
Simulium damnosum complex, 11, 502
Sinicorussus luhzhongensis, 414
Siphlonuridae, 628
Siphonaptera, 12, 147, 468, 469, 480–490, 640: and plague, 480; classification, 485; cocoon, 484; Cretaceous stem-group, 487, 488; Cretaceous, 485; ctenidia, 486; distributions, 480, 484; eggs, 485, 489; families, 480, 485; features, 482, 483; feeding, 482; fossil record, 486–8; hosts,

482; in amber, 485, 486; larva, 482, 484; life histories, 482; monophyly, 482; mouthparts, 483; neosomy, 485; pygidium, 482; sensillum, 482, 486; species diversity, 480; stem-group, 486; subcutaneous, 485
 sirenians, as hosts, 272
Siricidae, 409, 411, 412
Siricinae, 412
Siricoidea, 412
 sister groups, 40, 237: general, 40; immediate, 40; living, 40, 237
Sisyridae, 336, 341, 348, 352
Skaracarda, 99
 skippers, 592
 sleeping sickness, 545
Sminthuridae, 115, 628
 Smith, William, 23
 Smith, John B., 140
 Smithsonian Institution, 86, 91
 snakeflies (see Raphidoptera)
 Sneath, Peter, 26
 Snodgrass, Robert E., 140, 142
 social behavior, 397
 social insect ectosymbionts, 533
 sociality, in Aphidoidea, and galls, 293
Sojanoraphidiidae, 338
 Sokal, Robert, 26
Solenopsis, 443
Solenoptilidae, 336
Solfugida, 99, 103, 104, 437
 Solnhofen, Germany (Jurassic), 42, 43, 74, 184, 187
 Soyana, Russia (Permian), 69
Spania, 522
Spaniidae, 522
 Spanish (Alava) amber (Cretaceous), 79, 271, 518, 527
 Spanish fly, 390
Spargotermes limai, 92, 244
 specialized pollinators, evolution of speciation rates, ultimate causes, 573
 species concept: biological, 6, 9; evolutionary, 6; phylogenetic, 6
 species diversity in insects, 3–4, 6–15: and age, 1–3, 646; and amber fossils, 642; and design, 646; extinction rates, 636, 637; flight, 646; larva, 646; metamerism, 646; prospects, 650; rate of increase, 647; redundant appendages, 646; significance of exoskeleton, 646
 species lifespans in insects, 642–43
 species, 6–15: concepts, 6–11, 15; cryptic, 8; defined, 6, 9; described numbers (Table 1-1), 12; named hexapods, 12; rarity, 14; semi-, 8; sibling, 8; undescribed numbers, 12–14
 sperm pump, 468
Sphaeropsocidae, 270
Sphaerotermes, 248
Sphaerotermitinae, 248
Sphecidae, 451
 spheciiform wasps (see Apoidea)
Sphecodorydaloididae, 172
Sphecoidea, 454
Sphecomyrma, 40, 82, 446: *canadensis*, 446; *freyi*, 446, 448; reconstruction, 448
Sphecomyrmina, 36, 446
 sphecophiles, 377
Sphecopteridae, 172

INDEX

753

Sphenisciformes, as hosts, 278
Sphex, 454
 Sphingidae, 583, 584, 586
Sphyracephala: fossils, 88
 Spicipalpia, 550
 spiders (see Araneae)
 spiderweb cleptoparasites, 590
Spiloconis, 352
 spiracles, 125, 178
 spirochetes, 240, 272
 spittle bugs (see Cercopoidea)
 split-footed lacewings (see Nymphidae)
 spongillaflies (see Sisyridae)
 Spongiphoridae, 220
 spoon-winged lacewings (see Nemopteridae, Nemopterinae)
 springtails (see Collembola)
Srokalarva, 53, 335
 St. Helena earwig, 648
 Staatlichen Museum für Naturkunde in Stuttgart, 91
 stalk-eyed flies, 30, 541, 543
Stantoniella, 235
 Staphylinidae, 71, 358, 363, 372–377: classification, 374; diversity, 374; fossils, 375–7; habits, 377; in amber, 375; Jurassic, 374; oldest, 376; Triassic, 374; world catalogue, 374
 Staphyliniformia, 371–77: diversity, 372; ectoparasites, 373
 Staphylinoidea, 361, 372, 399, 630
 Statiomyidae, 618
 Stebliidae, 543
Steingelia cretacea, 299
 Steingeliidae, 298, 299
 stem group, 39, 51, 73, 81, 94, 96, 100, 107, 137, 163, 176; general, 40
 stemmata, 124, 331
Stenodictya lobata, 168
 Stenogastrinae, 437, 438
 Stenopelmatidae, 207, 208, 209, 628
 Stenopelmatoidea, 208
 Stenotritidae (see Colletidae)
 Stensham, Forthampton, UK (Triassic), 71
 Stephanidae, 417
 Stephanocircidae, 481, 485
 Stephanocircinae, 485
Stephanogaster magna, 416
 Stephanoidea, 414–415
 Sternorrhyncha, 288, 289–303, 338, 341; and angiosperm radiations, 302–3; and ants, 300–302; and domatia, 302; and myrmecophytes, 302; classification, 289; defense by ants, 301; early fossils, 289; features, 289; fossilized symbioses, 302; infraordinal features, 304; mouthparts, 289; population growth, 289; prepupae, 290; pupae, 290; tending by ants, 289; see also: honeydew
 sternum, 121
Stichotrema, 406
 stick-insects (see Phasmatodea)
 Stigmaphronidae, 415, 417, 426, 637: in Cretaceous ambers, 427
 sting, 429–30, 457
 stingless bees (see Meliponini)
 stipes, 124
Stizoides, 453
Stolotermites, 228, 240, 247, 628

Stomoxys, 489, 545: *calcitrans*, 542
 stoneflies (see Plecoptera)
 Stormberg Series, 72
Strashila incredibilis, 473, 474
 stratigraphic-clade rank correlation, 38
 Stratiomyidae, 492, 493, 516
 Stratiomyomorpha, 516, 519
 Streblidae, 545, 640
 Strepsiptera, 12, 201, 331, 390, 399–406, 494: adult mandibles, 469; and long-branch attraction, 402; biology, 399; cephalotheca, 399, 401; classification, 402; Cretaceous male, 404; diversity, 402; features, 400; females, 399, 401; fossils, 403–6; high DNA substitution rates, 402; hosts, 400; larval internal defense, 400; male halteres, 399; relationships, 399, 402; reproduction, 400; structure, 402, 403; triungula, 400
 stridulation, 367: in Lepidoptera, 599; in Orthoptera, 203–4; in Titanoptera, 215–6
 Sturtevant, Alfred, 26
 styli, 117, 131, 132, 149
 Stylopidae, 402
 Stylopida, 402, 406
Stylotermes, 247
 subalare, 128
 subcostal vein
 subcoxal theory, 125
 subimago, 128
 subsocial behavior, 283, 464
 subspecies, 9
 Subulicornia, 167
Succinapis, 466
 sulcus, 120, 122
 sunbirds, 615
 Sunchal, 92
 Sundance Formation, Wyoming (Jurassic), 76
 Suntrana, Alaska (Pliocene), 90
 suture, 120–122
 swallowtails, 595
 Swammerdam, Jan, 17, 18
 swiftlets as hosts, 278
 Switzerland, 89
 Swofford, David, 33
 Sycorinae, 490
Syllepte, 578
Sylvacoleus sharovi, 362
Sylvicola, 494, 517
Symporomyia, 489, 522
 Symphrasinae, 353
 Symphyta, 99, 108, 109
 Symphypleona, 115
 Symphyta, 409, 411
 symplesiomorphy, 27
Symplocarpus, 237
Sympycnites primaevus, 529
Synalpheus, 467
Synanthonedon, 579
 synapomorphy, 27, 29
Synchlora, 587
Syndixa, 505
 synonyms (taxonomic), 14
 Syntermiteinae, 248
Syntexis libocedrii, 412
 Syntonopteridae, 163, 164
 Syntonopteroidea, 163
 Syrphidae, 426, 492, 493, 537, 540, 618: fossil, 541

Syssphinx, 582
Systelloderes, 316
 systematics: defined, 15; history of 16–27; molecular, 289–32
Systolosoma, 366

T
 Tabanidae, 489, 490, 492, 522, 523: Cretaceous, 523; fossil, 523, 524
 Tabanomorpha, 519, 522, 523
 Tachinidae, 492, 543
Tachinus caelatus, 61
Tachyglossus aculeatus, 638
 Taeniopterygidae, 196
Taeniothrips, 283
 tagmata, 119
 tagmosis, 97
 Taldycupedidae, 362
 Tallahatta Formation, Mississippi (Eocene), 86
Tamandua, 638
 Tanaoceridae, 211
 Tanaoceroidea, 210
Tanaorrhinus, 582
 Tanaostigmatidae, 424
 Tantulocarida, 99
Tanychora, 429
 Tanyderidae, 498, 499: Cretaceous, 499
Tanypyscha, 499
 taphonomy, 42
Tapinoma, 443
 tarantulas, 102
 Tardigrada, 45, 93, 94, 96, 97: Cambrian, 96; cryptobiosis, 97; fossils, 96, 97; general, 96; in Cretaceous amber, 97
 tarsomeres, 127
Tarsophlebia, 183
 Tarsophlebiidae, 183
Tarsophlebioptera, 183, 184
 tarsus, 125
Tarwinia australis, 485–488
 toxic analyses, 37, 623, 636: criticism, 636
 Taxodiaceae, 76
Taxodium, 60
 taxon cycle, 631
 taxonomy, 33: general, 35
 Taymyraphididae, 295, 302
Tegeticula, 572, 573, 616: *corruptrix*, 573; *intermedia*, 572
 tegula, 129, 130
Telchin, 578
Telmatogeton, 504
 telotrophic ovarioles, 335
 telson, 131, 132
 Tenebrionidae, 358, 359: fossil, 389, 391; in arid habitats, 389; parental behavior, 389
 Tenrecidae, 638
 tenrecks, 638
 tent caterpillars, 583
 Tenthredinidae, 409, 410
 Tenthredinoidea, 410–412
 tentorium, 123, 137
 Tephritisidae, 492, 541, 624
 Tephritoidea, 541, 623: Eocene, 546
 Teratombiidae, 198
 Terebrantia, 284
 Termitaphididae, 326, 329: fossil, 329
Termitaradus: fossil, 329
 termite inquilines, 326

- Termitella*, 377
 termites (see Isoptera)
 Termitidae, 238, 239, 240, 242, 247, 250, 251, 377, 467, 638; classification, 248; diet, 238, 244; diversity, 247; ectosymbionts of, 377; nasutes, 249; nests, 51, 240, 242; soldiers, 239, 249; symbiotic fungi of, 248
 Termitinae, 248
Termitomyces, 248, 249
 termitophiles, 377
 Termopsidae, 238, 240, 242, 244, 247, 250, 251, 628; distribution, 242; diversity, 247; fossils, 247
Termopsis, 247
 terrestrialization, 109
Terrindusia, 554
 Tertiary radiations, 528, 539, 637
Testajapyx thomasi, 118
Tetraphalerites, 365
Tetraphalerus, 363, 365
 Tetrigidae, 205, 210
 Tetrigoidea, 210
Tettigarcta, 308, 630
 Tettigarctidae, 308
 Tettigometridae, 312
 Tettigoniidae, 127, 178, 205–7, 208: camouflage, 206, 207; nuptial feeding, 203; Tettigonioidea, 209
Thasus, 327
 Thaumaleidae, 504, 505
Thaumastobombus, 466
 Thaumastocoridae, 326, 329, 330
Thaumatoxena, 377
 Thecostraca, 99
Theope, 599
Therea, 237
Thereus, 598
 Therevidae, 521
 Theridiidae, 103
Thermobia domestica, 151
 Thomisidae, 103
Thomomydoecus, 280
 thoracic structure, 125–131
 thorny lacewings (see Rhachiberothidae)
 Thraupidae, 615
 thread-winged lacewings (see Nemopteridae, Crocinae)
 Thripidae, 284, 285, 286
 thrips (see Thysanoptera)
 Throscidae, 382, 383
 Thuringia, Germany (Triassic), 71
Thuringopteryx, 163, 165
 Thyrididae, 583
Thysania agripinna, 178
 Thysanoptera, 12, 147, 199, 280–87, 261, 262, 263, 623, 642; austral distributions in, 284; castes, 283; classification, 284; cleptoparasites, 283; communal behavior, 283; development, 282; diets, 283; diversity, 284; feeding, 282, 283; fossils, 285–287; gall forming, 283; haplodiploidy, 284; phylogeny, 286; polymorphism, 280, 283; prepupa, 282; pretarsus, 280, 281; pupa, 282; relationships, 284, 287; social behavior, 283; structure, 280, 281; wings, 286
 Thysanura, 150
 tibia, 125
 Tien Shan mountains, 71
 Tillyard, Robin J., 139, 140
Tillyardembia antennaeplana, 223
 Tillyardembiidae, 223
 Tillyardipteridae, 499
Timema, 189, 213, 215, 630
 Timematidae, 213
 Timematodea, 195, 213: defining features, 213
 tinamous, as hosts, 277
Tinaphis, 295
Tinea, 574
 Tineidae: fossil cases, 51
 Tineoidea, 574–575
Tineola, 574
 Tingidae, 319, 326, 329, 330
 Tingoidea, 328
 Tiphidae, 434, 630; Cretaceous, 434, 498
 Tipulidae, 492; fossils, 71, 498, 620
 Tipuloidea, 494, 496, 498, 517; Cretaceous, 470, 472, 475, 476, 485; fossils, 469, 470
 tissue preservation in fossil insects, 59, 60
 titanic crawlers (see Titanoptera)
 Titanoptera, 147, 191, 193, 215–216; stridulatory structures, 216
Titanus, 379
 Toldito Formation, New Mexico (Jurassic), 76
 Tolgoy Formation, Kazakhstan, 71
 Tongchuan Formation, 71
 Torridincolidae, 370
 Tortricidae, 572; fossil, 581
 Torymidae, 418, 420
Toxorhynchites, 506
 Toxorhynchitinae, 506
Trabutina mannipara, as source of manna, 297
 trace fossils, 50, 54, 92, 380, 551, 637: burrows, 50, 51, 54, 55, 92; coprolites, 50; frass, 50, 52, 53, 54; larval cases, 51; nests, 50
 tracheae, 125, 178
 Tracheata, 99, 107
Trachymyrmex, 278
 Trachypachidae, 363, 366, 367, 368, 630: Jurassic, 367
Trachypachus, 366
 Traditional, 334–35
 traps (Pleistocene/Holocene), 61
 tree hoppers (see Membracoidea)
 Tremembé Formation, Brazil (Oligocene), 92
 Tremicinae, 412
 Trench Fever, 272
 Triadophlebiidae, 183
 Triadophlebiomorpha, 183; defined, 183
 Triadotypidae, 183
 Triaplidae, 367
 Triassic, 4, 51, 52, 54, 70–73, 369: fossil insect deposits, 70–73
 Triassocoridae, 319
 Triassolestidae, 184
 Triassomachilidae, 148
Triassomachilis uralensis, 149
Triassothrips virginicus, 285, 286
 Triatominae, 323, 490
Tribolium, 389
 trichobothria, 326, 327
 Trichoceridae, 498
 Trichodectidae, 273, 275, 276, 277, 280
Tricholepidion gertschi, 151
 Tricholomataceae, 248, 278, 444
 Trichomatinae, 356
 trichomes, 377
 Trichomonadidae, 240
Trichonympha, 240, 241
 Trichonymphidae, 240
Trichophaga, 574
 Trichophilopteridae, 275
Trichoprosopon, 507
 Trichoptera, 12, 51, 147, 469, 548–555: adult structure, 549; biogeography, 552; bipolar distribution, 555; burrowing, 549; case building behavior, 549; classification, 550; corema, 550; earliest fossil, 550; ecological groups, 548; features, 560, 563; fossil cases, 51, 76, 86, 552, 555; fossils, 51, 554; free-living, 549; gills, 548; haustellum, 549; in Cretaceous amber, 555; larval features, 552; larval cases, 548, 550; life histories, 548, 555, 569; marine, 548; net spinners, 549; number living species, 545; oldest, 555; phylogeny, 553; predators, 549; prolegs, 548; purse case, 549; retreat makers, 549; saddle-case makers, 549; tube-case makers, 549; world species, 548
 Tricoleidae, 363
 Tridactylidae, 205, 210, 211; Cretaceous, 211
 Tridactyloidea, 203, 211
 Trigonalyidae, 417, 418
 Trigonalyoidea, 415, 417
 Trigonopterygidae, 211
 Trigonopterygoidea, 211
 Trigonotarbida, 99, 101
 Trilobita, 49, 98–100, 124
 Trimenoponidae, 273, 274
Triops cancriformis, 97
 Trioza, 291
Triplosoba pulchella, 164
 Triplosobiidae, 164
 Tristiridae, 210
 Tritocerebra, 94
 trochantellus, 126
 trochanter, 125, 126
 Trochilidae, 615
 Troctomorpha, 266
 Trogidae, 270, 378
 Trogiomorpha, 266
 troglobites, 230
Troides, 556, 595
 Troidinae, 597, 604
 tropical forest canopies, 12
 true bugs, 314
Trypanosoma: brucei, 545; *cruzi*, 326; *gambiense*, 545
Trypoxylon politum, 437
 tsetse flies (see Glossinidae)
 Tshekarda, Russia (Permian: Koshelevka Form), 69
 Tshekardocoleidae, 361, 362
 Tubulidentata, as hosts, 274
 Tubulifera, 284
Tunga, 485
 Tungara frog (see *Physalaemus pustulosus*)
Tupus gracilis, 175
 Turanodermatidae, 218

INDEX

755

- Turanophlebia*, 184
Turanothemistidae, 187
Turanoviidae, 218
Turonempis styx, 531
 twisted-winged parasites (see Strepsiptera)
Tychtodelopteridae, 341
 tympanum (auditory), 203; in
 Auchenorrhyncha, 304–308; in
 Lepidoptera, 583–585; in Mantodea, 257;
 in Orthoptera, 204; in Psocoptera, 266
Tymanogaster, 371
 type specimens, 35, 584; holotype, 35, 153;
 lectotype, 35; name-bearing, 35; neotype,
 35; paratype, 35
Typhoctes, 437
- U**
Ulmaceae, 410, 413
Ulmeriella, 244
Ulomites, 391
Ultracoelostoma, 300; honeydew, 300
Ululodes, 347, 348
 Umbelliferae, 624
Umenocoleidae, 220, 229, 234, 235
Unicalcarida, 409
 uniformitarianism, 24
Uniramia, 98
 University of California Museum of
 Paleontology, Berkeley, 91
 University of Kansas, 26
Upiga virescens, 573
Uraloptysmatidae, 548
Urania, 585
UranIIDAE, 569, 583, 585
Urodus parvula, 576, 577
Uropsylla tasmanica, 485
Uropygida, 99, 105
 Usinger, Robert, 26
Uskatelytridae, 332
 Ussher, Bishop, 23
Uzelothropidae, 284, 285
Uzelothrips scabrosus, 284
- V**
Vachonisia, 98
Valditermes, 244; *acutipennis*, 250; *brenanae*,
 250
Valeseguya, 500; *disjuncta*, 501
 valifers, 132, 134, 229
 valvulae, 132
 Van Houten cycles, 72
Vanhorniidae, 418, 421
Veliidae, 317, 321
Velocipedidae, 329
 velvet worms (see Onychophora)
Vermileonidae, 516
Vermipsyllidae, 480, 481, 485
 Vertebrata, 467
 vertebrate parasitism, 543
Vespidae, 353, 437, 438, 466, 467; in amber,
 439; nests, 439; relationships, 439
Vespina (see Euhymenoptera)
Vespinae, 437, 438
Vespoidea, 430, 433–451
- vestibulum, 229
 vicariance biogeography: criticsms, 631;
 defined, 631; geographic extinctions,
 631–34
 viruses, symbiotic, 427–428
Vitimotauliidae, 551
 viviparity, 230, 231, 543
Vladipteridae, 498
 vomer, 191, 211, 226
Vorkutiidae, 172
 Vosges, France, 71
- W**
 walking-sticks (see Phasmatodea)
 Wallace, Alfred Russel, 24, 25
 walnut (see Juglandaceae)
 wasps (see Apocrita)
 water bears (see Tardigrada)
 water hyacinth, 210
 water striders (see Gerrymorpha)
 wax secretion, 289, 290, 299
 Wealden deposits (Cretaceous), 78
 webspinners (see Embioidea)
 weevils (see Curculionoidea)
 Wegener, Alfred, 627
 Wellington Formation (Permian), 69
Welwitschia mirabilis, 608
 wetas (see Orthoptera, Ensifera)
Whalfera venatrix, 355, 356
 whip spiders (see Amblypygida)
 white flies (see Aleyrodoidea)
 Wigglesworth, Sir Vincent B., 142
 Wilcox Group (North America: Eocene), 87
 Willmann, Rainer, 144
 Wilson, E. O., 14, 631
 wing articulation, 130
 wing cells, 131
 wing coupling, 262, 265, 313, 315, 594
 wing function, 156–157
 wing scales, 556
 wing structure, 128–131, 617, 618
 wing veins, 128–129; naming, 190, 217, 231,
 259, 268, 521; structure, 158, 178
 wing venation, 9, 43, 159, 128, 184
 wings, 128–131, 155, 156, 158, 159; function,
 156; gill theory, 159; origin of, 154,
 158–160; paranotal lobes, 158; pronotal
 lobe theory, 159
 Winteraceae, 613, 621
 wood borings: fossil, 54
 wood roach (see *Cryptocercus*)
 wood wasps (see *Sympyta*, Siricoidea,
 Xiphidiocioidea)
 Woodworth, Charles, 143
 worker polymorphism, 444
 Wright, Sewell, 26
 Wurthiinae, 581
Wyeomyia, 502
- X**
Xamenophlebiidae, 183
Xanlothrips, 283
Xanthopan, 584; *morgani praedicta*, 585
Xenopsylla, 480; *cheopis*, 480, 482
- Xenopteridae**, 172
Xenozorotypus burmiticus, 201
Xenusia, 94
Xeris spectrum, 412
Xiphiosyllidae, 480
Xiphopterum, 215
Xiphosura, 97, 99; defined, 100; fossils, 74,
 97, 100
Xiphydriidae, 409, 411, 417, 630
Xiphydrioidae, 413
Xyelidae, 71, 409, 410, 616; Cretaceous, 410;
 Triassic, 409–410
Xyloidea, 409
Xyelotomidae, 409, 411
Xyelydidae, 409, 412
Xyleborini, 397
 xylococcid, 299
Xylotopa, 461, 456; *virginica*, 456
Xylocopinae, 467
Xylocopini, 435
Xylomyidae, 516
Xylophagidae, 493, 516, 520
Xylophagomorpha, 516
Xyloryctidae, 576
Xyronotidae, 211
- Y**
Ycaploca, 630
Ycaploca evansi, 431
 yellowjackets (see Vespidae)
 Yixian Formation (Cretaceous), 470, 478,
 523, 611, 616, 618
Yponomeuta, 557
Yponomeutidae, 577
Yponomeutoidea, 576
 yucca moths (see Prodoxidae)
- Z**
Zacryptocerus, 441; colony, 442; in amber,
 443
Zelmiarcha lebanensis, 510
Zeuzera, 579
 Zingerberaceae, 53
Zootermopsis, 247
 Zoraptera, 12, 147, 189, 191, 199–201;
 classification, 199; defining features, 199,
 200; dimorphism, 199; fossils, 201;
 phylogenetic position, 199–201
 zorapterans (see Zoraptera)
Zoropsocus, 286, 287; *itschetuensis*, 287;
 stanleyi, 285
Zorotypidae, 199, 200, 201
Zorotypus, 199, 201; *goeleti*, 201; *hubbardi*,
 200; *nascimbenei*, 201
Zygaena, 578
Zygaenidae, 579
Zygaenoidea, 579, 580
 Zygentoma, 12, 147, 149, 150–152, 160, 628;
 classification, 150; courtship, 151; defining
 features, 150; fossils, 152, 154
Zygophlebiidae, 183
 Zygoptera, 179, 180, 182, 183, 184; defined,
 183; fossil, 185; in amber, 185; mating
 wheel, 182