

## Fever Medication: Dosage Chart by Weight

\*The information contained on this Web site should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstance

### Tylenol® Dosage Table for Fever and Pain

#### Acetaminophen Dosage Table for Fever and Pain\*

Child's Weight (pounds)	6-11	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+	
Syrup: 160 mg/5 mL	1.25	2.5	3.75	5	7.5	10	12.5	15	20	mL
Chewable 80 mg Tablets	--	--	1 ½	2	3	4	5	6	8	tabs
Chewable 160 mg Tablets	--	--	--	1	1 ½	2	2 ½	3	4	tabs
Adult 325 mg Tablets	--	--	--	--	--	1	1	1 ½	2	tabs
Adult 500 mg Tablets	--	--	--	--	--	--	--	1	1	tabs

\*Read table notes below.

### Motrin®, Advil® Dosage Table for Fever and Pain

#### Ibuprofen Dosage Table for Fever and Pain\*

Child's Weight (pounds)	12-17	18-23	24-35	36-47	48-59	60-71	72-95	96+	
Infant Drops 50 mg/1.25 mL	1.25	1.875	2.5	3.75	5	--	--	--	mL
Liquid 100mg/5 mL	2.5	4	5	7.5	10	12.5	15	20	mL
Chewable 50 mg tablets	--	--	2	3	4	5	6	8	tabs
Junior strength 100 mg tablets	--	--	--	--	2	2 ½	3	4	tabs
Adult 200 mg tablets	--	--	--	--	1	1	1 ½	2	tabs

\*Read table notes below.

## Table Notes:

- **Age limit:** Don't use under 6 months of age unless directed by your child's pediatrician, because safety has not established and doesn't have FDA approval. Avoid multi-ingredient products in children under 6 years of age.
- **Dosage:** Determine by finding child's weight in the top row of the dosage table.
- **Measuring the dosage (should be in metric units):** Dosing syringes are more accurate than household utensils. Use the syringe or device that comes with the medication. If not available with the medication, medicine syringes are available at pharmacies. Household spoons are not reliable. **Note:** 1 level teaspoon equals 5 mL and that ½ teaspoon equals 2.5 mL.
- **Ibuprofen drops:** Ibuprofen infant drops come with a measuring syringe
- **Adult dosage:** 400 mg
- **Frequency:** Repeat every 6-8 hours as needed.

\*For more information, go to [www.healthychildren.org](http://www.healthychildren.org).