

ECOLOGY OF PLANT HUMMINGBIRD INTERACTIONS IN MAQUIPUCUNA, ECUADOR

Tatiana Santander, Esteban Guevara, Francisco Tobar, Holger Beck, Nicole Büttner, Andrea Nieto, Andrés Marcapata, Friederike Richter, María José Gavilanes, Cristian Poveda, Bryan Rojas, Rafael Wüest, Carolina Bello and Catherine H. Graham



October 22, 2020



Alaspungo



Contents

1. Introduction and project overview	1
2. Methodological Approach	2
Field transects	2
Time-lapse cameras	5
3. Resulting patterns	5
Plant-hummingbird interactions	5
Plants information and phenology	9
The Network of Interactions	12
4. Conclusions:	19
Acknowledgements	20

1. Introduction and project overview

One of the main hypotheses for how so many related species can co-occur is resource-partitioning where species use different resources, which limits competition among species and allows them to co-exist. In the case of hummingbirds and plants, each hummingbird species forages on a distinct set of flowers and each flowering plant species is visited by a subset of hummingbirds. Interactions between plants and hummingbirds are mutually beneficial. These mutualistic hummingbird-plant interactions are important from a hummingbird perspective because hummingbirds require nectar to fuel their high-energy lifestyles where they often hover – an energetically costly behavior – to take nectar. From a plant perspective most hummingbirds pollinate flowers as they forage on nectar, though some hummingbirds take nectar from the base of the flower, cheating the flower from this service of pollination. The intricate web of interactions between hummingbirds and their food plants evolved over millennia as a result of diffuse co-evolution which yielded a remarkable array of morphological forms and functions. On-going human activities, such as deforestation and climate change threaten these interaction webs, yet little is known as to how hummingbirds and their food plants will respond. To understand the influence of humans on this complex relationship, accurate, high quality data on hummingbird and flowering plant occurrence and hummingbird-plant interactions are required across broad regions and over an elevation range.

The Northwest slope of the Andes of Ecuador is an ideal place to study plant-hummingbird interactions because it is among the most biodiverse places on earth where multiple co-occurring species rely on each other for survival. There are ~360 species of hummingbirds on earth with the highest diversity in the Andes where up to 30 species can be found at a single site and ~1600 vascular plant species have been recorded in the region. Our study region was in the Pichincha Province (latitude 0°12' N to 0°10' S, longitude 78°59' W to 78°27' W) and covers 107 square kilometers with an elevation range from 800 to 3500 meters. Our sampling location in Maquipucuna reserve lies between 1534 and 1726 meters along this gradient.

The goal of the project was to determine the abiotic and biotic factors driving variation in hummingbird-plant interaction networks across elevation and land-use gradients. By evaluating these mutualistic interactions we are able to predict how diversity of both hummingbirds and plants will be influenced by elevation and anthropogenic activities. The project is led by Dr. Catherine Graham from the Swiss Federal Research Institute and executed by Aves y Conservación/BirdLife in Ecuador, Santa Lucía, Maquipucuna, and Un Poco del Chocó with collaboration of several reserves including Mashpi, Las Grallarias, Amagusa, Sachatamia, Yanacocha (Fundación Jocotoco), Verdecocha, Puyucunapi (Mindo Cloud Forest), Rumisitana, Pontificia Universidad Católica del Ecuador, and Alaspungo community. At Maquipucuna we collaborated with Rebeca Justicia, and Rodrigo Ontaneda from Maquipucuna Foundation.

2. Methodological Approach

To monitor abundance patterns, flowering phenology and hummingbird flower visitation we used a combination of field transects and time-lapse cameras. These transects were 1.5 km in length and were spread across the elevation and land-use gradient with 1 to 2 transects per site. We visited each of the 18 transects (11 in forest and 7 in disturbed sites) one time per month during a two year period. In Maquipucuna we sampled the transects from April 2017 to June 2019.




Figure 1: Location of the site in the elevation gradient.

Field transects

In Maquipucuna we have 1 transect of 1.5 km. The transect starts about 2,5 km from the Maquipucuna lodge and follows a part of the main trail where tourists hardly ever go. Following the soft slope through a mature secondary forest, the transect reaches a plateau at an elevation of 1600 masl. From here on the trail is characterized by a steep climb and several sharp bends. The forest matures as the transect continues. Natural disturbances such as landslides and tree falls are common in this part. Following a ridge where trees of the Melastomataceae family dominate, the trail reaches its last part with primary forests (sightings of Andean Bears are not uncommon in this area). Passing a vantage point with a beautiful view at around 1750 m of elevation the trail descends for about 150 meters. Here - surrounded by giant fig trees and dense primary forest vegetation - the transect ends (Figure 2).




Figure 2: Elevation gradient of the transect.

Along each transect, four to five kinds of data were taken:

- **Flower counts:** Any plant with hummingbird syndrome flowers within a distance of ~5 meters of the transect was counted and identified to species. Characteristics of a flower with the hummingbird syndrome include brightly colored flowers (purple, red, orange or yellow) with medium to long corollas. While most species hummingbirds use have these characteristics we were conservative and monitored any questionable species or plants we have seen hummingbirds feeding. For each plant either all flowers were counted or in the case of bushes with more than ~100 flowers, total flowers on 5 representative branches were counted and used to extrapolate the number of flowers on the plant. Each species was collected once and pressed in order to archive our work and/or verify identification with an expert. Plant specimens were deposited at the Herbarium of Catholic University in Quito and Ibarra.
- **Interaction observations:** During the flower census, any interaction of a hummingbird with a flower was noted.
- **Hummingbird counts:** Any hummingbird heard or seen at a distance of 20 meters was also noted.
- **Flower morphology:** Several flower morphological features were measured on at least three individuals per species wherever possible. The Flower traits included were: a) flower corolla length, the distance from the flower opening to the back of corolla, b) effective corolla distance by cutting open flowers and measuring the corolla length extending back to the flower nectarines, c) corolla opening, d) stigma and anther length.
- **Nectar concentration:** This data was taken only at three sites corresponding to low, medium and high transects. Sugar concentration was collected at flowering species for up to 12 flowers per species using a refractometer (a capillary tube is used to extract nectar).



Figure 3: Team researcher, Andreas Nieto, counts flowers along a transect.

Time-lapse cameras

We used time-lapse cameras to monitor hummingbird-plant interactions. Time-lapse cameras, which take a picture every second, were placed at individual flowers along the above described transects to capture visitation by hummingbird species. We placed cameras on all flowering plants along the transect roughly proportional to their abundance. The cameras turn on at dawn and record an image every second for several days, resulting in a dataset of millions of images. These images are efficiently processed using Motion Meerkat or Deep Meerkat which can be used to sort out images with hummingbirds which can be manually identified (in the past we have been able to identify 95% of birds in images). This approach minimizes reliance on time-consuming human flower observations, greatly increasing data collection in time and space permitting a rigorous test of network theory.



Figure 4: Team researcher Holger Beck shows how a camera is set up in order to film a flower.

3. Resulting patterns

Plant-hummingbird interactions

Maquipucuna reserve is one of the largest private properties that protects 66 plant species used by hummingbirds according to our project results (Annex 1). However, in our cameras we recorded 164 different interactions between 12 hummingbirds and 47 plants (Figure 5).



Figure 5: Examples of some of the hummingbirds and plants we caught in cameras.

Table 1: List of hummingbirds and number of interactions.

<i>Hummingbird</i>	No of interactions	No plants interacting
<i>Agelaiocercus coelestis</i>	518	28
<i>Phaethornis syrmatorphorus</i>	478	27
<i>Phaethornis yaruqui</i>	175	24
<i>Phaethornis striigularis</i>	410	23
<i>Coeligena wilsoni</i>	306	21
<i>Ocreatus underwoodii</i>	272	20
<i>Urosticte benjamini</i>	32	8
<i>Doryfera ludovicae</i>	31	4
<i>Heliodoxa jacula</i>	7	4
<i>Thalurania colombica</i>	4	3
<i>Colibri coruscans</i>	1	1
<i>Heliodoxa rubinoides</i>	4	1

The most common hummingbird recorded was *Agelaiocercus coelestis* and the most common plant was *Renealmia sessilifolia*. Although they are the most common species, they are not necessarily the species that interact with more species. The hummingbird that interacts more is *Agelaiocercus coelestis* and the plant that has more interactions is *Cavendishia grandifolia*. In table 1 and 2 we can observe the number of interaction for each species.

Table 2: List of plants and number of interactions.

Plant	No of interactions	No hummingbirds interacting
<i>Cavendishia grandifolia</i>	103	9
<i>Centropogon solanifolius</i>	136	7
<i>Gasteranthus quitensis</i>	165	7
<i>Heliconia impudica</i>	86	7
<i>Columnea kuczyniakii</i>	33	6
<i>Palicourea demissa</i>	29	6
<i>Psammisia ulbrichiana</i>	42	6
<i>Renealmia sessilifolia</i>	313	6
<i>Burmeistera multiflora</i>	59	5
<i>Columnea ciliata</i>	49	5
<i>Columnea eburnea</i>	134	5
<i>Costus pulverulentus</i>	87	5
<i>Glossoloma purpureum</i>	99	5
<i>Guzmania jaramilloi</i>	136	5
<i>Palicourea sodiroi</i>	29	5
<i>Burmeistera crispiloba</i>	6	4
<i>Columnea sp.</i>	37	4
<i>Guzmania rhonhofiana</i>	58	4
<i>Macleania smithiana</i>	15	4
<i>Psammisia sodiroi</i>	43	4
<i>Renealmia dolichocalyx</i>	231	4
<i>Bomarea pardina</i>	43	3
<i>Columnea picta</i>	17	3
<i>Erythrina megistophylla</i>	37	3
<i>Guzmania lehmanniana</i>	7	3
<i>Guzmania xanthobractea</i>	24	3
<i>Heliconia sclerotricha</i>	9	3
<i>Heliconia sp.</i>	7	3
<i>Psammisia aberrans</i>	18	3
<i>Besleria solanoides</i>	8	2
<i>Dicliptera scabra</i>	35	2
<i>Drymonia tenuis</i>	7	2
<i>Elleanthus robustus</i>	86	2
<i>Heliconia virginalis</i>	7	2
<i>Kohleria spicata</i>	4	2
<i>Podandroyne sp1</i>	5	2
<i>Tillandsia cyanea</i>	5	2
<i>Wercklea ferox</i>	4	2

<i>Centropogon nigricans</i>	5	1
<i>Cuatresia riparia</i>	4	1
<i>Drymonia brochidodroma</i>	3	1
<i>Fuchsia macrostigma</i>	1	1
<i>Justicia secunda</i>	1	1
<i>Kohleria villosa</i>	2	1
<i>Microchilus sp.</i>	4	1
<i>Psammisia cordifolia</i>	3	1
<i>Sobralia tamboana</i>	2	1

Plants information and phenology

We recorded the abundance of flowers from April 2017 to June 2019. The months with higher abundance of flowers are November and May (Figure 6).




Figure 6: Abundance of flowers by month. Points represent the sum of flowers at each month and the black line represents the mean trend.

However, not all plant produces flowers at the same time. In figure 7 we can observe the phenology of the four most common plant species.




Figure 7: Phenology of most common flowers by month. Points represent the number of flowers counted in each month and the line represents the mean trend. Each color represents a different plant species.

Below we describe the most representative plant families present in Maquipucuna.

GESNERIACEAE

Gesneriaceae, the African violet family has around 3000 species, distributed mainly in Central and South America, East and South Asia, Europe and Oceania. In Ecuador there are 200 species grouped in 25 genera. They could be herbs (*Kohleria*, *Diastema*), shrubs (*Glossoloma*, *Columnea*) or very rarely small trees (*Shuaria*, *Besleria*). Gesneriaceae usually have opposite leaves, axillary or terminal inflorescence (cyme, raceme or fascicles), flowers with five petals joined to form a colorful tube with 4 or 5 lobes. Four didynamous stamens (two longer and two shorter) generally fused together and located at the dorsal part of the flower, a simple elongated style with the stigma usually bilobed. In the Pichincha province 15 genera and 89 species have been reported. In our study 64 species were registered, 12 are endemic, 6 are endangered (EN), and 6 are vulnerable (VU). Additionally, we found 3 species that were not previously reported for Pichincha, 2 new records for Ecuador, and 5 new species. Maquipucuna has 15 species being *Columnea* the most representative with six species. Additionally, *Drymonia collegarum* is endemic and vulnerable of extinction (VU), and there is also a new *Columnea* species shared with Las Gralarias, Puyucunapi and Sachatamia.

ERICACEAE

Ericaceae also known as the blueberry family as “mortiño” is represented by 125 genera and 4000 species, widely distributed in temperate, subarctic, and also at high elevations in tropical regions. In Ecuador 21 genus and 240 species have been reported. Life forms include woody shrubs (*Cavendishia*, *Macleania*), trees (*Bejaria*, *Thibaudia*), or suffrutex (small plants with woody stems and soft branch as *Gaultheria*, *Disterigma*). Plants could be erect, prostrate or climbers with coriaceous leaves. Flowers are perfect (containing anther and stigma), mostly tubular with 4 to 7 lobes, anthers in twice number than the petals, often enlarger in one or two terminal tubes. Fruit usually is a capsule, berry or drupe. In Pichincha province there are 13 genus and 73 species. During EPHI project 45 species were registered and 18 are endemic: one is critically endangered (CR), four are endangered (EN), and 10 species are vulnerable (VU). *Macleania tropica* is the first record for Pichincha area, it was only known from Esmeraldas and Colombia. *Antoptherus ecuadorensis*, and *Macleania alata* are the first records made since the type collection in 1979 and 1986 respectively (these two species were collected nearby the study transects).. Maquipucuna has 15 species being *Psammisia* and *Macleania* the genus with more species. Only *Macleania recumbens* is endemic and vulnerable (VU).

The Network of Interactions

The interaction data we collected can be used to explore how the interactions network is organized at Maquipucuna. In figure 8 we show the structure of the network.

By analyzing the network structure, we found that the plant *Cavendishia grandifolia* and the hummingbird *Aglaiocercus coelestis* are the key species that holds the network together. If they are lost, the network will become less stable. By contrast, *Tillandsia cyanea* and *Heliodoxa rubinoides* are very specialized species which means they interact with a small group of specialized species. In table 3 we can observe the plants interacting with each hummingbird.




Figure 8: Network of interactions. Blue represents hummingbirds and green plants. Each line represents an interaction between a hummingbird and a plant obtained from our camera observations. Thicker lines indicate that the interaction was common while very thin lines indicate that the interaction occurred rarely. The size of the colored bar shows the number of interactions of a hummingbird or plant participated in an interaction.

Table 3: List of Hummingbirds with the plant species they visited.

<i>Hummingbird Species</i>	<i>Visited plant species</i>
	<i>Besleria solanoides</i>
	<i>Burmeistera crispiloba</i>
	<i>Burmeistera multiflora</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon nigricans</i>
	<i>Centropogon solanifolius</i>
	<i>Columnea ciliata</i>
	<i>Columnea eburnea</i>
	<i>Columnea kuczyniakii</i>
	<i>Columnea sp.</i>
	<i>Cuatresia riparia</i>
	<i>Drymonia tenuis</i>
	<i>Elleanthus robustus</i>
	<i>Erythrina megistophylla</i>
	<i>Gasteranthus quitensis</i>
	<i>Glossoloma purpureum</i>
	<i>Guzmania jaramilloi</i>
	<i>Guzmania lehmanniana</i>
	<i>Guzmania rhonhofiana</i>
	<i>Guzmania xanthobractea</i>
	<i>Heliconia impudica</i>
	<i>Kohleria spicata</i>
	<i>Palicourea demissa</i>
	<i>Palicourea sodiroi</i>
	<i>Podandrogyne sp1</i>
	<i>Psammisia ulbrichiana</i>
	<i>Renealmia dolichocalyx</i>
<i>Agelaiocercus coelestis</i>	<i>Renealmia sessilifolia</i>

	<i>Bomarea pardina</i>
	<i>Burmeistera crispiloba</i>
	<i>Burmeistera multiflora</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon solanifolius</i>
	<i>Columnea ciliata</i>
	<i>Columnea eburnea</i>
	<i>Columnea kuczyniakii</i>
	<i>Columnea picta</i>
	<i>Columnea sp.</i>
	<i>Costus pulverulentus</i>
	<i>Fuchsia macrostigma</i>
	<i>Gasteranthus quitensis</i>
	<i>Glossoloma purpureum</i>
	<i>Guzmania jaramilloi</i>
	<i>Guzmania lehmanniana</i>
	<i>Guzmania rhonhofiana</i>
	<i>Heliconia impudica</i>
	<i>Heliconia sp.</i>
	<i>Heliconia virginalis</i>
	<i>Macleania smithiana</i>
	<i>Palicourea demissa</i>
	<i>Psammisia aberrans</i>
	<i>Psammisia ulbrichiana</i>
	<i>Renealmia dolichocalyx</i>
	<i>Renealmia sessilifolia</i>
<i>Phaethornis syrmatorphorus</i>	<i>Wercklea ferox</i>
	<i>Bomarea pardina</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon solanifolius</i>

	<i>Columnea ciliata</i>
	<i>Columnea eburnea</i>
	<i>Columnea kucyniakii</i>
	<i>Columnea picta</i>
	<i>Columnea sp.</i>
	<i>Costus pulverulentus</i>
	<i>Dicliptera scabra</i>
	<i>Drymonia brochidodroma</i>
	<i>Gasteranthus quitensis</i>
	<i>Glossoloma purpureum</i>
	<i>Heliconia impudica</i>
	<i>Heliconia sclerotricha</i>
	<i>Heliconia sp.</i>
	<i>Heliconia virginalis</i>
	<i>Macleania smithiana</i>
	<i>Palicourea demissa</i>
	<i>Psammisia ulbrichiana</i>
	<i>Renealmia dolichocalyx</i>
	<i>Renealmia sessilifolia</i>
	<i>Sobralia tamboana</i>
<i>Phaethornis yaruqui</i>	<i>Wercklea ferox</i>
	<i>Burmeistera crispiloba</i>
	<i>Burmeistera multiflora</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon solanifolius</i>
	<i>Columnea ciliata</i>
	<i>Columnea eburnea</i>
	<i>Columnea picta</i>
	<i>Columnea sp.</i>
	<i>Costus pulverulentus</i>

	<i>Dicliptera scabra</i>
	<i>Gasteranthus quitensis</i>
	<i>Glossoloma purpureum</i>
	<i>Guzmania rhonhofiana</i>
	<i>Heliconia impudica</i>
	<i>Heliconia sclerotricha</i>
	<i>Justicia secunda</i>
	<i>Kohleria villosa</i>
	<i>Macleania smithiana</i>
	<i>Palicourea demissa</i>
	<i>Psammisia cordifolia</i>
	<i>Psammisia sodiroi</i>
	<i>Renealmia dolichocalyx</i>
<i>Phaethornis striigularis</i>	<i>Renealmia sessilifolia</i>
	<i>Bomarea pardina</i>
	<i>Burmeistera crispiloba</i>
	<i>Burmeistera multiflora</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon solanifolius</i>
	<i>Columnea ciliata</i>
	<i>Columnea eburnea</i>
	<i>Columnea kucyniakii</i>
	<i>Costus pulverulentus</i>
	<i>Gasteranthus quitensis</i>
	<i>Glossoloma purpureum</i>
	<i>Guzmania jaramilloi</i>
	<i>Guzmania lehmanniana</i>
	<i>Guzmania xanthobractea</i>
	<i>Heliconia sp.</i>
	<i>Palicourea sodiroi</i>

	<i>Podandrogyne sp1</i>
	<i>Psammisia aberrans</i>
	<i>Psammisia sodiroi</i>
	<i>Psammisia ulbrichiana</i>
<i>Coeligena wilsoni</i>	<i>Renealmia sessilifolia</i>
	<i>Besleria solanoides</i>
	<i>Burmeistera multiflora</i>
	<i>Cavendishia grandifolia</i>
	<i>Centropogon solanifolius</i>
	<i>Columnea kuczyniakii</i>
	<i>Costus pulverulentus</i>
	<i>Drymonia tenuis</i>
	<i>Elleanthus robustus</i>
	<i>Erythrina megistophylla</i>
	<i>Gasteranthus quitensis</i>
	<i>Guzmania jaramilloi</i>
	<i>Guzmania rhonhofiana</i>
	<i>Heliconia impudica</i>
	<i>Kohleria spicata</i>
	<i>Macleania smithiana</i>
	<i>Microchilus sp.</i>
	<i>Palicourea demissa</i>
	<i>Palicourea sodiroi</i>
	<i>Psammisia sodiroi</i>
<i>Ocreatus underwoodii</i>	<i>Renealmia sessilifolia</i>
	<i>Cavendishia grandifolia</i>
	<i>Columnea kuczyniakii</i>
	<i>Guzmania jaramilloi</i>
	<i>Heliconia impudica</i>
	<i>Palicourea sodiroi</i>

	<i>Psammisia sodiroi</i>
	<i>Psammisia ulbrichiana</i>
<i>Urosticte benjamini</i>	<i>Tillandsia cyanea</i>
	<i>Cavendishia grandifolia</i>
	<i>Erythrina megistophylla</i>
	<i>Guzmania xanthobracteata</i>
<i>Heliodoxa jacula</i>	<i>Palicourea demissa</i>
	<i>Cavendishia grandifolia</i>
	<i>Palicourea sodiroi</i>
	<i>Psammisia aberrans</i>
<i>Doryfera ludovicae</i>	<i>Psammisia ulbrichiana</i>
	<i>Gasteranthus quitensis</i>
	<i>Heliconia impudica</i>
<i>Thalurania colombica</i>	<i>Heliconia sclerotricha</i>
<i>Heliodoxa rubinoides</i>	<i>Tillandsia cyanea</i>
<i>Colibri coruscans</i>	<i>Centropogon solanifolius</i>

4. Conclusions:

- Many similar species can occur in the same place because they use different resources.
- Conservation efforts should consider not only species but interactions among species.
- Key hummingbird plants such as *Cavendishia grandifolia* and *Centropogon solanifolius* can be used in restoration in Maquipucuna. These species offer resources to more hummingbirds than the other plants where we recorded hummingbirds foraging (10 species).
- *Heliodoxa rubinoides* is the most specialized hummingbird. Species such as *Tillandsia cyanea* is key to maintaining this hummingbird in Maquipucuna.
- In Maquipucuna we recorded one new species of *Columnea sp.nov.* This species is also present in Las Gralarias, Puyucunapi and Sachatamia.
- The hummingbird *Colibri coruscans*, typical of disturbed areas, was only recorded in Maquipucuna and Santa Lucía.

- Maquipucuna does not have a clear flowering peak. However, some years more flowers are present from April to June.

Acknowledgements

We thank the European Research Council (EU grant agreement 787638), the Swiss National Science Foundation (grant No. 173342), and National Geographic Society (grant agreement 9952-16) for financial support. We are also grateful with Maquipucuna Foundation for their support with the project at the reserve. Ministry of Environment in Ecuador provided the research permit N° 016-2019-IC-FLO-FAU-DNB/MAE required to conduct field work.