

## *Gwynia capsula* (Jeffreys, 1859) and other Recent brachiopods from submarine caves in Croatia

by Eric SIMON & Gregory WILLEMS

### Abstract

Recent brachiopods were collected in submarine caves along the Croatian coast of the Adriatic Sea. In a karstic cave at U Privlaka, Lošinj Island, the micromorphic brachiopod *Gwynia capsula* (JEFFREYS, 1859) has been found for the first time in the Mediterranean Sea. In the same cave, the rhynchonellid brachiopod, *Tethyrhynchia mediterranea* LOGAN, 1994, is also abundant. The geographic distribution of these two species is now considerably extended.

**Key words:** Brachiopoda, Mediterranean, taxonomy, *Gwynia*.

### Résumé

Des brachiopodes Récents ont été collectés dans des grottes sous-marines au large des côtes croates de la Mer Adriatique. *Gwynia capsula* (JEFFREYS, 1859) a été découverte pour la première fois en Méditerranée dans une grotte située à U Privlaka, île de Lošinj. L'unique brachiopode rhynchonellide connu en Méditerranée, *Tethyrhynchia mediterranea* LOGAN, 1994, est également abondant dans cette même caverne. L'extension géographique observée pour ces deux espèces est importante.

**Mots-clefs:** Brachiopoda, Méditerranée, taxonomie, *Gwynia*.

### Introduction

Eleven Recent brachiopod species from the Mediterranean Sea were described and illustrated more than a century ago by DAVIDSON (1886-1888). A taxonomic guide to Mediterranean brachiopods, by LOGAN (1979), deals with the same eleven species. In 1994, however, a new rhynchonellid brachiopod, collected from submarine caves along the Mediterranean coasts of France and Tunisia, was described by LOGAN (*in* LOGAN & ZIBROWIUS, 1994) as *Tethyrhynchia mediterranea*, and represented the first record of a rhynchonellid in the Mediterranean Sea. This representative of a new brachiopod family, has considerable taxonomical and ecological interest. The fauna of submarine caves, particularly the micromorphic brachiopods, have not been completely investigated and attention must be paid to this special biocoenosis.

### Material and methods

Bioclasts of brachiopods were collected by one of us (G.W.) using SCUBA along the Croatian coast. Sediment samples were collected from various depths (3 - 60 m), at different stations, including several submarine caves, near Lošinj Island, northern Adriatic (Figure 1). To avoid overlooking extremely small brachiopods, the samples were passed through a 0.3 mm sieve. The material is preserved in the collections of the Institut royal des Sciences naturelles de Belgique in Brussels.

### Results

The most interesting brachiopod assemblage was collected at a depth of 35 m from a karstic, semi-dark cave situated in U Privlaka (Figure 1). Its one and only entrance is small, and its volume reaches  $\pm 30$  m<sup>3</sup>. Inside, the movement of water is reduced. The water temperature varied from 11°C in November to 13°C in July 1997. This slight difference of temperature does not favour water movement. The walls and roof of the cave are full of cavities. The surface available for brachiopod attachment is increased by the permanent activity of the rock-boring mollusc, *Lithophaga lithophaga* (L., 1758). Light attenuation leads to a presumed reduction of food resources, limiting the development of algae, bivalves and other sessile animals. Sponges and bryozoans are rare and quite small. Cirripedes, worm tubes and solitary cnidarians are more abundant. Under such conditions, micromorphic brachiopods, with low nutritional requirements, have a competitive advantage for surviving in cryptic habitats (LOGAN & ZIBROWIUS, 1994, p. 87). In the cave of U Privlaka, reduced competition for space still exists between brachiopods and two fixed bivalves which are not uncommon: *Arca tetragona* POLI, 1795 and *Modiolula phaseolina* (PHILIPPI, 1844).

A rich brachiopod fauna has been observed (Fig. 2). Six brachiopod species were collected from the U Privlaka cave: one Craniidae: *Neocrania anomala* (MÜLLER, 1776); one Tethyrhynchiidae: *Tethyrhynchia mediterranea* LOGAN, 1994; three Megathyrididae: *Argyrotheca cuneata* (RISSO, 1826), *A. cordata* (RISSO, 1826), and *A. cistellula* (SEARLES-WOOD, 1841) and one *ex* Megathyrididae:

*Gwynia capsula* (JEFFREYS, 1859), recently removed from this family by LOGAN, MACKINNON & PHORSON (1997, p. 250). The relative abundance of these six brachiopods is indicated in Figure 2.

*N. anomala* and *A. cuneata* are clearly visible, in living position, on the roof of the cave. *N. anomala* is the most common brachiopod in all caves from the Lošinj area. *A. cuneata*, although common in caves, is also observed on the drop off, attached to various small substrates. Adults of *A. cordata* are uncommon in caves, but frequent on the drop off, colonising a lot of small, hard substrates. Above 30 m, *A. cordata* is rarer. *A. cistellula* was considered as a rare species in the Mediterranean (LOGAN, 1979, p. 43) but it has probably been overlooked, because of its small size. A few specimens were found in the area of Lošinj, at the base of the drop off. However, in the U Privlaka cave, *A. cistellula* is clearly the dominant micromorphic species (Figure 2). Specimens collected are typical and their shell morphology fits the emended description and illustrations recently published by LOGAN, MACKINNON & PHORSON (1997, p. 245, pl. 3, figs. 3-10, p. 246) perfectly.

Micromorphic species as *Tethyrhynchia mediterranea* and *Gwynia capsula*, were found only collected in the semi-dark karstic cave of U Privlaka. This new discovery of *T. mediterranea* in the Adriatic extends its geographic distribution to the eastern Mediterranean basin, suggesting that this rhynchonellid brachiopod is probably distributed in the whole Mediterranean region. Its origin remains unknown but investigations in submarine caves along the Atlantic coasts could clarify this situation.

*G. capsula*, an eastern Atlantic brachiopod, is found around the coasts of the British Isles, northern France, the Netherlands, Norway, and north-western Spain (JEFFREYS, 1878; DUPREY, 1883; DAVIDSON, 1887; BOADEN, 1963; SWEDMARK, 1967, 1971; RIOULT, 1971; BRUNTON & CURRY, 1979; BESTEIRO & URGORRI, 1984; PHORSON, 1988; HARPER *et al.*, 1996; LOGAN *et al.*, 1997). Its occurrence in the northern Adriatic, and thus for the first time in the Mediterranean, considerably extends its geographical distribution. This is another argument for considering Recent Mediterranean brachiopods as immigrant species from the Atlantic, via Gibraltar, at the beginning of the Pliocene (HSÜ, 1974; HSÜ *et al.*, 1973). Other caves explored in the area of Lošinj Island yielded numerous *N. anomala* and *A. cuneata* and a few *A. cordata* and *A. cistellula*, but *T. mediterranea* and *G. capsula* were not observed. In these caves, situated at a lesser depth (from 3 to 10 m), fresh water springs may modify the local salinity.

### Systematic descriptions

Phylum Brachiopoda DUMÉRIL, 1806

Subphylum Rhynchonelliformea WILLIAMS *et al.*, 1996

Class Rhynchonellata WILLIAMS *et al.*, 1996

Superfamily Rhynchonelloidea D'ORBIGNY, 1847

Nomenclative note: the family-group names based on *Rhynchonella* should be attributed to D'ORBIGNY, 1847

and not to GRAY, 1848 (MANCENIDO, OWEN & MORRIS, 1993 p. 197).

Family Tethyrhynchiidae LOGAN, 1994

Genus *Tethyrhynchia* LOGAN, 1994

Type species: *Tethyrhynchia mediterranea* LOGAN, 1994

*Tethyrhynchia mediterranea* LOGAN, 1994

Plate 1, Figures 3 a-c, 4 a-d.

\* 1994 *Tethyrhynchia mediterranea* LOGAN *n. sp.* - LOGAN & ZIBROWIUS, pp. 77-88, plate 1, figs. 1-14, plate 2, figs. 1-9.

• 1997 *Tethyrhynchia mediterranea* LOGAN - LOGAN, MACKINNON & PHORSON, p. 240.

Material: 128 complete shells, 66 dorsal valves and 72 ventral valves collected from a submarine cave at U Privlaka (-35 m).

The specimens are extremely small, their size rarely exceeding 1 mm (largest individual: 1.40 mm long and 1.15 mm wide). Externally, the shell is smooth. The beak is slightly curved with an elongate-triangular, hypothyrid foramen limited by two small, disjunct, deltidial plates. In the ventral valve, an incomplete pedicle collar is observed and the teeth are relatively strong. Dental plates are absent. The dorsal valve has no cardinal process. Long and typical crura of the lunifer type support a trocholophous lophophore. These Croatian specimens fit perfectly with the original description given by LOGAN (1994).




Fig. 1. – Map illustrating the area of Lošinj Island (northern Adriatic, Croatia) and a detail of Lošinj Island with situation of the submarine cave at U Privlaka.

Order Terebratulida WAAGEN, 1883  
 Suborder Terebratellidina MUIR-WOOD, 1955  
 Superfamily Uncertain  
 Family Uncertain

Nomenclative note:

*Gwynia capsula* has been removed from the family Megathyrididae by LOGAN, MACKINNON & PHORSON (1997, pp. 249-250). These authors drew attention to similarities between *Gwynia* and the Jurassic genus *Zellania* MOORE, 1855 (Family uncertain), which exhibits several features similar to some characters observed in *Gwynia capsula* (see also BEECHER, 1893, pl. 1). The presence of submarginal ridges in the dorsal valve is certainly the most striking character which they have in common.

Genus *Gwynia* KING, 1859

Type species: *Terebratula capsula* JEFFREYS, 1859

***Gwynia capsula* (JEFFREYS, 1859)**

Plate 1, Figures 1 a-e, 2 a-e.

- \* 1859 *Terebratula capsula* n. s. - JEFFREYS, p. 43, pl. 2, fig. 7a-b.
- . 1859 *Gwynia capsula* - KING, 1, p. 258, figs. 1-5.
- . 1861 *Gwynia* - DAVIDSON, p. 28.
- . 1861 Subgenus? uncertain (*Gwynia*, King). *Terebratula ? capsula*, Jeffreys. - DAVIDSON, p. 39.
- . 1863 *Argiope capsula*, Jeffreys. - JEFFREYS, p. 21.
- . 1863 *Argiope Neapolitana* (probably *T. cordata* of Risso) - JEFFREYS, p. 22.
- . 1865 *Gwynia capsula* - SARS, pp. 96-97, 127, 134, pl. 4, figs. 121-122.
- . 1874 *Gwynia capsula*, Gwyn Jeffreys, sp. - DAVIDSON, 4, p. 5, pl. I, figs. 3-4.
- . 1877 *Gwynia (Argiope) capsula* Jeff. - FRIELE, 2(4), pp. 385-386, pl. 3, fig. 7b.
- . 1878 *Argiope capsula*, Jeffreys [placed in subgenus *Gwynia*] - JEFFREYS, p. 410-411.
- . 1880 *Gwynia (Terebratula) capsula*, Jeff. - DAVIDSON, p. 14.
- . 1883 *Argiope capsula*, Jeffr. - DUPREY, pp. 185-186.
- . 1887 *Gwynia capsula*, Gwyn Jeffreys, sp. - DAVIDSON, pp. 150-152, pl. 21, figs. 28a-c, 29.
- non 1891 *Gwynia capsula*, Gwyn Jeffreys, sp. - FISCHER & OEHLERT, pp. 125-128, fig. 12a-f.
- pp. 1893 *Gwynia capsula* - BEECHER, pp. 377, 382-384, 386, 387, 391, pl. 1, figs. Aa, non fig. A.
- . 1920 *Gwynia capsula* - DALL, pp. 324-325.
- . 1925 *Gwynia capsula* (Jeff.) - MASSY, pp. 38-39, 42.
- pp. 1927 *Gwynia capsula* - THOMSON, pp. 207-208, non fig. 61a-f, p. 208 (= "*Gwynia capsula*" sensu FISCHER & OEHLERT, 1891).
- . 1933 *Gwynia capsula* (Gwyn Jeffreys 1859) - CORI, p. 148, text-fig. 19.
- . 1933 *Gwynia capsula* (JEFFREYS) - STIASNY, p. 138.
- . 1959 *Gwynia* - HYMAN, 5(2), p. 1417.
- . 1960 *Gwynia* - BEAUCHAMP, pp. 1406, 1417.
- . 1963 *Gwynia capsula* Jeffries (*sic*) - BOADEN, p. 91.
- pp. 1965 *Gwynia capsula* (JEFFREYS, 1859) - HATAI & ELLIOTT in MOORE, p. H832, non fig. 715,1a-e, p. H831 (= "*Gwynia capsula*" sensu FISCHER & OEHLERT, 1891).
- . 1967 *Gwynia capsula* (Jeffreys) - SWEDMARK, pp. 1151-1152, fig. 1, p. 1151, fig. 2 (larva), p. 1152.
- . 1971 *Gwynia capsula* Jeffreys - SWEDMARK, fig. 1-E, p. 43, p. 44.
- . 1971 *Gwynia capsula* (Jeffreys) - RIOULT, pp. 26, 28.
- . 1971 *Gwynia capsula* (Jeffreys) - HULINGS & GRAY, p. 43.
- non 1975 *Gwynia capsula* Jeffreys, 1859 - ZEZINA, p. 193.
- . 1976 *Gwynia capsula* Jeffreys, 1859 - ZEZINA, table 3, p. 17, pp. 21, 23, 111.
- . 1978 *Gwynia capsula* (Jeffreys) - WILLIAMS & MACKAY, pp. 192, 193, 199, 200, pl. 4, figs. 17-19, pl. 5, fig. 20.
- . 1979 *Gwynia* - WILLIAMS & MACKAY, p. 734.
- . 1979 *Gwynia capsula* (Jeffreys) - BRUNTON & CURRY, pp. 21, 24, 46, fig. 23, p. 47.
- non 1983 *Gwynia capsula* - LOGAN, table 1, p. 169 (= "*Gwynia capsula*" sensu FISCHER & OEHLERT, 1891).
- . 1984 *Gwynia capsula* (Jeffreys, 1859) - BESTEIRO & URGORRI, pp. 219-224, pl. 1, figs. 5-10.
- . 1985 *Gwynia capsula* (Jeffreys, 1859) - ZEZINA, p. 193.
- . 1988 *Gwynia capsula* (Jeffreys) - NIELSEN, p. 445, fig. 50.1, p. 446.
- . 1988 *Gwynia capsula* (Jeffreys) - PHORSON, pp. 32-34, fig. 2, p. 33.
- non 1988 *Gwynia capsula* - LOGAN, table 1, p. 63 (= "*Gwynia capsula*" sensu FISCHER & OEHLERT, 1891).
- . 1989 *Gwynia capsula* (Jeffreys, 1859) - SAIZ SALINAS, p. 146.
- pp. 1993 *Gwynia capsula* - LOGAN, table 3, p. 232 (non "*Gwynia capsula*" sensu FISCHER & OEHLERT, 1891).
- . 1996 *Gwynia capsula* (JEFFREYS, 1859) - HARPER, DONOVAN & VELTKAMP, pp. 331-333, fig. 1(1-4), p. 332.
- . 1997 *Gwynia capsula* (JEFFREYS, 1859) - LOGAN, MACKINNON & PHORSON, pp. 239-252, fig. 1, p. 242, fig. 2 (1-14), p. 244, fig. 3 (1-2), p. 246, fig. 4 (1-4), p. 248.

Material: 90 complete shells, 47 dorsal valves and 18 ventral valves collected from a submarine cave at a depth of 35 m in U Privlaka, Lošinj Island, Croatia. Externally, specimens from U Privlaka have a biconvex, subcircular to slightly elongate shell, with a smooth surface except for faint growth lines. Some dissymmetrical shells are observed. The dorsal valve is a little larger than the ventral valve, the dorsal umbo being slightly longer than the ventral one. This character is unusual in articulate brachiopods and probably induced some misidentifications of brachial and ventral valves (HARPER *et al.*, 1996, p. 332, fig. 1). The amphithyrid foramen is large and rudimentary deltidial plates are present. The shell is endopunctate. A short pedicle collar is hardly visible in the ventral valve, which possesses small, pointed teeth, without dental plates. The dorsal valve has strong inner and outer socket ridges but is devoid of a cardinal process, hinge plates and septal pillar. The development, in the dorsal valve, of two postero-lateral submarginal ridges, arising from the valve floor, is a typical feature. These ridges

support a trochlophous lophophore (LOGAN, MACKINNON & PHORSON, 1997, p. 243).

This extremely small-sized brachiopod (largest individual found in U Privlaka: 1.02 mm. long and 0.84 mm. wide), living in a cryptic habitat, was probably overlooked in the western part of the Mediterranean. The Croatian population of *G. capsula* suggests a larger development for this species in the whole Mediterranean Sea.

Although *G. capsula* was described from Northern Ireland since 1859, very little was known about this species. FISCHER & OEHLERT (1891, p. 126, fig. 12a-f) reported the presence of "*Gwynia capsula*" at a great depth (822 and 4060 m) in the material collected by the Talisman (see also ZEZINA, 1975, pp. 193-194). They described and illustrated this specimen which, obviously, is not similar to the real *Gwynia capsula* (JEFFREYS, 1859). The beak of the ventral valve is clearly longer and pointed. Its triangular foramen is hypothyriddid. ZEZINA (1976, p. 111) already pointed out the confusion in FISCHER & OEHLERT that produced further inaccurate representations for this species (HATAI, 1965 p. H831, fig. 1a-f and THOMSON, 1927 p. 208, fig. 61 a-f). The first photograph of *G. capsula* was surprisingly published only three years ago (HARPER *et al.*, 1996, pp. 331-332, figs. 1-1 & 1-2) but the dorsal and the ventral valve were confused. A recent reappraisal of the morphology, distribution, life habits and phylogenetic affinities of *G. capsula* establishes the taxonomic status and definitive characters of this brachiopod (LOGAN *et al.*, 1997, pp. 239-252).




Fig. 2. – Relative abundance (as a % of the total number of individuals) of the different brachiopod species collected in the sediment of a submarine cave at U Privlaka, Lošinj Island, Croatia. N: total number of individuals found in 3 litres of dried bioclasts.

## Conclusion

A study of a collection of Croatian brachiopods identified one new species for the Mediterranean and two new species for the Adriatic area. At shallow depths, caves offer ecological conditions similar to those observed in the bathyal zone. But they also represent specific environments for the development of specialised shallow water micro-brachiopods such as *Tethyrhynchia mediterranea*, *Argyrotheca cistellula* and *Gwynia capsula*.

In Croatia, *G. capsula* has not been found outside caves or in cavities situated on the drop off. If the original Atlantic ecological conditions of *G. capsula* are taken into account, Mediterranean karstic caves are ideal refuges for this species, which probably does not prefer higher water temperatures or strong fluctuations in water temperature. A further development of this study will describe more precisely on the living conditions of these micro-morphic brachiopods.

## Acknowledgments

We thank A. LOGAN (Saint John, Canada) and D.I. MACKINNON (Christchurch, New Zealand) for advice. I thank Annie DHONDT (Brussels, Belgium) for reviewing the manuscript. Julien CILLIS (Brussels) is gratefully acknowledged for the photographs taken with the Scanning Electron Microscope.

## References

- DE BEAUCHAMP, P., 1960. Classe des Brachiopodes in P.P. GRASSÉ (Ed.). *Traité de Zoologie*, 5, 1380-1430.
- BEECHER, C.E., 1893. Revision of the Families of Loop-bearing Brachiopoda. *Transactions of the Connecticut Academy of Arts and Sciences*, 9, 376-399.
- BESTEIRO, C. & URGORRI, V., 1984. Sobre la presencia de *Argyrotheca cistellula* (SEARLES-WOOD, 1841), *Gwynia capsula* (JEFFREYS, 1859) (Brachiopoda, Terebratulida) y *Leptosynapta minuta* (BECHER, 1906) (Holothuroidea, Apoda) en las costas Gallegas. *Actas do IV Simposio Iberica de Estudos do Benthos Marinho*, 3, 219-226.
- BOADEN, P.J.S., 1963. The interstitial fauna of some North Wales beaches. *Journal of the Marine Biological Association of the United Kingdom*, 43, 79-96.
- BRUNTON, C.H.C. & CURRY, G.B., 1979. British Brachiopods. *Synopses of the British Fauna (New Series)*, 17, 1-64.
- CORI, C.I., 1933. Brachiopoda in G. GRIMPE & N. WAGLER (Eds.). *Die Tierwelt der Nord- und Ostsee*, 7, 133-150.
- DALL, W.H., 1920. Annotated list of the Recent Brachiopoda in the collection of the United States National Museum, with descriptions of thirty-three new forms. *Proceedings of the United States National Museum*, 57, 261-377.
- DAVIDSON, T. A., 1861. On Recent Terebratulidae. *The Annals and Magazine of Natural History*, Ser. 3, 3, 24-40.

- DAVIDSON, T. A., 1874. A monograph of British Fossil Brachiopoda, Supplement to the Recent, Tertiary, and Cretaceous species. *Palaeontographical Society (London) Monograph*, 4, part 1: 1-72.
- DAVIDSON, T. A., 1880. Report on the Brachiopoda dredged by H.M.S. Challenger during the years 1873-1876 in Report on the scientific results of the voyage of H.M.S. Challenger, during the years 1873-1876, under command of Captain George S. Nares and Captain Frank Tourie Thompson. Prepared under the superintendence of Sir C. Wyville Thompson (and of John Murray). *Challenger Zoological Reports*, 1, 1-67.
- DAVIDSON, T. A., 1886-1888. A Monograph of Recent Brachiopoda. *The Transactions of the Linnean Society of London*, Ser.2, 4, 3 parts, 248 pp.
- DUMÉRIL, A.M.C., 1806. Zoologie analytique ou méthode naturelle de classification des animaux. Allais, Paris, XXIV + 344 pp.
- DUPREY, E., 1883. Shells of the Littoral Zone in Jersey. *The Annals and Magazine of Natural History*, Ser. 5, 9, 185-190.
- FISCHER, P. & OEHLERT, D.P., 1891. Expéditions scientifiques du TRAVAILLEUR et du TALISMAN pendant les années 1880-1883. Brachiopodes. G. Masson Ed., Paris, 139 pp.
- FRIELE, H., 1877. The development of the skeleton in the genus *Waldheimia*. *Archiv for Mathematik og Naturvidenskab*, 2(4): 381-386.
- GRAY, J.E., 1848. On the arrangement of the Brachiopoda. *The Annals and Magazine of Natural History*, (2) 2(12): 435-440.
- HARPER, D.A.T, DONOVAN, S.K. & VELTKAMP, C.J., 1996. The micromorphic articulate brachiopod *Gwynia* from the Western Approaches, U.K. *Journal of Paleontology*, 70, 331-333.
- HATAI, K. in HATAI, K. & ELLIOTT, G.F., 1965. Family Megathyrididae DALL, 1870 in MOORE, R.C. (Ed.), Treatise on Invertebrate Paleontology, Part H, Brachiopoda. Geological Society of America and University of Kansas Press. Lawrence, Kansas: 830-832.
- HSÜ, K.J., 1974. The Miocene dessication of the Mediterranean and its climatological and zoogeographical implications. *Naturwissenschaften*, 61(4): 137-142.
- HSÜ, K.J., RYAN, W.B.F. & CITA, M.B., 1973. Late Miocene dessication of the Mediterranean. *Nature*, 242, n°5395, 240-244.
- HULINGS, N.C. & GRAY, J.S., 1971. A Manual for the Study of Meiofauna. *Smithsonian Contributions to Zoology*, 78, 1-83.
- HYMAN, L.H., 1959. The Invertebrates. Smaller Coelomate Groups, 5, McGraw-Hill book Company, New-York, London, Toronto, 546 pp.
- JEFFREYS, J.G., 1859. Further gleanings in British conchology. *The Annals and Magazine of Natural History*, 3(3): 30-43.
- JEFFREYS, J.G., 1863. British Conchology or an account of the Mollusca which now inhabit the British Isles and the surrounding seas, 2, Class Brachiopoda. John Van Voorst, London. 1-26.
- JEFFREYS, J.G., 1878. On the Mollusca procured during the "Lightning" and "Porcupine" Expeditions, 1868-1870. Part I. *Proceedings of the scientific meetings of the Zoological Society of London*, 46(1): 393-416.
- KING, W., 1859. On *Gwynia*, *Dielasma* and *Macandrevia*, three new genera of palliobranchiate Mollusca, one of which has been dredged in the Strangford Lough. *Proceedings of the Dublin University Zoological and Botanical Association*, 1, 256-262.
- VON LINNÉ, C., 1758. *Systema naturae*, 10th. ed., Stockholm, 823 pp.
- LOGAN, A., 1979. The Recent Brachiopoda of the Mediterranean Sea. *Bulletin de l'Institut Océanographique de Monaco*, 72, n°1434, 1-112.
- LOGAN, A., 1983. Brachiopoda collected by CANCAP I-III Expeditions to the South-East North Atlantic. 1976-1978. *Zoologische Mededelingen*, 57(18): 165-189.
- LOGAN, A., 1988. Brachiopoda collected by CANCAP IV and VI Expeditions to the South-East North Atlantic. 1980-1982. *Zoologische Mededelingen*, 62(5): 59-74.
- LOGAN, A., 1993. Recent Brachiopods from the Canarian-Cape Verdean Region: Diversity, Biogeographic Affinities, Bathymetric Range and Life Habits. *Courier Forschungs - Institut Senckenberg*, 159, 229-233.
- LOGAN, A., MACKINNON, D.I. & PHORSON, J.E., 1997. Morphology, Distribution, Life Habits and Phylogenetic Affinities of the Recent Brachiopod *Gwynia capsula* (JEFFREYS). *Pubblicazioni della Stazione Zoologica di Napoli: Marine Ecology*, 18(3): 239-252.
- LOGAN, A. & ZIBROWIUS, H., 1994. A new Genus and Species of Rhynchonellid (Brachiopoda, Recent) from Submarine Caves in the Mediterranean Sea. *Pubblicazioni della Stazione Zoologica di Napoli: Marine Ecology*, 15(1): 77-88 (1994).
- MANCEÑIDO, M.O., OWEN, E.F. & MORRIS, N.J., 1993. The dating of certain of d'Orbigny's brachiopod and bivalve nominal taxa. *Bulletin of Zoological Nomenclature*, 50(3): 196-199.
- MASSY, A.L., 1925. The Brachiopoda of the coasts of Ireland. *Proceedings of the Royal Irish Academy*, B, 37, 37-46.
- MOORE, C., 1855. On new Brachiopoda, from the Inferior Oolithe of Dundry. *Proceedings of the Archaeological and Natural History Society*, 5, 107-128.
- MUIR-WOOD, H., 1955. A history of the classification of the phylum Brachiopoda. British Museum (Natural history), London, 124 pp.
- MÜLLER, O.F., 1776. *Zoologiae Danicae prodromus seu animalium Daniae et Norvegiae indigenarum characteres, nomina, et synonyma imprimis popularium*. Havniae (Copenhagen), XXXII -282 pp.
- NIELSEN, C., 1988. Brachiopoda in R.P. HIGGINS and H. THIEL (eds.). Introduction to the Study of the Meiofauna. Smithsonian Institution Press, Washington, pp. 445-446.
- D'ORBIGNY, A., 1847. Considérations zoologiques et géologiques sur les Brachiopodes ou Palliobranches (parts 1-2). *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences*, 25(5): 193-195; 25(7): 266-269.
- PHILIPPI, R.A., 1844. *Enumeratio Molluscorum Siciliae*, 2, Eduardi Anton, Halle, 303 pp.
- PHORSON, J.E., 1988. Small Brachiopods. *Porcupine Newsletter*, 4(2): 32-34.
- POLI, I.X., 1795. *Testacea utriusque Siciliae*, 2, Regio Typographeio, Parmae, 264 pp.

- RIOULT, M., 1971. *Argyrotheca cistellula* (SEARLES-WOOD) (Brachiopoda, Terebratulacea) au large des côtes du Calvados. *Bulletin de la Société linnéenne de Normandie*, **102**, 26-28.
- RISSO, A., 1826. Histoire naturelle des principales productions de l'Europe méridionale et particulièrement de celles des environs de Nice et des Alpes maritimes, **4**, F.G. Levrault, Paris, VII - 439 pp.
- SAIZ SALINAS, J.I., 1989. Verzeichnis der rezenten Brachiopoden (Brachiopoda) von den iberischen Küsten und den angrenzenden Meeren. *Bonner Zoologische Beiträge*, **40**(2): 141-154.
- SARS, M., 1865. Om de I Norge forekommende fossile Dyrlevninger frå Quartaerperioden. Brøgger & Christie's, Christiania, 134 pp.
- SEARLES-WOOD, V., 1841. Mr. V. Searles-Wood's catalogue of the Crag mollusca. *The Annals and Magazine of Natural History*, **6**, 243-262.
- STIASNY, G., 1933. Verzeichnis der Brachiopoden-Sammlung des naturhistorischen Reichsmuseums in Leiden. *Zoologische Mededeelingen*, **15**(3-4): 129-147.
- SWEDMARK, B., 1964. The interstitial fauna of marine sand. *Biological Reviews*, **39**, 1-42.
- SWEDMARK, B., 1967. *Gwynia capsula* (Jeffreys), an articulate brachiopod with brood protection. *Nature*, **213**, 1151-1152.
- SWEDMARK, B., 1971. A review of gastropoda, Brachiopoda and Echinodermata in Marine Meiobenthos. *Smithsonian Contributions to Zoology*, **76**, 41-45.
- THOMSON, J. A., 1927. Brachiopod morphology and genera (Recent and Tertiary). New Zealand Board of Science and Art, Wellington, N.Z., Manual n° 7, 338 pp.
- WAAGEN, W.H., 1882-1885. Salt Range Fossils, Part 4(2), Brachiopoda. *Memoirs of the Geological Survey of Palaeontologica Indica*, **13**(1): 391-546.
- WILLIAMS, A., CARLSON, S.J., BRUNTON, C.H.C., HOLMER, L.E. & POPOV, L., 1996. A supra-ordinal classification of the Brachiopoda. *Philosophical Transactions of the Royal Society of London*, B, **351**(4): 1171-1193.
- WILLIAMS, A. & MACKAY, S., 1978. Secretion and ultrastructure of the periostracum of some terebratulide brachiopods. *Proceedings of the Royal Society of London*, B, **202**, 191-209.
- WILLIAMS, A. & MACKAY, S., 1979. Differentiation of the brachiopod periostracum. *Palaeontology (London)*, **22**, 721-736.
- ZEZINA, O.N., 1975. Recent Caribbean Deep-Sea Brachiopod Fauna, the Sources and Conditions of its Formation. *Akademia Nauk SSSR, Okeanologii Institut*, **100**, 188-195. [in Russian].
- ZEZINA, O.N., 1976. Ecology and Distribution of the Recent Brachiopods. *Akademia Nauk SSSR, Nauka, Moscow*, 138 pp. [in Russian].
- ZEZINA, O.N., 1985. Recent brachiopods and Problems of the Bathyal Oceanic Zone. *Nauka, Moscow*, 248 pp. [in Russian].

Eric SIMON.

Département de Paléontologie (Invertébrés),  
Institut royal des Sciences naturelles de Belgique,  
Rue Vautier, 29,  
B-1000 Bruxelles,  
Belgique.

Gregory WILLEMS.

Chaussée de la Gare, 9,  
B-5080 Rhisnes,  
Belgique.

Correspondence and requests for materials should be  
addressed to  
E. S. (e-mail: ericsimon98brach@yahoo.fr).

#### Plate 1

*Gwynia capsula* (JEFFREYS, 1859) from a semi-dark, submarine cave (depth: -35 m) in U Privlaka, Lošinj Island, Croatia.

Fig. 1 - Complete adult specimen (x47). 1a- ventral view, 1b- dorsal view, 1c- lateral view, 1d- posterior view and 1e- anterior view (Specimen IRSNB-IST N°6392).

Fig. 2 - Another complete, opened specimen (IRSNB-IST N°6393). 2a- dorsal valve in ventral view (x47), 2b- ventral valve in dorsal view (x47), 2c- dorsal valve in posterior view showing the postero-lateral submarginal ridges (x80), 2d- dorsal valve, a detail of sockets and socket ridges (x 128), and 2e- ventral valve, a detail of the teeth (x128).

*Tethyrhynchia mediterranea* LOGAN, 1994 from a semi-dark, submarine cave (depth: -35 m) in U Privlaka, Lošinj Island, Croatia.

Fig. 3 - Complete adult specimen (IRSNB-IST n° 6390). 3a- dorsal view (x35), 3b- ventral view (x32), 3c- lateral view (x35).

Fig. 4 - Another complete, opened specimen (IRSNB-IST n°6391). 4a- ventral valve in dorsal view (x35), 4b- dorsal valve in ventral view (x35), 4c- same ventral valve, detail of teeth and pedicle collar (x96) and 4d- same dorsal valve, detail of sockets, socket ridges and lunifer crura (x96).

