

On a collection of spiders (Arachnida: Araneae) from the island of Seili, with a new record for the Finnish fauna and a list of species from the island

Authors: Zamani, Alireza, Österman, Emil M., and Koponen, Seppo

Source: Arachnologische Mitteilungen: Arachnology Letters, 64(1) : 29-33

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit6404>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

On a collection of spiders (Arachnida: Araneae) from the island of Seili, with a new record for the Finnish fauna and a list of species from the island

Alireza Zamani, Emil M. Österman & Seppo Koponen



doi: 10.30963/aramit6404

Abstract. The results of a recent collection effort on the island of Seili (located in the Archipelago Sea, off the southwest coast of Finland in the Baltic Sea, Swedish name Själö) are reported herein. Out of 709 specimens collected at two sites using four different collection methods, 173 were adults (26 males, 147 females) and identifiable to species-level, resulting in 33 species from 13 families, including five species newly reported from the island. A male of *Cheiracanthium punctorum* (Villers, 1789), separately collected at a different place, represents the first record of this species in Finland and its northernmost record across its whole known range. An updated list of spider species reported from Seili (215 species from 25 families) is included as an appendix.

Keywords: Archipelago Sea, Baltic Sea, *Cheiracanthium punctorum*, Finland, list of species, Själö

Zusammenfassung. Spinnen (Arachnida: Araneae) von der Insel Seili, mit einem Neunachweis für Finnland und einer Artenliste für die Insel. Die Ergebnisse einer kürzlich durchgeföhrten Sammelaktion auf der Insel Seili (im Schärenmeer südwestlich der finnischen Südostküste in der Ostsee gelegen, schwedischer Name Själö) werden vorgestellt. Von 709 gesammelten Exemplaren von zwei Orten, gesammelt mit verschiedenen Methoden, waren 173 adult (26 Männchen, 147 Weibchen) und auf Artniveau bestimmbar. 33 Arten aus 13 Familien, darunter fünf Erstnachweise für die Insel, wurden erfasst. Ein Männchen von *Cheiracanthium punctorum* (Villers, 1789), gesondert an einer anderen Stelle gesammelt, stellt den Erstnachweis für Finnland und den nördlichsten Nachweis der Art insgesamt dar. Eine aktualisierte Artenliste der Spinnen von Seili (215 Arten aus 25 Familien) wird in einem Appendix präsentiert.

Abstrakti. Hämähäkkikeräys (Arachnida: Araneae) Seilin saarelta, mukaan lukien uusi löytö Suomen faunalle ja saaren lajiluettelo. Saaristomerellä sijaitsevan Seilin (Själö) saarella hiljattain tehdyn hämähäkkikeräyksen tulokset esitetään tässä. Kahdelta tutkimuspaikalta kerättiin neljällä eri menetelmällä 709 hämähäkkikysilöä, joista 173 aikuista (26 koirasta ja 147 naarasta) voitiin määrittää lajittolle. Tuloksena oli 33 lajia 13 heimosta; näistä viisi lajia on uusia Seilin saarelle. Erikseen saarelta kerätty lajin *Cheiracanthium punctorum* (Villers, 1789) koirasyksilöön ensimmäinen havainto lajista Suomesta ja pohjoisin löytö koko leveinnejysalueelta. Julkaisun liitteenä on Seilin saarelta löydettyjen hämähäkkilajien luettelo (215 lajia 25 heimosta).

The island of Seili (Swedish name Själö) is a small Finnish island (ca. 1.6 km² in area) situated in the middle zone of the Archipelago Sea (part of the Baltic Sea) and belongs to the hemiboreal (“oak forest”) area of Finland. Since 1964, the Archipelago Research Institute of the University of Turku has been active here. Although its main research theme is long-term monitoring and modelling of the marine environment, it also provides a base for terrestrial research and teaching.

Although the spider fauna of Seili has not been thoroughly investigated, data from scattered studies – of which research on cliff spiders by Häkkilä (1986) is worth mentioning – and from various collecting efforts is accessible from the spider database of the Zoological Museum of the University of Turku. Miettinen (1997) compiled a report on the wildlife and natural properties of the island of Seili in which he provided a list of spider species, mainly on the basis of data from the above-mentioned work by Häkkilä (1986) and museum collections. The known Seili spider fauna includes some rare and relatively interesting species, including the wasp spider *Argiope bruennichi* (Scopoli, 1772), the rare linyphiid *Abisko abiskoensis* (Holm, 1945) and a few species included in the Red List of Finnish spiders, namely *Aulonia albimana* (Walckenaer, 1805), *Brommella falcigera* (Balogh, 1935), *Jacksonella falconeri* (Jackson, 1908), *Micaria formicaria* (Sundevall, 1831), *Titanoeca spominima* (Taczanowski, 1866) and *Zelotes exiguus* (Müller & Schenkel, 1895) (Pajunen et al. 2019, also see ‘Appendix’).

In this paper, we present the results of a recent collection effort in the island of Seili in 2022, report a species new to the Finnish fauna and provide an updated list of spider species known from Seili, including five species new to the fauna of the island.

Material and methods

Spiders, and other invertebrates, were collected at two locations on the island of Seili during 18.–20. Jul. and 25.–27. Jul. 2022, as a part of the Terrestrial Invertebrate Ecology field course held at the University of Turku. The first location was a one-hectare area in a heath forest (60.2428°N, 21.9592°E; 10–15 m a.s.l.), in which the principal vascular plant species were blueberry (*Vaccinium myrtillus*), pine (*Pinus sylvestris*) and spruce (*Picea abies*). The second location was a one-hectare area in a broadleaf woodland (60.2347°N, 21.9664°E; 10–15 m a.s.l.), in which the principal vascular plant species were common hazel (*Corylus avellana*), small-leaved linden (*Tilia cordata*) and wood bluegrass (*Poa nemoralis*).

Apart from a single hand-collected specimen, the rest of the material was collected using the methods detailed below:

- 1) Heath forest (18.–20. Jul.): Six sets of ten sweeps of the herbaceous layer with sweeping nets, five sets of shaking ten branches inside the sweeping net, eight sets of sieving detritus, five pitfall traps.
- 2) Broadleaf woodland (18.–20. Jul.): Six sets of shaking ten branches inside the sweeping net, seven sets of sieving detritus, five pitfall traps.
- 3) Heath forest (25.–27. Jul.): Six sets of ten sweeps of the herbaceous layer with sweeping nets, six sets of shaking ten branches inside the sweeping net, four sets of sieving detritus, six pitfall traps.

Alireza ZAMANI, Emil M. ÖSTERMAN, Seppo KOPONEN, Zoological Museum, Biodiversity Unit, FI-20014 University of Turku, Finland;
E-mails: alireza.zamani@utu.fi, emil.m.osterman@utu.fi, sepkopo@utu.fi

Academic Editor: Petr Dolejš

submitted 26.8.2022, accepted 3.12.2022, online 28.12.2022

- 4) Broadleaf woodland (25.–27. Jul.): Six sets of shaking ten branches inside the sweeping net, four sets of sieving detritus, five pitfall traps.

The collecting was systematically randomized within the area to cover it more evenly. Sweeping nets were medium-sized standard nets. One branch was shaken per tree (pine or spruce in the heath forest, common hazel or small-leaved linden in the broadleaf woodland). A square of 0.25 m² was used for each set of sieving and the detritus was sieved three times through a medium-sized sieve. The pitfall traps were planted on the starting date and removed on ending date; each cup, with an upper diameter of 6.5 cm, was accompanied with a rain cover and filled with 70% denatured alcohol. All of the specimens were preserved in 70% denatured alcohol and deposited in the Zoological Museum of the University of Turku (ZMUT). The specimens were identified by the second author, using the keys and illustrations provided in Nentwig et al. (2022).

The specimen of *Cheiracanthium punctorum* was photographed using a Canon EOS 7D camera attached to an Olympus SZX16 stereomicroscope. Digital images were montaged using CombineZP and edited using Photoshop.

Results

A total 709 specimens were collected at the two aforementioned sites (Tab. 1), out of which 173 were adults (26 males, 147 females) and identifiable to the species-level, resulting in 33 species from 13 families (Tab. 2).

Additionally, a single specimen was collected separately and at a different location from those listed above, and represents a new record for the Finnish fauna:

Tab. 1: The total numbers of specimens collected using each method;
A = Heath forest (18.–20. Jul. 2022); **B** = Broadleaf woodland (18.–20. Jul. 2022); **C** = Heath forest (25.–27. Jul. 2022); **D** = Broadleaf woodland (25.–27. Jul. 2022)

	A	B	C	D
Pitfall	3	9	–	29
Sieving	313	25	52	58
Shaking	56	18	99	19
Sweeping	14	not conducted	23	not conducted

Cheiracanthiidae

Cheiracanthium punctorum (Villers, 1789) (Fig. 1a-d)

Identification. Sterghiu (1985), Almquist (2006)

Material examined. FINLAND, Varsinais-Suomi: 1 ♂ (ZMUT), Parainen, Seili, 60.2393°N, 21.9608°E, 10–15 m a.s.l., 16.–17. Jul. 2022 (leg. K. Ruohomäki, det. E. M. Österman, conf. A. Zamani).

Description. See Almquist (2006).

Collection method. This specimen was collected as a by-catch from one of several light traps for moths that had been set up for approximately 24 h. The light traps were located within one hectare in which all but one were located in a grassy heath forest, with the remaining light trap located on a nearby rocky meadow.

Distribution. Western Palaearctic (World Spider Catalog 2022). In northern Europe it has been recorded from Sweden (the island of Öland) and the Baltic states (Almquist 2006, Biteniektyte & Relys 2011, Cera 2018). The current material

Tab. 2: List of species recorded at the two collection sites (heath forest and broadleaf woodland) on the island of Seili. Species new to the fauna of the island are marked with an asterisk

Taxa	Heath forest	Broadleaf woodland
Anypheidae		
<i>Anyphaena accentuata</i> (Walckenaer, 1802)	2 ♀♀	3 ♂♂, 6 ♀♀
Araneidae		
<i>Cyclosa conica</i> (Pallas, 1772)*	4 ♀♀	–
Clubionidae		
<i>Clubiona comta</i> C. L. Koch, 1839	3 ♀♀	1 ♀
Dictynidae		
<i>Dictyna arundinacea</i> (Linnaeus, 1758)	2 ♀♀	–
<i>Lathys heterophthalma</i> Kulczyński, 1891	3 ♀♀	–
Liocranidae		
<i>Agroeca proxima</i> (O. P.-Cambridge, 1871)	2 ♂♂	–
Linyphiidae		
<i>Ceratinella brevis</i> (Wider, 1834)	1 ♂, 1 ♀	1 ♂, 1 ♀
<i>Diplocephalus picinus</i> (Blackwall, 1841)	–	1 ♂
<i>Linyphia triangularis</i> (Clerck, 1757)	8 ♂♂, 17 ♀♀	1 ♂
<i>Maso sundevalli</i> (Westring, 1851)	1 ♂, 1 ♀	–
<i>Minyriolus pusillus</i> (Wider, 1834)	3 ♂♂, 35 ♀♀	–
<i>Panamomops mengei</i> Simon, 1926*	1 ♀	15 ♀♀
<i>Tapinocyba pallens</i> (O. P.-Cambridge, 1873)	9 ♀♀	1 ♀
<i>Tenuiphantes mengei</i> (Kulczyński, 1887)	1 ♀	–
<i>Tenuiphantes tenebricola</i> (Wider, 1834)	3 ♀♀	–
<i>Troxochrus scabriculus</i> (Westring, 1851)	2 ♂♂, 2 ♀♀	–
<i>Walckenaeria antica</i> (Wider, 1834)	1 ♂, 2 ♀♀	–
<i>Walckenaeria cucullata</i> (C. L. Koch, 1836)	–	1 ♀
Lycosidae		
<i>Alopecosa aculeata</i> (Clerck, 1757)	2 ♀♀	–
Miturgidae		
<i>Zora spinimana</i> (Sundevall, 1833)	1 ♀	–
Philodromidae		
<i>Philodromus collaris</i> C. L. Koch, 1835*	1 ♀	–
Salticidae		
<i>Evarcha falcata</i> (Clerck, 1757)	2 ♀♀	–
<i>Heliophanus dubius</i> C. L. Koch, 1835	1 ♀	–
<i>Neon reticulatus</i> (Blackwall, 1853)	7 ♀♀	1 ♀
Tetragnathidae		
<i>Tetragnatha obtusa</i> C. L. Koch, 1837*	3 ♀♀	–
Theridiidae		
<i>Crustulina guttata</i> (Wider, 1834)	1 ♀	–
<i>Enoplognatha ovata</i> (Clerck, 1757)	–	5 ♀♀
<i>Euryopis flavomaculata</i> (C. L. Koch, 1836)	5 ♀♀	–
<i>Platnickina tincta</i> (Walckenaer, 1802)	3 ♀♀	–
<i>Robertus lividus</i> (Blackwall, 1836)	–	1 ♀
<i>Theridion varians</i> Hahn, 1833	1 ♀	–
Thomisidae		
<i>Diae dorsata</i> (Fabricius, 1777)*	1 ♀	–
<i>Ozyptila praticola</i> (C. L. Koch, 1837)	–	2 ♂♂, 1 ♀




Fig. 1: Male of *Cheiracanthium punctorum*. **a.** habitus, dorsal view; palp in **b.** prolateral; **c.** ventral; **d.** retrolateral views

is the first record of the species from Finland, and its northernmost record across its known range.

Medical significance. Although *C. punctorum* is capable of causing very painful human envenomations, often resulting in an immediate burning sensation that may last for a few hours, reports of such cases are rare and often written in local languages, for local people and local practitioners (e.g. Krumpálová 1997, Chalupský 2011), despite the widespread and expanding distribution of this species in Europe (Nentwig et al. 2013, Varl et al. 2017). In most cases only local symptoms (pain, swelling, redness) have been reported, with a mild dermonecrosis developing on very rare occasions (Vetter et al. 2006).

Conclusions

The material collected during this field course contained mainly common species. Besides *C. punctorum*, which is a new record for Finland, only the record of the crab spider *Diaeaa dorsata* is somewhat interesting, as it is a southern, rarely collected and apparently spreading species in Finland that was recently removed from the Red List of Finnish spiders (Koponen et al. 2013, Pajunen et al. 2019). However, apart from these two species, *Cyclosa conica*, *Panamomops mengei*, *Philodromus collaris* and *Tetragnatha obtusa* are also new for the list of Seili spiders, which now comprises 215 species from 25 families (Appendix). This, as well as the large number of common species absent from the list, shows the need for further arachnological research in Seili. The number of reported species is about a third of what is currently known for the whole of Finland (Koponen et al. 2016).

Acknowledgments

We thank Kai Ruohomäki (Turku, Finland) for providing the data presented in Tab. 1, participants of the field course for their active efforts in collecting, identifying and sorting the specimens, and Petr Dolejš (Prague, Czech Republic), Jan Dolanský (Pardubice, Czech Republic), Varpu Vahtera (Turku, Finland) and an anonymous reviewer for their constructive comments and suggestions on the manuscript.

References

- Almquist S 2006 Swedish Araneae, part 2 – families Dictynidae to Salticidae. – Insect Systematics & Evolution, Supplement 63: 285–601
- Biteniektyte M & Relys V 2011 The checklist of Lithuanian spiders (Arachnida: Araneae). – Biologija 57: 148–158
- Cera I 2018 The checklist of Latvian spiders (Arachnida: Araneae). – Environmental and Experimental Biology 16: 139–152
- Chalupský J 2011 „Zápřednice jedovatá“ Doplněk k článku (DDD 03/10, 104) [“*Cheiracanthium punctorum*”, addendum to the article (DDD 03/10, 104)]. – Zpravodaj DDD 20: 47 [in Czech]
- Häkkilä P 1986 Lounaisuomalaisten kalliojyrkänteiden hämähäkifauna [Spider fauna on cliffs in southwestern Finland]. – Master thesis, University of Turku. 29 + 15 pp. [in Finnish]
- Koponen S, Fritzén NR & Pajunen T 2016 Checklist of spiders in Finland (Araneae), 6th version. – University of Turku. – Internet: http://biolcoll.utu.fi/arach/checklist_of_spiders_in_Finland.htm (9. Oct. 2022)
- Koponen S, Pajunen T & Fritzén NR 2013 Atlas of the Araneae of Finland. – Finnish Expert Group on Araneae. – Internet: <http://biolcoll.utu.fi/arach/aran2013/aranmaps.htm> (9. Oct. 2022)
- Kronestedt T 1990 Separation of two species standing as *Alopecosa aculeata* (Clerck) by morphological, behavioural and ecological characters, with remarks on related species in the *pulverulenta* group (Araneae, Lycosidae). – Zoologica Scripta 19: 203–225 – doi: [10.1111/j.1463-6409.1990.tb00256.x](https://doi.org/10.1111/j.1463-6409.1990.tb00256.x)
- Krumpálová Z 1997 Pozor, je krásny, ale hryzie! [Beware, it is beautiful, but it bites!]. – Živa 45: 35–36 [in Slovak]
- Miettinen M (ed.) 1997 Seilin saariston luonto – Yhteenveto kolmen vuosikymmenen tutkimuksista [Nature of the Seili archipelago – an overview of thirty years studies]. Metsähallituksen luonnonsuojelujulkaisuja, sarja A 80. Oy Edita Ab, Helsinki. 94 pp. [in Finnish]
- Nentwig W, Blick T, Bosmans R, Gloor D, Hänggi A & Kropf C 2022 Spiders of Europe. Version 07.2022. – Internet: <https://www.araneae.nmbe.ch> (17. Aug. 2022) – doi: [10.24436/1](https://doi.org/10.24436/1)
- Nentwig W, Gnädingen M, Fuchs J & Ceschi A 2013 A two year study of verified spider bites in Switzerland and a review of the European spider bite literature. – Toxicon 73: 104–110 – doi: [10.1016/j.toxicon.2013.07.010](https://doi.org/10.1016/j.toxicon.2013.07.010)
- Pajunen T, Koponen S, Fritzén NR & Lehtinen PT 2019 Hämähäkit. Spiders. Araneae. – In: Hyväriinen E, Juslén A, Kemppainen E, Uddström A & Liukko UM (eds) The 2019 Red List of Finnish Species. Ministry of the Environment & Finnish Environment Institute, Helsinki. pp. 329–335
- Sterghiu C 1985 Fam. Clubionidae. – Fauna Republicii Socialiste România (Arachnida) 5: 1–168
- Varl T, Grenc D, Kostanjšek R & Brvar M 2017 Yellow sac spider (*Cheiracanthium punctorum*) bites in Slovenia: case series and review. – Wiener Klinische Wochenschrift 129: 630–633 – doi: [10.1007/s00508-017-1217-8](https://doi.org/10.1007/s00508-017-1217-8)
- Vetter RS, Isbister GK, Bush SP & Boutin LJ 2006 Verified bites by yellow sac spiders (genus *Cheiracanthium*) in the United States and Australia: where is the necrosis? – The American Journal of Tropical Medicine and Hygiene 74: 1043–1048 – doi: [10.4269/ajtmh.2006.74.1043](https://doi.org/10.4269/ajtmh.2006.74.1043)
- World Spider Catalog 2022 World spider catalog. Version 23.5. Natural History Museum, Bern. – Internet: <http://wsc.nmbe.ch> (17. Aug. 2022) – doi: [10.24436/2](https://doi.org/10.24436/2)

Appendix. List of spider species known from the island of Seili, compiled from the spider database of the Zoological Museum of the University of Turku, Kronestedt (1990), Miettinen (1997) and the results of the present paper.

Agelenidae

Tegenaria domestica (Clerck, 1757)

Amaurobiidae

Amaurobius fenestralis (Ström, 1768)

Anyphaenidae

Anyphaena accentuata (Walckenaer, 1802)

Araneidae

Araneus diadematus Clerck, 1757

Araneus marmoreus Clerck, 1757

Araneus quadratus Clerck, 1757

Araneus sturmi (Hahn, 1831)

Araniella cucurbitina (Clerck, 1757)

Argiope bruennichi (Scopoli, 1772)

Cercidia prominens (Westring, 1851)

Cyclosa conica (Pallas, 1772)

Gibbaranea omoeda (Thorell, 1870)

Larinoides cornutus (Clerck, 1757)

Larinoides patagiatus (Clerck, 1757)

Leviellus stroemi (Thorell, 1870)

Cheiracanthiidae

Cheiracanthium punctatum (Villers, 1789)

Clubionidae

Clubiona comta C. L. Koch, 1839

Clubiona frutetorum L. Koch, 1867

Clubiona lutescens Westring, 1851

Clubiona neglecta O. P.-Cambridge, 1862

Clubiona pallidula (Clerck, 1757)

Clubiona stagnatilis Kulczyński, 1897

Clubiona subsultans Thorell, 1875

Cybaeidae

Cryphoeca silvicola (C. L. Koch, 1834)

Dictynidae

Brommella falcigera (Balogh, 1935)

Dictyna arundinacea (Linnaeus, 1758)

Dictyna pusilla Thorell, 1856

Lathys heterophthalma Kulczyński, 1891

Gnaphosidae

Callilepis nocturna (Linnaeus, 1758)

Drassodes cupreus (Blackwall, 1834)

Drassodes pubescens (Thorell, 1856)

Drassodes villosus (Thorell, 1856)

Drassyllus praeficus (L. Koch, 1866)

Drassyllus pumilus (C. L. Koch, 1839)

Gnaphosa bicolor (Hahn, 1833)

Gnaphosa montana (L. Koch, 1866)

Gnaphosa muscorum (L. Koch, 1866)

Haplodrassus signifer (C. L. Koch, 1839)

Haplodrassus soerrenseni (Strand, 1900)

Haplodrassus umbratilis (L. Koch, 1866)

Micaria formicaria (Sundevall, 1831)

Micaria nivosa L. Koch, 1866

Micaria pulicaria s.str. (Sundevall, 1831)

Micaria silesiaca L. Koch, 1875

Zelotes clivicola (L. Koch, 1870)

Zelotes exiguus (Müller & Schenkel, 1895)

Zelotes latreillei (Simon, 1878)

Zelotes longipes (L. Koch, 1866)

Zelotes petrensis (C. L. Koch, 1839)
Zelotes subterraneus (C. L. Koch, 1833)

Hahniidae

Hahnia nava (Blackwall, 1841)

Hahnia ononidum Simon, 1875

Hahnia pusilla C. L. Koch, 1841

Linyphiidae

Abacoproces saltuum (L. Koch, 1872)

Abisko abiskoensis (Holm, 1945)

Agyneta conigera (O. P.-Cambridge, 1863)

Agyneta ramosa Jackson, 1912

Agyneta rurestris (C. L. Koch, 1836)

Agyneta subtilis (O. P.-Cambridge, 1863)

Anguliphantes angulipalpis (Westring, 1851)

Bathyphantes nigrinus (Westring, 1851)

Bathyphantes parvulus (Westring, 1851)

Bolyphantes alticeps (Sundevall, 1833)

Centromerus arcarius (O. P.-Cambridge, 1873)

Centromerus incilium (L. Koch, 1881)

Centromerus sylvaticus (Blackwall, 1841)

Ceratinella brevis (Wider, 1834)

Dicylbium nigrum (Blackwall, 1834)

Diplocentria bidentata (Emerton, 1882)

Diplocephalus picinus (Blackwall, 1841)

Diplostyla concolor (Wider, 1834)

Dismodicus elevatus (C. L. Koch, 1838)

Drapetisca socialis (Sundevall, 1833)

Entelecara acuminata (Wider, 1834)

Entelecara congenera (O. P.-Cambridge, 1879)

Erigone dentigera O. P.-Cambridge, 1874

Erigonella hiemalis (Blackwall, 1841)

Evansia merens O. P.-Cambridge, 1901

Gongylidium rufipes (Linnaeus, 1758)

Helophora insignis (Blackwall, 1841)

Improphanes decolor (Westring, 1861)

Incestophantes crucifer (Menge, 1866)

Jacksonella falconeri (Jackson, 1908)

Kaestneria dorsalis (Wider, 1834)

Leptophantes leprosus (Ohlert, 1865)

Leptophantes minutus (Blackwall, 1833)

Linyphia tenuipalpis Simon, 1884

Linyphia triangularis (Clerck, 1757)

Macrargus carpenteri (O. P.-Cambridge, 1895)

Macrargus rufus (Wider, 1834)

Maro minutus O. P.-Cambridge, 1907

Maso sundevalli (Westring, 1851)

Metopobactrus prominulus (O. P.-Cambridge, 1873)

Micrargus apertus (O. P.-Cambridge, 1871)

Micrargus herbigradus (Blackwall, 1854)

Microlinyphia pusilla (Sundevall, 1830)

Microneta viaria (Blackwall, 1841)

Minyriolus pusillus (Wider, 1834)

Moebelia penicillata (Westring, 1851)

Neriene clathrata (Sundevall, 1830)

Neriene montana (Clerck, 1757)

Obscuriphantes obscurus (Blackwall, 1841)

Oedothorax agrestis (Blackwall, 1853)

Oedothorax apicatus (Blackwall, 1850)

Oedothorax gibbosus (Blackwall, 1841)

Palliduphantes pallidus (O. P.-Cambridge, 1871)

Panamomops mengei Simon, 1926

Pelecopsis elongata (Wider, 1834)

Pityohyphantes phrygianus (C. L. Koch, 1836)

Poeciloneta variegata (Blackwall, 1841)

Porrhomma pallidum Jackson, 1913

Porrhomma pygmaeum (Blackwall, 1834)

Silometopus ambiguus (O. P.-Cambridge, 1906)

Silometopus incurvatus (O. P.-Cambridge, 1873)
Stemonyphantes lineatus (Linnaeus, 1758)
Tapinocyba pallens (O. P.-Cambridge, 1873)
Tapinocyboides pygmaeus (Menge, 1869)
Tapinopa longidens (Wider, 1834)
Tenuiphantes alacris (Blackwall, 1853)
Tenuiphantes cristatus (Menge, 1866)
Tenuiphantes mengei (Kulczyński, 1887)
Tenuiphantes tenebricola (Wider, 1834)
Thyreosthenius parasiticus (Westring, 1851)
Trichoncus backmani Millidge, 1955
Trichopterna cito (O. P.-Cambridge, 1873)
Troxochrota scabra Kulczyński, 1894
Troxochrus scabriculus (Westring, 1851)
Typhochrestus digitatus (O. P.-Cambridge, 1873)
Walckenaeria antica (Wider, 1834)
Walckenaeria atrotibialis (O. P.-Cambridge, 1878)
Walckenaeria capito (Westring, 1861)
Walckenaeria cucullata (C. L. Koch, 1836)
Walckenaeria cuspidata Blackwall, 1833
Walckenaeria dysderoides (Wider, 1834)
Walckenaeria nudipalpis (Westring, 1851)

Liocranidae

Agroeca cuprea Menge, 1873
Agroeca proxima (O. P.-Cambridge, 1871)
Scotina gracilipes (Blackwall, 1859)

Lycosidae

Acantholycosa lignaria (Clerck, 1757)
Alopecosa aculeata (Clerck, 1757)
Alopecosa fabrilis (Clerck, 1757)
Alopecosa pinetorum (Thorell, 1856)
Alopecosa taeniata (C. L. Koch, 1835)
Arctosa leopardus (Sundevall, 1833)
Aulonia albimana (Walckenaer, 1805)
Pardosa amentata (Clerck, 1757)
Pardosa lugubris s. str. (Walckenaer, 1802)
Pardosa monticola (Clerck, 1757)
Pardosa palustris (Linnaeus, 1758)
Pardosa prativaga (L. Koch, 1870)
Pardosa pullata (Clerck, 1757)
Pirata piraticus (Clerck, 1757)
Trochosa ruricola (De Geer, 1778)
Trochosa terricola Thorell, 1856
Xerolycosa miniata (C. L. Koch, 1834)
Xerolycosa nemoralis (Westring, 1861)

Mimetidae

Ero furcata (Villers, 1789)

Miturgidae

Zora armillata Simon, 1878
Zora spinimana (Sundevall, 1833)

Philodromidae

Philodromus aureolus (Clerck, 1757)
Philodromus cespitum (Walckenaer, 1802)
Philodromus collinus C. L. Koch, 1835
Philodromus margaritatus (Clerck, 1757)
Thanatus formicinus (Clerck, 1757)
Thanatus striatus C. L. Koch, 1845
Tibellus maritimus (Menge, 1875)
Tibellus oblongus (Walckenaer, 1802)

Phrurolithidae

Phrurolithus festivus (C. L. Koch, 1835)

Pisauridae

Pisaura mirabilis (Clerck, 1757)

Salticidae

Aelurillus v-insignitus (Clerck, 1757)
Attulus pubescens (Fabricius, 1775)
Attulus terebratus (Clerck, 1757)
Dendryphantes rufus (Sundevall, 1833)
Evarcha falcata (Clerck, 1757)
Heliophanus dubius C. L. Koch, 1835
Myrmarachne formicaria (De Geer, 1778)
Neon reticulatus (Blackwall, 1853)
Pseudeuophrys erratica (Walckenaer, 1826)
Salticus cingulatus (Panzer, 1797)
Salticus scenicus (Clerck, 1757)
Talavera petrensis (C. L. Koch, 1837)

Segestriidae

Segestria senoculata (Linnaeus, 1758)

Sparassidae

Micrommata virescens (Clerck, 1757)

Tetragnathidae

Metellina mengei (Blackwall, 1869)
Metellina merianae (Scopoli, 1763)
Metellina segmentata (Clerck, 1757)
Pachygnatha clercki Sundevall, 1823
Pachygnatha degeeri Sundevall, 1830
Pachygnatha listeri Sundevall, 1830
Tetragnatha dear mata Thorell, 1873
Tetragnatha extensa (Linnaeus, 1758)
Tetragnatha obtusa C. L. Koch, 1837

Theridiidae

Crustulina guttata (Wider, 1834)
Cryptachaea riparia (Blackwall, 1834)
Enoplognatha ovata (Clerck, 1757)
Episinus angulatus (Blackwall, 1836)
Euryopis flavomaculata (C. L. Koch, 1836)
Neottiura bimaculata (Linnaeus, 1767)
Parasteatoda lunata (Clerck, 1757)
Pholcomma gibbum (Westring, 1851)
Phylloneta impressa (L. Koch, 1881)
Phylloneta sisyphus (Clerck, 1757)
Platnickina tincta (Walckenaer, 1802)
Robertus arundineti (O. P.-Cambridge, 1871)
Robertus lividus (Blackwall, 1836)
Robertus scoticus Jackson, 1914
Steatoda bipunctata (Linnaeus, 1758)
Theridion varians Hahn, 1833
Thymoites bellissimus (L. Koch, 1879)

Thomisidae

Diae dorsata (Fabricius, 1777)
Misumena vatia (Clerck, 1757)
Ozyptila atomaria (Panzer, 1801)
Ozyptila praticola (C. L. Koch, 1837)
Ozyptila trux (Blackwall, 1846)
Xysticus audax (Schrank, 1803)
Xysticus cristatus (Clerck, 1757)
Xysticus ulmi (Hahn, 1831)

Titanoecidae

Titanoeca spominima (Taczanowski, 1866)