

## **Big trees of small baskets: phylogeny of the Australian genus *Spyridium* (Rhamnaceae: Pomaderreae), focusing on biogeographic patterns and species circumscriptions**

Authors: Clowes, Catherine, Fowler, Rachael M., Fahey, Patrick S., Kellermann, Jürgen, Brown, Gillian K., et al.

Source: Australian Systematic Botany, 35(2) : 95-119

Published By: CSIRO Publishing

URL: <https://doi.org/10.1071/SB21034>

---


BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).



Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



# Big trees of small baskets: phylogeny of the Australian genus *Spyridium* (Rhamnaceae: Pomaderreae), focusing on biogeographic patterns and species circumscriptions

Catherine Clowes<sup>A,\*</sup> , Rachael M. Fowler<sup>A</sup>, Patrick S. Fahey<sup>A</sup>, Jürgen Kellermann<sup>B,C</sup>, Gillian K. Brown<sup>D</sup> and Michael J. Bayly<sup>A</sup> 

For full list of author affiliations and declarations see end of paper

**\*Correspondence to:**

Catherine Clowes  
School of Biosciences, The University of Melbourne, Vic. 3010, Australia  
Email: [cclowes@student.unimelb.edu.au](mailto:cclowes@student.unimelb.edu.au)

**Handling Editor:**

Russell Barrett

**Received:** 29 September 2021

**Accepted:** 10 March 2022

**Published:** 18 May 2022

**Cite this:**

Clowes C *et al.* (2022)  
*Australian Systematic Botany*  
35(2), 95–119. doi:[10.1071/SB21034](https://doi.org/10.1071/SB21034)

© 2022 The Author(s) (or their employer(s)). Published by CSIRO Publishing.

This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License ([CC BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/))

OPEN ACCESS

## ABSTRACT

*Spyridium* Fenzl is a genus of ~45 species endemic to south-western and south-eastern Australia. This study provides the most comprehensive phylogenies of *Spyridium* to date, analysing both entire chloroplast genomes and the nuclear ribosomal array (18S–5.8S–26S). There was substantial incongruence between the chloroplast and nuclear phylogenies, creating phylogenetic uncertainty, but some clear relationships and biogeographic patterns could be established. Analyses support the monophyly of *Spyridium*, identifying an early east–west split at the base of the nuclear phylogeny and deep divergences of New South Wales and Tasmanian endemic clades. We also found evidence of more recent dispersal events between eastern and western Australia and between Tasmania and the mainland. Eleven taxa were found to be monophyletic in the nrDNA phylogeny and two were clearly polyphyletic (*S. eriocephalum* Fenzl and *S. phyllicoides* Reissek). Although the polyphyly of *S. eriocephalum* correlates with the two varieties, suggesting distinct taxa, further research is required on *S. phyllicoides*.

**Keywords:** biogeography, chloroplast genome, molecular phylogeny, next-generation sequencing, nuclear ribosomal DNA, Rhamnaceae, species delimitation, *Spyridium*.

## Introduction


*Spyridium* Fenzl is a member of the tribe Pomaderreae Reissek ex Endl., which includes 10 genera and ~230 species endemic to Australia and New Zealand (Kellermann *et al.* 2008). *Spyridium* is derived from the Greek word *spyridion* meaning ‘a small basket’; a reference to the flowerheads surrounded by leafy bracts (Perrin 2018). The tribe is easily distinguished within the family Rhamnaceae and is characterised by species with stellate hairs on at least some of the vegetative or floral parts.

Generic boundaries within Pomaderreae have been difficult to define (Thiele and West 2004). According to Bentham (1863, p. 410):

...most [genera], even the most natural ones, are difficult to characterize. The differences in their flowers and fruits are very trifling; they often pass into one another by the finest gradations, and habit, foliage and inflorescence must often be relied upon for fixing generic limits.

Consequently, numerous species within the tribe have been transferred from one genus to another (and in some cases back), including species from *Cryptandra* Sm. and *Spyridium*.

The distinction between *Cryptandra* and *Spyridium* had traditionally been the presence of a floral tube in the former and its absence (or close to it) in the latter. However, for some species from both genera, the floral tube grades from effectively absent to very much present, blurring these traditional generic boundaries. Recently, more distinct



**Fig. 1.** Distribution of *Spyridium* in Australia. Dots represent filtered records accessed from [Atlas of Living Australia \(2020\)](#). States and territories are also highlighted as follows: WA, Western Australia; SA, South Australia; NT, Northern Territory; Qld, Queensland; NSW, New South Wales; ACT, Australian Capital Territory; Vic., Victoria and Tas., Tasmania.

characters between the two genera have been described (including disc, stipule and inflorescence characters), resulting in the transfer of several species of *Cryptandra* to *Spyridium*, including *S. scortechinii*<sup>1</sup> and *S. buxifolium* (Thiele and West 2004). In one extreme example, *S. waterhousei* has been moved to several genera within the tribe (including *Cryptandra* and *Stenanthemum* Reissek) before being re-instated as *Spyridium waterhousei* (Kellermann 2007) on the basis of its clear placement in molecular phylogenetic analyses (Kellermann et al. 2005).

The genus *Spyridium* currently includes ~45 species (Kellermann and Barker 2012) distributed in semi-arid to temperate regions of southern Australia (Fig. 1), including at least six potentially undescribed species identified as part of work towards a *Flora of Australia* treatment (Kellermann et al. 2022). The genus has two hotspots of local endemism, including ~20 species in southern South Australia (SA) and ~15 species in south-western Western Australia (WA). While the majority of species endemic to the south-west are confined to that region, two species have disjunct distributions across the Nullarbor Plain (*S. tricolor* and *S. subochreatum*). In south-eastern Australia, many of the species have narrow distributions, but several taxa (*S. parvifolium*, *S. vexilliferum* var. *vexilliferum*, *S. eriocephalum* var. *eriocephalum*) occur widely in this area (Coates and Kirkpatrick 1999; Kellermann and Barker 2012).

The monophyly of *Spyridium* has been confirmed by multiple molecular phylogenies (Richardson et al. 2004; Kellermann et al. 2005; Kellermann and Udovicic 2007; Hauenschild et al. 2016, 2018). The first nuclear DNA-based phylogeny (ITS) to focus on the tribe Pomaderreae was published by Kellermann et al. (2005), including 15 species of *Spyridium*. *Spyridium* was strongly supported as

monophyletic, and four geographically based clades were identified, with one being endemic to the eastern mainland, one from Tasmania, one being endemic to the south-west, and one including species from south-eastern Australia more generally. In addition, two species were moved from other genera to *Spyridium* (*S. daltonii* from *Trymalium* and *S. waterhousei* from *Cryptandra*). A chloroplast DNA phylogeny (*trnL-trnF*) of the tribe, which included the same (now) 17 species of *Spyridium* (Kellermann and Udovicic 2007), did not resolve monophyly of the genus, instead species of *Spyridium* formed part of a large polytomy at the base of the tree. But, like the findings of the ITS phylogeny, species endemic to the east and Tasmania formed separate clades. Finally, Hauenschild et al. (2018) included 18 species of *Spyridium* in their study on Gondwanan biogeography of ziziphoid Rhamnaceae, on the basis of nuclear, plastid and mitochondrial markers, and found that eastern endemics diverged first, followed by Tasmanian endemics and, finally, those endemic to the south-west. These studies have provided valuable information about relationships in *Spyridium* and suggest early divergence of endemic groups from the south-west, east and Tasmania. However, more than half of the species of *Spyridium* have not yet been included in a molecular phylogeny.

The aim of this study is to produce a comprehensive molecular phylogeny for the genus *Spyridium*, by including all described species and analysing the full chloroplast genome (cpDNA) and the 18S–5.8S–26S array of nuclear rDNA (nrDNA). We use these phylogenies to investigate broad biogeographic patterns within *Spyridium*, to explore vicariance, dispersal and diversification in the genus and to assess the monophyly and relationships within and among currently accepted species, as well as some proposed but so far undescribed taxa.

<sup>1</sup>Author names for all taxa at species level and lower are shown in Table 1; vouchers and authors of phrase-name taxa are in Table 2.

## Materials and methods

### Taxon sampling

In total, 143 samples of *Spyridium* were analysed in this study (Table 1). All species, subspecies and varieties of *Spyridium* recognised in the Australian Plant Census (APC; CHAH 2020) were included with at least one sample, except for *S. bifidum* var. *bifidum*, which was excluded because of sequencing issues associated with low DNA yield and quality. Of the 56 taxa included in this study, 45 (when counted at both specific and intraspecific levels) were represented by more than one accession. For species with wide distributions, multiple samples from across the geographic range were included where possible. Samples of six proposed but so far undescribed taxa were also included (Table 2). Four samples from three other genera (*Pomaderris* Labill., *Cryptandra* and *Trymalium*) from the tribe Pomadereae were included as outgroups.

For most samples, fresh leaf material was collected and dried in silica gel, along with a voucher specimen. Eight samples were obtained from existing herbarium collections at the WA Herbarium (PERTH). Herbarium voucher details are given in Table 1; many of the vouchers deposited at the University of Melbourne Herbarium (MELU) also have duplicates deposited variously at the National Herbarium of Victoria (MEL), the Tasmanian Herbarium (HO), PERTH or the State Herbarium of South Australia (AD).

### DNA extraction

Total genomic DNA was extracted from ~60 mg of silica-dried leaf material, or ~30 mg of herbarium material, following a modified cetyl trimethylammonium bromide (CTAB) protocol (Shepherd and McLay 2011; McLay 2017) based on Doyle and Doyle (1987). Where possible, young leaf material from stem tips or dried floral leaves was selected for extraction. DNA quality and quantity were recorded using a Nanodrop 2000 (NanoDrop Products) and Qubit 2.0 fluorometer (Invitrogen) and used to inform library preparation.

### Library preparation and DNA sequencing

Genomic DNA was prepared for multiplexed sequencing by using the library preparation and sequencing protocol of Schuster *et al.* (2018), with a few modifications. In total, 100 µL of each sonicated sample was transferred to a PCR plate and cleaned with solid-phase reversible immobilisation (SPRI) beads, by using a beads:sample ratio targeted to retain fragments of >300 bp (Rohland and Reich 2012). Following incubation and bead capture on a 96S super magnet plate (Alpaqua), the supernatant-free sample was washed with 100 µL of 80% ethanol. All subsequent 80% ethanol washes were also performed with 100 µL. Final q-PCRs were performed with 20 µL per reaction.

### Sequence assembly

Quality filtering and base calling was conducted at Walter and Eliza Hall Institute of Medical Research (WEHI) with Illumina pipeline software (ver. 1.7) and pre-processed with custom scripts, as in Schuster *et al.* (2018). De-multiplexed reads were imported into Geneious (ver. 10.2, Dotmatics, see <https://www.geneious.com/>; Kears *et al.* 2012) and trimmed using an error probability limit of 0.05. Paired-end reads were set by name and contigs were built in CLC Genomics Workbench (ver. 10.0.1, QIAGEN, see <https://digitalinsights.qiagen.com/>) by using default *de novo* settings. Contigs less than 1800 bp were discarded and remaining contigs were trimmed, removing 150 bp from each end.

Nuclear rDNA (18S–5.8S–26S, including both internal transcribed spacers and partial external and non-transcribed spacers) sequences were initially assembled by mapping contigs to the reference sequence for *Helianthus annuus* L. (GenBank number: KF767534), because there was no suitable extended (5'ETS–3'ETS + NTS) nrDNA reference for *Spyridium* or close relatives. Although *Helianthus* is not closely related to *Spyridium*, the highly conserved nature of sections of nrDNA (Jobs and Thien 1997), and success building long contigs spanning the region, resulted in successful mapping. Contigs were separated where required to assist mapping and, once completed, annotations were transferred from the reference and a draft consensus sequence was generated, with gaps preserved and the reference sequence used where data were missing. Paired reads were mapped to the draft nrDNA sequence for quality-control purposes, with a base calling threshold of 75% employed. The consensus sequence was manually adjusted where required and a final consensus sequence generated. Once the first nrDNA sequence of *Spyridium* was finalised (CC211; Table 1), this new sequence was used as the reference for subsequent sequence building.

The cpDNA genome for each sample was assembled using the same methodology as for the nrDNA sequences but using a different reference sequence (*S. parvifolium* var. *parvifolium*, GenBank accession MH234313; Clowes *et al.* 2018) and with the threshold for base calls in the final consensus sequence set at 50%. In addition, contigs and paired reads could map to multiple best-fit locations to enable the assembly of the inverted repeats.

### Phylogenetic analyses

Sequences were aligned using the MAFFT (ver. 7.308, see <https://mafft.cbrc.jp/alignment/software/>; Katoh *et al.* 2002; Katoh and Standley 2013) Geneious plugin under default settings. Aligned sequences were reviewed in Geneious, base pairs were re-aligned by eye where required and ambiguous regions were excluded from the final alignment. In addition, inverted repeat A (IRA) was excluded from the cpDNA alignment at this stage.

Aligned sequences were partitioned before model testing and phylogenetic analyses. For the nrDNA, the alignment

**Table 1.** Voucher information for samples included in this study.

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Cryptandra amara</i> Sm.	CC377	MELU	Cultivated; origin south-eastern Australia <sup>A</sup>	OK481389	OK624202
<i>Pomaderris rotundifolia</i> <sup>B</sup> (F.Muell.) Rye	L.S.J. Sweedman 7822	PERTH08703434	Cape Arid NP; Mallee; WA	OK481390	OK624203
<i>Pomaderris vacciniifolia</i> Reissek	CC378	MELUD117648a	Cultivated; South Eastern Highlands; origin Vic.	OK481391	OK624204
<i>Spyridium bifidum</i> subsp. <i>wanillae</i> Kellermann & W.R.Barker	JK587	AD272790	Wanilla; Eyre Yorke Block; SA	OK481392	OK624205
<i>Spyridium bifidum</i> subsp. <i>wanillae</i>	RMF374	MELUD122295a	Wanilla; Eyre Yorke Block; SA	OK481393	OK624206
<i>Spyridium burragorang</i> K.R.Thiele	CANB606 176 <sup>C</sup>	CBG9611161	Cultivated; origin NSW	OK481394	OK624346
<i>Spyridium burragorang</i>	CC567 <sup>D</sup>	MELUD122292a	Cessnock; Sydney Basin; NSW	OK481395	OK624347
<i>Spyridium buxifolium</i> (Fenzl) K.R.Thiele	CC504	MELUD117685a	Denman; Sydney Basin IBRA; NSW	OK481396	OK624207
<i>Spyridium buxifolium</i>	CC568 <sup>C</sup>	MELUD122291a	Goulburn River NP; Sydney Basin; NSW	OK481397	OK624208
<i>Spyridium cinereum</i> N.A.Wakef.	CC428	MELUD117664a	Grampians NP; Victorian Midlands; Vic.	OK481398	OK624209
<i>Spyridium cinereum</i>	CC498	MELUD117683a	Croajingolong NP; South East Corner; Vic.	OK481399	OK624210
<i>Spyridium coactilifolium</i> Reissek	CC322	MELUD117633a	Victor Harbor; Kanmantoo; SA	OK481400	OK624211
<i>Spyridium coactilifolium</i>	CC562	MELUD122276a	Newland Head CP; Kanmantoo; SA	OK481401	OK624212
<i>Spyridium coalitum</i> Kellermann & W.R.Barker	CC554	MELUD122269a	Kangaroo Island; Kanmantoo; SA	OK481402	OK624213
<i>Spyridium coalitum</i>	JK521	AD232557	Kangaroo Island; Kanmantoo; SA	OK481403	OK624214
<i>Spyridium cordatum</i> (Turcz.) Benth.	CC528	MELUD117699a	Gairdner, Esperance Plains; WA	OK481404	OK624215
<i>Spyridium cordatum</i>	CC530	MELUD117701a	Jerdacuttup; Esperance Plains; WA	OK481405	OK624216
<i>Spyridium cordatum</i>	GFC10247	MELUD122286a	Ravensthorpe; Esperance Plains; WA	OK481406	OK624217
<i>Spyridium cordatum</i>	JK370	AD213501	Ravensthorpe; Esperance Plains; WA	OK481408	OK624219
<i>Spyridium cordatum</i>	JK361	AD213497	Lake King; Mallee; WA	OK481407	OK624218

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium daltonii</i> (F.Muell.) Kellermann	CC513	MELUD117686a (CC505)	Grampians NP; Victorian Midlands; Vic.	OK481409	OK624220
<i>Spyridium daphnoides</i> (Reissek) Kellermann	CC550	MELUD122265a	Kangaroo Island; Kanmantoo; SA	OK481410	OK624221
<i>Spyridium daphnoides</i>	CC561	MELUD122275a	Deep Creek CP; Kanmantoo; SA	OK481411	OK624222
<i>Spyridium daphnoides</i>	JK721	AD283745	Lock; Eyre Yorke Block; SA	OK481412	OK624223
<i>Spyridium eriocephalum</i> Fenzl var. <i>eriocephalum</i>	CC515	MELUD117687a	Long Forest Nature Conservation Reserve; Victorian Midlands; Vic.	OK481413	OK624224
<i>Spyridium eriocephalum</i> var. <i>eriocephalum</i>	WAP01	MELUD122301a	East Risdon NR; Tasmanian South East; Tas.	OK481414	OK624225
<i>Spyridium eriocephalum</i> var. <i>glabrisepalum</i> J.M.Black	CC551	MELUD122266a	Kangaroo Island; Kanmantoo; SA	OK481415	OK624226
<i>Spyridium eriocephalum</i> var. <i>glabrisepalum</i>	JK484	AD232614	Kangaroo Island; Kanmantoo; SA	OK481416	OK624227
<i>Spyridium erymnocladum</i> W.R.Barker	JK729	AD283738	E of Hincks Wilderness Protection Area; Eyre Yorke Block; SA	OK481417	OK624228
<i>Spyridium fontis-woodii</i> Kellermann & W.R.Barker	CC563	MELUD122277a	Coorong; Naracoorte Coastal Plain; SA	OK481418	OK624229
<i>Spyridium fontis-woodii</i>	JK441	AD213808	Coorong; Naracoorte Coastal Plain; SA	OK481419	OK624230
<i>Spyridium furculentum</i> W.R.Barker & Kellermann	MJB2558	MELUD122313a	Little Desert; MDD; Vic.	OK481420	OK624231
<i>Spyridium glaucum</i> Rye	CC534	MELUD117705a	Ravensthorpe; Esperance Plains; WA	OK481421	OK624232
<i>Spyridium globulosum</i> (Labill.) Benth.	CC517	MELUD117688a	Woodman Point Recreation Reserve; Swan Coastal Plain; WA	OK481422	OK624233
<i>Spyridium globulosum</i>	EMS2413	MELUD122282a	Torndirrup NP; Warren; WA	OK481423	OK624234
<i>Spyridium globulosum</i>	GFC10245	MELUD122284a	Hopetoun; Esperance Plains; WA	OK481424	OK624235
<i>Spyridium gunnii</i> (Hook.f.) Benth.	WAP12	MELUD122305a	Renison Bell; Tasmanian West; Tas.	OK481425	OK624236
<i>Spyridium gunnii</i>	WAP23	MELUD122323a	Molesworth Conservation Area; Tasmanian South East; Tas.	OK481426	OK624237

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium halmaturinum</i> F.Muell. ex Benth.	CC548	MELUD122263a	Kangaroo Island; Kanmantoo; SA	OK481427	OK624238
<i>Spyridium halmaturinum</i>	JK497	AD232543	Kangaroo Island; Kanmantoo; SA	OK481428	OK624239
<i>Spyridium halmaturinum</i>	JK533	AD232599	Kangaroo Island; Kanmantoo; SA	OK481429	OK624240
<i>Spyridium halmaturinum</i>	JK536	AD232602	Kangaroo Island; Kanmantoo; SA	OK481430	OK624241
<i>Spyridium lawrencei</i> (Hook.f.) Benth.	CC237	MELUD117616a	Royal George; Tasmanian Northern Midlands; Tas.	OK481431	OK624242
<i>Spyridium lawrencei</i>	WAP20	MELUD122311a	Three Thumbs State Reserve; Tasmanian South East; Tas.	OK481432	OK624243
<i>Spyridium leucopogon</i> (F.Muell. ex Reissek) F.Muell.	JK572	AD272772	Ungarra; Eyre Yorke Block; SA	OK481433	OK624244
<i>Spyridium majoranifolium</i> (Fenzl) Rye	CC521	MELUD117692a	Lake Pleasant View NR; Jarrah Forest; WA	OK481434	OK624245
<i>Spyridium majoranifolium</i>	CC529	MELUD117700a	Stirling Range NP; Esperance Plains; WA	OK481435	OK624246
<i>Spyridium majoranifolium</i>	CC538	MELUD117709a	Ravensthorpe; Esperance Plains; WA	OK481436	OK624247
<i>Spyridium majoranifolium</i>	CC540	MELUD117711a	Jerdacuttup Lakes NR; Esperance Plains; WA	OK481437	OK624248
<i>Spyridium majoranifolium</i>	EMS2412	MELUD122281a	Torndirrup NP; Warren; WA	OK481438	OK624249
<i>Spyridium microcephalum</i> (Turcz.) Benth.	CC527	MELUD117698a	Gairdner; Esperance Plains; WA	OK481439	OK624250
<i>Spyridium microcephalum</i>	KRM826	MELUD122289a	Cape Arid NP; Esperance Plains; WA	OK481440	OK624251
<i>Spyridium minutum</i> Rye	CC542	MELUD117713a	Scaddan; Mallee; WA	OK481441	OK624252
<i>Spyridium minutum</i>	G.J. Keighery & N. Gibson 4183	PERTH07624697	Grass Patch; Mallee; WA	OK481442	OK624253
<i>Spyridium montanum</i> Rye	DAR 206	PERTH08742278	Stirling Range NP; Esperance Plains; WA	OK481443	OK624254
<i>Spyridium montanum</i>	S. Barrett 962	PERTH06045588	Stirling Range NP; Esperance Plains; WA	OK481444	OK624255

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium mucronatum</i> Rye subsp. <i>mucronatum</i>	CC522	MELUD117693a	Kojaneerup South; Esperance Plains; WA	OK481445	OK624256
<i>Spyridium mucronatum</i> subsp. <i>mucronatum</i>	CC526	MELUD117697a	Needilup; Mallee; WA	OK481446	OK624257
<i>Spyridium mucronatum</i> subsp. <i>recurvum</i> Rye	CC523	MELUD117694a	Kojaneerup South; Esperance Plains; WA	OK481448	OK624259
<i>Spyridium mucronatum</i> subsp. <i>recurvum</i>	CC532	MELUD117703a	Cheadanup; Mallee; WA	OK481449	OK624260
<i>Spyridium mucronatum</i> subsp. <i>recurvum</i>	GFC10244	MELUD122283a	Ravensthorpe; Esperance Plains; WA	OK481450	OK624261
<i>Spyridium mucronatum</i> subsp. <i>multiflorum</i> Rye	S. Barrett 1427	PERTH07354827	Fitzgerald River NP; Esperance Plains; WA	OK481447	OK624258
<i>Spyridium nitidum</i> N.A.Wakef.	JK477	AD232608	Kangaroo Island; Kanmantoo; SA	OK481451	OK624262
<i>Spyridium nitidum</i>	JK584	AD272784	Wanilla; Eyre Yorke Block; SA	OK481452	OK624263
<i>Spyridium obcordatum</i> (Hook.f.) W.M.Curtis	CC239	MELUD117618a	Greens Beach; Furneaux; Tas.	OK481453	OK624264
<i>Spyridium obcordatum</i>	CC284	MELUD117626a	Shearwater; Furneaux; Tas.	OK481454	OK624265
<i>Spyridium obovatum</i> (Hook.) Benth. var. <i>obovatum</i>	CC235	MELUD117614a	Apsley Conservation Area; Tasmanian South East; Tas.	OK481455	OK624266
<i>Spyridium obovatum</i> var. <i>obovatum</i>	WAP17	MELUD122309a	Douglas-Apsley NP; Tasmanian South East; Tas.	OK481456	OK624267
<i>Spyridium obovatum</i> var. <i>velutinum</i> (F.Muell. ex Reissek) Benth.	CC236	MELUD117615a	Friendly Beaches; Tasmanian South East; Tas.	OK481457	OK624268
<i>Spyridium obovatum</i> var. <i>velutinum</i>	WAP08	MELUD122303a	Snug; Tasmanian South East; Tas.	OK481458	OK624269
<i>Spyridium obovatum</i> var. <i>velutinum</i>	WAP15	MELUD122306a	Tasman NP; Tasmanian South East; Tas.	OK481459	OK624270
<i>Spyridium obovatum</i> var. <i>velutinum</i>	WAP16	MELUD122307a	Orford; Tasmanian South East; Tas.	OK481460	OK624271
<i>Spyridium obovatum</i> var. <i>velutinum</i>	WAP19	MELUD122310a	Sandspit River Forest Reserve; Tasmanian South East; Tas.	OK481461	OK624272
<i>Spyridium oligocephalum</i> (Turcz.) Benth.	CC536	MELUD117707a	Hopetoun; Esperance Plains; WA	OK481462	OK624273

(Continued on next page)



Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium parvifolium</i> (Hook.) F.Muell.	CC057	MELUD117595a	Brisbane Ranges; Victorian Midlands; Vic.	OK481463	OK624274
<i>Spyridium parvifolium</i>	CC174	MELUD117609a	Dandenong Ranges NP; South Eastern Highlands; Vic.	OK481464	OK624275
<i>Spyridium parvifolium</i>	CC211	MELUD155066a	Rocky Cape NP; King; Tas.	OK481465	MH234313
<i>Spyridium parvifolium</i>	CC273	MELUD117623a (CC265)	Flinders Island; Furneaux; Tas.	OK481466	OK624276
<i>Spyridium parvifolium</i>	CC433	MELUD117665a	Grampians NP; Victorian Midlands; Vic.	OK481467	OK624277
<i>Spyridium parvifolium</i>	CC490	MELUD117682a (CC484)	Cann River; South East Corner; Vic.	OK481468	OK624278
<i>Spyridium parvifolium</i>	MJB2239E	MELUD118621a	Mount Buffalo NP; South Eastern Highlands; Vic.	OK481469	OK624279
<i>Spyridium parvifolium</i>	MJB2288A	MELUD118616a	Burrinjuck Waters State Park; Southern Highlands; NSW	OK481470	OK624280
<i>Spyridium phlebophyllum</i> (F.Muell.) F.Muell.	RMF388	MELUD122296a	Ikara–Flinders Ranges NP; Flinders Lofty Block; SA	OK481471	OK624281
<i>Spyridium phylloides</i> Reissek	CC557	MELUD122272a	Kangaroo Island; Kanmantoo; SA	OK481472	OK624282
<i>Spyridium phylloides</i>	CC565	MELUD122279a	Coorong; Naracoorte Coastal Plain; SA	OK481473	OK624283
<i>Spyridium phylloides</i>	JK487A2	AD232583	Kangaroo Island; Kanmantoo; SA	OK481474	OK624284
<i>Spyridium phylloides</i>	JK593	AD273231	Coffin Bay NP; Eyre Yorke Block; SA	OK481475	OK624285
<i>Spyridium polycephalum</i> (Turcz.) Rye	CC525	MELUD117696a	Needilup; Mallee; WA	OK481476	OK624286
<i>Spyridium polycephalum</i>	CC535	MELUD117706a	Ravensthorpe; Esperance Plains; WA	OK481477	OK624287
<i>Spyridium riparium</i> Rye	CC518	MELUD117689a	Mount Lindesay NP; Jarrah Forest; WA	OK481478	OK624288
<i>Spyridium riparium</i>	CC519	MELUD117690a	Kentdale; Warren; WA	OK481479	OK624289
<i>Spyridium scabridum</i> (Tate) Kellermann & W.R.Barker	CC553	MELUD122268a	Kangaroo Island; Kanmantoo; SA	OK481480	OK624290

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium scabridum</i>	JK502	AD232548	Kangaroo Island; Kanmantoo; SA	OK481481	OK624291
<i>Spyridium scabridum</i>	JK527	AD232565	Kangaroo Island; Kanmantoo; SA	OK481482	OK624292
<i>Spyridium scortechinii</i> (F.Muell.) K.R.Thiele	CC503	MELUD117684a	Tilba Tilba; South East Corner; NSW	OK481483	OK624293
<i>Spyridium scortechinii</i>	CC569 <sup>E</sup>	MELUD122290a	Kurrajong Heights; Sydney Basin; NSW	OK481484	OK624294
<i>Spyridium scortechinii</i>	JRH3933A	MELUD122298a	Woodsreef; Nandewar; NSW	OK481485	OK624295
<i>Spyridium</i> sp. Dwarf (J.Kellermann 579) Kellermann	JK489	AD232593	Seal Bay CP, Kangaroo Island; Kanmantoo; SA	OK481486	OK624296
<i>Spyridium</i> sp. Dwarf (J.Kellermann 579)	JK579	AD272779	Lincoln NP; Eyre Yorke Block; SA	OK481487	OK624297
<i>Spyridium</i> sp. Dwarf (J.Kellermann 579)	JK740	AD283757	Kangaroo Island; Kanmatoo; SA	OK481488	OK624298
<i>Spyridium</i> sp. Finniss (J.Kellermann 653 & F. Nge) Kellermann	JK653	AD283669	Bullock Hill CP; Kanmantoo; SA	OK481489	OK624299
<i>Spyridium</i> sp. Finniss (J.Kellermann 653 & F. Nge)	JK661	AD283668	Currency Creek; Kanmantoo; SA	OK481490	OK624300
<i>Spyridium</i> sp. Jerdacuttup (A.Williams 332) WA Herbarium	CC533	MELUD117704a	Jerdacuttup; Esperance Plains; WA	OK481491	OK624301
<i>Spyridium</i> sp. Jerdacuttup (A.Williams 332)	KRM738	MELUD122287a	WA36183 NR; Esperance Plains; WA	OK481492	OK624302
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560) Kellermann	JK510	AD232574	Kangaroo Island; Kanmantoo; SA	OK481495	OK624305
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560)	JK511	AD232575	Kangaroo Island; Kanmantoo; SA	OK481496	OK624306
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560)	JK518	AD232582	Kangaroo Island; Kanmantoo; SA	OK481497	OK624307
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560)	JK504A	AD232552	Kangaroo Island; Kanmantoo; SA	OK481494	OK624304
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560)	JK526A	AD232561	Kangaroo Island; Kanmantoo; SA	OK481498	OK624308

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium</i> sp. Kangaroo Island (W.R.Barker 7560)	CC556	MELUD122271a	Kangaroo Island; Kanmantoo; SA	OK481493	OK624303
<i>Spyridium</i> sp. Red dots (J.Kellermann 689) Kellermann	JK689	AD283725	Warramboo; Eyre Yorke Block; SA	OK481499	OK624309
<i>Spyridium</i> sp. Wollar (E.F.Constable s.n., NSW 16590) Kellermann	CC566 <sup>F</sup>	MELUD122293a	Wollar; Sydney Basin; NSW	OK481500	OK624310
<i>Spyridium spadiceum</i> (Fenzl) Benth.	CC520	MELUD117691a	Gull Rock NP; Jarrah Forest; WA	OK481501	OK624311
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i> Kellermann & W.R.Barker	CC345	MELUD117636a	Cowell; Eyre Yorke Block; SA	OK481503	OK624313
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	CC348	MELUD117638a	Campoona; Eyre Yorke Block; SA	OK481504	OK624314
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	RMF256	MELUD122299a	Wharminda; Eyre Yorke Block; SA	OK481507	OK624317
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	BSB838–173	AD261711	Hiltaba NR; Gawler; SA	OK481502	OK624312
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	CC546	MELUD117717a	Kimbra; Eyre Yorke Block; SA	OK481505	OK624315
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	JK561	AD273243	Cowell; Eyre Yorke Block; SA	OK481506	OK624316
<i>Spyridium stenophyllum</i> subsp. <i>renovatum</i>	RMF372	MELUD122294a	Rudall; Eyre Yorke Block; SA	OK481508	OK624318
<i>Spyridium stenophyllum</i> (Reissek) Kellermann & W.R.Barker subsp. <i>stenophyllum</i>	PSF46	MELUD122297a	Mangalo; Eyre Yorke Block; SA	OK481509	OK624319
<i>Spyridium subochreatum</i> (F.Muell.) Reissek	JK601	AD273246	Campoona; Eyre Yorke Block; SA	OK481514	OK624324
<i>Spyridium subochreatum</i>	CC371	MELUD117643a	Billiatt Wilderness Protection Area; MDD; SA	OK481511	OK624321
<i>Spyridium subochreatum</i>	MJB2557C	MELUD122314a	Little Desert NP; MDD; Vic.	OK481515	OK624325
<i>Spyridium subochreatum</i>	E.D. Adams 21/0907	PERTH08067805	Nuytsland NR; Esperance Plains; WA	OK481513	OK624323
<i>Spyridium subochreatum</i>	CC349	MELUD117639a	Campoona; Eyre Yorke Block; SA	OK481510	OK624320

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium subochreatum</i>	CC372	MELUD117644a	Ngarkat CP; MDD; SA	OK481512	OK624322
<i>Spyridium thymifolium</i> Reissek	CC298	MELUD117627a (CC285)	Mount Billy Conservation Reserve; Kanmantoo; SA	OK481516	OK624326
<i>Spyridium tricolor</i> W.R.Barker & Rye	CC545	MELUD117716a	Watraba; Eyre Yorke Block; SA	OK481517	OK624327
<i>Spyridium ulicinum</i> (Hook.) Benth.	WAP03C	MELUD122304a	Glenlusk; Tasmanian South East; Tas.	OK481519	OK624329
<i>Spyridium ulicinum</i>	WAP04C	MELUD122315a	Collinsvale; Tasmanian South East; Tas.	OK481520	OK624330
<i>Spyridium ulicinum</i>	WAP05B	MELUD122316a	Neika; Tasmanian South East; Tas.	OK481521	OK624331
<i>Spyridium ulicinum</i>	WAP06	MELUD122317a	Lower Longley; Tasmanian South East; Tas.	OK481522	OK624332
<i>Spyridium ulicinum</i>	WAP09B	MELUD122318a	Snug Tiers Nature Recreation Area; Tasmanian Southern Ranges; Tas.	OK481523	OK624333
<i>Spyridium ulicinum</i>	WAP10B	MELUD122319a	Ouse; Tasmanian Southern Ranges; Tas.	OK481524	OK624334
<i>Spyridium ulicinum</i>	WAP11C	MELUD122320a	Ouse; Tasmanian Southern Ranges; Tas.	OK481525	OK624335
<i>Spyridium ulicinum</i>	CC283	MELUD117625a (CC282)	Cataract Gorge Reserve; Tasmanian Northern Midlands; Tas.	OK481518	OK624328
<i>Spyridium vexilliferum</i> var. <i>latifolium</i> Benth.	CC464	MELUD117670a	Kingston SE; Naracoorte Coastal Plain; SA	OK481526	OK624336
<i>Spyridium vexilliferum</i> var. <i>latifolium</i>	CC473	MELUD117675a	Desert Camp Conservation Reserve; Naracoorte Coastal Plain; SA	OK481527	OK624337
<i>Spyridium vexilliferum</i> (Hook.) Reissek var. <i>vexilliferum</i>	DJM738	MEL	Grampians NP; Victorian Midlands; Vic.	OK481528	OK624338
<i>Spyridium vexilliferum</i> var. <i>vexilliferum</i>	WAP07G	MELUD118619a	Tom Gibson NR; Tasmanian Northern Midlands; Tas.	OK481529	OK624339
<i>Spyridium vexilliferum</i> var. <i>vexilliferum</i>	WAP13	MELUD122321a	Arthur Pieman Conservation Reserve; King; Tas.	OK481530	OK624340

(Continued on next page)

Table 1. (Continued)

Species	Collecting number	Voucher	Location	Nuclear rDNA GenBank accession number	Chloroplast genome GenBank accession number
<i>Spyridium vexilliferum</i> var. <i>vexilliferum</i>	WAPI4	MELUD122322a	Freycinet NP; Tasmanian South East; Tas.	OK481531	OK624341
<i>Spyridium villosum</i> (Turcz.) Benth.	S. Barrett 2200	PERTH08773874	Stirling Range NP; Esperance Plains; WA	OK481533	OK624343
<i>Spyridium villosum</i>	S. Barrett 950	PERTH06045596	Stirling Range NP; Esperance Plains; WA	OK481532	OK624342
<i>Spyridium waterhousei</i> F.Muell.	CC558	MELUD122273a	Kangaroo Island; Kanmantoo; SA	OK481534	OK624344
<i>Trymalium elachophyllum</i> Rye	CC531	MELUD117702a	Cheadanup; Mallee; WA	OK481535	OK624345

Names of described taxa included in this study follow the Australian Plant Census (CHAH 2020), the exception being *S. daphnoides*, which is a recent name change not yet updated in the Census (Kellermann 2021). Phrase names were formed according to Barker (2005). Authorities have been provided with the first reference to each taxon. Collectors are as follows: CC, Catherine Clowes; JK, Juergen Kellerman; L.S.J. Sweedman, Luke Sweedman; RMS, Rachael Fowler; CANB, Australian National Botanic Gardens (living collection number); GFC, Gillian Craig; WAP, Mark Wapstra; MJB, Michael Bayly; EMS, Libby Sandiford; KRM, Kenneth Mills; G.J. Keighery, Gregory Keighery; N. Gibson, Neil Gibson; DAR, Damien Rathbone; Barrett, Sarah Barrett; BSB, Juergen Kellerman; JRH, John Hosking; PSF, Patrick Fahey. To standardise location details, for samples collected from mainland Australia, either the reserve name is provided or the nearest town (where the sample was collected outside a reserve system). Location details provided for samples collected on islands include the island name. States have also been provided for all samples. For all samples, Interim Biogeographic Regionalisation for Australia 7 (IBRA 7) regions have also been provided in location information (Department of Agriculture Water and the Environment 2020). Location abbreviations are as follows: SA, South Australia; WA, Western Australia; Vic., Victoria; Tas. Tasmania; NSW, New South Wales; NP, National Park; NR, Nature Reserve; CP, Conservation Park. Nuclear rDNA and chloroplast genome GenBank accession numbers are provided. A voucher has not yet been lodged for *S. vexilliferum* var. *vexilliferum* (DJM738), but it is accessible at the National Herbarium of Victoria upon request from Daniel J. Murphy.

<sup>A</sup>The state of origin for this sample was unknown, but the distribution of this species is known to be south-eastern Australia.

<sup>B</sup>This sample was lodged with PERTH as *Spyridium tricolor*, but was determined by Juergen Kellerman to be *Pomaderris rotundifolia* on receipt from PERTH.

<sup>C</sup>This number is the accession for the plant in the Australian National Botanic Gardens Living Collections.

<sup>D</sup>This sample was collected by Dr Stephen Bell (Eastcoast Flora Survey). A unique collecting number was not provided, so this collecting number was generated by Catherine Clowes.

<sup>E</sup>This sample was collected by Dr Mathew Dell (Southeast Botanical Consulting). A unique collecting number was not provided, so this collecting number was generated by Catherine Clowes.

<sup>F</sup>This sample was collected by Dr Stephen Bell (Eastcoast Flora Survey). A unique collecting number was not provided, so this collecting number was generated by Catherine Clowes.

**Table 2.** Distribution and affinities of phrase-name taxa included in this study.

Phrase name	Distribution	Morphological affinities
<i>Spyridium</i> sp. Kangaroo Island (W.R. Barker 7560) Kellermann <sup>A</sup>	Sandy soils on Kangaroo Island	Morphologically similar to <i>S. vexilliferum</i> var. <i>vexilliferum</i> and <i>S. thymifolium</i>
<i>Spyridium</i> sp. Dwarf (J. Kellermann 579) Kellermann <sup>A</sup>	Southern Eyre Peninsula (SA) and Kangaroo Island on sand over limestone	Prostrate to upright shrub with similarities to <i>S. phyllicoides</i>
<i>Spyridium</i> sp. Red Dots (J. Kellermann 689) Kellermann <sup>A</sup>	Eyre Yorke Block on limestone	Like <i>S. phyllicoides</i> , but with a much smaller habit
<i>Spyridium</i> sp. Finnis (J. Kellermann 653 & F. Nge) Kellermann <sup>A</sup>	North-eastern Fleurieu Peninsula	A robust shrub which has similarities to <i>S. eriocephalum</i> var. <i>eriocephalum</i> and var. <i>glabrisepalum</i>
<i>Spyridium</i> sp. Wollar (E.F. Constable s.n.; NSW16590) Kellermann <sup>A</sup>	Only found in an area north-west of Sydney near Dubbo; on sandy soils	Similar to both <i>S. eriocephalum</i> var. <i>eriocephalum</i> and <i>S. scortechinii</i>
<i>Spyridium</i> sp. Jerdacuttup (A. Williams 332) WA Herbarium <sup>B</sup>	Jerdacuttup region of WA	Similar to <i>S. cordatum</i> (Wilkins <i>et al.</i> 2011)

<sup>A</sup>Phrase name used as part of collaborative work towards a *Flora of Australia* treatment (Kellermann *et al.* 2022); published here for the first time.

<sup>B</sup>Phrase name currently used in Australian Plant Census (CHAH 2020).

was partitioned as follows: partial external transcribed spacer (5'ETS), 18S, internal transcribed spacer 1 (ITS1), 5.8S, internal transcribed spacer 2 (ITS2), 26S, and partial non-transcribed spacer (3'ETS + NTS). The four cpDNA partitions, based on annotations from the reference sequence of *S. parvifolium* var. *parvifolium* (GenBank accession MH234313) were as follows: gene-coding sequence (CDS); transfer ribonucleic acid (tRNA); ribosomal ribonucleic acid (rRNA); and all remaining sequences, including introns and intergenic spacers (referred to in the partition as spacers).

Both nrDNA and cpDNA alignments were analysed using Bayesian inference (BI) and maximum likelihood (ML) methods. For the BI analyses, model testing was performed for each partition following Akaike's information criterion (AIC) using MrModeltest2 (ver. 2.4, J. A. A. Nylander, see <https://github.com/nylander/MrModeltest2>) for nrDNA (selected models: 5'ETS GTR + G, 18S GTR + I + G, ITS1 SYM + G, ITS2 K80 + G, 26S GTR + I + G and 3'ETS + NTS HKY + G; with 5.8S alignment being excluded from analyses because all sequences were identical) and cpDNA (selected models: CDS GTR + I + G, tRNA K80 + I, rRNA HKY and spacers GTR + I + G). Bayesian inference analyses were undertaken in MrBayes XSEDE (ver. 3.2.6, C. Zhang, J. Huelsenbeck, P. van der Mark, F. Ronquist and M. Teslenko, see <https://github.com/NBISweden/MrBayes/releases>; Ronquist and Huelsenbeck 2003) by using the CIPRES portal (Miller *et al.* 2010). For the nrDNA alignment (6367 bp), two independent analyses with four chains (Markov-chain Monte Carlo) were run for 5 000 000 generations, sampling every 1000 steps, with a burnin of 25%. For the cpDNA alignment (168 343 bp), chains were run for 2 500 000 generations, sampling every 500 steps. Output files were viewed in Tracer (ver. 1.6, A. Rambaut, A. J. Drummond and M. Suchard, see <https://github.com/beast-dev/tracer>), checking convergence (<0.01 standard deviation of split frequencies). The 50% majority-rule consensus

trees were visualised in FigTree (ver. 1.4.2, A. Rambaut and A. J. Drummond, see <https://github.com/rambaut/figtree>) with posterior probabilities (PP) of  $\geq 0.95$  being viewed as fully supported, and those lower considered unsupported.

Maximum likelihood analysis was performed with IQ-Tree using default settings (ver. 1.6.12, L. T. Nguyen, H. A. Schmidt, A. von Haeseler and B. Q. Minh, see <https://iqtree.org/>; Nguyen *et al.* 2015). IQ-Tree automatically employs ModelFinder (Kalyanamoorthy *et al.* 2017) to select models for each partition by using Bayesian information criterion (BIC) as default. For nrDNA analysis, the models were as follows: 5'ETS TPM2 + F + G4, 18S K2P + I + G4, ITS1 TIM2e + G4, 5.8S JC, ITS2 TNe + G4, 26S TIM3 + F + I and 3'ETS + NTS TN + F + G4; for cpDNA analysis, the models were as follows: CDS TVM + F + I, tRNA K2P, rRNA HKY + F and spacers TVM + F + I + G4. In total, 261 parsimony-informative characters were reported for the nrDNA ML analysis and 3129 were reported for the cpDNA ML analyses. The bootstrap consensus trees were viewed in FigTree and exported to TreeGraph 2 (ver. 2.14.0-771 beta, Bio10, see <https://treegraph.bioinfweb.info/>; Stöver and Müller 2010) where branches with <50% ultrafast bootstrap (UFBS) support where collapsed to allow for easier comparison with the BI tree. UFBS values of  $\geq 95\%$  were viewed as supported and those below 95% were considered unsupported.

The results of the nrDNA phylogeny were mapped using QGIS (ver. 3.18.3, QGIS Development Team, see <http://qgis.org/en/site/index.html>). Species distributions were accessed from Atlas of Living Australia (2020) and these data filtered removing samples lacking collection dates. Outliers were reviewed and removed where errors in location or identification were suspected. Taxa were mapped by clades according to the nrDNA phylogeny.

All sequences included in this study were uploaded to GenBank (Table 1). Text files containing the alignment,




Fig. 2. (Caption on next page)

**Fig. 2.** Nuclear rDNA (nrDNA) phylogeny of *Spyridium*, based on Bayesian inference (BI) analysis. Bayesian posterior probabilities (PP) and ultrafast bootstrap (UFBS) values are shown at nodes when <95%; values ≥95% are not shown. Where one value for a node is supported (≥95%) and the other for that node is unsupported (<95%), only the unsupported value is shown. Where a hyphen (-) is provided at a node, this node varied in resolution in the ML tree and was therefore not transferable to the BI phylogeny. Colour coding of clades and taxa in the bar to the right of the tree matches that used on maps in Fig. 3, 4. Labels are given for some clades (A–J) and subclades (A1–J3) discussed in text. Species polyphyletic across clades are highlighted in red text. Monophyletic taxa with supported nodes are highlighted in green text. Note: *S. eriocephalum* is polyphyletic, but var. *eriocephalum* is monophyletic (and therefore coloured half red and half green). Dashed lines associated with *S. tricolor*, *S. glaucum*, *S. phlebophyllum* and *S. subochreatum* E.D.Adams 21/0907 are provided as reference points connecting taxa to the sidebar.

partitions, BI and ML trees for each dataset (nrDNA and cpDNA) were submitted to TreeBase (accession number 28815).

## Results

### Nuclear rDNA phylogeny

The topologies of BI and ML nrDNA trees were largely congruent, and only the BI tree is shown here, with ML bootstrap values mapped onto it (Fig. 2). A key supported difference in the ML tree was in the placement of *S. daltonii* as divergent from the *S. parvifolium* clade (UFBS = 95), with *S. obcordatum* and *S. cinereum* being unsupported as successive sisters to a clade of *S. parvifolium* samples (UFBS = 85 for the former and UFBS = 65 for the latter). Another supported difference was that *S. burragorang* formed a clade, sister to *S. buxifolium* (UFBS = 98).

*Spyridium* was resolved as monophyletic (PP = 1.0, UFBS = 100), with a distinct split at the base of the tree between largely south-western Australian endemics (Clade A) and largely south-eastern Australian species (Clade B). Clade A (PP = 1.0, UFBS = 96) grouped species endemic to south-western Australia together with one species (*S. tricolor*) that includes outlying populations disjunct across the Nullarbor Plain (Fig. 3a), whereas Clade B (PP = 0.93, UFBS = 94) included species from south-eastern Australia, one that occurs in both eastern and western Australia (*S. subochreatum*, Clade J; Fig. 2, 3j), plus two south-western endemics (*S. oligocephalum* and *S. polycephalum*, Clade H; Fig. 2, 3h) nested among the south-eastern taxa. Within the largely eastern clade (B), two early diverging lineages were a clade (PP = 1.0, UFBS = 100) of primarily New South Wales (NSW and southern Queensland<sup>2</sup>) endemics (Clade C; Fig. 2, 3b) and a clade (PP = 1.0, UFBS = 100) of Tasmanian endemics (Clade D; Fig. 2, 3b). The only other Tasmanian endemic, *S. obcordatum*, was placed in Clade E with *S. cinereum*, *S. parvifolium* and *S. daltonii* (PP = 0.73; Fig. 2, 3d).

<sup>2</sup>*Spyridium scortechinii* has a distribution that extends approximately 20 km into southern Queensland. However, this species is largely restricted to NSW. For ease of reading, for the remainder of the text, we will refer to this clade (and the equivalent clade in the cpDNA phylogeny) as NSW endemics.

Of the 45 species, subspecies or varieties represented by more than one accession, 11 (24.4%) were monophyletic and 11 (24.4%) were polyphyletic (Fig. 2, Table 3). One species polyphyletic across disparate clades was *S. eriocephalum*, although var. *eriocephalum* was monophyletic (PP = 1.0, UFBS = 100; Clade F), whereas var. *glabrisepalum* samples were placed in a polytomy (PP = 100, UFBS = 97; Clade I). The other example of a species polyphyletic across disparate clades was *S. phyllicoides*, with the four accessions of this species divided among two clades (two samples in Clade I and two samples in Clade J2; Fig. 2, 4). Owing to a lack of support (BI or ML) for many branches at the tips of the nrDNA phylogeny, the remaining 23 taxa represented by more than one accession (51.1% of included taxa), were unresolved, i.e. with their monophyly neither supported nor strongly refuted.

### Chloroplast genome phylogeny

The topologies of BI and ML cpDNA trees were largely congruent, and only the BI tree is shown here, with ML bootstrap values being mapped onto it (Fig. 5). One unsupported difference in the ML tree was that *S. burragorang* formed a clade (UFBS = 92), sister to *S. buxifolium* (UFBS = 100). *Spyridium* was resolved as monophyletic (PP = 1.0, UFBS = 100). Clades dominated by Tasmanian endemics (K; PP = 1.0, UFBS = 100) and NSW endemics (L; PP = 1.0, UFBS = 100) diverged early in the cpDNA phylogeny (Fig. 5). *Spyridium phlebophyllum* also diverged early on a branch between Clades K and L (PP = 1.0, UFBS = 98). Clade N predominantly contained taxa endemic to WA, as well as a SA sample of *S. tricolor* (PP = 1.0, UFBS = 100), a species found in WA and in SA that is disjunct on either side of the Nullarbor Plain (Fig. 3a, 5). One Tasmanian endemic species (*S. obcordatum*) was sister to *S. parvifolium* and *S. daltonii* in Clade O (PP = 1.0, UFBS = 100; Fig. 5). *Spyridium polycephalum* and *S. oligocephalum*, both WA endemics, were resolved in Clade M6 (PP = 1.0, UFBS = 100; Fig. 3h, 5) as sister to a WA sample of the widespread, polyphyletic *S. subochreatum* (Fig. 3j, 5, Table 1). These samples were found nested within the






Fig. 3. (Caption on next page)

**Fig. 3.** Distributions of nrDNA clades of *Spyridium*, colour-coded to match groups shown in Fig. 2. Clade distributions are based on those of included species, using records in the *Atlas of Living Australia* (2020). Distributions of *S. phylloides*, *S. sp. Dwarf* (J.Kellermann 579) and *S. sp. Red Dots* (J.Kellermann 689) have been omitted from these maps and are provided in Fig. 4. (a) Distribution of Clade A1 (mid blue), Clade A2 (royal blue), *S. tricolor* (light blue) and *S. glaucum* (dark blue). The location of sample CC545 (*S. tricolor*) is highlighted. (b) Distribution of Clade C. The general location of the southern transition zone is also highlighted. (c) Distribution of Clade D. (d) Distribution of Clade E. (e) Distribution of *S. phlebophyllum*. (f) Distribution of *S. eriocephalum* var. *eriocephalum* from Clade F. (g) Distribution of Clade G. The location of sample CC566 (*S. sp. Wollar*) is highlighted. (h) Distribution of Clade H. (i) Distribution of Clade I, excluding *S. phylloides* and *S. sp. Dwarf* (J.Kellermann 579). (j) Distribution of Clade J1 (bright pink), Clade J2 (deep pink) and Clade J3 (light pink). The location of sample E.D.Adams 21/0907 (*S. subochreatum*) is highlighted (dark grey). *Spyridium phylloides* and *S. sp. Red Dots* (J.Kellermann 689) have been excluded from this map.

eastern Australian-dominated Clade M (PP = 1.0, UFBS = 100; Fig. 5). Of the 45 taxa that included more than one accession, nine (20.0%) were monophyletic and 30 (66.7%) were polyphyletic (Fig. 5, Table 3), two (4.4%) were paraphyletic (*S. parvifolium* with respect to *S. daltonii* in Clade O1 and *S. polycephalum* with respect to *S. oligocephalum* in Clade M6) and four (8.8%) were unresolved.

### Incongruence between the nrDNA and cpDNA phylogenies

Although the nrDNA and cpDNA trees included some nodes in common, there was substantial incongruence between the two phylogenies (Fig. 2, 5). This included differences in the order of divergence, in species relationships (within and between clades) and between the support for species circumscriptions (e.g. monophyly).

Key clades in common between the two trees were the clade of Tasmanian endemic species (nrDNA Clade D; cpDNA Clade K), and the clade of NSW endemic species (nrDNA Clade C; cpDNA Clade L); cpDNA Clade N also included most members of the WA endemic nrDNA Clade A, but excluding *S. globulosum*, *S. spadiceum* and one sample of *S. majoranifolium* that were placed in Clade M4 (Fig. 5). In the clade of NSW endemics, both trees showed similar relationships among species; however, within the Tasmanian and WA clades, relationships varied substantially.

Apart from this, there were many differences in the order of divergence between the two phylogenies, with these differences being particularly apparent at the base of each tree. For example, the base of the nrDNA phylogeny showed an early east–west split (Clade A–Clade B; Fig. 2), a split not supported by the cpDNA phylogeny (Fig. 5). The cpDNA phylogeny, instead, showed the Tasmanian (Clade K) and NSW (Clade L) endemics diverging first (Fig. 5).

There were many differences in species relationships between the two phylogenies, with this incongruence being particularly apparent from Clade M to Clade Q in the cpDNA phylogeny (Fig. 5). For example, *S. cinereum* is found in the *S. parvifolium*-dominated clade in the nrDNA phylogeny (Clade E; Fig. 2), but is nested within the SA endemic Clade M1 in the cpDNA tree (Fig. 5). In another example, *S. daphnoides* and *S. nitidum* were found to be

sister to *Spyridium* sp. Wollar (E.F.Constable s.n., NSW 16590) Kellermann and *S. eriocephalum* var. *eriocephalum* in the nrDNA phylogeny (Clades F and G), but were distantly related in the cpDNA tree (in Clades M1, M5 and P; Fig. 5).

There were also many differences relating to support for species circumscriptions between the two trees (Fig. 2, 5; Table 3). For example, *S. majoranifolium* was supported as monophyletic in the nrDNA phylogeny (PP = 1.0, UFBS = 100; clade A2) but found to be polyphyletic across Clades M5, N1 and N2 in the cpDNA tree. In another example, *S. sp. Kangaroo Island* (W.R.Barker 7560) was unresolved in the nrDNA phylogeny (Clade I; PP = 1.0, UFBS = 97; Fig. 2), but polyphyletic across Clades M2 and Q2 in the cpDNA tree (Fig. 5).

### Discussion

In this study, we have presented the most comprehensive molecular phylogeny of *Spyridium* to date, including all recognised species, subspecies and varieties (excluding *S. bifidum* var. *bifidum*), as well as six undescribed, phrase-name taxa. These results provide an understanding of biogeographic patterns within the genus, support some species circumscriptions, call others into question, and provide support for recognition of some currently undescribed taxa. However, incongruence between the nrDNA and cpDNA phylogenies means that some relationships within the genus were unable to be resolved by these results alone.

### Incongruencies: introgression or incomplete lineage sorting

Although the nrDNA and cpDNA phylogenies contained similarities, there were many differences between the two trees. These were apparent both in the relationships among major clades and when comparing implications for the monophyly, paraphyly or polyphyly of species and infra-specific taxa (Table 3). If the results of the cpDNA tree were viewed alone, as an indication of phylogenetic relationships, revising circumscriptions of half of the species in the genus might seem required. However, conversely, many of these polyphyletic taxa were found to be monophyletic or

unresolved in the nrDNA tree (Table 3), providing at least some support for, or not contradicting, most of the current circumscriptions of species.

Two key processes potentially leading to incongruence between nuclear and chloroplast gene trees are introgression, including chloroplast capture (Rieseberg and Soltis 1991; Tsitrone *et al.* 2003), and incomplete lineage sorting (ILS; Wiley and Lieberman 2011). Distinguishing the relative influence of these processes can be difficult, a problem commonly discussed in plant phylogenetic studies (e.g. Meudt and Bayly 2008; French *et al.* 2016; Barrett *et al.* 2018; Schuster *et al.* 2018). Other explanations for nuclear and chloroplast incongruence could include incomplete taxon sampling, which might result in artefacts of long branch attraction in one dataset over another; however, this is unlikely, given the almost complete taxon sampling in our study.

In our analyses, we have not attempted to distinguish potential instances of introgression from ILS, or the relative importance of each, but we expect, as outlined below, that both processes have potentially had a greater influence on the cpDNA results than on the nrDNA. As such, much of the species polyphyly reported in the cpDNA phylogeny might result from either chloroplast capture or ILS, and the nrDNA tree could better match the species tree for the genus.

In angiosperms, introgression of chloroplast genomes can be more apparent than nrDNA introgression (Rieseberg and Soltis 1991; Tsitrone *et al.* 2003). This is because chloroplasts are (generally) maternally inherited, not recombinant or subject to concerted evolution that can obscure signals of introgression in nrDNA (Álvarez and Wendel 2003), and have lower mutation rates (Palmer 1987), meaning that, in the absence of selective pressure or stochastic events, the signal of past cpDNA introgression is preserved. Potential introgression has been inferred in other studies of Rhamnaceae, including in *Pomaderris* (Nge *et al.* 2021c), *Discaria* Hook. (Aagesen *et al.* 2005; Medan *et al.* 2012) and *Ceanothus* L. (Hardig *et al.* 2002; Burge *et al.* 2013). One possible example of introgression contributing to incongruent results in the current study may be *S. sp.* Kangaroo Island (W.R.Barker 7560) Kellermann, which was unresolved in the nrDNA phylogeny (Clade I; Fig. 2), but was polyphyletic across disparate clades (M2 and Q2) in the cpDNA tree (Fig. 5). *Spyridium sp.* Kangaroo Island (W.R.Barker 7560) has an overlapping distribution, with several species found in the same subclades in the cpDNA phylogeny (e.g. *S. coalitum*, clade M2; and *S. thymifolium*, clade Q2); therefore, chloroplast capture through introgression is a possible explanation of this incongruence.

Another potential explanation for differences between cpDNA and nrDNA phylogenies is ILS. This occurs when the coalescence for alleles at a locus present in different species pre-dates the speciation event that gave rise to these species, and it can result in a gene tree that is incongruent with the species tree (Wiley and Lieberman 2011) or with other

gene trees. The usual expectation is that ILS has a greater confounding influence on phylogenetic analyses of nuclear data than of chloroplast data, because the effective population size of haploid chloroplast markers is only one quarter that of diploid nuclear markers, and this should lead to quicker coalescence of chloroplast sequences than of nuclear alleles (Birky *et al.* 1989). However, for nrDNA markers, as used in our study, the processes of concerted evolution (Arnheim 1983), which homogenise rDNA sequences within the genomes of organisms, can reduce the effective population size of rDNA markers relative to other nuclear sequences, by as much as 200-fold (Buckler and Holtsford 1996); this leads to substantially shorter coalescence times, and reduces the influence of ILS on nrDNA phylogenies.


For the reasons outlined above, in the following discussions of biogeography and taxonomy we place most emphasis on the nrDNA results but also incorporate additional details from cpDNA trees where relevant.

There is substantial scope to further explore the potential influence of introgression and ILS in the history of *Spyridium*. Multi-locus nuclear datasets, for instance, would be more amenable to statistical analyses of introgression (Joly *et al.* 2009; Joly 2012; García *et al.* 2017) than is nrDNA, and could give a stronger phylogenetic signal to assess incongruence between nuclear and plastid gene trees. Nonetheless, the data presented here highlight that nuclear and chloroplast incongruence is conspicuous in the genus.

## Broad biogeographic patterns in *Spyridium*

This study has provided the first comprehensive (all species) phylogenetic assessment of biogeographic patterns in *Spyridium*. Our inferences (below) are based on the molecular phylogenies and insight from other studies of biogeography in southern Australian. We focus on broad (continent scale) patterns in general terms. The work presented here could be extended by additional analyses, such as, for example, using dated trees or probabilistic biogeographic modelling (e.g. Ree and Smith 2008; Matzke 2013), although these are not explored here.

An early split in *Spyridium* has been identified in the nrDNA phylogeny, between the east and west of Australia across the Nullarbor Plain (Clades A and B; Fig. 2, 3a–j). There are many examples of east–west divergences in plants distributed across southern Australia, including *Phebalium* Vent. (Mole *et al.* 2004), *Eucalyptus* L'Hér. subgenus *Eucalyptus* (Ladiges *et al.* 2012), Goodeniaceae R.Br. (Jabaily *et al.* 2014), *Xanthorrhoea* Sol. ex Sm. (McLay *et al.* 2021), *Adenanthos* Labill. (Nge *et al.* 2021a) and *Pomaderris* (Nge *et al.* 2021c). The Nullarbor Plain disjunction in a range of plant groups has been related to vicariance associated with uplifting and climatic cooling in the mid-Miocene (Crisp and Cook 2007), including in *Pomaderris* (Nge *et al.* 2021c), a close relative of



**Fig. 4.** Distributions of samples of *S. phyllicoides*, *S. sp. Red Dots* (J.Kellermann 689) and *S. sp. Dwarf* (J.Kellermann 579) used in this study. For *S. phyllicoides*, samples are coloured by the clades in which they are placed in the nrDNA tree (Fig. 2), with the distribution of the species, on the basis of the records in the *Atlas of Living Australia* (2020), also shown (grey dots).

*Spyridium* in the tribe Pomaderreae. Nge *et al.* (2021c) concluded that *Pomaderris* was widespread throughout southern and eastern Australia until c. 14 Ma, when the Nullarbor Plain uplift occurred, with subsequent rapid ‘within region’ diversification in eastern Australia from c. 10 Ma, and little movement across biomes since. Although we did not use dated trees to determine diversification dates (as per Nge *et al.* 2021c), a similar explanation for the early east–west divergence in *Spyridium* could be inferred from our results.

Assuming the deep east–west divergence in *Spyridium* relates to formation of the Nullarbor Plain, the nrDNA tree suggests that up to three lineages in the genus have potentially dispersed across the plain subsequent to this early east–west divergence (Fig. 2, 3a, h, j). An east-to-west dispersal of *S. subochreatum* is inferred from both nrDNA and cpDNA trees, because the species is nested within eastern taxa and its distribution extends to just west of the Nullarbor Plain (Fig. 3j). Conversely, for *S. tricolor*, a west-to-east dispersal could be inferred, because the SA sample of this species (CC545; Fig. 3a) groups with western taxa. However, because it is sister to other western taxa in Clade A in the nrDNA phylogeny (Fig. 2), a reverse scenario could not be ruled out. Like *S. subochreatum*, *S. tricolor* grows on sandy soils and limestone (FloraBase – the Western Australian Flora, see <https://florabase.dpaw.wa.gov.au/>, accessed 7 May 2021), suggesting it could have made use of land connections south of the Nullarbor Plain that have been exposed at times of lower sea level since the late Pliocene (Nelson 1974; Wright and Ladiges 1997). An alternative explanation, that *S. tricolor* was widespread across this region and became disjunct during the Nullarbor Plain uplift, would require retention of morphological resemblance, such that it is recognised as a single species, despite a considerable geographic disjunction, for a substantial period of time since the mid-Miocene. Population-level sampling of *S. tricolor* using variable genomic markers could provide a greater insight into geographic history of the species and be used to further test its taxonomic circumscription.

The third lineage for which dispersal over the Nullarbor Plain might possibly be inferred is that including *S. polycephalum* and *S. oligocephalum* (Fig. 2, 3h, 5). A deep east-to-west dispersal event, before the diversification of the two species, is inferred from both the nrDNA and cpDNA trees, because this clade was nested within species from east of the Nullarbor Plain. Evidence of early east–west vicariance across southern Australia, followed by subsequent dispersal events such as these have been inferred in studies of other plant groups, such as, for example, in *Eucalyptus* subgenus *Eucalyptus* (Wright and Ladiges 1997), *Thelymitra* J.R.Forst. & G.Forst. (Nauheimer *et al.* 2018), *Calytrix tetragona* Labill. (Nge *et al.* 2021b) and *Pomaderris* (Nge *et al.* 2021c). Despite this, an alternative explanation of vicariance to account for the Western Australian distribution of the *S. polycephalum*–*S. oligocephalum* clade cannot be immediately discounted on the basis of our data. Although a vicariance explanation is less parsimonious because it would infer extinction of multiple lineages in western Australia, such reasoning assumes that multiple extinctions are less probable than is a single dispersal, which might not be true (Sanmartín and Meseguer 2016), for example, in the face of substantial climatic change in Australia since the mid-Miocene. A robust time-calibrated phylogeny for *Spyridium* could help corroborate one of these alternative scenarios.

Within the eastern Australian branch of the nrDNA phylogeny (Clade B, Fig. 2), an early NSW divergence is inferred (Fig. 3b). This deep divergence of NSW endemics from other south-eastern Australian taxa occurs near a broadly defined area that has been termed the southern transition zone (STZ; Fig. 3b) by Milner *et al.* (2012). The STZ is found east of the Great Dividing Range (GDR) and north of the Victoria–NSW border and is identified as a region where genetic or distributional discontinuities are seen in a range of taxa, but with the exact position of the discontinuities being dependent on habitat requirements of individual species and potentially different timescales of divergence (Milner *et al.* 2012). Other plant taxa showing genetic breaks across the STZ include *Hardenbergia violacea* (Schneev.) Stearn (Larcombe *et al.*




Fig. 5. (Caption on next page)

**Fig. 5.** Chloroplast genome (cpDNA) phylogeny of *Spyridium*, based on Bayesian inference (BI) analysis. Bayesian posterior probabilities (PP) <0.95 and ultrafast bootstrap (UFBS) values are shown at nodes when <95%; values ≥95% are not shown. Where one value for a node is supported (≥95%) and the other for that node is unsupported (<95%), only the unsupported value is shown. Where a hyphen (-) is provided at a node, this node varied in resolution in the ML tree and was therefore not transferable to the BI phylogeny. Coloured bar to the right of the tree indicates placement of samples in the nrDNA phylogeny (i.e. matching the coloured bar on Fig. 2). Labels are given for some clades (K–Q) and subclades (M1–Q2) discussed in text. Species polyphyletic across clades are highlighted in red text. Monophyletic taxa with >0.95 PP support are highlighted in green. Dashed lines associated with *S. tricolor*, *S. glaucum*, *S. phlebophyllum* and *S. subochreatum* E.D.Adams 21/0907 are provided as reference points connecting taxa to the sidebar.

2011), *Lomatia* R.Br. (Milner *et al.* 2012), *Callitris rhomboidea* R.Br. ex Rich. & A.Rich. (Worth *et al.* 2018) and *Xanthorrhoea* (McLay *et al.* 2021). *Spyridium* provides a further example of this pattern, although potential drivers of the divergence are unclear in this case.

Tasmanian endemics (excluding *S. obcordatum*) were found in a single, early diverging clade separate from their mainland counterparts in the nrDNA tree (Clade D, Fig. 2, 3c), suggesting early vicariance of *Spyridium* across Bass Strait. This early divergence and diversification of Tasmanian endemics is also supported by the cpDNA phylogeny (Clade K, Fig. 5) and the findings of Kellermann *et al.* (2005) and Hauenschild *et al.* (2018). The continued barriers to dispersal and gene-flow are likely to be the inundation of Bass Strait during interglacial periods (Galloway and Kemp 1981) and the semi-arid climate of the land-bridge exposed during glacial periods (Kirkpatrick and Fowler 1998). Major glacial and interglacial fluctuations occurred throughout the Quaternary (c. 2.2 Ma to c. 10 000 years ago; Hope 1994; Quilty 1994) and their resulting climatic extremes have been inferred to contribute to the limited distribution of narrow-range endemism in *Spyridium* in Tasmania (Coates and Kirkpatrick 1999).

Recent dispersal or gene flow between Victoria and Tasmania are here inferred for the lineage represented by *S. obcordatum*, the only endemic Tasmanian species not placed in Clades D or K (Fig. 2, 3d, 5), and several widespread taxa, including *S. eriocephalum* var. *eriocephalum*, *S. vexilliferum* var. *vexilliferum* and *S. parvifolium* (Fig. 2, 3d, f, j). Accessions of each of these widespread taxa collected from Tasmania (and Flinders Island for *S. parvifolium*) were found within the same clade as samples of the same taxa from Victoria (Table 1). Similar patterns of recent gene-flow between Victoria and Tasmania have been inferred in other plant groups, including *Eucalyptus globulus* Labill. (Freeman *et al.* 2001), *Hardenbergia violacea* (Larcombe *et al.* 2011), *Correa* Andrews (French *et al.* 2016), *Zieria veronicea* Sm. (Neal *et al.* 2019), *Xanthorrhoea* (McLay *et al.* 2021) and a range of other species (Worth *et al.* 2017). Evidence suggests that at least some areas of the Bassian Plain were covered in eucalypt woodland habitat (Hope 1978, 1994; Kirkpatrick and Fowler 1998) which could have been suitable for *S. parvifolium*, *S. vexilliferum* var. *vexilliferum* and *S. eriocephalum* var. *eriocephalum* (VicFlora 2018), i.e. potentially facilitating over-land rather than over-water dispersal between Victoria and Tasmania.

## Review of circumscriptions of species

The molecular phylogenies support the circumscriptions of several *Spyridium* species, but raise questions about others. A quarter of the taxa represented by more than one accession were identified as monophyletic in the nrDNA phylogeny, with approximately one-third being resolved as polyphyletic, and the remainder being unresolved (Fig. 2, Table 3). Of the monophyletic taxa resolved in the nrDNA tree, several were also found to be monophyletic in the cpDNA phylogeny (e.g. *S. obcordatum*, *S. scortechinii* and *S. montanum*), providing additional support for these circumscriptions of species (Fig. 5, Table 3). Of the polyphyletic taxa in the nrDNA phylogeny, two of the most notable were distributed across disparate clades, namely, *S. eriocephalum* and *S. phyllicoides* (Fig. 2). Given that both of these species were also resolved in separate clades in the cpDNA tree (Fig. 5), they are discussed in more detail below, along with several associated phrase-name taxa.

### *Spyridium eriocephalum*

*Spyridium eriocephalum* is polyphyletic and requires taxonomic revision, because its two varieties were found in distinct clades in both nrDNA and cpDNA phylogenies (Clades F and I, Fig. 2; Clades M1 and Q2, Fig. 5). *Spyridium eriocephalum* var. *eriocephalum* is monophyletic (albeit with limited sampling, but from separated localities) and geographically distinct from other taxa in the nrDNA tree (Fig. 2, 3f). *Spyridium eriocephalum* var. *glabrisepalum* is unresolved in a polytomy in the nrDNA phylogeny (Clade I, Fig. 3) with several other taxa (Fig. 2, 3i). The two varieties of *S. eriocephalum* are for the most part geographically distinct, with the exception being some overlap on Kangaroo Island (J. Kellermann, unpubl. data). The typical variety is widespread in south-eastern Australia (SA, Victoria, NSW and Tasmania), whereas var. *glabrisepalum* is restricted to Kangaroo Island. The two varieties are also morphologically distinguished by the presence of woolly sepal hairs (var. *eriocephalum*) versus hairless sepals that are instead glabrous-viscid (var. *glabrisepalum*; Canning 1986). Although the two taxa appear distinct (from each other) in both phylogenies, given that the two samples of var. *glabrisepalum* are placed with some other Kangaroo Island endemic taxa (e.g. *S. coalitum*) in the nrDNA tree, it

**Table 3.** Summary of resolution of taxa comparing the nrDNA phylogeny to the cpDNA trees.

Taxona	nrDNA phylogeny (Fig. 2)	cpDNA phylogeny (Fig. 5)	Congruent (in both trees)
<i>S. burragorang</i>	Unresolved	Unresolved	Unresolved
<i>S. buxifolium</i>	Unresolved	Monophyletic	–
<i>S. cinereum</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. coactilifolium</i>	Monophyletic	Unresolved	–
<i>S. coalitum</i>	Unresolved	Polyphyletic	–
<i>S. cordatum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. daphnoides</i>	Unresolved	Polyphyletic	–
<i>S. eriocephalum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. eriocephalum</i> var. <i>eriocephalum</i>	Monophyletic	Polyphyletic	–
<i>S. eriocephalum</i> var. <i>glabrisepalum</i>	Unresolved	Polyphyletic	–
<i>S. fontis-woodii</i>	Unresolved	Polyphyletic	–
<i>S. globulosum</i>	Unresolved	Polyphyletic	–
<i>S. gunnii</i>	Unresolved	Polyphyletic	–
<i>S. halmaturinum</i>	Unresolved	Polyphyletic	–
<i>S. lawrencei</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. majoranifolium</i>	Monophyletic	Polyphyletic	–
<i>S. microcephalum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. minutum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. montanum</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. mucronatum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. mucronatum</i> subsp. <i>mucronatum</i>	Unresolved	Unresolved	Unresolved
<i>S. mucronatum</i> subsp. <i>recurvum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. nitidum</i>	Unresolved	Polyphyletic	–
<i>S. obcordatum</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. obovatum</i>	Unresolved	Polyphyletic	–
<i>S. obovatum</i> var. <i>obovatum</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. obovatum</i> var. <i>velutinum</i>	Unresolved	Polyphyletic	–
<i>S. parvifolium</i>	Unresolved	Paraphyletic (with respect to <i>S. daltonii</i> )	–
<i>S. phylloides</i>	Polyphyletic	Polyphyletic	Polyphyletic

(Continued on next column)

**Table 3.** (Continued)

Taxona	nrDNA phylogeny (Fig. 2)	cpDNA phylogeny (Fig. 5)	Congruent (in both trees)
<i>S. polycephalum</i>	Unresolved	Paraphyletic (with respect to <i>S. oligocephalum</i> )	–
<i>S. riparium</i>	Monophyletic	Polyphyletic	–
<i>S. scabridum</i>	Unresolved	Polyphyletic	–
<i>S. scortechinii</i>	Monophyletic	Monophyletic	Monophyletic
<i>S. sp. Dwarf</i> (J. Kellermann 579)	Unresolved	Polyphyletic	–
<i>S. sp. Finnis</i> (J. Kellermann 653 & F. Nge)	Unresolved	Unresolved	Unresolved
<i>S. sp. Jerdacuttup</i> (A. Williams 332)	Unresolved	Monophyletic	–
<i>S. sp. Kangaroo Island</i> (W.R. Barker 7560)	Unresolved	Polyphyletic	–
<i>S. stenophyllum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. stenophyllum</i> subsp. <i>renovatum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. subochreatum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. ulicinum</i>	Monophyletic	Polyphyletic	–
<i>S. vexilliferum</i>	Unresolved	Polyphyletic	Polyphyletic
<i>S. vexilliferum</i> var. <i>latifolium</i>	Unresolved	Monophyletic	–
<i>S. vexilliferum</i> var. <i>vexilliferum</i>	Polyphyletic	Polyphyletic	Polyphyletic
<i>S. villosum</i>	Unresolved	Polyphyletic	–
Totals (45 taxa total)			
Monophyletic	11	9	6
Paraphyletic	0	2	1
Polyphyletic	11	30	12
Unresolved	23	4	4

Only taxa represented by more than one accession are included (=45 taxa). Only supported branches were considered when determining the resolution of the taxa (e.g. *S. gunnii* was unresolved in the nrDNA phylogeny because branches between the two accessions were not supported, resulting in an unresolved polytomy for the species). In the column 'Congruent (in both trees)', a dash (–) indicates that the resolution between the phylogenies was incongruent.

is possible that introgression may be influencing this placement (Fig. 2). However, placement of var. *glabrisepalum* in the cpDNA phylogeny is somewhat incongruent although supported, with samples being placed with accessions representing other taxa collected from a range of sites from SA to Tasmania (Fig. 5). Additional morphological or molecular work is recommended to re-assess these taxa and their relationships.

### *Spyridium phyllicoides*, S. sp. Dwarf (J.Kellermann 579) and S. sp. Red Dots (J.Kellermann 689)

*Spyridium phyllicoides* is polyphyletic and requires taxonomic revision. In both phylogenies, samples of *S. phyllicoides* were found in two clades (Clades I and J2, Fig. 2; Clades M1, M3 and Q1, Fig. 5). There is no biogeographic pattern to these (Fig. 4) and perhaps further unidentified forms exist within *S. phyllicoides*, in addition to the two forms already given phrase names, namely, *S. sp. Dwarf* (J.Kellermann 579) Kellermann and *S. sp. Red Dots* (J.Kellermann 689) Kellermann (Table 2). *Spyridium sp. Dwarf* (J.Kellermann 579) is distinguished from *S. phyllicoides* by generally smaller leaves and low-growing, almost prostrate habit (J. Kellermann, unpubl. data). However, the distribution of samples of *S. sp. Dwarf* (J.Kellermann 579; Fig. 5) overlaps with that of many species found in SA (Fig. 3f, g, i, j); therefore, it is possible that the incongruent result for this taxon (unresolved in the nrDNA phylogeny and polyphyletic in the cpDNA tree; Fig. 2, 5) may be attributed to chloroplast capture or ILS. Only one accession of *S. sp. Red Dots* (J.Kellermann 689) was included in this study and support for this taxon as distinct from *S. phyllicoides* in either phylogeny is limited (Clade J2, Fig. 2; Clade M3, Fig. 5). We recommend further investigation into both phrase-name taxa, particularly *S. sp. Dwarf* (J.Kellermann 579), and a more detailed investigation into the circumscription of *S. phyllicoides*.

### Conclusions

Here we have presented the first comprehensive phylogenies of the genus *Spyridium*, representing all described species and utilising both nrDNA and whole chloroplast genomes. Most incongruencies between the two trees could relate to introgression and chloroplast capture or ILS.

We found evidence of an early east–west split at the base of the nrDNA phylogeny and early diverging clades dominated by Tasmanian and NSW endemics. Our trees provide evidence of two subsequent within-species dispersal events across the Nullarbor Plain (*S. subochreatum* and *S. tricolor*) as well as a possible dispersal and diversification of a lineage including the south-western Australian endemics *S. polycephalum* and *S. oligocephalum* (although a vicariance explanation is also plausible in that case). In Tasmania, we

found *S. obcordatum* to be the result of a recent dispersal and subsequent diversification event, and evidence of recent gene-flow between Tasmania and Victoria in several wide-spread taxa (including *S. vexilliferum* var. *vexilliferum*).

Eleven taxa were supported as monophyletic in the nrDNA phylogeny and the following two were polyphyletic across disparate clades, requiring taxonomic review: *S. eriocephalum* (with two genetically distinct varieties) and *S. phyllicoides*.

### References

- Aagesen L, Medan D, Kellermann J, Hilger HH (2005) Phylogeny of the tribe Colletieae (Rhamnaceae) – a sensitivity analysis of the plastid region *trnL-trnF* combined with morphology. *Plant Systematics and Evolution* **250**, 197–214. doi:10.1007/s00606-004-0204-5
- Álvarez I, Wendel JF (2003) Ribosomal ITS sequences and plant phylogenetic inference. *Molecular Phylogenetics and Evolution* **29**, 417–434. doi:10.1016/S1055-7903(03)00208-2
- Arnheim N (1983) Concerted evolution of multigene families. In ‘Evolution of genes and proteins’. (Eds M Nei, R Koehn) pp. 38–61. (Sinauer: Sunderland, MA, USA)
- Atlas of Living Australia (2020) Occurrence records download on 2020-06-26: taxon\_name: ‘Spyridium’. Available at <https://doi.ala.org.au/doi/02d2462a-77d9-4f0b-9457-caf67834d693> [Verified 26 June 2020]
- Barker WR (2005) Standardising informal names in Australian publications. *Australian Systematic Botany Society Newsletter* **122**, 11–12.
- Barrett RA, Bayly MJ, Duretto MF, Forster PI, Ladiges PY, Cantrill DJ (2018) Phylogenetic analysis of *Zieria* (Rutaceae) in Australia and New Caledonia based on nuclear ribosomal DNA shows species polyphyly, divergent paralogues and incongruence with chloroplast DNA. *Australian Systematic Botany* **31**, 16–47. doi:10.1071/SB16034
- Bentham G (1863) Rhamnaceae. In ‘Flora Australiensis, a description of the plants of the Australian territory. Vol. 1: Ranunculaceae to Anacardiaceae’. pp. 409–445. (Lovell Reeve and Co: London, UK)
- Birky CW Jr, Fuerst P, Maruyama T (1989) Organelle gene diversity under migration, mutation, and drift: equilibrium expectations, approach to equilibrium, effects of heteroplasmic cells, and comparison to nuclear genes. *Genetics* **121**, 613–627. doi:10.1093/genetics/121.3.613
- Buckler ES IV, Holsford TP (1996) *Zea* systematics: ribosomal ITS evidence. *Molecular Biology and Evolution* **13**, 612–622. doi:10.1093/oxfordjournals.molbev.a025621
- Burge DO, Hopkins R, Tsai Y-HE, Manos PS (2013) Limited hybridization across an edaphic disjunction between the gabbro-endemic shrub *Ceanothus roderickii* (Rhamnaceae) and the soil-generalist *Ceanothus cuneatus*. *American Journal of Botany* **100**, 1883–1895. doi:10.3732/ajb.1200604
- Canning EM (1986) *Spyridium*. In ‘Flora of South Australia’, Vol. 2. (Eds JP Jessop, HR Toelken) (Government Printer: Adelaide, SA, Australia)
- CHAH (2020) *Spyridium* Fenzl. In ‘Australian Plant Census’. (Council of Heads of Australian Herbaria) Available at [https://biodiversity.org.au/nsi/services/search?product=APNI&tree.id=&name=Spyridium&inc\\_scientific=&inc\\_cultivar=&inc\\_other=&max=100&display=&search=true](https://biodiversity.org.au/nsi/services/search?product=APNI&tree.id=&name=Spyridium&inc_scientific=&inc_cultivar=&inc_other=&max=100&display=&search=true) [Verified 17 November 2020]
- Clowes C, Fowler RM, Brown GK, Bayly MJ (2018) The complete chloroplast genome sequence of *Spyridium parvifolium* var. *parvifolium* (family Rhamnaceae; tribe Pomaderreae). *Mitochondrial DNA. Part B, Resources* **3**, 807–809. doi:10.1080/23802359.2018.1483776
- Coates F, Kirkpatrick JB (1999) Is geographic range correlated with climatic range in Australian *Spyridium* taxa? *Australian Journal of Botany* **47**, 755–767. doi:10.1071/BT97066
- Crisp MD, Cook LG (2007) A congruent molecular signature of vicariance across multiple plant lineages. *Molecular Phylogenetics and Evolution* **43**, 1106–1117. doi:10.1016/j.ympev.2007.02.030
- Department of Agriculture Water and the Environment (2020) Interim Biogeographic Regionalisation for Australia (Regions – States and Territories) v. 7 (IBRA). Available at <https://www.environment.gov.au/fed/catalog/search/resource/details.page?uuid=%7B1273FBE2-F266-4F3F-895D-C1E45D77CAF5%7D> [Verified 26 July 2020]



- Doyle J, Doyle JL (1987) Genomic plant DNA preparation from fresh tissue-CTAB method. *Phytochemical Bulletin* **19**, 11–15.
- Freeman JS, Jackson HD, Steane DA, McKinnon GE, Dutkowski GW, Potts BM, Vaillancourt RE (2001) Chloroplast DNA phylogeography of *Eucalyptus globulus*. *Australian Journal of Botany* **49**, 585–596. doi:10.1071/BT00094
- French PA, Brown GK, Bayly MJ (2016) Incongruent patterns of nuclear and chloroplast variation in *Correa* (Rutaceae): introgression and biogeography in south-eastern Australia. *Plant Systematics and Evolution* **302**, 447–468. doi:10.1007/s00606-016-1277-7
- Galloway RW, Kemp EM (1981) Late Cainozoic environments of Australia. In 'Ecological Biogeography of Australia', Vol. 1. (Ed. A Keast) pp. 53–80. (Junk: The Hague, Netherlands)
- García N, Folk RA, Meerow AW, Chamala S, Gitzendanner MA, de Oliveira RS, Soltis DE, Soltis PS (2017) Deep reticulation and incomplete lineage sorting obscure the diploid phylogeny of rain-lilies and allies (Amaryllidaceae tribe Hippeastreae). *Molecular Phylogenetics and Evolution* **111**, 231–247. doi:10.1016/j.ympev.2017.04.003
- Hardig TM, Soltis PS, Soltis DE, Hudson RB (2002) Morphological and molecular analysis of putative hybrid speciation in *Ceanothus* (Rhamnaceae). *Systematic Botany* **27**, 734–746.
- Hauenschild F, Matuszak S, Muellner-Riehl AN, Favre A (2016) Phylogenetic relationships within the cosmopolitan buckthorn family (Rhamnaceae) support the resurrection of *Sarcomphalus* and the description of *Pseudoziziphus* gen. nov. *Taxon* **65**, 47–64. doi:10.12705/651.4
- Hauenschild F, Favre A, Michalak I, Muellner-Riehl AN (2018) The influence of the Gondwanan breakup on the biogeographic history of the ziziphoids (Rhamnaceae). *Journal of Biogeography* **45**, 2669–2677. doi:10.1111/jbi.13448
- Hope GS (1978) The late Pleistocene and Holocene vegetational history of Hunter Island, north-western Tasmania. *Australian Journal of Botany* **26**, 493–514. doi:10.1071/BT9780493
- Hope GS (1994) Quaternary vegetation. In 'History of the Australian vegetation: Cretaceous to recent'. (Ed. GS Hope) pp. 368–389. (University of Adelaide Press: Adelaide, SA, Australia)
- Jabaily RS, Shepherd KA, Gardner AG, Gustafsson MHG, Howarth DG, Motley TJ (2014) Historical biogeography of the predominantly Australian plant family Goodeniaceae. *Journal of Biogeography* **41**, 2057–2067. doi:10.1111/jbi.12363
- Jobs DV, Thien LB (1997) A conserved motif in the 5.8 S ribosomal RNA (rRNA) gene is a useful diagnostic marker for plant internal transcribed spacer (ITS) sequences. *Plant Molecular Biology Reporter* **15**, 326–334. doi:10.1023/A:1007462330699
- Joly S (2012) JML: testing hybridization from species trees. *Molecular Ecology Resources* **12**, 179–184. doi:10.1111/j.1755-0998.2011.03065.x
- Joly S, McLenachan PA, Lockhart PJ (2009) A statistical approach for distinguishing hybridization and incomplete lineage sorting. *American Naturalist* **174**, E54–E70. doi:10.1086/600082
- Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermini LS (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587–589. doi:10.1038/nmeth.4285
- Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Research* **30**, 3059–3066. doi:10.1093/nar/gkf436
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* **30**, 772–780. doi:10.1093/molbev/mst010
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C (2012) Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647–1649. doi:10.1093/bioinformatics/bts199
- Kellermann J (2007) Re-instatement of the name *Spyridium waterhousei* from Kangaroo Island, South Australia, with a short history of the tribe Pomaderreae (Rhamnaceae). *Journal of the Adelaide Botanic Gardens* **21**, 55–62.
- Kellermann J (2021) The importance of the 'h' – parahomonymy in *Trymalium* (Rhamnaceae: Pomaderreae). *Swainsona* **35**, 23–29.
- Kellermann J, Barker WR (2012) Revision of the *Spyridium bifidum*-*S. halmatrinum* complex (Rhamnaceae: Pomaderreae) from South Australia and Victoria. *Muelleria* **30**, 26–58.
- Kellermann J, Udovicic F (2007) Large indels obscure phylogeny in analysis of chloroplast DNA (*trnL-F*) sequence data: Pomaderreae (Rhamnaceae) revisited. *Telopea* **12**, 1–22.
- Kellermann J, Udovicic F, Ladiges PY (2005) Phylogenetic analysis and generic limits of the tribe Pomaderreae (Rhamnaceae) using internal transcribed spacer DNA sequences. *Taxon* **54**, 619–631. doi:10.2307/25065419
- Kellermann J, Rye BL, Thiele KR (2008) Nomenclatural notes, typifications and name changes in *Trymalium* (Rhamnaceae: Pomaderreae). *Transactions of the Royal Society of South Australia* **132**, 18–28. doi:10.1080/03721426.2008.10887089
- Kellermann J, Thiele KR, Udovicic F, Walsh NG (2022) Rhamnaceae. In 'Flora of Australia'. (Ed. PG Kodala) (Australian Biological Resources Study: Canberra, ACT, Australia) Available at <https://profiles.ala.org.au/opus/foa/profile/Rhamnaceae> [Verified 10 May 2022]
- Kirkpatrick JB, Fowler M (1998) Locating likely glacial forest refugia in Tasmania using palynological and ecological information to test alternative climatic models. *Biological Conservation* **85**, 171–182. doi:10.1016/S0006-3207(97)00144-4
- Ladiges PY, Bayly MJ, Nelson G (2012) Searching for ancestral areas and artifactual centers of origin in biogeography: with comment on east-west patterns across southern Australia. *Systematic Biology* **61**, 703–708. doi:10.1093/sysbio/sys005
- Larcombe MJ, McKinnon GE, Vaillancourt RE (2011) Genetic evidence for the origins of range disjunctions in the Australian dry sclerophyll plant *Hardenbergia violacea*. *Journal of Biogeography* **38**, 125–136. doi:10.1111/j.1365-2699.2010.02391.x
- Matzke NJ (2013) Probabilistic historical biogeography: new models for founder-event speciation, imperfect detection, and fossils allow improved accuracy and model-testing. *Frontiers of Biogeography* **5**, 242–247. doi:10.21425/F55419694
- McLay TG (2017) High quality DNA extraction protocol from recalcitrant plant tissue. Available at <https://www.protocols.io/view/high-quality-dna-extraction-protocol-from-recalcit-i8jchun> [Verified 12 July 2018]
- McLay TGB, Ladiges PY, Doyle SR, Bayly MJ (2021) Phylogeographic patterns of the Australian grass trees (*Xanthorrhoea* Asphodelaceae) shown using targeted amplicon sequencing. *Australian Systematic Botany* **34**, 206–225. doi:10.1071/SB20013
- Medan D, Arbetman M, Chaia EE, Premoli AC (2012) Interspecific and intergeneric hybridization in south American Rhamnaceae-Colletieae. *Plant Systematics and Evolution* **298**, 1425–1435. doi:10.1007/s00606-012-0646-0
- Meudt HM, Bayly MJ (2008) Phylogeographic patterns in the Australasian genus *Chionohebe* (*Veronica* s.l., Plantaginaceae) based on AFLP and chloroplast DNA sequences. *Molecular Phylogenetics and Evolution* **47**, 319–338. doi:10.1016/j.ympev.2007.12.019
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In 'Proceedings of the Gateway Computing Environments Workshop (GCE)', 14 November 2010, New Orleans, LA, USA. INSPEC Accession Number 11705685. (IEEE) doi:10.1109/GCE.2010.5676129
- Milner ML, Rossetto M, Crisp MD, Weston PH (2012) The impact of multiple biogeographic barriers and hybridization on species-level differentiation. *American Journal of Botany* **99**, 2045–2057. doi:10.3732/ajb.1200327
- Mole BJ, Udovicic F, Ladiges PY, Duretto MF (2004) Molecular phylogeny of *Phebalium* (Rutaceae: Boronieae) and related genera based on the nrDNA regions ITS 1 + 2. *Plant Systematics and Evolution* **249**, 197–212. doi:10.1007/s00606-004-0218-z
- Nauheimer L, Schley RJ, Clements MA, Micheneau C, Nargar K (2018) Australasian orchid biogeography at continental scale: molecular phylogenetic insights from the sun orchids (*Thelymitra*, Orchidaceae). *Molecular Phylogenetics and Evolution* **127**, 304–319. doi:10.1016/j.ympev.2018.05.031
- Neal WC, James EA, Bayly MJ (2019) Phylogeography, classification and conservation of pink zieria (*Zieria veronicea*; Rutaceae): influence of changes in climate, geology and sea level in south-eastern Australia. *Plant Systematics and Evolution* **305**, 503–520. doi:10.1007/s00606-019-01589-z
- Nelson EC (1974) Disjunct plant distributions on the south-western Nullarbor Plain, Western Australia. *Journal of the Royal Society of Western Australia* **57**, 105–117.

- Nge FJ, Biffin E, Thiele KR, Waycott M (2021a) Reticulate evolution, ancient chloroplast haplotypes, and rapid radiation of the Australian plant genus *Adenanthos* (Proteaceae). *Frontiers in Ecology and Evolution* **8**, 616741.
- Nge FJ, Biffin E, Waycott M, Thiele KR (2021b) Phylogenomics and continental biogeographic disjunctions: insight from the Australian starflowers (*Calytrix*). *American Journal of Botany* **109**, 291–308. doi:10.1002/ajb2.1790.
- Nge FJ, Kellermann J, Biffin E, Waycott M, Thiele KR (2021c) Historical biogeography of Pomaderris (Rhamnaceae): continental vicariance in Australia and repeated independent dispersals to New Zealand. *Molecular Phylogenetics and Evolution* **158**, 107085. doi:10.1016/j.ympev.2021.107085
- Nguyen L-T, Schmidt HA, Von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **32**, 268–274. doi:10.1093/molbev/msu300
- Palmer JD (1987) Chloroplast DNA evolution and biosystematic uses of chloroplast DNA variation. *American Naturalist* **130**, S6–S29. doi:10.1086/284689
- Perrin D (2018) 'Dictionary of botanical names: Australian plant names, meanings, derivation and application.' (JT Press)
- Quilty PG (1994) The background: 144 million years of Australian palaeoclimate and palaeogeography. In 'History of the Australian vegetation: Cretaceous to recent'. (Ed. RS Hill) Vol. 14, pp. 14–43. (University of Adelaide Press: Adelaide, SA, Australia)
- Ree RH, Smith SA (2008) Maximum likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. *Systematic Biology* **57**, 4–14. doi:10.1080/10635150701883881
- Richardson JE, Chatrou LW, Mols JB, Erkens RHJ, Pirie MD (2004) Historical biogeography of two cosmopolitan families of flowering plants: annonaceae and Rhamnaceae. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **359**, 1495–1508. doi:10.1098/rstb.2004.1537
- Rieseberg LH, Soltis DE (1991) Phylogenetic consequences of cytoplasmic gene flow in plants. *Evolutionary Trends in Plants* **5**, 65–84.
- Rohland N, Reich D (2012) Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Research* **22**, 939–946. doi:10.1101/gr.128124.111
- Ronquist F, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**, 1572–1574. doi:10.1093/bioinformatics/btg180
- Sanmartín I, Meseguer AS (2016) Extinction in phylogenetics and biogeography: from timetrees to patterns of biotic assemblage. *Frontiers in Genetics* **7**, 35. doi:10.3389/fgene.2016.00035
- Schuster TM, Setaro SD, Tibbitts JFG, Batty EL, Fowler RM, McLay TGB, Wilcox S, Ades PK, Bayly MJ (2018) Chloroplast variation is incongruent with classification of the Australian bloodwood eucalypts (genus *Corymbia*, family Myrtaceae). *PLoS One* **13**, e0195034. doi:10.1371/journal.pone.0195034
- Shepherd LD, McLay TGB (2011) Two micro-scale protocols for the isolation of DNA from polysaccharide-rich plant tissue. *Journal of Plant Research* **124**, 311–314. doi:10.1007/s10265-010-0379-5
- Stöver BC, Müller KF (2010) TreeGraph 2: combining and visualizing evidence from different phylogenetic analyses. *BMC Bioinformatics* **11**, 7. doi:10.1186/1471-2105-11-7
- Thiele KR, West JG (2004) *Spyridium burragorang* (Rhamnaceae), a new species from New South Wales, with new combinations for *Spyridium buxifolium* and *Spyridium scortechinii*. *Telopea* **10**, 823–829.
- Tsitrona A, Kirkpatrick M, Levin DA (2003) A model for chloroplast capture. *Evolution* **57**, 1776–1782. doi:10.1111/j.0014-3820.2003.tb00585.x
- VicFlora (2018) *Spyridium*. In 'Flora of Victoria'. (Royal Botanic Gardens Foundation Victoria) Available at <https://vicflora.rbg.vic.gov.au/flora/taxon/829f1202-25c7-429c-97f7-fc7a9ba62a2d> [Verified 23 April 2018]
- Wiley EO, Lieberman BS (2011) 'Phylogenetics: theory and practice of phylogenetic systematics.' (Wiley: Hoboken, NJ, USA)
- Wilkins C, Kern S, Rathborne D, Markey A (2011). 'Rare and poorly known flora of the Ravenhorpe Range and Bandalup Hill.' (Government of Western Australia: Perth, WA, Australia)
- Worth JRP, Holland BR, Beeton NJ, Schönfeld B, Rossetto M, Vaillancourt RE, Jordan GJ (2017) Habitat type and dispersal mode underlie the capacity for plant migration across an intermittent seaway. *Annals of Botany* **120**, 539–549. doi:10.1093/aob/mcx086
- Worth JRP, Sakaguchi S, Harrison PA, Brüniche-Olsen A, Janes JK, Crisp MD, Bowman DMJS (2018) Pleistocene divergence of two disjunct conifers in the eastern Australian temperate zone. *Biological Journal of the Linnean Society* **125**, 459–474. doi:10.1093/biolinnean/bly127
- Wright IJ, Ladiges PY (1997) Geographic variation in *Eucalyptus diversifolia* (Myrtaceae) and the recognition of new subspecies *E. diversifolia* subsp. *hesperia* and *E. diversifolia* subsp. *megacarpa*. *Australian Systematic Botany* **10**, 651–680. doi:10.1071/SB96019

**Data availability.** The data that support this study are available in TreeBASE at <https://www.treebase.org/treebase-web/home.html> (Accession number 28815).

**Conflicts of interest.** Michael J. Bayly is an Associate Editor for *Australian Systematic Botany*. Despite this relationship, he did not at any stage have Associate Editor-level access to this manuscript while in peer review, as is the standard practice when handling manuscripts submitted by an editor to this journal. *Australian Systematic Botany* encourages its editors to publish in the journal and they are kept totally separate from the decision-making process for their manuscripts. The authors declare that they have no further conflicts of interest.

**Declaration of funding.** This project was supported by a Holsworth Wildlife Research Endowment administered by The Ecological Society of Australia, and the Hansjörg Eichler Scientific Research Fund provided by the Australasian Systematic Botany Society. Support was also provided by a Megan Klemm Postgraduate Research Award, a Sophie Ducker Postgraduate Scholarship (both administered by The University of Melbourne Botany Foundation) and a Dame Margaret Blackwood Soroptimist Scholarship (managed by Soroptimist International). The revision of the family Rhamnaceae for the *Flora of Australia* (Kellermann *et al.* 2022) has been supported by several grants from the Australian Biological Resources Study (Canberra, ACT, Australia).

**Acknowledgements.** We thank Matthew Dell, Rose Barrett, Mark Wapstra, Stephen Bell, Libby Sandiford, Gillian Craig, Kenneth Mills, John Hosking and Daniel Murphy for collecting silica samples and vouchers. Thanks go to Barbara Rye for advice related to WA *Spyridium*. We acknowledge the Western Australian Herbarium for providing destructive samples for molecular analysis. Thanks go to Todd McLay, Erin Batty and Stephen Wilcox for assistance with molecular work, and Matthew Dell and Francis Nge for general support for the project. We appreciate the assistance Joanne Birch, Will Neal and Margaret Brookes have provided while lodging samples in the University of Melbourne Herbarium. We thank Andrew Drinnan, Tanja Schuster and Peter Vesik for providing suggestions related to the project direction during committee meetings. Finally, we thank the Halls Gap Outdoor Community Facebook Group for helping retrieve a lost field book.

#### Author affiliations

<sup>A</sup>School of Biosciences, The University of Melbourne, Vic. 3010, Australia.

<sup>B</sup>State Herbarium of South Australia, Botanic Gardens and State Herbarium, Adelaide, SA 5000, Australia.

<sup>C</sup>School of Biological Sciences, The University of Adelaide, Adelaide, SA 5005, Australia.

<sup>D</sup>Department of Environment and Science, Queensland Herbarium, Toowong, Qld 4066, Australia.