

Species Composition and Community Structure of Dung Beetles Attracted to Dung of Gaur and Elephant in the Moist Forests of South Western Ghats

Authors: Vinod, K.V., and Sabu, Thomas K.

Source: Journal of Insect Science, 7(56) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.007.5601>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Species composition and community structure of dung beetles attracted to dung of gaur and elephant in the moist forests of South Western Ghats

K.V. Vinod^b and Thomas K. Sabu^a

Litter Entomology Research Unit, P.G. & Research Department of Zoology, St. Joseph's College, Devagiri, Calicut, Kerala, India 673008

Abstract

The community structure of dung beetles attracted to dung of gaur, *Bos gaurus* (H. Smith) (Artiodactyla: Bovidae) and Asian elephant, *Elephas maximus* Linnaeus (Proboscidea: Elephantidae), is reported from the moist forests of Western Ghats, in South India. The dominance of dwellers over rollers, presence of many endemic species, predominance of regional species and higher incidence of the old world roller, *Ochicanthon laetum*, make the dung beetle community in the moist forests of the region unusual. The dominance of dwellers and the lower presence of rollers make the functional guild structure of the dung beetle community of the region different from assemblages in the moist forests of south East Asia and Neotropics, and more similar to the community found in Ivory Coast forests. The ability of taxonomic diversity indices to relate variation in dung physical quality with phylogenetic structure of dung beetle assemblage is highlighted. Comparatively higher taxonomic diversity and evenness of dung beetle assemblage attracted to elephant dung rather than to gaur dung is attributed to the heterogeneous nature of elephant dung. Further analyses of community structure of dung beetles across the moist forests of Western Ghats are needed to ascertain whether the abundance of dwellers is a regional pattern specific to the transitional Wayanad forests of south Western Ghats.

Keywords: *Caccobius gallinus*, *Drepanocerus setosus*, *Liatongus indicus*, *Onthophagus lemniscatus*, *Onthophagus andreweisi*, *Onthophagus madoqua*, *Onthophagus devagiriensis*, *Onthophagus ensifer*, *Onthophagus castetsi*, *Onthophagus elongates*, *Onthophagus vladimiri*, *Onthophagus laevis*, *Onthophagus brutus*, *Ochicanthon laetum*, *Sisyphus neglectes*, *Elephas maximus*, *Bos gaurus*

Correspondence: ^asabukthomas@gmail.com, ^bvinukv@gmail.com

Received: 8 May 2006 | **Accepted:** 9 March 2007 | **Published:** 14 November 2007

Copyright: This is an open access paper. We use the Creative Commons Attribution 2.5 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Volume 7, Number 56

Cite this paper as:

Vinod KV, Sabu TK. 2007. Species composition and community structure of dung beetles attracted to dung of gaur and elephant in the moist forests of South Western Ghats. 14pp. *Journal of Insect Science* 7:56, available online: insectscience.org/7.56

Introduction

Dung beetles are a conspicuous component of the diversity of insects in Afrotropical rain forests (Hanski 1983; Hanski and Cambefort 1991; Hanski and Krikken 1991; Davis 2000a; Escobar 2000; Feer 2000; Estrada and Coates-Estrada 2002; Scheffler 2005). They use dung produced by forest vertebrates, particularly mammals and occasionally that of birds and reptiles (Howden and Young 1981; Hanski and Cambefort 1991; Estrada and Coates-Estrada 2002; Krell et al. 2003) as food and as a substrate for oviposition (Halffter and Edmonds 1982; Hanski 1989; Gill 1991). The presence of a variety of dung-producing mammals has effects on the relative abundance and diversity of dung beetles (Cambefort and Walter 1991; Hanski 1991; Estrada et al. 1999). Since, such resources can be extremely patchy in space and time, resource partitioning and competition between co-occurring species plays a major role in structuring dung beetle communities (Hanski 1991; Feer and Pincebourde 2005). Based on their nesting strategies, dung beetles are divided broadly into three functional groups *viz.*, rollers (telecoprid nesters), tunnelers (paracoprid nesters) and dwellers (endocoprid nesters) (Cambefort and Hanski 1991). Rollers form food balls from a dung pat, which are rolled away, build a tunnel and bury it for use in feeding and breeding. Tunnelers create underground chambers beneath dung pat and construct nests using dung from the pat whereas dwellers breed in the dung pat itself. This functional stratification allows dung beetles to minimize the intense competition for limited food and space and also to protect the food from adverse environmental conditions (Halffter and Edmonds 1982; Cambefort and Hanski 1991).

Dung beetles have a variety of effects on the ecosystem. By burying dung and carrion as food for their offspring, dung beetles may increase the rate of soil nutrient cycling (Halffter and Mathews 1966; Bornemissa and Williams 1970; Nealis 1977) and reduce egg and larval populations of parasitic flies present in fresh dung of mammals (Bergstrom et al. 1976). Many act as important secondary dispersal agents for seeds of several tree species defecated by frugivorous vertebrates, thus participating in the natural process of forest regeneration (Estrada and Coates-Estrada 1991; Feer 1999; Vulinec 2000; Andresen 2001, 2002, 2003, 2006; Andresen and Levey 2004). In addition, they are good indicators of the impact of

large herbivore and human induced change in forest habitats (Howden and Nealis 1975; Klein 1989; Favila and Halffter 1997; Davis et al. 2000; Davis 2000b; Davis et al. 2001; McGoch et al. 2002; van Rensburg et al. 1999; Davis et al. 2004; Botes et al. 2006).

Organization of dung beetle communities is very sensitive to changes in abundance of food resources, vegetation structure, microclimatic variables and soil characteristics (Nealis 1977; Halffter et al. 1992; Lumaret et al. 1992; Osberg et al. 1994; Davis 1996; Estrada et al. 1999; Escobar 2000; Davis 2002). Changes in community organization of dung beetles include changes in species richness, species composition, abundance and guild structure (*e.g.*, according to their diet and their resource-relocation behavior). Dung beetle communities are strongly influenced by dung type and they change in relation to the availability of different dung types (Lumaret et al. 1992; Davis 1994; Davis 2002). Though many dung beetles are generalists and do not show any dung preferences, some are strict specialists with some, or various, degrees of specialization. Some dung beetles preferably select coarse fibred dung of non-ruminants, while others prefer the more fluid and fine dung of ruminants, or the odoriferous dung of omnivores (Davis 1994; Davis 2002; Holter et al. 2002; Krell et al. 2003). Dung of howler and woolly monkeys (*Alouatta* spp.; *Lagothrix* sp.) and elephants is the preferred resource for several dung beetle species (Howden and Young 1981; Peck and Forsyth 1982; Halffter and Edmonds 1982; Cambefort and Walter 1991; Estrada and Coates-Estrada 1991; Estrada et al. 1993; Estrada et al. 1999).

Structure of vegetation is believed to be another main factor determining the organization of dung beetle communities in tropical rainforests (Hanski and Cambefort 1991; Escobar 1997; Hill 1996; Davis and Sutton 1998; Davis et al. 2000; Halffter and Arellano 2002; Scheffler 2005). From African savannahs to Neotropical forests, dung beetles are highly habitat specific and there are distinct guilds of beetles associated with forests, edges and pasture habitats. Although some species can utilize more than a single habitat type, certain species may never be found outside their preferred habitat (Scheffler 2002).

However, in total contrast to the well documented data on the composition, community structure and habitat preference of dung beetle communities from forests of Afrotropical regions,




Figure 1. (a) Map of India showing the location of Western Ghats, (b) Western Ghats and (c) study site in Wayanad region of Western Ghats.

there exist no records of the dung beetle communities of moist forests in Western Ghats despite the fact that it is a global hot spot of biodiversity (Mayers et al. 2000; Bossuyt et al. 2004). The Western Ghats is the only tropical forest ecoregion of the Indian peninsula and is well known for regional variation in vegetation, rainfall patterns, topography and high levels of endemism across its entire stretch. (Nair 1991; WWF 2001). Though mammalian diversity is lower here than in other tropical hotspots, moist forests of the region support important populations of many endemic and non-endemic mammalian species displaying different degrees of feeding habits (WWF 2001), adding opportunities for the coexistence of various dung beetle species. The Asian elephant, *Elephas maximus* Linnaeus (Proboscidea: Elephantidae) and gaur, *Bos gaurus* (H. Smith) (Artiodactyla: Bovidae), are the major mega-mammalian herbivores in the moist forests of Western Ghats (Joy 1991; Sukumar 2003; WWF 2001). The main goal of this study is to gain knowledge of the composition and guild structure of the dung beetle community attracted to gaur dung in a well protected moist forest area in Western Ghats, and to compare its community structure with the beetle assemblage attracted to dung of the Asian elephant, the other major megaherbivore in the region about which data is available from earlier studies (Sabu et al. 2006).

Materials and Methods

Study area

The study was carried out in Thirunelly forests (900m amsl, 20.55 km²) (11° 53' N latitude and 76° 01' E longitude), 100 km North of Calicut, Kerala state (Figure 1), located in the northern boundary of south Western Ghats (WG) forests in Wayanad in the Nilgiri Biosphere region [5520.4 km²]. Biogeographically, Wayanad is a transition area between the moist and dry deciduous forests in south Western Ghats moist deciduous ecoregion. It harbors habitat restricted, endemic species as well as disjunct populations of species that are found in both regions (Pascal 1988; Rodgers and Panwar 1988; WWF 2001). Moist forests of the region are the summer refuge for herds of elephants and gaurs from the dry eastern side as the open grasslands, and streams originating from the upper ranges, together with the abundance of bamboo culms (*Bambusa sp.*) provide a wide choice of resource materials for grazers and browsers (Joy 1991; Nair 1991; WWF 2006). Temperature varies annually between 24–32°C. Relative humidity is in the range of 40–80%. Rainfall averages between 3,000 and 3,250 mm per year and occurs mostly in the wet months of June to November. June, July and August have the most rain (KSEB rainfall data 2002–2004). Occasional summer showers occur in April and May. Topographic variation is moderate with hills rising gently from the lower river valleys and slopes reaching 35–60°.




Figure 2. Dendrogram showing phylogenetic relationship of dung beetles attracted to gaur (A) and elephant (B) dung in the moist forests of south Western Ghats in Wayanad region.

Methodology

The study was conducted after the rainy season between November 25, 2003 to February 2, 2004. Dung beetles were collected using Cebo-Suspendido-Rejilla pit fall traps (Lobo et al. 1988; Veiga et al. 1989; Errouissi et al. 2004). Each trap consisted of a plastic basin (210 mm in diameter and 150 mm in depth), buried to its rim in soil and containing a water-formalin-liquid soap mixture. One litre of fresh dung was placed on a strip of wire grid (2.5 cm x 2.5 cm) at the top of the basin. Each trap was topped with a dark plastic plate supported on iron bars to prevent desiccation and inundation during periods of rain. A set of four replicate traps, with each replicate at each corner of a 100 m² plot was placed in the

study site. The traps were collected after one week of exposure and sampling was repeated 15 times (4 traps x 15 samples). Earlier studies on the succession pattern of dung beetles showed that dung pats that were 3–7 days older attracted a subset of species that were not attracted to fresh dung (Sabu et al. 2006; personal field observation) and the gaur and elephant dung pats in the humid study region remained moist and wet for 5–7 days. Hence, sample retrieval and bait replacement was done at weekly intervals. Beetles were identified to species level using Arrow (1931) and Balthasar (1963 a, b). Identification of specimens was done by the authors and confirmed with the assistance of specialists (see acknowledgements). Beetles measuring ≥ 13 mm were considered as large (Cambefort 1991).

Table 1. Abundance and guild structure of dung beetles attracted to gaur dung in the moist forests of Wayanad in Western Ghats.

Sl No.	Species	Nesting guild	Mean body length (mm)	Size	Abundance
1	<i>Ochicanthon laetum</i> Boucomont	R	4.9 ± 0.23	S	44
2	<i>Sisyphus neglectus</i> Gory	R	7.1 ± 0.32	S	83
3	<i>Drepanocerus setosus</i> Arrow	D	5.1 ± 0.35	S	246
4	<i>Liatonqus indicus</i> Arrow	D	10.5 ± 0.65	S	136
5	<i>Oniticellus cinctus</i> Fabricius	D	11.5 ± 0.76	S	7
6	<i>Caccobius gallinus</i> Arrow*	T	4.7 ± 0.38	S	35
7	<i>Caccobius meridionalis</i> Boucomont	T	4.2 ± 0.32	S	120
8	<i>Caccobius unicornis</i> Fabricius	T	2.9 ± 0.25	S	4
9	<i>Catharsius sagax</i> Quenstedt	T	34.2 ± 1.48	L	5
10	<i>Copris davisoni</i> Waterhouse	T	13.6 ± 0.55	L	28
11	<i>Copris repertus</i> Walker	T	21.3 ± 0.64	L	51
12	<i>Onitis subopacus</i> Arrow	T	21.1 ± 0.66	L	39
13	<i>Onitis virens</i> Lansberge	T	21.9 ± 1.15	L	35
14	<i>Onthophagus rectecornutus</i> Lansberge	T	9.2 ± 0.48	S	144
15	<i>Onthophagus bronzeus</i> Arrow	T	13.3 ± 0.67	L	106
16	<i>Onthophagus bifasciatus</i> Fabricius	T	6.6 ± 0.43	S	23
17	<i>Onthophagus centricornis</i> Fabricius	T	2.9 ± 0.25	S	4
18	<i>Onthophagus lemniscatus</i> Gillet	T	7.00	S	1
19	<i>Onthophagus quadridentatus</i> Fabricius	T	7.50	S	1
20	<i>Onthophagus andrewesi</i> Arrow	T	6.5 ± 0.41	S	436
21	<i>Onthophagus cervus</i> Fabricius	T	6.4 ± 0.43	S	180
22	<i>Onthophagus turbatus</i> Walker	T	7.6 ± 0.43	S	160
23	<i>Onthophagus negligens</i> Walker	T	5.7 ± 0.32	S	171
24	<i>Onthophagus brutus</i> Arrow*	T	8.6 ± 0.42	S	5
25	<i>Onthophagus madoqua</i> Arrow	T	4.6 ± 0.4	S	71
26	<i>Onthophagus laevis</i> Harold*	T	9.4 ± 0.65	S	44
27	<i>Onthophagus favrei</i> Boucomont	T	6.4 ± 0.68	S	34
28	<i>Onthophagus devagiriensis</i> Schoolmeesters & Sabu*	T	6.7 ± 0.32	S	27
29	<i>Onthophagus ensifer</i> Boucomont	T	6.5 ± 0.31	S	25
30	<i>Onthophagus castetsi</i> Lansberge	T	10.5 ± 0.54	S	23
31	<i>Onthophagus falsus</i> Gillet	T	7 ± 0.57	S	20
32	<i>Onthophagus urellus</i> Boucomont	T	9.5 ± 0.5	S	9
33	<i>Onthophagus usurpator</i> Balthasar	T	8.8 ± 0.2	S	8
34	<i>Onthophagus elongates</i> Frey*	T	3.8 ± 0.26	S	6
35	<i>Onthophagus fasciatus</i> Boucomont	T	6.1 ± 0.59	S	6
36	<i>Onthophagus vladimiri</i> Frey*	T	6.6 ± 0.38	S	6
37	<i>Onthophagus pacificus</i> Lansberge	T	7.4 ± 0.65	S	5

R – rollers, D – dwellers, T – tunnelers. S – small beetles (< 13 mm), L – large beetles (≥13mm).

* endemic.

first record from Western Ghats.

Voucher specimens are temporarily deposited in the insect collections of St. Joseph's College, Devagiri, Calicut, and will be transferred to the national insect collections of Zoological Survey of India (ZSI), Calicut and Indian Agricultural Research Institute (IARI), New Delhi.

Rainfall data was collected from the records of Kerala State Electricity Board at Thirunelly. Humidity and forest floor temperature were assessed with thermo-hygrometer. Slope of the terrain was calculated using the trigonometric formula 'tanθ' (where 'θ' is the angle of inclination).

Data analysis

The species diversity of the assemblage of dung beetles attracted to gaur dung pats was calculated using Fisher's alpha diversity (Fisher et al. 1943) and Simpson's dominance and evenness (Simpson 1949) indices. Beta diversity was analysed with incidence based on the Bray Curtis similarity index (Bray and Curtis 1957) as the

sampling methodology employed for the collection of elephant dung beetle assemblage (Sabu et al. 2006) varied. Taxonomic diversity was analysed using non-parametric average taxonomic distinctness (Δ^+) and variation in taxonomic distinctness (Λ^+) indices (Clarke and Warwick 2001; Warwick et al. 2002). A regional master list of forest dung beetles from Wayanad was compiled from Sabu (2005), Sabu et al. (2006), Sabu and Vinod (2005) and the present study. A randomization test was done to detect differences in average taxonomic distinctness and variation in taxonomic distinctness, for any observed set of species, from the 'expected' Δ^+ and Λ^+ values derived from regional master species list (Clarke and Warwick 1998). Five taxonomic levels namely, species, genus, tribe, subfamily and family were considered. Branch lengths between taxonomic classes were defined following the standardization proposed by Warwick and Clarke (2001). Equal step lengths were assumed between each successive taxonomic level, setting path length ω to 100 for two species




Figure 3. Endemic dung beetle species (A-K) and dominant dweller species (L) recorded from gaur dung baited traps from moist forests of Wayanad region of Western Ghats. (A) *Liatongus indicus* Arrow, (B) *Caccobius gallinus* Arrow, (C) *Onthophagus andrewesi* Arrow, (D) *O. brutus* Arrow, (E) *O. madoqua* Arrow, (F) *O. laevis* Harold, (G) *O. ensifer* Boucomont, (H) *O. castetsi* Lansberge, (I) *O. elongates* Frey, (J) *O. vladimiri* Frey, (K) *O. devagiriensis* Schoolmeesters & Sabu and (L) *Drepanoceros setosus* Arrow. Magnification for each image is given in parenthesis.

connected at the highest (taxonomically closest) possible level. So the weights used were $\omega = 20$ (species in the same genus), $\omega = 40$ (same tribe but different genus), $\omega = 60$ (same subfamily but different tribe) and $\omega = 80$ (same family but different subfamily).

Simpson's diversity index was calculated with EstimateS 7.5 program (Colwell 2005). All other diversity analysis was done with Primer 5 software version 5.2.9. Variances of qualitative taxonomic diversity indices values (Δ^+ and Δ^+) with respect to the master list values were estimated by drawing 95% confidence funnels using Primer package (Clarke and Gorley 2002). Variations in abundances among samples were

analysed with one-way ANOVA test (Zar 2003). Megastat, version 10.0 (Orris 2005), was used for all statistical analysis.

Results

As shown in Figure 2, 37 species of dung beetles representing 10 genera and six tribes were recorded. The assemblage consisted of 10 endemics (*Liatongus indicus*, *Caccobius gallinus*, *Onthophagus lemniscatus*, *O. andrewesi*, *O. madoqua*, *O. devagiriensis*, *O. ensifer*, *O. castetsi*, *O. elongates* and *O. vladimiri*) and two first reports from the Western Ghats (*Onthophagus laevis* and *O. brutus*) (Table 1, Figure 3). Beetles belonging to all three major

Table 2. Overall species richness, guild wise abundance (%) and richness, guild structure and abundance (%) of dominant species of dung beetles in moist forests.

Locality	Species richness (% in parenthesis)				Abundance**			Dominant species			
	Total	Roller	Dweller/others*	Tunneler	Roller	Dweller/others*	Tunneler	Species	Guild	Abundance**	Total abundance**
South India ¹	37	2 (5.4)	3 (8.1)	32 (86.5)	5.4	16.6	78	<i>Onthophagus andrewesi</i>	T	18.6	29.1
								<i>Drepanocerus setosus</i>	D	10.5	
South India ²	44	4 (9.1)	3 (6.8)	37 (84.1)	-	-	-	-	-	-	-
Malaysia ³	87	12 (13.8)	3 (3.4)	72 (82.8)	-	-	-	-	-	-	-
Malaysia ⁴	59	6 (10.1)	1 (1.7)	52 (88.1)	39.9	0.3	59.8	<i>Sisypus thoracicus</i>	R	39	53.4
								<i>Onthophagus cervicapra</i>	T	14.4	
Ivory coast ⁵	66	5 (7.6)	6 (9.1)	55 (83.3)	7.8	20	72.2	<i>Oniticellus pseudoplanatus</i>	D	13.1	25.3
								<i>Caccobius elephantinus</i>	T	12.2	
Makoku ⁵	55	4 (7.3)	3 (5.5)	48 (87.2)	8.7	0.5	90.8	<i>Onthophagus fuscidorsis</i>	T	25.5	33
								<i>Sisypus arboreus</i>	R	7.5	
Australia ⁶	22	9 (40.9)	0	13 (59.1)	66	34		<i>Temnoplectron aeneopictum</i>	R	29.9	56.7
								<i>Coctodactyla depressa</i>	T	26.8	
Bolivia ⁷	53	12 (22.6)	8 (15.1)	33 (62.3)	31.7	6.6	61.7	<i>Dichotomius lucasi</i>	T	16.1	29.5
								<i>Deltochilum amazonicum</i>	R	13.4	
Brasil ⁸	46	11(23.9)	3 (6.5)	32 (69.6)	4.9	1.4	93.8	<i>Uroxys pygmaeus</i>	T	21.1	34
								<i>Ateuchus sp 2</i>	T	12.9	
Columbia ⁹	39	13(33.3)	4 (10.3)	22 (56.4)	25.3	9.9	64.8	<i>Ateuchus sp.1</i>	T	33.3	40.5
								<i>A. muriayi</i>	T	7.2	
Columbia ¹⁰	52	10(19.2)	6 (11.5)	36 (69.2)	15.4	53.8	30.6	<i>Eurysternus sp</i>	-	44.5	50
								<i>Onthophagus sp.1</i>	T	5.5	
Guyana ¹¹	76	18(23.7)	8 (10.5)	50 (65.8)	27.1	14.8	58	<i>Ateuchus simplex</i>	T	15.6	29.4
								<i>Hansreia affinis</i>	R	13.8	
Mexico ¹²	30	8(26.7)	3 (10)	19 (63.3)	45	1.1	46	<i>Canthon femoralis</i>	R	28.1	42.7
								<i>Onthophagus batesi</i>	T	14.6	
Panama ¹³	59	13(22)	3 (5.1)	43 (72.9)	40	1.1	58.9	<i>Canthon aequinoctialis</i>	R	30.2	48.2
								<i>Canthidium aurifex</i>	T	18.2	
Peru ¹⁴	87	27(31)	5 (5.7)	55 (63.2)	-	-	-	-	-	-	-
								-	-	-	

¹ Present study, ² Combined data from present study and Sabu et al. 2006, ³ Davis 2000a, ⁴ Davis et al. 2000, ⁵ Cambefort and Walter 1991, ⁶ Howden et al. 1991, ⁷ Spector 2003, ⁸ Andresen 2002, ⁹ Escobar 2000, ¹⁰ Howden and Nealis 1975, ¹¹ Feer 2000, ¹² Estrada et al. 1999, ¹³ Gill 1991, ¹⁴ Valencia 2001

* Others – *Eurysternus* species categorised as intermediate between dwellers and tunnelers

** abundance data in percentages

T- tunneler, D- dweller, R- roller

functional guilds were present. *Onthophagus andrewesi*, a tunneler (18.6%), and *Drepanocerus setosus*, a dweller (10.5%), dominated the assemblage (Table 2). Tunnelers were the most speciose (32 species, 86.5 %) and abundant (78 %) functional guild. Rollers were represented by two species, *Ochicanthon laetum* and *Sisypus neglectes*, and was the least abundant functional group (5.4 % of total abundance) (Table 1). Smaller beetles dominated the assemblage in terms of species richness (83.8%) and abundance (88.8.4%). The assemblage was moderately diverse ($\alpha = 6.63$) and highly even ($1-\lambda = 0.92$). Variation in abundance among samples was not significant ($df = 14$, $f = 1.54$, $P = 0.09$). 21 species

belonging to 10 genera, six tribes and three nesting guilds were collected from elephant dung (Figure 2B). Bray Curtis similarity index illustrated moderate similarity (48.28) between gaur and elephant dung beetle assemblages. Taxonomic diversity and evenness of gaur dung beetle assemblage ($\Delta^+ = 42.91$, $\Lambda^+ = 471.4$) were lower in comparison to elephant dung beetle assemblage ($\Delta^+ = 58.48$, $\Lambda^+ = 344.3$). Values of both taxonomic diversity indices fell within the 95% limits of the probability funnel indicating that taxonomic diversity of both the assemblages did not vary significantly from the regional species pool.

Discussion

A taxonomic diversity index is a measure of biodiversity that indicates how different the species in a habitat are from each other (Harper and Hawksworth 1994). The taxonomic relatedness diversity indices have appealing sampling properties: non-dependence on quantitative data and consideration of the relatedness of species in an assemblage that are of great practical utility in diversity analysis and are considered as being most promising for biodiversity assessments (Warwick and Clarke 2001; Price 2002; Warwick et al. 2002; Magurran 2004). All dung beetle diversity assessments have been done so far with the conventional species richness and evenness-based diversity indices. However, such over reliance on patterns of dung beetle richness alone can be seriously misleading and community level data are important in dung beetle studies (Spector 2001). Hence, taxonomic relatedness based diversity properties of dung beetles attracted to dung of gaur and elephant were also used. Unfortunately, the IndVal methods of Dufrene and Legendre (1997) useful in detecting indicator species characterizing habitat types and groups of samples could not be used as the requisite data were not available.

The first report of the community structure and diversity of dung beetles in a moist forest locality in Western Ghats and South Asian region is provided. Most conspicuous is the difference observed in the guild structure of the community, when compared to dung beetle assemblages from other moist forests of the Afrotropical region. Dwellers are the dominant functional guild after tunnelers, and rollers are lower in richness and abundance. Such high abundance of dwellers is reported previously only from the moist forests of Ivory Coast in Africa.

Combining the 37 species recorded from the present study along with 7 species reported exclusively from elephant dung (Sabu et al. 2006) leads to an overall richness of 44 species in the region. Species richness is comparatively lower compared to the 87 species reported from Malaysia (Davis 2000a) and Peru (Valencia 2001), 76 from French Guyana (Feer 2000) and 66 species from African rain forests (Cambefort and Walter 1991). However, rain forests of Mexico (Estrada and Coates-Estrada 2002), Colombia (Escobar 2000) and Australia (Howden et al. 1991) have, on average, lower local richness of dung beetles. Though our sampling is limited to a

relatively short period of two months, no additional species were added from our two year study with bimonthly random sampling from the same region with more sampling effort (Sabu 2005). Hence, we consider that present study did successfully sample most, if not all, species of dung beetles that could be trapped with baited pitfall traps from the region. Two first reports (*Onthophagus laevis* and *O. brutus*) from Western Ghats and high abundance of endemics (32.6 %) indicate that further characterization of the dung beetle faunal diversity of other forests of Western Ghats down to more local scales may reveal more details of the regional variation in endemism and localised distribution patterns.

Comparatively high abundance of old world roller, *O. laetum*, whose overall abundance is very low in south east Asian forests (Davis et al. 2000), and dominance of *D. setosus* and the endemic species *L. indicus*, are most likely a regional phenomenon. *D. setosus* and *L. indicus* are prominent dwellers in both fresh and old dung pats of elephant and gaur in the region (unpublished observations). Dominance by a few tunneler, roller or guild unspecified species (personal communications, Fernando Vaz-de-Mello) in the range of 56.7 % to 29.4% or tunneler species alone in the range of 34% - 40.5%, is a general pattern of tropical moist forest dung beetle communities. Moist forests of the Ivory Coast (Cambefort and Walter 1991) and Wayanad are the only exceptions where the dominant species are distributed between tunneler and dweller guilds (29.1 % to 25.3%), and rollers are the least abundant guild.

Substantially high abundance of *D. setosus*, and *L. indicus* leads to the dominance of dwellers (Oniticellini). A similar situation exists in the moist forests of Ivory Coast in Western Africa with the abundance of *Oniticellus pseudoplanatus* (Oniticellini) and is attributed to the availability of undisturbed elephant dung pats in the region (Cambefort and Walter 1991). Dwellers are strongly associated with larger herbivore dung pats and breed successfully only in undisturbed dung pats with little competition from competitively superior tunnelers and rollers (Cambefort 1991; Hanski and Krikken 1991; Krell et al. 2003). Apparently, a similar situation prevails in the north Wayanad region with the presence of large amounts of undisturbed dung pats of elephant and gaur (unpublished observations), probably in excess of consumption

by dung beetles in these forests. The moist forests of the north Wayanad region merge gently with the drier forests on the eastern slope and are a summer refuge for herds of elephants and gaurs (Joy 1991; Nair 1991). Hence we attribute the high dweller abundance in the region to the abundance and seasonal movement of large herbivorous mammals and ready availability of large dung pats.

Although dwellers are dominant over rollers in moist forests of large herbivore rich Ivory Coast and Wayanad region (Joy 1991; Nair 1991; Krell et al. 2003), we are unaware of how much the massive slaughtering of African elephants which peaked during 1980–1989 (Sukumar 2003) in Ivory Coast might have affected the availability of elephant dung and dung beetle guild structure in the region. Columbian rainforests, described earlier with high dweller abundance (Howden and Nealis 1975), showed an entirely different guild structure in more recent reports with low presence of dwellers (Escobar 2000), which is probably related to the extensive deforestation of Amazonian forests (Anderson 1990; Skole and Tucker 1993).

The low abundance of rollers is in contrast to their high abundance and richness in South East Asian forests of Borneo (Davis et al. 2000). Analysis of diversity of forest floor arthropods including dung beetles along the altitudes of Wayanad forests revealed a general low incidence of rollers and absence of large rollers above mid elevations (800m amsl) whereas both small (*Sisyphus* and *Ochicanthon*) and large rollers (*Gymnopleurus*) are abundant in the middle and low elevation (600, 300m amsl) moist forests (Sabu 2005; Sabu et al. 2006). This indicates that low presence of rollers is a regional pattern and is not a sampling error arising from the more seasonal study as in the present case. Delay in drying dung pats in shady cool forests makes dung ball making and rolling an energetically costly activity for thermophilic rollers and makes them competitively inferior to other guilds (Krell et al. 2003). Hence, the low forest floor temperature and high humidity in these shady high humid forests which keeps elephant (5–7 days) and gaur dung pats (10–15 days) moist and wet for a longer period, as was found during succession studies with elephant dung (Sabu et al. 2006) and as observed in field conditions (unpublished observations), are likely to be the major reason for the lower abundance and richness of rollers in the region.

Although, variations in sampling effort restricts our ability to interpret the data, comparison of the assemblages suggests that gaur dung attracts a more highly speciose dung beetle assemblage than elephant dung. The dominance of small sized dung beetles in both elephant (Sabu et al. 2006) and gaur dung baited traps and dung pats (unpublished observations) indicate that availability of large voluminous dung pats do not lead to an abundance of large dung beetles in the study region. Dung of the non-ruminating elephant is more fibrous and coarse than gaur dung, but they are similar in the sense that they are both herbivore in origin, moist and non-pelleted (Botes et al. 2006; Doube 1991). Moderate similarity values indicate that beetle assemblages attracted to either elephant or gaur dung do not constitute entirely dissimilar communities, but rather one community with more generalists that can use both dung types and a few specialists as well. The presence of 7 species exclusively in elephant dung baited traps, along with the categorization of dung beetles into coarse and fine dung feeders (Davis 2002; Holter et al. 2002), suggest that they are elephant dung specialists. Absence of 23 dung beetle species attracted to gaur dung in elephant dung baited traps may be related to the fluid dung preference of these species. However, variations in the sampling methodology necessitate more empirical studies to reach conclusions.

Although species richness was higher in the dung beetle assemblages attracted to gaur dung pats, high Λ^+ (low taxonomic evenness) values indicate the presence of a phylogenetically closely related dung beetle assemblage. Analysis of taxonomic evenness by truncating the tree at various places and by removing the speciose genera showed that both taxonomic evenness and diversity of gaur dung beetle assemblage equaled that of elephant when species distribution under the genera *Onthophagus* and *Caccobius* were made even in both assemblages. High unevenness in taxonomic structure of the gaur dung beetle assemblage arises from the overrepresentation of *Onthophagus* and *Caccobius* species. The presence of 24 species of *Onthophagus* and 3 species of *Caccobius* in gaur dung (65% of the species attracted to gaur dung from genus *Onthophagus* and 73% from genus *Onthophagus* and *Caccobius*) compared to the presence of 7 *Onthophagus* species (33.3%) and the absence of *Caccobius* in elephant dung, reduced the taxonomic evenness of gaur dung beetle assemblage. This variation is distinctly




Figure 4. Physical structure of gaur and elephant dung from moist forests of south Western Ghats in Wayanad region. a). Fine fluid dung of gaur and b). Coarse fibrous dung of elephant.

shown by Λ^+ , as the variation in taxonomic distinctness index is sensitive to variations in taxonomic evenness of the assemblage and the presence of speciose genera reduces the taxonomic evenness of the assemblage which is reflected as higher Λ^+ values.

The overrepresentation of closely related species, and the resulting high unevenness of the taxonomic structure of dung beetles attracted to gaur dung in comparison to elephant dung, we relate to the coarse and fine dung preferences of dung beetles (Davis 2002; Holter et al. 2002), and to variation in the physical properties of the two dung types (Figure 4). For all groups of organisms, specific taxa attain their highest diversity in particular habitats, and when certain habitat types are absent from an area some groups of species become underrepresented while others become overrepresented compared to the regional picture (Warwick and Clarke 2001), resulting in a more uneven distribution across the phylogenetic tree. Dung pats are patchily distributed and ephemeral minor habitats for dung beetles (Elton 1949; Hanski 1991). Though many dung beetles are generalists and do not show any dung preferences, some preferably select coarse fibred dung of non-ruminants while others prefer the more fluid and fine dung of ruminants, and some others the odoriferous dung of omnivores (Davis 1994; 2002; Holter et al. 2002; Krell et al. 2003). Hence, two structurally different and contrasting

dung types (i.e. two minor habitats), the homogenous, fine, fluidy dung of the ruminant gaur and the heterogeneous dung of the elephant with both fibrous and fine dung particles, are readily available for the dung beetle community in the study region. Homogenous, fine gaur dung pats attract species with similar (fine) dung resource requirements and hence more closely related species belonging to specific genera or tribes. Whereas, heterogeneous elephant dung attracts both coarse and fine dung feeders and generalists from different tribes and genera (i.e. less related species) leading to the higher taxonomic evenness that is distinct in the dendrogram. The average taxonomic distinctness Δ^+ of the assemblages showed lesser variations than Λ^+ , as Δ^+ considers only the relatedness between individual member species involved and not the taxonomic evenness properties of the assemblage.

In summary, the present study provides for the first time data about community structure of dung beetles from moist forests of Western Ghats, as well from a South Asian region. Though with low species richness, elephant dung attracts a more taxonomically diverse and even dung beetle assemblage than gaur dung that is likely to be related to the more heterogeneous physical nature of elephant dung with both fluid and fibrous dung particles. The presence of many endemics (27%), predominance of *O. andrewesi*, an endemic of the

Western Ghats, and *D. setosus* recorded only from the Indian continent, and the higher incidence of the old world roller *O. laetum*, makes dung beetle assemblage in the moist forests of this region unusual. The dominance of dwellers (Oniticellini) over rollers makes the functional guild structure of dung beetle assemblage of the Wayanad forests more similar to the dung beetle community of the Ivory Coast forests of Western Africa and different from those of south East Asian (Borneo) and Neotropical forests. Furthermore, the current study reiterates that the abundance of dwellers is an indicator of the availability of undisturbed dung pats and herbivore abundance in moist forests. However, not enough data exists to establish that the predominance of dwellers, and the low abundance and species richness of rollers, is a general pattern applicable to entire moist forests of Western Ghats. Further studies are necessary to ascertain whether it is a regional pattern specific to the transitional Wayanad forests of Western Ghats alone.

Acknowledgements

Financial assistance provided by the Ministry of Environment and Forests (MoEF), Govt. of India is gratefully acknowledged. We are grateful to the Kerala Forest and Wild Life Department for providing us the permission and facilities in the study region. We are indebted to K.T. Thomachan, Devagiri College, Calicut, for statistical analysis. This manuscript greatly benefited from comments and suggestions by two anonymous reviewers. We thank Fernando Vaz-de-Mello, Paul Schoolmeesters, E. Andresen, A.J Davis, F.Feer, A. Estrada, F.S. Escobar, V.V. Ramamurthy and T.N. Ananthkrishnan for their generous help in numerous ways towards the study of dung beetles in the region.

References

- Anderson AB. Deforestation in Amazonia: Dynamics, causes, and alternatives. In: Anderson AB, editor. 1990. *Alternatives to Deforestation: Steps toward sustainable use of the Amazon rain forest* 3-23. Columbia University Press.
- Andresen E. 2006. Dispersion Primaria de Semillas por Primates y Dispersion Secundaria por Escarabajos Coprofagos en Tikal, Guatemala. *Biotropica* 38: 390-397.
- Andresen E. 2003. Effect of forest fragmentation on dung beetle communities and functional consequences for plant regeneration. *Ecography* 26: 87-97.
- Andresen E. 2002. Dung beetles in a Central Amazonian rainforest and their ecological role as secondary seed dispersers. *Ecological Entomology* 27: 257-270.
- Andresen E. 2001. Effect of dung presence, dung amount, and secondary dispersal by dung beetles on the fate of *Micropholis guyanensis* (Sapotaceae) seeds in Central Amazonia. *Journal of Tropical Ecology* 17: 61-78.
- Andresen E, Levey DJ. 2004. Effects of dung and seed size on secondary dispersal, seed predation, and seedling establishment of rain forest trees. *Oecologia* 139: 145-154.
- Arrow GJ. 1931. *The Fauna of British India including Ceylon and Burma, Coleoptera: Lamellicornia (Coprinae)*. Taylor and Francis, London.
- Balthasar V. 1963a. *Monographic der Scarabaeidae und Aphodiidae der Palaearktischen und Orientalischen Region (Coleoptera: Lamellicornia)*. Verlag der Tschechoslowakischen Akademie der Wissenschaften Prag.
- Balthasar V. 1963b. *Monographic der Scarabaeidae und Aphodiidae der Palaearktischen und Orientalischen Region (Coleoptera: Lamellicornia)*. Verlag der Tschechoslowakischen Akademie der Wissenschaften Prag.
- Bergstrom BC, Maki RL, Werner BA. 1976. Small dung beetles as biological control agents: laboratory studies of beetle action on trichostongylid eggs in sheep and cattle feces. *Proceedings of the Helminthology Society of Washington* 43: 171-174.
- Bossuyt F, Meegaskumbura M, Beenaerts N, Gower JD, Pethiyagoda R, Roelants K, Mannaert A, Wilkinson M, Bahir MM, Arachchi MK, Peter KLN, Schneider JC, Oommen VO, Milinkovitch CM. 2004. Local endemism within the Western Ghats, Srilanka biodiversity hotspot. *Science* 306: 479-481.
- Bornemissa GF, Williams CH. 1970. An effect of dung beetle activity on plant yield. *Pedobiologia* 10: 1-7.
- Botes A, McGeoch MA, van Rensburg BJ. 2006. Elephant- and human-induced changes to dung beetle (Coleoptera: Scarabaeidae) assemblages in the Maputaland Centre of Endemism. *Biological Conservation* 130: 573 -583.
- Bray JR, Curtis CT. 1957. An ordination of the upland forest communities of Southern Wisconsin. *Ecological Monographs* 27: 325-349.
- Cambeftor Y, Hanski I, Cambeftor Y. 1991. Biogeography and evolution. *Dung beetle ecology*, 51-67. Princeton University Press.
- Cambeftor Y, Hanski I. Dung beetle population biology. In: Hanski I, Cambeftor Y, editors. 1991. *Dung beetle ecology*, 36-50. Princeton University Press.
- Cambeftor Y, Walter P. Dung beetles in Tropical Forests in Africa. In: Hanski I, Cambeftor Y, editors. *Dung beetle ecology*, 198-210. Princeton University Press.

- Clarke KR, Gorley RN. 2002. *Primer v5 Primer-E-Ltd* Plymouth Marine Laboratory. Prospect place, West Hoe. United Kingdom.
- Clarke KR, Warwick RM. 2001. *Change in marine communities: an approach to statistical analysis and interpretation*, 2nd edition. Plymouth Marine Laboratory, UKPRIMER-E Ltd.
- Clarke KR, Warwick RM. 1998. A taxonomic distinctness index and its statistical properties. *Journal of Applied Ecology* 35: 523-531.
- Colwell RK. 2005. *EstimateS: statistical estimation of species richness and shared species from samples*. Version 6. User's Guide and application available at: <http://viceroy.eeb.uconn.edu/estimates>
- Davis AJ. 2000a. Species richness of dung-feeding beetles (Coleoptera: Aphodiidae, Scarabaeidae, Hybosoridae) in Tropical rainforest at Danum Valley, Sabah, Malaysia. *The Coleopterists Bulletin* 54: 221-231.
- Davis AJ. 2000b. Does reduced-impact logging help preserve biodiversity in tropical rainforests? A case study from Borneo using dung beetles as indicators. *Environmental Entomology* 29: 467-475.
- Davis AJ, Sutton SL. 1998. The effects of rainforest canopy loss on arboreal dung beetles in Borneo: implications for the measurement of biodiversity in derived tropical ecosystems. *Diversity and Distributions* 4: 167-173.
- Davis AJ, Holloway JD, Huijbregts H, Kirk-Spriggs AH, Sutton SL. 2001. Dung beetles as indicators of change in the forests of northern Borneo. *Journal of Applied Ecology* 38: 593-616.
- Davis AJ, Huijbregts J, Krikken J. 2000. The role of local and regional processes in shaping dung beetle communities in tropical forest plantations in Borneo. *Global Ecology and Biogeography* 9: 281-292.
- Davis ALV. 2002. Dung beetle diversity in South Africa: influential factors, conservation status, data inadequacies and survey design. *African Entomology* 10: 53-65.
- Davis ALV. 1996. Community organization in dung beetles (Coleoptera: Scarabaeidae): differences in body size and functional group structure between habitats. *African Journal of Ecology* 34: 258-275.
- Davis ALV. 1994. Associations of Afrotropical Coleoptera (Scarabaeidae, Aphodiidae, Staphylinidae, Hydrophilidae, Histeridae) with dung and decaying matter: implications for selection of fly-control agents for Australia. *Journal of Natural History* 28: 383-399.
- Davis ALV, Scholtz CH, Dooley PW, Bham N, Kryger U. 2004. Scarabaeinae dung beetles as indicators of biodiversity, habitat transformation and pest control chemical in agro-ecosystems. *South African Journal of Science* 100: September/October.
- Doube BM. Dung beetles of South Africa. In: Hanski I, Cambefort Y, editors. 1991. *Dung beetle ecology* 133-155. Princeton University Press.
- Dufrène M, Legendre P. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* 67: 345-366.
- Elton C. 1949. Population interspersion: An essay on animal community patterns. *Journal of Ecology* 37: 1-23.
- Errouissi F, Jay-Robert P, Lumaret JP, Plau P. 2004. Composition and structure of dung beetle (Coleoptera: Aphodiidae, Geotrupidae, Scarabaeidae) assemblages in mountain grasslands of the Southern Alps. *Annals of the Entomological Society of America* 97: 701-709.
- Escobar FS. 2000. Diversidad de colepteros coprofagos (Scarabaeidae: Scarabaeinae) en un mosaico de habitats en la reserva natural Nukak, Guiviare, Colombia. *Acta Zoologica Mexicana* 79: 103-121.
- Escobar FS. 1997. Estudio de la comunidad de coleópteros coprófagos (Scarabaeidae) en un remanente de bosque seco al norte del Tolima, Colombia. *Caldasia* 19: 419-430.
- Estrada A, Coates-Estrada R. 2002. Dung beetles in continuous forest, forest fragments and in an agricultural mosaic habitat island at Los Tuxtlas, Mexico. *Biodiversity and Conservation* 11: 1903-1918.
- Estrada A, Coates-Estrada R. 1991. Howling monkeys (*Alouatta palliata*), dung beetles (Scarabaeidae) and seed dispersal: ecological interactions in the tropical rain forest of Los Tuxtlas, Veracruz, Mexico. *Journal of Tropical Ecology* 7: 459-474.
- Estrada A, Anzures AD, Coates-Estrada R. 1999. Tropical rain forest fragmentation, howler monkeys (*Alouatta palliata*) and dung beetles at Los Tuxtlas, Mexico. *American Journal of Primatology* 48: 253-262.
- Estrada A, Halffter G, Coates-Estrada R, Meritt D. 1993. Dung beetles attracted to mammalian herbivore (*Alouatta palliata* Gray) and omnivore (*Nasua narica* Linnaeus) dung in the tropical rain forest of Los Tuxtlas, Mexico. *Journal of Tropical Ecology* 9: 45-54.
- Favila ME, Halffter G. 1997. The use of indicator groups for measuring biodiversity as related to community structure and function. *Acta Zoologica Mexicana (N.S.)* 72: 1-25.
- Feer F. 2000. Les Coléoptères Coprophages Et Nérophages (Scarabaeidae S. Str. Et Aphodiidae) De La Forêt De Guyane Française: Composition Spécifique Et Structure Des Peuplements. *Annales de la Société Entomologique de France (N. S.)* 36: 29-43.
- Feer F. 1999. Effects of dung beetles (Scarabaeidae) on seeds dispersed by howler monkeys (*Alouatta seniculus*) in the French Guianan rainforest. *Journal of Tropical Ecology* 15: 129-142.
- Feer F, Pincebourde S. 2005. Diel flight activity and ecological segregation with in an assemblage of tropical forest dung and carrion beetles. *Journal of Tropical Ecology* 21: 21-30.
- Fisher RA, Corbet AS, Williams CB. 1943. The relation between the number of species and the number of individuals in a random sample of an animal population. *Journal of Animal Ecology* 12: 42-58.

- Gill BD. Dung beetles in tropical American forests. In: Hanski I, Cambefort Y, editors. 1991. *Dung beetle ecology*, 211-229. Princeton University Press.
- Halffter G, Arellano L. 2002. Response of dung beetle diversity to human-induced changes in a tropical landscape. *Biotropica* 34: 144-154.
- Halffter G, Edmonds WD. 1982. *The Nesting Behavior of Dung Beetles (Scarabaeinae): An Ecological and Evolutionary Approach*. Instituto de Ecología, AC, Mexico, DF.
- Halffter G, Mathews EG. 1966. The natural history of dung beetles of the sub family Scarabaeinae (Coleoptera, Scarabaeidae). *Folia Entomologica Mexicana* 12-14: 1-132.
- Halffter G, Favila ME, Halffter V. 1992. A comparative study of the structure of the scarab guild in Mexican tropical rain forest and derived ecosystems. *Folia Entomologica Mexicana* 84: 131-156.
- Hanski I. The dung insect community. Hanski I, Cambefort Y, editors. 1991. *Dung beetle ecology*, 5-21. Princeton University Press.
- Hanski I. 1989. Metapopulation dynamics: does it help to have more of the same?. *Trends in Ecology and Evolution* 4: 113-114.
- Hanski I. 1983. Distributional ecology and abundance of dung and carrion-feeding beetles (Scarabaeidae) in tropical rain forests in Sarawak, Borneo. *Acta Zoologica Fennica* 167: 1-45.
- Hanski I, Cambefort Y. Competition in dung beetles. In: Hanski I, Cambefort Y, editors. 1991. *Dung beetle ecology*, 305-329. Princeton University Press.
- Hanski I, Krikken J. Dung beetles in the Tropical forests of South- East Asia. In: Hanski I, Cambefort Y, editors. 1991. *Dung beetle ecology*, 179-197. Princeton University Press.
- Harper JL, Hawksworth DL. 1994. Biodiversity: measurement and estimation. *Philosophical Transactions of the Royal Society of London. Series B. Biological Sciences* 345: 5-12.
- Hill CJ. 1996. Habitat specificity and food preferences of an assemblage of tropical Australian dung beetles. *Journal of Tropical Ecology* 12: 449-460.
- Holter P, Scholtz CH, Wardhaugh KG. 2002. Dung feeding in adult Scarabaeines (tunnellers and dwellers): even large dung beetles eat small particles. *Ecological Entomology* 27: 169-176.
- Howden HF, Nealis VG. 1975. Effects of clearing in a tropical rain forest on the composition of the coprophagous scarab beetle fauna (Coleoptera). *Biotropica* 7: 77-83.
- Howden HF, Young OP. 1981. Panamanian Scarabaeidae. *Contributions of the American Entomological Institute* 18: 1-204.
- Howden HF, Howden AT, Storey RI. 1991. Nocturnal perching of Scarabaeidae dung beetles (Coleoptera, Scarabaeidae) in an Australian tropical rain forest. *Biotropica* 23: 51-57.
- Joy MS. 1991. *Keralathile Vanyajeevisankethangal* (Wild Life Reserves in Kerala). State Institute of Languages, Kerala.
- Klein BC. 1989. Effects of forest fragmentation on dung and carrion beetle communities in central Amazonia. *Ecology* 70: 1715-1725.
- Krell FT, Westerwalbesloh SK, Weib I, Eggleton P, Linsenmair KE. 2003. Spatial separation of Afrotropical dung beetle guilds: a trade-off between competitive superiority and energetic constraints (Coleoptera: Scarabaeidae). *Ecography* 26: 210-222.
- Lobo JM, Martin Piera F, Veiga CM. 1988. Las trampas pitfall con cebo, sus posibilidades en el estudio de Scarabaeoidea (Col.) I. Características determinantes de su capacidad de captura. *Revue D Ecologie Et De Biologie Du Sol* 25: 77-100.
- Lumaret JP, Kadiri N, Bertrand M. 1992. Changes in resources: consequences for the dynamics of dung beetle communities. *Journal of Applied Ecology* 29: 349-356.
- Magurran AE. 2004. *Measuring biological diversity* Blackwell Publishing.
- Mayers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853-858.
- McGeoch MA, van Rensburg BJ, Botes A. 2002. The Verification and application of bioindicators: a case study of dung beetles in a savanna ecosystem. *Journal of Applied Ecology* 39: 661-672.
- Nair SC. 1991. *The Southern Western Ghats- a biodiversity conservation plan*. Indian National Trust for Art and Cultural Heritage, New Delhi.
- Nealis VG. 1977. Habitat associations and community analysis of South Texas dung beetles (Coleoptera: Scarabaeinae). *Canadian Journal of Zoology* 55: 138-147.
- Orris JB. 2005. Megastat version 10.0, Butler University, College of Business Administration, 4600 Sunset Ave, Indianapolis. Distributed by McGraw-Hill. <http://www.mhhe.com/support>
- Osberg DC, Doube BM, Hanrahan SA. 1994. Habitat specificity in African dung beetles: the effect of soil type on the survival of dung beetle immatures (Coleoptera: Scarabaeidae). *Tropical Zoology* 7: 1-10.
- Pascal JP. 1988. *Wet Evergreen Forests of the Western Ghats of India: Ecology, Structure, Floristic Composition and Succession*. French Institute, Pondicherry.
- Peck SB, Forsyth A. 1982. Composition, structure, and competitive behaviour in a guild of Ecuadorian rain forest dung beetles (Coleoptera, Scarabaeidae). *Canadian Journal of Zoology* 60: 1624-1634.

- Price ARG. 2002. Simultaneous 'hotspots' and 'cold spots' of marine biodiversity and implications for global conservation. *Marine Ecology Progress Series* 241: 23-27.
- Rodgers WA, Panwar HS. 1988. *Planning a wildlife protected areas network in India* Department of Environment, Forests, and Wildlife/Wildlife Institute of India report, Wildlife Institute of India 1-2.
- Sabu TK. 2005. *Litter insect dynamics with special reference to ecological succession and chemical ecology along varying altitudes in the Wynad and Coorg forests of Western Ghats*. Project report submitted to Ministry of Environment and Forests, Government of India.
- Sabu TK, Vinod KV. 2005. Comparative assessment of the guild structure and taxonomic diversity of two beetle (Coleoptera: Scarabaeinae) assemblages in the Wayanad region of Western Ghats. 47-52. *Proceedings of the National Conference safe environment for the future generations*. Dept. Zoology, Auxilium College, Vellore, Tamil Nadu.
- Sabu TK, Vinod KV, Vineesh PJ. 2006. Guild structure, diversity and succession of dung beetles associated with Indian elephant dung in South Western Ghats forests. *Journal of Insect Science* 6: 17, available online: <http://insectscience.org/6.17>
- Scheffler P. 2005. Dung beetle (Coleoptera: Scarabaeidae) diversity and community structure across three disturbance regimes in eastern Amazonia. *Journal of Tropical Ecology* 21: 9-19.
- Scheffler PY. 2002. Dung beetle (Coleoptera: Scarabaeidae) ecology in the intact and modified landscape of Eastern Amazonia. *PhD Thesis* Pennsylvania State University.
- Simpson EH. 1949. Measurement of diversity. *Nature* 163:688.
- Skole D, Tucker CJ. 1993. Tropical deforestation and habitat fragmentation in the Amazon: satellite data from 1978 to 1988. *Science* 260: 1905-1910.
- Spector SH. 2001. Conserving insects in the tropics: describing and predicting insect-habitat relationships at local to regional scales. *Ph D Thesis* University of Connecticut.
- Sukumar R. 2003. *The living elephants*. Oxford University Press.
- Valencia G. Diversity and Trophic Relationships of Dung Beetles of the Lower Urubamba Region, Peru. In: Alonso A, Dallmeier F, Campbell P, editors. 2001. *Urubamba: The Biodiversity of a Peruvian Rainforest*, 121-128. SI/MAB Series #7, Smithsonian Institution, Washington, DC.
- van Rensburg BJ, McGeoch MA, Chown SL, van Jaarsveld AS. 1999. Conservation of heterogeneity among dung beetles in the Maputaland Centre of Endemism, South Africa. *Biological Conservation* 7: 945-965.
- Veiga CM, Lobo JM, Piera FM. 1989. Las trampas pitfall con cebo, sus posibilidades en el estudio de las comunidades de Scarabaeoidea (Col.). II. Analisis de efectividad. *Revista de Ecología et de Biología du Sol* 26: 91-109.
- Vulinec K. 2000. Dung beetles (Coleoptera: Scarabaeidae), monkeys, and conservation in Amazonia. *Florida Entomologist* 83: 229-241.
- Warwick RM, Clarke KR. 2001. Practical measures of marine biodiversity based on relatedness of species. *Oceanography and Marine Biology. An Annual Review* 39: 207-231.
- Warwick RM, Ashman CM, Brown AR, Clarke KR, Dowell B, Hart B, Lewis RE, Shillabeer N, Somerfield PJ, Tapp JF. 2002. Inter-annual changes in the biodiversity and community structure of the macrobenthos in Tees Bay and Tees estuary, UK, associated with local and regional environmental events. *Marine ecology progress series* 234: 1-13.
- WWF 2001. Wild world, WWF full report, South Western Ghats montane rain forests (IMO151). Available online http://worldwildlife.org/wildworld/profiles/terrestrial/im/im0151_full.html
- Zar JH. 2003. *Biostatistical Analysis*. Pearson Education (Singapore) Pte. Ltd., Indian branch, Delhi, India.