

TEMPORAL OCCURRENCE OF THIRD-STAGE LARVAE OF ELAEOPHORA SCHNEIDERI IN TABANUS LINEOLA HINELLUS ON SOUTH ISLAND, SOUTH CAROLINA

Authors: Couvillion, C. E., Nettles, V. F., Sheppard, D. C., Joyner, R. L., and Bannaga, O. M.

Source: Journal of Wildlife Diseases, 22(2) : 196-200

Published By: Wildlife Disease Association

URL: <https://doi.org/10.7589/0090-3558-22.2.196>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

TEMPORAL OCCURRENCE OF THIRD-STAGE LARVAE OF *ELAEOPHORA SCHNEIDERI* IN *TABANUS LINEOLA HINELLUS* ON SOUTH ISLAND, SOUTH CAROLINA

C. E. Couvillion,^{1,5} V. F. Nettles,¹ D. C. Sheppard,² R. L. Joyner,³ and O. M. Bannaga⁴

ABSTRACT: The occurrence of third-stage larvae of the arterial worm (*Elaeophora schneideri*) in populations of the horse fly, *Tabanus lineola hinellus*, was studied on South Island, Georgetown County, South Carolina. Over the period from mid-April to mid-October 1982, horse flies had a bimodal pattern of abundance with peak populations in June and September. During individual bimonthly horse fly collections, the prevalence of infection of *T. l. hinellus* ranged from 0 to 1.23%. Prevalence of infected horse flies was biphasic with peaks in mid-May (1.23%) and mid-August (1.22%), 2 to 4 wk prior to peaks in horse fly populations. The intensity of infection of *T. l. hinellus* increased as time progressed and was highest in late summer. This increase resulted in greater recovery of infective larvae during August–September (mean intensity = 25) than in May–June (mean intensity = 4). Both prevalence and intensity of infection of *T. l. hinellus* were low compared to primary intermediate hosts of *E. schneideri* in the western U.S.

INTRODUCTION

Records of intermediate hosts of the arterial worm in the southeastern United States are restricted to a single report. Couvillion et al. (1984) found two species of horse flies, *Tabanus lineola hinellus* and *T. nigrovittatus*, infected with third-stage larvae of *E. schneideri*; and *T. l. hinellus* was considered the primary intermediate host.

The transmission of arterial worms is seasonal since it depends on the patterns of activity of horse flies, as well as seasonal aspects of the life history of the parasite in horse flies. Clark (1972) and Clark and Hibler (1973) provided data on the sea-

sonal occurrence of horse flies and larval *E. schneideri* in the western United States. The objective of this paper was to describe the temporal patterns of occurrence of *E. schneideri* in *T. l. hinellus* on South Island, South Carolina.

MATERIALS AND METHODS

The study was conducted on the Tom Yawkey Wildlife Center, an island complex comprised of South, Cat, and North islands in Georgetown County, South Carolina. It was selected as the study site because white-tailed deer (*Odocoileus virginianus*) infected with *E. schneideri* have been found on the South Island portion of the area (Hibler and Prestwood, 1981) and also because potential intermediate hosts were abundant. This study was conducted on a 48-ha hammock on the northern section of South Island. Detailed description of the area has been given by Epstein (1983).

In 1981, female *T. l. hinellus* were trapped for consecutive 3-day periods at monthly intervals from July through September. In 1982, trapping was done for consecutive 2-day periods every 2 wk from mid-April through mid-October. Six canopy traps and two malaise traps (Roberts, 1976) were placed in openings adjacent to woodlands alongside a north-south road. Traps were set and baited with dry ice at daylight. Horse flies were removed alive at midday and again when trapping was terminated at dusk. Following transport to the laboratory, live horse flies were dissected in search of third-stage larvae of *E. schneideri* (Couvillion et al., 1984) and larvae were counted.

Received for publication 26 October 1984.

¹ Southeastern Cooperative Wildlife Disease Study, Department of Parasitology, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA.

² Coastal Plain Experiment Station, Department of Entomology, University of Georgia, Tifton, Georgia 31793, USA.

³ South Carolina Wildlife and Marine Resources Department, Tom Yawkey Wildlife Center, South Island Plantation, Route 2, Georgetown, South Carolina 29440, USA.

⁴ Department of Parasitology, College of Veterinary Medicine, University of Georgia, Athens, Georgia 30602, USA.

⁵ Present address: College of Veterinary Medicine, Drawer V, Mississippi State University, Mississippi State, Mississippi 39762, USA.

TABLE 1. Numbers of female *Tabanus lineola hinellus* collected, examined, and found positive for third-stage larvae of *Elaeophora schneideri* and intensity of infection.

Date	Number of horse flies			Number of larvae in infected horse flies	
	Collected	Examined	Positive (%)	Average	Total
1981					
6 Jul ^a	130	130	0 (0)	0	0
4 Aug	69	66	1 (1.50)	15	15
8 Sep	745	517	0 (0)	0	0
29 Sep	1,293	1,276	4 (0.31)	24	95
Totals/means	2,237	1,989	5 (0.25)	22	110
1982					
13 Apr ^b	0	0	0 (0)	0	0
21 Apr	0	0	0 (0)	0	0
30 Apr	0	0	0 (0)	0	0
4 May	61	51	0 (0)	0	0
13 May	170	163	2 (1.23)	5	10
22 May	321	314	0 (0)	0	0
8 Jun	1,970	1,940	13 (0.67)	4	55
22 Jun	477	456	1 (0.22)	1	1
6 Jul	173	158	0 (0)	0	0
20 Jul	525	518	0 (0)	0	0
3 Aug	86	81	0 (0)	0	0
17 Aug	253	246	3 (1.22)	15	45
31 Aug	2,086	2,072	2 (0.10)	3	6
14 Sep	1,492	1,473	4 (0.27)	38	152
28 Sep	648	637	1 (0.16)	51	51
12 Oct	452	442	0 (0)	0	0
Totals/means	8,714	8,551	26 (0.30)	12	320
Overall totals/means	10,951	10,540	31 (0.29)	14	430

^a First day of consecutive 3-day collection period.

^b First day of consecutive 2-day collection period except for 21 April through 22 May which were 1-day collection periods.

To confirm the presence of infected deer on the study area, 12 deer were examined at necropsy in search of *E. schneideri*. Samples of forehead skin were examined histologically for microfilariae. Representative specimens of adult *E. schneideri* and histologic sections of skin with microfilariae were deposited in the U.S. National Parasite Collection, Beltsville, Maryland (Accession Nos. 78487, adults; 78488, histologic sections).

RESULTS

Data on *T. l. hinellus* collected and examined for third-stage larvae of *E. schneideri* are summarized in Table 1. Overall, 10,951 *T. l. hinellus* were captured, and 10,540 were dissected for third-stage larvae (Couvillion et al., 1984). Sim-

ilar proportions of infected horse flies were captured at different ($P > 0.05$) traps.

During 1982, *T. l. hinellus* were present during all trapping periods over 22 wk from May through mid-October (Table 1; Fig. 1). Following the first appearance of *T. l. hinellus* on 4 May, the population increased rapidly, reaching a peak on 8 June (Fig. 1). Abundance then decreased until late August/early September, when there was another pronounced increase in numbers of horse flies captured. Horse flies were not trapped during May and June in 1981; therefore a complete assessment of temporal changes in abundance cannot be made for that year.

Overall, 0.29% of *T. l. hinellus* har-




FIGURE 1. Temporal distribution of the abundance of *T. l. hinellus*, percent infected with third-stage larvae, and total number of third-stage larvae.

bored third-stage larvae (Couvillion et al., 1984). Average prevalence of infected *T. l. hinellus* was 0.25% for 1981 and 0.30% for 1982. Prevalence of infected *T. l. hinellus* varied among collection periods and ranged from 0 to 1.5% and 0 to 1.23% during 1981 and 1982, respectively (Fig. 1). Prevalence of infected *T. l. hinellus* was similar among collection periods for 1981 ($P > 0.05$, $\chi^2 = 6.028$); however, highly significant differences ($P < 0.01$, $\chi^2 = 22.69$) were noted during 1982. In 1982, prevalence of infected horse flies, like horse fly abundance, was biphasic with peaks occurring in mid-May (1.23%) and mid-August (1.22%) immediately preceding peaks in *T. l. hinellus* populations.

The mean number of third-stage larvae in infected *T. l. hinellus* was 14 for the two seasons combined, ranging from one to 47 ($\bar{x} = 22$) in 1981 and from one to 64 ($\bar{x} = 12$) in 1982 (Couvillion et al., 1984) (Table 1). During 1982 the intensity of infection of *T. l. hinellus* increased as time progressed ($r = 0.644$). A quadratic regression model fit the data more precisely ($r = 0.728$) and indicated that the intensity of infection varied non-linearly with time and increased to the highest level in late summer. This increase resulted in greater recovery of infective larvae during August–September (mean intensi-

ty = 25) than in May–June (mean intensity = 4) (Fig. 1).

Examinations for adult *E. schneideri* were done on 12 white-tailed deer over the 2-yr period. Of 10 deer collected in May and June of 1981, eight harbored adult *E. schneideri* and two were positive for microfilariae on histologic examination. In 1982, two deer were examined for adult *E. schneideri* in June and October. Microfilariae were found only in the deer examined in June; however, only a single dead adult worm was found in the carotid artery of the deer examined in October. Overall, infected deer harbored one to 31 ($\bar{x} = 6.5$) adult nematodes. The intensities of infection with adult *E. schneideri* in deer with and without microfilariae were nine to 31 ($\bar{x} = 18$) and one to three ($\bar{x} = 1.5$), respectively. Few microfilariae were seen in deer that were positive. Four deer without microfilariae harbored one nematode each.

DISCUSSION

The production of two generations of adult *T. l. hinellus* on South Island in 1982 was similar to the situation for *T. l. hinellus* in coastal Texas (Thompson, 1973) and *T. l. lineola* in Alabama (Burnett and Hays, 1977) and South Carolina (Sheppard, 1972). This is in contrast to most

temperate climate species of horse flies that produce only one generation per year (Harwood and James, 1979).

Differences among trapping periods in prevalence of infected horse flies were attributed in part to fluctuations in the ratio of blood-fed (potentially infected) to unfed horse flies. For instance, during peaks in *T. l. hinellus* populations, the ratio of blood-fed to unfed horse flies would be small because large numbers of newly emerged *T. l. hinellus* were attracted to traps while seeking their first blood meal.

Although deer were not examined to determine whether there was a seasonal change in numbers of microfilariae in forehead skin, the temporal change in the intensity of infection of horse flies raises the question of a circannual rhythm of microfilariae in deer, with highest levels in late summer. Circannual rhythms of microfilariae have been reported for *Onchocerca gutturosa*, a filarial parasite of cattle (*Bos taurus*) in which microfilariae reside in skin (Eichler, 1973).

Compared to primary intermediate hosts of the arterial worm in the western United States, the prevalence and intensity of infection of *T. l. hinellus* on South Island were low. The average prevalence of infection for *Hybomitra* spp. was as high as 19.1% in one area of New Mexico (Hibler et al., 1971). Average prevalences of *H. laticornis* and *H. aatos* with third-stage larvae in New Mexico were 8.8% and about 2.0%, respectively (Clark and Hibler, 1973; Davies, 1979). The mean intensity of infection of *T. l. hinellus* with third-stage larvae was less than half of that reported for infected horse flies in the western United States by Clark and Hibler (1973), and only about one-fourth of that of *H. aatos* (Davies, 1979). The low prevalence and intensity of infection of *T. l. hinellus* probably were due to the low prevalence and few microfilariae of patently infected deer and/or to biological and behavioral characteristics of horse flies

that affect uptake and development of microfilariae.

ACKNOWLEDGMENTS

Special thanks are extended to Dr. K. H. Pollock, Southeastern Fish and Game Statistics Project, Institute of Statistics, North Carolina State University, for assistance with statistical analysis. The assistance of our colleagues, especially Mr. E. A. Rollor, during the field phase of this study was greatly appreciated. This study was supported by an appropriation from the Congress of the United States. Funds were administered and research coordinated under the Federal Aid in Wildlife Restoration Act (50 Stat. 917) and through Contract Numbers 14-16-0009-82-015 and 14-16-0004-83-004, Fish and Wildlife Service, U.S. Department of the Interior. Further support was provided by the Yawkey Foundation, Tom Yawkey Wildlife Center, South Island, South Carolina.

LITERATURE CITED

- BURNETT, A. M., AND K. L. HAYS. 1977. Seasonal and diurnal distributions of adult female horse flies (Diptera, Tabanidae) at Gold Hill, Alabama. Ala. Exp. Stn. Circ. 237, 28 pp.
- CLARK, G. G. 1972. The role of horse flies (Diptera: Tabanidae) in the transmission of *Elaeophora schneideri* Wehr and Dikmans, 1935, in the Gila National Forest, New Mexico. M.S. Thesis. Colorado State Univ., Fort Collins, 169 pp.
- , AND C. P. HIBLER. 1973. Horse flies and *Elaeophora schneideri* in the Gila National Forest, New Mexico. J. Wildl. Dis. 9: 21–25.
- COUVILLION, C. E., D. C. SHEPPARD, V. F. NETTLES, AND O. M. BANNAGA. 1984. Intermediate hosts of *Elaeophora schneideri* Wehr and Dikmans, 1935 on South Island, South Carolina. J. Wildl. Dis. 20: 59–61.
- DAVIES, R. B. 1979. The ecology of *Elaeophora schneideri* in Vermejo Park, New Mexico. Ph.D. Thesis. Colorado State Univ., Fort Collins, 216 pp.
- EICHLER, D. A. 1973. Studies on *Onchocerca gutturosa* (Neumann, 1910) and its development in *Simulium ornatum* (Meigen, 1818). 3. Factors affecting the development of the parasite in its vector. J. Helminthol. 47: 73–88.
- EPSTEIN, M. B. 1983. Mortality and movements of white-tailed deer fawns on two coastal islands in South Carolina. M.S. Thesis. Frostburg State College, Frostburg, Maryland, 89 pp.
- HARWOOD, R. F., AND M. T. JAMES. 1979. Entomology in Human and Animal Health. Macmillan Publ. Co., New York, 548 pp.

- HIBLER, C. P., G. H. GATES, R. WHITE, AND B. R. DONALDSON. 1971. Observations on horseflies infected with larvae of *Elaeophora schneideri*. *J. Wildl. Dis.* 7: 43-45.
- , AND A. K. PRESTWOOD. 1981. Filial nematodes of white-tailed deer. *In Diseases and Parasites of White-tailed Deer*, W. R. Davidson et al. (eds.). Tall Timbers Research Station, Tallahassee, Florida, pp. 351-362.
- ROBERTS, R. H. 1976. The comparative efficiency of six trap types for the collection of Tabanidae (Diptera). *Mosq. News* 36: 530-535.
- SHEPPARD, D. C. 1972. Distribution and seasonal occurrence of Tabanidae along a transect of South Carolina. M.S. Thesis. Clemson Univ., Clemson, South Carolina, 130 pp.
- THOMPSON, P. H. 1973. Tabanidae (Diptera) of Texas. I. Coastal marsh species, West Galveston Bay; incidence, frequency, abundance and seasonal distribution. *Proc. Entomol. Soc. Wash.* 75: 359-364.