



Crassula campestris (Eckl. & Zeyh.) Endl. (Crassulaceae), a new record for the Italian flora

Authors: Brullo, Salvatore, Campo, Giovanni, Marcenò, Cosimo, Romano, Salvatore, and Siracusa, Giuseppe

Source: Willdenowia, 28(1/2) : 53-58

Published By: Botanic Garden and Botanical Museum Berlin (BGBM)

URL: <https://doi.org/10.3372/wi.28.2804>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

SALVATORE BRULLO, GIOVANNI CAMPO, COSIMO MARCENÒ, SALVATORE ROMANO & GIUSEPPE SIRACUSA

***Crassula campestris* (Eckl. & Zeyh.) Endl. (*Crassulaceae*), a new record for the Italian flora**

Abstract

Brullo, S., Campo, G., Marcenò, C., Romano, S. & Siracusa, G.: *Crassula campestris* (Eckl. & Zeyh.) Endl. (*Crassulaceae*), a new record for the Italian flora. – Willdenowia 28: 53-58. 1998. – ISSN 0511-9618.

Crassula campestris, a species widespread in S Africa, was discovered in Sicily near Palermo, which is a new record for the Italian flora. A description and illustration of *C. campestris* is provided and the species is compared with *C. tillaea* and *C. basaltica*, the two other *Crassula* species in Sicily.

In the course of a floristic investigation in N Sicily some populations of a peculiar *Crassula* species were found in 1990 and recollected in 1996. According to the literature (Bolòs & Vigo 1984, Webb & Akeroyd 1993, Fernandes 1997) these specimens were identified as *C. campestris* (Eckl. & Zeyh.) Endl., which is widespread in S Africa (Fernandes 1978, Toelken 1985). Up to now, this species had been recorded in the Mediterranean area only from some localities in Spain (Vigo & Terradas 1969, Cardona & al. 1976, Franquesa 1985, Velayos & al. 1990, Mateo Sanz & Aguilera Palasi 1990, Fernandez 1997). The occurrence of *C. campestris* in Sicily thus represents a new record for the Italian flora.

Crassula campestris (Eckl. & Zeyh.) Endl. in Walpers, Repert. Bot. Syst. 2: 253. 1843. – Fig. 1
≡ *Tetraphyle campestre* Eckl. & Zeyh., Enum. Pl. Afric. Austral. 3: 294. 1837.
Type: Cape, Swartkops River, *Ecklon & Zeyher 1873* (FI, G, K, S, SAM, TCD, fide Toelken 1985)
= *Crassula pentandra* subsp. *catalaunica* Vigo & Terradas in Acta Geobot. Barcinon. 4: 21. 1969.

Description

Annual herb, with erect to ascending branches, 2-8(12) cm high, glabrous greenish-yellow to purplish-green. Stem simple to branched, with apex sometimes curved. Leaves fleshy, lanceolate to lanceolate-subulate, 3-6 × 1-2 mm, bluntly acute, sessile, smooth, opposite, abruptly constricted at the base, connate, the fused portion 0.25 mm wide. Inflorescence thyrsoid with flowers arranged in sessile or subsessile dichasia. Bracts 1.5-2 mm long, fleshy, ovate-lanceolate, apiculate. Flowers 5-merous, subsessile or shortly pedicellate. Calyx ovoid, with lobes narrowly triangular to lanceola-




Fig. 1. *Crassula campestris* (Eckl. & Zeyh.) Endl. – A: habit; B: detail of the inflorescence; C: closed flower; D: opened flower; E: leaves; F: sepal; G: petal; H: follicle; I: seed; J: epipetalous nectarial squama; K: stamen; L: bract.




Fig. 2. Geographical distribution of *Crassula campestris* in the Mediterranean area.

te, 1.5-1.8 mm long, fleshy, green, often dorsally lightly tuberculate, fused at the base 0.2-0.3 mm high, bordered by a narrow hyaline margin, acute and drawn into an awn. Petals triangular-lanceolate, free or rarely shortly connate, violet-purplish, 0.7-1.2 mm long, acuminate, recurved above, protruding laterally between the sepals. Stamens 5, 0.4-0.6 mm long, filaments white to purplish, anthers 0.15 mm long, ovoid, yellow to purplish yellow. Epipetalous nectarial squamae 0.2-0.3 mm long, subcuneate below the apex. Carpels 5, erect, with a short style. Follicle yellowish with two seeds lightly striate, brown, ellipsoid, 0.3×0.2 mm.

Specimina visa

SICILY: Pendiçi nord-orientali di Monte Catalfano, in contrada Vignazza (Bagheria), 10.5.1990, Marcenò (PAL); *ibid.*, 24.6.1996, Marcenò, Romano & Campo (B, CAT, FI, PAL).

Ecology and distribution

Crassula campestris was collected on the NE slope of Mount Catalfano near Palermo (N Sicily), where it grows along the shoulder of paths near the sea on sandy soils mixed with minute calcareous debris. In this man-made habitat, *C. campestris* is a member of an ephemeral community characterized by numerous microphytes, such as *Trifolium suffocatum* L., *Crassula tillaea* Lest.-Garl., *Campanula erinus* L., *Centaurium pulchellum* Sw., *Tuberaria guttata* (L.) Fourr., *Vulpia ciliata* Dumort., *Valantia muralis* L., *Galium divaricatum* Lam. and *Oglifa gallica* (L.) Chrtk & Holub. Phytosociologically, this vegetation is comparable with the Polycarpo-Crassuletum *campestris* association of the Tuberarietea *guttatae* described by Franquesa (1995) from Cap de Creus in Spain. Presently, the Sicilian record represents the easternmost locality of this species (Fig. 2). According to Greuter & al. (1986) and Webb & Akeroyd (1993), *C. campestris* must be considered in the Mediterranean area as a xenophyte, which – in view of its scattered distribution – is probably naturalized for a long time.

Identification

Crassula campestris belongs to *C. sect. Glomeratae* Haw. due to its herbaceous annual habit, the sessile, connate leaves, the thyrsoid inflorescence with several contracted dichasia subten-




Fig. 3. Morphological differences between *Crassula campestris* (A₁-G₁), *C. basaltica* (A₂-G₂) and *C. tillaea* (A₂-G₃). – A: closed flower, B: opened flower; C: leaf; D: sepal; E: petal; F: stamen and epipetalous nectarial squama; G: seed.

ded by leaf-like bracts, the calyx longer than the corolla and 0.1-0.15 mm long anthers (Toelken 1977). Within the section it can be grouped together with, among others, *C. tillaea* Lester-Garland and *C. basaltica* Brullo & Siracusa, which both occur in the Mediterranean area.

The closest resemblance is with *C. basaltica* (Fig. 3), which is also characterized by 5-merous flowers and fleshy leaves. From this species *C. campestris*, however, differs markedly by the following features: its leaves being green to purplish-green, sessile, smooth, abruptly constricted at the base, lanceolate to lanceolate-subulate, connate at the base for 0.25 mm *versus* glaucous-green, petiolate, papillose-verrucose, gradually constricted towards base, spatulate-lanceolate, connate at the base for 0.1-0.2 mm (Fig. 3C); its calyx being ovoid, with narrowly triangular to lanceolate, 1.5-1.8 mm long, often dorsally lightly tuberculate lobes fused at the base and bordered by a narrow hyaline margin *versus* urceolate, with ovate-lanceolate, 1.2-1.3 mm long, above papillose, free lobes without hyaline margin (Fig. 3A-B, D); the petals being triangular-lanceolate, 0.7-1.2 mm long, curved above, protruding laterally between the sepals *versus* lanceolate, 0.6-0.7 mm long, erect and included (Fig. 3A-B, E); the stamens being 0.4-0.6 mm long, the anthers ovate, yellow to yellow-purplish and the epipetalous nectarial squamae subcuneate at the apex *versus* the stamens 0.3-0.4 mm long, the anthers suborbicular and purplish and the epipetalous nectarial squamae suborbicular at the apex (Fig. 3F); the seeds being slightly striate, 0.3 mm long *versus* smooth and 0.25 mm long (Fig. 3G).

Crassula campestris also resembles *C. tillaea* but differs in both leaf shape and flower features (Fig. 3). The latter species can easily be recognized by the triangular-lanceolate, non-fleshy leaves, which are 2-2.4 mm long and connate for 0.5-0.6 mm, the 3-4-merous flowers, cylindrical calyx with ovate-lanceolate, free and up to 1.5 mm long sepals, the ovate-lanceolate and apiculate petals, only 0.1 mm long anthers, transversely elliptical epipetalous nectarial squamae and the minutely striate seeds.

The Sicilian *Crassula* species can be keyed out as follows:

1. Flowers usually 3-merous (rarely 4-merous); leaves not fleshy, connate for 0.5-0.6 mm; anthers 0.1 mm long; nectarial squamae absent or, if present, transversely elliptical at the apex; seeds minutely striate *C. tillaea*
- Flowers 5-merous; leaves fleshy, slightly tuberculate, connate for 0.1-0.25 mm; anthers 0.15 mm long; nectarial squamae present, suborbicular or subcuneate at the apex; seeds smooth or slightly striate 2
2. Leaves smooth, lanceolate to lanceolate-subulate, abruptly throttled at the base; calyx ovoid, 1.5-1.8 mm long with lobes fused at the base; petals 0.7-1.2 mm long, protruding laterally; anthers ovate; nectarial squamae subcuneate at the apex *C. campestris*
- Leaves papillose-verrucose, spatulate-lanceolate, petiolate, gradually restricted towards the base; calyx urceolate, 1.2-1.3 mm long with free lobes; petals 0.6-0.7 mm long, included; anthers suborbicular, nectarial squamae suborbicular at the apex *C. basaltica*

Acknowledgements

Financial support by the Italian M.U.R.S.T. (40%, 60%) is gratefully acknowledged.

References

- Bolòs, O. & Vigo, J. 1984: Flora dels països Catalans **1**. – Barcelona.
- Brullo, S. & Siracusa, G. 1994: *Crassula basaltica* (*Crassulaceae*), a new species from Mt. Etna (Sicily). – *Fl. Medit.* **4**: 175-178.
- Cardona, M. A., Terradas, J. & Vigo, J. 1976: Sobre la planta del genero *Crassula descubierta* en el Baix Llobregat. – *Collect. Bot. (Barcelona)* **10(6)**: 143-146.

- Fernandes, R. B. 1978: *Crassulaceae africanae novae vel minus cognitae*. – Bol. Soc. Brot., Sér. 2, **52**: 165-220.
- 1997: *Crassula* L. – Pp. 99-103 in: Castroviejo, S., Aedo, C., Laínz, M., Morales, R., Muñoz Garmendia, F., Nieto Feliner, G., Paiva, J. (ed.), *Flora iberica* **5**. – Madrid.
- Franquesa, T. 1985: Notes floristiques II: Aportacions a la flora catalana. – Collect. Bot. (Barcelona) **16(1)**: 239-240.
- 1995: El paisatge vegetal de la península del Cap de Creus. – Arxius Secc. Ci. Inst. Estud. Catalans **109**.
- Greuter, W., Burdet, H. M. & Long, G. 1986: Med-checklist **3**. – Genève & Berlin.
- Mateo Sanz, G. & Aguilera Palasi, A. 1990: Aportación al conocimiento fitogeográfico de la Sierra del Espadán (Castellón). – Folia Bot. Misc. **7**: 67-80.
- Toelken, H. R. 1977: A revision of the genus *Crassula* in Southern Africa. – Contrib. Bolus Herb. **8**.
- 1985: *Crassulaceae*. – In: Leistner, O. A. (ed.), *Flora of Southern Africa* **14**. – Pretoria.
- Velajos, M., Carrasco, M. A. & Monge, C. 1990: Dos *Crassulaceae* de Ciudad Real (España). – Anales Jard. Bot. Madrid **47**: 53-58.
- Vigo, J. & Terradas, J. 1969: Sobre la vegetación de la zona de Acontillados triásicos del Baix Llobregat. – Acta Geobot. Barcinon. **4**.
- Webb, D. A. & Akeroyd, J. R. 1993: *Crassula* L. – Pp. 422-423 in: Tutin, T. G., Burges, N. A., Chater, A. O., Edmondson, J. R., Heywood, V. H., Moore, D. M., Valentine, D. H., Walters, S. M., Webb, D. A. with assist. of Akeroyd, J. R. & Newton, M. E. (ed.), *Flora europaea*, ed. 2, **1**. – Cambridge, etc.

Addresses of the authors:

- Prof. S. Brullo & Dr G. Siracusa, Dipartimento di Botanica, Università di Catania, Via A. Longo, 19, I-95125 Catania, Italy; fax: +39 95 441209.
- Dr G. Campo, Prof. G. C. Marcenò & Dr S. Romano, Dipartimento di Scienze Botaniche, Università di Palermo, Via Archirafi, 38, I-90123 Palermo, Italy; fax +39 91 6176089.