

PLATE 1598.

MELIOSMA SQUAMULATA, Hance.

SABIACEAE.

M, squamulata, Hance in *Journ. Bot.* 1876, p. 364; foliis simplicibus longe petiolatis ovaKbas ellipticis obtuse caudato-acuminatis basi in petiolum cuneatim angustatis integerrimis, supra glabris, subitus opacis pallidioribus minutissime lepidoto-canis, venis primariis utrinqne 3-5 arcuatim anastomosantibus, paniculis terminalibus solitariis v. 3-4-nis ferrugineo-pubescentibus, bracteis parvis ovatis hirsutis, sepalis ovato-rotundatis ciliolatis, petalis glabratis, filamentis antheriferis dorso squama bifida lobis divaricatis acutis adnatis, ovario glabro.

HAB. Hongkong, Wongneichung, Lamont ex Hance, Victoria Peak, O. Ford (No. 106).

Folia 2^{^-}3 poll, longa; petiolus ^-1[^] poll, longus.—D. OLIVER.

Fig. 1. Bud. 2. Antheriferous stamen, front and back. 3. Staminode, inserted on base of petal. 4. Ovary and disk. 5. Transverse section of same. Enlarged.

PLATE 1599.

HUTCHINSONIA PERPUSILLA, ffenul

CRUCIFER E. Tribe THLASPIDEA.


H. perpusilla, *Semsl.* (*sp. nov.*) ; herba annua, glaborrima, gracillima, sajpiissime triplo minor quam in icone, foliis radicalibus **rosulatis** oblongia vel rotundatis **integris** dentatis vel anlobatis, fioribns paucis minntissimis, petalis spathulatis filamentis filiformibus, capsulis ovoidia vel obovoideis, cellulis G-spermis, Bemnibua biseriatis ptmdulia.

HAB. Gilgic : Killa Panja, at 9000 feet, Dr. Giles.

Li the 'Genera Plantarum*' **HvtckinBta** ia limited to *H. petrcea*, which is usually branched from the base, lias pinnatetr divided leaves and two seeds in each cell of the capsule. Perhaps it would be better to refer *Iberidell-a Andersonii*, Hook, f., and T. Thorns, hero, and the whole of Boissier's species of *Iberidella* to *Jblthiunema*[^] as he has done.—W. B. HEMSLV.

Fig. 1. Upper part of inflorescence. 2. Sspal. i. Petal. 4. Stamens, ft and 6. Capsules. 7. Capsalo with one of the valves removed. 8. An immature seed.
Phlifirfd.





M.S.del.

Beruiettia longipes, Oliv

PLATE 1597.

EMBELIA KIBES, *Burm.* var. *penangiana*.

MYRSINACEAE.

E. *JLihes, Bwm. (DO. Prodr Mn, j). 85).* vtt, *pmangiana*; foliis oblongo-oblanceolatis breviter acuminatis basi cuneatim angustatis tenuiter coriaceis glabris, pauculis purpurasceni-puberulis.

HAB. Penang, *C. Curtis* (No. 306).

Folia 4-4¹ poll, longa, 1¹-1¹ poll, lata; petiolas \ poll, longus. *Panicula terminalis ampla* 6-10 poll, longa. *Baccm* obovoideo-globosae corrugato-tuberculatae 1¹ lin. longae; pedicelli fructiferi bacca duplo breviores.

At first sight this plant looks very different from the common *E. Ribes*, *Burm.*, in the attenuation of the leaf lamina into the petiole, but it scarcely merits specific separation as I at first thought it might.—
D. OLIVER.

Fig. 1. ^Estivation of corolla-lobes. 2. Bud. 3. Expanded flower. 4. Part of corolla laid open. 5. Pistil. 6. Longitudinal, and 7. transverse, section of ovary. 8. Fruit. 9. Section of same. 10. Embryo. *Enlarged*.



MS, del.

Embelia Ribes, Burm. var. penaiiana.

VOL.	TAD.	VOL.	TAB.
BOLAX G-lebaria, <i>Comm.</i> . . .	y. 492	CEANOTHJS papillosum, <i>Torr.</i> <i>et Gr.</i> . . .	iii. 272
.BOLETUS Emodeusis, <i>Berk.</i> . . .	ix. 870-IA	CELASTBUS Magellanicus, * <i>De Cand.</i> . . .	vi. 0 537
BOBONIA nana, <i>Hook.</i> . . .	iii. 270	subspicatus, <i>Hook.</i>	v. 482
rhomboidea, <i>Hook.</i> . . .	viii. 722	CENTBOSOLENIA glabra, <i>Hook.</i>	ix. 873
BOW&ANNIA verbascifolia, <i>Gardn.</i> . . .	vi. 519-20	CEBCOCABPUS betulsfolius, <i>Nutt.</i> . . .	iv. 322
BRACHYHYNITJM acuminatum, <i>Harv.</i> . . .	i. 19jf. 3	<i>ledifolius</i> , <i>Nutt.</i> . . .	9 324
microstomum, <i>Harv.</i> / . 4	<i>parvifolius</i> , <i>Nutt.</i> . . .	m 323
Bplachnoides, <i>Harv.</i> , "f. 2	CEBOPEGIA attenuata, <i>Hook.</i>	ix. 867
Weissia, <i>Hook.</i> f. 1	CESTBUM vestitum, ² <i>Hook.</i>	iv. 381
BBACHYBIS microcepala, <i>De Cand.</i> . . .	ii. 147	CHABBAEA suaveolens, <i>De Cand.</i>	v. 496
ramosissima, <i>Hook.</i> 142	CHAILLETIA cymosa, <i>Hook.</i>	vi. 591
BBOOKIA brevipes, <i>Harv.</i> . . .	iii. 231	floribunda, <i>Planch.</i>	viii. 792
BBUGUIEBIA Malabarica, <i>Am.</i> iv.	398	CHABA latifolia, <i>Willd.</i> . . .	vi. 532
<i>Bhaedei</i> , <i>Arn.</i> 3 9 7	CHEILANTHES chrysophylla, <i>Hook.</i> . . . f	x. 901
BBYUM nitens, <i>JfboA.</i> . . .	i. 19 / . 6	<i>monticola</i> , <i>Gardn.</i> . . .	v. 487
pellucens, <i>Hook.</i> 34	<i>nitidula</i> , <i>Hook.</i> . . .	x. 912
teretiusculum, <i>Hook.</i> 20 / . 1	<i>ochracea</i> , <i>Hook.</i> 904
BUBKEA Africana, > <i>Hook.</i> . . .	vi. 593-4	CHEIBANTHEBA linearis, <i>A. Cunn.</i> . . .	i. 47
CABOMBA Caroliniana, <i>A. Gray.</i> vii.	642	CHILIOTBICHUM amelloides, <i>Cass.</i> . . .	v. 485
<i>Piaubyensis</i> , <i>Gardn.</i> 641	CHBISTISONIA Stocksii, <i>Hook.</i>	ix. 836
CADABA beterotricba, <i>Stocks.</i> ix.	839	CUBTSOBACTBON HookeH, <i>Colenso.</i> . . .	• * 817
CALAKDBINA calyprata, <i>J. D. Hook.</i> . . .	iii. 296	CLADOCAUXON Brasihense, <i>Gardn.</i> . . .	vi. 528
CALCEOLARIA Sinolairii, <i>Hook.</i> vi.	561	CLADONIA perfolata, <i>Hook.</i> . . .	ii. 192
CALDASIA argentea, <i>J. D. Hook.</i> iii.	300	CLAYIJA spatimlata, <i>It. et P.</i> . . .	• 140
CALLIXENE parviflora, <i>Hook.</i> <i>fil.</i> . . .	vii. 632	CLAYTONIA Australasica, <i>J. D. Hook</i>	iii. 293
<i>polyphylla</i> , <i>Hook.</i> 674	CLEISTANTHUS polystachyus, <i>Hook. fil. %</i> . . .	viii. 779
CAMPANULA Jacobaea, <i>Ch. Sm.</i> viii.	772	CLEISTES speciosa, <i>Gardn?</i> . . %	v. 474
<i>Vidalii</i> , <i>Watson.</i> . . .	vii. 684	CLEMATIS anetbifolia, <i>Hook.</i> <i>Bojeri</i> , <i>Hook.</i> . . .	i. 78
CAMPYLANTHEBA Fraseri, ¹ <i>Hook.</i> . . .	i. 82	<i>oligophylla</i> , <i>I&olc.</i> . . .	• 10
CAMPYLANTHUS Benthami, <i>Webb.</i> . . .	viii. W6	<i>pimpinellifolia</i> , <i>Hook.</i> . . .	80
^JAPPAEIB brevispina, <i>W. et Arn.</i> ii.	126	<i>Staitieyii</i> , <i>Hook.</i> . . .	77
<i>diversifolia</i> , <i>TT. et Arn.</i> . . .	181	<i>trifida</i> , <i>Hook.</i> . . .	i. 589
<i>incanescens</i> , <i>JDe Cand.</i> . . .	123	CLEOMELLA Mexicana, <i>D.C.</i> t	28
<i>pedunculosa</i> , <i>Wall.</i> . . .	128	CLITOBLIA viridiflora, <i>Bout.</i> . .	152
CABDAMINE corymbosa, <i>Hook.</i> <i>fil.</i> . . .	vii. 686	COCOINIA Indica, <i>W. et Arn.</i> ii.	JL38
<i>heterophylla</i> , ² <i>Hook*</i>	i. 58	COCCULUS macranthus, <i>Hook. fil.</i> viii.	^59
<i>intermedia</i> , <i>fTooA.</i> . . .	iii. 258	CoCHLEABIA flava ⁴ , <i>Buck.</i> . .	ix. 805
<i>radicata</i> , <i>Hook. fil.</i>	ix. 882	CODONAKTHUS alternifoHa, <i>Planclu</i> . . .	viii. 796
CABEX filifolia, <i>Nult.</i> . . .	v. 448	COLLETIA discolor, <i>Hook.</i> . .	vi. 538
CAPPODETUS serratus, <i>Forst.</i> vi.	564	COMBBETUM salicifolium, <i>E. Afey.</i> . .	592
CASSEBEEBA gleichenioides, <i>Gardn.</i> 507	CONDALIA obovata, <i>Hook.</i> . .	iii. 287
CASSINE Maurocenia, <i>JFilld.</i> . . .	552	COHOHOBIA castaneoefolia, <i>A. St. Hil.</i> . . .	i. 63
CASSIPOUBEA elliptica, <i>Poir.</i> iii.	280	CONOSTYLIS vaginata, <i>Endl.</i> . .	ix. \$74
CATASETUM cristatum, var. <i>monstrosum</i> , <i>Hook.</i> . . .	ii. 177		
CAVALLIA einuata, ³ <i>Hook.</i> . . .	iii. 252		

¹ Is *Pronaya elegant*, Hugel.² Probably a form of *C. hirsuta*, Linn, which seems to occur all over the world.³ This undoubtedly belongs to the Nat. Ord. Loasae.¹ Is *Maytenus Magellanicus*, Hook. fil.² Is *Sessca vscita*, Miers.

	VOL. TAB.		VOL. TAB.
CONBADIA calycosa, <i>Hook.</i> .	vii. 689-90	DOUGKLASIA nivalis, <i>Lindl.</i> .	ii. 150
neglecta, <i>Hook.</i> .	ix. 874	DBABA deutata, <i>Hook. et Arn.</i> .	i. 31
COPTOPHYLLUM buniifolium,		violacea. <i>X>. C.</i> . . .	35
<i>Gardn.</i> . . . v. 477		DBACOPHYLLUM Milligani,	
millefolium, <i>Gardn.</i> .	478	<i>Hook.</i> . . . ix. 845	
COBOKIA buddlaeoides, -4. CWw. .	424	PBAFETES ericoides, <i>Hook. fil.</i> .	895
CoBB-EA Backhousiana, . . . i. 2		DBIMYS axillaris, <i>JPorst.</i> . . vi. 576	
fexruginea, <i>Backh.</i> .	3	piperita, <i>Hook. fil.</i> . ix. 896	
COTTILA myriophylloides, <i>J5for.</i> iv.	335	DBOSEBA Arcturi, <i>Hook.</i> . . i. 56	
CBATJSVA Koxburghii, <i>Br.</i> . . . ii. 178		bulbosa, <i>Hook.</i> . . iv. 375	
CEEMOLOBTJS Peruvianus, <i>D.C.</i> . . . i.	48	lunata, <i>Buchau.</i> . . i. 54	
pinnatifidus, <i>Hook.</i> .	100	xnacrophylla, <i>IAndl.</i> . iv. 376	
pubescens, <i>Hook.</i> . . .	81	Menziesii, <i>Br.</i> . . i. 53 *	
rhomboideus, <i>Hook.</i> .	32	Btolonifera, <i>Endl.</i> . iv. 389	
sinuatus, <i>Hook.</i> . . .	99	DEYMOGLOSSUM rigidum, <i>Hook.</i> x. 996	
CBOSSOLEPIS? pusilla, <i>Hugel.</i> . . . v.	413	EABINA mucronata, <i>IAmll.</i> . . v. 431	
CBOTAXABIA bupleurifolia,		ECHITES pulchella, <i>Gardn.</i> . . .	470
<i>Schlecht.</i> . . . ir. 372		EcHitiM stenosyphon, <i>Webb.</i> viii. 772	
crassipe9, <i>Hook.</i> . . . ix. 830		EILSOCABPUS coriaceus, <i>Hook.</i> ii. 154	
Cunninghami, <i>JBr.</i> . . .	829	Hinau, A. <i>Cunn.</i> . . vii. 602	
oblongifolia, <i>Hook.</i> .	-830	pubescens, <i>Hook.</i> . . ii. 155	
CRYPTOMEEIA Japonica, <i>Don.</i> . . . vii.	668	ENTOSTHODON latifolius,	
CBYPTONEMIA Forbesii, <i>Haru.</i> . . .	679	<i>J. J.</i> <i>Hook.</i> . . . iii. 245 c	
CYATHEA Beyrichiana, <i>Presl.</i> . . .	623	Mathewsi, <i>J. JD.</i> <i>Hook.</i> . . . 245 B	
<i>Cunninghami, Hook. fil.</i> s. 985		obtueifolius, <i>J. JD.</i>	
<i>c. integra, J. Sm.</i> . . . vii. 838		<i>Hook.</i> 245 A	
<i>Walker®, Hook.</i> . . .	647	EPIDENDBITM microbulbon,	
<i>CYMBIDIITM plicatum, jifar®.</i> . . . ii. 104		<i>Hook.</i> iv. 347	
CYPBIPEDIITM caudatum, <i>IAndl.</i> . . . vii. 658-9		EPIDENDBUM rigidum, <i>Jacg.</i> 314	
CTBTANDBA pendula, <i>JZ.</i> . . . viii. 735-6		EPILOBIUM confertifolium, <i>Hook.</i>	
CYSTOPTERIS Tasmanica, <i>Hook.</i> x. 959		<i>fil.</i> vii. 685	
<i>Douglasii, *Iook.</i> . . .	955	<i>m&cranthon, J. JD.</i>	
CYTINTTS dioicus, <i>Juss.</i> iv. 336		<i>Hook.</i> iii. 297	
DACPBYDIUM Oolensoi, <i>Hook.</i> . . .	548	<i>macropus, Hook.</i> . . . ix. 812	
<i>laxifolium, Hook. fit.</i> ix. 815		<i>melanocaulon, Hook.</i> . . . 813	
PALIBABDIA lobata, <i>Baldiv.</i> . . . i.	76	EEIOGONUM multiflorum,	
DAVALLIA serrsefolia. <i>Wall.</i> . . .	93	<i>Benth.</i> iii. 250*	
DEXUBOMEKOIT rigi" Benth. . .	37	EEIOPE macrostachya, <i>Mart.</i> . . . v. 46 #	
DjiSFONTAINIA spinosa, <i>JZ.</i> < f P. . . ^	33	EBIOSTEMON obcordatum,	
DICBANOI.EPIS distjcha, <i>Planch.</i> viii- 798		<i>A. Cunn.</i> i. 60	
DICBANTJM Jamesoni, <i>Hook.</i> . . . il.	179	EBYNGIUM: humile, <i>Cav.</i> . . . iii. 216	
macrodon, <i>Hook.</i> iv. 319		ESCALLONIA Organensis, <i>Gardn.</i> yi. 514	
DIDISCUS humilis, <i>J. D. Hook.</i> . . .	304	<i>polifolia, Hook.</i> . . . ii. 114	
<i>pilosus, Hugel.</i> . . .	307	<i>serrata, Srn.</i> . . . vi. 540	
DIDYMODON cirrhifolium, Saru. . . i. 18 f. 5		ETABAXLIA G-uineensis, <i>Benth.</i> v. 453-4	
<i>Tortula, -ffar.</i> . . . / . 2		ETJCAKYPTTIS macrocarpa, <i>Hook.</i> . . . 405—7	
<i>vaginatum, Hook.</i> . . . / . 4		<i>maculata, Hook.</i> . . . vii. 619	
DIMOEPOLEPIS australis,		<i>platypus, Hook.</i> . . . ix. 849	
^, Oray. ix. 856		<i>populifolia, Hook.</i> 879	
DIOSCOBABA pusilla, <i>Hook.</i> . . . vii. 678		ETIPHORBIA alata, <i>Hook.</i> . . . vii. 700	
DIOTOSPEBMA Drummondii, A. . .		<i>aphylla, Brouss.</i> . . . ii. 182	
<i>Gray.</i> ix. 855		<i>sphterorhiza, Benth.</i> . iv. 346	
DIPLAZITTM cordioliolum,		ETJPLOCA convolvulacea, <i>Nutt.</i> vii. 651	
<i>Illume.</i> ii. 184		EUTHEMIS leucocarpa, <i>Jack.</i> . viii. 711	
<i>cordifolium, Bl.</i> . . . x. 936		ETTOCA aretioides, <i>Hook. et</i>	
<i>elegans, Hook.</i> 939-40		<i>Arn.</i> iv. 355	
DIPLICOSIA ciliolata, <i>Hook. fil.</i> ix.	894		
DISA ferruginea, <i>Thunb.</i> . . . iii. 214			
DONATIA Magellanica, <i>Lam.</i> . . . i. 16			
BORSTENIA asaroides, <i>Gard.</i> . . . iv. 399			
<i>elata, Gard.</i> . . . iii. 220			

1 Not of Brown. It is now *D. Planchonii* *Hook. fil.*

2 *E. verructntm*, Richard in Voy. de l'As-trolabe, t. 26.

	VOL.	TAD.	VOL.	TAB.
EUTOCA lutea, <i>Hook, et Arn.</i>	iv.	354	GTTATTERIA Maypurensis, <i>Hook.</i>	iii. 227
FABIANA imbricata, <i>Ruiz et Pavon.</i>	iv.	340	GUNNERA Falklandica, <i>Hook.</i>	v. 489-90
FABBONIA tomentosa, <i>Hook, et 9 Wils.</i>	viii.	739 c	GYMNOGRAMMA aspidioides, <i>Hook.</i>	x. . 950
FAGUS cliffortioides, <i>Hook.fil.</i>	vii.	673	aureo-nitens, <i>Hook.</i>	ix. 820
• cliffortioidee, <i>Hook.fil.</i>	ix.	816 B	aurita [^] <i>Hook.</i>	x. 974
fusca, <i>Hook.fil.</i>	vii.	630-1	var /3. <i>Hook.</i>	. . 989
Gunni, <i>Hook, fil.</i>	ix.	881	microphylla, <i>Hook.</i>	. . 916
Menziesii, <i>Hook. fil.</i>	vii.	652	rutsefolia, var. Hispanica, <i>Hook.</i>	. . 935
Solandri, <i>Hook.fil.</i>	. .	639	Schomburgkiana, <i>Kze.</i>	. . 992
Solandri, <i>Hook. fil.</i>	ix.	816 A	vestita, <i>Wall.</i>	. . ii. 115
FABSETIA linearis, <i>Done.</i>	. .	808	GYMNOPSIS uniserialis, <i>Hook.</i>	. . 145
FENZLIA dianthiflora, <i>Benth.</i>	ii.	199	GYMNOPTERIS queratifolia, <i>J.Bernh.</i>	x. . 905
FOBSTERIA bellidifolia, <i>Hook.</i>	ix.	851	semipinnatifida, <i>Fée.</i>	. . 971-2
FRANKEVIA cymbifolia, <i>Hook.</i>	iii.	265	trilobata, <i>J. Sm.</i>	. . 907
FBIESIA racemosa, <i>A. Cunn.</i>	vii.	601	GYMINOSTOMUM cylindricum, <i>Hook.</i>	i. 17 / 2
FBITILLABIA Roylei, <i>Hook.</i>	ix.	860	• rufescens, <i>Hook.</i>	. . f*§
FUCHSIA cordifolia, <i>Benth. /3</i>	v.	450	Betifolium, <i>Hook, et Am.</i>	ii. ^135
dependens, <i>Hook.</i>	i.	65	spatulatum, <i>Harv.</i>	i. 17 f. 1
procumbens, <i>R. Cunn.</i>	v.	421	vernicosum, <i>Hook.</i>	. . f. 4
FUMABIA micrantha, <i>Lag.</i>	iv.	363	GYRINOPS Walla, <i>Gaertn.</i>	. . 5
GARDENIA Vogelii, <i>Hook.fil.</i>	viii.	782-3	ILEMODORUM distichophyllum, <i>Hook.</i>	ix. . 866
GABBYA Fadyenii, <i>Hook.</i>	iv.	333	HAKEA Baxteri, <i>Br.</i>	y. 429-40
GAULTHEBIÄ lanigera, <i>Hook.</i>	i.	66	conchifolia, <i>Hook.</i>	. . 430
tomentosa, <i>H. B. K.</i>	iv.	393	cristata, <i>Br.</i>	. . * 443
GENIOSPOBUM strobilifem, <i>Wall.</i>	v.	462	cucullata, <i>Br.</i>	. . 441
GENIOTOMA ligustrifolium, <i>A. Cunn.</i>	. .	430	heterophylla, <i>\$IooJc.</i>	. . 437
GENTIANA bellidifolia, <i>Hook, fit.</i>	yii.	635	incrassata, <i>Br.</i>	. . 442
detonaa, <i>Fries, var.</i>	ix.	857	intermedia, <i>Hook.</i>	. . 445
Jamcsoni, <i>Hook.</i>	i.	61	pandanicarpa, <i>Br.</i>	. . 434
GERANIUM -neafatum, <i>Hook.</i>	ii.	198	platysperma, <i>Hook.</i>	. . 433
GILIA congesta, <i>Hook.</i>	iii.	235	stenocarpa, <i>Br. T.</i>	. % . 444
GLEICHENIA simplex, <i>Hook.</i>	i.	t£	tricostata, <i>Br.</i>	. . . 435-6
• LOXINIA ichthyostoma, <i>Gardn.</i>	v.	472	undulata, <i>Br.</i>	. . . 447
Sarmientiana, <i>Gardn.</i>	iv.	378	HALQ [^] AGIS cordigera, <i>Jfug.</i>	yi. 598
GLYCYCABFUS racemosa, <i>Date.</i>	ix.	842	HAMAD [^] KJTAS Andicola, <i>Hook.</i>	ii. 137
GLYPH.EA grevioides, <i>Hook.</i>	fil. . . viii	760	HABTIGHSIA spectabilis, <i>Ad. Juss.</i>	vii. 615-16
GLYPHOCABPA Eoylii, <i>Hook.fil.</i>	ii.	1940	HABTHASTNTA puDgens, <i>Hook.</i>	et Am. iv. 334
GOMPHIA Sumatrana, <i>Jack.</i>	viii.	712	HABYEYA Capensis, <i>Hook.</i>	ii. 118
GONIOCABPUS serpyllifolius, <i>J. D. Hook.</i>	iii.	290	purpurea, <i>Harv.</i>	iv. * 3§1
vernicosus, <i>J. J. Hook.</i>	iv.	311	HEIMIA grandiflors [^] <i>Hook.</i>	vi. 554
GOSSYPIANTHUS rigidiflorus, <i>Hook.</i>	iii.	251	HELIOPHYLUM Gunnianum, <i>Hook.</i>	iv. 320
GRAMMITIS Ascensionis, <i>Hook.</i>	x.	967	HELIPTEBUM incanum, <i>De Cand.</i>	318
bisulcata, <i>Hook.</i>	^	998	HEMITELIA alternans, <i>Hook.</i>	vii. 622
Organensis, <i>Gardn.</i>	vi.	509	% Guianensis, <i>Hook.</i>	. . 648
GRATIA polygaloides, <i>Hook, et Am.</i>	iii.	271	Hostmanni, <i>Hook.</i>	. . 646
GBIMMIA Grisebachii, <i>Hook.fil.</i>	vii.	636	Imrayana, <i>Hook.</i>	. . 669
laxifolia, <i>Hook.fil.</i>	ii.	194D	Lindeni, <i>Hook.</i>	. . viii. 706
GTTABA (?) microphylla, <i>Hook.</i>	. .	129	Parkeri, <i>Hook.</i>	. . vii. 643
1 Is Wilsonia humilis, Br.			HEBHANNIA boraginiflora, <i>Hook.</i>	vi. 597
1 Is Axilaya purpurea, Benth.				

	VOL.	TAB.	VOL.	TAB.	
HESPEBOMELES heterophylla, <i>Hook.</i>	ix.	846	IPOMAEA nerifolia, <i>Gardn.</i>	v.	471
HEWABDIA Tasmanica, <i>Hook.</i>		858	IBESIXE grandiflora, <i>Hook.</i>	ii.	102
HIBBERTIA virgata, <i>Br.</i>	iii.	267	ISOPOGOX axillaris, <i>Br.</i>	v.	438
HIBISCUS Scindicus, <i>Stocks.</i>	ix.	802	IXEBBA brexioides, <i>A. Cunn.</i>	vi.	577-8
<i>tulipiflorus, Hook.</i>	viii.	707-8	JAMESONIA cinnamomea, <i>Kze.</i>	viii.	713
HOHEBIA populnea, <i>A. Cunn.</i>	vi.	565-6	JASMINUM lineare, <i>Br.</i>	ix.	831
HOLOTHBYX Harveyana, <i>Lindl.</i> ii.	103 A		JUNCUS Andicola, <i>Hook.</i>	viii.	71*
<i>parvifolia, Lindl.</i>	103 B		KA^DELIA Bheedei, <i>W. et Am.</i>	iv.	362
HOMOANTHTJS echinulatus, <i>Cass.</i> v.	491		KEEBLIA skirrhobasis, <i>De Cand.</i>	iii.	240
HOOXEBA fissidentoides, <i>Hook, et Wils.</i>	viii.	706 A	KBIGIA nervosa, <i>Hook.</i>	m	257
<i>obtusifolia, Harv.</i>	i.	24 / 11	LACHNOSTACHYS aibicans, <i>Hook.</i> v.		414
<i>prostrata, Harv.</i>		20 / 5	<i>ferruginosa, Hook.</i>	m	415
HOSTMANNIA elvasioides, <i>Planch.</i>	viii.	709	LAMBEBTIA illicifolia, <i>Hook.</i>	vi.	553
HUGELIA virgata, <i>Benth.</i>	ii.	200	LANGSDOFPFIA Indica, <i>Wight.</i>	iii.	205-6
HTTGONIA Flanchoni, <i>Hook.fil.</i>	viii.	777	LATUYBUS gladiatus, <i>Hook.</i>	i.	72
HUTCHINSIA Tasmanica, <i>Hook.</i>	ix~	848	LATOISIEBIA lycopodioides, <i>Gardn.</i>	vi.	502
<i>Tibetica, Thorns.</i>		900	LAWBENCIA glomerata, <i>Hook.</i>	v.	417
HYBBOCOTYLE cordifolia, <i>J. D.</i> <i>Hook.</i>	iv.	303	<i>spicata, Hook.</i>	iii.	261-2
<i>tripartite, Br.</i>		312	LEDOCAEPUM Beynoldsii, <i>Hook.</i>	i.	14
HYMENOLEPIS platyrhynchos, <i>Kze.</i>	x.	999	LEIANTHUS umbellatus, <i>Griseb.</i>	vii.	687-8
HYMENOLOBITS divaricatus, <i>Nutt.</i> iii.	277		LEPIDIUM latipes, <i>Hook.</i>	i.	41
HYMIESOPHYLLUM asplenoides, <i>Sw.</i>		956	<i>rotundum, De Cand.</i>	vii.	609
<i>fletosum, All. Cunn.</i>		962	LEPTADENIA Jacquemontiana, <i>Dene.</i>	ix.	838
<i>iucoides, Sw.</i>		963	LEPTONEMA Lindeni, <i>Hook.</i>	vii.	692
<i>undulatui&i Sw.</i>		964	LEPTOPTEBIS superba, <i>Hook.</i>	x.	910
HYMENOXIS (?) linearifolia, <i>Hook.</i>	ii.	146	LEPTOSPEBUM recurvtum, « <i>Hook. fil.</i>	ix.	893
HYPNUM alopecuroides, <i>Hook.</i>	i.	24 / 5	<i>rupestre, J. D. &</i>	iv.	308
<i>ambiguum, Harv.</i>		4	LEPTOTHECA speciosa, <i>Hook, et Wils.</i>	viii.	748
<i>CGu'd&tiim, Harv.</i>		7	LESKEA curvirostra, <i>Harv*</i>	i.	20 / 4
<i>curvulum, Hook.</i>		7	<i>fulva, Harv.</i>		23 / 2
<i>cyperoides, Hook.</i>		6	<i>polyantha, Hedw. var.</i>		/ 3
<i>humile, Harv.</i>		^	<i>pterogonoides, Harv.</i>		24 / 8
<i>inflexum, Harv.</i>		6	<i>secunda, Hook.</i>		23 / 1
<i>Kamounense, Hart.</i>		10	LEUCODON bartramioides, <i>Hook.</i>		nm
<i>xnicrocarpum, Hook.</i>		4	LEUCOLAENA peltigera, <i>Hook.</i>		45
<i>nervosum, Hook.</i>		3	LEUCOMPHALUS capparideus, <i>Benth.</i>	viii.	784
<i>papillatum, Harv.</i>		8	LEUCOPOGON lancifolius, <i>Hook.</i> <i>fil.</i>	ix.	898
<i>punctulatum, Harv.</i>		10	<i>suaveolens, Hook.fil.</i>		898
<i>propinquum, Harv.</i>		9	LINDENBEBGIA urticifolia, <i>Lehm.</i>		
<i>retroflexum, Kook.</i>		6	LINDENIA acutiflora, <i>Benth.</i>	v.	875
<i>Tavoyense, Hook.</i>		1	<i>rivalis, Benth.</i>	m	475
<i>vagans, Harv.</i>		2	LIXDEHNIA Capensis, <i>Th.</i>	ii.	476
HYPOCREA grossa, <i>Berk.</i>	ix.	872 B	LINJ>8JEA caudata, <i>Hook.</i>	x.	151
HrhoDEBBis Brownii, <i>J. Sm.</i>	vii.	675-6	<i>elegans, Hook.</i>	i.	958
HYPTIS Salzmanni, <i>Benth.</i>	v.	463	<i>macrophylla, Kaulf.</i>	ii.	98
<i>verticillata, Jacq.</i>		458	<i>media, Br.</i>	x.	193
ILEX affinis, <i>Gardn.</i>		465	LiSiAtfTHUS daturoides, <i>Griseb.</i>	ii.	195
<i>crassifolia, Hook.</i>		149	<i>frigidus, Sw.</i>		195
<i>cuneifolia, Hook.</i>		294	LOASA rupestris, <i>Gardn.</i>	vii.	663
<i>microphylla, Hook.</i>		139			
INDIGOFEBIA aspalathoides, <i>Vahl.</i>		188			
IFOMASA Goyazensis, <i>Gardn.</i> ^	v.	479			

1 Is *Monoploca linifolia*, Bunge, in PI. Preiss.2 Is *Adiantum dolotum*, Kze., & Hook. Gen. ct. Sp. Fil.

	VOL.	TTB.		VOL.	TAD.
LOBELIA physaloides, A. Cnnn.	vi.	555-6	MABSIPPOSPEBMUM grandiflorum, ¹ Hook.	. . . vi.	533
trigona, Rox.	iy.	358	MABTENSIA elegans, Herring.	•ii.	697
LOEFFLINGIA Texana, Hook.	iii.	275	MASTOSTIGMA varians, Stocks.	ix.	863
LLOYDIA longiscapa, Hook.	ix.	834	MAXILLABIA angusiifolia, Hook.	. . . iv.	348
LOGAVIA campanulata, Br.	. . .	832	curtipes, Hook.	. . .	384
LO^LABIA Colensoi, Hook. fil.	vii.	627-8	nana, Hook.	. . .	315
Eraseri, A. Cunn.	ii.	185	MAZUS pumilio, Br.	. . . vi.	557
lanceolata, Spr.	v.	*29	MECOCKELLA Origana, Nutt.	: iv.	360
nigra, Col.	x.	960	MECONOPSIS heterophylla, Benth.	. . . viii.	732
procera, Spr. ft	v.	427-8	MEIONECTES Brownii, J. D. Hook.	. . . iv.	306
vulcanica, Bl.	x.	979	MELASMA (?) Zeyheri, ² Hook.	iii.	255
LONICEBA diversifolia, Wall.	ix.	807	MELICOPE simplex, A. Cunn.	vi.	585
Loureiri, Bl.	. . .	806	ternata, Forst.	. . . Tii.	603
LOBANTHUS albiflorus, Hook.	vii.	683	MELLHANIA Leprieurii, Webb.	viii.	753
Colensoi, Hook. fil.	. . .	633	& 755		
lageniflorus, Wight.	iii.	229-30	MENISCITTM salicifolium, Wall.	x.	990
longifolius, Hook.	ix.	880	MENODOEA Africana, Hook.	. vi.	586
macranthus, Hook.	. . .	743-4	MERIMIA Texana, Hook.	. . . iii.	* 278
Schiedeanup, Schlecht.	iv.	374	MEBOSTACHYS (?) capitata, Hook.	. . .	273-4
LOTUS Brunneri, Webb.	. . .	754	METBOSIDEBOS diffPusa, ³ Sm.	vi.	569
purpleus, Webb.	. . .	757	MICBODESMIS puberula, Hook. fil. . . viii.	. . .	758
LUPINTTS arenarius, Gardn.	vi.	511	MICBOSPEBMA lobatum, Hook.	iii.	234
parvifolius, Gardn.	. . .	521	MILLETIA macrophylla, Bent ft % viii.	788-9	
LUXEMBEBG-IA ciliosa, Gardn.	. . .	516	MILLIGANIA cordifolift, ⁴ J. D. H&ok.	. iii.	299
LYCIUM Quitense, ¹ Hook.	viii.	723	MIMOSA floribunda, H. B. K.	iv.	373
LYCOPODIUM biforme, Hook.	iii.	228	rubicaulis, Lam.	. ii.	156
casparinoides, Spring,	x.	968	MITREMTCES viridis, Berk.	. ix.	869 B
compactum, Hook.	iii.	244	MITBEOLA oldenlandioides, Wall.	. 9 . o .	827
Jussieui, ² Desv.	ii.	186	paniculata. Wall.	. . .	828
lepidophyllum, Hook, et Gr..	. . .	162-3	MNIABUM fasciculatum, Hook.	iii.	293
Mathewsi, Hook.	i.	26	Myrsinum ramosum, Hook.	. . i.	20 / 2
pendulinum, Hook.	. . .	90	"rhynchosarpum", Hook.	. . .	/. 3
Pichinchense, Hook.	. . .	85	MONOLOPIA" nijnor, De Cand.	iv.	844
rufescens, Hook.	. . .	36	minor, De Cand.	. . .	343
scariosum, ³ Hook.	. . .	89	MOUBIBIA arborea, Gardn.	. vi.	515
scariosum, Forst. var.			MOZTNNA spathulata, Ort. j3	. iv.	357
decurrens, J. D. Hook.	x.	966	MYEOPHYLLTJM variifolium, J. D. Hook.	. iii. e £89	
LYSIMACHIA repens, D'Urv.	vi.	536	MYBSLNE MyrtilluY, Hook.	. ix.	825
MACABTHUBIA australis, Hugel.	v.	408	myrtoides, Hook.	. . .	877
MACLEANIA floribunda, Hook.	ii.	109	MYBTUS bullata, Sol.	. . . vi.	557
MACBOMITRION brachiatum, Hook, et WiU.	viii.	746	pedunculata, Hook. fil.	vii.	629
MAOBOSTIGMA australe,* Hook.	v.	412	NAPOLEONA Yogelii, Hook, et Planch.	. . . viii.	799
MADABAGLOSSA heterotricha, De Cand.	. . .	326	—800		
MANIHOT Grabami, Hook.	vi.	530	NATALIA paullinioides, Planch.	viii.	780
MABCHANTIA trichocephala, Hook.	. . .	158			
MABSILEA macropus, Hook.	x.	909			
MABSYPIANTHUS hypoides, Mart.	. . .	457			

¹ Is *Pceciolochroma Quitense*, Miers.² For *L. Justieuf* Desv. read *L. scariosum*, Forst.³ *L. scariosum*, Hook, not of Forster. It is *L. vestitum*, Desv.⁴ Is *Stylobarium lineare*, "Sees, in Plant. Preitss.1 Is *Rostkovia grandiflora*, Hook. fil. >2 Is *Graderia scabra*, Benth.3 Is *M. albiflora*, Sol.* Is a species of *Gunner a*; *G. cordifolia*, Hook. fil.

	VOL.	TAB.		VOL.	TAB.
NAUCLEA platanocarpa, <i>Hook.</i>			PALEFOXIA Texana, <i>Hook.</i>	ii.	148
<i>fil.</i>		787	PAPPEA Capensis, <i>JEckl. et Zeyh.</i>	iv.	352
NEOKEBA blanda, <i>Harv.</i>	i.	221/1. 1	PABCOTYCHIA illecebrioides, <i>Wohlb.</i>	Till.	756
<i>cordata, Hook.</i>		1/2	PASSIPLOBA Lescbenaultii, <i>De Cand.</i>	iv.	353
<i>crenulata, Harv.</i>		211/6	PEREOMIA G-aleottiana, <i>Hook.</i>	328	
<i>fimbriata, Harv.</i>		1/4	<i>lancifolia, Hook.</i>	i.	332
<i>flexuosa, Harv.</i>		1/3	PEBNETTIA pumila, <i>Hook.</i>	91	
<i>lancifolia, Harv.</i>		1/5	PEBSOONIA laureola, <i>Lindl.</i>	v.	9
<i>squarrosa, 2Zbo#.</i>		221/3	<i>quinquenervis, Hook.</i>	426	
<i>snbseriata, Hook.</i>		211/7	PETBOSELINTTM prostratum, <i>De Cand.</i>	425	
NEPENTHES villosa, 2foofr. <i>JZ</i>	is.	888	PHAOA densifolia, <i>Hook.</i>	iy.	305
NEUBACANTHTJS spbserostachys,			<i>Vogelii, Webb.</i>	iii.	282
<i>Dais.</i>		835	PHAGNALON melanoleucum, <i>Webb.</i>	viii.	763
NEUBADA procumbens, <i>L.</i>		840	PHABNACETTM semiquinquefidum, <i>Hook.</i>	i.	83
NETTBOCALYX Zeylanicus, <i>Hook.</i>	ii.	174	PHASCUM distatum, <i>Hook. et r Wils.</i>	viii.	737
NIEBUHBIA linearis, <i>W. et Am.</i>		124	<i>eoüguum, Hook. et Wils.</i>		737
<i>oblongifolia, W. et Am.</i>		127	PHEBALITJM grandiflorum, <i>Hook.</i>		
NIPHOBOLTJS Penangianus, <i>Hook.</i>		203	<i>montanum, Hook.</i>	i.	724
NIS>LIA fruticosa, <i>Jacq.</i>	vi.	599	<i>nudum, Hook.</i>	yi.	59
NOMAPHILA pinnatifida, <i>Dalz.</i>	is.	843	<i>retusum, Hook.</i>	i.	568
NOTHOCHILSNA distans, <i>Br.</i>	x.	980	<i>squamuligerura, Hook.</i>	viii.	57
OCHNA pulchrum, <i>Hook.</i>	vi.	588	PHILAGONIA fraxinifolia, <i>Hook.</i>		727
OCHTOCOSHUS Africanus, <i>Hook.</i>			PHOLISMA arenarium, <i>Nutt.</i>	vii.	710
<i>fil.</i>		773	PHYLLOCLADITS hypophylla, <i>Hook. fil.</i>	ix.	626
OOYMTTM bracteosum, <i>Benth.</i>	v.	455	<i>trichomanoides, Don.</i>	vi.	889
OENOTHEBA, alyssoides, <i>H. et Am.</i>			PHYIXOGLOSSUM Drummondii, <i>Kse.</i>	x.	549-51
<i>graciliflora H. et Am.</i>		339	PHYSOBHYNCHUS Brahmicus, <i>Hook.</i>		908
OESIA Conradi, <i>Tuck.</i>	vi.	338	<i>Hook.</i>	ix.	821-2
OLEABIA grandiflora, <i>Hook.</i>	ix.	862	PHYSOSYPHON Loddigesii, <i>Hook.</i>	vi.	508
<i>pannosa, Hook.</i>		862	PHYSTTBUS vaginatus, <i>Hook.</i>	v.	449
OLIGOGYKE Burchellii, <i>Hook.</i>	ii.	101	PIMELEA pbyodes, <i>Hook.</i>	ix.	865
ONYCHITTM mejolepis, <i>Desv.</i>	x.	902	PITTOSPORU3 rhombifoliuru, <i>A. Cunn.</i>	vii.	
OPHIOGLOSSUM Bergianum, <i>Cham.</i>		203	PLATYSTIGHA lineare, <i>Benth.</i>	i.	621
<i>intermedium, Hook.</i>		995	PLECTBANTHITS scrophula-		38
<i>palmaturff, Plum.</i>		4	<i>rioides, Wall.</i>	v.	
OFLOTHEGA Floridana, <i>Nutt.</i>	iii.	276	<i>ternifolue, Don.</i>	464	
OBTHODONTIUM Bulcatum, <i>Hook et WUs.</i>		viii. 739 B	PLEUBOTHALLIS cauliflora, <i>Hook.</i>	i.	460
OBTHOSYPHON rubicundus, <i>Benth.</i>		459	<i>tnlncata, Lindl.</i>		
OrMnDND Vacbellii, <i>Hook.</i>	i.	15	PODOCABPUS biformis, ¹ <i>Hook.</i>	vi.	765
OTJBISIA macropbylla, -JlooJc.	vi.	545-6	<i>Dieffenbachii,² Hook.</i>	544	
OXALIS Cataract®, <i>A. Cunn.</i>	v.	418	<i>ferruginea, Don.</i>	547	
<i>Cratensis, Hook.</i>	iv.	361	<i>nivalis, Hook.</i>	542	
<i>enneaphylla, Cav.</i>	v.	449	<i>Purdieana, Hook.</i>	582	
<i>lotoides, H. B. K.</i>	vii.	661		vii.	624
OXYANTHUS formosui, <i>Hook.</i>					
<i>fil.</i>		785-6			
OXYLOBIUH Batillum, <i>Hook.</i>	vii.	612			
OXYBIA elatior, <i>Br.</i>	v.	483			
P/OHYPHYLLFM: parvifolium, <i>Idndl.</i>	ii.	117			
PACHYSTIGMA pteleoides, <i>Hook.</i>	vii.	698-9			

1 Is *Froelickia Florida***, Moquin.1 Is *Dacrydium Colentoi*, (Tab. 548), Hook. fil.2 Is a sterile plant of *Veronica tetragona*. See Tab. 580.

	VOL.	TAB.	VUL.	TAD.
PODOCARPUS spicata, Br.	. . .	>i.	513	
PoEFALANTiūs compactus,				
<i>Gardn.</i>	. . .	527	PfcUNUS eamydoides, <i>Ch.</i> et	
distichopliyllus, <i>Mart.</i>	. . .	510	<i>Schlecht.</i>	iv. 371
flaccidus, <i>Kth.</i>	. . .	526	PSOBOSPEB&LUM tenuiflorum,	
luricifolius, <i>Gardn.</i>	. . .	524	<i>Hook. fil.</i>	viii. 771
rupcBris, <i>Gardn.</i>	. . .	525	PTEBIS Endliciieriana, <i>Ag.</i>	x. 973
speciosus, <i>Gardn.</i>	. . .	512	flexnosa, <i>Hook.</i>	ii. 119
uncinatus, <i>Gardn.</i>	. . .	523	geraniifolia, <i>Raddi.</i>	x. 915
POHLIA flexuoaa, <i>Hook.</i>	. . .	i. 19 / 5	rotundifolia, <i>Font.</i>	v. 422
POLYGONATUM graminifblium,			eeticaulis, <i>Hook.</i>	iii. 207
<i>Hook.</i>	. . .	ix. 833	PTEEOGONIUM microcarpum,	
POLPODIUM attenuatum, <i>Br.</i>	. . .	v. 409	<i>Harv.</i>	L 24 / 12
biauriculatum, <i>Hook.</i>	ii. . .	121	PTEBOSPEBMTJM reticuktum,	
chrysolepis, <i>Hook.</i>	. . .	721	<i>Wight.</i>	ii. 125
contiguum, <i>Wall*</i>	. . .	x. 987	PTJNEEBIA coagulans, ¹ <i>Stocks</i> ,	ix. 801
discolor, <i>Hook.</i>	. . .	386	QuEttCUS agrifolia, <i>Née.</i>	. . . iv. 377
eriophorum, <i>Hook.</i>	. . .	991	corrugata, <i>Hook.</i>	v. 403-4
^arinosum, <i>Hook.</i>	. . .	947	denaiflora, <i>H. et Am.</i>	iv. 380
Griffithianum, <i>Hook.</i>	. . .	951	Douglasii, <i>H. et Arn.</i>	m. 382-3
Hartwegianum* <i>Hook.</i>	iv. . .	390	lusitanica, <i>Lam.</i>	vü 562
Khasyanum, <i>Hook.</i>	x. . .	949	Mackiana, <i>Hook.</i>	iii. 224
lacknopus, <i>Wall.</i>	JIB	952	Skinneri, <i>Benth.</i>	v. 402
leucosorum, <i>Boj.</i>	. . .	942	QUINTINIA Berrata, <i>A. Cunn.</i>	vi. 558
(status senilis)	■■■	943	KANDIA speciosa, <i>Hook.</i>	. . . ix. 324
Inacrocarpum, <i>JPr.</i>	. ■■	934	RANUNCULUS binternatus, <i>Sm.</i>	v. 497
inurorum, <i>Hook.</i>	. i. . .	70	macropua, <i>Hook. fil.</i>	vii. 634
myriocarpum, <i>Hook.</i>	. . .	84	uiviculus, <i>Hook.</i>	vi. 571-2
onustum, <i>Hook.</i>	. . .	viii. 749	Peruvianus, <i>Pers.</i>	. . . viii. J 745
papillosum, <i>BL</i>	. . .	946	pilulifer,* <i>Hook.</i>	vi. \$00
pellucidum, <i>Kaulf.</i>	■■■	944	pimpinellifoliB, <i>Hook.</i>	iii. 260
var. pinnatifidum,	■■■	945	Bagittifolius, <i>Hook.</i>	•ii. 173
piligerum, <i>Hook.</i>	. . .	321	Bteuopctalus, <i>Hook.</i>	vii. 677
pilipes, <i>Hook.</i>	. . .	221	RAUWOLFIA heterophyle^ <i>Willd.</i>	iv. 366
punctulatum, <i>Hook.</i>	. . .	720	RHABDIA viminea, <i>Wall.</i>	. . . ix. 823
rhynchophyllum,			RHABDOTHECA picridiooides,	
<i>Hook.</i>	. . .	954	<i>Webb.</i>	. . . viii. 766
rostratum, <i>Hook.</i>	. . .	953	RHODODENDRON acuminatumi	
Bemiadnatum, <i>Hook.</i>	. . .	948	<i>Hook. fil.</i> . . * . qx.	886
setigerum, <i>Bl.</i>	. . .	941	.buxifolium, <i>Low.</i>	. . . 980
Bubcronatum, <i>Hook.</i>	. . .	719	ericoides, <i>Low.</i>	. . . 887
subptiolatum, <i>Hook.</i>	. . .	391-2	Lowii, <i>Hook. fil.</i>	. . . 883
Surucuchense, <i>Hook.</i>	i. . .	69	^agosum, <i>Low.</i>	. . . 885
Tweedieanum, <i>Hook.</i>	. . .	86	verticittatutn, <i>Low.</i>	. . . 884
POLYTICHUM dendroides,			RHODOPLEXIA Priccii, <i>Harv.</i>	vii. 613
<i>Hedw.</i>	25	RHUS Thunbergii, <i>Hook.</i>	. . . vi. 595
patulum, <i>Harv.</i>	. . .	18 / 1	RHYNOCHOSIA rufescens, <i>De</i>	
Bemiamclatum, <i>Hook.</i>	. . .		<i>Cand.</i>	. . . ii. 189
<i>fil.</i>	. . .	ii. 194 A	RICCIA velutina, <i>Wilson.</i>	. . . iii. E4i?
POPULUS trichocfirpa, <i>Torr.</i> et			RICHEA Bcoparia, <i>HJok. fiL</i>	. . . ix. 850
<i>Gray.</i>	. . .	878	RITCHIEA erecta, <i>Hook. fil.</i>	. . . viii. 769
POBTEBIA bractescens, <i>Hook.</i>	. . .	864		- 770
POTHOS remotiflorus, <i>Hook.</i>	. . .	133	RITINA inaequalis, <i>Hook.</i>	. . . ii. 13b
Bcandens, <i>L.</i>	. . .	175	RUBUS geoides, <i>Sm.</i>	. . . v. 495
POZOPSIS cordifolia, <i>Hook.</i>	. . .	859	Gunnianus, <i>Hook.</i>	. . . iii. 291
PBEPUBA connata, <i>Gardn.</i>	. . .	225-6	Lobbiinus, <i>Hook.</i>	. . . viii. 741-2
PBIONOTES Americaua, ¹ <i>Hook.</i>	i. . .	30	pulclierrimus, <i>Hook.</i>	. . . 729-30
PBOGKIA completa, <i>Hook.</i>	. . .	94	rosaoflorus, <i>Hook.</i>	. . . i. 46
			rosacfolius, <i>Sm.</i>	. . . iv. 340

¹ This constitutes the genus *Allodape* of Bndlicher,

1 Dunal refers this to the Genoa *Withania*.
9 fs *R. leptocauli**

	VOL.	TAB.		VOL.	TAB.
SAMTDEEA Indies, <i>Gatrtn.</i>	i.	7	STENOPIETALUM draboides, ¹ <i>Hook.</i>	vii.	617
SANTALUM Mida, <i>Hook.</i>	vi.	575	lineare, <i>JBr.</i>	•	618
var. <i>f3 Hook.</i>	•	563	procumbens, <i>Hook.</i>	•	610
SAPOTA marginata, <i>Dccsne.</i>	viii.	761	robustum, <i>JEndl.</i>	vii.	620
SABCOSTEMMA brachyetigina, <i>Wight.</i>	ix.	861	STEECTTLIA colorata, <i>Eoxb.</i>	ii.	143
<i>Daltoni, Dccsne.</i>	viii.	768	STBEPTANTHUS flavesrens, <i>Hook.</i>	i.	44
SAURAUJA barbigera, <i>Hook.</i>	iv.	331	• glandulosus, <i>Hook.</i>	•	43
pedunculate, <i>Hook.</i>	•	341-2	STBOBILOMYCES montosus, <i>Berk.</i>	*i.	872 A
SAUVAGESIA deflxifolia, <i>Gardn.</i>	v.	484	STBYCHNOS toxifera, <i>Schomb.</i>	iv.	364-5
SOAPHTGLOTTIS (?) fasciculata, <i>Hook.</i>	iv.	317	SYBBHOPODON ropsens, <i>Harv.</i>	i.	22 / . 4
<i>rosea, Hook.</i>	•	313	T^NITIS obtusa, <i>Hook.</i>	x.	994
SCHIEDEA Nuttallii, <i>Hook.</i>	vii.	649-50	TAPhR ^A villosa, <i>Gardn.</i>	v.	4G ^{>}
SCHISTLDIUM arcuatum, <i>Hook.</i> <i>et Wits.</i>	Tiii.	738	TAPUBA ciliata, <i>Gardn.</i>	•	4G6
<i>pulchellum, Hook, et</i> <i>Wits.</i>	•	738	TAXODITJM sempervirens, ² <i>Lamb. (?)</i>	iv.	379
SOHMIDELIA monophylla, <i>Hook.</i> <i>fil.</i>	•	776	TELOICA fragrantissima, <i>Hook.</i>	xii.	208-12
SOLEBODONTIUC secundum, <i>Harv.</i>	i.	21 / . 1	TEPHBOSIA suberosa, <i>Be Cand. u.</i>	•	120
<i>etrichtum, Harv.</i>	•	f. 2	TETEACAEPJEA Tasmanica, <i>Hook.</i>	•	•
SCOLOPENDBIUM Lindenii, <i>Hook.</i>	v.	488	TETRATHECA ciliata, <i>Lindl.</i>	•	268
SCYPHOOOBONIS viscosa, <i>A. Gray.</i>	ix.	854	TETRONIUM Magellanicum, <i>Willd.</i>	vi.	534
SCYTANTHTS Currori, <i>Hook.</i>	vii.	605-6	THALI ^A RUM Dulzelii, <i>Hook.</i>	ix.	8
<i>Gordoni, Hook.</i>	•	625	THAMNOCABPU ^S Guuuiaus, <i>Hook.</i>	vii.	6 ^A
SEDTJM Wallichianum, <i>Hook.</i>	•	604	THAUMASIA Cunninghami, <i>Hook.</i>	yi.	517-i
SEMOMVILLEA fenestrata, <i>Feizl.</i>	vi.	587	THIBAJJ ^A acuminata, <i>Hook.</i>	ii.	11
SXNECIO littoralis, <i>Gaudich.</i>	v.	493	^augustifolia, <i>Hook.</i>	•	Hi
SEBICOSTOMA pauciiloruin, <i>Stocks.</i>	ix.	804	<i>elliptica, E. et P.</i>	•	108
OERIS polynprpha, <i>Less.</i>	vi.	501	THYSANOCARPUS e'jgans, <i>Msch.</i>	4- if.	i.
SHUTEIU ^A glabrata, <i>W. et Am.</i>	iL	144	<i>pusillr</i> , <i>Hook.</i>	•	39
SIDA parnassisefolia, <i>Hook.</i>	iv.	384	THYSANOMITETON uYicinatum, <i>Harry.</i>	•	42
SINAPIDENDBON gracile, <i>Webb.</i>	viii.	751	THYBANTHA subulata, <i>Hook.</i>	vi.	22
<i>Vogelii, Webb.</i>	•	752	TILLAEA macrantha, <i>J. D. Hook.</i>	iv.	500
SINCLAIEIA discolor, <i>Hook.</i> <i>et Arn.</i>	v.	451-2	moschata, <i>Be Cand.</i>	vi.	310
SIPHOOAMPTLua giganteua, <i>Cav.</i>	viii.	716	<i>vcriticillaris</i> , ⁴ <i>Br.</i>	iii.	535
SISYEINCHIUM inourvatum, <i>Gardn.</i>	vi.	513	TODEA pellucida, ⁵ <i>Carm.</i>	i.	295
SLOANEA Jamaicensis, <i>Hook.</i>	vii.	693-6	TOFIELDIA sessiliflora, <i>Hook.</i>	vii.	8
SMLLACINA flexuosa, <i>Hook.</i>	vi.	529	TOBBEYA taxifolia, <i>Am.</i>	•	691
SONCHUS Baltoni, <i>Webb.</i>	viii.	765	TOVABIA pendula, <i>R.etP.</i>	vii.	232-3
SpHACELABJi. hordaceae, <i>Harv.</i>	viL	614	TBADESCANTIA gracilis, <i>H.B.K.</i>	•	664
SPHAEBA % obertsii, <i>Hook.</i>	i.	11	<i>birsuta, H.B.K.</i>	•	654
SPHYBOSFEBKUM myrtifolium, <i>Hook.</i>	ii.	112	TBICHANTITA major, <i>Hook.</i>	•	665
STATICE Stocksii, <i>Soiss.</i>	ix.	837	<i>minor, Hook.</i>	•	667
STELIS lamellata, <i>JAndl.</i>	i.	62	<i>GGG</i>	•	•
-SXELLABIA decipiens, <i>Hook. fil.</i>	vii.	680	-----	•	•
STENOGLOSSUM Bubulatum, <i>Lindl.</i>	•	51	1 Is MAJtea awtralis, Lehm. Cat. Sem. 1849	•	•
			2 Is a scicle branch of Aides btacUata, <i>Doug.</i>	•	•
			3 This proves to have hypoffynous scales, and therefore Bulliarda moschata, IJ'Ur ^A	•	•
			4 Not of De Candolle. It is UuU*arda re- curva, <i>Hook. fil.</i>	•	•
			5 Now Leptopteris pellucida, Presl.	•	•

VOL.	TAB.	VOL.	TAB.
TBICHINTTJM remotiflorum, <i>Hook.</i> vi.	596	VACCINITOC cereum, <i>Forst.</i> i.	87
TBICSOMANES bicorne, <i>Hook.</i> x.	992	<i>coriaceum, Hook. fil.</i> ix.	892
<i>Colensoi, Hook. fil.</i> 979		<i>Emirnense, Hook.</i> ii.	131
<i>elongatum, A. Cunn.</i> viii.	701	<i>Forbesii, Hook.</i> iv.	345
<i>giganteum, Borg.</i> 702		<i>Imrayi, Hook.</i> iii.	292
<i>melanorhizon, Hook.</i> 705		<i>Javanicum, Hook.</i> viii.	740
<i>montanum, Hook.</i> ii.	187	<i>muscicola, Hook.</i> 717	
<i>Petersii, A. Gray.</i> x.	986	<i>secundiflorum, Hook.</i> ii.	134
<i>Pluma, Hook.</i> 997		VALLABIS <i>Pergularia, Br.</i>	153
<i>polyanthos, Hook.</i> viii.	703	VEBONICA <i>Bidwilli, Hook.</i> ix.	814
<i>Smithii, Hook.</i> 704		<i>diffusa, Hook. fil.</i> vii.	645
<i>Spruceanum, Hook.</i> x.	981	<i>nivea, Hook. fil.</i> 640	
TBICHOBTOMUM subsecundum, <i>Hook. \$ Grev.</i> L	17/.5	<i>tetragona, Hook.</i> vi.	580
TRIDONTIXTM Tasmanicum, <i>J. D. Hook.</i> m.	248	VIGNA <i>hirta, Hook.</i> vii.	637
TBIPOLITJM obtusiflorum, <i>Hook.</i>	281	VILLABSIA <i>involucrata, Hook.</i> viii.	725
<i>macrocylis, Hook.</i>	285	VIOLA <i>balsaminoides, Gardn.</i> iii.	217
TBIGLOCHIU calcaratum, <i>Hook.</i> v.	416	<i>chrysantha, Hook.</i> i.	49
<i>Calcitrapa, Hook.</i> Tiii.	731	<i>maculata, Cav.</i> v.	499
<i>centrocarpum, Hook.</i>	728	VISCIJM <i>falcatum, Ch. et</i>	
<i>filifolium, ² Hook.</i> vi.	579	<i>*</i> <i>Schlecht.</i> , , , iv.	368
TBOPJBOLUM cirrbipes, <i>Hook.</i> t.	411	<i>incanum, Hook.</i> i.	73
<i>tuberosum, M. et P.</i> vii.	653	VIBNIA <i>Mocanera, Juss.</i> iii.	253
TBOPIDOCABPUM gracile, <i>Hook.</i> i.	43	VITEX <i>littoralis, A. Cunn.</i> v.	419-20
<i>Bcabriuscum, Hook.</i>	52	WAHLENBEBGIA <i>albonarginata,</i>	
TURNERA dichotoma, <i>Gardn.</i> vi.	522	<i>Hook.</i> ix.	818
U TIC ARIA Africana, <i>Don.</i> viii.	781	WEINMANNIA <i>biglandulosa,</i>	
USTEBIA Guineensis, <i>WilUL</i>	795	<i>A. Cunn.</i> iv.	301
UTBICULABIA nelumbifolia, <i>Gardn.</i> vi.	505-6	WEIBSIA <i>campylocarpa, Hook.</i>	*
UVABIA Vogelii, <i>Hook. fil.</i> viii.	767	<i>et Am.</i> ii.	136
VACCINIUM buxifolium, <i>Hook.</i> <i>fil.</i> ix.	891	<i>flaccida, Harv.</i> i.	18/.3
		<i>pallens, Hook. et WiU.</i> viii.	739 A
		WILSONIA <i>rotundifoila, Hook.</i> v.	410
		WIMMEBIA <i>concolor, Ch. et</i>	
		<i>Schlecht.</i> iv.	356
		XANTHOSIA <i>ciliata, Hook.</i> viii.	726
		<i>dissecta, J. D. Hook.</i> iv.	303
		XIMENIA <i>parviflora, Benih.</i>	350
		XYLOMELON <i>occidentale, Br.</i> v.	446

1 Is *T. quercifolium*. Hook et Gr.2 Is *T. triandrum*, Mich.

rfcccc.


CHEILANTHES CHRYSOPHYLLA, *Hook.*

Caespitosa, radice fibrosa, stipitibus 2-5 purpureo-ebeneis pilis longis subulatis paleaceis inferne prsecipue hispidis, frondibus pinnatis subcoriaceis dcltoideo-acuminatis 3-4-uncialibus supra glabris subtus furfuraceis aureo-flavis, pinnis lanceolatis paribus infimis deltoideis omnibus pinnatifidis, segmentis ovatis crenulatis inferioribus pinnarum infimarum oblongis subsinuatis, involucris e margine reflexo frondis continuis usque ad apices crenulatis, marginibus membranaceis nudis (non ciliatis).

Cheilanthes chrysophylla, *Hook. Gen. et Spec. Fit. 2. p. 13.*
HAB. Bare rocks, Kala-Panec, Khasya, Drs. *Hooker and Thomson.*

Of this rare species we have only seen specimens of Messrs. Hooker and Thomson from the above locality. It has much general affinity with *Cheilanthes farinosa*, Kaulf (Ehook* et Grev. Ic Fil. t. 134), but, independently of the colour of the powdery substance on the underside of the frond, (white in *Ch. farinosa*), the involucres are very different; in *Ch. farinosa* wholly membranaceous, nearly orbicular, and more or less united at the base, (in that sense often continuous), whereas here the involucres seem to be formed of the reflexed margin of the frond itself, of the same texture, altogether continuous, and though crenulated, never ciliated at the edge. The crenulated involucres may be considered to justify the placing of this species in *Cheilanthes*, rather than in *Jpteris* or *Allosorus*.

*Fig. 1. Upper side of a futile lobe of a lower pinna.
f. 2. Fertile pinna seen from beneath:—magnified.*



DCCCCII.

ONYCHIUM MEI.ANOI.EPIS, *Dene.*

Caudice brevi subrepente squamis rigidis appressis nigris paleaceo, stipitibus cespitosis 3-6 uncialibus gracilibus rachibusque subflexuosis canaliculatis glabris nitide stramineis (in sicco), fronde membranacea 3-6-unciali costata glabra pallide virente ovata triplicato-pinnatisecta, laciniis sterilibus angusto-cuneatis 3-5 fidis incisis, laciniis sublinearibus acutis, fertilibus oblongis cuspidatis, involucris pallidis membranaceis linearibus longitudine fere laciniarum, apicibus sterilibus.

Onychium melanolepis, *Dene*. *PL de VArabie Heur. Archiv. du Mus.* 2 p. 189. *Kze. in Schkuhr, FU. Suppl.* 2. p. 9. t. 104-/. 2. *Hook. Gen. et Sp. FU.* 2. p. 124.

Cheilanthes leptophylla, *Br. in Salt's Abyss. App.* iv. p. lxv.9 (*name only.*)

Allosorus cuspidatus, *Jaub. et Spack. Illustr. PL Or. S.* p. 1. t. 201,

HAB. Abyssinia, *Salt, H. Schimper*, n. 1672. South Persia, rock of Mount Pire-zend, between Shiraz and Kazeroum, *Aucl;er-Eloi* n. 5488. Caves in the Island of Karch, Persian Gulph, and at Dalechy, *Kotschy*, n. 10 and 198.

Mj? first knowledge of this plant wad from the specimens of Auchér-Eloi in.z*ly possession. On finding it to be identical with Mr. Brown's *Cheilanthes leptophylla* of Salt's plants in the Banksian Herbarium, I was anxious to figure it; which done, and not till then, I discovered that it was, under other names, represented in Kunze supplement to Schkuhr's FiHces, as well as in the fine work of Jaubert and Spach on Oriental plants. It is a Fern of very delicate texture, like some slender specimens of *Gymnogramme leptophylla** and it appears to me that Decaisne and Kunze have correctly referred it to *Onychium*; thp specific name is derived from the black scales on the caudex, jvwhich is wanting in the specimen from which our drawing was made.

Fig. 1. Sterile portion of a frond, f. 2. Fertile portion of ditto, seen from beneath'.—magnified.

Purdieanc.

N. O. Schizaeacerc.

DCCCCIII.

ANEMIA (*Coptophyllum*) AURITA, SW.

Caudic repente, frondibus sterilibus longe stipitatis deltoideo-oblongis subcoriaccis glabris lucidis pinnatis inferne bipinnatis, pinnulis subrotundis rhomboidisve petiolulatis flabellato-venosis crenulato-dentatis margine superiori ssepe auriculatis, fertili (radicali) solitaria bipinnata lobis glomeratis, stipitibus predunculoque hirsutis.

Anemia aurita, Swartz, *Syn. Fil* p. 157. *Willd. Sp. PL* 5. p. 95. *Presly SuppU Tent. Pteridogr.* p. 80.

HAB. Jamaica, Swartz; Dolphin feak, Wesfmorland, Purdie, Wilson.

Those species of *Anemia* which have the fertile fronds solitary and distinct from the sterile ones, have been separated by the late Mr. Gardner into a genus which he called? *Coptophyllum*, a separation which we are glad to find is not adopted by Presl in his work above quoted. Mr. J. Smith is disposed to refer it to *Mohria*. The present eqms to be one of the rarest of the group, and, for half a century from its discovery by Swartz, only to have been known to that author. Presl had only seen sterile fronds. Mr. Purdie detected the species in one spot in Jamaica, but could find no fructification. At length Mr. Wilson sent us, with many sterile plants, one fertile one, which we have here ^presented. It does not appear to have been met with any where but in Jamaica.

Fig. 1. Pinnule. /. 2. Cluster of fructifications, f. 3. Capsule:—magnified.

CCCCIV.

CHEILANTHES OCHRACEA, *Hook.*

C&spitosa, radice fibrosa, stipitibus brevibus squamis oblongis obtusis patentibus paleaceis una cum rachi pupureo-ebeneis, frondibus submembranaceis lato-lanceolatis sub-spithameis pilosulis subtus dense ochraceis furfuraceis pinnatis, pinnis plerisque oppositis lanceolatis obtusis infimis subdeltoides omnibus fere ad rachin profunde pinnatifidis, segmentis ovalibus obtusis vix crenulatis ciliatis, involucris continuis angustis e margine reflexo formatis, marginibus membranaceis crenulatis non ciliatis.

Cheilanthes ochracea, *Hook. Gen. et Spec. Fil. 2. p. 144.*

Allosorus ochraceus. *Hook, in Benth. PL Hartweg. p. 55.*

HAB. Moist shady places, at Morelia, Mexico, *Hartweg*, n. 418.

In this plant the involucre is so narrow, as may almost be overlooked, when the genus would be *Nothochlcena*; and when more evident, it is so continuous as to justify its being referred to *Allosorus*. I have placed it in *Cheilanthes* for the same reasons as har[^] actuated me in placing *Ch. chrysophylla* (Tab. 901) there, and on account of its affinity in other respects with that species. Here this pulverulent substance is very dense, of a singularly ochraceous colour, and the fronds are of a thinner texture than the last mentioned species; it differs further in *ihp* villous upper side of the fronds, in the ciliated margin[^] in the very short stipites, and is remarkable for the large spreading blunt chaffy scales which clothe the latter.

*Fig. 1. Lower segment of q[^]pinnule, seen from above.
/. 2. Fertile pinna, seen from beneath:—magnified.*

DCCCCV.

GYMNOPTERIS QUERCIFOLIA, *Bernh.*

Frondibus ternatis, pinnis sterilibus membranaceis subciliatis lateralibus subcordato-lobatis sessilibus in & quilateralibus intermedia maxima petiolata oblongo-ovata sinuata lobata, fertili longe stipitata, stipite graciflmo, pinnis linearibus, stipitibus inferne praecipue squamosis, caudice repente.

Gymnopteris quercifolia.—*Bernh.-Fresl. Tent. Pterid. p. 244.*
Acrostichum quercifolium, *Retz. Obs. Bot. 6. p. 39. Sw. Syn.*
Fil. p. 12. Schkh. Fil 2. t. 3.

Osmunda trifida, *Jacq. Coll. 3. p. 281. t. 20. /. 3.*

Leptochilus quercifolius, *Féé, Mem. sur la Fam. des Fougères.*
p. 88.

HAB. Ceylon, (Vahl.) Mrs. Genl Walker, *Gardner, n, 1170.
Madras Peninsula, Dr. Wight, *Herb. Propr. Cryptog.* n. 46.
China and Cochinchina, Moreau, Turanne, and Gaudichot,
(according to Presl).

A very peculiar looking and very distinct plant, represented long ago by Jacquin and by Schkuhr; but we cannot pay those authors the compliment to say they have figured it well or accurately. As to its genus, the most conservative Botanist will scarcely consider it can, with propriety, remain in *Acrostichum*. Bernhardi and Presl placed it in *Gymnopteris*, but M. Fée refers this, with several other genera, to *Leptochilus* of Kaulfuss, distinguished according to M. Fée, "sporangiis universalibus, Jn duobus sulcis angustis nascentibus," characters not very intelligible to us. Our present plant is remarkable for the length and slenderness of the fertile frond, and small size of the latter in comparison with the sterile one.

Fig. 1. Portion of a sterile frond. /. 2. Portion of a fertile frond:—magnified.

DCCCCVI.

ANEMIA (*Anemidictyon*) TWEEDIEANA, *Hook.*

Humilis, frondibus glaberrimis pinnatis, pinnis 3-7 subrotundo-ovalibus obtusis dentatis reticulatim venosis tenuicostatis, terminali inasquilateralis, spicis compositis bipinnatis una cum pedunculis fronde brevioribus, stipitibus basi squamosis rachique decidue fusco-Villosis.

HAB. Tucuman and Uruguay, S. Brazil, *Tweedie.*

This is one of the few Anemias distinguished by the reticulated (not forked) veins, of which Mr. John Smith has constituted the genus *Anemidictyon*, and in which he has been followed by Presl. We feel it safer, in the present state of the science, at least, to consider it rather a section of, than a distinct genus from, *Anemia*; for there is no natural character to distinguish it, nor any other mark beyond the mere anastomosing of the veins. As in *Anemia* proper, the species of this section are exceedingly variable. Presl enumerates 12, which he divides into groups, the one having the terminal lobe of the frond coadunate with the nearest lateral ones, and then lobed or cut, and the other^v into those which have the terminal lobe free. Among the former he places the *A. Phyllitidis* of J. Smith in Hook. &en. Fil. which he considers identical with *A. fraxinifolia* of Raddi, Fil. Bras, t 8. bis. Smith's plant, however, is represented with a free terminal lobe, and it becomes a question whether it be distinct from the *A. Phyllitidis* of Swartz, noi¹ can we point out any character of our present species, other than the smaller size, the few very obtuse pinnae to each frond, together with their freedom from all pubescence, combined with the short peduncle and spikes.

Fig. 1. Pinnule. /. 2. Lobes of the spikes. /. 3. Capsule: magnified.

DCCCCVIL

GYMNOPTERIS TRILOBATA, 8m.

Frondibus sterilibus hastato-trilobis seu pinnatifidis basi in petio}um alatum decurrentibus, lobis oblongis obtuse acuminatis intermedio nunc sinuato subpinnatifidis, fertilibus profunde tri-quinquefidis segmentis linearibus acuminatis infra lobos non raro sterilibus, stipitibus paleis longis subulatis patentibus fuscis squamosis, radice csespitsa.

Gymnopteris trilobata, J. Smith, in Hook. *Journ. of Bot.* 3.
/* 403, (*name only*).

Leptochilus subquinquefidus, Féé, *Acrost* p. 88. t. 49.

Gymnopteris subquinquefida. Presl. *JEpm. Bot.* p. 151.

HAB. Luzon, Cuming, (n. 3), Mr. Thus. Lobb, in *Herb. Nostr.*

A plant evidently of the same genus with that figured at our Tab. 905; and for similar reasons, as in that case, M. F6e refers it to *Leptochilus*. Our plants have a short stout horizontal very scaly caudex, and the fronds, including the stipes, are a foot to a foot and a half long. Strangely enough M. F6e quotes J. Smith's *Gymnopteris trilobata* under his *fierenoneuron diversifolium* (*Acrostichum BL* and *Cyrtogonium* of J. Sm.) an extremely different species, and No. 32 of Mr. Cuming's collection. Our species varies with the leaves simply oblong and subhastate to pinnatifid with nine and ten segments, and with the lower segments auriculate.* *Gymnopteris tacccefolia*, J. Sm. (Cuming, n. 357. *Leptochilus*, Féé, *Acrost.* t. 50), is closely allied to the more highly developed, form of this, but it is larger, more membranaceus, decidedly pinnate, with the lower pinna? bipartite, and has a creeping caudex. Smith's name is anterior to M. F6e's, but neither is appropriate to so very variable a species.

Fig. 1. Portion of the sterile frond. */. 2.* Portion of the fertile frond:—*magnified*.

DCCCCVIIIL

PHYLLOGLOSSUM DRUMMONDI, Kze.

Gen Char. Spica pedunculata, brevis, oblongo-ovata, capsulifera, bracteata. JBracteae plurimas, spiraliter dispositae, subimbricatæ, cordato-rotundatae, integerrimæ, crassius culse, subcarnosæ, mucronatim acuminatas, basi producta libera. Capsule solitariæ axillares (in bractearum axillis) reniformes, erectæ, bivalves, verticaliter usque ad basin de-hiacentes. Sporca numerosæ subtetrahedrae, pellucidæ.—Herbaceum pusillum, australasicum. Caudex brevissimus, subnulus. Radix ejfibris 1-3 simplicibus, crassiusculis, albis tuberosa; tuberibus oblongis basi attenuatis. Folia radicalia, vel subradicalia, Unearia, semiteretia, acuta, scapo sen pedunculo breviora. Scapus teres, crassus, solitarius, spiciferus.

*Phylloglossum Drummondi, Kze. in JBotanische Zeitung, 1843.
p. "(24c. cum Ic. xylogr. (1843.)*

*Lycopodium Sanguisorba, Spring Monogr des Lycop. JP. II.
p. 36 (1849.)*

HAB. Australia, Swan River, *Drummond*, n. 993. George Town, Van Diemen's Land, *R. Gunn*, Esq. n. 1560. New Zealand, peaty soil near the Wytangi River, *Dr. Hooker*^ *Rev. W. Colenso* (n. 325.), *Dr. Sinclair*.

In 1843 a woodcut and description of this very remarkable plant were published by Professor Xunze, in Mold's Botanische Zeitung for 1843, as a new genus and even a new natural group of the Linnsean Filices. It is a matter of surprise then^ that these should appear to be unknown to Dr. Spring, who presented his "Monographie des Lycopodiacees" to the Académie Royale de Belgique in 1849, and there described it as a new *Lycopodium*, "*L. Sanguisorba*," remarking, however, that "d'après son facies on croirait avoir affaire à une espèce du genre *Isoetes*." Professor Kunze's view we consider more correct; "Planta memorabilis, habitu fere *Plantaginem' pusillam* referens, seu inter Filices *Ophioglosso Sergiano*, Schlect. (Hooker, Ic. PI. p. 263.) non assimilis." It is of the size, and with the general habit, of that plant, but the spike is that of *Lycopodium*. It may well form a distinct order between *Ophioglossaceæ* and *Lycopodiaceæ* :— and we have here had little more to do than improve Kunze's accurate characters, by means of our more copious specimens.

Fig. 1. Entire plant, f. 2. Spike of fructification, f. 3, Inside view of a bractea with a capsule, f. 4. Outside view of ditto, f. 5. Spores:—all more or less magnified.

DCCCCIX.

MARSILEA MACROPUS.

Foliis peltatis quaternis petiolisque elongatis sericeo-tomentosis, foliola lato-cuneatis apice erosionis, pedunculis subradicalibus elongatis biuncialibus, capsulis oblique ovatis dense sericeo-strigosis transversim lineatis hinc basi gibbosis, caudice repente ramoso.

HAB. Australia, low inundated grounds; Lachlan river and Liverpool plains, All. Cunningham. Severn river, S. W. Australia, Drummond.

Our finest specimens are sent from Swan river, among the later collections of Mr. Drummond. It is a species of *Marsilea*, as far as I can find, hitherto quite undescribed ; remarkable for the very sericeously tomentose leaves, (especially the underside), and petiole and capsules, and for the great length of the peduncles of the latter.

The caudex creeps for some length, and is scarcely so thick as a crow's-quill, rooting, branched, and knotty; the knots are densely woolly with ferruginous hair, and seem to be, the rudiments of a new cluster of fronds. *Fronds* or *leaves* from the apex of a woolly knot or branch, two to four from one point. *Petioles* from four inches to a span long, erect, flexuous slender, silky, bearing at the point four spreading broadly cuneate leaflets, finely and radiately veined, the veins here and there anastomosing, villous with dense silky hairs, especially beneath: the hairs often deciduous above, and occasionally beneath, subulate, articulated, tawny. From the very base, among the cluster of petioles, arise one or two erect peduncles about two inches long, in other respects resembling the petioles; these are terminated each by an obliquely erect, ovate, compressed *capsule*, transversely striated, with a gibbosity on one side at the base, densely clothed with imbricating, subulate, jointed hairs.

Fig. 1. Leaflet, *f. 2.* Capsule, *f. 3.* The same cut through transversely, *f. 4.* Hairs from the Capsule :—all more or less magnified.

DCCCCX.

LEPTOPTERIS SUPEKBA.

Fronde brevi-petiolata lanceolata inferne longe attenuata bipinnata, pinnis primariis oblongo-linearibus acuminatis patentibus inferiopribus deflexis, pinnulis pinnatifidis laciniis lanceolatis obtusiusculis costatis integris vel furcatis, petiolo rachique primario inferne proecipue robusto rachibusque partialibus omnibus fusco-setoso-tomentosis.

Todea superba, Colenso, *Ferns of N. Zeal.* p. 28. *Herb. N. Zeal.* n. 2306, 793, 113, 1672, and 281.

Trichomanes, Banks, in *Herb. Jacq. mem. Herb. Imp. Palat. Vien. {Presl.}*

HAB. Northern Island, N. Zealand. *Forster in Herb. Hook.*; Dells, in shaded forests; on the mountain range near "Waikare lake, in decomposed sandstone soil, Deer. 1841; Tongarou mountain, 1838. Rev. W. Colenso. Dr. Sinclair, Upper Hill, Port Nicholson, Dr. Lyall.

I possess a specimen of this plant, gathered by Forster in N. Zealand, during Captain Cook's voyage, and in all probability it is the same plant named "*Trichomanes*" in *Jacq. Herb.*, and not the *L. pellucida* to which Presl refers it, communicated, by Sir Joseph Banks, to the Austrian Botanist: to Forster therefore is due its discovery. It seems to have remained unknown to any other Botanist till the Rev. Mr. Colenso detected it in 1838 and 1841 in the above localities. Its affinities are with the N. Zealand *Todea pellucida*, (see Tab. 8. of *Icones Plantarum*), and the Australian *Todea Fraseri*, Hook, et Grev. Ic. Fil. t. 101. The three we think are well separated from *Todea*, and Presl has determined the character of his Genus *Leptopteris* in his Suppl. Tent. Pteridogr. p. 70. As a species, it is abundantly distinguished by the outline of the frond tapering at the base, the stout stipes and main rachis, and the copious bristly tomentum of the underside of the main and partial rachis. "Some of the fronds," Mr. Colenso says, "are upwards of 'four feet in length. The old fronds spread outwards, while the younger ones, generally three in number, circinate and of a lighter green, rise in the most graceful suberect manner from the centre."

Fig. 1. Fertile pinnule as seen from the underside, *f. 2.* Small portion of the same. */. 3. 4.* Capsules :—*magnified*,

Mooreance.

N. O. Filices.

DCCCCXI.

ASPLENIUM NoViE-CALEDONIiE,

Fronde rachiformi deltoidea glaberrima 3-4-pinnata, rachibus pinnulisque linearibus subcoriaceis uninerviis, soris angusto-linearibus elongatis solitariis (qualibet pinnula) ad marginem dehiscentibus, stipite elongato inferne terete.

HAB. Crevices of rocks in very dry exposed situations, New Caledonia, *C. Moore, Esq.*

A very distinct species of *Asplenium*, which will assuredly rank in habit and structure with the *Darea* group, which is usually acknowledged to be inseparable, generically, from true *Asplenium*.

The present species is remarkable in the length and narrowness of the segments of the frond. Of the ultimate segments nearly the whole of one side of the costa is often occupied by the sorus: so that the species cannot well be confounded with any other of the genus.

Fig. 1. 2. Fertile portions of the frond:—magnified.

DCCCCXIL

CHELLANTHES NITIBULA.

Caudice brevi repente, stipitibus 2-5-uncialibus cespitosis rachibusque (supra puberulis) ebeneis squamis subulatis fuscis paleaceis deciduis hispidis, frondibus 3—4 y. 5-uncialibus subdeltoideo-oblongis acuminatis (sterili latioribus) coriaceo-membranaceis pallide virentibus glabris pinnato-pinnatifidis inferne subbipinnatis, pinnis approximatis ssepissime oppositis ovato-lanceolatis subdimidiatis (dimidio inferiore latiore) profunde pinnatifidis usque ad rachin, infimis iterum subpinnatis, pinnis secundariis pinnatifidis, lobis oblongis integris vel sinuatis sensim acutis, involucris subintramarginalibus continuis raro interruptis latis planis membranaceis fuscis appressis ssepissime lobatis crenatisque transversim rugosis.

Cheilanthes nitidula, *Hook* Gen. et Sp. Fih 2. p. 112.*

Pteris nitidula, *Wall. Cat, n. 89.*

Allosorus nitidulus, *Fresh*

HAB. Northern India. Kamaoun, JDr. *Wallich*. Rocks, Simla,
Dr. T. Thomson. Pundkester, N. India, *Mr. Edgeworth*.

One of Dr. "Wallich's many Indian discoveries, made either by himself in Northern India, or through the medium of the many excellent collectors -he was privileged to employ ; and whom he so successfully employed, especially in the search after Ferns, that very little of novelty has been left for succeeding Botanists. In the arrangement of the Genus *Cheilanthes* in the Genera and Species Filicum, we have placed this in the **<PterijloidecB*" group, having, as we there observed, nearly as strong a claim to be placed in *Pteris* or *Allosorus* as in *Cheilanthes*. Presl considers it an associate of *Cryptogramma*^ Br., for he refers it to the section of his *Allosorus*^ which includes that genus; erroneously we think. As a species it is very distinct.

Fig. 1. Upper side of a sterile pinna, f. 2. Under side of a fertile pinna; magnified.

DCCCCXIL

ASPLENIUM (*Athyriupi?*) GRAMMITOIDES, *Hook.*

Humile, fronde herbacea firma parce setulosa oblongo-lanceolata longe acuminata pinnata superne pinnatifida, pinnis lanceolatis subfalcatis obtusis basi superiore auriculatis omnibus lobato-vel dentato-pinnatifidis, stipite brevi paleaceo, paleis setaceis patentibus crispatis, soris satis magnis, involucris oblongis planiusculis ciliatis, inferioribus saepe diplazoideis.

Diplazium grammitoides, *JPresI*, *JSpimel*. *JBot*. p. 84.

HAB. Luzon, *Owning*, n. 56. Java, *Thos. Lobby* n. 258.

Mr. J. Smith in his *Enumeratio Filicum*, after the names often species of *Diplazium*, observes that "Mr. Cuming's n. 56 is probably one of them in a young state." Our plants, however, from that collector, have every appearance of being perfect, and this view is confirmed by copious specimens, identical with these, having been found in Java, by Mr. Thomas Lobb, and which are here figured. The late Dr. Presl entertained the same opinion, and described our plant in the work above quoted, as a new species, *Diplazium grammitoides*. The specific name I willingly adopt, though I doubt of the propriety of referring it to the Genus *Diplazium*, unless *Diplazium* were to receive every *Asplenium* which had a double involucre ever so sparingly mingled" with the single ones. Here a displazioid involucre is only found towards the base of the pinna, and where a lobe or pinnule may be considered to be combined with the pinna, each half opening towards its own rachis or costa. With more propriety it might be placed in *Athyrium*, as F6e has referred the *Diplazium brevisorum* of J. Smith, the species immediately after which Smith notices this, and from its affinity with *Athyrium Hohenackerianum* of Kunze(Schk. Fil. Suppl. tab. 126);—but which, nevertheless, M. F&e places in *Asplenium*. So different are the views of Botanists[^] in regard to the Genera of Ferns: Botanists, too, alike anxious to fix them upon a firm and solid basis.

Fig. 1. Upper portion of pinnae. */. 2.* Upper side of a fertile pinna, *f.* *3.* Involucre:—magnified.

DCCCCXIV.

ASPLENIUM ATTENUATUM, *BT.*

Frondibus linear-i-oblongis longe tenuiterque attenuatis (apice esepe proliferis)lobato-pinnatifidis basi subpinnatis apice integrimis, pinnis lobisque subrotundatis omnibus serrato-dentatis, stipitibus paleaceo-hirsutis, caudice repente.

Asplenium attenuatum. *Br. Prodr. Fl. Nov. Holl. p. 50.
Hook, et Grev. Ic. Fil. Tab. 209.*

HAB. New Holland; Port Jackson, *Brown, Fraser*; dry shady woods, Brisbane river, *Allan Cunningham*.

This species, though not new, seems of rare occurrence, and in the specimens before us, from Brisbane river, gathered by the late Allan Cunningham, shows itself under a new form, that of a prolific plant, throwing out roots and young fronds from the long attenuated apices whenever they touch the soil. The scales of the stipes, though, when seen with the naked eye, looking like chaffy hair, are, as represented accurately in *Icones Filicum*, above quoted, subulate, dark brown, membranaceous, coarsely reticulated, with four divaricated subulate segments at the base. As a species, this is very distinct from any other ye are acquainted with. The sori occupy nearly the centre of the lobes or pinnae, between the costa and the margin.

Fig. 1. Portion of a fructified frond, seen from beneath:—magnified.

DCCCCXV.

PTEBIS (Pellea) GERANIIFOLIA, *Raddi.*

Csespitosa glabra, frondibus cordiformibus profundissime quinquelobo-palmatis subcoriaceis opacis, laciniis lanceolatis pinnatifidis bipinnatifidisve, lobo primario intermedio basi cuneato, lobis ultimis ovato-lanceolatis integerrimis acutiusculis, sinubus acutis₃ stipite elongato rachibusque primariis nigro-ebeneis.

Ptens geraniifolia, *Raddi*, *Syn. Fil. Brasil* n. 110. *FiL Brasil*, p. 46. *Gardn. FiL Brasil, exsicc.* n. 39.

Pteris Pohliana, *Presl, Tent Pterid.* p. 145.

Pteris pedata, *Kunze, in Lincea*, 10. p. 522. *not Linn.*

Pteris pedatoides, *Desv.*

Pteris Mysurensis, *Heyne, in Wall. Cat.* n. 87.

Pellea pedata and geraniifolia, *Fee. Gen. Fil.* p. 130.

HAB. Brazil; *Hio, Raddi, Gardner, Armstrong, Mrs. Graham;* S. Brazil, *Sellow. Santa Martha, Columbia, Purdie. Cape of Good Hope, Menzies. Drége. Madagascar and Mauritius, Bqjer. Neelgherry, E. Indies, (WallicK). Volcano, Owhyhee, Macrae. Gallapagos, Douglas.*

Probably there are few Fern Herbaria, containing various suites of specimens, in which the present species of *Pteris* (with free or only forked veins), will not be found mixed and confounded with the original *Pteris pedata* of Linnaeus, a species having anastomosing veins, and therefore a *Litobrochia*, jPred. (*Doryopteris, J. Sm., Fee.*); so closely are they allied in other respects: and in the old and opaque specimens, of either kind, the venation is very difficult to be seen. The present is, indeed, usually a more compound species; but is often less so than is here represented. Desvaux was probably the first to distinguish the two, but Raddi's name is surely to be preferred to his. The above localities may be entirely depended on, as they are derived from specimens existing in my Herbarium.

Fig. 1. Fertile segments:—magnified.

Griffithiante.

N. O. Filices.

DCCCCXVL

GYMNOGRAMME MICROPHYLLA, *Hook.*

Pusilla densissime caespitosa glaberima, caudice gracili subrepente, frondibus triangulari-ovatis membranaceis subdianaphis pinnatis, pinnis profunde bi-tripinnatifidis, laciniis ovali-lanceolatis acutiusculis uninerviis monosoratis, stipitibus gracillimis fusco-purpureis nitidis omnino nudis, soris oblongis terminalibus, (nervo ante apicem evanescente).

H A B. Surureen, Khasiya, *Griffith.*

A delicate, quite new *Gymnogramme*, detected as far as we know only by the late Mr. Griffith, in the locality above given. It is evidently of the same group of the Genus as *G. leptophylla*, which the late Professor Link formed into a separate genus under the name of *Anogramme* in 1824, and in which he has been followed by M. Fée. *G. leptophylla* had already been referred to seven different genera.

The present plant grows on trees, and forms dense tufts; its short surculose caudices firmly entangled. The [^]tipites are extremely slender, filiform, from two to four inches long; the fronds from two to three inches in length, extremely delicate, first pinnated, the pinnae for the most part bipinnatifid, the lowest pinnae tripinnatifid, all cut into narrow segments, of which the ultimate ones are oblong-lanceolate, rather acute. The nerve occupies the centre of these, but disappears before the point, and it is the apex of this solitary nerve that bears the oblong prominent sorus.

Fig. 1. Portion of a fertile pinna, seen from above, *f.* 2. Lesser portion, seen from beneath. */. 3.* Ultimate segment of the same, with a sorus:—*magnified.*

DCCCCXVII.

ASPLENIUM MUCRONATUM, *Pr.*

Humile pendens (?) tenerimum glaberrimum, frondibus subsessilibus caspitosis linear-lanceolatis pinnatis, pinnis approximatis cordato-ovatis acuminatis refractis pinnatifidis, lobis ovatis mucronatis duobus inferioribus majoribus bitrlobis divaricatis, soris oblongis, stipitibus perbrevis fuscis nudis, rachi viridi gracili anguste alata.

Asplenium mucronatum, *PresL Delic. Prag.* 1. p. 178.

A, retortum, *Kaulf. En. Fil.* p. 171.

A. pteropus, *Kaulf. En. Fil.* p. 170. (*fide Preslii*).

A. laxum, *Raddi, Fil Bras.* p. 37. t. 22. bis. / 4.

HAB. South Brazil, on the rough trunks, of trees, *Chamisso, Pohl, Tweedie, Raddi.*

A very delicate and distinctly marked species. The tallest of the fronds is not a span long, extremely delicate, membranaceous, pale green, pinnated to the very apex with closely placed, refracted, cordato-ovate pinnae; each of which is rather deeply pinnatifid with ovate apiculated simple lobes; the two lower ones only being again divided, bifid, or more rarely trifid, and the lower and usually larger lobe laps over the pale coloured rachis, so as, on the underside of the Fern, to conceal a considerable portion. Sori rather large in proportion to the size of the plant, placed on the nerve which occupies the centre of the side lobes, the germinal and generally the lower lobes being sterile. The stipites rise several together, apparently from a tufted root, (our specimens exhibit no caudex): they scarcely exceed half an inch in length, and are dark coloured, while all the rest of the plant is a very pale subdiaphaneus green.

Fig. 1. Fertile pinna, seen from above. /. 2. The same seen from beneath:—magnified.

DCCCCXVIII

ASPLENIUM DELICATULUM, *Pr.*

Parvulum, caudice gracillimo longe repente filiformi nudo, frondibus sparsis brevi-stipitatis glaberrimis lanccolatis vel subtriangulari-ovatis acuminatis bipinnatis, pinnis remotis pinnulisve lanceolatis inciso-lobatis, laciniis angustis acutis submonosorosis.

Asplenium delicatulum, *Presl, Reliq. Hcenk. p. 47. t. 7.f. 3.*
Linncea. v. 9. p. 70.

HAB. Cordillera of Quito, and Trunks of Trees, Cuchaos,
Pceppig. Casapi, Peru, Mathetcs, n. 1785.

One of the smallest and most delicate of the *Asplenia*, not very unlike some species of *Hymenophyllum* or *Trichomanes* in general habit. Presl's figure in the *Reliquiae Haenkeanæ*, is extremely unsatisfactory, especially in the tufted fibrous roots. Afterwards the author describes in the *Linnæa*, from more perfect specimens, the real nature of the caudex. It is singularly long, slender, and creeping. In both those works the affinity of the species with *A. faniculaceum*, H. B. K. (Hook, et Grev. Ic. Fih t. 92), is noticed, and we ourselves have intimated that it is probably a starved state of that species. The long creeping filiform root or caudex, would also appear to confirm this suspicion. The caudex, here, however, is naked and wiry; in *A. faniculaceum* tomentose. This may vary, perhaps, according to situation and moisture. Poeppi^'s specimens, and those of Mr. Mathews, both in our possession, show no disposition to vary from the form here represented. Some of the specimens are indeed smaller, but none larger. It may be referred to the *Darea* or *Camopteris* group of *Asplenium*, having the segments and lobes so narrow as rarely to bear more than one sorus on each.

Fig. 1. Portion of the upper side of a pinna, f. 2. Fructified portion of the same, seen from beneath:—magnified.

DCCCCXIX.

ASPIDIUM (*Nephrodium*) SIMPLICIFOLIUM, *J. Sm.*

Frondibus pinnatis hirsutulis, pinnis paucis elliptico-oblongis integris subserratisve obtusis sessilibus, terminali maxima oblongo-lanceolata acuminata basi in&quali obtusa, arcubus venarum pluribus supra se positis soriferis, sonis non raro confluentibus meniscoideis, involucro obsolete subnullo, stipitibus paleaceo-setosis, rachibus venis venulisque hirsutis, caudice subrepente.

Nephrodium simplicifolium, *J. Sm. in Hook. Journ. of Bot. 3. p. 411, (name only). Presl, Epim. Bot. p. 50, (name only).*
Abacopteris simplicifolia, *Fée. Gen. Fit. p. 310.*
HAB. Samar, Phillipine Islands, *Cuming, n. 315.*

To this plant Presl refers the *Nephrodium acrostichoides* of J. Smith, (Cunning's n. 149, from Luzon), which to us appears abundantly distinct; and which J. Smith himself has subsequently referred to another Genus, *Cyclodium*, on acc^cin^a of his having observed it to have orbicular, not reniform, involucres. M. Fée goes further, and separates the *N. simplicifolium* generically both from *Nephrodium* and from *Cydonium*, and refers it to his new Genus *Abacopteris*, by characters which do not appear to us very valid or intelligible. Indeed, while he describes the indusium "reniforme, sinu affixum," he represents it most distinctly as *orbicular and jixed by the centre*. The indusium or involucre is however so indistinct, that we have failed to observe it in our specimens:—and we have further found the sori to be sometimes so confluent as to bring the genus very near to *Menisium*: and it is not impossible, that it may prove some undeveloped state of a more completely pinnated Fern.

Fig. 1. Portion of fertile pinna:—magnified.

DCCCCXX.

ASPIDIUM (*Nephrodium*) HETEBOPHYLLUM.

Frondibus csespitosis pubescenti-hirsutis simplicibus lingulato-oblongis brevi-acuminatis, sterilibus venis anastomosantibus, fertilibus angustioribus venis simplicibus, ambabus lobato-pinnatifidis lobis obtusis, involucris parvis deciduis, stipitibus brevibus hirsutis sparsimque squamosis, caudice repente.

Nephrodium Blumei, J. 8m. in Hook. Journ. of Bot. 3.p. 411.
exclus. syn.

Haplodictyum heterophyllum, Presl Epimel. Hot. p. 51. Féé,
Gen. Fil. p. 309.

HAB. Samar, Phillipine Islands, Cuming, n. 322.

Mr. J. Smith was the first to notice this Fern, and he referred it, in his account of the Ferns of the Phillipine Islands, collected by Mr. Cuming, to *Nephrodium*[^] being aware of course of the presence of an involucre to the sorus: and, believing it to be identical with the *Gymnogramme cane&cens*, of *Bluln. Fil. Java*, tab. 93, he called it *Nephrodium Blumei*. The fertile fronds (supposing the involucre to exist in an early stage) are quite those of *Nephrodium*, but the sterile ones are remarkable for the anastomosing of the veins at a little distance from the primary vein. Hence JPresl has formed of this Fern a new genus, between *Pleocnemia* and *Nephrodium*[^] in which he has been followed by M. F6e. Presl is assuredly correct in excluding Mr. Smith's reference to the *Gymnogramme canetcens*, Blume, (Gonophlebium, Presl); that plant has a long stipes, very acuminate and deeply pinnatifid (below pinnate) fronds, and apparently never any involucre; in shape very much resembling our *Aspidium* (*Nephrodium*) *Skinneri*, Tab. 924.

Fig. 1. Portion of a fertile frond. /. 2. Portion of a sterile ditto:—magnified.

DCCCCXXI.

ASPIDIUM (*Lastrea*) VOGELII.

Humile pilosulum, fronde triangulari-ovata acuminata membranacea pinnata, pinnis oblongo-lanceolatis basi decurrentibus lobato-pinnatifidis obtusis infimis subpinnatis, lobis pinnulatis ovatis obtusis, involucris reniformi-orbicularibus paucisetosis, stipite gracili brevi basi setuloso, caudice gracili repente.

HAB. Fernando Po, Dr. Vogel, (Niger Expedition).

Several specimens of this little species of *Lastrea* are in the Niger collection formed in Fernando Po, after the Expedition had left the Niger. One of the largest, and one among the smallest of the series are here represented. The frond is thin and membranaceous, semipellucid; the largest scarcely exceeding four inches in length, the upper extremity pinqatifid, ending in a long tapering point; the lower portions pinnate, even bipinnate at the base. The rachis and primary veins, and even the surface, especially beneath; are pilose. The pinnae are more or less deeply lobed, decurrent with tta rachis. at the base. Son generally one to each lobe or segment. Involucre moderately large, orbiculari-reniform, denticulate at the margin, and bearing on its superior surface four or five spreading, jointed, hairs. The stipites are about half the length of the frond, slender, with a few bristly hairs at the base. Caudex slender, almost filiform, creeping.

Fig. L Pinna with fructifications, seen from beneath. f_m 2.
Involucre:—*magnified*.

DCCCCXXII.

ASPIDIUM (*Nephrodium*) HOOKERI, *Wall*

Caespitosum bipedale, fronde subcoriacea lanceolatoacuminata inferne longe attenuata pinnata, pinnis subapproximatis lineari-oblongis brevi-acuminatis Berratis basi cuneatis sessilibus inferioribus plurimis paribus nanis suborbicularibus, siccitate venis inconspicuis, costa venisque subtus pubescentibus, sois parvis subbitrinalibut, involucris integrerimis, stipitibus brevibus incrassatis.

Aspidium Hookeri, *Wall. Cat. n. 338. ad p. 64,*
HAB. Dindigul, 4,000 feet elev., Madras Presidency, Dr.
Wight, Herb. Wight, propr. Crypt, n. 116; Ceylon, Mrs.
Geril. Walker.

This we consider a very distinct *Nephrodium*, somewhat indeed allied to *N. unitum*, Schott., but having nothing of the harshness and rigidity of that species, and with pinnae much less deeply lobed, rather serrated than lobed; and the lower pinnae, unusually small and cordate or rotundate, are extended down to the base of the stipes; these lesser pinnae are always barren.

Fig. 1. Pinna from a barren specimen; nat. size. f. 2. Portion of a fertile pinna, seen from beneath. /. 3. Involucre:—magnified.

Sellowiana.

N. (J. Filiccs.

Dcqccxxm.

ASPIDIUM (*Lastrea*) KLOTZSCHII, *Hook.*

Humile cscspitatum, frondibus digitalibus et ultra deltoideo-ovatis acuminatis subcoriaceis glabris, pinnulis oblongo-ovatis basi oblique cuneatis pinnatifidis, laciniis ovatis subspinuloso-acutis simplicibus vel inciso-dentatis, soris in singulo lobo solitarii reniformi-orbicularibus membranaceis fuscis, stipitibus gracilibus frondes superantibus inferno squamosis, squamis subulatis patentibus intense fuscis.

Polystichum aspidioideum, *Klotzsch*, in *Herb. Nostr.*

HAB. South Brazil, *Selloic.*

Our specimen of this Fern is from Dr. Klotzsch of the Royal Berlin Herbarium, with his name attached; but I am ignorant whether the species be any where published or not. Finding the involucre to be reniform, and not orbicular and peltate, as in *Polystichum*, I venture to place it in the *Lastrea* group of *Aspidium* or, according to the views of M. F6e, so unsettled is the nomenclature of the genera of Ferns, in *Aspidium* proper. It has, however, unquestionably, very much the general aspect of *Polystichum*, the texture of the frond being firm and somewhat coriaceous and glossy, and the teeth and lobules of the pinnules almost spinulose. It appears to be a distinct species from any hitherto described.

Fig. 1. Upper side of a fertile pinnule. /. 2. Under side of ditto, with fructifications.—magnified.

DCCCCXXIV.

ASPIDIUM (*Nephrodium*) SKINNERI, *Hook.*

Subhumile glaberrimum, fronde lanceolata tenui-acuminata pinnatifida basi pinnata, lobis approximatis pinnisque oblongis sursum subfalcatis acutiusculis, sonis copiosis, involucris longe ciliatis, stipite brevi nudo, caudice crasso repente squamoso.

HAB. Guatemala, G. U. Skinner, Esq.

Half a score specimens of this pretty Fern, for which I am indebted to Mr. Skinner, uniform in their character, would lead to the inference that such is the normal condition of the plant, and not an imperfectly developed state of some more compound *Nephrodium*. If I am right in this conjecture, it is a very distinct species, and not likely to be confounded with any other. The caudex is stout, repent, and scaly. The stipes, several arising from near the apex of this caudex, short, two inches or more long, rather slender, naked, except a few scales where the base joins on to the caudex. Frond six to eight inches long, somewhat membranous, not opaque, very much tapering at -the point, and then the lobes gradually become serrature³: the base is not much contracted, but it is truly pinnate with about two pairs of rather remote pinnae, above them the frond is of one piece, lanceolate, cut into many approximate lobes more than half way down to the costa; lobed like the pinnae, oblong, with a slight curvature upwards, rather acute. Main rachis slightly pubescent. Fructifications, when present, copious on the centre of every vein. Venation, that common to *Nephrodium*. Involucrum reniform, membranous, reticulated, and bearing long cilia on the margin.

Fig. 1. Upper side of a portion of the fertile frond. */. 2.* Underside of ditto, *f.* *3.* Involucrum:—*magnified.*

Simonsiana.

N. O. Filices.

DCCCCXXV.

ASPLENIUM (*Thamnopteris*) SIMONSIANUM, *Hook.*

Caespitosum glaberrimum subcoriaceo-membranaceum, frondibus (sesquipedalibus) elongato-lanceolatis abrupte anguste acuminatis, basi in stipitem perbrevem longe attenuatis angusto-marginatis, venis crebris ante marginem apicibus coalitis in lineam continuam marginalem, sōns copiosissimis erecto-patentibus totam paginam fere occupantibus.

HAB. E. Indies, Khasya hills, *Simons*, n. 432.

This is a true *Neottopteris* of J. Smith, (*Thamnopteris, Prest*) as far as the union of the apices of the veins into a continued *intramarginal* line is concerned; but such a genus is most unnaturally separated from several simple-fronded *Asplenia*. As a species its affinity is with *Neottopteris Phyl-HtidiSj* J, Sm. (*Asplenium, Ddt[†]* and no less with *N. stipitata*, J. Sm. (which I cannot distinguish from the preceding), but the fronds are here much narrower, more suddenly and narrowly acuminate, and at the base very much more attenuated. My only specimens are from Mr. Simons.

Fig. 1. Fertile portion of the frond, seen from beneath:—magnified.

Spruceatue.

N. O- Filices.

DCCCCXXVL

ASPLENIUM LORIFORME, *Hook.*

C&spitosum parce deciduo-squamulosum membranaceum, frondibus longissime lanceolatis loriformibus obscure serratis angustissime longe acuminatis inferneque in stipitem per brevem attenuatis, soris copiosis approximatis oblique patentibus.

HAB. On trees in forests at Tanaii, and other places near Par&, Amazon River, *Spruce*, n. 18.

An unusually narrow and elongated simple-fronded species of *Asplenium*, of a very membranaceous texture, pale and bright green colour, remarkably and gradually acuminate at the apex, and no less attenuate at the base. It belongs to the same group as *Aspl. serratum*, along with our *Aspl. Griffithianum* (v. Tab. 928); and is easily distinguished by the above characters.

Fig. 1. Portion of a fertile frond, seen from beneath:—magnified.

NuttalliatKB.

N. O. Filices.

DCOCCXXVIL

ASPLENITJM PINNATIPIDUM, Nutt.

Caespitosum glabrum, frondibus parvis divaricatis subcoriaceis hastato-lanceolatis longe acuminatis subtus setuloso-squamulosis inferne profunde fere ad rachin pinnatifidis, lobis cordatis sinuatolobatis dentatisve, medio semipinnatifidis lobis obtusis dentatis, apice subintegerrimis, soris paucis in singulo lobo linear-i-oblongis demum confluentibus, stipite inferne ebeneo, caudice brevi repente.

Asplenium pinnatifidum, Nutt. in Gen. of N. Am. PI. 2.p. 251.

**HAB. U. States, Banks[^]f the Schuylkill, near Philadelphia,
Nuttall, Dr. Bromfield, Mr. J. Me. Nab. Tennessee, Nut-
tall. Sandstone rocks, "Mine la Motte," Southern Missouri.
(Herb, nostr.)**

This appears to be a rare species, and gathered by very few Botanists. Muhlenberg, according to Nuttall, seems to have confounded it *with Aspleridum (Camptosorus, ZA.) rhizophyllum*, of which it has a good deal the habit; but, besides the very different venation peculiar to the latter plant, its fronds are entire (not pinnatifid), rooting and proliferous at the extremity.

Fig. 1. Portion of a frond, seen from above. /. 2. Fertile lobe, seen from beneath:—magnified.

DCCCCXXVIIIL

ASPLENIUM GRIFFITHIANUM, *Hook.*

Csespitosum subcoriaceum minute deciduo-squamulosum, frondibus lanceolatis breviter acuminatis basi longe in stipitem perbreuem attenuatis serratis acumine integerrimo, sonis totam fere paginam occupantibus, venis obscuris remotiusculis.

HAB. Mishmee, E. Indies, W. *Griffith, Esq.*

In the general form of the frond, in colour, texture, in the acuminated point, and in the attenuated base, this has a very close affinity with our *Asplenium Simonsianum*: it differs, however, in the shorter fronds, in the serrated margins, and ~~<in~~ the more remote and more patent lines of fructifications. If^ however, the frond be held up between the eye and the light, the nerves will be found to be all free, each of them terminating 9. little below the apex of a serrature, whereas in *A. Simonsianum*, they unite so as to form an intramarginal nerve, characteristic of the Genus *Neottopteris*, J. Sm:—*Thamnopterus*, of FresL

Fig. 1. Portion of the underside of a fertile frond:—*magnified.*

Miqueliancc.

N. O. Filiccs.

DCCCCXXIX.

ASPLENIUM SUBHASTATUM, *Hook.*

Frondibus cespitosis subcoriaceo-membranaceis opacis glaberrimis lanceolatis acutis basi subtruncato-cuneatis utrinque lobo obtuso subhastatis, stipite nudo longitudine frondis, sonis linearibus, venis erecto-patentibus remotiusculis,

HAB. Caraccas, (en Herb. *Miguel.*)

My specimen of this, as it appears to me, very distinct species of *Asplenium*, I owe to the kindness of Professor Miquel, who sent it to me with many other plants from Caraccas. I can find it nowhere described, and I venture to name it *subhastatum*, from the disposition of the majority of the fronds in my possession to have a spreading lobe on each side of the base, giving the frond a hastate form. It has the habit, rather of a *Scolopendrium* than of a single fronded *Asplenium*; but the fructifications are truly those of the latter Genus, as is the venation; each vein terminating in a slightly clavate apex, and free from any union with the adjacent veins.

Fig. 1. Portion of a fertile frond, seen from beneath:—magnified.

DCCCCXXX.

ASPLENIUM SCOLOPENDRIOIDES, *J.* 8m.

Fronde simplici lanceolata subito caudato-acuminata integer-rima basi in stipitem perbrevem longe attenuata, sonis linearibus, involucris subgeminatis superiore angustissimo reniformi, venis approximatis patentibus.

Asplenium scolopendrioides, *J. Sm. in Hook. Journ. of Bot.* 3.
p. 408. (name only).

HAB. Leyte, in the Phillipine Islands, *Cuming*, n. 318.

This is another *Asplenium*, nearly allied to our *A. biforme* t. 926, and *A. Griffithianum*, t. 928, remarkable for the sudden, narrow, tail-like apex of the frond; and I find a peculiarity also in the involucres, for on the inner or upper side, a "nerve-like" line will be seen to run parallel with and of the same length as the involucre, leaving a narrow area between it and the contracted involucre attached to the nerve. This line is caused by a slender membrane, evidently a portion of the involucre, remaining after the dehiscence. It gives the appearance of a double involucre, or of one opening in the middle; ahd it was probably this which suggested the name of *scolopendroides* to Mr. J. Smith.

Fig. 1. Portion of the underside of a fertile frond:—magnified.

DCCCCXXXL

ASPIDIUM (*Lastrea*) BOUTONIANUM, *Hook.*

Fronde glaberrima majuscula membranaceo-subcoriacea oblonga acuminata pinnata, pinnis sessilibus oblongo-linearibus obtusis usque ad apicem lobato-pinnatifidis, lobis ovalibus obtusis sinuato-serratis, yenis pinnatis apicibus clavatis supra albo-punctatis, soris venuias terminantibus, involucris ciliatis, rachi lsevissima supra pubescente, stipite breviusulo versus basin articulato deciduo nudo, caudice longissimo squamis peltatis erosionis tecto.

HAB. Eastern summit of the "Montagne longue," Mauritius,
M. Bouton.

I can no where find that this very distinct *Aspidium* (or *Lastrea*) is described, and yet it is one of the best marked of the family. It appears to have been only collected off the spot just mentioned, and is there said by my correspondent to be extremely rare in fructification. It is remarkable for the white spot or dots on the upper side of the pinnae where the veins terminate, which appear to be due to a resinous substance exuding from the vein, and forming a small scale which may be easily removed entire,—for the articulated stipes, but, above all, for the great length of the caudex, about the thickness of a writing pen, "which takes root in the rocks and climbs upon the adjacent trees;" this is densely clothed with chaffy peltate scales, torn and erode at the margin.

Fig. 1. Lobe of a pinna, showing the clavate apices to the veins. */>. 2.* Fertile lobe, seen from beneath. */>. 3.* Involucrum: —*magnified.*

DCCCCXXXil.

ASPLENIUM FRAGILE, *Fresh*

Pusillum csespitosum fragile, frondibus submembranaceis linearibus acutis pinnatis, pinnis oblique deltoideo-ovatis rhomboideisve acutis subflabellatis inciso-lobatis dentatisve, soris involucrisque oblongis, rachibus viridibus superne fuscis, stipitibus atro-fuscis non raro bulbilliferis, bulbillis viviparis,

Asplenium fragile, *Presl*, *Tent. Pterid.* p. 108. *Klotzsch*, in *Linncea*, v. 20. p. 355.

A. *stoloniferum*, *Presl*, *Reliq. Hank?* 1. p. 44. t. 6.f. 4. *excl Syn.*

HAB. Mountains of Peru, *Haenke*, Chimborazo, *Humboldt*, on rocks at an elev. of 14,000 feet, *Jameson*. Paramo de Mucuchies, Columbia, *Momitz*. n. 326.

W&e it not for the curious little viviparous bulbilli seen upon the stipes of the present plant, it would be difficult to say in what respect this species differs from *Aspl. viride* of the European Alps: and -it is possible that this peculiarity may originate in its elevated locality, which is no doubt very considerable at all times on the Andes of Columbia. Haenke's countries are always doubtful, for his plants, as given by P[^]esl. Dr. Jameson's are stated with much exactness. The accurate Kunze in *Linnaea*, v. 13. p. 140, has given this Fern as an inhabitant of Mexico ("ad muros Hacienda de Begia legit C. JEKrenberg"); but as he describes the largest fronds (frondes maximsc) to be only two inches long, and the stipes (short) as "glandular, paleaceo^is and green/" the probability is, I think, that the two plants are different.

Fig. 1. 2. Fertile pinna, seen from beneath:—magnified.

Dccccxxxin.

,

ACROSTICHUM (*Neurocallis*) AUREONITENS, *Hook.*

Cespitosum, frondibus carnosso-crassis subcoriaceis biformibus subtus (junioribusque supra) squamis imbricatis copiosis aureonitentibus tectis interne reticulatim venosis, sterilibus spathulatis integris integerrimisque costatis in stipitem brevem attenuatis, fertilibus longe stipitatis pinnatis, pinnis oblongis obtusis costatis, stipitibus aureo-squamosis.

HAB. Galapagos, *Owning*, n. 109. Chatham Island, Galapagos, *Capt. Wood, JR. N.*

A very beautiful and remarkable Fern, which I possess from two travellers, but both specimens probably gathered in the same locality, Chatham Island, of the Galapagos. From apparently a small fibrous root, many fronds arise, the whole underside and stipites of which are densely clothed with toothed lanceolate and much acuminate scales, of a glossy golden colour. The upper sides of the fronds have only a few scattered chaffy hairs. The sterile fronds are simple (undivided), but the fertile ones are pinnated with 5-9 rather small, oblong, ob*usp, sessile pinnae, the underside of which is uniformly clothed with capsules mixed with the chaffy scales:—the venation, very difficult to be accurately distinguished, consists of anastomosing veins, which form oblong, hexagonal areoles: so that this would be a true *Acrostichum* of Presl, probably a *Neurocatys* or *Cheilolepton* of M. F6e (if the two genera are in any way really distinct): yet having a habit very different from any known species of them.

Fig. 1. Portion of a sterile frond, showing the anastomosing of the veins. */. 2.* Portion of a fertile pinna, seen from above. */. 3.* Portion of the same, seen from beneath, with the scales in part, and the capsule in part, removed to show the reticulations. *f.* 4. A scale from the frond:—*magnified.*

DCCCCXXXIV.

POLYPODIUM MACROCARPUM, *Presl.*

Humile coriaceum, frondibus oblongo-ovatis obtusis profunde pinnatifidis subtus stipiteque equilongo squamis ovatis acuminatis peltatis serratis paleaceis, soris rotundatis majusculis demum confluentibus, caudice longe repente squamoso.

Polypodium macrocarpum, *Presl*, *Reliq. Hank.* p. 23. t.l.f.4.
Pleopeltis pinnatifida, *Gill*, in *Hook, and Grev. Ic. Fit.* t. 157.
HAB. Peru, *Hcenke*, (*Presl*), Rocks, Puruchucu, Peru, *Matthews*, n. 600. Andes above Ticicaca, Bolivia, *Mr. Pentland*. Massa Fuera, S. Pacific, lat. 34°, *Cuming*, n. 1352.

When this figure was drawn from a specimen gathered at Massa Fuera, South Pacific, in a latitude corresponding with Valparaiso, I little suspected it would prove the same as *Pleopeltis pinnatifida*^ Gillies in *Icones Filicum*; but such proves to be the case, and further that this is identical with the *Polypodium macrocarpum* of *Presl*, a name which should unquestionably, on every account, be preferred. It is a true *Polypodium*^ of the same group as our *Polypodium vulgare*, and we are now able to give some additional habitats for this widely extended species. Some of our specimens, from the Andes of Ticicaca, (14,000 feet above the level of the sea), have the fronds exactly spathulate, a little lobed at the base: but this is to be considered as an abnormal form, and from this we have aH the intermediate states to the many lobed and elongated fonn given in the *Icones Filicum*.

Fig. 1. Portion of a frond, upper side. *f. 2.* Fertile lobe, seen from beneath. *gf.* 3. Portion of a lobe with a sorus. */. 4.* Scale from the back of a frond:—all more or less magnified.

DCCCCXXXV.

GYMNOGRAMME RUT2EFOLIA, var Hispanica.

Humilis, pilis articulatis undique longe villosa, frondibus oblongis obtusis pinnatis, pinnis subchartaceis trapezoideo-ovatis obovatisve basi cuneatis subpetiolatis integris inciso-lobatisque inferioribus nunc iterum subpinnatis, lobis cuneatis, soris linearibus brevinsculis liberis, stipite frondem eubaequante, caudice brevi horizontali crasso, radicibus dense fibroso-caespitosis.

Grammitis rutaefolia, *JBr. Prodr. Fl. Nov. Holl* p. 2.

Gymnogramme rutsefolia, *Hook, et Grev. Ic. Fil. t. 90. JLehm. PL Preiss. 2. p. 110.*

Pleurosorus rutaefolius, *F  e, Gen. Fil. p. 180.*

6. *major*; pinnis omnibus pinnatifidis, pilis brevioribus magis glandulosis.

Gymnogramme subglandulosa, *Hook, et Grev. Ic. Fil. t. 9.*

Plerosorus cuneatus, *F  e. Gen. Fil. p. 180.*

7. *Hispanica*; frondibus tenuioribus magis herbaceis. {*Tab. nostr. DCCCCXXXV*}.

Grammitis Hispanica, *Cosson, Notes sur quelques Plantes Nouv. du Midi de VJEspagne*, p. 48.

HAB. 7. South of Spain; Puerto del yiento, in the Sierra de Honda, 1849, (n. 543), and in the Sierra Nevada, in clefts of rocks at Cortijo de la Vibora, 1851, Bourgeau.

We would gladly, if we could, with propriety, have kept the European *Gymnogramme* distinct, as M. Cosson has done, from the Australian *G. rutaefolia*. But the utmost we can make of it is a variety, of a thinner texture and greener colour. It is, however, a remarkable fact, that a Fern common enough in Australia, to which country it was, till now, supposed to be peculiar, should prove to be an inhabitant of the elevated Sierras in the South of Spain! —We are also satisfied that we do right in uniting our *Gymnogramme subglandulosa* with the present species.

Fig. 1. Sterile pinna, f. 2. Fertile pinna, seen from beneath :—magnified.

DCCCCXXXVI.

DIPLAZIUM (*Oxygenium*) CORDIFOLIUM, *Bl.*

Frondibua simplicibus elliptico-ovatis acuminatis integerrimis basi cordatis v. subsagittatis (non raro proliferis) lobato-pinnatifidisve vel pinnatis, pinnis paucis (3-5) lateribus ovato-lanceolatis. sessilibus terminali maxima basi cordata v. sagittata, involucris plerisque elongatis angustissimis, venis marginem versus anastomosantibus, stipitibus glabris nudis basi solummodo parce squamosis.

1. Frondibus integris v. basi lobato-sagittatis.

Diplazium cordifolium, *Bl. En. Fit. Jav. p. 190. Hook. Ic. PL t. 184.*

Asplenium ovatum, *Wall. Cat. n. 195.*

2. Frondibus integris pinnatisve.

Oxygenium ovatum, *JPresl, Tent. Pterid. p. 118. J. Smith, in Hook. Journ. Bot. 4. p. 178. (1842).*

Callipteris ovata, *J. Smith, in Hook. Journ. Bot. 3. p. 409. (1841). Fée. Gen. Fil p. 219.*

Anisogonium integrifolium, *Presl, Tent. Pterid.p. 116. Ejusd. Epimeh Bot. p. 92.*

Diplazium integrifolium, *Blume. En. Jav. p. 190. (JidePresl.) HAB. Java, Blume. Singapore, Wallich, Sir Wm. Norris. Isle of Leyte, Cuming. n. 307. Penang, Sir W. Norris.*

We had already represented the simple fronded state of this plant in the *Icones Plantarum* above quoted, and we now give the pinnated form of the same, which has, by some, been mistaken for a different species. We cannot agree with those Botanists who refer it to the genus *Callipteris*, of which *C prolifera*, *Borry*, is the type. *Presl* is quite mistaken at p. 92 of his *Epimelia Botanica*, in stating that Mr. J. Smith's *Callipteris* (or *Oxygenium*) *ovata*, is not the *Asplenium ovatum* of Wallich. The simple fronded state of the plant is identical with that, as is *Diplazium tordifolium* of Blume, which name, as the oldest published, we adopt. Some of our simple fronded specimens are nearly a foot long.

Fig. 1. Portion of the underside of a fertile pinna:— magnified.

DCCCCXXXVIL

ASPLENITJM (Asplenidictyon) FINLAYSONIANUM, *Wall.*

Frondibus subcoriaceis opacis oblongis pinnatis, pinnis 5-7 remotis oblique patentibus ovato-lanceolatis longe acuminate subintegerrimis margine superiore infra medium productis angulatis vel acute auriculatis basi cuneatis in petiolum brevem attenuates, terminali subrhomboides siccis triloba lobis inaequalibus, venis marginem versus anastomosantibus apiceque arcu conjunctis, stipite rachique compressis deciduo-subpaleaceis, caudice crasso squamoso fibroso. *JhosPei** yxu-rrt*

Asplenium Fiplyoonianum, *Wall.* Cat. n. 2682. V. /*?*-
Asplenium integerrimum, *Wall.*, in *Hook, et Grev. Ic. Fil.* t. 136, (*inaccurate in the venation*).

HAB. Nepal and Kamaoun, *Wallich.* Mishmee, *Griffith.* Gowhatta, Assam, *Simons.*

Our figure of the venation of the present East Indian Fern, in *Icones Filicum*, is inaccurate, and it is a plant that deserves to be better illustrated; the more so as our next plate will represent a Fern with very similar habit, and exactly similar venation, from the West Indies. Of the two Mr. J. Smith proposes to constitute a genus "*Asplenidictyon*!" Trusting, probably, to the figure in the *Icones Filicum*, Presl has placed *Aspl. Firilaysonium* in his group of *Asplenium*^ "Venis venuisque apice liberis." In the "venulae apice¹ arcu transverso conjunctae," our plant approaches Presl's *Thamnopteris* section of *Asplenium*; but, more than this, the veins anastomose, more or less, below the apices, as in *Omgonium*, where however the involucres are *diplozoid*. It borders too upon *Hemidictyon*, differing in wanting the marginal vein which unites the lateral veins in *Asplenium*(*Hemidictyon*)*marginatum*. *Allantodia Brunonis*, *Wall.*, again, has the venation anastomosing, like our species under consideration, but *the ultimate veinlets are all free and clavate within the margin.. Such a structure in the venation affords beautiful sectional characters, but does not necessarily afford sound generic distinctions.

Fig. 1. Portion of a fertile piuua, seen from beneath:— magnified.

DCCCCXXXVIII

ASPLENIUM (Hemidictyon) PURDIEANTTM, Hook.

Frondibus subcoriaceis opacis cordato-rotundatis pinnatis, pinnis 5-7 patentibus submembranaceis lateralibus oppositis . subsessilibus ovato-lanceolatis breviter acuminatis basi oblique cuneatis deorsum obtuse productis obtusangulis (seu pinnis inasquilateris), terminali subrhomboido-ovata acuminata sequilatera nunc subhastata lobis inscqualibus, venis marginem versus anastomosantibus, stipite rachique compressis subulatis fusco-paleaceis, radice fibrosa.

HAB. Moist woods, La Fundacion, Jamaica, *Purdie*.

This appears to be an entirely undescribed Fern of peculiar habit, having the fructification of an *Asplenium* (verum) and the venation of *Hemidictyon*[^] except that it wants the marginal vein which unites the terminal veinlets in the latter genus. This will consequently rank with the *Asplenium Finlaysonianum*, represented on our preceding plate; differing however in the shorter broader frond, the paleaceo-setose stipes and rachis, the less acuminated pinnae, which have the lowest half the broadest and most disposed to form an angle or auricle, (not the upper one). The root here too is simply fibrous, not forming a knob-like caudex.

Fig. 1. Portion of a fertile pinna:—magnified.

D C C C C X X X I X - XI.,

DIPLAZIUM (*Oxygenium*) ELEGANS.

Frondibus simplicibus v. pinnatis, pinnis lato-lanceolatis sessilibus anguste acuminatis serratis, lateralibus 2-9 suboppositis intermedia vix majore nunc basi hastato-biloba, involucris plurimis elongatis angustis, venis hie illic plerumque margineni versus anastomosantibus, stipitibus glabris nudis basi solummodo parce squamosis.

Oxygenium elegans, &. Smith in Hook. Journ. of Hot. 4. p. 178. {name only}.

Callipteris elegans, in Hook. Journ. Bot. 3. p. 409. Fée. Gen. Fil. p. 219.

Anisogonium elegans, Fresl, Epimeh Bot. p. 293.

A. grossum, Fresl, Epimel. Bot. p. 293. forma grandis, (name only, sub *A. elegante*).

HAB. Luzon, Cuming, n. 276, and at Leyte. n. 305. (larger form.)

I cannot but think that Mr. John Smith is correct in removing this to *Oxygenium* of Presl, rather than retaining it in *Callipteris* as M. Fée has done, though it has a stronger claim to a place there than our *Diplazium cordifolium* has; yet the venation of the present plant is not that of *Callipteris prolifera*, the original species, where all the lateral veinlets of a fascicle combine with the adjacent ones, and form as it were a secondary vein, parallel with the main vein. Here the union of the long parallel veinlets is comparatively rare, as shown in our figure 1. It is unquestionably a Fern closely allied to our *Diplazium cordifolium*, (see Tab. 936), and even more liable to vary, from a simple entire frond to pinnated* with seven or eight pinnae; but these pinnae are always serrated, narrower, and the ultimate pinna nearly uniform with the others, and the venation is more simple. Our large state of th§ plant Mr. Presl seems to have made a jiistinct species, (under the name of *grossum*), but we find all intermediate gradations in our own or Mr. Smith's Herbarium. Nearly allied to the large state of this is the pinnated form of *Diplazium alismcejblum*, Presl, of which the simple fronded form is figured in the Reliq. Hsenk. t. 8. f. 3. (Cuming. n. 116); but that has larger dark coloured fronds, with copious, coarse, black scales on the stipites and rachis, and venation as in our *Dipt. cordifolium*. Of this, however, Presl has made the Genus *Ochlogramma* (Epimel. Bot. p. 93), and Fee the Genus *JPteriglyphis*, Fée, Gen. Fil. p. 219. t. 18. B.:—each author giving a new and different specific name.

Fig. 1. Portion of a fertile pinna, seen from beneath:—
maanified.

DCCCCXLL

POLTPODIUM SETIGEBUM, *BL*

Frondibus fasciculatis longe sitipitatis linear-lanceolatis siccitate membranaceis subpellucidis, stipite ferrugineo-setosis, soris orbicularibus costam approximatis utrinque uniserialibus, venula infima superiore medio sorifera, radice caespitosa.

Polypodium setigerum, *Blume*, *En. FiL Jav.p.* 123.

Grammitis fasciculata, *Blume'*, *Fl. Jav.*, *FiL p.* 112. *t 47. l. 2.*
HAB. Epiphytal[^] on the trunks and branches of trees, summit of Mount Gede, Java, *Blume[^]Thos. Lobb.*

This is a very beautiful species, with an aureo-fuscous tint, increased by the deeper golden brown of the copious patent sefse., Blume first referred it to *Polypodium*, and we think rightly, and afterwards removed it to *Grammitis*, and he both figures and describes the *sori* as "*rotundi?*" Our specimens from Mr. Lobb precisely accord with one from Blume himself in our Herbarium. Thin and membranous as the texture appears when dry, it becomes thick and somewhat pulpy when soaked. Held between the eye and the light the frond, when magnified, is seen to be full of pellucid points or minute areoles.

Fig. 1. Portion of a fertile frond, seen from beneath—• magnified.

DCCCCXLIL

POLYPODIUM LEUCOSORUM, *By.*

Frondibus pendulislanceolatis subcoriaceis acuminatis(acumine subintegro) profunde pinnatifidis glabris subtus flavofuscis opacis, lobis oblongis e lata basi sensim attenuatis obtusis integerrimis vel leviter sinnato-crenatis, soris biserialibus in singula lacinia inter costam et marginem junioribus pulverulento-albis, venis internis, venuis primariis superioribus apice soriferis, stipite glaberrimo nudo gracili fronde multo^{ies} breviore, caudice crasso setaceo-paleaceo.

Polypodium leucosorum, Bojer, Hort. Maurit. p. 417. (name only.)

HAB. Old mossy trees, forests of Nouvelle Decouverte and Grand Port, Mauritius, *Bouton. Bourbon, Carmichael.*

I possess a specimen of this plant from Mauritius, sent by Mr. Bouton, with young fructification, exactly as he observes, "couverte d'une poussiere blanche," and another from Bourbon, gathered by the late Capt. Carmichael. M. Bouton justly notices its affinity with *Polypodium rigescens* of Bory, figured at Tab. 216 of Hook and Greville's *Icones Pilicium*. Its very much larger size, different colour, differently shaped lobes, and segments, and, especially, the young white son, are sufficiently characteristic of the present, as a species. A comparison of the figures of the respective species will satisfy any unprejudiced mind of the differences between the two.

Fig. 1. Lobe of a fertile frond, seen from beneath:—magnified.

N. O. Filices.

DCCCCXLIII

POLYPODIUM LEUCOSORUM, *Boj.*

(status senilis)

See description under Tab, 942.

HAB. Isle of Bourbon, "Polypodium, No. 3, from the Museum de PHistoire Naturelle."

The entire absence of white son, the very copious fructifications* and the usually broader and more obtuse segments of the frond misled me as to the identity of the species of this Fern; and it is only since the ljj;ograph was executed and all the impressions worked off, that I have satisfied myself it is the old state of *Polypodium leucosorum*, every segment loaded with brown son. Some of our specimens, too, prove that the sori vary to elliptical as shown at fig. 2. and the receptacles of the capsules are probably invariably oblong as shown fit fig. 1.

Fig. 1. Segment of a fertile frond, from which the sori are removed. / 2. fertile segment seen from beneath, with the sori more elliptical than usual:—magnified.

DCCCCXLIV-

POLYPODIUM (Ctenoptpris) PELLUCIDUM, *Kaulf.*

Glabrum subcoriaceum, frondibus ovatis oblongisve profunde pinnatifidis, laciniis valde approximatis patentibus oblongis crenato-dentatia vel serratis vel erecto-patentibus remotis marginatis pinnatifidis, soris copiosis rotundatis v. ovalibus saspe confluentibus, venis dichotomis striisque interstitialibus pellucidis, stipite rachique validis fuscis nitidis, caudice crasso repente squamoso.

- a. fronde lata laciniis acutiusculis serratiq,
Polypodium pellucidum, *Kaulf.* En. *FiL* p. 101. *Gaudichaud*,
in Freyc. Voy. p. 356. Hook, et Arn in JBot. of Beech. Vby.
p. 103. (TAB. NOSTB. *larger figure*).
 - /3. fronde àugustiore coriacea, venis vix pellucidis, laciniis numerosis approximatis erecto-patentibus obtusis crenato-serratis.
 - 7. fronde elliptica, laciniis valde approximatis obtusissimis obscure crenatis. (TAB. NOSTR *smaller figured*)
 - 8. fronde elongata bipinnatifida, laciniis primariis remotis.
Polypodium myriocarpum, *Hook. Ic. FiL t. 84.* (*See also the next plate, t. 945.*)
- HAB. Oahu, Sandwich Islands, *Chamisso, Douglas, Beechey, Dr. Diell.*

Without a coloured figure of a magnified portion of this Fern it would be difficult to give an idea of the peculiar venation of the plant. The frond is opaque, but the veins are pellucid and of a rich tawny colour, when seen between the eye and the light; and besides the free (clavate at the extremity) veinlets, of which the upper one bears the sorus, there is, what I here call, an interstitial pellucid stria always communicating with a crenature or sinus between the teeth of the margin, usually the most conspicuous of the pellucid lines. I am satisfied that the ^bove mentioned varieties belong to one and the same species, and that they pass the one into the other.

Tab. 944 exhibits vars. and (the larger specimen) 7. not size.—Fig. 1. Portion of the underside of a fertile lobe seen from beneath :—magnified.

Douglasianae.

N. O. 'Filices.

DCCCCXLV.

POLYPODIUM (Ctenopteris) PELLUCIDUM, *Kaulf.*

Vat. bipinnatifidum.

We have noticed this plant under the description of the preceding Plate (*Tab. 944*), and have there inserted the synonym of our *Pol. myriocarpum*. *Tab. 84* of the present work. Our representation is now accompanied by a magnified figure, showing the character of the venation* and proving that it is of the same nature as that of the true *P. pellucidum*. It is, indeed, an exceedingly variable species: but our Herbarium exhibits specimens which clearly show the passage from "one to the other, and the present can only be considered a bipinnatifid variety of that species." All the varieties appear to be found in the same island, Oahu—but they are probably by no means confined to it.

Fig. 1. Fertile lobe, seen from beneath, showing the venation and the clavate tips to the veinlets:—*magnified.*

Lobbiana.

N. O. Filices.

DCCCCXLVL

POLYPODIUM PAPILLOSUM, Bl.

Elatum glaberrimum, frondibus membranaceis elongato-oblongis acuminatis profunde fere ad rachin pectinato-pinnatifidis, lobis lineari-oblongis obtusis horizontaliter patentibus apicibus solummodo serratis, venis unifurcatis, soris uniserialibus inter costam et marginem profunde immersk* stipite rachique valida nitidissimis fuscis, caudice elongate repente nudo.

Polypodium papillosum, Bl. En. Fil. Jav.p. 131. J. Sm. En. FiL Philip, in Hook. Journ. of Bot. p, 394. Br. in Horsf. Plant. Jav. Bar. t. 2.

HAB. Mountain woods, on trees; Java, *Blume, Horsfield, Thos. Lobb.* South Camarines, Philippine Isles, *Cuming*, n. 185.

Blume compares this remarkable species with *P. vulgare*: but in form and texture it is widely different. Its sori, toq, are sunk into a deep poin^or bag, forming so many prominent papillae on the upper side of the frond, the mouth of which is contracted. Blume notices a var. "*B. frondibus brevioribus et laciniis acutiusculis.*"

Fig. 1. Portions of a fertile lobe, seen from beneath, f. 2. lesser portion of the same seen from above, f. 3. Soriferous sack, cut through vertically :—magnified.

Jamesonianiae.

N. O. Filices.

DCCCCXLVH.

POLYPODIUM FARINOSUM, *Hook.*

Parvum submembraneum fronde elliptico-lanceolata pinnata
utrinque pulveraceo-farinosa albida, pinnis approximatis
lineari-oblongis patentibus sessilibus subdecurrentibus ob-
tusis sinuato-crenatis basi sursum obtuse auriculatis, venis
unifurcatis apice clavatis, venula superiore apice sorifera,
sonis medio intra costam et marginem uniseriatis magnis
aureis, stipite gracili brevi nudo.

HAB. On the trunk of an old tree at the eastern ascent of
the Cordillera of Quito, where the forests commence; rare,
and very few specimens were gathered, *Prof. W. Jameson.*

Independent of the white pulverulent or subfarinaceous
substance, which invests both sides of the half dozen specimens
in my possession of this rare Fern, and which has the
appear^nce of being quite natural, not adventitious, there is a
something in the general outline or shape, and in the form of
the pinse, which, taken in conjunction with the simply furcate
veins and the large orange-coloured sori, renders this Fern
unlike any species with which I am acquainted. The pinnae
are very compact, or closely placed; the base of each is very
unequal, the upper base broad and dilated into a lobe or
obtuse auricle, while the lower is narrow, yet decurrent; so as,
In the upper half at least, to form rather a frond pinnatifid to
the very base than a truly and entirely pinnated one- I
have seen no specimen, nor any species like the present from
any Botanist save Professor Jameson.

Fig. 1. Three pinna, upp'er surface, f* 2. Fertile pinna,
seen from beneath:—magnified.*

Jamesoniana.

N. O. Filices.

DCCCCXLVHL

POLYPODIUM SEMIADNATUM, *Hook.*

Frondibus linearibus acuminatis longissimis petiolatis pinnatis, pinnis approximatis rigido-membranaceis suboblique ovatis obtusis crenato-lobatis basi superiore subauriculatis longe ciliatis basi costae adnatis, subtus petiolo gracili epaleaceo rachiqucfiliformibus nigris patenti-hirsutis, venulis furcatis, venula superiore apice sorifera.

HAB. On trunks of trees, Pilzham, 12,000 of elevation on the Quitinian Andes, *Prof. Jameson.*

When Ferns are pinnated there is commonly a contraction or constriction as far as the midrib, which even in a sessile pinna is usually the only point of attachment to the rachis, unless in the cases where the pinna* are decurrent. Here, on the other hand, the pinnas are not decurrent, yet is there a considerable portion of the base of the pinnae both above and below the midrib *adnate* with the rachis: in no portion of the very elongated frond are the pinnas confluent or decurrent. Our specimens are afoot, and a foot and a half long, the stipes is from two to four inches long.

Fig. 1. Pinnae, seen from above, /. 2. Fertile pinnae seen from beneath:—magnified.

DCCCCXLIX.

POLYPODIUM (Ctenopteris) KHASYANUM, *Hook.*

Pendulum, frondibus caespitosis subcoriaceo-membranaceis elongatis lineari-lanceolatis acuminatis basi attenuatis profunde pinnatifidis, lobis oblongis acutiusculis subsinuato-crenatis ciliatis rachi paginaque subtus hispido-hirsutis, venis simplicibus apice marginem versus soriferig, soris sub-elliptico-rotundatis immersis, stipite brevissimo subnullo, radice fibroso-caespitosa.

HAB. On trees, Khasya, *Drs. Hooker and Thomson.*

Fronds a foot to a foot and a half or more long, nearly sessile, for the leafy lobed portion extends nearly to the rGot/ The breadth in the widest part is two inches, tapering towards the base and towards the point. The margin and the surface and rachis beneath (the latter sometimes above) are hairy, and ciliated. The sori, occupying the apex of a single and simple vein, is always sunk in a depression, which depression occasions a roundish swelling or tubercle on the opposite side, only, however, slightly prominent, and very unlike the sack or bag represented in our *Polypodium papillosum*, Tab. 946.

Fig. 1. Fertile segment of a frond, seen from above. /. 2. the same, seen from beneath:—magnified

DCCCCLIL

POLYPODIUM (Marginaria) LACHNOPUS, *Wall*

Frondibus glaberrimis membranaceis firmis oblongo-lanceolatis acuminatis profunde fere ad costam pectinatim pinnatifidis, lobis linearibus acutiusculis obtuse serratis ad costam subtus minute nigro-squamulosis, squamulis cordatis ciliatis longe cuspidatis nigris, soris uniserialibus, venulis prope costam areolas fibrmantibus reliquis liberis furcatis, stipite brevi nudō caudice longe repente nigro-paleaceo.

Polypodium lachnopus, Wall. Cat n. 310.

HAB. Nepal and Kamaon, Wattich. Khasya, Drs. Hooker and Thomson.

Mr* J. Smith brings this into the genus *Goniophlebium*, Presl, and this he unites with *Marginaria*, Bory and Presl.: but the venation of PresFs *Goniophlebium* is very different from that of *Marginalia*, which is what is represented at the base of the lobe or segment of our plant (see fig. 1.) while all the rest of the lobe has the venation of true *Polypodium*[^] with free veins clubbed at the apex. Our plant is thus intermediate between the two genera now mentioned. The minute scales at and near the main costa present a beautiful appearance under the microscope: they are nearly black, almost orbicular, finely reticulated and have a long cuspidate or setiform point, (f. 2.) Under a lens too the upper side of the costa will be found pubescent.

Fig. 1. Fertile lobe, seen from beneath, and showing the venation. /. 2. Scale from the rachis beneath:—magnified.

DCCCCLni.

POLYPODIUM (*Drynaria*) ROSTRATUM, *Hook.*

Frondibus elliptico-lanceolatis rostrato-acuminatis basi attenuatis chartaceis opacis integerrimis glabris, soris magnis costam versus subuniseriatim dispositis, venis omnibus uniformibus reticulatis, areolis hexagonis oblongis, venuulis liberis simplicibus furcatis, stipite gracili fronde breviore, caudice gracili elongate filiformi.

HAB. Khasya, Drs. Hooker and Thomson.

A *Polypodium* of the *Drynaria* or *Phymatodes* group, which does not appear to be known hitherto to Botanists. Unlike that at our Tab. 951, it has no costulae, in other words the costa is not pinnated with strong strait parallel primary veins: the whole surface is reticulated, with oblong hexagons lying in an oblique direction, united by an intramarginal vein, and within each areole is generally one simple or forked, ultimate, free veinlet, the branches are often much divaricated. The frond is of a parchmenty or chartaceous texture, between coriaceous and membranaceous, opaque. The apex runs out into a long, narrow beak, whence the specific name.

Fig. 1. Sterile portion of a frond. */. 2.* Fertile portion seen from beneath:—*magnified.*

DCCCCLIV.

POLYPODIUM (Drynaria) RHYNCHOPHYLLUM, *Hook.*

Frondibus chartaceis glaberrimis nitidis sterilibus oblongo-ovatis obtusis, fertilibus lanceolatis longe acuminatis costulatis costis elevatis, areolis transversim oblongo-tetragonis, acumine sorifero, sterili venatione simpliciore appendiculis subnullis, stipite fronde breviore nudo, caudice elongato repente gracili ramoso copiose paleaceo.

HAB. Assam, Mrs. Mack, Griffith. Khasya, Drs. Hooker and Thomson.

Closely allied in general appearance to *Polypodium cuspidiflorum** of Reinwardt, of which we have authentic specimens in our possession from Java, and from Luzon, gathered by Cuming (n. 109,) This latter is, however, a much smaller plant, very opaque, with no visible costulae; the fronds have a broader base, more suddenly tapering into a beak, almost quite sessile, upon an exceedingly slender, filiform caudex, destitute of scales, but throwing out copious woolly fibres. The venation of our present species* is seen with difficulty, being quite sunk in the very opaque frond, but from what I do see of it, I should judge I am correct in referring it to *Drynaria*, Presl. M. F6e, however, places it in the genus *Craspedaria*, Link, which corresponds with the first section of Presl's *Marginaria*, though I suspect the venation is very different

Fig. 1. Sterile portion of a frond, showing the venation.
/. 2. Fertile portion of a frond, showing its peculiar venation:
—magnified.

* A species nowhere described as far as I can find.

DCCCCLV.

CYSTOPTERIS DOUGLASII, *Hook.*

Parva, frondibus membranaceis finnis glabris oblongo-lanceolatis pinnatis, pinnis inferioribus latioribus subovatis acuminatis acutiusculis pinnatifidis, lobis lato-oblongis ovatisve obtusis obtusedentatis, superioribus oblongo-lanceolatis obtusis lobato-pinnatifidis decurrentibus coadunatisque, soris majusculis, involucris reticulatis suborbicularibus dentatis cito capsularum pressione reflexis, stipite brevi.

*Cystopteris Douglasii, Hook. Gen. Fd. 1, p. 200,
HAI>. Sandwich Islands, D. Douglas.*

Stipes short, naked. Frond firm, rather rigid, of a dark, lurid green colour in the dried state. The form of the pinnae and segments approaches that of the N. American *C. bulbifera*, but they are broader and much less compound. The fructification, though much advanced in our specimens, is clearly that of the present genus.

Fig. 1. Upper pinnae, seen from above. /. 2. lower pinnae, fertile, seen from beneath. /. 3. Sorus:—magnified.

DCCCCLVL

HYMENOPHYLLUM ASPLETFLOIDES, SIV.

Frflndibus oblongis obtusis pendulis (siccate fusco-brunneis) profunde pinnatifidis, lobis ovatis rarius integris plerumque bi-tri-quadrifidis nunc subpalmatis, lobulis obtusis apice soriferis, involucris solitariis (singulo lobulo) fere orbicularibus liberis integerrimis basi cuneata solummodo immersa profunde bivalvibus, receptaculo brevi • onmino incluso, stipite tenui fronde breviore.

Hymenophyllum asplenioides, Sw. *Syn. Fil* p. 145. *Hook. Gen. e% Spec. Fil.* 1.p. 87.

H, *palmatum*, *Klotzsch*[^] *in Herb. Reg. Berol. et in Herb. Hook. HAB.* On the trunks of trees, Jamaica, {*Herb Nostr.*) Brazil, Sellow.

None of the great Fern Family more needs illustration by good figures than the species of *Hymenophyllum* and *Trichomanes*. One cannot say much in praise of the representations of this plant by Lamarck and Hedwig, quoted in the *Ænera et Species Filicum*. It is, indeed, a distinct and well marked plant, and yet, Dr. Klotzsch, did not recognise his *Hymenophyllum palmatum* to be identical with it. When the primary segments of the fronds bear three or four lobes with the sori upon them, they are necessarily broader and sub-palmate:—so that one cannot really consider such a character worthy of being recorded as constituting even a variety. The fewer or greater number of lobules may be seen on one and the same plant. The caudex is long, creeping, filiform. The texture of the frond is firm, composed of closely placed minute areoles. The venation consists of a central costa, and veins to each primary segment, sending off free branches to the apex of every lobe.

Fig. 1. A primary segment or lobe, with fructifications:—*magnified*, *f. 2.* A lobule with sorus, one of the valves of the involucrum being removed:—*more magnified*.

DCCCCLVII.

LINDSiEA MEDIA, *Br.*

Subelata glaberrima, frondibus subcoriaceis ovatolanceolatis acuminatis bi-tripinnatis, pinnulis oblique rhombeis obovato-cuneatisve inferioribus sublobatis reliquis integris, sterilibus serratis, venis flabellatis ramis saepe anastomosantibus, sonis dimidio superiore continuis, stipite elongato tetragono, rachibus tenui-alatis.

Lindsaea media, *Brown*, *Prodr. Fl. Nov. Holl* p| 156. *Hook. Gen. et Sp. Fil.* 1.p . 212.

HAB. Tropical shores of New Holland, *Brown*. East Coast of Tropical New Holland, *All. Cunningham*. Fitzroy Islands, *J. MacGillivray, Esq., {Voy. of the Rattlesnake, n. 256. ft.)*

A very rare and little known *Lindsaa*, belonging to the same group or division as *L. fidbellulata*, and *L. tenera* of Dryander, but very distinct as a species. Mr. MacGillivray has put us in possession of fine specimens, which have enabled us to give the accompanying figure. The primary pinnae are all very much acuminate, the ultimate ones extremely narrow, minute, and coadunate into a lobed point. The rachises are margined with a narrow wing. Of the fertile pinnules some have the flabellate dichotomous veins free, others anastomosing; in the sterile ones I believe always free, with the apices of the veins clavate. Some authors who feel the propriety of retaining the genus *Isoloma*, would perhaps place it there.

Fig. 1. Sterile pinnae with its free veins. /. 2. Fertile pinnae, with anastomosing veins:—magnified.

DCCCLVIII

LINDSJBA CAUDATA, Hook.

Stipite tereti rachique purpureo-ebeneis nitidis, fronde ampla bipinnata, pinnis angustis numerosis (11-17) lineari-oblongis apice longe caudato-acuminatis, pinnulis semiovalibus lunulato-falcatis decurvis obtusis membranaceis approxigiatis, basi superiore truncatis, margine superiore fere eemicircularibus integerrimis, terminalibus sensim minoribus demum in acumen confluentibus, soris omnino marginalibus ad apicem continuis.

Lindssea caudata, Hook. Gen. et 8p. Fil. l.p. 215.

HAB. Adam's Peak, Ceylon, Mrs* Genl Walker.

Nearly allied to *L. trapeziformis*, Dryander, which species is found in the East as well as the West Indies: but the present plant has more numerous pinnae on the frond, tapering to a tail-like point, blunted pinnules, exactly marginal sori; terete and darker coloured stipes;—and these marks are found in four fine specimens received from Ceylon at different periods. This dries to almost a black colour;—*L. trapeziformis*, retains its bright green in the Herbarium.

Fig. 1. 2. Fertile pinnules. /. 3. Portion of a fertile pinnule, showing the position of t1[^] involucre at the very edge of the pinnule:—magnified.

Gunniance.

N. O. Filices.

DCCCCLIX.

CYSTOPTEKIS TASMANICA, *Hook.*

Parva gracilis, stipite rachique capillaribus, frondibus oblongis pinnatis, pinnis late ovatis obtusis inciso-lobatis superioribus decurrentibus inferioribus subpetiolatis pinnatifidis, lobis ovatis obtusis integris vel subdentatis, soris paucis minutis, involucris ovatis acuminatis, rachi superne alata,

Gystopteris Tasmanica, *Hook. Gen. et Sp. Fil.* 1. p. 199.

HAB. Van Dieman's Land, R. *Gunn, Esq.*

As I have already observed elsewhere, I was at one time disposed to refer this to some of the states of the var. *dentata* of • *C. fragiliSy* but the fact of Mr. J. Smith having received from the same country, though from a different source, an exactly similar plant, together with the delicate habit, large (comparatively) and broad, sparingly divided pinnae and small fructifications, induce me to keep it distinct. Caudex slightly creeping in one of our specimens. Whole plant including the stipes, 4-6 inches high.

Fig. 1. Fertile pinna, seen from beneath. /• 2< Sorus and involucre:—magnified.

DCCCCLX.

LOMARIA NIGRA, *Col.*

Parvula luride viridis, frondibus sterilibus linearibus sublyrato-pinnatifidis, lobis (inferioribus nunc discretis) oblongis obtusis sinuato-crenatis glaberrimis infimis marginibus costis rariusque paginis inferioribus tomentosis, terminali maxima basi lobata, fertilibus pinnatis, pinnis paucis distantibus suberectis anguste linearibus apice" subulatis acuminatis v. apiculatis, terminali elongata* rachi parce stipitibusque sparse paleaceis.

Lomaria nigra, *Colenso in Tasm. JPhil. Journ.*

IIAB. New Zealand, dense forests between Tauranga and Roturua,*East Coast: *Rev. JV. Colenso*, n. 299. 331*

This is a very anomalous plant, and has rather an unnatural appearance, the fronds being blackish or lurid, brittle when dry, the pinnules often erose, sinuate and irregular in size and outline; as if it had grown in an ungenial locality. Mr. Colenso's specimens are good ones, however, and uniformly display the peculiar characters which separate it from its allies. In habit, general appearance, paleaceous rachis, and other points, it clearly resembles *L. fluviatilis*, Br* Prod. (*Li. rotundifolia*, *Raoul.*) a denizen of similar localities; in its typical state that plant has pinnate barren fronds, with the lower pinnules stipitate, (a tendency to which may be seen in the lower pinnae of our plate of *L. nigra*)> the terminal one does not run out into a long lobe, and the fertile pinnules are shorter and blunter. To the infinitely variable *JL lanceolata* again (see Plate t 429) the *L. nigra* is allied by the pinnatifid fronds, and narrow fertile pinnae with subulate tips. The pubescence consists, in the dried state, of a short rufous tomentum, sometimes spread over the under surface of the pinnae, at others confined to the margins and costae, and sometimes it is wholly absent.—**J.D.H.**

Fig. 1. Pubescent pinna, f. 2. Fertile pinna. J\ 3. Transverse section of the same, with the involucres reflexed showing the sori:—all magnified.

DCCCCLX1.

ADIANTUM GLAUCOPHYLLUM, *Hook,*

Elatum, frondibus ovatis sujpradecompositim pinnatis, pinnis eubchartaceo-membranaceis glabris utrinque glaucis longiuscule petiolulatis oblique cuneatis margine Superiore rotundato 2-5 lobo, lobis obtusis emarginatis v, bifidis, lobulis fructiferis subincurvis, sinubus soriferis, soris mediocribus, involucris orbiculari-reniformibus demum subcoriaceis, stipi te rachique gracili ebeneis nitidis.

Adiantum glaucophyllum, *Hook. Gen. et Sp. Fil. 2. p. 40.*

A. cuneatum, var. *angustifolium*, *Mart, et Galeot. Fil. Mex. p. 70.*

HAB. Cordillera of Mexico, inhabiting the cold region, at an elevation of 6,000 to 10,500 feet above the level of the sea, *Galeotti n. 6,266. and 6,359.* (the latter with narrower, more rigid, and almost coriaceous fronds) and *n. 6566;* Pic Orizaba, 9750 feet; *Linden, n. 48. Jurgensen, w, 322. Mr. JParki7ison, « Teapisca (Chiapas),*" *Linden, n. 1550. Baqueti, i Vvsraguas, Central America, Seemann.*

M.M.Martens and Galeotti referred their *Adianta* from the stations above alluded to, in Mexico, to the S. Brazilian *A. cuneatum* of Langsdorff and Fischer, as narrow pinnuled varieties; but I have other specimens from Linden and Seemann, and all agree in the characters above given. All accord, too, in being of a larger size and more compound than that species, with usually narrower, more rigid, always more glaucous pmnules, together with a less deep and narrow notch for the reception of the sori; and not only do they thus agree in the respective characters, but I have not seen anything to correspond with the present, except from Mexico and the adjacent Isthmus; certainly no such appearances are exhibited by' any true Brazilian *A. cuneatum*. It is of the *Capillus-Veneris* group, and is I think more nearly allied to the East Indian *A. venustum*, Don, than to any other.

Fig. 1. Sterile pinnule. /. 2. fertile pinnule :—magnified.

Colensoance.

N. O. Filices.

DCCCCLXIL

HYMENOPHYLLUM FLEXUOSUM, *AIL Cunn.*

Frondibus erectis subrigidis glaberrimis deltoideo-ovatis tri-quadripinnatifidis, segmentis lincaribus obtusis undulatis integerrimis ultimis plerumque furcatis, involucris terminilibus segmento latioribus orbicularibus integerrimis liberis (omnino exsertis) ad basin bivalvibus, valvis convexis, receptaculis iticlusis, stipite rachique alata crispata marginatis.

Hymenophyllum flexuosum, *All. Cunn. FL Nov. Zel. in Hook**
Comp. to Bot. Mag. 2. p. 369. Hook. Gen. et Sp. FiL
\p. 105.

HAB. New Zealand, Northern Island, *AIL Cunningham,*
Colenso, Dr^{} Logan, J. D. Hooker.*

Apparently a common species, on the trunks of trees, in woods, in the northern island of Neyr Zealand. It bears the orbicular involucres of *H. Javanicum*, but they are not confined to short lateral segments as in that[#]species:—nor is the shape of the involucres like that of *H. crispatum*. The fronds are more deeply and copiously divided than either, more crised, especially in the wings of the rachis and stipes, and the general habit is different, though difficult to be described in words. It seems to hold a middle rank between the two East Indian species just jnentioned.

Fig. 1. Portion of a fertile plant. /. 2. Sorus:—magnified.

DCCCCLXIII.

HYMENOPHYLLUM FUCOIDES, SIV.

ITrondibus oblongis acumirfatis inferne pinnatis, pinnis ovato-oblongis bipinnatifidis superne bitripinnatifidis segmentis linear-i-oblongis obtusis acutissime serratis, costis non raro spinosis, soris in ramos breves lateralibus, involucris ovatis lato-obovatisve magis minusve spinuloso-dentatis infra medium bilabiatis, receptaculis inclusis.

*Hymenophyllum fucoides** Sw. *FL Ind. Occ.* p. 1747. *Syn. Fa.* p. 146. Willd. *Sp. PL* 5. p. 523. Hook. *Gen. et Sp. Fil.* I. p. 101.

Trichomanes fucoides. Hedw. *Fil. cum Ic.*

HAB. Jamaica, Swartz, Purdie. Martinique, Sieber. Summit of the Organ mountains, Brazil, Gardner, ?i. 5951. Peru, Poeppig. Forests of the Columbian Andes, Jameson. Mexico, (Kunze). Caraccas, Linden, n. 57.

Some of our specimens are nearly twice the size of that here represented; especially those from Prof. W. Jameson: and it is certain that the form of the involucre is very variable, sometimes ovate and nearly entire, sometimes, and, more frequently, broadly obovate and more or less deeply spinuloso-serrai^e. The rachis is not winged below by the decurrent side branches, and hence the frond is there pinnated; while above, the rachis is so winged as to constitute a bi- or tripinnatifid frond. The stipes is wingless too, and often frougely setose, with short harsh hairs. The root is a creeping, slender, filiform, downy caudex, and runs extensively over rocks and trees.

Fig. 1. Pinna or primary lobe, with an ovate in^Alucro:—*magnified*, f. 2. Portion of a fertile frond with a broader obovate involucre:—*less magnified*. .f. 3. Lesser portion of a frond, with broadly obovate involucre, and spines on the rachis:—*magnified*.

Menziesiana.

N. O. Filices.

DCCCCLXIV. .

HYMENOPHYLLUM UNDULATUM, 8W.

Humile pendens, frondibus oblongis ovatisve tri-quadripinnatifidis inferne nunc pinnatis, lobis primariis patentibus, ultimis brevibus subpatentibus oblongis integerrimis obtusis vel emarginatis omnibus undulato-crispatis, rachibus alatis undulatis valde crispatis, involucris rarissimis in lobos breves terminalibus liberis orbiculari-ovatis usque ad basin bivalvibus, valvis convexis integerrimis, stipitibus gracilibus nudis.

Hymenophyllum undulatum, Sw. *Fl. Ind. Occ.* p. 1751* *Syn.*
Fil p. 148. *mild.* *Sp. PL* 5. p. 533. *Hedw.* *Fil cum Ic.*
Hook. Gen. et Sp. Fil. I. p. 105. .

HAB. Jamaica, *Swartz*, *Menzies*. Mountains of Andinarca,
Peru, *Mathews*.

A very distinct species. The stipes varies from one to three inches long. Fronds three four or five inches long, ii^ every part beautifully and regularly undulato-crispate.

Fig. i^ Portion of a fertile frond:—*magnified.* /• 2. Involucrum from which one valve is Removed: —*magnified.*

DCCCCLXV.

ADIANTUM FRAGILE, SW.

Frondibus caespitosis ovato-lanceolatis tri-quadripinnatis, pin-nulis brevi-petiolulatis (petiolulis gracillimis) tenui-membra-naceis obovato-cuneatis basi in petiolulum articulatis apice rotundatis, sterilibus serratis, fertilibus 3-4-lobatis omnibus cito caducis, lobis fertilibus retusis sinu soriferis, involucris oblongis rectis, stipite brevissimo vix ullo rachique nigro-ebeneis nitidis, radicibus cespitosis fibris rigidis lanuginosis.

Adiantum fragile, Sw. *FL Ind. Occ.* 3. p. 1721. *Syn. fit p. 125.*

A. cuneatum, Kunze. *PI. Exsicc. Poepp. in Herb. Nostr.; in IAnncea, 9. p. 82?*

HAB. Calcareous Rocks, Jamaica, Swartz, Dr. Wright, Otto, Wilson, Purdie. Cuba, Poeppig*

I_m have received West Indian specimens from different persons of this singular and well marked plant, all of them exhibiting the same peculiarity of shedding almost every leaflet or pinnule, a circumstance due to the presence of an articulation where the leaf is inserted upon the petiole: so that the specimens, when they reach me, are mere skeletons, consisting of wiry stipites, with the exceedingly slender and wiry rachis, very much branched, and the petiolules from which the pinnules had fallen, the latter all lying apart from the plant. There are ample characters for recognising this species.] know no *Adiantum* with so peculiar a-growth, so tufted, so very short in the stipes, and with such caducous pinnules. In other respects the latter a good deal resemble *A^c cuneatum* of Brazil, and *A. venustum* of Northern India.

Fig. 1. Sterile pinnae, f. 2. Fertile pinnae:—magnified.

DCCCCLXVI.

LYCOPODIUM SCARIOSUM, *Forst.*;*var. decurrens.*

Caule repente prostrato vase dichotome subflabellatum ramoso, (amis complanatis, foliis majoribus bifariis subimbricatis oblongo-falcatis acutis decurrentibus minoribus stipulae-formibus cauli appressis subulatis acutis obtusisvc omnibus apice coriaceis rarius scariosis, spicis terminalibus sessilibus v. breve pedunculatis, squamis subsexfariam imbricatis rhombeis rhombeo-ovatisve acutis obtusisve v. apice lamina scariosa appendiculatis marginibus plerumque scariosis.

L. scariosum, var. decurrens. Hook. Jih Flor. Nov. Zeland. ined.

L. decurrens, Br. Prod. p. 165.*

HAB. Tasmania; summit of the Western mountains, JR. W. Lawrence, Esq., Mount Wellington, R. C. Gufln, Esq., J. & J. Ifooker.

The *Lycopodium decurrens* of Mr. Brown was first referred to *L. scariosum*, Forst. by M. Spring, (Monogr, Lycopod. Pars 1. p. 108.); and after a careful examination of very many specimens with a large suite of those of Forster's plant from various parts of New Zealand and others from Lord Auckland's group, we have no hesitation in adopting that opinion. The length of the peduncle, and the presence or absence of the sgarigse white tips to the larger and stipulary leaves, are most variable characters in the New Zealand plant; and though these tips are generally absent in the Tasmaniaii plants, some of the leaves show them. The form of the scales of the spike, the breadth of their white margin, and the length and form oT their white tips are no less variable; the latter arc more or less recurved. The large common state of *L. scariosum* has not hitherto been found in Tasmania: the climate of the better explored districts would appear to be too dry for it. In New Zealand it is common in alpine woods of the Northern Island, and throughout the Middle and Southern, and it is also found in Chili, among the Andes of South America, and in the lofty mountains of Jamaica; it is the *L. Jussieui*, of Oesvaux and of this work, t. 185. X. *Haenpii*, Presl, and *L. Lessonianum* of A. liich. (Flor. Nov. Zeland.)—J.D.H*

*Fig. 1. Top of branch and spike, f. 2. and 3. Scales.
f. 4. Back view of branch with larger and*stipulary leaves:
—all magnified.*

Hookeriana.

N. O. Filices.

DCCCCLXVII.

GRAMMITIS ASCENSIONIS, *Hook.*

Casspítosa, humilis, glabra, frondibus triangulari-ovatis membranaceis pinnatis, pinnis profunde pinnatifidis inferioribus bipinnatifidis, laciniis oblongis obtusissimis integris y. bifidis apicibus soriferis, rachi compressa subulata, stipitibus fronde longioribus nudis, radice fibrosa.

HAB. Bocks and Banks on the Green Mountain, Ascension Island, elev. 1,200 to 1,800 feet. Dr. J. D. Hooker, Dr. Curror.

In size and texture this pretty *Grammitis* resembles the *G. leptophylla* of Swartz, but the pinnae and pinnules have nothing of the broad and cuneate form of that species. In ramification it more nearly approaches the *Gymnogramme chcerophylla*, but that species is much larger, and the ultimate segments are longer and narrower and almost acute, and the fructifications are more copious and continuous. The present plant is, as far as I know, wholly confined to the Island of Ascension.

Fig. 1. Fertile pinna seen from beneath. /. 2. Sorus:—magnified.

DCCCCLXVIII.

LYCOPODIUM (§Complanata) CASTJARINOIDES, Spring*

Longissime scandens flexuosum, caulis teretibus nudis vel subulato-squamosis, ramis copiosissimis dichotome partitis pendentibus rubris rubro-fuscisve comosis anguste linearibus compressis, foliis stipulisque conformibus subulatis sublonge vaginatis apice longe diaphaois albis raro serrulatis, vaginis cum ramo decurrentibus coadunatisque, fertilibus in eadem stirpe dichotome paniculatis omnibus apice spicigeris, spicis sursum curvatis cylindraceis uncialibus, bracteis cordatis denticulatis acumine subulato diaphano.

1. Ramis longissimis nunc pedalibus rectis rubris, foliis appressis brevibus apicibus piliformibus (fig. nostr. 1. 2. 3.) *Lycopodium casuarinoides*, *Spring*, *Monogr. Lycop. JP.* 1. p. 94. *JP. II.* p. 45. *L. rubellum*, *Presl*, *Sot. Bern.* p. 153. *L. comans*, *Hook. Jil. Fl. Antarct.* 2. p. 112. (*in note*). *L. leucolepis*, *Junghuhn et de Vrzes*, *in litt.* (*Spring*.)
 2. Ramis longissimis rectis rubris, foliis appressis late subulatis totis fere diaphanis scariosis serratis (fig. nostr. Q. 7.)
 3. Ramis brevioribus subflexuosis rufo-fulvis, foliis ut in 1. *Lycopodium filicaule*, *Hook. Jil. Fl. Antarct.* 2. p. 102. (*in note*.)
 4. Ramis longissimis flexuosis rufo-fulvis ultimis laxe foliosis, foliis patentibus apice piliformibus subulatis (fig. nostr. 4. 5.)
- HAB. Var. 1. Malacca (probably on Mount Ophir), *Cuming*, n. 2346. Var. 2. Mount Ophir, *TJws. Lobb*, *Sir Wm. Norris*. Var. 3. Moflong, Khasya hills, in Fir forests. *Griffith*. Var. 4. Malacca, *Griffiths*.—*Spring* gives Sumatra, (*Junghuhn*) *M%*, another locality for this species, on the authority of *Dr. de Vries*.

It is hard to say which of the above published names for this remarkable species should have the preference: three of them are given by different Botanists to the same plant of Cuming (his n. 2346.) *Spring's* work bears the date of 1842-1849, *PresPs* 1844, and *Dr. Hooker's* 1845. *Spring's* is unquestionably the most expressive, particularly for that form which Mr. Cuming detected, which almost exactly resembles the pendent branches of some slender *Casuarina*. But like too many species of the genus it takes other forms, and our n. 2. has the leaves of the ultimate or younger branches very much spreading, giving quite a different character to the plant. *Dr. Hooker's* *L. filicaule* is a pale coloured state, with shorter and flexuous branches.

Fig. 1. Small portion of a plant, var. 1. *not. size.* f. 2. Portion of a branch. /. 3. Capsule and scale:—*magnified*. f. 4. Small branch of var. 2.:—*nat size.* f. 5. Leaves of ditto:—*magnified*, f. 6. Small branch of var. 4.:—*nat size.* f. 7. Leaves:—*magnified*.

DCCCCLXIX.

LOMARIA VULCANICA, *Blume.*

Rhizomate crasso paleis setaceis crinito, frondibus sterilibus sublonge stipitatis coriaceis lanceolatis ovatis deltoidcisve pinnatifidis basi pinnatis, pinnulis approximatis (infimis deflexis) oblongo-ensiformibus subfalcatis acuminatis obtusive integerrimis v. obscure crenulatis glaberrimis v. costa nervisque pilosiusculis subtus pallidioribus, pinnis fertilibus lanceolatis longe acuminatis pinnatis, pinnulis patentibus linear-i-elongatis obtusis acutisve basi dilatatis, involucris laceris, stipite basi crinito.

L. vulcanica, *Blume*, *Enum. Plant. Jav. fasc. 2. p. 202.*

L. deltoides, et *L. deflexa*, *Colenso in Tasm. Phil. Journ.*

H A B. Tasmania, in mountainous districts, *M. C. Gumi, Esq.*, n. 28 and 1522. New Zealand; Auckland, *Sinclair*; Tarawera, *Rev. W. Colenso*. Mt, Gedei, Java, *Blume*.

A very distinct species, common to New Zealand, Tasmania, and the lofty mountains of Java. It is generally readily recognised by the falcate deflexed lower pinnules and deltoid glabrous pale coriaceous frond, with patent linear pinnules; but the frond is sometimes lanceolate, more or less pilose on the under surface, and the lower pinnules are sometimes not at all deflexed. In some of Mr. Colenso's specimens, gathered on dry rocks, the fronds are very small, coriaceous and rigid, and the rhizome an inch in diameter, covered with rigid black shining setaceous palese, and with pale scars where old fronds have fallen.—J.D.H.

I?ig. 1. Under surface or portion of a sterile frond, f. 2. Fertile pinnule, both from a hairy variety :—*magnified*.

Seemanniance.

N. O. Filices.

DCCCCLXX/

BLECHXUM LANCEOLA, SW.

Parvum, caudice perbrevi fibroso non raro stolonifero, frondibus simplicibus linear-lanceolatis anguste acuminatis minute ciliato-scabris nunc basi utrinque auriculatis vel pinnis 2-4 ovatis parvis pinnatis, stipite frondem fertilem subaequante gracillimo sparsim paleaceo. •

Blechnum Lspceola, Sw. *K. Vetenskaps Academ. Handl.* 1/1.p. 72. t S.f. 2. *Hook. Bot. Mag.* t. 3240. Kunze, *Flip.* 126. *. 57./. 1.

Blechnum lanceolatum, Raddi, *Fih Bras.* p. 52. t. 69./. 3.

Blechnum trifoliatum. Kaulf. *Enum. Fl* p. 157.

HAB. Brazil, about Rio, Raddi, Sellow, Gardner, (n. 50.^) J. D. Hooker; Boqueta, Veraguas, Seemann, n. 1556.

Notwithstanding the figures above mentioned, this plant has not been satisfactorily and fully represented, because authors have been unacquainted with the pinnated state of it, and even Kunze only knew the divided frond as arising from cultivation, and only as laciniated and auriculated. Mr. Seemann's specimens, however, are mostly pinnated, as represented in our left-hand figure. It is an extremely pretty species, the smallest of the Genus, and was supposed to be peculiar to Brazil till Mr. Seemann detected it in Veraguas, Isthmus of Panama, in 1849.

Fig. 1. Portion of a stejile frond or pinna. f_m 2. Portion of a fertile do:—magnified.

DCCCCLXXI-DCCCLXXII.

GYMNOPTERIS SEMIPINNATIFIDA, Fée.

Fronde subtriangulari-ovata pinnatifida basi nunc pinnata, lobis pinnisve oblongis obtusis vel acuminatis superioribus in lobum acuminatum magnum sinuatum confluentibus integerrimis v. subdentatis, fertilibus conformibus sed contractis, stipite (fertilis praecipue) longissimo decidue paleaceo.

Gymnopteris semipinnatifida, Féé, Acrost. p. 84. t. 44.

(3. froflde basi vix pinnata, lobis infimis decurrentibus, TAB. NOSTR. 971, 972.

HAB. French Guiana, Leprieur. ft. Rocky rivulet in tBe Serra de Santo Gabriel, Amazon, Spruce, (n. 2121.)

" Creeping rhizoma, much entangled among stones, so that it was often impossible to extract barren and fertile fronds in conjunction."—The large-fronded species of *Gymnopteris* are extremely difficult to determine without the aid of good figures. We are consequently thankful for the figure, as well as its accompanying description, of *G. semipinnatifida* that M. Féé has given us in his valuable work on *Acrosticaceae*. That species, however, he considered remarkable "en ce qu'elle offre sur un même petiole une paire de pinnules très-courtement pétiolée, qui la fait pinnae, deux ou trois paires intermediaresj soudeés vers la base, qui la font pinnatifide; et un sommet settlement ondulé ou sinueux, qui permet de la regarder comme entifere." Now in all that concerns this form* our plant agrees; except that the two lowest lobes are not separated from the rest by a wingless rachis; and further, instead of any of the pinnules being shortly petiolate, they are • decurrent upon the stipes. Our plant has a near approach to, but not quite, a pinnated frond at the base. As such I can only consider it a*form of M. Fee's *G. semipinnatifida*.

Sinclairianæ.

N. O. Filices.

DCCCCLXXIIL

PTEBIS (§*Litobrochia*) ENDLICHERIANA, Ag.

Frondibus (nunc amplis) bi-tripinnatis rigido-membranaceis,
pinnis subsessilibus lato-lanceolatis superioribus decurren-
tibus, supremis coadunatis profunde pinnatifidis, laciniis
oblongis acutis subfalcatis apice serratis, sinibus acutis,
"venis basalibus (prope costam) monoarcuatis, & reis sub-
quartaae marginalibus, stipite elongato rachibusque
stramineis." Ag.

Pteris Endlicheriana, Agardh, *Recens. Gen. Pterid. p. 66.*
Hook. Gen. Fil Tab. 65. B.

HAB. Norfolk Island, All. Cunningham. New Zealand^
Northern Island, Dr. Sinclair. Van Diemen's Land, i?.
Gunn, Esq.

Our friend Agardh has well distinguished this handsome *Pteris*, (*Litobrochia* of Presl), and has founded its main specific character upon the peculiar venation: there is only a single arch between the costule, next the costa; and the fourth (or generally the third) areola from the costal one is that which reaches the margin. Agardh's description was made from Norfolk Island specimens in our Herbarium; but the plant has since been found in New Zealand and in Van Diemen's Land.

Fig. 1.-2. Portion of a fertile pinna:—magnified.

Griffithianae.

N. O. Filices.

DCCCCLXXIV.

GYMNOGRAMME AURITA, *Hook.*

Elata, fronde ovata oblonga acuminata rigido-membranacea pinnata, pinnis remotis sessilibus oppositis e lata basi oblongis acuminatis profundè pinnatifidis, lobis oblongis acuminatis patentibus subcurvatis crenato-lobatis paribus infimis subliberis duplo longioribus subpinnatifidis, venulis simplicibus, soris oblongis breviusculis, stipite elongate, rachibus hirtellis.

HAB. Khasya, *Griffith*, Drs. *Hooker and Tjiomson*, TJtos.
Lobb.

A variable species in its size and in the length and breadth of the pinnules. Fronds one and a half to two feet long, stipes nearly as much, and, as well as the main rachis, rich, glossy, mahogany brown. Barely I have seen the frond bipinnate, with the pinnae very much elongated, as if unnaturally drawn up among other plants. The remarkable feature of this species is the large size of the lowest pair of segments or pinnules of the pinnae, and their being so closely attached to the rachis that their base laps over it. The veins are free, as of the true *Gymnogramme* (*Lctogramme*, J Sm.) The sori are oblong, obtuse, rather short.

Fig. 1. Portion of a fertile pinna :—magnified.

DCCCCLXXV.

ACTINIOPTERIS RADIATA, *Link.*

Actiniopteris radiata, *Link*, *FiL Sp. Hort. JBerol.* p. 80.
Asplenium radiatum, *Sw.* *Syn. FiL* p. 75. *Sp. PL* 5. p. 308.
Acrostichum radiatum, *König*.
Acrostichum australe, *Vahl*, *Symb.* 1. p. 84. t. 25.
Acrostichum dichotomum, *Fb'rsk. FL* ^*ZEgypt.-Arab.* p. 184.
Acropteris radiata, *Fée*, *Gen. FiL* p. 76.
Blechnum radiatum, *JPresl, Tent. Pterid.* p. 102.
/3. frondibus magis elongatis, segmentis paucioribus vix radiatis, apicibus fere omnibus integerrimis subulatis. (See TAB. 976, for synonyms and remarks).
HAB. Arabia, *Forskal*. Upper Egypt and Cordofan, *Kotschy** South Africa, Macalisberg, *Burke* (in Herb. Nostr.) Madagascar, Bourbon, (*SwartZy*) *Carmichael* Scinde, *Dr** Stocks. Bombay, *Dr. Gibson*. Old walls, Madras, common, *Dr. Wight*, (w. 109.) *Mr. Gideon TJiomson*. Old w^wll, foot of the Limestone hills, near Segain, Northern India, *Mr. JEdgeworth*. Moradabad, *Dr. Thos. Thomson*. Old walls, Agra, and at Sikaan in Ava, *Dr. Wallich*, (*Cat. n₉* 137.)

One of the most curious of Ferns, with flabellate leaves like a minute Palm, and these leaves or fronds are often seen quite drooping and pressed down upon the stipes (more so than any of our figures represent them), as if occasioned by a joint, or probably by the effect of drought. The stipit& are densely tufted from a fibrous root and often clothed with rather large ferruginous scales at the very base, slender, wiry, two to five or six inches long, with small, patent, deciduous scales. Fronds (in the more common form of which we are now speaking) one to two inches long, and more than that broad, sometimes forming a half circle, many times dichotomous, firm, rigid, glabrous, of a glaucous or pale lurid colour, their ultimate segments linear, 2-3 toothed: the fertile fronds usually but not always longer, less spreading, with fewer but more deeply divided segments tapering to a sharp point and entire or nearly so. The involucre is narrow, linear, fixed to a marginal nerve (which bears the sorus and opening inwards). Nerves strong, dichotomous, following the divisions of the frond,

Fig. 1.-2. Plants, nat. size. f. 3. Segments of a fertile frond:—*magnified*.

DCCCCLXXVL

ACTINIOPTERIS RADIATA, *Link.* j3.

(For figure and description of the usual form of this plant, see our preceding plate, TAB. 975, and description).

f}. frondibus magis elongatis, segmentis paucioribus vix radiatis₅ apicibus fere omnibus integerrimis subulatis.
(TAB. NOSTR. 976).

Actiniopteris australis, *Link*, *FiL Sp. Hort. BeroL p. 80.*

Asplenium australe, *Sw. Syn. FiL p. 76. and 258. t. 3.*
fVzlld. Sp. PL 5. p. 308.

Acrostichum australe, *Linn. SuppL p. 444.*

Pteris australis, *Hook, et Grev. Ic. FiL t. 8.*

Acropteris australis, *F  e. Gen. FiL p. 76.*

Blechnum flabellatum, *Presl, Tent. Pterid. p. 103.*

HAB. Mauritius and Bourbon, *Sonneratj Carmichael*, and others. Schoata, near Enderder, Abyssinia, *Schimper*, n. 577.

It requires an extensive suite of specimens, such as perhaps our own Herbarium alone possesses, of this plant, to satisfy oneself that the state of the elegant Fern here figured and that in the preceding plate belong to one and the same species. The extreme forms known to us are given in the two plates now mentioned. At TAB. 975. f. 2. we have one extreme, at our TAB. 976. f. 4. the other extreme; but these are surely connected by the other examples on the same plates. We cannot agree with those Botanists who would retain so remarkable a plant as the present in *Asplenium*[^] nor yet with those who would unite it with the *Asplenium septentrionale*, (*Acropteris* of Link). The habit and texture and ramification are totally different.

TAB. 976. *Actiniopteris radiata*, /3.—Fig. 1. Abyssinian specimen from Dr. Schimner :—not. size. f. 2.. Segments of a fertile frond :—magnified, f. 3. Smaller portion :—more highly magnified, f. 4. Specimen from Bourbon, communicated by the Paris Museum, f. 5. Portion of a fertile segment:—magnified.

DCCCCLXXVIL

ASPLENIUM ADIANTOIDES, *Raoul*;
var. Richardi.

Frondibus oblongis oblongo-lanceolatisve flaccidis rarius rigidis pinnatis bipinnatisve, pinnulis oblongis obovatis rhombeisve basi cuneatis sessilibus v. inferioribus stipitatis varie inciso-lobatis, soris brevibus in pinnulis incisis marginibus lobulorum approximatis marginalibusve, in pinnulis subintegris margine remotis, rachi nuda v. sparse squamata, stipite squamoso basi dense paleacco.

Asplenium adiantoides, *Raoul*, *Choix des PL Nouv. Zel. Tab. i.*

Var. *Richard*[^] Hook, fil.; fronde spitham&a erecta subcoriacea bipinnata lanccolata v. linear-lanceolata, pinnis linear-lanceolatis eret-topatentibus, pinnulis approximatis*brevi-stipitatis sessilibusvc obovatis basi cuneatis varie inciso-dentatis lftbatis obtusis, soris margine approximatis, rachi nuda v. basi stipiteque sparse paleacea. TAB. NOSTR. 977.

HAB. (*var. Richardi*). New River in the Southern Island of New Zealand, *Hb. bead A. Richard*.

The fern here figured was given to us by our late friend Professor A. Richard, it formed part of a small collection made at the South extremity of New Zealand by the Captain of a French Whaler. Although we have seen no olhef specimens we have no hesitation in pronouncing it a state of the variable *A. adiantoides*, from the yar. *Colensoi*. of which (see Plate 984) it differs chiefly in the more coriaceous fronds and robust habit." In these respects it approaches small varieties of *AsplJlaccidum* and others of *A. bulbiferum*. It is impossible to convey to our readers by a limited number of plates, any idea of the amount of varieties that several of the New Zealand Ferns display. We have selected *A. adiantoides* as an example of a species, all the states of which are rare in herbaria, though most abundant in the cold damp woods of New Zealand. We have received very many specimens of it, and from many localities, and can truly say, both of this and of other species of Ferns, that with the materials, so do the difficulties of discriminating the species increase, and this to so great a degree, that we may confidently predict a great revolution in the whole Order, when complete collections of all states of the widely spread species are studied.—J.D. H.

Fig. 1. Portion of a pinna and pinnules :—magnified.

DCCCCLXXVIII.

ASPLENIUM (Allautodia) BROWNH, *Hook.*

Frondibus amplis glabris deltoideis membranaceis bipinnatis, pinnis superioribus oblongis pinnatifidis reliquis iterum pinnatis, pinnulis ovato-oblongis serrato-dentatis, involucris oblongis demum cylindraceo-inflatis tenuibus fragilibus, stipite elongato nudo, rachi flexuosa.

Allantodia australis, *Br. Prod. p.* 149.

Athyrium australe, *Presl, Tent. Pterid.* 98. *Fée, Gen. Fil. p.* 186.

HAB. Van Diemen's Land, *Brown, J. JD. Hoohei, R. Gunn, Esq.* New Zealand* Northern Island, *Rev. W. Colenso, Allan Cunningham.*

We believe this to be identical, as to generic structure, with the *Polyodium umbrosum*, of Hort. Kew. on which Mr. Brown founded his genus *Allantodia*; and the excellent Link and Kaulfuss unite with him in considering it distinct from *Asplenium*, "Involucro fornicato, e vena lateraliter orto, eaque utroque margine inserto, interiore dehiscente." "Presl and Fée, however, unite it with *Athyrium*, and that view is adopted in our Genera Filicum, TAB. XVI. Mr. Bauer's figure there given exactly represents the more mature son; but I have failed to detect the insertion of the two margins of the involucrum on the vein. In a young state the involucrum is nearly flat, and the outer margin appears to me to be free from any union with the vein. With regard to *Athyrium*, as it stands in Presl and Fée, it is made to contain some species with involucres so little inflated, that it would puzzle any one to say whether they should be referred to that genus or to *Asplenium.*, "ⁱ Le port et Phabitude generale" M. Fée observes "les distinguent des *Asplenium*, mais ils s'en rapprochent beaucoup par la nature de *Tindusium*, surtout lorsque ce tegument est court (abbreviatum)."

Fig. Fertile segment:—*magnified.*

DCCCCLXXIX.

TRICHOMANES COLENSOI, *Hook.Jil.*

Caudice gracili filiformi elongato, frondibus oblongis acuminate laxe pinnatis, pinnis subpinnatifidis laciniis brevibus linearibus angustis acutis erecto-patentibus integris vel incisis, involucris solitariis basin versus singular pinnse insertis infundbuliformibus stipitatis liberis, columella longissime exserta flexuosa.

Trichomanes Colensoi, *Hook.fil. mst.*

HAB. Interior of the Northern Islands, New Zealand, near Waikare Lake, *Rev. W. Colenso*, n. 104.

A very elegant, and as it appears to Mr. Colenso and to me, as well as to Dr. Hooker, an entirely new species. It is very slender in every part, in the caudex, in the rather short stipes, and in the ramifications of the frond. I find only one involucre near the superior base of each pinna, infundibuliform, stipitate, and remarkable for the very long flexuose receptacle or columella, seven or eight times longer than the involucre.

Mr. Colenso writes, "This species of *Trichomanes* was discovered on the deep sides of a dark ravine on the banks of a small rivulet, which meandered through the dense and ever humid forests of the mountainous region between Waikare Lake and Rua Tahuna, in the interior of the Northern Island of New Zealand, where it grows very profusely."

Fig. 1. Fertile pinna of a frond. /. 2. Involucre and base of the receptacle with capsules:—magnified.

DCCCCLXXX,

NOTHOCHLJENA DISTANS, *Br.*

Casspitosa, frondibus erectis rigidis subcoriaceis ferrugineo-hirsutissimis linear-i-oblongis bipinnatis, pinnis oppositis remotis brevibus subsessilibus erecto-patentibus deltoideo-ovatis obtusis, pinnulis ovato-oblongis inferioribus pinna-tifidis superioribus subintegerrimis confluentibus marginibus recurvis subinvolucratis, stipite rachi costisque subtus ferrugineo-paleaceis.

Nothochlsena distans, *Brown*, *Prod. Fl. Nov. UoLL* p. 146.
Labill Sert. Austr.-Caled. p. 5. t. 8. *Kunze₇ in Plant. Preiss.* 2. p. 110.

HAB. New Holland; Port Jackson* *Brown*. Logan River, and granite rocks in the Bathurst country, *Fraser*. Port Stephens, *Capt. P. King*. Swan River, *Preiss*, *Drummond*, 7L 8. and 666. New Caledonia, *Lahillardizre*. New Zealand, Northern Island, *Colenso*, n. 732. 1109. 1914., *J. D. Hooker*.

Kunze justly describes the figure of this species of *Nothochlcena* in Labillardiere as "mediocris," the magnified representation being especially erroneous. When a fertile pinna is carefully examined, the margin, pale and almost scariose, will be seen to be recurved and to represent an imperfect involucrum, indicating a near approach to some *Cheilanthes*; between which two genera we have already had occasion to observe that the line of distinction is not easy to be drawn. Judging from the copious specimens we have received from New Zealand, from the northern islands at least, the species seems to be of as frequent occurrence there as it is in New Holland; and it is now ascertained to have a very extensive range, from Swan River in the West to New Caledonia and New Zealand in the East.

Fig. 1. Upper side of a pinna. /. 2. Under side of a fertile ditto, f. 3. Portion of a sorus.—magnified.

DCCCCLXXXI.

TRICHOMANES SPRUCEANUM, *Hook.*

Frondibus difformibus rigido-membranaceis bipinnatifidis, lobis primariis oblongo-ovatis obtusis secundariis bi-trifidis, rachibus subtus longe hirsutis sterilibus brevi-stipitatis latissime ovatis lobis imbricatis, fertilibus longissime stipitatis contractis oblongis obtusis, involucris cylindraceis in loborum apices omnino immersis, columna inclusa, stipitibus alatis, caudice longe repente setaceo-squamoso.

HAB. San Gabriel-catingas, Rio Negro, a tributary of the Amazon? June 1852, *Spruce*, n. 2334.*

No one can fail to see a great similarity between the *Trichomanes heterophyllum*, H.J3.K., figured by Kunze, Fil. 2. t. 109. and this; but they are truly distinct: our plant is nowhere pinnate, and the lobes both of the sterile and fertile fronds are very different in form and much more divided; the involucres are different in shape, and the columella is here included. Still the two unquestionably belong to the same natural group, differing from what we call the subgenus *Hymenostachys*" (see Gen. et Sp- Fil. 1. p. 114), which has a reticulated venation, the involucres arranged in a simple distichous spike, and connate; and from the subgenus *Feea* (/. c.) which has the involucres also arranged in a simple distichous spike and the veins all free:—whereas in our plant and in TV. *heterophyllum*] Humb. the fertile frond is not converted into a spike, but is merely a contracted form of the common frond. In all, the fertile stipes is very much elongated (here beautifully winged), and the sterile one very short.

Fig. 1. Portion of a sterile frond. /. 2. Portion of a fertile ditto, f. 3. Involucre:—magnified.

DCCCCLXXXIL

TRICHOMANES BICORNE, *Hook.*

Caudice crassiusculo radicante, frondibus caespitosis rigidiusculis ovatis seu oblongo-ovatis tripinnatifidis subtus in costam longe sparse setosis, laciiniis oblongis integris vel bifidis, involucris copiosis terminalibus obovatis in sinibus bifidis bicornibus segmentorum profunde immersis, receptaculo columellaque longiuscule exsertis, rachi alte alata.

HAB. Barra do Eio Negro, on logs or roots of t*ees, in dense moist forests, Amazon, (n. 1178.) and at St. Gabriel, Bio Negro, frequent on decayed logs on the Catingas, R. Spruce, Esq., n. 2334.

This we cannot but consider an entirely new species, in some respects allied to *T. alatum*, Sw., yet considerably different. It is rather short, and firm and compact in its mode of growth, and varies from two to four or five inches high. The costa beneath is beset with dark strongish hairs or setae. But the remarkable feature is the involucre, which is sunk between the two horns as it were of the apex of a segment, and the apex of the segment and the sides of the mouth of the involucre, forming together, almost the letter V. Both the receptacle of the capsules and the columella are much exserted.

Fig. 1. Fertile portiori of a frond. /. 2. Involucr:magnified.

DCCCCLXXXIIIL

ASPLENIUM ADIANTOIDES, *RaOUh*

var. minus. Hook.jiL

Flaccidum parvuluni, frondibus ovatis ovato-lanceolatisve laxe bipinnatis rarius pinnatis, pinnulis paucis distantibus inferioribus saspius longe stipitatis rhombeis basi oblique cuneatis crenatis varic lobatisve glaberrimis, soris raargine remotis, rachibus gracilibus parce paleaceis glabratissive, stipite ba^t squamoso.

A. Hookerianum, *Colenso in Tasm. Phil. Journ.*

HAB. Northern and Middle Islands of New Zealand, Kerikeri River, *Cunningham*. Banks Peninsula, *Raoul*, Dr. *Lyal*. Bay of Plenty, East Coast and Interior; Rev. W. *Colenso*, n. 78. 295. 1941.

A very beautiful, but like all its congeners, an extremely variable fern, allied to the elegant *A. Magellanicum* of Fuëgia? A very much larger state, a foot long, of this species is figured in M. Raoul's fine work, "Choix de Plantes de la Nouvelle Zelande," TAB. 1. We have taken the opportunity of figuring two other states of this interesting plant in the present work, (Plate 984 and 977), premising that we have little doubt but that they will be raised to specific rank by some Pteridologists, and would have been by ourselves, had we not ample proof amongst our extensive suites of specimens from many parts of the Island of their all being fronds of one species, which is almost as protean as the well known *Asplenium Ilaccidum*, Forst. Mr. Cunningham's specimens are much smaller than these figured here, simply pinnate, with stalked, 3-lobed or three-partite, very broad pinnae, *J-J inch across.--J^r.D.H.

[^] Fig. 1. Pinnule of pinnate variety, and f. 2. Pinnule of bipinnate state of *A. adiantoides* var. *minus* i—both magnified.

Colensoance.

N. O. Filices.

DCCCCLXXXIV.

ASPLENIUM ADIANTOIDES, *Raoul*

var. Colensoi, Hook, fit

Parvulum suberectum flaccidum v. subcoriaceum, fronde lanceolata laxe bipinnata, pinnis subremotis erecto-patentibus, pinnulis approximatis obovatis inferioribus breve stipitatis varie inciso-lobatis, lobis furcatis integrisve subacutis, superioribus sessilibus inciso-dentatis, soris margine approximatis, rachi costaque glaberrimis, v. costa stipite'que sparce paleacea basi squamata.

A. Colensoij Col. in Tasm. Phil. Journ.

HAB. Northern Island, New Zealand, Waikare Lake, and Tuki tuki river, *Rev. W. Colenso, n. 270.*

For other states of this beautiful and variable little Fern we must refer to Plates 977 and 983 of this volume; the present is intermediate in size between the original *A. adiantoides* of Raoul (*Choix de Plantes de la Nouvelle Zelande TAB. 1*) and his *A. triste*, (which are undoubtedly varieties of one plant), and in the outline and incision of the fronds it is intermediate between our var. *minor* (t. 983) and var. *Richardi* (t. 977).—*J. D. H.*

Fig. 1. and 2. Pinnules :—magnified.

DCCCCLXXXV.

CYATHEA CUNNINGHAMI, *Hook, fil*

Rachi submuricata, fronde tripinnata flaccida, pinnulis sessili-
bus lineari-elongatis oblongisve obtusis crenato-v. pinna-
tifido-lobatis, involucris demum laceris v. oblique cyathi-
formibus, costa submuricata superne strigoso-pubescente
inferne tomentosa et paleacea v. glabrata.

HAB. New Zealand, *Forster*. Wycari river, Bay of Islands,
A. Cunningham. Mountains of the East Coast and Interior;
Rev. W. Colenso.

The finest specimens of this beautiful fern that we have seen are Cunningham's, in our friend Mr. Heward's herbarium: Forster⁷'s consists of a single pinna in the Hookerian herbarium, and Mr. Colenso's are also single pinnæ. As a species it is intermediate in character between *C. medullaris* and *C. Smithii* (Fl. Nov. Zealand, ined.) differing conspicuously from the former in the flaccid, membranous, pale green fronds, which are more or less strigose along the costa above, ⁿan & tomentose with mixed palea along both costa and partial rachis below: from *C. Smithii* it is distinguished by the long narrow pinnules. The involucre covers the sorus as a delicate membrane in its youngest state and then bursts irregularly all around, or becoming detached on one side only turns over as a shallow cup which finally becomes reflexed and appressed to the frond, exactly as in *C. Smithii* and in many species of *Hemitelm*: both forms of involucre occur on the same frond. Main rachis rough with minute scattered raised points. Native name "Punui," according to Mr. Colenso.—J.D.H.

Fig. 1. Portion of rachis and pinnule with reflexed in-
volucres, *f.* % Involucre and receptacle. */. 3.* Very young
frond and sori:—*all magr^fied*.

D C C C L X X V I .

T B I C H O M A N E S P E T E R S I I , A. Gray.

Pusillum, caudicibus filiformibus tomentosis intricato-caespitosis, frond. 13-, 6 lineas longis oblongo-lanceolatis ovato-oblongis v. minoribus late obovatis cuneatisve undulatis majoribus subpinnatifido-sinuatis obtusis glaberrimis (junioribus margine hinc inde pilis nigris 2-3 furcatis ciliatis) penninervia in stipitem gracilem attenuatis, venis furcatis v. pinnato-ramosis liberis (intramarginali nulla), soro solitario terminali, indusio immerso tubuloso-infundibuliformi, ore dilatato libero leviter bilabiato. A. Gray.

Trichomanes Petersii, Asa Gray, in Sillim. Am. Journ. Sc. and Arts, 2d. Ser. May, 1853. p. 326.

HAB. United States; Hancock County* Alabama, not far from Sipsey river, found only on the face of an insulate sandstone rock, within the reach of the spray of a waterfall, &c'. M. Peters, Esq.

"The fronds, as in *T. muscoides*, are very diverse in shape : the dilated cuneate ones might be taken for the sterile form; but I observe that, more frequently than the narrower fronds, they bear a terminal indusium, which is sterile and empty. There is a peculiarity about the venation, some of the branches of the primary veins being evanescent Ifowards their base, so as apparently to lie free and independent in the frond" (as shown in our figures). "The slender stipes is as long as the frond itself.—It belongs to Hooker's first section of true *Trichomanes*, and of his 2nd sub-division, which contains *T. muscoides*, *T. pusillum*, *T. erosum*, and *T. apodium*. Our new Fern is most nearly related to the first of them (chiefly a West Indian species), having the involucre equally immersed in the body of the frond, (which is not the case with the other species) and the receptacle very short. But the shape of the fronds, their slender stipes and the total absence of an intramarginal vein are abundantly distinctive characters. In the shape of the broader fronds, and in the stellate hairs which sparsely beset the margins, our plant may be likened to *T. reptans*; but that species has a close and flabellate venation, and a cylindrical, exserted involucre, with a deeply two-lipped orifice.*"

In the above Dr. Asa Gray has left nothing for us to notice, save that the cellular structure of the frond is exceedingly compact, constituting very minute areoles.

Fig. 1. Fronds magnified; and f. 2, portion of a fertile frond: more highly magnified.

DCCCCLXXXVIL

POLYPODIUM (*Drynaria*) CONTIGUUM, *Wall*

Frondibus elongatis linear-lanceolatis acuminatis marginibus revolutis inferne in stipitem brevem attenuatis subcoriaceis glabris nudis rigidis reticulatim venosis, venis internis obscuris, areolis elongatis subhexagonoideis, yenis ultimis liberis divaricatis apice clavatis, soris marginalibus oblongis seriatim dispositis contiguis immersis, capsulis longe stipitatis squamulis stipitatis immixtis, caudice repente squamoso.

Polyodium contiguum, Wall. Cat. n. 285.

Drynaria revoluta, J. Smith, JEnum. Fil. JPhilipp. in Hook. Jburn. qffBot. 3. p. 491.

HAB. Kamoun, Wallich. Mergui, Griffith. Luzon, ^mCuming, Fil. Phillipm. n. 247. Singapore, Low. China, Fortune*. n. 21.

The natural affinity of this plant is with the *Pleopeltis ensifoliq* Carm. (Hook. Exot. Bot. 1. 62). and still more with *Pleopellis nuda*, Hook. Exot. Bot. 1. 63, now generally placed in *Drynaria* (Phymatodes). §*Lepisorus*, J, Sm, and we believe that the venation of these will be found sufficiently to correspond : specifically, however, our present plant differs, not only in the great length of the fronds and the marginal situation of the sori, but further in the shape of the sori, approaching to oval indeed in *Pleop. nuda*, here invariably oblong, so as properly to belong to the tribe *Grammitacece* of Presl, the receptacle of the sorus being in reality linear. In that groupe I would willingly have ranked it, but that I know of no genus there in which it could with propriety be placed, especially if the venation is considered. The difficulty would be readily solved with some botanists, by making a new genus of it; but we think it of much more importance to illustrate structure with the view to the future consideration of genera, than hastily to form a new genus, which, on further investigation, may be found unsound.

Fig. 1. Portion of a fertile frond, seen from beneath. /. 2. Capsule. /. 3. Peltate stipitate scale from among the capsules :—magnified.

Wrightiana.

N. O. Filices.

DCCCCLXXXVIII.

ANEMIA (Euanemia) MEXICANA, *Klotzsch.*

Elata, frondibus (sterilibus) pinnatis, pinnis. sub-13 ovato-acuminatis basi obtusis inasqualibus nunc auriculatis paten-tibus serratis brevi-petiolatis glabris penniveniis venuliis dichotomis liberis hinc auriculatis, spicis longe petiolatis pinnatis, pinnis pinnatifidis apice confluentibus (nunc omni-bus confluentibus) lobis oblongis capsuliferis, caudice repente stipitique basi squamoso-hispidis.

Anemia Mexicana, *Klotzsch*, in *Linncea*. 18. p. 526. *Kunze* in *Schkuhrj Fil. Supph* 2. p. 75. t. 131. et in *Linncea*, 23. p. 223.

A. speciosum, *JPresl, Supph JPterid.* p. 89.

HAB. Western Mexico, Née, Aschenborn. Texas, *Lindheimer* in *Herb. Nostr.* n. 572. Between Western Texas and El Paso, New Mexico, C. Wright, n. 826,

Our specimens are from a foot to a foot and a half long, with so much of the habit of *A. Phyttitudis*, that a casual inspection only would induce many Botanists to refer it to that species; but a more careful examination will exhibit a difference in the broad and obtuse base of the pinnae, and a more remarkable one in the pinnae being here furnished with dichotomous free veins, whereas in *A. phylltidis* they are reticulated, as in *A. Tioeediana* (Tab. 906.)

Texas is the most northern limit we are yet acquainted with of this genus.

Fig. 1. Portion of a sterile pinna. */. 2.* Segment of a fertile spike with capsules, seen from beneath:—*magnified** *f. 3.* Variety or monstrosity of a fertile spike, seen from above (from Texas, *Lindheimer*):—*not size.*

Griffithiana.

N. O. Filices.

DCCCCLXXXIX.

GYMNOGRAMME AURITA, *Hook.*

(Vide TAK. 974).

/3. frondibqp bipinnatis, pinnis magis elongatis, venulis furcatis
ramis ambobus soriferis.

HAB. Khasya; Mumbree, and Nuncklow, Eastern Bengal,
Griffith.

This is a variety to which we alluded in our description under Tab. 974, and which at first sight has the appearance of being different* from that species, an opinion that some wouM consider confirmed by the venules there being • simple, here forked, and each branch soriferous, so that there are as it were twin sori:—but after our figure was engraved a more careful examination has led to the conviction that the two are but varieties of one and the same species, of which intermediate grades exist in our extensive suites of specimens. Capsules hispid.

Fig. 1. Fertile segment, seen from beneath, *f. 2.* Capsule:
—*magnified* *

Wallichiana.

N. O. Filices.

*
DCCCCXC.

MENISCIUM SALICIFOLIUM, *Wall.*

Elatum glaberrimum coriaceo-submembranacemn, fronde pinnata, pinnis brevi-petiolatis linear-lanceolatis alternis tenui-acuminatis basi attenuatis, costa subtus pallida venisque prominentibus, stipite (spithameo et ultra) costaque nudis.

Meniscium salicifolium, Wall. Cat n. 63.

HAB. Penang, 1822, Dr. Wallich, Sir William Norris. Singapore, Thos. Lobb.

A fern, so far as I know, peculiar to the two localities above mentioned. It is remarkable for the narrow and very tapering willow-like pinnae of the frond, certainly in the specific character as given by Willdenow, of *M. angustifolium*, ("frondibus pinnatis pinnis linear-lanceolatis acuminatis integer-riinis, basi angustatis,") a near ally of that species; but that is a native of South America, from Caracas, and the whole height of the frond is described to be nine inches, the pinnae three inches long, the stipes 2*3 inches: whereas our plant is 2-3 feet long, the pinna 8-9 inches in length, the stipes more than nine inches. Each transverse veinlet bears two oval sori which soon become confluent.

Fig. 1. Portion of a fertile pinnule:—magnified.

DCCCCXCI.

POLYPODIUM ERIOPHOBUM, *Hook.*

Frondibus cordatis profunde 3-lobis utrinque molliter lanosis,
lobis lateralibus semicordatis pinnatifidis lobulis infimis
sinuato-lobatis, intermedio triangulari pinnatifido, venulis
bifidis apicibus soriferis, soris marginalibus, stipitibus fronde
3-4-plo longioribus lanosis rachibusque nigris, caudice elon-
gato repente nigro-squamoso.

Nothochlsea criophora, *Fée*, *Gen. Fit. p. 159. t. 13. y.^3.*

HAB. Shad\$ clefts, near the city of Oeiras, Brazil, *Gardner*•,
n. 2390.

M. F6e has examined and figured and described a specimen of this plant; but he has arrived at a different conclusion from myself in regard to the genus. To me it appears to be a true *Polypodium*, bearing as it does globose naked sori at the extremity of the forked veinlets, near the margin it is true, but not "continui, et limbum angustum constituentes."* li is clear, however, from his figure above quoted, that M. F6e had only a small and imperfectly developed frond before him. He describes it indeed, "sporangiis remotis," and figures only one or two capsules at the extremity of each veinlet; whereas the capsules in our specimens are very copious from each receptacle, forming approximate sori, but not a continuous line, so copious in the older specimens, so much extended over the disk among the copious, lax, woolly hairs, (partially concealing them), that our lamented friend Mr. Gardner distributed his specimens under the name of *Hemionitidis* sp. The wool is of a tawny hue.

Fig. 1. Upper side of a lobe, atid f. 2. Under side of a fertile lobe, the wool having been removed :—magnified.

Schomburgkianæ.

N. O. Filices.

DCCCCXII.

GYMNOGRAMME SCHOMBURGKIANA, *Kze.*

Parva fasciculata, frondibus oblongis obtusiusculis chartaccis nitidis bipinnatis inferne subtripinnatis, pinnulis plerisque obovato-cuneatis ovalibusve inciso-lobatis convexis, sterilibus pilis paucissimis solitariis in venas, fertilibus subtus in venas copiose pilis longis hispidis, veriulis dichotomis apicibus clavatis, soris oblongis laxis, stipite clongito rachique primario cbencis nitidis, rachibus secundariis alatis, caudice nodoso parvo.

Gymnogramme Schomburgkiana, *Kunze*, (*Klotzsch in Herb. Nostr. n. 1196*).

Anogramme Schomburgkiana, *Fée*, *Gen. Fil. p. 184.* (*name only*).

* This pretty species is only known to me by the specimens here figured, kindly sent by Dr. Klotzsch. M, Fée quotes Kunze MSS, as his authority for the name, and I presume it has not been yet anywhere published. The whole plant is very glossy: the texture is firm, and in a dry state at least the anterior surface of the pinnae is convex. Only a very few scattered solitary hairs are seen on the sterile frond, but the underside of the fertile ones is copiously hispid with long hairs on the veins beneath and among the capsules. The veinlets all terminate considerably within the margin and are there clavate.

Fig. 1. Sterile pinnae, seen from above. /. 2. Fertile pinnae, seen from beneath:—magnified.

Lobbiana.

N. O. Filices.

DCCCCXCIII.

LASTREA BORNEENSIS, *Hook.*

Fronde pinnato-pinnatifida (an bipinnata?) acuminata membranacea glabra, pinnis lanceolatis inferioribus longe angustè acuminatis profunde pinnatifidis, lobis oblongis acutis[^] pinnatifido-serratis, lobulis serraturisve obtusiusculis, soris in medio venularum, involucris reniformi-orbicularibus venam centralem spectantibus, rachibus costisque pubescentibus.

HAB. Borneo, near Sarawak, *Thos. Lobb.*

I possess only a portion, it would appear, of this frond, at any rate no stipes nor caudex; and in general form this has affinity with many species of this extensive genus. There* is however, a peculiarity in the direction of the involucres, which strikes me as being unusual. The veinlet on which the involucrum is placed forms an angle at or near the middle, and from that angle the sorus and involucrum appear to spring, and the latter is not directed towards the apex of the lobule following: the direction of the veinlet, but is free from it and has an oblique direction towards the main or central vein.

Fig. 1, Under side of a lobe with sori. /. 2. Single sorus: —magnified.

Lobbiance.

N. O. Filices.

DCCCCXCIV.

T\ffinITIS 0BTU8A, *Hook.*

Glabra coriacea nitida, frondibus simplicibus oblongo-ovatis obtusis brevissime mucronatis, soris intra marginem in-
crassatum et costam continuis (infra apicem. etiam) vel
interruptis, stipite gracili filiformi basi piloso, caudice
repente crinito.

HAB. .Borneo, near Sarawak, *Mr. Thos. Lobb.*

The genus *Tcenitis* has been of late properly considered to be confined to one species, *T. blechnitis*, and that is a pinnated species. Our present plant is a true *Tcenitis*, with a simple frond, and quite entire at the margin, except from the effect of injury. In all our fronds too, where the apex is uninjured, the two lines of sori are united or continuous within the apex, the only portion invariably free from sorus being at the very base. The plant is peculiarly rigid and coriaceous, glossy, the margin thickened. The caudex is creeping, but rather slender, and clothed with dark brown, glossy, crinite scales.

Fig. 1. Portion of a fertile frond: —magnified.

Lobbiance.

N. 0. Ophioglossaceae.

DCCCCXCV.

OPHIOGLOSSUM INTERMEDIUM, *Hook.*

Frondibus erectis membranaceis reticulatis elongato-linearispathulatis ift petiolum longiusculum sensim attenuatis indivisis, pedunculo spicam aequante apicem versus frondis medio affixo.

HAB. Borneo, near Sarawak, Mr. Titos. *Lobb.*

My only specimen of this *Ophioglossum* is that here represented, where three fronds arise from one small but somewhat tuberous root, and the plant seems to be terrestrial and to grow erect. Were it an epiphyte and a pendent" frond, the species might be supposed to be a form of the *O. pendulum* (sec Hook, et Grev. Ic. FiL Tab. 19); but here is no disposition in the frond to be dichotomous; and the narrow stipes of our present plant, the elongated peduncle and the situation of the peduncle (itself as long as the spike) forbid such a supposition. The membranaceous texture is in favour of its being the same as *O. pendulum*; and it may be considered to hold an intermediate rank, between the terrestrial erect and the epiphytal pendent species.

Fig. 1. Apex of a frond. /. 2: Portion of a spike;—magnified.

Lobbianc.

N. O. Filices.

DCCCCXCVL

DRYMOGLOSSUM RIGIDUM, *Hook.*

Glaberrimum, frondibus sterilibus obovatis in stipitem sequi-longum attenuatis costatis marginibus incrassatis nitidis, fertilibus longissimis linearibus crassis, stipite subseque longis, sors profunde immersis, caudice repente elongato squamoso.

HAB. Borneo, near Sarawak, *Mr. Thomas Lobb.*

A species of *Drymoglossum* extremely different from any hitherto described, remarkable for its thick, firm, coriaceous shining, obovate sterile fronds, incrassated at the margin, tapering into a rigid stipes as long as the fronds. The fertile frond, including the stipes, exceeds by many times the length of the sterile fronds with their stipites, and is also peculiar in being very narrow and very thick (almost semiterete), having a deep furrow on each side the costa, in which the sori are sunk. The veins anastomose, but the areoles are not appendiculated.

Fig. 1. Portion of a sterile frond, f. 2. Portion of a fertile frond; magnified.

DCCCCXCVIL

TRICHOMANES PLUMA, *Hook.*

Hispidulo-scabra, frondibus lanceolatis sublongo-stipitatis nigris decomposito-pinnatifidis, laciniis teretibus filiformi-subulatis confervoideis reticulatis, involucris versus apicem frondis prrecipue sitis parvis cyathiformibus in ramulis brevibus terminalibus, receptaculo longissimc exerto, soris inclusis, caudicc repente setaceo-squamoso.

HAB. Borneo, near Sarawak, Mr. Thomas Lobb.

This is only one of the many extremely interesting plants in a collection, for which I am indebted to Mr. Veitch of the Exotic Nurseries, Exeter and Chelsea, made in Borneo by Mr. Thomas Lobb. The first aspect of this plant is quite that of a *Sertularia*[^] especially of the well known *Sertularia Pluma*. A more close inspection will show it to be a true *TrichomaneS*) with the ramification so fine that when, magnified it resembles the branches of some confervoid marine Alga, particularly the Genus *Ceramium*: the branches are minutely reticulated and here and there beset with short bristles or rigid hairs : the pinnae or segments (for it is difficult to say whether this ramification should be called pinnate or pinnatifid) do not spread in two opposite directions (distichous), but stand out as it were, on all sides so as to form a thickened mass, in that respect also resembling the *Sertularia* in question and certain *Ceramia*. The involucres are particularly small for the size of the plant, and the columella or receptacle is unusually long and stout.

Fig. 1. Portion of a fertile tyranch. f. 2 Involucre and receptacle:—magnified.*

Lobbiance.

N. O. Filices.

DCCCCXCVIIL

GRAMMITIS BISULCATA, *Hook.*

Frondibus sparsis anguste linear-i-acuminatis curvatis rigidis
infeine in ptiolum attenuatis subscmiterctibus enerviis
antice profunde bisulcatis sulcis sorifcris,, soris ovali-
oblongis prominentibus, caudice clongato filiformi repcntc
squamoso.

HAB. Borneo; near Sarawak, TJios. *Lobb.*

There is no appearance of venation in the harsh, rigid fronds of this Fern. A transverse section (fig. 1. and 2.) shows them to be nearly scmiterete; the flatter side is marked with two deep furrows in which are situated the sori, the convex side has three shallow grooves. The caudex is long and creeping, everywhere clothed with scales, throwing out roots from below, and bearing the distinctly placed fronds on the upper side.

Fig. 1. 2. Transverse sections of a fertile frond:—magnified.

DCCCCXCIX.

HYMENOLEPIS PLATYRHYNCHOS, *Kze.*

Frondibus cacspitosis elongatis sublorato-lanccolatis coriaceo-membranaceis costatis basi in pctiolum brevem attenuatis, appendice terminali fructifera oblonga plana, soro oblongo amplo elcvato discum occupante, caudice crasso descendente squamoso.

*Hymenolepis platyrhynchos. Kze. in Schkh. Fil. SuppLp. 102.
Gymnopteris platyrhynchus, J. Smith, in Hook. Journ. Bot. 3.
p. 403.*

*Macroplethus platyrhynchus, Presl, JEpmel. Bot p. 142.
HAB. Luzon*, Cuminff, n. 196.*

This fine and apparently very rare Fern (for hitherto it has been found only by Mr. Cuming) was first named by Mr. J. Smith *Gymnopteris platyrhynchus*[^] a genus from which it differs widely in habit, in the monomorphous fronds, and in the limited mass of fructification. Kunze has, as it appears to me with more propriety, referred it to *Ilymenolepis*, which indeed Presl united with *Gymnopteris* in his "Tentameif Ptcridographiac," but which it may be presumed he now retains, since in his remarks upon *Macroplethus*, a genus destined for our present plant, he says, "^{ei} differt ab *Ilymenolepide*, quacum venis venulisque accedit, soro sub apice frondis centrali solitario elliptico, margine frondis piano undique cincto, receptaculo elliptico non solum parenchymati sed quoque costae insidenti." Even M. F6e cannot assent to these views, and he arranges the species with *Ilymenolepis*.

Fig. 1. Sterile portion of the frond, showing the venation.—magnified.

Mooreance.

N, O. Filices.

M.


ASPLENIUM (Darea) OBTUSILOBUM, *Hook.*

Parvum caespitosum sparsim stellato-pilosum, frondibus erectis ovato-lanceolatis pinnatis, pinnis petiolatis bipinnatifidie lobis linear-i-cuneatis obtusis integris vel bifidis costatis (seu univeniis), vena ante apicem evanescente, involucris solitariis omnino marginalibus exterius dehiscentibus, stipite rachique compressis alatis, radice fibrosa stolonifera.

HAB. On the ground in shady places. Island of Tanna, New Hebrides, Mr. C Moore.

The scattered, branched, stellated hairs of this little plant are invariably confined to the rachis and veins of the frond. As a species it appears very distinct.


Fig. 1. Pinna, with sorl f. 2. Single sorus:—magnified.









K9V4




















Engr. del et lith.


Cheilanthes vidula -

Burm. nom.


Pinna L.















MOM
















Etch. del et lith.

Baumgarten Augs.


Nephrolepis Rinconianum























2



3



2

1



rlois <W st II - k.


siflag. cadipluma


1 hair $\frac{1}{10}$ in mm


Singapore





























Flisch. del. et lit.


Ctenopteris Mayorum

n. we.




Such little bkh.

Synecodon fm.
Cone name Khan aridides
Pamplin amp




Fitch del et lith.
(Plumbeum Sm.) Guttiferae
Kharia Polyph. Organica




Etch. J. L. et lith.

Polyphoides
sepal. latifolius -
Khaem

Tenglin map






Scot. Nat. Hist.


Scot. Nat. Hist.

Pteridium aquilinum

r











2

// SSS



























Fitch del. & engr.

Lycopodium camineoides

Papilla 135

(Uj)



Vl&W



Fitch del. et lith.

Ethn itnp.

Ft. 371,972.




ft 974.



Pitch del et lith.

Pampin imp.


Synopteris aurita - Kuhn




Fitch del et lith.



Bentham imp.


Actinopteris radiata





















Fisch. del. et. lith.


M/J- adiantides


Pumpkin imp.














Fitch del. et lith.

Pumpkin Imp

A monogramme austral-

Kumbas

TL990.



Pamplin imp.

Fitch dn! .<< lith.

Mercurialis sclerophylla

